

INTERMEDIATES AND METHODS FOR THE SYNTHESIS OF HALICHONDRIN B ANALOGS

Related Applications

- 5 This application claims priority to U.S.S.N. 60/997,625, filed October 3, 2007, the entire contents of which are hereby incorporated herein by reference.

Background of the Invention

- 10 The invention relates to a method for the synthesis of halichondrin B and analogs thereof having pharmaceutical activity, such as anticancer or antimitotic (mitosis-blocking) activity. B-1939 (also known as E7389 or eribulin), a halichondrin B analog, has been reported to be useful for treating cancer and other proliferative disorders including melanoma, fibrosarcoma, leukemia, colon carcinoma, ovarian carcinoma, breast carcinoma, osteosarcoma, prostate carcinoma, lung carcinoma, and ras-
15 transformed fibroblasts.


- Halichondrin B is a structurally complex marine natural product containing multiple chiral centers on an extended carbon framework. Due to the limited availability of halichondrin B from natural sources, methods for the synthesis of halichondrin B have value for the purposes of developing the full medicinal potential of halichondrin B
20 analog. A method for the synthesis of halichondrin B analogs was published in 1992 (Aicher, T. D. et al., *J. Am. Chem. Soc.* 114:3162-3164). A method for the synthesis of Halichondrin B analogs, including B-1939, was described in WO 2005/118565 (EISAI COMPANY, LTD.). The method described in WO 2005/118565 has several practical advantages over the method disclosed by Aicher, including but not limited to the
25 discovery of several crystalline intermediates which enabled increased quality control, reproducibility, and throughput. Notwithstanding these advantages, several throughput limiting chromatographic purifications remained particularly relating to the C14-C26 fragment. For example, the C14-C26 fragment contains 4 chiral centers at C17, C20, C23, and C25 which require chromatography to control the quality of this fragment.
30 More specifically, installment of the C25 chiral center does not occur with high selectivity and could not be practically enhanced due to a lack of crystalline intermediates late in the C14-C26 synthesis.

- What is needed is a more efficient, less costly, more practical method for the synthesis of halichondrin B analogs, in particular B-1939.
35

Summary

The current disclosure relates to a method for the synthesis of Halichondrin B analogs, such as B-1939, from (-)-quinic acid according to the process illustrated in Scheme 1, below. The method introduces a number of new and crystalline intermediates which greatly improve the stereochemical quality of the compounds synthesized and reduces the need for chromatographic steps. Unlike the previously described methods, the presently claimed method is substantially more appropriate for pharmaceutical manufacturing.

The disclosure also pertains to the novel intermediates disclosed herein.



Scheme 1

WO 2005/118565 disclosed a method for making Halichondrin B analogs, such as B-1939, that included synthetic routes for (1) producing the compound of formula Ia from (-)-quinic acid, and for (2) producing the B-1939 from Compound AG. Both synthetic routes are suitable for use in the method of the present invention, and are
5 incorporated by reference herein.

The method of the present disclosure differs from the method disclosed in WO 2005/118565 in the process of synthesizing Compound AH from Compound AA. In particular, the present disclosure discloses highly efficient methods for generating the C25 chiral center, marked with an asterisk(*) in the relevant compounds in Scheme 1,
10 by a process of equilibrating and selectively crystallizing the desired C25 isomer *via* an alpha-methylated nitrile. In the method described in WO 2005/118565, Compound AH is synthesized by adding a methyl group to Compound AG, as shown above. This reaction generates the C25 chiral center.. The product of that reaction is a mixture of diastereomers with each possible configuration around that chiral center.
15 Chromatography can be used to partially isolate Compound AH from the mixture of diastereomers, as disclosed in WO2005/118565; however, the remaining diastereomers of Compound AH result in undesired impurities in subsequent reaction steps, impurities which can only be removed through additional purification procedures.

Unlike the methods of syntheses of halichondrin B analogs previously described,
20 the method of the present disclosure involves the formation of the C25 chiral center at an earlier stage in the synthesis of Compound AH. Several of the methylated intermediates, including Compound AD and Compound AF are crystallizable. By crystallizing one or more of the methylated intermediates in accordance with the methods of the present disclosure, one can produce a composition of comprising
25 Compound AH that is substantially diastereomerically pure. For example, Compound AC can be methylated to produce Compound AD. When Compound AD is produced, a the C25 chiral center is produced, the same chiral center discussed with respect to Compound AH. When this reaction occurs, a diastereomeric mixture is produced with each possible stereomeric configuration around that chiral center. Although the
30 methylation itself occurs with low stereoselectivity, surprisingly, the desired diastereomer of Compound AD stereoselectively crystallizes. Moreover, the undesired C25 stereoisomer can be epimerized under conditions from which the desired C25 stereoisomer crystallizes. Thus, the yield and quality of the C25 stereoisomer can be enhanced by crystallization induced dynamic resolution (CIDR).


35 Several other intermediates produced in the synthetic route from Compound AD to Compound AH can also be crystallized from reaction mixtures, resulting in an even higher purity composition of Compound AH than could be produced by previously

disclosed methods. In particular, Compound AF is a crystalline compound, while the corresponding non-methylated Compound AE requires chromatography for purification. Compound AF may be synthesized from Compound AD or it can be synthesized by methylating Compound AE.

5 Removal of chromatography steps from the processes used to synthesize halichondrin B analogs dramatically increases the product yield and reproducibility, while decreasing cost and production time. The present method also enables one to resolve difficult to resolve chiral centers at a considerably earlier points in the process, even as early as the production of Compound AH and Compound AI. B-1939 is
 10 suitably synthesized from Compound AI using methods such as those described in WO/2005/118565.

In an aspect, there is provided a method of obtaining a substantially diastereomerically pure composition of a compound of formula (I), comprising:

15 crystallizing said compound of formula (I) from a mixture of diastereomers under appropriate crystallization conditions, wherein said compound of formula (I) is:



wherein:

z is a single or double bond, provided that when z is a double bond, X² is C and
 20 Y¹ is hydrogen; and provided that when z is a single bond, X² is CH or O;


X¹ is O, S, or CN, provided that when X¹ is CN or S, X² is O;

Y¹ is a halide, hydrogen or O-L², or absent when X² is O;

L¹ and L² are independently selected from hydrogen and a protecting group, or
 L¹ and L² together are a protecting group, provided that when X¹ is CN, L¹ is absent; or
 25 a salt thereof,

wherein the protecting group of L¹ and L² is silyl, C₁-C₁₂ alkylcarbonyl, C₁-C₆ alkyl, aryl (C₁-C₆) alkyl, carbonate, C₁-C₆ alkoxy-(C₁-C₆) alkyl, 3,4-dimethoxybenzyl, p-methoxybenzyl, benzoyl, or C₁-C₆ ester, or L¹ and L² together are a protecting group selected from benzylidene, cyclic C₁-C₆ acetal, cyclic C₃-C₆ ketal, cyclic carbonate, and
 30 a C₄-C₇ alkyl ring with one member of the ring covalently linked to both the O of Y¹ and to X¹; or when X¹ and X² are O, L¹ and L² together are cyclic C₃-C₇ ketal.


In one embodiment, the disclosure pertains, at least in part, to a method of obtaining a substantially diastereomerically pure composition comprising a compound of formula (I). The method includes crystallizing the compound of formula (I) from a mixture of diastereomers under appropriate crystallization conditions, such that a substantially diastereomerically pure composition comprising a compound of formula (I) is formed. The compound of formula (I) is:



wherein:


- z is a single or double bond, provided that when z is a double bond, X² is C and Y¹ is hydrogen; and provided that when z is single bond, X² is CH or O;
 - X¹ is O, S, or CN, provided that when X¹ is CN or S, X² is O;
 - Y¹ is a halide, hydrogen or O-L², or absent when X² is O; and
 - L¹ and L² are independently selected from hydrogen and a protecting group, or L¹ and L² together are a protecting group, provided that when X¹ is CN, L¹ is absent;
- and salts thereof. The disclosure also pertains to compositions of compounds of formula (I) that are substantially free of diastereomers, as well as compounds of formula (I).

In another embodiment, the disclosure also pertains to a method of making a diastereomerically pure composition of a compound of formula (Ib) from a compound of formula (Ia), wherein the compound of formula (Ia) is:



(Ia)

and the compound of formula (Ib) is:




(Ib)

- 5 wherein L^{1a} and L^{1b} are independently selected from hydrogen and a protecting group, or L^{1a} and L^{1b} together are a divalent protecting group, provided that L^{1a} of formulae (Ia) and (Ib) are the same and L^{1b} of formulae (Ia) and (Ib) are the same. When L^{1a} or L^{1b} is a protecting group, it is preferably selected from the group consisting of C_1 - C_6 alkyl ethers, aryl (C_1 - C_6) alkyl ethers, silyl (C_1 - C_{10}) ethers, C_1 - C_6 alkyl esters, cyclic C_1 - C_6 acetals, cyclic C_2 - C_7 ketals, and cyclic carbonates. The method includes reacting the compound of formula (Ia) under alkylating conditions to form a mixture comprising the compound of formula (Ib) and diastereomers thereof; and crystallizing the compound of formula (Ib) from the mixture, under appropriate crystallization conditions.


15

In another aspect, there is provided a method of making a substantially diastereomerically pure composition of a compound of formula (Ib) from a compound of formula (Ia), wherein the compound of formula (Ia) is:



(Ia)

- 20 and the compound of formula (Ib) is:



(Ib)


wherein L^{1a} and L^{1b} are independently selected from hydrogen and a protecting group selected from C_1 - C_6 alkyl, aryl (C_1 - C_6) alkyl, silyl (C_1 - C_{10}), and C_1 - C_6 alkyl ester,

or L^{1a} and L^{1b} together are a divalent protecting group selected from cyclic C_1 - C_6 acetal, cyclic C_3 - C_7 ketal, and cyclic carbonate, provided that L^{1a} of formulae (Ia) and (Ib) are the same and L^{1b} of formulae (Ia) and (Ib) are the same, said method comprising:

- reacting the compound of formula (Ia) under alkylating conditions to form a
- 5 mixture comprising the compound of formula (Ib) and diastereomers thereof; and
- crystallizing the compound of formula (Ib) from the mixture, under appropriate crystallization conditions.

In another embodiment, the disclosure pertains, at least in part, to a method of

- 10 obtaining a substantially diastereomerically pure composition comprising a compound of formula (II). The method includes crystallizing the compound of formula (II) from a mixture of diastereomers under second appropriate crystallization conditions, such that a substantially diastereomerically pure composition comprising a compound of formula (II) is formed. The compound of formula (II) is:



(II)


wherein:

- c is a single or double bond, provided that when c is a double bond m is 0 and Y^3 is O or $CHCO_2-L^3$, and provided that when c is a single bond m is 0 or 1 and Y^3 is CH_2O-L^3 , $CH_2CO_2-L^3$ or $CH_2CH_2O-L^3$;
- 20 Y^2 is C_1 - C_7 sulfonate, $O-L^4$ or a halide;
- L^4 is hydrogen or a protecting group; and

L^3 and L^5 are each independently hydrogen or a protecting group, or L^3 and L^5 together are a protecting group, or a salt thereof. The disclosure also pertains to compositions of compounds of formula (II) that are substantially free of diastereomers, as well as compounds of formula (II).

5

In yet another embodiment, the disclosure also pertains to compounds of formula (III):




(III)

wherein: L^6 is hydrogen or a protecting group; and salts thereof.

10

In yet another embodiment, the disclosure also pertains to a composition comprising a compound of formula (IIIa):



(IIIa)


15 L^{6a} , L^{6b} , and L^{6c} are each protecting groups, or a salt thereof, and wherein the composition is substantially free of diastereomers.

Furthermore, the disclosure also pertains to a composition comprising a compound selected from the group consisting of formula (I), (Ia), (Ib), (Ic), (Id), (Ie), (II), (IIa), (IIb), (III) and (IIIa). The disclosure also pertains to each of the compounds

20

described in herein.

In another aspect, there is provided a compound of formula (I):



(I)

wherein:

z is a single or double bond, provided that when z is a double bond, X^2 is C and Y^1 is hydrogen; and provided that when z is single bond, X^2 is CH or O;

X^1 is O, S, or CN, provided that when X^1 is CN or S, X^2 is O;


5 Y^1 is a halide, hydrogen or $O-L^2$, or absent when X^2 is O; and

L^1 and L^2 are independently selected from hydrogen and a protecting group, or L^1 and L^2 together are a protecting group, provided that when X^1 is CN, L^1 is absent; or a salt thereof,

wherein the protecting group of L^1 and L^2 is silyl, C_1 - C_{12} alkylcarbonyl, C_1 - C_6 alkyl, aryl (C_1 - C_6) alkyl, carbonate, C_1 - C_6 alkoxy-(C_1 - C_6) alkyl, 3,4-dimethoxybenzyl, p-methoxybenzyl, benzoyl, or C_1 - C_6 ester, or L^1 and L^2 together are a protecting group selected from benzylidene, cyclic C_1 - C_6 acetal, cyclic C_3 - C_6 ketal, cyclic carbonate, and a C_4 - C_7 alkyl ring with one member of the ring covalently linked to both the O of Y^1 and to X^1 ; or when X^1 and X^2 are O, L^1 and L^2 together are cyclic C_3 - C_7 ketal.

15


In another aspect, there is provided a compound, wherein said compound is:



Compound AF.


20

In another aspect, there is provided a method for producing compound B-1939



which method comprises:

(i) reducing a compound of formula 8-20:




8-20

wherein:

Y^2 is a halide; and

5 L^5 is C_1 - C_{10} alkyl, C_4 - C_{10} aryl- C_1 - C_6 alkyl, or C_4 - C_{10} aryl


and converting the product to a compound of formula (III):



(III)

wherein: L^6 is hydrogen or silyl C_1 - C_{10} ; or a salt thereof;

(ii) converting the compound of formula (III) or salt thereof to produce compound AV




Compound AV

and converting compound AV

10

into compound AI




Compound AI

; and


(iii) converting compound AI into compound B-1939.

In another aspect, there is provided a compound having the formula:



Compound AU.


In another aspect, there is provided a compound having the formula:



5

Compound AR.

In another aspect, there is provided a composition comprising a compound having the formula:



10

wherein:

z is a single or double bond, provided that when z is a double bond, X^2 is C and Y^1 is hydrogen; and provided that when z is a single bond, X^2 is CH or O;

X^1 is O, S, or CN, provided that when X^1 is CN or S, X^2 is O;

Y^1 is a halide, hydrogen or $O-L^2$, or absent when X^2 is O;

15

L^1 and L^2 are independently selected from hydrogen and a protecting group, or L^1 and L^2 together are a protecting group, provided that when X^1 is CN, L^1 is absent; or a salt thereof,

wherein the protecting group of L^1 and L^2 is silyl, C_1 - C_{12} alkylcarbonyl, C_1 - C_6 alkyl, aryl (C_1 - C_6) alkyl, carbonate, C_1 - C_6 alkoxy-(C_1 - C_6) alkyl, 3,4-dimethoxybenzyl, p-methoxybenzyl, benzoyl, or C_1 - C_6 ester, or L^1 and L^2 together are a protecting group selected from benzylidene, cyclic C_1 - C_6 acetal, cyclic C_3 - C_6 ketal, cyclic carbonate, and a C_4 - C_7 alkyl ring with one member of the ring covalently linked to both the O of Y^1 and to X^1 ; or when X^1 and X^2 are O, L^1 and L^2 together are cyclic C_3 - C_7 ketal.

20

Detailed Description

The current invention pertains, at least in part, to methods and intermediates for the preparation and crystallization of intermediates and other compounds useful in the synthesis of halichondrin B and its analogs.

5

A. Definitions

In order that the present invention may be more readily understood, certain terms are first defined. Additional definitions are set forth throughout the detailed description.

The term “acetyl” refers to both acyl groups (e.g., -C(=O)-CH_3) and $\text{C}_1\text{-C}_8$ alkyl substituted carbonyls (e.g., $\text{-C(=O)-(C}_1\text{-C}_7\text{)alkyl}$). Preferably, the acetyl group is acyl.

The term “alkyl” refers to saturated hydrocarbons having one or more carbon atoms, including straight-chain alkyl groups (e.g., methyl, ethyl, propyl, butyl, pentyl, 5 hexyl, etc.), cyclic alkyl groups (or “cycloalkyl” or “alicyclic” or “carbocyclic” groups) (e.g., cyclopropyl, cyclopentyl, cyclohexyl, etc.), branched-chain alkyl groups (isopropyl, *tert*-butyl, *sec*-butyl, isobutyl, etc.), and alkyl-substituted alkyl groups (e.g., alkyl-substituted cycloalkyl groups and cycloalkyl-substituted alkyl groups). The terms “alkenyl” and “alkynyl” refer to unsaturated aliphatic groups analogous to alkyls, but 10 which contain at least one double or triple carbon-carbon bond respectively.

The term “alkoxy” refers to alkyl groups linked to the remainder of the molecule through an oxygen atom. Examples of alkoxy groups include, but are not limited to, methoxy, ethoxy, isopropoxy, propoxy, butoxy, and pentoxy groups. The alkoxy groups can be straight-chain or branched. Preferable alkoxy groups include methoxy.

15 The term “heterocyclic group” refers to closed ring structures analogous to carbocyclic groups in which one or more of the carbon atoms in the ring is an element other than carbon, for example, nitrogen, sulfur, or oxygen. Heterocyclic groups may be saturated or unsaturated. Additionally, heterocyclic groups (such as pyrrolyl, pyridyl, isoquinolyl, quinolyl, purinyl, and furyl) may have aromatic character, in which case 20 they may be referred to as “heteroaryl” or “heteroaromatic” groups. Exemplary heterocyclic groups include, but are not limited to pyrrole, furan, thiophene, thiazole, isothiazole, imidazole, triazole, tetrazole, pyrazole, oxazole, isooxazole, pyridine, pyrazine, pyridazine, pyrimidine, benzoxazole, benzodioxazole, benzothiazole, benzoimidazole, benzothiophene, methylenedioxyphenyl, quinoline, isoquinoline, 25 naphthridine, indole, benzofuran, purine, benzofuran, deazapurine, or indolizine.

The term “amine” or “amino,” refers to unsubstituted or substituted moiety of the formula $\text{-NR}^a\text{R}^b$, in which R^a and R^b are each independently hydrogen, alkyl, aryl, or heterocyclyl, or R^a and R^b , taken together with the nitrogen atom to which they are 30 attached, form a cyclic moiety having from 3 to 8 atoms in the ring. Thus, the term amino includes cyclic amino moieties such as piperidinyl or pyrrolidinyl groups, unless otherwise stated.

Regarding connectivity, an “arylalkyl” group, for example, is an alkyl group substituted with an aryl group (e.g., phenylmethyl (*i.e.*, benzyl)). An “alkylaryl” moiety is an aryl group substituted with an alkyl group (e.g., *p*-methylphenyl (*i.e.*, *p*-tolyl)). 35 Thus, the term imidazolyl-alkyl refers to an alkyl group substituted with an imidazolyl moiety.

The term “sulfonate” refers to moieties of the formula: $\text{R-SO}_2\text{-O-}$, wherein R is $\text{C}_1\text{-C}_4$ alkyl or $\text{C}_6\text{-C}_8$ aryl. Examples of sulfonates include, methanesulfonate (mesylate),

trifluoromethanesulfonate (triflate), *p*-toluenesulfonate (tosylate), and benzenesulfonate (bensylate).

As used in the description and drawings, an optional single/double bond is represented by a solid lines together with a second dashed line, and refers to a covalent
5 linkage between two carbon atoms which can be either a single bond or a double bond. For example, the structure:



can represent either butane or butene.

The term “protecting group” refers to moieties which may be cleaved from the
10 compound to yield a hydroxy group, a thiol group, a carboxylic acid group, or another functional group which a person of skill in the art desires to protect. Generally, protecting groups are selected such that they resist cleavage during reactions focused on other portions of the molecule. Protecting groups can be selected such that they are acid labile (e.g., cleavable in the presence of acid), base labile (e.g., cleavable in the presence
15 of base), or otherwise selectively cleavable. Protecting groups are well known to those of skill in the art. Examples of suitable protecting groups can be found, for examples in “Protective Groups in Organic Synthesis,” 3rd edition, John Wiley & Sons, Inc.

Examples of protecting groups, include, but are not limited to C₁-C₁₂ alkylcarbonyls, C₁-C₆ alkyls, C₁-C₁₅ alkyl silyl moieties (e.g., moieties which form alkyl
20 silyl ethers when bonded to an adjacent oxygen), aryl(C₁-C₆) alkyls, carbonates, and C₁-C₆ alkoxy-(C₁-C₆)alkyls (e.g., methoxymethyl).


Examples of C₁-C₁₀ alkyl silyl protecting groups include, but are not limited to, trimethylsilyl, triethylsilyl, t-butyldimethylsilyl, t-butyldiphenylsilyl, or triisopropylsilyl (e.g., trimethylsilyl ether, triethylsilyl ether, t-butyldimethylsilyl ether, t-
25 butyldiphenylsilyl ether, or triisopropylsilyl ether when taken together with an adjacent oxygen). Preferably, the alkyl silyl protecting group is t-butyldimethylsilyl ether.

Examples of C₁-C₆ alkyl protecting groups include methyl and t-butyl (e.g., methyl ethers and t-butyl ethers when taken together with an adjacent oxygen).

Examples of aryl (C₁-C₆) alkyl protecting groups include is 3,4-
30 dimethoxybenzyl, *p*-methoxybenzyl, benzyl, or trityl (e.g., 3,4-dimethoxybenzyl ether, *p*-methoxybenzyl ether, benzyl ether or trityl ether when taken together with an adjacent oxygen).

Compounds with two or more groups to be protected (e.g., hydroxy and/or thiol groups) may be protected together using a protecting group which attaches to both of the
35 hydroxy and/or thiol groups for which protection is desired. These protecting groups are also referred to herein as “divalent protecting groups.” Examples of divalent protecting groups which protect two hydroxy and/or thiol groups include, but are not limited to C₁-C₆ acetals, C₂-C₆ ketals, and cyclic carbonates. Examples of cyclic protecting groups

include, but are not limited to, acetonide, benzylidene, and, preferably, cyclohexylidene. Examples of protecting groups which protect two hydroxy and or thiol groups include those shown below. The arrows designate where the moiety is attached to the hydroxy or thiol groups on the compound:



- 5 The term “acceptable salts” refers to salts of the compounds of the invention which are acceptable for the methods of the invention, e.g., the synthesis of intermediates of halichondrin B analogs.

- 10 The compounds of the invention that are acidic in nature are capable of forming a wide variety of base salts. The chemical bases that may be used as reagents to prepare those that form base salts with such compounds. Such base salts include, but are not limited to those derived from such pharmaceutically acceptable cations such as alkali metal cations (*e.g.*, potassium and sodium) and alkaline earth metal cations (*e.g.*, calcium and magnesium), ammonium or water-soluble amine addition salts such as N-methylglucamine-(meglumine), and the lower alkanolammonium and other base salts of pharmaceutically acceptable organic amines. The base addition salts of compounds of the invention that are acidic in nature may be formed with cations by conventional methods.

- 20 The term “anti-solvent” includes organic solvents in which the compound of interest is not substantially soluble in. Examples of anti-solvents for the compounds of the present invention of formula (II) include non-polar organic solvents, such as heptane.

- 25 The term “alkylating reagent” refers to a reagent which is capable of adding an alkyl group, preferably a methyl group, to particular organic compounds described herein including, but not limited to, compounds of formula (Ia). Preferably, the alkylating reagent is a C₁-C₄ alkyl halide (preferably MeI) or a sulfonate.

- 30 The term “appropriate alkylating condition” refers to conditions which are selected such that an alkylating reaction is able to be performed. These conditions include an aprotic solvent (*e.g.*, tetrahydrofuran, toluene, or *t*-butyl methyl ether) and a base (*e.g.*, a metal amide or a metal alkoxide). Examples of bases which may be used in the alkylating conditions include, but are not limited to, LDA, KHMDS, and potassium *t*-butoxide.

The language "appropriate crystallization conditions" refers to conditions which are selected such that the desired diastereomer of a particular compound is crystallized, preferably a compound of formula (I) or (Ib). Examples of solvent systems that may be used to perform this crystallization include, but are not limited to, heptane and mixtures of heptane with one or more co-solvents, such as, but not limited to *tert*-butyl methyl ether and isopropanol. The ratio of heptane to *tert*-butyl methyl ether or isopropanol is selected such that the desired diastereomer is crystallized. The ratio may range from about 5:1 to about 3:1, and is preferably about 4:1. The appropriate conditions may also include the addition of a base. Examples of such bases include C₁-C₆ alkoxides (e.g., *t*-butyl oxide or isopropoxide). Alternatively, other solvent systems may also be used, such as, combinations of a protic solvent and an anti-solvent.

The language "second appropriate crystallization conditions" refers to conditions which are selected such that the desired diastereomer of a particular compound is crystallized, preferably a compound of formula (II) or (IIa). Examples of second appropriate crystallization conditions for the crystallization of compounds of formula (II) and/or (IIa) include dissolving the compound in a polar solvent (e.g., MTBE) and optionally adding an anti-solvent to precipitate the compound.

The term "contacting" refers to any interaction between two or more compounds which results in a chemical reaction, such as, but not limited to, the creation or the cleavage of one or more chemical bonds.

The language "mixture of diastereomers" refers to compositions which comprise two or more diastereomers.

The term "protic solvent" refers to a solvent which contains a dissociable H⁺ or a group capable of forming hydrogen bonds (e.g., hydroxyl or amine group). Examples are water, methanol, ethanol, formic acid, hydrogen fluoride and ammonia. Preferred protic solvents include alcohols, such as isopropanol.

The language "substantially diastereomerically pure composition" refers to compositions which the ratio of a particular compound to the compound with the opposite stereochemistry at the chiral center indicated with an asterisk in Scheme 1 is at least about 8:1 or greater, at least about 10:1 or greater, at least about 15:1 or greater, at least about 20:1 or greater, or, preferably, at least about 30:1 or greater. Diastereomeric purity can be enhanced using multiple kinetic or crystallization induced dynamic resolutions. It also can be enhanced by repeated recrystallizations.


The language "substantially no chromatography" refers to methods of synthesis which use 4 or fewer, 3 or fewer, 2 or fewer, 1 or fewer, or no chromatography steps. Preferably, the term refers to methods of synthesis which do not require preparative HPLC steps.

Certain abbreviations and acronyms are used herein. Definitions for these abbreviations and acronyms are listed below:

ACN	Acetonitrile
AcOH	Acetic Acid
CIDR	Crystallization induced dynamic resolution
DBU	Diazabicycloundecane
DCM	Dichloromethane
DIBAL	Diisobutylaluminium hydride
DME	Dimethoxyethane
DMF	Dimethylformamide
ESI	Electron spin injection
Et ₃ N	Triethylamine
EtOAc	Ethyl acetate
EtOH	Ethanol
FDA	Food and Drug Administration
HPLC	High pressure liquid chromatography
IPA	Isopropanol
ⁱ Pr ₂ NEt	Diisopropylethylamine
KHMDS	Potassium-Hexamethyldisilazane
KO ^t Bu	Potassium tert-butoxide
LDA	Lithium diisopropyl amide
LRMS	Low resolution mass spectrometry
MeI	Methyl iodide
MeOH	Methanol
MsCl	Mesyl chloride (methanesulfonyl chloride; CH ₃ SO ₂ Cl)
MTBE	Methyl tert-butyl ether
MsO-	Mesylate (methanesulfonate)
NaOEt	Sodium ethoxide
NaOMe	Sodium methoxide
NBS	N-bromosuccinimide
NIS	N-iodosuccinimide
NMR	Nuclear magnetic resonance
Ph ₃ P	Triphenyl phosphine
TBDPSCI	tert-Butyl diphenyl silyl chloride
TBME	tert-Butyl methyl ether
TBS	tert-Butyldimethyl silyl
TBSCl	tert-Butyldimethyl silyl chloride
TBSOTf	tert-Butyldimethylsilyl trifluoromethanesulphonate
^t BuOK	Potassium tert-butoxide
TEA	Triethylamine
TESOTf	Triethylsilyl trifluoromethanesulfonate
TsCl	Tosyl chloride (p-toluenesulfonyl chloride)
TfO-	Triflate (trifluoromethanesulfonate)
Tf ₂ O	Triflic anhydride (CF ₃ SO ₂) ₂ O
TsO-	Tosylate (p-toluenesulfonate)
THF	Tetrahydrofuran
TsOH	p-Toluene sulfonic acid
TosMIC	Toluenesulfonylmethyl isocyanide
Trt	Trityl (Triphenylmethyl)

B. Compounds

In one embodiment, the invention pertains to a compound of formula (I):



5 wherein:

z is a single or double bond, provided that when z is a double bond, X² is C and Y¹ is hydrogen; and provided that when z is single bond, X² is CH or O;

X¹ is O, S, or CN, provided that when X¹ is CN or S, X² is O;

Y¹ is a halide, hydrogen or O-L², or absent when X² is O; and

10 L¹ and L² are independently selected from hydrogen and a protecting group, or L¹ and L² together are a protecting group, provided that when X¹ is CN, L¹ is absent; and salts thereof. The invention also pertains to compounds of formula (I).

In an embodiment, L¹ and/or L² are each independently a silyl ether, a C₁-C₈ alkyl ether, an acyl (-C(=O)CH₃), or acetyl group. Preferably, X¹ is oxygen.

15 Preferably, L¹ and L² may represent the same protecting group attached to the molecule through both the O of X² when Y¹ is O-L² and X¹. Examples of such protecting groups include, but are not limited to, cyclic C₁-C₆ acetals, cyclic C₂-C₆ ketals, and cyclic carbonates. In a further embodiment, L¹ and L² are linked to a single divalent protecting group. Examples of divalent protecting groups include acetonides, 20 benzylidines, and preferably, cyclohexylidene. In certain embodiments, when both L¹ and L² are protecting groups, L¹ and L² when taken together may form a pentane, hexane, or pyran ring and link to X¹ to X² through a single carbon atom. Preferably, when Y¹ is O-L²; X¹ is O or S; L¹ and L² together form protecting group which is a C₄-C₇ alkyl ring with one member of the ring covalently linked to the O of O-L² and to X¹.

25 In one embodiment, X² is CH, Y¹ is O-L², and X¹ is O.


In another embodiment, when Y¹ is a halide, it is fluoride, chloride, iodide, or, preferably, bromide. In another further embodiment, L¹ is acetyl.

In another embodiment, when z is a double bond, Y is hydrogen, and X² is C. In another further embodiment, X¹ is oxygen and L¹ is a protecting group (when taken 30 together with X¹) selected from the group consisting of C₁-C₆ alkyl ether, aryl (C₁-C₆) alkyl ether, C₁-C₆ ester, and a silyl (C₁-C₁₀) ether.

In another further embodiment, X² is oxygen, when z is a single bond. In another further embodiment, L¹ is hydrogen. In another further embodiment, L¹ is a protecting group selected from a glycoside, C₁-C₆ alkyl, C₁-C₆ acetyl, and a C₁-C₆ ester.

35

Preferably, the compound of formula (I) is a compound of formula (Ib):




wherein L^{1a} and L^{1b} are hydrogen, independently selected protecting groups, or together a single divalent protecting group.


- 5 In a further embodiment, L^{1a} and L^{1b} are each protecting groups selected from C₁-C₆ alkyl ethers, aryl (C₁-C₆) alkyl ethers, silyl (C₁-C₁₀) ethers, C₁-C₆ alkyl esters, cyclic C₁-C₆ acetals, cyclic C₂-C₇ ketals, and cyclic carbonates.

- 10 In a further embodiment, the invention pertains to a composition comprising a compound of formula (Ib), wherein the composition is substantially diastereomerically pure. In a further embodiment, the ratio of compounds of formula (Ib) to the compounds with the opposite stereochemistry at the chiral center marked with the asterisk is at least about 8:1 or greater, at least about 20:1 or greater, or, preferably, at least about 30:1 or greater.


- 15 In a further embodiment, the compound of formula (I) is selected from the group consisting of:




Compound AD




Compound AJ



Compound AK




Compound AL



Compound AM

or a salt thereof.

In another embodiment, the invention pertains to a compound of formula (II):



wherein:

c is a single or double bond, provided that when c is a double bond m is 0 and Y^3 is O or $CHCO_2-L^3$, and provided that when c is a single bond m is 0 or 1 Y^3 is CH_2O-L^3 , $CH_2CO_2-L^3$ or $CH_2CH_2O-L^3$;

Y^2 is C_1-C_7 sulfonate, $O-L^4$ or a halide;

5 L^4 is hydrogen or a protecting group; and

L^3 and L^5 are each independently hydrogen or a protecting group, or L^3 and L^5 together are a protecting group, or a salt thereof.



Examples of Y^2 include halides, e.g., fluoride, chloride, bromide, or preferably, iodide. In another embodiment, Y^2 is $O-L^4$. Examples of L^4 include hydrogen. In
10 another embodiment, c is a double bond. Examples of Y^3 when c is a double bond include $CHCO_2-L^3$. Examples of L^3 groups include C_1-C_6 alkyl, e.g., methyl.

In another embodiment, c is a single bond. Examples of Y^3 when c is a single bond include $CH_2CH_2OL^3$. In a further embodiment, L^3 and L^5 may be linked to form a cyclic C_1-C_6 acetal or a cyclic C_2-C_7 ketal.


15 In a further embodiment, Y^3 is $CH_2CO_2-L^3$ and L^3 is C_1-C_{10} alkyl, C_4-C_{10} aryl- C_1-C_6 alkyl, or C_4-C_{10} aryl. In another further embodiment, Y^2 is a halide, e.g., iodide.

In a further embodiment, the invention pertains to a composition comprising a compound of formula (II), wherein the composition is substantially diastereomerically pure. In a further embodiment, the ratio of compounds of formula (II) to the compounds
20 with the opposite stereochemistry at the chiral center marked with the asterisk is at least about 8:1 or greater, at least about 20:1 or greater, or, preferably, at least about 30:1 or greater.

In another further embodiment, the compound of formula (II) is selected from the
25 group consisting of:



Compound AP, and




Compound AQ

Compound AF

or a salt thereof.

The invention also pertains to compositions comprising the compounds shown above substantially free of diastereomers.


In a further embodiment, the invention also pertains to a compound of formula (IIa):



In a further embodiment, the compound of formula (IIa) is substantially free of diastereomers, e.g., a compound with the opposite stereochemistry at the chiral carbon indicated with an asterisk in the formula above. In an embodiment, the invention pertains to a substantially diastereomerically pure composition comprising a compound of formula (IIa), wherein the ratio of compounds of formula (IIa) to compounds with the opposite stereochemistry at the chiral center marked with the asterisk is at least about 8:1 or greater, at least about 20:1 or greater, or, preferably, at least about 30:1 or greater.

The compound of formula (IIa) is particularly important because while it is crystalline, the corresponding non-methylated intermediate is not crystalline and requires purification via chromatography. The invention also pertains to a compounds of formula (IIa) in crystalline form.


In another embodiment, the invention pertains to a compound of formula (III):




wherein: L^6 is hydrogen or a protecting group; or an acceptable salt thereof. In an embodiment, the invention pertains to a substantially diastereomerically pure composition comprising a compound of formula (III), wherein the ratio of compounds of formula (III) to compounds with the opposite stereochemistry at the chiral center marked with the asterisk is at least about 8:1 or greater, at least about 20:1 or greater, or, preferably, at least about 30:1 or greater.

In a further embodiment L^6 is hydrogen or, when taken together with the oxygen to which it is bound, a silyl C_1 - C_{10} ether. Examples of such silyl C_1 - C_{10} ethers include, but are not limited to, trimethylsilyl ether, triethylsilyl ether, t-butyldimethylsilyl ether, t-butyldiphenylsilyl ether, or triisopropylsilyl ether.

In a further embodiment, the compound of formula (III) is:




Compound AR



Compound AH

The invention also pertains to compositions comprising the compounds shown above substantially free of diastereomers.


In another embodiment, the invention pertains to compounds of formula (IIIa):



(IIIa)

wherein L^{6a} , L^{6b} , and L^{6c} are each protecting groups, or a salt thereof. In a further embodiment, the invention pertains to a composition comprising the compound of formula (IIIa) wherein the composition is substantially free of diastereomers (e.g., compounds with the opposite stereochemistry at the chiral center indicated with an asterisk in formula (IIIa) above).

The invention also pertains, at least in part, to compounds of formula (Id):




(Id)

wherein L^{1a} and L^{1b} are independently selected from hydrogen and a protecting group, or L^{1a} and L^{1b} together are a divalent protecting group, or a salt thereof.

C. Methods

In one embodiment, the invention pertains to a method of obtaining a substantially diastereomerically pure composition comprising a compound of formula (I). The method includes crystallizing the compound of formula (I) from a mixture of diastereomers under appropriate crystallization conditions, such that a substantially diastereomerically pure composition comprising a compound of formula (I) is formed.

The mixture of diastereomers is preferably a mixture of compounds of formula (I) with compounds of formula (Ie), wherein said compounds of formula (Ie) is:




(Ie).

In one embodiment, the substantially diastereomerically pure composition comprises a ratio of compounds of formula (I) to compounds of formula (Ie) of at least about 8:1 or greater, of at least about 10:1 or greater, of at least about 20:1 or greater, or, preferably, at least about 30:1 or greater. In order to increase the diastereomeric purity of the compound of formula (I), additional recrystallizations of the compound under similar appropriate conditions may be conducted.

The appropriate crystallization conditions are selected such that the desired diastereomer is crystallized. Examples of solvent systems that may be used to perform this crystallization include, but are not limited to, heptane/ *tert*-butyl methyl ether and heptane/ isopropanol. The appropriate conditions may also include the addition of a base. Examples of such bases include C₁-C₆ alkoxides (e.g., *t*-butyl oxide or isopropoxide).


Alternatively, other solvent systems may also be used, such as, combinations of a protic solvent (e.g., an alcohol, e.g., isopropanol) and an anti-solvent (e.g., non-polar organic solvent, e.g., heptane).

In a further embodiment, the invention also pertains to a method of synthesizing the compound of formula (Ib) from a compound of formula (Ia) by contacting a compound of formula (Ia) with an alkylating reagent under appropriate alkylating conditions. The compound of formula (Ia) is:



(Ia)

and the compound of formula (Ib) is:



(Ib)

wherein L^{1a} and L^{1b} are independently selected from hydrogen and a protecting group, or L^{1a} and L^{1b} together are a divalent protecting group, provided that L^{1a} of formulae (Ia) and (Ib) are the same and L^{1b} of formulae (Ia) and (Ib) are the same. The method includes reacting the compound of formula (Ia) under alkylating conditions to form a mixture comprising the compound of formula (Ib) and diastereomers thereof; and

crystallizing the compound of formula (Ib) from the mixture, under appropriate crystallization conditions.

5 In order to increase the diastereomeric purity of the compound of formula (Ib), additional recrystallizations of the compound under similar appropriate conditions may be conducted. Preferably, the mixture of diastereomers after two or more crystallizations results in a ratio of compounds of formula (Ib) to compounds with the opposite stereochemistry around the chiral center indicated with the asterisk above in formula (Ib) to be at least about 8:1 or greater, at least about 10:1 or greater, at least about 20:1 or greater, or at least about 30:1 or greater.

10 In yet another embodiment, the invention also pertains, at least in part, to a method of obtaining a substantially diastereomerically pure composition comprising a compound of formula (I). The method includes contacting a mixture of diastereomers with a base at an appropriate temperature, such that a substantially diastereomerically pure composition comprising a compound of formula (I) is formed.

15 Examples of bases which may be used in the method include bases known in the art, such as amide bases, metal alkoxides and KHMDS. The base may be present in any amount such that the desired diastereomer is formed. Preferably, the base is present in sub stoichiometric amounts (e.g., less than one equivalent). In another further embodiment, the appropriate temperature is less than about -30 °C. In a further
20 embodiment, the compound of formula (I) is a compound of formula (Ib).


If kinetic resolution of the stereocenter is desired, the compound of formula (I) or (II) may be treated with sub-stoichiometric amounts of a strong base (e.g., an amide base, e.g., KHMDS) at low temperatures (e.g., less than about -30 °C). Once the reaction has taken place, compounds of formula (I) or (II) may be isolated from an
25 appropriate crystallization solvent system and recrystallized. Examples of solvent systems which may be used include, but are not limited to, heptane, heptane/*t*-butyl methyl ether and heptane/ isopropanol.

Alternatively, crystallization induced dynamic resolution (CIDR) may also be used to enhance the diastereomeric purity of compounds of formula (I) and/or (II). For
30 example, compounds of formula (I) and/or (II) may be treated with a weak base, such as an alkoxide, (e.g., potassium *t*-butyl oxide or potassium isopropoxide) in an appropriate crystallization solvent system. Examples of appropriate crystallization solvent systems include combinations of a protic solvent (e.g., isopropanol) and an anti-solvent (e.g., heptane) at non-cryogenic temperatures to provide purified compounds of formula (I) or
35 (II).

In another embodiment, the invention pertains to a method of obtaining a substantially diastereomerically pure composition comprising a compound of formula (II). The method includes crystallizing the compound of formula (II) from a mixture of

diastereomers under appropriate crystallization conditions, such that a substantially diastereomerically pure composition comprising a compound of formula (II) is formed.

In one embodiment, the composition comprises a ratio of compounds of formula (II) to compounds of formula (IIb) of at least about 8:1 or greater, at least about 10:1 or greater, at least about 20:1 or greater, or, preferably, at least about 30:1 or greater. The compound of formula (IIb) is:



In order to increase the diastereomeric purity of the compound of formula (II), additional recrystallizations of the compound under similar appropriate conditions may be conducted.

In another embodiment, the invention also pertains to a method of synthesizing a compound of formula (IIa) from a compound of formula (Ib). The method includes selectively crystallizing a compound of formula (Ib) under appropriate crystallization conditions; and reacting the compound of formula (Ib), under appropriate conditions, such that a compound of formula (IIa) is formed. Preferably, the compound of formula (IIa) is formed using substantially no chromatography. The compound of formula (Ib) may be reacted under appropriate conditions to form a compound of formula (IIa), after having had been diastereomerically purified using recrystallization. Furthermore, the appropriate conditions may comprise dissolving a crystallized compound of formula (Ib) in a solvent before reacting it under appropriate conditions to form a compound of formula (IIa).

Appropriate conditions for the synthesis of compounds of formula (IIa) from compounds of formula (Ib) are described, for example, in Schemes 5, 6, 8, 9, and 10. Methods for selectively crystallizing a compound of formula (I) or (Ib) from a mixture of diastereomers under appropriate crystallization conditions has been described above.

The invention also pertains, at least in part, to a method of synthesizing a compound of formula (IIIa) from a compound of formula (IIa). The method includes crystallizing a compound of formula (IIa) under second appropriate crystallization conditions; reacting the compound of formula (IIa) under appropriate conditions such that a compound of formula (IIIa) is formed.


Examples of second appropriate crystallization conditions for the crystallization of compounds of formula (IIa) include dissolving the compound in a polar solvent (e.g., MTBE) and optionally adding an anti-solvent to precipitate the compound. Examples of anti-solvents which may be used include heptane. Preferably, the compound of formula

(IIa) is reacted under appropriate conditions to form a compound of formula (IIIa), after having had been crystallized.

In another embodiment, the invention also pertains to a method of synthesizing a compound of formula (IIIa) from a compound of formula (Ib). The method includes
5 selectively crystallizing a compound of formula (Ib) under appropriate crystallization conditions; and reacting the compound of formula (Ib) under appropriate conditions such that a compound of formula (IIIa) is formed. Preferably, the compound of formula (IIIa) is formed using substantially no chromatography.

The compound of formula (Ib) may be reacted under appropriate conditions to
10 form a compound of formula (IIIa), after having had been diastereomerically purified using recrystallization. Furthermore, the appropriate conditions may comprise dissolving a crystallized compound of formula (Ib) in a solvent before reacting it under appropriate conditions to form a compound of formula (IIIa).

In another embodiment, the invention pertains to a method of synthesizing
15 compounds of formula (IV). The method includes crystallizing a compound of formula (Ib) from a mixture of diastereomers under appropriate crystallization conditions, as described above; reacting the selectively crystallized compound of formula (Ib) with appropriate reagents, such that a compound of formula (IV) is synthesized. The compound of formula (IV) is:



20 wherein each of L^{7a} , L^{7b} , L^{7c} , L^{7d} , and L^{7e} are each a protecting group or hydrogen. Examples of L^{7a} include phenyl. Examples of L^{7b} include methyl. Examples of L^{7c} and L^{7d} include TBS. Examples of L^{7e} includes hydrogen.

Examples of appropriate reagents which may be used to synthesize compounds
25 of formula (IV) from a compound of formula (Ib) include those described in Schemes 5, 6, 8, and 9 to form a compound of formula (IIIa). Methods which may be used to convert a compound of formula (IIIa) to a compound of formula (IV) are described in greater detail in WO/2005/118565, incorporated herein by reference in its entirety.


The compound of formula (Ib) may be reacted under appropriate conditions to form a compound of formula (IV), after having had been diastereomerically purified using recrystallization. Furthermore, the appropriate conditions may comprise dissolving a crystallized compound of formula (Ib) in a solvent before reacting it under appropriate conditions to form a compound of formula (IV).

In a further embodiment, the compound of formula (IV) is formed in greater than about 50% yield, greater than about 60% yield, or greater than about 70% yield from a compound of formula (Ib).

In a further embodiment, the invention also pertains to compounds of formula (I), (Ia), (Ib), (Ic), (Id), (Ie), (II), (IIa), (IIb), (III), (IIIa), (IV), (V), or otherwise described herein. The invention also pertains to compositions comprising compounds of any one of these formulae, substantially free of diastereomers. The invention also pertains to each of the intermediates and processes described herein.

In a further embodiment, the invention pertains to compositions comprising the compounds described herein substantially free of diastereomers, e.g., compounds with the opposite stereochemistry at the chiral carbon indicated with the asterisk in Scheme 1. The invention also pertains to methods of using these compounds to synthesize compounds of formula (IV), B-1939, or other halichondrin B analogs.


The invention pertains, at least in part, to methods and intermediates for the conversion of compounds of formula (I) to compounds of formula (III). Compounds of formula (III) may further be converted to compounds of formula (IV) and/or halichondrin B or analogs thereof.



The compounds of formula (III) can be synthesized by methods described herein. The invention pertains, at least in part, to all compounds and intermediates described herein and the processes of synthesizing the compounds and intermediates.

Conversion of Compound 2-1 to a Compound of Formula (Ib)

Compounds of formula (Ib) can be synthesized from compounds of formula 2-1, as shown in Scheme 2:



Scheme 2

Compound **2-1** may be converted to compound (**Ib**). In Scheme 2, L^{1a} and L^{1b} are protecting groups. Examples of protecting groups include, but are not limited to C_1 - C_6 alkyl ethers, aryl (C_1 - C_6) alkyl ethers, silyl (C_1 - C_{10}) ethers, C_1 - C_6 alkyl esters, cyclic C_1 - C_6 acetals, cyclic C_2 - C_7 ketals, and cyclic carbonates. Examples of R^2 include hydrogen, C_1 - C_6 alkyl (e.g., methyl, *t*-butyl, etc.), C_4 - C_{10} aryl (e.g., phenyl), and C_4 - C_{10} aryl- C_1 - C_6 alkyl groups (e.g., benzyl). Examples of R^3 and R^4 include CH_3 and OCH_3 respectively, or R^3 and R^4 taken together can be $(-CH_2CH_2)_2O$.

Compound **2-1** can be converted to compound **2-4** through the use of an appropriate reducing agent. Examples of such reducing agents include, but are not limited to, aluminum hydrides and borohydrides (e.g. BH_3 , AlH_3 , $LiBH_4$, $LiAlH_4$, $NaBH_4$, $NaAlH_4$, $ZnBH_4$).

The hydroxyl group of compound **2-4** may be transformed to a leaving group such as but not limited to a sulfonate (e.g., $MsO-$, $TsO-$, $TfO-$) or halide by methods described in the literature. Subsequent treatment with a cyanide source (e.g., KCN or $NaCN$) results in the formation of compound **2-5**.

Alternatively, compound **2-4** may be transformed to compound **2-5** by oxidation of the hydroxyl group to the aldehyde by methods described in the literature. Conversion of the aldehyde to the nitrile may be achieved with appropriate reagents such as, but not limited to, dimethyl phosphorocyanidate/ samarium iodide. Compound **2-5** may be alkylated in an appropriate solvent, e.g., an aprotic solvent such as tetrahydrofuran, toluene, TBME and subsequently treated with a strong base such as a


metal amide or metal alkoxides (e.g., LDA, KHMDS, or KO^tBu) and an appropriate alkyl halide (e.g., X-Me) or sulfonate to provide a compound of formula (Ib).

Alternatively, compound 2-1 may be converted to compound 2-7 by methods known in the art. Examples of such methods include, but not limited to, treatment with N,O-dimethylhydroxylamine hydrochloride/ trimethylaluminum. Compound 2-7 can be converted into the compound of formula (Id), by treatment with an appropriate carbon-nucleophile. Examples of such nucleophiles include, but are not limited to, alkyl Grignard reagents.

Alternatively, oxidation of compound 2-4 to the aldehyde followed by addition of an alkyl Grignard or other carbon-nucleophile provides a secondary alcohol. Oxidation using known methods results in the formation of the compound of formula (Id). Compounds of formula (Ib) may be synthesized, for example, by the treatment of the compound of formula (Id) with TosMIC in the presence of metal alkoxides, such as NaOEt and KO^tBu (J. Org. Chem. 42(19), 3114-3118, (1977)). Alternatively, the compound of formula (Id) may be transformed to the compound of formula (Ib) using reagents such as but not limited to dimethyl phosphorocyanidate/ samarium iodide.

Conversion of (-)-Quinic Acid to a Compound of Formula (Id)

Alternatively a compound of formula (Id) may also be synthesized as shown in Scheme 3.



Scheme 3

In Scheme 3, L^{1a} and L^{1b} are protecting groups, as described in Scheme 2. L^{2b} and L^{2c} are also protecting groups such as, but not limited to, cyclic acetals (X = O and/or S), cyclic ketals (X = O and/or S), and cyclic carbonates (X=O).


The synthesis of compound 3-9 from commercially available (-)-quinic acid has been described previously (WO/2005/118565). Compound 3-9 may be reduced with DIBAL or other reagents known in the art, such as aluminum hydrides and borohydrides, to provide lactol 3-10. The lactol 3-10 can be transformed using a Wittig

or Julia olefination to provide compound **3-11**. Deprotection followed by double bond migration and Michael addition generate compound (**Id**).

More specifically, a Wittig olefination may be carried out using $\text{MeC}(\text{OL}^{2b})(\text{OL}^{2c})\text{CH}_2\text{CH}_2\text{PPh}_3$ (prepared *in situ*), in a polar solvent (e.g., THF, MeOH, or DMF) at a temperature ranging from 0 °C to 50 °C. Acid-catalyzed sequential reactions (e.g., deprotection, migration, and Michael addition) may be carried out with an acid such as TsOH or HCl in a polar solvent (e.g. THF, or acetone) at a temperature ranging from 10 to 30 °C for about two to four hours. Alternatively, the migration and Michael addition may also be carried out with a base such as NaOMe in a polar solvent (e.g., THF or MeOH).

Conversion of Compound 4-1 to a Compound of Formula (I)

As shown in Scheme 4, compounds of formula **4-1** (diastereomeric mix) may undergo isomerization and crystallization to provide compounds of formula (**I**).




Scheme 4

In Scheme 4, z is a single or double bond, provided that when z is a double bond, X^2 is C and Y^1 is hydrogen; and provided that when z is single bond, X^2 is CH or O; X^1 is O, S, or CN, provided that when X^1 is CN or S, X^2 is O; Y^1 is a halide, hydrogen or O-L^2 , or absent when X^2 is O; and L^1 and L^2 are independently selected from hydrogen and a protecting group, or L^1 and L^2 together are a protecting group, provided that when X^1 is CN, L^1 is absent.

Diastereomers of formula **4-1** may be converted to compounds of formula (**I**) via treatment with sub-stoichiometric amounts of amide bases (e.g., KHMDS) at low temperatures (e.g., less than about -30 °C). Once quenched, compounds of formula (**I**) may be isolated and recrystallized from a suitable crystallization solvent systems, such as, but not limited to, heptane/*t*-butyl methyl ether and heptane/ isopropanol.

Alternatively a crystallization induced dynamic resolution (CIDR) may also be used to selectively crystallize compounds of formula (**I**). For example, diastereomers of formula **4-1** may be treated with a base, such as an alkoxide, (e.g., *t*-butyl oxide or isopropoxide) in an appropriate crystallization solvent system. Examples of appropriate crystallization solvent systems include combinations of a protic solvent (e.g., isopropanol) and an anti-solvent (e.g., heptane) at non-cryogenic temperatures to provide purified compounds of formula (**I**).

Conversion of a Compound of Formula (Ib) to Compound 5-13



Scheme 5

5


In Scheme 5, L^{1a} and L^{1b} are as described above in Scheme 2. L^{1d} is a suitable protecting group, e.g., C_1 - C_6 alkyl ether, aryl (C_1 - C_6) alkyl ether, C_1 - C_6 ester, or a silyl (C_1 - C_{10}) ether.

Compounds of formula (Ib) may be deprotected by various methods known in the art, depending on the nature of L^{1a} and L^{1b} . Examples of deprotecting reactions include, but are not limited to hydrogenation, reduction, oxidation, base induced deprotection, and acid induced deprotection. One of ordinary skill in the art would be able to choose an appropriate technique based on art recognized techniques (see, e.g., Protective Groups in Organic Synthesis, 3rd edition, John Wiley & Sons, Inc).

Once L^{1a} and L^{1b} have been removed, the deprotected compound 5-12 may be converted to compound 5-13 by treatment of compound 5-12 with 2-acetoxy-2-methylpropionyl bromide, catalytic water, in a polar aprotic solvent such as acetonitrile. The resulting intermediate may be treated with a base (e.g., diazabicycloundecane (DBU)) to provide compound 5-13.

Alternatively, compound 5-12 may be transformed to compound 5-13 using a multi-step process. The process involves selectively activating of one hydroxyl group as a halide, MsO^- , TsO^- , or TfO^- and protecting of the remaining hydroxyl group. Examples of suitable protecting groups for this step include L^{1d} groups such as C_1 - C_6 alkyl ethers, aryl (C_1 - C_6) alkyl ethers, C_1 - C_6 esters, and silyl (C_1 - C_{10}) ethers. The intermediate can be transformed to compound 5-13 using methods described previously.

Conversion of Compound 5-13 to Compound 6-16



Scheme 6


In Scheme 6, L^{1d} is hydrogen or a protecting group, C_1 - C_6 alkyl ether, aryl (C_1 - C_6) alkyl ether, C_1 - C_6 ester, or a silyl (C_1 - C_{10}) ether). L^{1c} may be hydrogen or a protecting group, such as, but not limited to a glycoside, C_1 - C_6 alkyl, or a C_1 - C_6 ester.

5 X^2 and X^3 may each be oxygen or hydroxy.

Oxidative cleavage of the olefin of compound **5-13** may be accomplished using ozone in a suitable solvent (e.g., methanol) at temperatures below 0 °C. The ozone adduct may be worked up using literature methods to provide compound **6-14**, wherein X^2 and X^3 are each carbonyl or hydroxy. Alternatively, a metal oxide (e.g., osmium tetroxide or potassium permanganate and sodium periodate) may also be used to provide compound **6-14**, wherein X^2 and X^3 are each carbonyl.

When X^2 and X^3 are each carbonyl, they can be reduced to provide compound **6-14**, wherein X^2 and X^3 are each hydroxy. Deprotection of L^1 may be achieved using literature methods (e.g., potassium carbonate in methanol) to provide compound **6-15**.
 15 Compound **6-15** may be treated with $NaIO_4$ to provide compound **6-16**, wherein L^{1c} is H. Alternatively, compound **6-16** may comprise a glycoside (e.g., L^{1c} is C_1 - C_3 alkyl, e.g., methyl) protecting group which can be added using methods known in the art, such as methanol in the presence of an acid catalyst.

20 Conversion of Compound 2-1 to Compound 6-16




Scheme 7

In Scheme 7, an alternate route to compound **6-16** is shown. Examples of R^2 include hydrogen, C_1 - C_6 alkyl (e.g., methyl, *t*-butyl, etc.), C_4 - C_{10} aryl (e.g., phenyl), and C_4 - C_{10} aryl- C_1 - C_6 alkyl groups (e.g., benzyl). L^{1a} and L^{1b} are protecting groups as described above. Examples of L^{1c} include hydrogen and protecting groups such as glycosides, C_1 - C_6 alkyl and C_1 - C_6 acetyl.

Compound **2-1** may be transformed to compound **7-17d** as described in Schemes 5 and 6. Treatment of compound **7-17d** with TosMIC and isomerization/ crystallization provides compound **6-16**, as shown in Schemes 2 and 4.

Conversion of Compound 5-16 to Compound 7-20



Scheme 8

5

Compound **8-20** may be prepared from **6-16** as shown in Scheme 8. In Scheme 8, L^{1c} is hydrogen or a protecting group as described previously; L^5 is C_1 - C_{10} alkyl, C_4 - C_{10} aryl- C_1 - C_6 alkyl, or C_4 - C_{10} aryl; and Y^2 is sulfonate or halide.

When L^{1c} is not hydrogen, the ether is hydrolyzed using literature methods to provide lactol (**6-16**, $L^{1c} = H$). The lactol (**6-16**, $L^{1c} = H$) is converted to compound **8-18** by an olefination reaction such as a stabilized Wittig reaction, a Wadsworth-Horner-Emmons reaction, or a Julia olefination. The Wittig olefination may be carried out with a stabilized ylide such as $Ph_3PCHCO_2L^5$ in polar solvent (e.g., THF, MeOH, or DMF) at an appropriate temperature (e.g., $-78^\circ C$ to $50^\circ C$). The Wadsworth-Horner-Emmons olefination may be carried out using a stabilized ylide (e.g. $(MeO)_2POCH_2CO_2L^5$) in a polar aprotic solvent (e.g., THF, or ACN) at an appropriate temperature (e.g., $-78^\circ C$ to $25^\circ C$) in the presence of a suitable base (e.g., $tBuOK$, NaH, or LiCl/tertiary amines (e.g. DBU, Pr_2NEt , Et_3N)).


Variations on these conditions are known in the art. For example, for variations on the Wadsworth-Horner-Emmons olefination see Org. React. 25, 73-253, (1977) and Tetrahedron Lett. 25, 2183 (1984). Furthermore, the Julia olefination may be carried out in a polar aprotic solvent (e.g., THF, DME or a halogenated solvent, e.g., CH_2Cl_2) in the presence of alkyl sulfone (e.g., alkyl(benzothiazol-2-ylsulfonyl)acetate) and suitable base (e.g., BuLi, LDA, KHMDS, or DBU) at an appropriate temperature (e.g., $-78^\circ C$ to $25^\circ C$). Variations on these conditions will be apparent from the literature on Julia olefination (see, Org. Biomol. Chem. 3, 1365-1368, (2005); Synlett, 26-28, (1998)).

Compound **8-19** can be obtained from compound **8-18** via catalytic hydrogenation, which may be carried out in the presence of a metal catalyst (e.g., palladium (Pd/C) or platinum (PtO_2)) in a polar solvent (e.g. EtOAc, MeOH). Preferably, the reaction is carried out under a hydrogen atmosphere, with a pressure ranging from 0.04 bar to 1.10 bar.

The hydroxyl group of **8-19** may be converted to a leaving group (e.g., MsO-, TsO-, TfO-) providing **8-20**, using a suitable sulfonyl anhydride or sulfonyl chloride (e.g., MsCl, TsCl, or Tf₂O) in a polar aprotic solvent (e.g., THF or a halogenated solvent (e.g., CH₂Cl₂)), in the presence of a suitable base (e.g., Et₃N).

5 Optionally, the leaving group of **8-20** may be converted to a halide. This reaction may be carried out in the presence of a halogenating reagent (e.g., NaI, or NaBr) in a polar solvent (e.g., DMF or acetone). Alternatively, the transformation of hydroxyl group to halide may be carried out using a halogenating reagent (e.g., NIS, or NBS) in a polar solvent (e.g., THF) in the presence of Ph₃P and a suitable base such as
10 pyridine.

Conversion of Compound 6-15 to Compound 8-20




Scheme 9

15

In Scheme 9, another method of converting compound **6-15** to compound **8-20** is shown. In Scheme 9, R^{9a} and R^{9b} are hydrogen, C₁-C₆ alkyl or taken together are a carbonyl group; Y² is sulfonate or halide; and Y³ is O, OL³, or CHCO₂-L³ wherein L³ is hydrogen or a protecting group.

20 As shown in Scheme 9, compound **8-20** can be prepared from compound **6-15** using literature methodology for the protection of 1,2 diols. Treatment of the neopentyl-hydroxyl of compound **9-21** (using methods described in Scheme 8) provides compound **9-22**. Deprotection of the compound **9-22** using literature methods provides a diol. Treatment of this diol with a reagent such as sodium periodate provides aldehyde **9-23**
25 (Y³ = O). Treatment of the aldehyde **9-23** as described in Scheme 8 provides compound **8-20**.

Conversion of Compound 7-17b to 8-20




5

Alternatively, compound **7-17b** may be converted to compound **8-20** as shown in Scheme 10. In Scheme 10, R^2 includes C_1 - C_6 alkyl such as methyl, ethyl, and *tert*-butyl; R^{10a} and R^{10b} are hydrogen, C_1 - C_6 alkyl or taken together are a carbonyl group; Y^2 is sulfonate or halide; and Y^3 is O, OL^3 , or $CHCO_2-L^3$. L^3 is hydrogen or a protecting group.

Compound **7-17b** may be transformed into compound **10-17f** by selectively protection of the 1,2-diol and a subsequent functional group transformation of the neopentyl hydroxyl group to a sulfonate or halide. Selective protection of 1,2-diol may be carried out with an aldehyde, ketone, acetal, or a carboxyl chloride (e.g. DMP, cyclohexanone, MeOPhCHO, or Ph_3P) in the presence of acid catalyst. The functional group transformation of the neopentyl hydroxyl group to a sulfonate or halide has been previously described above in Scheme 8. Compound **10-17g** and **10-22** may be prepared in a similar manner previously described in Scheme 2. Deprotection of the diol protecting group using literature procedures followed by treatment with sodium periodate provides **10-23** ($Y = O$). A Wittig, Horner-Wadsworth-Emmons, or Peterson type olefination is then followed with hydrogenation to provide compound **8-20**.


D. Chemical Examples

Example 1: Synthesis of Compound AD from Compound AC



- Compound AC (1 Wt, 1 V, 1 eq) was dissolved in THF (1.80 V) and cooled to -75 °C. KHMDS (0.50M solution in toluene, 6.60 V, 1.10 eq) was added at a rate such that internal temperature did not exceed -65 °C. Upon complete addition, stirring was continued at -75 °C for 30 minutes. A solution of MeI (0.188 V, 1.01 eq) in THF (0.50 V) was added at a rate such that internal temperature did not exceed -65 °C. Upon complete addition, stirring was continued at -75 °C for 1 hour. KHMDS (0.50M solution in toluene, 0.60 V, 0.10 eq) was added at a rate such that internal temperature did not exceed -70 °C and stirring was continued at -75 °C for additional 2.5 hours. Under vigorous stirring, 20 wt% NH₄Cl aq (1.50 Wt, 1.9 eq) was added at a rate such that internal temperature did not exceed -55 °C. Upon complete addition, the resultant mixture was allowed to warm to -20 °C. Water (1.50 V) was added and the mixture was further warmed to 0 °C. The biphasic mixture was transferred to a work-up vessel (the reactor was washed with MTBE (0.40 V)) and vigorous stirring was continued for 2 minutes. The aqueous layer was set aside and the organic layer was washed with water (2.0 V). The organic layer was concentrated and residual solvents and water were azeotropically removed with heptane (1.50 V x 2) to give the crude product as a yellow solid (1.1 Wt, dr=4.4:1).
- 25 The crude (1.1 Wt) was suspended in heptane-MTBE (4:1 v/v, 5.0 V) and heated to 80 °C. The resultant solution was: 1) cooled to 70 °C over 1 hour; 2) held at 70 °C for 0.5 hour; 3) cooled to 65 °C over 0.5 hour (precipitation started); 4) held at 65 °C for 0.5 hour; 5) cooled to 60 °C over 0.5 hour; 6) held at 50 °C for 0.5 hour; 7) cooled to room temperature and stirring was continued for 40 hours. Crystals were collected by
- 30 filtration, washed with heptane (1 V x 2) and dried under N₂/vacuum to give Compound AD as light tan powder (0.69 Wt, 0.66 eq, dr= 34:1). The mother liquor was concentrated to give an epimeric mixture (Compound AS) as yellow solid (epimeric mixture, 0.38 Wt, dr Compound AD: epimer =1:2.2).

Example 2: Diastereomeric Purification of Compound AD from Compound AS



Scheme 12

Each of the following methods was used in the reaction shown in Scheme 12 to
5 convert the undesired C25 epimer to the desired C25 isomer, using either stereoselective deprotonation-protonation or crystallization induced dynamic resolution (CIDR).

Method 1: Compound AS (1 Wt, 1 V, dr= 1:2.2) was dissolved in toluene (2.6 V) and cooled to -20 °C. KHMDS (0.50 M solution in toluene, 3.4 V, 0.60 eq) was added at a rate such that internal temperature did not exceed -16 °C. Upon complete addition,
10 stirring was continued at -20 °C for 15 minutes. Under vigorous stirring, 20 wt% NH₄Cl aq (1.0 Wt, 1.3 eq) was added at a rate such that internal temperature did not exceed -15 °C. After 5 minutes, the mixture was allowed to warm to 0 °C. The organic layer was separated, washed with water (2.0 V) and concentrated. The residual solvents and water were azeotropically removed with heptane (3.0 V x 2) to give the crude product as
15 yellow solid-oil mixture (dr= 2.6:1). The crude was suspended in Heptane-MTBE (5:1, v/v, 3.0 V) and heated to 80 °C. The resultant clear solution was cooled to room temperature (23 °C) over 3 hours (precipitation started at 45 °C). The crystals were collected by filtration, washed with: 1) heptane-MTBE (5:1 v/v, 1.0 V); 2) heptane (1.0 V) and dried under N₂/vacuum to give Compound AD as white powder (0.31 Wt, 0.31
20 eq, 0.08 eq). The mother liquor was concentrated to give Compound AS (0.69 Wt, dr= 1:1).

Method 2: Compound AS (1 Wt, 1 V, dr= 1:1) was dissolved in heptane-MTBE (5:1 v/v, 2.0 V) and KHMDS (0.50 M solution in toluene, 0.40 V, 0.07 eq) was added at 23 °C. Stirring was continued for 10 minutes and the mixture was cooled to 0 °C.
25 Compound AD (0.0001 Wt, 0.0001 eq) was added and stirring was continued for an additional 30 minutes (precipitation increased). 20 wt% NH₄Cl aq (0.20 Wt, 0.26 eq) was added under vigorous stirring. The resultant mixture was diluted with EtOAc (2.0 V) to dissolve Compound AD precipitation. The organic layer was separated, washed with water (1.0 V) and concentrated. The residual solvents and water were
30 azeotropically removed with heptane (5 V x 2) to give crude product as yellow solid-oil mixture (dr= 2.3:1). The crude was suspended in heptane-MTBE (3:1 v/v, 1.5 V) and heated to 80 °C. The resultant clear solution was cooled to 20 °C over 3 hours (precipitation started at 50 °C). The crystals were collected by filtration, washed with

heptane-MTBE (4:1 v/v, 1 V), and dried under N₂/vacuum to give Compound AD as white powder (0.22 Wt, 0.22 eq, 0.04 eq).

Method 3 (CIDR): Compound AS (1 Wt, 1 V, dr= 1:5) was dissolved in heptane (5 V) at 23 °C. t-BuOK (1.0 M solution in THF, 0.29 V, 0.10 eq) was added and stirring was continued for 10 minutes. The precipitations were collected by filtration, washed with heptane (10 V), and dried to give Compound AD as light tan powder (0.36 Wt, 0.36 eq, dr= 7.3:1, filtrate dr= 3.7:1).

Method 4: Compound AS (1 Wt, 1 V, 1 eq, dr=1:1.7) was dissolved in toluene (5.0V) and cooled to -70~-75 °C. KHMDS (0.5M solution in toluene, 0.500 eq, 2.88 V, 2.53 Wts) was added while maintaining an internal temperature below -65 °C. The resultant mixture was cooled to -70~-75 °C again and stirring was continued at -70~-75 °C for 4 hours. 20 wt% NH₄Cl (aqueous solution, 2.00 Wts) was added while maintaining an internal temperature below -60 °C. Upon complete addition, the mixture was allowed to warm to 0 °C over a period of 1.5~2 hours. MTBE (4.00 V, 2.96 Wt) and water (4.00 V, 4.00Wt) were added under stirring and the resultant biphasic mixture was allowed to partition. Organic layer (dr=6.5:1) was separated, sequentially washed with: 1) 20 wt% citric acid (aqueous solution, 1.0 Wt); 2) water (3.00 V); 3) water (3.00 V) and partially concentrated to ~2V under vacuum. The residue was subjected to solvent exchange with heptane (6.00 V x 2, partial concentration to ~2V each time, under vacuum) and diluted with heptane-IPA (6:1 v/v, 3.5V). The mixture was heated to 60 °C, cooled to 15~20 °C over 4 hours, and further stirred at 15~20 °C overnight. Crystals were collected by filtration, rinsed with heptane-IPA (9:1 v/v, 2.0 V) and dried under N₂/vacuum to give Compound AD (0.4 Wt, 0.4 eq, dr=57:1) as light tan powder.


¹H NMR (500 MHz, CDCl₃)
δ 4.40-4.44 (1H, m), 4.30 (1H, dd, J= 6.5, 3.5 Hz), 4.09 (1H, dd, J= 6.5, 3.0 Hz), 3.72-3.77 (1H, m); 3.37 (1H, dd, J= 10.0, 6.5 Hz), 2.91-2.99 (1H, m), 2.35-2.39 (1H, m), 2.07-2.12 (1H, m), 1.97-2.03 (1H, m), 1.96 (1H, dd, J= 14.0, 4.0 Hz), 1.82 (1H, d, J= 12.0 Hz), 1.58-1.70 (5H, m), 1.50-1.58 (6H, m), 1.42-1.49 (1H, m), 1.32-1.40 (2H, m), 1.29 (3H, d, J= 7.0 Hz), 1.11-1.20 (1H, m)

¹³C NMR (125 MHz, CDCl₃)
δ 122.95, 110.58, 78.29, 76.28, 75.92, 75.81, 72.16, 68.34, 43.80, 40.51, 37.61, 34.52, 29.85, 28.92, 27.24, 25.33, 24.24, 23.84, 22.50, 18.55

LRMS (ESI) m/z found 370.15 [M+Na]⁺

Melting Point 123 °C


Example 3: Synthesis of Compound AJ from Compound AD



LRMS (ESI) m/z found 289.95 [M+Na]⁺

Melting Point 189 °C

5 **Example 4: Synthesis of Compound AK from Compound AJ**



Scheme 14

- Compound AJ (1 Wt, 1 V, 1 eq) was suspended in acetonitrile (5.00 V) and
- 10 cooled to 0 °C. 2-acetoxy-2-methylpropionyl bromide (0.938 Wt, 0.656 V) was added at a rate such that the internal temperature did not exceed 7 °C. Upon complete addition, water (0.002 V, 3 mol%) was added and stirring was continued at 0 °C for an additional hour. The reaction mixture was diluted with MTBE (5.0 V). After internal temperature dropped to 0 °C, 10 wt% NaHCO₃ aq (5.0 V, 3.4 eq) was carefully added under vigorous
- 15 stirring maintaining internal temperature below 7°C and the resultant mixture was allowed to partition. The organic layer was set aside and the aqueous layer was extracted with MTBE (5.0 V). All of the organic layers were combined, sequentially washed with: 1) 10 wt% NaHCO₃ aq (2.0 V, 1.4 eq); 2) water (2.0 V); 3) brine (2.0 V), and concentrated to give crude Compound AK as light brown oil (1.47 Wt, 1.04 eq).
- 20 The crude product was azeotropically dried with toluene (4 V x 3) and used for next reaction without purification.

¹H NMR (500 MHz, CDCl₃)


- δ 5.20 (1H, br), 4.38 (1H, dd, J= 6.5, 3.5 Hz), 4.21-4.23 (1H, m), 4.04 (1H, dd, J= 10.0, 7.0 Hz), 3.79-3.83 (1H, m), 2.90-2.98 (1H, m), 2.51-2.56 (2H, m), 2.30-2.34 (1H, m), 2.11-2.15 (1H, m), 2.07 (3H, s), 1.65-1.71 (1H, m), 1.57-1.62 (3H, m), 1.49-1.55 (1H, m), 1.32 (3H, d, J= 6.5 Hz), 1.21-1.30 (1H, m)
- 25

¹³C NMR (125 MHz, CDCl₃)

- δ 169.39, 122.79, 78.13, 75.49, 75.42, 73.76, 68.45, 44.66, 43.48, 40.11, 29.48, 28.88, 28.38, 22.40, 21.12, 18.46
- 30

LRMS (ESI) m/z found 393.96 [M+Na]⁺

Example 5: Synthesis of Compound AL from Compound AJ



Scheme 15


5

Compound AJ (1 Wt, 1 V, 1 eq) was suspended in acetonitrile (3.0 V) and cooled to 0 °C. 2-acetoxy-2-methylpropionyl bromide (1.02 Wt, 1.30 eq) was added at a rate such that the internal temperature did not exceed 2 °C. Upon complete addition, an acetonitrile-water mixture (water (0.0020 V, 0.030 eq) and acetonitrile (0.020 V)) were added and stirring was continued at 0 °C for 2 hours. Under vigorous stirring, 10 wt% NaHCO₃ aq (5.0 V) was added at a rate such that the internal temperature did not exceed 8 °C (CO₂ evolution). Toluene (4.3 Wt, 5.0 V) was added and vigorous stirring was continued for 3 minutes. The mixture was allowed to partition and the organic layer was set aside. The aqueous layer was extracted with toluene (2.6 Wt, 3.0 V). All of the organic layers were combined and sequentially washed with: 1) 10 wt% NaHCO₃ aq (3.0 V); 2) water (2.0 V).

The organic layer was transferred to a reactor and subjected to distillation under atmospheric pressure to remove 5 Wt of solvent. The distillation included heating the organic layer to 90 °C to remove the acetonitrile and then heating the mixture to about 110 °C to remove the toluene. After cooling to 80 °C, toluene (2.50 Wt, 3 V) was added followed by DBU (1.12 V, 1.14 Wt, 2.00 eq). The mixture was re-heated to 100 °C and vigorously stirred for 17 hours. The reaction mixture was cooled to 0 °C and 1.00 M HCl aq (4.5 V, 1.2 eq) was added at a rate such that the internal temperature did not exceed 8 °C. The resultant mixture was allowed to partition. The organic layer was set aside and aqueous layer was extracted with toluene (1.73 Wt, 2.0 V). All of the organic layers were combined, sequentially washed with: 1) 1.00 M HCl aq (0.50 V, 0.13 eq); 2) 10 wt% NaHCO₃ aq (1.0 V); 3) water (2.0 Wt, 2.0 V), and concentrated. The residual toluene was azeotropically removed with IPA (2.0 V) to give the crude product as yellow solid. Crude Compound AL was suspended in IPA (5.0 V) and heated to 80 °C. The resultant solution was cooled to 0 °C over 2 hours and stirring was continued at 0 °C for an additional 30 minutes. Crystals were collected by filtration, washed with cold IPA (1 V) followed by heptane (1 V), and dried to give Compound AL as white powder (0.64 Wt, 0.59 eq). The mother liquor was concentrated and diluted with IPA-heptane (1:1 v/v, 1.0 V). A white precipitation was formed, collected by filtration, washed with:

1) IPA-heptane (1:1 v/v, 0.4 V); 2) heptane (0.4 V), and dried to give additional Compound AL (0.043 Wt, 0.040 eq).

Example 6: Synthesis of Compound AL from Compound AK



Compound AK (1 Wt, 1 V, 1 eq) was dissolved in toluene (5.0 V). DBU (0.818 Wt, 0.803 V, 2.0 eq) was added at 23°C and the mixture was heated to 100 °C. Upon
 10 complete consumption of Compound AK, the reaction mixture was cooled to 10°C and 1M HCl (3.5 V, 1.3 eq) was added. The resultant mixture was vigorously stirred for 5 minutes and allowed to partition. The organic layer was set aside and the aqueous layer was extracted with MTBE (5.0 V). All organic layers were combined, sequentially washed with: 1) water (2.0 V); 2) 10 wt% NaHCO₃ solution (2.0 V); 3) water (2.0 V),
 15 and concentrated to give a mixture of light brown oil and water. The residual water was azeotropically removed with heptane (3.0 V x 3) to give crude Compound AL as yellow solid (0.65 Wt, 0.83 eq)


¹H NMR (500 MHz, CDCl₃)
 20 δ 6.16 (1H, d, J= 10 Hz), 5.60-5.63 (1H, m), 5.01-5.02 (1H, m), 4.34-4.36 (1H, m), 3.80-3.85 (1H, m), 3.42 (1H, dd, J= 5.0, 2.0 Hz), 2.93-3.01 (1H, m), 2.53-2.57 (1H, m), 2.07-2.12 (1H, m) 2.03 (3H, s), 1.56-1.72 (4H, m), 1.49-1.55 (1H, m), 1.32 (3H, d, J= 6.5 Hz), 1.22-1.30 (1H, m)

25 ¹³C NMR (125 MHz, CDCl₃)
 δ 170.10, 142.16, 122.82, 122.42, 79.43, 75.26, 74.81, 69.52, 68.48, 40.36, 29.62, 28.90, 28.77, 22.49, 21.26, 18.54

LRMS (ESI) m/z found 314.04 [M+Na]⁺
 30

Melting Point 92 °C

Example 7: Synthesis of Compound AM from Compound AL



Scheme 17

5

Compound AL (1 Wt, 1 V, 1 eq) was dissolved in MeOH-DCM (5:3 v/v, 8.0 V) and cooled to -47 °C. O₃ was bubbled into the mixture maintaining an internal temperature below -42 °C. Upon complete consumption of Compound AL, excess O₃ was purged by N₂ bubbling until the peroxide test for the reactor outlet was negative.

10 The reaction mixture was then allowed to warm to -25 °C and NaBH₄ (0.0753 Wt, 0.580 eq) was added while maintaining an internal temperature below -17 °C. Upon complete addition, the mixture was stirred at -20 °C for 1 hour and then allowed to warm to 0 °C. NaBH₄ (granules, 0.0753 Wt, 0.580 eq) was added (while maintaining an internal temperature below 3 °C) and stirring was continued at 0 °C for one hour.

15 K₂CO₃ (0.712 Wt, 1.50 eq) was added at 0 °C and the reaction was allowed to warm to 20 °C. Upon complete consumption of the acetate intermediate (approx 4 hours), the reaction mixture was cooled to 0 °C and 10 wt% HCl aq (5.1 Wt, 4.1 eq) was added under vigorous stirring to adjust the pH to 6-7.

The resultant biphasic mixture was partially concentrated (to approx 5.6 Wt) for removal volatiles, re-diluted with water-THF (1:1 v/v, 4.0 V), and cooled to 15 °C. NaIO₄ (1.47 Wt, 2.00 eq) was added and the resultant slurry was stirred at 20 °C until complete consumption of triol (approx. 3 hours). The reaction mixture was then diluted with EtOAc (6.0 V), stirred vigorously for 5 minutes, and filtered through a pad of Celite (2 Wt). The filtrate (F-1) was separated and set aside and the filter cake was washed with EtOAc-EtOH (9:1 v/v, 4.0 V) (filtrate: F-2). NaCl (1.0 Wt) was added to F-1 and the resultant mixture was stirred vigorously for 5 minutes and allowed to partition. The organic layer was set aside and the aqueous layer was extracted with F-2. All of the organic layers were combined, sequentially washed with: 1) 10 wt% Na₂S₂O₃ aq (1.0 Wt); 2) water (1.0 V); 3) water (1.0 V) and concentrated to give a white solid.

25 The residual water and solvents were azeotropically removed with EtOAc (6.0 V x 3) to give the crude product as white solid (0.84 Wt, 0.96 eq). The crude was suspended in heptane-EtOAc (1:1 v/v, 3.5 V) and heated to 80 °C. The resultant solution was cooled to room temperature over 2 hours (the precipitation started at ~65 °C). The mixture was further cooled to 0 °C and stirring was continued for an additional hour. The crystals

were collected by filtration, washed with cold heptane-EtOAc (1:1 v/v, 1.8 V), and dried under N₂/vacuum to give Compound AM as white powder (0.58 Wt, 0.67 eq). The mother liquor was concentrated, suspended in heptane-EtOAc (4:3 v/v, 0.9 V), and heated to 80 °C. The resultant clear solution was cooled to 20 °C over 2 hours. The mixture was further cooled to 0 °C and stirring was continued for an additional hour. The crystals were collected by filtration, washed with cold heptane-EtOAc (4:3 v/v, 0.50 V) and dried under N₂/vacuum to give additional Compound AM as white powder (0.068 Wt, 0.08 eq).


¹H NMR (for major anomer, 500 MHz, CDCl₃)
 δ 4.96 (1H, s), 4.17 (1H, dd, J=6.0, 3.5 Hz), 3.90 (1H, d, J=9.5 Hz), 3.82-3.74 (2H, m), 3.41 (1H, dd, J= 10, 3.0 Hz), 3.01 (1H, s), 2.95-2.85 (1H, m), 2.51-2.45 (1H, m), 2.22-2.15 (1H, m), 1.72-1.64 (1H, m), 1.63-1.48 (3H, m), 1.29 (3H, d, J=13 Hz), 1.30-1.18 (1H, m)

¹³C NMR (for major anomer, 125 MHz, CDCl₃)
 δ 122.81, 92.46, 77.17, 75.70, 72.43, 71.18, 68.36, 40.28, 29.82, 28.70, 28.40, 22.42, 18.52

LRMS (ESI) m/z found 307.99 [M+MeOH+Na]⁺

Melting Point 116 °C

Example 8: Synthesis of Compound AN from Compound AM



Compound AM

Compound AN

Scheme 18

Compound AM (1 Wt, 1 V, 1 eq) was suspended in acetonitrile (4.0 V) and cooled to 10 °C. LiCl (0.184 Wt, 1.10 eq) was added followed by N,N-diisopropylethylamine (0.825 V, 1.20 eq). After the internal temperature dropped to 10 °C, trimethyl phosphonoacetate (0.703 V, 1.10 eq) was added at a rate such that the internal temperature did not exceed 13 °C. Upon complete addition, the reaction was stirred at 10 °C for one hour and was then allowed to warm to 20 °C. Stirring was continued at 20 °C until complete consumption of Compound AM. The reaction

¹H NMR (500 MHz, CDCl₃)


δ 4.18-4.23 (1H, m), 3.82-3.91 (2H, m), 3.67 (3H, s), 3.53 (2H, d, J= 6.5 Hz), 2.86-2.93 (1H, m), 2.40-2.46 (1H, m), 2.31-2.38 (2H, m), 2.17 (1H, t, J= 7.0 Hz), 1.85-1.92 (1H, m), 1.59-1.84 (6H, m), 1.49 (1H, dd, J= 14.0, 5.5 Hz), 1.32 (3H, d, J= 7.5 Hz), 1.23-1.30 (1H, m)

¹³C NMR (125 MHz, CDCl₃)

δ 173.90, 123.10, 84.23, 74.90, 73.28, 68.31, 67.73, 51.81, 40.28, 35.99, 31.75, 30.78, 27.12, 24.03, 22.27, 18.32

LRMS (ESI) m/z found 334.08 [M+Na]⁺

Example 10: Synthesis of Compound AF from Compound AO



Scheme 20

Compound AO (1 Wt, 1 V, 1 eq) was dissolved in DCM (4.50 V). TEA (1.16 V, 0.84 Wt, 2.60 eq) was added and the mixture was cooled to -70°C . A solution of Tf_2O (0.702 V, 1.30 eq) in DCM (1.50 V) was added at a rate such that the internal temperature did not exceed -65°C . Upon complete addition, the reaction was stirred at -73°C for 1.5 hours, allowed to warm to -20°C , and stirred at -20°C for an additional 30 minutes.

DMF (3.0 V) was added and the mixture was allowed to warm to 0°C . NaI (0.674 Wt, 1.40 eq) was added and the reaction was further warmed to 23°C . Upon complete consumption of the triflate (Compound AT), the reaction mixture was diluted with heptane (8.0 V) and cooled to 0°C . Water (9.0 V) was added while maintaining an internal temperature below 10°C . The resultant biphasic mixture was stirred vigorously for 3 minutes and then allowed to partition. The organic layer was set aside and the aqueous layer was extracted with MTBE (6.0 V). All of the organic layers were combined, sequentially washed with: 1) 1.00 M HCl aq (5.00 V, 1.56eq); 2) 10 wt%

NaHCO₃ aq (2.0 V); 3) 10 wt% Na₂S₂O₃ aq (2.0 V), 4) water (2.0 V); 5) water (2.0 V) and concentrated. The residue was dissolved in MTBE (6.0 V) and silica gel (1.0 Wt) was added. The resultant slurry was stirred at 22 °C for 5 minutes and then filtered. The silica gel on the filter was washed with MTBE (8.0 V) and the filtrate was concentrated to give crude product as reddish solid (1.35 Wt, 1.00 eq).

Compound AF (1.35 Wt, 1.00 eq) was suspended in MTBE (1.4 V) and heated to 45 °C. Heptane (2.8 V) was added while maintaining an internal temperature between 40 °C and 45 °C. The resultant clear solution was cooled to 22 °C over 1 hour and then stirred at 22 °C for 2 hours. The mixture was cooled to 0 °C and stirring was continued for an additional 2 hours. The precipitations were collected by filtration, washed with pre-cooled (0 °C) heptane-MTBE (1:3 v/v, 2.8 V) and dried under N₂/vacuum for one hour to give Compound AF as light tan powder (0.98 Wt, 0.72 eq). The mother liquor was concentrated and re-dissolved in MTBE (0.33 V). Heptane (0.33 V) was added and the resultant clear solution was cooled to 0 °C. A very small amount of Compound AF crystal (from the 1st crop) was added for seeding and stirring was continued at 0 °C for 15 hours. The precipitations were collected by filtration, washed with pre-cooled (0 °C) heptane-MTBE (1:2 v/v, 0.33 V), and dried under N₂/vacuum for 1 hour to give additional Compound AF as light tan powder (0.046 Wt, 0.034 eq).

20 Compound AT

¹H NMR (500 MHz, CDCl₃)

δ 4.46 (1H, d, J= 10.5 Hz), 4.38 (1H, d, J= 10.5 Hz), 4.21-4.26 (1H, m), 3.89 (1H, dd, J= 8.5, 6.0 Hz), 3.81-3.86 (1H, m), 3.68 (3H, s), 2.93-3.00 (1H, m), 2.41-2.50 (2H, m), 2.33-2.39 (1H, m), 1.91-1.97 (1H, m), 1.64-1.92 (6H, m), 1.45 (1H, dd, J= 14.5, 5.5 Hz), 1.25-1.35 (1H, m), 1.32 (3H, d, J= 7.0 Hz)

¹³C NMR (125 MHz, CDCl₃)

δ 173.62, 122.86, 117.51, 81.84, 78.54, 74.57, 73.08, 68.63, 51.94, 40.16, 35.28, 31.77, 30.64, 27.13, 23.95, 22.33, 18.42

LRMS (ESI) m/z found 446.12 [M+Na]⁺

Compound AF

¹H NMR (500 MHz, CDCl₃)

δ 4.21-4.26 (1H, m), 3.78-3.83 (2H, m), 3.67 (3H, s), 3.44 (1H, d, J= 10.0 Hz), 3.37 (1H, d, J= 10.0 Hz), 2.99-3.03 (1H, m), 2.49 (1H, dd, J= 9.0, 8.5 Hz), 2.42-2.47 (1H, m), 2.32-2.38 (1H, m), 1.80-1.89 (3H, m), 1.63-1.75 (5H, m), 1.33 (3H, d, J= 7.5 Hz), 1.24-1.30 (1H, m)

¹³C NMR (125 MHz, CDCl₃)


δ 173.74, 122.89, 81.75, 76.07, 75.10, 68.24, 51.86, 40.52, 39.00, 31.78, 30.75, 27.09, 24.36, 22.53, 18.72, 18.51

5

LRMS (ESI) m/z found 444.02 [M+Na]⁺

Melting Point 69.5 °C

10 **Example 11: Synthesis of Compound AP from Compound AF**



Scheme 21

Compound AF (1 Wt, 1 V, 1 eq) was dissolved in toluene (5.0 V) and cooled to
15 10 °C. LiBH₄ (2.0 M solution in THF, 2.4 V, 2.0 eq) was added and stirring was
continued at 20 °C for 18 hours. The reaction mixture was cooled to 0 °C and slowly
poured into a pre-cooled (0 °C) mixture of EtOAc (6 V) and 1.0 M HCl aq (6.0 V, 2.5
eq) under vigorous stirring. The reactor was rinsed with EtOAc (2 V) and the resultant
wash was combined with the biphasic mixture. The organic layer was set aside and the
20 aqueous layer was extracted with EtOAc (5.0 V). All of the organic layers were
combined, sequentially washed with: 1) 10 wt% NaHCO₃ aq (2 V); 2) water (2 V) and
concentrated. The residual water was azeotropically removed with toluene (5 V x 2) to
give Compound AP (0.93 Wt, 0.89 eq).

25 ¹H NMR (500 MHz, CDCl₃)

δ 4.24-4.30 (1H, m), 3.86 (1H, dd, J= 8.5, 6.0 Hz), 3.78-3.83 (1H, m), 3.62-3.68 (2H, m), 3.44 (1H, d, J= 10.5 Hz), 3.38 (1H, d, J= 10.5 Hz), 2.99-3.04 (1H, m), 2.51 (1H, dd, J= 14.0, 8.5 Hz), 2.06 (1H, t, J= 6.0 Hz), 1.86-1.92 (1H, m), 1.59-1.78 (9H, m), 1.33 (3H, d, J= 7.0 Hz), 1.24-1.31 (1H, m)


30

¹³C NMR (125 MHz, CDCl₃)

δ 122.94, 82.70, 76.27, 76.25, 68.42, 62.77, 40.50, 39.08, 33.72, 29.67, 27.25, 24.59, 22.55, 19.08, 18.51

35 LRMS (ESI) m/z found 416.02 [M+Na]⁺

Example 12: Synthesis of Compound AU from Compound AP



Scheme 22

5

An inert reactor was charged with Zn powder (2.5 Wt, 15 eq) at 23 °C. MeOH (5.0 V) was added followed by AcOH (2.0 V, 14 eq). The resultant slurry was stirred at 23 °C for 20 minutes and then cooled to 0 °C. A solution of Compound AP (1 Wt, 1 V, 1 eq) in MeOH (5.0 V) was added and vigorous stirring was continued at 0 °C for 3 hours and at 23 °C for 1.5 hours. The reaction mixture was diluted with EtOAc (20 V). Excess Zn powder was removed by filtration and rinsed with EtOAc (10 V). The filtrate was washed with 1.00 M HCl aq (10 V). The organic layer was set aside and the aqueous layer was extracted with EtOAc (20 V). All of the organic layers were combined, sequentially washed with: 1) 10 wt% NaHCO₃ aq (20 V); 2) 10 wt% Na₂S₂O₃ aq (8 V); 3) brine (8 V), and concentrated to give the crude product as pale yellow oil. The crude was purified by flash column chromatography (Biotage, heptane-EtOAc 3:7→2:8→0:10) to give Compound AU (0.62 Wt, 0.90 eq) as pale yellow oil.

¹H NMR (500 MHz, CDCl₃)


20 δ 5.01 (1H, s), 4.85 (1H, s), 4.41 (1H, br), 4.08-4.12 (1H, m), 3.93 (1H, br), 3.60-3.68 (2H, m), 3.12 (1H, br), 2.97-3.05 (1H, m), 2.69-2.73 (1H, m), 2.45 (1H, br), 2.29-2.33 (1H, m), 1.53-1.80 (10H, m), 1.33 (3H, d, J= 7.5 Hz)

¹³C NMR (125 MHz, CDCl₃)

25 δ 150.95, 123.29, 105.49, 79.66, 77.69, 68.79, 62.84, 41.83, 39.03, 34.33, 32.11, 30.89, 29.80, 22.93, 18.61

LRMS (ESI) m/z found 289.96 [M+Na]⁺

Example 13: Synthesis of Compound AR from Compound AU



Scheme 23

5


Compound AU (1 Wt, 1 V, 1 eq) was dissolved in MeOH (2.0 V). The mixture was cooled to 0 °C and HCl (6 M solution in IPA, 2.0 V, 13 eq) was added. The reaction was allowed to warm to 23 °C and stirring was continued until complete consumption of Compound AU (approx. 20 hours). The reaction mixture was diluted with toluene (8.0 V) and water (4.0 V) and the resultant biphasic mixture was heated at 60°C for 3 hours. After cooling down, organic layer was set aside and aqueous layer was extracted with EtOAc (8.0 V). All organic layers were combined, sequentially washed with: 1) 10 wt% NaHCO₃ aq (2.0 V); 2) brine (2.0 V); 3) water (2.0 V), and concentrated to give crude Compound AR (0.93 Wt, 0.93 eq) as pale yellow oil. The crude product was azeotropically dried with toluene (8 V x 2) and used for next reaction without purification.

¹H NMR (500 MHz, CDCl₃)
 δ 4.98-4.99 (1H, m), 4.84-4.85 (1H, m), 4.49-4.54 (1H, m), 4.39 (1H, d, J=10.5 Hz),
 20 4.00-4.05 (1H, m), 3.59-3.68 (2H, m), 2.63-2.72 (2H, m), 2.56-2.62 (1H, m), 2.25-2.30 (1H, m), 2.08-2.14 (1H, m), 1.97-2.02 (1H, m), 1.52-1.82 (7H, m), 1.26 (3H, d, J= 7.5 Hz), 1.24-1.34 (1H, m)

¹³C NMR (125 MHz, CDCl₃)
 25 δ 180.34, 151.14, 105.39, 79.71, 78.84, 77.54, 62.72, 39.02, 35.69, 34.15, 32.19, 32.16, 31.50, 29.64, 16.02

LRMS (ESI) m/z found 290.99 [M+Na]⁺

Example 14: Synthesis of Compound AH from Compound AR



Scheme 24

5

Compound AR (1 Wt, 1 V, 1 eq) was dissolved in DMF (2.0 vols) and imidazole (0.330 Wt, 1.30 eq) was added at 23 °C (endothermic). Upon complete dissolution of imidazole, the mixture was cooled to 10 °C and tert-butylchlorodiphenylsilane (TBDPSCI, 0.969 V, 1.02 Wt, 1.00 eq) was added. The reaction mixture was stirred at 10 °C for 1 hour, allowed to warm to 23 °C, and stirred until complete consumption of Compound AR (approx. 3 hours). The reaction mixture was diluted with Heptane-MTBE 1:1 (8.0 V) and cooled to 10 °C. Water (8.0 V) was added under vigorous stirring and the resultant mixture was allowed to partition. The aqueous layer was set aside. The organic layer was further washed with water (1.0 V) and concentrated.

Residual water and solvents were azeotropically removed with toluene (8.0 V x 2) to give Compound AH as colorless oil (1.98 Wt, 100%). The crude product was used for next reaction without purification

- ¹H NMR (500 MHz, CDCl₃)
- δ 7.65-7.67 (4H, m), 7.36-7.44 (6H, m), 4.99 (1H, dd, J= 4.0, 2.5 Hz), 4.84 (1H, dd, J= 4.0, 2.5 Hz), 4.50-4.55 (1H, m), 4.35 (1H, d, J= 9.0 Hz), 3.97-4.02 (1H, m), 3.66-3.70 (2H, m), 2.66-2.71 (1H, m), 2.61-2.66 (1H, m), 2.22-2.27 (1H, m), 2.08-2.14 (1H, m), 1.97-2.03 (1H, m), 1.50-1.81 (8H, m), 1.28 (3H, d, J= 7.5 Hz), 1.04 (9H, s)
- ¹³C NMR (125 MHz, CDCl₃)
- δ 180.18, 151.68, 135.79 (4C), 134.21 (2C), 129.84 (2C), 127.89 (4C), 105.27, 79.58, 78.83, 77.38, 64.02, 39.08, 35.78, 34.20, 32.29, 31.76, 31.60, 29.31, 27.16 (3C), 19.48, 16.15
- LRMS (ESI) m/z found 529.26 [M+Na]⁺

Compound AH

1) $(\text{CH}_3\text{O})\text{NHCH}_3 \cdot \text{HCl}$
 Me_3Al , CH_2Cl_2
 0°C

2) TBSCl, imidazole
 DMF, rt

Compound AV

5

20

25


EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures described
5 herein. Such equivalents are considered to be within the scope of the present invention and are covered by the following claims. The contents of all references, patents, and patent applications cited throughout this application are hereby incorporated by reference. The appropriate components, processes, and methods of those patents,
10 applications and other documents may be selected for the present invention and embodiments thereof.

CLAIMS

1. A method of obtaining a substantially diastereomerically pure composition of a compound of formula (I), comprising:

crystallizing said compound of formula (I) from a mixture of diastereomers under appropriate crystallization conditions, wherein said compound of formula (I) is:



wherein:

z is a single or double bond, provided that when z is a double bond, X² is C and Y¹ is hydrogen; and provided that when z is a single bond, X² is CH or O;

X¹ is O, S, or CN, provided that when X¹ is CN or S, X² is O;

Y¹ is a halide, hydrogen or O-L², or absent when X² is O;


L¹ and L² are independently selected from hydrogen and a protecting group, or L¹ and L² together are a protecting group, provided that when X¹ is CN, L¹ is absent; or a salt thereof,

wherein the protecting group of L¹ and L² is silyl, C₁-C₁₂ alkylcarbonyl, C₁-C₆ alkyl, aryl (C₁-C₆) alkyl, carbonate, C₁-C₆ alkoxy-(C₁-C₆) alkyl, 3,4-dimethoxybenzyl, p-methoxybenzyl, benzoyl, or C₁-C₆ ester, or L¹ and L² together are a protecting group selected from benzylidene, cyclic C₁-C₆ acetal, cyclic C₃-C₆ ketal, cyclic carbonate, and a C₄-C₇ alkyl ring with one member of the ring covalently linked to both the O of Y¹ and to X¹; or when X¹ and X² are O, L¹ and L² together are cyclic C₃-C₇ ketal.

2. The method of claim 1, wherein at least one of L¹ and L² is a protecting group, and is independently selected from the group consisting of C₁-C₆ silyl, C₁-C₆ alkyl, benzoyl, and C₁-C₈ alkyl substituted carbonyl.

3. The method of claim 1, wherein Y¹ is O-L²; and L¹ and L² together form a protecting group which is a C₄-C₇ alkyl ring with one member of the ring covalently linked to both the O of Y¹ and to X¹.


4. The method of claim 1, wherein said compound of formula (I) is of formula (Ib):




wherein L^{1a} and L^{1b} are independently selected from hydrogen and a protecting group, or L^{1a} and L^{1b} together are a divalent protecting group.

5. The method of claim 4, wherein said divalent protecting group is a cyclohexylidene protecting group.


6. The method of claim 1, wherein said compound of formula (I) is selected from the group consisting of:




Compound AD




Compound AJ



Compound AL



Compound AM and



Compound AK.


7. The method of claim 1, wherein said appropriate crystallization conditions comprise heptane as an anti-solvent.

8. The method of claim 7, wherein appropriate crystallization conditions further comprise a co-solvent selected from *tert*-butyl methyl ether and isopropanol.

9. The method of claim 1, wherein said appropriate crystallization conditions comprise the addition of a base.

10. The method of claim 9, wherein said base is potassium *tert*-butoxide.

11. The method of claim 1, wherein said composition comprises a ratio of compound of formula (I) to compound of formula (Ie) of at least about 20:1, wherein said compound of formula (Ie) is:



(Ie).

12. The method of claim 11, wherein said ratio is at least 30:1.

13. The method of claim 1, wherein L^1 and L^2 are independently a protecting group selected from C_1 - C_{12} alkylcarbonyl, C_1 - C_6 alkyl, C_1 - C_{15} alkyl silyl, aryl (C_1 - C_6) alkyl, carbonate and C_1 - C_6 alkoxy- $(C_1$ - $C_6)$ alkyl groups.


14. The method of claim 1, wherein L^1 and L^2 together are cyclic C_1 - C_6 acetal, cyclic C_3 - C_6 ketal, or cyclic carbonate.

15. The method of claim 1, wherein at least one of L^1 and L^2 is C_1 - C_6 ester.

16. The method of claim 1, wherein at least one of L^1 and L^2 is a protecting group selected from methoxymethyl, trimethylsilyl, triethylsilyl, t-butyldimethylsilyl, t-butyldiphenylsilyl, triisopropylsilyl, methyl, t-butyl, 3,4-dimethoxybenzyl, p-methoxybenzyl, benzyl, and trityl, or L^1 and L^2 together are cycloheptylidene.


17. The method of claim 1, wherein at least one of L^1 and L^2 is C_1 alkyl carbonyl, or L^1 and L^2 together are acetonide, benzylidene, cyclohexylidene, or cyclopentylidene.

18. A method of making a substantially diastereomerically pure composition of a compound of formula (Ib) from a compound of formula (Ia), wherein the compound of formula (Ia) is:



(Ia)

and the compound of formula (Ib) is:



(Ib)


wherein L^{1a} and L^{1b} are independently selected from hydrogen and a protecting group selected from C_1 - C_6 alkyl, aryl (C_1 - C_6) alkyl, silyl (C_1 - C_{10}), and C_1 - C_6 alkyl ester, or L^{1a} and L^{1b} together are a divalent protecting group selected from cyclic C_1 - C_6 acetal, cyclic C_3 - C_7 ketal, and cyclic carbonate, provided that L^{1a} of formulae (Ia) and (Ib) are the same and L^{1b} of formulae (Ia) and (Ib) are the same, said method comprising:

reacting the compound of formula (Ia) under alkylating conditions to form a mixture comprising the compound of formula (Ib) and diastereomers thereof; and

crystallizing the compound of formula (Ib) from the mixture, under appropriate crystallization conditions.


19. The method of claim 18, wherein said alkylating conditions comprise an aprotic solvent, a base, and a C_1 alkyl halide or C_1 - C_7 methyl sulfonate.

20. The method of claim 19, wherein the C₁-C₇ sulfonate is of formula R-SO₂-O-Me, in which R is C₁-C₄ alkyl or C₆ or C₇ aryl.
21. The method of claim 19, wherein said aprotic solvent is tetrahydrofuran, toluene, or t-butyl methyl ether.
22. The method of claim 18, wherein said appropriate crystallization conditions comprise the addition of a base.
23. The method of claim 22, wherein said base is lithium diisopropyl amide, potassium-hexamethyldisilazane, or potassium t-butoxide.
24. The method of claim 18, wherein said composition comprises a ratio of at least about 20:1 of compound of formula (1a) to compound of formula:



25. The method of claim 18, wherein L^{1a} and L^{1b} are each protecting groups selected from C₁-C₆ alkyl, aryl (C₁-C₆) alkyl, silyl (C₁-C₁₀), and C₁-C₆ alkyl ester, or L^{1a} and L^{1b} together are cyclic C₁-C₆ acetal, cyclic C₃-C₇ ketal, or cyclic carbonate.
26. The method of claim 18, wherein L^{1a} and L^{1b} together are a cyclohexylidene protecting group.

27. A compound of formula (I):



wherein:

z is a single or double bond, provided that when z is a double bond, X² is C and Y¹ is hydrogen;
and provided that when z is single bond, X² is CH or O;

X¹ is O, S, or CN, provided that when X¹ is CN or S, X² is O;

Y¹ is a halide, hydrogen or O-L², or absent when X² is O; and


L¹ and L² are independently selected from hydrogen and a protecting group, or L¹ and L² together are a protecting group, provided that when X¹ is CN, L¹ is absent; or a salt thereof,

wherein the protecting group of L¹ and L² is silyl, C₁-C₁₂ alkylcarbonyl, C₁-C₆ alkyl, aryl (C₁-C₆) alkyl, carbonate, C₁-C₆ alkoxy-(C₁-C₆) alkyl, 3,4-dimethoxybenzyl, p-methoxybenzyl, benzoyl, or C₁-C₆ ester, or L¹ and L² together are a protecting group selected from benzyldiene, cyclic C₁-C₆ acetal, cyclic C₃-C₆ ketal, cyclic carbonate, and a C₄-C₇ alkyl ring with one member of the ring covalently linked to both the O of Y¹ and to X¹; or when X¹ and X² are O, L¹ and L² together are cyclic C₃-C₇ ketal.

28. The compound of claim 27, wherein at least one of L¹ and L² is a protecting group and is silyl, C₁-C₆ alkyl, or C₁-C₆ alkyl substituted carbonyl.


29. The compound of claim 27, wherein Y¹ is O-L²; and L¹ and L² together form protecting group which is a C₄-C₇ alkyl ring with one member of the ring covalently linked to both the O of Y¹ and to X¹.

30. The compound of claim 27, wherein said compound is:




Compound AD.


31. The compound of claim 27, wherein said compound is selected from the group consisting of:




Compound AJ



Compound AK



Compound AL and




Compound AM.

32. The compound of claim 27, wherein at least one of L^1 and L^2 is a protecting group selected from methoxymethyl, trimethylsilyl, triethylsilyl, t-butyldimethylsilyl, t-butyldiphenylsilyl, triisopropylsilyl, methyl, t-butyl, 3,4-dimethoxybenzyl, p-methoxybenzyl, benzyl, and trityl, or L^1 and L^2 together are cycloheptylidene.

33. The compound of claim 27, wherein at least one of L^1 and L^2 is C_1 alkyl carbonyl, or L^1 and L^2 together are acetonide, benzylidene, cyclohexylidene, or cyclopentylidene.


34. The compound of claim 27, wherein said compound of formula (I) is of formula (Ib):



(Ib)

wherein L^{1a} and L^{1b} are independently selected from hydrogen and a protecting group selected from C_1 - C_6 alkyl, aryl (C_1 - C_6) alkyl, silyl (C_1 - C_{10}), and C_1 - C_6 alkyl ester, or L^{1a} and L^{1b} together are a divalent protecting group selected from cyclic C_1 - C_6 acetal, cyclic C_3 - C_7 ketal, and cyclic carbonate.

35. A compound of the formula (II):



wherein:

c is a single or double bond, provided that when c is a double bond, m is 0, and Y^3 is O or $CHCO_2-L^3$, and provided that, when c is a single bond, m is 0 or 1, and Y^3 is CH_2O-L^3 , $CH_2CO_2-L^3$ or $CH_2CH_2O-L^3$;

Y^2 is C_1 - C_7 sulfonate, $O-L^4$ or a halide;


L^4 is hydrogen or a protecting group selected from C_1 - C_{12} alkylcarbonyl, C_1 - C_6 alkyl, C_1 - C_{15} alkyl silyl, *t*-butyldiphenylsilyl, aryl (C_1 - C_6) alkyl, 3,4-dimethoxybenzyl, *p*-methoxybenzyl, carbonate, and C_1 - C_6 alkoxy-(C_1 - C_6) alkyl groups; and

L^3 and L^5 are each independently hydrogen or a protecting group selected from C_1 - C_{12} alkylcarbonyl, C_1 - C_6 alkyl, C_1 - C_{15} alkyl silyl, *t*-butyldiphenylsilyl, aryl (C_1 - C_6) alkyl, 3,4-dimethoxybenzyl, *p*-methoxybenzyl, carbonate, and C_1 - C_6 alkoxy-(C_1 - C_6) alkyl groups, or L^3 and L^5 together are a protecting group selected from cyclic C_1 - C_6 acetal, cyclic C_3 - C_7 ketal, and cyclic carbonate, or, when Y^3 is $CH_2CO_2-L^3$, L^3 is C_1 - C_{10} alkyl, C_6 - C_{10} aryl- C_1 - C_6 alkyl, or C_6 - C_{10} aryl; or a salt thereof.


36. The compound of claim 35, wherein Y^2 is a halide.

37. The compound of claim 36, wherein said halide is iodide.


38. The compound of claim 35, wherein said compound is selected from the group consisting of:




Compound AN




Compound AO




Compound AF



Compound AP,




Compound AQ, and




Compound AT

39. The compound of claim 35, wherein Y^2 is trifluoromethyl sulfonate or p-toluyyl sulfonate.
40. The compound of claim 35, wherein L^3 , L^4 , and L^5 are each independently hydrogen or a protecting group selected from C_1 - C_{12} alkylcarbonyl, C_1 - C_6 alkyl, C_1 - C_{15} alkyl silyl, aryl (C_1 - C_6) alkyl, carbonate, and C_1 - C_6 alkoxy- $(C_1$ - $C_6)$ alkyl groups, or L^3 and L^5 together are a protecting group selected from cyclic C_1 - C_6 acetal, cyclic C_3 - C_7 ketal, cyclic carbonate, or, when Y^3 is $CH_2CO_2-L^3$, L^3 is C_1 - C_{10} alkyl, C_6 - C_{10} aryl- C_1 - C_6 alkyl, or C_6 - C_{10} aryl.
41. The compound of claim 35, wherein L^3 , L^4 , or L^5 is a protecting group selected from methoxymethyl, trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl, triisopropylsilyl, methyl, t-butyl, 3,4-dimethoxybenzyl, p-methoxybenzyl, benzyl, and trityl, or L^3 and L^5 together are acetone, benzylidene, cycloheptylidene, cyclohexylidene, or cyclopentylidene.
42. The compound of claim 35, wherein L^3 is present and is methyl.
43. A compound, wherein said compound is:




Compound AF.

44. A method for producing compound B-1939



which method comprises:

(i) reducing a compound of formula 8-20:




wherein:

Y^2 is a halide; and


L^5 is C_1 - C_{10} alkyl, C_4 - C_{10} aryl- C_1 - C_6 alkyl, or C_4 - C_{10} aryl

and converting the product to a compound of formula (III):




wherein: L^6 is hydrogen or silyl C_1 - C_{10} ; or a salt thereof;

(ii) converting the compound of formula (III) or salt thereof to produce compound AV




and converting compound AV

into compound AI




(iii) converting compound AI into compound B-1939.

45. The method of claim 44, further comprising reacting a compound of formula (Ib):




wherein:

L^{1a} and L^{1b} are independently selected from hydrogen and a protecting group selected from C_1 - C_6 alkyl, aryl (C_1 - C_6) alkyl, silyl (C_1 - C_{10}), and C_1 - C_6 alkyl ester, or L^{1a} and L^{1b} together are a protecting group selected from cyclic C_1 - C_6 acetal, cyclic C_3 - C_7 ketal, and cyclic carbonate under conditions suitable to produce a compound of formula 5-13:




wherein L^{1d} is hydrogen or a protecting group selected from C_1 - C_6 alkyl, aryl (C_1 - C_6) alkyl, C_1 - C_6 ester, and a silyl (C_1 - C_{10}); subjecting the compound of formula 5-13 to oxidative cleavage to form a compound of formula 6-16:




wherein L^{1c} is hydrogen or a protecting group selected from C_1 - C_6 alkyl, or a C_1 - C_6 ester; and subjecting the compound of formula 6-16 to Horner-Wadsworth-Emmons reaction, hydrogenation, and halide functionalization to

produce the compound of formula 8-20.

46. The method of claim 45, wherein the compound of formula (Ib) is




Compound AD or



Compound AJ


47. The method of claim 46, wherein the compound of formula (Ib) is provided at a ratio of at least 8:1 relative to the compound with the opposite stereochemistry at the chiral center marked with the asterisk.

48. The method of claim 46, wherein the compound of formula 5-13 is




Compound AL.

49. The method of claim 46, wherein the compound of formula 6-15 is




Compound AM.

50. The method of claim 46, wherein the compound of formula (Ib)




Compound AK.

51. The method of claim 45, wherein the compound of formula 8-20 is




Compound AF.

52. The method of claim 45, wherein the compound of formula (III) is




Compound AR or



Compound AH.


53. The method of claim 45, wherein the compound of formula 8-20 is provided at a ratio of at least 8:1 relative to the compound with the opposite stereochemistry at the chiral center marked with the asterisk.

54. A compound having the formula:




Compound AU.

55. A compound having the formula:



Compound AR.

56. A composition comprising a compound having the formula:



wherein:

z is a single or double bond, provided that when z is a double bond, X² is C and Y¹ is hydrogen; and provided that when z is a single bond, X² is CH or O;


X¹ is O, S, or CN, provided that when X¹ is CN or S, X² is O;

Y¹ is a halide, hydrogen or O-L², or absent when X² is O;


L¹ and L² are independently selected from hydrogen and a protecting group, or L¹ and L² together are a protecting group, provided that when X¹ is CN, L¹ is absent; or a salt thereof,

wherein the protecting group of L¹ and L² is silyl, C₁-C₁₂ alkylcarbonyl, C₁-C₆ alkyl, aryl (C₁-C₆) alkyl, carbonate, C₁-C₆ alkoxy-(C₁-C₆) alkyl, 3,4-dimethoxybenzyl, p-methoxybenzyl, benzoyl, or C₁-C₆ ester, or L¹ and L² together are a protecting group selected from benzylidene, cyclic C₁-C₆ acetal, cyclic C₃-C₆ ketal, cyclic carbonate, and a C₄-C₇ alkyl ring with one member of the ring covalently linked to both the O of Y¹ and to X¹; or when X¹ and X² are O, L¹ and L² together are cyclic C₃-C₇ ketal.

57. The composition of claim 56, wherein the compound 4-1 has the formula:



58. The composition of claim 56, wherein a diastereomer of the compound 4-1 has the formula:



(Ib)

wherein L^{1a} and L^{1b} are independently selected from hydrogen and a protecting group, or L^{1a} and L^{1b} together are a divalent protecting group.

59. The composition of claim 58, wherein L^{1a} and L^{1b} are a protecting group, which is optionally selected from C₁-C₆ alkyl, aryl (C₁-C₆) alkyl, silyl (C₁-C₁₀), and C₁-C₆ ester, or L^{1a} and L^{1b} together are a divalent protecting group.

60. The composition of claim 59, wherein the divalent protecting group is cyclic C₁-C₆ acetal, cyclic C₃-C₇ ketal, or cyclic carbonate.


61. The composition of claim 56, wherein the protecting group of L^1 and L^2 is independently selected from silyl, C_1 - C_{12} alkylcarbonyl, C_1 - C_6 alkyl, aryl (C_1 - C_6) alkyl, carbonate, C_1 - C_6 alkoxy- $(C_1$ - $C_6)$ alkyl, 3,4-dimethoxybenzyl, p-methoxybenzyl, and C_1 - C_6 ester, or L^1 and L^2 together are a protecting group selected from benzylidene, cycloheptylidene, cyclic C_1 - C_6 acetal, cyclic C_3 - C_6 ketal, cyclic carbonate, and a C_4 - C_7 alkyl ring with one member of the ring covalently linked to both the O of Y^1 and to X^1 ; or when X^1 and X^2 are O, L^1 and L^2 together are cyclic C_3 - C_7 ketal.

62. The composition of claim 56, wherein at least one of L^1 and L^2 is a protecting group and is silyl, C_1 - C_6 alkyl, or C_1 - C_8 alkyl substituted carbonyl.


63. The composition of claim 56, wherein at least one of L^1 and L^2 is a protecting group selected from methoxymethyl, trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl, triisopropylsilyl, methyl, t-butyl, 3,4-dimethoxybenzyl, p-methoxybenzyl, benzyl, and trityl, or L^1 and L^2 together are cycloheptylidene.

64. The composition of claim 56, wherein at least one of L^1 and L^2 is C_1 alkyl carbonyl, or L^1 and L^2 together are acetonide, benzylidene, cyclohexylidene, or cyclopentylidene.


65. The composition of claim 56, wherein a diastereomer of the compound 4-1 is:




Compound AD; or a diastereomer of the compound 4-1 is selected from:




Compound AJ



Compound AK




Compound AL




Compound AM.

66. The composition of claim 56, wherein the ratio of the diastereomer having the formula:



to the diastereomer having the formula:



is at least 8:1, 20:1, or 30:1.