

1,3-Oxazoline and 1,3-thiazoline derivatives, their preparation, and their use as pesticides

5 Description

The invention relates to diaryl-1,3-oxazolines and -1,3-thiazolines, to processes for their preparation, to compositions comprising them, and to their use for controlling animal pests, in particular insects, spider mites,
10 ectoparasites and helminths.

Because of their biological activity, certain 1,3-oxazolines and 1,3-thiazolines are suitable for controlling animal pests (see, for example, EP-A-0 345 775 and EP-A-0 432 661; WO-A-97/06153 ; WO-A-93/24470
15 and WO-A-95/04726).


However, the level of action and/or duration of action of these prior-art compounds is not entirely satisfactory in all fields of application, in particular against certain organisms or when low concentrations are
20 applied.

Since modern pesticides must meet a wide range of demands, for example regarding level, duration and spectrum of action, use spectrum, toxicity, combination with other active substances, combination with formulation
25 auxiliaries or synthesis, and since the occurrence of resistances is possible, the development of such substances can never be regarded as concluded, and there is constantly a high demand for novel compounds which are advantageous over the known compounds, at least as far as some aspects are concerned.

30

It was an object of the present invention to provide compounds which widen the spectrum of the pesticides in various aspects.

This object, and other objects which are not mentioned explicitly and which
35 can be deduced or concluded from the ideas discussed herein, is solved by 1,3-oxazoline and 1,3-thiazoline derivatives of the formula (I)



in which the symbols have the following meanings:

5

A is phenyl, pyridyl, pyrimidinyl, pyrazinyl, pyrazolyl or thienyl, each of the abovementioned groups optionally being substituted by one or more, preferably one, two or three, radicals X;


10 X is identical or different

a) halogen, cyano, nitro;

b) (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₂-C₄)alkynyl, (C₁-C₄)alkoxy, (C₁-C₄)alkylthio, (C₁-C₄)alkylsulfinyl, (C₃-C₈)cycloalkyl, (C₄-C₈)cycloalkenyl, phenyl,

15 the radicals of group b optionally being substituted by one or more, preferably one, two or three, radicals selected from the group consisting of halogen, cyano, (C₁-C₄)alkyl, (C₁-C₄)haloalkyl, (C₁-C₄)alkoxy and (C₁-C₄)haloalkoxy;20 E is a single bond, (C₁-C₄)alkylene, -O-CH₂- or -CH₂-O-;

G is a radical selected from the group consisting of:



25

Z is oxygen or sulfur;

- R^1 , R^2 and R^3 are identical or different hydrogen, halogen, (C₁-C₄)haloalkyl, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)haloalkoxy or cyano;
- 5 R^4 is hydrogen or a group (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkenyl, (C₄-C₁₀)cycloalkylalkyl or (C₇-C₁₂)phenylalkyl, each of the abovementioned groups optionally being substituted by one or more substituents selected from the group consisting of halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl, (C₁-C₄)alkoxy, (C₁-C₃)haloalkoxy, (C₁-C₃)alkylthio, (C₃-C₈)cycloalkyl, (C₇-C₁₂)phenylalkoxy, (C₂-C₄)alkylcarbonyl, alkoxycarbonyl, (C₂-C₆)monoalkyl- and (C₃-C₉)dialkylamino-carbonyl, cyano and tri(C₁-C₄)alkylsilyl;
- 10 R^5 and R^6 are identical or different
- a) hydrogen, halogen, cyano, formyl, (C₂-C₅)alkylcarbonyl, (C₂-C₈)alkoxycarbonyl, (C₃-C₉)monoalkyl- and dialkylamino-carbonyl,
- 20 b) (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, (C₄-C₈)cycloalkenyl, (C₁-C₈)alkoxyalkyl, (C₁-C₈)alkylthio, (C₁-C₈)alkylsulfinyl, (C₁-C₈)alkylsulfonyl, NR^7R^8 , phenyl, phenoxy, (C₇-C₁₂)phenylalkyl, (C₇-C₁₂)phenylalkoxy, heterocyclyl, preferably (C₄-C₇)oxacycloalkyl, (C₄-C₇)oxacycloalkenyl, (C₄-C₇)thiacycloalkyl, (C₄-C₇)thiacycloalkenyl, pyridyl, pyrimidinyl, pyrazinyl, furanyl, thienyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, thiazolyl, oxadiazolyl, heterocyclyloxy derivatives, heterocyclyl(C₁-C₄)alkoxy, (C₃-C₁₂)heterocyclylalkyl, each of the groups b optionally being substituted by one or more, preferably one to three, radicals selected from the group consisting of halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl, (C₁-C₃)alkoxy, (C₁-C₃)haloalkoxy, (C₁-C₃)alkylthio, NR^7R^8 , alkylcarbonyl (C₂-C₅), alkoxycarbonyl (C₂-C₈), monoalkyl- and dialkylaminocarbonyl (C₃-C₉), cyano and tri(C₁-C₄)alkylsilyl;
- 30 c) (C₁-C₈)alkoxy, each of the groups c optionally being substituted by one or more, preferably one to three, radicals selected from the group consisting of halogen, (C₁-C₃)alkylthio, NR^7R^8 ,
- 35

alkylcarbonyl (C₂-C₅), alkoxycarbonyl (C₂-C₈), monoalkyl- and dialkylaminocarbonyl (C₃-C₉), cyano and tri(C₁-C₄)alkylsilyl; or

- 5 d) R⁵ and R⁶ together form a preferably monocyclic ring which originates by linking two of the groups mentioned under b), preferably (C₃-C₅)alkylene, (C₂-C₄)alkyleneoxy, (C₁-C₃)alkylene-dioxy, (C₂-C₄)alkyleneamino or alkyleneethio(C₂-C₄);

R⁷ and R⁸ are identical or different

- 10 a) hydrogen;
 b) (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, (C₇-C₁₂)phenylalkyl, (C₁-C₆)alkyl-carbonyl, (C₁-C₆)alkylsulfonyl which are optionally substituted by one or more, preferably one to three, radicals selected from the group consisting of halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl,
 15 (C₁-C₄)alkoxy, (C₁-C₃)haloalkoxy, (C₁-C₃)alkylthio, (C₁-C₃)alkyl-amino, (C₁-C₃)dialkylamino, (C₃-C₈)cycloalkyl, cyano and tri(C₁-C₄)alkylsilyl; or
 c) R⁷ and R⁸ together are alkylene (C₃-C₇), (C₃-C₇)oxa-, aza- or thia-alkylene, (C₃-C₇)alkylenecarbonyl or (C₃-C₇)alkylenesulfonyl;

20

their pure isomers (optical and geometric isomers) and isomer mixtures of these, their N-oxides and their salts which are suitable for use as pesticides.

- 25 Surprisingly, compounds of the formula (I) have a better acaricidal and insecticidal activity regarding the spectrum of action and the potency than known 1,3-oxazoline and 1,3-thiazoline derivatives.

The symbols in the formula (I) preferably have the following meanings:

30

A is preferably phenyl or pyridyl, especially preferably phenyl.

X is preferably

- a) halogen, cyano, nitro or
 35 b) (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₁-C₄)alkoxy, (C₁-C₄)alkylthio, (C₁-C₄)alkylsulfinyl, (C₃-C₇)cycloalkyl, (C₄-C₈)cycloalkenyl,


the radicals of group b optionally being substituted by one or more, preferably one, two or three, radicals selected from the group consisting of halogen, cyano, (C₁-C₄)alkoxy and (C₁-C₄)haloalkoxy;

5 X is especially preferably halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl, (C₁-C₄)alkoxy or (C₁-C₃)haloalkoxy.

E is preferably a single bond or -CH₂-, especially preferably a single bond.

10

G is preferably



15 especially preferably 3-pyridyl.

Z is preferably oxygen.

20 R¹, R², R³ are preferably H, halogen, (C₁-C₄)haloalkyl, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)haloalkoxy or cyano, especially preferably H, halogen, (C₁-C₄)haloalkyl, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)haloalkoxy,

25 R⁴ is preferably H or (C₁-C₈)alkyl which is optionally substituted by one or more, preferably 1 to 3, radicals selected from the group consisting of halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl, (C₁-C₄)alkoxy, (C₁-C₃)haloalkoxy, (C₁-C₃)alkylthio, (C₃-C₈)cycloalkyl, cyano or tri(C₁-C₄)alkylsilyl.

R⁵, R⁶ are preferably

- 30 a) H, halogen, cyano,
 b) (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, (C₄-C₈)cycloalkenyl, (C₁-C₈)alkylthio, (C₁-C₈)alkylsulfinyl, (C₁-C₈)alkylsulfonyl, (C₇-C₁₂)phenylalkyl, (C₇-C₁₂)phenylalkoxy, (C₄-C₇)oxycycloalkyl or (C₄-C₇)oxacycloalkenyl,

the groups b optionally being substituted by one or more, preferably 1 to 3, radicals selected from the group consisting of halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl, (C₁-C₃)alkoxy, (C₁-C₃)haloalkoxy, (C₁-C₃)alkylthio and cyano,

- 5 c) (C₁-C₈)alkoxy which can optionally be substituted by one or more, preferably 1 to 3, radicals selected from the group consisting of halogen, (C₁-C₃)alkylthio and cyano.

R⁵, R⁶ are especially preferably

- 10 a) H, halogen, cyano,
 b) (C₁-C₈)alkyl, (C₁-C₈)alkoxy, (C₃-C₈)cycloalkyl, (C₄-C₈)cycloalkenyl, (C₁-C₈)alkylthio, (C₁-C₈)alkylsulfinyl, (C₁-C₈)alkylsulfonyl, (C₇-C₁₂)phenylalkyl or (C₇-C₁₂)phenylalkoxy,

- 15 the groups b being substituted by one or more, preferably 1 to 3, radicals selected from the group consisting of halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl and cyano.

20 In the above formula, "halogen" is to be understood as meaning a fluorine, chlorine, bromine or iodine atom;

the term "(C₁-C₄)alkyl" an unbranched or branched hydrocarbon radical having 1 to 4 carbon atoms such as, for example, the methyl, ethyl, propyl, isopropyl, 1-butyl, 2-butyl, 2-methylpropyl or tert-butyl radical;

- 25 the term "(C₁-C₈)alkyl" the abovementioned alkyl radicals and, for example, the pentyl, 2-methylbutyl, 1,1-dimethylpropyl, hexyl, heptyl, octyl or the 1,1,3,3-tetramethylbutyl radical;

- the term "(C₁-C₄)haloalkyl" an alkyl group mentioned under the term "(C₁-C₄)alkyl" in which one or more hydrogen atoms are replaced by the
 30 abovementioned halogen atoms, preferably chlorine or fluorine, such as, for example, the trifluoromethyl group, the 1-fluoroethyl group, the 2,2,2-trifluoroethyl group, the chloromethyl, fluoromethyl group, the difluoromethyl group or the 1,1,2,2-tetrafluoroethyl group;

- the term "(C₃-C₈)cycloalkyl", for example, the cyclopropyl, cyclobutyl or cyclopentyl group; and the cyclohexyl, cycloheptyl or cyclooctyl radical;

- 35 the term "(C₃-C₈)halocycloalkyl" one of the abovementioned (C₃-C₅)cycloalkyl radicals in which one or more, in the case of fluorine optionally also all, hydrogen atoms are replaced by halogen, preferably

- fluorine or chlorine, such as, for example, the 2,2-difluoro or 2,2-dichlorocyclopropane group or the fluorocyclopentane radical;
the term "(C₂-C₄)alkenyl", for example, the vinyl, allyl, 2-methyl-2-propenyl or 2-butenyl group;
- 5 the term "(C₂-C₄)haloalkenyl" a (C₂-C₄)alkenyl group in which some or, in the case of fluorine also all, of the hydrogen atoms are replaced by halogen, preferably fluorine or chlorine;
the term "(C₂-C₄)alkynyl", for example, the ethynyl, propargyl, 2-methyl-2-propynyl or 2-butyryl group;
- 10 the term "(C₂-C₈)alkynyl" for example, the abovementioned radicals and, for example, the 1-pentynyl, 2-pentynyl, 3-pentynyl, 4-pentynyl or the 1-octynyl group;
the term "(C₂-C₄)haloalkynyl" a (C₂-C₄)alkynyl group in which some, in the case of fluorine also all, of the hydrogen atoms are replaced by halogen
- 15 atoms, preferably fluorine or chlorine;
the term "(C₁-C₄)hydroxyalkyl", for example, the hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 1-hydroxy-1-methylethyl or the 1-hydroxypropyl group;
the term "(C₁-C₄)alkanoyl-(C₁-C₄)alkyl", for example, an acetylmethyl, propionylmethyl, 2-acetylethyl or a butyrylmethyl group;
- 20 the term "(C₁-C₄)alkanoyl", for example, the formyl, acetyl, propionyl, 2-methylpropionyl or butyryl group;
the term "(C₁-C₈)alkanoyl" the abovementioned radicals and, for example, the valeroyl, pivaloyl, hexanoyl, heptanoyl or octanoyl group;
- 25 the term "(C₁-C₁₂)alkanoyl", for example, the abovementioned radicals and, for example, the nonanoyl, decanoyl or the dodecanoyl group;
the term "(C₂-C₄)haloalkanoyl" a (C₁-C₄)alkanoyl group in which some, in the case of fluorine also all, of the hydrogen atoms are replaced by halogen atoms, preferably fluorine or chlorine;
- 30 the term "(C₂-C₁₂)haloalkanoyl" a (C₁-C₂₀)alkanoyl group in which some, in the case of fluorine also all, of the hydrogen atoms are replaced by halogen atoms, preferably fluorine or chlorine;
the term "cyano(C₁-C₄)alkyl" a cyanoalkyl group whose hydrocarbon radical has the meanings given for the term "(C₁-C₄)alkyl";
- 35 the terms "(C₁-C₄)nitroalkyl" or "(C₁-C₄)thiocyanoalkyl" one of the abovementioned (C₁-C₄)alkyl groups which are substituted by a nitro or a thiocyno group;

- the term "(C₁-C₄)alkoxycarbonyl", for example, the methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl or tert-butoxycarbonyl group;
- the term "(C₁-C₈)alkoxycarbonyl", for example, the abovementioned
- 5 radicals and, for example, the pentyloxycarbonyl, hexyloxycarbonyl or the octyloxycarbonyl group;
- the term "(C₁-C₁₂)alkoxycarbonyl" the abovementioned radicals and, for example, the nonyloxycarbonyl, 2-methyloctyloxycarbonyl, decyloxycarbonyl or dodecyloxycarbonyl group;
- 10 the term "(C₁-C₄)alkoxycarbonyl-(C₁-C₄)alkyl", for example, a methoxycarbonylmethyl, ethoxycarbonylmethyl or methoxycarbonylethyl group;
- the term "(C₁-C₄)haloalkoxycarbonyl" a (C₁-C₄)alkoxycarbonyl group in which one or more, in the case of fluorine optionally also all, hydrogen
- 15 atoms are replaced by halogen, preferably fluorine or chlorine;
- the term "(C₁-C₄)alkylthio" an alkylthio group whose hydrocarbon radical has the meaning given for the term "(C₁-C₄)alkyl";
- the term "(C₁-C₈)alkylthio" an alkylthio group whose alkyl radical has the meaning given for the term "(C₁-C₈)alkyl";
- 20 the term "(C₁-C₄)haloalkylthio" a (C₁-C₄)alkylthio group in which one or more, in the case of fluorine optionally also all, hydrogen atoms of the hydrocarbon moiety are replaced by halogen, in particular chlorine or fluorine;
- the term "(C₁-C₄)alkylsulfinyl", for example, the methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl- or tert-butylsulfinyl group;
- 25 the term "(C₁-C₈)alkylsulfinyl" one of the abovementioned alkylsulfinyl groups and, for example, the pentylsulfinyl, 2-methylbutylsulfinyl, hexylsulfinyl or octylsulfinyl group;
- the term "(C₁-C₄)alkylsulfonyl", for example, the methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl or tert-butylsulfonyl group;
- 30 the terms "(C₁-C₄)haloalkylsulfinyl" and "(C₁-C₄)haloalkylsulfonyl" (C₁-C₄)alkylsulfinyl and -sulfonyl radicals having the abovementioned meanings where one or more, in the case of fluorine optionally also all, hydrogen atoms of the hydrocarbon moiety are replaced by halogen, in
- 35 particular chlorine or fluorine;
- the terms "fluoromethylsulfinyl" and "fluoromethylsulfonyl" the mono-, di- and trifluoromethylsulfinyl and -sulfonyl group;

- the term "(C₁-C₄)alkoxy" an alkoxy group whose hydrocarbon radical has the meaning given for the term "(C₁-C₄)alkyl";
- the term "(C₁-C₈)alkoxy" an alkoxy group whose hydrocarbon radical has the meaning given for the term "(C₁-C₈)alkyl";
- 5 the term "(C₁-C₈)alkylsulfonyl" one of the abovementioned alkylsulfonyl groups and, for example, the pentyl-, 2-methylbutyl-, hexyl-, heptyl- or octylsulfonyl group,
- the term "(C₁-C₄)alkylamino", for example, the methylamino, ethylamino, propylamino, isopropylamino, butylamino, isobutylamino, sec-butylamino or
- 10 the tert-butylamino group,
- the term "(C₁-C₈)alkylamino" one of the abovementioned (C₁-C₄)alkylamino groups and, for example, the pentylamino, hexylamino, heptylamino or octylamino group;
- the term "(C₁-C₄)dialkylamino", for example, the dimethylamino,
- 15 methylethylamino, diethylamino, dipropylamino or the dibutylamino group, but also cyclic systems such as, for example, the pyrrolidino or piperidino group,
- the term "(C₁-C₈)dialkylamino" one of the abovementioned (C₁-C₄)dialkylamino groups and, for example, the dipentyl, dihexyl or the
- 20 dioctylamino group;
- the term "(C₁-C₄)haloalkoxy" a haloalkoxy group whose halohydrocarbon radical has the meaning given for the term "(C₁-C₄)haloalkyl";
- the term "(C₁-C₄)alkoxy-(C₁-C₄)alkyl", for example, a 1-methoxyethyl group, a 2-methoxyethyl group, a 2-ethoxyethyl group, a methoxymethyl or
- 25 ethoxymethyl group, a 3-methoxypropyl group or a 4-butoxybutyl group;
- the terms "(C₁-C₄)haloalkoxy-(C₁-C₄)alkyl", "(C₁-C₄)alkoxy-(C₁-C₄)haloalkyl" and "(C₁-C₄)haloalkoxy-(C₁-C₄)haloalkyl" (C₁-C₄)alkoxy-(C₁-C₄)alkyl radicals having the abovementioned meanings where one or
- 30 more, in the case of fluorine optionally also all, hydrogen atoms of the relevant hydrocarbon moieties are replaced by halogen, preferably chlorine or fluorine;
- the term "(C₁-C₄)alkylthio-(C₁-C₄)alkyl", for example, methylthiomethyl, ethylthiomethyl, propylthiomethyl, 2-methylthioethyl, 2-ethylthioethyl or 3-methylthiopropyl;
- 35 the term "heterocyclyl" a heteroaromatic or heteroaliphatic ring system where "heteroaromatic ring system" is to be understood as meaning an aryl radical in which at least one CH group is replaced by N and/or at least two adjacent CH groups are replaced by S, NH or O, for example a radical of

thiophene, furan, pyrrole, thiazole, oxazole, imidazole, isothiazole, isoxazole, pyrazole, 1,3,4-oxadiazole, 1,3,4-thiadiazole, 1,3,4-triazole, 1,2,4-oxadiazole, 1,2,4-thiadiazole, 1,2,4-triazole, 1,2,3-triazole, 1,2,3,4-tetrazole, benzo[b]thiophene, benzo[b]furan, indole, 5 benzo[c]thiophene, benzo[c]furan, isoindole, benzoxazole, benzothiazole, benzimidazole, benzisoxazole, benzisothiazole, benzopyrazole, benzothiadiazole, benzotriazole, dibenzofuran, dibenzothiophene, carbazole, pyridine, pyrazine, pyrimidine, pyridazine, 1,3,5-triazine, 1,2,4-triazine, 1,2,4,5-triazine, quinoline, isoquinoline, quinoxaline, 10 quinazoline, cinnoline, 1,8-naphthyridine, 1,5-naphthyridine, 1,6-naphthyridine, 1,7-naphthyridine, phthalazine, pyridopyrimidine, purine, pteridine or 4H-quinolizine;

the term "heteroaliphatic ring system" a (C₃-C₈)cycloalkyl radical in which at least one carbon unit is replaced by O, S or a group NR¹¹ and R¹¹ is 15 hydrogen, (C₁-C₄)alkyl, (C₁-C₄)alkoxy or aryl;

the term "heterocyclyloxy" or "heterocyclylthio" one of the abovementioned heterocyclic radicals which are linked via an oxygen or sulfur atom;

the term (C₃-C₈)cycloalkoxy" or "(C₃-C₈)cycloalkylthio" one of the abovementioned (C₃-C₈)cycloalkyl radicals which are linked via an oxygen 20 or sulfur atom;

the term "aryl-(C₁-C₄)alkanoyl", for example, the phenylacetyl, 3-phenylpropionyl, 2-phenylpropionyl, 2-methyl-2-phenylpropionyl, 4-phenylbutyryl or the naphthylacetyl group;

the term "(C₃-C₈)cycloalkyl-(C₁-C₄)alkanoyl", for example, the 25 cyclopropylcarbonyl, cyclobutylcarbonyl, cyclopentylcarbonyl, cyclohexylcarbonyl, cyclohexylacetyl or the cyclohexylbutyryl group;

the term "heterocyclyl-(C₁-C₄)alkanoyl", for example, the thenoyl, furoyl, nicotinoyl, thienylacetyl or the pyridinepropionyl group;

the term "(C₃-C₈)cycloalkoxycarbonyl", for example, the 30 cyclobutyloxycarbonyl, cyclopentyloxycarbonyl, cyclohexyloxycarbonyl or the cycloheptyloxycarbonyl group;


the term "(C₃-C₈)cycloalkyl-(C₁-C₄)alkoxycarbonyl", for example, the cyclopropylmethoxycarbonyl, cyclobutylmethoxycarbonyl, cyclopentyloxymethoxycarbonyl, cyclohexyloxymethoxycarbonyl, 35 1-(cyclohexyl)ethoxycarbonyl or the 2-(cyclohexyl)ethoxycarbonyl group;

the explanation given above applies analogously to homologs or radicals derived therefrom.


The present invention relates to the compounds of the formula (I) in the form of the free base or of an acid addition salt. Acids which can be used for salt formation are inorganic acids such as hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid or organic acids such as formic acid, acetic acid, propionic acid, malonic acid, oxalic acid, fumaric acid, adipic acid, stearic acid, oleic acid, methane sulfonic acid, benzene sulfonic acid or toluene sulfonic acid.

Some of the compounds of the formula (I) have one or more asymmetric carbon atoms or stereoisomers on double bonds. Enantiomers or diastereomers may therefore occur. The invention encompasses not only the pure isomers, but also their mixtures. The diastereomer mixtures can be resolved into the components by customary methods, for example by selective crystallization from suitable solvents or by chromatography. Racemates can be resolved into the enantiomers by customary methods, for example by salt formation with a chiral, enantiomerically pure acid, separation of the diastereomeric salts and liberation of the pure enantiomers by means of a base.



Very especially preferred are the following groups of compounds of the formula (Ia) to (Ie):




25



12




5





10 In these formulae X^1 , X^2 , Z , R^4 , R^5 and R^6 each have the meanings given in the formula (I) ($X^1, X^2 \triangleq X$).

Particularly preferred are compounds of the formula (Ia1) to (Ia4):




15





5



10 Z, R^5 and R^6 each have in this case the meanings given in the formula (I).

The compounds according to the invention are prepared by methods which are known per se from the literature as they are described in standard works on organic synthesis, for example Houben-Weyl, Methoden der
 15 Organischen Chemie [Methods in Organic Chemistry], Georg-Thieme-Verlag, Stuttgart.


The preparation is carried out under reaction conditions which are known and suitable for the abovementioned reactions. Other variants which are
 20 known per se, but not illustrated here in greater detail, may also be made use of.

If desired, the starting materials may also be formed in situ, in such a way that they are not isolated from the reaction mixture but immediately reacted
 25 further to give the compounds of the formula (I).

The general chemistry of the 1,3-oxazolines is described, for example, in Tetrahedron, 1994, 50, 2297-2360 and in Nachr. Chem. Tech. Lab. 1996, 44, 744-750.

30

The invention also relates to a process for the preparation of compounds of the formula (I) by reacting known 1,3-oxazolines and 1,3-thiazolines of the formula (II) (see, for example, EP-A-0 345 775) (suitably substituted by R^1 , R^2 , R^3) with metalloïd-pyridine compounds of the formula (III) (suitably substituted by R^4 , R^5 , R^6), which comprises reacting a halogen and perfluoroalkylsulfonate compound of the formula (II)




in which
Y is Cl, Br, I or perfluoroalkylsulfonate and
A, Z and E have the meanings given in formula (I)

with an organometallic compound of the formula (III)

G-M (III)
in which
M is a B-, Sn- or Zn-containing leaving group and
G has the meaning given in formula (I),
with palladium catalysis.

The methods of A to H for the synthesis of a variety of subgroups of compounds of the formula (I) may be mentioned by way of example:

Method A




G = 3-pyridyl, 4-pyridyl, 2-pyridon-5-yl, 2-pyridon-3-yl (cf. formula (I)).

Linkage of the phenyl ring with a pyridine ring is effected, for example, by palladium catalysis using Suzuki coupling, Stille coupling or Negishi coupling (see, for example, P. Knochel, Chem. Review 1993, 93, 2117-

2188 or Jiro Tsuji, *Palladium Reagents and Catalysts*, John Wiley & Sons, 1996).


It is also possible for the functions required for the linkage to be swapped between (II) and (III):

Method B



It is also possible first to construct a pyridyl-phenyl structural unit and to convert these precursors into 1,3-oxazolines and 1,3-thiazolines of the formula (I).


15 Method C



The oxazoline precursors of the formula (II) are prepared, for example, by the following routes, some of which are known (EP-A-0 432 661; G. Helmchen, *Tetrahedron* 1996, 52, 7547-7583)):

Activated carboxylic acid derivatives are reacted with aminoalcohols (VI) (suitably substituted by R^1 , R^2 , R^3) (Synthesis K. Drauz, *J. Org. Chem.* 1993, 58, 3568-3571) to give the amidoalcohols (VII):

Method D (Ex. VP3)




- 5 The chlorides (VIII) (Y=Cl) are obtained from the amidoalcohols (VII) for example using thionyl chloride; when sulfone halides are used, the sulfonates (VIII) (Y= for example mesylate, tosylate) are formed.

The oxazolines (II) are formed from (VIII) by the action of bases.


10

Method E (Ex. VP4)




- 15 Bases which are suitable are, for example, basic salts such as alkali metal hydroxides, alkali metal carbonates, hydrides, alkoxides and amines. Starting from the 4-hydroxyphenyl derivatives of (II), the triflates may be obtained by sulfonation with, for example, trifluoromethanesulfonyl chloride.
- 20 The oxazoline precursors of the formula (IV) which contain a tin group are accessible by Stille coupling with distannane:

Method F (Ex. VP5)



- 5 The pyridine precursors of the formula (III) are formed, for example, by generating metalated pyridine derivatives with the suitable substituents R^4 , R^5 , R^6 as per formula (I) and their reaction with, for example, boric esters, tin halides or zinc halides.

10 Method G (Ex. VP7, VP8)




$R = \text{alkyl, alkylene}$

15

The aminoalcohol precursors (VI) can be prepared by reducing known 2-hydroxyacetophenone oximes by means of hydride.

Method H (Ex. VP2)

20



- 25 Collections of compounds of the formula (I) which can be synthesized by the abovementioned scheme may also be prepared in a parallel manner, and this may be effected manually or in a semiautomated or fully automated manner. In this case, it is possible, for example, to automate the procedure of the reaction, work-up or purification of the products or of the intermediates. In total, this is to be understood as meaning a procedure as

it is described, for example, by S.H. DeWitt in "Annual Reports in Combinatorial Chemistry and Molecular Diversity: Automated synthesis", Volume 1, Verlag Escom 1997, pages 69 to 77.

- 5 A series of commercially available apparatuses as they are offered by, for example, Stem Corporation, Woodrolfe Road, Tollesbury, Essex, CM9 8SE, England or H+P Labortechnik GmbH, Bruckmannring 28, 85764 Oberschleißheim, Germany, may be used for the parallel procedure of the reaction and work-up. For the parallel purification of compounds of the
10 formula (I), or of intermediates obtained during the preparation, use may be made, inter alia, of chromatography apparatuses, for example those by ISCO, Inc., 4700 Superior Street, Lincoln, NE 68504, USA.

- The apparatuses mentioned lead to a modular procedure in which the
15 individual process steps are automated, but manual operations must be performed between the process steps. This can be prevented by employing semi-integrated or fully integrated automation systems where the automation modules in question are operated by, for example, robots. Such automation systems can be obtained, for example, from Zymark
20 Corporation, Zymark Center, Hopkinton, MA 01748, USA.

- In addition to what has been described here, compounds of the formula (I) may be prepared in part or fully by solid-phase-supported methods. For this purpose, individual intermediate steps or all intermediate steps of the
25 synthesis or of a synthesis adapted to suit the procedure in question are bound to a synthetic resin. Solid-phase-supported synthesis methods are described extensively in the specialist literature, for example Barry A. Bunin in "The Combinatorial Index", Academic Press, 1998.

- The use of solid-phase-supported synthesis methods permits a series of
30 protocols which are known from the literature and which, in turn, can be performed manually or in an automated manner. For example, the "tea-bag method" (Houghten, US 4,631,211; Houghten et al., Proc. Natl. Acad. Sci, 1985, 82, 5131-5135), in which products by IRORI, 11149 North Torrey Pines Road, La Jolla, CA 92037, USA, are employed, may be
35 semiautomated. The automation of solid-phase-supported parallel syntheses is performed successfully, for example, by apparatuses by Argonaut Technologies, Inc., 887 Industrial Road, San Carlos, CA 94070, USA or MultiSynTech GmbH, Wullener Feld 4, 58454 Witten, Germany.

The preparation of the processes described herein yields compounds of the formula (I) in the form of substance collections which are termed libraries. The present invention also relates to libraries which comprise at least two compounds of the formula (I).

5

- The compounds of the formula (I) are suitable for controlling animal pests, in particular insects, arachnids, helminths and mollusks, very especially preferably for controlling insects and arachnids which are encountered in agriculture, in livestock breeding, in forests, in the protection of stored
- 10 goods and materials, and in the hygiene sector, and have good plant tolerance and favorable toxicity to warm-blooded species. They are active against normally sensitive and resistant species and against all or individual developmental stages. The abovementioned pests include:
- from the order of the Acarina, for example, *Acarus siro*, *Argas* spp.,
- 15 *Ornithodoros* spp., *Dermanyssus gallinae*, *Eriophyes ribis*, *Phyllocoptruta oleivora*, *Boophilus* spp., *Rhipicephalus* spp., *Amblyomma* spp., *Hyalomma* spp., *Ixodes* spp., *Psoroptes* spp., *Chorioptes* spp., *Sarcoptes* spp., *Tarsonemus* spp., *Bryobia praetiosa*, *Panonychus* spp., *Tetranychus* spp., *Eotetranychus* spp., *Oligonychus* spp., *Eutetranychus* spp.
- 20 From the order of the Isopoda, for example, *Oniscus aselus*, *Armadium vulgare*, *Porcellio scaber*.
- From the order of the Diplopoda, for example, *Blaniulus guttulatus*.
- From the order of the Chilopoda, for example, *Geophilus carpophagus*, *Scutigera* spp.
- 25 From the order of the Symphyla, for example, *Scutigera immaculata*.
- From the order of the Thysanura, for example, *Lepisma saccharina*.
- From the order of the Collembola, for example, *Onychiurus armatus*.
- From the order of the Orthoptera, for example, *Blatta orientalis*, *Periplaneta americana*, *Leucophaea maderae*, *Blattella germanica*, *Acheta domesticus*,
- 30 *Gryllotalpa* spp., *Locusta migratoria migratorioides*, *Melanoplus differentialis*, *Schistocerca gregaria*.
- From the order of the Isoptera, for example, *Reticulitermes* spp.
- From the order of the Anoplura, for example, *Phylloera vastatrix*, *Pemphigus* spp., *Pediculus humanus corporis*, *Haematopinus* spp.,
- 35 *Linognathus* spp.
- From the order of the Mallophaga, for example, *Trichodectes* pp., *Damalinea* spp.
- From the order of the Thysanoptera, for example, *Hercinothrips femoralis*, *Thrips tabaci*.

From the order of the Heteroptera, for example, *Eurygaster* spp., *Dysdercus intermedius*, *Piesma quadrata*, *Cimex lectularius*, *Rhodnius prolixus*, *Triatoma* spp.

- From the order of the Homoptera, for example, *Aleurodes brassicae*,
 5 *Bemisia tabaci*, *Trialeurodes vaporariorum*, *Aphis gossypii*, *Brevicoryne brassicae*, *Cryptomyzus ribis*, *Doralis fabae*, *Doralis pomi*, *Eriosoma lanigerum*, *Hyalopterus arundinis*, *Macrosiphum avenae*, *Myzus* spp., *Phorodon humuli*, *Rhopalosiphum padi*, *Empoasca* spp., *Euscelus bilobatus*, *Nephotettix cincticeps*, *Lecanium corni*, *Saissetia oleae*,
 10 *Laodelphax striatellus*, *Nilaparvata lugens*, *Aonidiella aurantii*, *Aspidiotus hederae*, *Pseudococcus* spp., *Psylla* spp.

- From the order of the Lepidoptera, for example, *Pectinophora gossypiella*, *Bupalus piniarius*, *Cheimatobia brumata*, *Lithocolletis blancardella*, *Hyponomeuta padella*, *Plutella maculipennis*, *Malacosoma neustria*,
 15 *Euproctis chrysorrhoea*, *Lymantria* spp., *Bucculatrix thurberiella*, *Phyllocnistis citrella*, *Agrotis* spp., *Euxoa* spp., *Feltia* spp., *Earias insulana*, *Heliothis* spp., *Laphygma exigua*, *Mamestra brassicae*, *Panolis flammea*, *Prodenia litura*, *Spodoptera* spp., *Trichoplusia ni*, *Carpocapsa pomonella*, *Pieris* spp., *Chilo* spp., *Pyrausta nubilalis*, *Ephestia kuehniella*, *Galleria mellonella*,
 20 *Cacoecia podana*, *Capua reticulana*, *Choristoneura fumiferana*, *Clysia ambiguella*, *Homona magnanima*, *Tortrix viridana*.

- From the order of the Coleoptera, for example, *Anobium punctatum*, *Rhizopertha dominica*, *Bruchidius obtectus*, *Acanthoscelides obtectus*,
 25 *Hylotrupes bajulus*, *Agelastica alni*, *Leptinotarsa decemlineata*, *Phaedon cochleariae*, *Diabrotica* spp., *Psylliodes chrysocephala*, *Epilachna varivestis*, *Atomaria* spp., *Oryzaephilus surinamensis*, *Anthonomus* spp., *Sitophilus* spp., *Otiorrhynchus sulcatus*, *Cosmopolites sordidus*, *Ceuthorrhynchus assimilis*, *Hypera postica*, *Dermestes* spp., *Trogoderma*,
 30 *Anthrenus* spp., *Attagenus* spp., *Lyctus* spp., *Meligethes aeneus*, *Ptinus* spp., *Niptus hololeucus*, *Gibbium psyllioides*, *Tribolium* spp., *Tenebrio molitor*, *Agriotes* spp., *Conoderus* spp., *Melolontha melolontha*, *Amphimallon solstitialis*, *Costelytra zealandica*.

- From the order of the Hymenoptera, for example, *Diprion* spp.,
 35 *Hoplocampa* spp., *Lasius* spp., *Monomorium pharaonis*, *Vespa* spp.

From the order of the Diptera, for example, *Aedes* spp., *Anopheles* spp., *Culex* spp., *Drosophila melanogaster*, *Musca* spp., *Fannia* spp., *Calliphora erythrocephala*, *Lucilia* spp., *Chrysomyia* spp., *Cuterebra* spp., *Gastrophilus* spp., *Hypobosca* spp., *Stomoxys* spp., *Oestrus* spp.,

Hypoderma spp., Tabanus spp., Tannia spp., Bibio hortulanus, Oscinella frit, Phorbia spp., Pegomyia hyoscyami, Ceratitis capitata, Dacus oleae, Tipula paludosa.

- From the order of the Siphonaptera, for example, Xenopsylla cheopsis,
5 Ceratophyllus spp.

From the order of the Arachnida, for example, Scorpio maurus, Latrodectus mactans.

- From the class of the helminths, for example, Haemonchus, Trichostrongylus, Ostertagia, Cooperia, Chabertia, Strongyloides,
10 Oesophagostomum, Hyostrongylus, Ancylostoma, Ascaris and Heterakis and also Fasciola.

- From the class of the Gastropoda, for example, Deroceras spp., Arion spp., Lymnaea spp., Galba spp., Succinea spp., Biomphalaria spp., Bulinus spp.,
15 Oncomelania spp.

From the class of the Bivalva, for example, Dreissena spp.

- The plant-parasitic nematodes which can be controlled in accordance with the invention include, for example, the root-parasitic soil-dwelling
20 nematodes such as, for example, those of the genera Meloidogyne (root knot nematodes, such as Meloidogyne incognita, Meloidogyne hapla and Meloidogyne javanica), Heterodera and Globodera (cyst-forming nematodes, such as Globodera rostochiensis, Globodera pallida, Heterodera trifolii) and of the genera Radopholus, such as Radopholus
25 similis, Pratylenchus such as Pratylenchus neglectus, Pratylenchus penetrans and Pratylenchus curvatus;

- Tylenchulus such as Tylenchulus semipenetrans, Tylenchorhynchus, such as Tylenchorhynchus dubius and Tylenchorhynchus claytoni, Rotylenchus such as Rotylenchus robustus, Heliocotylenchus such as Haliocotylenchus
30 multicinctus, Belonoaimus such as Belonoaimus longicaudatus, Longidorus such as Longidorus elongatus, Trichodorus such Trichodorus primitivus and Xiphinema such as Xiphinema index.

- Other nematode genera which can be controlled using the compounds
35 according to the invention are Ditylenchus (stem parasites, such as Ditylenchus dipsaci and Ditylenchus destructor), Aphelenchoides (foliar nematodes, such as Aphelenchoides ritzemabosi) and Anguina (seed nematodes, such as Anguina tritici).

The invention also relates to compositions, for example crop protection compositions, preferably insecticidal, acaricidal, ixodidical, nematocidal, molluskicidal or fungicidal, especially preferably insecticidal and acaricidal, compositions which comprise one or more compounds of the formula (I) in addition to suitable formulation auxiliaries.

In general, the compositions according to the invention comprise 1 to 95% by weight of the active substances of the formula (I).

To prepare the compositions according to the invention, the active substance and the other additives are combined and brought into suitable use form.

They can be formulated in various ways, depending on the biological and/or chemical-physical parameters which prevail. The following are examples of possible formulations:

Wettable powders (WP), emulsifiable concentrates (EC), aqueous solutions (SL), emulsions, sprayable solutions, oil- or water-based dispersions (SC), suspoemulsions (SE), dusts (DP), seed-dressing products, granules in the form of microgranules, spray granules, coated granules and absorption granules, water-dispersible granules (WG), ULV formulations, microcapsules, waxes or baits.

These individual types of formulations are known in principle and are described, for example, in:

Winnacker-Küchler, "Chemical Technology", Volume 7, C. Hauser Verlag Munich, 4th Edition 1986; van Falkenberg, "Pesticides Formulations", Marcel Dekker N.Y., 2nd Ed. 1972-73; K. Martens, "Spray Drying Handbook", 3rd Ed. 1979, G. Goodwin Ltd. London.

The necessary formulation auxiliaries such as inert materials, surfactants, solvents and other additives, are also known and described, for example, in Watkins, "Handbook of Insecticide Dust Diluents and Carriers", 2nd Ed., Darland Books, Caldwell N.J.; H. v. Olphen, "Introduction to Clay Colloid Chemistry", 2nd Ed., J. Wiley & Sons, N.Y.; Marsden, "Solvents Guide", 2nd Ed., Interscience, N.Y. 1950; McCutcheon's, "Detergents and Emulsifiers Annual", MC Publ. Corp., Ridgewood N.J.; Sisley and Wood, "Encyclopedia of Surface Active Agents", Chem. Publ. Co. Inc., N.Y. 1964;

Schönfeldt, "Grenzflächenaktive Äthylenoxidaddukte", [Surface-active ethylene oxide adducts] Wiss. Verlagsgesell., Stuttgart 1967; Winnacker-Küchler, "Chemische Technologie", Volume 7, C. Hauser Verlag Munich, 4th Edition 1986.

5

Based on these formulations, it is also possible to prepare combinations with other pesticidally active materials, fertilizers and/or growth regulators, for example in the form of a ready-mix formulation or a tank mix. Wettable powders are preparations which are uniformly dispersible in water which, besides the active substance, also comprise wetters, for example polyoxyethylated alkylphenols, polyoxyethylated fatty alcohols, alkylsulfonates or alkylphenolsulfonates and dispersants, for example sodium lignosulfonate or sodium 2,2'-dinaphthylmethane-6,6'-disulfonate, in addition to a diluent or inert material.

15

Emulsifiable concentrates are prepared by dissolving the active substance in an organic solvent, for example butanol, cyclohexanone, dimethylformamide, xylene or else higher-boiling aromatics or hydrocarbons, with addition of one or more emulsifiers. As emulsifiers, the following can be used, for example: calcium alkylarylsulfonates such as calcium dodecylbenzenesulfonate, or nonionic emulsifiers such as fatty acid polyglycol esters, alkylaryl polyglycol ethers, fatty alcohol polyglycol ethers, propylene oxide/ethylene oxide condensates, alkyl polyethers, sorbitan fatty acid esters, polyoxyethylene sorbitan fatty acid esters or polyoxyethylene sorbitol esters.

25

Dusts are obtained, for example, by grinding the active substance with finely divided solid materials, for example talc or natural clays, such as kaolin, bentonite, pyrophyllite or diatomaceous earth. Granules can be prepared either by atomizing the active substance onto adsorptive, granulated inert material or by applying active substance concentrates onto the surface of carrier materials such as sand or kaolinites, or of granulated inert material, by means of adhesives, for example polyvinyl alcohol or sodium polyacrylate, or else mineral oils. Suitable active substances can also be granulated in the manner which is customary for the preparation of fertilizer granules, if desired as a mixture with fertilizers.

30

35

The active substance concentration in wettable powders is, for example, approximately 10 to 90% by weight, the remainder to 100% by weight is

composed of customary formulation auxiliaries. In the case of emulsifiable concentrates, the active substance concentration may be approximately 5 to 80% by weight. Formulations in the form of dusts usually comprise 5 to 20% by weight of active substance, sprayable solutions approximately 2 to 20% by weight. In the case of granules, the active substance content depends partly on whether the active compound is in liquid or solid form and on which granulation auxiliaries, fillers and the like are being used.

Besides, the abovementioned active substance formulations comprise, if appropriate, the tackifiers, wetters, dispersants, emulsifiers, penetrants, solvents, fillers or carriers which are conventional in each case.

For use, the concentrates, which are present in commercially available form, are, if desired, diluted in the customary manner, for example in the case of wettable powders, emulsifiable concentrates, dispersions and in some cases also microgranules using water. Preparations in the form of dusts and granules and sprayable solutions are usually not diluted any further with other inert substances prior to use.

The application rate required varies with the external conditions such as, inter alia, temperature and humidity. It may vary within wide limits, for example between 0.0005 and 10.0 kg/ha or more of active substance, but it is preferably between 0.001 and 5 kg/ha.

The active substances according to the invention, in their commercially available formulations and in the use forms prepared from these formulations (see the abovementioned compositions) may be present in mixtures with other active substances such as insecticides, attractants, sterilants, acaricides, nematocides, fungicides, molluskicides, growth regulatory substances or herbicides.

The pesticides include, for example, phosphoric esters, carbamates, carboxylic esters, formamidines, tin compounds and materials produced by microorganisms.

35

Preferred components in mixtures are:

1. from the group of the phosphorus compounds

acephate, azamethiphos, azinphos-ethyl, azinphos-methyl, bromophos,
 bromophos-ethyl, cadusafos (F-67825), chlorethoxyphos, chlorfenvinphos,
 chlormephos, chlorpyrifos, chlorpyrifos-methyl, demeton, demeton-S-
 methyl, demeton-S-methyl sulfone, dialifos, diazinon, dichlorvos,
 5 dicrotophos, dimethoate, disulfoton, EPN, ethion, ethoprophos, etrimfos,
 famphur, fenamiphos, fenitriothion, fensulfothion, fenthion, fonofos,
 formothion, fosthiazate (ASC-66824), heptenophos, isazophos, isothioate,
 isoxathion, malathion, methacrifos, methamidophos, methidathion,
 salithion, mevinphos, monocrotophos, naled, omethoate, oxydemeton-
 10 methyl, parathion, parathion-methyl, phenthoate, phorate, phosalone,
 phosolan, phosphocarb (BAS-301), phosmet, phosphamidon, phoxim,
 pirimiphos, pirimiphos-ethyl, pirimiphos-methyl, profenofos, propaphos,
 proetamphos, prothiofos, pyraclofos, pyridapenthion, quinalphos, sulprofos,
 temephos, terbufos, tebupirimfos, tetrachlorvinphos, thiometon, triazophos,
 15 trichlorphon, vamidothion;

2. from the group of the carbamates

alanycarb (OK-135), aldicarb, 2-sec-butylphenyl methylcarbamate (BPMC),
 carbaryl, carbofuran, carbosulfan, cloethocarb, benfuracarb, ethiofencarb,
 20 furathiocarb, HCN-801, isoprocarb, methomyl, 5-methyl-m-cumenylbutyryl
 (methyl)carbamate, oxamyl, pirimicarb, propoxur, thiodicarb, thiofanox, 1-
 methylthio(ethylideneamino)-N-methyl-N-(morpholinothio)carbamate (UC
 51717), triazamate;

25 3. from the group of the carboxylic esters

acrinathrin, allethrin, alphametrin, 5-benzyl-3-furylmethyl (E)-(1R)-cis-
 2,2-dimethyl-3-(2-oxothioloan-3-ylidenemethyl)cyclopropanecarboxylate,
 beta-Cyfluthrin, beta-cypermethrin, bioallethrin, bioallethrin
 ((S)-cyclopentylisomer), bioresmethrin, bifenthrin, (RS)-1-cyano-
 30 1-(6-phenoxy-2-pyridyl)methyl (1RS)-trans-3-(4-tert-butylphenyl)-
 2,2-dimethylcyclopropanecarboxylate (NCI 85193), cycloprothrin, cyfluthrin,
 cyhalothrin, cythithrin, cypermethrin, cyphenothrin, deltamethrin,
 empenthrin, esfenvalerate, fenfluthrin, fenpropathrin, fenvalerate,
 flucythrinate, flumethrin, fluvalinate (D isomer), imiprothrin (S-41311),
 35 lambda-cyhalothrin, permethrin, phenothrin ((R) isomer), prallethrin,
 pyrethrins (natural products), resmethrin, tefluthrin, tetramethrin,
 theta-Cypermethrin (TD-2344), tralomethrin, transfluthrin,
 zeta-cypermethrin (F-56701);

4. from the group of the amidines
amitraz, chlordimeform;
5. from the group of the tin compounds
5 cyhexatin, fenbutatin oxide;
6. others
abamectin, ABG-9008, acetamiprid, anagrapha falcitera, AKD-1022, AKD-3059, ANS-118, bacillus thuringiensis, beauveria bassiana, bensultap,
10 bifenazate (D-2341), binapacryl, BJL-932, bromopropylate, BTG-504, BTG-505, buprofezin, camphechlor, cartap, chlorobenzilate, chlorfenapyr, chlorfluazuron, 2-(4-chlorophenyl)-4,5-diphenylthiophene (UBI-T 930), chlorfentezine, chromafenozide (ANS-118), CG-216, CG-217, CG-234, A-184699, 2-naphthylmethyl cyclopropanecarboxylate (Ro12-0470),
15 cyromazin, diaclofen (thiamethoxam), diafenthiuron, ethyl 2-chloro-N-(3,5-dichloro-4-(1,1,2,3,3,3-hexafluoro-1-propyloxy)phenyl)carbamoyl)-2-carboximidate, DDT, dicofol, diflubenzuron, N-(2,3-dihydro-3-methyl-1,3-thiazol-2-ylidene)-2,4-xylidine, dinobuton, dinocap, diofenolan, DPX-062, emamectin benzoate (MK-244), endosulfan, ethiprole (sulfethiprole),
20 ethofenprox, etoxazole (YI-5301), fenazaquin, fenoxycarb, fipronil, fluazuron, flumite (flufenazine, SZI-121), 2-fluoro-5-(4-(4-ethoxyphenyl)-4-methyl-1-pentyl)diphenyl ether (MTI 800), granulosis and nuclear polyhedrosis viruses, fenpyroximate, fenthioncarb, flubenzimine, flucycloxuron, flufenoxuron, flufenprox (ICI-A5683), fluproxyfen, gamma-
25 HCH, halofenozide (RH-0345), halofenprox (MTI-732), hexaflumuron (DE_473), hexythiazox, HOI-9004, hydramethylnon (AC 217300), lufenuron, imidacloprid, indoxacarb (DPX-MP062), kanemite (AKD-2023), M-020, MTI-446, ivermectin, M-020, methoxyfenozide (intrepid, RH-2485), milbemectin, NC-196, neemgard, nitenpyram (TI-304), 2-nitromethyl-4,5-dihydro-6H-thiazine (DS 52618), 2-nitromethyl-3,4-dihydrothiazole (SD 35651), 2-nitromethylene-1,2-thiazinan-3-ylcarbamaldehyde (WL 108477), Pyriproxyfen (S-71639), NC-196, NC-1111, NNI-9768, Novaluron (MCW-275), OK-9701, OK-9601, OK-9602, propargite, pymethrozine, pyridaben, pyrimidifen (SU-8801), RH-0345, RH-2485, RYI-210, S-1283, S-1833,
35 SB7242, SI-8601, silafluofen, silomadine (CG-177), spinosad, SU-9118, tebufenozide, tebufenpyrad (MK-239), teflubenzuron, tetradifon, tetrasul, thiacloprid, thiocyclam, TI-435, tolfenpyrad (OMI-88), triazamate (RH-7988), triflumuron, verbutin, vertalec (mykotal), YI-5301.

The active substance content of the use forms prepared from the commercially available formulations may range from 0.00000001 up to 95% by weight of active substance, preferably between 0.00001 and 1% by weight. Application is effected in a customary manner adapted to suit the use forms.

The invention also relates to a method of controlling harmful insects, Acarina, mollusks and/or nematodes in which an effective amount of a compound according to the invention or of a composition according to the invention is applied to these or to the plants, areas or substrates infested with them.

The use of a compound according to the invention or of a composition according to the invention for controlling harmful insects, Acarina, mollusks and/or nematodes is also the subject of the invention.

The active substances according to the invention are also suitable for use in the veterinary medicine sector, preferably for controlling endoparasites and ectoparasites, and in the field of animal keeping.

The active substances according to the invention may be applied in the known manner, such as by oral administration in the form of, for example, tablets, capsules, drinks or granules, by dermal application in the form of, for example, dipping, spraying, pouring on and spotting on, and dusting, and by parenteral administration in the form of, for example, an injection.

In addition, the compounds according to the invention are also suitable for use in technology, for example as wood preservative, as preservative in paints, in cooling lubricants for metalworking, or as preservative in drilling and cutting oils.

Accordingly, the compounds of the formula (I) according to the invention can also be employed particularly advantageously in livestock keeping (for example cattle, sheep, pigs and poultry such as chickens, geese and the like). In a preferred embodiment of the invention, the novel compounds, if appropriate in suitable formulations (cf. above), are administered orally to the animals, if appropriate together with the drinking water or feed. Since excretion in the feces is highly efficient, the development of insects in the animals' feces can be prevented very easily in this manner. The dosages

and formulations which are suitable in each case depend, in particular, on the species and the developmental stage of the productive livestock and also on the risk of infestation and can be determined readily and established by customary methods. For example, the compounds can be employed in cattle at dosages of 0.01 to 1 mg/kg bodyweight.

The compounds of the formula (I) according to the invention are also distinguished by an outstanding fungicidal action. Fungal pathogens which have already penetrated the plant tissue can be controlled successfully in a curative manner. This is especially important and advantageous in the case of those fungal diseases which can no longer be controlled effectively by the fungicides which are otherwise customary once infection has taken place. The spectrum of action of the claimed compounds encompasses a variety of economically important phytopathogenic fungi such as, for example, *Plasmopara viticola*, *Phytophthora infestans*, *Erysiphe graminis*, *Pyricularia oryzae*, *Pyrenophora teres*, *Leptosphaeria nodorum* and *Pellicularia sasakii* and *Puccinia recondita*.

These, but also predominantly insecticidally, acaricidally, molluskicidally or nematocidally active compounds of the formula (I), can be applied, in their commercially available formulations, either alone or in combination with other fungicides known from the literature.

Fungicides which are known from the literature and which can be combined in accordance with the invention with the compounds of the formula (I) are, for example, the following products:

aldimorph, andoprim, anilazine, BAS 480F, BAS 450F, benalaxyl, benodanil, benomyl, binapacryl, bitertanol, bromuconazole, buthiobate, captafol, captan, carbendazim, carboxin, CGA 173506, cyprofuram, dichlofluanid, dichlomezine, diclobutrazole, diethofencarb, difenconazole (CGA 169374), difluconazole, dimethirimol, dimethomorph, diniconazole, dinocap, dithianon, dodemorph, dodine, edifenfos, ethirimol, etridiazol, fenarimol, fenfuram, fenpiclonil, fenpropidin, fenpropimorph, fentin acetate, fentin hydroxide, ferimzone (TF164), fluazinam, fluobenzimine, fluquinconazole, fluorimide, flusilazole, flutolanil, flutriafol, folpet, fosetyl aluminum, fuberidazole, fulsulfamide (MT-F 651), furalaxyl, furconazole, furmecyclox, guazatine, hexaconazole, ICI A5504, imazalil, imibenconazole, iprobenfos, iprodione, isoprothiolane, KNF 317, copper compounds such as copper oxychloride, oxine-copper, copper oxide,

mancozeb, maneb, mepanipyrim (KIF 3535), metconazole, mepronil, metalaxyl, methasulfocarb, methfuroxam, MON 24000, myclobutanil, nabam, nitrothalidopropyl, nuarimol, ofurace, oxadixyl, oxycarboxin, penconazole, pencycuron, PP 969, probenazole, propineb, prochloraz, 5 procymidon, propamocarb, propiconazole, prothiocarb, pyracarbolid, pyrazophos, pyrifenox, pyroquilon, rabenzazole, RH7592, sulfur, tebuconazole, TF 167, thiabendazole, thicyofen, thiofanate-methyl, thiram, tolclofos-methyl, tolylfluanid, triadimefon, triadimenol, tricyclazole, tridemorph, triflumizol, triforine, validamycin, vinchlozolin, XRD 563, zineb, 10 sodium dodecylsulfonate, sodium dodecyl sulfate, sodium C13/C15-alcohol ether sulfonate, sodium cetostearyl phosphate ester, dioctylsodium sulfosuccinate, sodium isopropyl naphthalenesulfonate, sodium methylenebisnaphthalenesulfonate, cetyltrimethylammonium chloride, salts of long-chain primary, secondary or tertiary amines, alkylpropyleneamines, 15 laurylpyrimidinium bromide, ethoxylated quarternized fatty amines, alkyldimethylbenzylammonium chloride and 1-hydroxyethyl-2-alkylimidazoline.

The abovementioned components for combinations are known active 20 substances, many of which are described in C.D.S. Tomlin, S.B. Walker, The Pesticide Manual, 11th Edition, British Crop Protection Council, Farnham 1997.

The active substance content of the use forms prepared from the 25 commercially available formulations can be varied within wide ranges, the active substance concentration of the use forms may range from 0.0001 up to 95% by weight of active substance, preferably from 0.0001 to 1% by weight. They are applied in a customary manner adapted to suit the use forms.

30 They are used in a customary manner adapted to suit the use forms, for example by applying, in order to control pathogenic fungi, a fungicidally effective amount of a compound according to the invention or of a composition according to the invention to these fungi or to the plants, areas 35 of substrates infested with them, or to seed.

The invention also relates to seed, treated or coated with an effective amount of a compound according to the invention or of a composition according to the invention.

The compounds of the formula (I) can also be employed for controlling harmful organisms in crops of known genetically engineered plants or genetically engineered plants yet to be developed. As a rule, the transgenic plants are distinguished by especially advantageous properties, for example by resistances to particular crop protection agents, resistances to plant diseases or pathogens of plant diseases, such as particular insects or microorganisms such as fungi, bacteria or viruses. Other particular properties concern, for example, the harvested material with regard to quantity, quality, storage properties, composition and specific constituents. Thus, transgenic plants are known where the starch content is increased, or the starch quality is altered, or where the harvested material has a different fatty acid composition.

The use in economically important transgenic crops of useful plants and ornamentals is preferred, for example of cereals such as wheat, barley, rye, oats, millet, rice, cassava and maize or else crops of sugar beet, cotton, soya, oilseed rape, potatoes, tomatoes, peas and other types of vegetables.

When used in transgenic crops, in particular those which have resistances to insects, effects are frequently observed, in addition to the effects against harmful organisms to be observed in other crops, which are specific for application in the transgenic crop in question, for example an altered or specifically widened spectrum of pests which can be controlled, or altered application rates which may be employed for application.

The invention therefore also relates to the use of compounds of the formula (I) for controlling harmful organisms in transgenic crop plants.

The use according to the invention of compounds of the formula (I) or of compositions comprising them, for example as insecticide, acaricide, molluskicide, nematocide or fungicide, also includes the case where the compound of the formula (I) or its salt is formed from a precursor only after application, for example in the insect, in a plant or in the soil.

The contents of the German Patent Application 198 15 026.1, whose priority is claimed by the present application, and the contents of the

appended summary are therefore expressly incorporated by reference; they are part of the present description by way of quotation.

5 The examples which follow are intended to illustrate the invention without restricting it thereto.

A. Preparation Examples

Precursor VP1: 4'-bromo-2-hydroxyacetophenone O-methyloxime

10

Sodium formate (244.6 g, 3597 mmol) was introduced into ethanol/water 7:3 (1 l). 2,4'-dibromoacetophenone (200 g, 719 mmol) was added portionwise at approx. 60°C, and the mixture was then refluxed for 7 hours. Water (1 l) was then admixed. Upon cooling, the hydroxy ketone
15 crystallized and was isolated by filtration with suction. This gave the hydroxy ketone intermediate (143.6 g, = 93% yield) in its pure form. It was mixed with O-methylhydroxylamine hydrochloride (59.0 g, 706 mmol) and sodium acetate (57.9 g, 706 mmol) in dioxane/water 9:1 (700 ml) and the mixture was refluxed for 3 hours. After stirring with water (1.3 l) and
20 extraction with heptane/ethyl acetate 1:1, 4'-bromo-2-hydroxyacetophenone O-methyloxime, 174.3 g, was obtained as an oil, purity approx. 90%, 2 isomers.

25 Precursor VP2: 2-amino-2-(4-bromophenyl)ethanol (amino alcohol of the formula VI)

4'-bromo-2-hydroxyacetophenone O-methyloxime (54.8 g, 224 mmol) were added dropwise at 20°C under nitrogen as a solution in THF to a mixture of lithium aluminum hydride (20 g, 505 mmol) and aluminum chloride (15.3 g,
30 112 mmol) in THF (350 ml). After 4 hours, the mixture was hydrolyzed (40 ml methanol; 200 ml 2N NaOH) and extracted with heptane/ethyl acetate. The crude product (42 g) was recrystallized from heptane/ethyl acetate 2:1. This gave 2-amino-2-(4-bromophenyl)ethanol, 29.3 g, colorless crystals, m.p. 95°C.

35

Precursor VP3: N-[1-(4'-bromophenyl)-2-hydroxyethyl]-2,6-difluorobenzamide (formula VII)

- 2,6-difluorobenzoyl chloride (13.8 ml, 110 mmol) was added dropwise at 20°C to 2-amino-2-(4-bromophenyl)ethanol (25.0 g, 116 mmol) and triethylamine (19.5 ml, 139 mmol) in THF (150 ml). After 15 hours, the mixture was stirred with water and extracted with heptane/ethyl acetate.
- 5 This gave the amido alcohol product, 38.6 g, pale yellow solid, m.p. 141°C .

Precursor VP4: 2-(2,6-difluorophenyl)-4-(4-bromophenyl)oxazoline (formula III)

Synthesis using the thionyl chloride/alkali metal hydroxide method

- 10 The amido alcohol VP3 (38.5 g, 108 mmol) was mixed in dichloromethane with thionyl chloride (9.7 ml, 130 mmol) and heated for 5 hours at 40°C. The mixture was concentrated and the residue (= amidochloride, 36.5 g) was refluxed with 6N NaOH (32.5 ml) in dioxane (180 ml) (6 hours). The mixture was stirred with water and extracted with ethyl acetate. The crude
- 15 product (oil, 31.8 g) was recrystallized from heptane/ethyl acetate 9:1. This gave 2-(2,6-difluorophenyl)-4-(4-bromophenyl)oxazoline, 17.8 g, colorless crystals, m.p. 98°C; b.p. 365°C (GC).

Synthesis using the tosyl chloride/amine method

- 20 The amido alcohol VP3 (5.0 g, 14.0 mmol) was mixed in dichloromethane (40 ml) with triethylamine (5.67 g, 56 mmol) and tosyl chloride (2.94 g, 15.5 mmol) and refluxed for 5 hours. The mixture was concentrated, and the residue was stirred with water and extracted with ethyl acetate. The crude product (4.6 g) was recrystallized with ethanol/water 8:2. This gave
- 25 2-(2,6-difluorophenyl)-4-(4-bromophenyl)oxazoline, 3.4 g, colorless crystals, m.p. 97°C; ¹H NMR (CDCl₃, ppm): 4.24, 4.81, 5.43, oxazoline; 7.00, 7.43, F₂C₆H₃; 7.23, 7.51, BrC₆H₄ .

- Precursor VP5: 2-(2,6-difluorophenyl)-4-(4-trimethylstannylphenyl)-oxazoline (formula IX)
- 30

- VP3 2-(2,6-difluorophenyl)-4-(4-bromophenyl)oxazoline (16.4 g, 48.8 mmol) was introduced into dioxane under nitrogen, hexamethyldistannane (20 g, 61.2 mmol), Pd(PPh₃)₄ (1.4 g, 1.2 mmol) and lithium chloride (0.38 g, 4.4 mmol) were added and the mixture was refluxed for 7 hours. The reaction mixture was stirred with water and extracted with heptane/ethyl acetate. The crude product (19.6 g, oil) contained approximately 70% of the Sn product. Separation by chromatography (eluent: heptane/ethyl acetate
- 35

2:1) yielded 2-(2,6-difluorophenyl)-4-(4-trimethylstannylphenyl)oxazoline, 13.4 g, oil, turned crystalline, DC Rf 0.59 (VP4 Rf 0.49).

5 Precursor VP6: 5-bromo-2-trifluoroethoxypyridine

Sodium hydride (18.6 g, 60%, 464 mmol) was introduced into DMF (500 ml) under nitrogen. Trifluoroethanol (32.2 ml, 443 mmol) was added dropwise at 10°C. After 1 hour, 2,5-dibromopyridine (100 g, 422 mmol) was added portionwise, and the mixture was stirred in a water bath for 24 hours at 20°C. It was then stirred with water and extracted with heptane/ethyl acetate 9:1. Distillation of the crude product (105 g) yielded 5-bromo-2-trifluoroethoxypyridine, 82.1 g (purity 93%), colorless fluid, b.p. 97-104°C at 18 mbar.

15 Other alkoxypyridines and alkylthiopyridines were prepared analogously.

Precursor VP7: 5,5-dimethyl-2-(2-trifluoroethoxypyridin-5-yl)-1,3,2-dioxaborinane

20 5-Bromo-2-trifluoroethoxypyridine (30.0 g, 117 mmol) was introduced into THF/diethyl ether 2:1 (250 ml) under nitrogen and cooled. An n-butyllithium solution (2.5 M, 53.9 ml, 135 mmol) was added at approx. -85°C using a syringe. After 10 minutes, isopropyl borate (33.1 ml, 141 mmol) was added dropwise at approx. -80°C. The temperature of the mixture was allowed to climb to -10°C, and acetic acid (10.1 ml, 176 mmol) and 2,2-dimethylpropanediol (15.9 g, 152 mmol) were added. After 15 hours at 20°C, the mixture was stirred with water and extracted with heptane/ethyl acetate. The crude product (31.5 g) was recrystallized using heptane/ethyl acetate 95:5. This gave 5,5-dimethyl-2-(2-trifluoroethoxypyridin-5-yl)-1,3,2-dioxaborinane, 20.3 g, white solid; ¹H NMR (CDCl₃, ppm): 1.02, Me; 3.75, OCH₂; 4.77, PyOCH₂; 6.82, 7.97, 8.50, PyH.

Other 2-pyridyl-1,3,2-dioxaborinanes were prepared analogously.

35 Precursor VP8: 2-Ethoxy-5-trimethylstannylpyridine

5-Bromo-2-ethoxypyridine (14.9 g, 68 mmol) were introduced into THF/diethyl ether 2:1 (120 ml) under nitrogen. n-Butyllithium/hexane solution (2.5 M, 33 ml, 82 mmol) was added at -80°C. After 15 minutes,

chlorotrimethyltin (16.4 g, 82 mmol), dissolved in THF, was added dropwise. The temperature of the mixture was allowed to rise to 0°C, and the mixture was stirred with water and extracted with heptane/ethyl acetate. The crude product (15.7 g) was distilled in vacuo. This gave 2-ethoxy-5-trimethylstannylpyridine, 11.2 g, b.p. 66-75°C at 1.2 mbar. Other stannylpyridines were prepared analogously.

Precursor VP9: 5-[4-1-Amino-2-hydroxyethyl]-2-trifluoroethoxypyridine

2-Amino-2-(4-bromophenyl)ethanol VP2 (3.8 g, 17.1 mmol) and 5,5-dimethyl-2-(2-trifluoroethoxypyridin-5-yl)-1,3,2-dioxaborinane VP7 (6.55 g, 20.4 mmol) were refluxed together with tetrakis(triphenylphosphine)palladium (0.63 g, 3 mol%), sodium carbonate (3.7 g, 35.4 mmol) in toluene/ethanol/water (8:2:1, 50 ml) for 7 hours. After extractive work-up and purification by column chromatography, 5-[4-(1-amino-2-hydroxyethyl)phenyl]-2-trifluoroethoxypyridine, 2.2 g, m.p. 129°C were obtained.

Oxazoline products:

2-(2,6-Difluorophenyl)-4-[4-(2-ethoxypyridin-5-yl)phenyl]oxazoline (Ex. No. 2)

2-(2,6-Difluorophenyl)-4-(4-bromophenyl)oxazoline (VP4) (5.00 g, 14.8 mmol) was mixed with 2-ethoxy-5-trimethylstannylpyridine (VP8) (5.06 g, 17.8 mmol), Pd(PPh₃)₄ (0.68 g, 0.59 mmol) and lithium chloride (0.12 g, 2.9 mmol) in dioxane (40 ml) and the mixture was refluxed under nitrogen (7 hours). After extraction and chromatography, 2-(2,6-difluorophenyl)-4-[4-(2-ethoxypyridin-5-yl)phenyl]oxazoline was obtained, 3.86 g, m.p. 119°C ; ¹H NMR (CDCl₃, ppm): 1.42, 4.35, OCH₂CH₃; 4.40, 4.84, 5.51, oxazoline; 6.79, 7.77, 8.36, pyridine; 7.00, 7.41, 7.54, phenyl .

2-(2,6-Difluorophenyl)-4-[4-(2-trifluoroethoxypyridin-5-yl)phenyl]oxazoline (Ex. No. 3)

2-(2,6-Difluorophenyl)-4-(4-bromophenyl)oxazoline (VP4) (2.0 g, 5.9 mmol) was mixed with 2-trifluoroethoxy-5-trimethylstannylpyridine (2.61 g, 7.7 mmol), tetrakis(triphenylphosphine)palladium (0.22 g, 0.18 mmol) and

lithium chloride (0.38 g, 8.9 mmol) in dioxane (30 ml) and refluxed for 9 hours. After extractive work-up and column chromatography, 2-(2,6-difluorophenyl)-4-[4-(2-trifluoroethoxypyridin-5-yl)phenyl]oxazoline was obtained, 1.37 g, colorless crystals.

5 ^{19}F NMR (CDCl_3): -75.05 (CF_3), -110.45 (ArF) .

2-(2,6-Difluorophenyl)-4-[4-(2-n-propyloxypyridin-5-yl)phenyl]oxazoline (Ex. No. 4)

10 2-(2,6-Difluorophenyl)-4-(4-bromophenyl)oxazoline (VP4) (2.0 g, 2.9 mmol) was mixed with 2-n-propyloxy-5-trimethylstannylpyridine (4.6 g, 50%, 7.7 mmol), $\text{Pd}(\text{PPh}_3)_4$ (0.35 g, 0.3 mmol) and lithium chloride (50 mg) in dioxane and refluxed under nitrogen (7 hours). After chromatography and recrystallization in heptane/ethyl acetate 95:5, 2-(2,6-difluorophenyl)-
15 4-[4-(2-n-propyloxypyridin-5-yl)phenyl]oxazoline was obtained, 1.21 g, colorless crystals;
 ^1H NMR (CDCl_3 , ppm): 1.05, 1.82, 4.29, OC_3H_7 ; 4.32, 4.86, 5.51, oxazoline; 6.80, 7.78, 8.37, pyridine; 7.02, 7.42, 7.54, phenyl.

20 2-(2,6-Difluorophenyl)-4-[4-(2-isobutyloxypyridin-5-yl)phenyl]oxazoline (Ex. No. 8)

2-(2,6-Difluorophenyl)-4-(4-bromophenyl)oxazoline (1.00 g, 2.96 mmol) was mixed with 5,5-dimethyl-2-(2-isobutyloxypyridin-5-yl)-
25 1,3,2-dioxaborinane (1.4 g, purity 80%, 4.3 mmol), $\text{Pd}(\text{PPh}_3)_4$ (0.17 g, 0.15 mmol) and sodium carbonate (0.62 g, 5.9 mmol) in toluene/ethanol/water 8:2:1 (15 ml) and refluxed for 8 hours under nitrogen. After extraction and chromatography, 2-(2,6-difluorophenyl)-
30 4-[4-(2-isobutyloxypyridin-5-yl)phenyl]oxazoline was obtained, 1.14 g, colorless oil.
 ^1H NMR (CDCl_3 , ppm): 1.04, 2.11, 4.10, o-isobutyl; 4.33, 4.84, 5.51, oxazoline, 7.00, 7.42, 7.55, phenyl; 6.81, 7.77, 8.37, pyridine.

35 2-(2,6-Difluorophenyl)-4-[4-(2-trifluoroethoxypyridin-5-yl)phenyl]thiazoline (Ex. No. 19)

2-(2,6-Difluorophenyl)-4-(4-bromophenyl)thiazoline (0.90 g, 2.5 mmol, obtained from precursor VP3 and Lawesson's Reagent in toluene) was mixed with 2-trifluoroethoxy-5-trimethylstannylpyridine (1.21 g, 3.6 mmol),

$\text{Pd(PPh}_3)_4$ (0.15 g) and lithium chloride (0.16 g) in dioxane (10 ml) and the mixture was refluxed for 8 hours. After extraction and chromatography, 2-(2,6-difluorophenyl)-4-[4-(2-trifluoroethoxypyridin-5-yl)phenyl]thiazoline was obtained, 0.46 g, beige solid; MS: M^+ 450.

- 5 $^1\text{H NMR}$ (CDCl_3 , ppm): 3.45, 3.94, 5.83, thiazoline; 4.82, OCH_2CF_3 ; 6.95, 7.87, 8.36, pyridine; 7.00, 7.40, 7.53, phenyl.

2-(2,6-Difluorophenyl)-4-[4-(2-n-propylpyridin-5-yl)phenyl]oxazoline (Ex. No. 34)

10

2-(2,6-Difluorophenyl)-4-(4-bromophenyl)oxazoline (0.8 g, 2.4 mmol) was refluxed for 8 hours with dimethyl-2-(2-n-propylpyridin-5-yl)-1,3,2-dioxaborinane (0.72 g, 3.1 mmol), $\text{Pd(PPh}_3)_4$ (0.14 g, 0.12 mmol) and sodium carbonate (0.5 g) in toluene/ethanol/water 8:2:1 (11 ml). After extraction and chromatography, 2-(2,6-difluorophenyl)-4-[4-(2-n-propylpyridin-5-yl)phenyl]oxazoline was obtained, 0.20 g, pale solid.

15

$^1\text{H NMR}$ (CDCl_3 , ppm): 1.01, 1.80, 2.81, n-propyl; 4.34, 4.85, 5.53, oxazoline; 7.00, 7.44, 7.59, phenyl; 7.21, 7.79, 8.76, pyridyl.

- 20 2-(2,6-Difluorophenyl)-4-[4-(2-ethylsulfinylpyridin-5-yl)phenyl]oxazoline (Ex. No. 40)

2-(2,6-Difluorophenyl)-4-[4-(2-ethylthiopyridin-5-yl)phenyl]oxazoline (Ex. No. 39; 0.60 g, 1.5 mmol) was stirred for 5 hours at 20°C with 3-chloroperbenzoic acid (0.41 g, 1.7 mmol) in dichloroethane (10 ml). After extraction and chromatography, 2-(2,6-difluorophenyl)-4-[4-(2-ethylsulfinylpyridin-5-yl)phenyl]oxazoline was obtained, 0.21 g, colorless oil.

25

$^1\text{H NMR}$ (CDCl_3 , ppm): 1.24, 2.98, 3.21, $\text{SO-CH}_2\text{CH}_3$; 4.33, 4.87, 5.56, oxazoline; 7.01, 7.48, 7.64, phenyl; 8.03, 8.11, 8.84, pyridine.

30

2-(2,6-Difluorophenyl)-4-[4-(2-(N-acetylethylamino)pyridin-5-yl)phenyl]oxazoline (Ex. No. 51)

- 35 2-(2,6-Difluorophenyl)-4-(4-trimethylstannylphenyl)oxazoline (0.84 g, 2.0 mmol) was mixed with 2-(N-acetylethylamino)-5-iodopyridine (0.93 g, 3.2 mmol), $\text{Pd(PPh}_3)_4$ (0.12 g) and lithium chloride (0.15 g) in dioxane (10 ml) and the mixture was refluxed for 8 hours. After separation by chromatography (eluent: ethyl acetate), 2-(2,6-difluorophenyl)-4-[4-(2-(N-

acetylethylamino)pyridin-5-yl)phenyl]oxazoline was obtained, 0.31 g, pale oil.

¹H NMR (CDCl₃, ppm): 1.19, 3.94, NC₂H₅; 2.08, COCH₃; 4.33, 4.87, 5.54, oxazoline; 7.00, 7.43, 7.48, 7.61, phenyl; 7.32, 7.96, 8.75, pyridine.

5

2-(2,6-Difluorophenyl)-4-[4-(1-n-propyl-2-pyridon-5-yl)phenyl]oxazoline (Ex. No. 152)

2-(2,6-Difluorophenyl)-4-(4-trimethylstannylphenyl)oxazoline (0.5 g, 1.2 mmol) was mixed with 1-n-propyl-5-bromo-2-pyridone (0.51 g, 2.4 mmol), Pd(PPh₃)₄ (70 mg) and lithium chloride (0.20 g) in dioxane (10 ml) and refluxed for 8 hours under nitrogen. After extractive work-up and chromatography (eluent ethyl acetate), 2-(2,6-difluorophenyl)-4-[4-(1-n-propyl-2-pyridon-5-yl)phenyl]oxazoline was obtained, 0.20 g, oil.

¹H NMR (CDCl₃, ppm): 1.00, 1.81, 3.96 NC₃H₇; 4.30, 4.83, 5.49, oxazoline; 6.64, pyridone; 7.00, 7.4-7.7, phenyl and pyridone.

2-(2-Chlorophenyl)-4-[4-(2-trifluoroethoxypyridin-5-yl)phenyl]oxazoline (Ex. No. 25)

20

2-(2-Chlorophenyl)-4-(4-bromophenyl)oxazoline (prepared in analogy to VP4) (0.80 g, 2.4 mmol) and 5,5-dimethyl-2-(2-trifluoroethoxypyridin-5-yl)-1,3,2-dioxaborinane VP7 (1.03 g, 3.6 mmol) were refluxed with Pd(PPh₃)₄ (0.1 g) and sodium carbonate (0.5 g) in toluene/ethanol/water (8:2:1, 11 ml) (8 hours). After extraction and column chromatography, 2-(2-chlorophenyl)-4-[4-(2-trifluoroethoxypyridin-5-yl)phenyl]oxazoline was obtained as a pale oil, 0.86 g; ¹H NMR (CDCl₃): 4.31, 4.80, 5.50, oxazoline; 4.80 OCH₂CF₃; 6.93, 7.3 - 7.5, 7.84, 8.35, pyridine and phenyl.

2-(2,6-Difluorophenyl)-4-[2-ethoxy-4-(2-trifluoroethoxypyridin-5-yl)phenyl]-oxazoline (Ex. No. 106)

Ex. 106 was prepared in analogy to Ex. No. 25.

¹H NMR (CDCl₃): 1.45, 4.15 OEt; 4.45, 4.92, 5.73, oxazoline, 4.81 OCH₂CF₃; 6.94, 7.84, 8.35, pyridine; 6.98, 7.12, 7.41, 7.53, phenyl.

2-(2-Fluorophenyl)-4-[4-(2-trifluoroethoxypyridin-5-yl)phenyl]oxazoline (Ex. No. 402)


Ex. No. 402 was prepared in analogy to Ex. No. 25.

¹H NMR (CDCl₃): 4.31, 4.82, 5.48, oxazoline; 4.80, OCH₂CF₃; 6.92, 7.84, 8.35, pyridine; 7.20, 7.40, 7.50, 8.00, phenyl.

A. Chemical Examples (Tables 1 – 8)

Table 1

Oxazolines and thiazolines of the formula (I), G= 3-pyridyl



Ex. No.	Ar	Z	R ⁵	R ⁶	Physical. data (m.p., MS)
1	Ar ¹	O	OCH ₃	H	m.p. 120°C
2	Ar ¹	O	OC ₂ H ₅	H	m.p. 118°C, NMR
3	Ar ¹	O	OCH ₂ CF ₃	H	NMR
4	Ar ¹	O	OnC ₃ H ₇	H	NMR
5	Ar ¹	O	OiC ₃ H ₇	H	
6	Ar ¹	O	OCH ₂ CF ₂ CF ₂ H	H	
7	Ar ¹	O	OnC ₄ H ₉	H	
8	Ar ¹	O	OCH ₂ CH(CH ₃) ₂	H	NMR
9	Ar ¹	O	OsecButyl	H	
10	Ar ¹	O	OnC ₅ H ₁₁	H	
11	Ar ¹	O	OCH(CH ₃)C ₃ H ₇	H	
12	Ar ¹	O	OnC ₆ H ₁₃	H	
13	Ar ¹	O	OnC ₈ H ₁₇	H	
14	Ar ¹	O	OC ₂ H ₅	Cl	
15	Ar ¹	O	OCH ₂ CF ₃	Cl	
16	Ar ¹	O	OC ₂ H ₅	CH ₃	
17	Ar ¹	O	OCH ₂ CF ₃	CH ₃	
18	Ar ¹	S	OC ₂ H ₅	H	
19	Ar ¹	S	OCH ₂ CF ₃	H	MS 450, NMR
20	Ar ¹	S	OnC ₃ H ₇	H	
21	Ar ²	O	OC ₂ H ₅	H	
22	Ar ²	O	OCH ₂ CF ₃	H	

23	Ar ²	O	OnC ₃ H ₇	H
24	Ar ³	O	OC ₂ H ₅	H
25	Ar ³	O	OCH ₂ CF ₃	H
26	Ar ³	O	OnC ₃ H ₇	H
27	Ar ⁴	O	OC ₂ H ₅	H
28	Ar ⁴	O	OCH ₂ CF ₃	H

m.p.: 136-137°C

¹H-NMR (CDCl₃, TMS: δ =
 2.46 (s, 6H, CH₃); 4.33, 4.88,
 5.57 (each dd, 1H, oxazoline);
 4.85 8q, 2H, CH₂CF₃); 6.97
 7.90, 8.39 (each dd, 1H, PyH);
 7.59, 7.51 (each m, 2H, C₆H₄),
 7.08-7.28 (m, 3H, C₆H₃).

¹⁹F-NMR (CDCl₃, CFCl₃):
 δ = - 74.3

29	Ar ⁴	O	OnC ₃ H ₇	H
30	Ar ¹	O	OCH(CF ₃)CH ₂ OCH ₃	H
31	Ar ¹	O	CH ₃	H
32	Ar ¹	O	CF ₃	H
33	Ar ¹	O	C ₂ H ₅	H
34	Ar ¹	O	nC ₃ H ₇	H
35	Ar ¹	O	nC ₄ H ₉	H
36	Ar ¹	O	nC ₅ H ₁₁	H
37	Ar ¹	O	nC ₆ H ₁₃	H
38	Ar ¹	O	SCF ₃	H
39	Ar ¹	O	SC ₂ H ₅	H
40	Ar ¹	O	SO-C ₂ H ₅	H
41	Ar ¹	O	SO ₂ -C ₂ H ₅	H
42	Ar ¹	O	SCH ₂ CF ₃	H
43	Ar ¹	O	S-nC ₃ H ₇	H
44	Ar ¹	O	SO-nC ₃ H ₇	H
45	Ar ¹	O	SO ₂ -nC ₃ H ₇	H
46	Ar ¹	O	S-nC ₄ H ₉	H
47	Ar ¹	O	Br	H
48	Ar ¹	O	Cl	Cl
49	Ar ¹	O	NH-COCH ₃	H

m.p. 117°C

m.p. 100°C

m.p. 90°C


MS 414, 416

50	Ar ¹	O	NH-COC ₂ H ₅	H	NMR m.p. 170°C
51	Ar ¹	O	N(C ₂ H ₅)COCH ₃	H	
52	Ar ¹	O	1-Morpholinyl	H	
53	Ar ¹	O	CH ₂ OCH ₃	H	
54	Ar ¹	O	CH ₂ OC ₂ H ₅	H	
55	Ar ¹	O	C ₂ H ₄ OCH ₃	H	m.p. 120°C
56	Ar ¹	O	OCH ₂ C ₆ H ₅	H	
57	Ar ¹	O	OCH ₂ CON(CH ₃) ₂	H	
58	Ar ¹	O	OC ₂ H ₄ CH(CH ₃) ₂	H	
59	Ar ¹	O	OCH ₂ CH(C ₂ H ₅)CH ₃	H	
60	Ar ¹	O	OCH ₂ -COOCH ₃	H	¹⁹ F NMR: -90.1, d; -110.5, ArF
61	Ar ¹	O	OCHF ₂	H	
62	Ar ¹	O	OCH ₂ CHF ₂	H	
63	Ar ¹	O	OCH(CH ₃)CF ₃	H	
64	Ar ¹	O	OC ₃ H ₆ CF ₃	H	
65	Ar ¹	O	OCH ₂ C ₂ F ₅	H	
66	Ar ¹	O	C ₂ H ₄ OC ₂ H ₅	H	
67	Ar ¹	O	N(CH ₃)C ₂ H ₅	H	
68	Ar ¹	O	N(CH ₃)nC ₃ H ₇	H	
69	Ar ¹	O	NHnC ₃ H ₇	H	
70	Ar ¹	O	NHnC ₄ H ₉	H	
71	Ar ¹	O	N(C ₂ H ₅) ₂	H	
72	Ar ¹	O	1-Piperidiny	H	
73	Ar ¹	O	SO-CF ₃	H	
74	Ar ¹	O	SCHF ₂	H	
75	Ar ¹	O	SO-CHF ₂	H	
76	Ar ¹	O	SO-CH ₂ CF ₃	H	
77	Ar ¹	O	SO ₂ -CH ₂ CF ₃	H	
78	Ar ¹	O	SO ₂ -CF ₃	H	
79	Ar ¹	O	SO ₂ -CHF ₂	H	
80	Ar ¹	O	SCH ₃	H	
81	Ar ¹	O	SO-CH ₃	H	
82	Ar ¹	O	SO ₂ -CH ₃	H	
83	Ar ¹	O	C ₂ H ₄ CF ₃	H	
84	Ar ¹	O	CH ₂ CF ₃	H	
85	Ar ¹	O	cC ₅ H ₉	H	
86	Ar ¹	O	CH ₂ cC ₆ H ₁₁	H	

87	Ar ¹	O	CN	H
88	Ar ¹	O	OC ₆ H ₅	H
89	Ar ¹	O	CH ₂ C ₆ H ₅	H
90	Ar ¹	O	4(OCF ₃)-C ₆ H ₄	H
91	Ar ¹	O	OCH ₂ CF ₃	CN
92	Ar ¹	O	OCH ₂ CF ₃	SCH ₃
93	Ar ¹	O	R ⁵ +R ⁶ = -C ³ H ₆ O-	
94	Ar ¹	O	R ⁵ +R ⁶ = -OC ₂ H ₄ O-	
95	Ar ³	O	SCF ₃	H
96	Ar ³	O	OCHF ₂	H
97	Ar ³	O	OCF ₂ CF ₂ H	H

Table 2

Oxazolines of the formula (I), G= 3-pyridyl




Ex. No.	X ¹	X ²	Z	E	R ¹	R ⁵	physical data
100	F	F	O	CH ₂	H	OC ₂ H ₅	m.p. 105°C
101	F	F	O	CH ₂	H	OCH ₂ CF ₃	
102	F	F	O	CH ₂	H	OnC ₃ H ₇	
103	F	F	O	-	2-F	OC ₂ H ₅	
104	F	F	O	-	2-Cl	OCH ₂ CF ₃	
105	F	F	O	-	2-OCH ₃	OC ₂ H ₅	
106	F	F	O	-	2-OC ₂ H ₅	OCH ₂ CF ₃	NMR
107	F	F	O	-	3-Cl	OnC ₃ H ₇	
108	F	F	O	-	2-F	OCH ₂ CF ₃	
109	F	F	O	-	2-Cl	OC ₂ H ₅	
110	F	F	O	-	2-OCH ₃	OCH ₂ CF ₃	
111	F	F	O	-	2-OC ₂ H ₅	OC ₂ H ₅	
112	F	F	O	-	2-OCH ₃	OnC ₃ H ₇	
113	F	F	O	-	2-OC ₂ H ₅	OnC ₃ H ₇	
114	F	F	O	-	3-F	OC ₂ H ₅	
115	F	Cl	O	-	2-OCH ₃	OC ₂ H ₅	
116	Br	H	O	-	H	OC ₂ H ₅	
117	Br	H	O	-	H	OCH ₂ CF ₃	
118	I	H	O	-	H	OnC ₃ H ₇	
119	F	H	O	-	H	OCH ₂ CF ₃	
120	CF ₃	H	O	-	H	OC ₂ H ₅	
121	OCH ₃	H	O	-	H	OnC ₃ H ₇	
122	CN	H	O	-	H	OC ₂ H ₅	
123	OCHF ₂	H	O	-	H	OC ₂ H ₅	
124	C ₂ H ₅	H	O	-	H	OC ₂ H ₅	
125	OCH ₃	F	O	-	H	OC ₂ H ₅	
126	F	H	O	-	H	OC ₂ H ₅	
127	OCH ₃	F	O	-	H	OCH ₂ CF ₃	

128	CF ₃	H	O	-	H	OCH ₂ CF ₃
129	CH ₃	H	O	-	H	OC ₂ H ₅
130	CH ₃	H	O	-	H	OCH ₂ CF ₃
131	F	F	O	-	CH ₃	OC ₂ H ₅
132	F	F	O	-	C ₂ H ₅	OC ₂ H ₅
133	F	F	O	-	OCHF ₂	OC ₂ H ₅
134	F	F	O	CH ₂ O	H	OC ₂ H ₅

Table 3


Oxazolines of the formula (I), G= 2-pyridon-5-yl



Ex.No.	X ¹	X ²	R ⁴	R ⁵	physical data
150	F	F	H	H	m.p. 197°C NMR
151	F	F	C ₂ H ₅	H	
152	F	F	nC ₃ H ₇	H	
153	F	F	CH ₂ -CH=CH ₂	H	
154	F	F	CH ₂ -CH=CH-CH ₃	H	
155	F	F	CH ₂ CON(CH ₃) ₂	H	
156	F	F	CHF ₂	H	
157	F	F	CH ₂ CF ₃	H	
158	F	F	CH ₂ C ₆ H ₅	H	
159	F	F	CH ₂ CH(CH ₃) ₂	H	
160	F	F	CH ₂ -CCH	H	
161	F	F	CH ₂ -CC-CH ₃	H	
162	F	F	cC ₆ H ₁₁	H	
162	F	F	cC ₅ H ₉	H	
163	F	F	CH ₂ cC ₆ H ₁₁	H	
164	F	F	C ₂ H ₄ OC ₂ H ₅	H	
165	F	F	CH ₂ OCH ₂ C ₆ H ₅	H	
166	F	F	C ₂ H ₅	3-Cl	
167	F	F	nC ₃ H ₇	3-Cl	
168	F	F	C ₂ H ₅	3-CH ₃	
169	F	F	nC ₃ H ₇	3-CH ₃	

Table 4

Oxazolines of the formula (I), G= 4-pyridyl





Ex. No.	X ¹	X ²	Z	R ⁵	physical data
170	F	F	O	H	m.p. 137°C
171	F	F	O	OC ₂ H ₅	
172	F	F	O	Cl	
173	F	Cl	O	OC ₂ H ₅	
174	F	F	O	C ₂ H ₅	
175	F	F	O	nC ₄ H ₉	
176	F	F	O	OCH ₂ CF ₃	
177	F	F	O	CF ₃	
178	F	F	O	OnC ₃ H ₇	
179	F	F	O	SC ₂ H ₅	
180	F	F	O	SO-C ₂ H ₅	
181	F	F	O	SO ₂ -C ₂ H ₅	
182	F	F	S	OC ₂ H ₅	
183	F	F	S	OnC ₃ H ₇	
184	F	F	O	OCH ₂ -3-Pyridyl	
185	F	F	O	1-Pyrrolyl	
186	F	F	O		

Table 5


Oxazolines of the formula (I), G= 2-Pyridon-3-yl



Ex. No.	X ¹	X ²	Z	R ⁴	R ⁵	physical data
200	F	F	O	C ₂ H ₅	H	
201	F	F	O	C ₂ H ₅	6-OC ₂ H ₅	
202	F	F	O	C ₂ H ₅	6-OCH ₂ CF ₃	
203	F	F	O	C ₂ H ₅	6-OnC ₃ H ₇	
204	F	F	O	CH ₃	H	
205	F	F	O	CH ₃	6-OC ₂ H ₅	
206	F	F	O	CH ₃	6-OCH ₂ CF ₃	
207	F	F	O	CH ₃	6-OnC ₃ H ₇	
208	F	F	O	CH ₃	5-Cl	
209	F	F	O	CH ₃	5-Br	
210	F	F	O	CH ₂ CF ₃	H	
211	F	F	O	CH ₂ CF ₃	6-OC ₂ H ₅	

Table 6

Oxazolines of the formula (I), G= 2-pyridon-4-yl




Ex. No.	X ¹	X ²	R ⁴	R ⁵	physical data
250	F	F	CH ₃	H	
251	F	F	C ₂ H ₅	H	
252	F	F	CH ₂ CF ₃	H	
253	F	F	nC ₃ H ₇	H	
254	F	F	iC ₃ H ₇	H	
255	F	F	nC ₄ H ₉	H	
256	F	F	CH ₂ CH(CH ₃) ₂	H	
257	F	F	CH ₂ C ₆ H ₅	H	
258	F	F	CH ₂ cC ₆ H ₁₁	H	
259	F	F	C ₂ H ₄ OCH ₃	H	
260	F	F	C ₂ H ₄ OC ₂ H ₅	H	
261	F	F	C ₂ H ₅	5-Cl	









Table 7

Oxazolines of the formula (I), G= 3-pyridinyl,


Ar= heteroaryl




Ex. No.	Ar	Z	R ⁵	R ⁶	physical data
300		O	OC ₂ H ₅	H	
301		O	OCH ₂ CF ₃	H	
302		O	OC ₂ H ₅	H	
303		O	OCH ₂ CF ₃	H	
304		O	OC ₂ H ₅	H	
305		O	OC ₂ H ₅	H	

306		O	OC ₂ H ₅	H
307		O	OCH ₂ CF ₃	H
308		O	OC ₂ H ₅	H
309		O	OCH ₂ CF ₃	H
310		O	OC ₂ H ₅	H
311		O	OCH ₂ CF ₃	H
312		O	OCH ₂ CF ₃	H
313		O	OC ₂ H ₅	H

51



O OC₂H₅ H




O OCH₂CF₃ H

¹⁹F NMR: -62.3, s, -74.3, tr

Table 8















Oxazolines of the formula I, G= 3-pyridinyl



R⁵ = heterocyclyl,
heterocycloxy,
heterocyclalkyl,
heterocyclalkoxy

Ex.No.	X ¹	X ²	R ⁶	R ⁵	physical data
350	F	F	H		
351	F	F	H		
352	F	F	H		
353	F	F	H		
354	F	F	H		
355	F	F	H		
356	F	F	H		
357	F	F	H		
358	F	F	H		
359	F	F	H		

53

				
360	F	F	H	
				
361	F	F	H	
				
362	F	F	H	
				
363	F	F	H	
				
364	F	F	H	
				
365	F	F	H	
				
366	F	F	H	
				
367	F	F	H	
				
369	F	F	H	
				
370	F	F	H	
				
371	F	F	H	
				
372	F	F	H	
				
373	F	F	H	
				
374	F	F	H	











375	F	F	H	
376	F	F	H	
377	F	F	H	
378	F	F	H	
379	F	F	H	
380	F	F	H	
381	F	F	H	
382	F	F	H	
383	F	F	H	

Table 9

Oxazolines of the formula (I), G = 3-pyridinyl



Ex. No.	X ¹	X ²	R ⁶	R ⁵	physical data
400	2-F	4-F	H	OCH ₂ CF ₃	¹⁹ F NMR: -75.1, tr, CF ₃ , -105.4, m, ArF
401	2-F	5-F	H	OCH ₂ CF ₃	¹⁹ F NMR: -74.3, CF ₃ ; -115.1, -118.7, ArF
402	2-F	H	H	OCH ₂ CF ₃	NMR, m.p. 101°C
403	2-CH ₃	H	H	OCH ₂ CF ₃	Oil, ¹ H-NMR (CDCl ₃ , TMS): δ = 2.67 (s, 3H, CH ₃); 4.24, 4.78, 5.48 (each dd, 1H, oxazoline), 4.81 (q, 2H, CH ₂ CF ₃), 6.93, 8.35 (each d, 1H, PyH), 7.84 (dd, 1H, PyH); 7.52, 7.42 (each m, 2H, C ₆ H ₄); 7.91 (d, 1H, MeC ₆ H ₄), 7.2 - 7.4 (m, 3H, MeC ₆ H ₄). ¹⁹ F-NMR (CDCl ₃ , CFCl ₃): δ = -74.3
404	2-CH ₃	H	H	OCH ₂ CF ₃	
405	2-CH ₃	H	H	OnC ₃ H ₇	
406	2-Et	H	H	OCH ₂ CF ₃	
407	2-CH ₃	4-CH ₃	H	OCH ₂ CF ₃	
408	2-F	4-CH ₃	H	OCH ₂ CF ₃	
409	2-F	4-Cl	H	OCH ₂ CF ₃	
410	2-SCH ₃	H	H	OCH ₂ CF ₃	
411	2-OC ₂ H ₅	H	H	OCH ₂ CF ₃	

412	2-OCH ₃	H	H	OCH ₂ CF ₃	Oil, ¹ H-NMR (CDCl ₃ , TMS): δ = 3.97 (s, 3H, OCH ₃); 4.30, 4.80, 5.50 (each dd, 1H, oxazoline); 4.84 (q, CH ₂ CF ₃); 6.95, 7.90, 8.37 (each dd, 1H, PyH); 7.46, 7.54 (each M, 2H, C ₆ H ₄); 7.0 - 7.08 (m, 2H, MeOC ₆ H ₄), 7.43 - 7.58 (m, 1H, MeOC ₆ H ₄), 7.87 (dd, 1H, MeOC ₆ H ₄). ¹⁹ F-NMR (CDCl ₃ , CFCl ₃): δ = -74.3
2H,					

B. Formulation examples

- 5 a) A dust is obtained by mixing 10 parts by weight of active substance and 90 parts by weight of talc as inert material and comminuting the mixture in a hammer mill.
- 10 b) A wettable powder which is readily dispersible in water is obtained by mixing 25 parts by weight of active substance, 65 parts by weight of kaolin-containing quartz as inert material, 10 parts by weight of potassium lignosulfonate and 1 part by weight of sodium oleoylmethyltaurinate as wetter and dispersant and grinding the mixture in a pinned-disk mill.
- 15 c) A dispersion concentrate which is readily dispersible in water is prepared by mixing 40 parts by weight of active substance with 7 parts by weight of a sulfosuccinic monoester, 2 parts by weight of a sodium lignosulfonate and 51 parts by weight of water and grinding the mixture in a ball mill to a fineness of below 5 microns.
- 20 d) An emulsifiable concentrate can be prepared from 15 parts by weight of active substance, 75 parts by weight of cyclohexanone as solvent and 10 parts by weight of oxyethylated nonylphenol (10 EO) as emulsifier.
- 25 e) Granules can be prepared from 2 to 15 parts by weight of active substance and an inert granule carrier material such as attapulgite, pumice granules and/or quartz sand. It is expedient to use a suspension of the wettable powder of Example b) with a solids content of 30%, which is sprayed onto the surface of attapulgite granules, and these are dried and mixed intimately. The wettable powder amounts to approx. 5% by weight and the inert carrier material to approx. 95% by weight of the finished granules.
- 30

C. Biological examples

Example 1: Effect on the spider mites *Tetranychus urticae*

- 5 Bean plants (*Phaseolus vulgaris*) which were severely infested with a complete population of spider mites (*Tetranychus urticae*) were sprayed to runoff point with an aqueous solution of the formulated preparation to be tested. The mortality of all mite stages was determined after 7 days. At a concentration of 300 ppm (based on active substance content), a 90-100% mortality was shown by the preparations of Examples 1, 2, 3, 4, 7, 8, 9, 14, 18, 21, 31, 40, 41, 43, 44, 47, 51, 106, 152 and 153.

Example 2: Effect on the aphid *Aphis fabae*

- 15 Field beans (*Vicia faba*) which were densely populated with the black bean aphid (*Aphis fabae*) were sprayed to runoff point with an aqueous solution of the formulated preparation to be tested. The mortality of the aphids was determined after 3 days. At a concentration of 300 ppm (based on the active substance content), a 90-100% mortality was shown by the preparations of Examples 1, 2, 3, 4, 7, 8, 152.

Example 3: Effect on the butterfly larva *Spodoptera littoralis*

- 25 Ten L4 larvae of the Egyptian cotton leafworm (*Spodoptera littoralis*) were introduced into a Petri dish whose bottom was covered with filter paper and contained approx. 5 ml of nutrient medium. Filter paper, nutrient paper and the larvae which had been introduced were then sprayed with an aqueous solution of the formulated preparation to be tested. The Petri dish was subsequently sealed with a lid. After 4 days' storage at approx. 23°C, the effect of the preparation on the larvae was determined. At a concentration of 300 ppm (based on the active substance content), a 90-100% larval mortality was caused by the preparations of Examples 3, 4, 22, 105, 106.

Example 4: Effect on the egg-larval stage of *Heliothis virescens*

- 35 A Petri dish whose bottom was covered with filter paper and which contained approx. 5 ml of nutrient medium was prepared. Filter paper sections containing approx. 30 24-hour-old eggs of the tobacco budworm (*Heliothis virescens*) were immersed for 5 seconds into an aqueous

solution of the formulated preparation to be tested and subsequently placed in the Petri dish. A further 200 μ l of the aqueous solution were distributed over the nutrient medium. After the Petri dish had been sealed, it was stored in a controlled-environment cabinet at approx. 25°C. After 6 days' storage, the mortality of the preparation on the eggs and any larvae hatched from them was determined. At a concentration of 300 ppm (based on the active substance content), a 90-100% mortality was caused by the preparations of Examples 1, 2, 3, 4, 5, 6, 7, 8, 14, 17, 21, 22, 23, 34, 35, 39, 40, 43, 44, 106.

Example 5: Effect on the corn budworm *Diabrotica undecimpunctata*

Corn seed was pregerminated for 6 hours while submerged in water, then placed into 10-ml glass tubes and covered with 2 ml of soil in each case. After 1 ml of water had been added, the plants remained in the glass tubes at 21°C until plant length was approx. 3 cm. 10 L2 larvae of the corn budworm (*Diabrotica undecimpunctata*) were then introduced into each tube on the soil. Two hours later, 1 ml of an aqueous solution of the formulated preparation to be tested were pipetted onto the soil surface in the tubes. After storage for 5 days under laboratory conditions at 21°C, the soil and root sections were examined for live *Diabrotica* larvae and the mortality was determined. At a concentration of 300 ppm (based on the active substance content), a 100% mortality of the test animals employed was shown by the preparations of Example 2, 3, 5, 9, 19, 39.

Example 6: Effect on the larvae of the copperbottle *Lucilia cuprina*

In a sample container, 20 recently hatched larvae of the copperbottle *Lucilia cuprina* were placed on a larval nutrient medium of ground mutton which contained the test substance in a concentration of 100 ppm. Larval growth on the nutrient medium is monitored over 72 hours up to larva 3. At a concentration of 100 ppm (based on the active substance content), a 100% mortality in the fly larvae employed was shown by the preparations of Examples 2, 3, 4, 5, 105, 106.

Example 7: Effect on developmental stages of the cat flea *Ctenocephalides felis*


5 The larval nutrient medium of the cat flea, composed of equal parts of blood meal and quartz sand, was admixed with the test substance at a concentration of 1000 ppm. Approx. 30 flea eggs freshly obtained from a breeding colony were transferred onto the medium. To assess the affect of the preparation, hatching of the flea larvae and their development into pupae and adult fleas was observed. A mortality of 100% was caused by
10 the test substances of Examples 2, 3 and 4.



Patent claims:


1. A 1,3-Oxazoline derivative and a 1,3-thiazoline derivative of the formula (I)

5



in which the symbols have the following meanings:

- 10 A is phenyl, pyridyl, pyrimidinyl, pyrazinyl, pyrazolyl or thienyl, each of the abovementioned groups optionally being substituted by one or more radicals X;
- 15 X is identical or different
- a) halogen, cyano, nitro;
- b) (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₂-C₄)alkynyl, (C₁-C₄)alkoxy, (C₁-C₄)alkylthio, (C₁-C₄)alkylsulfinyl, (C₃-C₈)cycloalkyl, (C₄-C₈)cycloalkenyl, phenyl,
- 20 the radicals of group b optionally being substituted by one or more radicals selected from the group consisting of halogen, cyano, (C₁-C₄)alkyl, (C₁-C₄)haloalkyl, (C₁-C₄)alkoxy and (C₁-C₄)haloalkoxy;
- E is a single bond, (C₁-C₄)alkylene, -O-CH₂- or -CH₂-O-;
- 25 G is a radical selected from the group consisting of:



Z is oxygen or sulfur;

5 R¹, R² and R³ are identical or different hydrogen, halogen, (C₁-C₄)haloalkyl, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)haloalkoxy or cyano;

10 R⁴ is hydrogen or a group (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkenyl, (C₄-C₁₀)cycloalkylalkyl or (C₇-C₁₂)phenylalkyl, each of the abovementioned groups optionally being substituted by one or more substituents selected from the group consisting of halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl, (C₁-C₄)alkoxy, (C₁-C₃)haloalkoxy, (C₁-C₃)alkylthio, (C₃-C₈)cyclo-
15 alkyl, (C₇-C₁₂)phenylalkoxy, (C₂-C₄)alkylcarbonyl, alkoxycarbonyl, (C₂-C₆)monoalkyl- and (C₃-C₉)dialkylaminocarbonyl, cyano and tri(C₁-C₄)alkylsilyl;

R⁵ and R⁶ are identical or different

20 a) hydrogen, halogen, cyano, formyl, (C₂-C₅)alkylcarbonyl, (C₂-C₈)alkoxycarbonyl, (C₃-C₉)monoalkyl- and dialkylamino-carbonyl,

25 b) (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, (C₄-C₈)cycloalkenyl, (C₁-C₈)alkoxy-alkyl, (C₁-C₈)alkylthio, (C₁-C₈)alkylsulfinyl, (C₁-C₈)alkylsulfonyl, NR⁷R⁸, phenyl, phenoxy, (C₇-C₁₂)phenylalkyl, (C₇-C₁₂)phenyl-alkoxy, heterocyclyl, heterocyclyloxy derivatives, hetero-cyclyl(C₁-C₄)alkoxy, (C₃-C₁₂)heterocyclylalkyl, each of the groups b optionally being substituted by one or more radicals selected from
30 the group consisting of halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl, (C₁-C₃)alkoxy, (C₁-C₃)haloalkoxy, (C₁-C₃)alkylthio, NR⁷R⁸,



alkylcarbonyl (C₂-C₅), alkoxycarbonyl (C₂-C₈), monoalkyl- and dialkylaminocarbonyl (C₃-C₉), cyano and tri(C₁-C₄)alkylsilyl;

- 5 c) (C₁-C₈)alkoxy, each of the groups c optionally being substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₃)alkylthio, NR⁷R⁸, alkylcarbonyl (C₂-C₅), alkoxycarbonyl (C₂-C₈), monoalkyl- and dialkylaminocarbonyl (C₃-C₉), cyano and tri(C₁-C₄)alkylsilyl;
- 10 d) R⁵ and R⁶ together form a ring which originates by linking two of the groups mentioned under b), preferably (C₃-C₅)alkylene, (C₂-C₄)alkyleneoxy, (C₁-C₃)alkylenedioxy, (C₂-C₄)alkyleneamino or alkylenethio(C₂-C₄);
- 15 R⁷ and R⁸ are identical or different
- a) hydrogen;
- b) (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, (C₇-C₁₂)phenylalkyl, (C₁-C₆)alkylcarbonyl, (C₁-C₆)alkylsulfonyl which are optionally substituted by one or more radicals selected from the group consisting of halogen,
- 20 (C₁-C₄)alkyl, (C₁-C₃)haloalkyl, (C₁-C₄)alkoxy, (C₁-C₃)haloalkoxy, (C₁-C₃)alkylthio, (C₁-C₃)alkylamino, (C₁-C₃)dialkylamino, (C₃-C₈)cycloalkyl, cyano and tri(C₁-C₄)alkylsilyl; or
- c) R⁷ and R⁸ together are alkylene (C₃-C₇), (C₃-C₇)oxa-, aza- or thiaalkylene, (C₃-C₇)alkylenecarbonyl or (C₃-C₇)alkylenesulfonyl.
- 25

2. A compound of the formula (I) as claimed in claim 1, where the symbols have the following meanings:

30 A is phenyl or pyridyl;

X is


- a) halogen, cyano, nitro or
- 35 b) (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₁-C₄)alkoxy, (C₁-C₄)alkylthio, (C₁-C₄)alkylsulfinyl, (C₃-C₇)cycloalkyl, (C₄-C₈)cycloalkenyl,



the radicals of group b optionally being substituted by one or more radicals selected from the group consisting of halogen, cyano, (C₁-C₄)alkoxy and (C₁-C₄)haloalkoxy;

5 E is a single bond or -CH₂-;

G is



10

Z is oxygen;

R¹, R², R³ are H, halogen, (C₁-C₄)haloalkyl, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)haloalkoxy or cyano;

15

R⁴ is H or (C₁-C₈)alkyl which is optionally substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl, (C₁-C₄)alkoxy, (C₁-C₃)haloalkoxy, (C₁-C₃)alkylthio, (C₃-C₈)cycloalkyl, cyano or tri(C₁-C₄)alkylsilyl;

20

R⁵, R⁶ are

- a) H, halogen, cyano;
- b) (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, (C₄-C₈)cycloalkenyl, (C₁-C₈)alkylthio, (C₁-C₈)alkylsulfinyl, (C₁-C₈)alkylsulfonyl, (C₇-C₁₂)phenylalkyl, (C₇-C₁₂)phenylalkoxy, (C₄-C₇)oxycycloalkyl or (C₄-C₇)oxacycloalkenyl,

25

the groups b optionally being substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl, (C₁-C₃)alkoxy, (C₁-C₃)haloalkoxy, (C₁-C₃)alkylthio and cyano; or

30


- c) (C₁-C₈)alkoxy which is optionally substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₃)alkylthio and cyano.




3. A compound of the formula (I) as claimed in claim 1 or 2, where the symbols have the following meanings:

- 5 A is phenyl;
- X is halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl, (C₁-C₄)alkoxy or (C₁-C₃)haloalkoxy;
- 10 E is a single bond;
- G is 3-pyridyl;
- Z is oxygen;
- 15 R¹, R², R³ are H, halogen, (C₁-C₄)haloalkyl, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)haloalkoxy;
- R⁴ is H or (C₁-C₈)alkyl which is optionally substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl, (C₁-C₄)alkoxy, (C₁-C₃)haloalkoxy, (C₁-C₃)alkylthio, (C₃-C₈)cycloalkyl, cyano or tri(C₁-C₄)alkylsilyl;
- 20 R⁵, R⁶ are
- a) H, halogen, cyano,
- 25 b) (C₁-C₈)alkyl, (C₁-C₈)alkoxy, (C₃-C₈)cycloalkyl, (C₄-C₈)cycloalkenyl, (C₁-C₈)alkylthio, (C₁-C₈)alkylsulfinyl, (C₁-C₈)alkylsulfonyl, (C₇-C₁₂)phenylalkyl or (C₇-C₁₂)phenylalkoxy,
- the groups b optionally being substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₄)alkyl, (C₁-C₃)haloalkyl and cyano.
- 30

4. A compound of the formula (I) as claimed in one or more of claims 1 to 3 selected from the groups consisting of (Ia) to (Ie):




(Ia)




(Ib)

5




(Ic)

10



(Id)



(Ie)



15

where the symbols have the meanings given in the formula (I) in claims 1 to 3.




5. A compound of the formula (I) as claimed in claim 4, selected from the group consisting of (Ia1) to (Ia4):


5



10




15



20 where the symbols have the meanings given in the formula (I) in claims 1 to 3.

6. A process for the preparation of compounds of the formula (I) as claimed in one or more of claims 1 to 5, which comprises reacting a
25 halogen and perfluoroalkylsulfonate compound of the formula (II)



in which

- 5 Y is Cl, Br, I or perfluoroalkylsulfonate and
 A, Z and E have the meanings given in the formula (I)

with an organometallic compound of the formula (III)



- 10 in which

M is a B-, Sn- or Zn-containing leaving group and

G has the meanings given in the formula (I) in claim 1, with palladium catalysis.

15

7. A crop protection composition which comprises at least one compound as claimed in any of claims 1 to 5 and at least one formulation auxiliary.

20

8. A fungicidal composition as claimed in claim 7 which comprises a fungicidally effective amount of at least one compound as claimed in any of claims 1 to 7 together with the additives or auxiliaries conventionally used for this application.

25

9. An insecticidal, acaricidal or nematocidal composition as claimed in claim 7 which comprises an effective amount of at least one compound as claimed in any of claims 1 to 5 together with the additives or auxiliaries conventionally used for this application.

30

10. A crop protection composition which comprises a fungicidally, insecticidally, acaricidally or nematocidally effective amount of at least one compound as claimed in any of claims 1 to 5 and at least one further active substance together with the auxiliaries and additives conventionally used for this application.

35

11. A composition for use in the protection of timber or as preservative in sealants, in paints, in cooling lubricants for metalworking or in drilling and



cutting oils, comprising an effective amount of at least one compound as claimed in any of claims 1 to 5 together with the auxiliaries and additives conventionally used for these applications.

- 5 12. A compound as claimed in any of claims 1 to 5 or a composition as claimed in claim 7, 9 or 10 for use as a veterinary medicament.
13. A process for the preparation of a composition as claimed in any of claims 7 to 11, which comprises combining the active substance and the
10 other additives and bringing the mixture into a suitable use form.
14. The use of a compound as claimed in any of claims 1 to 5 or of a composition as claimed in any of claims 7, 8, 10 and 11 as a fungicide.
- 15 15. The use of a compound as claimed in any of claims 1 to 5 or of a composition as claimed in any of claims 7, 8 and 11 as timber preservative or as preservative in sealants, in paints, in cooling lubricants for metalworking or in drilling and cutting oils.
- 20 16. The use of a compound as claimed in any of claims 1 to 5 or of a composition as claimed in any of claims 7, 9 or 10 for controlling harmful insects, Acarina, mollusks and nematodes.
- 25 17. A method of controlling phytopathogenic fungi, which comprises applying a fungicidally effective amount of a compound as claimed in any of claims 1 to 5 or of a composition as claimed in any of claims 7, 8, 10 and 11 to these fungi or to the plants, areas or substrates infested with them, or to seed.
- 30 18. A method of controlling harmful insects, Acarina, mollusks and nematodes, in which an effective amount of a compound as claimed in any of claims 1 to 5 or of a composition as claimed in any of claims 7, 9 or 10 is applied to these or to the plants, areas or substrates infested with them.
- 35 19. Seed, treated or coated with an effective amount of a compound as claimed in any of claims 1 to 5 or of a composition as claimed in any of claims 7, 9 or 10.