

MATERIALS AND METHODS FOR THE SYNTHESIS OF ERROR-MINIMIZED NUCLEIC ACID MOLECULES

This application claims the benefit of U.S. provisional application Serial No. 61/593,813, filed February 1, 2012, which is hereby incorporated by reference in its entirety, including all Tables, Figures, and Claims.

FIELD OF THE INVENTION

[0001] The present invention relates generally to molecular biology and genetics, and to the synthesis of genes and other nucleic acid molecules.

INCORPORATION OF SEQUENCE LISTING

[0002] The material in the accompanying Sequence Listing is hereby incorporated by reference into this application. The accompanying sequence listing text file, name SGI-XXX.XXPCT_Sequence Listing, was created on DATE and is XX KB. The file can be assessed using Microsoft Word on a computer that uses Windows OS.

BACKGROUND

[0003] In modern molecular biology and genetic engineering, many molecular techniques that involve the use of molecules of nucleic acid often require the generation of a supply of nucleic acid molecules by synthetic methods. For example, to test hypotheses in the field of metabolic engineering or genomics, and to synthesize designed proteins and organisms with tailored genomes, cost-effective methods for synthesizing nucleic acid molecules with a high degree of fidelity to an intended nucleotide sequence are often required. Common methods of nucleic acid synthesis, *e.g.* synthesis of double-stranded DNA, include polymerase chain reaction methods and ligation chain reaction methods. Often, ensuring that a synthetic DNA molecule contains the correct nucleotide sequence is important, if not essential, for the success of the molecular technique in which the synthesized DNA is to be used. For example, the synthesis of a DNA coding sequence for use in gene expression of functional polypeptides requires a precise DNA sequence;

because even one nucleotide substitution, insertion or deletion can have significant consequences for the polypeptide that is ultimately produced. Thus, the process of minimizing DNA molecules having incorrect DNA sequences from a synthetic DNA population is widely considered to be essential in providing error-free synthetic DNA produced by a *de novo* gene synthesis method.

[0004] Recently, efforts to synthesize nucleic acid molecules accurately while controlling costs have yielded methods including microchip-based gene synthesis and PCR-based gene assembly technologies. While these conventional technologies provide the capability to synthesize multiple genes, reducing errors introduced into the desired gene-sequence remains challenging. To avoid the problems with sequence errors inherent in gene synthesis, some have focused on purifying the oligonucleotides that are used at the early stages of the synthesis process. However, these oligonucleotide purification approaches are costly, and sequence errors persist and propagate through the subsequent steps of the synthesis process.

[0005] Thus, there exists a need for alternative methods for reducing sequence errors within a population of DNA molecules. What is desired is a way to synthesize genes and other nucleic acid molecules with a greater yield of molecules having a desired nucleotide sequence. An approach that can correct sequence errors at a much later step in the synthesis process makes the desired increase in nucleotide sequence accuracy possible, while allowing the process to be cost-effective.

SUMMARY

[0006] The present invention provides methods and materials for error correction in the replication and amplification of nucleic acid molecules. In one embodiment of the invention a first plurality of double-stranded nucleic acid molecules having a nucleotide mismatch are fragmented by exposure to a unidirectional mismatch endonuclease. The nucleic acid molecules are cut at or near the mismatch site with an endonuclease, leaving a double-stranded nucleic acid molecule having a mismatch at or near the end of the molecule. In one embodiment the nucleic acid molecule is then exposed to an exonuclease having a unidirectional activity in the 5' to 3' or 3' to 5' direction, which therefore removes the mismatched nucleotide. A second plurality of double-stranded nucleic acid molecules is

assembled from the nucleic acids with the mismatched nucleotides removed. The missing nucleotides can then be filled in by the action of, *e.g.*, a DNA polymerase either directly or at a subsequent amplification step, and these steps can be repeated as many times as necessary. The result is double-stranded nucleic acid molecules with a decreased frequency of nucleotide mismatches versus the first plurality of nucleic acid molecules.

[0007] Thus, in one aspect the present invention provides a method for error correction of nucleic acid molecules. The method involves (a) obtaining a first plurality of double-stranded nucleic acid molecules having at least one nucleotide mismatch; (b) fragmenting the plurality of double-stranded nucleic acid molecules having a mismatch by reacting the nucleic acid molecules having a mismatch with at least one molecule having a unidirectional mismatch endonuclease activity; (c) removing the nucleotide mismatch by reacting the fragmented double-stranded nucleic acid molecules having a mismatch of (b) with at least one molecule having unidirectional exonuclease activity of the same directionality as the unidirectional mismatch endonuclease activity of (b) to provide a fragmented error-free double-stranded nucleic acid molecule; and (d) assembling a second plurality of double-stranded nucleic acid molecules having the fragmented error-free double-stranded nucleic acid molecule of (c). The second plurality of double-stranded nucleic acid molecules has a decreased frequency of nucleotide mismatches as compared to the first plurality of double-stranded nucleic acid molecules.

[0008] In one embodiment, the first plurality of nucleotide acid molecules can contain one or more synthetic nucleotide sequences. The first plurality of nucleotide acid molecules can contain a mixture of one or more naturally occurring gene sequences and one or more synthetic nucleotide sequences. The first plurality of nucleic acid molecules can be obtained by synthesizing the nucleic acid molecules in one embodiment, or by assembling the nucleic acid molecules from subsets and/or oligonucleotides in another embodiment.

[0009] In one embodiment of the method, steps (b) and (c) recited above are performed as separate reactions, but in another embodiment steps (b) and (c) are performed as a simultaneous or one-step reaction. In one embodiment of the method, the unidirectional mismatch endonuclease activity cuts 5' to the mismatch and the unidirectional exonuclease activity removes the nucleotide mismatch from the 5' end of the fragmented nucleic acid molecule. But in another embodiment the unidirectional mismatch

endonuclease activity cuts 3' to the mismatch and the unidirectional exonuclease activity removes the nucleotide mismatch from the 3' end of the fragmented nucleic acid molecule. Examples of the molecule having unidirectional mismatch endonuclease activity include, but are not limited to, RES I, CEL I, CEL II, an SP endonuclease, SP I endonuclease, T7 endonuclease, T4 endonuclease, endonuclease V, a Mut protein, a variant of any thereof, and a combination of any two or more thereof. In a preferred embodiment, CEL I, CEL II, or a combination of CEL I and CEL II is utilized. In another preferred embodiment, the molecule having a unidirectional mismatch endonuclease activity is encoded by a nucleic acid molecule comprising a nucleotide sequence which hybridizes under low, moderate, or high stringency conditions to a nucleic acid sequence selected from the group consisting of a) a nucleic acid sequence hybridizing under low, moderate, or high stringency conditions to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 01, SEQ ID NO: 03, SEQ ID NO: 05, SEQ ID NO: 07, SEQ ID NO: 09, SEQ ID NO: 12, SEQ ID NO: 15, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, a complement of any, and a fragment of any; b) a nucleic acid sequence exhibiting 70% or greater identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 01, SEQ ID NO: 03, SEQ ID NO: 05, SEQ ID NO: 07, SEQ ID NO: 09, SEQ ID NO: 12, SEQ ID NO: 15, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, a complement of any, and a fragment of any; and c) a nucleic acid sequence encoding a polypeptide exhibiting 60% or greater identity to an amino acid sequence selected from the group consisting of SEQ ID NO: 02, SEQ ID NO: 04, SEQ ID NO: 06, SEQ ID NO: 08, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 28, and SEQ ID NO: 29.

[0010] Examples of the molecule having unidirectional exonuclease activity include, but are not limited to, exonuclease III, a DNA polymerase, lambda exonuclease, T7 exonuclease, and T5 exonuclease, and variants thereof. In one embodiment the molecule having unidirectional exonuclease activity is a DNA polymerase with proofreading activity (e.g., 3' exonuclease proofreading activity). Examples of polymerases with proofreading

activity include, but are not limited to, T4 polymerase, T7 polymerase, and phi29 polymerase.

[0011] In a specific embodiment of the methods of the present invention, the at least one molecule having unidirectional mismatch endonuclease activity is selected from: CEL I, CEL II, variants of any thereof, and a combination of any two or more thereof; and the at least one molecule having unidirectional exonuclease activity selected from the group consisting of exonuclease III, a variant thereof, and a combination of any two or more thereof.

[0012] In one aspect of the invention, the present disclosure provides isolated nucleic acid molecules comprising nucleic acid sequences hybridizing under low, moderate, or high stringency conditions: a) a nucleic acid sequence hybridizing under low, moderate, or high stringency conditions to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 09, SEQ ID NO: 12, SEQ ID NO: 15, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, a complement thereof or a fragment of either; or b) a nucleic acid sequence exhibiting 70% or greater identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 09, SEQ ID NO: 12, SEQ ID NO: 15, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, a complement thereof or a fragment of either; or c) a nucleic acid sequence encoding a polypeptide exhibiting 50% or greater identity to an amino acid sequence selected from the group consisting of SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 28, and SEQ ID NO: 29.

[0013] In another aspect of the invention, the present disclosure provides recombinant nucleic acid constructs, such as recombinant nucleic acid vectors, which include a nucleic acid molecule of the invention as recited herein that is operably linked to a heterologous nucleic acid. In some embodiments the heterologous nucleic acid is a heterologous transcription control element. In some preferred embodiments, any of the above recombinant nucleic acid constructs can comprise a heterologous nucleic acid encoding a polypeptide sequence. The polypeptide sequence may include a secretion signal

or an epitope tag. In particular embodiments the nucleic acid constructs can comprise SEQ ID NO: 31 or SEQ ID NO: 33, or complements or variants thereof or comprise sequences that hybridize under low, medium, or high stringency conditions to either of SEQ ID NO: 31 or 33 or their complements or variants thereof.

[0014] In yet another aspect of the invention, the invention provides a recombinant host cell that includes a nucleic acid construct of the invention as disclosed herein. The recombinant host cell can be an insect cell, a mammalian cell, a microbial cell, or a plant cell. In some other embodiments, the disclosure also provides biological samples, biomass, and progeny derived from a host organism as described above. In yet other embodiments, the disclosure further provides biomaterials derived from a host organism as described above.

[0015] In another aspect of the present invention, the invention further provides isolated polypeptides. In some embodiments, such isolated polypeptides are expressed by a nucleic acid molecule of the invention as disclosed herein. The nucleic acid molecule expressing the polypeptides can be introduced into a host cell. In some embodiments the amino acid sequence of the polypeptide can comprise an amino acid sequence selected from the group consisting of SEQ ID NO: 11, amino acid residues 1 to 297 of SEQ ID NO: 11, amino acid residues 22 to 308 of SEQ ID NO: 11, SEQ ID NO: 17, amino acid residues 1 to 320 of SEQ ID NO: 17, and amino acid residues 22 to 331 of SEQ ID NO: 17.

[0016] In another aspect, the present invention discloses compositions comprising: (i) a molecule having a unidirectional mismatch endonuclease activity; and (ii) a molecule having unidirectional exonuclease activity of the same directionality as the unidirectional mismatch endonuclease activity in (i). In various embodiments the molecule of (i) is selected from the group consisting of RES I, CEL I, CEL II, T7 endonuclease, T4 endonuclease, endonuclease V, a Mut protein, a variant of any thereof, and a combination of any two or more thereof; and the molecule of (ii) is selected from the group consisting of exonuclease III, a DNA polymerase, a variant of any thereof, and a combination of any two or more thereof.

[0017] In yet another aspect, the present disclosure further provides a kit comprising (i) a molecule having a unidirectional mismatch endonuclease activity; and (ii) a molecule having unidirectional exonuclease activity of the same directionality as the unidirectional

mismatch endonuclease activity in (i). In other embodiments the kit can also have instructions for conducting a method for error correction as described herein and/or provide a link to a website that provides information on a method of error correction as described herein.

[0018] These and other objects, aspects, and features of the invention will become more fully apparent to those of ordinary skill in the art upon review of the following detailed description of the invention and the claims in conjunction with the accompanying figures.

BRIEF DESCRIPTION OF THE DRAWINGS

[0019] FIGURE 1 provides schematic illustration of an embodiment of the methods of the present invention.

[0020] FIGURE 2 provides a schematic illustration of steps of one embodiment of the present invention.

[0021] FIGURE 3 provides a flow chart illustrating steps taken in one embodiment of the invention.

[0022] FIGURE 4 is an alignment of a *Selaginella lepidophylla* CEL I endonuclease (SEQ ID NO: 02), a celery CEL I endonuclease (SEQ ID NO: 04), an *Apium* sp. CEL II endonuclease (SEQ ID NO: 06), another *Apium* sp. CEL II endonuclease (SEQ ID NO: 08), *Mimulus guttatus* CEL I endonuclease (SEQ ID NO: 10), a *Solanum tuberosum* CEL I endonuclease (SEQ ID NO: 13), a *Vitis vinifera* CEL II endonuclease (SEQ ID NO: 16), a *Solanum tuberosum* CEL II endonuclease (SEQ ID NO: 25), a *Medicago* sp. CEL II endonuclease (SEQ ID NO: 27). The sequence alignment of FIGURE 4 was generated using the program AlignX of the Vector NTI Advance™ 11.5 package (Invitrogen, Carlsbad, Calif.) with default settings. As discussed in detail elsewhere herein, several polypeptide domains and motifs with high degree of conservation have been identified from this sequence comparison analysis. In the alignment figure shown herein, a dash in an aligned sequence represents a gap, *i.e.*, a lack of an amino acid at that position. Black boxes and gray boxes identify identical amino acids and conserved amino acids, respectively, among aligned sequences.

[0023] FIGURE 5 depicts SDS polyacrylamide gel analysis of purified MimmulusC-His CEL I protein (FIG. 5A) and Western Blot results using anti-polyHistidine antibody (FIG. 5B). Lane 1: Fermentas Marker (5 μ L); Lane 2: MimmulusC-His Pre-Dialysis (12 μ L); Lane 4: Fermentas Marker (12 μ L); Lane 5: MimmulusC-His Post-Dialysis (12 μ L); Lane 7: Fermentas Marker (5 μ L); Lane 8: MimmulusC-His Post-Dialysis (6 μ L).

DETAILED DESCRIPTION OF THE INVENTION

[0024] The present application relates to compositions, methods and related materials useful for the production of error-minimized nucleic acid molecules.

[0025] In one aspect, the present disclosure provides materials and methods that can be used to reduce mismatch errors in a population of nucleic acid molecules. For example, nucleic acid molecules that encode mismatch endonucleases are disclosed as well as methods for using such nucleic acid molecules and polypeptides encoded thereby to reduce nucleotide mismatches in a nucleic acid population. The disclosure also provides recombinant nucleic acid molecules, and recombinant cells as well as recombinant organisms comprising such nucleic acid molecules and methods for using the same.

[0026] The singular form "a", "an", and "the" include plural references unless the context clearly dictates otherwise. For example, the term "a cell" includes one or more cells, including mixtures thereof.

[0027] Domain: "Domains" are groups of substantially contiguous amino acids in a polypeptide that can be used to characterize protein families and/or parts of proteins. Such domains typically have a "fingerprint", "motif", or "signature" that can comprise conserved primary sequence, secondary structure, and/or three-dimensional conformation. Generally, domains are correlated with specific *in vitro* and/or *in vivo* activities. A domain can have a length of from 4 amino acids to 400 amino acids, *e.g.*, 4 to 50 amino acids, or 4 to 20 amino acids, or 4 to 10 amino acids, or 4 to 8 amino acids, or 25 to 100 amino acids, or 35 to 65 amino acids, or 35 to 55 amino acids, or 45 to 60 amino acids, or 200 to 300 amino acids, or 300 to 400 amino acids.

[0028] Expression: As used herein, "expression" refers to the process of converting genetic information of a polynucleotide into RNA through transcription, which is typically

catalyzed by an enzyme, RNA polymerase, and into protein, through translation of mRNA on ribosomes.

[0029] The term "epitope", "tag", "tag sequence", or "protein tag" as used herein refers to a chemical moiety, either a nucleotide, oligonucleotide, polynucleotide or an amino acid, peptide or protein or other chemical, that when added to another sequence, provides additional utility or confers useful properties, particularly in the detection or isolation, to that sequence. Thus, for example, a homopolymer nucleic acid sequence or a nucleic acid sequence complementary to a capture oligonucleotide may be added to a primer or probe sequence to facilitate the subsequent isolation of an extension product or hybridized product. In the case of protein tags, histidine residues (e.g., 4 to 8 consecutive histidine residues) may be added to either the amino- or carboxy-terminus of a protein to facilitate protein isolation by chelating metal chromatography. Alternatively, amino acid sequences, peptides, proteins or fusion partners representing epitopes or binding determinants reactive with specific antibody molecules or other molecules (e.g., FLAG epitope, c-myc epitope, transmembrane epitope of the influenza A virus hemagglutinin protein, protein A, cellulose binding domain, calmodulin binding protein, maltose binding protein, chitin binding domain, glutathione S-transferase, and the like) may be added to proteins to facilitate protein isolation by procedures such as affinity or immunoaffinity chromatography. Chemical tag moieties include such molecules as biotin, which may be added to either nucleic acids or proteins and facilitates isolation or detection by interaction with avidin reagents, and the like. Numerous other tag moieties are known to, and can be envisioned by, the trained artisan, and are contemplated to be within the scope of this definition.

The polynucleotides of the invention and polypeptides encoded thereby

[0030] In one aspect of the present invention, the disclosure provides novel isolated nucleic acid molecules, nucleic acid molecules that hybridize to these nucleic acid molecules (e.g., complements), and nucleic acid molecules that encode the same protein due to the degeneracy of the DNA code. Additional embodiments of the present application further include the polypeptides encoded by the nucleic acid molecules of the present invention.

[0031] The polynucleotides and polypeptides of the present invention disclosed in the sequence listing or otherwise disclosed herein (and their fragments or variants) are

"biologically active" with respect to either a structural attribute, such as the capacity of a nucleic acid to hybridize to another nucleic acid molecule, or the ability of a polypeptide to be bound by an antibody (or to compete with another molecule for such binding). Alternatively, such an attribute may be catalytic and thus involve the capacity of the molecule to mediate a chemical reaction or response.

[0032] In some embodiments the polynucleotides and polypeptides of the present invention are recombinant. A recombinant polynucleotide or polypeptide is one derived from human manipulation of the polynucleotide or polypeptide and an organism using laboratory methods resulting in nucleic acid sequences (or polypeptides) that would not otherwise be found in (or produced by) the manipulated organism.

[0033] Nucleic acid molecules or fragments thereof of the present invention are capable of specifically hybridizing to other nucleic acid molecules under certain circumstances. "Specifically hybridize" refers to a process whereby complementary nucleic acid strands anneal to each other under appropriately stringent conditions. Nucleic acid molecules are said to exhibit "complete complementarity" if every nucleotide of one of the molecules is complementary to a nucleotide of the other and the nucleotide pairs form Watson-Crick base pairs. Two nucleic acid molecules are said to be "minimally complementary" if they can anneal to one another with sufficient stability to remain annealed under at least conventional "low-stringency" conditions. Similarly, the molecules are said to be "complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under conventional "high-stringency" conditions. Conventional stringency conditions are described by Sambrook *et al.* in *Molecular Cloning, A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Press, Cold Spring Harbor, NY (1989), and by Haymes *et al.* In: *Nucleic Acid Hybridization, A Practical Approach*, IRL Press, Washington, D.C. (1985). Departures from complete complementarity are therefore permissible, as long as such departures do not completely preclude the capacity of the molecules to form a double-stranded structure. Thus, in order for a nucleic acid molecule or fragment thereof of the present invention to serve as a primer or probe it needs only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed.

[0034] Appropriate stringency conditions which promote DNA hybridization include, for example, 6.0× sodium chloride/sodium citrate (SSC) at about 45°C, followed by a wash of 2.0×SSC at about 50°C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22°C, to high stringency conditions at about 65°C. Both temperature and salt may be varied, or either the temperature or the salt concentration may be held constant while the other variable is changed. These conditions are known to those skilled in the art, or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1- 6.3.6. For example, low stringency conditions may be used to select nucleic acid sequences with lower sequence identities to a target nucleic acid sequence. One may wish to employ conditions such as about 0.15 M to about 0.9 M sodium chloride, at temperatures ranging from about 20°C to about 55°C. High stringency conditions may be used to select for nucleic acid sequences with higher degrees of identity to the disclosed nucleic acid sequences (Sambrook *et al.*, 1989, *supra*). High stringency conditions typically involve nucleic acid hybridization in about 2×SSC to about 10×SSC (diluted from a 20×SSC stock solution containing 3 M sodium chloride and 0.3 M sodium citrate, pH 7.0 in distilled water), about 2.5× to about 5× Denhardt's solution (diluted from a 50× stock solution containing 1% (w/v) bovine serum albumin, 1% (w/v) ficoll, and 1% (w/v) polyvinylpyrrolidone in distilled water), about 10 mg/mL to about 100 mg/mL fish sperm DNA, and about 0.02% (w/v) to about 0.1% (w/v) SDS, with an incubation at about 50°C to about 70°C for several hours to overnight. High stringency conditions are preferably provided by 6×SSC, 5× Denhardt's solution, 100 mg/mL fish sperm DNA, and 0.1% (w/v) SDS, with incubation at 55°C for several hours. Hybridization is generally followed by several wash steps. The wash compositions generally comprise 0.5×SSC to about 10×SSC, and 0.01% (w/v) to about 0.5% (w/v) SDS with a 15-min incubation at about 20°C to about 70°C. Preferably, the nucleic acid segments remain hybridized after washing at least one time in 0.1×SSC at 65°C.

[0035] In one embodiment, a subset of the nucleic acid molecules of this invention includes fragments of the disclosed polynucleotides consisting of oligonucleotides of at least 12, at least 15, at least 16, at least 17, at least 18, at least 19, and at least 20 consecutive nucleotides of the disclosed polynucleotide. Such oligonucleotides are

fragments of the larger polynucleotide molecules disclosed in the sequence listings or otherwise described herein and find use, for example, as interfering molecules, probes and primers for detection of the polynucleotides of the present invention.

[0036] Nucleic acid molecules of the invention can include a sequence sufficient to encode a biologically active fragment of a domain of a mismatch endonuclease, an entire mismatch endonuclease, or several domains within an open reading frame encoding a mismatch endonuclease.

[0037] In another embodiment, the present disclosure specifically provides nucleotide sequences comprising regions that encode polypeptides. The encoded polypeptides may be the complete polypeptide encoded by the gene represented by the protein or polynucleotide, or may be fragments of the encoded protein. Preferably, polynucleotides provided herein encode polypeptides constituting a substantial portion of the complete protein, and more preferentially, constituting a sufficient portion of the complete protein to provide the relevant biological activity, *e.g.*, mismatch endonuclease activity.

[0038] Of particular interest are polynucleotides of the present invention that encode a mismatch endonuclease. Such polynucleotides may be expressed in recombinant cells or recombinant organisms to produce molecules having mismatch endonuclease activity. In some embodiments, nucleic acid molecules that are fragments of these mismatch endonuclease-encoding nucleotide sequences are also encompassed by the present invention. A "mismatch endonuclease fragment", as used herein, is intended to be a fragment of a nucleotide sequence that encodes a mismatch endonuclease. A fragment of a nucleotide sequence may encode a biologically active portion of a mismatch endonuclease, or it may be a fragment that can be used as a hybridization probe or PCR primer using methods disclosed herein. Fragments of nucleic acid molecules or polypeptides comprise at least 10, 25, 50, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1050, 1100, 1150, 1200, 1250, 1300, 1350, 1400, 1450, 1500, 1550, 1600, 1650, 1700, 1750, 1800, 1850, 1900, 1950, 2000, 2050, 2100, 2150, 2200, 2250, 2300, 2350, 2400, 2450, 2500, 2550, 2600, 2650, 2700, 2750, 2800, 2850, 2900, 2950, 3000, 3050, 3100, 3150, 3200, 3250, 3300, 3350 contiguous nucleotides or amino acids, or up to the number of nucleotides or amino acids present in a full-length nucleotide sequence or polypeptide sequence disclosed herein.

Fragments of the nucleotide sequences of the present invention include those that encode protein fragments that retain the biological activity of a mismatch endonuclease. By "retains activity" is intended that the fragment will have at least 30%, at least 50%, at least 70%, at least 80%, at least 90%, or at least 95% of the endonuclease activity of the full-length mismatch endonuclease protein. Methods for measuring endonuclease, including mismatch endonuclease activity are well known in the art. See, for example, U.S. Pat. No. 6,391,557; U.S. Pat. No. 7,129,075. Mismatch endonuclease activity refers to an activity of sufficient level to perform the step of fragmenting the dsDNA molecules (or removing the nucleotide mismatch) in the method within a convenient time period for conducting the assay. In different embodiments the activity is sufficient to perform the fragmenting or removal within 2 hours or within 4 hours or within 6 hours, or within 10 hours or within 12 hours or within 24 hours.

[0039] In different embodiments a fragment of a mismatch endonuclease-encoding nucleotide sequence that encodes a biologically active portion of a polypeptide of the invention will encode at least 15, 25, 30, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350 contiguous amino acids, or up to the total number of amino acids present in a full-length mismatch endonuclease protein disclosed in the sequence listing or otherwise disclosed herein. For example, a mismatch endonuclease fragment in accordance with the present invention may have an N-terminal or a C-terminal truncation of at least 20 amino acids, at least 50, at least 75, at least 90, at least 100, or at least 150 amino acids relative to any one of the mismatch endonuclease amino acid sequences disclosed in the sequence listing or otherwise disclosed herein.

[0040] Also of interest in the present invention are variants of the polynucleotides disclosed in the sequence listing or otherwise disclosed herein. Such variants may be naturally-occurring, including homologous polynucleotides from the same or a different species, or may be non-natural variants, for example polynucleotides synthesized using chemical synthesis methods, or generated using recombinant DNA techniques. Variants can be generated having modified nucleic acid molecules in which nucleotides have been inserted, deleted, and/or substituted, and such modifications can provide a desired effect on the endonuclease biological activity as described herein. Degeneracy of the genetic code provides the possibility to substitute at least one base of the protein encoding sequence of a

gene with a different base without causing the amino acid sequence of the polypeptide produced from the gene to be changed. Hence, the nucleic acid molecules of the present invention may also have any base sequence that has been changed from any one of the polynucleotide sequences disclosed herein by substitution in accordance with degeneracy of the genetic code.

[0041] The skilled artisan will further appreciate that changes can be introduced by mutation of the nucleotide sequences of the invention, thereby leading to changes in the amino acid sequence of the encoded endonuclease proteins, without altering the biological activity of the proteins. Thus, variant isolated nucleic acid molecules can be created by introducing one or more nucleotide substitutions, additions, or deletions into the corresponding nucleotide sequence disclosed herein, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Such variant nucleotide sequences are also encompassed by the present invention.

[0042] For example, conservative amino acid substitutions may be made at one or more predicted nonessential amino acid residues. A "nonessential" amino acid residue, as used herein, is a residue that can be altered from the wild-type sequence of a mismatch endonuclease protein without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been well defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine).

[0043] As discussed above, it will be appreciated by one skilled in the art that amino acid substitutions may be made in non-conserved regions that retain function. In general,

such substitutions would not be made for conserved amino acid residues, or for amino acid residues residing within a conserved motif, where such residues are essential for protein activity. Conserved residues, domains and motifs of mismatch endonuclease sequences are reported in the art. Examples of residues that are conserved and that may be essential for protein activity include, for example, residues that are identical between all proteins contained in an alignment of the amino acid sequences of the present invention and known mismatch endonuclease sequences. Examples of residues that are conserved but that may allow conservative amino acid substitutions and still retain activity include, for example, residues that have only conservative substitutions between all proteins contained in an alignment of the amino acid sequences of the present invention and known mismatch endonuclease sequences. However, one of skill in the art would understand that functional variants may have minor conserved or non-conserved alterations in the conserved residues.

[0044] In some embodiments of the present invention, such mismatch endonuclease variants include proteins having an amino acid sequence that differs from any one of the polypeptides disclosed herein, by an amino acid deletion, insertion, or substitution at one or more of the positions corresponding to the conserved amino acid residues as identified in Figure 4. In some preferred embodiments, such mismatch endonuclease variants include proteins having an amino acid sequence that differs from the polypeptide sequence of SEQ ID NO: 11 or SEQ ID NO: 17 or a fragment of either, by an amino acid deletion, insertion, or substitution at one or more of the positions corresponding to the conserved amino acid residues as identified in Figure 4, and combinations of any thereof.

[0045] Alternatively, variant nucleotide sequences can be made by introducing mutations randomly along all or part of the coding sequence, such as by saturation mutagenesis, and the resultant mutants can subsequently be screened for ability to confer mismatch endonuclease activity in order to identify mutants that retain mismatch endonuclease activity. For example, following mutagenesis, the encoded protein can be expressed recombinantly, and the activity of the protein can be determined using standard assay techniques. Methods for assaying endonuclease activity and particularly mismatch endonuclease activity are well known in the art. See, for example, U.S. Pat. No. 6,391,557; U.S. Pat. No. 7,129,075.

[0046] In addition, using sequence-based methods such as PCR, hybridization, and the like corresponding mismatch endonuclease sequences can be identified, such sequences having substantial identity to the sequences of the invention. See, for example, Sambrook and Russell (2001, *supra*.)

[0047] Polynucleotides and polypeptides that are variants of the polynucleotides and polypeptides provided herein will generally demonstrate significant identity with the polynucleotides and polypeptides provided herein. Of particular interest are polynucleotide and polypeptide homologs having at least about 50% sequence identity, preferably at least about 60%, preferably at least about 70%, more preferably at least about 75%, more preferably at least about 80%, more preferably at least about 85%, more preferably at least about 90%, even more preferably at least about 95%, and most preferably at least about 96%, 97%, 98% or 99% sequence identity with any one of the polynucleotide or polypeptide sequences described in the sequence listing or otherwise described herein. For example, the invention provides polynucleotide homologs having the recited percent sequence identities to polynucleotides of any of SEQ ID NOs: 1, 3, 5, 7, 9, 12, 15, 18, 20, 22, 24, 26, 29, 30, and 32, as well as to constructs SEQ ID NOs: 31 and 33. The invention also provides polypeptides that are encoded by any of the polynucleotides disclosed herein. The invention also provides polypeptide variants having the recited percent sequence identities to polypeptides of any of SEQ ID NO: 2, 4, 6, 8, 10, 11, 13, 14, 16, 17, 19, 21, 23, 25, 27, 28, and 29. The invention also provides fragments of the polynucleotides and polypeptides disclosed herein.

[0048] "Sequence identity" refers to the extent to which two optimally aligned polynucleotide or peptide sequences are invariant throughout a window of alignment of components, *e.g.*, nucleotides or amino acids. An "identity fraction" for aligned segments of a test sequence and a reference sequence is the number of identical components which are shared by the two aligned sequences divided by the total number of components in reference sequence segment, *i.e.*, the entire reference sequence or a smaller defined part of the reference sequence.

[0049] "Percentage of sequence identity" or "percent sequence identity", as used herein with reference to polynucleotides, refers to the percentage of identical nucleotides or amino acids in a linear polynucleotide sequence of a reference ("query") polynucleotide

molecule (or its complementary strand) as compared to a test ("subject") polynucleotide molecule (or its complementary strand) when the two sequences are optimally aligned. The terms are used in the same way with reference to polypeptide sequences and their corresponding amino acid residues. As is known in the art, when calculating percentage sequence identity for the polynucleotide and/or polypeptide sequences described herein, any leader sequences or sequence tags or other such sequences included for a purpose such as, for example, ease of purification, expression, secretion, etc. are not included in the sequence for the purpose of such calculation.

[0050] Percent sequence identity is determined by comparing two optimally locally aligned sequences over a comparison window defined by the length of the local alignment between the two sequences. The polynucleotide sequences in the comparison window may comprise additions or deletions (*e.g.*, gaps or overhangs) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Local alignment between two sequences only includes segments of each sequence that are deemed to be sufficiently similar according to a criterion that depends on the algorithm used to perform the alignment (*e.g.* BLAST). The percentage identity is calculated by determining the number of positions at which the identical nucleic acid base (or polypeptide amino acid) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100. Optimal alignment of sequences for aligning a comparison window are well known to those skilled in the art and any may be used in the invention such as the local homology algorithm of Smith and Waterman (*Add. APL. Math.* 2:482, 1981), by the global homology alignment algorithm of Needleman and Wunsch (*J Mol. Biol.* 48:443, 1970), by the search for similarity method of Pearson and Lipman (*Proc. Natl. Acad. Sci. (USA)* 85: 2444, 1988), by heuristic implementations of these algorithms such as, GAP, BESTFIT, FASTA, and TFASTA available as part of the GCG™ Wisconsin Package™ (Genetics Computer Group, Accelrys Inc., Burlington, Mass.), by heuristic implementations of these algorithms such as NCBI BLAST, WU-BLAST, BLAT, SIM, BLASTZ, or by manual inspection. As described above, an "identity fraction" for aligned segments of a test sequence and a reference sequence is the number of identical components which are shared by the two aligned sequences divided by the total

number of components in the reference sequence segment, *i.e.*, the entire reference sequence or a smaller defined part of the reference sequence. Percent sequence identity is represented as the identity fraction multiplied by 100. The comparison of one or more polynucleotide sequences may be to a full-length polynucleotide sequence or a portion thereof, or to a longer polynucleotide sequence. For purposes of this invention "percent identity" may also be determined using BLASTX version 2.0 for translated nucleotide sequences and BLASTN version 2.0 for polynucleotide sequences.

[0051] For purposes of this invention, "percent identity" may also be determined using BLASTX version 2.0 for translated nucleotide sequences and BLASTN version 2.0 for polynucleotide sequences (or BLASTp for polypeptide sequences). In a preferred embodiment of the present invention, the presently disclosed gene regulatory sequences comprise protein, peptide, nucleic acid molecules or fragments having a BLAST score of more than 200, preferably a BLAST score of more than 300, and even more preferably a BLAST score of more than 400 with their respective homologues.

[0052] When two sequences have been identified for comparison, GAP and BESTFIT programs can be employed to determine their optimal alignment. For this purpose, the percent of sequence identity is preferably determined using the BESTFIT or GAP program of the Sequence Analysis Software PackageTM (Version 10; Genetics Computer Group, Inc., Madison, Wis.). GAP utilizes the algorithm of Needleman and Wunsch (Needleman and Wunsch, *J. Mol. Biol.* 48:443-453, 1970) to find the alignment of two sequences that maximizes the number of matches and minimizes the number of gaps. BESTFIT performs an optimal alignment of the best segment of similarity between two sequences and inserts gaps to maximize the number of matches using the local homology algorithm of Smith and Waterman (Smith and Waterman, *Adv. Applied Math.*, 2:482-489, 1981, Smith *et al.*, *Nucl. Acids Res.* 11:2205-2220, 1983). The percent identity is most preferably determined using the BESTFIT program. Typically, the default values of 5.00 for gap weight and 0.30 for gap weight length are used. The term "substantial sequence identity" between polynucleotide or polypeptide sequences refers to polynucleotide or polypeptide comprising a sequence that has at least 50% sequence identity, preferably at least about 70%, preferably at least about 80%, more preferably at least about 85%, more preferably at least about 90%, even more preferably at least about 95%, and most preferably

at least about 96%, 97%, 98% or 99% sequence identity compared to a reference sequence using the programs. Thus, according to one embodiment of the invention are protein, peptide, or polynucleotide molecules that have at least about 50% sequence identity, preferably at least about 70%, preferably at least about 80%, more preferably at least about 85%, more preferably at least about 90%, even more preferably at least about 95%, and most preferably at least about 96%, 97%, 98% or 99% sequence identity with a protein, peptide, or polynucleotide sequence described herein. Polynucleotide molecules that are capable of regulating transcription of operably linked transcribable polynucleotide molecules and have a substantial percent sequence identity to the polynucleotide sequences of the polynucleotide molecules provided herein are encompassed within the scope of this invention.

[0053] In one aspect of the invention, the present disclosure also provides polypeptides that are encoded by any of the polynucleotides of the invention described herein. Thus, the invention provides polypeptides of SEQ ID NOs: 2, 4, 6, 8, 10, 11, 13, 14, 16, 17, 19, 21, 23, 25, 27, 28, and 29. The invention also provides variants or fragments of the polynucleotides disclosed herein, and polypeptides encoded by any of the polynucleotide variants or fragments disclosed herein.

[0054] The endonuclease polypeptides of the present invention, including full-length polypeptides and biologically active fragments and fusion polypeptides, can be produced in genetically engineered host cells according to conventional techniques. Suitable host cells are those cell types that can be transformed or transfected with exogenous DNA and grown in culture, and include bacteria, insect cells, plant cells, fungal cells, and cultured higher eukaryotic cells. Eukaryotic cells, particularly cultured cells of multicellular organisms, are preferred. Techniques for manipulating cloned DNA molecules and introducing exogenous DNA into a variety of host cells are disclosed by Sambrook et al., 1989, *supra*; and Ausubel et al., eds., *Current Protocols in Molecular Biology*, John Wiley and Sons, Inc., NY, 1987.

[0055] In general, a nucleic acid sequence encoding an endonuclease polypeptide is operably linked to other genetic elements required for its expression, generally including a transcription promoter and terminator, within an expression vector or construct. The vector or construct will also commonly contain one or more selectable markers and one or more origins of replication, although those skilled in the art will recognize that within certain

systems selectable markers may be provided on separate vectors, and replication of the exogenous DNA may be provided by integration into the host cell genome. Selection of promoters, terminators, selectable markers, vectors and other elements is a matter of routine design within the level of ordinary skill in the art. Many such elements are described in the literature and are available through commercial suppliers.

[0056] To direct an endonuclease polypeptide into the secretory pathway of a host cell, a secretory signal sequence (also known as a leader sequence, pre sequence, or prepro sequence) can be included in the expression vector. The secretory signal sequence may be that of the native endonuclease polypeptide, or may be derived from another secreted protein or synthesized de novo. The secretory signal sequence is operably linked to the endonuclease-encoding DNA sequence, *i.e.*, the two sequences are joined in the correct reading frame and positioned to direct the newly synthesized polypeptide into the secretory pathway of the host cell. Secretory signal sequences are commonly positioned 5' to the DNA sequence encoding the polypeptide of interest, although certain secretory signal sequences may be positioned elsewhere in the DNA sequence of interest (see, *e.g.*, U.S. Pat. Nos. 5,037,743 and 5,143,830).

[0057] A variety of prokaryotic and eukaryotic cells are suitable host cells for the present invention, including but are not limited to microbial cells, algal cells, fungal cells, insect cells, mammalian cells, and plant cells. For example, when plants cells are used as hosts, the use of *Agrobacterium rhizogenes* as a vector for expressing genes in plant cells is well known in the field of plant biotechnology. Transformation of insect cells and production of foreign polypeptides therein is described extensively in, for example, U.S. Pat. No. 5,162,222 and WIPO publication WO 94/06463. Insect cells can be infected with recombinant baculovirus, commonly derived from *Autographa californica* nuclear polyhedrosis virus (AcNPV). See, *e.g.*, D. R. *et al.*, *Baculovirus Expression Vectors: A Laboratory Manual*, New York, Oxford University Press., 1994; and Richardson, Ed., *Baculovirus Expression Protocols. Methods in Molecular Biology*, Totowa, N.J., Humana Press, 1995. The second method of making recombinant baculovirus utilizes a transposon-based system described by Luckow *et al.* (*J Virol* 67:4566-79, 1993), Bac-to-Bac® Kit (Life Technologies, Inc., Carlsbad, CA). This system utilizes a transfer vector, pFastBac1™ (Life Technologies, Inc., Carlsbad, CA) containing a Tn7 transposon to move

the DNA encoding a polypeptide of interest into a baculovirus genome maintained in *E. coli* as a large plasmid called a "bacmid." The pFastBac1TM transfer vector utilizes the AcNPV polyhedrin promoter to drive the expression of the gene of interest, in this case a mismatch endonuclease. Further, pFastBac1TM (Life Technologies, Inc., Carlsbad, CA) can be modified to a considerable degree. The polyhedrin promoter can be removed and substituted with the baculovirus basic protein promoter (also known as Pcor, p6.9 or MP promoter) which is expressed earlier in the baculovirus infection, and has been shown to be advantageous for expressing secreted proteins. See, *e.g.*, Hill-Perkins and Possee *J. Gen. Virol.* 71:971-6, 1990; Bonning *et al.*, *J. Gen. Virol.* 75:1551-6, 1994; and, Chazenbalk and Rapoport, *J. Biol. Chem.* 270:1543-9, 1995. In such transfer vector constructs, a short or long version of the basic protein promoter can be used. Moreover, transfer vectors can be constructed to include secretory signal sequences derived from insect proteins. For example, a secretory signal sequence from Ecdysteroid Glucosyltransferase (EGT), honey bee melittin, or baculovirus gp67 can be used in recombinant nucleic acid constructs in accordance with the present invention. In addition, transfer vectors can include an in-frame fusion with DNA encoding an epitope tag at the C- or N-terminus of the expressed endonuclease polypeptide. Using techniques known in the art, a transfer vector containing an endonuclease of the present invention may be transformed into *E. coli*, and screened for bacmids which contain an interrupted lacZ gene indicative of recombinant baculovirus. The bacmid DNA containing the recombinant baculovirus genome can be isolated, using common techniques, and used to transfect *Spodoptera frugiperda* insect cells, *e.g.* Sf9 cells. Recombinant virus that expresses recombinant endonuclease can be subsequently produced. Recombinant viral stocks can be made by methods commonly used the art.

[0058] Fungal cells, including yeast cells are suitable as hosts for the present invention. Yeast species of particular interest in this regard include *Saccharomyces cerevisiae*, *Pichia pastoris*, and *Pichia methanolica*. Methods for transforming cells of these yeast species with exogenous DNA and producing recombinant polypeptides therefrom are well known in the art. See, for example, U.S. Pat. Nos. 4,599,311; 4,931,373; 4,870,008; 5,037,743; and 4,845,075. Transformed cells are selected by phenotype determined by the selectable marker, commonly drug resistance or the ability to grow in the absence of a particular nutrient (*e.g.*, adenine or leucine). Suitable promoters and

terminators for use in yeast include those from glycolytic enzyme genes (see, *e.g.*, U.S. Pat. Nos. 4,599,311; 4,615,974; and 4,977,092) and alcohol dehydrogenase genes. See also U.S. Pat. Nos. 4,990,446; 5,063,154; 5,139,936 and 4,661,454. The use of *Pichia methanolica* as host for the production of recombinant proteins is well known (see, *e.g.*, PCT Publication Nos. WO199717450, WO199717451, WO199802536, and WO 91998/902565).

Transformation systems for other yeasts, including *Hansenula polymorpha*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Kluyveromyces fragilis*, *Ustilago maydis*, *Pichia pastoris*, *Pichia guilliermondii* and *Candida maltosa* are also known in the art. See, for example, Gleeson *et al.*, *J. Gen. Microbiol.* 132:3459-65, 1986 and U.S. Pat. No. 4,882,279. *Aspergillus* cells may be used as recombinant host cells according to a variety of known methods described in, for example, U.S. Pat. No. 4,935,349. Methods for transforming *Acremonium chrysogenum* and *Neurospora* sp. are also well known (see, *e.g.*, U.S. Pat. Nos. 5,162,228; 4,486,533).

[0059] Prokaryotic host cells, including strains of the bacteria *Escherichia coli*, *Bacillus* and other genera are also useful host cells within the present invention. Techniques for transforming these hosts and expressing foreign DNA sequences cloned therein are well known in the art (see, *e.g.*, Sambrook *et al.*, *Ibid.*). When expressing an endonuclease polypeptide in bacteria such as *E. coli*, the polypeptide may be directed to the periplasmic space by a bacterial secretion sequence, or may be retained in the cytoplasm, typically as insoluble granules. In the former case, the polypeptide can be recovered from the periplasmic space in a soluble and functional form by disrupting the cells (by, for example, sonication or osmotic shock) to release the contents of the periplasmic space and recovering the protein, thereby obviating the need for denaturation and refolding. In the latter case, the cells are lysed, and the granules are recovered and denatured using, for example, guanidine isothiocyanate or urea. The denatured polypeptide can then be refolded and dimerized by diluting the denaturant, such as by dialysis against a solution of urea and a combination of reduced and oxidized glutathione, followed by dialysis against a buffered saline solution.

[0060] In addition, cultured mammalian cells are also suitable hosts for the present invention. Methods for introducing exogenous DNA into mammalian host cells are well known and include, but are not limited to, liposome-mediated transfection (Hawley-Nelson *et al.*, *Focus* 15:73, 1993; Ciccarone *et al.*, *Focus* 15:80, 1993); calcium phosphate-mediated transfection (Wigler *et al.*, *Cell* 14:725, 1978; Corsaro and Pearson, *Somatic Cell*

Genetics 7:603, 1981; Graham and Van der Eb, *Virology* 52:456, 1973); electroporation (Neumann *et al.*, *EMBO J.* 1:841-5, 1982), DEAE-dextran mediated transfection (Ausubel *et al.*, *ibid.*), and viral vectors (Miller and Rosman, *BioTechniques* 7:980-90, 1989; Wang and Finer, *Nature Med.* 2:714-6, 1996). The production of recombinant polypeptides in cultured mammalian cells is described extensively in scientific literature and patent literature (see, *e.g.*, U.S. Pat. Nos. 4,713,339; 4,784,950; 4,579,821; and 4,656,134). Suitable cultured mammalian cells include, but are not limited to, BHK (ATCC No. CRL 1632), BHK 570 (ATCC No. CRL 10314), COS-1 (ATCC No. CRL 1650), COS-7 (ATCC No. CRL 1651), 293 (ATCC No. CRL 1573; Graham *et al.*, *J. Gen. Virol.* 36:59-72, 1977) and Chinese hamster ovary (*e.g.* CHO-K1; ATCC No. CCL 61) cell lines. Additional suitable cell lines are known in the art and available from public depositories such as the American Type Culture Collection, Manassas, Va. In general, strong transcription promoters are preferred, such as promoters from SV-40 or cytomegalovirus. See, *e.g.*, U.S. Pat. No. 4,956,288. Other suitable promoters include those from metallothionein genes (U.S. Pat. Nos. 4,579,821 and 4,601,978) and the adenovirus major late promoter.

Methods for correcting errors in nucleic acid molecules

[0061] In one aspect, embodiments or methods of the present invention provide a process for error correction in nucleic acid molecules. Errors arise in the replication, amplification, and/or synthesis of nucleic acid molecules. An "error" is a deviation from the nucleotide sequence that the nucleic acid molecules are intended to have, *e.g.* the desired sequence resulting from replication and/or amplification and/or synthesis procedures. Errors include deletions from, substitutions in, and additions to the desired nucleotide sequence, and may arise at any point in the synthesis by any mechanism.

[0062] The chemical synthesis of oligonucleotides is inherently subject to the occurrence of errors in nucleotide insertion due to the limitations of the chemistry involved, which generally has involved some type of solid-phase synthesis involving sequential addition of nucleotides to the 3' end of the growing molecule. The occurrence of incomplete reactions or side reactions places an upper limit on the length of nucleotides that can be synthesized, but even shorter nucleotides incorporate some rate of unintended or erroneous nucleotides.

[0063] The assembled nucleic acid molecules are double-stranded by default. Double-stranded nucleic acid molecules can be denatured and annealed by conventional methods. For example, heat denaturation of double-stranded nucleic acid molecules separates the double-stranded molecules into pairs of corresponding single-stranded molecules. Cooling the single-stranded molecules promotes their annealing into double-stranded molecules as individual nucleotides comprising the nucleic acid molecules coalesce into nucleotide base pairs along complementary stretches of nucleotide sequence. The kinetics or other physical or chemical parameters of denaturation and annealing may be controlled to promote mixing of the single-stranded molecules, so that the single-stranded molecules change partners. For example, if a double-stranded DNA molecule had a sequence error in both strands at the 400th nucleotide from one end, after denaturation and annealing, the single strands of that molecule may be paired with other single-stranded molecules lacking an error at that position, resulting in a nucleotide mismatch at that position. Thus, the denaturation and annealing process can produce double-stranded nucleic acid molecules with mismatches between nucleotide bases at sites of error. These mismatches can be targeted for removal, for example, by reacting annealed molecules with endonucleases having certain characteristics under appropriate conditions. A mismatch site or nucleotide mismatch site is a site on a double-stranded nucleic acid molecule where non-complementary base pairs are situated opposite each other. Nucleotide mismatch is caused by the erroneous insertion, deletion or mis-incorporation of bases that can arise during DNA replication or amplification. Examples of mismatched bases are G/T or A/C pairing or other deviations from standard G/C and A/T Watson-Crick base pairing. Mismatches can also be caused by tautomerization of bases during synthesis.

[0064] An aspect of the invention may be practiced to correct and reduce errors in double-stranded nucleic acid molecules. A first set of double-stranded nucleic acid molecules, which are intended to have a desired nucleotide sequence and a desired length, are reacted with one or more endonucleases. In one embodiment the endonuclease is a mismatch endonuclease that cuts the nucleic acids at or near the mismatch site, resulting in a nucleic acid molecule having a mismatched end. This can be accomplished by the selection of one or more appropriate endonucleases. The nucleic acid molecules having a mismatch are thus cut into smaller fragments that have a nucleotide mismatch at or near the

end. The nucleic acids having the mismatch end are then treated with an endonuclease which cuts into the mismatch end and thus removes the mismatch. The resulting overhangs are then filled by the action of another enzyme, *e.g.* a DNA polymerase or other molecule having polymerase activity, and made into a fully double-stranded nucleic acid molecule. In one embodiment the process of denaturing, annealing, cutting with an appropriate endonuclease, and filling an overhang if necessary, can be repeated until the mismatch rate in the sample is insignificant, meaning that the rate of error is so low that it has no material effect on the outcome of the study for which the nucleic acids are being used. Nucleic acids having a mismatch at or near the end refers to a nucleic acid molecule having a mismatched nucleotide pair within 10 or less nucleotides of the end of the nucleic acid molecule. In other embodiments the mismatched nucleotide pair is present within 9 or less nucleotides of the end of the nucleic acid molecule, or within 8 or less, or within 7 or less, or within 6 or less, or within 5 or less, or within 4 or less, or within 3 or less, or within 2 or less, or within 1 nucleotide of the nucleotide mismatch.

[0065] FIGURE 1 is a diagram showing a general illustration of a method of the invention. In this embodiment, error-free dsDNA molecules 101 are contained in a solution with dsDNA molecules having an error 103. Error-free dsDNA molecules are those having the “correct” desired nucleotide sequence, *i.e.* the desired sequence, whereas those dsDNA molecules having an error have an “incorrect” nucleotide in their sequence, *i.e.*, that deviates from the desired sequence. The dsDNA molecules are denatured and then annealed, producing double-stranded nucleic acid molecules (dsDNA) with one or more errors or mismatches 105 in one of the single strands of the dsDNA. The dsDNA molecules are then exposed to the action of an endonuclease, thus cleaving the dsDNA into fragments. In some embodiments the endonuclease is a mismatch endonuclease that produces a dsDNA fragment having a mismatch at or near the end of the molecule 107. In the embodiment depicted in Fig. 1 the mismatch occurs at the 3’ end of one of the single strands, but the mismatch can also occur at the 5’ end depending on which endonuclease(s) are selected. In the embodiment depicted the nucleic acids are then exposed to the action of a 3’ exonuclease, which “chews” away the 3’ end of the molecule and thus removes the mismatch error by removing the incorrect nucleotide. The nucleic acid molecules are then again denatured, annealed, exposed to mismatch endonuclease, filled if desirable, and

optionally amplified to produce error-free dsDNA molecules 111. The process can be repeated several times until the rate of error in the strands is insignificant. In some embodiments a ligase can be used to ligate the strands if desired.

[0066] Double-stranded nucleic acid molecules in the invention can be fragmented by reacting them with a unidirectional mismatch endonuclease. Unidirectional mismatch endonuclease refers to any molecule or combination of molecules having unidirectional mismatch endonuclease activity. In one embodiment the molecule is an enzyme, but the molecule can also be another molecule that is not an enzyme but that nevertheless has unidirectional endonuclease activity. The unidirectional mismatch endonucleases can be used either as a single endonuclease or as a mixture of endonucleases and other molecules. In one embodiment a single mismatch endonuclease enzyme is used, and in other embodiments a single mismatch endonuclease enzyme can be used in combination with another non-enzymatic molecule that is necessary for or enhances mismatch endonuclease activity.

[0067] As used herein, the term "mismatch endonuclease" refers to an enzyme activity that is able to both recognize a mismatch in a heteroduplex polynucleotide (*e.g.*, a double-stranded nucleic acid molecule containing a nucleotide mismatch) and cut one or both strands of the heteroduplex at or near the mismatch. A molecule having unidirectional mismatch endonuclease activity consistently cuts on one side of the mismatch, either the 5' or 3' side, and not on the other side. In one embodiment the mismatch endonuclease is substantially unidirectional, meaning that at least 90% of the cuts are on one side of the mismatch, either the 5' or 3' side, but allowing for up to 10% of the cuts being on the opposing side. In various embodiments the mismatch endonucleases of the invention can recognize a nucleotide mismatch in the heteroduplex polynucleotide and cut both strands of the heteroduplex. In various embodiments the cut is introduced within 10 or less nucleotides of the nucleotide mismatch. In other embodiments the cut is introduced within 9 or less nucleotides, or 8 or less, or 7 or less, or 6 or less, or 5 or less, or 4 or less, or 3 or less, or 2 or less, or within 1 nucleotide of the nucleotide mismatch. In another embodiment the cut is introduced at the nucleotide mismatch site, thus leaving at least one of the mismatched nucleotides at the terminal end of the nucleic acid molecule. In one embodiment the mismatch endonuclease leaves a blunt end cut over both strands of the

heteroduplex. When a blunt end is left at a nucleotide mismatch site the mismatched nucleotide pair is present at the end of the nucleic acid molecule. But in other embodiments the cuts can produce an overhang of one or more nucleotides such as, for example, 1 nucleotide or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or more than 10 nucleotides. Conventional methods are used to assemble these fragments into a second set of double-stranded nucleic acid molecules 111, which are overwhelmingly more likely to have the desired nucleotide sequence and desired length than were the first set of molecules.

[0068] A variety of mismatch endonucleases will find use in the present invention. RES I, CEL I, CEL II, SP nuclease, SP I endonuclease, T7 endonuclease, T4 endonuclease, endonuclease V, a Mut protein, are all useful mismatch endonucleases. One unidirectional mismatch endonuclease useful in the invention is commercially available as SURVEYOR® nuclease (Transgenomic, Inc., Omaha, NE), which uses CEL II as a principal component. It is also advantageous to utilize combinations of more than one of any of these. In a specific embodiment the mismatch endonuclease utilized is a combination of CEL I and CEL II mismatch endonucleases. Some of these have also been expressed recombinantly (CEL I and SP I) (Pimkin et al. *BMC Biotechnology*, 7:29 (2007)). SP nuclease has been described by Doetsch et al. *Nucleic Acids Res.*, Vol. 16, No. 14 (1988).

[0069] In various embodiments a component can also be added to the reaction mixture to increase the action of endonucleases to create a double-stranded break in the nucleic acid molecule. Some endonucleases, *e.g.* endonuclease V, cleave only one strand of the nucleic acid molecule. But cleaving of both strands can be promoted by the inclusion in the medium of manganese ions, Mn^{+2} at an appropriate concentration. In one embodiment the reaction medium includes about 10nM Mn^{+2} in a convenient form, *e.g.* $MnCl_2$. In another embodiment the additional step is taken to exclude magnesium Mg^{+2} from the reaction medium.

[0070] Variants of the mismatch endonucleases disclosed herein are also useful in the invention. With reference to this disclosure the person of ordinary skill will realize many more endonucleases that will be useful in the present invention. It is also likely that new endonucleases having the required activity will be discovered or developed, and those can also find use in the application of the present invention. Thus, variants and homologs of the mismatch endonucleases disclosed herein are also useful in the invention. In various

embodiments a protein having at least 70% sequence identity, or at least 75% sequence identity, or at least 80% sequence identity, or at least 85% sequence identity, or at least 90% sequence identity, or at least 95% or 96% or 97% or 98% or 99% sequence identity to any endonuclease disclosed herein is also useful in the invention. Thus, in two separate embodiments a protein having any of the above sequence identities to either CEL I or CEL II can be used in the invention.

[0071] In an alternative embodiment, the protein, peptide, or nucleic acid molecules useful in the invention comprise a protein, peptide, or nucleic acid sequence that exhibits 70% or greater identity, and more preferably at least 75% or greater, 80% or greater, 85% or greater, 87% or greater, 88% or greater, 89% or greater, 90% or greater, 91% or greater, 92% or greater, 93% or greater, 94% or greater, 95% or greater, 96% or greater, 97% or greater, 98% or greater, or 99% or greater identity to a protein, peptide, or nucleic acid molecule selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 33 in the Sequence Listing, any complements thereof, any fragments thereof, or any functional domain thereof. Thus, all variants and homologs of all endonucleases and exonuclease disclosed herein will be useful in this invention, and such variants and homologs can be discovered or designed using the principles disclosed herein.

[0072] For nucleic acids and polypeptides, the term "variant" is used herein to denote a polypeptide, protein or polynucleotide molecule with some differences, generated synthetically or naturally, in their base or amino acid sequences as compared to a reference polypeptide or polynucleotide, respectively. For example, these differences include substitutions, insertions, deletions or any desired combinations of such changes in a reference polypeptide or polynucleotide. Polypeptide and protein variants can further consist of changes in charge and/or post-translational modifications (such as glycosylation, methylation, phosphorylation, *etc.*). Biologically active variants of the polynucleotide sequences are also encompassed by the compositions of the present invention. Biologically active variants of the invention may be created by site-directed mutagenesis, induced mutation, or may occur as allelic variants (polymorphisms). Synthetic nucleotide sequences are those that are made through chemical processes in a laboratory environment. An example of a synthetic nucleotide is an oligonucleotide made using the known phosphoramidite chemical synthesis. This chemical method joins nucleotides in the 3' to 5'

direction using phosphoramidite building blocks derived from protected 2'-deoxynucleosides (dA, dC, dG, and T). Another process for creating synthetic nucleic acids is the polymerase chain reaction. Naturally produced oligonucleotides are synthesized either in Nature or in the laboratory using natural processes, such as the synthesis of an oligonucleotide by a microorganism. Synthetic nucleotide sequences can differ from naturally produced nucleotide sequences because the naturally produced sequences may have been processed through some post-transcriptional modification to result in a chemically changed nucleotide sequence. Synthetically produced oligonucleotides can be assembled from smaller oligonucleotides or subsets to form larger synthetic oligonucleotides.

[0073] The term “functional homolog” as used herein describes those proteins or polypeptides that have at least one characteristic in common. Such characteristics include sequence similarity, biochemical activity, transcriptional pattern similarity and phenotypic activity. Typically, a functional homolog is a polypeptide that has sequence similarity to a reference polypeptide, and that carries out one or more of the biological activities of the reference polypeptide. Functional homologs will typically give rise to the same characteristics to a similar, but not necessarily the same, degree. Typically, functionally homologous proteins give the same characteristics where the quantitative measurement due to one of the homologs is at least 20% of the other; more typically, between 30 to 40%; more typically, between 50-60%; even more typically, between 70 to 80%; even more typically, between 90 to 95%; even more typically, between 98 to 100% of the other.

[0074] A functional homolog and the reference polypeptide may be naturally occurring polypeptides, and the sequence similarity may be due to convergent or divergent evolutionary events. As such, functional homologs are sometimes designated in the literature as homologs, orthologs, or paralogs. Variants of a naturally-occurring functional homolog, such as polypeptides encoded by mutants or a wild-type coding sequence, may themselves be functional homologs. As used herein, functional homologs can also be created via site-directed mutagenesis of the coding sequence for a polypeptide, or by combining domains from the coding sequences for different naturally-occurring polypeptides. The term “functional homolog” sometimes applied to the nucleic acid that encodes a functionally homologous polypeptide.

[0075] Functional homologs can be identified by analysis of nucleotide and polypeptide sequence alignments. For example, performing a query on a database of nucleotide or polypeptide sequences can identify homologs of polypeptides. Sequence analysis can involve BLAST, Reciprocal BLAST, or PSI-BLAST analysis of non-redundant databases using amino acid sequence of a polypeptide as the reference sequence. Amino acid sequence is, in some instances, deduced from the nucleotide sequence. Typically, those polypeptides in the database that have greater than 40% sequence identity are candidates for further evaluation for suitability as a polypeptide. Amino acid sequence similarity allows for conservative amino acid substitutions, such as substitution of one hydrophobic residue for another or substitution of one polar residue for another. If desired, manual inspection of such candidates can be carried out in order to narrow the number of candidates to be further evaluated. Manual inspection can be performed by selecting those candidates that appear to have domains present in the polypeptide of interest, *e.g.*, conserved functional domains.

[0076] Conserved regions can be identified by locating a region within the primary amino acid sequence of a polypeptide that is a repeated sequence, forms some secondary structure (*e.g.*, helices and beta sheets), establishes positively or negatively charged domains, or represents a protein motif or domain. See, *e.g.*, the Pfam web site describing consensus sequences for a variety of protein motifs and domains on the World Wide Web at [sanger.ac.uk/Software/Pfam/and pfam.janelia.org/](http://sanger.ac.uk/Software/Pfam/and_pfam.janelia.org/). A description of the information included at the Pfam database is described in, for example, Sonnhammer *et al.* (*Nucl. Acids Res.*, 26:320-322, 1998), Sonnhammer *et al.* (*Proteins*, 28:405-420, 1997); and Bateman *et al.* (*Nucl. Acids Res.*, 27:260-262, 1999). Conserved regions also can be determined by aligning sequences of the same or related polypeptides from closely related species. Closely related species preferably are from the same family. In some embodiments, alignment of sequences from two different species is adequate. Examples of domains indicative of the activity of the polypeptide of interest can be found in various literature sources and species, such as plants, algae, fungi, bacteria, and animals, which can be investigated.

[0077] Typically, polypeptides that exhibit at least 40% amino acid sequence identity are useful to identify conserved regions. Conserved regions of related polypeptides exhibit at least 45% amino acid sequence identity, *e.g.*, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90% amino acid sequence identity. In some embodiments, a

conserved region exhibits at least 92%, 94%, 96%, 98%, or 99% amino acid sequence identity.

[0078] As used herein, the term “exonuclease” refers to an enzyme activity that removes nucleotides from one or more termini of a polynucleotide. In some embodiments the polynucleotide is bound to a second polynucleotide to form a double-stranded nucleic acid molecule. A molecule having unidirectional exonuclease activity proceeds to remove nucleotides in a 5' to 3' direction or a 3' to 5' direction in a stepwise manner. The molecule can be an enzyme, or another molecule that has exonuclease activity. A molecule having a substantially unidirectional exonuclease activity indicates that at least 90% of the nucleotides removed are on the 5' or 3' side of the nucleic acid molecule, and from 0% up to 10% are removed on the opposite side of the nucleic acid molecule. In one embodiment the exonuclease used in the method of the invention has the same directionality as the mismatch endonuclease used in the method.

[0079] A variety of exonucleases will be useful in the present invention. Exonuclease refers to any molecule having exonuclease activity, including both enzyme and non-enzyme molecules. In various embodiments the exonuclease is exonuclease III, a DNA polymerase having exonuclease activity, lambda exonuclease, T7 exonuclease, and T5 exonuclease, and variants thereof. The exonucleases can also be utilized in a combination of two or more of the exonucleases. DNA polymerases often have a 3' exonuclease activity, and one such DNA polymerase that is commercially available is the PHUSION® DNA polymerase (Finnzymes Oy, Espoo, Sweden). Other DNA polymerases with exonuclease activity include T4 DNA polymerase, phi29 polymerase. Variants and homologs of exonucleases disclosed herein can also find use in the present invention, and can be identified or discovered as disclosed herein.

[0080] Figure 2 depicts five distinct oligos 202 for the purpose of illustration, but any number of oligos may be used. The oligos may be obtained in any manner, including purchase from an industrial supplier and/or independent synthesis. Any number of oligos may be obtained in a manner different from that in which one or more other oligos are obtained. Any oligo may or may not be sequenced to determine whether it comprises enough molecules with the desired nucleotide sequence. Any of the oligos may optionally be further purified to reduce the number of any nucleotide-sequence errors they may bear.

[0081] In some embodiments, the oligos are obtained for both strands of the nucleic acid molecule that is intended to have a desired nucleotide sequence. Oligos 202 can be obtained for only a single strand of DNA that is intended to have a desired nucleotide sequence. In some embodiments oligos 202 may be obtained for both strands of DNA so that a set of oligos 202 comprises some oligos 202 having overlapping fragments of a full-length desired nucleotide sequence. A set of oligos 202 with such sequence overlaps can be used to assemble a full-length molecule intended to have a desired nucleotide sequence more efficiently than is otherwise possible. This increase in efficiency means that a smaller amount of full-length molecules (or even no full-length molecules) intended to have a desired nucleotide sequence may be used in order to obtain more full-length molecules intended to have a desired nucleotide sequence. This efficiency allows better control of the costs of nucleic acid molecule synthesis.

[0082] Referring to Figs. 1 and 2, the oligos 202 are amplified into the oligos 204, increasing the number of molecules comprising each oligo 202. Each amplified oligo 204 is represented by a double arrow. The double arrow is merely a representational device: the number of molecules of each oligo 204 after amplification is not necessarily twice the number of molecules of each oligo 202 present before amplification, and is likely orders of magnitude greater. Any amplified oligo 204 may or may not be sequenced to determine whether it comprises enough molecules with the desired nucleotide sequence. Any amplified oligo 204 optionally may be further purified to reduce the number of any nucleotide sequence errors they may bear.

[0083] The amplified oligos 204 are used to assemble a first set of full length molecules 206 that are intended to have a desired nucleotide sequence. Double, parallel line-segments represent a full-length, double-stranded DNA molecule 206. Within a set of such full-length molecules 206, however, it is expected that there may be one or more molecules with one or more sequence errors 208. Sequence errors are denoted with a short slash along the full-length molecule 208. There may be many molecules 208 with one or more sequence errors at different points in the sequence. Within a set of such full-length molecules 206, it may also be expected that there are one or more molecules without any sequence errors 210.

[0084] The first set of dsDNA molecules 206 is denatured, so that the two strands of each molecule separate. The set of denatured, single-stranded, full-length molecules 212 thus may comprise one or more molecules without sequence errors 214, and one or more molecules with one or more sequence errors 216. There may be many molecules 216 with one or more sequence errors at different points in the sequence. The set of full-length molecules 206 may be denatured in any manner, for example, by heating the molecules 206.

[0085] The set of denatured molecules 212 is then annealed to obtain double-stranded DNA (dsDNA) molecules 218 that are intended to have a desired nucleotide sequence. Within a set of such dsDNA molecules, it may be expected that there are one or more molecules with one or more sequence errors, or mismatches in the dsDNA (220 and 105), and one or more molecules without any sequence errors or mismatches (not shown). The denatured set of single-stranded (ssDNA) molecules 212 may be annealed in any manner, for example, by cooling the molecules.

[0086] There may be many molecules (220 and 105) with one or more sequence errors or mismatches in dsDNA at different points in the sequence. The distribution of sequence errors over the second set of molecules 218 will most likely be different from that over the first set of molecules 206, since one or more single-stranded molecules 214 and 216 will anneal to other single-stranded molecules 214 and 216 different from those to which they were bound before denaturation. For example, a double-stranded molecule 208 in the first set of molecules 206 may have two sequence errors, one in each strand, are directly across from each other. During denaturation, a single strand 216 from the molecule 208 may move near a single-stranded molecule without errors 214. During annealing, a second full length molecule 220 may form that has an error in only one of its two strands.

[0087] The second set of full-length molecules 218 may be cut (e.g., by a mismatch endonuclease) to form a third set of molecules (not shown, but graphically depicted in Fig. 1 107), so that two or more molecules in the third set of molecules are shorter than full-length molecules 206 or 218. The cuts can occur wherever there is a mismatch in dsDNA. In one embodiment the cuts leave blunt ends and in other embodiments can leave sticky ends or overhangs. In one embodiment the endonuclease is one or more of the endonucleases that are unidirectional mismatch endonucleases disclosed herein. These endonucleases will cut dsDNA where there is a mismatch between the two strands of the dsDNA nucleic acid

molecule. This will thus leave dsDNA having a mismatch at the end of the nucleic acid molecule, such as in nucleic acid 107 (shown as a blunt end cut embodiment). These dsDNA can then be digested with an exonuclease, for example a unidirectional exonuclease, that will "chew off" the end of the molecule and eliminate the incorrect nucleotide, resulting in a dsDNA with overhangs (109 in Fig. 1). These dsDNA can then be annealed and amplified and any gaps filled in with a polymerase. In some embodiments nicks can be repaired with a ligase if necessary.

[0088] With reference to this disclosure it can be seen that with each cycle of denaturation, annealing, and cutting with endonucleases and exonucleases, the number of sequence errors in the set of molecules is much lower than in the starting set of molecules. By providing a unique and powerful error-correction process operating late in the nucleic acid molecule synthesis process, the exemplary method for error correction of nucleic acid molecules yields a set of full-length molecules 111 intended to have a desired nucleotide sequence that has remarkably fewer errors than can be otherwise obtained.

[0089] FIG. 3 is a flow chart depicting one embodiment of the method for synthesis of error-minimized nucleic acid molecules. At step 302, oligos 101 (Fig. 1) of a length smaller than that of the full-length desired nucleotide sequence (*i.e.*, "oligonucleotide fragments" of the full-length desired nucleotide sequence) are obtained. Each oligo 101 is intended to have a desired nucleotide sequence that comprises a part of the full length desired nucleotide sequence. In various embodiments each oligo 101 may also be intended to have a desired nucleotide sequence that comprises an adapter primer for PCR amplification of the oligo 101, a tethering sequence for attachment of the oligo to a DNA microchip, or any other nucleotide sequence determined by any experimental purpose or other intention. The oligos may be obtained in any of one or more ways, for example, through synthesis, purchase, *etc.*

[0090] At step 304, the oligos 101 obtained are amplified to obtain more of each oligo. The amplification may be accomplished by any method, for example, by PCR. Introduction of additional errors into the nucleotide sequences of any of the oligos 103 may occur during amplification. The distinct amplified oligos result from the amplification at step 304. Oligos may be amplified by adapter primers, and the adapter sequence may be cleaved off by means of type IIS restriction endonucleases.

[0091] At step 306 the amplified oligos are assembled into a first plurality of double-stranded nucleic acid, which in one embodiment are intended to have a desired length that is the full length of the desired nucleotide sequence intended to be synthesized. Assembly of amplified oligos into full-length molecules may be accomplished in any way, for example, by using a PCR-based method. One or more of the double-stranded nucleic acid molecules (or full-length molecules) may be a double-stranded nucleic acid molecule containing at least one nucleotide mismatch (105), caused by one or more sequence errors in one or both of its strands. And one or more of the double-stranded nucleic acid molecules (or full length molecules) may be a double-stranded nucleic acid molecule containing no nucleotide mismatches or sequence errors in one of its single strands 105 (FIG. 1).

[0092] At step 308 the first plurality of double-stranded nucleic acid molecules are reacted with one or more endonucleases. In some embodiments the endonuclease is a unidirectional mismatch endonuclease, which fragments the double-stranded nucleic acid molecules having at least one nucleotide mismatch by cutting at or near the mismatched nucleotide pair. The one or more endonucleases thus cut the double-stranded nucleic acid molecules having a mismatch into shorter molecules that have a mismatch nucleotide pair at or near the terminal nucleotide of the nucleic acid molecule. In the case of a sticky end or overhang, the terminal nucleotide is the last nucleotide of the single stranded overhang. In the case of a blunt end cut, the terminal nucleotide will be either of the terminal pair of nucleotides of the nucleic acid molecule.

[0093] At step 310 of the embodiment of the methods the nucleotide mismatch is removed from the double-stranded nucleic acid molecules having a mismatch at or near the terminal nucleotide of the nucleic acid molecule by the action of a molecule having unidirectional exonuclease activity. In one embodiment the unidirectional exonuclease activity is of the same directionality as the unidirectional mismatch endonuclease activity. Thus, if the unidirectional mismatch endonuclease cleaves to the 3' side of the nucleotide pair mismatch, then the unidirectional exonuclease can chew away nucleotides at the 3' end of the nucleic acid strands comprising the nucleic acid molecule. The nucleotides that are chewed away by the exonuclease can then be replaced by the action of a polymerase either immediately or during the next optional replication/amplification phase. This then provides a fragmented error-free double-stranded nucleic acid molecule.

[0094] Finally in step 312 a second plurality of double-stranded nucleic acid molecules having the fragmented error-free double-stranded nucleic acid molecules 111 is assembled. This second plurality of double-stranded nucleic acid molecules has a decreased frequency of nucleotide mismatches as compared to the first plurality of double-stranded nucleic acid molecules.

[0095] In various embodiments of the invention the steps of the methods can be varied. For example in one embodiment the steps of fragmenting double-stranded nucleic acid molecules having a nucleotide mismatch, and the step of removing incorrect nucleotides with a matching unidirectional exonuclease can be performed sequentially as separate steps in a two-step reaction. But in other embodiments, one or more of the steps involved in the methods can be performed simultaneously. Thus, in one embodiment the reaction with endonuclease and exonuclease (*e.g.*, steps 308 and 310 depicted in the embodiment in Fig. 2) can be performed simultaneously as a one-step reaction. This embodiment involves identifying reaction parameters where both the mismatch endonuclease and unidirectional exonuclease can perform their reactions in the same reaction. In a one-step reaction the reaction container does not need to be opened once all of the components necessary for the reaction have been added until the reaction is complete. In some embodiments the fragmentation step can be performed with SURVEYOR[®] nuclease (Transgenomic, Inc., Omaha, NE) and the exonuclease/error correction step performed with exonuclease III, and this combination of enzymes can be used in either a two-step or one-step procedure, as detailed in Example 3.

[0096] The present invention also provides compositions useful for performing the methods of the invention. The compositions can comprise (i) a molecule having a unidirectional mismatch endonuclease activity; and (ii) a molecule having unidirectional exonuclease activity of the same directionality as the unidirectional mismatch endonuclease activity in (i). In various embodiments the molecule having unidirectional mismatch endonuclease activity can be any described herein. The molecule having unidirectional exonuclease activity can also be any molecule having said activity described herein. The molecules can be combined in any combinations. In various examples the molecule having unidirectional mismatch endonuclease activity can be any of RES I, CEL I, CEL II, T7 endonuclease, T4 endonuclease, endonuclease V, a Mut protein, a variant of any thereof,

and a combination of any two or more thereof. This molecule can be combined in a composition with a molecule having unidirectional exonuclease activity, for example, any of exonuclease III, a DNA polymerase, a variant of any thereof, or a combination of any two or more thereof. In different embodiments the composition can be provided in a dried form, or in a suitable buffer. The composition can also be provided in a tube, vial, or other suitable container. In one embodiment the molecules of (i) and (ii) are provided in a purified form in the container.

[0097] The present invention also provides kits useful in performing the methods of the invention. The kits may include any two or more of the following components: a mismatch endonuclease, a 5' or 3' exonuclease, a DNA polymerase (with or without 3' exonuclease proofreading activity), suitable buffers for performing the method, instructions for performing one or more methods of the invention, and information identifying a website that contains information about an error correction method of the invention. The information may be instructions for performing a method of the invention. In one embodiment the kit contains a unidirectional mismatch endonuclease and exonuclease III, each in a quantity sufficient to perform a method of the invention. The kits of the invention can also comprise any composition described herein. The endonucleases, exonucleases and DNA polymerases included in the kits can be any disclosed herein, such as SURVEYOR® nuclease (Transgenomic, Inc., Omaha, NE) or a generic substitute, exonuclease III, and PHUSION® DNA polymerase (Finnzymes Oy, Finland) or a generic substitute of any. The components of the kits can be provided in individual suitable containers, or can be provided with one or more components in a single container. The components of the kit can be provided in a purified form in the containers. The components of the kit can also be provided in dried form, or within a suitable buffer.

[0098] The present invention can also be utilized in conjunction with various techniques for the manipulation of nucleic acids. For example, in one embodiment the error correction techniques of the present invention can be utilized following or in conjunction with methods of joining nucleic acid molecules to ensure correct replication of nucleic acid molecules during the procedure. Examples include methods disclosed in US Patent Application No. 2010/0035768. Although any methods of assembly of nucleic acids would benefit from the addition of error correction methods as disclosed herein. The kits of the

invention can therefore contain components for performing additional procedures and manipulations of DNA either before or after performing an error correction method of the invention. Therefore, in addition to the kit components described above, a kit may also include one or more of any of the following components: a non-thermostable 5' to 3' exonuclease that lacks 3' exonuclease activity (e.g., T5 exonuclease), a crowding agent (e.g., polyethylene glycol, a Ficoll), a thermostable non-strand displacing DNA polymerase with 3' exonuclease activity, a mixture of a said DNA polymerase and another DNA polymerase that lacks 3' exonuclease activity, and a thermostable ligase.

[0099] Throughout this disclosure, various information sources are referred to and incorporated by reference. The information sources include, for example, scientific journal articles, patent documents, textbooks, and World Wide Web browser-inactive page addresses. The reference to such information sources is solely for the purpose of providing an indication of the general state of the art at the time of filing. While the contents and teachings of each and every one of the information sources can be relied on and used by one of skill in the art to make and use embodiments of the invention, any discussion and comment in a specific information source should in no way be considered as an admission that such comment was widely accepted as the general opinion in the field.

[0100] The discussion of the general methods given herein is intended for illustrative purposes only. Other alternative methods and embodiments will be apparent to those of skill in the art upon review of this disclosure, and are to be included within the spirit and purview of this application.

[0101] It should also be understood that the following examples are offered to illustrate, but not limit, the invention.

EXAMPLE 1: Assembly of synthetic HA and NA genes from oligonucleotides.

[0102] Synthetic gene products were assembled from a plurality of oligonucleotides with overlapping sequences at their terminal ends using Gibson Assembly™ (Synthetic Genomics, Inc., San Diego, CA) as previously described (see, e.g., US Patent Application No. 2010/0035768). Genes included representative hemagglutinin (HA) genes and representative neuraminidase (NA) genes.

Reagents5X Isothermal (ISO) Buffer:

[0103] The 5X ISO buffer contains 25% PEG-8000, 500 mM Tris-HCl pH 8.0, 50 mM MgCl₂, 50 mM DTT, 1 mM each of the 4 dNTPs, and 5 mM NAD.

Six ml of this buffer can be prepared by combining the following:

3 ml of 1 M Tris-HCl pH 8.0

300 µl of 1 M MgCl₂

600 µl of 10 mM dNTPs

300 µl of 1 M DTT (1.54 g dissolved in dH₂O up to 10 ml)

1.5 g PEG-8000

300 µl of 100 mM NAD (Sigma; 0.66 g dissolved in dH₂O up to 10 ml; resuspend by heating at 50°C followed by continuous vortexing)

Add water to 6 ml, aliquot 1 ml and store at -80°C.

2X Assembly Master Mix:

[0104] The 2X Assembly Master Mix contains the ISO reaction buffer and the enzymatic activities required for assembly of the component oligonucleotides into the gene products: 5X isothermal (ISO) reaction buffer) T5 exonuclease (Epicentre), PHUSION[®] DNA polymerase (Finnzymes Oy, Vantaa, Finland) and Taq DNA ligase.

800 µl of the 2X assembly master mix, sufficient for 80 reactions, can be prepared by combining the following:

320 µl 5X ISO buffer as prepared above

6.4 µl of 1 U/ µl T5 exo (diluted 1:10 from enzyme stock in 1x T5 exo buffer)

20 µl of 2 U/µl PHUSION[®] polymerase (Finnzymes Oy, Vantaa, Finland)

80 µl of 40 U/µl Taq ligase

374 µl dH₂O

Mix well and store at -20°C, or on ice if to be used immediately.

The assembly mixture can be stored at -20°C for at least one year. The enzymes remain active following at least 10 freeze-thaw cycles. The mixture is ideal for the assembly of DNA molecules with 20-150 bp overlaps.

Oligonucleotides:

[0105] Standard oligonucleotides were purchased at a concentration of 10,000 nM each. The entire HA gene sequence was covered using 52 oligonucleotides (oligos), and the entire NA gene sequence was covered using 44 NA oligonucleotides.

Assembly of Genes

[0106] For the assembly reactions 10 µl of each oligo was pooled, for a concentration of 192 nM per oligo (10,000 nM/52) for HA and 227nM per oligo (10,000 nM/44) for NA. Oligo lengths were on average 60 bases with 30 bp overlaps.

[0107] Once pooled, 10 µl of this oligo mix was added to 10 µl of the 2X assembly master mix prepared as above. Reactions were incubated for 50°C for 1 hour. Following assembly reactions, the gene products were amplified by PCR as follows:

5 µl assembly reaction

20 µl 5X PHUSION® HF Buffer (Finnzymes Oy, Vantaa, Finland)

2 µl 10mM dNTPs

71 µl water

1 µl Hot Start PHUSION® Polymerase (Finnzymes Oy, Vantaa, Finland)

0.5µl 100uM RC-Univ-PKS10-F primer (universal forward primer for cloning vector)

0.5µl 100uM RC-Univ-PKS10-R primer (universal reverse primer for cloning vector)

Cycling reaction was as follows:

98°C for 1minute,

98°C for 10 seconds, 60°C for 30 seconds, 72°C for 1.5 minutes,

Repeated for 24 additional cycles, 72°C for 5 minutes, and then kept at 4°C.

EXAMPLE 2: Cloning of assembled genes into cloning vector.

[0108] In order to clone the assembled gene products into the PKS10 cloning vector, the plasmid was first amplified with primers to create matching overlapping sequences and the termini for a subsequent assembly reaction with the assembled gene product.

Preparation of Cloning Vector

The universal PKS10 cloning vector as amplified by PCR as follows:

20 µl 5X PHUSION[®] HF PCR Buffer

2 µl 10mM dNTPs

75 µl water

1 µl Hot Start PHUSION[®] Polymerase (Finnzymes Oy, Vantaa, Finland)

1 µl 6ng/µl PKS10 plasmid template

0.5 µl 100uM Univ-PKS10-F primer

0.5 µl 100uM Univ-PKS10-R primer

Cycling reaction was as follows:

98°C for 30 seconds,

98°C for 10 seconds, 60°C for 30 seconds, 72°C for 3 minutes,

Repeated for 29 additional cycles, 72°C for 5 minutes, and then kept at 4°C.

[0109] The resulting PCR product was then gel purified using the QIAGEN[®] (Qiagen, GmbH, Hilden, Germany) gel purification kit. The typical yield for this PCR reaction was about 50 ng/µl.

Assembly of Synthetic Genes into Cloning Vector

[0110] PCR products of amplified synthetic genes or error-corrected synthetic genes were gel purified with the QIAGEN[®] gel purification kit, and used for assembly with the gel purified universal PKS10 vector PCR product as follows:

0.3 µl vector

4.7 µl HA or NA

5 µl 2X assembly master mix

[0111] Reactions were incubated for 50°C for 1 hour, and then 20 µl of water was added and mixed with the assembly reaction. This diluted assembly reaction mix was then used to transform *E. coli* by standard electroporation methods (Epicentre Epi300 cells). 1/1000th of the 1 ml SOC outgrowth was plated onto LB Carbenicillin plates to obtain individual colonies.

[0112] A number of cultures of individual colonies were then grown, plasmid DNA was prepared, and then the sequence of the synthetic genes was determined using standard

Sanger sequencing protocols. The percentage of clones containing the desired sequence was determined, and error rates were determined by multiplying the number of clones sequenced by the number of base pairs (bp) of DNA that was synthesized, then dividing this number by the total number of errors.

EXAMPLE 3: Error correction of synthetic HA and NA genes using a mismatch endonuclease together with an exonuclease.

[0113] Assembled HA and NA gene products were subjected to various error correction methods using combinations of an endonuclease enzyme and an exonuclease enzyme to remove inherent mismatches, primarily due to incorrect sequences within oligonucleotides incorporated in the assembled gene products. The result is a simple, high-fidelity, high efficiency gene synthesis of the HA and NA genes. Error rates were determined by multiplying the number of clones sequenced by the number of base pairs (bp) of DNA that was synthesized, then dividing this number by the total number of errors.

A. Two-Step Reactions

[0114] In the two-step reactions, the endonuclease enzyme reaction was performed first, followed by the exonuclease enzyme reaction. Following PCR of the assembled gene product as indicated in Example 1, the following reactions were performed.

[0115] SURVEYOR[®] nuclease/Exonuclease III

8 µl of the PCR product was denatured and annealed as follows:

98°C for 2 minutes, slow cool to 85°C at 2°C/second, hold at 85°C for 2 minutes, slow cool to 25°C at 0.1°C/second, hold at 25°C for 2 minutes, and then hold at 10°C.

2 µl SURVEYOR[®] nuclease (Transgenomic, Inc., Omaha, NE) (a mismatch endonuclease derived from the celery plant that cleaves all types of mismatches) was added and incubated at 42°C for 1 hour.

1 µl Exonuclease III (diluted 1 to 4000 in 1X HF Buffer) was added and incubated at 37°C for 1 hour.

2 µl of this reaction mix was then amplified by PCR as described above.

Optionally, steps 1-3 were repeated to increase fidelity (to further reduce errors in the final gene product).

Results of Error Correction using Two-Step Reactions

[0116] A synthetic HA gene was assembled from oligonucleotides and then subjected to error correction using SURVEYOR[®] nuclease alone, or SURVEYOR[®] nuclease followed by Exonuclease III in the two-step reaction above. The resultant error rates were as follows:

Gene	Error correction Method	Error Rate	Number of correct clones
HA	None	1/ 1,791 bp	11 out of 23 (48%)
HA	SURVEYOR [®] nuclease	1/ 3,710 bp	18 out of 29 (62%)
HA	SURVEYOR [®] nuclease + Exonuclease III	1/ 5,572 bp	21 out of 28 (75%)

[0117] Thus, error correction of the synthetic HA gene increased the number of correct sequences obtained from 48% to 62% using SURVEYOR[®] nuclease treatment alone with an error rate improvement of 1/1,791 bp to 1/3,710 bp; and from 48% to 75% with an error rate improvement of 1/1,791 bp to 1/5,572 bp using the two-step error correction method combining an endonuclease enzyme with an exonuclease enzyme as disclosed.

[0118] In another experiment, both HA and NA synthetic genes were assembled from oligonucleotides and then subjected to error correction using SURVEYOR[®] nuclease alone, or SURVEYOR[®] nuclease followed by Exonuclease III in the two-step reaction above. The resultant error rates were as follows:

Gene	Error correction Method	Error Rate	Number of correct clones
HA	None	1/ 1,635 bp	2 out of 21 (9.5%)
HA	SURVEYOR [®] nuclease	1/ 2,828 bp	16 out of 30 (58.3%)
HA	SURVEYOR [®] nuclease + Exonuclease III	1/ 6,169 bp	22 out of 31 (71%)
NA	None	1/ 1,850 bp	13 out of 28 (46.4%)
NA	SURVEYOR [®] nuclease	1/ 2,480 bp	18 out of 31 (58.1%)
NA	SURVEYOR [®] nuclease + Exonuclease III	1/ 5,314 bp	24 out of 30 (80%)

[0119] Thus, error correction of the synthetic HA gene increased the number of correct sequences obtained from 9.5% to 58.3% using SURVEYOR[®] nuclease treatment alone with an error rate improvement of 1/1635 bp to 1/2828 bp; and from 9.5% to 71% with an error rate improvement of 1/1635 bp to 1/6169 bp using the two-step error correction method combining an endonuclease enzyme with an exonuclease enzyme as disclosed.

[0120] Error correction of the synthetic NA gene increased the number of correct sequences obtained from 46.4% to 58.1% using SURVEYOR[®] nuclease (Transgenomic, Inc., Omaha, NE) treatment alone with an error rate improvement of 1/1850 bp to 1/2480 bp; and from 46.4% to 80% with an error rate improvement of 1/1850 bp to 1/5314 bp using the two-step error correction method combining an endonuclease enzyme with an exonuclease enzyme as disclosed.

Alternative endonucleases + Exonuclease III

[0121] Reactions were performed as in the two-step procedure above but varying the endonuclease and conditions in step 2 as follows:

[0122] T4 endonuclease was substituted for SURVEYOR[®] nuclease, and incubated at 37°C for 1 hour.

[0123] Endonuclease V was substituted for SURVEYOR[®] nuclease, and incubated at 37°C for 1 hour.

Results of Error Correction using two-step reactions with alternative endonucleases

Gene	Error correction Method	Error Rate	Number of correct clones
HA	None	1/ 876 bp	3 out of 23 (13%)
HA	SURVEYOR [®] nuclease + Exonuclease III	1/ 16,716 bp	25 out of 28 (89%)
HA	T4 endonuclease + Exonuclease III	1/ 2,239 bp	11 out of 30 (37%)
HA	Endonuclease V + Exonuclease III	1/ 1,327 bp	3 out of 20 (15%)

[0124] Thus, other alternative nucleases were able to increase the number of correct sequences obtained and improve the error rate to varying degrees in combination with an exonuclease enzyme as disclosed.

B. One-step reactions

[0125] In the one-step reactions, the endonuclease enzyme reaction was performed simultaneously (in the same reaction mixture at the same time) with the exonuclease enzyme reaction. Following PCR of the assembled gene product as indicated in Example 1, the following reactions were performed.

8 μ l of the PCR product was denatured and annealed as follows:

98°C for 2 minutes, slow cool to 85°C at 2°C/second, hold at 85°C for 2 minutes, slow cool to 25°C at 0.1°C/second, hold at 25°C for 2 minutes, and then hold at 10°C.

2 μ l SURVEYOR[®] nuclease (Transgenomic, Inc., Omaha, NE) and 1 μ l Exonuclease III, (diluted 1 to 4000 in 1X HF Buffer) were added and incubated at 42°C for 1 hour.

2 μ l of this reaction mix was then then amplified by PCR as described above.

[0126] Optionally, steps 1 and 2 were repeated to increase fidelity.

Results of Error Correction using one-step reactions

[0127] A synthetic HA gene was assembled from oligonucleotides and then subjected to error correction using SURVEYOR[®] nuclease (Transgenomic, Inc., Omaha, NE) together with Exonuclease III in the one-step reaction above. After the reaction components were added the reaction vessel was not opened again until the reaction had finished. The temperature of the incubation was varied from 4°C to 50°C to determine the optimal reaction conditions for the error correction. The resultant error rates were as follows:

Gene	Temperature (°C)	Error Rate	Number of correct clones
HA	4	1/ 1,357 bp	10 out of 25 (40%)
HA	25	1/ 1,866 bp	11 out of 25 (44%)

HA	30	1/ 6,716 bp	22 out of 30 (73%)
HA	37	1/ 3,582 bp	19 out of 30 (63%)
HA	42	1/ 6,716 bp	23 out of 30 (77%)
HA	50	1/ 5,572 bp	21 out of 28 (75%)

[0128] Thus, the one-step error correction of the synthetic HA gene using the using SURVEYOR[®] nuclease together with Exonuclease III increased the number of correct sequences obtained and the error rates at various temperatures from 30°C to 50°C. One-step error correction methods can be readily performed at 42°C.

EXAMPLE 4: Error correction of synthetic HA and NA genes using an endonuclease enzyme alone or together with an exonuclease enzyme or a polymerase enzyme having exonuclease activity.

[0129] Assembled HA gene products were subjected to various error correction methods using combinations of various endonuclease enzymes alone or together with an exonuclease enzyme or a polymerase enzyme having exonuclease activity to remove inherent mismatches, primarily due to incorrect sequences within oligonucleotides incorporated in the assembled gene products.

[0130] SURVEYOR[®] nuclease (Transgenomic, Inc., Omaha, NE) (cuts 3' to mismatch) was used to perform error correction alone or in a two-step reaction as described above together with PHUSION[®] DNA polymerase (Finnzymes, Oy, Finland), which has 3' to 5' exonuclease activity. Reaction conditions were 42°C for 20 minutes with SURVEYOR[®] nuclease, followed by 37°C for 20 minutes with PHUSION[®] DNA polymerase.

[0131] T7 endonuclease (cuts 5' to mismatch) was used to perform error correction alone or in a two-step reaction as described above together with T5 exonuclease having 5' to 3' exonuclease activity. Reaction conditions were 37°C for 20 minutes with T7 endonuclease, followed by 37°C for 20 minutes with T5 exonuclease.

Results of Error Correction using two-step reactions with alternative endonucleases and alternative exonuclease activities

Gene	Error correction Method	Error Rate	Number of correct clones
HA	None	1/ 791 bp	2 out of 19 (10.5%)
HA	SURVEYOR [®] nuclease	1/ 1,725 bp	8 out of 26 (30.8%)
HA	SURVEYOR [®] nuclease + PHUSION [®] DNA polymerase	1/ 2,217 bp	13 out of 26 (50%)
HA	T7 endonuclease	1/ 973 bp	5 out of 25 (20%)
HA	T7 endonuclease + T5 exonuclease	1/ 1,504 bp	6 out of 21 (28.6%)

[0132] Thus, other alternative nucleases were able to increase the number of correct sequences obtained and improve the error rate to varying degrees in combination with an exonuclease enzyme or another enzyme having exonuclease activity as disclosed.

EXAMPLE 5: Identification and isolation of genes encoding novel mismatch endonucleases.

[0133] Several novel genes encoding novel mismatch endonucleases were identified and isolated. The nucleotide sequences of these genes together with the deduced amino acid sequences are provided in the accompanying Sequence Listing.

[0134] In a BLASTX homology analysis, the nucleotide sequence of each of the novel genes was determined to encode a protein having homology to known mismatch endonucleases. A homology search for the nucleotide sequences of the genes and the deduced amino acid sequences was also conducted using the DDBJ/GenBank/EMBL database. Additionally, sequence identity and similarity were also determined using GENOMEQUEST[™] software (GenomeQuest, Inc., Westborough, MA) (Gene-IT, Worcester, Mass.). As reported in Table 1, the deduced amino acid sequence of each of the genes exhibited high sequence similarity with the amino acid sequences of known mismatch endonucleases CEL I and CEL II isolated from celery and *Selaginella lepidophylla* (US Pat. Nos. 6,391,557; 7,078,211; and 7,560,261).

[0135] TABLE 1. Amino acid sequence homology to known endonucleases was calculated using the AlignX[®] (Life Technologies, Carlsbad, CA) tool of the Vector NTI[®] package (Life Technologies, Carlsbad, CA).

Source Organism	RES I (SEQ ID NO: 02)	CEL I (SEQ ID NO: 04)	CEL II (SEQ ID NO: 06)	CEL II (SEQ ID NO: 08)
Mimulus guttatus (SEQ ID NO: 10)	51%	75%	48%	51%
Solanum tuberosum (SEQ ID NO: 13)	50%	74%	46%	50%
Vitis vinifera (SEQ ID NO: 16)	52%	51%	73%	74%
Vitis vinifera (SEQ ID NO: 23)	50%	50%	68%	72%
Solanum tuberosum (SEQ ID NO: 25)	50%	48%	68%	73%
Medicago sp. (SEQ ID NO: 27)	51%	51%	67%	71%

[0136] FIGURE 4 is an alignment of a *Selaginella lepidophylla* RES I endonuclease (SEQ ID NO: 02), a celery CEL I endonuclease (SEQ ID NO: 04), an *Apium* sp. CEL II endonuclease (SEQ ID NO: 06), another *Apium* sp. CEL II endonuclease (SEQ ID NO: 08), *Mimulus guttatus* CEL I endonuclease (SEQ ID NO: 10), a *Solanum tuberosum* CEL I endonuclease (SEQ ID NO: 13), a *Vitis vinifera* CEL II endonuclease (SEQ ID NO: 16), a *Solanum tuberosum* CEL II endonuclease (SEQ ID NO: 25), a *Medicago* sp. CEL II endonuclease (SEQ ID NO: 27). In this example the sequence alignment of FIGURE 4 was generated using the program AlignX[®] (Life Technologies, Carlsbad, CA) of the Vector NTI Advance[®] (Invitrogen Corp., Carlsbad, CA) 11.5 package (Invitrogen, Carlsbad, Calif.) with default settings. As discussed in detail elsewhere herein, several polypeptide domains and motifs with high degree of conservation have been identified from this sequence comparison analysis. In the alignment figure shown herein, a dash in an aligned sequence

represents a gap, *i.e.*, a lack of an amino acid at that position. Black boxes and gray boxes identify identical amino acids and conserved amino acids, respectively, among aligned sequences.

[0137] In addition, using the program SignalP 4.0, a proteolytic cleavage site was predicted between the A30 and W31 of the full-length polypeptide of *Mimulus guttatus* CEL I endonuclease (SEQ ID NO: 10). As a result, the mature core region of *Mimulus guttatus* CEL I endonuclease, which was predicted to correspond to residues 31 to 306 of the amino acid sequence of SEQ ID NO: 10, was subsequently used for the production of recombinant of *Mimulus guttatus* CEL I endonuclease in insect cells as described in detail below, e.g., Examples 6, 7, and 8.

[0138] Similarly, the mature core region of *Vitis vinifera* CEL II endonuclease, predicted to correspond to residues 25 to 323 of the amino acid sequence of SEQ ID NO: 23, was subsequently used for the production of recombinant *Vitis vinifera* CEL II endonuclease in insect cells as described in Examples 6, 7, and 8 below.

EXAMPLE 6: Construction of recombinant expression cassettes suitable for heterologous enzyme production in recombinant insect cells.

[0139] This Example describes the construction of two recombinant expression cassettes to enable the heterologous expression of the mismatch endonucleases isolated from *Mimulus guttatus* and *Vitis vinifera* in insect cells by utilizing the Bac-to-Bac[®] Baculovirus Expression System (Life Technologies, Inc., Carlsbad, Calif.).

[0140] Two chimeric expression cassettes were designed for the recombinant expression of chimeric polypeptides containing the mature core region of either *Mimulus guttatus* CEL I endonuclease (SEQ ID NO: 14) or *Vitis vinifera* CEL II endonuclease (residues 25 to 323 of SEQ ID NO: 23). Each chimeric polypeptide contained a coding sequence of a mature core region that was operably linked to an N-terminal secretion signal for honeybee melittin (Tesier *et al.*, *Gene* 98, 177-183), and a C-terminal 8X poly-Histidine epitope tag with linkers. The amino acid sequences of the chimeric proteins are disclosed in the Sequence Listing as SEQ ID NO: 11 and SEQ ID NO: 17.

[0141] The amino acid sequences of SEQ ID NO: 11 and SEQ ID NO: 17 were then used to generate expression cassettes having their DNA sequences codon optimized for

expression in insect cells. For this purpose, the codon bias of *Spodoptera frugiperda* was used. The nucleotide sequences of the two codon-optimized expression cassettes are disclosed herein in the Sequence Listing as SEQ ID NO: 31 and SEQ ID NO: 33. Each of the recombinant expression cassettes were subsequently cloned into the expression vector pFastbac1 at the two cloning sites 5' *EcoRI* and 3' *NotI*. The resulting plasmids, which were named Mimmulus-C-His-pFastbac1 and Vitis-C-His-pFastbac1 respectively, were used to infect Sf9 insect cells. P1 baculovirus stocks were generated in Sf9 insect cells using the BAC-TO-BAC® System (Life Technologies, Inc., Carlsbad, CA) according to manufacturer's specifications.

EXAMPLE 7: Preparation of solubilized membrane extracts of insect cells expressing recombinant chimeric endonucleases.

[0142] This Example describes details of the production of the recombinantly-expressed Mimmulus-C-His and Vitis-C-His endonucleases in cultures of insect cells by utilizing the BAC-TO-BAC® Baculovirus Expression System (Life Technologies, Inc., Carlsbad, Calif.). Briefly, P1 virus generation and heterologous production of the recombinant expression cassettes were performed according to the manufacturer's specifications. Cell lysates from P1 virus stock cultures for expression of the recombinant endonucleases were analyzed by Western blot assay using anti-His tag antibody as described below.

Preparation of crude solubilized membrane extracts from insect cell cultures:

[0143] Preparation of membranes: Cell pellet of each insect cell culture was resuspended in IMAC A buffer (20 mM Tris, 500 mM NaCl, 0.0125% Brij-35, 0.01% Triton X-100, 0.005% Tween-20, pH 8.0). The cell suspension was then sonicated (3 × 30 seconds, pulsing) and centrifuged at 18,000 rpm for 60 minutes, using a bench top centrifuge. The supernatant was decanted and the pellet was resuspended in 20 mM Tris-HCl, 150 mM NaCl, 5% glycerol. The protein concentration was quantitated and then diluted down to 10 mg/ml into the final buffer (50 mM Tris-HCl, 300 mM NaCl, 10 µM ZnCl₂, 20% glycerol). The protein extract was aliquoted into 1 ml fractions and snap frozen using liquid nitrogen and stored at -80°C. To monitor cell lysis efficiency, SDS-PAGE gel assays (CRITERION™ Stain-Free precast PAGE system) (BioRad Laboratories,

Inc., Hercules, CA) and Western Blot analysis (anti-HIS epitope antibody) were typically performed on whole-cell and resuspended pellet samples.

[0144] Both recombinant Mimmulus-C-His and Vitis-C-His endonucleases were found soluble in the following solubilization study. Insect cell pellets were resuspended 1:10 in 20 mM Tris-HCl, 150 mM NaCl. The resuspended protein was then broken into 4 equal tubes to create 4 conditions as follows:

- I. 20 mM Tris-HCl, 150 mM NaCl
- II. 20 mM Tris-HCl, 150 mM NaCl, 0.0125% Brij-35, 0.01% Triton X-100, 0.005% Tween-20
- III. 20 mM Tris-HCl, 150 mM NaCl, 8M Urea
- IV. 20 mM Tris-HCl, 150 mM NaCl, 0.0125% Brij-35, 0.01% Triton X-100, 0.005% Tween-20, 8 M Urea

[0145] The four samples above were sonicated (3 x 15 seconds, on ice), and centrifuged at 18,000 rpm for 60 minutes using the bench top Allegra centrifuge. SDS-PAGE gel assays and Western Blot analysis were typically performed on whole-cell and resuspended pellet samples. Primary antibody was monoclonal anti-polyHistidine antibody produced in mouse, dilution: 1:3000; Secondary antibody was goat anti-mouse Peroxidase-conjugated; dilution: 1:20,000. Detection was performed with SUPERSIGNAL™ West Pico Chemiluminescent Substrate (Pierce Chemical Co., Rockford, IL).

Purification of recombinant Mimmulus-C-His endonuclease expressed heterologously in insect cells

[0146] Each of the expression cassettes described in Example 6 contained a nucleotide sequence encoding a secretion signal for honeybee melittin which was operably linked to the nucleic acid sequence encoding the mature core region of the endonuclease. This feature allowed for the recombinant protein to be secreted into the culture media once it was produced in the cytoplasm of the insect cells.

[0147] 1 L of insect cell culture in conditioned media was batch-bound overnight to 5 ml of Ni-SEPHAROSE® 6 FF resin (Pharmacia Fine Chemicals, Piscataway, NJ). Resin was then collected by centrifugation, packed in a 5 ml column, and connected to an AKTA® Explorer (GE Health Care Biosciences, Inc., Uppsala, Sweden). The column was washed

with 10 CV of IMAC Buffer A (20 mM Tris 500 mM NaCl 5 mM Imidazole pH 7.5). Bound protein was eluted with a linear 4% to 100% gradient of IMAC Buffer B over 30 CV (20 mM Tris, 500 mM NaCl, 1 M Imidazole, pH 7.5), collecting 2.5 ml fractions. The following samples were typically analyzed by SDS PAGE CRITERION™ Stain-Free (BioRad Laboratories, Inc., Hercules, CA) and Western blot with an anti-His antibody: (1) Elution fractions, (2) Load, (3) Flow Through (FT), and (4) Wash samples were. Protein containing fractions were pooled and dialyzed against final formulation buffer. (50 mM Tris, 30 mM NaCl, 10 μ M ZnCl₂, 20% glycerol, pH 7.6). Dialyzed pool was filtered (0.45 μ L) and analyzed by SDS PAGE CRITERION™ Stain Free and Western blot with an anti-His antibody. Protein concentration in protein samples was determined by UV spectrophotometry. The pool was dispensed into 1 ml aliquots, snap-frozen using liquid nitrogen and stored at -80°C. Final concentration of MimmulusC-His protein was 0.75 mg/ml. Total amount of protein from 1 L of cell culture was 9 mg. Formulation buffer was as follows: 50 mM Tris-HCl, 300 mM NaCl, 10 μ M ZnCl₂; 20% glycerol, pH 7.6.

[0148] Final concentration of MimmulusC-His protein was 0.75 mg/ml. Total amount of protein from 1 L of cell culture was 9 mg. Formulation buffer was as follows: 50 mM Tris-HCl, 300 mM NaCl, 10 μ M ZnCl₂; 20% glycerol, pH 7.6

[0149] FIGURE 5 depicts SDS polyacrylamide gel analysis of purified MimmulusC-His CEL I protein (FIG. 5A) and Western Blot results using anti-polyHistidine antibody (FIG. 5B). Lane 1: Fermentas Marker (5 μ L); Lane 2: MimmulusC-His Pre-Dialysis (12 μ L); Lane 4: Fermentas Marker (12 μ L); Lane 5: MimmulusC-His Post-Dialysis (12 μ L); Lane 7: Fermentas Marker (5 μ L); Lane 8: MimmulusC-His Post-Dialysis (6 μ L). Primary antibody was monoclonal Anti-polyHistidine antibody produced in mouse, Dilution: 1:3000 ; Secondary antibody was goat anti-mouse Peroxidase-conjugated. Dilution: 1:20,000. Detection was performed with SUPERSIGNAL® West Pico Chemiluminescent Substrate (Pierce Chemical Co., Rockford, IL).

EXAMPLE 8: Error correction of a synthetic gene using purified *Mimulus* C-His chimeric endonuclease enzymes.

[0150] The purified recombinant *Mimulus* C-His chimeric endonuclease isolated as described in Example 7 above was subjected to various two-step error correction assays as described in Example 3, *i.e.*, an endonuclease enzyme reaction was performed first,

followed by an exonuclease enzyme reaction. Error rates were determined by multiplying the number of clones sequenced by the number of base pairs (bp) of DNA that was synthesized, then dividing this number by the total number of errors.

[0151] Briefly, a synthetic NA gene was assembled from oligonucleotides as indicated in Example 1 and then subjected to error correction using unpurified recombinant *Mimulus* C-His endonuclease, or purified recombinant *Mimulus* C-His endonuclease, followed by an exonuclease reaction as described in the two-step reaction above. In this experiment, either T5 exonuclease or Exonuclease III was used in the exonuclease treatment step. The resultant error rates were as follows:

[0152] TABLE 2: Error correction assays performed with unpurified recombinant *Mimulus* CEL I mature core derived from solubilized membrane extracts.

Gene	Error correction Method	Error Rate	Number of correct clones
NA	None	1/ 1,572 bp	12 out of 30 (40%)
NA	Unpurified recombinant <i>Mimulus</i> CEL I mature core	1/ 2,801 bp	23 out of 42 (55%)
NA	Unpurified recombinant <i>Mimulus</i> CEL I mature core + T5 exonuclease	1/ 2,517bp	10 out of 20 (50%)
NA	Unpurified recombinant <i>Mimulus</i> CEL I mature core + Exonuclease III	1/ 10,131 bp	19 out of 23 (83%)

[0153] Thus, error correction of the synthetic HA gene using unpurified recombinant *Mimulus* CEL I treatment alone provided an error rate improvement of 1/ 1,572 bp to 1/ 2,801 bp. Error rates were greatly improved using the two-step error correction method combining an endonuclease enzyme with an exonuclease enzyme as disclosed in TABLE 2. In particular, a combination of unpurified recombinant *Mimulus* CEL I mature core and Exonuclease III increased the number of correct sequences obtained from 40% to 83%, and provided an error rate improvement of 1/ 1,572 bp to 1/ 10,131 bp.

[0154] In another experiment, both unpurified and purified *Mimulus* CEL I endonucleases were tested in two-step error correction assays. In this experiment, an HA synthetic gene was assembled from oligonucleotides and then subjected to error correction using *Mimulus* CEL I endonuclease (42°C, 1 hour), followed by Exonuclease III treatment

(55°C, 1 hour) in a two-step reaction as described in Example 3. The resultant error rates were as follows:

[0155] TABLE 3: Error correction assays performed with recombinant *Mimulus* CEL I mature core purified by Ni-SEPHAROSE® column chromatography (Pharmacia Fine Chemicals, Piscataway, NJ).

Gene	Error correction Method	Error Rate	Number of correct clones
NA	None	1/1,204 bp	8 out of 32 (25%)
NA	Unpurified recombinant <i>Mimulus</i> CEL I mature core + Exonuclease III	1/ 3,081 bp	25 out of 42 (59.5%)
NA	Purified recombinant <i>Mimulus</i> CEL I mature core + Exonuclease III	1/ 13,570 bp	33 out of 37 (89%)

[0156] Thus, error correction of the synthetic HA gene increased the number of correct sequences obtained from 25% to 59.5% using unpurified recombinant *Mimulus* CEL I mature core alone with an error rate improvement of 1/1,204 bp to 1/ 3,081 bp; and from 25% to 89% with an error rate improvement of 1/1,204 bp to 1/ 13,570 bp using the two-step error correction method combining an endonuclease enzyme with Exonuclease III treatment.

SEQUENCES

Exemplary Endonuclease - RES I

US7078211- SEQ ID NO: 01 - Nucleic Acid Sequence

> RES I_US7078211_SEQIDNO_01

ATGGCAACGACCAAGACGAGCGGGATGGCGCTGGCTTTGCTCCTCGTCGCCGCCCTGGCCG
TGGGAGCTGCGGCCTGGGGGAAAGAGGGCCATCGCCTCACTTGTATGGTCGCCGAGCCCTT
TCTAAGCTCTGAATCCAAGCAAGCTGTGGAGGAGCTTCTCTCTGGAAGAGATCTCCCGGAC
TTGTGTTTCATGGGCCGATCAGATTCAAGATCGTATAAGTTTAGATGGACTGGTCCTTTGC
ACTACATCGATACTCCAGACAACCTCTGCACCTATGACTATGATCGTGACTGCCACGATTC
CCATGGGAAGAAGGACGTGTGTGTCGCTGGTGGGATCAACAATTACTCGTCGCAGCTGGAA
ACGTTTCTAGATTCAAGAGCTCGTCGTATAACTTGACCGAGGCGCTGCTCTTCTGGCTC
ACTTTGTGCGGGATATACACCAGCCCTTGCACGTAGCATTTACGAGTGATGCCGAGGCAA

TGGCGTGCACGTCCGCTGGTTTGGACGAAAGGCCAACTTGCATCACGTCTGGGATACAGAA
TTTATTTCTAGAGCCAAATCGTGTGTACTACCACGACATTTCCAAGATGCTCCGGAACATTA
CCAGGAGCATAACTAAGAAGAATTTCAATAGTTGGAGCAGATGTAAGACTGATCCGGCGGC
TTGTATTGATAGTTATGCGACAGAAAGTATAGATGCTTCTTGCAACTGGGCATACAAAGAC
GCACCCGACGGAAGCTCTCTAGATGATGATTACTTCTTTCACGCCTTCCAATTGTTGAGC
AGCGTCTTGCTCAAGGGGGCGTCAGGCTGGCGTCAATACTCAACAGGATTTTTGGAGGAGC
AAAGTCGAACAGGTCCAGTCGCTCAAGCATGTAG

US7078211- SEQ ID NO: 02 - Amino Acid Sequence

>RES I_US7078211_SEQIDNO_02

MATTKTSGMALALLLVAALAVGAAAWGKEGHRLTCMVAEPFLSSESKQAVEELLSGRDLPD
LCSWADQIRRSYKFRWTGPLHYIDTPDNLCTYDYDRDCHDSHGKKDVCVAGGINNYSSQLE
TFLDSESSSYNLTEALLFLAHFVGDIHQPLHVAFTSDAGGNGVHVRWFGRKANLHHVWDTE
FISRANRVYYHDISKMLRNITRSITKKNFNSWSRCKTDPAACIDSYATESIDASCNWAYKD
APDGSSLDDDYFSSRLPIVEQRLAQGGVRLASILNRIFGGAКСNRSSRSSM

Exemplary Endonuclease - CEL I

US6391557- SEQ ID NO: 03 - Nucleic Acid Sequence

>CEL I_US6391557_SEQIDNO_03

TACTCACTATAGGGCTCGAGCGCCCGCCCGGGCAGGTATAATATTAGACTTGTACTCAATG
ACAAGCGCCATCTATGAGTTTCATCATGCCTATATATAAACACATGAACCTGTCATTGTTT
ATTTATGCATTATTGTTGTATTAGCTGAAAAATTTCTGGCAAATGACGCGATTATATTCTG
TGTTCTTTCTTTTGTGGCTCTTGAGTTGAACCGGGTGTTAGAGCCTGGAGCAAAGAAGG
CCATGTCATGACATGTCAAATTGCGCAGGATCTGTTGGAGCCAGAAGCAGCACATGCTGTA
AAGATGCTGTTACCGGACTATGCTAATGGCAACTTATCGTCGCTGTGTGTGTGGCCTGATC
AAATTCGACACTGGTACAAGTACAGGTGGACTAGCTCTCTCCATTTTCATCGATACACCTGA
TCAAGCCTGTTTCAATTTGATTACCAGAGAGACTGTCATGATCCACATGGAGGGAAGGACATG
TGTGTTGCTGGAGCCATTCAAAATTTACATCTCAGCTTGGACATTTCCGCCATGGAACAT
CTGATCGTCGATATAATATGACAGAGGCTTTGTTATTTTTATCCCACTTCATGGGAGATAT
TCATCAGCCTATGCATGTTGGATTTACAAGTGATATGGGAGGAAACAGTATAGATTTGCGC
TGGTTTCGCCACAAATCCAACCTGCACCATGTTTGGGATAGAGAGATTATTCTTACAGCTG

CAGCAGATTACCATGGTAAGGATATGCACTCTCTCCTACAAGACATACAGAGGAACTTTAC
AGAGGGTAGTTGGTTGCAAGATGTTGAATCCTGGAAGGAATGTGATGATATCTCTACTTGC
GCCAATAAGTATGCTAAGGAGAGTATAAACTAGCCTGTAAGTGGGGTTACAAAGATGTTG
AATCTGGCGAAACTCTGTCAGATAAATACTTCAACACAAGAATGCCAATTGTCATGAAACG
GATAGCTCAGGGTGGAATCCGTTTATCCATGATTTTGAACCGAGTTCTTGAAGCTCCGCA
GATCATTCTTTGGCATGAATTTAGATACTGATATTCGCATTTCTCATGACACCCTTCTCTT
ATGCAATTTGCAGATCAGCTGTGATTCACTAATTGAA

US6391557- SEQ ID NO: 04 - Amino Acid Sequence

>CEL I_US6391557_SEQIDNO_04

MTRLYSVFLLLLALVVEPGVRAWSKEGHVMTQCIAQDLLEPEAAHAVKMLLPDYANGNLSS
LCVWPDQIRHWYKYRWTSSLHFIDTPDQACSFYQRDCHDPHGGKDMCVAGAIQNFTSQLG
HFRHGTSDDRYNMTEALLFLSHFMGDIHQPMHVGFTSDMGGNSIDLRFWRHKSNLHHVWDR
EIIILTAAADYHGKDMHSLQLDIQRNFTEGSWLQDVESWKECDDISTCANKYAKESIKLACN
WGYKDVESGETLSDKYFNTRMPIVMKRIAQGGIRLSMILNRVLGSSADHSLA

Exemplary Endonucleases - CEL II

US7560261- SEQ ID NO: 05 - Nucleic Acid Sequence

>CEL II_US7560261_SEQIDNO_05

ATGGGTATGTTGACTTATACTGGAATTTATTTTCTGCTATTACTTCCAAGTGTTTTCTGTT
GGGGAAAACAAGGACATTTTGCAATTTGTAAAATTGCCAGGGGTTCCCTAGTAAAGATGC
ACTGACTGCAGTGAAAGCATTGCTCCCAGAATATGCAGATGGTGATCTAGCAGCTGTTTGC
TCCTGGGCTGACGAGGTTTCGATTTTCATATGCGTTGGAGTAGCCATTACATTATGTGGACA
CGCCTGATTTTCAGGTGTAAGTATAAATACTGTAGAGATTGCCATGATTCTGTTGGACGGAA
AGACCGGTGTGTTACTGGAGCAATTCACAACTACACAGAGCAACTTCTATTGGGTGTTTCAT
GACTTGAATTCAAAAATGAATAACAACCTTGACGGAGGCACCTTATGTTCTTATCACATTTTCG
TTGGTGATGTCCATCAGCCTCTACATGTTGGCTTCCTTGGCGATGAAGGAGGAAACACAAT
CACCGTCCGCTGGTATCGGAGGAAAACCAATTTGCATCATGTATGGGACACAATGATGATT
GAATCCTCCTTGAAGACATTCTACAATTCAGATCTTTCTAGCTTAATACAAGCTATTTCAGA
GCAATATTACAGGTGTCTGGCTTACCGACAGCTTATCTTGGAGCAATTGCACTGCTGATCA
TGTGGTTTGTCCAGACCCGTATGCTTCTGAAAGCATTGAGTTGGCCTGCAAGTTTGCCTAC

AGAAATGCCACACCTGGGACCACTTTAGGAGATGAGTACTTCCTCTCTCGGTTGCCTGTTG
CGGAGAAGAGGTTGGCTCAGGCTGGGGTCCGTTTGGCTGCTACTCTTAACCGAATCTTCAC
TTCAAACCCCAGCGATCTCACAAGATTGAATATGCATAATGGTGGACATAGAAGCAGTAAC
AATATTGAAATAGTGTA

US7560261- SEQ ID NO: 06 - Amino Acid Sequence

>CEL II_US7560261_SEQIDNO_06

MGMLTYTGIYFLLLLPSVFCWGKQGHFAICKIAQGFLSKDALTAVKALLPEYADGDLAAVC
SWADEVRFHMRWSSPLHYVDTPDFRCNYKYCRDCHDSVGRKDRCVTGAIHNYTEQLLLGVH
DLNSKMNNNLTEALMFLSHFVGDVHQPLHVGFLGDEGGNTITVRWYRRKTNLHHVWDTMMI
ESSLKTFYNSDLSSLIQAIQSNITGVWLTDLSLSWSNCTADHVVCPPYASESIELACKFAY
RNATPGTTLGDEYFLSRLPVAEKRLAQAGVRLAATLNRIFTSNPSDLTRLNMHNGGHRSSN
NIEIV

US7560261- SEQ ID NO: 07 - Nucleic Acid Sequence

>CEL II_US7560261_SEQIDNO_07

TGGGGAAAACAAGGACATTTTGCAATTTGTAAAATTGCCCAGGGGTTTCCTTAGTAAAGATG
CACTGACTGCAGTGAAAGCATTGCTCCCAGAATATGCAGATGGTGATCTAGCAGCTGTTTG
CTCCTGGGCTGACGAGGTTGATTTTCATATGCGTTGGAGTAGCCATTACATTATGTGGAC
ACGCCTGATTTCAGGTGTAATACTATAAATACTGTAGAGATTGCCATGATTCTGTTGGACGGA
AAGACCGGTGTGTTACTGGAGCAATTCACAATACTACACAGAGCAACTTCTATTGGGTGTTCA
TGACTTGAATTCAAAAATGAATAACAACCTTGACGGAGGCACTTATGTTCTTATCACATTTT
GTTGGTGATGTCCATCAGCCTCTACATGTTGGCTTCCTTGGCGATGAAGGAGGAAACACAA
TCACCGTCCGCTGGTATCGGAGGAAAACCAATTTGCATCATGTATGGGACACAATGATGAT
TGAATCCTCCTTGAAGACATTCTACAATTCAGATCTTTCTAGCTTAATACAAGCTATTCAG
AGCAATATTACAGGTGTCTGGCTTACCGACAGCTTATCTTGGAGCAATTGCACTGCTGATC
ATGTGGTTTGTCCAGACCCGTATGCTTCTGAAAGCATTGAGTTGGCCTGCAAGTTTGCCTA
CAGAAATGCCACACCTGGGACCACTTTAGGAGATGAGTACTTCCTCTCTCGGTTGCCTGTT
GCGGAGAAGAGGTTGGCTCAGGCTGGGGTCCGTTTGGCTGCTACTCTTAACCGAATCTTCA
CTTCAAACCCCAGCGATCTCACAAGATTGAATATGCATAATGGTGGACATAGAAGCAGTAA
CAATATTGAAATAGTGTA

US7560261- SEQ ID NO: 08 - Amino Acid Sequence**>CEL II_US7560261_SEQIDNO_08**

WGKQGHFAICKIAQGFLSKDALTAVKALLPEYADGDLAAVCSWADEVRFHMRWSSPLHYVD
TPDFRCNYKYCRDCHDSVGRKDRCVTGAIHNYTEQLLLGVHDLNSKMNNNLTEALMFLSHF
VGDVHQPLHVGFLGDEGGNTITVRWYRRKTNLHHVWDTMMIESSLKTFYNSDLSSLIQAIQ
SNITGVWLTDLSWSNCTADHVVCDPYASESIELACKFAYRNATPGTTLGDEYFLSRLPV
AEKRLAQAGVRLAATLNRIFTSNPSDLTRLNMHNGGHRSSNNIEIV

Exemplary Endonucleases - CEL I Variant - *Mimulus guttatus***Nucleic Acid Sequence SEQ ID NO: 09**

ATGCAGATGTCGATTTACGAGGAATTTTTGTTTCTTATTTTGCTTTATTTCTTTGTGTTT
GTGTTGTTTATGAACCTTGTGTCCAGGCATGGAGTAAAGAAGGTCATTCCATGACATGCAA
AATTGCTCAGGATTTGCTGGGACCAGAGGCGAAGCATGCTGTCCAAATGCTGTTACCTGAA
AATGTTAATGGTGATTTATCGGCACCTTAGCGTGTGGCCTGACCAAGTAAGACACTGGTATA
AGTACCGTTGGACGAGCCCTCTTCACTTCATAGACACACCAGATCAAGCCTGTAATTTCAA
TTATCAGAGGGATTGCCATGATCCACATGGTGTTAAGGGTATGTGTGTAGCGGGGGCAATT
CAGAACTTCACCAATCAGCTTTTCGCATTATCGGCACGGAACCTCTGATCGACGCTATAATA
TGACAGAGGCCTTGTTGTTCTTGGCACACTTCATGGGAGATATTCATCAGCCACTGCATGT
TGGATTACGAGTGACGAAGGAGGAAACACTATAGACTTGCGCTGGTTCAGACACAAGTCA
AATCTGCACCATGTATGGGACAGAGAGATAATTCTTACAGCTGCAGCAGATTACTACGGAA
AGGACATTGACCTCCTGCAAGAAGACATTAAGGGAAACTTCACTGATGGAATCTGGTCTGG
TGATCTTGCCCTCTTGGAGGGAATGCAGTGATATATTTTCTTGTGTCAACAAGTATGCTGCT
GAGAGTATAAACATGGCCTGCAAAATGGGGTTACAAAGATGTTAAATCAGGGGACACTCTTT
CAGATGATTACTTTAATTCAAGATTGCCGATTGTTATGAAACGCATAGCTCAGGGTGGAGT
CCGTTTAGCTATGATTTTGAACCGGGTTTTTCGGTGATAGCAAAGAGGATTCCTTAATTGCT
ACTTAA

Amino Acid Sequence SEQ ID NO: 10

MQMSISRGIFVSYFALFLCVCVVYEPVQAWSKEGHSMTCKIAQDLLGPEAKHAVQMLLPE
NVNGDLSALSVWPDQVRHWYKYRWTSPHLFIDTPDQACNFNYQRDCHDPHGKGMCVAGAI
QNFTNQLSHYRHGTSDDRYNMTEALLFLAHFMGDIHQPLHVGFSTDEGGNTIDLRFWRHKS

NLHHVWDREIILTAAADYYGKDIDLLQEDIKGNFTDGIWSGDLASWRECSDFSCVNKYAA
 ESINMACKWGYKDVKSGDTLSDDYFNSRLPIVMKRIAQGGVRLAMILNRVFGDSKEDSLIA
 T

Amino Acid Sequence for insect cell expression SEQ ID NO: 11

MKFLVNVALVFMVVYISYIYAWSKEGHSMTCKIAQDLLGPEAKHAVQMLLPENVNGDLSAL
 SVWPDQVRHWYKYRWTSPLHFIDTPDQACNFNYQRDCHDPHGVKGMCVAGAIQNFTNQLSH
 YRHGTSDDRYNMTEALLFLAHFMGDIHQPLHVGFTSDEGGNTIDLRWFRHKS NLHHVWDRE
 IILTAAADYYGKDIDLLQEDIKGNFTDGIWSGDLASWRECSDFSCVNKYAAESINMACKW
 GYKDVKSGDTLSDDYFNSRLPIVMKRIAQGGVRLAMILNRVFGDSKEDSLIATGSHHHHHH
HHG

Underlined - honeybee melittin secretion signal;
 polyhistidine tag with linkers

Exemplary Endonucleases - CEL I Variant - Solanum tuberosum

Nucleic Acid Sequence SEQ ID NO: 12

ATGTTGAGGTTAACTTCATTAAGCATTATTTTCTTTCTCTGTCTTGCTTTTATCAACCATC
 ATGGTGCTGAAGCATGGAGCAAAGAGGGGCATATGATGACATGTCGCATCGCGCAGGGCTT
 GTTGAATGATGAGGCAGCTCATGCAGTCAAGATGTTGTTGCCGGAATATGTTAACGGCGAC
 TTATCGGCCCTCTGTGTGTGGCCGGATCAAGTCCGGCACTGGTATAAGTATAAATGGACAA
 GCCCTCTACACTTCATTGATACACCAGATAAAGCTTGCAACTTTGATTATGAAAGGGACTG
 TCATGATCAACATGGAGTGAAGGATATGTGTGTTGCTGGTGCAATTCAGAACTTTACTACT
 CAACTCTCTCATTACAGAGAGGGAACCTTCTGATCGTCGATATAATATGACAGAGGCCTTGC
 TGTTCCTTGTCACATTTTATGGGAGATATCCATCAACCAATGCATGTTGGCTTTACAAGTGA
 TGCTGGAGGAAATAGTATTGATTTACGCTGGTTTAGGCATAAATCGAACTTGCACCATGTG
 TGGGATAGGGAGATAATTCTAACAGCTGCTAAAGACTACTATGCAAAGGATGTAAACCTCC
 TTGAAGAAGACATTGAAGGAACTTCACTGACGGAATTTGGTCTGATGATCTTGCTTCTTG
 GAGAGAATGTGGCAATGTCTTTTCTTGTGTAAACAAGTTTGCAACGGAAGTATAAATATA
 GCATGCAAATGGGGATACAAAAGTGTTGAAGCTGGTGAACTTTATCAGATGATTATTTCA
 ATTCAAGACTTCCAATAGTGATGAAACGAGTAGCACAAGGTGGAATACGATTAGCCATGCT
 TTAAACAACGTTTTTGGAGTTTCTCAACAAGAAGATTGAGTTGCTGCAACTTAA

Amino Acid Sequence SEQ ID NO: 13

MLRLTSLSIIFFLCCLAFINHHGAEAWSKEGHMMTCRIAQGLLNDEAAHAVKMLLPEYVNGD
 LSALCVWPDQVRHWYKYKWTSPHLHFIDTPDKACNFDYERDCHDQHGVKDMCVAGAIQNFTT
 QLSHYREGTSDRRYNMTEALLFLSHFMGDIHQPMHVGFSDAGGNSIDLRWFRHKSNLHHV
 WDREIILTAAKDYYAKDVNLLEEDIEGNFTDGIWSDDLASWRECGNVFSCVNKFATESINI
 ACKWGYKSVEAGETLSDDYFNSRLPIVMKRVAQGGIRLAMLNNVFGVSQQEDSVAAT

Exemplary Endonucleases - CEL I Mature Core Sequence**Amino Acid Sequence SEQ ID NO: 14**

WSKEGHSMTCKIAQDLLGPEAKHAVQMLLPENVNGDLSALSVWPDQVRHWYKYRWTSPLHF
 IDTPDQACNFDYERDCHDQPHGVKGMVAGAIQNFTNQLSHYRHGTSDRRYNMTEALLFLAH
 FMGDIHQPLHVGFSDDEGGNTIDLRWFRHKSNLHHVWDREIILTAAADYYGKDIDLLQEDI
 KGNFTDGIWSGDLASWRECSDIFSCVNKYAAESINMACKWGYKDVKSGDTLSDDYFNSRLP
 IVMKRIAQGGVRLAMILNRVFGDSKEDSLIAT

Exemplary Endonucleases - CEL II Variant - Vitis vinifera**Nucleic Acid Sequence SEQ ID NO: 15**

ATGTGGGGAAAGGAAGGACACTATGCAGTTTGTAAAATAGCTGAGGGGTTCTTTCTGAAG
 ATGCATTAGGAGCAGTGAAAGGATTGCTTCCAGATTATGCTGATGGTGATCTGGCTGCCGT
 TTGCTCCTGGGCTGATGAGATTCGTCACTTCCATTGGCGATGGAGTGGCCCTTTACAT
 TATGTAGATACACCAGATTACAGGTGTAATTATGAATACTGCAGAGACTGCCATGACTTCA
 GAGGACACAAAGATATATGTGTAAGTGGAGCAATTTACAACTACACAAAGCAACTCACTTC
 TGGTTATCACAATTCAGGTTCAGAAATAAGATACAATTTGACAGAGGCCCTCATGTTCTTA
 TCAGATTTTATTGGGGATGTCCATCAGCCCTACATGTTGGTTTTACTGGAGATGAAGGTG
 GGAACACAATAATAGTCCGTTGGTACCGGAGGAAGACTAATTTGCATCATATATGGGATGA
 CATGATCATTGATTCCGCCCTGAAGACATATTACAATTCAGATATTGCAATCATGATACAA
 GCCATTCAAAGAAATATTACAGGTGACTGGTCCTTTGATATCTCATCATGGAATAATTGTG
 CATCTGATGATACGGCTTGTCCAAACCTGTATGCGTCTGAAGGCATTAGTTTAGCTTGCAA
 GTTTGCTTACAGAAATGCCACACCAGGAAGCACTCTAGGAGATGATTACTTCTGTCTCGG

CTACCAATTGTGGAGAAGAGGCTAGCCCCGAGTGGGATCCGCCTGGCTGCCACCCTTAACC
GTATCTTTGCTTCTCAAGGCAAGAGAGCTAAAGCATGA

Amino Acid Sequence SEQ ID NO: 16

MWGKEGHYAVCKIAEGFLSEDALGAVKLLPDYADGDAAVCSWADEIRHNFHWRWSGPLH
YVDTDPDYRCNYEYCRDCHDFRGHKDICTGAIYNYTKQLTSGYHNSGSEIRYNLTEALMFL
SDFIGDVHQPLHVGFTGDEGGNTIIVRWYRRKTNLHHIWDDMIIDSALKTYNSDIAIMI
QAIQRNITGDWSFDISSWKNCASDDTACPNLYASEGISLACKFAYRNATPGSTLGDDYFLSR
LPIVEKRLAPSGIRLAATLNRI FASQ GKRAKA

Amino Acid Sequence for insect cell expression SEQ ID NO: 17

MKFLVNVALVFMVVIISYIYAWGKEGHYAVCKIAEGFLSEDALGAVKALLPDYAEGLAAV
CSWADEIRHNFHWRWSGPLHYVDTDPDYRCNYEYCRDCHDFRGHKDICTGAIYNYTKQLTS
GYHNSGSEIRYNLTEALMFLSHFIGDVHQPLHVGFTGDEGGNTIIVRWYRRKTNLHHIWDN
MIIDSALKTYNSDLAIMIQAIQRNITGDWSFDISSWKNCASDDTACPNLYASESISLACK
FAYRNATPGSTLGDDYFLSRLPIVEKRLAQGGIRLAATLNRI FASQ PKISLK HEDKRVEKT
TPVDYIEWSPLOQFSGSHHHHHHHHG

Underlined - honeybee melittin secretion signal;
polyhistidine tag with linkers

Exemplary Endonucleases - CEL II Variant - Chocolate pots

Nucleic Acid Sequence SEQ ID NO: 18

ATGACTTGGGGATTTTGGGCACATCGGCAAATACATCGCCAAGCCGTTTATCTTATGCCTT
CGCCCGTGGCAGAGTTCTTTCGCGCAAATGTTCAAGAACTTGTCGACCGCTCGGTTGAAGC
CGATGAACGCCGACGCATAGACCCCAACGAAGCTCCGCAACACTTCATTGATTTAGACCGC
TACGGTGCCTATCCTTTTGAACAACTTCCGAGAGATTATGAAAAAGCCGTTGAGAAATTGC
GCTATGAGCGGCTGAAAGAAAATGGACTTGTGCCGTGGCGCATTGCCGCCTTTGCCGATAG
CCTCACCAACGCATTTCCGGGAGCAGAACC GCGAAAAAATTTTATACTTCGCCGCAAATTTA
GGGCATTATGTCGCCGATGCTAACGTGCCACTTCATGCCACCGAAAAC TACGACGGACAAC
TCACAGGGCAAAAAGGATTGCACGCACGTTGGGAAACTATTTATCCTCAAAAGTTTATGCT
CCCACGAGAAACCACCTATCTCGAAAACGGGAGCATCTTTATCATTGACAACATCACCGAA

GAAGCCTTCAACTGGTCATTAGAAAAGTTATGTATTGAGCCAACAAGTTTGGCGATTGATA
 AGCAAATTCGAATCGGAATTGTCAGAAGAAGAATTGTATGAGTTAAATTCATCAGACGCGCC
 GCCATTTTCGTCGCGATTTTTCACAACGCTATTATGAAAACTCAAAGAAAAATTGAATCAA
 ATGGTTGAAAAATGCTTTGAGTTAAGCGTCATTAGGGTAGCGTCAGTTTGGTATTTTCTT
 GGTTAAAAGCAGAAAAACCGAATTTATTTAACTTATTAAAAAATTGA

Amino Acid Sequence SEQ ID NO: 19

MTWGFWAHRQIHRQAVYLMPSPVAEFFRANVQELVDRSVEADERRRIDPNEAPQHFIDLDR
 YGAYPFEQLPRDYEKAVEKFGYERLKENGLVPWRIAFAFADSLTNAFREQNREKILYFAANL
 GHYVADANVPLHATENYDGQLTGQKGLHARWETIYPQKFMLPRETTYLENGSIFIIDNITE
 EAFNWSLESYVLSQQVLAIKQIQSELSEEEELYELNSSDAPPFRRDFSQRYYEKLKEKLNQ
 MVEKCFELSVIRVASVWYFSWLKAEKPNLFNLLKN

Exemplary Endonucleases - CEL II Variant - Obsidian pool

Nucleic Acid Sequence SEQ ID NO: 20

ATGTTTTGGGCACATCAAAAAGTCAACGAGCATGCCATTGATTTATTACCCGAGCCACTCC
 GCAGTTTTTATGAACAAAATAAGGAATACATAGTTAAGGAGTCGGTCGCCCCCTGATCTCAG
 GCGTGCAGAAAACAAGGAAGAAGGTTATTATCACTATATGGATCTCGATAAATATGGTGAA
 TATCCGTTCAAGAATTTGCCAGAAAACCTACGACGACGCAGTAAAAAGGTTTGGTTACGATA
 CTGTTCTCAAGAACGGAATTGTGCCGTGGAAGGTAAAATGGTTGACAGACAGTTTGAGTCA
 AGCTATGGAGAGAAAGGATGTGCCACAGGTCTTAAGACTTTCAGCCGACCTTGGTCATTAT
 GTTGCTGACATGCATGTTCCATTTTATTTCGACAGAAAATTATGATGGACAGCTGACAGGCA
 ACATAGGAATACACTTCAGATGGGAAAGCGGCATTCCAGAACATTTTGAACAAATTACAA
 CTATGAGGGAATAGAGCCCGCTGTTTACTTCAAGCATCCTGATAAAAAGGCATTTGAGATA
 CTGACTATGAGTTACAAGTTGATTCTACCTTCTCTCAAGGCTGATAGTCTTGCAAAAGTTG
 GATTGAATGGAAAGAGACTTTATAAAGTTGAGAGAGAAGACGGTAAAAAAGTTTACGTTTA
 TTCAAACGAGTATTATGAGAAGTTCAACAAAAACCTTGGTGGTATTGTAGAATCGCAGATG
 AGGCTGGCAATCCATGATGTTGCAAGCTACTGGTATACTGCATGGGTAAATGCCGGTAAAC
 CAAAGTTTTGGTAA

Amino Acid Sequence SEQ ID NO: 21

MFWAHQKVNEHAIDLLPEPLRSFYEQNKEYIVKESVAPDLRRAENKEEGYYHYMDLDKYGE
 YPFKNLPENYDDAVKRFGYDTVLKNGIVPWVKWLTDSLSQAMERKDVPQVLRLSADLGHY
 VADMHVPPFHSTENYDGQLTGNIGIHFRWESGIPEHFGTNYNYEGIEPAVYFKHPDKKAFEI
 LTMSYKLILPSLKADSLAKVGLNGKRLYKVEREDGKKVYVYSNEYYEKFNKNLGGIVESQM
 RLAIHDVASYWYTAWVNAGKPKFW

Exemplary Endonucleases - CEL II Variant - Vitis vinifera**Nucleic Acid Sequence SEQ ID NO: 22**

ATGGCTTGGTCTGGGGTCTTGTTGATTGTGAGGGCACTTGTTCTTCTGCAATTGATTCCTG
 GAATTCTGAGTTGGGGAAAGGAAGGACACTATGCAGTTTGTAATAAGCTGAGGGGTTCCT
 TTCTGAAGATGCATTAGGAGCAGTGAAAGCATTGCTTCCAGATTATGCTGAAGGTGATCTG
 GCTGCGGTTTGCTCCTGGGCTGATGAGATTTCGTCACAACTTCCATTGGCGATGGAGTGGCC
 CTTTACATTATGTAGATACGCCAGATTACAGGTGTAACCTATGAATACTGCAGAGACTGCCA
 TGACTTCAGAGGACACAAAGATATATGTGTAACCTGGAGCAATTTACAATTACACAAAGCAA
 CTCACTTCTGGTTATCACAATTCAGGTTTCAGAAATAAGATACAATTTGACAGAGGCACTCA
 TGTTCTTATCACATTTTATTGGGGATGTCCATCAGCCCCTACATGTTGGTTTTACTGGAGA
 TGAAGGTGGGAACACAATAATAGTCCGTTGGTACCGGAGGAAGACTAATTTGCATCATATA
 TGGGATAACATGATCATTGATTCCGCCCTGAAGACATATTACAATTCAGATCTTGCAATCA
 TGATACAAGCCATTCAAAGAAATATTACGGGTGATTGGTCCTTTGATATCTCATCATGGAA
 AAATTGTGCATCTGATGATACGGCTTGTCCAAACCTGTATGCTTCTGAAAGCATTAGTTTA
 GCTTGCAAGTTTGCTTACAGAAATGCCACACCAGGAAGCACTCTAGGAGATGATTACTTCC
 TGTCTCGGCTACCAATTGTGGAGAAGAGGCTAGCCCAAGGTGGGATCCGCCTGGCTGCCAC
 CCTTAACCGTATCTTTGCTTCTCAACCAAAAATCTCTCTCAAGCATGAAGATAAAAGGGTA
 GAGAAAACAACCTCCAGTGGATTATATAGAGTGGAGCCCACTGCAACAATTTTCATAA

Amino Acid Sequence SEQ ID NO: 23

MAWSGVLLIVRALVLLQLIPGILSWGKEGHYAVCKIAEGFLSEDALGAVKALLPDYAEGDL
 AAVCSWADEIRHNFHWRWSGPLHYVDTPDYRCNYEYCRDCHDFRGHKDICVTGAIYNYTKQ

LTSGYHNSGSEIRYNLTEALMFLSHFIGDVHQPLHVGFTGDEGGNTIIVRWYRRKTNLHHI
WDNMIIDSALKTYYNSDLAIMIQAIQRNITGDWSFDISSWKNCASDDTACPNLYASESISL
ACKFAYRNATPGSTLGDDYFLSRLPIVEKRLAQGGIRLAATLNRI FASQPKISLKHEDKRV
EKTTTPVDYIEWSPLQQFS

Exemplary Endonucleases - CEL II Variant - Solanum

Nucleic Acid Sequence SEQ ID NO: 24

ATGGGTGGGTTTGAGCTCAAATGGTTTGTAGGAGTAGCTGTTGTTCTGATGATGGTTCAAA
ATATTCTTGGTTGGGGGAAAGAGGGACACTATATTATCTGCAAAATTGCTGAGGAATATCT
AACAGAAGATGCTTTAGCTGCAGTCAAAGCATTACTCCCAGATCAAGCCGAAGGTGATCTT
GCAGCTGTCTGCTCCTGGCCTGATGAGGTTTCGGCGCCACTACCACTACCGCTGGAGCTCTC
CATTACATTATGTAGATACACCTGATTTCTTGTGCAATTACAAATATTGCCGAGACTGCCA
TGACGGGCATGGGCTCAAGGACAGGTGTGTTACGGGAGCAATATACAACTACTCAATGCAA
CTTTCGCAGGGATATTATGATTTGAATTCAGAAAAATACAACCTGACTGAAGCACTTATGT
TCTTGTCTCATTTTGTGTTGGTGACGTACATCAGCCTCTCCATGTTGGTTTCACTGGAGATCT
TGGTGGAACAGTATAATTGTTTCGTTGGTACAGGAGGAAGACTAATTTGCACCATGTATGG
GATAACATGATTATTGAATCTGCGTTGAAGACATACTACAAATCTGATATAATGTTAATGA
CACAAGTTCTTCTGAAAAACATCACTCATGAATGGTCCGATGATGTTCCATCTTGGGAAGA
TTGCAAGGAGATGGTTTGTCTGACCCATATGCTTCTGAAAGTATCCGTTTGGCCTGCAAA
TTTGCCTACAGAAATGCAACCCCGGGAAGCACTTTAACAGACGATTACTTCCTCTCTCGTC
TTCCTGTTGTGGAGAAGAGGTTGGCACAAGGTGGGGTCCGCTTGGCCGAAGTTCTCAACAG
AATTTTCACTAAAAAACCATCAGATGCTGCACAATGA

Amino Acid Sequence SEQ ID NO: 25

MGGFELKWFVGVAVVLMMVQNILGWGKEGHYIICKIAEEYLTEDALAAVKALLPDQAEGL
AAVCSWPDEVRRHYHYRWSSPLHYVDTPDFLCNYKYCRDCHDGHGLKDRCVGTGAIYNYSMQ
LSQGYIDLNSEKYNLTEALMFLSHFVGDVHQPLHVGFTGDLGGNSIIVRWYRRKTNLHHVW
DNMIIESALKTYKSDIMLMTQVLLKNITHEWSDDVPSWEDCKEMVCPDPYASESIRLACK
FAYRNATPGSTLTDDYFLSRLPVVEKRLAQGGVRLAEVLNRIFTKKPSDAAQ

Exemplary Endonucleases - CEL II Variant - Medicago**Nucleic Acid Sequence SEQ ID NO: 26**

ATGATCACGCTCTTAGTTCCGTTGCTGCTATCACTCGCGTTGCCAAATGTTCTGGCTTGGG
GAAAAGATGGTCACTATGCAATTTGTAAAATTTACAGGAGTATCTTAGTGAAGATGCTCT
ATTTGCAGTCAAACAATTACTTCCAGATTCTGCTCAAGCTGATCTTGCTTCAGTTTGCTCT
TGGCCTGATGAGATTCGCCATAATTACCATTATCGTTGGAGTAGTCCTTTACATTATATTG
ATACACCAGATTTCAAATGTAAGTATCAATATTGCAGAGACTGTCATGATTCTTATGGACA
TAAGCATAGATGCGTTACTGGAGCAATATACAATTATACAATGCAATTAAAATTAGCTAAC
GCCGATGCTTCATCTGAATTAAAATATAACTTGACAGAGGCACTTATGTTCTTGTCACATT
TTGTTGGAGATGTTTCATCAGCCCCTACATGTTGGTTTTACTGGAGACCTAGGTGGAAACTC
AATAACAGTTCGTTGGTACAGGAGGAAAACAAATCTTCATCACGTATGGGATAACATGATT
ATTGAGTCTGCTCTGAAAAAGTTCTATGGTTCAGATCTTTCAACTATGATACAGGCTATTC
AAAGGAATATTAGTGATATTTGGTCAAATGATGTATCTATTTGGGAACATTGTGCACACAA
CCACACAGCATGTCCAGACCGGTATGCTTCTGAGAGTATTAGCTTGGCATGCAAGTTTGCG
TATAAGAATGCTACACCGGGAAGCACTTTGGAAGATGACTACTTCCTTTCTCGGTTGCCTA
TTGTGGAGAAAAGGCTGGCTCAAGGTGGTGTGCGACTTGCAGCTATCCTCAACCACATTTT
CACTCCGAAGACCAGAATAGCTCAAGCTTAA

Amino Acid Sequence SEQ ID NO: 27

MITLLVPLLLSLALPNVLAWGKDGHYAICKISQEYLSEDALFAVKQLLPDSAQADLASVCS
WPDEIRHNYHYRWSSPLHYIDTPDFKCNYQYCRDCHDSYGHKHRCVTGAIYNYTMQLKLAN
ADASSELKYNLTEALMFLSHFVGDVHQPLHVGFTGDLGGNSITVRWYRRKTNLHHVWDNMI
IESALKKFYGS DLSTMIQAIQRNISDIWSNDVSIWEHCAHNHTACPDRYASESISLACKFA
YKNATPGSTLEDDYFLSRLPIVEKRLAQGGVRLAAILNHIFTPKTRIAQA

Exemplary Endonucleases – CEL II Variant Mature Core Sequence**Amino Acid Sequence SEQ ID NO: 28**

WGKEGHYAVCKIAEGFLSEDALGAVKGLLPDYADGDLAAVCSWADEIRHNFHWRWSGPLHY
 VDTPDYRCNIEYCRDCHDFRGHKDICTGAIYNYTKQLTSGYHNSGSEIRYNLTEALMFLS
 DFIGDVHQPLHVGFTGDEGGNTIIVRWYRRKTNLHHIWDDMIIDSALKTYYNSDIAIMIQA
 IQRNITGDWSFDISSWKNCASDDTACPNLYASEGISLACKFAYRNATPGSTLGDDYFLSRL
 PIVEKRLAPSGIRLAATLNRIFASQ GK

Exemplary Endonucleases – CEL II Variant Mature Core Sequence**Amino Acid Sequence SEQ ID NO: 29**

WGKEGHYAVCKIAEGFLSEDALGAVKALLPDYAEGDLAAVCSWADEIRHNFHWRWSGPLHY
 VDTPDYRCNIEYCRDCHDFRGHKDICTGAIYNYTKQLTSGYHNSGSEIRYNLTEALMFLS
 HFIGDVHQPLHVGFTGDEGGNTIIVRWYRRKTNLHHIWDNMIIDSALKTYYNSDLAIMIQA
 IQRNITGDWSFDISSWKNCASDDTACPNLYASESISLACKFAYRNATPGSTLGDDYFLSRL
 PIVEKRLAQGGIRLAATLNRIFASQPK

Codon-Optimized Mature Core Region of *Mimulus guttatus* CEL I**Nucleic acid Sequence SEQ ID NO: 30**

TGGAGTAAGGAGGGACATAGCATGACATGTAAGATAGCCCAGGACTTGTGTTGGGTCCCGAAG
 CCAAACACGCCGTGCAAATGTTGTTGCCTGAAAATGTGAACGGCGACCTAAGCGCCTTGTC
 GGTGTGGCCGGACCAAGTGAGACACTGGTACAAATACAGATGGACCTCCCCTTTGCACTTC
 ATTGACACCCCCGATCAGGCTTGCAACTTTAACTACCAGAGAGACTGCCATGACCCGCACG
 GTGTAAAAGGCATGTGCGTTGCCGGTGCCATTCAAAATTTACGAACCAATTGTCGCACTA
 CAGACACGGCACGTGCGACAGACGTTACAACATGACGGAGGCCTTGTTGTTTTTGGCCCCAC
 TTTATGGGCGATATTCATCAGCCGTTGCACGTGGGCTTCACGTCAGACGAAGGCGGCAACA
 CGATTGACTTGAGATGGTTTCGCCACAAGAGCAACTTGCATCACGTATGGGATCGAGAAAT
 TATCCTAACTGCCGCTGCGGACTACTACGGAAAGGACATCGACCTACTCCAGGAGGATATC
 AAAGGCAATTTTACTGACGGCATCTGGTCGGGCGATTTGGCCTCGTGGAGAGAATGTTCCG
 ACATTTTTTTCGTGTGTGAACAAGTACGCTGCCGAATCCATAAACATGGCTTGTAATGGGG

CTACAAGGATGTGAAATCGGGTGACACGCTCTCGGACGACTATTTCAACAGTCGTCTCCCG
ATCGTAATGAAAAGAATCGCTCAAGGAGGCGTTTCGCTTAGCAATGATTCTCAACAGAGTAT
TCGGTGATAGCAAAGAGGACAGCTTGATTGCCACG

Codon-Optimized Expression Cassette for Insect Cell

Expression of *Mimulus guttatus* CEL I

Nucleic acid Sequence SEQ ID NO: 31

ACCATGAAGTTCTTGGTCAACGTAGCACTGGTTTTTATGGTAGTCTATATCAGCTACATTT
ACGCGTGGAGTAAGGAGGGACATAGCATGACATGTAAGATAGCCCAGGACTTGTTGGGTCC
CGAAGCCAAACACGCCGTGCAAATGTTGTTGCCTGAAAATGTGAACGGCGACCTAAGCGCC
TTGTTCGGTGTGGCCGGACCAAGTGAGACACTGGTACAAATACAGATGGACCTCCCCCTTTGC
ACTTCATTGACACCCCCGATCAGGCTTGCAACTTTAACTACCAGAGAGACTGCCATGACCC
GCACGGTGTAAAAGGCATGTGCGTTGCCGGTGCCATTCAAATTTACGAACCAATTGTTCG
CACTACAGACACGGCACGTTCGGACAGACGTTACAACATGACGGAGGCCTTGTTGTTTTTGG
CCCCTTTATGGGCGATATTCATCAGCCGTTGCACGTGGGCTTCACGTCAGACGAAGGCGG
CAACACGATTGACTTGAGATGGTTTCGCCACAAGAGCAACTTGCATCACGTATGGGATCGA
GAAATTATCCTAACTGCCGCTGCGGACTACTACGGAAAGGACATCGACCTACTCCAGGAGG
ATATCAAAGGCAATTTTACTGACGGCATCTGGTCGGGCGATTTGGCCTCGTGGAGAGAATG
TTCGGACATTTTTTTCGTGTGTGAACAAGTACGCTGCCGAATCCATAAACATGGCTTGTA
TGGGGCTACAAGGATGTGAAATCGGGTGACACGCTCTCGGACGACTATTTCAACAGTCGTC
TCCCGATCGTAATGAAAAGAATCGCTCAAGGAGGCGTTTCGCTTAGCAATGATTCTCAACAG
AGTATTCGGTGATAGCAAAGAGGACAGCTTGATTGCCACGGGCTCGCACCATCACCACCAT
CACCACCACGGTTGATAA

Codon-Optimized Mature Core Region of *Vitis vinifera* CEL II

Nucleic acid Sequence SEQ ID NO: 32

TGGGGCAAAGAAGGCCACTACGCCGTGTGTAAGATTGCGGAGGGCTTTTTGTTCGGAAGACG
CATTGGGAGCGGTCAAAGCCTTGTTGCCGGACTACGCGGAAGGCGACTTGGCAGCCGTATG
TAGCTGGGCCGACGAGATCAGACACAACCTTTCACTGGAGATGGTCGGGCCCCACTGCATTAC
GTCGACACGCCGGATTACAGATGCAACTACGAGTACTGCCGCGACTGTCACGACTTCAGAG

GCCACAAAGACATTTGCGTCACGGGCGCGATATACAACCTACACGAAACAATTGACGTGGG
 CTACCACAACAGTGGCTCCGAGATTCGATAACAACCTCACGGAGGCCTTGATGTTCTCTCG
 CATTTTCATTGGCGACGTGCACCAACCGCTGCATGTGGGCTTTACGGGCGATGAAGGCGGAA
 ATACGATCATTGTCCGTTGGTACCGCAGAAAGACCAACCTCCACCACATATGGGACAACAT
 GATCATCGACTCGGCGTTGAAGACCTACTACAACAGCGACCTGGCCATAATGATCCAGGCG
 ATTCAAAGAAACATCACCGGCGATTGGTCCTTTGACATCAGCAGCTGGAAGAAGTGTGCCA
 GTGACGACACTGCTTGTCCGAACCTATACGCGTCGGAGAGCATCTCGTTGGCCTGTAAATT
 TGCCTACAGAAATGCCACCCCCGGTTCGACGCTGGGCGACGACTACTTCTTGTGCGGATTG
 CCGATTGTTGAAAAACGCCTCGCCCAAGGCGGTATTAGATTGGCCGCCACCTTGAACCGTA
 TTTTTCCTCGCAACCGAAAATCTCGCTGAAACACGAAGACAAGAGAGTCGAGAAGACGAC
 GCCGGTAGACTACATCGAGTGGTCGCCATTGCAACAGTTCAGC

Codon-Optimized Expression Cassette for Insect Cell

Expression of *Mimulus guttatus* CEL I

Nucleic acid Sequence SEQ ID NO: 33

ACCATGAAGTTCTTGGTGAACGTGGCGCTGGTGTTCATGGTCGTGTACATCTCCTACATTT
 ACGCGTGGGGCAAAGAAGGCCACTACGCCGTGTGTAAGATTGCGGAGGGCTTTTTGTTCGA
 AGACGCATTGGGAGCGGTCAAAGCCTTGTTGCCGGAAGGCGACTTGGCAGCC
 GTATGTAGCTGGGCCGACGAGATCAGACACAACCTTTCCTGGAGATGGTCGGGCCCCACTGC
 ATTACGTCGACACGCCGGATTACAGATGCAACTACGAGTACTGCCGCGACTGTCACGACTT
 CAGAGGCCACAAAGACATTTGCGTCACGGGCGCGATATACAACCTACACGAAACAATTGACG
 TCGGGCTACCACAACAGTGGCTCCGAGATTCGATAACAACCTCACGGAGGCCTTGATGTTCC
 TCTCGCATTTTCATTGGCGACGTGCACCAACCGCTGCATGTGGGCTTTACGGGCGATGAAGG
 CGGAAATACGATCATTGTCCGTTGGTACCGCAGAAAGACCAACCTCCACCACATATGGGAC
 AACATGATCATCGACTCGGCGTTGAAGACCTACTACAACAGCGACCTGGCCATAATGATCC
 AGGCGATTCAAAGAAACATCACCGGCGATTGGTCCTTTGACATCAGCAGCTGGAAGAAGT
 TGCCAGTGACGACACTGCTTGTCCGAACCTATACGCGTCGGAGAGCATCTCGTTGGCCTGT
 AAATTTGCCTACAGAAATGCCACCCCCGGTTCGACGCTGGGCGACGACTACTTCTTGTGCG
 GATTGCCGATTGTTGAAAAACGCCTCGCCCAAGGCGGTATTAGATTGGCCGCCACCTTGAA
 CCGTATTTTTGCCTCGCAACCGAAAATCTCGCTGAAACACGAAGACAAGAGAGTCGAGAAG

ACGACGCCGGTAGACTACATCGAGTGGTCGCCATTGCAACAGTTCAGCGGAAGCCACCACC
ATCACCACCATCATCACGGCTGATAA

CLAIMS

What is claimed is:

1. A method for error correction of nucleic acid molecules, said method comprising:
 - (a) obtaining a first plurality of double-stranded nucleic acid molecules comprising at least one nucleotide mismatch;
 - (b) fragmenting said plurality of double-stranded nucleic acid molecules having a mismatch by reacting said nucleic acid molecules having a mismatch with at least one molecule having unidirectional mismatch endonuclease activity;
 - (c) removing said nucleotide mismatch by reacting said fragmented double-stranded nucleic acid molecules having a mismatch of (b) with at least one molecule having unidirectional exonuclease activity of the same directionality as the unidirectional mismatch endonuclease activity of (b) to provide a fragmented error-free double-stranded nucleic acid molecule; and
 - (d) assembling a second plurality of double-stranded nucleic acid molecules comprising said fragmented error-free double-stranded nucleic acid molecule of (c), wherein said second plurality of double-stranded nucleic acid molecules has a decreased frequency of nucleotide mismatches as compared to said first plurality of double-stranded nucleic acid molecules.
2. A method according to claim 1, wherein said first plurality of nucleotide acid molecules comprises one or more synthetic nucleotide sequences.
3. A method according to claim 1, wherein said first plurality of nucleotide acid molecules comprises a mixture of one or more naturally occurring gene sequences and one or more synthetic nucleotide sequences.
4. A method according to claim 1, wherein obtaining a first plurality of nucleic acid molecules comprises synthesizing the nucleic acid molecules.

5. A method according to claim 1, wherein obtaining a first plurality of nucleic acid molecules comprises assembling the nucleic acid molecules from subsets and/or oligonucleotides.
6. A method according to claim 1, wherein step (b) and step (c) are performed as separate reactions.
7. A method according to claim 1, wherein step (b) and step (c) are performed as a one-step, simultaneous reaction.
8. A method according to claim 1, wherein said unidirectional mismatch endonuclease activity cuts 5' to said mismatch and said unidirectional exonuclease activity removes said nucleotide mismatch from the 5' end of said fragmented nucleic acid molecule.
9. A method according to claim 1, wherein said unidirectional mismatch endonuclease activity cuts 3' to said mismatch and said unidirectional exonuclease activity removes said nucleotide mismatch from the 3' end of said fragmented nucleic acid molecule.
10. A method according to claim 1, wherein said at least one molecule having unidirectional mismatch endonuclease activity is selected from the group consisting of RES I, CEL I, CEL II, an SP endonuclease, SP I, T7 endonuclease, T4 endonuclease, endonuclease V, a Mut protein, a variant of any thereof, and a combination of any two or more of the above.
11. A method according to claim 10, wherein said at least one molecule having unidirectional mismatch endonuclease activity is selected from the group consisting of: CEL I, CEL II, a variant of any thereof, and a combination of any two or more of the above.
12. A method according to claim 1, wherein said at least one molecule having unidirectional mismatch endonuclease activity is encoded by a nucleic acid sequence selected from the group consisting of:

- a) a nucleic acid sequence hybridizing under low, moderate, or high stringency conditions to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 01, SEQ ID NO: 03, SEQ ID NO: 05, SEQ ID NO: 07, SEQ ID NO: 09, SEQ ID NO: 12, SEQ ID NO: 15, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, a complement of any, and a fragment of any;
 - b) a nucleic acid sequence exhibiting 70% or greater identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 01, SEQ ID NO: 03, SEQ ID NO: 05, SEQ ID NO: 07, SEQ ID NO: 09, SEQ ID NO: 12, SEQ ID NO: 15, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, a complement of any, and a fragment of any; and
 - c) a nucleic acid sequence encoding a polypeptide exhibiting 60% or greater identity to an amino acid sequence selected from the group consisting of SEQ ID NO: 02, SEQ ID NO: 04, SEQ ID NO: 06, SEQ ID NO: 08, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 28, and SEQ ID NO: 29.
13. A method according to claim 1, wherein said at least one molecule having unidirectional exonuclease activity is selected from the group consisting of exonuclease III, a DNA polymerase, lambda exonuclease, T7 exonuclease, T5 exonuclease, and a variant of any thereof.
 14. A method according to claim 1, wherein said at least one molecule having unidirectional exonuclease activity is a polymerase with proofreading activity.
 15. A method according to claim 14, wherein said polymerase with proofreading activity is selected from the group consisting of T4 polymerase, T7 polymerase, and phi29 polymerase.
 16. A method according to claim 1, wherein

said at least one molecule having unidirectional mismatch endonuclease activity is selected from the group consisting of CEL I, CEL II, a variant of any thereof, and a combination of any two or more of the above; and

said at least one molecule having unidirectional exonuclease activity is selected from the group consisting of exonuclease III and a variant thereof.

17. An isolated nucleic acid molecule comprising:
 - a) a nucleic acid sequence hybridizing under low, moderate, or high stringency conditions to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 09, SEQ ID NO: 12, SEQ ID NO: 15, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, a complement thereof or a fragment of either; or
 - b) a nucleic acid sequence exhibiting 70% or greater identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 09, SEQ ID NO: 12, SEQ ID NO: 15, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, a complement thereof or a fragment of either; or
 - d) a nucleic acid sequence encoding a polypeptide exhibiting 50% or greater identity to an amino acid sequence selected from the group consisting of SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 28, and SEQ ID NO: 29.
18. A nucleic acid molecule according to claim 17, wherein said nucleic acid sequence encodes a molecule having mismatch endonuclease activity.
19. A recombinant nucleic acid construct comprising a nucleic acid molecule according to claim 17 operably linked to a heterologous nucleic acid.
20. A recombinant nucleic acid construct according to claim 19, wherein said heterologous nucleic acid is a heterologous transcriptional control element.

21. A recombinant nucleic acid construct according to claim 19, wherein said heterologous nucleic acid comprises a nucleic acid sequence encoding a polypeptide sequence.
22. A recombinant nucleic acid construct according to claim 21, wherein said polypeptide sequence comprises a secretion signal or an epitope tag.
23. A recombinant host cell comprising a nucleic acid construct according to claim 19.
24. A recombinant host cell according to claim 23, wherein said host cell is an insect cell, a mammalian cell, a microbial cell, or a plant cell.
25. An isolated polypeptide, wherein said polypeptide is expressed by a nucleic acid molecule comprising a nucleic acid sequence according to claim 17 introduced into a host cell.
26. An isolated polypeptide according to claim 25, wherein said polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11, amino acid residues 1 to 297 of SEQ ID NO: 11, amino acid residues 22 to 308 of SEQ ID NO: 11, SEQ ID NO: 17, amino acid residues 1 to 320 of SEQ ID NO: 17, and amino acid residues 22 to 331 of SEQ ID NO: 17.
27. A composition comprising: (i) a molecule having a unidirectional mismatch endonuclease activity; and (ii) a molecule having unidirectional exonuclease activity of the same directionality as the unidirectional mismatch endonuclease activity of (i).
28. A composition according to claim 27, wherein the molecule of (i) is selected from the group consisting of RES I, CEL I, CEL II, T7 endonuclease, T4 endonuclease, endonuclease V, a Mut protein, a variant of any thereof, and a combination of any two or more of the above; and the molecule of (ii) is selected from the group consisting of exonuclease III, a DNA polymerase, a variant of any thereof, and a combination of any two or more of the above.

29. A composition according to claim 27, wherein the molecule of (i) is selected from the group consisting of CEL I, CEL II, a variant of any thereof, and a combination of any two or more of the above; and the molecule of (ii) is selected from the group consisting of exonuclease III and a variant thereof.
30. A composition according to claim 27, wherein the molecule of (i) is selected from the group consisting of RES I, CEL I, CEL II, T7 endonuclease, T4 endonuclease, endonuclease V, a Mut protein, a variant of any thereof, and a combination of any two or more of the above; and the molecule of (ii) is exonuclease III or a variant thereof.
31. A kit comprising a composition according to claim 27.
32. The kit of claim 31 wherein the molecule having unidirectional mismatch endonuclease activity is selected from the group consisting of: RES I, CEL I, CEL II, T7 endonuclease, T4 endonuclease, endonuclease V, a Mut protein, a variant of any thereof, and a combination of any two or more of the above; and the molecule having unidirectional exonuclease activity of the same directionality as the unidirectional mismatch endonuclease is selected from the group consisting of: exonuclease III, a DNA polymerase, a variant of any thereof, and a combination of any two or more of the above.
33. The kit of claim 32 wherein
the molecule having unidirectional exonuclease activity of the same directionality as the unidirectional mismatch endonuclease activity is selected from the group consisting of: RES I, CEL I, CEL II, T7 endonuclease, T4 endonuclease, endonuclease V, a Mut protein, a variant of any thereof, and a combination of any two or more of the above; and
the molecule having unidirectional exonuclease activity of the same directionality as the unidirectional mismatch endonuclease is selected from the group consisting of: exonuclease III or a variant thereof.

SEQUENCE LISTING

<110> SYNTHETIC GENOMICS, INC.
 GIBSON, Daniel
 CAIAZZA, Nicky
 RICHARDSON, Toby

<120> MATERIALS AND METHODS FOR THE SYNTHESIS OF ERROR-MINIMIZED
 NUCLEIC ACID MOLECULES

<130> SGI1550-1WO

<140> PCT/US2013/24496

<141> 2013-02-01

<150> US 61/593,813

<151> 2012-02-01

<160> 33

<170> PatentIn version 3.5

<210> 1

<211> 888

<212> DNA

<213> Selaginella lepidophylla

<400> 1

atggcaacga ccaagacgag cgggatggcg ctggctttgc tcctcgtcgc cgccctggcc	60
gtgggagctg cggcctgggg gaaagagggc catcgctca cttgtatggt cgccgagccc	120
tttctaagct ctgaatccaa gcaagctgtg gaggagcttc tctctggaag agatctccc	180
gacttgtgtt catgggccga tcagattcga agatcgtata agtttagatg gactggctct	240
ttgcactaca tcgatactcc agacaacctc tgcacctatg actatgatcg tgactgccac	300
gattcccatg ggaagaagga cgtgtgtgtc gctggtggga tcaacaatta ctcgctgcag	360
ctggaaacgt ttctagattc agagagctcg tcgtataact tgaccgaggc gctgctcttc	420
ctggctcact ttgtcgggga tatacaccag cccttgacg tagcatttac gagtgatgcc	480
ggaggcaatg gcgtgcacgt ccgctggttt ggacgaaagg ccaacttgca tcacgtctgg	540
gatacagaat ttatttctag agccaatcgt gtgtactacc acgacatttc caagatgctc	600
cggaacatta ccaggagcat aactaagaag aatttcaata gttggagcag atgtaagact	660
gatccggcgg cttgtattga tagttatgcg acagaaagta tagatgcttc ttgcaactgg	720
gcatacaaag acgcacccga cggaagctct ctagatgatg attacttctc ttcaagcctt	780
ccaattgttg agcagcgtct tgctcaaggg ggcgtcaggc tggcgtcaat actcaacagg	840
atTTTTTggag gagcaaagtc gaacaggtcc agtcgctcaa gcatgtag	888

<210> 2

<211> 295

<212> PRT

<213> Selaginella lepidophylla

<400> 2

Met Ala Thr Thr Lys Thr Ser Gly Met Ala Leu Ala Leu Leu Val
 1 5 10 15

Ala Ala Leu Ala Val Gly Ala Ala Ala Trp Gly Lys Glu Gly His Arg
 20 25 30

Leu Thr Cys Met Val Ala Glu Pro Phe Leu Ser Ser Glu Ser Lys Gln
 35 40 45

Ala Val Glu Glu Leu Leu Ser Gly Arg Asp Leu Pro Asp Leu Cys Ser
 50 55 60

Trp Ala Asp Gln Ile Arg Arg Ser Tyr Lys Phe Arg Trp Thr Gly Pro
 65 70 75 80

Leu His Tyr Ile Asp Thr Pro Asp Asn Leu Cys Thr Tyr Asp Tyr Asp
 85 90 95

Arg Asp Cys His Asp Ser His Gly Lys Lys Asp Val Cys Val Ala Gly
 100 105 110

Gly Ile Asn Asn Tyr Ser Ser Gln Leu Glu Thr Phe Leu Asp Ser Glu
 115 120 125

Ser Ser Ser Tyr Asn Leu Thr Glu Ala Leu Leu Phe Leu Ala His Phe
 130 135 140

Val Gly Asp Ile His Gln Pro Leu His Val Ala Phe Thr Ser Asp Ala
 145 150 155 160

Gly Gly Asn Gly Val His Val Arg Trp Phe Gly Arg Lys Ala Asn Leu
 165 170 175

His His Val Trp Asp Thr Glu Phe Ile Ser Arg Ala Asn Arg Val Tyr
 180 185 190

Tyr His Asp Ile Ser Lys Met Leu Arg Asn Ile Thr Arg Ser Ile Thr
 195 200 205

Lys Lys Asn Phe Asn Ser Trp Ser Arg Cys Lys Thr Asp Pro Ala Ala
 210 215 220

Cys Ile Asp Ser Tyr Ala Thr Glu Ser Ile Asp Ala Ser Cys Asn Trp
 225 230 235 240

Ala Tyr Lys Asp Ala Pro Asp Gly Ser Ser Leu Asp Asp Asp Tyr Phe
 245 250 255

Ser Ser Arg Leu Pro Ile Val Glu Gln Arg Leu Ala Gln Gly Gly Val
 260 265 270

Arg Leu Ala Ser Ile Leu Asn Arg Ile Phe Gly Gly Ala Lys Ser Asn
 275 280 285

Arg Ser Ser Arg Ser Ser Met
 290 295

<210> 3
 <211> 1135
 <212> DNA
 <213> Celery

<400> 3
 tactcactat agggctcgag cgcccgcccg ggcaggtata atattagact tgtactcaat 60
 gacaagcgcc atctatgagt ttcatcatgc ctatatataa acacatgaac ctgtcattgt 120
 tcatttatgc attattgttg tattagctga aaaatttctg gcaaattgacg cgattatatt 180
 ctgtgttctt tcttttgttg gctctttag ttgaaccggg tgtagagcc tggagcaaag 240
 aaggccatgt catgacatgt caaattgcgc aggatctgtt ggagccagaa gcagcacatg 300
 ctgtaaagat gctgttaccg gactatgcta atggcaactt atcgtcgctg tgtgtgtggc 360
 ctgatcaaat tcgacactgg tacaagtaca ggtggactag ctctctccat ttcatcgata 420
 cacctgatca agcctgttca tttgattacc agagagactg tcatgatcca catggaggga 480
 aggacatgtg tggtgctgga gccattcaaa atttcacatc tcagcttgga catttccgcc 540
 atggaacatc tgatcgtcga tataatatga cagaggcttt gttattttta tcccacttca 600
 tgggagatat tcatcagcct atgcatgttg gatttacaag tgatatggga ggaaacagta 660
 tagatttgcg ctggtttcgc cacaatcca acctgcacca tgtttgggat agagagatta 720
 ttcttacagc tgcagcagat taccatggta aggatatgca ctctctccta caagacatac 780
 agaggaactt tacagagggt agttggttgc aagatgttga atcctggaag gaatgtgatg 840
 atatctctac ttgcgccaat aagtatgcta aggagagtat aaaactagcc tgtaactggg 900
 gttacaaaga tggtgaatct ggcgaaactc tgtcagataa atacttcaac acaagaatgc 960
 caattgtcat gaaacggata gtcaggggtg gaatccgttt atccatgatt ttgaaccgag 1020
 ttcttggaag ctccgcagat cattctttgg catgaattta gatactgata ttgcgatttc 1080

tcattgacacc cttctcttat gcaatttgca gatcagctgt gattcactaa ttgaa

1135

<210> 4
 <211> 296
 <212> PRT
 <213> Celery

<400> 4

Met Thr Arg Leu Tyr Ser Val Phe Phe Leu Leu Leu Ala Leu Val Val
 1 5 10 15

Glu Pro Gly Val Arg Ala Trp Ser Lys Glu Gly His Val Met Thr Cys
 20 25 30

Gln Ile Ala Gln Asp Leu Leu Glu Pro Glu Ala Ala His Ala Val Lys
 35 40 45

Met Leu Leu Pro Asp Tyr Ala Asn Gly Asn Leu Ser Ser Leu Cys Val
 50 55 60

Trp Pro Asp Gln Ile Arg His Trp Tyr Lys Tyr Arg Trp Thr Ser Ser
 65 70 75 80

Leu His Phe Ile Asp Thr Pro Asp Gln Ala Cys Ser Phe Asp Tyr Gln
 85 90 95

Arg Asp Cys His Asp Pro His Gly Gly Lys Asp Met Cys Val Ala Gly
 100 105 110

Ala Ile Gln Asn Phe Thr Ser Gln Leu Gly His Phe Arg His Gly Thr
 115 120 125

Ser Asp Arg Arg Tyr Asn Met Thr Glu Ala Leu Leu Phe Leu Ser His
 130 135 140

Phe Met Gly Asp Ile His Gln Pro Met His Val Gly Phe Thr Ser Asp
 145 150 155 160

Met Gly Gly Asn Ser Ile Asp Leu Arg Trp Phe Arg His Lys Ser Asn
 165 170 175

Leu His His Val Trp Asp Arg Glu Ile Ile Leu Thr Ala Ala Ala Asp
 180 185 190

Tyr His Gly Lys Asp Met His Ser Leu Leu Gln Asp Ile Gln Arg Asn
 195 200 205

Phe Thr Glu Gly Ser Trp Leu Gln Asp Val Glu Ser Trp Lys Glu Cys
 210 215 220

Asp Asp Ile Ser Thr Cys Ala Asn Lys Tyr Ala Lys Glu Ser Ile Lys
 225 230 235 240

Leu Ala Cys Asn Trp Gly Tyr Lys Asp Val Glu Ser Gly Glu Thr Leu
 245 250 255

Ser Asp Lys Tyr Phe Asn Thr Arg Met Pro Ile Val Met Lys Arg Ile
 260 265 270

Ala Gln Gly Gly Ile Arg Leu Ser Met Ile Leu Asn Arg Val Leu Gly
 275 280 285

Ser Ser Ala Asp His Ser Leu Ala
 290 295

<210> 5
 <211> 933
 <212> DNA
 <213> *Apium* sp.

<400> 5
 atgggtatgt tgacttatac tggaatttat tttctgctat tacttccaag tgttttctgt 60
 tggggaaaac aaggacattt tgcaatttgt aaaattgccc aggggttcct tagtaaagat 120
 gcactgactg cagtgaagc attgctcca gaatatgcag atggtgatct agcagctgtt 180
 tgctcctggg ctgacgaggt tcgatttcatt atgcgttgga gtagccatt acattatgtg 240
 gacacgcctg atttcaggtg taactataaa tactgtagag attgccatga ttctgttgga 300
 cggaaagacc ggtgtgttac tggagcaatt cacaactaca cagagcaact tctattgggt 360
 gttcatgact tgaattcaaa aatgaataac aacttgacgg aggcacttat gttcttatca 420
 catttcgttg gtgatgtcca tcagcctcta catgttggtt tccttggcga tgaaggagga 480
 aacacaatca ccgtccgctg gtatcggagg aaaaccaatt tgcattcatgt atgggacaca 540
 atgatgattg aatcctcctt gaagacattc tacaattcag atctttctag cttaatacaa 600
 gctattcaga gcaatattac aggtgtcttg cttaccgaca gcttatcttg gagcaattgc 660
 actgctgata atgtggtttg tccagaccgg tatgcttctg aaagcattga gttggcctgc 720
 aagtttgctt acagaaatgc cacacctggg accactttag gagatgagta ctctctctct 780
 cggttgcctg ttgcggagaa gaggttggtt caggctgggg tccgtttggc tgctactctt 840
 aaccgaatct tcacttcaaa cccagcgat ctcaagat tgaatatgca taatggtgga 900

catagaagca gtaacaatat tgaaatagtg taa

933

<210> 6
 <211> 310
 <212> PRT
 <213> Apium sp.

<400> 6

Met Gly Met Leu Thr Tyr Thr Gly Ile Tyr Phe Leu Leu Leu Leu Pro
 1 5 10 15

Ser Val Phe Cys Trp Gly Lys Gln Gly His Phe Ala Ile Cys Lys Ile
 20 25 30

Ala Gln Gly Phe Leu Ser Lys Asp Ala Leu Thr Ala Val Lys Ala Leu
 35 40 45

Leu Pro Glu Tyr Ala Asp Gly Asp Leu Ala Ala Val Cys Ser Trp Ala
 50 55 60

Asp Glu Val Arg Phe His Met Arg Trp Ser Ser Pro Leu His Tyr Val
 65 70 75 80

Asp Thr Pro Asp Phe Arg Cys Asn Tyr Lys Tyr Cys Arg Asp Cys His
 85 90 95

Asp Ser Val Gly Arg Lys Asp Arg Cys Val Thr Gly Ala Ile His Asn
 100 105 110

Tyr Thr Glu Gln Leu Leu Leu Gly Val His Asp Leu Asn Ser Lys Met
 115 120 125

Asn Asn Asn Leu Thr Glu Ala Leu Met Phe Leu Ser His Phe Val Gly
 130 135 140

Asp Val His Gln Pro Leu His Val Gly Phe Leu Gly Asp Glu Gly Gly
 145 150 155 160

Asn Thr Ile Thr Val Arg Trp Tyr Arg Arg Lys Thr Asn Leu His His
 165 170 175

Val Trp Asp Thr Met Met Ile Glu Ser Ser Leu Lys Thr Phe Tyr Asn
 180 185 190

Ser Asp Leu Ser Ser Leu Ile Gln Ala Ile Gln Ser Asn Ile Thr Gly
 195 200 205

Val Trp Leu Thr Asp Ser Leu Ser Trp Ser Asn Cys Thr Ala Asp His
210 215 220

Val Val Cys Pro Asp Pro Tyr Ala Ser Glu Ser Ile Glu Leu Ala Cys
225 230 235 240

Lys Phe Ala Tyr Arg Asn Ala Thr Pro Gly Thr Thr Leu Gly Asp Glu
245 250 255

Tyr Phe Leu Ser Arg Leu Pro Val Ala Glu Lys Arg Leu Ala Gln Ala
260 265 270

Gly Val Arg Leu Ala Ala Thr Leu Asn Arg Ile Phe Thr Ser Asn Pro
275 280 285

Ser Asp Leu Thr Arg Leu Asn Met His Asn Gly Gly His Arg Ser Ser
290 295 300

Asn Asn Ile Glu Ile Val
305 310

<210> 7
<211> 873
<212> DNA
<213> Apium sp.

<400> 7
tggggaaaac aaggacattt tgcaatttgt aaaattgccc aggggttcct tagtaaagat 60
gcactgactg cagtgaaagc attgctccca gaatatgcag atggtgatct agcagctgtt 120
tgctcctggg ctgacgaggt tcgatttcat atgcgttgga gtagccatt acattatgtg 180
gacacgcctg atttcaggtg taactataaa tactgtagag attgccatga ttctgttgga 240
cggaaagacc ggtgtgttac tggagcaatt cacaactaca cagagcaact tctattgggt 300
gttcatgact tgaattcaaa aatgaataac aacttgacgg aggcacttat gttcttatca 360
catttcgttg gtgatgtcca tcagcctcta catgttggct tccttggcga tgaaggagga 420
aacacaatca ccgtccgctg gtatcgagg aaaaccaatt tgcattcatgt atgggacaca 480
atgatgattg aatcctcctt gaagacattc tacaattcag atctttctag cttataataca 540
gctattcaga gcaatattac aggtgtctgg cttaccgaca gcttatcttg gagcaattgc 600
actgctgata atgtggtttg tccagaccog tatgcttctg aaagcattga gttggcctgc 660
aagtttgctt acagaaatgc cacacctggg accactttag gagatgagta cttcctctct 720
cggttgcttg ttgcggagaa gaggttggct caggctgggg tccgtttggc tgctactctt 780

aaccgaatct tcacttcaaa cccagcgat ctcaagaat tgaatatgca taatggtgga 840

catagaagca gtaacaatat tgaaatagtg taa 873

<210> 8

<211> 290

<212> PRT

<213> Apium sp.

<400> 8

Trp Gly Lys Gln Gly His Phe Ala Ile Cys Lys Ile Ala Gln Gly Phe
1 5 10 15

Leu Ser Lys Asp Ala Leu Thr Ala Val Lys Ala Leu Leu Pro Glu Tyr
20 25 30

Ala Asp Gly Asp Leu Ala Ala Val Cys Ser Trp Ala Asp Glu Val Arg
35 40 45

Phe His Met Arg Trp Ser Ser Pro Leu His Tyr Val Asp Thr Pro Asp
50 55 60

Phe Arg Cys Asn Tyr Lys Tyr Cys Arg Asp Cys His Asp Ser Val Gly
65 70 75 80

Arg Lys Asp Arg Cys Val Thr Gly Ala Ile His Asn Tyr Thr Glu Gln
85 90 95

Leu Leu Leu Gly Val His Asp Leu Asn Ser Lys Met Asn Asn Asn Leu
100 105 110

Thr Glu Ala Leu Met Phe Leu Ser His Phe Val Gly Asp Val His Gln
115 120 125

Pro Leu His Val Gly Phe Leu Gly Asp Glu Gly Gly Asn Thr Ile Thr
130 135 140

Val Arg Trp Tyr Arg Arg Lys Thr Asn Leu His His Val Trp Asp Thr
145 150 155 160

Met Met Ile Glu Ser Ser Leu Lys Thr Phe Tyr Asn Ser Asp Leu Ser
165 170 175

Ser Leu Ile Gln Ala Ile Gln Ser Asn Ile Thr Gly Val Trp Leu Thr
180 185 190

Asp Ser Leu Ser Trp Ser Asn Cys Thr Ala Asp His Val Val Cys Pro
 195 200 205

Asp Pro Tyr Ala Ser Glu Ser Ile Glu Leu Ala Cys Lys Phe Ala Tyr
 210 215 220

Arg Asn Ala Thr Pro Gly Thr Thr Leu Gly Asp Glu Tyr Phe Leu Ser
 225 230 235 240

Arg Leu Pro Val Ala Glu Lys Arg Leu Ala Gln Ala Gly Val Arg Leu
 245 250 255

Ala Ala Thr Leu Asn Arg Ile Phe Thr Ser Asn Pro Ser Asp Leu Thr
 260 265 270

Arg Leu Asn Met His Asn Gly Gly His Arg Ser Ser Asn Asn Ile Glu
 275 280 285

Ile Val
 290

<210> 9
 <211> 921
 <212> DNA
 <213> Mimmulus guttatus

<400> 9
 atgcagatgt cgatttcacg aggaattttt gtttcttatt ttgctttatt tctttgtgtt 60
 tgtgtttgttt atgaaccttg tgtccaggca tggagtaaag aaggtcattc catgacatgc 120
 aaaattgctc aggatttgct gggaccagag gcgaagcatg ctgtccaaat gctgttacct 180
 gaaaatgtta atggtgattt atcggcactt agcgtgtggc ctgaccaagt aagacactgg 240
 tataagtacc gttggacgag ccctcttcac ttcatagaca caccagatca agcctgtaat 300
 ttcaattatc agagggattg ccatgatcca catgggtgta agggatatgtg tgtagcgggg 360
 gcaattcaga acttcaccaa tcagctttcg cattatcggc acggaacctc tgatcgacgc 420
 tataatatga cagaggcctt gttgttcttg gcacacttca tgggagatat tcatcagcca 480
 ctgcatgttg gattcacgag tgacgaagga ggaaacacta tagacttgcg ctggttcaga 540
 cacaagtcaa atctgcacca tgtatgggac agagagataa ttcttacagc tgcagcagat 600
 tactacggaa aggacattga cctcctgcaa gaagacatta agggaaactt cactgatgga 660
 atctgggtctg gtgatcttgc ctcttgaggg gaatgcagtg atatattttc ttgtgtcaac 720
 aagtatgctg ctgagagtat aaacatggcc tgcaaattggg gttacaaaga tgttaaataca 780
 ggggacactc tttcagatga ttactttaat tcaagattgc cgattgttat gaaacgcata 840

gctcaggggtg gagtccggtt agctatgatt ttgaaccggg ttttcggtga tagcaaagag 900
gattccttaa ttgctactta a 921

<210> 10
<211> 306
<212> PRT
<213> Mimulus guttatus

<400> 10

Met Gln Met Ser Ile Ser Arg Gly Ile Phe Val Ser Tyr Phe Ala Leu
1 5 10 15

Phe Leu Cys Val Cys Val Val Tyr Glu Pro Cys Val Gln Ala Trp Ser
20 25 30

Lys Glu Gly His Ser Met Thr Cys Lys Ile Ala Gln Asp Leu Leu Gly
35 40 45

Pro Glu Ala Lys His Ala Val Gln Met Leu Leu Pro Glu Asn Val Asn
50 55 60

Gly Asp Leu Ser Ala Leu Ser Val Trp Pro Asp Gln Val Arg His Trp
65 70 75 80

Tyr Lys Tyr Arg Trp Thr Ser Pro Leu His Phe Ile Asp Thr Pro Asp
85 90 95

Gln Ala Cys Asn Phe Asn Tyr Gln Arg Asp Cys His Asp Pro His Gly
100 105 110

Val Lys Gly Met Cys Val Ala Gly Ala Ile Gln Asn Phe Thr Asn Gln
115 120 125

Leu Ser His Tyr Arg His Gly Thr Ser Asp Arg Arg Tyr Asn Met Thr
130 135 140

Glu Ala Leu Leu Phe Leu Ala His Phe Met Gly Asp Ile His Gln Pro
145 150 155 160

Leu His Val Gly Phe Thr Ser Asp Glu Gly Gly Asn Thr Ile Asp Leu
165 170 175

Arg Trp Phe Arg His Lys Ser Asn Leu His His Val Trp Asp Arg Glu
180 185 190

Ile Ile Leu Thr Ala Ala Ala Asp Tyr Tyr Gly Lys Asp Ile Asp Leu
195 200 205

Leu Gln Glu Asp Ile Lys Gly Asn Phe Thr Asp Gly Ile Trp Ser Gly
210 215 220

Asp Leu Ala Ser Trp Arg Glu Cys Ser Asp Ile Phe Ser Cys Val Asn
225 230 235 240

Lys Tyr Ala Ala Glu Ser Ile Asn Met Ala Cys Lys Trp Gly Tyr Lys
245 250 255

Asp Val Lys Ser Gly Asp Thr Leu Ser Asp Asp Tyr Phe Asn Ser Arg
260 265 270

Leu Pro Ile Val Met Lys Arg Ile Ala Gln Gly Gly Val Arg Leu Ala
275 280 285

Met Ile Leu Asn Arg Val Phe Gly Asp Ser Lys Glu Asp Ser Leu Ile
290 295 300

Ala Thr
305

<210> 11
<211> 308
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic construct: Insect Cell Expression

<400> 11

Met Lys Phe Leu Val Asn Val Ala Leu Val Phe Met Val Val Tyr Ile
1 5 10 15

Ser Tyr Ile Tyr Ala Trp Ser Lys Glu Gly His Ser Met Thr Cys Lys
20 25 30

Ile Ala Gln Asp Leu Leu Gly Pro Glu Ala Lys His Ala Val Gln Met
35 40 45

Leu Leu Pro Glu Asn Val Asn Gly Asp Leu Ser Ala Leu Ser Val Trp
50 55 60

Pro Asp Gln Val Arg His Trp Tyr Lys Tyr Arg Trp Thr Ser Pro Leu
65 70 75 80

His Phe Ile Asp Thr Pro Asp Gln Ala Cys Asn Phe Asn Tyr Gln Arg
85 90 95

Asp Cys His Asp Pro His Gly Val Lys Gly Met Cys Val Ala Gly Ala
100 105 110

Ile Gln Asn Phe Thr Asn Gln Leu Ser His Tyr Arg His Gly Thr Ser
115 120 125

Asp Arg Arg Tyr Asn Met Thr Glu Ala Leu Leu Phe Leu Ala His Phe
130 135 140

Met Gly Asp Ile His Gln Pro Leu His Val Gly Phe Thr Ser Asp Glu
145 150 155 160

Gly Gly Asn Thr Ile Asp Leu Arg Trp Phe Arg His Lys Ser Asn Leu
165 170 175

His His Val Trp Asp Arg Glu Ile Ile Leu Thr Ala Ala Ala Asp Tyr
180 185 190

Tyr Gly Lys Asp Ile Asp Leu Leu Gln Glu Asp Ile Lys Gly Asn Phe
195 200 205

Thr Asp Gly Ile Trp Ser Gly Asp Leu Ala Ser Trp Arg Glu Cys Ser
210 215 220

Asp Ile Phe Ser Cys Val Asn Lys Tyr Ala Ala Glu Ser Ile Asn Met
225 230 235 240

Ala Cys Lys Trp Gly Tyr Lys Asp Val Lys Ser Gly Asp Thr Leu Ser
245 250 255

Asp Asp Tyr Phe Asn Ser Arg Leu Pro Ile Val Met Lys Arg Ile Ala
260 265 270

Gln Gly Gly Val Arg Leu Ala Met Ile Leu Asn Arg Val Phe Gly Asp
275 280 285

Ser Lys Glu Asp Ser Leu Ile Ala Thr Gly Ser His His His His His
290 295 300

His His His Gly
305

<210> 12
 <211> 909
 <212> DNA
 <213> Solanum tuberosum

<400> 12
 atgttgaggt taacttcatt aagcattatt ttctttctct gtcttgcttt tatcaaccat 60
 catggtgctg aagcatggag caaagagggg catatgatga catgtcgcat cgcgcagggc 120
 ttgttgaatg atgaggcagc tcatgcagtc aagatgttgt tgccggaata tgttaacggc 180
 gacttatcgg ccctctgtgt gtggccggat caagtccggc actggtataa gtataaatgg 240
 acaagccctc tacacttcatt tgatacacca gataaagctt gcaactttga ttatgaaagg 300
 gactgtcatg atcaacatgg agtgaaggat atgtgtgttg ctggtgcaat tcagaacttt 360
 actactcaac tctctcatta cagagagggg acttctgata gtcgatataa tatgacagag 420
 gccttgctgt tcttgtcaca ttttatggga gatatccatc aaccaatgca tgttggcttt 480
 acaagtgatg ctggaggaaa tagtattgat ttacgctggg ttaggcataa atcgaacttg 540
 caccatgtgt gggataggga gataattcta acagctgcta aagactacta tgcaaaggat 600
 gtaaaccctc ttgaagaaga cattgaagga aacttcactg acggaatttg gtctgatgat 660
 cttgcttctt ggagagaatg tggcaatgtc ttttcttggt taaacaagtt tgcaacggaa 720
 agtataaata tagcatgcaa atggggatac aaaagtgttg aagctggtga aactttatca 780
 gatgattatt tcaattcaag acttccaata gtgatgaaac gagtagcaca aggtggaata 840
 cgattagcca tgctttttaa caacgttttt ggagtttctc aacaagaaga ttcagttgct 900
 gcaacttaa 909

<210> 13
 <211> 302
 <212> PRT
 <213> Solanum tuberosum

<400> 13

Met Leu Arg Leu Thr Ser Leu Ser Ile Ile Phe Phe Leu Cys Leu Ala
 1 5 10 15

Phe Ile Asn His His Gly Ala Glu Ala Trp Ser Lys Glu Gly His Met
 20 25 30

Met Thr Cys Arg Ile Ala Gln Gly Leu Leu Asn Asp Glu Ala Ala His
 35 40 45

Ala Val Lys Met Leu Leu Pro Glu Tyr Val Asn Gly Asp Leu Ser Ala
 50 55 60

Leu Cys Val Trp Pro Asp Gln Val Arg His Trp Tyr Lys Tyr Lys Trp
65 70 75 80

Thr Ser Pro Leu His Phe Ile Asp Thr Pro Asp Lys Ala Cys Asn Phe
85 90 95

Asp Tyr Glu Arg Asp Cys His Asp Gln His Gly Val Lys Asp Met Cys
100 105 110

Val Ala Gly Ala Ile Gln Asn Phe Thr Thr Gln Leu Ser His Tyr Arg
115 120 125

Glu Gly Thr Ser Asp Arg Arg Tyr Asn Met Thr Glu Ala Leu Leu Phe
130 135 140

Leu Ser His Phe Met Gly Asp Ile His Gln Pro Met His Val Gly Phe
145 150 155 160

Thr Ser Asp Ala Gly Gly Asn Ser Ile Asp Leu Arg Trp Phe Arg His
165 170 175

Lys Ser Asn Leu His His Val Trp Asp Arg Glu Ile Ile Leu Thr Ala
180 185 190

Ala Lys Asp Tyr Tyr Ala Lys Asp Val Asn Leu Leu Glu Glu Asp Ile
195 200 205

Glu Gly Asn Phe Thr Asp Gly Ile Trp Ser Asp Asp Leu Ala Ser Trp
210 215 220

Arg Glu Cys Gly Asn Val Phe Ser Cys Val Asn Lys Phe Ala Thr Glu
225 230 235 240

Ser Ile Asn Ile Ala Cys Lys Trp Gly Tyr Lys Ser Val Glu Ala Gly
245 250 255

Glu Thr Leu Ser Asp Asp Tyr Phe Asn Ser Arg Leu Pro Ile Val Met
260 265 270

Lys Arg Val Ala Gln Gly Gly Ile Arg Leu Ala Met Leu Leu Asn Asn
275 280 285

Val Phe Gly Val Ser Gln Gln Glu Asp Ser Val Ala Ala Thr
290 295 300

<210> 14
 <211> 276
 <212> PRT
 <213> Mimmulus guttatus

<400> 14

Trp Ser Lys Glu Gly His Ser Met Thr Cys Lys Ile Ala Gln Asp Leu
 1 5 10 15

Leu Gly Pro Glu Ala Lys His Ala Val Gln Met Leu Leu Pro Glu Asn
 20 25 30

Val Asn Gly Asp Leu Ser Ala Leu Ser Val Trp Pro Asp Gln Val Arg
 35 40 45

His Trp Tyr Lys Tyr Arg Trp Thr Ser Pro Leu His Phe Ile Asp Thr
 50 55 60

Pro Asp Gln Ala Cys Asn Phe Asn Tyr Gln Arg Asp Cys His Asp Pro
 65 70 75 80

His Gly Val Lys Gly Met Cys Val Ala Gly Ala Ile Gln Asn Phe Thr
 85 90 95

Asn Gln Leu Ser His Tyr Arg His Gly Thr Ser Asp Arg Arg Tyr Asn
 100 105 110

Met Thr Glu Ala Leu Leu Phe Leu Ala His Phe Met Gly Asp Ile His
 115 120 125

Gln Pro Leu His Val Gly Phe Thr Ser Asp Glu Gly Gly Asn Thr Ile
 130 135 140

Asp Leu Arg Trp Phe Arg His Lys Ser Asn Leu His His Val Trp Asp
 145 150 155 160

Arg Glu Ile Ile Leu Thr Ala Ala Ala Asp Tyr Tyr Gly Lys Asp Ile
 165 170 175

Asp Leu Leu Gln Glu Asp Ile Lys Gly Asn Phe Thr Asp Gly Ile Trp
 180 185 190

Ser Gly Asp Leu Ala Ser Trp Arg Glu Cys Ser Asp Ile Phe Ser Cys
 195 200 205

Val Asn Lys Tyr Ala Ala Glu Ser Ile Asn Met Ala Cys Lys Trp Gly
 210 215 220

Tyr Lys Asp Val Lys Ser Gly Asp Thr Leu Ser Asp Asp Tyr Phe Asn
225 230 235 240

Ser Arg Leu Pro Ile Val Met Lys Arg Ile Ala Gln Gly Gly Val Arg
245 250 255

Leu Ala Met Ile Leu Asn Arg Val Phe Gly Asp Ser Lys Glu Asp Ser
260 265 270

Leu Ile Ala Thr
275

<210> 15
<211> 831
<212> DNA
<213> Vitis vinifera

<400> 15
atgtggggaa aggaaggaca ctatgcagtt tgtaaaatag ctgaggggtt cctttctgaa 60
gatgcattag gagcagtgaaggattgctt ccagattatg ctgatggtga tctggctgcc 120
gtttgctcct gggctgatga gattcgctcac aacttcatt ggcatggag tggcccttta 180
cattatgtag atacaccaga ttacaggtgt aattatgaat actgcagaga ctgccatgac 240
ttcagaggac acaagatat atgtgtaact ggagcaattt acaactacac aaagcaactc 300
acttctggtt atcacaattc aggttcagaa ataagataca atttgacaga ggccctcatg 360
ttcttatcag attttattgg ggatgtccat cagcccctac atgttggtt tactggagat 420
gaaggtggga acacaataat agtccgttgg taccggagga agactaattt gcatcatata 480
tgggatgaca tgatcattga ttccgccttg aagacatatt acaattcaga tattgcaatc 540
atgatacaag ccattcaaag aaatattaca ggtgactggt cctttgatat ctcatcatgg 600
aaaaattgtg catctgatga tacggcttgt ccaaacctgt atgogtctga aggcattagt 660
ttagcttgca agtttgctta cagaaatgcc acaccaggaa gcaacttagg agatgattac 720
ttcctgtctc ggctaccaat tgtggagaag aggctagccc cgagtgggat ccgcctggct 780
gccaccctta accgtatctt tgcttctcaa ggcaagagag ctaaagcatg a 831

<210> 16
<211> 276
<212> PRT
<213> Vitis vinifera

<400> 16

Met Trp Gly Lys Glu Gly His Tyr Ala Val Cys Lys Ile Ala Glu Gly

1	5	10	15
Phe	Leu	Ser	Glu
	20	Asp	Ala
Leu	Gly	Ala	Val
	25	Lys	Gly
Leu	Leu	Pro	Asp
	30		
Tyr	Ala	Asp	Gly
	35	Asp	Leu
Ala	Ala	Val	Cys
	40	Ser	Trp
Ala	Asp	Glu	Ile
	45		
Arg	His	Asn	Phe
	50	His	Trp
Arg	Trp	Ser	Gly
	55	Pro	Leu
His	Tyr	Val	Asp
	60		
Thr	Pro	Asp	Tyr
	65	Arg	Cys
Asn	Tyr	Glu	Tyr
	70	Cys	Arg
Asp	Cys	His	Asp
	75		80
Phe	Arg	Gly	His
	85	Lys	Asp
Ile	Cys	Val	Thr
	90	Gly	Ala
Ile	Tyr	Asn	Tyr
	95		
Thr	Lys	Gln	Leu
	100	Thr	Ser
Gly	Tyr	His	Asn
	105	Ser	Gly
Ser	Glu	Ile	Arg
	110		
Tyr	Asn	Leu	Thr
	115	Glu	Ala
Leu	Met	Phe	Leu
	120	Ser	Asp
Phe	Ile	Gly	Asp
	125		
Val	His	Gln	Pro
	130	Leu	His
Val	Gly	Phe	Thr
	135	Gly	Asp
Glu	Gly	Gly	Asn
	140		
Thr	Ile	Ile	Val
	145	Arg	Trp
Tyr	Arg	Arg	Lys
	150	Thr	Asn
Leu	His	His	Ile
	155		160
Trp	Asp	Asp	Met
	165	Ile	Ile
Asp	Ser	Ala	Leu
	170	Lys	Thr
Tyr	Tyr	Tyr	Asn
	175		Ser
Asp	Ile	Ala	Ile
	180	Met	Ile
Gln	Ala	Ile	Gln
	185	Arg	Asn
Ile	Thr	Gly	Asp
	190		
Trp	Ser	Phe	Asp
	195	Ile	Ser
Ser	Ser	Trp	Lys
	200	Asn	Cys
Ala	Ser	Asp	Asp
	205		Thr
Ala	Cys	Pro	Asn
	210	Leu	Tyr
Ala	Ser	Glu	Gly
	215	Ile	Ser
Leu	Ala	Cys	Lys
	220		
Phe	Ala	Tyr	Arg
	225	Asn	Ala
Thr	Pro	Gly	Ser
	230	Thr	Leu
Gly	Asp	Asp	Tyr
	235		240
Phe	Leu	Ser	Arg
	245	Leu	Pro
Ile	Val	Glu	Lys
	250	Arg	Leu
Ala	Pro	Ser	Gly
	255		

Ile Arg Leu Ala Ala Thr Leu Asn Arg Ile Phe Ala Ser Gln Gly Lys
 260 265 270

Arg Ala Lys Ala
 275

<210> 17
 <211> 331
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic construct: Insect Cell Expression

<400> 17

Met Lys Phe Leu Val Asn Val Ala Leu Val Phe Met Val Val Tyr Ile
 1 5 10 15

Ser Tyr Ile Tyr Ala Trp Gly Lys Glu Gly His Tyr Ala Val Cys Lys
 20 25 30

Ile Ala Glu Gly Phe Leu Ser Glu Asp Ala Leu Gly Ala Val Lys Ala
 35 40 45

Leu Leu Pro Asp Tyr Ala Glu Gly Asp Leu Ala Ala Val Cys Ser Trp
 50 55 60

Ala Asp Glu Ile Arg His Asn Phe His Trp Arg Trp Ser Gly Pro Leu
 65 70 75 80

His Tyr Val Asp Thr Pro Asp Tyr Arg Cys Asn Tyr Glu Tyr Cys Arg
 85 90 95

Asp Cys His Asp Phe Arg Gly His Lys Asp Ile Cys Val Thr Gly Ala
 100 105 110

Ile Tyr Asn Tyr Thr Lys Gln Leu Thr Ser Gly Tyr His Asn Ser Gly
 115 120 125

Ser Glu Ile Arg Tyr Asn Leu Thr Glu Ala Leu Met Phe Leu Ser His
 130 135 140

Phe Ile Gly Asp Val His Gln Pro Leu His Val Gly Phe Thr Gly Asp
 145 150 155 160

Glu Gly Gly Asn Thr Ile Ile Val Arg Trp Tyr Arg Arg Lys Thr Asn

165	170	175
Leu His His Ile Trp Asp Asn Met Ile Ile Asp Ser Ala Leu Lys Thr 180 185 190		
Tyr Tyr Asn Ser Asp Leu Ala Ile Met Ile Gln Ala Ile Gln Arg Asn 195 200 205		
Ile Thr Gly Asp Trp Ser Phe Asp Ile Ser Ser Trp Lys Asn Cys Ala 210 215 220		
Ser Asp Asp Thr Ala Cys Pro Asn Leu Tyr Ala Ser Glu Ser Ile Ser 225 230 235 240		
Leu Ala Cys Lys Phe Ala Tyr Arg Asn Ala Thr Pro Gly Ser Thr Leu 245 250 255		
Gly Asp Asp Tyr Phe Leu Ser Arg Leu Pro Ile Val Glu Lys Arg Leu 260 265 270		
Ala Gln Gly Gly Ile Arg Leu Ala Ala Thr Leu Asn Arg Ile Phe Ala 275 280 285		
Ser Gln Pro Lys Ile Ser Leu Lys His Glu Asp Lys Arg Val Glu Lys 290 295 300		
Thr Thr Pro Val Asp Tyr Ile Glu Trp Ser Pro Leu Gln Gln Phe Ser 305 310 315 320		
Gly Ser His His His His His His His His Gly 325 330		

<210> 18
 <211> 840
 <212> DNA
 <213> Chocolate pots

<400> 18
 atgacttgagg gattttgggc acatcggaac atacatcgcc aagccgttta tcttatgcct 60
 tcgcccgtgg cagagttctt tcgcgcaaatt gttcaagaac ttgtcgaccg ctcggttgaa 120
 gccgatgaac gccgacgcat agacccaac gaagctccgc aacacttcat tgatttagac 180
 cgctacggtg cctatccttt tgaacaactt ccgagagatt atgaaaaagc cgttgagaaa 240
 ttccgctatg agcggctgaa agaaaatgga cttgtgccgt ggcgcattgc cgcctttgcc 300
 gatagcctca ccaacgcatt tcgggagcag aaccgcgaaa aaattttata cttcgccgca 360

aatttagggc attatgtcgc cgatgctaac gtgccacttc atgccaccga aaactacgac 420
ggacaactca cagggcaaaa aggattgcac gcacgttggg aaactattta tcctcaaaag 480
tttatgtccc cacgagaaac cacctatctc gaaaacggga gcatctttat cattgacaac 540
atcaccgaag aagccttcaa ctggtcatta gaaagttatg tattgagcca acaagttttg 600
gcgattgata agcaaattca atcggaattg tcagaagaag aattgtatga gttaaattca 660
tcagacgcgc cgccatttcg tcgcgatttt tcacaacgct attatgaaaa actcaaagaa 720
aaattgaatc aaatggttga aaaatgcttt gagttaagcg tcattagggg agcgtcagtt 780
tggtattttt cttgggttaaa agcagaaaaa ccgaatttat ttaacttatt aaaaaattga 840

<210> 19

<211> 279

<212> PRT

<213> Chocolate pots

<400> 19

Met Thr Trp Gly Phe Trp Ala His Arg Gln Ile His Arg Gln Ala Val
1 5 10 15

Tyr Leu Met Pro Ser Pro Val Ala Glu Phe Phe Arg Ala Asn Val Gln
20 25 30

Glu Leu Val Asp Arg Ser Val Glu Ala Asp Glu Arg Arg Arg Ile Asp
35 40 45

Pro Asn Glu Ala Pro Gln His Phe Ile Asp Leu Asp Arg Tyr Gly Ala
50 55 60

Tyr Pro Phe Glu Gln Leu Pro Arg Asp Tyr Glu Lys Ala Val Glu Lys
65 70 75 80

Phe Gly Tyr Glu Arg Leu Lys Glu Asn Gly Leu Val Pro Trp Arg Ile
85 90 95

Ala Ala Phe Ala Asp Ser Leu Thr Asn Ala Phe Arg Glu Gln Asn Arg
100 105 110

Glu Lys Ile Leu Tyr Phe Ala Ala Asn Leu Gly His Tyr Val Ala Asp
115 120 125

Ala Asn Val Pro Leu His Ala Thr Glu Asn Tyr Asp Gly Gln Leu Thr
130 135 140

Gly Gln Lys Gly Leu His Ala Arg Trp Glu Thr Ile Tyr Pro Gln Lys

145 150 155 160
 Phe Met Leu Pro Arg Glu Thr Thr Tyr Leu Glu Asn Gly Ser Ile Phe
 165 170 175
 Ile Ile Asp Asn Ile Thr Glu Glu Ala Phe Asn Trp Ser Leu Glu Ser
 180 185 190
 Tyr Val Leu Ser Gln Gln Val Leu Ala Ile Asp Lys Gln Ile Gln Ser
 195 200 205
 Glu Leu Ser Glu Glu Glu Leu Tyr Glu Leu Asn Ser Ser Asp Ala Pro
 210 215 220
 Pro Phe Arg Arg Asp Phe Ser Gln Arg Tyr Tyr Glu Lys Leu Lys Glu
 225 230 235 240
 Lys Leu Asn Gln Met Val Glu Lys Cys Phe Glu Leu Ser Val Ile Arg
 245 250 255
 Val Ala Ser Val Trp Tyr Phe Ser Trp Leu Lys Ala Glu Lys Pro Asn
 260 265 270
 Leu Phe Asn Leu Leu Lys Asn
 275

<210> 20
 <211> 807
 <212> DNA
 <213> Obsidian pool

<400> 20
 atgttttggg cacatcaaaa agtcaacgag catgccattg atttattacc cgagccactc 60
 cgcagttttt atgaacaaaa taaggaatac atagttaagg agtcggtcgc ccoctgatctc 120
 aggcggtcag aaaacaagga agaaggttat tatcactata tggatctcga taaatatggt 180
 gaatatccgt tcaagaattt gccagaaaac tacgacgacg cagtaaaaag gtttggttac 240
 gatactgttc tcaagaacgg aattgtgccg tggaaggtaa aatggttgac agacagtttg 300
 agtcaagcta tggagagaaa ggatgtgcc aaggtcttaa gactttcagc cgaccttggt 360
 cattatgttg ctgacatgca tgttccattt cattcgacag aaaattatga tggacagctg 420
 acaggcaaca taggaataca cttcagatgg gaaagcggca ttccagaaca ttttgaaca 480
 aattacaact atgaggggaat agagcccgt gtttacttca agcatcctga taaaaaggca 540
 tttgagatac tgactatgag ttacaagttg attctacctt ctctcaaggc tgatagtctt 600

gcaaaagtgtg gattgaatgg aaagagactt tataaagtgtg agagagaaga cggtaaaaaa 660
 gtttacgttt attcaaacga gtattatgag aagttcaaca aaaaccttgg tggatttgta 720
 gaatcgcaga tgaggctggc aatccatgat gttgcaagct actgggtatac tgcattgggta 780
 aatgccggta aaccaaagtt ttggttaa 807

<210> 21
 <211> 268
 <212> PRT
 <213> Obsidian pool

<400> 21

Met Phe Trp Ala His Gln Lys Val Asn Glu His Ala Ile Asp Leu Leu
 1 5 10 15

Pro Glu Pro Leu Arg Ser Phe Tyr Glu Gln Asn Lys Glu Tyr Ile Val
 20 25 30

Lys Glu Ser Val Ala Pro Asp Leu Arg Arg Ala Glu Asn Lys Glu Glu
 35 40 45

Gly Tyr Tyr His Tyr Met Asp Leu Asp Lys Tyr Gly Glu Tyr Pro Phe
 50 55 60

Lys Asn Leu Pro Glu Asn Tyr Asp Asp Ala Val Lys Arg Phe Gly Tyr
 65 70 75 80

Asp Thr Val Leu Lys Asn Gly Ile Val Pro Trp Lys Val Lys Trp Leu
 85 90 95

Thr Asp Ser Leu Ser Gln Ala Met Glu Arg Lys Asp Val Pro Gln Val
 100 105 110

Leu Arg Leu Ser Ala Asp Leu Gly His Tyr Val Ala Asp Met His Val
 115 120 125

Pro Phe His Ser Thr Glu Asn Tyr Asp Gly Gln Leu Thr Gly Asn Ile
 130 135 140

Gly Ile His Phe Arg Trp Glu Ser Gly Ile Pro Glu His Phe Gly Thr
 145 150 155 160

Asn Tyr Asn Tyr Glu Gly Ile Glu Pro Ala Val Tyr Phe Lys His Pro
 165 170 175

Asp Lys Lys Ala Phe Glu Ile Leu Thr Met Ser Tyr Lys Leu Ile Leu

180

185

190

Pro Ser Leu Lys Ala Asp Ser Leu Ala Lys Val Gly Leu Asn Gly Lys
 195 200 205

Arg Leu Tyr Lys Val Glu Arg Glu Asp Gly Lys Lys Val Tyr Val Tyr
 210 215 220

Ser Asn Glu Tyr Tyr Glu Lys Phe Asn Lys Asn Leu Gly Gly Ile Val
 225 230 235 240

Glu Ser Gln Met Arg Leu Ala Ile His Asp Val Ala Ser Tyr Trp Tyr
 245 250 255

Thr Ala Trp Val Asn Ala Gly Lys Pro Lys Phe Trp
 260 265

<210> 22
 <211> 972
 <212> DNA
 <213> Vitis vinifera

<400> 22
 atggccttggt ctgggggtctt gttgattgtg agggcacttg ttcttctgca attgattcct 60
 ggaattctga gttggggaaa ggaaggacac tatgcagttt gtaaaatagc tgaggggttc 120
 ctttctgaag atgcattagg agcagtgaag gcattgcttc cagattatgc tgaaggtgat 180
 ctggctgctg tttgctcctg ggctgatgag attcgtcaca acttccattg gcgatggagt 240
 ggccctttac attatgtaga tacgccagat tacaggtgta actatgaata ctgcagagac 300
 tgccatgact tcagaggaca caaagatata tgtgtaactg gagcaattta caattacaca 360
 aagcaactca cttctgggta tcacaattca gggtcagaaa taagatacaa tttgacagag 420
 gcaactcatgt tcttatcaca ttttattggg gatgtccatc agcccctaca tggttggtttt 480
 actggagatg aaggtgggaa cacaataata gtccgttggt accggaggaa gactaatttg 540
 catcatatat gggataacat gatcattgat tccgccctga agacatatta caattcagat 600
 cttgcaatca tgataaagc cattcaaaga aatattacgg gtgattggtc ctttgatata 660
 tcatcatgga aaaattgtgc atctgatgat acggcttggt caaacctgta tgcttctgaa 720
 agcattagtt tagcttgcaa gtttgcttac agaaatgcca caccaggaag cactctagga 780
 gatgattact tcctgtctcg gctaccaatt gtggagaaga ggctagccca aggtgggatac 840
 cgcttggtg ccacccttaa ccgtatcttt gcttctcaac caaaaatctc tctcaagcat 900
 gaagataaaa gggtagagaa aacaactcca gtggattata tagagtggag cccactgcaa 960

caattttcat aa

972

<210> 23
 <211> 323
 <212> PRT
 <213> Vitis vinifera

<400> 23

Met Ala Trp Ser Gly Val Leu Leu Ile Val Arg Ala Leu Val Leu Leu
 1 5 10 15

Gln Leu Ile Pro Gly Ile Leu Ser Trp Gly Lys Glu Gly His Tyr Ala
 20 25 30

Val Cys Lys Ile Ala Glu Gly Phe Leu Ser Glu Asp Ala Leu Gly Ala
 35 40 45

Val Lys Ala Leu Leu Pro Asp Tyr Ala Glu Gly Asp Leu Ala Ala Val
 50 55 60

Cys Ser Trp Ala Asp Glu Ile Arg His Asn Phe His Trp Arg Trp Ser
 65 70 75 80

Gly Pro Leu His Tyr Val Asp Thr Pro Asp Tyr Arg Cys Asn Tyr Glu
 85 90 95

Tyr Cys Arg Asp Cys His Asp Phe Arg Gly His Lys Asp Ile Cys Val
 100 105 110

Thr Gly Ala Ile Tyr Asn Tyr Thr Lys Gln Leu Thr Ser Gly Tyr His
 115 120 125

Asn Ser Gly Ser Glu Ile Arg Tyr Asn Leu Thr Glu Ala Leu Met Phe
 130 135 140

Leu Ser His Phe Ile Gly Asp Val His Gln Pro Leu His Val Gly Phe
 145 150 155 160

Thr Gly Asp Glu Gly Gly Asn Thr Ile Ile Val Arg Trp Tyr Arg Arg
 165 170 175

Lys Thr Asn Leu His His Ile Trp Asp Asn Met Ile Ile Asp Ser Ala
 180 185 190

Leu Lys Thr Tyr Tyr Asn Ser Asp Leu Ala Ile Met Ile Gln Ala Ile
 195 200 205

Gln Arg Asn Ile Thr Gly Asp Trp Ser Phe Asp Ile Ser Ser Trp Lys
 210 215 220

Asn Cys Ala Ser Asp Asp Thr Ala Cys Pro Asn Leu Tyr Ala Ser Glu
 225 230 235 240

Ser Ile Ser Leu Ala Cys Lys Phe Ala Tyr Arg Asn Ala Thr Pro Gly
 245 250 255

Ser Thr Leu Gly Asp Asp Tyr Phe Leu Ser Arg Leu Pro Ile Val Glu
 260 265 270

Lys Arg Leu Ala Gln Gly Gly Ile Arg Leu Ala Ala Thr Leu Asn Arg
 275 280 285

Ile Phe Ala Ser Gln Pro Lys Ile Ser Leu Lys His Glu Asp Lys Arg
 290 295 300

Val Glu Lys Thr Thr Pro Val Asp Tyr Ile Glu Trp Ser Pro Leu Gln
 305 310 315 320

Gln Phe Ser

<210> 24
 <211> 891
 <212> DNA
 <213> Solanum sp.

<400> 24
 atgggtgggt ttgagctcaa atggtttgta ggagtagctg ttgttctgat gatggttcaa 60
 aatattcttg gttgggggaa agaggacac tatattatct gcaaaattgc tgaggaatat 120
 ctaacagaag atgcttttagc tgcagtcaaa gcattactcc cagatcaagc cgaagggtgat 180
 cttgcagctg tctgctcctg gcctgatgag gttcggcgcc actaccacta ccgctggagc 240
 tctccattac attatgtaga tacacctgat ttcttgtgca attacaaata ttgccgagac 300
 tgccatgacg ggcatgggct caaggacagg tgtgttacgg gagcaatata caactactca 360
 atgcaacttt cgcagggata ttatgatttg aattcagaaa aatacaactt gactgaagca 420
 cttatgtttc tgtctcattt tgttggtgac gtacatcagc ctctccatgt tggtttccact 480
 ggagatcttg gtggaaacag tataattgtt cgttggtaca ggaggaagac taatttgcac 540
 catgtatggg ataacatgat tattgaatct gcgttgaaga catactacaa atctgatata 600
 atgttaatga cacaagttct tctgaaaaac atcactcatg aatggtccga tgatgttcca 660

tcttggaag attgcaagga gatggtttgt cctgacccat atgcttctga aagtatccgt 720
 ttggcctgca aatttgccta cagaaatgca accccgggaa gcactttaac agacgattac 780
 ttccctctctc gtcttctctgt tgtggagaag aggttggcac aaggtgggggt ccgcttggcc 840
 gaagttctca acagaatttt cactaaaaaa ccatcagatg ctgcacaatg a 891

<210> 25
 <211> 296
 <212> PRT
 <213> Solanum tuberosum

<400> 25

Met Gly Gly Phe Glu Leu Lys Trp Phe Val Gly Val Ala Val Val Leu
 1 5 10 15

Met Met Val Gln Asn Ile Leu Gly Trp Gly Lys Glu Gly His Tyr Ile
 20 25 30

Ile Cys Lys Ile Ala Glu Glu Tyr Leu Thr Glu Asp Ala Leu Ala Ala
 35 40 45

Val Lys Ala Leu Leu Pro Asp Gln Ala Glu Gly Asp Leu Ala Ala Val
 50 55 60

Cys Ser Trp Pro Asp Glu Val Arg Arg His Tyr His Tyr Arg Trp Ser
 65 70 75 80

Ser Pro Leu His Tyr Val Asp Thr Pro Asp Phe Leu Cys Asn Tyr Lys
 85 90 95

Tyr Cys Arg Asp Cys His Asp Gly His Gly Leu Lys Asp Arg Cys Val
 100 105 110

Thr Gly Ala Ile Tyr Asn Tyr Ser Met Gln Leu Ser Gln Gly Tyr Tyr
 115 120 125

Asp Leu Asn Ser Glu Lys Tyr Asn Leu Thr Glu Ala Leu Met Phe Leu
 130 135 140

Ser His Phe Val Gly Asp Val His Gln Pro Leu His Val Gly Phe Thr
 145 150 155 160

Gly Asp Leu Gly Gly Asn Ser Ile Ile Val Arg Trp Tyr Arg Arg Lys
 165 170 175

Thr Asn Leu His His Val Trp Asp Asn Met Ile Ile Glu Ser Ala Leu

180

185

190

Lys Thr Tyr Tyr Lys Ser Asp Ile Met Leu Met Thr Gln Val Leu Leu
 195 200 205

Lys Asn Ile Thr His Glu Trp Ser Asp Asp Val Pro Ser Trp Glu Asp
 210 215 220

Cys Lys Glu Met Val Cys Pro Asp Pro Tyr Ala Ser Glu Ser Ile Arg
 225 230 235 240

Leu Ala Cys Lys Phe Ala Tyr Arg Asn Ala Thr Pro Gly Ser Thr Leu
 245 250 255

Thr Asp Asp Tyr Phe Leu Ser Arg Leu Pro Val Val Glu Lys Arg Leu
 260 265 270

Ala Gln Gly Gly Val Arg Leu Ala Glu Val Leu Asn Arg Ile Phe Thr
 275 280 285

Lys Lys Pro Ser Asp Ala Ala Gln
 290 295

<210> 26
 <211> 885
 <212> DNA
 <213> Medicago sp.

<400> 26
 atgatcacgc tcttagttcc gttgctgcta tcaactcgcgt tgccaaatgt tctggcttgg 60
 ggaaaagatg gtcactatgc aatttgtaaa atttcacagg agtatcttag tgaagatgct 120
 ctatttgcag tcaaacaatt acttccagat tctgctcaag ctgatcttgc ttcagtttgc 180
 tcttggcctg atgagattcg ccataattac cattatcggt ggagtagtcc ttacattat 240
 attgatacac cagatttcaa atgtaactat caatattgca gagactgtca tgattcttat 300
 ggacataagc atagatgcgt tactggagca atatacaatt atacaatgca attaaaatta 360
 gctaacgccg atgcttcatc tgaattaaaa tataacttga cagaggcact tatgttcttg 420
 tcacatcttg ttggagatgt tcatcagccc ctacatgttg gttttactgg agacctaggt 480
 ggaaactcaa taacagttcg ttggtacagg aggaaaacaa atcttcatca cgtatgggat 540
 aacatgatta ttgagtctgc tctgaaaaag ttctatgggt cagatctttc aactatgata 600
 caggctattc aaaggaatat tagtgatatt tggtaaagt atgtatctat ttgggaacat 660
 tgtgcacaca accacacagc atgtccagac cggtatgctt ctgagagtat tagcttggca 720

tgcaagtttg cgtataagaa tgctacaccg ggaagcactt tggaagatga ctacttcctt 780
 tctcggttgc ctattgtgga gaaaaggctg gctcaaggtg gtgtgcgact tgcagctatc 840
 ctcaaccaca ttttcactcc gaagaccaga atagctcaag cttaa 885

<210> 27
 <211> 294
 <212> PRT
 <213> Medicago sp.
 <400> 27

Met Ile Thr Leu Leu Val Pro Leu Leu Leu Ser Leu Ala Leu Pro Asn
 1 5 10 15

Val Leu Ala Trp Gly Lys Asp Gly His Tyr Ala Ile Cys Lys Ile Ser
 20 25 30

Gln Glu Tyr Leu Ser Glu Asp Ala Leu Phe Ala Val Lys Gln Leu Leu
 35 40 45

Pro Asp Ser Ala Gln Ala Asp Leu Ala Ser Val Cys Ser Trp Pro Asp
 50 55 60

Glu Ile Arg His Asn Tyr His Tyr Arg Trp Ser Ser Pro Leu His Tyr
 65 70 75 80

Ile Asp Thr Pro Asp Phe Lys Cys Asn Tyr Gln Tyr Cys Arg Asp Cys
 85 90 95

His Asp Ser Tyr Gly His Lys His Arg Cys Val Thr Gly Ala Ile Tyr
 100 105 110

Asn Tyr Thr Met Gln Leu Lys Leu Ala Asn Ala Asp Ala Ser Ser Glu
 115 120 125

Leu Lys Tyr Asn Leu Thr Glu Ala Leu Met Phe Leu Ser His Phe Val
 130 135 140

Gly Asp Val His Gln Pro Leu His Val Gly Phe Thr Gly Asp Leu Gly
 145 150 155 160

Gly Asn Ser Ile Thr Val Arg Trp Tyr Arg Arg Lys Thr Asn Leu His
 165 170 175

His Val Trp Asp Asn Met Ile Ile Glu Ser Ala Leu Lys Lys Phe Tyr
 180 185 190

Gly Ser Asp Leu Ser Thr Met Ile Gln Ala Ile Gln Arg Asn Ile Ser
195 200 205

Asp Ile Trp Ser Asn Asp Val Ser Ile Trp Glu His Cys Ala His Asn
210 215 220

His Thr Ala Cys Pro Asp Arg Tyr Ala Ser Glu Ser Ile Ser Leu Ala
225 230 235 240

Cys Lys Phe Ala Tyr Lys Asn Ala Thr Pro Gly Ser Thr Leu Glu Asp
245 250 255

Asp Tyr Phe Leu Ser Arg Leu Pro Ile Val Glu Lys Arg Leu Ala Gln
260 265 270

Gly Gly Val Arg Leu Ala Ala Ile Leu Asn His Ile Phe Thr Pro Lys
275 280 285

Thr Arg Ile Ala Gln Ala
290

<210> 28
<211> 271
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic construct

<400> 28

Trp Gly Lys Glu Gly His Tyr Ala Val Cys Lys Ile Ala Glu Gly Phe
1 5 10 15

Leu Ser Glu Asp Ala Leu Gly Ala Val Lys Gly Leu Leu Pro Asp Tyr
20 25 30

Ala Asp Gly Asp Leu Ala Ala Val Cys Ser Trp Ala Asp Glu Ile Arg
35 40 45

His Asn Phe His Trp Arg Trp Ser Gly Pro Leu His Tyr Val Asp Thr
50 55 60

Pro Asp Tyr Arg Cys Asn Tyr Glu Tyr Cys Arg Asp Cys His Asp Phe
65 70 75 80

Arg Gly His Lys Asp Ile Cys Val Thr Gly Ala Ile Tyr Asn Tyr Thr
85 90 95

Lys Gln Leu Thr Ser Gly Tyr His Asn Ser Gly Ser Glu Ile Arg Tyr
 100 105 110

Asn Leu Thr Glu Ala Leu Met Phe Leu Ser Asp Phe Ile Gly Asp Val
 115 120 125

His Gln Pro Leu His Val Gly Phe Thr Gly Asp Glu Gly Gly Asn Thr
 130 135 140

Ile Ile Val Arg Trp Tyr Arg Arg Lys Thr Asn Leu His His Ile Trp
 145 150 155 160

Asp Asp Met Ile Ile Asp Ser Ala Leu Lys Thr Tyr Tyr Asn Ser Asp
 165 170 175

Ile Ala Ile Met Ile Gln Ala Ile Gln Arg Asn Ile Thr Gly Asp Trp
 180 185 190

Ser Phe Asp Ile Ser Ser Trp Lys Asn Cys Ala Ser Asp Asp Thr Ala
 195 200 205

Cys Pro Asn Leu Tyr Ala Ser Glu Gly Ile Ser Leu Ala Cys Lys Phe
 210 215 220

Ala Tyr Arg Asn Ala Thr Pro Gly Ser Thr Leu Gly Asp Asp Tyr Phe
 225 230 235 240

Leu Ser Arg Leu Pro Ile Val Glu Lys Arg Leu Ala Pro Ser Gly Ile
 245 250 255

Arg Leu Ala Ala Thr Leu Asn Arg Ile Phe Ala Ser Gln Gly Lys
 260 265 270

<210> 29
 <211> 271
 <212> PRT
 <213> Vitis vinifera

<400> 29

Trp Gly Lys Glu Gly His Tyr Ala Val Cys Lys Ile Ala Glu Gly Phe
 1 5 10 15

Leu Ser Glu Asp Ala Leu Gly Ala Val Lys Ala Leu Leu Pro Asp Tyr
 20 25 30

Ala Glu Gly Asp Leu Ala Ala Val Cys Ser Trp Ala Asp Glu Ile Arg
 35 40 45

His Asn Phe His Trp Arg Trp Ser Gly Pro Leu His Tyr Val Asp Thr
 50 55 60

Pro Asp Tyr Arg Cys Asn Tyr Glu Tyr Cys Arg Asp Cys His Asp Phe
 65 70 75 80

Arg Gly His Lys Asp Ile Cys Val Thr Gly Ala Ile Tyr Asn Tyr Thr
 85 90 95

Lys Gln Leu Thr Ser Gly Tyr His Asn Ser Gly Ser Glu Ile Arg Tyr
 100 105 110

Asn Leu Thr Glu Ala Leu Met Phe Leu Ser His Phe Ile Gly Asp Val
 115 120 125

His Gln Pro Leu His Val Gly Phe Thr Gly Asp Glu Gly Gly Asn Thr
 130 135 140

Ile Ile Val Arg Trp Tyr Arg Arg Lys Thr Asn Leu His His Ile Trp
 145 150 155 160

Asp Asn Met Ile Ile Asp Ser Ala Leu Lys Thr Tyr Tyr Asn Ser Asp
 165 170 175

Leu Ala Ile Met Ile Gln Ala Ile Gln Arg Asn Ile Thr Gly Asp Trp
 180 185 190

Ser Phe Asp Ile Ser Ser Trp Lys Asn Cys Ala Ser Asp Asp Thr Ala
 195 200 205

Cys Pro Asn Leu Tyr Ala Ser Glu Ser Ile Ser Leu Ala Cys Lys Phe
 210 215 220

Ala Tyr Arg Asn Ala Thr Pro Gly Ser Thr Leu Gly Asp Asp Tyr Phe
 225 230 235 240

Leu Ser Arg Leu Pro Ile Val Glu Lys Arg Leu Ala Gln Gly Gly Ile
 245 250 255

Arg Leu Ala Ala Thr Leu Asn Arg Ile Phe Ala Ser Gln Pro Lys
 260 265 270

<210> 30

<211> 828
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Synthetic construct: Codon Optimized Mature Core Mimulus DNA
 Encodes SEQ ID NO:14

<400> 30

tggagtaagg agggacatag catgacatgt aagatagccc aggacttggt ggggtcccgaa	60
gccaaacacg ccgtgcaaat gttgttgccct gaaaatgtga acggcgacct aagcgacctg	120
tcggtgtggc cggaccaagt gagacactgg taaaaataga gatggacctc ccctttgcac	180
ttcattgaca ccccgatca ggcttgcaac tttaactacc agagagactg ccatgacccg	240
cacggtgtaa aaggcatgtg cgttgccggg gccattcaaa atttcacgaa ccaattgtcg	300
cactacagac acggcacgtc ggacagacgt tacaacatga cggaggcctt gttgtttttg	360
gccacttta tgggcgatat tcatcagccg ttgcacgtgg gcttcacgtc agacgaaggc	420
ggcaacacga ttgacttgag atggtttcgc cacaagagca acttgcatca cgtatgggat	480
cgagaaatta tcctaactgc cgctgcccgc tactacggaa aggacatcga cctactccag	540
gaggatatca aaggcaattt tactgacggc atctggctcg gcgatttggc ctctgtggaga	600
gaatgttcgg acattttttc gtgtgtgaac aagtacgctg ccgaatccat aaacatggct	660
tgtaaatggg gctacaagga tgtgaaatcg ggtgacacgc tctcggaaga ctatttcaac	720
agtcgtctcc cgatcgtaat gaaaagaatc gctcaaggag gcgttcgctt agcaatgatt	780
ctcaacagag tattcgggtga tagcaaagag gacagcttga ttgccacg	828

<210> 31
 <211> 933
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Synthetic construct: Blue Sky Report Mimulus DNA Encodes SEQ ID
 NO:11

<400> 31

accatgaagt tcttgggtcaa cgtagcactg gtttttatgg tagtctatat cagctacatt	60
tacgcgtgga gtaaggaggg acatagcatg acatgtaaga tagcccagga cttgttgggt	120
cccgaagcca aacacgccgt gcaaatgttg ttgcctgaaa atgtgaacgg cgacctaacg	180
gccttgctcg tgtggccgga ccaagtgaga cactggtaca aatacagatg gacctcccct	240
ttgcacttca ttgacacccc cgatcagggt tgcaacttta actaccagag agactgccat	300
gacccgcacg gtgtaaaagg catgtgcgtt gccgggtgcca ttcaaaattt cacgaaccaa	360
ttgtcgcact acagacacgg cacgtcggac agacgttaca acatgacgga ggccttggtg	420

```

tttttggccc actttatggg cgatattcat cagccgttgc acgtgggctt cacgtcagac      480
gaaggcggca acacgattga cttgagatgg ttctgccaca agagcaactt gcatcacgta      540
tgggatcgag aaattatcct aactgccgct gcggactact acggaaagga catcgaccta      600
ctccaggagg atatcaaagg caattttact gacggcatct ggtcgggcga tttggcctcg      660
tggagagaat gttcggacat tttttcgtgt gtgaacaagt acgtgccga atccataaac      720
atggcttgta aatggggcta caaggatgtg aaatcgggtg acacgctctc ggacgactat      780
ttcaacagtc gtctcccgat cgtaatgaaa agaatcgctc aaggaggcgt tcgcttagca      840
atgattctca acagagtatt cggatgtagc aaagaggaca gcttgattgc cacgggctcg      900
caccatcacc accatcacca ccacggttga taa                                  933

```

```

<210> 32
<211> 897
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Synthetic construct: Codon Optimized Vitis DNA Encodes Residues
      XXX YYY of SEQ ID NO:23

```

```

<400> 32
tggggcaaag aaggccacta cgccgtgtgt aagattgcgg agggcttttt gtcggaagac      60
gcattgggag cggatcaaagc cttgttgccg gactacgcgg aaggcgactt ggcagccgta      120
tgtagctggg ccgacgagat cagacacaac ttctactgga gatggtcggg cccactgcat      180
tacgtcgaca cgccggatta cagatgcaac tacgagtact gccgcgactg tcacgacttc      240
agaggccaca aagacatttg cgtcacgggc gcgatataca actacacgaa acaattgacg      300
tcgggctacc acaacagtgg ctccgagatt cgatacaacc tcacggaggc cttgatgttc      360
ctctcgcat tcatggcgga cgtgcaccaa ccgctgcatg tgggctttac gggcgatgaa      420
ggcgaaata cgatcattgt ccgttggtac cgcagaaaga ccaacctcca ccacatatgg      480
gacaacatga tcatcgactc ggcgttgaag acctactaca acagcgacct ggccataatg      540
atccaggcga ttcaaagaaa catcacgggc gattggctct ttgacatcag cagctggaag      600
aactgtgcca gtgacgacac tgcttgtccg aacctatacg cgtcggagag catctcgttg      660
gcctgtaa at ttgcctacag aaatgccacc cccggttcga cgctgggcga cgactacttc      720
ttgtcgcgat tgccgattgt tgaaaaacgc ctgcaccaag gcggtattag attggccgcc      780
acctgaacc gtatTTTTTgc ctgcgaaccg aaaatctcgc tgaaacacga agacaagaga      840
gtcgagaaga cgacgccggt agactacatc gagtggctgc cattgcaaca gttcagc      897

```

<210> 33
 <211> 1002
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Synthetic construct: Blue Sky Report Vitis DNA Encodes SEQ ID NO:17

<400> 33

accatgaagt tcttggtgaa cgtggcgctg gtgttcattg tcgtgtacat ctctacatt	60
tacgcgtggg gcaaagaagg ccactacgcc gtgtgtaaga ttgcggaggg ctttttgtcg	120
gaagacgcat tgggagcggc caaagccttg ttgccggact acgcggaagg cgacttggca	180
gccgtatgta gctgggcccga cgagatcaga cacaactttc actggagatg gtcgggcccga	240
ctgcattacg tcgacacgcc ggattacaga tgcaactacg agtactgccg cgactgtcac	300
gacttcagag gccacaaaga catttgcgtc acgggcgcga tatacaacta cacgaaacaa	360
ttgacgtcgg gctaccacaa cagtggctcc gagattcgat acaacctcac ggaggccttg	420
atgttctctc cgcatttcat tggcgacgtg caccaaccgc tgcattgtgg ctttacgggc	480
gatgaaggcg gaaatacgat cattgtcgtg ttgtaccgca gaaagaccaa cctccaccac	540
atatgggaca acatgatcat cgactcggcg ttgaagacct actacaacag cgacctggcc	600
ataatgatcc aggcgattca aagaacatc accggcgatt ggtcctttga catcagcagc	660
tggaagaact gtgccagtga cgacactgct tgtccgaacc tatacgcgtc ggagagcatt	720
tcgttggcct gtaaatttgc ctacagaaat gccacccccg gttcgacgct gggcgacgac	780
tacttcttgt cgcgattgcc gattgttgaa aaacgcctcg cccaaggcgg tattagattg	840
gccgccacct tgaaccgtat ttttgcctcg caaccgaaaa tctcgtgaa acacgaagac	900
aagagagtcg agaagacgac gccggtagac tacatcgagt ggtcgccatt gcaacagttc	960
agcgggaagcc accaccatca ccaccatcat cagggctgat aa	1002