Claims

1. A method for producing a cellular structure comprising hypothalamic tissue, the method comprising the steps of:

(1) culturing an aggregate of human pluripotent stem cells in suspension in a medium containing a bone morphogenetic protein signal transduction pathway activating substance and of a substance acting on the Shh signaling pathway;

(2) further culturing the cell aggregate obtained in the step (1) in a medium containing a substance acting on the Shh signaling pathway, but not containing a bone morphogenetic protein signal transduction pathway activating substance; and

(3) recovering the cell aggregate obtained in the step (2) and subjecting the cells constituting the cell aggregate to dissociation culture,

wherein the bone morphogenetic protein signal transduction pathway activating substance in the step (1) is selected from the group consisting of BMP2, BMP4, BMB7 and GDF5, and the concentration thereof in the medium is 0.1 nM to 5.0 nM; and

wherein the substance acting on the Shh signaling pathway in the steps (1) and (2) is selected from the group consisting of proteins belonging to the Hedgehog family, Shh receptors, Shh receptor agonists, Purmorphamine, SAG and Hh-Ag 1.5, and the concentration thereof in the medium is 0.1 μM to 2.0 μM.

2. The producing method according to claim 1, wherein the step (2) is carried out under an oxygen partial pressure condition of 30 to 60 %.

3. The producing method according to claim 1 or 2, wherein the step (3) is carried out under an oxygen partial pressure condition of 30 to 60 %.

4. The producing method according to any one of claims 1 to 3,

wherein the bone morphogenetic protein signal transduction pathway activating substance in the step (1) is BMP4,

wherein the substance acting on the Shh signaling pathway in the steps (1) and (2) is SAG, and

wherein the differentiation into dorsal hypothalamic tissue is induced by the steps (1) and (2).

5. The producing method according to claim 4, wherein the dorsal hypothalamic tissue comprises one or more neurons selected from the group consisting of vasopressin

neurons, oxytocin neurons, thyrotropin releasing hormone neurons, corticotropin releasing hormone neurons and neuropeptide Y neurons.

6. The producing method according to any one of claims 1 to 3,

wherein the bone morphogenetic protein signal transduction pathway activating substance in the step (1) is BMP4, and the concentration thereof in the medium is 0.1 nM to 3.0 nM,

wherein the substance acting on the Shh signaling pathway in the steps (1) and (2) is SAG, and the concentration thereof in the medium is 0.2 μM to 1.5 μM,

wherein the medium further comprises an Akt inhibitor, and

wherein the differentiation into ventral hypothalamus tissue is induced by the steps (1) and (2).

7. The producing method according to claim 6, wherein the ventral hypothalamic tissue comprises one or more neurons selected from the group consisting of agouti-related protein neurons, proopiomelanocortin neurons, melanin-concentrating hormone neurons and Orexin neurons.

8. The producing method according to any one of claims 1 to 7, wherein the suspension culture is carried out in the absence of feeder cells.

9. The producing method according to any one of claims 1 to 8, wherein the aggregate in the step (1) is formed by culturing dispersed human pluripotent stem cells in suspension.

10. The producing method according to claim 9, wherein the suspension culture is carried out by SFEBq (Serum-free Floating culture of Embryoid Body-like aggregates with quick reaggregation) method.

11. A method for producing a cellular structure comprising hypothalamic tissue and pituitary tissue, the method comprising the steps of:

(i) culturing an aggregate of human pluripotent stem cells in suspension in a medium containing a bone morphogenetic protein signal transduction pathway activating substance and a substance acting on the Shh signaling pathway,

(ii) further culturing the cell aggregate formed in the step (i) in suspension in a medium containing a bone morphogenetic protein signal transduction pathway

activating substance and the substance acting on the Shh signaling pathway, and

(iii) culturing the cell aggregate obtained in the step (ii) in suspension in a medium suitable for simultaneous induction of pituitary and hypothalamus, wherein said medium is obtained by mixing the medium used in step (ii) in a volume ratio of 1:1 with a medium having a ciliary neurotrophic factor (CNTF), a brain-derived neurotrophic factor (BDNF), a neurotrophin 3 (NT-3), a fetal bovine serum, an N2 supplement and a B27 supplement;

wherein the bone morphogenetic protein signal transduction pathway activating substance is selected from the group consisting of BMP2, BMP4, BMB7 and GDF5 and the substance acting on the Shh signaling pathway is selected from the group consisting of proteins belonging to the Hedgehog family, Shh receptors, Shh receptor agonists, Purmorphamine, SAG and Hh-Ag 1.5.

12. The producing method according to claim 11, wherein the steps (ii) and (iii) are carried out under an oxygen partial pressure condition of 30 to 60 %.

13. The producing method according to claim 11or 12,

wherein the following step (a) is performed between the steps (ii) and (iii), and wherein, in the step (iii), the cell aggregate obtained in the step (a) is cultured in suspension:

(a) further culturing the cell aggregate obtained in the step (ii) in suspension in a medium containing the substance acting on the Shh signaling pathway,

wherein the substance acting on the Shh signaling pathway is selected from the group consisting of proteins belonging to the Hedgehog family, Shh receptors, Shh receptor agonists, Purmorphamine, SAG and Hh-Ag 1.5.

14. The producing method according to any one of claims 11 to 13, wherein the step (a) is carried out under an oxygen partial pressure condition of 30 to 60 %.

15. The producing method according to any one of claims 11 to 14, wherein the suspension culture is carried out in the absence of feeder cells.

16. The producing method according to any one of claims 11 to 15, wherein the aggregate in the step (i) is formed by culturing dispersed human pluripotent stem cells in suspension.

17. The producing method according to any one of claims 11 to 16, wherein the bone morphogenetic protein signal transduction pathway activating substance is BMP4.

18. The producing method according to any one of claims 11 to 17, wherein the substance acting on the Shh signaling pathway is SAG.

19. A cellular structure obtained by the producing method according to any one of claims 1 to 18, wherein

(a) the hypothalamic tissue is corticotropin-releasing hormone (CRH) positive;

(b) the dorsal hypothalamic tissue is vasopressin neuron (AVP) positive, Rx positive, Pax6 positive and Nkx2.1 negative;

(c) the ventral hypothalamic tissue is positive for melanin concentrating hormone neurons (MCH) positive, Rx positive, Pax6 negative and Nkx2.1 positive; and

(d) the pituitary tissue is Adrenocorticotropic hormone (ACTH) positive.