

QUINAZOLINE COMPOUNDS

FIELD OF THE INVENTION

The present invention relates to compounds that inhibit TNNI3K and methods of making and using the same. Specifically, the present invention relates to quinazolines as TNNI3K inhibitors.

BACKGROUND OF THE INVENTION

Cardiac troponin I-interacting kinase (TNNI3K), also known as CARK (for cardiac ankyrin repeat kinase), is a protein kinase that exhibits highly selective expression for cardiac tissues and has been shown to interact with components of the sarcomere, including troponin I (Zhao, Y. et al., *J. Mol. Med.*, 2003, 81, 297-304; Feng, Y. et al., *Gen. Physiol. Biophys.*, 2007, 26, 104-109; Wang, H. et al., *J. Cell. Mol. Med.*, 2008, 12, 304-315). Although substrates for TNNI3K have not been identified to date, recent reports suggest that this protein does play a role in the development of pressure-induced cardiomyocyte hypertrophy and contractile dysfunction (Wheeler, F. C. et al., *Mamm. Genome*, 2005, 16, 414-423; Wang, X. et al. "TNNI3K, a cardiac-specific kinase, promotes cardiac hypertrophy in vivo", Poster presentation at the 2006 Scientific Sessions of the American Heart Association, Chicago, IL, Wheeler, F. C. et al., *PLos Genet*, 2009, 5(9), e1000647; and Pu, W.T., *PLos Genet*, 2009, 5(9), e1000643). Inhibition of the kinase activity of TNNI3K may disrupt these signaling pathways, and enable the mitigation and/or reversal of cardiac hypertrophy seen in patients with progressively worsening heart failure.

In response to mechanical, neurohormonal, and genetic stimuli, the heart will undergo hypertrophy, or muscle growth and remodeling, in order to maintain sufficient cardiac output to meet tissue oxygen demands. While these structural changes are initially seen as compensatory, sustained dysregulation of hypertrophic signaling can lead to heart failure, the pathophysiological state in which the heart can no longer adequately function as a pump (Mudd, J. O. and Kass, D. A., *Nature*, 2008, 451, 919-928).

Prevention or reversal of pathological cardiac hypertrophy has the potential to delay or prevent the development of congestive heart failure (McKinsey, T. A. and Kass, D. A., *Nat. Rev. Drug Discov.*, 2007, 6, 617-635; Kaye, D. M. and Krum, H., *Nat. Rev. Drug Discov.*, 2007, 6, 127-139).


Heart failure is responsible for a reduced quality of life and premature death in a significant proportion of sufferers, and is characterized by impaired cardiac function either due to reduced pump function (systolic dysfunction) or reduced filling (diastolic dysfunction). Congestive heart failure (CHF) is characterized by impaired left ventricular

function, increased peripheral and pulmonary vascular resistance and reduced exercise tolerance and dyspnea. The prevalence of heart failure is anticipated to increase with ageing populations, prompting a need for new and improved methods of treating heart failure.

5

SUMMARY OF THE INVENTION

The invention is directed to novel quinazolines. Specifically, the invention is directed to a compound according to Formula I:



10

I

wherein:

R¹ is (C₁-C₄)alkyl;

R² is H or halogen;

15 R³ is H, halogen, (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, hydroxyl, hydroxy(C₁-C₈)alkyl-, (C₁-C₈)alkoxy, (C₃-C₈)cycloalkyloxy, (C₁-C₈)haloalkoxy, (C₁-C₈)alkylthio-, (C₁-C₈)haloalkylthio-, (C₃-C₈)cycloalkylthio-, (C₁-C₈)alkylsulfonyl, (C₁-C₈)haloalkylsulfonyl, (C₃-C₈)cycloalkylsulfonyl, phenyl, 5-membered heteroaryl, or -N(Rᵃ)(Rᵇ),

wherein said phenyl or heteroaryl is optionally substituted one to three times,

20 independently, by halogen, (C₁-C₆)alkyl, (C₁-C₄)haloalkyl, (C₁-C₆)alkoxy, (C₁-C₄)haloalkoxy, hydroxy(C₁-C₄)alkyl-, or -N(Rᵃ)(Rᵇ);

each Rᵃ is independently (C₁-C₄)alkyl, wherein said (C₁-C₄)alkyl is optionally substituted one to three times, independently, by halogen, hydroxyl, (C₁-C₆)alkoxy, amino, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)((C₁-C₆)alkyl)amino, -CO₂H, -CO₂(C₁-C₆)alkyl, -CONH₂,

25 -CONH(C₁-C₆)alkyl, or -CON((C₁-C₆)alkyl)((C₁-C₆)alkyl), and

Rᵇ is (C₁-C₄)alkyl;

or Rᵃ and Rᵇ taken together with the nitrogen atom to which they are attached form a 5-membered or 6-membered heterocyclic ring, optionally containing one additional heteroatom selected from nitrogen, oxygen and sulfur, wherein said ring is optionally

30 substituted one or two times, independently, by halogen, (C₁-C₄)alkyl, (C₁-C₄)haloalkyl,

amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, hydroxyl, hydroxy(C₁-C₄)alkyl-, oxo, (C₁-C₄)alkoxy, (C₁-C₄)haloalkoxy, or (C₁-C₄)alkoxy(C₁-C₄)alkyl;

R⁴ is H;

R⁵, R⁶, and R⁷ are each, independently, H, halogen, nitro, amino,

5 (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, hydroxyl, or OR^c; and

each R^c is, independently, (C₁-C₈)alkyl or a 5-6 membered heterocycloalkyl, wherein said (C₁-C₈)alkyl is optionally substituted one to three times, independently, by halogen, hydroxyl, (C₁-C₄)alkoxy, amino, (C₁-C₄)alkylamino, or -N(R^a)(R^b);

or R⁶ and R⁷ taken together represent -O(C₁-C₂)alkylo-;

10 or a salt, particularly a pharmaceutically acceptable salt, thereof.

The compounds of the invention are inhibitors of TNNI3K and can be useful for the treatment of cardiac diseases and disorders, particularly heart failure. Accordingly, the invention is further directed to pharmaceutical compositions comprising a compound of the
15 invention. The invention is still further directed to methods of inhibiting TNNI3K and treatment of conditions associated therewith using a compound of the invention or a pharmaceutical composition comprising a compound of the invention.

DETAILED DESCRIPTION OF THE INVENTION

20 As used herein, the term "alkyl" represents a saturated, straight or branched hydrocarbon moiety, which may be unsubstituted or substituted by one, or more of the substituents defined herein. Exemplary alkyls include, but are not limited to methyl (Me), ethyl (Et), propyl, isopropyl, butyl, isobutyl, *t*-butyl and pentyl. The term "C₁-C₄" refers to an alkyl containing from 1 to 4 carbon atoms.

25 When the term "alkyl" is used in combination with other substituent groups, such as "haloalkyl" or "hydroxyalkyl" or "arylalkyl", the term "alkyl" is intended to encompass a divalent straight or branched-chain hydrocarbon radical. For example, "arylalkyl" is intended to mean the radical -alkylaryl, wherein the alkyl moiety thereof is a divalent straight or branched-chain carbon radical and the aryl moiety thereof is as defined herein,
30 and is represented by the bonding arrangement present in a benzyl group (-CH₂-phenyl).

As used herein, the term "cycloalkyl" refers to a non-aromatic, saturated, cyclic hydrocarbon ring. The term "(C₃-C₈)cycloalkyl" refers to a non-aromatic cyclic hydrocarbon ring having from three to eight ring carbon atoms. Exemplary
35 "(C₃-C₈)cycloalkyl" groups useful in the present invention include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl.

"Alkoxy" refers to a group containing an alkyl radical attached through an oxygen linking atom. The term "(C₁-C₄)alkoxy" refers to a straight- or branched-chain hydrocarbon radical having at least 1 and up to 4 carbon atoms attached through an oxygen linking atom. Exemplary "(C₁-C₄)alkoxy" groups useful in the present invention include, but are not limited to, methoxy, ethoxy, *n*-propoxy, isopropoxy, *n*-butoxy, *s*-butoxy, and *t*-butoxy.

"Alkylthio-" refers to a group containing an alkyl radical attached through a sulfur linking atom. The term "(C₁-C₄)alkylthio-" refers to a straight- or branched-chain hydrocarbon radical having at least 1 and up to 4 carbon atoms attached through a sulfur linking atom. Exemplary "(C₁-C₄)alkylthio-" groups useful in the present invention include, but are not limited to, methylthio-, ethylthio-, *n*-propylthio-, isopropylthio-, *n*-butylthio-, *s*-butylthio-, and *t*-butylthio-.

"Alkylsulfonyl" refers to a group containing an alkyl radical attached through a sulfonyl radical (i.e. -S(=O)₂-). The term "(C₁-C₄)alkylsulfonyl" refers to a straight- or branched-chain hydrocarbon radical having at least 1 and up to 4 carbon atoms attached through a sulfonyl radical. Exemplary "(C₁-C₄)alkylsulfonyl" groups useful in the present invention include, but are not limited to, methanesulfonyl, ethanesulfonyl, propanesulfonyl, and butanesulfonyl.

"Cycloalkyloxy" and "cycloalkylthio" refers to a group containing a saturated carbocyclic ring atoms attached through an oxygen or sulfur linking atom, respectively. Examples of "cycloalkyloxy" moieties include, but are not limited to, cyclopropoxy, cyclobutoxy, cyclopentyloxy, cyclohexyloxy, and the like.

"Aryl" represents a group or moiety comprising an aromatic, monovalent monocyclic or bicyclic hydrocarbon radical containing from 6 to 10 carbon ring atoms, which may be unsubstituted or substituted by one or more of the substituents defined herein, and to which may be fused one or more cycloalkyl rings, which may be unsubstituted or substituted by one or more substituents defined herein.

Generally, in the compounds of this invention, aryl is phenyl.

Heterocyclic groups may be heteroaryl or heterocycloalkyl groups.

"Heterocycloalkyl" represents a group or moiety comprising a non-aromatic, monovalent monocyclic or bicyclic radical, which is saturated or partially unsaturated, containing 3 to 10 ring atoms, which includes 1 to 4 heteroatoms selected from nitrogen, oxygen and sulfur, and which may be unsubstituted or substituted by one or more of the substituents defined herein. Illustrative examples of heterocycloalkyls include, but are not limited to, azetidiny, pyrrolidyl (or pyrrolidinyl), piperidiny, piperazinyl, morpholinyl, tetrahydro-4*H*-1,4-thiazinyl, tetrahydrofuryl (or tetrahydrofuranyl), dihydrofuryl, oxazolinyl, thiazolinyl, pyrazolinyl, tetrahydropyranyl, dihydropyranyl, 1,3-dioxolanyl, 1,3-dioxanyl, 1,4-

dioxanyl, 1,3-oxathiolanyl, 1,3-oxathianyl, 1,3-dithianyl, azabicyclo[3.2.1]octyl, azabicyclo[3.3.1]nonyl, azabicyclo[4.3.0]nonyl, oxabicyclo[2.2.1]heptyl and 1,5,9-triazacyclododecyl.

Generally, in the compounds of this invention, heterocycloalkyl groups are 5-membered and/or 6-membered heterocycloalkyl groups, such as pyrrolidyl (or pyrrolidinyl), tetrahydrofuryl (or tetrahydrofuranyl), tetrahydrothienyl, dihydrofuryl, oxazolinyl, thiazolinyl, pyrazolinyl, piperidyl (or piperidinyl), piperazinyl, morpholinyl, tetrahydropyranyl, dihydropyranyl, 1,3-dioxanyl, tetrahydro-4*H*-1,4-thiazinyl, 1,4-dioxanyl, 1,3-oxathianyl, and 1,3-dithianyl.

"Heteroaryl" represents a group or moiety comprising an aromatic monovalent monocyclic or bicyclic radical, containing 5 to 10 ring atoms, including 1 to 4 heteroatoms selected from nitrogen, oxygen and sulfur, which may be unsubstituted or substituted by one or more of the substituents defined herein. This term also encompasses bicyclic heterocyclic-aryl compounds containing an aryl ring moiety fused to a heterocycloalkyl ring moiety, containing 5 to 10 ring atoms, including 1 to 4 heteroatoms selected from nitrogen, oxygen and sulfur, which may be unsubstituted or substituted by one or more of the substituents defined herein. Illustrative examples of heteroaryls include, but are not limited to, thienyl, pyrrolyl, imidazolyl, pyrazolyl, furyl (or furanyl), isothiazolyl, isoxazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridyl (or pyridinyl), pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, tetrazinyl, triazolyl, tetrazolyl, benzo[b]thienyl, isobenzofuryl, 2,3-dihydrobenzofuryl, chromenyl, chromanyl, indolizinyl, isoindolyl, indolyl, indazolyl, purinyl, isoquinolyl, quinolyl, phthalazinyl, naphthridinyl, quinazolinyl, benzothiazolyl, benzimidazolyl, tetrahydroquinolyl, cinnolyl, pteridinyl, and isothiazolyl.

Generally, the heteroaryl groups present in the compounds of this invention are 5-membered and/or 6-membered monocyclic heteroaryl groups. Selected 5-membered heteroaryl groups contain one nitrogen, oxygen or sulfur ring heteroatom, and optionally contain 1, 2 or 3 additional nitrogen ring atoms. Selected 6-membered heteroaryl groups contain 1, 2, 3 or 4 nitrogen ring heteroatoms. Selected 5- or 6-membered heteroaryl groups include thienyl, pyrrolyl, imidazolyl, pyrazolyl, furyl, isothiazolyl, isoxazolyl, oxazolyl, oxadiazolyl, thiazolyl, triazolyl, tetrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, and triazinyl.

"Oxo" represents a double-bonded oxygen moiety; for example, if attached directly to a carbon atom forms a carbonyl moiety (C=O).

The terms "halogen" and "halo" represent chloro, fluoro, bromo or iodo substituents.

"Hydroxy" or "hydroxyl" is intended to mean the radical -OH.

As used herein, the term "compound(s) of the invention" means a compound of formula (I) (as defined above) in any form, i.e., any salt or non-salt form (e.g., as a free acid or base form, or as a pharmaceutically acceptable salt thereof) and any physical form thereof (e.g., including non-solid forms (e.g., liquid or semi-solid forms), and solid forms (e.g., amorphous or crystalline forms, specific polymorphic forms, solvates, including hydrates (e.g., mono-, di- and hemi- hydrates)), and mixtures of various forms.

As used herein, the term "optionally substituted" means that the groups may be either unsubstituted or substituted with one or more of the specified substituents.

The alternative definitions for the various groups and substituent groups of Formula I provided throughout the specification are intended to particularly describe each compound species disclosed herein, individually, as well as groups of one or more compound species. The scope of this invention includes any combination of these group and substituent group definitions.

The invention is further directed to a compound according to Formula I, wherein:

R¹ is (C₁-C₄)alkyl;

R² is H or halogen;

R³ is H, halogen, (C₁-C₄)alkoxy, (C₅-C₆)cycloalkyloxy, (C₁-C₄)haloalkoxy, (C₁-C₄)alkylthio-, (C₁-C₄)haloalkylthio-, (C₅-C₆)cycloalkylthio-, (C₁-C₄)alkylsulfonyl, (C₁-C₄)haloalkylsulfonyl, (C₅-C₆)cycloalkylsulfonyl, or -N(R^a)(R^b);

each R^a is independently (C₁-C₄)alkyl, wherein said (C₁-C₄)alkyl is optionally substituted one to three times, independently, by halogen, hydroxyl, (C₁-C₄)alkoxy, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, -CO₂H, -CO₂(C₁-C₄)alkyl, -CONH₂, -CONH(C₁-C₄)alkyl, or -CON((C₁-C₄)alkyl)((C₁-C₄)alkyl), and

R^b is (C₁-C₄)alkyl;

or R^a and R^b taken together with the nitrogen atom to which they are attached form a 5-membered or 6-membered heterocyclic ring, optionally containing one additional heteroatom selected from nitrogen, oxygen and sulfur, wherein said ring is optionally substituted one or two times, independently, by halogen, (C₁-C₄)alkyl, (C₁-C₄)haloalkyl, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, hydroxyl, hydroxy(C₁-C₄)alkyl-, oxo, (C₁-C₄)alkoxy, (C₁-C₄)haloalkoxy, or (C₁-C₄)alkoxy(C₁-C₄)alkyl;

R⁴ is H;

R⁵, R⁶, and R⁷ are each, independently, H, halogen, nitro, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, hydroxyl, or OR^c; and

each R^c is, independently, (C₁-C₄)alkyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, piperidinyl, piperazinyl, morpholinyl, tetrahydropyranyl, tetrahydro-4*H*-1,4-thiazinyl, or 1,4-dioxanyl, wherein said (C₁-C₄)alkyl is optionally substituted by

halogen, hydroxyl, trifluoromethyl, (C₁-C₄)alkoxy, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, piperidinyl, piperazinyl, morpholinyl, tetrahydropyranyl, tetrahydro-4*H*-1,4-thiazinyl, or 1,4-dioxanyl;

- 5 or R⁶ and R⁷ taken together represent -O(C₁-C₂)alkylO-;
or a salt, particularly a pharmaceutically acceptable salt, thereof.

In another embodiment of this invention:

R¹ is (C₁-C₄)alkyl;

- 10 R² is H;

R³ is H, halogen, (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, hydroxyl, hydroxy(C₁-C₈)alkyl-, (C₁-C₈)alkoxy, (C₃-C₈)cycloalkyloxy, (C₁-C₈)haloalkoxy, (C₁-C₈)alkylthio-, (C₁-C₈)haloalkylthio-, (C₃-C₈)cycloalkylthio-, phenyl, 5-membered heteroaryl, or -N(R^a)(R^b), wherein said phenyl or heteroaryl is optionally substituted one to three times,

- 15 independently, by halogen, (C₁-C₆)alkyl, (C₁-C₄)haloalkyl, (C₁-C₆)alkoxy, (C₁-C₄)haloalkoxy, hydroxy(C₁-C₄)alkyl-, or -N(R^a)(R^b);

each R^a is independently (C₁-C₄)alkyl, wherein said (C₁-C₄)alkyl is optionally substituted one to three times, independently, by halogen, hydroxyl, (C₁-C₆)alkoxy, amino, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)((C₁-C₆)alkyl)amino, -CO₂H, -CO₂(C₁-C₆)alkyl, -CONH₂,
20 -CONH(C₁-C₆)alkyl, or -CON((C₁-C₆)alkyl)((C₁-C₆)alkyl), and

R^b is (C₁-C₄)alkyl;

or R^a and R^b taken together with the nitrogen atom to which they are attached form a 5-membered or 6-membered heterocyclic ring, optionally containing one additional heteroatom selected from nitrogen, oxygen and sulfur, wherein said ring is optionally
25 substituted one or two times, independently, by halogen, (C₁-C₄)alkyl, (C₁-C₄)haloalkyl, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, hydroxyl, hydroxy(C₁-C₄)alkyl-, oxo, (C₁-C₄)alkoxy, (C₁-C₄)haloalkoxy, or (C₁-C₄)alkoxy(C₁-C₄)alkyl;

R⁴ is H;

R⁵ is H;

- 30 R⁶ and R⁷ are each OR^c; and

each R^c is, independently, (C₁-C₈)alkyl, optionally substituted one to three times, independently, by halogen, hydroxyl, (C₁-C₄)alkoxy, amino, (C₁-C₄)alkylamino, or ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino;

or a salt, particularly a pharmaceutically acceptable salt, thereof.

35

In another embodiment of this invention:

R¹ is (C₁-C₃ alkyl);

R² is H;

R³ is H, halogen, C₁-C₆ alkyl, C₁-C₆ haloalkyl, hydroxyC₁-C₆ alkyl-, C₁-C₆ alkoxy, (C₃-C₆)cycloalkyloxy, C₁-C₆ haloalkoxy, C₁-C₆ alkylthio-, (C₃-C₆)cycloalkylthio-,

5 C₁-C₆ haloalkylthio-, phenyl, 5-membered heteroaryl, or -N(R^a)(R^b),

wherein said heteroaryl contains one heteroatom selected from N, O and S, or contains one nitrogen atom and optionally contains 1 additional heteroatom selected from N, O and S, or contains two nitrogen atoms and optionally contains 1 additional heteroatom selected from N, O and S; and said phenyl or heteroaryl is optionally
10 substituted one to three times, independently, by halogen, (C₁-C₆)alkyl, (C₁-C₄)haloalkyl, or -N(R^a)(R^b);

each R^a is independently (C₁-C₄)alkyl, wherein said (C₁-C₄)alkyl is optionally substituted by hydroxyl, trifluoromethyl, (C₁-C₆)alkoxy, amino, (C₁-C₆)alkylamino, or ((C₁-C₆)alkyl)((C₁-C₆)alkyl)amino, and

15 R^b is (C₁-C₄)alkyl;

or R^a and R^b taken together with the nitrogen atom to which they are attached form a 5-membered or 6-membered heterocyclic ring, optionally containing one additional heteroatom selected from nitrogen, oxygen and sulfur, wherein said ring is optionally substituted one or two times, independently, by (C₁-C₄)alkyl, (C₁-C₄)haloalkyl,
20 hydroxy(C₁-C₄)alkyl-, oxo, or (C₁-C₄)alkoxy(C₁-C₄)alkyl;

R⁴ is H;

R⁵ is H;

R⁶ and R⁷ are each OR^c; and

each R^c is, independently, (C₁-C₄)alkyl, optionally substituted one to three times,
25 independently, by halogen;

or a salt, particularly a pharmaceutically acceptable salt, thereof.

In a specific embodiment of this invention, R¹ is -CH₃.

In another specific embodiment of this invention, R² is H or F. In a further specific
30 embodiment of this invention, R² is H.

In another embodiment of the invention, R³ is H, halogen, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, (C₁-C₆)alkoxy, (C₅-C₆)cycloalkyloxy, (C₁-C₆)haloalkoxy, (C₁-C₆)alkylthio-, (C₅-C₆)cycloalkylthio-, (C₁-C₆)haloalkylthio-, (C₁-C₆)alkylsulfonyl, (C₁-C₆)haloalkylsulfonyl, (C₅-C₆)cycloalkylsulfonyl, phenyl, 5-membered heteroaryl, or -N(R^a)(R^b), wherein said
35 heteroaryl contains one heteroatom selected from N, O and S, or contains one nitrogen atom and optionally contains 1 additional heteroatom selected from N, O and S, or

contains two nitrogen atoms and optionally contains 1 additional heteroatom selected from N, O and S; and said phenyl or heteroaryl is optionally substituted one to three times, independently, by halogen, (C₁-C₆)alkyl, (C₁-C₄)haloalkyl, or -N(R^a)(R^b).

In a further embodiment of this invention, R³ is H, halogen, (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, hydroxyl, hydroxy(C₁-C₈)alkyl-, (C₁-C₈)alkoxy, (C₃-C₈)cycloalkyloxy, (C₁-C₈)haloalkoxy, (C₁-C₈)alkylthio-, (C₁-C₈)haloalkylthio-, (C₃-C₈)cycloalkylthio-, phenyl, 5-membered heteroaryl, or -N(R^a)(R^b), wherein said phenyl or heteroaryl is optionally substituted one to three times, independently, by halogen, (C₁-C₆)alkyl, (C₁-C₄)haloalkyl, (C₁-C₆)alkoxy, (C₁-C₄)haloalkoxy, hydroxy(C₁-C₄)alkyl-, or -N(R^a)(R^b).

In yet a further embodiment of this invention, R³ is H, halogen, C₁-C₆ alkyl, C₁-C₆ haloalkyl, hydroxyC₁-C₆ alkyl-, C₁-C₆ alkoxy, (C₃-C₆)cycloalkyloxy, C₁-C₆ haloalkoxy, C₁-C₆ alkylthio-, (C₃-C₆)cycloalkylthio-, C₁-C₆ haloalkylthio-, phenyl, 5-membered heteroaryl, or -N(R^a)(R^b), wherein said heteroaryl contains one heteroatom selected from N, O and S, or contains one nitrogen atom and optionally contains 1 additional heteroatom selected from N, O and S, or contains two nitrogen atoms and optionally contains 1 additional heteroatom selected from N, O and S; and said phenyl or heteroaryl is optionally substituted one to three times, independently, by halogen, (C₁-C₆)alkyl, (C₁-C₄)haloalkyl, or -N(R^a)(R^b).

In yet a further embodiment of this invention, R³ is H, halogen, (C₁-C₄)alkoxy, (C₅-C₆)cycloalkyloxy, (C₁-C₄)haloalkoxy, (C₁-C₄)alkylthio-, (C₁-C₄)haloalkylthio-, (C₅-C₆)cycloalkylthio-, (C₁-C₄)alkylsulfonyl, (C₁-C₄)haloalkylsulfonyl, (C₅-C₆)cycloalkylsulfonyl, or -N(R^a)(R^b).

In a specific embodiment of this invention, each R^a is independently -CH₃, -CH₂CH₃, or -CH₂CF₃, and R^b is -CH₃; or R^a and R^b taken together with the nitrogen to which they are attached represent pyrrolidin-1-yl, piperidin-1-yl, or morpholin-4-yl, wherein said pyrrolidin-1-yl, piperidin-1-yl, or morpholin-4-yl is optionally substituted one or two times, independently, by F, -CH₃, or -CF₃.

In another specific embodiment of this invention, R³ is H, F, Cl, -OCH₃, -OCH(CH₃)₂, -OCF₃, -OCH₂CF₃, -OCH(CH₃)CF₃, -SCH₃, -SCH₂CF₃, -SO₂CH₃, -N(CH₃)₂, -N(CH₃)CH₂CH₃, -N(CH₃)CH₂CF₃, 2-(trifluoromethyl)pyrrolidin-1-yl, 2,5-(dimethyl)pyrrolidin-1-yl, piperidin-1-yl, 4,4-difluoro-piperidin-1-yl, or morpholin-4-yl.

In a further embodiment of this invention, R⁵, R⁶, and R⁷ are each, independently, H, halogen, nitro, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, hydroxyl, or OR^c, and each R^c is, independently, (C₁-C₄)alkyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, piperidinyl, piperazinyl, morpholinyl, tetrahydropyranyl, tetrahydro-4*H*-1,4-thiazinyl, or 1,4-dioxanyl, wherein said (C₁-C₄)alkyl is optionally substituted by

halogen, hydroxyl, trifluoromethyl, (C₁-C₄)alkoxy, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, piperidinyl, piperazinyl, morpholinyl, tetrahydropyranyl, tetrahydro-4*H*-1,4-thiazinyl, or 1,4-dioxanyl; or R⁶ and R⁷ taken together represent -O(C₁-C₂)alkylO-.

5 In a specific embodiment of this invention, R⁵ is H, Cl, -OCH₃, -OCH₂CH₃, -OCH(CH₃)₂, -O(CH₂)₂OCH₃, or -O(tetrahydropyran-4-yl).

In another specific embodiment of this invention, R⁵ is H.

In another embodiment of this invention, R⁶ and R⁷ are each OR^c; and each R^c is, independently, (C₁-C₈)alkyl, optionally substituted one to three times, independently, by
10 halogen, hydroxyl, (C₁-C₄)alkoxy, amino, (C₁-C₄)alkylamino, or ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino.

In a further embodiment of this invention, R⁶ and R⁷ are each OR^c; and each R^c is, independently, (C₁-C₄)alkyl, optionally substituted one to three times, independently, by halogen.

15 In yet a further embodiment of this invention, R⁶ and R⁷ are each OR^c; and each R^c is, independently, (C₁-C₄)alkyl, optionally substituted by (C₁-C₄)alkoxy.

In a specific embodiment of this invention, R⁶ is H, Cl, I, -OCH₃, -OCH₂CH₃, -OCH(CH₃)₂, -O(CH₂)₂OCH₃, -O(CH₂)₃-morpholin-4-yl, -NH₂, -NHCH₃, or -N(CH₃)₂.

In another specific embodiment of this invention, R⁶ is H.

20 In another specific embodiment of this invention, R⁷ is H, -OH, -OCH₃, -OCH₂CH₃, -OCH(CH₃)₂, -O(CH₂)₃Cl, -O(CH₂)₂OCH₃, -O(CH₂)₃OCH₃, -O(CH₂)₃-morpholin-4-yl, -NH₂, -NHCH₃, or -NO₂.

In another specific embodiment of this invention, R⁶ and R⁷ taken together represent -O(CH₂)₂O-.

25 Specific compounds of this invention are:

3-{[6,7-bis(methyloxy)-4-quinazolinyl]amino}-4-(dimethylamino)-*N*-methylbenzenesulfonamide;

3-{[6,7-bis(methyloxy)-4-quinazolinyl]amino}-*N*-methylbenzenesulfonamide;

30 3-{[6,7-bis(methyloxy)-4-quinazolinyl]amino}-*N*-methyl-4-(4-morpholinyl)benzenesulfonamide;

3-{[6,7-bis(methyloxy)-4-quinazolinyl]amino}-4-[ethyl(methyl)amino]-*N*-methylbenzenesulfonamide;

3-{[6,7-bis(methyloxy)-4-quinazolinyl]amino}-*N*-methyl-4-[(1-methylethyl)oxy]benzenesulfonamide;

35 3-{[6,7-bis(methyloxy)-4-quinazolinyl]amino}-*N*-methyl-4-(methyloxy)benzenesulfonamide;

3-{{6,7-bis(methyloxy)-4-quinazoliny]amino}-N-methyl-4-(methylthio)benzenesulfonamide;

4-(dimethylamino)-N-methyl-3-(4-quinazolinylamino)benzenesulfonamide;

3-[(6-chloro-4-quinazoliny]amino]-4-(dimethylamino)-N-methylbenzenesulfonamide;

4-(dimethylamino)-N-methyl-3-{{6-(methyloxy)-4-quinazoliny]amino}benzenesulfonamide;

3-{{6,7-bis(methyloxy)-4-quinazoliny]amino}-N-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

3-[(6-chloro-4-quinazoliny]amino]-N-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

3-(7,8-dihydro[1,4]dioxino[2,3-g]quinazolin-4-ylamino)-4-(dimethylamino)-N-methylbenzenesulfonamide;

3-{{6,7-bis(methyloxy)-4-quinazoliny]amino}-4-(4,4-difluoro-1-piperidiny)-N-methylbenzenesulfonamide;

3-[(6-chloro-4-quinazoliny]amino]-4-(4,4-difluoro-1-piperidiny)-N-methylbenzenesulfonamide;

3-{{7-[(3-chloropropyl)oxy]-4-quinazoliny]amino}-4-(dimethylamino)-N-methylbenzenesulfonamide;

3-{{6,7-bis[[2-(methyloxy)ethyl]oxy]-4-quinazoliny]amino}-4-(dimethylamino)-N-methylbenzenesulfonamide;

3-{{6,7-bis[[2-(methyloxy)ethyl]oxy]-4-quinazoliny]amino}-N-methyl-4-(methyloxy)benzenesulfonamide;

3-{{6,7-bis[[2-(methyloxy)ethyl]oxy]-4-quinazoliny]amino}-N-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

3-{{6,7-bis[[2-(methyloxy)ethyl]oxy]-4-quinazoliny]amino}-N-methyl-4-(methylthio)benzenesulfonamide;

3-{{6,7-bis(methyloxy)-4-quinazoliny]amino}-N-methyl-4-[(2R)-2-(trifluoromethyl)-1-pyrrolidiny]benzenesulfonamide;

3-[(6-chloro-4-quinazoliny]amino]-N-methyl-4-[(2R)-2-(trifluoromethyl)-1-pyrrolidiny]benzenesulfonamide;

5-{{6,7-bis(methyloxy)-4-quinazoliny]amino}-2-fluoro-N-methylbenzenesulfonamide;

3-{{6,7-bis(methyloxy)-4-quinazoliny]amino}-N-methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-(1-piperidinyl)benzenesulfonamide;

3-[(6-chloro-4-quinazoliny)amino]-*N*-methyl-4-(1-piperidinyl)benzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-4-(2,5-dimethyl-1-pyrrolidinyl)-*N*-methylbenzenesulfonamide;

3-[(6-chloro-4-quinazoliny)amino]-4-(2,5-dimethyl-1-pyrrolidinyl)-*N*-methylbenzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)thio]benzenesulfonamide;

3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methylbenzenesulfonamide;

3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-4-(dimethylamino)-*N*-methylbenzenesulfonamide;

3-(5-chloro-6,7-dimethoxyquinazolin-4-ylamino)-*N*-methylbenzenesulfonamide;

4-methoxy-3-(7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-ylamino)-*N*-methylbenzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[methyl(2,2,2-trifluoroethyl)amino]benzenesulfonamide;

3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[methyl(2,2,2-trifluoroethyl)amino]benzenesulfonamide;

3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-(methyloxy)benzenesulfonamide;

3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-4-fluoro-*N*-methylbenzenesulfonamide;

3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-4-chloro-*N*-methylbenzenesulfonamide;

3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-(4-morpholinyl)benzenesulfonamide;

5-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-2-fluoro-*N*-methyl-4-(methyloxy)benzenesulfonamide;

3-[[6-(ethyloxy)-7-(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

4-(dimethylamino)-3-(7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-ylamino)-*N*-methylbenzenesulfonamide;

N-methyl-3-{{6-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazoliny]amino}-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

3-{{7-[(3-chloropropyl)oxy]-4-quinazoliny]amino}-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

5 3-{{5-(ethyloxy)-7-(methyloxy)-4-quinazoliny]amino}-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

N-methyl-3-{{5-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazoliny]amino}-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

10 *N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]-3-{{5,6,7-tris(methyloxy)-4-quinazoliny]amino}benzenesulfonamide;

3-{{5,7-bis(methyloxy)-4-quinazoliny]amino}-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide;

5-{{5,7-bis(methyloxy)-4-quinazoliny]amino}-2-fluoro-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

15 5-{{5,7-bis(methyloxy)-4-quinazoliny]amino}-2-fluoro-*N*-methyl-4-(methyloxy)benzenesulfonamide;

5-{{6,7-bis(methyloxy)-4-quinazoliny]amino}-2-fluoro-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

20 3-{{5,7-bis(methyloxy)-4-quinazoliny]amino}-*N*-methyl-4-(methylthio)benzenesulfonamide;

3-{{5,7-bis(methyloxy)-4-quinazoliny]amino}-*N*-methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide;

N-methyl-4-[(trifluoromethyl)oxy]-3-{{5,6,7-tris(methyloxy)-4-quinazoliny]amino}benzenesulfonamide;

25 5-{{5,7-bis(methyloxy)-4-quinazoliny]amino}-4-(dimethylamino)-2-fluoro-*N*-methylbenzenesulfonamide;

5-{{6,7-bis(methyloxy)-4-quinazoliny]amino}-4-(dimethylamino)-2-fluoro-*N*-methylbenzenesulfonamide;

30 2-fluoro-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]-5-{{5,6,7-tris(methyloxy)-4-quinazoliny]amino}benzenesulfonamide;

4-chloro-*N*-methyl-3-{{5,6,7-tris(methyloxy)-4-quinazoliny]amino}benzenesulfonamide;

N-methyl-3-{{7-(methyloxy)-5-[[2-(methyloxy)ethyl]oxy]-4-quinazoliny]amino}-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

35 *N*-methyl-3-{{7-(methyloxy)-5-(tetrahydro-2*H*-pyran-4-yloxy)-4-quinazoliny]amino}-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

N-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]-3-[[5,6,7-tris(methyloxy)-4-quinazolinyl]amino]benzenesulfonamide;

N-methyl-4-(methylsulfonyl)-3-[[5,6,7-tris(methyloxy)-4-quinazolinyl]amino]benzenesulfonamide;

5 3-[[5-(ethyloxy)-7-(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide;

N-methyl-3-[[5-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazolinyl]amino]-4-[(trifluoromethyl)oxy]benzenesulfonamide;

10 3-[[5-(ethyloxy)-7-(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide;

N-methyl-3-[[5-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazolinyl]amino]-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-(methylsulfonyl)benzenesulfonamide;

15 3-[[6,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide;

3-[[6-iodo-7-(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

20 3-[[6,7-bis(ethyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

4-(dimethylamino)-*N*-methyl-3-((7-nitroquinazolin-4-yl)amino)benzenesulfonamide;

4-(dimethylamino)-*N*-methyl-3-[(7-[[3-(4-morpholinyl)propyl]oxy]-4-quinazolinyl]amino]benzenesulfonamide;

25 *N*-methyl-3-[(7-[[3-(4-morpholinyl)propyl]oxy]-4-quinazolinyl]amino)-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

N-methyl-3-[(7-[[3-(methyloxy)propyl]oxy]-4-quinazolinyl]amino)-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

4-(dimethylamino)-*N*-methyl-3-[(7-[[3-(methyloxy)propyl]oxy]-4-quinazolinyl]amino]benzenesulfonamide;

30 3-[[5,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-(methylsulfonyl)benzenesulfonamide;

3-(7-hydroxy-6-methoxyquinazolin-4-ylamino)-*N*-methyl-4-(2,2,2-trifluoroethoxy)benzenesulfonamide;

35 3-(7-isopropoxy-6-methoxyquinazolin-4-ylamino)-*N*-methyl-4-(2,2,2-trifluoroethoxy)-benzenesulfonamide;

3-(7-ethoxy-6-methoxy-quinazolin-4-ylamino)-*N*-methyl-4-(2,2,2-trifluoro-ethoxy)-benzenesulfonamide;

3-[6-methoxy-7-(2-methoxy-ethoxy)-quinazolin-4-ylamino]-*N*-methyl-4-(2,2,2-trifluoro-ethoxy)-benzenesulfonamide;

5 3-[[5,7-bis(methoxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide;

3-[[5,7-bis(methoxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide;

10 3-((6-aminoquinazolin-4-yl)amino)-4-(dimethylamino)-*N*-methylbenzenesulfonamide;

3-((7-aminoquinazolin-4-yl)amino)-4-(dimethylamino)-*N*-methylbenzenesulfonamide;

4-(dimethylamino)-3-((6-(dimethylamino)quinazolin-4-yl)amino)-*N*-methylbenzenesulfonamide;

15 4-(dimethylamino)-*N*-methyl-3-((6-(methylamino)quinazolin-4-yl)amino)benzenesulfonamide; and

4-(dimethylamino)-*N*-methyl-3-((7-(methylamino)quinazolin-4-yl)amino)benzenesulfonamide;

and salts, particularly pharmaceutically acceptable salts, thereof.

20

Representative compounds of this invention include the compounds of Examples 1-88.

The compounds according to Formula I may contain one or more asymmetric center (also referred to as a chiral center) and may, therefore, exist as individual
25 enantiomers, diastereomers, or other stereoisomeric forms, or as mixtures thereof. Chiral centers, such as chiral carbon atoms, may also be present in a substituent such as an alkyl group. Where the stereochemistry of a chiral center present in Formula I, or in any chemical structure illustrated herein, is not specified the structure is intended to encompass all individual stereoisomers and all mixtures thereof. Thus, compounds
30 according to Formula I containing one or more chiral center may be used as racemic mixtures, enantiomerically enriched mixtures, or as enantiomerically pure individual stereoisomers.

Individual stereoisomers of a compound according to Formula I which contain one or more asymmetric center may be resolved by methods known to those skilled in the art.
35 For example, such resolution may be carried out (1) by formation of diastereoisomeric salts, complexes or other derivatives; (2) by selective reaction with a stereoisomer-specific

reagent, for example by enzymatic oxidation or reduction; or (3) by gas-liquid or liquid chromatography in a chiral environment, for example, on a chiral support such as silica with a bound chiral ligand or in the presence of a chiral solvent. The skilled artisan will appreciate that where the desired stereoisomer is converted into another chemical entity by one of the separation procedures described above, a further step is required to liberate the desired form. Alternatively, specific stereoisomers may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer to the other by asymmetric transformation.

When a disclosed compound or its salt is named or depicted by structure, it is to be understood that the compound or salt, including solvates (particularly, hydrates) thereof, may exist in crystalline forms, non-crystalline forms or a mixture thereof. The compound or salt, or solvates (particularly, hydrates) thereof, may also exhibit polymorphism (i.e. the capacity to occur in different crystalline forms). These different crystalline forms are typically known as "polymorphs." It is to be understood that when named or depicted by structure, the disclosed compound, or solvates (particularly, hydrates) thereof, also include all polymorphs thereof. Polymorphs have the same chemical composition but differ in packing, geometrical arrangement, and other descriptive properties of the crystalline solid state. Polymorphs, therefore, may have different physical properties such as shape, density, hardness, deformability, stability, and dissolution properties. Polymorphs typically exhibit different melting points, IR spectra, and X-ray powder diffraction patterns, which may be used for identification. One of ordinary skill in the art will appreciate that different polymorphs may be produced, for example, by changing or adjusting the conditions used in crystallizing/recrystallizing the compound.

For solvates of the compounds of the invention, or salts thereof, that are in crystalline form, the skilled artisan will appreciate that pharmaceutically-acceptable solvates may be formed wherein solvent molecules are incorporated into the crystalline lattice during crystallization. Solvates may involve nonaqueous solvents such as ethanol, isopropanol, DMSO, acetic acid, ethanolamine, and ethyl acetate, or they may involve water as the solvent that is incorporated into the crystalline lattice. Solvates wherein water is the solvent that is incorporated into the crystalline lattice are typically referred to as "hydrates." Hydrates include stoichiometric hydrates as well as compositions containing variable amounts of water. The invention includes all such solvates.

Because of their potential use in medicine, the salts of the compounds of Formula I are preferably pharmaceutically acceptable. The compounds of this invention are bases, wherein a desired salt form may be prepared by any suitable method known in the art,

including treatment of the free base with an inorganic acid, such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like, or with an organic acid, such as acetic acid, trifluoroacetic acid, maleic acid, succinic acid, mandelic acid, fumaric acid, malonic acid, pyruvic acid, oxalic acid, glycolic acid, salicylic acid, pyranosidyl acid, such as glucuronic acid or galacturonic acid, alpha-hydroxy acid, such as citric acid or tartaric acid, amino acid, such as aspartic acid or glutamic acid, aromatic acid, such as benzoic acid or cinnamic acid, sulfonic acid, such as p-toluenesulfonic acid, methanesulfonic acid, ethanesulfonic acid or the like. Examples of pharmaceutically acceptable salts include sulfates, pyrosulfates, bisulfates, sulfites, bisulfites, phosphates, chlorides, bromides, iodides, acetates, propionates, decanoates, caprylates, acrylates, formates, isobutyrate, caproates, heptanoates, propiolates, oxalates, malonates succinates, suberates, sebacates, fumarates, maleates, butyne-1,4-dioates, hexyne-1,6-dioates, benzoates, chlorobenzoates, methylbenzoates, dinitrobenzoates, hydroxybenzoates, methoxybenzoates, phthalates, phenylacetates, phenylpropionates, phenylbutyrates, citrates, lactates, γ -hydroxybutyrates, glycollates, tartrates mandelates, and sulfonates, such as xylenesulfonates, methanesulfonates, propanesulfonates, naphthalene-1-sulfonates and naphthalene-2-sulfonates.

If an inventive basic compound is isolated as a salt, the corresponding free base form of that compound may be prepared by any suitable method known to the art, including treatment of the salt with an inorganic or organic base, suitably an inorganic or organic base having a higher pK_a than the free base form of the compound.


General Methods of Preparation

The compounds of Formula I may be obtained by using synthetic procedures illustrated in the Schemes below or by drawing on the knowledge of a skilled organic chemist. The synthesis provided in these Schemes are applicable for producing compounds of the invention having a variety of different R^1 and R^2 groups employing appropriate precursors, which are suitably protected if needed, to achieve compatibility with the reactions outlined herein. Subsequent deprotection, where needed, affords compounds of the nature generally disclosed. While the Schemes are shown with compounds only of Formula I, they are illustrative of processes that may be used to make the compounds of the invention.

Compound names were generated using the software naming program ACD/Name Pro V6.02 available from Advanced Chemistry Development, Inc., 110 Yonge Street, 14th Floor, Toronto, Ontario, Canada, M5C 1T4 (<http://www.acdlabs.com/>).

As shown in Scheme 1, the compounds of Formula I can be prepared under a variety of conditions by reaction of an arylamine (e.g., Ar-NH-R², specifically, Ar-NH₂) with an activated quinazoline.


5 *Scheme 1*



a) Isopropanol, μ w (6-15 bar), 150 °C, 15 min; b) Isopropanol, μ w (1-2 bar), 100 °C, 30 min

10 As shown in Schemes 2-5, the compounds of Formula I can be prepared *via* common functional group transformations on advanced intermediates.


Scheme 2



a) morpholine, MeOH, 90 °C, 3 days; b) NaOMe, MeOH, μ w, 100 °C, 10 min

15


Scheme 3



a) NaBO₃, AcOH, 50 °C, 2 hours

20


Scheme 4



a) H_2 , Pd/C, AcOH, EtOAc, MeOH, DMF, 25 °C, 2 hours; b) RBr, K_2CO_3 , DMF, 60°C, 16 hours

5

Scheme 5



10

a) H_2 , Pd/C, MeOH, 25 °C, 2.5 hours; b) H_2CO , $\text{NaBH}(\text{OAc})_3$, AcOH, CH_2Cl_2 , 25 °C, 2 hours

The invention also includes various deuterated forms of the compounds of Formula I. Each available hydrogen atom attached to a carbon atom may be independently replaced with a deuterium atom. A person of ordinary skill in the art will know how to synthesize deuterated forms of the compounds of Formula I. For example, deuterated alkyl groups (e.g., *N*-(deuteriomethyl)amines or R^a/R^b alkyls) may be prepared by conventional techniques (see for example: methyl- d_3 -amine available from Aldrich Chemical Co., Milwaukee, WI, Cat. No. 489,689-2). Employing such compounds according to Scheme 1 will allow for the preparation of compounds of Formula I in which various hydrogen atoms of the *N*-methyl or pyrimidinyl groups are replaced with a deuterium atom.

Methods of Use

25

The present invention is directed to a method of inhibiting TNNI3K which comprises contacting the kinase with a compound of Formula I or a salt thereof, particularly a pharmaceutically acceptable salt thereof. This invention is also directed to a method of treatment of a TNNI3K-mediated disease or disorder comprising administering

an effective amount of the compound of Formula I or a salt thereof, particularly a pharmaceutically acceptable salt thereof, to a patient, specifically a human, in need thereof. As used herein, "patient" refers to a human or other mammal. Specifically, this invention is directed to a method of inhibiting TNNI3K activity, comprising contacting the
5 kinase with an effective amount of a compound of Formula I or a pharmaceutically acceptable salt thereof. For example, TNNI3K activity may be inhibited in mammalian cardiac tissue by administering to a patient in need thereof, an effective amount a compound of Formula I or a pharmaceutically acceptable salt thereof.

The compounds of this invention may be particularly useful for treatment of
10 TNNI3K-mediated diseases or disorders, specifically by inhibition of TNNI3K activity, where such diseases or disorders are selected from heart failure, particularly congestive heart failure; cardiac hypertrophy; and heart failure or congestive heart failure resulting from cardiac hypertrophy. The compounds of this invention may also be useful for the treatment of heart failure or congestive heart failure resulting from myocardial ischemia or
15 myocardial infarction.

A therapeutically "effective amount" is intended to mean that amount of a compound that, when administered to a patient in need of such treatment, is sufficient to effect treatment, as defined herein. Thus, e.g., a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof, is a quantity of an
20 inventive agent that, when administered to a human in need thereof, is sufficient to modulate or inhibit the activity of TNNI3K such that a disease condition which is mediated by that activity is reduced, alleviated or prevented. The amount of a given compound that will correspond to such an amount will vary depending upon factors such as the particular compound (e.g., the potency (pXC_{50}), efficacy (EC_{50}), and the biological half-life of the
25 particular compound), disease condition and its severity, the identity (e.g., age, size and weight) of the patient in need of treatment, but can nevertheless be routinely determined by one skilled in the art. Likewise, the duration of treatment and the time period of administration (time period between dosages and the timing of the dosages, e.g., before/with/after meals) of the compound will vary according to the identity of the mammal
30 in need of treatment (e.g., weight), the particular compound and its properties (e.g., pharmaceutical characteristics), disease or condition and its severity and the specific composition and method being used, but can nevertheless be determined by one of skill in the art.

"Treating" or "treatment" is intended to mean at least the mitigation of a disease
35 condition in a patient, where the disease condition is caused or mediated by TNNI3K. The methods of treatment for mitigation of a disease condition include the use of the

compounds in this invention in any conventionally acceptable manner, for example for prevention, retardation, prophylaxis, therapy or cure of a disease. The compounds of Formula I of this invention may be useful for the treatment of heart failure, particularly congestive heart failure. The compounds of Formula I of this invention may be useful for the treatment of cardiac hypertrophy, and heart failure or congestive heart failure resulting from cardiac hypertrophy, myocardial ischemia or myocardial infarction.

The compounds of the invention may be administered by any suitable route of administration, including both systemic administration and topical administration.

Systemic administration includes oral administration, parenteral administration,

transdermal administration, rectal administration, and administration by inhalation.

Parenteral administration refers to routes of administration other than enteral, transdermal, or by inhalation, and is typically by injection or infusion. Parenteral administration includes intravenous, intramuscular, and subcutaneous injection or infusion. Inhalation refers to administration into the patient's lungs whether inhaled through the mouth or through the nasal passages. Topical administration includes application to the skin.

The compounds of the invention may be administered once or according to a dosing regimen wherein a number of doses are administered at varying intervals of time for a given period of time. For example, doses may be administered one, two, three, or four times per day. Doses may be administered until the desired therapeutic effect is achieved or indefinitely to maintain the desired therapeutic effect. Suitable dosing regimens for a compound of the invention depend on the pharmacokinetic properties of that compound, such as absorption, distribution, and half-life, which can be determined by the skilled artisan. In addition, suitable dosing regimens, including the duration such regimens are administered, for a compound of the invention depend on the condition being treated, the severity of the condition being treated, the age and physical condition of the patient being treated, the medical history of the patient to be treated, the nature of concurrent therapy, the desired therapeutic effect, and like factors within the knowledge and expertise of the skilled artisan. It will be further understood by such skilled artisans that suitable dosing regimens may require adjustment given an individual patient's response to the dosing regimen or over time as individual patient needs change.

Treatment of TNNI3K-mediated disease conditions may be achieved using the compounds of this invention as a monotherapy, or in dual or multiple combination therapy, such as in combination with other cardiovascular agents, for example, in combination with one or more of the following agents: a beta-blocker, an ACE inhibitor, an angiotensin receptor blocker (ARB), a calcium channel blocker, a diuretic, a renin inhibitor, a centrally acting antihypertensive, a dual ACE/NEP inhibitor, an aldosterone synthase inhibitor, and

an aldosterone-receptor antagonist, which are administered in effective amounts as is known in the art.

Examples of suitable beta blockers include timolol (such as BLOCARDENTM), carteolol (such as CARTROLTM), carvedilol (such as COREGTM), nadolol (such as CORGARDTM), propanolol (such as INNOPRAN XLTM), betaxolol (such as KERLONETM),

5 penbutolol (such as LEVATOLTM), metoprolol (such as LOPRESSORTM and TOPROL-XLTM), atenolol (such as TENORMINTM), pindolol (such as VISKENTM), bisoprolol, bucindolol, esmolol, acebutolol, labetalol, nebivolol, celiprolol, sotalol, and oxprenolol.

Examples of suitable ACE inhibitors include alacepril, benazepril, benazaprilat, captopril,

10 ceronapril, cilazapril, delapril, enalapril, enalaprilat, fosinopril, lisinopril, moexipiril, moveltopril, perindopril, quinapril, quinaprilat, ramipril, ramiprilat, spirapril, temocapril, trandolapril, and zofenopril. Preferred ACE inhibitors are benazepril, enalapril, lisinopril, and ramipril. Examples of suitable angiotensin receptor blockers include candesartan, eprosartan, irbesartan, losartan, olmesartan, tasosartan, telmisartan, and valsartan.

15 Examples of suitable calcium channel blockers include dihydropyridines (DHPs) and non-DHPs. Suitable DHPs include amlodipine, felodipine, ryosidine, isradipine, lacidipine, nicardipine, nifedipine, nigulpidine, niludipine, nimodipine, nisoldipine, nitrendipine, and nivaldipine, and their pharmaceutically acceptable salts. Suitable non-DHPs are flunarizine, prenylamine, diltiazem, fendiline, gallopamil, mibefradil, anipamil, tiapamil, and

20 verampimil, and their pharmaceutically acceptable salts. A suitable diuretic is a thiazide derivative selected from amiloride, chlorothiazide, hydrochlorothiazide, methylchlorothiazide, and chlorothalidon. A suitable renin inhibitor is aliskiren. Examples of suitable centrally acting antihypertensives include clonidine, guanabenz, guanfacine and methyldopa. Examples of suitable dual ACE/NEP inhibitors include omapatrilat,

25 fasidotril, and fasidotrilat. Examples of suitable aldosterone synthase inhibitors include anastrozole, fadrozole, and exemestane. Examples of suitable aldosterone-receptor antagonists include spironolactone and eplerenone.

The invention further includes the use of compounds of the invention as an active therapeutic substance, in particular in the treatment of diseases mediated by TNNI3K.

30 Specifically, the invention includes the use of compounds of the invention in the treatment of heart failure, particularly congestive heart failure; cardiac hypertrophy; heart failure or congestive heart failure resulting from cardiac hypertrophy; and heart failure or congestive heart failure resulting from myocardial ischemia or myocardial infarction.

In another aspect, the invention includes the use of compounds of the invention in

35 the manufacture of a medicament for use in the treatment of the above disorders.

Compositions

The compounds of the invention will normally, but not necessarily, be formulated into a pharmaceutical composition prior to administration to a patient. Accordingly, in another aspect the invention is directed to pharmaceutical compositions comprising a
5 compound of the invention and a pharmaceutically-acceptable excipient.

The pharmaceutical compositions of the invention may be prepared and packaged in bulk form wherein an effective amount of a compound of the invention can be extracted and then given to the patient such as with powders, syrups, and solutions for injection. Alternatively, the pharmaceutical compositions of the invention may be prepared and
10 packaged in unit dosage form. For oral application, for example, one or more tablets or capsules may be administered. A dose of the pharmaceutical composition contains at least a therapeutically effective amount of a compound of this invention (i.e., a compound of Formula I or a salt, particularly a pharmaceutically acceptable salt, thereof). When prepared in unit dosage form, the pharmaceutical compositions may contain from 1 mg to
15 1000 mg of a compound of this invention.

The pharmaceutical compositions of the invention typically contain one compound of the invention. However, in certain embodiments, the pharmaceutical compositions of the invention contain more than one compound of the invention. In addition, the pharmaceutical compositions of the invention may optionally further comprise one or more
20 additional pharmaceutically active compounds.

As used herein, "pharmaceutically-acceptable excipient" means a material, composition or vehicle involved in giving form or consistency to the composition. Each excipient must be compatible with the other ingredients of the pharmaceutical composition when commingled such that interactions which would substantially reduce the efficacy of
25 the compound of the invention when administered to a patient and interactions which would result in pharmaceutical compositions that are not pharmaceutically-acceptable are avoided. In addition, each excipient must of course be of sufficiently high purity to render it pharmaceutically-acceptable.

The compounds of the invention and the pharmaceutically-acceptable excipient or excipients will typically be formulated into a dosage form adapted for administration to the
30 patient by the desired route of administration. Conventional dosage forms include those adapted for (1) oral administration such as tablets, capsules, caplets, pills, troches, powders, syrups, elixirs, suspensions, solutions, emulsions, sachets, and cachets; (2) parenteral administration such as sterile solutions, suspensions, and powders for
35 reconstitution; (3) transdermal administration such as transdermal patches; (4) rectal administration such as suppositories; (5) inhalation such as aerosols and solutions; and

(6) topical administration such as creams, ointments, lotions, solutions, pastes, sprays, foams, and gels.

Suitable pharmaceutically-acceptable excipients will vary depending upon the particular dosage form chosen. In addition, suitable pharmaceutically-acceptable excipients may be chosen for a particular function that they may serve in the composition. For example, certain pharmaceutically-acceptable excipients may be chosen for their ability to facilitate the production of uniform dosage forms. Certain pharmaceutically-acceptable excipients may be chosen for their ability to facilitate the production of stable dosage forms. Certain pharmaceutically-acceptable excipients may be chosen for their ability to facilitate the carrying or transporting the compound or compounds of the invention once administered to the patient from one organ, or portion of the body, to another organ, or portion of the body. Certain pharmaceutically-acceptable excipients may be chosen for their ability to enhance patient compliance.

Suitable pharmaceutically-acceptable excipients include the following types of excipients: diluents, fillers, binders, disintegrants, lubricants, glidants, granulating agents, coating agents, wetting agents, solvents, co-solvents, suspending agents, emulsifiers, sweeteners, flavoring agents, flavor masking agents, coloring agents, anti-caking agents, humectants, chelating agents, plasticizers, viscosity increasing agents, antioxidants, preservatives, stabilizers, surfactants, and buffering agents. The skilled artisan will appreciate that certain pharmaceutically-acceptable excipients may serve more than one function and may serve alternative functions depending on how much of the excipient is present in the formulation and what other ingredients are present in the formulation.

Skilled artisans possess the knowledge and skill in the art to enable them to select suitable pharmaceutically-acceptable excipients in appropriate amounts for use in the invention. In addition, there are a number of resources that are available to the skilled artisan which describe pharmaceutically-acceptable excipients and may be useful in selecting suitable pharmaceutically-acceptable excipients. Examples include Remington's Pharmaceutical Sciences (Mack Publishing Company), The Handbook of Pharmaceutical Additives (Gower Publishing Limited), and The Handbook of Pharmaceutical Excipients (the American Pharmaceutical Association and the Pharmaceutical Press).

The pharmaceutical compositions of the invention are prepared using techniques and methods known to those skilled in the art. Some of the methods commonly used in the art are described in Remington's Pharmaceutical Sciences (Mack Publishing Company).

In one aspect, the invention is directed to a solid oral dosage form such as a tablet or capsule comprising an effective amount of a compound of the invention and a diluent or

filler. Suitable diluents and fillers include lactose, sucrose, dextrose, mannitol, sorbitol, starch (e.g. corn starch, potato starch, and pre-gelatinized starch), cellulose and its derivatives (e.g. microcrystalline cellulose), calcium sulfate, and dibasic calcium phosphate. The oral solid dosage form may further comprise a binder. Suitable binders include starch (e.g. corn starch, potato starch, and pre-gelatinized starch), gelatin, acacia, sodium alginate, alginic acid, tragacanth, guar gum, povidone, and cellulose and its derivatives (e.g. microcrystalline cellulose). The oral solid dosage form may further comprise a disintegrant. Suitable disintegrants include crospovidone, sodium starch glycolate, croscarmellose, alginic acid, and sodium carboxymethyl cellulose. The oral solid dosage form may further comprise a lubricant. Suitable lubricants include stearic acid, magnesium stearate, calcium stearate, and talc.

EXAMPLES

The following examples illustrate the invention. These examples are not intended to limit the scope of the present invention, but rather to provide guidance to the skilled artisan to prepare and use the compounds, compositions, and methods of the present invention. While particular embodiments of the present invention are described, the skilled artisan will appreciate that various changes and modifications can be made without departing from the spirit and scope of the invention.

In the following experimental descriptions, the following abbreviations may be used:


Abbreviation	Meaning
aq	aqueous
brine	saturated aqueous sodium chloride
CH ₂ Cl ₂	methylene chloride
CH ₃ CN or MeCN	acetonitrile
CH ₃ SNa	sodium methyl mercaptide
d	day
DMF	<i>N,N</i> -dimethylformamide
DMSO	dimethylsulfoxide
equiv	equivalents
Et	ethyl
Et ₃ N or TEA	triethylamine
Et ₂ O	diethyl ether
EtOAc	ethyl acetate
h, hr	hour
HCl	hydrochloric acid
<i>i</i> -Pr ₂ NEt	<i>N,N'</i> -diisopropylethylamine

LCMS	liquid chromatography-mass spectroscopy
Me	methyl
MeOH or CH ₃ OH	methanol
MgSO ₄	magnesium sulfate
min	minute
MS	mass spectrum
μw	microwave
NaH	sodium hydride
NaHCO ₃	sodium bicarbonate
Na ₂ SO ₄	sodium sulfate
NH ₄ Cl	ammonium chloride
Pd/C	palladium on carbon
Ph	phenyl
rt	room temperature
satd	saturated
SPE	solid phase extraction
TFA	trifluoroacetic acid
THF	tetrahydrofuran
<i>t</i> _R	retention time

PREPARATION 1

4-fluoro-*N*-methyl-3-nitrobenzenesulfonamide

5



Step 1. 4-fluoro-3-nitrobenzenesulfonyl chloride


1-Fluoro-2-nitrobenzene (50.0 g, 0.354 mol) was added to chlorosulfonic acid (91 g, 0.778 mol) at 65 °C. The resulting mixture was then heated to 100 °C for 18 h. The mixture was cooled to room temperature, poured over ice and extracted with methylene chloride. The combined organic layers were then washed with NaHCO₃, then brine, dried over MgSO₄, filtered and concentrated *in vacuo* to afford 4-fluoro-3-nitrobenzenesulfonyl chloride (55.3 g, 65% yield) as a brown oil.

15

Step 2. 4-fluoro-*N*-methyl-3-nitrobenzenesulfonamide

To a solution of 4-fluoro-3-nitrobenzenesulfonyl chloride (43 g, 179.5 mmol) in THF (500 mL), was added triethylamine (150 mL, 1.08 mol). The mixture was cooled to -35 °C and methylamine hydrochloride (14.5 g, 215.4 mmol) in water was added dropwise. After 1 h, the mixture was warmed to room temperature and diluted with 1:1 water/ethyl acetate. The organic layer was separated and washed with saturated aqueous sodium bicarbonate, then brine, dried over MgSO₄, filtered and concentrated *in vacuo*. The crude residue was purified *via* column chromatography (20% ethyl acetate/petroleum ether) to give 4-fluoro-*N*-methyl-3-nitrobenzenesulfonamide (38 g, 90% yield) as a yellow solid.

PREPARATION 2

3-amino-*N*-methyl-4-[(1-methylethyl)oxy]benzenesulfonamideStep 1. *N*-methyl-4-[(1-methylethyl)oxy]-3-nitrobenzenesulfonamide

NaH (0.440 g, 11 mmol) was added to 20 mL of isopropanol and the resulting mixture stirred at room temperature. After 30 minutes, 4-fluoro-*N*-methyl-3-nitrobenzenesulfonamide (2.34 g, 10 mmol) was added. The reaction mixture was then stirred at room temperature overnight. The mixture was poured into EtOAc and water. The organic phase was separated, dried over Na₂SO₄, and concentrated *in vacuo* to give the crude product. Purification *via* column chromatography (1:1 petroleum ether/ethyl acetate) afforded *N*-methyl-4-[(1-methylethyl)oxy]-3-nitrobenzenesulfonamide (1.6 g, 58% yield) as a yellow solid. MS (*m/z*) 274.7 (*M*+H⁺)

Step 2. 3-amino-*N*-methyl-4-[(1-methylethyl)oxy]benzenesulfonamide

To a mixture of *N*-methyl-4-[(1-methylethyl)oxy]-3-nitrobenzenesulfonamide (1.6 g, 5.8 mmol) in ethanol (20 mL) under nitrogen, Pd/C (0.160 g) was added. The flask was then evacuated and recharged with hydrogen three times. The resulting mixture was allowed to stir under a hydrogen atmosphere overnight at room temperature. The mixture was then filtered and concentrated to afford 3-amino-*N*-methyl-4-[(1-methylethyl)oxy]benzenesulfonamide (1.1 g, 77%) as a white solid. ¹H NMR (400 MHz,


DMSO- d_6) δ 7.01 - 7.10 (m, 2H), 6.87 - 6.98 (m, 2H), 5.08 (br. s., 2H), 4.63 (dt, J = 5.93, 11.98 Hz, 1H), 2.34 - 2.41 (m, 3H), 1.29 (d, J = 6.02 Hz, 6H); MS (m/z) 244.7 ($M+H^+$)

- The following anilines were prepared from 4-fluoro-*N*-methyl-3-nitrobenzenesulfonamide using procedures analogous to those described in Preparation 2:

Aniline Product	Conditions for Step 1	MS (m/z)	^1H NMR
3-amino- <i>N</i> -methyl-4-(methoxy)benzene sulfonamide	sodium methoxide, methanol	217.0 ($M+H^+$)	^1H NMR (400 MHz, DMSO- d_6) δ 7.09 (q, J = 4.85 Hz, 1H), 7.03 (s, 1H), 6.94 (s, 2H), 5.18 (s, 2H), 3.83 (s, 3H), 2.36 (d, J = 5.02 Hz, 3H)
3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide	NaH, 2,2,2-trifluoroethanol	285.0 ($M+H^+$)	^1H NMR (400 MHz, DMSO- d_6) δ ppm 7.18 (q, J =5.02 Hz, 1 H), 7.05 - 7.13 (m, 2 H), 6.93 (dd, J =8.53, 2.26 Hz, 1 H), 5.25 (s, 2 H), 4.81 (q, J =8.87 Hz, 2 H), 2.38 (d, J =4.77 Hz, 3 H)
3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide	NaH, 1,1,1-trifluoro-2-propanol	299.0 ($M+H^+$)	^1H NMR (400 MHz, DMSO- d_6) δ ppm 7.14 - 7.22 (m, 2 H), 7.11 (d, J =2.26 Hz, 1 H), 6.92 (dd, J =8.41, 2.38 Hz, 1 H), 5.19 - 5.30 (m, 3 H), 2.39 (d, J =5.02 Hz, 3 H), 1.45 (d, J =6.27 Hz, 3 H)

10

PREPARATION 3

3-amino-*N*-methyl-4-(methylthio)benzenesulfonamide

Step 1. *N*-methyl-4-(methylthio)-3-nitrobenzenesulfonamide

To a solution of 4-fluoro-*N*-methyl-3-nitrobenzenesulfonamide (15 g, 64.01 mmol) in THF (150 mL), was added 20% CH₃SNa (22.4 g, 64.01 mmol) dropwise. The resulting mixture was then stirred overnight. In the morning, the mixture was poured into ethyl acetate and water, the organic phase separated, dried over Na₂SO₄, filtered and concentrated. The crude material was then purified *via* column chromatography (1:1 ethyl acetate/petroleum ether) to afford *N*-methyl-4-(methylthio)-3-nitrobenzenesulfonamide (3.29 g, 19%) as a yellow solid. MS (m/z) 262.7 (M+H⁺)


Step 2. 3-amino-*N*-methyl-4-(methylthio)benzenesulfonamide

To a solution of *N*-methyl-4-(methylthio)-3-nitrobenzenesulfonamide (1.0 g, 3.81 mmol) in 10 mL of ethanol and 10 mL of NH₄Cl, zinc dust (2.5 g, 3.81 mmol) was added. The reaction mixture was stirred overnight at room temperature. The mixture was then filtered and diluted with ethyl acetate and water. The organic phase was separated, washed with water and brine, dried over MgSO₄, filtered and concentrated to afford 3-amino-*N*-methyl-4-(methylthio)benzenesulfonamide (0.500 g, 56%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.06 (d, *J* = 8.03 Hz, 1H), 6.86 (s, 1H), 6.67 - 6.76 (m, 1H), 5.28 (br. s., 2H), 2.17 (s, 3H), 2.21 (s, 3H); MS (m/z) 232.7 (M+H⁺)

The following anilines were prepared from 4-fluoro-*N*-methyl-3-nitrobenzenesulfonamide using procedures analogous to those described in Preparation 3:

Aniline Product	Conditions	MS (m/z)	¹ H NMR
3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)thio]benzenesulfonamide	Step 1: 2,2,2-trifluoroethane thiol, NaH Step 2: NiCl ₂ , NaBH ₄ , MeOH	300.7 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 7.48 (d, <i>J</i> =7.94 Hz, 1 H), 7.33 (q, <i>J</i> =5.00 Hz, 1 H), 7.14 (d, <i>J</i> =1.98 Hz, 1 H), 6.85 (dd, <i>J</i> =8.05, 2.09 Hz, 1 H), 5.87 (s, 2 H), 3.72 (q, <i>J</i> =10.36 Hz, 2 H), 2.39 (d, <i>J</i> =5.07 Hz, 3 H)

PREPARATION 4

3-amino-*N*-methyl-4-(4-morpholinyl)benzenesulfonamide

5

Step 1. *N*-methyl-4-(4-morpholinyl)-3-nitrobenzenesulfonamide

To a solution of 4-fluoro-*N*-methyl-3-nitrobenzenesulfonamide (2.00 g, 8.54 mmol) and morpholine (0.744 g, 8.54 mmol) in THF (100 mL), was added *N,N*-diisopropylethylamine (2.21 g, 17.08 mmol). The resulting solution was stirred at 50 °C overnight. In the morning, the reaction mixture was cooled to room temperature and concentrated to dryness *in vacuo*. The residue was dissolved in ethyl acetate and washed with water and brine, dried over MgSO₄, filtered and concentrated *in vacuo* to obtain *N*-methyl-4-(4-morpholinyl)-3-nitrobenzenesulfonamide (2.5 g, 97%) as a red oil. MS (m/z) 302.0 (M+H⁺)

15

Step 2. 3-amino-*N*-methyl-4-(4-morpholinyl)benzenesulfonamide

To a mixture of *N*-methyl-4-(4-morpholinyl)-3-nitrobenzenesulfonamide (2.5 g, 8.30 mmol) in THF (100 mL) under nitrogen, Pd/C (0.8 g) was added. The flask was then evacuated and recharged with hydrogen three times. The resulting mixture was allowed to stir under a hydrogen atmosphere at 50 °C overnight. The mixture was then filtered and concentrated to afford 3-amino-*N*-methyl-4-(4-morpholinyl)benzenesulfonamide (1.98 g, 88%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.07 - 7.17 (m, 2H), 7.01 (d, *J* = 8.28 Hz, 1H), 6.94 (dd, *J* = 1.88, 8.16 Hz, 1H), 5.20 (s, 2H), 3.72 - 3.81 (m, 4H), 2.80 - 2.89 (m, 4H), 2.38 (d, *J* = 4.77 Hz, 3H); MS (m/z) 272.2 (M+H⁺)


25

The following anilines were prepared from 4-fluoro-*N*-methyl-3-nitrobenzenesulfonamide and the indicated amine using the procedures described in Preparation 4:

30

Aniline Product	Amine / Conditions in Step 1	MS (m/z)	¹ H NMR
3-amino-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide	dimethyl amine	230.2 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.03 - 7.10 (m, 2H), 7.00 (d, <i>J</i> = 8.28 Hz, 1H), 6.93 (dd, <i>J</i> = 2.13, 8.16 Hz, 1H), 5.13 (s, 2H), 2.62 (s, 6H), 2.38 (d, <i>J</i> = 5.02 Hz, 3H)
3-amino-4-[ethyl(methyl)amino]- <i>N</i> -methylbenzenesulfonamide	ethyl(methyl) amine	244.1 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.06 - 7.13 (m, 2H), 7.02 (d, <i>J</i> = 8.28 Hz, 1H), 6.93 (dd, <i>J</i> = 1.76, 8.03 Hz, 1H), 5.11 (s, 2H), 2.89 (q, <i>J</i> = 7.03 Hz, 2H), 2.60 (s, 3H), 2.39 (d, <i>J</i> = 5.02 Hz, 3H), 1.03 (t, <i>J</i> = 7.03 Hz, 3H)
3-amino-4-(4,4-difluoro-1-piperidinyl)- <i>N</i> -methylbenzenesulfonamide	4,4-difluoro-piperidine	306.1 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 7.15 (br. s., 1 H), 7.10 (d, <i>J</i> =2.01 Hz, 1 H), 7.05 (d, <i>J</i> =8.28 Hz, 1 H), 6.92 (dd, <i>J</i> =8.16, 2.13 Hz, 1 H), 5.29 (s, 2 H), 2.95 (t, <i>J</i> =4.77 Hz, 4 H), 2.39 (s, 3 H), 2.10 - 2.25 (m, 4 H)
3-amino- <i>N</i> -methyl-4-[(2 <i>R</i>)-2-(trifluoromethyl)-1-pyrrolidinyl]benzene sulfonamide	(2 <i>R</i>)-2-(trifluoromethyl) pyrrolidine (conditions: dioxane, 110 °C, sealed tube)	324.1 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 7.31 (d, <i>J</i> =8.28 Hz, 1 H), 7.13 - 7.22 (m, 1 H), 7.10 (d, <i>J</i> =2.26 Hz, 1 H), 6.91 (dd, <i>J</i> =8.28, 2.26 Hz, 1 H), 5.20 (s, 2 H), 4.40 - 4.53 (m, 1 H), 3.49 - 3.59 (m, 1 H), 2.64 - 2.78 (m, 1 H), 2.42 (s, 3 H), 2.27 - 2.37 (m, 1 H), 1.84 - 2.04 (m, 3 H)
3-amino- <i>N</i> -methyl-4-(1-piperidinyl)benzenesulfonamide	piperidine	270.1 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 7.11 (br. s., 1 H), 7.08 (d, <i>J</i> =2.01 Hz, 1 H), 6.99 (d, <i>J</i> =8.28 Hz, 1 H), 6.91 - 6.96 (m, 1 H), 5.09 (s, 2 H), 2.74 - 2.86 (m, 4 H), 2.38 (s, 3 H), 1.64 - 1.72 (m, 4 H), 1.48 - 1.58 (m, 2 H)
3-amino-4-(2,5-dimethyl-1-pyrrolidinyl)- <i>N</i> -methylbenzenesulfonamide	2,5-dimethylpyrrolidine	284.1 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 7.29 (d, <i>J</i> =8.03 Hz, 1 H), 7.19 (br. s., 1 H), 7.09 (d, <i>J</i> =2.26 Hz, 1 H), 6.92 (dd, <i>J</i> =8.16, 2.13 Hz, 1 H), 5.38 (s, 2 H), 3.04 - 3.14 (m, 2 H), 2.41 (s, 3 H), 1.95 - 2.07 (m, 2 H), 1.42 - 1.55 (m, 2 H), 0.88 (d, <i>J</i> =6.02 Hz, 6 H)

PREPARATION 5


3-amino-4-fluoro-*N*-methylbenzenesulfonamide

5

To a mixture of 4-fluoro-*N*-methyl-3-nitrobenzenesulfonamide (1.6 g, 6.83 mmol) in THF (50 mL) under nitrogen, Pd/C (0.600 g) was added. The flask was then evacuated and recharged with hydrogen. The resulting mixture was allowed to stir under a hydrogen atmosphere overnight at 50 °C. The mixture was then filtered and concentrated to afford 3-amino-4-fluoro-*N*-methylbenzenesulfonamide (1.25 g, 89%) as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.26 (q, *J* = 4.85 Hz, 1H), 7.13 - 7.22 (m, 2H), 6.90 (ddd, *J* = 2.38, 4.27, 8.41 Hz, 1H), 5.63 (s, 2H), 2.40 (d, *J* = 5.02 Hz, 3H); MS (*m/z*) 205.1 (*M*+H⁺)

10

PREPARATION 6

3-amino-4-chloro-*N*-methylbenzenesulfonamide

15

Step 1. 4-chloro-*N*-methyl-3-nitrobenzenesulfonamide

A solution of 4-chloro-3-nitrobenzenesulfonyl chloride (10 g, 39.1 mmol) in THF (100 mL) was cooled to -40 °C before being treated with a solution of methylamine hydrochloride (2.64 g, 39.1 mmol) in 10 ml of water followed by TEA (5.44 mL, 39.1 mmol). The reaction mixture was stirred and allowed to warm to room temperature over 1 hour before being partitioned between 350 mL EtOAc and 30 mL brine. The organic layer was washed twice with brine, dried over MgSO₄ and subjected to flash chromatography (330 g silica gel, 0-40% EtOAc/hexane) to afford 4-chloro-*N*-methyl-3-nitrobenzenesulfonamide (6.38 g, 65%) as a light yellow solid. MS (*m/z*) 251.0 (*M*+H⁺).

25

Step 2. 3-amino-4-chloro-*N*-methylbenzenesulfonamide


A solution of 4-chloro-*N*-methyl-3-nitrobenzenesulfonamide (6.35 g, 25.3 mmol) in EtOH (150 mL) and water (50.0 mL) was treated with iron (14.15 g, 253 mmol) and NH₄Cl (13.55 g, 253 mmol) and heated at 90 °C for 4 hours before being cooled and filtered

30

through celite. The filter cake was washed with EtOAc and the combined filtrate was filtered again to remove precipitated NH_4Cl before being concentrated. The resulting crude material was partitioned between 350 ml EtOAc and 50 ml saturated aqueous NaHCO_3 . The organic layer was washed with brine, dried over MgSO_4 , concentrated and subjected to flash column chromatography (330 g silica gel, 0-15% EtOAc/DCM) to afford 3-amino-4-chloro-*N*-methylbenzenesulfonamide (5.60 g, 100%) as a light yellow crystalline solid. ^1H NMR (400 MHz, MeOD) δ ppm 7.39 (d, $J=8.28$ Hz, 1 H), 7.27 (d, $J=2.26$ Hz, 1 H), 7.03 (dd, $J=8.28, 2.26$ Hz, 1 H), 2.54 (s, 3 H). MS 221.0 ($\text{M}+\text{H}^+$).

10

PREPARATION 7

3-amino-*N*-methyl-4-[methyl(2,2,2-trifluoroethyl)amino]benzenesulfonamide

15 Step 1. phenylmethyl [(4-fluoro-3-nitrophenyl)sulfonyl]methylcarbamate

A solution of 4-fluoro-*N*-methyl-3-nitrobenzenesulfonamide (2 g, 8.54 mmol) in THF (20 mL) was treated with triethylamine (2.380 mL, 17.08 mmol) followed by dropwise addition of benzyl chloroformate (3.75 mL, 11.10 mmol). The mixture was stirred at 25 °C for 5 hours before being concentrated. The residue was treated with water and extracted with CH_2Cl_2 . The organic extracts were washed (brine), dried (sodium sulfate), concentrated, and subjected to flash chromatography (25-50% EtOAc-hexanes) to give a yellow solid, which was suspended in EtOAc-hexanes, collected by filtration, and washed with hexanes to give phenylmethyl [(4-fluoro-3-nitrophenyl)sulfonyl]methylcarbamate (1 g, 32%) as a white solid. MS (m/z) 391.0 ($\text{M}+\text{Na}^+$).

25

Step 2. phenylmethyl methyl({3-nitro-4-[(2,2,2-trifluoroethyl)amino]phenyl}sulfonyl)carbamate

A solution of phenylmethyl [(4-fluoro-3-nitrophenyl)sulfonyl]methylcarbamate (1 g, 2.71 mmol) in THF (10 mL) at 25 °C was treated with 2,2,2-trifluoroethylamine (0.592 g, 5.97 mmol) and stirred for 20 hours before being concentrated to give a yellow oil, which

30

was dissolved in EtOAc / hexanes. A yellow precipitate formed, which was collected by filtration and washed with hexanes to give phenylmethyl methyl({3-nitro-4-[(2,2,2-trifluoroethyl)amino]phenyl}sulfonyl)carbamate (1.07 g, 88%) as a yellow solid. MS (m/z) 448.1 (M+H⁺).

5

Step 3. phenylmethyl methyl({4-[methyl(2,2,2-trifluoroethyl)amino]-3-nitrophenyl}sulfonyl)carbamate


A solution of phenylmethyl methyl({3-nitro-4-[(2,2,2-trifluoroethyl)amino]phenyl}sulfonyl)carbamate (1 g, 2.24 mmol) in DMF (1 mL) at 25 °C was treated with sodium hydride (0.179 g, 4.47 mmol) and stirred for 2 minutes before being treated with
 10 iodomethane (0.42 mL, 6.71 mmol). After 1 hour, the mixture was diluted with water and extracted with EtOAc. The organic extract was washed (brine), dried (sodium sulfate), concentrated, and subjected to flash chromatography (10-35% EtOAc-hexanes) to give phenylmethyl methyl({4-[methyl(2,2,2-trifluoroethyl)amino]-3-nitrophenyl}sulfonyl)
 15 carbamate (539 mg, 52%) as a yellow oil. MS (m/z) 462.1 (M+H⁺).

Step 4. 3-amino-N-methyl-4-[methyl(2,2,2-trifluoroethyl)amino]benzenesulfonamide

A solution of phenylmethyl methyl({4-[methyl(2,2,2-trifluoroethyl)amino]-3-nitrophenyl}sulfonyl)carbamate (539 mg, 1.17 mmol) in MeOH (10 mL) at 25 °C was
 20 treated with 10% palladium on carbon (124 mg, 0.117 mmol) and stirred under an atmosphere of hydrogen (balloon) overnight before being filtered through celite. The filtrate was again filtered through a 0.45 micron syringe filter and concentrated to give 3-amino-N-methyl-4-[methyl(2,2,2-trifluoroethyl)amino]benzenesulfonamide (320 mg, 92%) as a brown oil. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.14 - 7.20 (m, 2 H), 7.12 (d, *J*=2.26 Hz, 1 H), 6.95 (dd, *J*=8.28, 2.26 Hz, 1 H), 5.23 (s, 2 H), 3.82 (q, *J*=9.87 Hz, 2 H), 2.83 (s, 3 H), 2.39 (d, *J*=5.02 Hz, 3 H). MS (m/z) 298.0 (M+H⁺).

PREPARATION 8

5-amino-2-fluoro-N-methylbenzenesulfonamide



30

Step 1. 2-fluoro-5-nitrobenzenesulfonyl chloride

A mixture of 1-fluoro-4-nitrobenzene (3.0 g, 21.3 mmol) in chlorosulfonic acid (5.5 mL, 84 mmol) was stirred at 90-100 °C for 8 hours before being cooled to room temperature and slowly poured into ice water and extracted with EtOAc. The organic
5 extract was washed with saturated aq. NaHCO₃ and water, dried (sodium sulfate), and concentrated to give 2-fluoro-5-nitrobenzenesulfonyl chloride (3.2 g, 63%) as a colorless oil, which was used directly in the next step.

Step 2. 2-fluoro-*N*-methyl-5-nitrobenzenesulfonamide

10 A solution of 2-fluoro-5-nitrobenzenesulfonyl chloride (3.2 g, 12.6 mmol) in THF (30 mL) at -45 °C was treated with methylamine hydrochloride (1.0 g, 15.1 mmol) and triethylamine (2.1 mL, 15.1 mmol) and stirred for 30 minutes. The mixture was then treated with 6M aqueous HCl to adjust the pH to 3 and warmed to room temperature before being diluted with water and extracted with EtOAc. The organic extract was dried
15 (sodium sulfate), concentrated, and subjected to flash chromatography (5-20% EtOAc-petroleum ether) to give 2-fluoro-*N*-methyl-5-nitrobenzenesulfonamide as a yellow solid (3.0 g, 93%). MS (m/z) 235.1 (M+H⁺).

Step 3. 5-amino-2-fluoro-*N*-methylbenzenesulfonamide


20 A solution of 2-fluoro-*N*-methyl-5-nitrobenzenesulfonamide (3.0 g, 12.8 mmol) in MeOH (40 mL) was treated with 10% palladium on carbon (300 mg, 0.28 mmol) and stirred under hydrogen (40 psi) for 8 hours before being filtered through celite and concentrated to give 5-amino-2-fluoro-*N*-methylbenzenesulfonamide (2.5 g, 96%) as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.40 - 7.49 (m, 1 H), 7.01 - 7.09 (m, 1
25 H), 6.94 (dd, *J*=5.95, 2.87 Hz, 1 H), 6.71 - 6.77 (m, 1 H), 5.49 (br. s., 2 H), 2.45 (d, *J*=4.85 Hz, 3 H). MS (m/z) 205.1 (M+H⁺).

The following anilines were prepared from the indicated nitrobenzenes using procedures analogous to those described in Preparation 8:

Aniline Product	Nitrobenzene in Step 1	MS (m/z)	¹ H NMR
3-amino- <i>N</i> -methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide	1-nitro-2-[(trifluoromethyl)oxy]benzene	271.0 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 7.39 (q, <i>J</i> =4.77 Hz, 1 H), 7.31 (dd, <i>J</i> =8.53, 1.51 Hz, 1 H), 7.24 (d, <i>J</i> =2.26 Hz, 1 H), 6.92 (dd, <i>J</i> =8.41, 2.38 Hz, 1 H), 5.92 (s, 2 H), 2.43 (d, <i>J</i> =4.77 Hz, 3 H)
5-amino-2-fluoro- <i>N</i> -methyl-4-(methoxy)benzenesulfonamide	4-fluoro-2-(methoxy)-1-nitrobenzene	235.1 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 7.31 (br. s., 1 H), 6.96 (d, <i>J</i> =7.28 Hz, 1 H), 6.90 (d, <i>J</i> =11.91 Hz, 1 H), 4.97 (s, 2 H), 3.82 (s, 3 H), 2.40 (d, <i>J</i> =3.75 Hz, 3 H)

5

PREPARATION 9

5-amino-4-(dimethylamino)-2-fluoro-*N*-methylbenzenesulfonamide

10

Step 1. 2,4-difluoro-5-nitrobenzenesulfonyl chloride

A mixture of 2,4-difluoro-1-nitrobenzene (20 g, 126 mmol) in chlorosulfonic acid (44 g, 378 mmol) was stirred at 100 °C for 48 hours before being poured into ice-water and extracted with EtOAc. The organic extract was dried (Na₂SO₄) and concentrated, and the residue was triturated with 10% EtOAc-petroleum ether to give 2,4-difluoro-5-nitrobenzenesulfonyl chloride as a brown oil (21 g, 81%) which was used directly in the next step.

Step 2. 2,4-difluoro-*N*-methyl-5-nitrobenzenesulfonamide

A solution of 2,4-difluoro-5-nitrobenzenesulfonyl chloride (21 g, 81 mmol) in THF (400 mL) at -60 °C was treated with methylamine hydrochloride (6.6 g, 97 mmol) and then

treated dropwise with triethylamine (22.6 mL, 162 mmol). After stirring for 6 hours at -60 to -40 °C the mixture was adjusted to pH 3 with the addition of 15% aqueous HCl, diluted with water, and extracted with EtOAc. The organic extracts were dried (Na₂SO₄), concentrated, and subjected to flash chromatography (17% EtOAc-petroleum ether) to give 2,4-difluoro-*N*-methyl-5-nitrobenzenesulfonamide (8 g, 38%) as a brown solid.

¹H NMR (400 MHz, CDCl₃) δ ppm 8.66 - 8.74 (m, 1 H), 7.20 - 7.25 (m, 1 H), 4.81 - 4.91 (m, 1 H), 2.78 - 2.81 (m, 3 H).

Step 3. 4-(dimethylamino)-2-fluoro-*N*-methyl-5-nitrobenzenesulfonamide


A solution of 2,4-difluoro-*N*-methyl-5-nitrobenzenesulfonamide (8.0 g, 31.6 mmol) in CH₂Cl₂ (200 mL) at -20 °C was treated with dimethylamine hydrochloride (2.56 g, 31.6 mmol). The resulting mixture was treated dropwise with triethylamine and stirred for 1 hour before being treated with 15% aqueous HCl to adjust the pH to 3, diluted with water, and extracted with EtOAc. The organic extract was dried (sodium sulfate), concentrated, and subjected to flash chromatography (20-50% EtOAc-petroleum ether) to give 4-(dimethylamino)-2-fluoro-*N*-methyl-5-nitrobenzenesulfonamide (4.0 g, 46%) as a yellow solid. MS (m/z) 278.1 (M+H⁺).

Step 4. 5-amino-4-(dimethylamino)-2-fluoro-*N*-methylbenzenesulfonamide

A solution of 4-(dimethylamino)-2-fluoro-*N*-methyl-5-nitrobenzenesulfonamide (4.0 g, 14.3 mmol) in MeOH (100 mL) was treated with 10% Pd/C (400 mg) and stirred under H₂ (50 psi) for 16 hours before being filtered, concentrated, and subjected to flash chromatography (33-50% EtOAc-petroleum ether) to give 5-amino-4-(dimethylamino)-2-fluoro-*N*-methylbenzenesulfonamide as a white solid (2.5 g, 71%). ¹H NMR (400 MHz, CDCl₃) δ ppm 7.13 (d, *J*=7.28 Hz, 1 H), 6.75 (d, *J*=11.69 Hz, 1 H), 4.58 (q, *J*=4.85 Hz, 1 H), 3.87 (br. s., 2 H), 2.66 (d, *J*=5.51 Hz, 3 H). MS (m/z) 248.1 (M+H⁺).

PREPARATION 10

5-amino-2-fluoro-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide



Step 1. 4-fluoro-1-nitro-2-[(2,2,2-trifluoroethyl)oxy]benzene

A mixture of 2,4-difluoro-1-nitrobenzene (10 g, 62.9 mmol) and 2,2,2-trifluoroethanol (6.29 g, 62.9 mmol) in THF (100 mL) at 25 °C was treated with Cs₂CO₃ (20.5 g, 62.9 mmol) and stirred for 8 hours before being diluted with the addition of water and extracted with EtOAc. The organic extract was dried (Na₂SO₄), concentrated, and subjected to flash chromatography (3% EtOAc-petroleum ether) to give 4-fluoro-1-nitro-2-[(2,2,2-trifluoroethyl)oxy]benzene (10 g, 67%) as a yellow solid. MS (m/z) 240.0 (M+H⁺).

Step 2. 2-fluoro-5-nitro-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonyl chloride

A mixture of 4-fluoro-1-nitro-2-[(2,2,2-trifluoroethyl)oxy]benzene (10 g, 41.8 mmol) in chlorosulfonic acid (82 mL, 125.5 mmol) was stirred at 50 °C for 8 hours before being poured into ice and extracted with EtOAc. The organic extracts were dried (Na₂SO₄) and concentrated to give 2-fluoro-5-nitro-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonyl chloride (15 g, crude) as a brown oil, which was used directly in the next step.


Step 3. 2-fluoro-*N*-methyl-5-nitro-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide

A mixture of 2-fluoro-5-nitro-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonyl chloride (15 g, crude) in THF (150 mL) at -45 °C was treated with methylamine hydrochloride (5.96 g, 89 mmol) and then treated dropwise with triethylamine (12.4 mL, 89 mmol). After stirring for 1 hour at -45 °C, the mixture was adjusted to pH 3 by the addition of aqueous 3M HCl, warmed to room temperature, diluted with water, and extracted with EtOAc. The organic extract was dried (Na₂SO₄), concentrated, and subjected to flash chromatography (9-17% EtOAc-petroleum ether) to give 2-fluoro-*N*-methyl-5-nitro-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide (10 g, 72% for two steps) as a yellow solid. MS (m/z) 333.0 (M+H⁺).

Step 4. 5-amino-2-fluoro-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide

A mixture of 2-fluoro-*N*-methyl-5-nitro-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide (10 g, 30.1 mmol) in MeOH (150 mL) was treated with 10% Pd/C (1 g) and stirred under H₂ (45 psi) at 45 °C for 10 hours before being filtered. The filtrate was concentrated to give 5-amino-2-fluoro-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide (8 g, 88%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.40 (q, *J*=5.07 Hz, 1 H), 7.10 (d, *J*=11.69 Hz, 1 H), 7.05 (d, *J*=7.28 Hz, 1 H), 5.04 (s, 2 H), 4.83 (q, *J*=8.82 Hz, 2 H), 2.42 (d, *J*=4.41 Hz, 3 H). MS (m/z) 303.0 (M+H⁺).

PREPARATION 11

3-amino-*N*-methyl-4-(methylsulfonyl)benzenesulfonamide

5

Step 1. *N*-methyl-4-(methylsulfonyl)-3-nitrobenzenesulfonamide

A mixture of *N*-methyl-4-(methylthio)-3-nitrobenzenesulfonamide (1.0 g, 3.81 mmol) in AcOH (15 mL) was treated dropwise with 30% aqueous H₂O₂ (5 mL, 44 mmol) and stirred at 117 °C for 2 hours before being cooled to room temperature and diluted with water (100 mL). The precipitate that formed was collected by filtration and dried to give *N*-methyl-4-(methylsulfonyl)-3-nitrobenzenesulfonamide (1.5 g, crude) as a yellow solid, which was used directly in the next step.


Step 2. 3-amino-*N*-methyl-4-(methylsulfonyl)benzenesulfonamide

A solution of *N*-methyl-4-(methylsulfonyl)-3-nitrobenzenesulfonamide (1.5 g, crude) in EtOH (20 mL) and saturated aqueous NH₄Cl (20 mL) was treated with Zn dust (3.3 g, 51 mmol) and stirred at 50 °C overnight before being filtered (EtOAc wash). The filtrate was diluted with water and extracted with EtOAc. The organic extract was washed with water and brine, dried (MgSO₄), concentrated, and subjected to flash chromatography (33% EtOAc-petroleum ether) to give 3-amino-*N*-methyl-4-(methylsulfonyl)benzenesulfonamide (950 mg, 94% 2 steps) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.73 (d, *J*=8.28 Hz, 1 H), 7.58 (q, *J*=4.94 Hz, 1 H), 7.34 (d, *J*=1.76 Hz, 1 H), 7.01 (dd, *J*=8.53, 1.76 Hz, 1 H), 6.48 (s, 2 H), 3.18 (s, 3 H), 2.47 (d, *J*=5.02 Hz, 3 H). MS (*m/z*) 265.0 (M+H⁺).

25

PREPARATION 12

4,5-dichloro-6,7-bis(methoxy)quinazoline



30


A mixture of 5-chloro-6,7-bis(methyloxy)-4(1*H*)-quinazolinone (100 mg, 0.42 mmol, prepared according to *J. Med. Chem.* (1999) **42** 3860) in POCl₃ (2 mL) was heated at 80-90 °C for 8 hours before being concentrated. The residue was treated with saturated aqueous NaHCO₃ and extracted with EtOAc. The organic extract was dried (Na₂SO₄) and concentrated to give 4,5-dichloro-6,7-bis(methyloxy)quinazoline (70 mg, 65%) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.94 (s, 1 H), 7.56 (s, 1 H), 4.05 (s, 3 H), 3.87 (s, 3 H). MS (m/z) 258.9 (M+H⁺).

The following 4-chloroquinazolines were prepared from the indicated commercially available or known quinazolinones using procedures analogous to those described in Preparation 12:

Quinazoline Product	Quinazolinone SM	MS (m/z)	¹ H NMR
4-chloro-6,7-bis{[2-(methyloxy)ethyl]oxy}quinazoline	6,7-bis{[2-(methyloxy)ethyl]oxy}-4(1 <i>H</i>)-quinazolinone	313.0 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 8.88 (s, 1 H), 7.50 (s, 1 H), 7.46 (s, 1 H), 4.32 - 4.42 (m, 4 H), 3.75 - 3.81 (m, 4 H), 3.37 (s, 3 H), 3.36 (s, 3 H)
4-chloro-6-iodo-7-(methyloxy)quinazoline	6-iodo-7-(methyloxy)-4(1 <i>H</i>)-quinazolinone (prepared as in US 6225318 B1 (2001))	320.7 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.03 (s, 1 H), 8.67 (s, 1 H), 7.48 (s, 1 H), 4.08 (s, 3 H)

PREPARATION 13

4-chloro-6-(ethyloxy)-7-(methyloxy)quinazoline



Step 1. 6-(ethyloxy)-7-(methyloxy)-4(1*H*)-quinazolinone

A mixture of 6-hydroxy-7-(methyloxy)-4(1*H*)-quinazolinone (1 g, 5.2 mmol) in water (12 mL), MeCN (6 mL), and MeOH (1 mL) at 25 °C was treated with Cs₂CO₃ (1.7 g, 5.2 mmol). After the mixture was stirred for 30 minutes, iodoethane (0.42 mL, 5.2 mmol) was

added and the mixture was stirred at 60 °C for 40 hours before being cooled to room temperature. The solid was collected by filtration and washed with MeOH to give 6-(ethyloxy)-7-(methyloxy)-4(1*H*)-quinazolinone (560 mg, 49%). MS (*m/z*) 221.1 (*M*+*H*⁺).

5 Step 2. 4-chloro-6-(ethyloxy)-7-(methyloxy)quinazoline

A mixture of 6-(ethyloxy)-7-(methyloxy)-4(1*H*)-quinazolinone (560 mg, 2.54 mmol) in POCl₃ (5 mL, 53.6 mmol) was treated with 1 drop of DMF and heated at 100 °C for 2 hours before being concentrated. The residue was treated with saturated aqueous NaHCO₃ and extracted with CH₂Cl₂. The organic extracts were dried (sodium sulfate) and concentrated to give 4-chloro-6-(ethyloxy)-7-(methyloxy)quinazoline (416 mg, 1.74 mmol, 69%) as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.88 (s, 1 H), 7.46 (s, 1 H), 7.39 (s, 1 H), 4.26 (q, *J*=6.94 Hz, 2 H), 4.02 (s, 3 H), 1.44 (t, *J*=7.03 Hz, 3 H). MS (*m/z*) 239.0 (*M*+*H*⁺).


15 The following quinazolines were prepared using procedures analogous to those described in Preparation 13 by substituting the indicated alkyl halide in the place iodoethane in Step 1:

Quinazoline Product	Alkyl Halide in Step 1	MS (<i>m/z</i>)	¹ H NMR
4-chloro-6-[(1-methylethyl)oxy]-7-(methyloxy)quinazoline	2-bromopropane	253.0 (<i>M</i> + <i>H</i> ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 8.88 (s, 1 H), 7.46 (s, 1 H), 7.41 (s, 1 H), 4.86 - 4.97 (m, 1 H), 4.01 (s, 3 H), 1.38 (d, <i>J</i> =6.02 Hz, 6 H)

20

PREPARATION 14

4-chloro-5-(ethyloxy)-7-(methyloxy)quinazoline



25

Step 1. 5-(ethyloxy)-7-(methyloxy)-4(1*H*)-quinazolinone

A mixture of [5-hydroxy-7-(methyloxy)-4-oxo-3(4*H*)-quinazolinyl]methyl 2,2-dimethylpropanoate (1 g, 3.26 mmol, prepared according to *J. Med. Chem.* (2006) **49**

6465), triphenylphosphine (1.370 g, 5.22 mmol), and ethanol (0.210 mL, 3.59 mmol) in DCM (10 mL) at 0 °C was treated dropwise over 2 minutes with a 40% solution of DEAD in toluene (2.379 mL, 5.22 mmol). After addition, the mixture was warmed to room temperature and stirred for 3 hours before being concentrated. The residue was dissolved in 7 M ammonia in methanol (10 mL, 70.0 mmol) and the mixture was stirred for 3 days. The solid was collected by filtration and washed with MeOH and CH₂Cl₂ to give 5-(ethyloxy)-7-(methyloxy)-4(1*H*)-quinazolinone (286 mg, 40%) as a white solid. MS (m/z) 221.1 (M+H⁺).

10 Step 2. 4-chloro-5-(ethyloxy)-7-(methyloxy)quinazoline

A mixture of 5-(ethyloxy)-7-(methyloxy)-4(1*H*)-quinazolinone (285 mg, 1.29 mmol) in POCl₃ (10 mL, 107 mmol) was treated with 1 drop of DMF and stirred at 100 °C for 1 hour before being concentrated. The residue was treated with saturated aqueous NaHCO₃ and extracted with CH₂Cl₂. The organic extracts were dried (sodium sulfate), concentrated, and subjected to flash chromatography (1-3% MeOH-CH₂Cl₂) to give 4-chloro-5-(ethyloxy)-7-(methyloxy)quinazoline (260 mg, 84%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.84 (s, 1 H), 7.03 (d, *J*=2.51 Hz, 1 H), 6.84 (d, *J*=2.26 Hz, 1 H), 4.22 (q, *J*=6.86 Hz, 2 H), 3.96 (s, 3 H), 1.47 (t, *J*=6.90 Hz, 3 H). MS (m/z) 239.0 (M+H⁺).

20 The following quinazolines were prepared using procedures analogous to those described in Preparation 14 by substituting the indicated alcohol in the place of ethanol in Step 1:


Quinazoline Product	Alcohol in Step 1	MS (m/z)	¹ H NMR
4-chloro-5-[(1-methylethyl)oxy]-7-(methyloxy)quinazoline	2-propanol	253.0 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 8.82 (s, 1 H), 7.02 (d, <i>J</i> =2.26 Hz, 1 H), 6.88 (d, <i>J</i> =2.26 Hz, 1 H), 4.85 - 4.96 (m, 1 H), 3.96 (s, 3 H), 1.40 (d, <i>J</i> =6.02 Hz, 6 H)
4-chloro-7-(methyloxy)-5-(tetrahydro-2 <i>H</i> -pyran-4-yloxy)quinazoline	tetrahydro-2 <i>H</i> -pyran-4-ol	295.0 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 8.69 (br. s., 1 H), 6.78 (s, 2 H), 4.81 - 4.91 (m, 1 H), 3.86 - 3.93 (m, 5 H), 3.49 - 3.56 (m, 2 H), 1.89 - 2.01 (m, 2 H), 1.68 (m, <i>J</i> =13.49, 6.93, 6.93, 3.51 Hz, 2 H)

4-chloro-7-(methoxy)-5- {[2-(methoxy)ethyl]oxy}quinazoline	2-(methoxy)ethanol	269.0 (M+H ⁺)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 8.84 (s, 1 H), 7.04 (d, <i>J</i> =2.26 Hz, 1 H), 6.88 (d, <i>J</i> =2.26 Hz, 1 H), 4.26 - 4.33 (m, 2 H), 3.96 (s, 3 H), 3.77 - 3.83 (m, 2 H), 3.36 (s, 3 H)
---	--------------------	------------------------------	---

PREPARATION 15

4-chloro-5,6,7-tris(methoxy)quinazoline

5

Step 1. 4,5,6-tris(methoxy)-1*H*-indole-2,3-dione

10 A solution of 3,4,5-trimethoxyaniline (10 g, 54.6 mmol) in Et₂O (100 mL) was treated with a solution of 2M HCl in Et₂O (40.9 mL, 82 mmol) and the resulting light gray solid was collected and washed with ether to give the HCl salt of the aniline (11.5 g), which was treated with oxalyl chloride (16.72 mL, 191 mmol) and the mixture was heated at 170 °C for 1.5 hours before being cooled to room temperature and treated with MeOH. The resulting mixture was heated to boiling and filtered while hot to give a red solid, which
15 was washed with MeOH and dried in air to give 4,5,6-tris(methoxy)-1*H*-indole-2,3-dione (10.5 g, 81%) as a red-orange solid. MS (*m/z*) 238.0 (M+H⁺).

Step 2. 6-amino-2,3,4-tris(methoxy)benzoic acid

20 A solution of 4,5,6-tris(methoxy)-1*H*-indole-2,3-dione (10 g, 42.2 mmol) in 33% aqueous NaOH (70.5 mL, 582 mmol) at 100 °C was treated with 30% aqueous hydrogen peroxide (12.9 mL, 126 mmol) dropwise over 30 min, at a rate to prevent the foaming mixture from overflowing the flask. After addition was complete, the mixture was stirred for 10 minutes before being cooled and diluted by the addition of ice. The mixture was adjusted to pH 8 by addition of concentrated HCl (70 mL) and then acidified to pH 3 with

the addition of glacial acetic acid. The solid that formed was collected by filtration, washed with water, and dried under vacuum to give 6-amino-2,3,4-tris(methyloxy)benzoic acid (2.7 g, 28%) as a tan solid. MS (m/z) 228.0 (M+H⁺).

5 Step 3. methyl 6-amino-2,3,4-tris(methyloxy)benzoate

A solution of 6-amino-2,3,4-tris(methyloxy)benzoic acid (2.7 g, 11.88 mmol) in THF (20 mL) and MeOH (20 mL) at 25 °C was treated with a solution of 2M TMS-diazomethane in Et₂O (14.85 mL, 29.7 mmol) and stirred for 10 minutes before being concentrated to give methyl 6-amino-2,3,4-tris(methyloxy)benzoate (2.87 g, 99%) as a
10 brown solid. MS (m/z) 242.1 (M+H⁺).

Step 4. 5,6,7-tris(methyloxy)-4(1*H*)-quinazolinone


A solution of methyl 6-amino-2,3,4-tris(methyloxy)benzoate (2.87 g, 11.90 mmol) and formamidine acetate (3.72 g, 35.7 mmol) in 2-methoxyethanol (25 mL) was stirred at
15 125 °C for 2 hrs before being concentrated. The residue was suspended in water and the solid was collected by filtration, washed with water, and dried under vacuum (45 °C) to give 5,6,7-tris(methyloxy)-4(1*H*)-quinazolinone (1 g, 34%) as a light brown solid. MS (m/z) 237.1 (M+H⁺).

20 Step 5. 4-chloro-5,6,7-tris(methyloxy)quinazoline

5,6,7-tris(methyloxy)-4(1*H*)-quinazolinone (0.8 g, 3.39 mmol) was treated with POCl₃ (10 mL, 107 mmol) and one drop of DMF, and stirred at 100 °C for 1 hour before being concentrated. The residue was treated with saturated aqueous NaHCO₃ and extracted with CH₂Cl₂. The organic extracts were dried (sodium sulfate) and concentrated
25 to give 4-chloro-5,6,7-tris(methyloxy)quinazoline (770 mg, 89%) as a tan solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.88 (s, 1 H), 7.37 (s, 1 H), 4.04 (s, 3 H), 3.94 (s, 3 H), 3.91 (s, 3 H). MS (m/z) 255.0 (M+H⁺).

PREPARATION 16

N-methyl-3-({6-(methyloxy)-7-[(phenylmethyl)oxy]-4-quinazolinyl}amino)-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide



5

A solution of 3-amino-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide (3.0 g, 10.7 mmol) in *i*-PrOH (100 mL) was treated with 4-chloro-6-(methyloxy)-7-[(phenylmethyl)oxy] quinazoline (3.2 g, 10.7 mmol, prepared according to *J. Med. Chem.* (1999) **42** 5369) and stirred at 80 °C for 2 hours before being concentrated. The residue was washed with petroleum ether and dried to give *N*-methyl-3-({6-(methyloxy)-7-[(phenylmethyl)oxy]-4-quinazolinyl}amino)-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide (4.2 g, 72%) as a brown solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 11.42 (br. s., 1 H), 8.75 (s, 1 H), 8.20 (s, 1 H), 7.80 - 7.84 (m, 2 H), 7.49 - 7.57 (m, 4 H), 7.35 - 7.49 (m, 4 H), 5.34 (s, 2 H), 4.91 (q, *J*=8.82 Hz, 2 H), 3.99 (s, 3 H), 2.45 (d, *J*=5.07 Hz, 3 H). MS (*m/z*) 549.1 (*M*+*H*⁺).


15

The following compound was prepared with procedures analogous to that described in Preparation 16 using the specified quinazoline and aniline starting materials:

20

Name	Structure	Quinazoline	Aniline
4-(dimethylamino)- <i>N</i> -methyl-3-[(6-nitro-4-quinazolinyl)amino]benzenesulfonamide		4-chloro-6-nitroquinazoline	3-amino-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide

EXAMPLE 1

3-[[6,7-bis(methoxy)-4-quinazoliny]amino]-4-(dimethylamino)-*N*-methylbenzenesulfonamide hydrochloride

To a mixture of 3-amino-4-(dimethylamino)-*N*-methylbenzenesulfonamide (0.100g, 0.434mmol) and 4-chloro-6,7-bis(methoxy)quinazoline (0.075 g, 0.334 mmol) in isopropanol (2 mL) was added HCl (0.334 mL, 0.334 mmol). The resulting mixture was then subjected to microwave irradiation (150 °C, 6 bar) for 15 minutes. The mixture was then concentrated and purified *via* column chromatography (5-10% MeOH/CH₂Cl₂) to give an orange solid that was triturated with EtOAc and dried to afford 3-[[6,7-bis(methoxy)-4-quinazoliny]amino]-4-(dimethylamino)-*N*-methylbenzenesulfonamide (54 mg, 35% yield) as orange crystals.


The following compounds were prepared with procedures analogous to that described in Example 1 using 4-chloro-6,7-bis(methoxy)quinazoline and the specified aniline as starting materials:

Ex.	Name	Structure	Aniline
2	3-[[6,7-bis(methoxy)-4-quinazoliny]amino]- <i>N</i> -methylbenzenesulfonamide hydrochloride		3-amino- <i>N</i> -methylbenzenesulfonamide
3	3-[[6,7-bis(methoxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-(4-morpholinyl)benzenesulfonamide trifluoroacetate		3-amino- <i>N</i> -methyl-4-(4-morpholinyl)benzenesulfonamide

4	3-[[6,7-bis(methoxy)-4-quinazoliny]amino]-4-[ethyl(methyl)amino]- <i>N</i> -methylbenzenesulfonamide		3-amino-4-[ethyl(methyl)amino]- <i>N</i> -methylbenzene sulfonamide
5	3-[[6,7-bis(methoxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(1-methylethyl)oxy]benzenesulfonamide		3-amino- <i>N</i> -methyl-4-[(1-methylethyl)oxy]benzenesulfonamide
6	3-[[6,7-bis(methoxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-(methoxy)benzenesulfonamide		3-amino- <i>N</i> -methyl-4-(methoxy)benzene sulfonamide
7	3-[[6,7-bis(methoxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-(methylthio)benzenesulfonamide trifluoroacetate		3-amino- <i>N</i> -methyl-4-(methylthio)benzene sulfonamide

EXAMPLE 8

4-(dimethylamino)-*N*-methyl-3-(4-quinazolinylamino)benzenesulfonamide trifluoroacetate




A mixture of 3-amino-4-(dimethylamino)-*N*-methylbenzenesulfonamide (136 mg, 0.59 mmol) and 4-chloroquinazoline (75 mg, 0.46 mmol) in *i*-PrOH (2 mL) was subjected to microwave irradiation (150 °C, 6 bar) for 30 minutes before being filtered and subjected to reverse phase HPLC (10-50% MeCN/water/0.1%TFA; Sunfire C18 OBD 5 μM , 30x75 mm) to give an oil, which was dissolved in water and lyophilized to give 4-(dimethylamino)-*N*-methyl-3-(4-quinazolinylamino)benzenesulfonamide trifluoroacetate (30 mg, 13%) as an orange solid.

The following compounds were prepared with procedures analogous to that described in Example 8 using the specified quinazoline and aniline starting materials:

Ex.	Name	Structure	Quinazoline	Aniline
9	3-[(6-chloro-4-quinazolinyl)amino]-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide		4,6-dichloroquinazoline	3-amino-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide
10	4-(dimethylamino)- <i>N</i> -methyl-3-[[6-(methoxy)-4-quinazolinyl]amino]benzene sulfonamide trifluoroacetate		4-chloro-6-(methoxy)quinazoline	3-amino-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide

EXAMPLE 11







- 5 3-[[6,7-bis(methoxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide hydrochloride










- 10 A mixture of 4-chloro-6,7-bis(methoxy)quinazoline (100 mg, 0.445 mmol) and 3-amino-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide (139 mg, 0.490 mmol) in *i*-PrOH (2 mL) was subjected to microwave irradiation (100 °C, 2 bar) for 30 minutes before being cooled to room temperature. The solid that formed was collected by filtration and washed with isopropanol to give 3-[[6,7-bis(methoxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide hydrochloride (160 mg, 70%) as a pale yellow solid.
- 15

The following compounds were prepared with procedures analogous to that described in Example 11 using the specified quinazoline and aniline starting materials:


Ex.	Name	Structure	Quinazoline	Aniline
12	3-[(6-chloro-4-quinazoliny)amino]- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide		4,6-dichloroquinazoline	3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide
13	3-(7,8-dihydro[1,4]dioxino[2,3-g]quinazolin-4-ylamino)-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide bis(trifluoroacetate)		4-chloro-7,8-dihydro[1,4]dioxino[2,3-g]quinazoline	3-amino-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide
14	3-[[6,7-bis(methoxy)-4-quinazoliny]amino]-4-(4,4-difluoro-1-piperidiny)- <i>N</i> -methylbenzenesulfonamide trifluoroacetate		4-chloro-6,7-bis(methoxy)quinazoline	3-amino-4-(4,4-difluoro-1-piperidiny)- <i>N</i> -methylbenzenesulfonamide
15	3-[(6-chloro-4-quinazoliny)amino]-4-(4,4-difluoro-1-piperidiny)- <i>N</i> -methylbenzenesulfonamide trifluoroacetate		4,6-dichloroquinazoline	3-amino-4-(4,4-difluoro-1-piperidiny)- <i>N</i> -methylbenzenesulfonamide
16	3-[(7-[(3-chloropropyl)oxy]-4-quinazoliny)amino]-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide		4-chloro-7-[(3-chloropropyl)oxy]quinazoline	3-amino-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide
17	3-[[6,7-bis{2-(methoxy)ethyl}oxy]-4-quinazoliny]amino]-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide trifluoroacetate		4-chloro-6,7-bis{2-(methoxy)ethyl}oxy}quinazoline	3-amino-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide

18	3-[(6,7-bis{[2-(methyloxy)ethyl]oxy}-4-quinazoliny]amino]- <i>N</i> -methyl-4-(methyloxy)benzenesulfonamide hydrochloride		4-chloro-6,7-bis{[2-(methyloxy)ethyl]oxy}quinazoline	3-amino- <i>N</i> -methyl-4-(methyloxy)benzenesulfonamide
19	3-[(6,7-bis{[2-(methyloxy)ethyl]oxy}-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide		4-chloro-6,7-bis{[2-(methyloxy)ethyl]oxy}quinazoline	3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide
20	3-[(6,7-bis{[2-(methyloxy)ethyl]oxy}-4-quinazoliny]amino]- <i>N</i> -methyl-4-(methylthio)benzenesulfonamide		4-chloro-6,7-bis{[2-(methyloxy)ethyl]oxy}quinazoline	3-amino- <i>N</i> -methyl-4-(methylthio)benzenesulfonamide
21	3-[(6,7-bis(methyloxy)-4-quinazoliny]amino)- <i>N</i> -methyl-4-[(2 <i>R</i>)-2-(trifluoromethyl)-1-pyrrolidinyl]benzenesulfonamide trifluoroacetate		4-chloro-6,7-bis(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-[(2 <i>R</i>)-2-(trifluoromethyl)-1-pyrrolidinyl]benzenesulfonamide
22	3-[(6-chloro-4-quinazoliny]amino)- <i>N</i> -methyl-4-[(2 <i>R</i>)-2-(trifluoromethyl)-1-pyrrolidinyl]benzenesulfonamide trifluoroacetate		4,6-dichloroquinazoline	3-amino- <i>N</i> -methyl-4-[(2 <i>R</i>)-2-(trifluoromethyl)-1-pyrrolidinyl]benzenesulfonamide
23	5-[(6,7-bis(methyloxy)-4-quinazoliny]amino)-2-fluoro- <i>N</i> -methylbenzenesulfonamide hydrochloride		4-chloro-6,7-bis(methoxy)quinazoline	5-amino-2-fluoro- <i>N</i> -methylbenzenesulfonamide

24	3-{[6,7-bis(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide trifluoroacetate		4-chloro-6,7-bis(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide
25	3-{[6,7-bis(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-(1-piperidiny)benzenesulfonamide trifluoroacetate		4-chloro-6,7-bis(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-(1-piperidiny)benzenesulfonamide
26	3-[(6-chloro-4-quinazoliny]amino)- <i>N</i> -methyl-4-(1-piperidiny)benzenesulfonamide trifluoroacetate		4,6-dichloroquinazoline	3-amino- <i>N</i> -methyl-4-(1-piperidiny)benzenesulfonamide
27	3-{[6,7-bis(methyloxy)-4-quinazoliny]amino}-4-(2,5-dimethyl-1-pyrrolidiny)- <i>N</i> -methylbenzenesulfonamide trifluoroacetate		4-chloro-6,7-bis(methoxy)quinazoline	3-amino-4-(2,5-dimethyl-1-pyrrolidiny)- <i>N</i> -methylbenzenesulfonamide
28	3-[(6-chloro-4-quinazoliny]amino)-4-(2,5-dimethyl-1-pyrrolidiny)- <i>N</i> -methylbenzenesulfonamide trifluoroacetate		4,6-dichloroquinazoline	3-amino-4-(2,5-dimethyl-1-pyrrolidiny)- <i>N</i> -methylbenzenesulfonamide
29	3-{[6,7-bis(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)thio]benzenesulfonamide hydrochloride		4-chloro-6,7-bis(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)thio]benzenesulfonamide
30	3-{[5,7-bis(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methylbenzenesulfonamide hydrochloride		4-chloro-5,7-bis(methoxy)quinazoline	3-amino- <i>N</i> -methylbenzenesulfonamide

EXAMPLE 31

3-[[5,7-bis(methoxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide hydrochloride









5







A mixture of 4-chloro-5,7-bis(methoxy)quinazoline (75 mg, 0.334 mmol, prepared according to *J. Med. Chem.* (2006) **49** 6465) and 3-amino-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide (95 mg, 0.334 mmol) in *i*-PrOH (2 mL) was subjected to microwave irradiation (100 °C, 1 bar) for 30 minutes before being cooled to room temperature. The solid was collected by filtration, washed with *i*-PrOH, and dried under vacuum to give 3-[[5,7-bis(methoxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide hydrochloride (130 mg, 76%) as a yellow solid.







15








The following compounds were prepared with procedures analogous to that described in Example 31 using the specified quinazoline and aniline starting materials:







Ex.	Name	Structure	Quinazoline	Aniline
32	3-[[5,7-bis(methoxy)-4-quinazolinyl]amino]-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide		4-chloro-5,7-bis(methoxy)quinazoline	3-amino-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide
33	3-(5-chloro-6,7-dimethoxyquinazolin-4-ylamino)- <i>N</i> -methylbenzenesulfonamide trifluoroacetate		4,5-dichloro-6,7-bis(methoxy)quinazoline	3-amino- <i>N</i> -methylbenzenesulfonamide

34	4-methoxy-3-(7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-ylamino)- <i>N</i> -methylbenzenesulfonamide bis(trifluoroacetate)		4-chloro-7-(methoxy)-6-{{3-(4-morpholinyl)propyl}oxy}quinazoline (<i>J. Med. Chem.</i> (2004) 47 871)	3-amino- <i>N</i> -methyl-4-(methoxy)benzenesulfonamide
35	3-{{6,7-bis(methoxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-[methyl(2,2,2-trifluoroethyl)amino]benzenesulfonamide trifluoroacetate		4-chloro-6,7-bis(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-[methyl(2,2,2-trifluoroethyl)amino]benzenesulfonamide
36	3-{{5,7-bis(methoxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-[methyl(2,2,2-trifluoroethyl)amino]benzenesulfonamide hydrochloride		4-chloro-5,7-bis(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-[methyl(2,2,2-trifluoroethyl)amino]benzenesulfonamide
37	3-{{5,7-bis(methoxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-(methoxy)benzenesulfonamide hydrochloride		4-chloro-5,7-bis(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-(methoxy)benzenesulfonamide
38	3-{{5,7-bis(methoxy)-4-quinazoliny]amino}-4-fluoro- <i>N</i> -methylbenzenesulfonamide hydrochloride		4-chloro-5,7-bis(methoxy)quinazoline	3-amino-4-fluoro- <i>N</i> -methylbenzenesulfonamide
39	3-{{5,7-bis(methoxy)-4-quinazoliny]amino}-4-chloro- <i>N</i> -methylbenzenesulfonamide hydrochloride		4-chloro-5,7-bis(methoxy)quinazoline	3-amino-4-chloro- <i>N</i> -methylbenzenesulfonamide

40	3-{{[5,7-bis(methoxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-(4-morpholinyl)benzenesulfonamide hydrochloride		4-chloro-5,7-bis(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-(4-morpholinyl)benzenesulfonamide
41	5-{{[6,7-bis(methoxy)-4-quinazoliny]amino}-2-fluoro- <i>N</i> -methyl-4-(methoxy)benzenesulfonamide hydrochloride		4-chloro-6,7-bis(methoxy)quinazoline	5-amino-2-fluoro- <i>N</i> -methyl-4-(methoxy)benzenesulfonamide
42	3-{{[6-(ethoxy)-7-(methoxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide hydrochloride		4-chloro-6-(ethoxy)-7-(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide
43	4-(dimethylamino)-3-(7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-ylamino)- <i>N</i> -methylbenzenesulfonamide bis(formate)		4-chloro-7-(methoxy)-6-{{[3-(4-morpholinyl)propyl]oxy}quinazoline	3-amino-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide
44	<i>N</i> -methyl-3-{{[6-[(1-methylethyl)oxy]-7-(methoxy)-4-quinazoliny]amino}-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide hydrochloride		4-chloro-6-[(1-methylethyl)oxy]-7-(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide
45	3-{{[7-[(3-chloropropyl)oxy]-4-quinazoliny]amino}- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide		4-chloro-7-[(3-chloropropyl)oxy]quinazoline	3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide

46	3-{{[5-(ethyloxy)-7-(methyloxy)-4-quinazoliny]amino}-N-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide		4-chloro-5-(ethyloxy)-7-(methyloxy)quinazoline	3-amino-N-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide
47	N-methyl-3-{{[5-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazoliny]amino}-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide		4-chloro-5-[(1-methylethyl)oxy]-7-(methyloxy)quinazoline	3-amino-N-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide
48	N-methyl-4-[(2,2,2-trifluoroethyl)oxy]-3-{{[5,6,7-tris(methyloxy)-4-quinazoliny]amino}benzenesulfonamide hydrochloride		4-chloro-5,6,7-tris(methyloxy)quinazoline	3-amino-N-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide
49	3-{{[5,7-bis(methyloxy)-4-quinazoliny]amino}-N-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide trifluoroacetate		4-chloro-5,7-bis(methyloxy)quinazoline	3-amino-N-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide
50	5-{{[5,7-bis(methyloxy)-4-quinazoliny]amino}-2-fluoro-N-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide trifluoroacetate		4-chloro-5,7-bis(methyloxy)quinazoline	5-amino-2-fluoro-N-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide
51	5-{{[5,7-bis(methyloxy)-4-quinazoliny]amino}-2-fluoro-N-methyl-4-(methyloxy)benzenesulfonamide trifluoroacetate		4-chloro-5,7-bis(methyloxy)quinazoline	5-amino-2-fluoro-N-methyl-4-(methyloxy)benzenesulfonamide

52	5-[[6,7-bis(methoxy)-4-quinazoliny]amino]-2-fluoro- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide hydrochloride		4-chloro-6,7-bis(methoxy)quinazoline	5-amino-2-fluoro- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide
53	3-[[5,7-bis(methoxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-(methylthio)benzenesulfonamide hydrochloride		4-chloro-5,7-bis(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-(methylthio)benzenesulfonamide
54	3-[[5,7-bis(methoxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide hydrochloride		4-chloro-5,7-bis(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide
55	<i>N</i> -methyl-4-[(trifluoromethyl)oxy]-3-[[5,6,7-tris(methoxy)-4-quinazoliny]amino]benzenesulfonamide trifluoroacetate		4-chloro-5,6,7-tris(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide
56	5-[[5,7-bis(methoxy)-4-quinazoliny]amino]-4-(dimethylamino)-2-fluoro- <i>N</i> -methylbenzenesulfonamide hydrochloride		4-chloro-5,7-bis(methoxy)quinazoline	5-amino-4-(dimethylamino)-2-fluoro- <i>N</i> -methylbenzenesulfonamide
57	5-[[6,7-bis(methoxy)-4-quinazoliny]amino]-4-(dimethylamino)-2-fluoro- <i>N</i> -methylbenzenesulfonamide hydrochloride		4-chloro-6,7-bis(methoxy)quinazoline	5-amino-4-(dimethylamino)-2-fluoro- <i>N</i> -methylbenzenesulfonamide
58	2-fluoro- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]-5-[[5,6,7-tris(methoxy)-4-quinazoliny]amino]benzenesulfonamide trifluoroacetate		4-chloro-5,6,7-tris(methoxy)quinazoline	5-amino-2-fluoro- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide


59	4-chloro- <i>N</i> -methyl-3-{{5,6,7-tris(methoxy)-4-quinazoliny]amino}benzene sulfonamide trifluoroacetate		4-chloro-5,6,7-tris(methoxy)quinazoline	3-amino-4-chloro- <i>N</i> -methylbenzene sulfonamide
60	<i>N</i> -methyl-3-[[7-(methoxy)-5-[[2-(methoxy)ethyl]oxy]-4-quinazoliny]amino]-4-[[2,2,2-trifluoroethyl]oxy]benzenesulfonamide hydrochloride		4-chloro-7-(methoxy)-5-[[2-(methoxy)ethyl]oxy]quinazoline	3-amino- <i>N</i> -methyl-4-[[2,2,2-trifluoroethyl]oxy]benzenesulfonamide
61	<i>N</i> -methyl-3-[[7-(methoxy)-5-(tetrahydro-2 <i>H</i> -pyran-4-yloxy)-4-quinazoliny]amino]-4-[[2,2,2-trifluoroethyl]oxy]benzenesulfonamide		4-chloro-7-(methoxy)-5-(tetrahydro-2 <i>H</i> -pyran-4-yloxy)quinazoline	3-amino- <i>N</i> -methyl-4-[[2,2,2-trifluoroethyl]oxy]benzenesulfonamide
62	<i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]-3-{{5,6,7-tris(methoxy)-4-quinazoliny]amino}benzene sulfonamide trifluoroacetate		4-chloro-5,6,7-tris(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide
63	<i>N</i> -methyl-4-(methylsulfonyl)-3-{{5,6,7-tris(methoxy)-4-quinazoliny]amino}benzene sulfonamide trifluoroacetate		4-chloro-5,6,7-tris(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-(methylsulfonyl)benzenesulfonamide
64	3-[[5-(ethoxy)-7-(methoxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-[[trifluoromethyl]oxy]benzenesulfonamide		4-chloro-5-(ethoxy)-7-(methoxy)quinazoline	3-amino- <i>N</i> -methyl-4-[[trifluoromethyl]oxy]benzenesulfonamide

65	<i>N</i> -methyl-3-{{5-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazoliny]amino}-4-[(trifluoromethyl)oxy]benzenesulfonamide		4-chloro-5-[(1-methylethyl)oxy]-7-(methyloxy)quinazoline	3-amino- <i>N</i> -methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide
66	3-{{5-(ethyloxy)-7-(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide		4-chloro-5-(ethyloxy)-7-(methyloxy)quinazoline	3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide
67	<i>N</i> -methyl-3-{{5-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazoliny]amino}-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide		4-chloro-5-[(1-methylethyl)oxy]-7-(methyloxy)quinazoline	3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide
68	3-{{6,7-bis(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-(methylsulfonyl)benzenesulfonamide trifluoroacetate		4-chloro-6,7-bis(methyloxy)quinazoline	3-amino- <i>N</i> -methyl-4-(methylsulfonyl)benzenesulfonamide
69	3-{{6,7-bis(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide		4-chloro-6,7-bis(methyloxy)quinazoline	3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide
70	3-{{6-iodo-7-(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide		4-chloro-6-iodo-7-(methyloxy)quinazoline	3-amino- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide

71	3-{{[6,7-bis(ethyloxy)-4-quinazoliny]amino}-N-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide		4-chloro-6,7-bis(ethyloxy)quinazoline (<i>J. Med. Chem.</i> (2005) 48 7445)	3-amino-N-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide
72	4-(dimethylamino)-N-methyl-3-((7-nitroquinazolin-4-yl)amino)benzenesulfonamide hydrochloride		4-chloro-7-nitroquinazoline	3-amino-4-(dimethylamino)-N-methylbenzenesulfonamide


EXAMPLE 73

4-(dimethylamino)-N-methyl-3-[(7-{[3-(4-morpholinyl)propyl]oxy}-4-quinazoliny]amino]benzenesulfonamide hydrochloride



A mixture of 3-[(7-{[3-(4-chloropropyl)oxy]-4-quinazoliny]amino}-N-methyl-4-(dimethylamino)phenyl]benzenesulfonamide (50 mg, 0.111 mmol) and morpholine (0.048 mL, 0.556 mmol) in MeOH (1 mL) was stirred at 90 °C for 3 days before being cooled and concentrated. The residue was triturated with Et₂O-EtOAc and the resulting solid was collected by filtration, washed with Et₂O, and dried to give 4-(dimethylamino)-N-methyl-3-[(7-{[3-(4-morpholinyl)propyl]oxy}-4-quinazoliny]amino]benzenesulfonamide hydrochloride (33 mg, 48%) as an off-white solid.


The following compound was prepared with procedures analogous to that described in Example 73 using the specified quinazoline starting material:

Ex.	Name	Structure	Quinazoline SM
74	<i>N</i> -methyl-3-[(7-{[3-(4-morpholinyl)propyl]oxy}-4-quinazolinyl)amino]-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide trifluoroacetate		3-({7-[(3-chloropropyl)oxy]-4-quinazolinyl}amino)- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide

5

EXAMPLE 75

N-methyl-3-[(7-{[3-(methyloxy)propyl]oxy}-4-quinazolinyl)amino]-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide




10

A mixture of 3-({7-[(3-chloropropyl)oxy]-4-quinazolinyl}amino)-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide (150 mg, 0.297 mmol) in MeOH (3 mL) was treated with a solution of 25% sodium methoxide in methanol (3 g, 13.88 mmol) and subjected to microwave irradiation (100 °C) for 10 minutes before being cooled and neutralized with the addition of concentrated HCl. The resulting solution was diluted with saturated aqueous NaHCO₃ and extracted with CH₂Cl₂. The organic extracts were dried (sodium sulfate), concentrated, and subjected to reverse phase HPLC (30-70% MeCN/aq. NH₄OH pH 10, XBridge C18 OBD 30x150 mm, 5 μm) to give *N*-methyl-3-[(7-{[3-(methyloxy)propyl]oxy}-4-quinazolinyl)amino]-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide (37 mg, 25%) as a white solid.

20


The following compound was prepared with procedures analogous to that described in Example 75 using the specified quinazoline starting material:

Ex.	Name	Structure	Quinazoline SM
76	4-(dimethylamino)- <i>N</i> -methyl-3-[(7-[(3-(methoxy)propyl)oxy]-4-quinazolinyl)amino]benzene sulfonamide		3-({7-[(3-chloropropyl)oxy]-4-quinazolinyl}amino)-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide

5

EXAMPLE 77

3-[[5,7-bis(methoxy)-4-quinazolinyl]amino]-*N*-methyl-4-(methylsulfonyl)benzenesulfonamide trifluoroacetate




10

A mixture of 3-[[5,7-bis(methoxy)-4-quinazolinyl]amino]-*N*-methyl-4-(methylthio)benzenesulfonamide (200 mg, 0.476 mmol) and sodium perborate tetrahydrate (220 mg, 1.427 mmol) in AcOH (1.902 ml) was heated at 50 °C for 2 hours before being filtered and extracted with EtOAc. The organic extract was washed with water, saturated aqueous NaHCO₃, and brine, then dried (sodium sulfate), concentrated, and subjected to reverse phase HPLC (Sunfire prep C18 OBD, 30x 150 mm, 10-50 % Water/CH₃CN+0.1%TFA) to give 3-[[5,7-bis(methoxy)-4-quinazolinyl]amino]-*N*-methyl-4-(methylsulfonyl)benzenesulfonamide trifluoroacetate (28 mg, 10%) as a white solid.

20

EXAMPLE 78

3-(7-hydroxy-6-methoxyquinazolin-4-ylamino)-*N*-methyl-4-(2,2,2-trifluoroethoxy)benzenesulfonamide




5

A mixture of *N*-methyl-3-({6-(methyloxy)-7-[(phenylmethyl)oxy]-4-quinazolinyl}amino)-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide (4.2 g, 7.66 mmol) in EtOAc (120 mL), MeOH (40 mL), and DMF (40 mL) was treated with 60% AcOH (0.4 mL) and Pd/C (400 mg) was stirred under H₂ (balloon) for 2 hours before being filtered and concentrated to give 3-(7-hydroxy-6-methoxyquinazolin-4-ylamino)-*N*-methyl-4-(2,2,2-trifluoroethoxy) benzenesulfonamide (3.2 g, 91%) as a yellow solid.

15

EXAMPLE 79

3-(7-isopropoxy-6-methoxy-quinazolin-4-ylamino)-*N*-methyl-4-(2,2,2-trifluoro-ethoxy)-benzenesulfonamide





20

A solution of 3-[[7-hydroxy-6-(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide (170 mg, 0.37 mmol) in DMF (5 mL) was treated with 2-bromopropane (181 mg, 1.48 mmol) and K₂CO₃ (154 mg, 1.11 mmol) and stirred at 60 °C for 16 hours before being poured into water. The precipitate that formed was collected by filtration, washed with MeOH, and dried to give 3-(7-isopropoxy-6-methoxy-quinazolin-4-ylamino)-*N*-methyl-4-(2,2,2-trifluoro-ethoxy)-benzenesulfonamide (80 mg, 43%) as an off-white solid.

25

The following compounds were prepared with procedures analogous to that described in Example 79 using the indicated alkyl halide:


Ex.	Name	Structure	Alkyl Halide
80	3-(7-ethoxy-6-methoxy-quinazolin-4-ylamino)- <i>N</i> -methyl-4-(2,2,2-trifluoroethoxy)-benzenesulfonamide trifluoroacetate		bromoethane
81	3-[6-methoxy-7-(2-methoxyethoxy)-quinazolin-4-ylamino]- <i>N</i> -methyl-4-(2,2,2-trifluoroethoxy)-benzenesulfonamide trifluoroacetate		1-bromo-2-(methyloxy)ethane

5

EXAMPLES 82 & 83

3-[[5,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide hydrochloride (enantiomer 1) & 3-[[5,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide hydrochloride (enantiomer 2)


10



A racemic mixture of 3-[[5,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide hydrochloride (528 mg) was subjected to chiral SFC (Chiralpak OJ-H, 15% MeOH cosolvent) to provide 3-[[5,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide hydrochloride (unassigned enantiomer 1, 33mg) & 3-[[5,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide hydrochloride (unassigned enantiomer 2, 49 mg).

20

EXAMPLE 84

3-((6-aminoquinazolin-4-yl)amino)-4-(dimethylamino)-*N*-methylbenzenesulfonamide

5

A mixture of 4-(dimethylamino)-*N*-methyl-3-((6-nitroquinazolin-4-yl)amino)benzenesulfonamide (715 mg, 1.78 mmol) in MeOH (50 mL) was treated with 10% Pd/C (189 mg, 0.178 mmol) and stirred under H₂ (balloon) at 25 °C for 2.5 hours before being filtered through celite and concentrated. The residue was dissolved in CH₂Cl₂/MeOH, washed with saturated aqueous NaHCO₃, dried (Na₂SO₄), and concentrated to give 3-((6-aminoquinazolin-4-yl)amino)-4-(dimethylamino)-*N*-methylbenzenesulfonamide (433 mg, 59%) as a yellow solid.

10

The following compound was prepared with procedures analogous to that described in Example 84:

15


Ex.	Name	Structure	Quinazoline SM
85	3-((7-aminoquinazolin-4-yl)amino)-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide		4-(dimethylamino)- <i>N</i> -methyl-3-[(7-amino-4-quinazolinyl)amino]benzenesulfonamide

EXAMPLE 86

4-(dimethylamino)-3-((6-(dimethylamino)quinazolin-4-yl)amino)-*N*-methylbenzenesulfonamide bis(trifluoroacetate) &

EXAMPLE 87

5 4-(dimethylamino)-*N*-methyl-3-((6-(methylamino)quinazolin-4-yl)amino)benzenesulfonamide bis(trifluoroacetate)



A mixture of 3-((6-aminoquinazolin-4-yl)amino)-4-(dimethylamino)-*N*-methylbenzenesulfonamide (129 mg, 0.346 mmol) in CH_2Cl_2 (3 mL) was treated with AcOH (0.022 mL, 0.381 mmol) and 37% aqueous formaldehyde (0.028 mL, 0.381 mmol) and stirred at 25 °C. After 5 minutes, sodium triacetoxyborohydride (110 mg, 0.520 mmol) was added and the mixture was stirred for 2 hours before being quenched with the addition of saturated aqueous NaHCO_3 and extracted with a $\text{CH}_2\text{Cl}_2/\text{MeOH}$ mixture. The organic extract was dried (Na_2SO_4), concentrated, and subjected to reverse phase HPLC (10-40% MeCN/ H_2O /0.1% TFA on a Sunfire TM Prep C18 OBD TM 5 μm 30x150 mm column) to give 4-(dimethylamino)-*N*-methyl-3-((6-(methylamino)quinazolin-4-yl)amino)benzenesulfonamide (3 mg, 1.4%) as a yellow solid and 4-(dimethylamino)-3-((6-(dimethylamino)quinazolin-4-yl)amino)-*N*-methylbenzenesulfonamide (3.6 mg, 1.6%) as a yellow solid.

The following compound was prepared with procedures analogous to that described in Examples 86 & 87:

Ex.	Name	Structure	Quinazoline SM
88	4-(dimethylamino)- <i>N</i> -methyl-3-((7-(methylamino)quinazolin-4-yl)amino)benzenesulfonamide		3-((7-aminoquinazolin-4-yl)amino)-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide

Spectroscopic data for Examples 1-88:

Ex.	Name	t _R (min)	MS (m/z)	¹ H NMR
1	3-{[6,7-bis(methyloxy)-4-quinazolinyl]amino}-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide hydrochloride	2.06 ^a	418	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 11.38 (br. s., 1H), 8.81 (s, 1H), 8.29 (s, 1H), 7.59 - 7.71 (m, 2H), 7.31 - 7.39 (m, 2H), 7.23 (d, <i>J</i> = 9.54 Hz, 1H), 4.00 (s, 3H), 4.01 (s, 3H), 2.82 (s, 6H), 2.43 (d, <i>J</i> = 4.27 Hz, 3H)
2	3-{[6,7-bis(methyloxy)-4-quinazolinyl]amino}- <i>N</i> -methylbenzenesulfonamide hydrochloride	1.81 ^a	375.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 10.98 (br. s., 1H), 8.85 (s, 1H), 8.05 - 8.18 (m, 3H), 7.73 (t, <i>J</i> = 7.91 Hz, 1H), 7.67 (d, <i>J</i> = 7.78 Hz, 1H), 7.59 (q, <i>J</i> = 4.43 Hz, 1H), 7.29 (s, 1H), 4.02 (d, <i>J</i> = 3.76 Hz, 6H), 2.48 (s, 3H)
3	3-{[6,7-bis(methyloxy)-4-quinazolinyl]amino}- <i>N</i> -methyl-4-(4-morpholinyl)benzene sulfonamide trifluoroacetate	1.78 ^a	459.9	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 10.66 (br. s., 1H), 8.76 (s, 1H), 8.05 (s, 1H), 7.92 - 7.98 (m, 1H), 7.71 (dd, <i>J</i> = 2.01, 8.53 Hz, 1H), 7.44 (q, <i>J</i> = 5.02 Hz, 1H), 7.37 (d, <i>J</i> = 8.53 Hz, 1H), 7.28 (s, 1H), 3.99 (s, 3H), 4.01 (s, 3H), 3.50 - 3.60 (m, 4H), 2.96 - 3.03 (m, 4H), 2.45 (d, <i>J</i> = 4.77 Hz, 3H)
4	3-{[6,7-bis(methyloxy)-4-quinazolinyl]amino}-4-[ethyl(methyl)amino]- <i>N</i> -methylbenzenesulfonamide	1.91 ^a	432.2	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 11.29 (br. s., 1H), 8.80 (s, 1H), 8.29 (s, 1H), 7.62 - 7.71 (m, 2H), 7.32 - 7.39 (m, 2H), 7.26 (d, <i>J</i> = 8.53 Hz, 1H), 3.99 (s, 3H), 4.01 (s, 3H), 3.07 (q, <i>J</i> = 6.94 Hz, 2H), 2.79 (s, 3H), 2.43 (d, <i>J</i> = 4.77 Hz, 3H), 0.96 (t, <i>J</i> = 7.03 Hz, 3H)
5	3-{[6,7-bis(methyloxy)-4-quinazolinyl]amino}- <i>N</i> -methyl-4-(methyloxy)benzenesulfonamide	1.93 ^a	433.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 11.09 (br. s., 1H), 8.76 (s, 1H), 8.12 (s, 1H), 7.78 - 7.85 (m, 2H), 7.40 - 7.49 (m, 2H), 7.30 (s, 1H), 4.00 (d, <i>J</i> = 7.55 Hz, 6H), 3.89 (s, 3H), 2.45 (d, <i>J</i> = 4.78 Hz, 3H)
6	3-{[6,7-bis(methyloxy)-4-quinazolinyl]amino}- <i>N</i> -methyl-4-[(1-methylethyl)oxy]benzene sulfonamide	1.75 ^a	405	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 10.93 (br. s., 1H), 8.77 (s, 1H), 8.05 (s, 1H), 7.80 - 7.85 (m, 1H), 7.76 (dd, <i>J</i> = 2.27, 8.81 Hz, 1H), 7.39 - 7.49 (m, 2H), 7.29 (s, 1H), 4.78 (quin, <i>J</i> = 5.98 Hz, 1H), 4.01 (s, 3H), 3.99 (s, 3H), 2.45 (d, <i>J</i> = 5.04 Hz, 3H), 1.21 (s, 3H), 1.19 (s, 3H)

7	3-[[6,7-bis(methyloxy)-4-quinazolinyl]amino]- <i>N</i> -methyl-4-(methylthio)benzenesulfonamide trifluoroacetate	1.80 ^a	421.1	¹ H NMR (400 MHz, methanol-d ₄) δ 7.08 (s, 1H), 6.41 (s, 1H), 6.32 - 6.37 (m, 1H), 6.30 (d, <i>J</i> = 1.76 Hz, 1H), 6.10 (d, <i>J</i> = 8.53 Hz, 1H), 5.73 (s, 1H), 2.54 (s, 3H), 2.56 (s, 3H), 1.05 (s, 3H), 1.02 (s, 3H)
8	4-(dimethylamino)- <i>N</i> -methyl-3-(4-quinazolinylamino)benzenesulfonamide trifluoroacetate	1.67 ^a	358.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.77 (br. s., 1 H), 8.78 - 8.87 (m, 1 H), 8.75 (s, 1 H), 8.06 (dd, <i>J</i> =8.91, 1.88 Hz, 1 H), 7.87 (d, <i>J</i> =9.04 Hz, 1 H), 7.74 (d, <i>J</i> =2.01 Hz, 1 H), 7.63 (dd, <i>J</i> =8.78, 2.26 Hz, 1 H), 7.32 (q, <i>J</i> =5.02 Hz, 1 H), 7.23 (d, <i>J</i> =8.78 Hz, 1 H), 2.80 (s, 6 H), 2.43 (d, <i>J</i> =5.02 Hz, 3 H)
9	3-[(6-chloro-4-quinazolinyl)amino]-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide	1.89 ^a	392.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.32 (br. s., 1 H), 8.85 (s, 1 H), 8.19 (d, <i>J</i> =2.51 Hz, 1 H), 7.87 (d, <i>J</i> =9.29 Hz, 1 H), 7.77 (dd, <i>J</i> =9.29, 2.51 Hz, 1 H), 7.72 (d, <i>J</i> =2.26 Hz, 1 H), 7.67 (dd, <i>J</i> =8.53, 2.26 Hz, 1 H), 7.33 (q, <i>J</i> =4.94 Hz, 1 H), 7.25 (d, <i>J</i> =8.78 Hz, 1 H), 3.97 (s, 3 H), 2.82 (s, 6 H), 2.39 - 2.47 (m, 3 H)
10	4-(dimethylamino)- <i>N</i> -methyl-3-[[6-(methyloxy)-4-quinazolinyl]amino]benzenesulfonamide trifluoroacetate	1.89 ^a	388.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.41 (s, 1 H), 8.36 (s, 1 H), 7.97 (d, <i>J</i> =2.26 Hz, 1 H), 7.73 (s, 1 H), 7.68 (dd, <i>J</i> =8.78, 2.26 Hz, 1 H), 7.40 - 7.50 (m, 2 H), 7.20 (s, 1 H), 4.88 (q, <i>J</i> =8.78 Hz, 2 H), 3.95 (s, 3 H), 3.94 (s, 3 H), 2.46 (d, <i>J</i> =4.77 Hz, 3 H)
11	3-[[6,7-bis(methyloxy)-4-quinazolinyl]amino]- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide hydrochloride	2.10 ^a	473.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.91 (br. s., 1 H), 8.57 (br. s., 1 H), 8.49 (s, 1 H), 7.77 - 7.95 (m, 3 H), 7.67 - 7.77 (m, 1 H), 7.47 (d, <i>J</i> =8.28 Hz, 2 H), 4.88 (q, <i>J</i> =8.70 Hz, 2 H), 2.46 (d, <i>J</i> =4.77 Hz, 3 H)
12	3-[(6-chloro-4-quinazolinyl)amino]- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide	2.14 ^a	447.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.05 (br. s., 1 H), 8.78 (s, 1 H), 8.24 (s, 1 H), 7.69 (d, <i>J</i> =2.26 Hz, 1 H), 7.63 - 7.68 (m, 1 H), 7.29 - 7.35 (m, 2 H), 7.23 (d, <i>J</i> =8.78 Hz, 1 H), 4.41 - 4.59 (m, 4 H), 2.80 (s, 6 H), 2.43 (d, <i>J</i> =5.02 Hz, 3 H)

13	3-(7,8-dihydro[1,4]dioxino[2,3-g]quinazolin-4-ylamino)-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide bis(trifluoroacetate)	1.87 ^a	416.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.63 (br. s., 1 H), 8.77 (s, 1 H), 8.07 (s, 1 H), 7.97 (d, J=2.01 Hz, 1 H), 7.69 (dd, J=8.53, 2.01 Hz, 1 H), 7.38 - 7.51 (m, 2 H), 7.30 (s, 1 H), 4.02 (s, 3 H), 3.98 (s, 3 H), 3.12 (t, J=5.14 Hz, 4 H), 2.45 (d, J=5.02 Hz, 3 H), 1.87 - 2.05 (m, 4 H)
14	3-{{6,7-bis(methoxy)-4-quinazoliny]amino}-4-(4,4-difluoro-1-piperidiny)- <i>N</i> -methylbenzenesulfonamide trifluoroacetate	2.14 ^a	494.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.57 (br. s., 1 H), 8.67 - 8.86 (m, 2 H), 8.06 (dd, J=8.91, 2.13 Hz, 1 H), 7.93 - 8.01 (m, 1 H), 7.89 (d, J=9.04 Hz, 1 H), 7.67 (dd, J=8.53, 2.26 Hz, 1 H), 7.36 - 7.49 (m, 2 H), 3.12 (t, J=5.27 Hz, 4 H), 2.46 (d, J=4.77 Hz, 3 H), 1.89 - 2.03 (m, 4 H)
15	3-[(6-chloro-4-quinazoliny]amino)-4-(4,4-difluoro-1-piperidiny)- <i>N</i> -methylbenzenesulfonamide trifluoroacetate	2.24 ^a	468.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.45 (br. s., 1 H), 8.44 (s, 1 H), 8.38 (d, J=9.03 Hz, 1 H), 7.88 - 7.95 (m, 1 H), 7.54 (dd, J=8.53, 2.01 Hz, 1 H), 7.13 - 7.36 (m, 4 H), 4.28 (t, J=6.02 Hz, 2 H), 3.85 (t, J=6.53 Hz, 2 H), 2.77 (s, 6 H), 2.43 (s, 3 H), 2.26 (quin, J=6.21 Hz, 2 H)
16	3-({7-[(3-chloropropyl)oxy]-4-quinazoliny]amino)-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide	2.15 ^a	450.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.95 (br. s., 1 H), 8.76 (br. s., 1 H), 8.60 (d, J=9.29 Hz, 1 H), 7.76 (d, J=2.01 Hz, 1 H), 7.64 (dd, J=8.66, 2.13 Hz, 1 H), 7.44 (d, J=8.78 Hz, 1 H), 7.33 (q, J=5.19 Hz, 1 H), 7.20 - 7.29 (m, 2 H), 4.30 (t, J=5.77 Hz, 2 H), 4.01 (br. s., 2 H), 3.68 (t, J=12.55 Hz, 2 H), 3.57 (s, 3 H), 3.52 (br. s., 2 H), 3.34 (br. s., 2 H), 3.14 (br. s., 2 H), 2.80 (s, 6 H), 2.40 - 2.46 (m, 3 H), 2.23 (d, J=9.03 Hz, 2 H)
17	3-[(6,7-bis[{2-(methoxy)ethyl}oxy]-4-quinazoliny]amino)-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide trifluoroacetate	1.96 ^a	506.2	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.12 (br. s., 1 H), 8.75 (s, 1 H), 8.19 (s, 1 H), 7.78 - 7.84 (m, 2 H), 7.41 - 7.49 (m, 2 H), 7.36 (s, 1 H), 4.35 (q, J=4.85 Hz, 4 H), 3.89 (s, 3 H), 3.76 - 3.83 (m, 4 H), 3.37 (s, 3 H), 3.37 (s, 3 H), 2.46 (d, J=5.02 Hz, 3 H)

18	3-[(6,7-bis{[2-(methyloxy)ethyl]oxy}-4-quinazoliny]amino]- <i>N</i> -methyl-4-(methyloxy)benzenesulfonamide hydrochloride	1.90 ^a	493.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.39 (br. s., 1 H), 8.35 (s, 1 H), 7.96 (d, J=2.01 Hz, 1 H), 7.77 (s, 1 H), 7.68 (dd, J=8.53, 2.26 Hz, 1 H), 7.39 - 7.51 (m, 2 H), 7.23 (s, 1 H), 4.88 (q, J=8.70 Hz, 2 H), 4.22 - 4.32 (m, 4 H), 3.70 - 3.82 (m, 4 H), 3.37 (s, 3 H), 3.36 (s, 3 H), 2.46 (s, 3 H)
19	3-[(6,7-bis{[2-(methyloxy)ethyl]oxy}-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide	2.08 ^a	561.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.58 (br. s., 1 H), 8.29 (s, 1 H), 7.85 (s, 1 H), 7.62 - 7.74 (m, 2 H), 7.56 (d, J=8.28 Hz, 1 H), 7.42 - 7.51 (m, 1 H), 7.22 (s, 1 H), 4.22 - 4.33 (m, 4 H), 3.71 - 3.81 (m, 4 H), 3.38 (s, 3 H), 3.36 (s, 3 H), 2.46 (s, 6 H)
20	3-[(6,7-bis{[2-(methyloxy)ethyl]oxy}-4-quinazoliny]amino]- <i>N</i> -methyl-4-(methylthio)benzenesulfonamide	1.97 ^a	509.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.06 (br. s., 1 H), 8.74 (s, 1 H), 8.06 (s, 1 H), 7.69 (d, J=2.01 Hz, 1 H), 7.64 (dd, J=8.78, 2.26 Hz, 1 H), 7.54 (d, J=9.03 Hz, 1 H), 7.36 (q, J=4.94 Hz, 1 H), 7.31 (s, 1 H), 4.91 - 5.04 (m, 1 H), 4.01 (s, 3 H), 3.98 (s, 3 H), 3.32 - 3.43 (m, 1 H), 3.15 - 3.24 (m, 1 H), 2.44 (d, J=5.02 Hz, 3 H), 2.10 - 2.23 (m, 1 H), 1.70 - 1.94 (m, 3 H)
21	3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(2 <i>R</i>)-2-(trifluoromethyl)-1-pyrrolidiny]benzenesulfonamide trifluoroacetate	2.14 ^a	512.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.85 (br. s., 1 H), 8.73 (br. s., 2 H), 8.01 - 8.10 (m, 1 H), 7.87 (d, J=9.03 Hz, 1 H), 7.71 (br. s., 1 H), 7.60 (dd, J=8.78, 2.01 Hz, 1 H), 7.52 (d, J=8.78 Hz, 1 H), 7.33 (q, J=4.94 Hz, 1 H), 4.96 (br. s., 1 H), 3.36 - 3.47 (m, 1 H), 3.14 - 3.25 (m, 1 H), 2.44 (d, J=5.02 Hz, 3 H), 2.09 - 2.23 (m, 1 H), 1.68 - 1.95 (m, 3 H)
22	3-[(6-chloro-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(2 <i>R</i>)-2-(trifluoromethyl)-1-pyrrolidiny]benzenesulfonamide trifluoroacetate	2.25 ^a	486.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.53 (br. s., 1 H), 8.89 (s, 1 H), 8.34 (s, 1 H), 8.06 - 8.21 (m, 2 H), 7.89 (q, J=4.77 Hz, 1 H), 7.61 (t, J=9.29 Hz, 1 H), 7.37 (s, 1 H), 4.04 (s, 3 H), 4.01 (s, 3 H), 2.57 (d, J=4.77 Hz, 3 H)

23	5-{{[6,7-bis(methyloxy)-4-quinazoliny]amino}-2-fluoro- <i>N</i> -methylbenzenesulfonamide hydrochloride	1.85 ^a	393.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.86 (br. s., 1 H), 8.72 (s, 1 H), 8.05 (d, J=2.01 Hz, 1 H), 7.98 (s, 1 H), 7.89 (dd, J=8.66, 2.13 Hz, 1 H), 7.79 - 7.86 (m, 1 H), 7.73 (q, J=5.10 Hz, 1 H), 7.29 (s, 1 H), 4.00 - 4.04 (m, 3 H), 3.97 - 4.00 (m, 3 H), 2.5 (d, 3H, obscured by DMSO)
24	3-{{[6,7-bis(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide trifluoroacetate	2.05 ^a	459.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.72 (br. s., 1 H), 8.79 (s, 1 H), 8.09 (s, 1 H), 7.92 (d, J=1.76 Hz, 1 H), 7.69 (dd, J=8.53, 2.26 Hz, 1 H), 7.41 (q, J=5.19 Hz, 1 H), 7.34 (d, J=8.78 Hz, 1 H), 7.29 (s, 1 H), 4.02 (s, 3 H), 3.98 (s, 3 H), 2.95 (br. s., 4 H), 2.45 (d, J=5.02 Hz, 3 H), 1.46 (br. s., 6 H)
25	3-{{[6,7-bis(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-(1-piperidiny)benzenesulfonamide trifluoroacetate	2.16 ^a	458.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.68 (br. s., 1 H), 8.78 (s, 2 H), 8.07 (dd, J=8.91, 2.13 Hz, 1 H), 7.95 (d, J=2.01 Hz, 1 H), 7.89 (d, J=9.03 Hz, 1 H), 7.67 (dd, J=8.41, 2.13 Hz, 1 H), 7.40 (q, J=4.94 Hz, 1 H), 7.33 (d, J=8.53 Hz, 1 H), 2.95 (br. s., 4 H), 2.45 (d, J=5.02 Hz, 3 H), 1.47 (br. s., 6 H)
26	3-[(6-chloro-4-quinazoliny)amino]- <i>N</i> -methyl-4-(1-piperidiny)benzenesulfonamide trifluoroacetate	2.30 ^a	432.1	¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ ppm 10.97 (br. s., 1 H), 8.83 (s, 1 H), 7.97 (br. s., 1 H), 7.72 (br. s., 1 H), 7.59 (dd, J=9.03, 2.20 Hz, 1 H), 7.28 (s, 1 H), 7.18 - 7.26 (m, 1 H), 7.15 (d, J=9.03 Hz, 1 H), 4.01 (s, 3 H), 3.96 (s, 3 H), 3.77 - 3.84 (m, 2 H), 2.40 (d, J=4.88 Hz, 3 H), 1.88 - 2.01 (m, 2 H), 1.56 - 1.67 (m, 2 H), 1.12 (d, J=6.10 Hz, 6 H)
27	3-{{[6,7-bis(methyloxy)-4-quinazoliny]amino}-4-(2,5-dimethyl-1-pyrrolidiny)- <i>N</i> -methylbenzenesulfonamide trifluoroacetate	2.21 ^a	472.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.96 (br. s., 1 H), 8.83 (s, 1 H), 8.64 (s, 1 H), 8.09 (dd, J=8.91, 2.13 Hz, 1 H), 7.89 (d, J=9.03 Hz, 1 H), 7.80 (d, J=2.01 Hz, 1 H), 7.58 (dd, J=8.91, 2.38 Hz, 1 H), 7.24 (d, J=4.77 Hz, 1 H), 7.18 (d, J=9.03 Hz, 1 H), 3.82 (br. s., 2 H), 2.42 (d, J=4.77 Hz, 3 H), 1.89 - 2.03 (m, 2 H), 1.57 - 1.69 (m, 2 H), 1.12 (d, 6 H)

28	3-[(6-chloro-4-quinazoliny)amino]-4-(2,5-dimethyl-1-pyrrolidiny)- <i>N</i> -methylbenzenesulfonamide trifluoroacetate	2.38 ^a	446.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.83 (br. s., 1 H), 8.61 (br. s., 1 H), 8.03 (s, 1 H), 7.96 (d, J=8.03 Hz, 1 H), 7.72 - 7.80 (m, 2 H), 7.61 (q, J=4.94 Hz, 1 H), 7.29 (s, 1 H), 4.14 (q, J=10.21 Hz, 2 H), 3.99 (s, 3 H), 3.99 (s, 3 H), 2.47 (d, J=5.02 Hz, 3 H)
29	3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)thio]benzenesulfonamide hydrochloride	2.08 ^a	489.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.97 (s, 1 H), 8.48 (s, 1 H), 8.39 (t, J=1.76 Hz, 1 H), 7.94 - 8.00 (m, 1 H), 7.55 - 7.65 (m, 1 H), 7.45 - 7.55 (m, 2 H), 6.83 (d, J=2.26 Hz, 1 H), 6.75 (d, J=2.26 Hz, 1 H), 4.10 (s, 3 H), 3.92 (s, 3 H), 2.44 - 2.49 (m, 3 H)
30	3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]- <i>N</i> -methylbenzenesulfonamide hydrochloride	1.86 ^a	375.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.79 (s, 1 H), 8.83 (s, 1 H), 8.60 (br. s., 1 H), 7.73 (dd, J=8.66, 2.13 Hz, 1 H), 7.48 - 7.60 (m, 2 H), 6.98 (d, J=2.26 Hz, 1 H), 6.92 (d, J=2.26 Hz, 1 H), 5.08 (q, J=8.78 Hz, 2 H), 4.14 (s, 3 H), 3.99 (s, 3 H), 2.47 (d, J=5.02 Hz, 3 H)
31	3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide hydrochloride	2.08 ^a	473	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.92 (s, 1 H), 9.30 (d, J=2.01 Hz, 1 H), 8.57 (s, 1 H), 7.42 - 7.49 (m, 2 H), 7.40 (q, J=5.02 Hz, 1 H), 6.85 (d, J=2.26 Hz, 1 H), 6.79 (d, J=2.26 Hz, 1 H), 4.08 - 4.13 (m, 3 H), 3.93 (s, 3 H), 2.77 (s, 6 H), 2.46 (d, J=5.02 Hz, 3 H)
32	3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide	2.01 ^a	418.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.87 (br. s., 1 H), 8.57 (s, 1 H), 8.17 (s, 1 H), 7.92 (d, J=8.82 Hz, 1 H), 7.57 - 7.64 (m, 1 H), 7.48 - 7.56 (m, 2 H), 7.34 (s, 1 H), 4.01 (s, 3 H), 3.85 (s, 3 H), 2.45 (d, J=4.85 Hz, 3 H)
33	3-(5-chloro-6,7-dimethoxyquinazolin-4-ylamino)- <i>N</i> -methylbenzenesulfonamide trifluoroacetate	0.97 ^b	409.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.87 (br. s., 1 H), 10.24 (br. s., 1 H), 8.70 (s, 1 H), 8.10 (br. s., 1 H), 7.73 - 7.84 (m, 2 H), 7.35 - 7.48 (m, 2 H), 7.32 (d, J=2.21 Hz, 1 H), 4.25 (t, J=5.51 Hz, 2 H), 3.93 - 4.04 (m, 5 H), 3.85 (s, 3 H), 3.65 - 3.74 (m, 2 H), 3.48 (d, J=6.62 Hz, 2 H), 3.30 (br. s., 2 H), 3.11 (br. s., 2 H), 2.42 (d, J=4.85 Hz, 3 H), 2.23 - 2.34 (m, 2 H)

34	4-methoxy-3-(7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-ylamino)- <i>N</i> -methylbenzenesulfonamide bis(trifluoroacetate)	0.86 ^b	518.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.93 (br. s., 1 H), 8.78 (s, 1 H), 8.05 (s, 1 H), 7.78 (d, J=2.01 Hz, 1 H), 7.72 (dd, J=8.66, 2.13 Hz, 1 H), 7.51 (d, J=8.78 Hz, 1 H), 7.44 (q, J=5.02 Hz, 1 H), 7.28 (s, 1 H), 4.02 (s, 3 H), 3.98 (s, 3 H), 3.88 - 3.97 (m, 2 H), 2.96 (s, 3 H), 2.44 (d, J=5.02 Hz, 3 H)
35	3-{{6,7-bis(methyloxy)-4-quinazolinyl}amino}- <i>N</i> -methyl-4-[methyl(2,2,2-trifluoroethyl)amino]benzene sulfonamide trifluoroacetate	1.94 ^a	486.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.02 - 11.20 (m, 1 H), 8.85 (s, 1 H), 8.79 (br. s., 1 H), 7.61 - 7.77 (m, 2 H), 7.52 (q, J=5.02 Hz, 1 H), 6.98 (d, J=1.76 Hz, 1 H), 6.88 (d, J=2.26 Hz, 1 H), 4.16 (s, 3 H), 4.07 (q, J=9.54 Hz, 2 H), 3.99 (s, 3 H), 2.92 (s, 3 H), 2.48 (d, J=5.02 Hz, 3 H)
36	3-{{5,7-bis(methyloxy)-4-quinazolinyl}amino}- <i>N</i> -methyl-4-[methyl(2,2,2-trifluoroethyl)amino]benzene sulfonamide hydrochloride	2.05 ^a	486.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.09 (s, 1 H), 8.88 (s, 1 H), 8.78 (br. s., 1 H), 7.69 (dd, J=8.66, 2.13 Hz, 1 H), 7.47 (q, J=4.94 Hz, 1 H), 7.41 (d, J=8.78 Hz, 1 H), 6.98 (d, J=1.76 Hz, 1 H), 6.91 (d, J=2.01 Hz, 1 H), 4.18 (s, 3 H), 4.04 (s, 3 H), 3.99 (s, 3 H), 2.45 (d, J=5.02 Hz, 3 H)
37	3-{{5,7-bis(methyloxy)-4-quinazolinyl}amino}- <i>N</i> -methyl-4-(methyloxy)benzenesulfonamide hydrochloride	1.86 ^a	405.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.73 (s, 1 H), 8.78 (s, 1 H), 8.21 (br. s., 1 H), 7.78 - 7.87 (m, 1 H), 7.62 - 7.73 (m, 2 H), 6.96 (d, J=2.26 Hz, 1 H), 6.90 (d, J=2.26 Hz, 1 H), 4.13 (s, 3 H), 3.99 (s, 3 H), 2.49 (d, J=5.02 Hz, 3 H)
38	3-{{5,7-bis(methyloxy)-4-quinazolinyl}amino}-4-fluoro- <i>N</i> -methylbenzenesulfonamide hydrochloride	1.84 ^a	393.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.92 (s, 1 H), 8.79 (s, 1 H), 8.41 (br. s., 1 H), 7.92 (d, J=8.28 Hz, 1 H), 7.70 - 7.77 (m, 2 H), 6.97 (d, J=2.26 Hz, 1 H), 6.92 (d, J=2.26 Hz, 1 H), 4.16 (s, 3 H), 3.99 (s, 3 H), 2.49 (s, 3 H)
39	3-{{5,7-bis(methyloxy)-4-quinazolinyl}amino}-4-chloro- <i>N</i> -methylbenzenesulfonamide hydrochloride	1.92 ^a	409.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.81 (br. s., 1 H), 8.85 (s, 1 H), 8.52 (s, 1 H), 7.67 (dd, J=8.41, 2.13 Hz, 1 H), 7.44 - 7.53 (m, 2 H), 7.00 (d, J=2.01 Hz, 1 H), 6.92 (d, J=2.01 Hz, 1 H), 4.23 (s, 3 H), 4.00 (s, 3 H), 3.67 - 3.75 (m, 4 H), 2.95 - 3.02 (m, 4 H), 2.46 (d, J=5.02 Hz, 3 H)

40	3-{{[5,7-bis(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-(4-morpholinyl)benzenesulfonamide hydrochloride	1.84 ^a	460.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.07 (br. s., 1 H), 8.76 (s, 1 H), 8.13 (s, 1 H), 7.78 (d, J=7.78 Hz, 1 H), 7.74 (q, J=4.77 Hz, 1 H), 7.43 (d, J=12.05 Hz, 1 H), 7.31 (s, 1 H), 4.01 (s, 3 H), 3.99 (s, 3 H), 3.90 (s, 3 H), 2.53 (d, J=5.27 Hz, 3 H)
41	5-{{[6,7-bis(methyloxy)-4-quinazoliny]amino}-2-fluoro- <i>N</i> -methyl-4-(methyloxy)benzenesulfonamide hydrochloride	1.72 ^a	423.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.19 (br. s., 1 H), 8.76 (s, 1 H), 8.10 (s, 1 H), 7.79 - 7.91 (m, 2 H), 7.50 - 7.59 (m, 2 H), 7.34 (s, 1 H), 4.93 (q, J=8.78 Hz, 2 H), 4.26 (q, J=6.86 Hz, 2 H), 4.02 (s, 3 H), 2.47 (d, J=5.02 Hz, 3 H), 1.46 (t, J=7.03 Hz, 3 H)
42	3-{{[6-(ethyloxy)-7-(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide hydrochloride	1.97 ^a	487.1	¹ H NMR (400 MHz, METHANOL- <i>d</i> ₄) δ ppm 8.44 (br. s., 2 H), 8.38 (s, 1 H), 8.12 (d, J=2.21 Hz, 1 H), 7.69 (s, 1 H), 7.63 (dd, J=8.38, 2.21 Hz, 1 H), 7.27 (d, J=8.38 Hz, 1 H), 7.20 (s, 1 H), 4.27 (t, J=5.73 Hz, 2 H), 4.01 (s, 3 H), 3.77 (t, J=4.63 Hz, 4 H), 2.79 - 2.85 (m, 8 H), 2.74 (br. s., 4 H), 2.57 (s, 3 H), 2.13 - 2.22 (m, 2 H)
43	4-(dimethylamino)-3-(7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-ylamino)- <i>N</i> -methylbenzenesulfonamide bis(formate)	0.89 ^b	531.2	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.27 (br. s., 1 H), 8.77 (s, 1 H), 8.15 (s, 1 H), 7.88 (d, J=2.26 Hz, 1 H), 7.84 (dd, J=8.66, 2.38 Hz, 1 H), 7.53 - 7.59 (m, 2 H), 7.35 (s, 1 H), 4.84 - 5.00 (m, 3 H), 4.01 (s, 3 H), 2.47 (d, J=5.02 Hz, 3 H), 1.40 (d, J=6.02 Hz, 6 H)
44	<i>N</i> -methyl-3-{{[6-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazoliny]amino}-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide hydrochloride	2.04 ^a	501.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.58 (s, 1 H), 8.19 - 8.38 (m, 2 H), 7.95 (br. s., 1 H), 7.56 (br. s., 1 H), 7.37 (br. s., 2 H), 7.11 (br. s., 2 H), 4.86 (q, J=8.95 Hz, 2 H), 4.26 (t, J=5.77 Hz, 2 H), 3.85 (t, J=6.53 Hz, 2 H), 2.45 (s, 3 H), 2.25 (quin, J=6.27 Hz, 2 H)
45	3-{{[7-[(3-chloropropyl)oxy]-4-quinazoliny]amino}- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide	2.09 ^a	505.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.18 (s, 1 H), 9.14 (d, J=2.01 Hz, 1 H), 8.53 (s, 1 H), 7.48 - 7.56 (m, 2 H), 7.45 (q, J=4.85 Hz, 1 H), 6.86 (d, J=2.26 Hz, 1 H), 6.80 (d, J=2.51 Hz, 1 H), 5.12 (q, J=8.95 Hz, 2 H), 4.44 (q, J=6.94 Hz, 2 H), 3.92 (s, 3 H), 2.47 (d, J=5.02 Hz, 3 H), 1.46 (t, J=6.90 Hz, 3 H)

46	3-[[5-(ethyloxy)-7-(methyloxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide	2.08 ^a	487.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.00 (s, 1 H), 8.91 (s, 1 H), 8.48 (s, 1 H), 7.51 - 7.59 (m, 2 H), 7.44 (q, J=5.02 Hz, 1 H), 6.84 (d, J=2.26 Hz, 1 H), 6.81 (d, J=2.26 Hz, 1 H), 5.10 (q, J=8.87 Hz, 2 H), 5.00 (dt, J=12.17, 5.96 Hz, 1 H), 3.92 (s, 3 H), 2.48 (d, J=5.02 Hz, 3 H), 1.44 (d, J=6.02 Hz, 6 H)
47	<i>N</i> -methyl-3-[[5-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazoliny]amino]-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide	2.13 ^a	501.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.59 (s, 1 H), 8.40 (s, 1 H), 8.30 (d, J=9.03 Hz, 1 H), 7.94 (d, J=2.26 Hz, 1 H), 7.69 (dd, J=8.78, 2.26 Hz, 1 H), 7.45 (d, J=8.78 Hz, 2 H), 7.26 (dd, J=9.16, 2.38 Hz, 1 H), 7.17 (d, J=2.51 Hz, 1 H), 4.88 (q, J=8.78 Hz, 2 H), 4.21 (t, J=6.27 Hz, 2 H), 3.52 (t, J=6.27 Hz, 2 H), 3.28 (s, 3 H), 2.46 (d, J=4.52 Hz, 3 H), 2.03 (quin, J=6.34 Hz, 2 H)
48	<i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]-3-[[5,6,7-tris(methyloxy)-4-quinazoliny]amino]benzenesulfonamide hydrochloride	2.23 ^a	503.1	¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ ppm 10.59 (br. s., 1 H), 8.75 (s, 2 H), 7.63 - 7.70 (m, 1 H), 7.52 - 7.63 (m, 1 H), 7.47 (q, J=4.64 Hz, 1 H), 6.93 (s, 1 H), 6.86 (d, J=1.95 Hz, 1 H), 5.49 - 5.68 (m, 1 H), 4.12 (s, 3 H), 3.97 (s, 3 H), 2.47 (d, J=4.88 Hz, 3 H), 1.52 (d, J=6.35 Hz, 3 H)
49	3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide trifluoroacetate	2.08 ^a	487.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.14 (s, 1 H), 8.76 (s, 1 H), 8.6 (s, 1 H), 8.21 (d, J=8.28 Hz, 1 H), 7.83 - 7.93 (m, 2 H), 6.94 (d, J=2.01 Hz, 1 H), 6.90 (d, J=2.01 Hz, 1 H), 4.11 (s, 3 H), 3.99 (s, 3 H), 3.35 (s, 3 H), 2.54 (d, J=4.77 Hz, 3 H)
50	5-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-2-fluoro- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide trifluoroacetate	2.03 ^a	491.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.81 (br. s., 1 H), 8.77 (s, 2 H), 7.65 - 7.74 (m, 1 H), 7.39 (d, J=11.80 Hz, 1 H), 6.92 (d, J=2.01 Hz, 1 H), 6.83 (d, J=2.26 Hz, 1 H), 4.15 (s, 3 H), 4.04 (s, 3 H), 3.97 (s, 3 H), 2.52 (br. s., 3 H)
51	5-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-2-fluoro- <i>N</i> -methyl-4-(methyloxy)benzenesulfonamide trifluoroacetate	1.89 ^a	423.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.23 (br. s., 1 H), 8.78 (s, 1 H), 8.12 (s, 1 H), 7.81 - 7.88 (m, 2 H), 7.61 (d, J=11.80 Hz, 1 H), 7.33 (s, 1 H), 4.96 (q, J=8.78 Hz, 2 H), 4.02 (s, 3 H), 4.00 (s, 3 H), 2.54 (d, J=4.77 Hz, 3 H)

52	5-{{6,7-bis(methyloxy)-4-quinazoliny]amino}-2-fluoro- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide hydrochloride	1.90 ^a	491.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.80 (s, 1 H), 8.74 (s, 1 H), 7.91 (d, J=1.51 Hz, 1 H), 7.77 (dd, J=8.41, 1.88 Hz, 1 H), 7.57 - 7.66 (m, 2 H), 6.94 - 6.98 (m, 2 H), 4.14 (s, 3 H), 3.99 (s, 3 H), 2.54 (s, 3 H), 2.47 (d, J=5.02 Hz, 3 H)
53	3-{{5,7-bis(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-(methylthio)benzenesulfonamide hydrochloride	1.92 ^a	421.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.84 (s, 1 H), 8.84 (s, 1 H), 8.37 - 8.44 (m, 1 H), 7.86 - 7.91 (m, 1 H), 7.79 - 7.86 (m, 2 H), 6.98 - 7.03 (m, 2 H), 4.14 (s, 3 H), 3.99 (s, 3 H), 2.49 - 2.53 (m, 3 H)
54	3-{{5,7-bis(methyloxy)-4-quinazoliny]amino}- <i>N</i> -methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide hydrochloride	2.06 ^a	459.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.75 (br. s., 1 H), 8.92 (br. s., 1 H), 8.76 (s, 1 H), 7.68 - 7.88 (m, 3 H), 7.17 (s, 1 H), 4.18 (s, 3 H), 4.03 (s, 3 H), 3.91 (s, 3 H), 2.53 (br. s., 3 H)
55	<i>N</i> -methyl-4-[(trifluoromethyl)oxy]-3-{{5,6,7-tris(methyloxy)-4-quinazoliny]amino}benzenesulfonamide trifluoroacetate	2.14 ^a	489.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.95 (s, 1 H), 8.83 (s, 1 H), 8.20 (d, J=7.78 Hz, 1 H), 7.63 (q, J=4.77 Hz, 1 H), 7.17 (d, J=12.55 Hz, 1 H), 6.97 (d, J=2.01 Hz, 1 H), 6.91 (d, J=2.26 Hz, 1 H), 4.14 (s, 3 H), 3.98 (s, 3 H), 2.85 (s, 6 H), 2.50 (br. s., 3 H)
56	5-{{5,7-bis(methyloxy)-4-quinazoliny]amino}-4-(dimethylamino)-2-fluoro- <i>N</i> -methylbenzenesulfonamide hydrochloride	1.93 ^a	436.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.21 (br. s., 1 H), 8.79 (s, 1 H), 8.22 (s, 1 H), 7.50 - 7.63 (m, 2 H), 7.32 (s, 1 H), 7.00 (d, J=13.30 Hz, 1 H), 4.01 (s, 3 H), 3.99 (s, 3 H), 2.85 (s, 6 H), 2.49 (d, J=4.77 Hz, 3 H)
57	5-{{6,7-bis(methyloxy)-4-quinazoliny]amino}-4-(dimethylamino)-2-fluoro- <i>N</i> -methylbenzenesulfonamide hydrochloride	1.78 ^a	436.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.69 (s, 1 H), 8.97 (br. s., 1 H), 8.75 (s, 1 H), 7.77 (q, J=4.77 Hz, 1 H), 7.58 (d, J=11.54 Hz, 1 H), 7.13 (s, 1 H), 5.12 (q, J=8.70 Hz, 2 H), 4.13 (s, 3 H), 4.02 (s, 3 H), 3.90 (s, 3 H), 2.54 (d, J=4.77 Hz, 3 H)
58	2-fluoro- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]-5-{{5,6,7-tris(methyloxy)-4-quinazoliny]amino}benzenesulfonamide trifluoroacetate	2.09 ^a	521.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.76 (s, 1 H), 9.03 (br. s., 1 H), 8.69 (s, 1 H), 7.87 (d, J=8.28 Hz, 1 H), 7.64 - 7.72 (m, 1 H), 7.60 (dd, J=8.28, 1.76 Hz, 1 H), 7.15 (s, 1 H), 4.21 (s, 3 H), 4.02 (s, 3 H), 3.90 (s, 3 H), 2.50 (br. s., 3 H)

59	4-chloro- <i>N</i> -methyl-3-[[5,6,7-tris(methyloxy)-4-quinazoliny]amino]benzene sulfonamide trifluoroacetate	2.01 ^a	439.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.34 (br. s., 1 H), 8.32 (s, 1 H), 7.97 (d, J=1.32 Hz, 1 H), 7.68 (s, 1 H), 7.64 (dd, J=8.49, 1.87 Hz, 1 H), 7.43 (d, J=8.60 Hz, 2 H), 7.16 (s, 1 H), 4.74 - 4.94 (m, 4 H), 3.90 (s, 3 H), 2.44 (s, 3 H), 1.34 (d, J=5.95 Hz, 6 H)
60	<i>N</i> -methyl-3-[[7-(methyloxy)-5-[[2-(methyloxy)ethyl]oxy]-4-quinazoliny]amino]-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide hydrochloride	2.04 ^a	517.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.03 (br. s., 1 H), 10.26 (br. s., 1 H), 8.47 (s, 1 H), 7.88 (s, 2 H), 7.71 (dd, J=8.71, 2.32 Hz, 1 H), 7.38 - 7.54 (m, 2 H), 7.16 (s, 1 H), 4.87 (q, J=8.75 Hz, 2 H), 3.93 (s, 3 H), 2.42 (d, J=4.85 Hz, 3 H)
61	<i>N</i> -methyl-3-[[7-(methyloxy)-5-(tetrahydro-2 <i>H</i> -pyran-4-yloxy)-4-quinazoliny]amino]-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide	2.03 ^a	543.1	¹ H NMR (400 MHz, METHANOL- <i>d</i> ₄) δ ppm 9.02 (d, J=2.26 Hz, 1 H), 8.69 (s, 1 H), 7.66 (dd, J=8.78, 2.26 Hz, 1 H), 7.40 (d, J=9.03 Hz, 1 H), 7.00 (s, 1 H), 5.32 (dt, J=12.74, 6.31 Hz, 1 H), 4.18 (s, 3 H), 4.00 (s, 3 H), 3.89 (s, 3 H), 2.49 (s, 3 H), 1.53 (d, J=6.27 Hz, 3 H)
62	<i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]-3-[[5,6,7-tris(methyloxy)-4-quinazoliny]amino]benzene sulfonamide trifluoroacetate	2.14 ^a	517.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.93 (s, 1 H), 8.61 - 8.73 (m, 2 H), 8.19 (d, J=8.28 Hz, 1 H), 7.75 - 7.91 (m, 2 H), 7.16 (s, 1 H), 4.19 (s, 3 H), 4.03 (s, 3 H), 3.89 (s, 3 H), 3.36 (s, 3 H), 2.55 (d, J=5.02 Hz, 3 H)
63	<i>N</i> -methyl-4-(methylsulfonyl)-3-[[5,6,7-tris(methyloxy)-4-quinazoliny]amino]benzene sulfonamide trifluoroacetate	1.94 ^a	483.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.83 (br. s., 1 H), 8.71 (br. s., 1 H), 8.48 (s, 1 H), 7.85 - 7.93 (m, 1 H), 7.80 (dd, J=8.66, 1.88 Hz, 1 H), 7.48 - 7.60 (m, 2 H), 7.24 - 7.31 (m, 1 H), 4.92 (q, J=8.70 Hz, 2 H), 4.30 (t, J=5.77 Hz, 2 H), 4.02 (br. s., 2 H), 3.68 (br. s., 2 H), 3.51 (br. s., 2 H), 3.34 (br. s., 2 H), 3.15 (br. s., 2 H), 2.47 (d, J=5.02 Hz, 3 H), 2.23 (d, J=9.03 Hz, 2 H)
64	3-[[5-(ethyloxy)-7-(methyloxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide	2.15 ^a	473.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.16 (s, 1 H), 9.04 (d, J=2.26 Hz, 1 H), 8.53 (s, 1 H), 7.76 (dd, J=8.53, 1.51 Hz, 1 H), 7.68 (br. s., 1 H), 7.63 (dd, J=8.66, 2.38 Hz, 1 H), 6.88 (d, J=2.26 Hz, 1 H), 6.85 (d, J=2.26 Hz, 1 H), 5.05 (dt, J=12.05, 6.02 Hz, 1 H), 3.93 (s, 3 H), 2.52 (br. s., 3 H), 1.46 (d, J=6.27 Hz, 6 H)

65	<i>N</i> -methyl-3-{[5-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazoliny]amino}-4-[(trifluoromethyl)oxy]benzenesulfonamide	2.21 ^a	487.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.05 (s, 1 H), 9.01 (d, J=2.01 Hz, 1 H), 8.51 (s, 1 H), 7.49 - 7.60 (m, 2 H), 7.44 (q, J=5.02 Hz, 1 H), 6.86 (d, J=2.26 Hz, 1 H), 6.80 (d, J=2.26 Hz, 1 H), 5.53 (dt, J=12.99, 6.43 Hz, 1 H), 4.44 (q, J=6.94 Hz, 2 H), 3.92 (s, 3 H), 2.48 (d, J=5.02 Hz, 3 H), 1.54 (d, J=6.27 Hz, 3 H), 1.46 (t, J=7.03 Hz, 3 H)
66	3-[[5-(ethyloxy)-7-(methyloxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide	2.14 ^a	501.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.84 (s, 1 H), 8.73 (d, J=1.76 Hz, 1 H), 8.45 (s, 1 H), 7.50 - 7.60 (m, 2 H), 7.40 - 7.49 (m, 1 H), 6.84 (d, J=2.26 Hz, 1 H), 6.80 (d, J=2.26 Hz, 1 H), 5.41 - 5.55 (m, 1 H), 4.92 - 5.02 (m, 1 H), 3.93 (s, 3 H), 2.49 (d, J=5.02 Hz, 3 H), 1.52 (d, J=6.27 Hz, 3 H), 1.44 (dd, J=6.02, 2.26 Hz, 6 H)
67	<i>N</i> -methyl-3-[[5-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazoliny]amino]-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide	2.20 ^a	515.1	¹ H NMR (400 MHz, METHANOL- <i>d</i> ₄) δ ppm 8.81 (s, 1 H), 8.72 (s, 1 H), 8.35 (d, J=8.28 Hz, 1 H), 8.02 (dd, J=8.28, 1.76 Hz, 1 H), 7.78 (s, 1 H), 7.30 (s, 1 H), 4.12 (s, 3 H), 4.10 (s, 3 H), 3.24 (s, 3 H), 2.67 (s, 3 H)
68	3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-(methylsulfonyl)benzenesulfonamide trifluoroacetate	1.78 ^a	453.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.62 (br. s., 1 H), 8.43 (s, 1 H), 8.00 (d, J=2.01 Hz, 1 H), 7.78 (s, 1 H), 7.69 (dd, J=8.78, 2.26 Hz, 1 H), 7.53 (d, J=8.78 Hz, 1 H), 7.42 - 7.50 (m, 1 H), 7.22 (s, 1 H), 5.39 (dt, J=12.80, 6.40 Hz, 1 H), 3.96 (s, 3 H), 3.95 (s, 3 H), 2.47 (d, J=5.02 Hz, 3 H), 1.36 (d, J=6.27 Hz, 3 H)
69	3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide	1.96 ^a	487.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.45 (br. s., 1 H), 8.44 (s, 1 H), 8.37 (d, J=9.03 Hz, 1 H), 7.90 (d, J=1.76 Hz, 1 H), 7.54 (dd, J=8.53, 2.26 Hz, 1 H), 7.15 - 7.37 (m, 4 H), 4.20 (t, J=6.27 Hz, 2 H), 3.52 (t, J=6.27 Hz, 2 H), 3.29 (s, 3 H), 2.77 (s, 6 H), 2.43 (s, 3 H), 2.03 (quin, J=6.27 Hz, 2 H)

70	3-[[6-iodo-7-(methyloxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide	2.06 ^a	568.7	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.38 (br. s., 1 H), 8.34 (s, 1 H), 7.97 (d, J=2.26 Hz, 1 H), 7.72 (s, 1 H), 7.67 (dd, J=8.66, 2.13 Hz, 1 H), 7.39 - 7.52 (m, 2 H), 7.17 (s, 1 H), 4.88 (q, J=8.95 Hz, 2 H), 4.08 - 4.29 (m, 4 H), 2.46 (s, 3 H), 1.43 (q, J=7.03 Hz, 6 H)
71	3-[[6,7-bis(ethyloxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide	2.02 ^a	500.9	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 8.91 (d, J=9.04 Hz, 1 H), 8.77 (s, 1 H), 8.59 (d, J=2.26 Hz, 1 H), 8.46 (dd, J=9.04, 2.26 Hz, 1 H), 7.77 (d, J=2.01 Hz, 1 H), 7.63 (dd, J=8.53, 2.26 Hz, 1 H), 7.34 (q, J=4.77 Hz, 1 H), 7.24 (d, J=8.78 Hz, 1 H), 2.79 (s, 6 H), 2.43 (d, J=4.77 Hz, 3 H)
72	4-(dimethylamino)- <i>N</i> -methyl-3-((7-nitroquinazolin-4-yl)amino)benzenesulfonamide hydrochloride	2.05 ^a	403.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.03 (s, 1 H), 8.28 (s, 1 H), 8.02 - 8.09 (m, 2 H), 7.50 (dd, J=8.53, 2.26 Hz, 1 H), 7.30 (q, J=4.68 Hz, 1 H), 7.22 (d, J=8.53 Hz, 1 H), 6.91 (dd, J=8.78, 2.26 Hz, 1 H), 6.72 (d, J=2.26 Hz, 1 H), 6.07 (s, 2 H), 2.76 (s, 6 H), 2.43 (d, J=4.77 Hz, 3 H)
73	4-(dimethylamino)- <i>N</i> -methyl-3-[[7-[[3-(4-morpholinyl)propyl]oxy]-4-quinazoliny]amino]benzenesulfonamide hydrochloride	1.68 ^a	501.2	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.00 (br. s., 1 H), 8.79 (s, 1 H), 8.15 (s, 1 H), 7.69 (d, J=2.26 Hz, 1 H), 7.65 (dd, J=8.78, 2.26 Hz, 1 H), 7.27 - 7.35 (m, 2 H), 7.24 (d, J=8.78 Hz, 1 H), 4.29 - 4.40 (m, 4 H), 3.75 - 3.83 (m, 4 H), 3.37 (s, 3 H), 3.37 (s, 3 H), 2.81 (s, 6 H), 2.43 (d, J=5.02 Hz, 3 H)
74	<i>N</i> -methyl-3-[[7-[[3-(4-morpholinyl)propyl]oxy]-4-quinazoliny]amino]-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide trifluoroacetate	1.63 ^a	556.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.20 (s, 1 H), 9.13 (d, J=2.26 Hz, 1 H), 8.56 (s, 1 H), 7.75 (dd, J=8.66, 1.63 Hz, 1 H), 7.68 (s, 1 H), 7.62 (dd, J=8.53, 2.26 Hz, 1 H), 6.89 (d, J=2.26 Hz, 1 H), 6.83 (d, J=2.26 Hz, 1 H), 4.43 (q, J=7.03 Hz, 2 H), 3.93 (s, 3 H), 2.52 (s, 3 H), 1.50 (t, J=6.90 Hz, 3 H)
75	<i>N</i> -methyl-3-[[7-[[3-(methyloxy)propyl]oxy]-4-quinazoliny]amino]-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide	1.97 ^a	501.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.99 (s, 1 H), 8.82 - 8.92 (m, 2 H), 7.70 (dd, J=8.78, 2.26 Hz, 1 H), 7.50 - 7.61 (m, 2 H), 7.22 (s, 1 H), 5.12 (q, J=8.78 Hz, 2 H), 4.17 (s, 3 H), 4.04 (s, 3 H), 3.92 (s, 3 H), 2.47 (d, J=4.77 Hz, 3 H)

76	4-(dimethylamino)- <i>N</i> -methyl-3-[(7-[[3-(methyloxy)propyl]oxy]-4-quinazoliny]amino]benzene sulfonamide	1.38 ^a	446.2	¹ H NMR (400 MHz, METHANOL- <i>d</i> ₄) δ ppm 9.02 (d, J=2.01 Hz, 1 H), 8.54 (s, 1 H), 7.59 (dd, J=8.78, 2.26 Hz, 1 H), 7.36 (d, J=8.78 Hz, 1 H), 6.79 (d, J=2.01 Hz, 1 H), 6.70 - 6.76 (m, 1 H), 5.28 (dt, J=12.55, 6.27 Hz, 1 H), 4.08 (s, 3 H), 3.90 (s, 3 H), 2.50 (s, 3 H), 1.52 (d, J=6.27 Hz, 3 H)
77	3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]- <i>N</i> -methyl-4-(methylsulfonyl)benzenesulfonamide trifluoroacetate	1.87 ^a	453.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.57 (br. s., 1 H), 8.76 (s, 1 H), 8.63 (br. s., 1 H), 7.79 (q, J=4.68 Hz, 1 H), 7.59 (d, J=11.54 Hz, 1 H), 6.93 (d, J=2.01 Hz, 1 H), 6.85 (d, J=2.26 Hz, 1 H), 5.08 (q, J=8.78 Hz, 2 H), 4.11 (s, 3 H), 3.98 (s, 3 H), 2.54 (d, J=4.77 Hz, 3 H)
78	3-(7-hydroxy-6-methoxyquinazolin-4-ylamino)- <i>N</i> -methyl-4-(2,2,2-trifluoroethoxy)benzenesulfonamide	0.80 ^b	458.9	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.66 (br. s., 1 H), 8.64 (s, 1 H), 7.91 (br. s., 1 H), 7.86 (br. s., 1 H), 7.77 (d, J=10.58 Hz, 1 H), 7.45 - 7.57 (m, 2 H), 7.23 - 7.32 (m, 1 H), 4.81 - 4.96 (m, 2 H), 4.17 - 4.31 (m, 2 H), 3.95 (s, 3 H), 2.44 (d, J=4.63 Hz, 3 H), 1.42 (t, J=6.73 Hz, 3 H)
79	3-(7-isopropoxy-6-methoxyquinazolin-4-ylamino)- <i>N</i> -methyl-4-(2,2,2-trifluoroethoxy)benzenesulfonamide	0.94 ^b	501.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.64 (br. s., 1 H), 8.80 (s, 1 H), 8.34 (br. s., 1 H), 7.76 (dd, J=8.66, 2.13 Hz, 1 H), 7.48 - 7.61 (m, 2 H), 7.09 (d, J=2.01 Hz, 1 H), 6.91 (d, J=2.01 Hz, 1 H), 5.05 (q, J=8.78 Hz, 2 H), 4.55 - 4.63 (m, 2 H), 3.98 (s, 3 H), 3.77 - 3.83 (m, 2 H), 3.23 (s, 3 H), 2.47 (d, J=5.02 Hz, 3 H), 1.04 (d, J=6.02 Hz, 3 H)
80	3-(7-ethoxy-6-methoxyquinazolin-4-ylamino)- <i>N</i> -methyl-4-(2,2,2-trifluoroethoxy)benzenesulfonamide trifluoroacetate	1.03 ^b	487.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.17 (br. s., 1 H), 8.75 (s, 1 H), 8.02 (s, 1 H), 7.76 - 7.89 (m, 2 H), 7.51 - 7.58 (m, 2 H), 7.39 (s, 2 H), 4.90 (q, J=8.60 Hz, 2 H), 4.26 - 4.34 (m, 2 H), 3.97 (s, 3 H), 3.72 - 3.78 (m, 2 H), 3.32 (s, 3 H), 2.45 (d, J=4.85 Hz, 3 H)

81	3-[6-methoxy-7-(2-methoxy-ethoxy)-quinazolin-4-ylamino]- <i>N</i> -methyl-4-(2,2,2-trifluoro-ethoxy)-benzenesulfonamide trifluoroacetate	1.00 ^b	517.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.91 (s, 1 H), 8.87 (d, J=1.76 Hz, 1 H), 8.48 (s, 1 H), 7.48 - 7.58 (m, 2 H), 7.44 (q, J=4.77 Hz, 1 H), 6.95 (d, J=2.26 Hz, 1 H), 6.87 (d, J=2.26 Hz, 1 H), 5.09 (q, J=8.78 Hz, 2 H), 4.98 (dt, J=9.54, 5.02 Hz, 1 H), 3.86 - 3.95 (m, 5 H), 3.45 - 3.56 (m, 2 H), 2.48 (d, J=4.77 Hz, 3 H), 2.09 - 2.19 (m, 2 H), 1.69 - 1.85 (m, 2 H)
82	3-{{[5,7-bis(methyloxy)-4-quinazolinyl]amino}- <i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide hydrochloride	2.07 ^a	487.1	¹ H NMR (400 MHz, METHANOL- <i>d</i> ₄) δ ppm 9.23 (d, J=2.26 Hz, 1 H), 8.46 (s, 1 H), 7.52 (dd, J=8.78, 2.26 Hz, 1 H), 7.32 (d, J=8.78 Hz, 1 H), 6.71 (dd, J=9.16, 2.13 Hz, 2 H), 5.27 (dt, J=12.61, 6.37 Hz, 1 H), 4.04 (s, 3 H), 3.86 (s, 3 H), 2.50 (s, 3 H), 1.53 (d, J=6.27 Hz, 3 H)
83	3-{{[5,7-bis(methyloxy)-4-quinazolinyl]amino}- <i>N</i> -methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide hydrochloride	2.07 ^a	487.1	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.78 (s, 1 H), 8.93 (s, 1 H), 8.42 (s, 1 H), 7.85 (d, J=2.26 Hz, 1 H), 7.70 (dd, J=8.66, 2.13 Hz, 1 H), 7.45 (d, J=8.53 Hz, 2 H), 7.21 (s, 1 H), 4.87 (q, J=8.78 Hz, 2 H), 4.00 (s, 3 H), 2.46 (d, J=4.77 Hz, 3 H)
84	3-((6-aminoquinazolin-4-yl)amino)-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide	1.54 ^a	372.9	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 9.05 (br. s., 1 H), 8.31 (s, 1 H), 8.03 - 8.11 (m, 2 H), 7.50 (dd, J=8.66, 1.88 Hz, 1 H), 7.22 (d, J=8.53 Hz, 1 H), 6.94 (dd, J=8.91, 1.88 Hz, 1 H), 6.60 - 6.69 (m, 1 H), 6.55 (d, J=2.01 Hz, 1 H), 2.80 (d, J=4.77 Hz, 3 H), 2.76 (s, 6 H), 2.43 (s, 3 H)
85	3-((7-aminoquinazolin-4-yl)amino)-4-(dimethylamino)- <i>N</i> -methylbenzenesulfonamide	1.61 ^a	373.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 8.95 (s, 1 H), 8.46 (d, J=2.26 Hz, 1 H), 8.33 (s, 1 H), 7.56 (d, J=8.78 Hz, 1 H), 7.50 (dd, J=8.28, 2.26 Hz, 1 H), 7.29 - 7.39 (m, 2 H), 7.26 (dd, J=8.78, 2.26 Hz, 1 H), 7.14 (d, J=2.26 Hz, 1 H), 5.76 (s, 2 H), 2.79 (s, 6 H), 2.45 (d, J=5.02 Hz, 3 H)
86	4-(dimethylamino)-3-((6-(dimethylamino)quinazolin-4-yl)amino)- <i>N</i> -methylbenzenesulfonamide bis(trifluoroacetate)	1.80 ^a	401.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.97 (br. s., 1 H), 8.65 (s, 1 H), 7.59 - 7.71 (m, 3 H), 7.46 (dd, J=9.03, 2.26 Hz, 1 H), 7.38 (s, 1 H), 7.28 - 7.35 (m, 1 H), 7.21 - 7.26 (m, 1 H), 2.85 (s, 3 H), 2.81 (s, 6 H), 2.42 (d, J=5.27 Hz, 3 H)

87	4-(dimethylamino)- <i>N</i> -methyl-3-((6-(methylamino)quinazolin-4-yl)amino)benzenesulfonamide bis(trifluoroacetate)	1.71 ^a	387.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 10.77 (br. s., 1 H), 8.78 - 8.87 (m, 1 H), 8.75 (s, 1 H), 8.06 (dd, J=8.91, 1.88 Hz, 1 H), 7.87 (d, J=9.04 Hz, 1 H), 7.74 (d, J=2.01 Hz, 1 H), 7.63 (dd, J=8.78, 2.26 Hz, 1 H), 7.32 (q, J=5.02 Hz, 1 H), 7.23 (d, J=8.78 Hz, 1 H), 2.80 (s, 6 H), 2.43 (d, J=5.02 Hz, 3 H)
88	4-(dimethylamino)- <i>N</i> -methyl-3-((7-(methylamino)quinazolin-4-yl)amino)benzenesulfonamide	1.71 ^a	387.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 11.14 (br. s., 1 H), 8.71 (s, 1 H), 7.72 - 7.77 (m, 1 H), 7.60 - 7.72 (m, 4 H), 7.33 (q, J=5.19 Hz, 1 H), 7.24 (d, J=8.53 Hz, 1 H), 3.13 (s, 6 H), 2.81 (s, 6 H), 2.42 (d, J=5.27 Hz, 3 H)

^a LCMS Method: Agilent 1100 Series LC/MSD SL or VL using electrospray positive [ES+ve to give M+H⁺] equipped with a Sunfire C18 5.0 μm column (3.0 mm x 50 mm, i.d.), eluting with 0.05% TFA in water (solvent A) and 0.05% TFA in acetonitrile (solvent B), using the following elution gradient: 10% – 100% (solvent B) over 2.5 minutes and holding at 100% for 1.7 minutes at a flow rate of 1.0 ml/minutes.

^b LCMS Method: Agilent 1200 Series LC/MSD VL using electrospray positive [ES+ve to give M+H⁺] equipped with a shim-pack XR-ODS 2.2 μm column (3.0 mm x 30 mm, 3.0 mm i.d.) eluting with 0.0375% TFA in water (solvent A) and 0.01875% TFA in acetonitrile (solvent B), using the following elution gradient 10-80% (solvent B) over 0.9 minutes and holding at 80% for 0.6 minutes at a flow rate of 1.2 mL/minutes.

Pharmaceutical Compositions

Example A

Tablets are prepared using conventional methods and are formulated as follows:

Ingredient	Amount per tablet
Compound of Example I	5mg
Microcrystalline cellulose	100mg
Lactose	100mg
Sodium starch glycollate	30mg
Magnesium stearate	2mg
Total	237mg

Example B

Capsules are prepared using conventional methods and are formulated as follows:

Ingredient	Amount per tablet
Compound of Example 3	15mg
Dried starch	178mg
Magnesium stearate	2mg
Total	195mg

Biological Assay(s)

Materials: His-MBP-TEV-Full length human TNNI3K (hTNNI3K) was expressed in *Baculokinase* system and purified from amylose affinity column followed by Superdex200. The fluorescent ligand 5-([2-([3-([4-[(5-hydroxy-2-methylphenyl)amino]-2-pyrimidinyl)amino)phenyl]carbonyl)amino)ethyl]amino)carbonyl)-2-(6-hydroxy-3-oxo-3*H*-xanthen-9-yl)benzoic acid was used. The preparation of this fluorescent ligand is disclosed in U.S. Provisional Patent Application No. 61/237,815 filed August 28, 2009, the disclosure of which is incorporated by reference herein. The other buffer components, including MgCl₂ (Catalog Number M1028), Bis-Tris (Catalog Number B7535), DTT (Catalog Number D9779) and Chaps (Catalog Number C3023) were purchased from Sigma-Aldrich.

Biological Assay Method I:

A fluorescent polarization assay was used to determine does response of compound inhibition on hTNNI3K ATP binding. The binding of 5-([2-([3-([4-[(5-hydroxy-2-methylphenyl)amino]-2-pyrimidinyl)amino)phenyl]carbonyl)amino)ethyl]amino)carbonyl)-2-(6-hydroxy-3-oxo-3*H*-xanthen-9-yl)benzoic acid to the hTNNI3K ATP binding pocket results in increase of fluorescent polarization and the displacement of 5-([2-([3-([4-[(5-hydroxy-2-methylphenyl)amino]-2-pyrimidinyl)amino)phenyl]carbonyl)amino)ethyl]amino)carbonyl)-2-(6-hydroxy-3-oxo-3*H*-xanthen-9-yl)benzoic acid by a competitive compound leads to fluorescent polarization decrease.

Solution 1: Ten (10) mL of a 5 nM 5-([2-([3-([4-[(5-hydroxy-2-methylphenyl)amino]-2-pyrimidinyl)amino)phenyl]carbonyl)amino)ethyl]amino)carbonyl)-2-(6-hydroxy-3-oxo-3*H*-xanthen-9-yl)benzoic acid solution (Solution 1) was prepared by mixing 5 μ L of 1M DTT and 80 μ L of 10% (w/v) Chaps and 5 μ L of a 10 μ M 5-([2-([3-([4-[(5-hydroxy-2-methylphenyl)amino]-2-pyrimidinyl)amino)phenyl]carbonyl)amino)ethyl]amino)carbonyl)-

2-(6-hydroxy-3-oxo-3*H*-xanthen-9-yl)benzoic acid stock solution into 9910 μ L buffer (20 mM Tris, 15 mM MgCl₂, pH 7.5). (Stock solution: 10 μ M solution of 5-({[2-({[3-({4-[(5-hydroxy-2-methylphenyl)amino]-2-pyrimidinyl)amino]phenyl}carbonyl)amino]ethyl]amino}carbonyl)-2-(6-hydroxy-3-oxo-3*H*-xanthen-9-yl)benzoic acid in 100% DMSO)


5 Solution 2 was formed by mixing 53.8 μ L of 2.6 μ M hTNNI3K with a 6946.2 μ L aliquot of Solution 1 (the above 5-({[2-({[3-({4-[(5-hydroxy-2-methylphenyl)amino]-2-pyrimidinyl)amino]phenyl}carbonyl)amino]ethyl]amino}carbonyl)-2-(6-hydroxy-3-oxo-3*H*-xanthen-9-yl)benzoic acid solution) to make up a 7 mL of mixture of hTNNI3K and 5-({[2-({[3-({4-[(5-hydroxy-2-methylphenyl)amino]-2-pyrimidinyl)amino]phenyl}carbonyl)amino]ethyl]amino}carbonyl)-2-(6-hydroxy-3-oxo-3*H*-xanthen-9-yl)benzoic acid (Solution 2).

10 Fifty (50) nL of inhibitors in DMSO (or DMSO controls) were stamped into a 384-well low volume Greiner black plate, followed by addition of 5 μ L of Solution 1 to column 18 and 5 μ L Solution 2 to columns 1-17 and 19-24 of the plate. The plate was then spun at 500 rpm for 30 seconds and incubated at room temperature for 60 minutes. After that, the fluorescent polarization was measured on Analyst (ex/em: 485/530 nm, Dichroic: 505). For dose response experiments, normalized data were fit by ABASE/XC₅₀ and $pXC_{50} = (\log((b-y)/(y-a)))/d - \log(x)$, where x is the compound concentration and y is the % activity at specified compound concentration, a is the minimum % activity, b is the maximum % activity, and d is the Hill slope.

20 The pXC₅₀s are averaged to determine a mean value, for a minimum of 2 experiments. As determined using the above method, the compounds of Examples 1-88 exhibited a pXC₅₀ greater than or equal to 6.0. Preferred compounds of the invention, including the compounds of Examples 1, 3, 4, 5, 10, 11, 14, 16, 17, 18, 19, 20, 22, 26, 28, 30, 36, 38, 39, 40, 42, 44, 45, 47, 50, 51, 53, 55, 58, 59, 60, 61, 64, 65, 67, 69, 70, 74, 75, 77, 80, 81, 84, 85, 86, 87, and 88, exhibited a pXC₅₀ of between approximately 7.0 and approximately 8.0. For instance, the compound of Example 3 inhibited hTNNI3K in the above method with a mean pXC₅₀ of approximately 7.1. More preferred compounds of the invention, including the compounds of Examples 21, 25, 27, 31, 32, 35, 37, 46, 48, 49, 54, 56, 62, 66, 68, 71, 73, 76, 82, and 83, exhibited a pXC₅₀ of between approximately 8.0 and approximately 9.0. For instance, the compound of Example 68 inhibited hTNNI3K in the above method with a mean pXC₅₀ of approximately 8.4.

What is claimed is:

1. A compound according to Formula I:



I

wherein:

R¹ is (C₁-C₄)alkyl;

R² is H or halogen;

R³ is H, halogen, (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, hydroxyl, hydroxy(C₁-C₈)alkyl-, (C₁-C₈)alkoxy, (C₃-C₈)cycloalkyloxy, (C₁-C₈)haloalkoxy, (C₁-C₈)alkylthio-, (C₁-C₈)haloalkylthio-, (C₃-C₈)cycloalkylthio-, (C₁-C₈)alkylsulfonyl, (C₁-C₈)haloalkylsulfonyl, (C₃-C₈)cycloalkylsulfonyl, phenyl, 5-membered heteroaryl, or -N(R^a)(R^b),

wherein said phenyl or heteroaryl is optionally substituted one to three times, independently, by halogen, (C₁-C₆)alkyl, (C₁-C₄)haloalkyl, (C₁-C₆)alkoxy, (C₁-C₄)haloalkoxy, hydroxy(C₁-C₄)alkyl-, or -N(R^a)(R^b);

each R^a is independently (C₁-C₄)alkyl, wherein said (C₁-C₄)alkyl is optionally substituted one to three times, independently, by halogen, hydroxyl, (C₁-C₆)alkoxy, amino, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)((C₁-C₆)alkyl)amino, -CO₂H, -CO₂(C₁-C₆)alkyl, -CONH₂, -CONH(C₁-C₆)alkyl, or -CON((C₁-C₆)alkyl)((C₁-C₆)alkyl), and

R^b is (C₁-C₄)alkyl;

or R^a and R^b taken together with the nitrogen atom to which they are attached form a 5-membered or 6-membered heterocyclic ring, optionally containing one additional heteroatom selected from nitrogen, oxygen and sulfur, wherein said ring is optionally substituted one or two times, independently, by halogen, (C₁-C₄)alkyl, (C₁-C₄)haloalkyl, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, hydroxyl, hydroxy(C₁-C₄)alkyl-, oxo, (C₁-C₄)alkoxy, (C₁-C₄)haloalkoxy, or (C₁-C₄)alkoxy(C₁-C₄)alkyl;

R⁴ is H;

R⁵, R⁶, and R⁷ are each, independently, H, halogen, nitro, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, hydroxyl, or OR^c; and

each R^c is, independently, (C₁-C₈)alkyl or a 5-6 membered heterocycloalkyl, wherein said (C₁-C₈)alkyl is optionally substituted one to three times, independently, by halogen, hydroxyl, (C₁-C₄)alkoxy, amino, (C₁-C₄)alkylamino, or -N(R^a)(R^b);

or R⁶ and R⁷ taken together represent -O(C₁-C₂)alkylo-;

5 or a salt thereof.

2. The compound or salt according to claim 1, wherein R³ is H, halogen, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, (C₁-C₆)alkoxy, (C₅-C₆)cycloalkyloxy, (C₁-C₆)haloalkoxy, (C₁-C₆)alkylthio-, (C₅-C₆)cycloalkylthio-, (C₁-C₆)haloalkylthio-, (C₁-C₆)alkylsulfonyl, 10 (C₁-C₆)haloalkylsulfonyl, (C₅-C₆)cycloalkylsulfonyl, phenyl, 5-membered heteroaryl, or -N(R^a)(R^b), wherein said heteroaryl contains one heteroatom selected from N, O and S, or contains one nitrogen atom and optionally contains 1 additional heteroatom selected from N, O and S, or contains two nitrogen atoms and optionally contains 1 additional heteroatom selected from N, O and S; and said phenyl or heteroaryl is optionally 15 substituted one to three times, independently, by halogen, (C₁-C₆)alkyl, (C₁-C₄)haloalkyl, or -N(R^a)(R^b).

3. The compound or salt according to claim 1, wherein R³ is H, halogen, (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, hydroxyl, hydroxy(C₁-C₈)alkyl-, (C₁-C₈)alkoxy, 20 (C₃-C₈)cycloalkyloxy, (C₁-C₈)haloalkoxy, (C₁-C₈)alkylthio-, (C₁-C₈)haloalkylthio-, (C₃-C₈)cycloalkylthio-, phenyl, 5-membered heteroaryl, or -N(R^a)(R^b), wherein said phenyl or heteroaryl is optionally substituted one to three times, independently, by halogen, (C₁-C₆)alkyl, (C₁-C₄)haloalkyl, (C₁-C₆)alkoxy, (C₁-C₄)haloalkoxy, hydroxy(C₁-C₄)alkyl-, or -N(R^a)(R^b).

25 4. The compound or salt according to claim 1, wherein R³ is H, halogen, C₁-C₆ alkyl, C₁-C₆ haloalkyl, hydroxyC₁-C₆ alkyl-, C₁-C₆ alkoxy, (C₃-C₆)cycloalkyloxy, C₁-C₆ haloalkoxy, C₁-C₆ alkylthio-, (C₃-C₆)cycloalkylthio-, C₁-C₆ haloalkylthio-, phenyl, 5-membered heteroaryl, or -N(R^a)(R^b), wherein said heteroaryl contains one heteroatom 30 selected from N, O and S, or contains one nitrogen atom and optionally contains 1 additional heteroatom selected from N, O and S, or contains two nitrogen atoms and optionally contains 1 additional heteroatom selected from N, O and S; and said phenyl or heteroaryl is optionally substituted one to three times, independently, by halogen, (C₁-C₆)alkyl, (C₁-C₄)haloalkyl, or -N(R^a)(R^b).

35

5. The compound or salt according to claim 1, wherein R³ is H, halogen, (C₁-C₄)alkoxy, (C₅-C₆)cycloalkyloxy, (C₁-C₄)haloalkoxy, (C₁-C₄)alkylthio-, (C₁-C₄)haloalkylthio-, (C₅-C₆)cycloalkylthio-, (C₁-C₄)alkylsulfonyl, (C₁-C₄)haloalkylsulfonyl, (C₅-C₆)cycloalkylsulfonyl, or -N(R^a)(R^b).

5

6. The compound or salt according to any one of claims 1-5, wherein:

each R^a is independently -CH₃, -CH₂CH₃, or -CH₂CF₃; and

R^b is -CH₃;

or R^a and R^b taken together with the nitrogen to which they are attached represent

10 pyrrolidin-1-yl, piperidin-1-yl, or morpholin-4-yl, wherein said pyrrolidin-1-yl, piperidin-1-yl, or morpholin-4-yl is optionally substituted one or two times, independently, by F, -CH₃, or -CF₃.

7. The compound or salt according to any one of claims 1-6, wherein:

15 R⁵, R⁶, and R⁷ are each, independently, H, halogen, nitro, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, hydroxyl, or OR^c, and each R^c is, independently, (C₁-C₄)alkyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, piperidinyl, piperazinyl, morpholinyl, tetrahydropyranyl, tetrahydro-4*H*-1,4-thiazinyl, or 1,4-dioxanyl, wherein said (C₁-C₄)alkyl is optionally substituted by halogen, hydroxyl, trifluoromethyl, 20 (C₁-C₄)alkoxy, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, piperidinyl, piperazinyl, morpholinyl, tetrahydropyranyl, tetrahydro-4*H*-1,4-thiazinyl, or 1,4-dioxanyl;

or R⁶ and R⁷ taken together represent -O(C₁-C₂)alkylO-.

25 8. The compound or salt according to any one of claims 1-7, wherein R⁶ is H.

9. The compound or salt according to any one of claims 1-7, wherein:

R⁶ and R⁷ are each OR^c; and

30 each R^c is, independently, (C₁-C₈)alkyl, optionally substituted one to three times, independently, by halogen, hydroxyl, (C₁-C₄)alkoxy, amino, (C₁-C₄)alkylamino, or ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino.

10. The compound or salt according to claim 1, wherein:

R¹ is (C₁-C₄)alkyl;

35 R² is H or halogen;

R³ is H, halogen, (C₁-C₄)alkoxy, (C₅-C₆)cycloalkyloxy, (C₁-C₄)haloalkoxy, (C₁-C₄)alkylthio-, (C₁-C₄)haloalkylthio-, (C₅-C₆)cycloalkylthio-, (C₁-C₄)alkylsulfonyl, (C₁-C₄)haloalkylsulfonyl, (C₅-C₆)cycloalkylsulfonyl, or -N(R^a)(R^b);

each R^a is independently (C₁-C₄)alkyl, wherein said (C₁-C₄)alkyl is optionally substituted one to three times, independently, by halogen, hydroxyl, (C₁-C₄)alkoxy, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, -CO₂H, -CO₂(C₁-C₄)alkyl, -CONH₂, -CONH(C₁-C₄)alkyl, or -CON((C₁-C₄)alkyl)((C₁-C₄)alkyl), and

R^b is (C₁-C₄)alkyl;

or R^a and R^b taken together with the nitrogen atom to which they are attached form a 5-membered or 6-membered heterocyclic ring, optionally containing one additional heteroatom selected from nitrogen, oxygen and sulfur, wherein said ring is optionally substituted one or two times, independently, by halogen, (C₁-C₄)alkyl, (C₁-C₄)haloalkyl, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, hydroxyl, hydroxy(C₁-C₄)alkyl-, oxo, (C₁-C₄)alkoxy, (C₁-C₄)haloalkoxy, or (C₁-C₄)alkoxy(C₁-C₄)alkyl;

R⁴ is H;

R⁵, R⁶, and R⁷ are each, independently, H, halogen, nitro, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, hydroxyl, or OR^c; and

each R^c is, independently, (C₁-C₄)alkyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, piperidinyl, piperazinyl, morpholinyl, tetrahydropyranyl, tetrahydro-4H-1,4-thiazinyl, or 1,4-dioxanyl, wherein said (C₁-C₄)alkyl is optionally substituted by halogen, hydroxyl, trifluoromethyl, (C₁-C₄)alkoxy, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, piperidinyl, piperazinyl, morpholinyl, tetrahydropyranyl, tetrahydro-4H-1,4-thiazinyl, or 1,4-dioxanyl;

or R⁶ and R⁷ taken together represent -O(C₁-C₂)alkylo-.

11. The compound or salt according to claim 1, wherein:

R¹ is (C₁-C₃)alkyl;

R² is H;

R³ is H, halogen, C₁-C₆ alkyl, C₁-C₆ haloalkyl, hydroxyC₁-C₆ alkyl-, C₁-C₆ alkoxy, (C₃-C₆)cycloalkyloxy, C₁-C₆ haloalkoxy, C₁-C₆ alkylthio-, (C₃-C₆)cycloalkylthio-, C₁-C₆ haloalkylthio-, phenyl, 5-membered heteroaryl, or -N(R^a)(R^b),

wherein said heteroaryl contains one heteroatom selected from N, O and S, or contains one nitrogen atom and optionally contains 1 additional heteroatom selected from N, O and S, or contains two nitrogen atoms and optionally contains 1 additional heteroatom selected from N, O and S; and said phenyl or heteroaryl is optionally

substituted one to three times, independently, by halogen, (C₁-C₆)alkyl, (C₁-C₄)haloalkyl, or -N(R^a)(R^b);

each R^a is independently (C₁-C₄)alkyl, wherein said (C₁-C₄)alkyl is optionally substituted by hydroxyl, trifluoromethyl, (C₁-C₆)alkoxy, amino, (C₁-C₆)alkylamino, or

5 ((C₁-C₆)alkyl)((C₁-C₆)alkyl)amino, and

R^b is (C₁-C₄)alkyl;

or R^a and R^b taken together with the nitrogen atom to which they are attached form a 5-membered or 6-membered heterocyclic ring, optionally containing one additional heteroatom selected from nitrogen, oxygen and sulfur, wherein said ring is optionally

10 substituted one or two times, independently, by (C₁-C₄)alkyl, (C₁-C₄)haloalkyl, hydroxy(C₁-C₄)alkyl-, oxo, or (C₁-C₄)alkoxy(C₁-C₄)alkyl;

R⁴ is H;

R⁵ is H;

R⁶ and R⁷ are each OR^c; and

15 each R^c is, independently, (C₁-C₄)alkyl, optionally substituted one to three times, independently, by halogen.

12. A compound selected from:

20 3-[[6,7-bis(methyloxy)-4-quinazolinyl]amino]-4-(dimethylamino)-*N*-methylbenzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methylbenzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-(4-morpholinyl)benzenesulfonamide;

25 3-[[6,7-bis(methyloxy)-4-quinazolinyl]amino]-4-[ethyl(methyl)amino]-*N*-methylbenzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(1-methylethyl)oxy]benzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-(methyloxy)benzenesulfonamide;

30 3-[[6,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-(methylthio)benzenesulfonamide;

4-(dimethylamino)-*N*-methyl-3-(4-quinazolinylamino)benzenesulfonamide;

3-[(6-chloro-4-quinazolinyl)amino]-4-(dimethylamino)-*N*-methylbenzenesulfonamide;

35 4-(dimethylamino)-*N*-methyl-3-[[6-(methyloxy)-4-quinazolinyl]amino]benzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

3-[(6-chloro-4-quinazoliny)amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

5 3-(7,8-dihydro[1,4]dioxino[2,3-*g*]quinazolin-4-ylamino)-4-(dimethylamino)-*N*-methylbenzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-4-(4,4-difluoro-1-piperidiny)-*N*-methylbenzenesulfonamide;

10 3-[(6-chloro-4-quinazoliny)amino]-4-(4,4-difluoro-1-piperidiny)-*N*-methylbenzenesulfonamide;

3-({7-[(3-chloropropyl)oxy]-4-quinazoliny}amino)-4-(dimethylamino)-*N*-methylbenzenesulfonamide;

3-[[6,7-bis{[2-(methyloxy)ethyl]oxy}-4-quinazoliny]amino]-4-(dimethylamino)-*N*-methylbenzenesulfonamide;

15 3-[[6,7-bis{[2-(methyloxy)ethyl]oxy}-4-quinazoliny]amino]-*N*-methyl-4-(methyloxy)benzenesulfonamide;

3-[[6,7-bis{[2-(methyloxy)ethyl]oxy}-4-quinazoliny]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

20 3-[[6,7-bis{[2-(methyloxy)ethyl]oxy}-4-quinazoliny]amino]-*N*-methyl-4-(methylthio)benzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[(2*R*)-2-(trifluoromethyl)-1-pyrrolidiny]benzenesulfonamide;

3-[(6-chloro-4-quinazoliny)amino]-*N*-methyl-4-[(2*R*)-2-(trifluoromethyl)-1-pyrrolidiny]benzenesulfonamide;

25 5-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-2-fluoro-*N*-methylbenzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide;

30 3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-(1-piperidiny)benzenesulfonamide;

3-[(6-chloro-4-quinazoliny)amino]-*N*-methyl-4-(1-piperidiny)benzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-4-(2,5-dimethyl-1-pyrrolidiny)-*N*-methylbenzenesulfonamide;

35 3-[(6-chloro-4-quinazoliny)amino]-4-(2,5-dimethyl-1-pyrrolidiny)-*N*-methylbenzenesulfonamide;

3-{{6,7-bis(methyloxy)-4-quinazolinyl}amino}-*N*-methyl-4-[(2,2,2-trifluoroethyl)thio]benzenesulfonamide;

3-{{5,7-bis(methyloxy)-4-quinazolinyl}amino}-*N*-methylbenzenesulfonamide;

3-{{5,7-bis(methyloxy)-4-quinazolinyl}amino}-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

3-{{5,7-bis(methyloxy)-4-quinazolinyl}amino}-4-(dimethylamino)-*N*-methylbenzenesulfonamide;

3-(5-chloro-6,7-dimethoxyquinazolin-4-ylamino)-*N*-methylbenzenesulfonamide;

4-methoxy-3-(7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-ylamino)-*N*-methylbenzenesulfonamide;

3-{{6,7-bis(methyloxy)-4-quinazolinyl}amino}-*N*-methyl-4-[methyl(2,2,2-trifluoroethyl)amino]benzenesulfonamide;

3-{{5,7-bis(methyloxy)-4-quinazolinyl}amino}-*N*-methyl-4-[methyl(2,2,2-trifluoroethyl)amino]benzenesulfonamide;

3-{{5,7-bis(methyloxy)-4-quinazolinyl}amino}-*N*-methyl-4-(methyloxy)benzenesulfonamide;

3-{{5,7-bis(methyloxy)-4-quinazolinyl}amino}-4-fluoro-*N*-methylbenzenesulfonamide;

3-{{5,7-bis(methyloxy)-4-quinazolinyl}amino}-4-chloro-*N*-methylbenzenesulfonamide;

3-{{5,7-bis(methyloxy)-4-quinazolinyl}amino}-*N*-methyl-4-(4-morpholinyl)benzenesulfonamide;

5-{{6,7-bis(methyloxy)-4-quinazolinyl}amino}-2-fluoro-*N*-methyl-4-(methyloxy)benzenesulfonamide;

3-{{6-(ethyloxy)-7-(methyloxy)-4-quinazolinyl}amino}-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

4-(dimethylamino)-3-(7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-ylamino)-*N*-methylbenzenesulfonamide;

N-methyl-3-{{6-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazolinyl}amino}-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

3-{{7-[(3-chloropropyl)oxy]-4-quinazolinyl}amino}-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

3-{{5-(ethyloxy)-7-(methyloxy)-4-quinazolinyl}amino}-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

N-methyl-3-{{5-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazolinyl}amino}-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

N-methyl-4-[(2,2,2-trifluoroethyl)oxy]-3-[[5,6,7-tris(methyloxy)-4-quinazolinyl]amino]benzenesulfonamide;

3-[[5,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide;

5 5-[[5,7-bis(methyloxy)-4-quinazolinyl]amino]-2-fluoro-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

5-[[5,7-bis(methyloxy)-4-quinazolinyl]amino]-2-fluoro-*N*-methyl-4-(methyloxy)benzenesulfonamide;

10 5-[[6,7-bis(methyloxy)-4-quinazolinyl]amino]-2-fluoro-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

3-[[5,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-(methylthio)benzenesulfonamide;

3-[[5,7-bis(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide;

15 *N*-methyl-4-[(trifluoromethyl)oxy]-3-[[5,6,7-tris(methyloxy)-4-quinazolinyl]amino]benzenesulfonamide;

5-[[5,7-bis(methyloxy)-4-quinazolinyl]amino]-4-(dimethylamino)-2-fluoro-*N*-methylbenzenesulfonamide;

20 5-[[6,7-bis(methyloxy)-4-quinazolinyl]amino]-4-(dimethylamino)-2-fluoro-*N*-methylbenzenesulfonamide;

2-fluoro-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]-5-[[5,6,7-tris(methyloxy)-4-quinazolinyl]amino]benzenesulfonamide;

4-chloro-*N*-methyl-3-[[5,6,7-tris(methyloxy)-4-quinazolinyl]amino]benzenesulfonamide;

25 *N*-methyl-3-[[7-(methyloxy)-5-[[2-(methyloxy)ethyl]oxy]-4-quinazolinyl]amino]-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

N-methyl-3-[[7-(methyloxy)-5-(tetrahydro-2*H*-pyran-4-yloxy)-4-quinazolinyl]amino]-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

30 *N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]-3-[[5,6,7-tris(methyloxy)-4-quinazolinyl]amino]benzenesulfonamide;

N-methyl-4-(methylsulfonyl)-3-[[5,6,7-tris(methyloxy)-4-quinazolinyl]amino]benzenesulfonamide;

3-[[5-(ethyloxy)-7-(methyloxy)-4-quinazolinyl]amino]-*N*-methyl-4-[(trifluoromethyl)oxy]benzenesulfonamide;

35 *N*-methyl-3-[[5-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazolinyl]amino]-4-[(trifluoromethyl)oxy]benzenesulfonamide;

3-[[5-(ethyloxy)-7-(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide;

N-methyl-3-[[5-[(1-methylethyl)oxy]-7-(methyloxy)-4-quinazoliny]amino]-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide;

5 3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-(methylsulfonyl)benzenesulfonamide;

3-[[6,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide;

10 3-[[6-iodo-7-(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

3-[[6,7-bis(ethyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

4-(dimethylamino)-*N*-methyl-3-((7-nitroquinazolin-4-yl)amino)benzenesulfonamide;

15 4-(dimethylamino)-*N*-methyl-3-[(7-{[3-(4-morpholiny)propyl]oxy}-4-quinazoliny]amino]benzenesulfonamide;

N-methyl-3-[(7-{[3-(4-morpholiny)propyl]oxy}-4-quinazoliny]amino]-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

N-methyl-3-[(7-{[3-(methyloxy)propyl]oxy}-4-quinazoliny]amino]-4-[(2,2,2-trifluoroethyl)oxy]benzenesulfonamide;

20 4-(dimethylamino)-*N*-methyl-3-[(7-{[3-(methyloxy)propyl]oxy}-4-quinazoliny]amino]benzenesulfonamide;

3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-(methylsulfonyl)benzenesulfonamide;

25 3-(7-hydroxy-6-methoxyquinazolin-4-ylamino)-*N*-methyl-4-(2,2,2-trifluoroethoxy)benzenesulfonamide;

3-(7-isopropoxy-6-methoxy-quinazolin-4-ylamino)-*N*-methyl-4-(2,2,2-trifluoroethoxy)-benzenesulfonamide;

3-(7-ethoxy-6-methoxy-quinazolin-4-ylamino)-*N*-methyl-4-(2,2,2-trifluoro-ethoxy)-benzenesulfonamide;

30 3-[6-methoxy-7-(2-methoxy-ethoxy)-quinazolin-4-ylamino]-*N*-methyl-4-(2,2,2-trifluoro-ethoxy)-benzenesulfonamide;

3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide;

35 3-[[5,7-bis(methyloxy)-4-quinazoliny]amino]-*N*-methyl-4-[(2,2,2-trifluoro-1-methylethyl)oxy]benzenesulfonamide;

3-((6-aminoquinazolin-4-yl)amino)-4-(dimethylamino)-*N*-methylbenzenesulfonamide;

3-((7-aminoquinazolin-4-yl)amino)-4-(dimethylamino)-*N*-methylbenzenesulfonamide;

5 4-(dimethylamino)-3-((6-(dimethylamino)quinazolin-4-yl)amino)-*N*-methylbenzenesulfonamide;

4-(dimethylamino)-*N*-methyl-3-((6-(methylamino)quinazolin-4-yl)amino)benzenesulfonamide; and

10 4-(dimethylamino)-*N*-methyl-3-((7-(methylamino)quinazolin-4-yl)amino)benzenesulfonamide;

or a pharmaceutically acceptable salt thereof.

13. A pharmaceutical composition comprising the compound or salt according to any one of claims 1-12 and one or more pharmaceutically-acceptable excipients.

15

14. A method of inhibiting TNNI3K comprising contacting TNNI3K with the compound or salt according to any one of claims 1-12.

15. A method for treating congestive heart failure comprising administering to a patient in need thereof an effective amount of the compound or salt according to any one of claims 1-12.

20

16. A method for treating congestive heart failure comprising administering to a patient in need thereof the pharmaceutical composition according to claim 13.

25