ADENO-ASSOCIATED VIRUS VECTOR DELIVERY OF MUSCLE SPECIFIC MICRO-DYSTROPHIN TO TREAT MUSCULAR DYSTROPHY

[0001] This application claims priority to U.S. Provisional Patent Application No. 62/473,148, filed March 17, 2017 which is incorporated by reference herein in its entirety.

INCORPORATION BY REFERENCE OF MATERIAL SUBMITTED ELECTRONICALLY

[0002] This application contains, as a separate part of the disclosure, a Sequence Listing in computer-readable form which is incorporated by reference in its entirety and identified as follows: Filename: 51475_Seqlisting.txt; Size: 29,519 bytes, created; March 13, 2018.

FIELD OF INVENTION

[0003] The invention provides gene therapy vectors, such as adeno-associated virus (AAV) vectors, expressing a miniaturized human micro-dystrophin gene and method of using these vectors to express micro-dystrophin in skeletal muscles including diaphragm and cardiac muscle and to protect muscle fibers from injury, increase muscle strength and reduce and/or prevent fibrosis in subjects suffering from muscular dystrophy.

BACKGROUND

[0004] The importance of muscle mass and strength for daily activities, such as locomotion and breathing, and for whole body metabolism is unequivocal. Deficits in muscle function produce muscular dystrophies (MDs) that are characterized by muscle weakness and wasting and have serious impacts on quality of life. The most well-characterized MDs result from mutations in genes encoding members of the dystrophin-associated protein complex (DAPC). These MDs result from membrane fragility associated with the loss of sarcolemmal-cytoskeleton tethering by the DAPC. Duchenne Muscular Dystrophy (DMD) is one of the most devastating muscle disease affecting 1 in 5000 newborn males.

[0005] DMD is caused by mutations in the DMD gene leading to reductions in mRNA and the absence of dystrophin, a 427 kD sarcolemmal protein associated with the dystrophin-associated protein complex (DAPC) (Hoffman *et al., Cell* 51(6):919-28, 1987). The DAPC is composed of multiple proteins at the muscle sarcolemma

that form a structural link between the extra-cellular matrix (ECM) and the cytoskeleton via dystrophin, an actin binding protein, and alpha-dystroglycan, a laminin-binding protein. These structural links act to stabilize the muscle cell membrane during contraction and protect against contraction-induced damage. With dystrophin loss, membrane fragility results in sarcolemmal tears and an influx of calcium, triggering calcium-activated proteases and segmental fiber necrosis (Straub *et al., Curr Opin. Neurol.* 10(2): 168-75, 1997). This uncontrolled cycle of muscle degeneration and regeneration ultimately exhausts the muscle stem cell population (Sacco *et al., Cell*, 2010. 143(7): p. 1059-71; Wallace *et al., Annu Rev Physiol,* 2009. 71: p. 37-57), resulting in progressive muscle weakness, endomysial inflammation, and fibrotic scarring.

[0006] Without membrane stabilization from dystrophin or a micro-dystrophin, DMD will manifest uncontrolled cycles of tissue injury and repair ultimately replace lost muscle fibers with fibrotic scar tissue through connective tissue proliferation. Fibrosis is characterized by the excessive deposits of ECM matrix proteins, including collagen and elastin. ECM proteins are primarily produced from cytokines such as TGFβ that is released by activated fibroblasts responding to stress and inflammation. Although the primary pathological feature of DMD is myofiber degeneration and necrosis, fibrosis as a pathological consequence has equal repercussions. The over-production of fibrotic tissue restricts muscle regeneration and contributes to progressive muscle weakness in the DMD patient. In one study, the presence of fibrosis on initial DMD muscle biopsies was highly correlated with poor motor outcome at a 10-year follow-up (Desguerre *et al., J Neuropathol Exp Neurol*, 2009. 68(7): p. 762-7). These results point to fibrosis as a major contributor to DMD muscle dysfunction and highlight the need for early intervention prior to overt fibrosis.

[0007] Adeno-associated virus (AAV) is a replication-deficient parvovirus, the single-stranded DNA genome of which is about 4.7 kb in length including 145 nucleotide inverted terminal repeat (ITRs). There are multiple serotypes of AAV. The nucleotide sequences of the genomes of the AAV serotypes are known. For example, the nucleotide sequence of the AAV serotype 2 (AAV2) genome is presented in Srivastava *et al., J Virol,* 45: 555-564 (1983) as corrected by Ruffing *et al., J Gen Virol,* 75: 3385-3392 (1994). As other examples, the complete genome of

AAV-1 is provided in GenBank Accession No. NC_002077; the complete genome of AAV-3 is provided in GenBank Accession No. NC_1829; the complete genome of AAV-4 is provided in GenBank Accession No. NC_001829; the AAV-5 genome is provided in GenBank Accession No. AF085716; the complete genome of AAV-6 is provided in GenBank Accession No. NC_00 1862; at least portions of AAV-7 and AAV-8 genomes are provided in GenBank Accession Nos. AX753246 and AX753249, respectively (see also U.S. Patent Nos. 7,282,199 and 7,790,449 relating to AAV-8); the AAV-9 genome is provided in Gao *et al., J. Virol., 78*: 6381-6388 (2004); the AAV-10 genome is provided in *Mol. Ther.,* 13(1): 67-76 (2006); and the AAV-11 genome is provided in *Virology,* 330(2): 375-383 (2004). Cloning of the AAVrh.74 serotype is described in Rodino-Klapac., *et al. Journal of translational medicine* **5**, 45 (2007). *Cis*-acting sequences directing viral DNA replication (rep), encapsidation/packaging and host cell chromosome integration are contained within the ITRs. Three AAV promoters (named p5, p19, and p40 for their relative map locations) drive the expression of the two AAV internal open reading frames encoding rep and cap genes. The two rep promoters (p5 and p19), coupled with the differential splicing of the single AAV intron (*e.g.*, at AAV2 nucleotides 2107 and 2227), result in the production of four rep proteins (rep 78, rep 68, rep 52, and rep 40) from the rep gene. Rep proteins possess multiple enzymatic properties that are ultimately responsible for replicating the viral genome. The cap gene is expressed from the p40 promoter and it encodes the three capsid proteins VP1, VP2, and VP3. Alternative splicing and non-consensus translational start sites are responsible for the production of the three related capsid proteins. A single consensus polyadenylation site is located at map position 95 of the AAV genome. The life cycle and genetics of AAV are reviewed in Muzyczka, *Current Topics in Microbiology and Immunology, 158*: 97-129 (1992).

[0008] AAV possesses unique features that make it attractive as a vector for delivering foreign DNA to cells, for example, in gene therapy. AAV infection of cells in culture is noncytopathic, and natural infection of humans and other animals is silent and asymptomatic. Moreover, AAV infects many mammalian cells allowing the possibility of targeting many different tissues *in vivo*. Moreover, AAV transduces slowly dividing and non-dividing cells, and can persist essentially for the lifetime of those cells as a transcriptionally active nuclear episome (extrachromosomal element).

The AAV proviral genome is infectious as cloned DNA in plasmids which makes construction of recombinant genomes feasible. Furthermore, because the signals directing AAV replication, genome encapsidation and integration are contained within the ITRs of the AAV genome, some or all of the internal approximately 4.3 kb of the genome (encoding replication and structural capsid proteins, rep-cap) may be replaced with foreign DNA such as a gene cassette containing a promoter, a DNA of interest and a polyadenylation signal. The rep and cap proteins may be provided *in trans*. Another significant feature of AAV is that it is an extremely stable and hearty virus. It easily withstands the conditions used to inactivate adenovirus (56°C to 65°C for several hours), making cold preservation of AAV less critical. AAV may even be lyophilized. Finally, AAV-infected cells are not resistant to superinfection.

[0009] Multiple studies have demonstrated long-term (> 1.5 years) recombinant AAV-mediated protein expression in muscle. See, Clark *et al., Hum Gene Ther, 8*: 659-669 (1997); Kessler *et al., Proc Nat. Acad Sc. USA, 93*: 14082-14087 (1996); and Xiao *et al., J Virol, 70*: 8098-8108 (1996). See also, Chao *et al., Mol Ther, 2*:619-623 (2000) and Chao *et al., Mol Ther, 4*:217-222 (2001). Moreover, because muscle is highly vascularized, recombinant AAV transduction has resulted in the appearance of transgene products in the systemic circulation following intramuscular injection as described in Herzog *et al., Proc Natl Acad Sci USA, 94*: 5804-5809 (1997) and Murphy *et al., Proc Natl Acad Sci USA, 94*: 13921-13926 (1997). Moreover, Lewis *et al., J Virol, 76*: 8769-8775 (2002) demonstrated that skeletal myofibers possess the necessary cellular factors for correct antibody glycosylation, folding, and secretion, indicating that muscle is capable of stable expression of secreted protein therapeutics.

[0010] Functional improvement in patients suffering from DMD and other muscular dystrophies requires gene restoration at an early stage of disease. There is a need for treatments that increase muscle strength and protect against muscle injury in patients suffering from DMD.

SUMMARY OF INVENTION

[0011] The present invention is directed to gene therapy vectors, e.g. AAV, expressing the micro-dystrophin gene to skeletal muscles including diaphragm and cardiac muscle to protect muscle fibers from injury, increase muscle strength and reduce and/or prevent fibrosis

[0012] The invention provides for therapies and approaches for increasing muscular force and/or increasing muscle mass using gene therapy vectors to deliver micro-dystrophin to address the gene defect observed in DMD. As shown in Example 2, treatment with micro-dystrophin gene therapy resulted in a greater muscle force *in vivo*. Furthermore, delivery of micro-dystrophin gene therapy intramuscularly and systemically showed delivery of dystrophin to the muscles in mice models *in vivo*.

[0013] In one embodiment, the invention provides for a rAAV vector comprising a muscle specific control element nucleotide sequence, and a nucleotide sequence encoding the micro-dystrophin protein. For example, the nucleotide sequence encodes a functional micro-dystrophin protein, wherein the nucleotide is, e.g., at least 65%, at least 70%, at least 75%, at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, or 89%, more typically at least 90%, 91%, 92%, 93%, or 94% and even more typically at least 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to SEQ ID NO: 1, wherein the protein retains micro-dystrophin activity. The micro-dystrophin protein provides stability to the muscle membrane during muscle contraction, e.g. micro-dystrophin acts as a shock absorber during muscle contraction.

[0014] The invention also provides for rAAV vectors wherein the nucleotide sequence encodes a functional micro-dystrophin protein comprising a nucleotide sequence that hybridizes under stringent conditions to the nucleic acid sequence of SEQ ID NO: 1, or compliments thereof, and encodes a functional micro-dystrophin protein.

[0015] In one embodiment, the rAAV vector is a non-replicating, recombinant adeno-associated virus (AAV) termed rAAVrh74.MHCK7.micro-dystrophin. This vector genome contains minimal elements required for gene expression, including AAV2 inverted terminal repeats (ITR), the micro-dystrophin, SV40 intron (SD/SA), and synthetic polyadenylation (Poly A) signal, all under the control of the MHCK7 promoter/enhancer. The schematic of the vector genome and expression cassette is shown **Error! Reference source not found.**. The AAVrh74 serotype can be employed to achieve efficient gene transfer in skeletal and cardiac muscle following IV administration.

[0016] The term "stringent" is used to refer to conditions that are commonly understood in the art as stringent. Hybridization stringency is principally determined

by temperature, ionic strength, and the concentration of denaturing agents such as formamide. Examples of stringent conditions for hybridization and washing are 0.015 M sodium chloride, 0.0015 M sodium citrate at 65-68°C or 0.015 M sodium chloride, 0.0015M sodium citrate, and 50% formamide at 42°C. See Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory, (Cold Spring Harbor, N.Y. 1989). More stringent conditions (such as higher temperature, lower ionic strength, higher formamide, or other denaturing agent) may also be used, however, the rate of hybridization will be affected. In instances wherein hybridization of deoxyoligonucleotides is concerned, additional exemplary stringent hybridization conditions include washing in 6x SSC 0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos).

[0017] Other agents may be included in the hybridization and washing buffers for the purpose of reducing non-specific and/or background hybridization. Examples are 0.1% bovine serum albumin, 0.1% polyvinyl-pyrrolidone, 0.1% sodium pyrophosphate, 0.1% sodium dodecylsulfate, NaDodSO4, (SDS), ficoll, Denhardt's solution, sonicated salmon sperm DNA (or other non-complementary DNA), and dextran sulfate, although other suitable agents can also be used. The concentration and types of these additives can be changed without substantially affecting the stringency of the hybridization conditions. Hybridization experiments are usually carried out at pH 6.8-7.4, however, at typical ionic strength conditions, the rate of hybridization is nearly independent of pH. See Anderson *et al.*, Nucleic Acid Hybridisation: A Practical Approach, Ch. 4, IRL Press Limited (Oxford, England). Hybridization conditions can be adjusted by one skilled in the art in order to accommodate these variables and allow DNAs of different sequence relatedness to form hybrids.

[0018] The term "muscle specific control element" refers to a nucleotide sequence that regulates expression of a coding sequence that is specific for expression in muscle tissue. These control elements include enhancers and promoters. The invention provides for constructs comprising the muscle specific control elements MCKH7 promoter, the MCK promoter and the MCK enhancer.

[0019] The term "operably linked" refers to the positioning of the regulatory element nucleotide sequence, e.g. promoter nucleotide sequence, to confer expression of said nucleotide sequence by said regulatory element.

[0020] In one aspect, the invention provides for a rAAV vector wherein the muscle specific control element is a human skeletal actin gene element, cardiac actin gene element, myocyte-specific enhancer binding factor (MEF), muscle creatine kinase (MCK), truncated MCK (tMCK), myosin heavy chain (MHC), hybrid α-myosin heavy chain enhancer-/MCK enhancer-promoter (MHCK7), C5-12, murine creatine kinase enhancer element, skeletal fast-twitch troponin c gene element, slow-twitch cardiac troponin c gene element, the slow-twitch troponin i gene element, hypoxia-inducible nuclear factors, steroid-inducible element or glucocorticoid response element (GRE).

[0021] For example, the muscle specific control element is the MHCK7 promoter nucleotide sequence SEQ ID NO: 2 or the muscle specific control element is MCK nucleotide sequence SEQ ID NO: 4. In addition, in any of the rAAV vectors of the invention, the muscle specific control element nucleotide sequence, e.g. MHCK7 or MCK nucleotide sequence, is operably linked to the nucleotide sequence encoding the micro-dystrophin protein. For example, the MHCK7 promoter nucleotide sequence (SEQ ID NO: 2) is operably linked to the human micro-dystrophin coding sequence (SEQ ID NO: 1) as set out in the construct provided in Figure 1 or Figure 10 (SEQ ID NO: 3). In another example, the MCK promoter (SEQ ID NO: 4) is operably linked to the human micro-dystrophin coding sequence (SEQ ID NO: 1) as set out in the construct provided in Figure 7 or Figure 11 (SEQ ID NO: 5). In another aspect, the invention provides for a rAAV vector comprising the nucleotide sequence of SEQ ID NO: 1 and SEQ ID NO: 2. The invention also provides for a rAAV vector comprising the nucleotide sequence of SEQ ID NO: 1 and SEQ ID NO: 4.

[0022] In a further aspect, the invention provides for a rAAV vector comprising the nucleotide sequence of SEQ ID NO: 3 or SEQ ID NO: 5. For example, the rAAVrh74.MHCK7.microdystrophin vector comprises the nucleotide sequence of SEQ ID NO: 3 and shown in Figure 10. This rAAV vector comprises the MHCK7 promoter, a chimeric intron sequence, the coding sequence for the human micro-dystrophin gene, polyA, ampicillin resistance and the pGEX plasmid backbone with pBR322 origin or replication.

[0023] The invention provides for a recombinant AAV vector comprising the human micro-dystrophin nucleotide sequence of SEQ ID NO: 1 and the MHCK7 promoter nucleotide sequence of SEQ ID NO: 3. This rAAV vector is the AAV serotype AAVrh.74.

[0024] The invention also provides for a recombinant AAV vector comprising the pAAV.MHCK7.micro-dystrophin construct nucleotide sequence of SEQ ID NO: 3. This rAAV vector is the AAV serotype AAVrh.74.

[0025] The rAAV vectors of the invention may be any AAV serotype, such as the serotype AAVrh.74, AAV1, AAV2, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12 or AAV13.

[0026] The invention also provides for pharmaceutical compositions (or sometimes referred to herein as simply "compositions") comprising any of the rAAV vectors of the invention.

[0027] In another embodiment, the invention provides for methods of producing a rAAV vector particle comprising culturing a cell that has been transfected with any rAAV vector of the invention and recovering rAAV particles from the supernatant of the transfected cells. The invention also provides for viral particles comprising any of the recombinant AAV vectors of the invention.

[0028] The invention provides for methods of treating muscular dystrophy comprising administering a therapeutically effective amount of any of the recombinant AAV vectors of the invention expressing human micro-dystrophin.

[0029] The invention provides for methods of treating muscular dystrophy comprising administering a therapeutically effective amount of a recombinant AAV vector comprising the human micro-dystrophin nucleotide sequence of SEQ ID NO: 1 and the MHCK7 promoter nucleotide sequence of SEQ ID NO: 2.

[0030] The invention also provides for methods of treating muscular dystrophy comprising administering a therapeutically effective amount of a recombinant AAV vector comprising the pAAV.MHCK7.micro-dystrophin construct nucleotide sequence of SEQ ID NO: 3.

[0031] "Fibrosis" refers to the excessive or unregulated deposition of extracellular matrix (ECM) components and abnormal repair processes in tissues upon injury,

including skeletal muscle, cardiac muscle, liver, lung, kidney, and pancreas. The ECM components that are deposited include fibronectin and collagen, e.g. collagen 1, collagen 2 or collagen 3.

[0032] The invention also provides for methods of reducing or preventing fibrosis in a subject suffering from muscular dystrophy comprising administering a therapeutically effective amount of any recombinant AAV vector of the invention.

[0033] In another embodiment, the invention provides for methods of preventing fibrosis in a subject in need thereof, comprising administering a therapeutically effective amount of a recombinant AAV vector of the invention. For example, any of the rAAV of the invention can be administered to subjects suffering from muscular dystrophy to prevent fibrosis, e.g. the rAAV of the invention expressing a human micro-dystrophin protein administered before fibrosis is observed in the subject. In addition, the rAAV of the invention expressing a human micro-dystrophin gene can be administered to a subject at risk of developing fibrosis, such as those suffering or diagnosed with muscular dystrophy, e.g. DMD. The rAAV of the invention can be administered to the subject suffering from muscular dystrophy in order to prevent new fibrosis in these subjects.

[0034] The invention contemplates administering any of the AAV vectors of the invention before fibrosis is observed in the subject. In addition, the rAAV of the invention can be administered to a subject at risk of developing fibrosis, such as those suffering or diagnosed with muscular dystrophy, e.g. DMD. The rAAV of the invention can be administered to the subject suffering from muscular dystrophy who already has developed fibrosis in order to prevent new fibrosis in these subjects.

[0035] The invention also provides for methods of increasing muscular force and/or muscle mass in a subject suffering from muscular dystrophy comprising administering a therapeutically effective amount of a rAAV vector of the invention expressing a human micro-dystrophin gene. These methods can further comprise the step of administering a rAAV expressing micro-dystrophin.

[0036] The invention contemplates administering any of the AAV vectors of the invention to patients diagnosed with DMD before fibrosis is observed in the subject or before the muscle force has been reduced or before the muscle mass has been reduced.

[0037] The invention also contemplates administering a AAV of the invention to a subject suffering from muscular dystrophy who already has developed fibrosis, in order to prevent new fibrosis in these subjects or to reduce fibrosis in these patients. The invention also provides for administering any of the rAAV of the invention to the patient suffering from muscular dystrophy who already has reduced muscle force or has reduced muscle mass in order to protect the muscle from further injury.

[0038] In any of the methods of the invention, the subject may be suffering from muscular dystrophy such as DMD or any other dystrophin-associated muscular dystrophy.

[0039] In another aspect, a rAAV vector expressing the micro-dystrophin protein comprises the coding sequence of the micro-dystrophin gene operably linked to a muscle-specific control element other than MHCK7 or MCK. For example, whrein the muscle-specific control element is human skeletal actin gene element, cardiac actin gene element, myocyte-specific enhancer binding factor MEF, tMCK (truncated MCK), myosin heavy chain (MHC), C5-12 (synthetic promoter), murine creatine kinase enhancer element, skeletal fast-twitch troponin C gene element, slow-twitch cardiac troponin C gene element, the slow-twitch troponin I gene element, hypozia-inducible nuclear factors, steroid-inducible element, or glucocorticoid response element (GRE).

[0040] In any of the methods of the invention, the rAAV vector or composition can be administered by intramuscular injection or intravenous injection.

[0041] In addition, in any of the methods of the invention, the rAAV vector or composition can be administered systemically. For example, the rAAV vector or composition can be parenterally administration by injection, infusion, or implantation.

[0042] In another embodiment, the invention provides a composition comprising any of the rAAV vectors of the invention for reducing fibrosis in a subject in need thereof.

[0043] In addition, the invention provides a composition comprising any of the recombinant AAV vectors of the invention for preventing fibrosis in a patient suffering from muscular dystrophy.

[0044] The invention provides for compositions comprising any of the recombinant AAV vectors of the invention for treating muscular dystrophy.

[0045] The invention provides for compositions comprising a recombinant AAV vector comprising the human micro-dystrophin nucleotide sequence of SEQ ID NO: 1 and the MHCK7 promoter sequence of SEQ ID NO: 2 for treatment of muscular dystrophy.

[0046] The invention provides for a composition comprising a recombinant AAV vector comprising the pAAV.MHCK7.micro-dystrophin construct comprising the nucleotide sequence of SEQ ID NO: 3 for treatment of muscular dystrophy.

[0047] The invention also provides for compositions comprising any of the rAAV vectors of the invention for increasing muscular force and/or muscle mass in a subject suffering from muscular dystrophy. In a further embodiment, the invention provides for compositions comprising any of the rAAV vectors of the invention for treatment of muscular dystrophy.

[0048] The compositions of the invention can be formulated for intramuscular injection or intravenous injection. The composition of the invention is also formulated for systemic administration, such as parenterally administration by injection, infusion or implantation.

[0049] In addition, any of the compositions can be formulated for administration to a subject suffering from muscular dystrophy such as DMD or any other dystrophin associated muscular dystrophy.

[0050] In a further embodiment, the invention provides for use of any of the rAAV vectors of the invention for preparation of a medicament for reducing fibrosis in a subject in need thereof. For example, the subject in need can be suffering from muscular dystrophy, such as DMD or any other dystrophin associated muscular dystrophy.

[0051] In another embodiment, the invention provides for use of a rAAV vector of the invention for the preparation of a medicament to prevent fibrosis in a subject suffering from muscular dystrophy.

[0052] In addition, the invention provides for use of a recombinant AAV vector of the invention to preparation of a medicament to increase muscular strength and/or muscle mass in a subject suffering from muscular dystrophy.

[0053] The invention also provides for use of the rAAV vectors of the invention for the preparation of a medicament for treatment of muscular dystrophy.

[0054] The invention provides for use of a recombinant AAV vector comprising the human micro-dystrophin nucleotide sequence of SEQ ID NO: 1 and the MHCK7 promoter nucleotide sequence of SEQ ID NO: 2 for preparation of a medicament for the treatment of muscular dystrophy.

[0055] The invention provides for use of a recombinant AAV vector comprising the pAAV.MHCK7.micro-dystrophin construct nucleotide sequence of SEQ ID NO: 3 for treatment of muscular dystrophy.

[0056] In any of the uses of the invention, the medicament can be formulated for intramuscular injection or intravenous injection. In addition, in any of the uses of the invention, the medicament can be formulated for systemic administration such as parenteral administration by injection, infusion, or implantation.

[0057] Any of the medicaments can be prepared for administration to a subject suffering from muscular dystrophy such as DMD or any other dystrophin associated muscular dystrophy.

BRIEF DESCRIPTION OF THE DRAWINGS

[0058] **Figure 1** illustrates the pAAV.MHCK7.micro-dystrophin construct. In this construct, the cDNA expression cassette is flanked by AAV2 inverted terminal repeat sequences (ITR). The construct is characterized by an in-frame rod deletion (R4–R23), while hinges 1, 2 and 4 (H_1, H_2 and H_4) and the cysteine rich domain remain producing a 138 kDa protein. The expression of the micro-dystrophin protein (3579 bp) is guided by a MHCK7 promoter (795 bp). The intron and 5' UTR are derived from plasmid pCMVß (Clontech). The micro-dystrophin cassette had a consensus Kozak immediately in front of the ATG start and a small 53 bp synthetic polyA signal for mRNA termination. The human micro-dystrophin cassette contained the (R4–

R23/Δ71–78) domains as previously described by Harper *et al.* (*Nature Medicine* 8, 253-261 (2002)).

[0059] **Figure 2** demonstrates dystrophin protein expression following intramuscular delivery of AAVrh74.MHCK7 construct. The tibialis anterior muscle of mdx mice was injected with 1 x 10^{11} vg (n=5 per group). Six weeks later the muscles were harvested and stained for dystrophin expression with an N-terminal antibody for dystrophin and hematoxylin and eosin staining.

[0060] **Figures 3A-3C** provide skeletal muscle force measurements and quantification of micro-dystrophin expression following intramuscular injection of AAVrh74.MHCK7 construct. (A) The tibialis anterior muscle of mdx mice was injected with1 x 10^{11} vg (n=5) with AAVrh74.MHCK7 construct. Six weeks later the tibialis anterior muscles were harvested and subjected to in vivo force measurements. The dosed cohort had significantly greater force production than untreated mdx controls.

[0061] **Figures 4A-4C** demonstrate widespread transduction of skeletal, diaphragm and cardiac muscle fibers after systemic administration of the AAVrh.74.MHCK7.micro-dys construct. (A) Mdx mice were treated systemically at 6 weeks of age via the tail vein with 6x10^{12} vg (2 x 10^{14} vg/kg) of AAVrh.74.MHCK7.micro-dystrophin following 12 weeks of treatment. (B) Staining for micro-dystrophin demonstrates quantification of the percentage of muscle fibers expressing micro-dystrophin in each tissue. (C) Shows the specific force measured in the diaphragm at the low and high (planned clinical) dose. No significant difference was seen at low dose; however there was significant improvement at the high dose.

[0062] **Figure 5** demonstrates dystrophin protein expression following systemic delivery of AAVrh.74.MHCK7.micro-dystrophin construct. Mdx mice (n=5) were treated systemically starting at 6 weeks of age via the tail vein with 6 x 10^{12}vg of AAVrh.74.MHCK7.micro-dystrophin. Following 12 weeks of treatment, all muscles were harvested and stained for dystrophin and restoration of DAPC components (beta-sarcoglycan shown).

[0063] **Figures 6A-6D** demonstrate the toxicology/safety of AAVrh.74.MHCK7. Hematoxylin and eosin (H&E) staining was performed on the following muscle tissues to analyze toxicity: Tibialis anterior (TA), Gastrocnemius (GAS), Quadriceps

(QD), Psoas (PSO), Triceps (TRI), and Diaphragm (DIA) (Figure 6A). No toxicity was noted. As an indicator of efficacy, the number of muscle fibers with centrally placed nuclei (CN) was quantified (Figure 6B). CN are indicative of cycles of muscle degeneration and regeneration and thus reduction in CN demonstrates treatment effect. (Figure 6C) demonstrates the total number of fibers is unchanged with treatment. The amount of creatine kinase is provided in (D) showing improvement at high dose. Independent t-tests were used to locate differences ($p<.05$); Data are reported as means ± SEM.

[0064] **Figure 7** illustrates the pAAV.MCK.micro-dystrophin plasmid construct.

[0065] **Figure 8** provides the results of a rAAVrh74.MCK. micro-dystrophin (human) potency assay. The tibialis anterior muscle of mdx mice was injected with 3 x 10^9, 3 x 10^{10}, or 1 x 10^{11} vg (n=3 per group). Four weeks later the muscles were harvested and stained for dystrophin expression with the N-terminal Dys3 antibody. There was a linear correlation between expression and dose with very little expression (no effect level) at 3 x 10^9 vg and 89% expression at 1 x 10^{11} vg.

[0066] **Figures 9A-9C** demonstrate that Human micro-dystrophin improves force generation and protection from eccentric contraction induced injury. (A) Dystrophin protein immunostaining in the extensor digitorum longus (EDL) and TA shows expression in a mdx myofibers following rAAVrh.74-MCK-micro-dystrophin (human) injection via the femoral artery. Mock-infected muscle was stained in an identical manner and exposures are time matched. (B) rAAVrh.74-MCK-micro-dystrophin significantly increased normalized specific force relative to mock-treated mdx muscles ($P<0.05$ vs. mdx). (C) mdx muscles infected with rAAVrh.74-MCK-Micro-dys(human) were compared with mock-infected contralateral mdx EDL muscles and WT (WT C57Bl/10) EDL muscles for force drop during repetitive eccentric contractions at 12 weeks post gene transfer. rAAVrh.74-MCK-microdystrophin (Micro-dys) treatment significantly protected against loss of force compared with mock-treated mdx muscles ($P< 0.001$ vs. mdx). Errors are SEMs.

[0067] **Figure 10** provides the nucleic acid sequence (SEQ ID NO: 3 rAAVrh74.MHCK7. micro-dystrophin).

[0068] **Figure 11** provides the nucleic acid sequence (SEQ ID NO: 5) rAAVrh74.MCK.micro-dystrophin.

[0069] **Figures 12A – 12B** provides the immunological response to systemic delivery of AAVrh74.MHCK7.micro-dystrophin in the non-human primate. (A) ELISpot response to AAV capsid and micro-dystrophin peptide pools. ConA is the positive control and DMSO is the negative control. There were three pools to AAVrh74 and four peptide pools specific to micro-dystrophin. (B) ELISA positive titers of circulating neutralizing antibodies to vector capsid. Serum was isolated from primates biweekly and analyzed for antibody titer. Titer reported corresponds to last dilution at which ratio of response ≥ 2.

[0070] **Figure 13A -B** demonstrates systemic delivery in rhesus macaque with AAVrh74.MHCK7.microdystrophin. Anti-FLAG immunofluorescence staining in the left side muscles demonstrated robust micro-dystrophin expression.

[0071] **Figure 14** demonstrates the effect of systemic treatment with rAAVrh74.MHCK7.micro-dystrophin on transgene expression. Immunofluorescence staining for micro-dystrophin using an N-terminal dystrophin antibody in the heart, diaphragm, psoas, and tibialis anterior (TA) demonstrates robust expression in the mid (6e12 vg; 2e14 vg/kg) and high dose (1.2e13 vg; 6e14 vg/kg) treated animals 3 months post-injection. 20x images are shown.

[0072] **Figure 15** demonstrates the effect of systemic treatment with rAAVrh74.MHCK7.micro-dystrophin on transgene expression. Immunofluorescence staining for micro-dystrophin using an N-terminal dystrophin antibody in the gastrocnemius, quadriceps, tricep and gluteus demonstrates robust expression in the mid (6e12 vg; 2e14 vg/kg) and highest dose (1.2e13 vg; 6e14 vg/kg) treated animals 3 months post-injection. 20x images are shown.

[0073] **Figure 16** demonstrates the effect of systemic treatment with rAAVrh74.MHCK7.micro-dystrophin on muscle pathology. (A) H&E stain of diaphragm, tibialis anterior, gastrocnemius, and quadricep muscle from C57BL/6 WT, mdx, and rAAVrh74.MHCK7.micro-dystrophin treated mice (Mid dose-2e14vg/kg; high dose-6e14vg/kg), (B) Quantification of average fiber size demonstrated a normalization of fiber size across all tissue. ****$p<0.001$, one-way ANOVA; Data are reported as means ± SEM. 20x images are shown.

[0074] **Figure 17** demonstrates the effect of systemic treatment with rAAVrh74.MHCK7.micro-dystrophin on muscle pathology. (A) H&E stain of tricep,

gluteal and psoas muscle from C57BL/6 WT, mdx, and rAAVrh74.MHCK7.micro-dystrophin treated mice (mid dose-2e14vg/kg; high dose-6e14vg/kg), (B) Quantification of average fiber size demonstrated larger fibers in a dose dependent manner. ****p<0.001, one-way ANOVA; Data are reported as means ± SEM. 20x images are shown.

[0075] **Figure 18** demonstrates the effect of systemic treatment with rAAVrh74.MHCK7.micro-dystrophin on central nucleation. Dose escalation illustrates reductions in central nucleation in all skeletal muscles and diaphragm. Two-way ANOVA were used to locate differences ($p<0.05$). Data are reported as means ± SEM.

[0076] **Figure 19** demonstrates the effect of systemic treatment with rAAVrh74.MHCK7.micro-dystrophin on collagen deposition. Dose escalation illustrates reductions in collagen accumulation (%) in the diaphragm. *p<0.05, one-way ANOVA; Data are reported as means ± SEM. 20x images are shown.

[0077] **Figure 20** demonstrates correction of force deficits in the diaphragm. Following 3 or 6 months of treatment, diaphragm muscle strips were harvested to measure specific force (normalized to cross sectional area). Treatment restored force to WT levels. *p<0.05. One-way ANOVA was utilized to determine differences from mdx-LR mice.

[0078] **Figure 21** demonstrates correction of force deficits in the TA. (A) Following 3-6 months of treatment TA muscles were harvested (both Left and Right) to measure specific force (normalized to TA weight). Treatment restored force to WT levels. (B) Treatment rescued TA muscles from fatigue after rigorous protocol of eccentric contractions. *p<0.05. One-way ANOVA was utilized to determine differences from mdx-LR mice.

[0079] **Figure 22** provides distribution of average vg copies in various tissues from three mdx mice after IV delivery of rAAVrh74.MHCK7.micro-dystrophin.

[0080] **Figure 23.** Serum chemistries for ssAAVrh74.MHCK7.micro-dystrophin systemically injected mice and age matched control groups serum chemistries were analyzed by an independent CRO (Charles River Laboratories) which indicate normal values across all chemistries analyzed. The only abnormal values were elevated AST and ALT noted in MDX vehicle treated animals [MDX-LR (lactated ringers)] that

was normalized with treatment. AST and ALT are known to be elevated in DMD. ALT = alanine aminotransferase, ALP/K = alkaline phosphatase, AST = aspartate aminotransferase, BUN = blood urea nitrogen, B/C = Blood to creatinine ratio, CREAT = creatine, GLU = glucose, TP= total protein, TBIL = total bilirubin, DBIL = direct bilirubin

[0081] **Figure 24** provides biodistribution western blots on muscles and organs from rAAVrh74.MHCK7.micro-dystrophin systemically injected mdx mice.

[0082] **Figure 25** provides the pNLREP2-Caprh74 AAV helper plasmid map.

[0083] **Figure 26** provides the Ad Helper plasmid pHELP.

DETAILED DESCRIPTION

[0084] The present invention provides for gene therapy vectors, e.g. rAAV vectors, overexpressing human micro-dystrophin and methods of reducing and preventing fibrosis in muscular dystrophy patients. Muscle biopsies taken at the earliest age of diagnosis of DMD reveal prominent connective tissue proliferation. Muscle fibrosis is deleterious in multiple ways. It reduces normal transit of endomysial nutrients through connective tissue barriers, reduces the blood flow and deprives muscle of vascular-derived nutritional constituents, and functionally contributes to early loss of ambulation through limb contractures. Over time, treatment challenges multiply as a result of marked fibrosis in muscle. This can be observed in muscle biopsies comparing connective tissue proliferation at successive time points. The process continues to exacerbate leading to loss of ambulation and accelerating out of control, especially in wheelchair-dependent patients.

[0085] Without early treatment including a parallel approach to reduce fibrosis it is unlikely that the benefits of exon skipping, stop-codon read-through, or gene replacement therapies can ever be fully achieved. Even small molecules or protein replacement strategies are likely to fail without an approach to reduce muscle fibrosis. Previous work in aged *mdx* mice with existing fibrosis treated with AAV.micro-dystrophin demonstrated that we could not achieve full functional restoration (Liu, M., *et al., Mol Ther* 11, 245-256 (2005)). It is also known that progression of DMD cardiomyopathy is accompanied by scarring and fibrosis in the ventricular wall.

[0086] As used herein, the term "AAV" is a standard abbreviation for adeno-associated virus. Adeno-associated virus is a single-stranded DNA parvovirus that grows only in cells in which certain functions are provided by a co-infecting helper virus. There are currently thirteen serotypes of AAV that have been characterized. General information and reviews of AAV can be found in, for example, Carter, 1989, Handbook of Parvoviruses, Vol. 1, pp. 169-228, and Berns, 1990, Virology, pp. 1743-1764, Raven Press, (New York). However, it is fully expected that these same principles will be applicable to additional AAV serotypes since it is well known that the various serotypes are quite closely related, both structurally and functionally, even at the genetic level. (See, for example, Blacklowe, 1988, pp. 165-174 of Parvoviruses and Human Disease, J. R. Pattison, ed.; and Rose, Comprehensive Virology 3:1-61 (1974)). For example, all AAV serotypes apparently exhibit very similar replication properties mediated by homologous rep genes; and all bear three related capsid proteins such as those expressed in AAV2. The degree of relatedness is further suggested by heteroduplex analysis which reveals extensive cross-hybridization between serotypes along the length of the genome; and the presence of analogous self-annealing segments at the termini that correspond to "inverted terminal repeat sequences" (ITRs). The similar infectivity patterns also suggest that the replication functions in each serotype are under similar regulatory control.

[0087] An "AAV vector" as used herein refers to a vector comprising one or more polynucleotides of interest (or transgenes) that are flanked by AAV terminal repeat sequences (ITRs). Such AAV vectors can be replicated and packaged into infectious viral particles when present in a host cell that has been transfected with a vector encoding and expressing rep and cap gene products.

[0088] An "AAV virion" or "AAV viral particle" or "AAV vector particle" refers to a viral particle composed of at least one AAV capsid protein and an encapsidated polynucleotide AAV vector. If the particle comprises a heterologous polynucleotide (i.e. a polynucleotide other than a wild-type AAV genome such as a transgene to be delivered to a mammalian cell), it is typically referred to as an "AAV vector particle" or simply an "AAV vector". Thus, production of AAV vector particle necessarily includes production of AAV vector, as such a vector is contained within an AAV vector particle.

AAV

[0089] Recombinant AAV genomes of the invention comprise nucleic acid molecule of the invention and one or more AAV ITRs flanking a nucleic acid molecule. AAV DNA in the rAAV genomes may be from any AAV serotype for which a recombinant virus can be derived including, but not limited to, AAV serotypes AAVrh.74, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-6, AAV-7, AAV-8, AAV-9, AAV-10, AAV-11, AAV-12 and AAV-13. Production of pseudotyped rAAV is disclosed in, for example, WO 01/83692. Other types of rAAV variants, for example rAAV with capsid mutations, are also contemplated. See, for example, Marsic *et al., Molecular Therapy*, 22(11): 1900-1909 (2014). As noted in the Background section above, the nucleotide sequences of the genomes of various AAV serotypes are known in the art. To promote skeletal muscle specific expression, AAV1, AAV6, AAV8 or AAVrh.74 can be used.

[0090] DNA plasmids of the invention comprise rAAV genomes of the invention. The DNA plasmids are transferred to cells permissible for infection with a helper virus of AAV (e.g., adenovirus, E1-deleted adenovirus or herpesvirus) for assembly of the rAAV genome into infectious viral particles. Techniques to produce rAAV particles, in which an AAV genome to be packaged, rep and cap genes, and helper virus functions are provided to a cell are standard in the art. Production of rAAV requires that the following components are present within a single cell (denoted herein as a packaging cell): a rAAV genome, AAV rep and cap genes separate from (*i.e.*, not in) the rAAV genome, and helper virus functions. The AAV rep and cap genes may be from any AAV serotype for which recombinant virus can be derived and may be from a different AAV serotype than the rAAV genome ITRs, including, but not limited to, AAV serotypes AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-6, AAV-7, AAVrh.74, AAV-8, AAV-9, AAV-10, AAV-11, AAV-12 and AAV-13. Production of pseudotyped rAAV is disclosed in, for example, WO 01/83692 which is incorporated by reference herein in its entirety.

[0091] A method of generating a packaging cell is to create a cell line that stably expresses all the necessary components for AAV particle production. For example, a plasmid (or multiple plasmids) comprising a rAAV genome lacking AAV rep and cap genes, AAV rep and cap genes separate from the rAAV genome, and a selectable marker, such as a neomycin resistance gene, are integrated into the genome of a cell.

AAV genomes have been introduced into bacterial plasmids by procedures such as GC tailing (Samulski *et al.*, 1982, *Proc. Natl. Acad. S6. USA*, 79:2077-2081), addition of synthetic linkers containing restriction endonuclease cleavage sites (Laughlin *et al.*, 1983, Gene, 23:65-73) or by direct, blunt-end ligation (Senapathy & Carter, 1984, J. Biol. Chem., 259:4661-4666). The packaging cell line is then infected with a helper virus such as adenovirus. The advantages of this method are that the cells are selectable and are suitable for large-scale production of rAAV. Other examples of suitable methods employ adenovirus or baculovirus rather than plasmids to introduce rAAV genomes and/or rep and cap genes into packaging cells.

[0092] General principles of rAAV production are reviewed in, for example, Carter, 1992, Current Opinions in Biotechnology, 1533-539; and Muzyczka, 1992, Curr. Topics in Microbial. and Immunol., 158:97-129). Various approaches are described in Ratschin *et al.*, *Mol. Cell. Biol.* 4:2072 (1984); Hermonat *et al., Proc. Natl. Acad. Sci. USA*, 81:6466 (1984); Tratschin et al., *Mol. Cell. Biol.* 5:3251 (1985); McLaughlin *et al., J. Virol.*, 62:1963 (1988); and Lebkowski *et al., Mol. Cell. Biol.,* 7:349 (1988). Samulski *et al.*, J. Virol., 63:3822-3828 (1989); U.S. Patent No. 5,173,414; WO 95/13365 and corresponding U.S. Patent No. 5,658.776 ; WO 95/13392; WO 96/17947; PCT/US98/18600; WO 97/09441 (PCT/US96/14423); WO 97/08298 (PCT/US96/13872); WO 97/21825 (PCT/US96/20777); WO 97/06243 (PCT/FR96/01064); WO 99/11764; Perrin *et al. Vaccine* 13:1244-1250 (1995); Paul *et al. Human Gene Therapy* 4:609-615 (1993); Clark *et al. Gene Therapy* 3:1124-1132 (1996); U.S. Patent. No. 5,786,211; U.S. Patent No. 5,871,982; and U.S. Patent. No. 6,258,595. The foregoing documents are hereby incorporated by reference in their entirety herein, with particular emphasis on those sections of the documents relating to rAAV production.

[0093] The invention thus provides packaging cells that produce infectious rAAV. In one embodiment packaging cells may be stably transformed cancer cells such as HeLa cells, 293 cells and PerC.6 cells (a cognate 293 line). In another embodiment, packaging cells are cells that are not transformed cancer cells, such as low passage 293 cells (human fetal kidney cells transformed with E1 of adenovirus), MRC-5 cells (human fetal fibroblasts), WI-38 cells (human fetal fibroblasts), Vero cells (monkey kidney cells) and FRhL-2 cells (rhesus fetal lung cells).

[0094] Recombinant AAV (*i.e.*, infectious encapsidated rAAV particles) of the invention comprise a rAAV genome. In exemplary embodiments, the genomes of both rAAV lack AAV rep and cap DNA, that is, there is no AAV rep or cap DNA between the ITRs of the genomes. Examples of rAAV that may be constructed to comprise the nucleic acid molecules of the invention are set out in International Patent Application No. PCT/US2012/047999 (WO 2013/016352) incorporated by reference herein in its entirety.

[0095] In an exemplary embodiment, the recombinant AAV vector of the inveiton is produced by the triple transfection method (Xiao et al. , *J Virol* **72**, 2224-2232 (1998) using the AAV vector plasmids pAAV.MHCK7.micro-dystrophin, pNLRep2-Caprh74 and pHelp, pAAV contains the micro-dystrophin gene expression cassette flanked by AAV2 inverted terminal repeat sequences (ITR). It is this sequence that is encapsidated into AAVrh74 virions. The plasmid contains the micro-dystrophin sequence and the MHCK7 enhancer and core promoter elements of the muscle specific promoter to drive gene expression. The expression cassette also contains an SV40 intron (SD/SA) to promote high-level gene expression and the bovine growth hormone polyadenylation signal is used for efficient transcription termination.

[0096] The pNLREP2-Caprh74 is an AAV helper plasmid that encodes the 4 wild-type AAV2 rep proteins and the 3 wild-type AAV VP capsid proteins from serotype rh74. A schematic map of the pNLREP2-Caprh74 plasmid is shown in **Error! Reference source not found.**25.

[0097] The pHELP adenovirus helper plasmid is 11,635 bp and was obtained from Applied Viromics. The plasmid contains the regions of adenovirus genome that are important for AAV replication, namely E2A, E4ORF6, and VA RNA (the adenovirus E1 functions are provided by the 293 cells). The adenovirus sequences present in this plasmid only represents ~40% of the adenovirus genome, and does not contain the *cis* elements critical for replication such as the adenovirus terminal repeats. Therefore, no infectious adenovirus is expected to be generated from such a production system. . A schematic map of the pHELP plasmid is shown in Figure 26.

[0098] The rAAV may be purified by methods standard in the art such as by column chromatography or cesium chloride gradients. Methods for purifying rAAV vectors from helper virus are known in the art and include methods disclosed in, for

example, Clark *et al., Hum. Gene Ther., 10*(6): 1031-1039 (1999); Schenpp and Clark, *Methods Mol. Med., 69* 427-443 (2002); U.S. Patent No. 6,566,118 and WO 98/09657.

[0099] In another embodiment, the invention contemplates compositions comprising rAAV of the present invention. Compositions of the invention comprise rAAV and a pharmaceutically acceptable carrier. The compositions may also comprise other ingredients such as diluents and adjuvants. Acceptable carriers, diluents and adjuvants are nontoxic to recipients and are preferably inert at the dosages and concentrations employed and include buffers and surfactants such as pluronics.

[00100] Titers of rAAV to be administered in methods of the invention will vary depending, for example, on the particular rAAV, the mode of administration, the treatment goal, the individual, and the cell type(s) being targeted, and may be determined by methods standard in the art. Titers of rAAV may range from about $1x10^6$, about $1x10^7$, about $1x10^8$, about $1x10^9$, about $1x10^{10}$, about $1x10^{11}$, about $1x10^{12}$, about $1x10^{13}$ to about $1x10^{14}$ or more DNase resistant particles (DRP) per ml. Dosages may also be expressed in units of viral genomes (vg).

[00101] Methods of transducing a target cell with rAAV, *in vivo* or *in vitro*, are contemplated by the invention. The *in vivo* methods comprise the step of administering an effective dose, or effective multiple doses, of a composition comprising a rAAV of the invention to an animal (including a human being) in need thereof. If the dose is administered prior to development of a disorder/disease, the administration is prophylactic. If the dose is administered after the development of a disorder/disease, the administration is therapeutic. In embodiments of the invention, an effective dose is a dose that alleviates (eliminates or reduces) at least one symptom associated with the disorder/disease state being treated, that slows or prevents progression to a disorder/disease state, that slows or prevents progression of a disorder/disease state, that diminishes the extent of disease, that results in remission (partial or total) of disease, and/or that prolongs survival. An example of a disease contemplated for prevention or treatment with methods of the invention is DMD.

[00102] Combination therapies are also contemplated by the invention. Combination as used herein includes both simultaneous treatment and sequential

treatments. Combinations of methods of the invention with standard medical treatments (*e.g.,* corticosteroids) are specifically contemplated, as are combinations with novel therapies.

[00103] Administration of an effective dose of the compositions may be by routes standard in the art including, but not limited to, intramuscular, parenteral, intravenous, oral, buccal, nasal, pulmonary, intracranial, intraosseous, intraocular, rectal, or vaginal. Route(s) of administration and serotype(s) of AAV components of the rAAV (in particular, the AAV ITRs and capsid protein) of the invention may be chosen and/or matched by those skilled in the art taking into account the infection and/or disease state being treated and the target cells/tissue(s) that are to express the micro-dystrophin protein.

[00104] The invention provides for local administration and systemic administration of an effective dose of rAAV and compositions of the invention. For example, systemic administration is administration into the circulatory system so that the entire body is affected. Systemic administration includes enteral administration such as absorption through the gastrointestinal tract and parenteral administration through injection, infusion or implantation.

[00105] In particular, actual administration of rAAV of the present invention may be accomplished by using any physical method that will transport the rAAV recombinant vector into the target tissue of an animal. Administration according to the invention includes, but is not limited to, injection into muscle and injection into the bloodstream. Simply resuspending a rAAV in phosphate buffered saline has been demonstrated to be sufficient to provide a vehicle useful for muscle tissue expression, and there are no known restrictions on the carriers or other components that can be co-administered with the rAAV (although compositions that degrade DNA should be avoided in the normal manner with rAAV). Capsid proteins of a rAAV may be modified so that the rAAV is targeted to a particular target tissue of interest such as muscle. See, for example, WO 02/053703, the disclosure of which is incorporated by reference herein. Pharmaceutical compositions can be prepared as injectable formulations or as topical formulations to be delivered to the muscles by transdermal transport. Numerous formulations for both intramuscular injection and transdermal transport have been previously developed and can be used in the practice of the

invention. The rAAV can be used with any pharmaceutically acceptable carrier for ease of administration and handling.

[00106] The dose of rAAV to be administered in methods disclosed herein will vary depending, for example, on the particular rAAV, the mode of administration, the treatment goal, the individual, and the cell type(s) being targeted, and may be determined by methods standard in the art. Titers of each rAAV administered may range from about $1x10^6$, about $1x10^7$, about $1x10^8$, about $1x10^9$, about $1x10^{10}$, about $1x10^{11}$, about $1x10^{12}$, about $1x10^{13}$, about $1x10^{14}$, or to about $1x10^{15}$ or more DNase resistant particles (DRP) per ml. Dosages may also be expressed in units of viral genomes (vg) (*i.e.*, $1x10^7$ vg, $1x10^8$ vg, $1x10^9$ vg, $1x10^{10}$ vg, $1x10^{11}$ vg, $1x10^{12}$ vg, $1x10^{13}$ vg, $1x10^{14}$ vg, $1x10^{15}$ respectively). Dosages may also be expressed in units of viral genomes (vg) per kilogram (kg) of bodyweight (*i.e.*, $1x10^{10}$ vg/kg, $1x10^{11}$ vg/kg, $1x10^{12}$ vg/kg, $1x10^{13}$ vg/kg, $1x10^{14}$ vg/kg, $1x10^{15}$ vg/kg respectively). Methods for titering AAV are described in Clark *et al., Hum. Gene Ther., 10*: 1031-1039 (1999).

[00107] In particular, actual administration of rAAV of the present invention may be accomplished by using any physical method that will transport the rAAV recombinant vector into the target tissue of an animal. Administration according to the invention includes, but is not limited to, injection into muscle and injected into the bloodstream. Simply resuspending a rAAV in phosphate buffered saline has been demonstrated to be sufficient to provide a vehicle useful for muscle tissue expression, and there are no known restrictions on the carriers or other components that can be co-administered with the rAAV (although compositions that degrade DNA should be avoided in the normal manner with rAAV). Capsid proteins of a rAAV may be modified so that the rAAV is targeted to a particular target tissue of interest such as muscle. See, for example, WO 02/053703, the disclosure of which is incorporated by reference herein. Pharmaceutical compositions can be prepared as injectable formulations or as topical formulations to be delivered to the muscles by transdermal transport. Numerous formulations for both intramuscular injection and transdermal transport have been previously developed and can be used in the practice of the invention. The rAAV can be used with any pharmaceutically acceptable carrier for ease of administration and handling.

[00108] For purposes of intramuscular injection, solutions in an adjuvant such as sesame or peanut oil or in aqueous propylene glycol can be employed, as well as

sterile aqueous solutions. Such aqueous solutions can be buffered, if desired, and the liquid diluent first rendered isotonic with saline or glucose. Solutions of rAAV as a free acid (DNA contains acidic phosphate groups) or a pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxpropylcellulose. A dispersion of rAAV can also be prepared in glycerol, liquid polyethylene glycols and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms. In this connection, the sterile aqueous media employed are all readily obtainable by standard techniques well-known to those skilled in the art.

[00109] The pharmaceutical carriers, diluents or excipients suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating actions of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of a dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal and the like. In many cases it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by use of agents delaying absorption, for example, aluminum monostearate and gelatin.

[00110] Sterile injectable solutions are prepared by incorporating rAAV in the required amount in the appropriate solvent with various other ingredients enumerated above, as required, followed by filter sterilization. Generally, dispersions are prepared by incorporating the sterilized active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the

freeze drying technique that yield a powder of the active ingredient plus any additional desired ingredient from the previously sterile-filtered solution thereof.

[00111] Transduction with rAAV may also be carried out *in vitro*. In one embodiment, desired target muscle cells are removed from the subject, transduced with rAAV and reintroduced into the subject. Alternatively, syngeneic or xenogeneic muscle cells can be used where those cells will not generate an inappropriate immune response in the subject.

[00112] Suitable methods for the transduction and reintroduction of transduced cells into a subject are known in the art. In one embodiment, cells can be transduced *in vitro* by combining rAAV with muscle cells, *e.g.*, in appropriate media, and screening for those cells harboring the DNA of interest using conventional techniques such as Southern blots and/or PCR, or by using selectable markers. Transduced cells can then be formulated into pharmaceutical compositions, and the composition introduced into the subject by various techniques, such as by intramuscular, intravenous, subcutaneous and intraperitoneal injection, or by injection into smooth and cardiac muscle, using *e.g.*, a catheter.

[00113] Transduction of cells with rAAV of the invention results in sustained expression of the micro-dystrophin protein. The present invention thus provides methods of administering/delivering rAAV which express micro-dystrophin protein to an animal, preferably a human being. These methods include transducing tissues (including, but not limited to, tissues such as muscle, organs such as liver and brain, and glands such as salivary glands) with one or more rAAV of the present invention. Transduction may be carried out with gene cassettes comprising tissue specific control elements. For example, one embodiment of the invention provides methods of transducing muscle cells and muscle tissues directed by muscle specific control elements, including, but not limited to, those derived from the actin and myosin gene families, such as from the myoD gene family (See Weintraub *et al., Science, 251*: 761-766 (1991)), the myocyte-specific enhancer binding factor MEF-2 (Cserjesi and Olson, *Mol Cell Biol* 11: 4854-4862 (1991)), control elements derived from the human skeletal actin gene (Muscat *et al., Mol Cell Biol, 7*: 4089-4099 (1987)), the cardiac actin gene, muscle creatine kinase sequence elements (See Johnson *et al., Mol Cell Biol, 9*:3393-3399 (1989)) and the murine creatine kinase enhancer (mCK) element, control elements derived from the skeletal fast-twitch troponin C gene, the

slow-twitch cardiac troponin C gene and the slow-twitch troponin I gene: hypoxia-inducible nuclear factors (Semenza *et al., Proc Natl Acad Sci USA, 88*: 5680-5684 (1991)), steroid-inducible elements and promoters including the glucocorticoid response element (GRE) (See Mader and White, *Proc. Natl. Acad. Sci. USA* 90: 5603-5607 (1993)), and other control elements.

[00114] Muscle tissue is an attractive target for *in vivo* DNA delivery, because it is not a vital organ and is easy to access. The invention contemplates sustained expression of microdystrophin from transduced myofibers.

[00115] By "muscle cell" or "muscle tissue" is meant a cell or group of cells derived from muscle of any kind (for example, skeletal muscle and smooth muscle, *e.g.* from the digestive tract, urinary bladder, blood vessels or cardiac tissue). Such muscle cells may be differentiated or undifferentiated, such as myoblasts, myocytes, myotubes, cardiomyocytes and cardiomyoblasts.

[00116] The term "transduction" is used to refer to the administration/delivery of the coding region of the micro-dystrophin to a recipient cell either *in vivo* or *in vitro*, via a replication-deficient rAAV of the invention resulting in expression of micro-dystrophin by the recipient cell.

[00117] Thus, the invention provides methods of administering an effective dose (or doses, administered essentially simultaneously or doses given at intervals) of rAAV that encode micro-dystrophin to a patient in need thereof.

EXAMPLES

Example 1

Generation of the pAAV.MHCK7.micro-dystrophin construct

[00118] The pAAV.MHCK7.micro-dystrophin plasmid contains a human micro-dystrophin cDNA expression cassette flanked by AAV2 inverted terminal repeat sequences (ITR) (see Fig. 1). The micro-dystrophin construct was characterized by an in-frame rod deletion (R4–R23), while hinges 1, 2 and 4 and cysteine rich domain remain producing a 138 kDa protein. The expression of the micro-dystrophin protein (3579 bp) was guided by a MHCK7 promoter (795 bp). The plasmid was constructed from the pAAV.MCK.micro-dystrophin plasmid by removing the MCK promoter and inserting the MHCK7 promoter. After the core promoter, the 53 bp endogenous

mouse MCK Exon1 (untranslated) is present for efficient transcription initiation, followed by the SV40 late 16S/19S splice signals (97 bp) and a small 5'UTR (61 bp). The intron and 5' UTR are derived from plasmid pCMVß (Clontech). The micro-dystrophin cassette had a consensus Kozak immediately in front of the ATG start and a small 53 bp synthetic polyA signal for mRNA termination. The human micro-dystrophin cassette contained the (R4–R23/Δ71–78) domains as previously described by Harper *et al.* (*Nature Medicine* **8**, 253-261 (2002)). The complementary DNA was codon optimized for human usage and synthesized by GenScript (Piscataway, NJ) (*Mol Ther* **18**, 109-117 (2010)). The only viral sequences included in this vector were the inverted terminal repeats of AAV2, which are required for both viral DNA replication and packaging. The micro-dystrophin cassette has a small 53 bp synthetic polyA signal for mRNA termination.

[00119] Previous studies have validated cardiac expression using MHCK7 promoter (Salva *et al. Mol Ther* **15**, 320-329 (2007) and AAVrh74 achieving skeletal, diaphragm, and cardiac muscle expression (Sondergaard *et al. Annals of clinical and Transl Neurology* ***2***, 256-270 (2015)), The sequence of construct of Fig. 1 was encapsidated into AAVrh.74 virions. The molecular clone of the AAVrh.74 serotype was cloned from a rhesus macaque lymph node and is described in in Rodino-Klapac *et al. Journal of Translational medicine* **5**, 45 (2007).

Table 1 shows the molecular features of the plasmid pAAV.MHCK7.micro-dystrophin (SEQ ID NO: 3)

Table 1. Molecular Features of plasmid pAAV.MHCK7.micro-dystrophin

TYPE	START	END	NAME	DESCRIPTION
REGION	7	116	5' ITR	Wild-type AAV2 inverted terminal repeat
REGION	236	1036	MHCK7	Mouse myosin heavy chain complex – E box muscle creatine kinase fusion enhancer/promoter
REGION	1046	1195	Chimeric intron	5' donor site from human β-globin gene and the branchpoint and 3' splice acceptor site from IgG heavy chain variable region
GENE	1206	4786	huDys cDNA	Human micro-dystrophin cDNA
REGION	4787	4842	PolyA	Synthetic PolyA
REGION	4933	5042	3' ITR	Wild-type AAV2 inverted terminal repeat
GENE	6808	7668	AmpR	β-lactamase gene
REGION	7823	8442	Ori	Plasmid origin of replication

Example 2

Intramuscular Expression Studies Using rAAV.MHCK7.micro-dystrophin

[00120] Expression studies were conducted with the human micro-dystrophin construct (rAAVrh74.MHCK7. micro-dystrophin; described in Example 1) by intramuscular injection. The tibialis anterior muscle of mdx mice (spontaneous Dmd^{mdx} mutant mice that do not express dystrophin) were injected with 1 x 10^{11} vg of the cassette (n=5 per group). Six weeks later the muscles were harvested and stained for dystrophin (Dys3) expression with an N-terminal antibody for dystrophin and hematoxylin and eosin (HE) staining. Figure 2 shows diffuse gene expression and reduction in centrally located nuclei with 1 x 10^{11} vg dose compared to the untreated muscle. Furthermore, a decrease in central nucleation with an increase in average fibers/frame was observed following treatment with micro-dystrophin construct. Expression levels of the rAAVrh74.MHCK7. micro-dystrophin construct were quantified at about 73%.

[00121] In addition to measuring micro-dystrophin localization and expression levels, skeletal muscle force was measured following intramuscular injection of the cassette. Intramuscular expression of pAAV.MHCK7.micro-dystrophin construct resulted in significantly greater absolute and specific force production compared with untreated controls (Figures 3A and 3B, respectfully).

Example 3

Systemic Delivery of rAAVrh.74.MHCK7.micro-dystrophin to mdx mice

[00122] Cohorts of mdx mice were injected via tail vein with either 2 x10^{12} vg (8x10^{13} vg/kg) or high dose (planned clinical dose) 6 x10^{12} vg (2x10^{14} vg/kg) of rAAVrh.74.MHCK7.micro-dystrophin at 6 weeks of age. Following 12 weeks of treatment, all muscles were harvested and stained for dystrophin and restoration of DAPC components. Systemically injected (tail vein) mice showed high levels of staining of dystrophin throughout all muscles. Figure 4A represents the widespread transduction of skeletal, diaphragm and cardiac muscle fibers after a 6x10^{12} vg (2 x 10^{14} vg/kg) systemic dose. Figure 4B shows quantification of the percentage of muscle fibers expressing micro-dystrophin in each tissue. Finally the diaphragm was tested for functional improvement (Figure 4C). No significant difference was seen at low dose; however there was significant improvement at the high dose. Importantly,

Figure 5 demonstrates other components of the DAPC were completely restored following micro-dystrophin delivery. Shown is Beta-sarcoglycan (B-SG).

[00123] The toxicology/safety of AAVrh.74.MHCK7.micro-dystrophin are were evaluated by administering the vector via intravenous (i.v.) injection to the tail vein of mdx mice per Table 2. There was no evidence of toxicity in any of the muscle tissues analyzed including: Tibialis anterior (TA), Gastrocnemius (GAS), Quadriceps (QD), Psoas (PSO), Triceps (TRI), and Diaphragm (DIA) (Figure 6A and 6B). The number of centrally placed nuclei were decreased with the high dose $6x10^{12}$ vg (2×10^{14} vg/kg). Historically, central nucleation of skeletal muscles in untreated age matched mdx mice are on average ~80%. Finally, the preliminary data from a small sample size (n=3) demonstrates a decreased level of CK release (U/L) in serum of high dose treated mice (D). Independent t-tests were used to locate differences ($p<.05$); Data are reported as means ± SEM.

Table 2. Outline of toxicology/safety study of rAAVrh.74.MHCK7.micro-dystrophin in mice.

Cohort Number	Study Agent	Dose (vg/kg)	Treatment Day 0	Follow-up Day 1	Sacrificial End-Point	
					Week 6	Extra
(1) Low Dose	AAVrh.74.MHCK7.Micro-dys	8.0 $x10^{13}$	Single *i.v.* injection to the tail vein of mdx mice	24 h Weight, Clinical Observations	5M	+2
(2) High Dose	AAVrh.74.MHCK7.Micro-dys	2.0 $x10^{14}$			5M	+2
(3) Control	Vehicle (LRS)	0			5M	+2
TOTAL MICE						**N = 21**

.Example 4

Generation of the pAAV.MCK.micro-dystrophin construct

[00124] The pAAV.MCK.micro-dystrophin plasmid was constructed by inserting the MCK expression cassette driving a codon optimized human micro-dystrophin cDNA sequence into the AAV cloning vector psub201 (Samulski *et al., J. Virol.* 61(10):3096-3101). A muscle-specific regulatory element was included in the construct to drive muscle-specific gene expression. This regulatory element

comprised the mouse MCK core enhancer (206 bp) fused to the 351 bp MCK core promoter (proximal). After the core promoter, the construct comprises the 53 bp endogenous mouse MCK Exon1 (untranslated) for efficient transcription initiation, followed by the SV40 late 16S/19S splice signals (97 bp) and a small 5'UTR (61 bp). The intron and 5' UTR was derived from plasmid pCMVß (Clontech). The micro-dystrophin cassette has a consensus Kozak immediately in front of the ATG start and a small 53 bp synthetic polyA signal for mRNA termination. The human micro-dystrophin cassette contains the (R4–R23/Δ71–78) domains as previously described by Harper *et al. Nat. Med.* 8(3):253-61, 2002

[00125] The pAAV.MCK.micro-dystrophin plasmid contained the human micro-dystrophin cDNA expression cassette flanked by AAV2 inverted terminal repeat sequences (ITR) (see Fig. 7). This sequence was encapsidated into AAVrh.74 virions. The molecular clone of the AAVrh.74 serotype was cloned from a rhesus macaque lymph node and is described in Rodino-Klapac *et al. Journal of Tran. Med.* 45 (2007).

Example 5

Potency and Dose Analysis Using rAAV.MCK.micro-dystrophin

[00126] Expression studies were conducted with the human micro-dystrophin construct (rAAV.MCK.micro-dystrophin; described in Example 1) by intramuscular injection. The tibialis anterior (TA) muscle of mdx mice (spontaneous Dmd^{mdx} mutant mice that do not express dystrophin) were injected with 3×10^9, 3×10^{10}, or 1×10^{11} vg (n=3 per group). Four weeks later the muscles were harvested and stained for dystrophin expression using an antibody specific for the N-terminal Dys3 and hematoxylin and eosin (HE) staining. Figure 8 show a linear correlation between expression and dose where very little expression (no effect level) at 3×10^9 vg and 89% expression at 1×10^{11} vg.

Example 6

Vascular Delivery of rAAV.MCK.micro-dystrophin to mdx Mice

[00127] Using an isolated limb perfusion model (Rodino-Klapac *et al., J. Trans. Med.* 5(45): 1-11, 2007), mdx mice (n=10) were injected with 1×10^{11} vg of rAAVrh.74.MCK.micro-dystrophin via the femoral artery and performed outcomes analysis was carried out. Three months post gene transfer, lower limb muscles were

harvested and efficacy studies demonstrated significant improvement in both force and resistance to eccentric contraction induced injury (Figure 9).

[00128] Dystrophin protein immunostaining in the extensor digitorum longus (EDL) muscle and TA muscle shows expression in a mdx myofibers following rAAVrh.74-MCK-micro-dystrophin treatment (Fig. 9A). Mock-infected muscle was stained in an identical manner and exposures were time matched. Fig. 9B demonstrates that rAAVrh.74-MCK-micro-dystrophin significantly increased normalized specific force relative to mock-treated mdx muscles ($P<0.05$ vs. mdx). In addition, the mdx muscles infected with rAAVrh.74-MCK-micro-dystrophin (human) were compared with mock-infected contralateral mdx EDL muscles (blue) and Wild Type (WT C57Bl/10) EDL muscles for force drop during repetitive eccentric contractions at 12 weeks post gene transfer (Fig. 9C). It was found that rAAVrh.74-MCK-microdystrophin (Micro-dys) treatment significantly protected against loss of force compared with mock-treated mdx muscles ($P< 0.001$ vs. mdx).

Example 7

Primate Studies

[00129] In order to apply pre-clinical finds in mice to a clinical paradigm, a non-human primate (NHP) was dosed systemically in order to evaluate safety and efficacy for future clinical trials. The effect of $2x10^{14}$ vg total dose of AAVrh74.MHCK7.micro-dystrophin.FLAG delivered intravenously through the cephalic vein was studied in a non-human primate. This dose was proportional (based on animal weight) to the systemic dose given to mice and corresponded to the mid-dose ($6.0x10^{12}$ vg Total Dose) given to mice.

[00130] Baseline chemistries and immunological studies including enzyme-linked immunosorbent spot assay (ELISpot) analysis were carried out to measure T cells against AAVrh.74 capsid and micro-dystrophin as well as anti-AAV antibody titers. Three peptide pools were used for the AAVrh.74 capsid protein (Genemed Synthesis, San Antonio, TX) containing 34-36 peptides, each 18 amino acids long and overlapping by 11 residues. Four peptide pools encompassing the micro-dystrophin.FLAG protein (Genemed Synthesis) each 18 amino acids long and overlapping by 11 residues. Concanavalin A (ConA) (Sigma, 1μg/mL) served as a positive control and 0.25% dimethylsulfoxide (DMSO) as a negative control. These

studies were repeated every two weeks for the entire study. At 3 months following treatment, animals were euthanized to obtain a full tissue necropsy. Immunological assays did not show any unexpected responses to the capsid or transgene by ELISpot (Figure 12A) and no unexpected antibody responses to the AAVrh74 capsid by ELISA (Figure 12B).

[00131] In addition full complete blood count and chemistry panels showed slight elevation of liver enzymes which were normalized back to baseline with no intervention or treatment necessary, as shown in Table 3 below.

Table 3

Blood Chemistry							
13-176	Baseline	24 hours	2 week	4 week	6 week	8 week	12 week
Total Protein (6.4-7 mg/dL)	7	7	6.7	6.6	6.6	6.6	6.7
Billirubin, Total (0.15-0.23 mg/dL)	0.2	0.4	0.3	0.2	0.3	0.4	0.4
ALT (31-50U/L)	39	38	75	104	182	172	65
AST (19-38 U/L)	35	63	50	92	98	121	64
Alkaline Phosphatease (504-821 U/L)	417	396	332	383	598	608	578
GGT	77	77	120	106	131	156	134
CK	109	504	164	183	137	126	123

[00132] There were no other unexpected chemistry values throughout the duration of the study. Finally, a full analysis of all skeletal muscles demonstrated widespread expression in muscle fibers through immunofluorescence staining with a FLAG specific antibody and western blot detection using a mouse monoclonal antibody to dystrophin (Figure 14A, B).

[00133] The data together demonstrates that systemic delivery of AAVrh74.MHCK7.micro-dystrophin.FLAG established safety and efficacy with widespread expression across all skeletal muscles in a non-human primate.

Example 8
Pre-Clinical Study to Demonstrate Efficacy

[00134] A pre-clinical study was carried out to demonstrate efficacy of systemic delivery of rAAVrh74.MHCK7.micro-dystrophin in treating skeletal and cardiac muscle deficits in mdx mice. The AAVrh74 vector containing a codon optimized human micro dystrophin transgene driven by a muscle and cardiac specific promoter, MHCK7 as described in Example 1 was used for this study.

[00135] Systemic injections of the rAAVrh74.MHCK7.micro-dystrophin via the tail vein in mdx (dystrophin null) mice were used for a dose response study. The results of this study demonstrated that systemic injections in mdx mice was effective in normalizing histologic and functional outcomes measured in limb and diaphragm in a dose dependent manner. Additionally, no significant vector-associated toxicity was reported following formal histopathology review by a board-certified veterinary pathologist.

[00136] The vector for this study was produced by the Nationwide Children's Hospital Viral Vector Core utilizing a triple-transfection method of HEK293 cells, under research grade conditions. Characterization of the vector following production included titer determination by qPCR with a supercoiled standard, endotoxin level determination (EU/mL) and a sterility assessment. The produced vector was analyzed by SDS-PAGE to verify banding pattern consistency with expected rAAV. The vector was produced using plasmid containing the microdystrophin construct, a muscle specific MHCK7 promoter to drive expression, a consensus Kozak sequence (CCACC), an SV40 chimeric intron, synthetic polyadenylation site (53 bp) (**Error! Reference source not found.**). The microdystrophin expression cassette was cloned between AAV2 ITRs packaged into an AAVrh74 vector for enhanced transduction of skeletal and cardiac tissue.

[00137] Potency determination of the rAAVrh74.MHCK7.micro-dystrophin test article was achieved by performing intramuscular injections of the vector into mdx mice. Wild type mice serve as a positive control and injection of sterile lactated ringers into mdx mice serve as a negative control.

Table 4: Overview of rAAVrh74.MHCK7.micro-dystrophin Study Design

Delivery Route	Animal Strain	Total Dose (vg)	# Mice	Treatment Endpoint	Analysis
IM (Potency)	mdx	1E+11	3	1 mo	IF, H&E
IV (Efficacy)	mdx	2E+12	5	3 mo	IF, H&E, Diaph Phys
IV (Efficacy)	mdx	6E+12	8	3 mo	IF, H&E, Diaph Phys, TA Phys, Path, Biodistribution, Western Blot
IV (Efficacy)	mdx	1.2E+13	8	3 mo	IF, H&E, Diaph Phys, TA Phys, Path, Biodistribution, Western Blot
IV (Efficacy)	C57BL/6	6E+12	5	3 mo	IF, H&E, Diaph Phys, TA Phys, Path
IV (Efficacy)	mdx	6E+12	5	6 mo	IF, H&E, Diaph Phys, TA Phys, Path
IV (Efficacy)	mdx	--	8	3 mo	IF, H&E, Diaph Phys, Path,
IV (Efficacy)	C57BL/6	--	6	3 mo	IF, H&E, Diaph Phys, Path,

IF: immunofluoresence; H&E: hematoxylin & eosin staining; Diaph/TA Phys: specific force measurements in the diaphragm and TA muscle; Path: formal histopathology; '--': uninjected
All injected animals were treated at 4-5 weeks of age and necropsied 3 or 6 months post-injection. Control mice were necropsied at 4 months of age and 7 months of age.

[00138] The animals indicated in Table 4 were dosed at the age indicated (4-5 weeks of age) with a tail vein injection for systemic delivery. To perform accurate dosing with an intramuscular injection, animals were briefly anesthetized by isoflurane inhalation. Doses were administered by direct injection into the tibialis anterior muscle of the lower hind limb. No anesthesia was required for accurate dosing with systemic delivery. Doses were administered by the vasculature through the tail vein. Care was taken to accurately deposit the entire vector dose into the vessel. After the dosing was performed, animals were placed on a heating pad until spontaneous movement was regained, and then returned to the cage. Observations of each animal were performed weekly for the whole duration of the study.

[00139] At the appropriate age as listed in Table 4, mice were overdosed with Ketamine/Xylazine mixture (200mg/kg/20mg/kg). Blood was collected via heart puncture and whole blood was sent for complete blood count (CBC) analysis and serum was stored at -80°C till serum chemistries were analyzed by Charles Rivers Laboratory. Tissues were then collected and sent for analysis by an independent veterinary histopathologist and in house.

[00140] Intramuscular delivery of rAAVrh74.MHCK7.micro-dystrophin to dystrophin null mice at $1x10^{11}$ vg total dose resulted in ~70% expression of dystrophin in the injected TA muscles. Immunofluorescence imaging of the vector dosed mouse confirmed expression of the micro-dystrophin gene.

Restoration of Dystrophin Expression Following Systemic Treatment with rAAVrh74.MHCK7.micro-dystrophin

[00141] Efficacy determination of the rAAVrh74.MHCK7.micro-dystrophin test article was achieved by performing systemic injections in mdx mice (*genotype: C57BL/10ScSn-Dmd^{mdx}/J*) using dose escalation at low, mid and high dose ($2.0x10^{12}$ vg Total Dose; $6.0x10^{12}$ vg Total Dose; $1.2x10^{13}$ vg Total Dose) to assess transgene expression and efficacy of the vector when delivered systemically at the time points of 3 and 6 months post injection. Mice were injected at 4-5 weeks of age and a full necropsy was performed at both 3 and 6 months post-injection. Based on mean animal weights per group, these doses equal: $8x10^{13}$ vg/kg, $2x10^{14}$ vg/kg and $6x10^{14}$ vg/kg. Injection of an equal volume of lactated ringers served as a negative control.

Injection of an equal volume of lactated ringers into C57BL/6 mice served as a positive control. Safety was determined by performing systemic injections in WT mice at a dose of 6.0 $x10^{12}$ vg Total Dose (denoted as WT TX-mid Dose group). Immunofluorescence stain of skeletal muscles tibialis anterior (TA), gastrocnemius (GAS), quadriceps (QUAD), gluteus (GLUT), psoas, tricep (TRI), diaphragm (DIA), and heart was carried out to determine restoration of dystrophin and to ensure efficacy of viral vector of rAAVrh74.MHCK7.micro-dystrophin.

[00142] The skeletal muscles (TA, QUAD, GLUT, TRI) were extracted, along with the heart and diaphragm, for analysis. Organs were also removed for toxicology and biodistribution studies. Micro-dystrophin transgene expression remained high following 3-6 months treatment. This was accompanied by improved muscle histopathology and improved function with no adverse effects in off target organs.

Reversal of Dystrophic Phenotype in rAAVrh74.MHCK7.micro-dystrophin Systemically Treated mdx Mice

[00143] Hematoxylin & Eosin (H&E) stain of skeletal muscles, diaphragm and heart was carried out to determine reversal and improvement of dystrophic pathology following systemic injection of rAAVrh74.MHCK7.micro-dystrophin at $2x10^{12}$ vg total dose (Low Dose; n=1), $6x10^{12}$ vg total dose (Mid Dose; n=8), and $1.2x10^{13}$ vg total dose (High Dose; n=8) for each dose with euthanasia 12 weeks post-injection. At 24 weeks post-injection, a second cohort of animals treated with mid dose ($6x10^{12}$ vg total dose) were evaluated for reversal and improvement of dystrophin pathology (n=5).

[00144] Immunofluorescence staining for the human micro-dystrophin protein was used to determine micro-dystrophin transgene expression in both left and right sides of six skeletal muscles (TA, GAS, QUAD, GLUT, psoas, TRI), as well as the diaphragm and the heart in all dystrophin null mice injected with the micro-dystrophin vector. This was carried out to determine restoration of dystrophin and to ensure efficacy of viral vector of rAAVrh74.MHCK7.micro-dystrophin at $2x10^{12}$ vg total dose (Low Dose; n=2), $6x10^{12}$ vg total dose (Mid Dose; n=8), and $1.2x10^{13}$ vg total dose (High Dose; n=8) for each dose with euthanasia 12 weeks post-injection.

[00145] In order to evaluate expression and transduction efficiency, images from all three dosing cohorts and both left and right sides of each muscle were utilized for

quantification. Four 20X images were taken of each muscle and the percent of micro-dystrophin positive fibers was determined for each image resulting in the average percent transduction for each muscle. Figures 14 and 15 present representative images from treated mice from the mid dose ($6x10^{12}$ vg; $2x10^{14}$vg/kg) and the high dose ($1.2x10^{13}$ vg; $6x10^{14}$vg/kg). Dystrophin null mice that were injected with lactated ringers and age matched were included for negative control and wild-type mice injected with lactated ringers were included for positive controls. The heart demonstrated ≥75% in all animals analyzed.

[00146] The muscles from untreated animals exhibited widespread myopathy including fatty infiltration, central nucleation, fibrosis and focal areas of necrosis. H&E staining in **Error! Reference source not found.**16 and **Error! Reference source not found.**17 illustrates this dystrophic phenotype in dystrophin null mice when compared to normal WT mice and the improvement of muscle pathology following treatment at either the mid dose ($6x10^{12}$ vg; $2x10^{14}$ vg/kg) or the high dose ($1.2x10^{13}$ vg; $6x10^{14}$ vg/kg). Quantification of histological parameters showed a reduction in central nucleation (Figure 18) and a normalization of average fiber diameters (**Error! Reference source not found.**16 and 17) in treated mice in all muscles in a dose dependent manner. Sirius Red staining demonstrated a reduction in collagen deposition in the diaphragm in both the mid and high dose cohorts compared to untreated (mdx LR) cohorts (Figure 19).

Functional Assessment of Systemic Treatment with rAAVrh74.MHCK7.micro-dystrophin

[00147] To determine whether micro-dystrophin gene transfer provided a functional strength benefit to diseased muscle, the functional properties of both the diaphragm and the tibialis anterior from mdx mice, WT mice, and vector dosed mice at three dose levels were assessed. The dose escalation included low dose ($8x10^{13}$ vg/kg), mid dose ($2x10^{14}$ vg/kg), and high dose ($6x10^{14}$ vg/kg). Functional assessment of systemic treatment with rAAVrh74.MHCK7.micro-dystrophin using ex-vivo assessment of specific force and decrease in force output following eccentric contractions in the TA was utilized 24 weeks post-injection in animals systemically injected with rAAVrh74.MHCK7.micro-dystrophin at $6x10^{12}$ vg total dose (Mid Dose). Additionally, specific force output in the diaphragm was assessed in the same animals.

[00148] As outlined in the previous figures, histopathology exhibited a more normalized environment with improvements in central nucleation, collagen deposition, and fiber size in the mid and high doses. Tail vein delivery of rAAVrh74.MHCK7.micro-dystrophin led to a stepwise improvement in specific force output in the diaphragm (176.9 mN/mm^2 in the mid dose group versus 227.78 mN/mm^2 in the high dose group). Additionally, the long-term treated cohort represents mice 6 months post injection (mid dose $2x10^{14}$ vg/kg) and there was no deviation in diaphragm force output long-term (176.9 mN/mm^2 vs 194.9 mN/mm^2) (**Error! Reference source not found.**20).

[00149] Furthermore, functional deficits in tibialis anterior muscle in mdx mice were observed compared to WT mice. Mdx mice demonstrated 50% decrease in force output compared to WT mice (171.3 mN/mm^2 vs. 291.65 mN/mm^2) and greater loss of force following eccentric contractions (32% loss in mdx; 5% loss in WT). Systemic delivery of the mid dose level of rAAVrh74.MHCK7.micro-dystrophin resulted in 65.5% dystrophin in the tibialis anterior muscle and restoration of specific force output which improved to 235.4 mN/mm2 and protected the muscle from repeated eccentric contraction damage with only a 25% decrease in force (Figure 21). The WT Mid Dose group represents a wild-type treated cohort in order to demonstrate absence of toxicity and maintenance of functional outcome measures after vector treatment.

Summary

[00150] After the initial demonstration of biopotency by intramuscular injection, comparable or increased restoration of micro-dystrophin was achieved with vascular delivery while transducing skeletal muscles, diaphragm and the heart. The efficacy demonstrated reversal of dystrophic features in a dose dependent manner by reduction of inflammation, fewer degenerating fibers, and improved functional recovery by protecting against eccentric contractions in the tibialis anterior and diaphragm. The functional benefits of the vector include a stepwise improvement to wild-type levels in force generation of the diaphragm and the TA.

Example 9
Toxicology and Biodistribution of Systemic Treatment with rAAVrh74.MHCK7.micro-dystrophin

[00151] Organs and tissues from mdx mice given systemic injection of rAAVrh74.MHCK7.micro-dystrophin were collected for real-time quantitative PCR to detect specific sequences of vector DNA. Protein extracted from all collected organs and tissues were run on Western blot to detect micro-dystrophin in off-target organs.

[00152] Test article was given at three dose levels: low (2×10^{12} vg; 8×10^{13} vg/kg), mid (6×10^{12} vg; 8×10^{14} vg/kg) and high dose (1.2×10^{13} vg; 6×10^{14} vg/kg) by intravenous route at 4-5 weeks of age. To assess the safety of the vector, H&E staining was performed on cryosections of muscle tissue and all major organs harvested from the same cohorts of mice previously described. Also included were organs and muscles from C57BL6 WT mice treated systemically with the vector at the mid dose. Lactated ringers treated mdx and WT mice were also included for histopathology analysis. These sections were formally reviewed for toxicity by a third-party, board-certified, veterinary pathologist and no adverse effects were detected in any sample from any of the mice; results are summarized below.

[00153] Group details and study design are shown in 4 below.

Table 5: rAAVrh74.MHCK7.micro-dystrophin Safety Study Design

Delivery Route	Animal Strain	Total Dose (vg)	No. of Mice	Treatment Endpoint	Pathology Report Number
IV	mdx	$2x10^{12}$	5	3 mo	AAVrh74-mdx-MOUSE-001.1
IV	mdx	$6x10^{12}$	7	3 mo	AAVrh74-mdx-MOUSE-001.1/001.2
IV	mdx	$1.2x10^{13}$	8	3 mo	AAVrh74-mdx-MOUSE-001.2
IV	C57BL/6	$6x10^{12}$	5	3 mo	AAVrh74-mdx-MOUSE-001.2
IV	mdx	$6x10^{12}$	5	6 mo	AAVrh74-mdx-MOUSE-001.3
IV	mdx	--	8	3 mo	AAVrh74-mdx-MOUSE-001.2
IV	C57BL/6	--	6	3 mo	AAVrh74-mdx-MOUSE-001.2

Histopathological Review of Vector Transduced Tissue

[00154] IV injection of rAAVrh74.MHCK7.micro-dystrophin did not elicit any microscopic changes in myofibers of any skeletal muscles examined. In addition, no treatment-related lesions were seen in any of the tissues evaluated histologically. Any changes noted were seen in both treated and control mice and were considered incidental findings. Taken together, these data indicate that this test article was well tolerated by the test subjects. Furthermore, relative to reference specimens from age-matched, untreated mdx mice, administration of rAAVrh74.MHCK7.micro-dystrophin decreased myofiber atrophy in treated mdx mice, thus showing that the test article can ameliorate the degree of myopathy associated with deficiencies of mdx.

[00155] In addition to review of diseased mdx mice systemically treated with vector, rAAVrh74.MHCK7.micro-dystrophinwas delivered systemically to five C57BL/6 WT mice at a dose identical to the minimally efficacious dose (MED) established in the studies above in mdx mice, $6x10^{12}$ vg total dose ($2x10^{14}$ vg/kg). This allowed for the study of intravenous delivery of the test article in healthy WT mice to determine more definitively if any adverse effects result solely due to the treatment. Here again a variety of skeletal muscles including the diaphragm, along with heart, and five other organs were harvested and H&E sections of each tissue were formally reviewed by an independent veterinary pathologist.

Vector Genome Biodistribution

[00156] The presence of test article-specific DNA sequences was examined using a real time, quantitative PCR assay (qPCR). Biodistribution analysis was performed on tissue samples collected from three vector dosed mdx animals per dose level. A positive signal was anything equal to or greater than 100 single-stranded DNA copies/µg genomic DNA detected. Tissues were harvested at necropsy and vector specific primer probe sets specific for sequences of the MHCK7 promoter were utilized. Figure 22 and Table 6 below depicts the vector genome copies detected in each tissue sample from rAAVrh74.MHCK7.micro-dystrophin injected mice.

Table 6: Vector Genome copy numbers in organs and muscles from three vector dosed mdx mice per dose level. Values are shown in vg/µg genomic DNA.

Tissue	2.00E+12 (average vg copies/ug)	6.00E+12 (average vg copies/ug)	1.20E+13 (average vg copies/ug)
Hrt	2.84E+04	7.65E+05	5.35E+06
Lng	3.14E+04	2.52E+05	1.49E+06
Liv	4.36E+04	1.11E+07	1.80E+07
Kid	1.96E+04	3.27E+05	1.06E+06
Spl	5.69E+04	5.27E+05	5.78E+05
Gon	5.74E+04	3.68E+04	3.50E+05
Dia	2.22E+04	3.55E+05	2.32E+06
Pso	1.28E+05	1.60E+05	1.57E+06
Tri	1.60E+05	5.45E+05	2.50E+06
Qd	2.66E+06	6.57E+05	2.29E+06
Gas	1.69E+05	5.80E+05	2.93E+06
TA	5.86E+05	1.25E+05	1.32E+06

[00157] rAAVrh74.MHCK7.micro-dystrophin transcript was detected at varying levels in all collected tissues. As expected, the highest levels were seen in skeletal muscle and the heart. The lowest levels were detected in gonad, lung, kidney, and spleen. These data indicate that the test article was efficiently delivered into all investigated tissues of vector dosed mice.

[00158] As the qPCR results above indicate, intravenous delivery of rAAVrh74.MHCK7.micro-dystrophin resulted in distribution of vector transcript to varying levels in most tissues, with the highest levels occurring in the liver, heart, and quadriceps muscle (mid dose) and the liver, heart and gastrocnemius muscle (high dose). Therefore, the objective of this portion of the study was to determine the protein expression of the human micro-dystrophin transgene in these tissues to ensure the functionality of the muscle specific MHCK7 promoter. Western blotting was used to detect micro-dystrophin expression in the tissue samples.

[00159] Protein expression and vector biodistribution were also assessed using qPCR and western blotting (Figure 23), and these data indicate normal levels of vector in off-site organs and minimal detection of micro-dystrophin protein in the high dose treated livers. These results were correlated with no toxicity as determined by the pathologist in the liver. Additionally, serum chemistries were analyzed by an

independent CRO (Charles River Laboratories) which indicate normal values across all chemistries analyzed. There were three abnormal values in the liver enzyme AST, 2 of which were demonstrated in the mdx-LR group and 1 of which in the mid-dose group (Figure 23). A subset of animals underwent creatine kinase analysis (CK), however, samples were analyzed pre and post physiology evaluation. Analysis of serum corroborates the lack of toxicity after test article delivery.

[00160] Micro-dystrophin protein expression was observed in varying amounts in all skeletal muscle samples as well as heart samples (Figure 24). However, there was minimal protein detected in the high dosed cohorts in the liver. This is believed to be a benign result and it might be that the presence in the liver is due to expression in smooth muscle of the liver. Importantly, there were no adverse histopathologic effects denoted by the independent pathologist report in the liver.

Summary

[00161] Histopathology review concluded that the mdx-LR cohort exhibited widespread myopathy affecting all seven skeletal muscles evaluated as well as the right ventricular wall of the heart. The principal findings of the histopathology review included pronounced and widespread myofiber atrophy (30-75% of normal myofiber size), minimal to mild mononuclear cell inflammations, increased interstitial space, and increased cytoplasmic mineral deposits. The diaphragm exhibited the most marked changes in mononuclear cell infiltration and myofiber atrophy. The heart exhibited a few small foci of minimal mononuclear cell accumulation in the ventricular myocardium. Vector dosed cohorts had substantially reduced myopathy in all skeletal tissues and the heart. The reductions in histopathologic findings were in a dose dependent manner with the high dose group having substantially less degeneration and inflammation. There were no adverse effects due to vector treatment with rAAVrh74.MHCK7.micro-dystrophin as was documented in the WT treated cohort and the vector dosed mdx cohorts. There were incidental findings in the liver and lung of mdx and WT mice, regardless of treatment, in which the mice exhibited mild vacuolation of hepatocyte cytoplasm. Therefore, the test article was safe, efficacious and the protective effect was dose-dependent.

REFERENCES

1. Hoffman, E.P., Brown, R.H., Jr. & Kunkel, L.M. Dystrophin: the protein product of the Duchenne muscular dystrophy locus. *Cell* **51**, 919-928 (1987).
2. Straub, V. & Campbell, K.P. Muscular dystrophies and the dystrophin-glycoprotein complex. *Curr Opin Neurol* **10**, 168-175 (1997).
3. Sacco, A., *et al.* Short telomeres and stem cell exhaustion model Duchenne muscular dystrophy in mdx/mTR mice. *Cell* **143**, 1059-1071 (2010).
4. Wallace, G.Q. & McNally, E.M. Mechanisms of muscle degeneration, regeneration, and repair in the muscular dystrophies. *Annu Rev Physiol* **71**, 37-57 (2009).
5. Zhou, L. & Lu, H. Targeting fibrosis in Duchenne muscular dystrophy. *J Neuropathol Exp Neurol* **69**, 771-776 (2010).
6. Desguerre, I., *et al.* Endomysial fibrosis in Duchenne muscular dystrophy: a marker of poor outcome associated with macrophage alternative activation. *J Neuropathol Exp Neurol* **68**, 762-773 (2009).
7. DiPrimio, N., McPhee, S.W. & Samulski, R.J. Adeno-associated virus for the treatment of muscle diseases: toward clinical trials. *Curr Opin Mol Ther* **12**, 553-560 (2010).
8. Mendell, J.R., *et al.* Sustained alpha-sarcoglycan gene expression after gene transfer in limb-girdle muscular dystrophy, type 2D. *Ann Neurol* **68**, 629-638 (2010).
9. Mendell, J.R., *et al.* Limb-girdle muscular dystrophy type 2D gene therapy restores alpha-sarcoglycan and associated proteins. *Ann Neurol* **66**, 290-297 (2009).
10. Mendell, J.R., *et al.* A phase 1/2a follistatin gene therapy trial for becker muscular dystrophy. *Molecular therapy : the journal of the American Society of Gene Therapy* **23**, 192-201 (2015).
11. Carnwath, J.W. & Shotton, D.M. Muscular dystrophy in the mdx mouse: histopathology of the soleus and extensor digitorum longus muscles. *J Neurol Sci* **80**, 39-54 (1987).

12. Coulton, G.R., Morgan, J.E., Partridge, T.A. & Sloper, J.C. The mdx mouse skeletal muscle myopathy: I. A histological, morphometric and biochemical investigation. *Neuropathol Appl Neurobiol* **14**, 53-70 (1988).

13. Cullen, M.J. & Jaros, E. Ultrastructure of the skeletal muscle in the X chromosome-linked dystrophic (mdx) mouse. Comparison with Duchenne muscular dystrophy. *Acta Neuropathol* **77**, 69-81 (1988).

14. Dupont-Versteegden, E.E. & McCarter, R.J. Differential expression of muscular dystrophy in diaphragm versus hindlimb muscles of mdx mice. *Muscle Nerve* **15**, 1105-1110 (1992).

15. Stedman, H.H., *et al.* The mdx mouse diaphragm reproduces the degenerative changes of Duchenne muscular dystrophy. *Nature* **352**, 536-539 (1991).

16. Deconinck, A.E., *et al.* Utrophin-dystrophin-deficient mice as a model for Duchenne muscular dystrophy. *Cell* **90**, 717-727 (1997).

17. Grady, R.M., *et al.* Skeletal and cardiac myopathies in mice lacking utrophin and dystrophin: a model for Duchenne muscular dystrophy. *Cell* **90**, 729-738 (1997).

18. Love, D.R., *et al.* An autosomal transcript in skeletal muscle with homology to dystrophin. *Nature* **339**, 55-58 (1989).

19. Tinsley, J.M., *et al.* Primary structure of dystrophin-related protein. *Nature* **360**, 591-593 (1992).

20. Tinsley, J., *et al.* Expression of full-length utrophin prevents muscular dystrophy in mdx mice. *Nat Med* **4**, 1441-1444 (1998).

21. Squire, S., *et al.* Prevention of pathology in mdx mice by expression of utrophin: analysis using an inducible transgenic expression system. *Hum Mol Genet* **11**, 3333-3344 (2002).

22. Rafael, J.A., Tinsley, J.M., Potter, A.C., Deconinck, A.E. & Davies, K.E. Skeletal muscle-specific expression of a utrophin transgene rescues utrophin-dystrophin deficient mice. *Nat Genet* **19**, 79-82 (1998).

23. Zhou, L., *et al.* Haploinsufficiency of utrophin gene worsens skeletal muscle inflammation and fibrosis in mdx mice. *J Neurol Sci* **264**, 106-111 (2008).

24. Gutpell, K.M., Hrinivich, W.T. & Hoffman, L.M. Skeletal Muscle Fibrosis in the mdx/utrn+/- Mouse Validates Its Suitability as a Murine Model of Duchenne Muscular Dystrophy. *PloS one* **10**, e0117306 (2015).

25. Rodino-Klapac, L.R., *et al.* Micro-dystrophin and follistatin co-delivery restores muscle function in aged DMD model. *Human molecular genetics* **22**, 4929-4937 (2013).

26. Nevo, Y., *et al.* The Ras antagonist, farnesylthiosalicylic acid (FTS), decreases fibrosis and improves muscle strength in dy/dy mouse model of muscular dystrophy. *PloS one* **6**, e18049 (2011).

27. Rodino-Klapac, L.R., *et al.* A translational approach for limb vascular delivery of the micro-dystrophin gene without high volume or high pressure for treatment of Duchenne muscular dystrophy. *J Transl Med* **5**, 45 (2007).

28. Mulieri, L.A., Hasenfuss, G., Ittleman, F., Blanchard, E.M. & Alpert, N.R. Protection of human left ventricular myocardium from cutting injury with 2,3-butanedione monoxime. *Circ Res* **65**, 1441-1449 (1989).

29. Rodino-Klapac, L.R., *et al.* Persistent expression of FLAG-tagged micro dystrophin in nonhuman primates following intramuscular and vascular delivery. *Molecular therapy : the journal of the American Society of Gene Therapy* **18**, 109-117 (2010).

30. Grose, W.E., *et al.* Homologous recombination mediates functional recovery of dysferlin deficiency following AAV5 gene transfer. *PloS one* **7**, e39233 (2012).

31. Liu, M., *et al.* Adeno-associated virus-mediated microdystrophin expression protects young mdx muscle from contraction-induced injury. *Mol Ther* **11**, 245-256 (2005).

32. Harper, S.Q., *et al.* Modular flexibility of dystrophin: implications for gene therapy of Duchenne muscular dystrophy. *Nature medicine* **8**, 253-261 (2002).

33. Rodino-Klapac, L.R., *et al.* Persistent expression of FLAG-tagged micro dystrophin in nonhuman primates following intramuscular and vascular delivery. *Mol Ther* **18**, 109-117 (2010).

34. Salva, M.Z., *et al.* Design of tissue-specific regulatory cassettes for high-level rAAV-mediated expression in skeletal and cardiac muscle. *Mol Ther* **15**, 320-329 (2007).

35. Sondergaard, P.C., *et al.* AAV.Dysferlin Overlap Vectors Restore Function in Dysferlinopathy Animal Models. *Annals of clinical and translational neurology* **2**, 256-270 (2015).

36. De, B.P., *et al.* High levels of persistent expression of alpha1-antitrypsin mediated by the nonhuman primate serotype rh.10 adeno-associated virus despite preexisting immunity to common human adeno-associated viruses. *Mol Ther* **13**, 67-76 (2006).

37. Rodino-Klapac, L.R., *et al.* A translational approach for limb vascular delivery of the micro-dystrophin gene without high volume or high pressure for treatment of Duchenne muscular dystrophy. *Journal of translational medicine* **5**, 45 (2007).

38. Bulfield *et al.,* X chromosome-linked muscular dystrophy (mdx) in the mouse. *Proc Natl Acad Sci U S A.* 1984; **81**(4): 1189–1192.

39. Sicinski *et al.,* The molecular basis of muscular dystrophy in the mdx mouse: a point mutation. *Science.* 1989 **30**;244(4912):1578-80

SEQUENCE LISTING

<110> RESEARCH INSTITUTE AT NATIONWIDE CHILDREN'S HOSPITAL

<120> ADENO-ASSOCIATED VIRUS VECTOR DELIVERY OF MUSCLE SPECIFIC MICRO-DYSTROPHIN TO TREAT MUSCULAR DYSTROPHY

<130> 28335/51475A PCT

<150> US 62/473,148
<151> 2017-03-17

<160> 5

<170> PatentIn version 3.5

<210> 1
<211> 3579
<212> DNA
<213> Homo sapiens

<400> 1
atgctgtggt gggaggaggt ggaggattgt tatgaaaggg aggacgtgca gaagaagact 60
tttaccaagt gggtgaacgc tcagttcagc aaatttggga agcagcacat cgagaatctg 120
ttttccgacc tgcaggatgg gagacggctg ctggatctgc tggaaggact gactggccag 180
aagctgccca aagagaaggg gagcactagg gtgcacgccc tgaacaacgt gaacaaagct 240
ctgagagtgc tgcagaacaa caacgtggat ctggtgaata ttggcagtac tgatatcgtg 300
gacgggaacc acaaactgac actgggcctg atctggaaca ttattctgca ctggcaggtg 360
aaaaatgtga tgaagaacat catggccggg ctgcagcaga ccaattccga gaagatcctg 420
ctgtcttggg tgcggcagag cacccgcaac tatccccagg tgaacgtgat taacttcact 480
acatcctgga gcgacgggct ggccctgaat gctctgattc acagccacag gcctgatctg 540
ttcgactgga atagcgtggt gtgccagcag tctgccacac agcgcctgga acatgccttc 600
aatatcgctc ggtaccagct ggggatcgaa aaactgctgg acccagagga tgtggacact 660
acatacccag ataaaaagtc tattctgatg tacattacta gcctgttcca ggtgctgcca 720
cagcaggtgt ctattgaagc cattcaggag gtggaaatgc tgccccgccc ccccaaagtg 780
actaaagagg agcattttca gctgcatcat cagatgcatt acagccagca gattaccgtg 840
agcctggctc agggatatga gcgcaccagt agtccaaaac cacggttcaa gtcctacgct 900
tatacccagg ctgcctacgt gacaactagc gaccctacta gatccccctt tccatcccag 960
cacctggagg ccccagagga caagagcttt gggtccagcc tgatggaaag cgaggtgaat 1020
ctggatcggt accagacagc cctggaggag gtgctgagct ggctgctgag tgctgaagac 1080
acactgcagg cccagggcga aatttccaat gacgtggaag tggtgaagga tcagttccac 1140
acacacgagg gctatatgat ggacctgaca gctcaccagg ggcgcgtggg caatatcctg 1200
cagctgggct ctaaactgat cggcaccggg aaactgagtg aggacgagga aacagaagtg 1260
caggagcaga tgaacctgct gaacagccgc tgggagtgtc tgagagtggc tagtatggag 1320
aagcagtcca acctgcaccg ggtgctgatg gacctgcaga accagaaact gaaagagctg 1380
aacgactggc tgacaaagac tgaggaacgc acaaggaaga tggaggagga gccactggga 1440

cccgacctgg aggatctgaa gagacaggtg cagcagcata aggtgctgca ggaggatctg 1500
gaacaggagc aggtgcgggt gaactccctg acacatatgg tggtggtggt ggacgaatct 1560
agtggagatc acgccaccgc cgccctggag gaacagctga aggtgctggg ggaccggtgg 1620
gccaacattt gccggtggac cgaggacagg tgggtgctgc tgcaggacat cctgctgaaa 1680
tggcagaggc tgaccgagga gcagtgtctg tttagtgctt ggctgagcga gaaagaggac 1740
gccgtgaaca agatccacac aaccggcttt aaggatcaga acgaaatgct gtctagcctg 1800
cagaaactgg ctgtgctgaa ggccgatctg gagaaaaaga agcagagcat gggcaaactg 1860
tatagcctga aacaggacct gctgagcacc ctgaagaaca agagcgtgac ccagaagaca 1920
gaagcctggc tggataactt tgcccgctgc tgggacaacc tggtgcagaa actggagaaa 1980
agtacagctc agatctctca ggctgtgacc acaacccagc ctagcctgac ccagacaacc 2040
gtgatggaaa ccgtgaccac cgtgacaacc cgcgaacaga tcctggtgaa acatgcccag 2100
gaagagctgc cacctccacc tccccagaag aagagaaccc tggagcggct gcaggagctg 2160
caggaagcca ctgacgaact ggacctgaag ctgaggcagg ccgaagtgat taaggggtct 2220
tggcagcctg tgggcgatct gctgattgat tccctgcagg accacctgga aaaggtgaag 2280
gctctgagag gcgaaattgc tccactgaag gagaacgtga gtcatgtgaa cgatctggct 2340
agacagctga caacactggg catccagctg agcccataca atctgagcac actggaggac 2400
ctgaatacca ggtggaagct gctgcaggtg gctgtggaag accgggtgcg gcagctgcat 2460
gaggcccatc gcgacttcgg accagccagc cagcactttc tgagcacatc cgtgcagggg 2520
ccctgggaga gggccatttc tcccaacaag gtgccctact atattaatca cgagacccag 2580
accacttgtt gggaccatcc caagatgaca gaactgtacc agtccctggc cgatctgaac 2640
aacgtgaggt ttagcgctta cagaaccgct atgaagctga gacggctgca gaaggccctg 2700
tgcctggatc tgctgtccct gtccgccgcc tgcgatgccc tggatcagca taatctgaag 2760
cagaacgatc agccaatgga tatcctgcag atcatcaact gcctgaccac tatctacgac 2820
aggctggagc aggagcacaa caacctggtg aacgtgcctc tgtgcgtgga tatgtgcctg 2880
aactggctgc tgaacgtgta tgacactggg cgcaccggcc ggatcagagt gctgagtttt 2940
aaaactggga ttatctccct gtgtaaggcc cacctggagg acaagtacag gtacctgttc 3000
aagcaggtgg ctagtagcac tggattttgt gaccagcgcc gcctgggact gctgctgcat 3060
gatagtatcc agattcctag acagctggga gaggtggcta gtttcggagg atctaacatc 3120
gaacccagcg tgcgcagctg tttccagttt gccaataaca aacctgaaat cgaggctgct 3180
ctgttcctgg attggatgcg cctggaacca cagagcatgg tgtggctgcc tgtgctgcac 3240
agagtggctg ccgccgaaac tgccaagcac caggctaaat gcaacatctg caaggaatgt 3300
cccattatcg gctttcgcta caggagtctg aaacatttta actacgatat ttgccagagc 3360
tgcttctttt ccggaagagt ggccaaagga cacaagatgc actaccctat ggtggaatat 3420
tgcaccccaa ctacatctgg cgaagatgtg cgcgattttg ccaaggtgct gaagaataag 3480
tttcggacta agaggtactt cgccaagcac ccccgcatgg ggtatctgcc agtgcagaca 3540
gtgctggaag gagacaatat ggagaccgat acaatgtga 3579

<210> 2
<211> 810
<212> DNA
<213> Adeno-associated virus

<400> 2
gtttaaacaa gcttgcatgt ctaagctaga cccttcagat taaaaataac tgaggtaagg 60
gcctgggtag gggaggtggt gtgagacgct cctgtctctc ctctatctgc ccatcggccc 120

tttggggagg aggaatgtgc ccaaggacta aaaaaaggcc atggagccag aggggcgagg 180
gcaacagacc tttcatgggc aaaccttggg gccctgctgt ctagcatgcc ccactacggg 240
tctaggctgc ccatgtaagg aggcaaggcc tggggacacc cgagatgcct ggttataatt 300
aacccagaca tgtggctgcc ccccccccc caacacctgc tgcctctaaa aataaccctg 360
tccctggtgg atcccctgca tgcgaagatc ttcgaacaag gctgtggggg actgagggca 420
ggctgtaaca ggcttggggg ccagggctta tacgtgcctg ggactcccaa agtattactg 480
ttccatgttc ccggcgaagg gccagctgtc ccccgccagc tagactcagc acttagttta 540
ggaaccagtg agcaagtcag cccttggggc agcccataca aggccatggg gctgggcaag 600
ctgcacgcct gggtccgggg tgggcacggt gcccgggcaa cgagctgaaa gctcatctgc 660
tctcaggggc ccctccctgg ggacagcccc tcctggctag tcacaccctg taggctcctc 720
tatataaccc aggggcacag gggctgccct cattctacca ccacctccac agcacagaca 780

gacactcagg agccagccag cggcgcgccc 810

<210> 3
<211> 8611
<212> DNA
<213> Adeno-associated virus

<400> 3
gcccaatacg caaaccgcct ctccccgcgc gttggccgat tcattaatgc agctggcgcg 60
ctcgctcgct cactgaggcc gcccgggcaa agcccgggcg tcgggcgacc tttggtcgcc 120
cggcctcagt gagcgagcga gcgcgcagag agggagtggc caactccatc actaggggtt 180
ccttgtagtt aatgattaac ccgccatgct aattatctac gtagccatgt ctagagttta 240
aacaagcttg catgtctaag ctagaccctt cagattaaaa ataactgagg taagggcctg 300
ggtaggggag gtggtgtgag acgctcctgt ctctcctcta tctgcccatc ggccctttgg 360
ggaggaggaa tgtgcccaag gactaaaaaa aggccatgga gccagagggg cgagggcaac 420
agacctttca tgggcaaacc ttggggccct gctgtctagc atgccccact acgggtctag 480
gctgcccatg taaggaggca aggcctgggg acacccgaga tgcctggtta taattaaccc 540
agacatgtgg ctgccccccc ccccccaaca cctgctgcct ctaaaaataa ccctgtccct 600
ggtggatccc ctgcatgcga agatcttcga acaaggctgt gggggactga gggcaggctg 660
taacaggctt gggggccagg gcttatacgt gcctgggact cccaaagtat tactgttcca 720
tgttcccggc gaagggccag ctgtcccccg ccagctagac tcagcactta gtttaggaac 780
cagtgagcaa gtcagccctt ggggcagccc atacaaggcc atggggctgg gcaagctgca 840
cgcctgggtc cggggtgggc acggtgcccg ggcaacgagc tgaaagctca tctgctctca 900
ggggcccctc cctggggaca gcccctcctg gctagtcaca ccctgtaggc tcctctatat 960
aacccagggg cacaggggct gccctcattc taccaccacc tccacagcac agacagacac 1020
tcaggagcca gccagcggcg cgcccaggta agtttagtct ttttgtcttt tatttcaggt 1080
cccggatccg gtggtggtgc aaatcaaaga actgctcctc agtggatgtt gcctttactt 1140
ctaggcctgt acggaagtgt tacttctgct ctaaaagctg cggaattgta cccgcggccg 1200
ccaccatgct gtggtgggag gaggtggagg attgttatga aagggaggac gtgcagaaga 1260
agacttttac caagtgggtg aacgctcagt tcagcaaatt tgggaagcag cacatcgaga 1320
atctgttttc cgacctgcag gatgggagac ggctgctgga tctgctggaa ggactgactg 1380
gccagaagct gcccaaagag aaggggagca ctagggtgca cgccctgaac aacgtgaaca 1440
aagctctgag agtgctgcag aacaacaacg tggatctggt gaatattggc agtactgata 1500

tcgtggacgg gaaccacaaa ctgacactgg gcctgatctg gaacattatt ctgcactggc 1560
aggtgaaaaa tgtgatgaag aacatcatgg ccgggctgca gcagaccaat tccgagaaga 1620
tcctgctgtc ttgggtgcgg cagagcaccc gcaactatcc ccaggtgaac gtgattaact 1680
tcactacatc ctggagcgac gggctggccc tgaatgctct gattcacagc cacaggcctg 1740
atctgttcga ctggaatagc gtggtgtgcc agcagtctgc cacacagcgc ctggaacatg 1800
ccttcaatat cgctcggtac cagctgggga tcgaaaaact gctggaccca gaggatgtgg 1860
acactacata cccagataaa aagtctattc tgatgtacat tactagcctg ttccaggtgc 1920
tgccacagca ggtgtctatt gaagccattc aggaggtgga aatgctgccc cgccccccca 1980
aagtgactaa agaggagcat tttcagctgc atcatcagat gcattacagc cagcagatta 2040
ccgtgagcct ggctcaggga tatgagcgca ccagtagtcc aaaaccacgg ttcaagtcct 2100
acgcttatac ccaggctgcc tacgtgacaa ctagcgaccc tactagatcc ccctttccat 2160
cccagcacct ggaggcccca gaggacaaga gctttgggtc cagcctgatg gaaagcgagg 2220
tgaatctgga tcggtaccag acagccctgg aggaggtgct gagctggctg ctgagtgctg 2280
aagacacact gcaggcccag ggcgaaattt ccaatgacgt ggaagtggtg aaggatcagt 2340
tccacacaca cgagggctat atgatggacc tgacagctca ccaggggcgc gtgggcaata 2400
tcctgcagct gggctctaaa ctgatcggca ccgggaaact gagtgaggac gaggaaacag 2460
aagtgcagga gcagatgaac ctgctgaaca gccgctggga gtgtctgaga gtggctagta 2520
tggagaagca gtccaacctg caccgggtgc tgatggacct gcagaaccag aaactgaaag 2580
agctgaacga ctggctgaca aagactgagg aacgcacaag gaagatggag gaggagccac 2640
tgggacccga cctggaggat ctgaagagac aggtgcagca gcataaggtg ctgcaggagg 2700
atctggaaca ggagcaggtg cgggtgaact ccctgacaca tatggtggtg gtggtggacg 2760
aatctagtgg agatcacgcc accgccgccc tggaggaaca gctgaaggtg ctggggacc 2820
ggtgggccaa catttgccgg tggaccgagg acaggtgggt gctgctgcag gacatcctgc 2880
tgaaatggca gaggctgacc gaggagcagt gtctgtttag tgcttggctg agcgagaaag 2940
aggacgccgt gaacaagatc cacacaaccg gctttaagga tcagaacgaa atgctgtcta 3000
gcctgcagaa actggctgtg ctgaaggccg atctggagaa aaagaagcag agcatgggca 3060
aactgtatag cctgaaacag gacctgctga gcaccctgaa gaacaagagc gtgacccaga 3120
agacagaagc ctggctggat aactttgccc gctgctggga caacctggtg cagaaactgg 3180
agaaaagtac agctcagatc tctcaggctg tgaccacaac ccagcctagc ctgacccaga 3240
caaccgtgat ggaaaccgtg accaccgtga caacccgcga acagatcctg gtgaaacatg 3300
cccaggaaga gctgccacct ccacctcccc agaagaagag aaccctggag cggctgcagg 3360
agctgcagga agccactgac gaactggacc tgaagctgag gcaggccgaa gtgattaagg 3420
ggtcttggca gcctgtgggc gatctgctga ttgattccct gcaggaccac ctggaaaagg 3480
tgaaggctct gagaggcgaa attgctccac tgaaggagaa cgtgagtcat gtgaacgatc 3540
tggctagaca gctgacaaca ctgggcatcc agctgagccc atacaatctg agcacactgg 3600
aggacctgaa taccaggtgg aagctgctgc aggtggctgt ggaagaccgg gtgcggcagc 3660
tgcatgaggc ccatcgcgac ttcggaccag ccagccagca ctttctgagc acatccgtgc 3720
aggggccctg ggagagggcc atttctccca acaaggtgcc ctactatatt aatcacgaga 3780
cccagaccac ttgttgggac catcccaaga tgacagaact gtaccagtcc ctggccgatc 3840
tgaacaacgt gaggtttagc gcttacagaa ccgctatgaa gctgagacgg ctgcagaagg 3900
ccctgtgcct ggatctgctg tccctgtccg ccgcctgcga tgccctggat cagcataatc 3960
tgaagcagaa cgatcagcca atggatatcc tgcagatcat caactgcctg accactatct 4020
acgacaggct ggagcaggag cacaacaacc tggtgaacgt gcctctgtgc gtggatatgt 4080
gcctgaactg gctgctgaac gtgtatgaca ctgggcgcac cggccggatc agagtgctga 4140
gttttaaaac tgggattatc tccctgtgta aggcccacct ggaggacaag tacaggtacc 4200
tgttcaagca ggtggctagt agcactggat tttgtgacca gcgccgcctg ggactgctgc 4260

tgcatgatag tatccagatt cctagacagc tgggagaggt ggctagtttc ggaggatcta 4320
acatcgaacc cagcgtgcgc agctgtttcc agtttgccaa taacaaacct gaaatcgagg 4380
ctgctctgtt cctggattgg atgcgcctgg aaccacagag catggtgtgg ctgcctgtgc 4440
tgcacagagt ggctgccgcc gaaactgcca agcaccaggc taaatgcaac atctgcaagg 4500
aatgtcccat tatcggcttt cgctacagga gtctgaaaca ttttaactac gatatttgcc 4560
agagctgctt cttttccgga agagtggcca aaggacacaa gatgcactac cctatggtgg 4620
aatattgcac cccaactaca tctggcgaag atgtgcgcga ttttgccaag gtgctgaaga 4680
ataagtttcg gactaagagg tacttcgcca agcacccccg catggggtat ctgccagtgc 4740
agacagtgct ggaaggagac aatatggaga ccgatacaat gtgagcggcc gcaataaaag 4800
atctttattt tcattagatc tgtgtgttgg ttttttgtgt gtctagagca tggctacgta 4860
gataagtagc atggcgggtt aatcattaac tacaaggaac ccctagtgat ggagttggcc 4920
actccctctc tgcgcgctcg ctcgctcact gaggccgggc gaccaaaggt cgcccgacgc 4980
ccgggctttg cccgggcggc ctcagtgagc gagcgagcgc gccagctggc gtaatagcga 5040
agaggcccgc accgatcgcc cttcccaaca gttgcgcagc ctgaatggcg aatggaagtt 5100
ccagacgatt gagcgtcaaa atgtaggtat ttccatgagc gtttttcctg ttgcaatggc 5160
tggcggtaat attgttctgg atattaccag caaggccgat agtttgagtt cttctactca 5220
ggcaagtgat gttattacta atcaaagaag tattgcgaca acggttaatt tgcgtgatgg 5280
acagactctt ttactcggtg gcctcactga ttataaaaac acttctcagg attctggcgt 5340
accgttcctg tctaaaatcc ctttaatcgg cctcctgttt agctcccgct ctgattctaa 5400
cgaggaaagc acgttatacg tgctcgtcaa agcaaccata gtacgcgccc tgtagcggcg 5460
cattaagcgc ggcgggtgtg gtggttacgc gcagcgtgac cgctacactt gccagcgccc 5520
tagcgcccgc tcctttcgct ttcttccctt cctttctcgc cacgttcgcc ggctttcccc 5580
gtcaagctct aaatcggggg ctccctttag ggttccgatt tagtgattta cggcacctcg 5640
accccaaaaa acttgattag ggtgatggtt cacgtagtgg gccatcgccc tgatagacgg 5700
tttttcgccc tttgacgttg gagtccacgt tctttaatag tggactcttg ttccaaactg 5760
gaacaacact caaccctatc tcggtctatt cttttgattt ataagggatt ttgccgattt 5820
cggcctattg gttaaaaaat gagctgattt aacaaaaatt taacgcgaat tttaacaaaa 5880
tattaacgtt tacaatttaa atatttgctt atacaatctt cctgtttttg gggcttttct 5940
gattatcaac cggggtacat atgattgaca tgctagtttt acgattaccg ttcatcgatt 6000
ctcttgtttg ctccagactc tcaggcaatg acctgatagc ctttgtagag acctctcaaa 6060
aatagctacc ctctccggca tgaatttatc agctagaacg gttgaatatc atattgatgg 6120
tgatttgact gtctccggcc tttctcaccc gtttgaatct ttacctacac attactcagg 6180
cattgcattt aaaatatatg agggttctaa aaatttttat ccttgcgttg aaataaaggc 6240
ttctcccgca aaagtattac agggtcataa tgtttttggt acaaccgatt tagctttatg 6300
ctctgaggct ttattgctta attttgctaa ttctttgcct tgcctgtatg atttattgga 6360
tgttggaagt tcctgatgcg gtattttctc cttacgcatc tgtgcggtat ttcacaccgc 6420
atatggtgca ctctcagtac aatctgctct gatgccgcat agttaagcca gccccgacac 6480
ccgccaacac ccgctgacgc gccctgacgg gcttgtctgc tcccggcatc cgcttacaga 6540
caagctgtga ccgtctccgg gagctgcatg tgtcagaggt tttcaccgtc atcaccgaaa 6600
cgcgcgagac gaaagggcct cgtgatacgc ctatttttat aggttaatgt catgataata 6660
atggtttctt agacgtcagg tggcactttt cggggaaatg tgcgcggaac ccctatttgt 6720
ttatttttct aaatacattc aaatatgtat ccgctcatga gacaataacc ctgataaatg 6780
cttcaataat attgaaaaag gaagagtatg agtattcaac atttccgtgt cgcccttatt 6840
ccctttttttg cggcattttg ccttcctgtt tttgctcacc cagaaacgct ggtgaaagta 6900
aaagatgctg aagatcagtt gggtgcacga gtgggttaca tcgaactgga tctcaacagc 6960
ggtaagatcc ttgagagttt tcgccccgaa gaacgttttc caatgatgag cacttttaaa 7020

gttctgctat gtggcgcggt attatcccgt attgacgccg ggcaagagca actcggtcgc 7080
cgcatacact attctcagaa tgacttggtt gagtactcac cagtcacaga aaagcatctt 7140
acggatggca tgacagtaag agaattatgc agtgctgcca taaccatgag tgataacact 7200
gcggccaact tacttctgac aacgatcgga ggaccgaagg agctaaccgc ttttttgcac 7260
aacatggggg atcatgtaac tcgccttgat cgttgggaac cggagctgaa tgaagccata 7320
ccaaacgacg agcgtgacac cacgatgcct gtagcaatgg caacaacgtt gcgcaaacta 7380
ttaactggcg aactacttac tctagcttcc cggcaacaat taatagactg gatggaggcg 7440
gataaagttg caggaccact tctgcgctcg gcccttccgg ctggctggtt tattgctgat 7500
aaatctggag ccggtgagcg tgggtctcgc ggtatcattg cagcactggg gccagatggt 7560
aagccctccc gtatcgtagt tatctacacg acggggagtc aggcaactat ggatgaacga 7620
aatagacaga tcgctgagat aggtgcctca ctgattaagc attggtaact gtcagaccaa 7680
gtttactcat atatacttta gattgattta aaacttcatt tttaatttaa aaggatctag 7740
gtgaagatcc tttttgataa tctcatgacc aaaatccctt aacgtgagtt ttcgttccac 7800
tgagcgtcag accccgtaga aaagatcaaa ggatcttctt gagatccttt ttttctgcgc 7860
gtaatctgct gcttgcaaac aaaaaaacca ccgctaccag cggtggtttg tttgccggat 7920
caagagctac caactctttt tccgaaggta actggcttca gcagagcgca gataccaaat 7980
actgtccttc tagtgtagcc gtagttaggc caccacttca agaactctgt agcaccgcgt 8040
acatacctcg ctctgctaat cctgttacca gtggctgctg ccagtggcga taagtcgtgt 8100
cttaccgggt tggactcaag acgatagtta ccggataagg cgcagcggtc gggctgaacg 8160
ggggttcgt gcacacagcc cagcttggag cgaacgacct acaccgaact gagataccta 8220
cagcgtgagc tatgagaaag cgccacgctt cccgaaggga gaaaggcgga caggtatccg 8280
gtaagcggca gggtcggaac aggagagcgc acgagggagc ttccaggggg aaacgcctgg 8340
tatctttata gtcctgtcgg gtttcgccac ctctgacttg agcgtcgatt tttgtgatgc 8400
tcgtcagggg ggcggagcct atggaaaaac gccagcaacg cggccttttt acggttcctg 8460
gccttttgct ggccttttgc tcacatgttc tttcctgcgt tatcccctga ttctgtggat 8520
aaccgtatta ccgggtttga gtgagctgat accgctcgcc gcagccgaac gaccgagcgc 8580
agcgagtcag tgagcgacca agcggaagag c 8611

<210> 4
<211> 564
<212> DNA
<213> Adeno-associated virus

<400> 4
cagccactat gggtctaggc tgcccatgta aggaggcaag gcctggggac acccgagatg 60
cctggttata attaacccag acatgtggct gctccccccc cccaacacct gctgcctgag 120
cctcaccccc accccggtgc ctgggtctta ggctctgtac accatggagg agaagctcgc 180
tctaaaaata accctgtccc tggtgggctg tggggactg agggcaggct gtaacaggct 240
tgggggccag ggcttatacg tgcctgggac tcccaaagta ttactgttcc atgttcccgg 300
cgaagggcca gctgtccccc gccagctaga ctcagcactt agtttaggaa ccagtgagca 360
agtcagccct tggggcagcc catacaaggc catggggctg ggcaagctgc acgcctgggt 420
ccggggtggg cacggtgccc gggcaacgag ctgaaagctc atctgctctc aggggcccct 480
ccctggggac agcccctcct ggctagtcac accctgtagg ctcctctata taacccaggg 540
gcacaggggc tgcccccggg tcac 564

<210> 5
<211> 8409
<212> DNA
<213> Adeno associated virus

<400> 5

gcccaatacg caaaccgcct ctccccgcgc gttggccgat tcattaatgc agctggcgcg 60
ctcgctcgct cactgaggcc gcccgggcaa agcccgggcg tcgggcgacc tttggtcgcc 120
cggcctcagt gagcgagcga gcgcgcagag agggagtggc caactccatc actaggggtt 180
ccttgtagtt aatgattaac ccgccatgct aattatctac gtagccatgt ctagacagcc 240
actatgggtc taggctgccc atgtaaggag gcaaggcctg gggacacccg agatgcctgg 300
ttataattaa cccagacatg tggctgctcc ccccccccaa cacctgctgc ctgagcctca 360
cccccacccc ggtgcctggg tcttaggctc tgtacaccat ggaggagaag ctcgctctaa 420
aaataaccct gtccctggtg ggctgtgggg gactgagggc aggctgtaac aggcttgggg 480
gccagggctt atacgtgcct gggactccca aagtattact gttccatgtt cccggcgaag 540
ggccagctgt cccccgccag ctagactcag cacttagttt aggaaccagt gagcaagtca 600
gcccttgggg cagcccatac aaggccatgg ggctgggcaa gctgcacgcc tgggtccggg 660
gtgggcacgg tgcccgggca acgagctgaa agctcatctg ctctcagggg cccctccctg 720
gggacagccc ctcctggcta gtcacaccct gtaggctcct ctatataacc caggggcaca 780
ggggctgccc ccgggtcacc accacctcca cagcacagac agacactcag gagccagcca 840
gccaggtaag tttagtcttt ttgtctttta tttcaggtcc cggatccggt ggtggtgcaa 900
atcaaagaac tgctcctcag tggatgttgc ctttacttct aggcctgtac ggaagtgtta 960
cttctgctct aaaagctgcg gaattgtacc cgcggccgcc accatgctgt ggtgggagga 1020
ggtggaggat tgttatgaaa gggaggacgt gcagaagaag acttttacca agtgggtgaa 1080
cgctcagttc agcaaatttg ggaagcagca catcgagaat ctgttttccg acctgcagga 1140
tgggagacgg ctgctggatc tgctggaagg actgactggc cagaagctgc ccaaagagaa 1200
ggggagcact agggtgcacg ccctgaacaa cgtgaacaaa gctctgagag tgctgcagaa 1260
caacaacgtg gatctggtga atattggcag tactgatatc gtggacggga accacaaact 1320
gacactgggc ctgatctgga acattattct gcactggcag gtgaaaaatg tgatgaagaa 1380
catcatggcc gggctgcagc agaccaattc cgagaagatc ctgctgtctt gggtgcggca 1440
gagcacccgc aactatcccc aggtgaacgt gattaacttc actacatcct ggagcgacgg 1500
gctggccctg aatgctctga ttcacagcca caggcctgat ctgttcgact ggaatagcgt 1560
ggtgtgccag cagtctgcca cacagcgcct ggaacatgcc ttcaatatcg ctcggtacca 1620
gctggggatc gaaaaactgc tggacccaga ggatgtggac actacatacc cagataaaaa 1680
gtctattctg atgtacatta ctagcctgtt ccaggtgctg ccacagcagg tgtctattga 1740
agccattcag gaggtggaaa tgctgccccg ccccccaaa gtgactaaag aggagcattt 1800
tcagctgcat catcagatgc attacagcca gcagattacc gtgagcctgg ctcagggata 1860
tgagcgcacc agtagtccaa aaccacggtt caagtcctac gcttataccc aggctgccta 1920
cgtgacaact agcgacccta ctagatcccc ctttccatcc cagcacctgg aggccccaga 1980
ggacaagagc tttgggtcca gcctgatgga aagcgaggtg aatctggatc ggtaccagac 2040
agccctggag gaggtgctga gctggctgct gagtgctgaa gacacactgc aggcccaggg 2100
cgaaatttcc aatgacgtgg aagtggtgaa ggatcagttc cacacacacg agggctatat 2160
gatggacctg acagctcacc aggggcgcgt gggcaatatc ctgcagctgg gctctaaact 2220
gatcggcacc gggaaactga gtgaggacga ggaaacagaa gtgcaggagc agatgaacct 2280
gctgaacagc cgctgggagt gtctgagagt ggctagtatg gagaagcagt ccaacctgca 2340

ccgggtgctg atggacctgc agaaccagaa actgaaagag ctgaacgact ggctgacaaa 2400
gactgaggaa cgcacaagga agatggagga ggagccactg ggacccgacc tggaggatct 2460
gaagagacag gtgcagcagc ataaggtgct gcaggaggat ctggaacagg agcaggtgcg 2520
ggtgaactcc ctgacacata tggtggtggt ggtggacgaa tctagtggag atcacgccac 2580
cgccgccctg gaggaacagc tgaaggtgct gggggaccgg tgggccaaca tttgccggtg 2640
gaccgaggac aggtgggtgc tgctgcagga catcctgctg aaatggcaga ggctgaccga 2700
ggagcagtgt ctgtttagtg cttggctgag cgagaaagag gacgccgtga acaagatcca 2760
cacaaccggc tttaaggatc agaacgaaat gctgtctagc ctgcagaaac tggctgtgct 2820
gaaggccgat ctggagaaaa agaagcagag catgggcaaa ctgtatagcc tgaaacagga 2880
cctgctgagc accctgaaga acaagagcgt gacccagaag acagaagcct ggctggataa 2940
ctttgcccgc tgctgggaca acctggtgca gaaactggag aaaagtacag ctcagatctc 3000
tcaggctgtg accacaaccc agcctagcct gacccagaca accgtgatgg aaaccgtgac 3060
caccgtgaca acccgcgaac agatcctggt gaaacatgcc caggaagagc tgccacctcc 3120
acctccccag aagaagagaa ccctggagcg gctgcaggag ctgcaggaag ccactgacga 3180
actggacctg aagctgaggc aggccgaagt gattaagggg tcttggcagc ctgtgggcga 3240
tctgctgatt gattccctgc aggaccacct ggaaaaggtg aaggctctga gaggcgaaat 3300
tgctccactg aaggagaacg tgagtcatgt gaacgatctg gctagacagc tgacaacact 3360
gggcatccag ctgagcccat acaatctgag cacactggag gacctgaata ccaggtggaa 3420
gctgctgcag gtggctgtgg aagaccgggt gcggcagctg catgaggccc atcgcgactt 3480
cggaccagcc agccagcact ttctgagcac atccgtgcag gggccctggg agagggccat 3540
ttctcccaac aaggtgccct actatattaa tcacgagacc cagaccactt gttgggacca 3600
tcccaagatg acagaactgt accagtccct ggccgatctg aacaacgtga ggtttagcgc 3660
ttacagaacc gctatgaagc tgagacggct gcagaaggcc ctgtgcctgg atctgctgtc 3720
cctgtccgcc gcctgcgatg ccctggatca gcataatctg aagcagaacg atcagccaat 3780
ggatatcctg cagatcatca actgcctgac cactatctac gacaggctgg agcaggagca 3840
caacaacctg gtgaacgtgc ctctgtgcgt ggatatgtgc ctgaactggc tgctgaacgt 3900
gtatgacact gggcgcaccg gccggatcag agtgctgagt tttaaaactg ggattatctc 3960
cctgtgtaag gcccacctgg aggacaagta caggtacctg ttcaagcagg tggctagtag 4020
cactggattt tgtgaccagc gccgcctggg actgctgctg catgatagta tccagattcc 4080
tagacagctg ggagaggtgg ctagtttcgg aggatctaac atcgaaccca gcgtgcgcag 4140
ctgtttccag tttgccaata acaaacctga aatcgaggct gctctgttcc tggattggat 4200
gcgcctggaa ccacagagca tggtgtggct gcctgtgctg cacagagtgg ctgccgccga 4260
aactgccaag caccaggcta aatgcaacat ctgcaaggaa tgtcccatta tcggctttcg 4320
ctacaggagt ctgaaacatt ttaactacga tatttgccag agctgcttct tttccggaag 4380
agtggccaaa ggacacaaga tgcactaccc tatggtggaa tattgcaccc caactacatc 4440
tggcgaagat gtgcgcgatt ttgccaaggt gctgaagaat aagtttcgga ctaagaggta 4500
cttcgccaag caccccccgca tggggtatct gccagtgcag acagtgctgg aaggagacaa 4560
tatggagacc gatacaatgt gagcggccgc aataaaaagat ctttatttttc attagatctg 4620
tgtgttggtt ttttgtgtgt ctagagcatg gctacgtaga taagtagcat ggcgggttaa 4680
tcattaacta caaggaaccc ctagtgatgg agttggccac tccctctctg cgcgctcgct 4740
cgctcactga ggccgggcga ccaaaggtcg cccgacgccc gggctttgcc cgggcggcct 4800
cagtgagcga gcgagcgcgc cagctggcgt aatagcgaag aggcccgcac cgatcgccct 4860
tcccaacagt tgcgcagcct gaatggcgaa tggaagttcc agacgattga gcgtcaaaat 4920
gtaggtattt ccatgagcgt ttttcctgtt gcaatggctg gcggtaatat tgttctggat 4980
attaccagca aggccgatag tttgagttct tctactcagg caagtgatgt tattactaat 5040
caaagaagta ttgcgacaac ggttaatttg cgtgatggac agactctttt actcggtggc 5100

ctcactgatt ataaaaacac ttctcaggat tctggcgtac cgttcctgtc taaaatccct 5160
ttaatcggcc tcctgtttag ctcccgctct gattctaacg aggaaagcac gttatacgtg 5220
ctcgtcaaag caaccatagt acgcgccctg tagcggcgca ttaagcgcgg cgggtgtggt 5280
ggttacgcgc agcgtgaccg ctacacttgc cagcgcccta gcgcccgctc ctttcgcttt 5340
cttcccttcc tttctcgcca cgttcgccgg ctttccccgt caagctctaa atcgggggct 5400
ccctttaggg ttccgattta gtgatttacg gcacctcgac cccaaaaaac ttgattaggg 5460
tgatggttca cgtagtgggc catcgccctg atagacggtt tttcgccctt tgacgttgga 5520
gtccacgttc tttaatagtg gactcttgtt ccaaactgga acaacactca accctatctc 5580
ggtctattct tttgatttat aagggatttt gccgatttcg gcctattggt taaaaaatga 5640
gctgatttaa caaaaattta acgcgaattt taacaaaata ttaacgttta caatttaaat 5700
atttgcttat acaatcttcc tgtttttggg gcttttctga ttatcaaccg gggtacatat 5760
gattgacatg ctagttttac gattaccgtt catcgattct cttgtttgct ccagactctc 5820
aggcaatgac ctgatagcct ttgtagagac ctctcaaaaa tagctaccct ctccggcatg 5880
aatttatcag ctagaacggt tgaatatcat attgatggtg atttgactgt ctccggcctt 5940
tctcacccgt ttgaatcttt acctacacat tactcaggca ttgcatttaa aatatatgag 6000
ggttctaaaa atttttatcc ttgcgttgaa ataaaggctt ctcccgcaaa agtattacag 6060
ggtcataatg tttttggtac aaccgattta gctttatgct ctgaggcttt attgcttaat 6120
tttgctaatt ctttgccttg cctgtatgat ttattggatg ttggaagttc ctgatgcggt 6180
attttctcct tacgcatctg tgcggtattt cacaccgcat atggtgcact ctcagtacaa 6240
tctgctctga tgccgcatag ttaagccagc cccgacaccc gccaacaccc gctgacgcgc 6300
cctgacgggc ttgtctgctc ccggcatccg cttacagaca agctgtgacc gtctccggga 6360
gctgcatgtg tcagaggttt tcaccgtcat caccgaaacg cgcgagacga aagggcctcg 6420
tgatacgcct atttttatag gttaatgtca tgataataat ggtttcttag acgtcaggtg 6480
gcacttttcg gggaaatgtg cgcggaaccc ctatttgttt atttttctaa atacattcaa 6540
atatgtatcc gctcatgaga caataaccct gataaatgct tcaataatat tgaaaaagga 6600
agagtatgag tattcaacat ttccgtgtcg cccttattcc ctttttgcg gcattttgcc 6660
ttcctgtttt tgctcaccca gaaacgctgg tgaaagtaaa agatgctgaa gatcagttgg 6720
gtgcacgagt gggttacatc gaactggatc tcaacagcgg taagatcctt gagagttttc 6780
gccccgaaga acgttttcca atgatgagca cttttaaagt tctgctatgt ggcgcggtat 6840
tatcccgtat tgacgccggg caagagcaac tcggtcgccg catacactat tctcagaatg 6900
acttggttga gtactcacca gtcacagaaa agcatcttac ggatggcatg acagtaagag 6960
aattatgcag tgctgccata accatgagtg ataacactgc ggccaactta cttctgacaa 7020
cgatcggagg accgaaggag ctaaccgctt ttttgcacaa catgggggat catgtaactc 7080
gccttgatcg ttgggaaccg gagctgaatg aagccatacc aaacgacgag cgtgacacca 7140
cgatgcctgt agcaatggca acaacgttgc gcaaactatt aactggcgaa ctacttactc 7200
tagcttcccg gcaacaatta atagactgga tggaggcgga taaagttgca ggaccacttc 7260
tgcgctcggc ccttccggct ggctggttta ttgctgataa atctggagcc ggtgagcgtg 7320
ggtctcgcgg tatcattgca gcactggggc cagatggtaa gccctcccgt atcgtagtta 7380
tctacacgac ggggagtcag gcaactatgg atgaacgaaa tagacagatc gctgagatag 7440
gtgcctcact gattaagcat tggtaactgt cagaccaagt ttactcatat atactttaga 7500
ttgatttaaa acttcatttt taatttaaaa ggatctaggt gaagatcctt tttgataatc 7560
tcatgaccaa aatcccttaa cgtgagtttt cgttccactg agcgtcagac cccgtagaaa 7620
agatcaaagg atcttcttga gatccttttt ttctgcgcgt aatctgctgc ttgcaaacaa 7680
aaaaaccacc gctaccagcg gtggtttgtt tgccggatca agagctacca actcttttc 7740
cgaaggtaac tggcttcagc agagcgcaga taccaaatac tgtccttcta gtgtagccgt 7800
agttaggcca ccacttcaag aactctgtag caccgcgtac atacctcgct ctgctaatcc 7860

tgttaccagt ggctgctgcc agtggcgata agtcgtgtct taccgggttg gactcaagac 7920
gatagttacc ggataaggcg cagcggtcgg gctgaacggg gggttcgtgc acacagccca 7980
gcttggagcg aacgacctac accgaactga gatacctaca gcgtgagcta tgagaaagcg 8040
ccacgcttcc cgaagggaga aaggcggaca ggtatccggt aagcggcagg gtcggaacag 8100
gagagcgcac gagggagctt ccagggggaa acgcctggta tctttatagt cctgtcgggt 8160
ttcgccacct ctgacttgag cgtcgatttt tgtgatgctc gtcaggggggg cggagcctat 8220
ggaaaaacgc cagcaacgcg gccttttac ggttcctggc cttttgctgg ccttttgctc 8280
acatgttctt tcctgcgtta tcccctgatt ctgtggataa ccgtattacc gggtttgagt 8340
gagctgatac cgctcgccgc agccgaacga ccgagcgcag cgagtcagtg agcgaccaag 8400
cggaagagc 8409

CLAIMS

What is claimed:

1. A recombinant AAV vector comprising a muscle specific control element nucleotide sequence and a nucleotide sequence encoding the micro-dystrophin protein.

2. The recombinant AAV vector of claim 1 wherein the nucleotide sequence encoding the micro-dystrophin protein comprises

a) a nucleotide sequence that is at least 85% identical to the nucleotide sequence SEQ ID NO: 1 and encodes a functional micro-dystrophin protein, or

b) the nucleotide sequences of SEQ ID NO: 1.

3. The recombinant AAV vector of claim 1 or 2, wherein the muscle specific control element is human skeletal actin gene element, cardiac actin gene element, myocyte-specific enhancer binding factor (MEF), muscle creatine kinase (MCK), truncated MCK (tMCK), myosin heavy chain (MHC), hybrid α-myosin heavy chain enhancer-/MCK enhancer-promoter (MHCK7), C5-12, murine creatine kinase enhancer element, skeletal fast-twitch troponin c gene element, slow-twitch cardiac troponin c gene element, slow-twitch troponin i gene element, an hypoxia-inducible nuclear factor, steroid-inducible element, or glucocorticoid response element (GRE).

4. The recombinant AAV vector of any one of claims 1- 3, wherein the muscle specific control element is muscle creatine kinase (MCK) or hybrid α-myosin heavy chain enhancer-/MCK enhancer-promoter (MHCK7).

5. The recombinant AAV vector of any one of claims 1- 4, wherein the muscle specific control element is muscle creatine kinase (MCK) comprising the nucleotide sequence SEQ ID NO: 4 or hybrid α-myosin heavy chain enhancer-/MCK enhancer-promoter (MHCK7) comprising the nucleotide sequence SEQ ID NO: 2.

6. The recombinant AAV vector of any one of claims 1-5 comprising nucleotide sequences of SEQ ID NOS: 1 and 2.

7. The recombinant AAV vector of any one of claims 1-6 comprising the nucleotide sequence of SEQ ID NO: 3.

8. A recombinant AAV vector comprising the human micro-dystrophin nucleotide sequence of SEQ ID NO: 1 and the MHCK7 promoter sequence of SEQ ID NO: 2.

9. A recombinant AAV vector comprising the pAAV.MHCK7.micro-dystrophin construct nucleotide sequence of SEQ ID NO: 3.

10. The recombinant AAV vector of any one of claims 1-5 comprising the nucleotide sequences of SEQ ID NOS: 1 and 4.

11. The recombinant AAV vector of any one of claims 1-5 or 10 comprising the nucleotide sequence of SEQ ID NO: 5.

12. The recombinant AAV vector of any one of claims 1-11, wherein the vector is of the serotype AAVrh.74, AAV1, AAV2, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12 or AAV13.

13. The recombinant AAV vector of any one of claims 1-12, wherein the muscle specific control element nucleotide sequence is operably linked to the micro-dystrophin nucleotide sequence.

14. A composition comprising the recombinant AAV vector of any one of claims 1-13 and a pharmaceutically acceptable carrier.

15. A method of increasing muscular force or muscle mass in a subject suffering from muscular dystrophy comprising administering a therapeutically effective amount of the recombinant AAV vector of any one of claims 1-13 or the composition of claim 14.

16. A method of reducing or preventing fibrosis in a subject suffering from muscular dystrophy comprising administering a therapeutically effective amount of the recombinant AAV vector of any one of claims 1-13 or the composition of claim 14.

17. A method of treating muscular dystrophy comprising administering a therapeutically effective amount of the recombinant AAV vector of any one of claims 1-13 or the composition of claim 14.

18. A method of treating muscular dystrophy comprising administering a therapeutically effective amount of a recombinant AAV vector comprising the human micro-dystrophin nucleotide sequence of SEQ ID NO: 1 and the MHCK7 promoter nucleotide sequence of SEQ ID NO: 2.

19. A method of treating muscular dystrophy comprising administering a therapeutically effective amount of a recombinant AAV vector comprising the pAAV.MHCK7.micro-dystrophin construct nucleotide sequence of SEQ ID NO: 3.

20. The method of any one of claims 15-19, wherein the muscular dystrophy is Duchenne muscular dystrophy.

21. The method of any one of claims 15-19, wherein the recombinant AAV vector or the composition is administered by intramuscular injection or intravenous injection.

22. The method of any one of claims 15-19, wherein the recombinant AAV vector or the composition is administered systemically.

23. The method of claim 22, where the recombinant AAV vector or the composition is parenterally administered by injection, infusion or implantation.

24. A composition comprising the recombinant AAV vector of any one of claims 1-13 for increasing muscular force or muscle mass in a subject suffering from muscular dystrophy.

25. A composition comprising the recombinant AAV vector of any one of claims 1-13 for the treatment of muscular dystrophy.

26. A composition comprising a recombinant AAV vector comprising the human micro-dystrophin nucleotide sequence of SEQ ID NO: 1 and the MHCK7 promoter sequence of SEQ ID NO: 2 for the treatment of muscular dystrophy.

27. A composition comprising a recombinant AAV vector comprising the pAAV.MHCK7.micro-dystrophin construct nucleotide sequence of SEQ ID NO: 3 for the treatment of muscular dystrophy.

28. The composition of any one of claims 24-27 wherein the muscular dystrophy is Duchenne muscular dystrophy.

29. The composition of any one of claims 24-28 formulated for intramuscular injection or intravenous injection.

30. The method of any one of claims 24-28 wherein the recombinant AAV vector or the composition is administered systemically.

31. The method of claim 30, where the recombinant AAV vector or the composition is parenterally administration by injection, infusion, or implantation.

32. Use of the recombinant AAV vector of any one of claims 1-13, or the composition of claim 14 for the preparation of a medicament for increasing muscular strength or muscle mass in a subject suffering from muscular dystrophy.

33. Use of the recombinant AAV vector of any one of claims 1-13, or the composition of claim 14, for the preparation of a medicament for the treatment of muscular dystrophy.

34. Use of the recombinant AAV vector of any one of claims 1-13 or the composition of claim 14, for the preparation of a medicament for reducing or preventing fibrosis in a subject suffering from muscular dystrophy.

35. Use of a recombinant AAV vector comprising the human micro-dystrophin nucleotide sequence of SEQ ID NO: 1 and the MHCK7 promoter nucleotide sequence of SEQ ID NO: 2 for preparation of a medicament for the treatment of muscular dystrophy.

36. Use of a recombinant AAV vector comprising the pAAV.MHCK7.micro-dystrophin construct nucleotide sequence of SEQ ID NO: 3 for the treatment of muscular dystrophy.

37. The use of any one of any one of claims 32 – 36 wherein the muscular dystrophy is Duchenne muscular dystrophy.

38. The use of any one of claims 32-37 wherein the medicament is formulated for intramuscular or intravenous administration.

39. The use of any one of claims 32-37 wherein the medicament is formulated for systemic delivery.

40. The use of claim 39, where the medicament is formulated for parenteral administration by injection, infusion or implantation.