

COMPOSITIONS OF ENANTIOMERIC DIACYL TARTARIC ANHYDRIDES AND PREPARATION THEREOF

The present invention relates to compositions comprising mixtures of two enantiomeric forms of diacyl tartaric anhydride and mixtures obtainable by reacting an organic compound with a mixture comprising two enantiomeric forms of diacyl tartaric anhydride. The invention also relates to a method for the preparation of such enantiomeric mixtures, starting from a mixture comprising D-tartaric acid and L-tartaric acid.

Diacyl tartaric anhydride compounds (such as diacetyl tartaric anhydride, herein after referred to as "DATA"), are versatile chemical intermediates. Such compounds can be used, inter alia, for the manufacture of a large group of emulsifiers for food use, for example by reacting DATA with a fatty acid monoglyceride ester or the corresponding diester or any mixture thereof.

Diacyl tartaric anhydride compounds are generally prepared by reacting tartaric acid with a fatty acid anhydride (having preferably between 1 to 4 carbon atoms). Tartaric acid exists in three basic structures: L-(+)-tartaric acid (natural tartaric acid) and D-(-)-tartaric acid, which are both optical active, and the meso-form, which is not optical active. Mixtures of these forms also exist, e.g. the racemic mixture D,L-tartaric acid. Tartaric acid is a white solid material (at ambient conditions).

At present, diacetyl tartaric anhydride (DATA) is prepared by reacting tartaric acid with acetic anhydride. Both L-tartaric acid, D-tartaric acid as well as D,L-tartaric acid are not well soluble in acetic anhydride. However, once L-tartaric acid starts to react, it is solubilized in acetic anhydride very quickly. On the other hand, D-tartaric acid and D,L-tartaric acid are not well solubilized. For this reason, DATA is prepared up till now starting from L-(+)-

tartaric acid, and consequently, diacetyl-L-tartaric anhydride (L-DATA) is obtained.

Thus, in the conventional reactions for the preparation of diacyl tartaric anhydride starting from tartaric acid and a fatty acid anhydride (having preferably between 1 to 4 carbon atoms), D-tartaric acid and D,L-tartaric acid (the racemic mixture) cannot be used. This is a disadvantage, since it limits the choice of starting materials.

10 Additionally, when L-DATA is used in the preparation of emulsifiers for application of foodstuffs (as set out above), such emulsifiers can generally not be used in applications where all foodproducts and ingredients need to be kosher. Therefore, there is a need for a process in

15 which it would be possible to prepare diacyl tartaric anhydrides by reacting tartaric acid with fatty acid anhydrides (having preferably between 1 to 4 carbon atoms), and wherein it is possible to use equally well L-tartaric acid, D-tartaric acid as well as all mixtures thereof, such

20 as racemic mixtures.

It has now been found that such a process is possible, when at least a part of the reaction between tartaric acid and a fatty acid anhydride (having preferably between 1 and 4 carbon atoms) is carried out under dielectric heating conditions. Preferably, at least the first part of the reaction should be carried out under dielectric heating conditions, and more preferably dielectric heating conditions should be employed during the entire reaction

25 time. As an alternative, the dielectric heating may be employed intermittently.

In Synthetic Communications, 23(4), 419-424 (1993), it is reported that organic reactions can be carried out under dielectric heating (by a microwave). However, its use in connection to the preparation of diacyl tartaric anhydrides is neither mentioned nor suggested, and additionally, its

35

use in connection to reactions involving stereo isomers is also not reported therein.

A preferred fatty acid anhydride for the above purpose is acetic anhydride, yielding upon reaction with tartaric acid diacetyl tartaric anhydride (DATA).

It is possible to prepare compositions comprising diacyl tartaric anhydride, which diacyl tartaric anhydride is a mixture of L-diacyl tartaric anhydride and D-diacyl tartaric anhydride, in any desired ratio, depending upon the starting ratio of the L-tartaric acid and D-tartaric acid. As a consequence of this invention, it is now possible to prepare a composition comprising diacyl-L-tartaric anhydride and diacyl-D-tartaric anhydride, wherein the molar ratio of said forms of tartaric anhydride in the composition is between 1:100 and 100:1. For example, the obtained ratio between L-diacyl tartaric anhydride and D-diacyl tartaric anhydride can be any ratio between 1:20 and 20:1. Likewise, the ratio can be around 1:1, as is the case in a racemic mixture.

The compositions according to the above comprising one or more enantiomers of diacyl tartaric anhydride can be used as a starting material in further reactions. Accordingly, compositions are obtainable by reacting:

- i. an organic compound, with
- ii. a mixture comprising diacyl-L-tartaric anhydride and diacyl-D-tartaric anhydride,

wherein the molar ratio of the enantiomers obtained by said reaction of the organic compound (i) with said tartaric anhydrides (ii) is between 1:100 and 100:1, depending on the molar ratio of L- and D- tartaric acid in the starting mixture. Such compositions are also subject of the invention, as well as methods for the preparation of such compositions. Likewise, the obtained ratio can be any ratio between 1:20 and 20:1. The ratio can also be around 1:1, as

is the case in a racemic mixture. In the latter case, a racemic mixture of L-diacyl tartaric anhydride and D-diacyl tartaric anhydride can be used for said reaction. The organic compound referred to above which can be used in
5 such a reaction can be e.g. an alcohol. Preferred alcohols are e.g. glycerol monoesters or diesters of a fatty acid and any mixture thereof.

The invention is further exemplified by the following
10 example, but should not be interpreted as limiting the invention.

Example

In a three-necked flask, the following ingredients were
15 introduced:

27.6 gram (0.18 mmol) tartaric acid

56 ml (0.59 mmol) acetic anhydride

10 μ l (180 ppm) H₂SO₄ (concentrated, 96-98%, p.a.)

10 grams of glass beads.

20

Reactions according to the above recipe were carried out using L-tartaric acid as well as D,L-tartaric acid along the following lines.

25 The three-necked flask was put in a microwave oven. The temperature regulation was carried out through monitoring the temperature via a fibre optic thermometer (by Luxtron), which was connected to a computer, controlling the temperature. The temperature programme employed ensured
30 quick heating up to at least 100°C (in about 1-2 minutes), whereafter the temperature was maintained between about 100°C and about 110°C. Reactions were carried out for 15, 30 45 and 60 minutes. After dielectric heating, the volatile compounds were evaporated under reduced pressure.
35 The product obtained was in all cases an off-white crystalline compound. Both NMR and IR analysis indicated that free OH-groups were substantially absent in the

product, from which it can be concluded that the conversion was quantitative.

CLAIMS

1. A composition comprising diacyl-L-tartaric anhydride and diacyl-D-tartaric anhydride, wherein the molar ratio of said forms of tartaric anhydride is between 1:100 and 100:1.
2. A composition according to claim 1, characterized in that said ratio is between 1:20 and 20:1.
3. A composition according to claim 1 or 2, wherein the diacyl tartaric anhydride is present in the form of a racemic mixture.
4. A composition according to any of claims 1-3, characterized in that the acyl group has between 1 and 4 carbon atoms.
5. A composition according to claim 4, characterized in that the acyl group is an acetyl group.
6. A composition obtainable by reacting:
 - i. an organic compound,
 - ii. a mixture comprising diacyl-L-tartaric anhydride and diacyl-D-tartaric anhydride,wherein the molar ratio of the obtained reaction products of the organic compound (i) with said tartaric anhydrides (ii) is between 1:100 and 100:1.
7. A composition according to claim 6, characterized in that said ratios are both between 1:20 and 20:1.
8. A composition according to claim 6 or 7 comprising a racemic mixture of compounds obtained by reacting an organic compound with a mixture comprising diacyl-L-tartaric anhydride and diacyl-D-tartaric anhydride.



9. A composition according to any of claims 6-8, characterized in that the acyl group has between 1 and 4 carbon atoms.
- 5 10. A composition according to claim 9, characterized in that the acyl group is an acetyl group.
11. A composition according to any of claims 6-10, wherein said organic compound (i) is an alcohol.
- 10 12. A composition according to claim 11, wherein the alcohol is a glycerol monoester or diester of a fatty acid or any mixture thereof.
- 15 13. Method for the preparation of a mixture of diacyl tartaric anhydride compounds by reacting:
- i. a fatty acid anhydride,
- ii. a mixture comprising L-tartaric acid and D-tartaric acid, wherein the molar ratio of said
- 20 forms of tartaric acid is between 1:100 and 100:1,
- wherein at least part of the reaction is carried out under dielectric heating.
- 25 14. Method according to claim 13, wherein said ratio is between 1:20 and 20:1.
15. Method according to claim 13 or 14, wherein the tartaric acid is present in the form of a racemic
- 30 mixture.
16. Method according to any of claims 13-15, wherein the fatty acid anhydride has between 1 and 4 carbon atoms.
- 35 17. Method according to claim 16, wherein the fatty acid anhydride is acetic anhydride.





PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07C 59/347, 51/567	A1	(11) International Publication Number: WO 96/35658 (43) International Publication Date: 14 November 1996 (14.11.96)
(21) International Application Number: PCT/EP96/01934 (22) International Filing Date: 6 May 1996 (06.05.96) (30) Priority Data: 95201233.4 12 May 1995 (12.05.95) HP (34) Countries for which the regional or international application was filed: NL et al. (71) Applicant (for all designated States except US): QUEST INTERNATIONAL B.V. [NL/NL]; Huizerstraatweg 28, NL-1411 GP Naarden (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): COENDERS, Peter, Joannes [NL/NL]; Pearl Buckerf 98, NL-3315 BB Dordrecht (NL). GALEMA, Saskia, Alexandra [NL/NL]; Bernweg 173, NL-3137 NC Vlaardingen (NL). GOMBERT, Johannes, Durk [NL/NL]; Kerkpad 10, NL-6883 AA Velp (NL). MEER, Anna, Sophie, E., ter [NL/NL]; Aalscholverlaan 266, NL-3136 KL Vlaardingen (NL).	(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIP patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: COMPOSITIONS OF ENANTIOMERIC DIACYL TARTARIC ANHYDRIDES AND PREPARATION THEREOF (57) Abstract <p>The present invention relates to compositions comprising mixtures of two enantiomeric forms of diacyl tartaric anhydride and mixtures obtainable by reacting an organic compound with a mixture comprising two enantiomeric forms of diacyl tartaric anhydride. The invention also relates to a method for the preparation of such enantiomeric mixtures, starting from a mixture comprising D-tartaric acid and L-tartaric acid. The method involves reacting tartaric acid with a fatty acid anhydride under dielectric heating conditions.</p>		