



Abstract**1,1'-Bis(phosphino)ferrocene ligands for alkoxy carbonylation**

Diastereomer mixture comprising diastereomers of the formulae (I.1) and (I.2)



where

R^2 , R^4 are each independently selected from $-(C_1-C_{12})$ -alkyl, $-(C_3-C_{12})$ -cycloalkyl, $-(C_3-C_{12})$ -heterocycloalkyl, $-(C_6-C_{20})$ -aryl;

the R^1 , R^3 radicals are each a $-(C_3-C_{20})$ -heteroaryl radical;

R^1 , R^3 may each independently be substituted by one or more substituents selected from $-(C_1-C_{12})$ -alkyl, $-(C_3-C_{12})$ -cycloalkyl, $-(C_3-C_{12})$ -heterocycloalkyl, $-O-(C_1-C_{12})$ -alkyl, $-O-(C_1-C_{12})$ -alkyl- $-(C_6-C_{20})$ -aryl, $-O-(C_3-C_{12})$ -cycloalkyl, $-S-(C_1-C_{12})$ -alkyl, $-S-(C_3-C_{12})$ -cycloalkyl, $-COO-(C_1-C_{12})$ -alkyl, $-COO-(C_3-C_{12})$ -cycloalkyl, $-CONH-(C_1-C_{12})$ -alkyl, $-CONH-(C_3-C_{12})$ -cycloalkyl, $-CO-(C_1-C_{12})$ -alkyl, $-CO-(C_3-C_{12})$ -cycloalkyl, $-N-[(C_1-C_{12})$ -alkyl] $_2$, $-(C_6-C_{20})$ -aryl, $-(C_6-C_{20})$ -aryl- $-(C_1-C_{12})$ -alkyl, $-(C_6-C_{20})$ -aryl- $O-(C_1-C_{12})$ -alkyl, $-(C_3-C_{20})$ -heteroaryl, $-(C_3-C_{20})$ -heteroaryl- $-(C_1-C_{12})$ -alkyl, $-(C_3-C_{20})$ -heteroaryl- $O-(C_1-C_{12})$ -alkyl, $-COOH$, $-OH$, $-SO_3H$, $-NH_2$, halogen;

and

R^2 , R^4 , if they are $-(C_1-C_{12})$ -alkyl, $-(C_3-C_{12})$ -cycloalkyl, $-(C_3-C_{12})$ -heterocycloalkyl or $-(C_6-C_{20})$ -aryl, may each independently be substituted by one or more substituents selected from $-(C_1-C_{12})$ -alkyl, $-(C_3-C_{12})$ -cycloalkyl, $-(C_3-C_{12})$ -heterocycloalkyl, $-O-(C_1-C_{12})$ -alkyl, $-O-(C_1-C_{12})$ -alkyl- $-(C_6-C_{20})$ -aryl, $-O-(C_3-C_{12})$ -cycloalkyl, $-S-(C_1-C_{12})$ -alkyl, $-S-(C_3-C_{12})$ -cycloalkyl, $-COO-(C_1-C_{12})$ -alkyl, $-COO-(C_3-C_{12})$ -cycloalkyl, $-CONH-(C_1-C_{12})$ -alkyl, $-CONH-(C_3-C_{12})$ -cycloalkyl, $-CO-(C_1-C_{12})$ -alkyl, $-CO-(C_3-C_{12})$ -cycloalkyl, $-N-[(C_1-C_{12})$ -alkyl] $_2$, $-(C_6-C_{20})$ -aryl, $-(C_6-C_{20})$ -aryl- $-(C_1-C_{12})$ -alkyl, $-(C_6-C_{20})$ -aryl- $O-(C_1-C_{12})$ -alkyl, $-(C_3-C_{20})$ -heteroaryl, $-(C_3-C_{20})$ -heteroaryl- $-(C_1-C_{12})$ -alkyl, $-(C_3-C_{20})$ -heteroaryl- $O-(C_1-C_{12})$ -alkyl, $-COOH$, $-OH$, $-SO_3H$, $-NH_2$, halogen.

The invention further relates to Pd complex mixtures comprising the diastereomers according to the invention, and to the use thereof in an alkoxy carbonylation process.