

BICYCLIC PEPTIDE LIGANDS SPECIFIC FOR EphA2

FIELD OF THE INVENTION

The present invention relates to polypeptides which are covalently bound to non-aromatic molecular scaffolds such that two or more peptide loops are subtended between attachment points to the scaffold. In particular, the invention describes peptides which are high affinity binders of the Eph receptor tyrosine kinase A2 (EphA2). The invention also includes drug conjugates comprising said peptides, conjugated to one or more effector and/or functional groups, to pharmaceutical compositions comprising said peptide ligands and drug conjugates and to the use of said peptide ligands and drug conjugates in preventing, suppressing or treating a disease or disorder characterised by overexpression of EphA2 in diseased tissue (such as a tumour).

BACKGROUND OF THE INVENTION

Cyclic peptides are able to bind with high affinity and target specificity to protein targets and hence are an attractive molecule class for the development of therapeutics. In fact, several cyclic peptides are already successfully used in the clinic, as for example the antibacterial peptide vancomycin, the immunosuppressant drug cyclosporine or the anti-cancer drug octreotide (Driggers *et al.* (2008), Nat Rev Drug Discov 7 (7), 608-24). Good binding properties result from a relatively large interaction surface formed between the peptide and the target as well as the reduced conformational flexibility of the cyclic structures. Typically, macrocycles bind to surfaces of several hundred square angstrom, as for example the cyclic peptide CXCR4 antagonist CVX15 (400 Å²; Wu *et al.* (2007), Science 330, 1066-71), a cyclic peptide with the Arg-Gly-Asp motif binding to integrin αVβ3 (355 Å²) (Xiong *et al.* (2002), Science 296 (5565), 151-5) or the cyclic peptide inhibitor upain-1 binding to urokinase-type plasminogen activator (603 Å²; Zhao *et al.* (2007), J Struct Biol 160 (1), 1-10).

Due to their cyclic configuration, peptide macrocycles are less flexible than linear peptides, leading to a smaller loss of entropy upon binding to targets and resulting in a higher binding affinity. The reduced flexibility also leads to locking target-specific conformations, increasing binding specificity compared to linear peptides. This effect has been exemplified by a potent and selective inhibitor of matrix metalloproteinase 8, (MMP-8) which lost its selectivity over other MMPs when its ring was opened (Cherney *et al.* (1998), J Med Chem 41 (11), 1749-51). The favorable binding properties achieved through macrocyclization are even more pronounced in multicyclic peptides having more than one peptide ring as for example in vancomycin, nisin and actinomycin.

Different research teams have previously tethered polypeptides with cysteine residues to a synthetic molecular structure (Kemp and McNamara (1985), J. Org. Chem; Timmerman *et al.* (2005), ChemBioChem). Meloen and co-workers had used tris(bromomethyl)benzene and related molecules for rapid and quantitative cyclisation of multiple peptide loops onto synthetic scaffolds for structural mimicry of protein surfaces (Timmerman *et al.* (2005), ChemBioChem). Methods for the generation of candidate drug compounds wherein said compounds are generated by linking cysteine containing polypeptides to a molecular scaffold as for example TATA (1,1',1''-(1,3,5-triazinane-1,3,5-triyl)triprop-2-en-1-one, Heinis *et al.* Angew Chem, Int Ed. 2014; 53:1602–1606).

10

Phage display-based combinatorial approaches have been developed to generate and screen large libraries of bicyclic peptides to targets of interest (Heinis *et al.* (2009), Nat Chem Biol 5 (7), 502-7 and WO 2009/098450). Briefly, combinatorial libraries of linear peptides containing three cysteine residues and two regions of six random amino acids (Cys-(Xaa)₆-Cys-(Xaa)₆-Cys) were displayed on phage and cyclised by covalently linking the cysteine side chains to a small molecule scaffold.

15

SUMMARY OF THE INVENTION

According to a first aspect of the invention, there is provided a peptide ligand specific for EphA2 comprising a polypeptide comprising at least three cysteine residues, separated by at least two loop sequences, and a non-aromatic molecular scaffold which forms covalent bonds with the cysteine residues of the polypeptide such that at least two polypeptide loops are formed on the molecular scaffold.

20

According to a further aspect of the invention, there is provided a drug conjugate comprising a peptide ligand as defined herein conjugated to one or more effector and/or functional groups.

25

According to a further aspect of the invention, there is provided a pharmaceutical composition comprising a peptide ligand or a drug conjugate as defined herein in combination with one or more pharmaceutically acceptable excipients.

30

According to a further aspect of the invention, there is provided a peptide ligand or drug conjugate as defined herein for use in preventing, suppressing or treating a disease or disorder characterised by overexpression of EphA2 in diseased tissue (such as a tumour).

35

BRIEF DESCRIPTION OF THE FIGURES

Figure 1: Body weight changes after administering BCY6031 to female Balb/C nude mice bearing LU-01-0046 tumor. Data points represent group mean body weight.

Figure 2: Tumor volume trace after administering BCY6031 to female Balb/C nude mice bearing LU-01-0046 tumor. Data points represent group mean. The treatment was
5 ceased from day 28.

Figure 3: General schematic demonstrating the concept of preparing Bicycle drug conjugates (BDCs).

Figure 4: Plot of mean tumour volume versus time for BCY6136 in HT1080 xenograft mice. Doses (2, 3 and 5 mg/kg) were administered on days 0 and 7. Body weight
10 changes during treatment indicative of tumour burden, drug-associated toxicology and overall animal health are illustrated in the top right inset.

Figure 5: Plot of mean tumour volume versus time for BCY6136 in NCI-H1975 xenograft mice. Doses (1, 2 and 3 mg/kg) were administered on days 0, 7, 14, 21, 28 and
15 35. Body weight changes during treatment indicative of tumour burden, drug-associated toxicology and overall animal health are illustrated in the top right inset.

Figure 6: Plot of mean tumour volume versus time for BCY6136 in MDA-MB-231 xenograft mice. Doses (1, 2 and 3 mg/kg) were administered on day 0, 7, 14, 21, 28, 35
and 45. Body weight changes during treatment indicative of tumour burden, drug-associated toxicology and overall animal health are illustrated in the top right inset.

Figures 7 to 9: Body weight changes and tumor volume traces after administering BCY6136 (Figure 7), ADC (Figure 8) and BCY6033 (Figure 9) to female
20 BALB/c nude mice bearing PC-3 xenograft. Data points represent group mean body weight.

Figure 10: Body weight changes and tumor volume traces after administering BCY6136, EphA2-ADC or Docetaxel to male Balb/c nude mice bearing PC-3 xenograft. Data
25 points represent group mean body weight.

Figures 11 to 13: Body weight changes and tumor volume trace after administering BCY6033 (Figure 11), BCY6136 (Figure 12) and BCY6082 (Figure 13) to female
Balb/c nude mice bearing NCI-H1975 xenograft. Data points represent group mean tumor
volume and body weight.

Figures 14 and 15: Body weight changes and tumor volume traces after administering BCY6136 and ADC to female Balb/c nude mice bearing LU-01-0251 xenograft.
Data points represent group mean body weight.

Figure 16: Body weight changes and tumor volume traces after administering BCY6033, BCY6136, BCY6082 and BCY6031 to female Balb/c nude mice bearing LU-01-
35 0046. Data points represent group mean body weight.

Figure 17: Body weight changes and tumor volume traces after administering BCY6136 or ADC to female Balb/c nude mice bearing LU-01-0046 NSCLC PDX model. Data points represent group mean body weight.

Figures 18 to 22: Body weight changes and tumor volume traces after administering BCY6033 (Figure 18), BCY6136 (Figure 19), BCY6082 (Figure 20), BCY6173 (Figure 21) and BCYs 6175 and 6031 (Figure 22) to female Balb/c nude mice bearing LU-01-0046. Data points represent group mean body weight.

Figure 23: Body weight changes and tumor volume traces after administering BCY6136 (referred to in Figure 23 as BT5528), BCY8245 or BCY8781 to female BALB/c nude mice bearing LU-01-0412 xenograft. Data points represent group mean tumor volume (left panel) and body weight (right panel).

Figure 24: Body weight changes and tumor volume traces after administering BCY6136 to female Balb/c nude mice bearing LU-01-0486 xenograft. Data points represent group mean body weight.

Figures 25 to 27: Body weight changes and tumor volume trace after administering BCY6033 (Figure 25), BCY6136 (Figure 26) and BCY6082 (Figure 27) to female Balb/c nude mice bearing MDA-MB-231-luc xenograft. Data points represent group mean tumor volume and body weight.

Figure 28: Body weight changes and tumor volume traces after administering BCY6136 to female BALB/c mice bearing EMT-6 syngeneic. Data points represent group mean body weight. The dosage of group 3 and group 4 was changed to 5 mpk and 3 mpk from Day 14.

Figure 29: Body weight changes and tumor volume traces after administering BCY6136 to female Balb/c nude mice bearing NCI-N87 xenograft. Data points represent group mean body weight.

Figure 30: Body weight changes and tumor volume traces after administering BCY6136 to female Balb/c nude mice bearing SK-OV-3 xenograft. Data points represent group mean body weight.

Figure 31: Body weight changes and tumor volume traces after administering BCY6136 to female Balb/c nude mice bearing OE21 xenograft. Data points represent group mean body weight.

Figure 32: Body weight changes and tumor volume traces after administering BCY6136 to female CB17-SCID mice bearing MOLP-8 xenograft. Data points represent group mean body weight.

Figure 33: Body weight changes and Tumor volume traces after administering BCY6082 to female CB17-SCID mice bearing MOLP-8 xenograft. Data points represent group mean body weight.

Figures 34 to 42: Body weight changes and tumor volume traces after administering BCY6082 (Figure 34), BCY6031 (Figure 35), BCY6173 (Figure 36), BCY6135 (Figure 37), BCY6033 (Figure 38), BCY6136 (Figure 39), BCY6174 (Figure 40), BCY6175 (Figure 41) and ADC (Figure 42) to female BALB/c nude mice bearing HT1080 xenograft. Data points represent group mean body weight. Where error bars are present in the above Figures, these represent standard error of the mean (SEM).

DETAILED DESCRIPTION OF THE INVENTION

10 In one embodiment, said loop sequences comprise 2, 3, 5, 6 or 7 amino acid acids.

In a further embodiment, said loop sequences comprise three cysteine residues separated by two loop sequences one of which consists of 2 amino acids and the other of which consists of 7 amino acids (such as those listed in Table 4).

15

In a further embodiment, said loop sequences comprise three cysteine residues separated by two loop sequences both of which consist of 5 amino acids (such as those listed in Tables 3 and 4).

20 In a further embodiment, said loop sequences comprise three cysteine residues separated by two loop sequences both of which consist of 6 amino acids (such as those listed in Tables 3 to 5).

25 In a further embodiment, said loop sequences comprise three cysteine residues separated by two loop sequences both of which consist of 6 amino acids (such as those listed in Table 10).

30 In a further embodiment, said loop sequences comprise three cysteine residues separated by two loop sequences one of which consists of 7 amino acids and the other of which consists of 3 amino acids (such as those listed in Table 4).

35 In a further embodiment, said loop sequences comprise three cysteine residues separated by two loop sequences one of which consists of 6 amino acids and the other of which consists of 7 amino acids (such as those listed in Table 5).

In one embodiment, the peptide ligand comprises an amino acid sequence selected from:



wherein X_1 and X_2 represent the amino acid residues between the cysteine residues listed in Tables 3 to 5 and C_i , C_{ii} and C_{iii} represent first, second and third cysteine residues, respectively or a pharmaceutically acceptable salt thereof.

- 5 In a further embodiment, the peptide ligand comprises an amino acid sequence selected from one or more of the peptide ligands listed in one or more Tables 3 to 5.

In a further embodiment, the peptide ligand comprises an amino acid sequence selected from:

10 $C_i-X_1-C_{ii}-X_2-C_{iii}$

wherein X_1 and X_2 represent the amino acid residues between the cysteine residues listed in Table 10 and C_i , C_{ii} and C_{iii} represent first, second and third cysteine residues, respectively or a pharmaceutically acceptable salt thereof.

- 15 In a further embodiment, the peptide ligand comprises an amino acid sequence selected from one or more of the peptide ligands listed in Table 10.

In one embodiment, said loop sequences comprise three cysteine residues separated by two loop sequences both of which consist of 6 amino acids and the peptide ligand has an amino acid sequence selected from:

$C_i(\text{HyP})\text{LVNPLC}_{ii}\text{LHP}(\text{D-Asp})\text{W}(\text{HArg})\text{C}_{iii}$ (SEQ ID NO: 1); and

$C_i\text{PLVNPLC}_{ii}\text{LHPGWTC}_{iii}$ (SEQ ID NO: 97);

wherein HyP is hydroxyproline, HArg is homoarginine and C_i , C_{ii} and C_{iii} represent first, second and third cysteine residues, respectively or a pharmaceutically acceptable salt thereof.

In a further embodiment, said loop sequences comprise three cysteine residues separated by two loop sequences both of which consist of 6 amino acids and the peptide ligand has the following amino acid sequence:

30 $C_i(\text{HyP})\text{LVNPLC}_{ii}\text{LHP}(\text{D-Asp})\text{W}(\text{HArg})\text{C}_{iii}$ (SEQ ID NO: 1);

wherein HyP is hydroxyproline, HArg is homoarginine and C_i , C_{ii} and C_{iii} represent first, second and third cysteine residues, respectively or a pharmaceutically acceptable salt thereof.

- 35 In one embodiment, the peptide ligand of the invention is a peptide ligand which is other than the amino acid sequence:

$C_i(\text{HyP})\text{LVNPLC}_{ii}\text{LHP}(\text{D-Asp})\text{W}(\text{HArg})\text{C}_{iii}$ (SEQ ID NO: 1).

In one embodiment, said loop sequences comprise three cysteine residues separated by two loop sequences both of which consist of 6 amino acids and the peptide ligand has an amino acid sequence selected from:

5 (β-Ala)-Sar₁₀-A(HArg)D-C_i(HyP)LVNPLC_{ii}LHP(D-Asp)W(HArg)C_{iii} (SEQ ID NO: 2) (BCY6099; Compound 66); and

(β-Ala)-Sar₁₀-A(HArg)D-C_iPLVNPLC_{ii}LHPGWTC_{iii} ((β-Ala)-Sar₁₀-(SEQ ID NO: 11)) (BCY6014; Compound 67);

wherein Sar is sarcosine, HArg is homoarginine and HyP is hydroxyproline.

10

In a further embodiment, said loop sequences comprise three cysteine residues separated by two loop sequences both of which consist of 6 amino acids and the peptide ligand has the following amino acid sequence:

(β-Ala)-Sar₁₀-A(HArg)D-C_i(HyP)LVNPLC_{ii}LHP(D-Asp)W(HArg)C_{iii} (SEQ ID NO: 2) (BCY6099; Compound 66);

15

wherein Sar is sarcosine, HArg is homoarginine and HyP is hydroxyproline.

In one embodiment, the peptide ligand of the invention is a peptide ligand which is other than the amino acid sequence:

20 (β-Ala)-Sar₁₀-A(HArg)D-C_i(HyP)LVNPLC_{ii}LHP(D-Asp)W(HArg)C_{iii} (SEQ ID NO: 2) (BCY6099; Compound 66).

In one embodiment, the molecular scaffold is selected from 1,1',1''-(1,3,5-triazinane-1,3,5-triyl)triprop-2-en-1-one (TATA) and the peptide ligand is selected from any one of the peptide

25 ligands listed in Tables 3 to 5.

In an alternative embodiment, the molecular scaffold is selected from 1,1',1''-(1,3,5-triazinane-1,3,5-triyl)triprop-2-en-1-one (TATA) and the peptide ligand is selected from any one of the peptide ligands listed in Table 10.

30

In one embodiment, the molecular scaffold is selected from 1,1',1''-(1,3,5-triazinane-1,3,5-triyl)triprop-2-en-1-one (TATA) and the peptide ligand is selected from:

(β-Ala)-Sar₁₀-A(HArg)D-C_i(HyP)LVNPLC_{ii}LHP(D-Asp)W(HArg)C_{iii} (SEQ ID NO: 2) (BCY6099; Compound 66); and

35 (β-Ala)-Sar₁₀-A(HArg)D-C_iPLVNPLC_{ii}LHPGWTC_{iii} ((β-Ala)-Sar₁₀-(SEQ ID NO: 11)) (BCY6014; Compound 67);

wherein Sar is sarcosine, HArg is homoarginine and HyP is hydroxyproline.

In a further embodiment, the molecular scaffold is selected from 1,1',1''-(1,3,5-triazinane-1,3,5-triyl)triprop-2-en-1-one (TATA) and the peptide ligand is:

(β -Ala)-Sar₁₀-A(HArg)D-C_i(HyP)LVNPLC_{ii}LHP(D-Asp)W(HArg)C_{iii} (SEQ ID NO: 2);

5 wherein Sar is sarcosine, HArg is homoarginine and HyP is hydroxyproline.

In one embodiment, the peptide ligand is selected from any one of Compounds 1-113 or a pharmaceutically acceptable salt thereof.

10 In a further embodiment, the peptide ligand is Compound 66 (BCY6099) or Compound 67 (BCY6014) or a pharmaceutically acceptable salt thereof.

In a yet further embodiment, the peptide ligand is Compound 66 (BCY6099) or a pharmaceutically acceptable salt thereof.

15

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the art, such as in the arts of peptide chemistry, cell culture and phage display, nucleic acid chemistry and biochemistry. Standard techniques are used for molecular biology, genetic and biochemical methods (see
20 Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, 3rd ed., 2001, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; Ausubel *et al.*, Short Protocols in Molecular Biology (1999) 4th ed., John Wiley & Sons, Inc.), which are incorporated herein by reference.

Nomenclature

25 Numbering

When referring to amino acid residue positions within the peptides of the invention, cysteine residues (C_i, C_{ii} and C_{iii}) are omitted from the numbering as they are invariant, therefore, the numbering of amino acid residues within the peptides of the invention is referred to as below:

30 -C_i-HyP₁-L₂-V₃-N₄-P₅-L₆-C_{ii}-L₇-H₈-P₉-(D-Asp)₁₀-W₁₁-(HArg)₁₂-C_{iii}- (SEQ ID NO: 1).

For the purpose of this description, all bicyclic peptides are assumed to be cyclised with 1,1',1''-(1,3,5-triazinane-1,3,5-triyl)triprop-2-en-1-one (TATA) yielding a tri-substituted 1,1',1''-(1,3,5-triazinane-1,3,5-triyl)tripropan-1-one structure. Cyclisation with TATA occurs on C_i, C_{ii},
35 and C_{iii}.

Molecular Format

N- or C-terminal extensions to the bicycle core sequence are added to the left or right side of the sequence, separated by a hyphen. For example, an N-terminal (β -Ala)-Sar₁₀-Ala tail would be denoted as:

(β -Ala)-Sar₁₀-A-(SEQ ID NO: X).

5

Inversed Peptide Sequences

In light of the disclosure in Nair *et al* (2003) J Immunol 170(3), 1362-1373, it is envisaged that the peptide sequences disclosed herein would also find utility in their retro-inverso form. For example, the sequence is reversed (i.e. N-terminus become C-terminus and *vice versa*) and their stereochemistry is likewise also reversed (i.e. D-amino acids become L-amino acids and *vice versa*).

10

Peptide Ligands

A peptide ligand, as referred to herein, refers to a peptide, peptidic or peptidomimetic covalently bound to a molecular scaffold. Typically, such peptides, peptidics or peptidomimetics comprise a peptide having natural or non-natural amino acids, two or more reactive groups (i.e. cysteine residues) which are capable of forming covalent bonds to the scaffold, and a sequence subtended between said reactive groups which is referred to as the loop sequence, since it forms a loop when the peptide, peptidic or peptidomimetic is bound to the scaffold. In the present case, the peptides, peptidics or peptidomimetics comprise at least three cysteine residues (referred to herein as C_i, C_{ii} and C_{iii}), and form at least two loops on the scaffold.

15

20

Advantages of the Peptide Ligands

Certain bicyclic peptides of the present invention have a number of advantageous properties which enable them to be considered as suitable drug-like molecules for injection, inhalation, nasal, ocular, oral or topical administration. Such advantageous properties include:

25

- Species cross-reactivity. This is a typical requirement for preclinical pharmacodynamics and pharmacokinetic evaluation;

30

- Protease stability. Bicyclic peptide ligands should in most circumstances demonstrate stability to plasma proteases, epithelial ("membrane-anchored") proteases, gastric and intestinal proteases, lung surface proteases, intracellular proteases and the like. Protease stability should be maintained between different species such that a bicyclic peptide lead candidate can be developed in animal models as well as administered with confidence to humans;

35

- Desirable solubility profile. This is a function of the proportion of charged and hydrophilic versus hydrophobic residues and intra/inter-molecular H-bonding, which is important for formulation and absorption purposes;
- 5 - An optimal plasma half-life in the circulation. Depending upon the clinical indication and treatment regimen, it may be required to develop a bicyclic peptide with short or prolonged *in vivo* exposure times for the management of either chronic or acute disease states. The optimal exposure time will be governed by the requirement for sustained exposure (for maximal therapeutic efficiency) versus the requirement for short exposure times to minimise
10 toxicological effects arising from sustained exposure to the agent;
- Selectivity. Certain peptide ligands of the invention demonstrate good selectivity over other Eph receptor tyrosine kinases, such as EphA1, EphA3, EphA4, EphA5, EphA6, EphA7 and EphB1 and factor XIIA, carbonic anhydrase 9 and CD38 (selectivity data for selected
15 peptide ligands of the invention may be seen in Tables 7 and 14). It should also be noted that selected peptide ligands of the invention exhibit cross reactivity with other species (eg mouse and rat) to permit testing in animal models (Tables 3 to 6 and 15); and
- Safety. Bleeding events have been reported in pre-clinical *in vivo* models and clinical
20 trials with EphA2 Antibody Drug Conjugates. For example, a phase 1, open-label study with MEDI-547 was halted due to bleeding and coagulation events that occurred in 5 of 6 patients (Annunziata *et al*, Invest New Drugs (2013) 31:77–84). The bleeding events observed in patients were consistent with effects on the coagulation system observed in rat and monkey pre-clinical studies: increased activated partial thromboplastin time and increased
25 fibrinogen/fibrin degradation product (Annunziata *et al* /IBID). Overt bleeding events were reportedly seen in toxicology studies in monkeys (Annunziata *et al*, IBID). Taken together these results imply that MEDI-547 causes Disseminated Intravascular Coagulation (DIC) in both preclinical species and patients. The BDCs reported here have short *in vivo* half lives (< 30 minutes) and are therefore intrinsically less likely to give rise to DIC in patients. Results
30 shown here (see BIOLOGICAL DATA sections 5 and 6 and Table 20) demonstrate that selected Bicycle Drug Conjugates of the invention have no effect on coagulation parameters and gave rise to no bleeding events in pre-clinical studies.

Pharmaceutically Acceptable Salts

- 35 It will be appreciated that salt forms are within the scope of this invention, and references to peptide ligands include the salt forms of said ligands.

The salts of the present invention can be synthesized from the parent compound that contains a basic or acidic moiety by conventional chemical methods such as methods described in *Pharmaceutical Salts: Properties, Selection, and Use*, P. Heinrich Stahl (Editor), Camille G. Wermuth (Editor), ISBN: 3-90639-026-8, Hardcover, 388 pages, August 2002.

- 5 Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with the appropriate base or acid in water or in an organic solvent, or in a mixture of the two.

- Acid addition salts (mono- or di-salts) may be formed with a wide variety of acids, both
- 10 inorganic and organic. Examples of acid addition salts include mono- or di-salts formed with an acid selected from the group consisting of acetic, 2,2-dichloroacetic, adipic, alginic, ascorbic (e.g. L-ascorbic), L-aspartic, benzenesulfonic, benzoic, 4-acetamidobenzoic, butanoic, (+) camphoric, camphor-sulfonic, (+)-(1S)-camphor-10-sulfonic, capric, caproic, caprylic, cinnamic, citric, cyclamic, dodecylsulfuric, ethane-1,2-disulfonic, ethanesulfonic, 2-
- 15 hydroxyethanesulfonic, formic, fumaric, galactaric, gentisic, glucoheptonic, D-gluconic, glucuronic (e.g. D-glucuronic), glutamic (e.g. L-glutamic), α -oxoglutaric, glycolic, hippuric, hydrohalic acids (e.g. hydrobromic, hydrochloric, hydriodic), isethionic, lactic (e.g. (+)-L-lactic, ($\pm$)-DL-lactic), lactobionic, maleic, malic, (-)-L-malic, malonic, ($\pm$)-DL-mandelic, methanesulfonic, naphthalene-2-sulfonic, naphthalene-1,5-disulfonic, 1-hydroxy-2-naphthoic,
- 20 nicotinic, nitric, oleic, orotic, oxalic, palmitic, pamoic, phosphoric, propionic, pyruvic, L-pyroglutamic, salicylic, 4-amino-salicylic, sebacic, stearic, succinic, sulfuric, tannic, (+)-L-tartaric, thiocyanic, *p*-toluenesulfonic, undecylenic and valeric acids, as well as acylated amino acids and cation exchange resins.

- 25 One particular group of salts consists of salts formed from acetic, hydrochloric, hydriodic, phosphoric, nitric, sulfuric, citric, lactic, succinic, maleic, malic, isethionic, fumaric, benzenesulfonic, toluenesulfonic, sulfuric, methanesulfonic (mesylate), ethanesulfonic, naphthalenesulfonic, valeric, propanoic, butanoic, malonic, glucuronic and lactobionic acids. One particular salt is the hydrochloride salt. Another particular salt is the acetate salt.

- 30 If the compound is anionic, or has a functional group which may be anionic (e.g., -COOH may be -COO⁻), then a salt may be formed with an organic or inorganic base, generating a suitable cation. Examples of suitable inorganic cations include, but are not limited to, alkali metal ions such as Li⁺, Na⁺ and K⁺, alkaline earth metal cations such as Ca²⁺ and Mg²⁺, and
- 35 other cations such as Al³⁺ or Zn²⁺. Examples of suitable organic cations include, but are not limited to, ammonium ion (i.e., NH₄⁺) and substituted ammonium ions (e.g., NH₃R⁺, NH₂R₂⁺, NHR₃⁺, NR₄⁺). Examples of some suitable substituted ammonium ions are those derived

from: methylamine, ethylamine, diethylamine, propylamine, dicyclohexylamine, triethylamine, butylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine, benzylamine, phenylbenzylamine, choline, meglumine, and tromethamine, as well as amino acids, such as lysine and arginine. An example of a common quaternary ammonium ion is $N(CH_3)_4^+$.

5

Where the peptides of the invention contain an amine function, these may form quaternary ammonium salts, for example by reaction with an alkylating agent according to methods well known to the skilled person. Such quaternary ammonium compounds are within the scope of the peptides of the invention.

10

Modified Derivatives

It will be appreciated that modified derivatives of the peptide ligands as defined herein are within the scope of the present invention. Examples of such suitable modified derivatives include one or more modifications selected from: N-terminal and/or C-terminal modifications; replacement of one or more amino acid residues with one or more non-natural amino acid residues (such as replacement of one or more polar amino acid residues with one or more isosteric or isoelectronic amino acids; replacement of one or more non-polar amino acid residues with other non-natural isosteric or isoelectronic amino acids); addition of a spacer group; replacement of one or more oxidation sensitive amino acid residues with one or more oxidation resistant amino acid residues; replacement of one or more amino acid residues with one or more replacement amino acids, such as an alanine, replacement of one or more L-amino acid residues with one or more D-amino acid residues; N-alkylation of one or more amide bonds within the bicyclic peptide ligand; replacement of one or more peptide bonds with a surrogate bond; peptide backbone length modification; substitution of the hydrogen on the alpha-carbon of one or more amino acid residues with another chemical group, modification of amino acids such as cysteine, lysine, glutamate/aspartate and tyrosine with suitable amine, thiol, carboxylic acid and phenol-reactive reagents so as to functionalise said amino acids, and introduction or replacement of amino acids that introduce orthogonal reactivities that are suitable for functionalisation, for example azide or alkyne-group bearing amino acids that allow functionalisation with alkyne or azide-bearing moieties, respectively.

In one embodiment, the modified derivative comprises an N-terminal and/or C-terminal modification. In a further embodiment, wherein the modified derivative comprises an N-terminal modification using suitable amino-reactive chemistry, and/or C-terminal modification using suitable carboxy-reactive chemistry. In a further embodiment, said N-terminal or C-terminal modification comprises addition of an effector group, including but not limited to a cytotoxic agent, a radiochelator or a chromophore.

35

In a further embodiment, the modified derivative comprises an N-terminal modification. In a further embodiment, the N-terminal modification comprises an N-terminal acetyl group. In this embodiment, the N-terminal residue is capped with acetic anhydride or other appropriate reagents during peptide synthesis leading to a molecule which is N-terminally acetylated. This embodiment provides the advantage of removing a potential recognition point for aminopeptidases and avoids the potential for degradation of the bicyclic peptide.

In an alternative embodiment, the N-terminal modification comprises the addition of a molecular spacer group which facilitates the conjugation of effector groups and retention of potency of the bicyclic peptide to its target.

In a further embodiment, the modified derivative comprises a C-terminal modification. In a further embodiment, the C-terminal modification comprises an amide group. In this embodiment, the C-terminal residue is synthesized as an amide during peptide synthesis leading to a molecule which is C-terminally amidated. This embodiment provides the advantage of removing a potential recognition point for carboxypeptidase and reduces the potential for proteolytic degradation of the bicyclic peptide.

In one embodiment, the modified derivative comprises replacement of one or more amino acid residues with one or more non-natural amino acid residues. In this embodiment, non-natural amino acids may be selected having isosteric/isoelectronic side chains which are neither recognised by degradative proteases nor have any adverse effect upon target potency.

Alternatively, non-natural amino acids may be used having constrained amino acid side chains, such that proteolytic hydrolysis of the nearby peptide bond is conformationally and sterically impeded. In particular, these concern proline analogues, bulky sidechains, C α -disubstituted derivatives (for example, aminoisobutyric acid, Aib), and cyclo amino acids, a simple derivative being amino-cyclopropylcarboxylic acid.

In one embodiment, the modified derivative comprises the addition of a spacer group. In a further embodiment, the modified derivative comprises the addition of a spacer group to the N-terminal cysteine (C_i) and/or the C-terminal cysteine (C_{iii}).

In one embodiment, the modified derivative comprises replacement of one or more oxidation sensitive amino acid residues with one or more oxidation resistant amino acid residues. In a further embodiment, the modified derivative comprises replacement of a tryptophan residue

with a naphthylalanine or alanine residue. This embodiment provides the advantage of improving the pharmaceutical stability profile of the resultant bicyclic peptide ligand.

In one embodiment, the modified derivative comprises replacement of one or more charged
5 amino acid residues with one or more hydrophobic amino acid residues. In an alternative
embodiment, the modified derivative comprises replacement of one or more hydrophobic
amino acid residues with one or more charged amino acid residues. The correct balance of
charged versus hydrophobic amino acid residues is an important characteristic of the bicyclic
peptide ligands. For example, hydrophobic amino acid residues influence the degree of
10 plasma protein binding and thus the concentration of the free available fraction in plasma,
while charged amino acid residues (in particular arginine) may influence the interaction of the
peptide with the phospholipid membranes on cell surfaces. The two in combination may
influence half-life, volume of distribution and exposure of the peptide drug, and can be tailored
according to the clinical endpoint. In addition, the correct combination and number of charged
15 versus hydrophobic amino acid residues may reduce irritation at the injection site (if the
peptide drug has been administered subcutaneously).

In one embodiment, the modified derivative comprises replacement of one or more L-amino
acid residues with one or more D-amino acid residues. This embodiment is believed to
20 increase proteolytic stability by steric hindrance and by a propensity of D-amino acids to
stabilise α -turn conformations (Tugyi *et al* (2005) PNAS, 102(2), 413–418).

In one embodiment, the modified derivative comprises removal of any amino acid residues
and substitution with alanines, such as D-alanines. This embodiment provides the advantage
25 of identifying key binding residues and removing potential proteolytic attack site(s).

It should be noted that each of the above mentioned modifications serve to deliberately
improve the potency or stability of the peptide. Further potency improvements based on
modifications may be achieved through the following mechanisms:

- 30 - Incorporating hydrophobic moieties that exploit the hydrophobic effect and lead to
lower off rates, such that higher affinities are achieved;
- Incorporating charged groups that exploit long-range ionic interactions, leading to
faster on rates and to higher affinities (see for example Schreiber *et al*, *Rapid, electrostatically*
35 *assisted association of proteins* (1996), Nature Struct. Biol. 3, 427-31); and
- Incorporating additional constraint into the peptide, by for example constraining side

chains of amino acids correctly such that loss in entropy is minimal upon target binding, constraining the torsional angles of the backbone such that loss in entropy is minimal upon target binding and introducing additional cyclisations in the molecule for identical reasons.

- 5 (for reviews see Gentilucci *et al*, Curr. Pharmaceutical Design, (2010), 16, 3185-203, and Nestor *et al*, Curr. Medicinal Chem (2009), 16, 4399-418).

Isotopic variations

- The present invention includes all pharmaceutically acceptable (radio)isotope-labeled
10 peptide ligands of the invention, wherein one or more atoms are replaced by atoms having the same atomic number, but an atomic mass or mass number different from the atomic mass or mass number usually found in nature, and peptide ligands of the invention, wherein metal chelating groups are attached (termed "effector") that are capable of holding relevant (radio)isotopes, and peptide ligands of the invention, wherein certain functional groups are
15 covalently replaced with relevant (radio)isotopes or isotopically labelled functional groups.

- Examples of isotopes suitable for inclusion in the peptide ligands of the invention comprise isotopes of hydrogen, such as ^2H (D) and ^3H (T), carbon, such as ^{11}C , ^{13}C and ^{14}C , chlorine, such as ^{36}Cl , fluorine, such as ^{18}F , iodine, such as ^{123}I , ^{125}I and ^{131}I , nitrogen, such as ^{13}N and
20 ^{15}N , oxygen, such as ^{15}O , ^{17}O and ^{18}O , phosphorus, such as ^{32}P , sulfur, such as ^{35}S , copper, such as ^{64}Cu , gallium, such as ^{67}Ga or ^{68}Ga , yttrium, such as ^{90}Y and lutetium, such as ^{177}Lu , and Bismuth, such as ^{213}Bi .

- Certain isotopically-labelled peptide ligands of the invention, for example, those
25 incorporating a radioactive isotope, are useful in drug and/or substrate tissue distribution studies, and to clinically assess the presence and/or absence of the EphA2 target on diseased tissues. The peptide ligands of the invention can further have valuable diagnostic properties in that they can be used for detecting or identifying the formation of a complex between a labelled compound and other molecules, peptides, proteins, enzymes or
30 receptors. The detecting or identifying methods can use compounds that are labelled with labelling agents such as radioisotopes, enzymes, fluorescent substances, luminous substances (for example, luminol, luminol derivatives, luciferin, aequorin and luciferase), etc. The radioactive isotopes tritium, *i.e.* ^3H (T), and carbon-14, *i.e.* ^{14}C , are particularly useful for this purpose in view of their ease of incorporation and ready means of detection.

- 35 Substitution with heavier isotopes such as deuterium, *i.e.* ^2H (D), may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in*

vivo half-life or reduced dosage requirements, and hence may be preferred in some circumstances.

- 5 Substitution with positron emitting isotopes, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , can be useful in Positron Emission Topography (PET) studies for examining target occupancy.

- 10 Isotopically-labeled compounds of peptide ligands of the invention can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the accompanying Examples using an appropriate isotopically-labeled reagent in place of the non-labeled reagent previously employed.

Non-Aromatic Molecular scaffold

- 15 References herein to the term “non-aromatic molecular scaffold” refer to any molecular scaffold as defined herein which does not contain an aromatic (i.e. unsaturated) carbocyclic or heterocyclic ring system.

Suitable examples of non-aromatic molecular scaffolds are described in Heinis *et al* (2014) *Angewandte Chemie, International Edition* 53(6) 1602-1606.

- 20 As noted in the foregoing documents, the molecular scaffold may be a small molecule, such as a small organic molecule.

- 25 In one embodiment the molecular scaffold may be a macromolecule. In one embodiment the molecular scaffold is a macromolecule composed of amino acids, nucleotides or carbohydrates.

In one embodiment the molecular scaffold comprises reactive groups that are capable of reacting with functional group(s) of the polypeptide to form covalent bonds.

- 30 The molecular scaffold may comprise chemical groups which form the linkage with a peptide, such as amines, thiols, alcohols, ketones, aldehydes, nitriles, carboxylic acids, esters, alkenes, alkynes, azides, anhydrides, succinimides, maleimides, alkyl halides and acyl halides.

- 35 An example of an α unsaturated carbonyl containing compound is 1,1',1''-(1,3,5-triazinane-1,3,5-triyl)triprop-2-en-1-one (TATA) (*Angewandte Chemie, International Edition* (2014), 53(6), 1602-1606).

Effector and Functional Groups

According to a further aspect of the invention, there is provided a drug conjugate comprising a peptide ligand as defined herein conjugated to one or more effector and/or functional groups.

5

Effector and/or functional groups can be attached, for example, to the N and/or C termini of the polypeptide, to an amino acid within the polypeptide, or to the molecular scaffold.

Appropriate effector groups include antibodies and parts or fragments thereof. For instance, an effector group can include an antibody light chain constant region (CL), an antibody CH1 heavy chain domain, an antibody CH2 heavy chain domain, an antibody CH3 heavy chain domain, or any combination thereof, in addition to the one or more constant region domains. An effector group may also comprise a hinge region of an antibody (such a region normally being found between the CH1 and CH2 domains of an IgG molecule).

15

In a further embodiment of this aspect of the invention, an effector group according to the present invention is an Fc region of an IgG molecule. Advantageously, a peptide ligand-effector group according to the present invention comprises or consists of a peptide ligand Fc fusion having a $t_{1/2}$ half-life of a day or more, two days or more, 3 days or more, 4 days or more, 5 days or more, 6 days or more or 7 days or more. Most advantageously, the peptide ligand according to the present invention comprises or consists of a peptide ligand Fc fusion having a $t_{1/2}$ half-life of a day or more.

25

Functional groups include, in general, binding groups, drugs, reactive groups for the attachment of other entities, functional groups which aid uptake of the macrocyclic peptides into cells, and the like.

The ability of peptides to penetrate into cells will allow peptides against intracellular targets to be effective. Targets that can be accessed by peptides with the ability to penetrate into cells include transcription factors, intracellular signalling molecules such as tyrosine kinases and molecules involved in the apoptotic pathway. Functional groups which enable the penetration of cells include peptides or chemical groups which have been added either to the peptide or the molecular scaffold. Peptides such as those derived from such as VP22, HIV-Tat, a homeobox protein of *Drosophila* (*Antennapedia*), e.g. as described in Chen and Harrison, *Biochemical Society Transactions* (2007) Volume 35, part 4, p821; Gupta *et al.* in *Advanced Drug Discovery Reviews* (2004) Volume 57 9637. Examples of short peptides which have been shown to be efficient at translocation through plasma membranes include

35

the 16 amino acid penetratin peptide from *Drosophila Antennapedia* protein (Derossi *et al* (1994) J Biol. Chem. Volume 269 p10444), the 18 amino acid 'model amphipathic peptide' (Oehlke *et al* (1998) Biochim Biophys Acts Volume 1414 p127) and arginine rich regions of the HIV TAT protein. Non peptidic approaches include the use of small molecule mimics or
5 SMOCs that can be easily attached to biomolecules (Okuyama *et al* (2007) Nature Methods Volume 4 p153). Other chemical strategies to add guanidinium groups to molecules also enhance cell penetration (Elson-Scwab *et al* (2007) J Biol Chem Volume 282 p13585). Small molecular weight molecules such as steroids may be added to the molecular scaffold to enhance uptake into cells.

10

One class of functional groups which may be attached to peptide ligands includes antibodies and binding fragments thereof, such as Fab, Fv or single domain fragments. In particular, antibodies which bind to proteins capable of increasing the half-life of the peptide ligand *in vivo* may be used.

15

In one embodiment, a peptide ligand-effector group according to the invention has a $t_{1/2}$ half-life selected from the group consisting of: 12 hours or more, 24 hours or more, 2 days or more, 3 days or more, 4 days or more, 5 days or more, 6 days or more, 7 days or more, 8 days or more, 9 days or more, 10 days or more, 11 days or more, 12 days or more, 13 days
20 or more, 14 days or more, 15 days or more or 20 days or more. Advantageously a peptide ligand-effector group or composition according to the invention will have a $t_{1/2}$ range 12 to 60 hours. In a further embodiment, it will have a $t_{1/2}$ half-life of a day or more. In a further embodiment still, it will be in the range 12 to 26 hours.

25

In one particular embodiment of the invention, the functional group is selected from a metal chelator, which is suitable for complexing metal radioisotopes of medicinal relevance.

Possible effector groups also include enzymes, for instance such as carboxypeptidase G2 for use in enzyme/prodrug therapy, where the peptide ligand replaces antibodies in ADEPT.


30

In one particular embodiment of the invention, the functional group is selected from a drug, such as a cytotoxic agent for cancer therapy. Suitable examples include: alkylating agents such as cisplatin and carboplatin, as well as oxaliplatin, mechlorethamine, cyclophosphamide, chlorambucil, ifosfamide; Anti-metabolites including purine analogs
35 azathioprine and mercaptopurine or pyrimidine analogs; plant alkaloids and terpenoids including vinca alkaloids such as Vincristine, Vinblastine, Vinorelbine and Vindesine; Podophyllotoxin and its derivatives etoposide and teniposide; Taxanes, including paclitaxel,


originally known as Taxol; topoisomerase inhibitors including camptothecins: irinotecan and topotecan, and type II inhibitors including amsacrine, etoposide, etoposide phosphate, and teniposide. Further agents can include antitumour antibiotics which include the immunosuppressant dactinomycin (which is used in kidney transplantations), doxorubicin,
 5 epirubicin, bleomycin, calicheamycins, and others.

In one further particular embodiment of the invention, the cytotoxic agent is selected from maytansinoids (such as DM1) or monomethyl auristatins (such as MMAE).

- 10 DM1 is a cytotoxic agent which is a thiol-containing derivative of maytansine and has the following structure:



Monomethyl auristatin E (MMAE) is a synthetic antineoplastic agent and has the following structure:



In one yet further particular embodiment of the invention, the cytotoxic agent is selected from maytansinoids (such as DM1). Data is presented herein in Table 6 which demonstrates the effects of peptide ligands conjugated to toxins containing DM1.

- 5 In one embodiment, the cytotoxic agent is linked to the bicyclic peptide by a cleavable bond, such as a disulphide bond or a protease sensitive bond. In a further embodiment, the groups adjacent to the disulphide bond are modified to control the hindrance of the disulphide bond, and by this the rate of cleavage and concomitant release of cytotoxic agent.
- 10 Published work established the potential for modifying the susceptibility of the disulphide bond to reduction by introducing steric hindrance on either side of the disulphide bond (Kellogg *et al* (2011) Bioconjugate Chemistry, 22, 717). A greater degree of steric hindrance reduces the rate of reduction by intracellular glutathione and also extracellular (systemic) reducing agents, consequentially reducing the ease by which toxin is released, both inside
- 15 and outside the cell. Thus, selection of the optimum in disulphide stability in the circulation (which minimises undesirable side effects of the toxin) versus efficient release in the intracellular milieu (which maximises the therapeutic effect) can be achieved by careful selection of the degree of hindrance on either side of the disulphide bond.
- 20 The hindrance on either side of the disulphide bond is modulated through introducing one or more methyl groups on either the targeting entity (here, the bicyclic peptide) or toxin side of the molecular construct.

In one embodiment, the drug conjugate additionally comprises a linker between said peptide
25 ligand and said cytotoxic agents.

In one embodiment, the cytotoxic agent and linker is selected from any combinations of those described in WO 2016/067035 (the cytotoxic agents and linkers thereof are herein
30 incorporated by reference).

In one embodiment the cytotoxic agent is selected from DM1 or MMAE.


In one embodiment, the linker between said cytotoxic agent and said bicyclic peptide comprises one or more amino acid residues. Thus, in one embodiment, the cytotoxic agent
35 is MMAE and the linker is selected from: -Val-Cit-, -Trp-Cit-, -Val-Lys-, -D-Trp-Cit-, -Ala-Ala-Asn-, D-Ala-Phe-Lys- or -Glu-Pro-Cit-Gly-hPhe-Tyr-Leu- (SEQ ID NO: 98). In a further embodiment, the cytotoxic agent is MMAE and the linker is selected from: -Val-Cit-, -Trp-Cit-,

-Val-Lys- or -D-Trp-Cit-. In a yet further embodiment, the cytotoxic agent is MMAE and the linker is -Val-Cit- or -Val-Lys-. In a still yet further embodiment, the cytotoxic agent is MMAE and the linker is -Val-Cit-.

- 5 In an alternative embodiment, the linker between said cytotoxic agent comprises a disulfide bond, such as a cleavable disulfide bond. Thus, in a further embodiment, the cytotoxic agent is DM1 and the linker is selected from: -S-S-, -SS(SO₃H)-, -SS-(Me)-, -(Me)-SS-(Me)-, -SS-(Me₂)- or -SS-(Me)-SO₃H-. In a further embodiment, the cytotoxic agent is DM1 and the linker comprises an -S-S- moiety, such as (N-succinimidyl 3-(2-pyridyldithio)propionate
- 10 (SPDB), or an -SS(SO₃H)- moiety, such as SO₃H-SPDB.

- In an alternative embodiment, the cytotoxic agent comprises a non-cleavable cytotoxic agent. Thus, in one embodiment the cytotoxic agent is non-cleavable MMAE (such as the cytotoxic agent within BCY6063) or non-cleavable DM1 (such as the cytotoxic agent within
- 15 BCY6064).

In one embodiment, the cytotoxic agent is DM1 and the drug conjugate comprises a compound of formula (A):




20

(A)

wherein said bicycle is selected from any one of BCY6099 and BCY6014 as defined herein.

- In an alternative embodiment, the cytotoxic agent is DM1 and the drug conjugate comprises
- 25 a compound of formula (B):



(B)

wherein said bicycle is selected from any one of BCY6099 and BCY6014 as defined herein.

- 5 In an alternative embodiment, the cytotoxic agent is DM1 and the drug conjugate comprises a compound of formula (A), wherein said bicycle is selected from BCY6099 as defined herein. This BDC is known herein as BCY6027. Data is presented herein which demonstrates excellent competition binding for BCY6027 in the EphA2 competition binding assay as shown in Table 6.

10

In an alternative embodiment, the cytotoxic agent is DM1 and the drug conjugate comprises a compound of formula (B), wherein said bicycle is selected from BCY6099 as defined herein. This BDC is known herein as BCY6028. Data is presented herein which demonstrates excellent competition binding for BCY6028 in the EphA2 competition binding

15 assay as shown in Table 6.

In an alternative embodiment, the cytotoxic agent is DM1 and the drug conjugate comprises a compound of formula (A), wherein said bicycle is selected from BCY6014 as defined herein. This BDC is known herein as BCY6031. Data is presented herein which demonstrates

20 excellent competition binding for BCY6031 in the EphA2 competition binding assay as shown in Table 6. Data is also presented herein in Table 11 and Figures 1 and 2 which demonstrate that BCY6031 treatment completely eradicated non-small cell lung carcinomas from day 32 and no tumour regrowth occurred following dosing suspension on day 28.

- 25 In an alternative embodiment, the cytotoxic agent is DM1 and the drug conjugate comprises a compound of formula (B), wherein said bicycle is selected from BCY6014 as defined herein. This BDC is known herein as BCY6032. Data is presented herein which


demonstrates excellent competition binding for BCY6032 in the EphA2 competition binding assay as shown in Table 6.

5 In an alternative embodiment, the cytotoxic agent is MMAE or DM1 and the drug conjugate is selected from any of the BDCs listed in Table 11. Data is presented herein which shows that these BDCs exhibited excellent cross reactivity between human, mouse and rodent EphA2 as shown in Table 11.


10 In a further embodiment, the cytotoxic agent is MMAE or DM1 and the drug conjugate is selected from any of the BDCs listed in Table 13.


15 In a further embodiment, the cytotoxic agent is MMAE or DM1 and the drug conjugate is selected from BCY6033, BCY6082, BCY6136 and BCY6173. Data is presented herein which shows that these four Bicycle Drug Conjugates exhibited no significant binding to: closely related human homologs EphA1, EphA3, EphA4, EphA5, EphA6, EphA7 and EphB4; mouse EphA3 and EphA4; and rat EphA3 and EphB1 as shown in Tables 14 and 15.


In a yet further embodiment, the drug conjugate is selected from any one of: BCY6031, BCY6033, BCY6082, BCY6135, BCY6136, BCY6173, BCY6174 and BCY6175:
20





BCY6031

















In one embodiment, the drug conjugate is other than BCY6027, BCY6028, BCY6135, BCY6136, BCY6173, BCY6174 and BCY6175.

In a still yet further embodiment, the drug conjugate is BCY6136. Data is presented herein in
5 Studies 7 and 8 which show that BCY6136 showed significant and potent anti-tumor activity in the PC-3 xenograft prostate cancer model (see Figures 7 to 10 and Tables 21 to 24). Data is also provided herein which show that BCY6136 demonstrated potent antitumor activity in the NCI-H1975 xenograft lung cancer (NSCLC) model (see Figures 11 to 13 and Tables 25 to 30). Data is also presented herein in Studies 10 and 11 which show that BCY6136
10 demonstrated potent anti-tumor effect in both large and small tumour size LU-01-0251 PDX lung cancer (NSCLC) models (see Figures 14 and 15 and Tables 31 to 34) wherein complete tumor regression was observed. Data is also presented herein in Study 12 which show that BCY6136 demonstrated significant anti-tumor effect in the LU-01-0046 PDX lung cancer (NSCLC) model (see Figure 16 and Tables 35 and 36) wherein complete tumor
15 regression was observed for BCY6136. Data is also presented herein in Study 13 which show that BCY6136 demonstrated dose dependent anti-tumor activity in the LU-01-0046 PDX lung cancer (NSCLC) model (see Figure 17 and Tables 37 and 38). Data is also presented herein in Study 14 which show BCY6136 eradicated tumors in the LU-01-0046 PDX lung cancer (NSCLC) model (see Figures 18 to 22 and Tables 39 to 42). Data is also
20 presented herein in Studies 15 and 16 which demonstrate the effects of BCY6136 in two models which make use of cell lines with low/negligible EphA2 expression (namely Lu-01-0412 and Lu-01-0486). This data is shown in Figures 23 and 24 and Tables 43 to 46 and demonstrate that BCY6136 had no effect upon tumor regression in either cell line but BCYs BCY8245 and BCY8781, which bind to a target highly expressed in the Lu-01-0412 cell line,
25 completely eradicated the tumour. Data is presented herein in Study 17 which show that BCY6136 demonstrated potent antitumor activity in the MDA-MB-231 xenograft breast cancer model (see Figures 25 to 27 and Tables 47 to 50). Data is also presented herein in Study 18 which demonstrates the effects of BCY6136 in a breast cancer model which makes use of a cell line with low/negligible EphA2 expression (namely EMT6). This data is shown in
30 Figure 28 and Tables 51 and 52 and demonstrates that BCY6136 had no effect upon tumor regression in this cell line. Data is also presented herein in Study 19 which show that BCY6136 demonstrated significant antitumor activity in the NCI-N87 xenograft gastric cancer model (see Figure 29 and Tables 53 and 54). Data is also presented herein in Study 20 which show that BCY6136 demonstrated significant antitumor activity in the SK-OV-3
35 xenograft ovarian cancer model (see Figure 30 and Tables 55 and 56) compared with the ADC MEDI-547 which demonstrated moderate antitumour activity. Data is also presented herein in Study 21 which show that BCY6136 demonstrated significant antitumor activity in

the OE-21 xenograft oesophageal cancer model (see Figure 31 and Tables 57 and 58). Data is also presented herein in Study 22 which show that BCY6136 demonstrated dose-dependent antitumor activity in the MOLP-8 xenograft multiple myeloma model and BCY6082 demonstrated significant antitumor activity (see Figures 32 and 33 and Tables 59 and 60). Data is also presented herein in Study 23 which show that BCY6136 demonstrated potent antitumor activity in the HT-1080 xenograft fibrosarcoma model (see Figures 34 to 41 and Tables 61 and 62).

Synthesis

The peptides of the present invention may be manufactured synthetically by standard techniques followed by reaction with a molecular scaffold *in vitro*. When this is performed, standard chemistry may be used. This enables the rapid large scale preparation of soluble material for further downstream experiments or validation. Such methods could be accomplished using conventional chemistry such as that disclosed in Timmerman *et al* (*supra*).

Thus, the invention also relates to manufacture of polypeptides or conjugates selected as set out herein, wherein the manufacture comprises optional further steps as explained below. In one embodiment, these steps are carried out on the end product polypeptide/conjugate made by chemical synthesis.

Optionally amino acid residues in the polypeptide of interest may be substituted when manufacturing a conjugate or complex.

Peptides can also be extended, to incorporate for example another loop and therefore introduce multiple specificities.

To extend the peptide, it may simply be extended chemically at its N-terminus or C-terminus or within the loops using orthogonally protected lysines (and analogues) using standard solid phase or solution phase chemistry. Standard (bio)conjugation techniques may be used to introduce an activated or activatable N- or C-terminus. Alternatively additions may be made by fragment condensation or native chemical ligation e.g. as described in (Dawson *et al*. 1994. Synthesis of Proteins by Native Chemical Ligation. Science 266:776-779), or by enzymes, for example using subtiligase as described in (Chang *et al* Proc Natl Acad Sci U S A. 1994 Dec 20; 91(26):12544-8 or in Hikari *et al* Bioorganic & Medicinal Chemistry Letters Volume 18, Issue 22, 15 November 2008, Pages 6000-6003).

Alternatively, the peptides may be extended or modified by further conjugation through disulphide bonds. This has the additional advantage of allowing the first and second peptide to dissociate from each other once within the reducing environment of the cell. In this case, the molecular scaffold could be added during the chemical synthesis of the first peptide so
5 as to react with the three cysteine groups; a further cysteine or thiol could then be appended to the N or C-terminus of the first peptide, so that this cysteine or thiol only reacted with a free cysteine or thiol of the second peptide, forming a disulfide –linked bicyclic peptide-peptide conjugate.

10 Similar techniques apply equally to the synthesis/coupling of two bicyclic and bispecific macrocycles, potentially creating a tetraspecific molecule.

Furthermore, addition of other functional groups or effector groups may be accomplished in the same manner, using appropriate chemistry, coupling at the N- or C-termini or via side
15 chains. In one embodiment, the coupling is conducted in such a manner that it does not block the activity of either entity.

Pharmaceutical Compositions

According to a further aspect of the invention, there is provided a pharmaceutical composition
20 comprising a peptide ligand or a drug conjugate as defined herein in combination with one or more pharmaceutically acceptable excipients.

Generally, the present peptide ligands will be utilised in purified form together with pharmacologically appropriate excipients or carriers. Typically, these excipients or carriers
25 include aqueous or alcoholic/aqueous solutions, emulsions or suspensions, including saline and/or buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride and lactated Ringer's. Suitable physiologically-acceptable adjuvants, if necessary to keep a polypeptide complex in suspension, may be chosen from thickeners such as carboxymethylcellulose, polyvinylpyrrolidone, gelatin and
30 alginates.

Intravenous vehicles include fluid and nutrient replenishers and electrolyte replenishers, such as those based on Ringer's dextrose. Preservatives and other additives, such as antimicrobials, antioxidants, chelating agents and inert gases, may also be present (Mack
35 (1982) Remington's Pharmaceutical Sciences, 16th Edition).

- The peptide ligands of the present invention may be used as separately administered compositions or in conjunction with other agents. These can include antibodies, antibody fragments and various immunotherapeutic drugs, such as cyclosporine, methotrexate, adriamycin or cisplatin and immunotoxins. Pharmaceutical compositions can include
- 5 "cocktails" of various cytotoxic or other agents in conjunction with the protein ligands of the present invention, or even combinations of selected polypeptides according to the present invention having different specificities, such as polypeptides selected using different target ligands, whether or not they are pooled prior to administration.
- 10 The route of administration of pharmaceutical compositions according to the invention may be any of those commonly known to those of ordinary skill in the art. For therapy, the peptide ligands of the invention can be administered to any patient in accordance with standard techniques. The administration can be by any appropriate mode, including parenterally, intravenously, intramuscularly, intraperitoneally, transdermally, via the pulmonary route, or
- 15 also, appropriately, by direct infusion with a catheter. Preferably, the pharmaceutical compositions according to the invention will be administered by inhalation. The dosage and frequency of administration will depend on the age, sex and condition of the patient, concurrent administration of other drugs, counterindications and other parameters to be taken into account by the clinician.
- 20 The peptide ligands of this invention can be lyophilised for storage and reconstituted in a suitable carrier prior to use. This technique has been shown to be effective and art-known lyophilisation and reconstitution techniques can be employed. It will be appreciated by those skilled in the art that lyophilisation and reconstitution can lead to varying degrees of activity
- 25 loss and that levels may have to be adjusted upward to compensate.
- The compositions containing the present peptide ligands or a cocktail thereof can be administered for prophylactic and/or therapeutic treatments. In certain therapeutic applications, an adequate amount to accomplish at least partial inhibition, suppression,
- 30 modulation, killing, or some other measurable parameter, of a population of selected cells is defined as a "therapeutically-effective dose". Amounts needed to achieve this dosage will depend upon the severity of the disease and the general state of the patient's own immune system, but generally range from 0.005 to 5.0 mg of selected peptide ligand per kilogram of body weight, with doses of 0.05 to 2.0 mg/kg/dose being more commonly used. For
- 35 prophylactic applications, compositions containing the present peptide ligands or cocktails thereof may also be administered in similar or slightly lower dosages.

A composition containing a peptide ligand according to the present invention may be utilised in prophylactic and therapeutic settings to aid in the alteration, inactivation, killing or removal of a select target cell population in a mammal. In addition, the peptide ligands described herein may be used extracorporeally or *in vitro* selectively to kill, deplete or otherwise
5 effectively remove a target cell population from a heterogeneous collection of cells. Blood from a mammal may be combined extracorporeally with the selected peptide ligands whereby the undesired cells are killed or otherwise removed from the blood for return to the mammal in accordance with standard techniques.

10 ***Therapeutic Uses***

The bicyclic peptides of the invention have specific utility as EphA2 binding agents.

Eph receptor tyrosine kinases (Ephs) belong to a large group of receptor tyrosine kinases (RTKs), kinases that phosphorylate proteins on tyrosine residues. Ephs and their membrane
15 bound ephrin ligands (ephrins) control cell positioning and tissue organization (Poliakov *et al.* (2004) *Dev Cell* 7, 465-80). Functional and biochemical Eph responses occur at higher ligand oligomerization states (Stein *et al.* (1998) *Genes Dev* 12, 667-678).

Among other patterning functions, various Ephs and ephrins have been shown to play a role
20 in vascular development. Knockout of EphB4 and ephrin-B2 results in a lack of the ability to remodel capillary beds into blood vessels (Poliakov *et al.*, *supra*) and embryonic lethality. Persistent expression of some Eph receptors and ephrins has also been observed in newly-formed, adult micro-vessels (Brantley-Sieders *et al.* (2004) *Curr Pharm Des* 10, 3431-42; Adams (2003) *J Anat* 202, 105-12).

25 The de-regulated re-emergence of some ephrins and their receptors in adults also has been observed to contribute to tumor invasion, metastasis and neo-angiogenesis (Nakamoto *et al.* (2002) *Microsc Res Tech* 59, 58-67; Brantley-Sieders *et al.*, *supra*). Furthermore, some Eph family members have been found to be over-expressed on tumor cells from a variety of
30 human tumors (Brantley-Sieders *et al.*, *supra*); Marme (2002) *Ann Hematol* 81 Suppl 2, S66; Booth *et al.* (2002) *Nat Med* 8, 1360-1).

EPH receptor A2 (ephrin type-A receptor 2) is a protein that in humans is encoded by the
35 *EPHA2* gene.

EphA2 is upregulated in multiple cancers in man, often correlating with disease progression, metastasis and poor prognosis e.g.: breast (Zelinski *et al* (2001) *Cancer Res.* 61, 2301–

2306; Zhuang *et al* (2010) *Cancer Res.* 70, 299–308; Brantley-Sieders *et al* (2011) *PLoS One* 6, e24426), lung (Brannan *et al* (2009) *Cancer Prev Res (Phila)* 2, 1039–1049; Kinch *et al* (2003) *Clin Cancer Res.* 9, 613–618; Guo *et al* (2013) *J Thorac Oncol.* 8, 301–308), gastric (Nakamura *et al* (2005) *Cancer Sci.* 96, 42–47; Yuan *et al* (2009) *Dig Dis Sci* 54, 2410–2417),
 5 pancreatic (Mudali *et al* (2006) *Clin Exp Metastasis* 23, 357–365), prostate (Walker-Daniels *et al* (1999) *Prostate* 41, 275–280), liver (Yang *et al* (2009) *Hepatol Res.* 39, 1169–1177) and glioblastoma (Wykosky *et al* (2005) *Mol Cancer Res.* 3, 541–551; Li *et al* (2010) *Tumour Biol.* 31, 477–488).

- 10 The full role of EphA2 in cancer progression is still not defined although there is evidence for interaction at numerous stages of cancer progression including tumour cell growth, survival, invasion and angiogenesis. Downregulation of EphA2 expression suppresses tumour cancer cell propagation (Binda *et al* (2012) *Cancer Cell* 22, 765–780), whilst EphA2 blockade inhibits VEGF induced cell migration (Hess *et al* (2001) *Cancer Res.* 61, 3250–3255),
 15 sprouting and angiogenesis (Cheng *et al* (2002) *Mol Cancer Res.* 1, 2–11; Lin *et al* (2007) *Cancer* 109, 332–40) and metastatic progression (Brantley-Sieders *et al* (2005) *FASEB J.* 19, 1884–1886).

An antibody drug conjugate to EphA2 has been shown to significantly diminish tumour
 20 growth in rat and mouse xenograft models (Jackson *et al* (2008) *Cancer Research* 68, 9367–9374) and a similar approach has been tried in man although treatment had to be discontinued for treatment related adverse events (Annunziata *et al* (2013) *Invest New drugs* 31, 77–84).

- 25 Polypeptide ligands selected according to the method of the present invention may be employed in *in vivo* therapeutic and prophylactic applications, *in vitro* and *in vivo* diagnostic applications, *in vitro* assay and reagent applications, and the like. Ligands having selected levels of specificity are useful in applications which involve testing in non-human animals, where cross-reactivity is desirable, or in diagnostic applications, where cross-reactivity with
 30 homologues or paralogues needs to be carefully controlled. In some applications, such as vaccine applications, the ability to elicit an immune response to predetermined ranges of antigens can be exploited to tailor a vaccine to specific diseases and pathogens.

Substantially pure peptide ligands of at least 90 to 95% homogeneity are preferred for
 35 administration to a mammal, and 98 to 99% or more homogeneity is most preferred for pharmaceutical uses, especially when the mammal is a human. Once purified, partially or to homogeneity as desired, the selected polypeptides may be used diagnostically or

therapeutically (including extracorporeally) or in developing and performing assay procedures, immunofluorescent stainings and the like (Lefkovite and Pernis, (1979 and 1981) Immunological Methods, Volumes I and II, Academic Press, NY).

- 5 According to a further aspect of the invention, there is provided a peptide ligand or a drug conjugate as defined herein, for use in preventing, suppressing or treating a disease or disorder characterised by overexpression of EphA2 in diseased tissue (such as a tumour).

- 10 According to a further aspect of the invention, there is provided a method of preventing, suppressing or treating a disease or disorder characterised by overexpression of EphA2 in diseased tissue (such as a tumour), which comprises administering to a patient in need thereof an effector group and drug conjugate of the peptide ligand as defined herein.

- 15 In one embodiment, the EphA2 is mammalian EphA2. In a further embodiment, the mammalian EphA2 is human EphA2.

In one embodiment, the disease or disorder characterised by overexpression of EphA2 in diseased tissue is selected from cancer.

- 20 Examples of cancers (and their benign counterparts) which may be treated (or inhibited) include, but are not limited to tumours of epithelial origin (adenomas and carcinomas of various types including adenocarcinomas, squamous carcinomas, transitional cell carcinomas and other carcinomas) such as carcinomas of the bladder and urinary tract, breast, gastrointestinal tract (including the esophagus, stomach (gastric), small intestine, colon, rectum and anus), liver (hepatocellular carcinoma), gall bladder and biliary system, exocrine pancreas, kidney, lung (for example adenocarcinomas, small cell lung carcinomas, non-small cell lung carcinomas, bronchioalveolar carcinomas and mesotheliomas), head and neck (for example cancers of the tongue, buccal cavity, larynx, pharynx, nasopharynx, tonsil, salivary glands, nasal cavity and paranasal sinuses), ovary, fallopian tubes, peritoneum, vagina, vulva, penis, cervix, myometrium, endometrium, thyroid (for example thyroid follicular carcinoma), adrenal, prostate, skin and adnexae (for example melanoma, basal cell carcinoma, squamous cell carcinoma, keratoacanthoma, dysplastic naevus); haematological malignancies (i.e. leukemias, lymphomas) and premalignant haematological disorders and disorders of borderline malignancy including haematological malignancies and related conditions of lymphoid lineage (for example acute lymphocytic leukemia [ALL], chronic lymphocytic leukemia [CLL], B-cell lymphomas such as diffuse large B-cell lymphoma [DLBCL], follicular lymphoma, Burkitt's lymphoma, mantle cell lymphoma, T-cell lymphomas

and leukaemias, natural killer [NK] cell lymphomas, Hodgkin's lymphomas, hairy cell leukaemia, monoclonal gammopathy of uncertain significance, plasmacytoma, multiple myeloma, and post-transplant lymphoproliferative disorders), and haematological malignancies and related conditions of myeloid lineage (for example acute

5 myelogenousleukemia [AML], chronic myelogenousleukemia [CML], chronic myelomonocytic leukemia [CMML], hypereosinophilic syndrome, myeloproliferative disorders such as polycythaemia vera, essential thrombocythaemia and primary myelofibrosis, myeloproliferative syndrome, myelodysplastic syndrome, and promyelocytic leukemia);

10 tumours of mesenchymal origin, for example sarcomas of soft tissue, bone or cartilage such as osteosarcomas, fibrosarcomas, chondrosarcomas, rhabdomyosarcomas, leiomyosarcomas, liposarcomas, angiosarcomas, Kaposi's sarcoma, Ewing's sarcoma, synovial sarcomas, epithelioid sarcomas, gastrointestinal stromal tumours, benign and malignant histiocytomas, and dermatofibrosarcomaprotuberans; tumours of the central or peripheral nervous system (for example astrocytomas, gliomas and glioblastomas,

15 meningiomas, ependymomas, pineal tumours and schwannomas); endocrine tumours (for example pituitary tumours, adrenal tumours, islet cell tumours, parathyroid tumours, carcinoid tumours and medullary carcinoma of the thyroid); ocular and adnexal tumours (for example retinoblastoma); germ cell and trophoblastic tumours (for example teratomas, seminomas, dysgerminomas, hydatidiform moles and choriocarcinomas); and paediatric and

20 embryonal tumours (for example medulloblastoma, neuroblastoma, Wilms tumour, and primitive neuroectodermal tumours); or syndromes, congenital or otherwise, which leave the patient susceptible to malignancy (for example Xeroderma Pigmentosum).

In a further embodiment, the cancer is selected from: breast cancer, lung cancer, gastric

25 cancer, pancreatic cancer, prostate cancer, liver cancer, glioblastoma and angiogenesis.

In a further embodiment, the cancer is selected from: prostate cancer, lung cancer (such as non-small cell lung carcinomas (NSCLC)), breast cancer (such as triple negative breast cancer), gastric cancer, ovarian cancer, oesophageal cancer, multiple myeloma and

30 fibrosarcoma.

In a yet further embodiment, the cancer is prostate cancer. Data is presented herein in Studies 7 and 8 which show that BCY6033 and BCY6136 showed significant and potent anti-tumor activity in the PC-3 xenograft prostate cancer model (see Figures 7 to 10 and

35 Tables 21 to 24).

In a yet further embodiment, the drug conjugate is useful for preventing, suppressing or treating solid tumours such as fibrosarcomas and breast, and non-small cell lung carcinomas.

- 5 In a yet further embodiment, the cancer is selected from lung cancer, such as non-small cell lung carcinomas (NSCLC). Data is presented herein which demonstrates that a BDC of the invention (BCY6031) completely eradicated non-small cell lung carcinomas from day 32 and no tumour regrowth occurred following dosing suspension on day 28. This data clearly demonstrates the clinical utility of the BDCs of the present invention in cancers such as lung
- 10 cancers, in particular non-small cell lung carcinomas. Data is also presented herein in Study 9 which show that BCY6033 demonstrated dose dependent anti-tumor activity, BCY6082 demonstrated significant antitumor activity and BCY6136 demonstrated potent antitumor activity in the NCI-H1975 xenograft lung cancer (NSCLC) model (see Figures 11 to 13 and Tables 25 to 30). Data is also presented herein in Studies 10 and 11 which show that
- 15 BCY6136 demonstrated potent anti-tumor effect in both large and small tumour size LU-01-0251 PDX lung cancer (NSCLC) models (see Figures 14 and 15 and Tables 31 to 34) wherein complete tumor regression was observed. Data is also presented herein in Study 12 which show that BCY6033, BCY6136, BCY6082 and BCY6031 demonstrated significant anti-tumor effect in the LU-01-0046 PDX lung cancer (NSCLC) model (see Figure 16 and
- 20 Tables 35 and 36) wherein complete tumor regression was observed for BCY6033 and BCY6136. Data is also presented herein in Study 13 which show that BCY6136 demonstrated dose dependent anti-tumor activity in the LU-01-0046 PDX lung cancer (NSCLC) model (see Figure 17 and Tables 37 and 38). Data is also presented herein in Study 14 which show that BCY6082 demonstrated dose dependent antitumor activity,
- 25 BCY6031 and BCY6173 demonstrated antitumor activity and BCY6033, BCY6136 and BCY6175 eradicated tumors in the LU-01-0046 PDX lung cancer (NSCLC) model (see Figures 18 to 22 and Tables 39 to 42). Data is also presented herein in Studies 15 and 16 which demonstrate the effects of BCY6136 in two models which make use of cell lines with low/negligible EphA2 expression (namely Lu-01-0412 and Lu-01-0486). This data is shown
- 30 in Figures 23 and 24 and Tables 43 to 46 and demonstrate that BCY6136 had no effect upon tumor regression in either cell line but BCYs BCY8245 and BCY8781, which bind to a target highly expressed in the Lu-01-0412 cell line, completely eradicated the tumour. In a further embodiment, the cancer is breast cancer. In a yet further embodiment, the breast cancer is triple negative breast cancer. Data is presented herein in Study 17 which show that
- 35 BCY6082 demonstrated anti-tumor activity, BCY6033 demonstrated dose dependent antitumor activity and BCY6136 demonstrated potent antitumor activity in the MDA-MB-231 xenograft breast cancer model (see Figures 25 to 27 and Tables 47 to 50). Data is also

presented herein in Study 18 which demonstrates the effects of BCY6136 in a breast cancer model which makes use of a cell line with low/negligible EphA2 expression (namely EMT6). This data is shown in Figure 28 and Tables 51 and 52 and demonstrates that BCY6136 had no effect upon tumor regression in this cell line. In an alternative embodiment, the breast cancer is Herceptin resistant breast cancer. Without being bound by theory, EphA2 is believed to be implicated in the resistance to Herceptin, therefore, an EphA2-targeting entity has potential utility in patients who have failed to respond to Herceptin.

In a further embodiment, the cancer is gastric cancer. Data is presented herein in Study 19 which show that BCY6136 demonstrated significant antitumor activity in the NCI-N87 xenograft gastric cancer model (see Figure 29 and Tables 53 and 54).

In a further embodiment, the cancer is ovarian cancer. Data is presented herein in Study 20 which show that BCY6136 demonstrated significant antitumor activity in the SK-OV-3 xenograft ovarian cancer model (see Figure 30 and Tables 55 and 56) compared with the ADC MEDI-547 which demonstrated moderate antitumour activity.

In a further embodiment, the cancer is oesophageal cancer. Data is presented herein in Study 21 which show that BCY6136 demonstrated significant antitumor activity in the OE-21 xenograft oesophageal cancer model (see Figure 31 and Tables 57 and 58).

In a further embodiment, the cancer is multiple myeloma. Data is presented herein in Study 22 which show that BCY6136 demonstrated dose-dependent antitumor activity in the MOLP-8 xenograft multiple myeloma model and BCY6082 demonstrated significant antitumor activity (see Figures 32 and 33 and Tables 59 and 60).

In a further embodiment, the cancer is fibrosarcoma. Data is presented herein in Study 23 which show that BCY6173, BCY6135, BCY6174 and BCY6175 demonstrated dose dependent antitumor activity and BCY6082, BCY6031, BCY6033 and BCY6136 demonstrated potent antitumor activity in the HT-1080 xenograft fibrosarcoma model (see Figures 34 to 41 and Tables 61 and 62).

References herein to the term "prevention" involves administration of the protective composition prior to the induction of the disease. "Suppression" refers to administration of the composition after an inductive event, but prior to the clinical appearance of the disease. "Treatment" involves administration of the protective composition after disease symptoms become manifest.

Animal model systems which can be used to screen the effectiveness of the peptide ligands in protecting against or treating the disease are available. The use of animal model systems is facilitated by the present invention, which allows the development of polypeptide ligands
5 which can cross react with human and animal targets, to allow the use of animal models.

Furthermore, data is presented herein which demonstrates an association between copy number variation (CNV) and gene expression for EphA2 from multiple tumor types. Thus, according to a further aspect of the invention, there is provided a method of preventing,
10 suppressing or treating cancer, which comprises administering to a patient in need thereof an effector group and drug conjugate of the peptide ligand as defined herein, wherein said patient is identified as having an increased copy number variation (CNV) of EphA2.

In one embodiment, the cancer is selected from those identified herein as having increased
15 CNV of EphA2. In a further embodiment, the cancer is breast cancer.

The invention is further described below with reference to the following examples.

Examples

20

Abbreviations	Name	Precursor Name	Precursor or CAS	Supplier
1Nal	1-Naphthylalanine	Fmoc-3-(1-naphthyl)-L-alanine	96402-49-2	Fluorochem
2FuAla	2-Furylalanine	Fmoc-L-2-furylalanine	159611-02-6	Combi Blocks
2Nal	2-Naphthylalanine	Fmoc-3-(2-naphthyl)-L-alanine	112883-43-9	Alfa Aesar
3,3-DPA	3,3-Diphenylalanine	fmoc-3,3-diphenylalanine	189937-46-0	Alfa Aesar
3,4-DCPhe	3,4-Dichlorophenylalanine	Fmoc-3,4-dichloro-L-phenylalanine	17766-59-5	PolyPeptide
3Pal	3-(3-Pyridyl)-Alanine	N-Fmoc-3-(3-pyridyl)-L-βnine	175453-07-3	Fluorochem
4,4-BPA	4,4'-Biphenylalanine	Fmoc-L-4, 4'-Biphenylalanine	199110-64-0	Alfa Aesar
4BenzylPro	4-Benzyl-pyrrolidine-2-carboxylic acid	Fmoc-4-Benzyl-pyrrolidine-2-carboxylic acid		PolyPeptide
4BrPhe	4-Bromophenylalanine	Fmoc-4-Bromo-L-phenylalanine	198561-04-5	PolyPeptide
4FIPro	4-Fluoro-pyrrolidine-2-carboxylic acid	Fmoc-4-fluoro-pyrrolidine-2-carboxylic acid	203866-19-7	PolyPeptide
4MeoPhe	4-Methoxyphenylalanine	Fmoc-4-Methoxyphenylalanine	77128-72-4	Iris Biotech
4Pal	3-(4-Pyridyl)-Alanine	N-Fmoc-3-(4-pyridyl)-L-alanine	169555-95-7	Fluorochem
4PhenylPro	4-Phenyl-pyrrolidine-2-carboxylic acid	Fmoc-4-phenyl-pyrrolidine-2-carboxylic acid	269078-71-9	Cambridge Bioscience
Ac	Acetyl			

AC3C	1-Aminocyclopropane-1-carboxylic acid	1-(Fmoc-amino)cyclopropanecarboxylic acid	126705-22-4	Iris Biotech
AC4C	1-Amino-1-cyclobutanecarboxylic acid	1-(Fmoc-amino)-cyclobutylcarboxylic acid	885951-77-9	Fluorochem
AC5C	1-Amino-1-cyclopentanecarboxylic acid	1-(Fmoc-amino)cyclopentanecarboxylic acid	117322-30-2	Iris Biotech
AF488	AlexaFluor488	AlexaFluor488-NHS Ester		Fisher Scientific
Aib	2-Aminoisobutyric acid	Fmoc- α -aminoisobutyric acid	94744-50-0	Fluorochem
Aza-Gly	Azaglycine			
Aze	Azetidine	Fmoc-L-azetidine-2-carboxylic acid	136552-06-2	Combi Blocks
β -Ala	β -Alanine	Fmoc- β -alanine	35737-10-1	Fluorochem
β -AlaSO ₃ H	β -Alanine(SO ₃ H)	Fmoc-alpha-sulfo-beta-Alanine	1005412-03-2	Iris Biotech
C5g	Cyclopentylglycine	Fmoc-L-cyclopentylglycine	220497-61-0	Fluorochem
Cba	β -Cyclobutylalanine	Fmoc- β -cyclobutyl-L-alanine	478183-62-9	IRIS Biotech GmbH
Cpa	β -Cyclopropylalanine	Fmoc- β -cyclopropyl-L-alanine	214750-76-2	Fluorochem
Cpg	Cyclopropylglycine	Fmoc-L-cyclopropylglycine	1212257-18-5	Apollo Scientific
Cya	Cysteic acid	Fmoc-L-cysteic acid	320384-09-6	
D-3,3-DPA	3,3-diphenyl-D-alanine	Fmoc-3,3-diphenyl-D-alanine	189937-46-0	Chem-Impex international

D-Arg	D-Arginine	Fmoc-D-Arginine(Pbf)	187618-60-6	Iris Biotech
D-Asp	D-Aspartic acid	Fmoc-D-aspartic acid 4-tert-butyl ester	112883-39-3	Sigma aldrich
D-Cya	D-cysteic acid	Fmoc-D-cysteic acid		Costom synthesis
D-K	D-Lysine	Fmoc-D-Lysine(Boc)	92122-45-7	Sigma Aldrich
DOTA	1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid			
Fl	5(6)-carboxyfluorescein			Sigma
HArg	HomoArginine	Fmoc-L-HomoArg(Pbf)-OH	401915-53-5	Fluorochem
HPhe	HomoPhenylalanine	Fmoc-L-Homophenylalanine	132684-59-4	Iris Biotech
HyP	Hydroxyproline	Fmoc-Hydroxyproline(tBu)-OH	122996-47-8	Sigma
hSerMe	HomoSerine(methyl)	Fmoc-O-methyl-L-homoserine	173212-86-7	Iris Biotech
Lys(Dde)	Lysine(Dde)	N- α -Fmoc-N- ϵ -1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethyl-L-lysine	150629-67-7	Sigma Aldrich
NO2Phe	4-Nitrophenylalanine	Fmoc-4-nitro-L-phenylalanine	95753-55-2	PolyPeptide
Phg	Phenylglycine	Fmoc-L-phenylglycine	102410-65-1	Combi Blocks
Pip	Pipecolic acid	Fmoc-L-Pipecolic acid	86069-86-5	Peptech
Sar	Sarcosine, such that Sar _x represents x Sar residues	Fmoc-Sarcosine-OH	77128-70-2	Sigma
tBuGly	Tert-leucine	Fmoc-L-tert-leucine	132684-60-7	Fluorochem

Thi	2-Thienylalanine	Fmoc-2-Thienylalanine	130309-35-2	Novabiochem
ThiAz	3-(1,2,4-triazol-1-yl)-Alanine	Fmoc-3-(1,2,4-triazol-1-yl)-Ala-OH	1217449-37-0	Sigma
ΨAla	Reduced amide on backbone			

Materials and Methods

Peptide Synthesis

- 5 Peptides were synthesized by solid phase synthesis. Rink Amide MBHA Resin was used. To a mixture containing Rink Amide MBHA (0.4-0.45 mmol/g) and Fmoc-Cys(Trt)-OH (3.0 eq) was added DMF, then DIC (3 eq) and HOAt (3 eq) were added and mixed for 1 hour. 20% piperidine in DMF was used for deblocking. Each subsequent amino acid was coupled with 3 eq using activator reagents, DIC (3.0 eq) and HOAT (3.0 eq) in DMF. The reaction was
- 10 monitored by ninhydrin color reaction or tetrachlor color reaction. After synthesis completion, the peptide resin was washed with DMF x 3, MeOH x 3, and then dried under N₂ bubbling overnight. The peptide resin was then treated with 92.5% TFA/2.5% TIS/2.5% EDT/2.5% H₂O for 3h. The peptide was precipitated with cold isopropyl ether and centrifuged (3 min at 3000 rpm). The pellet was washed twice with isopropyl ether and the crude peptide was dried under
- 15 vacuum for 2 hours and then lyophilised. The lyophilised powder was dissolved in of ACN/H₂O (50:50), and a solution of 100 mM TATA in ACN was added, followed by ammonium bicarbonate in H₂O (1M) and the solution mixed for 1 h. Once the cyclisation was complete, the reaction was quenched with 1M aq. Cysteine hydrochloride (10 eq relative to TATA), then mixed and left to stand for an hour. The solution was lyophilised to afford crude product. The
- 20 crude peptide was purified by Preparative HPLC and lyophilized to give the product

All amino acids, unless noted otherwise, were used in the L- configurations.

Chemical structure of a long peptide chain, likely a model for a protein or a polymer. The structure is labeled "Solid phase synthesis" with an arrow pointing to the right. The peptide backbone is shown with various side chains, including a long aliphatic chain, a cyclic peptide, and a long aliphatic chain with a terminal amine. The structure is highly detailed, showing the stereochemistry of the amino acids and the connectivity of the side chains.

BCY6099

Sequence: (β-Ala)-Sar₁₀-(SEQ ID NO: 2)-CONH₂

8.0 g of resin was used to generate 2.1 g BCY6099 (99.2% purity; 16.3% yield) as a white solid.

BCY6099 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	15-45% B over 20 minutes, then 3 min 95% B
Retention Time:	11.31 min
LCMS (ESI):	m/z 1061.8 [M+3H] ³⁺ , 796.5 [M+4H] ⁴⁺
Peptide mw	3183.68


BCY6014 (Compound 67)



4.79 g of resin was used to generate 1.07g BCY6014 (Q1: 131.9 mg, 97.99% purity; Q2: 141.7 mg, 99.04% purity; Q3: 800.7 mg, 92.35% purity; 16.9% yield) as white a solid.

BCY6014 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	9.95 min
LCMS (ESI):	m/z 1013.8 [M+3H] ³⁺ , 760.4 [M+4H] ⁴⁺
Peptide mw	3039.53

BCY6104 (Compound 99)




Sequence: (β-Ala)-Sar₁₀-A(HArg)DC(Hyp)(Hse(Me))VNPLCLHP(D-Asp)W(HArg)C ((β-Ala)-Sar₁₀-(SEQ ID NO: 85))

4.44 g of resin was used to generate 700 mg BCY6104 (95.87% purity, 10.5% yield) as white a solid.

BCY6104 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	7.06 min
LCMS (ESI):	m/z 1062.1 [M+3H] ³⁺ , 796.8 [M+4H] ⁴⁺
Peptide mw	3185.65

BCY6103 (Compound 100)



WO 2019/122863

BCY6103


Sequence: (β-Ala)-Sar₁₀-A(HArg)DC(HyP)(Hse(Me))VNPLCLHP(D-Asp)WTC ((β-Ala)-Sar₁₀-(SEQ ID NO: 86))

4.44 g of resin was used to generate 700 mg BCY6103 (98.9% purity, 11.1% yield) as white a solid.

BCY6103 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	8.02 min
LCMS (ESI):	m/z 1039.1 [M+3H] ³⁺ , 779.5 [M+4H] ⁴⁺
Peptide mw	3117.55


BCY6101 (Compound 101)

Sequence: (β-Ala)-Sar₁₀-A(HArg)DC(HyP)LVNPLCLHP(D-Ala)WTC ((β-Ala)-Sar₁₀-(SEQ ID NO: 87))



BCY6101 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H2O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	9.79 min
LCMS (ESI):	m/z 1023.6 [M+3H] ³⁺ , 768.0 [M+4H] ⁴⁺
Peptide mw	3069.55

BCY6102 (Compound 102)




BCY6102 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1% TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	9.89 min
LCMS (ESI):	m/z 1018 [M+3H] ³⁺ , 763.9 [M+4H] ⁴⁺
Peptide mw	3053.56

BCY6139 (Compound 103)

5

Sequence: (β-Ala)-Sar₁₀-ARDC(Hyp)LVNPLCLHPGWTC ((β-Ala)-Sar₁₀-(SEQ ID NO: 89))




BCY6139

4.44 g of resin was used to generate 900 mg BCY6139 (97.4% purity, 11.2% yield) as white a solid.

BCY6139 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	8.95 min
LCMS (ESI):	m/z 1014.6 [M+3H] ³⁺ , 761.2 [M+4H] ⁴⁺
Peptide mw	3042.51

5 BCY6138 (Compound 104)


Sequence: (β-Ala)-Sar₁₀-ARDCPLVNPLCL(D-3,3-DPA)PGWTC ((β-Ala)-Sar₁₀-(SEQ ID NO: 90))



1.11 g of resin was used to generate 200 mg BCY6138 (95.2% purity, 12.2% yield) as white a solid.

BCY6138 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 20 minutes, then 3 min 95% B
Retention Time:	14.46 min
LCMS (ESI):	m/z 1037.6 [M+3H] ³⁺
Peptide mw	3111.63


5 BCY6137 (Compound 105)
 Sequence: (β-Ala)-Sar₁₀-ARDCPLVNPLCLHPGWTCLH ((β-Ala)-Sar₁₀-(SEQ ID NO: 91))



4.44 g of resin was used to generate 600 mg BCY6137 (98.9% purity, 9.06% yield) as white a solid.

BCY6137 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	14.46 min
LCMS (ESI):	m/z 1092.7 [M+3H] ³⁺ , 819.6 [M+4H] ⁴⁺
Peptide mw	3275.8

5 **BCY6042 (Compound 91)**

Sequence: Ac-(SEQ ID NO: 14)-Sar₆-(D-K)

1.11 g of resin was used to generate 99.2 mg BCY6042 (99.2% purity, 7.0% yield) as white a solid.

BCY6042 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	9.12 min
LCMS (ESI):	m/z 943.5 [M+3H] ³⁺
Peptide mw	2825.31

BCY6019 (Compound 77)


Sequence: Ac-(SEQ ID NO: 12)-Sar₆-(D-K)

4.79 g of resin was used to generate 732.0 mg BCY6019 (92.82% purity, 12.2% yield) as white a solid.

BCY6019 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H2O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	11.36 min
LCMS (ESI):	m/z 935.5 [M+3H] ³⁺
Peptide mw	2805.32

5

BCY6059 (Compound 106)



Sequence: Ac-(SEQ ID NO: 12)-Sar₆-(D-K[Ac])

To a solution of BCY6019 (0.05 g, 17.82 μmol , 1.00 eq) in H₂O (3 mL) was adjusted PH=11 by Na₂CO₃ (aq) and added acetyl acetate (5.46 mg, 53.46 μmol , 5.01 μL , 3.00 eq). The mixture was stirred at 15 °C for 1 hr. LC-MS showed BCY6019 was consumed completely and one main peak with desired MS was detected. The reaction was adjusted PH=7 by 1 N HCl and directly purified by prep-HPLC (TFA condition). Compound **BCY6059** (18.1 mg, 6.36 μmol , 35.67% yield) was obtained as a white solid.

BCY6059 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	6.67 min
LCMS (ESI):	m/z 949.8 [M+3H] ³⁺
Peptide mw	2848.36

BCY6160 (Compound 107)





Sequence: (β-AlaSO₃H)-Sar₄-(Cya)-Sar₄-(Cya)-A(HArg)DCPLVNPLCLHP(D-Cya)WTC ((β-AlaSO₃H)-Sar₄-(Cya)-Sar₄-(Cya)-92))

5 1.11 g of resin was used to generate 45.2 mg BCY6160 (95.5% purity, 2.5% yield) as white a solid.

BCY6160 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	11.38 min
LCMS (ESI):	m/z 1124.9 [M+3H] ³⁺
Peptide mw	3376.83

BCY6009 (Compound 108)




Sequence: ((β-Ala)-Sar₁₀-ARDCPLVNPLCLHPGWTC ((β-Ala)-Sar₁₀-(SEQ ID NO: 10))

4.79 g of resin was used to generate 2.42 g BCY6009 (>88.92% purity, 36.0% yield) as white a solid.

BCY6009 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	10.16 min
LCMS (ESI):	m/z 1008.9 [M+3H] ³⁺ , 756.9 [M+4H] ⁴⁺
Peptide mw	3025.5

BCY6017 (Compound 109)




BCY-6017

Sequence: A(HArg)DCPLVNPLCLHPGWTC (SEQ ID NO: 11)

1.19 g of resin was used to generate 189.9 mg BCY6017 (95.05% purity, 16.8% yield) as white a solid.

BCY6017 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H2O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	10.01 min
LCMS (ESI):	m/z 1129.1 [M+2H] ²⁺ , 753.0 [M+3H] ³⁺
Peptide mw	2257.67

BCY6018 (Compound 110)




Sequence: (β-Ala)-Sar₅-(SEQ ID NO: 11)

1.19 g of resin was used to generate 289.1 mg BCY6018 (97.92% purity, 21.0% yield) as white a solid.

BCY6018 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	9.77 min
LCMS (ESI):	m/z 1342.9 [M+2H] ²⁺ , 895.3 [M+3H] ³⁺
Peptide mw	2684.14


BCY6152 (Compound 111)



Sequence: (β-AlaSO₃H)-Sar₁₀-(SEQ ID NO: 11)

1.11 g of resin was used to generate 150.0 mg BCY6152 (98.75% purity; 9.5% yield) as white a solid.

BCY6152 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	10.09 min
LCMS (ESI):	m/z 1040.3 [M+3H] ³⁺
Peptide mw	3119.59



5

BCY6141 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	11.41 min
LCMS (ESI):	m/z 1085.5 [M+3H] ³⁺
Peptide mw	3255.78

BCY6026 (Compound 87)




Sequence: ACPLVNPLCLHPGWSCRGQ (SEQ ID NO: 77)

1.11 g of resin was used to generate 285.0 mg BCY6026 (97.7% purity; 24.2% yield) as white a solid.

BCY6026 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	9.31 min
LCMS (ESI):	m/z 1150 [M+2H] ²⁺ , 767.0 [M+3H] ³⁺
Peptide mw	2299.71

5

BCY6153 (Compound 113)



Sequence: (β-AlaSO₃H)-Sar₅-(SEQ ID NO: 11)

1.11 g of resin was used to generate 140.0 mg BCY6153 (98.59% purity, 9.9% yield) as white a solid.

BCY6153 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	20-50% B over 20 minutes, then 3 min 95% B
Retention Time:	10.33 min
LCMS (ESI):	m/z 1382.6 [M+2H] ²⁺ , 922.0 [M+3H] ³⁺
Peptide mw	2764.2

Preparation of Bicyclic Peptide Drug Conjugates

The general schematic for preparing Bicycle drug conjugates (BDCs) is shown in Figure 3 and Table A describes the component targeting bicycle and linker/toxin within each BDC.


5 **Table A**

BDC (BCY No)	Targeting Bicycle (BCY No)	Linker/Toxin	6048	6017	
6136	6099	ValCit-MMAE	6036	6019	
6033	6014		6028	6009	
6029	6009		6039	6014	DM1-(Me)-SS-(Me)-
6122	6104		6055	6014	DM1-SS-(Me ₂)-
6053	6018		6077	6014	DM1-SS-(Me)-SO ₃ H-
6049	6017				Non-cleavable (MMAE)
6037	6019		6063	6014	Non-cleavable (DM1)
6030	6009	TrpCit-MMAE	6064	6014	MMAE-Ala-Ala-Asn
6034	6014		6105	6014	MMAE-D-Ala-Phe-
6050	6017		6106	6014	Lys-
6054	6018		6175	6099	MMAE-Glu-Pro-Cit-
6038	6019		6107	6014	Gly-hPhe-Tyr-Leu-
6061	6014	ValLys-MMAE			
6174	6099				
6062	6014	D-TrpCit-MMAE			
6135	6099	DM1-SS-			
6031	6014				
6134	6104				
6027	6009				
6047	6017				
6035	6019				
6051	6018				
6154	6152				
6155	6153				
6173	6099	DM1-SS(SO ₃ H)-			
6082	6014				
6150	6018				
6151	6104				
6162	6138				
6161	6137				
6032	6014	DM1-SS-(Me)-			
6052	6018				


The synthesis of Bicyclic Peptide Drug Conjugates BCY6027, BCY6028, BCY6031 and BCY6032 listed in Table 6 were performed using the protocol disclosed in WO 2016/067035.

Activated bicycle peptides with formula (C) and (D):

5



(C)



(D)

were synthesised by reacting the free amino group of the bicycle precursors with,
 10 respectively, SPP (N-succinimidyl 4-(2-pyridyldithio)pentanoate, Annova Chem) and SPDB
 (N-succinimidyl 3-(2-pyridyldithio)propionate, Annova Chem) in DMSO. Concentrations of
 bicycle precursors were 10 mM or higher, with a 1.3-fold excess of SPP or SPDB, and a 20-
 fold excess of diisopropylethylamine, at room temperature. The reaction was judged
 complete after 1 hour, as judged by LCMS. Purification was performed by reverse phase as
 15 described above. Appropriate fractions were lyophilised.


Activated bicycle peptides with formula (C) and (D) were disulphide exchanged with 1.15
 equivalents of DM1 (as the free thiol), in semi aqueous conditions (50 % dimethylacetamide
 and 50% 100mM sodium acetate pH 5.0 supplemented with 2mM EDTA) for 21 hours at
 20 room temperature under a nitrogen gas blanket. Concentrations of activated bicycle peptides
 with structure C and D in the reaction were at 10 mM or higher.

This was followed by standard reverse phase purification using a C18 column. Fractions at
 purity greater than 95% were isolated and lyophilised. The materials did not contain
 25 measurable quantities of free toxin.

MMAE Series

Val-Cit-MMAE Series

Val-Cit-MMAE Linker




Compound 2

- 5 The peptide was synthesized by solid phase synthesis. 50g CTC Resin (sub: 1.0 mmol/g) was used. To a mixture containing CTC Resin (50 mmol, 50 g, 1.0 mmol/g) and Fmoc-Cit-OH (19.8 g, 50 mmol, 1.0 eq) was added DCM (400 mL), then DIEA (6.00 eq) was added and mixed for 3 hours. And then MeOH (50 mL) was added and mixed for 30 min for capping. 20% piperidine in DMF was used for deblocking. Boc-Val-OH (32.5g, 150mmol, 3eq) was coupled with 3 eq using HBTU (2.85 eq) and DIPEA (6.0 eq) in DMF (400 mL). The reaction was monitored by ninhydrin colour reaction test. After synthesis completion, the peptide resin was washed with DMF X 3, MeOH X 3, and then dried under N₂

bubbling overnight. After that the peptide resin was treated with 20% HFIP/DCM for 30 min for 2 times. The solution was removed on a rotary evaporator to give the crude. The crude peptide was dissolved in ACN/H₂O, then lyophilized twice to give the peptide product (17.3g crude).

LCMS (ESI):	m/z 374.9 [M+H] ⁺
Molecular weight	374.44

Compound 3




5

A solution of Compound 2 (4.00 g, 10.68 mmol, 1.00 eq) in DCM (40.00 mL) and MeOH (20.00 mL) was stirred at room temperature, then (4-aminophenyl)methanol (1.58 g, 12.82 mmol, 1.20 eq) and EEDQ (5.28 g, 21.37 mmol, 2.00 eq) were added and the mixture stirred in the dark for 9 hrs. TLC (dichloromethane/methanol= 5/1, R_f = 0.56) indicated one new spot had formed. The reaction mixture was concentrated under reduced pressure to remove solvent. The resulting residue was purified by flash silica gel chromatography (ISCO®; 120 g SepaFlash® Silica Flash Column, Eluent of 0~20% MeOH/DCM @ 80 mL/min). Compound 3 (3.00 g, 6.26 mmol, 58.57% yield) was obtained as a white solid.

10

LCMS (ESI):	m/z 480.1 [M+H] ⁺
Molecular weight	479.58


Compound 4



To a solution of Compound **3** (3.00 g, 6.26 mmol, 1.00 eq) in anhydrous THF (35.00 mL) and anhydrous DCM (15.00 mL) was added (4-nitrophenyl) chloroformate (6.31 g, 31.30 mmol, 5.00 eq) and pyridine (2.48 g, 31.30 mmol, 2.53 mL, 5.00 eq), and the mixture was stirred at 25 °C for 5 hrs. TLC (dichloromethane/methanol= 10/1, R_f = 0.55) indicated a new spot had formed. The reaction mixture was filtered, and the filtrate was concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 120 g SepaFlash® Silica Flash Column, Eluent of 0~10% DCM/MeOH@ 80 mL/min). Compound **4** (2.00 g, 3.10 mmol, 49.56% yield) was obtained as a white solid.

LCMS (ESI):	m/z 667.3 [M+Na] ⁺
Molecular weight	644.68


Compound 5



A mixture of Compound **4** (278.43 mg, 387.80 μmol , 1.00 eq) and DIEA (501.19 mg, 3.88 mmol, 677.29 μL , 10.00 eq) in DMF (5.00 mL) was stirred under nitrogen for 10 min. MMAE (250.00 mg, 387.80 μmol , 1.00 eq) and HOBt (52.40 mg, 387.80 μmol , 1.00 eq) were added and the mixture was stirred at 0 °C under nitrogen for 20 min and stirred at 30 °C for additional 18 hrs. LC-MS showed one main peak with desired mass was detected. The resulting mixture was purified by flash C18 gel chromatography (ISCO®; 130 g SepaFlash® C18 Flash Column, Eluent of 0~50% MeCN/H₂O @ 75 mL/min). Compound **5** (190.00 mg, 155.29 μmol , 40.04% yield) was obtained as a white solid.


LCMS (ESI):	m/z 1223.4 [M+H] ⁺
Molecular weight	1223.57

Compound 6



To a solution of Compound **5** (170.00 mg, 138.94 μmol , 1.00 eq) in DCM (2.70 mL) was added 2,2,2-trifluoroacetic acid (413.32 mg, 3.62 mmol, 268.39 μL , 26.09 eq), and the mixture was stirred at 25 °C for 1 hr. LC-MS showed Compound **5** was consumed completely. The mixture was concentrated under reduced pressure to give a residue. The residue was dissolved in THF (10.00 mL) and was added K₂CO₃ (192.03 mg, 1.39 mmol, 10.00 eq), the mixture was stirred at room temperature for additional 3 hrs. LC-MS showed one main peak with desired mass was detected. The resulting reaction mixture was concentrated under reduced pressure to remove solvent to give a residue. The residue was purified by flash C18 gel chromatography (ISCO®; 130 g SepaFlash® C18 Flash Column, Eluent of 0~50% MeCN/H₂O @ 75 mL/min). Compound **6** (110.00 mg, 97.92 μmol , 70.48% yield) was obtained as a white solid.


LCMS (ESI):	m/z 1123.4 [M+H] ⁺
Molecular weight	1123.45

Compound 7

To a solution of Compound **6** (110.00 mg, 97.92 μmol , 1.00 eq) in DMA (5 mL), DIEA (25.31 mg, 195.83 μmol , 2.00 eq) and tetrahydropyran-2,6-dione (22.34 mg, 97.92 μmol , 1.00 eq). The mixture was stirred at room temperature for 18 hrs. LC-MS showed Compound **6** was consumed completely and one main peak with desired mass was detected. The reaction mixture was purified by flash C18 gel chromatography (ISCO®; 130 g SepaFlash® C18 Flash Column, Eluent of 0~50% MeCN/H₂O @ 75 mL/min). Compound **7** (100.00 mg, 80.81 μmol , 82.53% yield) was obtained as a white solid.

LCMS (ESI):	m/z 1237.4 [M+H] ⁺
Molecular weight	1236.74

10 **Compound 8 (MMAE-PABC-Cit-Val-Glutarate-NHS)**



To a solution of Compound **7** (100.00 mg, 80.81 μmol , 1.00 eq) in DMA (4.5 mL) and DCM (1.5 mL) was added 1-hydroxypyrrolidine-2,5-dione (27.90 mg, 242.42 μmol , 3.00 eq) under N₂, the mixture was stirred at 0 °C for 30 min. EDCI (46.47 mg, 242.43 μmol , 3.00 eq) was added in the mixture, and the mixture was stirred at 25 °C for additional 16 hrs. LC-MS showed Compound **7** was consumed completely and one main peak


with desired mass was detected. The reaction mixture was purified by flash C18 gel chromatography (ISCO®; 130 g SepaFlash® C18 Flash Column, Eluent of 0~50% MeCN/H₂O @ 75 mL/min). Compound **8** (90.00 mg, 60.69 μmol, 75.11% yield) was obtained as a white solid.

LCMS (ESI):	m/z 1334.5 [M+H] ⁺
Molecular weight	1334.62

General Procedure for Coupling MMAE-PABC-Cit-Val-Glutarate-NHS with Targeting Bicycles

- To a solution of Bicycle (1.0 – 1.3 eq) in DMA was added DIEA (3 eq) and MMAE-PABC-Cit-Val-Glutarate-NHS (1 eq). The mixture was stirred at 25 °C for 18 hr. The reaction was monitored by LC-MS and once complete, was directly purified by preparative HPLC.

BCY6136




- BCY6099 (71.5 mg, 22.48 μmol) was used as the bicycle reagent. Compound BCY6136 (40.9 mg, 9.05 μmol, 40.27% yield, 97.42% purity) was obtained as a white solid.

BCY6136 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B

Retention Time:	11.35 min
LCMS (ESI):	m/z 1468.1 [M+3H] ³⁺ , 1101.2 [M+4H] ⁴⁺ , 881.3 [M+5H] ⁵⁺
Peptide mw	4404.2


BCY6033



BCY6014 (70.00 mg, 22.47 μmol , 1.00 eq) was used as the bicycle reagent. Compound BCY6033 (33.90 mg, 7.96 μmol , 34.57% yield) was
5 obtained as a white solid.

BCY6033 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	35-65% B over 20 minutes, then 3 min 95% B
Retention Time:	7.47 min
LCMS (ESI):	m/z 1065.2 [M+4H] ⁴⁺ , 852.2 [M+5H] ⁵⁺
Peptide mw	4259.04


BCY6029



BCY6009 (70.0 mg, 22.47 μmol , 1 eq) was used as the bicycle reagent. Compound **BCY6029** (32.9 mg, 7.75 μmol , 33.49% yield) was obtained as a white solid.

BCY6029 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	35-65% B over 20 minutes, then 3 min 95% B
Retention Time:	7.46 min
LCMS (ESI):	m/z 1061.7 [M+4H] ⁴⁺
Peptide mw	4245.02


5 BCY6122



BCY6104 (71.59 mg, 22.48 μmol , 1.00 eq) was used as the bicycle reagent. Compound **BCY6122** (38.30 mg, 8.57 μmol , 38.14% yield, 98.58% purity) was obtained as a white solid.

BCY6122 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	10.72 min
LCMS (ESI):	m/z 1101.8 [M+4H] ⁴⁺ , 881.5 [M+5H] ⁵⁺
Peptide mw	4406.18

BCY6053




BCY6018 (72.40 mg, 26.97 μmol , 1.2 eq) was used as the bicycle reagent. Compound **BCY6053** (38.3 mg, 9.81 μmol , 43.65% yield) was obtained as a white solid.

BCY6053 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min

Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	12.95 min
LCMS (ESI):	m/z 1301.7 [M+3H] ³⁺ , 976.5 [M+4H] ⁴⁺
Peptide mw	3905.67

BCY6049




BCY6017 (50.75 mg, 22.48 μmol , 1.2 eq) was used as the bicycle reagent. Compound BCY6049 (22.5 mg, 6.47 μmol , 34.54% yield) was obtained as a white solid.

BCY6049 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	14.28 min
LCMS (ESI):	m/z 1159.6 [M+3H] ³⁺ , 869.8 [M+4H] ⁴⁺

Peptide mw	3479.2
------------	--------

BCY6037





BCY6019 (65.00 mg, 22.47 μmol , 1.00 eq) was used as the bicycle reagent. Compound **BCY6037** (26.80 mg, 6.66 μmol , 28.74% yield) was

5 obtained as a white solid.

BCY6037 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	35-65% B over 20 minutes, then 3 min 95% B
Retention Time:	8.79 min
LCMS (ESI):	m/z 1342.1 [M+3H] ³⁺ , 1006.6 [M+4H] ⁴⁺
Peptide mw	4025.84


Trp-Cit-MMAE Series
Trp-Cit-MMAE Linker



General procedure for preparation of 3

To a solution of compound 2 (4.00 g, 8.67 mmol, 1.00 eq), DIC (1.61 g, 12.78 mmol, 1.97 mL, 9.00 eq) and HOBt (10.54 g, 78.00 mmol, 9.00 eq) in DMF (30.00 mL) was added (4-aminophenyl)methanol (9.61 g, 78.00 mmol, 9.00 eq). The mixture was stirred at 15 °C for 1 hour. LC-MS showed compound 2 was consumed completely and one main peak with desired MS was detected. The mixture was purified by prep-HPLC. Compound 3 (4.20 g, 7.41 mmol, 85.49% yield) was obtained as a white solid.

LCMS (ESI):	m/z 566.9 [M+H] ⁺
Molecular weight	566.66


General procedure for preparation of 4

To a solution of compound 3 (4.20 g, 6.30 mmol, 1.00 eq), DIPEA (1.09 g, 8.40 mmol, 1.47 mL, 7.00 eq) in DMF (30.00 mL) was added bis(4-nitrophenyl) carbonate (11.50 g, 37.79 mmol, 6.00 eq) in one part. The mixture was stirred at 0-15 °C for 1.5 hour. LC-MS showed compound 3

was consumed completely and one main peak with desired MS was detected. Directly purified by prep-HPLC (TFA condition). Compound **4** (2.00 g, 2.40 mmol, 38.16% yield) was obtained as a white solid.

LCMS (ESI):	m/z 732.0 [M+H] ⁺
Molecular weight	731.76


General procedure for preparation of **5**



To a solution of compound **4** (300.00 mg, 360.63 μ mol, 1.00 eq), DIEA (93.22 mg, 721.27 μ mol, 125.97 μ L, 3.00 eq) in DMF (10.00 mL) was added MMAE (233.03 mg, 324.57 μ mol, 0.90 eq) and HOBt (48.73 mg, 360.63 μ mol, 1.00 eq) at 0 °C. The mixture was stirred at 30 °C for 18 hour. LC-MS showed compound **4** was consumed completely and one main peak with desired MS was detected. Directly purified by prep-HPLC (neutral condition). Compound **5** (250.00 mg, 190.75 μ mol, 52.89% yield) was obtained as a yellow solid.

LCMS (ESI):	m/z 1310.5 [M+H] ⁺
Molecular weight	1310.65


General procedure for preparation of **6**



To a solution of compound 5 (240.00 mg, 183.12 μmol , 1.00 eq) in DCM (10.00 mL) was added TFA (1.54 g, 13.51 mmol, 1.00 mL, 73.76 eq). The mixture was stirred at 15 °C for 2 hour. And the mixture was concentrated under reduced pressure to remove solvent to give a residue, the residue was dissolved in THF and added K_2CO_3 and stirred at 15 °C for 2 h. LC-MS showed compound 5 was consumed completely and one main peak with desired MS was detected. The reaction mixture was concentrated under reduced pressure to remove solvent to give a residue. The residue was purified by prep-HPLC (neutral condition). The crude product **6** (125.00 mg, 94.37 μmol , 51.53% yield, TFA) was used into the next step without further purification.

LCMS (ESI):	m/z 1210.4 $[\text{M}+\text{H}]^+$
Molecular weight	1209.53

General procedure for preparation of 7




10

To a solution of compound 6 (125.00 mg, 94.37 μmol , 1.00 eq, TFA) in DMA (5.00 mL) was added DIEA (24.39 mg, 188.75 μmol , 32.96 μL , 2.00 eq), tetrahydropyran-2,6-dione (21.54 mg, 188.75 μmol , 2.00 eq). The mixture was stirred at 15 °C for 2 hour. LC-MS showed compound **6**

was consumed completely and one main peak with desired MS was detected. Directly purified by prep-HPLC (neutral condition). Compound **7** (100.00 mg, 75.49 μmol , 80.00% yield) was obtained as a white solid.

LCMS (ESI):	m/z 1324.5 [M+H] ⁺
Molecular weight	1324.63

General procedure for preparation of **8** (MMAE-PABC-Cit-Trp-Glutarate-NHS)



To a solution of compound **7** (100.00 mg, 75.49 μmol , 1.00 eq), 1-hydroxypropylidene-2, 5-dione (26.07 mg, 226.48 μmol , 3.00 eq) in DMA (3.00 mL) and DCM (1.00 mL) was added EDCI (43.42 mg, 226.48 μmol , 3.00 eq). The mixture was stirred at 15 °C for 4 hour. LC-MS showed compound **7** was consumed completely and one main peak with desired MS was detected. The DCM was removed. Directly was purified by prep-HPLC (neutral condition). Compound **8** (60.00 mg, 42.20 μmol , 55.91% yield) was obtained as a white solid.

LCMS (ESI):	m/z 711.2 [M+2H] ²⁺
Molecular weight	1421.7

General Procedure for Coupling MMAE-PABC-Cit-Trp-Glutarate-NHS with Targeting Bicycles


To a solution of Bicycle (1.0-1.3 eq) in DMA was added DIEA (3 eq) and MMAE-PABC-Cit-Trp-Glutarate-NHS (1 eq). The mixture was stirred at 25 °C for 18 hr. The reaction was monitored by LC-MS and once complete, was directly purified by preparative HPLC.

15 BCY6030




5 BCY6034

BCY6030 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	35-65% B over 20 minutes, then 3 min 95% B
Retention Time:	7.90 min
LCMS (ESI):	m/z 1083.7 [M+4H] ⁴⁺
Peptide mw	4332.17




BCY6014 (88.21 mg, 23.21 μmol , 1.10 eq) was used as the bicycle reagent. Compound **BCY6034** (27.70 mg, 6.05 μmol , 28.70% yield, 95.02% purity) was obtained as a white solid.

BCY6034 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	30-60% B over 20 minutes, then 3 min 95% B
Retention Time:	11.49 min
LCMS (ESI):	m/z 1449.3 [M+3H] ³⁺ , 1087.4 [M+4H] ⁴⁺
Peptide mw	4346.13




BCY6017 (57.17 mg, 25.32 μmol , 1.2 eq) was used as the bicycle reagent. Compound **BCY6050** (0.0519 g, 14.56 μmol , 69.01% yield) was obtained as a white solid.

BCY6050 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1% TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	13.55 min
LCMS (ESI):	m/z 1188.5 [M+3H] ³⁺ , 891.7 [M+4H] ⁴⁺
Peptide mw	3564.25




BCY6018 (67.97 mg, 25.32 μ mol, 1.2 eq) was used as the bicycle reagent. Compound **BCY6054** (40.10 mg, 10.05 μ mol, 47.62% yield) was obtained as a white solid.

BCY6054 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1% TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	13.73 min
LCMS (ESI):	m/z 1330.4 [M+3H] ³⁺ , 998.1 [M+4H] ⁴⁺
Peptide mw	3990.72




BCY6019 (81.39 mg, 23.21 μ mol, 1.10 eq) was used as the bicycle reagent. Compound **BCY6038** (34.10 mg, 8.02 μ mol, 38.00% yield, 96.68% purity) was obtained as a white solid.

BCY6038 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	35-65% B over 20 minutes, then 3 min 95% B
Retention Time:	9.56 min
LCMS (ESI):	m/z 1371.0 [M+3H] ³⁺ , 1028.3 [M+4H] ⁴⁺
Peptide mw	4111.9




General procedure for preparation of Compound 2



To a mixture of Compound **1** (3.00 g, 5.89 mmol, 1 eq) and (4-aminophenyl)methanol (869.93 mg, 7.06 mmol, 1.2 eq) in DCM (35 mL) and MeOH (18 mL) was added EEDQ (2.91 g, 11.77 mmol, 2 eq) in the dark under nitrogen, the mixture was stirred at 25 °C for 5 hr. LC-MS showed Compound **1** was consumed completely and one main peak with desired MS was detected. The resulting reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 120 g SepaFlash® Silica Flash Column, Eluent of 0~20% MeOH/DCM @ 80mL/min). Compound **2** (2.2 g, 3.58 mmol, 60.79% yield) was obtained as a white solid.

LCMS (ESI):	m/z 615.0 [M+H] ⁺
Molecular weight	614.78

General procedure for preparation of Compound 3




To a solution Compound **2** (500 mg, 813.31 μmol , 1 eq) in THF (10 mL) was added DIEA (630.69 mg, 4.88 mmol, 849.98 μL , 6 eq) at 0 °C under nitrogen with stirring for 30 mins. Then bis(4-nitrophenyl) carbonate (1.48 g, 4.88 mmol, 6 eq) was added thereto, the mixture was stirred at 25 °C under nitrogen for additional 21 hr. LC-MS showed one main peak with desired MS was detected. The resulting reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 40 g SepaFlash®) Silica Flash Column, Eluent of 0~20% MeOH/DCM @ 40 mL/min). Compound **3** (500 mg, 641.13 μmol , 78.83% yield) was obtained as a yellow solid.

5

LCMS (ESI):	m/z 780.0 [M+H] ⁺
Molecular weight	779.89

General procedure for preparation of Compound 4




To a mixture of Compound **3** (500 mg, 512.90 μmol , 1.23 eq) in DMF (8 mL) was added DIEA (135.01 mg, 1.04 mmol, 181.95 μL , 2.5 eq) with stirring at 0 °C for 30 mins. Then MMAE (300 mg, 417.84 μmol , 1 eq) and HOBT (84.69 mg, 626.76 μmol , 1.5 eq) was added thereto, and the mixture was stirred at 40 °C for 15 hr. LC-MS showed compound **3** was consumed completely and one main peak with desired MS was detected. The residue was purified by flash C18 gel chromatography (ISCO®; 130 g SepaFlash® C18 Flash Column, Eluent of 0~60% MeCN/H₂O @ 75 mL/min). Compound **4** (330 mg, 242.87 μmol , 58.13% yield) was obtained as a white solid.

10

LCMS (ESI):	m/z 679.7 [M+2H] ²⁺
-------------	--------------------------------

Molecular weight	1358.77
------------------	---------


General procedure for preparation of Compound 5



To a solution of Compound **4** (325 mg, 239.19 μ mol, 1 eq) in DCM (18 mL) was added TFA (3.03 g, 26.60 mmol, 1.97 mL, 111.22 eq) at 0 °C, the mixture was stirred at 25 °C for 2 hr. LC-MS showed compound **4** was consumed completely. Then the reaction mixture was concentrated under reduced pressure to give a residue, the residue was dissolved in THF (10 mL) and K₂CO₃ (661.16 mg, 4.78 mmol, 20 eq) was added thereto. The mixture was stirred at 25 °C for 15 hrs. LC-MS showed one main peak with desired MS was detected. The resulting reaction mixture was filtered and the filtrate was concentrated under reduced pressure to give a residue. The residue was purified by flash C18 gel chromatography (ISCO®; 130 g SepaFlash® C18 Flash Column, Eluent of 0~60% MeCN/H₂O @ 75 mL/min). Compound **5** (170 mg, 135.07 μ mol, 56.47% yield) was obtained as a white solid.

LCMS (ESI):	m/z 629.7 [M+2H] ²⁺
Molecular weight	1258.65


General procedure for preparation of Compound 6



A round bottle containing a solution of compound **5** (140 mg, 111.23 μmol , 1 eq) in DMA (5 mL) was purged using a nitrogen balloon and added DIEA (28.75 mg, 222.46 μmol , 38.75 μL , 2 eq) at 0 °C with stirring for 10 mins, tetrahydropyran-2,6-dione (25.38 mg, 222.46 μmol , 2 eq) was added as a solution in DMA. The mixture was stirred at 25 °C for 12 hr. LC-MS showed compound **5** was consumed completely and one main peak with desired MS was detected. The resulting reaction mixture was purified by flash C18 gel chromatography (ISCO®; 43 g SepaFlash® C18 Flash Column, Eluent of 0~60% MeCN/H₂O @ 40 mL/min). Compound **6** (120 mg, 87.42 μmol , 78.59% yield) was obtained as a white solid.

LCMS (ESI):	m/z 686.7 [M+2H] ²⁺
Molecular weight	1372.75


General procedure for preparation of Compound **7** (MMAE-PABC-Lys(Dde)-Val-Glutamate-NHS)



To a solution of compound **6** (120 mg, 87.42 μmol , 1 eq) in DMA (9 mL) and DCM (3 mL) was added 1-hydroxypyrrolidine-2,5-dione (30.18 mg, 262.25 μmol , 3 eq) with stirring, and EDCI (50.27 mg, 262.25 μmol , 3 eq) was added thereto, the mixture was stirred at 0 °C for 30 mins and at 25 °C for additional 19 hr. LC-MS showed compound **6** was consumed completely and one main peak with desired MS was detected. The resulting reaction mixture was concentrated under reduced pressure to remove DCM. The mixture was purified by flash C18 gel chromatography (ISCO®; 43 g SepaFlash® C18 Flash Column, Eluent of 0-60% MeCN/H₂O @ 40 mL/min). Compound **7** (60 mg, 40.82 μmol , 46.70% yield) was obtained as a white solid.


LCMS (ESI):	m/z 735.3 [M+2H] ²⁺
Molecular weight	1469.83

General Procedure for Coupling MMAE-PABC-Lys(Dde)-Val-Glutarate-NHS with Targeting Bicycles




To a solution of Bicyclic (1.0-1.3 eq) in DMA was added DIEA (3 eq) and MMAE-PABC-Lys(Dde)-Val-Glutarate-NHS (1 eq). The mixture was stirred at 25 °C for 18 hr. The reaction was monitored by LC-MS and once complete, was directly purified by preparative HPLC.

General Procedure for Dde Deprotection



To a solution of Dde protected peptide (1 eq) in DMF was added hydrazine hydrate (6500 eq), and the mixture was stirred at 25 °C for 1 hr. LC-MS was used to monitor the reaction, and once complete, the mixture was purified by preparative HPLC and the clean fractions lyophilised.

5 BCY6061




BCY6014 (124.12 mg, 40.82 μmol , 1.2 eq) was used as the bicycle reagent. Dde-BCY6038 (80 mg, 18.20 μmol , 53.51% yield) was obtained as a white solid.

LCMS (ESI):	m/z 1099.0 [M+4H] ⁴⁺ , 879.4 [M+5H] ⁵⁺
Molecular weight	4395.24

5 Dde-BCY6061 (78 mg, 17.75 μmol) was deprotected using hydrazine according to the general procedure to give **BCY6061** (47.1 mg, 11.13 μmol , 62.73% yield) as a white solid.

BCY6061 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1% TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	12.01 min
LCMS (ESI):	m/z 1058.1 [M+4H] ⁴⁺ , 846.5 [M+5H] ⁵⁺
Peptide mw	4230.03

BCY6174



BCY6099 (389.77 mg, 122.47 μmol , 1.2 eq) was used as the bicycle reagent. Dde-BCY6174 (0.250 g, 55.10 μmol , 53.99% yield) was obtained as a white solid.


LCMS (ESI):	m/z 1513.0 $[\text{M}+3\text{H}]^{3+}$, 1135.0 $[\text{M}+4\text{H}]^{4+}$, 908.2 $[\text{M}+5\text{H}]^{5+}$
Molecular weight	4538.38

- 5 Dde-BCY6174 (0.250 g, 55.10 μmol , 1.0 eq) was deprotected using hydrazine according to the general procedure to give BCY6174 (0.1206 g, 27.45 μmol , 49.82% yield) as a white solid.


BCY6174 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1% TFA in ACN

Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	9.85 min
LCMS (ESI):	m/z 1458.5 [M+3H] ³⁺ , 1094.1 [M+4H] ⁴⁺ , 875.4 [M+5H] ⁵⁺
Peptide mw	4373.17

D-Trp-Cit-MMAE Series
D-Trp-Cit-MMAE Linker




General procedure for preparation of Compound 3



To a solution of compound 1 (2 g, 4.33 mmol, 1.00 eq), DIC (4.92 g, 39.00 mmol, 6.00 mL), HOBt (5.27 g, 39.00 mmol, 9.00 eq) in DMF (30.00 mL) was added (4-aminophenyl)methanol (4.80 g, 39.00 mmol, 9.00 eq). The mixture was stirred at 15 °C for 1 hour. LC-MS showed compound 1 was consumed completely and one main peak with desired MS was detected. Directly purified by prep-HPLC (neutral condition). Compound 2 (2 g, 3.53 mmol, 81.45% yield) was obtained as a white solid.

LCMS (ESI):	m/z 666.9 [M+H] ⁺
Molecular weight	666.78


General procedure for preparation of Compound 4



To a solution of compound 2 (2 g, 3.00 mmol, 1 eq), DIEA (2.71 g, 21.00 mmol, 3.66 mL, 7 eq) in DMF (20 mL) was added bis(4-nitrophenyl) carbonate (5.48 g, 18.00 mmol, 6 eq) in one part at 0°C. The mixture was stirred at 0-15 °C for 2 hr. LC-MS showed compound 2 was consumed completely and one main peak with desired MS was detected. Directly purified by prep-HPLC (neutral condition). Compound 3 (0.9 g, 1.08 mmol, 36.07% yield) was obtained as a yellow solid.

LCMS (ESI):	m/z 832.0 [M+H] ⁺
Molecular weight	831.88


General procedure for preparation of Compound 5



To a solution of compound 3 (350 mg, 420.74 μmol, 1.00 eq), HOBt (56.85 mg, 420.74 μmol, 1 eq) and DIEA (163.13 mg, 1.26 mmol, 219.86 μL, 3 eq) in DMF (10 mL) was added MMAE (302.08 mg, 420.74 μmol, 1 eq) at 0°C. The mixture was stirred at 40 °C for 18 hour. LC-MS showed compound 4 was consumed completely and one main peak with desired MS was detected. Directly purified by prep-HPLC (neutral condition). Compound 4 (0.22 g, 155.95 μmol, 37.06% yield) was obtained as a yellow solid.

LCMS (ESI):	m/z 1410.5 [M+H] ⁺ , 705.7 [M+2H] ²⁺
Molecular weight	1410.76


General procedure for preparation of Compound 6



To a solution of compound 4 (0.21 g, 148.86 μmol , 1 eq) in DCM (9 mL) was added TFA (1.54 g, 13.51 mmol, 1 mL, 90.73 eq). The mixture was stirred at 15 °C for 4 h, and concentrated under reduced pressure to give a residue, dissolved in THF, then added K_2CO_3 (s) and stirred at 15 °C for 16 h. LC-MS showed compound 4 was consumed completely and one main peak with desired MS was detected. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (neutral condition). Compound 5 (0.13 g, 102.02 μmol , 68.54% yield, 95% purity) was obtained as a white solid.

LCMS (ESI):	m/z 1210.4 $[\text{M}+\text{H}]^+$, 605.8 $[\text{M}+2\text{H}]^{2+}$
Molecular weight	1210.53

General procedure for preparation of Compound 7




To a solution of compound 5 (0.12 g, 99.13 μmol , 1 eq) in DMA (5 mL) was added DIEA (38.44 mg, 297.40 μmol , 51.80 μL , 3 eq) and tetrahydropyran-2,6-dione (22.62 mg, 198.26 μmol , 2 eq). The mixture was stirred at 15 °C for 16 h. LC-MS showed compound 5 was consumed

completely and one main peak with desired MS was detected. Directly purified by prep-HPLC (neutral condition). Compound 6 (0.09 g, 67.94 μmol , 68.54% yield) was obtained as a white solid.

LCMS (ESI):	m/z 662.7 $[\text{M}+2\text{H}]^{2+}$
Molecular weight	1324.63

General procedure for preparation of Compound 8



5

To a solution of compound 6 (0.09 g, 67.95 μmol , 1 eq), HOSu (23.46 mg, 203.84 μmol , 3 eq) in DMA (6 mL) and DCM (2 mL) was added EDCI (39.08 mg, 203.84 μmol , 3 eq). The mixture was stirred at 15 °C for 16 h. LC-MS showed compound 6 was consumed completely and one main peak with desired MS was detected. DCM was removed and directly purified by prep-HPLC (neutral condition). Compound 7 (0.06 g, 40.09 μmol , 59.01% yield, 95% purity) was obtained as a white solid.

LCMS (ESI):	m/z 711.2 $[\text{M}+2\text{H}]^{2+}$
Molecular weight	1421.7

10


BCY6062


To a solution of BCY6014 (76.99 mg, 25.32 μmol , 1.2 eq) in DMA (5 mL) was added DIEA (8.18 mg, 63.31 μmol , 11.03 μL , 3 eq), compound 7 (0.03 g, 21.10 μmol , 1 eq). The mixture was stirred at 15 °C for 16 hr. The reaction was monitored by LC-MS and once complete, the mixture was purified by preparative HPLC. **BCY6062** (0.0255 g, 5.70 μmol , 27.01% yield, 97.15% purity) was obtained as a white solid.

BCY6062 Analytical Data

Mobile Phase:	A: 0.1% TFA in H2O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	13.15 min
LCMS (ESI):	m/z 1449.2 [M+3H] ³⁺ , 1087.0 [M+4H] ⁴⁺
Peptide mw	4346.13

DM1 Series
DM1-SS- Series
DM1-SPDP-TFP Linker





To a solution of 2-(2-pyridyldisulfanyl)pyridine (12.37 g, 56.18 mmol, 1.50 eq) in EtOH (100.00 mL) was added 4-sulfanylbutoanoic acid (4.50 g, 37.45 mmol, 1.00 eq). The mixture was stirred at 15 °C for 18 hours under N₂. LC-MS showed compound **1** was consumed completely and one main peak with desired mass was detected. The reaction mixture was concentrated under reduced pressure to remove solvent to give a residue.

5 The residue was purified by preparative HPLC (C-18 360 g, neutral condition). Compound SPDB (1.9 g, 8.29 mmol, 22.12% yield) was obtained as a yellow solid.


¹H NMR: ES6446-8-P1A 400 MHz CDCl₃

δ ppm 1.98 (q, J=7.09 Hz, 2 H), 2.45 (t, J=7.15 Hz, 2 H), 2.79 (t, J=7.03 Hz, 2 H), 7.03 (dd, J=7.15, 4.89 Hz, 1 H), 7.19 (s, 1 H), 7.56 - 7.65 (m, 2 H), 8.41 (d, J=4.52 Hz, 1 H).

LCMS (ESI):	m/z 230.0 [M+H] ⁺
Molecular weight	229.31

10


DM1-SPDB



A mixture of DM1 (250.00 mg, 338.62 μmol , 1.00 eq) and 4-(2-pyridyldisulfanyl)butanoic acid (100.95 mg, 440.21 μmol , 1.30 eq) was added under nitrogen in a 50 mL of flask with DMF (10.00 mL) purged with N_2 for 30 mins. The mixture was stirred at room temperature for 1 hr. LC-MS showed that the DM1 was consumed completely and one main peak with desired mass was detected. The residue was purified by flash C18 gel chromatography (ISCO®; 120 g SepaFlash® C18 Flash Column, Eluent of 0~60% MeCN/ H_2O @ 85 mL/min). DM1-SPDB (120.00 mg, 140.11 μmol , 41.38% yield) was obtained as a white solid.

LCMS (ESI):	m/z 838.0 $[\text{M}+\text{H}-\text{H}_2\text{O}]^+$
Molecular weight	856.44

DM1-SPDB-TFP




To a solution of DM1-SPDB (120.00 mg, 140.11 μmol , 1.00 eq) and 2,3,5,6-tetrafluorophenol (69.81 mg, 420.34 μmol , 3.00 eq) in DCM (1.00 mL) and DMA (3.00 mL) was added EDCI (80.58 mg, 420.34 μmol , 3.00 eq). The mixture was stirred at 15 °C for 4 hours. LC-MS showed DM1-SPDB was consumed completely and one main peak with desired mass was detected. The DCM was removed and the residue The mixture was directly purified by preparative HPLC (neutral condition). Compound DM1-SPDB-TFP (60.00 mg, 59.73 μmol , 42.63% yield) was obtained as a white solid.

LCMS (ESI):	m/z 985.9 [M+H-H ₂ O] ⁺
Molecular weight	1004.5

General Procedure for Coupling DM1-SPDB-TFP with Targeting Bicycles


To a solution of targeting Bicycle (1.1-1.3 eq) in DMA was added DIEA (3 eq) and DM1-SPDB-TFP (1 eq). The mixture was stirred at 25 °C for 18 hr. The reaction was monitored by LC-MS and once complete, the mixture was directly purified by preparative HPLC.

BCY6135




BCY6099 (114.1 mg, 35.84 μmol) was used as the bicycle reagent. 22.4 mg Compound BCY6135 (5.30 μmol , 17.74% yield, 95.14% purity) was obtained as a white solid.

BCY6135 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	9.81
LCMS (ESI):	m/z 1341.5 [M+3H] ³⁺ , 805.0 [M+5H] ⁵⁺
Peptide mw	4021.08




BYC6014 (121.07 mg, 39.82 μmol) was used as the bicycle reagent. 59.90 mg compound BYC6031 (14.67 μmol , 36.85% yield, 95.02% purity) was obtained as a white solid.

BYC6031 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	35-65% B over 20 minutes, then 3 min 95% B
Retention Time:	6.284 min
LCMS (ESI):	m/z 1286.4 [M+3H-H ₂ O] ³⁺ , 965.6 [M+4H-H ₂ O] ⁴⁺
Peptide mw	3877.96




BYC6104 (95.11 mg, 29.87 μmol , 1 eq) was used as the bicycle reagent. BYC6134 (0.0232 g, 5.64 μmol , 18.89% yield, 97.82% purity) was obtained as a white solid.

BYC6134 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	9.10 min
LCMS (ESI):	m/z 1001.8 [M+4H-H ₂ O] ⁴⁺
Peptide mw	4026.1




BCY6009 (60.24 mg, 19.91 μmol , 1.00 eq) was used as the bicyclic reagent. BCY6027 (20.40 mg, 5.11 μmol , 25.69% yield, 96.88% purity) was obtained as a white solid.

BCY6027 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	35-65% B over 20 minutes, then 3 min 95% B
Retention Time:	5.97 min
LCMS (ESI):	m/z 1932.1 [M+2H] ²⁺ , 1282.5 [M+3H-H ₂ O] ³⁺
Peptide mw	3863.99




BCY6017 (61.81 mg, 27.38 μmol , 1.1 eq) was used as the bicycle reagent. BCY6047 (0.032 g, 10.34 μmol , 41.53% yield) was obtained as a white solid.

BCY6047 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	38-68% B over 30 minutes, then 3 min 95% B
Retention Time:	12.28 min
LCMS (ESI):	m/z 1026.3 [M+3H-H ₂ O] ³⁺
Peptide mw	3096.1




BCY6019 (115.22 mg, 32.86 μmol , 1.10 eq) was used as the bicycle reagent. BCY6035 (37.80 mg, 10.37 μmol , 34.73% yield) was obtained as a white solid.

BCY6035 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	35-65% B over 20 minutes, then 3 min 95% B
Retention Time:	12.28 min
LCMS (ESI):	m/z 1208.8 [M+3H-H ₂ O] ³⁺ , 911.5 [M+4H] ⁴⁺
Peptide mw	3643.73




BCY6018 (73.48 mg, 27.38 μmol , 1.1 eq) was used as the bicycle reagent. BCY6051 (0.0582 g, 16.52 μmol , 66.39% yield) was obtained as a white solid.

BCY6051 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28 - 68% B over 30 minutes, then 3 min 95% B
Retention Time:	11.37 min
LCMS (ESI):	m/z 880.5 [M+4H] ⁴⁺
Peptide mw	3522.57



BCY6152 (93.17 mg, 29.87 μmol , 1 eq) was used as the bicycle reagent. BCY6154 (40.10 mg, 9.93 μmol , 33.27% yield, 98.06% purity) was obtained as a white solid.



BCY6154 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28 - 68% B over 30 minutes, then 3 min 95% B
Retention Time:	11.94 min
LCMS (ESI):	m/z 1313.8 [M+3H-H ₂ O] ³⁺ , 985.8 [M+4H-H ₂ O] ⁴⁺
Peptide mw	3958.02



BCY6153 (82.55 mg, 29.87 μmol , 1 eq) was used as the bicycle reagent. BCY6155 (0.0312 g, 8.55 μmol , 28.62% yield, 98.69% purity) was obtained as a white solid.

BCY6155 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28 - 68% B over 30 minutes, then 3 min 95% B
Retention Time:	12.93 min
LCMS (ESI):	m/z 897.1 [M+4H-H ₂ O] ⁴⁺
Peptide mw	3602.63

5 DM1-SS(SO₃H)- Series
DM1-SPDP(SO₃H)-NHS Linker




To a solution of 4-sulfanylbutoanoic acid (2.0 g, 16.64 mmol, 1 eq) and 2-(2-pyridyldisulfanyl) pyridine (11.0 g, 49.93 mmol, 3 eq) in EtOH (50 mL) was added AcOH (1.05 g, 17.48 mmol, 1 mL, 1.05 eq). The mixture was stirred at 40 °C for 16 hr under N₂. LC-MS showed one main peak with desired mass was detected and TLC indicated 4-sulfanylbutoanoic acid was consumed completely. The reaction mixture was concentrated under reduced pressure to remove solvent to give a residue. The residue was purified by preparative HPLC (neutral condition). Compound **2** (2.0 g, 8.72 mmol, 52.4% yield) was obtained as yellow solid.

¹H NMR: 400 MHz CDCl₃

δ ppm 2.03-2.11 (m, 2 H), 2.54 (t, J=7.20 Hz, 2 H), 2.88 (t, J=7.20 Hz, 2 H), 7.11-7.14 (m, 1 H), 7.67-7.74 (m, 2 H), 8.50 (d, J=4.80 Hz, 1 H).

LCMS (ESI):	230 [M+H] ⁺
Molecular weight	229.31

Compound 3




To a solution of compound **2** (0.5 g, 2.18 mmol, 1 eq) in DCE (5 mL) was added chlorosulfonic acid (1.5 g, 13.08 mmol, 6 eq) in three portions and DIEA (1.13 g, 8.72 mmol, 4 eq) in two portions. The mixture was stirred at 75 °C for 2 hr. LC-MS showed compound **2** was consumed completely and one main peak with desired mass was detected. The reaction mixture was quenched by addition 3 mL of H₂O and the DCE was removed. The residue was directly purified by preparative HPLC (neutral conditions). Compound **3** (0.68 g, 1.76 mmol, 80.6% yield, 80% purity) was obtained as light yellow oil.

¹H NMR: 400 MHz CDCl₃

δ ppm 2.49-2.54 (m, 2 H), 3.63-3.67 (m, 2 H), 3.90 (t, J=6.60 Hz, 2 H), 7.09-7.12 (m, 1 H), 7.66-7.76 (m, 2 H), 8.47 (dd, J=4.80 Hz, 0.80 Hz, 1 H), 8.56 (s, 1 H).


LCMS (ESI):	310.0 [M+H] ⁺
Molecular weight	309.37

DM1-SO₃H-SPDB**DM1-SO₃H-SPDB**

To a solution of **DM1** (1.0 g, 1.35 mmol, 1 eq) and compound **3** (502.9 mg, 1.63 mmol, 1.2 eq) in DMF (10 mL) was added NaHCO₃ (aq) until the pH reached 8. The mixture was stirred at 25 °C for 1 hr. LC-MS showed DM1 was consumed completely and one main peak with desired mass was detected. The residue was directly purified by preparative HPLC (neutral condition). Compound **DM1-SO₃H-SPDB** (0.28 g, 299.0 μmol, 22.1% yield) was obtained as a white solid.

LCMS (ESI):	918.2 [M+H ₂ O] ⁺
Molecular weight	936.50

DM1-SO₃H-SPDB-NHS




To a solution of **DM1-SO₃H-SPDB** (103.2 mg, 896.95 μ mol, 3 eq), 1-Hydroxypropylidene -2,5-dione (103.2 mg, 896.95 μ mol, 3 eq) in DMA (6 mL) and DCM (2 mL) was added EDCI (171.9 mg, 896.95 μ mol, 3 eq). The mixture was stirred at 25 °C for 16 hr. LC-MS showed DM1-SO₃H-SPDB was consumed completely and one main peak with desired mass was detected. DCM was removed. The residue was The mixture was directly purified by preparative HPLC (neutral condition). Compound **DM1-SO₃H-SPDB-NHS** (0.22 g, 212.85 μ mol, 71.2% yield) was obtained as a white solid.

LCMS (ESI):	1015.2 [M+H-H ₂ O] ⁺
Molecular weight	1033.57

General Procedure for Coupling DM1-SO₃H-SPDB-NHS with Targeting Bicycles


To a solution of targeting Bicycle (1.1-1.3 eq) in DMA was added DIEA (3 eq) and DM1-SO₃H-SPDB-NHS (1 eq). The mixture was stirred at 25 °C for 16 hr. The reaction was monitored by LC-MS and once complete, the mixture was directly purified by preparative HPLC.

BCY6173




BCY6099 (200.15 mg, 62.89 μmol) was used as the bicycle reagent. 57.1 mg compound BCY6173 (3.40 μmol , 22.79% yield, 95.80% purity) was obtained as a white solid.

BCY6173 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	10.30 min
LCMS (ESI):	m/z 1361.9 [M+3H-H ₂ O] ³⁺ , 1021.8 [M+4H-H ₂ O] ⁴⁺
Peptide mw	4101.15




BCY6014 (711.9 mg, 234.14 μmol) was used as the bicycle reagent. 308 mg compound BCY6082 (74.97 μmol , 35.2% yield, 96.36% purity) was obtained as a white solid.

BCY6082 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	11.95 min
LCMS (ESI):	m/z 1299.3 [M+3H-H ₂ O] ³⁺ , 975.0 [M+4H-H ₂ O] ⁴⁺
Peptide mw	3911.04




BCY6018 (77.91 mg, 29.03 μmol , 1 eq) was used as the bicycle reagent. BCY6150 (0.0249 g, 6.61 μmol , 22.78% yield) was obtained as a white solid.

BCY6150 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H2O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	12.31 min
LCMS (ESI):	m/z 1195.4 [M+3H-H2O] ³⁺
Peptide mw	3602.63




BCY6104 (120.17 mg, 37.73 μmol , 1.3 eq) was used as the bicycle reagent. BCY6151 (0.0256 g, 6.16 μmol , 21.22% yield) was obtained as a white solid.

BCY6151 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H2O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	8.68 min
LCMS (ESI):	m/z 1362.3 [M+3H-H2O] ³⁺
Peptide mw	4105.16



BCY6138 (82.80 mg, 26.61 μmol , 1.1 eq) was used as the bicycle reagent. BCY6162 (0.0362 g, 8.98 μmol , 37.13% yield) was obtained as a white solid.



BCY6162 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	21.09 min
LCMS (ESI):	m/z 1323.5 [M+3H-H ₂ O-44] ³⁺
Peptide mw	4026.74



BCY6137 (79.67 mg, 24.48 μmol , 1.1 eq) was used as the bicycle reagent. BCY6161 (0.0232 g, 5.26 μmol , 21.76% yield) was obtained as a white solid.

BCY6161 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1% TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	10.22 min
LCMS (ESI):	m/z 1392 [M+3H-H ₂ O] ³⁺
Peptide mw	4192.33

5 DM1-SS-Me Series
DM1-SS-Me Linker


**SPP****1****SPP**

To a solution of 2-(2-pyridyldisulfanyl)pyridine (2.46 g, 11.18 mmol, 1.50 eq) and AcOH (1.05 g, 17.49 mmol, 1.00 mL, 2.35 eq) in EtOH (50.00 mL) was added 4-sulfanylpentanoic acid (1.00 g, 7.45 mmol, 1.00 eq). The mixture was stirred at 40 °C for 18 hours under N₂. LC-MS showed compound **1** was consumed completely and one main peak with desired mass was detected. The reaction mixture was concentrated under reduced pressure to remove solvent to give a residue. The residue was purified by preparative HPLC (neutral condition). Compound **SPP** (1.61 g, 6.62 mmol, 88.81% yield) was obtained as a yellow solid.

¹H NMR: 400 MHz DMSO-d₆

δ ppm 1.36 (d, J=6.78 Hz, 3 H), 1.88 - 2.07 (m, 2 H), 2.56 (td, J=7.53, 1.76 Hz, 2 H), 3.00 - 3.09 (m, 1 H), 7.11 (ddd, J=7.34, 4.96, 1.00 Hz, 1 H), 7.66 (td, J=7.78, 1.76 Hz, 1 H), 7.73 - 7.77 (m, 1 H), 8.48 (dt, J=4.02, 0.88 Hz, 1 H).


LCMS (ESI):	243.8 [M+H] ⁺
Molecular weight	243.34

DM1-SPP

A solution of DM1 (200 mg, 270.90 μmol , 1.00 eq), 4-(2-pyridyl)disulfanyl)pentanoic acid (98.89 mg, 406.35 μmol , 1.50 eq) in H_2O (5.00 mL) was adjusted $\text{pH} = 8$ using NaHCO_3 (aq). The mixture was stirred at 15°C for 1 hour. LC-MS showed DM1 was consumed completely and one main peak with desired mass was detected (main MS was $\text{M}+1-18$). The mixture was directly purified by preparative HPLC (neutral condition). Compound **DM1-SPP** (120 mg, 137.86 μmol , 50.89% yield) was obtained as a white solid.

LCMS (ESI):	852.0 $[\text{M}+\text{H}-\text{H}_2\text{O}]^+$
Molecular weight	870.47

DM1-SPP-TFP




To a solution of DM1-SPP (0.175 g, 201.04 μmol , 1.0 eq), 2,3,5,6-tetrafluorophenol (100.16 mg, 603.13 μmol , 3.0 eq) in DCM (1.0 mL) and DMA (3.0 mL) was added EDCI (115.62 mg, 603.13 μmol , 3.0 eq). The mixture was stirred at 15 °C for 12 hour. LC-MS showed DM1-SPP was consumed completely and one main peak with desired MS was detected. The DCM was removed and the residue was purified by prep-HPLC (neutral condition). Compound **DM1-SPP-TFP** (0.123 g, 120.76 μmol , 60.07% yield) was obtained as a white solid.

LCMS (ESI):	999.9 [M+H-H ₂ O] ⁺
Molecular weight	1018.53

General Procedure for Coupling DM1-SPP-TFP with Targeting Bicycles

To a solution of targeting Bicycle (1.1-1.3 eq) in DMA was added DIEA (3 eq) and DM1-SPP-TFP (1 eq). The mixture was stirred at 25 °C for 16 hr. The reaction was monitored by LC-MS and once complete, the mixture was directly purified by preparative HPLC.


BCY6032



To a solution of DM1-SPP (30.00 mg, 34.46 μmol , 1.00 eq) in DMF (5.00 mL) was added DIEA (13.36 mg, 103.38 μmol , 3.00 eq) and HATU (13.10 mg, 34.46 μmol , 1.00 eq). After 1 h, BYC6014 (104.79 mg, 34.46 μmol , 1.00 eq) was added and the mixture was stirred at 15 °C for 2 hours. LC-MS showed 40% of DM1-SPP was remained. Several new peaks were observed on LC-MS and 20% of desired compound was detected. The mixture was directly purified by preparative HPLC (TFA condition). Compound **BYC6032** (10.00 mg, 2.57 μmol , 7.45% yield) was obtained as a white solid.


BYC6032 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	25-55% B over 20 minutes, then 3 min 95% B
Retention Time:	13.38 min
LCMS (ESI):	m/z 1292.1 [M+3H-H ₂ O] ³⁺ , 969.0 [M+4H-H ₂ O] ⁴⁺
Peptide mw	3892.94

BCY6052



BCY6018 (86.96 mg, 32.40 μmol , 1.1 eq) was used as the bicycle reagent. BCY6052 (32.30 mg, 9.13 μmol , 31.01% yield) was obtained as a white solid.


BCY6052 Analytical Data	
Mobile Phase:	A: 0.1% Formic acid in H ₂ O B: ACN
Flow:	1.0ml/min
Column:	Eclipse XDB-Phenyl 3.5 μm 100*3.0mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	35-65% B over 20 minutes, then 3 min 95% B
Retention Time:	6.96 min
LCMS (ESI):	m/z 1173.4 [M+3H-H ₂ O] ³⁺ , 884.6 [M+4H] ⁴⁺
Peptide mw	3536.58



BCY6017 (66.50 mg, 29.45 μmol, 1.2 eq) was used as the bicycle reagent. BCY6048 (40.80 mg, 13.12 μmol, 53.45% yield) was obtained as a white solid.

BCY6048 Analytical Data	
Mobile Phase:	A: 0.1% Formic acid in H2O B: ACN
Flow:	1.0ml/min
Column:	Eclipse XDB-Phenyl 3.5um 100*3.0mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	35-65% B over 20 minutes, then 3 min 95% B
Retention Time:	7.56 min
LCMS (ESI):	m/z 1031.0 [M+3H-H2O] ³⁺ , 884.6 [M+4H] ⁴⁺
Peptide mw	3110.13


BCY6036



BCY6019 (113.60 mg, 32.40 μmol , 1.10 eq) was used as the bicycle reagent. BCY6036 (53.20 mg, 14.00 μmol , 47.54% yield, 96.26% purity) was obtained as a white solid.

BCY6036 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1% TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	35-65% B over 20 minutes, then 3 min 95% B
Retention Time:	8.19 min
LCMS (ESI):	m/z 1213.6 [M+3H-H ₂ O] ³⁺ , 914.7 [M+4H] ⁴⁺
Peptide mw	3657.76


BCY6028




BCY6009 (99.00 mg, 29.45 μmol , 1.00 eq) was used as the bicycle reagent. BCY6028 (24.30 mg, 6.05 μmol , 20.56% yield, 96.61% purity) was obtained as a white solid.

BCY6028 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1% TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	35-65% B over 20 minutes, then 3 min 95% B
Retention Time:	6.43 min
LCMS (ESI):	m/z 965.6 [M+4H-H ₂ O] ⁴⁺
Peptide mw	3877.96

5 Disulfide Linkers (Various Hindrances)
BCY6039 (DM1-Me-SS-Me-Bicycle)




Compound 2A



To a solution of 2-(2-pyridyldisulfany)pyridine (2.46 g, 11.18 mmol, 1.50 eq) and AcOH (1.05 g, 17.49 mmol, 1.00 mL, 2.35 eq) in EtOH (50.00 mL) was added 4-sulfanylpentanoic acid (**1A**) (1.00 g, 7.45 mmol, 1.00 eq). The mixture was stirred at 40 °C for 18 hours under N₂. LC-MS showed **1A** was consumed completely and one main peak with the desired mass was detected. The reaction mixture was concentrated under reduced pressure to give a residue, which was purified by preparative HPLC (neutral condition). Compound **2A** (1.61 g, 6.62 mmol, 88.81% yield) was obtained as a light yellow solid.

LCMS (ESI):	243.9 [M+H] ⁺
Molecular weight	243.34


10 Compound 3A



To a solution of **2A** (0.01 g, 41.09 μmol , 1.00 eq), 1-hydroxypyrrolidine-2,5-dione (14.19 mg, 123.28 μmol , 3.00 eq) in DMA (1 mL) was added EDCI (23.63 mg, 123.28 μmol , 3.00 eq). The mixture was stirred at 15 °C for 16 hr. LC-MS showed **2A** was consumed completely and one main peak with desired mass was detected. The residue was purified by preparative HPLC (neutral condition). Compound **3A** (0.011 g, 32.31 μmol , 78.63% yield) was obtained as a white solid.

LCMS (ESI):	340.8 [M+H] ⁺
Molecular weight	340.41


Compound 4A



To a solution of BCY6014 (98.25 mg, 32.31 μmol , 1.00 eq) in DMA (3 mL) was added DIEA (8.26 mg, 64.62 μmol , 11.26 μL , 2.00 eq) and 3A (0.011 g, 32.31 μmol , 1.00 eq). The mixture was stirred at 15 °C for 18 hr. LC-MS showed 3A was consumed completely and one main peak with desired mass was detected. The mixture was directly purified by preparative HPLC (neutral condition). Compound **4A** (0.04 g, 12.25 μmol , 37.90% yield) was obtained as a white solid.

LCMS (ESI):	1088.7 [M+3H] ³⁺ , 816.5 [M+4H] ⁴⁺
Molecular weight	3264.88

Compound 5A




To a solution of 4A (0.04 g, 12.25 μmol , 1.00 eq) in MeCN (4 mL) and H_2O (2 mL) was added TCEP (4.21 mg, 14.70 μmol , 4.05 μL , 1.20 eq). The mixture was stirred at 15 $^\circ\text{C}$ for 1 hr. LC-MS showed 4A was consumed completely and one main peak with the desired mass was detected.

The residue was purified by preparative HPLC (neutral condition). Compound **5A** (0.035 g, 11.09 μmol , 90.53% yield) was obtained as a white solid.

LCMS (ESI):	1052.2 $[\text{M}+3\text{H}]^{3+}$
Molecular weight	3155.73

DM3-SPy




5

To a solution of 4-(2-pyridyldisulfanylpentanoic acid (**2A**) (22.46 mg, 92.29 μmol , 1.20 eq), HATU (35.09 mg, 92.29 μmol , 1.20 eq), DIEA (29.82 mg, 230.71 μmol , 40.19 μL , 3.00 eq) in DMF (5 mL) was added **1B** (0.05 g, 76.90 μmol , 1.00 eq). The mixture was stirred at 15 °C for 1 hr. LC-MS showed **1B** was consumed completely and one main peak with the desired mass was detected. The residue was purified by preparative HPLC (neutral condition). Compound **DM3-SPy** (0.025 g, 28.56 μmol , 37.13% yield) was obtained as a white solid.

LCMS (ESI):	875.1 $[\text{M}+\text{H}]^+$
Molecular weight	875.49

10


BCY6039




A solution of DM3-SPy (0.015 g, 17.13 μmol , 1.00 eq) and 5A (54.08 mg, 17.13 μmol , 1.00 eq) in DMF (3 mL) was adjusted to pH=8 using $\text{NaHCO}_3(\text{aq})$. The mixture was stirred at 15 $^\circ\text{C}$ for 1 hr. LC-MS showed DM3-SPy was consumed completely and one main peak with desired mass was detected. The mixture was directly purified by preparative HPLC (TFA condition). Compound **BCY6039** (0.0263 g, 6.58 μmol , 38.39% yield) was obtained as a white solid.

BCY6039 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H2O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE (1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	13.01 min
LCMS (ESI):	m/z 976.1 $[\text{M}+4\text{H}-\text{H}_2\text{O}]^{4+}$
Peptide mw	3921.01


BCY6055 (DM1-SS-Me₂-Bicycle)



Compound 2

To a solution of compound **1** (0.045 g, 126.96 μmol , 1 eq) in H_2O (1 mL) was adjusted pH = 13 using 1 N NaOH solution. The mixture was stirred at 15 °C for 16 hr. LC-MS showed compound **1** was consumed completely and one main peak with the desired mass was detected. The residue was purified by preparative HPLC (neutral condition). Compound **2** (0.03 g, 116.56 μmol , 91.81% yield) was obtained as a yellow solid.


LCMS (ESI):	257.9 $[\text{M}+\text{H}]^+$
Molecular weight	257.37

Compound 3

A solution of compound **2** (0.03, 116.56 μmol , 1.0 eq) and DM1 (111.87 mg, 151.53 μmol , 1.3 eq) in DMF (5 mL) was stirred at 15 °C for 2 hours. LC-MS showed DM1 was consumed completely and one main peak with desired mass was detected. The mixture was directly purified by preparative HPLC (NH_4HCO_3 condition). Compound **3** (0.05 g, 56.53 μmol , 48.50% yield) was obtained as a white solid.

LCMS (ESI):	866.0 $[\text{M}+\text{H}-\text{H}_2\text{O}]^+$
Molecular weight	884.49


5 Compound 4



To a solution of compound **3** (0.05 g, 56.53 μmol , 1.0 eq) and 2,3,5,6-tetrafluorophenol (28.16 mg, 169.59 μmol , 3.0 eq) in DMA (3 mL) and DCM (1 mL) was added EDCI (32.51 mg, 169.59 μmol , 3 eq). The mixture was stirred at 15 °C for 16 hr. LC-MS showed compound **3** was consumed completely and one main peak with desired mass was detected. DCM was removed and the mixture was directly purified by preparative HPLC (neutral condition). Compound **4** (0.03 g, 29.05 μmol , 51.40% yield) was obtained as a white solid.

LCMS (ESI):	1014.0 $[\text{M}+\text{H}-\text{H}_2\text{O}]^+$
Molecular weight	1032.55


BCY6055



To a solution of BCY6014 (106.01 mg, 34.87 μmol , 1.2 eq) in DMA (3 mL) was added DIEA (11.27 mg, 87.16 μmol , 15.18 μL , 3.0 eq) and compound **4** (0.03 g, 29.05 μmol , 1.0 eq). The mixture was stirred at 15 °C for 16 hr. LC-MS showed compound **4** was consumed completely and one main peak with desired mass was detected. The mixture was directly purified by preparative HPLC (TFA condition). Compound **BCY6055** (0.0352 g, 9.01 μmol , 31.01% yield) was obtained as a white solid.

BCY6055 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE (1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	12.68 min
LCMS (ESI):	m/z 1296.1 [M+3H-H ₂ O] ³⁺ , 972.4 [M+4H-H ₂ O] ⁴⁺
Peptide mw	3906.98

BCY6077 (DM1-SS-Me-(SO₃H)-Bicycle)



General procedure for preparation of compound 2



1


2

To a solution of compound 1 (0.1 g, 410.94 μmol , 1 eq) in 1,2-dichloroethane (3 mL) was added sulfurochloric acid (0.86 g, 7.38 mmol, 491.43 μL , 17.96 eq) on three parts and DIEA (318.67 mg, 2.47 mmol, 429.47 μL , 6 eq) was added on two parts. The mixture was stirred at 75 °C for 16 hr. LC-MS showed compound 1 was consumed completely and one main peak with desired MS was detected MS324, one main peak of byproduct MS 221 was PySSPy. The solvent was removed and dissolved in H₂O/MeCN=15/1. Directly purified by prep-HPLC (neutral condition: MeCN/H₂O). Compound 2 (0.055 g, 170.06 μmol , 41.38% yield) was obtained as a yellow oil.

5

LCMS (ESI):	323.6 [M+H] ⁺
Molecular weight	323.4

General procedure for preparation of DM1-SO₃H-SPP




To a solution of DM1 (113.00 mg, 153.06 μmol , 1.1 eq), compound 2 (0.045 g, 139.14 μmol , 1 eq) in DMF (2 mL) was adjusted PH=8 used for NaHCO₃(aq). The mixture was stirred at 15 °C for 1 hr. LC-MS showed DM1 was consumed completely and one main peak with desired MS was detected. Directly purified by prep-HPLC (neutral condition). Compound DM1-SO₃H-SPP (0.075 g, 78.90 μmol , 56.71% yield) was obtained as a white solid.

10

LCMS (ESI):	931.9 [M+H-H ₂ O] ⁺
-------------	---

Molecular weight	950.52
------------------	--------

General procedure for preparation of DM1-SO3H-SPP-NHS



To a solution of DM1-SO3H-SPP (0.06 g, 63.12 μmol , 1 eq), 1-hydroxypropylidene-2,5-dione (7.99 mg, 69.43 μmol , 1.1 eq) in DMA (1.5 mL) and DCM (0.5 mL) was added EDCI (13.31 mg, 69.43 μmol , 1.1 eq). The mixture was stirred at 15 °C for 18 hr. LC-MS showed DM1-SO3H-SPP was consumed completely and one main peak with desired MS was detected. Directly purified by prep-HPLC (neutral condition: MeCN/H₂O). Compound DM1-SO3H-SPP-NHS (0.045 g, 42.96 μmol , 68.05% yield) was obtained as a white solid.


LCMS (ESI):	984 [M-NHS+K] ⁺
Molecular weight	1047.6

General procedure for preparation of BCY6077


To a solution of BCY6014 (101.58 mg, 33.41 μmol , 1 eq) in DMA (1 mL) was added DIEA (12.95 mg, 100.23 μmol , 17.46 μL , 3 eq) and DM1-SO3H-SPP-TFP (0.035 g, 33.41 μmol , 1 eq). The mixture was stirred at 15 °C for 16 hr. LC-MS showed DM1-SO3H-SPP-TFP was consumed completely and one main peak with desired MS was detected. Directly purified by prep-HPLC (TFA condition). Compound BCY6077 (41.30 mg, 10.03 μmol , 30.01% yield, 96.44% purity) was obtained as a white solid.

BCY6077 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H2O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE (1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	11.80 min
LCMS (ESI):	m/z 978.2 [M+4H-18-44] ⁴⁺
Peptide mw	3972.06

Non-Cleavable Series
BCY6063 (MMAE)




5 Glutarate-MMAE



To a solution of MMAE (0.2 g, 278.56 μmol , 1.0 eq) in DMA (3 mL) was added DIEA (108.01 mg, 835.68 μmol , 145.56 μL , 3.0 eq) and tetrahydropyran-2,6-dione (63.57 mg, 557.12 μmol , 2.0 eq). The mixture was stirred at 15 °C for 16 hr. LC-MS showed MMAE was consumed completely and one main peak with desired mass was detected. The mixture was directly purified by preparative HPLC (neutral condition). Compound Glutarate-MMAE (0.12 g, 144.22 μmol , 51.77% yield) was obtained as a white solid.

LCMS (ESI):	832.3 [M+H] ⁺
Molecular weight	832.09


Glutarate-MMAE-NHS



To a solution of Glutarate-MMAE (0.12 g, 144.22 μmol , 1.0 eq), 1-hydroxypyrrolidine-2, 5-dione (49.79 mg, 432.65 μmol , 3.0 eq) in DMA (3 mL) and DCM (1 mL) was added EDCI (82.94 mg, 432.65 μmol , 3.0 eq). The mixture was stirred at 15 °C for 16 hr. LC-MS showed Glutarate-MMAE was consumed completely and one main peak with desired mass was detected. The mixture was directly purified by preparative HPLC (TFA condition). Compound Glutarate-MMAE-NHS (0.055 g, 59.19 μmol , 41.04% yield) was obtained as a white solid.

LCMS (ESI):	929.2 [M+H] ⁺
Molecular weight	929.17


BCY6063




To a solution of BCY6014 (98.17 mg, 32.29 μmol , 1.2 eq) in DMA (2 mL) were added DIEA (10.43 mg, 80.72 μmol , 14.06 μL , 3 eq) and Glutarate-MMAE-NHS (0.025 g, 26.91 μmol , 1 eq). The mixture was stirred at 15 $^{\circ}\text{C}$ for 16 hr. LC-MS showed Glutarate-MMAE-NHS was consumed completely and one main peak with desired mass was detected. The mixture was directly purified by preparative HPLC (TFA condition). Compound **BCY6063** (32.10 mg, 8.33 μmol , 30.95% yield) was obtained as a white solid.

BCY6063 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min
Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	10.86 min
LCMS (ESI):	m/z 963.8 [M+4H] ⁴⁺ , 771.1 [M+5H] ⁵⁺
Peptide mw	3854.56

BCY6064 (DM1)






To a solution of DM1 (0.1 g, 135.45 μmol , 1 eq), 3-[(2-bromoacetyl)amino]propanoic acid (34.14 mg, 162.54 μmol , 1.2 eq) in DMF (5 mL) was added TEA (41.12 mg, 406.35 μmol , 56.56 μL , 3 eq). The mixture was stirred at 15 °C for 1 hr. LC-MS showed DM1 was consumed completely and one main peak with the desired mass was detected. The mixture was directly purified by preparative HPLC (neutral conditions). Compound 1 (0.08 g, 92.23 μmol , 68.09% yield) was obtained as a white solid.

LCMS (ESI):	849.1 [M+H-H ₂ O] ⁺
Molecular weight	867.41


Compound 2



To a solution of compound **1** (0.08 g, 92.23 μmol , 1 eq), 2,3,5,6-tetrafluorophenol (45.95 mg, 276.69 μmol , 3 eq) in DMA (3 mL) and DCM (1 mL) was added EDCI (53.04 mg, 276.69 μmol , 3 eq). The mixture was stirred at 15 °C for 4 hr. LC-MS showed compound **1** was consumed completely and one main peak with desired mass was detected. The mixture was directly purified by preparative HPLC (neutral condition). Compound **2** (0.06 g, 59.09 μmol , 64.06% yield) was obtained as a white solid.

LCMS (ESI):	997.0 [M+H-H ₂ O] ⁺
Molecular weight	1015.46

5

BCY6064**2**


To a solution of BCY6014 (107.79 mg, 35.45 μmol , 1.2 eq) in DMA (3 mL) was added DIEA (11.45 mg, 88.63 μmol , 15.44 μL , 3.0 eq) and compound **2** (0.030 g, 29.54 μmol , 1 eq). The mixture was stirred at 15 °C for 16 hr. LC-MS showed compound **2** was consumed completely and one main peak with desired mass was detected. The mixture was directly purified by preparative HPLC (TFA condition). Compound **BCY6064** (28.40 mg, 7.30 μmol , 24.71% yield) was obtained as a white solid.

10

BCY6064 Analytical Data	
Mobile Phase:	A: 0.1% TFA in H ₂ O B: 0.1%TFA in ACN
Flow:	1.0ml/min

Column:	Gemini-NX C18 5um 110A 150*4.6mm
Instrument:	Agilent 1200 HPLC-BE(1-614)
Method:	28-68% B over 30 minutes, then 3 min 95% B
Retention Time:	10.26 min
LCMS (ESI):	m/z 968.4 [M+4H-H2O] ⁴⁺
Peptide mw	3889.89

BCY6105



General procedure for preparation of Compound 2

To a solution of compound **1** (3.5 g, 5.68 mmol, 1.0 eq) in DCM (20 mL) and MeOH (10 mL), (4-aminophenyl)methanol (978.5 mg, 7.95 mmol, 1.4 eq) and EEDQ (2.81 g, 11.35 mmol, 2.0 eq) were added in the dark, and the mixture was stirred at 25 °C for 18 hr. LC-MS showed compound **1** was consumed completely and one main peak with desired MS was detected ($[M+H]^+=722.0$). The resulting reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 220g SepaFlash®

Silica Flash Column, Eluent of 0~10% Methanol/Dichloromethane @ 80 mL/min). Compound **2** (3.0 g, 4.16 mmol, 73.2% yield) was obtained as a yellow solid.

General procedure for preparation of Compound 3

To a solution of compound **2** (2.5 g, 3.46 mmol, 1.0 eq) in THF (30 mL) was added DIEA (2.69 g, 20.78 mmol, 3.62 mL, 6.0 eq) and bis(4-nitrophenyl) carbonate (6.32 g, 20.78 mmol, 6.0 eq), and the mixture was stirred at 25 °C for 16 hr. TLC indicated compound **2** was consumed completely and one new spot formed. The reaction was clean according to TLC. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 220 g SepaFlash® Silica Flash Column, Eluent of 0~5% Methanol/Dichloromethane @ 100 mL/min). Compound **3** (2.2 g, 2.48 mmol, 71.6% yield) was obtained as a yellow solid.

10

General procedure for preparation of Compound 4

To a solution of compound **3** (0.3 g, 338.24 μmol, 1.0 eq) in DMF (5 mL), HOBt (50.3 mg, 372.06 μmol, 1.1 eq), DIEA (131.1 mg, 1.01 mmol, 176.7 μL, 3.0 eq), and MMAE (218.6 mg, 304.42 μmol, 0.9 eq) were added. The mixture was stirred at 40 °C for 16 hr. LC-MS showed one peak with desired MS ($[M+H]^+$)=1466.4, $[M+2H]^+/2=733.2$). The reaction mixture was then directly purified by prep-HPLC (neutral condition), and compound **4** (0.2 g, 136.44 μmol, 40.3% yield) was obtained as a white solid.

15

General procedure for preparation of Compound 5

Compound **4** (0.175 g, 119.39 μmol, 1.0 eq) was first dissolved in TFA (1.8 mL), and then triisopropylsilane (13.5 g, 85.20 mmol, 17.5 mL, 713.7 eq) was added. The mixture was stirred at 0 °C for 30 min. LC-MS showed one peak with desired MS ($[M+H]^+$)=1123.4, $[M+2H]^+/2=562.2$). The reaction mixture was concentrated under reduced pressure to remove solvent to give a residue. The residue was purified by prep-HPLC (neutral condition). Compound **5** (0.1 g, 89.02 μmol, 74.6% yield) was obtained as a yellow solid.

20

General procedure for preparation of Compound 6

To a solution of compound **5** (0.07 g, 62.31 μmol , 1.0 eq) in DMA (1.0 mL), DIEA (24.2 mg, 186.94 μmol , 32.6 μL , 3.0 eq) and tetrahydropyran-2,6-dione (14.2 mg, 124.62 μmol , 2.0 eq) were added. The mixture was stirred at 25 °C for 2 hr. LC-MS showed compound **5** was consumed completely and one main peak with desired MS was detected ($[\text{M}+\text{H}]^+=1237.4$, $[\text{M}+2\text{H}]^+/2=619.3$). The reaction mixture was then directly purified by prep-HPLC (neutral condition), and compound **6** (0.04 g, 32.32 μmol , 51.8% yield) was obtained as a light yellow solid.

5

General procedure for preparation of Compound 7

To a solution of compound **6** (0.04 g, 32.32 μmol , 1.0 eq), 1-hydroxypyrrolidine-2,5-dione (11.2 mg, 96.97 μmol , 3.0 eq) in DMA (3.0 mL) and DCM (1.0 mL), EDCI (18.6 mg, 96.97 μmol , 3.0 eq) was added. The mixture was stirred at 25 °C for 18 hr. LC-MS showed compound **6** was consumed completely and one main peak with desired MS was detected ($[\text{M}+\text{H}]^+=1334.5$, $[\text{M}+2\text{H}]^+/2=667.7$). The reaction mixture was then directly purified by prep-HPLC (TFA condition), and compound **7** (0.025 g, 18.73 μmol , 57.9% yield) was obtained as a white solid.


10

General procedure for preparation of BCY6105

To a solution of BCY6014 (82.0 mg, 26.98 μmol , 1.2 eq) in DMA (4 mL), DIEA (8.7 mg, 67.44 μmol , 11.7 μL , 3.0 eq) and compound **7** (0.03 g, 22.48 μmol , 1.0 eq) were added. The mixture was stirred at 25 °C for 18 hr. LC-MS showed compound **7** was consumed completely and one main peak with desired MS was detected ($[\text{M}+4\text{H}]^+/4=1065.2$). The reaction mixture was then directly purified by prep-HPLC (TFA condition). Compound **BCY6105** (0.024 g, 5.41 μmol , 24.1% yield, 96.06% purity) was obtained as a white solid.

15

BCY6106



General procedure for preparation of Compound 2

To a solution of compound **1** (5.0 g, 10.67 mmol, 1.0 eq) in DCM (30 mL) and MeOH (10 mL), EEDQ (5.28 g, 21.34 mmol, 2.0 eq) and (4-aminophenyl)methanol (2.63 g, 21.34 mmol, 2.0 eq) were added. The mixture was stirred at 20 °C for 18 hr. LC-MS showed compound **1** was consumed completely and one main peak with desired MS was detected (desired m/z=574, while Boc group falling off and partially falling off corresponded to m/z=474 and 518, respectively). The reaction mixture was concentrated under reduced pressure to remove solvent to give a residue. The residue was purified by prep-HPLC (neutral condition). Compound **2** (3.7 g, 6.45 mmol, 60.4% yield) was obtained as a yellow solid.

General procedure for preparation of Compound 3

To a solution of compound **2** (3.4 g, 5.93 mmol, 1.0 eq) in DMF (20 mL) was added DIEA (5.36 g, 41.49 mmol, 7.23 mL, 7.0 eq) and bis(4-nitrophenyl) carbonate (10.82 g, 35.56 mmol, 6.0 eq) in one part. The mixture was stirred at 25 °C for 2 hr. LC-MS showed one peak with desired MS (m/z=639 corresponded to the mass with Boc group falling off during ESI). The reaction mixture was directly purified by prep-HPLC (neutral condition). Compound **3** (3.0 g, 4.06 mmol, 68.5% yield) was obtained as a yellow solid.

General procedure for preparation of Compound 4

To a solution of compound **3** (707.4 mg, 957.55 umol, 1.0 eq) in DMF (15 mL), HOBt (155.3 mg, 1.15 mmol, 1.2 eq), DIEA (371.3 mg, 2.87 mmol, 500.4 µL, 3.0 eq), and MMAE (0.55 g, 766.04 umol, 0.8 eq) were added. The mixture was stirred at 30 °C for 16 hr. LC-MS showed one peak with desired MS (desired m/z=1317, and m/z=609 corresponded to the mass with two protons and Boc group falling off during ESI). The reaction mixture was then directly purified by prep-HPLC (neutral condition). Compound **4** (0.53 g, 402.23 umol, 42.0% yield) was obtained as a yellow solid.

General procedure for preparation of Compound 5

To a solution of compound **4** (0.526 g, 399.20 umol, 1.0 eq) in DMF (4 mL), piperidine (862.2 mg, 10.13 mmol, 1.0 mL, 25.4 eq) was added. The mixture was stirred at 25 °C for 30 min. LC-MS showed compound **4** was consumed completely and one main peak with desired MS was

detected (desired $m/z=1095$, and $m/z=265$ corresponded to Fmoc-piperidine adduct). The reaction mixture was then directly purified by prep-HPLC (neutral condition). Compound **5** (0.230 g, 209.97 μmol , 52.6% yield) was obtained as a white solid.

General procedure for preparation of Compound 6

5 To a solution of Fmoc-(D-Ala)-Phe-OH (125.6 mg, 273.87 μmol , 1.2 eq) in DMF (10 mL), EDCI (52.5 mg, 273.87 μmol , 1.2 eq), HOBt (37.0 mg, 273.87 μmol , 1.2 eq), and compound **5** (0.25 g, 228.23 μmol , 1 eq) were added. The mixture was stirred at 25 °C for 3 hr. LC-MS showed compound **5** was consumed completely and one peak with desired MS was detected ($m/z=718$ corresponded to the mass with two protons and Boc group falling off during ESI). The reaction mixture was then directly purified by prep-HPLC (neutral condition). Compound **6** (0.18 g, 117.20 μmol , 51.3% yield) was obtained as a white solid.

10

General procedure for preparation of Compound 7

To a solution of compound **6** (0.18 g, 117.20 μmol , 1.0 eq) in DMF (8 mL), piperidine (1.72 g, 20.25 mmol, 2.0 mL, 172.8 eq) was added. The mixture was stirred at 25 °C for 1 hr. LC-MS showed compound **7** was consumed completely and one main peak with desired MS was detected ($m/z=1314$ and 657 corresponded to the desired mass, and $m/z=265$ corresponded to Fmoc-piperidine adduct). The reaction mixture was then directly purified by prep-HPLC (neutral condition). Compound **7** (0.13 g, 98.96 μmol , 84.4% yield) was obtained as a white solid.

15

General procedure for preparation of Compound 8

To a solution of compound **7** (0.105 g, 79.93 μmol , 1.0 eq) in DMA (4 mL), DIEA (31.0 mg, 239.79 μmol , 41.8 μL , 3.0 eq) and tetrahydropyran-2,6-dione (27.4 mg, 239.79 μmol , 3.0 eq) were added. The mixture was stirred at 25 °C for 2 hr. LC-MS showed compound **7** was consumed completely and one main peak with desired MS was detected ($m/z=664.5$ corresponded to the mass with two protons and Boc group falling off during ESI). The reaction mixture was then directly purified by prep-HPLC (neutral condition), and compound **8** (0.09 g, 63.04 μmol , 78.8% yield) was obtained as a white solid.

20

General procedure for preparation of Compound 9

To a solution of compound **8** (0.09 g, 63.04 μmol , 1.0 eq), 1-hydroxypyrrolidine-2,5-dione (21.7 mg, 189.11 μmol , 3.0 eq) in DMA (3 mL) and DCM (1 mL), EDCI (36.2 mg, 189.11 μmol , 3.0 eq) dissolved in 1 mL DCM was added. The mixture was stirred at 25 °C for 18 hr. LC-MS showed compound **8** was consumed completely and one main peak with desired MS was detected (desired $m/z=1524$ (one proton) and 763 (two protons), while $m/z=713$ corresponded to the mass with Boc group falling off during ESI). The reaction mixture was directly purified by prep-HPLC (neutral condition). Compound **9** (0.07 g, 45.91 μmol , 72.8% yield) was obtained as a white solid.


General procedure for preparation of Compound 10

To a solution of BCY6014 (167.5 mg, 55.09 μmol , 1.2 eq) in DMF (3 mL), DIEA (11.8 mg, 91.81 μmol , 16.0 μL , 2.0 eq) and compound **9** (0.07 g, 45.91 μmol , 1.0 eq) were added. The mixture was stirred at 25 °C for 16 hr. LC-MS showed compound **9** was consumed completely and one main peak with desired MS was detected ($[M+4H]^+/4=1112.9$, $[M+5H]^+/5=890.5$). The reaction mixture was concentrated under reduced pressure to remove solvent to give a residue. The crude product **10** (0.220 g, crude) was used in the next step without further purification.

General procedure for preparation of BCY6106

To a solution of compound **10** (0.200 g, 44.95 μmol , 1.0 eq) in DCM (4 mL), 1 mL TFA was added. The mixture was stirred at 25 °C for 1 hr. LC-MS showed one main peak with desired MS ($[M+4H]^+/4=1088.0$, $[M+5H]^+/5=870.8$). The reaction mixture was concentrated under reduced pressure to give a residue, which was then directly purified by prep-HPLC (TFA condition). Compound **BCY6106** (0.0297 g, 20.06 μmol , 14.5% yield, 95.46% purity) was obtained as a white solid.

BCY6175



General procedure for preparation of Compound 9

The synthesis of **Compound 9** was performed in an analogous manner to that described in BCY6106.


General procedure for preparation of Compound 10A

To a solution of **BCY6099** (195.15 mg, 61.32 μmol , 1.1 eq) in DMA (3 mL) were added DIEA (21.61 mg, 167.23 μmol , 29.13 μL , 3 eq) and compound 9 (0.085 g, 55.74 μmol , 1.0 eq). The mixture was stirred at 25 °C for 16 hr. LC-MS showed compound 9 was consumed completely and one main peak with desired m/z was detected. The reaction mixture was concentrated under reduced pressure to remove solvent to afford a residue (light yellow oil). The reaction was directly purified by prep-HPLC (neutral condition). Compound **10A** (0.160 g, 34.84 μmol , 62.50% yield) was obtained as a white solid.

General procedure for preparation of BCY6175

To a solution of compound **10A** in DCM (4.5 mL) was added TFA (4.5 mL). The mixture was stirred at 0 °C for 30 min. LC-MS showed compound **10A** was consumed completely and one main peak with desired m/z was detected. The reaction mixture was concentrated under reduced pressure to remove solvent to afford a residue, which was purified by prep-HPLC (TFA condition). Compound **BCY6175** (61.40 mg, 13.56 μmol , 31.13% yield) was obtained as a white solid.

BCY6107



General procedure for preparation of Compound 2

To a solution of compound 1 (3.0 g, 8.49 mmol, 1.0 eq) in DCM (30 mL), EEDQ (2.52 g, 10.19 mmol, 1.2 eq) and (4-aminophenyl)methanol (1.25 g, 10.19 mmol, 1.2 eq) were added. The mixture was stirred at 25 °C for 16 hr. LC-MS showed compound 1 was consumed completely and one main peak with desired MS was detected ($[M+H]^+$ 459.5). In addition, TLC indicated compound 1 was consumed completely and new spots formed. The reaction mixture was concentrated under reduced pressure to remove solvent to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 120 g SepaFlash® Silica Flash Column, Eluent of 0~60% Ethylacetate/Petroleum ether gradient @ 80 mL/min). Compound 2 (3.5 g, 7.63 mmol, 89.9% yield) was obtained as a yellow solid.

General procedure for preparation of Compound 3

To a solution of compound **2** (3.3 g, 7.20 mmol, 1.0 eq) in THF (100 mL), DIEA (4.65 g, 35.98 mmol, 6.27 mL, 5.0 eq) and bis(4-nitrophenyl) carbonate (8.76 g, 28.79 mmol, 4.0 eq) were added. The mixture was stirred at 25 °C for 16 hr. LC-MS showed compound **2** was consumed completely and one main peak with desired MS was detected ($[M+H]^+$ 624.0). In addition, TLC indicated compound **2** was consumed completely and new spots formed. The reaction mixture was concentrated under reduced pressure to remove solvent to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 120 g SepaFlash® Silica Flash Column, Eluent of 0~15% Ethylacetate/Petroleum ether gradient @ 80 mL/min). Compound **3** (3.0 g, 4.81 mmol, 66.8% yield) was obtained as a yellow solid.

General procedure for preparation of Compound 4

To a solution of compound **3** (124.09 mg, 198.97 μ mol, 1.0 eq) in DMF (5 mL), HOBt (32.3 mg, 238.77 μ mol, 1.2 eq), DIEA (77.1 mg, 596.92 μ mol, 103.9 μ L, 3.0 eq), and MMAE (0.1 g, 139.28 μ mol, 0.7 eq) were added. The mixture was stirred at 25°C for 1 hr. LC-MS showed compound **3** was consumed completely and one main peak with desired MS was detected ($[M+H]^+$ 1202.5, $[M+Na]^+$ 1224.5). The reaction mixture was then directly purified by prep-HPLC (neutral condition). After lyophilization, compound **4** (0.08 g, 66.53 μ mol, 33.4% yield) was obtained as a white solid.

General procedure for preparation of Compound 5

To a solution of compound **4** (0.08 g, 66.53 μ mol, 1.0 eq) in DMF (4 mL), piperidine (862.2 mg, 10.13 mmol, 1 mL, 152.2 eq) was added. The mixture was stirred at 25 °C for 1 hr. LC-MS showed compound **4** was consumed completely and one main peak with desired MS was detected ($[M+H]^+$ 981.5, $[M+Na]^+$ 1003.5, while $m/z=264.0$ corresponded to Fmoc-piperidine adduct). The reaction mixture was directly purified by prep-HPLC (neutral condition). Compound **5** (0.055 g, 56.11 μ mol, 84.3% yield) was obtained as a white solid.

General procedure for preparation of Compound 6

To a solution of Fmoc-Glu(t-Bu)-Pro-Cit-Gly-HPhe-Tyr(t-Bu)-OH (74.1 mg, 66.31 μ mol, 1.3 eq) in DMF (4 mL), EDCI (12.7 mg, 66.31 μ mol, 1.3 eq), HOBt (8.9 mg, 66.31 μ mol, 1.3 eq), and compound **5** (0.05 g, 51.01 μ mol, 1.0 eq) were added. The mixture was stirred at 25 °C for 30 min. LC-MS indicated 20% of compound **5** remained, several new peaks formed, and 60% of the reaction mixture was desired product ($[M+2H^+]/2=1040.4$). The reaction mixture was directly purified by prep-HPLC (neutral condition). Compound **6** (0.07 g, 33.66 μ mol, 66.0% yield) was obtained as a white solid.

General procedure for preparation of Compound 7

To a solution of compound **6** (0.07 g, 33.66 μmol , 1.0 eq) in DMF (4 mL), piperidine (2.9 mg, 33.66 μmol , 3.3 μL , 1.0 eq) was added. The mixture was stirred at 25 °C for 15 min. LC-MS showed compound **6** was consumed completely and one main peak with desired MS was detected ($[\text{M}+2\text{H}^+]/2=929.1$, while $m/z=264.2$ corresponded to Fmoc-piperidine adduct). The reaction mixture was directly purified by prep-HPLC (neutral condition). Compound **7** (0.045 g, 24.23 μmol , 72.0% yield) was obtained as a white solid.

General procedure for preparation of Compound 8

To a solution of compound **7** (0.04 g, 21.54 μmol , 1.0 eq) in DMA (1 mL), DIEA (8.3 mg, 64.61 μmol , 11.2 μL , 3.0 eq) and tetrahydropyran-2,6-dione (7.4 mg, 64.61 μmol , 3.0 eq) were added. The mixture was stirred at 25 °C for 1 hr. LC-MS showed compound **7** was consumed completely and one main peak with desired MS was detected ($[\text{M}+2\text{H}^+]/2=986.4$). The reaction mixture was then directly purified by prep-HPLC (neutral condition). Compound **8** (0.035 g, 17.75 μmol , 82.4% yield) was obtained as a white solid.

General procedure for preparation of Compound 9

To a solution of compound **8** (0.035 g, 17.75 μmol , 1.0 eq), 1-hydroxypyrrolidine-2,5-dione (6.1 mg, 53.26 μmol , 3.0 eq) in DMA (3 mL) and DCM (1 mL), EDCI (10.2 mg, 53.26 μmol , 3.0 eq) was added. The mixture was stirred at 25 °C for 16 hr. LC-MS showed compound **8** was partially remained and one peak with desired MS was detected ($[\text{M}+2\text{H}^+]/2=1034.7$). DCM was then removed, following by mixture being purified by prep-HPLC (neutral condition). Compound **9** (0.03 g, 14.50 μmol , 81.7% yield) was obtained as a white solid.

General procedure for preparation of Compound 10

To a solution of BCY6014 (52.9 mg, 17.40 μmol , 1.59 μL , 1.2 eq) in DMF (2 mL), DIEA (5.6 mg, 43.51 μmol , 7.6 μL , 3.0 eq) and compound **9** (0.03 g, 14.50 μmol , 1.0 eq) were added. The mixture was stirred at 25 °C for 16 hr. LC-MS showed one main peak with desired MS ($[\text{M}+4\text{H}^+]/4=1249.2$, $[\text{M}+5\text{H}^+]/5=999.3$). The solvent was then removed and the resulting crude product **10** (0.06 g, crude) was used into the next step without further purification.

General procedure for preparation of BCY6107

To a solution of compound **10** (0.055 g, 11.01 μmol , 1.0 eq) in DCM (1 mL), 1 mL TFA was added. The mixture was stirred at 0 °C for 15 min. LC-MS showed compound **10** was consumed completely and one main peak with desired MS was detected ($[\text{M}+4\text{H}^+]/4=1221.0$, $[\text{M}+5\text{H}^+]/5=977.0$). The reaction mixture was concentrated under reduced pressure to remove solvent, resulting a residue which was then directly purified by

prep-HPLC (TFA condition). Compound **BCY6107** (20.4 mg, 4.03 μ mol, 36.6% yield, 96.36% purity) was obtained as a white solid.

BIOLOGICAL DATA

5 **Study 1: Fluorescence polarisation measurements**

(a) Direct Binding Assay

Peptides with a fluorescent tag (either fluorescein, SIGMA or Alexa Fluor488™, Fisher Scientific) were diluted to 2.5nM in PBS with 0.01% tween 20 or 50mM HEPES with 100mM NaCl and 0.01% tween pH 7.4 (both referred to as assay buffer). This was combined with a
10 titration of protein in the same assay buffer as the peptide to give 1nM peptide in a total volume of 25 μ L in a black walled and bottomed low bind low volume 384 well plates, typically 5 μ L assay buffer, 10 μ L protein (Table 1) then 10 μ L fluorescent peptide. One in two serial dilutions were used to give 12 different concentrations with top concentrations ranging from 500nM for known high affinity binders to 10 μ M for low affinity binders and selectivity assays.
15 Measurements were conducted on a BMG PHERAstar FS equipped with an "FP 485 520 520" optic module which excites at 485nm and detects parallel and perpendicular emission at 520nm. The PHERAstar FS was set at 25°C with 200 flashes per well and a positioning delay of 0.1 second, with each well measured at 5 to 10 minute intervals for 60 minutes. The gain used for analysis was determined for each tracer at the end of the 60 minutes where there
20 was no protein in the well. Data was analysed using Systat Sigmaplot version 12.0. mP values were fit to a user defined quadratic equation to generate a Kd value: $f = y_{min} + (y_{max} - y_{min}) / \text{Lig} * ((x + \text{Lig} + K_d) / 2 - \sqrt{(((x + \text{Lig} + K_d) / 2)^2 - (\text{Lig} * x))})$. "Lig" was a defined value of the concentration of tracer used.

25 (b) Competition Binding Assay

Peptides without a fluorescent tag were tested in competition with a peptide with a fluorescent tag and a known Kd (Table 2). Reference Compound A has the sequence FI-G-Sar₅-ACPWGPWCPVNRPGCA (FI-G-Sar₅-(SEQ ID NO: 94)). Reference Compound B has the sequence FI-G-Sar₅-ACPWGPFWCPVNRPGCA (FI-G-Sar₅-(SEQ ID NO: 95)). Reference
30 Compound C has the sequence FI-G-Sar₅-ADVTCWGPFWCPVNRPGCA (FI-G-Sar₅-(SEQ ID NO: 96)). Each of Reference Compounds A, B and C contain a TBMB molecular scaffold. Peptides were diluted to an appropriate concentration in assay buffer as described in the direct binding assay with a maximum of 5% DMSO, then serially diluted 1 in 2. Five μ L of diluted peptide was added to the plate followed by 10 μ L of human or mouse EphA2 (Table 1) at a
35 fixed concentration which was dependent on the fluorescent peptide used (Table 2), then 10 μ L fluorescent peptide added. Measurements were conducted as for the direct binding assay, however the gain was determined prior to the first measurement. Data analysis was in Systat

Sigmaplot version 12.0 where the mP values were fit to a user defined cubic equation to generate a K_i value:

$$f = y_{min} + (y_{max} - y_{min}) / \text{Lig} * ((\text{Lig} * ((2 * ((\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c)^2 - 3 * (\text{Kcomp} * (\text{Lig} - \text{Prot} * c) + \text{Klig} * (\text{Comp} - \text{Prot} * c) + \text{Klig} * \text{Kcomp}))^{0.5} * \text{COS}(\text{ARCCOS}((-2 * (\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c)^3 + 9 * (\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c) * (\text{Kcomp} * (\text{Lig} - \text{Prot} * c) + \text{Klig} * (\text{Comp} - \text{Prot} * c) + \text{Klig} * \text{Kcomp}) - 27 * (-1 * \text{Klig} * \text{Kcomp} * \text{Prot} * c)) / (2 * (((\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c)^2 - 3 * (\text{Kcomp} * (\text{Lig} - \text{Prot} * c) + \text{Klig} * (\text{Comp} - \text{Prot} * c) + \text{Klig} * \text{Kcomp}))^{0.5}))) / 3)) - (\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c))) / ((3 * \text{Klig}) + ((2 * ((\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c)^2 - 3 * (\text{Kcomp} * (\text{Lig} - \text{Prot} * c) + \text{Klig} * (\text{Comp} - \text{Prot} * c) + \text{Klig} * \text{Kcomp}))^{0.5} * \text{COS}(\text{ARCCOS}((-2 * (\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c)^3 + 9 * (\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c) * (\text{Kcomp} * (\text{Lig} - \text{Prot} * c) + \text{Klig} * (\text{Comp} - \text{Prot} * c) + \text{Klig} * \text{Kcomp}) - 27 * (-1 * \text{Klig} * \text{Kcomp} * \text{Prot} * c)) / (2 * (((\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c)^2 - 3 * (\text{Kcomp} * (\text{Lig} - \text{Prot} * c) + \text{Klig} * (\text{Comp} - \text{Prot} * c) + \text{Klig} * \text{Kcomp}))^{0.5}))) / 3)) - (\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c))))). \\ \text{"Lig", "Klig" and "Prot" were all defined values relating to: fluorescent peptide concentration, the } K_d \text{ of the fluorescent peptide and EphA2 concentration respectively.}$$

Table 1: Ephrin receptors and source

<i>Receptor (domain)</i>	<i>Species</i>	<i>Format/tag</i>	<i>Supplier</i>	<i>Catalogue number</i>
EphA1 (Ecto)	Human	Fc fusion	R&D systems	7146-A1
EphA2 (Ecto)	Human	C-terminal polyHis	R&D systems	3035-A2
EphA2 (Ecto)	Human	C-terminal polyHis	In-house	N/A
EphA2 (Ecto)	Mouse	Fc fusion	R&D Systems	639-A2
EphA2 (Ecto)	Mouse	C-terminal polyHis	Sino Biological	50586-M08H
EphA2 (ligand binding)	Rat	C-terminal polyHis	In-house	N/A
EphA2 (ligand binding)	Dog	C-terminal polyHis	In-house	N/A
EphA3 (Ecto)	Human	Fc fusion	R&D systems	6444-A3
EphA3 (Ecto)	Human	N-terminal polyHis	In-house	N/A

EphA3 (Ecto)	Rat	C-terminal polyHis	Sino Biological	80465-R08H
EphA4 (Ecto)	Human	Fc fusion	R&D systems	6827-A4
EphA4 (Ecto)	Human	C-terminal polyHis	Sino Biological	11314-H08H
EphA4 (Ecto)	Rat	C-terminal polyHis	Sino Biological	80123-R08H
EphA6 (Ecto)	Human	Fc fusion	R&D systems	5606-A6
EphA7 (Ecto)	Human	Fc fusion	R&D systems	6756-A7
EphB1 (Ecto)	Rat	Fc fusion	R&D systems	1596-B1
EphB4 (Ecto)	human	C-terminal polyHis	R&D systems	3038-B4

Table 2: Final concentrations of fluorescent peptide and EphA2 as used with Competition Binding Assays

<i>Fluorescent peptide</i>	<i>Concentration of fluorescent peptide (nM)</i>	<i>Concentration of Human EphA2 (nM)</i>	<i>Concentration of Mouse EphA2 (nM)</i>
Reference Compound A	10	75	
Reference Compound B	1	30	
Reference Compound C	0.8 (human) 1 (mouse)	2.4	50

- 5 Certain peptide ligands of the invention were tested in the above mentioned assays and the results are shown in Tables 3-7:

Table 3: Biological Assay Data for Peptide Ligands of the Invention (TATA peptides, Direct Binding Assay)

Bicycle Compound Number	Sequence	Human EphA2 (K_D, nM $\pm$ 95% CI)
1	ACMNDWWCAMGWKCA-Sar ₆ -K(FI) ((SEQ ID NO: 3)-Sar ₆ -K(FI))	304 $\pm$ 91.99
2	ACVPDRRCAYMNVCA-Sar ₆ -K(FI) ((SEQ ID NO: 4)-Sar ₆ -K(FI))	74.91 $\pm$ 6.6
3	ACVWDGRRCAYMNVCA-Sar ₆ -K(FI) ((SEQ ID NO: 5)-Sar ₆ -K(FI))	129.8 $\pm$ 80.75
4	ACWVDSRCAYMNVCA-Sar ₆ -K(FI) ((SEQ ID NO: 6)-Sar ₆ -K(FI))	124.6 $\pm$ 51.74
5	ACVPDSRCAYMNVCA-Sar ₆ -K(FI) ((SEQ ID NO: 7)-Sar ₆ -K(FI))	93.95 $\pm$ 23.62
6	ACYVGKECAIRNVCA-Sar ₆ -K(FI) ((SEQ ID NO: 8)-Sar ₆ -K(FI))	168.5 $\pm$ 20.58
7	ACYVGKECAYMNVCA-Sar ₆ -K(FI) ((SEQ ID NO: 9)-Sar ₆ -K(FI))	149.73 $\pm$ 39.2
8	FI-G-Sar ₅ -ACYVGKECAYMNVCA (FI-G-Sar ₅ -(SEQ ID NO: 9))	218.33 $\pm$ 10.51
9	FI-(β -Ala)-Sar ₁₀ -ARDCPLVNPLCLHPGWTC (FI-(β -Ala)-Sar ₁₀ -(SEQ ID NO: 10))	6.43 $\pm$ 1.15
10	FI-(β -Ala)-Sar ₁₀ -A(HArg)DCPLVNPLCLHPGWTC (FI-(β -Ala)-Sar ₁₀ -(SEQ ID NO: 11))	9.07 $\pm$ 2.49
11	Ac-CPLVNPLCLHPGWTC(LHG-Sar ₆ -(D-K[FII])) (Ac-(SEQ ID NO: 12)-Sar ₆ -(D-K[FII]))	3.08 $\pm$ 0.43
12	Ac-CPLVNPLCLHPGWTC(L(D-His)G-Sar ₆ -(D-K[FII])) (Ac-SEQ ID NO: 13)-Sar ₆ -(D-K[FII]))	10.56 $\pm$ 0.77
13	Ac-CPLVNPLCLHPGWSCRGQ-Sar ₆ -(D-K[FII])) (Ac-(SEQ ID NO: 14)-Sar ₆ -(D-K[FII]))	5.29 $\pm$ 0.79
14	Ac-CPLVNPLCLHPGWSC(HArg)GQ-Sar ₆ -(D-K[FII])) (Ac-(SEQ ID NO: 15)-Sar ₆ -(D-K[FII]))	9.96 $\pm$ 0.55

Table 4: Biological Assay Data for Peptide Ligands of the Invention (TATA peptides, Competition Binding Assay)

Bicycle Compound Number	Sequence	K _i , nM ± 95% CI				
		Human EphA2		Mouse EphA2		
		Fluorescent Peptide				
		Reference Compound C	Reference Compound B	Reference Compound A	Reference Compound C	
15	ACMNDWWCAGWKCA (SEQ ID NO: 3)	277.5 ± 38.22				
16	ACVPDRRCAYMNVCA (SEQ ID NO: 4)	69.97 ± 8.67				
17	(β-Ala)-Sar ₁₀ -ACVPDRRCAYMNVCA ((β-Ala)-Sar ₁₀ -(SEQ ID NO: 16))	85.05 ± 1.08				
18	DLRCGGDPRCAYMNVCA (SEQ ID NO: 17)	70.8 ± 2.35				
19	SRPCVIDSRCAYMNVCA (SEQ ID NO: 18)	94.75 ± 24.01				
20	ESRCSPDARCAVMNVCA (SEQ ID NO: 19)	57.05 ± 4.61				
21	HSGCRPDPRCAYMNVCA (SEQ ID NO: 20)	62.15 ± 4.61				
22	GSGCKPDSPRCAYMNVCA (SEQ ID NO: 21)	63.25 ± 13.82				
23	ETVCLPDSPRCAYMNVCA (SEQ ID NO: 22)	130 ± 15.68				
24	GQVCIVDARCAVMNVCA (SEQ ID NO: 23)	168.5 ± 16.66				

Bicycle Compound Number	Sequence	Ki, nM ± 95% CI				
		Human EphA2		Mouse EphA2		
		Fluorescent Peptide				
		Reference Compound C	Reference Compound B	Reference Compound A	Reference Compound C	
25	ACVPDRRCAFENVCDH (SEQ ID NO: 24)	97.3 ± 3.33				
26	ACVPDRRCAFNMVCEDR (SEQ ID NO: 25)	39.05 ± 10.29				
27	ACVPDRRCAFQDVCDE (SEQ ID NO: 26)	159 n=1				
28	ACVPDRRCAFRDVCLTG (SEQ ID NO: 27)	1700 n=1				
29	ACYVGKECAYMNVCA (SEQ ID NO: 9)	209.5 ± 110.74	106.65 ± 24.94	87.7 n=1		
30	ACQPSNHCAFMYNCA (SEQ ID NO: 28)	293 n=1	186.53 ± 86.86	137 n=1		
31	ACSPTPACAVQNLCA (SEQ ID NO: 29)	223 n=1	177 ± 60.76			
32	ACTSCWAYPDSFCA (SEQ ID NO: 30)	232 ± 52.19		151 n=1		
33	ACTKPTGFCAYPDTICA (SEQ ID NO: 31)	268.5 ± 16.66				
34	ACRGEWGYCAYPDTICA	347.5 ± 57.82				

Bicycle Compound Number	Sequence	K _i , nM ± 95% CI				
		Human EphA2		Mouse EphA2		
		Fluorescent Peptide				
		Reference Compound C	Reference Compound B	Reference Compound A	Reference Compound C	
	(SEQ ID NO: 32)					
35	ACRNWGMYCAYPDTICA (SEQ ID NO: 33)	282.5 ± 65.66				
36	ACPDWGGKYCAYPDTICA (SEQ ID NO: 34)	160 ± 1.96				
37	ACRVYGPYCAYPDTICA (SEQ ID NO: 35)	294.5 ± 20.58				
38	ACSSCWAYPDSVCA (SEQ ID NO: 36)	400.33 ± 205.19				
39	ACQSCWAYPDTYCA (SEQ ID NO: 37)	321.33 ± 119.53				
40	ACGFMGLEPCETFCA (SEQ ID NO: 38)	187.5 ± 20.58				
41	ACGFMGLVPCEVHCA (SEQ ID NO: 39)	155 ± 9.8				
42	ACGFMGLEPCCEMVCA (SEQ ID NO: 40)	320.5 ± 14.7				
43	ACGFMGLEPCVTYCA	233.5 ± 20.58				

Bicycle Compound Number	Sequence	K _i , nM ± 95% CI				
		Human EphA2		Mouse EphA2		
		Fluorescent Peptide				
		Reference Compound C	Reference Compound B	Reference Compound A	Reference Compound C	
	(SEQ ID NO: 41)					
44	ACGFMGLEPCELVCA (SEQ ID NO: 42)	126.8 ± 21.17				
45	ACGFMGLVPCNVFCA (SEQ ID NO: 43)	142 ± 41.16				
46	ACGFMGLEPCELFCA (SEQ ID NO: 44)	81.7 ± 7.06				
47	ACGFMGLEPCELCMPK (SEQ ID NO: 45)	185 ± 74.48				
48	ACGFMGLEPCELYCA (SEQ ID NO: 46)	127.5 ± 14.7				
49	ACGFMGLEPCELYCAHT (SEQ ID NO: 47)	144 ± 17.64				
50	ACGFMGLEPCEMYCA (SEQ ID NO: 48)	140 ± 45.08				
51	ACGFMGLVPCELYCADN (SEQ ID NO: 49)	84.4 ± 36.46				
52	ACPLVNPLCLTSGWKCA	115.33 ± 11.33				

Bicycle Compound Number	Sequence	K _i , nM ± 95% CI			
		Human EphA2		Mouse EphA2	
		Fluorescent Peptide			
		Reference Compound C	Reference Compound B	Reference Compound A	Reference Compound C
	(SEQ ID NO: 50)				
53	ACPMVNPLCLHPGWICA (SEQ ID NO: 51)	15.4 ± 3.17			
54	ACPLVNPLCLHPGWICA (SEQ ID NO: 52)	15.25 ± 2.84			
55	ACPLVNPLCLHPGWRCA (SEQ ID NO: 53)	20.55 ± 0.88			
56	ACPLVNPLCNLPGWTCA (SEQ ID NO: 54)	184 ± 115.64			
57	ACPLVNPLCLVPGWSCA (SEQ ID NO: 55)	35.4 ± 10			
58	ACPLVNPLCLLDGWTCA (SEQ ID NO: 56)	38.35 ± 5.39			
59	ACPLVNPLCLMPGWGCA (SEQ ID NO: 57)	114.5 ± 10.78			
60	ACPLVNPLCMIGNWTCA (SEQ ID NO: 58)	96.2 ± 0.59			
61	ACPLVNPLCLMTGWSCA	241.5 ± 44.1			

Bicycle Compound Number	Sequence	K _i , nM ± 95% CI				
		Human EphA2		Mouse EphA2		
		Fluorescent Peptide				
		Reference Compound C	Reference Compound B	Reference Compound A	Reference Compound C	
	(SEQ ID NO: 59)					
62	ACPLVNPLCMMGGWKCA (SEQ ID NO: 60)	67.1 ± 19.21				
63	ACPLVNPLCLYGSWKCA (SEQ ID NO: 61)	59.05 ± 28.32				
64	ACPLVNPLCLHPGWTCA (SEQ ID NO: 62)	30 n=1				
65	ARDCPLVNPLCLHPGWTCA (SEQ ID NO: 63)	6.05 ± 1.38			39.1 ± 0.39	
66 (BCY6099)	(β-Ala)-Sar ₁₀ -A(HArg)DC(HyP)LVNPLCLHP(D- Asp)W(HArg)C (SEQ ID NO: 2)	4.94 ± 1.41			57.6 ± 24.86	
67 (BCY6014)	(β-Ala)-Sar ₁₀ -A(HArg)DCPLVNPLCLHPGWTCA ((β-Ala)-Sar ₁₀ -(SEQ ID NO: 11)	8.51 ± 0.17			61.7 ± 15.48	
68	Ac-ARDCPLVNPLCLHPGWTCA-Sar ₆ -(D-K) (Ac-(SEQ ID NO: 63)-Sar ₆ -(D-K))	19.3 ± 4.92			166.5 ± 30.38	
69	Ac-A(HArg)DCPLVNPLCLHPGWTCA- Sar ₆ -(D-K) (Ac-(SEQ ID NO: 11)-A-Sar ₆ -(D-K))	17.5 ± 0.98			164.5 ± 2.94	
70	RPACPLVNPLCLHPGWTCA	10.06 ± 2.96				

Bicycle Compound Number	Sequence	K _i , nM ± 95% CI				
		Human EphA2		Mouse EphA2		
		Fluorescent Peptide				
		Reference Compound C	Reference Compound B	Reference Compound A	Reference Compound C	
	(SEQ ID NO: 64)					
71	RPPCPLVNPLCLHPGWTCA (SEQ ID NO: 65)	11.11 ± 2.25				
72	KHSCPLVNPLCLHPGWTCA (SEQ ID NO: 66)	11.92 ± 6.04				
73	ACPLVNPLCLHPGWTCLHG (SEQ ID NO: 67)	1.98 ± 0.49			7.27 ± 1.09	
74	Ac-CPLVNPLCLHPGWTCLHG (Ac-(SEQ ID NO: 12))	1.76 ± 0.54				
75	(β-Ala)-Sar ₁₀ -ACPLVNPLCLHPGWTCLHG ((β-Ala)-Sar ₁₀ -(SEQ ID NO: 67))	2.48 ± 0.27			18 ± 1.18	
76	(β-Ala)-Sar ₁₀ -ACPLVNPLCLHPGWTCL(D-His)G ((β-Ala)-Sar ₁₀ -(SEQ ID NO: 68))	10.01 ± 1.55			75.15 ± 14.41	
77 (BCY6019)	Ac-CPLVNPLCLHPGWTCLHG-Sar ₆ -(D-K) (Ac-(SEQ ID NO: 12)-Sar ₆ -(D-K))	5.41 ± 0.86			48.23 ± 15.72	
78	Ac-CPLVNPLCLHPGWTCL(D-His)G-Sar ₆ -(D-K) (Ac-(SEQ ID NO: 13)- Sar ₆ -(D-K))	15.6 ± 4.7			115.03 ± 41.16	
79	ACPLVNPLCLHPG(2NaI)TCLHG (SEQ ID NO: 69)	162 ± 17.64				

Bicycle Compound Number	Sequence	K _i , nM ± 95% CI				
		Human EphA2		Mouse EphA2		
		Fluorescent Peptide				
		Reference Compound C	Reference Compound B	Reference Compound A	Reference Compound C	
80	RHDCPLVNPLCLLPGWTCA (SEQ ID NO: 70)	7.11 ± 0.72				
81	TPRCPLVNPLCLMPGWTC (SEQ ID NO: 71)	9.8 ± 2.61				
82	ACPLVNPLCLAPGWTC (SEQ ID NO: 72)	46.2 n=1				
83	ACPLVNPLCLAPGWTCRS (SEQ ID NO: 73)	7.05 ± 1.11				
84	ACPLVNPLCLEPGWTCA (SEQ ID NO: 74)	53.9 n=1				
85	ACPLVNPLCLEPGWTCAR (SEQ ID NO: 75)	10.95 ± 1.6				
86	ACPLVNPLCLHPGWSCA (SEQ ID NO: 76)	56.15 ± 11.27				
87 (BCY6026)	ACPLVNPLCLHPGWSCRQ (SEQ ID NO: 77)	2.57 ± 0.63			18.6 ± 0.59	
88	Ac-CPLVNPLCLHPGWSCRQ (Ac-(SEQ ID NO: 14))	1.64 ± 0.75				

Bicycle Compound Number	Sequence	K _i , nM ± 95% CI				
		Human EphA2		Mouse EphA2		
		Fluorescent Peptide				
		Reference Compound C	Reference Compound B	Reference Compound A	Reference Compound C	
89	(β-Ala)-Sar ₁₀ -ACPLVNPLCLHPGWSCRGQ ((β-Ala)-Sar ₁₀ -(SEQ ID NO: 77))	2.86 ± 1.29			29.55 ± 4.61	
90	(β-Ala)-Sar ₁₀ -ACPLVNPLCLHPGWSC(HArg)GQ ((β-Ala)-Sar ₁₀ -(SEQ ID NO: 78))	5.41 ± 0.67			47.05 ± 11.47	
91 (BCY6042)	Ac-CPLVNPLCLHPGWSCRGQ-Sar ₆ -(D-K) (Ac-(SEQ ID NO: 14)-Sar ₆ -(D-K))	5.98 ± 1.42			49.87 ± 14.44	
92	Ac-CPLVNPLCLHPGWSC(HArg)GQ-Sar ₆ -(D-K) (Ac-(SEQ ID NO: 15)-Sar ₆ -(D-K))	10.56 ± 6.56			75.27 ± 21.72	
93	ACPLVNPLCLHPG(2NaI)SCRGQ (SEQ ID NO: 79)	228 ± 103.88				
94	ACPLVNPLCLTPGWTCTNT (SEQ ID NO: 80)	13.25 ± 4.05				
95	ACPMVNPLCLHPGWKCA (SEQ ID NO: 81)	11.91 ± 3.73				
96	ACPMVNPLCLTPGWICA (SEQ ID NO: 82)	16.07 ± 4.58				
97	ACPMVNPLCLHPGWTCA (SEQ ID NO: 83)	20 ± 1.02				

Table 5: Biological Assay Data for TATA Peptide Ligands of the Invention (Competition Binding Assay)

Bicycle Compound Number	Sequence	K _i , nM ± 95% CI	
		Human EphA2	Mouse EphA2
		Fluorescent peptide	
		Reference Compound C	Reference Compound C
98	(β-Ala)-Sar ₁₀ -H(D-Asp)VT-C(Aib)(1NaI)G(Aib)F(1NaI)CP(tBuGly)N(HArg)P(D-Asp)C ((β-Ala)-Sar ₁₀ -(SEQ ID NO: 84))	251.5 ± 73.5	

Table 6: Biological Assay Data for Peptide Ligands of the Invention (BDC competition binding data with TATA Scaffolds)

BDC Compound Number	Bicycle precursor	General Formula	K _i , nM	
			Human EphA2	Mouse EphA2
			Fluorescent Peptide	
			Reference Compound C	Reference Compound C
BCY6027	BCY6099	Formula (A)	10.23	
BCY6028	BCY6099	Formula (B)	13.04	
BCY6031	BCY6014	Formula (A)	12.62	34.70
BCY6032	BCY6014	Formula (B)	11.42	35.90

Table 7: Selectivity Data for Peptide Ligands of the Invention (Selectivity Direct Binding Assay)

Bicycle Compound Number	mouse EphA2	rat EphA2	dog EphA2	Human & mouse EphA3	rat EphA3	human EphA4	rat & mouse EphA4	rat EphB1	Human EphB4	Human EphA7	Human EphA6	Human EphA1	Human Factor Xlla	human Carbonic anhydrase 9	human CD38
2	516.5 ± 236.18	210 ± 1.96		>1000	>100	>1000	10890 n=1								>6000
7	216	252.5 ± 6.86													
9						>3000									
11						>3000									
12						>3000									
13						>3000									
14						>3000									

Study 2: Fluorescence polarisation measurements (Alternative Protocol)**(a) Competition binding**

Peptides without a fluorescent tag were tested in competition with a peptide with a fluorescent tag and a known K_d (Table 9). Five μL of increasing (2 fold) concentrations of test compound was added to the plate followed by 10 μL of EphA2 protein (Table 8) at a fixed concentration which was dependent on the fluorescent peptide used (Table 9), then 10 μL fluorescent peptide added. Buffer was assay buffer as above with DMSO <1%. Measurements were conducted on a BMG PHERAstar FS equipped with an "FP 485 520 520" optic module which excites at 485nm and detects parallel and perpendicular emission at 520nm. The PHERAstar FS was set at 25°C with 200 flashes per well and a positioning delay of 0.1 second, with each well measured at 5 to 10 minute intervals for 60 minutes. Alternatively, measurements were done on at similar time intervals on a Perkin Elmer Envision equipped with FITC FP Dual Mirror, FITC FP 480 excitation filter and FITC FP P-pol 535 and FITC FP S-pol emission filters with 30 flashes and a G-Factor of 1.2. Data analysis was in Systat Sigmaplot version 12.0 or 13.0 where the mP values at 60 minutes were fit to a user defined cubic equation to generate a K_i value:

$$f = y_{\min} + (y_{\max} - y_{\min}) / \text{Lig} * ((\text{Lig} * ((2 * ((\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c)^2 - 3 * (\text{Kcomp} * (\text{Lig} - \text{Prot} * c) + \text{Klig} * (\text{Comp} - \text{Prot} * c) + \text{Klig} * \text{Kcomp}))^{0.5} * \cos(\arccos((-2 * (\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c)^3 + 9 * (\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c) * (\text{Kcomp} * (\text{Lig} - \text{Prot} * c) + \text{Klig} * (\text{Comp} - \text{Prot} * c) + \text{Klig} * \text{Kcomp})) - 27 * (-1 * \text{Klig} * \text{Kcomp} * \text{Prot} * c)) / (2 * (((\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c)^2 - 3 * (\text{Kcomp} * (\text{Lig} - \text{Prot} * c) + \text{Klig} * (\text{Comp} - \text{Prot} * c) + \text{Klig} * \text{Kcomp}))^{0.5}))) / 3)) - (\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c))) / ((3 * \text{Klig}) + ((2 * ((\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c)^2 - 3 * (\text{Kcomp} * (\text{Lig} - \text{Prot} * c) + \text{Klig} * (\text{Comp} - \text{Prot} * c) + \text{Klig} * \text{Kcomp}))^{0.5} * \cos(\arccos((-2 * (\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c)^3 + 9 * (\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c) * (\text{Kcomp} * (\text{Lig} - \text{Prot} * c) + \text{Klig} * (\text{Comp} - \text{Prot} * c) + \text{Klig} * \text{Kcomp})) - 27 * (-1 * \text{Klig} * \text{Kcomp} * \text{Prot} * c)) / (2 * (((\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c)^2 - 3 * (\text{Kcomp} * (\text{Lig} - \text{Prot} * c) + \text{Klig} * (\text{Comp} - \text{Prot} * c) + \text{Klig} * \text{Kcomp}))^{0.5}))) / 3)) - (\text{Klig} + \text{Kcomp} + \text{Lig} + \text{Comp} - \text{Prot} * c))))).$$

"Lig", "Klig" and "Prot" were all defined values relating to: fluorescent peptide concentration, the K_d of the fluorescent peptide and EphA2 concentration respectively.

Table 8: Eph receptors and source

Receptor (domain)	Species	Format/tag	Supplier	Catalogue number
EphA2 (Ecto)	Human	C-terminal polyHis	R&D systems	3035-A2

EphA2 (Ecto)	Human	C-terminal polyHis	In-house	N/A
EphA2 (Ecto)	Mouse	C-terminal polyHis	Sino Biological	50586-M08H
EphA2 (ligand binding)	Rat	C-terminal polyHis	In-house	N/A

Table 9: Final concentrations of fluorescent peptide and EphA2 as used with competition binding assays

Fluorescent peptide	Concentration of fluorescent peptide (nM)	Concentration of human EphA2 (nM)	Concentration of mouse EphA2 (nM)	Concentration of rat EphA2 (nM)
Reference Compound C	0.8	2.4 or 25	50 or 15nM	25

- 5 Certain peptide ligands and bicycle drug conjugates of the invention were tested in the above mentioned competition binding assay and the results are shown in Tables 10 to 11:

Table 10: Competition Binding with Selected Bicyclic Peptides

<i>Bicycle No.</i>	<i>Human Ki (nM)</i>	<i>Mouse Ki (nM)</i>	<i>Rat Ki (nM)</i>
BCY6009 (Compound 108)	12.7	26.7	18.0
BCY6014 (Compound 67)	14.5	39.6	24.4
BCY6017 (Compound 109)	8.3		
BCY6018 (Compound 110)	13.1		
BCY6019 (Compound 77)	6.4	16.0	
BCY6026 (Compound 87)	4.4		
BCY6042 (Compound 91)	6.7		
BCY6059	43.2		

(Compound 106)			
BCY6099 (Compound 66)	2.7	4.5	1.9
BCY6101 (Compound 101)	9.7	6.9	
BCY6102 (Compound 102)	14.6	25.1	
BCY6103 (Compound 100)	14.8	20.8	
BCY6104 (Compound 99)	5.1	19.8	
BCY6137 (Compound 105)	2.2		
BCY6138 (Compound 104)	566.0		
BCY6139 (Compound 103)	5.7		
BCY6141 (Compound 112)	90.4		
BCY6152 (Compound 111)	23.3		
BCY6153 (Compound 113)	18.2		
BCY6160 (Compound 107)	14.0		
BCY6039	9.4		
BCY6105	8.86		
BCY6106	12.9		
BCY6175	1		
BCY6107	19.18		

The results from the competition binding assay in Table 10 show that Bicycle peptides targeting human EphA2 (BCY6014 and BCY6099) bind with high affinity to mouse and rat EphA2. Similarly, BCY6019 binds to both human and mouse EphA2. These results show that certain peptides of the invention can be used in *in vivo* mouse and rat efficacy and toxicology models.

Table 11: Competition Binding with Selected Bicycle Drug Conjugates (BDCs)

<i>Bicycle ID</i>	<i>Human Ki (nM)</i>	<i>Mouse Ki (nM)</i>	<i>Rat Ki (nM)</i>
BCY6061	12.0	32.3	14.2
BCY6174	1.7	3.9	3.0
BCY6029	2.3		
BCY6033	9.9	34.2	13.4
BCY6037	7.3		
BCY6049	8.8	28.1	
BCY6053	48.2	29.7	
BCY6122	13.7	10.4	
BCY6136	1.9	5.5	3.2
BCY6030	5.6		
BCY6034	5.9	35.9	
BCY6038	2.8		
BCY6050	168.1	62.2	
BCY6054	53.6	73.6	
BCY6027	10.2		
BCY6031	12.5	35.1	20.0
BCY6035	15.2		
BCY6047	53.2	34.2	
BCY6051	54.0	43.6	
BCY6134	7.4	12.6	
BCY6135	2.4	5.0	2.9
BCY6154	8.0		
BCY6155	12.5		
BCY6063	7.8	66.8	
BCY6028	13.0		
BCY6032	11.4	35.9	
BCY6036	18.6		
BCY6048	120.7	87.2	
BCY6052	30.5	27.1	
BCY6064	12.5	40.7	

BCY6162	44.9		
BCY6082	10.5	34.1	13.9
BCY6150	17.9		
BCY6151	9.0		
BCY6161	2.1		
BCY6173	1.7	4.3	2.5
BCY6077	6.5	25.3	
BCY6055	15.8		
BCY6062	12.9	20.3	

Table 11 shows that certain Bicycle Drug Conjugates of the invention exhibit excellent cross reactivity between human, mouse and rodent EphA2. Peptides of the invention can therefore be used in mouse and rat efficacy and toxicology *in vivo* models.

5

(b) SPR Measurements

Non-Fc fusion proteins were biotinylated with EZ-Link™ Sulfo-NHS-LC-Biotin for 1 hour in 4mM sodium acetate, 100mM NaCl, pH 5.4 with a 3x molar excess of biotin over protein. The degree of labelling was determined using a Fluorescence Biotin Quantification Kit (Thermo) after dialysis of the reaction mixture into PBS. For analysis of peptide binding, a Biacore T200 instrument was used utilising a XanTec CMD500D chip. Streptavidin was immobilized on the chip using standard amine-coupling chemistry at 25°C with HBS-N (10 mM HEPES, 0.15 M NaCl, pH 7.4) as the running buffer. Briefly, the carboxymethyl dextran surface was activated with a 7 min injection of a 1:1 ratio of 0.4 M 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (EDC)/0.1 M N-hydroxy succinimide (NHS) at a flow rate of 10 µl/min. For capture of streptavidin, the protein was diluted to 0.2 mg/ml in 10 mM sodium acetate (pH 4.5) and captured by injecting 120µl onto the activated chip surface. Residual activated groups were blocked with a 7 min injection of 1 M ethanolamine (pH 8.5):HBS-N (1:1). Buffer was changed to PBS/0.05% Tween 20 and biotinylated EphA2 was captured to a level of 500-1500 RU using a dilution of protein to 0.2µM in buffer. A dilution series of the peptides was prepared in this buffer with a final DMSO concentration of 0.5% with a top peptide concentration was 50 or 100nM and 6 further 2-fold dilutions. The SPR analysis was run at 25°C at a flow rate of 90µl/min with 60 seconds association and 900-1200 seconds dissociation. Data were corrected for DMSO excluded volume effects. All data were double-referenced for blank injections and reference surface using standard processing procedures and data processing and kinetic fitting were performed using Scrubber software, version 2.0c (BioLogic Software).

25

Data were fitted using simple 1:1 binding model allowing for mass transport effects where appropriate.

For binding of Bicycle Drug Conjugates a Biacore 3000 instrument was used. For biotinylated proteins immobilisation levels were 1500 RU and the top concentration was 100nM. Otherwise the method was the same as described above using either the CMD500D or a CM5 chip (GE Healthcare). For the Fc-tagged proteins, a CM5 chip was activated as described above and then goat anti-human IgG antibody (Thermo-Fisher H10500) was diluted to 20µg/ml in 10mM sodium acetate pH5.0 and captured to approximately 3000 RU. The surface was then blocked as described above. Subsequent capture of the Fc-tagged proteins was carried out to obtain approximately 200-400 RU of the target protein. The proteins used are described below. All proteins were reconstituted as per manufacturer's suggested buffers and concentrations and captured using 5-10µg/ml protein in PBS/0.05% Tween 20.

Table 12

Receptor	Species	Format/tag	Supplier	Catalogue number
EphA1	Human	Fc fusion	Sino Biologics	15789-H02H
EphA2	Human	0.95mol biotin/monomer	In house	N/A
EphA2	Mouse	Fc fusion	R&D Systems	639-A2
EphA2	Rat	1.4mol biotin/monomer	In house	N/A
EphA3	Human	Fc fusion	R&D Systems	6444-A3
EphA3	Mouse	Fc fusion	Sino Biologics	51122-M02H
EphA3	Rat	Fc fusion	Sino Biologics	80465-R02H
EphA4	Human	Fc fusion	Sino Biologics	11314-H03H
EphA4	Mouse	Fc fusion	Sino Biologics	50575-M02H

EphA4	Rat	Fc fusion	Sino Biologics	80123-R02H
EphA5	Human	3.1mol biotin/monomer	R&D Systems	3036-A5
EphA6	Human	Fc fusion	R&D Systems	5606-A6
EphA7	Human	Fc fusion	R&D Systems	6756-A7
EphB1	Rat	Fc fusion	R&D Systems	1596-B1
EphB4	Human	Fc fusion	Sino Biologics	10235-H02H

Certain peptide ligands and bicycle drug conjugates of the invention were tested in the above mentioned competition binding assay and the results are shown in Tables 13 to 15:

Table 13: SPR Binding Analysis with Selected Bicyclic Peptides and Bicycle Drug Conjugates of the Invention

Bicycle/BDC No.	Human				Mouse				Rat			
	K _D (nM)	K _{off} (s ⁻¹)	t _{1/2} (min)	K _{on} (M ⁻¹ s ⁻¹)	K _D (nM)	K _{off} (s ⁻¹)	t _{1/2} (min)	K _{on} (M ⁻¹ s ⁻¹)	K _D (nM)	K _{off} (s ⁻¹)	t _{1/2} (min)	K _{on} (M ⁻¹ s ⁻¹)
BCY6026	1.02	1.02E-03	11.3	9.92E+05								
BCY6031	1.99	4.95E-03	2.3	2.49E+06								
BCY6032	2.10	5.27E-03	2.2	2.52E+06								
BCY6033	3.41	3.43E-03	3.5	9.99E+05	21.8	6.37E-03	1.8	2.92E+05	166	4.42E-03	2.6	2.67E+04
BCY6034	1.64	3.65E-03	3.2	2.23E+06								
BCY6082	2.42	2.42E-03	4.8	9.87E+05	18.3	5.97E-03	1.9	3.27E+05	28.8	3.64E-03	3.2	1.26E+05
BCY6136	1.17	1.15E-03	10.0	9.86E+05	2.53	1.11E-03	10.4	4.37E+05	2.96	9.11E-04	12.6	3.07E+05
BCY6173	0.73	1.24E-03	9.3	1.69E+06	2.95	1.14E-03	10.1	3.86E+05	1.10	9.60E-04	12.0	8.81E+05

Table 13 details binding affinities and kinetic parameters (Koff and Kon) for binding of selected Bicycle Drug Conjugates to human EphA2 determined using the SPR assay.

Table 14: SPR Binding Analysis with Selected Bicycle Drug Conjugates of the Invention with Human Eph Homologs

<i>BDC No.</i>	<i>EphA1</i>	<i>EphA3</i>	<i>EphA4</i>	<i>EphA5</i>	<i>EphA6</i>	<i>EphA7</i>	<i>EphB4</i>
BCY6033	no binding @ 5µM	no binding @ 5µM	no binding @ 5µM	no binding @ 25µM	no binding @ 20µM	no binding @ 20µM	no binding @ 20µM
BCY6082	no binding @ 5µM	no binding @ 5µM	no binding @ 5µM	no binding @ 25µM	no binding @ 20µM	no binding @ 20µM	no binding @ 20µM
BCY6136	no binding @ 5µM	no binding @ 5µM	no binding @ 5µM	no binding @ 25µM	no binding @ 20µM	no binding @ 20µM	no binding @ 20µM
BCY6173	no binding @ 5µM	no binding @ 5µM	no binding @ 5µM	no binding @ 25µM	no binding @ 20µM	no binding @ 20µM	no binding @ 20µM

Table 14 illustrates binding results with four Bicycle Drug Conjugates (BCY6033, BCY6082, BCY6136 and BCY6173) in the SPR assay with closely related human Ephrin homologs. The results show that compounds of the invention exhibit no significant binding to closely related human homologs: EphA1, EphA3, EphA4, EphA5, EphA6, EphA7 and EphB4.

Table 15: SPR Binding Analysis with Selected Bicycle Drug Conjugates of the Invention with Mouse and Rat Eph Orthologs

BDC No.	Mouse EphA3	Mouse EphA4	Rat EphA3	Rat EphB1
BCY6033	no binding @ 20μM	no binding @ 20μM	no binding @ 20μM	no binding @ 20μM
BCY6082	no binding @ 20μM	no binding @ 20μM	no binding @ 20μM	no binding @ 20μM
BCY6136	no binding @ 20μM	no binding @ 20μM	no binding @ 20μM	no binding @ 20μM
BCY6173	no binding @ 20μM	no binding @ 20μM	no binding @ 20μM	no binding @ 20μM

The results in Table 15 show that certain Bicycle Drug Conjugates of the invention (BCY6033, BCY6082, BCY6136 and BCY6173) are also selective for mouse and rat EphA2 and exhibit no significant binding to closely related homologs: mouse EphA3 and EphA4; and rat EphA3 and EphB1.

Studies 3 and 7-23

In each of Studies 3 and 7-23, the following methodology was adopted for each study:

(a) Materials

(i) Animals and Housing Condition

Animals

Species: *Mus Musculus*

Strain: Balb/c nude or CB17-SCID

Age: 6-8 weeks

Body weight: 18-22 g

Number of animals: 9-90 mice

Animal supplier: Shanghai Lingchang Biotechnology Experimental Animal Co. Limited

Housing condition

The mice were kept in individual ventilation cages at constant temperature and humidity with 3-5 animals in each cage.

- Temperature: 20~26 °C.
- Humidity 40-70%.

Cages: Made of polycarbonate. The size is 300 mm x 180 mm x 150 mm. The bedding material is corn cob, which is changed twice per week.

Diet: Animals had free access to irradiation sterilized dry granule food during the entire study period.

Water: Animals had free access to sterile drinking water.

Cage identification: The identification labels for each cage contained the following information: number of animals, sex, strain, the date received, treatment, study number, group number and the starting date of the treatment.

- 5 Animal identification: Animals were marked by ear coding.

(ii) Test and Postitive Control Articles

Number	Physical Description	Molecular Weight	Purity	Storage Condition
BCY6031	Lyophilised powder	3878.92	97.99%	Stored at -80°C
BCY6033	Lyophilised powder	4260.01	99.12%	Stored at -80°C
BCY6082	Lyophilised powder	3911.04	96.8%	Stored at -80°C
BCY6135	Lyophilised powder	4021	95.14%	Stored at -80°C
BCY6136	Lyophilised powder	4402.23	97.5-98.6%	Stored at -80°C
BCY6173	Lyophilised powder	4101.15	95.80%	Stored at -80°C
BCY6174	Lyophilised powder	4537	99.50%	Stored at -80°C
BCY6175	Lyophilised powder	4492.29	96.20%	Stored at -80°C
BCY8245	Lyophilised powder	4173.85	99.30%	Stored at -80°C
BCY8781	Lyophilised powder	4173.83	99.00%	Stored at -80°C
ADC (MEDI-547) ¹	Solution (10.47 mg/ml concentration)	-	> 99.00%	Stored at -80°C

¹Full details of MEDI-547 (a fully human monoclonal antibody 1C1 (recognizing both human and murine EphA2) conjugated to MMAF via an mc linker) are described in Jackson *et al* (2008) Cancer Res 68, 9367–74.

10

(b) *Experimental Methods and Procedures*

(i) Observations

All the procedures related to animal handling, care and the treatment in the study were performed according to the guidelines approved by the Institutional Animal Care and Use Committee (IACUC) of WuXi AppTec, following the guidance of the Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC). At the time of routine monitoring, the animals were daily checked for any effects of tumor growth and treatments on normal behavior such as mobility, food and water consumption (by looking only), body weight gain/loss, eye/hair matting and any other abnormal effect as stated in the protocol. Death and

15

observed clinical signs were recorded on the basis of the numbers of animals within each subset.

(ii) Tumor Measurements and the Endpoints

The major endpoint was to see if the tumor growth could be delayed or mice could be cured.

- 5 Tumor volume was measured three times weekly in two dimensions using a caliper, and the volume was expressed in mm³ using the formula: $V = 0.5 a \times b^2$ where a and b are the long and short diameters of the tumor, respectively. The tumor size was then used for calculations of T/C value. The T/C value (in percent) is an indication of antitumor effectiveness; T and C are the mean volumes of the treated and control groups, respectively, on a given day.
- 10 TGI was calculated for each group using the formula: $TGI (\%) = [1 - (T_i - T_0) / (V_i - V_0)] \times 100$; T_i is the average tumor volume of a treatment group on a given day, T_0 is the average tumor volume of the treatment group on the day of treatment start, V_i is the average tumor volume of the vehicle control group on the same day with T_i , and V_0 is the average tumor volume of the vehicle group on the day of treatment start.

15 (iii) Sample Collection

At the end of study the tumors of all groups were collected for FFPE.

(iv) Statistical Analysis

Summary statistics, including mean and the standard error of the mean (SEM), are provided for the tumor volume of each group at each time point.

- 20 Statistical analysis of difference in tumor volume among the groups was conducted on the data obtained at the best therapeutic time point after the final dose.
- A one-way ANOVA was performed to compare tumor volume among groups, and when a significant F-statistics (a ratio of treatment variance to the error variance) was obtained, comparisons between groups were carried out with Games-Howell test. All data were analyzed
- 25 using GraphPad Prism 5.0. $P < 0.05$ was considered to be statistically significant.

Study 3: In vivo efficacy in the LU-01-0046 PDX model

- Cancer cell lines (CCL) are originally derived from patient tumors, but acquire the ability to proliferate within *in vitro* cell cultures. As a result of *in vitro* manipulation, CCL that have
- 30 been traditionally used in cancer research undergo genetic transformations that are not restored when cells are allowed to grow *in vivo*. Because of the cell culturing process cells that are better adapted to survive in culture are selected, tumor resident cells and proteins that interact with cancer cells are eliminated, and the culture becomes phenotypically homogeneous. Researchers are beginning to attribute the reason that only 5% of anti-cancer
 - 35 agents are approved by the Food and Drug Administration after pre-clinical testing to the lack of tumor heterogeneity and the absence of the human stromal microenvironment. Specifically, CCL-xenografts often are not predictive of the drug response in the primary

tumors because CCL do not follow pathways of drug resistance or the effects of the microenvironment on drug response found in human primary tumors. To overcome these problems, the inventors have used PDX models to improve the predictive power of pre-clinical models.

5

PDX are created when cancerous tissue from a patient's primary tumor is implanted directly into an immunodeficient mouse. PDX can maintain patient histology, including the presence of non-tumor cells (eg stromal cells) and thus better mimic the tumor microenvironment. In general PDX are therefore more reflective of the heterogeneity and histology of primary

10 tumors than CCL-xenografts.

BCY6031 was screened in a primary adenocarcinoma PDX xenograft (LU-01-0046) derived from a patient with non-small cell lung carcinomas (NSCLC). LU-01-0046 has been shown to express high levels of EphA2 using RNA sequencing. BCY6031 exhibited excellent efficacy in the LU-01-0046 model and is therefore a promising novel therapy for the treatment of non-small cell lung cancer.

15

(a) Treatment Arms

The experiment was designed to compare tumour growth in vehicle treated animals and animals treated with BCY6031 at 5 mg/kg qw for four weeks.

20 **Table 16**

<i>Gr</i>	<i>n</i>		<i>Treatment</i>	<i>Dose (mg/kg)</i>	<i>Dosing Volume (μl/g)</i>	<i>Dosing Route</i>	<i>Schedule</i>
1	6		Vehicle	-	10	i.v.	biw*1week
2	3		BCY6031	5	10	i.v.	qw*4weeks

Note: n: animal number; Dosing volume: adjust dosing volume based on body weight.

(b) Experimental Method

(i) PDX information

Table 17

<i>Model Name</i>	<i>Cancer Type</i>	<i>Tumor growth speed</i>	<i>Array</i>	<i>RSQ</i>	<i>EPH2 expression</i>
LU-01-0046	NSCLC	Tumor size can reach 1000 mm ³ in 40 days after tumor inoculation	6.790	31.312	High

25 **(ii) Tumor Inoculation**

Each mouse was inoculated subcutaneously in the right flank with an approximately 30 mm³ LU-01-0046 tumor fragment. Drug treatment was started when the average tumor volume

reached 943 mm³. The test article, route of administration, dosing frequency and the animal numbers in each group are described above.

(iii) Testing Article Formulation Preparation

Table 18

Test article	Dose(mg/kg)	Formulation
Vehicle	--	50 mM Acetate, 10% Sucrose pH5 (without DMSO)
BCY6031	5	Dissolve 4.59 mg BCY6031 into 4.498 ml formulation buffer to get the 1 mg/ml BCY6031 stock solution; Dilute 450ul 1 mg/ml BCY6031 with 450 µl formulation buffer.

5 (c) Results

(i) Mortality, Morbidity, and Body Weight Gain or Loss

Animal body weight was monitored regularly as an indirect measure of toxicity. Body weight change in female Balb/C nude mice bearing LU-01-0046 tumor dosed with BCY6031 is shown in Figure 1.

10 (ii) Tumor Growth Curve

The tumor growth curve is shown in Figure 2.

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for BCY6031 in the PDX model LU-01-0046 was calculated based on tumor volume measurements on day 7 after the start of treatment.

15 Table 19: Tumor growth inhibition analysis (T/C and TGI) on Day 7

Group	Treatment	Tumor Volume (mm³)^a	T/C^b (%)	TGI (%)	P value
1	Vehicle, biw	2191±473	--	--	--
2	BCY6031, 5 mpk, qw	463±158	21.1	138.6	<i>p</i> <0.05

a. Mean ± SEM.

b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

20 (e) Discussion

The study evaluated the therapeutic efficacy of BCY6031 in the LU-01-0046 PDX model. The measured body weights are shown in the Figure 1. Tumor volumes of the treatment group at various time points are shown in Table 19 and Figure 2.

The mean tumor size of vehicle treated mice reached 2191 mm³ on day 7. BCY6031 at 5 mg/kg produced potent antitumor activity with tumor measured as 463 mm³ (TGI=138.6%, $p<0.05$) by day 7. Furthermore, the BCY6031 treatment completely eradicated the tumors from day 32 and no tumour regrowth occurred following dosing suspension on day 28. BCY6031 gave rise to no significant body weight loss (Figure 1) and there were no adverse clinical observations on drug treated mice throughout the study.

Study 4: In vivo efficacy of BCY6136 in CDX xenograft models

The study evaluated the therapeutic efficacy of BCY6136 in three Cancer Cell Line Derived (CDX) models: the HT1080 fibrosarcoma line, the MDA-MB-231 triple negative breast cancer line and the NCI-H1975 non-small cell lung cancer (NSCLC) line.

(a) Experimental method

Balb/c mice were inoculated subcutaneously with tumour cells at the right flank and drug treatment started when the average the average tumour volume reached between 150 and 200 mm³. Tumour measurements and statistical analysis were performed as described above. Tumour bearing animals were treated once weekly with BCY6136 or vehicle.

(b) Discussion

Figures 4-6 show that BCY6136 is effective in breast, lung and fibrosarcoma xenograft models following once weekly dosing.

The HT1080 fibrosarcoma model:

In the HT1080 model complete regression of tumour growth was achieved by day 14 following once weekly dosing with BCY6136 on days 0 and 7 at 3 and 5 mg/kg (Figure 4). Once weekly dosing with BCY6136 at 2 mg/kg on days 0 and 7 gave rise to tumour stasis (partial regression) (Figure 4). BCY6136 treatment gave rise to no significant body weight loss (Figure 4 inset) and there were no adverse clinical observations on drug treated mice throughout the study.

The NCI-H1975 NSCLC model:

Complete regression of tumour growth in the NCI-H1975 model was observed by around day 28 following 2 and 3 mg/kg once weekly dosing with BCY6136 (Figure 5). Following dosing cessation on day 35 no tumour regrowth was observed in the 3 mg/kg treated animals from day 35 to day 72 when the 3 mg/kg arm measurements ended (Figure 5). Dosing with BCY6136 at 2 mg/kg gave rise to complete regression in this model from around day 28. Following dosing cessation on day 35 there was no tumour regrowth until around day 51 at

the 2 mg/kg dose. At this dose level moderate tumour re-growth was observed from around day 51 until study termination on day 77. 1 mg/kg treatment with BCY6136 gave rise to tumour stasis (partial regression) (Figure 5). BCY6136 treatment gave rise to no significant body weight loss (Figure 5 inset) and there were no adverse clinical observations on drug treated mice throughout the study.

The MDA-MB-231 breast model:

Tumour stasis (partial regression) was observed in the MDA-MB231 model following once weekly dosing at 2 and 3 mg/kg from days 0 to day 45 (Figure 6). Some body weight loss (attributed to tumour burden) was observed in the 2 mg/kg treated animals (Figure 6 inset).

These results demonstrate that BCY6136 gives rise to profound tumour growth inhibition in mice implanted with fibrosarcoma, breast and lung CDX xenografts following once daily dosing.

Study 5: Safety studies in the rat

Six (6) female rats were randomly assigned to 3 groups of 2 rats/group to determine the toxicity of BCY6136, following administered by IV bolus injection at 5, 7.5 and 10 mg/kg on days 1 and 8. The study was terminated on day 15.

No significant effects on coagulation parameters (Prothrombin time (sec), Activated partial thromboplastin time (sec) or Fibrogen levels (g/L) were observed on days 2, 12 and 15 (data not shown). No in-life bleeding events were reported and no evidence of internal bleeding was detected following pathology examination.

Study 6: Safety studies in the cynomologous monkeys

Twenty eight day toxicology studies with BCY6136 we conducted in cynomologous monkeys. BCY6136 was dosed at 1.0 and 2.0 mg/kg on days 1, 8, 15 and 22. Animals were euthanised and necropsied on day 29 (7 days after the final dose).

No significant effects on coagulation parameters relative to baseline were observed on days 18, 22 and 25 (data not shown) and day 29 (Table 20). No in-life bleeding events were reported and no evidence of internal bleeding was detected following pathology examination.

Table 20: Day 29 coagulation parameters following 1.0 and 2.0 mg/kg BCY6136 dosing to cynomolgus monkeys

	1.0 mg/kg x 4		2.0 mg/kg x 4	
	Baseline	Day 29	Baseline	Day 29
PT(s)	13.4	11.7	9.4	9.7
PT(s)	11	9.2	11.2	11.0
APTT(s)	18.9	19.4	19.4	20.9
APTT(s)	16.1	15.7	18.7	18.2
FIB(g/L)	2.08	2.42	1.86	6.1
FIB(g/L)	2.28	2.35	1.82	3.1

5 **Study 7: In vivo efficacy study of BCY6033 and BCY6136 and ADC in treatment of PC-3 xenograft in Balb/c nude mice**

(a) **Study Objective**

The objective of the research is to evaluate the *in vivo* anti-tumor efficacy of BCY6033 and BCY6136 in treatment of PC-3 xenograft.

10 (b) **Experimental Design**

Group	Treatment	n	Dose (mg/kg)	Dosing Volume (μl/g)	Dosing Route	Schedule
1	Vehicle	3	-	10	<i>iv</i>	qw
2	BCY6136	3	1	10	<i>iv</i>	qw
3	BCY6136	3	2	10	<i>iv</i>	qw
4	BCY6136	3	3	10	<i>iv</i>	qw
5	ADC	3	3	10	<i>iv</i>	qw
6	BCY6033	3	3	10	<i>iv</i>	qw

(c) **Experimental Methods and Procedures**

(i) **Cell Culture**

15 The PC-3 tumor cells will be maintained in F12K medium supplemented with 10% heat inactivated fetal bovine serum at 37°C in an atmosphere of 5% CO₂ in air. The tumor cells will be routinely subcultured twice weekly. The cells growing in an exponential growth phase will be harvested and counted for tumor inoculation.

(ii) **Tumor Inoculation**

Each mouse will be inoculated subcutaneously at the right flank with PC-3 (10×10^6) tumor cells for tumor development. The animals will be randomized and treatment will be started when the average tumor volume reaches approximately 150 mm^3 . The test article administration and the animal numbers in each group are shown in the following experimental design table.

5 (iii) Testing Article Formulation Preparation

Test article	Con. (mg/ml)	Formulation
Vehicle	-	50 mM Acetate/acetic acid pH 5 10% sucrose
BCY6136	0.1	Dilute 90 μl 1 mg/ml BCY6136 stock with 810 μl vehicle buffer
	0.2	Dilute 180 μl 1 mg/ml BCY6136 stock with 720 μl vehicle buffer
	0.3	Dilute 270 μl 1 mg/ml BCY6136 stock with 630 μl vehicle buffer
ADC	0.3	Dilute 26 μl 10.47 mg/ml ADC stock with 874 μl ADC buffer
BCY6033	0.3	Dilute 270 μl 1 mg/ml BCY6033 stock with 630 μl vehicle buffer

(d) Results

(i) Body Weight change and Tumor Growth Curve

Body weight and tumor growth curve are shown in Figures 7 to 9.

(ii) Tumor Volume Trace

- 10 Mean tumor volume over time in female Balb/c nude mice bearing PC-3 xenograft is shown in Table 21.

Table 21: Tumor volume trace over time

Gr	Treatment	0	2	4	7	9	11	14	16	18	21
1	Vehicle, qw	149±9	235±9	377±9	718±30	1126±41	1431±79	1792±69	2070±152		
2	BCY6136, 1mpk, qw	150±11	185±25	228±31	201±17	183±23	153±38	137±33	107±32	64±28	45±23
3	BCY6136, 2 mpk, qw	149±18	179±28	158±22	137±16	122±15	114±20	101±16	79±20	57±19	42±17
4	BCY6136 3 mpk, qw	149±2	155±8	144±16	132±20	107±28	94±23	83±22	70±27	38±16	35±17
5	ADC 3 mpk, qw	151±27	203±10	210±12	189±11	185±16	190±37	158±36	124±35	103±27	74±14
6	BCY6033, 3mpk, qw	151±33	214±53	204±51	192±53	163±43	151±40	141±39	116±36	83±28	63±32

Gr	Treatment	23	25	28	30	32	35	37	39	42
1	Vehicle, qw									
2	BCY6136, 1mpk, qw	35±18	28±14	37±19	34±17	42±21	42±23	43±21	28±14	18±9
3	BCY6136, 2 mpk, qw	21±11	22±12	22±12	24±12	33±16	22±11	26±14	22±12	16±9

4	BCY6136 3 mpk, qw	21±10	23±12	27±14	22±11	24±12	20±11	27±14	12±6	12±6
5	ADC 3 mpk, qw	53±16	50±22	46±23	70±35	78±39	53±27	60±30	53±27	40±22
6	BCY6033, 3mpk, qw	59±31	44±27	39±24	40±29	47±32	41±27	41±30	34±24	33±27

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for test articles in the PC-3 xenograft model was calculated based on tumor volume measurements at day 16 after the start of treatment.

5 Table 22: Tumor growth inhibition analysis

Gr	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	P value compare with vehicle
1	Vehicle, qw	2070±152	--	--	--
2	BCY6136, 1mpk, qw	107±32	5.2	102.2	<i>p</i> <0.001
3	BCY6136, 2 mpk, qw	79±20	3.8	103.6	<i>p</i> <0.001
4	BCY6136, 3 mpk, qw	70±27	3.4	104.1	<i>p</i> <0.001
5	ADC, 3 mpk, qw	124±35	6.0	101.4	<i>p</i> <0.001

6	BCY6033, 3mpk, qw	116±36	5.6	101.8	p<0.001
---	----------------------	--------	-----	-------	---------

a. Mean ± SEM.

b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

5 **(e) Results Summary and Discussion**

In this study, the therapeutic efficacy of test articles in the PC-3 xenograft model was evaluated. The measured body weights and tumor volumes of all treatment groups at various time points are shown in the Figures 7 to 9 and Tables 21 and 22.

The mean tumor size of vehicle treated mice reached 2070 mm³ on day 16. BCY6136 at 1 mg/kg, qw (TV=107 mm³, TGI=102.2%, p<0.001), BCY6136 at 2 mg/kg, qw (TV=79 mm³, TGI=103.6%, p<0.001) and BCY6136 at 3 mg/kg, qw (TV=70 mm³, TGI=104.1%, p<0.001) showed potent anti-tumor effect.

BCY6033 at 3 mg/kg, qw (TV=116 mm³, TGI=101.8%, p<0.001) and ADC at 3 mg/kg, qw (TV=124 mm³, TGI=101.4%, p<0.001) showed comparable anti-tumor effect.

In this study, animal body weight was monitored regularly. All mice maintained their body weight well.

Study 8. In vivo efficacy study of BCY6136 in treatment of PC-3 xenograft in Balb/c nude mice

(a) Study Objective

The objective of the research is to evaluate the *in vivo* anti-tumor efficacy of BCY6136 in treatment of PC-3 xenograft in Balb/c nude mice.

(b) Experimental Design

Group	Treatment	Dose (mg/kg)	N ^a	Dosing Route	Schedule
1	Vehicle	--	4	i.v.	qw x4 weeks
2	BCY6136	0.167	4	i.v.	qw x4 weeks
3 ^b	BCY6136	0.5	4	i.v.	qw x4 weeks
4	BCY6136	1.5	4	i.v.	qw x4 weeks
5 ^b	BCY6136	0.5	4	i.v.	q2w x2 weeks
6 ^b	BCY6136	1.5	4	i.v.	q2w x2 weeks
7	EphA2-ADC	0.33	4	i.v.	qw x4 weeks
8	EphA2-ADC	1	4	i.v.	qw x4 weeks
9	EphA2-ADC	3	4	i.v.	qw x4 weeks
10 ^c	Docetaxel	15	4	i.v.	qw x4 weeks

a. N, the number of animals in each group.

b. After 4 weeks' treatment demonstrated in the experimental design table, the mice of group 3, 5 and 6 were treated with BCY6136 1.5 mg/kg qw from day 52 during the monitoring schedule.

c. Due to the severe body weight loss of the Docetaxel treated mice after the first dosing, the treatment was suspended for 2 weeks, then a lower dosage (Docetaxel, 10 mg/kg) was performed on day 28. After that, the mice were treated with BCY6136 1.5 mg/kg qw from day 42 to day 70.

(c) Experimental Methods and Procedures

(i) Cell Culture

The tumor cells were maintained in F-12K medium supplemented with 10% heat inactivated fetal bovine serum at 37°C in an atmosphere of 5% CO₂ in air. The tumor cells were routinely subcultured twice weekly. The cells growing in an exponential growth phase were harvested and counted for tumor inoculation.

(ii) Tumor Inoculation

Each mouse was inoculated subcutaneously at the right flank with PC-3 tumor cells (10 × 10⁶) in 0.2 ml of PBS for tumor development. 52 animals were randomized when the average tumor volume reached 454 mm³. The test article administration and the animal numbers in each group were shown in the experimental design table.

(iii) Testing Article Formulation Preparation

Test article	Purity	Conc. (mg/ml)	Formulation
Vehicle	-	-	25 mM Histidine pH 7 10% sucrose
BCY6136	98.6%	-	50 mM Acetate 10% sucrose pH 5
		1	Dissolve 2.70 mg BCY6136 in 2.662 ml Acetate buffer
		0.3	Dilute 300µl 1 mg/ml BCY6136 stock with 700 µl Acetate buffer ¹
		0.15	Dilute 600 µl 0.3 mg/ml BCY6136 stock with 600 µl Acetate buffer
		0.05	Dilute 200 µl 0.3 mg/ml BCY6136 stock with 1000 µl Acetate buffer
		0.0167	Dilute 66.7 µl 0.3 mg/ml BCY6136 stock with 1133.3 µl Acetate buffer
EphA2-ADC	-	-	25 mM Histidine pH 5.5
		0.033	Dilute 9.3 µl 4.24 mg/ml EphA2-ADC stock with 1191 µl His buffer
		0.1	Dilute 28 µl 4.24 mg/ml EphA2-ADC stock with 1172 µl His buffer
		0.3	Dilute 84.9 µl 4.24 mg/ml EphA2-ADC stock with 1115 µl His buffer
Docetaxel	-	10	Mix 0.5 ml 20mg Docetaxel with 1.5 ml buffer
		1.5	Dilute 180 µl 10 mg/ml Docetaxel stock with 1020 µl saline buffer
1. 50 mM Acetate 10% sucrose pH 5 3. 25 mM Histidine pH 5.5			

(c) Results(i) Body Weight change and Tumor Growth Curve

Body weight and tumor growth curve is shown in Figure 10.

5 (ii) Tumor Volume Trace

Mean tumor volume over time in male Balb/c nude mice bearing PC-3 xenograft is shown in Table 23.

Table 23: Tumor volume trace over time (Day 0 to day 20)

Gr.	Treatment	Days after the start of treatment									
		0	2	4	6	8	10	13	15	17	20
1	Vehicle, qw	456± 25	648± 50	880 ±23	1022±2 9	1178±1 18	1327±1 33	1631±9 3	1868±9 0	2052±1 39	2364±1 02
2	BCY6136 0.167 mpk, qw	450± 33	631± 55	695 ±78	739 ±39	850 ±68	904 ±73	975 ±47	1089±7 4	1124±9 2	1188±1 11
3	BCY6136 0.5 mpk, qw	451± 47	622± 96	519 ±70	460 ±55	398 ±50	329 ±38	260 ±33	249 ±33	231 ±38	234 ±42
4	BCY6136 1.5 mpk, qw	458± 49	587± 63	494 ±54	363 ±32	283 ±32	237 ±24	192 ±13	164 ±16	155 ±20	131 ±19
5	BCY6136 0.5 mpk, q2w	454± 37	643± 25	531 ±37	458 ±33	411 ±32	382 ±49	430 ±88	522 ±124	560 ±129	530 ±147
6	BCY6136 1.5 mpk, q2w	452± 42	590± 75	457 ±49	375 ±44	328 ±47	242 ±63	206 ±61	197 ±62	182 ±55	128 ±36
	1.5 mpk, qw										
7	EphA2-ADC 0.33 mpk, qw	457± 43	636± 57	712 ±70	792 ±78	870 ±87	900 ±58	1049±6 6	1242±1 23	1443±1 29	1637±1 81
8	EphA2-ADC 1 mpk, qw	450± 49	617± 48	673 ±50	721 ±61	782 ±78	755 ±67	840 ±93	913 ±91	978 ±100	981 ±100
9	EphA2-ADC			643	593	433	290	268	232	225	184

10	3 mpk, qw	452± 60	593± 98	±141	±106	±103	±81	±64	±60	±66	±62
	Docetaxel	453±	584±	632	636	568	408	374	388	361	419
	15 mpk, qw	62	72	±56	±48	±50	±31	±26	±36	±25	±31

(iii) Tumor Growth Inhibition Analysis
Tumor growth inhibition rate for test articles in the PC-3 xenograft model was calculated based on tumor volume measurements at day 20 after the start of the treatment.

Table 24: Tumor growth inhibition analysis

Gr	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	P value compared with vehicle
1	Vehicle, qw	2364±102	--	--	--
2	BCY6136, 0.167 mpk, qw	1188±111	50.2	61.4	<i>p</i> <0.001
3	BCY6136, 0.5 mpk, qw	234±42	9.9	111.4	<i>p</i> <0.001
4	BCY6136, 1.5 mpk, qw	131±19	5.5	117.2	<i>p</i> <0.001
5	BCY6136, 0.5 mpk, q2w	530±147	22.4	96.0	<i>p</i> <0.001
6	BCY6136, 1.5 mpk, q2w	128±36	5.4	117.0	<i>p</i> <0.001
7	EphA2-ADC, 0.33 mpk, qw	1637±181	69.2	38.1	<i>p</i> <0.001
8	EphA2-ADC, 1 mpk, qw	981±100	41.5	72.2	<i>p</i> <0.001
9	EphA2-ADC, 3 mpk, qw	184±62	7.8	114.0	<i>p</i> <0.001
10	Docetaxel, 15 mpk, qw	419±31	17.7	101.8	<i>p</i> <0.001

a. Mean ± SEM.

b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

5 (d) Results Summary and Discussion

In this study, the therapeutic efficacy of test articles in the PC-3 xenograft model was evaluated. The measured body weights and tumor volumes of all treatment groups at various time points are shown in the Figure 10 and Tables 23 and 24.

- 10 The mean tumor size of vehicle treated mice reached 2364 mm³ on day 20. BCY6136 at 0.167 mg/kg, qw (TV=1188 mm³, TGI=61.4%, *p*<0.001), 0.5 mg/kg, q2w (TV=530 mm³, TGI=96.0%, *p*<0.001), 0.5 mg/kg, qw (TV=234 mm³, TGI=111.4%, *p*<0.001) and 1.5 mg/kg, qw (TV=131 mm³, TGI=117.2%, *p*<0.001) produced significant anti-tumor activity in dose or dose-frequency dependent manner on day 20. BCY6136 at 1.5 mg/kg, q2w (TV=128 mm³, TGI=117.0%, *p*<0.001) produced comparable anti-tumor activity with BCY6136 1.5 mg/kg qw.
- 15 Among them, the mice treated with BCY6136, 0.5 mg/kg qw or BCY6136, 0.5 mg/kg q2w showed obvious tumor relapse after ceasing the treatment, further treatment with BCY6136,

1.5 mg/kg qw from day 52 worked well on the tumor regression. The mice treated with BCY6136, 1.5 mg/kg q2w also showed tumor relapse after ceasing the treatment, but further dosing didn't work on complete tumor regression. The mice treated with BCY6136, 1.5 mpk qw didn't show any tumor relapse until day 48.

- 5 EphA2-ADC at 0.33 mg/kg, qw (TV=1637 mm³, TGI=38.1%, $p<0.001$), 1 mg/kg, qw (TV=981 mm³, TGI=72.2%, $p<0.001$) and 3 mg/kg, qw (TV=184 mm³, TGI=114.0%, $p<0.001$) produced significant anti-tumor activity in dose dependent manner on day 20. The mice treated with EphA2-ADC, 3 mg/kg qw didn't show any tumor relapse until day 59.

- Docetaxel at 15 mg/kg, qw (TV=419 mm³, TGI=101.8%, $p<0.001$) produced significant anti-tumor activity but caused severe animal body weight loss. After ceasing the treatment, the mice showed obvious tumor relapse. The treatment with BCY6136, 1.5 mg/kg qw from day 42 worked well on tumor regression of these mice.

Study 9. In vivo efficacy test of BCY6033, BCY6136 and BCY6082 in treatment of

15 **NCI-H1975 xenograft in Balb/c nude mice**

(a) Study Objective

The objective of the research was to evaluate the *in vivo* anti-tumor efficacy of BCY6033, BCY6136 and BCY6082 in treatment of NCI-H1975 xenograft model in Balb/c nude mice.

(b) Experimental Design

Group	Treatment	n	Dose (mg/kg)	Dosing Volume (μl/g)	Dosing Route	Schedule
1	Vehicle	3	--	10	<i>iv</i>	qw
2	BCY6033	3	1	10	<i>iv</i>	qw
3	BCY6033	3	2	10	<i>iv</i>	qw
4	BCY6033	3	3	10	<i>iv</i>	qw
5	BCY6136	3	1	10	<i>iv</i>	qw
6	BCY6136	3	2	10	<i>iv</i>	qw
7	BCY6136	3	3	10	<i>iv</i>	qw
8	BCY6082	3	2	10	<i>iv</i>	qw
9	BCY6082	3	5	10	<i>iv</i>	qw

20

(c) Experimental Methods and Procedures

(i) Cell Culture

The cells growing in an exponential growth phase were harvested and counted for tumor inoculation.

25 **(ii) Tumor Inoculation**

Each mouse was inoculated subcutaneously at the right flank with NCI-H1975 tumor cells (10×10^6) in 0.2ml of PBS for tumor development. 36 animals were randomized when the average tumor volume reached 149 mm³. The test article administration and the animal numbers in each group were shown in the experimental design table.

5 (iii) Testing Article Formulation Preparation

Treatment	Dose (mg/ml)	Formulation
Vehicle		50 mM Acetate, 10% sucrose pH=5
BCY6033	1	Dissolve 6.71 mg BCY6033 in 6.710 ml formulation buffer
	0.3	Dilute 270 µl 1 mg/ml BCY6033 with 630 µl formulation buffer
	0.2	Dilute 180 µl 1 mg/ml BCY6033 with 720 µl formulation buffer
	0.1	Dilute 90 µl 1 mg/ml BCY6033 with 810 µl formulation buffer
BCY6136	1	Dissolve 3.79 mg BCY6136 in 3.695ml formulation buffer
	0.3	Dilute 270 µl 1 mg/ml BCY6136 with 630 µl formulation buffer
	0.2	Dilute 180 µl 1 mg/ml BCY6136 with 720 µl formulation buffer
	0.1	Dilute 90 µl 1 mg/ml BCY6136 with 810 µl formulation buffer
BCY6082	1	Weigh and dissolve 4.30 mg BCY6082 in 4.162 ml formulation buffer
	0.5	Dilute 450 µl 1 mg/ml BCY6082 with 450 µl formulation buffer
	0.2	Dilute 180 µl 1 mg/ml BCY6082 with 720 µl formulation buffer

(iv) Sample Collection

On PG-D23, we fixed the tumors of Group 1 for FFPE.

On PG-D44, we fixed the tumors of Group 2 and 5 for FFPE.

10 At the end of study, we the tumors of Group 6 for FFPE.

(d) Results

(i) Body Weight change and Tumor Growth Curve

Body weight and tumor growth are shown in Figures 11 to 13.

(ii) Tumor Volume Trace

15 Mean tumor volume over time in female Balb/c nude mice bearing NCI-H1975 xenograft is shown in Table 25 to 29.

Table 25: Tumor volume trace (PG-D0~PG-D17)

Gr	Treatment	Days after the start of treatment							
		0	2	4	7	9	11	14	17
.									

1	Vehicle, qw	148 ±4	195±1 1	297±3 3	466±6 4	732±1 07	1028±1 92	1278±2 52	1543±2 98
2	BCY6033, 1 mpk, qw	149 ±10	160±4	207±1 3	259±4 9	330±6 9	365±83	341±59	336±54
3	BCY6033, 2 mpk, qw	149 ±10	183±1 1	276±2 4	365±4 2	405±2 0	364±19	319±32	304±33
4	BCY6033, 3 mpk, qw	149 ±6	161±4	207±2 6	260±2 1	270±4 2	243±52	187±53	131±43
5	BCY6136, 1 mpk, qw	150 ±6	178±2 0	232±4 9	336±4 3	400±2 4	407±42	299±11 3	261±12 7
6	BCY6136, 2 mpk, qw	150 ±14	181±2 6	237±2 7	277±3 6	297±3 7	306±55	256±53	218±49
7	BCY6136, 3 mpk, qw	148 ±9	168±1 0	231±6	365±1 6	390±1 3	423±42	319±26	228±16
8	BCY6082, 2 mpk, qw	148 ±5	157±4	223±1 9	370±8 4	447±1 02	658±18 8	906±33 2	1123±4 10
9	BCY6082, 5 mpk, qw	148 ±6	176±1 2	235±1 9	378±5 9	436±6 8	510±82	484±78	491±10 3

Table 26: Tumor volume trace (PG-D18~PG-D35)

Gr.	Treatment	Days after the start of treatment							
		18	21	23	25	28	30	33	35
1	Vehicle, qw	186 4±3 95	2371± 470	--	--	--	--	--	--
2	BCY6033, 1 mpk, qw	278 ±71	306±8 1	343±8 6	366±8 9	466±1 15	481±1 12	619±1 70	780±2 36
3	BCY6033, 2 mpk, qw	172 ±25	95±12	61±6	39±4	13±1	12±1	6±3	6±3
4	BCY6033, 3 mpk, qw	75± 15	29±4	20±6	13±2	6±0	4±0	1±0	2±1
5	BCY6136, 1 mpk, qw	215 ±11 3	205±1 17	197±1 13	200±1 05	202±1 12	202±1 17	230±1 42	241±1 27

6	BCY6136, 2 mpk, qw	149 ±31	99±30	69±22	42±13	30±10	16±8	20±9	4±2
7	BCY6136, 3 mpk, qw	149 ±17	94±30	50±15	41±21	21±8	6±6	10±6	3±1
8	BCY6082, 2 mpk, qw	119 9±4 08	1528± 604	1978± 792	2499± 931	--	--	--	--
9	BCY6082, 5 mpk, qw	471 ±14 3	390±1 33	368±1 22	295±1 02	227±8 6	--	--	--

Table 27: Tumor volume trace (PG-D37~PG-D53)

Gr.	Treatment	Days after the start of treatment							
		37	39	42	44	46	49	51	53
2	BCY6033, 1 mpk, qw	877±188	945±145	1258±173	--	--	--	--	--
3	BCY6033, 2 mpk, qw	3±1	1±0	1±0	1±0	1±0	1±0	1±0	1±0
4	BCY6033, 3 mpk, qw	0±0	0±0	0±0	0±0	1±0	0±0	1±0	1±0
5	BCY6136, 1 mpk, qw	277±149	294±159	351±188	--	--	--	--	--
6	BCY6136, 2 mpk, qw	7±4	2±1	1±0	3±1	2±1	3±2	6±3	14±10
7	BCY6136, 3 mpk, qw	3±3	2±1	1±0	0±0	0±0	0±0	1±0	1±0

Table 28: Tumor volume trace (PG-D56~PG-D74)

Gr.	Treatment	Days after the start of treatment								
		56	58	60	63	65	67	70	72	74
3	BCY6033, 2 mpk, qw	1±0	1±0	1±0	1±0	1±0	2±1	4±3	7±6	--
4	BCY6033,	1±0	1±0	0±0	0±0	0±0	0±0	0±0	0±0	--

	3 mpk, qw									
6	BCY6136, 2 mpk, qw	16± 11	27±18	34±23	45±31	63±40	71±47	95±70	111±7 3	122±7 5
7	BCY6136, 3 mpk, qw	1±0	1±0	1±0	0±0	0±0	0±0	0±0	0±0	--

Table 29: Tumor volume trace (PG-D77~PG-D98)

Gr.	Treatment	Days after the start of treatment						
		77	81	84	88	91	95	98
6	BCY6136, 2 mpk, qw	208±112	337±123	501±172	626±182	856±245	1035±169	1266±39

(iii) Tumor Growth Inhibition Analysis

- 5 Tumor growth inhibition rate for BCY6033, BCY6136 and BCY6082 in the NCI-H1975 xenograft model was calculated based on tumor volume measurements at day 21 after the start of treatment.

Table 30: Tumor growth inhibition analysis

Gr	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	P value
1	Vehicle, qw	2371±470	--	--	--
2	BCY6033, 1 mpk, qw	306±81	12.9	92.9	<i>p</i> <0.001
3	BCY6033, 2 mpk, qw	95±12	4.0	102.5	<i>p</i> <0.001
4	BCY6033, 3 mpk, qw	29±4	1.2	105.4	<i>p</i> <0.001
5	BCY6136, 1 mpk, qw	205±117	8.6	97.5	<i>p</i> <0.001
6	BCY6136, 2 mpk, qw	99±30	4.2	102.3	<i>p</i> <0.001
7	BCY6136, 3 mpk, qw	94±30	4.0	102.4	<i>p</i> <0.001

8	BCY6082, 2 mpk, qw	1528±604	64.4	37.9	$p>0.05$
9	BCY6082, 5 mpk, qw	390±133	16.4	89.1	$p<0.001$

a. Mean $\pm$ SEM.

b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

(e) Results Summary and Discussion

- 5 In this study, the therapeutic efficacy of BCY6033, BCY6136 and BCY6082 in the NCI-H1975 xenograft model was evaluated. The measured body weights and tumor volumes of all treatment groups at various time points are shown in the Figures 11 to 13 and Tables 25 to 30.

The mean tumor size of vehicle treated mice reached 2371 mm³ on day 21. BCY6033 at 1 mg/kg (TV=306 mm³, TGI=92.9%, $p<0.001$), 2 mg/kg (TV=95 mm³, TGI=102.5%, $p<0.001$) and 3 mg/kg (TV=29 mm³, TGI=105.4%, $p<0.001$) produced dose-dependent antitumor activity. BCY6033 at 2 mg/kg and 3 mg/kg eradicated the tumors or regressed the tumor to small size, the treatments was suspended from day 35, and the tumors didn't show obvious re-growth in following 5-6 weeks monitoring.

- 15 BCY6136 at 1 mg/kg (TV=205 mm³, TGI=97.5%, $p<0.001$), 2 mg/kg (TV=99 mm³, TGI=102.3%, $p<0.001$) and 3 mg/kg (TV=94 mm³, TGI=102.4%, $p<0.001$) produced potent antitumor activity. BCY6136 at 2 mg/kg and 3 mg/kg eradicated the tumors or regressed the tumor to small size. The treatments was suspended from day 35, and the tumors in 3 mg/kg group didn't show obvious re-growth in following 5-6 weeks monitoring, however tumors in 2 mg/kg group showed obvious regrowth and didn't show significant tumor inhibition when resuming the dosing.

BCY6082 at 2 mg/kg (TV=1528 mm³, TGI=37.9%, $p>0.05$) didn't show obvious antitumor activity, BCY6082 at 5 mg/kg (TV=390 mm³, TGI=89.1%, $p<0.001$) produced significant antitumor activity.

- 25 In this study, one mouse treated with BCY6033 3mg/kg lost over 15% bodyweight during the monitoring, other mice maintained the bodyweight well.

Study 10. In vivo efficacy study of BCY6136 in the LU-01-0251 PDX model in Balb/c nude mice

30 (a) Study Objective

The objective of the research is to evaluate the *in vivo* anti-tumor efficacy of BCY6136 in the LU-01-0251 PDX model in Balb/c nude mice.

(b) Experimental Design

Group	Treatment	n	Dose (mg/kg)	Dosing Volume (µl/g)	Dosing Route	Schedule
1	Vehicle	5	--	10	iv	qw
2	BCY6136	5	1	10	iv	qw
3	BCY6136	5	2	10	iv	qw
4	BCY6136	5	3	10	iv	qw
5	ADC	5	3	10	iv	qw

(c) Experimental Methods and Procedures**(i) Tumor Inoculation**

Each mouse was inoculated subcutaneously at the right flank with LU-01-0251 of tumor fragment (~30 mm³) for tumor development. The treatment was started when the average tumor volume reached 174 mm³ for efficacy study. The test article administration and the animal number in each group are shown in the experimental design table.

(ii) Testing Article Formulation Preparation

Test article	Conc. (mg/ml)	Formulation
Vehicle	--	50 mM Acetate 10% sucrose pH 5
BCY6136	0.3	Dissolve 6.11 mg BCY6136 in 20 ml Acetate buffer ¹
	0.2	Dilute 940 µl 0.3 mg/ml BCY6136 stock with 470 µl Acetate buffer
	0.1	Dilute 470 µl 0.3 mg/ml BCY6136 stock with 940 µl Acetate buffer
ADC	0.3	Dilute 43 µl 10.47 mg/ml ADC stock with 1457 µl ADC buffer ²
1. Acetate buffer: 50 mM Acetate 10% sucrose pH 5		
2. ADC buffer: 20 mM Histidine pH 5.5		

(d) Results**(i) Body Weight change and Tumor Growth Curve**

Body weight and tumor growth curve are shown in Figure 14.

(ii) Tumor Volume Trace

Mean tumor volume on day 28 after the start of treatment in female Balb/c nude mice bearing LU-01-0251 xenograft is shown in Table 31.

Table 31: Tumor volume trace over time

	Group 1	Group 2	Group 3	Group 4	Group 5
Day	Vehicle	BCY6136, 1 mpk, qw	BCY6136, 2 mpk, qw	BCY6136, 3 mpk, qw	ADC, 3 mpk, qw

0	174±17	175±15	174±17	175±14	174±16
3	264±33	230±29	205±21	187±19	227±12
7	403±68	281±55	154±21	118±13	239±42
10	562±83	370±104	111±19	72±12	241±46
14	777±163	362±104	62±17	30±5	191±47
17	1021±246	437±136	46±13	17±3	139±39
21	1472±342	526±167	30±18	4±3	101±31
24	1790±417	491±132	32±24	1±1	70±23
28	2208±512	499±128	32±30	0±0	39±14

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for BCY6136 and ADC in the LU-01-0251 PDX model was calculated based on tumor volume measurements at day 28 after the start of the treatment.

5 **Table 32: Tumor growth inhibition analysis**

Group	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	<i>P</i> value
1	Vehicle, qw	2208±512	--	--	--
2	BCY6136, 1 mpk, qw	499±128	22.6	84.0	<i>p</i> <0.001
3	BCY6136, 2 mpk, qw	32±30	1.4	107.0	<i>p</i> <0.001
4	BCY6136, 3 mpk, qw	0±0	0.0	108.6	<i>p</i> <0.001
5	ADC, 3 mpk, qw	39±14	1.8	106.6	<i>p</i> <0.001

a. Mean ± SEM; b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

(e) Results Summary and Discussion

10 In this study, the therapeutic efficacy of BCY6136 and ADC in LU-01-0251 PDX model was evaluated. The measured body weight and tumor volume of all treatment groups at various time points are shown in the Figure 14 and Tables 31 and 32.

15 In this study, the mean tumor volume of vehicle treated mice reached 2208 mm³ on day 28 after the start of treatment. BCY6136 at 1 mg/kg, qw (TV=499 mm³, TGI=84.0%, *p*<0.001), 2 mg/kg, qw (TV=32 mm³, TGI=107.0%, *p*<0.001) and 3 mg/kg, qw (TV=0 mm³, TGI=108.6%, *p*<0.001) produced dose-dependent anti-tumor activity. ADC at 3 mg/kg, qw (TV=39 mm³, TGI=106.6%, *p*<0.001) showed significant anti-tumor activity.

Study 11: In vivo efficacy study of BCY6136 in the LU-01-0251 PDX model in**Balb/c nude mice****(a) Study Objective**

- The objective of the research is to evaluate the *in vivo* anti-tumor efficacy of BCY6136 in the LU-01-0251 PDX model in Balb/c nude mice.

(b) Experimental Design

Group	Treatment	n	Dose (mg/kg)	Dosing Volume (μl/g)	Dosing Route	Schedule
1	Vehicle	5	--	10	iv	Qw*21
2	BCY6136	5	1	10	iv	Qw*28
3 ^a	BCY6136	5	2	10	iv	Qw*70
4 ^b	BCY6136	5	3	10	iv	Qw*56
5 ^c	ADC	5	3	10	iv	Qw*70

- a. The dosing schedule was kept from day 0 to day 70 for all the mice of this group, then the mouse 3-2 and mouse 3-4 were further dosed with BCY6136 3 mg/kg qw from day 77 while the treatment of the other 3 mice was suspended. The dosing schedule was kept from day 0 to day 56 for all the mice of this group.
- b. The dosing schedule was kept from day 0 to day 70 for all the mice of this group.

(c) Experimental Methods and Procedures**(i) Tumor Inoculation**

- Each mouse was inoculated subcutaneously at the right flank with LU-01-0251 of tumor fragment (~30 mm³) for tumor development. The treatment was started when the average tumor volume reached 960 mm³ for efficacy study. The test article administration and the animal number in each group are shown in the experimental design table.

(ii) Testing Article Formulation Preparation

Test article	Conc. (mg/ml)	Formulation
Vehicle	--	25 mM Histidine 10% sucrose pH 7
BCY6136	0.3	0.3 mg/ml BCY6136 was prepared as in Study 10 hereinbefore
	0.2	Dilute 940 μl 0.3 mg/ml BCY6136 stock with 470 μl His-buffer ¹
	0.1	Dilute 470 μl 0.3 mg/ml BCY6136 stock with 940 μl His-buffer
ADC	0.3	Dilute 43 μl 10.47 mg/ml ADC stock with 1457 μl ADC-buffer ²
1. His-buffer: 25 mM Histidine 10% sucrose pH 7		
2. ADC-buffer: 20 mM Histidine pH 5.5		

(iii) Sample Collection

Tumor of mouse #3-2 was collected for FFPE on Day94. Tumors of mice #5-2 and 5-3 were collected and embed into 1 FFPE block on Day140.

(d) Results

(i) Body Weight change and Tumor Growth Curve

- 5 Body weight and tumor growth curve are shown in Figure 15.

(ii) Tumor Volume Trace

Mean tumor volume on day 0 to day 28 after the start of treatment in female Balb/c nude mice bearing LU-01-0251 xenograft is shown in Table 33.

Table 33: Tumor volume trace over time

Day	Group 1	Group 2	Group 3	Group 4	Group 5
	Vehicle	BCY6136, 1 mpk, qw	BCY6136, 2 mpk, qw	BCY6136, 3 mpk, qw	ADC, 3 mpk, qw
0	962±102	963±97	962±137	960±103	959±124
3	1176±108	1003±121	973±105	989±128	1043±158
7	1351±142	1056±151	873±125	890±98	1100±156
10	1591±179	1122±139	722±157	674±96	1172±188
14	1951±225	1417±191	503±151	342±64	1228±174
17	2301±344	1672±262	398±160	216±43	1143±186
21		1794±328	307±169	94±26	996±187
24		1867±408	261±168	62±14	867±178
28		2120±483	217±167	45±16	713±178

- 10 **(iii) Tumor Growth Inhibition Analysis**

Tumor growth inhibition rate for BCY6136 and ADC in the LU-01-0251 PDX model was calculated based on tumor volume measurements at day 17 after the start of the treatment.

Table 34: Tumor growth inhibition analysis

Group	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	P value
1	Vehicle, qw	2301±344	--	--	--
2	BCY6136, 1 mpk, qw	1672±262	72.7	47.0	$p > 0.05$
3	BCY6136, 2 mpk, qw	398±160	17.3	142.1	$p < 0.001$
4	BCY6136, 3 mpk, qw	216±43	9.4	155.6	$p < 0.001$
5	ADC, 3 mpk, qw	1143±186	49.7	86.3	$p < 0.01$

a. Mean ± SEM;

b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

(e) Results Summary and Discussion

In this study, the therapeutic efficacy of BCY6136 and ADC in LU-01-0251 PDX model was evaluated. The measured body weight and tumor volume of all treatment groups at various time points are shown in the Figure 15 and Tables 33 and 34.

In this study, the treatment was started when the average tumor volume reached 960 mm³. On day 17 after the start of treatment, the mean tumor volume of vehicle treated mice reached 2301 mm³. BCY6136 at 1 mg/kg qw (TV=1672 mm³, TGI=47.0%, $p > 0.05$) didn't show obvious antitumor activity; BCY6136 at 2 mg/kg qw (TV=398 mm³, TGI=142.1%, $p < 0.001$) and 3 mg/kg qw (TV=216 mm³, TGI=155.6%, $p < 0.001$) produced dose-dependent anti-tumor activity on day 17.

After 70 days' treatment with BCY6136 at 2 mg/kg qw, 3 in 5 of these mice showed complete tumor regression, the other 2 mice showed obvious tumor relapse from day 42 to day 77. Then further treatment with BCY6136 3 mg/kg qw was performed to the two relapse tumors from day 7, one of tumor showed obvious tumor regress while another one showed resistance to the treatment.

After 56 days' treatment with BCY6136 at 3 mg/kg qw, all the mice of this group showed complete tumor regression.

ADC at 3 mg/kg qw (TV=1143 mm³, TGI=86.3%, $p < 0.01$) showed obvious anti-tumor activity on day 17, after another 53 day' treatment, these mice showed further but not complete tumor regression.

In this study, there were some mice showed sudden bodyweight loss, this may have the relationship with the long term feeding of the immune-deficiency mice.

Study 12: In vivo efficacy study of BCY6033, BCY6136, BCY6082 and BCY6031 in the LU-01-0046 NSCLC PDX model in Balb/c nude mice

(a) Study Objective

The objective of the research is to evaluate the *in vivo* anti-tumor efficacy of BCY6033, BCY6136, BCY6082 and BCY6031 in large LU-01-0046 PDX tumors in Balb/c nude mice.

(b) Experimental Design

Group	Treatment	n	Dose (mg/kg)	Dosing Volume (μl/g)	Dosing Route	Schedule
BCYs						
1	Vehicle	5	--	10	iv	qw
2	BCY6082	5	1	10	iv	qw

3	BCY6082	5	3	10	iv	qw
4	BCY6033	5	1	10	iv	qw
5	BCY6033	5	3	10	iv	qw
6	BCY6136	5	1	10	iv	qw
7	BCY6136	5	3	10	iv	qw
8	ADC	5	3	10	iv	qw
9	BCY6031	5	3	10	iv	qw

(c) Experimental Methods and Procedures

(i) Tumor Inoculation

- Each mouse was inoculated subcutaneously at the right flank with LU-01-0046 of tumor fragment (~30 mm³) for tumor development. The treatment was started when the average tumor volume reaches 955 mm³ for BT17BDCs study and 1039 mm³ for BCYs study. The test article administration and the animal numbers in each group are shown in the experimental design table.

(ii) Testing Article Formulation Preparation

Test article	Conc. (mg/ml)	Formulation
Vehicle	-	50 mM Acetate 10% sucrose pH 5
BCY6033	0.1	Dilute 150 µl 1 mg/ml BCY6033 stock with 1350 µl Acetate buffer ¹
	0.3	Dilute 450 µl 1 mg/ml BCY6033 stock with 1050 µl Acetate buffer
BCY6136	0.1	Dilute 150 µl 1 mg/ml BCY6136 stock with 1350 µl Acetate buffer
	0.3	Dilute 450 µl 1 mg/ml BCY6136 stock with 1050 µl Acetate buffer
BCY6082	0.1	Dilute 150 µl 1 mg/ml BCY6082 stock with 1350 µl Acetate buffer
	0.3	Dilute 450 µl 1 mg/ml BCY6082 stock with 1050 µl Acetate buffer
BCY6031	0.3	Dissolve 5.72 mg BCY6031 in 5.6 ml Acetate buffer to make 1 mg/ml stock. Dilute 450 µl 1 mg/ml BCY6031 with 1050 µl Acetate buffer
ADC	0.3	Dilute 43 µl 10.47 mg/ml ADC stock solution into 1457 µl with buffer ²
1. Acetate buffer: 50 mM Acetate 10% sucrose pH5		
2. Dissolve 0.419 g His. hydrochloride in 100ml water, use 1M HCl adjust PH to 5.5		

(d) Results**(i) Body Weight change and Tumor Growth Curve**

Body weight and tumor growth curve are shown in Figure 16.

(ii) Tumor Volume Trace

- 5 Mean tumor volume over time in female Balb/c nude mice bearing LU-01-0046 is shown in Table 35.

Table 35: Tumor volume trace over time (BCY's Section)

Group	Treatment	Days after the start of treatment							
		0	4	8	11	15	18	22	
1	Vehicle, qw	1044±115	1762±178	2404±262	--	--	--	--	--
2	BCY6082, 1 mpk, qw	1049±133	1592±178	2279±168	--	--	--	--	--
3	BCY6082, 3 mpk, qw	1033±111	1040±124	1294±182	1298±101	1849±189	2052±168	1999±425	
4	BCY6033, 1 mpk, qw	1030±124	1173±227	1791±324	2408±484	--	--	--	--
5	BCY6033, 3 mpk, qw	1046±128	555±85	441±144	182±76	163±94	114±54	88±76	
6	BCY6136, 1 mpk, qw	1037±130	1163±146	1927±283	2483±530	--	--	--	--
7	BCY6136, 3 mpk, qw	1036±100	784±146	548±107	362±110	325±122	275±152	233±187	
8	ADC, 3 mpk, qw	1033±114	1155±230	2200±505	--	--	--	--	--
9	BCY6031, 3 mpk, qw	1042±117	820±149	1319±233	901±188	672±198	522±315	515±323	

Note: the tumor volume trace didn't show after the day22 for the group3, 5, 7 and 9.

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for test articles in the LU-01-0046 PDX model was calculated based on tumor volume measurements at day 22 and day 28 respectively for the two section studies after the start of the treatment.

5 Table 36: Tumor growth inhibition analysis (BCYs section on day 22)

Group	Treatment	Tumor Volume	T/C ^b (%)	TGI (%)	P value
1	Vehicle, qw	6186±596*	--	--	--
2	BCY6082, 1 mpk, qw	5805±428*	93.8	7.5	<i>p</i> >0.05
3	BCY6082, 3 mpk, qw	1999±425	32.3	81.2	<i>p</i> <0.01
4	BCY6033, 1 mpk, qw	4384±881*	70.9	34.8	<i>p</i> >0.05
5	BCY6033, 3 mpk, qw	88±76	1.4	118.6	<i>P</i> <0.001
6	BCY6136, 1 mpk, qw	4564±981*	73.8	31.4	<i>p</i> >0.05
7	BCY6136, 3 mpk, qw	233±187	3.8	115.6	<i>p</i> <0.001
8	ADC, 3 mpk, qw	5446±1250*	88.0	14.2	<i>p</i> >0.05
9	BCY6031, 3 mpk, qw	515±323	8.3	110.2	<i>p</i> <0.001

a. Mean ± SEM;

b. Tumor Growth Inhibition is calculated by dividing the average tumor volume of the treated group by the average tumor volume of the control group (T/C).

*Some groups was terminated before day 22, and the tumor size was calculated by exponential growth equation acquisition as below:

Vehicle group: $Y = 995.4 \times \exp(0.1134 \times X)$.

BCY6082, 1mpk group: $Y = 939.1 \times \exp(0.1128 \times X)$.

BCY6033, 1mpk group: $Y = 846.6 \times \exp(0.0945 \times X)$.

BCY6136, 1mpk group: $Y = 855.0 \times \exp(0.0974 \times X)$.

ADC, 3mpk group: $Y = 757.4 \times \exp(0.1312 \times X)$.

(e) Results Summary and Discussion

In this study, the therapeutic efficacy of test articles in large LU-01-0046 tumors was evaluated. The measured body weights and tumor volumes of all treatment groups at various time points are shown in the Figure 16 and Tables 35 and 36.

In BCYs study, the mean tumor size of vehicle treated mice was calculated as 6186 mm³ on day 22. BCY6082, BCY6033, BCY6136 at 1 mg/kg and ADC at 3mg/kg didn't show obvious anti-tumor activity when starting treatment from tumor size of 1000mm³.

BCY6082 (TV=1999 mm³, TGI=81.2%, $p<0.01$), BCY6033 (TV=88 mm³, TGI=118.6%, $p<0.001$), BCY6136 (TV=233 mm³, TGI=115.6%, $p<0.001$) and BCY6031 (TV=115 mm³, TGI=110.2%, $p<0.001$) at 3 mg/kg produced significant anti-tumor activity. Among them, BCY6033 and BCY6136 eradicated 2/5 and 4/5 tumors completely.

Study 13: In vivo efficacy of BCY6136 in Balb/c nude mice bearing LU-01-0046

NSCLC PDX model

(a) Study Objective

The objective of the research was to evaluate the *in vivo* therapeutic efficacy of BCY6136 in Balb/c nude mice bearing LU-01-0046 NSCLC PDX model.

(b) Experimental Design

Group	Treatment	n	Dose (mg/kg)	Dosing Route	Schedule
1	Vehicle	5	--	i.v.	qw*2 w
2	BCY6136	5	1	i.v.	qw*3 w
3	BCY6136	5	2	i.v.	qw*4 w
4	BCY6136	5	3	i.v.	qw*4 w
5	ADC	5	3	i.v.	qw*3 w
6	ADC	5	5	i.v.	qw*3 w

Note: Groups were terminated when average tumor volume reached over 2000 mm³ and tumors were harvested for FFPE: Group 1 on PG-D14, group 5 on PG-D18, group 2 & 6 on PG-D21 and group 3 & 4 on PG-D31.

(c) Experimental Methods and Procedures

(i) Tumor Inoculation

Each mouse was inoculated subcutaneously at the right flank with certain kind of tumor fragment (~30 mm³) for tumor development. The treatments were started when the average tumor volume reached approximately 198 mm³. The test article administration and the animal numbers in each group are shown in the experimental design table.

(ii) Testing Article Formulation Preparation

Gr	Compounds	Dose (mg/kg)	Con. (mg/ml)	Formulation
1	Vehicle	-	-	50 mM Acetate, 10% Sucrose pH 5 (without DMSO)
2	BCY6136	1	0.1	Dissolve 10.93 mg BCY6136 in 10.766 ml vehicle, ultrasonic simply to make the 1 mg/ml BCY6136 stock solution Dilute 150 μ l 1 mg/ml BCY6136 stock solution with 1350 μ l vehicle
3	BCY6136	2	0.2	Dilute 300 μ l 1 mg/ml BCY6136 stock solution with 1200 μ l vehicle
4	BCY6136	3	0.3	Dilute 450 μ l 1 mg/ml BCY6136 stock solution with 1050 μ l vehicle
Buffer 2: Dissolve 0.419 g His. hydrochloride in 100 ml water, use 1 M HCl adjust pH to 5.5				
5	ADC	3	0.3	Dilute 43 μ l 10.47 mg/ml ADC stock solution with 1457 μ l with buffer 2
6	ADC	5	0.5	Dilute 71.6 μ l 10.47 mg/ml ADC stock solution with 1428.4 μ l with buffer 2

Note: The dosing formulation frequently is fresh prepared timely.

(iii) Sample Collection

- Groups were terminated when average tumor volume reached over 2000 mm³ and tumors were harvested for FFPE after the last measurement: Group 1 on PG-D14, group 5 on PG-D18, group 2 & 6 on PG-D21 and group 3 & 4 on PG-D31.

(d) Results

(i) Body Weight change and Tumor Growth Curve

Body weight and tumor growth curve are shown in Figure 17.

(ii) Tumor Volume Trace

- 10 Mean tumor volume over time in female Balb/c nude mice bearing LU-01-0046 NSCLC PDX model is shown in Table 37.

Table 37: Tumor volume trace over time (mm³)

Gr	1	2	3	4	5	6
Treatment	Vehicle qw	BCY6136 1 mpk, qw	BCY6136 2 mpk, qw	BCY6136 3 mpk, qw	ADC 3 mpk, qw	ADC 5 mpk, qw

	201 ±					195 ±
0	37	198 ± 39	201 ± 40	200 ± 46	195 ± 28	40
	441 ±					389 ±
3	82	310 ± 59	283 ± 77	155 ± 40	418 ± 99	68
	927 ±		423 ±			596 ±
7	171	547 ± 88	132	74 ± 19	643 ± 159	116
	1546 ±	747 ±	321 ±			882 ±
10	377	121	108	31 ± 8	938 ± 230	134
	2307 ±	1058 ±				1215
14	594	140	264 ± 95	26 ± 11	1475 ± 466	± 193
		1390 ±				1576
17	-	205	127 ± 41	26 ± 13	2281 ± 556	± 228
		2138 ±				2049
21	-	301	118 ± 34	64 ± 42	-	± 242
24	-	-	101 ± 40	99 ± 63	-	-
			255 ±	276 ±		
28	-	-	140	176	-	-
			582 ±	477 ±		
31	--	-	346	283	-	-

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for test articles in Balb/c nude mice bearing LU-01-0046 PDX model was calculated based on tumor volume measured on PG-D14.

Table 38: Tumor growth inhibition analysis

Gr	Treatment	Tumor Volume (mm ³) ^a	T/C (%) ^b	TGI (%) ^c	<i>P</i> value compared with vehicle
1	Vehicle qw	2307 ± 594	--	--	--
2	BCY6136 1 mpk, qw	1058 ± 140	45.9	59.1	<i>p</i> <0.05
3	BCY6136 2 mpk, qw	264 ± 95	11.4	97.0	<i>p</i> <0.001
4	BCY6136 3 mpk, qw	26 ± 11	1.1	108.3	<i>p</i> <0.001
5	ADC	1475 ± 466	63.9	39.2	<i>p</i> >0.05

3 mpk, qw					
6	ADC	1215 ± 193	52.7	51.6	$p > 0.05$
5 mpk, qw					

a. Mean ± SEM.

b. Tumor Growth Inhibition was calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

c. TGI was calculated for each group using the formula: $TGI (\%) = [1 - (T_i - T_0) / (V_i - V_0)] \times 100$

5 (e) Results Summary and Discussion

In the present study, the therapeutic efficacy of test articles in the LU-01-0046 PDX model was evaluated. The measured body weights and tumor volumes of all treatment groups at various time points were shown in the Figure 17 and Tables 37 and 38.

The mean tumor size of vehicle treated mice reached 2307 mm³ on PG-D14. BCY6136 at 1 mg/kg (TV=1058 mm³, TGI=59.1%, $p < 0.05$), at 2 mg/kg (TV=264 mm³, TGI=97.0%, $p < 0.001$) and at 3 mg/kg (TV=26 mm³, TGI=108.3%, $p < 0.001$) produced dose-dependent antitumor activity. ADC at 3 mg/kg and 5 mg/kg did not show obvious antitumor activity ($p > 0.05$).

In this study, all of the group's animals maintained the body weight well.

15 Study 14: In vivo efficacy study of BCY6033, BCY6136, BCY6082, BCY6173, BCY6175 and BCY6031 in the LU-01-0046 NSCLC PDX model in Balb/c nude mice

(a) Study Objective

The objective of the research is to evaluate the *in vivo* anti-tumor efficacy of test articles in the LU-01-0046 NSCLC PDX model in Balb/c nude mice.

20 (b) Experimental Design

Group	Treatment	n	Dose (mg/kg)	Dosing Volume (μl/g)	Dosing Route	Schedule
Part 1						
1	Vehicle	5	--	10	iv	qw
2	BCY6033	5	1/2	10	iv	qw
3	BCY6033	5	3	10	iv	qw
4	BCY6136	5	1/2	10	iv	qw
5	BCY6136	5	3	10	iv	qw
6	BCY6082	5	1	10	iv	qw
7	BCY6082	5	3	10	iv	qw
Part 2						
8	Vehicle	5	--	10	iv	qw
9	BCY6173	5	1	10	iv	qw
10	BCY6173	5	3	10	iv	qw

11	BCY6175	5	3	10	iv	qw
12	BCY6031	5	3	10	iv	qw

(c) Experimental Methods and Procedures

(i) Tumor Inoculation

- Each mouse was inoculated subcutaneously at the right flank with LU-01-0046 of tumor fragment (~30 mm³) for tumor development. The treatment was started when the average tumor volume reaches 200 mm³ for part 1 study and 192 mm³ for part 2 study. The test article administration and the animal numbers in each group are shown in the experimental design table.

(ii) Testing Article Formulation Preparation

Test article	Conc. (mg/ml)	Formulation
Vehicle	-	50mM Acetate 10% sucrose pH 5
BCY6033	0.1	Dilute 150 µl 1 mg/ml BCY6033 stock with 1350 µl Acetate buffer ¹
	0.3	Dilute 450 µl 1 mg/ml BCY6033 stock with 1050 µl Acetate buffer
BCY6136	0.1	Dilute 150 µl 1 mg/ml BCY6136 stock with 1350 µl Acetate buffer
	0.3	Dilute 450 µl 1 mg/ml BCY6136 stock with 1050 µl Acetate buffer
BCY6082	0.1	Dilute 150 µl 1 mg/ml BCY6082 stock with 1350 µl Acetate buffer
	0.3	Dilute 450 µl 1 mg/ml BCY6082 stock with 1050 µl Acetate buffer
BCY6173	0.1	Dissolve 3.65 mg BCY6173 in 3.5 ml Acetate buffer to make 1 mg/ml stock. Dilute 150 µl 1 mg/ml BCY6173 with 1350 µl Acetate buffer
	0.3	Dilute 450 µl 1 mg/ml BCY6173 stock with 1050 µl Acetate buffer
BCY6175	0.3	Dissolve 3.02 mg BCY6175 in 2.9 ml Acetate buffer to make 1mg/ml stock. Dilute 450 µl 1 mg/ml BCY6175 with 1050 µl Acetate buffer
BCY6031	0.3	Dissolve 5.72 mg BCY6031 in 5.6 ml Acetate buffer to make 1mg/ml stock. Dilute 450 µl 1 mg/ml BCY6031 with 1050 µl Acetate buffer

1. Acetate buffer: 50 mM Acetate 10% sucrose pH 5

(d) Results

(i) Body Weight change and Tumor Growth Curve

Body weight and tumor growth curve are shown in Figures 18 to 22.

(ii) Tumor Volume Trace

- 5 Mean tumor volume on day 21 after the start of treatment in female Balb/c nude mice bearing LU-01-0046 is shown in Tables 39 and 40.

Table 39: Tumor volume trace over time (Part 1)

Gr	Treatment	Days after the start of treatment						
		0	3	6	10	14	17	21
1	Vehicle, qw	202±2	328±4	536±6	953±10	1386±97	1833±13	2551±24
		6	8	8	7		2	2
2	BCY6033, 1 mpk, qw	201±2	285±4	449±8	623±11	891±196	967±228	1285±23
		3	7	7	2			4
3	BCY6033, 3 mpk, qw	201±2	187±4	91±34	37±14	3±3	0±0	0±0
		6	3					
4	BCY6136, 1 mpk, qw	200±3	293±5	426±9	682±15	964±194	976±258	1285±23
		3	6	1	1			4
5	BCY6136, 3 mpk, qw	201±3	194±3	135±2	52±18	13±9	4±4	0±0
		3	1	7				
6	BCY6082, 1 mpk, qw	201±2	295±4	466±6	877±80	1201±10	1502±10	1826±22
		9	3	5		6	8	4
7	BCY6082, 3 mpk, qw	201±3	235±3	310±4	398±65	634±136	729±184	1042±29
		4	6	4				0

Table 40: Tumor volume trace over time (Part 2)

G	Treatment	Days after the start of treatment						
		0	3	7	10	14	17	21

8	Vehicle, qw	192±3 0	311±8 3	562±14 6	830±23 0	1320±44 4	1652±52 8	2342±65 1
BCY6173								
9	, 1 mpk, qw	191±3 3	318±5 8	553±88	817±16 5	1314±27 6	1546±27 6	2151±26 2
BCY6173								
10	, 3 mpk, qw	192±3 7	259±5 1	400±53	455±28	636±92	646±138	890±260
BCY6175								
11	, 3 mpk, qw	192±4 2	186±5 7	92±38	19±11	0±0	0±0	0±0
BCY6031								
12	, 3 mpk, qw	191±3 8	207±4 6	387±70	355±11 0	544±159	643±185	874±281

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for test articles in the LU-01-0046 PDX model was calculated based on tumor volume measurements at day 21 after the start of the treatment.

Table 41: Tumor growth inhibition analysis (Part 1)

Group	Treatment	Tumor Volume	T/C ^b (%)	TGI (%)	P value
1	Vehicle, qw	2551±242	--	--	--
2	BCY6033, 1 mpk, qw	1285±234	50.4	53.9	$p < 0.001$
3	BCY6033, 3 mpk, qw	0±0	0.0	108.6	$p < 0.001$
4	BCY6136, 1 mpk, qw	1285±234	50.4	53.9	$p < 0.001$
5	BCY6136, 3 mpk, qw	0±0	0.0	108.5	$p < 0.001$
6	BCY6082, 1 mpk, qw	1826±224	71.6	30.8	$p < 0.05$

7	BCY6082, 3 mpk, qw	1042±290	40.8	64.2	$p < 0.001$
---	-----------------------	----------	------	------	-------------

a. Mean $\pm$ SEM; b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

Table 42: Tumor growth inhibition analysis (Part 2)

Group	Treatment	Tumor Volume	T/C ^b (%)	TGI (%)	P value
8	Vehicle, qw	2342±651	--	--	--
9	BCY6173, 1 mpk, qw	2151±262	91.8	8.9	$p > 0.05$
10	BCY6173, 3 mpk, qw	890±260	38.0	67.5	$p < 0.05$
11	BCY6175, 3 mpk, qw	0±0	0.0	108.9	$p < 0.001$
12	BCY6031, 3 mpk, qw	874±281	37.3	68.2	$p < 0.05$

a. Mean $\pm$ SEM; b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

(e) Results Summary and Discussion

In this study, the therapeutic efficacy of test articles in the LU-01-0046 PDX model was evaluated. The measured body weights and tumor volumes of all treatment groups at various time points are shown in the Figures 18 to 22 and Tables 39 to 42.

- 10 In part 1 study, the mean tumor size of vehicle treated mice reached 2551 mm³ on day 21 after the start of treatment.
- BCY6033 at 1/2 mg/kg, qw (TV=1285 mm³, TGI=53.9%, $p < 0.001$) and BCY6136 at 1/2 mg/kg, qw (TV=1285 mm³, TGI=53.9%, $p < 0.001$) produced significant anti-tumor activity, but didn't exhibit any tumor regression. BCY6033 at 3 mg/kg, qw (TV=0 mm³, TGI=108.6%, $p < 0.001$) and BCY6136 at 3 mg/kg, qw (TV=0 mm³, TGI=108.5%, $p < 0.001$) completely eradicated the tumors, 1 of 5 tumors respectively in BCY6033 and BCY6136 3 mg/kg groups showed regrowth after the dosing suspension and the tumors were resistant to BCY6033 or BCY6136 treatment when resuming the dosing. The remaining tumors in the BCY6033 and BCY6136 groups (4/5 for each group) showed no regrowth after 80 days of dosing suspension.
- 20 BCY6082 at 1 mg/kg, qw (TV=1826 mm³, TGI=30.8%, $p < 0.05$) and 3 mg/kg, qw (TV=1042 mm³, TGI=64.2%, $p < 0.001$) produced dose-dependent anti-tumor activity, but didn't show tumor regression.

In part 2 study, the mean tumor size of vehicle treated mice reached 2342 mm³ on day 21 after the start of treatment. BCY6173 at 1 mg/kg, qw (TV=2151 mm³, TGI=8.9%, $p > 0.05$) did not show anti-tumor activity. BCY6173 at 3 mg/kg, qw (TV=890 mm³, TGI=67.5%, $p < 0.05$) produced obvious anti-tumor activity.

- 5 BCY6175 at 3 mg/kg, qw (TV=0 mm³, TGI=108.9%, $p < 0.001$) completely eradicated 4/5 tumors on day 14. BCY6031 at 3 mg/kg, qw (TV=874 mm³, TGI=68.2%, $p < 0.05$) produced obvious anti-tumor activity, but didn't show any tumor regression.

Study 15: In vivo efficacy study of BCY6136 in the LU-01-0412 NSCLC PDX model in Balb/c nude mice

(a) Study Objective

The objective of the project is to evaluate the *in vivo* therapeutic efficacy of BCY6136 in the LU-01-0412 NSCLC PDX model in BALB/c nude mice.

(b) Experimental Design

Gr	Treatment	n	Dose (mg/kg)	Dosing Volume (μl/g)	Dosing Route	Schedule
1	Vehicle	6	--	10	iv	Qw, 4
2	BCY6136	6	1	10	iv	Qw, 4
3	BCY6136	6	3	10	iv	Qw, 4
4	BCY8245	6	3	10	iv	Qw, 4
5	BCY8781	6	3	10	iv	Qw, 4

(c) Experimental Methods and Procedures

(i) Tumor Inoculation

Each mouse was inoculated subcutaneously at the right flank with LU-01-0412 tumor fragment (~30 mm³) for tumor development. Animals were randomized when the average tumor volume reached 159 mm³. The test article administration and the animal numbers in each group were shown in the experimental design table.

(ii) Testing Article Formulation Preparation

Test article	Conc. (mg/ml)	Formulation
Vehicle	-	25 mM Histidine 10% sucrose pH7
BCY6136	1	Dissolve 6.06 mg BCY6136 in 5.969 ml 50 mM Acetate/acetic acid pH5 10% sucrose
	0.1	Dilute 180 μl 1 mg/ml BT5528 with 1620 μl 50 mM Acetate/acetic acid pH5 10% sucrose

	0.3	Dilute 540 μ l 1 mg/ml BT5528 with 1260 μ l 50 mM Acetate/acetic acid pH5 10% sucrose
BCY8245	1	Dissolve 4.15 mg BCY8245 powder in 4.121 ml vehicle buffer
	0.3	Dilute 540 μ l 1 mg/ml BCY8245 with 1260 μ l vehicle buffer
BCY8781	1	Dissolve 4.08 mg BCY8781 powder in 80.8 μ l DMSO, then dilute to 1 mg/ml with 3.958 vehicle buffer
	0.3	Dilute 540 μ l 1 mg/ml BCY8781 with 1260 μ l vehicle buffer

(iii) Sample Collection

Plasma from vehicle and 3 extra mice treated with BCY6136, BCY8245 and BCY8781 were collected at 30 min and 24 h post dosing. Tumor from vehicle and 3 extra mice treated with BCY6136, BCY8245 and BCY8781 were collected at 24 h post dosing.

5 (d) **Results**(i) Body Weight change and Tumor Growth Curve

Body weight and tumor growth curves are shown in Figure 23.

(ii) Tumor Volume Trace

Mean tumor volume over time in female BALB/c nude mice bearing LU-01-0412 xenograft is shown in Table 43.

Table 43: Tumor volume trace over time

	Group 1	Group 2	Group 3	Group 4	Group 5
Days	Vehicle	BCY6136	BCY6136	BCY8245	BCY8781
	Qw*4	1 mpk, Qw*4	3 mpk, Qw*4	3 mpk, Qw*4	3 mpk, Qw*4
0	159 $\pm$ 11	159 $\pm$ 13	159 $\pm$ 11	159 $\pm$ 12	159 $\pm$ 11
4	255 $\pm$ 12	214 $\pm$ 16	197 $\pm$ 16	168 $\pm$ 18	176 $\pm$ 21
7	309 $\pm$ 20	237 $\pm$ 16	195 $\pm$ 16	132 $\pm$ 10	167 $\pm$ 13
11	395 $\pm$ 31	246 $\pm$ 19	156 $\pm$ 18	78 $\pm$ 4	107 $\pm$ 15
14	464 $\pm$ 31	300 $\pm$ 18	177 $\pm$ 29	45 $\pm$ 5	72 $\pm$ 12
18	521 $\pm$ 26	369 $\pm$ 32	210 $\pm$ 32	21 $\pm$ 2	44 $\pm$ 8
21	611 $\pm$ 33	470 $\pm$ 46	225 $\pm$ 32	11 $\pm$ 1	31 $\pm$ 6
25	737 $\pm$ 68	632 $\pm$ 47	252 $\pm$ 37	6 $\pm$ 1	20 $\pm$ 6
28	788 $\pm$ 80	664 $\pm$ 52	299 $\pm$ 37	2 $\pm$ 1	14 $\pm$ 5
32	1104 $\pm$ 142	758 $\pm$ 70	416 $\pm$ 52	1 $\pm$ 1	12 $\pm$ 5

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for BCY6136, BCY8245 and BCY8781 in the LU-01-0412 xenograft model was calculated based on tumor volume measurements on day 32 after the start of the treatment.

Table 44: Tumor growth inhibition analysis

Group	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	P value
1	Vehicle, qw*4	1104±142	--	--	--
2	BCY6136, 1mpk, qw*4	758±70	68.6	36.7	<i>p</i> <0.05
3	BCY6136, 3mpk, qw*4	416±52	37.6	72.9	<i>p</i> <0.001
4	BCY8245, 3 mpk, qw*4	1±1	0.1	116.8	<i>p</i> <0.001
5	BCY8781, 3 mpk, qw*4	12±5	1.0	115.6	<i>p</i> <0.001

5 a. Mean ± SEM;

b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

(e) Results Summary and Discussion

10 In this study, the therapeutic efficacy of BCY6136, BCY8245 and BCY8781 in the LU-01-0412 xenograft model was evaluated. The measured body weight and tumor volume of all treatment groups at various time points are shown in Figure 23 and Tables 43 and 44.

The mean tumor volume of vehicle treated mice reached 1104 mm³ on day 32 after the start of treatment. BCY6136 at 1 mg/kg, qw *4 (TV=758 mm³, TGI=36.7%, *p*<0.05) and 3 mg/kg, qw*4 (TV=416 mm³, TGI=72.9%, *p*<0.001) produced dose-dependent antitumor activity, but 15 didn't show any tumor regression. BCY8245 at 3 mg/kg, qw*4 (TV=1 mm³, TGI=116.8%, *p*<0.001) and BCY8781 at 3 mg/kg, qw*4 (TV=12 mm³, TGI=115.6%, *p*<0.001) regressed the tumors obviously. Among them, 5 of 6 tumor treated with BCY8245 3 mg/kg and 2 of 6 tumor treated with d BCY8781 3 mg/kg were completely eradicated on day 32.

In this study, animals in all groups maintained the body weight well.

20

Study 16: In vivo efficacy study of BCY6136 in treatment of LU-01-0486 PDX model in Balb/c nude mice

(a) Study Objective

25 The objective of the research is to evaluate the *in vivo* anti-tumor efficacy of BCY6136 in the LU-01-0486 PDX model in Balb/c nude mice.

(b) Experimental Design

Gr	Treatment	n	Dose (mg/kg)	Dosing Volume (µl/g)	Dosing Route	Schedule
1	Vehicle	5	--	10	iv	qw
2	BCY6136	5	1	10	iv	qw
3	BCY6136	5	2	10	iv	qw
4	BCY6136	5	3	10	iv	qw

(c) Experimental Methods and Procedures**(i) Tumor Inoculation**

Each mouse was inoculated subcutaneously at the right flank with LU-01-0486 of tumor fragment (~30 mm³) for tumor development. The treatment was started when the average tumor volume reached 180 mm³ for efficacy study. The test article administration and the animal number in each group are shown in the experimental design table.

(ii) Testing Article Formulation Preparation

Test article	Conc. (mg/ml)	Formulation
Vehicle	--	50 mM Acetate 10% sucrose pH 5
BCY6136	0.3	0.3 mg/ml BCY6136 was prepared as described in Study 10
	0.2	Dilute 940 µl 0.3 mg/ml BCY6136 with 470 µl Acetate buffer ¹
	0.1	Dilute 470 µl 0.3 mg/ml BCY6136 with 940 µl Acetate buffer
<i>1. Acetate buffer: 50 mM Acetate 10% sucrose pH 5</i>		

(d) Results**(i) Body Weight change and Tumor Growth Curve**

Body weight and tumor growth curve are shown in Figure 24.

(ii) Tumor Volume Trace

Mean tumor volume on day 14 after the start of treatment in female Balb/c nude mice bearing LU-01-0486 xenograft is shown in Table 45.

Table 45: Tumor volume trace over time

Group	Treatment	Days after the start of treatment				
		0	3	7	10	14
1	Vehicle, qw	179±20	232±30	358±45	450±47	651±112
2	BCY6136, 1 mpk, qw	180±23	221±20	326±34	420±34	638±71

3	BCY6136, 2 mpk, qw	179±27	222±26	365±44	459±82	645±105
4	BCY6136, 3 mpk, qw	180±25	209±37	304±51	348±77	449±115

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for BCY6136 in the LU-01-0486 PDX model was calculated based on tumor volume measurement at day 14 after the start of the treatment.

Table 46: Tumor growth inhibition analysis

Group	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	P value
1	Vehicle, qw	651±112	--	--	--
2	BCY6136, 1 mpk, qw	638±71	98.0	3.0	$p > 0.05$
3	BCY6136, 2 mpk, qw	645±105	99.1	1.2	$p > 0.05$
4	BCY6136, 3 mpk, qw	449±115	68.9	43.1	$p > 0.05$

- 5 a. Mean ± SEM; b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

(e) Results Summary and Discussion

- In this study, the therapeutic efficacy of BCY6136 in LU-01-0486 PDX model was evaluated. The measured body weight and tumor volume of all treatment groups at various time points are shown in the Figure 24 and Tables 45 and 46.

- 10 In this study, the mean tumor volume of vehicle treated mice reached 651 mm³ on day 14 after the start of treatment. BCY6136 at 1 mg/kg, qw (TV=638 mm³, TGI=3.0%, $p > 0.05$) and 2 mg/kg, qw (TV=645 mm³, TGI=1.2%, $p > 0.05$) didn't show any anti-tumor activity. BCY6136 at 3 mg/kg, qw (TV=449 mm³, TGI=43.1%, $p > 0.05$) produced slight anti-tumor activity without
15 statistical significance.

Study 17: In vivo efficacy test of BCY6033, BCY6136 and BCY6082 in treatment of MDA-MB-231-luc xenograft in Balb/c nude mice

(a) Study Objective

- 20 The objective of the research was to evaluate the *in vivo* anti-tumor efficacy of BCY6033, BCY6136 and BCY6082 in treatment of MDA-MB-231-luc xenograft model in Balb/c nude mice.

(b) Experimental Design

Gr	Treatment	n	Dose (mg/kg)	Dosing Volume (μl/g)	Dosing Route	Schedule
1	Vehicle	3	--	10	<i>iv</i>	qw
2	BCY6033	3	1	10	<i>iv</i>	qw
3	BCY6033	3	2	10	<i>iv</i>	qw
4	BCY6033	3	3	10	<i>iv</i>	qw
5	BCY6136	3	1	10	<i>iv</i>	qw
6	BCY6136	3	2	10	<i>iv</i>	qw
7	BCY6136	3	2	10	<i>iv</i>	qw
8	BCY6082	3	2	10	<i>iv</i>	qw
9	BCY6082	3	5	10	<i>iv</i>	qw

(c) Experimental Methods and Procedures

(i) Cell Culture

The cells growing in an exponential growth phase were harvested and counted for tumor inoculation.

5 (ii) Tumor Inoculation

Each mouse was inoculated subcutaneously at the right flank with MDA-MB-231-luc tumor cells (10×10^6) in 0.1ml of PBS with 0.1 ml matrigel for tumor development. 36 animals were randomized when the average tumor volume reached 159 mm³. The test article administration and the animal numbers in each group were shown in the experimental

10 design table.

(iii) Testing Article Formulation Preparation

Treatment	Dose (mg/ml)	Formulation
Vehicle		50 mM Acetate, 10% sucrose pH=5
BCY6033	1	Dissolve 6.71 mg BCY6033 into 6.710 ml formulation buffer
	0.3	Dilute 270 μl 1 mg/ml BCY6033 into 630 μl formulation buffer
	0.2	Dilute 180 μl 1 mg/ml BCY6033 into 720 μl formulation buffer
	0.1	Dilute 90 μl 1 mg/ml BCY6033 into 810 μl formulation buffer
BCY6136	1	Dissolve 3.79 mg BCY6136 into 3.695ml formulation buffer
	0.3	Dilute 270 μl 1 mg/ml BCY6136 into 630 μl formulation buffer

BCY6082	0.2	Dilute 180 µl 1 mg/ml BCY6136 into 720 µl formulation buffer
	0.1	Dilute 90 µl 1 mg/ml BCY6136 into 810 µl formulation buffer
	1	Weigh and dissolve 4.30 mg BCY6082 into 4.162 ml formulation buffer
BCY6082	0.5	Dilute 450 µl 1 mg/ml BCY6082 into 450 µl formulation buffer
	0.2*	Dilute 180 µl 1 mg/ml BCY6082 into 720 µl formulation buffer

(iv) Sample Collection

On PG-D24, we collected and fixed the tumors of Group 1, 8 and 9 for FFPE.

On PG-D33, we collected and fixed the tumors of Group 2 and 5 for FFPE.

At the end of study, we collected and fixed the tumors of Group 3, 4, 6 and 7 for FFPE.

5 (d) **Results**(i) Body Weight change and Tumor Growth Curve

Body weight and tumor growth are shown in Figures 25 to 27.

(ii) Tumor Volume Trace

Mean tumor volume over time in female Balb/c nude mice bearing MDA-MB-231-luc

10 xenograft is shown in Tables 47 to 49.

Table 47: Tumor volume trace (PG-D0~PG-D17)

Gr	Treatme nt	Days after the start of treatment							
		0	2	4	7	9	11	14	17
1	Vehicle, qw	159±1 4	269±8	306±1 9	425±5 2	688±5 4	908±5 4	1064±9 8	1315±9 5
BCY6033									
2	, 1 mpk, qw	159±6	219±1 9	221±5 5	296±7 6	329±6 4	421±7 7	479±84	609±12 2
BCY6033									
3	, 2 mpk, qw	159±1 0	240±7 3	215±5 7	201±4 7	109±3 6	84±34	64±32	59±35
4	BCY6033 ,	158±7	189±2 7	147±3 2	109±2 6	79±11	66±7	41±5	31±6

3 mpk, qw									
BCY6136									
5	,	159±1	226±3	221±5	310±7	416±8	526±7	636±92	809±13
	1 mpk, qw	0	6	4	2	9	7		5
BCY6136									
6	,	159±1	218±1	182±2	182±2	101±2	77±24	36±4	41±10
	2 mpk, qw	6	7	2	6	0			
BCY6136									
7	,	158±5	241±1	259±6	325±1	258±1	246±1	162±19	178±10
	3 mpk, qw		2		4	2	5		
BCY6082									
8	,	159±1	210±1	242±1	305±1	445±5	611±7	734±13	926±10
	2 mpk, qw	3	0	6	9	8	6	9	5
BCY6082									
9	,	159±7	227±3	247±4	250±6	276±7	241±6	220±56	184±85
	5 mpk, qw		1	7	5	9	1		

Table 48: Tumor volume trace (PG-D19~PG-D33)

Gr	Treatme nt	Days after the start of treatment						
		19	21	24	26	28	31	33
1	Vehicle, qw	1453±12 8	1661±17 3	--	--	--	--	--
BCY603								
2	3, 1 mpk, qw	724±162	880±156	1069±18 9	1182±16 4	1342±16 6	1647±11 3	--
3	BCY603 3,	61±35	67±44	100±76	133±96	163±106	221±143	257±15 2

	2 mpk, qw							
	BCY603							
4	3, 3 mpk, qw	29±7	22±12	22±8	21±9	21±10	43±20	57±29
	BCY613							
5	6, 1 mpk, qw	879±190	994±213	1253±31 3	1431±35 3	1507±25 3	2181±60 9	--
	BCY613							
6	6, 2 mpk, qw	35±9	33±9	31±17	41±32	59±45	82±59	87±71
	BCY613							
7	6, 3 mpk, qw	171±21	132±19	108±19	85±15	81±8	87±14	92±18
	BCY608							
8	2, 2 mpk, qw	1034±17 8	1287±94	--	--	--	--	--
	BCY608							
9	2, 5 mpk, qw	214±120	218±146	--	--	--	--	--

Table 49: Tumor volume trace (PG-D35~PG-D47)

Gr.	Treatment	Days after the start of treatment					
		35	38	40	42	45	47
3	BCY6033, 2 mpk, qw	352±210	456±271	525±302	683±400	738±429	853±476
4	BCY6033, 3 mpk, qw	79±47	118±71	139±82	220±125	312±176	423±222
6	BCY6136,	124±106	156±120	179±142	239±197	285±239	350±298

2 mpk, qw							
7	BCY6136, 3 mpk, qw	129±38	173±65	181±65	269±113	293±114	371±128

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for BCY6033, BCY6136 and BCY6082 in the MDA-MB-231-luc xenograft model was calculated based on tumor volume measurements at day 21 after the start of treatment.

Table 50: Tumor growth inhibition analysis

Gr	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	P value
1	Vehicle, qw	1661±173	--	--	--
2	BCY6033, 1 mpk, qw	880±156	53.0	52.0	<i>p</i> <0.001
3	BCY6033, 2 mpk, qw	67±44	4.1	106.1	<i>p</i> <0.001
4	BCY6033, 3 mpk, qw	22±12	1.3	109.1	<i>p</i> <0.001
5	BCY6136, 1 mpk, qw	994±213	59.8	44.4	<i>p</i> <0.01
6	BCY6136, 2 mpk, qw	33±9	2.0	108.4	<i>p</i> <0.001
7	BCY6136, 3 mpk, qw	132±19	8.0	101.7	<i>p</i> <0.001
8	BCY6082, 2 mpk, qw	1287±94	77.5	24.9	<i>p</i> >0.05
9	BCY6082, 5 mpk, qw	218±146	13.1	96.1	<i>p</i> <0.001

a. Mean ± SEM.

b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

(e) Results Summary and Discussion

In this study, the therapeutic efficacy of BCY6033, BCY6136 and BCY6082 in the MDA-MB-231-luc xenograft model was evaluated. The measured body weights and tumor volumes of

all treatment groups at various time points are shown in the Figures 25 to 27 and Tables 47 to 50.

The mean tumor size of vehicle treated mice reached 1661 mm³ on day 21. BCY6033 at 1 mg/kg (TV=880 mm³, TGI=52.0%, $p<0.001$), 2 mg/kg (TV=67 mm³, TGI=106.1%, $p<0.001$) and 3 mg/kg (TV=22 mm³, TGI=109.1%, $p<0.001$) produced dose-dependent antitumor activity. BCY6033 at 2 mg/kg and 3 mg/kg regressed the tumors potently, but the tumors showed obvious re-growth from day 21.

BCY6136 at 1 mg/kg (TV=994 mm³, TGI=44.4%, $p<0.01$) showed moderate antitumor activity, BCY6136 at 2 mg/kg (TV=33 mm³, TGI=108.4%, $p<0.001$) and 3 mg/kg (TV=132 mm³, TGI=101.1%, $p<0.001$) produced potent antitumor activity, but the tumors showed obvious re-growth from day 28.

BCY6082 at 2 mg/kg (TV=1287 mm³, TGI=24.9%, $p>0.05$) didn't show obvious antitumor activity, BCY6082 at 5 mg/kg (TV=218 mm³, TGI=96.1%, $p<0.001$) produced significant antitumor activity.

In this study, one mouse treated with BCY6136 2 mg/kg lost over 15% bodyweight during the treatment schedule, other mice maintained the bodyweight well.

Study 18: In vivo efficacy test of BCY6136 in treatment of EMT-6 syngeneic model in BALB/c mice

(a) Study Objective

The objective of the research was to evaluate the *in vivo* anti-tumor efficacy of BCY6136 in treatment of EMT-6 syngeneic model in BALB/c mice.

(b) Experimental Design

Group	Treatment	Dose (mg/kg)	N	Dosing Route	Schedule	Sample Collection
1	Vehicle	--	5	iv	qw*4	tumors from spare
2	BCY6136	3	5	iv	qw*4	mice will be
3	BCY6136	1/5 ^b	5	iv	qw*4	collected for FACS
4	BCY6136	0.3/3 ^b	5	iv	qw*4	

a. The injection volume of each mouse is 10 ml/kg.

b. The dosage of group 3 and group 4 was changed to 5 mpk and 3 mpk from Day 14.

(c) Experimental Methods and Procedures

(i) Cell Culture

The EMT-6 tumor cells were maintained *in vitro* as a monolayer culture in EMEM medium supplemented with 10% heat inactivated fetal bovine serum at 37°C in an atmosphere of 5% CO₂ in air. The tumor cells were routinely subcultured twice weekly by trypsin-EDTA treatment.

The cells growing in an exponential growth phase were harvested and counted for tumor inoculation.

(ii) Tumor Inoculation

- Each mouse was inoculated subcutaneously at the right flank with EMT-6 tumor cells (5×10^6) in 0.1 ml of PBS for tumor development. 44 animals were randomized when the average tumor volume reached 75 mm^3 . The test article administration and the animal numbers in each group were shown in the experimental design table.

(iii) Testing Article Formulation Preparation

BCY6136 formulation		
Treatment	Conc.(mg/ml)	Formulation
Vehicle/buffer	--	50 mM Acetate, 10% sucrose pH=5
BCY6136	1	Dissolve 6.2 mg BCY6136 with 6113 μl buffer
BCY6136	0.3	Dilute 450 μl 1 mg/ml BCY6136 stock with 1050 μl buffer
BCY6136	0.1	Dilute 150 μl 1 mg/ml BCY6136 stock with 1350 μl buffer
BCY6136	0.03	Dilute 45 μl 1 mg/ml BCY6136 stock with 1455 μl buffer

BCY6136 formulation		
Treatment	Conc.(mg/ml)	Formulation
Vehicle/buffer	--	50 mM Acetate, 10% sucrose pH=5
BCY6136	1	stock
BCY6136	0.3	Dilute 420 μl 1 mg/ml BCY6136 stock with 980 μl buffer
BCY6136	0.3	Dilute 420 μl 1 mg/ml BCY6136 stock with 980 μl buffer
BCY6136	0.5	Dilute 700 μl 1 mg/ml BCY6136 stock with 700 μl buffer

10 (iv) Sample Collection

3 tumors from spare mice were collected for FACS on day 11. The data was supplied by biology team.

(d) Results

(i) Body Weight change and Tumor Growth Curve

- 15 Body weight and tumor growth curve are shown in Figure 28.

(ii) Tumor Volume Trace

Mean tumor volume over time in female BALB/c mice bearing EMT-6 syngeneic is shown in Table 51.

Table 51: Tumor volume trace over time

Gr	Treatment	Days after the start of treatment									
		0	3	5	7	10	12	14	17	19	21
1	Vehicle, qw	82±4	141±11	260±24	443±90	557±99	703±119	812±139	948±191	1129±248	1499±340
2	BCY6136, 3 mpk, qw	82±4	58±1	59±2	125±18	240±23	322±23	374±22	431±37	486±50	561±61
3	BCY6136, 1/5 ^a mpk, qw	82±4	108±18	204±27	350±57	426±49	588±72	691±65	850±98	1018±115	1272±140
4	BCY6136, 0.3/3 ^a mpk, qw	82±4	130±16	255±35	358±34	450±67	607±94	731±112	872±119	1082±133	1394±161

5 The dosage of group 3 and group 4 was changed to 5 mpk and 3 mpk from Day 14.

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for BCY6136 in EMT-6 syngeneic model was calculated based on tumor volume measurements on day 21 after the start of treatment.

Table 52: Tumor growth inhibition analysis

Gr	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	P value compare with vehicle
1	Vehicle, qw	1499±340	--	--	--
2	BCY6136, 3 mpk, qw	561±61	37.4	66.2	<i>p</i> <0.05
3	BCY6136, 1/5 ^c mpk, qw	1272±140	84.8	16.1	ns
4	BCY6136, 0.3/3 ^c mpk, qw	1394±161	93.0	7.4	ns

10 a. Mean ± SEM.

b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

c. The dosage of group 3 and group 4 was changed to 5 mpk and 3 mpk from Day 14.

(e) Results Summary and Discussion

In this study, the therapeutic efficacy of BCY6136 in EMT-6 syngeneic model was evaluated. The measured body weights and tumor volumes of all treatment groups at various time points are shown in the Figure 28 and Tables 51 and 52.

The mean tumor size of vehicle treated mice reached 1499 mm³ on day 21. BCY6136 at 3 mg/kg, qw (TV=561 mm³, TGI=66.2%, $p<0.05$) showed obvious antitumor activity. BCY6136 at 1/5 mg/kg, qw (TV=1272 mm³, TGI=16.1%, $p>0.05$) and BCY6136 at 0.3/3 mg/kg, qw (TV=1394 mm³, TGI=7.4%, $p>0.05$) didn't show any antitumor activity.

The dosage of group 3 and group 4 was changed to 5 mpk and 3 mpk from day 14. Tumor ulceration was found in mouse 3-5 on Day 14, and the mice was deal with antibiotic cream. In this study, all mice maintained the bodyweight well.

Study 19: In vivo efficacy study of BCY6136 in treatment of NCI-N87 xenograft in Balb/c nude mice

(a) Study Objective

The objective of the research is to evaluate the *in vivo* anti-tumor efficacy of BCY6136 in treatment of NCI-N87 xenograft in Balb/c nude mice.

(b) Experimental Design

Group	Treatment	n	Dose (mg/kg)	Dosing Volume (μl/g)	Dosing Route	Schedule
1	Vehicle	3	--	10	<i>iv</i>	Qw
2	BCY6136	3	1	10	<i>iv</i>	Qw
3	BCY6136	3	2	10	<i>iv</i>	Qw
4	BCY6136	3	3	10	<i>iv</i>	Qw

(c) Experimental Methods and Procedures

(i) Cell Culture

The NCI-N87 tumor cells were maintained in RPMI-1640 medium supplemented with 10% heat inactivated fetal bovine serum at 37°C in an atmosphere of 5% CO₂ in air. The tumor cells were routinely subcultured twice weekly. The cells growing in an exponential growth phase were harvested and counted for tumor inoculation.

(ii) Tumor Inoculation

Each mouse was inoculated subcutaneously at the right flank with NCI-N87 tumor cells (10 x 10⁶) with matrigel (1:1) in 0.2 ml of PBS for tumor development. The animals were randomized and treatment was started when the average tumor volume reached approximately 176 mm³. The test article administration and the animal number in each group are shown in the experimental design table.

(iii) Testing Article Formulation Preparation

Test article	Conc. (mg/ml)	Formulation
Vehicle	-	50 mM Acetate 10% sucrose pH 5
BCY6136	1	Dissolve 4.295 mg BCY6136 in 4.214 ml Acetate buffer ¹
	0.1	Dilute 90 µl 1 mg/ml BCY6136 stock with 810 µl Acetate buffer
	0.2	Dilute 180 µl 1 mg/ml BCY6136 stock with 720 µl Acetate buffer
	0.3	Dilute 270 µl 1 mg/ml BCY6136 stock with 630 µl Acetate buffer
1. Acetate buffer: 50 mM Acetate 10% sucrose pH 5		

(d) Results**(i) Body Weight change and Tumor Growth Curve**

Body weight and tumor growth curve is shown in Figure 29.

(ii) Tumor Volume Trace

- 5 Mean tumor volume over time in female Balb/c nude mice bearing NCI-N87 xenograft is shown in Table 53.

Table 53: Tumor volume trace over time

Gr.	Treatme nt	Days after the start of treatment														
		0	2	4	7	9	11	14	16	18	21	23	25	28	30	
1	Vehicle,	17	21	26	42	53	59	73	82	91						
	qw	4± 7	3± 5	6± 6	1± 10	7± 17	8± 30	4± 46	1± 55	8± 91	1024 ±83	1151 ±68	1305 ±57	1407 ±64	1465 ±90	
	BCY613	17	20	21	22	23	18	24	27	30						
2	6,	6± 7	0± 8	0± 14	4± 27	8± 21	4± 18	4± 23	6± 35	8± 44	343 ±37	390 ±43	406 ±48	422 ±42	425 ±47	
	1 mpk,															
	qw															
3	BCY613	17	19	16	17	16	96	13	15	16						
	6,	6± 18	7± 25	8± 25	0± 26	5± 34	±2 7	3± 35	0± 52	0± 49	190 ±63	203 ±65	218 ±66	201 ±53	210 ±60	
	2 mpk,															
4	qw															
	BCY613	17	19	16	15	14	95	14	14	16						
	6,	7± 8	7± 9	9± 7	8± 3	8± 8	±1 6	1± 12	5± 24	4± 28	202 ±28	205 ±30	201 ±16	196 ±21	201 ±22	
	3 mpk,															
	qw															

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for BCY6136 in the NCI-N87 xenograft was calculated based on tumor volume measurements at day 30 after the start of treatment.

Table 54: Tumor growth inhibition analysis

Group	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	P value
1	Vehicle, qw	1465±90	--	--	--
2	BCY6136, 1 mpk, qw	425±47	29.0	80.7	$p < 0.001$
3	BCY6136, 2 mpk, qw	210±60	14.3	97.4	$p < 0.001$
4	BCY6136, 3 mpk, qw	201±22	13.7	98.1	$p < 0.001$

a. Mean ± SEM.

b. Tumor growth inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

5 (e) **Results Summary and Discussion**

In this study, the therapeutic efficacy of BCY6136 in the NCI-N87 model was evaluated. The measured body weight and tumor volume of all treatment groups at various time points are shown in the Figure 29 and Tables 53 and 54.

10 The mean tumor size of vehicle treated mice reached 1465 mm³ on day 30. BCY6136 at 1 mg/kg, qw (TV=425 mm³, TGI=80.7%, $p < 0.001$) and 2 mg/kg, qw (TV=210 mm³, TGI=97.4%, $p < 0.001$) produced significant antitumor activity in a dose-dependent manner, BCY6136 at 3 mg/kg, qw (TV=201 mm³, TGI=98.1%, $p < 0.001$) showed comparable antitumor activity with BCY6136 at 2 mpk.

15 In this study, no obvious body weight loss was found in all the groups during the treatment schedule.

Study 20: In vivo efficacy study of BCY6136 in treatment of SK-OV-3 xenograft in Balb/c nude mice

(a) Study Objective

20 The objective of the research is to evaluate the *in vivo* anti-tumor efficacy of BCY6136 in treatment of SK-OV-3 xenograft in Balb/c nude mice.

(b) Experimental Design

Group	Treatment	n	Dose (mg/kg)	Dosing Volume (ul/g)	Dosing Route	Schedule
1	Vehicle	3	--	10	<i>iv</i>	Qw
2	ADC	3	3	10	<i>iv</i>	Qw
3	BCY6136	3	1	10	<i>iv</i>	Qw
4	BCY6136	3	2	10	<i>iv</i>	Qw
5	BCY6136	3	3	10	<i>iv</i>	Qw

(c) Experimental Methods and Procedures**(i) Cell Culture**

The SK-OV-3 tumor cells were maintained in McCoy's 5a medium supplemented with 10% heat inactivated fetal bovine serum at 37°C in an atmosphere of 5% CO₂ in air. The tumor cells were routinely subcultured twice weekly. The cells growing in an exponential growth phase were harvested and counted for tumor inoculation.

(ii) Tumor Inoculation

Each mouse was inoculated subcutaneously at the right flank with SK-OV-3 tumor cells (10 x 10⁶) with matrigel (1:1) in 0.2 ml of PBS for tumor development. The animals were randomized and treatment was started when the average tumor volume reached approximately 186 mm³. The test article administration and the animal number in each group are shown in the experimental design table.

(iii) Testing Article Formulation Preparation

Test article	Purity	Conc. (mg/ml)	Formulation
Vehicle	-	-	50 mM Acetate 10% sucrose pH5
BCY6136	98.5%	1	Dissolve 3.65 mg BCY6136 in 3.60 ml 50 mM Acetate buffer ¹
		0.1	Dilute 90 µl 1 mg/ml BCY6136 stock with 810 µl Acetate buffer ¹
		0.2	Dilute 180 µl 1 mg/ml BCY6136 stock with 720 µl Acetate buffer ¹
		0.3	Dilute 270 µl 1 mg/ml BCY6136 stock with 630 µl Acetate buffer ¹
ADC	ADC	0.3	Dilute 69 µl 10.47 mg/ml ADC stock with 2331 µl ADC buffer ²
1. Acetate buffer: 50 mM Acetate 10% sucrose pH5 2. ADC buffer: 25 mM Histidine 10% sucrose pH5.5			

(d) Results**(i) Body Weight change and Tumor Growth Curve**

Body weight and tumor growth curve is shown in Figure 30.

(ii) Tumor Volume Trace

Mean tumor volume over time in female Balb/c nude mice bearing SK-OV-3 xenograft is shown in Table 55.

Table 55: Tumor volume trace over time

Gr.	Treatment	Days after the start of treatment													
		0	2	5	7	9	12	14	16	19	21	23	26	28	
1	Vehicle, qw	18	24	31	39	47	60	742	891	1076	1173	1340	1490	1560	
		7±	3±	3±	9±	0±	6±	±10	±133	±185	±214	±236	±273	±305	
		16	24	28	37	23	61	3							
2	ADC, 3 mpk, qw	18	18	21	26	26	33	353	392	449	481	573	647	684	
		7±	1±	2±	3±	8±	5±	±18	±63	±4	±27	±33	±26	±111	
		16	15	16	35	14	23								
3	BCY6136, 2 mpk, qw	18	22	29	33	35	44	503	587	702	752	893	1002	1035	
		6±	2±	3±	1±	6±	0±	±28	±33	±43	±26	±34	±68	±67	
		23	19	34	21	23	8								
4	BCY6136, 2 mpk, qw	18	17	16	18	18	20	200	230	229	231	236	240	277	
		6±	0±	4±	8±	0±	2±	±29	±46	±48	±58	±49	±48	±58	
		23	18	28	33	34	29								
5	BCY6136, 3 mpk, qw	18	16	15	16	15	18	187	212	208	204	205	227	254	
		4±	8±	0±	4±	8±	0±	±4	±17	±29	±12	±17	±31	±48	
		24	18	12	12	8	8								

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for BCY6136 in the SK-OV-3 xenograft was calculated based on tumor volume measurements at day 28 after the start of treatment.

5 Table 56: Tumor growth inhibition analysis

Group	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	P value
1	Vehicle, qw	1560±305	--	--	--
2	ADC, 3 mpk, qw	684±111	43.9	63.8	<i>p</i> <0.01
3	BCY6136, 1 mpk, qw	1035±67	66.4	38.1	<i>p</i> >0.05
4	BCY6136, 2 mpk, qw	277±58	17.8	93.3	<i>p</i> <0.001
5	BCY6136, 3 mpk, qw	254±48	16.3	95.0	<i>p</i> <0.001

a. Mean ± SEM.

b. Tumor growth inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

(e) Results Summary and Discussion

5 In this study, the therapeutic efficacy of BCY6136 in the SK-OV-3 model was evaluated. The measured body weight and tumor volume of all treatment groups at various time points are shown in the Figure 30 and Tables 55 and 56.

The mean tumor size of vehicle treated mice reached 1560 mm³ on day 28. ADC at 3 mg/kg, qw (TV=684 mm³, TGI=63.8%, *p*<0.01) showed moderate anti-tumor efficacy. BCY6136 at 1 mg/kg, qw (TV=1035 mm³, TGI=38.1%, *p*>0.05) didn't show obvious anti-tumor activity. BCY6136 at 2 mg/kg, qw (TV=277 mm³, TGI=93.3%, *p*<0.001) and 3 mg/kg, qw (TV=254 mm³, TGI=95.0%, *p*<0.001) produced significant anti-tumor activity.

In this study, no obvious body weight loss was found in all the groups during the treatment schedule.

Study 21: In vivo efficacy study of BCY6136 in treatment of OE21 xenograft in**Balb/c nude mice****(a) Study Objective**

The objective of the research is to evaluate the *in vivo* anti-tumor efficacy of BCY6136 in treatment of OE21 xenograft in Balb/c nude mice.

(b) Experimental Design

Group	Treatment	n	Dose (mg/kg)	Dosing Volume (μl/g)	Dosing Route	Schedule
1	Vehicle	3	--	10	<i>iv</i>	qw
2	BCY6136	3	1	10	<i>iv</i>	qw
3	BCY6136	3	2	10	<i>iv</i>	qw
4	BCY6136	3	3	10	<i>iv</i>	qw

(c) Experimental Methods and Procedures**(i) Cell Culture**

The OE21 tumor cells were maintained in RPMI-1640 medium supplemented with 10% heat inactivated fetal bovine serum at 37°C in an atmosphere of 5% CO₂ in air. The tumor cells were routinely subcultured twice weekly. The cells growing in an exponential growth phase were harvested and counted for tumor inoculation.

(ii) Tumor Inoculation

Each mouse was inoculated subcutaneously at the right flank with OE21 tumor cells (5×10^6) with matrigel (1:1) in 0.2 ml of PBS for tumor development. The animals were randomized and treatment was started when the average tumor volume reached approximately 157 mm³. The test article administration and the animal number in each group are shown in the experimental design table.

(iii) Testing Article Formulation Preparation

Test article	Conc. (mg/ml)	Formulation
Vehicle	-	50 mM Acetate 10% sucrose pH 5
BCY6136	1	Dissolve 4.295 mg BCY6136 in 4.214 ml Acetate buffer ¹
	0.1	Dilute 90 μl 1 mg/ml BCY6136 stock with 810 μl Acetate buffer
	0.2	Dilute 180 μl 1 mg/ml BCY6136 stock with 720 μl Acetate buffer
	0.3	Dilute 270 μl 1 mg/ml BCY6136 stock with 630 μl Acetate buffer
1. Acetate buffer: 50 mM Acetate 10% sucrose pH 5		

(d) Results**(i) Body Weight change and Tumor Growth Curve**

Body weight and tumor growth curve is shown in Figure 31.

(ii) Tumor Volume Trace

- 5 Mean tumor volume over time in female Balb/c nude mice bearing OE21 xenograft is shown in Table 57.

Table 57: Tumor volume trace over time

Gr.	Treatment	Days after the start of treatment											
		0	2	4	7	9	11	14	16	18	21	23	
1	Vehicle, qw	155±9	211±16	291±16	379±14	456±32	539±13	828±42	955±40	1035±58	1250±46	1586±57	
2	BCY6136, 1 mpk, qw	159±14	202±28	251±29	282±6	331±19	392±35	609±56	694±44	777±68	1083±85	1155±98	
3	BCY6136, 2 mpk, qw	157±19	197±13	219±6	235±27	268±35	243±37	346±78	371±98	396±109	515±94	537±122	
4	BCY6136, 3 mpk, qw	155±19	200±16	197±7	209±11	229±26	211±14	289±38	318±53	330±40	474±42	489±51	

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for BCY6136 in the OE21 xenograft was calculated based on tumor volume measurements at day 23 after the start of treatment.

Table 58: Tumor growth inhibition analysis

Group	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	P value
1	Vehicle, qw	1586±57	--	--	--
2	BCY6136, 1 mpk, qw	1155±98	72.8	30.4	<i>p</i> <0.05
3	BCY6136, 2 mpk, qw	537±122	33.9	73.4	<i>p</i> <0.001
4	BCY6136, 3 mpk, qw	489±51	30.8	76.7	<i>p</i> <0.001

a. Mean $\pm$ SEM.

b. Tumor growth inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

(e) Results Summary and Discussion

- 5 In this study, the therapeutic efficacy of BCY6136 in the OE21 model was evaluated. The measured body weight and tumor volume of all treatment groups at various time points are shown in the Figure 31 and Tables 57 and 58.

The mean tumor size of vehicle treated mice reached 1586 mm³ on day 23. BCY6136 at 1 mg/kg, qw (TV= 1155 mm³, TGI = 30.4% $p < 0.05$) showed slight anti-tumor activity. BCY6136
10 at 2 mg/kg, qw (TV=537 mm³, TGI=73.4%, $p < 0.001$) and 3 mg/kg, qw (TV=489 mm³, TGI=76.7%, $p < 0.001$) produced significant anti-tumor activity.

In this study, no obvious body weight loss was found in all the groups during the treatment schedule.

15 **Study 22: In vivo efficacy test of BCY6136 and BCY6082 in treatment of MOLP-8 xenograft in CB17-SCID mice**

(a) Study Objective

The objective of the research is to evaluate the *in vivo* anti-tumor efficacy of BCY6136 and BCY6082 in treatment of MOLP-8 xenograft in CB17-SCID mice.

20 **(b) Experimental Design**

Group	Treatment	n	Dose (mg/kg)	Dosing Volume (μl/g)	Dosing Route	Schedule
1	Vehicle	3	--	10	<i>iv</i>	qw
2	BCY6136	3	1	10	<i>iv</i>	qw
3	BCY6136	3	2	10	<i>iv</i>	qw
4	BCY6136	3	3	10	<i>iv</i>	qw
5	BCY6082	3	1	10	<i>iv</i>	qw
6	BCY6082	3	2	10	<i>iv</i>	qw
7	BCY6082	3	3	10	<i>iv</i>	qw

(c) Experimental Methods and Procedures

(i) Cell Culture

- The MOLP-8 tumor cells were maintained in vitro as a monolayer culture in RMPI-1640 supplemented with 20% heat inactivated fetal bovine serum at 37°C in an atmosphere of 5%
25 CO₂ in air. The tumor cells were routinely subcultured by trypsin-EDTA treatment. The cells growing in an exponential growth phase were harvested and counted for tumor inoculation.

(ii) Tumor Inoculation

Each mouse was inoculated subcutaneously at the right flank with MOLP-8 tumor cells (10×10^6) in 0.2 ml PBS with 50% matrigel for tumor development. 36 animals were randomized when the average tumor volume reached 141 mm^3 . The test article administration and the

5 animal numbers in each group were shown in the experimental design table.

(iii) Testing Article Formulation Preparation

Treatment	Concentration (mg/ml)	Formulation
Vehicle	--	50 mM Acetate, 10% sucrose pH=5
BCY6136	0.1	Dilute 90 μl 1 mg/ml BCY6136 stocks* with 810 μl buffer***
	0.2	Dilute 180 μl 1 mg/ml BCY6136 stocks* with 720 μl buffer***
	0.3	Dilute 270 μl 1 mg/ml BCY6136 stocks* with 630 μl buffer***
BCY6082	0.1	Dilute 90 μl 1 mg/ml BCY6082 stocks** with 810 μl buffer***
	0.2	Dilute 180 μl 1 mg/ml BCY6082 stocks** with 720 μl buffer***
	0.3	Dilute 270 μl 1 mg/ml BCY6082 stocks** with 630 μl buffer***
<p>* BCY6136 stocks: 10.93 mg BCY6136 dissolved to 10.93 mL 50 mM Acetate, 10% sucrose ,pH=5, and separated into individual tubes and stored at -80°C.</p> <p>** BCY6082 stocks: 2.43 mg BCY6136 dissolved to 2.43 mL 50 mM Acetate, 10% sucrose ,pH=5, and separated into individual tubes and stored at -80°C.</p> <p>***Buffer: 50 mM Acetate, 10% sucrose pH=5</p>		

(d) **Results**(i) Body Weight change and Tumor Growth Curve

Body weight and tumor growth curve are shown in Figures 32 and 33.

10 (ii) Tumor Volume Trace

Mean tumor volume over time in female CB17-SCID mice bearing MOLP-8 xenograft is shown in Table 59.

Table 59: Tumor volume trace over time

Gr	Treatment	Days after the start of treatment						
		0	2	4	7	9	11	14
.								

1	Vehicle, qw	139±2	375±3 6	604±2 8	984±8 8	1451±1 33	1981±1 96	2528±2 95
2	BCY6136, 1 mpk, qw	143±1 3	299±6	444±4 9	576±3 1	806±85	1132±1 70	1446±2 34
3	BCY6136, 2 mpk, qw	140±1 5	271±4 3	250±2	509±2 3	662±78	873±49	1218±1 44
4	BCY6136, 3 mpk, qw	142±1 9	239±6 7	197±2 0	342±7 8	425±90	693±13 3	938±15 5
5	BCY6082, 1 mpk, qw	142±4	303±4 9	456±8 3	809±1 69	1365±2 77	1708±1 90	2296±5 11
6	BCY6082, 2 mpk, qw	139±5	273±4 6	428±1 8	682±5 0	945±73	1240±8 5	1554±8 4
7	BCY6082, 3 mpk, qw	142±4	219±7	369±7 7	471±8 1	656±11 5	997±21 2	1321±3 36

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for BCY6136 and BCY6082 in the MOLP-8 xenograft model was calculated based on tumor volume measurements at day 14 after the start of treatment.

Table 60: Tumor growth inhibition analysis

Gr	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	<i>P</i> value compared with vehicle
1	Vehicle, qw	2528±295	--	--	--
2	BCY6136, 1 mpk, qw	1446±234	57.2	45.5	<i>p</i> >0.05
3	BCY6136, 2 mpk, qw	1218±144	48.2	54.9	<i>p</i> <0.05
4	BCY6136, 3 mpk, qw	938±155	37.1	66.7	<i>p</i> <0.01
5	BCY6082, 1 mpk, qw	2296±511	90.8	9.8	<i>p</i> >0.05
6	BCY6082, 2 mpk, qw	1554±84	61.5	40.8	<i>p</i> >0.05

7	BCY6082, 3 mpk, qw	1321±336	52.3	50.6	$p<0.05$
---	-----------------------	----------	------	------	----------

a. Mean $\pm$ SEM.

b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

(e) Results Summary and Discussion

- 5 In this study, the therapeutic efficacy of BCY6136 and BCY6082 in the MOLP-8 xenograft model was evaluated. The measured body weights and tumor volumes of all treatment groups at various time points are shown in the Figures 32 and 33 and Tables 59 and 60.

The mean tumor size of vehicle treated mice reached 2528 mm³ on day 14. BCY6136 at 1 mg/kg (TV=1146 mm³, TGI=45.5%, $p>0.05$), 2 mg/kg (TV=1218 mm³, TGI=54.9%, $p<0.05$) and 3 mg/kg (TV=938 mm³, TGI=66.7%, $p<0.01$) produced dose-dependent antitumor activity, but all of dosage didn't regress the tumors in MOLP-8 xenografts.

- 10 BCY6082 at 1 mg/kg (TV=2296 mm³, TGI=9.8%, $p>0.05$) and 2 mg/kg (TV=1554 mm³, TGI=40.8%, $p>0.05$) didn't show obvious anti-tumor activity. BCY6082 at 3 mg/kg inhibited the tumor growth significantly (TV=1321 mm³, TGI=50.6%, $p<0.05$), but didn't regress the tumors in MOLP-8 xenografts.

15 In this study, all of mice maintained the bodyweight well.

Study 23: In vivo efficacy test of BCYs in treatment of HT1080 xenograft in BALB/c nude mice

- 20 **(a) Study Objective**

The objective of the research was to evaluate the *in vivo* anti-tumor efficacy of BCYs in treatment of HT1080 xenograft model in BALB/c nude mice.

(b) Experimental Design

Group	Treatment	n	Dose (mg/kg)	Dosing Volume (μl/g)	Dosing Route	Schedule
1	Vehicle	3	--	10	<i>iv</i>	qw
2	BCY6082	3	2	10	<i>iv</i>	qw
3	BCY6031	3	2	10	<i>iv</i>	qw
4	BCY6173	3	1	10	<i>iv</i>	qw
5	BCY6173	3	2	10	<i>iv</i>	qw
6	BCY6173	3	3	10	<i>iv</i>	qw
7	BCY6135	3	1	10	<i>iv</i>	qw
8	BCY6135	3	2	10	<i>iv</i>	qw
9	BCY6135	3	3	10	<i>iv</i>	qw
10	BCY6033	3	3	10	<i>iv</i>	qw

11	BCY6033	3	5	10	<i>iv</i>	qw
12	BCY6136	3	2	10	<i>iv</i>	qw
13	BCY6136	3	3	10	<i>iv</i>	qw
14	BCY6136	3	5	10	<i>iv</i>	qw
15	BCY6174	3	1	10	<i>iv</i>	qw
16	BCY6174	3	2	10	<i>iv</i>	qw
17	BCY6174	3	3	10	<i>iv</i>	qw
18	BCY6175	3	1	10	<i>iv</i>	qw
19	BCY6175	3	2	10	<i>iv</i>	qw
20	BCY6175	3	3	10	<i>iv</i>	qw
21	ADC	3	3	10	<i>iv</i>	qw

Note: n: animal number; Dosing volume: adjust dosing volume based on body weight 10 µl/g.

(c) Experimental Methods and Procedures

(i) Cell Culture

The HT1080 tumor cells will be maintained in medium supplemented with 10% heat inactivated fetal bovine serum at 37°C in an atmosphere of 5% CO₂ in air. The tumor cells will be routinely subcultured twice weekly. The cells growing in an exponential growth phase will be harvested and counted for tumor inoculation.

(ii) Tumor Inoculation

Each mouse will be inoculated subcutaneously at the right flank with HT1080 tumor cells (5*10⁶) for tumor development. The animals will be randomized and treatment will be started when the average tumor volume reaches approximately 150-200 mm³. The test article administration and the animal numbers in each group are shown in the following experimental design table.

(iii) Testing Article Formulation Preparation

Treatment	Dose (mg/ml)	Formulation
Vehicle	--	50 mM Acetate/acetic acid pH5 10% sucrose
BCY6082	0.2	Dilute 160 µl 1 mg/ml BCY6082 stock with 640 µl buffer
BCY6031	0.2	Dilute 180 µl 1 mg/ml BCY6031 stock with 720 µl buffer
BCY6173	1	Dissolve 2.13 mg BCY6173 with 2.04 ml buffer
	0.1	Dilute 90 µl 1 mg/ml BCY6173 stock with 810 µl buffer
	0.2	Dilute 180 µl 1 mg/ml BCY6173 stock with 720 µl buffer
	0.3	Dilute 270 µl 1 mg/ml BCY6173 stock with 630 µl buffer
BCY6135	1	Dissolve 2 mg BCY6135 with 1.9 ml buffer
	0.1	Dilute 90 µl 1 mg/ml BCY6135 stock with 810 µl buffer
	0.2	Dilute 180 µl 1 mg/ml BCY6135 stock with 720 µl buffer

	0.3	Dilute 270 μ l 1 mg/ml BCY6135 stock with 630 μ l buffer
BCY6033	0.3	Dilute 270 μ l 1 mg/ml BCY6033 stock with 630 μ l buffer
	0.5	Dilute 450 μ l 1 mg/ml BCY6033 stock with 450 μ l buffer
BCY6136	0.2	Dilute 200 μ l 1 mg/ml BCY6136 stock with 800 μ l buffer
	0.3	Dilute 300 μ l 1 mg/ml BCY6136 stock with 700 μ l buffer
	0.5	Dilute 500 μ l 1 mg/ml BCY6136 stock with 500 μ l buffer
BCY6174	1	Dissolve 2.69 mg BCY6174 with 2.677 ml buffer
	0.1	Dilute 90 μ l 1 mg/ml BCY6174 stock with 810 μ l buffer
	0.2	Dilute 180 μ l 1 mg/ml BCY6174 stock with 720 μ l buffer
	0.3	Dilute 270 μ l 1 mg/ml BCY6174 stock with 630 μ l buffer
BCY6175	1	Dissolve 2 mg BCY6175 with 1.924 ml buffer
	0.1	Dilute 90 μ l 1 mg/ml BCY6175 stock with 810 μ l buffer
	0.2	Dilute 180 μ l 1 mg/ml BCY6175 stock with 720 μ l buffer
	0.3	Dilute 270 μ l 1 mg/ml BCY6175 stock with 630 μ l buffer
ADC	0.3	Dilute 25.78 μ l 10.47 mg/ml ADC stock with 874.22 μ l 25 mM Histidine pH 7 10% sucrose

(d) Results**(i) Body Weight change and Tumor Growth Curve**

Body weight and tumor growth curve are shown in Figures 34 to 42.

(ii) Tumor Volume Trace

- 5 Mean tumor volume over time in female Balb/c nude mice bearing HT1080 xenograft is shown in Table 61.

Table 61: Tumor volume trace over time

Gr	Treatment	Days after the start of treatment						
		0	2	4	7	9	11	14
1	Vehicle,	179 $\pm$ 2	312 $\pm$ 8	529 $\pm$ 13	886 $\pm$ 20	1185 $\pm$ 17	1467 $\pm$ 22	1737 $\pm$ 25
	qw	2	4	5	7	2	4	8
BCY6082								
2	2 mpk,	177 $\pm$ 1	183 $\pm$ 3					
	qw	6	1	99 $\pm$ 27	61 $\pm$ 17	33 $\pm$ 10	12 $\pm$ 5	5 $\pm$ 3
3	BCY6031	177 $\pm$ 2	215 $\pm$ 3					
	,	4	5	133 $\pm$ 37	63 $\pm$ 31	53 $\pm$ 37	45 $\pm$ 36	71 $\pm$ 67

2 mpk, qw								
BCY6173								
4	1 mpk, qw	178±2						1074±15
		6	276±8	328±73	594±62	745±22	960±53	0
BCY6173								
5	2 mpk, qw	178±2	277±6	262±12	309±23			
		8	1	5	8	425±334	436±323	480±347
BCY6173								
6	3 mpk, qw	179±4	182±7					
		3	1	133±88	87±68	77±65	60±54	47±42
BCY6135								
7	1 mpk, qw	178±2	267±6					
		2	6	262±58	436±67	599±89	703±36	871±28
BCY6135								
8	2 mpk, qw	178±2	176±4					
		3	8	117±43	70±23	67±23	52±21	62±7
BCY6135								
9	3 mpk, qw	177±3	178±7					
		9	9	92±67	62±46	62±51	57±51	44±40
BCY6033								
10	3 mpk, qw	178±2	186±3					
		6	4	79±30	29±15	12±8	6±4	9±7
BCY6033								
11	5 mpk, qw	178±3	117±2					
		6	0	41±10	12±4	6±2	4±0	0±0
BCY6136								
12	2mpk, qw	178±1	249±2					
		9	2	115±8	126±53	158±71	140±89	245±116
BCY6136								
13	3 mpk, qw	178±3	168±2					
		6	1	72±18	22±7	21±15	8±6	3±2
BCY6136								
14	5 mpk, qw	178±2	165±3					
		6	3	52±10	18±7	9±4	5±2	2±1

BCY6174								
15	1 mpk, qw	180±3 5	231±1 9	226±29	432±37	602±63	742±62	1066±13 0
BCY6174								
16	2mpk, qw	178±3 1	203±5 0	123±29	216±47	291±40	326±68	532±91
BCY6174								
17	3 mpk, qw	178±3 3	195±1 3	110±39	58±23	34±17	21±11	11±7
BCY6175								
18	1 mpk, qw	178±2 7	248±6 2	244±74	347±18	435±18	558±38	769±26
BCY6175								
19	2 mpk, qw	178±2 2	223±4 2	158±59	116±35	156±52	166±51	295±88
BCY6175								
20	3 mpk, qw	179±3 9	189±4 8	116±50	43±18	33±18	25±13	11±9
ADC								
21	3 mpk, qw	180±2 6	158±3 0	58±8	18±2	7±1	2±2	0±0

(iii) Tumor Growth Inhibition Analysis

Tumor growth inhibition rate for BCYs in the HT1080 xenograft model was calculated based on tumor volume measurements at day 14 after the start of treatment.

Table 62: Tumor growth inhibition analysis

Gr	Treatment	Tumor Volume (mm ³) ^a	T/C ^b (%)	TGI (%)	<i>P</i> value compare
1	Vehicle, qw	1737±258	--	--	--
2	BCY6082, 2 mpk,	5±3	0.3	111.1	<i>p</i> <0.01
3	BCY6031, 2 mpk,	71±67	4.1	106.8	<i>p</i> <0.01
4	BCY6173, 1 mpk,	1074±150	61.8	42.5	<i>p</i> >0.05
5	BCY6173, 2 mpk,	480±347	27.6	80.6	<i>p</i> <0.05
6	BCY6173, 3 mpk,	47±42	2.7	108.4	<i>p</i> <0.01
7	BCY6135, 1 mpk,	871±28	50.1	55.5	<i>p</i> <0.01
8	BCY6135, 2 mpk,	62±7	3.5	107.5	<i>p</i> <0.001

9	BCY6135, 3 mpk,	44±40	2.5	108.6	$p<0.001$
10	BCY6033, 3 mpk,	9±7	0.5	110.8	$p<0.001$
11	BCY6033, 5 mpk,	0±0	0.0	111.4	$p<0.001$
12	BCY6136, 2mpk, qw	245±116	14.1	95.7	$p<0.001$
13	BCY6136, 3 mpk,	3±2	0.2	111.2	$p<0.001$
14	BCY6136, 5 mpk,	2±1	0.1	111.3	$p<0.001$
15	BCY6174, 1 mpk,	1066±130	61.4	43.1	$p<0.05$
16	BCY6174, 2mpk, qw	532±91	30.6	77.3	$p<0.01$
17	BCY6174, 3 mpk,	11±7	0.6	110.7	$p<0.001$
18	BCY6175, 1 mpk,	769±26	44.3	62.1	$p<0.01$
19	BCY6175, 2 mpk,	295±88	17.0	92.5	$p<0.001$
20	BCY6175, 3 mpk,	11±9	0.6	110.8	$p<0.001$
21	ADC, 3 mpk, qw	0±0	0.0	111.5	-

a. Mean ± SEM.

b. Tumor Growth Inhibition is calculated by dividing the group average tumor volume for the treated group by the group average tumor volume for the control group (T/C).

(e) Results Summary and Discussion

- 5 In this study, the therapeutic efficacy of BCYs in the HT1080 xenograft model was evaluated. The measured body weights and tumor volumes of all treatment groups at various time points are shown in the Figures 34 to 42 and Tables 61 and 62.

The mean tumor size of vehicle treated mice reached 1737 mm³ on day 14.

- 10 BCY6082 at 2 mg/kg, qw (TV=5 mm³, TGI=111.1%, $p<0.01$) and BCY6031 at 2 mg/kg qw (TV=7 mm³, TGI=106.8%, $p<0.01$) showed potent anti-tumor activity.

BCY6173 at 1 mg/kg, qw (TV=1074 mm³, TGI=42.5%, $p>0.05$), 2 mg/kg, qw (TV=480 mm³, TGI=80.6%, $p<0.05$) and 3 mg/kg, qw (TV=7 mm³, TGI=108.4%, $p<0.01$) produced dose-dependent antitumor activity.

- 15 BCY6135 at 1 mg/kg, qw (TV=871 mm³, TGI=55.5%, $p<0.01$), 2 mg/kg, qw (TV=62 mm³, TGI=107.5%, $p<0.001$) and 3 mg/kg, qw (TV=44 mm³, TGI=108.6%, $p<0.001$) produced dose-dependent antitumor activity.

BCY6033 at 3 mg/kg, qw (TV=9 mm³, TGI=110.8%, $p<0.001$) and 5 mg/kg, qw (TV=0 mm³, TGI=111.4%, $p<0.001$) showed potent anti-tumor activity, and completely eradicated the tumors by day 14 at 5 mg/kg.

BCY6136 at 2 mg/kg, qw (TV=345 mm³, TGI=95.7%, $p<0.001$), 3 mg/kg, qw (TV=3 mm³, TGI=111.2%, $p<0.001$) and 5 mg/kg, qw (TV=2 mm³, TGI=111.3%, $p<0.001$) showed potent anti-tumor activity.

BCY6174 at 1 mg/kg, qw (TV=1066 mm³, TGI=43.1%, $p<0.05$), 2 mg/kg, qw (TV=532 mm³, TGI=77.3%, $p<0.01$) and 3 mg/kg, qw (TV=11 mm³, TGI=110.7%, $p<0.001$) produced dose-dependent antitumor activity.

BCY6175 at 1 mg/kg, qw (TV=769 mm³, TGI=62.1%, $p<0.01$), 2 mg/kg, qw (TV=295 mm³, TGI=92.5%, $p<0.001$) and 3 mg/kg, qw (TV=11 mm³, TGI=110.8%, $p<0.001$) produced dose-dependent antitumor activity.

ADC at 3 mg/kg, qw (TV=0 mm³, TGI=111.5%) completely eradicated the tumors by day 14.

Study 24: Investigation of Association between Copy Number Variation (CNV) and gene expression for EphA2 from multiple tumour types

Methods

1. Select all studies in cBioPortal (<http://www.cbioportal.org/>) and search for EPHA2.
 - (a) Remove provisional studies.
 - (b) Deselect studies with overlapping samples to prevent sample bias (based on warning in cBioPortal)- always keep PanCancer study if this is an option.
 - (c) Studies selected for analysis (Table 63).

20

Table 63: Studies analysed from cBioPortal and units in study

Study Name	Units
Breast Invasive Carcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Lung Squamous Cell Carcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Kidney Renal Papillary Cell Carcinoma (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Kidney Renal Clear Cell Carcinoma (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Colon Adenocarcinoma (TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Head and Neck Squamous Cell Carcinoma (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Bladder Urothelial Carcinoma (TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)

Uveal Melanoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Lung Adenocarcinoma (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Ovarian Serous Cystadenocarcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Breast Cancer (METABRIC, Nature 2012 & Nat Commun 2016)	mRNA expression (microarray)
Mesothelioma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Colorectal Adenocarcinoma (TCGA, Nature 2012)	RNA Seq RPKM
Cervical Squamous Cell Carcinoma (TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Sarcoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Cancer Cell Line Encyclopedia (Novartis/Broad, Nature 2012)	mRNA expression (microarray)
Rectum Adenocarcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Liver Hepatocellular Carcinoma (TCGA, PanCancer Atlas)	EPHA2: mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Stomach Adenocarcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Uterine Corpus Endometrial Carcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Skin Cutaneous Melanoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Prostate Adenocarcinoma (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Kidney Chromophobe (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Pediatric Wilms' Tumor (TARGET, 2018)	Epha2: mRNA expression (RNA-Seq RPKM)

Pheochromocytoma and Paraganglioma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Thyroid Carcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Esophageal Adenocarcinoma (TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Cholangiocarcinoma (TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Brain Lower Grade Glioma (TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Thymoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Pediatric Acute Lymphoid Leukemia - Phase II (TARGET, 2018)	Epha2: mRNA expression (RNA-Seq RPKM)
Diffuse Large B-Cell Lymphoma (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Glioblastoma Multiforme (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Metastatic Prostate Cancer, SU2C/PCF Dream Team (Robinson et al., Cell 2015)	mRNA expression / capture (RNA Seq RPKM)
Acute Myeloid Leukemia (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Testicular Germ Cell Tumors (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Adrenocortical Carcinoma (TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNASeqV2)
Uterine Carcinosarcoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Pancreatic Adenocarcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/Merged from Illumina HiSeq_RNASeqV2 syn4976369
Prostate Adenocarcinoma (MSKCC, Cancer Cell 2010)	mRNA Expression
Prostate Adenocarcinoma (Fred Hutchinson CRC, Nat Med 2016)	mRNA expression

2. Export CNV and RNA expression data from cBioPortal.
3. Test if CNVs are statistically significantly associated with changes in mRNA expression for EphA2 (log2 not applied).
- 5 (a) Run non-parametric Kruskal-Wallis test in GraphPad Prism (7.04) and R/R studio (threshold for significance: $p < 0.01$).
 - (i) GraphPad Prism: set up column table, run non-parametric test with no matching or pairing and do not assume Gaussian distribution.
 - (ii) Packages used in R:
 - 10 1. XLConnect
 2. dplyr
 3. Kruskal-Wallis Rank Sum Test: `Kruskal.test`.
4. Adjust for multiple comparisons (include all possible comparisons even if $n=1$ within a group) in R/Rstudio using Dunn's test (threshold for significance: $p < 0.025$).
- 15 (a) `dunn.test` with multiple comparison method= "bonferonni".

Results

- The results are shown in Table 64 below. Across 41 publicly available datasets compiled in cBioPortal that report both Copy Number Variation (CNV) and mRNA gene expression for
- 20 EphA2, there are numerous cancer types where cases have been reported with EphA2 shallow-deletions (< 2 copies). Although less common, in these same cancer types a subset of tumors harbored EphA2 deep deletions (> 1 copy loss or biallelic loss), EphA2 gains (2-3 copies) or EphA2 amplifications (> 3 copies). Indications where $> 33\%$ of tumors had either shallow-deletions or deep deletions in EphA2 included: kidney chromophobe,
- 25 cholangiocarcinoma, pheochromocytoma and paraganglioma, lung squamous cancer, breast, rectum, brain lower grade glioma, liver, adrenocortical carcinoma, mesothelioma, esophageal adenocarcinoma and colon cancer. In contrast, there were no studies where $> 33\%$ of samples had either gains or amplification in EphA2. Taken together these results demonstrate that deletions in EphA2 DNA are found across a variety of indications.
- 30 Approximately one third of all samples analyzed in the 41 studies harbored EphA2 CNVs. Based on this high percentage of CNVs across studies, and the high percentage of shallow deletions within specific tumor types, statistical testing was performed to identify possible associations between copy number changes and RNA expression. Tumors per indication
- 35 were allocated to 1 of 5 classes:
- a) Deep deletion;
 - b) Shallow deletion;

- c) Diploid;
- d) Gain; or
- e) Amplification.

5 Kruskal-Wallis testing was then performed to detect if the distributions of mRNA expression values per classes differed between classes ($P < 0.01$). For those TCGA data sets with $P < 0.01$ and to identify which classes were different to one another post-hoc testing was performed by calculating Z-statistics with adjusted P-values calculated (Bonferroni). For simplicity of interpretation pair-wise comparisons vs. diploid per indication were reviewed
10 (although all pair-wise P-values were calculated). 19/41 of these studies had a Kruskal-Wallis p-value of <0.01 demonstrating that copy number is statistically significantly associated with RNA expression. Of these 19 studies, 17 of them had a Bonferroni adjusted $P < 0.025$ for Diploid vs. Shallow Deletion indicating an association of decreased EphA2 mRNA expression with decreased EphA2 copy number. Only 2 of these 19 studies had a
15 Bonferroni adjusted $P < 0.025$ for Diploid vs. Gain and both were breast cancer studies. Furthermore, one of these breast cancer studies (Breast Invasive Carcinoma (TCGA, PanCancer Atlas)) had a Bonferroni adjusted $P < 0.025$ for both Diploid vs. Shallow Deletion and Diploid vs. Gain suggesting that copy number alterations may have a strong impact on EphA2 RNA expression in breast cancer.

20 The central dogma of genetics suggests that reduced copy number in EphA2 lead to reduced RNA and protein expression. Therefore, the observed associations between copy number loss of EphA2 and reduced mRNA expression in a variety of tumor types suggest that EphA2 protein expression may also be reduced. Similarly, copy number gains of EphA2
25 in breast cancer that were associated with increased mRNA expression may also suggest increased EphA2 protein expression. Moreover, higher EphA2 protein expression (measured by FACS) is associated with increased efficacy of certain EphA2 bicyclic drug conjugates of the invention (measured by tumor volume) in preclinical *in vivo* models. Taken together if copy number alterations that are associated with mRNA expression changes do predict
30 protein expression levels then patients with tumors containing copy number deletions of EphA2 may be less likely to respond to EphA2 bicyclic drug conjugates of the invention. Similarly, if patients with tumor copy number gains in EphA2 (e.g. breast cancer) it is possible that these patients would be more likely to respond to EphA2 bicyclic drug conjugates of the invention. Therefore, if patients were stratified by EphA2 copy number
35 status, then this information could be used to both exclude and select patients for treatment with EphA2 bicyclic drug conjugates of the invention to increase efficacy.

Table 64: Results of Investigation of Association between Copy Number Variation (CNV) and gene expression for EphA2

		Number of samples/group (n=X)					Kruskal-wallis test	Pairwise comparison, Z statistic (adjusted p-value), Bonferonni				
Study name	Units	Deep deletion	Shallow deletion	Diploid	Gain	Amplification	Kruskal-wallis statistic	p-value	Deep Deletion n	Diploid - Shallow deletion n	Diploid-Gain	Amplification - Diploid
Breast Invasive Carcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369	5	415	511	61	2	80.816	< 2.2e-16	0.17618 (1.0000)	6.460580 (0.0000)*	-4.603180 (0.0000)*	0.713978 (1.0000)
Lung Squamous Cell Carcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/ Merged from Illumina	3	207	201	55	0	52.942	1.89E-11	-1.584610 (0.3392)	6.786501 (0.0000)*	-0.019607 (1.0000)	N/A

[illegible]

Head and Neck Squamous Cell Carcinoma (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNAS eqV2)	3	86	345	54	0	32.72	3.69E- 07	- 2.4449 14 (0.0435))	4.6807 89 (0.0000))	- 1.5306 70 (0.3776))	N/A
Bladder Urothelial Carcinoma (TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNAS eqV2)	0	73	245	80	4	28.906	2.34E- 06	N/A	5.2032 51 (0.0000))	0.2117 44 (1.0000))	0.5817 04 (1.0000))
Uveal Melanoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369	0	24	56	0	0	21.051	4.47E- 06	N/A	4.5880 95 (0.0000))	N/A	N/A

Lung Adenocarcinoma (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNAS eqV2)	1	115	263	121	3	28.874	8.29E-06	- 0.6904 60 (1.0000))	4.2801 00 (0.0001))	- 0.6267 07 (1.0000))	2.2764 58 (0.1141))
Ovarian Serous Cystadenocarcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369	0	59	78	60	4	25.349	1.31E-05	N/A	4.3900 97 (0.0000))	- 0.2392 49 (1.0000))	0.2405 43 (1.0000))
Breast Cancer (METABRIC, Nature 2012 & Nat Commun 2016)	mRNA expression (microarray)	1	491	1349	25	0	23.875	2.65E-05	0.5689 37 (1.0000))	2.2745 64 (0.0688))	- 4.1152 88 (0.0001))	N/A

Mesothelioma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369	0	29	50	3	0	18.866	8.00E- 05	N/A	4.3194 25 (0.0000) *)	0.1704 78 (1.0000)	N/A
Colorectal Adenocarcinoma (TCGA, Nature 2012)	RNA Seq RPKM	0	53	138	2	0	18.847	8.08E- 05	N/A	4.2980 92 (0.0000) *)	- 0.3389 75 (1.0000)	N/A
Cervical Squamous Cell Carcinoma (TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNAS eqV2)	1	31	167	76	0	19.435	2.22E- 04	- 1.6182 48 (0.3168))	- 3.4296 09 (0.0018) *)	- 1.4463 39 (0.4442)	N/A
Sarcoma (TCGA,	mRNA Expression Batch	0	43	113	70	4	19.389	2.27E- 04	N/A	3.6669 49	- 0.8524 54	0.9530 27

PanCancer Atlas)	Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369	17	279	418	150	13	20.977	0.00032	- 2.0848 79 (0.1854)	(0.0007))*	(1.0000)	(1.0000)
Cancer Cell Line Encyclopedia (Novartis/Broad, Nature 2012)	mRNA expression (microarray)	17	279	418	150	13	20.977	0.00032	- 2.0848 79 (0.1854)	- 3.6159 35 (0.0015))*	- 2.0070 04 (0.2237)	- 0.1088 80 (1.0000)
Rectum Adenocarcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369	1	54	78	3	0	18.215	0.00039 71	- 1.9265 19 (0.1621)	3.8771 66 (0.0003))*	1.1674 00 (0.7291)	N/A
Liver Hepatocellular Carcinoma	EPHA2: mRNA Expression,	1	130	194	21	2	15.514	0.00374 5	0.3023 41	3.6972 48	- 0.3366 59	0.4544 54

(TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNAS eqV2)								(1.0000)	(0.0011)*	(1.0000)	(1.0000)
Stomach Adenocarcinoma (TCGA, PanCancer Atlas)	2	90	264	44	7	13.966	0.007404	-	2.072978	1.606072	1.750466	1.602806
	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369								(0.1909)	(0.5413)	(0.4002)	(0.5449)
Uterine Corpus Endometrial Carcinoma (TCGA, PanCancer Atlas)	3	61	395	43	5	12.916	0.0117	-	1.905863	1.039307	1.597383	2.268798
	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369								(0.2833)	(1.0000)	(0.5509)	(0.1164)

Skin Cutaneous Melanoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369	2	70	216	72	3	12.242	0.01564	1.0945 26 (1.0000)	2.6744 93 (0.0374)	0.0959 66 (1.0000)	1.6926 28 (0.4526)
Prostate Adenocarcinoma (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNAS eqV2)	0	44	438	4	1	10.112	0.01764	N/A	2.9055 02 (0.0110) [*]	1.3746 09 (0.5078)	- 0.0827 90 (1.0000)
Kidney Chromophobe (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNAS eqV2)	0	52	12	1	0	7.8781	0.01947	N/A	2.4983 40 (0.0187) [*]	1.8631 69 (0.0937)	N/A

Pediatric Wilms' Tumor (TARGET, 2018)	Epha2: mRNA expression (RNA-Seq RPKM)	0	22	74	5	0	7.4912	0.02362	N/A	2.6907 66 (0.0107) *)	- 0.1732 74 (1.0000)	N/A
Pheochromocytoma and Paraganglioma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369	4	96	60	1	0	8.8074	0.03196	- 1.4115 67 (0.4742)	2.2013 44 (0.0831)	1.9461 34 (0.1549)	N/A
Thyroid Carcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369	0	4	474	2	0	5.1773	0.08	N/A	2.2218 84 (0.0394)	0.5035 77 (0.9218)	N/A

Esophageal Adenocarcinoma (TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNAS eqV2)	1	64	83	32	1	7.6886	0.1037	- 1.4626 79 (0.7178)	0.9109 90 (1.0000)	- 1.6823 11 (0.4625)	- 0.3622 98 (1.0000)
Cholangiocarcinoma (TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNAS eqV2)	2	27	7	0	0	4.1691	0.1244	- 2.0378 40 (0.0623)	0.9721 00 (0.4965)	N/A	N/A
Brain Lower Grade Glioma (TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNAS eqV2)	0	191	303	13	0	4.0473	0.1322	N/A	0.7223 83 (0.7051)	- 1.7715 14 (0.1147)	N/A
Thymoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS	0	8	110	1	0	4.0322	1.33E-01	N/A	1.9823 34 (0.0712)	0.3691 15 (1.0000)	N/A

[illegible]


Metastatic Prostate Cancer, SU2C/PCF Dream Team (Robinson et al., Cell 2015)	mRNA expression / capture (RNA Seq RPKM)	2	21	87	7	0	4.069	0.254	- 1.8126 13 (0.2097)	0.9925 71 (0.9628)	0.3140 89 (1.0000)	N/A
Acute Myeloid Leukemia (TCGA, PanCancer Atlas)	mRNA Expression, RSEM (Batch normalized from Illumina HiSeq_RNAS eqV2)	0	1	160	4	0	2.4016	0.301	N/A	- 1.5391 42 (0.1857)	- 0.1995 32 (1.0000)	N/A
Testicular Germ Cell Tumors (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369	1	29	92	22	0	3.3144	0.3456	0.5748 46 (1.0000)	- 0.4431 10 (1.0000)	- 1.7511 61 (0.2398)	N/A

Adrenocortical Carcinoma (TCGA, PanCancer Atlas)	RSEM (Batch normalized from Illumina HiSeq_RNAS eqV2)	0	28	47	1	0	2.0003	0.3678	N/A	1.3463 97 (0.2673)	0.5501 03 (0.8734)	N/A
Uterine Carcinosarcoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369	0	16	22	16	2	2.44	0.4862	N/A	0.4760 71 (1.0000)	- 0.5502 92 (1.0000)	1.2151 02 (0.6730)
Pancreatic Adenocarcinoma (TCGA, PanCancer Atlas)	mRNA Expression Batch Normalized/ Merged from Illumina HiSeq_RNAS eqV2 syn4976369	2	50	106	9	1	3.3833	4.96E-01	- 1.1950 82 (1.0000)	0.1594 42 (1.0000)	- 0.6025 58 (1.0000)	1.2176 97 (1.0000)

Prostate Adenocarcinoma (MSKCC, Cancer Cell 2010)	mRNA Expression	0	5	77	3	0	1.3139	0.5184	N/A	- 0.4065 79 (1.0000)	- 1.0899 48 (0.4136)	N/A
Prostate Adenocarcinoma (Fred Hutchinson CRC, Nat Med 2016)	mRNA expression	0	39	84	10	0	0.0283 51	0.9859	N/A	0.1604 04 (1.0000)	0.0797 85 (1.0000)	N/A

CLAIMS

1. A peptide ligand specific for EphA2 comprising a polypeptide comprising at least three cysteine residues, separated by at least two loop sequences, and a non-aromatic molecular scaffold which forms covalent bonds with the cysteine residues of the polypeptide such that at least two polypeptide loops are formed on the molecular scaffold.
2. The peptide ligand as defined in claim 1, wherein said loop sequences comprise 2, 3, 5, 6 or 7 amino acid acids.
3. The peptide ligand as defined in claim 1 or claim 2, wherein said loop sequences comprise three cysteine residues separated by two loop sequences one of which consists of 2 amino acids and the other of which consists of 7 amino acids (such as those listed in Table 4).
4. The peptide ligand as defined in claim 1 or claim 2, wherein said loop sequences comprise three cysteine residues separated by two loop sequences both of which consist of 5 amino acids (such as those listed in Tables 3 and 4).
5. The peptide ligand as defined in claim 1 or claim 2, wherein said loop sequences comprise three cysteine residues separated by two loop sequences both of which consist of 6 amino acids (such as those listed in Tables 3 to 5 and 10).
6. The peptide ligand as defined in claim 1 or claim 2, wherein said loop sequences comprise three cysteine residues separated by two loop sequences one of which consists of 7 amino acids and the other of which consists of 3 amino acids (such as those listed in Table 4).
7. The peptide ligand as defined in claim 1 or claim 2, wherein said loop sequences comprise three cysteine residues separated by two loop sequences one of which consists of 6 amino acids and the other of which consists of 7 amino acids (such as those listed in Table 5).
8. The peptide ligand as defined in any one of claims 1 to 7, wherein the peptide ligand comprises an amino acid sequence selected from:



wherein X_1 and X_2 represent the amino acid residues between the cysteine residues listed in Tables 3 to 5 and 10 and C_i , C_{ii} and C_{iii} represent first, second and third cysteine residues, respectively or a pharmaceutically acceptable salt thereof.

5 9. The peptide ligand as defined in any one of claims 1 to 8, wherein the peptide ligand comprises an amino acid sequence selected from one or more of the peptide ligands listed in one or more Tables 3 to 5 and 10.

10 10. The peptide ligand as defined in any one of claims 1 to 8, wherein said loop sequences comprise three cysteine residues separated by two loop sequences both of which consist of 6 amino acids and the peptide ligand has the following amino acid sequence:

$C_i(\text{HyP})\text{LVNPLC}_{ii}\text{LHP}(\text{D-Asp})\text{W}(\text{HArg})\text{C}_{iii}$ (SEQ ID NO: 1); and

$C_i\text{PLVNPLC}_{ii}\text{LHPGWTC}_{iii}$ (SEQ ID NO: 97);

15 such as:

$C_i(\text{HyP})\text{LVNPLC}_{ii}\text{LHP}(\text{D-Asp})\text{W}(\text{HArg})\text{C}_{iii}$ (SEQ ID NO: 1);

wherein HyP is hydroxyproline, HArg is homoarginine and C_i , C_{ii} and C_{iii} represent first, second and third cysteine residues, respectively or a pharmaceutically acceptable salt thereof.

20

11. The peptide ligand as defined in claim 10, wherein said loop sequences comprise three cysteine residues separated by two loop sequences both of which consist of 6 amino acids and the peptide ligand has the following amino acid sequence:

$(\beta\text{-Ala})\text{-Sar}_{10}\text{-A}(\text{HArg})\text{D-}C_i(\text{HyP})\text{LVNPLC}_{ii}\text{LHP}(\text{D-Asp})\text{W}(\text{HArg})\text{C}_{iii}$ (SEQ ID NO: 2)

25 (BCY6099; Compound 66); and

$(\beta\text{-Ala})\text{-Sar}_{10}\text{-A}(\text{HArg})\text{D-}C_i\text{PLVNPLC}_{ii}\text{LHPGWTC}_{iii}$ (($\beta\text{-Ala})\text{-Sar}_{10}\text{-}$ (SEQ ID NO: 11))

(BCY6014; Compound 67);

such as:

$(\beta\text{-Ala})\text{-Sar}_{10}\text{-A}(\text{HArg})\text{D-}C_i(\text{HyP})\text{LVNPLC}_{ii}\text{LHP}(\text{D-Asp})\text{W}(\text{HArg})\text{C}_{iii}$ (SEQ ID NO: 2)

30 (BCY6099; Compound 66);

wherein Sar is sarcosine, HArg is homoarginine and HyP is hydroxyproline.

12. The peptide ligand as defined in any one of claims 1 to 11, wherein the molecular scaffold is selected from 1,1',1''-(1,3,5-triazinane-1,3,5-triyl)triprop-2-en-1-one (TATA) and
35 the peptide ligand is selected from any one of the peptide ligands listed in Tables 3, 4, 5 and 10.

13. The peptide ligand as defined in any one of claims 1 to 11, wherein the molecular scaffold is selected from 1,1',1''-(1,3,5-triazinane-1,3,5-triyl)triprop-2-en-1-one (TATA) and the peptide ligand is

(β -Ala)-Sar₁₀-A(HArg)D-C_i(HyP)LVNPLC_{ii}LHP(D-Asp)W(HArg)C_{iii} (SEQ ID NO: 2)

5 (BCY6099; Compound 66); and

(β -Ala)-Sar₁₀-A(HArg)D-C_iPLVNPLC_{ii}LHPGWT C_{iii} ((β -Ala)-Sar₁₀-(SEQ ID NO: 11))

(BCY6014; Compound 67);

such as:

(β -Ala)-Sar₁₀-A(HArg)D-C_i(HyP)LVNPLC_{ii}LHP(D-Asp)W(HArg)C_{iii} (SEQ ID NO: 2)

10 (BCY6099; Compound 66);

wherein Sar is sarcosine, HArg is homoarginine and HyP is hydroxyproline.

14. The peptide ligand as defined in any one of claims 1 to 13, which is selected from any one of Compounds 1-113 or a pharmaceutically acceptable salt thereof.

15

15. The peptide ligand as defined in any one of claims 1 to 13, which is Compound 66 (BCY6099) or Compound 67 (BCY6014) or a pharmaceutically acceptable salt thereof, such as Compound 66 (BCY6099) or a pharmaceutically acceptable salt thereof.

20 16. The peptide ligand as defined in any one of claims 1 to 15, wherein the pharmaceutically acceptable salt is selected from the free acid or the sodium, potassium, calcium, ammonium salt.

17. The peptide ligand as defined in any one of claims 1 to 16, wherein the EphA2 is human
25 EphA2.

18. A drug conjugate comprising a peptide ligand as defined in any one of claims 1 to 17, conjugated to one or more effector and/or functional groups.

30 19. The drug conjugate as defined in claim 18, wherein said cytotoxic agent is selected from DM1 or MMAE.

20. The drug conjugate as defined in claim 18 or claim 19, which additionally comprises a linker between said peptide ligand and said cytotoxic agents.

35

21. The drug conjugate as defined in claim 20, wherein said cytotoxic agent is MMAE and the linker is selected from: -Val-Cit-, -Trp-Cit-, -Val-Lys-, -D-Trp-Cit-, -Ala-Ala-Asn-, D-Ala-Phe-Lys- or -Glu-Pro-Cit-Gly-hPhe-Tyr-Leu- (SEQ ID NO: 98).

5 22. The drug conjugate as defined in claim 21, wherein said cytotoxic agent is MMAE and the linker is -Val-Cit-.

23. The drug conjugate as defined in claim 20, wherein said cytotoxic agent is DM1 and the linker is selected from: -S-S-, -SS(SO₃H)-, -SS-(Me)-, -(Me)-SS-(Me)-, -SS-(Me₂)- or -SS-(Me)-SO₃H-.

10

24. The drug conjugate as defined in any one of claims 18 to 23, which is selected from any one of BCY6027, BCY6028, BCY6031 and BCY6032; the BDCs listed in Tables 11 or 13; or BCY6033, BCY6082, BCY6136 and BCY6173.

15

25. The drug conjugate as defined in claim 24, which is selected from any one of: BCY6031, BCY6033, BCY6082, BCY6135, BCY6136, BCY6173, BCY6174 and BCY6175.

26. The drug conjugate as defined in claim 24 or claim 25, which is BCY6136.

20

27. A pharmaceutical composition which comprises the peptide ligand of any one of claims 1 to 17 or the drug conjugate of any one of claims 18 to 26, in combination with one or more pharmaceutically acceptable excipients.

25 28. The drug conjugate as defined in any one of claims 18 to 26, for use in preventing, suppressing or treating a disease or disorder characterised by overexpression of EphA2 in diseased tissue.

29. The drug conjugate as defined in any one of claims 18 to 26, for use in preventing, suppressing or treating cancer.

30

30. The drug conjugate for use as defined in claim 29, wherein the cancer is selected from: prostate cancer, lung cancer (such as non-small cell lung carcinomas (NSCLC)), breast cancer (such as triple negative breast cancer), gastric cancer, ovarian cancer, oesophageal cancer, multiple myeloma and fibrosarcoma.

35

31. A method of preventing, suppressing or treating cancer, which comprises administering to a patient in need thereof a drug conjugate as defined in any one of claims 18 to 26, wherein said patient is identified as having an increased copy number variation (CNV) of EphA2.

5