

What is claimed is:

1. A method for treating or reducing the risk of developing a fetal and neonatal alloimmune and/or autoimmune disorder comprising administering to a pregnant women a composition comprising an antibody
5 comprising a light chain having the amino acid sequence of SEQ ID NO:19 and a heavy chain having the amino acid sequence of SEQ ID NO:24, wherein the composition is administered to the pregnant woman from about gestational week 13 to about gestational week 35.
2. The method of claim 1, wherein the antibody is administered at a dose per administration and is based on
10 the weight of the pregnant women at first dosing and is not adjusted upward based on weight gain by the pregnant women.
3. The method of claim 1, wherein the antibody is administered at a dose per administration and is based on the weight of the pregnant women at first dosing and is adjusted upward based on weight gain by the
15 pregnant women.
4. The method of claim 1, wherein the composition is administered at least every other week.
5. The method of claim 1, wherein the composition is administered every other week.
20
6. The method of claim 1, wherein the composition is administered at least every week.
7. The method of claim 1, wherein IVIG is administered to the pregnant women after cessation of administration of the antibody and prior to birth.
25
8. The method of claim 7, wherein IVIG is administered to the pregnant women 40 – 100 hrs prior to birth.
9. The method of claim 7, wherein the IVIG is administered at a dose of 200 mg/kg – 1000 mg/kg based on the weight of the pregnant women.
30
10. The method of claim 1, wherein the composition is administered from gestational week 13 to gestational week 35.
11. The method of claim 1, wherein the composition is administered from gestational week 14 to gestational
35 week 35.
12. The method of claim 1, wherein the composition is administered from gestational week 15 to gestational week 35.
- 40 13. The method of claim 1, wherein the composition is administered from gestational week 16 to gestational week 35.