

NONAUTOLOGOUS MULTI-STRESSED CANCER CELLS AND USES THEREOF FOR VACCINATING AND TREATING CANCERS

FIELD OF INVENTION

5 The present invention relates to the field of advanced therapy medicinal products (AMTPs) for cell therapy. In particular, it relates to a composition comprising stressed HT-29, HCT-116 and LoVo cells, and immunogenic stress proteins produced by these cells in response to stresses applied *in vitro*. The composition allows to simultaneously counteract multiple cell resistance mechanisms observed in cancer cells, and is therefore
10 suitable for vaccinating and treating cancers in human patients.

BACKGROUND OF INVENTION

The immune system is based on two defense mechanisms, namely, innate immunity, which is rapid but nonspecific; and acquired immunity, which is slower, but specific and
15 has a memory. These two complementary mechanisms provide the ability to fight against external and internal “aggressions” by mobilizing the cells, either directly, during the cell-mediated immune response, or through the secretion of active molecules (*e.g.*, immunoglobulins, cytokines, etc.) during the humoral immune response.

Immunity not only protects the host against development of primary nonviral cancers but
20 also sculpts tumor immunogenicity. Cancer immunoediting is a process consisting of three phases: elimination (*i.e.*, cancer immunosurveillance), equilibrium, and escape. In the first phase, the immune system fights against tumor proliferation, entailing the involvement of tissue and environmental changes that are associated with the tumor (with the mobilization of non-specific cells such as macrophage, NK cells, DC cells, etc., the
25 secretion of anti-proliferative and/or apoptotic molecules, the production of cytokines, and the mobilization and activation of CD4⁺ and CD8⁺ T cells). During the second phase, sensitive cancer cells are eliminated and an immune selection of the most resistant cells operates. The mechanisms of resistance that are set in motion are resistance to apoptosis, secretion of inhibiting cytokines (such as TGF- β , IL-10, PGE2 or IDO), alteration of the

antigen presentation (with a partial or complete loss of expression of class I major histocompatibility complex (MHC)), secretion of neutralizing molecules, and expression of MICA or MICB transcripts to counter-attack the cell-mediated immune system. During this phase, the number of cancer cells destroyed is in equilibrium with the number of resistant cells, hence the name of this phase. It corresponds *inter alia* to the remission phase observed during cancer-patient treatments. During the third phase, cancer cells that are resistant to the various protective mechanisms of the immune system proliferate. These cancer cells then develop a tumor mass, that is the clinical manifestation of the physiological escape phenomenon. A related escape phenomenon is also observed in advanced and metastatic stages of cancers.

Anti-cancer treatments include surgery, which objective is primarily to remove or reduce the tumor mass, but without any real inhibitory effect on the process of carcinogenesis, hence surgery is generally supplemented with various treatment therapies to fully eliminate cancer cells.

Radiation therapy is intended to bring about an alteration in the DNA of rapidly proliferating cells, which is the case with the cancer cells. The side effects of radiotherapy are twofold: first, not only cancer cells are irradiated but also healthy cells, which can cause the “cancerization” of healthy cells; second, cancer cells develop resistance to radiation-induced apoptosis by expressing chaperone proteins such as HSP, GRP, etc., resulting in an escape phenomenon.

Chemotherapies are intended to eliminate cancer cells, by acting either on the cancer cells themselves, or by inhibiting specific metabolic pathways. Chemotherapy agents include agents directly interacting with DNA such as electrophilic agents, intercalating agents, or splitting agents, agents indirectly interacting with DNA such as inhibitors of DNA synthesis (antimetabolites, topoisomerase inhibitors, inhibitors of spindle formation, etc.), neovascularization inhibitors, proteasome inhibitors, and the like. Here again, resistance mechanisms are developed by cancer cells and despite polychemotherapy strategies, a relapse phenomenon can be observed with massive tumor proliferation.

Immunotherapies comprises passive immunotherapy, based on the injection of antibodies or cytokines that block a receptor, induce cell lysis, stimulate cytotoxicity, or lift the inhibition of apoptosis; and active immunotherapy, also called “immunization by vaccination”, in which patients are immunized against cancer using “advanced therapy medicinal products” (ATMPs), a new class of medicines defined in Europe by
5 EU Regulation 1394/2007.

A wide variety of ATMPs for gene therapy (called GTMPs) have been used in clinical trials in a wide variety of cancers; as few examples, adenoviral vectors were used to express p53 in head and neck carcinoma or to express CD40L in bladder carcinoma;
10 oncolytic herpes virus encoding GM-CSF was used in patients with melanoma; and T cells engineered with chimeric antigen receptors (CARs) redirected against various tumor-associated antigens have also been recently used.

AMTPs for cell therapy (called CTMPs) have also been used, including autologous dendritic cells (DCs) loaded with tumor antigens, in the attempt to elicit clinically relevant
15 immune responses. For example, the FDA and EMA approved in 2010 and 2013, respectively, a DC-based therapeutic vaccine for prostate cancer (Sipuleucel-T, trade name “Provenge”, developed by Dendreon Pharmaceuticals, LLC) – although the European Commission withdrew the marketing authorization for Sipuleucel-T in the European Union in 2015.

20 However, with AMTPs as well, escape phenomena are observed, that are similar to the natural escape phenomena described in 3-phase cancer immunoediting involving the immune system.

There is therefore still a need for therapeutic strategies for the treatment of cancer, which can provide an inherent effectiveness and/or contribute to a multitherapeutic strategy.

25 A number of studies have been carried out to better understand cancer resistance phenomena, and in particular the role of chaperone proteins in the resistance to apoptosis. In particular, heat shock proteins (HSPs), glucose regulated proteins (GRPs) and multidrug resistance proteins (MRP) are known to be resistance factors and to have a protective effect. These proteins are produced by cancer cells as a mechanism of

resistance, as a result of metabolic attacks such as hypoxia, low carbohydrate concentration, etc., of physical attacks such as thermal stress, radiations, etc., or of chemical attacks such as medication.

As a consequence, studies have been carried out to develop treatment strategies based on inhibition of these chaperone proteins, but these are not without side effects, including a high toxicity level for overall mixed results. Studies have also been carried out to use chaperone proteins, in particular HSP70, as factors of immunization against cancer cells in mice. This approach however focuses on a specific chaperone protein and does not take into account the entire set of protection mechanisms developed by cancer cells.

To address this issue, the Inventors have described an *in vitro* process for obtaining pharmaceutical or vaccine compositions capable of counteracting cancer cell resistance mechanisms, a process which is otherwise adapted to the *in vitro* production of autologous cancer cells with particular resistance mechanism[s] identical to that developed *in situ* by cancer cells subjected *in vivo* to a specific stress (or stresses) in the course of a treatment protocol applied to a patient (U.S. Pat. 11,096,995; European Pat. 3 057 981). Following administration of these compositions in mice *in vivo*, data showed a favorable impact on tumor volume and weight, validating the proof of concept.

Here, the Inventors go further, and provide with a new composition comprising a selection of nonautologous multi-stressed cells, simultaneously counteracting multiple resistance mechanisms, suitable for vaccinating and treating various cancer in human patients.

SUMMARY

The present invention is as disclosed hereafter, and in particular in the appended claims.

In particular, the present invention relates to a composition comprising (i) stressed HT-29, HCT-116 and LoVo cells, and (ii) immunogenic stress proteins produced by these cells in response to a stress applied *in vitro*.

In one embodiment, stressed HT-29, HCT-116 and LoVo cells have developed resistance mechanism in response to one or several stress[es] applied *in vitro*, selected from the group comprising radiations, thermal stress, chemical stress, metabolic stress and any combinations thereof, leading to the production of the stress proteins.

- 5 In one embodiment, stressed HT-29, HCT-116 and LoVo cells are non-proliferative.

In one embodiment, immunogenic stress proteins are haptenated. In one embodiment, immunogenic stress proteins are haptenated with an hapten selected from the group comprising 2,4-dinitrophenyl (DNP); 2,4-dinitrofluorobenzene; sulfanilic acid; *N*-iodoacetyl-*N'*-(5-sulfonic-naphthyl)ethylene diamine; anilin; *p*-amino benzoic acid; biotin; fluorescein and derivatives thereof (including FITC, TAMRA, and Texas Red);
 10 digoxigenin; 5-nitro-3-pyrazolecarbamide; 4,5-dimethoxy-2-nitrocinnamide; 2-(3,4-dimethoxyphenyl)-quinoline-4-carbamide; 2,1,3-benzoxadiazole-5-carbamide; 3-hydroxy-2-quinoxalinecarbamide, 4-(dimethylamino)azobenzene-4'-sulfonamide (DABSYL); rotenone
 15 isooxazolinI(*E*)-2-(2-(2-oxo-2,3-dihydro-1*H*-benzo[*b*][1,4]diazepin-4-yl)phenoxy)aceta-
 mide; 7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylic acid; and
 2-acetamido-4-methyl-5-thiazolesulfonamide; and
p-methoxyphenylpyrazopodophyllamide. In one embodiment, immunogenic stress proteins are haptenated with 2,4-dinitrophenyl (DNP).

- 20 In one embodiment, the composition is a pharmaceutical composition or a vaccine composition, and further comprises at least one pharmaceutically acceptable excipient.

In one embodiment, the composition comprises from about 10^5 to about 10^8 stressed HT-29, HCT-116 and LoVo cells.

- 25 The present invention further relates to this composition, for use in treating cancer in a subject in need thereof. It also relates to a method of treating cancer in a subject in need thereof, comprising administering this composition to the subject.

The present invention further relates to an intermediate composition comprising (i) one of stressed HT-29 cells, stressed HCT-116 cells and stressed LoVo cells, and (ii) stress

proteins, wherein the one of stressed HT-29 cells, stressed HCT-116 cells or stressed LoVo cells have developed resistance mechanism in response to (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*, leading to the production of the stress proteins.

- 5 The present invention further relates to an intermediate composition comprising (i) one of stressed HT-29 cells, stressed HCT-116 cells and stressed LoVo cells, and (ii) stress proteins, wherein the one of stressed HT-29 cells, stressed HCT-116 cells or stressed LoVo cells have developed resistance mechanism in response to (i) a metabolic stress, and (ii) a chemical stress applied *in vitro*, leading to the production of the stress proteins.
- 10 The present invention further relates to a method of manufacturing said intermediate compositions, comprising the following steps:
 - a) cultivating HT-29, HCT-116 or LoVo cells in a suitable culture medium;
 - b) subjecting the HT-29, HCT-116 or LoVo cells cultured in step a) to one or several stress[es] *in vitro*, wherein these HT-29, HCT-116 or LoVo cells develop resistance
 - 15 mechanisms in response to the one or several stress[es] and thereby produce stress proteins,
 - c) recovering the stressed HT-29, HCT-116 or LoVo cells together with the stress proteins they have produced in step b), and
 - d) treating the stressed HT-29, HCT-116 or LoVo cells and the stress proteins they
 - 20 have produced, all together recovered in step c), with a molecule or by a process capable of rendering the stress proteins immunogenic.

In one embodiment, step c) is carried out at least several hours after completion of step b), preferably at least 12 hours or more after completion of step b).

- 25 In one embodiment, step d) comprises linking the stress proteins to or complexing the stress proteins with a means capable to confer immunogenicity. In one embodiment, the means capable to confer immunogenicity is an hapten. In one embodiment, the means capable to confer immunogenicity is an hapten selected from the group comprising 2,4-dinitrophenyl (DNP); 2,4-dinitrofluorobenzene; sulfanilic acid; *N*-iodoacetyl-*N'*-(5-sulfonic-naphthyl)ethylene diamine; anilin; *p*-amino benzoic acid;

biotin; fluorescein and derivatives thereof (including FITC, TAMRA, and Texas Red);
 digoxigenin; 5-nitro-3-pyrazolecarbamide; 4,5-dimethoxy-2-nitrocinnamide;
 2-(3,4-dimethoxyphenyl)-quinoline-4-carbamide; 2,1,3-benzoxadiazole-5-carbamide;
 3-hydroxy-2-quinoxalinecarbamide, 4-(dimethylamino)azobenzene-4'-sulfonamide
 5 (DABSYL); rotenone isooxazoline;
 (E)-2-(2-(2-oxo-2,3-dihydro-1H-benzo[b][1,4]diazepin-4-yl)phenoxy)acetamide;
 7-(diethylamino)-2-oxo-2H-chromene-3-carboxylic acid;
 2-acetamido-4-methyl-5-thiazolesulfonamide; and
 p-methoxyphenylpyrazopodophyllamide. In one embodiment, the means capable to
 10 confer immunogenicity is 2,4-dinitrophenyl (DNP).

In one embodiment – where the method is for manufacturing an intermediate composition
 “DS-A” comprising (i) one of stressed HT-29 cells, stressed HCT-116 cells and stressed
 LoVo cells, and (ii) stress proteins, wherein the one of stressed HT-29 cells, stressed
 HCT-116 cells or stressed LoVo cells have developed resistance mechanism in response
 15 to (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*, leading
 to the production of the stress proteins – , step b) of said method comprises subjecting the
 HT-29, HCT-116 or LoVo cells cultured in step a) to the following stresses *in vitro*,
 applied concomitantly or successively:

- (i) an *in vitro* culture in a depleted medium, under hypoxia, and/or at low pH;
 20 preferably an *in vitro* culture in low serum culture conditions in a 2 % FBS culture
 medium,
- (ii) an *in vitro* radiation with a total dose ranging from about 0.25 to about 25 Gy,
 preferably from about 1 to about 15 Gy, preferably for a period ranging from about 1 to
 about 20 minutes; preferably an *in vitro* radiation with a total dose of 10 Gy for a period
 25 of about 1 to about 5 minutes, and
- (iii) an *in vitro* thermic choc at a temperature ranging from about 38°C to about 45°C,
 applied to the cells for a period ranging from about 15 minutes to about 4 hours;
 preferably an *in vitro* thermic choc at a temperature of about 42°C for a period of about
 60 minutes.

In one embodiment – where the method is for manufacturing an intermediate composition “DS-B” comprising (i) one of stressed HT-29 cells, stressed HCT-116 cells and stressed LoVo cells, and (ii) stress proteins, wherein the one of stressed HT-29 cells, stressed HCT-116 cells or stressed LoVo cells have developed resistance mechanism in response to (i) a metabolic stress, and (ii) a chemical stress applied *in vitro*, leading to the production of the stress proteins – step b) of said method comprises subjecting the HT-29, HCT-116 or LoVo cells cultured in step a) to the following stresses *in vitro*, applied concomitantly or successively:

- (i) an *in vitro* culture in a depleted medium, under hypoxia, and/or at low pH; preferably an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium,
- (ii) an *in vitro* exposition to at least one or several chemotherapeutic agents and/or alcohols, for a period ranging from about 6 hours to about 120 hours.

According to the latest embodiment, the *in vitro* exposition to at least one or several chemotherapeutic agents and/or alcohols may be as follows:

- when the cells are HT-29 cells, the *in vitro* exposition at (ii) is to about 13 μ M oxaliplatin for a period of about 72 hours; or
- when the cells are HCT-116 cells, the *in vitro* exposition at (ii) is to about 315 nM SN-38 (7-ethyl-10-hydroxy-camptothecin) for a period of about 48 hours; or
- when the cells are LoVo cells, the *in vitro* exposition at (ii) is to about 5 μ M fluorouracil (5-FU) for a period of about 48 hours.

The present invention further relates to a method of manufacturing the composition of the invention, comprising the following steps:

- a. obtaining six intermediate compositions, wherein the six intermediate compositions are:
 - 1) an intermediate composition “DS-A” comprising stressed HT-29 cells and stress proteins,
 - 2) an intermediate composition “DS-A” comprising stressed HCT-116 cells and stress proteins,

- 3) an intermediate composition “DS-A” comprising stressed LoVo cells and stress proteins,
 - 4) an intermediate composition “DS-B” comprising stressed HT-29 cells and stress proteins,
 - 5 5) an intermediate composition “DS-B” comprising stressed HCT-116 cells and stress proteins,
 - 6) an intermediate composition “DS-B” comprising stressed LoVo cells and stress proteins,
 - b. mixing these six intermediate compositions together.
- 10 In one embodiment, the six intermediate compositions are mixed together in an equal ratio of stressed HT-29, HCT-116 and LoVo cells.

DEFINITIONS

In the present invention, the following terms have the following meanings:

- 15 “**HT-29 cells**”, as used herein, refers to a human colon adenocarcinoma cell line isolated from a primary tumor in 1964 from a 44-year-old woman. It comprises, *inter alia*, the following sequence variations:
- heterozygous for *APC* p.Glu853Ter (c.2557G>T),
 - heterozygous for *APC* p.Thr1556Asnfs*3 (c.4666dupA),
 - 20 - heterozygous for *BRAF* p.Val600Glu (c.1799T>A),
 - heterozygous for *PIK3CA* p.Pro449Thr (c.1345C>A),
 - homozygous for *SMAD4* p.Gln311Ter (c.931C>T), and
 - homozygous for *TP53* p.Arg273His (c.818G>A).

- This cell line is referenced under accession number “CVCL_0320” on the Cellosaurus database and “HTB-38™” in the ATCC repository collection, from which it is commercially available. Other accession numbers for this cell line include “0111” in the Banco Células do Rio de Janeiro (BCRJ); “ACC 299” in the Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures GmbH; “91072201” in the General Collection of the European Collection of Authenticated Cell Cultures
- 25

(ECACC); “30038” in the Korean Cell Line Bank (KCLB); and “GDC0149” in the China Center for Type Culture Collection (CCTCC). The content of these collections, in particular the description of the genotype, HLA typing, STR profile and phenotype of this cell line, is hereby incorporated by reference.

5 “**HCT-116 cells**”, as used herein, refers to a human colon carcinoma cell line isolated from a primary tumor in 1981 from a 48-year-old man. It comprises, *inter alia*, the following sequence variations:

- homozygous for *ACVR2A* p.Lys437Argfs*5 (c.1310delA),
- heterozygous for *BRCA2* p.Ile2675Aspfs*6 (c.8021dupA),
- 10 - heterozygous for *CDKN2A* p.Arg24Serfs*20 (c.68dupG),
- heterozygous for *CDKN2A* p.Asp74fs*21 (c.220delG),
- heterozygous for *CDKN2A* p.Glu33Argfs*20 (c.97delG),
- heterozygous for *CTNNA1* p.Ser45del (c.133_135delTCT),
- heterozygous for *EP300* p.Met1470Cysfs*22 (c.4408delA),
- 15 - heterozygous for *EP300* p.Asn1700Thrfs*9 (c.5099delA),
- heterozygous for *KRAS* p.Gly13Asp (c.38G>A),
- heterozygous for *PIK3CA* p.His1047Arg (c.3140A>G), and
- homozygous for *TGFBR2* p.Lys128Serfs*35 (c.383delA).

This cell line is referenced under accession number “CVCL_0291” on the Cellosaurus
 20 database and “CCL-247TM” in the ATCC repository collection, from which it is commercially available. Other accession numbers for this cell line include “0288” in the Banco Células do Rio de Janeiro (BCRJ); “ACC 581” in the Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures GmbH; “91091005” in the General Collection of the European Collection of Authenticated Cell Cultures
 25 (ECACC); and “10247” in the Korean Cell Line Bank (KCLB). The content of these collections, in particular the description of the genotype, HLA typing, STR profile and phenotype of this cell line, is hereby incorporated by reference.

“**LoVo cells**”, as used herein, refers to a human colorectal adenocarcinoma cell line isolated from a fragment of a metastatic tumor nodule in the left supraclavicular region

in 1971 from a 56-year-old man. It comprises, *inter alia*, the following sequence variations:

- heterozygous for *ACVR2A* p.Lys437Argfs*5 (c.1310delA),
- heterozygous for *APC* p.Arg1114Ter (c.3340C>T),
- 5 - heterozygous for *APC* p.Met1431fs*42 (c.4289delC),
- heterozygous for *APC* p.Arg2816Gln (c.8447G>A),
- heterozygous for *B2M* p.Leu15Phefs*41 (c.43_44delCT),
- heterozygous for *FBXW7* p.Arg505Cys (c.1513C>T),
- heterozygous for *KRAS* p.Gly13Asp (c.38G>A),
- 10 - heterozygous for *SMAD2* p.Ala292Val (c.875C>T), and
- homozygous for *TGFBR2* p.Lys128Serfs*35 (c.383delA).

This cell line is referenced under accession number “CVCL_0399” on the Cellosaurus database and “CCL-229TM” in the ATCC repository collection, from which it is commercially available. Other accession numbers for this cell line include “0332” in the
 15 Banco Células do Rio de Janeiro (BCRJ); “ACC 350” in the Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures GmbH; “87060101” in the General Collection of the European Collection of Authenticated Cell Cultures (ECACC); and “10229” in the Korean Cell Line Bank (KCLB). The content of these collections, in particular the description of the genotype, HLA typing, STR profile and
 20 phenotype of this cell line, is hereby incorporated by reference

“**Chemotherapeutic agent**”, as used herein, refers to any molecule that is effective in inhibiting tumor growth. Examples of chemotherapeutic agents include those described under subgroup L01 of the Anatomical Therapeutic Chemical (ATC) Classification System. Further examples of chemotherapeutic agents include, but are not limited to:

- 25 i. alkylating agents, such as, *e.g.*:
 - nitrogen mustards, including chlormethine, cyclophosphamide, ifosfamide, trofosfamide, chlorambucil, melphalan, prednimustine, bendamustine, uramustine, chlornaphazine, cholophosphamide, estrarnustine, mechlorethamine, mechlorethamine oxide hydrochloride, novembichin,
 - 30 phenesterine, uracil mustard and the like;

- nitrosoureas, including carmustine, lomustine, semustine, fotemustine, nimustine, ranimustine, streptozocin, chlorozotocin, and the like;
- alkyl sulfonates, including busulfan, mannosulfan, treosulfan, and the like;
- aziridines, including carboquone, thiotepa, triaziquone, triethylenemelamine, benzodopa, meturedopa, uredopa, and the like; hydrazines, including procarbazine, and the like;
- triazenes, including dacarbazine, temozolomide, and the like; ethylenimines and methylamelamines, including altretamine, triethylenemelamine, triethylenephosphoramidate, triethylenethiophosphoramide, trimethylolomelamine and the like;
- and others, including mitobronitol, pipobroman, actinomycin, bleomycin, mitomycins (including mitomycin C, and the like), plicamycin, and the like;
- ii. acetogenins, such as, *e.g.*, bullatacin, bullatacinone, and the like;
- iii. benzodiazepines, such as, *e.g.*, 2-oxoquazepam, 3-hydroxyphenazepam, bromazepam, camazepam, carburazepam, chlordiazepoxide, cinazepam, cinolazepam, clonazepam, cloniprazepam, clorazepate, cyprazepam, delorazepam, demoxepam, desmethylflunitrazepam, devazepide, diazepam, diclazepam, difludiazepam, doxefazepam, elfazepam, ethyl carfluzepate, ethyl dirazepate, ethyl loflazepate, flubromazepam, fletazepam, fludiazepam, flunitrazepam, flurazepam, flutemazepam, flutoprazepam, fosazepam, gidazepam, halazepam, iclazepam, irazepine, kenazepine, ketazolam, lorazepam, lormetazepam, lufuradom, meclonazepam, medazepam, menitrazepam, metaclazepam, motrazepam, *N*-desalkylflurazepam, nifoxipam, nimetazepam, nitemazepam, nitrazepam, nitrazepate, nordazepam, nortetrazepam, oxazepam, phenazepam, pinazepam, pivoxazepam, prazepam, proflazepam, quazepam, QH-II-66, reclazepam, RO4491533, Ro5-4864, SH-I-048A, sulazepam, temazepam, tetrazepam, tifluadom, tolufazepam, triflunordazepam, tuclazepam, uldazepam, arfendazam, clobazam, CP-1414S, lofendazam, triflubazam, girisopam, GYKI-52466, GYKI-52895, nerisopam, talampanel, tofisopam, adinazolam, alprazolam, bromazolam, clonazolam, estazolam, flualprazolam, flubromazolam, flunitrazolam, nitrazolam, pyrazolam, triazolam, bretazenil, climazolam, EVT-201, FG-8205, flumazenil, GL-II-73, imidazenil, ¹²³I-iomazenil, L-655,708, loprazolam,

- midazolam, PWZ-029, remimazolam, Ro15-4513, Ro48-6791, Ro48-8684, Ro4938581, sarmazenil, SH-053-R-CH₃-2'F, cloxazolam, flutazolam, haloxazolam, mexazolam, oxazolam, bentazepam, clotiazepam, brotizolam, ciclotizolam, deschloroetizolam, etizolam, fluclotizolam, israpafant, JQ1, metizolam, olanzapine, telenzepine, lopirazepam, zapizolam, razobazam, ripazepam, zolazepam, zomebazam, zometapine, premazepam, clazolam, anthramycin, avizafone, rilmazafone, and the like;
- 5 iv. antimetabolites, such as, *e.g.*:
- 10 ■ antifolates, including aminopterin, methotrexate, pemetrexed, pralatrexate, pteropterin, raltitrexed, denopterin, trimetrexate, pemetrexed, and the like;
- 15 ■ purine analogues, including pentostatin, cladribine, clofarabine, fludarabine, nelarabine, tioguanine, mercaptopurine, and the like;
- 15 ■ pyrimidine analogues, including fluorouracil (5-FU), capecitabine, doxifluridine, tegafur, tegafur/gimeracil/oteracil, carmofur, floxuridine, cytarabine, gemcitabine, azacytidine, decitabine, and the like; and
- 15 ■ hydroxycarbamide;
- v. androgens, such as, *e.g.*, calusterone, dromostanolone propionate, epitiostanol, mepitiothane, testolactone, and the like;
- vi. anti-adrenals, such as, *e.g.*, aminoglutethimide, mitotane, trilostane, and the like;
- 20 vii. folic acid replenishers, such as, *e.g.*, frolinic acid, and the like;
- viii. maytansinoids, such as, *e.g.*, maytansine, ansamitocins, and the like;
- ix. platinum analogs, such as, *e.g.*, platinum, carboplatin, cisplatin, dicycloplatin, nedaplatin, oxaliplatin, satraplatin, and the like;
- x. antihormonal agents, such as, *e.g.*:
- 25 ■ anti-estrogens, including tamoxifen, raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, toremifene, and the like;
- 25 ■ anti-androgens, including flutamide, nilutamide, bicalutamide, leuprolide, goserelin, and the like;
- 30 xi. trichothecenes, such as, *e.g.*, T-2 toxin, verracurin A, roridinA, anguidine and the like;

- xii. toxoids, such as, *e.g.*, cabazitaxel, docetaxel, larotaxel, ortataxel, paclitaxel, tesetaxel, and the like;
- xiii. others, such as, *e.g.*, camptothecin (including its derivatives: belotecan, cositecan, etirinotecan, pegol, exatecan, gimatecan, irinotecan, lurtotecan, rubitecan, silatecan, SN-38 (7-ethyl-10-hydroxy-camptothecin), and topotecan); bryostatin; callostatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); cryptophycins (including cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including its synthetic analogues: KW-2189 and CBI-TMI); eleutherobin; pancratistatin; sarcodictyin; spongistatin; aclacinomysins; authramycin; azaserine; bleomycin; cactinomycin; carabycin; canninomycin; carzinophilin; chromomycins; dactinomycin; daunorubicin; detorubicin; 6-diazo-5-oxo-L-norleucine; doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin, deoxydoxorubicin, and the like); epirubicin; esorubicin; idanrbicin; marcellomycin; mycophenolic acid; nogalamycin; olivomycins; peplomycin; potfiromycin; puromycin; quelamycin; rodorubicin; streptomgrin; streptozocin; tubercidin; ubenimex; zinostatin; zorubicin; aceglatone; aldophospharnide glycoside; aminolevulinic acid; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfornithine; elliptinium acetate; epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; PSK[®]; razoxane; rhizoxin; sizofiran; spirogennanium; tenuazonic acid; 2,2',2''-trichlorotriethylarnine; urethan; vindesine; dacarbazine; mannomustine; mitobromtol; mitolactol; pipobroman; gacytosine; arabinoside; 6-thioguanine; vinblastine; etoposide; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; xeloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; topoisomerase I inhibitor SN38; difluoromethylornithine; retinoic acid; and the like.

“**Detectable levels**”, as used herein in the context of a protein, in particular of a stress protein, means that said protein is present in the composition at stake in an amount or concentration that can be detected by means and methods classically used by the skilled

artisan to detect proteins. Such means and methods are well-known in the art, and include, without limitation, mass spectrometry, such as liquid chromatography coupled to tandem mass spectrometry (LC-MS/MS), as detailed in Examples 3-5 below.

“**Hapten**”, as used herein, refers to a small molecule that elicits an immune response only when attached to a large carrier such as a protein. Once the patient has generated antibodies to a hapten-carrier conjugate, the hapten may be able to bind to the antibodies, but it will usually not initiate an immune response; usually, only the hapten-carrier conjugate can do this. Examples of haptens include, but are not limited to, 2,4-dinitrophenyl (DNP); 2,4-dinitrofluorobenzene; sulfanilic acid; *N*-iodoacetyl-*N'*-(5-sulfonic-naphthyl)ethylene diamine; anilin; *p*-amino benzoic acid; biotin; fluorescein and derivatives thereof (including FITC, TAMRA, and Texas Red); digoxigenin; 5-nitro-3-pyrazolecarbamide; 4,5-dimethoxy-2-nitrocinnamide; 2-(3,4-dimethoxyphenyl)-quinoline-4-carbamide; 2,1,3-benzoxadiazole-5-carbamide; 3-hydroxy-2-quinoxalinecarbamide, 4-(dimethylamino)azobenzene-4'-sulfonamide (DABSYL); rotenone isooxazoline; (*E*)-2-(2-(2-oxo-2,3-dihydro-1*H*-benzo[*b*][1,4]diazepin-4-yl)phenoxy)acetamide; 7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylic acid; 2-acetamido-4-methyl-5-thiazolesulfonamide; and *p*-methoxyphenylpyrazopodophyllamide.

“**Overexpression**”, and declinations thereof, refers herein to a relative or absolute expression and/or abundance of a marker or protein which is at least 10 % higher, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more higher in a subject cell or subject cell population, as compared to a reference cell or reference cell population. Alternatively or additionally, overexpression may also refer to a mean fluorescence intensity (MFI) which is at least 10 % higher, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more higher in a subject cell or subject cell population, as compared to a reference cell or reference cell population. In particular, the reference cell is a cell (or the reference cell population is a population comprising or preferably consisting of those cells) from the same cell line as the subject cell, but which has not been subject to the method of manufacturing the composition or intermediate

- compositions described herein; in particular a cell (or a population comprising or preferably consisting of those cells) from the same cell line as the subject cell, which is cultured in classical conditions such as, *e.g.*, in 10 % FBS, and is not subjected to any of a metabolic stress, radiations, a thermal stress and/or a chemical stress applied *in vitro*, as described hereafter; or alternatively that is cultured in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, or at low pH (*e.g.*, below pH 6.5) (*i.e.*, subjected to a metabolic stress), but not subjected to any other stress among radiations, a thermal stress and/or a chemical stress applied *in vitro*, as described hereafter.
- 10 “**Underexpression**”, and declinations thereof, refers herein to a relative or absolute expression and/or abundance of a marker or protein which is at least 10 % lower, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more lower in a subject cell or subject cell population, as compared to a reference cell or reference cell population. Alternatively or additionally, overexpression may also refer to a mean
- 15 fluorescence intensity (MFI) which is at least 10 % lower, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more lower in a subject cell or subject cell population, as compared to a reference cell or reference cell population. In particular, the reference cell (or the reference cell population is a population comprising or preferably consisting of those cells) is a cell from the same cell line as the subject cell, but which
- 20 has not been subject to the method of manufacturing the composition or intermediate compositions described herein; in particular a cell (or a population comprising or preferably consisting of those cells) from the same cell line as the subject cell, which is cultured in classical conditions such as, *e.g.*, in 10 % FBS, and is not subjected to any of a metabolic stress, radiations, a thermal stress and/or a chemical stress applied *in vitro*, as described hereafter; or alternatively that is cultured in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, or at low pH (*e.g.*, below pH 6.5) (*i.e.*, subjected to a metabolic stress), but not subjected to any other stress among radiations, a thermal stress and/or a chemical stress applied *in vitro*, as described hereafter.
- 25

“**Vaccine composition**”, as used herein, refers to a composition which comprises at least one antigen or immunogen (such as, *e.g.*, immunogenic stress and/or resistance proteins) and optionally an adjuvant, in a pharmaceutically acceptable excipient, and which is useful for inducing an immune response in a patient upon administration.

- 5 “**Adjuvant**”, as used herein, refers to a substance that enhances, increases and/or potentiates an immune response to an antigen (such as, *e.g.*, immunogenic stress and/or resistance proteins) in a subject upon administration.

“**Pharmaceutically acceptable excipient**”, as used herein, refers to a solid, semi-solid or liquid component of a pharmaceutical composition or a vaccine composition that is not
10 an active ingredient (*i.e.*, that is neither the stressed HT-29, HCT-116 and LoVo cells nor the immunogenic stress and/or resistance proteins), and that does not produce an adverse, allergic or other untoward reaction when administered to a patient. The most of these pharmaceutically acceptable excipients are described in detail in, *e.g.*, Allen (Ed.), **2017**. *Ansel’s pharmaceutical dosage forms and drug delivery systems* (11th ed.). Philadelphia,
15 PA: Wolters Kluwer; Remington, Allen & Adeboye (Eds.), **2013**. *Remington: The science and practice of pharmacy* (22nd ed.). London: Pharmaceutical Press; and Sheskey, Cook & Cable (Eds.), **2017**. *Handbook of pharmaceutical excipients* (8th ed.). London: Pharmaceutical Press.

20 **DETAILED DESCRIPTION**

Drug Product (DP / STC-1010)

The present invention relates to a composition comprising or consisting of (i) at least one of stressed HT-29, HCT-116 or LoVo cells, and (ii) immunogenic stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these at least one
25 of HT-29, HCT-116 or LoVo cells in response to a stress that was applied *in vitro*.

In one embodiment, the composition comprises or consists of (i) stressed HT-29 cells, and (ii) immunogenic stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these HT-29 cells in response to a stress that was applied *in vitro*.

In one embodiment, the composition comprises or consists of (i) stressed HCT-116 cells, and (ii) immunogenic stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these HCT-116 cells in response to a stress that was applied *in vitro*.

- 5 In one embodiment, the composition comprises or consists of (i) stressed LoVo cells, and (ii) immunogenic stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these LoVo cells in response to a stress that was applied *in vitro*.

- 10 In one embodiment, the composition comprises or consists of (i) stressed HT-29 and HCT-116 cells, and (ii) immunogenic stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these HT-29 and HCT-116 cells in response to a stress that was applied *in vitro*.

- 15 In one embodiment, the composition comprises or consists of (i) stressed HT-29 and LoVo cells, and (ii) immunogenic stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these HT-29 and LoVo cells in response to a stress that was applied *in vitro*.

- 20 In one embodiment, the composition comprises or consists of (i) stressed HCT-116 and LoVo cells, and (ii) immunogenic stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these HCT-116 and LoVo cells in response to a stress that was applied *in vitro*.

In one embodiment, the composition comprises or consists of (i) stressed HT-29, HCT-116 and LoVo cells, and (ii) immunogenic stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these HT-29, HCT-116 and LoVo cells in response to a stress that was applied *in vitro*.

- 25 In the following, any reference to the composition comprises HT-29, HCT-116 and LoVo cells is intended to encompass compositions comprising one, two or the three of HT-29, HCT-116 and LoVo cells.

According to the invention, the composition comprises HT-29, HCT-116 and LoVo cells which are stressed. By “**stressed**”, it is meant that these cells have developed [a] resistance mechanism[s] in response to [a] stress[es] applied *in vitro*. As a consequence, these cells have produced stress and/or resistance proteins which form part of the composition.

In one embodiment, the stress is selected from the group comprising or consisting of radiations, thermal stress, chemical stress, metabolic stress and any combinations thereof.

In one embodiment, the stress includes a combination, whether concomitant or successive, of two, three or more of radiations, thermal stress, chemical stress, and metabolic stress.

In one embodiment, the composition of the invention comprises or consists of:

- stressed HT-29 cells and stress and/or resistance proteins produced by these HT-29 cells in response to (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*;
- 15 - stressed HCT-116 cells and stress and/or resistance proteins produced by these HCT-116 cells in response to (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*; and
- stressed LoVo cells and stress and/or resistance proteins produced by these LoVo cells in response to (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*.

In one embodiment, the composition of the invention comprises or consists of:

- stressed HT-29 cells and stress and/or resistance proteins produced by these HT-29 cells in response to (i) a metabolic stress, and (ii) a chemical stress applied *in vitro*;
- stressed HCT-116 cells and stress and/or resistance proteins produced by these HCT-116 cells in response to (i) a metabolic stress, and (ii) a chemical stress applied *in vitro*; and
- 25 - stressed LoVo cells and stress and/or resistance proteins produced by these LoVo cells in response to (i) a metabolic stress, and (ii) a chemical stress applied *in vitro*.

In one embodiment, the composition of the invention comprises or consists of:

- stressed HT-29 cells and stress and/or resistance proteins produced by these HT-29 cells in response to (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*;
- stressed HCT-116 cells and stress and/or resistance proteins produced by these
5 HCT-116 cells in response to (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*;
- stressed LoVo cells and stress and/or resistance proteins produced by these LoVo cells in response to (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*;
- 10 - stressed HT-29 cells and stress and/or resistance proteins produced by these HT-29 cells in response to (i) a metabolic stress, and (ii) a chemical stress applied *in vitro*;
- stressed HCT-116 cells and stress and/or resistance proteins produced by these HCT-116 cells in response to (i) a metabolic stress, and (ii) a chemical stress applied *in vitro*; and
- 15 - stressed LoVo cells and stress and/or resistance proteins produced by these LoVo cells in response to (i) a metabolic stress, and (ii) a chemical stress applied *in vitro*.

In one embodiment, the stress is radiation. Radiation is preferably at a dose sufficiently low not to kill or inactivate the cells, but sufficiently high to induce the production of stress and/or resistance proteins. In one embodiment, radiation comprises or consists of
20 irradiating the cells with a total dose ranging from about 0.25 to about 25 Gy, preferably from about 1 to about 15 Gy, such as, *e.g.*, with a total dose of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 Gy. In one embodiment, the irradiation period ranges from about 1 to about 20 minutes, such as, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 minutes, preferably from about 1 to about 5 minutes. In one embodiment,
25 radiation comprises or consists of irradiating the cells with a total dose of 10 Gy for a period of about 1.5 to 2 minutes, *i.e.*, with a dose of about 5 to about 6.6 Gy/minute. In one embodiment, radiation comprises or consists of irradiating the cells with a total dose of 10 Gy for a period of about 5 minutes, *i.e.*, with a dose of about 2 Gy/minute.

In one embodiment, the stress is a thermal stress. Thermal stress is preferably at a
30 temperature sufficiently low not to kill or inactivate the cells, but sufficiently high to

- induce the production of stress and/or resistance proteins. In one embodiment, thermal stress comprises or consists of cultivating the cells at a temperature greater than 37°C, preferably ranging from about 38°C to about 45°C, such as, *e.g.*, at a temperature of about 38°C, 39°C, 40°C, 41°C, 42°C, 43°C, 44°C, or 45°C. In one embodiment, thermal stress is applied to the cells for a period ranging from about 15 minutes to about 4 hours, preferably from about 30 minutes to about 2 hours, such as, *e.g.*, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, or 120 minutes. In one embodiment, thermal stress comprises or consists of cultivating the cells at a temperature of about 42°C for a period of about 60 minutes.
- 10 In one embodiment, the stress is a chemical stress. Chemical stress is carried out by exposing the cells to at least one or several chemical agents preferably at doses sufficiently low not to kill or inactivate the cells, but sufficiently high to induce the production of stress and/or resistance proteins. In one embodiment, chemical stress comprises or consists of exposing the cells to at least one or several chemotherapeutic agents and/or alcohols. In one embodiment, chemical stress is applied to the cells for a period ranging from about 6 hours to about 120 hours, preferably from about 24 hours to about 96 hours, such as, *e.g.*, 24, 30, 36, 42, 48, 54, 60, 66, 72, 78, 84, 90, or 96 hours. In one embodiment, chemical stress comprises or consists of exposing the cells, preferably HT-29 cells, to oxaliplatin, preferably at a concentration ranging from about 1 µM to about 20 µM, such as, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 µM, more preferably at a concentration of about 13 µM, for a period of about 60 hours to about 84 hours, preferably for a period of about 72 hours. In one embodiment, chemical stress comprises or consists of exposing the cells, preferably HCT-116 cells, to SN-38 (7-ethyl-10-hydroxy-camptothecin), preferably at a concentration ranging from about 20 nM to about 400 nM, such as, *e.g.*, 20, 40, 60, 80, 100, 120, 140, 160, 180, 200, 220, 240, 260, 280, 300, 320, 340, 360, 380, or 400 nM, more preferably at a concentration of about 31.5 nM, 100 nM or 315 nM, for a period of about 36 hours to about 60 hours, preferably for a period of about 48 hours. In one embodiment, chemical stress comprises or consists of exposing the cells, preferably LoVo cells, to fluorouracil (5-FU), preferably at a concentration ranging from about 0.5 µM to about 15 µM, such as, *e.g.*, 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5, 9.0,

9.5, 10.0, 10.5, 11.0, 11.5, 12.0, 12.5, 13.0, 13.5, 14.0, 14.5, or 15 μ M, more preferably at a concentration of about 5 μ M, for a period of about 36 hours to about 60 hours, preferably for a period of about 48 hours.

In one embodiment, the stress is a metabolic stress. Metabolic stress is carried out by
5 cultivating the cells in a non-physiological cell culture medium that does not kill or inactivate the cells, but induces the production of stress and/or resistance proteins. In one embodiment, metabolic stress comprises or consists of cultivating the cells in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, at low pH (*e.g.*,
10 below pH 6.5). In one embodiment, metabolic stress comprises or consists of cultivating the cells in low serum culture conditions, such as, *e.g.*, in a 2 % FBS culture medium (instead of a 10 % FBS culture medium as classically used).

In one embodiment, the composition of the invention comprises or consists of:

- stressed HT-29 cells and stress and/or resistance proteins produced by these HT-29
15 cells in response to (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium, (ii) an *in vitro* radiation with a total dose of 10 Gy for a period of about 1 to about 5 minutes, and (iii) an *in vitro* thermic choc at a temperature of about 42°C for a period of about 60 minutes;
- stressed HCT-116 cells and stress and/or resistance proteins produced by these
20 HCT-116 cells in response to (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium, (ii) an *in vitro* radiation with a total dose of 10 Gy for a period of about 1 to about 5 minutes, and (iii) an *in vitro* thermic choc at a temperature of about 42°C for a period of about 60 minutes;
- stressed LoVo cells and stress and/or resistance proteins produced by these LoVo
25 cells in response to (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium, (ii) an *in vitro* radiation with a total dose of 10 Gy for a period of about 1 to about 5 minutes, and (iii) an *in vitro* thermic choc at a temperature of about 42°C for a period of about 60 minutes;
- stressed HT-29 cells and stress and/or resistance proteins produced by these HT-29
30 cells in response to (i) an *in vitro* culture in low serum culture conditions in a 2 %

- FBS culture medium, and (ii) an *in vitro* exposition to about 13 μ M oxaliplatin for a period of about 72 hours;
- stressed HCT-116 cells and stress and/or resistance proteins produced by these HCT-116 cells in response to (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium, and (ii) an *in vitro* exposition to about 31.5 nM, 100 nM or 315 nM SN-38 (7-ethyl-10-hydroxy-camptothecin) for a period of about 48 hours; and
 - stressed LoVo cells and stress and/or resistance proteins produced by these LoVo cells in response to (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium, and (ii) an *in vitro* exposition to about 5 μ M fluorouracil (5-FU) for a period of about 48 hours.

In one embodiment, the composition comprises HT-29, HCT-116 and LoVo cells which are non-proliferative. By “**non-proliferative**” or “**inactive**”, it is meant that these cells are not capable of cell proliferation, that is, the process by which a cell grows and divides to produce two daughter cells.

The skilled artisan is well aware of means and methods for rendering cells non-proliferative. By way of examples, cells can be rendered non-proliferative by radiation with a dose sufficiently high to kill or inactivate the cells. In one embodiment, radiation – to render the cells non-proliferative – comprises or consists of irradiating the cells with a total dose of 25 Gy or above. Alternatively or additionally, cells can be rendered non-proliferative by ethanol fixation, *e.g.*, using from about 10 % to about 50 % v/v of ethanol. Alternatively or additionally, cells can be rendered non-proliferative by at least one freeze-thaw cycle. Alternatively or additionally, cells can be rendered non-proliferative by linkage to or by complexation with a means capable to confer immunogenicity, such as, *e.g.*, by haptentation.

The skilled artisan is also familiar with means and methods to determine whether a cell is non-proliferative or not, *e.g.*, by carrying out viability tests by cell culture (to assess the total lack of proliferation) and/or propidium iodide (which distinguishes between living cells and dead cells).

In one embodiment, non-proliferative cells are structurally intact, *i.e.*, they display an intact plasma membrane. In one embodiment, non-proliferative cells are non-structurally intact, *i.e.*, they do not display an intact plasma membrane. In the latter case, cells are presents in the composition in the form of membrane fragments, organelles and other
5 cytoplasm constituents.

In one embodiment, cells have been rendered non-proliferative after having developed their resistance mechanism[s] in response to the stress[es] applied *in vitro* and therefore, after having produced stress and/or resistance proteins. In one embodiment, cells have been rendered non-proliferative several hours after having been stressed, preferably more
10 than 12, 24, 36, 48, 60, 72, 84, 96 hours or more after having been stressed.

In one embodiment, the composition comprises stressed HT-29, HCT-116 and LoVo cells which overexpress at least one, at least two, at least three or the four following markers: Cmhsp70.1 (HSP70), CD227 (MUC1), CD95 (FAS receptor) and/or CD243 (MDR-1).

In one embodiment, the composition comprises stressed HT-29, HCT-116 and LoVo cells
15 which express at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more of at least one, at least two, at least three or the four following markers: Cmhsp70.1 (HSP70), CD227 (MUC1), CD95 (FAS receptor) and/or CD243 (MDR-1); as compared to HT-29, HCT-116 and/or LoVo cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress,
20 (ii) radiations, (iii) a thermal stress and (iv) a chemical stress applied *in vitro*; and/or as compared to HT-29, HCT-116 and/or LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations, (ii) a thermal stress and (iii) a chemical stress applied *in vitro*. Additionally or alternatively, at least one, at least two, at least three or the four markers Cmhsp70.1 (HSP70), CD227 (MUC1), CD95 (FAS
25 receptor) and/or CD243 (MDR-1) are expressed in at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more cells in the composition than in a population of HT-29, HCT-116 and/or LoVo cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations, (iii) a thermal stress and (iv) a chemical stress applied *in vitro*; and/or than
30 in a population of HT-29, HCT-116 and/or LoVo cells subjected to a metabolic stress,

e.g., cultured in 2 % FBS but not subjected to any of (i) radiations, (ii) a thermal stress and (iii) a chemical stress applied *in vitro*. Additionally or alternatively, the mean fluorescence intensity (MFI) for at least one, at least two, at least three or the four markers Cmhsp70.1 (HSP70), CD227 (MUC1), CD95 (FAS receptor) and/or CD243 (MDR-1) is
5 at least twice higher, preferably at least 3 times, 4 times, 5 times, 6 times or more higher in the composition than in a population of HT-29, HCT-116 and/or LoVo cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations, (iii) a thermal stress and (iv) a chemical stress applied *in vitro*; and/or than in a population of HT-29, HCT-116 and/or LoVo cells subjected to
10 a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations, (ii) a thermal stress and (iii) a chemical stress applied *in vitro*.

According to the invention, the composition comprises stress and/or resistance proteins.

In one embodiment, these stress and/or resistance proteins are associated with the membrane of the HT-29, HCT-116 and LoVo cells, *i.e.*, exposed at the surface, whether
15 outer or inner surface, of the HT-29, HCT-116 and LoVo cells; and/or are contained within the HT-29, HCT-116 and LoVo cells, *i.e.*, contained within their cytoplasm or any of their organelles; and/or are in a free state in the composition, *i.e.*, not associated with the membrane of nor contained within the HT-29, HCT-116 and LoVo cells, *e.g.*, because they were secreted by these cells.

20 According to the invention, the composition comprises stress and/or resistance proteins which are immunogenic. By “**immunogenic**” or “**immunocompetent**”, it is meant that the stress and/or resistance proteins are capable of eliciting an immune response (*e.g.*, the production of antibodies) in a patient when administered to said patient.

In one embodiment, the stress and/or resistance proteins have been rendered
25 immunogenic in the presence of a molecule or by a process capable of rendering them immunogenic. In one embodiment, the stress and/or resistance proteins have been rendered immunogenic by linkage to or by complexation with a means capable to confer immunogenicity.

Means and processes capable of conferring immunogenicity are well known to the skilled artisan. In one embodiment, the means capable of conferring immunogenicity comprise or consist of one or several molecule[s] that is/are not naturally present in the HT-29, HCT-116 and LoVo cells or in their environment. Such means are commonly referred to as “antigens”. Some antigens can trigger a humoral or cell-mediated immune response by themselves, and are then considered as “immunogens”. Some examples of such antigens include, but are not limited to, adjuvants. The skilled artisan is well aware of adjuvants suitable for use in cancer vaccines, which, when complexed with the stress and/or resistance proteins, render them immunogenic. See, *e.g.*, Dubensky & Reed, **2010** (5) (*Semin Immunol.* **22**(3):155-61) or Cuzzubbo *et al.*, **2021** (*Front Immunol.* **11**:615240). 10

However, some antigens cannot initiate an immune response by themselves and require to be conjugated beforehand to a carrier, *e.g.*, the stress and/or resistance proteins, to become immunogenic. The latter antigens are sometimes referred to as “incomplete antigens” and include, but are not limited to, haptens.

15 Some examples of haptens include, but are not limited to, 2,4-dinitrophenyl (DNP); 2,4-dinitrofluorobenzene; sulfanilic acid; *N*-iodoacetyl-*N'*-(5-sulfonic-naphthyl)ethylene diamine; anilin; *p*-amino benzoic acid; biotin; fluorescein and derivatives thereof (including FITC, TAMRA, and Texas Red); digoxigenin; 5-nitro-3-pyrazolecarbamide; 4,5-dimethoxy-2-nitrocinnamide; 2-(3,4-dimethoxyphenyl)-quinoline-4-carbamide; 20 2,1,3-benzoxadiazole-5-carbamide; 3-hydroxy-2-quinoxalinecarbamide, 4-(dimethylamino)azobenzene-4'-sulfonamide (DABSYL); rotenone isooxazoline; (*E*)-2-(2-(2-oxo-2,3-dihydro-1*H*-benzo[*b*][1,4]diazepin-4-yl)phenoxy)acetamide; 7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylic acid; 2-acetamido-4-methyl-5-thiazolesulfonamide; and 25 *p*-methoxyphenylpyrazopodophyllamide.

In one embodiment, stress and/or resistance proteins are haptenated. In one embodiment, stress and/or resistance proteins are 2,4-dinitrophenylated (DNP).

The skilled artisan is well aware of means and methods to haptinize proteins such as stress and/or resistance proteins.

Another means capable of conferring immunogenicity includes opsonization, which is a well-known process involving the binding of an opsonin to the stress and/or resistance proteins.

In one embodiment, stress and/or resistance proteins are selected from the group comprising or consisting of radiation resistance proteins, thermal stress resistance
5 proteins, chemical stress resistance proteins, metabolic stress resistance proteins, and combination thereof.

In one embodiment, the composition specifically comprises, to detectable levels, at least one or several of the following proteins, as compared to any of HT-29, HCT-116 or LoVo
10 cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations (iii) a thermal stress and (iv) a chemical stress applied *in vitro*; and/or as compared to any of HT-29, HCT-116 or LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations, (ii) a thermal stress and (iii) a chemical stress applied *in vitro*: tyrosine-protein kinase
15 HCK, polypyrimidine tract-binding protein 3, E3 ubiquitin-protein ligase RNF213, serine/arginine-rich splicing factor 8, EH domain-containing protein 4, LIM domain only protein 7, DNA-directed RNA polymerase I subunit RPA2, 2'-5'-oligoadenylate synthase 3, WD repeat and HMG-box DNA-binding protein 1, beta-2-glycoprotein 1, serine/threonine-protein phosphatase PP1-gamma catalytic subunit, anillin,
20 unconventional myosin-Ib, AP-2 complex subunit alpha-2, cyclin-dependent kinase 2, signal transducer and activator of transcription 1-alpha/beta, pumilio homolog 1, ATP-binding cassette sub-family F member 1, Rac GTPase-activating protein 1, cingulin, syntaxin-binding protein 3, mitochondrial carnitine/acylcarnitine carrier protein, importin subunit alpha-7, ribosomal protein S6 kinase alpha-4, Ras-related protein Rab-5A,
25 ribonucleoside-diphosphate reductase large subunit, low molecular weight phosphotyrosine protein phosphatase, ribonucleoside-diphosphate reductase subunit M2, ADP-ribosylation factor-like protein 1, dynamin-2, Ras-related protein Rab-13, IST1 homolog, Forkhead box protein K1, sorbitol dehydrogenase, Bcl-2-like protein 1, tripartite motif-containing protein 29, kinesin-like protein KIF22, methylsterol
30 monooxygenase 1, caveolae-associated protein 1, BRCA1-associated ATM activator 1,

- protein FAM83H, protein O-mannosyl-transferase TMTC3, inhibitor of nuclear factor kappa-B kinase-interacting protein, zinc finger CCCH-type antiviral protein 1, nucleolar protein 9, leucine zipper protein 1, polyhomeotic-like protein 2, tensin-4, LEM domain-containing protein 2, importin-4, Rho guanine nucleotide exchange factor 1, PDZ
- 5 and LIM domain protein 5, UAP56-interacting factor, MMS19 nucleotide excision repair protein homolog, kinesin-like protein KIF2C, ADP-ribose glycohydrolase MACROD1, kinesin-like protein KIFC1, ATP-dependent RNA helicase DHX33, echinoderm microtubule-associated protein-like 4, fanconi anemia group I protein, EH domain-containing protein 3, opioid growth factor receptor, gamma-adducin, DNA
- 10 dC-dU-editing enzyme APOBEC-3B, neuroplastin, zinc transporter 1, glutaredoxin-3, cytosolic thymidine kinase, myristoylated alanine-rich C-kinase substrate, UMP-CMP kinase, macrophage-capping protein, high mobility group protein HMGI-C, smoothelin, platelet-activating factor acetylhydrolase IB subunit gamma, transcription initiation factor TFIID subunit 9, inactive hydroxysteroid dehydrogenase-like protein 1, borealin,
- 15 la-related protein 4, NADH dehydrogenase [ubiquinone] 1 alpha subcomplex assembly factor 2, aurora kinase B, golgi-associated plant pathogenesis-related protein 1, ATP-binding cassette sub-family F member 2, armadillo repeat-containing X-linked protein 3, regulator complex protein LAMTOR3, probable ATP-dependent RNA helicase DDX20, V-type proton ATPase subunit H, and calcium-binding protein 39.
- 20 In one embodiment, the composition overexpresses at least one or several of the following proteins, as compared to any of HT-29, HCT-116 or LoVo cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations (iii) a thermal stress and (iv) a chemical stress applied *in vitro*; and/or as compared to any of HT-29, HCT-116 or LoVo cells subjected to a metabolic stress, *e.g.*,
- 25 cultured in 2 % FBS but not subjected to any of (i) radiations, (ii) a thermal stress and (iii) a chemical stress applied *in vitro*: HLA class I histocompatibility antigen B alpha chain, HLA class I histocompatibility antigen A alpha chain, HLA class I histocompatibility antigen C alpha chain, CD9 antigen, nuclear autoantigen Sp-100, ATP-binding cassette sub-family D member 3, ATP-binding cassette sub-family E
- 30 member 1, ATP-binding cassette sub-family F member 1, ATP-binding cassette sub-family F member 2, Bcl-2-like protein 1, Bcl-2-associated transcription factor 1,

- cytochrome c oxidase subunit 2, mitochondrial cytochrome c oxidase subunit 4 isoform 1, mitochondrial cytochrome c oxidase subunit 5A, peroxisomal acyl-coenzyme A oxidase 1, heat shock protein beta-1, heat shock protein HSP 90-beta, heat shock cognate 71 kDa protein, heat shock 70 kDa protein 6, heat shock 70 kDa protein 4, ribonuclease inhibitor, inhibitor of nuclear factor kappa-B kinase-interacting protein, Ras-related protein Rab-5A, Ras-related protein Rab-6A, Ras-related protein Rab-5C, Ras-related protein Rab-7a, Ras-related protein Rab-13, Ras-related protein Rab-25, Ras-related protein Rab-15, Ras-related protein Rab-8A, Ras-related protein Rab-10, Ras-related protein Rap-1b, Ras-related protein Rab-1A, Ras-related protein Rap-1A, Ras-related C3 botulinum toxin substrate 1, Ras-related protein Rab-8B, Ras-related protein Rab-18, Ras-related protein Rap-2c, X-ray repair cross-complementing protein 6, DNA mismatch repair protein Msh6, MMS19 nucleotide excision repair protein homolog, protein transport protein Sec16A, sodium/potassium-transporting ATPase subunit alpha-1, facilitated glucose transporter member 1 solute carrier family 2, mitochondrial tricarboxylate transport protein, monocarboxylate transporter 1, protein transport protein Sec61 subunit beta, protein transport protein Sec61 subunit alpha isoform 1, transport and Golgi organization protein 1 homolog, adenosine 3'-phospho 5'-phosphosulfate transporter 1, transportin-1, and zinc transporter 1.
- 20 In one embodiment, the composition overexpresses at least one or several of the following membrane proteins, as compared to any of HT-29, HCT-116 or LoVo cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations (iii) a thermal stress and (iv) a chemical stress applied *in vitro*; and/or as compared to any of HT-29, HCT-116 or LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations, (ii) a thermal stress and (iii) a chemical stress applied *in vitro*: HLA class I histocompatibility antigen B alpha chain, HLA class I histocompatibility antigen A alpha chain, HLA class I histocompatibility antigen C alpha chain, CD9 antigen, ATP-binding cassette sub-family D member 3, Bcl-2-like protein 1, cytochrome c oxidase subunit 2, mitochondrial cytochrome c oxidase subunit 4 isoform 1, inhibitor of nuclear factor kappa-B kinase-interacting protein, sodium/potassium-transporting ATPase subunit

alpha-1, facilitated glucose transporter member 1 solute carrier family 2, mitochondrial tricarboxylate transport protein, monocarboxylate transporter 1, protein transport protein Sec61 subunit beta, protein transport protein Sec61 subunit alpha isoform 1, transport and Golgi organization protein 1 homolog, adenosine 3'-phospho 5'-phosphosulfate transporter 1, and zinc transporter 1.

In one embodiment, the composition specifically comprises, to detectable levels, at least one or several of the following proteins, as compared to any of HT-29, HCT-116 or LoVo cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations (iii) a thermal stress and (iv) a chemical stress applied *in vitro*; and/or as compared to any of HT-29, HCT-116 or LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations, (ii) a thermal stress and (iii) a chemical stress applied *in vitro*: ATP-binding cassette sub-family F member 1, ATP-binding cassette sub-family F member 2, Bcl-2-like protein 1, inhibitor of nuclear factor kappa-B kinase-interacting protein, Ras-related protein Rab-5A, Ras-related protein Rab-13, MMS19 nucleotide excision repair protein homolog, and zinc transporter 1.

In one embodiment, the composition specifically comprises, to detectable levels, at least one or several of the following membrane proteins, as compared to any of HT-29, HCT-116 or LoVo cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations (iii) a thermal stress and (iv) a chemical stress applied *in vitro*; and/or as compared to any of HT-29, HCT-116 or LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations, (ii) a thermal stress and (iii) a chemical stress applied *in vitro*: Bcl-2-like protein 1, inhibitor of nuclear factor kappa-B kinase-interacting protein, and zinc transporter 1.

In one embodiment, the composition may further comprise tumor-associated antigens (TAA) and/or tumor-specific antigens (TSA).

By “**tumor antigen**”, it is meant an antigenic substance or molecule produced by cancer cells. Tumor antigens are classified into two categories: “**tumor-specific antigens**” or

“TSA”, which are present only on cancer cells but not on non-cancer cells; and “**tumor-associated antigens**” or “TAA”, which are present on cancer and non-cancer cells.

Among common TSA are, in particular, neoantigens. As used herein, a “**neoantigen**”
5 refers to an aberrant tumor-specific antigen which is encoded in cancer cells by genes comprising one or several mutations, *e.g.*, caused by genetic instability during carcinogenesis. As a consequence, the amino acid sequence encoded by this mutated gene may itself comprise mutations leading to the production of abnormal proteins that are not
10 found in normal cells. These mutated proteins, which may be considered as non-self protein or foreign proteins, can then be recognized by neoantigen-specific T cell receptors, activate the immune system, and lead to the immune system’s attack on cancer cells. Neoantigens can also be produced by viral infection, alternative splicing and/or gene rearrangement. See, *e.g.*, Zhang *et al.* (2021. *Front Immunol.* **12**:672356) or Jiang *et al.* (2019. *J Hematol Oncol.* **12**(1):93).

15 In one embodiment, TAA and/or TSA are specific of the HT-29, HCT-116 and LoVo cells.

In one embodiment, TAA and/or TSA are naturally immunogenic. In one embodiment, TAA and/or TSA may be rendered immunogenic, or their immunogenicity may be increased, by a molecule or a process capable of rendering them immunogenic.

20 Means capable to confer immunogenicity are well known to the skilled artisan and have been described above. These include, *inter alia* but without limitation, haptens.

In one embodiment, these TAA and/or TSA are associated with the membrane of the HT-29, HCT-116 and LoVo cells, *i.e.*, exposed at the surface, whether outer or inner surface, of the HT-29, HCT-116 and LoVo cells; and/or are contained within the HT-29,
25 HCT-116 and LoVo cells, *i.e.*, contained within their cytoplasm or any of their organelles; and/or are in a free-state in the composition, *i.e.*, not associated with the membrane of nor contained within the HT-29, HCT-116 and LoVo cells, *e.g.*, because they were secreted by these cells.

In one embodiment, the composition is a pharmaceutical composition or a vaccine composition, and further comprises at least one pharmaceutically acceptable excipient.

Pharmaceutically acceptable excipients include, but are not limited to, water, saline, Ringer's solution, dextrose solution, and solutions of ethanol, glucose, sucrose, dextran, mannose, mannitol, sorbitol, polyethylene glycol (PEG), phosphate, acetate, gelatin, collagen, Carbopol[®], vegetable oils, and the like. One may additionally include suitable preservatives, stabilizers, antioxidants, antimicrobials, and buffering agents, such as, *e.g.*, BHA, BHT, citric acid, ascorbic acid, tetracycline, and the like.

Other examples of pharmaceutically acceptable excipients that may be used in the composition of the invention include, but are not limited to, ion exchangers, alum such as aluminum phosphate or aluminium hydroxide, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

In addition, some pharmaceutically acceptable excipients may include surfactants (*e.g.*, hydroxypropylcellulose); carriers, such as, *e.g.*, solvents and dispersion media containing, *e.g.*, water, ethanol, polyol (*e.g.*, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils, such as, *e.g.*, peanut oil and sesame oil; isotonic agents, such as, *e.g.*, sugars or sodium chloride; coating agents, such as, *e.g.*, lecithin; agents delaying absorption, such as, *e.g.*, aluminum monostearate and gelatin; preservatives, such as, *e.g.*, benzalkonium chloride, benzethonium chloride, chlorobutanol, thimerosal and the like; buffers, such as, *e.g.*, boric acid, sodium and potassium bicarbonate, sodium and potassium borates, sodium and potassium carbonate, sodium acetate, sodium biphosphate and the like; tonicity agents, such as, *e.g.*, dextran 40, dextran 70, dextrose, glycerin, potassium chloride, propylene glycol, sodium chloride; antioxidants and stabilizers, such as, *e.g.*, sodium

bisulfite, sodium metabisulfite, sodium thiosulfite, thiourea and the like; nonionic wetting or clarifying agents, such as, *e.g.*, polysorbate 80, polysorbate 20, poloxamer 282 and tyloxapol; viscosity modifying agents, such as, *e.g.*, dextran 40, dextran 70, gelatin, glycerin, hydroxyethylcellulose, hydroxymethylpropylcellulose, lanolin, methylcellulose, petrolatum, polyethylene glycol, polyvinyl alcohol, polyvinylpyrrolidone, carboxymethylcellulose; and the like.

Examples of adjuvants include, but are not limited to, helper peptide, aluminum salts (such as, *e.g.*, aluminum hydroxide (also named alum), aluminum phosphate, and the like), Freund's incomplete adjuvant, Freund's complete adjuvant, saponin, Merck adjuvant 65, Smith-Kline Beecham adjuvant AS-2, Aquilla adjuvant QS-21, MPL™ immunostimulant, 3d-MPL, LEIF, calcium salts, iron salts, zinc salts, acylated tyrosine, acylated sugars, cationically-derivatized polysaccharides, anionically-derivatized polysaccharides, polyphosphazenes, biodegradable microspheres, monophosphoryl lipid A, muramyl tripeptide phosphatidyl ethanolamine, cytokines (such as, *e.g.*, interleukin-2, interleukin-12, interleukin-4, interleukin-7 and the like), CpG-containing oligonucleotide, and combinations thereof.

In one embodiment, the composition comprises from about 10^5 to about 10^8 stressed HT-29, HCT-116 and LoVo cells, preferably from about 10^6 to about 10^7 stressed HT-29, HCT-116 and LoVo cells, such as, about 1×10^6 , 2×10^6 , 3×10^6 , 4×10^6 , 5×10^6 , 6×10^6 , 7×10^6 , 8×10^6 , 9×10^6 , or 1×10^7 stressed HT-29, HCT-116 and LoVo cells. In one embodiment, the composition comprises about 3×10^6 stressed HT-29, HCT-116 and LoVo cells.

In one embodiment, the composition comprises from about 10^6 to about 10^9 stressed HT-29, HCT-116 and LoVo cells per mL of composition, preferably from about 10^7 to about 10^8 stressed HT-29, HCT-116 and LoVo cells per mL of composition, such as, about 1×10^7 , 2×10^7 , 3×10^7 , 4×10^7 , 5×10^7 , 6×10^7 , 7×10^7 , 8×10^7 , 9×10^7 , or 1×10^8 stressed HT-29, HCT-116 and LoVo cells per mL of composition. In one embodiment, the composition comprises about 3×10^7 stressed HT-29, HCT-116 and LoVo cells per mL of composition.

In one embodiment, the composition comprises an equal ratio of stressed HT-29, HCT-116 and LoVo cells (*i.e.*, about 1:1:1). In one embodiment, the composition comprises at least 1.5, 2 or 2.5 times more stressed HT-29 than stressed HCT-116 or LoVo cells. In one embodiment, the composition comprises at least 1.5, 2 or 2.5 times more stressed HCT-116 than stressed HT-29 or LoVo cells. In one embodiment, the composition comprises at least 1.5, 2 or 2.5 times more stressed LoVo than stressed HT-29 or HCT-116 cells.

In one embodiment, the composition comprises:

- from about 10^5 to about 10^8 stressed HT-29 cells, preferably from about 10^6 to about 10^7 stressed HT-29 cells, such as, about 1×10^6 , 2×10^6 , 3×10^6 , 4×10^6 , 5×10^6 , 6×10^6 , 7×10^6 , 8×10^6 , 9×10^6 , or 1×10^7 stressed HT-29 cells,
- from about 10^5 to about 10^8 stressed HCT-116 cells, preferably from about 10^6 to about 10^7 stressed HCT-116 cells, such as, about 1×10^6 , 2×10^6 , 3×10^6 , 4×10^6 , 5×10^6 , 6×10^6 , 7×10^6 , 8×10^6 , 9×10^6 , or 1×10^7 stressed HCT-116 cells, and
- from about 10^5 to about 10^8 stressed LoVo cells, preferably from about 10^6 to about 10^7 stressed LoVo cells, such as, about 1×10^6 , 2×10^6 , 3×10^6 , 4×10^6 , 5×10^6 , 6×10^6 , 7×10^6 , 8×10^6 , 9×10^6 , or 1×10^7 stressed LoVo cells.

Intermediate products

The present invention also relates to intermediate compositions useful in the preparation of the composition described above.

In one embodiment, the intermediate compositions comprise at least one pharmaceutically acceptable excipient, as defined above.

In one embodiment, the intermediate composition comprises or consists of (i) stressed HT-29, HCT-116 or LoVo cells, and (ii) stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these HT-29, HCT-116 or LoVo cells in response to (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*. Radiations, thermal stresses and metabolic stresses have been described above.

In one embodiment, the stress and/or resistance proteins are immunogenic. Means and methods for rendering stress and/or resistance proteins immunogenic have been described above.

5 In one embodiment, the intermediate composition may further comprise tumor-associated antigens (TAA) and/or tumor-specific antigens (TSA), as described above. In one embodiment, TAA and/or TSA are specific of the HT-29, HCT-116 or LoVo cells. In one embodiment, TAA and/or TSA are naturally immunogenic. In one embodiment, TAA and/or TSA may be rendered immunogenic, or their immunogenicity may be increased, by linkage to or by complexation with a means capable to confer immunogenicity. Means
10 capable to confer immunogenicity are well known to the skilled artisan and have been described above. These include, *inter alia* but without limitation, haptens.

In one embodiment, the intermediate composition comprises or consists of (i) stressed HT-29 cells and (ii) stress and/or resistance proteins produced by these HT-29 cells in response to:

- 15 (i) an *in vitro* culture in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, and/or at low pH (*e.g.*, below pH 6.5),
- (ii) an *in vitro* radiation with a total dose ranging from about 0.25 to about 25 Gy, preferably from about 1 to about 15 Gy, such as, *e.g.*, with a total dose of 1, 2, 3, 4,
20 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 Gy. In one embodiment, the irradiation period ranges from about 1 to about 20 minutes, such as, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 minutes, preferably from about 1 to about 5 minutes, and
- (iii) an *in vitro* thermic choc at a temperature ranging from about 38°C to about 45°C,
25 such as, *e.g.*, at a temperature of about 38°C, 39°C, 40°C, 41°C, 42°C, 43°C, 44°C, or 45°C, applied to the cells for a period ranging from about 15 minutes to about 4 hours, preferably from about 30 minutes to about 2 hours, such as, *e.g.*, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, or 120 minutes.

In one embodiment, the intermediate composition comprises or consists of (i) stressed HT-29 cells and (ii) stress and/or resistance proteins produced by these HT-29 cells in response to

- (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium,
- 5 (ii) an *in vitro* radiation with a total dose of 10 Gy for a period of about 1 to about 5 minutes, and
- (iii) an *in vitro* thermic choc at a temperature of about 42°C for a period of about 60 minutes.

This intermediate composition is herein referred to as “**HT-29 DS-A**”.

- 10 In one embodiment, HT-29 DS-A comprises stress and/or resistance proteins selected from the group comprising or consisting of radiation resistance proteins, thermal stress resistance proteins, metabolic stress resistance proteins, and combination thereof.

In one embodiment, HT-29 DS-A cells overexpress at least one, at least two or the three following markers: Cmhsp70.1 (HSP70), CD227 (MUC1) and/or CD107 (LAMP-1).

- 15 In one embodiment, HT-29 DS-A cells express at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more of at least one, at least two or the three following markers: Cmhsp70.1 (HSP70), CD227 (MUC1) and/or CD107 (LAMP-1); as compared to HT-29 cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*; and/or as compared to HT-29 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*. Additionally or alternatively, at least one, at least two or the three markers Cmhsp70.1 (HSP70), CD227 (MUC1) and CD107 (LAMP-1) are expressed in at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 %
20 or more cells in a population of HT-29 DS-A cells than in a population of HT-29 cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*; and/or than in a population of HT-29 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*.

Additionally or alternatively, the mean fluorescence intensity (MFI) for at least one, at least two or the three markers Cmhsp70.1 (HSP70), CD227 (MUC1) and/or CD107 (LAMP-1) is at least twice higher, preferably at least 3 times, 4 times, 5 times, 6 times or more higher in a population of HT-29 DS-A cells than in a population of HT-29 cells
5 cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*; and/or than in a population of HT-29 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*.

In one embodiment, HT-29 DS-A cells overexpress at least one or several of the following
10 proteins, as compared to HT-29 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*: poly(rC)-binding protein 2, rho guanine nucleotide exchange factor 1, golgin subfamily B member 1, beta-2-glycoprotein 1, plakophilin-3, protein FAM83H, protein transport protein Sec16A, inverted formin-2, anillin, protein ECT2, plectin, epiplakin, proliferation
15 marker protein Ki-67, vesicular integral-membrane protein VIP36, and lysosome-associated membrane glycoprotein 1.

In one embodiment, HT-29 DS-A cells specifically express, to detectable levels, at least one or several of the following proteins, as compared to HT-29 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and
20 (ii) a thermal stress applied *in vitro*: poly(rC)-binding protein 2, rho guanine nucleotide exchange factor 1, golgin subfamily B member 1, beta-2-glycoprotein 1, plakophilin-3, protein FAM83H, protein transport protein Sec16A, inverted formin-2, anillin, and protein ECT2.

In one embodiment, HT-29 DS-A cells overexpress at least one or several of the following
25 membrane and/or cell surface proteins, as compared to HT-29 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*: vesicular integral-membrane protein VIP36, lysosome-associated membrane glycoprotein 1, ribosome-binding protein 1, Kunitz-type protease inhibitor 2, lysophospholipid acyltransferase 5, emerin, protein LYRIC,

elongation of very long chain fatty acids protein 1, and protein transport protein Sec61 subunit gamma.

- In one embodiment, HT-29 DS-A cells overexpress at least one or several of the following cell surface proteins, as compared to HT-29 cells subjected to a metabolic stress, *e.g.*,
 5 cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*: heat shock-related 70 kDa protein, annexin, anoctamin-6, immunoglobulin superfamily member 3, serotransferrin, tumor necrosis factor receptor superfamily member 10B, clusterin, furin, HLA class II histocompatibility antigen gamma chain, CD109 antigen, chloride intracellular channel protein 4, protocadherin
 10 fat 1, Natural resistance-associated macrophage protein 2, tumor necrosis factor receptor superfamily member 10A, calpain-5, MHC class I polypeptide-related sequence A, high mobility group protein B1, tetraspanin-15, UL16-binding protein 2, integrin beta-7, sonic hedgehog protein, toll-like receptor 3, beta-2-glycoprotein 1, tissue factor, proprotein convertase subtilisin/kexin type 6, endothelial protein C receptor, volume-regulated
 15 anion channel subunit LRRC8A, cadherin EGF LAG seven-pass G-type receptor 3, zinc transporter ZIP6, HLA class II histocompatibility antigen DM alpha chain, cystine/glutamate transporter, lysophosphatidic acid receptor 2, syndecan-1, hyaluronidase-2, integrin alpha-4, histidine-rich glycoprotein, transforming growth factor beta-1 proprotein, and metalloproteinase inhibitor 2.
- 20 In one embodiment, the intermediate composition comprises or consists of (i) stressed HCT-116 cells and (ii) stress and/or resistance proteins produced by these HCT-116 cells in response to:
- (i) an *in vitro* culture in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival
 25 of cancer cells), under hypoxia, and/or at low pH (*e.g.*, below pH 6.5),
 - (ii) an *in vitro* radiation with a total dose ranging from about 0.25 to about 25 Gy, preferably from about 1 to about 15 Gy, such as, *e.g.*, with a total dose of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 Gy. In one embodiment, the irradiation period ranges from about 1 to about 20 minutes, such as, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10,

11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 minutes, preferably from about 1 to about 5 minutes, and

- (iii) an *in vitro* thermic choc at a temperature ranging from about 38°C to about 45°C, such as, *e.g.*, at a temperature of about 38°C, 39°C, 40°C, 41°C, 42°C, 43°C, 44°C, or 45°C, applied to the cells for a period ranging from about 15 minutes to about 4 hours, preferably from about 30 minutes to about 2 hours, such as, *e.g.*, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, or 120 minutes.

In one embodiment, the intermediate composition comprises or consists of (i) stressed HCT-116 cells and (ii) stress and/or resistance proteins produced by these HCT-116 cells in response to

- (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium,
 (ii) an *in vitro* radiation with a total dose of 10 Gy for a period of about 1 to about 5 minutes, and
 (iii) an *in vitro* thermic choc at a temperature of about 42°C for a period of about 60 minutes.

This intermediate composition is herein referred to as “**HCT-116 DS-A**”.

In one embodiment, HCT-116 DS-A comprises stress and/or resistance proteins selected from the group comprising or consisting of radiation resistance proteins, thermal stress resistance proteins, metabolic stress resistance proteins, and combination thereof.

- 20 In one embodiment, HCT-116 DS-A cells overexpress at least one or the two following markers: Cmhs70.1 (HSP70) and/or CD227 (MUC1).

In one embodiment, HCT-116 DS-A cells express at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more of at least one or the two following markers: Cmhs70.1 (HSP70) and/or CD227 (MUC1); as compared to
 25 HCT-116 cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*; and/or as compared to HCT-116 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*. Additionally or alternatively, at least one or the two markers Cmhs70.1

(HSP70) and CD227 (MUC1) are expressed in at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more cells in a population of HCT-116 DS-A cells than in a population of HCT-116 cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*; and/or than in a population of HCT-116 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*. Additionally or alternatively, the mean fluorescence intensity (MFI) for at least one or the two markers Cmhsp70.1 (HSP70) and CD227 (MUC1) is at least twice higher, preferably at least 3 times, 4 times, 5 times, 6 times or more higher in a population of HCT-116 DS-A cells than in a population of HCT-116 cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*; and/or than in a population of HCT-116 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*.

In one embodiment, HCT-116 DS-A cells overexpress at least one or several of the following proteins, as compared to HCT-116 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*: poly(rC)-binding protein 2, rho guanine nucleotide exchange factor 1, golgin subfamily B member 1, beta-2-glycoprotein 1, plakophilin-3, protein FAM83H, protein transport protein Sec16A, inverted formin-2, anillin, protein ECT2, plectin, epiplakin, proliferation marker protein Ki-67, vesicular integral-membrane protein VIP36, and lysosome-associated membrane glycoprotein 1.

In one embodiment, HCT-116 DS-A cells specifically express, to detectable levels, at least one or several of the following proteins, as compared to HCT-116 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*: poly(rC)-binding protein 2, rho guanine nucleotide exchange factor 1, golgin subfamily B member 1, beta-2-glycoprotein 1, plakophilin-3, protein FAM83H, protein transport protein Sec16A, inverted formin-2, anillin, and protein ECT2.

In one embodiment, HCT-116 DS-A cells overexpress at least one or several of the following membrane and/or cell surface proteins, as compared to HCT-116 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*: integrin alpha-6,

5 lysosome-associated membrane glycoprotein 1, ribosome-binding protein 1, delta(14)-sterol reductase LBR, mitochondrial proton/calcium exchanger protein, RRP12-like protein, extended syptotagmin-1, protein transport protein Sec61 subunit alpha isoform 1, solute carrier family 2, facilitated glucose transporter member 1, desmoglein-2, kinectin, protein LYRIC, sphingosine-1-phosphate lyase 1,

10 vesicle-associated membrane protein-associated protein A, lysophospholipid acyltransferase 7, Sigl recognition particle receptor subunit beta, torsin-1A-interacting protein 1, ER membrane protein complex subunit 1, decarboxylating sterol-4-alpha-carboxylate 3-dehydrogese, membrane-associated progesterone receptor component 1, CD44 antigen, MICOS complex subunit MIC26, transmembrane emp24

15 domain-containing protein 4, ORM1-like protein 2, dolichol-phosphate mannosyltransferase subunit 3, cytoskeleton-associated protein 4, transmembrane protein 43, retinol dehydrogese 11, Sigl peptidase complex subunit 2, serine palmitoyltransferase 2, coiled-coil domain-containing protein 47, very-long-chain enoyl-CoA reductase, protocadherin Fat 1, transmembrane emp24 domain-containing

20 protein 7, caveolin-1, NF-X1-type zinc finger protein NFXL1, receptor expression-enhancing protein 6, mitochondrial HIG1 domain family member 2A, small integral membrane protein 20, DH dehydrogese [ubiquinone] 1 alpha subcomplex subunit 13, DH dehydrogese [ubiquinone] 1 beta subcomplex subunit 3, mitochondrial thiamine pyrophosphate carrier, mitochondrial fission 1 protein, amine oxidase

25 [flavin-containing] B, mitochondrial inner membrane protein OXA1L, 2-hydroxyacyl-CoA lyase 2, nicalin, plasma membrane calcium-transporting ATPase 1, myoferlin, and cation-independent mannose-6-phosphate receptor.

In one embodiment, the intermediate composition comprises or consists of (i) stressed LoVo cells and (ii) stress and/or resistance proteins produced by these LoVo cells in

30 response to:

- (i) an *in vitro* culture in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, and/or at low pH (*e.g.*, below pH 6.5),
 - (ii) an *in vitro* radiation with a total dose ranging from about 0.25 to about 25 Gy, preferably from about 1 to about 15 Gy, such as, *e.g.*, with a total dose of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 Gy. In one embodiment, the irradiation period ranges from about 1 to about 20 minutes, such as, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 minutes, preferably from about 1 to about 5 minutes, and
 - (iii) an *in vitro* thermic choc at a temperature ranging from about 38°C to about 45°C, such as, *e.g.*, at a temperature of about 38°C, 39°C, 40°C, 41°C, 42°C, 43°C, 44°C, or 45°C, applied to the cells for a period ranging from about 15 minutes to about 4 hours, preferably from about 30 minutes to about 2 hours, such as, *e.g.*, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, or 120 minutes.
- 15 In one embodiment, the intermediate composition comprises or consists of (i) stressed LoVo cells and (ii) stress and/or resistance proteins produced by these LoVo cells in response to
- (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium,
 - (ii) an *in vitro* radiation with a total dose of 10 Gy for a period of about 1 to about 5 minutes, and
 - (iii) an *in vitro* thermic choc at a temperature of about 42°C for a period of about 60 minutes.

This intermediate composition is herein referred to as “**LoVo DS-A**”.

- 25 In one embodiment, LoVo DS-A comprises stress and/or resistance proteins selected from the group comprising or consisting of radiation resistance proteins, thermal stress resistance proteins, metabolic stress resistance proteins, and combination thereof.

In one embodiment, LoVo DS-A cells overexpress at least one, at least two or the three following markers: Cmhsp70.1 (HSP70), CD227 (MUC1) and/or CD107 (LAMP-1).

In one embodiment, LoVo DS-A cells express at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more of at least one, at least two or the three following markers: Cmhsp70.1 (HSP70), CD227 (MUC1) and/or CD107 (LAMP-1); as compared to LoVo cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*; and/or as compared to LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*. Additionally or alternatively, at least one, at least two or the three markers Cmhsp70.1 (HSP70), CD227 (MUC1) and CD107 (LAMP-1) are expressed in at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more cells in a population of LoVo DS-A cells than in a population of LoVo cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*; and/or than in a population of LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*. Additionally or alternatively, the mean fluorescence intensity (MFI) for at least one, at least two or the three markers Cmhsp70.1 (HSP70), CD227 (MUC1) and CD107 (LAMP-1) is at least twice higher, preferably at least 3 times, 4 times, 5 times, 6 times or more higher in a population of LoVo DS-A cells than in a population of LoVo cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*; and/or than in a population of LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*.

In one embodiment, LoVo DS-A cells overexpress at least one or several of the following proteins, as compared to LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*: poly(rC)-binding protein 2, rho guanine nucleotide exchange factor 1, golgin subfamily B member 1, beta-2-glycoprotein 1, plakophilin-3, protein FAM83H, protein transport protein Sec16A, inverted formin-2, anillin, protein ECT2, plectin, epiplakin, proliferation marker protein Ki-67, vesicular integral-membrane protein VIP36, and lysosome-associated membrane glycoprotein 1.

In one embodiment, LoVo DS-A cells specifically express, to detectable levels, at least one or several of the following proteins, as compared to LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*: poly(rC)-binding protein 2, rho guanine nucleotide exchange factor 1, golgin subfamily B member 1, beta-2-glycoprotein 1, plakophilin-3, protein FAM83H, protein transport protein Sec16A, inverted formin-2, anillin, and protein ECT2.

In one embodiment, LoVo DS-A cells overexpress at least one or several of the following membrane and/or cell surface proteins, as compared to LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to any of (i) radiations and (ii) a thermal stress applied *in vitro*: vesicular integral-membrane protein VIP36, nucleolar complex protein 4 homolog, SigI peptidase complex catalytic subunit SEC11A, SigI recognition particle receptor subunit beta, microsomal glutathione S-transferase 1, DPH--cytochrome P450 reductase, very-long-chain (3R)-3-hydroxyacyl-CoA dehydratase 3, mannosyl-oligosaccharide glucosidase, oxysterol-binding protein-related protein 8, receptor expression-enhancing protein 5, MICOS complex subunit MIC26, beta-1,3-galactosyl-O-glycosyl-glycoprotein beta-1,6-N-acetylglucosaminyltransferase 3, mitochondrial ATP synthase membrane subunit DAPIT, mitochondrial ATP synthase subunit f, CDGSH iron-sulfur domain-containing protein 2, vitamin K epoxide reductase complex subunit 1-like protein 1, protein FAM162A, erlin-1, long-chain-fatty-acid--CoA ligase 1, CYB5B_HUMAN Cytochrome b5 type B, retinol dehydrogese 11, transmembrane emp24 domain-containing protein 2, very-long-chain enoyl-CoA reductase, 3-beta-hydroxysteroid-Delta(8),Delta(7)-isomerase, monocarboxylate transporter 1, HLA class I histocompatibility antigen B alpha chain, erlin-2, CD9 antigen, ATP synthase protein 8, mitochondrial fission process protein 1, cytochrome c oxidase subunit NDUFA4, DH dehydrogese [ubiquinone] 1 alpha subcomplex subunit 13, DH dehydrogese [ubiquinone] 1 beta subcomplex subunit 4, DH dehydrogese [ubiquinone] 1 beta subcomplex subunit 3, mitochondrial fission 1 protein, and BRI3-binding protein.

In one embodiment, the intermediate composition comprises or consists of (i) stressed HT-29, HCT-116 or LoVo cells, and (ii) stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these HT-29, HCT-116 or LoVo cells in response to (i) a metabolic stress, and (ii) a chemical stress applied *in vitro*. Chemical stresses and metabolic stresses have been described above.

In one embodiment, the stress and/or resistance proteins are immunogenic. Means and methods for rendering stress and/or resistance proteins immunogenic have been described above.

In one embodiment, the intermediate composition may further comprise tumor-associated antigens (TAA) and/or tumor-specific antigens (TSA), as described above. In one embodiment, TAA and/or TSA are specific of the HT-29, HCT-116 or LoVo cells. In one embodiment, TAA and/or TSA are naturally immunogenic. In one embodiment, TAA and/or TSA may be rendered immunogenic, or their immunogenicity may be increased, by linkage to or by complexation with a means capable to confer immunogenicity. Means capable to confer immunogenicity are well known to the skilled artisan and have been described above. These include, *inter alia* but without limitation, haptens.

In one embodiment, the intermediate composition comprises or consists of (i) stressed HT-29 cells and (ii) stress and/or resistance proteins produced by these HT-29 cells in response to

- (i) an *in vitro* culture in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, and/or at low pH (*e.g.*, below pH 6.5),
- (ii) an *in vitro* exposition to at least one or several chemotherapeutic agents and/or alcohols, for a period ranging from about 6 hours to about 120 hours, preferably from about 24 hours to about 96 hours, such as, *e.g.*, 24, 30, 36, 42, 48, 54, 60, 66, 72, 78, 84, 90, or 96 hours.

In one embodiment, the intermediate composition comprises or consists of (i) stressed HT-29 cells and (ii) stress and/or resistance proteins produced by these HT-29 cells in response to

- (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium, and
- (ii) an *in vitro* exposition to about 13 μ M oxaliplatin for a period of about 72 hours.

This intermediate composition is herein referred to as “**HT-29 DS-B**”.

- 5 In one embodiment, HT-29 DS-B comprises stress and/or resistance proteins selected from the group comprising or consisting of radiation resistance proteins, thermal stress resistance proteins, metabolic stress resistance proteins, and combination thereof.

In one embodiment, HT-29 DS-B cells overexpress at least one, at least two or the three following markers: CD54 (ICAM-1), CD95 (FAS receptor) and/or CD107 (LAMP-1).

- 10 In one embodiment, HT-29 DS-B cells express at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more of at least one, at least two or the three following markers: CD54 (ICAM-1), CD95 (FAS receptor) and/or CD107 (LAMP-1); as compared to HT-29 cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, and (ii) a chemical stress
- 15 applied *in vitro* (*e.g.*, an *in vitro* exposition to about 13 μ M oxaliplatin for a period of about 72 hours); and/or as compared to HT-29 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 13 μ M oxaliplatin for a period of about 72 hours). Additionally or alternatively, at least one, at least two or the three markers CD54 (ICAM-1), CD95 (FAS
- 20 receptor) and/or CD107 (LAMP-1) are expressed in at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more cells in a population of HT-29 DS-B cells than in a population of HT-29 cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, and (ii) a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 13 μ M oxaliplatin for a period of about 72 hours); and/or than in a population of HT-29 cells subjected to a
- 25 metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 13 μ M oxaliplatin for a period of about 72 hours). Additionally or alternatively, the mean fluorescence intensity (MFI) for at least one, at least two or the three markers CD54 (ICAM-1), CD95 (FAS receptor) and/or

CD107 (LAMP-1) is at least twice higher, preferably at least 3 times, 4 times, 5 times, 6 times or more higher in a population of HT-29 DS-B cells than in a population of HT-29 cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, and (ii) a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 13 μ M oxaliplatin for a period of about 72 hours); and/or than in a population of HT-29 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 13 μ M oxaliplatin for a period of about 72 hours).

In one embodiment, HT-29 DS-B cells overexpress at least one or several of the following proteins, as compared to HT-29 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro*: beta-2-glycoprotein 1, histone H2A type 1, poly(rC)-binding protein 2, kinectin, plectin, and NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 7.

In one embodiment, HT-29 DS-B cells specifically express, to detectable levels, at least one or several of the following proteins, as compared to HT-29 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro*: beta-2-glycoprotein 1, histone H2A type 1, and poly(rC)-binding protein 2.

In one embodiment, HT-29 DS-B cells overexpress at least one or several of the following membrane and/or cell surface proteins, as compared to HT-29 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro*: ribosome-binding protein 1, and NADPH--cytochrome P450 reductase.

In one embodiment, HT-29 DS-B cells overexpress at least one or several of the following cell surface proteins, as compared to HT-29 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro*: CD109 antigen, HLA class II histocompatibility antigen gamma chain, HLA class I histocompatibility antigen alpha chain F, hyaluronan mediated motility receptor, integrin beta-8, integrin beta-3, proprotein convertase subtilisin/kexin type 6, clusterin, serotransferrin, natural resistance-associated macrophage protein 2, MHC class I polypeptide-related sequence A, tumor necrosis factor receptor superfamily

- member 10B, endothelial protein C receptor, cadherin EGF LAG seven-pass G-type receptor 3, tumor necrosis factor receptor superfamily member 10A, cystine/glutamate transporter, tissue factor, transforming growth factor beta-1 proprotein, immunoglobulin superfamily member 3, anoctamin-6, metalloproteinase inhibitor 2, toll-like receptor 3,
- 5 volume-regulated anion channel subunit LRRC8A, tetraspanin-15, zinc transporter ZIP6, furin, protocadherin fat 1, hyaluronidase-2, lysophosphatidic acid receptor 2, high mobility group protein B1, chloride intracellular channel protein 4, UL16-binding protein 2, calpain-5, annexin A9, histidine-rich glycoprotein, integrin alpha-4, and heat shock-related 70 kDa protein 2.
- 10 In one embodiment, the intermediate composition comprises or consists of (i) stressed HCT-116 cells and (ii) stress and/or resistance proteins produced by these HCT-116 cells in response to
- (i) an *in vitro* culture in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival
- 15 of cancer cells), under hypoxia, and/or at low pH (*e.g.*, below pH 6.5),
- (ii) an *in vitro* exposition to at least one or several chemotherapeutic agents and/or alcohols, for a period ranging from about 6 hours to about 120 hours, preferably from about 24 hours to about 96 hours, such as, *e.g.*, 24, 30, 36, 42, 48, 54, 60, 66, 72, 78, 84, 90, or 96 hours.
- 20 In one embodiment, the intermediate composition comprises or consists of (i) stressed HCT-116 cells and (ii) stress and/or resistance proteins produced by these HCT-116 cells in response to
- (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium, and
- 25 (ii) an *in vitro* exposition to about 31.5 nM, 100 nM or 315 nM SN-38 (7-ethyl-10-hydroxy-camptothecin) for a period of about 48 hours.

This intermediate composition is herein referred to as “**HCT-116 DS-B**”.

In one embodiment, HCT-116 DS-B comprises stress and/or resistance proteins selected from the group comprising or consisting of radiation resistance proteins, thermal stress resistance proteins, metabolic stress resistance proteins, and combination thereof.

5 In one embodiment, HCT-116 DS-B cells overexpress the following marker: CD66 (CEA).

In one embodiment, HCT-116 DS-B cells express at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more of the marker CD66 (CEA) as compared to HCT-116 cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, and (ii) a chemical stress applied *in vitro* (10 *e.g.*, an *in vitro* exposition to about 31.5 nM, 100 nM or 315 nM SN-38 (7-ethyl-10-hydroxy-camptothecin) for a period of about 48 hours); and/or as compared to HCT-116 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 31.5 nM, 100 nM or 315 nM SN-38 (7-ethyl-10-hydroxy-camptothecin) for a period of about 48 hours). Additionally or alternatively, the marker CD66 (CEA) is expressed in at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more cells in a population of HCT-116 DS-B cells than in a population of HCT-116 cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, and (ii) a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 31.5 nM, 100 nM or 315 nM SN-38 (7-ethyl-10-hydroxy-camptothecin) for a period of about 48 hours); and/or than in a population of HCT-116 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 31.5 nM, 100 nM or 315 nM SN-38 (7-ethyl-10-hydroxy-camptothecin) for a period of about 48 hours). Additionally or alternatively, the mean fluorescence intensity (MFI) for the marker CD66 (CEA) is at least twice higher, preferably at least 3 times, 4 times, 5 times, 6 times or more higher in a population of HCT-116 DS-B cells than in a population of HCT-116 cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, and (ii) a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 31.5 nM, 100 nM or 315 nM SN-38 (7-ethyl-10-hydroxy-camptothecin) for a period of

about 48 hours); and/or than in a population of HCT-116 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 31.5 nM, 100 nM or 315 nM SN-38 (7-ethyl-10-hydroxy-camptothecin) for a period of about 48 hours).

- 5 In one embodiment, HCT-116 DS-B cells overexpress at least one or several of the following proteins, as compared to HCT-116 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro*: beta-2-glycoprotein 1, histone H2A type 1, poly(rC)-binding protein 2, kinectin, plectin, and NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 7.
- 10 In one embodiment, HCT-116 DS-B cells specifically express, to detectable levels, at least one or several of the following proteins, as compared to HCT-116 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro*: beta-2-glycoprotein 1, histone H2A type 1, and poly(rC)-binding protein 2.

- 15 In one embodiment, HCT-116 DS-B cells overexpress at least one or several of the following membrane and/or cell surface proteins, as compared to HCT-116 cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro*: mitochondrial phosphate carrier protein, integrin beta-4, lysosome-associated membrane glycoprotein 1, CD44 antigen, ribosome-binding protein 1, protein transport protein Sec61 subunit alpha isoform 1, kinectin, HLA class I
- 20 histocompatibility antigen A alpha chain, lysophospholipid acyltransferase 7, membrane-associated progesterone receptor component 1, microsomal glutathione S-transferase 1, NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 4, desmoglein-2, integrin alpha-3, torsin-1A-interacting protein 1, plasma membrane calcium-transporting ATPase 1, sphingosine-1-phosphate lyase 1, V-type proton ATPase
- 25 116 kDa subunit a1, mitochondrial inner membrane protein OXA1L, mitochondrial NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 5, transmembrane protein 43, amine oxidase [flavin-containing] B, protein transport protein Sec61 subunit beta, secretory carrier-associated membrane protein 3, protein FAM162A, retinol dehydrogenase 11, ADP-ribosylation factor-like protein 8B, NADH dehydrogenase

[ubiquinone] 1 beta subcomplex subunit 3, dolichol-phosphate mannosyltransferase subunit 3, mitochondrial thiamine pyrophosphate carrier, and ORM1-like protein 2.

In one embodiment, the intermediate composition comprises or consists of (i) stressed LoVo cells and (ii) stress and/or resistance proteins produced by these LoVo cells in response to

- (i) an *in vitro* culture in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, and/or at low pH (*e.g.*, below pH 6.5),
- (ii) an *in vitro* exposition to at least one or several chemotherapeutic agents and/or alcohols, for a period ranging from about 6 hours to about 120 hours, preferably from about 24 hours to about 96 hours, such as, *e.g.*, 24, 30, 36, 42, 48, 54, 60, 66, 72, 78, 84, 90, or 96 hours.

In one embodiment, the intermediate composition comprises or consists of (i) stressed LoVo cells and (ii) stress and/or resistance proteins produced by these LoVo cells in response to

- (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium, and
- (ii) an *in vitro* exposition to about 5 μ M fluorouracil (5-FU) for a period of about 48 hours.

This intermediate composition is herein referred to as “**LoVo DS-B**”.

In one embodiment, LoVo DS-B comprises stress and/or resistance proteins selected from the group comprising or consisting of radiation resistance proteins, thermal stress resistance proteins, metabolic stress resistance proteins, and combination thereof.

In one embodiment, LoVo DS-B cells overexpress at least one or the two following markers: CD243 (MDR-1) and CD66 (CEA); preferably LoVo DS-B cells overexpress CD243 (MDR-1).

In one embodiment, LoVo DS-B cells express at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more of at least one or the two

markers CD243 (MDR-1) and CD66 (CEA) – preferably the marker CD243 (MDR-1) – as compared to LoVo cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, and (ii) a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 5 μ M fluorouracil (5-FU) for a period of about 48 hours); and/or as compared to LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 5 μ M fluorouracil (5-FU) for a period of about 48 hours). Additionally or alternatively, at least one or the two markers CD243 (MDR-1) and CD66 (CEA) – preferably the marker CD243 (MDR-1) – are expressed in at least 10 % more, preferably at least 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 % or more cells in a population of LoVo DS-B cells than in a population of LoVo cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, and (ii) a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 5 μ M fluorouracil (5-FU) for a period of about 48 hours); and/or than in a population of LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 5 μ M fluorouracil (5-FU) for a period of about 48 hours). Additionally or alternatively, the mean fluorescence intensity (MFI) for at least one or the two markers CD243 (MDR-1) and CD66 (CEA) – preferably the marker CD243 (MDR-1) – is at least twice higher, preferably at least 3 times, 4 times, 5 times, 6 times or more higher in a population of LoVo DS-B cells than in a population of LoVo cells cultured in classical conditions such as, *e.g.*, in 10 % FBS, and not subjected to any of (i) a metabolic stress, and (ii) a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 5 μ M fluorouracil (5-FU) for a period of about 48 hours); and/or than in a population of LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro* (*e.g.*, an *in vitro* exposition to about 5 μ M fluorouracil (5-FU) for a period of about 48 hours).

In one embodiment, LoVo DS-B cells overexpress at least one or several of the following proteins, as compared to LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro*: beta-2-glycoprotein 1, histone

H2A type 1, poly(rC)-binding protein 2, kinectin, plectin, and NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 7.

In one embodiment, LoVo DS-B cells specifically express, to detectable levels, at least one or several of the following proteins, as compared to LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro*: beta-2-glycoprotein 1, histone H2A type 1, and poly(rC)-binding protein 2.

In one embodiment, LoVo DS-B cells overexpress at least one or several of the following membrane and/or cell surface proteins, as compared to LoVo cells subjected to a metabolic stress, *e.g.*, cultured in 2 % FBS but not subjected to a chemical stress applied *in vitro*: nuclear pore membrane glycoprotein 210, reticulon-4, NADPH--cytochrome P450 reductase, long-chain-fatty-acid--CoA ligase 3, long-chain-fatty-acid--CoA ligase 4, vesicle-associated membrane protein-associated protein A, mitochondrial tricarboxylate transport protein, kinectin, vesicle-associated membrane protein-associated protein B/C, very-long-chain (3R)-3-hydroxyacyl-CoA dehydratase 3, vesicular integral-membrane protein VIP36, very-long-chain enoyl-CoA reductase, microsomal glutathione S-transferase 1, cytochrome b5 type B, mitochondrial ATP synthase membrane subunit DAPIT, mitochondrial ATP synthase subunit f, cytochrome c oxidase subunit 2, mitochondrial heme protein cytochrome c1, HLA class I histocompatibility antigen B alpha chain, protein LYRIC, monocarboxylate transporter 1, cytochrome c oxidase subunit NDUF44, mannosyl-oligosaccharide glucosidase, signal peptidase complex catalytic subunit SEC11A, protein FAM162A, transmembrane emp24 domain-containing protein 2, retinol dehydrogenase 11, erlin-1, BRI3-binding protein, 3-beta-hydroxysteroid-Delta(8),Delta(7)-isomerase, receptor expression-enhancing protein 5, CDGSH iron-sulfur domain-containing protein 2, CD9 antigen, extended synaptotagmin-2, very long-chain acyl-CoA synthetase, Kunitz-type protease inhibitor 2, CD166 antigen, long-chain-fatty-acid--CoA ligase 1, leukocyte surface antigen CD47, tapasin, and beta-1,3-galactosyl-O-glycosyl-glycoprotein beta-1,6-N-acetylglucosaminyltransferase 3.

Method of manufacturing

The present invention also relates to a method of manufacturing the intermediate compositions described above.

In one embodiment, the method of manufacturing the intermediate compositions described above comprises the following steps:

- a) cultivating HT-29, HCT-116 or LoVo cells in a suitable culture medium;
- b) subjecting the HT-29, HCT-116 or LoVo cells cultured in step a) to one or several stress[es] *in vitro*, wherein these HT-29, HCT-116 or LoVo cells develop resistance mechanism[s] in response to the one or several stress[es] and thereby produce stress and/or resistance proteins, and
- c) recovering the stressed HT-29, HCT-116 or LoVo cells together with the stress and/or resistance proteins they have produced in step b).

In one embodiment, step a) is carried out in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, and/or at low pH (*e.g.*, below pH 6.5). In one embodiment, step a) is carried out in low serum culture conditions in a 2 % FBS culture medium. In one embodiment, this culture in a depleted medium is maintained during step b).

Stresses applied at step b) have been described above.

- In one embodiment, step c) is carried out at least several hours after the completion of step b), preferably more than 12, 24, 36, 48, 60, 72, 84, 96 hours or more after the completion of step b). This ensures that the HT-29, HCT-116 or LoVo cells have had sufficient time to develop their resistance mechanism[s] in response to the one or several stress[es] and thus, sufficient time to produce stress and/or resistance proteins.
- In one embodiment, the method further comprises a step d) of treating the stressed HT-29, HCT-116 or LoVo cells and the stress and/or resistance proteins they have produced, all together recovered in step c), with a molecule or by a means capable of rendering the stress and/or resistance proteins immunogenic.

In one embodiment, step d) comprises or consists of linking or complexing the stress and/or resistance proteins to/with a means capable to confer immunogenicity.

Means capable to confer immunogenicity are well known to the skilled artisan. In one embodiment, the means capable to confer immunogenicity comprise or consist of one or several molecule[s] that is/are not naturally present in the HT-29, HCT-116 and LoVo cells or in their environment. Such means have been detailed above, and include, but are not limited to, haptens.

Examples of haptens include, but are not limited to, 2,4-dinitrophenyl (DNP); 2,4-dinitrofluorobenzene; sulfanilic acid; *N*-iodoacetyl-*N'*-(5-sulfonic-naphthyl)ethylene diamine; anilin; *p*-amino benzoic acid; biotin; fluorescein and derivatives thereof (including FITC, TAMRA, and Texas Red); digoxigenin; 5-nitro-3-pyrazolecarbamide; 4,5-dimethoxy-2-nitrocinnamide; 2-(3,4-dimethoxyphenyl)-quinoline-4-carbamide; 2,1,3-benzoxadiazole-5-carbamide; 3-hydroxy-2-quinoxalinecarbamide, 4-(dimethylamino)azobenzene-4'-sulfonamide (DABSYL); rotenone isooxazoline; 15 (*E*)-2-(2-(2-oxo-2,3-dihydro-1*H*-benzo[*b*][1,4]diazepin-4-yl)phenoxy)acetamide; 7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylic acid; 2-acetamido-4-methyl-5-thiazolesulfonamide; and *p*-methoxyphenylpyrazopodophyllamide.

In one embodiment, step d) comprises or consists of haptenating the stress and/or resistance proteins. In one embodiment, step d) comprises or consists of 20 2,4-dinitrophenylating the stress and/or resistance proteins (*i.e.*, haptenating the stress and/or resistance proteins with 2,4-dinitrophenyl).

The skilled artisan is well aware of means and methods to haptenize proteins such as stress and/or resistance proteins.

25 In one embodiment where the intermediate compositions further comprise tumor-associated antigens (TAA) and/or tumor-specific antigens (TSA), specific of the HT-29, HCT-116 or LoVo cells, step d) also comprises linking or complexing these TAA and/or TSA to/with a means capable to confer immunogenicity, in the same conditions as described above for stress and/or resistance proteins.

In one embodiment, the method further comprises a step of inactivating the HT-29, HCT-116 or LoVo cells in order to render them non-proliferative. If applicable, this step occurs at any time after the HT-29, HCT-116 or LoVo cells have developed their resistance mechanism[s] in response to the stress[es] applied *in vitro* and therefore, after
5 the HT-29, HCT-116 or LoVo cells have produced stress and/or resistance proteins. In one embodiment, the step of inactivating the HT-29, HCT-116 or LoVo cells is carried out at least several hours after the completion of step b), preferably more than 12, 24, 36, 48, 60, 72, 84, 96 hours or more after the completion of step b). In one embodiment, the step of inactivating the HT-29, HCT-116 or LoVo cells is carried out after the completion
10 of step c). In one embodiment, the step of inactivating the HT-29, HCT-116 or LoVo cells is carried out after the completion of step d), if applicable.

The skilled artisan is well aware of means and methods for rendering cells non-proliferative. By way of examples, cells can be rendered non-proliferative by radiation with a dose sufficiently high to kill or inactivate the cells. In one embodiment,
15 radiation – to render the cells non-proliferative – comprises or consists of irradiating the cells with a total dose of 25 Gy or above. Alternatively or additionally, cells can be rendered non-proliferative by ethanol fixation, *e.g.*, using from about 10 % to about 50 % v/v of ethanol. Alternatively or additionally, cells can be rendered non-proliferative by at least one freeze-thaw cycle. Alternatively or additionally, cells can be rendered
20 non-proliferative by linkage to or by complexation with a means capable to confer immunogenicity, such as, *e.g.*, by haptation.

The skilled artisan is also familiar with means and methods to determine whether a cell is non-proliferative or not, *e.g.*, by carrying out viability tests by cell culture (to assess the total lack of proliferation) and/or propidium iodide (which distinguishes between
25 living cells and dead cells).

In one embodiment, the method is for manufacturing the intermediate composition “**HT-29 DS-A**” comprising or consisting of (i) stressed HT-29 cells, and (ii) stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these HT-29 cells in response to (i) a metabolic stress, (ii) radiations and (iii) a thermal stress
30 applied *in vitro*.

In this embodiment, step b) of the method comprises or consists of subjecting the HT-29 cells cultured in step a) of the method to the following stresses *in vitro*: a metabolic stress, radiations and a thermal stress. Radiations, thermal stresses and metabolic stresses have been described above.

- 5 In this embodiment, step b) of the method comprises or consists of subjecting the HT-29 cells cultured in step a) of the method to the following stresses *in vitro*:
- (i) an *in vitro* culture in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, and/or at low pH (*e.g.*, below pH 6.5),
 - 10 (ii) an *in vitro* radiation with a total dose ranging from about 0.25 to about 25 Gy, preferably from about 1 to about 15 Gy, such as, *e.g.*, with a total dose of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 Gy. In one embodiment, the irradiation period ranges from about 1 to about 20 minutes, such as, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 minutes, preferably from about 1 to about 15 minutes; and
 - (iii) an *in vitro* thermic choc at a temperature ranging from about 38°C to about 45°C, such as, *e.g.*, at a temperature of about 38°C, 39°C, 40°C, 41°C, 42°C, 43°C, 44°C, or 45°C, applied to the cells for a period ranging from about 15 minutes to about 4 hours, preferably from about 30 minutes to about 2 hours, such as, *e.g.*, 30, 35, 20 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, or 120 minutes.

In this embodiment, step b) of the method comprises or consists of subjecting the HT-29 cells cultured in step a) of the method to the following stresses *in vitro*:

- (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium,
- (ii) an *in vitro* radiation with a total dose of 10 Gy for a period of about 1 to about 25 5 minutes, and
- (iii) an *in vitro* thermic choc at a temperature of about 42°C for a period of about 60 minutes.

In one embodiment, the stress under (i) is applied throughout the whole period of step b).

- In one embodiment, the stress under (iii) is applied immediately or directly after the stress under (ii), such as, *e.g.*, less than 10 minutes, 20 minutes, 30 minutes, 45 minutes, 1 hour,
- 30

2 hours, or 3 hours after the end of the stress under (ii), while maintaining the stress under (i).

In one embodiment, the method is for manufacturing the intermediate composition “**HCT-116 DS-A**” comprising or consisting of (i) stressed HCT-116 cells, and (ii) stress
5 and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these HCT-116 cells in response to (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*.

In this embodiment, step b) of the method comprises or consists of subjecting the HCT-116 cells cultured in step a) of the method to the following stresses *in vitro*: a
10 metabolic stress, radiations and a thermal stress. Radiations, thermal stresses and metabolic stresses have been described above.

In this embodiment, step b) of the method comprises or consists of subjecting the HCT-116 cells cultured in step a) of the method to the following stresses *in vitro*:

- 15 (i) an *in vitro* culture in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, and/or at low pH (*e.g.*, below pH 6.5),
- (ii) an *in vitro* radiation with a total dose ranging from about 0.25 to about 25 Gy, preferably from about 1 to about 15 Gy, such as, *e.g.*, with a total dose of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 Gy. In one embodiment, the irradiation period
20 ranges from about 1 to about 20 minutes, such as, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 minutes, preferably from about 1 to about 5 minutes; and
- (iii) an *in vitro* thermic choc at a temperature ranging from about 38°C to about 45°C, such as, *e.g.*, at a temperature of about 38°C, 39°C, 40°C, 41°C, 42°C, 43°C, 44°C,
25 or 45°C, applied to the cells for a period ranging from about 15 minutes to about 4 hours, preferably from about 30 minutes to about 2 hours, such as, *e.g.*, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, or 120 minutes.

In this embodiment, step b) of the method comprises or consists of subjecting the HCT-116 cells cultured in step a) of the method to the following stresses *in vitro*:

- (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium,
- (ii) an *in vitro* radiation with a total dose of 10 Gy for a period of about 1 to about 5 minutes, and
- (iii) an *in vitro* thermic choc at a temperature of about 42°C for a period of about 5 60 minutes.

In one embodiment, the stress under (i) is applied throughout the whole period of step b).
 In one embodiment, the stress under (iii) is applied immediately or directly after the stress under (ii), such as, *e.g.*, less than 10 minutes, 20 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, or 3 hours after the end of the stress under (ii), while maintaining the stress under (i).
 10 under (i).

In one embodiment, the method is for manufacturing the intermediate composition “**LoVo DS-A**” comprising or consisting of (i) stressed LoVo cells, and (ii) stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these LoVo cells in response to (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*.
 15 applied *in vitro*.

In this embodiment, step b) of the method comprises or consists of subjecting the LoVo cells cultured in step a) of the method to the following stresses *in vitro*: a metabolic stress, radiations and a thermal stress. Radiations, thermal stresses and metabolic stresses have been described above.

20 In this embodiment, step b) of the method comprises or consists of subjecting the LoVo cells cultured in step a) of the method to the following stresses *in vitro*:

- (i) an *in vitro* culture in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, and/or at low pH (*e.g.*, below pH 6.5),
- 25 (ii) an *in vitro* radiation with a total dose ranging from about 0.25 to about 25 Gy, preferably from about 1 to about 15 Gy, such as, *e.g.*, with a total dose of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 Gy. In one embodiment, the irradiation period ranges from about 1 to about 20 minutes, such as, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10,

11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 minutes, preferably from about 1 to about 5 minutes; and

- (iii) an *in vitro* thermic choc at a temperature ranging from about 38°C to about 45°C, such as, *e.g.*, at a temperature of about 38°C, 39°C, 40°C, 41°C, 42°C, 43°C, 44°C, or 45°C, applied to the cells for a period ranging from about 15 minutes to about 4 hours, preferably from about 30 minutes to about 2 hours, such as, *e.g.*, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, or 120 minutes.

In this embodiment, step b) of the method comprises or consists of subjecting the LoVo cells cultured in step a) of the method to the following stresses *in vitro*:

- (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium,
 (ii) an *in vitro* radiation with a total dose of 10 Gy for a period of about 1 to about 5 minutes, and
 (iii) an *in vitro* thermic choc at a temperature of about 42°C for a period of about 60 minutes.

- In one embodiment, the stress under (i) is applied throughout the whole period of step b). In one embodiment, the stress under (iii) is applied immediately or directly after the stress under (ii), such as, *e.g.*, less than 10 minutes, 20 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, or 3 hours after the end of the stress under (ii), while maintaining the stress under (i).

- In one embodiment, the method is for manufacturing the intermediate composition “**HT-29 DS-B**” comprising or consisting of (i) stressed HT-29 cells, and (ii) stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these HT-29 cells in response to (i) a metabolic stress, and (ii) a chemical stress applied *in vitro*.

- In this embodiment, step b) of the method comprises or consists of subjecting the HT-29 cells cultured in step a) of the method to the following stresses *in vitro*: a metabolic stress and a chemical stress. Chemical stresses and metabolic stresses have been described above.

In this embodiment, step b) of the method comprises or consists of subjecting the HT-29 cells cultured in step a) of the method to the following stresses *in vitro*:

- (i) an *in vitro* culture in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, and/or at low pH (*e.g.*, below pH 6.5),
- (ii) an *in vitro* exposition to at least one or several chemotherapeutic agents and/or alcohols, for a period ranging from about 6 hours to about 120 hours, preferably from about 24 hours to about 96 hours, such as, *e.g.*, 24, 30, 36, 42, 48, 54, 60, 66, 72, 78, 84, 90, or 96 hours.

In this embodiment, step b) of the method comprises or consists of subjecting the HT-29 cells cultured in step a) of the method to the following stresses *in vitro*:

- (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium,
- (ii) an *in vitro* exposition to about 13 μ M oxaliplatin for a period of about 72 hours.

In one embodiment, the stress under (i) is applied throughout the whole period of step b).

In one embodiment, the stress under (ii) is applied while maintaining the stress under (i).

- In one embodiment, the method is for manufacturing the intermediate composition “**HCT-116 DS-B**” comprising or consisting of (i) stressed HCT-116 cells, and (ii) stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these HCT-116 cells in response to (i) a metabolic stress, and (ii) a chemical stress applied *in vitro*.

- In this embodiment, step b) of the method comprises or consists of subjecting the HCT-116 cells cultured in step a) of the method to the following stresses *in vitro*: a metabolic stress and a chemical stress. Chemical stresses and metabolic stresses have been described above.

In this embodiment, step b) of the method comprises or consists of subjecting the HCT-116 cells cultured in step a) of the method to the following stresses *in vitro*:

- (i) an *in vitro* culture in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, and/or at low pH (*e.g.*, below pH 6.5),
- (ii) an *in vitro* exposition to at least one or several chemotherapeutic agents and/or alcohols, for a period ranging from about 6 hours to about 120 hours, preferably

from about 24 hours to about 96 hours, such as, *e.g.*, 24, 30, 36, 42, 48, 54, 60, 66, 72, 78, 84, 90, or 96 hours.

In this embodiment, step b) of the method comprises or consists of subjecting the HCT-116 cells cultured in step a) of the method to the following stresses *in vitro*:

- 5 (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium,
- (ii) an *in vitro* exposition to about 31.5 nM, 100 nM or 315 nM SN-38 (7-ethyl-10-hydroxy-camptothecin) for a period of about 48 hours.

In one embodiment, the stress under (i) is applied throughout the whole period of step b).

In one embodiment, the stress under (ii) is applied while maintaining the stress under (i).

- 10 In one embodiment, the method is for manufacturing the intermediate composition “**LoVo DS-B**” comprising or consisting of (i) stressed LoVo cells, and (ii) stress and/or resistance proteins, wherein the stress and/or resistance proteins were produced by these LoVo cells in response to (i) a metabolic stress, and (ii) a chemical stress applied *in vitro*.

- 15 In this embodiment, step b) of the method comprises or consists of subjecting the LoVo cells cultured in step a) of the method to the following stresses *in vitro*: a metabolic stress and a chemical stress. Chemical stresses and metabolic stresses have been described above.

In this embodiment, step b) of the method comprises or consists of subjecting the LoVo cells cultured in step a) of the method to the following stresses *in vitro*:

- 20 (i) an *in vitro* culture in a depleted medium (*e.g.*, partially or totally lacking one or several substances that are useful or even essential for the growth and/or survival of cancer cells), under hypoxia, and/or at low pH (*e.g.*, below pH 6.5),
- (ii) an *in vitro* exposition to at least one or several chemotherapeutic agents and/or alcohols, for a period ranging from about 6 hours to about 120 hours, preferably
- 25 from about 24 hours to about 96 hours, such as, *e.g.*, 24, 30, 36, 42, 48, 54, 60, 66, 72, 78, 84, 90, or 96 hours.

In this embodiment, step b) of the method comprises or consists of subjecting the LoVo cells cultured in step a) of the method to the following stresses *in vitro*:

- (i) an *in vitro* culture in low serum culture conditions in a 2 % FBS culture medium,
- (ii) an *in vitro* exposition to about 5 μ M fluorouracil (5-FU) for a period of about 48 hours.

In one embodiment, the stress under (i) is applied throughout the whole period of step b).

- 5 In one embodiment, the stress under (ii) is applied while maintaining the stress under (i).

The present invention also relates to a method of manufacturing the composition comprising or consisting of (i) stressed HT-29, HCT-116 and LoVo cells, and (ii) immunogenic stress and/or resistance proteins, described above. This final composition is herein referred to as “**DP**”.

- 10 In one embodiment, the method of manufacturing the composition comprising or consisting of (i) stressed HT-29, HCT-116 and LoVo cells, and (ii) immunogenic stress and/or resistance proteins, described above, comprises the following steps:
- a) obtaining the HT-29 DS-A, HCT-116 DS-A, LoVo DS-A, HT-29 DS-B, HCT-116 DS-B and LoVo DS-B intermediate compositions using the method of
 - 15 manufacturing the intermediate compositions described above; and
 - b) mixing the HT-29 DS-A, HCT-116 DS-A, LoVo DS-A, HT-29 DS-B, HCT-116 DS-B and LoVo DS-B intermediate compositions together.

- In one embodiment, step b) comprises mixing the six intermediate compositions to a final concentration of about 10^6 to about 10^9 stressed HT-29, HCT-116 and LoVo cells per mL of composition, preferably from about 10^7 to about 10^8 stressed HT-29, HCT-116 and LoVo cells per mL of composition, such as, about 1×10^7 , 2×10^7 , 3×10^7 , 4×10^7 , 5×10^7 , 6×10^7 , 7×10^7 , 8×10^7 , 9×10^7 , or 1×10^8 stressed HT-29, HCT-116 and LoVo cells per mL of composition. In one embodiment, step b) comprises mixing the six intermediate compositions to a final concentration of about 3×10^7 stressed HT-29,
- 20 HCT-116 and LoVo cells per mL of composition.
- 25

In one embodiment, step b) comprises mixing the six intermediate compositions in an equal ratio of stressed HT-29, HCT-116 and LoVo cells (*i.e.*, about 1:1:1:1:1:1). In one embodiment, step b) comprises mixing the six intermediate compositions in a ratio comprising or consisting of at least 1.5, 2 or 2.5 times more stressed HT-29 than stressed

HCT-116 or LoVo cells. In one embodiment, step b) comprises mixing the six intermediate compositions in a ratio comprising or consisting of at least 1.5, 2 or 2.5 times more stressed HCT-116 than stressed HT-29 or LoVo cells. In one embodiment, step b) comprises mixing the six intermediate compositions in a ratio comprising or consisting of at least 1.5, 2 or 2.5 times more stressed LoVo than stressed HT-29 or HCT-116 cells.

In one embodiment, step b) comprises mixing together:

- from about 10^5 to about 10^8 stressed HT-29 cells (from HT-29 DS-A and HT-29 DS-B), preferably from about 10^6 to about 10^7 stressed HT-29 cells, such as, about 1×10^6 , 2×10^6 , 3×10^6 , 4×10^6 , 5×10^6 , 6×10^6 , 7×10^6 , 8×10^6 , 9×10^6 , or 1×10^7 stressed HT-29 cells,
- from about 10^5 to about 10^8 stressed HCT-116 cells (from HCT-116 DS-A and HCT-116 DS-B), preferably from about 10^6 to about 10^7 stressed HCT-116 cells, such as, about 1×10^6 , 2×10^6 , 3×10^6 , 4×10^6 , 5×10^6 , 6×10^6 , 7×10^6 , 8×10^6 , 9×10^6 , or 1×10^7 stressed HCT-116 cells, and
- from about 10^5 to about 10^8 stressed LoVo cells (from LoVo DS-A and LoVo DS-B), preferably from about 10^6 to about 10^7 stressed LoVo cells, such as, about 1×10^6 , 2×10^6 , 3×10^6 , 4×10^6 , 5×10^6 , 6×10^6 , 7×10^6 , 8×10^6 , 9×10^6 , or 1×10^7 stressed LoVo cells.

Therapeutic uses and methods

The present invention also relates to a method of treating cancer in a subject in need thereof, comprising administering to said subject the composition comprising or consisting of (i) stressed HT-29, HCT-116 and LoVo cells, and (ii) immunogenic stress and/or resistance proteins produced by these cells in response to a stress applied *in vitro*, as defined hereinabove.

The present invention also relates to the composition comprising or consisting of (i) stressed HT-29, HCT-116 and LoVo cells, and (ii) immunogenic stress and/or resistance proteins produced by these cells in response to a stress applied *in vitro*, as defined hereinabove, for use in treating cancer in a subject in need thereof.

The present invention also relates to the use of the composition comprising or consisting of (i) stressed HT-29, HCT-116 and LoVo cells, and (ii) immunogenic stress and/or resistance proteins produced by these cells in response to a stress applied *in vitro*, as defined hereinabove, for treating cancer in a subject in need thereof.

- 5 The present invention also relates to the use of the composition comprising or consisting of (i) stressed HT-29, HCT-116 and LoVo cells, and (ii) immunogenic stress and/or resistance proteins produced by these cells in response to a stress applied *in vitro*, as defined hereinabove, for the manufacture of a medicament for treating cancer in a subject in need thereof.

- 10 In one embodiment, treating cancer comprises eliciting an immune response against cancer cells from said cancer.

Examples of cancers include those listed in the 10th revision of the International Statistical Classification of Diseases and Related Health Problems (ICD), under chapter II, blocks C00 to D48.

- 15 Further examples of cancers include, but are not limited to, recurrent, metastatic or multi-drug resistant cancer.

- Further examples of cancers include, but are not limited to, adenofibroma, adenoma, agnogenic myeloid metaplasia, AIDS-related malignancies, ameloblastoma, anal cancer, angiofollicular mediastinal lymph node hyperplasia, angiokeratoma, angiolymphoid
 20 hyperplasia with eosinophilia, angiomas, anhidrotic ectodermal dysplasia, anterofacial dysplasia, apocrine metaplasia, apudoma, asphyxiating thoracic dysplasia, astrocytoma (including, *e.g.*, cerebellar astrocytoma and cerebral astrocytoma), atriadigital dysplasia, atypical melanocytic hyperplasia, atypical metaplasia, autoparenchymatous metaplasia, basal cell hyperplasia, benign giant lymph node
 25 hyperplasia, bile duct cancer (including, *e.g.*, extrahepatic bile duct cancer), bladder cancer, bone cancer, brain tumor (including, *e.g.*, brain stem glioma, cerebellar astrocytoma glioma, malignant glioma, supratentorial primitive neuroectodermal tumors, visual pathway and hypothalamic glioma, ependymoma, medulloblastoma, gestational trophoblastic tumor glioma, and paraganglioma), branchionia, female breast cancer, male

breast cancer, bronchial adenomas/carcinoids, bronchopulmonary dysplasia, cancer
growths of epithelial cells, pre-cancerous growths of epithelial cells, metastatic growths
of epithelial cells, carcinoid heart disease, carcinoid tumor (including, *e.g.*,
gastrointestinal carcinoid tumor), carcinoma (including, *e.g.*, carcinoma of unknown
5 primary origin, adrenocortical carcinoma, islet cells carcinoma, adeno carcinoma,
adeoncortical carcinoma, basal cell carcinoma, basosquamous carcinoma, bronchiolar
carcinoma, Brown-Pearce carcinoma, cystadenocarcinoma, ductal carcinoma,
hepatocarcinoma, Krebs carcinoma, papillary carcinoma, oat cell carcinoma, small cell
lung carcinoma, non-small cell lung carcinoma, squamous cell carcinoma, transitional
10 cell carcinoma, Walker carcinoma, Merkel cell carcinoma, and skin carcinoma),
cementoma, cementum hyperplasia, cerebral dysplasia, cervical cancer, cervical
dysplasia, cholangioma, cholesteatoma, chondroblastoma, chondroectodermal dysplasia,
chordoma, choristoma, chrondroma, cleidocranial dysplasia, colon adenocarcinoma,
colon cancer, colon carcinoma, colorectal adenocarcinoma, colorectal cancer, colorectal
15 carcinoma, local metastasized colorectal cancer, congenital adrenal hyperplasia,
congenital ectodermal dysplasia, congenital sebaceous hyperplasia, connective tissue
metaplasia, craniocarpotarsal dysplasia, craniodiaphysial dysplasia, craniometaphysial
dysplasia, craniopharyngioma, cylindroma, cystadenoma, cystic hyperplasia (including,
e.g., cystic hyperplasia of the breast), cystosarconia phyllodes, dentin dysplasia, denture
20 hyperplasia, diaphysial dysplasia, ductal hyperplasia, dysgenninoma, dysplasia
epiphysialis hemimelia, dysplasia epiphysialis multiplex, dysplasia epiphysialis punctate,
ectodermal dysplasia, Ehrlich tumor, enamel dysplasia, encephaloophthalmic dysplasia,
endometrial cancer (including, *e.g.*, ependymoma and endometrial hyperplasia),
ependymoma, epithelial cancer, epithelial dysplasia, epithelial metaplasia, esophageal
25 cancer, Ewing's family of tumors (including, *e.g.*, Ewing's sarcoma), extrahepatic bile
duct cancer, eye cancer (including, *e.g.*, intraocular melanoma and retinoblastoma),
faciodigitogenital dysplasia, familial fibrous dysplasia of jaws, familial white folded
dysplasia, fibroma, fibromuscular dysplasia, fibromuscular hyperplasia, fibrous dysplasia
of bone, florid osseous dysplasia, focal epithelial hyperplasia, gall bladder cancer,
30 ganglioneuroma, gastric cancer (including, *e.g.*, stomach cancer), gastrointestinal
carcinoid tumor, gastrointestinal tract cancer, gastrointestinal tumors, Gaucher's disease,
germ cell tumors (including, *e.g.*, extracranial germ cell tumors, extragonadal germ cell

tumors, and ovarian germ cell tumors), giant cell tumor, gingival hyperplasia, glioblastoma, glomangioma, granulosa cell tumor, gynandroblastoma, hamartoma, head and neck cancer, hemangioendothelioma, hemangioma, hemangiopericytoma, hepatocellular cancer, hepatoma, hereditary renal-retinal dysplasia, hidrotic ectodermal dysplasia, histiocytoma, histiocytosis, hypergammaglobulinemia, hypohidrotic ectodermal dysplasia, hypopharyngeal cancer, inflammatory fibrous hyperplasia, inflammatory papillary hyperplasia, intestinal cancers, intestinal metaplasia, intestinal polyps, intraocular melanoma, intravascular papillary endothelial hyperplasia, kidney cancer, laryngeal cancer, leiomyoma, leukemia (including, *e.g.*, acute lymphoblastic leukemia, acute lymphocytic leukemia, acute myeloid leukemia, acute myelogenous leukemia, acute hairy cell leukemia, acute B-cell leukemia, acute T-cell leukemia, acute HTLV leukemia, chronic lymphoblastic leukemia, chronic lymphocytic leukemia, chronic myeloid leukemia, chronic myelogenous leukemia, chronic hairy cell leukemia, chronic B-cell leukemia, chronic T-cell leukemia, and chronic HTLV leukemia), Leydig cell tumor, lip and oral cavity cancer, lipoma, liver cancer, lung cancer (including, *e.g.*, small cell lung cancer and non-small cell lung cancer), lymphangiomyoma, lymphangioma, lymphoma (including, *e.g.*, AIDS-related lymphoma, central nervous system lymphoma, primary central nervous system lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, Hodgkin's lymphoma during pregnancy, non-Hodgkin's lymphoma during pregnancy, mast cell lymphoma, B-cell lymphoma, adenolymphoma, Burkitt's lymphoma, cutaneous T-cell lymphoma, large cell lymphoma, and small cell lymphoma), lymphopenic thymic dysplasia, lymphoproliferative disorders, macroglobulinemia (including, *e.g.*, Waldenstrom's macroglobulinemia), malignant carcinoid syndrome, malignant mesothelioma, malignant thymoma, mammary dysplasia, mandibulofacial dysplasia, medulloblastoma, meningioma, mesenchymoma, mesonephroma, mesothelioma (including, *e.g.*, malignant mesothelioma), metaphyseal dysplasia, metaplastic anemia, metaplastic ossification, metaplastic polyps, metastatic squamous neck cancer (including, *e.g.*, metastatic squamous neck cancer with occult primary), Mondini dysplasia, monostotic fibrous dysplasia, mucoepithelial dysplasia, multiple endocrine neoplasia syndrome, multiple epiphyseal dysplasia, multiple myeloma/plasma cell neoplasm, mycosis fungoides, myelodysplastic syndrome, myeloid metaplasia, myeloproliferative disorders, chronic myeloproliferative disorders,

myoblastoma, myoma, myxoma, nasal cavity and paranasal sinus cancer, nasopharyngeal cancer, prostatic neoplasm, colon neoplasm, abdomen neoplasm, bone neoplasm, breast neoplasm, digestive system neoplasm, liver neoplasm, pancreas neoplasm, peritoneum neoplasm, endocrine glands neoplasm (including, *e.g.*, adrenal neoplasm, parathyroid neoplasm, pituitary neoplasm, testicles neoplasm, ovary neoplasm, thymus neoplasm, and thyroid neoplasm), eye neoplasm, head and neck neoplasm, nervous system neoplasm (including, *e.g.*, central nervous system neoplasm and peripheral nervous system neoplasm), lymphatic system neoplasm, pelvic neoplasm, skin neoplasm, soft tissue neoplasm, spleen neoplasm, thoracic neoplasm, urogenital tract neoplasm, neurilemmoma, neuroblastoma, neuroepithelioma, neurofibroma, neurofibromatosis, neuroma, nodular hyperplasia of prostate, nodular regenerative hyperplasia, oculoauriculovertebral dysplasia, oculodentodigital dysplasia, oculovertbral dysplasia, odontogenic dysplasia, odontoma, opthalmomandibulomelic dysplasia, oropharyngeal cancer, osteoma, ovarian cancer (including, *e.g.*, ovarian epithelial cancer and ovarian low malignant potential tumor), pancreatic cancer (including, *e.g.*, islet cell pancreatic cancer and exocrine pancreatic cancer), papilloma, paraganglioma, nonchromaffin paraganglioma, paranasal sinus and nasal cavity cancer, paraproteinemias, parathyroid cancer, periapical cemental dysplasia, pheochromocytoma (including, *e.g.*, penile cancer), pineal and supratentorial primitive neuroectodermal tumors, pinealoma, pituitary tumor, plasma cell neoplasm/multiple myeloma, plasmacytoma, pleuropulmonary blastoma, polyostotic fibrous dysplasia, polyps, pregnancy cancer, pre-neoplastic disorders (including, *e.g.*, benign dysproliferative disorders such as benign tumors, fibrocystic conditions, tissue hypertrophy, intestinal polyps, colon polyps, esophageal dysplasia, leukoplakia, keratoses, Bowen's disease, Farmer's skin, solar cheilitis, and solar keratosis), primary hepatocellular cancer, primary liver cancer, primary myeloid metaplasia, prostate cancer, pseudoachondroplastic spondyloepiphyseal dysplasia, pseudoepitheliomatous hyperplasia, purpura, rectal cancer, renal cancer (including, *e.g.*, kidney cancer, renal pelvis, ureter cancer, transitional cell cancer of the renal pelvis and ureter), reticuloendotheliosis, retinal dysplasia, retinoblastoma, salivary gland cancer, sarcomas (including, *e.g.*, uterine sarcoma, soft tissue sarcoma, carcinosarcoma, chondrosarcoma, fibrosarcoma, hemangiosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, myosarcoma, myxosarcoma, rhabdosarcoma,

sarcoidosis sarcoma, osteosarcoma, Ewing sarcoma, malignant fibrous histiocytoma of bone, and clear cell sarcoma of tendon sheaths), sclerosing angioma, secondary myeloid metaplasia, senile sebaceous hyperplasia, septooptic dysplasia, Sertoli cell tumor, Sezary syndrome, skin cancer (including, *e.g.*, melanoma skin cancer and non-melanoma skin cancer), small intestine cancer, spondyloepiphyseal dysplasia, squamous metaplasia (including, *e.g.*, squamous metaplasia of amnion), stomach cancer, supratentorial primitive neuroectodermal and pineal tumors, supratentorial primitive neuroectodermal tumors, symptomatic myeloid metaplasia, teratoma, testicular cancer, theca cell tumor, thymoma (including, *e.g.*, malignant thymoma), thyroid cancer, trophoblastic tumors (including, *e.g.*, gestational trophoblastic tumors), ureter cancer, urethral cancer, uterine cancer, vaginal cancer, ventriculoradial dysplasia, verrucous hyperplasia, vulvar cancer, Waldenstrom's macroglobulinemia, and Wilms' tumor.

In one embodiment, the cancer is selected from the group comprising or consisting of renal cancer, liver cancer, pancreatic cancer, endometrial cancer, colorectal cancer, ovarian cancer, breast cancer, lung cancer, head and neck cancer, cervical cancer, melanoma and glioma.

In one embodiment, the cancer is selected from the group comprising or consisting of colon and colorectal cancers, including, but not limited to, colon adenocarcinoma, colon cancer, colon carcinoma, colorectal adenocarcinoma, colorectal cancer, and colorectal carcinoma.

In one embodiment, the composition is administered or is to be administered once. In one embodiment, the composition is administered or is to be administered several times, such as 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 times or more; or indefinitely until the cancer is treated.

In one embodiment, the composition is administered or is to be administered every week, every two weeks, every three weeks, every month, every two months, every three months, every six months, every year.

In one embodiment, the composition is administered or is to be administered every week, every two weeks, every three weeks, every month, preferably every week, for the first

weeks, such as for the first 2, 3, 4, 5, 6, 7, 8, 9, 10 weeks or more, preferably for the first 6 to 8 weeks, as an induction or bolus regimen.

In one embodiment, the composition is administered or is to be administered every month, every two months, every three months, every six months, every year after the induction
5 or bolus regimen, as a boost regimen.

In one embodiment, the composition is administered or is to be administered every week for the first 6 to 8 weeks; then about one month later; then about 3 months later; then every 6 months until the cancer is treated.

In one embodiment, the composition is administered or is to be administered at a total
10 dose ranging from about 10^5 to about 10^8 stressed HT-29, HCT-116 and LoVo cells, preferably from about 10^6 to about 10^7 stressed HT-29, HCT-116 and LoVo cells, such as, about 1×10^6 , 2×10^6 , 3×10^6 , 4×10^6 , 5×10^6 , 6×10^6 , 7×10^6 , 8×10^6 , 9×10^6 , or 1×10^7 stressed HT-29, HCT-116 and LoVo cells.

In one embodiment, the composition is administered or is to be administered as a single
15 dose.

In one embodiment, the composition is administered or is to be administered in at least two doses, such as 2, 3, 4, 5 or more dose.

In one embodiment, the composition is administered or is to be administered at a total dose of about 3×10^6 stressed HT-29, HCT-116 and LoVo cells. In one embodiment, the
20 composition is administered or is to be administered at a total dose of about 6×10^6 stressed HT-29, HCT-116 and LoVo cells. In one embodiment, the composition is administered or is to be administered in a single dose of 6×10^6 stressed HT-29, HCT-116 and LoVo cells, or in two doses of 3×10^6 stressed HT-29, HCT-116 and LoVo cells.

In one embodiment, the composition is administered or is to be administered in a volume
25 ranging from about 10 μ L to about 1 mL, such as, about 10 μ L, 50 μ L, 100 μ L, 200 μ L, 300 μ L, 400 μ L, 500 μ L, 600 μ L, 700 μ L, 800 μ L, 900 μ L, or 1 mL, preferably in a volume of about 100 μ L.

In one embodiment, the composition is administered or is to be administered systemically.
In one embodiment, the composition is administered or is to be administered by injection, preferably by systemic injection.

Examples of systemic injections include, but are not limited to, intravenous (iv),
5 subcutaneous (sq), intradermal (id), intramuscular (im), intraarterial, intraparenteral, intranodal, intralymphatic, intraperitoneal (ip), intracranial, intracardiac, intralesional, intraprostatic, intravaginal, intrarectal, intrathecal, intranasal, intratumoral (it), intravesicular, and perfusion.

In one embodiment, the composition is administered or is to be administered
10 intradermally, subcutaneously, intratumorally or intravenously.

In one embodiment, the composition is administered or is to be administered intradermally.

It will be understood that other suitable routes of administration are also contemplated in the present invention, and the administration mode will ultimately be decided by the
15 attending physician within the scope of sound medical judgment.

In one embodiment, the composition is administered or is to be administered to the subject before, concomitantly with, or after administration of at least one additional therapy.

Examples of suitable additional therapies include chemotherapeutic agents as defined above, radiation therapy and immunostimulatory agents.

20 Suitable examples of radiation therapies include, but are not limited to, external beam radiotherapy (such as, *e.g.*, superficial X-rays therapy, orthovoltage X-rays therapy, megavoltage X-rays therapy, radiosurgery, stereotactic radiation therapy, cobalt therapy, electron therapy, fast neutron therapy, neutron-capture therapy, proton therapy, and the like); brachytherapy; unsealed source radiotherapy; tomotherapy; and the like.

25 Suitable examples of immunostimulatory agents include those described under subgroup L03 of the Anatomical Therapeutic Chemical Classification System. Further examples of immunostimulatory agents include, but are not limited to, cytokines (such as, *e.g.*,

filgrastim, pegfilgrastim, lenograstim, molgramostim, sargramostim, ancestim, albinterferon, interferon alfa natural, interferon alfa 2a, peginterferon alfa-2a, interferon alfa 2b, peginterferon alfa-2b, interferon alfa n1, interferon alfacon-1, interferon alpha-n3, interferon beta natural, interferon beta 1a, interferon beta 1b, interferon gamma,

5 aldesleukin, oprelvekin, and the like); immune checkpoint inhibitors (such as, *e.g.*, inhibitors of CTLA4, PD-1, PD-L1, LAG-3, B7-H3, B7-H4, TIM3, A2AR, and/or IDO, including nivolumab, pembrolizumab, pidilizumab, AMP-224, MPDL3280A, MDX-1105, MEDI-4736, arelumab, ipilimumab, tremelimumab, pidilizumab, IMP321, MGA271, BMS-986016, lirilumab, urelumab, PF-05082566, IPH2101, MEDI-6469,

10 CP-870,893, mogamulizumab, varlilumab, avelumab, galiximab, AMP-514, AUNP 12, indoximod, NLG-919, INCB024360, and the like); toll-like receptor agonists (such as, *e.g.*, buprenorphine, carbamazepine, ethanol, fentanyl, GS-9620, imiqimod, lefitolimod, levorphanol, methadone, morphine, (+)-morphine, morphine-3-glucuronide, oxcarbazepine, oxycodone, pethidine, resiquimod, SD-101, tapentadol, tilisotolimod,

15 VTX-2337, glucuronoxylomannan from *Cryptococcus*, MALP-2 from *Mycoplasma*, MALP-404 from *Mycoplasma*, OspA from *Borrelia*, porin from *Neisseria* or *Haemophilus*, hsp60, hemmagglutinin, LcrV from *Yersinia*, bacterial flagellin, lipopolysaccharide, lipoteichoic acid, lipomannan from *Mycobacterium*, glycosylphosphatidylinositol, lysophosphatidylserine, lipophosphoglycan from

20 *Leishmania*, zymosan from *Saccharomyces*, Pam2CGDPKHPKSF, Pam3CSK4, CpG oligodeoxynucleotides, poly(I:C) nucleic acid sequences, poly(A:U) nucleic acid sequences, double-stranded viral RNA, and the like); STING receptor agonists (such as, *e.g.*, those described in §0240 of WO2017100305, vadimezan, CL656, ADU-S100, 3'3'-cGAMP, 2'3'-cGAMP, ML RR-S2 CDG, ML RR-S2 cGAMP, cyclic di-GMP,

25 DMXAA, DiABZI, and the like); CD1 ligands; growth hormone; granulocyte-macrophage colony-stimulating factor (GM-CSF); immunocyanin; pegademase; prolactin; tasonermin; female sex steroids; histamine dihydrochloride; poly ICLC; vitamin D; lentinan; plerixafor; roquinimex; mifamurtide; glatiramer acetate; thymopentin; thymosin α 1; thymulin; polyinosinic:polycytidylic acid; pidotimod;

30 Bacillus Calmette-Guérin; melanoma vaccine; sipuleucel-T; and the like.

In one embodiment, the composition may be administered to the subject after administration of chemotherapeutic agent such as, *e.g.*, cyclophosphamide.

Additionally or alternatively, the composition may be administered or to the subject concomitantly with administration of an immunostimulatory agent, such as, *e.g.*,
5 granulocyte-macrophage colony-stimulating factor (GM-CSF).

Kit-of-parts

The present invention also relates to a kit-of-part.

The term “**kit-of-parts**”, also termed “**package**”, “**commercial package**”, or
10 “**pharmaceutical package**”, shall encompass an entity of physically separated components, which are intended for individual storage and/or use, but in functional relation to each other. In particular, the term “**kit-of-parts**” encompasses an entity of physically separated components which are to be used or administered separately, in any order, in a given time interval, typically but not necessarily ranging from hours to days,
15 weeks or more; or concomitantly, either as an extemporaneous formulation or individually, in any order, within a time interval ranging from seconds to minutes or hours.

In one embodiment, the kit-of-parts comprises:

- the intermediate composition named “**HT-29 DS-A**”, as defined above;
- 20 - the intermediate composition named “**HCT-116 DS-A**”, as defined above; and
- the intermediate composition named “**LoVo DS-A**”, as defined above.

In one embodiment, the kit-of-parts comprises:

- the intermediate composition named “**HT-29 DS-B**”, as defined above;
- the intermediate composition named “**HCT-116 DS-B**”, as defined above; and
- 25 - the intermediate composition named “**LoVo DS-B**”, as defined above.

In one embodiment, the kit-of-parts comprises:

- the intermediate composition named “**HT-29 DS-A**”, as defined above;
- the intermediate composition named “**HCT-116 DS-A**”, as defined above;

- the intermediate composition named “**LoVo DS-A**”, as defined above;
- the intermediate composition named “**HT-29 DS-B**”, as defined above;
- the intermediate composition named “**HCT-116 DS-B**”, as defined above; and
- the intermediate composition named “**LoVo DS-B**”, as defined above.

- 5 In one embodiment, the kit-of-parts further comprises instructions for mixing the intermediate compositions, according to the method of manufacturing the composition comprising or consisting of (i) stressed HT-29, HCT-116 and LoVo cells, and (ii) immunogenic stress and/or resistance proteins, named “**DP**”, described above.

In one embodiment, the kit-of-parts further comprises at least one additional therapeutic agent, such as, *e.g.*, at least one chemotherapeutic agent and/or immunostimulatory agent,
10 as defined above.

In one embodiment, the kit-of-parts comprises:

- the composition named “**DP**”, as defined above;
- at least one additional therapeutic agent, such as, *e.g.*, at least one chemotherapeutic
15 agent and/or immunostimulatory agent, as defined above.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-B are two Venn diagrams comparing the number of identified proteins in each of the 3 human cell lines (HT-29, HCT-116 and LoVo) for untreated samples.

20 **Figure 1A:** RCB samples.

Figure 1B: MCB samples.

Figure 2 is a graph showing the differential number of proteins expressed in the HCT-116 MCB, DS-A, DS-B and in the final product (DP) comprising all six DS described above (hence including the HCT-116 DS-A and DS-B), by comparison to the initial HCT-116
25 RCB cultured in classical conditions (with 10 % FBS). “+” indicates proteins that are newly expressed, “-” indicates the number of proteins which are no longer expressed at all.

Figure 3 is a graph showing the number of proteins over- and under-expressed in the HCT-116 MCB, DS-A and DS-B, by comparison to the initial HCT-116 RCB cultured in classical conditions (with 10 % FBS).

Figure 4 is a graph showing the differential number of proteins expressed in the HT-29 MCB, DS-A, DS-B and in the final product (DP) comprising all six DS described above (hence including the HT-29 DS-A and DS-B), by comparison to the initial HT-29 RCB cultured in classical conditions (with 10 % FBS). “+” indicates proteins that are newly expressed, “-” indicates the number of proteins which are no longer expressed at all.

Figure 5 is a graph showing the number of proteins over- and under-expressed in the HT-29 MCB, DS-A and DS-B, by comparison to the initial HT-29 RCB cultured in classical conditions (with 10 % FBS).

Figure 6 is a graph showing the differential number of proteins expressed in the LoVo MCB, DS-A, DS-B and in the final product (DP) comprising all six DS described above (hence including the LoVo DS-A and DS-B), by comparison to the initial LoVo RCB cultured in classical conditions (with 10 % FBS). “+” indicates proteins that are newly expressed, “-” indicates the number of proteins which are no longer expressed at all.

Figure 7 is a graph showing the number of proteins over- and under-expressed in the LoVo MCB, DS-A and DS-B, by comparison to the initial LoVo RCB cultured in classical conditions (with 10 % FBS).

Figure 8 is a graph showing the number of proteins identified (on the y-axis), whether membrane and/or cell surface proteins or others (*i.e.*, any other protein for which the terms “membrane” or “cell surface” were not reported as annotations in the “cellular component” field of the Thermo Protein Center database), in several compositions of HT-29 cells: HT-29 MCB, HT-29 DS-A, and HT-29 DS-B; as well as in the drug product (DP) comprising a mix of HT-29 DS-A, HT-29 DS-B, HCT-116 DS-A, HCT-116 DS-B, LoVo DS-A and LoVo DS-B. Experiments were performed in triplicates, indicated as “#1”, “#2” and “#3”.

Figures 9A-D are pie charts showing the protein distribution according to *p*-values and fold changes, between HT-29 DS-A *versus* HT-29 MCB (**Fig. 9A**), HT-29 DS-B *versus* HT-29 MCB (**Fig. 9B**), HT-29 DS-A *versus* HT-29 DS-B (**Fig. 9C**), and DP *versus* HT-29 MCB (**Fig. 9D**).

- 5 **Figure 10** is a graph showing the number of membrane and/or cell surface proteins (on the y-axis) which are either statistically significantly under-expressed (*i.e.*, with a $p\text{-value} \leq 0.05$ and an abundance ratio ≤ 0.5) or over-expressed (*i.e.*, with a $p\text{-value} \leq 0.05$ and an abundance ratio ≥ 2) between HT-29 DS-A *versus* HT-29 MCB, HT-29 DS-B *versus* HT-29 MCB, HT-29 DS-A *versus* HT-29 DS-B, and DP *versus*
- 10 HT-29 MCB. The number of under-expressed proteins is displayed as negative number.

Figures 11A-C are three graphs showing the secretion of IL-8 or IL-12 from dendritic cells, expressed in pg/mL. **Fig. 11A**: secretion of IL-8 in absence of CD40L; **Fig. 11B**: secretion of IL-8 in presence of 0.6 $\mu\text{g/mL}$ CD40L; **Fig. 11C**: secretion of IL-12 in presence of 0.6 $\mu\text{g/mL}$ CD40L.

- 15 **Figure 12** is a graph showing the IFN γ release in the supernatant collected from a co-culture of mDCs and autologous CD8⁺ T cells treated with various ratios of STC-1010. The hashed line represents the limit of detection of IFN γ .

- Figures 13A-C** are three graphs showing the expression of the DCs activation markers CD40, CD83 and CD86 assessed by RT-qPCR upon treatment with various doses of
- 20 STC-1010. Data are expressed as relative quantity normalized to the expression of chicken GAPDH. **Fig. 13A**: expression of CD40; **Fig. 13B**: expression of CD83; **Fig. 13C**: expression of CD86.

EXAMPLES

The present invention is further illustrated by the following examples.

Example 1

5 **Selection of cell lines for the human colorectal cancer vaccine**

We aimed at developing an anti-cancer treatment turning cancer cell resistance mechanisms against themselves, by artificially developing resistance mechanisms *in vitro* in cancer cell lines and rendering them immunogenic.

Several colorectal human cell lines were selected as suitable candidates for the
10 development of a human colorectal cancer vaccine.

The initial selection of cell lines was made based on their biological and genetic characteristics, so as to cover the largest panel of clinical typologies: microsatellite stability (MSS vs. MSI status); mutated vs. wild-type *BRAF*, *KRAS*, *PIK3CA*, *PTEN* and *TP53* genes; known resistance to common anticancer treatments (fluorouracil [5-FU],
15 oxaliplatin, SN-38 (7-ethyl-10-hydroxy-camptothecin), anti-VEGF antibodies).

Based on the literature and databases (SelTARbase, Cellosaurus and GDSC database), the following cell lines were selected:

- LS 174T (ATCC[®] ref.: CL-188[™]),
- SW620 (ATCC[®] ref.: CCL-227[™]),
- 20 - SW480 (ATCC[®] ref.: CCL-228[™]),
- LoVo (ATCC[®] ref.: CCL-229[™]),
- SW48 (ATCC[®] ref.: CCL-231[™]),
- HCT-116 (ATCC[®] ref.: CCL-247[™]), and
- HT-29 (ATCC[®] ref.: HTB-38[™]),

25 Pre-Research Cell Bank (RCB): these cell lines were acquired from ATCC and cultured with 10 % FBS to obtain seven RCB, but only six out of these seven had an expected growth profile and were viable after a freeze-thaw cycle: SW620 was discarded at this stage (**Table 1**).

Table 1: RCB cultures

Cell line	Medium	Freezing time	Viability at thawing	Doubling time
HCT-116	McCoy's 5A	P3	85-92 %	25 h
HT-29	McCoy's 5A	P3	83 %	34 h
LoVo	F12K	P3	83 %	45 h
LS 174T	EMEM	P2	47-63 %	32.4 h
SW48	Leibovitz's L15	P2	96 %	n.d.
SW480	Leibovitz's L15	P3	82 %	55 h
SW620	Leibovitz's L15	P3	n.d.	n.d.

Pre-RCB 2 %: these 6 RCBs, with expected growth profile, have been adapted to low serum culture condition (10 % FBS → 5 % FBS → 2 % FBS) to mimic the nutrients depletion action observed with anti-VEGF antibodies.

- 5 Only four cell lines were able to grow in low serum culture condition and were viable after a freeze-thaw cycle (HCT-116, HT-29, LoVo, and SW480) (**Table 2**).

Table 2: RCB 2 % cultures

Cell line	Freezing time	Viability at thawing	Doubling time
HCT-116	P9	37-55 %	25-30 h
HT-29	P6	52-78 %	35 h
LoVo	P4	28-49 %	40-45 h
LS 174T	P4	n.d.	53 h
SW480	P9	93-96 %	60-70 h
SW48	P4	n.d.	n.d.

These four cell lines were then exposed to two further types of stresses to obtain “DS-A” and “DS-B”.

- 10 DS-A: cells were exposed to a physical stress comprising a combination of (1) a low dose ionizing radiation (10 Gy) for 5 minutes, and (2) a thermal stress (1 hour at 42°C), in order to induce heat shock protein HSP70 expression.

DS-B: cells were exposed to a chemical stress comprising a chemical stimulation with chemotherapeutic agents commonly used for the colorectal cancer: 5-FU (Sigma, F6627), oxaliplatin (Sigma, Y0000271) or SN-38 (7-ethyl-10-hydroxy-camptothecin) (Sigma, H0165). In particular, the association of a chemotherapeutic agent with each cell line was based on literature assumptions (**Table 3**), and further on the degree of resistance of the cell line to each chemotherapeutic agent. For each cell line, this resistance level to each chemotherapeutic agent was assessed by determining the half maximal inhibitory concentration (IC₅₀) in a cytotoxic assay. IC₅₀ are given in **Table 4**.

Table 3: Chemoresistance of different colorectal human cell lines to 5-FU, oxaliplatin and SN-38, according to the literature.

Drug	Resistance cell lines
5-FU	HCT-116, HT-29, SW48, SW480
Oxaliplatin	HCT-116, HT-29, LoVo, SW48
SN-38	HCT-116, HT-29, SW48

Table 4: IC₅₀ of 5-FU, oxaliplatin and SN-38 in different colorectal human cell lines.

Cell line	IC ₅₀		
	5-FU	Oxaliplatin	SN-38
HT-29	3.6 µM	460 nM	130 nM
HCT-116	17 µM	600 nM	50 nM
LoVo	2.4 µM	1.1 µM	22 nM
SW480	23.1 µM	<i>n.d.</i>	<i>n.d.</i>

After stress exposure, whether physical or chemical, cells were haptenated, *i.e.*, rendered immunogenic through binding of a carrier molecule – capable of conferring immunogenicity – to the stress proteins expressed as a resistance mechanism in response to the stress, whether these stress proteins be free, bound at the surface or within the interior of the cancer cells. The carrier molecule is dinitrophenyl.

Finally, three cell lines with the best proliferative capacities in culture medium with reduced serum (2 % FBS) after stress exposure were selected: LoVo, HCT-116 and HT-29.

Master Cell Bank (MCB): these selected 3 cells lines were cultured in low serum culture condition (2 % FBS) to obtain master cell banks (MCB), mimicking a metabolic stress (nutriment depletion observed in treatment therapy with anti-VEGF antibodies).

5 **Example 2**

Human colorectal cancer vaccine manufacture process

Based on the preliminary results obtained in **Example 1**, we manufactured a vaccine composition comprising multi-stressed, haptenated and non-proliferative HT-29, HCT-116 and LoVo cells.

- 10 For each cell line, starting from the MCB, a DS-A and DS-B were manufactured.

For the DS-A, the MCB of each cell line was thawed (if frozen) and cultured *in vitro*. During the course of their growth phase or their plateau phase, a physical stress comprising a low dose of ionizing radiation (10 Gy) for 5 minutes together with a thermal stress (1 hour at 42°C) was applied. Stress proteins expressed in reaction to this stress
15 were then haptenated with dinitrophenyl. After formulation at 30 million cells/mL and freezing for storage purposes, these cells (comprising the haptenated stress molecules) were inactivated, *i.e.*, rendered non-proliferative, with a high of ionizing radiation (25 Gy), to inhibit cell proliferation while maintaining the cell structure intact.

For the DS-B, the MCB of each cell line was thawed (if frozen) and cultured *in vitro*.
20 During the course of their growth phase or their plateau phase, a chemical stress comprising a chemical stimulation with chemotherapeutic agents was applied: for HT-29 cells, 13 µM oxaliplatin was applied for 72 hours; for HCT-116 cells, 315 nM SN-38 (7-ethyl-10-hydroxy-camptothecin) was applied for 48 hours (or 100 nM SN-38 in most recent experiments with similar final results); and for LoVo cells, 5 µM 5-FU was applied
25 for 48 hours. Stress proteins expressed in reaction to this stress were then haptenated with dinitrophenyl. After formulation at 30 million cells/mL and freezing for storage purposes, these cells (comprising the haptenated stress molecules) were inactivated, *i.e.*, rendered non-proliferative, with a high of ionizing radiation (25 Gy), to inhibit cell proliferation while maintaining the cell structure intact.

A comparison of marker expression was carried out to validate each DS-A and DS-B.

For the three DS-A, the radiation and thermal stress applied to the three cell lines induced the same phenotypic changes in all three DS-A, with an overexpression of HSP70 and CD227 as compared to unstressed cells (**Table 5**). These data were confirmed with experiments performed on several other cell batches similarly treated, which all confirmed HSP70 and CD227 overexpression after radiation and thermal stress.

Table 5: phenotypic changes in HT-29, HCT-116 and LoVo cells, before and after DS-A treatment, in the RCB (10% FBS, non-treated), the MCB (2 % FBS, non-treated) and the DS-A (2 % FBS, treated). % indicate the percentage of cells expressing the marker out of the total number of cells. Values in parenthesis indicate the MFI (mean fluorescence intensity).

Cell line	Treatment		Cmhsp70.1 (HSP70)	CD227 (MUC1)	CD107 (LAMP-1)
HT-29	RCB	10 % FBS Non-treated	51 % (50421)	14 % (9985)	39 % (34915)
	MCB	2 % FBS Non-treated	38 % (131328)	18 % (19461)	97 % (25064)
	DS-A	2 % FBS 10 Gy + 42°C	90 % (> 350000)	60 % (> 13000)	98 % (31617)
HCT-116	RCB	10 % FBS Non-treated	76 % (46853)	16 % (11584)	53 % (19160)
	MCB	2 % FBS Non-treated	70 % (57775)	22 % (15945)	77 % (49348)
	DS-A	2 % FBS 10 Gy + 42°C	99 % (252504)	79 % (2241)	55 % (45097)
LoVo	RCB	10 % FBS Non-treated	75 % (58743)	29 % (13490)	36 % (29510)
	MCB	2 % FBS Non-treated	83 % (39879)	32 % (9023)	81 % (21491)
	DS-A	2 % FBS 10 Gy + 42°C	99 % (200960)	88 % (8246)	85 % (91816)

For the DS-B, treatment of HT-29 cells with oxaliplatin led to an overexpression of CD95, CD107 and CD54 markers as compared to unstressed cells; and treatment of HCT-116 cells with SN-38 (7-ethyl-10-hydroxy-camptothecin) or LoVo cells with 5-FU led to an overexpression of CD66 marker as compared to unstressed cells, and of CD243 in LoVo
5 treated by 5FU (**Table 6**). These data were confirmed with experiments performed on several other cell batches similarly treated, which all confirmed:

- overexpression of CD66 in HCT-116 cells after treatment with SN-38, regardless of the preliminary cells culture conditions (whether in T25 or T225 CellStacks);
- overexpression of CD54, CD95 and CD107 in HT-29 cells after treatment with
10 oxaliplatin; and
- overexpression of CD243 (and, to a lesser extent, CD66) in LoVo cells after treatment with 5-FU.

Finally, the vaccine composition (DP or “STC-1010”) could be formulated by pooling all six DS (3 DS-A – one for each cell line; and 3 DS-B – one for each cell line) in 100 µL
15 doses (each comprising 3 million cells). FACS analysis revealed a high expression (assessed as a percentage of cells expressing the marker out of the total number of cells and as mean fluorescence intensity) of HSP70, CD227, CD95 and CD243 in this final product (**Table 7**).

Table 6: phenotypic changes in HT-29, HCT-116 and LoVo cells, before and after DS-B treatment, in the RCB (10% FBS, non-treated), the MCB (2 % FBS, non-treated) and the DS-B (2 % FBS, treated). % indicate the percentage of cells expressing the marker out of the total number of cells. Values in parenthesis indicate the MFI (mean fluorescence intensity). Two columns for a given marker indicate duplicate results on two different cell batches.

Cell line	Treatment		CD54 (ICAM-1)		CD66 (CEA)		CD95 (FAS receptor)		CD107 (LAMP-1)		CD243 (MDR-1)
HT-29	RCB	10 % FBS	35 % (4472)	57 % (3927)	/	/	8 % (5023)	14 % (7582)	25 % (26420)	30 % (50787)	/
		Non-treated									
	MCB	2 % FBS	47 % (2910)	61 % (2906)	98 % (19255)	/	37 % (5286)	58 % (5277)	46 % (21115)	54 % (20446)	2 % (8310)
HCT-116	DS-B	2 % FBS	96 % (15566)	98 % (16010)	98 % (25417)	/	86 % (13912)	93 % (13121)	96 % (13689)	73 % (12107)	6 % (10837)
	RCB	10 % FBS	/	/	3 %	2 %	/	/	41 % (61194)	53 % (19160)	/
		Non-treated									
HCT-116	MCB	2 % FBS	/	/	5 %	5 %	/	/	33 % (45359)	28 % (29726)	7 % (10726)
	DS-B	Non-treated									
		2 % FBS	/	/	63 % (13130)	81 % (22844)	/	/	35 % (25461)	44 % (20675)	4 % (6693)

LoVo	RCB	10 % FBS Non-treated	/	/	72 % (40510)	72 % (42147)	96 % (21287)	/	41 % (52928)	53 % (29510)	/
	MCB	2 % FBS Non-treated	0 %	/	58 % (88799)	68 % (81155)	68 % (12648)	/	33 % (50995)	28 % (38006)	55 % (7711)
	DS-B	2 % FBS 5-FU	0 %	/	72 % (127233)	88 % (89793)	75 % (17183)	/	35 % (73187)	44 % (46862)	98 % (66072)

Table 7: phenotypic changes in STC-1010 (final product). % indicates the percentage of cells expressing the marker out of the total number of cells. MFI: mean fluorescence intensity. Ranges indicate results obtained for 5 different batches (including batches at 3×10^6 or 3×10^7 cells/mL). CD243 was assessed on a single batch only.

STC-1010		HSP70	CD227	CD54	CD95	CD66	CD243
	%	94-99%	65-88%	13-49%	82-95%	32-60%	96%
	MFI	191647-958843	15229-19809	10145-18830	14174-22645	21916-49224	162377

Example 3**LC-MS/MS identification and relative quantification of proteins in the six intermediate compositions**

During this study, different cell lines from different taxonomies with and without being
 5 subjected to radiations, thermal stress, chemical stress, metabolic stress or combinations thereof *in vitro* were analyzed.

Expressed proteins were identified in specific databases. An adapted sample preparation was performed to improve protein detection.

Individual amounts of detected proteins were evaluated.

10 The normalized signal obtains for each protein was compared to the others.

Material and methods

During the production, cell lines have to be collected according to gene expression after exposures to different stresses (radiations, thermal stress, chemical stress, metabolic stress or combinations thereof). These different stresses induced an overexpression of
 15 antigens. Immunogenicity has also been enhanced by chemically marking the surface proteins with haptens. Protein expression was compared between different cell samples after exposure to different stresses. A summary of the different compositions analyzed in this study is given in **Table 8**.

Table 8

Sample identification	Cell concentration	Viability (Yes/No)	Medium
HT-29 RCB	1 mL $\approx$ 6.10 ⁶ cells	Yes	McCoy's 5A, 2 % FBS, 10 % DMSO
HCT-116 RCB	1 mL $\approx$ 6.10 ⁶ cells	Yes	McCoy's 5A, 2 % FBS, 10 % DMSO
LoVo RCB	1 mL $\approx$ 6.10 ⁶ cells	Yes	F12K, 2 % FBS, 10 % DMSO
HT-29 MCB	1 mL $\approx$ > 6.10 ⁶ cells	Yes	McCoy's 5A, 2 % FBS, 10 % DMSO
HCT-116 MCB	1 mL $\approx$ > 6.10 ⁶ cells	Yes	McCoy's 5A, 2 % FBS, 10 % DMSO
LoVo MCB	1 mL $\approx$ > 6.10 ⁶ cells	Yes	F12K, 2 % FBS, 10 % DMSO

LoVo DS-A	0.5 mL $\approx 5.10^5$ cells	No	Earle's Balanced Salt Solution (EBSS), saccharose, 5 % DMSO
LoVo DS-B	0.5 mL $\approx 5.10^5$ cells	No	EBSS, saccharose, 5 % DMSO
HT-29 DS-A	0.5 mL $\approx 5.10^5$ cells	No	EBSS, saccharose, 5 % DMSO
HT-29 DS-B	0.5 mL $\approx 5.10^5$ cells	No	EBSS, saccharose, 5 % DMSO
HCT-116 DS-A	0.5 mL $\approx 5.10^5$ cells	No	EBSS, saccharose, 5 % DMSO
HCT-116 DS-B	0.5 mL $\approx 5.10^5$ cells	No	EBSS, saccharose, 5 % DMSO
DP	0.75 mL $\approx 3.10^6$ cells	No	EBSS, saccharose, 5 % DMSO

Sample preparation process

The sample preparation process applied to cells pellets in order to detect proteins comprised a chemical lysis of the cells followed by protein digestion. Each generated peptide was separated according to physicochemical property using a nanoflow chromatographic system coupled to a high-resolution mass spectrometer (NanoLC-MS/MS). The main advantage of using this technology is to improve the sensitivity of the instrument and increase the number of identified proteins.

Analytical method

NanoLC-MS/MS analysis were performed in Data Dependent Analysis (DDA) mode, also called shotgun proteomics or peptide mapping. This analytical tool allows acquiring MS and MS/MS spectrum of thousands of peptides through the whole chromatographic separation.

Protein identification

Once the experimental MS and MS/MS spectrum acquired, data process was performed using a software search engine that uses mass spectrometry data to identify proteins from proteome databases. In this case, experimental data was correlated to the full human proteome (SWISS Prot databases) for proteins identification.

To validate the identified proteins, some identification parameters were implemented to ensure the specificity and eliminate false positives. Protein identification was performed with 2 peptides with at least one protein-specific peptide in order to be very stringent about the identified proteins. The identified proteins were the ones present in higher

amount. Moreover, for the identification process, some frequently observed peptide modifications such as methionine oxidation or pyroglutamic acid formation from glutamine were added to the search engine.

A protein amount normalization was also done.

5 Individual quantification evaluation

This strategy comprised spiking a universal calibration curve of synthetic peptides accurately calibrated thanks to the Readybeads™ technology (Anaquant, Villeurbanne, France). This calibration curve allowed the conversion of peptide signal in quantity of all identified proteins. Peptides used in the calibration curve have been chosen for their
10 specificity; their sequences do not correspond to any of the proteome of the cell lines used in bioprocess.

Those standards represent an accurate calibration curve adapted to quantitate an amount of any proteins. The reproducibility and the stability of this calibration curve are ensured thanks to the Readybeads™ technology. All identified peptides and proteins can be
15 normalized using this calibration curve. The normalized signal allowed batch-to-batch comparison.

Results

The 13 samples were treated and analyzed in parallel. Some pellet samples were difficult to lyse, and in order to compare same injected protein quantity, protein quantitation was
20 performed using the Pierce™ BCA Protein Assay Kit before digestion. Despite this quantification step, after sample analysis, injected protein quantity seemed highly different between samples.

This difference could either come from a difference in protein quantity injected, or from a difference in protein dynamic range in the samples.

25 In order to evaluate those hypotheses, AQTBeads added to sample were used as quality control. Readybeads™ quality control showed a good linearity with slope and $r^2 > 0.9$ and quantification through the range 1 to 500 fmol injected protein quantity. Peptide digests were also quantified after LC-MS injection. Results are reported in **Table 9**.

Table 9: quality control parameters.

Sample identification	R ²	Slope	Intercept	QC N°1 (10 fmol)	QC N°2 (5 fmol)	Number of identified proteins	Number of quantified proteins	Peptide quantification (µg/mL)
HT-29 RCB	0.96	1	22.2	7	5	1841	745	420
HCT-116 RCB	0.97	0.97	22.	6	5	1667	491	<LOD
LoVo RCB	0.97	0.98	22.5	9	5	1652	555	376
HT-29 MCB	0.96	0.99	22.3	9	6	1881	790	475
HCT-116 MCB	0.96	0.98	22.4	5	5	1856	646	342
LoVo MCB	0.95	0.95	22.6	6	4	1466	394	698
LoVo DS-A	0.96	1	22.1	11	6	2002	1151	NA
LoVo DS-B	0.95	0.94	22.6	15	4	1102	105	498
HT-29 DS-A	0.95	0.94	22.7	12	4	1179	98	498
HT-29 DS-B	0.94	0.93	22.8	12	4	563	21	265
HCT-116 DS-A	0.95	0.99	22.1	10	6	2118	1369	787
HCT-116 DS-B	0.96	0.96	22.5	16	5	1341	208	465
DP	0.96	0.96	22.6	9	5	1556	333	365

Global protein comparison

Untreated cells

RCB (Research Cell Bank) samples were untreated samples in 10 % FBS medium.

MCB (Master Cell Bank) samples come from RCB with a medium adaptation (2 % FBS).

- 5 The proteins identified in the three human cell lines (HT-29, HCT-116 and LoVo) were compared (**Figs. 1A** and **1B**).

Almost 50 % of identified proteins were commonly identified in all three, untreated, human cell lines. This result was expected since these three cell lines are derived from colon or colorectal carcinomas.

10 *DS-A treatment*

DS-A treatment comprises exposure of the cells to a low dose of ionizing radiation (10 Gy) for 5 minutes together with a thermal stress (1 hour at 42°C). These cells were also subjected to a metabolic stress (*i.e.*, a medium adaptation from 10 % to 2 % FBS between RCB and MCB).

- 15 After this DS-A treatment, some proteins appeared to be over-expressed in all three cell lines (protein proportion ratio between DS-A samples and MCB sample > 2.5 with respect to global protein quantity). These proteins are reported in **Table 10**.

Table 10: overexpressed proteins after DS-A treatment. The ratio is between protein proportion (with respect to global protein quantity) measured in DS-A *versus* MCB samples. Highlighted in bold are proteins specifically identified only in the DS-A samples but not in the MCB samples.

20

Uniprot accession number	Protein	HT-29 DS-A vs. HT-29 MCB	HCT-116 DS-A vs. HCT-116 MCB	LoVo DS-A vs. LoVo MCB
Q15366	Poly(rC)-binding protein 2	590.0	908.0	1111.6
Q92888	Rho guanine nucleotide exchange factor 1	590.0	34.9	148.2
Q14789	Golgin subfamily B member 1	590.0	331.8	129.7
P02749	Beta-2-glycoprotein 1	295.0	52.4	240.8
Q9Y446	Plakophilin-3	295.0	279.4	37.1
Q6ZRV2	Protein FAM83H	295.0	139.7	55.6
O15027	Protein transport protein Sec16A	295.0	87.3	259.4
Q27J81	Inverted formin-2	295.0	52.4	111.2
Q9NQW6	Anillin	295.0	192.1	222.3
Q9H8V3	Protein ECT2	295.0	34.9	74.1
Q15149	Plectin	9.7	3.6	22.5
P58107	Epiplakin	7.7	3.9	1908.2
P46013	Proliferation marker protein Ki-67	5.8	3.3	3.2
Q12907	Vesicular integral-membrane protein VIP36	5.5	261.9	7.5
P11279	Lysosome-associated membrane glycoprotein 1	2.5	8.9	74.1

These ten proteins which are identified only in the DS-A samples but not in the MCB samples reflect that the DS-A treatment led the cells to develop a resistance mechanism by which they have produced stress proteins.

DS-B treatment

- 5 DS-B treatment comprises exposure of the cells to chemotherapeutic agents, namely 13 μ M oxaliplatin applied for 72 hours on HT-29 cells; 315 nM SN-38 (7-ethyl-10-hydroxy-camptothecin) applied for 48 hours on HCT-116 cells; and 5 μ M 5-FU applied for 48 hours on LoVo cells. These cells were also subjected to a metabolic stress (*i.e.*, a medium adaptation from 10 % to 2 % FBS between RCB and MCB).
- 10 After this DS-B treatment, some proteins appeared to be over-expressed (protein proportion ratio between DS-B samples and MCB sample > 2.5 with respect to global protein quantity). These proteins are reported in **Table 11**.

Table 11: overexpressed proteins after DS-B treatment. The ratio is between protein proportion (with respect to global protein quantity) measured in DS-B *versus* MCB samples. Highlighted in bold are proteins specifically identified only in the DS-B samples but not in the MCB samples.

Uniprot accession number	Protein	HT-29 DS-B vs. HT-29 MCB	HCT-116 DS-B vs. HCT-116 MCB	LoVo DS-B vs. LoVo MCB
P02749	Beta-2-glycoprotein 1	10.4	342.0	343.8
P0C0S8	Histone H2A type 1	16674.8	11284.4	21313.9
Q15366	Poly(rC)-binding protein 2	10.4	854.9	687.5
Q86UP2	Kinectin	10.4	2.5	4.4
Q15149	Plectin	35.0	19.7	5.0
O95182	NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 7	10.4	3.3	3.5

Overexpressed proteins seemed mainly associated to DNA repair. In order to go deeper in the results analysis, we decided to focus on membrane proteins.

Focus on membrane protein*Untreated cells*

Tables 12 to 14 represent the membrane proteins overexpressed (protein proportion ratio > 2.5 with respect to global protein quantity) in one human cell line (untreated RCB and MCB) compared with the two others.

Table 12: membrane proteins overexpressed in HT-29 RCB and MCB compared with HCT-116 and LoVo RCB and MCB samples.

Uniprot accession number	Protein
Q9NQC3	Reticulon-4
P04844	Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit 2
P23229	Integrin alpha-6
P51572	B-cell receptor-associated protein 31
O95573	Long-chain-fatty-acid--CoA ligase 3
Q15758	Neutral amino acid transporter B(0)
P53007	Tricarboxylate transport protein, mitochondrial
O60488	Long-chain-fatty-acid--CoA ligase 4
O75746	Calcium-binding mitochondrial carrier protein Aralar1
Q9Y6C9	Mitochondrial carrier homolog 2
Q9HDC9	Adipocyte plasma membrane-associated protein
P07099	Epoxide hydrolase 1
Q13724	Mannosyl-oligosaccharide glucosidase
O00264	Membrane-associated progesterone receptor component 1
A0FGR8	Extended synaptotagmin-2
Q15738	Sterol-4-alpha-carboxylate 3-dehydrogenase, decarboxylating
Q9Y5M8	Signal recognition particle receptor subunit beta
O15270	Serine palmitoyltransferase 2
Q8WY22	BRI3-binding protein
O43676	NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3
Q9Y3D6	Mitochondrial fission 1 protein

Table 13: membrane proteins overexpressed in HCT-116 RCB and MCB compared with HT-29 and LoVo RCB and MCB samples.

Uniprot accession number	Protein
Q96AG4	Leucine-rich repeat-containing protein 59
O43169	Cytochrome b5 type B

Table 14: membrane proteins overexpressed in LoVo RCB and MCB compared with HT-29 and HCT-116 RCB and MCB samples.

Uniprot accession number	Protein
Q13423	NAD(P) transhydrogenase, mitochondrial

5 *DS-A treatment*

DS-A treatment comprises exposure of the cells to a low dose of ionizing radiation (10 Gy) for 5 minutes together with a thermal stress (1 hour at 42°C). These cells were also subjected to a metabolic stress (*i.e.*, a medium adaptation from 10 % to 2 % FBS between RCB and MCB).

- 10 Membrane proteins appeared to be mainly overexpressed rather than underexpressed.

Tables 15 to 17 describe the membrane proteins overexpressed (protein proportion ratio > 2.5 with respect to global protein quantity) in each human cell line (DS-A sample) compared to its corresponding MCB sample.

- 15 **Table 15:** membrane proteins overexpressed in HT-29 DS-A compared with HT-29 MCB.

Uniprot accession number	Protein
Q12907	Vesicular integral-membrane protein VIP36
P11279	Lysosome-associated membrane glycoprotein 1
Q9P2E9	Ribosome-binding protein 1
O43291	Kunitz-type protease inhibitor 2
Q6P1A2	Lysophospholipid acyltransferase 5
P50402	Emerin
Q86UE4	Protein LYRIC
Q9BW60	Elongation of very long chain fatty acids protein 1

P60059	Protein transport protein Sec61 subunit gamma
--------	---

Table 16: membrane proteins overexpressed in HCT-116 DS-A compared with HCT-116 MCB.

Uniprot accession number	Protein
P23229	Integrin alpha-6
P11279	Lysosome-associated membrane glycoprotein 1
Q9P2E9	Ribosome-binding protein 1
Q14739	Delta(14)-sterol reductase LBR
O95202	Mitochondrial proton/calcium exchanger protein
Q5JTH9	RRP12-like protein
Q9BSJ8	Extended syptotagmin-1
P61619	Protein transport protein Sec61 subunit alpha isoform 1
P11166	Solute carrier family 2, facilitated glucose transporter member 1
Q14126	Desmoglein-2
Q86UP2	Kinectin
Q86UE4	Protein LYRIC
O95470	Sphingosine-1-phosphate lyase 1
Q9P0L0	Vesicle-associated membrane protein-associated protein A
Q96N66	Lysophospholipid acyltransferase 7
Q9Y5M8	Sigl recognition particle receptor subunit beta
Q5JTV8	Torsin-1A-interacting protein 1
Q8N766	ER membrane protein complex subunit 1
Q15738	Sterol-4-alpha-carboxylate 3-dehydrogese, decarboxylating
O00264	Membrane-associated progesterone receptor component 1
P16070	CD44 antigen
Q9BUR5	MICOS complex subunit MIC26
Q7Z7H5	Transmembrane emp24 domain-containing protein 4
Q53FV1	ORM1-like protein 2
Q9P2X0	Dolichol-phosphate mannosyltransferase subunit 3
Q07065	Cytoskeleton-associated protein 4
Q9BTV4	Transmembrane protein 43
Q8TC12	Retinol dehydrogese 11

Q15005	Sigl peptidase complex subunit 2
O15270	Serine palmitoyltransferase 2
Q96A33	Coiled-coil domain-containing protein 47
Q9NZ01	Very-long-chain enoyl-CoA reductase
Q14517	Protocadherin Fat 1
Q9Y3B3	Transmembrane emp24 domain-containing protein 7
Q03135	Caveolin-1
Q6ZNB6	NF-X1-type zinc finger protein NFXL1
Q96HR9	Receptor expression-enhancing protein 6
Q9BW72	HIG1 domain family member 2A, mitochondrial
Q8N5G0	Small integral membrane protein 20
Q9P0J0	DH dehydrogese [ubiquinone] 1 alpha subcomplex subunit 13
O43676	DH dehydrogese [ubiquinone] 1 beta subcomplex subunit 3
Q9HC21	Mitochondrial thiamine pyrophosphate carrier
Q9Y3D6	Mitochondrial fission 1 protein
P27338	Amine oxidase [flavin-containing] B
Q15070	Mitochondrial inner membrane protein OXA1L
A1L0T0	2-hydroxyacyl-CoA lyase 2
Q969V3	Nicalin
P20020	Plasma membrane calcium-transporting ATPase 1
Q9NZM1	Myoferlin
P11717	Cation-independent mannose-6-phosphate receptor

Table 17: membrane proteins overexpressed in LoVo DS-A compared with LoVo MCB.

Uniprot accession number	Protein
Q12907	Vesicular integral-membrane protein VIP36
Q9BVI4	Nucleolar complex protein 4 homolog
P67812	Sigl peptidase complex catalytic subunit SEC11A
Q9Y5M8	Sigl recognition particle receptor subunit beta
P10620	Microsomal glutathione S-transferase 1
P16435	DPH--cytochrome P450 reductase
Q9P035	Very-long-chain (3R)-3-hydroxyacyl-CoA dehydratase 3
Q13724	Mannosyl-oligosaccharide glucosidase

Q9BZF1	Oxysterol-binding protein-related protein 8
Q00765	Receptor expression-enhancing protein 5
Q9BUR5	MICOS complex subunit MIC26
O95395	Beta-1,3-galactosyl-O-glycosyl-glycoprotein beta-1,6-N-acetylglucosaminyltransferase 3
Q96IX5	ATP synthase membrane subunit DAPIT, mitochondrial
P56134	ATP synthase subunit f, mitochondrial
Q8N5K1	CDGSH iron-sulfur domain-containing protein 2
Q8N0U8	Vitamin K epoxide reductase complex subunit 1-like protein 1
Q96A26	Protein FAM162A
O75477	Erlin-1
P33121	Long-chain-fatty-acid--CoA ligase 1
O43169	CYB5B_HUMAN Cytochrome b5 type B
Q8TC12	Retinol dehydrogese 11
Q15363	Transmembrane emp24 domain-containing protein 2
Q9NZ01	Very-long-chain enoyl-CoA reductase
Q15125	3-beta-hydroxysteroid-Delta(8),Delta(7)-isomerase
P53985	Monocarboxylate transporter 1
P01889	HLA class I histocompatibility antigen, B alpha chain
O94905	Erlin-2
P21926	CD9 antigen
P03928	ATP synthase protein 8
Q9UDX5	Mitochondrial fission process protein 1
O00483	Cytochrome c oxidase subunit NDUF4
Q9P0J0	DH dehydrogese [ubiquinone] 1 alpha subcomplex subunit 13
O95168	DH dehydrogese [ubiquinone] 1 beta subcomplex subunit 4
O43676	DH dehydrogese [ubiquinone] 1 beta subcomplex subunit 3
Q9Y3D6	Mitochondrial fission 1 protein
Q8WY22	BRI3-binding protein

DS-B treatment

DS-B treatment comprises exposure of the cells to chemotherapeutic agents, namely 13 μ M oxaliplatin applied for 72 hours on HT-29 cells; 315 nM SN-38

(7-ethyl-10-hydroxy-camptothecin) applied for 48 hours on HCT-116 cells; and 5 μ M 5-FU applied for 48 hours on LoVo cells. These cells were also subjected to a metabolic stress (*i.e.*, a medium adaptation from 10 % to 2 % FBS between RCB and MCB).

Membrane proteins appeared to be mainly overexpressed rather than underexpressed.

- 5 **Tables 18 to 20** describe the membrane proteins overexpressed (protein proportion ratio > 2.5 with respect to global protein quantity) in each human cell line (DS-B sample) compared to its corresponding MCB sample.

Table 18: membrane proteins overexpressed in HT-29 DS-B compared with HT-29 MCB.

Uniprot accession number	Protein
Q9P2E9	Ribosome-binding protein 1
P16435	NADPH--cytochrome P450 reductase

- 10 **Table 19:** membrane proteins overexpressed in HCT-116 DS-B compared with HCT-116 MCB.

Uniprot accession number	Protein
Q00325	Phosphate carrier protein, mitochondrial
P16144	Integrin beta-4
P11279	Lysosome-associated membrane glycoprotein 1
P16070	CD44 antigen
Q9P2E9	Ribosome-binding protein 1
P61619	Protein transport protein Sec61 subunit alpha isoform 1
Q86UP2	Kinectin
P04439	HLA class I histocompatibility antigen, A alpha chain
Q96N66	Lysophospholipid acyltransferase 7
O00264	Membrane-associated progesterone receptor component 1
P10620	Microsomal glutathione S-transferase 1
O95168	NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 4
Q14126	Desmoglein-2
P26006	Integrin alpha-3

Q5JTV8	Torsin-1A-interacting protein 1
P20020	Plasma membrane calcium-transporting ATPase 1
O95470	Sphingosine-1-phosphate lyase 1
Q93050	V-type proton ATPase 116 kDa subunit a1
Q15070	Mitochondrial inner membrane protein OXA1L
O43674	NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 5, mitochondrial
Q9BTV4	Transmembrane protein 43
P27338	Amine oxidase [flavin-containing] B
P60468	Protein transport protein Sec61 subunit beta
O14828	Secretory carrier-associated membrane protein 3
Q96A26	Protein FAM162A
Q8TC12	Retinol dehydrogenase 11
Q9NVJ2	ADP-ribosylation factor-like protein 8B
O43676	NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3
Q9P2X0	Dolichol-phosphate mannosyltransferase subunit 3
Q9HC21	Mitochondrial thiamine pyrophosphate carrier
Q53FV1	ORM1-like protein 2

Table 20: membrane proteins overexpressed in LoVo DS-B compared with LoVo MCB.

Uniprot accession number	Protein
Q8TEM1	Nuclear pore membrane glycoprotein 210
Q9NQC3	Reticulon-4
P16435	NADPH--cytochrome P450 reductase
O95573	Long-chain-fatty-acid--CoA ligase 3
O60488	Long-chain-fatty-acid--CoA ligase 4
Q9P0L0	Vesicle-associated membrane protein-associated protein A
P53007	Tricarboxylate transport protein, mitochondrial
Q86UP2	Kinectin
O95292	Vesicle-associated membrane protein-associated protein B/C
Q9P035	Very-long-chain (3R)-3-hydroxyacyl-CoA dehydratase 3
Q12907	Vesicular integral-membrane protein VIP36
Q9NZ01	Very-long-chain enoyl-CoA reductase

P10620	Microsomal glutathione S-transferase 1
O43169	Cytochrome b5 type B
Q96IX5	ATP synthase membrane subunit DAPIT, mitochondrial
P56134	ATP synthase subunit f, mitochondrial
P00403	Cytochrome c oxidase subunit 2
P08574	Cytochrome c1, heme protein, mitochondrial
P01889	HLA class I histocompatibility antigen, B alpha chain
Q86UE4	Protein LYRIC
P53985	Monocarboxylate transporter 1
O00483	Cytochrome c oxidase subunit NDUFA4
Q13724	Mannosyl-oligosaccharide glucosidase
P67812	Signal peptidase complex catalytic subunit SEC11A
Q96A26	Protein FAM162A
Q15363	Transmembrane emp24 domain-containing protein 2
Q8TC12	Retinol dehydrogenase 11
O75477	Erlin-1
Q8WY22	BRI3-binding protein
Q15125	3-beta-hydroxysteroid-Delta(8),Delta(7)-isomerase
Q00765	Receptor expression-enhancing protein 5
Q8N5K1	CDGSH iron-sulfur domain-containing protein 2
P21926	CD9 antigen
A0FGR8	Extended synaptotagmin-2
O14975	Very long-chain acyl-CoA synthetase
O43291	Kunitz-type protease inhibitor 2
Q13740	CD166 antigen
P33121	Long-chain-fatty-acid--CoA ligase 1
Q08722	Leukocyte surface antigen CD47
O15533	Tapasin
O95395	Beta-1,3-galactosyl-O-glycosyl-glycoprotein beta-1,6-N-acetylglucosaminyltransferase 3

Example 4**LC-MS/MS identification and relative quantification of proteins in the final composition**

This study aimed at identifying differentially expressed protein in the final vaccine
5 composition (DP, comprising all six DS described above, and pooled together: 3 DS-A –
one for each cell line; and 3 DS-B – one for each cell line).

Fig. 2 shows the differential number of proteins expressed in the HCT-116 MCB, DS-A,
DS-B and in the final product (DP) comprising all six DS described above (hence
including the HCT-116 DS-A and DS-B), by comparison to the initial HCT-116 RCB
10 cultured in classical conditions (with 10 % FBS). “+” indicates proteins that are newly
expressed, “-” indicates the number of proteins which are no longer expressed at all.

Fig. 3 shows the number of proteins over- and under-expressed in the HCT-116 MCB,
DS-A and DS-B, by comparison to the initial HCT-116 RCB cultured in classical
conditions (with 10 % FBS).

15 **Fig. 4** shows the differential number of proteins expressed in the HT-29 MCB, DS-A,
DS-B and in the final product (DP) comprising all six DS described above (hence
including the HT-29 DS-A and DS-B), by comparison to the initial HT-29 RCB cultured
in classical conditions (with 10 % FBS). “+” indicates proteins that are newly expressed,
“-” indicates the number of proteins which are no longer expressed at all.

20 **Fig. 5** shows the number of proteins over- and under-expressed in the HT-29 MCB, DS-A
and DS-B, by comparison to the initial HT-29 RCB cultured in classical conditions (with
10 % FBS).

Fig. 6 shows the differential number of proteins expressed in the LoVo MCB, DS-A,
DS-B and in the final product (DP) comprising all six DS described above (hence
25 including the LoVo DS-A and DS-B), by comparison to the initial LoVo RCB cultured
in classical conditions (with 10 % FBS). “+” indicates proteins that are newly expressed,
“-” indicates the number of proteins which are no longer expressed at all.

Fig. 7 shows the number of proteins over- and under-expressed in the LoVo MCB, DS-A and DS-B, by comparison to the initial LoVo RCB cultured in classical conditions (with 10 % FBS).

A comparison of these LC/MS data is given in **Table 21**, indicating the number and percentage of proteins which are exclusively present in the final composition (DP) (*i.e.*, not identified in any of the three cell lines at RCB stage), which are over-expressed in the final composition (DP) (in comparison to the three cell lines at RCB stage), and which are similarly or less expressed in the final composition (DP) (in comparison to the three cell lines at RCB stage).

Table 21: comparison of the relative expression of proteins identified in the final composition (DP), in comparison to the three cell lines at RCB stage. Proteins identified in each RCB but not identified in the final composition (DP) are not counted. Comparison was done by applying a multiplication factor 3 for the final composition (DP), to take the dilution factor 3 due to the mixing of the 3 cell lines in this DP into account.

Differential expression DP versus RCBs	Number of proteins	%
Similarly or less expressed in the DP	1419	91%
Over-expressed in the DP	49	3%
Exclusively present in the DP	88	6%
	1556	100%

Table 22 shows the list of proteins which are exclusively found in the final composition (DP) but not in any of the RCBs. The presence of any of these proteins is therefore characteristic of the final composition having undergone all the different treatments described above.

Table 22: proteins exclusively found in the final composition (DP).

Uniprot accession number	Protein
P08631	Tyrosine-protein kinase HCK
O95758	Polypyrimidine tract-binding protein 3

Q63HN8	E3 ubiquitin-protein ligase RNF213
Q9BRL6	Serine/arginine-rich splicing factor 8
Q9H223	EH domain-containing protein 4
Q8WWI1	LIM domain only protein 7
Q9H9Y6	DNA-directed RNA polymerase I subunit RPA2
Q9Y6K5	2'-5'-oligoadenylate synthase 3
O75717	WD repeat and HMG-box DNA-binding protein 1
P02749	Beta-2-glycoprotein 1
P36873	Serine/threonine-protein phosphatase PP1-gamma catalytic subunit
Q9NQW6	Anillin
O43795	Unconventional myosin-Ib
O94973	AP-2 complex subunit alpha-2
P24941	Cyclin-dependent kinase 2
P42224	Signal transducer and activator of transcription 1-alpha/beta
Q14671	Pumilio homolog 1
Q8NE71	ATP-binding cassette sub-family F member 1
Q9H0H5	Rac GTPase-activating protein 1
Q9P2M7	Cingulin
O00186	Syntaxin-binding protein 3
O43772	Mitochondrial carnitine/acylcarnitine carrier protein
O60684	Importin subunit alpha-7
O75676	Ribosomal protein S6 kinase alpha-4
P20339	Ras-related protein Rab-5A
P23921	Ribonucleoside-diphosphate reductase large subunit
P24666	Low molecular weight phosphotyrosine protein phosphatase
P31350	Ribonucleoside-diphosphate reductase subunit M2
P40616	ADP-ribosylation factor-like protein 1
P50570	Dynamin-2
P51153	Ras-related protein Rab-13
P53990	IST1 homolog
P85037	Forkhead box protein K1
Q00796	Sorbitol dehydrogenase
Q07817	Bcl-2-like protein 1

Q14134	Tripartite motif-containing protein 29
Q14807	Kinesin-like protein KIF22
Q15800	Methylsterol monooxygenase 1
Q6NZI2	Caveolae-associated protein 1
Q6PJG6	BRCA1-associated ATM activator 1
Q6ZRV2	Protein FAM83H
Q6ZXV5	Protein O-mannosyl-transferase TMTC3
Q70UQ0	Inhibitor of nuclear factor kappa-B kinase-interacting protein
Q7Z2W4	Zinc finger CCCH-type antiviral protein 1
Q86U38	Nucleolar protein 9
Q86V48	Leucine zipper protein 1
Q8IXK0	Polyhomeotic-like protein 2
Q8IZW8	Tensin-4
Q8NC56	LEM domain-containing protein 2
Q8TEX9	Importin-4
Q92888	Rho guanine nucleotide exchange factor 1
Q96HC4	PDZ and LIM domain protein 5
Q96QD9	UAP56-interacting factor
Q96T76	MMS19 nucleotide excision repair protein homolog
Q99661	Kinesin-like protein KIF2C
Q9BQ69	ADP-ribose glycohydrolase MACROD1
Q9BW19	Kinesin-like protein KIFC1
Q9H6R0	ATP-dependent RNA helicase DHX33
Q9HC35	Echinoderm microtubule-associated protein-like 4
Q9NVI1	Fanconi anemia group I protein
Q9NZN3	EH domain-containing protein 3
Q9NZT2	Opioid growth factor receptor
Q9UEY8	Gamma-adducin
Q9UH17	DNA dC- dU-editing enzyme APOBEC-3B
Q9Y639	Neuroplastin
Q9Y6M5	Zinc transporter 1
O76003	Glutaredoxin-3
P04183	Thymidine kinase, cytosolic

P29966	Myristoylated alanine-rich C-kinase substrate
P30085	UMP-CMP kinase
P40121	Macrophage-capping protein
P52926	High mobility group protein HMGI-C
P53814	Smoothelin
Q15102	Platelet-activating factor acetylhydrolase IB subunit gamma
Q16594	Transcription initiation factor TFIID subunit 9
Q3SXM5	Inactive hydroxysteroid dehydrogenase-like protein 1
Q53HL2	Borealin
Q71RC2	La-related protein 4
Q8N183	NADH dehydrogenase [ubiquinone] 1 alpha subcomplex assembly factor 2
Q96GD4	Aurora kinase B
Q9H4G4	Golgi-associated plant pathogenesis-related protein 1
Q9UG63	ATP-binding cassette sub-family F member 2
Q9UH62	Armadillo repeat-containing X-linked protein 3
Q9UHA4	Ragulator complex protein LAMTOR3
Q9UHI6	Probable ATP-dependent RNA helicase DDX20
Q9UI12	V-type proton ATPase subunit H
Q9Y376	Calcium-binding protein 39

A further analysis of the 1556 proteins identified in the final composition (DP) has shown that 97 of them are of particular interest and can be categorized in 12 superfamilies based on biological, clinical, and cancer prognostic value. Out of these 97 proteins, 52 are overexpressed in the final composition (DP) in comparison to the three cell lines at RCB stage, and 8 are exclusively found in the final composition (DP) but not in any of the RCBs. **Table 23** summaries the number of proteins in each of these 12 superfamilies.

Table 23: summary of 97 proteins of biological, clinical, and cancer prognostic value.

Type	Total number	Over-expressed in the DP	Exclusively present in the DP
Adhesion	3		
Antigen	12	5	

ATP-binding cassette	4	4	2
BCL	2	2	1
COX	7	4	
EGFR	2		
HSP	7	5	
Inhibitor	7	2	1
MUC	1		
RAS-related	27	16	2
Repair	7	3	1
Transporter	18	11	1
	97	52	8

“Adhesion” corresponds to CAM proteins (cell adhesion molecules), including IgCAMs (such as ICAM1), cadherins, integrins, and selectins.

“ATP binding cassette” corresponds to transmembrane proteins of the transport system superfamily, which are linked with the drug resistance phenomena.

- 5 “BCL” corresponds to proteins that regulate cell death, being either pro-apoptotic (such as BAX, BAK1/Bcl-2 homologous antagonist killer, and Bcl-2-associated death promoter) or anti-apoptotic (such as Bcl-2, and Bcl-xL).

“COX” corresponds to cytochrome C oxidase proteins (also termed “complex IV”), which are proteins from the terminal component of the mitochondrial respiratory chain.

- 10 Mutations in cytochrome C oxidase is involved in cancer (in particular in cytochrome C oxidase subunit 4).

“EGFR” corresponds to epidermal growth factor, involved in the pathogenesis and progression of different carcinoma types.

- 15 “HSP” corresponds to heat shock proteins, which are a class of proteins overexpressed in a wide range of human cancers and implicated in tumor cell proliferation, differentiation, invasion, metastasis, death, and recognition by the immune system.

“Inhibitor” corresponds to proteins linked with pro- or anti-cancer proliferation.

“MUC” corresponds to mucin proteins, which are heavily glycosylated proteins. MUC13 in particular is frequently and aberrantly expressed in a variety of epithelial carcinomas, including gastric, colorectal, and ovarian cancers.

- 5 “RAS-related” corresponds to Rap GTP-binding proteins, a type of small GTPase. More than 30 % of all human cancers – including 95 % of pancreatic cancers and 45 % of colorectal cancers – are driven by mutations of the RAS family of genes.

“Repair” corresponds to proteins linked with tumor progression.

“Transporter” corresponds to transmembrane proteins with function in drug resistance.

- 10 **Table 24** identifies these 52 proteins that are overexpressed, among which 8 are exclusively expressed, in the final composition (DP) versus the RCBs.

Table 24

Protein class	Uniprot accession number	Protein	Membrane protein? (Y/N)	Exclusively expressed in DP? (Y/N)
Antigen	P01889	HLA class I histocompatibility antigen, B alpha chain	Y	N
Antigen	P04439	HLA class I histocompatibility antigen, A alpha chain	Y	N
Antigen	P10321	HLA class I histocompatibility antigen, C alpha chain	Y	N
Antigen	P21926	CD9 antigen	Y	N
Antigen	P23497	Nuclear autoantigen Sp-100	N	N
ATP-binding cassette	P28288	ATP-binding cassette sub-family D member 3	Y	N
ATP-binding cassette	P61221	ATP-binding cassette sub-family E member 1	N	N
ATP-binding cassette	Q8NE71	ATP-binding cassette sub-family F member 1	N	Y
ATP-binding cassette	Q9UG63	ATP-binding cassette sub-family F member 2	N	Y
BCL	Q07817	Bcl-2-like protein 1	Y	Y
BCL	Q9NYF8	Bcl-2-associated transcription factor 1	N	N
COX	P00403	Cytochrome c oxidase subunit 2	Y	N
COX	P13073	Cytochrome c oxidase subunit 4 isoform 1, mitochondrial	Y	N
COX	P20674	Cytochrome c oxidase subunit 5A, mitochondrial	N	N
COX	Q15067	Peroxisomal acyl-coenzyme A oxidase 1	N	N
HSP	P04792	Heat shock protein beta-1	N	N
HSP	P08238	Heat shock protein HSP 90-beta	N	N

HSP	P11142	Heat shock cognate 71 kDa protein	N	N	N
HSP	P17066	Heat shock 70 kDa protein 6	N		N
HSP	P34932	Heat shock 70 kDa protein 4	N		N
Inhibitor	P13489	Ribonuclease inhibitor	N		N
Inhibitor	Q70UQ0	Inhibitor of nuclear factor kappa-B kinase-interacting protein	Y		Y
RAS related	P20339	Ras-related protein Rab-5A	N		Y
RAS related	P20340	Ras-related protein Rab-6A	N		N
RAS related	P51148	Ras-related protein Rab-5C	N		N
RAS related	P51149	Ras-related protein Rab-7a	N		N
RAS related	P51153	Ras-related protein Rab-13	N		Y
RAS related	P57735	Ras-related protein Rab-25	N		N
RAS related	P59190	Ras-related protein Rab-15	N		N
RAS related	P61006	Ras-related protein Rab-8A	N		N
RAS related	P61026	Ras-related protein Rab-10	N		N
RAS related	P61224	Ras-related protein Rap-1b	N		N
RAS related	P62820	Ras-related protein Rab-1A	N		N
RAS related	P62834	Ras-related protein Rap-1A	N		N
RAS related	P63000	Ras-related C3 botulinum toxin substrate 1	N		N
RAS related	Q92930	Ras-related protein Rab-8B	N		N
RAS related	Q9NP72	Ras-related protein Rab-18	N		N

RAS related	Q9Y3L5	Ras-related protein Rap-2c	N	N	N
Repair	P12956	X-ray repair cross-complementing protein 6	N	N	N
Repair	P52701	DNA mismatch repair protein Msh6	N	N	N
Repair	Q96T76	MMS19 nucleotide excision repair protein homolog	N	N	Y
Transporter	O15027	Protein transport protein Sec16A	N	N	N
Transporter	P05023	Sodium/potassium-transporting ATPase subunit alpha-1	Y	N	N
Transporter	P11166	Solute carrier family 2, facilitated glucose transporter member 1	Y	N	N
Transporter	P53007	Tricarboxylate transport protein, mitochondrial	Y	N	N
Transporter	P53985	Monocarboxylate transporter 1	Y	N	N
Transporter	P60468	Protein transport protein Sec61 subunit beta	Y	N	N
Transporter	P61619	Protein transport protein Sec61 subunit alpha isoform 1	Y	N	N
Transporter	Q5JRA6	Transport and Golgi organization protein 1 homolog	Y	N	N
Transporter	Q8TB61	Adenosine 3'-phospho 5'-phosphosulfate transporter 1	Y	N	N
Transporter	Q92973	Transportin-1	N	N	N
Transporter	Q9Y6M5	Zinc transporter 1	Y	Y	Y

Table 25 shows the implication of these 97 proteins of biological, clinical, and cancer prognostic value in various types of cancers. As seen, the final composition (DP) expresses markers that are not only linked to colorectal cancer, as could be expected given that HT-29, HCT-116 and LoVo cells are colorectal cell lines, but also to other types of cancers. This suggests that the composition described herein could be useful for treating not only colorectal cancer but also a wide variety of other cancers.

Table 25

Cancer type	Number of identified markers
Breast	6
Colorectal	8
Liver	37
Renal	48
Pancreatic	15
Endometrial	13
Head and neck	4
Ovarian	8
Lung	6
Cervical	2
Melanoma	1
Glioma	1

Example 5

LC-MS/MS identification and relative quantification of surface proteins in 2 intermediate compositions and in the final product

This study aimed at characterizing the surface proteome of tumoral cells at different production steps and finally identifying differentially expressed surface proteins in one of the starting materials (HT-29 MCB), in the 2 corresponding intermediate compositions (HT-29 DS-A and HT-29 DS-B) and in the final product (DP) comprising all 6 DS described above (hence including the HT-29 DS-A and HT-29 DS-B).

Material and methods

Similar type of product (DS /DP) as those described in the previous examples 3 & 4 have been analyzed. The same steps have been applied specifically exposures to different stresses (radiations, thermal stress, chemical stress, metabolic stress or combinations thereof). These different stresses induced an overexpression of antigens. Immunogenicity has also been enhanced by chemically marking the surface proteins with haptens. Protein expression was compared between different cell samples after exposure to different stresses. A summary of the different compositions analyzed in this study is given in **Table 26**.

10 **Table 26**

Sample identification	Product	Volume (mL)	Cell concentration (cells/mL)
HT-29 MCB	HT-29 cultured in McCoy's 5A, 2 % FBS, 10 % DMSO	1	6×10^6
HT-29 DS-A	HT-29 in EBSS, saccharose, 5 % DMSO, stressed by radiation and thermic choc, then haptenated	0.5	3.88×10^7
HT-29 DS-B	HT-29 in EBSS, saccharose, 5 % DMSO, stressed by chemical stress, then haptenated	0.15	2.6×10^7
DP	Drug product comprising HT-29 DS-A and HT-29 DS-B, together with HCT-116 DS-A, HCT-116 DS-B, LoVo DS-A and LoVo DS-B	2	3×10^6

Samples treatment

When possible, an amount of 6×10^6 cells (otherwise, whole sample was used) were gently washed with PBS and were then biotinylated using Pierce™ Cell Surface Protein Biotinylation. Cell were then lysed and proteins were isolated using Pierce™ Cell Surface Protein Biotinylation and Isolation Kit (Thermo Scientific, Catalog Numbers A44390). Both preparations steps were performed according to manufacturer's instructions.

Proteins were then precipitated with methanol-chloroform, precipitates were washed with methanol, dried and solubilized in iST LYSE (PreOmics GmbH) buffer by micro-cavitation (Bioruptor Pico, Diagenode).

Proteins were digested using LysC and trypsin. Peptides were purified using a mixed-mode reverse phase cation exchanger SPE column (PreOmics GmbH), dried and solubilized in 100 µL of 3% acetonitrile 0.1% formic acid aqueous solution.

- Peptides concentration was determined using BCA method (**Table 27**). Despite low measured peptides concentrations, their quantities were sufficient for this feasibility study.

Table 27

Sample identification	Mean peptide concentration (µg/µL)
HT-29 MCB	0.029
HT-29 DS-A	0.050
HT-29 DS-B	0.049
DP	0.048

Analytical method

LC-MS/MS

- 250 ng of peptides were injected in triplicate for each sample.

Chromatography was performed using an Ultimate 3000 (Dionex) equipment using PepMap100 C18 (75 µm × 50 cm, 2 µm material) column applying a 2.5%-to-35% acetonitrile 120-minute gradient at a flow rate of 300 nL/minute after a 3-minute trapping step on precolumn.

- Data were acquired using a Q-Exactive (Thermo) mass spectrometer using experimental settings described in **Table 28**. MS/MS scan was performed on the 10 most intense ions of each cycle, 6545 cycles were performed, thus an average of 17 cycles per chromatographic peak.

Table 28

Source	Voltage	Sweep gas	Transfer tube temperature
	1900V	0 psi	275°C
Mass spectrometer	Resolution	Accumulation time	Normalized collision energy
MS scan	70 000	60 ms	---
MS/MS scan	17 500	60 ms	28

Protein identification

Data were processed with Proteome Discoverer 2.4.

Proteins were identified using the SEQUEST-HT algorithm against a database gathering
 5 human reference proteome mined from NeXtProt and cRAP contaminant database depleted of human proteins.

Search parameters were:

- enzyme = trypsin (full);
- allowed miscleavage = 2;
- 10 - precursor error tolerance = 10 ppm;
- fragment error tolerance = 0.02 Da;
- dynamic modification = oxidation (M), deamidation (N/Q), CAMthiopropionyl (K);
- protein terminus modification = acetylation, CAMthiopropionyl;
- 15 - static modification = carbamidomethyl (C).

False Discovery Rate (FDR) determination was made using Percolator algorithm.

All spectra reported with a confidence less than high by SEQUEST-HT, *i.e.*, considered as not identified, were processed a second time by the same algorithm against the same database than above, but using modified settings:

- 20 - enzyme = trypsin_R (semi);
- allowed miscleavage = 2;

- precursor error tolerance = 10 ppm;
- fragment error tolerance = 0.02 Da;
- dynamic modification = oxidation (M), deamidation (N/Q), CAMthiopropionyl (K);
- 5 - protein terminus modification = acetylation, CAMthiopropionyl;
- static modification = carbamidomethyl (C).

False Discovery Rate (FDR) determination was made using Percolator algorithm.

- Proteins were considered as part of “membrane” or “cell surface” when these keywords were reported as annotations in the “cellular component” field of the Thermo Protein
- 10 Center database. It has to be noted that several cellular components may be reported for the same protein. Furthermore, the term “membrane” can refer to membrane other than plasma membrane (*e.g.*, nucleus membrane, organelle membrane, vesicle membrane, etc.).

Protein quantification

- 15 Data were processed using Minora and feature mapper for Proteome Discoverer 2.4 software.

Peak integration parameters were:

- post-acquisition recalibration = true (fine parameters);
- minimum trace length = 5;
- 20 - max delta RT for isotope = 0.2 minutes;
- PSM confidence level for integration = high.

Chromatographic alignment parameters were:

- RT alignment = true;
- parameter tuning = fine;
- 25 - max RT shift = 5 minutes;
- mass tolerance = 10 ppm.

Feature mapping parameters were:

- RT tolerance = automatic;

- mass tolerance = automatic;
- S/N threshold = 2.

Statistical analyses were performed by using Precursors Ions quantifier node for Proteome Discoverer 2.4 software.

5 General quantification settings were:

- peptide to use = unique + RAZOR (unique meaning peptides that are not shared by different proteins or protein groups; RAZOR meaning peptides shared by multiple protein's groups but only used to quantify protein with the largest number of unique peptides and with the longest amino acid sequence);
- 10 - consider proteins groups for peptide uniqueness = true;
- reject quan results with missing channels = false.

Precursor quantification settings were:

- precursor abundance based on = area;
- min number replicate feature = 50% (peptides must be detected in at least 50% of
- 15 sample of one group for be use in quantification).

Normalization settings were:

- normalization mode = total Peptide amount (calculates the total sum of abundance values for each injection over all peptides identified, the injection with the highest total abundance is used as reference to correct abundance values in all other
- 20 injections by a constant factor per injection, so that at the end the total abundance is the same for all injections).

Quan rollup hypothesis testing settings were:

- ratio calculation = pairwise ratio-based (peptides ratios are calculated as geometric median of all combinations ratio from all replicates for selected study factor. The
- 25 proteins ratio is subsequently calculated as geometric median of peptides group ratio);
- Imputation mode = replicate-based resampling (missing values are replaced with random values sampled from distributions centered around medians of detected values of (technical, biological) replicates);

- hypothesis test = *t*-test (background-based).

The hypothesis test giving the *p*-value is a *t*-background test (or ANOVA background). This test has been based on the assumption that most protein abundances do not vary in response to stimulus, in proteomics. This method determines a rank of protein ratios considered mainly constant between conditions before testing each protein abundance ratio against median and variance of this constant population. This test is useful with studies having missing values and can be used only when hundreds of proteins are identified. It does not require technical replicate.

Results

10 Number of proteins detection

The number of proteins and peptides identified in each sample are displayed in **Table 29** and **Fig. 8**.

Table 29

Sample identification	PEP: number of peptides PRO: number of proteins
HT-29 MCB	EP : 5 828 RO : 1 418
HT-29 DS-A	EP : 15 368 RO : 2 373
HT-29 DS-B	EP : 13 928 RO : 2 290
DP	EP : 10 121 RO : 1 758

Relative quantification of proteins

- 15 Four comparisons were done to compare:
- HT-29 DS-A to HT-29 DS-B,
 - HT-29 DS-A to HT-29 MCB,
 - HT-29 DS-B to HT-29 MCB, and

- HT-29 MCB to DP.

Note 1: due to biological or technical variations and/or the stochastic nature of mass spectrometric acquisition of trace data, some protein abundance values may be lost. Thus, we consider as quantifiable, proteins whose quantifiable peptides are present in at least 50% of the injections in either group.

Note 2: for some proteins or peptides, the abundance ratios are equal to 1000 or 0.001. These values are arbitrary. In the first case, they mean that this protein or peptide has been quantified only in the numerator condition; in the second case, only in the denominator condition.

- 10 Note 3: proteins not identified on the basis of their peptides' MS/MS fragmentation spectra can still be quantified through "match between run". This quantification is based on the similarity of the XIC (MS1) chromatographic characteristics of their peptides (exact mass, retention time) with those of peptides identified in at least one acquisition of the same experiment.
- 15 Note 4: protein abundance is considered statistically and significantly different when the associated p -value is less than or equal to 0.05 with an abundance ratio less than or equal to 0.5 or greater than or equal to 2.

Fig. 9A-D show the proteins distribution according to p -values and fold changes in each of the four comparisons.

- 20 **Fig. 10** shows the number of proteins from membrane and/or cell surface with statistical and significant difference in each of the four comparisons.

For all injections, more than 65% of the proteins identified were associated to the terms "membrane" or "cell surface" in the "cellular component" field of the Thermo Protein Center database (**Fig. 8**). As a comparison, in non-biotinylated HeLa cells, a lower proportion of identified proteins (51%, *i.e.*, 816 proteins out of 3562) are associated to the terms "membrane" or "cell surface" in the same database. This tends to indicate successful enrichment of surface proteome in all four samples.

Relative quantification and comparisons between samples showed that abundances of about 29% to 34% of proteins significantly vary when stressed/haptenated cells (DS-A or DS-B) or drug product (DP) are compared to non-treated cells (MCB) (**Fig. 9A-C**, respectively). A lower proportion (15%) is observed when both stressed/haptenated cell samples (DS-A and DS-B) are compared to each other (**Fig. 9D**).

More than one half of significantly over- or under-expressed proteins are annotated as membrane and cell surface proteins (**Fig. 10**).

Comparison results

HT-29 DS-A/HT-29 MCB

- 10 Based on the raw data, an abundance ratio DS-A/MCB ≥ 1000 has been selected to sort the overexpressed membrane or cell surface proteins after a stress by radiation and thermic choc, followed by haptenation.

Considering the cells membranes proteins, 455 proteins were identified as overexpressed in the DS-A compared to the MCB with an abundance ratio ≥ 1000 .

- 15 Considering the specific cell surface proteins, 127 proteins were identified as overexpressed in the DS-A compared to the MCB with an abundance ratio ≥ 2 , and 38 of them with an abundance ratio of 1000 (**Table 30**).

Table 30

Uniprot accession number	Protein
P54652	Heat shock-related 70 kDa protein
O76027	Annexin
Q4KMQ2	Anoctamin-6
O75054	Immunoglobulin superfamily member 3
P02787	Serotransferrin
O14763	Tumor necrosis factor receptor superfamily member 10B
P10909	Clusterin
P09958	Furin
P04233	HLA class II histocompatibility antigen gamma chain

Q6YHK3	CD109 antigen
Q9Y696	Chloride intracellular channel protein 4
Q14517	Protocadherin Fat 1
P49281	Natural resistance-associated macrophage protein 2
O00220	Tumor necrosis factor receptor superfamily member 10A
O15484	Calpain-5
Q29983	MHC class I polypeptide-related sequence A
P09429	High mobility group protein B1
O95858	Tetraspanin-15
Q9BZM5	UL16-binding protein 2
P26010	Integrin beta-7
Q15465	Sonic hedgehog protein
O15455	Toll-like receptor 3
P02749	Beta-2-glycoprotein 1
P13726	Tissue factor
P29122	Proprotein convertase subtilisin/kexin type 6
Q9UNN8	Endothelial protein C receptor
Q8IWT6	Volume-regulated anion channel subunit LRRC8A
Q9NYQ7	Cadherin EGF LAG seven-pass G-type receptor 3
Q13433	Zinc transporter ZIP6
P28067	HLA class II histocompatibility antigen, DM alpha chain
Q9UPY5	Cystine/glutamate transporter
Q9HBW0	Lysophosphatidic acid receptor 2
P18827	Syndecan-1
Q12891	Hyaluronidase-2
P13612	Integrin alpha-4
P04196	Histidine-rich glycoprotein
P01137	Transforming growth factor beta-1 proprotein
P16035	Metalloproteinase inhibitor 2

HT-29 DS-A/HT-29 MCB

Based on the raw data, an abundance ratio DS-B/MCB ≥ 1000 has been selected to sort the overexpressed membrane or cell surface proteins after a chemical choc, followed by haptenation.

- 5 Considering the cells membranes proteins, 430 proteins were identified as overexpressed in the DS-B compared to the MCB with an abundance ratio ≥ 1000 .

Considering the specific cell surface proteins, 127 proteins were identified as overexpressed in the DS-B compared to the MCB with an abundance ratio ≥ 2 , and 37 of them with an abundance ratio of 1000 (**Table 31**).

10 **Table 31**

Uniprot accession number	Protein
Q6YHK3	CD109 antigen
P04233	HLA class II histocompatibility antigen gamma chain
P30511	HLA class I histocompatibility antigen, alpha chain F
O75330	Hyaluronan mediated motility receptor
P26012	Integrin beta-8
P05106	Integrin beta-3
P29122	Proprotein convertase subtilisin/kexin type 6
P10909	Clusterin
P02787	Serotransferrin
P49281	Natural resistance-associated macrophage protein 2
Q29983	MHC class I polypeptide-related sequence A
O14763	Tumor necrosis factor receptor superfamily member 10B
Q9UNN8	Endothelial protein C receptor
Q9NYQ7	Cadherin EGF LAG seven-pass G-type receptor 3
O00220	Tumor necrosis factor receptor superfamily member 10A
Q9UPY5	Cystine/glutamate transporter
P13726	Tissue factor

P01137	Transforming growth factor beta-1 proprotein
O75054	Immunoglobulin superfamily member 3
Q4KMQ2	Anoctamin-6
P16035	Metalloproteinase inhibitor 2
O15455	Toll-like receptor 3
Q8IWT6	Volume-regulated anion channel subunit LRRC8A
O95858	Tetraspanin-15
Q13433	Zinc transporter ZIP6
P09958	Furin
Q14517	Protocadherin Fat 1
Q12891	Hyaluronidase-2
Q9HBW0	Lysophosphatidic acid receptor 2
P09429	High mobility group protein B1
Q9Y696	Chloride intracellular channel protein 4
Q9BZM5	UL16-binding protein 2
O15484	Calpain-5
O76027	Annexin A9
P04196	Histidine-rich glycoprotein
P13612	Integrin alpha-4
P54652	Heat shock-related 70 kDa protein 2

DP/HT-29 MCB

Note: this comparison is not representative of the real protein overexpression due to the composition of the DP that gathers HT-29 DS-A and HT-29 DS-B, but also HCT-116 DS-A, HCT-116 DS-B, LoVo DS-A and LoVo DS-B; hence, a dilution factor would need to be taken account for the MCB comparison.

Based on the raw data, an abundance ratio $DP/MCB \geq 1000$ has been selected to sort the overexpressed membrane or cell surface proteins.

Considering the cells membranes proteins, 343 proteins were identified as overexpressed in the DP compared to the MCB with an abundance ratio ≥ 1000 .

Considering the specific cell surface proteins, 112 proteins were identified as overexpressed in the DP compared to the MCB with an abundance ratio ≥ 2 , and 34 of them with an abundance ratio of 1000 (**Table 32**).

Table 32

Uniprot accession number	Protein
P02749	Beta-2-glycoprotein 1
P28067	HLA class II histocompatibility antigen, DM alpha chain
Q6YHK3	CD109 antigen
P04233	HLA class II histocompatibility antigen gamma chain
P30511	HLA class I histocompatibility antigen, alpha chain F
O75330	Hyaluronan mediated motility receptor
P26012	Integrin beta-8
P05106	Integrin beta-3
P29122	Proprotein convertase subtilisin/kexin type 6
P10909	Clusterin
P02787	Serotransferrin
Q29983	MHC class I polypeptide-related sequence A
O14763	Tumor necrosis factor receptor superfamily member 10B
Q9UNN8	Endothelial protein C receptor
Q9NYQ7	Cadherin EGF LAG seven-pass G-type receptor 3
O00220	Tumor necrosis factor receptor superfamily member 10A
P13726	Tissue factor
O75054	Immunoglobulin superfamily member 3
Q4KMQ2	Anoctamin-6
O15455	Toll-like receptor 3
O95858	Tetraspanin-15
Q13433	Zinc transporter ZIP6
P09958	Furin
Q14517	Protocadherin Fat 1
Q9HBW0	Lysophosphatidic acid receptor 2
Q9Y696	Chloride intracellular channel protein 4
Q9BZM5	UL16-binding protein 2

O15484	Calpain-5
O76027	Annexin A9
P04196	Histidine-rich glycoprotein
P13612	Integrin alpha-4
P54652	Heat shock-related 70 kDa protein 2
P08648	Integrin alpha-5
P43003	Excitatory amino acid transporter 1

Proteome identification in the drug product (DP)

Table 33 describes the 132 cell surface proteins identified in 3 triplicates of the DP comprising a mix of HT-29 DS-A, HT-29 DS-B, HCT-116 DS-A, HCT-116 DS-B, LoVo DS-A and LoVo DS-B.

5 **Table 33**

Uniprot accession number	Protein	Gene symbol
P10809	60 kDa heat shock protein, mitochondrial	<i>HSPD1</i>
P16422	Epithelial cell adhesion molecule	<i>EPCAM</i>
P08238	Heat shock protein HSP 90-beta	<i>HSP90AB1</i>
P11279	Lysosome-associated membrane glycoprotein 1	<i>LAMP1</i>
P13688	Carcinoembryonic antigen-related cell adhesion molecule 1	<i>CEACAM1</i>
P05362	Intercellular adhesion molecule 1	<i>ICAM1</i>
P25445	Tumor necrosis factor receptor superfamily member 6	<i>FAS</i>
P06576	ATP synthase subunit beta, mitochondrial	<i>ATP5B</i>
P11021	Endoplasmic reticulum chaperone BiP	<i>HSPA5</i>
P13667	Protein disulfide-isomerase A4	<i>PDIA4</i>
P30101	Protein disulfide-isomerase A3	<i>PDIA3</i>
P27797	Calreticulin	<i>CALR</i>
P02786	Transferrin receptor protein 1	<i>TFRC</i>
P05187	Alkaline phosphatase, placental type	<i>ALPP</i>
P17301	Integrin alpha-2	<i>ITGA2</i>

P07355	Annexin A2	<i>ANXA2</i>
O15031	Plexin-B2	<i>PLXNB2</i>
P11717	Cation-independent mannose-6-phosphate receptor	<i>IGF2R</i>
P08195	4F2 cell-surface antigen heavy chain	<i>SLC3A2</i>
P06756	Integrin alpha-V	<i>ITGAV</i>
Q9HDC9	Adipocyte plasma membrane-associated protein	<i>APMAP</i>
Q9BS26	Endoplasmic reticulum resident protein 44	<i>ERP44</i>
P06733	Alpha-enolase	<i>ENO1</i>
P05556	Integrin beta-1	<i>ITGB1</i>
Q12907	Vesicular integral-membrane protein VIP36	<i>LMAN2</i>
P09525	Annexin A4	<i>ANXA4</i>
P30040	Endoplasmic reticulum resident protein 29	<i>ERP29</i>
P21587	5'-nucleotidase	<i>NT5E</i>
P26006	Integrin alpha-3	<i>ITGA3</i>
P00505	Aspartate aminotransferase, mitochondrial	<i>GOT2</i>
Q96JJ7	Protein disulfide-isomerase TMX3	<i>TMX3</i>
Q07021	Complement component 1 Q subcomponent-binding protein, mitochondrial	<i>C1QBP</i>
P04439	HLA class I histocompatibility antigen, A alpha chain	<i>HLA-A</i>
P50895	Basal cell adhesion molecule	<i>BCAM</i>
P18084	Integrin beta-5	<i>ITGB5</i>
Q8TCT9	Minor histocompatibility antigen H13	<i>HM13</i>
P35232	Prohibitin OS=Homo sapiens (Human)	<i>PHB</i>
Q99623	Prohibitin-2 OS=Homo sapiens (Human)	<i>PHB2</i>
P23229	Integrin alpha-6	<i>ITGA6</i>
Q92692	Nectin-2	<i>PV RL2;</i> <i>NECTIN2</i>
P27487	Dipeptidyl peptidase 4	<i>DPP4</i>
P04083	Annexin A1	<i>ANXA1</i>
P01893	Putative HLA class I histocompatibility antigen, alpha chain H	<i>HLA-H</i>
P16070	CD44 antigen	<i>CD44</i>

P10321	HLA class I histocompatibility antigen, C alpha chain	<i>HLA-C</i>
Q96PD2	Discoidin, CUB and LCCL domain-containing protein 2	<i>DCBLD2</i>
Q6UVK1	Chondroitin sulfate proteoglycan 4	<i>CSPG4</i>
P01889	HLA class I histocompatibility antigen, B alpha chain	<i>HLA-B</i>
P08582	Melanotransferrin	<i>MF12</i> ; <i>MELTF</i>
Q14114	Low-density lipoprotein receptor-related protein 8	<i>LRP8</i>
P16144	Integrin beta-4	<i>ITGB4</i>
P01130	Low-density lipoprotein receptor	<i>LDLR</i>
Q12913	Receptor-type tyrosine-protein phosphatase eta	<i>PTPRJ</i>
P30533	Alpha-2-macroglobulin receptor-associated protein	<i>LRPAP1</i>
Q99523	Sortilin	<i>SORT1</i>
Q15836	Vesicle-associated membrane protein 3	<i>VAMP3</i>
P00533	Epidermal growth factor receptor	<i>EGFR</i>
P05067	Amyloid-beta precursor protein	<i>APP</i>
P15151	Poliovirus receptor	<i>PVR</i>
Q13641	Trophoblast glycoprotein	<i>TPBG</i>
P11233	Ras-related protein Ral-A	<i>RALA</i>
P56199	Integrin alpha-1	<i>ITGA1</i>
P07225	Vitamin K-dependent protein S	<i>PROS1</i>
Q9Y639	Neuroplastin	<i>NPTN</i>
Q14126	Desmoglein-2	<i>DSG2</i>
P61769	Beta-2-microglobulin	<i>B2M</i>
P48960	CD97 antigen	<i>CD97</i> ; <i>ADGRE5</i>
P10909	Clusterin	<i>CLU</i>
O43490	Prominin-1	<i>PROM1</i>
O14763	Tumor necrosis factor receptor superfamily member 10B	<i>TNFRSF10B</i>
Q6YHK3	CD109 antigen	<i>CD109</i>
Q12846	Syntaxin-4	<i>STX4</i>
P09382	Galectin-1	<i>LGALS1</i>

P78536	Disintegrin and metalloproteinase domain-containing protein 17	<i>ADAM17</i>
Q9UBR2	Cathepsin Z	<i>CTSZ</i>
P19013	Keratin, type II cytoskeletal 4	<i>KRT4</i>
P14735	Insulin-degrading enzyme	<i>IDE</i>
P19075	Tetraspanin-8	<i>TSPAN8</i>
Q8WTV0	Scavenger receptor class B member 1	<i>SCARB1</i>
P17813	Endoglin	<i>ENG</i>
O14672	Disintegrin and metalloproteinase domain-containing protein 10	<i>ADAM10</i>
Q10589	Bone marrow stromal antigen 2	<i>BST2</i>
P43007	Neutral amino acid transporter A	<i>SLC1A4</i>
Q9P2B2	Prostaglandin F2 receptor negative regulator	<i>PTGFRN</i>
P08962	CD63 antigen	<i>CD63</i>
P15291	Beta-1,4-galactosyltransferase 1	<i>B4GALT1</i>
P13987	CD59 glycoprotein	<i>CD59</i>
O00220	Tumor necrosis factor receptor superfamily member 10A	<i>TNFRSF10A</i>
O94985	Calsyntenin-1	<i>CLSTN1</i>
O95297	Myelin protein zero-like protein 1	<i>MPZL1</i>
P08648	Integrin alpha-5	<i>ITGA5</i>
P51809	Vesicle-associated membrane protein 7	<i>VAMP7</i>
Q14517	Protocadherin Fat 1	<i>FAT1</i>
P32004	Neural cell adhesion molecule L1	<i>L1CAM</i>
Q9Y696	Chloride intracellular channel protein 4	<i>CLIC4</i>
Q29983	MHC class I polypeptide-related sequence A	<i>MICA</i>
Q9UII2	ATPase inhibitor, mitochondrial	<i>ATPIF1</i>
Q9H0X4	Protein FAM234A	<i>ITFG3;</i> <i>FAM234A</i>
Q9BZM5	UL16-binding protein 2	<i>ULBP2</i>
O15258	Protein RER1	<i>RER1</i>
P08174	Complement decay-accelerating factor	<i>CD55</i>
P29317	Ephrin type-A receptor 2	<i>EPHA2</i>

P19021	Peptidyl-glycine alpha-amidating monooxygenase	<i>PAM</i>
Q03405	Urokinase plasminogen activator surface receptor	<i>PLAUR</i>
P19256	Lymphocyte function-associated antigen 3	<i>CD58</i>
P02749	Beta-2-glycoprotein 1	<i>APOH</i>
P15529	Membrane cofactor protein	<i>CD46</i>
P02787	Serotransferrin	<i>TF</i>
Q13444	Disintegrin and metalloproteinase domain-containing protein 15	<i>ADAM15</i>
P49810	Presenilin-2	<i>PSEN2</i>
P04156	Major prion protein	<i>PRNP</i>
P13726	Tissue factor	<i>F3</i>
P21926	CD9 antigen	<i>CD9</i>
Q13433	Zinc transporter ZIP6	<i>SLC39A6</i>
Q4KMQ2	Anoctamin-6	<i>ANO6</i>
P08581	Hepatocyte growth factor receptor	<i>MET</i>
Q9UNN8	Endothelial protein C receptor	<i>PROCR</i>
P02788	Lactotransferrin	<i>LTF</i>
P19634	Sodium/hydrogen exchanger 1	<i>SLC9A1</i>
P09958	Furin OS=Homo sapiens (Human)	<i>FURIN</i>
P30511	HLA class I histocompatibility antigen, alpha chain F	<i>HLA-F</i>
P31431	Syndecan-4	<i>SDC4</i>
Q14210	Lymphocyte antigen 6D	<i>LY6D</i>
P14174	Macrophage migration inhibitory factor	<i>MIF</i>
O75054	Immunoglobulin superfamily member 3	<i>IGSF3</i>
Q9NYQ7	Cadherin EGF LAG seven-pass G-type receptor 3	<i>CELSR3</i>
P21810	Biglycan	<i>BGN</i>
P43003	Excitatory amino acid transporter 1	<i>SLC1A3</i>
P13612	Integrin alpha-4	<i>ITGA4</i>
Q13492	Phosphatidylinositol-binding clathrin assembly protein	<i>PICALM</i>
P15328	Folate receptor alpha	<i>FOLR1</i>
Q00839	Heterogeneous nuclear ribonucleoprotein U	<i>HNRNPU</i>

Example 6***In vitro* evaluation of the functional activity of STC-1010 through mixed lymphocyte reaction assay**

The aim was to assess *in vitro* the final product (DP or “STC-1010”), for its potential to induce an immunogenic profile of human monocytes-derived dendritic cells (DCs) when applied alone or in combination with CD40L (naturally present *in vivo*) and to evaluate, in a mixed lymphocyte reaction (MLR) assay, the T cell activation mediated by dendritic cells. The study was performed on a co-culture of HLA-matched monocyte-derived mDCs and CD8⁺ T cells, in which system i) the cytokine profile of DCs was measured through the quantitation of cytokines released in the culture supernatants (including IL-12 and IL-8), and ii) the DC-mediated T cell activation was measured through the assessment of the MLR response by the mean of the quantification of released IFN γ using specific homogeneous time-resolved fluorescence (HTRF)-based detection kits.

Materials and methods

Human PBMCs from healthy donors were used in this study in order to isolate freshly i) monocytes which were used to obtain mature DCs through differentiation/maturation protocols, as well as ii) CD8⁺ lymphocytes.

Briefly, human monocytes were freshly isolated from PBMCs and were differentiated into DCs under cultivation in the presence of GM-CSF and IL-4. At the end of the differentiation process, DCs were matured in a specific cocktail in the presence of LPS and IFN γ containing STC-1010 at three different ratios (1:1, 3:1 and 10:1, plus a condition without STC-1010 as negative control) in the presence and absence of 0.6 μ g/mL CD40L.

Upon maturation, mDCs were validated as CD209⁺ CD1a⁺ CD80⁺ CD83⁺ CD86⁺ and cell culture supernatants were retrieved for IL-12 and IL-8 quantitation by means of HTRF.

Then, DCs were co-cultured, at an appropriate stimulator:responder ratio of 1:4, with CD8⁺ T cells isolated from the same donor. Control conditions (untreated), as well as T cells alone and mDCs alone, were included in the experiment. 72 hours following co-culture, supernatants were collected and effects of STC-1010 were evaluated on CD8⁺

T cell activation by mean of the quantification of released IFN γ levels that was used as a key representative surrogate of T cell activation. Cytokine level quantification was performed by HTRF.

Results

5 Effects of STC-1010 on mDC cytokines

STC-1010 was able to limit in a ratio-dependent manner the secretion of IL-8 by mDCs, both in absence and in presence of CD40L (**Fig. 11A** and **11B**, respectively). STC-1010 was also able to enhance in a ratio-dependent manner the secretion of IL-12 by mDCs in presence of CD40L (**Fig. 11C**). This ratio-dependent response confirms the action of
10 STC-1010 on the maturation of DCs.

Two other cytokines (IL-10 and TNF α) were also evaluated in the course of this study: STC-1010 demonstrated a trend to limit IL-10 secretion in the presence of CD40L; while TNF α secretion was slightly decreased upon STC-1010 exposure with an effect culminating at the highest ratio, both in absence or presence of CD40L (data not shown).

15 Effects of STC-1010 on the MLR response through IFN γ quantification

mDCs co-cultured with autologous CD8⁺ T cells displayed an effective MLR response when previously exposed to CD40L during the maturation period. This effect was evidenced through the evaluation of IFN γ release in the supernatant collected from the co-culture, which showed a significant up-regulation compared to control and vehicle
20 (**Fig. 12**).

Conclusion

In the light of these data, STC-1010 has demonstrated its ability to modulate DC maturation as evidenced through the cytokine profile, with an immunogenic and ratio-dependent activity of STC-1010.

25 This immunogenic profile was confirmed through MLR where CD8⁺ T cells co-cultured with STC-1010-primed DCs had an improved functional activity.

Example 7

***In ovo* evaluation of the functional activity of STC-1010 through chorioallantoic membrane assay**

The aim was to assess *in ovo* the final product (DP or “STC-1010”), for its potential to
5 induce an immune response in a chorioallantoic membrane (CAM) assay.

Materials and methods

Preparation of chicken embryos

Fertilized White Leghorn eggs were incubated at 37.5°C with 50 % relative humidity for 9 days. At that moment (E9), the CAM was dropped down by drilling a small hole through
10 the eggshell into the air sac, and a 1-cm² window was cut in the eggshell above the CAM. At least 20 eggs were opened for each study group (but because eggshell opening is an invasive surgical act, some death can occur during the first hours after opening, hence data may have been collected with 15-20 eggs per group).

Treatment

15 Before treatment, the viability of each egg was checked and surviving eggs were randomized in groups. All eggs of a group were treated with a volume of 100 µL of STC-1010, with three test conditions:

- test condition “STC1010 [1]”: 10⁵ cells/mL, *i.e.*, 10⁴ cells/embryo;
- test condition “STC1010 [2]”: 5 × 10⁵ cells/mL, *i.e.*, 5 × 10⁴ cells/embryo; and
- 20 - test condition “STC1010 [3]”: 10⁶ cells/mL, *i.e.*, 10⁵ cells/embryo.

A negative control (“Neg Ctrl”) was performed in parallel, in absence of STC-1010.

Quantitative evaluation of chicken immune cells

On day E18, peripheral blood was collected and treated with heparin to prevent blood clotting.

25 For dendritic cells (DCs) evaluation, 100 µL of individual samples (n = 8 per group) was recovered, from which RNA was extracted, reverse-transcribed, pre-amplified and analyzed by qPCR with specific primers for chicken CD40, CD83 and CD86 sequences.

For all points done in qPCR, expression of chicken GAPDH was also analyzed, as reference gene expression, and used to normalize immune biomarker expression between samples. Calculation of C_q for each sample, mean C_q and relative amounts of immune cells for each group were directly managed by the Bio-Rad® CFX Maestro software.

- 5 For T lymphocytes evaluation, the remaining blood samples were pooled within group. Then, blood samples were processed with Hypaque-Ficoll (HF) separation for peripheral blood mononuclear cells (PBMCs) isolation. After that, purified PBMCs were labelled with anti-chicken CD45 (Thermofisher, Ref.: MA5-28679), anti-chicken CD3 (Southern Biotech, Ref.: 8200-26), anti-chicken CD4 (Thermo Fisher, Ref.: MA5-28686) and
- 10 anti-chicken CD8 (Thermo Fisher, Ref.: MA5-28686) for T lymphocytes evaluation, through flow cytometry analysis.

Quantitative evaluation of immune cytokines

- On day E18, peripheral blood was individually collected (n = 5 per group) and treated with heparin to prevent blood clotting. Then, blood samples were centrifuged for plasma
- 15 collection, which was followed by ELISA analysis for IL-12 and IFN γ expression, each plasma sample being evaluated at three dilutions. All ELISA kits were ordered from Cusabio (chicken IL-12 Elisa kit, Ref.: CSB-E12836C; chicken IFN ELISA kit, Ref.: CSB-E08550Ch).

Statistical analysis and significance

- 20 For all quantitative data, the outlier identification and the one-way ANOVA (with post-tests between each couple of groups) were done using Prism® (GraphPad Software).

Results

Quantitative evaluation of chicken immune cells by FACS

- Leukocyte activation was evaluated in groups “Neg Ctrl” and “STC-1010 [3]” by FACS
- 25 quantification of CD45⁺, CD3⁺, CD4⁺ and CD8⁺ cells in PBMCs purified at E18. **Table 34** shows the FACS analysis data of these different cell subsets included in peripheric leukocytes (as a % in peripheric CD45⁺ leucocytes).

Table 34

	CD3/CD4 staining			
	CD3 ⁻ /CD4 ⁻	CD3 ⁺ /CD4 ⁻	CD3 ⁻ /CD4 ⁺	CD3 ⁺ /CD4 ⁺
Neg Ctrl	97.5	0.6	1.7	0.3
STC-1010 [3]	86.4 (↓)	0.3 (↓)	12.5 (↑)	0.5 (↑)
	CD3/CD8 staining			
	CD3 ⁻ /CD8 ⁻	CD3 ⁺ /CD8 ⁻	CD3 ⁻ /CD8 ⁺	CD3 ⁺ /CD8 ⁺
Neg Ctrl	98.5	0.4	0.8	0.3
STC-1010 [3]	97.6 (↓)	0.1 (↓)	2.0 (↑)	0.3 (=)

These data show an increase of CD4⁺ and CD8⁺ leukocytes, in particular a high increase of CD3⁻/CD4⁺ leukocytes.

Quantitative evaluation of immune cytokines

- IL-12 and IFN γ expression level in peripheral blood was estimated in all groups at E18. To pertinently evaluate cytokines secretion, each plasma sample was evaluated at three dilution¹₂1/2, 1/10 and 1/50.

Table 35 shows the data analysis of peripheral IL12 secretion.

Table 35

	IL-12 analysis			<i>p</i> -value versus STC-1010 [..]		
	n	C° (pg/mL)	% increase	[1]	[2]	[3]
Dilution ¹ ₂ 1/2						
Neg Ctrl	5	1.54	<i>n.a.</i>	<i>n.a.</i>	<i>n.a.</i>	<i>n.a.</i>
STC-1010 [1]	5	2.3	49.94	0.6896	<i>n.a.</i>	<i>n.a.</i>
STC-1010 [2]	5	2.88	87.41	0.2510	0.8387	<i>n.a.</i>
STC-1010 [3]	5	3.99	159.84	0.0128	0.1093	0.4021
Dilution at 1/10						
Neg Ctrl	5	3.24	<i>n.a.</i>	<i>n.a.</i>	<i>n.a.</i>	<i>n.a.</i>

STC-1010 [1]	5	6.63	104.74	0.4535	<i>n.a.</i>	<i>n.a.</i>
STC-1010 [2]	5	8.23	154.23	0.1582	0.8899	<i>n.a.</i>
STC-1010 [3]	5	17.1	427.99	<0.0001	0.0013	0.0056
Dilution at 1/50						
Neg Ctrl	5	9.48	<i>n.a.</i>	<i>n.a.</i>	<i>n.a.</i>	<i>n.a.</i>
STC-1010 [1]	5	27	184.81	0.2099	<i>n.a.</i>	<i>n.a.</i>
STC-1010 [2]	5	19.18	102.36	0.6723	0.7964	<i>n.a.</i>
STC-1010 [3]	5	56.93	500.55	0.0002	0.0139	0.0022

As seen in these results, a net increase of IL-12 secretion induced by STC-1010 was observed when compared to the negative control, at all three sample dilutions. No significant increase of IFN γ was however observed following STC-1010 treatment (data not shown).

5 Quantitative evaluation of chicken immune cells by RT-qPCR

The expression of three DCs activation markers (CD40, CD83, CD86) was evaluated by RT-qPCR. Results show an upregulation of these markers induced by STC-1010, in particular at the highest dose for CD40 and CD86, and at all doses and more significantly at the lowest dose for CD83 (**Fig. 13A-C**).

10 **Conclusion**

Out of the embryos, the vast majority survived throughout this study, confirming the absence of toxicity of STC-1010:

- negative control: n = 50, 3 embryos died (6% lethality);
- group "STC1010 [1]": n = 48, 2 embryos died (4.17% lethality);
- 15 - group "STC1010 [2]": n = 33, 0 embryo died (0% lethality);
- group "STC1010 [3]": n = 29, 1 embryo died (3.45% lethality).

STC-1010 was shown to be able to boost the immune system and activate innate and/or adaptive immune responses using this CAM assay.

Based on these promising results, a new study will be performed to evaluate the *in ovo* efficacy of STC-1010 on human colorectal adenocarcinoma-grafted chicken embryos.

CLAIMS

1. A composition comprising (i) stressed HT-29, HCT-116 and LoVo cells, and
 5 (ii) immunogenic stress proteins produced by these cells in response to a stress applied *in vitro*.
2. The composition according to claim 1, wherein stressed HT-29, HCT-116 and LoVo cells have developed resistance mechanism in response to one or several stress[es] applied *in vitro*, selected from the group comprising radiations, thermal stress, chemical stress, metabolic stress and any combinations thereof, leading to
 10 the production of the stress proteins.
3. The composition according to claim 1 or 2, wherein stressed HT-29, HCT-116 and LoVo cells are non-proliferative.
4. The composition according to any one of claims 1 to 3, wherein immunogenic stress proteins are haptenated;
 15 preferably with a hapten selected from the group comprising 2,4-dinitrophenyl (DNP); 2,4-dinitrofluorobenzene; sulfanilic acid; *N*-iodoacetyl-*N'*-(5-sulfonic-naphthyl)ethylene diamine; anilin; *p*-amino benzoic acid; biotin; fluorescein and derivatives thereof (including FITC, TAMRA, and Texas Red); digoxigenin; 5-nitro-3-pyrazolecarbamide;
 20 4,5-dimethoxy-2-nitrocinnamide;
 2-(3,4-dimethoxyphenyl)-quinoline-4-carbamide;
 2,1,3-benzoxadiazole-5-carbamide; 3-hydroxy-2-quinoxalinecarbamide,
 4-(dimethylamino)azobenzene-4'-sulfonamide (DABSYL); rotenone isooxazolinI(*E*)-2-(2-(2-oxo-2,3-dihydro-1*H*-benzo[*b*][1,4]diazepin-4-yl)phenozy)
 25 acetamide; 7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylic acid;
 2-acetamido-4-methyl-5-thiazolesulfonamide; and
p-methoxyphenylpyrazopodophyllamide;
 more preferably with 2,4-dinitrophenyl (DNP).

5. The composition according to any one of claims 1 to 4, being a pharmaceutical composition or a vaccine composition, and further comprising at least one pharmaceutically acceptable excipient.
6. The composition according to any one of claims 1 to 5, comprising from about 10^5 to about 10^8 stressed HT-29, HCT-116 and LoVo cells.
7. The composition according to any one of claims 1 to 6, for use in treating cancer in a subject in need thereof.
8. An intermediate composition comprising (i) one of stressed HT-29 cells, stressed HCT-116 cells and stressed LoVo cells, and (ii) stress proteins,
wherein the one of stressed HT-29 cells, stressed HCT-116 cells or stressed LoVo cells have developed resistance mechanism in response to (i) a metabolic stress, (ii) radiations and (iii) a thermal stress applied *in vitro*, leading to the production of the stress proteins.
9. An intermediate composition comprising (i) one of stressed HT-29 cells, stressed HCT-116 cells or stressed LoVo cells, and (ii) stress proteins,
wherein the one of stressed HT-29 cells, stressed HCT-116 cells or stressed LoVo cells have developed resistance mechanism in response to (i) a metabolic stress, and (ii) a chemical stress applied *in vitro*, leading to the production of the stress proteins.
10. A method of manufacturing the intermediate compositions according to claim 8 or 9, comprising the following steps:
 - a) cultivating HT-29, HCT-116 or LoVo cells in a suitable culture medium;
 - b) subjecting the HT-29, HCT-116 or LoVo cells cultured in step a) to one or several stress[es] *in vitro*, wherein these HT-29, HCT-116 or LoVo cells develop resistance mechanisms in response to the one or several stress[es] and thereby produce stress proteins,
 - c) recovering the stressed HT-29, HCT-116 or LoVo cells together with the stress proteins they have produced in step b), and

- d) treating the stressed HT-29, HCT-116 or LoVo cells and the stress proteins they have produced, all together recovered in step c), with a molecule or by a process capable of rendering the stress proteins immunogenic.
11. The method according to claim 10, wherein step c) is carried out at least several hours after completion of step b), preferably at least 12 hours or more after completion of step b).
12. The method according to claim 10 or 11, wherein step d) comprises linking the stress proteins to or complexing the stress proteins with a means capable to confer immunogenicity.
13. The method according to claim 12, wherein the means capable to confer immunogenicity is an hapten;
 preferably the hapten is selected from the group comprising 2,4-dinitrophenyl (DNP); 2,4-dinitrofluorobenzene; sulfanilic acid; *N*-iodoacetyl-*N'*-(5-sulfonic-naphthyl)ethylene diamine; anilin; *p*-amino benzoic acid; biotin; fluorescein and derivatives thereof (including FITC, TAMRA, and Texas Red); digoxigenin; 5-nitro-3-pyrazolecarbamide; 4,5-dimethoxy-2-nitrocinnamide; 2-(3,4-dimethoxyphenyl)-quinoline-4-carbamide; 2,1,3-benzoxadiazole-5-carbamide; 3-hydroxy-2-quinoxalinecarbamide, 4-(dimethylamino)azobenzene-4'-sulfonamide (DABSYL); rotenone isooxazoline; (*E*)-2-(2-(2-oxo-2,3-dihydro-1*H*-benzo[*b*][1,4]diazepin-4-yl)phenoxy)acetamide; 7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylic acid; 2-acetamido-4-methyl-5-thiazolesulfonamide; and *p*-methoxyphenylpyrazopodophyllamide;
 more preferably the hapten is 2,4-dinitrophenyl (DNP).
14. The method according to any one of claims 10 to 13, for manufacturing the intermediate compositions according to claim 8, wherein step b) comprises subjecting the HT-29, HCT-116 or LoVo cells cultured in step a) to the following stresses *in vitro*, applied concomitantly or successively:

- (i) an *in vitro* culture in a depleted medium, under hypoxia, and/or at low pH;
preferably an *in vitro* culture in low serum culture conditions in a 2 % FBS
culture medium,
- 5 (ii) an *in vitro* radiation with a total dose ranging from about 0.25 to about 25 Gy,
preferably from about 1 to about 15 Gy, preferably for a period ranging from
about 1 to about 20 minutes;
preferably an *in vitro* radiation with a total dose of 10 Gy for a period of about
1 to about 5 minutes, and
- 10 (iii) an *in vitro* thermic choc at a temperature ranging from about 38°C to about
45°C, applied to the cells for a period ranging from about 15 minutes to about
4 hours;
preferably an *in vitro* thermic choc at a temperature of about 42°C for a period
of about 60 minutes.
- 15 **15.** The method according to any one of claims **10** to **13**, for manufacturing the
intermediate compositions according to claim **9**, wherein step b) comprises
subjecting the HT-29, HCT-116 or LoVo cells cultured in step a) to the following
stresses *in vitro*, applied concomitantly or successively:
- 20 (i) an *in vitro* culture in a depleted medium, under hypoxia, and/or at low pH;
preferably an *in vitro* culture in low serum culture conditions in a 2 % FBS
culture medium,
- (ii) an *in vitro* exposition to at least one or several chemotherapeutic agents
and/or alcohols, for a period ranging from about 6 hours to about 120 hours.
- 16.** The method according to claim **15**, wherein:
- 25 - the cells are HT-29 cells and the *in vitro* exposition at (ii) is to about 13 µM
oxaliplatin for a period of about 72 hours; or
- the cells are HCT-116 cells and the *in vitro* exposition at (ii) is to about
315 nM SN-38 (7-ethyl-10-hydroxy-camptothecin) for a period of about
48 hours; or
- 30 - the cells are LoVo cells and the *in vitro* exposition at (ii) is to about 5 µM
fluorouracil (5-FU) for a period of about 48 hours.

17. A method of manufacturing the composition according to any one of claims 1 to 6, comprising the following steps:
- a. obtaining six intermediate compositions according to claims 8 and 9, wherein the six intermediate compositions are:
- 5 1) an intermediate composition according to claim 8 comprising stressed HT-29 cells and stress proteins,
 - 2) an intermediate composition according to claim 8 comprising stressed HCT-116 cells and stress proteins,
 - 3) an intermediate composition according to claim 8 comprising stressed
10 LoVo cells and stress proteins,
 - 4) an intermediate composition according to claim 9 comprising stressed HT-29 cells and stress proteins,
 - 5) an intermediate composition according to claim 9 comprising stressed HCT-116 cells and stress proteins,
 - 15 6) an intermediate composition according to claim 9 comprising stressed LoVo cells and stress proteins,
- b. mixing these six intermediate compositions together;
preferably wherein these six intermediate compositions are mixed together in an equal ratio of stressed HT-29, HCT-116 and LoVo cells.