

ALKANESULFONANILIDE DERIVATIVES AND
PHARMACEUTICALLY ACCEPTABLE ACID ADDITION
SALTS THEREOF

The invention herein described is concerned with increasing the growth rate, enhancing the lean meat deposition and/or improving the lean meat to fat ratio in farm and domestic animals (ie swine, poultry, cattle, sheep, goats, rabbits, cats, dogs, etc). This is achieved by either oral or parenteral administration of various 3-amino-4-hydroxy(or alkoxy)phenethanolamine derivatives, including novel derivatives presently revealed.

By way of background, sulfonanilide derivatives, their acid addition and metal salts are disclosed in United States Patent Nos. 3,341,584 and 3,801,631. These patents reveal that the above-mentioned compounds are useful as adrenergic neurotransmitters, vasopressors, vasodepressors, analgesics, bronchodilators, α -receptor stimulants, β -receptor stimulants, α -receptor blocking agents, β -receptor blocking agents, papaverine-like smooth muscle depressants and anti-inflammatory agents.


In addition to the above-said sulfonanilide derivatives, related 3-amino-4-hydroxyphenethanolamine derivatives are described with similar pharmacological activities in the following publications: Belgium patents numbered 784,105; 762,504 and 765,988; United States patents numbered 3,763,232; 3,917,847 and 3,711,545; Journal of Medicinal Chemistry, 10, 762-772 (1967), ibid., 17, 49-57 (1974) and ibid., 19, 626-633 (1976).

In recent years, the cost of raising meat animals has increased markedly in response to economic fluctuations in the areas of energy resources, ancillary products, and consumer markets. The necessity of providing adequate meat protein supplies to an expanding population is self-evident. A method for increasing the quantity and/or


16
2-

quality of animal protein supplies while maintaining ordinary feed requirements would facilitate delivery of required food supplies.

5 In light of the foregoing summary of some demands and limitations of conventional methods for the production of meat products, an improved method for quantitative and/or qualitative improvement in animal yields is highly desirable.



It has been unexpectedly discovered that oral or parenteral administration of selected 3-amino-4-hydroxy(or alkoxy)phenethanolamine derivatives not only will increase the growth rates of certain warm-blooded animals but also will increase lean meat deposition and improve the lean meat to fat ratio in the bodies of these animals. In this disclosure, the term "lean meat" is used interchangeably with the amount of muscle or protein present in referenced animals. The above-mentioned effects are achieved by oral or parenteral administration to appropriate animals of an effective amount of a compound represented by the following structural formula:



(I)

wherein, R₁ is hydrogen or C₁-C₄ alkyl; R₂ is hydrogen, C₁-C₄ alkyl, cycloalkyl C₃-C₅, benzyl, phenethyl, α-methylphenethyl or α,α-dimethylphenethyl; R₃ is OH, OR₆ or SR₇; R₄ is hydrogen, C₁-C₄ alkyl, benzyl or C₂-C₅ alkanoyl; R₅ is hydrogen, methyl or ethyl; R₆ is C₁-C₆ alkyl, benzyl, phenyl or allyl; R₇ is hydrogen, C₁-C₆ alkyl, benzyl, phenyl or allyl; X is NHSO₂R₈, N(SO₂R₈)₂,


$N(R_1)_2$, $NHCO R_1$, $NH-CO-N(R_1)_2$, $NHSO_2 N(R_1)_2$, $NHCOOR_9$, or $NHCOO$ -benzyl; R_8 is C_1-C_4 alkyl; R_9 is C_1-C_5 alkyl; with the provisos that when R_2 is cycloalkyl C_3-C_5 , benzyl, phenethyl, α -methylphenethyl or α,α -dimethylphenethyl, R_1 of $NR_1 R_2$ is hydrogen; and when R_3 is OR_6 or SR_7 , X is only $NHSO_2 R_8$; and further that the R_1 group in $(R_1)_2$ may or may not be identical substituents; and the optically active isomers and pharmacologically acceptable acid addition and metal salts thereof.

10 The present invention also provides an animal feed composition which, for increasing the growth and lean meat-to-fat ratio of domestic and farm animals, contains a pharmaceutically acceptable diluent and a compound of formula I above.

15 A preferred group of compounds for use in this invention, as depicted by formula I above, are those wherein R_4 is hydrogen; R_3 is OH ; R_5 is hydrogen; R_1 is hydrogen or C_1-C_4 alkyl; R_2 is hydrogen, C_1-C_4 alkyl, α,α -dimethylphenethyl or α -methylphenethyl; X is
20 $NHSO_2 CH_3$, $N(R_1)_2$, $NHCO R_1$, $NH-CO-N(R_1)_2$, $NHSO_2 N(R_1)_2$ or $NHCOOR_9$, where R_9 is C_1-C_5 alkyl; with the provisos that when R_2 is α -methylphenethyl or α,α -dimethylphenethyl, R_1 in $NR_1 R_2$ is hydrogen; and both R_1 's in $(R_1)_2$ may be the same or different substituents; and the optically active
25 isomers and pharmacologically acceptable acid addition and metal salts thereof.

A still more preferred group of compounds of the present invention have the structure depicted by Formula II:

30




(II)

-5-

wherein R_1 is hydrogen or C_1 - C_4 alkyl; R_2 is C_1 - C_4 alkyl or α , α -dimethylphenethyl; with the proviso that when R_2 is α , α -dimethylphenethyl, R_1 of NR_1R_2 is hydrogen; and the optically active isomers and pharmacologically acceptable acid addition and metal salts thereof.

Among the pharmacologically acceptable acid addition salts useful in the present invention are the hydrochloride, sulfate, phosphate, gluconate, succinate, propionate, oleate, linoleate, linolenate, fumarate and abiate salts.

Novel compounds of the present invention are depicted by the structure shown as Formula III below:




(III)


wherein X is $NHSO_2R_8$ or $N(SO_2R_8)_2$; R_1 is hydrogen or C_1 - C_4 alkyl; R_2 is hydrogen, C_1 - C_4 alkyl, benzyl, phenethyl, α -methylphenethyl or α , α -dimethylphenethyl; R_3 is OR_6 or SR_7 ; R_6 is C_1 - C_4 alkyl, benzyl, phenyl or allyl; R_7 is hydrogen, C_1 - C_6 alkyl, benzyl, phenyl or allyl; R_8 is C_1 - C_4 alkyl; with the provisos that when R_2 is benzyl, phenethyl, α -methylphenethyl or α , α -dimethylphenethyl, R_1 of NR_1R_2 is hydrogen; and when X is $N(SO_2R_8)_2$, R_3 is OH; and the optically active isomers and pharmacologically acceptable acid addition and metal salts thereof.

A preferred group of these novel formula III compounds are those wherein R_3 is OR_6 and R_1 , R_2 , R_6 , R_8 , and X are as described above for formula III compounds.

Advantageously, the compounds of this invention,
5 as delineated by formulas I, II, and III above, generally can be prepared by one or more of the synthetic routes described in the U.S. Patent Nos. 3,341,584 and 3,801,631 referred to above, and graphically illustrated below.




-7-




In the above reactions, R_1 in the function $(R_1)_2$ may represent the same or different substituents selected from hydrogen and C_1-C_4 alkyl and R_4 is as defined for the formula I compounds above.

5 The above-illustrated acetophenone compounds may be converted by the reactions, graphically illustrated below, to compounds of the invention having the formula I structure in which R_3 is OH, R_1 is hydrogen and X, R_2 , R_4 , and R_5 are as described for formula I compounds depicted above.



OR



In the above reaction, X, R₁, R₂, R₄, and R₅ are as defined with respect to formula I


5 Additionally, the novel 2'-hydroxy-5'-[1-alkoxy-2-(alkyl-, dialkyl- or aralkylamino)ethyl]-alkanesulfonanilides of this invention can be prepared by the following steps:

(1) reacting a 2'-hydroxy-5'-[1-hydroxy-2-(alkylamino)ethyl]alkanesulfonanilide or the dialkylamino or aralkylamino derivative thereof, with
10 an equivalent amount or slight excess of thionyl chloride to obtain the 2'-hydroxy-5'-[1-chloro-2-(alkyl-, dialkyl- or aralkylamino)ethyl]alkanesulfonanilide hydrochloride;


(2) converting the above-formed alkane-sulfonanilide hydrochloride salt to the 2'-hydroxy-5'-[1-alkoxy-2-(alkyl-, dialkyl- or aralkylamino)ethyl]-alkanesulfonanilide hydrochloride by reacting the hydrochloride salt with the appropriate C₁-C₆ alcohol at temperature between about 0° to 150°C; and

20 (3) when a free base of an above-named hydrochloride-salt compound is desired, it is obtained by neutralization of the hydrochloride salt with a suitable aqueous base (i.e., sodium hydroxide, potassium hydroxide etc.).


25 The 2'-hydroxy-5'-[1-alkoxy-2-(alkyl-, dialkyl- or aralkylamino)ethyl]alkanesulfonanilide derivatives and salts as well as 2'-hydroxy-5'-[1-benzyloxy-2-(alkyl-, dialkyl- or aralkylamino)-ethyl]alkanesulfonanilide derivatives and salts are
30 readily prepared by the above-described procedures and further illustrated in the following equations:



5



10



15


wherein R_1 is hydrogen or C_1 - C_4 alkyl; R_2 is hydrogen, C_1 - C_4 alkyl, benzyl, phenethyl, α -methylphenethyl or α , α -dimethylphenethyl; R_8 is C_1 - C_4 alkyl and R_6 is C_1 - C_6 alkyl, benzyl, phenyl or allyl; with the condition that when R_2 is benzyl, phenethyl, α -methylphenethyl or α , α -dimethylphenethyl, R_1 of NR_1R_2 is hydrogen.

The 2'-hydroxy-5'-[1-phenoxy-2-(alkyl-, dialkyl- or aralkylamino)ethyl]alkanesulfonanilide compounds can be prepared by reacting an appropriate 2'-hydroxy-5'-[1-chloro-2-(alkyl-, dialkyl- or aralkylamino)ethyl]alkanesulfonanilide compounds with an excess amount (i.e., preferably 10 to 15 equivalents) of an alkali metal phenoxide in the presence of an aprotic solvent (i.e., tetrahydrofuran, ether, toluene, benzene, etc.). This reaction is carried out at a temperature between about -5 and +10°C under a blanket of an inert gas.

35

The 2'-hydroxy-5'-[1-alkyl-, benzyl-, or allylthio)-2-(alkyl-, dialkyl-, or aralkylamino)ethyl]-alkanesulfonanilide compounds of the invention can be prepared by reacting an appropriate 2'-hydroxy-5'-[1-chloro-2-(alkyl-, dialkyl-, or aralkylamino)ethyl]-alkanesulfonanilide hydrochloride with an equimolar or excess amount of an alkyl, benzyl or allylmercaptan (i.e., R_6SH).

This reaction is illustrated by the following equation:



wherein R_1 is hydrogen or C_1 - C_4 alkyl, R_2 is hydrogen, C_1 - C_4 alkyl, benzyl, phenethyl, α -methylphenethyl or α,α -dimethylphenethyl; R_8 is C_1 - C_4 alkyl; R_7 is hydrogen, C_1 - C_6 alkyl, benzyl, or allyl. This reaction is generally conducted in the presence of an aprotic solvent (i.e., a chlorinated hydrocarbon such as ether or other similar material) at a temperature ranging from -5 to +50°C under a blanket of an inert gas.

The 2'-hydroxy-5'-[1-phenylthio-2-(alkyl-, dialkyl-, or aralkylamino)ethyl]alkanesulfonanilides can be prepared by reacting an appropriate 2'-hydroxy-5'-[1-chloro-2-(alkyl-, dialkyl-, or aralkylamino)ethyl]-alkanesulfonanilide compound with an excess amount of an alkali metal thiophenoxide (i.e., sodium or potassium thiophenoxide) in an aprotic solvent (i.e., tetrahydrofuran, benzene, toluene or ether) at a temperature ranging from about -5 to +10°C under a blanket of an inert gas.

The formula I compounds of this invention can generally be administered either orally or parenterally to domestic or farm animals with resultant increases in growth rates and enhancement of the lean meat to fat ratio in these animals. In actual practice, the formula I alkanesulfonanilides may be directly mixed with animal feeds or prepared in the form of an animal-feed premix, concentrate, or supplement which can be blended with or applied as a top dressing to animal feeds. Regardless of the procedure selected, the active compound should be present at levels from about 0.05 to 500 ppm and preferably 0.1 to 100 ppm in the final feed.

Animal-feed premixes, supplements or concentrates can be prepared by mixing on a weight basis about 0.5 to 50% of a suitable formula I alkanesulfonanilide with about 50 to 99.5% of an edible diluent. Diluents suitable for use in the manufacture of animal-feed supplements, concentrates, and premixes include the following: corn meal, soybean meal, bone meal, alfalfa meal, cottonseed oil meal, urea, molasses, and other similar materials. Use of the diluents in feed supplements, concentrates, and premixes improves uniformity of distribution of the active ingredient in the finished feed.

Feeds for swine, cattle, sheep, and goats preferably contain about 0.05 to 400 grams of active ingredient per ton of feed with an optimum level of about 0.25 to 100 grams per ton. Preferred poultry and domestic-pet feeds range from about 0.05 to 400 grams and most preferably 0.2 to 100 grams of active ingredient per ton of feed.

13-

When parenteral administration is desired, formula I alkanesulfonanilides may be formulated as pastes or pellets and administered to the animals by subcutaneous injection. This procedure involves

5 injection of a formula I alkanesulfonanilide derivative in an amount sufficient to provide the animal with 0.001 to 100 mg/kg of body weight/day of the active compound. The preferred dosage for swine, cattle, sheep, and goats ranges from about 0.0001 to 50 mg/day/kg of body weight

10 of the active alkanesulfonanilide derivative. The preferred dosage for poultry and domestic pets ranges from about 0.001 to 10 mg/day/kg of body weight.

Paste formulations suitable for subcutaneous injection can be prepared by dispersing a formula I

15 alkanesulfonanilide derivative in pharmacologically-acceptable oil (ie, peanut oil, corn oil, sesame oil, etc). Pellets for subcutaneous injection can be prepared by mixing a formula I alkanesulfonanilide derivative with a suitable diluent (eg "Carbowax", carnauba wax, etc -

20 "Carbowax" is a registered Trade Mark). A lubricant (ie magnesium or calcium stearate) can be added to improve the pelleting process.

In order to obtain the dosage levels necessary to achieve desired results (ie increase in growth rates and

25 improvement in lean meat-to-fat rates), it may be necessary to administer multiple pellets. Also, implants may be made periodically during treatment periods in order to maintain proper animal drug levels.

In addition to effects on growth rates and lean meat to fat ratios, administration of formula I alkanesulfonanilides to meat-producing animals frequently results in enhanced efficiency of feed utilization. With the use of materials and methods revealed in the present invention, producers can market superior quality meat animals in a short period of time while incurring minimum feed costs.

The following non-limiting examples further serve to illustrate the invention.

EXAMPLE 1

Preparation of 2'-hydroxy-5'-[1-chloro-2-(isopropylamino)ethyl]methanesulfonanilide hydrochloride

A 1-g sample of 2'-hydroxy-5'-[1-hydroxy-2-(isopropylamino)ethyl]methanesulfonanilide is added to 2 mL of ice-cold thionyl chloride. The mixture is then stirred at ambient temperature until the reaction is completed. Excess thionyl chloride is removed in vacuo, and the residue is washed with ethyl ether thus yielding the product compound.

The compound 2'-hydroxy-5'-[1-chloro-2-(isopropylamino)ethyl]methanesulfonanilide hydrochloride can also be prepared by heating 1 g of 2'-hydroxy-5'-[1-hydroxy-2-(isopropylamino)ethyl]methanesulfonanilide in 10 mL of dry acetonitrile or dry dimethoxyethane containing 2 mL of thionyl chloride at reflux temperature for 10 minutes following the procedure described in the Journal of Medicinal Chemistry, 19, 632 (1976).

EXAMPLE 2Preparation of 2'-hydroxy-5'-[1-ethoxy-2-(isopropylamino)ethyl]methanesulfonanilide hydrochloride


5 A 1-g sample of 2'-hydroxy-5'-[1-chloro-2-(isopropylamino)ethyl]methanesulfonanilide hydrochloride is added to 10 mL of absolute ethanol and heated at reflux for 30 minutes. The mixture is evaporated to dryness, and the residue is washed with ethyl ether thus yielding the product compound. The same procedure
10 is used to prepare 2'-hydroxy-5'-[1-isopropoxy-2-(isopropylamino)ethyl]methanesulfonanilide hydrochloride, m.p. 121-123, by substituting isopropanol for ethanol in the above reaction.

Neutralization of the 2'-hydroxy-5'-[1-ethoxy-2-(isopropylamino)ethyl]methanesulfonanilide hydrochloride
15 salt with aqueous sodium hydroxide yields the free base of this compound which can be separated from the aqueous solution and readily converted to other acid salts by titration.

20

EXAMPLE 3Preparation of chloro intermediates

Several chloro intermediates can be prepared by the method of Example 1 as illustrated in the
25 following equation:



35

wherein substituents are:

	<u>R₁</u>	<u>R₂</u>	<u>R₈</u>
5	H	<u>t</u> -butyl	methyl
	H	<u>n</u> -propyl	methyl
	H	<u>i</u> -propyl	<u>n</u> -butyl
	methyl	methyl	methyl
10	H	benzyl	methyl
	H	<u>t</u> -butyl	n-propyl
	ethyl	ethyl	methyl
	H	phenethyl	methyl
	H	α,α -dimethylphenethyl	methyl

15


EXAMPLE 4

Preparation of ethers

Several ethers can be prepared according to the method of Example 2 as illustrated in the following equation:

20

25



30

35

-17-

wherein substituents are:

	<u>R₂</u>	<u>R₆</u>	<u>M.P. °C</u>
5	isopropyl	methyl	
	isopropyl	benzyl	
	isopropyl	allyl	
	<u>t</u> -butyl	ethyl	
	isopropyl	isopropyl	121-123
10	phenethyl	ethyl	
	α,α -dimethylphenethyl	methyl	
	α,α -dimethylphenethyl	ethyl	
	isopropyl	<u>n</u> -butyl	
	isopropyl	<u>n</u> -hexyl	
15	α,α -dimethylphenethyl	isopropyl	210-212

EXAMPLE 5

Preparation of 2'-hydroxy-5'-[1-phenoxy-2-(isopropylamino)-ethyl]methanesulfonanilide

20 A 3.25 g sample of 2'-hydroxy-5'-[1-chloro-2-(isopropylamino)ethyl]methanesulfonanilide hydrochloride is slowly added to 45 g of sodium phenoxide in 150 mL of dry tetrahydrofuran at 0 to 5°C under N₂ atmosphere. The mixture is stirred for 24 hours at ambient temperature and then poured on ice whereafter the crude product compound is separated.

EXAMPLE 6

Preparation of 2'-hydroxy-5'-[1-phenylthio-2-(isopropylamino)ethyl]methanesulfonanilide


30 The compound 2'-hydroxy-5'-[1-phenylthio-2-(isopropylamino)ethyl]methanesulfonanilide can be prepared following the method described in Example 5 with the substitution of sodium thiophenoxide for sodium phenoxide.

EXAMPLE 7Preparation of 2'-hydroxy-5'-[1-methylthio-2-(isopropyl-amino)ethyl]methanesulfonanilide

5 A 1-g sample of 2'-hydroxy-5'-[1-chloro-2-(isopropylamino)ethyl]methanesulfonanilide hydrochloride is slowly added to 5 mL of methylmercaptan in 25 mL of ethylene dichloride at 0 to -10°C under N₂ atmosphere. This mixture is first stirred several hours at ice-bath temperature and then stirred at ambient temperature for 24 hours. The mixture is filtered. The filter cake is washed with ethylene dichloride, added to water, and basified with 10% NaOH solution to pH 10 in an ice bath. The aqueous mixture is extracted several times with methylene chloride. The combined extracts are dried over MgSO₄ and evaporated to dryness thus yielding the crude product compound.

EXAMPLE 8

20 Several thioethers can be prepared using the method of Example 7 as illustrated in the following equation:




wherein substituents are:

	<u>R₂</u>	<u>R₇</u>
5	isopropyl	isopropyl
	isopropyl	benzyl
	<u>t</u> -butyl	methyl
	phenethyl	methyl
10	α,α -dimethylphenyl	methyl
	isopropyl	ethyl
	isopropyl	allyl

Use of NaSH in place of R₇SH without further
basification yields:

15



20

25

30

35

EXAMPLE 9

2'-Acetoxy-5'-[2-(isopropylamino)-1-hydroxyethyl]aniline
hydrochloride


5 A mixture containing 36.2 g (0.2 mole) of 4'-
hydroxy-3'-nitroacetophenone, 15.7 g of acetyl chloride,
22 mL of 50% KOH solution, 200 mL of H₂O and 300 mL of
95% ethanol is stirred and heated to reflux for 18 hours.
10 The mixture is evaporated to remove ethanol, and the
aqueous mixture is extracted with CHCl₃ to solubilize
the product. The CHCl₃ extract is evaporated to dryness
to afford 4'-acetoxy-3'-nitroacetophenone, which is
brominated with an equimolar quantity of Br₂ in CHCl₃
15 (100 mL) to afford 2-bromo-4'-acetoxy-3-nitroaceto-
phenone. This phenacyl bromide is then reacted in 100 mL
of ethanol with 2 equivalents of isopropylamine to
afford 4'-acetoxy-3'-nitro-2-isopropylaminoacetophenone,
which isolated as the hydrochloride by the procedure
20 reported by Larson et al. in Journal of Medicinal
Chemistry, 10, 467 (1967) for a related compound. This
ketoamine is then hydrogenated in 90% aqueous ethanol
with 0.5 g of 10% Pd-C/3.5 kg/cm² H₂ pressure in a Parr
hydrogenator until the uptake of H₂ is completed. The
25 mixture is filtered, and the filtrate is evaporated to
dryness to afford the crude title compound.

30

35

-21-

Similarly, substitution of pivaloyl chloride, butyryl chloride, propionyl chloride, isobutyryl chloride, and valeryl chloride in place of acetyl chloride in the above sequence affords the corresponding 2'-pivaloxy-, 2'-butyryloxy-2'-propionyloxy-, 2'-isobutyryloxy-, and 2'-valeryloxy-5'-[2-(isopropylamino)-1-hydroxyethyl]aniline hydrochlorides. Similarly, substitution of the isopropylamine by t-butylamine, 2-butylamine, ethylamine, n-propylamine, cyclopentylamine and α,α -dimethylphenethylamine affords the following compounds:




$\text{R}_1 = \text{t-butyl}, 2\text{-butyl}, \phi\text{CH}_2\text{C}(\text{CH}_3)-, \text{ethyl}, \text{n-propyl}, \text{cyclopentyl}.$

EXAMPLE 102'-Pivaloyloxy-5-[1-hydroxy-2-(α,α -dimethylphenethyl-amino)ethyl]methanesulfonanilide

2'-Pivaloyloxy-3'-nitroacetophenone (10 g), which is prepared by the method described in Example 9, in 200 mL of MeOH is reduced in a Parr hydrogenator with 0.2 g of PtO₂. The mixture is filtered, and the solvent is removed to afford crude 2'-pivaloyoxy-3'-aminoacetophenone. This material is stirred in 75 mL of pyridine and 4.32 g of methanesulfonyl chloride is added dropwise at 25°C. After three hours, the mixture is poured into an ice water mixture, and the precipitate is collected and washed with H₂O, dilute HCl and H₂O successively to afford 2'-pivaloyloxy-5'-acetylmethanesulfonanilide. This material is then brominated by the procedure of Example 9 to afford 2'-pivaloyloxy-5'-(2-bromoacetyl)-methanesulfonanilide, which is then reacted with isopropylamine according to the procedure of Example 9 to afford 2'-pivaloyloxy-5'-(2-isopropylaminoacetyl)-methanesulfonanilide hydrochloride. This intermediate is then reduced by the procedure of Example 9 to yield 2'-pivaloyloxy-5'-[(2-isopropylamino)-1-hydroxyethyl]-methanesulfonanilide hydrochloride.

Similarly, the following compounds are prepared by the above sequence with or without different acyl chlorides (R₄Cl) for pivaloyl chloride and with or without substitution of R₁R₂NH for isopropylamine.



-23-

	<u>R₄</u>	<u>R₁</u>	<u>R₂</u>
	CH ₃ CO	H	isopropyl
	CH ₃ CO	H	α,α -dimethylphenethyl
5	CH ₃ CO	H	<u>t</u> -butyl
	C ₂ H ₅ CO	H	isopropyl
	C ₂ H ₅ CO	H	α,α -dimethylphenethyl
	C ₂ H ₅ CO	H	<u>t</u> -butyl
	<u>n</u> -C ₃ H ₇ CO	H	isopropyl
10	<u>n</u> -C ₃ H ₇ CO	H	<u>t</u> -butyl
	<u>n</u> -C ₃ H ₇ CO	H	α,α -dimethylphenethyl
	<u>i</u> -C ₃ H ₇ CO	H	isopropyl
	<u>i</u> -C ₃ H ₇ CO	H	α,α -dimethylphenethyl
	<u>i</u> -C ₃ H ₇ CO	H	<u>t</u> -butyl
15	pivaloyl	H	<u>t</u> -butyl
	pivaloyl	H	α,α -dimethylphenethyl
	pivaloyl	H	cyclopentyl
	pivaloyl	methyl	methyl

EXAMPLE 11

20

N-{2-Hydroxy-5-[1-hydroxy-2-(isopropylamino)ethyl]-phenyl}dimethanesulfonamide

25


In 250 mL of CH₂Cl₂ containing 26 mL of triethylamine, 20 g of 3'-amino-4'-benzyloxyacetophenone is stirred at 10°C and 19 g of methanesulfonyl chloride is added gradually. The mixture is then stirred at room temperature for 24 hours and extracted with 10% aqueous NaOH solution. The CH₂Cl₂ solution is further washed with H₂O, dried over MgSO₄, and evaporated in vacuo until

30

crystallization begins to occur. The dimethanesulfonamide derivative (A) is then collected. In 100 mL of dioxane containing 7 mL of H₂O, and 7 mL of H₂O, and

8.2 g of selenium dioxide, compound A is added gradually at 70°C. Heating is continued for an overnight period, and the mixture is heated at reflux for an hour. The mixture is decolorized with activated carbon, filtered, and evaporated to dryness in vacuo. The residual oil is mixed with 500 mL of CH₂Cl₂ and washed twice with 100 mL of H₂O. The CH₂Cl₂ solution is dried over MgSO₄ and evaporated to dryness to afford the crude glyoxal, which is further dissolved in 100 mL of CH₂Cl₂. A half of this solution is stirred, and 3.2 mL of isopropyl amine is gradually added. After one hour, the mixture is evaporated to dryness; the residue is triturated with ether and filtered. This solid is stirred in 250 mL of ethanol at 15°C and 2.4 g of NaBH₄ is added in one portion. The mixture is stirred at room temperature overnight and evaporated to dryness. The residual solid is heated in 200 mL of boiling CH₂Cl₂ and filtered. This washing procedure is repeated twice. The remaining residue is partially dissolved in 30 mL of CH₂Cl₂, and the mixture is washed with H₂O (3 X 200 mL). The CH₂Cl₂ solutions are combined, washed with H₂O (3 X 300 mL), dried over MgSO₄, and evaporated to dryness to afford a yellow oil. This oil is triturated with Et₂O to afford 5.5 g of pale-yellow solid (B) after filtration. A 5-gram sample of B is added to 200 mL of MeOH containing 1 g of 5% Pd-C and hydrogenated at 50-45 p.s.i.g. until uptake of H₂ is completed. The mixture is filtered; the catalyst is washed with 50 mL of MeOH, and the methanol solutions are evaporated to dryness to afford the crude title compound, which is washed with ether; m.p. 174°C.

The following compounds are prepared in the above manner with or without substitution of methane-sulfonyl chloride with the appropriate R_8SO_2Cl and with or without substitution of isopropylamine with R_2NH_2 .



10

<u>R₈</u>	<u>R₂</u>
C ₂ H ₅	isopropyl
methyl	<u>t</u> -butyl
15 methyl	α,α-dimethylphenethyl
isopropyl	isopropyl
n-butyl	isopropyl
n-butyl	<u>t</u> -butyl
n-butyl	α,α-dimethylphenethyl
20 isopropyl	<u>t</u> -butyl
isopropyl	α,α-dimethylphenethyl
methyl	cyclopentyl
methyl	α-methylphenethyl
methyl	ethyl

25

30

35

EXAMPLE 12

Evaluation of test compounds for increasing the growth rate of animals and enhancing the lean meat-to-fat ratio thereof by reducing fat deposition in said animals and increasing the lean meat thereof

- 5 CIF female mice from Carworth Farms are received when they are six weeks old. They are housed ten to a cage in air-conditioned rooms (22°C to 25°C) with automatic diurnal illumination (14 hours light and 10 hours dark). Compounds of the invention are added to a
- 10 basic diet of "Purina" (registered Trade Mark) Laboratory Chow which is supplied ad libitum and contains the following ingredients:

TABLE IDescription of Diet

15	<u>Guaranteed Analysis</u>	<u>%</u>
	Crude protein not less than	23.0
	Crude fat not less than	4.5
	Crude Fiber not more than	6.0
	Ash not more than	9.0

20 Ingredients

- Meat and bone meal, dried skimmed milk, wheat germ meal, fish meal, animal liver meal, dried beet pulp, ground extruded corn, ground oat groats, soybean meal, dehydrated alfalfa meal, cane molasses, animal fat preserved with BHA,
- 25 vitamin B₁₂ supplement, calcium pantothenate, choline chloride, folic acid, riboflavin supplement, brewer's dried yeast, thiamin, niacin, vitamin A supplement, D-activated plant sterol, vitamin E supplement, calcium carbonate, dicalcium phosphate, iodized salt, ferric
- 30 ammonium citrate, iron oxide, manganous oxide, cobalt carbonate, copper oxide, zinc oxide. Water is also allowed ad libitum.

Various experimental treatments are randomly assigned to mice cages. Each treatment is tested with three replicates, i.e., in three cages of ten mice each.

5 There are ten control cages each of which contains ten mice. Drugs are mixed in the diet at the dosage level indicated. Feed and water are offered ad libitum over a 12-day test period. Feed spilled is collected during the test period. At the end of the experiment the collected
10 feed is weighed and the mean feed consumption per cage of ten mice is determined for each treatment. The mice are weighed as a group of ten and the weight gain is determined. The mice are sacrificed by cervical dislocation. The right uterine fat pad of each mouse is removed. The
15 fat pads for each cage of ten mice are weighed as a unit.

A correlation between the reduction in fat pad weights of treated animals and reduction in total body fat of treated animals has been previously established. This relationship was established using several treatment groups
20 in which the total body fat of treated animals was determined and found to correlate closely with reduction in fat pad weights of animals receiving the same treatment.

Results of representative experiments are presented in Table II. These data indicate that compounds
25 of the invention cause increases in growth as measured by weight gain and concomitant decreases in body fat of experimental animals.

30

35

TABLE II
Evaluation of test compounds for growth enhancement and reduction of fat pad weight in mice



Compound	Level in diet (ppm)	Gain (g)	% ± Control	Fat pad wt % ± control
	200	15.3	+ 27.5	-26.3
	100	17.0	+ 41.7	-24.5
	50	15.5	+ 29.2	- 7.1
	25	11.7	+ 2.5	-19.9
	12	12.2	+ 1.7	-12.8
	6	11.7	+ 2.5	-11.3
	100	13.1	+ 23.6	-13.5

TABLE II (Continued)





Compound	Level in diet (ppm)	Gain (g)	% Control	Fat pad wt % ± control
	200	17.3	+ 44.3	-51.1
	100	12.9	+ 7.5	-36.9
	50	15.5	+ 29.2	-25.5
	25	12.1	+ 0.8	-13.2
	12	13.5	+ 12.5	+ 4.2
	6	15.1	+ 25.8	- 7.0
-29-				
	50	18.7	+ 77.4	- 5.1

TABLE II (Continued)

Compound	Level in diet (ppm)	Gain (g)	% ± Control	Fat pad wt % ± control
 <chem>CCN(C)CCOC(=O)Nc1ccc(cc1)COCc2ccccc2</chem>	200	21.4	+ 46.6	- 7.1
 <chem>CCN(C)CCOC(=O)Nc1ccc(cc1)COCc2ccccc2</chem>	200	24.5	+ 67.8	+ 0.6
	50	22.9	+ 56.8	+ 1.8
 <chem>CCN(C)CCOC(=O)Nc1ccc(cc1)COCc2ccccc2</chem>	200	18.4	+ 41.5	+ 3.8
	50	16.9	+ 30.0	- 2.4

#4

TABLE II (Continued)


Compound	Level in diet (ppm)	Gain (g)	% ± Control	Fat pad wt % ± control
$ \begin{array}{c} \text{CH}_3 \\ \\ \text{HO}-\text{C}_6\text{H}_4-\text{CHCH}_2-\text{NH}-\text{C}-\text{CH}_2-\text{C}_6\text{H}_4-\text{CH}_2-\text{C}(\text{CH}_3)_2-\text{CH}_3 \\ \\ \text{O}-\text{CH}(\text{CH}_3)_2 \\ \text{CH}_3-\text{SO}_2-\text{NH} \end{array} \cdot \text{HCl} $	50	13.6	+ 3.8	- 7.0
$ \begin{array}{c} \text{HO}-\text{C}_6\text{H}_4-\text{CHCH}_2-\text{NH}-\text{CH}(\text{CH}_3)_2 \\ \\ \text{O}-\text{CH}(\text{CH}_3)_2 \\ \text{CH}_3-\text{SO}_2-\text{NH} \end{array} $	200	15.0	+ 41.5	+ 4.7
$ \begin{array}{c} \text{HO}-\text{C}_6\text{H}_4-\text{CHCH}_2-\text{NH}-\text{C}(\text{CH}_3)_2-\text{CH}_2-\text{C}_6\text{H}_4-\text{CH}_2-\text{C}(\text{CH}_3)_2-\text{CH}_3 \\ \\ \text{O}-\text{CH}(\text{CH}_3)_2 \\ \text{CH}_3-\text{SO}_2-\text{NH} \end{array} $	200	17.5	+ 65.1	- 15.5

TABLE II (Continued)

Compound	Level in diet (ppm)	Gain (g)	% ± Control	Fat pad wt % ± control
$ \begin{array}{c} \text{HO}-\text{C}_6\text{H}_4-\text{CH}(\text{OH})-\text{CH}_2-\text{NHC}(\text{CH}_3)_2 \cdot \frac{1}{2} \text{HOOC}-\text{C}(\text{H})=\text{C}(\text{H})-\text{COOH} \\ \\ \text{NHSO}_2\text{N}(\text{CH}_3)_2 \end{array} $ (m.p. 198-200°C)				
$ \begin{array}{c} \text{HO}-\text{C}_6\text{H}_4-\text{CH}(\text{OH})-\text{CH}_2-\text{NHC}(\text{CH}_3)_2\text{CH}_2-\text{C}_6\text{H}_4 \cdot \text{HCl} \\ \\ \text{NHSO}_2\text{N}(\text{CH}_3)_2 \end{array} $				
$ \begin{array}{c} \text{HO}-\text{C}_6\text{H}_4-\text{CH}(\text{OH})-\text{CH}_2-\text{NHC}(\text{CH}_3)_3 \cdot \text{HCl} \\ \\ \text{NHSO}_2\text{CH}_3 \end{array} $ (m.p. 236-237°C)				

46


TABLE II (Continued)

Compound	Level in diet (ppm)	Gain (g)	% ± Control	Fat pad wt % ± control
				
				



CLAIMS:

1. A compound having the structure:




wherein X is NHSO_2R_8 or $\text{N}(\text{SO}_2\text{R}_8)_2$; R_1 is hydrogen or $\text{C}_1\text{-C}_4$ alkyl; R_2 is hydrogen, $\text{C}_1\text{-C}_4$ alkyl, benzyl, phenethyl, α -methylphenethyl, or α,α -dimethylphenethyl; R_3 is OR_6 or SR_7 ; R_8 is $\text{C}_1\text{-C}_4$ alkyl; R_6 is $\text{C}_1\text{-C}_6$ alkyl, benzyl, phenyl or allyl; R_7 is hydrogen, $\text{C}_1\text{-C}_6$ alkyl, benzyl, phenyl or allyl; with the provisos that when R_2 is benzyl, phenethyl α -methylphenethyl or α,α -dimethylphenethyl, R_1 of NR_1R_2 is hydrogen; and when X is $\text{N}(\text{SO}_2\text{R}_8)_2$, R_3 is OH; and the optically active isomers, and the pharmacologically acceptable acid addition and metal salts thereof.

2. A compound according to Claim 1, wherein R_3 is OR_6 .

3. A compound according to Claim 1, N-{5-[2-[(α,α -dimethylphenethyl)amino]-1-hydroxyethyl]-2'-hydroxyphenyl}dimethanesulfonamide, and the optical isomers and acid addition salts thereof.

4. A method for increasing the growth rate and the lean meat-to-fat ratio of meat-producing animals, comprising administering either orally or parenterally to meat-producing animals a growth-promoting and lean meat-enhancing amount of a compound having the formula





wherein, R_1 is hydrogen or C_1-C_4 alkyl; R_2 is hydrogen, C_1-C_4 alkyl, cycloalkyl C_3-C_5 , benzyl, phenethyl, α -methylphenethyl or α,α -dimethylphenethyl; R_3 is OH, OR_6 or SR_7 ; R_4 is hydrogen, C_1-C_4 alkyl, benzyl or C_2-C_5 alkanoyl; R_5 is hydrogen, methyl or ethyl; R_6 is C_1-C_6 alkyl, benzyl, phenyl or allyl; R_7 is hydrogen, C_1-C_6 alkyl, benzyl, phenyl or allyl; X is $NHSO_2R_8$, $N(SO_2R_8)_2$, $N(R_1)_2$, $NHCO_2R_1$, $NH-CO-N(R_1)_2$, $NHSO_2N(R_1)_2$, $NHCOOR_9$, or $NHCOO$ -benzyl; R_8 is C_1-C_4 alkyl; R_9 is C_1-C_5 alkyl; with the provisos that when R_2 is cycloalkyl C_3-C_5 , benzyl, phenethyl, α -methylphenethyl or α,α -dimethylphenethyl, R_1 of NR_1R_2 is hydrogen; and when R_3 is OR_6 or SR_7 , X is only $NHSO_2R_8$; and further that the R_1 group in $(R_1)_2$ may or may not be identical substituents; and the optically active isomers and pharmacologically acceptable acid addition and metal salts thereof.

5. A method according to Claim 4 wherein R_4 is hydrogen; R_3 is OH; R_5 is hydrogen; R_1 is hydrogen or C_1-C_4 alkyl; R_2 is hydrogen, C_1-C_4 alkyl, α,α -dimethylphenyl or α -methylphenethyl; X is $NHSO_2CH_3$, $N(R_1)_2$, $NHCO_2R_1$, $NHCO_2R_1$, $NH-CO-N(R_1)_2$, $NHSO_2N(R_1)_2$ or $NHCOOR_9$, where R_9 is C_1-C_5 alkyl; with the provisos that when R_2 is α -methylphenethyl or α,α -dimethylphenethyl, R_1 of NR_1R_2 is hydrogen; and both R_1 in $(R_1)_2$ may be the same or different substituents; and the optically active isomers and pharmacologically acceptable acid addition and metal salts thereof.


6. A method according to Claim 4 wherein the compound is 2'-hydroxy-5'-[1-hydroxy-2-(isopropylamino)ethyl]methanesulfonanilide hydrochloride.

7. A method according to Claim 4 wherein the compound is 2'-hydroxy-5'-(2-[(α,α -dimethylphenethyl)-amino]-1-hydroxyethyl)methanesulfonanilide hydrochloride.



8. A method according to Claim 4 wherein the compound is 2'-hydroxy-5'-[1-hydroxy-2-(t-butylamino)-ethyl]methanesulfonanilide hydrochloride.

9. An animal feed composition which, for increasing the growth and lean meat-to-fat ratio of domestic and farm animals, contains a compound having the formula:



wherein, R_1 is hydrogen or C_1 - C_4 alkyl; R_2 is hydrogen, C_1 - C_4 alkyl, cycloalkyl C_3 - C_5 , benzyl, phenethyl, α -methylphenethyl or α,α -dimethylphenethyl; R_3 is OH , OR_6 or SR_7 ; R_4 is hydrogen, C_1 - C_4 alkyl, benzyl or C_2 - C_5 alkanoyl; R_5 is hydrogen, methyl or ethyl; R_6 is C_1 - C_6 alkyl, benzyl, phenyl or allyl; R_7 is hydrogen, C_1 - C_6 alkyl, benzyl, phenyl or allyl; X is NHSO_2R_8 , $\text{N}(\text{SO}_2\text{R}_8)_2$, $\text{N}(\text{R}_1)_2$, NHCOR_1 , $\text{NH-CO-N}(\text{R}_1)_2$, $\text{NHSO}_2\text{N}(\text{R}_1)_2$, NHCOOR_9 , or NHCOO-benzyl ; R_8 is C_1 - C_4 alkyl; R_9 is C_1 - C_5 alkyl; with the provisos that when R_2 is cycloalkyl C_3 - C_5 , benzyl, phenethyl, α -methylphenethyl or α,α -dimethylphenethyl, R_1 of NR_1R_2 is hydrogen; and when R_3 is OR_6 or SR_7 , X is only NHSO_2R_8 ; and further that the R_1 group in $(\text{R}_1)_2$ may or may not be identical substituents; and the optically active isomers and pharmacologically acceptable acid addition and metal salts thereof.

10. An animal feed composition according to Claim 9, wherein said compound is as defined in any one of Claims 5-8.

11. An animal feed composition according to Claim 9 or Claim 10, which is an animal feed premix, concentrate or supplement.



12. A compound according to Claim 1, substantially as described in any one of Examples 2, 4-8 or 11 herein.

13. A method for increasing the growth rate and the lean meat-to-fat ratio of meat-producing animals, according to Claim 4 and substantially as hereinbefore described.

14. An animal feed composition, according to Claim 9 and substantially as hereinbefore described.
