

BIPARATOPIC POLYPEPTIDES ANTAGONIZING WNT SIGNALING IN TUMOR CELLS

5

FIELD OF THE INVENTION

The present invention relates to novel low-density lipoprotein receptor-like protein 5 (LRP5) and low-density lipoprotein receptor-like protein 6 (LRP6) binding polypeptides. The invention also relates to nucleic acids encoding such polypeptides; to methods for preparing such polypeptides; to host cells expressing or capable of expressing such polypeptides; to compositions comprising such polypeptides; and to uses of such polypeptides or such compositions, in particular for therapeutic purposes in the field of cancer diseases.

15

BACKGROUND OF THE INVENTION

Activation of the Wnt signaling pathway requires binding of extracellular Wnt ligands to the Frizzled receptor and to the co-receptor LRP5 (Accession number: UniProtKB -O75197 / LRP5_HUMAN) or its closely related homologue LRP6 (Accession number: UniProtKB - O75581 / LRP6_HUMAN). There are 19 Wnt proteins and 10 Frizzled receptors in mammalian cells. In the absence of Wnt ligand, cytoplasmic beta-catenin is phosphorylated by a protein complex consisting of the scaffolding proteins Axin and APC and the kinases GSK3beta and CK1a. Subsequent recognition by the ubiquitin ligase beta-TrcP leads to ubiquitin-mediated degradation of beta-catenin. In the presence of Wnt ligand, binding of Wnt to Frizzled and LRP5 or LRP6 leads to recruitment of the cytoplasmic effector protein Dvl and phosphorylation of the LRP5 or LRP6 cytoplasmic tail, which provides the docking site for Axin. Axin sequestration by LRP5 or LRP6 leads to the inactivation of the Axin-APC-GSK3beta complex and, therefore, intracellular beta-catenin stabilization and accumulation. Hence,

20
25
30

cytoplasmic levels of beta-catenin rise, and beta-catenin migrates to the nucleus and complexes with members of the T-cell factor (TCF)/Lymphoid enhancer-binding factor (LEF) family of transcription factors. Basal transcription machinery and transcriptional co-activators are then recruited, including cAMP response element-binding protein (CREB)-binding protein (CBP) or its homolog p300,
5 leading to expression of various target genes, including Axin2, cyclin D1 and c-Myc.

An additional level of ligand-dependent Wnt pathway regulation is mediated by
10 the E3 ligase RNF43, and its closely related homologue ZNRF3, and by the secreted R-Spondin proteins (de Lau *et al.* "The R-spondin/Lgr5/Rnf43 module: regulator of Wnt signal strength". *Genes Dev.* 2014; 28(4):305-16). RNF43 mediates the ubiquitination of the Frizzled/LRP5 or LRP6 receptor complex at the cell surface, leading to its degradation and, thereby, inhibiting ligand-dependent
15 Wnt pathway activity. The activity of RNF43 is counteracted by the R spondin family members (R-spondin 1 to 4 ligands). When R-Spondin ligand is present, it removes RNF43 from the cell surface, allowing Frizzled/LRP5 or LRP6 complex accumulation and enhancement of Wnt signaling in the presence of Wnt ligands.

LRP5 and LRP6 function as gatekeepers of ligand dependent Wnt signaling activation and, therefore, may be considered as targets to achieve complete blockade of the pathway mediated by all 19 Wnt ligands and 10 Frizzled receptors and enhanced by R-spondin ligands. In particular, Wnt ligands can be divided into a Wnt1 class and a Wnt3a class, each binding to different
20 epitopes/regions of LRP5 and LRP6 for signaling. The ectodomain of LRP5 and LRP6 comprises four repeating units of a beta-propeller connected to an EGF-like domain, followed by three LDLR-type A repeats. Combined structural and functional analyses of LRP5 and LRP6 suggest that Wnt1 (Wnt1-class ligand)
25 binds to a fragment containing beta-propeller 1 and 2 and Wnt3a binds to a fragment containing beta-propeller 3 and 4 of LRP6. So far, only a low-resolution
30 picture of LRP6 ectodomain containing beta-propeller from 1 to 4 regions is

reported (Ahn *et al.* "Structural basis of Wnt signaling inhibition by Dickkopf binding to LRP5/6". *Dev Cell*. 2011; 21(5):862-73). However, the uncertainties of these low-resolution reconstructions (40 Å) and the absence of structural data of the LRP6 ectodomain in complex with the Wnt ligands do not allow defining the exact epitopes involved in Wnt1 or Wnt3a ligand binding.

Hyperactivation of Wnt signaling is involved in the pathogenesis of various types of cancer. In some cancer types frequent mutations in downstream signaling molecules contribute to constitutively activated Wnt pathway (e.g. APC mutations in colorectal cancer; beta-catenin activating mutation in hepatocellular carcinoma). In contrast, in Triple Negative Breast Cancer (TNBC), Non Small Cell Lung Cancer (NSCLC), pancreatic adenocarcinoma and in a subset of Colorectal Cancer (CRC) and endometrial cancers, Wnt signaling activation is driven by a ligand dependent mechanism (i.e. by an autocrine/paracrine Wnt activation), as detected by beta-catenin intracellular accumulation. In NSCLC, TNBC and pancreatic adenocarcinoma, ligand dependent Wnt activation is mediated by multiple mechanisms, including increased expression of the Wnt ligands and/or of LRP5 and LRP6 receptors, or silencing of LRP5 and LRP6 negative regulator DKK1 (TNBC: Liu *et al.* "LRP6 overexpression defines a class of breast cancer subtype and is a target for therapy". *Proc Natl Acad Sci U S A* 2010; 107(11):5136-41; Khramtsov *et al.* "Wnt/beta-catenin pathway activation is enriched in basal-like breast cancers and predicts poor outcome". *Am J Pathol*. 2010; 176(6): 2911-20; NSCLC: Nakashima *et al.* "Wnt1 overexpression associated with tumor proliferation and a poor prognosis in non-small cell lung cancer patients". *Oncol Rep*. 2008; 19(1):203-9; Pancreatic cancer: Zhang *et al.* "Canonical wnt signaling is required for pancreatic carcinogenesis". *Cancer Res*. 2013; 73(15):4909-22). In particular, published data have shown that in healthy tissues (e.g. mammary and lung epithelium), beta-catenin is localized solely at the plasma membrane. In contrast, the majority of TNBC, NSCLC and pancreatic adenocarcinoma primary clinical samples showed beta-catenin intracellular accumulation (i.e. in the cytoplasm/ nucleus; biomarker of Wnt signaling

activation), due to aberrant Wnt signaling. Recent publications have shown that ligand dependent Wnt signaling activation is mediated by mutated/inactivated RNF43 (Giannakis *et al.* "RNF43 is frequently mutated in colorectal and endometrial cancers". *Nat Genet.* 2014; 46(12):1264-6) or by activating R-Spondin fusion transcripts (encoding R-spondin2 or R-spondin3 proteins driven by constitutively active strong promoters; Seshagiri *et al.* "Recurrent R-spondin fusions in colon cancer". *Nature* 2012; 488(7413):660-4) in a subset of CRC and endometrial cancers. Inactivating RNF43 mutations and R-Spondin fusion transcripts have both been shown to augment ligand dependent Wnt signaling *in vitro* by increasing the abundance of Frizzled on the cell surface. Ligand dependent Wnt activation in tumors was shown to drive tumor growth and resistance to chemotherapy or immunotherapy, and is linked to recurrence in pre-clinical models.

Some LRP5 or LRP6 binding molecules, able to modulate the Wnt signaling pathway, are known in the art:

Dickkopf-1 (DKK1) is a LRP5 and LRP6 inhibitor. DKK1 associates with both the Wnt co-receptors, LRP5 and 6, and the transmembrane protein, Kremen, inhibits Wnt signaling and leads to rapid LRP5 and LRP6 internalization. It is shown that DKK1 inhibits both Wnt1 and Wnt3a mediated signaling. Structural modeling studies show that a single DKK1 molecule cooperatively binds to an extended region of the LRP6 ectodomain (from beta-propeller 1 to 3). The structural analyses suggest a DKK1 cooperative binding-interaction with LRP6 with an initial binding to the beta-propeller 3 region that facilitates the interaction/binding to the beta-propeller 1 and 2 region via a conformation change of the LRP6 ectodomain. However, elucidation of the defined epitopes within the beta-propeller 1, 2 and 3 domains involved in the DKK1 binding to LRP6 is lacking due to the low resolution of the structural reconstructions of the full LRP6 ectodomain bound to DKK1, as mentioned.

It was shown that DKK1 treatment *in vivo* causes severe toxicity in the gastrointestinal tract. In particular, it was shown that adenovirus mediated expression of DKK1 in adult mice markedly inhibited proliferation in small intestine and colon, accompanied by progressive architectural
5 degeneration, severe body weight loss and mortality from colitis and systemic infection. In particular, LRP5 and LRP6 are expressed in the intestine in the proliferative epithelial cells and are required for proliferation of the intestinal epithelium, suggesting that LRP5 and LRP6 inhibition may be toxic for this and other normal tissues (Zhong *et al.* "Lrp5 and Lrp6 play compensatory roles in
10 mouse intestinal development". *J Cell Biochem.* 2012; 113(1):31-8). This makes it doubtful whether agents which inhibit LRP5 and LRP6, or which inhibit the Wnt (Wnt1 and Wnt3a) signaling pathway in general, can be used for therapeutic purposes, e.g. can be developed as anti-cancer drugs.

15 WO2009/056634 refers to LRP6 binding molecules that may either interact with the Wnt1 signaling pathway or with the Wnt3/3a signaling pathway, which may be antagonistic or agonistic, and which may be used for diagnostic purposes or to treat "Wnt signaling-related disorders", such as osteoarthritis, polycystic kidney disease, or cancer. No specific examples for such binding molecules, defined by
20 their amino acid sequence, are provided in this document.

WO2011/138391 and WO2011/138392 are disclosing multivalent LRP6 binding antibodies. WO2011/138391 is claiming antibodies which are blocking one Wnt signaling pathway (Wnt1 or Wnt3) without potentiating the other pathway (Wnt3
25 or Wnt1, respectively). WO2011/138392 i.a. provides antibodies or antibody fragments which potentiate Wnt signaling by LRP6 receptor clustering.

WO2011/138391 explains that for achieving the desired effect, LRP6 binding molecules need to be formatted into full length IgG antibodies. Examples of LRP6
30 biparatopic molecules are provided which include an IgG molecule, having a first binding specificity, coupled to a single chain Fv portion, having the second

binding specificity. Some formats are described as having significantly reduced thermal stability (T_m of 50 to 52°C). An Fc portion may impart effector functions on an IgG molecule, such as complement-dependent cytotoxicity (CDC) or antibody-dependent cellular toxicity (ADCC).

5

WO2013/067355 discloses half-life extended biparatopic LRP6 binding scFv immunoglobulin constructs, derived from IgG molecules disclosed in WO2011/138391.

10

WO2011/119661 discloses antibodies that bind to LRP6 and inhibit the signaling induced by a first Wnt isoform, esp. by Wnt3 or Wnt3a, but potentiate signaling induced by a second Wnt isoform, which may be a Wnt1, 2, 2b, 4, 6, 7a, 7b, 8a, 9a, 9b, 10a or 10b isoform. Bispecific molecules are disclosed which bind to the E1-E2 region of LRP6 as well as to the E3-E4 region of LRP6. The knob-in-hole technique was used to generate bispecific antibodies.

15

Identification of the binding epitopes (defined amino acid residues within the LRP6 ectodomain / beta-propeller regions) involved in the binding of the LRP6 antibodies is not provided in WO2009/056634, nor in WO2011/138391 or WO2013/067355, and only partially in WO2011/119661. In particular, LRP6 binding antibodies can inhibit Wnt signaling via alternative mechanisms according to binding to different regions of LRP6, including competing with Wnts directly or inhibiting formation of ternary receptor complexes (Wnt-LRP6-Frizzled), whereas others enhance signaling, possibly by receptor clustering (Ahn *et al.* "Structural basis of Wnt signaling inhibition by Dickkopf binding to LRP5/6". *Dev Cell*. 2011; 21(5):862-73).

20

25

However, none of the binding molecules described in the art has so far been authorized by health authorities for the use as a medicament to treat any disease. Specifically, such use requires very specific binding properties, the right specificity, so that such molecules does or does not bind, activate or inhibit other

30

targets (e.g. resulting in undesired activation or inhibition of other signaling pathways, or lack of activation or inhibition with respect to target isoforms), in the case of bi- or multispecific agents the right balance between the two or more binding specificities, suitable pharmacokinetic and –dynamic properties, an acceptable toxicological profile, and of course *in vivo* efficacy.

In view of the above, there is a need for novel therapeutic agents that allow an efficient treatment of several types of cancer diseases and tumors. It is thus an object of the invention to provide such pharmacologically active agents that can be used in the treatment of several cancer diseases, including NSCLC and TNBC.

In particular, it is an object of the invention to provide such pharmacologically active agents, compositions and/or methods of treatment that provide certain advantages compared to the agents, compositions and/or methods currently used and/or known in the art. These advantages include *in vivo* efficacy, improved therapeutic and pharmacological properties, less side effects, and other advantageous properties such as improved ease of preparation or reduced costs of goods, especially as compared to candidate drugs already known in the art.

BRIEF SUMMARY

According to a first aspect of the disclosure, the present disclosure provides polypeptides which specifically bind to LRP5 or LRP6, wherein such polypeptide of the disclosure comprises a first immunoglobulin single variable domain (a) selected from the group of immunoglobulin single variable domains (i) to (iii) defined by having the following CDR sequences:

(i):

CDR1: TYTVG (= SEQ ID NO:1)

CDR2: AIRRRGSSTYYADSVKG (= SEQ ID NO:2)

CDR3: DTRTVALLQYRYDY (= SEQ ID NO:3)

(ii):

CDR1: SYAMG (= SEQ ID NO:4)

CDR2: AIRRSGRRTYYADSVKG (= SEQ ID NO:5)

CDR3: ARRVRSSSTRYNTGTWWWEY (= SEQ ID NO:6)

(iii):

CDR1: RYTMG (= SEQ ID NO:7)

CDR2: AIVRSGGSTYYADSVKG (= SEQ ID NO:8)

CDR3: DRRGRGENYILLYSSGRYEY (= SEQ ID NO:9),

and a second immunoglobulin single variable domain (b) selected from the group of immunoglobulin single variable domains (iv) and (v) defined by having the following CDR sequences:

(iv):

CDR1: SYAMG (= SEQ ID NO:10)

CDR2: AISWSGGSTYYADSVKG (= SEQ ID NO:11)

CDR3: SPIPYGSLLRRRNNDY (= SEQ ID NO:12)

(v):

CDR1: SYAMG (= SEQ ID NO:13)

CDR2: AISWRSGSTYYADSVKG (= SEQ ID NO:14)

CDR3: DPRGYGVAYVSAYYEY (= SEQ ID NO:15).

The terms “first” and “second” with respect to such immunoglobulin single variable domains is solely intended to indicate that these domains are two different domains (as they will at least include different CDR sequences). Thus, these terms shall not be understood to refer to the exact order or sequence of the domains within such polypeptide chain.

The polypeptides of the disclosure optionally comprise a third immunoglobulin single variable domain, such as especially an albumin binding immunoglobulin single variable domain, such as the Alb11 domain, comprising the following CDRs:

CDR1(Alb11): SFGMS (= SEQ ID NO:16)

CDR2(Alb11): SISGSGSDTLYADSVKG (= SEQ ID NO:17)

CDR3(Alb11): GGSLSR (= SEQ ID NO:18).

According to a more specific example, the polypeptides of the disclosure include immunoglobulin single variable domains which are VHH domains, and preferably humanized VHH domains.

According to an even more specific example, the polypeptides of the disclosure include a first immunoglobulin single variable domain (a) selected from the group of immunoglobulin single variable domains (i) to (iii) having the following sequences:

(i):

AVQLVESGGGLVQPGGSLRLSCAASGRTFSTYTVGWFRQAPGKEREF
VAAIRRRGSSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYY
CAADTRTVALLQYRYDYWGQGTLVTVSS

(= SEQ ID NO:19)

(ii):

AVQLVESGGGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAPGKEREF
FVAAIRRRSGRRTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYY
CAAARRVRSSTRYNTGTWWWEYWGQGTLVTVSS

(= SEQ ID NO:20)

(iii):

AVQLVESGGGLVQPGGSLRLSCAASGLTFSRYTMGWFRQAPGKEREF
VAAIVRSGGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYY
CAADRRGRGENYILLYSSGRYEWGQGTLVTVSS

(= SEQ ID NO:21),

and a second immunoglobulin single variable domain (b) selected from the group consisting of immunoglobulin single variable domains (iv) and (v) having the following sequences:

(iv):

EVQLVESGGGLVQPGGSLRLSCAASGRTFSSYAMGWFRQAPGKERE
FVAAISWSGGSTYYADSVKGRFTISRDNKNTVYLQMNSLRPEDTAVY
YCAASPIPYGSLLRRRNNDYWGQGTLVTVSS

5 (= SEQ ID NO:22), and

(v):

EVQLVESGGGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAPGKERE
FVAAISWRSGSTYYADSVKGRFTISRDNKNTVYLQMNSLRPEDTAVY
YCAADPRGYGVAYVSAYEYWGQGTLVTVSS

10 (= SEQ ID NO:23).

According to a specifically preferred example, the polypeptides of the disclosure additionally include a half-life extending moiety, wherein said half-life extending moiety is covalently linked to said polypeptide and is optionally selected from the group consisting of an albumin binding moiety, such as an albumin binding peptide or an albumin binding immunoglobulin domain, preferably an albumin binding immunoglobulin single variable domain, more preferably the Alb11 domain, a transferrin binding moiety, such as an anti-transferrin immunoglobulin domain, a polyethylene glycol molecule, human serum albumin, and a fragment of human serum albumin.

Specifically preferred are polypeptides which include, in addition to the two immunoglobulin single variable domains (a) and (b) as outlined above, an Alb11 domain having the following sequence:

EVQLVESGGGLVQPGNSLRLSCAASGFTFSSFGMSWVRQAPGKGLEW
VSSISGSGSDTLYADSVKGRFTISRDNKNTTLYLQMNSLRPEDTAVYYCTI
GGSLSRSSQGTLVTVSS

(= SEQ ID NO:24)

According to a further example, the disclosure specifically includes polypeptides comprising or consisting of any of the following three polypeptide chains:

F13500575, having the sequence SEQ ID NO:25,

F13500571, having the sequence SEQ ID NO:26, and

5 F13500720, having the sequence SEQ ID NO:27.

According to another aspect of the disclosure, there is provided a polypeptide selected from the group of polypeptides comprising or consisting of SEQ ID NO:25, SEQ ID NO:26, and SEQ ID NO:27.

10

The disclosure also relates to nucleic acid molecules, expression vectors, host cells, and methods of manufacturing used in the production of a polypeptide of the disclosure. Nucleic acid molecules encoding the polypeptides of the disclosure can be used, in an isolated form, for constructing respective
15 expression vectors, which then may be transfected into host cells used for biopharmaceutical production of the polypeptides of the disclosure. Such method of manufacturing typically comprises the steps of culturing the host cell under conditions that allow expression of the polypeptide, recovering the polypeptide and purifying it according to methods known in the art.

20

According to another aspect of the disclosure, there is provided an expression vector comprising a nucleic acid encoding a polypeptide as defined herein.

According to another aspect of the disclosure, there is provided a host cell
25 carrying an expression vector as defined herein.

According to another aspect of the disclosure, there is provided a method of manufacturing a polypeptide as defined herein, comprising the steps of culturing a host cell as defined herein under conditions that allow expression of a polypeptide as defined herein; and recovering and optionally purifying said polypeptide.

According to another aspect of the disclosure, there is provided a pharmaceutical composition comprising (i) as the active ingredient, the polypeptide as defined herein, and (ii) a pharmaceutically acceptable carrier, and optionally (iii) a diluent, excipient, adjuvant and/or stabilizer.

Further aspects, embodiments, examples, uses and methods involving the polypeptides of the disclosure will become clear from the following detailed description of the disclosure and from the appended claims.

The disclosure provides for novel molecules that allow a more efficient treatment of several cancer types, such as TNBC, CRC, and NSCLC, with less side effects. The polypeptides of the disclosure provide for a surprising therapeutic effect (i.e. efficacy) in the treatment of cancer patients, in that they may induce tumor regression resulting in pathological complete response (pCR). This, in turn, is expected to result in significant improvement of progression free survival and overall survival, especially in high unmet medical need indications such as e.g. in breast cancer. Thus, the polypeptides of the disclosure provide for novel therapeutic options in the treatment of several cancer types, esp. those showing a deregulated Wnt signaling pathway and beta-catenin accumulation.

According to another aspect of the disclosure, there is provided the polypeptide as defined herein for use as a medicament in a method for the treatment, prevention or alleviation of a disease, disorder or condition in a human being or an animal, preferably for use in a method (a) for the treatment of cancer, such as breast cancer, lung cancer, pancreatic cancer, colorectal cancer, sarcomas,

ovarian cancer or hepatocellular carcinoma, or (b) for the treatment of idiopathic pulmonary disease, or (c) for the treatment of a retinopathy caused by abnormal Wnt signaling.

- 5 According to another aspect, there is provided a polypeptide as defined herein for use in the treatment of cancer in combination with an immune checkpoint inhibitor selected from the group consisting of anti-PD1 antibodies, anti-PDL1 antibodies, anti-CTLA4 antibodies, anti-BTLA antibodies, anti-LAG3 antibodies, and anti-TIM3 antibodies, or in combination with a cancer vaccine.

10

Furthermore, the polypeptides of the disclosure are easy to manufacture, have a high stability and low antigenicity, and offer a variety of options regarding administration routes, in addition to injection and infusion.

5 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a schematic representation of biparatopic polypeptides antagonizing Wnt1 and Wnt3a signaling. They consist of three domains, with two domains binding to distinct epitopes of LRP5 and LRP6 (Wnt1 and Wnt3a
10 blocker) and one domain for half-life extension (human serum albumin binder).

Figure 2 shows the lack of correlation between binding FACS and ELISA assays for a representative number of LRP6 binding VHHs derived from the llama immunization. The number “1” panel of VHHs is characterized by high affinity to
15 cells expressing LRP6 on the plasma membrane, as detected by FACS binding assays (on the y axis MCF values are reported). The number “2” panel of VHHs is characterized by high affinity to recombinant human LRP6 ectodomain (rhLRP6-Fc), as detected by ELISA binding assays (on the x axis OD405 values are reported).

20 **Figure 3** shows the binding of three half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs to human LRP5 (Figure 3A) and human LRP6 (Figure 3B) overexpressing HEK293 cell lines when compared to the negative control consisting of a non-targeting binder (VHH construct that binds to a
25 bacterial protein which is not expressed in HEK293 cells).

Figure 4 shows complete DKK1 competition of three half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs for binding to both human LRP5 (Figure 4A) and human LRP6 (Figure 4B) overexpressing HEK293 cell
30 lines, as detected by a FACS-based DKK1 competition assay.

Figure 5 shows complete inhibition of Wnt1 and Wnt3a pathway of three half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs (Figure 5A) and comparison to other LRP6 binding molecules (Figure 5B: Knob HC YW210.09 and MOR08168IgG1LALA 6475 scfv; Figure 5C: 802T) in combined Wnt1 and Wnt3a reporter assay.

Figure 6 shows inhibition of Wnt signaling in cancer cells, as detected by inhibition of relative Axin2 mRNA expression (Figure 6A), and cell proliferation (Figure 6B) as detected by decreased percentage (%) of viable cells, after treatment with three half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs (final concentration of 1mM) and by treatment with 802T, compared to F013500571 and untreated (control) cells (left-hand side / right-hand side diagram of Figure 6B); dose-response curves are shown for treatment with one half-life extended biparatopic LRP5/LRP6 construct (Figure 6C) and 802T (Figure 6D).

Figure 7 shows *in vivo* efficacy of the half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs (F013500571 in Figure 7A and F013500720 in Figure 7B) and of the of Knob HC YW210.09 (Figure 7C) in a Wnt driven tumor models (MMTV-Wnt1 xenograft model).

Figure 8 shows Wnt pathway inhibition in tumors treated with the half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs F013500571 and F013500720, as detected by reduction of Axin2 mRNA expression relative to the control group.

Figure 9 shows the effect of Wnt3a driven signaling inhibition on pro-inflammatory cytokine TNFalpha release by dendritic cells (Figure 9A) and the effect on T-cell activation, as determined by interferon-gamma release (Figure 9B) upon treatment with a half-life extended biparatopic LRP5/LRP6

cross-reactive VHH construct. Each symbol represents a unique dendritic cell (DC) donor. Data shown are normalized to TNFalpha levels of the untreated control (Figure 9A) and each symbol represents a unique donor pair for DC and T cells (Panel B).

5

DETAILED DESCRIPTION OF THE INVENTION

Definitions

10 The above and other aspects and embodiments of the invention will become clear from the further description herein, in which:

a) Unless indicated or defined otherwise, all terms used have their usual meaning in the art, which will be clear to the skilled person. Reference is for example made
15 to the standard handbooks, such as Sambrook et al, "Molecular Cloning: A Laboratory Manual" (2nd Ed.), Vols. 1-3, Cold Spring Harbor Laboratory Press (1989); Lewin, "Genes IV", Oxford University Press, New York, (1990), and Roitt et al., "Immunology" (2nd Ed.), Gower Medical Publishing, London, New York (1989), as well as to the general background art cited herein. Furthermore, unless
20 indicated otherwise, all methods, steps, techniques and manipulations that are not specifically described in detail can be performed and have been performed in a manner known *per se*, as will be clear to the skilled person. Reference is for example again made to the standard handbooks, to the general background art referred to above and to the further references cited therein.

25

b) Unless indicated otherwise, the terms "*immunoglobulin*" and "*immunoglobulin sequence*" - whether used herein to refer to a *heavy chain antibody* or to a *conventional 4-chain antibody* - are used as general terms to include both the full-size antibody, the individual chains thereof, as well as all parts, domains or
30 fragments thereof (including but not limited to antigen-binding domains or fragments such as VHH domains or VH/VL domains, respectively). In addition,

the term "sequence" as used herein (for example in terms like "immunoglobulin sequence", "antibody sequence", "(single) variable domain sequence", "VHH sequence" or "protein sequence"), should generally be understood to include both the relevant amino acid sequence as well as nucleic acid sequences or nucleotide sequences encoding the same, unless the context requires a more limited interpretation;

c) The term "*domain*" (of a polypeptide or protein) as used herein refers to a folded protein structure which has the ability to retain its tertiary structure independently of the rest of the protein. Generally, domains are responsible for discrete functional properties of proteins, and in many cases may be added, removed or transferred to other proteins without loss of function of the remainder of the protein and/or of the domain.

d) The term "*immunoglobulin domain*" as used herein refers to a globular region of an antibody chain (such as e.g. a chain of a conventional 4-chain antibody or of a heavy chain antibody), or to a polypeptide that essentially consists of such a globular region. Immunoglobulin domains are characterized in that they retain the immunoglobulin fold characteristic of antibody molecules, which consists of a 2-layer sandwich of about 7 antiparallel beta-strands arranged in two beta-sheets, optionally stabilized by a conserved disulphide bond.

e) The term "*immunoglobulin variable domain*" as used herein means an immunoglobulin domain essentially consisting of four "framework regions" which are referred to in the art and hereinbelow as "framework region 1" or "FR1"; as "framework region 2" or "FR2"; as "framework region 3" or "FR3"; and as "framework region 4" or "FR4", respectively; which framework regions are interrupted by three "complementarity determining regions" or "CDRs", which are referred to in the art and hereinbelow as "complementarity determining region 1" or "CDR1"; as "complementarity determining region 2" or "CDR2"; and as "complementarity determining region 3" or "CDR3", respectively. Thus, the general structure or sequence of an immunoglobulin variable domain can be

indicated as follows: FR1 - CDR1 - FR2 - CDR2 - FR3 - CDR3 - FR4. It is the immunoglobulin variable domain(s) that confer specificity to an antibody for the antigen by carrying the antigen-binding site.

5 f) The term "*immunoglobulin single variable domain*" as used herein means an immunoglobulin variable domain which is capable of specifically binding to an epitope of the antigen without pairing with an additional variable immunoglobulin domain. One example of immunoglobulin single variable domains in the meaning of the present invention are "*domain antibodies*", such as the immunoglobulin
10 single variable domains VH and VL (VH domains and VL domains). Another important example of immunoglobulin single variable domains are "*VHH domains*" (or simply "VHHs") from camelids, as defined hereinafter.

In view of the above definition, the antigen-binding domain of a conventional
15 4-chain antibody (such as an IgG, IgM, IgA, IgD or IgE molecule; known in the art) or of a Fab fragment, a F(ab')₂ fragment, an Fv fragment such as a disulphide linked Fv or a scFv fragment, or a diabody (all known in the art) derived from such conventional 4-chain antibody, would normally not be regarded as an immunoglobulin single variable domain, as, in these cases, binding to the
20 respective epitope of an antigen would normally not occur by one (single) immunoglobulin domain but by a pair of (associating) immunoglobulin domains such as light and heavy chain variable domains, i.e. by a VH-VL pair of immunoglobulin domains, which jointly bind to an epitope of the respective antigen.

25 f1) "*VHH domains*" , also known as VHHs, V_HH domains, VHH antibody fragments, and VHH antibodies, have originally been described as the antigen binding immunoglobulin (variable) domain of "heavy chain antibodies" (i.e. of "antibodies devoid of light chains"; Hamers-Casterman C, Atarhouch T, Muyldermans S, Robinson G, Hamers C, Songa EB, Bendahman N, Hamers R.:
30 "Naturally occurring antibodies devoid of light chains"; Nature 363, 446-448

(1993)). The term "VHH domain" has been chosen in order to distinguish these variable domains from the heavy chain variable domains that are present in conventional 4-chain antibodies (which are referred to herein as "V_H domains" or "VH domains") and from the light chain variable domains that are present in
5 conventional 4-chain antibodies (which are referred to herein as "V_L domains" or "VL domains"). VHH domains can specifically bind to an epitope without an additional antigen binding domain (as opposed to VH or VL domains in a conventional 4-chain antibody, in which case the epitope is recognized by a VL domain together with a VH domain). VHH domains are small, robust and efficient
10 antigen recognition units formed by a single immunoglobulin domain.

In the context of the present invention, the terms VHH domain, VHH, V_HH domain, VHH antibody fragment, VHH antibody, as well as "Nanobody[®]" and "Nanobody[®] domain" ("Nanobody" being a trademark of the company Ablynx
15 N.V.; Ghent; Belgium) are used interchangeably and are representatives of immunoglobulin single variable domains (having the structure: FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4 and specifically binding to an epitope without requiring the presence of a second immunoglobulin variable domain), and which may also be distinguished from VH domains by the so-called "hallmark residues", as defined
20 in e.g. WO2009/109635, Fig. 1.

The amino acid residues of a VHH domain are numbered according to the general numbering for V_H domains given by Kabat *et al.* ("Sequence of proteins of immunological interest", US Public Health Services, NIH Bethesda, MD,
25 Publication No. 91), as applied to VHH domains from Camelids, as shown e.g. in Figure 2 of Riechmann and Muyldermans, J. Immunol. Methods 231, 25-38 (1999). According to this numbering,

- FR1 comprises the amino acid residues at positions 1-30,
- CDR1 comprises the amino acid residues at positions 31-35,
- 30 - FR2 comprises the amino acids at positions 36-49,
- CDR2 comprises the amino acid residues at positions 50-65,

- FR3 comprises the amino acid residues at positions 66-94,
- CDR3 comprises the amino acid residues at positions 95-102, and
- FR4 comprises the amino acid residues at positions 103-113.

However, it should be noted that - as is well known in the art for V_H domains and for VHH domains - the total number of amino acid residues in each of the CDRs may vary and may not correspond to the total number of amino acid residues indicated by the Kabat numbering (that is, one or more positions according to the Kabat numbering may not be occupied in the actual sequence, or the actual sequence may contain more amino acid residues than the number allowed for by the Kabat numbering). This means that, generally, the numbering according to Kabat may or may not correspond to the actual numbering of the amino acid residues in the actual sequence.

Alternative methods for numbering the amino acid residues of V_H domains, which methods can also be applied in an analogous manner to VHH domains, are known in the art. However, in the present description, claims and figures, the numbering according to Kabat and applied to VHH domains as described above will be followed, unless indicated otherwise.

The total number of amino acid residues in a VHH domain will usually be in the range of from 110 to 120, often between 112 and 115. It should however be noted that smaller and longer sequences may also be suitable for the purposes described herein.

Further structural characteristics and functional properties of VHH domains and polypeptides containing the same can be summarized as follows:

VHH domains (which have been "designed" by nature to functionally bind to an antigen without the presence of, and without any interaction with, a light chain variable domain) can function as a single, relatively small, functional antigen-binding structural unit, domain or polypeptide. This distinguishes the VHH

domains from the VH and VL domains of conventional 4-chain antibodies, which by themselves are generally not suited for practical application as single antigen-binding proteins or immunoglobulin single variable domains, but need to be combined in some form or another to provide a functional antigen-binding unit (as in for example conventional antibody fragments such as Fab fragments; in scFv's, which consist of a VH domain covalently linked to a VL domain).

Because of these unique properties, the use of VHH domains - either alone or as part of a larger polypeptide - offers a number of significant advantages over the use of conventional VH and VL domains, scFv's or conventional antibody fragments (such as Fab- or F(ab')₂-fragments):

- only a single domain is required to bind an antigen with high affinity and with high selectivity, so that there is no need to have two separate domains present, nor to assure that these two domains are present in the right spacial conformation and configuration (i.e. through the use of especially designed linkers, as with scFv's);
- VHH domains can be expressed from a single gene and require no post-translational folding or modifications;
- VHH domains can easily be engineered into multivalent and multispecific formats (as further discussed herein);
- VHH domains are highly soluble and do not have a tendency to aggregate (as with the mouse-derived antigen-binding domains described by Ward *et al.*, Nature 341: 544-546 (1989));
- VHH domains are highly stable to heat, pH, proteases and other denaturing agents or conditions and, thus, may be prepared, stored or transported without the use of refrigeration equipments, conveying a cost, time and environmental savings;
- VHH domains are easy and relatively cheap to prepare, even on a scale required for production. For example, VHH domains and polypeptides containing the same can be produced using microbial fermentation (e.g. as further described

below) and do not require the use of mammalian expression systems, as with for example conventional antibody fragments;

- VHH domains are relatively small (approximately 15 kDa, or 10 times smaller than a conventional IgG) compared to conventional 4-chain antibodies and

antigen-binding fragments thereof, and therefore

- show high(er) penetration into tissues and

- can be administered in higher doses

than such conventional 4-chain antibodies and antigen-binding fragments thereof;

- VHH domains can show so-called cavity-binding properties (inter alia due to

their extended CDR3 loop, compared to conventional VH domains) and can

therefore also access targets and epitopes not accessible to conventional 4-chain antibodies and antigen-binding fragments thereof.

Methods of obtaining VHH domains binding to a specific antigen or epitope have

been described earlier, e.g. in WO2006/040153 and WO2006/122786. As also

described therein in detail, VHH domains derived from camelids can be

"humanized" by replacing one or more amino acid residues in the amino acid sequence of the original VHH sequence by one or more of the amino acid

residues that occur at the corresponding position(s) in a VH domain from a

conventional 4-chain antibody from a human being. A humanized VHH domain

can contain one or more fully human framework region sequences, and, in an

even more specific embodiment, can contain human framework region

sequences derived from DP-29, DP-47, DP-51, or parts thereof, optionally

combined with JH sequences, such as JH5.

f2) "*Domain antibodies*", also known as "Dab"s, "Domain Antibodies", and

"dAbs" (the terms "Domain Antibodies" and "dAbs" being used as trademarks by

the GlaxoSmithKline group of companies) have been described in e.g. Ward,

E.S., *et al.*: "Binding activities of a repertoire of single immunoglobulin variable

domains secreted from *Escherichia coli*"; Nature 341: 544-546 (1989); Holt, L.J.

et al.: "Domain antibodies: proteins for therapy"; TRENDS in Biotechnology 21(11): 484-490 (2003); and WO2003/002609.

Domain antibodies essentially correspond to the VH or VL domains of non-camelid mammals, in particular human 4-chain antibodies. In order to bind an epitope as a single antigen binding domain, i.e. without being paired with a VL or VH domain, respectively, specific selection for such antigen binding properties is required, e.g. by using libraries of human single VH or VL domain sequences.

Domain antibodies have, like VHHs, a molecular weight of approximately 13 to approximately 16 kDa and, if derived from fully human sequences, do not require humanization for e.g. therapeutical use in humans. As in the case of VHH domains, they are well expressed also in prokaryotic expression systems, providing a significant reduction in overall manufacturing cost.

Domain antibodies, as well as VHH domains, can be subjected to affinity maturation by introducing one or more alterations in the amino acid sequence of one or more CDRs, which alterations result in an improved affinity of the resulting immunoglobulin single variable domain for its respective antigen, as compared to the respective parent molecule. Affinity-matured immunoglobulin single variable domain molecules of the invention may be prepared by methods known in the art, for example, as described by Marks *et al.*, 1992, Biotechnology 10:779-783, or Barbas, *et al.*, 1994, Proc. Nat. Acad. Sci, USA 91: 3809-3813.; Shier *et al.*, 1995, Gene 169:147-155; Yelton *et al.*, 1995, Immunol. 155: 1994-2004; Jackson *et al.*, 1995, J. Immunol. 154(7):3310-9; and Hawkins *et al.*, 1992, J. Mol. Biol. 226(3): 889 896; KS Johnson and RE Hawkins, "Affinity maturation of antibodies using phage display", Oxford University Press 1996.

f3) Furthermore, it will also be clear to the skilled person that it is possible to "graft" one or more of the CDRs mentioned above onto other "scaffolds", including but not limited to human scaffolds or non-immunoglobulin scaffolds. Suitable scaffolds and techniques for such CDR grafting are known in the art.

g) The terms "*epitope*" and "*antigenic determinant*", which can be used interchangeably, refer to the part of a macromolecule, such as a polypeptide, that is recognized by antigen-binding molecules, such as conventional antibodies or the polypeptides of the invention, and more particularly by the antigen-binding site of said molecules. Epitopes define the minimum binding site for an immunoglobulin, and thus represent the target of specificity of an immunoglobulin.

The part of an antigen-binding molecule (such as a conventional antibody or a polypeptide of the invention) that recognizes the epitope is called a *paratope*.

h) The term "*biparatopic*" (antigen-)binding molecule or "*biparatopic*" polypeptide as used herein shall mean a polypeptide comprising a first immunoglobulin single variable domain and a second immunoglobulin single variable domain as herein defined, wherein these two variable domains are capable of binding to two different epitopes of one antigen, which epitopes are not normally bound at the same time by one monospecific immunoglobulin, such as e.g. a conventional antibody or one immunoglobulin single variable domain. The biparatopic polypeptides according to the invention are composed of variable domains which have different epitope specificities, and do not contain mutually complementary variable domain pairs which bind to the same epitope. They do therefore not compete with each other for binding to LRP5 or LRP6.

i) A polypeptide (such as an immunoglobulin, an antibody, an immunoglobulin single variable domain, a polypeptide of the invention, or generally an antigen binding molecule or a fragment thereof) that can "*bind*", "*bind to*", "*specifically bind*", or "*specifically bind to*", that "*has affinity for*" and/or that "*has specificity for*" a certain epitope, antigen or protein (or for at least one part, fragment or epitope thereof) is said to be "*against*" or "*directed against*" said epitope, antigen or protein or is a "*binding*" molecule with respect to such epitope, antigen or protein.

k) Generally, the term "*specificity*" refers to the number of different types of antigens or epitopes to which a particular antigen-binding molecule or antigen-binding protein (such as an immunoglobulin, an antibody, an immunoglobulin single variable domain, or a polypeptide of the invention) can bind. The specificity of an antigen-binding protein can be determined based on its affinity and/or avidity. The affinity, represented by the equilibrium constant for the dissociation of an antigen with an antigen-binding protein (K_D), is a measure for the binding strength between an epitope and an antigen-binding site on the antigen-binding protein: the lesser the value of the K_D , the stronger the binding strength between an epitope and the antigen-binding molecule (alternatively, the affinity can also be expressed as the affinity constant (K_A), which is $1/K_D$). As will be clear to the skilled person (for example on the basis of the further disclosure herein), affinity can be determined in a manner known *per se*, depending on the specific antigen of interest. Avidity is the measure of the strength of binding between an antigen-binding molecule (such as an immunoglobulin, an antibody, an immunoglobulin single variable domain, or a polypeptide of the invention) and the pertinent antigen. Avidity is related to both the affinity between an epitope and its antigen binding site on the antigen-binding molecule and the number of pertinent binding sites present on the antigen-binding molecule.

Typically, antigen-binding proteins (such as the polypeptides of the invention) will bind with a dissociation constant (K_D) of $10E-5$ to $10E-14$ moles/liter (M) or less, and preferably $10E-7$ to $10E-14$ moles/liter (M) or less, more preferably $10E-8$ to $10E-14$ moles/liter, and even more preferably $10E-11$ to $10E-13$ (as measured e.g. in a Kinexa assay; known in the art), and/or with an association constant (K_A) of at least $10E7$ ME^{-1} , preferably at least $10E8$ ME^{-1} , more preferably at least $10E9$ ME^{-1} , such as at least $10E11$ ME^{-1} . Any K_D value greater than $10E-4$ M is generally considered to indicate non-specific binding. Preferably, a polypeptide of the invention will bind to the desired antigen with a K_D less than 500 nM, preferably less than 200 nM, more preferably less than 10 nM, such as less than 500 pM. Specific binding of an antigen-binding protein to an antigen or epitope

can be determined in any suitable manner known *per se*, including, for example, the assays described herein, Scatchard analysis and/or competitive binding assays, such as radioimmunoassays (RIA), enzyme immunoassays (EIA) and sandwich competition assays, and the different variants thereof known *per se* in the art.

l) The term "*cross-reactive*" in connection with binding molecules which are able to bind to LRP5 as well as to LRP6 ("*LRP5/LRP6 cross-reactive*") is intended to mean that such binding molecules can specifically bind to an epitope comprised in the LRP5 molecule, and can, alternatively, also specifically bind to an epitope comprised in the LRP6 molecule. Usually, such cross-reactivity may arise in case that the epitopes of the different proteins bound by such binding molecule have a similar structure and/or sequence, e.g. represent conserved epitopes, e.g. are shared by proteins belonging to the same protein family (e.g. LRP5 and LRP6, belonging to the LRP protein family).

m) Amino acid residues will be indicated according to the standard three-letter or one-letter amino acid code, as generally known and agreed upon in the art.

When comparing two amino acid sequences, the term "*amino acid difference*" refers to insertions, deletions or substitutions of the indicated number of amino acid residues at a position of the reference sequence, compared to a second sequence. In case of substitution(s), such substitution(s) will preferably be conservative amino acid substitution(s), which means that an amino acid residue is replaced with another amino acid residue of similar chemical structure and which has little or essentially no influence on the function, activity or other biological properties of the polypeptide. Such conservative amino acid substitutions are well known in the art, for example from WO 98/49185, wherein conservative amino acid substitutions preferably are substitutions in which one amino acid within the following groups (i) - (v) is substituted by another amino acid residue within the same group: (i) small aliphatic, nonpolar or slightly polar residues: Ala, Ser, Thr, Pro and Gly; (ii) polar, negatively charged residues and

their (uncharged) amides: Asp, Asn, Glu and Gln; (iii) polar, positively charged residues: His, Arg and Lys; (iv) large aliphatic, nonpolar residues: Met, Leu, Ile, Val and Cys; and (v) aromatic residues: Phe, Tyr and Trp. Particularly preferred conservative amino acid substitutions are as follows:

- 5 Ala into Gly or into Ser;
 Arg into Lys;
 Asn into Gln or into His;
 Asp into Glu;
 Cys into Ser;
- 10 Gln into Asn;
 Glu into Asp;
 Gly into Ala or into Pro;
 His into Asn or into Gln;
 Ile into Leu or into Val;
- 15 Leu into Ile or into Val;
 Lys into Arg, into Gln or into Glu;
 Met into Leu, into Tyr or into Ile;
 Phe into Met, into Leu or into Tyr;
 Ser into Thr;
- 20 Thr into Ser;
 Trp into Tyr;
 Tyr into Trp or into Phe;
 Val into Ile or into Leu.

- 25 n) A nucleic acid or polypeptide molecule is considered to be "*(in) essentially isolated (form)*" - for example, when compared to its native biological source and/or the reaction medium or cultivation medium from which it has been obtained - when it has been separated from at least one other component with which it is usually associated in said source or medium, such as another nucleic
- 30 acid, another protein/polypeptide, another biological component or macromolecule or at least one contaminant, impurity or minor component. In

particular, a nucleic acid or polypeptide molecule is considered "essentially isolated" when it has been purified at least 2-fold, in particular at least 10- fold, more in particular at least 100-fold, and up to 1000-fold or more. A nucleic acid or polypeptide molecule that is "in essentially isolated form" is preferably essentially
5 homogeneous, as determined using a suitable technique, such as a suitable chromatographical technique, such as polyacrylamide-gel electrophoresis;

o) "*Sequence identity*" between e.g. two immunoglobulin single variable domain sequences indicates the percentage of amino acids that are identical between
10 these two sequences. It may be calculated or determined as described in paragraph f) on pages 49 and 50 of WO2008/020079. "*Sequence similarity*" indicates the percentage of amino acids that either are identical or that represent conservative amino acid substitutions.

15 *Target specificity*

The polypeptides of the invention have specificity for LRP5 as well as LRP6, in that they comprise immunoglobulin single variable domains specifically binding to epitopes included in both of these molecules (LRP5/LRP6 cross-reactive binding
20 molecules).

The molecules of the invention shall bind to the human forms of LRP5 and LRP6, and preferably also to counterparts in other species relevant for drug development, i.e. cynomolgus and mouse LRP5 and LRP6.

25 *Polypeptides of the invention*

In its broadest sense, the invention provides novel pharmacologically active agents for the treatment of cancer diseases. The agents according to the invention belong to a novel class of binding molecules, namely LRP5/LRP6
30 cross-reactive biparatopic polypeptides, comprising two or more immunoglobulin single variable domains binding to LRP5 and/or LRP6 at different epitopes. The

terms “cross-reactive” and “biparatopic” are explained above, so that LRP5/LRP6 cross-reactive biparatopic molecules can be defined as molecules being able to bind to LRP5 at two different epitopes comprised in the LRP5 protein, and also being able to bind to LRP6 at the corresponding two epitopes comprised in the LRP6 protein.

More specifically, the polypeptides of the invention include:

- a first immunoglobulin single variable domain which is able to specifically bind to LRP5 as well as to LRP6 (LRP5/LRP6 cross-reactive) via an epitope / in a manner that results in inhibition of the Wnt1 signaling pathway, so that Wnt1-driven target gene transcription is inhibited, and
- a second immunoglobulin single variable domain which is able to specifically bind to LRP5 as well as to LRP6 (LRP5/LRP6 cross-reactive) via an epitope / in a manner that results in inhibition of the Wnt3a signaling pathway, so that Wnt3a-driven target gene transcription is inhibited.

Due to the two immunoglobulin single variable domains present in such polypeptide, wherein the two domains are binding to different epitopes (Wnt1 / Wnt3a signaling related), these molecules are biparatopic binding molecules. This biparatopic binding mode is schematically shown in Figure 1.

In this context, it should be noted that it is assumed that the polypeptides of the invention can bind to one single LRP5 or LRP6 molecule via both of its LRP5/LRP6 binding domains, as shown in Fig. 1 (intramolecular binding mode). However, other binding modes may occur as well.

Finally, it is assumed that the polypeptides of the invention are able to compete with DKK1 – a natural ligand of LRP5 and LRP6, and interfering with Wnt1 and Wnt3a signaling – for binding to LRP5 and LRP6, thereby inhibiting the Wnt1 as well as the Wnt3a signaling pathway. However, also this theory should not be understood as limiting the scope of the invention.

More specifically, the polypeptides according to the invention specifically bind to LRP5 or LRP6, wherein such polypeptides comprise a first immunoglobulin single variable domain (a) selected from the group of immunoglobulin single variable domains (i) to (iii) defined by having the following CDR sequences:

(i):

CDR1: TYTVG (= SEQ ID NO:1)
 CDR2: AIRRRGSSTYYADSVKG (= SEQ ID NO:2)
 CDR3: DTRTVALLQYRYDY (= SEQ ID NO:3)
 [= CDRs of the Wnt1-333E06mod domain]

(ii):

CDR1: SYAMG (= SEQ ID NO:4)
 CDR2: AIRRSGRRTYYADSVKG (= SEQ ID NO:5)
 CDR3: ARRVRSSSTRYNTGTWWWEY (= SEQ ID NO:6)
 [= CDRs of the Wnt1-333G06 domain]

(iii):

CDR1: RYTMG (= SEQ ID NO:7)
 CDR2: AIVRSGGSTYYADSVKG (= SEQ ID NO:8)
 CDR3: DRRGRGENYILLYSSGRYEY (= SEQ ID NO:9)
 [= CDRs of the Wnt1-332D03mod domain],

and a second immunoglobulin single variable domain (b) selected from the group of immunoglobulin single variable domains (iv) and (v) defined by having the following CDR sequences:

(iv):

CDR1: SYAMG (= SEQ ID NO:10)
 CDR2: AISWSGGSTYYADSVKG (= SEQ ID NO:11)
 CDR3: SPIPYGSLLRRRNNDY (= SEQ ID NO:12)
 [= CDRs of the Wnt3a-093A01 domain]

(v):

CDR1: SYAMG (= SEQ ID NO:13)
 CDR2: AISWRSGSTYYADSVKG (= SEQ ID NO:14)
 CDR3: DPRGYGVAYVSAYYEY (= SEQ ID NO:15)

[= CDRs of the Wnt3a-367B10 domain].

The use of the terms “first” and “second” with respect to such immunoglobulin single variable domains is solely intended to indicate that these domains are different domains, as they will include different CDR sequences and will bind to different epitopes. These terms shall, however, not be understood to refer to the exact order or sequence of the domains within such polypeptide chain. In other words, the above immunoglobulin single variable domains (a) and (b) may either be arranged in the order (a) – (b) or in the order (b) – (a) within such polypeptide of the invention.

The term “specifically bind to LRP5 or LRP6” is intended to mean that the immunoglobulin single variable domains (a) and (b) are cross-reactive with respect to LRP5 and LRP6. Of course, the binding properties of such molecules are determined by their (CDR) sequences, so that the feature “specifically binding to LRP5 or LRP6” set out above and in the claims is only intended to illustrate the utility of the invention, and not to limit the scope of this invention.

Immunoglobulin single variable domains typically essentially consist of four framework regions (FR1 to FR4, respectively) and three complementarity determining regions (CDR1 to CDR3, respectively). To be located within one polypeptide, or polypeptide chain, said first and said second immunoglobulin single variable domains need to be covalently linked, either directly or by a linker peptide.

Thus, the general structure of the molecules of the invention can also be depicted as follows:

FR(a)1 - CDR(a)1 - FR(a)2 - CDR(a)2 - FR(a)3 - CDR(a)3 - FR(a)4 – [linker peptide] - FR(b)1 - CDR(b)1 - FR(b)2 - CDR(b)2 - FR(b)3 - CDR(b)3 - FR(b)4
wherein

FR(a) denotes a framework region of the first immunoglobulin single variable domain,

FR(b) denotes a framework region of the second immunoglobulin single variable domain,

5 CDR(a) denotes a CDR of the first immunoglobulin single variable domain,
 CDR(b) denotes a CDR of the second immunoglobulin single variable domain,
 [linker peptide] denotes a linker peptide that may optionally be present,
 wherein the CDRs are having the sequences as set out above.

10 Again, it shall be understood that (a) and (b) can be exchanged, i.e. that molecules having the general structure
 FR(b)1 - CDR(b)1 - FR(b)2 - CDR(b)2 - FR(b)3 - CDR(b)3 - FR(b)4 – [linker peptide] - FR(a)1 - CDR(a)1 - FR(a)2 - CDR(a)2 - FR(a)3 - CDR(a)3 - FR(a)4 shall also be encompassed by the present invention.

15 The linker peptide optionally comprises or consists of a third domain, such as e.g. an albumin binding immunoglobulin single variable domain, such as the Alb11 domain, comprising the following CDRs:

CDR(Alb11)1: SFGMS (= SEQ ID NO:16)
 20 CDR(Alb11)2: SISGSGSDTLYADSVKG (= SEQ ID NO:17)
 CDR(Alb11)3: GGSLSR (= SEQ ID NO:18)

This results in a group of polypeptides of the invention having the following general structure:

25 FR(a)1 - CDR(a)1 - FR(a)2 - CDR(a)2 - FR(a)3 - CDR(a)3 - FR(a)4 – [linker peptide] - FR(Alb11)1 - CDR(Alb11)1 - FR(Alb11)2 - CDR(Alb11)2 - FR(Alb11)3 - CDR(Alb11)3 - FR(Alb11)4 - [linker peptide] - FR(b)1 - CDR(b)1 - FR(b)2 - CDR(b)2 - FR(b)3 - CDR(b)3 - FR(b)4.

Again, the order of the three immunoglobulin single variable domains (a), (b), and Alb11 is not fixed but polypeptides in which the above domains are arranged in the order:

(b) – Alb11 – (a)

5 shall be encompassed as well.

Furthermore, polypeptides having the Alb11 domain at the N- or C-terminal end of the polypeptide (e.g. Alb11 – (a) – (b), Alb11 – (b) – (a), (a) – (b) – Alb11, or (b) – (a) – Alb11) shall also be encompassed by the invention.

10 In three preferred embodiments, the polypeptides of the invention include immunoglobulin single variable domains defined as follows:

First preferred embodiment: Polypeptides comprising a first immunoglobulin single variable domain having the following CDR sequences:

15 CDR1: TYTVG (= SEQ ID NO:1)

CDR2: AIRRRGSSTYYADSVKG (= SEQ ID NO:2)

CDR3: DTRTVALLQYRYDY (= SEQ ID NO:3)

and a second immunoglobulin single variable domain having the following CDR sequences:

20 CDR1: SYAMG (= SEQ ID NO:10)

CDR2: AISWSGGSTYYADSVKG (= SEQ ID NO:11)

CDR3: SPIPYGSLLRRRNNDY (= SEQ ID NO:12).

Second preferred embodiment: Polypeptides comprising a first immunoglobulin single variable domain having the following CDR sequences:

25 CDR1: SYAMG (= SEQ ID NO:4)

CDR2: AIRRSGRRTYYADSVKG (= SEQ ID NO:5)

CDR3: ARRVRSSSTRYNTGTWWWEY (= SEQ ID NO:6)

and a second immunoglobulin single variable domain having the following CDR sequences:

30 CDR1: SYAMG (= SEQ ID NO:13)

CDR2: AISWRSGSTYYADSVKG (= SEQ ID NO:14)

CDR3: DPRGYGVAYVSAYYEY (= SEQ ID NO:15).

Third preferred embodiment: Polypeptides comprising a first immunoglobulin
5 single variable domain having the following CDR sequences:

CDR1: RYTMG (= SEQ ID NO:7)

CDR2: AIVRSGGSTYYADSVKG (= SEQ ID NO:8)

CDR3: DRRGRGENYILLYSSGRYEY (= SEQ ID NO:9)

and a second immunoglobulin single variable domain having the following CDR
10 sequences:

CDR1: SYAMG (= SEQ ID NO:13)

CDR2: AISWRSGSTYYADSVKG (= SEQ ID NO:14)

CDR3: DPRGYGVAYVSAYYEY (= SEQ ID NO:15).

15 Of course, the variants as set out above – i.e. optionally including linker peptides
and/or further domains, esp. including an Alb11 domain, different orders of the
immunoglobulin single variable domains – shall apply to these three preferred
embodiments as well.

20 In a specifically preferred embodiment, the albumin binding immunoglobulin
single variable domain is located between the two LRP5/LRP6 binding
immunoglobulin single variable domains. Thus, three specifically preferred
embodiments can be envisaged as follows:

25 First specifically preferred embodiment: Polypeptides comprising a first
(LRP5/LRP6 binding) immunoglobulin single variable domain having the following
CDR sequences:

CDR1: TYTVG (= SEQ ID NO:1)

CDR2: AIRRRGSSTYYADSVKG (= SEQ ID NO:2)

30 CDR3: DTRTVALLQYRYDY (= SEQ ID NO:3)

an albumin binding immunoglobulin single variable domain having the following CDR sequences:

CDR1: SFGMS (= SEQ ID NO:16)

CDR2: SISGSGSDTLYADSVKG (= SEQ ID NO:17)

5 CDR3: GGSLSR (= SEQ ID NO:18);

and a second (LRP5/LRP6 binding) immunoglobulin single variable domain having the following CDR sequences:

CDR1: SYAMG (= SEQ ID NO:10)

CDR2: AISWSGGSTYYADSVKG (= SEQ ID NO:11)

10 CDR3: SPIPYGSLLRRRNNDY (= SEQ ID NO:12);

either in this order, or the order of the above domains being changed.

Second specifically preferred embodiment: Polypeptides comprising a first (LRP5/LRP6 binding) immunoglobulin single variable domain having the following

15 CDR sequences:

CDR1: SYAMG (= SEQ ID NO:4)

CDR2: AIRRSGRRTYYADSVKG (= SEQ ID NO:5)

CDR3: ARRVRSSSTRYNTGTWWWEY (= SEQ ID NO:6);

an albumin binding immunoglobulin single variable domain having the following

20 CDR sequences:

CDR1: SFGMS (= SEQ ID NO:16)

CDR2: SISGSGSDTLYADSVKG (= SEQ ID NO:17)

CDR3: GGSLSR (= SEQ ID NO:18);

and a second (LRP5/LRP6 binding) immunoglobulin single variable domain

25 having the following CDR sequences:

CDR1: SYAMG (= SEQ ID NO:13)

CDR2: AISWRSGSTYYADSVKG (= SEQ ID NO:14)

CDR3: DPRGYGVAYVSAYYEY (= SEQ ID NO:15);

either in this order, or the order of the above domains being changed.

30

Third specifically preferred embodiment: Polypeptides comprising first (LRP5/LRP6 binding) immunoglobulin single variable domain having the following CDR sequences:

CDR1: RYTMG (= SEQ ID NO:7)

CDR2: AIVRSGGSTYYADSVKG (= SEQ ID NO:8)

CDR3: DRRGRGENYILLYSSGRYEY (= SEQ ID NO:9);

an albumin binding immunoglobulin single variable domain having the following CDR sequences:

CDR1: SFGMS (= SEQ ID NO:16)

CDR2: SISGSGSDTLYADSVKG (= SEQ ID NO:17)

CDR3: GGSLSR (= SEQ ID NO:18);

and a second (LRP5/LRP6 binding) immunoglobulin single variable domain having the following CDR sequences:

CDR1: SYAMG (= SEQ ID NO:13)

CDR2: AISWRSGSTYYADSVKG (= SEQ ID NO:14)

CDR3: DPRGYGVAYVSAYYEY (= SEQ ID NO:15);

either in this order, or the order of the above domains being changed.

The CDR sequences mentioned above are summarized in Tables IA, IB, and IC:

TABLE IA: CDR sequences of immunoglobulin single variable domains interfering with Wnt1 signaling:

	Wnt1-333E06mod	Wnt1-333G06	Wnt1-332D03mod
CDR1	TYTVG (SEQ ID NO:1)	SYAMG (SEQ ID NO:4)	RYTMG (SEQ ID NO:7)
CDR2	AIRRRGSSTYYADS VKG (SEQ ID NO:2)	AIRRSGRRRTYYADSVK G (SEQ ID NO:5)	AIVRSGGSTYYADS VKG (SEQ ID NO:8)
CDR3	DTRTVALLQYRYDY (SEQ ID NO:3)	ARRVRSSTRYNTGTW WWEY (SEQ ID NO:6)	DRRGRGENYILLYS SGRYEY (SEQ ID NO:9)

TABLE IB: CDR sequences of immunoglobulin single variable domains interfering with Wnt3a signaling:

	Wnt3a-093A01	Wnt3a-367B10
CDR1	SYAMG (SEQ ID NO:10)	SYAMG (SEQ ID NO:13)
CDR2	AISWSGGSTYYADSVKG (SEQ ID NO:11)	AISWRSGSTYYADSVKG (SEQ ID NO:14)
CDR3	SPIPYGSLLRRRNNDY (SEQ ID NO:12)	DPRGYGVAYVSAYYEY (SEQ ID NO:15)

5 **TABLE IC:** CDR sequences of immunoglobulin single variable domain binding to serum albumin (Alb11 domain):

	Alb11 domain
CDR1	SFGMS (SEQ ID NO:16)
CDR2	SISGSGSDTLYADSVKG (SEQ ID NO:17)
CDR3	GGSLSR (SEQ ID NO:18)

In addition to the CDR sequences as set out above, the immunoglobulin single variable domains comprised in the polypeptides of the invention include
 10 immunoglobulin framework region (FR) sequences. These sequences are preferably not immunogenic in humans, and are therefore preferably human or humanized FR sequences. Suitable human or humanized FR sequences are known in the art. Specifically preferred FR sequences can be taken from the embodiments shown below, disclosing the complete immunoglobulin single
 15 variable domains and thereby CDR sequences as well as FR sequences.

According to a more specific embodiment, the polypeptides of the invention include immunoglobulin single variable domains which are VHH domains, and preferably humanized VHH domains.

According to an even more specific embodiment, the polypeptides of the invention include a first immunoglobulin single variable domain (a) selected from the group consisting of immunoglobulin single variable domains (i) to (iii) having the following sequences:

(i)

AVQLVESGGGLVQPGGSLRLSCAASGRTFSTYTVGWFRQAPGKEREFVA
AIRRRGSSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAA
DTRTVALLQYRYDYWGQGTLVTVSS

[= Wnt1-333E06mod domain; = SEQ ID NO:19]

(ii)

AVQLVESGGGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAPGKEREFVA
AIRRSGRRTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAA
ARRVRSSTRYNTGTWWWEYWGQGTLVTVSS

[= Wnt1-333G06 domain; = SEQ ID NO:20], and

(iii)

AVQLVESGGGLVQPGGSLRLSCAASGLTFSRYTMGWFRQAPGKEREFVA
AIVRSGGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAA
DRRGRGENYILLYSSGRYEYWGQGTLVTVSS

[= Wnt1-332D03mod domain; = SEQ ID NO:21],

and a second immunoglobulin single variable domain (b) selected from the group consisting of immunoglobulin single variable domains (iv) and (v) having the following sequences:

(iv)

EVQLVESGGGLVQPGGSLRLSCAASGRTFSSYAMGWFRQAPGKEREFVA
AISWSGGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAA
SPIPYGSLRRRRNNYDYWGQGTLVTVSS

[= Wnt3a-093A01 domain; = SEQ ID NO:22], and

(v)

EVQLVESGGGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAPGKEREFVA
 AISWRSGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAA
 DPRGYGVAYVSAYYEWGQGT LTVSS

5 [= Wnt3a-367B10 domain; = SEQ ID NO:23].

Preferred embodiments are polypeptides comprising

- a first immunoglobulin single variable domain having the amino acid sequence as shown in SEQ ID NO:19 and a second immunoglobulin single variable domain
- 10 having the amino acid sequence as shown in SEQ ID NO:22; or
- a first immunoglobulin single variable domain having the amino acid sequence as shown in SEQ ID NO:20 and a second immunoglobulin single variable domain having the amino acid sequence as shown in SEQ ID NO:23; or
- a first immunoglobulin single variable domain having the amino acid sequence
- 15 as shown in SEQ ID NO:21 and a second immunoglobulin single variable domain having the amino acid sequence as shown in SEQ ID NO:23.

Thus, the above embodiments can schematically be presented as

isvd(a) – [linker peptide] – isvd(b),

- 20 wherein “isvd” denotes the respective immunoglobulin single variable domain, and wherein otherwise the same definitions and variants shall apply as set out above, esp. with respect to the presence of optional linker peptides and/or further domains, esp. an Alb11 domain, and with respect to different orders of the immunoglobulin single variable domains.

25

According to specific embodiments of the invention, the above polypeptides may additionally include a half-life extending moiety, wherein said half-life extending moiety is covalently linked to said polypeptide and is optionally selected from the group consisting of an albumin binding moiety, such as an albumin binding

30 peptide or an albumin binding immunoglobulin domain, preferably an albumin binding immunoglobulin single variable domain, more preferably the Alb11

domain, a transferrin binding moiety, such as an anti-transferrin immunoglobulin domain, a polyethylene glycol molecule, serum albumin, preferably human serum albumin, and a fragment of (human) serum albumin.

- 5 The sequence of the above-mentioned Alb11 immunoglobulin single variable domain is as follows:

EVQLVESGGGLVQPGNSLRSLCAASGFTFSSFGMSWVRQAPGKGLEW
VSSISGSGSDTLYADSVKGRFTISRDNAKTTLYLQMNSLRPEDTAVYYCTI
GGSLSRSSQGTLVTVSS

10 (= Alb11 domain; = SEQ ID NO:24)

Further examples of immunoglobulin single variable domains binding to human serum albumin are known in the art, and are described in further detail in e.g. international patent publications WO2006/122787 and WO2008/028977. Other
 15 peptides binding to human serum albumin are described e.g. in WO2008/068280, WO2009/127691, and WO2011/095545.

Thus, three preferred specific embodiments of the invention are as follows:

- 20 First preferred specific embodiment: Polypeptides comprising
- a first (LRP5/LRP6 binding) immunoglobulin single variable domain having the amino acid sequence as shown in SEQ ID NO:19;
 - an albumin binding immunoglobulin single variable domain having the amino acid sequence as shown in SEQ ID NO:24;
 - 25 - a second (LRP5/LRP6 binding) immunoglobulin single variable domain having the amino acid sequence as shown in SEQ ID NO:22;
- either in this order, or the order of the above three domains being changed.

Second preferred specific embodiment: Polypeptides comprising

- 30 - a first (LRP5/LRP6 binding) immunoglobulin single variable domain having the amino acid sequence as shown in SEQ ID NO:20;

- an albumin binding immunoglobulin single variable domain having the amino acid sequence as shown in SEQ ID NO:24;
 - a second (LRP5/LRP6 binding) immunoglobulin single variable domain having the amino acid sequence as shown in SEQ ID NO:23;
- 5 either in this order, or the order of the above three domains being changed.

Third preferred specific embodiment: Polypeptides comprising

- a first (LRP5/LRP6 binding) immunoglobulin single variable domain having the amino acid sequence as shown in SEQ ID NO:21;
 - 10 - an albumin binding immunoglobulin single variable domain having the amino acid sequence as shown in SEQ ID NO:24;
 - a second (LRP5/LRP6 binding) immunoglobulin single variable domain having the amino acid sequence as shown in SEQ ID NO:23;
- either in this order, or the order of the above three domains being changed.

15 In even more specifically preferred embodiments, the albumin binding immunoglobulin single variable domain is located between the two LRP5/LRP6 binding immunoglobulin single variable domains.

20 The sequences of the immunoglobulin single variable domains mentioned above are summarized in Tables IIA, IIB, and IIC:

TABLE IIA: Sequences of immunoglobulin single variable domains interfering with Wnt1 signaling:

VHH ID SEQ ID NO:	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4
Wnt1- 333E06 mod SEQ ID NO:19	AVQLVESG GGLVQPG GSLRLSCAA SGRTFS	TYTVG	WFRQAPG KEREFVA	AIRRRGS STYYADS VKG	RFTISRDNKNTVYL QMNSLRPEDTAVYY CAA	DTRTVA LLQYRY DY	WGQGTL VTVSS

Wnt1-333G06 SEQ ID NO:20	AVQLVESG GGLVQPG GSLRLSCAA SGGTFS	SYAMG	WFRQAPG KEREFVA	AIRRSGR RTYYADS VKG	RFTISRDNSKNTVYL QMNSLRPEDTAVYY CAA	ARRVRS STRYNT GTWW WEY	WGQGT LTVSS
Wnt1-332D03 mod SEQ ID NO:21	AVQLVESG GGLVQPG GSLRLSCAA SGLTFS	RYTMG	WFRQAPG KEREFVA	AIVRSGG STYYADS VKG	RFTISRDNSKNTVYL QMNSLRPEDTAVYY CAA	DRRGR GENYILL YSSGRY EY	WGQGT LTVSS

TABLE IIB: Sequences of immunoglobulin single variable domains interfering with Wnt3a signaling:

VHH ID SEQ ID NO:	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4
Wnt3a-093A01 SEQ ID NO:22	EVQLVESG GGLVQPG GSLRLSCAA SGRTFS	SYAMG	WFRQAPG KEREFVA	AISWSGG STYYADS VKG	RFTISRDNSKNTVYL QMNSLRPEDTAVYY CAA	SPIPYGS LLRRRN NYDY	WGQGT LTVSS
Wnt3a-367B10 SEQ ID NO:23	EVQLVESG GGLVQPG GSLRLSCAA SGGTFS	SYAMG	WFRQAPG KEREFVA	AISWRSG STYYADS VKG	RFTISRDNSKNTVYL QMNSLRPEDTAVYY CAA	DPRGY GVAYVS AYYEEY	WGQGT LTVSS

TABLE IIC: Sequence of immunoglobulin single variable domain binding to serum albumin (Alb11 domain):

VHH ID SEQ ID NO:	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4
Alb11 SEQ ID NO:24	EVQLVESG GGLVQPG NSLRLSCA ASGFTFS	SFGMS	WVRQAPG KGLEWVS	SISGSGSD TLYADSV KG	RFTISRDNAKTTLYL QMNSLRPEDTAVYY CTI	GGSLSR	SSQGT LTVSS

As set out above, the (at least two) immunoglobulin single variable domains present in a polypeptide of the invention can be linked to each other either directly, without use of a linker, or via a linker. The linker is preferably a linker peptide and will, according to the invention, be selected so as to allow binding of the at least two different immunoglobulin single variable domains to each of their target epitopes.

Suitable linkers will *inter alia* depend on the epitopes and, specifically, the distance between the epitopes on the target molecules to which the immunoglobulin single variable domains shall bind. This will be clear to the skilled person based on the disclosure herein, optionally after some limited degree of routine experimentation.

Thus, suitable linkers may comprise an amino acid sequence, e.g. having a length of 9 or more amino acids, preferably at least 17 amino acids, such as about 20 to 40 amino acids. The linker sequence may be a naturally occurring sequence or a non-naturally occurring sequence. If used for therapeutic purposes, the linker is preferably non-immunogenic in the subject to which the polypeptide of the invention is administered.

One useful group of linker sequences are linkers derived from the hinge region of heavy chain antibodies as described in WO1996/34103 and WO1994/04678. Other examples are poly-alanine linker sequences such as Ala-Ala-Ala.

Further preferred examples of linker sequences are Gly/Ser linkers of different length such as $(\text{gly}_x\text{ser}_y)_z$ linkers, including e.g. $(\text{gly}_4\text{ser})_3$, $(\text{gly}_4\text{ser})_5$, $(\text{gly}_4\text{ser})_7$, $(\text{gly}_3\text{ser})_3$, $(\text{gly}_3\text{ser})_5$, $(\text{gly}_3\text{ser})_7$, $(\text{gly}_3\text{ser}_2)_3$, $(\text{gly}_3\text{ser}_2)_5$, and $(\text{gly}_3\text{ser}_2)_7$.

Alternatively, or in addition, to a polypeptide linker, the at least two immunoglobulin single variable domains present in a polypeptide of the invention may be linked to each other via another moiety, such as another polypeptide

which, in a preferred but non-limiting embodiment, may be a further immunoglobulin single variable domain as already described above. Such moiety may either be essentially inactive or may have a biological effect such as improving the desired properties of the polypeptide or may confer one or more additional desired properties to the polypeptide. As already set out above, a preferred additional polypeptide domain will increase the half-life of the polypeptide, such as a (human) serum albumin binding domain, such as the Alb11 domain.

Thus, according to a further embodiment, the invention specifically includes polypeptides comprising any of the following sequences, wherein the exact amino acid sequences can be taken from Table III below:

SEQ ID NO:25 (= sequence of polypeptide F013500575),

SEQ ID NO:26 (= sequence of polypeptide F013500571),

and

SEQ ID NO:27 (= sequence of polypeptide F013500720).

According to an even more specific embodiment, the polypeptides of the invention are selected from the following group of molecules:

Polypeptide F013500575, having the sequence SEQ ID NO:25,

Polypeptide F013500571, having the sequence SEQ ID NO:26, and

Polypeptide F013500720, having the sequence SEQ ID NO:27.

Table III: Sequences of three specific embodiments of polypeptides of the invention

ID SEQ ID NO:	Amino Acid Sequence (CDR sequences underlined)
F013500575 SEQ ID NO:25	AVQLVESGGGLVQPGGSLRLSCAASGRTFSTYTVGWFRQAPGKEREFVAAIRRRGSSTYYA DSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAADTRTVALLQYRYDYWGQGLTVTV SSGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLVESGGGLVQP GNSLRL SCAASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYADSVKGRFTISRDN AKTTLYL QMNSLRPEDTAVYYCTIGGSLRSSQGLTVTVSSGGGSGGGGSGGGGSGGGGSGGGGS GGGGSGGGGSEVQLVESGGGLVQPGGSLRLSCAASGRTFSSYAMGWFRQAPGKEREFVA AISWSGGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAASPIPYGSLRRRN NYDYWGQGLTVTVSSA
F013500571 SEQ ID NO:26	AVQLVESGGGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAPGKEREFVAAIRRSGRRTYY ADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAAARRVRSSTRYNTGTWWWWEYW GQGLTVTVSSGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLVESGGGLV QPGNSLRLSCAASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYADSVKGRFTISRDN NAKTTLYLQMNSLRPEDTAVYYCTIGGSLRSSQGLTVTVSSGGGSGGGGSGGGGSGGGG GSGGGSGGGGSGGGGSEVQLVESGGGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAP GKEREFVAAISWRSGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAADPRG YGVAYVSAYEYWGQGLTVTVSSA
F013500720 SEQ ID NO:27	AVQLVESGGGLVQPGGSLRLSCAASGLTFSRYTMGWFRQAPGKEREFVAAIVRSGGSTYY ADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAADRRGRGENYILLYSSGRYEWGQ GTLTVTVSSGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLVESGGGLVQP GNSLRLSCAASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYADSVKGRFTISRDN A KTTLYLQMNSLRPEDTAVYYCTIGGSLRSSQGLTVTVSSGGGSGGGGSGGGGSGGGGS GGGGSGGGGSGGGGSEVQLVESGGGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAPGK EREFVAAISWRSGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAADPRGYG VAYVSAYEYWGQGLTVTVSSA

As explained before, unless indicated otherwise, the polypeptides of the invention
 5 may include further moieties and/or additional polypeptide domains, as long as
 their binding to LRP5/LRP6 will not be prevented by such additional moiety or
 domain.

The polypeptides of the invention can additionally contain modifications such as
 10 glycosyl residues or modified amino acid side chains, and they may be
 PEGylated in order to increase half-life and other properties of such molecule.
 Techniques and reagents useful for PEGylating biparatopic immunoglobulin
 single variable domain constructs may be taken e.g. from WO2011/107507.

The polypeptides of the invention may have a modified N-terminal sequence, e.g. a deletion of one or more of the N-terminal amino acids, or an exchange of e.g. the first, N-terminal amino acid (e.g. glutamate to alanine), to optimize the molecule for being expressed by using certain expression systems (such as
5 specific vectors or host cells), or for being expressed as inclusion bodies or in soluble form, or for being secreted into the medium or the periplasmic space or for being contained within the cell, or for yielding a more homogenous product. The polypeptides of the invention may have a modified C-terminal sequence, such as an additional alanine (as shown for the three embodiments shown in
10 Table III above), and/or further amino acid exchanges in the C-terminal part or at other defined positions within any of the framework regions, as explained e.g. in WO2012/175741, WO2011/075861, or WO2013/024059, in order to e.g. further enhance stability or reduce immunogenicity of such polypeptides.

15 Furthermore, half-life of the polypeptides of the invention may be enhanced by adding an albumin domain, i.e. by converting them into albumin fusion proteins. Examples of useful albumin moieties, and how to add them to binding molecules, are e.g. provided in WO2001/079271 and WO2003/059934.

20 Preferably, the polypeptides of the invention are having binding (EC_{50}) values, measured in a FACS binding assay as described in Example 7.1 below, in the range of 10^{-6} moles/liter or less, more preferably 10^{-9} moles/liter or less, and even more preferably in the range of from 10^{-10} to 10^{-13} moles/liter, or are having an IC_{50} value as measured in a combined Wnt1 and Wnt3a reporter assay as set
25 out in Example 7.3 below, of 10^{-9} moles/liter or below, and preferably in the range of from 5×10^{-10} moles/liter to 10^{-12} moles/liter.

The polypeptides of the invention allow a more efficient treatment of several cancer types, such as TNBC, CRC, and NSCLC. They are having improved *in vitro*
30 characteristics (i.e. higher efficacy of Wnt pathway inhibition), cf. e.g. Examples 7 and 8 below, and significant *in vivo* tumor growth inhibition

properties, leading to higher *in vivo* efficacy as compared to other LRP6 binding molecules described in the art, as shown in e.g. Examples 9 and 10 below.

In particular, as shown *in vivo* in a Wnt driven tumor model, the LRP5/LRP6 cross-reactive half-life extended biparatopic humanized VHH constructs could inhibit Wnt signaling and tumor growth *in vivo*, and even provided for substantial tumor shrinkage (i.e. tumor growth inhibition higher than 100%), which could not be achieved, under the same experimental settings, with a LRP6 binder known in the art. Tumor shrinkage (i.e. tumor regression) is of course the desired therapeutic effect (i.e. efficacy) for treatment of cancer patients. Furthermore, tumor regression, resulting in pathological complete response (pCR), is an acknowledged clinical endpoint, indicating significant improvement of progression free survival and overall survival.

In the same *in vivo* experiments, no significant body weight changes were observed (< 10%), and results from gastrointestinal histopathological analyses did not indicate any toxic effects of the above polypeptides of the invention. This is specifically surprising in view of the *in vivo* DKK1 expression studies discussed above (i.e. resulting in intestinal mucosa ulceration and body weight loss).

Thus, the polypeptides of the invention are in fact providing for novel therapeutic options for the treatment of cancer diseases, and especially even for use in high unmet medical need indications such as (triple negative) breast cancer.

Surprisingly, the inventors arrived at this solution not by going the conventional route, i.e. trying to develop inhibitors or binding molecules having a high specificity / selectivity for one given target, such as LRP6 (mentioned as a target for the treatment of "Wnt signaling mediated diseases" e.g. in WO2009/056634). In contrast thereto, the inventors developed molecules targeting two related proteins at the same time – LRP6 and LRP5 – and thereby achieved the above significantly improved *in vitro* and *in vivo* effects which could

not be expected from the prior art.

The underlying molecular mechanism for this superiority is not fully clear, but it may be speculated – without wishing to be bound to a specific theory – that such cross-reactive molecules may have additional, and thereby stronger, effects on the highly complicated signaling cascade of the Wnt signaling pathway.

The above advantageous effects will further be illustrated in the Examples below and by way of the comparative data included therein.

Moreover, the polypeptides of the invention are easy to manufacture and are more soluble, which means that they may be stored and/or administered in higher concentrations compared to conventional antibodies. They are stable at room temperature and have prolonged stability even at extremes of pH, so that they may be prepared, stored and/or transported without the use of refrigeration equipment, conveying cost, time and environmental savings. Due to the above and due to their low immunogenicity, they furthermore offer a variety of options regarding administration routes other than injection and infusion, as well as regarding administration regimens and uses of specific devices.

Nucleic acids, vectors, host cells

According to further aspects, the invention relates to nucleic acid molecules and expression vectors encoding the polypeptides of the invention, as well as to host cells expressing the same. These nucleic acids, vectors, and host cells are useful for manufacturing the polypeptides of the invention, and further aspects and embodiments thereof will be described further below in connection with an outline of methods of manufacturing the polypeptides of the invention.

Therapeutic use

Due to their biological properties, the polypeptides of the invention are suitable for treating diseases characterised by excessive or abnormal cell proliferation, such as cancer and idiopathic pulmonary fibrosis (IPF).

For example, the following cancers, tumors, and other proliferative diseases may be treated with polypeptides according to the invention, without being restricted thereto:

Cancers of the head and neck; Cancers of the lung, such as e.g. non-small cell lung cancer (NSCLC) and small cell lung cancer (SCLC); Neoplasms of the mediastinum, such as e.g. neurogenic tumors and mesenchymal tumors; Cancers of the gastrointestinal (GI) tract, such as e.g. cancers of the esophagus, stomach (gastric cancer), pancreas, liver and biliary tree (including e.g. hepatocellular carcinoma (HCC)), and the small and large intestine (including e.g. colorectal cancer); Cancers of the prostate; Cancers of the testis; Gynecologic cancers, such as e.g. cancers of the ovary; Cancers of the breast, such as e.g. mammary carcinoma, hormone receptor positive breast cancer, Her2 positive breast cancer, and triple negative breast cancer; Cancers of the endocrine system; Sarcomas of the soft tissues, such as e.g. fibrosarcoma, rhabdomyosarcoma, angiosarcoma, Kaposi's sarcoma; Sarcomas of the bone, such as e.g. myeloma, osteosarcoma, Ewing's tumor, fibrosarcoma, osteochondroma, osteoblastoma, and chondroblastoma; Mesotheliomas; Cancers of the skin, such as e.g. basal cell carcinoma, squamous cell carcinoma, Merkel's cell carcinoma, and melanoma; Neoplasms of the central nervous system and brain, such as e.g. astrocytoma, glioblastoma, gliomas neuroblastomas, and retinoblastomas; Lymphomas and leukemias such as e.g. B-cell non-Hodgkin lymphomas (NHL), T-cell non-Hodgkin lymphomas, chronic B-cell lymphocytic leukemia (B-CLL), chronic T-cell lymphocytic leukemia (T-CLL), Hodgkin's disease (HD), large granular lymphocyte leukemia (LGL),

chronic myelogenous leukemia (CML), acute myelogenous/myeloid leukemia (AML), acute lymphatic/lymphoblastic leukemia (ALL), multiple myeloma (MM), plasmacytoma, and myelodysplastic syndromes (MDS); and cancers of unknown primary site.

5

All cancers, tumors, neoplasms, etc., mentioned above which are characterized by their specific location/origin in the body are meant to include both the primary tumors and the metastatic tumors derived therefrom.

10 More specifically, the polypeptides of the invention are useful for the treatment of diseases, and esp. cancer diseases in which abnormal cell proliferation is caused by, or involves, abnormal (activated) Wnt signaling.

Thus, the polypeptides of the invention are specifically useful for the treatment of
15 solid tumors, and more specifically for the treatment of lung, liver, colon, brain, thyroid, pancreas, breast, ovary and prostate cancers, and even more specifically for the treatment of non-small cell lung cancer (NSCLC), triple-negative breast cancer (TNBC), and colo-rectal cancer (CRC). In particular, the polypeptides of the invention may be used to treat patients with locally advanced or metastatic
20 TNBC, patients with metastatic NSCLC or locally advanced or metastatic CRC, as single agent or in combination, to prolong progression free survival (PFS) and overall survival (OS). Furthermore, the polypeptides of the invention may be used as neoadjuvant treatment for breast cancer patients to achieve pathological complete response (pCR; defined as the absence of residual invasive and *in situ*
25 cancer by histo-pathological evaluation of the complete resected breast specimen and all sampled regional lymph nodes following completion of neoadjuvant systemic therapy).

The polypeptides of the invention may be used in therapeutic regimens in the
30 context of first line, second line, or any further line treatments.

The polypeptides of the invention may be used for the prevention, short-term or long-term treatment of the above-mentioned diseases, optionally also in combination with radiotherapy and/or surgery.

5 Likewise, the polypeptides of the invention are specifically useful for the treatment of other diseases caused by abnormal cell proliferation involving the Wnt signaling pathway, such as idiopathic pulmonary fibrosis (IPF) (Königshoff *et al.* "Functional Wnt signaling is increased in idiopathic pulmonary fibrosis". *PLoS One* 2008;3(5):e2142; Lam *et al.* "Wnt coreceptor Lrp5 is a driver of idiopathic
10 pulmonary fibrosis". *Am J Respir Crit Care Med.* 2014;190(2):185-95).

Furthermore, the polypeptides of the invention are specifically useful for the treatment of retinopathies, and esp. for the treatment of diabetic retinopathy, due to abnormal Wnt activation in the inner retina cells, causing the increase in
15 abnormal new retinal vessel formation leading to development and progression of diabetic retinopathy (Chen, Y., *et al.* "Activation of the Wnt pathway plays a pathogenic role in diabetic retinopathy in humans and animal models" *The Am J Pathol.* 2009; 175(6):2676-85., Gao *et al.* "Elevated LRP6 levels correlate with vascular endothelial growth factor in the vitreous of proliferative diabetic
20 retinopathy" *Mol Vis.* 2015;21:665-72).

Finally, as it could be shown that inhibition of Wnt1/Wnt3a signaling pathways can also have an effect on dendritic cells (DCs) and dendritic cell function, the polypeptides of the invention may also be useful in the treatment of immunologic
25 and infectious diseases, as well as for influencing the tumor microenvironment in the various cancer diseases already set out above. Tumors actively suppress antitumor immunity, and DCs play an important role in the cancer immunoescape mechanism. In particular, studies have shown that Wnt ligands in the tumor microenvironment can also initiate paracrine signaling within the immune cells
30 and regulate host antitumor immunity (Hong *et al.* "beta-catenin promotes

regulatory T-cell responses in tumors by inducing vitamin A metabolism in dendritic cells". *Cancer Res.* 2015;75(4):656-65).

Of course, the above also includes the use of the polypeptides of the invention in various methods of treating the above diseases by administering a therapeutically effective dose to a patient in need thereof, as well as the use of these polypeptides for the manufacture of medicaments for the treatment of such diseases, as well as pharmaceutical compositions including such polypeptides of the inventions, as well as the preparation and/or manufacture of medicaments including such polypeptides of the invention, and the like.

Combinations with other active substances

The polypeptides of the invention may be used on their own or in combination with other pharmacologically active substances, such as state-of-the-art or standard-of-care compounds, such as e.g. cytostatic or cytotoxic substances, cell proliferation inhibitors, anti-angiogenic substances, steroids, immune modulators / checkpoint inhibitors, and the like.

Cytostatic and/or cytotoxic active substances which may be administered in combination with the compounds according to the invention include, without being restricted thereto, hormones, hormone analogues and antihormones, aromatase inhibitors, LHRH agonists and antagonists, inhibitors of growth factors (growth factors such as for example platelet derived growth factor (PDGF), fibroblast growth factor (FGF), vascular endothelial growth factor (VEGF), epidermal growth factor (EGF), insuline-like growth factors (IGF), human epidermal growth factor (HER, e.g. HER2, HER3, HER4) and hepatocyte growth factor (HGF)), inhibitors are for example (anti-)growth factor antibodies, (anti-)growth factor receptor antibodies and tyrosine kinase inhibitors, such as for example cetuximab, gefitinib, afatinib, nintedanib, imatinib, lapatinib, bosutinib and trastuzumab; antimetabolites (e.g. antifolates such as methotrexate, raltitrexed, pyrimidine

analogues such as 5-fluorouracil (5-FU), capecitabine and gemcitabine, purine and adenosine analogues such as mercaptopurine, thioguanine, cladribine and pentostatin, cytarabine (ara C), fludarabine); antitumor antibiotics (e.g. anthracyclins); platinum derivatives (e.g. cisplatin, oxaliplatin, carboplatin);

5 alkylation agents (e.g. estramustin, meclorethamine, melphalan, chlorambucil, busulphan, dacarbazin, cyclophosphamide, ifosfamide, temozolomide, nitrosoureas such as for example carmustin and lomustin, thiotepa); antimitotic agents (e.g. Vinca alkaloids such as for example vinblastine, vindesin, vinorelbin and vincristine; and taxanes such as paclitaxel, docetaxel); angiogenesis

10 inhibitors, tubuline inhibitors; DNA synthesis inhibitors, PARP inhibitors, topoisomerase inhibitors (e.g. epipodophyllotoxins such as for example etoposide and etopophos, teniposide, amsacrin, topotecan, irinotecan, mitoxantrone), serine/threonine kinase inhibitors (e.g. PDK1 inhibitors, Raf inhibitors, A-Raf inhibitors, B-Raf inhibitors, C-Raf inhibitors, mTOR inhibitors, mTORC1/2

15 inhibitors, PI3K inhibitors, PI3K α inhibitors, dual mTOR/PI3K inhibitors, STK33 inhibitors, AKT inhibitors, PLK1 inhibitors (such as volasertib), inhibitors of CDKs, Aurora kinase inhibitors), tyrosine kinase inhibitors (e.g. PTK2/FAK inhibitors), protein protein interaction inhibitors, MEK inhibitors, ERK inhibitors, FLT3 inhibitors, BRD4 inhibitors, IGF-1R inhibitors, TRAILR2 agonists, Bcl-xL

20 inhibitors, Bcl-2 inhibitors, Bcl-2/Bcl-xL inhibitors, ErbB receptor inhibitors, BCR-ABL inhibitors, ABL inhibitors, Src inhibitors, rapamycin analogs (e.g. everolimus, temsirolimus, ridaforolimus, sirolimus), androgen synthesis inhibitors, androgen receptor inhibitors, DNMT inhibitors, HDAC inhibitors, ANG1/2 inhibitors, CYP17 inhibitors, radiopharmaceuticals, immunotherapeutic agents

25 such as immune checkpoint inhibitors (e.g. CTLA4, PD1, PD-L1, LAG3, and TIM3 binding molecules / immunoglobulins, such as ipilimumab, nivolumab, pembrolizumab), cancer vaccines such as traditional tumor vaccine (cell based vaccines, e.g. Sipuleucel-T for treatment of prostate cancer), personalized neoantigen vaccines, and oncolytic viruses, and various chemotherapeutic

30 agents such as amifostin, anagrelid, clodronat, filgrastin, interferon, interferon

alpha, leucovorin, rituximab, procarbazine, levamisole, mesna, mitotane, pamidronate and porfimer.

Particularly preferred are methods of treatment including the use of the polypeptides of the invention in combination with a drug selected from the group consisting of:

(i) anti-VEGF antibodies (bevacizumab and other anti-angiogenic substances) with or without chemotherapy combination (including doxorubicin / cyclophosphamide combination and/or capecitabine / docetaxel combination in neoadjuvant setting; taxane / platinum regimen for first and later line treatment) in breast cancer patients;

(ii) EGFR TKIs for EGFR mutant NSCLC or crizotinib for ALK translocated NSCLC with or without chemotherapy combination (platinum based cytotoxic combination therapy, including gemcitabine / cisplatin in first line treatment; docetaxel or pemetrexed in second line treatment in lung cancer patients;

(iii) anti-EGFR antibodies (cetuximab and panitumumab in *KRAS* wild-type tumors) with or without chemotherapy combination (including irinotecan), anti-VEGF antibody combination (bevacizumab and other anti-angiogenic substances) or regorafenib combination, e.g. for the treatment of CRC patients.

(iv) immunotherapeutic agents, including anti-PD-1 agents, such as pembrolizumab and nivolumab, anti-PD-L1 agents, anti-CTLA4 agents, anti-BTLA agents, anti-LAG3 agents, and anti-TIM3 agents, such as anti-PDL1 antibodies etc., e.g. for treatment of breast cancer, lung cancer and CRC patients

(v) chemotherapeutic agents, such as platinum-based antineoplastic agents, or in a combination with a FOLFOX chemotherapy regimen, including folinic acid, 5'-fluorouracil, and oxaliplatin, or in a combination with a FOLFOXIRI chemotherapy regimen, including folinic acid, 5'-fluorouracil, oxaliplatin, and irinotecan, e.g. for treatment of breast cancer or CRC patients.

When two or more substances or principles are to be used as part of a combined treatment regimen, they can be administered via the same route of administration

or via different routes of administration, at essentially the same time (i.e. simultaneously, concurrently) or at different times (e.g. sequentially, successively, alternately, consecutively, or according to any other sort of alternating regime).

5 When the substances or principles are to be administered simultaneously via the same route of administration, they may be administered as different pharmaceutical formulations or compositions or as part of a combined pharmaceutical formulation or composition. Also, when two or more active substances or principles are to be used as part of a combined treatment regimen,
10 each of the substances or principles may be administered in the same amount and according to the same regimen as used when the compound or principle is used on its own, and such combined use may or may not lead to a synergistic effect. However, when the combined use of the two or more active substances or principles leads to a synergistic effect, it may also be possible to reduce the
15 amount of one, more or all of the substances or principles to be administered, while still achieving the desired therapeutic action. This may for example be useful for avoiding, limiting or reducing any unwanted side-effects that are associated with the use of one or more of the substances or principles when they are used in their usual amounts, while still obtaining the desired pharmacological
20 or therapeutic effect.

Of course, the above includes the preparation, and methods of preparing, the polypeptides of the invention for the combined use with the above combination partners. Also included are the preparation, and methods of preparing, the above-
25 mentioned combination partners for the combined use with the polypeptides of the invention. Thus, the invention hereby e.g. provides methods of using, or preparing for use, an immune modulator / checkpoint inhibitor, such as an anti-PD1 antibody, such as pembrolizumab or nivolumab, for the administration in combination with a polypeptide of the invention, and more specifically for the
30 administration in a combination therapy regimen with a polypeptide of the invention.

Furthermore, the invention also encompasses kits comprising at least one polypeptide of the invention and one or more other components selected from the group consisting of other drugs used for the treatment of the diseases and disorders as described above, and devices as described below.

Pharmaceutical compositions, methods of administration, dosages

It will be clear to the skilled person that the above methods of treatment of a disease include the preparation of a medicament for the treatment of said disease. Thus, the invention further relates to pharmaceutical compositions for the treatment of the diseases mentioned hereinabove, wherein such compositions comprise at least one polypeptide of the invention.

The polypeptides of the invention and/or the compositions comprising the same can be administered to a patient in need thereof in any suitable manner, depending on the specific pharmaceutical formulation or composition to be used. Thus, the polypeptides of the invention and/or the compositions comprising the same can for example be administered intravenously (i.v.), subcutaneously (s.c.), intramuscularly (i.m.), intraperitoneally (i.p.), transdermally, orally, sublingually (e.g. in the form of a sublingual tablet, spray or drop placed under the tongue and adsorbed through the mucus membranes into the capillary network under the tongue), (intra-)nasally (e.g. in the form of a nasal spray and/or as an aerosol), topically, by means of a suppository, by inhalation, or any other suitable manner in an effective amount or dose.

The polypeptides of the invention and/or the compositions comprising the same are administered according to a regimen of treatment that is suitable for treating and/or alleviating the disease, disorder or condition to be treated or alleviated.

The clinician will generally be able to determine a suitable treatment regimen, depending on factors such as the disease, disorder or condition to be treated or

alleviated, the severity of the disease, the severity of the symptoms thereof, the specific polypeptide of the invention to be used, the specific route of administration and pharmaceutical formulation or composition to be used, the age, gender, weight, diet, general condition of the patient, and similar factors well known to the clinician. Generally, the treatment regimen will comprise the administration of one or more polypeptides of the invention, or of one or more compositions comprising the same, in therapeutically effective amounts or doses.

Generally, for the treatment and/or alleviation of the diseases, disorders and conditions mentioned herein and depending on the specific disease, disorder or condition to be treated, the potency of the specific polypeptide of the invention to be used, the specific route of administration and the specific pharmaceutical formulation or composition used, the polypeptides of the invention will generally be administered in an amount between 0.005 and 20.0 mg per kilogram of body weight and dose, preferably between 0.05 and 10.0 mg/kg/dose, and more preferably between 0.5 and 10 mg/kg/dose, either continuously (e.g. by infusion) or more preferably as single doses (such as e.g. twice a week, weekly, or monthly doses; cf. below), but can significantly vary, especially, depending on the before-mentioned parameters. Thus, in some cases it may be sufficient to use less than the minimum dose given above, whereas in other cases the upper limit may have to be exceeded. When administering large amounts it may be advisable to divide them up into a number of smaller doses spread over the day.

Depending on the specific polypeptide of the invention and its specific pharmacokinetic and other properties, it may be administered daily, every second, third, fourth, fifth or sixth day, weekly, monthly, and the like. An administration regimen could include long-term, weekly treatment. By "long-term" is meant at least two weeks and preferably months, or years of duration.

The efficacy of the polypeptides of the invention, and of compositions comprising the same, can be tested using any suitable *in vitro* assay, cell-based assay,

in vivo assay and/or animal model known *per se*, or any combination thereof, depending on the specific disease involved. Suitable assays and animal models will be clear to the skilled person, and for example include the assays and animal models used in the Examples below.

5

Preferably, the polypeptides of the invention are having better characteristics than conventional antibodies known in the art (such as the LRP6 binders described in the "Background of the Invention" section above) in at least one of these assays or models, and preferably in one or more of the *in vivo* models.

10

Formulations

For pharmaceutical use, the polypeptides of the invention may be formulated as a pharmaceutical preparation comprising (i) at least one polypeptide of the invention and (ii) at least one pharmaceutically acceptable carrier, diluent, excipient, adjuvant, and/or stabilizer, and (iii) optionally one or more further pharmacologically active polypeptides and/or compounds. By "pharmaceutically acceptable" is meant that the respective material does not show any biological or otherwise undesirable effects when administered to an individual and does not interact in a deleterious manner with any of the other components of the pharmaceutical composition (such as e.g. the pharmaceutically active ingredient) in which it is contained. Specific examples can be found in standard handbooks, such as e.g. Remington's Pharmaceutical Sciences, 18th Ed., Mack Publishing Company, USA (1990). For example, the polypeptides of the invention may be formulated and administered in any manner known *per se* for conventional antibodies and antibody fragments and other pharmaceutically active proteins. Thus, according to a further embodiment, the invention relates to a pharmaceutical composition or preparation that contains at least one polypeptide of the invention and at least one pharmaceutically acceptable carrier, diluent, excipient, adjuvant and/or stabilizer, and optionally one or more further pharmacologically active substances.

30

Pharmaceutical preparations for parenteral administration, such as intravenous, intramuscular, subcutaneous injection or intravenous infusion may for example be sterile solutions, suspensions, dispersions, emulsions, or powders which
5 comprise the active ingredient and which are suitable, optionally after a further dissolution or dilution step, for infusion or injection. Suitable carriers or diluents for such preparations for example include, without limitation, sterile water and pharmaceutically acceptable aqueous buffers and solutions such as physiological phosphate-buffered saline, Ringer's solutions, dextrose solution, and Hank's
10 solution; water oils; glycerol; ethanol; glycols such as propylene glycol, as well as mineral oils, animal oils and vegetable oils, for example peanut oil, soybean oil, as well as suitable mixtures thereof.

Solutions of the polypeptides of the invention may also contain a preservative to
15 prevent the growth of microorganisms, such as antibacterial and antifungal agents, for example, p-hydroxybenzoates, parabens, chlorobutanol, phenol, sorbic acid, thiomersal, (alkali metal salts of) ethylenediamine tetraacetic acid, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride. Optionally, emulsifiers and/or
20 dispersants may be used. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance of the required particle size in the case of dispersions or by the use of surfactants. Other agents delaying absorption, for example, aluminum monostearate and gelatin, may also be added. The solutions may be filled into injection vials, ampoules, infusion bottles,
25 and the like.

In all cases, the ultimate dosage form must be sterile, fluid and stable under the conditions of manufacture and storage. Sterile injectable solutions are prepared by incorporating the active compound in the required amount in the appropriate
30 solvent with various of the other ingredients enumerated above, as required, followed by filter sterilization. In the case of sterile powders for the preparation of

sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

5

Usually, aqueous solutions or suspensions will be preferred. Generally, suitable formulations for therapeutic proteins such as the polypeptides of the invention are buffered protein solutions, such as solutions including the protein in a suitable concentration (such as from 0.001 to 400 mg/ml, preferably from 0.005 to

10 200 mg/ml, more preferably 0.01 to 200 mg/ml, more preferably 1.0 - 100 mg/ml, such as 1.0 mg/ml (i.v. administration) or 100 mg/ml (s.c. administration) and an aqueous buffer such as:

- phosphate buffered saline, pH 7.4,
- other phosphate buffers, pH 6.2 to 8.2,
- 15 - acetate buffers, pH 3.2 to 7.5, preferably pH 4.8 to 5.5
- histidine buffers, pH 5.5 to 7.0,
- succinate buffers, pH 3.2 to 6.6, and
- citrate buffers, pH 2.1 to 6.2,

and, optionally, salts (e.g. NaCl) and/or sugars (such as e.g. sucrose and

20 trehalose) and/or other polyalcohols (such as e.g. mannitol and glycerol) for providing isotonicity of the solution.

Preferred buffered protein solutions are solutions including about 0.05 mg/ml of the polypeptide of the invention dissolved in 25 mM phosphate buffer, pH 6.5,

25 adjusted to isotonicity by adding 220 mM trehalose. In addition, other agents such as a detergent, e.g. 0.02 % Tween-20 or Tween-80, may be included in such solutions. Formulations for subcutaneous application may include significantly higher concentrations of the polypeptide of the invention, such as up to 100 mg/ml or even above 100 mg/ml. However, it will be clear to the person

30 skilled in the art that the ingredients and the amounts thereof as given above do only represent one, preferred option. Alternatives and variations thereof will be

immediately apparent to the skilled person, or can easily be conceived starting from the above disclosure.

Also, compared to conventional antibodies or antibody fragments, one major
5 advantage of the use of the polypeptides of the invention is that they can also be easily administered via routes other than parenteral administration and can be easily formulated for such administration. For example, as described in international patent application WO2004/041867, such polypeptides may be formulated for oral, intranasal, intrapulmonary and transdermal administration.

10 According to a further aspect of the invention, a polypeptide of the invention may be used in combination with a device useful for the administration of the polypeptide, such as a syringe, injector pen, micropump, or other device.

15 *Methods of manufacture and purification*

The invention further provides methods of manufacturing a polypeptide of the invention, such methods generally comprising the steps of:

- culturing host cells comprising a nucleic acid encoding a polypeptide of the
20 invention (hereinafter: "nucleic acid of the invention") under conditions that allow expression of the polypeptide of the invention; and,
- recovering or isolating the polypeptide expressed by the host cells from the culture; and
- optionally further purifying and/or modifying and/or formulating the polypeptide
25 of the invention.

A nucleic acid of the invention can e.g. be a DNA molecule comprising coding sequences as well as regulatory sequences and optionally natural or artificial introns, or can be a cDNA molecule. It may have its original codons or may have
30 an optimized codon usage that has been specifically adapted for expression in the intended host cell or host organism. According to one embodiment of the

invention, the nucleic acid of the invention is in essentially isolated form, as defined above.

The nucleic acid of the invention may also be in the form of, be present in and/or be part of a vector, such as for example a plasmid, cosmid or YAC, which again may be in essentially isolated form. The vector may especially be an expression vector, i.e. a vector that can provide for expression of the polypeptide *in vitro* and/or *in vivo* (e.g. in a suitable host cell, host organism and/or expression system). Such expression vector generally comprises at least one nucleic acid of the invention that is operably linked to one or more suitable regulatory element(s), such as promoter(s), enhancer(s), terminator(s), and the like. Specific examples of such regulatory elements and other elements, such as integration factor(s), selection marker(s), signal or leader sequence(s), reporter gene(s), and the like, useful or necessary for expressing polypeptides of the invention, are disclosed e.g. on pp. 131 to 133 of WO2006/040153.

The nucleic acids of the invention can be prepared or obtained in a manner known *per se* (e.g. by automated DNA synthesis and/or recombinant DNA technology), based on the information on the amino acid sequences for the polypeptides of the invention given herein.

According to another embodiment, the invention relates to a host or host cell that expresses or is capable of expressing a polypeptide of the invention; and/or that contains a nucleic acid encoding a polypeptide of the invention. According to a particularly preferred embodiment, said host cells are bacterial cells, yeast cells, fungal cells or mammalian cells.

For production on industrial scale, preferred heterologous hosts for the (industrial) production of immunoglobulin single variable domain polypeptides and protein therapeutics containing them include strains of *E. coli*, *Pichia pastoris*, and *S. cerevisiae* that are suitable for large scale expression, production and

fermentation, and in particular for large scale (bio-)pharmaceutical expression, production and fermentation.

Polypeptides of the invention produced in a cell as set out above can be
5 produced either intracellularly (e.g. in the cytosol, in the periplasma or in inclusion bodies) and can then be isolated from the host cells and optionally be further purified; or they can be produced extracellularly (secreted into the medium in which the host cells are cultured) and then isolated from the culture medium and optionally further purified.

10 Further methods and reagents used for the recombinant production of polypeptides, such as suitable expression vectors, transformation or transfection methods, selection markers, methods of induction of protein expression, culture conditions, and the like, are known in the art. Similarly, protein isolation and
15 purification techniques useful in a method of manufacture of a polypeptide of the invention are well known to the skilled person.

Production of the polypeptides of the invention through fermentation in convenient recombinant host organisms such as *E. coli* and yeast is cost-
20 effective, as compared to conventional antibodies which usually require expensive mammalian cell culture facilities. Furthermore, achievable levels of expression are high and yields of the polypeptides of the invention are in the range of 1 to 10 g/l (*E. coli*) and up to 10 g/l (yeast) and more.

EXAMPLES

Example 1: Immunization of llamas with LRP5 and LRP6 for the induction of humoral immune responses

5

Several protocols for the immunization of llamas needed to be worked out and implemented for identification of LRP5/LRP6 cross-reactive binding VHH domains: Llamas were initially immunized with recombinant extracellular domains of LRP6 and LRP5 proteins (human and mouse). However, functional

10 characterization of the above mentioned LRP5 recombinant protein revealed that only the Wnt1 class binding epitope was properly folded. In contrast, there was no indication for proper folding of the LRP5-Wnt3a class binding domain. Thus, further work was required to develop suitable antigens for immunization. As a work-around, llamas were immunized with HEK293 cells stably transfected with

15 human LRP5 or human LRP6. However, also then, only very low expression of human LRP5 could be achieved, by transient or by stable transfection, and using different cell lines (HEK293, CHO and NIH-3T3 cells). Thus, even further work was required to achieve sufficient expression of LRP5. In the end, and after some unsuccessful trial and error, this could be achieved by developing a protocol

20 involving stable co-transfection of HEK293 cells with MesDC-2, a chaperone intended to increase exogenous LRP5 expression. Even then, i.e. upon co-expression of MesDC-2 during the generation of the LRP5 stable transfected cell line, instability of protein expression was repeatedly observed. This resulted in the problem that LRP5 expression could be lost during immunization and

25 selection. To solve this further problem, passaging of the cells expressing LRP5 was limited as much as possible and additional cell sorting was performed to enrich for cells expressing LRP5.

30

Llamas were additionally immunized with LRP5-coding DNA and LRP6-coding DNA with and without the hMesDC-2 chaperone in opposite flanks. Additional boosts were delivered to several llamas in an attempt to enhance the cross-

reactive immune response, with the objective to increase chances of identifying LRP5/LRP6 cross-reactive VHH domains.

Immune blood (PBL) samples were taken at regular intervals, serum responses were determined, and total RNA was prepared from the isolated PBL. Medium serum responses to LRP6 were observed, in contrast to low serum responses to LRP5 upon immunization with recombinant protein. A medium LRP5 immune response was observed for llamas immunized with DNA. In contrast, very low immune response was observed for the cell immunizations. Additionally, synthetic libraries were also explored. Nevertheless, finally, sufficient diversity of the repertoire could be achieved for continuing with the next steps, as outlined in Example 2.

Example 2: Isolation of LRP5 and LRP6 binding monovalent VHH domains (VHHs)

Library construction:

Total RNA was extracted immediately following collection of the immune tissues, and RNA integrity and concentration was verified. cDNA samples were made from these RNA preparations. Nucleotide sequences encoding VHHs were amplified from the cDNA samples in a one-step RT-PCR reaction. The 700bp amplicons specifically amplified from the IgG2 and IgG3 cDNAs in the sample were isolated from agarose gel and subsequently used as template in a nested PCR reaction. The PCR products were subsequently digested with *Sfi*I and *Bst*EII and ligated into the corresponding restriction sites of phagemid vector pAX50. The ligation mixtures were electroporated into *Escherichia coli* TG-1. The resulting pool of transformants constituted the genetic diversity of the phage display library.

pAX50 is an expression vector derived from pUC119, which contains a resistance gene for ampicillin and the *lac* promoter followed by the coding sequence of the pIII protein signal peptide in frame with a downstream VHH domain cloning site. In frame with the VHH domain coding sequence, the vector codes for a C-terminal Myc and hexa-histidine tag and a coli phage pIII protein. After infection of the *E. coli* TG-1 library clones with helper phage, the presence of pAX50 allows for production of phage particles from these clones, displaying the individual VHH domains as a fusion protein with the pIII protein.

Selection:

The VHH domain-phagemid libraries were constructed and used for selections. Given the very high species homology across species (between llama and human LRP5 and LRP6), it was uncertain whether the immune response raised in the llamas would raise sufficient diversity of the VHH domains. Therefore, two synthetic libraries were used in parallel with the immune libraries during selections.

Different strategies were used during selections as follows:

- Alternation of LRP5 and LRP6 derived tools to enhance chances of identifying LRP5/LRP6 cross-reactive VHH domains, e.g. selections on libraries from LRP5 immunized llamas with LRP6 derived proteins or the use of both LRP5 and LRP6 proteins during selection on synthetic libraries.
- Alternation of species source to select for human/mouse LRP5/LRP6 cross-reactive VHH domains (mouse cross-reactivity of such LRP5 and LRP6 antagonist allowing to evaluate efficacy, i.e. tumor growth inhibition, and safety profiles, required for assessment of the therapeutic window, in the same pre-clinical models (i.e. in xenograft tumor-mouse models)).
- "In solution" selections with recombinant proteins to keep the epitopes in their native conformation: As an additional obstacle, LRP5 and LRP6 recombinant proteins were found to lose their proper folding if directly coated on ELISA binding plates. Therefore, the recombinant proteins were biotinylated and, after

confirming proper folding in functional assays, were used for selection “in solution”.

- Selections using cells overexpressing LRP5 or LRP6, to have native conformation of the receptors. This surprisingly turned out to be an important
5 ruse, especially needed to improve selection of binders to the Wnt3a–class binding domain of LRP5, since functional data of recombinant protein showed lack of proper folding of the Wnt3a binding epitope.

Example 3: Screening of the monovalent VHHs

10 After selection, clones were grown in 96 deep well plates (1 mL volume), and VHH expression was induced by adding IPTG. Periplasmic extracts of single clones were prepared according to standard methods, as e.g. reported in WO2011/107507, and screened for binding to human LRP6 and LRP5. Initially,
15 the periplasmic extracts were screened in binding ELISA assays using recombinant LRP5 and LRP6, which represent sensitive, robust and high-throughput assays, when compared to FACS-based binding assays. After purification, VHHs identified in ELISA assays were further characterized using binding FACS assay to confirm binding of purified VHHs to the LRP5 and LRP6
20 receptors in their native conformation.

Usually, a good correlation between ELISA and FACS binding assays is expected. However, in the present case, the best LRP5 and LRP6 binding VHHs in ELISA assays (i.e. with high affinity to recombinant LRP5 or LRP6 ectodomain)
25 did not show any binding or very weak binding to human LRP5 and LRP6 in a FACS binding assay, as shown in Figure 2 for the panel “2” binders. The use of different coating buffers (dPBS vs. bicarbonate buffer) and blocking solutions in the ELISA set-up (Marvel vs. BSA) did not resolve the observed discrepancy. Instead, it was found out that very weak binders in ELISA showed high affinity to
30 LRP5 and LRP6 in binding FACS assays using LRP5 and LRP6 expressing cells, as shown in Figure 2 for the panel “1” binders. These further data and

experiments thus allowed for selection of high affinity binders recognizing the native conformation of the two receptors. Additionally, confirmation was thereby obtained that these high affinity binders are recognizing a conformation dependent epitope, and not a linear epitope in LRP5 and LRP6 proteins. These
5 further non-routine data and experiments thus allowed for selection of therapeutically relevant LRP5 and LRP6 binders, which should have high affinity towards LRP5 and LRP6 expressed on the plasma membrane in their native conformation.

10 Thus, despite (i) the low throughput of FACS binding assays, (ii) the less robust assay set-up, and (iii) the above-described difficulties encountered due to the loss of recombinant protein expression upon passaging of the cells over-expressing LRP5, these assays were subsequently used for further selection and characterization of high affinity VHH binders. Briefly, cells were incubated with
15 purified VHH dilutions (1:5 serial dilutions from 1 μ M to 1 pM, final concentration) for 1.5 hours at 4 °C on a plate shaker. After washing the cells 5 x with FACS buffer, consisting of 1x phosphate buffered saline (PBS) + 10% fetal bovine serum (FBS) + 0,05% sodium azide, they were incubated for 30 minute up to 1 hour at 4°C with a polyclonal mouse antibody which binds to the framework
20 regions of the VHHs and, therefore, binds to all the LRP5 and/or LRP6 binders tested. After washing the cells 3X with FACS buffer, the cells were incubated for 30 minutes up to 1 hour at 4 °C with the labeled secondary antibody (anti-mouse PE), followed by 3X washing step with FACS buffer. Fluorescence was measured using FACS Array (BD).

25 A total of hundred LRP5/LRP6 cross-reactive VHHs families/clusters were identified from immune libraries and from synthetic origin, based on binding FACS data and sequence analysis. Representative examples thereof are shown, and defined by their sequence, further below. The VHHs were expressed in *E.coli*
30 and purified. In case expression in *E.coli* proved insufficient, VHHs were

produced in *Pichia pastoris*. A brief description of expression and purification of VHHs is reported further below.

Generic expression of VHHs in *E.coli*:

5

The encoding sequences were cloned into the pAX100 expression vector and expressed in *E.coli* as c-Myc hexa-histidine-tagged proteins. *E.coli* TG-1 cells containing the VHH constructs of interest were grown (37 °C, 250 rpm) in shake flasks in TB medium supplemented with kanamycin and induced by addition of 1mM IPTG for expression. After spinning the cell cultures, periplasmic extracts were prepared by freeze-thawing the pellets and resuspending in dPBS.

10

Generic expression of VHHs in *Pichia (P.) pastoris*:

15

The encoding sequences were cloned into the pAX159 expression vector and expressed in *P. pastoris* as c-Myc hexa-histidine-tagged proteins. *P. pastoris* X-33 cells containing VHH constructs of interest were grown (30 °C, 250 rpm) in BGCM (Buffered Glycerol-Complex Medium; Invitrogen). On the third day, the medium was switched to BMCM (Buffered Methanol-Complex Medium; Invitrogen) and the culture was grown further and was regularly induced by addition of 0.5 vol% methanol (100%). After spinning the cell culture, supernatant (containing the secreted VHH) was collected.

20

VHH purification:

25

Hexa-histidine-tagged VHHs were purified on the Tecan EVO150 by immobilised metal affinity chromatography (RoboColumns 100ul Nickel Sepharose™ 6 FF, Atoll), eluted from the column with 250mM imidazole and subsequently desalted towards dPBS. The purity and integrity of VHHs was verified by SDS-PAGE and/or western blot using anti-Myc and anti-VHH detection.

30

Example 4: *In vitro* characterization of purified monovalent VHHs

After VHH screening, the purified VHHs having high affinity to cells expressing LRP5 and LRP6 were characterized by using several functional and biophysical assays as described below:

4.1 LRP5 and LRP6 binding potency and cross-reactivity: FACS-based DKK1 competition assay

During characterization of LRP5/LRP6 cross-reactive monovalent VHHs, it was observed that data obtained in binding FACS assay did not always correlate with the potency observed in Wnt1 and Wnt3a reporter assays, most likely due to fast off-rate of some of the VHHs. Therefore, this additional assay needed to be established (i.e. a DKK1 competition FACS), which proved to be more reliable for selectivity and binding potency determination and comparison between LRP5 and LRP6 binding. The aim was to select functional VHHs which bind to LRP5 and LRP6 with a similar potency, in order to achieve blockade of both receptors at the same concentration. The identified Wnt1 and Wnt3a functional VHHs were thus characterized in DKK-1 competition FACS as follows:

For FACS-based DKK1 competition assay, HEK293 cells with stable overexpression of human LRP5 or human LRP6 were used. Human recombinant DKK1 (rhDKK1 - R&D Systems, Cat 5439-DK/CF) was added to the cells at the constant final concentration of 1nM. Cells were incubated with rhDKK1 and LRP5 and/or LRP6 binder dilutions (1:5 serial dilution of the purified VHHs) for 1.5 hours at 4 °C on a plate shaker. After washing the cells three times with FACS buffer, they were incubated with biotinylated goat anti-human DKK1 (R&D Systems, Cat BAF1096) for 30 minutes at 4 °C on a plate shaker. After washing the cells three times with FACS buffer, they were incubated with Streptavidin PE (BD Biosciences, Cat 554061) for 30 minute to 1 hour at 4 °C on a plate shaker in the dark. Cells were washed twice with FACS buffer and fluorescence was measured using FACS Array (BD) and MCF values were reported.

LRP5/LRP6 cross-reactive VHHs are expected to compete with human DKK1 for binding to HEK293 overexpressing human LRP5, as well as for binding to HEK293 overexpressing human LRP6. In contrast, an LRP5 specific VHH would compete with human DKK1 for binding to HEK293 cells overexpressing human LRP5 but not, or with very low potency ($>200\text{nM}$) for binding to HEK293 cells overexpressing human LRP6 (and the same would apply, vice versa, to LRP6 specific VHHs). As a result of this experiment, it could be shown that the present LRP5/LRP6 cross-reactive VHHs competed with human DKK1 for binding to HEK293 cells overexpressing human LRP5, as well as to those overexpressing human LRP6 (i.e. decrease in MCF value with increasing concentration of the binder with complete inhibition of DKK1 binding corresponded to ≤ 60 MCF values at the highest concentration tested).

4.2 Species cross-reactivity: mouse and cynomolgus monkey

To determine whether the selected panel of LRP5/LRP6 cross-reactive VHHs was able to bind to LRP5 and LRP6 of mouse and of cynomolgus monkey origin, DKK1 competition FACS was performed as follows:

Serial dilutions of VHHs were incubated with HEK293 cells stably expressing mouse LRP5, cyno LRP5, mouse LRP6, or cyno LRP6 in the presence of 1 and 0.3 nM hDKK1 (concentration below EC50 value of mouse and cyno, respectively). Binding of DKK1 to the cells was detected using a biotinylated anti-DKK1 antibody with Streptavidin-PE as secondary detection as described above. As a result, such cross-reactivity could be demonstrated.

4.3 Epitope binning

Binning experiments were performed for the most potent Wnt1 signaling blocking LRP5/LRP6 cross-reactive VHHs, to identify different epitope bins. In particular,

individual VHHs were analyzed for their ability to compete with other biotinylated VHHs (called reference VHHs) for LRP5 and LRP6 receptor binding using FACS based assays. Serial dilutions of the individual VHHs were incubated on HEK293 stably expressing human LRP5 or LRP6 together with 200pM or 500pM biotinylated reference VHH (concentrations below EC50 value). The binding of biotinylated reference VHH to the cells was detected using streptavidin-PE. A VHH competing with the reference VHHs for binding to LRP5 and LRP6 shows a decrease in the fluorescence measured using FACS Array.

As a result of these experiments, the Wnt1 blockers could be categorized into three bins. For the Wnt3a blockers, the lower affinity of the VHHs did not allow to perform epitope binning experiments.

4.4 Wnt1 and Wnt3a reporter assays

The ability of the LRP5/LRP6 cross-reactive VHHs to inhibit Wnt signaling was tested in functional Wnt1 and Wnt3a assays. Also in this regard, no established protocol could be used, but several attempts needed to be tried to establish biochemical functional assays, such as Wnt1/Wnt3a – LRP5 / LRP6 blocking assays: In addition to the difficulties encountered with recombinant LRP5 and LRP6 proteins (see Example 1), a functional recombinant Wnt1 ligand (including commercially available) is not available. Wnt proteins contain many conserved cysteines and are modified by mono-unsaturated fatty acid (palmitoleic acid), attached to a conserved serine. These post-translational modifications are required for efficient signaling and for Wnt secretion. Structural analyses show that one of the domains, containing the palmitoleic acid lipid, is required for binding to the Frizzled receptors, leading to a conformational change that allows the interaction of the Wnt ligands with LRP5 and LRP6 on the cell surface. Thus, it turned out that such post-translational modification is required for functional studies involving this protein, but at the same time such lipid-based post-translational modifications render these proteins very difficult to express and

purify (low solubility). Thus, this turned out to be a major hurdle for biochemical assays.

Thus, a cell based functional assay was developed for characterization of the purified VHHs: A Wnt beta-lactamase reporter gene assay. In particular for the Wnt1 pathway inhibition, CellSensor LEF/TCF-bla FreeStyle 293F cells (Invitrogen, Cat. K1677) were transfected with human Wnt1, and clones with stable overexpression of human Wnt1 were selected. For testing the Wnt3a pathway inhibition, CellSensor LEF/TCF-bla FreeStyle 293F cells with stable overexpression of human Wnt3a were generated. The CellSensor® LEF/TCF - bla FreeStyle™ 293 cell line contains a beta-lactamase reporter gene under control of the Wnt inducible LEF/TCF promoter, which is stably integrated into FreeStyle™ 293 cells (Invitrogen). The expression of Wnt1 or Wnt3a in these cells thus results in the constitutive expression and, therefore, enzymatic activity of the beta-lactamase. The treatment with LRP5/LRP6 cross-reactive functional VHHs is thus expected to lead to inhibition of Wnt1 and Wnt3a pathway leading to inhibition of beta-lactamase enzymatic activity.

For the assay, 1E06 / ml cells with overexpression of Wnt1 or Wnt3a were seeded into a 384 well tissue culture plate and incubated overnight at 37 °C. The following day, serial dilutions of various LRP5/LRP6 cross-reactive VHH solutions were prepared and added to the cells in the presence of LiCl at the final concentration of 10nM. DKK1, as positive control, was added to the cells at the final concentration of 200nM. DKK1 treatment resulted in a complete inhibition of Wnt1 and Wnt3a pathway, and therefore a complete inhibition of beta-lactamase enzymatic activity. The cells were incubated overnight at 37 °C. The following day, beta-lactamase enzymatic activity was measured according to the manufacturer's instructions (Invitrogen, Cat K1085). For fluorescence emission, values at 460 nm and 530 nm were obtained using a standard fluorescence plate reader and the 460/530 nm emission ratios plotted for the indicated treatment.

Efficacy was calculated against the positive control (DKK1; 200nM final concentration).

A total of twelve LRP5/LRP6 cross-reactive Wnt1 blockers with full efficacy and potency better than 50nM were selected. Fourteen cross-reactive Wnt3a blockers, mainly with low potencies, were identified. Only one Wnt3a blocker showed good potency (below 5nM).

4.5 Wnt1 and Wnt3a phosphorylation assays

The most potent and efficacious leads from each bin for the Wnt1 blockers and the most potent and efficacious Wnt3a blockers were subsequently tested in Wnt1 and Wnt3a dependent LRP5 and LRP6 phosphorylation assays. Cellsensor LEF/TCF 293F cells from Invitrogen (cat K1677), co-transfected with expression vectors coding for either Wnt1 or Wnt3a, were used in the phosphorylation assays. Since Wnt-Frizzled-LRP5 or -LRP6 complex formation results in LRP5 or LRP6 phosphorylation and subsequent downstream signaling, quantification of phosphorylation can be used to measure such signaling. To obtain a LRP5 and LRP6 specific read-out, the cells were lysed and immuno-precipitation was performed with an LRP6 or LRP5 selective antibody (directed against the intracellular domain of the two receptors). In Western blot, phosphorylated LRP6 or LRP5 was detected using a polyclonal anti-phospho-LRP6 (Ser1490) antibody (Cell Signaling Technology), which detects both LRP6 and LRP5 phosphorylated protein. A selected panel of purified Wnt1 and Wnt3a blocking VHHs, containing at least one representative VHH from each bin, were tested at final concentrations of 10 to 100nM. In particular, cells were incubated overnight in the presence of the blocking VHHs prior cell lysis and LRP5 and LRP6 immuno-precipitation. Efficacy of the Wnt1 and Wnt3a blocking VHHs in blocking LRP5 and LRP6 phosphorylation was calculated via quantification of the Western blot bands against the positive control (DKK1, final concentration of 1 μ M).

4.6 Biophysical characterization

The LRP5/LRP6 cross-reactive VHHs were further characterized for expression and purification in *E. coli* and in *Pichia pastoris*, as reported in Example 3. In particular, expression yields for the monovalent lead panel VHHs was considered acceptable if they were above 0.1 mg/L. The selected LRP5/LRP6 cross-reactive VHHs showed expression ranging between 0.1 and 8.2 mg/L in *E.coli* and higher in *Pichia pastoris* (> 1mg/L). The expression was evaluated by SDS-PPAGE analysis.

Thermal stability of the monovalent LRP5/LRP6 cross-reactive VHHs was determined in a fluorescence-based thermal shift assay (TSA) using the Lightcycler (Roche). VHHs were incubated at different pH values in the presence of Sypro Orange and a temperature gradient was applied. Upon heat-induced unfolding, hydrophobic patches of the proteins are exposed to which the Sypro Orange binds resulting in an increase in fluorescence intensity (Ex/Em = 465/580nm). The inflection point of the first derivative of the fluorescence intensity curve serves as a measure of the melting temperature (T_m). For all VHHs, T_m increased with increasing pH and levels off at pH 6, a typical T_m pattern seen for VHHs. An average of 82 °C at pH 7 was obtained for the LRP5/LRP6 cross-reactive Wnt1 blocker VHHs and Wnt3a blocker VHHs.

Potential occurrence of aggregation and multimerisation for LRP5 and LRP6 VHHs was investigated by analytical size exclusion chromatography (SEC). To this end, 8ug of purified VHH sample at 0,5 mg/mL were injected via the Dionex Ultimate 3000 equipment on an Agilent SEC-3 column. L-arginine buffer (10mM phosphate, 300mM Arg-HCl, pH 6.0) was used as mobile phase and a flow rate of 1mL/min was applied. None of the LRP5/LRP6 VHHs showed major aggregation issues during SEC analysis: profiles indicated more than 95% monomer for most samples.

Example 5: Generation and characterization of half-life extended biparatopic constructs

The LRP5/LRP6 cross-reactive Wnt1 and Wnt3a VHHs were used as building blocks to generate a biparatopic construct as depicted in Figure 1. A genetic fusion to a serum albumin binding VHH was used as a half-life extension methodology. The three building blocks (Wnt1 blocker, Wnt3a blocker, and the albumin binder) were linked via a flexible linker. VHHs were produced in *P. pastoris* and purified as described in Example 3. The resulting constructs, i.e. biparatopic, half-life extended LRP5/LRP6 cross-reactive VHH constructs, were cloned in *P. pastoris* expression vector pAX159, in the form of C-terminally cMyc-hexa-histidine tagged VHH constructs, according to standard procedures as e.g. reported in WO2012/131078. Different orientation of the building blocks and different linkers, esp. GS-linkers, were explored. A relatively long GS-linker was selected, based on modeling data reflecting an extended surface area between the potential Wnt1 and Wnt3a binding sites in LRP6 (i.e. beta-propellers 1 and 2 not being in close proximity to beta-propeller 3). Best results regarding potency in the combined Wnt1 and Wnt3a reporter assay were obtained by putting the human serum albumin/HSA binding VHH in the middle. A 35 GS linker was used, and Wnt1 and Wnt3a VHH blockers arranged in a preferred order.

For selection of optimal VHH binders and binder combinations, a library was generated where the human serum albumin (HSA) binding VHH was placed in between the LRP5/LRP6 Wnt1-Wnt3a blockers. In particular, the panel of high affinity binders with high potency and efficacy in Wnt1 or Wnt3a assays (reporter and phosphorylation assays) were used in the library to generate half-life extended biparatopic constructs designed as depicted in Figure 1. After expression in *Pichia pastoris* (as reported in Example 3) followed by purification, the half-life extended biparatopic constructs were subsequently screened in the Wnt1 and Wnt3a reporter assays (described in Example 4) in the presence of

30uM HSA at three dilutions (1/100, 1/1000, 1/7000), to assess efficacy and relative potency. In general, a good correlation was observed between the data in the Wnt1 and Wnt3a reporter assays and high efficacies were measured for numerous formats. A total of 11 half-life extended biparatopic LRP5 and LRP6 constructs were selected for further characterization, taking into account the efficacy in both reporter assays and diversity of the Wnt1 and Wnt3a blockers. This further characterization assays are described below.

Wnt1 / Wnt3a reporter assays:

Wnt1 and Wnt3a reporter assays were performed as described in Example 4.4, in the presence of 30uM HSA final concentration. The purified biparatopic LRP5/LRP6 constructs were tested at 12 dilutions, starting from 2.5 uM. The majority of the constructs showed high potency - ranging between 1.7 nM and 0.16 nM - and full efficacy in both reporter assays, as shown in Table IV below. The sequences of the individual Wnt1 and Wnt3a inhibiting VHH domains set out therein are provided in Table V below:

Table IV: Potency and efficacy of selected half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs in the reporter assays in the presence of HSA

Construct	N-term	C-term	Wnt1 reporter assay		Wnt3a reporter assay	
			IC ₅₀ (M)	% inh.	IC ₅₀ (M)	% inh.
F013500053	F0129093A01	F0130333G06	2,3E-10	102	4,1E-10	97
F013500039	F0129093A03	F0130333E06	1,7E-09	102	1,6E-09	82
F013500046	F0130332D03	F0129093A01	5,1E-10	102	6,0E-10	96
F013500016	F0130332D03	F0130367B10	3,1E-10	102	2,7E-10	99
F013500018	F0130332D03	F0130378B05	7,4E-10	102	1,3E-09	98
F013500047	F0130333E06	F0129093A01	8,8E-10	102	9,1E-10	94
F013500026	F0130333E06	F0130367B10	3,3E-10	102	2,6E-10	100
F013500021	F0130333G06	F0130367B10	1,6E-10	102	1,8E-10	102

F013500032	F0130378A04	F0130333G06	2,4E-10	102	4,3E-10	94
F013500033	F0130378B05	F0130333G06	1,8E-10	102	2,5E-10	88
F013500030	F0130378B05	F0130372C08	1,4E-09	102	1,3E-09	96

Table V: Sequences of the VHH domains set out in Table IV

VHH ID Specificity	Amino Acid Sequence	SEQ ID NO:
F0129093A01 Wnt3a	EVQLVESGGGLVQPGGSLRLSCAASGRTFSSYAMGWFRQAPGKER EFVAAISWSGGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDT AVYYCAASPIPYGSLLRRRNNDYWGQGTLVTVSS <u>AAAEQKLISEED</u> <u>LNGAAHHHHHH</u>	28
F0130333G06 Wnt1	EVQLVESGGGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAPGKER EFVAAIRRSGRRTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTA VYYCAAARRVRSSTRYNTGTWWWEYWGQGTLVTVSS <u>AAAEQKLI</u> <u>SEEDLNGAAHHHHHH</u>	29
F0129093A03 Wnt3a	EVQLVESGGGLVQPGGSLRLSCAASGRTFSTYVMGWFRQAPGKER EFVAAINWSGSRTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDT AVYYCAASRSSYAGRTYYELYDYWGQGTLVTVSS <u>AAAEQKLISEEDL</u> <u>NGAAHHHHHH</u>	30
F0130333E06 Wnt1	EVQLVESGGGLVQPGGSLRLSCAASGRTFSTYTVGWFRQAPGKER FVAAIRRRGSSTYYSDSVKGRFTISRDN SKNTVYLQMNSLRPEDTAV YYCAADTRTVALLQYRYDYWGQGTLVTVSS <u>AAAEQKLISEEDLNGA</u> <u>AHHHHHH</u>	31
F0130332D03 Wnt1	EVQLVESGGGLVQPGGSLRLSCAASGLTFSRYTMGWFRQAPGKER EFVAAINRSGGSTYYSDSVKGRFTISRDN SKNTVYLQMNSLRPEDTA VYYCAADRRGRGENYSLLYSSNRYEYWGQGTLVTVSS <u>AAAEQKLISE</u> <u>EDLNGAAHHHHHH</u>	32
F0130367B10 Wnt3a	EVQLVESGGGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAPGKER EFVAAISWRSGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEGT AVYYCAADPRGYGVAYVSAYYEWGQGTLVTVSS <u>GAAEQKLISEED</u> <u>LNGAAHHHHHH</u>	33
F0130378B05 Wnt3a	EVQLVESGGGLVQPGGSLRLSCVASGRTFSSYAMGWFRQAPGKER EFVAAISRSGGRTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTA VYYCAADRRVYSTLPPTTSRYNYWGQGTLVTVSS <u>AAAEQKLISEEDL</u> <u>NGAAHHHHHH</u>	34

F0130378A04 Wnt3a	EVQLVESGGGLVQPGGSLRLSCAASGRTFSSYAMGWFRQAPGKER EFVAAITRTGRRYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTA VYYCAADRRGYYYYDSSFYDWGQGTLTVTVSSAAAEQKLISEEDLN <u>GAAHHHHHH</u>	35
F0130372C08 Wnt1	EVQLVESGGGLVQPGGSLRLSCAASGRTFSDYGMGWFRQAPGKE REFVAAISWSGGRTYYADSVKGRFTISRDN SKNTVYLQMNSLRPED TAVYYCAAKRRGRGSVSPNSSRYNYWGQGTLTVTVSSAAAEQKLIS <u>EEDLNGAAHHHHHH</u>	36
F013500016	EVQLVESGGGLVQPGGSLRLSCAASGLTFSRYTMGWFRQAPGKER EFVAAINRSGGSTYYSDSVKGRFTISRDN SKNTVYLQMNSLRPEDTA VYYCAADRRGRGENYSLLYSSNRYEYWGQGTLTVTVSSGGGGSGGG GSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLVESGGGLVQPGN SLRLSCAASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYAD SVKGRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLRSSQGT LTVTVSSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSEV QLVESGGGLVQPGGSLRLSCAASGRTFSSYAMGWFRQAPGKEREF VAAISWRSGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEGTAV YYCAADPRGYGVAYVSAYYEWGQGTLTVTVSSGAAEQKLISEEDLN <u>GAAHHHHHH</u>	37
F013500018	EVQLVESGGGLVQPGGSLRLSCAASGLTFSRYTMGWFRQAPGKER EFVAAINRSGGSTYYSDSVKGRFTISRDN SKNTVYLQMNSLRPEDTA VYYCAADRRGRGENYSLLYSSNRYEYWGQGTLTVTVSSGGGGSGGG GSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLVESGGGLVQPGN SLRLSCAASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYAD SVKGRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLRSSQGT LTVTVSSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSEV QLVESGGGLVQPGGSLRLSCVASGRTFSSYAMGWFRQAPGKEREF VAAISRSGRTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVY YCAADRRVYSTLPPTTSRYNYWGQGTLTVTVSSGAAEQKLISEEDLN <u>GAAHHHHHH</u>	38
F013500021	EVQLVESGGGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAPGKER EFVAAIRRSRRYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTA VYYCAAARRVRSSTRYNTGTWWWEYWGQGTLTVTVSSGGGGSGGG GGSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLVESGGGLVQPG NSLRLSCAASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYA DSVKGRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLRSSQG TLTVTVSSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSE VQLVESGGGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAPGKEREF FVAAISWRSGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEGTA VYYCAADPRGYGVAYVSAYYEWGQGTLTVTVSSGAAEQKLISEEDL <u>NGAAHHHHHH</u>	39

F013500026	EVQLVESGGGLVQPGGSLRLSCAASGRTFSTYTVGWFRQAPGKER FVAAIRRRGSSTYYSDSVKGRFTISRDN SKNTVYLQMNSLRPEDTAV YYCAADTRTVALLQYRYDYWGQGLTVTVSSGGGGSGGGGSGGGG SGGGSGGGGSGGGGSGGGGSEVQLVESGGGLVQPGNSLRLSCA ASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYADSVKGRFT ISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLRSSQGLTVTVSSG GGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLVESG GGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAPGKEREFVAAISW RSGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEGTAVYYCAAD PRGYGVAYVSAYEYWGQGLTVTVSSGAAEQKLISEEDLN GAAHH HHHH	40
F013500030	EVQLVESGGGLVQPGGSLRLSCVASGRTFSSYAMGWFRQAPGKER EFVAAISRSGRTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTA VYYCAADRRVYSTLPPTTSRYNYWGQGLTVTVSSGGGGSGGGGSG GGGSGGGGSGGGGSGGGGSGGGGSEVQLVESGGGLVQPGNSLR LSCAASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYADSVK GRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLRSSQGLTVTV SSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLVE SGGGLVQPGGSLRLSCAASGRTFSDYGMGWFRQAPGKEREFVAAI SWSGGRTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCA AKRRGRGSVSPNSSRYNYWGQGLTVTVSSGAAEQKLISEEDLN GA AHHHHH	41
F013500032	EVQLVESGGGLVQPGGSLRLSCAASGRTFSSYAMGWFRQAPGKER EFVAAITRTGRRYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTA VYYCAADRRGYYYDYSSFYDYWGQGLTVTVSSGGGGSGGGGSGG GGSGGGGSGGGGSGGGGSGGGGSEVQLVESGGGLVQPGNSLRLS CAASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYADSVKG RFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLRSSQGLTVTVS SGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLVES GGGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAPGKEREFVAAIR RSGRRTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAA ARRVRSSTRYNTGTWWWEYWGQGLTVTVSSGAAEQKLISEEDLN GAAHHHHH	42

F013500033	EVQLVESGGGLVQPGGSLRLSCVASGRTFSSYAMGWFRQAPGKER EFVAAISRSGGRITYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTA VYYCAADRRVYSTLPPTTSRYNYWGQGLTVTVSSGGGGSGGGGSG GGGSGGGGSGGGGSGGGGSGGGGSEVQLVESGGGLVQPGNSLR LSCAASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYADSVK GRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLRSSQGLTVTV SSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLVE SGGGLVQPGGSLRLSCAASGFTFSSYAMGWFRQAPGKEREFVAAI RRSGRRTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCA AARRVRSSTRYNTGTWWWEYWGQGLTVTVSSGAAEQKLISEEDL <u>NGAAHHHHHH</u>	43
F013500039	EVQLVESGGGLVQPGGSLRLSCAASGRTFSTYVMGWFRQAPGKER EFVAAINWWSRITYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDT AVYYCAASRSSYAGRTYYELYDYWGQGLTVTVSSGGGGSGGGGSG GGGSGGGGSGGGGSGGGGSGGGGSEVQLVESGGGLVQPGNSLR LSCAASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYADSVK GRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLRSSQGLTVTV SSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLVE SGGGLVQPGGSLRLSCAASGRTFSTYTVGWFRQAPGKEREFVAAIR RRGSSTYYSDSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAAD TRTVALLQYRYDYWGQGLTVTVSSGAAEQKLISEEDLNGAAHHHH <u>HH</u>	44
F013500046	EVQLVESGGGLVQPGGSLRLSCAASGLTFSRYTMGWFRQAPGKER EFVAAINRSGGSTYYSDSVKGRFTISRDN SKNTVYLQMNSLRPEDTA VYYCAADRRGRGENYSLLYSSNRYEYWGQGLTVTVSSGGGGSGGG GSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLVESGGGLVQPGN SLRLSCAASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYAD SVKGRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLRSSQGL TVTVSSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSEV QLVESGGGLVQPGGSLRLSCAASGRTFSSYAMGWFRQAPGKEREF VAAISWSGGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAV YYCAASPIPYGSLRRRRNNYDYWGQGLTVTVSSGAAEQKLISEEDLN <u>GAAHHHHHH</u>	45

F013500047	EVQLVESGGGLVQPGGSLRLSCAASGRTFSTYTVGWFRQAPGKER FVAAIRRRGSSTYYSDSVKGRFTISRDN SKNTVYLQMNSLRPEDTAV YYCAADTRTVALLQYRYDYWGQGLTVTVSSGGGGSGGGGSGGGG SGGGSGGGGSGGGGSGGGGSEVQLVESGGGLVQPGNSLRLSCA ASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYADSVKGRFT ISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLRSSQGLTVTVSSG GGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLVESG GGLVQPGGSLRLSCAASGRTFSSYAMGWFRQAPGKEREFVAAISW SGGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCAAS PIPYGSLLRRRNNDYWGQGLTVTVSS <u>GAAEQKLISEEDLNGAAHH</u> <u>HHHH</u>	46
F013500053	EVQLVESGGGLVQPGGSLRLSCAASGRTFSSYAMGWFRQAPGKER EFVAAISWSGGSTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDT AVYYCAASPIPYGSLLRRRNNDYWGQGLTVTVSSGGGGSGGGGS GGGGSGGGGSGGGGSGGGGSGGGGSEVQLVESGGGLVQPGNSL RLSCAASGFTFSSFGMSWVRQAPGKGLEWVSSISGSGSDTLYADSV KGRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLRSSQGLTVT VSSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSEVQLV ESGGGLVQPGGSLRLSCAASGGTFSSYAMGWFRQAPGKEREFVAA IRRSGRRTYYADSVKGRFTISRDN SKNTVYLQMNSLRPEDTAVYYCA AARRVRSSTRYNTGTWWWEYWGQGLTVTVSS <u>GAAEQKLISEE</u> <u>DLNGAAHHHHHH</u>	47

(Note: The molecules described in this Table V include a myc-hexa-histidine tag (underlined), to allow for easier purification of the recombinantly expressed polypeptides; this tag is not required for - nor does it usually interfere with - the binding of the molecules to their targets)

5

Example 6: Sequence optimization of VHHs and VHH constructs

Sequence optimization is a process in which the parental sequence is mutated to make it more identical to the human IGHV3-IGHJ germline consensus sequence. For example, specific amino acids in the framework regions (with the exception of the so-called hallmark residues) are exchanged for their human counterparts, in a way that the protein structure, activity and stability shall be preserved.

15

These mutations may be categorized as follows:

1. Standard: Sequence optimization of these positions is not expected to dramatically change the stability or activity or affinity of the VHH and they are therefore altered all at once, yielding a basic variant.
- 5 2. Unique: It is not known if sequence optimization of these positions affects the stability or activity or affinity of the VHH and therefore they are investigated on an individual basis on top of the basic variant.

10 Hallmark residues are known to be critical for the stability, activity and affinity of the VHH and are therefore not mutated.

In addition, the amino acids present in the CDRs for which there is experimental evidence that they are sensitive to post-translational modifications (PTM) were altered in such a way that the PTM site is inactivated while the protein structure,
15 activity and stability shall remain intact. The most common post-translational modifications described for antibodies and VHHs are listed in Table VI below. The sensitivity of the VHHs for post-translational modifications was analysed in accelerated stress studies, applying several standard conditions including H₂O₂ treatment to analyze methionine oxidation, high temperature, high pH and long
20 storage to study asparagine deamidation and aspartate isomerization. The percentage of oxidation, deamidation and isomerization was measured according to standard procedures and compared to the reference samples (VHHs stored at -20°C). Whole protein analysis in Reverse Phase Chromatography (RPC) and peptide mapping using Mass Spectrometry (MS) were performed to identify
25 potentially sensitive residues. In case of post-translational modifications observed in the VHHs after stress test, the corresponding amino acid(s) were mutated.

Table VI: Potential post-translational modifications, and motifs potentially triggering them

Motif	Modification
M	Met oxidation
N-S/G/H/N/A	Asn deamidation
D-S/G/H	Asp isomerisation
N-X-S/T-X (X ≠ P)	Asn glycosylation
Q/E	pyroglutamate

5 As a result, several mutations were introduced in the constructs set out above, resulting i.a. in the three constructs shown in Table III above, which were chosen for further *in vitro* and *in vivo* characterization, as set out in the Examples further below.

10 **Example 7: *In vitro* characterization of three half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs; Comparison to other LRP6 binding molecules**

After VHH sequence optimization, three half-life extended biparatopic
15 LRP5/LRP6 cross-reactive VHH constructs were recombinantly expressed and purified, and were characterized by using several functional and biophysical assays as described below.

7.1 FACS binding assays

20

Binding to human LRP5 and LRP6 was determined on cells by FACS analysis, as reported in Figures 3A and 3B. In particular, binding to human LRP5 was tested on HEK293 cells with stable overexpression of human LRP5. For human LRP6

binding, HEK293 cells with stable overexpression of human LRP6 were used. The cells were incubated with the LRP5 and LRP6 binder dilutions (1:5 serial dilution of the binders corresponding to the final concentrations indicated in Figure 3A and Figure 3B) for 1.5 hours at 4°C on a plate shaker. After washing the cells 5 times with FACS buffer (1x PBS (Invitrogen cat.no.141190-094) + 10% FBS (Sigma cat.no. F7524) + 0,05% sodium azide), they were incubated for 1 hour at 4°C with a polyclonal mouse antibody which binds to the framework regions of the VHHs. After washing the cells 3 times with FACS buffer, the cells were incubated for 1 hour at 4°C with the labelled secondary antibody (anti-mouse PE (115-116-071), followed by 3 times washing step with FACS buffer. Fluorescence was measured using FACS Array (BD). Binding to human LRP5 and LRP6 corresponds to ≥ 600 MCF values at the highest concentration tested. Negative control consisted of a non-targeting binder (VHH construct that binds to a bacterial protein which is not expressed in HEK293 cells). As shown in Figure 3A and Figure 3B, respectively, binding to human LRP5 and LRP6 corresponds to ≥ 600 MCF and to ≥ 1600 MCF values, respectively, at the highest tested concentrations of the three half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs. These data confirm that the formatted, biparatopic, and sequence-optimized binding molecules bind to both human LRP5 and human LRP6 in their native conformation in a cellular assay system. EC₅₀ values of binding to hLRP5 and hLRP6 are reported in Table VII below.

Table VII: EC₅₀ values of binding to human LRP5 and LRP6 determined by FACS binding assays

FACS based binding assay	F013500571	F013500575	F013500720
hLRP5, EC ₅₀ (nM)	0.01	0.001	0.02
hLRP6, EC ₅₀ (nM)	0.02	0.01	0.02

7.2 FACS - DKK1 competition assay

Potency and efficacy of the three LRP5/LRP6 cross-reactive VHH constructs were further analysed using the FACS-based DKK1 competition assay, as described in Example 4.1. HEK293 cells with stable overexpression of human LRP5 or human LRP6 were incubated with serial dilution of the LRP5/LRP6 cross-reactive VHH constructs (1:5 serial dilution corresponding to the final concentrations indicated in Figure 4A and Figure 4B). The LRP5/LRP6 cross-reactive VHHs competed with human DKK1 for binding to HEK293 cells overexpressing human LRP5, as well as for binding to HEK293 cells overexpressing human LRP6, as shown in Figures 4A and 4B, respectively. Complete inhibition of DKK1 binding was achieved at the highest tested concentrations (≥ 10 nM) and corresponded to MCF values ≤ 60 . In contrast, an LRP5 specific VHH would compete with human DKK1 for binding to HEK293 cells overexpressing human LRP5 but not, or with very low potency (> 200 nM) for binding to HEK293 cells overexpressing human LRP6 (and the same would apply, vice versa, to LRP6 specific VHHs). As a result of this experiment, it could be shown that the present LRP5/LRP6 cross-reactive VHHs competed with human DKK1 for binding to HEK293 cells overexpressing human LRP5, as well as to those overexpressing human LRP6 (i.e. decrease in MCF value with increasing concentration of the binder with complete inhibition of DKK1 binding corresponded to ≤ 60 MCF values at the highest concentration tested). IC₅₀ values of DKK1 competition for binding to hLRP5 and hLRP6 of the three LRP5/LRP6 cross-reactive VHH constructs are reported in Table VIII below. These data confirmed the binding of the three LRP5/LRP6 cross-reactive VHH constructs to human LRP5 and human LRP6 and showed very similar affinity (defined here by the IC₅₀ potency values in the DKK1 competition assay) between the two receptors. Furthermore, the data strengthen the notion that the formatted, biparatopic, and sequence-optimized binding molecules are binding to human LRP5 as well as to human LRP6 in their native conformation.

Table VIII: IC₅₀ values of DKK1 competition for binding to human LRP5 and LRP6 determined by FACS binding assays

DKK1 competition assay	F013500571	F013500575	F013500720
hLRP5, IC ₅₀ (nM)	0.03	0.03	0.3
hLRP6, IC ₅₀ (nM)	0.01	0.1	0.2

5

7.3 Combined Wnt1 and Wnt3a reporter assay

Potency and efficacy of the formatted, biparatopic, and sequence-optimized binding molecules were analysed using a combined Wnt1 and Wnt3a reporter assay to allow functional testing of both Wnt1 and Wnt3a blockers in the same assay. The combined Wnt1 and Wnt3a reporter assay is based on the assay described in Example 4.4 with the following changes in the protocol. 1E06 / ml cells with overexpression of Wnt1, seeded into a 384 well tissue culture plate were treated with recombinant human Wnt3a (rec.human Wnt3a: R&D #5036-WN/CF) at the final concentration of 500ng/ml and then the cells were incubated overnight at 37 °C. The following day, serial dilutions of various LRP5/LRP6 biparatopic cross-reactive VHH solutions were prepared and added to the cells in the presence of LiCl at the final concentration of 10nM. DKK1, as positive control, was added to the cells at the final concentration of 200nM. DKK1 treatment resulted in a complete inhibition of the combined Wnt1 and Wnt3a pathway, and therefore a complete inhibition of beta-lactamase enzymatic activity. The cells were incubated overnight at 37 °C. The following day, beta-lactamase enzymatic activity was measured according to the manufacturer's instructions. As reported in Example 4.4., for fluorescence emission, values at 460 nm and 530 nm were obtained using a standard fluorescence plate reader and the 460/530 nm emission ratios plotted for the indicated treatment. The value of the ratio of fluorescence [460/535nm] reported as "baseline" in Figure 5A corresponds to complete inhibition of Wnt1 and Wnt3a pathway, determined by the treatment with the positive control (DKK1; 200nM final concentration). The value of the ratio

25

of fluorescence [460/535nm] reported as “Wnt1” corresponds to activation of Wnt1 pathway only, determined from the Wnt1 overexpressing cells (i.e. without treatment with recombinant human Wnt3a). The value of the ratio of fluorescence [460/535nm] reported as “Wnt1 + Wnt3a” corresponds to the combined activation of Wnt1 and Wnt3a pathways, determined by treatment of the Wnt1 overexpressing cells with recombinant human Wnt3a. As shown in Figure 5A, complete inhibition (i.e. ratio of fluorescence [460/535nm] corresponding to the baseline) is achieved by treatment with the three LRP5/LRP6 cross-reactive, formatted, biparatopic, and sequence-optimized binding molecules. Furthermore, high potency is also reported, as shown in Table IX below by the IC₅₀ values.

Table IX: IC₅₀ values of Wnt1 and Wnt3a pathway inhibition in combined Wnt1 and Wnt3a reporter assay

Combined Wnt1 and Wnt3a reported assay	F013500571	F013500575	F013500720
IC ₅₀ (nM)	0.05	0.2	0.06

Next, potency and efficacy of the LRP5/LRP6 cross-reactive, formatted, biparatopic, and sequence-optimized binding molecules were compared to previously disclosed LRP6 binding molecules reported in WO2011/138391 and WO2011/119661:

In WO2011/138391, multivalent antibodies binding to LRP6 and inhibiting both propeller 1 (e.g., Wnt1) and propeller 3 (e.g., Wnt 3) ligand interaction were disclosed. These multivalent LRP6 binding antibodies are biparatopic LRP6 binding molecules consisting of an IgG antibody as a first receptor binding domain and of an scFv fragment as a second receptor binding domain, where the IgG antibody and scFv fragment are linked together by a linker. In WO2011/138391 it is reported that all the LRP6 binding molecules have roughly the same potency in Wnt1 and Wnt3a reporter assay (Fig. 18 of WO2011/138391). Therefore, any of those multivalent LRP6 binding molecules

could be chosen for comparative experiments. Thus, it was decided to use the "901" construct (referred to as MOR08168IgG1LALA 6475 scfv; also shown in Fig. 27 of WO2011/138391) as a first comparative compound.

5 Derivatives of this "901" construct are shown in WO2013/067355. Specifically, compounds named 801T and 802T are disclosed (cf. disclosure on p.132 of the specification), both having two LRP6 binding scFv domains, plus a half-life extending moiety. As 801T and 802T appear to have the same in vitro potency and biophysical characteristics, only one of them - variant 802T - was included in
10 the experiments described in the following.

In WO2011/119661, bispecific antibodies binding to LRP6 and inhibiting signaling by multiple Wnt isoforms were disclosed. These bispecific anti-LRP6 antibodies bind to two different regions of LRP6 and inhibit signaling induced by Wnt
15 isoforms, among them Wnt1 and Wnt3a. Knobs-into-holes engineering (Atwell *et al.* "Stable heterodimers from remodeling the domain interface of a homodimer using a phage display library". *J Mol Biol.* 1997; 270(1):26-35) was used to build these bispecific anti-LRP6 antibodies. Example 11 of WO2011/119661 discloses an IgG hybrid with YW211.31.62 and YW210.09 heavy chain heterodimers. Thus,
20 for comparative purposes, two bispecific IgG hybrid antibodies with YW211.31.62 and YW210.09 heavy chain heterodimers were generated with the knobs-into-holes engineering technology, i.e. with amino acid changes engineered to create a knob on the CH3 of the heavy chain of YW210.09 and a hole on the CH3 of the heavy chain of YW211.31.62 or vice versa, and referred here as Knob HC
25 YW210.09 and Knob HC YW211.31.62, respectively. These two constructs were characterized in Wnt1 and Wnt3a reporter assays according to Example 4.4. As expected, the two bispecific IgG hybrids with YW211.31.62 and YW210.09 heavy chain heterodimers showed similar potencies in Wnt1 and Wnt3a assays, as reported in Table X below. Thus, for the comparative examples shown further
30 below, Knob HC YW210.09 was selected as a second comparative compound.

Table X: IC₅₀ values of Wnt1 and Wnt3a pathway inhibition in Wnt1 and Wnt3a reporter assays

Wnt reporter assay	Knob HC YW210.09	Knob HC YW211.31.62
Wnt1, IC ₅₀ (nM)	4.8	4.6
Wnt3a, IC ₅₀ (nM)	0.7	0.9

Potency and efficacy of the half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs were compared to MOR08168IgG1LALA 6475 scfv biparatopic LRP6 binding molecule, to Knob HC YW210.09 bispecific anti-LRP6 molecule and to 802T bispecific anti-LRP6 molecule using the combined Wnt1 and Wnt3a reporter assay. As shown in Figure 5B, complete inhibition (i.e. ratio of fluorescence [460/535nm] corresponding to the baseline in Figure 5B) was achieved by treatment with Knob HC YW210.09 (the bispecific IgG hybrid antibody with YW211.31.62 and YW210.09 heavy chain heterodimers) similar to the LRP5/LRP6 cross-reactive, formatted, biparatopic, and sequence-optimized binding molecule F013500571. However, F013500571 showed higher potency, as reported in Table XI below. Instead, MOR08168IgG1LALA 6475 scfv and 802T biparatopic LRP6 binding molecules showed lack of complete Wnt1 and Wnt3a inhibition (i.e. ratio of fluorescence [460/535nm] significantly higher than the baseline in Figure 5B and in Figure 5C, respectively). These data indicate that both MOR08168IgG1LALA 6475 scfv and 802T biparatopic LRP6 binding molecules have significant lower efficacy in Wnt1 and Wnt3a pathway inhibition when compared to F013500571. Therefore, Knob HC YW210.09 was selected as a comparative compound for *in vivo* experiments as described further below (Example 9; *in vivo* efficacy)

Table XI: IC₅₀ values of Wnt1 and Wnt3a pathway inhibition in combined Wnt1 and Wnt3a reporter assay

Combined Wnt1 and Wnt3a reported assay	F013500571	Knob HC YW210.09	MOR08168IgG1LALA 6475 scfv	802T
IC ₅₀ (nM)	0.05	1.15	0.19	0.01

Example 8: Effects of three half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs on Wnt Signaling and viability in cancer cell lines

The ability of the half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs to inhibit active Wnt signaling was further characterized using cancer cell lines with active Wnt signaling, as previously described (Bafico *et al.* "An autocrine mechanism for constitutive Wnt pathway activation in human cancer cells". *Cancer Cell* 2004;6(5):497-506; DeAlmeida *et al.* "The soluble wnt receptor Frizzled8CRD-hFc inhibits the growth of teratocarcinomas in vivo". *Cancer Res.* 2007;67(11):5371-9; Akiri *et al.* "Wnt pathway aberrations including autocrine Wnt activation occur at high frequency in human non-small-cell lung carcinoma". *Oncogene.* 2009; 28(21):2163-72). Briefly, the cancer cell lines with active Wnt signaling, PA-1 and PA-TU-8988S, were seeded in 12-well plates and treated for two days with the LRP5/LRP6 cross-reactive VHH constructs at the final concentration of 1 uM. The ability of inhibiting Wnt signaling was detected by inhibition of mRNA expression of Axin2, the endogenous Wnt target gene. qPCR expression analysis was performed using standard RNA techniques: RNA isolation was performed using the QIAGEN RNeasy Mini Kit according to the protocol provided by QIAGEN; cDNA synthesis using SuperScript VILO cDNA Synthesis Kit (Invitrogen, Cat. No. 11754050) and qPCR using TaqMan Gene Expression Assay with Axin2 TaqMan primers/probe (Hs00610344_m1 AXIN2 FAM, Life Technologies) and with eukaryotic 18s endogenous control VIC-MGB (4319413E-1307061, Applied Biosystems).

As shown in Figure 6A, both PA-TU8988S and PA-1 cancer cells treated with three half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs showed significantly reduced Axin2 relative mRNA levels (i.e. normalized to the endogenous control) when compared to untreated (Control) cells. These data demonstrated the ability of the half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs to inhibit Wnt signaling in cancer cell lines with active Wnt signaling. Furthermore, the effect of Wnt signaling blockade on cell viability was investigated in PA-TU8988S and YAPC cancer cell lines, whose proliferation was previously reported to be dependent on active Wnt signaling (Jiang *et al.*

“Inactivating mutations of RNF43 confer Wnt dependency in pancreatic ductal adenocarcinoma”. *Proc Natl Acad Sci U S A.* 2013; 110(31):12649-54). Cell viability was measured by performing an Alamar Blue assay (Invitrogen, Cat.# DAL1100) after ten days of treatment with the half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs (final concentration of 1 μ M) or with comparative compound 802T (final concentration 1 μ M). As shown in Figure 6B, PA-TU8988S cancer cells treated with three half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs showed significantly reduced percentage of viable cells cell ($\geq 75\%$ reduction) when compared to untreated (control) cells. Upon treatment with 802T, no effect on cell viability was detected (Figure 6B, diagram on the right-hand side). These data demonstrate the ability of the half-life extended biparatopic LRP5/LRP6 cross-reactive VHH constructs to inhibit cell proliferation of cancer cells which are dependent on active Wnt signaling, and also show a superior effect of these constructs when compared to 802T.

Furthermore, dose dependency of cell viability was assessed for F013500571 in comparison to 802T, in PA-TU8988S (Figure 6C) and YAPC (Figure 6D) cancer cell lines, as described above. Dose dependent decrease in cell viability was detected upon treatment with F013500571. In contrast, no effect on cell viability was shown upon treatment with the comparative compound 802T in both PA-TU8988S and YAPC cancer cell lines. These data show that the half-life

extended biparatopic LRP5/LRP6 cross-reactive VHH constructs have a superior effect when compared to 802T.

Example 9: *In vivo* Efficacy

5

The LRP5/LRP6 cross-reactive biparatopic half-life extended VHH constructs / binding molecules were further characterized *in vivo* in a Wnt driven tumor model. Experiments were conducted to determine if these binding molecules inhibit tumor growth *in vivo*. Efficacy of the comparative compound Knob HC YW210.09 was also determined using the same Wnt driven tumor model. Transgenic expression of the Wnt ligands using a mouse mammary tumor virus LTR enhancer (MMTV promoter) leads in mice to extensive ductal hyperplasia followed by mammary adenocarcinomas in transgenic (TG) mice by 6 months of age. These mammary tumors are driven by glucocorticoid induced overexpression of Wnt ligands and have characteristics similar to TNBC tumors, including expression of epithelial and mesenchymal markers (basal-like phenotype) and active Wnt signaling as assessed by intracellular beta-catenin localization. In particular, mammary tumors derived from MMTV-Wnt-1 transgenic mice are Wnt1 dependent. Indeed, blocking of Wnt activity using a soluble Wnt receptor comprising the Frizzled8 cysteine-rich domain (CRD) fused to the human Fc domain (F8CRDhFc) (DeAlmeida *et al.* "The soluble wnt receptor Frizzled8CRD-hFc inhibits the growth of teratocarcinomas *in vivo*". *Cancer Res.* 2007;67(11):5371-9) was reported to inhibit tumor growth *in vivo*. Therefore, tumors isolated from MMTV-Wnt1 transgenic mice were passaged subcutaneously as tumor pieces in nude mice for 2 to 5 passages prior to initiation of the efficacy experiment. Between 14 to 21 days post-implant, when tumors reached a mean volume of approximately 150 to 250 mm³, mice were randomized into groups with 7 mice per group and dosed *i.v.* with the compounds. The LRP5/LRP6 cross-reactive biparatopic half-life extended VHH constructs were administered to the mice *i.v.* twice per week, with the dosages shown in Figure 7A for F013500571 and in Figure 7B for F013500720. Comparative compound Knob HC YW210.09 was also dosed *i.v.* twice per week,

10

15

20

25

30

but at higher doses, i.e. 30 and 45mg/kg (Figure 7C), due to the data obtained with this compound in the *in vitro* experiments explained above. Tumor volume and body weight were monitored during the efficacy experiment and the median tumor volumes are reported in Figures 7A to 7C. Tumor growth inhibition (TGI) was determined at the end of the efficacy experiment. In particular, TGI was determined for each treatment group as compared to the control group (treatment of mice with histidine buffer - 20mM Histidine pH6.5 buffer - in the experiment shown in Figures 7A and 7B, or with citrate buffer for the experiment shown in Figure 7C). Furthermore, gastro-intestinal (GI) histo-pathological analysis (via H&E staining of sections of the GI tract from the duodenum to the rectum) was performed at the end of the efficacy experiment to evaluate potential toxicity of the LRP5 and LRP6 antagonists. Tumor growth inhibition (TGI), outcome of the GI histo-pathological analysis at the end of the *in vivo* efficacy study, mortality which corresponds to the number of mice that needed to be sacrificed due to significant loss of body weight (>18% loss of body weight compared to start of the efficacy experiment) and number of tumor regressions (tumor volume at the end of the experiment smaller than tumor volume measure at the start of the treatment) are reported for each treatment group in Tables XIIA, XIIB, and XIIC, relating to the experiments and data shown also in Figures 7A, 7B and 7C, respectively.

Table XIIA: *In vivo* efficacy of F013500571 administered *i.v.* twice per week.

Results of this experiment are also shown in Figure 7A

	Dose [mg/kg]	TGI [%]	Regressions [x/7]	Mortality [x/7]	GI Histopathological evaluation
Control	Histidine buffer	-	-	-	-
F013500571	10	128	7	0	No findings
	4	113	7	0	No findings
	2	61	0	0	n.d.

Table XIIB: *In vivo* efficacy of F013500720 administered *i.v.* twice per week.

Results of this experiment are also shown in Figure 7B

	Dose [mg/kg]	TGI [%]	Regressions [x/7]	Mortality [x/7]	GI Histopathological evaluation
Control	Histidine buffer	-	-	-	-
F013500720	1	128	7	0	No findings
	0.4	60	0	0	No findings
	0.2	17	0	0	n.d.

5 **Table XIIC:** *In vivo* efficacy of Knob HC YW210.09 administered *i.v.* twice per week. Results of this experiment are also shown in Figure 7C

<i>i.v.</i>	Dose [mg/kg]	TGI [%]	Regressions [x/7]	Mortality [x/7]	GI Histopathological evaluation
Control	Citrate buffer	-	-	-	-
Knob HC YW210.09	30	80	0	0	-
	45	85	0	0	No findings

As can be taken from Figures 7A to 7C and Tables XIIA to XIIC above, treatment with the LRP5/LRP6 cross-reactive half-life extended biparatopic VHH constructs (F013500571 at 4 and 10 mg/kg and F013500720 at 1mg/kg in a 2x/week
 10 schedule) indeed resulted in tumor regression (i.e. tumor growth inhibition (TGI) > 100% which corresponds to tumor shrinkage; decrease of the tumor volume at the end of the efficacy experiment compared to the tumor volume at the start of the experiment), with no significant body weight changes (< 10%) and no findings

reported after GI histopathological analysis. Importantly, in contrast thereto, no tumor regression could be observed upon treatment with the LRP6 specific binder Knob HC YW210.09, even at the maximum administrable *i.v.* dose/schedule in mice, which correspond to 90 mg/kg with 3x/week schedule.

5

To further investigate the differences in *in vivo* efficacy as observed in the experiment explained above, a further experiment was set up allowing more frequent administration (three times/week) of an even higher dose of the comparative compound to the mice. In brief, in order to achieve such higher exposure, the comparative compound was administered *i.p.*, as indicated in the following Table XIID:

10

Table XIID: *In vivo* efficacy of Knob HC YW210.09 administered *i.p.* twice or three times per week, as indicated.

<i>i.p.</i>	Dose [mg/kg]	Schedule	TGI [%]	Regressions [x/5]	Mortality [x/5]
Control	Citrate buffer	Three times/week	-	-	-
Knob HC YW210.09	45	Twice/ week	55	1	0
	90	Twice/ week	85	0	0
	90	Three times/ week	87	0	0

15

As can be seen from the data shown in Table XIID, also in this setting no significantly stronger effect was obtained in terms of TGI. In other words, these experiments, data and results clearly indicate a higher efficacy of the half-life extended biparatopic VHH constructs when compared to Knob HC YW210.09, and an unprecedented capability of the polypeptides of the invention to not only reduce tumor growth, but to even induce tumor shrinkage. Of course, tumor shrinkage (i.e. tumor regression) is the desired therapeutic effect (i.e. efficacy) for treatment of cancer patients. Indeed, in clinical studies treatments that induce

20

tumor regression resulting in pathological complete response (pCR) positively lead to significant improvement of progression free survival and overall survival in high unmet medical need indications such as in breast cancer.

5 The above comparative examples also show that the LRP5/LRP6 cross-reactive biparatopic half-life extended VHH constructs are not only superior in terms of their binding characteristics, such as affinity or K_D values, but that they have highly advantageous and superior properties in an *in vivo* setting.

10 Next, it was investigated whether compound MOR08168IgG1LALA 6475 scfv may provide for a similar advantageous effect. For this purpose an *in vivo* tolerability study was performed in mice as follows: MOR08168IgG1LALA 6475 scfv compound was administered *i.v.* at 3 mg/kg, twice per week (2qw); the same dose/regimen at which *in vivo* efficacy was detected in a xenograft tumor model
15 in WO2011/138391, as described in Fig. 22 therein. First treatment with MOR08168IgG1LALA 6475 scfv was performed at day 1 and starting from day 6 significant loss of body weight was detected in mice. At day 10, some of the mice treated with MOR08168IgG1LALA 6475 scfv compound showed significant body weight loss (>10%). At day 11 mice were sacrificed and gastro-intestinal (GI)
20 histopathological analysis revealed inflammation with erosion in the colon and in the cecum of the mice. These data suggest that MOR08168IgG1LALA 6475 scfv is not tolerated at the efficacious dose/regimen. Therefore, LRP5/LRP6 cross-reactive biparatopic half-life extended VHH constructs are superior with regard to therapeutic window; i.e. they induce tumor regression with no significant body
25 weight changes (< 10%) and no findings reported after GI histopathological analysis.

Example 10: *In vivo* Wnt pathway inhibition

30 To further characterize the effect of the LRP5/LRP6 cross-reactive biparatopic half-life extended VHH constructs / binding molecules on Wnt signaling, tumors

were isolated at the end of the efficacy experiment, described in Example 9. In particular, tumors were isolated 16 hours after the last injection with the compounds or with the control treatment. Wnt signaling inhibition was determined by reduction of mRNA expression of Axin2 in tumors, analysed as described in Example 8. The fold change of Axin2 mRNA expression relative to the control group is reported in Figure 8A for the *in vivo* efficacy with F013500571 and in Figure 8B for the *in vivo* efficacy with F013500720. Quantification of Axin2 mRNA reduction for each treatment group is reported in Tables XIII A and XIII B below.

Table XIII A: Reduction of Axin2 mRNA expression in tumors by treatment with F013500571. The data refers to Figure 7A and 8A.

	Dose [mg/kg]	TGI [%]	Axin2 reduction (%)
Control	Histidine buffer	-	
F013500571	10	128	83
	4	113	77
	2	61	35

Table XIII B: Reduction of Axin2 mRNA expression in tumors by treatment with F013500720. The data refers to Figure 7B and 8B.

	Dose [mg/kg]	TGI [%]	Axin2 reduction (%)
Control	Histidine buffer	-	
F013500720	1	128	92
	0,4	60	-21
	0,2	17	35

As can be seen from Figures 8A and 8B and Tables XIII A and XIII B, a significant reduction and dose dependent decrease (in particular for treatment with F013500571) of Axin2 mRNA expression was observed in tumors treated with the LRP5/LRP6 cross-reactive binding molecules, compared to the control group.

5 These results suggest that the LRP5/LRP6 cross-reactive binding molecules are indeed able to inhibit tumor growth by suppressing Wnt signaling in the tumor cells.

Example 11: Industrial manufacturing process

10
11.1 *Fermentation*: Any of the polypeptides set out in Tables III and V above can be expressed in the cytoplasm of different E. coli strains like W3110, TG1, BL21, BL21(DE3), HMS174, HMS174(DE3), MM294 under control of an inducible promoter. This promoter can be chosen from lacUV5, tac, T7, trp, T5, araB. The
15 cultivation media are preferably fully defined according to Wilms *et al.*, 2001 (Wilms, B., Hauck, A., Reuss, M., Syltschik, C., Mattes, R., Siemann, M., and Altenbuchner, J.: High-Cell-Density Fermentation for Production of L-N-Carbamoylase Using an Expression System Based on the Escherichia coli rhaBAD Promoter. *Biotechnology and Bioengineering*, 73: 95-103 (2001)), DeLisa
20 *et al.*, 1999 (DeLisa, M. P., Li, J. C., Rao, G., Weigand, W. A., and Bentley, W. E.: Monitoring GFP-operon fusion protein expression during high cell density cultivation of Escherichia coli using an on-line optical sensor. *Biotechnology and Bioengineering*, 65: 54–64.(1999)) or equivalent. However, supplementation of the medium with amino acids like isoleucine, leucine, lysine, methionine,
25 phenylalanine, threonine, tryptophan and valin or complex media components such as soy peptone or yeast extract may be beneficial. The process for fermentation is performed in a fed-batch mode. Conditions: Temperature 30 – 40 °C, pH 6 – 7.5, dissolved oxygen is kept above 20%. After consumption of the initial C-source the culture is fed with the feed media stated above (or equivalent).
30 When a dry cell weight in the fermenter of 40 to 90 g/L is reached the culture is induced with an appropriate inducer corresponding to the used promoter system

(e.g. IPTG, lactose, arabinose). The induction can either be performed as a pulsed full induction or as a partial induction by feeding the respective inducer into the fermenter over a prolonged time. The production phase should last 4 hours at least. The cells are recovered by centrifugation in bowl centrifuges, tubular bowl centrifuges or disc stack centrifuges, the culture supernatant is discarded.

11.2 Purification: The E. coli cell mass is resuspended in 6- to 8-fold amount of lysis buffer (phosphate or Tris buffer, pH 7-8.5). Cell lysis is preferably performed by high pressure homogenization followed by removing of the cell debris by centrifugation in bowl, tubular bowl or disc stack centrifuges. Supernatant containing the target protein is optionally filtrated using a 0.22-10 µm filter and separated via cation exchange chromatography (e.g. Toyopearl MegaCap® II SP-550EC, Toyopearl GigaCap S-650M, SP Sepharose BB, SP Sepharose FF or S HyperCel™) at pH 7-8.5. Elution is performed by a linear increasing NaCl gradient at pH 7-8.5. Fractions containing the target protein are pooled and subsequently incubated with 5-10 mM DTT in order to prevent dimerization or aggregation mediated by free cysteine residues. After further addition of 0.8-1 M ammonium sulfate or 2-3 M NaCl, solution is separated via hydrophilic interaction chromatography (e.g. Phenyl Sepharose HP, Phenyl Sepharose FF, Butyl Sepharose HP, Butyl Sephrose FF, Butyl Toyopearl 650 (S,M,C), Phenyl Toyopearl 650 (S,M,C)) at pH 7-8.5. Elution is carried out at pH 7-8.5 by a linear decreasing ammonium sulfate or NaCl gradient in presence of 5 mM DTT. Fractions containing the target protein with a purity level of minimally 90% are pooled and desalted by diafiltration in presence of 5 mM DTT followed by concentration to approximately 5 mg/ml. Subsequent refolding is performed by diluting the protein solution 1:5-1:20 with 50 mM Tris, 150 mM NaCl, 4 mM Cystamin, 10 mM CHAPS at pH 8.5 to a final protein concentration of 0.25-1 mg/ml. Refolding solution is incubated under stirring for 12-36 h at room temperature and then separated by cation exchange chromatography (e.g. SP Sepharose FF, SP Sepharose HP, Toyopearl SP-650 (S, M, C)) at pH 7-8.5.

Elution is performed by a linear increasing NaCl gradient at pH 7-8.5. Fractions containing monomeric target protein are pooled and formulated in 25 mM Na-phosphate, 220 mM endotoxin free trehalose, pH 7.5 via diafiltration. The solution is sterilized by filtration and stored at 2 to 8 °C.

5

Example 12: Pharmaceutical formulation for s.c. administration

Any of the above biparatopic polypeptide constructs of the invention can be selected for the manufacture of a pharmaceutical formulation for subcutaneous application having a composition as follows:

10

Drug substance: 100 mg/ml (1 to 3 nmol/ml)

Acetate buffer: 25 mM

Trehalose: 220 mM

Tween-20: 0.02 %

15

Drug substance is formulated in a solution having the above composition, sterilized and stored at 2 to 8 °C.

Example 13: Pharmaceutical use in humans

20

The solution as prepared in Example 11.2 above is applied to a patient in need thereof, such as a human being suffering from a cancer sensitive to Wnt signaling inhibitors, by intravenous infusion (dosage of 100 to 200 mg) every two to four weeks.

25

Example 14: Effect of Wnt3a-signaling inhibition on pro-inflammatory cytokine release by dendritic cells in an ex-vivo assay

PBMCs were obtained from healthy donors with informed consent. Human monocyte-derived dendritic cells (Mo-DCs) were generated as follows: PBMCs were cultured in X-VIVO medium supplemented with 50 ng/mL GM-CSF and 50 ng/mL IL-4. After 24 h of culture, the supernatant was carefully removed and

30

replaced with X-VIVO medium supplemented with the same GM-CSF and IL-4.

On the fourth day, the supernatant was carefully removed and replaced with

X-VIVO medium in the presence of LPS only or in combination with human

Wnt3a or with human Wnt3a and LRP5/LRP6 cross-reactive binding molecules.

5 On the following day, the supernatants were collected and subjected to analysis of TNF-alpha through ELISA according to the manufacturer's instructions. As

previously reported (Oderup *et al.* "Canonical and noncanonical Wnt proteins program dendritic cell responses for tolerance". *J Immunol.* 2013;190(12):

6126-34), and as shown in Figure 9A, Wnt3a directly inhibits pro-inflammatory

10 cytokine secretion (i.e. TNF-alpha release) by differentiated dendritic cells (DCs).

Wnt3a driven suppression of TNF-alpha release from DCs was restored by

addition of LRP5/LRP6 cross-reactive binding molecules.

These data show that the formatted, biparatopic, and sequence-optimized binding

15 molecules are able to restore TNFalpha secretion by Wnt3a treated dendritic

cells, thereby suppressing the Wnt inhibitory effect on dendritic cells.

It is important to notice that blocking the Wnt pathway in dendritic cells in the

tumor microenvironment might represents a potential therapeutic approach

20 towards breaking tumor-mediated immune suppression and augmenting anti-tumor immunity.

To investigate the effects of DCs on T cells (effector T cells), DCs pre-treated

with Wnt3a with or without LRP5/LRP6 cross-reactive binding molecules were co-

25 cultured with T cells, isolated from PBMCs, as previously described (Oderup et al.

"Canonical and noncanonical Wnt proteins program dendritic cell responses for

tolerance". *J Immunol.* 2013;190(12): 6126-34). After 3 days of DC/T cells co-

culture, the supernatants were collected and subjected to analysis of IFN gamma through ELISA, according to the manufacturer's instructions.

IFNgamma secretion is a marker of T cell activation. As shown in Figure 9B, Wnt3a mediated DC inhibition leads to reduced IFNgamma secretion by T cells (inhibition of T cell function), which is completely restored by treatment with LRP5/LRP6 cross-reactive binding molecules.

5

In summary, these data show that LRP5/LRP6 cross-reactive binding molecules suppress the Wnt inhibitory effect on dendritic cells, leading to restoration of T cell function.

10

It is known that continuous activation/stimulation of T cells induces terminal differentiation, resulting in an exhausted T cell phenotype, a progressive loss of T-cell function. Therefore, it is envisaged that the effect of LRP5/LRP6 cross-reactive binding molecules on T cells, mediated by activation of DCs, might be limited by T cell exhaustion. Combination treatments, combining the

15

administration of LRP5/LRP6 cross-reactive binding molecules with the administration of an immune checkpoint inhibitor which blocks T cell exhaustion, are therefore expected to help to activate and maintain T cell function, thereby changing the tumor microenvironment, and thereby supporting the therapeutic effect of the molecules of the invention.

20

Claims:

1. A polypeptide which specifically binds to LRP5 or LRP6, comprising
 - a first immunoglobulin single variable domain selected from the group of immunoglobulin single variable domains (i) to (iii) defined by comprising the following CDR sequences:

(i):

CDR1: TYTVG (= SEQ ID NO:1)

CDR2: AIRRRGSSTYYADSVKG (= SEQ ID NO:2)

CDR3: DTRTVALLQYRYDY (= SEQ ID NO:3)

(ii):

CDR1: SYAMG (= SEQ ID NO:4)

CDR2: AIRRSGRRTYYADSVKG (= SEQ ID NO:5)

CDR3: ARRVRSSSTRYNTGTWWWEY (= SEQ ID NO:6)

(iii):

CDR1: RYTMG (= SEQ ID NO:7)

CDR2: AIVRSGGSTYYADSVKG (= SEQ ID NO:8)

CDR3: DRRGRGENYILLYSSGRYEY (= SEQ ID NO:9),

and

- a second immunoglobulin single variable domain selected from the group of immunoglobulin single variable domains (iv) and (v) defined by comprising the following CDR sequences:

(iv):

CDR1: SYAMG (= SEQ ID NO:10)

CDR2: AISWSGGSTYYADSVKG (= SEQ ID NO:11)

CDR3: SPIPYGSLRRRRNNYDY (= SEQ ID NO:12)

(v):

CDR1: SYAMG (= SEQ ID NO:13)

CDR2: AISWRSGSTYYADSVKG (= SEQ ID NO:14)

CDR3: DPRGYGVAYVSAYYEY (= SEQ ID NO:15).

2. The polypeptide of claim 1 wherein
- said first immunoglobulin single variable domain comprises the following CDR sequences:
CDR1: TYTVG (= SEQ ID NO:1)
CDR2: AIRRRGSSTYYADSVKG (= SEQ ID NO:2)
CDR3: DTRTVALLQYRYDY (= SEQ ID NO:3)
and wherein
 - said second immunoglobulin single variable domain comprises the following CDR sequences:
CDR1: SYAMG (= SEQ ID NO:10)
CDR2: AISWSGGSTYYADSVKG (= SEQ ID NO:11)
CDR3: SPIPYGSLLRRRNNDY (= SEQ ID NO:12).
3. The polypeptide of claim 1 wherein
- said first immunoglobulin single variable domain comprises the following CDR sequences:
CDR1: SYAMG (= SEQ ID NO:4)
CDR2: AIRRSGRRYYADSVKG (= SEQ ID NO:5)
CDR3: ARRVRSSSTRYNTGTWWWEY (= SEQ ID NO:6)
and wherein
 - said second immunoglobulin single variable domain comprises the following CDR sequences:
CDR1: SYAMG (= SEQ ID NO:13)
CDR2: AISWRSGSTYYADSVKG (= SEQ ID NO:14)
CDR3: DPRGYGVAYVSAYYDY (= SEQ ID NO:15).
4. The polypeptide of claim 1 wherein
- said first immunoglobulin single variable domain comprises the following CDR sequences:
CDR1: RYTMG (= SEQ ID NO:7)
CDR2: AIVRSGGSTYYADSVKG (= SEQ ID NO:8)
CDR3: DRRGRGENYILLYSSGRYDY (= SEQ ID NO:9)
and wherein

- said second immunoglobulin single variable domain comprises the following CDR sequences:

CDR1: SYAMG (= SEQ ID NO:13)

CDR2: AISWRSGSTYYADSVKG (= SEQ ID NO:14)

CDR3: DPRGYGVAYVSAYYEY (= SEQ ID NO:15).

5. The polypeptide of any of claims 1 to 4, wherein said immunoglobulin single variable domains are VHH domains, and preferably humanized VHH domains.
6. The polypeptide of claim 1 or claim 2, wherein
 - said first immunoglobulin single variable domain has the amino acid sequence as shown in SEQ ID NO:19 and
 - said second immunoglobulin single variable domain has the amino acid sequence as shown in SEQ ID NO:22.
7. The polypeptide of claim 1 or claim 3, wherein
 - said first immunoglobulin single variable domain has the amino acid sequence as shown in SEQ ID NO:20 and
 - said second immunoglobulin single variable domain has the amino acid sequence as shown in SEQ ID NO:23.
8. The polypeptide of claim 1 or claim 4, wherein
 - said first immunoglobulin single variable domain has the amino acid sequence as shown in SEQ ID NO:21 and
 - said second immunoglobulin single variable domain has the amino acid sequence as shown in SEQ ID NO:23.
9. The polypeptide according to any of claims 1 to 8, wherein said first and said second immunoglobulin single variable domains are covalently linked by a linker peptide, wherein said linker peptide optionally comprises or consists of a third immunoglobulin single variable domain.

10. The polypeptide according to claim 9, wherein said third immunoglobulin single variable domain is an albumin binding immunoglobulin single variable domain, and preferably the Alb11 domain, defined by SEQ ID NO:24.
11. A polypeptide selected from the group of polypeptides comprising or consisting of SEQ ID NO:25, SEQ ID NO:26, and SEQ ID NO:27.
12. An expression vector comprising a nucleic acid encoding a polypeptide according to any of claims 1 to 11.
13. A host cell carrying an expression vector according to claim 12.
14. A method of manufacturing a polypeptide according to any of claims 1 to 11, comprising the steps of
 - culturing a host cell according to claim 13 under conditions that allow expression of a polypeptide according to any of claims 1 to 11; and
 - recovering and optionally purifying said polypeptide.
15. A pharmaceutical composition comprising (i) as the active ingredient, the polypeptide according to any of claims 1 to 11, and (ii) a pharmaceutically acceptable carrier, and optionally (iii) a diluent, excipient, adjuvant and/or stabilizer.
16. The polypeptide according to any of claims 1 to 11 for use as a medicament in a method for the treatment, prevention or alleviation of a disease, disorder or condition in a human being or an animal, preferably for use in a method (a) for the treatment of cancer, such as breast cancer, lung cancer, pancreatic cancer, colorectal cancer, sarcomas, ovarian cancer or hepatocellular carcinoma, or (b) for the treatment of idiopathic pulmonary disease, or (c) for the treatment of a retinopathy caused by abnormal Wnt signaling.

17. A polypeptide of any of claims 1 to 11 for use in the treatment of cancer in combination with an immune checkpoint inhibitor selected from the group consisting of anti-PD1 antibodies, anti-PDL1 antibodies, anti-CTLA4 antibodies, anti-BTLA antibodies, anti-LAG3 antibodies, and anti-TIM3 antibodies, or in combination with a cancer vaccine.