

1/182




Figure 1

2/182




Figure 2

Multiplexed PCR & RT-PCR; LDR w/Tags, 24 samples.
Distribute LDR products from each sample into 16 wells.




Figure 3

Multiplexed PCR & RT-PCR; LDR w/Tags, 24 samples





Figure 4

(Multiplexed PCR & RT-PCR; LDR w/Tags, 24 samples)



Real-Time PCR results, 24 samples

Figure 5

6/182




Figure 6

7/182

Distribute 1 sample into 24 tubes (wells), then Multiplexed PCR; LDR w/Tags
Distribute LDR products (24 x 16 wells).




Figure 7

Distribute 1 sample into 24 tubes (wells), then Multiplexed PCR; LDR w/Tags





Figure 8

9/182

(Distribute 1 sample into 24 tubes (wells), then Multiplexed PCR; LDR w/Tags)



Real-Time PCR results, 1 sample
w/ "Pixel" Enumeration

Figure 9

10/182




Figure 10

11/182

Dilute 1 sample into 24 tubes (wells), then Multiplexed RT-PCR; LDR w/Tags
Distribute LDR products (24 x 16 wells).




Figure 11

12/182

Dilute 1 sample into 24 tubes (wells), then Multiplexed RT-PCR; LDR w/Tags





Figure 12

(Dilute 1 sample into 24 tubes (wells), then Multiplexed RT-PCR; LDR w/Tags)



Real-Time PCR results, 1 sample
w/ "Pixel" Enumeration

Figure 13

14/182




Figure 14

15/182

Multiplexed PCR & RT-PCR; 24 samples.
Distribute PCR products from each sample into 16 wells.





Figure 15

Multiplexed PCR & RT-PCR; 24 samples



PCR and RT-PCR Products Captured on Solid Support

Figure 16

Multiplexed PCR & RT-PCR; 24 samples





Figure 17

18/182

(Multiplexed PCR & RT-PCR; 24 samples)



LDR-FRET results, 24 samples

Figure 18

19/182




Figure 19

20/182





Figure 20

Distribute 1 sample into 24 tubes (wells), then Multiplexed PCR



PCR Products Captured on Solid Support

Figure 21

Distribute 1 sample into 24 tubes (wells), then Multiplexed PCR





Figure 22

23/182

(Distribute 1 sample into 24 tubes (wells), then Multiplexed PCR)



LDR-FRET results, 1 sample
w/ "Pixel" Enumeration

Figure 23

24/182




Figure 24

25/182





Figure 25

26/182

Dilute 1 sample into 24 tubes (wells), then Multiplexed RT-PCR



RT-PCR Products Captured on Solid Support

Figure 26

27/182

Dilute 1 sample into 24 tubes (wells), then Multiplexed RT-PCR





Figure 27

28/182

(Dilute 1 sample into 24 tubes (wells), then Multiplexed RT-PCR)



LDR-FRET results, 1 sample
w/ "Pixel" Enumeration

Figure 28

29/182

Poisson Distribution of 6 to 48 molecules in 24 wells

Wells with	6	12	18	24	30	36	42	48
0	19	15	11	9	7	5	4	3
1	5	7	9	9	9	8	7	6
2	1	2	3	4	5	6	6	6
3	0	0	1	1	2	3	4	4
4	0	0	0	0	1	1	2	2
5	0	0	0	0	0	0	1	1
6	0	0	0	0	0	0	0	0
7	0	0	0	0	0	0	0	0
8	0	0	0	0	0	0	0	0
9	0	0	0	0	0	0	0	0
10	0	0	0	0	0	0	0	0
11	0	0	0	0	0	0	0	0
12	0	0	0	0	0	0	0	0
13	0	0	0	0	0	0	0	0
14	0	0	0	0	0	0	0	0
15	0	0	0	0	0	0	0	0
16	0	0	0	0	0	0	0	0
17	0	0	0	0	0	0	0	0
18	0	0	0	0	0	0	0	0
19	0	0	0	0	0	0	0	0
20	0	0	0	0	0	0	0	0
21	0	0	0	0	0	0	0	0
22	0	0	0	0	0	0	0	0
23	0	0	0	0	0	0	0	0




Figure 29

Poisson Distribution of 12 to 96 molecules in 24 wells

Wells with	12	24	36	48	60	72	84	96
0	15	9	5	3	2	1	1	0
1	7	9	8	6	5	4	3	2
2	2	4	6	6	6	5	4	4
3	0	1	3	4	5	5	5	5
4	0	0	1	2	3	4	5	5
5	0	0	0	1	2	2	3	4
6	0	0	0	0	1	1	2	3
7	0	0	0	0	0	1	1	1
8	0	0	0	0	0	0	0	1
9	0	0	0	0	0	0	0	0
10	0	0	0	0	0	0	0	0
11	0	0	0	0	0	0	0	0
12	0	0	0	0	0	0	0	0
13	0	0	0	0	0	0	0	0
14	0	0	0	0	0	0	0	0
15	0	0	0	0	0	0	0	0
16	0	0	0	0	0	0	0	0
17	0	0	0	0	0	0	0	0
18	0	0	0	0	0	0	0	0
19	0	0	0	0	0	0	0	0
20	0	0	0	0	0	0	0	0
21	0	0	0	0	0	0	0	0
22	0	0	0	0	0	0	0	0
23	0	0	0	0	0	0	0	0




Figure 30

Poisson Distribution of 12 to 96 molecules in 48 wells

Wells with	12	24	36	48	60	72	84	96
0	37	29	23	18	14	11	8	6
1	9	15	17	18	17	16	15	13
2	1	4	6	9	11	12	13	13
3	0	1	2	3	4	6	7	9
4	0	0	0	1	1	2	3	4
5	0	0	0	0	0	1	1	2
6	0	0	0	0	0	0	0	1
7	0	0	0	0	0	0	0	0
8	0	0	0	0	0	0	0	0
9	0	0	0	0	0	0	0	0
10	0	0	0	0	0	0	0	0
11	0	0	0	0	0	0	0	0
12	0	0	0	0	0	0	0	0
13	0	0	0	0	0	0	0	0
14	0	0	0	0	0	0	0	0
15	0	0	0	0	0	0	0	0
16	0	0	0	0	0	0	0	0
17	0	0	0	0	0	0	0	0
18	0	0	0	0	0	0	0	0
19	0	0	0	0	0	0	0	0
20	0	0	0	0	0	0	0	0
21	0	0	0	0	0	0	0	0
22	0	0	0	0	0	0	0	0
...	...	...	...	...	...	...	...	...
47	0	0	0	0	0	0	0	0




Figure 31

Poisson Distribution of 24 to 192 molecules in 48 wells

Wells with	24	48	72	96	120	144	168	192
0	29	18	11	6	4	2	1	1
1	15	18	16	13	10	7	5	4
2	4	9	12	13	12	11	9	7
3	1	3	6	9	10	11	10	9
4	0	1	2	4	6	8	9	9
5	0	0	1	2	3	5	6	8
6	0	0	0	1	1	2	4	5
7	0	0	0	0	0	1	2	3
8	0	0	0	0	0	0	1	1
9	0	0	0	0	0	0	0	1
10	0	0	0	0	0	0	0	0
11	0	0	0	0	0	0	0	0
12	0	0	0	0	0	0	0	0
13	0	0	0	0	0	0	0	0
14	0	0	0	0	0	0	0	0
15	0	0	0	0	0	0	0	0
16	0	0	0	0	0	0	0	0
17	0	0	0	0	0	0	0	0
18	0	0	0	0	0	0	0	0
19	0	0	0	0	0	0	0	0
20	0	0	0	0	0	0	0	0
21	0	0	0	0	0	0	0	0
22	0	0	0	0	0	0	0	0
...	...	...	...	...	...	...	...	...
47	0	0	0	0	0	0	0	0




Figure 32

33/182

Poisson Distribution of 1 to 8 molecules in 8 wells

Wells with	1	2	3	4	5	6	7	8
0	7	6	5	5	4	4	3	3
1	1	2	2	2	3	3	3	3
2	0	0	0	1	1	1	1	1
3	0	0	0	0	0	0	0	0
4	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	0
7	0	0	0	0	0	0	0	0




Figure 33

34/182

Poisson Distribution of 2 to 16 molecules in 8 wells

Wells with	2	4	6	8	10	12	14	16
0	6	5	4	3	2	2	1	1
1	2	2	3	3	3	3	2	2
2	0	1	1	1	2	2	2	2
3	0	0	0	0	1	1	1	1
4	0	0	0	0	0	0	1	1
5	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	0
7	0	0	0	0	0	0	0	0




Figure 34

Poisson Distribution of 4 to 32 molecules in 8 wells

Wells with	4	8	12	16	20	24	28	32
0	5	3	2	1	1	0	0	0
1	2	3	3	2	2	1	1	1
2	1	1	2	2	2	2	1	1
3	0	0	1	1	2	2	2	2
4	0	0	0	1	1	1	2	2
5	0	0	0	0	1	1	1	1
6	0	0	0	0	0	0	1	1
7	0	0	0	0	0	0	0	0
8	0	0	0	0	0	0	0	0
9	0	0	0	0	0	0	0	0
10	0	0	0	0	0	0	0	0
11	0	0	0	0	0	0	0	0
12	0	0	0	0	0	0	0	0
13	0	0	0	0	0	0	0	0
14	0	0	0	0	0	0	0	0
15	0	0	0	0	0	0	0	0
16	0	0	0	0	0	0	0	0
17	0	0	0	0	0	0	0	0
18	0	0	0	0	0	0	0	0
19	0	0	0	0	0	0	0	0
20	0	0	0	0	0	0	0	0
21	0	0	0	0	0	0	0	0
22	0	0	0	0	0	0	0	0
23	0	0	0	0	0	0	0	0




Figure 35

Poisson Distribution of 8 to 64 molecules in 8 wells

Wells with	8	16	24	32	40	48	56	64
0	3	1	0	0	0	0	0	0
1	3	2	1	1	0	0	0	0
2	1	2	2	1	1	0	0	0
3	0	1	2	2	1	1	0	0
4	0	1	1	2	1	1	1	0
5	0	0	1	1	1	1	1	1
6	0	0	0	1	1	1	1	1
7	0	0	0	0	1	1	1	1
8	0	0	0	0	1	1	1	1
9	0	0	0	0	0	1	1	1
10	0	0	0	0	0	0	1	1
11	0	0	0	0	0	0	0	1
12	0	0	0	0	0	0	0	0
13	0	0	0	0	0	0	0	0
14	0	0	0	0	0	0	0	0
15	0	0	0	0	0	0	0	0
16	0	0	0	0	0	0	0	0
17	0	0	0	0	0	0	0	0
18	0	0	0	0	0	0	0	0
19	0	0	0	0	0	0	0	0
20	0	0	0	0	0	0	0	0
21	0	0	0	0	0	0	0	0
22	0	0	0	0	0	0	0	0
23	0	0	0	0	0	0	0	0




Figure 36

Poisson Distribution of 16 to 128 molecules in 8 wells

Wells with	16	32	48	64	80	96	112	128
0	1	0	0	0	0	0	0	0
1	2	1	0	0	0	0	0	0
2	2	1	0	0	0	0	0	0
3	1	2	1	0	0	0	0	0
4	1	2	1	0	0	0	0	0
5	0	1	1	1	0	0	0	0
6	0	1	1	1	1	0	0	0
7	0	0	1	1	1	0	0	0
8	0	0	1	1	1	1	0	0
9	0	0	1	1	1	1	0	0
10	0	0	0	1	1	1	1	0
11	0	0	0	1	1	1	1	0
12	0	0	0	0	1	1	1	1
13	0	0	0	0	1	1	1	1
14	0	0	0	0	0	1	1	1
15	0	0	0	0	0	1	1	1
16	0	0	0	0	0	0	1	1
17	0	0	0	0	0	0	1	1
18	0	0	0	0	0	0	0	1
19	0	0	0	0	0	0	0	1
20	0	0	0	0	0	0	0	0
21	0	0	0	0	0	0	0	0
22	0	0	0	0	0	0	0	0
23	0	0	0	0	0	0	0	0




Figure 37

38/182

- A. PCR-LDR-qPCR carryover prevention reaction to detect mutation. Isolate genomic or cfDNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, and amplify mutation containing regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR step.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Mutation-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 38

39/182

- A. PCR-qLDR carryover prevention reaction to detect mutation. Isolate genomic or cfDNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, aliquot into 12, 24, 48, or 96 wells, and amplify mutation containing regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Mutation-specific ligation oligonucleotides contain complementary tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Wash away unligated primers. Denature product from target, allowing tails to hybridize and bringing donor (D) and fluorescent group (F1) in close proximity. Then detect, and enumerate FRET signal.

Or

- F. Mutation-specific ligation oligonucleotides contain complementary tails for subsequent fluorescent detection. Ligase covalently seals the two oligonucleotides together.
- G. Denature product from target, allowing tails to hybridize and separate quencher (Q) from fluorescent reporter (F1). Then detect, and enumerate un-quenched fluorescent signal.





Figure 39

40/182

A. PCR-LDR-qPCR carryover prevention reaction to detect mutation. Isolate genomic or cfDNA.

B. Treat with UDG (uracil DNA glycosylase) for carryover prevention. Locus-specific primers are unblocked with RNaseH2 only when bound to target. Amplify mutation containing regions using PCR (and dUTP). Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)

C. PCR products incorporate dU, allowing for carryover prevention.

D. Mutation-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)

E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.

F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 40

41/182

- A. PCR-LDR-qPCR carryover prevention reaction to detect mutation. Isolate genomic or cfDNA.
- B. Treat with UDG for carryover prevention, and amplify mutation containing regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Mutation-specific ligation oligonucleotides (Mt) contain tags (Ai, Ci') for subsequent PCR amplification, while wild-type probe (Wt) does not. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target.
- E. Ligase covalently seals the unblocked upstream and phosphorylated downstream oligonucleotides together. Aliquot into separate wells for detection.
- F. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci'). Each Taqman probe spans the ligation junction.
- G. The 5'->3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 41

42/182

- A. PCR-qLDR carryover prevention reaction to detect mutation. Isolate genomic or cfDNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, aliquot into 12, 24, 48, or 96 wells, and amplify mutation containing regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Mutation-specific ligation oligonucleotides (Mt) contain tails for subsequent FRET detection, while wild-type probe (Wt) does not. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target.
- E. Ligase covalently seals the unblocked upstream and phosphorylated downstream oligonucleotides together.
- F. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 42

43/182

- A. PCR-qLDR carryover prevention reaction to detect mutation. Isolate genomic or cfDNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, aliquot into 12, 24, 48, or 96 wells, and amplify mutation containing regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Mutation-specific ligation oligonucleotides (Mt) overlap at the mutation base, and contain tails for subsequent FRET detection, while wild-type probe (Wt) does not. 5'-nuclease activity of polymerase cleaves off only matching 5'-overlapping base and additional flap, leaving ligation-competent 5-phosphate.
- E. Ligase covalently seals the upstream and unblocked phosphorylated downstream oligonucleotides together.
- F. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 43

44/182

- A. PCR-LDR-qPCR carryover prevention reaction to detect mutation. Isolate genomic or cfDNA.
- B. Treat with UDG for carryover prevention, and amplify mutation containing regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Mutation-specific ligation oligonucleotides contain tags (Ai, Bi'-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- F. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'->3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 44

45/182

A. PCR-LDR-qPCR carryover prevention reaction to detect methylation. Isolate genomic or cfDNA. Optional: enrich for methylated DNA using antibodies. Treat with methyl-sensitive restriction endonucleases Bsh1236I (CG[^]CG) and HinP1I (G[^]CGC), and UNG (37°C, 30-60 minutes) to completely digest unmethylated DNA and prevent carryover.

B. Amplify methyl containing regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)

C. PCR products incorporate dU, allowing for carryover prevention. (Products lack methyl groups, providing additional protection.)

D. Methyl region-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.

E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.

F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 45

- 46/182
- PCR-LDR-qPCR carryover prevention reaction to detect methylation. Isolate genomic or cfDNA. Treat with Bsh1236I (CG[^]CG) and HinfI (G[^]CGC), and UNG (37°C, 30-60 minutes) to completely digest unmethylated DNA and prevent carryover.
 - Amplify methyl containing regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
 - PCR products incorporate dU, allowing for carryover prevention.
 - Methyl region-specific ligation oligonucleotides contain tags (Ai, Bi'-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
 - Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
 - Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
 - After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 46

47/182

- A. PCR-qLDR carryover prevention reaction to detect methylation. Isolate genomic or cfDNA. Optional: enrich for methylated DNA using antibodies. Treat with methyl-sensitive restriction endonucleases Bsh1236I (CG[^]CG) and HinP1I (G[^]CGC), and UNG (37°C, 30-60 minutes) to completely digest unmethylated DNA and prevent carryover.
- B. Aliquot into 12, 24, 48, or 96 wells, and amplify methyl containing regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. (Products lack methyl groups, providing additional protection.) Separate product from primers and capture biotinylated product on solid support.
- D. Methyl region-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together.
- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 47

48/182

- A. Nuclease-Ligation-PCR-qPCR carryover prevention reaction to detect methylation. Isolate genomic or cfDNA. Treat with HaeIII (GG[^]CC) and methyl-sensitive restriction endonucleases Bsh1236I (CG[^]CG) and HinP1I (G[^]CGC), and UNG (37°C, 30-60 minutes) to completely digest unmethylated DNA and prevent carryover.
- B. Ligate hairpin oligonucleotide containing tag sequence (Ai') to newly liberated phosphate of template strand.
- C. Treat with HinP1I and Bsh1236I at 37°C, then heat kill endonucleases while activating Taq polymerase. Locus-specific primers (containing Ci tag sequence) are unblocked with RNaseH2 only when bound to target, and extended with polymerase, whose 5' nuclease activity digests 5' portion of hairpin to generate target complement with Ai' sequence. Unligated hairpin extends on itself.
- D. Amplify methyl containing regions using PCR (and dUTP) with tag-primers Ai and Ci. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- E. PCR products incorporate dU, allowing for carryover prevention. (Products lack methyl groups, providing additional protection.) Aliquot into separate wells for Taqman detection using locus-specific primers and Taqman probe.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 48

49/182

- A. Nuclease-Ligation-PCR-qPCR carryover prevention reaction to detect methylation. Isolate genomic or cfDNA. Treat with HaeIII (GG[^]CC), Bsh1236I (CG[^]CG) and HinP1I (G[^]CGC), and UNG.
- B. Ligate hairpin oligonucleotide containing tag sequence (Ai') to newly liberated phosphate of template strand.
- C. Treat with HinP1I and Bsh1236I at 37°C, then heat kill endonucleases while activating Taq polymerase. Locus-specific primers (containing Ci tag sequence) are unblocked with RNaseH2 only when bound to target, and extended with polymerase, whose 5' nuclease activity digests 5' portion of hairpin to generate target complement with Ai' sequence.
- D. Amplify methyl containing regions using PCR (and dUTP) with tag-primers Ai and Ci. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- E. PCR products incorporate dU, allowing for carryover prevention. Aliquot into separate wells for UniTaq detection, and amplify using locus-specific primers tailed with Aj, and Bj-Cj, as well as Unitaq-specific primers (F1-Bj-Q-Aj, and Cj).
- F. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bj and Bj'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 49

50/182

- A. PCR-LDR-qPCR carryover prevention reaction to detect methylation. Isolate genomic or cfDNA. Treat with Bsh1236I (CG[^]CG) and UNG (37°C, 30-60 minutes) to completely digest unmethylated DNA and prevent carryover. Treat with bisulfite, which renders the strands non-complementary.
- B. Hybridize locus-specific primers in presence of BstU1 (CG[^]CG), will cleave carryover DNA. Primers are unblocked with RNaseH2 only when bound to target. Amplify methyl containing regions using PCR (and dUTP). Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Blocking oligo may be used to limit amplification of originally un-methylated DNA (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention. (Products lack methyl groups, providing additional protection.)
- D. Methyl region-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 50

- A. PCR-LDR-qPCR carryover prevention reaction to detect methylation. Isolate genomic or cfDNA. Treat with Bsh1236I (CG[^]CG) and UNG (37°C, 30-60 minutes) to completely digest unmethylated DNA and prevent carryover. Treat with bisulfite, which renders the strands non-complementary.
- B. Hybridize locus-specific primers in presence of BstU1 (CG[^]CG), will cleave carryover DNA. Primers are unblocked with RNaseH2 only when bound to target. Amplify methyl containing regions using PCR (and dUTP). Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Blocking oligo may be used to limit amplification of originally unmethylated DNA (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Methyl region-specific ligation oligonucleotides contain tags (Ai, Bi-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- F. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal.




Figure 51

52/182

A. PCR-qLDR carryover prevention reaction to detect methylation. Isolate genomic or cfDNA. Treat with Bsh1236I (CG[^]CG) and UNG (37°C, 30-60 minutes) to completely digest unmethylated DNA and prevent carryover. Treat with bisulfite, which renders the strands non-complementary.

B. Hybridize locus-specific primers in presence of BstU1 (CG[^]CG), will cleave carryover DNA. Primers are unblocked with RNaseH2 only when bound to target. Aliquot into 12, 24, 48, or 96 wells, and amplify methyl containing regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin. Blocking oligo may be used to limit amplification of originally unmethylated DNA.

C. PCR products incorporate dU, allowing for carryover prevention. (Products lack methyl groups, providing additional protection.) Separate product from primers and capture biotinylated product on solid support.

D. Methyl region-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together.

E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 52

53/182

- A. PCR-qPCR carryover prevention reaction to detect methylation. Isolate genomic or cfDNA. Treat Bsh1236I (CG[^]CG) and UNG (37°C, 30-60 minutes) to completely digest unmethylated DNA and prevent carryover. Treat with bisulfite, which renders the strands non-complementary.

- B. Hybridize locus-specific primers in presence of BstU1 (CG[^]CG), will cleave carryover DNA. Primers are unblocked with RNaseH2 only when bound to target. Amplify methyl containing regions using PCR with dNTPs. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Blocking oligos may be used to limit amplification of wild-type DNA. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)

- C. PCR products are unmethylated providing carryover protection.

- D. Aliquot into separate wells for Taqman detection using locus-specific primers and Taqman probe. Optional blocking oligonucleotide is used to limit amplification of wild-type DNA

- E. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)






Figure 53

54/182


- A. Overview: RT-PCR-LDR-qPCR carryover prevention reaction to detect translocation at the mRNA level. Illustration of translocations between two genes shown at the DNA level.



- B. Examples of fusion junctions between exons 1-b, 2-b, and 3-b in mRNAs are illustrated. Isolate mRNA from whole blood cells, exosomes, or CTCs. Generate cDNA using reverse transcriptase and primer complementary to exon b. PCR amplify using forward primers to exons 1, 2, and 3. Independent of translocation breakpoint, the primers will amplify the smallest fragment containing the exon junction region.



- C. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci' for individual Taqman probes; or Ai, Bi-Ci for UniTaq probes) for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together, if there is perfect complementarity at the junction. Aliquot into separate wells for detection using real-time PCR.



- D. When using tag-specific primers (Ai, Ci) for amplifying LDR products of the upper set, each Taqman probe spans the ligation junction, and can be scored individually. When using UniTaq-specific primers (F1-Bi-Q-Ai, Ci), for amplifying LDR products of the lower set, the same primer set scores for the given translocation, independent of the specific exon junction.




Figure 54

55/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to detect translocation at the mRNA level. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'->3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 55

56/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to detect translocation at the mRNA level. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Bi-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- F. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal.




Figure 56

57/182

- A. RT-PCR-qLDR carryover prevention reaction to detect translocation at the mRNA level. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, aliquot into 12, 24, 48, or 96 wells, and heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Exon junction-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.






Figure 57

58/182


- A. Overview: RT-PCR-LDR-qPCR carryover prevention reaction to detect alternative splicing. Illustration of gene with 5 exons shown at the DNA level.



- B. Examples of normal (1-2-3a-4), alternative splice variant (1-2-3b-4) mRNAs are illustrated. Isolate mRNA from whole blood cells, exosomes, or CTCs. Generate cDNA using reverse transcriptase and primer complementary to exon 4. PCR amplify using forward primer to exon 2, generating amplicons of both splice variants.



- C. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci' for individual Taqman probes; or Ai, Bi-Ci for UniTaq probes) for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together, if there is perfect complementarity at the junction. Aliquot into separate wells for detection using real-time PCR.



- D. When using tag-specific primers (Ai, Ci) for amplifying LDR products of the upper set, each Taqman probe spans the ligation junction, and can be scored individually. When using UniTaq-specific primers (F1-Bi-Q-Ai, Ci), for amplifying LDR products of the lower set, a given primer set scores for a given ligation junction.






Figure 58

59/182


- A. RT-PCR-LDR-qPCR carryover prevention reaction to quantify wt and alternatively spliced (3a vs. 3b) transcript. Isolate mRNA.




- B. Treat with UDГ (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)




- C. PCR products incorporate dU, allowing for carryover prevention.



- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)



- E. Treat with UDГ for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.



- F. The 5'->3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. The relative Ct values reflects the relative amount of each transcript.






Figure 59

60/182


- A. RT-PCR-LDR-qPCR carryover prevention reaction to quantify wt and alternatively spliced (3a vs. 3b) transcript. Isolate mRNA.




- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)




- C. PCR products incorporate dU, allowing for carryover prevention.




- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Bi-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.



- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).



- F. Double-stranded DNA products of PCR (not shown); the polymerase blocker stops extension of the bottom strand.




- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'->3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal. The relative Ct values reflects the relative amount of each transcript.



Figure 60

61/182


- A. RT-PCR-qLDR carryover prevention reaction to quantify wt and alternatively spliced (3a vs. 3b) transcript. Isolate mRNA.




- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, aliquot into 12, 24, 48, or 96 wells, and heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.



- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.



- D. Exon junction-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)



- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 61

62/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to detect low-level alternatively spliced (e.g. 3b) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Exon-specific primer does not amplify wt transcript. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 62

63/182

- A. PCR-LDR carryover prevention reaction to detect low-level alternatively spliced (e.g. 3b) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Exon-specific primer does not amplify wt transcript. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Bi-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- F. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal.




Figure 63

64/182

- A. PCR-LDR carryover prevention reaction to detect low-level alternatively spliced (e.g. 3b) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, aliquot into 12, 24, 48, or 96 wells, and heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Exon junction-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.






Figure 64

65/182


- A. Overview: RT-PCR-LDR-qPCR carryover prevention reaction to detect alternative splicing. Illustration of gene with 3 exons, and an alternative start site and first exon, shown at the DNA level.



- B. Examples of normal (1-2-3), alternative splice variant (1a-2-3) mRNAs are illustrated. Isolate mRNA from whole blood cells, exosomes, or CTCs. Generate cDNA using reverse transcriptase and primer complementary to exon 2. PCR amplify using forward primers to exon 1 and 1a, generating amplicons of both splice variants.



- C. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci' for individual Taqman probes; or Ai, Bi-Ci for UniTaq probes) for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together, if there is perfect complementarity at the junction. Aliquot into separate wells for detection using real-time PCR.



- D. When using tag-specific primers (Ai, Ci) for amplifying LDR products of the upper set, each Taqman probe spans the ligation junction, and can be scored individually. When using UniTaq-specific primers (F1-Bi-Q-Ai, Ci), for amplifying LDR products of the lower set, a given primer set scores for a given ligation junction.




Figure 65

66/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to quantify wt and alternative transcript start site (1 vs. 1a). Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. The relative Ct values reflects the relative amount of each transcript.




Figure 66

67/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to quantify wt and alternative transcript start site (1 vs. 1a). Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. The relative Ct values reflects the relative amount of each transcript.




Figure 67

68/182

- A. RT-PCR-qLDR carryover prevention reaction to quantify wt and alternative transcript start site (1 vs. 1a). Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, aliquot into 12, 24, 48, or 96 wells, and heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Exon junction-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 68

69/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to detect low level of alternative transcript start site (1 vs. 1a). Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Exon-specific primer does not amplify wt transcript. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 69

70/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to detect low level of alternative transcript start site (1 vs. 1a). Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Exon-specific primer does not amplify wt transcript. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Bi-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- F. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal.




Figure 70

71/182

- A. RT-PCR-qLDR carryover prevention reaction to detect low level of alternative transcript start site (1 vs. 1a). Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, aliquot into 12, 24, 48, or 96 wells, and heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Exon junction-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.






Figure 71

72/182


- A. Overview: RT-PCR-LDR-qPCR carryover prevention reaction to detect exon deletion. Illustration of gene with 5 exons and 4 introns shown at the DNA level.



- B. Example of normal (e1-e2-e3-e4-e5) and exon deletion (e1-e2-e3-e5) mRNAs are illustrated. Isolate mRNA from whole blood cells, exosomes, or CTCs. Generate cDNA using reverse transcriptase and primer complementary to exon e5. PCR amplify using forward primers to exons e3, and e4, generating amplicons of both splice variants.



- C. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci' for individual Taqman probes; or Ai, Bi-Ci for UniTaq probes) for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together, if there is perfect complementarity at the junction. Aliquot into separate wells for detection using real-time PCR.



- D. When using tag-specific primers (Ai, Ci) for amplifying LDR products of the upper set, each Taqman probe spans the ligation junction, and can be scored individually. When using UniTaq-specific primers (F1-Bi-Q-Ai, Ci), for amplifying LDR products of the lower set, a given primer set scores for a given ligation junction.




Figure 72

73/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to quantify wt and alternatively spliced (exon deletion) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. The relative Ct values reflects the relative amount of each transcript.




Figure 73

74/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to quantify wt and alternatively spliced (exon deletion) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Bi-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- F. Double-stranded DNA products of PCR (not shown); the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'->3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal. The relative Ct values reflects the relative amount of each transcript.




Figure 74

75/182

- A. RT-PCR-qLDR carryover prevention reaction to quantify wt and alternatively spliced (exon deletion) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, aliquot into 12, 24, 48, or 96 wells, and heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Exon junction-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 75

76/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to detect low-level alternatively spliced (exon deletion) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Blocking oligo suppresses amplification of wt transcript. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 76

77/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to detect low-level alternatively spliced (exon deletion) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Blocking oligo suppresses amplification of wt transcript. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Bi-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- F. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal.




Figure 77

78/182

- A. RT-PCR-qLDR carryover prevention reaction to detect low-level alternatively spliced (exon deletion) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, aliquot into 12, 24, 48, or 96 wells, and heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Blocking oligo suppresses amplification of wt transcript. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Exon junction-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.






Figure 78

79/182


- A. Overview: RT-PCR-LDR-qPCR carryover prevention reaction to detect alternative splicing with intron insertion. Illustration of gene with 5 exons and 4 introns shown at the DNA level.



- B. Examples of normal (e1-e2-e3-e4-e5) and intron insertion (e1-i1-e2-e3-e4-e5) mRNAs are illustrated. Isolate mRNA from whole blood cells, exosomes, or CTCs. Generate cDNA using reverse transcriptase and primer complementary to exon e2. PCR amplify using forward primers to exon e1, generating amplicons of both splice variants.



- C. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci' for individual Taqman probes; or Ai, Bi-Ci for UniTaq probes) for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together, if there is perfect complementarity at the junction. Aliquot into separate wells for detection using real-time PCR.



- D. When using tag-specific primers (Ai, Ci) for amplifying LDR products of the upper set, each Taqman probe spans the ligation junction, and can be scored individually. When using UnTaq-specific primers (F1-Bi-Q-Ai, Ci), for amplifying LDR products of the lower set, a given primer set scores for a given ligation junction.




Figure 79

80/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to quantify wt and alternatively spliced (intron insertion) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. The relative Ct values reflects the relative amount of each transcript.




Figure 80

81/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to quantify wt and alternatively spliced (intron insertion) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Bi-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- F. Double-stranded DNA products of PCR (not shown); the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal. The relative Ct values reflects the relative amount of each transcript.




Figure 81

82/182

- A. RT-PCR-qLDR carryover prevention reaction to quantify wt and alternatively spliced (intron insertion) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, aliquot into 12, 24, 48, or 96 wells, and heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Exon junction-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 82

83/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to detect low-level alternatively spliced (intron insertion) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Intron-specific primer does not amplify wt transcript. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 83

84/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to detect low-level alternatively spliced (intron insertion) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Intron-specific primer does not amplify wt transcript. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Exon junction-specific ligation oligonucleotides contain tags (Ai, Bi-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- F. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal.




Figure 84

85/182

- A. PCR-LDR carryover prevention reaction to detect low-level alternatively spliced (intron insertion) transcript. Isolate mRNA.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, aliquot into 12, 24, 48, or 96 wells, and heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Intron-specific primer does not amplify wt transcript. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Exon junction-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 85

86/182

- A. PCR-LDR-qPCR carryover prevention reaction to enumerate DNA copy number. Isolate tumor DNA from CTCs or tumor-specific exosomes.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, and amplify chromosomal regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. For accurate enumeration, aliquot into 12, 24, 48, or 96 wells prior to PCR.
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Locus-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. Determine copy number based on signal from wells where original distribution was one copy/well.




Figure 86

87/182

- A. PCR-LDR-qPCR carryover prevention reaction to enumerate DNA copy number. Isolate tumor DNA from CTCs or tumor-specific exosomes.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, and amplify chromosomal regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. For accurate enumeration, aliquot into 12, 24, 48, or 96 wells prior to PCR.
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Locus-specific ligation oligonucleotides contain tags (Ai, Bi'-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- F. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'->3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal. Determine copy number based on signal from wells where original distribution was one copy/well.




Figure 87

88/182

- A. PCR-qLDR carryover prevention reaction to enumerate DNA copy number. Isolate tumor DNA from CTCs or tumor-specific exosomes.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, aliquot into 12, 24, 48, or 96 wells, and amplify mutation containing regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Locus-specific ligation: oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal. Determine copy number based on signal from wells where original distribution was one copy/well.




Figure 88

89/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to enumerate RNA copy number. Isolate RNA from whole blood cells, exosomes, or CTCs..
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. For accurate enumeration, aliquot into 12, 24, 48, or 96 wells prior to PCR.
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Locus-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. Determine copy number based on signal from wells where original distribution was one copy/well.




Figure 89

90/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to enumerate RNA copy number. Isolate RNA from whole blood cells, exosomes, or CTCs.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. For accurate enumeration, aliquot into 12, 24, 48, or 96 wells prior to PCR.
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Locus-specific ligation oligonucleotides contain tags (Ai, Bi'-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- F. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'->3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal. Determine copy number based on signal from wells where original distribution was one copy/well.




Figure 90

91/182

- A. RT-PCR-qLDR carryover prevention reaction to enumerate RNA copy number. Isolate RNA from whole blood cells, exosomes, or CTCs.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, and aliquot into 12, 24, 48, or 96 wells. Use reverse-transcriptase to make cDNA copy (using dUTP) with 3' transcript-specific primers. Activate Taq polymerase and perform limited cycle PCR amplification (12-20) to maintain relative ratios of different amplicons. Primers contain universal tails (with identical 8-11 bases to prevent primer dimers), also enabling universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Locus-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal. Determine copy number based on signal from wells where original distribution was one copy/well.




Figure 91

92/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to quantify miRNA. Isolate miRNA from exosomes.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Hybridize probe complementary to 3' end of target miRNA, containing a stem-loop, tag (Tj), and blocking group. Extend 3' end of loop primer with reverse transcriptase (using dUTP). Activate Taq polymerase and perform limited cycle PCR amplification (12-20), using bridge and tag primers (Ti, Tj), to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. miRNA sequence-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 92

93/182

- A. RT-PCR-LDR-qPCR carryover prevention reaction to quantify miRNA. Isolate miRNA from exosomes.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Hybridize probe complementary to 3' end of target miRNA, containing a stem-loop, tag (Tj), and blocking group. Extend 3' end of loop primer with reverse transcriptase (using dUTP). Activate Taq polymerase and perform limited cycle PCR amplification (12-20), using bridge and tag primers (Ti, Tj), to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. miRNA sequence-specific ligation oligonucleotides contain tags (Ai, Bi-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- F. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal.




Figure 93

94/182

- A. RT-PCR-qLDR carryover prevention reaction to quantify miRNA. Isolate miRNA from exosomes.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, and aliquot into 12, 24, 48, or 96 wells. Hybridize probe complementary to 3' end of target miRNA, containing a stem-loop, tag (Tj), and blocking group. Extend 3' end of loop primer with reverse transcriptase (using dUTP). Activate Taq polymerase and perform limited cycle PCR amplification (12-20), using bridge and tag primers (Ti, Tj), to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers, with one primer containing a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. miRNA-specific ligation
oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 94

95/182

- A. Ligation-RT-PCR-LDR-qPCR carryover prevention reaction to quantify miRNA. Isolate miRNA from exosomes.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Ligate probe complementary to 3' end of target miRNA, containing a stem-loop, Tag (Tj'), and blocking group. Extend with primer Tj and reverse transcriptase (using dUTP). Activate Taq polymerase and perform limited cycle PCR amplification (12-20), using bridge and tag primers (Ti, Tj), to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. miRNA sequence-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- F. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 95

96/182

- A. Ligation-RT-PCR-LDR-qPCR carryover prevention reaction to quantify miRNA. Isolate miRNA from exosomes.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Ligate probe complementary to 3' end of target miRNA, containing a stem-loop, Tag (Tj'), and blocking group. Extend with primer Tj and reverse transcriptase (using dUTP). Activate Taq polymerase and perform limited cycle PCR amplification (12-20), using bridge and tag primers (Ti, Tj), to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. miRNA sequence-specific ligation oligonucleotides contain tags (Ai, Bi-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- E. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- F. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- G. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal.




Figure 96

97/182

- A. Ligation-RT-PCR-qLDR carryover prevention reaction to quantify miRNA. Isolate miRNA from exosomes.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, and aliquot into 12, 24, 48, or 96 wells. Ligate probe complementary to 3' end of target miRNA, containing a stem-loop, Tag (Tj'), and blocking group. Extend with primer Tj and reverse transcriptase (using dUTP). Activate Taq polymerase and perform limited cycle PCR amplification (12-20), using bridge and tag primers (Ti, Tj), to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers, with one primer containing a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. miRNA-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- E. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 97

98/182

- A. Ligation-RT-PCR-LDR-qPCR carryover prevention reaction to quantify miRNA. Isolate miRNA from exosomes.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Ligate probe complementary to 3' end of target miRNA, containing a stem-loop, Tag (Tj'), and blocking group. Extend with primer Tj and reverse transcriptase (using dUTP).
- C. miRNA-specific bridge primers are unblocked with RNaseH2 only when bound to target. Perform limited cycle PCR amplification (12-20), using tag primers (Ti, Tj), to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- D. PCR products incorporate dU, allowing for carryover prevention.
- E. miRNA sequence-specific ligation oligonucleotides contain tags (Ai, Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- F. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci). Each Taqman probe spans the ligation junction.
- G. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 98

99/182

- A. Ligation-RT-PCR-LDR-qPCR carryover prevention reaction to quantify miRNA. Isolate miRNA from exosomes.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, heat kill. Ligate probe complementary to 3' end of target miRNA, containing a stem-loop, Tag (Tj'), and blocking group. Extend with primer Tj and reverse transcriptase (using dUTP).
- C. miRNA-specific bridge primers are unblocked with RNaseH2 only when bound to target. Perform limited cycle PCR amplification (12-20), using tag primers (Ti, Tj), to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- D. PCR products incorporate dU, allowing for carryover prevention.
- E. miRNA sequence-specific ligation oligonucleotides contain tags (Ai, Bi-Ci') for subsequent PCR amplification. Ligase covalently seals the two oligonucleotides together. Aliquot into separate wells for detection.
- F. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- G. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- H. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal.




Figure 99

100/182

- A. Ligation-RT-PCR-qLDR carryover prevention reaction to quantify miRNA. Isolate miRNA from exosomes.
- B. Treat with UDG (uracil DNA glycosylase) for carryover prevention, and aliquot into 12, 24, 48, or 96 wells. Ligate probe complementary to 3' end of target miRNA, containing a stem-loop, Tag (Tj'), and blocking group. Extend with primer Tj and reverse transcriptase (using dUTP).
- C. miRNA-specific bridge primers are unblocked with RNaseH2 only when bound to target. Perform limited cycle PCR amplification (12-20), using tag primers (Ti, Tj), to maintain relative ratios of different amplicons. Primers contain identical 8-11 base tails to prevent primer dimers, with one primer containing a 5' biotin.
- D. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- E. miRNA-specific ligation oligonucleotides contain tails for subsequent FRET detection. Ligase covalently seals the two oligonucleotides together. (LDR products are not amplified by either original PCR or LDR primers, providing additional protection.)
- F. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 100

101/182




Figure 101

102/182




Figure 102

103/182





Figure 103

104/182





Figure 104



106/182




Figure 106




Figure 107

108/182




Figure 108




Figure 109

110/182




Figure 110




Figure 111

112/182




Figure 112




Figure 113

114/182




Figure 114




Figure 115

116/182




Figure 116




Figure 117

118/182




Figure 118




Figure 119

120/182




Figure 120




Figure 121

122/182




Figure 122




Figure 123

124/182




Figure 124




Figure 125

126/182




Figure 126




Figure 127

128/182





Figure 128



130/182




Figure 130




Figure 131

132/182




Figure 132




Figure 133

134/182




Figure 134




Figure 135

136/182




Figure 136




Figure 137

138/182




Figure 138




Figure 139

140/182




Figure 140




Figure 141

142/182




Figure 142

143/182




Figure 143

144/182




Figure 144

145/182

- A. PCR-LDR-qPCR carryover prevention reaction to detect low-level mutation. Isolate genomic or cfDNA.
- B. Treat with UDG for carryover prevention. Upstream locus primers contain mutation-specific ribose bases (mr), which are removed with RNaseH2 only when bound to target, thus unblocking primer. Primers with Wt sequence remain blocked. Preferentially amplify mutation containing regions using PCR (and dUTP). Downstream primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Mutation-specific ligation oligonucleotides (Mt) contain tags (Ai, Ci') for subsequent PCR amplification, while wild-type probe (Wt) does not. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target.
- E. Ligase covalently seals the unblocked mutation-specific upstream and phosphorylated downstream oligonucleotides together. Aliquot into separate wells for detection.
- F. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci'). Each Taqman probe spans the ligation junction.
- G. The 5'->3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 145

146/182

- A. PCR-LDR-qPCR carryover prevention reaction to detect low-level mutation. Isolate genomic or cfDNA.
- B. Treat with UDG for carryover prevention. Upstream locus primers contain mutation-specific ribose bases (mr), which are removed with RNaseH2 only when bound to target, thus unblocking primer. Primers with Wt sequence remain blocked. Preferentially amplify mutation containing regions using PCR (and dUTP). Downstream primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Mutation-specific ligation oligonucleotides (Mt) contain tags (Ai, Bi', Ci') for subsequent PCR amplification, while wild-type probe (Wt) does not. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target.
- E. Ligase covalently seals the unblocked mutation-specific upstream and phosphorylated downstream oligonucleotides together. Aliquot into separate wells for detection.
- F. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- G. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- H. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 146

147/182

- A. PCR-qLDR carryover prevention reaction to detect low-level mutation. Isolate genomic or cfDNA.
- B. Treat with UDG for carryover prevention. Upstream locus primers contain mutation-specific ribose bases (mr), which are removed with RNaseH2 only when bound to target, thus unblocking primer. Primers with Wt sequence remain blocked. Preferentially amplify mutation containing regions using PCR (and dUTP). Downstream primers contain universal tails to enable universal primer amplification to append a 5' biotin.
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Mutation-specific ligation oligonucleotides (Mt) contain tails for subsequent FRET detection, while wild-type probe (Wt) does not. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target.
- E. Ligase covalently seals the unblocked upstream and phosphorylated downstream oligonucleotides together.
- F. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 147

148/182

- A. PCR-LDR-qPCR carryover prevention reaction to detect low-level mutation. Isolate genomic or cfDNA.
- B. Treat with UDG for carryover prevention. Upstream locus primers contain ribose bases (r), which are removed with RNaseH2 only when bound to target, thus unblocking primer. Blocking LNA or PNA Wt probes facilitate preferential amplification of mutation containing regions using PCR (and dUTP). Downstream primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Mutation-specific ligation oligonucleotides (Mt) contain tags (Ai, Ci') for subsequent PCR amplification. Blocking LNA or PNA Wt probes suppress ligation to Wt sequence. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target.
- E. Ligase covalently seals the unblocked mutation-specific upstream and phosphorylated downstream oligonucleotides together. Aliquot into separate wells for detection.
- F. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci'). Each Taqman probe spans the ligation junction.
- G. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 148

149/182

- A. PCR-LDR-qPCR carryover prevention reaction to detect low-level mutation. Isolate genomic or cfDNA.
- B. Treat with UDG for carryover prevention. Upstream locus primers contain ribose bases (r), which are removed with RNaseH2 only when bound to target, thus unblocking primer. Blocking LNA or PNA Wt probes facilitate preferential amplification of mutation containing regions using PCR (and dUTP). Downstream primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Mutation-specific ligation oligonucleotides (Mt) contain tags (Ai, Bi', Ci') for subsequent PCR amplification. Blocking LNA or PNA Wt probes suppress ligation to Wt sequence. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target.
- E. Ligase covalently seals the unblocked mutation-specific upstream and phosphorylated downstream oligonucleotides together. Aliquot into separate wells for detection.
- F. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- G. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- H. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'->3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 149

150/182

- A. PCR-qLDR carryover prevention reaction to detect low-level mutation. Isolate genomic or cfDNA.
- B. Treat with UDG for carryover prevention. Upstream locus primers contain ribose bases (r), which are removed with RNaseH2 only when bound to target, thus unblocking primer. Blocking LNA or PNA Wt probes facilitate preferential amplification of mutation containing regions using PCR (and dUTP). Downstream primers contain universal tails to enable universal primer amplification to append a 5' biotin. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Mutation-specific ligation oligonucleotides (Mt) contain tails for subsequent FRET detection. Blocking LNA or PNA Wt probes suppress ligation to Wt sequence. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target.
- E. Ligase covalently seals the unblocked upstream and phosphorylated downstream oligonucleotides together.
- F. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 150

151/182

- A. PCR-LDR-qPCR carryover prevention reaction to detect low-level methylation. Isolate genomic or cfDNA. Treat with Bsh1236I (CG[^]CG) and UNG (37°C, 30-60 minutes) to completely digest un-methylated DNA and prevent carryover. Treat with bisulfite, which renders the strands non-complementary.
- B. Locus primers contain ribose bases (r), which are removed with RNaseH2 only when bound to target, thus unblocking primer. Blocking LNA or PNA probes may be used to limit amplification of originally un-methylated DNA using PCR (and dUTP). Downstream primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Methyl region-specific ligation oligonucleotides (m) contain tags (Ai, Ci') for subsequent PCR amplification. Blocking LNA or PNA probes suppress ligation to originally un-methylated DNA sequence. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target.
- E. Ligase covalently seals the unblocked methyl region-specific upstream and phosphorylated downstream oligonucleotides together. Aliquot into separate wells for detection.
- F. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci'). Each Taqman probe spans the ligation junction.
- G. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 151

152/182

- A. PCR-LDR-qPCR carryover prevention reaction to detect low-level methylation. Isolate genomic or cfDNA. Treat with Bsh1236I (CG[^]CG) and UNG (37°C, 30-60 minutes) to completely digest un-methylated DNA and prevent carryover. Treat with bisulfite, which renders the strands non-complementary.
- B. Locus primers contain ribose bases (r), which are removed with RNaseH2 only when bound to target, thus unblocking primer. Blocking LNA or PNA probes may be used to limit amplification of originally un-methylated DNA using PCR (and dUTP). Downstream primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention.
- D. Methyl region-specific ligation oligonucleotides (m) contain tags (Ai, Bi', Ci') for subsequent PCR amplification. Blocking LNA or PNA probes suppress ligation to originally un-methylated DNA sequence. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target.
- E. Ligase covalently seals the unblocked methyl region-specific upstream and phosphorylated downstream oligonucleotides together. Aliquot into separate wells for detection.
- F. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with Unitaq-specific primers (F1-Bi-Q-Ai, Ci).
- G. Double-stranded DNA products of PCR; the polymerase blocker stops extension of the bottom strand.
- H. After the denaturation step, as the temperature decreases, a hairpin forms between Bi and Bi'. The 5'→3' nuclease activity of Taq polymerase extends primer Ci and liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 152

153/182

- A. PCR-LDR-qPCR carryover prevention reaction to detect low-level methylation. Isolate genomic or cfDNA. Treat with Bsh1236I (CG[^]CG) and UNG (37°C, 30-60 minutes) to completely digest un-methylated DNA and prevent carryover. Treat with bisulfite, which renders the strands non-complementary.
- B. Locus primers contain ribose bases (r), which are removed with RNaseH2 only when bound to target, thus unblocking primer. Blocking LNA or PNA probes may be used to limit amplification of originally un-methylated DNA using PCR (and dUTP). Downstream primers contain universal tails to enable universal primer amplification to append a 5' biotin. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support.
- D. Methyl region-specific ligation oligonucleotides (m) contain tails for subsequent FRET detection. Blocking LNA or PNA probes suppress ligation to originally un-methylated DNA sequence. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target.
- E. Ligase covalently seals the unblocked upstream and phosphorylated downstream oligonucleotides together.
- F. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 153

155/182

- F. (PCR products incorporate dU, allowing for carryover prevention. Optional: wild-type sequence on end may be clipped off with restriction endonuclease.)

- G. Mutation-specific ligation oligonucleotides (Mt) contain tags (Ai, Ci') for subsequent PCR amplification, while (optional) wild-type probe (Wt) does not. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target. (Optional: wild-type sequence on 3' end may be clipped off with exonuclease III)

- H. Ligase covalently seals the unblocked upstream and phosphorylated downstream oligonucleotides together. Aliquot into separate wells for detection.

- I. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci'). Each Taqman probe spans the ligation junction.

- J. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 154 (Cont.)

156/182

- A. Loop-PCR-qLDR carryover prevention reaction to detect low-level mutation. Isolate genomic or cfDNA.
- B. Treat with UDG for carryover prevention. Upstream locus-specific primers also contain 5' sequences complementary to Wt sequence of top strand, and are unblocked with RNaseH2 only when bound to target. Amplify mutation containing regions using PCR (and dUTP). Downstream primers contain universal tails to enable universal primer amplification to append a 5' biotin. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. (PCR products incorporate dU, allowing for carryover prevention.)
- D. In detail: PCR is performed with a polymerase lacking 5' nuclease, 3' nuclease, and strand-displacement activity. (i) Denaturation of Wt bottom strand results in hairpin with perfect match at 3' end, which is extended by polymerase. (ii) Denaturation of Mt bottom strand results in hairpin with mismatch at 3' end, which generally is not extended by polymerase. (iii) Denaturation of top strand results in hairpin on 5' side, which denatures during the extend step of PCR (72oC).
- E. (i) After extension of hairpin on wild-type DNA, extended hairpin sequence does not denature at 72oC and prevents upstream primer from generating full-length top strand. (ii) Hairpin sequence of mutant DNA does not extend, and thus denatures at 72oC, enabling upstream primer to generate full-length top strand. (iii) Likewise, top strand product denatures at 72oC, allowing polymerase to generate full-length bottom strand. The difference in hairpin extension preference of upstream primers with (i) wild-type and (ii) mutant template results in preferential amplification of mutant DNA.




Figure 155

157/182

- F. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support. Optional: wild-type sequence on end may be clipped off with restriction endonuclease.
- G. Mutation-specific ligation oligonucleotides (Mt) contain tails for subsequent FRET detection, while (optional) wild-type probe (Wt) does not. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target. (Optional: wild-type sequence on 3' end may be clipped off with exonuclease III)
- H. Ligase covalently seals the unblocked upstream and phosphorylated downstream oligonucleotides together.
- I. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.




Figure 155 (Cont.)

158/182

- A. Loop-PCR-LDR-qPCR carryover prevention reaction to detect low-level methylation. Isolate genomic or cfDNA. Treat with Bsh1236I (CG[^]CG) and UDG (37°C, 30-60 minutes) to completely digest unmethylated DNA and prevent carryover. Treat with bisulfite, which renders the strands non-complementary.
- B. Upstream locus-specific primers also contain 5' sequences complementary to bisulfite-treated unmethylated sequence of top strand, and are unblocked with RNaseH2 only when bound to target. Amplify methylation containing regions using PCR (and dUTP). Downstream primers contain identical 8-11 base tails to prevent primer dimers. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. (PCR products incorporate dU, allowing for carryover prevention.)
- D. In detail: PCR is performed with a polymerase lacking 5' nuclease, 3' nuclease, and strand-displacement activity. (i) Denaturation of unmethylated bottom strand results in hairpin with perfect match at 3' end, which is extended by polymerase. (ii) Denaturation of methylated bottom strand results in hairpin with mismatches, which generally is not extended by polymerase. (iii) Denaturation of top strand results in hairpin on 5' side, which denatures during the extend step of PCR (72°C).
- E. (i) After extension of hairpin on unmethylated DNA, extended hairpin sequence does not denature at 72°C and prevents upstream primer from generating full-length top strand. (ii) Hairpin sequence of methylated DNA does not extend, and thus denatures at 72°C, enabling upstream primer to generate full-length top strand. (iii) Likewise, top strand product denatures at 72°C, allowing polymerase to generate full-length bottom strand. The difference in hairpin extension preference of upstream primers with (i) unmethylated and (ii) methylated template results in preferential amplification of methylated DNA.




Figure 156

159/182

- F. PCR products incorporate dU, allowing for carryover prevention. Optional: sequence on end complementary to unmethylated region may be clipped off with restriction endonuclease.
- G. Methylation-specific ligation oligonucleotides (Meth) contain tags (Ai, Ci') for subsequent PCR amplification, while (optional) unmethylated probe (Umet) does not. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target. (Optional: wild-type sequence on 3' end may be clipped off with exonuclease III)
- H. Ligase covalently seals the unblocked upstream and phosphorylated downstream oligonucleotides together. Aliquot into separate wells for detection.
- I. Treat with UDG for carryover prevention, which also destroys original target amplicons. Only authentic LDR products will amplify, when using PCR (and dUTP) with tag-specific primers (Ai, Ci'). Each Taqman probe spans the ligation junction.
- J. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 156 (Cont.)

160/182

- A. Loop-PCR-qLDR carryover prevention reaction to detect low-level methylation. Isolate genomic or cfDNA. Treat with Bsh1236I (CG[^]CG) and UDG (37°C, 30-60 minutes) to completely digest unmethylated DNA and prevent carryover. Treat with bisulfite, which renders the strands non-complementary.
- B. Upstream locus-specific primers also contain 5' sequences complementary to bisulfite-treated unmethylated sequence of bottom strand, and are unblocked with RNaseH2 only when bound to target. Amplify methylation containing regions using PCR (and dUTP). Downstream primers contain universal primer tails to enable universal primer amplification to append a 5' biotin. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. (PCR products incorporate dU, allowing for carryover prevention.)
- D. In detail: PCR is performed with a polymerase lacking 5' nuclease, 3' nuclease, and strand-displacement activity. (i) Denaturation of unmethylated bottom strand results in hairpin with perfect match at 3' end, which is extended by polymerase. (ii) Denaturation of methylated bottom strand results in hairpin with mismatches, which generally is not extended by polymerase. (iii) Denaturation of top strand results in hairpin on 5' side, which denatures during the extend step of PCR (72°C).
- E. (i) After extension of hairpin on unmethylated DNA, extended hairpin sequence does not denature at 72°C and prevents upstream primer from generating full-length top strand. (ii) Hairpin sequence of methylated DNA does not extend, and thus denatures at 72°C, enabling upstream primer to generate full-length top strand. (iii) Likewise, top strand product denatures at 72°C, allowing polymerase to generate full-length bottom strand. The difference in hairpin extension preference of upstream primers with (i) unmethylated and (ii) methylated template results in preferential amplification of methylated DNA.







Figure 157

161/182

- F. PCR products incorporate dU, allowing for carryover prevention. Separate product from primers and capture biotinylated product on solid support. Optional: sequence on end complementary to unmethylated region may be clipped off with restriction endonuclease.



- G. Methyion-specific ligation oligonucleotides (Meth) contain tails for subsequent FRET detection, while (optional) unmethylated probe (Umet) does not. Upstream ligation primer contains block, which may be removed by cleaving ribose base with RNaseH2 only when hybridized to target. (Optional: wild-type sequence on 3' end may be clipped off with exonuclease III)



- H. Ligase covalently seals the unblocked upstream and phosphorylated downstream oligonucleotides together.
- I. Wash away unligated primers. Denature product from target, then detect, and enumerate FRET signal.



Figure 157 (Cont.)

162/182

- A. PCR-qPCR carryover prevention reaction to detect low-level methylation. Isolate genomic or cfDNA. Optional: enrich for methylated DNA using antibodies. Treat with methyl-sensitive restriction endonucleases Bsh1236I (CG[^]CG) and HinP1I (G[^]CGC), and UNG (37°C, 30-60 minutes) to completely digest unmethylated DNA and prevent carryover.
- B. Amplify methyl containing regions using PCR (and dUTP) with locus-specific primers. Perform limited cycle amplification (12-20) to maintain relative ratios of different amplicons. (Optional: aliquot into 12, 24, 48, or 96 wells prior to PCR.)
- C. PCR products incorporate dU, allowing for carryover prevention. (Products lack methyl groups, providing additional protection.)
- D. Optional: Aliquot into separate wells for detection. Amplify methyl containing regions using PCR (and dUTP) with nested locus-specific primers and internal Taqman probes.
- E. The 5'→3' nuclease activity of Taq polymerase liberates the fluorescent group to generate signal. (PCR products incorporate dU, allowing for carryover prevention.)




Figure 158

Detection of BRAF V600E by PCR-LDR-qPCR





Figure 159

Single molecule detection of BRAF V600E by PCR-LDR-qPCR



Single molecule detection of BRAF V600E by PCR-LDR-qPCR in Plasma




Figure 161

Detection of TP53 R248Q by PCR-LDR-qPCR




Figure 162

Detection of TP53 R248Q by PCR-LDR-qPCR




Figure 163

Single molecule detection of TP53 R248Q by PCR-LDR-qPCR




Figure 164

Detection of KRAS G12C by PCR-LDR-qPCR




Figure 166

171/182

Detection of KRAS G12S by PCR-LDR-qPCR




Figure 167

Single molecule detection of KRAS G12C by PCR-LDR-qPCR in Plasma




Figure 168

Detection of KRAS G12D by PCR-LDR-qPCR




Figure 169

Detection of KRAS G12A by PCR-LDR-qPCR




Figure 170

Detection of KRAS G12V by PCR-LDR-qPCR




Figure 171

Single molecule detection of KRAS G12V by PCR-LDR-qPCR in Plasma




Figure 172

Detection of methylation at VIM-S3 by PCR-LDR-qPCR




Figure 173

Single molecule detection of methylation of VIM_S3 by PCR-LDR-qPCR




Figure 174

Methylation Detection VIM_S3 Top strand using PCR-LDR-qPCR
and Taqman Probe version "A"




Figure 175

Methylation Detection VIM_S3 Bottom strand using PCR-LDR-qPCR
and Taqman Probe version “A”




Figure 176

Methylation Detection VIM_S3 Top strand using PCR-LDR-qPCR
and Taqman Probe version “B”




Figure 177

Methylation Detection VIM_S3 Bottom strand using PCR-LDR-qPCR
and Taqman Probe version “B”




Figure 178