

HETERODIMERIC ANTIBODIES THAT BIND CD3 AND TUMOR ANTIGENS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority under 35 U.S.C. §119(e) to U.S. Provisional Patent Application No. 62/085,117, filed November 26, 2014, U.S. Provisional Patent Application No. 62/084,908, filed November 26, 2014, U.S. Provisional Patent Application No. 62/085,027, filed November 26, 2014, U.S. Provisional Patent Application No. 62/085,106, filed November 26, 2014, U.S. Provisional Patent Application No. 62/159,111, filed May 8, 2015, U.S. Provisional Patent Application No. 62/251,005, filed November 4, 2015 and U.S. Provisional Patent Application No. 62/250,971, filed November 4, 2015, all of which are expressly incorporated herein by reference in their entirety, with particular reference to the figures, legends and claims therein.

BACKGROUND OF THE INVENTION

[0002] Antibody-based therapeutics have been used successfully to treat a variety of diseases, including cancer and autoimmune/inflammatory disorders. Yet improvements to this class of drugs are still needed, particularly with respect to enhancing their clinical efficacy. One avenue being explored is the engineering of additional and novel antigen binding sites into antibody-based drugs such that a single immunoglobulin molecule co-engages two different antigens. Such non-native or alternate antibody formats that engage two different antigens are often referred to as bispecifics. Because the considerable diversity of the antibody variable region (Fv) makes it possible to produce an Fv that recognizes virtually any molecule, the typical approach to bispecific generation is the introduction of new variable regions into the antibody.

[0003] A number of alternate antibody formats have been explored for bispecific targeting (Chames & Baty, 2009, mAbs 1[6]:1-9; Holliger & Hudson, 2005, Nature Biotechnology 23[9]:1126-1136; Kontermann, mAbs 4(2):182 (2012), all of which are expressly incorporated herein by reference). Initially, bispecific antibodies were made by fusing two cell lines that each produced a single monoclonal antibody (Milstein et al., 1983, Nature 305:537-540). Although the resulting hybrid hybridoma or quadroma did produce bispecific antibodies, they were only a minor population, and extensive purification was required to isolate the

desired antibody. An engineering solution to this was the use of antibody fragments to make bispecifics. Because such fragments lack the complex quaternary structure of a full length antibody, variable light and heavy chains can be linked in single genetic constructs. Antibody fragments of many different forms have been generated, including diabodies, single chain diabodies, tandem scFv's, and Fab₂ bispecifics (Chames & Baty, 2009, mAbs 1[6]:1-9; Holliger & Hudson, 2005, Nature Biotechnology 23[9]:1126-1136; expressly incorporated herein by reference). While these formats can be expressed at high levels in bacteria and may have favorable penetration benefits due to their small size, they clear rapidly *in vivo* and can present manufacturing obstacles related to their production and stability. A principal cause of these drawbacks is that antibody fragments typically lack the constant region of the antibody with its associated functional properties, including larger size, high stability, and binding to various Fc receptors and ligands that maintain long half-life in serum (i.e. the neonatal Fc receptor FcRn) or serve as binding sites for purification (i.e. protein A and protein G).

[0004] More recent work has attempted to address the shortcomings of fragment-based bispecifics by engineering dual binding into full length antibody -like formats (Wu et al., 2007, Nature Biotechnology 25[11]:1290-1297; USSN12/477,711; Michaelson et al., 2009, mAbs 1[2]:128-141; PCT/US2008/074693; Zuo et al., 2000, Protein Engineering 13[5]:361-367; USSN09/865,198; Shen et al., 2006, J Biol Chem 281[16]:10706-10714; Lu et al., 2005, J Biol Chem 280[20]:19665-19672; PCT/US2005/025472; expressly incorporated herein by reference). These formats overcome some of the obstacles of the antibody fragment bispecifics, principally because they contain an Fc region. One significant drawback of these formats is that, because they build new antigen binding sites on top of the homodimeric constant chains, binding to the new antigen is always bivalent.

[0005] For many antigens that are attractive as co-targets in a therapeutic bispecific format, the desired binding is monovalent rather than bivalent. For many immune receptors, cellular activation is accomplished by cross-linking of a monovalent binding interaction. The mechanism of cross-linking is typically mediated by antibody/antigen immune complexes, or via effector cell to target cell engagement. For example, the low affinity Fc gamma receptors (FcγRs) such as FcγRIIa, FcγRIIb, and FcγRIIIa bind monovalently to the antibody

Fc region. Monovalent binding does not activate cells expressing these FcγRs; however, upon immune complexation or cell-to-cell contact, receptors are cross-linked and clustered on the cell surface, leading to activation. For receptors responsible for mediating cellular killing, for example FcγRIIIa on natural killer (NK) cells, receptor cross-linking and cellular activation occurs when the effector cell engages the target cell in a highly avid format (Bowles & Weiner, 2005, *J Immunol Methods* 304:88-99, expressly incorporated by reference).. Similarly, on B cells the inhibitory receptor FcγRIIb downregulates B cell activation only when it engages into an immune complex with the cell surface B-cell receptor (BCR), a mechanism that is mediated by immune complexation of soluble IgG's with the same antigen that is recognized by the BCR (Heyman 2003, *Immunol Lett* 88[2]:157-161; Smith and Clatworthy, 2010, *Nature Reviews Immunology* 10:328-343; expressly incorporated by reference). As another example, CD3 activation of T-cells occurs only when its associated T-cell receptor (TCR) engages antigen-loaded MHC on antigen presenting cells in a highly avid cell-to-cell synapse (Kuhns et al., 2006, *Immunity* 24:133-139). Indeed nonspecific bivalent cross-linking of CD3 using an anti-CD3 antibody elicits a cytokine storm and toxicity (Perruche et al., 2009, *J Immunol* 183[2]:953-61; Chatenoud & Bluestone, 2007, *Nature Reviews Immunology* 7:622-632; expressly incorporated by reference). Thus for practical clinical use, the preferred mode of CD3 co-engagement for redirected killing of targets cells is monovalent binding that results in activation only upon engagement with the co-engaged target.

[0006] CD38, also known as cyclic ADP ribose hydrolase, is a type II transmembrane glycoprotein with a long C-terminal extracellular domain and a short N-terminal cytoplasmic domain. Among hematopoietic cells, an assortment of functional effects have been ascribed to CD38 mediated signaling, including lymphocyte proliferation, cytokine release, regulation of B and myeloid cell development and survival, and induction of dendritic cell maturation. CD38 is unregulated in many hematopoeitic malignancies and in cell lines derived from various hematopoietic malignancies including non-Hodgkin's lymphoma (NHL), Burkitt's lymphoma (BL), multiple myeloma (MM), B chronic lymphocytic leukemia (B-CLL), B and T acute lymphocytic leukemia (ALL), T cell lymphoma (TCL), acute myeloid leukemia (AML), hairy cell leukemia (HCL), Hodgkin's

Lymphoma (HL), and chronic myeloid leukemia (CML). On the other hand, most primitive pluripotent stem cells of the hematopoietic system are CD38-. In spite of the recent progress in the discovery and development of anti-cancer agents, many forms of cancer involving CD38-expressing tumors still have a poor prognosis. Thus, there is a need for improved methods for treating such forms of cancer.

[0007] B-cell antigen CD19 (CD19, also known as B-cell surface antigen B4, Leu-12) is a human pan-B-cell surface marker that is expressed from early stages of pre-B cell development through terminal differentiation into plasma cells. CD 19 promotes the proliferation and survival of mature B cells. It associates in a complex with CD21 on the cell surface. It also associates with CD81 and Leu-13 and potentiates B cell receptor (BCR) signaling. Together with the BCR, CD19 modulates intrinsic and antigen receptor-induced signaling thresholds critical for clonal expansion of B cells and humoral immunity. In collaboration with CD21 it links the adaptive and the innate immune system. Upon activation, the cytoplasmic tail of CD19 becomes phosphorylated which leads to binding by Src-family kinases and recruitment of PI-3 kinase. It is an attractive immunotherapy target for cancers of lymphoid origin since it is also expressed on the vast majority of NHL cells as well as some leukemias.

[0008] A number of antibodies or antibody conjugates that target CD19 have been evaluated in pre-clinical studies or in clinical trials for the treatment of cancers. These anti-CD19 antibodies or antibody conjugates include but are not limited to MT-103 (a single-chain bispecific CD19/CD3 antibody; Hoffman et al, 2005 Int J Cancer 115:98-104; Schlereth et al, 2006 Cancer Immunol Immunother 55:503-514), a CD19/CD16 diabody (Schlenzka et al, 2004 Anti-cancer Drugs 15:915-919; Kipriyanov et al, 2002 J Immunol 169:137-144), BU12-saporin (Flavell et al, 1995 Br J Cancer 72:1373-1379), and anti-CD19-idarubicin (Rowland et al, 1993 Cancer Immunol Immunother 55:503-514); all expressly incorporated by reference.

[0009] CD123, also known as interleukin-3 receptor alpha (IL-3R α), is expressed on dendritic cells, monocytes, eosinophils and basophils. CD123 is also constitutively expressed by committed hematopoietic stem/progenitor cells, by most of the myeloid lineage (CD13+, CD14+, CD33+, CD15^{low}), and by some CD19+ cells. It is absent from CD3+ cells.

[0010] Thus while bispecifics generated from antibody fragments suffer biophysical and pharmacokinetic hurdles, a drawback of those built with full length antibody -like formats is that they engage co-target antigens multivalently in the absence of the primary target antigen, leading to nonspecific activation and potentially toxicity. The present invention solves this problem by introducing novel bispecific antibodies directed to CD3 and CD38.

BRIEF SUMMARY OF THE INVENTION

[0011] Accordingly, in one aspect the present invention provides heterodimeric antibodies comprising: a) a first monomer comprising: i) a first heavy chain comprising: 1) a first variable heavy domain; 2) a first constant heavy chain comprising a first Fc domain; 3) a scFv comprising a scFv variable light domain, an scFv linker and a scFv variable heavy domain; wherein said scFv is covalently attached to the C-terminus of said Fc domain using a domain linker; b) a second monomer comprising a second heavy chain comprising a second variable heavy domain and a second constant heavy chain comprising a second Fc domain; and c) a common light chain comprising a variable light domain and a constant light domain.

[0012] In a further aspect, the invention provides heterodimeric antibodies comprising: a) a first monomer comprising: i) a first heavy chain comprising: 1) a first variable heavy domain; 2) a first constant heavy domain comprising a first Fc domain; and 3) a first variable light domain, wherein said first variable light domain is covalently attached to the C-terminus of said first Fc domain using a domain linker; b) a second monomer comprising: i) a second variable heavy domain; ii) a second constant heavy domain comprising a second Fc domain; and iii) a third variable heavy domain, wherein said second variable heavy domain is covalently attached to the C-terminus of said second Fc domain using a domain linker; c) a common light chain comprising a variable light domain and a constant light domain.

[0013] In an additional aspect, the invention provides heterodimeric antibodies comprising: a) a first monomer comprising: i) a first heavy chain comprising: 1) a first variable heavy domain; 2) a first constant heavy chain comprising a first CH1 domain and a first Fc domain; 3) a scFv comprising a scFv variable light domain, an scFv linker and a scFv variable heavy domain; wherein said scFv is covalently attached between the C-terminus of said CH1

domain and the N-terminus of said first Fc domain using domain linkers; b) a second monomer comprising a second heavy chain comprising a second variable heavy domain and a second constant heavy chain comprising a second Fc domain; and c) a common light chain comprising a variable light domain and a constant light domain.

[0014] In a further aspect, the invention provides heterodimeric antibodies comprising: a) a first monomer comprising: i) a first heavy chain comprising: 1) a first variable heavy domain; 2) a first constant heavy domain comprising a first Fc domain; and 3) a first variable light domain, wherein said second variable light domain is covalently attached between the C-terminus of the CH1 domain of said first constant heavy domain and the N-terminus of said first Fc domain using domain linkers; b) a second monomer comprising: i) a second variable heavy domain; ii) a second constant heavy domain comprising a second Fc domain; and iii) a third variable heavy domain, wherein said second variable heavy domain is covalently attached to the C-terminus of said second Fc domain using a domain linker; c) a common light chain comprising a variable light domain and a constant light domain.

[0015] In an additional aspect, the invention provides heterodimeric antibodies comprising: a) a first monomer comprising: i) a first heavy chain comprising: 1) a first variable heavy domain; 2) a first constant heavy chain comprising a first CH1 domain and a first Fc domain; 3) a scFv comprising a scFv variable light domain, an scFv linker and a scFv variable heavy domain; wherein said scFv is covalently attached between the C-terminus of said CH1 domain and the N-terminus of said first Fc domain using domain linkers; b) a second monomer comprising a second Fc domain; and c) a light chain comprising a variable light domain and a constant light domain.

[0016] In some aspects, the first and second Fc domains have a set of amino acid substitutions selected from the group consisting of S364K/E357Q : L368D/K370S; L368D/K370S : S364K; L368E/K370S : S364K; T411T/E360E/Q362E : D401K; L368D/K370S : S364K/E357L and K370S : S364K/E357Q. Furthermore, the variable heavy domain(s) and the variable light domain(s) bind a first target tumor antigen (TTA), the scFv binds a second TTA or human CD3. In some embodiments, the TTA is selected from the group consisting of CD19, CD20 and CD123.

[0017] In a further aspect, the invention provides anti-CD3 antigen binding domains have CDRs and/or the variable domains and/or the scFv sequences depicted in the Figures for H1.32_L1.47, H1.89_L1.47, H1.90_L1.47, H1.33_L1.47 and H1.31_L1.47. The invention further provides nucleic acid compositions, expression vector compositions and host cells.

[0018] In an additional aspect, the invention provides heterodimeric antibodies comprising a) a first monomer comprising: i) a first Fc domain; ii) an anti-CD3 scFv comprising a scFv variable light domain, an scFv linker and a scFv variable heavy domain; wherein said scFv is covalently attached to the N-terminus of said Fc domain using a domain linker; b) a second monomer comprising a heavy chain comprising: i) a heavy variable domain; and ii) a heavy chain constant domain comprising a second Fc domain; and c) a light chain comprising a variable light domain and a variable light constant domain; wherein the anti-CD3 scFv is selected from the group consisting of anti-CD3 H1.32_L1.47, anti-CD3 H1.89_L1.47, anti-CD3 H1.90_L1.47 and anti-CD3 H1.33_L1.47 (SEQ ID NO:XX). The heavy variable domain and the light variable domain bind a TTA (including, but not limited to CD19, Cd20, CD38 and CD123).

[0019] In an additional aspect, the invention provides anti-CD20 antibody binding domains comprising : a) a variable light domain comprising a vlCDR1 having the sequence RASWSVSYIH (SEQ ID NO:XX), a vlCDR2 having the sequence ATSNLAS (SEQ ID NO:XX), and a vlCDR3 having the sequence QQWTHNPPT (SEQ ID NO:XX); and b) a variable heavy domain comprises a vhCDR1 having the sequence SYNMH (SEQ ID NO:XX), a vhCDR2 having the sequence AIYPGNGATSYSQKFQG (SEQ ID NO:XX) and a vhCDR3 having the sequence SYYMGGDWYFDV (SEQ ID NO:XX). In some embodiments, the anti-CD20 antibody binding domains have the C2B8 H1.202_L1.113 sequences.

[0020] In an additional aspect, the invention provides anti-CD20 antibody binding domains comprising: a) a variable light domain comprising a vlCDR1 having the sequence RASSVSYIH (SEQ ID NO:XX), a vlCDR2 having the sequence ATSNLAS (SEQ ID NO:XX), and a vlCDR3 having the sequence QQWTSNPPT (SEQ ID NO:XX); and b) a variable heavy domain comprises a vhCDR1 having the sequence SYNMH (SEQ ID NO:XX), a vhCDR2 having the sequence AIYPGNGDTSYNQKFQG (SEQ ID NO:XX) and a vhCDR3 having the sequence STYYGGDWYFNV (SEQ ID NO:XX).

[0021] In some embodiments, the anti-CD20 antibody binding domains have the C2B8_H1L1 sequences.

[0022] In an additional aspect, the invention provides heterodimeric antibodies comprising a) a first monomer comprising: i) a first Fc domain; ii) an anti-CD3 scFv comprising a scFv variable light domain, an scFv linker and a scFv variable heavy domain; wherein said scFv is covalently attached to the N-terminus of said Fc domain using a domain linker; b) a second monomer comprising a heavy chain comprising: i) a heavy variable domain; and ii) a heavy chain constant domain comprising a second Fc domain; and c) a light chain comprising a variable light domain and a variable light constant domain; wherein the variable heavy and light chains form a C2B8 H1.202_L1.113 or C2B8_H1L1 binding domain.

[0023] In an additional aspect, the invention provides heterodimeric antibodies comprising a) a first monomer comprising: i) a first Fc domain; ii) an anti-CD3 scFv comprising a scFv variable light domain, an scFv linker and a scFv variable heavy domain; wherein said scFv is covalently attached to the N-terminus of said Fc domain using a domain linker; b) a second monomer comprising a heavy chain comprising: i) a heavy variable domain; and ii) a heavy chain constant domain comprising a second Fc domain; and c) a light chain comprising a variable light domain and a variable light constant domain. In this embodiment, the variable domains bind CD123 and can have the sequences of 7G3_H1.109_L1.47.

[0024] In additional aspects, the present invention provides heterodimeric antibodies selected from the group consisting of XENP15049, XENP15051; XENP15050, XENP13676, XENP14696, XENP15629, XENP15053, XENP15630, XENP15631, XENP15632, XENP15633, XENP15634, XENP15635, XENP15636, XENP15638, XENP15639, XENP13677, XENP14388, XENP14389, XENP14390, XENP14391, XENP14392, XENP14393, XENP16366, XENP16367, XENP16368, XENP16369, XENP16370, XENP16371, XENP16372, XENP16373, XENP16375, XENP16376 and XENP16377. Nucleic acids, expression vectors and host cells are all provided as well, in addition to methods of making these proteins and treating patients with them.

BRIEF DESCRIPTION OF THE DRAWINGS

[0025] Figures 1A and 1B depict several formats of the present invention. Two forms of the “bottle opener” format are depicted, one with the anti-CD3 antigen binding domain comprising a scFv and the anti-TTA antigen binding domain comprising a Fab, and one with these reversed. The mAb-Fv, mAb-scFv, Central-scFv and Central-Fv formats are all shown. In addition, “one-armed” formats, where one monomer just comprises an Fc domain, are shown, both a one arm Central-scFv and a one arm Central-Fv. A dual scFv format is also shown.

[0026] Figure 2 depicts the sequences of the “High CD3” anti-CD3_H1.30_L1.47 construct, including the variable heavy and light domains (CDRs underlined), as well as the individual vl and vhCDRs, as well as an scFv construct with a charged linker (double underlined). As is true of all the sequences depicted in the Figures, this charged linker may be replaced by an uncharged linker or a different charged linker, as needed.

[0027] Figure 3 depicts the sequences of the “High-Int #1” Anti-CD3_H1.32_L1.47 construct, including the variable heavy and light domains (CDRs underlined), as well as the individual vl and vhCDRs, as well as an scFv construct with a charged linker (double underlined). As is true of all the sequences depicted in the Figures, this charged linker may be replaced by an uncharged linker or a different charged linker, as needed.

[0028] Figure 4 depicts the sequences of the “High-Int #2” Anti-CD3_H1.89_L1.47 construct, including the variable heavy and light domains (CDRs underlined), as well as the individual vl and vhCDRs, as well as an scFv construct with a charged linker (double underlined). As is true of all the sequences depicted in the Figures, this charged linker may be replaced by an uncharged linker or a different charged linker, as needed.

[0029] Figure 5 depicts the sequences of the “High-Int #3” Anti-CD3_H1.90_L1.47 construct, including the variable heavy and light domains (CDRs underlined), as well as the individual vl and vhCDRs, as well as an scFv construct with a charged linker (double underlined). As is true of all the sequences depicted in the Figures, this charged linker may be replaced by an uncharged linker or a different charged linker, as needed.

[0030] Figure 6 depicts the sequences of the "Int" Anti-CD3_H1.90_L1.47 construct, including the variable heavy and light domains (CDRs underlined), as well as the individual vl and vhCDRs, as well as an scFv construct with a charged linker (double underlined). As is true of all the sequences depicted in the Figures, this charged linker may be replaced by an uncharged linker or a different charged linker, as needed.

[0031] Figure 7 depicts the sequences of the "Low" Anti-CD3_H1.31_L1.47 construct, including the variable heavy and light domains (CDRs underlined), as well as the individual vl and vhCDRs, as well as an scFv construct with a charged linker (double underlined). As is true of all the sequences depicted in the Figures, this charged linker may be replaced by an uncharged linker or a different charged linker, as needed.

[0032] Figure 8 depicts the sequences of the High CD38: OKT10_H1.77_L1.24 construct, including the variable heavy and light domains (CDRs underlined), as well as the individual vl and vhCDRs, as well as an scFv construct with a charged linker (double underlined).

[0033] Figure 9 depicts the sequences of the intermediate CD38: OKT10_H1L1.24 construct, including the variable heavy and light domains (CDRs underlined), as well as the individual vl and vhCDRs, as well as an scFv construct with a charged linker (double underlined).

[0034] Figure 10 depicts the sequences of the Low CD38: OKT10_H1L1 construct, including the variable heavy and light domains (CDRs underlined), as well as the individual vl and vhCDRs, as well as an scFv construct with a charged linker (double underlined).

[0035] Figure 11 depicts the sequences of XENP15331.

[0036] Figure 12 depicts the sequences of XENP13243.

[0037] Figure 13 depicts the sequences of XENP14702.

[0038] Figure 14 depicts the sequences of XENP15426.

[0039] Figure 15 depicts the sequences of XENP14701.

[0040] Figure 16 depicts the sequence of XENP14703.

[0041] Figure 17 depicts the sequence of XENP13243.

[0042] Figure 18 depicts the sequences of XENP18967.

[0043] Figure 19 depicts the sequences of XENP18971.

[0044] Figure 20 depicts the sequences of XENP18969.

[0045] Figure 21 depicts the sequences of XENP18970.

[0046] Figure 22 depicts the sequences of XENP18972.

[0047] Figure 23 depicts the sequences of XENP18973.

[0048] Figure 24 depicts the sequences of XENP15055.

[0049] Figure 25 depicts the sequences of XENP13544.

[0050] Figure 26 depicts the sequences of XENP13694.

[0051] Figure 27 depicts the sequence of human CD3 ϵ .

[0052] Figure 28 depicts the full length (SEQ ID NO:130) and extracellular domain (ECD; SEQ ID NO:131) of the human CD38 protein.

[0053] Figure 29A -29E depict useful pairs of heterodimerization variant sets (including skew and pI variants).

[0054] .Figure 30 depict a list of isosteric variant antibody constant regions and their respective substitutions. pI₋(-) indicates lower pI variants, while pI₊(+) indicates higher pI variants. These can be optionally and independently combined with other heterodimerization variants of the invention (and other variant types as well, as outlined herein).

[0055] Figure 31 depict useful ablation variants that ablate Fc γ R binding (sometimes referred to as "knock outs" or "KO" variants).

[0056] Figure 32 show two particularly useful embodiments of the invention.

[0057] Figure 33 depicts a number of charged scFv linkers that find use in increasing or decreasing the pI of heterodimeric antibodies that utilize one or more scFv as a component. A single prior art scFv linker with a single charge is referenced as "Whitlow", from Whitlow et al., Protein Engineering 6(8):989-995 (1993). It should be noted that this linker was used for reducing aggregation and enhancing proteolytic stability in scFvs.

[0058] Figure 34 depicts a list of engineered heterodimer-skewing Fc variants with heterodimer yields (determined by HPLC-CIEX) and thermal stabilities (determined by DSC). Not determined thermal stability is denoted by "n.d.".

[0059] Figure 35 Expression yields of bispecifics after protein A affinity purification.

[0060] Figure 36 Cationic exchange purification chromatograms.

[0061] Figure 37 Redirected T cell cytotoxicity assay, 24 h incubation, 10k RPMI8226 cells, 400k T cells. Test articles are anti-CD38 x anti-CD3 bispecifics. Detection was by LDH

[0062] Figure 38 Redirected T cell cytotoxicity assay, 24 h incubation, 10k RPMI8226 cells, 500k human PBMCs. Test articles are anti-CD38 x anti-CD3 bispecifics. Detection was by LDH.

[0063] Figure 39 depicts the sequences of XENP14419,

[0064] Figure 40 depicts the sequences of XENP14420.

[0065] Figure 41 depicts the sequences of XENP14421.

[0066] Figure 42 depicts the sequences of XENP14422.

[0067] Figure 43 depicts the sequences of XENP14423.

[0068] Figure 44 Redirected T cell cytotoxicity assay, 96 h incubation, 40k RPMI8226 cells, 400k human PBMC. Test articles are anti-CD38 x anti-CD3 Fab-scFv-Fcs. Detection was by flow cytometry, specifically the disappearance of CD38+ cells.

[0069] Figure 45 Further analysis of redirected T cell cytotoxicity assay described in Figure 1. The first row shows the Mean Fluorescence Intensity (MFI) of activation marker CD69 on CD4+ and CD8+ T cells as detected by flow cytometry. The second row shows the percentage of CD4+ and CD8+ T cells that are Ki-67+, a measure of cell proliferation. The third row shows the intracellular Mean Fluorescence Intensity (MFI) of granzyme B inhibitor PI-9 on CD4+ and CD8+ T cells as detected by flow cytometry.

[0070] Figure 46 Design of mouse study to examine anti-tumor activity of anti-CD38 x anti-CD3 Fab-scFv-Fc bispecifics.

[0071] Figure 47 Tumor size measured by IVIS® as a function of time and treatment

[0072] Figure 48 IVIS® bioluminescent images (Day 10)

[0073] Figure 49 Depletion of CD38⁺ cells in cynomolgus monkeys following single doses of the indicated test articles

[0074] Figure 50 T cell activation measured by CD69 Mean Fluorescence Intensity (MFI) in cynomolgus monkeys, color coding as in Figure 49.

[0075] Figure 51 Serum levels of IL-6, following single doses of the indicated test articles.

[0076] Figure 52 depicts the sequences of XENP15427.

[0077] Figure 53 depicts the sequences of XENP15428.

[0078] Figure 54 depicts the sequences of XENP15429.

[0079] Figure 55 depicts the sequences of XENP15430.

[0080] Figure 56 depicts the sequences of XENP15431.

[0081] Figure 57 depicts the sequences of XENP15432.

[0082] Figure 58 depicts the sequences of XENP15433.

[0083] Figure 59 depicts the sequences of XENP15434.

[0084] Figure 60 depicts the sequences of XENP15435.

[0085] Figure 61 depicts the sequences of XENP15436.

[0086] Figure 62 depicts the sequences of XENP15437.

[0087] Figure 63 depicts the sequences of XENP15438.

[0088] Figure 64 shows binding affinities in a Biacore assay.

[0089] Figure 65 shows the Heterodimer purity during stable pool generation using varied Light chain, Fab-Fc, and scFv-Fc ratios.

[0090] Figure 66 Human IgM and IgG2 depletion by anti-CD38 x anti-CD3 bispecifics in a huPBMC mouse model.

[0091] Figure 67 depicts stability-optimized, humanized anti-CD3 variant scFvs.

Substitutions are given relative to the H1_L1.4 scFv sequence. Amino acid numbering is Kabat numbering.

[0092] Figure 68. Amino acid sequences of stability-optimized, humanized anti-CD3 variant scFvs. CDRs are underlined. For each heavy chain/light chain combination, four sequences are listed: (i) scFv with C-terminal 6xHis tag, (ii) scFv alone, (iii) VH alone, (iv) VL alone.

[0093] Figure 69 Redirected T cell cytotoxicity assay, 24 h incubation, 10k RPMI8226 cells, 500k PBMC. Test articles are anti-CD38 (OKT10_H1L1, OKT10_H1.77_L1.24) x anti-CD3 Fab-scFv-Fcs. Detection was by LDH.

[0094] Figure 70 huPBL-SCID Ig-depletion study. Test articles were dosed 8 d after PBMC engraftment at 0.03, 0.3, or 3 mg/kg. Route of administration was intraperitoneal. Blood samples were taken 14 d after PBMC engraftment, processed to serum, and assayed for human IgM and IgG2.

[0095] Figure 71 depicts the sequences of XENP15049.

[0096] Figure 72 depicts the sequences of XENP15051.

[0097] Figure 73 depicts the sequences of XENP15050.

[0098] Figure 74 depicts the sequences of XENP13676.

[0099] Figure 75 depicts the sequences of XENP14696.

[00100] Figure 76 depicts the sequences of XENP15629.

[00101] Figure 77 depicts the sequences of XENP15053.

[00102] Figure 78 depicts the sequences of XENP15630.

[00103] Figure 79 depicts the sequences of XENP15631.

[00104] Figure 80 depicts the sequences of XENP15632.

[00105] Figure 81 depicts the sequences of XENP15633.

[00106] Figure 82 depicts the sequences of XENP15634.

[00107] Figure 83 depicts the sequences of XENP15635.

[00108] Figure 84 depicts the sequences of XENP15636.

[00109] Figure 85 depicts the sequences of XENP15638.

[00110] Figure 86 depicts the sequences of XENP15639.

- [00111] Figure 87 depicts the sequences of XENP13677.
- [00112] Figure 88 depicts the sequences of XENP14388.
- [00113] Figure 89 depicts the sequences of XENP14389.
- [00114] Figure 90 depicts the sequences of XENP14390.
- [00115] Figure 91 depicts the sequences of XENP14391.
- [00116] Figure 92 depicts the sequences of XENP14392.
- [00117] Figure 93 depicts the sequences of XENP14393.
- [00118] Figure 94 depicts the sequences of XENP16366.
- [00119] Figure 95 depicts the sequences of XENP16367.
- [00120] Figure 96 depicts the sequences of XENP16368.
- [00121] Figure 97 depicts the sequences of XENP16369.
- [00122] Figure 98 depicts the sequences of XENP16370.
- [00123] Figure 99 depicts the sequences of XENP16371.
- [00124] Figure 100 depicts the sequences of XENP16372.
- [00125] Figure 101 depicts the sequences of XENP16373.
- [00126] Figure 102 depicts the sequences of XENP16374.
- [00127] Figure 103 depicts the sequences of XENP16375.
- [00128] Figure 104 depicts the sequences of XENP16376.
- [00129] Figure 105 depicts the sequences of XENP16377.
- [00130] Figure 106 depicts the sequences of the CD20 and CD123 antigens.
- [00131] Figure 107 Surface plasmon resonance determination of CD3 affinity. Test articles are anti-CD20 (C2B8_H1.202_L1.113) x anti-CD3 Fab-scFv-Fcs. Human CD3 $\delta\epsilon$ -Fc (Sino Biological) was covalently bound to the chip surface. Test articles were passed over at 3.125, 12.5, 50, and 200 nM.

[00132] Figure 108 Surface plasmon resonance determination of CD3 affinity. Test articles are anti-CD20 (C2B8_H1.202_L1.113) x anti-CD3 Fab-scFv-Fcs. Cynomolgus monkey CD3 $\delta\epsilon$ -Fc (Sino Biological) was covalently bound to the chip surface. Test articles were passed over at 3.125, 12.5, 50, and 200 nM.

[00133] Figure 109 Surface plasmon resonance determination of CD3 affinity. Test articles are anti-CD20 (C2B8_H1.202_L1.113) x anti-CD3 Fab-scFv-Fcs. Human CD3 $\delta\epsilon$ -Fc (Sino Biological) was covalently bound to the chip surface. Test articles were passed over at 31.25, 125, 500, and 2000 nM.

[00134] Figure 110 Surface plasmon resonance determination of CD3 affinity. Test articles are anti-CD20 (C2B8_H1.202_L1.113) x anti-CD3 Fab-scFv-Fcs. Cynomolgus monkey CD3 $\delta\epsilon$ -Fc (Sino Biological) was covalently bound to the chip surface. Test articles were passed over at 31.25, 125, 500, and 2000 nM.

[00135] Figure 111 Surface plasmon resonance determination of CD3 affinity. Test articles are anti-CD20 (C2B8_H1.202_L1.113) x anti-CD3 Fab-scFv-Fcs. Cynomolgus monkey CD3 $\delta\epsilon$ -Fc (Sino Biological) was covalently bound to the chip surface. Test articles were passed over at 31.25, 125, 500, and 2000 nM.

[00136] Figure 112 Redirected T cell cytotoxicity assay, 24 h incubation, 10k Ramos cells, 250k PBMC. Test articles are anti-CD20 (C2B8_H1.202_L1.113) x anti-CD3 Fab-scFv-Fcs. Detection was by LDH.

[00137] Figure 113 Redirected T cell cytotoxicity assay, 24 h incubation, 20k Jeko cells, 200k PBMC (CD19-depleted). Test articles are anti-CD20 (C2B8_H1.202_L1.113) x anti-CD3 Fab-scFv-Fcs. Detection was by flow cytometry, specifically the disappearance of CD19⁺ cells.

[00138] Figure 114 IL-6 production after 24 h for the experiment described in Figure 113.

[00139] Figure 115 Redirected T cell cytotoxicity assay, 5 h incubation, 20k Jeko cells, 500k PBMC (CD19-depleted). Test articles are anti-CD20 (C2B8_H1L1) x anti-CD3 Fab-scFv-Fcs. Detection was by flow cytometry, specifically the disappearance of CD19⁺ cells.

[00140] Figure 116 Redirected T cell cytotoxicity assay, 24 h incubation, 20k Jeko cells, 500k PBMC (CD19-depleted). Test articles are anti-CD20 (C2B8_H1.202_L1.113) x anti-CD3 Fab-scFv-Fcs. Detection was by flow cytometry, specifically the disappearance of CD19⁺ cells.

[00141] Figure 117 IL-6 production after 24 h for the experiment described in Figure 113.

[00142] Figure 118 Redirected T cell cytotoxicity assay, 24 h incubation, 10k RPMI8226 cells, 500k PBMC. Test articles are anti-CD38 (OKT10_H1L1, OKT10_H1.77_L1.24) x anti-CD3 Fab-scFv-Fcs. Detection was by LDH.

[00143] Figure 119 huPBL-SCID Ig-depletion study. Test articles were dosed 1 and 8 d after PBMC engraftment at 5 mg/kg. Route of administration was intraperitoneal. Blood samples were taken 14 d after PBMC engraftment, processed to serum, and assayed for human IgM and IgG2.

[00144] Figure 120 huPBL-SCID Ig-depletion study. Test articles were dosed 8 d after PBMC engraftment at 0.03, 0.3, or 3 mg/kg. Route of administration was intraperitoneal. Blood samples were taken 14 d after PBMC engraftment, processed to serum, and assayed for human IgM and IgG2.

[00145] Figure 121 depicts the sequences of High CD20 C2B8_H1.202_L1.113.

[00146] Figure 122 depicts the sequences of Low CD20 C2B8_H1L1.

[00147] Figure 123 depicts the sequences of CD123 7G3_H1.109_L1.57.

[00148] Figure 124 shows a matrix of possible combinations for the invention. An "A" means that the CDRs of the referenced CD3 sequences can be combined with the CDRs of the TTA on the right hand side. That is, the vhCDRs from the variable heavy chain CD3 H1.30 sequence and the vlCDRs from the variable light chain of CD3 L1.57 sequence can be combined with the vhCDRs from the CD38 OKT10 H1.77 sequence and the vlCDRs from the OKT10L1.24 sequence. A "B" means that the CDRs from the CD3 constructs can be combined with the variable heavy and light domains from the TTA. That is, the vhCDRs from the variable heavy chain CD3 H1.30 sequence and the vlCDRs from the variable light

chain of CD3 L1.57 sequence can be combined with the variable heavy domain CD38 OKT10 H1.77 sequence and the OKT10L1.24 sequence. A "C" is reversed, such that the variable heavy domain and variable light domain from the CD3 sequences are used with the CDRs of the TTAs. A "D" is where both the variable heavy and variable light chains from each are combined. An "E" is where the scFv of the CD3 is used with the CDRs of the TTA, and an "F" is where the scFv of the CD3 is used with the variable heavy and variable light domains of the TTA antigen binding domain.

DETAILED DESCRIPTION OF THE INVENTION

I. Definitions

[00149] In order that the application may be more completely understood, several definitions are set forth below. Such definitions are meant to encompass grammatical equivalents.

[00150] By "ablation" herein is meant a decrease or removal of activity. Thus for example, "ablating FcγR binding" means the Fc region amino acid variant has less than 50% starting binding as compared to an Fc region not containing the specific variant, with less than 70-80-90-95-98% loss of activity being preferred, and in general, with the activity being below the level of detectable binding in a Biacore assay. Of particular use in the ablation of FcγR binding are those shown in Figure 16.

[00151] By "ADCC" or "antibody dependent cell-mediated cytotoxicity" as used herein is meant the cell-mediated reaction wherein nonspecific cytotoxic cells that express FcγRs recognize bound antibody on a target cell and subsequently cause lysis of the target cell. ADCC is correlated with binding to FcγRIIIa; increased binding to FcγRIIIa leads to an increase in ADCC activity.

[00152] By "ADCP" or antibody dependent cell-mediated phagocytosis as used herein is meant the cell-mediated reaction wherein nonspecific cytotoxic cells that express FcγRs recognize bound antibody on a target cell and subsequently cause phagocytosis of the target cell.

[00153] By "modification" herein is meant an amino acid substitution, insertion, and/or deletion in a polypeptide sequence or an alteration to a moiety chemically linked to a

protein. For example, a modification may be an altered carbohydrate or PEG structure attached to a protein. By "amino acid modification" herein is meant an amino acid substitution, insertion, and/or deletion in a polypeptide sequence. For clarity, unless otherwise noted, the amino acid modification is always to an amino acid coded for by DNA, e.g. the 20 amino acids that have codons in DNA and RNA.

[00154] By "amino acid substitution" or "substitution" herein is meant the replacement of an amino acid at a particular position in a parent polypeptide sequence with a different amino acid. In particular, in some embodiments, the substitution is to an amino acid that is not naturally occurring at the particular position, either not naturally occurring within the organism or in any organism. For example, the substitution E272Y refers to a variant polypeptide, in this case an Fc variant, in which the glutamic acid at position 272 is replaced with tyrosine. For clarity, a protein which has been engineered to change the nucleic acid coding sequence but not change the starting amino acid (for example exchanging CGG (encoding arginine) to CGA (still encoding arginine) to increase host organism expression levels) is not an "amino acid substitution"; that is, despite the creation of a new gene encoding the same protein, if the protein has the same amino acid at the particular position that it started with, it is not an amino acid substitution.

[00155] By "amino acid insertion" or "insertion" as used herein is meant the addition of an amino acid sequence at a particular position in a parent polypeptide sequence. For example, -233E or 233E designates an insertion of glutamic acid after position 233 and before position 234. Additionally, -233ADE or A233ADE designates an insertion of AlaAspGlu after position 233 and before position 234.

[00156] By "amino acid deletion" or "deletion" as used herein is meant the removal of an amino acid sequence at a particular position in a parent polypeptide sequence. For example, E233- or E233# or E233() designates a deletion of glutamic acid at position 233. Additionally, EDA233- or EDA233# designates a deletion of the sequence GluAspAla that begins at position 233.

[00157] By "variant protein" or "protein variant", or "variant" as used herein is meant a protein that differs from that of a parent protein by virtue of at least one amino acid

modification. Protein variant may refer to the protein itself, a composition comprising the protein, or the amino sequence that encodes it. Preferably, the protein variant has at least one amino acid modification compared to the parent protein, e.g. from about one to about seventy amino acid modifications, and preferably from about one to about five amino acid modifications compared to the parent. As described below, in some embodiments the parent polypeptide, for example an Fc parent polypeptide, is a human wild type sequence, such as the Fc region from IgG1, IgG2, IgG3 or IgG4, although human sequences with variants can also serve as "parent polypeptides", for example the IgG1/2 hybrid of Figure 19. The protein variant sequence herein will preferably possess at least about 80% identity with a parent protein sequence, and most preferably at least about 90% identity, more preferably at least about 95-98-99% identity. Variant protein can refer to the variant protein itself, compositions comprising the protein variant, or the DNA sequence that encodes it. Accordingly, by "antibody variant" or "variant antibody" as used herein is meant an antibody that differs from a parent antibody by virtue of at least one amino acid modification, "IgG variant" or "variant IgG" as used herein is meant an antibody that differs from a parent IgG (again, in many cases, from a human IgG sequence) by virtue of at least one amino acid modification, and "immunoglobulin variant" or "variant immunoglobulin" as used herein is meant an immunoglobulin sequence that differs from that of a parent immunoglobulin sequence by virtue of at least one amino acid modification. "Fc variant" or "variant Fc" as used herein is meant a protein comprising an amino acid modification in an Fc domain. The Fc variants of the present invention are defined according to the amino acid modifications that compose them. Thus, for example, N434S or 434S is an Fc variant with the substitution serine at position 434 relative to the parent Fc polypeptide, wherein the numbering is according to the EU index. Likewise, M428L/N434S defines an Fc variant with the substitutions M428L and N434S relative to the parent Fc polypeptide. The identity of the WT amino acid may be unspecified, in which case the aforementioned variant is referred to as 428L/434S. It is noted that the order in which substitutions are provided is arbitrary, that is to say that, for example, 428L/434S is the same Fc variant as M428L/N434S, and so on. For all positions discussed in the present invention that relate to antibodies, unless otherwise noted, amino acid position numbering is according to the EU index. The EU index or EU

index as in Kabat or EU numbering scheme refers to the numbering of the EU antibody (Edelman et al., 1969, Proc Natl Acad Sci USA 63:78-85, hereby entirely incorporated by reference.) The modification can be an addition, deletion, or substitution. Substitutions can include naturally occurring amino acids and, in some cases, synthetic amino acids. Examples include U.S. Pat. No. 6,586,207; WO 98/48032; WO 03/073238; US2004-0214988A1; WO 05/35727A2; WO 05/74524A2; J. W. Chin et al., (2002), Journal of the American Chemical Society 124:9026-9027; J. W. Chin, & P. G. Schultz, (2002), ChemBioChem 11:1135-1137; J. W. Chin, et al., (2002), PICALO United States of America 99:11020-11024; and, L. Wang, & P. G. Schultz, (2002), Chem. 1-10, all entirely incorporated by reference.

[00158] As used herein, "protein" herein is meant at least two covalently attached amino acids, which includes proteins, polypeptides, oligopeptides and peptides. The peptidyl group may comprise naturally occurring amino acids and peptide bonds, or synthetic peptidomimetic structures, i.e. "analogs", such as peptoids (see Simon et al., PNAS USA 89(20):9367 (1992), entirely incorporated by reference). The amino acids may either be naturally occurring or synthetic (e.g. not an amino acid that is coded for by DNA); as will be appreciated by those in the art. For example, homo-phenylalanine, citrulline, ornithine and noreucine are considered synthetic amino acids for the purposes of the invention, and both D- and L-(R or S) configured amino acids may be utilized. The variants of the present invention may comprise modifications that include the use of synthetic amino acids incorporated using, for example, the technologies developed by Schultz and colleagues, including but not limited to methods described by Cropp & Shultz, 2004, Trends Genet. 20(12):625-30, Anderson et al., 2004, Proc Natl Acad Sci USA 101 (2):7566-71, Zhang et al., 2003, 303(5656):371-3, and Chin et al., 2003, Science 301(5635):964-7, all entirely incorporated by reference. In addition, polypeptides may include synthetic derivatization of one or more side chains or termini, glycosylation, PEGylation, circular permutation, cyclization, linkers to other molecules, fusion to proteins or protein domains, and addition of peptide tags or labels.

[00159] By "residue" as used herein is meant a position in a protein and its associated amino acid identity. For example, Asparagine 297 (also referred to as Asn297 or N297) is a residue at position 297 in the human antibody IgG1.

[00160] By "Fab" or "Fab region" as used herein is meant the polypeptide that comprises the VH, CH1, VL, and CL immunoglobulin domains. Fab may refer to this region in isolation, or this region in the context of a full length antibody, antibody fragment or Fab fusion protein. By "Fv" or "Fv fragment" or "Fv region" as used herein is meant a polypeptide that comprises the VL and VH domains of a single antibody. As will be appreciated by those in the art, these generally are made up of two chains.

[00161] By "IgG subclass modification" or "isotype modification" as used herein is meant an amino acid modification that converts one amino acid of one IgG isotype to the corresponding amino acid in a different, aligned IgG isotype. For example, because IgG1 comprises a tyrosine and IgG2 a phenylalanine at EU position 296, a F296Y substitution in IgG2 is considered an IgG subclass modification.

[00162] By "non-naturally occurring modification" as used herein is meant an amino acid modification that is not isotypic. For example, because none of the IgGs comprise a serine at position 434, the substitution 434S in IgG1, IgG2, IgG3, or IgG4 (or hybrids thereof) is considered a non-naturally occurring modification.

[00163] By "amino acid" and "amino acid identity" as used herein is meant one of the 20 naturally occurring amino acids that are coded for by DNA and RNA.

[00164] By "effector function" as used herein is meant a biochemical event that results from the interaction of an antibody Fc region with an Fc receptor or ligand. Effector functions include but are not limited to ADCC, ADCP, and CDC.

[00165] By "IgG Fc ligand" as used herein is meant a molecule, preferably a polypeptide, from any organism that binds to the Fc region of an IgG antibody to form an Fc/Fc ligand complex. Fc ligands include but are not limited to FcγRIs, FcγRIIs, FcγRIIIIs, FcRn, C1q, C3, mannan binding lectin, mannose receptor, staphylococcal protein A, streptococcal protein G, and viral FcγR. Fc ligands also include Fc receptor homologs (FcRH), which are a family of Fc receptors that are homologous to the FcγRs (Davis et al., 2002, Immunological Reviews 190:123-136, entirely incorporated by reference). Fc ligands may include undiscovered molecules that bind Fc. Particular IgG Fc ligands are FcRn and Fc gamma receptors. By "Fc ligand" as used herein is meant a molecule, preferably a

polypeptide, from any organism that binds to the Fc region of an antibody to form an Fc/Fc ligand complex.

[00166] By "Fc gamma receptor", "FcγR" or "FcγammaR" as used herein is meant any member of the family of proteins that bind the IgG antibody Fc region and is encoded by an FcγR gene. In humans this family includes but is not limited to FcγRI (CD64), including isoforms FcγRIa, FcγRIb, and FcγRIc; FcγRII (CD32), including isoforms FcγRIIa (including allotypes H131 and R131), FcγRIIb (including FcγRIIb-1 and FcγRIIb-2), and FcγRIIc; and FcγRIII (CD16), including isoforms FcγRIIIa (including allotypes V158 and F158) and FcγRIIIb (including allotypes FcγRIIb-NA1 and FcγRIIb-NA2) (Jefferis et al., 2002, *Immunol Lett* 82:57-65, entirely incorporated by reference), as well as any undiscovered human FcγRs or FcγR isoforms or allotypes. An FcγR may be from any organism, including but not limited to humans, mice, rats, rabbits, and monkeys. Mouse FcγRs include but are not limited to FcγRI (CD64), FcγRII (CD32), FcγRIII (CD16), and FcγRIII-2 (CD16-2), as well as any undiscovered mouse FcγRs or FcγR isoforms or allotypes.

[00167] By "FcRn" or "neonatal Fc Receptor" as used herein is meant a protein that binds the IgG antibody Fc region and is encoded at least in part by an FcRn gene. The FcRn may be from any organism, including but not limited to humans, mice, rats, rabbits, and monkeys. As is known in the art, the functional FcRn protein comprises two polypeptides, often referred to as the heavy chain and light chain. The light chain is beta-2-microglobulin and the heavy chain is encoded by the FcRn gene. Unless otherwise noted herein, FcRn or an FcRn protein refers to the complex of FcRn heavy chain with beta-2-microglobulin. A variety of FcRn variants used to increase binding to the FcRn receptor, and in some cases, to increase serum half-life, are shown in the Figure Legend of Figure 83.

[00168] By "parent polypeptide" as used herein is meant a starting polypeptide that is subsequently modified to generate a variant. The parent polypeptide may be a naturally occurring polypeptide, or a variant or engineered version of a naturally occurring polypeptide. Parent polypeptide may refer to the polypeptide itself, compositions that comprise the parent polypeptide, or the amino acid sequence that encodes it. Accordingly, by "parent immunoglobulin" as used herein is meant an unmodified immunoglobulin polypeptide that is modified to generate a variant, and by "parent antibody" as used herein

is meant an unmodified antibody that is modified to generate a variant antibody. It should be noted that "parent antibody" includes known commercial, recombinantly produced antibodies as outlined below.

[00169] By "Fc" or "Fc region" or "Fc domain" as used herein is meant the polypeptide comprising the constant region of an antibody excluding the first constant region immunoglobulin domain and in some cases, part of the hinge. Thus Fc refers to the last two constant region immunoglobulin domains of IgA, IgD, and IgG, the last three constant region immunoglobulin domains of IgE and IgM, and the flexible hinge N-terminal to these domains. For IgA and IgM, Fc may include the J chain. For IgG, the Fc domain comprises immunoglobulin domains C γ 2 and C γ 3 (C γ 2 and C γ 3) and the lower hinge region between C γ 1 (C γ 1) and C γ 2 (C γ 2). Although the boundaries of the Fc region may vary, the human IgG heavy chain Fc region is usually defined to include residues C226 or P230 to its carboxyl-terminus, wherein the numbering is according to the EU index as in Kabat. In some embodiments, as is more fully described below, amino acid modifications are made to the Fc region, for example to alter binding to one or more Fc γ R receptors or to the FcRn receptor.

[00170] By "heavy constant region" herein is meant the CH1-hinge-CH2-CH3 portion of an antibody.

[00171] By "Fc fusion protein" or "immunoadhesin" herein is meant a protein comprising an Fc region, generally linked (optionally through a linker moiety, as described herein) to a different protein, such as a binding moiety to a target protein, as described herein. In some cases, one monomer of the heterodimeric antibody comprises an antibody heavy chain (either including an scFv or further including a light chain) and the other monomer is a Fc fusion, comprising a variant Fc domain and a ligand. In some embodiments, these "half antibody-half fusion proteins" are referred to as "Fusionbodies".

[00172] By "position" as used herein is meant a location in the sequence of a protein. Positions may be numbered sequentially, or according to an established format, for example the EU index for antibody numbering.

[00173] By "target antigen" as used herein is meant the molecule that is bound specifically by the variable region of a given antibody. A target antigen may be a protein, carbohydrate, lipid, or other chemical compound. A wide number of suitable target antigens are described below.

[00174] By "strandedness" in the context of the monomers of the heterodimeric antibodies of the invention herein is meant that, similar to the two strands of DNA that "match", heterodimerization variants are incorporated into each monomer so as to preserve the ability to "match" to form heterodimers. For example, if some pI variants are engineered into monomer A (e.g. making the pI higher) then steric variants that are "charge pairs" that can be utilized as well do not interfere with the pI variants, e.g. the charge variants that make a pI higher are put on the same "strand" or "monomer" to preserve both functionalities. Similarly, for "skew" variants that come in pairs of a set as more fully outlined below, the skilled artisan will consider pI in deciding into which strand or monomer that incorporates one set of the pair will go, such that pI separation is maximized using the pI of the skews as well.

[00175] By "target cell" as used herein is meant a cell that expresses a target antigen.

[00176] By "variable region" as used herein is meant the region of an immunoglobulin that comprises one or more Ig domains substantially encoded by any of the V.kappa., V.lamda., and/or VH genes that make up the kappa, lambda, and heavy chain immunoglobulin genetic loci respectively.

[00177] By "wild type or WT" herein is meant an amino acid sequence or a nucleotide sequence that is found in nature, including allelic variations. A WT protein has an amino acid sequence or a nucleotide sequence that has not been intentionally modified.

[00178] The antibodies of the present invention are generally isolated or recombinant. "Isolated," when used to describe the various polypeptides disclosed herein, means a polypeptide that has been identified and separated and/or recovered from a cell or cell culture from which it was expressed. Ordinarily, an isolated polypeptide will be prepared by at least one purification step. An "isolated antibody," refers to an antibody which is substantially free of other antibodies having different antigenic specificities. "Recombinant"

means the antibodies are generated using recombinant nucleic acid techniques in exogenous host cells.

[00179] “Specific binding” or “specifically binds to” or is “specific for” a particular antigen or an epitope means binding that is measurably different from a non-specific interaction. Specific binding can be measured, for example, by determining binding of a molecule compared to binding of a control molecule, which generally is a molecule of similar structure that does not have binding activity. For example, specific binding can be determined by competition with a control molecule that is similar to the target.

[00180] Specific binding for a particular antigen or an epitope can be exhibited, for example, by an antibody having a KD for an antigen or epitope of at least about 10^{-4} M, at least about 10^{-5} M, at least about 10^{-6} M, at least about 10^{-7} M, at least about 10^{-8} M, at least about 10^{-9} M, alternatively at least about 10^{-10} M, at least about 10^{-11} M, at least about 10^{-12} M, or greater, where KD refers to a dissociation rate of a particular antibody-antigen interaction. Typically, an antibody that specifically binds an antigen will have a KD that is 20-, 50-, 100-, 500-, 1000-, 5,000-, 10,000- or more times greater for a control molecule relative to the antigen or epitope.

[00181] Also, specific binding for a particular antigen or an epitope can be exhibited, for example, by an antibody having a KA or Ka for an antigen or epitope of at least 20-, 50-, 100-, 500-, 1000-, 5,000-, 10,000- or more times greater for the epitope relative to a control, where KA or Ka refers to an association rate of a particular antibody-antigen interaction.

II. Overview

[00182] Bispecific antibodies that co-engage CD3 and a tumor antigen target have been designed and used to redirect T cells to attack and lyse targeted tumor cells. Examples include the BiTE and DART formats, which monovalently engage CD3 and a tumor antigen. While the CD3-targeting approach has shown considerable promise, a common side effect of such therapies is the associated production of cytokines, often leading to toxic cytokine release syndrome. Because the anti-CD3 binding domain of the bispecific antibody engages all T cells, the high cytokine-producing CD4 T cell subset is recruited. Moreover, the CD4 T cell subset includes regulatory T cells, whose recruitment and expansion can potentially lead

to immune suppression and have a negative impact on long-term tumor suppression. In addition, these formats do not contain Fc domains and show very short serum half-lives in patients.

[00183] While the CD3-targeting approach has shown considerable promise, a common side effect of such therapies is the associated production of cytokines, often leading to toxic cytokine release syndrome. Because the anti-CD3 binding domain of the bispecific antibody engages all T cells, the high cytokine-producing CD4 T cell subset is recruited. Moreover, the CD4 T cell subset includes regulatory T cells, whose recruitment and expansion can potentially lead to immune suppression and have a negative impact on long-term tumor suppression. One such possible way to reduce cytokine production and possibly reduce the activation of CD4 T cells is by reducing the affinity of the anti-CD3 domain for CD3.

[00184] Accordingly, in some embodiments the present invention provides antibody constructs comprising anti-CD3 antigen binding domains that are “strong” or “high affinity” binders to CD3 (e.g. one example are heavy and light variable domains depicted as H1.30_L1.47 (optionally including a charged linker as appropriate)) and also bind to CD38. In other embodiments, the present invention provides antibody constructs comprising anti-CD3 antigen binding domains that are “lite” or “lower affinity” binders to CD3. Additional embodiments provides antibody constructs comprising anti-CD3 antigen binding domains that have intermediate or “medium” affinity to CD3 that also bind to CD38.

[00185] It should be appreciated that the “high, medium, low” anti-CD3 sequences of the present invention can be used in a variety of heterodimerization formats. While the majority of the disclosure herein uses the “bottle opener” format of heterodimers, these variable heavy and light sequences, as well as the scFv sequences (and Fab sequences comprising these variable heavy and light sequences) can be used in other formats, such as those depicted in Figure 2 of WO Publication No. 2014/145806, the Figures, formats and legend of which is expressly incorporated herein by reference.

[00186] Accordingly, the present invention provides heterodimeric antibodies that bind to two different antigens, e.g the antibodies are “bispecific”, in that they bind two

different target antigens, generally target tumor antigens (TTAs) as described below. These heterodimeric antibodies can bind these target antigens either monovalently (e.g. there is a single antigen binding domain such as a variable heavy and variable light domain pair) or bivalently (there are two antigen binding domains that each independently bind the antigen). The heterodimeric antibodies of the invention are based on the use different monomers which contain amino acid substitutions that “skew” formation of heterodimers over homodimers, as is more fully outlined below, coupled with “pI variants” that allow simple purification of the heterodimers away from the homodimers, as is similarly outlined below. For the heterodimeric bispecific antibodies of the invention, the present invention generally relies on the use of engineered or variant Fc domains that can self-assemble in production cells to produce heterodimeric proteins, and methods to generate and purify such heterodimeric proteins.

III. Antibodies

[00187] The present invention relates to the generation of bispecific antibodies that bind two different antigens, e.g. CD3 and a target tumor antigen such as CD20, CD38 and CD123, and are generally therapeutic antibodies. As is discussed below, the term “antibody” is used generally. Antibodies that find use in the present invention can take on a number of formats as described herein, including traditional antibodies as well as antibody derivatives, fragments and mimetics, described herein.

[00188] Traditional antibody structural units typically comprise a tetramer. Each tetramer is typically composed of two identical pairs of polypeptide chains, each pair having one “light” (typically having a molecular weight of about 25 kDa) and one “heavy” chain (typically having a molecular weight of about 50-70 kDa). Human light chains are classified as kappa and lambda light chains. The present invention is directed to the IgG class, which has several subclasses, including, but not limited to IgG1, IgG2, IgG3, and IgG4. Thus, “isotype” as used herein is meant any of the subclasses of immunoglobulins defined by the chemical and antigenic characteristics of their constant regions. It should be understood that therapeutic antibodies can also comprise hybrids of isotypes and/or subclasses. For example, as shown in US Publication 2009/0163699, incorporated by reference, the present invention covers pI engineering of IgG1/G2 hybrids.

[00189] The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition, generally referred to in the art and herein as the "Fv domain" or "Fv region". In the variable region, three loops are gathered for each of the V domains of the heavy chain and light chain to form an antigen-binding site. Each of the loops is referred to as a complementarity-determining region (hereinafter referred to as a "CDR"), in which the variation in the amino acid sequence is most significant. "Variable" refers to the fact that certain segments of the variable region differ extensively in sequence among antibodies. Variability within the variable region is not evenly distributed. Instead, the V regions consist of relatively invariant stretches called framework regions (FRs) of 15-30 amino acids separated by shorter regions of extreme variability called "hypervariable regions" that are each 9-15 amino acids long or longer.

[00190] Each VH and VL is composed of three hypervariable regions ("complementary determining regions," "CDRs") and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4.

[00191] The hypervariable region generally encompasses amino acid residues from about amino acid residues 24-34 (LCDR1; "L" denotes light chain), 50-56 (LCDR2) and 89-97 (LCDR3) in the light chain variable region and around about 31-35B (HCDR1; "H" denotes heavy chain), 50-65 (HCDR2), and 95-102 (HCDR3) in the heavy chain variable region; Kabat et al., SEQUENCES OF PROTEINS OF IMMUNOLOGICAL INTEREST, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991) and/or those residues forming a hypervariable loop (e.g. residues 26-32 (LCDR1), 50-52 (LCDR2) and 91-96 (LCDR3) in the light chain variable region and 26-32 (HCDR1), 53-55 (HCDR2) and 96-101 (HCDR3) in the heavy chain variable region; Chothia and Lesk (1987) J. Mol. Biol. 196:901-917. Specific CDRs of the invention are described below.

[00192] Throughout the present specification, the Kabat numbering system is generally used when referring to a residue in the variable domain (approximately, residues 1-107 of the light chain variable region and residues 1-113 of the heavy chain variable region) and the EU numbering system for Fc regions (e.g, Kabat et al., supra (1991)).

[00193] The present invention provides a large number of different CDR sets. In this case, a "full CDR set" comprises the three variable light and three variable heavy CDRs, e.g. a vLCDR1, vLCDR2, vLCDR3, vhCDR1, vhCDR2 and vhCDR3. These can be part of a larger variable light or variable heavy domain, respectfully. In addition, as more fully outlined herein, the variable heavy and variable light domains can be on separate polypeptide chains, when a heavy and light chain is used (for example when Fabs are used), or on a single polypeptide chain in the case of scFv sequences.

[00194] The CDRs contribute to the formation of the antigen-binding, or more specifically, epitope binding site of antibodies. "Epitope" refers to a determinant that interacts with a specific antigen binding site in the variable region of an antibody molecule known as a paratope. Epitopes are groupings of molecules such as amino acids or sugar side chains and usually have specific structural characteristics, as well as specific charge characteristics. A single antigen may have more than one epitope.

[00195] The epitope may comprise amino acid residues directly involved in the binding (also called immunodominant component of the epitope) and other amino acid residues, which are not directly involved in the binding, such as amino acid residues which are effectively blocked by the specifically antigen binding peptide; in other words, the amino acid residue is within the footprint of the specifically antigen binding peptide.

[00196] Epitopes may be either conformational or linear. A conformational epitope is produced by spatially juxtaposed amino acids from different segments of the linear polypeptide chain. A linear epitope is one produced by adjacent amino acid residues in a polypeptide chain. Conformational and nonconformational epitopes may be distinguished in that the binding to the former but not the latter is lost in the presence of denaturing solvents.

[00197] An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation. Antibodies that recognize the same epitope can be verified in a simple immunoassay showing the ability of one antibody to block the binding of another antibody to a target antigen, for example "binning."

[00198] The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function. Kabat et al. collected numerous primary sequences of the variable regions of heavy chains and light chains. Based on the degree of conservation of the sequences, they classified individual primary sequences into the CDR and the framework and made a list thereof (see SEQUENCES OF IMMUNOLOGICAL INTEREST, 5th edition, NIH publication, No. 91-3242, E.A. Kabat et al., entirely incorporated by reference).

[00199] In the IgG subclass of immunoglobulins, there are several immunoglobulin domains in the heavy chain. By "immunoglobulin (Ig) domain" herein is meant a region of an immunoglobulin having a distinct tertiary structure. Of interest in the present invention are the heavy chain domains, including, the constant heavy (CH) domains and the hinge domains. In the context of IgG antibodies, the IgG isotypes each have three CH regions. Accordingly, "CH" domains in the context of IgG are as follows: "CH1" refers to positions 118-220 according to the EU index as in Kabat. "CH2" refers to positions 237-340 according to the EU index as in Kabat, and "CH3" refers to positions 341-447 according to the EU index as in Kabat. As shown herein and described below, the pI variants can be in one or more of the CH regions, as well as the hinge region, discussed below.

[00200] It should be noted that the sequences depicted herein start at the CH1 region, position 118; the variable regions are not included except as noted. For example, the first amino acid of SEQ ID NO: 2, while designated as position "1" in the sequence listing, corresponds to position 118 of the CH1 region, according to EU numbering.

[00201] Another type of Ig domain of the heavy chain is the hinge region. By "hinge" or "hinge region" or "antibody hinge region" or "immunoglobulin hinge region" herein is meant the flexible polypeptide comprising the amino acids between the first and second constant domains of an antibody. Structurally, the IgG CH1 domain ends at EU position 220, and the IgG CH2 domain begins at residue EU position 237. Thus for IgG the antibody hinge is herein defined to include positions 221 (D221 in IgG1) to 236 (G236 in IgG1), wherein the numbering is according to the EU index as in Kabat. In some embodiments, for example in the context of an Fc region, the lower hinge is included, with the "lower hinge" generally

referring to positions 226 or 230. As noted herein, pI variants can be made in the hinge region as well.

[00202] The light chain generally comprises two domains, the variable light domain (containing the light chain CDRs and together with the variable heavy domains forming the Fv region), and a constant light chain region (often referred to as CL or C_κ).

[00203] Another region of interest for additional substitutions, outlined below, is the Fc region.

[00204] Thus, the present invention provides different antibody domains. As described herein and known in the art, the heterodimeric antibodies of the invention comprise different domains within the heavy and light chains, which can be overlapping as well. These domains include, but are not limited to, the Fc domain, the CH1 domain, the CH2 domain, the CH3 domain, the hinge domain, the heavy constant domain (CH1-hinge-Fc domain or CH1-hinge-CH2-CH3), the variable heavy domain, the variable light domain, the light constant domain, FAb domains and scFv domains.

[00205] Thus, the "Fc domain" includes the -CH2-CH3 domain, and optionally a hinge domain. The heavy chain comprises a variable heavy domain and a constant domain, which includes a CH1-optional hinge-Fc domain comprising a CH2-CH3. The light chain comprises a variable light chain and the light constant domain.

[00206] Some embodiments of the invention comprise at least one scFv domain, which, while not naturally occurring, generally includes a variable heavy domain and a variable light domain, linked together by a scFv linker. As shown herein, there are a number of suitable scFv linkers that can be used, including traditional peptide bonds, generated by recombinant techniques.

[00207] The linker peptide may predominantly include the following amino acid residues: Gly, Ser, Ala, or Thr. The linker peptide should have a length that is adequate to link two molecules in such a way that they assume the correct conformation relative to one another so that they retain the desired activity. In one embodiment, the linker is from about 1 to 50 amino acids in length, preferably about 1 to 30 amino acids in length. In one embodiment, linkers of 1 to 20 amino acids in length may be used, with from about 5 to

about 10 amino acids finding use in some embodiments. Useful linkers include glycine-serine polymers, including for example (GS)_n, (GSGGS)_n, (GGGGS)_n, and (GGGS)_n, where *n* is an integer of at least one (and generally from 3 to 4), glycine-alanine polymers, alanine-serine polymers, and other flexible linkers. Alternatively, a variety of nonproteinaceous polymers, including but not limited to polyethylene glycol (PEG), polypropylene glycol, polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol, may find use as linkers, that is may find use as linkers.

[00208] Other linker sequences may include any sequence of any length of CL/CH1 domain but not all residues of CL/CH1 domain; for example the first 5-12 amino acid residues of the CL/CH1 domains. Linkers can be derived from immunoglobulin light chain, for example C κ or C λ . Linkers can be derived from immunoglobulin heavy chains of any isotype, including for example C γ 1, C γ 2, C γ 3, C γ 4, C α 1, C α 2, C δ , C ϵ , and C μ . Linker sequences may also be derived from other proteins such as Ig-like proteins (e.g. TCR, FcR, KIR), hinge region-derived sequences, and other natural sequences from other proteins.

[00209] In some embodiments, the linker is a “domain linker”, used to link any two domains as outlined herein together. While any suitable linker can be used, many embodiments utilize a glycine-serine polymer, including for example (GS)_n, (GSGGS)_n, (GGGGS)_n, and (GGGS)_n, where *n* is an integer of at least one (and generally from 3 to 4 to 5) as well as any peptide sequence that allows for recombinant attachment of the two domains with sufficient length and flexibility to allow each domain to retain its biological function. . In some cases, and with attention being paid to “strandedness”, as outlined below, charged domain linkers, as used in some embodiments of scFv linkers can be used.

[00210] In some embodiments, the scFv linker is a charged scFv linker, a number of which are shown in Figure 33. Accordingly, the present invention further provides charged scFv linkers, to facilitate the separation in pI between a first and a second monomer. That is, by incorporating a charged scFv linker, either positive or negative (or both, in the case of scaffolds that use scFvs on different monomers), this allows the monomer comprising the charged linker to alter the pI without making further changes in the Fc domains. These charged linkers can be substituted into any scFv containing standard linkers. Again, as will be appreciated by those in the art, charged scFv linkers are used on the correct “strand” or

monomer, according to the desired changes in pI. For example, as discussed herein, to make triple F format heterodimeric antibody, the original pI of the Fv region for each of the desired antigen binding domains are calculated, and one is chosen to make an scFv, and depending on the pI, either positive or negative linkers are chosen.

[00211] Charged domain linkers can also be used to increase the pI separation of the monomers of the invention as well, and thus those included in Figure 33 can be used in any embodiment herein where a linker is utilized.

[00212] In some embodiments, the antibodies are full length. By “full length antibody” herein is meant the structure that constitutes the natural biological form of an antibody, including variable and constant regions, including one or more modifications as outlined herein, particularly in the Fc domains to allow either heterodimerization formation or the purification of heterodimers away from homodimers. Full length antibodies generally include Fab and Fc domains, and can additionally contain extra antigen binding domains such as scFvs, as is generally depicted in the Figures.

[00213] In one embodiment, the antibody is an antibody fragment, as long as it contains at least one constant domain which can be engineered to produce heterodimers, such as pI engineering. Other antibody fragments that can be used include fragments that contain one or more of the CH1, CH2, CH3, hinge and CL domains of the invention that have been pI engineered. For example, Fc fusions are fusions of the Fc region (CH2 and CH3, optionally with the hinge region) fused to another protein. A number of Fc fusions are known the art and can be improved by the addition of the heterodimerization variants of the invention. In the present case, antibody fusions can be made comprising CH1; CH1, CH2 and CH3; CH2; CH3; CH2 and CH3; CH1 and CH3, any or all of which can be made optionally with the hinge region, utilizing any combination of heterodimerization variants described herein.

[00214] In particular, the formats depicted in Figure 1 are antibodies, usually referred to as “heterodimeric antibodies”, meaning that the protein has at least two associated Fc sequences self-assembled into a heterodimeric Fc domain.

Chimeric and Humanized Antibodies

[00215] In some embodiments, the antibody can be a mixture from different species, e.g. a chimeric antibody and/or a humanized antibody. In general, both “chimeric antibodies” and “humanized antibodies” refer to antibodies that combine regions from more than one species. For example, “chimeric antibodies” traditionally comprise variable region(s) from a mouse (or rat, in some cases) and the constant region(s) from a human. “Humanized antibodies” generally refer to non-human antibodies that have had the variable-domain framework regions swapped for sequences found in human antibodies. Generally, in a humanized antibody, the entire antibody, except the CDRs, is encoded by a polynucleotide of human origin or is identical to such an antibody except within its CDRs. The CDRs, some or all of which are encoded by nucleic acids originating in a non-human organism, are grafted into the beta-sheet framework of a human antibody variable region to create an antibody, the specificity of which is determined by the engrafted CDRs. The creation of such antibodies is described in, e.g., WO 92/11018, Jones, 1986, Nature 321:522-525, Verhoeyen et al., 1988, Science 239:1534-1536, all entirely incorporated by reference. “Backmutation” of selected acceptor framework residues to the corresponding donor residues is often required to regain affinity that is lost in the initial grafted construct (US 5530101; US 5585089; US 5693761; US 5693762; US 6180370; US 5859205; US 5821337; US 6054297; US 6407213, all entirely incorporated by reference). The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region, typically that of a human immunoglobulin, and thus will typically comprise a human Fc region. Humanized antibodies can also be generated using mice with a genetically engineered immune system. Roque et al., 2004, Biotechnol. Prog. 20:639-654, entirely incorporated by reference. A variety of techniques and methods for humanizing and reshaping non-human antibodies are well known in the art (See Tsurushita & Vasquez, 2004, Humanization of Monoclonal Antibodies, Molecular Biology of B Cells, 533-545, Elsevier Science (USA), and references cited therein, all entirely incorporated by reference). Humanization methods include but are not limited to methods described in Jones et al., 1986, Nature 321:522-525; Riechmann et al., 1988; Nature 332:323-329; Verhoeyen et al., 1988, Science, 239:1534-1536; Queen et al., 1989, Proc Natl Acad Sci, USA 86:10029-33; He et al., 1998, J. Immunol. 160: 1029-1035; Carter et al., 1992, Proc Natl Acad Sci USA 89:4285-9,

Presta et al., 1997, *Cancer Res.* 57(20):4593-9; Gorman et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:4181-4185; O'Connor et al., 1998, *Protein Eng* 11:321-8, all entirely incorporated by reference. Humanization or other methods of reducing the immunogenicity of nonhuman antibody variable regions may include resurfacing methods, as described for example in Roguska et al., 1994, *Proc. Natl. Acad. Sci. USA* 91:969-973, entirely incorporated by reference. In one embodiment, the parent antibody has been affinity matured, as is known in the art. Structure-based methods may be employed for humanization and affinity maturation, for example as described in USSN 11/004,590. Selection based methods may be employed to humanize and/or affinity mature antibody variable regions, including but not limited to methods described in Wu et al., 1999, *J. Mol. Biol.* 294:151-162; Baca et al., 1997, *J. Biol. Chem.* 272(16):10678-10684; Rosok et al., 1996, *J. Biol. Chem.* 271(37): 22611-22618; Rader et al., 1998, *Proc. Natl. Acad. Sci. USA* 95: 8910-8915; Krauss et al., 2003, *Protein Engineering* 16(10):753-759, all entirely incorporated by reference. Other humanization methods may involve the grafting of only parts of the CDRs, including but not limited to methods described in USSN 09/810,510; Tan et al., 2002, *J. Immunol.* 169:1119-1125; De Pascalis et al., 2002, *J. Immunol.* 169:3076-3084, all entirely incorporated by reference.

IV. Heterodimeric Antibodies

[00216] Accordingly, in some embodiments the present invention provides heterodimeric antibodies that rely on the use of two different heavy chain variant Fc sequences, that will self-assemble to form heterodimeric Fc domains and heterodimeric antibodies.

[00217] The present invention is directed to novel constructs to provide heterodimeric antibodies that allow binding to more than one antigen or ligand, e.g. to allow for bispecific binding. The heterodimeric antibody constructs are based on the self-assembling nature of the two Fc domains of the heavy chains of antibodies, e.g. two "monomers" that assemble into a "dimer". Heterodimeric antibodies are made by altering the amino acid sequence of each monomer as more fully discussed below. Thus, the present invention is generally directed to the creation of heterodimeric antibodies which can co-engage antigens in several ways, relying on amino acid variants in the constant regions that are different on each chain

to promote heterodimeric formation and/or allow for ease of purification of heterodimers over the homodimers.

[00218] Thus, the present invention provides bispecific antibodies. An ongoing problem in antibody technologies is the desire for “bispecific” antibodies that bind to two different antigens simultaneously, in general thus allowing the different antigens to be brought into proximity and resulting in new functionalities and new therapies. In general, these antibodies are made by including genes for each heavy and light chain into the host cells. This generally results in the formation of the desired heterodimer (A-B), as well as the two homodimers (A-A and B-B (not including the light chain heterodimeric issues)). However, a major obstacle in the formation of bispecific antibodies is the difficulty in purifying the heterodimeric antibodies away from the homodimeric antibodies and/or biasing the formation of the heterodimer over the formation of the homodimers.

[00219] There are a number of mechanisms that can be used to generate the heterodimers of the present invention. In addition, as will be appreciated by those in the art, these mechanisms can be combined to ensure high heterodimerization. Thus, amino acid variants that lead to the production of heterodimers are referred to as “heterodimerization variants”. As discussed below, heterodimerization variants can include steric variants (e.g. the “knobs and holes” or “skew” variants described below and the “charge pairs” variants described below) as well as “pI variants”, which allows purification of homodimers away from heterodimers. As is generally described in WO2014/145806, hereby incorporated by reference in its entirety and specifically as below for the discussion of “heterodimerization variants”, useful mechanisms for heterodimerization include “knobs and holes” (“KIH”; sometimes herein as “skew” variants (see discussion in WO2014/145806), “electrostatic steering” or “charge pairs” as described in WO2014/145806, pI variants as described in WO2014/145806, and general additional Fc variants as outlined in WO2014/145806 and below.

[00220] In the present invention, there are several basic mechanisms that can lead to ease of purifying heterodimeric antibodies; one relies on the use of pI variants, such that each monomer has a different pI, thus allowing the isoelectric purification of A-A, A-B and B-B dimeric proteins. Alternatively, some scaffold formats, such as the “triple F” format,

also allows separation on the basis of size. As is further outlined below, it is also possible to “skew” the formation of heterodimers over homodimers. Thus, a combination of steric heterodimerization variants and pI or charge pair variants find particular use in the invention.

[00221] In general, embodiments of particular use in the present invention rely on sets of variants that include skew variants, that encourage heterodimerization formation over homodimerization formation, coupled with pI variants, which increase the pI difference between the two monomers.

[00222] Additionally, as more fully outlined below, depending on the format of the heterodimer antibody, pI variants can be either contained within the constant and/or Fc domains of a monomer, or charged linkers, either domain linkers or scFv linkers, can be used. That is, scaffolds that utilize scFv(s) such as the Triple F format can include charged scFv linkers (either positive or negative), that give a further pI boost for purification purposes. As will be appreciated by those in the art, some Triple F formats are useful with just charged scFv linkers and no additional pI adjustments, although the invention does provide pI variants that are on one or both of the monomers, and/or charged domain linkers as well. In addition, additional amino acid engineering for alternative functionalities may also confer pI changes, such as Fc, FcRn and KO variants.

[00223] In the present invention that utilizes pI as a separation mechanism to allow the purification of heterodimeric proteins, amino acid variants can be introduced into one or both of the monomer polypeptides; that is, the pI of one of the monomers (referred to herein for simplicity as “monomer A”) can be engineered away from monomer B, or both monomer A and B change be changed, with the pI of monomer A increasing and the pI of monomer B decreasing. As is outlined more fully below, the pI changes of either or both monomers can be done by removing or adding a charged residue (e.g. a neutral amino acid is replaced by a positively or negatively charged amino acid residue, e.g. glycine to glutamic acid), changing a charged residue from positive or negative to the opposite charge (aspartic acid to lysine) or changing a charged residue to a neutral residue (e.g. loss of a charge; lysine to serine.). A number of these variants are shown in the Figures.

[00224] Accordingly, this embodiment of the present invention provides for creating a sufficient change in pI in at least one of the monomers such that heterodimers can be separated from homodimers. As will be appreciated by those in the art, and as discussed further below, this can be done by using a “wild type” heavy chain constant region and a variant region that has been engineered to either increase or decrease its pI (wt A+B or wt A - B), or by increasing one region and decreasing the other region (A+ -B- or A- B+).

[00225] Thus, in general, a component of some embodiments of the present invention are amino acid variants in the constant regions of antibodies that are directed to altering the isoelectric point (pI) of at least one, if not both, of the monomers of a dimeric protein to form “pI antibodies”) by incorporating amino acid substitutions (“pI variants” or “pI substitutions”) into one or both of the monomers. As shown herein, the separation of the heterodimers from the two homodimers can be accomplished if the pIs of the two monomers differ by as little as 0.1 pH unit, with 0.2, 0.3, 0.4 and 0.5 or greater all finding use in the present invention.

[00226] As will be appreciated by those in the art, the number of pI variants to be included on each or both monomer(s) to get good separation will depend in part on the starting pI of the components, for example in the triple F format, the starting pI of the scFv and Fab of interest. That is, to determine which monomer to engineer or in which “direction” (e.g. more positive or more negative), the Fv sequences of the two target antigens are calculated and a decision is made from there. As is known in the art, different Fvs will have different starting pIs which are exploited in the present invention. In general, as outlined herein, the pIs are engineered to result in a total pI difference of each monomer of at least about 0.1 logs, with 0.2 to 0.5 being preferred as outlined herein.

[00227] Furthermore, as will be appreciated by those in the art and outlined herein, in some embodiments, heterodimers can be separated from homodimers on the basis of size. As shown in Figures 1 for example, several of the formats allow separation of heterodimers and homodimers on the basis of size.

[00228] In the case where pI variants are used to achieve heterodimerization, by using the constant region(s) of the heavy chain(s), a more modular approach to designing and

purifying bispecific proteins, including antibodies, is provided. Thus, in some embodiments, heterodimerization variants (including skew and purification heterodimerization variants) are not included in the variable regions, such that each individual antibody must be engineered. In addition, in some embodiments, the possibility of immunogenicity resulting from the pI variants is significantly reduced by importing pI variants from different IgG isotypes such that pI is changed without introducing significant immunogenicity. Thus, an additional problem to be solved is the elucidation of low pI constant domains with high human sequence content, e.g. the minimization or avoidance of non-human residues at any particular position.

[00229] A side benefit that can occur with this pI engineering is also the extension of serum half-life and increased FcRn binding. That is, as described in USSN 13/194,904 (incorporated by reference in its entirety), lowering the pI of antibody constant domains (including those found in antibodies and Fc fusions) can lead to longer serum retention in vivo. These pI variants for increased serum half life also facilitate pI changes for purification.

[00230] In addition, it should be noted that the pI variants of the heterodimerization variants give an additional benefit for the analytics and quality control process of bispecific antibodies, as the ability to either eliminate, minimize and distinguish when homodimers are present is significant. Similarly, the ability to reliably test the reproducibility of the heterodimeric antibody production is important.

Heterodimerization Variants

[00231] The present invention provides heterodimeric proteins, including heterodimeric antibodies in a variety of formats, which utilize heterodimeric variants to allow for heterodimeric formation and/or purification away from homodimers.

[00232] There are a number of suitable pairs of sets of heterodimerization skew variants. These variants come in “pairs” of “sets”. That is, one set of the pair is incorporated into the first monomer and the other set of the pair is incorporated into the second monomer. It should be noted that these sets do not necessarily behave as “knobs in holes” variants, with a one-to-one correspondence between a residue on one monomer and a

residue on the other; that is, these pairs of sets form an interface between the two monomers that encourages heterodimer formation and discourages homodimer formation, allowing the percentage of heterodimers that spontaneously form under biological conditions to be over 90%, rather than the expected 50% (25 % homodimer A/A:50% heterodimer A/B:25% homodimer B/B).

Steric Variants

[00233] In some embodiments, the formation of heterodimers can be facilitated by the addition of steric variants. That is, by changing amino acids in each heavy chain, different heavy chains are more likely to associate to form the heterodimeric structure than to form homodimers with the same Fc amino acid sequences. Suitable steric variants are included in Figure 29.

[00234] One mechanism is generally referred to in the art as “knobs and holes”, referring to amino acid engineering that creates steric influences to favor heterodimeric formation and disfavor homodimeric formation can also optionally be used; this is sometimes referred to as “knobs and holes”, as described in USSN 61/596,846, Ridgway et al., Protein Engineering 9(7):617 (1996); Atwell et al., J. Mol. Biol. 1997 270:26; US Patent No. 8,216,805, all of which are hereby incorporated by reference in their entirety. The Figures identify a number of “monomer A – monomer B” pairs that rely on “knobs and holes”. In addition, as described in Merchant et al., Nature Biotech. 16:677 (1998), these “knobs and hole” mutations can be combined with disulfide bonds to skew formation to heterodimerization.

[00235] An additional mechanism that finds use in the generation of heterodimers is sometimes referred to as “electrostatic steering” as described in Gunasekaran et al., J. Biol. Chem. 285(25):19637 (2010), hereby incorporated by reference in its entirety. This is sometimes referred to herein as “charge pairs”. In this embodiment, electrostatics are used to skew the formation towards heterodimerization. As those in the art will appreciate, these may also have an effect on pI, and thus on purification, and thus could in some cases also be considered pI variants. However, as these were generated to force heterodimerization and were not used as purification tools, they are classified as “steric

variants". These include, but are not limited to, D221E/P228E/L368E paired with D221R/P228R/K409R (e.g. these are "monomer corresponding sets) and C220E/P228E/368E paired with C220R/E224R/P228R/K409R.

[00236] Additional monomer A and monomer B variants that can be combined with other variants, optionally and independently in any amount, such as pI variants outlined herein or other steric variants that are shown in Figure 37 of US 2012/0149876, the figure and legend and SEQ ID NOs of which are incorporated expressly by reference herein.

[00237] In some embodiments, the steric variants outlined herein can be optionally and independently incorporated with any pI variant (or other variants such as Fc variants, FcRn variants, etc.) into one or both monomers, and can be independently and optionally included or excluded from the proteins of the invention.

[00238] A list of suitable skew variants is found in Figure 29, with Figure 34 showing some pairs of particular utility in many embodiments. Of particular use in many embodiments are the pairs of sets including, but not limited to, S364K/E357Q : L368D/K370S; L368D/K370S : S364K; L368E/K370S : S364K; T411T/E360E/Q362E : D401K; L368D/K370S : S364K/E357L and K370S : S364K/E357Q. In terms of nomenclature, the pair "S364K/E357Q : L368D/K370S" means that one of the monomers has the double variant set S364K/E357Q and the other has the double variant set L368D/K370S.

pI (Isoelectric point) Variants for Heterodimers

[00239] In general, as will be appreciated by those in the art, there are two general categories of pI variants: those that increase the pI of the protein (basic changes) and those that decrease the pI of the protein (acidic changes). As described herein, all combinations of these variants can be done: one monomer may be wild type, or a variant that does not display a significantly different pI from wild-type, and the other can be either more basic or more acidic. Alternatively, each monomer is changed, one to more basic and one to more acidic.

[00240] Preferred combinations of pI variants are shown in Figure 30. As outlined herein and shown in the figures, these changes are shown relative to IgG1, but all isotypes

can be altered this way, as well as isotype hybrids. In the case where the heavy chain constant domain is from IgG2-4, R133E and R133Q can also be used.

Antibody Heterodimers Light chain variants

[00241] In the case of antibody based heterodimers, e.g. where at least one of the monomers comprises a light chain in addition to the heavy chain domain, pI variants can also be made in the light chain. Amino acid substitutions for lowering the pI of the light chain include, but are not limited to, K126E, K126Q, K145E, K145Q, N152D, S156E, K169E, S202E, K207E and adding peptide DEDE at the c-terminus of the light chain. Changes in this category based on the constant lambda light chain include one or more substitutions at R108Q, Q124E, K126Q, N138D, K145T and Q199E. In addition, increasing the pI of the light chains can also be done.

Isotypic Variants

[00242] In addition, many embodiments of the invention rely on the “importation” of pI amino acids at particular positions from one IgG isotype into another, thus reducing or eliminating the possibility of unwanted immunogenicity being introduced into the variants. A number of these are shown in Figure 21 of US Publ. 2014/0370013, hereby incorporated by reference. That is, IgG1 is a common isotype for therapeutic antibodies for a variety of reasons, including high effector function. However, the heavy constant region of IgG1 has a higher pI than that of IgG2 (8.10 versus 7.31). By introducing IgG2 residues at particular positions into the IgG1 backbone, the pI of the resulting monomer is lowered (or increased) and additionally exhibits longer serum half-life. For example, IgG1 has a glycine (pI 5.97) at position 137, and IgG2 has a glutamic acid (pI 3.22); importing the glutamic acid will affect the pI of the resulting protein. As is described below, a number of amino acid substitutions are generally required to significantly affect the pI of the variant antibody. However, it should be noted as discussed below that even changes in IgG2 molecules allow for increased serum half-life.

[00243] In other embodiments, non-isotypic amino acid changes are made, either to reduce the overall charge state of the resulting protein (e.g. by changing a higher pI amino

acid to a lower pI amino acid), or to allow accommodations in structure for stability, etc. as is more further described below.

[00244] In addition, by pI engineering both the heavy and light constant domains, significant changes in each monomer of the heterodimer can be seen. As discussed herein, having the pIs of the two monomers differ by at least 0.5 can allow separation by ion exchange chromatography or isoelectric focusing, or other methods sensitive to isoelectric point.

Calculating pI

[00245] The pI of each monomer can depend on the pI of the variant heavy chain constant domain and the pI of the total monomer, including the variant heavy chain constant domain and the fusion partner. Thus, in some embodiments, the change in pI is calculated on the basis of the variant heavy chain constant domain, using the chart in the Figure 19 of US Pub. 2014/0370013. As discussed herein, which monomer to engineer is generally decided by the inherent pI of the Fv and scaffold regions. Alternatively, the pI of each monomer can be compared.

pI Variants that also confer better FcRn in vivo binding

[00246] In the case where the pI variant decreases the pI of the monomer, they can have the added benefit of improving serum retention in vivo.

[00247] Although still under examination, Fc regions are believed to have longer half-lives in vivo, because binding to FcRn at pH 6 in an endosome sequesters the Fc (Ghetie and Ward, 1997 Immunol Today. 18(12): 592-598, entirely incorporated by reference). The endosomal compartment then recycles the Fc to the cell surface. Once the compartment opens to the extracellular space, the higher pH, ~7.4, induces the release of Fc back into the blood. In mice, Dall'Acqua et al. showed that Fc mutants with increased FcRn binding at pH 6 and pH 7.4 actually had reduced serum concentrations and the same half life as wild-type Fc (Dall'Acqua et al. 2002, J. Immunol. 169:5171-5180, entirely incorporated by reference). The increased affinity of Fc for FcRn at pH 7.4 is thought to forbid the release of the Fc back into the blood. Therefore, the Fc mutations that will increase Fc's half-life in vivo will ideally increase FcRn binding at the lower pH while still allowing release of Fc at higher pH. The

amino acid histidine changes its charge state in the pH range of 6.0 to 7.4. Therefore, it is not surprising to find His residues at important positions in the Fc/FcRn complex.

[00248] Recently it has been suggested that antibodies with variable regions that have lower isoelectric points may also have longer serum half-lives (Igawa et al., 2010 PEDS. 23(5): 385-392, entirely incorporated by reference). However, the mechanism of this is still poorly understood. Moreover, variable regions differ from antibody to antibody. Constant region variants with reduced pI and extended half-life would provide a more modular approach to improving the pharmacokinetic properties of antibodies, as described herein.

Additional Fc Variants for Additional Functionality

[00249] In addition to pI amino acid variants, there are a number of useful Fc amino acid modification that can be made for a variety of reasons, including, but not limited to, altering binding to one or more FcγR receptors, altered binding to FcRn receptors, etc.

[00250] Accordingly, the proteins of the invention can include amino acid modifications, including the heterodimerization variants outlined herein, which includes the pI variants and steric variants. Each set of variants can be independently and optionally included or excluded from any particular heterodimeric protein.

FcγR Variants

[00251] Accordingly, there are a number of useful Fc substitutions that can be made to alter binding to one or more of the FcγR receptors. Substitutions that result in increased binding as well as decreased binding can be useful. For example, it is known that increased binding to Fc RIIIa generally results in increased ADCC (antibody dependent cell-mediated cytotoxicity; the cell-mediated reaction wherein nonspecific cytotoxic cells that express FcγRs recognize bound antibody on a target cell and subsequently cause lysis of the target cell). Similarly, decreased binding to FcγRIIb (an inhibitory receptor) can be beneficial as well in some circumstances. Amino acid substitutions that find use in the present invention include those listed in USSNs 11/124,620 (particularly Figure 41), 11/174,287, 11/396,495, 11/538,406, all of which are expressly incorporated herein by reference in their entirety and specifically for the variants disclosed therein. Particular variants that find use include, but

are not limited to, 236A, 239D, 239E, 332E, 332D, 239D/332E, 267D, 267E, 328F, 267E/328F, 236A/332E, 239D/332E/330Y, 239D, 332E/330L, 243A, 243L, 264A, 264V and 299T.

[00252] In addition, there are additional Fc substitutions that find use in increased binding to the FcRn receptor and increased serum half life, as specifically disclosed in USSN 12/341,769, hereby incorporated by reference in its entirety, including, but not limited to, 434S, 434A, 428L, 308F, 259I, 428L/434S, 259I/308F, 436I/428L, 436I or V/434S, 436V/428L and 259I/308F/428L.

Ablation Variants

[00253] Similarly, another category of functional variants are "FcγR ablation variants" or "Fc knock out (FcKO or KO)" variants. In these embodiments, for some therapeutic applications, it is desirable to reduce or remove the normal binding of the Fc domain to one or more or all of the Fcγ receptors (e.g. FcγR1, FcγRIIa, FcγRIIb, FcγRIIIa, etc.) to avoid additional mechanisms of action. That is, for example, in many embodiments, particularly in the use of bispecific antibodies that bind CD3 monovalently it is generally desirable to ablate FcγRIIIa binding to eliminate or significantly reduce ADCC activity. wherein one of the Fc domains comprises one or more Fcγ receptor ablation variants. These ablation variants are depicted in Figure 31, and each can be independently and optionally included or excluded, with preferred aspects utilizing ablation variants selected from the group consisting of G236R/L328R, E233P/L234V/L235A/G236del/S239K, E233P/L234V/L235A/G236del/S267K, E233P/L234V/L235A/G236del/S239K/A327G, E233P/L234V/L235A/G236del/S267K/A327G and E233P/L234V/L235A/G236del. It should be noted that the ablation variants referenced herein ablate FcγR binding but generally not FcRn binding.

Combination of Heterodimeric and Fc Variants

[00254] As will be appreciated by those in the art, all of the recited heterodimerization variants (including skew and/or pl variants) can be optionally and independently combined in any way, as long as they retain their "strandedness" or "monomer partition". In addition, all of these variants can be combined into any of the heterodimerization formats.

[00255] In the case of pI variants, while embodiments finding particular use are shown in the Figures, other combinations can be generated, following the basic rule of altering the pI difference between two monomers to facilitate purification.

[00256] In addition, any of the heterodimerization variants, skew and pI, are also independently and optionally combined with Fc ablation variants, Fc variants, FcRn variants, as generally outlined herein.

Useful Formats of the Invention

[00257] As will be appreciated by those in the art and discussed more fully below, the heterodimeric fusion proteins of the present invention can take on a wide variety of configurations, as are generally depicted in Figures 1. Some figures depict “single ended” configurations, where there is one type of specificity on one “arm” of the molecule and a different specificity on the other “arm”. Other figures depict “dual ended” configurations, where there is at least one type of specificity at the “top” of the molecule and one or more different specificities at the “bottom” of the molecule. Thus, the present invention is directed to novel immunoglobulin compositions that co-engage a different first and a second antigen.

[00258] As will be appreciated by those in the art, the heterodimeric formats of the invention can have different valencies as well as be bispecific. That is, heterodimeric antibodies of the invention can be bivalent and bispecific, wherein one target tumor antigen (e.g. CD3) is bound by one binding domain and the other target tumor antigen (e.g. CD20, CD19, CD38, CD123, etc.) is bound by a second binding domain. The heterodimeric antibodies can also be trivalent and bispecific, wherein the first antigen is bound by two binding domains and the second antigen by a second binding domain. As is outlined herein, when CD3 is one of the target antigens, it is preferable that the CD3 is bound only monovalently, to reduce potential side effects.

[00259] The present invention utilizes anti-CD3 antigen binding domains in combination with anti-target tumor antigen (TTA) antigen binding domains. As will be appreciated by those in the art, any collection of anti-CD3 CDRs, anti-CD3 variable light and variable heavy domains, Fabs and scFvs as depicted in any of the Figures (see particularly

Figures 2 through 7, and Figure 68) can be used. Similarly, any of the anti-TTA antigen binding domains can be used, e.g. anti-CD38, anti-CD20, anti-CD19 and anti-CD123 antigen binding domains, whether CDRs, variable light and variable heavy domains, Fabs and scFvs as depicted in any of the Figures can be used, optionally and independently combined in any combination.

Bottle opener format

[00260] One heterodimeric scaffold that finds particular use in the present invention is the “triple F” or “bottle opener” scaffold format as shown in Figure 1A, A and B. In this embodiment, one heavy chain of the antibody contains an single chain Fv (“scFv”, as defined below) and the other heavy chain is a “regular” FAb format, comprising a variable heavy chain and a light chain. This structure is sometimes referred to herein as “triple F” format (scFv-FAb-Fc) or the “bottle-opener” format, due to a rough visual similarity to a bottle-opener (see Figures 1). The two chains are brought together by the use of amino acid variants in the constant regions (e.g. the Fc domain, the CH1 domain and/or the hinge region) that promote the formation of heterodimeric antibodies as is described more fully below.

[00261] There are several distinct advantages to the present “triple F” format. As is known in the art, antibody analogs relying on two scFv constructs often have stability and aggregation problems, which can be alleviated in the present invention by the addition of a “regular” heavy and light chain pairing. In addition, as opposed to formats that rely on two heavy chains and two light chains, there is no issue with the incorrect pairing of heavy and light chains (e.g. heavy 1 pairing with light 2, etc.).

[00262] Many of the embodiments outlined herein rely in general on the bottle opener format that comprises a first monomer comprising an scFv, comprising a variable heavy and a variable light domain, covalently attached using an scFv linker (charged, in many but not all instances), where the scFv is covalently attached to the N-terminus of a first Fc domain usually through a domain linker (which, as outlined herein can either be un-charged or charged). The second monomer of the bottle opener format is a heavy chain, and the composition further comprises a light chain.

[00263] In general, in many preferred embodiments, the scFv is the domain that binds to the CD3, with the Fab of the heavy and light chains binding to the other TTA. In addition, the Fc domains of the invention generally comprise skew variants (e.g. a set of amino acid substitutions as shown in Figure 29 and Figure 34, with particularly useful skew variants being selected from the group consisting of S364K/E357Q : L368D/K370S; L368D/K370S : S364K; L368E/K370S : S364K; T411T/E360E/Q362E : D401K; L368D/K370S : S364K/E357L and K370S : S364K/E357Q), optionally ablation variants (including those shown in Figure 31), optionally charged scFv linkers (including those shown in Figure 33) and the heavy chain comprises pI variants (including those shown in Figure 30).

[00264] The present invention provides bottle opener formats where the anti-CD3 scFv sequences are as shown in Figure 2 to Figure 7 and Figure 68.

[00265] The present invention provides bottle opener formats with CD38 antigen binding domains wherein the anti-CD38 sequences are as shown in the Figures, including Figures 8 to 10.

[00266] The present invention provides bottle opener formats with CD20 antigen binding domains wherein the anti-CD20 sequences are as shown in the Figures.

[00267] The present invention provides bottle opener formats with CD19 antigen binding domains wherein the anti-CD19 sequences are as shown in the Figures.

[00268] The present invention provides bottle opener formats with CD123 antigen binding domains wherein the anti-CD123 sequences are as shown in the Figures.

mAb-Fv format

[00269] One heterodimeric scaffold that finds particular use in the present invention is the mAb-Fv format shown in Figure 1. In this embodiment, the format relies on the use of a C-terminal attachment of an "extra" variable heavy domain to one monomer and the C-terminal attachment of an "extra" variable light domain to the other monomer, thus forming a third antigen binding domain, wherein the Fab portions of the two monomers bind a TTA and the "extra" scFv domain binds CD3.

[00270] In this embodiment, the first monomer comprises a first heavy chain, comprising a first variable heavy domain and a first constant heavy domain comprising a first Fc domain, with a first variable light domain covalently attached to the C-terminus of the first Fc domain using a domain linker. The second monomer comprises a second variable heavy domain of the second constant heavy domain comprising a second Fc domain, and a third variable heavy domain covalently attached to the C-terminus of the second Fc domain using a domain linker. The two C-terminally attached variable domains make up a scFv that binds CD3. This embodiment further utilizes a common light chain comprising a variable light domain and a constant light domain, that associates with the heavy chains to form two identical Fabs that bind a TTA. . As for many of the embodiments herein, these constructs include skew variants, pl variants, ablation variants, additional Fc variants, etc. as desired and described herein.

[00271] The present invention provides mAb-Fv formats where the anti-CD3 scFv sequences are as shown in Figure 2 to Figure 7 and Figure 68.

[00272] The present invention provides mAb-Fv formats wherein the anti-CD38 sequences are as shown in Figures 8 to 10.

[00273] The present invention provides mAb-Fv formats with CD20 antigen binding domains wherein the anti-CD20 sequences are as shown in the Figures.

[00274] The present invention provides mAb-Fv formats with CD19 antigen binding domains wherein the anti-CD19 sequences are as shown in in the Figures.

[00275] The present invention provides mAb-Fv formats with CD123 antigen binding domains wherein the anti-CD123 sequences are as shown in in the Figures.

[00276] The present invention provides mAb-Fv formats comprising ablation variants as shown in Figure 31.

[00277] The present invention provides mAb-Fv formats comprising skew variants as shown in Figures 29 and 34.

mAb-scFv

[00278] One heterodimeric scaffold that finds particular use in the present invention is the mAb-Fv format shown in Figure 1. In this embodiment, the format relies on the use of a C-terminal attachment of a scFv to one of the monomers, thus forming a third antigen binding domain, wherein the Fab portions of the two monomers bind a TTA and the “extra” scFv domain binds CD3. Thus, the first monomer comprises a first heavy chain (comprising a variable heavy domain and a constant domain), with a C-terminally covalently attached scFv comprising a scFv variable light domain, an scFv linker and a scFv variable heavy domain. This embodiment further utilizes a common light chain comprising a variable light domain and a constant light domain, that associates with the heavy chains to form two identical Fabs that bind a TTA. As for many of the embodiments herein, these constructs include skew variants, pI variants, ablation variants, additional Fc variants, etc. as desired and described herein.

[00279] The present invention provides mAb-Fv formats where the anti-CD3 scFv sequences are as shown in Figure 2 to Figure 7 and Figure 68.

[00280] The present invention provides mAb-Fv formats wherein the anti-CD38 sequences are as shown in Figures 8 to 10.

[00281] The present invention provides mAb-Fv formats with CD20 antigen binding domains wherein the anti-CD20 sequences are as shown in in the Figures.

[00282] The present invention provides mAb-Fv formats with CD19 antigen binding domains wherein the anti-CD19 sequences are as shown in in the Figures.

[00283] The present invention provides mAb-Fv formats with CD123 antigen binding domains wherein the anti-CD123 sequences are as shown in in the Figures.

[00284] The present invention provides mAb-Fv formats comprising ablation variants as shown in Figure 31.

[00285] The present invention provides mAb-Fv formats comprising skew variants as shown in Figures 29 and 34.

Central scFv

[00286] One heterodimeric scaffold that finds particular use in the present invention is the Central-scFv format shown in Figure 1. In this embodiment, the format relies on the use of an inserted scFv domain thus forming a third antigen binding domain, wherein the Fab portions of the two monomers bind a TTA and the “extra” scFv domain binds CD3. The scFv domain is inserted between the Fc domain and the CH1-Fv region of one of the monomers, thus providing a third antigen binding domain.

[00287] In this embodiment, one monomer comprises a first heavy chain comprising a first variable heavy domain, a CH1 domain and Fc domain, with a scFv comprising a scFv variable light domain, an scFv linker and a scFv variable heavy domain. The scFv is covalently attached between the C-terminus of the CH1 domain of the heavy constant domain and the N-terminus of the first Fc domain using domain linkers. This embodiment further utilizes a common light chain comprising a variable light domain and a constant light domain, that associates with the heavy chains to form two identical Fabs that bind a TTA. As for many of the embodiments herein, these constructs include skew variants, pl variants, ablation variants, additional Fc variants, etc. as desired and described herein.

[00288] The present invention provides Central-scFv formats where the anti-CD3 scFv sequences are as shown in Figure 2 to Figure 7 and Figure 68.

[00289] The present invention provides Central-scFv formats wherein the anti-CD38 sequences are as shown in Figures 8 to 10.

[00290] The present invention provides Central-scFv formats with CD20 antigen binding domains wherein the anti-CD20 sequences are as shown in in the Figures.

[00291] The present invention provides Central-scFv formats with CD19 antigen binding domains wherein the anti-CD19 sequences are as shown in in the Figures.

[00292] The present invention provides Central-scFv formats with CD123 antigen binding domains wherein the anti-CD123 sequences are as shown in v

[00293] The present invention provides Central-scFv formats comprising ablation variants as shown in Figure 31.

[00294] The present invention provides Central-scFv formats comprising skew variants as shown in Figures 29 and 34.

Central-Fv format

[00295] One heterodimeric scaffold that finds particular use in the present invention is the Central-Fv format shown in Figure 1. In this embodiment, the format relies on the use of an inserted scFv domain thus forming a third antigen binding domain, wherein the Fab portions of the two monomers bind a TTA and the “extra” scFv domain binds CD3. The scFv domain is inserted between the Fc domain and the CH1-Fv region of the monomers, thus providing a third antigen binding domain, wherein each monomer contains a component of the scFv (e.g. one monomer comprises a variable heavy domain and the other a variable light domain).

[00296] In this embodiment, one monomer comprises a first heavy chain comprising a first variable heavy domain, a CH1 domain and Fc domain and an additional variable light domain. The light domain is covalently attached between the C-terminus of the CH1 domain of the heavy constant domain and the N-terminus of the first Fc domain using domain linkers. The other monomer comprises a first heavy chain comprising a first variable heavy domain, a CH1 domain and Fc domain and an additional variable heavy domain. The light domain is covalently attached between the C-terminus of the CH1 domain of the heavy constant domain and the N-terminus of the first Fc domain using domain linkers.

[00297] This embodiment further utilizes a common light chain comprising a variable light domain and a constant light domain, that associates with the heavy chains to form two identical Fabs that bind a TTA. As for many of the embodiments herein, these constructs include skew variants, pI variants, ablation variants, additional Fc variants, etc. as desired and described herein.

[00298] The present invention provides Central-Fv formats where the anti-CD3 scFv sequences are as shown in Figure 2 to Figure 7 and Figure 68.

[00299] The present invention provides Central-Fv formats wherein the anti-CD38 sequences are as shown in Figures 8 to 10.

[00300] The present invention provides Central-Fv formats with CD20 antigen binding domains wherein the anti-CD20 sequences are as shown in the Figures.

[00301] The present invention provides Central-Fv formats with CD19 antigen binding domains wherein the anti-CD19 sequences are as shown in the Figures.

[00302] The present invention provides Central-Fv formats with CD123 antigen binding domains wherein the anti-CD123 sequences are as shown in the Figures.

[00303] The present invention provides Central-Fv formats comprising ablation variants as shown in Figure 31.

[00304] The present invention provides Central-Fv formats comprising skew variants as shown in Figures 29 and 34.

One armed central-scFv

[00305] One heterodimeric scaffold that finds particular use in the present invention is the one armed central-scFv format shown in Figure 1. In this embodiment, one monomer comprises just an Fc domain, while the other monomer uses an inserted scFv domain thus forming the second antigen binding domain. In this format, either the Fab portion binds a TTA and the scFv binds CD3 or vice versa. The scFv domain is inserted between the Fc domain and the CH1-Fv region of one of the monomers.

[00306] In this embodiment, one monomer comprises a first heavy chain comprising a first variable heavy domain, a CH1 domain and Fc domain, with a scFv comprising a scFv variable light domain, an scFv linker and a scFv variable heavy domain. The scFv is covalently attached between the C-terminus of the CH1 domain of the heavy constant domain and the N-terminus of the first Fc domain using domain linkers. The second monomer comprises an Fc domain. This embodiment further utilizes a light chain comprising a variable light domain and a constant light domain, that associates with the heavy chain to form a Fab. As for many of the embodiments herein, these constructs include skew variants, pI variants, ablation variants, additional Fc variants, etc. as desired and described herein.

[00307] The present invention provides one armed central-scFv formats where the anti-CD3 scFv sequences are as shown in Figure 2 to Figure 7 and Figure 68.

[00308] The present invention provides one armed central-scFv formats wherein the anti-CD38 sequences are as shown in Figures 8 to 10.

[00309] The present invention provides one armed central-scFv formats with CD20 antigen binding domains wherein the anti-CD20 sequences are as shown in the Figures.

[00310] The present invention provides one armed central-scFv formats with CD19 antigen binding domains wherein the anti-CD19 sequences are as shown in the Figures.

[00311] The present invention provides one armed central-scFv formats with CD123 antigen binding domains wherein the anti-CD123 sequences are as shown in the Figures.

[00312] The present invention provides one armed central-scFv formats comprising ablation variants as shown in Figure 31.

[00313] The present invention provides one armed central-scFv formats comprising skew variants as shown in Figures 29 and 34.

Dual scFv formats

[00314] The present invention also provides dual scFv formats as are known in the art and shown in Figure 1.

[00315] The present invention provides dual scFv formats where the anti-CD3 scFv sequences are as shown in Figure 2 to Figure 7 and Figure 68.

[00316] The present invention provides dual scFv formats wherein the anti-CD38 sequences are as shown in Figures 8 to 10.

[00317] The present invention provides dual scFv formats with CD20 antigen binding domains wherein the anti-CD20 sequences are as shown in the Figures.

[00318] The present invention provides dual scFv formats with CD19 antigen binding domains wherein the anti-CD19 sequences are as shown in the Figures.

[00319] The present invention provides dual scFv formats with CD123 antigen binding domains wherein the anti-CD123 sequences are as shown in the Figures.

[00320] The present invention provides dual scFv formats comprising ablation variants as shown in Figure 31.

[00321] The present invention provides dual scFv formats comprising skew variants as shown in Figures 29 and 34.

Target Antigens

[00322] The bispecific antibodies of the invention have two different antigen binding domains: one that binds to CD3 (generally monovalently), and one that binds to a target tumor antigen (sometimes referred to herein as “TTA”). Suitable target tumor antigens include, but are not limited to, CD20, CD38, CD123; ROR1, ROR2, BCMA; PSMA; SSTR2; SSTR5, CD19, FLT3, CD33, PSCA, ADAM 17, CEA, Her2, EGFR, EGFR-vIII, CD30, FOLR1, GD-2, CA-IX, Trop-2, CD70, CD38, mesothelin, EphA2, CD22, CD79b, GPNMB, CD56, CD138, CD52, CD74, CD30, CD123, RON, ERBB2, and EGFR.

[00323] The “triple F” format is particularly beneficial for targeting two (or more) distinct antigens. (As outlined herein, this targeting can be any combination of monovalent and divalent binding, depending on the format). Thus the immunoglobulins herein preferably co-engage two target antigens. Each monomer’s specificity can be selected from the lists herein. Additional useful bispecific formats for use with an anti-CD3 binding domain are shown in Figure 1.

[00324] Particular suitable applications of the heterodimeric antibodies herein are co-target pairs for which it is beneficial or critical to engage each target antigen monovalently. Such antigens may be, for example, immune receptors that are activated upon immune complexation. Cellular activation of many immune receptors occurs only by cross-linking, achieved typically by antibody/antigen immune complexes, or via effector cell to target cell engagement. For some immune receptors, for example the CD3 signaling receptor on T cells, activation only upon engagement with co-engaged target is critical, as nonspecific cross-linking in a clinical setting can elicit a cytokine storm and toxicity. Therapeutically, by engaging such antigens monovalently rather than multivalently, using the immunoglobulins herein, such activation occurs only in response to cross-linking only in the microenvironment of the primary target antigen. The ability to target two different antigens

with different valencies is a novel and useful aspect of the present invention. Examples of target antigens for which it may be therapeutically beneficial or necessary to co-engage monovalently include but are not limited to immune activating receptors such as CD3, FcγRs, toll-like receptors (TLRs) such as TLR4 and TLR9, cytokine, chemokine, cytokine receptors, and chemokine receptors. In many embodiments, one of the antigen binding sites binds to CD3, and in some embodiments it is the scFv-containing monomer.

[00325] Virtually any antigen may be targeted by the immunoglobulins herein, including but not limited to proteins, subunits, domains, motifs, and/or epitopes belonging to the following list of target antigens, which includes both soluble factors such as cytokines and membrane-bound factors, including transmembrane receptors: 17-IA, 4-1BB, 4Dc, 6-keto-PGF1a, 8-iso-PGF2a, 8-oxo-dG, A1 Adenosine Receptor, A33, ACE, ACE-2, Activin, Activin A, Activin AB, Activin B, Activin C, Activin RIA, Activin RIA ALK-2, Activin RIB ALK-4, Activin RIIA, Activin RIIB, ADAM, ADAM10, ADAM12, ADAM15, ADAM17/TACE, ADAM8, ADAM9, ADAMTS, ADAMTS4, ADAMTS5, Addressins, aFGF, ALCAM, ALK, ALK-1, ALK-7, alpha-1-antitrypsin, alpha-V/beta-1 antagonist, ANG, Ang, APAF-1, APE, APJ, APP, APRIL, AR, ARC, ART, Artemin, anti-Id, ASPARTIC, Atrial natriuretic factor, av/b3 integrin, Axl, b2M, B7-1, B7-2, B7-H, B-lymphocyte Stimulator (BlyS), BACE, BACE-1, Bad, BAFF, BAFF-R, Bag-1, BAK, Bax, BCA-1, BCAM, Bcl, BCMA, BDNF, b-ECGF, bFGF, BID, Bik, BIM, BLC, BL-CAM, BLK, BMP, BMP-2 BMP-2a, BMP-3 Osteogenin, BMP-4 BMP-2b, BMP-5, BMP-6 Vgr-1, BMP-7 (OP-1), BMP-8 (BMP-8a, OP-2), BMPR, BMPR-IA (ALK-3), BMPR-IB (ALK-6), BRK-2, RPK-1, BMPR-II (BRK-3), BMPs, b-NGF, BOK, Bombesin, Bone-derived neurotrophic factor, BPDE, BPDE-DNA, BTC, complement factor 3 (C3), C3a, C4, C5, C5a, C10, CA125, CAD-8, Calcitonin, cAMP, carcinoembryonic antigen (CEA), carcinoma-associated antigen, Cathepsin A, Cathepsin B, Cathepsin C/DPPI, Cathepsin D, Cathepsin E, Cathepsin H, Cathepsin L, Cathepsin O, Cathepsin S, Cathepsin V, Cathepsin X/Z/P, CBL, CCL, CCK2, CCL, CCL1, CCL11, CCL12, CCL13, CCL14, CCL15, CCL16, CCL17, CCL18, CCL19, CCL2, CCL20, CCL21, CCL22, CCL23, CCL24, CCL25, CCL26, CCL27, CCL28, CCL3, CCL4, CCL5, CCL6, CCL7, CCL8, CCL9/10, CCR, CCR1, CCR10, CCR10, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CD1, CD2, CD3, CD3E, CD4, CD5, CD6, CD7, CD8, CD10, CD11a, CD11b, CD11c,

CD13, CD14, CD15, CD16, CD18, CD19, CD20, CD21, CD22, CD23, CD25, CD27L, CD28, CD29, CD30, CD30L, CD32, CD33 (p67 proteins), CD34, CD38, CD40, CD40L, CD44, CD45, CD46, CD49a, CD52, CD54, CD55, CD56, CD61, CD64, CD66e, CD74, CD80 (B7-1), CD89, CD95, CD123, CD137, CD138, CD140a, CD146, CD147, CD148, CD152, CD164, CEACAM5, CFTR, cGMP, CINC, Clostridium botulinum toxin, Clostridium perfringens toxin, CKb8-1, CLC, CMV, CMV UL, CNTF, CNTN-1, COX, C-Ret, CRG-2, CT-1, CTACK, CTGF, CTLA-4, CX3CL1, CX3CR1, CXCL, CXCL1, CXCL2, CXCL3, CXCL4, CXCL5, CXCL6, CXCL7, CXCL8, CXCL9, CXCL10, CXCL11, CXCL12, CXCL13, CXCL14, CXCL15, CXCL16, CXCR, CXCR1, CXCR2, CXCR3, CXCR4, CXCR5, CXCR6, cytokeratin tumor-associated antigen, DAN, DCC, DcR3, DC-SIGN, Decay accelerating factor, des(1-3)-IGF-I (brain IGF-1), Dhh, digoxin, DNAM-1, Dnase, Dpp, DPPIV/CD26, Dtk, ECAD, EDA, EDA-A1, EDA-A2, EDAR, EGF, EGFR (ErbB-1), EMA, EMMPRIN, ENA, endothelin receptor, Enkephalinase, eNOS, Eot, eotaxin1, EpCAM, Ephrin B2/ EphB4, EPO, ERCC, E-selectin, ET-1, Factor IIa, Factor VII, Factor VIIIc, Factor IX, fibroblast activation protein (FAP), Fas, FcR1, FEN-1, Ferritin, FGF, FGF-19, FGF-2, FGF3, FGF-8, FGFR, FGFR-3, Fibrin, FL, FLIP, Flt-3, Flt-4, Follicle stimulating hormone, Fractalkine, FZD1, FZD2, FZD3, FZD4, FZD5, FZD6, FZD7, FZD8, FZD9, FZD10, G250, Gas 6, GCP-2, GCSF, GD2, GD3, GDF, GDF-1, GDF-3 (Vgr-2), GDF-5 (BMP-14, CDMP-1), GDF-6 (BMP-13, CDMP-2), GDF-7 (BMP-12, CDMP-3), GDF-8 (Myostatin), GDF-9, GDF-15 (MIC-1), GDNF, GDNF, GFAP, GFRa-1, GFR-alpha1, GFR-alpha2, GFR-alpha3, GITR, Glucagon, Glut 4, glycoprotein IIb/IIIa (GP IIb/IIIa), GM-CSF, gp130, gp72, GRO, Growth hormone releasing factor, Hapten (NP-cap or NIP-cap), HB-EGF, HCC, HCMV gB envelope glycoprotein, HCMV) gH envelope glycoprotein, HCMV UL, Hemopoietic growth factor (HGF), Hep B gp120, heparanase, Her2, Her2/neu (ErbB-2), Her3 (ErbB-3), Her4 (ErbB-4), herpes simplex virus (HSV) gB glycoprotein, HSV gD glycoprotein, HGFA, High molecular weight melanoma-associated antigen (HMW-MAA), HIV gp120, HIV IIIB gp 120 V3 loop, HLA, HLA-DR, HM1.24, HMFG PEM, HRG, Hrk, human cardiac myosin, human cytomegalovirus (HCMV), human growth hormone (HGH), HVEM, I-309, IAP, ICAM, ICAM-1, ICAM-3, ICE, ICOS, IFNg, Ig, IgA receptor, IgE, IGF, IGF binding proteins, IGF-1R, IGFBP, IGF-I, IGF-II, IL, IL-1, IL-1R, IL-2, IL-2R, IL-4, IL-4R, IL-5, IL-5R, IL-6, IL-6R, IL-8, IL-9, IL-10, IL-12, IL-13, IL-15, IL-18, IL-18R, IL-23, interferon (INF)-alpha, INF-beta, INF-

gamma, Inhibin, iNOS, Insulin A-chain, Insulin B-chain, Insulin-like growth factor 1, integrin alpha2, integrin alpha3, integrin alpha4, integrin alpha4/beta1, integrin alpha4/beta7, integrin alpha5 (alphaV), integrin alpha5/beta1, integrin alpha5/beta3, integrin alpha6, integrin beta1, integrin beta2, interferon gamma, IP-10, I-TAC, JE, Kallikrein 2, Kallikrein 5, Kallikrein 6, , Kallikrein 11, Kallikrein 12, Kallikrein 14, Kallikrein 15, Kallikrein L1, Kallikrein L2, Kallikrein L3, Kallikrein L4, KC, KDR, Keratinocyte Growth Factor (KGF), laminin 5, LAMP, LAP, LAP (TGF- 1), Latent TGF-1, Latent TGF-1 bp1, LBP, LDGF, LECT2, Lefty, Lewis-Y antigen, Lewis-Y related antigen, LFA-1, LFA-3, Lfo, LIF, LIGHT, lipoproteins, LIX, LKN, Lptn, L-Selectin, LT-a, LT-b, LTB4, LTBP-1, Lung surfactant, Luteinizing hormone, Lymphotoxin Beta Receptor, Mac-1, MAdCAM, MAG, MAP2, MARC, MCAM, MCAM, MCK-2, MCP, M-CSF, MDC, Mer, METALLOPROTEASES, MGDF receptor, MGMT, MHC (HLA-DR), MIF, MIG, MIP, MIP-1-alpha, MK, MMAC1, MMP, MMP-1, MMP-10, MMP-11, MMP-12, MMP-13, MMP-14, MMP-15, MMP-2, MMP-24, MMP-3, MMP-7, MMP-8, MMP-9, MPIF, Mpo, MSK, MSP, mucin (Muc1), MUC18, Muellierian-inhibitin substance, Mug, MuSK, NAIP, NAP, NCAD, N-Cadherin, NCA 90, NCAM, NCAM, Neprilysin, Neurotrophin-3,-4, or -6, Neurturin, Neuronal growth factor (NGF), NGFR, NGF-beta, nNOS, NO, NOS, Npn, NRG-3, NT, NTN, OB, OGG1, OPG, OPN, OSM, OX40L, OX40R, p150, p95, PADPr, Parathyroid hormone, PARC, PARP, PBR, PBSF, PCAD, P-Cadherin, PCNA, PDGF, PDGF, PDK-1, PECAM, PEM, PF4, PGE, PGF, PGI2, PGJ2, PIN, PLA2, placental alkaline phosphatase (PLAP), PIGF, PLP, PP14, Proinsulin, Prorelaxin, Protein C, PS, PSA, PSCA, prostate specific membrane antigen (PSMA), PTEN, PTHrp, Ptk, PTN, R51, RANK, RANKL, RANTES, RANTES, Relaxin A-chain, Relaxin B-chain, renin, respiratory syncytial virus (RSV) F, RSV Fgp, Ret, Rheumatoid factors, RLIP76, RPA2, RSK, S100, SCF/KL, SDF-1, SERINE, Serum albumin, sFRP-3, Shh, SIGIRR, SK-1, SLAM, SLPI, SMAC, SMDF, SMOH, SOD, SPARC, Stat, STEAP, STEAP-II, TACE, TACI, TAG-72 (tumor-associated glycoprotein-72), TARC, TCA-3, T-cell receptors (e.g., T-cell receptor alpha/beta), TdT, TECK, TEM1, TEM5, TEM7, TEM8, TERT, testicular PLAP-like alkaline phosphatase, TfR, TGF, TGF-alpha, TGF-beta, TGF-beta Pan Specific, TGF-beta RI (ALK-5), TGF-beta RII, TGF-beta RIIB, TGF-beta RIII, TGF-beta1, TGF-beta2, TGF-beta3, TGF-beta4, TGF-beta5, Thrombin, Thymus Ck-1, Thyroid stimulating hormone, Tie, TIMP, TIQ, Tissue Factor,

TMEFF2, Tmpo, TMPRSS2, TNF, TNF-alpha, TNF-alpha beta, TNF-beta2, TNFc, TNF-RI, TNF-RII, TNFRSF10A (TRAIL R1 Apo-2, DR4), TNFRSF10B (TRAIL R2 DR5, KILLER, TRICK-2A, TRICK-B), TNFRSF10C (TRAIL R3 DcR1, LIT, TRID), TNFRSF10D (TRAIL R4 DcR2, TRUNDD), TNFRSF11A (RANK ODF R, TRANCE R), TNFRSF11B (OPG OCIF, TR1), TNFRSF12 (TWEAK R FN14), TNFRSF13B (TACI), TNFRSF13C (BAFF R), TNFRSF14 (HVEM ATAR, HveA, LIGHT R, TR2), TNFRSF16 (NGFR p75NTR), TNFRSF17 (BCMA), TNFRSF18 (GITR AITR), TNFRSF19 (TROY TAJ, TRADE), TNFRSF19L (RELT), TNFRSF1A (TNF RI CD120a, p55-60), TNFRSF1B (TNF RII CD120b, p75-80), TNFRSF26 (TNFRH3), TNFRSF3 (LTbR TNF RIII, TNFC R), TNFRSF4 (OX40 ACT35, TXGP1 R), TNFRSF5 (CD40 p50), TNFRSF6 (Fas Apo-1, APT1, CD95), TNFRSF6B (DcR3 M68, TR6), TNFRSF7 (CD27), TNFRSF8 (CD30), TNFRSF9 (4-1BB CD137, ILA), TNFRSF21 (DR6), TNFRSF22 (DcTRAIL R2 TNFRH2), TNFRST23 (DcTRAIL R1 TNFRH1), TNFRSF25 (DR3 Apo-3, LARD, TR-3, TRAMP, WSL-1), TNFSF10 (TRAIL Apo-2 Ligand, TL2), TNFSF11 (TRANCE/RANK Ligand ODF, OPG Ligand), TNFSF12 (TWEAK Apo-3 Ligand, DR3 Ligand), TNFSF13 (APRIL TALL2), TNFSF13B (BAFF BLYS, TALL1, THANK, TNFSF20), TNFSF14 (LIGHT HVEM Ligand, LTg), TNFSF15 (TL1A/VEGI), TNFSF18 (GITR Ligand AITR Ligand, TL6), TNFSF1A (TNF-a Conectin, DIF, TNFSF2), TNFSF1B (TNF-b LTa, TNFSF1), TNFSF3 (LTb TNFC, p33), TNFSF4 (OX40 Ligand gp34, TXGP1), TNFSF5 (CD40 Ligand CD154, gp39, HIGM1, IMD3, TRAP), TNFSF6 (Fas Ligand Apo-1 Ligand, APT1 Ligand), TNFSF7 (CD27 Ligand CD70), TNFSF8 (CD30 Ligand CD153), TNFSF9 (4-1BB Ligand CD137 Ligand), TP-1, t-PA, Tpo, TRAIL, TRAIL R, TRAIL-R1, TRAIL-R2, TRANCE, transferring receptor, TRF, Trk, TROP-2, TSG, TSLP, tumor-associated antigen CA 125, tumor-associated antigen expressing Lewis Y related carbohydrate, TWEAK, TXB2, Ung, uPAR, uPAR-1, Urokinase, VCAM, VCAM-1, VECAD, VE-Cadherin, VE-cadherin-2, VEGFR-1 (flt-1), VEGF, VEGFR, VEGFR-3 (flt-4), VEGI, VIM, Viral antigens, VLA, VLA-1, VLA-4, VNR integrin, von Willebrands factor, WIF-1, WNT1, WNT2, WNT2B/13, WNT3, WNT3A, WNT4, WNT5A, WNT5B, WNT6, WNT7A, WNT7B, WNT8A, WNT8B, WNT9A, WNT9A, WNT9B, WNT10A, WNT10B, WNT11, WNT16, XCL1, XCL2, XCR1, XCR1, XEDAR, XIAP, XPD, and receptors for hormones and growth factors.

[00326] Exemplary antigens that may be targeted specifically by the immunoglobulins of the invention include but are not limited to: CD20, CD19, Her2, EGFR, EpCAM, CD3, FcγRIIIa (CD16), FcγRIIa (CD32a), FcγRIIb (CD32b), FcγRI (CD64), Toll-like receptors (TLRs) such as TLR4 and TLR9, cytokines such as IL-2, IL-5, IL-13, IL-12, IL-23, and TNFα, cytokine receptors such as IL-2R, chemokines, chemokine receptors, growth factors such as VEGF and HGF, and the like. To form the bispecific antibodies of the invention, antibodies to any combination of these antigens can be made; that is, each of these antigens can be optionally and independently included or excluded from a bispecific antibody according to the present invention.

[00327] Particularly preferred combinations for bispecific antibodies are an antigen-binding domain to CD3 and an antigen binding domain selected from a domain that binds CD19, CD20, CD38 and CD123, the sequences of which are shown in the Figures.

Nucleic acids of the Invention

[00328] The invention further provides nucleic acid compositions encoding the bispecific antibodies of the invention. As will be appreciated by those in the art, the nucleic acid compositions will depend on the format and scaffold of the heterodimeric protein. Thus, for example, when the format requires three amino acid sequences, such as for the triple F format (e.g. a first amino acid monomer comprising an Fc domain and a scFv, a second amino acid monomer comprising a heavy chain and a light chain), three nucleic acid sequences can be incorporated into one or more expression vectors for expression. Similarly, some formats (e.g. dual scFv formats such as disclosed in Figure 1) only two nucleic acids are needed; again, they can be put into one or two expression vectors.

[00329] As is known in the art, the nucleic acids encoding the components of the invention can be incorporated into expression vectors as is known in the art, and depending on the host cells used to produce the heterodimeric antibodies of the invention. Generally the nucleic acids are operably linked to any number of regulatory elements (promoters, origin of replication, selectable markers, ribosomal binding sites, inducers, etc.). The expression vectors can be extra-chromosomal or integrating vectors.

[00330] The nucleic acids and/or expression vectors of the invention are then transformed into any number of different types of host cells as is well known in the art, including mammalian, bacterial, yeast, insect and/or fungal cells, with mammalian cells (e.g. CHO cells), finding use in many embodiments.

[00331] In some embodiments, nucleic acids encoding each monomer and the optional nucleic acid encoding a light chain, as applicable depending on the format, are each contained within a single expression vector, generally under different or the same promoter controls. In embodiments of particular use in the present invention, each of these two or three nucleic acids are contained on a different expression vector. As shown herein and in 62/025,931, hereby incorporated by reference, different vector ratios can be used to drive heterodimer formation. That is, surprisingly, while the proteins comprise first monomer:second monomer:light chains (in the case of many of the embodiments herein that have three polypeptides comprising the heterodimeric antibody) in a 1:1:2 ratio, these are not the ratios that give the best results. See **Error! Reference source not found..**

[00332] The heterodimeric antibodies of the invention are made by culturing host cells comprising the expression vector(s) as is well known in the art. Once produced, traditional antibody purification steps are done, including an ion exchange chromatography step. As discussed herein, having the pIs of the two monomers differ by at least 0.5 can allow separation by ion exchange chromatography or isoelectric focusing, or other methods sensitive to isoelectric point. That is, the inclusion of pI substitutions that alter the isoelectric point (pI) of each monomer so that such that each monomer has a different pI and the heterodimer also has a distinct pI, thus facilitating isoelectric purification of the "triple F" heterodimer (e.g., anionic exchange columns, cationic exchange columns). These substitutions also aid in the determination and monitoring of any contaminating dual scFv-Fc and mAb homodimers post-purification (e.g., IEF gels, cIEF, and analytical IEX columns).

Treatments

[00333] Once made, the compositions of the invention find use in a number of applications. CD20, CD38 and CD123 are all unregulated in many hematopoietic malignancies and in cell lines derived from various hematopoietic malignancies,

accordingly, the heterodimeric antibodies of the invention find use in treating cancer, including but not limited to, all B cell lymphomas and leukemias, including but not limited to non-Hodgkin's lymphoma (NHL), Burkitt's lymphoma (BL), multiple myeloma (MM), B chronic lymphocytic leukemia (B-CLL), B and T acute lymphocytic leukemia (ALL), T cell lymphoma (TCL), acute myeloid leukemia (AML), hairy cell leukemia (HCL), Hodgkin's Lymphoma (HL), chronic lymphocytic leukemia (CLL), non-Hodgkin's lymphoma, and chronic myeloid leukemia (CML).

[00334] Accordingly, the heterodimeric compositions of the invention find use in the treatment of these cancers.

Antibody Compositions for In Vivo Administration

[00335] Formulations of the antibodies used in accordance with the present invention are prepared for storage by mixing an antibody having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers (Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. [1980]), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein complexes); and/or non-ionic surfactants such as TWEENTM, PLURONICSTM or polyethylene glycol (PEG).

[00336] The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary

activities that do not adversely affect each other. For example, it may be desirable to provide antibodies with other specificities. Alternatively, or in addition, the composition may comprise a cytotoxic agent, cytokine, growth inhibitory agent and/or small molecule antagonist. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

[00337] The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980).

[00338] The formulations to be used for in vivo administration should be sterile, or nearly so. This is readily accomplished by filtration through sterile filtration membranes.

[00339] Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g. films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and .gamma. ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods.

[00340] When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the

aggregation mechanism is discovered to be intermolecular S--S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

Administrative modalities

[00341] The antibodies and chemotherapeutic agents of the invention are administered to a subject, in accord with known methods, such as intravenous administration as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation routes. Intravenous or subcutaneous administration of the antibody is preferred.

Treatment modalities

[00342] In the methods of the invention, therapy is used to provide a positive therapeutic response with respect to a disease or condition. By "positive therapeutic response" is intended an improvement in the disease or condition, and/or an improvement in the symptoms associated with the disease or condition. For example, a positive therapeutic response would refer to one or more of the following improvements in the disease: (1) a reduction in the number of neoplastic cells; (2) an increase in neoplastic cell death; (3) inhibition of neoplastic cell survival; (5) inhibition (i.e., slowing to some extent, preferably halting) of tumor growth; (6) an increased patient survival rate; and (7) some relief from one or more symptoms associated with the disease or condition.

[00343] Positive therapeutic responses in any given disease or condition can be determined by standardized response criteria specific to that disease or condition. Tumor response can be assessed for changes in tumor morphology (i.e., overall tumor burden, tumor size, and the like) using screening techniques such as magnetic resonance imaging (MRI) scan, x-radiographic imaging, computed tomographic (CT) scan, bone scan imaging, endoscopy, and tumor biopsy sampling including bone marrow aspiration (BMA) and counting of tumor cells in the circulation.

[00344] In addition to these positive therapeutic responses, the subject undergoing therapy may experience the beneficial effect of an improvement in the symptoms associated with the disease.

[00345] An improvement in the disease may be characterized as a complete response. By “complete response” is intended an absence of clinically detectable disease with normalization of any previously abnormal radiographic studies, bone marrow, and cerebrospinal fluid (CSF) or abnormal monoclonal protein in the case of myeloma.

[00346] Such a response may persist for at least 4 to 8 weeks, or sometimes 6 to 8 weeks, following treatment according to the methods of the invention. Alternatively, an improvement in the disease may be categorized as being a partial response. By “partial response” is intended at least about a 50% decrease in all measurable tumor burden (i.e., the number of malignant cells present in the subject, or the measured bulk of tumor masses or the quantity of abnormal monoclonal protein) in the absence of new lesions, which may persist for 4 to 8 weeks, or 6 to 8 weeks.

[00347] Treatment according to the present invention includes a “therapeutically effective amount” of the medicaments used. A “therapeutically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve a desired therapeutic result.

[00348] A therapeutically effective amount may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the medicaments to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the antibody or antibody portion are outweighed by the therapeutically beneficial effects.

[00349] A “therapeutically effective amount” for tumor therapy may also be measured by its ability to stabilize the progression of disease. The ability of a compound to inhibit cancer may be evaluated in an animal model system predictive of efficacy in human tumors.

[00350] Alternatively, this property of a composition may be evaluated by examining the ability of the compound to inhibit cell growth or to induce apoptosis by in vitro assays

known to the skilled practitioner. A therapeutically effective amount of a therapeutic compound may decrease tumor size, or otherwise ameliorate symptoms in a subject. One of ordinary skill in the art would be able to determine such amounts based on such factors as the subject's size, the severity of the subject's symptoms, and the particular composition or route of administration selected.

[00351] Dosage regimens are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. Parenteral compositions may be formulated in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier.

[00352] The specification for the dosage unit forms of the present invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

[00353] The efficient dosages and the dosage regimens for the bispecific antibodies used in the present invention depend on the disease or condition to be treated and may be determined by the persons skilled in the art.

[00354] An exemplary, non-limiting range for a therapeutically effective amount of an bispecific antibody used in the present invention is about 0.1-100 mg/kg, such as about 0.1-50 mg/kg, for example about 0.1-20 mg/kg, such as about 0.1-10 mg/kg, for instance about 0.5, about such as 0.3, about 1, or about 3 mg/kg. In another embodiment, the antibody is administered in a dose of 1 mg/kg or more, such as a dose of from 1 to 20 mg/kg, e.g. a dose of from 5 to 20 mg/kg, e.g. a dose of 8 mg/kg.

[00355] A medical professional having ordinary skill in the art may readily determine and prescribe the effective amount of the pharmaceutical composition required. For

example, a physician or a veterinarian could start doses of the medicament employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved.

[00356] In one embodiment, the bispecific antibody is administered by infusion in a weekly dosage of from 10 to 500 mg/kg such as of from 200 to 400 mg/kg. Such administration may be repeated, e.g., 1 to 8 times, such as 3 to 5 times. The administration may be performed by continuous infusion over a period of from 2 to 24 hours, such as of from 2 to 12 hours.

[00357] In one embodiment, the bispecific antibody is administered by slow continuous infusion over a long period, such as more than 24 hours, if required to reduce side effects including toxicity.

[00358] In one embodiment the bispecific antibody is administered in a weekly dosage of from 250 mg to 2000 mg, such as for example 300 mg, 500 mg, 700 mg, 1000 mg, 1500 mg or 2000 mg, for up to 8 times, such as from 4 to 6 times. The administration may be performed by continuous infusion over a period of from 2 to 24 hours, such as of from 2 to 12 hours. Such regimen may be repeated one or more times as necessary, for example, after 6 months or 12 months. The dosage may be determined or adjusted by measuring the amount of compound of the present invention in the blood upon administration by for instance taking out a biological sample and using anti-idiotypic antibodies which target the antigen binding region of the bispecific antibody.

[00359] In a further embodiment, the bispecific antibody is administered once weekly for 2 to 12 weeks, such as for 3 to 10 weeks, such as for 4 to 8 weeks.

[00360] In one embodiment, the bispecific antibody is administered by maintenance therapy, such as, e.g., once a week for a period of 6 months or more.

[00361] In one embodiment, the bispecific antibody is administered by a regimen including one infusion of an bispecific antibody followed by an infusion of an bispecific antibody conjugated to a radioisotope. The regimen may be repeated, e.g., 7 to 9 days later.

[00362] As non-limiting examples, treatment according to the present invention may be provided as a daily dosage of an antibody in an amount of about 0.1-100 mg/kg, such as 0.5, 0.9, 1.0, 1.1, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, 45, 50, 60, 70, 80, 90 or 100 mg/kg, per day, on at least one of day 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40, or alternatively, at least one of week 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 after initiation of treatment, or any combination thereof, using single or divided doses of every 24, 12, 8, 6, 4, or 2 hours, or any combination thereof.

[00363] In some embodiments the bispecific antibody molecule thereof is used in combination with one or more additional therapeutic agents, e.g. a chemotherapeutic agent. Non-limiting examples of DNA damaging chemotherapeutic agents include topoisomerase I inhibitors (e.g., irinotecan, topotecan, camptothecin and analogs or metabolites thereof, and doxorubicin); topoisomerase II inhibitors (e.g., etoposide, teniposide, and daunorubicin); alkylating agents (e.g., melphalan, chlorambucil, busulfan, thiotepa, ifosfamide, carmustine, lomustine, semustine, streptozocin, decarbazine, methotrexate, mitomycin C, and cyclophosphamide); DNA intercalators (e.g., cisplatin, oxaliplatin, and carboplatin); DNA intercalators and free radical generators such as bleomycin; and nucleoside mimetics (e.g., 5-fluorouracil, capecitabine, gemcitabine, fludarabine, cytarabine, mercaptopurine, thioguanine, pentostatin, and hydroxyurea).

[00364] Chemotherapeutic agents that disrupt cell replication include: paclitaxel, docetaxel, and related analogs; vincristine, vinblastin, and related analogs; thalidomide, lenalidomide, and related analogs (e.g., CC-5013 and CC-4047); protein tyrosine kinase inhibitors (e.g., imatinib mesylate and gefitinib); proteasome inhibitors (e.g., bortezomib); NF- κ B inhibitors, including inhibitors of I κ B kinase; antibodies which bind to proteins overexpressed in cancers and thereby downregulate cell replication (e.g., trastuzumab, rituximab, cetuximab, and bevacizumab); and other inhibitors of proteins or enzymes known to be upregulated, over-expressed or activated in cancers, the inhibition of which downregulates cell replication.

[00365] In some embodiments, the antibodies of the invention can be used prior to, concurrent with, or after treatment with Velcade® (bortezomib).

[00366] All cited references are herein expressly incorporated by reference in their entirety.

[00367] Whereas particular embodiments of the invention have been described above for purposes of illustration, it will be appreciated by those skilled in the art that numerous variations of the details may be made without departing from the invention as described in the appended claims.

EXAMPLES

[00368] Examples are provided below to illustrate the present invention. These examples are not meant to constrain the present invention to any particular application or theory of operation. For all constant region positions discussed in the present invention, numbering is according to the EU index as in Kabat (Kabat et al., 1991, Sequences of Proteins of Immunological Interest, 5th Ed., United States Public Health Service, National Institutes of Health, Bethesda, entirely incorporated by reference). Those skilled in the art of antibodies will appreciate that this convention consists of nonsequential numbering in specific regions of an immunoglobulin sequence, enabling a normalized reference to conserved positions in immunoglobulin families. Accordingly, the positions of any given immunoglobulin as defined by the EU index will not necessarily correspond to its sequential sequence.

[00369] General and specific scientific techniques are outlined in US Publications 2015/0307629, 2014/0288275 and WO2014/145806, all of which are expressly incorporated by reference in their entirety and particularly for the techniques outlined therein.

EXAMPLES

EXAMPLE 1: ALTERNATE FORMATS

Bispecifics Production

[00370] Cartoon schematics of anti-CD38 x anti-CD3 bispecifics are shown in Figures 1. Amino acid sequences for alternate format anti-CD38 x anti-CD3 bispecifics are listed in Figure 39 to Figure 43. DNA encoding the three chains needed for bispecific expression were generated by gene synthesis (Blue Heron Biotechnology, Bothell, Wash.) and were

subcloned using standard molecular biology techniques into the expression vector pTT5. Substitutions were introduced using either site-directed mutagenesis (QuikChange, Stratagene, Cedar Creek, Tex.) or additional gene synthesis and subcloning. DNA was transfected into HEK293E cells for expression and resulting proteins were purified from the supernatant using protein A affinity (GE Healthcare) and cation exchange chromatography. Yields following protein A affinity purification are shown in Figure 35. Cation exchange chromatography purification was performed using a HiTrap SP HP column (GE Healthcare) with a wash/equilibration buffer of 50 mM MES, pH 6.0 and an elution buffer of 50 mM MES, pH 6.0 + 1 M NaCl linear gradient (see Figure 36 for chromatograms).

Redirected T Cell Cytotoxicity

[00371] Anti-CD38 x anti-CD3 bispecifics were characterized in vitro for redirected T cell cytotoxicity (RTCC) of the CD38⁺ RPMI8266 myeloma cell line. 10k RPMI8266 cells were incubated for 24 h with 500k human PBMCs. RTCC was measured by LDH fluorescence as indicated (see Figure 37).

EXAMPLE 2

Redirected T Cell Cytotoxicity

[00372] Anti-CD38 x anti-CD3 Fab-scFv-Fc bispecifics were characterized in vitro for redirected T cell cytotoxicity (RTCC) of the CD38⁺ RPMI8266 myeloma cell line. 40k RPMI8266 cells were incubated for 96 h with 400k human PBMCs. RTCC was measured by flow cytometry as indicated (see Figure 44). CD4⁺ and CD8⁺ T cell expression of CD69, Ki-67, and PI-9 were also characterized by flow cytometry and are shown in Figure 45.

Mouse Model of Anti-Tumor Activity

[00373] Four groups of five NOD scid gamma (NSG) mice each were engrafted with 5x10⁶ RPMI8226TrS tumor cells (multiple myeloma, luciferase-expressing) by intravenous tail vein injection on Day -23. On Day 0, mice were engrafted intraperitoneally with 10x10⁶ human PBMCs. After PBMC engraftment on Day 0, test articles are dosed weekly (Days 0, 7) by intraperitoneal injection at dose levels indicated in Figure 4. Study design is further summarized in Figure 46. Tumor growth was monitored by measuring total flux per mouse

using an in vivo imaging system (IVIS®). Both XmAb13551 and XmAb15426 showed substantial anti-tumor effects (see Figure 47 and Figure 48).

Studies in Cynomolgus Monkey

[00374] Cynomolgus monkeys were given a single dose of anti-CD38 x anti-CD3 bispecifics. An anti-RSV x anti-CD3 bispecific control was also included. Dose levels were: 20 µg/kg XmAb13551 (n=2), 0.5 mg/kg XmAb15426 (n=3), 3 mg/kg XmAb14702 (n=3), or 3 mg/kg XmAb13245 (anti-RSV x anti-CD3 control, n=3) (in 3 independent studies). Anti-CD38 x anti-CD3 bispecifics rapidly depleted CD38+ cells in peripheral blood (see Figure 49). Anti-CD38 x anti-CD3 bispecifics resulted in T cell activation as measured by CD69 expression (see Figure 50). Serum levels of IL-6 were also measured (see Figure 51). Note that, compared to XmAb13551, XmAb15426 had an increased duration of CD38+ cell depletion and lower levels of T cell activation and IL-6 production.

[00375] XmAb15426 and XmAb14702 were tested at single doses of 0.5 mg/kg and 3 mg/kg respectively. Both antibodies were well-tolerated at these higher doses, consistent with the moderate levels of IL6 observed in serum from the treated monkeys. Moreover, XmAb15426, with intermediate CD3 affinity, more effectively depleted CD38+ cells at 0.5 mg/kg compared to the original high-affinity XmAb13551 dosed at 2, 5 or 20 µg/kg. Depletion by XmAb15426 was more sustained compared to the highest dose of of XmAb13551 in the previous study (7 vs. 2 days, respectively). Notably, although target cell depletion was greater for XmAb15426, T cell activation (CD69, CD25 and PD1 induction) was much lower in monkeys treated with XmAb15426 even dosed 25-fold higher than the 20 µg/kg XmAb13551 group. XmAb14702, with very low CD3 affinity, had little effect on CD38+ cells and T cell activation.

[00376] These results demonstrate that modulating T cell activation by attenuating CD3 affinity is a promising method to improve the therapeutic window of T cell-engaging bispecific antibodies. This strategy has potential to expand the set of antigens amenable to targeted T cell immunotherapy by improving tolerability and enabling higher dosing to overcome antigen sink clearance with targets such as CD38. We have shown that by

reducing affinity for CD3, XmAb 15426 effectively depletes CD38⁺ cells while minimizing the CRS effects seen with comparable doses of its high-affinity counterpart XmAb13551.

CLAIMS

1. A heterodimeric antibody comprising:
 - a) a first monomer comprising an anti-CD3 scFv-linker-CH2-CH3, wherein said anti-CD3 scFv comprises a first variable heavy domain comprising a VHCDR1 of SEQ ID NO: 2, a VHCDR2 of SEQ ID NO: 3, and VHCDR3 of SEQ ID NO: 4; and a first variable light domain comprising a VLCDR1 of SEQ ID NO: 6, a VLCDR2 of SEQ ID NO: 7, and a VLCDR3 of SEQ ID NO: 8, and wherein said CH2-CH3 is a first Fc domain;
 - b) a second monomer comprising a VH-CH1-hinge-CH2-CH3, wherein VH is a second variable heavy domain and CH2-CH3 is a second Fc domain; and
 - c) a light chain comprising a second variable light domain,wherein said second variable heavy domain and said second variable light domain from a CD20 binding domain;
wherein said second variable heavy domain comprises a VHCDR1 of SEQ ID NO:431, a VHCDR2 of SEQ ID NO:432, and a VHCDR3 of SEQ ID NO:433; and said second variable light domain comprises a VLCDR1 of SEQ ID NO:435, a VLCDR2 of SEQ ID NO:436, and a VLCDR3 of SEQ ID NO:437.
2. The heterodimeric antibody of claim 1, wherein said second variable heavy domain comprises SEQ ID NO:430.
3. The heterodimeric antibody of any one of claim 1 or claim 2, wherein said second variable light domain comprises SEQ ID NO:434.
4. The heterodimeric antibody of claim 1,
wherein said first monomer comprises SEQ ID NO: 324;
wherein said second monomer comprises SEQ ID NO: 323; and
wherein said light chain comprises SEQ ID NO: 325.
5. The heterodimeric antibody of claim 1, wherein the Fc domains comprise an ablation variant of Fig 31.

6. The heterodimeric antibody of claim 1, wherein the first or second Fc domain comprises a pl variant of FIG 30.
7. The heterodimeric antibody of claim 1, wherein the first and second Fc domain comprise a set of amino acid substitutions of figure 29.
8. The heterodimeric antibody of claim 1, wherein the first and second Fc domain have a set of amino acid substitutions selected from the group consisting of: L368D/K370S : S364K; L368D/K370S : S364K/E357L; L368D/K370S : S364K/E357Q; T411E/K360E/Q362E : D401K; L368E/K370S : S364K; K370S : S364K; L368E/K370S : S364K/E357Q; K370S : S364K/E357Q;; L368E/K370S : S364K/E357Q; and K370S : S364K/E357Q, wherein the numbering is according to the EU index as in Kabat.
9. The heterodimeric antibody of claim 1, wherein said first Fc domain comprises the amino acid substitutions S364K/E357Q and said second Fc domain comprises the amino acid substitutions L368D/K370S, wherein the numbering is according to the EU index as in Kabat.
10. The heterodimeric antibody of claim 1, wherein CH1-hinge-CH2-CH3 of said second monomer comprises the amino acid substitutions N208D/Q295E/N384D/Q418E/N421D, wherein the numbering is according to the EU index as in Kabat.
11. The heterodimeric antibody of claim 1, wherein said first and second Fc domains each comprise the amino acid substitutions E233P/L234V/L235A/G236del/S267K, wherein the numbering is according to the EU index as in Kabat.
12. The heterodimeric antibody of claim 1, wherein said anti-CD3 scFv comprises a linker selected from FIG 33.
13. The heterodimeric antibody of claim 1, wherein the scFv comprises a linker (GKPGS)₄.
14. A composition comprising:
- a) a first nucleic acid encoding the first monomer according to any one of claims 1-13;
 - b) a second nucleic encoding the second monomer according to any one of claims 1-13;
 - and

- c) a third nucleic acid encoding the light chain according to any one of claims 1- 13.
15. A composition comprising:
- a) a first expression vector comprising said first nucleic acid of claim 14;
 - b) a second expression vector comprising said second nucleic acid of claim 14; and
 - c) a third expression vector comprising said third nucleic acid of claim 14.
16. A cell comprising:
- a) a first nucleic acid encoding the first monomer according to any one of claims 1-13;
 - b) a second nucleic encoding the second monomer according to any one of claims 1-13;
and
 - c) a third nucleic acid encoding the light chain according to any one of claims 1-13.
17. A cell comprising the composition of claim 14 or 15.
18. A cell expressing the heterodimeric antibody of any of claims 1-13.
19. A method of making a heterodimeric antibody comprising culturing the cell of any of claims 16-18 and isolating said heterodimeric antibody.
20. A pharmaceutical composition comprising the heterodimeric antibody of any one of claims 1-13.
21. The heterodimeric antibody of any one of claims 1-13 or the pharmaceutical composition of claim 20 for use in treatment of hematopoietic malignancies.
22. The heterodimeric antibody of any one of claims 1-13 or pharmaceutical composition of claim 20 for use in treatment of a cancer selected from a group consisting of: B cell lymphomas and leukemias, non-Hodgkin's lymphoma (NHL), Burkitt's lymphoma (BL), multiple myeloma (MM), B chronic lymphocytic leukemia (B-CLL), B and T acute lymphocytic leukemia (ALL), T cell lymphoma (TCL), acute myeloid leukemia (AML), hairy cell leukemia (HCL), Hodgkin's Lymphoma (HL), chronic lymphocytic leukemia (CLL), non-Hodgkin's lymphoma, and chronic myeloid leukemia (CML).