

Melanocyte-regulating peptides

This application claims the benefit of European Patent Application EP19382674.0 filed on August 2nd, 2019.

5 Technical Field

This invention generally relates to the field of melanocyte-related disorders. In particular, the present invention refers to peptides with the ability to regulate melanogenesis, either promoting or reducing the production of melanin, as well as to inhibit the proliferation of malignant melanocytes; to pharmaceutical and cosmetic
10 compositions containing thereof as an active ingredients, as well as to their use in therapy and cosmetics.

Background Art

Through a process called melanogenesis, melanocytes produce melanin, which is a pigment found in the skin,
15 eyes, hair, nasal cavity, and inner ear. This melanogenesis leads to a long-lasting pigmentation, which is in contrast to the pigmentation that originates from oxidation of already-existing melanin.

There are both basal and activated levels of melanogenesis; in general, lighter-skinned people have low basal levels of melanogenesis. Exposure to UV-B radiation causes increased melanogenesis. The purpose of
20 melanogenesis is to protect the hypodermis, the layer under the skin, from damage by UV-B radiation. The color of the melanin is black, allowing it to absorb a majority of the UV-B light and block it from passing through the epidermis.

It is known that during the life of a person apparent changes in the coloring of his/her skin develop and thus,
25 for example, marks on the skin of the face, chest and hands of elderly people appear which are clear signs of aging. Furthermore, the continuous overexposure to UV radiation does not just cause accelerated aging of the skin, known as photoaging, which is characterized by the appearance of signs of skin aging at a much earlier age, among them the appearance of marks in those areas of the skin overexposed to UV radiation, but which can also give rise to the formation of hyperpigmented cancerous lesions or melanomas.

30

It frequently occurs that in an area of a person's skin the density of melanin within the melanocytes is greater than in the surrounding areas and as a consequence the color of the affected area on that person is darker than the rest. These areas are known as areas of hyperpigmentation. Among the causes of hyperpigmentation are hormonal changes, melasma, lentigo, piebaldism, Addison's disease, hypersensitivity
35 to ultraviolet radiation due to agents which favor the action of radiation (phototoxics), or hyperpigmentation as a consequence of an inflammatory lesion. The marks associated with acne, eczema, scars or hair removal belong to this last type of hyperpigmentation and are marks that can even last several years.

It is also possible for a person's skin areas which have lower melanin densities than that in surrounding areas.

Vitiligo, pityriasis alba, tinea versicolor, and postinflammatory effect are common causes of pigment loss. The attenuation of the irregularities of pigmentation are either due to aging and/or photoaging, to hormonal disorders or to post-inflammatory processes and, particularly, the re-establishment of the pigmentation in the areas affected by vitiligo with topical applications is, therefore, of interest to the cosmetic and pharmaceutical
5 sector.

Pigmentary disorders are a global problem. Despite being common, pigmentary disorders remain difficult to treat.

- 10 The main approaches for the treatment of hyperpigmentation disorders are mainly based on preventing the production of melanin by hyperactive melanocytes or promoting removal of melanin from the epidermis by increasing epidermal cell turnover. A number of agents that inhibit tyrosinase, the rate-limiting enzyme involved in melanin synthesis, are currently used to treat disorders of hyperpigmentation with moderate success. These agents include hydroquinone, kojic acid (also referred as "KA"), arbutin, mequinol and azelaic
15 acid. Contact dermatitis is a common side effect occurring in up to 15% of patients who use these agents, limiting their usage. The efficacy of these agents varies significantly from patient to patient. Topical retinoids reduce epidermal melanin via inhibition of tyrosinase and also increase epidermal turnover, thereby facilitating melanin dispersion and removal. However, irritant dermatitis can be a frequent adverse effect, and caution should be used when prescribing higher concentrations of topical retinoids in darker skin types, since resulting
20 irritation may result in erythema, desquamation and further postinflammatory hyperpigmentation. Laser therapy may sometimes be effective for hyperpigmentation. This poses, however, a significant problem in darker skinned patients, who have increased baseline epidermal melanin that is also targeted at these wavelengths.
- 25 Disorders of depigmentation also involve an inflammatory and autoimmune pathogenesis, so topical corticosteroids and immunomodulators may be useful in the treatment regimen. Because postinflammatory hypopigmentation is usually clinically apparent after the inflammatory process has resolved, these same topical agents may not be beneficial at this late stage. Light therapy, including but not limited to psoralens plus UVA (oral or topical), psoralens and natural sunlight, UVB, narrow-band UVB and 308 nm excimer lasers
30 have been used to treat both hypopigmentation and depigmentation. Disorders of depigmentation (e.g., vitiligo) may be particularly difficult to treat because there is complete absence of melanocytes in the interfollicular epidermis. Hence, therapy is aimed at recruiting or transplanting melanocytes to these sites.

In spite of the efforts made, therefore, there is still the need of developing further molecules (peptides) which
35 can efficiently regulate melanocyte activity and, consequently, can be efficient in the treatment of pigmentary disorders.

Summary of Invention

The present inventors have found that peptides of sequences SEQ ID NO: 1 and 2 efficiently regulate melanocyte activity.

5

On one hand, the present inventors have surprisingly found that the peptides of the invention are involved in melanogenesis pathway. Thus, some of the peptides stimulated the production of melanin, which can be an indicia of melanocyte activation, whereas others, such as F22 and F29, drastically reduced the amount of melanin. What is even more remarkable is the fact that the peptides of the invention were more efficient in
10 regulating melanin production when compared with other compounds known in the state of the art. In this regard, it was found that when KA, which is commonly used in hyperpigmentation, was used at a concentration of 1 μ M, the reduction was substantially lower than the reduction achieved with the peptides of the invention at the same concentration. Therefore, these data do not only support the effect of the peptides of the invention on melanin-production, but also illustrates that they are even more efficient than other
15 compounds available in the state of the art currently used with the same purpose.

On the other hand, the present inventors have also found that these peptides regulate melanocyte activity not only in terms of melanin formation, but also in terms of cell proliferation. As it is shown in Table 2, the peptides of the invention were active in inhibiting cancer cell proliferation, in particular melanoma cell lines, including
20 metastatic melanoma cell lines.

Thus, in a first aspect the present invention provides a peptide or a pharmaceutical or cosmetical acceptable salt thereof, the peptide having an amino acid sequence with at least 85% of identity with respect sequence SEQ ID NO: 1 or 2:

25

$(\text{Trp})_m-(\text{Asn})_n-\text{Lys-Gly-Thr-Ile-Leu-Lys-Ala-Ser-Val-Asp-Tyr-Ile-Arg-Lys}-(\text{Leu})_p-(\text{Gln})_q$ (SEQ ID NO: 1)

$(\text{Arg})_{m_x}-(\text{Arg})_{n_x}-\text{Arg-Arg-Phe-Asn-Ile-Asn-Asp-Arg-Ile-Lys-Glu-Leu-Gly-Thr-Leu}-(\text{Ile})_{p_x}-(\text{Pro})_{q_x}$ (SEQ ID NO: 2)

30 wherein

"m", "n", "p", and "q" represent integers and are selected from 0 and 1;

"m_x", "n_x", "p_x", and "q_x" represent integers and are selected from 0 and 1;

a C-terminal end corresponding to $-\text{C(O)}\text{R}_4$;


35 a N-terminal end corresponding to $-\text{NHR}_5$;

R₄ is a radical selected from the group consisting of $-\text{OH}$ and $-\text{NR}_{17}\text{R}_{18}$;

R₅ is a radical selected from the group consisting of $-\text{H}$, $(\text{C}_1-\text{C}_{20})\text{alkyl}$ and $(\text{C}_1-\text{C}_{10})\text{alkyl}$ substituted by one or more radicals selected from the group consisting of: halogen, $(\text{C}_1-\text{C}_{10})\text{alkyl}$, $-\text{OR}_6$, $-\text{NR}_7\text{R}_8$, $-\text{SR}_9$, $-\text{SOR}_{10}$, $-\text{SO}_2\text{R}_{11}$, and $-\text{CO}_2\text{R}_{12}$; and

R₁₇ and R₁₈ are radicals independently selected from the group consisting of: -H and (C₁-C₁₀)alkyl, and (C₁-C₁₀)alkyl substituted by one or more radicals selected from the group consisting of: halogen, (C₁-C₁₀)alkyl, -OR₆, -NR₇R₈, -SR₉, -SOR₁₀, -SO₂R₁₁, and -CO₂R₁₂; the peptide optionally including a linker biradical "L" of formula (I)

5



which is connecting an alpha carbon atom of an amino acid located at position "i" with an alpha carbon atom of an amino acid located at position "i+4" or "i+7" in the peptide sequence

10

wherein

"a" and "b" are the same or different and are 0 or 1;

"c" is comprised from 1 to 10;

15

R₁ and R₃ are biradicals independently selected from the group consisting of: (C₁-C₁₀)alkyl; (C₁-C₁₀)alkyl substituted by one or more radicals selected from the group consisting of: halogen, (C₁-C₁₀)alkyl, -OR₆, -NR₇R₈, -SR₉, -SOR₁₀, -SO₂R₁₁, and -CO₂R₁₂; (C₂-C₁₀)alkenyl; (C₂-C₁₀)alkenyl substituted by one or more radicals selected from the group consisting of: halogen, (C₁-C₁₀)alkyl, -OR₆, -NR₇R₈, -SR₉, -SOR₁₀, -SO₂R₁₁, and -CO₂R₁₂; (C₂-C₁₀)alkynyl; and (C₂-C₁₀)alkynyl substituted by one or more radicals selected from the group consisting of: halogen, (C₁-C₁₀)alkyl, -OR₆, -NR₇R₈, -SR₉, -SOR₁₀, -SO₂R₁₁, and -CO₂R₁₂;

20

25

R₂ is a biradical selected from the group consisting of: -O-, C(=O), C(=O)NR₁₃, C(=O)O, S(=O), S(=O)₂, NR₁₄, (C₁-C₁₀)alkyl, (C₂-C₁₀)alkenyl, (C₂-C₁₀)alkynyl, -NR₁₅-NR₁₆-, -N=N-, -S-S-, and a known ring system comprising from 3 to 14 members, the system comprising from 1 to 3 rings, where:

each one of the rings is saturated, partially unsaturated, or aromatic;

the rings are isolated, partially or totally fused,

30

each one of the members forming the known ring system is selected from the group consisting of: -CH-, -CH₂-, -NH-, -N-, -SH-, -S-, and -O-; and


the ring system is optionally substituted by one or more radicals independently selected from the group consisting of halogen, -OH, -NO₂, (C₁-C₁₀)alkyl, (C₁-C₁₀)haloalkyl, and (C₁-C₁₀)alkyl-O-; and

35

R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, and R₁₆ are radicals independently selected from the group consisting of: -H and (C₁-C₁₀)alkyl; and

the amino acids which are connected by the linker being of formula (II)

40



5

wherein

R_{19} is a monoradical selected from the group consisting of: (C_1-C_{10}) alkyl, (C_2-C_{10}) alkenyl, (C_2-C_{10}) alkynyl, and a known ring system comprising from 3 to 14 members, the system comprising from 1 to 3 rings, where:

- 10 each one of the rings is saturated, partially unsaturated, or aromatic;
 the rings are isolated, partially or totally fused,
 each one of the members forming the known ring system is selected from the group consisting of: -CH-, -CH₂-, -NH-, -N-, -SH-, -S-, and -O-.

15

In a second aspect, the present invention provides a cosmetical or pharmaceutical composition comprising a cosmetically or therapeutically effective amount of the peptide as defined in the first aspect of the invention together with acceptable cosmetical or pharmaceutical excipients and/or carriers.

- 20 In a third aspect, the present invention provides the peptide or pharmaceutical salt thereof as defined in the first aspect of the invention or the pharmaceutical composition of the second aspect of the invention, for use as a medicament. This aspect can be alternatively formulated as the use of a peptide or pharmaceutical salt thereof as defined in the first aspect of the invention or the pharmaceutical composition of the second aspect of the invention, in the manufacture of a medicament for the treatment of a disease. This aspect can also be
 25 alternatively formulated as a method for the treatment of a disease, the method comprising administering an effective therapeutic amount of a peptide or pharmaceutical salt thereof as defined in the first or the pharmaceutical composition of the second aspect of the invention, to a subject in need thereof.

- In a fourth aspect, the present invention provides the peptide or pharmaceutical salt thereof as defined in the
 30 first aspect of the invention or the pharmaceutical composition as defined in the second aspect of the invention for use in the treatment of cancer. This aspect can alternatively be formulated as the use of the peptide or pharmaceutical salt thereof as defined in the first aspect of the invention or the pharmaceutical composition as defined in the second aspect of the invention in the manufacture of a medicament for the treatment of cancer. This aspect can also be alternatively formulated as a method for the treatment of cancer,
 35 the method comprising administering an effective therapeutic amount of the peptide or pharmaceutical salt thereof as defined in the first aspect of the invention or the pharmaceutical composition as defined in the second aspect of the invention, to a subject in need thereof.

In a fifth aspect the present invention also provides the non-therapeutic use of a peptide or cosmetical salt

thereof as defined in the first aspect of the invention or a cosmetical composition as defined in the second aspect of the invention for regulating melanogenesis, particularly for reducing or increasing the production of melanin.

- 5 In a last aspect the present invention provides a cosmetical method for the treatment of the hyper- or hypopigmentation in a subject in need thereof, the method comprising the application of the peptide or cosmetical salt thereof as defined in the first aspect or the cosmetical composition as defined in the second aspect of the invention to the affected area suffering from the hyper- or hypopigmentation.
- 10 In further aspects the present invention provides: (a) a combination comprising the peptide or pharmaceutical salt thereof as defined in the first or second aspect of the invention; and a therapeutic agent, particularly an anti-cancer agent; (b) a combination comprising the peptide or pharmaceutical salt thereof as defined in the first or second aspect of the invention; and a therapeutic agent, particularly an anti-cancer agent, for use as a medicament, more particularly for use in the treatment of cancer; (c) a peptide or pharmaceutical salt thereof
- 15 as defined in the first or second aspect of the invention for use in combination therapy for the prevention or treatment of cancer, wherein the therapy comprises its administration to a subject simultaneously, sequentially or separately with an anti-cancer agent; and (d) an anti-cancer agent for use in combination therapy with the peptide or pharmaceutical salt thereof as defined in the first or second aspect of the invention, wherein the use comprises the prevention or treatment of cancer.

20

Brief Description of Drawings

FIG. 1 shows the percentage of melanin in treated melanoma cells with respect to that of untreated melanoma cells. As positive control Kojic Acid (KA) was used at different concentrations (0.1, 1 and 10 μ M).

25

Detailed description of the invention

- All terms as used herein in this application, unless otherwise stated, shall be understood in their ordinary meaning as known in the art. Other more specific definitions for certain terms as used in the present
- 30 application are as set forth below and are intended to apply uniformly through-out the specification and claims unless an otherwise expressly set out definition provides a broader definition.

For the purposes of the present invention, any ranges given include both the lower and the upper end-points of the range.

35

The present invention provides peptides comprising sequences with at least 85% degree identity with SEQ ID NO: 1 or 2, as it has been stated above.

In the present invention the term "identity" refers to the percentage of residues that are identical in the two

sequences when the sequences are optimally aligned. If, in the optimal alignment, a position in a first sequence is occupied by the same amino acid residue as the corresponding position in the second sequence, the sequences exhibit identity with respect to that position. The level of identity between two sequences (or "percent sequence identity") is measured as a ratio of the number of identical positions shared by the sequences with respect to the size of the sequences (i.e., percent sequence identity = (number of identical positions/total number of positions) x 100).

A number of mathematical algorithms for rapidly obtaining the optimal alignment and calculating identity between two or more sequences are known and incorporated into a number of available software programs. Examples of such programs include the MATCH-BOX, MULTAIN, GCG, FASTA, and ROBUST programs for amino acid sequence analysis, among others. Preferred software analysis programs include the ALIGN, CLUSTAL W, and BLAST programs (e.g., BLAST 2.1, BL2SEQ, and later versions thereof).

For amino acid sequence analysis, a weight matrix, such as the BLOSUM matrixes (e.g., the BLOSUM45, BLOSUM50, BLOSUM62, and BLOSUM80 matrixes), Gonnet matrixes, or PAM matrixes (e.g., the PAM30, PAM70, PAM120, PAM160, PAM250, and PAM350 matrixes), are used in determining identity.

The BLAST programs provide analysis of at least two amino acid sequences, either by aligning a selected sequence against multiple sequences in a database (e.g., GenSeq), or, with BL2SEQ, between two selected sequences. BLAST programs are preferably modified by low complexity filtering programs such as the DUST or SEG programs, which are preferably integrated into the BLAST program operations. If gap existence costs (or gap scores) are used, the gap existence cost preferably is set between about -5 and -15. Similar gap parameters can be used with other programs as appropriate. The BLAST programs and principles underlying them are further described in, e.g., Altschul et al., "Basic local alignment search tool", 1990, J. Mol. Biol. v. 215, pages 403-410.

For multiple sequence analysis, the CLUSTAL W program can be used. The CLUSTAL W program desirably is run using "dynamic" (versus "fast") settings. Amino acid sequences are evaluated using a variable set of BLOSUM matrixes depending on the level of identity between the sequences. The CLUSTAL W program and underlying principles of operation are further described in, e.g., Higgins et al., "CLUSTAL V: improved software for multiple sequence alignment", 1992, CABIOS, 8(2), pages 189-191.

As used herein, the term "pharmaceutical or cosmetical acceptable salt" refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutical and cosmetical acceptable salts are well known in the art. Examples of pharmaceutical and cosmetical acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, trifluoroacetic acid, oxalic acid, maleic acid,

tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutical and cosmetical acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Salts derived from appropriate bases include alkali metal, alkaline earth metal, and ammonium. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutical and cosmetical acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, lower alkyl sulfonate and aryl sulfonate.

15

The term (C₁-C₁₀)alkyl refers to a saturated straight or branched alkyl chain having from 1 to 10 carbon atoms. Illustrative non-limitative examples are: methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, neo-pentyl and n-hexyl.

20 The term (C₁-C₂₀)alkyl refers to a saturated straight or branched alkyl chain having from 1 to 20 carbon atoms.

The term (C₂-C₁₀)alkenyl refers to a saturated straight, or branched alkyl chain containing from 2 to 10 carbon atoms and also containing one or more double bonds. Illustrative non-limitative examples are ethenyl, propenyl, butenyl, 1-methyl-2-buten-1-yl, and the like.

25

The term (C₂-C₁₀)alkynyl refers to a saturated straight, or branched alkyl chain containing from 2 to 20 carbon atoms and also containing one or more triple bonds. Examples include, among others, ethynyl, 1-propynyl, 2-butylnyl, 1,3-butyndynyl, 4-pentylnyl, and 1-hexynyl.

30 The term "halogen" refers to the group in the periodic table consisting of five chemically related elements: fluorine (F), chlorine (Cl), bromine (Br), iodine (I), and astatine (At).

The term (C₁-C₁₀)haloalkyl refers to a group resulting from the replacement of one or more hydrogen atoms from a (C₁-C₁₀)alkyl group with one or more, preferably from 1 to 6, halogen atoms, which can be the same or different. Examples include, among others, trifluoromethyl, fluoromethyl, 1-chloroethyl, 2-chloroethyl, 1-fluoroethyl, 2-fluoroethyl, 2-bromoethyl, 2-iodoethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, 3-fluoropropyl, 3-chloropropyl, 2,2,3,3-tetrafluoropropyl, 2,2,3,3,3-pentafluoropropyl, heptafluoropropyl, 4-fluorobutyl, and nonafluorobutyl.

35

The term "known" ring system as used herein refers to a ring system which is chemically feasible and is known in the art and so intends to exclude those ring systems that are not chemically possible.

According to the present invention when the ring system is formed by "isolated" rings means that the ring system is formed by two, three or four rings and said rings are bound via a bond from the atom of one ring to the atom of the other ring. The term "isolated" also embraces the embodiment in which the ring system has only one ring. Illustrative non-limitative examples of known ring systems consisting of one ring are those derived from: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopropenyl, cyclobutenyl, cyclopentenyl, phenyl, and cycloheptenyl.

10

According to the present invention when the ring system has rings "totally fused", means that the ring system is formed by two, three or four rings in which two or more atoms are common to two adjoining rings. Illustrative non-limitative examples are 1,2,3,4-tetrahydronaphthyl, 1-naphthyl, 2-naphthyl, anthryl, or phenanthryl.

15

According to the present invention when the ring system is "partially fused" it means that the ring system is formed by three or four rings, being at least two of said rings totally fused (i.e. two or more atoms being common to the two adjoining rings) and the remaining ring(s) being bound via a bond from the atom of one ring to the atom of one of the fused rings.

20

Unless otherwise stated, the amino acids forming the peptides of the invention can have L- or D-configuration.

Amino acids used in the construction of peptides of the present invention may be prepared by organic synthesis, or obtained by other routes, such as, for example, degradation of or isolation from a natural source.

25

In one embodiment of the first aspect of the invention, the peptide is other than one of sequence SEQ ID NO: 24, 25 or 26.

In one embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide is one wherein m and n are the same. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide is one wherein p and q are the same. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, m and n represent 1 and p and q represent 0. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, m and n represent 0 and p and q represent 1.

In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide is a lineal peptide.

In the present invention the term "lineal peptide" means that it does not include the "L" biradical.

In one embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide is a lineal peptide consisting of SEQ ID NO: 1, wherein:

- 5 - when "m" and "n" are the same, then "p" is different from "q"; and
- when m is 0, and n=p= 1, then q is 1.

In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide is a lineal peptide consisting of SEQ ID NO: 1, and is

10 selected from the peptides wherein:

- at least three of m, n, p and q are the same; or, alternatively,
- when "m" is 0 and "n" is 1, then "p" is 0 and "q" is 0 or 1; or, alternatively
- when "m" is 0 and "n" is 1, then "p" is 0 or 1, and "q" is 1.

- 15 In one embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide is one wherein m_x and n_x are the same. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide is one wherein p_x and q_x are the same. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, m_x and n_x represent 1 and p_x
- 20 and q_x represent 0. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, m_x and n_x represent 0 and p_x and q_x represent 1.

In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one comprising a "L" linker wherein a= 1.

- 25 In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one comprising a "L" linker wherein b= 1.

In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one comprising a "L" linker wherein c= 1.

- 30 In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one comprising a "L" linker wherein a=b=c= 1.

In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one comprising a "L" linker wherein R_1

- 35 and R_3 are biradicals independently selected from the group consisting of: $(C_1-C_{10})alkyl$; $(C_2-C_{10})alkenyl$; and $(C_2-C_{10})alkynyl$. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one comprising a "L" linker wherein R_1 and R_3 are the same or different and represent $(C_1-C_{10})alkyl$.

In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one comprising a "L" linker wherein R₂ is a biradical selected from the group consisting: -O-, C(=O), C(=O)NR₁₃, C(=O)O, S(=O), S(=O)₂, NR₁₄, (C₁-C₁₀)alkyl, (C₂-C₁₀)alkenyl, (C₂-C₁₀)alkynyl, -NR₁₅-NR₁₆-, -N=N-, -S-S-, and a known ring system consisting of
 5 one ring from 3 to 6 members, the ring:

being saturated, partially unsaturated, or aromatic;
 each one of the members forming the known ring system being selected from the group consisting of: -CH-, -CH₂-, -NH-, -N-, -SH-, -S-, and -O-; and
 10 the ring system being optionally substituted by one or more radicals independently selected from the group consisting of halogen, -OH, -NO₂, (C₁-C₁₀)alkyl, (C₁-C₁₀)haloalkyl, and (C₁-C₁₀)alkyl-O-.

In another embodiment of the first aspect of the invention, optionally in combination with any of the
 15 embodiments provided above or below, the peptide or salt thereof is one comprising a "L" linker wherein R₂ is a biradical selected from the group consisting of: (C₁-C₁₀)alkyl, (C₂-C₁₀)alkenyl, and (C₂-C₁₀)alkynyl. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one comprising a "L" linker wherein R₂ is (C₂-C₁₀)alkenyl.

20 In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one comprising a "L" linker wherein R₁ and R₃ are the same or different and represent (C₁-C₁₀)alkyl; and R₂ is (C₂-C₁₀)alkenyl.

25 In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one comprising a "L" linker wherein R₁ and R₃ are the same or different and represent (C₁-C₁₀)alkyl; R₂ is (C₂-C₁₀)alkenyl; and a=b=c= 1.

In another embodiment of the first aspect of the invention, optionally in combination with any of the
 30 embodiments provided above or below, the peptide or salt thereof is one wherein R₁₉ is selected from the group consisting of: (C₁-C₁₀)alkyl, (C₂-C₁₀)alkenyl, and (C₂-C₁₀)alkynyl. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one wherein R₁₉ is a (C₁-C₁₀)alkyl monoradical. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the
 35 peptide or salt thereof is one comprising a "L" linker wherein R₁, R₃ and R₁₉ are the same or different and represent (C₁-C₁₀)alkyl; and R₂ is (C₂-C₁₀)alkenyl.

In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one wherein R₄ is -OH (i.e., the C-

terminal end is $-C(O)OH$). In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one wherein R_4 is $-NR_{17}R_{18}$, R_{17} and R_{18} meaning the same. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the N-terminal end
 5 corresponds to $-NH_2$. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the C-terminal and N-terminal ends of the peptide of the invention are, respectively, $-C(O)OH$ and $-NH_2$. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the C-terminal and N-terminal ends of the peptide of the invention are, respectively, $-C(O)NH_2$ and $-NH_2$.

10

In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the C-terminal and N-terminal ends of the peptide of the invention are, respectively, $-C(O)OH$ and $-NHR_5$, wherein R_5 is selected from (C_1-C_{20}) alkyl and (C_1-C_{10}) alkyl substituted by one or more radicals selected from the group consisting of: halogen, (C_1-C_{10}) alkyl, $-OR_6$, $-NR_7R_8$, $-SR_9$, $-$
 15 SOR_{10} , $-SO_2R_{11}$, and $-CO_2R_{12}$.


In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the C-terminal and N-terminal ends of the peptide of the invention are, respectively, $-C(O)R_4$ and $-NH_2$, wherein R_4 means $-NHR_{18}$, and R_{18} is a radical selected from the group
 20 consisting of: (C_1-C_{10}) alkyl and (C_1-C_{10}) alkyl substituted by one or more radicals selected from the group consisting of: halogen, (C_1-C_{10}) alkyl, $-OR_6$, $-NR_7R_8$, $-SR_9$, $-SOR_{10}$, $-SO_2R_{11}$, and $-CO_2R_{12}$.

In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the C-terminal and N-terminal ends of the peptide of the invention are,
 25 respectively, $-C(O)R_4$ and $-NH_2$, wherein R_4 means $-NR_{17}R_{18}$, and R_{17} and R_{18} are radicals independently selected from the group consisting of: (C_1-C_{10}) alkyl and (C_1-C_{10}) alkyl substituted by one or more radicals selected from the group consisting of: halogen, (C_1-C_{10}) alkyl, $-OR_6$, $-NR_7R_8$, $-SR_9$, $-SOR_{10}$, $-SO_2R_{11}$, and $-CO_2R_{12}$.


30 In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the linker biradical of formula (I) is between an alpha carbon atom of an amino acid located at position "i" in the peptide sequence of sequence SEQ ID NO: 1 and an alpha carbon atom of an amino acid located at position "i+7" in the peptide sequence SEQ ID NO: 1.

35 In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is selected from the group consisting of amino acid sequences SEQ ID NO: 3 to 8:


SEQ ID NO: 3




5 SEQ ID NO: 4




SEQ ID NO: 5




SEQ ID NO: 6



SEQ ID NO: 7



SEQ ID NO: 8




wherein "m", "n", "p", "q", "L" and "R₁₉" are as defined in any of the previous embodiments.


In another embodiment of the first aspect of the invention, optionally in combination with any of the
 30 embodiments provided above or below, the linker biradical of formula (I) is between an alpha carbon atom of an amino acid located at position "i" in the peptide sequence of SEQ ID NO: 2 and an alpha carbon atom of an amino acid located at position "i+7" in the peptide sequence of SEQ ID NO: 2.

In another embodiment of the first aspect of the invention, optionally in combination with any of the
 35 embodiments provided above or below, the peptide or salt thereof is selected from the group consisting of amino acid sequences SEQ ID NO: 9 to 16:


SEQ ID NO: 9




SEQ ID NO: 10




SEQ ID NO: 11




SEQ ID NO: 12




SEQ ID NO: 13




SEQ ID NO: 14



SEQ ID NO: 15




SEQ ID NO: 16




wherein "m_x", "n_x", "p_x", "q_x", "L" and "R₁₉" are as defined in any of the previous embodiments.

- In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one with at least 85% of identity with a sequence SEQ ID NO: 3 to 8, wherein R_1 , R_3 and R_{19} are the same or different and represent (C_1-C_{10}) alkyl; and R_2 is (C_2-C_{10}) alkenyl. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one with at least 85% of identity with a sequence SEQ ID NO: 3 to 8, wherein R_1 , R_3 and R_{19} are the same or different and represent (C_1-C_{10}) alkyl; R_2 is (C_2-C_{10}) alkenyl; the C-terminal is selected from $-C(O)OH$; and $-CONH_2$ and the N-terminal end is $-NH_2$.
- 10 In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one with at least 85% of identity with a sequence SEQ ID NO: 9 to 16, wherein R_1 , R_3 and R_{19} are the same or different and represent (C_1-C_{10}) alkyl; and R_2 is (C_2-C_{10}) alkenyl. In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide or salt thereof is one with at least 85% of identity with a sequence SEQ ID NO: 9 to 16, wherein R_1 , R_3 and R_{19} are the same or different and represent (C_1-C_{10}) alkyl; R_2 is (C_2-C_{10}) alkenyl; the C-terminal is selected from $-C(O)OH$ and $-CONH_2$ and the N-terminal end is $-NH_2$.
- 20 In another embodiment of the first aspect of the invention, the peptide is one having at least 85% of identity with an amino acid sequence, or pharmaceutical or cosmeceutical salt thereof, selected from the group consisting of sequence SEQ ID NO: 17 to 23:


SEQ ID NO: 17 (hereinafter also referred "F22")




SEQ ID NO: 18 (hereinafter also referred "F29")




SEQ ID NO: 19 (hereinafter also referred "F31")




SEQ ID NO: 20 (hereinafter also referred "F05")




SEQ ID NO: 21 (hereinafter also referred "F13")



SEQ ID NO: 22 (hereinafter also referred "F16")



SEQ ID NO: 23 (hereinafter also referred "F18")



In another embodiment of the first aspect of the invention, optionally in combination with any of the
 20 embodiments provided above or below, the peptide or salt thereof is a peptide which has an identity of 85%,
 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% with respect to any
 of the sequences SEQ ID NO: 1 to 23, and "m", "n", "p", "q", "m_x", "n_x", "p_x", "q_x", L and R₁₉ are as defined in
 any of the above embodiments. In another embodiment of the first aspect of the invention, optionally in
 combination with any of the embodiments provided above or below, the peptide or salt thereof is a peptide
 25 which has an identity of 100% with respect to any of the sequences SEQ ID NO: 1 to 23, and "m", "n", "p", "q",
 "m_x", "n_x", "p_x", "q_x", L and R₁₉ are as defined in any of the above embodiments. In one embodiment, optionally
 in combination with any of the embodiments provided above or below, the two aminoacidic residues
 connected by the linker are not considered when performing the alignment to determine the identical positions
 for confirming the identity degree.

30

In one embodiment of the first aspect of the invention, optionally in combination with any of the embodiments
 provided above or below, the peptide of the first aspect is conjugated to a moiety, such as a label, a drug or a
 cell penetration peptide. In another embodiment of the first aspect of the invention, optionally in combination
 with any of the embodiments provided above or below, the peptide is conjugated to a label. In another
 35 embodiment of the first aspect of the invention, optionally in combination with any of the embodiments

provided above or below, the label is conjugated to the N- or C-terminal of the peptide.

A "label" as used herein is a molecule or compound that can be detected by a variety of methods including fluorescence, electrical conductivity, radioactivity, size, and the like. The label may be intrinsically capable of emitting a signal, such as for example fluorescent label that emits light of a particular wavelength following excitation by light of another lower, characteristic wavelength. Alternatively, the label may not be capable of intrinsically emitting a signal but it may be capable of being bound by another compound that does emit a signal. An example of this latter situation is a label such as biotin which itself does not emit a signal but which when bound to labeled avidin or streptavidin molecules can be detected. Other examples of this latter kind of label are ligands that bind specifically to particular receptors. Detectably labeled receptors are allowed to bind to ligand labeled unit specific markers in order to visualize such markers.

Labels that may be used according to the invention include but are not limited to electron spin resonance molecule, a fluorescent molecule, a chemiluminescent molecule, a radioisotope, an enzyme substrate, an enzyme, a biotin molecule, an avidin molecule, an electrical charge transferring molecule, a semiconductor nanocrystal, a semiconductor nanoparticle, a colloid gold nanocrystal, a ligand, a microbead, a magnetic bead, a paramagnetic molecule, a quantum dot, a chromogenic substrate, an affinity molecule, a protein, a peptide, nucleic acid, a carbohydrate, a hapten, an antigen, an antibody, an antibody fragment, and a lipid. Radioisotopes can be detected with film or charge coupled devices (CCDs), ligands can be detected by binding of a receptor having a fluorescent, chemiluminescent or enzyme tag, and microbeads can be detected using electron or atomic force microscopy.

The conjugation of the label to the peptide can be performed following routine protocols well-known for the skilled in the art.

In another embodiment, the peptide of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, is conjugated to a drug. In one embodiment, the drug is conjugated to the N-terminal end of the peptide.

30


In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the peptide is conjugated to a cell penetrating peptide.

In present invention the term "cell penetrating peptide" ("CPP") refers to short peptides that facilitate cellular uptake of various molecular cargo (from nanosize particles to small chemical molecules and large fragments of DNA). The "cargo" is associated to peptides via the C(t) or N(t)-end, either through chemical linkage via covalent bonds or through non-covalent interactions. The function of the CPPs are to deliver the cargo into cells, a process that commonly occurs through endocytosis. Current use is limited by a lack of cell specificity in CPP-mediated cargo delivery and insufficient understanding of the modes of their uptake. CPPs typically

have an amino acid composition that either contains a high relative abundance of positively charged amino acids such as lysine or arginine or has sequences that contain an alternating pattern of polar/charged amino acids and non-polar, hydrophobic amino acids. These two types of structures are referred to as polycationic or amphipathic, respectively. A third class of CPPs are the hydrophobic peptides, containing only apolar residues, with low net charge or have hydrophobic amino acid groups that are crucial for cellular uptake. The conjugation of the CPP to the peptide provided in the present invention can be performed following well-known routine protocols, such as solid phase synthesis or solution selective capping. (cf. Copolovici D. M. et al., "Cell-Penetrating Peptides: Design, Synthesis, and Applications", 2014, ACS Nano, 2014, 8 (3), pp 1972–1994). In another embodiment of the first aspect of the invention, optionally in combination with any of the embodiments provided above or below, the cell penetrating peptide is a polycationic CPP, polyArg or, alternatively, penetratine.

The process for the preparation of the peptides according to the first aspect of the invention comprises:


(1.a) the coupling, by condensation, of the corresponding amino acids of the peptide with a compound of formula (III) and a compound of formula (IV), which correspond to the amino acids referred as "i" and "i+4" or "i+7". Compounds (III) and (IV) will be those suffering a subsequent cyclization step to generate the "L" biradical":



wherein R_{19} is as defined above, Z_1 and Z_2 are the same or different and represent $(\text{C}_2\text{-C}_{10})$ alkenyl; and

(1.b) a cyclization step comprising the ring-closed metathesis of Z_1 and Z_2 (cf. Kim Young-Woo et al., "Synthesis of all-hydrocarbon stapled α -helical peptides by ring-closing olefin metathesis", Nature Protocols, 2011, 6(6), p. 761-771; Scott J. M. et al., "Application of Ring-Closing Metathesis to the Synthesis of Rigidified Amino Acids and Peptides", J. Am. Chem. Soc., 1996, v.118 (40), pp 9606–9614) performed in solution with a Grubbs (I or II generation) catalyst; or, alternatively,

(2a) the coupling, by condensation, of the required amino acids, including a compound of formula (V) and a compound of formula (VI), which correspond to the amino acids referred as "i" and "i+4" or "i+7". Compounds (V) and (VI) will be those suffering a subsequent cyclization step to generate the "L" biradical":



wherein R_{19} is as defined above, Z_3 and Z_4 are the same or different, selected from the group consisting of: halogen –SH, –NHR₂₀, –OH, $(\text{C}_2\text{-C}_{10})$ alkyl-SH, $(\text{C}_1\text{-C}_{10})$ alkyl-OH, $(\text{C}_1\text{-C}_{10})$ alkyl-NHR₂₁, $\text{C}(=\text{O})\text{OH}$, $(\text{C}_1\text{-C}_{10})$ alkyl-

$C_{10})C(=O)OH$, $C(=O)NHR_{22}$, $(C_1-C_{10})alkylC(=O)NHR_{23}$, OR_{24} , $C(=O)$ -halogen, $C(=O)-OR_{25}$, $S(=O)$ -halogen, $S(=O)-OR_{26}$, $S(=O)_2R_{27}$ where R_{20} , R_{21} , R_{22} , R_{23} , R_{24} , R_{25} , R_{26} and R_{27} are monoradicals selected from the group consisting of: hydrogen, $(C_1-C_{10})alkyl$, $(C_2-C_{10})alkenyl$, and $(C_2-C_{10})alkynyl$; a known ring system comprising from 3 to 14 carbon atoms, the system comprising from 1 to 3 rings, where:

5

each one of the rings is saturated, partially unsaturated, or aromatic;

the rings are isolated, partially or totally fused,

each one of the members forming the known ring system is selected from the group consisting of: $-CH-$, $-CH_2-$, $-NH-$, $-N-$, $-SH-$, $-S-$, and $-O-$; and

10


the ring system is optionally substituted by one or more radicals independently selected from the group consisting of halogen, $-OH$, $-NH_2$, $-SH$, $C(=O)$ -halogen $(C_1-C_{10})haloalkyl$, and $(C_1-C_{10})alkyl-O-$; and

(2b) a cyclization step comprising the coupling reaction between Z_3 and Z_4 radicals; or, alternatively,

15

(3a) the coupling, by condensation, of the corresponding amino acids of the peptide with a compound of formula (VII) and a compound of formula (VIII), which correspond to the amino acids referred as "i" and "i+4" or "i+7". Compounds (VII) and (VIII) will be those suffering a subsequent cyclization step to generate the "L" biradical":

20



wherein R_{19} is as defined above, one of Z_5 and Z_6 is $(C_2-C_{10})alkynyl$ and the other is $(C_2-C_{10})alkylN_3$; and

25

(3.b) a cyclization step comprising the condensation of Z_5 and Z_6 radicals by well-known protocols such as the Cu(I)-mediated Huisgen 1,3-dipolar cycloaddition reaction (a.k.a. a "click" reaction) to generate a 1,4-substituted 1,2,3-triazole bridge (cf. Kolb H.C. et al., "The growing impact of click chemistry on drug discovery", 2003, Drug Discov Today, 8(24):1128-1137).

30

The process for the preparation of the peptide according to the second aspect of the invention comprises the coupling, by condensation, of the carboxylic group or C-terminus of one amino acid with the amino group or N-terminus of another, this coupling reaction being repeated the number of times required until the desired peptide is obtained.

35

The compounds of formula (III), (IV), (V), (VI), (VII), and (VIII) are commercially available and are coupled by condensation to the already formed portion of peptide sequence. These compounds can carry beads for the appropriate solid phase synthesis of the peptide, as well as protecting groups of the carboxy, amino or side-chain. Illustrative non-limitative examples of compounds are: 2-(2'-propenyl)alanine, 2-(3'-butenyl)glycine, 2-

(4'-pentenyl)alanine, 2-(6'-heptenyl) alanine, 2-(7'-octenyl)alanine, allyl-glycine, 5-azido-norvaline, and alpha-propargyl-alanine, among others.

The "coupling" step can be performed in solid phase, following the protocol "deprotection-wash-coupling-wash", by condensation of the carboxylic group of one amino acid with the amino group of another amino acid residue, using amino acids as defined above as well as alpha-alpha di-substituted amino acids of formula (II) to (VII).

The general principle of solid phase peptide synthesis is to repeat cycles of deprotection-wash-coupling-wash. The free N-terminal amine of a solid-phase attached peptide is coupled to a single N-protected amino acid unit. This unit is then deprotected, revealing a new N-terminal amine to which a further amino acid may be attached. Amino acids have reactive moieties at the N- and C-termini, which facilitates amino acid coupling during synthesis. Many amino acids also have reactive side chain functional groups, which can interact with free termini or other side chain groups during synthesis and peptide elongation and negatively influence yield and purity. To facilitate proper amino acid synthesis with minimal side chain reactivity, chemical groups have been developed to bind to specific amino acid functional groups and block, or protect, the functional group from nonspecific reactions. These protecting groups, while vast in nature, can be separated into three groups, as follows: N-terminal protecting groups, C-terminal protecting groups (mostly used in liquid-phase synthesis), and side chain protecting groups.

20

For coupling the peptides the carboxyl group is usually activated. This is important for speeding up the reaction. There are two main types of activating groups: carbodiimides and triazolols. However, the use of pentafluorophenyl esters (FDPP, PFPOH) and BOP-Cl are useful for cyclising peptides.

25 Purified, individual amino acids are reacted with these protecting groups prior to synthesis and then selectively removed during specific steps of peptide synthesis.

Exemplary resins which may be employed by the present invention include, but are not limited to: (1) alkenyl resins (e.g., REM resin, vinyl sulfone polymer-bound resin, vinyl-polystyrene resin); (2) amine functionalized resins (e.g., amidine resin, N-(4-Benzyloxybenzyl)hydroxylamine polymer bound, (aminomethyl)polystyrene, polymer bound (R)-(+)-a-methylbenzylamine, 2-Chlorotrityl Knorr resin, 2-N-Fmoc-Amino-dibenzocyclohepta-1,4-diene, polymer-bound resin, 4-[4-(1-Fmoc-aminoethyl)-2-methoxy-5-nitrophenoxy]butyramidomethyl-polystyrene resin, 4-Benzyloxybenzylamine, polymer-bound, 4-Carboxybenzenesulfonamide, polymer-bound, Bis(tert-butoxycarbonyl)thiopseudourea, polymer-bound, Dimethylaminomethyl-polystyrene, Fmoc-3-amino-3-(2-nitrophenyl)propionic acid, polymer-bound, N-Methyl aminomethylated polystyrene, PAL resin, Sieber amide resin, tert-Butyl N-(2-mercaptoethyl)carbamate, polymer-bound, Triphenylchloromethane-4-carboxamide polymer bound); (3) benzhydrylamine (BHA) resins (e.g., 2-Chlorobenzhydryl chloride, polymer-bound, HMPB-benzhydrylamine polymer bound, 4-Methylbenzhydryl, polymer-bound, Benzhydryl chloride, polymer-bound, Benzhydrylamine polymer-bound);

- (4) Br-functionalized resins (e.g., 4-(Benzyloxy)benzyl bromide polymer bound, 4-Bromopolystyrene, Brominated PPOA resin, Brominated Wang resin, Bromoacetal, polymer-bound, Bromopolystyrene, HypoGel® 200 Br, Polystyrene A-Br for peptide synthesis, Selenium bromide, polymer-bound, TentaGel HL-Br, TentaGel MB-Br, TentaGel S-Br); (5) Chloromethyl resins (e.g., 5-[4-(Chloromethyl)phenyl]pentyl]styrene, polymer-bound, 4-(Benzyloxy)benzyl chloride polymer bound, 4-Methoxybenzhydryl chloride, polymer-bound); (6) CHO-functionalized resins (e.g., (4-Formyl-3-methoxyphenoxy)methyl)polystyrene, (4-Formyl-3-methoxyphenoxy)methyl)polystyrene, 3-Benzyloxybenzaldehyde, polymer-bound, 4-Benzyloxy-2,6-dimethoxybenzaldehyde, polymer-bound, Formylpolystyrene, HypoGel® 200 CHO, Indole resin, Polystyrene A—CH(OEt)₂, TentaGel HL—CH(OEt)₂); (7) Cl-functionalized resins (e.g., Benzoyl chloride polymer bound, (chloromethyl)polystyrene, Merrifield's resin); (8) CO₂H functionalized resins (e.g., Carboxyethylpolystyrene, HypoGel® 200 COOH, Polystyrene AM—COOH, TentaGel HL—COOH, TentaGel MB—COOH, TentaGel S-COOH); (9) Hypo-Gel resins (e.g., HypoGel® 200 FMP, HypoGel® 200 PHB, HypoGel® 200 Trt-OH, HypoGel® 200 HMB); (10) I-functionalized resins (e.g., 4-Iodophenol, polymer-bound, Iodopolystyrene); Janda-Jels™ (JandaJelα-Rink amide, JandaJel-NH₂, JandaJel—Cl, JandaJel-4-Mercaptophenol, JandaJel-OH, JandaJel-1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide, JandaJel-1,3,4,6,7,8-hexahydro-2H-pyrimido-[1,2-a]pyrimidine, JandaJel-morpholine, JandaJel-polypyridine, JandaJel-Triphenylphosphine, JandaJel-Wang); (11) MBHA resins (3[4'-(Hydroxymethyl)phenoxy]propionic acid-4-methylbenzhydrylamine resin, 4-(Hydroxymethyl)phenoxyacetic acid polymer-bound to MBHA resin, HMBA-4-methylbenzhydrylamine polymer bound, 4-Methylbenzhydrylamine hydrochloride polymer bound Capacity (amine)); (12) NH₂ functionalized resins ((Aminomethyl)polystyrene, (Aminomethyl)polystyrene, HypoGel® 200 NH₂, Polystyrene AM-NH₂, Polystyrene Microspheres 2-aminoethylated, Polystyrol Microspheres 2-bromoethylated, Polystyrol Microspheres 2-hydroxyethylated, TentaGel HL-NH₂, Tentagel M Br, Tentagel M NH₂, Tentagel M OH, TentaGel MB-NH₂, TentaGel S-NH₂, TentaGel S-NH₂); (13) OH-functionalized resins (e.g., 4-hydroxymethylbenzoic acid, polymer-bound, Hydroxymethyl Resins, OH-functionalized Wang Resins); (14) oxime resins (e.g., 4—Chlorobenzophenone oxime polymer bound, Benzophenone oxime polymer bound, 4-Methoxybenzophenone oxime polymer bound); (15) PEG resins (e.g., ethylene glycol polymer bound); (16) Boc/Blz peptide synthesis resins (e.g., Boc-Lys(Boc)-Lys[Boc-Lys(Boc)]-Cys(Acm)-b-Ala-O-PAM resin, Boc-Lys(Fmoc)-Lys[Boc-Lys(Fmoc)]-b-Ala-O-Pam resin, Boc-Lys(Boc)-Lys[Boc-Lys(Boc)]-Lys[Boc-Lys(Boc)-Lys[Boc-Lys(Boc)]]-b-Ala-O-PAM resin, Boc-Lys(Fmoc)-Lys[Boc-Lys(Fmoc)]-Lys[Boc-Lys(Fmoc)-Lys[Boc-Lys(Fmoc)]]-b-Ala-O-PAM resin, Boc-Lys(Boc)-Lys[Boc-Lys(Boc)]-Lys[Boc-Lys(Boc)-Lys[Boc-Lys(Boc)]]—Cys(Acm)-b-Ala-O-PAM resin, Preloaded PAM resins); (17) Fmoc-/t-Bu peptide synthesis resins (e.g., Fmoc-Lys(Fmoc)-Lys[Fmoc-Lys(Fmoc)]-b-Ala-O-Wang resin, Fmoc-Lys(Fmoc)-Lys[Fmoc-Lys(Fmoc)]-Lys[Fmoc-Lys(Fmoc)-Lys[Fmoc-Lys(Fmoc)]]-b-Ala-O-Wang resin, Preloaded TentaGel® S Trityl Resins, Preloaded TentaGel® Resins, Preloaded Trityl Resins, Preloaded Wang Resins, Trityl Resins Preloaded with Amino Alcohols); (19) thiol-functionalized resins (e.g., HypoGel® 200 S-Trt, Polystyrene AM-S-Trityl, TentaGel HL-S-Trityl, TentaGel MB-S-Trityl, TentaGel S-S-Trityl); and (20) Wang resins (e.g., Fmoc-Ala-Wang resin, Fmoc-Arg(Pbf)-Wang resin, Fmoc-Arg(Pmc)-Wang resin, Fmoc-Asn(Trt)-Wang resin, Fmoc-Asp(OtBu)-Wang resin, Fmoc-Cys(Acm)-Wang resin, Fmoc-Cys(StBu)-Wang resin, Fmoc-Cys(Trt) Wang resin, Fmoc-Gln(Trt)-Wang resin, Fmoc-Glu(OtBu)-Wang resin, Fmoc-Gly-Wang resin, Fmoc-His(Trt)-Wang resin, Fmoc-Ile-Wang resin,

Fmoc-Leu-Wang resin, Fmoc-Lys(Boc)-Wang resin, Fmoc-Met-Wang resin, Fmoc-D-Met-Wang resin, Fmoc-Phe-Wang resin, Fmoc-Pro-Wang resin, Fmoc-Ser(tBu)-Wang resin, Fmoc-Ser(Trt)-Wang resin, Fmoc-Thr(tBu)-Wang resin, Fmoc-Trp(Boc) Wang resin, Fmoc-Trp-Wang resin, Fmoc-Tyr(tBu)-Wang resin, Fmoc-Val-Wang resin).

5

"Protecting group" (PG) refers to a grouping of atoms that when attached to a reactive group in a molecule masks, reduces or prevents that reactivity.

Suitable amino-protecting groups include methyl carbamate, ethyl carbamate, 9-fluorenylmethyl carbamate (Fmoc), 9-(2-sulfo)fluorenylmethyl carbamate, 9-(2,7-dibromo)fluorenylmethyl carbamate, 2,7-di-t-butyl-[9-(10,10-dioxo-10,10,10,10-tetrahydrothioxanthyl)]methyl carbamate (DBD-Tmoc), 4-methoxyphenacyl carbamate (Phenoc), 2,2,2-trichloroethyl carbamate (Troc), 2-trimethylsilylethyl carbamate (Teoc), 2-phenylethyl carbamate (hZ), 1-(1-adamantyl)-1-methylethyl carbamate (Adpoc), 1,1-dimethyl-2-haloethyl carbamate, 1,1-dimethyl-2,2-dibromoethyl carbamate (DB-t-BOC), 1,1-dimethyl-2,2,2-trichloroethyl carbamate (TCBOC), 1-methyl-1-(4-biphenyl)ethyl carbamate (Bpoc), 1-(3,5-di-t-butylphenyl)-1-methylethyl carbamate (t-Bumeoc), 2-(2'- and 4'-pyridyl)ethyl carbamate (Pyoc), 2-(N,N-dicyclohexylcarboxamido)ethyl carbamate, t-butyl carbamate (BOC), 1-adamantyl carbamate (Adoc), vinyl carbamate (Voc), allyl carbamate (Alloc), 1-isopropylallyl carbamate (Ipaoc), cinnamyl carbamate (Coc), 4-nitrocinnamyl carbamate (Noc), 8-quinolyl carbamate, N-hydroxypiperidyl carbamate, alkylthio carbamate, benzyl carbamate (Cbz), p-methoxybenzyl carbamate (Moz), p-nitobenzyl carbamate, p-bromobenzyl carbamate, p-chlorobenzyl carbamate, 2,4-dichlorobenzyl carbamate, 4-methylsulfinylbenzyl carbamate (MsZ), 9-anthrylmethyl carbamate, diphenylmethyl carbamate, 2-methylthioethyl carbamate, 2-methylsulfonylethyl carbamate, 2-(p-toluenesulfonyl)ethyl carbamate, [2-(1,3-dithianyl)]methyl carbamate (Dmoc), 4-methylthiophenyl carbamate (Mtpc), 2,4-dimethylthiophenyl carbamate (Bmpc), 2-phosphonioethyl carbamate (Peoc), 2-triphenylphosphonioisopropyl carbamate (Ppoc), 1,1-dimethyl-2-cyanoethyl carbamate, m-chloro-p-acyloxybenzyl carbamate, p-(dihydroxyboryl)benzyl carbamate, 5-benzisoxazolymethyl carbamate, 2-(trifluoromethyl)-6-chromonylmethyl carbamate (Troc), m-nitrophenyl carbamate, 3,5-dimethoxybenzyl carbamate, o-nitrobenzyl carbamate, 3,4-dimethoxy-6-nitrobenzyl carbamate, phenyl(o-nitrophenyl)methyl carbamate, phenothiazinyl-(10)-carbonyl derivative, N'-p-toluenesulfonylaminocarbonyl derivative, N'-phenylaminothiocabonyl derivative, t-amyl carbamate, S-benzyl thiocarbamate, p-cyanobenzyl carbamate, cyclobutyl carbamate, cyclohexyl carbamate, cyclopentyl carbamate, cyclopropylmethyl carbamate, p-decyloxybenzyl carbamate, 2,2-dimethoxycarbonylvinyl carbamate, o-(N,N-dimethylcarboxamido)benzyl carbamate, 1,1-dimethyl-3-(N,N-dimethylcarboxamido)propyl carbamate, 1,1-dimethylpropynyl carbamate, di(2-pyridyl)methyl carbamate, 2-furanylmethyl carbamate, 2-iodoethyl carbamate, isobornyl carbamate, isobutyl carbamate, isonicotinyl carbamate, p-(p'-methoxyphenylazo)benzyl carbamate, 1-methylcyclobutyl carbamate, 1-methylcyclohexyl carbamate, 1-methyl-1-cyclopropylmethyl carbamate, 1-methyl-1-(3,5-dimethoxyphenyl)ethyl carbamate, 1-methyl-1-(p-phenylazophenyl)ethyl carbamate, 1-methyl-1-phenylethyl carbamate, 1-methyl-1-(4-pyridyl)ethyl carbamate, phenyl carbamate, p-(phenylazo)benzyl carbamate, 2,4,6-tri-t-butylphenyl carbamate, 4-(trimethylammonium)benzyl carbamate, 2,4,6-trimethylbenzyl carbamate,

- formamide, acetamide, chloroacetamide, trichloroacetamide, trifluoroacetamide, phenylacetamide, 3-phenylpropanamide, picolinamide, 3-pyridylcarboxamide, N-benzoylphenylalanyl derivative, benzamide, p-phenylbenzamide, o-nitrophenylacetamide, o-nitrophenoxycetamide, acetoacetamide, (N'-dithiobenzyloxycarbonylamino)acetamide, 3-(p-hydroxyphenyl)propanamide, 3-(o-nitrophenyl)propanamide,
- 5 2-methyl-2-(o-nitrophenoxy)propanamide, 2-methyl-2-(o-phenylazophenoxy)propanamide, 4-chlorobutanamide, 3-methyl-3-nitrobutanamide, o-nitrocinnamide, N-acetylmethionine derivative, o-nitrobenzamide, o-(benzyloxymethyl)benzamide, 4,5-diphenyl-3-oxazolin-2-one, N-phthalimide, N-dithiasuccinimide (Dts), N-2,3-diphenylmaleimide, N-2,5-dimethylpyrrole, N-1,1,4,4-tetramethyldisilylazacyclopentane adduct (STABASE), 5-substituted 1,3-dimethyl-1,3,5-triazacyclohexan-2-
- 10 one, 5-substituted 1,3-dibenzyl-1,3,5-triazacyclohexan-2-one, 1-substituted 3,5-dinitro-4-pyridone, N-methylamine, N-allylamine, N-[2-trimethylsilyl]ethoxy]methylamine (SEM), N-3-acetoxypyrrolamine, N-(1-isopropyl-4-nitro-2-oxo-3-pyrroline-3-yl)amine, quaternary ammonium salts, N-benzylamine, N-di(4-methoxyphenyl)methylamine, N-5-dibenzosuberylamine, Ntriphenylmethylamine (Tr), N-[(4-methoxyphenyl)diphenylmethyl]amine (MMTr), N-9-phenylfluorenylamine (PhF), N-2,7-dichloro-9-
- 15 fluorenylmethyleneamine, N-ferrocenylmethylamino (Fcm), N-2-picolylamino N'-oxide, N-1,1-dimethylthiomethyleneamine, N-benzylideneamine, N-p-methoxybenzylideneamine, N-diphenylmethyleneamine, N-[(2-pyridyl)mesityl]methyleneamine, N-(N',N'-dimethylaminomethylene)amine, N,N'-isopropylidenediamine, N-p-nitrobenzylideneamine, N-salicylideneamine, N-5-chlorosalicylideneamine, N-(5-chloro-2-hydroxyphenyl)phenylmethyleneamine, N-cyclohexylideneamine, N-(5,5-dimethyl-3-oxo-1-
- 20 cyclohexenyl)amine, N-borane derivative, N-diphenylborinic acid derivative, N-[phenyl(pentacarbonylchromium- or tungsten)carbonyl]amine, N-copper chelate, N-zinc chelate, N-nitroamine, N-nitrosoamine, amine N-oxide, diphenylphosphinamide (Dpp), dimethylthiophosphinamide (Mpt), diphenylthiophosphinamide (Ppt), dialkyl phosphoramidates, dibenzyl phosphoramidate, diphenyl phosphoramidate, benzenesulfenamide, o-nitrobenzenesulfenamide (Nps), 2,4-dinitrobenzenesulfenamide,
- 25 pentachlorobenzenesulfenamide, 2-nitro-4-methoxybenzenesulfenamide, triphenylmethylsulfenamide, 3-nitropyridinesulfenamide (Npys), p-toluenesulfonamide (Ts), benzenesulfonamide, 2,3,6-trimethyl-4-methoxybenzenesulfonamide (Mtr), 2,4,6-trimethoxybenzenesulfonamide (Mtb), 2,6-dimethyl-4-methoxybenzenesulfonamide (Pme), 2,3,5,6-tetramethyl-4-methoxybenzenesulfonamide (Mte), 4-methoxybenzenesulfonamide (Mbs), 2,4,6-trimethylbenzenesulfonamide (Mts), 2,6-dimethoxy-4-
- 30 methylbenzenesulfonamide (iMds), 2,2,5,7,8-pentamethylchroman-6-sulfonamide (Pmc), methanesulfonamide (Ms), β -trimethylsilylethanesulfonamide (SES), 9-anthracenesulfonamide, 4-(4',8'-dimethoxynaphthylmethyl)benzenesulfonamide (DNMBS), benzylsulfonamide, trifluoromethylsulfonamide, and phenacylsulfonamide.
- 35 Examples of suitably protected carboxylic acids further include, but are not limited to, silyl-, alkyl-, alkenyl-, aryl-, and arylalkyl-protected carboxylic acids. Examples of suitable silyl groups include trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl, triisopropylsilyl, and the like. Examples of suitable alkyl groups include methyl, benzyl, p-methoxybenzyl, 3,4-dimethoxybenzyl, trityl, t-butyl, tetrahydropyran-2-yl. Examples of suitable alkenyl groups include allyl. Examples of suitable aryl groups include optionally

substituted phenyl, biphenyl, or naphthyl. Examples of suitable arylalkyl groups include optionally substituted benzyl (e.g., p-methoxybenzyl (MPM), 3,4-dimethoxybenzyl, O-nitrobenzyl, p-nitrobenzyl, p-halobenzyl, 2,6-dichlorobenzyl, p-cyanobenzyl), and 2- and 4-picoly.

5 In second aspect the present invention provides a pharmaceutical or cosmetical composition.

The expression "therapeutically effective amount" as used herein, refers to the amount of a compound that, when administered passes to blood stream and is sufficient to prevent development of, or alleviate to some extent, one or more of the symptoms of the disease which is addressed. The particular dose of the peptide
 10 administered according to this invention will of course be determined by the particular circumstances surrounding the case, including the compound administered, the route of administration, the particular condition being treated, and the similar considerations. In one embodiment of the second aspect of the invention, the pharmaceutical composition is a parenteral composition. In another embodiment, the pharmaceutical composition is for oral administration.

15 The expression "cosmetically effective amount", in the context of the present invention, refers to the amount of a compound that when it is applied (for example topically) is retained within the skin tissue (such as at subcutaneous level), without significantly reaching, or not reaching, the blood stream, thus exerting a local effect within skin structure. The particular dose of the peptide administered according to this invention will of
 20 course be determined by the particular circumstances surrounding the case. In one embodiment of the invention, the "cosmetically effective amount" of the peptide of the invention is higher, equal or lower than the amount needed to achieve a therapeutic effect when it is administered by parenteral route, particularly by intravenous route.

25 The term "pharmaceutically acceptable" refers to that excipients or carriers suitable for use in the pharmaceutical technology for preparing compositions with medical use.

The term "cosmetically acceptable" which is herein used interchangeably refers to that excipients or carriers suitable for use in contact with human skin without undue toxicity, incompatibility, instability, allergic response,
 30 among others.

The expression "excipients and/or carriers" refers to acceptable materials, compositions or vehicles. Each component must be pharmaceutically or cosmetically acceptable in the sense of being compatible with the other ingredients of the composition. It must also be suitable for use in contact with the tissue or organ of
 35 humans and non-human animals without excessive toxicity, irritation, allergic response, immunogenicity or other problems or complications commensurate with a reasonable benefit/risk ratio. Examples of suitable acceptable excipients are solvents, dispersion media, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like. Except insofar as any conventional excipient medium is incompatible with a

substance or its derivatives, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutical or cosmeceutical composition, its use is contemplated to be within the scope of this invention.

- 5 The formulations of the pharmaceutical and cosmeceutical compositions described herein may be prepared by any method known or hereafter developed in the art of pharmacology and cosmeceutics. In general, such preparatory methods include the step of bringing the active ingredient (the peptide) into association with a excipient and/or one or more other accessory ingredients, and then, if necessary and/or desirable, shaping and/or packaging the product into a desired single- or multi-dose unit.

10

A pharmaceutical composition of the invention may be prepared, packaged, and/or sold in bulk, as a single unit dose, and/or as a plurality of single unit doses. As used herein, a "unit dose" is discrete amount of the pharmaceutical composition comprising a predetermined amount of the active ingredient.

- 15 The relative amounts of the active ingredient (i.e., the peptide as defined in any of the previous aspects and embodiments), the acceptable excipients, and/or any additional ingredients in the composition of the invention will vary, depending upon the identity, size, and/or condition of the subject treated and further depending upon the route by which the composition is to be administered.

- 20 Acceptable excipients used in the manufacture of these compositions include, but are not limited to, inert diluents, dispersing and/or granulating agents, surface active agents and/or emulsifiers, disintegrating agents, binding agents, preservatives, buffering agents, lubricating agents, and/or oils. Such excipients may optionally be included in the inventive formulations. Excipients such as cocoa butter and suppository waxes, coloring agents, coating agents, sweetening, flavoring, and perfuming agents can be present in the composition,
25 according to the judgment of the formulator.

- Exemplary diluents include, but are not limited to, calcium carbonate, sodium carbonate, calcium phosphate, dicalcium phosphate, calcium sulfate, calcium hydrogen phosphate, sodium phosphate lactose, sucrose, cellulose, microcrystalline cellulose, kaolin, mannitol, sorbitol, inositol, sodium chloride, dry starch, cornstarch,
30 powdered sugar, and combinations thereof.

- Exemplary granulating and/or dispersing agents include, but are not limited to, potato starch, corn starch, tapioca starch, sodium starch glycolate, clays, alginic acid, guar gum, citrus pulp, agar, bentonite, cellulose and wood products, natural sponge, cation-exchange resins, calcium carbonate, silicates, sodium carbonate,
35 cross-linked polyvinylpyrrolidone (crospovidone), sodium carboxymethyl starch (sodium starch glycolate), carboxymethyl cellulose, cross-linked sodium carboxymethyl cellulose (croscarmellose), methylcellulose, pregelatinized starch (starch 1500), microcrystalline starch, water insoluble starch, calcium carboxymethyl cellulose, magnesium aluminum silicate (Veegum), sodium lauryl sulfate, quaternary ammonium compounds, and combinations thereof.

Exemplary surface active agents and/or emulsifiers include, but are not limited to, natural emulsifiers (e.g. acacia, agar, alginic acid, sodium alginate, tragacanth, chondrux, cholesterol, xanthan, pectin, gelatin, egg yolk, casein, wool fat, cholesterol, wax, and lecithin), colloidal clays (e.g. bentonite [aluminum silicate] and

5 Veegum [magnesium aluminum silicate]), long chain amino acid derivatives, high molecular weight alcohols (e.g. stearyl alcohol, cetyl alcohol, oleyl alcohol, triacetin monostearate, ethylene glycol distearate, glyceryl monostearate, and propylene glycol monostearate, polyvinyl alcohol), carbomers (e.g., carboxy polymethylene, polyacrylic acid, acrylic acid polymer, and carboxyvinyl polymer), carrageenan, cellulosic derivatives (e.g., carboxymethylcellulose sodium, powdered cellulose, hydroxymethyl cellulose, hydroxypropyl

10 cellulose, hydroxypropyl methylcellulose, methylcellulose), sorbitan fatty acid esters {e.g., polyoxyethylene sorbitan monolaurate [Tween 20], polyoxyethylene sorbitan [Tween 60], polyoxyethylene sorbitan monooleate [Tween 80], sorbitan monopalmitate [Span 40], sorbitan monostearate [Span 60], sorbitan tristearate [Span 65], glyceryl monooleate, sorbitan monooleate [Span 80]}, polyoxyethylene esters (e.g., polyoxyethylene monostearate [Myrj 45], polyoxyethylene hydrogenated castor oil, polyethoxylated castor oil,

15 polyoxymethylene stearate, and Solutol), sucrose fatty acid esters, polyethylene glycol fatty acid esters (e.g., Cremophor), polyoxyethylene ethers, (e.g., polyoxyethylene lauryl ether [Brij 30]), poly(vinyl-pyrrolidone), diethylene glycol monolaurate, triethanolamine oleate, sodium oleate, potassium oleate, ethyl oleate, oleic acid, ethyl laurate, sodium lauryl sulfate, Pluronic F 68, Poloxamer 188, cetrimonium bromide, cetylpyridinium chloride, benzalkonium chloride, docusate sodium, etc. and/or combinations thereof.

20

Exemplary binding agents include, but are not limited to, starch (e.g., cornstarch and starch paste); gelatin; sugars (e.g., sucrose, glucose, dextrose, dextrin, molasses, lactose, lactitol, mannitol); natural and synthetic gums (e.g., acacia, sodium alginate, extract of Irish moss, panwar gum, ghatti gum, mucilage of isapol husks, carboxymethylcellulose, methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropyl cellulose,

25 hydroxypropyl methylcellulose, microcrystalline cellulose, cellulose acetate, polyvinylpyrrolidone), magnesium aluminum silicate (Veegum), and larch arabogalactan); alginates; polyethylene oxide; polyethylene glycol; inorganic calcium salts; silicic acid; polymethacrylates; waxes; water; alcohol; and combinations thereof.

Exemplary preservatives may include antioxidants, chelating agents, antimicrobial preservatives, antifungal

30 preservatives, alcohol preservatives, acidic preservatives, and other preservatives. Exemplary antioxidants include, but are not limited to, alpha tocopherol, ascorbic acid, ascorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, monothioglycerol, potassium metabisulfite, propionic acid, propyl gallate, sodium ascorbate, sodium bisulfite, sodium metabisulfite, and sodium sulfite. Exemplary chelating agents include ethylenediaminetetraacetic acid (EDTA), citric acid monohydrate, disodium edetate, dipotassium edetate,

35 edetic acid, fumaric acid, malic acid, phosphoric acid, sodium edetate, tartaric acid, and trisodium edetate. Exemplary antimicrobial preservatives include, but are not limited to, benzalkonium chloride, benzethonium chloride, benzyl alcohol, bronopol, cetrimide, cetylpyridinium chloride, chlorhexidine, chlorobutanol, chlorocresol, chloroxylenol, cresol, ethyl alcohol, glycerin, hexetidine, imidurea, phenol, phenoxyethanol, phenylethyl alcohol, phenylmercuric nitrate, propylene glycol, and thimerosal. Exemplary antifungal

preservatives include, but are not limited to, butyl paraben, methyl paraben, ethyl paraben, propyl paraben, benzoic acid, hydroxybenzoic acid, potassium benzoate, potassium sorbate, sodium benzoate, sodium propionate, and sorbic acid. Exemplary alcohol preservatives include, but are not limited to, ethanol, polyethylene glycol, phenol, phenolic compounds, bisphenol, chlorobutanol, hydroxybenzoate, and

5 phenylethyl alcohol. Exemplary acidic preservatives include, but are not limited to, vitamin A, vitamin C, vitamin E, beta-carotene, citric acid, acetic acid, dehydroacetic acid, ascorbic acid, sorbic acid, and phytic acid. Other preservatives include, but are not limited to, tocopherol, tocopherol acetate, deteroxime mesylate, cetrimide, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), ethylenediamine, sodium lauryl sulfate (SLS), sodium lauryl ether sulfate (SLES), sodium bisulfite, sodium metabisulfite, potassium sulfite,

10 potassium metabisulfite, Glydant Plus, Phenonip, methylparaben, Germall 115, Germaben II, Neolone, Kathon, and Euxyl. In certain embodiments, the preservative is an anti-oxidant. In other embodiments, the preservative is a chelating agent.

Exemplary buffering agents include, but are not limited to, citrate buffer solutions, acetate buffer solutions,

15 phosphate buffer solutions, ammonium chloride, calcium carbonate, calcium chloride, calcium citrate, calcium gluconate, calcium gluceptate, calcium gluconate, D-gluconic acid, calcium glycerophosphate, calcium lactate, propanoic acid, calcium levulinate, pentanoic acid, dibasic calcium phosphate, phosphoric acid, tribasic calcium phosphate, calcium hydroxide phosphate, potassium acetate, potassium chloride, potassium gluconate, potassium mixtures, dibasic potassium phosphate, monobasic potassium phosphate, potassium

20 phosphate mixtures, sodium acetate, sodium bicarbonate, sodium chloride, sodium citrate, sodium lactate, dibasic sodium phosphate, monobasic sodium phosphate, sodium phosphate mixtures, tromethamine, magnesium hydroxide, aluminum hydroxide, alginic acid, pyrogen-free water, isotonic saline, Ringer's solution, ethyl alcohol, and combinations thereof.

25 Exemplary lubricating agents include, but are not limited to, magnesium stearate, calcium stearate, stearic acid, silica, talc, malt, glyceryl behenate, hydrogenated vegetable oils, polyethylene glycol, sodium benzoate, sodium acetate, sodium chloride, leucine, magnesium lauryl sulfate, sodium lauryl sulfate, and combinations thereof.

30 Exemplary oils include, but are not limited to, almond, apricot kernel, avocado, babassu, bergamot, black current seed, borage, cade, camomile, canola, caraway, carnauba, castor, cinnamon, cocoa butter, coconut, cod liver, coffee, corn, cotton seed, emu, eucalyptus, evening primrose, fish, flaxseed, geraniol, gourd, grape seed, hazel nut, hyssop, isopropyl myristate, jojoba, kukui nut, lavandin, lavender, lemon, litsea cubeba, macademia nut, mallow, mango seed, meadowfoam seed, mink, nutmeg, olive, orange, orange roughy, palm,

35 palm kernel, peach kernel, peanut, poppy seed, pumpkin seed, rapeseed, rice bran, rosemary, safflower, sandalwood, sasquana, savoury, sea buckthorn, sesame, shea butter, silicone, soybean, sunflower, tea tree, thistle, tsubaki, vetiver, walnut, and wheat germ oils. Exemplary oils include, but are not limited to, butyl stearate, caprylic triglyceride, capric triglyceride, cyclomethicone, diethyl sebacate, dimethicone 360, isopropyl myristate, mineral oil, octyldodecanol, oleyl alcohol, silicone oil, and combinations thereof.

Liquid dosage forms for parenteral administration include, but are not limited to, pharmaceutically acceptable liposomes emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredients, the liquid dosage forms may comprise inert diluents commonly used in the art such as, for
 5 example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. In certain embodiments for parenteral administration, the conjugates of the invention are mixed with
 10 solubilizing agents such as polyethoxylated castor oil (e.g. CREMOPHOR™), alcohols, oils, modified oils, glycols, polysorbates, cyclodextrins, polymers, and combinations thereof.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile
 15 injectable preparation may be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as
 20 oleic acid are used in the preparation of injectables. Alternatively, the preparation can be in the form of liposomes.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in
 25 sterile water or other sterile injectable medium prior to use.

In order to prolong the effect of a drug, it is often desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends
 30 upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

The peptides of the invention can be in micro-encapsulated form with one or more excipients as noted above.
 35 In one embodiment, the peptides of the invention are formulated in liposomes.

It will be appreciated that peptides and pharmaceutical compositions of the present invention can be employed in combination therapies. The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or

procedures and the desired therapeutic effect to be achieved. It will be appreciated that the therapies employed may achieve a desired effect for the same purpose (for example, an inventive conjugate useful for detecting tumors may be administered concurrently with another agent useful for detecting tumors), or they may achieve different effects (e.g., control of any adverse effects).

5

The pharmaceutical composition of the present invention may be administered either alone or in combination with one or more other therapeutic agents. By "in combination with," it is not intended to imply that the agents must be administered at the same time and/or formulated for delivery together, although these methods of delivery are within the scope of the invention. The compositions can be administered concurrently with, prior
10 to, or subsequent to, one or more other desired therapeutics or medical procedures. In general, each agent will be administered at a dose and/or on a time schedule determined for that agent. Additionally, the invention encompasses the delivery of the peptide or pharmaceutical compositions in combination with agents that may improve their bioavailability, reduce and/or modify their metabolism, inhibit their excretion, and/or modify their distribution within the body.

15

The particular combination of therapies to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and/or the desired therapeutic effect to be achieved. It will be appreciated that the therapies employed may achieve a desired effect for the same disorder (for example, a peptide of the invention may be administered concurrently with another biologically active agent used to treat
20 the same disorder), and/or they may achieve different effects (e.g., control of any adverse effects). It will further be appreciated that biologically active agents utilized in this combination may be administered together in a single composition or administered separately in different compositions.

The expression "in combination with" also encompasses the possibility of conjugating (by chemical-physical
25 interactions) the peptide of the invention to any of the further agents mentioned above and below, which can be either a therapeutic agent or an agent for improving the profile of the peptide (such as bioavailability), among others.

In one embodiment, the peptides of the invention are administered in combination with one or more anti-
30 cancer agents. An anti-cancer agent may be, for instance, methotrexate, vincristine, adriamycin, cisplatin, non-sugar containing chloroethylnitrosoureas, 5-fluorouracil, mitomycin C, bleomycin, doxorubicin, dacarbazine, taxol, fragyline, Meglamine GLA, valrubicin, carmustaine and poliferposan, MMI270, BAY 12-9566, RAS farnesyl transferase inhibitor, farnesyl transferase inhibitor, MMP, MTA/LY231514, LY264618/Lometexol, Glamolec, CI-994, TNP-470, Hycamtin/Topotecan, PKC412, Valspodar/PSC833,
35 Novantrone/Mitroxantrone, Metaret/Suramin, Batimastat, E7070, BCH-4556, CS-682, 9-AC, AG3340, AG3433, Incel/VX-710, VX-853, ZD0101, ISI641, ODN 698, TA 2516/Marmistat, BB2516/Marmistat, CDP 845, D2163, PD183805, DX895 if, Lemonal DP 2202, FK 317, Picibanil/OK-432, AD 32/Valrubicin, Metastron/strontium derivative, Temodal/Temozolomide, Evacet/liposomal doxorubicin, Yewtaxan/Paclitaxel, Taxol/Paclitaxel, Xeload/Capecitabine, Furtulon/Doxifluridine, Cyclopax/oral paclitaxel, Oral Taxoid, SPU-

077/Cisplatin, HMR 1275/Flavopiridol, CP-358 (774)/EGFR, CP-609 (754)/RAS oncogene inhibitor, BMS-182751/oral platinum, UFT (Tegafur/Uracil), Ergamisol/Levamisole, Eniluracil/776C85/5FU enhancer, Campto/Levamisole, Camptosar/Irinotecan, Tumodex/Ralitrexed, Leustatin/Cladribine, Paxex/Paclitaxel, Doxil/liposomal doxorubicin, Caelyx/liposomal doxorubicin, Fludara/Fludarabine, Pharmarubicin/Epirubicin,

5 DepoCyt, ZD1839, LU 79553/Bis-Naphtalimide, LU 103793/Dolastain, Caetyx/liposomal doxorubicin, Gemzar/Gemcitabine, ZD 0473/Anormed, YM 116, iodine seeds, CDK4 and CDK2 inhibitors, PARP inhibitors, D4809/Dexifosamide, Ifes/Mesnex/Ifosamide, Vumon/Teniposide, Paraplatin/Carboplatin, Plantinol/cisplatin, Vepeside/Etoposide, ZD 9331, Taxotere/Docetaxel, prodrug of guanine arabinoside, Taxane Analog, nitrosoureas, alkylating agents such as melphelan and cyclophosphamide, Aminoglutethimide, Asparaginase,

10 Busulfan, Carboplatin, Chlorombucil, Cytarabine HCl, Dactinomycin, Daunorubicin HCl, Estramustine phosphate sodium, Etoposide (VP16-213), Floxuridine, Fluorouracil (5-FU), Flutamide, Hydroxyurea (hydroxycarbamide), Ifosfamide, Interferon Alfa-2a, Alfa-2b, Leuprolide acetate (LHRH-releasing factor analogue), Lomustine (CCNU), Mechlorethamine HCl (nitrogen mustard), Mercaptopurine, Mesna, Mitotane (o.p.DDD), Mitoxantrone HCl, Octreotide, Plicamycin, Procarbazine HCl, Streptozocin, Tamoxifen citrate,

15 Thioguanine, Thiotepa, Vinblastine sulfate, Amsacrine (m-AMSA), Azacitidine, Erthropoietin, Hexamethylmelamine (HMM), Interleukin 2, Mitoguanzone (methyl-GAG; methyl glyoxal bis-guanyldrazone; MGBG), Pentostatin (2'-deoxycoformycin), Semustine (methyl-CCNU), Teniposide (VM-26) or Vindesine sulfate, signal transduction inhibitors (such as MEK, BRAF, AKT, her2, mTOR, and PI3K inhibitors), but it is not so limited.

20

As it is illustrated below, the peptides of the first aspect of the invention are useful in the therapeutic treatment of melanoma, particularly metastatic melanoma by reducing the proliferation of malignant melanocytes. In one embodiment, optionally in combination with any of the embodiments provided above or below, when the peptide or pharmaceutical composition is used with a therapeutic purpose, it is orally or parenterally

25 administered.

The present invention also provides the non-therapeutic use of the peptide or cosmetical salt thereof for regulating melanogenesis, particularly for reducing or increasing the production of melanin. All the embodiments provided above regarding the peptide and cosmetical composition are also embodiments of this

30 aspect. In one embodiment, optionally in combination with any of the embodiments provided above or below, the peptide is for use in the treatment of hyperpigmentation (by reducing the amount of melanin) or hypopigmentation (by increasing the amount of melanin). In another embodiment of the non-therapeutic use, optionally in combination with any of the embodiments provided above or below, it is regulated the melanogenesis in skin or hair, particularly in skin. In another embodiment, optionally in combination with any

35 of the embodiments provided above or below, the subject does not suffer skin cancer. In one embodiment, optionally in combination with any of the embodiments provided above or below, when the peptide or cosmetical composition is used with a non-therapeutic/cosmetic purpose, it is topically administered.

The present invention also provides a cosmetical method for the treatment of the hyper or hypopigmentation

comprising the application of the peptide or cosmetical salt thereof as defined in the first aspect or the cosmetical composition as defined in the second aspect of the invention. This cosmetic method does not encompass the therapeutic method. All the embodiments provided above regarding the peptide and cosmetic composition are also embodiments of this aspect. In one embodiment of the cosmetical method, optionally in combination with any of the embodiments provided above or below, the peptide or cosmetical composition is applied on skin and/or hair. In one embodiment of the cosmetical method, optionally in combination with any of the embodiments provided above or below, the peptide or cosmetical composition is applied on skin.

The present invention also provides a pigmenting composition, such as a topical cosmetical or pharmaceutical pigmenting composition, comprising the peptide or salt thereof as defined in any of embodiments provided above.

The invention also provides a depigmenting composition, such as a topical cosmetical or pharmaceutical depigmenting composition, comprising the peptide or salt thereof as defined in any of the embodiments provided above.

Throughout the description and claims the word "comprise" and variations of the word, are not intended to exclude other technical features, additives, components, or steps. Furthermore, the word "comprise" encompasses the case of "consisting of". Additional objects, advantages and features of the invention will become apparent to those skilled in the art upon examination of the description or may be learned by practice of the invention. The following examples are provided by way of illustration, and they are not intended to be limiting of the present invention. Furthermore, the present invention covers all possible combinations of particular and preferred embodiments described herein.

Examples

1. MATERIALS AND METHODS

Synthetic general procedure

30

Compounds IDP-F

Materials were purchased as following: Fmoc-protected α -amino acids (---); Rinkamide MBHA Resin (Tianjin Nankai HECHENG S&T Co., Ltd); HBTU ((2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate), GL Biochem); N-methyl morpholine (Sinopharm Chemical Reagent Co., Ltd.); Succinic anhydride (Aladdin); acetonitrile (Xingke Chemical); ninhydrin (Sinopharm Chemical Reagent Co., Ltd.); Piperidine (Vertellus); Dimethylformamide, DMF (Zhejiang jiangshan chemical co., Ltd); trifluoroacetic acid, TFA (Trifluoroacetic acid, Solvay), TIS (Thioanisole, Solvay)

Briefly, the linear polypeptides were synthesized manually using Fmoc based SPPS (solid phase peptides synthesis) on Rink amide MBHA resin as support.

The following protocol was used:

- 5 1. The Fmoc protective group was removed with 20% piperidine in DMF.
2. The resin was washed with DMF five times.
3. The subsequent Fmoc-protected amino acid was coupled for 45 min using Fmoc-AA (3 equiv.), HBTU (3 equiv.), and N-methyl morpholine (6 equiv.).
- 10 4. The resin was washed with DMF five times. Coupling was checked by ninhydrin test.
5. Repeat from step 1.
6. N-terminal was capped by reacting with succinic anhydride (10 equiv.) and N-methyl morpholine (10 equiv.).
- 15 The peptide was cleaved from the resin and deprotected by exposure to solution F (95% TFA, 2.5% water, 2.5% TIS) and lyophilized.

The lyophilized peptides were purified by reverse phase HPLC using a C18 column (see compounds characterization for details). The peptides were identified by LC-MS-ESI. All the mass spectral data for all the
20 compounds are shown below in Table 1.

Materials were purchased as following: Fmoc-protected α -amino acids (include than the olefinic amino acids Fmoc-[(S)-2-(4 pentenyl)alanine]OH, Fmoc-[(R)-2-(4 pentenyl)alanine]OH, Fmoc-[(S)-2-(7 octenyl)alanine]OH, Fmoc-[(R)-2-(4 pentenyl)alanine]OH, 2-(6-chloro-1-H-benzotriazole-1-yl)-1,1,3,3-tetramethylaminium
25 hexafluorophosphate (TBTU), resins, dimethylformamide (DMF), N,N-diisopropylethylamine (DIEA), trifluoroacetic acid (TFA), 1,2-dichloroethane (DCE), Grubbs Ru(IV) catalyst and piperidine were purchased from different suppliers.

Briefly, the linear polypeptides were synthesized with automatic synthesizer using Fmoc solid phase peptide
30 chemistry. Only the coupling with olefinic amino acids was performed manually after removing the resins from the reactor vessel, as disclosed in the previous section.

The ring-closing metathesis reaction was performed in solution with a first generation Grubbs catalyst after cleaving the linear peptide from the resin, as disclosed by Scott J. M. and colleagues (Scott J.M. et al.,
35 "Application of Ring-Closing Metathesis to the Synthesis of Rigidified Amino Acids and Peptides", 1996, J. Am. Chem. Soc., 1996, 118 (40), pp 9606–9614).

The deprotected peptide precipitated with methyl-tert-butyl ether at 4° C. and lyophilized.

The lyophilized peptides were purified by reverse phase HPLC using a C18 column (see compounds

characterization for details). The peptides were identified by LC-MS-ESI. All the mass spectral data for all the compounds are shown below in Table 1.

HPLC conditions:

5

IDP-F05. The compound was purified by HPLC-RP (C-18 column; Pump A: H₂O with 0,1% TFA; Pump B Acetonitrile with 0,1% TFA) using a linear gradient 30%-40% of B in 20 minutes (R. T. = 10,2). Purity grade 95,06% by HPLC;

10 IDP-F13. The compound was purified by HPLC-RP (C-18 column; Pump A: H₂O with 0,1% TFA; Pump B Acetonitrile with 0,1% TFA) using a linear gradient 32%-42% of B in 20 minutes (R. T. = 14,26). Purity grade 97,5% by HPLC;

IDP-F16. The compound was purified by HPLC-RP (C-18 column; Pump A: H₂O with 0,1% TFA; Pump B Acetonitrile with 0,1% TFA) using a linear gradient 35%-45% of B in 20 minutes (R. T. = 7.30). Purity grade 95,7% by HPLC;

15 IDP-F18. The compound was purified by HPLC-RP (C-18 column; Pump A: H₂O with 0,1% TFA; Pump B Acetonitrile with 0,1% TFA) using a linear gradient 51%-61% of B in 20 minutes (R. T. = 10.97-11.97). Purity grade 95,6% by HPLC;

IDP-F22. The compound was purified by HPLC-RP (C-18 column; Pump A: H₂O with 0,1% TFA; Pump B Acetonitrile with 0,1% TFA) using a linear gradient 53%-63% of B in 20 minutes (R. T. = 10.1-11.36). Purity grade 97,06% by HPLC;

IDP-F29. The compound was purified by HPLC-RP (C-18 column; Pump A: H₂O with 0,1% TFA; Pump B Acetonitrile with 0,1% TFA) using a linear gradient 30%-40% of B in 20 minutes (R. T. = 8.91). Purity grade ≤50% by HPLC;

25 IDP-F31. The compound was purified by HPLC-RP (C-18 column; Pump A: H₂O with 0,1% TFA; Pump B Acetonitrile with 0,1% TFA) using a linear gradient 38%-48% of B in 20 minutes (R. T. = 10,65). Purity grade 95,49% by HPLC;


TABLE 1: mass characterization

N. ID	Sequence		MW (1H)	Mass (2H)	Mass (3H)	Mass (4H)	Mass (5H)
F05	RRRRFNIXDRIKELXTL	calcul.	2277.51				
		found		1139,7			
F13	RRXRFNINDXIKELGTL	calcul.	2136.52				
		found		1068,6	712,8	534,5	
F16	RRFXINDRIKXLGTLP	calcul.	2103.58				
		found		1052,2	701,9	526,9	
F18	RRFNIXDRIKELXTLP	calcul.	2175.65				
		found		1088,2	725,9		

F22	WNXGTILKAXVDYIRK	calcul.	1968.37				
		found		984,7	656,9	493	
F29	KGTIXKASVDYXRKLQ	calcul.	1898.27				
		found		949,5	633,5		
F31	KGTILKAXVDYIRKXQ	calcul.	1924.36				
		found		962,5	642,2	481,9	

The X biradical represents, the compound of formula:

5



and a L biradical corresponding to $-(CH_2)_3-CH=CH-(CH_2)_3-$ or to $-(CH_2)_6-CH=CH-(CH_2)_3-$ links both Xs radicals
10 in each sequence.

Cell lines:

SK-MEL-28, skin, human malignant melanoma, ATCC-HTB-72™.

15 IGR-37, skin, human melanoma derived from metastatic site: groin lymph node.

IGR-39, skin, human melanoma derived from metastatic site: groin lymph node.

B16-F10, skin, mouse melanoma, ATCC-CRL-6475™.

501-MEL, skin, human malignant melanoma.

20 Cell culture

Cell lines SK-MEL-28, IGR-37, IGR-39, B16-F10, 501-MEL and A-375 were cultured in incubator at 37°C in Dulbecco's modified Eagle's medium (Invitrogen Life Technologies, Foster City, CA) containing 10% fetal bovine serum inactivated (FBS, Dutscher, France), penicillin (50 U/ml), and streptomycin sulphate (50 mg/ml)

25 (Gibco BRL, Germany). During the amplification step and the assays adherent cells were rinsed with DPBS (Dulbecco's Phosphate Buffered Saline, Sigma D1283) three times and afterward treated for 5 minutes with trypsin ([0,5 g / ml] / EDTA [0.2 g/ml]) (Gibco-BRL, 15400054) in solution of DPBS at 37 °C, and, once detached, transferred in the culturing medium. Cells were counted in a Neubauer chamber after labelling with Tripzan-Blue. Each assay was performed only when the viability was superior to 90%.

30

Viability assay

Cell lines SK-MEL-28, IGR-37, B16-F10, 501-MEL and A-375 were seeded at a density of 1,000 cells/well in

96 well plates. After 24h, the compounds to be tested were added to calculate the efficacy at the concentration of 10 μ M with serial dilutions. Controls are the untreated cells. Each experiment was performed in triplicate.

- 5 Cells were incubated during 72h in incubator under CO₂ atmosphere at 37°C. Cell viability was measured by colorimetric MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide) assay. MTT was incubated for a further 4 h, and the medium was then discarded. DMSO was added to dissolve the formazan product, and absorbance at 570 nm was measured with a microplate reader (MULTISKAN, Labsystems). Cell viability percentages were calculated by dividing the absorbance value of cells treated with a given compound by the
10 absorbance of untreated cells.

Statistics

- Data analysis was performed calculating the percentage of cell viability normalized vs. the values of negative
15 control, which was considered equal to 100%. The dose/response curve was fitted through the sigmoidal equation dose-response (variable slope) and the EC₅₀ values were calculated as follow:

$$Y = \text{Bottom} + (\text{Top} - \text{Bottom}) / (1 + 10^{\{[(\text{LogEC}_{50} - X) * \text{HillSlope}]\}}),$$

- 20 where: X is compound concentration (log scale) and Y is the response

Calculations and graphs were conducted using GraphPad Prism (Prism 6 for Windows).

2. RESULTS

25

2.1 Efficacy results

Tables 2a, 2b and 2c provide the anti-proliferative activity in several melanoma cell lines at 10 μ M.

Table 2a

30

Compounds	SK-MEL-28	501-MEL
F05	60	90
F13	40	40
F16	50	50
F18	70	70
F22	100	90
F29	30	30
F31	60	70

35

Table 2b

Compounds	IGR-37
F13	40
F16	50
F18	60
F22	80
F31	30

Table 2c

Compounds	B16-F10
F05	60
F18	40
F22	40
F31	50

5

All the peptides of the invention show a remarkable anticancer activity against the tested melanoma cell lines, even of aggressive phenotypes.

2.2 Measurement of melanin content in B16-F10 melanoma cells

10

B16-F10 melanoma cells were seeded into 25 cm² cell culture flask and incubated for 24 h. Then, 0.1 μ M α -MSH was added and cells were treated with the aforementioned peptides at doses lower than their respective IC₅₀ for a total of 48 h incubation. After washing twice with phosphate buffered saline, cells were dissolved in 1mL of 1N NaOH. For measurement of melanin content, 100 μ L aliquots of solution were then placed in 96-
15 well plates and the absorbance was measured at 450nm using microplate reader. Anti-melanin activity was expressed by the percentage of melanin to that of untreated melanoma cells. As positive control Kojic Acid (KA) was used at different concentrations (0.1, 1 and 10 μ M).

The results are summarized in FIG. 1. As it can be seen, the peptides of the invention are involved in
20 melanogenesis pathway. Thus, some of the peptides stimulate the production of melanin, which can be an indicia of melanocyte activation, whereas others, such as F22 and F29 drastically reduces the amount of melanin. What is even more remarkably is the fact that the peptides of the invention are more efficient in regulating melanocytes when compared with other compounds. In this regard, it was found that KA was used at a concentration of 1 μ M the reduction was substantially lower than the reduction achieved when the peptide
25 of the invention F22 was used at the same concentration.

Therefore, this data not only supports the effect of the peptides of the invention on melanocyte activity, but also illustrates that they are even more efficient than other compounds available in the state of the art currently used with the same aim.

5 2.3 Determinations of depigmenting activity in zebrafish embryos.

Pigment formation during zebrafish development starts at about 24hpf by the retinal pigment epithelium (RPE) and then progresses towards the dorsolateral skin, where melanocytes are distributed. Biochemical melanin quantification performed in untreated embryos to compare to embryos treated with compound F22. F22 was dissolved in 1X PBS to stock solution. To proceed to zebrafish embryos drug incubation, F22 was serially diluted in E3 medium to obtain: 2.5 μ M in a final volume of 1 ml.

Melanin Quantification Assay

50 embryos per triplicate were collected and dissolved in lysis buffer at 95°C for 30 minutes. 200 μ L of the solution was placed in a 96 well plate and total melanin content was quantified using spectrometer at 490 nm. Results were compared with synthetic melanin. Standard curve by plotting the average Blank-corrected 490 nm measurement for each Melanin standard vs its concentration in μ g/mL. Standard curve was used to determine the melanin protein concentration of each unknown sample.

20 Imaging analysis:

Embryos were placed in a robotic microfluidic system VAST (Union Biometrica), which allows the automatic aspiration, placement and rotation of the larvae under the microscope (Leica, DM6-B). Once embryos were positioned under the right angle (dorsal view), images are acquired with the VAST onboard camera. One dorsal image was acquired per larva. FIJI program was used for the quantification. RGB Images were converted to 8-bit and the orientation of the larvae was determined. A fixed threshold was applied for generating binary masks from the images. A specific region of interest ROI was selected depending on the orientation of the larva. The ROIs were generated such that the eyes would be excluded from the quantification. Segmented areas inside the ROIs were then measured using the Analyse Pixels command. The Level of pigmentation was provided as the sum of all pixels in those areas.

30

Results: Melanin Quantification

Table 1. Melanin quantification by biochemical analysis. The table shows the quantification of melanin as percentage compared to the negative control (E3 medium) at 72 hrs of incubation. Asterisks indicate statistical significance after a One-way ANOVA: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

	<u>Melanine concentration (% to control)</u>
<u>Untreated</u>	<u>100</u>
<u>F22 2.5 uM</u>	<u>61.41 **</u>

Table2. Melanin quantification by imaging analysis. Melanin content is calculated as the mean of the sum of pixels related to the body of each zebrafish larvae (E3 medium) at 96 hpf. Asterisks indicate statistical significance after a One-way ANOVA: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

	<u>Pigmented area (pixels)</u>
<u>Untreated</u>	<u>11200,00</u>
<u>F22 2.5 uM</u>	<u>6500,00 ***</u>

From both quantifications used, it can be concluded that there are evidences of depigmenting effect of compound F22 at 2.5 μ M when analyzing the biochemical melanin amount and, comparing with the analysis of the imaging study.

10

Citation List

Altschul et al., "Basic local alignment search tool", 1990, J. Mol. Biol, v. 215, pages 403-410;

15 Copolovici D. M. et al., "Cell-Penetrating Peptides: Design, Synthesis, and Applications", 2014, ACS Nano, 2014, 8 (3), pp 1972–1994;

Higgins et al., "CLUSTAL V: improved software for multiple sequence alignment", 1992, CABIOS, 8(2), pages 189-191;

20

Kim Young-Woo et al., "Synthesis of all-hydrocarbon stapled α -helical peptides by ring-closing olefin metathesis", Nature Protocols, 2011, 6(6), p. 761-771;

Kolb H.C. et al., "The growing impact of click chemistry on drug discovery", 2003, Drug Discov Today, 25 8(24):1128–1137); and

Scott J.M. et al., "Application of Ring-Closing Metathesis to the Synthesis of Rigidified Amino Acids and Peptides", 1996, J. Am. Chem. Soc., 1996, 118 (40), pp 9606–9614;

30 For reasons of completeness, various aspects of the invention are set out in the following numbered clauses:

Clauses

1. A peptide or a pharmaceutical or cosmetical acceptable salt thereof, the peptide having an amino acid sequence with at least 85% of identity with respect sequence SEQ ID NO: 1 or 2:


(Trp)_m-(Asn)_n-Lys-Gly-Thr-Ile-Leu-Lys-Ala-Ser-Val-Asp-Tyr-Ile-Arg-Lys-(Leu)_p-(Gln)_q (SEQ ID NO: 1)

5

(Arg)_{m_x}-(Arg)_{n_x}-Arg-Arg-Phe-Asn-Ile-Asn-Asp-Arg-Ile-Lys-Glu-Leu-Gly-Thr-Leu-(Ile)_{p_x}-(Pro)_{q_x} (SEQ ID NO: 2)

wherein

- 10 "m", "n", "p", and "q" represent integers and are selected from 0 and 1;
 "m_x", "n_x", "p_x", and "q_x" represent integers and are selected from 0 and 1;
 a C-terminal end corresponding to -C(O)R₄;
 a N-terminal end corresponding to -NHR₅;
 R₄ is a radical selected from the group consisting of -OH and -NR₁₇R₁₈;
 15 R₅ is a radical selected from the group consisting of -H, (C₁-C₂₀)alkyl and (C₁-C₁₀)alkyl substituted by one or more radicals selected from the group consisting of: halogen, (C₁-C₁₀)alkyl, -OR₆, -NR₇R₈, -SR₉, -SOR₁₀, -SO₂R₁₁, and -CO₂R₁₂; and
 R₁₇ and R₁₈ are radicals independently selected from the group consisting of: -H, (C₁-C₁₀)alkyl and (C₁-C₁₀)alkyl substituted by one or more radicals selected from the group consisting of: halogen, (C₁-C₁₀)alkyl, -OR₆, -NR₇R₈, -SR₉, -SOR₁₀, -SO₂R₁₁, and -CO₂R₁₂; the peptide optionally including a linker
 20 biradical "L" of formula (I)



- 25 which is connecting an alpha carbon atom of an amino acid located at position "i" with an alpha carbon atom of an amino acid located at position "i+4" or "i+7" in the peptide sequence

wherein

"a" and "b" are the same or different and are 0 or 1;

"c" is comprised from 1 to 10;

30


R₁ and R₃ are biradicals independently selected from the group consisting of: (C₁-C₁₀)alkyl; (C₁-C₁₀)alkyl substituted by one or more radicals selected from the group consisting of: halogen, (C₁-C₁₀)alkyl, -OR₆, -NR₇R₈, -SR₉, -SOR₁₀, -SO₂R₁₁, and -CO₂R₁₂; (C₂-C₁₀)alkenyl; (C₂-C₁₀)alkenyl substituted by one or more radicals selected from the group consisting of: halogen, (C₁-C₁₀)alkyl, -OR₆, -NR₇R₈, -SR₉, -SOR₁₀, -SO₂R₁₁, and -CO₂R₁₂; (C₂-C₁₀)alkynyl; and (C₂-C₁₀)alkynyl substituted by one or more radicals selected from the group consisting of: halogen, (C₁-C₁₀)alkyl, -OR₆, -NR₇R₈, -SR₉, -SOR₁₀, -SO₂R₁₁, and -CO₂R₁₂;
 35

R₂ is a biradical selected from the group consisting of: -O-, C(=O), C(=O)NR₁₃, C(=O)O, S(=O), S(=O)₂, NR₁₄,

(C₁-C₁₀)alkyl, (C₂-C₁₀)alkenyl, (C₂-C₁₀)alkynyl, -NR₁₅-NR₁₆-, -N=N-, -S-S-, and a known ring system comprising from 3 to 14 members, the system comprising from 1 to 3 rings, where:

- each one of the rings is saturated, partially unsaturated, or aromatic;
- 5 the rings are isolated, partially or totally fused,
- each one of the members forming the known ring system is selected from the group consisting of: -CH-, -CH₂-, -NH-, -N-, -SH-, -S-, and -O-; and
- the ring system is optionally substituted by one or more radicals independently selected from the group consisting of halogen, -OH, -NO₂, (C₁-C₁₀)alkyl, (C₁-C₁₀)haloalkyl, and (C₁-C₁₀)alkyl-O-; and
- 10 R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, and R₁₆ are radicals independently selected from the group consisting of: -H and (C₁-C₁₀)alkyl; and

- the amino acids which are connected by the linker being of formula (II)
- 15



- 20 wherein

- R₁₉ is a monoradical selected from the group consisting of: (C₁-C₁₀)alkyl, (C₂-C₁₀)alkenyl, (C₂-C₁₀)alkynyl, and a known ring system comprising from 3 to 14 members, the system comprising from 1 to 3 rings, where:
- each one of the rings is saturated, partially unsaturated, or aromatic;
- 25 the rings are isolated, partially or totally fused,
- each one of the members forming the known ring system is selected from the group consisting of: -CH-, -CH₂-, -NH-, -N-, -SH-, -S-, and -O-.

2. The peptide according to clause 1, which has an identity of 100% with respect to sequence SEQ ID NO: 1 or 2.
- 30

3. The peptide according to any of the previous clauses, wherein the linker biradical of formula (I) is between an alpha carbon atom of an amino acid located at position "i" in the peptide sequence of sequence SEQ ID NO: 1 and an alpha carbon atom of an amino acid located at position "i+7" in the peptide sequence SEQ ID NO: 1; or, alternatively, is between an alpha carbon atom of an amino acid located at position "i" in the peptide sequence of sequence SEQ ID NO: 2 and an alpha carbon atom of an amino acid located at position "i+7" in the peptide sequence SEQ ID NO: 2.
- 35

4. The peptide according to any one of the previous clauses, wherein a=b=c=1.

5. The peptide according to any one of the previous clauses, wherein
 R_1 and R_3 are biradicals independently selected from the group consisting of: (C_1-C_{10}) alkyl; (C_2-C_{10}) alkenyl;
 and (C_2-C_{10}) alkynyl;
- 5 R_2 is a biradical selected from the group consisting of: (C_1-C_{10}) alkyl, (C_2-C_{10}) alkenyl, and (C_2-C_{10}) alkynyl; and
 R_{19} is a monoradical selected from the group consisting of: (C_1-C_{10}) alkyl, (C_2-C_{10}) alkenyl, and (C_2-C_{10}) alkynyl.
6. The peptide according to any one of the previous clauses, wherein R_1 , R_3 and R_{19} are (C_1-C_{10}) alkyl; and R_2
 is (C_2-C_{10}) alkenyl.
- 10
7. The peptide according to any one of the previous clauses, wherein the C-terminal end corresponds
 to $-C(O)OH$ or $-C(O)NH_2$, and the N-terminal end corresponds to $-NH_2$.
8. The peptide according to any one of the previous clauses, which has at least a 85%, at least 90%, at least
 15 95% or it is 100% identical to a peptide sequence SEQ ID NO: 3 to 16, wherein "L", "R19", "m", "n",
 "p", "q", "mx", "nx", "px" and "qx" are as defined in any of the preceding clauses.
9. The peptide according to any one of the previous clauses, which is a peptide with an amino acid having at
 least 85%, at least 90%, at least 95% or is 100% identical to an amino acid sequence selected from
 20 the group consisting of: SEQ ID NO: 17 to 23.
10. A pharmaceutical or cosmetical composition comprising a therapeutically or cosmetically effective amount
 of the peptide as defined in any one of the clauses 1-9, together with acceptable pharmaceutical or
 cosmetical excipients and/or carriers.
- 25
11. A non-therapeutic use of a peptide or cosmetical salt thereof as defined in any one of the clauses 1-9 or a
 cosmetical composition as defined in clause 10 for regulating melanogenesis, particularly for
 reducing or increasing the production of melanin.
- 30 12. The non-therapeutic use of a peptide according to clause 11 for the treatment of hyperpigmentation or
 hypopigmentation.
13. A peptide or pharmaceutical salt thereof as defined in any one of the clauses 1-9 or a pharmaceutical
 composition as defined in clause 10 for use as a medicament.
- 35
14. A peptide or a pharmaceutical salt thereof as defined in any one of the clauses 1-9 or the pharmaceutical
 composition as defined in clause 10, for use in the treatment and/or prevention of cancer, particularly
 a melanoma cancer, such as a metastatic melanoma cancer.

15. A peptide or pharmaceutical salt thereof as defined in any one of the clauses 1-9 or the pharmaceutical composition as defined in clause 10, for use in the treatment and/or prevention of a disease related to the dysregulation of melanogenesis, particularly by inhibiting cancer melanocytes proliferation.

Claims

1. A peptide or a pharmaceutical or cosmetical acceptable salt thereof, the peptide having an amino acid sequence with at least 90% of identity with respect sequence SEQ ID NO: 1 or 2:

5

$(\text{Trp})_m-(\text{Asn})_n-\text{Lys}-\text{Gly}-\text{Thr}-\text{Ile}-\text{Leu}-\text{Lys}-\text{Ala}-\text{Ser}-\text{Val}-\text{Asp}-\text{Tyr}-\text{Ile}-\text{Arg}-\text{Lys}-(\text{Leu})_p-(\text{Gln})_q$ (SEQ ID NO: 1)

$(\text{Arg})_{mx}-(\text{Arg})_{nx}-\text{Arg}-\text{Arg}-\text{Phe}-\text{Asn}-\text{Ile}-\text{Asn}-\text{Asp}-\text{Arg}-\text{Ile}-\text{Lys}-\text{Glu}-\text{Leu}-\text{Gly}-\text{Thr}-\text{Leu}-(\text{Ile})_{px}-(\text{Pro})_{qx}$ (SEQ ID NO: 2)

10 wherein

"m", "n", "p", and "q" represent integers and are selected from 0 and 1;

"m_x", "n_x", "p_x", and "q_x" represent integers and are selected from 0 and 1;

a C-terminal end corresponding to $-\text{C}(\text{O})\text{R}_4$;

15 a N-terminal end corresponding to $-\text{NHR}_5$;


R₄ is a radical selected from the group consisting of $-\text{OH}$ and $-\text{NR}_{17}\text{R}_{18}$;

R₅ is a radical selected from the group consisting of $-\text{H}$, $(\text{C}_1-\text{C}_{20})$ alkyl and $(\text{C}_1-\text{C}_{10})$ alkyl substituted by one or more radicals selected from the group consisting of: halogen, $(\text{C}_1-\text{C}_{10})$ alkyl, $-\text{OR}_6$, $-\text{NR}_7\text{R}_8$, $-\text{SR}_9$, $-\text{SOR}_{10}$, $-\text{SO}_2\text{R}_{11}$, and $-\text{CO}_2\text{R}_{12}$; and

20 R₁₇ and R₁₈ are radicals independently selected from the group consisting of: $-\text{H}$, $(\text{C}_1-\text{C}_{10})$ alkyl and $(\text{C}_1-\text{C}_{10})$ alkyl substituted by one or more radicals selected from the group consisting of: halogen, $(\text{C}_1-\text{C}_{10})$ alkyl, $-\text{OR}_6$, $-\text{NR}_7\text{R}_8$, $-\text{SR}_9$, $-\text{SOR}_{10}$, $-\text{SO}_2\text{R}_{11}$, and $-\text{CO}_2\text{R}_{12}$;

the peptide further including a linker biradical "L" of formula (I)

25



which is connecting an alpha carbon atom of an amino acid located at position "i" with an alpha carbon atom of an amino acid located at position "i+4" or "i+7" in the peptide sequence

30 wherein

"a" and "b" are the same or different and are 0 or 1;

"c" is comprised from 1 to 10;

35

R₁ and R₃ are biradicals independently selected from the group consisting of: $(\text{C}_1-\text{C}_{10})$ alkyl; $(\text{C}_1-\text{C}_{10})$ alkyl substituted by one or more radicals selected from the group consisting of: halogen, $(\text{C}_1-\text{C}_{10})$ alkyl, $-\text{OR}_6$, $-\text{NR}_7\text{R}_8$, $-\text{SR}_9$, $-\text{SOR}_{10}$, $-\text{SO}_2\text{R}_{11}$, and $-\text{CO}_2\text{R}_{12}$; $(\text{C}_2-\text{C}_{10})$ alkenyl; $(\text{C}_2-\text{C}_{10})$ alkenyl substituted by one or more radicals selected from the group consisting of: halogen, $(\text{C}_1-\text{C}_{10})$ alkyl, $-\text{OR}_6$, $-\text{NR}_7\text{R}_8$, $-\text{SR}_9$, $-\text{SOR}_{10}$, $-\text{SO}_2\text{R}_{11}$, and $-\text{CO}_2\text{R}_{12}$; $(\text{C}_2-\text{C}_{10})$ alkynyl; and $(\text{C}_2-\text{C}_{10})$ alkynyl substituted


by one or more radicals selected from the group consisting of: halogen, (C₁-C₁₀)alkyl, -OR₆, -NR₇R₈, -SR₉, -SOR₁₀, -SO₂R₁₁, and -CO₂R₁₂;

5 R₂ is a biradical selected from the group consisting of: -O-, C(=O), C(=O)NR₁₃, C(=O)O, S(=O), S(=O)₂, NR₁₄, (C₁-C₁₀)alkyl, (C₂-C₁₀)alkenyl, (C₂-C₁₀)alkynyl, -NR₁₅-NR₁₆-, -N=N-, -S-S-, and a known ring system comprising from 3 to 14 members, the system comprising from 1 to 3 rings, where:

10 each one of the rings is saturated, partially unsaturated, or aromatic;
the rings are isolated, partially or totally fused,
each one of the members forming the known ring system is selected from the group consisting of: -CH-, -CH₂-, -NH-, -N-, -SH-, -S-, and -O-; and
the ring system is optionally substituted by one or more radicals independently selected from the group consisting of halogen, -OH, -NO₂, (C₁-C₁₀)alkyl, (C₁-C₁₀)haloalkyl, and (C₁-C₁₀)alkyl-
15 O-; and

R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, and R₁₆ are radicals independently selected from the group consisting of: -H and (C₁-C₁₀)alkyl; and

20 the amino acids which are connected by the linker being of formula (II)



25

wherein

R₁₉ is a monoradical selected from the group consisting of: (C₁-C₁₀)alkyl, (C₂-C₁₀)alkenyl, (C₂-C₁₀)alkynyl, and a known ring system comprising from 3 to 14 members, the system comprising from 1 to 3 rings, where:

30 each one of the rings is saturated, partially unsaturated, or aromatic;
the rings are isolated, partially or totally fused,

each one of the members forming the known ring system is selected from the group consisting of: -CH-, -CH₂-, -NH-, -N-, -SH-, -S-, and -O-.

35 2. The peptide according to claim 1, which has an identity of 100% with respect to sequence SEQ ID NO: 1 or 2.

3. The peptide according to any of the previous claims, wherein the linker biradical of formula (I) is between

an alpha carbon atom of an amino acid located at position "i" in the peptide sequence of sequence SEQ ID NO: 1 and an alpha carbon atom of an amino acid located at position "i+7" in the peptide sequence SEQ ID NO: 1; or, alternatively, is between an alpha carbon atom of an amino acid located at position "i" in the peptide sequence of sequence SEQ ID NO: 2 and an alpha carbon atom of an amino acid located at position "i+7" in the peptide sequence SEQ ID NO: 2.

4. The peptide according to any one of the previous claims, wherein $a=b=c=1$.

5. The peptide according to any one of the previous claims, wherein

10 R_1 and R_3 are biradicals independently selected from the group consisting of: (C_1-C_{10}) alkyl; (C_2-C_{10}) alkenyl; and (C_2-C_{10}) alkynyl;

R_2 is a biradical selected from the group consisting of: (C_1-C_{10}) alkyl, (C_2-C_{10}) alkenyl, and (C_2-C_{10}) alkynyl; and

R_{19} is a monoradical selected from the group consisting of: (C_1-C_{10}) alkyl, (C_2-C_{10}) alkenyl, and (C_2-C_{10}) alkynyl.

15 6. The peptide according to any one of the previous claims, wherein R_1 , R_3 and R_{19} are (C_1-C_{10}) alkyl; and R_2 is (C_2-C_{10}) alkenyl.

7. The peptide according to any one of the previous claims, wherein the C-terminal end corresponds to $-C(O)OH$ or $-C(O)NH_2$, and the N-terminal end corresponds to $-NH_2$.

20

8. The peptide according to any one of the previous claims, which has at least a 85%, at least 90%, at least 95% or it is 100% identical to a peptide sequence SEQ ID NO: 3 to 16, wherein "L", "R19", "m", "n", "p", "q", "mx", "nx", "px" and "qx" are as defined in any of the preceding claims.

25 9. The peptide according to any one of the previous claims, which is a peptide with an amino acid having at least 85%, at least 90%, at least 95% or is 100% identical to an amino acid sequence selected from the group consisting of: SEQ ID NO: 17 to 23.

10. The peptide according to any one of the previous claims, which is conjugated to a label, drug, a cell
30 penetrating peptide or a PROTAC.

11. A pharmaceutical or cosmetical composition comprising a therapeutically or cosmetically effective amount of a peptide or a pharmaceutical or cosmetical acceptable salt thereof as defined in any one of claims 1-10, together with acceptable pharmaceutical or cosmetical excipients and/or carriers.

35

12. The pharmaceutical or cosmetical composition according to claim 11, which is a topical composition.

13. Use of a peptide or a salt thereof as defined in any one of claims 1-10 or a composition as defined in claim

- 11 for the preparation of a medicament for regulating melanogenesis, particularly for reducing or increasing the production of melanin.
14. The use as claimed in claim 13, wherein the medicament is for the treatment of hyperpigmentation or
5 hypopigmentation.
15. A peptide or pharmaceutical salt thereof as defined in any one of claims 1-10, or a pharmaceutical composition as defined in claim 11, for use as a medicament.
- 10 16. Use of a peptide or a pharmaceutical salt thereof as defined in any one of claims 1-10, or the pharmaceutical composition as defined in claim 11, for the preparation of a medicament for the treatment and/or prevention of cancer, particularly a melanoma cancer, such as a metastatic melanoma cancer.
17. Use of a peptide or pharmaceutical salt thereof as defined in any one of claims 1-10, or the
15 pharmaceutical composition as defined in claim 11, for the preparation of a medicament for the treatment and/or prevention of a disease related to the dysregulation of melanogenesis, particularly by inhibiting cancer melanocytes proliferation.
18. A depigmenting composition comprising a peptide as defined in any one of claims 1-10
20 and one or more cosmetically acceptable excipients.
19. A pigmenting composition comprising a peptide as defined in any one of claims 1-10 and one or more cosmetically acceptable excipients.
- 25 20. The composition as claimed in any one of the claims 18-19, wherein the composition is a topical composition.