



PHOSPHINE LIGAND PROMOTED SYNTHESIS OF TETRAARYLOXYALKANE AND DIARYLCARBONATES

FIELD OF THE INVENTION

This invention is directed to methods for making tetraaryloxymethane and diaryl carbonates which are useful in the preparation of polycarbonate resins by melt transesterification technology.

5 Manufacture of polycarbonate resins by melt transesterification of
diaryl carbonates provides some important environmental advantages over
the standard commercial process which employs direct reaction of carbon
monoxide, phenol, and phosgene. In addition, the melt process produces a
product characterized by very low levels of contaminants which make them
particularly suitable for use in optical applications such as the manufacture of
10 compact discs.

Manufacture of polycarbonates by melt estrification of diaryl carbonate
is not used extensively on a commercial scale because, at the present time,
there is no satisfactory synthesis for the diaryl carbonate. Prior art methods
for the synthesis of the diaryl carbonate includes direct phosgenation of a
15 phenol in an organic medium. Another process involves a multi-step low
conversion rate process using a catalyst such as a titanium catalyst and a third
process involves the direct carbonylation of a phenol requiring a catalyst such
as pladium that is difficult to recycle. US 5, 663, 406 (King, Jr.et.al.) discloses a method of
producing carbonate ester by a metal induced reaction of a hydroxy compound with a sulfur
compoud.

20 The present invention provides an improved method for producing
tetraaryloxyalkanes such as tetraphenoxymethane from phenol and carbon
disulfide in the presence of cuprus oxide and an aliphatic, cycloaliphatic or
aryl phosphine ligand.





SUMMARY OF THE INVENTION


5 Addition of phosphine ligands can produce high yields of tetraphenoxymethane, TPM, and diphenylcarbonate, DPC, in short reaction times and improved reaction rates by a copper-induced condensation of phenol and carbon disulfide, CS₂. Either aromatic or aliphatic substituted phosphine ligands can be employed. The copper(I) oxide utilization is significantly improved in reactions containing phosphine ligands, such that yields exceeding 100% tetraphenoxymethane were observed. Some of the sulfur-containing byproduct is transferred to the phosphine in the form of phosphine sulfide.

10 The invention provides an enhanced method for making tetraaryloxyalkane and diaryl carbonates such as tetraphenoxymethane and diphenyl carbonate from cuprous oxide, phenol, and carbon disulfide. More particularly, the present invention provides a method for making tetraaryloxyalkane and diaryl carbonate comprising contacting at least one hydroxyaromatic compound with carbon disulfide cuprous oxide and from 0.1 to 10 mole percent based on the moles of cuprous oxide of a phosphine ligand. The presence of as little as 1-10 mole percent of an aromatic or aliphatic phosphine significantly increases the reaction rate and yield of tetraphenoxymethane and diphenyl carbonate. The reaction is shown below, where the term ligand indicates an aliphatic, cycloaliphatic or aromatic phosphine.


15



Tetraphenoxy-methane (TPM)



Diphenyl-carbonate (DPC)



Diphenylthio-carbonate (DPTC)


ligand = none or phosphine,
 where phosphine = [triphenylphosphine, triphenylphosphite,
 triphenylphosphine oxide, tricyclohexyl phosphine, tributyl phosphine,
 1,2-bis(diphenylphosphino)ethane (diphos), 1,2-bis(diphenylphosphino)butane]

DESCRIPTION OF THE INVENTION


The instant invention provides a method of producing carbonate ester comprising the step of mixing a hydroxy compound of the formula



- 5 wherein R is selected from the group consisting of substituted and unsubstituted alkyl radicals, and substituted and unsubstituted aryl radicals; a sulfur compound selected from the group consisting of the formulas



and



wherein R' is a divalent aliphatic radical, O or S, and R'' is R or OR; and a promoter comprising at least one metal source that is capable of reacting with said hydroxy compound and said sulphur compound.

5 It has been discovered that the addition of small amounts of aromatic or aliphatic phosphines to the fundamental reaction, shown in Scheme 1, improves the yield of tetraphenoxymethane and diphenyl carbonate, DPC, from cuprous oxide, phenol, and carbon disulfide. Significant improvements in the rate of reaction were observed when phosphine ligands are present
10 during the reaction. The melt reaction between cuprous oxide, phenol, carbon disulfide and low levels of methyl cyanide (CH₃CN), reaction, is compared with a similar reaction performed in the presence of about 10 mole percent triphenylphosphine, TPP, (reaction 131A) in table 1. Once the reaction containing 10 mole percent triphenylphosphine (131A) undergoes an initial
15 lag time of approximately 1.5 hours, the production of tetraphenoxymethane increases dramatically, with the total yield of tetraphenoxymethane, increasing from 5% to 105% within 5 hours. The initial induction period may be due to limited solubility and initial formation of a phosphine-associated cuprous phenoxide complex.

20 Table 1 shows the data for control reactions that are cuprous oxide based with and without added water in addition to the data for the reactions that included various types and molar levels of phosphine-containing ligands. The table lists the auxiliary reagents used, the yield at 3 hours, the total yield at 30 hours, and the distribution of products.

When the reaction is carried out in the presence of water, diphenyl carbonate is generated insitu and becomes the primary product.

A similar reaction that used 1/10th the level of triphenylphosphine (1 mole percent triphenylphosphine; 35A) worked almost as well (Total Yield, Table 1) as the reaction containing 10 mole percent triphenylphosphine (131A). In fact, the reaction containing 1 mole percent triphenylphosphine (35A) exhibited a much reduced lag time. The lag time was reduced to 15 minutes for the reaction using 1 mole percent triphenylphosphine compared to a lag time of 1.5 hours for the 10 mole percent triphenylphosphine rxn. A graph comparing the first 7 hours of both reactions is shown in Figure 2.

A reaction containing 10 mole percent triphenylphosphine with an added water phase (131B) showed an increase in the early rate of reaction, with a 75% yield by 3 hrs and a 95% yield by 4 hrs. The corresponding 1 mole percent triphenylphosphine reaction (35B) also demonstrated an enhanced rate of reaction of > 80% by 3 hours. A graph comparing these reactions is shown in Figure 3.

Aromatic bidentate phosphines as ligands have also been successfully employed to enhance the initial rate of tetraphenoxymethane and diphenyl carbonate production. These ligands have two aromatic phosphine groups linked by an aliphatic bridge. The two bidentate ligands screened were 1,2-bis(diphenylphosphino)ethane (also known as diphos) (reactions 919A & B, 1014A & B) and 1,4-bis(diphenylphosphino)butane (reactions 925A & B). The initial rate of reaction (see 3 hr. yield, Table 1) for these ligands is similar to that of triphenylphosphine.

Aliphatic phosphines as ligands have also been successful in increasing the rate and overall yield of tetraphenoxymethane and diphenyl carbonate from carbon disulfide, phenol and Cu₂O. Reactions utilizing tricyclohexyl

phosphine ((CyHex)₃-P) as the added ligand at both the 1 & 10 mole percent level (1024A & B; 927A & B) demonstrated a much improved reaction rate and yield. Surprisingly, even reaction 1029B, which used only 0.1 mole percent tricyclohexyl phosphine in combination with added water, gave a 3 hr
5 yield of 90%. The first seven hours of the reactions, with added water, that contain all three levels of tricyclohexyl phosphine (0.1, 1, and 10 mole percent) are shown in Figure 4.

The use of an alternate aliphatic phosphine at the 10 mole percent level, tributylphosphine (reactions 1017A & B), increased the initial rate only
10 slightly while depressing the overall yield.

In all of the reactions which contained high levels of phosphine ligands (10 mole percent) and achieved high total yields, significant levels of diphenylthiocarbonate (DPTC; half-product of the reaction) were observed as the reaction progressed. The levels of DPTC ranged from 25-50% by 3 to 7
15 hrs. In comparison, the fundamental Cu₂O-mediated reactions without ligands produce less than 1% of DPTC. In certain ligand-promoted cases, however, the final levels of DPTC decreased to less than 2%. Systems such as these (131A, 131B, 26B, 927B) demonstrate that some ligand-promoted reactions are capable of converting the half-product, DPTC, to
20 tetraphenoxymethane insitu.

Reactions using a 10 mole percent level of aromatic bidentate phosphines and tricyclohexyl phosphine as ligands show the same general trend of significant levels of DPTC being formed early. In all reactions with 10 mole percent ligand, the level of DPTC does not appear to be transient, and
25 once formed, remains at a level of 20-50%.

It has been noted, however, that lowering the level of ligand used in the reaction can have a positive effect on the rate. The phosphine ligand

containing reactions which gave the best yield/fastest rate at the 10 mole percent level were also screened at the reduced level of 1 mole percent ligand. For all three of these ligands, (triphenyl phosphine: 35A & B; DiPhos: 1014A & B, tricyclohexyl phosphine: 1024A & B), less than 1% DPTC was noted
5 throughout the reaction with little negative effect on the final yield of tetraphenoxymethane and diphenyl carbonate. Although diphenylthiocarbonate, DPTC, can be a precursor in the production of tetraphenoxymethane, this half-product is not always seen as a discrete intermediate on the path to tetraphenoxymethane. It is more efficient if the
10 diphenylthiocarbonate half-product goes on to directly form tetraphenoxymethane so that an additional conversion step is not necessary. Therefore, reactions and conditions which produce little or no diphenylthiocarbonate by the end of the reaction are preferred.

Gas chromatography-mass spectrometer (GC-MS) analysis of the
15 products of reactions which used high levels of phosphine ligands (10 mole percent) show that both the phosphine oxide and phosphine sulfide are produced during the reactions, concurrent with a decrease in the parent phosphine. The bidentate ligands themselves were too large to be observed by gas chromatography, therefore neither the oxide nor the sulfide were seen; it
20 is assumed, however, that phosphine oxides and sulfides would also be produced in these cases.

A set of control reactions were done that utilized only triphenylphosphine (at a 50 mole percent level) in conjunction with phenol and carbon disulfide; i.e. - no cuprous oxide, Cu_2O , was used. This set of
25 reactions tested the catalytic ability of triphenylphosphine itself to convert carbon disulfide and phenol to tetraphenoxymethane while also evaluating if the conversion of triphenylphosphine to the phosphine oxide and phosphine sulfide occurred under typical reaction conditions. No reaction to

tetraphenoxymethane occurred without cuprous oxide. Low levels of triphenylphosphine oxide, (TPP-O) and triphenylphosphine sulfide (TPP-S) were seen. However, when the amounts were compared to those in a cuprous oxide containing reaction, the level of TPP-O was one-fifth and the level of TPP-S was less than 1% without the use of cuprous oxide. This indicates that triphenylphosphine is capable of acting as a scavenger for sulfur, thus forming TPP-S during the conversion of carbon disulfide and phenol to DPTC and tetraphenoxymethane in the presence of cuprous oxide, so that not all of the sulfur is transferred directly to form cuprous sulfide, Cu_2S .

The metal oxide promoter is cuprous oxide Cu_2O . The use of small amounts of aromatic or aliphatic phosphines as a ligand significantly improves the reaction rate and overall yield of tetraphenoxymethane and ultimately diphenyl carbonate.

The instant invention relates to a method for manufacturing carbonate ester, which is subsequently used to produce polycarbonate. More particularly, it relates to a method whereby carbonate ester is obtained by the metal induced reaction of a hydroxy compound with a sulfur compound. Products of the present invention include, but are not limited to, diphenyl carbonate, di(O-phenyl) thiocarbonate, and diethyl carbonate.


The hydroxy compound comprises materials represented by the formula



wherein R is selected from the group consisting of substituted and unsubstituted C_{1-8} alkyl radicals, and substituted and substituted C_{6-13} aryl radicals. It includes cresols, bisphenol-A, xylenols, p-cumyl phenol, n-alkylated phenols, halogenated phenols, and alcohols. Preferably, the

hydroxy compound includes either substituted or unsubstituted phenol, or anhydrous ethanol.

For use in the instant invention, the sulfur compound comprises materials represented by formulas selected from the following:



wherein R' is a divalent aliphatic radical, O, or S; and R'' is R or OR; and wherein R is as previously defined. Suitable sulfur compounds include carbon disulfide and di (O-phenyl) thiocarbonate.

10

Addition of phosphine ligands can produce high yields of tetraphenoxymethane and diphenylcarbonate in short reaction times and with improved reaction rates by copper-induced condensation of phenol and carbon disulfide. Either aromatic or aliphatic substituted phosphine ligands can be employed. The copper(I) cuprous oxide utilization is significantly improved in reactions containing phosphine ligands, such that yields exceeding 100% tetraphenoxymethane are observed.

15

Water may be added to the reagents. Water may be added in a quantity sufficient to render it a co-solvent with the hydroxy compound, such as phenol, wherein a homogeneously mixed solution results. Water can also be added in an excess quantity sufficient to produce a two phase system with the hydroxy compound.

20

The materials can reacted in either a batch or continuous process. They can be heated to a temperature of about 20 - 250°C. Preferably, they are heated to a temperature of about 45° to about 180°C.

5 The reactor can be operated at reflux or under pressure. The pressure can range from atmospheric pressure to 20 Mpa. Preferably, the pressure is held in the range of atmospheric pressure to 1 Mpa. The reaction atmosphere can be comprises of air or an inert gas, such as argon or nitrogen. The reagents should be mixed sufficiently to make a heterogeneous slurry.

10 The total reaction time is dependent in part of temperature and on the desired degree of completion of the reaction. The required reaction time is determinable by one skilled in the art.

The materials produced by the present invention can be reacted under melt conditions with substantially no solvent present among the reagents. The process of the instant invention can also be run in the presence of a solvent. Suitable solvents are well known in the art. They include
15 hydrocarbons such as toluene, xylenes, hexanes, heptanes, and octanes; acetonitrile; benzonitrile; ethers such as tetrahydrofuran, ether, dibutyl ether, and methyl butyl ether; methyl ethyl ketone; acetone; methyl isobutyl ketone; chlorinated hydrocarbons such as chlorobenzenes and chlorobenzenes;
20 nitromethane; and nitrobenzene.

Alternatively, orthocarbonate can be generated by the method of the instant invention. The orthocarbonate comprises materials represented by the formula:



25 wherein each R is as previously defined.

When orthocarbonate is generated, the method of the instant invention comprises a further step of converting the orthocarbonate to a carbonate ester. When the method is used to generate orthocarbonate as a precursor to carbonate ester, it is preferable to remove water formed during the reaction of the promoter, hydroxy compound and sulfur compound. This eliminates any potential hydrolysis during preparation of the orthocarbonate. This orthocarbonate can subsequently be used to produce a carbonate ester by acidic hydrolysis.

Ligands include compounds derived from phosphine by replacement of hydrogen atoms with alkyl or aryl substituents.

Illustrative phosphines include triphenyl phosphine, triphenyl phosphite, triphenyl phosphine oxide, tricyclohexyl phosphine, tributyl phosphine, 1,2-bis(diphenylphosphino) ethane, and 1,2-bis(diphenylphosphino) butane.

DESCRIPTION OF FIGURES

Figure 1. Comparison Between Reactions using cuprous oxide, phenol and carbon disulfide to Form tetraphenoxymethane ; With (131A) and Without (116A) 10 mol% Triphenylphosphine (TPP).

Figure 2: Comparison Between Reactions using cuprous oxide, phenol and carbon disulfide along with the ligand Triphenylphosphine (TPP) to Form tetraphenoxymethane ; no water added 1 mol % triphenylphosphine (35A) vs. 10 mol % triphenylphosphine (131A)

Figure 3: Comparison Between Reactions using cuprous oxide, phenol and carbon disulfide along with the ligand Triphenylphosphine (TPP) to Form tetraphenoxymethane ; water added 1 mol % triphenylphosphine (35B) vs. 10 mol % TPP (131B)

Figure 4: Comparison Between Reactions using cuprous oxide , phenol and carbondisulfide along with the ligand Tricyclohexyl phosphine (tricyclohexyl phosphine) to form tetraphenoxymethane ; water added

ADDED; 0.1 mol % (CyHex)₃P (1029B), 1 mol % (CyHex)₃P (1024B),
5 and 10 mol % (CyHex)₃P (927B).

TABLE 1: Melt reactions of Cu₂O with phenol and CS₂; Control reactions and reactions containing aromatic and aliphatic phosphines as ligands

	Reaction [*]	ligand	ACN?	H ₂ O	3 hr yield	Total Yield (30h)	TPM/DPC ^{**}
10	116A (Control)	----	Y	N	34%	95%	93/2%
	116B (Control)	----	Y	Y	34%	78%	21/57%
15	131A	10 mol% TPPhosphine	Y	N	35%	135%	129/6%
	131B	10 mol% TPPhosphine	Y	Y	75%	105%	46/59%
	26A	10 mol% TPPhosphine	N	N	1.5%	95%	58/2%
	26B	10 mol% TPPhosphine	N	Y	34%	86%	38/23%
	114A	10 mol% TPPhosphine ^{***}	Y	N	57%	125%	70/5%
20	35A	1 mol% TPPhosphine	Y	N	65%	110%	106/4%
	35B	1 mol% TPPhosphine	Y	Y	84%	111%	69/42%
	919A	10 mol% diphos	Y	N	70%	113%	63/0%
	919B	10 mol% diphos	Y	Y	62%	93%	57/16%
	1014A	1 mol% diphos	Y	N	72%	86% (24h)	84/2%
	1014B	1 mol% diphos	Y	Y	42%	69% (24h)	36/33%
25	925A	10 mol% (Ø ₂ P-C ₄ -PØ ₂)	Y	N	34%	112%	32/2%
	925B	10 mol% (Ø ₂ P-C ₄ -PØ ₂)	Y	Y	34%	59%	20/12%
	927A	10 mol% (CyHex) ₃ -P	Y	N	75%	106% (70h)	53/3%
	927B	10 mol% (CyHex) ₃ -P	Y	Y	95%	100% (70h)	50/48%
	1024A	1 mol% (CyHex) ₃ -P	Y	N	117%	120%	117/3%
30	1024B	1 mol% (CyHex) ₃ -P	Y	Y	105%	114%	57/57%
	1029A	0.1 mol% (CyHex) ₃ -P	Y	N	40%	89%	84/5%
	1029B	0.1 mol% (CyHex) ₃ -P	Y	Y	90%	106%	33%/73%
	1017A	10 mol% Tributylphosphine	Y	N	55%	70%	56/6%
	1017B	10 mol% Tributylphosphine	Y	Y	45%	58%	24/22%
35	311A	10 mol% TPPhosphite	Y	N	22%	42%	19/19%
	311B	10 mol% TPPhosphite	Y	Y	36%	51%	12/39%
	102A	10 mol% TPPhosphine Oxide	Y	N	43%	53% (23h)	53/0%
	102B	10 mol% TPPhosphine Oxide	Y	Y	36%	36% (23h)	15/21%

40 * All reactions were run at 65°C under N₂, with mechanical stirring.

** Remainder of yield is DPTC.

*** Addition of 3 g of mol. sieves (3Å) to remove any water formed during reaction.

as a scavenger for sulfur, thus forming TPP-S during the conversion of CS₂ and phenol to DPTC and tetraphenoxymethane in the presence of Cu₂O, so that not all of the sulfur is transferred directly to form cuprous sulfide, Cu₂S.

EXPERIMENTAL

5 The melt reactions were run by combining phenol, cuprous oxide, phosphorus-containing ligand, internal standards, and a small amount of acetonitrile in a round bottom flask; some of the reactions contained water as a second phase. These reagents were stirred for 5 minutes at 45°C under a nitrogen blanket in order to melt the phenol and disperse the cuprous oxide. The carbon disulfide was added and the temperature was concurrently raised
10 to 65°C where it was kept for the duration of the reaction. Representative reactions and their charges are included below. The reagents were used as received with no drying or pre-purification steps.

Cuprous Oxide (Cu₂O), Phenol, CS₂ and 10 mole % Triphenylphosphine (TPP) as added Ligand, (CH₃CN, no H₂O) (131A)

15 Into a 250 mL, 3-neck round bottom flask, equipped with an overhead stir paddle, condenser and nitrogen blanket, were placed phenol (30.02g; 0.319 mol), cuprous oxide (1.5043; 0.01051 mol), triphenyl phosphine (0.555g; 0.002116 mol, 10 mol % based on Cu⁺), acetonitrile (2.5 mL; 0.04787 mol), and internal standards [3-methyl anisole (0.3123g) and biphenyl (0.3104g)]. The
20 flask was lowered into a 45°C oil bath and stirring was commenced. The reaction was mixed, while the phenol melted, for 5 minutes prior to the addition of the carbon disulfide, CS₂. The carbon disulfide was added (2 mL; 0.3325 mol) and the temperature of the oil bath was immediately raised to 65°C for the duration of the reaction (120 h). The stir speed was fast to ensure
25 good mixing of the heterogeneous mixture. During the course of the reaction, it changed from a dark red suspension to a well dispersed black suspension as

the dark red Cu_2O was converted to black Cu_2S . Samples were taken every 15-30 min. for the first three hours and intermittently thereafter. 0.4 mL samples were pipetted out, diluted with 1.5 mL CH_3CN , followed by filtration using Whatman Uniprep filters, containing a 0.45 μm PTFE membrane. Samples were analyzed using a Hewlett Packard 5890 Series II gas chromatograph equipped with a 30 meter DB-1 coated capillary column and a flame ionization detector (FID).

Cuprous Oxide (Cu_2O), Phenol, CS_2 and 10 mole % Triphenylphosphine (triphenylphosphine) as added Ligand, (CH_3CN , H_2O) (131B)

Into a 250 mL, 3-neck round bottom flask, equipped with an overhead stir paddle, condenser and nitrogen blanket, were placed phenol (30.04g; 0.319 mol), cuprous oxide (1.5025; 0.01050 mol), triphenyl phosphine (0.5575g; 0.002126 mol, 10 mol % based on Cu^+), acetonitrile (2.5 mL; 0.04787 mol), water (36 mL; 2.0mol), and internal standards [3-methyl anisole (0.3018g) and biphenyl (0.3018g)]. The flask was lowered into a 45°C oil bath and stirring was commenced. The reaction was mixed, while the phenol melted, for 5 minutes prior to the addition of the carbon disulfide, CS_2 . The carbon disulfide was added (2 mL; 0.3325 mol) and the temperature of the oil bath was immediately raised to 65°C for the duration of the reaction (47 h). The stir speed was fast to ensure good mixing of the heterogeneous mixture. During the course of the reaction, it changed from a dark red suspension to a well dispersed black suspension as the dark red Cu_2O was converted to black Cu_2S . Samples were taken every 15-30 min. for the first three hours and intermittently thereafter. 0.7 mL samples were pipetted out, diluted, filtered and analyzed as stated above.

Cuprous Oxide (Cu_2O), Phenol, CS_2 and 1 mole % Triphenylphosphine (triphenylphosphine) as added Ligand, (CH_3CN , H_2O) (35B)

Into a 250 mL, 3-neck round bottom flask, equipped with an overhead stir paddle, condenser and nitrogen blanket, were placed phenol (30.03g; 0.319 mol), cuprous oxide (1.5110; 0.01056 mol), triphenyl phosphine (0.0587g; 0.0002238 mol, 1 mol % based on Cu^+), acetonitrile (2.5 mL; 0.04787 mol),
5 water (36 mL; 2.0mol), and internal standards [3-methyl anisole (0.3196g) and biphenyl (0.3060g)]. The flask was lowered into a 45°C oil bath and stirring was commenced. The reaction was mixed, while the phenol melted, for 5 minutes prior to the addition of the carbon disulfide, CS_2 . The carbon disulfide was added (2 mL; 0.3325 mol) and the temperature of the oil bath
10 was immediately raised to 65°C for the duration of the reaction (47 h). The stir speed was fast to ensure good mixing of the heterogeneous mixture. During the course of the reaction, it changed from a dark red suspension to a well dispersed black suspension as the dark red Cu_2O was converted to black Cu_2S . Samples were taken every 15-30 min. for the first three hours and
15 intermittently thereafter. 0.7 mL samples were pipetted out, diluted, filtered and analyzed as stated above.

Cuprous Oxide (Cu_2O), Phenol, CS_2 and 10 mole % 1,2-bis(diphenylphosphino)ethane (DiPhos) as added Ligand, (CH_3CN , no H_2O) (919A)

Into a 250 mL, 3-neck round bottom flask, equipped with an overhead
20 stir paddle, condenser and nitrogen blanket, were placed phenol (29.95g; 0.318 mol), cuprous oxide (1.5071; 0.01053 mol), 1,2-bis(diphenylphosphino)ethane (DiPhos, 0.8351g; 0.002096 mol, 10 mol % based on Cu^+), acetonitrile (2.5 mL; 0.04787 mol), and internal standards [3-methyl anisole (0.3073g) and biphenyl (0.3048g)]. The flask was lowered into
25 a 45°C oil bath and stirring was commenced. The reaction was mixed, while the phenol melted, for 5 minutes prior to the addition of the carbon disulfide, CS_2 . The CS_2 was added (2 mL; 0.3325 mol) and the temperature of the oil bath was immediately raised to 65°C for the duration of the reaction (27 h).

The stir speed was fast to ensure good mixing of the heterogeneous mixture. During the course of the reaction, it changed from a dark red suspension to a well dispersed black suspension as the dark red Cu_2O was converted to black Cu_2S . Samples were taken every 15-30 min. for the first three hours and
5 intermittently thereafter. 0.4 mL samples were pipetted out, diluted, filtered and analyzed as stated above.

Cuprous Oxide (Cu_2O), Phenol, CS_2 and 10 mole % Tricyclohexyl phosphine ((CyHex) $_3$ -P) as added Ligand, (CH_3CN , no H_2O) (927A)

Into a 250 mL, 3-neck round bottom flask, equipped with an overhead
10 stir paddle, condenser and nitrogen blanket, were placed phenol (29.29g; 0.311 mol), cuprous oxide (1.5056g; 0.01052 mol), Tricyclohexyl phosphine (tricyclohexyl phosphine), 0.7562g; 0.002122 mol, 10 mol % based on Cu^+ , acetonitrile (2.5 mL; 0.04787 mol), and internal standards [3-methyl anisole (0.3132g) and biphenyl (0.3001g)]. The flask was lowered into a 45°C oil bath
15 and stirring was commenced. The reaction was mixed, while the phenol melted, for 5 minutes prior to the addition of the carbon disulfide, CS_2 . The carbon disulfide was added (2 mL; 0.0325 mol) and the temperature of the oil bath was immediately raised to 65°C for the duration of the reaction (70 h). The stir speed was fast to ensure good mixing of the heterogeneous mixture.
20 During the course of the reaction, it changed from a dark red suspension to a well dispersed black suspension as the dark red Cu_2O was converted to black Cu_2S . Samples were taken every 15-30 min. for the first three hours and intermittently thereafter. 0.4 mL samples were pipetted out, diluted, filtered and analyzed as stated above.

25 *Cuprous Oxide (Cu_2O), Phenol, CS_2 and 1 mole % Tricyclohexyl phosphine ((CyHex) $_3$ -P) as added Ligand, (CH_3CN , no H_2O) (927A)*

Into a 250 mL, 3-neck round bottom flask, equipped with an overhead stir paddle, condenser and nitrogen blanket, were placed phenol (30.36g; 0.322 mol), cuprous oxide (1.5163; 0.01060 mol), Tricyclohexyl phosphine (tricyclohexyl phosphine), 0.0766g; 0.0002149 mol, 1 mol % based on Cu⁺,
5 acetonitrile (2.5 mL; 0.04787 mol), and internal standards [3-methyl anisole (0.3132g) and biphenyl (0.3130g)]. The flask was lowered into a 45°C oil bath and stirring was commenced. The reaction was mixed, while the phenol melted, for 5 minutes prior to the addition of the carbon disulfide, CS₂. The carbon disulfide was added (2 mL; 0.3325 mol) and the temperature of the oil
10 bath was immediately raised to 65°C for the duration of the reaction (30 h). The stir speed was fast to ensure good mixing of the heterogeneous mixture. During the course of the reaction, it changed from a dark red suspension to a well dispersed black suspension as the dark red Cu₂O was converted to black Cu₂S. Samples were taken every 15-30 min. for the first three hours and
15 intermittently thereafter. 0.4 mL samples were pipetted out, diluted, filtered and analyzed as stated above.

CLAIMS:

1. A method for making tetraaryloxyalkane and diaryl carbonate comprising contacting at least one hydroxyaromatic compound with carbon disulfide, cuprous oxide and from 0.1 to 10 mole percent based on the mols of cuprous oxide of a phosphine ligand.
2. The method of claim 1 in which the ligand is an aliphatic, cycloaliphatic, or aromatic phosphine.
3. The method according to claim 1 in which the reaction mixture includes auxiliary reagents selected from the group consisting of water and methylcyanide.
4. The method according to claim 1 in which the hydroxyaromatic compound is selected from the group consisting of unsubstituted phenols, cresols, xylenols, n-alkylated phenols, halogenated phenols, bisphenol A, and p-cumyl phenol.
5. The method according to claim 1 wherein the phosphine ligand is triphenylphosphine, tricyclohexylphosphine, triphenylphosphite, 1,2-bis(diphenylphosphino)ethane, tributylphosphine, triphenylphosphine oxide, or 1,2-(diphenylphosphino) butane