WHAT IS CLAIMED IS:

1. An anti-B cell maturation antigen (BCMA) antibody, or an antigen-binding fragment thereof, comprising a heavy chain variable region comprising a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 167, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 168, and a CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 169; and a light chain variable region comprising a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 170, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 171, and a CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 172.

2. An anti-B cell maturation antigen (BCMA) antibody, or an antigen-binding fragment thereof, comprising a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:53, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:54.

3. The anti-BCMA antibody or antigen-binding fragment thereof of claim 1 or claim 2, wherein the heavy chain variable region comprises an amino acid sequence that has at least 90% sequence identity to the sequence set forth in SEQ ID NO:53, and the light chain variable region comprises an amino acid sequence that has at least 90% sequence identity to the sequence set forth in SEQ ID NO:54.

4. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-3, wherein the heavy chain variable region comprises the amino acid sequence set forth in SEQ ID NO:53, and the light chain variable region comprises the amino acid sequence set forth in SEQ ID NO:54.

5. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-4, wherein the antigen-binding fragment of the antibody is an antigen-binding fragment (Fab), a Fab', a $F(ab')_2$, a variable fragment (Fv) or a single chain variable fragment (scFv).

6. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-5, comprising a linker positioned between the heavy chain variable region and the light chain variable

region.

7. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-6, wherein the antigen-binding fragment thereof is a single chain variable fragment (scFv).

8. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-7, comprising the amino acid sequence set forth in SEQ ID NO:85.

9. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-8, wherein the antibody comprises a human variable region framework region.

10. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-9, which is a fully human or an antigen-binding fragment thereof.

11. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-9, which is a chimeric antibody or an antigen-binding fragment thereof.

12. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-9, which is a humanized antibody or an antigen-binding fragment thereof.

13. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-8, wherein the antigen-binding fragment is a human single chain variable fragment (scFv).

14. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-6 and 8-12, wherein the antigen-binding fragment is an antigen-binding fragment (Fab).

15. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-6 and 8-12, wherein the antigen-binding fragment is a $F(ab')_2$.

16. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 5-15, wherein the one or more of the scFv, the Fab and the $F(ab')_2$ are comprised in a fusion protein with a heterologous sequence to form the extracellular antigen-binding domain.

17. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-16, wherein the antibody or antigen-binding fragment binds to B cell maturation antigen (BCMA) with a dissociation constant (K_d) of from 1×10^{-9} M to 1×10^{-6} M.

18. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-17, wherein the antibody or antigen-binding fragment binds to B cell maturation antigen (BCMA) with a dissociation constant (K_D) of from 1×10^{-9} M to 1×10^{-8} M.

19. The anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-18, wherein the antibody or antigen-binding fragment binds to a non-linear epitope of BCMA.

20. A composition comprising the anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-19, and a pharmaceutically acceptable carrier.

21. An immunoconjugate comprising the anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-19, linked to a therapeutic agent.

22. The immunoconjugate of claim 21, wherein the therapeutic agent is a drug, a cytotoxin, or a radioactive isotope.

23. A composition comprising the immunoconjugate of claim 21 or 22, and a pharmaceutically acceptable carrier.

24. A bispecific molecule comprising the anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-19, linked to a second functional moiety.

25. The bispecific molecule of claim 24, wherein the second functional moiety has a different binding specificity than the antibody or antigen binding fragment thereof.

26. The bispecific molecule of claim 24 or claim 25, wherein the second functional moiety has a binding specificity for an immune cell.

27. The bispecific molecule of any one of claims 24-26, wherein the second functional moiety has a binding specificity for CD3.

28. A composition comprising the bispecific molecule of any one of claims 24-27, and a pharmaceutically acceptable carrier.

29. A nucleic acid molecule that encodes an anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-19.

30. An expression vector comprising the nucleic acid molecule of claim 29.

31. A host cell comprising the expression vector of claim 30.

32. A method for detecting BCMA in a whole cell or tissue, comprising:

contacting a cell or tissue with the anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-19, wherein the antibody or antigen-binding fragment thereof comprises a detectable label; and

determining the amount of the labeled antibody or antigen-binding fragment thereof bound to the cell or tissue by measuring the amount of detectable label associated with the cell or tissue, wherein the amount of bound antibody or antigen-binding fragment thereof indicates the amount of BCMA in the cell or tissue.

33. Use of a composition comprising the anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-19 for the manufacture of a medicament for treating a tumor in a subject.

34. The use of claim 33, wherein the medicament reduces the number of the tumor cells.

35. The use of claim 33 or claim 34, wherein the medicament reduces the tumor size.

36. The use of any one of claims 33-35, wherein the medicament eradicates the tumor in the subject.

37. The use of any one of claims 33-36, wherein the tumor is selected from the group consisting of multiple myeloma, Non-Hodgkin Lymphoma, Hodgkin Lymphoma, Chronic Lymphocytic Leukemia (CLL), glioblastoma, and Waldenstrom’s Macroglobulinemia.

38. The use of any one of claims 33-37, wherein the tumor is multiple myeloma.

39. The use of any one of claims 33-38, wherein the subject is a human.

40. A kit for treating a tumor, comprising the anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-19.

41. The kit of claim 40, wherein the kit comprises written instructions for using the antibody or antigen-binding fragment thereof for treating a subject having a tumor.

42. The kit of claim 40 or claim 41, wherein the tumor is multiple myeloma.

43. The composition comprising the anti-BCMA antibody or antigen-binding fragment thereof of any one of claims 1-19 for use as a therapy.