**WHAT IS CLAIMED IS:**

1. Use of 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid or a pharmaceutically salt thereof in the manufacture of a pharmaceutical composition for the treatment of T2DM, wherein:

the pharmaceutical composition is in an oral dosage form; and

the pharmaceutical composition is to be administered to a human in need thereof

2. The use of claim 1, wherein the pharmaceutical composition is in an immediate-release solid dosage form.

3. The use of claim 1 or 2, wherein the pharmaceutical composition is in an immediate-release tablet dosage form.

4. The use of any one of claims 1 to 3, wherein the pharmaceutical composition comprises 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid or a pharmaceutically salt thereof in an amount equivalent to about 10 mg to about 140 mg of 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid.

5. The use of any one of claims 1 to 3, wherein the 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid or pharmaceutically salt thereof in the pharmaceutical composition is tris salt of 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid.

6 The use of any one of claims 1 to 5, wherein the pharmaceutical composition is to be administered twice daily.

7. The use of claim 6, wherein the two daily administrations are separated by at least about 8 hours.

8. The use of any one of claims 1 to 7, wherein the treating D2TM comprises improving glycemic control.

9. Use of 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid or a pharmaceutically salt thereof in the manufacture of a

pharmaceutical composition for weight management control or for the treatment of obesity or overweight, wherein:

the pharmaceutical composition is in a solid oral dosage form; and

the pharmaceutical composition is to be administered to a human in need thereof.

10. The use of claim 9, wherein the pharmaceutical composition comprises 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid or a pharmaceutically salt thereof in an amount equivalent to about 10 mg to about 140 mg of 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid.

11. The use of claim 9 or 10, wherein the pharmaceutical composition is to be administered twice daily.

12. The use of claim 11, wherein the two daily administrations are separated by at least about 8 hours.

13. An immediate-release oral pharmaceutical composition comprising 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid or a pharmaceutically salt thereof.

14. An immediate-release oral pharmaceutical composition comprising:

2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, or a pharmaceutically acceptable salt thereof;

a filler;

a disintegrant; and

a lubricant.

15. The pharmaceutical composition of claim 14 wherein:

the 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid or a pharmaceutically acceptable salt thereof is tris salt of 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

the filler comprises microcrystalline cellulose, lactose, or a combination thereof;

the disintegrant is sodium starch glycolate; and

the lubricant is magnesium stearate or sodium stearyl fumarate.

16. The pharmaceutical composition of claim 14 wherein:

the 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid or a pharmaceutically acceptable salt thereof is tris salt of 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

the filler comprises microcrystalline cellulose, lactose, or a combination thereof;

the disintegrant is crospovidone; and

the lubricant is magnesium stearate or sodium stearyl fumarate.

17. The pharmaceutical composition of claim 14 or 15, wherein:

the 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid or a pharmaceutically acceptable salt thereof comprises about 1.0% to about 35.0% by weight;

the filler comprises about 60% to about 95% by weight;

the disintegrant comprises about 1.0% to about 5.0% by weight; and

lubricant comprises about 0.2% to about 2.5% by weight.

18. The pharmaceutical composition of any one of claims 13 to 17, wherein the composition comprise tris salt of 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid in an amount equivalent to about 10 mg to about 120 mg of 2-[(4-{6-[(4-cyano-2-fluorobenzyl)oxy]pyridin-2-yl}piperidin-1-yl)methyl]-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid.

19. The pharmaceutical composition of any one of claims 13 to 17 for use in the treatment of T2DM or in weight management control.

20. The pharmaceutical composition of claim 19 wherein the composition is used by twice daily administration.