

33375-00

ARYLPYRROLES FOR SUBTERRANEAN TERMITE CONTROL

BACKGROUND OF THE INVENTION

A broad class of arylpyrrole compounds and their insecticidal use in crop protection and animal health are described in U.S. 5,010,098; U.S. 5,310,938; and U.S. 5,455,263. It has now been found that a small select
5 group of these arylpyrrole compounds are particularly useful for the control of pestiferous subterranean termites.

Termites are known to occur in virtually every state in the U.S., except Alaska, in all U.S. territories, and
10 throughout the world in every continent except Antarctica. Termites cause extensive damage to wood, wood structures, wood products and cultivated plants and crops. They invade and damage poles, posts, timbers, lumber, buildings, shelters and the like. They are also
15 destructive to cultivated plants and crops where they burrow into stems, tunnel out stalks and roots, and girdle the bark of trees. They eat grass and injure or damage field crops such as rubber, tea, coffee, cocoa, citrus and sugarcane.

20 Conventional methods to control termites have a common mechanism of action. This mechanism employs a chemical zone through which termites will not travel. This zone acts as a barrier by, either directly or indirectly, repelling the termites. However, while
25 effective in repelling termites, conventional treatments cause very little, if any, effective control of the termite populations, as very few termites which encounter



the barrier actually expire. Therefore, due to the minimal termite mortality and impact on termite foraging activity of conventional termite control methods, there remain vast populations of termites that continue to
5 represent a significant threat to growing and harvested wood, wood products and wooden structures. In light of the limitations of conventional materials and methods to control termite populations, alternatives to the conventional methods for controlling these significantly
10 destructive insects is highly desirable.

Therefore, it is an object of this invention to provide an efficient, efficacious method for the control of pestiferous subterranean termite populations.


It is a feature of this invention that the pressure
15 and threat of destructive termite populations may be significantly reduced.

Further objects and features of the invention will become apparent in the detailed description set forth below.

20

SUMMARY OF THE INVENTION

The present invention provides a method for the control of pestiferous subterranean termite populations
25 which comprises applying to the soil a termiticidally effective amount of a select arylpyrrole compound of formula I.



(I)

wherein R is hydrogen or $\text{C}_1\text{-C}_4$ alkoxymethyl;

X is Cl or Br , and


5 L , M , and Q are each independently hydrogen, Cl , Br ,
 I , F or $\text{C}_1\text{-C}_4$ haloalkyl.

DETAILED DESCRIPTION OF THE INVENTION

Protecting growing and harvested wood, wood products
10 and wooden structures from damage and destruction caused
by termites while effectively controlling the populations
of said termites is a continuing scientific challenge.
Conventional methods and materials such as organo-
phosphates, pyrethroids, chlorinated hydrocarbons, and
15 the like employ a common mechanism of action which is to
essentially to form a chemical barrier which directly or
indirectly repels the targeted termites. Consequently,
the termite population size, and the pressure from said
population, is not significantly reduced. Therefore,
20 because these populations continue to persist in the
targeted ecological environment, they continue to forage
and, through trial and error, find gaps in the chemical
barrier to gain access to the wood, wood product, or
wooden structure meant to be protected. Further, in the
25 case of subterranean termite populations, post-treatment
activities such as gardening, paving, remodeling, and the
like may disturb the chemical barrier in the soil and
create gaps that may be breached by the targeted
subterranean termite populations. Moreover, as the

conventional materials degrade over time and become less effective as repellents, the largely uncontrolled populations of termites can return to attack and infest the susceptible wood, wood product or wooden structure.

5 It has now been found that a select group of arylpyrrole compounds of formula I



(I)

wherein R is hydrogen or C₁-C₄alkoxymethyl;

10 X is Cl or Br; and

L, M and Q are each independently hydrogen,
Cl, Br, I, F or C₁-C₄haloalkyl;

is particularly useful for the control of pestiferous subterranean termites. Advantageously, said compounds
15 provide a non-repellent, slow-acting toxin useful for the control of pestiferous subterranean termites which imbibe or ingest soil for excavation or nutrition. Said

subterranean termites exchange nutrients between colony members; therefore, a slow-acting toxin allows sufficient
20 time for one member to pass the toxin to another member before visible toxic effects are manifested. This slow action is particularly useful for the effective control of subterranean, social, termite populations. Further, since the select formula I arylpyrrole compounds are non-
25 repellent, the targeted pestiferous termite colony members will continually travel through, attack or infest material which has been treated according to the method of invention, resulting in significant mortality, and control

of said colony members and concomitant reduction in termite population pressure.

In the specification and claims, the term "subterranean termite" designates the order Isoptera and encompasses the families of *Rhinotermitidae*, *Termitidae*,
5 *Kalotermitidae*, and *Hodotermitidae*. Examples of species of subterranean termites controlled by the method of invention include *Reticulitermes flavipes*, *Reticulitermes virginicus*, *Reticulitermes hesperus*, *Coptotermes*
10 *formosanus* and the like.

Surprisingly, among the vast genre of arylpyrrole compounds known in the art, a small select group of formula I arylpyrroles are especially effective and particularly useful for the control of subterranean
15 termites. Those particular arylpyrrole compounds of formula I preferred for use in the method of invention are those formula I compounds wherein R is C₁-C₄alkoxymethyl.

Another group of particular arylpyrrole compounds preferred for use in the method of invention is that
20 group of compounds of formula I wherein L and Q are hydrogen and M is 4-Cl.

A third group of compounds of formula I especially useful in the method of the invention is that group wherein X is Br.

25 Exemplary of the select formula I arylpyrrole compounds which are useful in the inventive method are:
4-bromo-2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(p-chlorophenyl)-1-(ethoxymethyl)-5-(trifluoro-
30 methyl)pyrrole-3-carbonitrile;
4-chloro-2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-chloro-2-(p-chlorophenyl)-1-(ethoxymethyl)-5-(trifluoro-
methyl)pyrrole-3-carbonitrile;

- 4-chloro-2-(p-chlorophenyl)-1-(methoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(p-chlorophenyl)-1-(methoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
5 4-bromo-1-(ethoxymethyl)-5-(trifluoromethyl)-2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile;
4-bromo-2-(3,4-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(2,4-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
10 4-bromo-2-(3,5-difluorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(2,3-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
15 4-bromo-2-(3,4-difluorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(3,5-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(3,4,5-trifluorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
20 4-bromo-2-(3,4,5-trichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(2,5-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
25 4-bromo-2-(2,6-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-chloro-2-(3,4-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-chloro-2-(3,5-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile; or
30 4-chloro-1-(ethoxymethyl)-5-(trifluoromethyl)-2-(α,α,α -p-tolyl)pyrrole-3-carbonitrile.

Those compounds particularly useful in the method of the invention are: 4-bromo-2-(p-chlorophenyl)-1-(ethoxy

methyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-chloro-2-(p-chlorophenyl)-1-(ethoxymethyl)-5-(trifluoro-
methyl)pyrrole-3-carbonitrile;
4-bromo-2-(3,5-dichlorophenyl)-1-(ethoxymethyl)-5-
5 (trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-
carbonitrile; and
4-chloro-2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-
carbonitrile.

10 More preferably, 4-bromo-2-(p-chlorophenyl)-1-
(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile
is useful in the inventive method for controlling
pestiferous subterranean termite populations.

In accordance with the method of invention, effective
15 control of pestiferous subterranean termite populations
which imbibe or ingest soil for excavation or nutrition may
be obtained by treating the soil with a spray, dust,
powder, granule, solution, emulsion, suspension or
dispersion containing an effective amount of the select
20 arylpyrrole compound of formula I. Rates of application of
said arylpyrrole compound will vary according to prevailing
environmental conditions such as insect species, population
density, weather conditions, soil type, vegetation growth,
topographical characteristics, and the like. In general,
25 rates of application of active ingredient sufficient to
supply soil concentrations of about 1 ppm to 2,500 ppm,
preferably about 25 ppm to 1,000 ppm may be effective.

For a more clear understanding of the invention, the
following examples are set forth below. These examples
30 are merely illustrative and are not understood to limit
the scope or underlying principles of the invention in
any way. Indeed, various modifications of the invention,
in addition to those shown and described herein, will
become apparent to those skilled in the art from the

following examples and the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

5

EXAMPLE 1

Evaluation Of The Termiticidal Activity Of Chlorfenapyr Over Time

For this evaluation, ten termites (*Reticulitermes virginicus*) are placed in a 5 cm petri dish containing a thin layer of moist soil which has been treated with an acetone solution of chlorfenapyr at various concentrations. (The acetone is evaporated before introduction of insects.) Observations of insect mortality are made at 8 hr. and 24 hr. after treatment. After 24 hr., all termites are removed from the treated soil and placed in 5 cm petri dishes containing non-treated soil and a strip of moistened cellulose filter paper as food supply. Observations of continued insect mortality are made daily for 7 days after exposure to the treated soil. The treatments are replicated 3 times. The data obtained are averaged and shown in Table I, below.

25

Table I

Test Compound	Concentration (ppm)	% Mortality (Time After Treatment)		
		8 hr.	24 hr.	7 days
Chlorfenapyr	100	0	100	100
Chlorfenapyr	50	0	33.7	100
Chlorfenapyr	25	0	6.7	100
Chlorfenapyr	10	0	0	100
Chlorfenapyr	5	0	0	100
Untreated Control	0	0	0	0

DISCUSSION

In this Example, the delayed-action mortality in
5 termites exposed to the arylpyrrole compound is evident.
There is no mortality after 8 hours of exposure to
chlorfenapyr (Table I). However, after 24 hours the
percentage of mortality increases. Further, over the
next six days, even after the termites have been removed
10 to untreated soil, mortality continues to increase.
Ultimately, exposure to concentrations as low as 5 ppm
chlorfenapyr causes 100% mortality.


15

EXAMPLE 2

Evaluation Of The Termiticidal Activity Of Test Compounds Over Time

In this evaluation, samples of test compound are
20 dissolved in acetone and applied to a sandy loam soil in
quantities sufficient to supply concentrations of 1,000,
100, 10 and 1 ppm by weight. Treated soil is then
moistened with 5% by weight of water. The moistened,
treated soil (5 g) is placed uniformly over the bottom of
25 a 100 x 15 mm glass dish. Ten worker termites
(*Reticulitermes* spp.) are placed on the treated soil and
held for 7 days. Observations of moribundity are made
daily. In this evaluation, moribundity is defined as
ataxia, paralysis, or sluggishness. Moribund termites
30 are recorded and removed daily. Each treatment is
replicated 4 times. The data are averaged and shown in
Table II, below.

Table II



Test Compound					Rate	% Moribundity	
R	X	L	M	Q	(ppm)	1 DAT	7 DAT
H	Br	H	4-Cl	H	1,000	100	100
H	Br	H	4-Cl	H	100	46.7	100
H	Br	H	4-Cl	H	10	3.2	75.6
H	Br	H	4-Cl	H	1	10.0	16.7
CH ₂ OC ₂ H ₅	Br	H	4-Cl	H	1,000	98.3	100
CH ₂ OC ₂ H ₅	Br	H	4-Cl	H	100	19.2	100
CH ₂ OC ₂ H ₅	Br	H	4-Cl	H	10	6.7	91.7
CH ₂ OC ₂ H ₅	Br	H	4-Cl	H	1	5.0	49.4
CH ₂ OC ₂ H ₅	Br	H	4-Br	H	1,000	96.7	100
CH ₂ OC ₂ H ₅	Br	H	4-Br	H	100	98.3	100
CH ₂ OC ₂ H ₅	Br	H	4-Br	H	10	11.7	100
CH ₂ OC ₂ H ₅	Br	H	4-Br	H	1	10.0	43.3
CH ₂ OC ₂ H ₅	Cl	H	4-Cl	H	1,000	100	100
CH ₂ OC ₂ H ₅	Cl	H	4-Cl	H	100	100	100
CH ₂ OC ₂ H ₅	Cl	H	4-Cl	H	10	26.4	100
CH ₂ OC ₂ H ₅	Cl	H	4-Cl	H	1	1.7	86.7
CH ₂ OC ₂ H ₅	Br	3-Cl	H	5-Cl	1,000	100	100
CH ₂ OC ₂ H ₅	Br	3-Cl	H	5-Cl	100	90	100
CH ₂ OC ₂ H ₅	Br	3-Cl	H	5-Cl	10	3.3	96.8
CH ₂ OC ₂ H ₅	Br	3-Cl	H	5-Cl	1	3.3	36.7
Untreated Control					0	1.9	16.8

DISCUSSION

The unique delayed-action mortality of the formula I arylpyrrole compound is demonstrated by the data shown in Table II of this Example. For instance, exposure to arylpyrrole compounds at concentrations of 10 ppm causes little moribundity after 24 hours (3.2% to 26.4%). However, after 7 days exposure, significant moribundity (75.6% to 100%) results for all of these compounds.

EXAMPLE 3

Evaluation Of Termiticidal Activity Of Chlorfenapyr On A Variety Of Species

In this evaluation, moist soil is treated with a gradient dilution of a 2SC formulation of chlorfenapyr as described in U.S. Patent 5,496,845. Groups of 10 termites each from 4 different species of termites are placed in a 5 cm petri dish containing a thin layer of the treated soil. Observations of insect mortality are made at 8 hr. and 24 hr. after treatment. After 24 hr., all termites are removed from the treated soil and placed in petri dishes containing non-treated soil and a strip of moistened cellulose filter paper as food. Insect mortality observations are made daily, for 7 days after exposure to the treated soil. Each treatment is replicated 6 times. The data are averaged and shown in Table III, below.

Table III

Test	Rate	% Mortality at 7 DAT			
		<i>Reticulitermes</i>	<i>Reticulitermes</i>	<i>Reticulitermes</i>	<i>Coptotermes</i>
Compound	(ppm)	<i>flavipes</i>	<i>virginicus</i>	<i>hesperus</i>	<i>formosanus</i>
Chlorfenapyr	1000	96.7	86.7	100	96.7
Chlorfenapyr	500	90.0	78.3	100	98.3
Chlorfenapyr	100	18.3	16.7	22.5	43.3
Chlorfenapyr	50	8.3	20.0	20.0	36.7
Chlorfenapyr	10	3.3	16.7	10.0	23.3
Untreated Control	0	6.7	0	5.0	10.0

DISCUSSION

5

In this Example wherein all of the major species of subterranean termites in the U.S. are studied, the data in Table III clearly demonstrate that all termite species tested are equally susceptible to the lethal effects of the formula I arylpyrrole compound.

10

EXAMPLE 4

15 Comparative Evaluation Of Non-Repellency And Efficacy Of
Formula I Arylpyrrole Compounds And Pyrethroid Compounds
In The Soil

20 A sandy loam soil is treated with acetone solutions of either chlorfenapyr or permethrin to obtain concentrations of 100, 10, or 1 ppm. Untreated soil is used for the controls. After treatments have dried, enough treated soil is transferred into 15 cm lengths of transparent plastic tubing (1.5 cm diam.) to form a



column of treated soil 5 cm in length. A 1 cm thick 6% agar plug is fitted at each end of the 5 cm column of soil. Several pieces of woodflour (food source) are placed in one open end which, when the tube is inverted, is situated beneath the treated-soil column. Thirty (30) termites (*Reticulitermes flavipes*) are then introduced into the top open end of the plastic tubing opposite the food source. Both ends of the tubing are sealed, to form a "test unit". Test units are held at room temperature, in the dark, for the duration of the study. After 1 day and 3 days, measurements are taken of the distance the termites tunnel into the 5 cm column of treated soil. Each concentration is replicated 4 times for each test compound. At the conclusion of the experiment, the test units are destructively sampled to determine termite mortality.

Table IV

<u>Test Compound</u>	<u>Rate (ppm)</u>	<u>Depth in cm of termite penetration (% of 5 cm)</u>				<u>% Mortality</u>
		<u>1 dat</u>		<u>3 dat</u>		<u>3 dat</u>
chlorfenapyr	100	4.4	(88%)	5.0	(100%)	93.0%
	10	3.8	(76%)	5.0	(100%)	46.0%
	1	5.0	(100%)	5.0	(100%)	33.0%
permethrin	100	0.0	(0%)	0.0	(0%)	7.0%
	10	0.1	(2%)	0.2	(4%)	0.0%
	1	1.2	(24%)	2.6	(52%)	17.0%
untreated control	0	3.3	(66%)	5.0	(100%)	20.0%

Discussion

In Table IV, repellency is clearly demonstrated for the pyrethroid compound, permethrin. There is very little penetration by the termites into the permethrin treated soil column, especially at rates of 100 or 10 ppm. As can be seen, few termites died from exposure due to the repellency of permethrin. Even at 1 ppm concentration of permethrin when termites penetrated through more than 50% of the treated soil column, mortality did not differ from that observed in the untreated control. In contrast, the formula I arylpyrrole compound, chlorfenapyr, clearly demonstrated non-repellency. At all three concentrations of chlorfenapyr, the termites rapidly penetrated the treated soil column. Further, exposure to the chlorfenapyr in the treated soil columns proved lethal to the termites. Even at the lowest concentration tested (1 ppm), exposure to chlorfenapyr caused higher mortality in termites than did exposure to the same 1 ppm concentration of permethrin.

EXAMPLE 5

Evaluation Of Non-Repellency And Efficacy Of Test Compounds In The Soil

In this evaluation, 300 termites (*Reticulitermes flavipes*) are added to a 5 cm-round tubular container filled with a sand/vermiculite substrate and containing a small, pre-weighed pine block (unprotected wood). This container is connected with a 7 cm length of 3 mm plastic tubing to a second container filled with a sandy loam soil which has been treated with a test compound. At the opposite end of the second container is connected, by plastic tubing, a third container in which is placed a single pre-weighed block of pine wood (protected wood). Termites must tunnel through

the treated soil in order to access the "protected" block of pine wood. After 5 weeks, the termites from all containers are removed, % insect mortality is determined and the pine blocks are weighed. The treatments are replicated 6 times. The data is averaged and shown in Table V below.

Table V

<u>Test Compound</u>	<u>Rate (ppm)</u>	<u>% Mortality</u>	<u>Consumption of Unprotected Wood</u>	<u>Consumption of Protected Wood</u>
Chlorfenapyr	1000	100	36 mg	16 mg
Chlorfenapyr	500	99.4	52 mg	37 mg
Chlorfenapyr	100	93.4	308 mg	75 mg
Chlorfenapyr	50	87.7	625 mg	133 mg
Chlorfenapyr	10	44.5	833 mg	625 mg
Untreated Control	0	29.0	900 mg	842 mg

10

Discussion

In this Example, the data in Table V demonstrate the advantages of the non-repellence of an arylpyrrole compound, chlorfenapyr. In the absence of chlorfenapyr residues in the soil in the middle container, termites consume equal quantities of wood from both ends of the test chamber. However, in all tests where the soil in the middle container is treated with chlorfenapyr, significantly less wood is consumed from the protected pine block than is consumed from the unprotected pine block. In this study the percentage mortality is consistently $\geq 85\%$ at soil concentrations as low as 50 ppm.


20

33375-00



I CLAIM:

1. A method for the control of subterranean termite populations which comprises applying to the soil a termiticidally effective amount of an arylpyrrole compound of formula I



(I)

wherein R is hydrogen or C₁-C₄alkoxymethyl;

X is Cl or Br, and

L, M, and Q are each independently hydrogen, Cl, Br, I, F or C₁-C₄haloalkyl.

2. The method according to claim 1 wherein said arylpyrrole compound is applied at a rate sufficient to provide a soil concentration of about 1.0 ppm to 2500 ppm.

3. The method according to claim 2 wherein said arylpyrrole compound is applied at a rate sufficient to provide a soil concentration of about 5 ppm to 1000 ppm.

4. The method according to claim 3 wherein said arylpyrrole compound is applied at a rate sufficient to provide a soil concentration of about 25 ppm to 1000 ppm.



5. The method according to claim 1 having an arylpyrrole compound of formula I wherein R is C₁-C₄-alkoxymethyl.

6. The method according to claim 1 having an arylpyrrole compound wherein L and Q are hydrogen and M is 4-Cl.

7. The method according to claim 1 having an arylpyrrole compound wherein X is Br.

8. The method according to claim 1 wherein said arylpyrrole compound is selected from the group consisting of: 4-bromo-2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(p-chlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-chloro-2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-chloro-2-(p-chlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-chloro-2-(p-chlorophenyl)-1-(methoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(p-chlorophenyl)-1-(methoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-1-(ethoxymethyl)-5-(trifluoromethyl)-2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile;
4-bromo-2-(3,4-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(2,4-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(3,5-difluorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;



4-bromo-2-(2,3-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(3,4-difluorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(3,5-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(3,4,5-trifluorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(3,4,5-trichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(2,5-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(2,6-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-chloro-2-(3,4-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-chloro-2-(3,5-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile; and
4-chloro-1-(ethoxymethyl)-5-(trifluoromethyl)-2-(α,α,α -p-tolyl)pyrrole-3-carbonitrile.

9. The method according to claim 2 wherein said arylpyrrole compound is selected from the group consisting of: 4-bromo-2-(p-chlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-chloro-2-(p-chlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(3,5-dichlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
4-bromo-2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile; and
4-chloro-2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile.



-19-

10. The method according to claim 9 wherein said arylpyrrole compound is 4-bromo-2-(p-chlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile.