

NUCLEIC ACID LIBRARIES, PEPTIDE LIBRARIES AND USES THEREOF

DESCRIPTION

5 **[1]** The invention relates to libraries of nucleic acids that encode a plurality of peptides that represent fragments of naturally occurring proteins. Each peptide may be selected from, and collectively the plurality of peptides can be representative of, a proteome of a species or an organism, or such peptides may be selected from more than one proteome, and hence collectively the plurality can be representative of a metaproteome. The peptides may also be selected from proteins that are differentially expressed between cell or tissue types. The invention also relates to libraries of such peptides, to methods involving such libraries of nucleic acids and/or peptides, and/or to computer readable media or data-processing systems comprising information relating to such libraries.

10 **[2]** The identification of new therapeutic targets is a key starting point for drug discovery. Drug discovery efforts have traditionally been focused upon identifying classically-druggable targets, such as kinases, G-protein coupled receptors (GPCRs) and ion channels. However, such chemically facile targets do not always represent the most biologically relevant targets for therapeutic intervention. Drugging protein:protein interactions (PPIs) is of particular interest because these represent the predominant type of target involved in defective signalling pathways utilised by cancer cells and a large set of potentially actionable interfaces in human disease. Unfortunately, systematic attempts to drug PPIs and other 'undruggable' targets have been limited by technological restrictions, in large part due to limitations in current high-throughput DNA and RNA-based genomics technologies in being able to identify new druggable space at the proteome level.

15 **[3]** Current genomics-based technologies that can identify candidate drug targets linked to disease biology using unbiased 'phenotypic' assays, have typically been performed using gene knock-outs (e.g., CRISPR), or at the transcriptomic level using RNAi. These approaches yield important information on which targets may represent important nodes in disease progression and therapeutic intervention in disease, but suffer a serious limitation: because they screen at the genetic, rather than protein-level, they cannot identify how to drug those targets or determine if those represent druggable candidates as an inherent part of the process. This is because such genetic screens remove target proteins rather than inhibiting them. To gain such crucial additional information on druggability, a new high-throughput proteome-level screening technology would need to be used; one that can handle the higher complexity of screening protein function (>300,000 unique protein transcripts and millions of unique PPIs) compared to gene function (~30,000 genes and their splice variants).

20 **[4]** Recently, the systematic identification of novel drug target sites directly in the human proteome has gained a level of tractability and attention with the introduction of DNA-encoded, protein-fragment expression libraries that can be screened in high throughput in phenotypic assays (such as described in WO 2013/116903); often dubbed 'Protein-interference' (Protein-i). Such protein-fragment libraries, typically derived from diverse bacterial genomes, are composed of small self-folding sub-domains that form the evolutionary building blocks of larger proteins. When assembled into libraries for intracellular expression in mammalian cells, they represent a highly diverse collection of 3-dimensional shapes for docking to target proteins and exploring candidate novel druggable sites across the human proteome. Crucially, these protein fragments are small enough to describe discrete spatial sites in target proteins, and thus can be recapitulated with small-molecule drugs subsequently designed to match that shape. Moreover, because protein-fragment libraries describe many more shapes than current small-molecule libraries, this offers a more robust approach to informing the rational design of future small molecule drugs to novel validated targets.

30 **[5]** While bacterial-derived protein-fragment libraries have been shown to be successful in Protein-i screening and are straightforward to generate by fragmenting and cloning into expression libraries due to bacterial genomes

being composed mostly of coding sequence, they may, however, be under-powered in possessing a large fraction of protein-fragments that can functionally interact with mammalian (e.g. human) proteins, compared to using fragments of a mammalian or human proteome itself.

[6] However, creating protein-fragment libraries directly from a mammalian (e.g. human) genome is complicated by the fact that DNA of higher organisms contains mostly non-coding sequences (>95% of human DNA is estimated to be non-coding) and a much larger absolute number of coding sequences and thus generally require a large degree of manual bespoke cloning to assemble fragments thereof into expression libraries for phenotypic screening.

[7] Those bacterial-derived protein-fragment libraries described to date (e.g. in WO2013/116903) are obtained by mechanically shearing genomes and randomly inserting fragments into vectors. This leads to many fragments of random size that are either in frame (1:6 chance) or out of frame (5:6 chance) with the original gene in the bacterium. The same strategy would not work for eukaryotic organisms since most of their DNA is non-coding. In addition, bacterial-derived protein-fragment libraries such as these have no "inventory" i.e. because the sequences were randomly cloned it is not possible to say exactly what is contained within a given library other than by very deep sequencing.

[8] These practical limitations have led to significant inertia in mining a potentially rich alternative vein of directly relevant protein-fold structural diversity in target-identification and validation screens in human cells.

[9] Other screening approaches are described, for example, in WO 2001/86297. Here random short (40-mer and 20-mer) peptide phage display libraries are generated and used to find peptides that bind to a pre-selected target or a known, pre-identified consensus motif. This relies on existing disease targets being known/recognised and does not facilitate the identification of new targets. WO 2007/097923 discloses libraries, and means to produce such libraries, of peptide structures that are representative of the repertoire of protein structures existing in nature. However, such libraries are selected to comprise those peptides that are capable of folding or assuming their native conformations independently of artificial scaffolds or flanking sequences in the proteins from which they are derived.

[10] WO 2010/129310 describes libraries of nucleic acids encoding peptides from proteins comprising an entire natural proteome (or known bioactive peptides) that in each case are expressed and secreted to the outside of the cell. The use of such libraries to isolate biologically-active secreted peptides ("BASPs") is described therein, as well as how such libraries are constructed, starting from high-throughput oligonucleotide synthesis but without disclosure of the sequences synthesised or the peptides encoded in such libraries. Indeed, little information is described therein on the amino acid sequences or other particular (e.g. advantageous) features of the peptides encoded, or of the nucleic acids encoding such peptides, nor on the method by which such peptides are selected for inclusion in (or exclusion from) such libraries, or the design (and features, e.g., of the sequence) of the nucleic acids selected to be synthesised for such libraries. Little further information is provided on such important matters of library design (e.g. in-silico construction) in the corresponding scientific publication for this technology (Natarajan et al, 2014; PNAS 111:E474).

[11] There are also several known phage display libraries. WO 2015/095355 relates to detection of an antibody against a pathogen. It describes a phage display library comprising viral protein sequences. A related paper: Xu et al, 2015; Science 348, describes the VirScan technology and it is said to combine DNA microarray synthesis and bacteriophage display to create a uniform, synthetic representation of peptide epitopes comprising the human virome. An earlier publication by the same group, Larman et al, 2011, Nat Biotechnol 29:535, describes a similar approach but relates to a T7 "peptidome" phage display library which comprises peptides from a human genome (i.e. 36 amino acid peptides from approx. 24,000 unique ORFs from a human genome).

[12] Accordingly, it is one object of the present invention to provide a library which encodes protein fragments/peptides wherein such a library can be used in screening methods including but not limited to PPI screens. In other objects, the present invention provides alternative, improved, simpler, cheaper and/or integrated means or methods that address one or more of these or other problems. An object underlying the present invention is solved by the subject matter as disclosed or defined anywhere herein, for example by the subject matter of the attached claims.

[13] The figures show:

[14] **Figure 1:** depicts a screen for SEPs expressed in the HuPEX library that are able to overcome 6-thioguanine toxicity. Cells carrying a library of inserts that express SEPs are treated with 500nM 6-thioguanine for 6 days. Enrichment between 6-thioguanine treatment (n=3) and DMSO control (n=3) is shown.

[15] **Figure 2:** depicts a screen for SEPs across all three libraries (HuPEX, BugPEX, OmePEX) for SEPs able to selectively kill cells lacking the PTEN tumour suppressor gene.

[16] **Figure 3:** depicts the effect on MCF10A PTEN KO cells of a peptide (7-924) identified from the screen described in Example 4(2) expressed by pMOST25 compared to empty vector and a positive control (shRNA against NLK).

[17] **Figure 4:** depicts the experimental principle of the MNNG-induced parthanotos phenotypic screen of Examples 9 and 10.

[18] **Figure 5:** depicts the relative abundance of DNA sequences encoding for SEPs from the HuPEX library that are present in a control aliquot of HuPEX-expressing HeLa cells before (D0) treatment with 6.7uM MNNG, and in a treatment aliquot of such HuPEX-expressing HeLa cells after 8 days (D8) of such MNNG treatment. Peptides showing a significantly increased relative abundance at D8 are marked by triangles.

[19] **Figure 6:** depicts the relative abundance of DNA sequences encoding for SEPs from the BugPEX library that are present in a control aliquot of BugPEX expressing HeLa cells before (D0) treatment with 6.7uM MNNG, and in a treatment aliquot of such BugPEX expressing HeLa cells after 8 days (D8) of such MNNG treatment. Axes are as for Figure 5, and peptides showing a significantly increased relative abundance at D8 are marked by triangles.

[20] **Figure 7:** depicts the relative abundance of DNA sequences encoding for SEPs from the OmePEX library that are present in a control aliquot of OmePEX-expressing HeLa cells before (D0) treatment with 6.7uM MNNG, and in a treatment aliquot of such OmePEX-expressing HeLa cells after 8 days (D8) of such MNNG treatment. Axes are as for Figure 5, and peptides showing a significantly increased relative abundance at D8 are marked by triangles.

[21] **Figure 8 and Figure 9:** Phenotypic screen of PEX libraries for autophagy induction

[22] **Figure 8 (A-C):** HEK293FT Cells were engineered to stably express an GFP-LC3/RFP-LC3DG autophagy reporter (Kaizuka et al Molecular Cell 2016). Subsequently, autophagy reporter cells were infected with the pooled HuPEX (HPX), BugPEX (BPX) and OmePEX (OPX) libraries and selected on puromycin. After selection, cells enriched in the low GFP-LC3 gate, compared to unsorted controls, were flow-sorted and peptide sequences were amplified and sent to NGS analysis as described previously. The graphs (Figure 8A (HPX), Figure 8B (BPX) and Figure 8C (OPX)) show the population of selected hits (autophagy inducers) in the marked region compared to control.

[23] **Figure 9:** Hits were re-run individually in a flow-cytometry experiment after infection with lentivirus carrying either control sequences or putative hits. A selection of candidates is shown with BPX-497507 representing a strong and robust hit able to induce autophagy as measured by GFP-LC3 reduction. Torin1 (250nM) is shown as a positive control.

[24] The present invention, and particular non-limiting aspects and/or embodiments thereof, can be described in

more detail as follows:

5 [25] In a first aspect, the present invention provides a library of nucleic acids, each nucleic acid comprising a coding region of defined nucleic acid sequence encoding for a peptide having a length of between 25 and 110 amino acids, and having an amino acid sequence being a region of a sequence selected from the amino acid
10 sequence of a naturally occurring protein of one or more organisms; wherein the library comprises nucleic acids that encode for a plurality of at least about 10,000 (or 5,000) different such peptides, and wherein the amino acid sequence of each of at least 50 (or at least 25) of such peptides is a sequence region of the amino acid sequence of a different protein of a plurality of different such naturally occurring proteins (or wherein, for clarity, in respect of each of the different naturally occurring proteins, the library comprises one or more nucleic acids encoding for a
15 peptide having an amino acid sequence being a sequence region of the amino acid sequence of such naturally occurring protein.

[26] Suitably, a library in accordance with any aspect or embodiment of the invention comprises nucleic acids that encode a plurality of at least approximately 20,000, 50,000, 100,000, 200,000, 250,000, 300,000, 475,000 or 500,000 different such peptides. The library may also comprise nucleic acids that encode over 300,000 or 500,000
20 different such peptides. For example, in certain embodiments the library may comprise nucleic acids that encode for a plurality of at least 50,000 different such peptides, and wherein the amino acid sequence of each of at least 100 of such peptide is a sequence region of the amino acid sequence of at least 100 different naturally occurring proteins (or wherein, for clarity, in respect of each of the at least 100,000 different naturally occurring proteins, the library comprises one or more nucleic acids encoding for a peptide having an amino acid sequence being a sequence region
25 of the amino acid sequence of such naturally occurring protein); in particular of such embodiments the library may comprise nucleic acids that encode for a plurality of at least 100,000 different such peptides, and wherein the amino acid sequence of each of at least 150 of such peptide is a sequence region of the amino acid sequence of at least 150 different naturally occurring proteins (or wherein, in respect of each of the at least 150 different naturally occurring proteins, the library comprises one or more nucleic acids encoding for a peptide having an amino acid
30 sequence being a sequence region of the amino acid sequence of such naturally occurring protein). In another embodiment, the library may comprise nucleic acids that encode for a plurality of at least 10,000 different such peptides, and wherein the amino acid sequence of each of at least 1,000 of such peptides is a sequence region of the amino acid sequence of a different protein of such plurality of different naturally occurring proteins.

[27] In one embodiment, the library may comprise nucleic acids that encode for a plurality of at least 200,000
35 different such peptides, and wherein the amino acid sequence of each of at least 20,000 of such peptide is a sequence region of the amino acid sequence of at least 20,000 different naturally occurring proteins; in particular of such embodiments the library may comprise nucleic acids that encode for a plurality of at least 300,000 different such peptides, and wherein the amino acid sequence of each of at least 25,000 of such peptide is a sequence region of the amino acid sequence of at least 25,000 different naturally occurring proteins. In one embodiment, the nucleic acids are present in the mixture in an amount that is proportional to the complexity and size of the genome or transcriptome of the organism of interest. In other embodiments, the number of different nucleic acids in a library in accordance with the invention may depend on the desired screening application and on the number of sequences that may be workable in a particular application. This may include a consideration of whether the library is for use as a primary or secondary screen, for example.

40 [28] As used herein the term "different" in the context of the peptides encoded by the nucleic acids within the library of the invention means that any one peptide has at least one amino acid difference compared to any other peptide encoded within the library. In other words, each nucleic acid within the library encodes a unique peptide.

[29] In one embodiment, a "naturally occurring protein" is one which has a sequence found in a reference proteome. Examples of reference proteomes and how to use information from reference proteomes are described herein. As used herein the term "different" in the context of the naturally occurring proteins means that any one such protein has at least one amino acid difference compared to any other such protein. In other words, each nucleic acid within the library encodes a unique peptide. Suitably, "different" naturally occurring proteins have an amino acid sequence identity that is less than about 98%, 95% or 92% sequence identity, such as less than about 95% or 90% sequence identity. In one suitable embodiment, "different" naturally occurring proteins have different entry numbers (or other identifier) of a database, such as having different UniProt identifiers. For example, the cyclin-dependent kinases with the UniProt (www.uniprot.org) identifiers P24941 and P11802 (human CDK2 and CDK4, respectively) are, in such an embodiment, "different" naturally occurring proteins.

[30] Advantageously, each nucleic acid comprises a coding region of defined (or known) nucleic acid sequence encoding for a peptide. By "defined" (or "known") nucleic acid sequence is meant that the sequence of substantially all (e.g. each) nucleic acid sequence within the library is defined (or known). In particular, the library is non-random i.e. it does not represent a collection of random genomic sequences (which may or may not express peptide sequences), even if the genome sequence as a whole may have been determined (e.g. known), but the library has been designed, starting from protein sequences and, optionally filtered to generate a subset of nucleic acids encoding peptides with particular predicted features, in particular with specific and defined amino acid sequence. Thus, advantageously, the identity of substantially all (e.g. each) of the sequences in the library will be defined (or known) such that a library in accordance with the invention may have an inventory (that is, e.g., comprising or consisting of a pre-designated or pre-designed collection of individual members of defined (or known) sequences), even if may not be known which specific sequence is in which specific member of the library. This allows the sequences therein to be readily identified. Such libraries may be designed to have a desired complexity and/or to filter out undesired sequences, as described herein.

[31] In some embodiments the nucleic acids of the library are synthetic (e.g., they have been – at least initially – generated by chemical rather than biological processes). Thus, suitably, the library provides synthetic or non-natural nucleic acids (and/or comprises non-natural sequences of nucleic acids). Such nucleic acids are, suitably, designed according to any one of the methods as described herein and synthesised according to methods available to those skilled in the art, particularly those high volume/high throughput methods included those described elsewhere herein. Importantly, such synthesised nucleic acids comprise design features which distinguish them from naturally occurring nucleic acids. Such design features include, for example, use of codon frequency tables to generate the nucleic acids such that the sequence of nucleotides making up the codons within the nucleic acids encoding the peptides do not represent those codons which would be found at that position in the amino acid sequence of the naturally occurring protein. In addition, the nucleic acids for use in the libraries in accordance with the invention may comprise restriction sites which would not be present in the naturally occurring nucleic acid sequences and would generate peptide sequences comprising additional amino acids which would not occur in the naturally occurring protein sequence. Suitably, the nucleic acids (e.g., the sequences thereof) for use in the libraries in accordance with the invention may be generated using the design principles set out in the methods as described herein.

[32] In one embodiment of the library in accordance with the invention, the library provides multiple nucleic acid sequences encoding multiple peptides derived from any one protein. That is, in respect of each of the different naturally occurring proteins the library comprises a plurality of (i.e., more than one) nucleic acids encoding for a peptide having an amino acid sequence being a sequence region of the amino acid sequence of such naturally occurring protein. Accordingly, with this meaning of such embodiment, the first aspect of the present invention may alternatively be stated (e.g., for clarity, as indicated above) as relating to **a library of nucleic acids**, each nucleic

acid comprising a coding region of defined (or known) nucleic acid sequence encoding for a peptide having a length of between 25 and 110 amino acids, and having an amino acid sequence being a region of a sequence selected from the amino acid sequence of a naturally occurring protein of one or more organisms; wherein the library comprises nucleic acids that encode for a plurality of at least 10,000 different such peptides, and wherein in respect of each of the different naturally occurring proteins, the library comprises one or more nucleic acids encoding for a peptide having an amino acid sequence being a sequence region of the amino acid sequence of such naturally occurring protein.

[33] For example, in one such embodiment at least about 1% of the naturally occurring proteins a plurality of the nucleic acids encodes for different peptides from the amino acid sequences of such naturally occurring proteins. Suitably, in respect of at least about 5%, 10%, 25% or 50% of the naturally occurring proteins, a plurality of the nucleic acids encodes for different peptides from the amino acid sequences of such naturally occurring proteins. In other embodiments, a plurality of the nucleic acids encodes for different peptides from at least about 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100% of the naturally occurring proteins. In a particular embodiment, a plurality of the nucleic acids encodes for different peptides from the amino acid sequences of between about 90% and 100% of such naturally occurring proteins. Suitably a "different peptide" is one that has a different amino acid, such as differing by one or more (e.g. between about two and ten, five and 20, 15 and 40 or 30 and 50, or more than 50) amino acids.

[34] In another embodiment, the invention provides a library of (e.g. synthetic) nucleic acids, wherein the plurality of the nucleic acids encodes for different peptides, the amino acid sequences of which are sequence regions spaced along the amino acid sequence of the naturally occurring protein. Suitably this spacing is chosen so as to generate a library which encodes a workable number of peptides and may be varied according to the desired number of peptides represented by the library. By "workable number" is meant a suitable number which can be generated economically whilst providing a suitable number for use in a screening application or method of choice. For example, for primary cell-based selection screens, a "workable number" may be larger (say, 300,000, 500,000 or even millions) that for secondary screening or array-based or pull-down screening (using say, tens of thousands up to about 250,000). Suitably, an expression library of nucleic acids of the present invention may, suitably, be used at a complexity of between about 250,000 and 500,000 different peptides (such as about 300,000); and a library of peptides expressed therefrom may be used for solid-phase screening at a complexity of greater than 500,000, such as greater than 750,000 1,000,000, 1,500,000 or 2,000,000 (or more). In other embodiments, a "workable number" may be smaller than such numbers of peptides (such as, in those embodiments directed to the "focused" section of naturally occurring proteins described below). Suitably, in such embodiments, an expression library of nucleic acids of the present invention may be used at a complexity of between about 5,000 and 250,000 different peptides (such as at least about 10,000), for example between about 10,000 and 25,000, 20,000 and 50,000, 40,000 and 100,000 or 100,000 and 200,000 different peptides.

[35] Accordingly, in one embodiment, the sequence regions are spaced by a window of amino acids apart, or by multiples of such window, along the amino acid sequence of the naturally occurring protein wherein, the window is between 1, 5, 10, 15, 20, 25, 30, 35, 40 or 45 and about 55 amino acids; in particular wherein, the window is between about 2 and 40 amino acids, more particularly wherein, the window is between about 5 and about 20 amino acids; most particularly wherein the window of spacing is about 8, 10, 12 or 15 amino acids. In a suitable embodiment, the window used to space the sequence regions is the same number of amino acids (or multiples thereof) between each of the sequence regions along the amino acid sequence of the naturally occurring protein (and optionally, for each of the other naturally occurring proteins used to form the sequences of the peptides encoded by the nucleic acids of the library. In one further embodiment, the library may contain sequence regions

spaced by multiples of such windows as one or more of the intervening peptides/amino acid sequences may get dropped from the selection process in view of not conforming to one or more of the various filtering criteria.

[36] Suitably, the library is designed to be a complex library. By a "complex library" is meant one which has a multitude of different proteins and peptides represented therein, in particular one comprising sequence (or structurally) diverse peptides such as those defined from evolutionary diverse species. "Complexity" is therefore considered in terms of the numbers of proteins and peptides. Advantageously, the library of the present invention provides a much higher degree of complexity than those libraries available in the prior art. In one embodiment the library comprises (synthetic) nucleic acids encoding at least 5,000 different peptides from at least 5,000 different proteins. In another embodiment, the library of (synthetic) nucleic acids in accordance with the invention comprises nucleic acids encoding for at least 100,000 (or 20,000 or 50,000) different peptides from at least 10,000 (or 2,000 or 5,000, respectively) different naturally occurring proteins. It will be appreciated that the number of different peptides encoded by the library in accordance with the invention may be upwards of at least 10,000, 20,000 50,000 or 100,000 different peptides encoded by at least 50, 75, 100 or 150 10,000 different naturally occurring proteins; the size of the library may be over at least 50,000, 100,000 or 250,000 peptides, for example. The number of different peptides and the number of different naturally occurring proteins from which they are derived may be varied according to the specific application. Accordingly, and as described above, the library may encode for at least approximately 5,000, 10,000, 50,000, 100,000, 200,000, 250,000, 300,000, 475,000 or 500,000 different such peptides, over 300,000 or 500,000 different such peptides (or, e.g. suitably for focused libraries, over 10,000 or 50,000 different such peptides); suitably wherein on average two or more (such as about 5, 8, 10 or 15) peptides are derived from the amino acid sequence of the same naturally occurring protein.

[37] Given the complexity of certain embodiments of the library of the present invention - for example, those encoding for (or comprising) at least 10,000 (or 200,000 or 300,000) peptides from at least 1,000 or 10,000 (or 20,000 or 25,000, respectively) different naturally occurring proteins - the e.g. encoded peptides will in such embodiments, typically, represent those derived from a **diverse** selection of different naturally occurring proteins. Indeed, typically, the diverse selection of different naturally occurring proteins would (e.g. also or, optionally, only) comprise proteins that are non-secreted proteins and/or are not extracellular proteins, including those from a plurality of species. For example, the different naturally occurring proteins may comprise (e.g. also or, optionally, only) a set of proteins that include proteins other than (e.g. human and mouse) cytokines, chemokines, growth factors and their receptors (optionally, such other proteins as well as such cytokines, chemokines, growth factors and their receptors). In particular embodiments, the different naturally occurring proteins may comprise a set of proteins that include proteins that are cytoplasmic proteins (optionally, cytoplasmic protein as well as including non-cytoplasmic proteins such as also including secreted proteins and/or extracellular proteins).

[38] In additional or alternative embodiments, a library of the present invention encodes for (or comprises) peptides that are not (previously) known to be - e.g., are not pre-selected to be - bioactive peptides, and/or could putatively modulate cellular responses by interacting with cell surface receptors.

[39] In other embodiments of the library of the present invention - for example, those encoding for (or comprising) at least 50,000 (or 10,000 or 100,000) peptides from at least 100 (or 50 or 150, respectively) different naturally occurring proteins - the e.g. encoded peptides will in such embodiments, typically, represent those derived from a **focused** selection of different naturally occurring proteins. In such embodiment, the different naturally occurring proteins may be pre-selected from or a subset of a larger set of naturally occurring proteins (such as those from one or more reference proteomes) based on one more (e.g. pre-determined) criteria. In particular of such embodiments, all of the different naturally occurring proteins may fulfill such criteria, and each (e.g., all) of the (encoded) peptides of such a library is intended to have, or has, a sequence that is a sequence region of the amino

acid sequence of a naturally occurring protein that fulfills such (e.g. pre-determined) criteria. One non-limiting example of such criteria can be that the naturally occurring proteins are those being secreted proteins and/or extracellular proteins, such as that (e.g. all of) the naturally occurring proteins may be cytokines, chemokines, growth factors and their receptors. Alternatively, such criteria may comprise that the naturally occurring proteins are not secreted proteins and/or are not extracellular proteins, such as that (e.g. all of) the naturally occurring proteins may not be cytokines, chemokines, growth factors and their receptors.

[40] In alternative embodiments, criteria for (e.g., each and/or all of) the naturally occurring proteins from which the peptides encoded by (or comprised in) the library are derived, can include or consist of one or more of the following criteria; such protein/s is/are:

- 10 • From a subcellular compartment e.g. Cytoplasmic, nuclear, mitochondrial, cytoskeletal, or ribosomal
- One or more given enzymatic class. For example, a kinase, protease, esterase, or phosphatase;
- One or more given receptor type. For example, a G-coupled protein receptor or a nuclear hormone receptor;
- Membrane transport proteins and/or ion-channel proteins;
- Structural proteins;
- 15 • Transcription factors or DNA-binding proteins;
- DNA repair proteins. For example, a protein of the mismatch repair pathway;
- Involved in one or more (e.g. related or inter-related) cell-signaling/signal-transduction pathway. For example, the MAPK/ERK pathway, PI3K/Akt signaling, ErbB/HER signaling, mTOR signaling, NF-kappaB signaling or Jak/Stat (IL-6) receptor signaling;
- 20 • Interact (e.g., in-vivo, or as determined by laboratory procedures such as yeast 2-hybrid or affinity purification/mass spectrometry) with a given protein or at least one protein from a (e.g. functional) class of proteins. For example, interact with KRas or with a kinase (e.g., (ABL, BCL-ABL, SRC, KIT, PLK, CDKs, PLK, Aurora, MAPKs, JAKs, FLT3 or EGFR);
- Associated with a given disease (e.g. hits from genome-wide association studies GWAS), such as cancer. For example, as BRCA1 and/or BRCA2 is associated with breast cancer, or as JAK2 is associated myeloproliferative neoplasms (Stadler et al 2010, J Clin Oncol 28:4255);
- 25 • Hits from functional screens e.g. CRISPR, RNAi, gene-trapping, mutagenesis, cDNA screens, PROTEINi screens.

[41] In another embodiment, the invention provides a library of (e.g. synthetic) nucleic acids in accordance with the invention, wherein each nucleic acid encodes a different peptide. Suitably, the mean number of nucleic acids that encode a (e.g. different) peptide from the naturally occurring proteins is greater than 1; in particular between about 1.01 and 1.5 such nucleic acids (peptides) per such protein (such as about 1.02, 1.05, 1.1, 1.2, 1.3 or 1.4), or is at least about 10 (or 2 or 5) nucleic acids (peptides) per such protein, in particular wherein the mean is between about 5 and about 2,000 such nucleic acids (peptides) per such protein or is between about 5 and about 1,000 such nucleic acids (peptides) per such protein (or between about 100 and about 1,500, or between about 250 and about 1,000; 35 in particular between about 5 and about 100 or about 5 and about 50) such nucleic acids (peptides) per such protein. In some embodiments, the mean number of nucleic acids may be up to at least 500 nucleic acids, although it will be appreciated that the number of nucleic acids encoding different peptides for any particular protein will depend on the size of the protein as well as the size of the window by which the amino acid sequences are spaced. In a related embodiment, the invention provides a library of (e.g. synthetic) nucleic acids in accordance with the invention, wherein 95% of the naturally occurring proteins represented therein are represented by between about 2 40 and about 20 (such as between about 3 and about 35 or 40) peptide sequences.

[42] In another embodiment, there is provided a library of (e.g. synthetic) nucleic acids in accordance with the invention, wherein the amino acid sequence of the naturally occurring protein is one selected from the group of

amino acids sequences of proteins comprised in a reference proteome; in particular wherein the amino acid sequence of the naturally occurring protein is one selected from the group of amino acids sequences of (e.g., non-redundant) proteins comprised in a reference proteome.

5 [43] In another embodiment, the reference proteome is one or more of the reference proteomes selected from the group of reference proteomes listed in any of Table A and/or Table B herein, or any updated versions of the proteomes listed therein.

10 [44] In another embodiment, the amino acid sequences of the plurality of encoded peptides are sequence regions selected from amino acid sequences of naturally occurring proteins (or polypeptide chains or domains thereof) with a known three-dimensional structure; in particular wherein the naturally occurring protein (or polypeptide chain or domain thereof) is comprised in the Protein Data Bank (<https://www.wwpdb.org>), and optionally that has a Pfam annotation (<http://pfam.xfam.org>). In another embodiment, the sequence region selected from the amino acid sequence of the protein does not include an ambiguous amino acid of such amino acid sequence comprised in the reference proteome or the Protein Data Bank.

15 [45] Suitably, the organism or species from which the amino acid sequence of a naturally occurring protein is selected to generate the (synthetic) nucleic acid library in accordance with the invention is *Homo sapiens*.

20 [46] In another embodiment, the different naturally occurring proteins used to generate the library of (e.g. synthetic) nucleic acids in accordance with the invention are naturally occurring proteins of a plurality of different organisms or species. Suitably, a plurality of different species is selected from the group of (micro)organism species listed in Table A; in particular wherein the plurality of different species comprises at least 2, 3 or 5 (micro)organism species (such as at least about 10, 20, 25, or 50) across at least 2 (such as from at least about 3 or 5) of the phyla listed in Table A.

25 [47] In another embodiment, a plurality of different species is selected from the group of species listed in Table B, in particular wherein the plurality of different species comprises at least 2, 3, 5 or 10 (such as at least about 20, 50, 100, 200, 300 or over 400) species listed in Table B, across at least about 3, 5 or 5 (in particular, across at least 5 or 6) of the sections of such table described therein as: archaea, bacteria, fungi, invertebrates, plants, protozoa, mammals and non-mammalian vertebrates.

[48] Suitably, the plurality of different organisms or species is at least 10, 20, 50, 100 (in particular) or 250 different organisms or species. In some embodiments, the plurality may include up to about 20, 50, 100, 250 or 500 different organisms or species. Accordingly, a high diversity within the library may be achieved.

30 [49] In another embodiment, the different naturally occurring proteins used to generate the library of (e.g. synthetic) nucleic acids in accordance with the invention are naturally occurring proteins that are differentially expressed between two or more different cell populations (e.g. cell types) or tissue types. For example, the different naturally occurring proteins may be differentially expressed between a diseased and normal cell or tissue types; in particular wherein the different naturally occurring proteins are differentially expressed between human cancer cells and non-cancerous human cells, for example. In further embodiments, the different naturally occurring proteins are disease-specific; in particular wherein the different naturally occurring proteins are expressed by human cancer cells but not by non-cancerous human cells. In another embodiment, the different naturally occurring proteins may be differentially expressed between one cell population that has been infected by a pathogen or treated with a substance (such as a pathogenic substance or a drug) and a second cell population of the corresponding cell type that has not been so treated (or treated differently). For example, an inflammatory phenotype may be induced in a cell line by treatment with a pro-inflammatory substance such as phorbol myristate (PMA) and naturally occurring proteins that are differentially expressed in such treated population of cells compared to non-treated cells are considered. In another example differentially expressed naturally occurring proteins may be identified by comparison

35
40

of protein expression between immune cells that have become exhausted (T cell exhaustion) compared to proficient immune cells (e.g. T cells).

[50] In another embodiment, the library of (e.g. synthetic) nucleic acids in accordance with the invention is one in which the plurality of encoded peptides have diverse sequences; in particular wherein the plurality of encoded peptides differ in amino acid sequence from one another by at least 2 or 3 amino acids; in particular wherein the plurality of encoded peptides differ in amino acid sequence from one another by at least about 5, 8 or 10 amino acids. Suitably, the library of (synthetic) nucleic acids is one in which the diversity of the plurality of encoded peptides have a sequence similarity of less than about 90% or 80%; in particular have a sequence similarity of less than about 70%, 60% or 50%. Such sequence similarity may be determined using hierarchical clustering with CD-HIT (Fu et al 2012, Bioinformatics 28:3150; <http://weizhongli-lab.org/cd-hit/>). Suitable parameters for this are described in the Examples section herein.

[51] In another embodiment the peptides encoded by the library in accordance with the invention are predicted not to contain disordered regions; in particular wherein the prediction is as determined by SLIDER (Super-fast predictor of proteins with long intrinsically disordered regions; Peng et al, 2014; Proteins: Structure, Function and Bioinformatics 82:145; <http://biomine.cs.vcu.edu/servers/SLIDER>) and/or DISEMBL (Linding et al 2003, Structure 11:1453; <http://dis.embl.de>). Suitable parameters for this are described in the Examples section herein.

[52] In another embodiment, the peptides encoded by the library in accordance with the invention are predicted to have an isoelectric point (pI) of less than about 6 and greater than about 8. Suitable examples of using the "pI" function of "R Peptide package" are described in the Examples section, herein.

[53] In another embodiment, at least about 30%, 40% or 50% of the amino acid sequences of the peptides encoded by the library in accordance with the invention are identical to a peptide in a different organism; in particular between about 30% and 70% of the amino acid sequences of the peptides encoded by the library in accordance with the invention are identical to a peptide in a different organism.

[54] In a further embodiment, each nucleic acid within the library in accordance with the invention encodes for a peptide of the same length. It would be understood by the skilled person that this may include some limited variability. In other embodiments, the encoded peptide has a length of: between about 25 and about 100, 90, 85, 80, 75, 70, 65, 60, 55, 50 or 45; between about 30 and 110; between about 30 and about 100, 90, 85, 80, 75, 70, 65, 60, 55 or 50; between about 30 and about 100, 90, 85, 80, 75, 70, 65, 60 or 55; in particular wherein the encoded peptide has a length of between 30 and about 75; more particularly wherein the encoded peptide has a length of between about 35 and about 70 or between about 35 and about 50; and most particularly wherein the encoded peptide has a length of the encoded peptide that is between 35 and 60 amino acids, such as is 42, 43, 44, 45, 46, 47 or 48 amino acids in length. Suitably, the length of the peptides encoded by the library in accordance with the invention will be determined by practical considerations such as the maximum limit of the oligonucleotide synthesis technique used (and taking into account the additional features added, as described herein, for example). Suitably, in the library of (synthetic) nucleic acids in accordance with the invention, the coding region encoding for the peptide uses the human codon frequency table set forth in Table 1.1. However, alternative human codon frequency tables may be used or, and depending on the species of the expression system in which the nucleic acid library is intended to be expressed, codon frequency tables of other species may be used. In one embodiment, the most frequent human codon is used to encode amino acids of the peptide. Accordingly, at least a portion of the nucleic acid sequences will be not the natural genomic sequence. Such sequences may also be modified further. In further embodiments, an alternative human codon is used to encode one or more amino acids of the peptide.

[55] Suitably, "undesirable sequences or sub-sequences" are excluded from the library in accordance with the invention. Such "undesirable sequences or sub-sequences" may include, for example, those sequences made from

combinations of codons. Particular examples of such undesired sub-sequence include an internal Kozak sequence and/or a restriction enzyme site for a restriction enzyme intended to be used for the cloning of the resulting library, in each case generated from a combination of codons. Such undesirable sequences or sub-sequences can be avoided by using the next most commonly used (or another) codon in place of one or other of the codons in the combination.

5 Thus, suitably, in the library of (synthetic) nucleic acids in accordance with the invention, the coding region does not contain a combination of codons that forms an internal Kozak sequence. The person of ordinary skill will understand the term "Kozak sequence", and such meaning can include a sequence of nucleotide bases identified by the notation "(gcc)gccRccAUGG", where: (i) a lower-case letter denotes the most common base at a position where the base can nevertheless vary; (ii) upper-case letters indicate highly conserved bases, i.e. the "AUGG" sequence is constant or rarely, if ever, changes, with the exception being the IUPAC ambiguity code "R" which indicates that a purine (adenine or guanine) is always observed at this position (with adenine being believed to be more frequent); and (iii) the sequence in brackets "(gcc)" is of uncertain significance. In particular embodiments, a Kozak sequence is "CCATGG".

15 **[56]** In some embodiments of the library of nucleic acids of the invention, the following sequences are also undesirable sequences or subsequences: "GGATCC", "CTCGAG", "GGGGGG", "AAAAAA", "TTTTTT", "CCCCC", sequences that cause issues with oligonucleotide synthesis or sequencing or PCR amplification, hairpin sequences, an in-frame STOP codon (except at the terminus).

20 **[57]** In another embodiment, each nucleic acid in the library in accordance with the invention further comprises one or more nucleic acid sequences 5' and/or 3' of the coding region that include at least one restriction enzyme recognition sequence; optionally with a linker nucleic acid sequence between the coding region and the restriction enzyme recognition sequence(s). Suitably, the coding region does not contain a combination of codons that forms the restriction enzyme recognition sequence(s) that is(are) included in the 5' and/or 3' nucleic acid sequence(s). As described above, this can be avoided by using the next most commonly used codon in place of one or other of the codons in the combination.

25 **[58]** The nucleic acid library of the present invention (including also those embodiments where it is (e.g., initially) comprised by a synthetic nucleic acid library) may be amplified, propagated or otherwise maintained by non-chemical processes. For example, such library may be amplified by in-vitro enzymatic (e.g. biological) processes such as PCR or in-vitro transcription/translation; or may be replicated (and/or propagated/maintained) in-vivo processes such as being cloned into a vector which is replicated in a host cell (e.g., a bacterial or mammalian host cell).

30 **[59]** In another embodiment, each nucleic acid in the library in accordance with the invention further comprises additional sequences to enable amplification of the nucleic acid sequence (suitably, in each case not within the coding region that encodes for the peptide, such as 5' and/or 3' of the coding region and, optionally, the restriction enzyme recognition sequence(s)); in particular wherein such amplification is by PCR amplification.

35 **[60]** In another embodiment, each nucleic acid in the library in accordance with the invention is cloned into a vector. The term "vector" will be art recognized, and includes the meaning of a nucleic acid that can be used to propagate, produce, maintain or introduce a nucleic acid comprised within it in/into a cell, such as for expression of a peptide or polypeptide encoded by a sequence comprising said nucleic acid. One type of vector is a plasmid, which refers to a linear or circular double-stranded DNA molecule into which additional nucleic acid segments can be ligated. Another type of vector is a viral vector (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), wherein additional DNA segments can be introduced into the viral genome. Certain vectors are capable of autonomous replication in a cell, such as host cell, into which they are introduced (e.g., bacterial vectors comprising a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) integrate into the genome of a cell upon introduction into the cell and culturing under selective

pressure, and thereby are replicated along with the genome. A vector can be used to direct the expression of a chosen peptide or polynucleotide in a cell; in particular a peptide encoded by a nucleic acid comprised in the library of the present invention. Suitably, when used to express a peptide of the present invention in a eukaryotic cell (such as a mammalian cell), the vector is a lentiviral vector, or is retroviral vector.

5 [61] Suitably, each nucleic acid further comprises a start codon, a Kozak sequence, a stop codon and/or a nucleic acid sequence encoding a peptide tag, or wherein the vector comprises a start codon, a Kozak sequence, a stop codon and/or a nucleic acid sequence encoding a peptide tag operatively linked to the nucleic acid. Suitably, a peptide tag, if included, is one selected from the group consisting of: V4, FLAG, Strep/HA and GFP; in particular where the tag is a V4 tag or a FLAG tag.

10 [62] In one embodiment, each nucleic acid is suitable for or capable of expressing a polypeptide comprising the encoded peptide.

[63] In another embodiment, the polypeptide comprises the encoded peptide and further comprises an N-terminal methionine, one or more additional amino acids encoded by one or more restriction enzyme recognition sequence, and/or one or more peptide tags.

15 [64] In another embodiment, the individual members (or subsets of members) of the library of (e.g. synthetic) nucleic acids in accordance with the invention are in a pooled format (or form). Suitably, a library of the present invention in "pooled format" (or "pooled form") includes those where the individual members (or subset of members) thereof are in admixture with other members (or subsets); for example, a solution (or dried precipitate thereof) of such members contained in a single vessel, or a population of host cells containing recombinant vectors of the present invention.

20 [65] Suitably, the individual members (or sub-sets of members) in the library in accordance with the invention are spatially separated. A "spatially separated" library can be considered as a library in which a plurality of members (or sub-sets of members) of the library are physically separated, suitably in an ordered manner, from each other. Examples of a spatially separated library include those where individual members (or sub-sets of members) are comprised in individual wells of one or more microtitre plates, are arrayed on a solid surface or are bound (in an ordered manner) to a silicon wafer. In another embodiment, individual members (or sub-sets of members) in the library in accordance with the invention are each individually addressable; that is they can be retrieved (e.g. without undue searching or screening) from the library. Suitable methods for addressing or interrogating the library in accordance with the invention may include Next Generation Sequencing (NGS), PCR etc. Also, when the library is present in a spatially separated format (or form), the individual members (or sub-sets of members) may be "individually addressable" by knowing the spatial location of the applicable individual member (or sub-set). In either of these embodiments, use of a computer program, data file or database (such as those utilising a computer-readable medium or data-processing system, of the invention) can facilitate the retrieval of individual members (or sub-sets of members) that are comprised in an individually addressable library of the invention.

30 [66] Suitably, a library of the present invention is not generated from cDNA.

[67] **In another aspect, the invention provides a library of peptides** encoded by the library of (e.g. synthetic) nucleic acids in accordance with the invention. Suitably, the peptides are synthetic or recombinant. In one embodiment, the individual members (or subsets of members) thereof are in a pooled format (or form). In another embodiment, the individual members (or subsets of members) thereof are spatially separate and or individually addressable.

40 [68] In certain embodiments, any library of (synthetic) nucleic acids of the present invention may be comprised in an admixture with, may be an adjunct to (or otherwise combined with) or may be used together with, another library of nucleic acids that encode for peptides having a length of between about 25 and about 110 amino acids. For

example, such other library may be one described in co-pending application PCT/GB2016/054038 (the contents of which are incorporated herein by reference); in particular, such a library that encodes small Open Reading Frames (sORFs) such as that encodes at least 500, 1000, 1500 or about 2000 human sORFs.

[69] Accordingly: (1) any library of peptides of the present invention may also be comprised in an admixture with, may be an adjunct to (or otherwise combined with) or may be used together with, another library of peptides that is encoded by such other library of nucleic acids; and (2) the uses, methods and processes involving a nucleic acids or peptide library of the present invention, can also involve its use in admixture with, as an adjunct to (or otherwise combined with) or use together with such other library of nucleic acids or peptides (respectively). For example, Example 4 herein, describes a screen using both a "HuPEX" library of the present invention optionally with a human sORF library described in PCT/GB2016/054038.

[70] In some embodiments, the library of peptides is not a T7 display library, or is not a phage display library, or is not a display library. In some embodiments, the library is not a plurality of peptides derived from a plurality of (e.g. human) pathogens, such as from a plurality of viruses, bacteria or fungi that are, for example pathogenic to humans.

[71] In another aspect, there is provided a container or carrier comprising the library of (e.g. synthetic) nucleic acids in accordance with the invention and/or comprising the library of peptides in accordance with the invention. Suitable containers include a vessel, (such as an Eppendorf tube), a microtiter plate or a silicon carrier.

[72] In another aspect, the invention also relates to a method of identifying at least one binding partner of a binding interaction between a peptide comprised in a library of the present invention to a target (in particular, to a protein target), the method comprising the steps of

- Exposing a peptide library of the invention to the target under conditions permitting binding between the target and at least one peptide of the library; and
- Identifying the binding peptide or the bound target.

[73] In certain embodiments, the peptide library is exposed to the target by providing a library of nucleic acids of the invention in a form (e.g. in host cells) and under conditions such that the library of peptides is expressed by the library of nucleic acids.

[74] In another aspect (or embodiment of the above), the invention provides a method to identify the target and/or peptide; including, for example, elution of the peptide/nucleic acid, selection for cells expressing the peptide followed by e.g. PCR and sequencing identification. The target may be identified by, for example, pull-down mass-spectrometry. Such methods are described elsewhere herein, in PCT/GB2016/054038 and/or WO2013/116903.

[75] In a particular aspect, the invention also relates to a method of identifying a target protein that modulates a phenotype of a mammalian cell, said method comprising exposing a population of in vitro cultured mammalian cells capable of displaying said phenotype to a library of nucleic acids of the present invention (or a library of peptides of the present invention), identifying in said cell population an alteration in said phenotype following said exposure, selection of said cells undergoing the phenotypic change and identifying a peptide encoded by (or a peptide of) such library that alters the phenotype of the cell, providing said peptide and identifying the cellular protein that binds to said peptide, said cellular protein being a target protein that modulates the phenotype of the mammalian cell. Suitable methods and phenotypic screens are described, for example, in PCT/GB2016/054038, and the technical features described therein for such methods and screens, but using a library of the present invention, are incorporated herein by reference.

[76] In certain embodiments, such method includes a further step of identifying a compound that binds to said target protein and displaces or blocks binding of said peptide. Such further step, thereby identifying a compound which binds to a target protein and displaces or blocks binding of said peptide wherein the compound modulates the phenotype of a mammalian cell.

[77] Accordingly, in another particular aspect, the invention also relates to a method of identifying a compound which binds to a target protein and displaces or blocks binding of a peptide wherein the compound modulates a phenotype of a mammalian cell, said method comprising the steps:

- i. exposing a population of in vitro cultured mammalian cells capable of displaying said phenotype to a library of nucleic acids of the present invention or a library of peptides of the present invention;
- ii. identifying a cell in the population that displays an alteration in said phenotype following said exposure;
- iii. identifying a peptide encoded by (or of) such library that alters said phenotype of the cell;
- iv. identifying a cellular protein that binds to said peptide, said cellular protein being a target protein that modulates said phenotype of the mammalian cell;
- v. identifying a compound that binds to said target protein and displaces or blocks binding of said peptide.

[78] In one other aspect, the invention relates to a use of: (a) a library of nucleic acids according to the present invention; and/or (b) a library of peptides according to the present invention, **to identify a peptide that binds to a target** (in particular, a protein target). In certain embodiments, the identified peptide modulates a phenotype of a mammalian cell.

[79] In another other aspect, the invention relates to a use of: (a) a library of nucleic acids according to the present invention; and/or (b) a library of peptides according to the present invention, **to identify a target** (in particular, a protein target) that modulates a phenotype of a mammalian cell.

[80] In yet another other aspect, the invention relates to a use of: (a) a library of nucleic acids according to the present invention; and/or (b) a library of peptides according to the present invention, **to identify a compound** which binds to a target (in particular, a protein target) and, optionally, that displaces or blocks binding of a peptide to the target. In certain embodiments, the peptide and/or the compound modulates a phenotype of a mammalian cell.

[81] Suitably in such aspects, methodologies for performing phenotypic screens using libraries (or peptides) of the present invention can range from: (1) pathway-specific readouts that use heterologous reporters (for example GFP or Luciferase) to register either total protein levels, protein localisation or ultimate pathway activity at the level of gene transcription in live cells; (2) registering endogenous protein levels, or their localisation, using antibodies or other affinity reagents, or pathway-specific transcriptional outputs using qPCR or RNA-sequencing in fixed 'non-living' cells; (3) high-content, or 'holistic' based readouts in live cells that are capable of registering specific 'destination' phenotypic readouts of therapeutic relevance, such as differentiation, senescence and cell-death, all of which are coordinated and can be specifically modulated by a complex interplay of multiple cellular pathways. In some embodiments of the invention the assay readout method uses a GFP reporter, for instance as described in Kaizuka et al Molecular Cell 2016.

[82] In a specific aspect of the invention that covers 'holistic' phenotypic assays, Synthetic Lethality screening is of particular importance. Synthetic Lethality screening is an approach in which targets, for instance cancer targets, and candidate therapeutics are sought that can selectively impact tumour cells versus normal cells by exploiting unpredictable secondary points of weakness, which can occur in tumour cells as they heavily rewire their signalling pathways to support unrestrained cell proliferation. Such screens therefore must be performed in live cells and in an unbiased fashion by suppressing or modulating genes (using CRISPR), mRNA (using RNAi), or protein, or protein conformation (using Protein-i) in the cell and then determining whether a consistent negative impact on the overall growth or survival of a tumour cell type occurs; preferably one that harbours a specific genetic alteration(s) that occurs in a tumour situation versus a normal cell type. These direct 'holistic' cell-viability output based screens are performed using either large panels of genetically characterised tumour cells and normal cells to gain correlative information on tumour genotype- dependent responses, or more efficiently using specifically-engineered cell lines

that are isogenic for a chosen mutant versus normal genotype that exists in cancer cells versus normal cells, respectively.

[83] Suitably, in certain embodiments of uses and method of the present invention - that involve the inventive libraries described herein – related to phenotypes related to the modulation of cell-signalling pathways; in particular uses or method that involve the identification of peptides (e.g., from the libraries of the present invention) which modulate cell-signalling pathways and the identification of protein targets and surface sites on such proteins that participate in signal transduction and may be useful as drug targets to modulate cell-signalling pathways, in particular pathways which are active in cancer cells.

[84] A cell signalling pathway is, suitably, a series of interacting factors in a cell that transmit an intracellular signal within the cell in response to an extracellular stimulus at the cell surface and leading to changes in cell phenotype. Transmission of signals along a cell signalling pathway typically results in the activation of one or more transcription factors which alter gene expression. Preferred cell signalling pathways for uses, methods or screens related to the present invention display aberrant activity in disease models, for example activation, up-regulation or mis-regulation in diseased cells, such a cancer cells. For example, a pathway may be constitutively activated (i.e. permanently switched on) in a cancer cell, or inappropriately activated by an extracellular ligand, for example in an inflammatory cell in rheumatoid arthritis.

[85] A functional cell signalling pathway is typically considered as a pathway that is intact and capable of transmitting signals, if the pathway is switched on or activated, for example by an appropriate extracellular stimulus. An active cell-signalling pathway is typically considered a pathway that has been switched on, for example by an appropriate extracellular stimulus and is actively transmitting signals.

[86] Suitable cell signalling pathways include any signalling pathway that results in a transcriptional event in response to a signal received by a cell.

[87] Cell signalling pathways for investigation as described herein may include cell-signalling pathways that may be activated or altered in cancer cells, such as Ras/Raf, 20 Hedgehog, Fas, Wnt, Akt, ERK, TGF β , EGF, PDGF, Met, PI3K and Notch signalling pathways.

[88] **In yet another aspect, the invention relates to a computer-readable medium** (for example, one for use in – e.g., one specifically adapted for use in - a screening method described herein) having information stored thereon comprising: (a) the nucleic acid sequences comprised in a library of nucleic acids of the present invention; and/or (b) the amino acid sequences of the peptides encoded by said nucleic acids.

[89] **In a related aspect, the invention relates to a data-processing system** (for example, one for use in – e.g., one specifically adapted for use in - a screening method described herein) storing and/or processing information comprising: (a) the nucleic acid sequences comprised in a library of nucleic acids of the present invention; and/or (b) the amino acid sequences of the peptides encoded by said nucleic acids.

[90] In view of the above, it will be appreciated that the present invention also relates to the following items:

Item 1: A library of nucleic acids as described in the first aspect of the invention, wherein the species of the organism is Homo sapiens.

Item 2: The library of nucleic acids according to item 1, wherein the different naturally occurring proteins are naturally occurring proteins of a plurality of organisms of different species.

Item 3: The library of nucleic acids according to item 2, wherein a plurality of different species is selected from the group of (micro) organism species listed in Table A; in particular wherein the plurality of different species comprises at least 20 (micro)organism species across at least 5 of the phyla listed in Table A.

Item 4: The library of nucleic acids according to item 2, wherein a plurality of different species is selected from the

group of species listed in Table B, in particular wherein the plurality of different species comprises at least 50 of the species listed in Table B across at least 5 of the sections of Table B described therein as: archaea, bacteria, fungi, invertebrates, plants, protozoa, mammals and non-mammalian vertebrates.

Item 5: The library of nucleic acids according to any of items 2 to 4, wherein the plurality of different organisms is at least 100 different organisms.

Item 6: The library of nucleic acids according to any of items 1 to 4, wherein the different naturally occurring proteins are those that are differentially expressed between two or more different cell populations or tissue types.

Item 7: The library of nucleic acids according to item 6, wherein the different naturally occurring proteins are differentially expressed between a diseased and normal cell or tissue types; in particular wherein the different naturally occurring proteins are differentially expressed between human cancer cells and non-cancerous human cells.

Item 8: The library of nucleic acids according items 6 or 7, wherein the different naturally occurring proteins are disease-specific; in particular wherein the different naturally occurring proteins are expressed by human cancer cells but not by non-cancerous human cells.

Item 9: The library of nucleic acids according to any of items 1 to 8, wherein the plurality of encoded peptides have diverse sequences; in particular wherein the plurality of encoded peptides differ in amino acid sequence from one another by at least 2 or 3 amino acids; in particular wherein the plurality of encoded peptides differ in amino acid sequence from one another by at least about 5, 8 or 10 amino acids.

Item 10: The library of nucleic acids according to item 9, wherein the sequence similarity of the plurality of encoded peptides is less than about 80%.

Item 11: The library of nucleic acids according to any of items 1 to 10, wherein the peptides are predicted not to contain disordered regions.

Item 12: The library of nucleic acids according to any of items 1 to 11, wherein the peptides are predicted to have an isoelectric point (pI) of less than about 6 and greater than about 8.

Item 13: The library of nucleic acids according to any of items 1 to 12, wherein at least about 40% of the amino acid sequences of the peptide are identical to a peptide in a different organism.

Item 14: The library of nucleic acids according to any of items 1 to 13, wherein each nucleic acid encodes for a peptide of the same length.

Item 15: The library of nucleic acids according to any of items 1 to 14, wherein the encoded peptide has a length that is between 35 and 60 amino acids, such as is 42, 43, 44, 45, 46, 47 or 48 amino acids in length.

Item 16: The library of nucleic acids according to any of items 1 to 15, wherein the coding region encoding for the peptide uses the human codon frequency table set forth in Table 1.1.

Item 17: The library of nucleic acids of item 16, wherein the most frequent human codon is used to encode amino acids of the peptide.

Item 18: The library of nucleic acids of items 16 or 17, wherein an alternative human codon is used to encode one or more amino acids of the peptide.

Item 19: The library of nucleic acids according to any of items 1 to 18, wherein the coding region does not contain a combination of codons that forms an internal Kozak sequence; in particular wherein the Kozak sequence is ccatgg.

Item 20: The library of nucleic acids according to any of items 1 to 19, wherein each nucleic acid further comprises one or more nucleic acid sequences 5' and/or 3' of the coding region that include at least one restriction enzyme recognition sequence; optionally with a linker nucleic acid sequence between the coding region and the restriction

enzyme recognition sequence(s).

Item 21: The library of nucleic acids according to item 20, wherein the coding region does not contain a combination of codons that forms the restriction enzyme recognition sequence(s) that is (are) included in the 5' and/or 3' nucleic acid sequence(s).

- 5 Item 22: The library of nucleic acids according to any of items 1 to 21, wherein each nucleic acid further comprises additional sequences to enable amplification of the nucleic acid sequence; in particular wherein such amplification is by PCR amplification.

Item 23: The library of nucleic acids according to any of items 1 to 22, wherein each nucleic acid is cloned into a vector.

- 10 Item 24: The library of nucleic acids according to any of items 1 to 23, wherein each nucleic acid further comprises a start codon, a Kozak sequence, a stop codon and/or a nucleic acid sequence encoding a peptide tag, or wherein the vector comprises a start codon, a Kozak sequence, a stop codon and/or a nucleic acid sequence encoding a peptide tag operatively linked to the nucleic acid.

- Item 25: The library of nucleic acids of item 24, wherein each nucleic acid is suitable for or capable of expressing a polypeptide comprising the encoded peptide.

Item 26: The library of nucleic acids of item 25, wherein the polypeptide comprises the encoded peptide and further comprising an N-terminal methionine, one or more additional amino acids encoded by one or more restriction enzyme recognition sequence, and/or one or more peptide tags.

- Item 27: The library of nucleic acids according to any of items 1 to 26, wherein the individual members thereof are in a pooled format.

Item 28: The library of nucleic acids according to any of items 1 to 27, wherein the individual members thereof are spatially separated.

Item 29: A library of peptides, encoded by the library of nucleic acids as described in another aspect of the invention, suitably according to any of items 1 to 28.

- 25 Item 30: The library of peptides of item 29, wherein the peptides are synthetic or recombinant.

Item 31: The library of peptides of item 29, wherein the individual members thereof are in a pooled format.

Item 32: The library of peptides of item 29, wherein the individual members thereof are spatially separated.

Item 33: A container or carrier comprising the library of nucleic acids as described in another aspect of the invention, suitably according to any of items 1 to 33.

- 30 Item 33a: A container or carrier comprising the library of peptides as described in another aspect of the invention, suitably according to any of items 30 to 32.

Item 34: The container or carrier of item 33 or 33a that is a microliter plate or a silicon carrier.

Item 35: A method of identifying at least one binding partner of a binding interaction between a peptide that is: (a) comprised in a library as described in another aspect of the invention, suitably according to item 29 to 32, or (b) is expressed by a library of nucleic acids as described in another aspect of the invention, suitably according to any of items 1 to 28, to a protein target; the method comprising the steps of:

- exposing the invention to the protein target under conditions permitting binding between the target and at least one peptide of or expressed by the library; and
- identifying the binding peptide or the bound target.

- 40 Item 36: A use of a library of nucleic acids as described in another aspect of the invention, suitably according to any

of items 1 to 28, to identify a peptide that binds to a target protein; in particular wherein the peptide modulates a phenotype of a mammalian cell.

Item 37: A use of a library of nucleic acids as described in another aspect of the invention, suitably according to any of items 1 to 28, to identify a target protein that modulates a phenotype of a mammalian cell.

- 5 Item 38: A computer-readable medium having information stored thereon comprising: (a) the nucleic acid sequences comprised in a library of nucleic acids as described in another aspect of the invention, suitably according to any of items 1 to 28; and/or (b) the amino acid sequences of the peptides encoded by said nucleic acids.

- Item 39: A data-processing system storing and/or processing information comprising: (a) the nucleic acid sequences comprised in a library of nucleic acids as described in another aspect of the invention, suitably according to any of
10 items 1 to 28; and/or (b) the amino acid sequences of the peptides encoded by said nucleic acids.

[91] In view of the above, it will be appreciated that the present invention also relates to the following clauses:

- Clause 1: **A library of nucleic acids**, each nucleic acid comprising a coding region of defined nucleic acid sequence encoding for a peptide having a length of between 25 and 110 amino acids, and having an amino acid sequence being a region of a sequence selected from the amino acid sequence of a naturally occurring protein of one or more
15 organisms; wherein the library comprises nucleic acids that encode for a plurality of at least 10,000 different such peptides, and wherein the amino acid sequence of each of at least 50 of such peptides is a sequence region of the amino acid sequence of a different protein of a plurality of different such naturally occurring proteins.

Clause 2: The library of nucleic acids of clause 1, wherein each of the plurality of different naturally occurring proteins fulfils one or more pre-determined criteria.

- 20 Clause 3: The library of nucleic acids of clause 2, wherein each of the plurality of naturally occurring proteins is associated with a given disease, such as cancer.

Clause 4: The library of nucleic acids of clause 3, wherein the disease is breast cancer.

Clause 5: The library of nucleic acids of clause 2, wherein each of the plurality of naturally occurring proteins is a cytoplasmic protein.

- 25 Clause 6: The library of nucleic acids of clause 5, wherein each of the plurality of naturally occurring proteins is a cytoplasmic kinase.

Clause 7: The library of nucleic acids of clause 2, wherein each of the plurality of naturally occurring proteins interacts with a given protein or at least one protein from a (functional) class of proteins.

- Clause 8: The library of nucleic acids of clause 7, wherein each of the plurality of naturally occurring proteins
30 interacts with KRAs.

- Clause 9: The library of nucleic acids of any one of clauses 1 to 8, wherein the library comprises nucleic acids that encode for a plurality of at least 50,000 different such peptides, and wherein the amino acid sequence of each of at least 100 of such peptide is a sequence region of the amino acid sequence of at least 100 different naturally occurring proteins; in particular wherein the library comprises nucleic acids that encode for a plurality of at least
35 100,000 different such peptides, and wherein the amino acid sequence of each of at least 150 of such peptide is a sequence region of the amino acid sequence of at least 150 different naturally occurring proteins.

- Clause 10: The library of nucleic acids of any one of clauses 1 to 9, wherein the library comprises nucleic acids that encode for a plurality of at least 10,000 different such peptides, and wherein the amino acid sequence of each of at least 1,000 of such peptides is a sequence region of the amino acid sequence of a different protein of such plurality
40 of different naturally occurring proteins.

Clause 11: The library of nucleic acids of any one of clauses 1 to 10, wherein the library comprises nucleic acids that encode for a plurality of at least 200,000 different such peptides, and wherein the amino acid sequence of each of at least 20,000 of such peptide is a sequence region of the amino acid sequence of at least 20,000 different naturally

occurring proteins; in particular wherein the library comprises nucleic acids that encode for a plurality of at least 300,000 different such peptides, and wherein the amino acid sequence of each of at least 25,000 of such peptide is a sequence region of the amino acid sequence of at least 25,000 different naturally occurring proteins.

5 Clause 12: The library of nucleic acids of any one of clauses 1 or 11, wherein that in respect of at least about 1% of the naturally occurring proteins a plurality of the nucleic acids encodes for different peptides from the amino acid sequences of such naturally occurring proteins.

Clause 13: The library of nucleic acids of clause 12, wherein that in respect of at least about 50% of the naturally occurring proteins a plurality of the nucleic acids encodes for different peptides from the amino acid sequences of such naturally occurring proteins.

10 Clause 14: The library of nucleic acids of clause 13, wherein the plurality of the nucleic acids encodes for different peptides, and the amino acid sequences of which are sequence regions spaced along the amino acid sequence of the naturally occurring protein.

Clause 15: The library of nucleic acids of clause 14, wherein the sequence regions are spaced by a window of amino acids apart, or by multiples of such window, along the amino acid sequence of the naturally occurring protein
15 wherein, the window is between 1 and about 55 amino acids; in particular wherein the window is between about 5 and about 20 amino acids; most particularly wherein the window of spacing is about 8, 10, 12 or 15 amino acids.

Clause 16: The library of nucleic acids of any one of clauses 1 to 14 comprising nucleic acids encoding for at least 100,000 different peptides from at least 10,000 different naturally occurring proteins.

20 Clause 17: The library of nucleic acids of any one of clauses 1 to 16, wherein each nucleic acid encodes a different peptide.

Clause 18: The library of nucleic acids of any one of clauses 1 to 17, wherein the mean number of nucleic acids that encode a different peptide from the naturally occurring proteins is greater than 1; in particular between about 1.01 and 1.5 such nucleic acids (peptides) per such protein.

25 Clause 19: The library of nucleic acids of clause 18, wherein the mean number of nucleic acids that encode a different peptide from the naturally occurring proteins is at least about 5 such nucleic acids (peptides) per such protein, in particular wherein the mean is between about 5 and about 2,000 such nucleic acids (peptides) per such protein or is between about 5 and about 1,000 nucleic acids (peptides) per such protein.

Clause 20: The library of nucleic acids of clause 19, wherein the mean number of nucleic acids that encode a different peptide from the naturally occurring proteins is between about 100 and about 1,500 such nucleic acids (peptides) per such protein or is between about 250 and about 1,000 such nucleic acids (peptides) per such protein.
30

Clause 21: The library of nucleic acids of claim 20, wherein the mean number of nucleic acids that encode a different peptide from the naturally occurring proteins is between about 5 and about 100 such nucleic acids (peptides) per such protein or is between about 5 and about 50 such nucleic acids (peptides) per such protein.

35 Clause 22: The library of nucleic acids of any one of clauses 1 to 21, wherein the amino acid sequence of the naturally occurring protein is one selected from the group of amino acids sequences of non-redundant proteins comprised in a reference proteome, suitably, the reference proteome is one or more of the reference proteomes selected from the group of reference proteomes listed in Table A and/or Table B, or an updated version of such reference proteome.

40 Clause 23: The library of nucleic acids of any one of clauses 1 to 22, wherein the amino acid sequences of the plurality of encoded peptides are sequence regions selected from amino acid sequences of naturally occurring proteins (or polypeptide chains or domains thereof) with a known three-dimensional structure; in particular wherein the naturally occurring protein (or polypeptide chain or domain thereof) is comprised in the Protein Data Bank, and optionally that has a Pfam annotation.

Clause 24: The library of nucleic acids of any one of clauses 22 to 23, wherein the sequence region selected from the amino acid sequence of the protein does not include an ambiguous amino acid of such amino acid sequence comprised in the reference proteome or the Protein Data Bank.

Clause 25: **A library of peptides** encoded by the library of nucleic acids of any one of clauses 1 to 24.

- 5 [92] Suitably, the present invention is a library of nucleic acids, each nucleic acid comprising a coding region of defined nucleic acid sequence encoding for a peptide having a length of between 25 and 110 amino acids, and having an amino acid sequence being a region of a sequence selected from the amino acid sequence of a naturally occurring protein of one or more organisms; wherein the library comprises nucleic acids that encode for a plurality of at least 10,000 different such peptides, and wherein the amino acid sequence of each of at least 50 of such peptides is a sequence region of the amino acid sequence of a different protein of a plurality of different such naturally occurring proteins, and wherein the isoelectric point (pI) of each encoded amino acid sequence is greater than 7.4 or less than 6.0, and/or wherein the nucleic acid sequence does not contain the sequences "GGATCC", and/or "CTCGAG", and/or wherein the nucleic acid sequence does not contain the sequences "GGGGGG" and/or "AAAAAA" and/or "TTTTTT" and/or "CCCCCC", and/or wherein the nucleic acid sequence does not contain sequences that cause issues with oligonucleotide synthesis or sequencing or PCR amplification, and/or wherein the nucleic acid sequence does not contain a hairpin sequence, and/or wherein the nucleic acid sequence does not contain an in-frame STOP codon (except at the terminus) and/or wherein the nucleic acid sequence does not contain a KOZAK sequence (except at the start), i.e. does not contain an internal KOZAK sequence.

- 15 [93] The terms "of the [present] invention", "in accordance with the [present] invention", "according to the [present] invention" and the like, as used herein are intended to refer to all aspects and embodiments of the invention described and/or claimed herein.

- 20 [94] As used herein, the term "comprising" is to be construed as encompassing both "including" and "consisting of", both meanings being specifically intended, and hence individually disclosed embodiments in accordance with the present invention. Where used herein, "and/or" is to be taken as specific disclosure of each of the two specified features or components with or without the other. For example "A and/or B" is to be taken as specific disclosure of each of (i) A, (ii) B and (iii) A and B, just as if each is set out individually herein. In the context of the present invention, the terms "about" and "approximately" denote an interval of accuracy that the person skilled in the art will understand to still ensure the technical effect of the feature in question. The term typically indicates deviation from the indicated numerical value by $\pm 20\%$, $\pm 15\%$, $\pm 10\%$, and for example $\pm 5\%$. As will be appreciated by the person of ordinary skill, the specific such deviation for a numerical value for a given technical effect will depend on the nature of the technical effect. For example, a natural or biological technical effect may generally have a larger such deviation than one for a man-made or engineering technical effect. As will be appreciated by the person of ordinary skill, the specific such deviation for a numerical value for a given technical effect will depend on the nature of the technical effect. For example, a natural or biological technical effect may generally have a larger such deviation than one for a man-made or engineering technical effect. Where an indefinite or definite article is used when referring to a singular noun, e.g. "a", "an" or "the", this includes a plural of that noun unless something else is specifically stated.

- 30 [95] It is to be understood that application of the teachings of the present invention to a specific problem or environment, and the inclusion of variations of the present invention or additional features thereto (such as further aspects and embodiments), will be within the capabilities of one having ordinary skill in the art in light of the teachings contained herein.

- 35 [96] Unless context dictates otherwise, the descriptions and definitions of the features set out above are not limited to any particular aspect or embodiment of the invention and apply equally to all aspects and embodiments that are described.

[97] All references, patents, and publications cited herein are hereby incorporated by reference in their entirety.

Table A: Database source of reference proteomes of an evolutionary diverse set of microbes

NCBI ID	Species name	Domain	Phylum	Class
NC_000854.2	Aeropyrum pernix K1, Compl Gen	Archaea	Crenarchaeota	Thermoprotei
NC_002754.1	Sulfolobus solfataricus P2, Compl Gen	Archaea	Crenarchaeota	Thermoprotei
NC_000917.1	Archaeoglobus fulgidus DSM 4304, Compl Gen	Archaea	Euryarchaeota	Archaeoglobi
NC_006396.1	Haloarcula marismortui ATCC 43049 Chrom I, complete sequence	Archaea	Euryarchaeota	Halobacteria
NC_006397.1	Haloarcula marismortui ATCC 43049 Chrom II, complete sequence	Archaea	Euryarchaeota	Halobacteria
NC_006389.1	Haloarcula marismortui ATCC 43049 plasmid pNG100, complete sequence	Archaea	Euryarchaeota	Halobacteria
NC_006390.1	Haloarcula marismortui ATCC 43049 plasmid pNG200, complete sequence	Archaea	Euryarchaeota	Halobacteria
NC_006391.1	Haloarcula marismortui ATCC 43049 plasmid pNG300, complete sequence	Archaea	Euryarchaeota	Halobacteria
NC_006392.1	Haloarcula marismortui ATCC 43049 plasmid pNG400, complete sequence	Archaea	Euryarchaeota	Halobacteria
NC_006393.1	Haloarcula marismortui ATCC 43049 plasmid pNG500, complete sequence	Archaea	Euryarchaeota	Halobacteria
NC_006394.1	Haloarcula marismortui ATCC 43049 plasmid pNG600, complete sequence	Archaea	Euryarchaeota	Halobacteria
NC_006395.1	Haloarcula marismortui ATCC 43049 plasmid pNG700, complete sequence	Archaea	Euryarchaeota	Halobacteria
NC_001869.1	Halobacterium sp. NRC-1 plasmid pNRC100, complete sequence	Archaea	Euryarchaeota	Halobacteria
NC_002608.1	Halobacterium sp. NRC-1 plasmid pNRC200 Compl Gen	Archaea	Euryarchaeota	Halobacteria
NC_002607.1	Halobacterium sp. NRC-1, Compl Gen	Archaea	Euryarchaeota	Halobacteria
NC_001732.1	Methanocaldococcus jannaschii DSM 2661 plasmid pDSM2661_1, complete sequence	Archaea	Euryarchaeota	Methanococci
NC_000961.1	Pyrococcus horikoshii OT3 DNA, Compl Gen	Archaea	Euryarchaeota	Thermococci
NC_002689.2	Thermoplasma volcanium GSS1 DNA, Compl Gen	Archaea	Euryarchaeota	Thermoplasmata
NC_009525.1	Mycobacterium tuberculosis H37Ra, Compl Gen	Bacteria	Actinobacteria	Actinobacteria
NC_003155.5	Streptomyces avermitilis MA-4680 = NBRC 14893 DNA, Compl Gen	Bacteria	Actinobacteria	Actinobacteria
NC_004719.1	Streptomyces avermitilis MA-4680 = NBRC 14893 plasmid SAP1 DNA, complete sequence	Bacteria	Actinobacteria	Actinobacteria
NC_002935.2	Corynebacterium diphtheriae NCTC 13129, Compl Gen	Bacteria	Actinobacteria	Actinomycetales
NC_004663.1	Bacteroides thetaiotaomicron VPI-5482 Chrom, Compl Gen	Bacteria	Bacteroidetes	Bacteroidia
NC_004703.1	Bacteroides thetaiotaomicron VPI-5482 plasmid p5482, complete sequence	Bacteria	Bacteroidetes	Bacteroidia
NC_002932.3	Chlorobium tepidum TLS Chrom, Compl Gen	Bacteria	Chlorobi	Chlorobia
NC_001263.1	Deinococcus radiodurans R1 Chrom 1, complete sequence	Bacteria	Deinococcus-Thermus	Deinococci
NC_001264.1	Deinococcus radiodurans R1 Chrom 2, complete sequence	Bacteria	Deinococcus-Thermus	Deinococci
NC_000959.1	Deinococcus radiodurans R1 plasmid CP1, complete sequence	Bacteria	Deinococcus-Thermus	Deinococci
NC_000958.1	Deinococcus radiodurans R1 plasmid MP1, complete sequence	Bacteria	Deinococcus-Thermus	Deinococci
NC_001733.1	Methanocaldococcus jannaschii DSM 2661 plasmid pDSM2661_2, complete sequence	Bacteria	Euryarchaeota	Methanococci
NC_000909.1	Methanocaldococcus jannaschii DSM 2661, Compl Gen	Bacteria	Euryarchaeota	Methanococci

NC_005707.1	Bacillus cereus ATCC 10987 plasmid pBc10987, complete sequence	Bacteria	Firmicutes	Bacilli
NC_003909.8	Bacillus cereus ATCC 10987, Compl Gen	Bacteria	Firmicutes	Bacilli
NC_003210.1	Listeria monocytogenes EGD-e Chrom, Compl Gen	Bacteria	Firmicutes	Bacilli
NC_008226.1	Clostridioides difficile 630 plasmid pCD630, complete sequence	Bacteria	Firmicutes	Clostridia
NC_009089.1	Clostridioides difficile 630, Compl Gen	Bacteria	Firmicutes	Clostridia
NC_002696.2	Caulobacter crescentus CB15 Chrom, Compl Gen	Bacteria	Proteobacteria	Alpha Proteobacteria
NC_007493.1	Rhodobacter sphaeroides 2.4.1 Chrom 1, complete sequence	Bacteria	Proteobacteria	Alphaproteobacteria
NC_007494.1	Rhodobacter sphaeroides 2.4.1 Chrom 2, complete sequence	Bacteria	Proteobacteria	Alphaproteobacteria
NC_009007.1	Rhodobacter sphaeroides 2.4.1 plasmid A, partial sequence	Bacteria	Proteobacteria	Alphaproteobacteria
NC_007488.1	Rhodobacter sphaeroides 2.4.1 plasmid B, complete sequence	Bacteria	Proteobacteria	Alphaproteobacteria
NC_007489.1	Rhodobacter sphaeroides 2.4.1 plasmid C, complete sequence	Bacteria	Proteobacteria	Alphaproteobacteria
NC_007490.1	Rhodobacter sphaeroides 2.4.1 plasmid D, complete sequence	Bacteria	Proteobacteria	Alphaproteobacteria
NC_009008.1	Rhodobacter sphaeroides 2.4.1 plasmid E, partial sequence	Bacteria	Proteobacteria	Alphaproteobacteria
NC_008767.1	Neisseria meningitidis serogroup C FAM18 Compl Gen	Bacteria	Proteobacteria	Betaproteobacteria
NC_002937.3	Desulfovibrio vulgaris str. Hildenborough Chrom, Compl Gen	Bacteria	Proteobacteria	Deltaproteobacteria
NC_005863.1	Desulfovibrio vulgaris str. Hildenborough plasmid pDV, complete sequence	Bacteria	Proteobacteria	Deltaproteobacteria
NC_002939.5	Geobacter sulfurreducens PCA Chrom, Compl Gen	Bacteria	Proteobacteria	Deltaproteobacteria
NC_002163.1	Campylobacter jejuni subsp. jejuni NCTC 11168 = ATCC 700819 Chrom, Compl Gen	Bacteria	Proteobacteria	Epsilonproteobacteria
NC_000915.1	Helicobacter pylori 26695 Chrom, Compl Gen	Bacteria	Proteobacteria	Epsilonproteobacteria
NC_009085.1	Acinetobacter baumannii ATCC 17978 Chrom, Compl Gen	Bacteria	Proteobacteria	Gammaproteobacteria
NC_009083.1	Acinetobacter baumannii ATCC 17978 plasmid pAB1, complete sequence	Bacteria	Proteobacteria	Gammaproteobacteria
NC_009084.1	Acinetobacter baumannii ATCC 17978 plasmid pAB2, complete sequence	Bacteria	Proteobacteria	Gammaproteobacteria
NC_000907.1	Haemophilus influenzae Rd KW20 Chrom, Compl Gen	Bacteria	Proteobacteria	Gammaproteobacteria
NC_002942.5	Legionella pneumophila subsp. pneumophila str. Philadelphia 1 Chrom, Compl Gen	Bacteria	Proteobacteria	Gammaproteobacteria
NC_002516.2	Pseudomonas aeruginosa PAO1 Chrom, Compl Gen	Bacteria	Proteobacteria	Gammaproteobacteria
NC_002950.2	Porphyromonas gingivalis W83, Compl Gen	Bacteria	Proteobacteria	Porphyromonas gingivalis
NC_001318.1	Borrelia burgdorferi B31 Chrom, Compl Gen	Bacteria	Spirochaetes	Spirochaetales
NC_001903.1	Borrelia burgdorferi B31 plasmid cp26, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_000948.1	Borrelia burgdorferi B31 plasmid cp32-1, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_000949.1	Borrelia burgdorferi B31 plasmid cp32-3, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_000950.1	Borrelia burgdorferi B31 plasmid cp32-4, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_000951.1	Borrelia burgdorferi B31 plasmid cp32-6, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_000952.1	Borrelia burgdorferi B31 plasmid cp32-7, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_000953.1	Borrelia burgdorferi B31 plasmid cp32-8, complete	Bacteria	Spirochaetes	Spirochaetales

NC_000954.1	sequence Borrelia burgdorferi B31 plasmid cp32-9, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_001849.2	Borrelia burgdorferi B31 plasmid lp17, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_001851.2	Borrelia burgdorferi B31 plasmid lp28-1, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_001852.1	Borrelia burgdorferi B31 plasmid lp28-2, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_001853.1	Borrelia burgdorferi B31 plasmid lp28-3, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_001855.1	Borrelia burgdorferi B31 plasmid lp36, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_001857.2	Borrelia burgdorferi B31 plasmid lp54, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_000956.1	Borrelia burgdorferi B31 plasmid lp56, complete sequence	Bacteria	Spirochaetes	Spirochaetales
NC_000853.1	Thermotoga maritima MSB8 Chrom, Compl Gen	Bacteria	Thermotogae	Thermotogales

Table B: Database source of reference proteomes of an evolutionary diverse set of species

RefSeq assembly accession ID	Species name	Assembly level	ftp path (ftp://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/...)
Archaea:			
GCF_000213215.1	Acidianus hospitalis W1	Compl Gen	000/213/215/GCF_000213215.1_ASM21321v1
GCF_000144915.1	Acidilobus saccharovorans 345-15	Compl Gen	000/144/915/GCF_000144915.1_ASM14491v1
GCF_000025665.1	Aciduliprofundum boonei T469	Compl Gen	000/025/665/GCF_000025665.1_ASM2566v1
GCF_000327505.1	Aciduliprofundum sp. MAR08-339	Compl Gen	000/327/505/GCF_000327505.1_ASM32750v1
GCF_000008665.1	Archaeoglobus fulgidus DSM 4304	Compl Gen	000/008/665/GCF_000008665.1_ASM866v1
GCF_000025285.1	Archaeoglobus profundus DSM 5631	Compl Gen	000/025/285/GCF_000025285.1_ASM2528v1
GCF_000385565.1	Archaeoglobus sulfatocalidus PM70-1	Compl Gen	000/385/565/GCF_000385565.1_ASM38556v1
GCF_000194625.1	Archaeoglobus veneficus SNP6	Compl Gen	000/194/625/GCF_000194625.1_ASM19462v1
GCF_000317795.1	Caldisphaera lagunensis DSM 15908	Compl Gen	000/317/795/GCF_000317795.1_ASM31779v1
GCF_000018305.1	Caldvirga maquilensis IC-167	Compl Gen	000/018/305/GCF_000018305.1_ASM1830v1
GCF_000019605.1	Candidatus Korarchaeum cryptofilum OPF8	Compl Gen	000/019/605/GCF_000019605.1_ASM1960v1
GCF_000404225.1	Candidatus Methanomassiliicoccus intestinalis Isoire-Mx1	Compl Gen	000/404/225/GCF_000404225.1_ASM40422v1
GCF_000300255.2	Candidatus Methanomethylophilus alvus Mx1201	Compl Gen	000/300/255/GCF_000300255.2_ASM30025v2
GCF_000800805.1	Candidatus Methanoplasma termitum	Compl Gen	000/800/805/GCF_000800805.1_ASM80080v1
GCF_000812185.1	Candidatus Nitrosopelagicus brevis	Compl Gen	000/812/185/GCF_000812185.1_ASM81218v1
GCF_000299395.1	Candidatus Nitrosopumilus sp. AR2	Compl Gen	000/299/395/GCF_000299395.1_ASM29939v1
GCF_000955905.1	Candidatus Nitrosotenuis cloacae	Compl Gen	000/955/905/GCF_000955905.1_ASM95590v3
GCF_000231015.2	Desulfurococcus fermentans DSM 16532	Compl Gen	000/231/015/GCF_000231015.2_ASM23101v3
GCF_000020905.1	Desulfurococcus	Compl Gen	000/020/905/GCF_000020905.1_ASM2090v1

	kamchatkensis 1221n		
GCF_000186365.1	Desulfurococcus mucosus DSM 2162	Compl Gen	000/186/365/GCF_000186365.1_ASM18636v1
GCF_000025505.1	Ferroglobus placidus DSM 10642	Compl Gen	000/025/505/GCF_000025505.1_ASM2550v1
GCF_000152265.2	Ferroplasma acidarmanus fer1	Compl Gen	000/152/265/GCF_000152265.2_ASM15226v2
GCF_000789255.1	Geoglobus acetivorans	Compl Gen	000/789/255/GCF_000789255.1_ASM78925v1
GCF_001006045.1	Geoglobus ahangari	Compl Gen	001/006/045/GCF_001006045.1_ASM100604v1
GCF_000196895.1	Halalkalicoccus jeotgali B3	Compl Gen	000/196/895/GCF_000196895.1_ASM19689v1
GCF_001305655.1	Halanaeroarchaeum sulfurireducens	Compl Gen	001/305/655/GCF_001305655.1_ASM130565v1
GCF_000223905.1	Haloarcula hispanica ATCC 33960	Compl Gen	000/223/905/GCF_000223905.1_ASM22390v1
GCF_001488575.1	Halobacterium hubeiense	Compl Gen	001/488/575/GCF_001488575.1_Halobacterium_hubeiense_JI20-1
GCF_000069025.1	Halobacterium salinarum R1	Compl Gen	000/069/025/GCF_000069025.1_ASM6902v1
GCF_000230955.2	Halobacterium sp. DL1	Compl Gen	000/230/955/GCF_000230955.2_ASM23095v3
GCF_000306765.2	Haloferax mediterranei ATCC 33500	Compl Gen	000/306/765/GCF_000306765.2_ASM30676v2
GCF_000025685.1	Haloferax volcanii DS2	Compl Gen	000/025/685/GCF_000025685.1_ASM2568v1
GCF_000172995.2	Halogeometricum borinquense DSM 11551	Compl Gen	000/172/995/GCF_000172995.2_ASM17299v2
GCF_000023965.1	Halomicrobium mukohataei DSM 12286	Compl Gen	000/023/965/GCF_000023965.1_ASM2396v1
GCF_000224475.1	halophilic archaeon DL31	Compl Gen	000/224/475/GCF_000224475.1_ASM22447v1
GCF_000217715.1	Halopiger xanaduensis SH-6	Compl Gen	000/217/715/GCF_000217715.1_ASM21771v1
GCF_000237865.1	Haloquadratum walsbyi C23	Compl Gen	000/237/865/GCF_000237865.1_ASM23786v1
GCF_000470655.1	Halorhabdus tiamatea SARL4B	Compl Gen	000/470/655/GCF_000470655.1_HATI1
GCF_000023945.1	Halorhabdus utahensis DSM 12940	Compl Gen	000/023/945/GCF_000023945.1_ASM2394v1
GCF_000022205.1	Halorubrum lacusprofundi ATCC 49239	Compl Gen	000/022/205/GCF_000022205.1_ASM2220v1
GCF_000517625.1	Halostagnicola larsenii XH-48	Compl Gen	000/517/625/GCF_000517625.1_ASM51762v1
GCF_000025325.1	Haloterrigena turkmenica DSM 5511	Compl Gen	000/025/325/GCF_000025325.1_ASM2532v1
GCF_000015145.1	Hyperthermus butylicus DSM 5456	Compl Gen	000/015/145/GCF_000015145.1_ASM1514v1
GCF_000017945.1	Ignicoccus hospitalis KIN4/I	Compl Gen	000/017/945/GCF_000017945.1_ASM1794v1
GCF_000204925.1	Metallosphaera cuprina Ar-4	Compl Gen	000/204/925/GCF_000204925.1_ASM20492v1
GCF_000016605.1	Metallosphaera sedula DSM 5348	Compl Gen	000/016/605/GCF_000016605.1_ASM1660v1
GCF_000762265.1	Methanobacterium formicicum	Compl Gen	000/762/265/GCF_000762265.1_ASM76226v1
GCF_000191585.1	Methanobacterium lacus	Compl Gen	000/191/585/GCF_000191585.1_ASM19158v1
GCF_000214725.1	Methanobacterium paludis	Compl Gen	000/214/725/GCF_000214725.1_ASM21472v1
GCF_000499765.1	Methanobacterium sp. MB1	Compl Gen	000/499/765/GCF_000499765.1_Methanobacterium
GCF_001477655.1	Methanobrevibacter millerae	Compl Gen	001/477/655/GCF_001477655.1_ASM147765v1
GCF_000024185.1	Methanobrevibacter ruminantium M1	Compl Gen	000/024/185/GCF_000024185.1_ASM2418v1
GCF_000016525.1	Methanobrevibacter smithii ATCC 35061	Compl Gen	000/016/525/GCF_000016525.1_ASM1652v1
GCF_000739065.1	Methanocaldococcus bathoardescens	Compl Gen	000/739/065/GCF_000739065.1_ASM73906v1
GCF_000023985.1	Methanocaldococcus fervens AG86	Compl Gen	000/023/985/GCF_000023985.1_ASM2398v1

GCF_000092305.1	Methanocaldococcus infernus ME	Compl Gen	000/092/305/GCF_000092305.1_ASM9230v1
GCF_000025525.1	Methanocaldococcus sp. FS406-22	Compl Gen	000/025/525/GCF_000025525.1_ASM2552v1
GCF_000024625.1	Methanocaldococcus vulcanius M7	Compl Gen	000/024/625/GCF_000024625.1_ASM2462v1
GCF_000063445.1	Methanocella arvoryzae MRE50	Compl Gen	000/063/445/GCF_000063445.1_ASM6344v1
GCF_000251105.1	Methanocella conradii HZ254	Compl Gen	000/251/105/GCF_000251105.1_ASM25110v1
GCF_000013725.1	Methanococcoides burtonii DSM 6242	Compl Gen	000/013/725/GCF_000013725.1_ASM1372v1
GCF_000970325.1	Methanococcoides methylutens MM1	Compl Gen	000/970/325/GCF_000970325.1_ASM97032v1
GCF_000017185.1	Methanococcus aeolicus Nankai-3	Compl Gen	000/017/185/GCF_000017185.1_ASM1718v1
GCF_000017225.1	Methanococcus maripaludis C7	Compl Gen	000/017/225/GCF_000017225.1_ASM1722v1
GCF_000220645.1	Methanococcus maripaludis X1	Compl Gen	000/220/645/GCF_000220645.1_ASM22064v1
GCF_000017165.1	Methanococcus vannielii SB	Compl Gen	000/017/165/GCF_000017165.1_ASM1716v1
GCF_000006175.1	Methanococcus voltae A3	Compl Gen	000/006/175/GCF_000006175.1_ASM617v2
GCF_000015765.1	Methanocorpusculum labreanum Z	Compl Gen	000/015/765/GCF_000015765.1_ASM1576v1
GCF_000304355.2	Methanoculleus bourgensis MS2	Compl Gen	000/304/355/GCF_000304355.2_Mb_MS2
GCF_000015825.1	Methanoculleus marisnigri JR1	Compl Gen	000/015/825/GCF_000015825.1_ASM1582v1
GCF_000196655.1	Methanohalobium evestigatum Z-7303	Compl Gen	000/196/655/GCF_000196655.1_ASM19665v1
GCF_000025865.1	Methanohalophilus mahii DSM 5219	Compl Gen	000/025/865/GCF_000025865.1_ASM2586v1
GCF_000147875.1	Methanolacinia petrolearia DSM 11571	Compl Gen	000/147/875/GCF_000147875.1_ASM14787v1
GCF_000306725.1	Methanobolus psychrophilus R15	Compl Gen	000/306/725/GCF_000306725.1_ASM30672v1
GCF_000328665.1	Methanomethylovorans hollandica DSM 15978	Compl Gen	000/328/665/GCF_000328665.1_ASM32866v1
GCF_000017625.1	Methanoregula boonei 6A8	Compl Gen	000/017/625/GCF_000017625.1_ASM1762v1
GCF_000327485.1	Methanoregula formicica SMSP	Compl Gen	000/327/485/GCF_000327485.1_ASM32748v1
GCF_000204415.1	Methanoseta concilii GP6	Compl Gen	000/204/415/GCF_000204415.1_ASM20441v1
GCF_000235565.1	Methanoseta harundinacea 6Ac	Compl Gen	000/235/565/GCF_000235565.1_ASM23556v1
GCF_000217995.1	Methanosalsum zhilinae DSM 4017	Compl Gen	000/217/995/GCF_000217995.1_ASM21799v1
GCF_000007345.1	Methanosarcina acetivorans C2A	Compl Gen	000/007/345/GCF_000007345.1_ASM734v1
GCF_000970305.1	Methanosarcina barkeri 3	Compl Gen	000/970/305/GCF_000970305.1_ASM97030v1
GCF_000970025.1	Methanosarcina barkeri MS	Compl Gen	000/970/025/GCF_000970025.1_ASM97002v1
GCF_000970285.1	Methanosarcina horonobensis HB-1 = JCM 15518	Compl Gen	000/970/285/GCF_000970285.1_ASM97028v1
GCF_000970265.1	Methanosarcina lacustris Z-7289	Compl Gen	000/970/265/GCF_000970265.1_ASM97026v1
GCF_000970205.1	Methanosarcina mazei S-6	Compl Gen	000/970/205/GCF_000970205.1_ASM97020v1
GCF_000970085.1	Methanosarcina siciliae T4/M	Compl Gen	000/970/085/GCF_000970085.1_ASM97008v1
GCF_000970045.1	Methanosarcina sp. MTP4	Compl Gen	000/970/045/GCF_000970045.1_ASM97004v1
GCF_000969965.1	Methanosarcina sp. WWM596	Compl Gen	000/969/965/GCF_000969965.1_ASM96996v1
GCF_000969885.1	Methanosarcina thermophila TM-1	Compl Gen	000/969/885/GCF_000969885.1_ASM96988v1

GCF_000969905.1	Methanosarcina vacuolata Z-761	Compl Gen	000/969/905/GCF_000969905.1_ASM96990v1
GCF_000012545.1	Methanosphaera stadtmanae DSM 3091	Compl Gen	000/012/545/GCF_000012545.1_ASM1254v1
GCF_000021965.1	Methanosphaerula palustris E1-9c	Compl Gen	000/021/965/GCF_000021965.1_ASM2196v1
GCF_000013445.1	Methanospirillum hungatei JF-1	Compl Gen	000/013/445/GCF_000013445.1_ASM1344v1
GCF_000145295.1	Methanothermobacter marburgensis str. Marburg	Compl Gen	000/145/295/GCF_000145295.1_ASM14529v1
GCF_000008645.1	Methanothermobacter thermautotrophicus str. Delta H	Compl Gen	000/008/645/GCF_000008645.1_ASM864v1
GCF_000179575.2	Methanothermococcus okinawensis IH1	Compl Gen	000/179/575/GCF_000179575.2_ASM17957v2
GCF_000166095.1	Methanothermus fervidus DSM 2088	Compl Gen	000/166/095/GCF_000166095.1_ASM16609v1
GCF_000214415.1	Methanotorris igneus Kol 5	Compl Gen	000/214/415/GCF_000214415.1_ASM21441v1
GCF_000025625.1	Natrialba magadii ATCC 43099	Compl Gen	000/025/625/GCF_000025625.1_ASM2562v1
GCF_000230735.2	Natrinema pellirubrum DSM 15624	Compl Gen	000/230/735/GCF_000230735.2_ASM23073v3
GCF_000230715.2	Natronobacterium gregoryi SP2	Compl Gen	000/230/715/GCF_000230715.2_ASM23071v3
GCF_000328685.1	Natronococcus occultus SP4	Compl Gen	000/328/685/GCF_000328685.1_ASM32868v1
GCF_000591055.1	Natronomonas moolapensis 8.8.11	Compl Gen	000/591/055/GCF_000591055.1_ASM59105v1
GCF_000026045.1	Natronomonas pharaonis DSM 2160	Compl Gen	000/026/045/GCF_000026045.1_ASM2604v1
GCF_000725425.1	Palaeococcus pacificus DY20341	Compl Gen	000/725/425/GCF_000725425.1_ASM72542v1
GCF_000008265.1	Picrophilus torridus DSM 9790	Compl Gen	000/008/265/GCF_000008265.1_ASM826v1
GCF_000015205.1	Pyrobaculum islandicum DSM 4184	Compl Gen	000/015/205/GCF_000015205.1_ASM1520v1
GCF_000195935.2	Pyrococcus abyssi GE5	Compl Gen	000/195/935/GCF_000195935.2_ASM19593v1
GCF_000007305.1	Pyrococcus furiosus DSM 3638	Compl Gen	000/007/305/GCF_000007305.1_ASM730v1
GCF_000011105.1	Pyrococcus horikoshii OT3	Compl Gen	000/011/105/GCF_000011105.1_ASM1110v1
GCF_000211475.1	Pyrococcus sp. NA2	Compl Gen	000/211/475/GCF_000211475.1_ASM21147v1
GCF_000263735.1	Pyrococcus sp. ST04	Compl Gen	000/263/735/GCF_000263735.1_ASM26373v1
GCF_000215995.1	Pyrococcus yayanosii CH1	Compl Gen	000/215/995/GCF_000215995.1_ASM21599v1
GCF_001412615.1	Pyrodictium delaneyi	Compl Gen	001/412/615/GCF_001412615.1_ASM141261v1
GCF_000223395.1	Pyrolobus fumarii 1A	Compl Gen	000/223/395/GCF_000223395.1_ASM22339v1
GCF_000403645.1	Salinarchaeum sp. Harcht-Bsk1	Compl Gen	000/403/645/GCF_000403645.1_ASM40364v1
GCF_000092465.1	Staphylothermus hellenicus DSM 12710	Compl Gen	000/092/465/GCF_000092465.1_ASM9246v1
GCF_000015945.1	Staphylothermus marinus F1	Compl Gen	000/015/945/GCF_000015945.1_ASM1594v1
GCF_000012285.1	Sulfolobus acidocaldarius DSM 639	Compl Gen	000/012/285/GCF_000012285.1_ASM1228v1
GCF_000022385.1	Sulfolobus islandicus L.S.2.15	Compl Gen	000/022/385/GCF_000022385.1_ASM2238v1
GCF_000007005.1	Sulfolobus solfataricus P2	Compl Gen	000/007/005/GCF_000007005.1_ASM700v1
GCF_000011205.1	Sulfolobus tokodaii str. 7	Compl Gen	000/011/205/GCF_000011205.1_ASM1120v1
GCF_000151105.2	Thermococcus barophilus MP	Compl Gen	000/151/105/GCF_000151105.2_ASM15110v2
GCF_000265525.1	Thermococcus cleftensis	Compl Gen	000/265/525/GCF_000265525.1_ASM26552v1
GCF_000769655.1	Thermococcus eurythermalis	Compl Gen	000/769/655/GCF_000769655.1_ASM76965v1

GCF_000022365.1	<i>Thermococcus</i> <i>gammatolerans</i> EJ3	Compl Gen	000/022/365/GCF_000022365.1_ASM2236v1
GCF_000009965.1	<i>Thermococcus kodakarensis</i> KOD1	Compl Gen	000/009/965/GCF_000009965.1_ASM996v1
GCF_000246985.2	<i>Thermococcus litoralis</i> DSM 5473	Compl Gen	000/246/985/GCF_000246985.2_ASM24698v3
GCF_000585495.1	<i>Thermococcus nautili</i>	Compl Gen	000/585/495/GCF_000585495.1_ASM58549v1
GCF_000018365.1	<i>Thermococcus onnurineus</i> NA1	Compl Gen	000/018/365/GCF_000018365.1_ASM1836v1
GCF_000517445.1	<i>Thermococcus paralvinellae</i>	Compl Gen	000/517/445/GCF_000517445.1_ASM51744v1
GCF_000022545.1	<i>Thermococcus sibiricus</i> MM 739	Compl Gen	000/022/545/GCF_000022545.1_ASM2254v1
GCF_000151205.2	<i>Thermococcus</i> sp. AM4	Compl Gen	000/151/205/GCF_000151205.2_ASM15120v2
GCF_000813245.1	<i>Thermophilum</i> <i>carboxydiphosphus</i> 1505	Compl Gen	000/813/245/GCF_000813245.1_ASM81324v1
GCF_000015225.1	<i>Thermophilum pendens</i> Hrk 5	Compl Gen	000/015/225/GCF_000015225.1_ASM1522v1
GCF_000993805.1	<i>Thermophilum</i> sp. 1807-2	Compl Gen	000/993/805/GCF_000993805.1_ASM99380v1
GCF_000264495.1	<i>Thermogladius cellulolyticus</i> 1633	Compl Gen	000/264/495/GCF_000264495.1_ASM26449v1
GCF_000195915.1	<i>Thermoplasma acidophilum</i> DSM 1728	Compl Gen	000/195/915/GCF_000195915.1_ASM19591v1
GCF_000011185.1	<i>Thermoplasma volcanium</i> GSS1	Compl Gen	000/011/185/GCF_000011185.1_ASM1118v1
GCF_000350305.1	<i>Thermoplasmatales</i> archaeon BRNA1	Compl Gen	000/350/305/GCF_000350305.1_ASM35030v1
GCF_000253055.1	<i>Thermoproteus tenax</i> Kra 1	Compl Gen	000/253/055/GCF_000253055.1_ASM25305v1
GCF_000193375.1	<i>Thermoproteus uzoniensis</i> 768-20	Compl Gen	000/193/375/GCF_000193375.1_ASM19337v1
GCF_000092185.1	<i>Thermosphaera aggregans</i> DSM 11486	Compl Gen	000/092/185/GCF_000092185.1_ASM9218v1
GCF_000148385.1	<i>Vulcanisaeta distributa</i> DSM 14429	Compl Gen	000/148/385/GCF_000148385.1_ASM14838v1
GCF_000190315.1	<i>Vulcanisaeta moutnovskia</i> 768-28	Compl Gen	000/190/315/GCF_000190315.1_ASM19031v1
Bacteria:			
GCF_000020965.1	<i>Dictyoglomus thermophilum</i> H-6-12	Compl Gen	000/020/965/GCF_000020965.1_ASM2096v1
GCF_001676785.2	<i>Brachyspira hyodysenteriae</i> ATCC 27164	Compl Gen	001/676/785/GCF_001676785.2_ASM167678v2
GCF_000017685.1	<i>Leptospira biflexa</i> serovar Patoc strain 'Patoc 1 (Paris)'	Compl Gen	000/017/685/GCF_000017685.1_ASM1768v1
GCF_001399775.1	<i>Thermus aquaticus</i> Y51MC23	Compl Gen	001/399/775/GCF_001399775.1_ASM139977v1
GCF_000219605.1	<i>Pseudomonas stutzeri</i>	Compl Gen	000/219/605/GCF_000219605.1_ASM21960v1
GCF_000021685.1	<i>Thermomicrobium roseum</i> DSM 5159	Compl Gen	000/021/685/GCF_000021685.1_ASM2168v1
GCF_000024405.1	<i>Seibaldella termitidis</i> ATCC 33386	Compl Gen	000/024/405/GCF_000024405.1_ASM2440v1
GCF_000146505.1	<i>Fibrobacter succinogenes</i> subsp. <i>succinogenes</i> S85	Compl Gen	000/146/505/GCF_000146505.1_ASM14650v1
GCF_000158275.2	<i>Fusobacterium nucleatum</i> subsp. <i>animalis</i> 7_1	Compl Gen	000/158/275/GCF_000158275.2_ASM15827v2
GCF_000025305.1	<i>Acidaminococcus fermentans</i> DSM 20731	Compl Gen	000/025/305/GCF_000025305.1_ASM2530v1
GCF_000010785.1	<i>Hydrogenobacter</i> <i>thermophilus</i> TK-6	Compl Gen	000/010/785/GCF_000010785.1_ASM1078v1
GCF_000284095.1	<i>Selenomonas ruminantium</i> subsp. <i>lactilytica</i> TAM6421	Compl Gen	000/284/095/GCF_000284095.1_ASM28409v1
GCF_000317695.1	<i>Anabaena cylindrica</i> PCC 7122	Compl Gen	000/317/695/GCF_000317695.1_ASM31769v1
GCF_000179635.2	<i>Ruminococcus albus</i> 7 = DSM 20455	Compl Gen	000/179/635/GCF_000179635.2_ASM17963v2

GCF_001543175.1	<i>Aerococcus urinae</i>	Compl Gen	001/543/175/GCF_001543175.1_ASM154317v1
GCF_000953275.1	<i>Clostridioides difficile</i>	Compl Gen	000/953/275/GCF_000953275.1_CD630DERM
GCF_000145615.1	<i>Thermoanaerobacterium thermosaccharolyticum</i> DSM 571	Compl Gen	000/145/615/GCF_000145615.1_ASM14561v1
GCF_000013105.1	<i>Moorella thermoacetica</i> ATCC 39073	Compl Gen	000/013/105/GCF_000013105.1_ASM1310v1
GCF_000270085.1	<i>Erysipelothrix rhusiopathiae</i> str. Fujisawa	Compl Gen	000/270/085/GCF_000270085.1_ASM27008v1
GCF_000612055.1	<i>Trueperella pyogenes</i>	Compl Gen	000/612/055/GCF_000612055.1_ASM61205v1
GCF_001729525.1	<i>Brevibacterium linens</i>	Compl Gen	001/729/525/GCF_001729525.1_ASM172952v1
GCF_000734015.1	<i>Thermodesulfobacterium commune</i> DSM 2178	Compl Gen	000/734/015/GCF_000734015.1_ASM73401v1
GCF_000058485.1	<i>Frankia alni</i> ACN14a	Compl Gen	000/058/485/GCF_000058485.1_ASM5848v1
GCF_000092645.1	<i>Thermobispora bispora</i> DSM 43833	Compl Gen	000/092/645/GCF_000092645.1_ASM9264v1
GCF_000024385.1	<i>Thermomonospora curvata</i> DSM 43183	Compl Gen	000/024/385/GCF_000024385.1_ASM2438v1
GCF_000024985.1	<i>Sphaerobacter thermophilus</i> DSM 20745	Compl Gen	000/024/985/GCF_000024985.1_ASM2498v1
GCF_000269985.1	<i>Kitasatospora setae</i> KM-6054	Compl Gen	000/269/985/GCF_000269985.1_ASM26998v1
GCF_000008305.1	<i>Mesoplasma florum</i> L1	Compl Gen	000/008/305/GCF_000008305.1_ASM830v1
GCF_000021285.1	<i>Thermosiphon africanus</i> TCF52B	Compl Gen	000/021/285/GCF_000021285.1_ASM2128v1
GCF_000025205.1	<i>Gardnerella vaginalis</i> 409-05	Compl Gen	000/025/205/GCF_000025205.1_ASM2520v1
GCF_000009905.1	<i>Symbiobacterium thermophilum</i> IAM 14863	Compl Gen	000/009/905/GCF_000009905.1_ASM990v1
GCF_000015025.1	<i>Acidothermus cellulolyticus</i> 11B	Compl Gen	000/015/025/GCF_000015025.1_ASM1502v1
GCF_000144695.1	<i>Acetohalobium arabaticum</i> DSM 5501	Compl Gen	000/144/695/GCF_000144695.1_ASM14469v1
GCF_000024945.1	<i>Veillonella parvula</i> DSM 2008	Compl Gen	000/024/945/GCF_000024945.1_ASM2494v1
GCF_000022325.1	<i>Caldicellulosiruptor bescii</i> DSM 6725	Compl Gen	000/022/325/GCF_000022325.1_ASM2232v1
GCF_000020485.1	<i>Halothermothrix orenii</i> H 168	Compl Gen	000/020/485/GCF_000020485.1_ASM2048v1
GCF_000175575.2	<i>Acidithiobacillus caldus</i> ATCC 51756	Compl Gen	000/175/575/GCF_000175575.2_ASM17557v2
GCF_000022565.1	<i>Acidobacterium capsulatum</i> ATCC 51196	Compl Gen	000/022/565/GCF_000022565.1_ASM2256v1
GCF_000247605.1	<i>Acetobacterium woodii</i> DSM 1030	Compl Gen	000/247/605/GCF_000247605.1_ASM24760v1
GCF_000019165.1	<i>Heliobacterium modesticaldum</i> Ice1	Compl Gen	000/019/165/GCF_000019165.1_ASM1916v1
GCF_000020945.1	<i>Coprothermobacter proteolyticus</i> DSM 5265	Compl Gen	000/020/945/GCF_000020945.1_ASM2094v1
GCF_000619905.2	<i>Nitrosospora briensis</i> C-128	Compl Gen	000/619/905/GCF_000619905.2_ASM61990v2
GCF_000012745.1	<i>Thiobacillus denitrificans</i> ATCC 25259	Compl Gen	000/012/745/GCF_000012745.1_ASM1274v1
GCF_000661895.1	<i>Rubrobacter radiotolerans</i>	Compl Gen	000/661/895/GCF_000661895.1_ASM66189v1
GCF_000328625.1	<i>Halobacteroides halobius</i> DSM 5150	Compl Gen	000/328/625/GCF_000328625.1_ASM32862v1
GCF_000024605.1	<i>Ammonifex degensii</i> KC4	Compl Gen	000/024/605/GCF_000024605.1_ASM2460v1
GCF_000145035.1	<i>Butyrivibrio proteoclasticus</i> B316	Compl Gen	000/145/035/GCF_000145035.1_ASM14503v1
GCF_000092125.1	<i>Meiothermus silvanus</i> DSM 9946	Compl Gen	000/092/125/GCF_000092125.1_ASM9212v1
GCF_000024365.1	<i>Nakamurella multipartita</i> DSM 44233	Compl Gen	000/024/365/GCF_000024365.1_ASM2436v1
GCF_000023265.1	<i>Acidimicrobium ferrooxidans</i>	Compl Gen	000/023/265/GCF_000023265.1_ASM2326v1

	DSM 10331		
GCF_000317125.1	<i>Chroococcidiopsis thermalis</i> PCC 7203	Compl Gen	000/317/125/GCF_000317125.1_ASM31712v1
GCF_000317025.1	<i>Pleurocapsa</i> sp. PCC 7327	Compl Gen	000/317/025/GCF_000317025.1_ASM31702v1
GCF_000816145.1	<i>Pseudothermotoga hypogea</i> DSM 11164 = NBRC 106472	Compl Gen	000/816/145/GCF_000816145.1_ASM81614v1
GCF_000024205.1	<i>Desulfotomaculum acetoxidans</i> DSM 771	Compl Gen	000/024/205/GCF_000024205.1_ASM2420v1
GCF_000011905.1	<i>Dehalococcoides mccartyi</i> 195	Compl Gen	000/011/905/GCF_000011905.1_ASM1190v1
GCF_000008625.1	<i>Aquifex aeolicus</i> VF5	Compl Gen	000/008/625/GCF_000008625.1_ASM862v1
GCF_000191045.1	<i>Desulfurobacterium thermolithotrophum</i> DSM 11699	Compl Gen	000/191/045/GCF_000191045.1_ASM19104v1
GCF_000222305.1	<i>Borrelia</i> <i>bissetii</i> DN127	Compl Gen	000/222/305/GCF_000222305.1_ASM22230v1
GCF_000018605.1	<i>Petrogla mobilis</i> SJ95	Compl Gen	000/018/605/GCF_000018605.1_ASM1860v1
GCF_000184705.1	<i>Thermaerobacter marianensis</i> DSM 12885	Compl Gen	000/184/705/GCF_000184705.1_ASM18470v1
GCF_000025885.1	<i>Aminobacterium colombiense</i> DSM 12261	Compl Gen	000/025/885/GCF_000025885.1_ASM2588v1
GCF_000317065.1	<i>Pseudanabaena</i> sp. PCC 7367	Compl Gen	000/317/065/GCF_000317065.1_ASM31706v1
GCF_000237205.1	<i>Simkania negevensis</i> Z	Compl Gen	000/237/205/GCF_000237205.1_ASM23720v1
GCF_000023885.1	<i>Slackia heliotrinireducens</i> DSM 20476	Compl Gen	000/023/885/GCF_000023885.1_ASM2388v1
GCF_000305935.1	<i>Thermacetogenium phaeum</i> DSM 12270	Compl Gen	000/305/935/GCF_000305935.1_ASM30593v1
GCF_000092405.1	<i>Syntrophothermus lipocalidus</i> DSM 12680	Compl Gen	000/092/405/GCF_000092405.1_ASM9240v1
GCF_000317575.1	<i>Staniera cyanosphaera</i> PCC 7437	Compl Gen	000/317/575/GCF_000317575.1_ASM31757v1
GCF_000328705.1	<i>Saccharothrix espanaensis</i> DSM 44229	Compl Gen	000/328/705/GCF_000328705.1_ASM32870v1
GCF_000025645.1	<i>Thermoanaerobacter italicus</i> Ab9	Compl Gen	000/025/645/GCF_000025645.1_ASM2564v1
GCF_000014965.1	<i>Syntrophobacter fumaroxidans</i> MPOB	Compl Gen	000/014/965/GCF_000014965.1_ASM1496v1
GCF_000017805.1	<i>Roseiflexus castenholzii</i> DSM 13941	Compl Gen	000/017/805/GCF_000017805.1_ASM1780v1
GCF_000012865.1	<i>Carboxydotherrmus hydrogenoformans</i> Z-2901	Compl Gen	000/012/865/GCF_000012865.1_ASM1286v1
GCF_000017305.1	<i>Kineococcus radiotolerans</i> SRS30216 = ATCC BAA-149	Compl Gen	000/017/305/GCF_000017305.1_ASM1730v1
GCF_000281175.1	<i>Caldilinea aerophila</i> DSM 14535 = NBRC 104270	Compl Gen	000/281/175/GCF_000281175.1_ASM28117v1
GCF_000025605.1	<i>Thermocrinis albus</i> DSM 14484	Compl Gen	000/025/605/GCF_000025605.1_ASM2560v1
GCF_000021945.1	<i>Chloroflexus aggregans</i> DSM 9485	Compl Gen	000/021/945/GCF_000021945.1_ASM2194v1
GCF_000025005.1	<i>Thermobaculum terrenum</i> ATCC BAA-798	Compl Gen	000/025/005/GCF_000025005.1_ASM2500v1
GCF_000199675.1	<i>Anaerolinea thermophila</i> UNI-1	Compl Gen	000/199/675/GCF_000199675.1_ASM19967v1
GCF_000021025.1	<i>Laribacter hongkongensis</i> HLHK9	Compl Gen	000/021/025/GCF_000021025.1_ASM2102v1
GCF_000217795.1	<i>Thermodesulfatator indicus</i> DSM 15286	Compl Gen	000/217/795/GCF_000217795.1_ASM21779v1
GCF_000010305.1	<i>Gemmatimonas aurantiaca</i> T-27	Compl Gen	000/010/305/GCF_000010305.1_ASM1030v1
GCF_000299235.1	<i>Leptospirillum ferriphilum</i> ML-04	Compl Gen	000/299/235/GCF_000299235.1_ASM29923v1
GCF_000024345.1	<i>Kribbella flavida</i> DSM 17836	Compl Gen	000/024/345/GCF_000024345.1_ASM2434v1

GCF_000212395.1	<i>Thermodesulfobium narugense</i> DSM 14796	Compl Gen	000/212/395/GCF_000212395.1_ASM21239v1
GCF_000195335.1	<i>Marinithermus hydrothermalis</i> DSM 14884	Compl Gen	000/195/335/GCF_000195335.1_ASM19533v1
GCF_000754265.1	<i>Candidatus Baumannia cicadellinicola</i>	Compl Gen	000/754/265/GCF_000754265.1_ASM75426v1
GCF_000183745.1	<i>Oceanithermus profundus</i> DSM 14977	Compl Gen	000/183/745/GCF_000183745.1_ASM18374v1
GCF_000025265.1	<i>Conexibacter woesei</i> DSM 14684	Compl Gen	000/025/265/GCF_000025265.1_ASM2526v1
GCF_000194605.1	<i>Fluviicola taffensis</i> DSM 16823	Compl Gen	000/194/605/GCF_000194605.1_ASM19460v1
GCF_000494755.1	<i>Actinoplanes friuliensis</i> DSM 7358	Compl Gen	000/494/755/GCF_000494755.1_ASM49475v1
GCF_000016985.1	<i>Alkaliphilus metalliredigens</i> QYMF	Compl Gen	000/016/985/GCF_000016985.1_ASM1698v1
GCF_000185805.1	<i>Thermovibrio ammonificans</i> HB-1	Compl Gen	000/185/805/GCF_000185805.1_ASM18580v1
GCF_000020225.1	<i>Akkermansia muciniphila</i> ATCC BAA-835	Compl Gen	000/020/225/GCF_000020225.1_ASM2022v1
GCF_000317495.1	<i>Crinalium epipsammum</i> PCC 9333	Compl Gen	000/317/495/GCF_000317495.1_ASM31749v1
GCF_000021725.1	<i>Nautilia profundicola</i> AmH	Compl Gen	000/021/725/GCF_000021725.1_ASM2172v1
GCF_000213255.1	<i>Mahella australiensis</i> 50-1 BON	Compl Gen	000/213/255/GCF_000213255.1_ASM21325v1
GCF_000020505.1	<i>Chlorobaculum parvum</i> NCIB 8327	Compl Gen	000/020/505/GCF_000020505.1_ASM2050v1
GCF_000024545.1	<i>Streckbrandtia nassauensis</i> DSM 44728	Compl Gen	000/024/545/GCF_000024545.1_ASM2454v1
GCF_000144645.1	<i>Thermosediminibacter oceani</i> DSM 16646	Compl Gen	000/144/645/GCF_000144645.1_ASM14464v1
GCF_001688625.1	<i>Flavonifractor plautii</i>	Compl Gen	001/688/625/GCF_001688625.1_ASM168862v1
GCF_000024025.1	<i>Catenulispora acidiphila</i> DSM 44928	Compl Gen	000/024/025/GCF_000024025.1_ASM2402v1
GCF_000021565.1	<i>Persephonella marina</i> EX-H1	Compl Gen	000/021/565/GCF_000021565.1_ASM2156v1
GCF_000021545.1	<i>Sulfurihydrogenibium azorense</i> Az-Fu1	Compl Gen	000/021/545/GCF_000021545.1_ASM2154v1
GCF_000024165.1	<i>Candidatus Accumulibacter phosphatis</i> clade IIA str. UW-1	Compl Gen	000/024/165/GCF_000024165.1_ASM2416v1
GCF_000092425.1	<i>Truepera radiovictrix</i> DSM 17093	Compl Gen	000/092/425/GCF_000092425.1_ASM9242v1
GCF_000019905.1	<i>Exiguobacterium sibiricum</i> 255-15	Compl Gen	000/019/905/GCF_000019905.1_ASM1990v1
GCF_000283575.1	<i>Oscillibacter valericigenes</i> Sjm18-20	Compl Gen	000/283/575/GCF_000283575.1_ASM28357v1
GCF_000023705.1	<i>Methylothermobacter mobilis</i> JLW8	Compl Gen	000/023/705/GCF_000023705.1_ASM2370v1
GCF_000271665.2	<i>Pelosinus fermentans</i> JBW45	Compl Gen	000/271/665/GCF_000271665.2_ASM27166v2
GCF_000145255.1	<i>Gallionella capsiferiformans</i> ES-2	Compl Gen	000/145/255/GCF_000145255.1_ASM14525v1
GCF_000020005.1	<i>Natronaerobius thermophilus</i> JW/NM-WN-LF	Compl Gen	000/020/005/GCF_000020005.1_ASM2000v1
GCF_000020785.1	<i>Hydrogenobaculum</i> sp. Y04AAS1	Compl Gen	000/020/785/GCF_000020785.1_ASM2078v1
GCF_000265425.1	<i>Terriglobus roseus</i> DSM 18391	Compl Gen	000/265/425/GCF_000265425.1_ASM26542v1
GCF_000020145.1	<i>Elusimicrobium minutum</i> Pei191	Compl Gen	000/020/145/GCF_000020145.1_ASM2014v1
GCF_000226295.1	<i>Chloracidobacterium thermophilum</i> B	Compl Gen	000/226/295/GCF_000226295.1_ASM22629v1
GCF_000348785.1	<i>Ilumatobacter coccineus</i> YM16-304	Compl Gen	000/348/785/GCF_000348785.1_ASM34878v1

GCF_000306785.1	Modestobacter marinus	Compl Gen	000/306/785/GCF_000306785.1_ASM30678v1
GCF_000183405.1	Calditerrivibrio nitroreducens DSM 19672	Compl Gen	000/183/405/GCF_000183405.1_ASM18340v1
GCF_000328765.2	Tepidanaerobacter acetatoxydans Re1	Compl Gen	000/328/765/GCF_000328765.2_ASM32876v2
GCF_000284115.1	Phycisphaera mikurensis NBRC 102666	Compl Gen	000/284/115/GCF_000284115.1_ASM28411v1
GCF_000025965.1	Aromatoleum aromaticum EbN1	Compl Gen	000/025/965/GCF_000025965.1_ASM2596v1
GCF_000143165.1	Dehalogenimonas lykanthroporepellens BL-DC- 9	Compl Gen	000/143/165/GCF_000143165.1_ASM14316v1
GCF_000297055.2	Sulfuricella denitrificans skB26	Compl Gen	000/297/055/GCF_000297055.2_ASM29705v2
GCF_000166415.1	Halanaerobium hydrogeniformans	Compl Gen	000/166/415/GCF_000166415.1_ASM16641v1
GCF_000284335.1	Caldisericum exile AZM16c01	Compl Gen	000/284/335/GCF_000284335.1_ASM28433v1
GCF_000007085.1	Caldanaerobacter subterraneus subsp. tengcongensis MB4	Compl Gen	000/007/085/GCF_000007085.1_ASM708v1
GCF_000177635.2	Desulfurispirillum indicum S5	Compl Gen	000/177/635/GCF_000177635.2_ASM17763v2
GCF_000178955.2	Granulicella mallensis MP5ACTX8	Compl Gen	000/178/955/GCF_000178955.2_ASM17895v2
GCF_000192745.1	Polymorphum gilvum SL003B-26A1	Compl Gen	000/192/745/GCF_000192745.1_ASM19274v1
GCF_000724625.1	Fimbriimonas ginsengisoli Gsoil 348	Compl Gen	000/724/625/GCF_000724625.1_ASM72462v1
GCF_000279145.1	Melioribacter roseus P3M-2	Compl Gen	000/279/145/GCF_000279145.1_ASM27914v1
GCF_000147715.2	Mesotoga prima MesG1.Ag.4.2	Compl Gen	000/147/715/GCF_000147715.2_ASM14771v3
GCF_001443605.1	bacterium L21-Spi-D4	Compl Gen	001/443/605/GCF_001443605.1_ASM144360v1
GCF_001007995.1	'Deinococcus soli' Cha et al. 2014	Compl Gen	001/007/995/GCF_001007995.1_ASM100799v1
GCF_000484535.1	Gloeobacter kilauensis JS1	Compl Gen	000/484/535/GCF_000484535.1_ASM48453v1
GCF_000940805.1	Gynuella sunshinyii YC6258	Compl Gen	000/940/805/GCF_000940805.1_ASM94080v1
GCF_000828975.1	Burkholderiales bacterium GJ-E10	Compl Gen	000/828/975/GCF_000828975.1_ASM82897v1
GCF_000828655.1	Thermotoga caldifontis AZM44c09	Compl Gen	000/828/655/GCF_000828655.1_ASM82865v1
GCF_001281505.1	Lawsonella clevelandensis	Compl Gen	001/281/505/GCF_001281505.1_ASM128150v1
GCF_001023575.1	Actinobacteria bacterium IMCC26256	Compl Gen	001/023/575/GCF_001023575.1_ASM102357v1
GCF_001507665.1	Deinococcus actinosclerus	Compl Gen	001/507/665/GCF_001507665.1_ASM150766v1
GCF_000952975.1	Peptoniphilus sp. ING2-D1G	Compl Gen	000/952/975/GCF_000952975.1_D1G
Fungi:			
GCF_000002945.1	Schizosaccharomyces pombe	Chrom	000/002/945/GCF_000002945.1_ASM294v2
GCF_000315895.1	Millerozyma farinosa CBS 7064	Chrom	000/315/895/GCF_000315895.1_ASM31589v1
GCF_000209165.1	Scheffersomyces stipitis CBS 6054	Chrom	000/209/165/GCF_000209165.1_ASM20916v1
GCF_000146045.2	Saccharomyces cerevisiae S288c	Compl Gen	000/146/045/GCF_000146045.2_R64
GCF_000243375.1	Torulaspora delbrueckii	Chrom	000/243/375/GCF_000243375.1_ASM24337v1
GCF_000002525.2	Yarrowia lipolytica CLIB122	Chrom	000/002/525/GCF_000002525.2_ASM252v1
GCF_000026365.1	Zygosaccharomyces rouxii	Chrom	000/026/365/GCF_000026365.1_ASM2636v1
GCF_000006445.2	Debaryomyces hansenii CBS767	Chrom	000/006/445/GCF_000006445.2_ASM644v2
GCF_000182925.2	Neurospora crassa OR74A	Chrom	000/182/925/GCF_000182925.2_NC12

GCF_000091045.1	Cryptococcus neoformans var. neoformans JEC21	Chrom	000/091/045/GCF_000091045.1_ASM9104v1
GCF_000328475.2	Ustilago maydis 521	Chrom	000/328/475/GCF_000328475.2_Umaydis521_2.0
GCF_000182965.3	Candida albicans SC5314	Chrom	000/182/965/GCF_000182965.3_ASM18296v3
GCF_000002545.3	[Candida] glabrata	Chrom	000/002/545/GCF_000002545.3_ASM254v2
GCF_000149955.1	Fusarium oxysporum f. sp. lycopersici 4287	Chrom	000/149/955/GCF_000149955.1_ASM14995v2
GCF_000240135.3	Fusarium graminearum PH-1	Chrom	000/240/135/GCF_000240135.3_ASM24013v3
GCF_000091225.1	Encephalitozoon cuniculi GB-M1	Chrom	000/091/225/GCF_000091225.1_ASM9122v1
GCF_000237345.1	Naumovozyma castellii CBS 4309	Chrom	000/237/345/GCF_000237345.1_ASM23734v1
GCF_000227115.2	Naumovozyma dairenensis CBS 421	Chrom	000/227/115/GCF_000227115.2_ASM22711v2
GCF_000277815.2	Encephalitozoon hellem ATCC 50504	Chrom	000/277/815/GCF_000277815.2_ASM27781v3
GCF_000002515.2	Kluyveromyces lactis	Compl Gen	000/002/515/GCF_000002515.2_ASM251v1
GCF_000091025.4	Eremothecium gossypii ATCC 10895	Compl Gen	000/091/025/GCF_000091025.4_ASM9102v4
GCF_000226115.1	Thielavia terrestris NRRL 8126	Compl Gen	000/226/115/GCF_000226115.1_ASM22611v1
GCF_000185945.1	Cryptococcus gattii WM276	Chrom	000/185/945/GCF_000185945.1_ASM18594v1
GCF_000026945.1	Candida dubliniensis CD36	Compl Gen	000/026/945/GCF_000026945.1_ASM2694v1
GCF_000235365.1	Eremothecium cymbalariae DBVPG#7215	Chrom	000/235/365/GCF_000235365.1_ASM23536v1
GCF_000146465.1	Encephalitozoon intestinalis ATCC 50506	Chrom	000/146/465/GCF_000146465.1_ASM14646v1
GCF_000226095.1	Thermothelomyces thermophila ATCC 42464	Compl Gen	000/226/095/GCF_000226095.1_ASM22609v1
GCF_001672515.1	Colletotrichum higginsianum IMI 349063	Chrom	001/672/515/GCF_001672515.1_ASM167251v1
GCF_000303195.2	Fusarium pseudograminearum CS3096	Chrom	000/303/195/GCF_000303195.2_FP7
GCF_000236905.1	Tetrapisipora phaffii CBS 4417	Chrom	000/236/905/GCF_000236905.1_ASM23690v1
GCF_000149555.1	Fusarium verticillioides 7600	Chrom	000/149/555/GCF_000149555.1_ASM14955v1
GCF_000315875.1	Candida orthopsilosis	Chrom	000/315/875/GCF_000315875.1_ASM31587v1
GCF_000002495.2	Magnaporthe oryzae 70-15	Chrom	000/002/495/GCF_000002495.2_MG8
GCF_000142805.1	Lachancea thermotolerans CBS 6340	Chrom	000/142/805/GCF_000142805.1_ASM14280v1
GCF_000304475.1	Kazachstania africana CBS 2517	Chrom	000/304/475/GCF_000304475.1_Ka_CBS2517
GCF_000027005.1	Komagataella phaffii GS115	Chrom	000/027/005/GCF_000027005.1_ASM2700v1
GCF_000280035.1	Encephalitozoon romaleae SJ-2008	Chrom	000/280/035/GCF_000280035.1_ASM28003v2
GCF_000002655.1	Aspergillus fumigatus Af293	Chrom	000/002/655/GCF_000002655.1_ASM265v1
GCF_001640025.1	Sugiyamaella lignohabitans	Compl Gen	001/640/025/GCF_001640025.1_ASM164002v2
GCF_000187245.1	Ogataea parapolyomorpha DL-1	Chrom	000/187/245/GCF_000187245.1_Hansenula_2
GCF_000219625.1	Zymoseptoria tritici IPO323	Chrom	000/219/625/GCF_000219625.1_MYCGR_v2.0
GCF_000315915.1	Tetrapisipora blattae CBS 6284	Chrom	000/315/915/GCF_000315915.1_ASM31591v1
GCF_001298625.1	Saccharomyces eubayanus	Chrom	001/298/625/GCF_001298625.1_SEUB3.0
Invertebrates:			
GCF_000237925.1	Schistosoma mansoni	Chrom	000/237/925/GCF_000237925.1_ASM23792v2
GCF_000004555.1	Caenorhabditis briggsae	Chrom	000/004/555/GCF_000004555.1_ASM455v1

GCF_000002985.6	Caenorhabditis elegans	Compl Gen	000/002/985/GCF_000002985.6_WBcel235
GCF_000002335.3	Tribolium castaneum	Chrom	000/002/335/GCF_000002335.3_Tcas5.2
GCF_000005575.2	Anopheles gambiae str. PEST	Chrom	000/005/575/GCF_000005575.2_AgamP3
GCF_000001215.4	Drosophila melanogaster	Chrom	000/001/215/GCF_000001215.4_Release_6_plu s_ISO1_MT
GCF_000269505.1	Drosophila miranda	Chrom	000/269/505/GCF_000269505.1_DroMir_2.2
GCF_000001765.3	Drosophila pseudoobscura pseudoobscura	Chrom	000/001/765/GCF_000001765.3_Dpse_3.0
GCF_000754195.2	Drosophila simulans	Chrom	000/754/195/GCF_000754195.2_ASM75419v2
GCF_000005975.2	Drosophila yakuba	Chrom	000/005/975/GCF_000005975.2_dyak_caf1
GCF_000002325.3	Nasonia vitripennis	Chrom	000/002/325/GCF_000002325.3_Nvit_2.1
GCF_000002195.4	Apis mellifera	Chrom	000/002/195/GCF_000002195.4_Amel_4.5
GCF_000224145.2	Ciona intestinalis	Chrom	000/224/145/GCF_000224145.2_KH
GCF_001277935.1	Drosophila busckii	Chrom	001/277/935/GCF_001277935.1_ASM127793v1
GCF_000214255.1	Bombus terrestris	Chrom	000/214/255/GCF_000214255.1_Bter_1.0
Plants:			
GCF_000317415.1	Citrus sinensis	Chrom	000/317/415/GCF_000317415.1_Csi_valencia_1 .0
GCF_000987745.1	Gossypium hirsutum	Chrom	000/987/745/GCF_000987745.1_ASM98774v1
GCF_000208745.1	Theobroma cacao	Chrom	000/208/745/GCF_000208745.1_Criollo_cocoa_ genome_V2
GCF_000004075.2	Cucumis sativus	Chrom	000/004/075/GCF_000004075.2_ASM407v2
GCF_000002775.3	Populus trichocarpa	Chrom	000/002/775/GCF_000002775.3_Poptr2_0
GCF_000001735.3	Arabidopsis thaliana	Chrom	000/001/735/GCF_000001735.3_TAIR10
GCF_000686985.1	Brassica napus	Chrom	000/686/985/GCF_000686985.1_Brassica_napu s_assembly_v1.0
GCF_000309985.1	Brassica rapa	Chrom	000/309/985/GCF_000309985.1_Brapa_1.0
GCF_000695525.1	Brassica oleracea var. oleracea	Chrom	000/695/525/GCF_000695525.1_BOL
GCF_000148765.1	Malus domestica	Chrom	000/148/765/GCF_000148765.1_MalDomGD1.0
GCF_000340665.1	Cajanus cajan	Chrom	000/340/665/GCF_000340665.1_C.cajan_V1.0
GCF_000331145.1	Cicer arietinum	Chrom	000/331/145/GCF_000331145.1_ASM33114v1
GCF_000004515.4	Glycine max	Chrom	000/004/515/GCF_000004515.4_Glycine_max_v 2.0
GCF_001865875.1	Lupinus angustifolius	Chrom	001/865/875/GCF_001865875.1_LupAngTanjil_ v1.0
GCF_000219495.3	Medicago truncatula	Chrom	000/219/495/GCF_000219495.3_MedtrA17_4.0
GCF_000499845.1	Phaseolus vulgaris	Chrom	000/499/845/GCF_000499845.1_PhaVulg1_0
GCF_001190045.1	Vigna angularis	Chrom	001/190/045/GCF_001190045.1_Vigan1.1
GCF_001625215.1	Daucus carota subsp. sativus	Chrom	001/625/215/GCF_001625215.1_ASM162521v1
GCF_000710875.1	Capsicum annuum	Chrom	000/710/875/GCF_000710875.1_Pepper_Zunla_ 1_Ref_v1.0
GCF_000188115.3	Solanum lycopersicum	Chrom	000/188/115/GCF_000188115.3_SL2.50
GCF_000512975.1	Sesamum indicum	Chrom	000/512/975/GCF_000512975.1_S_indicum_v1. 0
GCF_001433935.1	Oryza sativa Japonica Group	Chrom	001/433/935/GCF_001433935.1_IRGSP-1.0
GCF_000231095.1	Oryza brachyantha	Chrom	000/231/095/GCF_000231095.1_Oryza_brachya ntha.v1.4b
GCF_000263155.2	Setaria italica	Chrom	000/263/155/GCF_000263155.2_Setaria_italica _v2.0
GCF_000003195.2	Sorghum bicolor	Chrom	000/003/195/GCF_000003195.2_Sorbi1
GCF_000005005.1	Zea mays	Chrom	000/005/005/GCF_000005005.1_B73_RefGen_v 3
GCF_001540865.1	Ananas comosus	Chrom	001/540/865/GCF_001540865.1_ASM154086v1

GCF_000313855.2	Musa acuminata subsp. malaccensis	Chrom	000/313/855/GCF_000313855.2_ASM31385v2
GCF_001876935.1	Asparagus officinalis	Chrom	001/876/935/GCF_001876935.1_Aspof.V1
GCF_000005505.2	Brachypodium distachyon	Chrom	000/005/505/GCF_000005505.2_Brachypodium_distachyon_v2.0
GCF_001406875.1	Solanum pennellii	Chrom	001/406/875/GCF_001406875.1_SPENNV200
GCF_000612285.1	Gossypium arboreum	Chrom	000/612/285/GCF_000612285.1_Gossypium_arboreum_v1.0
GCF_000327365.1	Gossypium raimondii	Chrom	000/327/365/GCF_000327365.1_Graimondii2_0
GCF_000003745.3	Vitis vinifera	Chrom	000/003/745/GCF_000003745.3_12X
GCF_000091205.1	Cyanidioschyzon merolae strain 10D	Compl Gen	000/091/205/GCF_000091205.1_ASM9120v1
GCF_001879085.1	Nicotiana attenuata	Chrom	001/879/085/GCF_001879085.1_NIATTr2
GCF_000442705.1	Elaeis guineensis	Chrom	000/442/705/GCF_000442705.1_EG5
GCF_000184155.1	Fragaria vesca subsp. vesca	Chrom	000/184/155/GCF_000184155.1_FraVesHawaii_1.0
GCF_000214015.2	Ostreococcus tauri	Chrom	000/214/015/GCF_000214015.2_version_050606
GCF_000633955.1	Camelina sativa	Chrom	000/633/955/GCF_000633955.1_Cs
GCF_000346735.1	Prunus mume	Chrom	000/346/735/GCF_000346735.1_P.mume_V1.0
GCF_000741045.1	Vigna radiata var. radiata	Chrom	000/741/045/GCF_000741045.1_Vradiata_ver6
GCF_000511025.2	Beta vulgaris subsp. vulgaris	Chrom	000/511/025/GCF_000511025.2_RefBeet-1.2.2
GCF_000092065.1	Ostreococcus lucimarinus CCE9901	Compl Gen	000/092/065/GCF_000092065.1_ASM9206v1
GCF_000090985.2	Micromonas commoda	Compl Gen	000/090/985/GCF_000090985.2_ASM9098v2
GCF_000826755.1	Ziziphus jujuba	Chrom	000/826/755/GCF_000826755.1_ZizJuj_1.1
Protoza:			
GCF_000150955.2	Phaeodactylum tricornutum CCAP 1055/1	Chrom	000/150/955/GCF_000150955.2_ASM15095v2
GCF_000002845.2	Leishmania braziliensis MHOM/BR/75/M2904	Chrom	000/002/845/GCF_000002845.2_ASM284v2
GCF_000227135.1	Leishmania donovani	Chrom	000/227/135/GCF_000227135.1_ASM22713v2
GCF_000002725.2	Leishmania major strain Friedlin	Compl Gen	000/002/725/GCF_000002725.2_ASM272v2
GCF_000234665.1	Leishmania mexicana MHOM/GT/2001/U1103	Chrom	000/234/665/GCF_000234665.1_ASM23466v4
GCF_000002875.2	Leishmania infantum JPCM5	Chrom	000/002/875/GCF_000002875.2_ASM287v2
GCF_000755165.1	Leishmania panamensis	Chrom	000/755/165/GCF_000755165.1_ASM75516v1
GCF_000210295.1	Trypanosoma brucei gambiense DAL972	Chrom	000/210/295/GCF_000210295.1_ASM21029v1
GCF_000165345.1	Cryptosporidium parvum Iowa II	Chrom	000/165/345/GCF_000165345.1_ASM16534v1
GCF_000006565.2	Toxoplasma gondii ME49	Chrom	000/006/565/GCF_000006565.2_TGA4
GCF_900002335.2	Plasmodium chabaudi chabaudi	Chrom	900/002/335/GCF_900002335.2_PCHAS01
GCF_000321355.1	Plasmodium cynomolgi strain B	Chrom	000/321/355/GCF_000321355.1_PcynB_1.0
GCF_000002765.3	Plasmodium falciparum 3D7	Chrom	000/002/765/GCF_000002765.3_ASM276v1
GCF_000006355.1	Plasmodium knowlesi strain H	Chrom	000/006/355/GCF_000006355.1_ASM635v1
GCF_001601855.1	Plasmodium reichenowi	Chrom	001/601/855/GCF_001601855.1_ASM160185v1
GCF_000002415.2	Plasmodium vivax	Chrom	000/002/415/GCF_000002415.2_ASM241v2
GCF_000165395.1	Babesia bovis	Chrom	000/165/395/GCF_000165395.1_ASM16539v1
GCF_000981445.1	Babesia bigemina	Chrom	000/981/445/GCF_000981445.1_Bbig001
GCF_000691945.1	Babesia microti strain RI	Chrom	000/691/945/GCF_000691945.1_ASM69194v1

GCF_000342415.1	Theileria equi strain WA	Chrom	000/342/415/GCF_000342415.1_JCVI-bewag-v1.1
GCF_000003225.2	Theileria annulata strain Ankara	Chrom	000/003/225/GCF_000003225.2_ASM322v1
GCF_000165365.1	Theileria parva	Chrom	000/165/365/GCF_000165365.1_ASM16536v1
GCF_000208865.1	Neospora caninum Liverpool	Chrom	000/208/865/GCF_000208865.1_ASM20886v2
GCF_000149405.2	Thalassiosira pseudonana CCMP1335	Chrom	000/149/405/GCF_000149405.2_ASM14940v2
GCF_000004695.1	Dictyostelium discoideum	Chrom	000/004/695/GCF_000004695.1_dicty_2.7
GCF_000740895.1	Theileria orientalis strain Shintoku	Compl Gen	000/740/895/GCF_000740895.1_ASM74089v1
GCF_001680005.1	Plasmodium coatneyi	Chrom	001/680/005/GCF_001680005.1_ASM168000v1
GCF_001602025.1	Plasmodium gaboni	Chrom	001/602/025/GCF_001602025.1_ASM160202v1
Mammals:			
GCF_000002275.2	Ornithorhynchus anatinus	Chrom	000/002/275/GCF_000002275.2_Ornithorhynchus_anatinus_5.0.1
GCF_000004665.1	Callithrix jacchus	Chrom	000/004/665/GCF_000004665.1_Callithrix_jacchus-3.2
GCF_000364345.1	Macaca fascicularis	Chrom	000/364/345/GCF_000364345.1_Macaca_fascicularis_5.0
GCF_000772875.2	Macaca mulatta	Chrom	000/772/875/GCF_000772875.2_Mmul_8.0.1
GCF_000264685.2	Papio anubis	Chrom	000/264/685/GCF_000264685.2_Panu_2.0
GCF_000151905.2	Gorilla gorilla gorilla	Chrom	000/151/905/GCF_000151905.2_gorGor4
GCF_000258655.2	Pan paniscus	Chrom	000/258/655/GCF_000258655.2_panpan1.1
GCF_000001515.7	Pan troglodytes	Chrom	000/001/515/GCF_000001515.7_Pan_tro_3.0
GCF_000001545.4	Pongo abelii	Chrom	000/001/545/GCF_000001545.4_P_pygmaeus_2.0.2
GCF_000001405.36	Homo sapiens	Chrom	000/001/405/GCF_000001405.36_GRCh38.p10
GCF_000002285.3	Canis lupus familiaris	Chrom	000/002/285/GCF_000002285.3_CanFam3.1
GCF_000181335.2	Felis catus	Chrom	000/181/335/GCF_000181335.2_Felis_catus_8.0
GCF_000002305.2	Equus caballus	Chrom	000/002/305/GCF_000002305.2_EquCab2.0
GCF_000003025.5	Sus scrofa	Chrom	000/003/025/GCF_000003025.5_Sscrofa10.2
GCF_000003055.6	Bos taurus	Chrom	000/003/055/GCF_000003055.6_Bos_taurus_UMD_3.1.1
GCF_000247795.1	Bos indicus	Chrom	000/247/795/GCF_000247795.1_Bos_indicus_1.0
GCF_001704415.1	Capra hircus	Chrom	001/704/415/GCF_001704415.1_ARS1
GCF_000298735.2	Ovis aries	Chrom	000/298/735/GCF_000298735.2_Oar_v4.0
GCF_000003625.3	Oryctolagus cuniculus	Chrom	000/003/625/GCF_000003625.3_OryCun2.0
GCF_000001635.25	Mus musculus	Chrom	000/001/635/GCF_000001635.25_GRCm38.p5
GCF_000001895.5	Rattus norvegicus	Chrom	000/001/895/GCF_000001895.5_Rnor_6.0
GCF_000002295.2	Monodelphis domestica	Chrom	000/002/295/GCF_000002295.2_MonDom5
GCF_000409795.2	Chlorocebus sabaeus	Chrom	000/409/795/GCF_000409795.2_Chlorocebus_sabaeus_1.1
GCF_000146795.2	Nomascus leucogenys	Chrom	000/146/795/GCF_000146795.2_Nleu_3.0
GCF_000317375.1	Microtus ochrogaster	Chrom	000/317/375/GCF_000317375.1_MicOch1.0
Non-mammalian vertebrates:			
GCF_000242695.1	Lepisosteus oculatus	Chrom	000/242/695/GCF_000242695.1_LepOcu1
GCF_000002035.5	Danio rerio	Chrom	000/002/035/GCF_000002035.5_GRCz10
GCF_000951615.1	Cyprinus carpio	Chrom	000/951/615/GCF_000951615.1_common_carp_genome
GCF_001660625.1	Ictalurus punctatus	Chrom	001/660/625/GCF_001660625.1_IpCoco_1.2
GCF_000721915.3	Esox lucius	Chrom	000/721/915/GCF_000721915.3_Eluc_V3

GCF_000233375.1	Salmo salar	Chrom	000/233/375/GCF_000233375.1_ICSASG_v2
GCF_000633615.1	Poecilia reticulata	Chrom	000/633/615/GCF_000633615.1_Guppy_female_1.0_MT
GCF_000313675.1	Oryzias latipes	Chrom	000/313/675/GCF_000313675.1_ASM31367v1
GCF_001858045.1	Oreochromis niloticus	Chrom	001/858/045/GCF_001858045.1_ASM185804v2
GCF_001663975.1	Xenopus laevis	Chrom	001/663/975/GCF_001663975.1_Xenopus_laevis_v2
GCF_000004195.3	Xenopus tropicalis	Chrom	000/004/195/GCF_000004195.3_Xenopus_tropicalis_v9.1
GCF_000241765.3	Chrysemys picta bellii	Chrom	000/241/765/GCF_000241765.3_Chrysemys_picta_bellii-3.0.3
GCF_000002315.4	Gallus gallus	Chrom	000/002/315/GCF_000002315.4_Gallus_gallus-5.0
GCF_000146605.2	Meleagris gallopavo	Chrom	000/146/605/GCF_000146605.2_Turkey_5.0
GCF_001522545.2	Parus major	Chrom	001/522/545/GCF_001522545.2_Parus_major1.1
GCF_000090745.1	Anolis carolinensis	Chrom	000/090/745/GCF_000090745.1_AnoCar2.0
GCF_000180615.1	Takifugu rubripes	Chrom	000/180/615/GCF_000180615.1_FUGU5
GCF_000151805.1	Taeniopygia guttata	Chrom	000/151/805/GCF_000151805.1_Taeniopygia_guttata-3.2.4
GCF_000247815.1	Ficedula albicollis	Chrom	000/247/815/GCF_000247815.1_FicAlb1.5
GCF_001577835.1	Coturnix japonica	Chrom	001/577/835/GCF_001577835.1_Coturnix_japonica_2.0
GCF_001465895.1	Nothobranchius furzeri	Chrom	001/465/895/GCF_001465895.1_Nfu_20140520
GCF_000523025.1	Cynoglossus semilaevis	Chrom	000/523/025/GCF_000523025.1_Cse_v1.0

[98] The examples show:

[99] **Example 1:** Design of a nucleic acid library of the invention that encodes short peptides based on amino acid sequences of proteins comprised in the human proteome ("HuPEX").

- 5 [100] First, a single "mega protein" amino acid sequence was generated by concatenating the amino acid sequences of a plurality of individual proteins (in this case, all 21,018 such proteins) comprised in the reference proteome of *Homo sapiens*. Hence, such amino acid sequences can be considered to be of naturally occurring proteins; that is they occur naturally in humans. The reference proteome used (UP000005640_9606.fasta) was obtained from the EMBL-EBI web-based resource: "Reference proteomes - Primary proteome sets for the Quest For
- 10 Orthologs" (http://www.ebi.ac.uk/reference_proteomes, accessed on 05-Feb-2017), Release 2017_01 based on UniProt Release 2017_01.

[101] Each join between the amino acid sequences of two concatenated proteins was marked by using a spacer symbol "_", such that (for example) two example proteins listed, in FASTA-format, in a human reference proteome:

15 >tr|A0A024R161|A0A024R161_HUMAN Guanine nucleotide-binding protein subunit gamma OS=Homo sapiens GN=DNAJC25-GNG10 PE=3 SV=1
MGAPLLSPGWGAGAAGRRWWMLLAPLLPALLLVLPAGALVEGLYCGTRDCYEVLGVSRSA
GKAEIARAYRQLARRYHPDRYRPQPGDEGPGRT PQSAEEAFLLVATAYETLKVVSQAAEL
QQYCMQNACKDALLVGVPA GSNPFREPRSCALL

20 >tr|A0A075B6F4|A0A075B6F4_HUMAN T-cell receptor beta variable 21/OR9-2 (pseudogene) (Fragment) OS=Homo sapiens GN=TRBV21OR9-2
PE=4 SV=1
XRFLSEPTRCLRLCCVALSFWGAASMDTKVTQRPRFLVKANEQKAKMDCVPPIKRHSYVY
WYHKTLEELKFFIYFQNEEIIQKAEI INERFSAQCPQNSPCTLEIQTESGDTARYFCA
NSK

- 25 [102] would be concatenated to form a single amino acid sequence, with the region around the join shown below:

[...] LVGVPA GSNPFREPRSCALL_XRFLSEPTRCLRLCCVALS [...]

[103] Second, a computer program selected all 46 amino acid-long regions that were spaced apart by a window of 10 amino acids from the “mega protein” amino acid sequence representing the plurality of proteins (in this case, all 21,018). However, in other embodiments, the computer program could be instructed to select amino acid regions having an alternative (predefined) length, for example any such length that is between 25 amino acids and 110 amino acids; and/or the computer program could be instructed to space such regions apart by an alternative (predefined) window, for example such a window of between about 2 and 40 amino acids. For example, a filing tiling of such a selection of 46 amino acid regions spaced by a 10 amino acid window from a concatenation of the first proteins above would generate the following amino acid sequences representing short peptides (only the first three such peptides sequences are shown):

10 Peptide 1: MGAPLLSPGWGAGAAGRWWMLLAPLLPALLLVRPAGALVEGLYCG
Peptide 2: GAGAAGRWWMLLAPLLPALLLVRPAGALVEGLYCGTRDCYEVLGV
Peptide 2: MLLAPLLPALLLVRPAGALVEGLYCGTRDCYEVLG [...]

[104] Third, the resulting set of 46 amino acid-long peptide sequences were filtered by removing from the set any sequences with any one (or more) of the following features:

- 15 (a) Any sequence having a length NOT equal to 46 amino acids (both, as a quality-control step for the computer program, and to remove any – short – sequences arising from the end of a protein sequence);
- (b) Any sequence comprising the spacer symbol (in this case, “_”) representing a join between the amino acid sequences of two proteins;
- 20 (c) Any sequence comprising a symbol for an ambiguous amino acid (for example, “B”, “J”, “X” or “Z”; where in certain databases, such ambiguity codes can have the following meanings: B = D or N, J = I or L, X = unknown, Z = E or Q; as an example, the first 46 amino acid-long peptide sequence of the second protein shown above – as it starts with an “X” – would hence be removed from the set); and/or
- 25 (d) Any sequence that is NOT unique (that is, any sequence that is 100% identical to any other sequence).

[105] Fourth, the isoelectric point (pI) of each resulting peptide sequence remaining in the set was predicted using generally available pI-prediction software (in this case, the “pI” function of the “R peptide Package”; Osorio et al 2015, The R Journal. 7:4; <https://cran.r-project.org/web/packages/Peptides/index.html>; using the argument pKscale = “EMBOSS”). Any peptide having an amino acid sequence predicted to have a pI of between 6 and 8 was also removed from the set. The isoelectric point of a peptide is the pH at which the peptide would have no net charge, and can often precipitate out of a solution of such pH. Accordingly, peptides with a predicted pI of around physiological pH (e.g. 6 to 7.4 or 6 to 8) are, in some embodiments, excluded from the set, as they may be those more likely to have unfavourable properties under such conditions (e.g., by precipitating upon expression).

35 **[106]** Fifth, the resulting set of amino acid sequences (after the filtering described in the third and second step) was subjected to a final filtering to positively select only those 46 amino acid sequences that showed 100% identity to an amino acid sequence present in the reference proteome of Mouse (*Mus musculus*; UP000000589_10090.fasta, obtained as described above).

[107] As will now be understood by the person of ordinary skill, the final set of (46-long) amino acid sequences will be spaced apart along the naturally occurring protein(s) by a window of (in this case) 10 amino acids or multiples of 10 amino acids. This is because one or more intervening amino acid sequences may have been omitted from the final set because it did not conform to one or other of the various filtering criteria.

[108] The final set of (46-long) amino acid sequences consisted, in this example, of about 300,000 individual sequences, with at least one such sequence representative of over 21,000 different proteins of the human proteome, and a mean of about 14.2 such sequences per protein and a standard deviation of 3.8 sequences (with 95% of the proteins will be represented by between 7.6 and 20.7 peptide sequences).

5 **[109]** As will now be apparent to the person of ordinary skill, the final set of amino acid sequences of peptides that represent the proteome(s) used for the selection may be more or less than this number, and the distribution of the number of peptides for each of the proteins may also differ. Not only will the number and nature of the final set of amino acid sequences depend on the sequences of the plurality of proteins first concatenated (for example, the proteome of a single species used, or those of multiple species), but other factors can affect the final set. For
10 example, the sequence regions initially selected may be shorter (or longer) than the 46 amino acids used herein, and/or spacing window may be shorter (or longer) than the 10 amino acids used herein. Furthermore, one or more of the filtering criteria described above may be omitted, or different filtering criteria may be applied instead or as well (examples of other filtering steps are described in other examples herein). Indeed, depending on the particular properties (such as size, diversity, coverage and/or solubility etc.) desired for a library of the present invention,
15 and/or the method of its physical preparation (such as described below), and in particular the limitations in size, complexity or cost of the method used for its physical preparation, the person of ordinary skill will now have the ability to select only those amino acid sequences for peptides to form any particular library suitable for their needs.

[110] Six, each 46 amino acid sequence was reverse translated using for each amino acid the most frequently used codon for such amino acid of the human codon frequency table shown in Table 1.1 below (Codon Usage Database; Nakamura et al, 2000, NAR 28:292; <http://www.kazusa.or.jp/codon/cgi-bin/showcodon.cgi?species=9606&aa=1&style=N>, accessed 04-Jun-2017).
20

[111] As will now be apparent to the person of ordinary skill, however, alternative human codon frequency tables may be used or, and depending on the species of the expression system in which the nucleic acid library is intended to be expressed, codon frequency tables of other species may be used.

25

Table 1.1: Human codon frequency table (codon | aa | fraction per codon per aa).

UUU	F	0.46	UCU	S	0.19	UAU	Y	0.44	UGU	C	0.46
UUC	F	0.54	UCC	S	0.22	UAC	Y	0.56	UGC	C	0.54
UUA	L	0.08	UCA	S	0.15	UAA	*	0.30	UGA	*	0.47
UUG	L	0.13	UCG	S	0.05	UAG	*	0.24	UGG	W	1.00
CUU	L	0.13	CCU	P	0.29	CAU	H	0.42	CGU	R	0.08
CUC	L	0.20	CCC	P	0.32	CAC	H	0.58	CGC	R	0.18
CUA	L	0.07	CCA	P	0.28	CAA	Q	0.27	CGA	R	0.11
CUG	L	0.40	CCG	P	0.11	CAG	Q	0.73	CGG	R	0.20
AUU	I	0.36	ACU	T	0.25	AAU	N	0.47	AGU	S	0.15
AUC	I	0.47	ACC	T	0.36	AAC	N	0.53	AGC	S	0.24
AUA	I	0.17	ACA	T	0.28	AAA	K	0.43	AGA	R	0.21
AUG	M	1.00	ACG	T	0.11	AAG	K	0.57	AGG	R	0.21
GUU	V	0.18	GCU	A	0.27	GAU	D	0.46	GGU	G	0.16
GUC	V	0.24	GCC	A	0.40	GAC	D	0.54	GGC	G	0.34
GUA	V	0.12	GCA	A	0.23	GAA	E	0.42	GGA	G	0.25
GUG	V	0.46	GCG	A	0.11	GAG	E	0.58	GGG	G	0.25

[112] Seven, the resulting nucleic acid sequences encoding for the peptides were then analysed for undesired sub-sequences resulting from combinations of codons. Particular examples of such undesired sub-sequence include an internal Kozak sequence and/or a restriction enzyme site for a restriction enzyme intended to be used for the cloning of the resulting library, in each case generated from a combination of codons. If an undesired sub-sequence was present in a nucleic acid sequence, then the next most commonly used codon was used in place of one or other of the codons in the combination such that undesired subsequence was no longer present in the nucleic acid sequence.

[113] In this example, a Kozak sequence of "CCATGG" was deemed undesired, and any codon combinations in the nucleic acid sequences that formed such a sub-sequence were adapted to use a less frequent codon so that such sub-sequence was no longer present.

[114] As the library of this example was to be cloned into expression vectors using certain restriction enzymes, any sub-sequences in nucleic acid sequences that were recognition sites for such restriction enzymes were also deemed undesirable. The restriction enzymes BamHI and XhoI were envisioned as restriction enzymes that may be used in such cloning procedures, and hence any of the following sub-sequences (the respective recognition site for such restriction enzymes) were deemed undesirable: 5'GGATCC and 5' CTCGAG. Any codon combinations in the nucleic acid sequences that formed such any sub-sequence that was such a (predefined) restriction enzymes recognition site were adapted to use a less frequent codon so that such sub-sequence was no longer present. As will be now apparent to the person of ordinary skill, depending on any restriction enzyme(s) planned to be used, the applicable sub-sequence consisting of a recognition site of such restriction enzyme(s) can be so removed by appropriate use of an alternative codon.

[115] As will also be apparent to the person of ordinary skill, as the nucleic acid sequences were generated by reverse translation of the (naturally occurring) amino acid sequences, the resulting library of nucleic acid sequences will comprise nucleic acids sequences that themselves are non-natural, as codons will be used (and/or combinations or codons) that are not those used by the naturally occurring genomic sequence to code one or more particular amino acids at that position in the protein. Accordingly, the nucleic acid libraries of the present invention will comprise a plurality of non-natural nucleic acid sequences.

[116] **Example 2:** Synthesis of the HuPEX nucleic acid library of the invention.

[117] First, each nucleic acid sequence in the library designed in Example 1 to encode for unique 46 amino acid long peptide "fragments" of naturally occurring human proteins was adapted with 5' and 3' sequences adapted to provide the sequence of an oligonucleotide enable cloning of the nucleic acid library and/or to enable expression of the peptide encoded therein.

[118] In this example, the general structure of each resulting nucleic acid sequence of such oligonucleotide (with the 5' and 3' regions) was:

35 Forward-amp_BamHI_Kozac_Gly_VARIABLE-REGION_Stop_XhoI_Reverse-amp,

[119] where "Forward-amp" and "Reverse-amp" represent nucleic acid sequences chosen so that the resulting oligo could be amplified by PCR using appropriate primers; "BamHI" and "XhoI" represent nucleic acid sequences of the respective restriction enzyme's recognition site; "Kozak" represents a Kozak sequence (including a start codon); "Gly" represents a codon for a single glycine acid-linker; "Stop" represents a stop codon; and "VARIABLE-REGION" represents an individual 138bp nucleic acid that encodes a given 46 amino acid long peptide (present in the set), from the library of nucleic acid designed in Example 1.

[120] In this example, the forward and reverse amplification sequences, Kozak sequence and stop codon used is set forth in Table 2.1 below.

Table 2.1: Common sequences used in the HuPEX library.

Feature	Sequence (5' to 3')	SEQ ID NO.:
Forward amplification	TGCCACCTGACGTCTAAGAA	1
Reverse amplification	GCTCACTCAAAGGCGGTAAT	2
Kozak sequence	CCATGG	N/A
Stop codon	TAA	N/A

[121] Accordingly, a nucleic acid sequence of an indicative oligonucleotide resulting from such design and encoding for the following indicative 46 amino acid sequence (**SEQ ID NO. 13**) present in the set:

5 LAQTACVVGR PGPHTQFLA AKERTKSHVP SLLDADVEGQ SRDYTV,

[122] has a complete nucleic acid sequence shown below (SEQ ID NO. 3), where the following features are marked as follows: 138bp region coding for the above 46 amino acid sequence in bold; forward and reverse amplification sequences in lower case; restriction enzyme sites boxed, Kozak sequence double underlined; and "*" is indicated above the first base of the start and stop codons.

10 **SEQ ID NO. 3:** Nucleic acid sequence of an indicative oligonucleotide

*

tgccacctga cgtctaagaa GGATCCATGG GACTCGCCCA GACCGCCTGT GTGGTGGGCA

GGCCCGGCCC CCACCCACC CAGTTCCTCG CCGCCAAGGA AAGGACCAAG AGCCACGTGC

*

15 CCAGCCTCCT CGACGCCGAC GTGGAAGGCC AGAGCAGGGA CTACACCGTG TAACTCGAGg

ctcactcaaa ggcggtaat

20 [123] As will now be apparent to the person of ordinary art, the nucleic acid sequence of the final oligonucleotides may be, for example, alternatively designed not to comprise a Kozak/start or stop codon(s). In this case, the Kozak/start and stop codon(s) can be provided by the host vector. Such a design (as demonstrated in other examples below) can allow the libraries to be produced in tagged format, by cloning them into a vector with, for example, a vector-encoded FLAG tag, N- or C-terminally to the encoded peptide.

25 [124] In alternative embodiments, two versions of a library of the invention may be constructed: one with and one without a start codon. Such an embodiment can be useful as an internal reference for data analysis of screens using the library, as it would allow effects of the peptide from effects of the vector/construct itself (which should be identified as false positives) to be distinguished. For example, to distinguish those cases where particular sequences are amplified better by PCR, DNA sequences that attract some cellular machinery and cause indirect effects etc.

30 [125] Second, the set of nucleic acid sequences of all approximately 300,000 oligonucleotides was used to chemically synthesis each of the oligonucleotides by conventional approaches.

[126] In this case, the semi-pooled 10,000-well silicon chip-based oligonucleotide synthesis approach of Twist Bioscience (San Francisco, California) was used to synthesise all approximately 300,000 oligonucleotides, and the oligonucleotides were therefore available in sub-pools each having a complexity of approximately 2,000 oligonucleotides. It should be noted here, that if each sub-pool-synthesised oligonucleotide were synthesised with a different combination of pairs of primer-pairs (for example, 45 different primer-pairs used in all combinations would provide $45 \times 45 = 2025$ different combinations), then it would then be possible to recover an *individual* sequence by conducting PCR with only using the applicable specific primer-pair combination. However, other synthetic oligonucleotide approaches could be used, such as array-based synthesis (e.g., Affymetrix) or in-situ synthesis printing (Agilent).

40

[127] **Example 3:** Cloning of the HuPEx nucleic acid library of the invention into expression vectors.

[128] The set of approximately 300,000 oligonucleotides were (in pooled or sub-pooled format) cloned into a lentiviral expression system briefly described as follows.

[129] First, a pool of oligonucleotides was subjected to PCR amplification (by standard procedures and using the primers having the sequences 5'-TGCCACCTGACGTCTAAGAA-3' (**SEQ ID NO. 4**) and 5'-ATTACCGCCTTTGAGTGAGC-3' (**SEQ ID NO. 5**), corresponding to the forward and reverse amplification sequences, respectively, in the oligonucleotides. Second, the resulting product was digested with the applicable restriction enzymes (in this case BamHI and XhoI) to provide sticky-ended constructs for cloning.

[130] Third, the sticky-ended constructs were ligated into a sample of a BamHI/XhoI-digested lentiviral vector (for example, pMOST25). Sixty (60) base pairs of the cloning site of pMOST25 is shown below by SEQ ID NO. 6, and the resulting recombinant construct with a BamHI/XhoI-digested amplification product of the indicative oligonucleotide shown in Example 2, would have the sequence shown by SEQ ID NO. 7. BamHI and XhoI restriction sites are boxed; the Kozak sequence is double underlined; the first base of the start/stop codons marked above with a "*"; and the 138bp region coding for the 46 amino acid sequence in bold.

15 **SEQ ID NO. 6:** 60bp of the cloning site of pMOST25 lentiviral vector.

AGTAGCATCG CATTAGCCG G GATCC AGCGC TGCTAC CTC GAG TAAGTGA CTAGGCAATC

SEQ ID NO. 7: Indicative recombinant construct.

20 AGTAGCATCG CATTAGCCG G GATCC ^{*}ATGGG **ACTCGCCCAG ACCGCCTGTG TGGTGGGCAG**
 GCCCGGCCCC CACCCACCC AGTTCCTCGC CGCCAAGGAA AGGACCAAGA GCCACGTGCC
 CAGCCTCCTC GACGCCGACG TGAAGGCCA GAGCAGGGAC TACACCGTGT AA ^{*}CTCGAGTA
 25 AGTGACTAGG CAATC

[131] Transformation/transfection of the recombinant vector into a host cell will enable propagation, expression and/or screening of the nucleic acid library of the invention. As will be apparent to the person of ordinary skill, the a nucleic acid library - after being amplified and/or propagated/maintained in host cells - may be considered to no longer be "synthetic", as the nucleic acid molecules will, by then, have been enzymatically (either in-vitro or in-vivo) produced. However, such a nucleic acid library is still considered to be one of the present invention.

[132] Expression of the construct shown in SEQ ID NO. 7 will result in the 48 amino acid peptide having a sequence as shown in SEQ ID NO. 8, where the sequence of the originally encoded 46 amino acid indicative peptide is shown in bold after the initial methionine and the linking glycine.

SEQ ID NO. 8: Indicative expressed peptide.

35 **MGLAQ**TACVV GRPGPHPTQF **LA**AKERTKSH VPSLLDADVE GQSRDYTV

[133] As the person of ordinary skill we note, such expressed comprises N- and C-terminal amino acids corresponding to the 5' and 3' restriction enzyme recognition sites (and, in this case, a single linking Val). Accordingly, even though the original 46 amino acid sequence is a fragment of a naturally occurring protein, the resulting 52 amino acid expressed peptide is a non-natural. Accordingly, the libraries of the present invention will encode for a plurality of non-natural peptides.

[134] The cloning of the library can be conducted in a pooled manner, or in a semi-pooled manner. Furthermore, the resulting pooled/sub-pooled clones can be individually picked/arrayed using automated cell-sorting technologies, such as the QPix (Molecular Devices), FACS or single-cell dispensing (e.g. VIPS™ Cell Dispenser or Solentim Ltd, UK, or the Single Cell Printer – SCP – or Cytena GmbH, Germany).

[135] **Example 4:** Phenotypic screening with the lentiviral-cloned HuPEX library.

[136] A HuPEX library, such as designed and constructed above, was used to screen for phenotypic alterations using the assay formats described above, either as pooled or arrayed libraries. Prior to such screen, the HuPEX library can, alternatively, be pooled with one or more other analogous libraries that expressed short peptides. For example, the HuPEX library can be screened in a pool together with a library that expresses human small Open Reading Frames (sORFs), such as the sORF library described in PCT/GB2016/054038 (in particular, in Example A) of PCT/GB2016/054038).

[137] (1) Pooled 6-thioguanine resistance screen

[138] Resistance to the chemotherapeutic drug 6-thioguanine has been previously demonstrated to be a fairly strict selection system; with a narrow group of proteins being able to mediate the phenotype (see Wang et al 2014, Science 343:80). The inventors sought to use this system to demonstrate how a library of the present invention can be utilised in identifying phenotype-modulating proteins even under such stringent conditions.

[139] HEK293 cells were transfected with the pooled HuPEX nucleic acid library, cloned into the lentiviral vector as above, which is designed to express a plurality of short-expressed-peptides (SEPs). Virus was harvested, titered, and a batch of KBM7 cells was infected with the SEP-expressing viruses. The library of virus-transduced KBM7 cells was subsequently exposed to a concentration of 6-thioguanine that was experimentally determined to kill 99.999% of KBM7 cells. Survivors, carrying inserts that express resistance-inducing SEPs, were isolated from the pool, expanded and genomic DNA was harvested. The inserts that express resistance-inducing SEPs were amplified using PCR and submitted to Next Generation Sequencing. After bioinformatics analysis of the data, SEPs that mediate the resistance to 6-thioguanine, and likely acting upon mismatch repair processes, were identified (**Figure 1**).

[140] (2) Pooled PTEN synthetic lethality screen

[141] To find SEPs able to selectively inhibit proliferation or kill cells lacking the PTEN tumour suppressor, SEPs were screened in an isogenic cell model pair (MCF10A WT and MCF10A PTEN knockout).

[142] As in (1) above, a pool of cells was infected with a library of HuPEX expressing SEPs. Cells were also infected with BugPEX and OmePEX libraries in a similar fashion. Under all conditions, the target cell lines, MCF10A and MCF10A PTEN KO, were infected in parallel. The cells were plated at low density in order to allow for cell growth over a period of five days. Samples from either cell population were then submitted to NGS as in (1). The relative abundances of SEP sequences in the wildtype-control control set (MCF10A) and the PTEN knockout set (MCF10A PTEN KO) were compared, and SEPs depleted in the knockout cells were identified (**Figure 2**).

[143] Identified hits were then validated in the same models by repeating the primary assay with additional controls. Hits were considered validated if they showed a significant reduction of cell growth in MCF10A PTEN KO cells compared to MCF10A cells in three biological replicates. Figure 3 shows an example validated hit being HuPEX sequence #30-325, a 46 amino acid sequence from human Tetraspanin-3 (the sequence of which, together with the two leading two amino acids - "MG" - as it is expressed by pMOST25 is shown as **SEQ ID NO. 15**), which inhibited growth > 60% in PTEN knockout cells compared to the control cell line, to an amount comparable with a positive control (shRNA against NLK, previously described to be synthetic lethal with PTEN knockout. Mendes-Pereira et. Al, PlosONE 2012). The nucleic acid sequence of the oligonucleotide that was synthesised to express such peptide is shown in **SEQ ID NO. 16**).

[144] **Example 5:** Design and construction of a nucleic acid library of the invention that encodes short peptides based on amino acid sequences of proteins comprised in the proteomes of evolutionary diverse microbiota ("BugPEX").

[145] A library of the invention was designed and constructed wherein the amino acid sequences of the source-proteins were those of naturally occurring proteins from a plurality of different species; in this example, the protein sequences comprised in the reference proteomes of an evolutionary diverse set of micro-organisms.

[146] First, a mega protein amino acid sequence was generated as described in Example 1, but by using all the protein sequences comprised in the set of reference proteomes set out Table A. The second, third and fourth steps of Example 1 were conducted by analogy, to generate a filtered set of several hundred thousand unique 46 amino acid long sequences.

[147] In this case however, the fifth step of Example 1 was replaced by an alternative filtering step to select, from such list of several hundred thousand amino acid sequences, 500,000 of such sequences that were predicted to be least likely to have a disordered segment. For example, the program DisEMBL (Linding et al 2003, Structure 11:1453; <http://dis.embl.de>) can be applied to consider and sort on the three intrinsically disordered protein parameters of loops/coils, hot-loops and Remark-465. The resulting number of amino acids present in a disordered stretch in a given peptide is counted, and such counts are ranked for each of the three parameters. The peptide sequences are then ranked by the mean of all three ranked-parameters so that peptides predicted to be highly disordered are ranked at the bottom of the list. Alternatively, peptides having a long disordered segment can be predicted using SLIDER (Super-fast predictor of proteins with long intrinsically disordered regions; Peng et al, 2014; Proteins: structure, Function and Bioinformatics 82:145; <http://biomine.cs.vcu.edu/servers/SLIDER>) using default parameters.

[148] The resulting 600,000 amino acid sequences in the filtered set were reverse-translated as described to nucleic acid sequences in the sixth step of Example 1, with alternative codons used to avoid undesired codon-combinations as described by the seventh step in Example 1. With an estimated 80,000 naturally occurring proteins represented by such 600,000 peptides, this suggests a mean coverage of about 7.5 nucleic acids (peptides) per naturally occurring protein

[149] For synthesis of oligonucleotides comprising these 600,000 nucleic acid sequences, the procedure described in Example 2 was followed except that in this case the general structure of the resulting oligonucleotide (with the 5' and 3' regions) was:

Forward-amp_BamHI_Val_VARIABLE-REGION_XhoI_Reverse-amp,

[150] where "Val" represents a codon for a single valine amino acid-linker, and the other features are as described in Example 2, such that or an indicative 46 amino acid sequence (**SEQ ID NO. 14**) of:

PRYLKGLKD VVQLSLRRPS FRASRQRPPII SLNERILEFN KRNITA,

[151] the resulting oligonucleotide sequence has a complete nucleic acid sequence shown below (SEQ ID NO. 9), where the following features are marked as follows: 138bp region coding for the above 46 amino acid sequence in bold; forward and reverse amplification sequences in lower case; restriction enzyme sites boxed.

SEQ ID NO. 9: Nucleic acid sequence of an indicative oligonucleotide

tgccacctga cgtctaagaa **GGATCC**GTGC **CTAGGTACCT GAAGGGCTGG CTGAAGGACG**
TGGTGCAGCT GAGCCTGAGG AGGCCTAGCT TCAGGGCCAG CAGGCAGAGG CCTATCATCA
GCCTGAACGA GAGGATTCTG GAGTTCAACA AGAGGAACAT CACAGCC**CTC GAG**gctcact
 caaaggcggt aat

[152] Note that in this case, the oligonucleotide does not comprise a Kozak/start or a stop codon, which in this example is provided by the expression vector into which the oligonucleotide is cloned, as described next.

[153] The oligonucleotides were amplified and digested as described in first and second steps of Example 3. In this case, however, the resulting digest products were cloned into a sample of a BamHI/XhoI-digested lentiviral vector comprising a Kozak/start and stop codon (eg pMOST25A). Sixty (60) base pairs of the cloning site of pMOST25A is shown below by SEQ ID NO. 10, and the resulting recombinant construct with a BamHI/ XhoI-digested amplification product of the indicative oligonucleotide shown in this Example 5, would have the sequence shown by SEQ ID NO. 11. BamHI and XhoI restriction sites are boxed; the first base of the start/stop codons marked above with a “*”; and the 138bp region coding for the 46 amino acid sequence in bold.

SEQ ID NO. 10: 60bp of the cloning site of pMOST25A lentiviral vector.

10 CATTAGCCAT GGGATCCAGC GCTGCTACC TCGAGTAAGT GACTAGGCCAA TCTAATCTAT

SEQ ID NO. 11: Indicative recombinant construct.

15
20

CATTAGCCAT GGGATCCGTG CCTAGGTACC TGAAGGGCTG GCTGAAGGAC GTGGTGCAGC
TGAGCCTGAG GAGGCCTAGC TTCAGGGCCA GCAGGCAGAG GCCTATCATC AGCCTGAACG
AGAGGATTCT GGAGTTCAAC AAGAGGAACA TCACAGCCCTCGAGTAAGTG ACTAGGCAAT

CTAATCT

[154] Expression of the construct shown in SEQ ID NO. 11 will result in the 52 amino acid peptide having a sequence as shown in SEQ ID NO. 12, where the sequence of the originally encoded 46 amino acid indicative peptide is shown in bold after the initial methionine, the two amino acids encoded by the BamHI site and the linking valine, followed by the two amino acids encoded by the XhoI site.

25 **SEO ID NO. 12:** Indicative expressed peptide.

MGSVPRYLKG WLKDVVQLSL RRPSFRASRQ RPIISLNERI LEFNKRNITA LE

[155] Example 6: Design and construction of a nucleic acid library of the invention that encodes short peptides based on amino acid sequences of proteins comprised in the proteomes of evolutionarily diverse organisms ("OmePEX").

30 **[156]** Another library of the invention was designed and constructed wherein the amino acid sequences of the source-proteins were those of naturally occurring proteins comprised in the reference proteomes of a yet more evolutionary diverse set of species, including a number of species from each of those being: archaea; bacteria, fungi, invertebrates, plants, protozoa, mammals and non-mammalian vertebrates.

[157] First, a mega protein amino acid sequence was generated as described in Example 1, but by using all the protein sequences comprised in 467 reference proteomes set out Table B. The second and third steps of Example 1 were conducted by analogy, to generate a pre-filtered set of over 1 million unique 46 amino acid long sequences.

[158] This set of pre-filtered sequences were subjected to hierarchical clustering using iterative runs of CD-HIT (Fu et al 2012, Bioinformatics 28:3150; <http://weizhongli-lab.org/cd-hit/>) with more stringent thresholds to obtain maximum diversity in the resulting peptides, described briefly as follows: Three rounds of clustering were performed; a first round used an 80% sequence similarity threshold parameter ("`-c 0.8`"; example command: `cdhit -i peptides.faa -o peptides_0.8.faa -c 0.8 -T 0 -M 32000 -n 5`); a second round, using the output of the first round, used a 60% sequence similarity threshold ("`-c 0.6`"; and a third and final run, using the output of the second round, used a sequence similarity threshold of 50% ("`-c 0.5`").

[159] The isoelectric point (pI) of each resulting peptide sequence remaining was predicted, and any having a predicted pI of between 6 and 8 was also removed from the set, as described in step four of Example 1.

[160] Analogously to as described in Example 2, the resulting set of amino acid sequences were further filtered to remove those predicted to have intrinsically disordered regions. In this case however, the top 475,000 sequences (ie, those least predicted to comprise intrinsically disordered regions) were selected to form a first set of unique 46 amino acid sequences.

[161] Separately, a second set of 25,000 unique 46 amino acid sequences were generated using the procedure described above in this Example 6, except that the amino acid sequences of the naturally occurring proteins first concatenated were those proteins having a known three-dimensional structure. In this case, about 10,150 polypeptide chains present in Protein Data Bank (<https://www.wwpdb.org>, on 5-Feb-2017) and that had a Pfam annotation (<http://pfam.xfam.org>, version 30.0).

[162] The first and second sets of amino acid sequences were combined, and 500,000 oligonucleotides, each encoding a separate of the amino acid sequences, was designed (having the same general structure) and synthesised as described in Example 5. The resulting synthesised oligonucleotides were PCR amplified, BamHI/XhoI digested and cloned into pMOST25A as described in Example 5 to form an expression library of the invention.

[163] **Example 7** [prophetic]: Design and construction of a nucleic acid library of the invention that encodes short peptides based on differentially expressed proteins ("DiffPEX").

[164] In this example, a query of a gene-expression database (e.g. the EMBL-EBI Expression Atlas; <http://www.ebi.ac.uk/gxa/home/>) is conducted to identify a sub-set of genes in the proteome that are differentially expressed between two tissue types. For example, the 5,000 most differentially expressed proteins between a human melanoma cell line (or patient sample) and a comparable but non-cancerous human cell line are identified by such a query.

[165] The reference amino acid sequences of this set of 5,000 proteins are used to generate a filtered set of over 20,000 unique 46 amino acid sequences and over 20,000 oligonucleotides, each encoding a separate of the amino acid sequences, are designed (having the same general structure), synthesised and cloned into pMOST25A as described in Example 5; *except* that in this case: (1) the window spacing along each protein's amino acid sequence is less than 10 amino acids so as to increase the density of tiling of the amino acid sequences across the sequence of these naturally occurring proteins; and (2) the SLIDER program (for prediction and filtering out of unstructured regions) is not used.

[166] **Example 8** [prophetic]: Generation of peptide libraries encoded by nucleic acid libraries of the invention.

[167] A library of peptides of the invention (eg, one encoded by a nucleic acid library of the invention) is generated as follows.

[168] First, the amplified and BamHI/XhoI-digested oligonucleotides of Example 7 are, instead, cloned into a CIS display construct (Odegrip et al 2003; PNAS 101:2806), having the general design:

Promoter_NUCLEIC_ACID_LIBRARY_repA_CIS_ori

[169] CIS display exploits the ability of a DNA replication initiator protein (RepA) to bind exclusively to the template DNA from which it has been expressed, a property called cis-activity. The peptide library is created by ligation of a nucleic acid of the invention to a DNA fragment that encodes RepA. After in vitro transcription and translation, a pool of protein-DNA complexes is formed where each protein is stably associated with the DNA that encodes it. These complexes are amenable to the affinity selection of ligands to targets of interest.

[170] CIS display, exploits the high-fidelity cis-activity that is exhibited by a group of bacterial plasmid DNA-replication initiation proteins typified by RepA of the R1 plasmid (Nikoletti et al 1988, J. Bacteriol. 170:1311). In this context, cis-activity refers to the property of the RepA family of proteins to bind exclusively to the template DNA from

which they have been expressed. R1 plasmid replication is initiated through the binding of RepA to the plasmid origin of replication (*ori*). *Ori* is separated from the RepA-coding sequence by a DNA element termed *CIS*. This element is thought to be critical in controlling the *cis*- activity of RepA (Masai & Arai 1988, Nucleic Acids Res. 16:6493). The consensus model for *cis*-activity is that the *CIS* element, which contains a rho- dependent transcriptional terminator, causes the host RNA polymerase to stall. This delay allows nascent RepA polypeptide emerging from translating ribosomes to bind transiently to *CIS*, which in turn directs the protein to bind to the adjacent *ori* site (Praszkier & Pittard 1999, J. Bacteriol. 181:2765).

[171] By genetically fusing peptide libraries to the N-terminus of the RepA protein, we can achieve a direct linkage of peptides to the DNA molecules that encode them; thus, the link between genotype to phenotype that is the common feature of display technologies is established.

[172] The peptide library is generated by in-vitro transcription and translation using an *E.coli* lysate system as described by Odegrip et al (2003), and solid phase selection for peptides (and hence the DNA sequence encoding them) that bind to immobilised target can be conducted (e.g., by one or more rounds of selection) also as described by Odegrip et al (2003).

[173] Example 9: Phenotypic screening with the lentiviral-cloned HuPEX library, and identification of SEP-binding targets.

[174] Using a phenotypic screen, the inventors were able to identify short-expressed-peptides (SEPs) from the HuPEX library described above that increased the survival of HeLa cells treated with methylnitronitrosoguanidine (MNNG), an inducer of cell death via parthanatos. Parthanatos, is a PARP-1 dependent form of programmed cell death (Yu et al, 2006; PNAS 103: 2653) that plays a role in neuronal cell death and is associated with diseases including Parkinson's disease, stroke, heart attack and diabetes.

[175] HEK293 cells were transfected with the pooled HuPEX nucleic acid library, cloned into the lentiviral vector as above, which is designed to express a plurality of SEPs. Virus was harvested, titered, and a batch of HeLa cells was infected with the SEP-expressing viruses and selected for 8 days. The library of virus-transduced HeLa cells was then exposed ("D0" time point) to a near-lethal dose (6.7uM) of MNNG. This dose was established as the maximum dose where co-incubation with the PARP-1 inhibitor olaparib was still able to rescue HeLa cells from parthanatos/cell-death.

[176] Genomic DNA was extracted from a control aliquot of such HeLa cells without MNNG treatment at D0, and large-scale amplicon DNA sequencing was performed to determine the relative abundance of HuPEX SEP-coding nucleic acids, and again from a second aliquot of such HeLa cells after culture for 8 days ("D8") in the presence of 6.7uM MNNG (Figure 4), so as to identify those expressed SEPS from the HuPEX library that showed an increased relative abundance at D8 compared to D0, and thus increased the survival of HeLa cells in the presence of MNNG.

[177] Of a predicted 300,000 different SEP-coding inserts represented in the HuPEX library, almost 288,000 were represented at least one member (as found by amplicon DNA sequencing) at D0; and by showing a significant increase in relative abundance at D8, at least 72 SEPs were identified from the HuPEX library as increasing the survival of HeLa cells in the presence of MNNG (Figure 5).

[178] Of these SEPs, a number were determined to be fragments of naturally occurring proteins that were associated with detoxification mechanisms. For example, a KEGG-pathway analysis (<http://www.kegg.jp/kegg/pathway.html>) determined that certain of such SEPs were fragments of naturally occurring proteins involved in "chemical carcinogenesis" and "metabolism of xenobiotics by cytochrome p450).

[179] Certain of the identified SEPs were used as a "bait" in yeast 2-hybrid screening technology (against a human cDNA library as prey) to identify the protein target(s) that the HuPEX SEP would bind to in HeLa cells.

[180] Example 10: Phenotypic screening with the lentiviral-cloned BugPEX and OmePEX libraries.

[181] The BugPEx and the OmePEx libraries were screened in a phenotypic screen of parthanatos, analogously to as described in Example 9, to identify SEPs expressed from each of these libraries that showed increased relative abundance at D8 compared to D0, and hence were able to increase the survival of HeLa cells in the presence of the parthanatos-inducing MNNG.

- 5 [182] Of a predicted 600,000 different SEP-coding inserts represented in the BugPEx library, almost 510,000 were represented at least one member (as found by amplicon DNA sequencing) at D0; and by showing a significant increase in relative abundance at D8, at least 58 SEPs were identified from the BugPEx library as increasing the survival of HeLa cells in the presence of MNNG (Figure 6).

- 10 [183] Of a predicted 500,000 different SEP-coding inserts represented in the OmePEx library, almost 490,000 were represented at least one member (as found by amplicon DNA sequencing) at D0; and by showing a significant increase in their relative abundance at D8, at least 64 SEPs were identified from the OmePEx library as increasing the survival of HeLa cells in the presence of MNNG (Figure 7).

[184] **Example 11:** Phenotypic screening with the lentiviral-cloned -PEx libraries

- 15 [185] Using a phenotypic screen, the inventors were able to identify a short-expressed-peptide (SEP) from the combined HuPEx (HPX), BugPEx (BPX) & OmePEx (OPX) libraries described above that decreased GFP-LC3 in HEK293FT cells engineered to express GFP-LC3/RFP-LC3DG Autophagic Flux Reporter (AFR cells, Kaizuka et al. Molecular Cell 2016).

- 20 [186] HEK293FT cells were transfected with the pooled HuPEx, BugPEx & OmePEx nucleic libraries, cloned into the lentiviral vector as described herein, for example pMOST25a as shown in SEQ ID No.10, which is designed to express a plurality of SEPs. Virus was harvested, titered, and a batch of HEK293FT-AFR cells was infected with the SEP-expressing viruses and selected for 4 days and SEP expressing cells were then expanded for a further 2 days without selection. The library of virus-transduced HEK293FT-AFR cells were then assessed by flow cytometry, SEP transduced HEK293FT-AFR cells enriched in the low GFP-LC3 gate, compared to unsorted controls, were flow-sorted and peptide sequences were amplified and sent to NGS analysis i.e. amplicon DNA sequencing as described in the previous example. Figures 8A-C show the population of selected hits (marked region) compared to control.

- 25 [187] SEPs sequences identified as being enriched in low GFP-LC3 gate were cloned into suitable lenti-viral expression and SEP-expressing virus generated in HEK293FT cells. Each SEP-expression virus population was individually assessed by flow cytometry in conditions described above. Each SEP expressing population was assessed in comparison to control-SEP expressing populations or uninfected HEK293FT-AFR cells treated with Torin1, an inducer of Autophagic Flux (Figure 9). A selection of candidates is shown with BPX-497507 representing a strong and robust hit able to induce Autophagy as measured by GFP-LC3 reduction. Torin1 (250nM) is shown as a positive control.
- 30

CLAIMS

1. **A library of nucleic acids**, each nucleic acid comprising a coding region of defined nucleic acid sequence encoding for a peptide having a length of between 25 and 110 amino acids, and having an amino acid sequence being a region of a sequence selected from the amino acid sequence of a naturally occurring protein of one or more organisms; wherein the library comprises nucleic acids that encode for a plurality of at least 10,000 different such peptides, and wherein the amino acid sequence of each of at least 50 of such peptides is a sequence region of the amino acid sequence of a different protein of a plurality of different such naturally occurring proteins, and wherein the isoelectric point (pI) of each encoded amino acid sequence is greater than 8.0 or less than 6.0.
2. The library of nucleic acids of any one of claims 1, wherein each of the plurality of different naturally occurring proteins fulfils one or more pre-determined criteria.
3. The library of nucleic acids of claim 2, wherein each of the plurality of naturally occurring proteins is associated with a given disease, such as cancer.
4. The library of nucleic acids of claim 3, wherein the disease is breast cancer.
5. The library of nucleic acids of claim 2, wherein each of the plurality of naturally occurring proteins is a cytoplasmic protein.
6. The library of nucleic acids of claim 5, wherein each of the plurality of naturally occurring proteins is a cytoplasmic kinase.
7. The library of nucleic acids of claim 2, wherein each of the plurality of naturally occurring proteins interacts with a given protein or at least one protein from a (functional) class of proteins.
8. The library of nucleic acids of claim 7, wherein each of the plurality of naturally occurring proteins interacts with KRas.
9. The library of nucleic acids of any one of claims 1 to 8, wherein the library comprises nucleic acids that encode for a plurality of at least 50,000 different such peptides, and wherein the amino acid sequence of each of at least 100 of such peptide is a sequence region of the amino acid sequence of at least 100 different naturally occurring proteins; in particular wherein the library comprises nucleic acids that encode for a plurality of at least 100,000 different such peptides, and wherein the amino acid sequence of each of at least 150 of such peptide is a sequence region of the amino acid sequence of at least 150 different naturally occurring proteins.
10. The library of nucleic acids of any one of claims 1 to 9, wherein the library comprises nucleic acids that encode for a plurality of at least 10,000 different such peptides, and wherein the amino acid sequence of each of at least 1,000 of such peptides is a sequence region of the amino acid sequence of a different protein of such plurality of different naturally occurring proteins.
11. The library of nucleic acids of any one of claims 1 to 10, wherein the library comprises nucleic acids that encode for a plurality of at least 200,000 different such peptides, and wherein the amino acid sequence of each of at least 20,000 of such peptide is a sequence region of the amino acid sequence of at least 20,000 different naturally occurring proteins; in particular wherein the library comprises nucleic acids that encode for a plurality of at least 300,000 different such peptides, and wherein the amino acid sequence of each of at

- least 25,000 of such peptide is a sequence region of the amino acid sequence of at least 25,000 different naturally occurring proteins.
12. The library of nucleic acids of any one of claims 1 or 11, wherein that in respect of at least about 1% of the naturally occurring proteins a plurality of the nucleic acids encodes for different peptides from the amino acid sequences of such naturally occurring proteins.
13. The library of nucleic acids of claim 12, wherein that in respect of at least about 50% of the naturally occurring proteins a plurality of the nucleic acids encodes for different peptides from the amino acid sequences of such naturally occurring proteins.
14. The library of nucleic acids of claim 13, wherein the plurality of the nucleic acids encodes for different peptides, and the amino acid sequences of which are sequence regions spaced along the amino acid sequence of the naturally occurring protein.
15. The library of nucleic acids of claim 14, wherein the sequence regions are spaced by a window of amino acids apart, or by multiples of such window, along the amino acid sequence of the naturally occurring protein wherein, the window is between 1 and about 55 amino acids; in particular wherein the window is between about 5 and about 20 amino acids; most particularly wherein the window of spacing is about 8, 10, 12 or 15 amino acids.
16. The library of nucleic acids of any one of claims 1 to 14 comprising nucleic acids encoding for at least 100,000 different peptides from at least 10,000 different naturally occurring proteins.
17. The library of nucleic acids of any one of claims 1 to 16, wherein each nucleic acid encodes a different peptide.
18. The library of nucleic acids of any one of claims 1 to 17, wherein the mean number of nucleic acids that encode a different peptide from the naturally occurring proteins is greater than 1; in particular between about 1.01 and 1.5 such nucleic acids (peptides) per such protein.
19. The library of nucleic acids of claim 18, wherein the mean number of nucleic acids that encode a different peptide from the naturally occurring proteins is at least about 5 such nucleic acids (peptides) per such protein, in particular wherein the mean is between about 5 and about 2,000 such nucleic acids (peptides) per such protein or is between about 5 and about 1,000 nucleic acids (peptides) per such protein.
20. The library of nucleic acids of claim 19, wherein the mean number of nucleic acids that encode a different peptide from the naturally occurring proteins is between about 100 and about 1,500 such nucleic acids (peptides) per such protein or is between about 250 and about 1,000 such nucleic acids (peptides) per such protein.
21. The library of nucleic acids of claim 19, wherein the mean number of nucleic acids that encode a different peptide from the naturally occurring proteins is between about 5 and about 100 such nucleic acids (peptides) per such protein or is between about 5 and about 50 such nucleic acids (peptides) per such protein.
22. The library of nucleic acids of any one of claims 1 to 21, wherein the amino acid sequence of the naturally occurring protein is one selected from the group of amino acids sequences of non-redundant proteins comprised in a reference proteome, suitably, the reference proteome is one or more of the reference proteomes selected from the group of reference proteomes listed in Table A and/or Table B, or an updated version of such reference proteome.

23. The library of nucleic acids of any one of claims 1 to 22, wherein the amino acid sequences of the plurality of encoded peptides are sequence regions selected from amino acid sequences of naturally occurring proteins (or polypeptide chains or domains thereof) with a known three-dimensional structure; in particular wherein the naturally occurring protein (or polypeptide chain or domain thereof) is comprised in the Protein Data Bank, and optionally that has a Pfam annotation.
24. The library of nucleic acids of any one of claims 22 to 23, wherein the sequence region selected from the amino acid sequence of the protein does not include an ambiguous amino acid of such amino acid sequence comprised in the reference proteome or the Protein Data Bank.
25. The library of nucleic acids according to any preceding claim, wherein the library is for expression in a mammalian cell, preferably a human cell.
26. The library of nucleic acids according to any preceding claim, wherein the library is cloned into a lentiviral vector or a retroviral vector.
27. **A library of peptides** encoded by the library of nucleic acids of any one of claims 1 to 26 .
28. **A method of identifying a target protein that modulates a phenotype of a mammalian cell**, said method comprising:
- exposing a population of in vitro cultured mammalian cells capable of displaying said phenotype to a library of nucleic acids according to any of claims 1-26 or a library of peptides according to claim 27,
 - identifying in said cell population an alteration in said phenotype following said exposure,
 - selection of said cells undergoing the phenotypic change and identifying a peptide encoded by (or a peptide of) such library that alters the phenotype of the cell,
 - providing said peptide and identifying the cellular protein that binds to said peptide, said cellular protein being a target protein that modulates the phenotype of the mammalian cell.
29. The method according to claim 28, wherein the method includes a further step of identifying a compound that binds to said target protein and displaces or blocks binding of said peptide, wherein the compound modulates the phenotype of a mammalian cell.
30. **Use** of: (a) a library of nucleic acids according to any of claims 1-26; and/or (b) a library of peptides according to claim 27, **to identify a peptide that binds to a target.**
31. Use according to claim 30, wherein the target is a protein target.
32. Use according to claim 30 or claim 31, wherein the identified peptide modulates a phenotype of a mammalian cell.
33. **Use** of: (a) a library of nucleic acids according to any of claims 1-26; and/or (b) a library of peptides according to claim 27, **to identify a compound** which binds to a target.
34. Use according to claim 33, wherein the target is a protein target.
35. Use according to claim 33 or claim 34 wherein, said compound displaces or blocks binding of a peptide to the target.

36. Use according to any of claims 33 to 35, wherein, the peptide and/or the compound modulates a phenotype of a mammalian cell.
37. A method according to claim 28 or claim 29, or Use according to claims 30 to 36, wherein the phenotype is a phenotype related to the modulation of a cell-signalling pathway.
- 5 38. A method or use according to claim 37, wherein the method or use comprises the identification of peptides which modulate cell-signalling pathways and the identification of protein targets and surface sites on such proteins that participate in signal transduction.
39. A method or use according to claim 37 or 38, wherein the cell-signalling pathway is active or altered in cancer cells.