

JAK2 INHIBITORS AND THEIR USE FOR THE TREATMENT OF MYELOPROLIFERATIVE DISEASES AND CANCER

FIELD OF THE INVENTION

5 **[0001]** This invention relates to novel compounds that are useful as anti-cancer/antiproliferative agents. This invention also relates to a method of using the compounds in the treatment of proliferative diseases, such as cancer, and to pharmaceutical compositions containing the compounds.

10 BACKGROUND OF THE INVENTION


[0002] Efforts to identify new therapeutic approaches to *Ph(-)* myeloproliferative disease have been bolstered by the observations of constitutive activation of the JAK-Stat signaling pathway MPD patients. In particular, a single valine to phenylalanine mutation at residue 617 (JAK2-V617F) in JAK2 has been observed in the majority of
15 PV (95%), ET (50-60%) and PMF (50-60%) patients (Table 2, Kralovics et al., 2005; Baxter et al., 2005; Tefferi et al., 2005). The V617F mutation resides in the region of the *JAK2* gene encoding the pseudokinase domain which is thought to function as an autoinhibitory domain to regulate JAK2 tyrosine kinase activity. Mutations in exon 12 of JAK2 which also result in constitutive JAK2 kinase activity are also observed
20 with a lesser frequency (<5% in PV and ET) and are mutually exclusive with JAK2V617F lesions (Pardanari et al., 2007; Scott et al., 2007). JAK2 is a member of a non-receptor tyrosine kinase family that also includes JAK1, Tyk2 and JAK3 and function as mediators of cytokine receptor signaling (for review see Murray, 2007). Upon cytokine binding to its cognate receptor, receptor-bound JAK family members
25 are activated and phosphorylate a STAT, a latent transcription factor, which upon JAK-mediated phosphorylation undergoes dimerization and translocation to the nucleus to regulate gene expression. Genetic and biochemical studies have established distinct combinations of engagement of a JAK family member with an individual cytokine receptor. For instance, erythropoietin (EPO), thrombopoietin
30 (TPO) and granulocyte colony stimulating factor (GM-CSF) receptor engagement results in the predominant activation of JAK2 to mediate downstream signaling. Consistent with the patho-physiology of MPD associated with the JAK2-V617F

- mutation, these cytokines promote the differentiation and expansion of the cell types underlying PV, ET and PMF, respectively. Unlike other genetic activation events, expression of JAK2-V617F is not sufficient to promote transformation in cell based model systems and has been shown to require the co-expression of the type I cytokine receptors, highlighting an important functional co-dependence of the JAK-cytokine receptor interaction (Lu et al., 2005). Interestingly, activating mutations in the TPO receptor (MPL, tryptophan to leucine substitution at residue 515) have been identified in MPD patients afflicted with JAK2-V617F *negative* PMF and ET (5% and 1%, respectively) resulting in constitutive JAK2-Stat activation (Pikman et al., 2006).
- These observations indicate that JAK-Stat signal transduction can be activated through mutation in MPD at multiple points in the pathway in a mutually exclusive manner and suggest the possible presence of additional pathway mutations in JAK2 – V617F and MPL-W515L negative MPD.
- [0003]** Important validation of JAK2 signaling as a driver of *Ph(-)* MPD emerged from rodent models where JAK2-V617F mutant signaling was reconstituted in the hematopoietic stem cell compartment. Several laboratories demonstrated that viral transduction of JAK2-V617F into mouse bone marrow and subsequent re-implantation into recipient mice reconstituted several aspects human MPD (Wernig et al. 2006, Lacout et al., 2006, Bumm et al., 2006, Zaeleskas et al., 2006). These features included elevated hematocrit, splenomegaly from extramedullary hematopoiesis, granulocytosis and bone marrow fibrosis all which are also manifested in polycythemia vera. Interestingly, unlike the human condition, thrombocytosis was not observed in these murine models and was suggested to be attributable to secondary genetic events that contribute to platelet expansion (Wernig et al., 2006).
- Similar reconstitution of the TPO receptor mutation (MPL-W515L) in rodent bone marrow resulted in a myeloproliferative disease with a more rapid onset than JAK2-V617F animals that was reminiscent of primary myelofibrosis including splenomegaly, hepatomegaly, and reticulin fibrosis of bone marrow (Pikman et al., 2006). Also unlike the JAK2-V617F model, mice expressing MPL-W515L displayed dramatic thrombocytosis perhaps indicating a more dominant function of the receptor activation compared to JAK2-V617 in the expansion of this lineage. Nonetheless,

these observations collectively underscore the role of both MPL-W515L and JAK2-V617F as driver mutations underlying the progression of human MPD.

[0004] A key question to the genetic basis of MPD is the role of additional genetic events that contribute to disease progression beyond JAK2 and MPL. Several lines of evidence suggest additional genetic alterations in MPD disease progression. In fact, mitotic recombination occurs frequently in MPD patients to generate two JAK2-V617F alleles indicating a selection for cell clones homozygous for the mutated kinase (Levine et al., 2005). In this regard it will be important to develop conditional JAK2-V617F knock-in animals and to determine the phenotypic consequences of the homozygous versus heterozygous JAK2-V617F burden. Additionally, there is evidence for an inherited germline allele that precedes and predisposes patients to acquire JAK2-V617F (Goerttler et al., 2005; Levine et al., 2006) as well as loss of chromosomal region 20q in some MPD patients. Although MPD conversion AML is observed clinically at moderate levels and activating JAK chromosomal translocations are observed in leukemia, epidemiological data suggest that it is questionable that JAK2-V617F is a genetic driver in this context suggesting additional genetic alterations are required for full leukemic transformation (Theocharides et al., 2007). These observations notwithstanding, JAK2 inhibition with small molecule inhibitors is sufficient to modulate disease progression in pre-clinical animal models suggesting that JAK2 activation is sufficient for maintaining MPD (Paradani et al., 2007). It will be important to identify these additional genetic alterations and decipher how these genetic changes contribute to the disease progression of PV, ET and PMF in the context of JAK2-V617F and MPL-W515L. It will also be important to implement approaches to identify if other JAK2 pathway components are mutated in MPD patients not associated with the acquisition of JAK2-V617F, JAK2 exon12 or MPL-W515L mutations.

[0005] The patent publication WO2006/122137 discloses compounds which are useful as IKK inhibitors. Example No. A171 discloses the compound of the formula




which was found to have weak activity against JAK2 in the assay described hereinbelow.

DETAILED DESCRIPTION OF THE INVENTION

5 [0006] The present invention provides for compounds of formula I, pharmaceutical compositions employing such compounds and for methods of using such compounds.

[0007] In accordance with the present invention, there are disclosed compounds of formula I




(I)

or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein:

Y is C₁₋₄ alkyl;

X is C₁₋₄ alkyl;

R is



any of which are optionally fused with a 5 or 6 membered carbocycle or heterocycle having one heteroatom selected from NR³ or S, said fused carbocycle or heterocycle being optionally substituted with 0-3 R¹.

[0008] R¹ is H, halo, CN, C₁₋₆ alkyl substituted with 0-3 R^c, CF₃, CONR^aR^a, NR^aR^a, COOR^b, SO₂-C₁₋₄alkyl, C(O)R^d, cycloalkyl substituted with 0-3 R^c, furanyl, tetrahydropyranyl, or pyridinyl;

R^2 is absent, H, C_{1-6} alkyl substituted with 0-3 R^c , $C(O)O-C_{1-4}$ alkyl, SO_2-C_{1-4} alkyl, cycloalkyl substituted with 0-3 R^c , or tetrahydropyranyl;

R^3 is absent, H, or $C(O)O-C_{1-4}$ alkyl;

R^a is H, C_{1-6} alkyl substituted with 0-3 R^c , C_{3-6} cycloalkyl substituted with 0-3 R^c , tetrahydropyranyl, or dioxotetrahydrothiophenyl;

R^b is H or C_{1-6} alkyl;

R^c is H, halo, CN, OH, $O-C_{1-4}$ alkyl, $O-C_{1-4}$ alkyl- $O-C_{1-4}$ alkyl, NH_2 , $N(C_{1-4}$ alkyl) $_2$, $C(O)N(C_{1-4}$ alkyl) $_2$, SO_2-C_{1-4} alkyl, or morpholinyl or piperazinyl, either of which are optionally substituted with 0-1 C_{1-4} alkyl;


R^d is C_{1-6} alkyl, or azetidiny, azetidiny, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, dioxidothiomorpholinyl or tetrahydropyranyl, any of which are substituted with 0-2 R^c ; and

R^e is H, halo, CN, C_{1-4} alkyl, OH, $O-C_{1-4}$ alkyl, SO_2-C_{1-4} alkyl, $NHC(O)-C_{1-4}$ alkyl, morpholinyl, $OC(O)-C_{1-4}$ alkyl, $C(O)N(C_{1-4}$ alkyl) $_2$, or $O-C_{1-4}$ alkyl- $O-C_{1-4}$ alkyl.

15

[0009] In another embodiment are compounds of Formula (I), wherein:

R is



any of which are optionally substituted with 0-3 R^1 .


[0010] In another embodiment are compounds of Formula (I), wherein:

- 5 Y is methyl; and
 X is ethyl.

[0011] In another embodiment are compounds of Formula (I), wherein:


R is

10




[0012] In another embodiment are compounds of Formula (I), wherein:

15 R is



any of which are optionally substituted with 0-2 R^1

[0013] In another embodiment are compounds of Formula (I), wherein



R¹ is H, halo, CN, C₁₋₆ alkyl substituted with 0-3 R^c, CF₃, CONR^aR^a, COOR^b,
5 SO₂-C₁₋₄alkyl, C(O)R^d, cycloalkyl substituted with 0-3 R^c, or pyridinyl;

R^a is H, C₁₋₆ alkyl substituted with 0-3 R^c, C₃₋₆ cycloalkyl substituted with 0-3 R^c, tetrahydropyranyl or dioxotetrahydrothiophenyl;


R^b is H or C₁₋₆ alkyl;

R^c is H, halo, OH, O-C₁₋₄alkyl, SO₂-C₁₋₄alkyl or morpholinyl;

10 R^d is C₁₋₆ alkyl, or azetidiny, pyrrolidinyl, morpholinyl, piperazinyl or dioxidothiomorpholinyl, any of which are substituted with 0-2 R^c;

R^c is H, halo, CN, OH, O-C₁₋₄alkyl, SO₂-C₁₋₄alkyl, NHC(O)-C₁₋₄alkyl or morpholinyl.

15 [0014] In another embodiment are compounds of Formula (I), wherein:



R¹ is H, halo, C₁₋₆ alkyl substituted with 0-3 R^c, CF₃, CONR^aR^a, COOR^b,
C(O)R^d, cycloalkyl substituted with 0-3 R^c or furanyl;

20 R² is H, C₁₋₆ alkyl substituted with 0-3 R^c, SO₂-C₁₋₄alkyl, cycloalkyl substituted with 0-3 R^c, or tetrahydropyranyl;

R^a is H, or C₁₋₆ alkyl substituted with 0-3 R^c;


R^b is H or C₁₋₆ alkyl;

25 R^c is H, halo, CN, OH, O-C₁₋₄alkyl, O-C₁₋₄alkyl-O-C₁₋₄alkyl, NH₂, N(C₁₋₄alkyl)₂, C(O)N(C₁₋₄alkyl)₂, SO₂-C₁₋₄alkyl, or morpholinyl or piperazinyl, either of which are optionally substituted with 0-1 C₁₋₄alkyl;

R^d is C₁₋₆ alkyl, or morpholinyl, piperazinyl or dioxidothiomorpholinyl, any of which are substituted with 0-2 R^c; and

R^c is H, C_{1-4} alkyl, CN, OH, $NHC(O)-C_{1-4}$ alkyl or morpholinyl.

[0015] In another embodiment are compounds of Formula (I), wherein:



- 5 R^1 is C_{1-6} alkyl substituted with 0-3 R^c ; and
 R^2 is C_{1-6} alkyl.

[0016] In another embodiment are compounds of Formula (I), wherein the compound of formula (I) is selected from the exemplified compounds.

- 10 **[0017]** In another embodiment the present invention provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and one or more of the compounds of the present invention or a stereoisomer, tautomer, pharmaceutically acceptable salt, solvate, or prodrug thereof.

- [0018]** In another embodiment, the present invention provides a pharmaceutical
 15 composition comprising a compound of formula I or pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

- [0019]** In another embodiment, the present invention provides a pharmaceutical composition, comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of one or more of the compounds of the present invention or a
 20 stereoisomer, tautomer, pharmaceutically acceptable salt, solvate or prodrug thereof.

- [0020]** In another embodiment, the present invention provides a method for treating proliferative disorders and/or cancer comprising administering to a patient in need of such treatment a therapeutically effective amount of one or more of the compounds of the present invention or a stereoisomer, tautomer, pharmaceutically acceptable salt, solvate
 25 or prodrug thereof, in combination with one or more other anti-cancer agent, or a stereoisomer, tautomer, pharmaceutically acceptable salt, solvate or prodrug thereof.

[0021] In another embodiment, the present invention provides a method for treating proliferative disorders and/or cancer comprising administering to a patient in need of such treatment a therapeutically effective amount of one or more of the compounds of the

present invention or a stereoisomer, tautomer, pharmaceutically acceptable salt, solvate or prodrug thereof.

[0022] In another embodiment, the present invention provides a method of treating a patient in need of proliferative disorder and/or cancer treatment, comprising
5 administering a compound of the present invention or a stereoisomer, tautomer, pharmaceutically acceptable salt, solvate, or prodrug form thereof in an amount effective to treat the proliferative disorder and/or cancer.

[0023] In another embodiment, the present invention provides a method of treating myeloproliferative diseases, such as polycythemia vera, essential thrombocytopenia and
10 primary myelofibrosis, solid tumors of the pancreas, prostate, lung, head and neck, breast, colon, ovary, gastric cancer as well as other tumor types including multiple myeloma, melanoma, neuroblastoma, glioblastoma, systemic mastocytosis, and hematological malignancies such as acute myelogenous leukemia (including refractory acute myeloid leukemia) and acute lymphoid leukemia.

15 [0024] In another embodiment, the present invention provides a novel method, comprising administering a compound of the present invention or a stereoisomer, tautomer, pharmaceutically acceptable salt, solvate, or prodrug form thereof in an amount effective to treat a proliferative disorder and/or cancer.

[0025] In another embodiment, the present invention provides a method for treating
20 proliferative disorders and/or cancer comprising administering to a patient in need of such treatment a therapeutically effective amount of a compound of formula (I) or a stereoisomer, tautomer, pharmaceutically acceptable salt, solvate or prodrug thereof, in combination with one or more other anti-cancer agent or antiproliferative agent and/or another agent, or a stereoisomer, tautomer, pharmaceutically acceptable salt, solvate or
25 prodrug thereof.

[0026] In another embodiment, the present invention provides a method for treating proliferative disorders and/or cancer comprising administering to a patient in need of such treatment a therapeutically effective amount of one or more of the compounds of the present invention or a stereoisomer, tautomer, pharmaceutically acceptable salt, solvate
30 or prodrug thereof, in combination with one or more other anti-cancer agent, or a stereoisomer, tautomer, pharmaceutically acceptable salt, solvate or prodrug thereof.

[0027] In another embodiment, the present invention provides a compound of the present invention for use in therapy.

[0028] In another embodiment, the present invention provides a compound of the present invention for use in therapy for treating a proliferative disorder and/or cancer.

5 [0029] In another embodiment, the present invention also provides the use of a compound of formula I of the present invention for the manufacture of a medicament for the treatment of a proliferative disorder and/or cancer.

[0030] The present invention may be embodied in other specific forms without departing from the spirit or essential attributes thereof. This invention encompasses all
10 combinations of preferred aspects and/or embodiments of the invention noted herein. It is understood that any and all embodiments of the present invention may be taken in conjunction with any other embodiment or embodiments to describe additional more preferred embodiments. It is also to be understood that each individual element of the preferred embodiments is its own independent preferred embodiment. Furthermore, any
15 element of an embodiment is meant to be combined with any and all other elements from any embodiment to describe an additional embodiment.

DEFINITIONS

[0031] The following are definitions of terms that may be used in the present
20 specification. The initial definition provided for a group or term herein applies to that group or term throughout the present specification individually or as part of another group, unless otherwise indicated.

[0032] Compounds of this invention may have one or more asymmetric centers. Unless otherwise indicated, all chiral (enantiomeric and diastereomeric) and racemic
25 forms of compounds of the present invention are included in the present invention. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds, and all such stable isomers are contemplated in the present invention. Cis and trans geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. The present
30 compounds can be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically active starting materials. All chiral, (enantiomeric and

diastereomeric) and racemic forms and all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomer form is specifically indicated.

[0033] The term “substituted,” as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided
5 that the designated atom’s normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced. Keto substituents are not present on aromatic moieties. Ring double bonds, as used herein, are double bonds that are formed between two adjacent ring atoms (e.g., C=C, C=N, or N=N).

10 [0034] When any variable (e.g., R^3) occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R^3 , then said group may optionally be substituted with up to two R^3 groups and R^3 at each occurrence is selected independently from the definition of R^3 . Also, combinations
15 of substituents and/or variables are permissible only if such combinations result in stable compounds.

[0035] When a bond to a substituent is shown to cross a bond connecting two atoms in a ring, then such substituent may be bonded to any atom on the ring. When a substituent is listed without indicating the atom via which such substituent is bonded to
20 the rest of the compound of a given formula, then such substituent may be bonded via any atom in such substituent. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

[0036] In cases wherein there are nitrogen atoms (e.g., amines) on compounds of the present invention, these can be converted to N-oxides by treatment with an oxidizing
25 agent (e.g., MCPBA and/or hydrogen peroxides) to afford other compounds of this invention. Thus, all shown and claimed nitrogen atoms are considered to cover both the shown nitrogen and its N-oxide ($N \rightarrow O$) derivative.

[0037] As used herein, the term “alkyl” or “alkylene” is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified
30 number of carbon atoms. For example, “ C_{1-10} alkyl” (or alkylene), is intended to include C_1 , C_2 , C_3 , C_4 , C_5 , C_6 , C_7 , C_8 , C_9 , and C_{10} alkyl groups. Additionally, for example, “ C_1 - C_6 alkyl” denotes alkyl having 1 to 6 carbon atoms. Alkyl groups can be unsubstituted or

substituted so that one or more of its hydrogens are replaced by another chemical group. Example alkyl groups include, but are not limited to, methyl (Me), ethyl (Et), propyl (*e.g.*, n-propyl and isopropyl), butyl (*e.g.*, n-butyl, isobutyl, t-butyl), pentyl (*e.g.*, n-pentyl, isopentyl, neopentyl), and the like.

- 5 **[0038]** “Alkenyl” or “alkenylene” is intended to include hydrocarbon chains of either straight or branched configuration and having one or more double carbon-carbon bonds that may occur in any stable point along the chain. For example, “C₂₋₆ alkenyl” (or alkenylene), is intended to include C₂, C₃, C₄, C₅, and C₆ alkenyl groups. Examples of alkenyl include, but are not limited to, ethenyl, 1-propenyl, 2-
10 propenyl, 2-butenyl, 3-butenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 2-methyl-2-propenyl, 4-methyl-3-pentenyl, and the like.

- 15 **[0039]** “Alkynyl” or “alkynylene” is intended to include hydrocarbon chains of either straight or branched configuration and having one or more triple carbon-carbon bonds that may occur in any stable point along the chain. For example, “C₂₋₆ alkynyl” (or
alkynylene), is intended to include C₂, C₃, C₄, C₅, and C₆ alkynyl groups; such as ethynyl, propynyl, butynyl, pentynyl, hexynyl and the like.

[0040] “Halo” or “halogen” includes fluoro, chloro, bromo, and iodo.

- 20 **[0041]** The term “cycloalkyl” refers to cyclized alkyl groups, including mono-, bi- or poly-cyclic ring systems. C₃₋₁₀ cycloalkyl is intended to include C₃, C₄, C₅, C₆, and C₇ cycloalkyl groups. Example cycloalkyl groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, norbornyl, and the like. As used
herein, “carbocycle” or “carbocyclic residue” is intended to mean any stable 3, 4, 5, 6, or 7-membered monocyclic or bicyclic or 7, 8, 9, 10, 11, 12, or 13-membered bicyclic or
25 tricyclic ring, any of which may be saturated, partially unsaturated, unsaturated or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclobutenyl, cyclopentyl, cyclopentenyl, cyclohexyl, cycloheptenyl, cycloheptyl, cycloheptenyl, adamantyl, cyclooctyl, cyclooctenyl, cyclooctadienyl,
30 [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane, [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, anthracenyl, and tetrahydronaphthyl (tetralin). As shown above, bridged rings are also included in the definition of carbocycle (*e.g.*, [2.2.2]bicyclooctane). Preferred carbocycles, unless otherwise specified,

are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, and indanyl. When the term “carbocycle” is used, it is intended to include “aryl”. A bridged ring occurs when one or more carbon atoms link two non-adjacent carbon atoms. Preferred bridges are one or two carbon atoms. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring may also be present on the bridge.

[0042] The term “aryl” refers to monocyclic or bicyclic aromatic hydrocarbon groups having 6 to 12 carbon atoms in the ring portion, such as phenyl, naphthyl, biphenyl and diphenyl groups, each of which may be substituted.

[0043] The term “substituted aryl” refers to an aryl group substituted by, for example, one to four substituents such as alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, arylalkyl, halo, trifluoromethoxy, trifluoromethyl, hydroxy, alkoxy, alkanoyl, alkanoyloxy, aryloxy, arylalkyloxy, amino, alkylamino, arylamino, arylalkylamino, dialkylamino, alkanoylamino, thiol, alkylthio, ureido, nitro, cyano, carboxy, carboxyalkyl, carbamyl, alkoxycarbonyl, alkylthiono, arylthiono, arylsulfonamide, sulfonic acid, alkylsulfonyl, sulfonamido, aryloxy and the like. The substituent may be further substituted by hydroxy, halo, alkyl, alkoxy, alkenyl, alkynyl, aryl or arylalkyl.

[0044] The term “heteroaryl” refers to an optionally substituted, aromatic group for example, which is a 4 to 7 membered monocyclic, 7 to 11 membered bicyclic, or 10 to 15 membered tricyclic ring system, which has at least one heteroatom and at least one carbon atom-containing ring, for example, pyridine, tetrazole, indazole.

[0045] As used herein, the term “heterocycle”, “heterocyclyl”, “heterocyclic system” or “heterocyclic group” is intended to mean a stable 5, 6, or 7- membered monocyclic or bicyclic or 7, 8, 9, 10, 11, 12, 13, or 14-membered bicyclic heterocyclic ring which is saturated, partially unsaturated or fully unsaturated, and which consists of carbon atoms and 1, 2, 3 or 4 heteroatoms independently selected from N, O and S; and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may optionally be oxidized (i.e., N→O and S(O)_p). The nitrogen atom may be substituted or unsubstituted (i.e., N or NR wherein R is H or another substituent, if defined). The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure. The

heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. A nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. When the term "heterocycle" is used, it is intended to include heteroaryl.

[0046] Examples of heterocycles include, but are not limited to, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzoxazolyl, benzoxazoliny, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazoliny, carbazolyl, 4*aH*-carbazolyl, carboliny, chromanyl, chromenyl, cinnoliny, decahydroquinoliny, 2*H*,6*H*-1,5,2-dithiaziny, dihydrofuro[2,3-*b*]tetrahydrofuran, furanyl, furazanyl, imidazolidiny, imidazoliny, imidazolyl, 1*H*-indazolyl, indolenyl, indoliny, indoliziny, indolyl, 3*H*-indolyl, isatinoyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindoliny, isoindolyl, isoquinoliny, isothiazolyl, isothiazolopyridiny, isoxazolyl, isoxazolopyridiny, methylenedioxyphenyl, morpholiny, naphthyridiny, octahydroisoquinoliny, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidiny, oxazolyl, oxindolyl, pyrimidiny, phenanthridiny, phenanthroliny, phenaziny, phenothiaziny, phenoxathiny, phenoxaziny, phthalaziny, piperaziny, piperidiny, piperidony, 4-piperidonyl, piperonyl, pteridiny, puriny, pyranly, pyraziny, pyrazolidiny, pyrazoliny, pyrazolyl, pyridaziny, pyridoazole, pyridoimidazole, pyridothiazole, pyridiny, pyrimidiny, pyrrolidiny, pyrroliny, 2-pyrrolidonyl, 2*H*-pyrrolyl, pyrrolyl, quinazoliny, quinoliny, 4*H*-quinoliziny, quinoxaliny, quinuclidiny, tetrahydrofuranyl, tetrahydroisoquinoliny, tetrahydroquinoliny, tetrazolyl, 6*H*-1,2,5-thiadiaziny, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triaziny, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, and xanthenyl. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

[0047] Preferred 5- to 10-membered heterocycles include, but are not limited to, pyridiny, furanyl, thienyl, pyrrolyl, pyrazolyl, pyraziny, piperaziny, piperidiny, imidazolyl, imidazolidiny, indolyl, tetrazolyl, isoxazolyl, morpholiny, oxazolyl,

oxadiazolyl, oxazolidinyl, tetrahydrofuranyl, thiadiazinyl, thiadiazolyl, thiazolyl, triazinyl, triazolyl, benzimidazolyl, 1*H*-indazolyl, benzofuranyl, benzothiofuranyl, benztetrazolyl, benzotriazolyl, benzisoxazolyl, benzoxazolyl, oxindolyl, benzoxazolinyl, benzthiazolyl, benzisothiazolyl, isatinoyl, isoquinolinyl, octahydroisoquinolinyl, 5 tetrahydroisoquinolinyl, tetrahydroquinolinyl, isoxazolopyridinyl, quinazolinyl, quinolinyl, isothiazolopyridinyl, thiazolopyridinyl, oxazolopyridinyl, imidazolopyridinyl, and pyrazolopyridinyl.

[0048] Preferred 5 to 6 membered heterocycles include, but are not limited to, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, pyrazinyl, piperazinyl, piperidinyl, 10 imidazolyl, imidazolidinyl, indolyl, tetrazolyl, isoxazolyl, morpholinyl, oxazolyl, oxadiazolyl, oxazolidinyl, tetrahydrofuranyl, thiadiazinyl, thiadiazolyl, thiazolyl, triazinyl, and triazolyl. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

[0049] In another embodiment, heterocycles include, but are not limited to, pyridyl, 15 pyridinyl, isoxazolyl, isoquinolinyl, thienyl, pyrazolyl, furanyl, pyrrolyl, thiazolyl, imidazolyl, pyrazinyl, thiadiazolyl, pyrimidinyl, pyridazinyl, oxazolyl, isothiazolyl, oxadiazolyl, indanonyl, piperazinyl, pyranyl, or pyrrolyl

[0050] Also included are smaller heterocycles, such as, epoxides and aziridines.

[0051] As used herein, the term “aromatic heterocyclic group” or “heteroaryl” is 20 intended to mean a stable monocyclic and polycyclic aromatic hydrocarbons that include at least one heteroatom ring member such as sulfur, oxygen, or nitrogen. Heteroaryl groups include, without limitation, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, furyl, quinolyl, isoquinolyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuryl, benzothienyl, benzthiazolyl, isoxazolyl, pyrazolyl, triazolyl, tetrazolyl, 25 indazolyl, 1,2,4-thiadiazolyl, isothiazolyl, purinyl, carbazolyl, benzimidazolyl, indolinyl, benzodioxolanyl, benzodioxane, and the like. Heteroaryl groups can be substituted or unsubstituted. The nitrogen atom may be substituted or unsubstituted (i.e., N or NR wherein R is H or another substituent, if defined). The nitrogen and sulfur heteroatoms may optionally be oxidized (i.e., N→O and S(O)_p). It is to be noted that total number of 30 S and O atoms in the aromatic heterocycle is not more than 1. Bridged rings are also included in the definition of heterocycle. A bridged ring occurs when one or more atoms (i.e., C, O, N, or S) link two non-adjacent carbon or nitrogen atoms. Preferred bridges

include, but are not limited to, one carbon atom, two carbon atoms, one nitrogen atom, two nitrogen atoms, and a carbon-nitrogen group. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring may also be present on the bridge.

5 **[0052]** The term “carbocyclic ring” or “carbocycle” refers to stable, saturated, partially saturated or unsaturated, mono or bicyclic hydrocarbon rings that contain 3-12 atoms. Particularly, this includes a monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms. Suitable values include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, dihydroindenyl and tetrahydronaphthyl. The term
10 “optionally substituted” as it refers to “carbocyclic ring” or “carbocycle” herein indicates that the carbocyclic ring may be substituted at one or more substitutable ring positions by one or more groups independently selected from alkyl (preferably lower alkyl), alkoxy (preferably lower alkoxy), nitro, monoalkylamino (preferably a lower alkylamino), dialkylamino (preferably a di[lower]alkylamino), cyano, halo, haloalkyl (preferably
15 trifluoromethyl), alkanoyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkyl amido (preferably lower alkyl amido), alkoxyalkyl (preferably a lower alkoxy[lower]alkyl), alkoxycarbonyl (preferably a lower alkoxycarbonyl), alkylcarbonyloxy (preferably a lower alkylcarbonyloxy) and aryl (preferably phenyl), said aryl being optionally substituted by halo, lower alkyl and lower
20 alkoxy groups.

[0053] The term “heteroatoms” shall include oxygen, sulfur and nitrogen.

[0054] The compounds of formula I may exist as a free form (with no ionization) or may form salts which are also within the scope of this invention. Pharmaceutically acceptable (i.e. non-toxic, physiologically acceptable) salts are preferred, although other
25 salts are also useful, e.g., in isolating or purifying the compounds of this invention.

[0055] The compounds of formula I may form salts with alkali metals such as sodium, potassium and lithium, with alkaline earth metals such as calcium and magnesium, with organic bases such as dicyclohexylamine, tributylamine, pyridine and amino acids such as arginine, lysine and the like. Such salts can be formed as known to
30 those skilled in the art.

[0056] The compounds for formula I may form salts with a variety of organic and inorganic acids. Such salts include those formed with hydrogen chloride, hydrogen

bromide, methanesulfonic acid, sulfuric acid, acetic acid, trifluoroacetic acid, oxalic acid, maleic acid, benzenesulfonic acid, toluenesulfonic acid and various others (e.g., nitrates, phosphates, borates, tartrates, citrates, succinates, benzoates, ascorbates, salicylates and the like). Such salts can be formed as known to those skilled in the art.

5 **[0057]** In addition, zwitterions (“inner salts”) may be formed.

[0058] All stereoisomers of the compounds of the instant invention are contemplated, either in admixture or in pure or substantially pure form. The definition of compounds according to the invention embraces all the possible stereoisomers and their mixtures. It very particularly embraces the racemic forms and the isolated optical isomers having the
10 specified activity. The racemic forms can be resolved by physical methods, such as, for example, fractional crystallization, separation or crystallization of diastereomeric derivatives or separation by chiral column chromatography. The individual optical isomers can be obtained from the racemates from the conventional methods, such as, for example, salt formation with an optically active acid followed by crystallization.

15 **[0059]** The present invention is intended to include all isotopes of atoms occurring in the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include deuterium and tritium. Isotopes of carbon include ¹³C and ¹⁴C. Isotopically-labeled compounds of the invention can generally
20 be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described herein, using an appropriate isotopically-labeled reagent in place of the non-labeled reagent otherwise employed.

[0060] Compounds of the formula I may also have prodrug forms. Since prodrugs are known to enhance numerous desirable qualities of pharmaceuticals (e.g., solubility,
25 bioavailability, manufacturing, etc.) the compounds of the present invention may be delivered in prodrug form. Thus, the present invention is intended to cover prodrugs of the presently claimed compounds, methods of delivering the same and compositions containing the same. “Prodrugs” are intended to include any covalently bonded carriers that release an active parent drug of the present invention *in vivo* when such prodrug is
30 administered to a mammalian subject. Prodrugs of the present invention are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent

compound. Prodrugs include compounds of the present invention wherein a hydroxy, amino, or sulfhydryl group is bonded to any group that, when the prodrug of the present invention is administered to a mammalian subject, it cleaves to form a free hydroxyl, free amino, or free sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate, and benzoate derivatives of alcohol and amine functional groups in the compounds of the present invention.

[0061] Various forms of prodrugs are well known in the art. For examples of such prodrug derivatives, see:

- a) *Design of Prodrugs*, edited by H. Bundgaard, (Elsevier, 1985) and *Methods in Enzymology*, Vol. 112, pp. 309-396, edited by K. Widder, et al. (Academic Press, 1985);
- b) *A Textbook of Drug Design and Development*, edited by Krosgaard-Larsen and H. Bundgaard, Chapter 5, "Design and Application of Prodrugs," by H. Bundgaard, pp. 113-191 (1991); and
- c) H. Bundgaard, *Advanced Drug Delivery Reviews*, 8, 1-38 (1992).

[0062] It should further be understood that solvates (e.g., hydrates) of the compounds of formula I are also within the scope of the present invention. Methods of solvation are generally known in the art.

[0063] "Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent. It is preferred that the presently recited compounds do not contain a N-halo, S(O)₂H, or S(O)H group.

[0064] As used herein, "treating" or "treatment" cover the treatment of a disease-state in a mammal, particularly in a human, and include: (a) preventing the disease-state from occurring in a mammal, in particular, when such mammal is predisposed to the disease-state but has not yet been diagnosed as having it; (b) inhibiting the disease-state, i.e., arresting its development; and/or (c) relieving the disease-state, i.e., causing regression of the disease state.

[0065] "Therapeutically effective amount" is intended to include an amount of a compound of the present invention that is effective when administered alone or in combination. "Therapeutically effective amount" is also intended to include an amount of the combination of compounds which is effective for the treatment of disease.

[0066] The present invention further includes compositions comprising one or more compounds of the present invention and a pharmaceutically acceptable carrier.

[0067] A “pharmaceutically acceptable carrier” refers to media generally accepted in the art for the delivery of biologically active agents to animals, in particular, mammals.

- 5 Pharmaceutically acceptable carriers are formulated according to a number of factors well within the purview of those of ordinary skill in the art. These include, without limitation: the type and nature of the active agent being formulated; the subject to which the agent-containing composition is to be administered; the intended route of administration of the composition; and the therapeutic indication being targeted.
- 10 Pharmaceutically acceptable carriers include both aqueous and non-aqueous liquid media, as well as a variety of solid and semi-solid dosage forms. Such carriers can include a number of different ingredients and additives in addition to the active agent, such additional ingredients being included in the formulation for a variety of reasons, *e.g.*, stabilization of the active agent, binders, etc., well known to those of ordinary skill
- 15 in the art. Descriptions of suitable pharmaceutically acceptable carriers, and factors involved in their selection, are found in a variety of readily available sources such as, for example, *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA, 1985, which is incorporated herein by reference in its entirety.

20

UTILITY

- [0068] The present invention is based on the discovery that certain compounds are inhibitors of protein kinases. More specifically, compounds such as those described in this invention inhibit the protein tyrosine kinase activity of members of the JAK family of receptors. These inhibitors will be useful in the treatment of proliferative diseases
- 25 that are dependent on signaling by one or more of these receptors. Such diseases include myeloproliferative diseases, solid tumors of the pancreatic, prostate, lung, head and neck, breast, colon, ovary, as well as other tumor types including multiple myeloma, melanoma, neuroblastoma, glioblastoma and hematological malignancies such as acute myelogenous leukemia.
- 30 [0069] The invention also relates to a pharmaceutical composition of compound of formula I, or pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier in the treatment of hyperproliferative disorder in mammal. In particular, said

pharmaceutical composition is expected to inhibit the growth and/or metastasis of those primary and recurrent solid tumors which are associated with Flt-3 (Fms-like kinase-3), JAK2, JAK3, and JAK1, especially those tumors which are significantly dependent on JAK2, for their growth and spread, including for example, cancers of the blood, thyroid, breast, colon, pancreas, or a variety of tumor types including multiple myeloma, melanoma, neuroblastoma and glioblastoma. In addition to JAK2 activity, inhibitory activity against JAK3/JAK1 or TYK2 may be useful in the treatment of certain cancers having an inflammatory component.

[0070] Thus according to a further aspect of the invention there is provided the use of a compound of the formula I, or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the production of an antiproliferative effect in a warm-blooded animal such as a human being.

[0071] According to a further feature of the invention there is provided a method for producing an antiproliferative effect in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof as defined herein before.

[0072] By virtue of their ability to inhibit Flt-3, JAK2, and JAK3, kinases, compounds of the present invention can be used for the treatment of proliferative diseases, including cancer.

[0073] Thus, the present invention provides methods for the treatment of a variety of cancers, including, but not limited to, the following:

carcinoma including that of the bladder (including accelerated and metastatic bladder cancer), breast, colon (including colorectal cancer), kidney, liver, lung (including small and non-small cell lung cancer and lung adenocarcinoma), ovary, prostate, testes, genitourinary tract, lymphatic system, rectum, larynx, pancreas (including exocrine pancreatic carcinoma), esophagus, stomach, gall bladder, cervix, thyroid, and skin (including squamous cell carcinoma);

hematopoietic tumors of lymphoid lineage including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma, histiocytic lymphoma, and Burkett's lymphoma;

hematopoietic tumors of myeloid lineage including acute and chronic myelogenous leukemias, myelodysplastic syndrome, myeloid leukemia, and promyelocytic leukemia;

5 tumors of the central and peripheral nervous system including astrocytoma, neuroblastoma, glioma, and schwannomas;

tumors of mesenchymal origin including fibrosarcoma, rhabdomyosarcoma, and osteosarcoma; and

other tumors including melanoma, xenoderma pigmentosum, keratoactanthoma, seminoma, thyroid follicular cancer, and teratocarcinoma.

10 **[0074]** The present invention provides methods for the treatment of leukemia, myeloproliferative diseases such as polycythemia vera, essential thrombocytopenia and myelofibrosis, multiple myeloma, colon cancer, breast cancer, and gastric cancer.

[0075] The anti-proliferative treatment defined herein before may be applied as a sole therapy or may involve, in addition to a compound of the invention, one or more other
15 substances and/or treatments. Such treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. The compounds of this invention may also be useful in combination with known anti-cancer and cytotoxic agents and treatments, including radiation. If formulated as a fixed dose, such combination products employ the compounds of this
20 invention within the dosage range described below and the other pharmaceutically active agent within its approved dosage range. Compounds of formula I may be used sequentially with known anticancer or cytotoxic agents and treatment, including radiation when a combination formulation is inappropriate.

[0076] The term “anti-cancer” agent includes any known agent that is useful for the
25 treatment of cancer including the following: 17 α -ethinylestradiol, diethylstilbestrol, testosterone, prednisone, fluoxymesterone, dromostanolone propionate, testolactone, megestrolacetate, methylprednisolone, methyl-testosterone, prednisolone, triamcinolone, chlorotrianisene, hydroxyprogesterone, aminoglutethimide, estramustine, medroxyprogesteroneacetate, leuprolide, flutamide, toremifene, Zoladex; matrix
30 metalloproteinase inhibitors; VEGF inhibitors, such as anti-VEGF antibodies (Avastin[®]) and small molecules such as ZD6474 and SU6668; Vatalanib, BAY-43-9006, SU11248, CP-547632, and CEP-7055; HER 1 and HER 2 inhibitors including anti-HER2

- antibodies (Herceptin); EGFR inhibitors including gefitinib, erlotinib, ABX-EGF, EMD72000, 11F8, and cetuximab; Eg5 inhibitors, such as SB-715992, SB-743921, and MKI-833; pan Her inhibitors, such as canertinib, EKB-569, CI-1033, AEE-788, XL-647, mAb 2C4, and GW-572016; kinase inhibitors, e.g. Gleevec[®] and dasatinib (Sprycel[®]);
- 5 Casodex[®] (bicalutamide, Astra Zeneca), Tamoxifen; MEK-1 kinase inhibitors, MAPK kinase inhibitors, PI3 kinase inhibitors; PDGF inhibitors, such as imatinib; anti-angiogenic and antivasular agents which, by interrupting blood flow to solid tumors, render cancer cells quiescent by depriving them of nutrition; castration, which renders androgen dependent carcinomas non-proliferative; inhibitors of non-receptor and
- 10 receptor tyrosine kinases; inhibitors of integrin signaling; tubulin acting agents such as vinblastine, vincristine, vinorelbine, vinflunine, paclitaxel, docetaxel, 7-O-methylthiomethylpaclitaxel, 4-desacetyl-4-methylcarbonatepaclitaxel, 3'-*tert*-butyl-3'-N-*tert*-butyloxycarbonyl-4-deacetyl-3'-dephenyl-3'-N-debenzoyl-4-O-methoxycarbonyl-paclitaxel, C-4 methyl carbonate paclitaxel, epothilone A, epothilone B, epothilone C,
- 15 epothilone D, desoxyepothilone A, desoxyepothilone B, [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7-11-dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-aza-17 oxabicyclo [14.1.0]heptadecane-5,9-dione (ixabepilone), [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-3-[2-[2-(aminomethyl)-4-thiazolyl]-1-methylethenyl]-7,11-dihydroxy-8,8,10,12,16-pentamethyl-
- 20 4-17-dioxabicyclo[14.1.0]-heptadecane-5,9-dione, and derivatives thereof; CDK inhibitors, antiproliferative cell cycle inhibitors, epidophyllotoxin, etoposide, VM-26; antineoplastic enzymes, e.g., topoisomerase I inhibitors, camptothecin, topotecan, SN-38; procarbazine; mitoxantrone; platinum coordination complexes such as cisplatin, carboplatin and oxaliplatin; biological response modifiers; growth inhibitors;
- 25 antihormonal therapeutic agents; leucovorin; tegafur; antimetabolites such as purine antagonists (e.g. 6-thioguanine and 6-mercaptopurine; glutamine antagonists, e.g. DON (AT-125; d-oxo-norleucine); ribonucleotide reductase inhibitors; mTOR inhibitors; and haematopoietic growth factors.
- [0077] Additional cytotoxic agents include cyclophosphamide, doxorubicin,
- 30 daunorubicin, mitoxanthrone, melphalan, hexamethyl melamine, thiotepa, cytarabin, idatrexate, trimetrexate, dacarbazine, L-asparaginase, bicalutamide, leuprolide, pyridobenzoindole derivatives, interferons, and interleukins.

[0078] In the field of medical oncology it is normal practice to use a combination of different forms of treatment to treat each patient with cancer. In medical oncology the other component(s) of such treatment in addition to the antiproliferative treatment defined herein before may be surgery, radiotherapy or chemotherapy. Such

5 chemotherapy may cover three main categories of therapeutic agent:

(i) antiangiogenic agents that work by different mechanisms from those defined hereinbefore (for example, linomide, inhibitors of integrin $\alpha v \beta 3$ function, angiostatin, razoxane);

(ii) cytostatic agents such as antiestrogens (for example, tamoxifen, toremifene, raloxifene, droloxifene, idoxifene), progestogens (for example, megestrol acetate), aromatase inhibitors (for example, anastrozole, letrozole, bora-zole, exemestane), antihormones, antiprogestogens, antiandrogens (for example, flutamide, nilutamide, bicalutamide, cyproterone acetate), LHRH agonists and antagonists (for example, gosereline acetate, leuprolide), inhibitors of testosterone 5 α -dihydroreductase (for
10 example, finasteride), farnesyltransferase inhibitors, anti-invasion agents (for example, metalloproteinase inhibitors such as marimastat and inhibitors of urokinase plasminogen activator receptor function) and inhibitors of growth factor function, (such growth factors include for example, EGF, FGF, platelet derived growth factor and hepatocyte growth factor, such inhibitors include growth factor antibodies, growth factor receptor antibodies
15 such as Avastin[®] (bevacizumab) and Erbitux[®] (cetuximab); tyrosine kinase inhibitors and serine/threonine kinase inhibitors); and

(iii) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as antimetabolites (for example, antifolates such as methotrexate, fluoropyrimidines such as 5-fluorouracil, purine and adenosine analogues, cytosine
25 arabinoside); Intercalating antitumour antibiotics (for example, anthracyclines such as doxorubicin, daunomycin, epirubicin and idarubicin, mitomycin-C, dactinomycin, mithramycin); platinum derivatives (for example, cisplatin, carboplatin); alkylating agents (for example, nitrogen mustard, melphalan, chlorambucil, busulphan, cyclophosphamide, ifosfamide nitrosoureas, thiotepa; antimitotic agents (for example,
30 vinca alkaloids like vincristine, vinorelbine, vinblastine and vinflunine) and taxoids such as Taxol[®] (paclitaxel), Taxotere[®] (docetaxel) and newer microtubule agents such as epothilone analogs (ixabepilone), discodermolide analogs, and eleutherobin analogs;

topoisomerase inhibitors (for example, epipodophyllotoxins such as etoposide and teniposide, amsacrine, topotecan, irinotecan); cell cycle inhibitors (for example, flavopyridols); biological response modifiers and proteasome inhibitors such as Velcade[®] (bortezomib).

5 **[0079]** As stated above, the formula I compounds of the present invention are of interest for their antiproliferative effects. Such compounds of the invention are expected to be useful in a wide range of disease states including cancer, psoriasis, and rheumatoid arthritis.

10 **[0080]** More specifically, the compounds of formula I are useful in the treatment of a variety of cancers, including (but not limited to) the following:

- carcinoma, including that of the prostate, pancreatic ductal adeno-carcinoma, breast, colon, lung, ovary, pancreas, and thyroid;
- tumors of the central and peripheral nervous system, including neuroblastoma, glioblastoma, and medulloblastoma;
- 15 – hematological malignancies such as acute myelogenous leukemia (AML), and
- other tumors, including melanoma and multiple myeloma.

20 **[0081]** Due to the key role of kinases in the regulation of cellular proliferation in general, inhibitors could act as reversible cytostatic agents which may be useful in the treatment of any disease process which features abnormal cellular proliferation, e.g., benign prostate hyperplasia, familial adenomatosis polyposis, neuro-fibromatosis, pulmonary fibrosis, arthritis, psoriasis, glomerulonephritis, restenosis following angioplasty or vascular surgery, hypertrophic scar formation and inflammatory bowel disease.

25 **[0082]** The compounds of formula I are especially useful in treatment of tumors having a high incidence of tyrosine kinase activity, such as prostate, colon, brain, thyroid and pancreatic tumors. By the administration of a composition (or a combination) of the compounds of this invention, development of tumors in a mammalian host is reduced.

[0083] Compounds of formula I may also be useful in the treatment of other cancerous diseases (such as acute myelogenous leukemia).

30 **[0084]** The pharmaceutical compositions of the present invention containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches,

lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs.

[0085] When a compound according to this invention is administered into a human subject, the daily dosage will normally be determined by the prescribing physician with the dosage generally varying according to the age, weight, sex and response of the individual patient, as well as the severity of the patient's symptoms.

[0086] If formulated as a fixed dose, such combination products employ the compounds of this invention within the dosage range described above and the other pharmaceutically active agent or treatment within its approved dosage range.

10 Compounds of formula I may also be administered sequentially with known anticancer or cytotoxic agents when a combination formulation is inappropriate. The invention is not limited in the sequence of administration; compounds of formula I may be administered either prior to or after administration of the known anticancer or cytotoxic agent(s).

[0087] The compounds may be administered in a dosage range of about 0.05 to 200 mg/kg/day, preferably less than 100 mg/kg/day, in a single dose or in 2 to 4 divided doses.

DOSAGE AND FORMULATION

[0088] The compounds of this invention can be administered in such oral dosage forms as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. They may also be administered in intravenous (bolus or infusion), intraperitoneal, subcutaneous, or intramuscular form, all using dosage forms well known to those of ordinary skill in the pharmaceutical arts. They can be administered alone, but generally will be administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

[0089] The dosage regimen for the compounds of the present invention will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration; the species, age, sex, health, medical condition, and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment; the frequency of treatment; the route of administration, the renal and hepatic function of the patient,

and the effect desired. A physician or veterinarian can determine and prescribe the effective amount of the drug required to treat the cancer.

- [0090]** By way of general guidance, the daily oral dosage of each active ingredient, when used for the indicated effects, will range between about 0.001 to 1000 mg/kg of body weight, preferably between about 0.001 to 100 mg/kg of body weight per day, and most preferably between about 0.001 to 20 mg/kg/day. Intravenously, the most preferred doses will range from about 0.1 to about 10 mg/kg. Compounds of this invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three, or four times daily.
- 10 **[0091]** Compounds of this invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using transdermal skin patches. When administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.
- 15 **[0092]** The compounds are typically administered in admixture with suitable pharmaceutical diluents, excipients, or carriers (collectively referred to herein as pharmaceutical carriers) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.
- 20 **[0093]** For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the like; for oral administration in liquid form, the oral drug components can be
- 25 combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents, and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as
- 30 acacia, tragacanth, or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium

chloride, and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, and the like.

[0094] The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large
5 unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

[0095] Compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include
polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol,
10 polyhydroxyethylaspartamidephenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid,
15 polyorthoesters, polyacetals, polydihydropyrans, polycyanoacylates, and crosslinked or amphipathic block copolymers of hydrogels.

[0096] Dosage forms (pharmaceutical compositions) suitable for administration may contain from about 1 milligram to about 1000 milligrams of active ingredient per dosage unit. In these pharmaceutical compositions the active ingredient will
20 ordinarily be present in an amount of about 0.1-95% by weight based on the total weight of the composition.

[0097] Gelatin capsules may contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and
25 capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.

[0098] Liquid dosage forms for oral administration can contain coloring and
30 flavoring to increase patient acceptance.

[0099] In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols

are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl-or propyl-paraben, and chlorobutanol.

[00100] Suitable pharmaceutical carriers are described in *Remington's Pharmaceutical Sciences*, Mack Publishing Company, a standard reference text in this field.

BIOLOGICAL ASSAYS

JAK2 Tyrosine Kinase Assay

[00101] The assays were performed in V-bottom 384-well plates. The final assay volume was 30 μ L prepared from 15 μ L additions of enzyme and substrates (fluoresceinated peptide and ATP) and test compounds in assay buffer (100 mM HEPES pH 7.4, 10 mM $MgCl_2$, 25 mM Beta-Glycerolphosphate, 0.015% Brij35 and 4 mM DTT). The reaction was initiated by the combination of JAK2 with substrates and test compounds. The reaction was incubated at room temperature for 60 min. and terminated by adding 45 μ L of 35 mM EDTA to each sample. The reaction mixture was analyzed on the Caliper LabChip 3000 by electrophoretic separation of the fluorescent substrate and phosphorylated product. Inhibition data were calculated by comparison to no enzyme control reactions for 100% inhibition and vehicle-only reactions for 0% inhibition. The final concentration of reagents in the assays is ATP, 30 μ M; JAK2 fluorescent peptide, 1.5 μ M; JAK2, 1 nM; and DMSO, 1.6%. Dose response curves were generated to determine the concentration required inhibiting 50% of kinase activity (IC_{50}). Compounds were dissolved at 10 mM in dimethylsulfoxide (DMSO) and evaluated at eleven concentrations, each in duplicate. IC_{50} values were derived by non-linear regression analysis

JAK3 Tyrosine Kinase Assay

[00102] The assays were performed in V-bottom 384-well plates. The final assay volume was 30 μ L prepared from 15 μ L additions of enzyme and substrates (fluoresceinated peptide and ATP) and test compounds in assay buffer (100 mM HEPES pH 7.4, 10 mM $MgCl_2$, 25 mM Beta-Glycerolphosphate, 0.015% Brij35 and 4 mM DTT). The reaction was initiated by the combination of JAK3 with substrates and test compounds. The reaction was incubated at room temperature for 60 min. and terminated by adding 45 μ L of 35 mM EDTA to each sample. The reaction mixture was analyzed on the Caliper LabChip 3000 by electrophoretic separation of the fluorescent substrate and phosphorylated product. Inhibition data were calculated by comparison to no enzyme control reactions for 100% inhibition and vehicle-only reactions for 0% inhibition. The final concentration of reagents in the assays is ATP, 8 μ M; JAK3 fluorescent peptide, 1.5 μ M; JAK3, 2.5 nM; and DMSO, 1.6%. Dose response curves were generated to determine the concentration required inhibiting 50% of kinase activity (IC_{50}). Compounds were dissolved at 10 mM in dimethylsulfoxide (DMSO) and evaluated at eleven concentrations, each in duplicate. IC_{50} values were derived by non-linear regression analysis.

Cell Proliferation Inhibition Assay

[00103] Compounds were evaluated for their ability to inhibit cell proliferation, using an assay that measures thymidine incorporation in the DNA of the dividing cell, which is directly correlated with cell numbers. SET-2 cells were plated at 10000 cells/well in 96-well plates and were cultured for 24 h in RPMI-1640 supplemented with 20% fetal bovine serum, before test compounds were added. Compounds were diluted in culture medium such that the final concentration of dimethyl sulfoxide never exceeded 0.1%. Following the addition of compounds, the cells were cultured for an additional 72 h before cell proliferation was determined by measuring DNA synthesis with 3H -thymidine incorporation.

[00104] Compounds described herein were tested in the JAK2 and Cell Proliferation Inhibition assay described above. The following results were obtained.

TABLE 1

Example#	JAK2 (IC ₅₀ , uM)	SET-2 (IC ₅₀ , uM)
3	1.50E-03	0.01
4	0.03	2.53
15	1.52E-03	0.49
29	0.01	10.00
33	6.37E-04	0.01
34	1.50E-03	0.24

Example#	JAK2 (IC50, μM)	SET-2 (IC50, μM)
35	5.36E-04	2.85
38	0.04	0.06
49	0.05	3.93
53	1.47E-03	0.10
55	7.48E-04	4.30E-03
61	7.51E-04	

Example#	JAK2 (IC50, μ M)	SET-2 (IC50, μ M)
70	7.33E-04	0.33
73	1.00E-03	0.06
81	0.01523	0.339
98	0.000695	0.007365
101	7.43E-04	10.00
109	1.52E-03	4.15

Example#	JAK2 (IC50, μ M)	SET-2 (IC50, μ M)
116	1.49E-03	0.17
125	0.05	1.27

[00105] Compounds described herein have been tested in one or more of the above identified assays and have been found to be active.

5


C. *In vivo* assay

[00106] Compound formulated in appropriate vehicle was administered to mice either as an oral bolus or injected intraperitoneally or subcutaneously to Balb/C mice. Blood samples were collected through orbital bleeding or cardiac puncture at various time points post compound administration. Aliquots (90 μ L) of the blood were placed in a 96 well plate. Diluted solution (5ul) of murine thrombopoietin (mTPO; Peprotech) (diluted 1:1000 in dPBS/BSA, 2 μ L in 2ml dPBS/BSA was delivered to each well for pathway stimulation. The plates were incubated at 37°C in a water bath for 10 minutes. 1.5ml of fix/lyse solution (from BD Biosciences) was added to each well. The samples were then incubated at 37° C for 15 minutes to ensure lysis of red blood cells (RBCs). The plates were centrifuged at 2000 rpm for 5 minutes and aspirated using the V&P 96 well aspirator. The cells were washed with dPBS, re-centrifuged, re-suspended and transferred to a regular 96 well plate. The cells were washed twice with dPBS followed by addition of CD61 FITC (from eBiosciences) for surface receptor staining for 45 min in the dark at room temperature. The cells were then washed in dPBS and permeabilized for pSTAT5 staining as follows: 200 μ L of

permeabilization buffer III (from BD Bioscience) was added and the samples were incubated on ice for 30 minutes and then samples were washed twice with dPBS/1%BSA and re-suspended in 100 μ L of dPBS/BSA containing Stat5 (pY695) Alexa 647 antibodies. The samples were incubated at room temperature in the dark
5 for 45 minutes, washed twice and re-suspended in 10 μ L of the Stat5 (pY695) Alexa 647 antibodies used were added to the sample. The samples were re-suspended in 200 μ L of dPBS/BSA and run on the FACSCanto on sample images were processed using Diva 6 and FlowJo 8.5.3 analysis software.


11495 PCT

Data for Example 26



11495 PCT

In vivo data for Example 73



ABBREVIATIONS


[00107] The following abbreviations may be employed in the methods of preparation and Examples:

- 5 h = hours
DCM = dichloromethane
THF = tetrahydrofuran
HPLC = high performance liquid chromatography
DIEA = diisopropylethyl amine
- 10 i-PrOH = isopropyl alcohol
TFA = trifluoroacetic acid
min = minutes
DMF = dimethylformamide
EDC = N-(3-Dimethylaminopropyl)N'-ethylcarbodiimide
- 15 HOBt = hydroxybenzotriazole
NMP = N-methylpyrrolidinone
EtOAc = ethyl acetate
AcOH = acetic acid
BOP reagent = benzotriazole-1-yl-oxy-tris-(dimethylamino)-
- 20 phasphoniumhexafluorophosphate
brine = saturated aqueous sodium chloride solution
Et₃N = triethylamine
t_R = retention time
rt – room temperature
- 25 NCS = N-chlorosuccinamide
NBS = N-bromosuccinamide
NIS = N-iodosuccinamide

30

Methods of Preparation


[00108] Compounds of general formula I in which the R group is thiazole (as in Ia1) and R¹ and R² groups are CF₃ or alkyl or cycloalkyl or combine to form a saturated carbocyclic or heterocyclic ring or where R² group is COOR^b could be prepared using the general method depicted in Scheme 1. Dichloro intermediate II (prepared using procedure reported in WO200612237) could be combined with a 2,4-dimethoxybenzyl and the resulting secondary amine is capped with suitable protective group (Boc) (III). The second chlorine atom could be converted into the corresponding amine (IV) through the benzophenone imine intermediate. The amino compound could be halogenated to intermediate V. V could be subjected to transition metal mediated indole ring formation and the resulting indole nitrogen is capped with ethyl iodide to afford VI. Ester hydrolysis followed by amide bond formation and cleavage of protective groups with acid treatment would yield amine VII. Amine VII could be converted into thiourea VIII by first coupling with benzoyl isothiocyanate followed treatment with aqueous base. Formation of thiazole could be achieved by condensation with an α -bromoketone derivative (R¹CHBrCOR²).



- a) 2,4-dimethoxybenzylamine, heat; b) NaHMDS, Boc₂O; c) (Ph)₂=NH; d) HCl;
 e) NIS; f) Pd₂(dba)₃, ethyl pyruvate; g) EtI, Cs₂CO₃; h) NaOH (aq); i)
 5 dicyclopropylamine HCl, HATU, DIPEA; j) TFA; k) Benzoyl isothiocyanate;
 l) NaOH (aq); m) R²CHBrCOR¹

Scheme 1

- 10 [00109] Compounds of general formula Ia2 in which the R¹ group is CONR^aR^a could be made using Scheme 2. Thiourea intermediate (VIII) could be combined with EtO₂CCHBrCOR¹ to afford the thiazole ester (IX). The ester could be hydrolyzed and the acid could be coupled with amine to afford thiazole amide derivative (Ia)




a) $\text{EtO}_2\text{CCHBrCOR}^1$; b) NaOH (aq) ; c) HNR^aR^a , HATU, DIPEA

5

Scheme 2

[00110] Similarly, compounds of general formula Ia3 in which the R^1 group is CONR^aR^a could be prepared using the general protocol depicted in Scheme 3.

10




a) $\text{R}^2\text{CHBrCOCO}_2\text{Me}$; b) NaOH (aq) ; c) HNR^aR^a , HATU, DIPEA

15

Scheme 3


[00111] Compounds of general formula Ia in which R^1 is halogen (Cl, Br or I) could be prepared by condensing an α,α' -dihaloketone as depicted in Scheme 4.



a) $R^2COCH(Hal)_2$

Scheme 4

- 5 **[00112]** Alternatively, thiourea derivative VIII could be converted to room temperature into C-5 un-substituted thiazole XI and then directly halogenated using electrophilic halogen source or through metallation followed by quenching with an electrophilic halogenating agent (Scheme 5).




10

a) $BrCH_2COR^2$; b) Selectfluor or NCS or NBS or NIS or tBuLi followed Selectfluor or NBS or NCS

Scheme 5

15

[00113] Compounds of general formula Ia5 in which R^1 is SO_2R^b could be synthesized using the general synthetic approach shown in Scheme 6




a) Br₂-acetic acid; b) EtOH, heat

Scheme 6

5

[00114] Compounds with general formula Ia in which R¹ and R² combine to form an aromatic or heteroaromatic ring could be prepared using Scheme 7.




10

a) Pd(0) catalyst, NaOtBu, phosphine ligand, heat

Scheme 7

15 **[00115]** Alternatively, these compounds could be made by first coupling aniline or heteroaniline (XVI) with the isothiocyanate (XV) followed by oxidative cyclization (Scheme 8).




a) 1,1'-Thiocarbonyldi-2(1H)-pyridone; b) NaH; c) NIS

Scheme 8

5

[00116] Compounds of general formula Ib1 could be prepared using the general synthetic approach depicted in Scheme 9. Aniline VII could be combined with γ -dithiomethylketone compound XVII, (prepared using the procedure reported at room temperature in *Synlett*, p 2331 (2008)) under basic condition to afford XVIII.


10 Stepwise condensation of the Boc-protected hydrazine derivative would give the required pyrazole Ib1.



15 a) NaH, THF; b) R¹N(Boc)NH₂, AcOH, 35-40°C; c) HCO₂H or TFA, 60°C

Scheme 9


[00117] Compounds of general formula Ib1 or If1 and 1f could also be prepared by coupling C-4 halo derivative (XIX) with an appropriately substituted 2-aminopyrazole derivative (XX) using a transition metal catalyzed reaction (Scheme 10).



a) isoamyl nitrite, CH_2I_2 or isoamyl nitrite, CH_2Br_2 ; b) $\text{Pd}_2(\text{dba})_3$, Xanphos, Cs_2CO_3

Scheme 10

[00118] Compounds of general formula Ib2 in which R^2 group is CONR^aR^a could be synthesized using Scheme 11. Aniline VII could be combined with γ -dithiomethylketone derivative XXII, (prepared using the procedure from *Tetrahedron*, p 2631 (2003)) to afford intermediate XXIII. Stepwise condensation of Boc-protected hydrazine derivative would give the required pyrazole aldehyde XXIV. Aldehyde could be oxidized using oxone or sodium hypochlorite to furnish carboxylic acid XXV. Coupling of acid XXV with amine would give pyrazole amide Ib2.




a) NaH, THF, heat; b) $R^1N(Boc)NH_2$, AcOH; c) TFA; d) oxone or sodium hypochlorite; e) HNR^aR^a , HATU, DIPEA

5

Scheme 11

[00119] Compounds of general formula Ic1 could be prepared using the general protocol as shown in Scheme 12. Aniline VII could be coupled with chloroacetyl
 10 chloride and the resulting amide could be treated with thioamide (R^2CSNH_2) to furnish thiazole Ic1.




a) chloroacetyl chloride, base; b) R^2CSNH_2

15


Scheme 12

[00120] Compounds of general formula 1d1 could be made as per Scheme 13. Previously described isothiocyanate derivative XV could be combined with amidine XXV under dehydrating reaction conditions to give 1,2,4-thiadiazole (1d1).




Scheme 13

[00121] Compounds of general formula 1e1 could be prepared using a synthetic approach as shown in Scheme 14. Isothiocyanate XV could be combined with azide XXVI in the presence of phosphine to yield 1,3-oxazole 1e1.




Scheme 14

[00122] Compounds of general formula 1g1 could be prepared using a synthetic approach as shown in Scheme 15. Amine VII could be combined with acyl isothiocyanate XXVII. The acylthiourea could be condensed with hydrazine derivative to yield the 1,2,4-triazol derivative 1g1.




Scheme 15

5

EXAMPLES**Example 1**

10 ***N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

1A Preparation of 6-chloro-*N*-(2,4-dimethoxybenzyl)-1-methyl-1*H*-imidazo[4,5-*c*]pyridin-4-amine



15

[00123] In a 200 mL round bottom flask, 4,6-dichloro-1-methyl-1*H*-imidazo[4,5-*c*]pyridine (15 g, 74.2 mmol) (prepared as per WO2006122137 example A1.5) was dissolved in 2,4-dimethoxybenzylamine (25 mL, 166 mmol). The reaction mixture


was heated to 110 °C for 3 h and cooled to room temperature. Water was added and a tan colored solid precipitated out. Vacuum filtration yielded a tan solid. The solid was washed with ethyl acetate and hexane to yield 6-chloro-*N*-(2,4-dimethoxybenzyl)-1-methyl-1*H*-imidazo[4,5-*c*]pyridin-4-amine (23.459 g, 95 % yield) as a cream colored solid.

[00124] MS (ESI) *m/z* 333.0 (M+H)

[00125] ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 8.05 (s, 1 H), 7.17 (t, 1 H, *J*=6.19 Hz), 7.06 (d, 1 H, *J*=8.25 Hz), 6.87 (s, 1 H), 6.55 (d, 1 H, *J*=2.47 Hz), 6.42 (dd, 1 H, *J*=8.52, 2.47 Hz), 4.54 (d, 2 H, *J*=5.77 Hz), 3.82 (s, 3 H), 3.74 (s, 3 H), 3.72 (s, 3 H)

10

1B Preparation of *tert*-butyl 6-chloro-1-methyl-1*H*-imidazo[4,5-*c*]pyridin-4-yl(2,4-dimethoxybenzyl)carbamate




[00126] To a solution of 6-chloro-*N*-(2,4-dimethoxybenzyl)-1-methyl-1*H*-imidazo[4,5-*c*]pyridin-4-amine (example 1A, 0.63 g, 1.893 mmol) in tetrahydrofuran (38 mL) cooled to -78 °C, was added NaHMDS (1.0 M THF, 2.366 mL, 2.366 mmol) slowly. After stirring at -78 °C for 30 min, BOC₂O (0.483 mL, 2.082 mmol) was added. The cold bath was removed and the reaction was allowed to warm to rt ON. By HPLC, ~50% conversion was observed. The solution was re-cooled to -78 °C. NaHMDS (1.0 M THF, 2.366 mL, 2.366 mmol) was added and the mixture was stirred 30 min before the addition of BOC₂O (0.483 mL, 2.082 mmol). The cold bath was removed and the reaction warmed to room temperature 5 h. The solvent was removed under reduced pressure and the residue partitioned between ethyl acetate and water. The organic layer was washed twice with water, dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo*. The crude residue was purified by flash chromatography using an Isco 40g column eluting with 50 - 100% ethyl acetate/hexane to yield *tert*-butyl 6-chloro-1-methyl-1*H*-imidazo[4,5-*c*]pyridin-4-yl(2,4-dimethoxybenzyl) carbamate (0.638 g, 78 % yield) as a foamy white solid.

[00127] MS (ESI) m/z 433.0 (M+H)

[00128] ^1H NMR (400 MHz, CHLOROFORM- d) δ ppm 7.91 (s, 1 H), 7.47 (d, 1 H, $J=8.31$ Hz), 7.19 (s, 1 H), 6.38 (dd, 1 H, $J=8.31$, 2.52 Hz), 6.32 (d, 1 H, $J=2.52$ Hz), 5.13 (s, 2 H), 3.80 (s, 3 H), 3.74 (s, 3 H), 3.59 (s, 3 H), 1.40 (s, 9 H)

5

1C Preparation of *tert*-butyl 2,4-dimethoxybenzyl(6-(diphenylmethyleneamino)-1-methyl-1*H*-imidazo[4,5-*c*]pyridin-4-yl)carbamate




10 [00129] Anhydrous dioxane (4 mL) was added to a mixture of *tert*-butyl 6-chloro-1-methyl-1*H*-imidazo[4,5-*c*]pyridin-4-yl(2,4-dimethoxybenzyl)carbamate (example **1B**, 0.638 g, 1.474 mmol), benzophenone imine (0.297 mL, 1.769 mmol), $\text{Pd}_2(\text{dba})_3$ (0.270 g, 0.295 mmol), 9,9-dimethyl-4,5-bis(diphenylphosphino)xanthene (0.256 g, 0.442 mmol), and cesium carbonate (0.672 g, 2.063 mmol). The reaction mixture was
15 heated to 90 °C overnight. After cooling to rt, solvent was removed *in vacuo*. The residue was partitioned between ethyl acetate and water. The organic layer was washed twice with water, dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo*. The crude residue was purified by flash chromatography using an Isco 40g column eluting with 50-100% ethyl acetate/hexane to provide *tert*-
20 butyl 2,4-dimethoxybenzyl(6-(diphenylmethyleneamino)-1-methyl-1*H*-imidazo[4,5-*c*]pyridin-4-yl)carbamate (0.757 g, 89 % yield) as a light yellow solid.

[00130] MS (ESI) m/z 414.0 (M-CPh $_2$ +2H)

[00131] ^1H NMR (400 MHz, DMSO- d_6) δ ppm 8.16 (s, 1 H), 7.67 (dd, 2 H, $J=8.56$, 1.51 Hz), 7.44 - 7.60 (m, 3 H), 7.25 (d, 1 H, $J=8.31$ Hz), 6.96 - 7.11 (m, 4 H),
25 6.87 (s, 1 H), 6.46 (d, 1 H, $J=2.52$ Hz), 6.33 (dd, 1 H, $J=8.56$, 2.27 Hz), 4.52 (s, 2 H), 3.70 (s, 3 H), 3.69 (s, 3 H), 3.67 (s, 3 H), 1.25 (s, 9 H)

1D Preparation of *tert*-butyl 6-amino-1-methyl-1*H*-imidazo[4,5-*c*]pyridin-4-yl(2,4-dimethoxybenzyl)carbamate




[00132] In a 1000 mL round bottom flask, *tert*-butyl 2,4-dimethoxybenzyl(6-(diphenylmethyleneamino)-1-methyl-1*H*-imidazo[4,5-*c*]pyridin-4-yl)carbamate (example **1C**, 9.2 g, 15.93 mmol) was stirred in THF (50 mL). 1 N HCl (33 mL) was added. After 2 min, the reaction was quenched with 1 N NaOH (65 mL) and ethyl acetate (100 mL). The organic layer was separated, dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The residue was dried under vacuum for 1 h and triturated with ether (4x). Isolated *tert*-butyl 6-amino-1-methyl-1*H*-imidazo[4,5-*c*]pyridin-4-yl(2,4-dimethoxybenzyl) carbamate (6.587 g, 15.93 mmol, 100 % yield) as a cream solid. Material carried forward with no further purification.

[00133] MS (ESI) *m/z* 413.0 (M+H)

[00134] ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.93 (s, 1 H), 7.44 (d, 1 H, *J*=8.31 Hz), 6.44 (d, 1 H, *J*=2.27 Hz), 6.35 (dd, 1 H, *J*=8.56, 2.27 Hz), 6.25 (s, 1 H), 5.58 (s, 2 H), 4.87 (s, 2 H), 3.67 (s, 6 H), 3.63 (s, 3 H), 1.30 (s, 9 H)

1E Preparation of *tert*-butyl 6-amino-7-iodo-1-methyl-1*H*-imidazo[4,5-*c*]pyridin-4-yl(2,4-dimethoxybenzyl)carbamate




[00135] In a round bottom flask, *tert*-butyl 6-amino-1-methyl-1*H*-imidazo[4,5-*c*]pyridin-4-yl(2,4-dimethoxybenzyl)carbamate (example **1D**, 7.887 g, 19.08 mmol) was dissolved in MeCN (200 mL) and cooled to 0 °C. *N*-Iodosuccinimide (4.51 g, 20.03 mmol) was dissolved in the remaining MeCN (50 mL) and added dropwise to the reaction mixture over 40 min. The reaction mixture was stirred at 0 °C for an

additional 10 min and was quenched with 2 M sodium hydrogensulfite (125 mL). Stirring and temperature were maintained for 50 min. The mixture was transferred to a separatory funnel. The aqueous layer was extracted with CH₂Cl₂ (3 x 100 mL). The combined organic layers were washed with water and brine, dried over Na₂SO₄, filtered and concentrated *in vacuo*. The crude residue was purified by flash chromatography using an Isco 120g column eluting with 20-100% ethyl acetate/CH₂Cl₂ to provide *tert*-butyl 6-amino-7-iodo-1-methyl-1H-imidazo[4,5-c]pyridin-4-yl(2,4-dimethoxybenzyl)carbamate (8.436 g, 82 % yield) as a pale yellow solid.

10 **[00136]** MS (ESI) *m/z* 539.0 (M)

[00137] ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.00 (s, 1 H), 7.40 (d, 1 H, *J*=8.06 Hz), 6.43 (d, 1 H, *J*=2.52 Hz), 6.31 - 6.39 (m, 1 H), 5.73 (s, 2 H), 4.86 (s, 2 H), 3.95 (s, 3 H), 3.67 (s, 3 H), 3.64 (s, 3 H), 1.30 (s, 9 H)

15 **1F** Preparation of ethyl 4-(*tert*-butoxycarbonyl(2,4-dimethoxybenzyl)amino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxylate



[00138] To a flask charged with *tert*-butyl 6-amino-7-iodo-1-methyl-1H-imidazo[4,5-c]pyridin-4-yl(2,4-dimethoxybenzyl)carbamate (example **1E**, 3.49 g, 6.47 mmol) in DMA (43.1 mL) was added Pd₂(dba)₃ (0.474 g, 0.518 mmol), ethyl pyruvate (7.22 mL, 64.7 mmol), and *N*-methyldicyclohexylamine (2.77 mL, 12.94 mmol). The reaction mixture was sparged with argon for 10 min and heated at 60 °C 6 h. The reaction was complete by LCMS. After cooling to room temperature, the reaction mixture was diluted with ethyl acetate and saturated aqueous NaHCO₃. The resultant emulsion was vacuum filtered through a pad of Celite and transferred to a separatory funnel. The aqueous layer was extracted with ethyl acetate. The combined organic layers were washed with 10% aqueous LiCl solution (2 x). The organic phase was dried over MgSO₄, filtered and concentrated *in vacuo*. The brown


residue was purified by flash chromatography using an Isco 120g column eluting with 40-80% ethyl acetate/hexane to yield ethyl 4-(*tert*-butoxycarbonyl(2,4-dimethoxybenzyl)amino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxylate (2.830 g, 86 % yield)

5 [00139] MS (ESI) m/z 510.0 (M+H)

[00140] ^1H NMR (400 MHz, CHLOROFORM- d) δ ppm 7.83 (s, 1 H), 7.57 (d, 1 H, $J=8.31$ Hz), 7.39 (d, 1 H, $J=2.27$ Hz), 6.40 (dd, 1 H, $J=8.44$, 2.39 Hz), 6.32 (d, 1 H, $J=2.52$ Hz), 5.16 (s, 2 H), 4.43 (q, 2 H, $J=7.05$ Hz), 4.09 (s, 3 H), 3.74 (s, 3 H), 3.56 (s, 3 H), 1.43 (t, 3 H, $J=7.18$ Hz), 1.40 (s, 9 H)

10

1G Preparation of ethyl 4-(*tert*-butoxycarbonyl(2,4-dimethoxybenzyl)amino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxylate




[00141] To a solution of ethyl 4-(*tert*-butoxycarbonyl(2,4-dimethoxybenzyl)amino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxylate (example 1F, 5.046 g, 9.90 mmol) in DMF (49.5 mL) was added ethyl iodide (1.601 mL, 19.81 mmol) and cesium carbonate (6.45 g, 19.81 mmol). The reaction mixture was heated at 60 °C 60 min and cooled to room temperature. The reaction mixture was diluted with ethyl acetate and vacuum filtered to remove cesium carbonate. Saturated aqueous sodium bicarbonate was added and the aqueous layer was extracted with ethyl acetate (2 x). The combined organic layers were washed with 10% lithium chloride (3 x), dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo*. Crude product was carried forward.

[00142] MS (ESI) m/z 538.1 (M+H)


25 [00143] ^1H NMR (400 MHz, CHLOROFORM- d) δ ppm 7.79 (s, 1 H), 7.62 (d, 1 H $J=8.28$ Hz), 7.43 (s, 1 H), 6.39 (dd, 1 H, $J=8.41$, 2.38 Hz), 6.32 (d, 1 H, $J=2.26$ Hz), 5.20 (s, 2 H), 4.79 (q, 2 H, $J=7.03$ Hz), 4.41 (q, 2 H, $J=7.28$ Hz), 4.07 (s, 3 H), 3.74 (s, 3 H), 3.61 (s, 3 H), 1.44 (t, 3 H, $J=7.15$ Hz), 1.41 (s, 9 H), 1.37 (t, 3 H, $J=7.03$ Hz)

1H Preparation of 4-(*tert*-butoxycarbonyl(2,4-dimethoxybenzyl)amino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxylic acid



- 5 **[00144]** To a solution of ethyl 4-(*tert*-butoxycarbonyl(2,4-dimethoxybenzyl)amino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxylate (example **1G**, 5.32 g, 9.9 mmol) in ethanol (49.5 mL) was added 1 N NaOH aqueous solution (49.5 mL). The reaction mixture was heated at 60 °C 3 h and cooled to rt. Ethanol was removed by concentration *in vacuo*. The
- 10 residue was acidified with 1 N HCl and the product was collected by vacuum filtration. 4-(*tert*-Butoxycarbonyl(2,4-dimethoxybenzyl)amino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxylic acid (4.3 g, 8.44 mmol, 85 % yield over two steps) was isolated as a white solid after drying on the vacuum.
- 15 **[00145]** MS (ESI) *m/z* 510.0 (M+H)

1I Preparation of *tert*-butyl 7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-yl(2,4-dimethoxybenzyl)carbamate




- 20 **[00146]** To a mixture of 4-(*tert*-butoxycarbonyl(2,4-dimethoxybenzyl)amino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxylic acid (example **1H**, 0.89 g, 1.747 mmol) in acetonitrile (17.47 mL) was added dicyclopropylamine hydrochloride (0.303 g, 2.271 mmol), *N*-methylmorpholine (0.499 mL, 4.54 mmol), DMAP (0.021 g, 0.175 mmol), and HATU (0.797 g, 2.096
- 25 mmol). The reaction mixture was stirred at rt 5 h and heated to 50 °C for 45 min.

The reaction was cooled to room temperature. Acetonitrile was removed *in vacuo* and the residue was partitioned between saturated aqueous sodium bicarbonate and dichloromethane. The aqueous layer was extracted with dichloromethane. The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The crude solid was purified by flash chromatography using an Isco 80 g column eluting with 1 - 4% MeOH/CH₂Cl₂ to yield *tert*-butyl 7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-yl(2,4-dimethoxybenzyl)carbamate (0.8 g, 78 % yield) as an off-white solid.

10 [00147] MS (ESI) *m/z* 589.0 (M+H)

[00148] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 7.78 (s, 1 H), 7.67 (d, 1 H, *J*=8.28 Hz), 6.85 (s, 1 H), 6.40 (dd, 1 H, *J*=8.41, 2.38 Hz), 6.33 (d, 1 H, *J*=2.51 Hz), 5.20 (s, 2 H), 4.58 (q, 2 H, *J*=7.03 Hz), 4.02 (s, 3 H), 3.74 (s, 3 H), 3.62 (s, 3 H), 2.77 - 2.85 (m, 2 H), 1.42 (s, 9 H), 1.39 (t, 3 H, *J*=7.40 Hz), 0.80 - 0.89 (m, 4 H), 0.72 - 0.78 (m, 4 H)

1J Preparation of 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide



20 [00149] To a solution of *tert*-butyl 7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-yl(2,4-dimethoxybenzyl)carbamate (example **II**, 2.66 g, 4.51 mmol) in dichloromethane (33.8 mL) was added TFA (11.28 mL). The reaction mixture was stirred at rt 45 min and was concentrated *in vacuo*. The crude residue was partitioned between ethyl acetate and saturated aqueous sodium bicarbonate. The aqueous layer was extracted with ethyl acetate (2 x) and the combined organics were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The crude solid was purified by flash chromatography using an Isco 40 g column eluting with 4 - 10% MeOH/CH₂Cl₂ to yield 4-amino-*N,N*-

dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (1.2 g, 3.55 mmol, 79 % yield) as a pale yellow solid.

[00150] MS (ESI) m/z 339.2 (M+H)

[00151] ^1H NMR (400 MHz, MeOD) δ ppm 7.95 (s, 1 H), 7.15 (s, 1 H), 4.51 (q, 2 H $J=7.13$ Hz), 4.04 (s, 3 H), 2.89 - 2.98 (m, 2 H), 1.35 (t, 3 H), 0.82 - 0.91 (m, 4 H), 0.70 - 0.81 (m, 4 H)


1 Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00152] To a mixture of 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example 1J, 30 mg, 0.089 mmol), 2-bromothiazole (11.98 μL , 0.133 mmol), $\text{Pd}_2(\text{dba})_3$ (12.18 mg, 0.013 mmol), BINAP (24.84 mg, 0.040 mmol) and sodium *tert*-butoxide (13.63 mg, 0.142 mmol) was added PhMe (887 μL). The mixture was sparged with argon gas for 5 min and heated to 85 °C for 5.5 h. The reaction mixture was filtered through a plug of Celite and concentrated. The crude residue was purified by preparative HPLC to provide *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (4 mg, 10.49 % yield) along with some recovered starting material.

[00153] MS (ESI) m/z 423.0 (M+H)


[00154] ^1H NMR (500 MHz, DMSO- d_6) δ ppm 8.14 (s, 1H), 7.47 (d, 1H, $J = 3.57$ Hz), 7.24 (s, 1H), 7.14 (d, 1H, $J = 3.57$ Hz), 4.63 (q, 2H, $J = 6.87$ Hz), 4.05 (s, 3H), 2.88 - 2.99 (m, 2H), 1.39 (t, 3H, $J = 7.01$ Hz), 0.73 - 0.81 (m, 4H), 0.64 - 0.70 (m, 4H)

Example 2




***N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-methylthiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

2A Preparation of 4-(3-benzoylthioureido)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide



- 5 **[00155]** A solution of 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **1J**, 234 mg, 0.69 mmol) and benzoyl isothiocyanate (0.1 ml, 0.83 mmol) in acetone (3 ml) was stirred at room temperature for 3h. The reaction mixture was cooled to 0°C and water was added. The resulting brown solid was collected by filtration and air-dried to give
- 10 4-(3-benzoylthioureido)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (306 mg, 88% yield)
- [00156]** MS (ESI) *rt* = 1.86 min, *m/z* 502 (M+H).
- [00157]** ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.44 (br. s., 1 H) 7.99 - 8.07 (m, 2 H) 7.64 - 7.74 (m, 1 H) 7.58 (t, *J*=7.65 Hz, 2 H) 7.34 (s, 1 H) 4.47 (br. s., 2 H) 4.11 (s,
- 15 3 H) 2.90 - 2.99 (m, 2 H) 2.08 (s, 1 H) 1.28 (br. s., 3 H) 0.64 - 0.81 (m, 8 H).

2B Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide



- 20 **[00158]** 4-(3-Benzoylthioureido)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo [4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **2A**, 306 mg, 0.61 mmol) and 1N NaOH (10 ml, 10.00 mmol) in 10 ml EtOH were heated at 60°C for 1h. The reaction was cooled to room temperature and concentrated. The remaining aqueous solution was extracted (3x) with ethyl acetate. The organic phase
- 25 was dried over Na₂SO₄, filtered, and concentrated. Purification by silica gel

chromatography (dichloromethane/methanol 0-4%) gave *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide as a yellow solid (199 mg, 82% yield).

[00159] MS (ESI) m/z 398 (M+H).

5 [00160] ^1H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 11.02 (br. s., 1 H) 9.27 (s, 2 H) 7.75 (s, 1 H) 6.89 (s, 1 H) 4.52 (q, $J=7.12$ Hz, 2 H) 4.06 (s, 3 H) 2.77 - 2.90 (m, 2 H) 1.45 (t, $J=7.15$ Hz, 3 H) 0.73 - 0.97 (m, 8 H).


2 Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-methylthiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide-

[00161] *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example 2B, 40 mg, 0.10 mmol) and
 15 chloroacetone (62.4 mg, 0.40 mmol) in EtOH (2 ml) were heated at 80°C for 20 min. The reaction was cooled to room temperature and the solvent removed *in vacuo*. Saturated NaHCO₃ solution was added and the aqueous layer extracted (3x) with dichloromethane. The organic layers were combined, dried over Na₂SO₄, filtered, and concentrated. The crude material was purified by silica gel chromatography
 20 (dichloromethane/methanol 0-7%) to provide *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-methylthiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (9 mg, 12 % yield) as a white solid.

[00162] MS (CI) m/z 436.1 (M+H)


[00163] ^1H NMR (400 MHz, DMSO-*d*₆) δ ppm 10.70 (br s, 1H), 8.12 (s, 1H), 7.23 (s, 1H), 6.69 (s, 1H), 4.61 (q, 2H, $J=7.19$ Hz), 4.04 (s, 3H), 2.88 - 2.98 (m, 2H),
 25 2.29 (s, 3H), 1.38 (t, 3H, $J=7.15$ Hz), 0.72 - 0.81 (m, 4H), 0.62 - 0.70 (m, 4H)

Example 3



***N,N*-dicyclopropyl-6-ethyl-4-(4-(fluoromethyl)thiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

3A Preparation of 1-bromo-3-fluoropropan-2-one



[00164] Bromine (0.36 ml, 7.02 mmol) was added dropwise to a solution of 1-fluoropropan-2-one (0.51 ml, 7.02 mmol) in carbon tetrachloride (25 ml). The solution was heated at 45°C for 3h. The reaction was cooled to room temperature and quenched with water. Upon extraction with dichloromethane (3x), the combined organic layers were dried over Na₂SO₄, filtered, and concentrated. Purification by silica gel chromatography (0-20% EtOAc/Hexanes) gave 1-bromo-3-fluoropropan-2-one as a yellow oil (118 mg, 65%).

[00165] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 5.14 (s, 1 H) 5.02 (s, 1 H) 4.10 (d, *J*=2.86 Hz, 2 H).


3 Preparation of *N,N*-dicyclopropyl-6-ethyl-4-(4-(fluoromethyl)thiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00166] Prepared from *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example 2B) and 1-bromo-3-fluoropropan-2-one (example 3A) using the same procedure for 2.

[00167] MS (ESI) *rt* = 1.94 min, *m/z* 454 (M+H).

[00168] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 9.34 (br. s., 1 H) 7.75 (s, 1 H) 6.96 (d, *J*=3.52 Hz, 1 H) 6.90 (s, 1 H) 5.27 - 5.53 (m, 2 H) 4.75 (q, *J*=7.12 Hz, 2 H) 3.98 - 4.11 (m, 3 H) 2.76 - 2.91 (m, 2 H) 1.54 (t, *J*=7.04 Hz, 3 H) 0.70 - 0.94 (m, 8 H).

Example 4



4-(4-*tert*-butylthiazol-2-ylamino)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide


5

[00169] Prepared from *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo [4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **2B**) and 1-chloro-3,3-dimethylbutan-2-one using the same procedure as reported for **2**.

[00170] MS (ESI) *rt* = 1.99 min, *m/z* 478 (*M*+*H*).

10 **[00171]** ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 7.74 (s, 1 H) 6.90 (s, 1 H) 6.48 (s, 1 H) 4.74 (q, *J*=7.03 Hz, 1 H) 4.04 (s, 3 H) 2.77 - 2.90 (m, 2 H) 1.53 (t, *J*=7.03 Hz, 3 H) 1.37 (s, 9 H) 0.70 - 0.96 (m, 8 H).

Example 5




15

***N,N*-dicyclopropyl-6-ethyl-4-(4-(isopropylsulfonylmethyl)thiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

5A Preparation of 1-chloro-1-(isopropylsulfonyl)propan-2-one

20



[00172] Sulfuryl chloride (0.248 mL, 3.05 mmol) was added dropwise to a solution of 1-(isopropylsulfonyl)propan-2-one (418 mg, 2.55 mmol) in 1:1 AcOH/DCM (16

ml) at 0°C. The reaction was stirred at rt for 4h. The solvent was removed *in vacuo*, then diluted with dichloromethane. The organic layer was washed (2x) with saturated NaHCO₃ solution, dried over Na₂SO₄, filtered, and concentrated to leave 800 mg of a colorless oil. The material was used without further purification.

- 5 **[00173]** ¹H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 5.19 (s, 1 H) 3.65 (dt, *J*=13.66, 6.90 Hz, 1 H) 2.56 (s, 3 H) 1.45 (dd, *J*=6.94, 3.05 Hz, 6 H).


5 Preparation of *N,N*-dicyclopropyl-6-ethyl-4-(4-(isopropylsulfonylmethyl)thiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-
 10 d]pyrrolo[2,3-*b*]pyridine-7-carboxamide

- [00174]** Prepared from *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo [4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (example 2B) and 1-chloro-1-(isopropylsulfonyl)propan-2-one (example 5^a) using same procedure as
 15 reported for **2**.

[00175] MS (ESI) *rt* = 1.84 min, *m/z* 542 (*M*+*H*).

- [00176]** ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 7.82 (s, 1 H) 7.01 (s, 1 H) 6.90 (s, 1 H) 4.73 (q, *J*=7.04 Hz, 2 H) 4.35 (s, 2 H) 4.06 (s, 3 H) 3.36 (quin, *J*=6.88 Hz, 1 H) 2.79 - 2.91 (m, 2 H) 1.53 (t, *J*=7.04 Hz, 3 H) 1.43 (d, *J*=6.82 Hz, 6 H) 0.74 -
 20 0.92 (m, 8 H).

Example 6




***N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-propylthiazol-2-ylamino)-1,6-**
 25 **dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

- [00177]** Prepared from *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo [4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (example 2B) and 1-bromopentan-2-one using the same procedure as reported for **2**.

[00178] MS (ESI) $rt = 1.94$ min, m/z 464 (M+H).

[00179] 1H NMR (400 MHz, CHLOROFORM- d) δ ppm 8.43 - 9.78 (m, 1 H) 7.71 (s, 1 H) 6.89 (s, 1 H) 6.48 (s, 1 H) 4.75 (q, $J=7.04$ Hz, 2 H) 4.04 (s, 3 H) 2.76 - 2.92 (m, 2 H) 2.68 (t, $J=7.37$ Hz, 2 H) 1.76 (dq, $J=14.94, 7.42$ Hz, 2 H) 1.53 (t, $J=7.04$ Hz, 3 H) 1.01 (t, $J=7.26$ Hz, 3 H) 0.69 - 0.94 (m, 8 H).

Example 7



10 ***N,N*-dicyclopropyl-6-ethyl-4-(4-ethylthiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**


[00180] Prepared from *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo [4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example 2B) and 1-bromobutan-2-one using the same procedure as reported for 2.

15 [00181] MS (ESI) m/z 450 (M+H).

[00182] 1H NMR (400 MHz, CHLOROFORM- d) δ ppm 7.71 (s, 1 H) 6.89 (s, 1 H) 6.49 (s, 1 H) 4.75 (q, $J=7.03$ Hz, 2 H) 4.04 (s, 3 H) 2.80 - 2.88 (m, 2 H) 2.75 (q, $J=7.45$ Hz, 1 H) 1.54 (t, $J=7.15$ Hz, 2 H) 1.33 (t, $J=7.40$ Hz, 2 H) 0.82 (br. s., 8 H).

20

Example 8




***N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-(pyridin-2-yl)thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

[00183] A solution of 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example 1J, 70 mg, 0.21 mmol) and benzoyl isothiocyanate (36 μ l, 0.27 mmol) in acetone (1 ml) was stirred at room temperature for 1.5h. The solvent was removed *in vacuo*, the residue was taken up in EtOH (1.5 ml), and K₂CO₃ (40 mg, 0.290 mmol) was added. The reaction was heated at 60°C for 3h. Upon cooling to room temperature, 2-bromo-1-(pyridin-2-yl)ethanone, hydrobromide (139.4 mg, 0.50 mmol) was added and the reaction heated at 80°C for 24h. The reaction mixture was cooled to room temperature and concentrated. The residue was purified by silica gel chromatography (0-7% MeOH/dichloromethane). Re-purification by preparative HPLC provided 10.2 mg (10% yield) of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-(pyridin-2-yl)thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide as a brown solid.

[00184] MS (ESI) *rt* = 1.78 min, *m/z* 499 (M+H).

[00185] ¹H NMR (DMSO-*d*₆) δ ppm 10.9 (s, 1 H), 8.2 (bs, 1 H), 7.81 (d, 1 H), 7.75 (s, 1H), 3.92 (m, 4H), 3.77 (q, 2H, *J* = 7.8 Hz), 2.52-2.81 (m, 7H).

Example 9




20 **Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-(trifluoromethyl)thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

[00186] This compound was prepared according to example 2 using 3-bromo-1,1,1-trifluoroacetone to provide *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-(trifluoromethyl) thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide.

[00187] MS (ESI) *m/z* 490.0 (M+H)

[00188] ^1H NMR (500 MHz, DMSO- d_6) δ ppm 11.71 (s, 1H), 8.16 (s, 1H), 7.86 (s, 1H), 7.27 (s, 1H), 4.63 (q, 2H), 4.05 (s, 3H), 2.89 – 2.98 (m, 2H), 1.40 (t, 3H, $J = 6.87$ Hz), 0.74 – 0.80 (m, 4H), 0.64 – 0.71 (m, 4H)

5

Example 10

Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-methylthiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

10 [00189] 4-Amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example 1J, 86.3 mg, 0.255 mmol) was dissolved in acetone and benzoyl isothiocyanate (37.8 μL , 0.281 mmol) was added. The reaction was stirred at room temperature for 2h. The solvent was removed *in vacuo*. The residue was taken up in ethanol (1500 μL) and K_2CO_3 (49.3 mg, 0.357


15 mmol) was added. The reaction mixture was warmed to 60 $^\circ\text{C}$ for 3 h. 2-Bromopropanal (45.4 mg, 0.332 mmol) was added. After 3 h, additional 2-bromopropanal (~100 mg) was added. Stirred at 60 $^\circ\text{C}$ for 16h. Additional 2-bromopropanal was added and the reaction mixture was stirred at 60 $^\circ\text{C}$ 3 h. The ethanol was removed by concentration *in vacuo*. The crude residue was purified by

20 flash chromatography using an Isco 12 g column eluting with 2 - 10% $\text{MeOH}/\text{CH}_2\text{Cl}_2$). Additional purification was accomplished via preparative HPLC. *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-methylthiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (20 mg, 17.47 % yield) was isolated as a white solid.

25 [00190] MS (ESI) m/z 436.0 ($\text{M}+\text{H}$)

[00191] ^1H NMR (500 MHz, DMSO- d_6) δ ppm 10.56 (br s, 1H), 8.12 (s, 1H), 7.23 (s, 1H), 7.12 (s, 1H), 4.62 (q, 2H, $J = 6.78$ Hz), 4.04 (s, 3H), 2.87 – 2.99 (m, 2H), 2.39 (s, 3H), 1.35 – 1.46 (m, 3H), 0.72 – 0.81 (m, 4H), 0.59 – 0.71 (m, 4H)

Example 11



***N,N*-dicyclopropyl-6-ethyl-4-(5-ethylthiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**


5

[00192] Prepared from 2-bromobutanal and 4-Amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example 1J) using the same procedure as that of example 8.

[00193] MS (ESI) *rt* = 1.86 min, *m/z* 450 (*M*+*H*).

10 **[00194]** ¹H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 8.91 - 9.22 (m, 1 H) 7.71 (s, 1 H) 7.14 (s, 1 H) 6.89 (s, 1 H) 4.75 (q, *J*=6.94 Hz, 2 H) 4.05 (s, 3 H) 2.80 - 2.89 (m, 4 H) 1.55 (t, *J*=5.00 Hz, 3 H) 1.37 (t, *J*=7.49 Hz, 3 H) 0.73 - 0.92 (m, 8 H).


Example 12



15


2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-*N,N*-dimethylthiazole-4-carboxamide

20 **12A** Preparation of 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-4-carboxylic acid



- [00195] 4-Amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (**1J**, 300 mg, 0.89 mmol) and benzoyl isothiocyanate (0.16 ml, 1.15 mmol) in acetone (4 ml) were stirred at rt for 1.5h. The solvent was removed *in vacuo* and the residue taken up in EtOH (6 ml). K₂CO₃ (172 mg, 1.241 mmol) was added and the reaction heated at 60°C for 3h. The reaction was cooled to rt and ethyl 3-bromo-2-oxopropanoate (692 mg, 3.55 mmol) was added. The reaction was heated at 70°C overnight. Upon completion of the cyclization step, 1N NaOH (6.3 ml, 6.30 mmol) was added to the reaction mixture and heated at 60°C for 4h. The reaction was cooled to room temperature and the solvent removed *in vacuo*. The remaining aqueous solution was acidified to pH 1 and the precipitate removed by vacuum filtration, washing with water, EtOAc, and then dichloromethane to afford 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-4-ylamino)thiazole-4-carboxylic acid (325 mg, 79% yield) as brown solid.
- [00196] MS (ESI) *rt* = 1.85 min, *m/z* 466 (M+H).
- [00197] ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.16 (s, 1 H) 7.94 (s, 1 H) 7.26 (s, 1 H) 4.63 (q, *J*=6.94 Hz, 2 H) 4.05 (s, 3 H) 3.36 (br. S., 1 H) 2.89 – 2.99 (m, 2 H) 1.40 (t, *J*=7.03 Hz, 3 H) 0.61 – 0.83 (m, 8 H).

20 **12** Preparation of 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-*N,N*-dimethylthiazole-4-carboxamide



- [00198] To 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d] pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-4-carboxylic acid (example **12A**, 44 mg, 0.095 mmol) in DMF (2 ml) was added HATU (108 mg, 0.284 mmol), DIPEA (0.066 mL, 0.378 mmol), and dimethylamine (0.013 mL, 0.189 mmol). The reaction was stirred at rt for 3h, then the solvent was removed *in vacuo*. The residue was purified by preparative HPLC to give 2-(7-


(dicycloppropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo [2,3-b]pyridine-4-ylamino)-*N,N*-dimethylthiazole-4-carboxamide (10 mg , 21% yield) as a brown solid.

[00199] MS (ESI) m/z 493 (M+H)

5 [00200] ^1H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 9.45 (br. S., 1 H) 7.81 (s, 1 H) 7.43 (s, 1 H) 6.90 (s, 1 H) 4.74 (q, $J=7.03$ Hz, 2 H) 4.05 (s, 3 H) 3.31 (br. S., 3 H) 3.13 (br. S., 3 H) 2.75 – 2.90 (m, 2 H) 1.54 (t, $J=7.15$ Hz, 3 H) 0.65 – 0.96 (m, 8 H).

10 Examples **13-17** were prepared using the procedure as described for example **12**.

Example 13




15

N-cyclopropyl-2-(7-(dicycloppropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-4-carboxamide

[00201] MS (ESI) $rt = 1.99$ min, m/z 505 (M+H).

20 [00202] ^1H NMR (400 MHz, MeOD) δ ppm 8.07 (s, 1 H) 7.74 (s, 1 H) 7.26 (s, 1 H) 4.73 (q, $J=7.03$ Hz, 1 H) 4.13 (s, 3 H) 2.94 – 3.04 (m, 2 H) 2.85 – 2.93 (m, 1 H) 1.51 (t, $J=7.03$ Hz, 2 H) 0.67 – 0.95 (m, 12 H).

Example 14




25

***N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-(morpholine-4-carbonyl)thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

[00203] MS (ESI) *rt* = 1.87 min, *m/z* 535 (M+H).

5 [00204] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 9.21 (br. s., 1 H) 7.76 (s, 1 H) 7.56 (s, 1 H) 6.81 - 6.94 (m, 1 H) 4.74 (q, *J*=7.11 Hz, 2 H) 4.07 (s, 3 H) 3.79 (br. s., 4 H) 3.50 (d, *J*=4.52 Hz, 2 H) 2.74 - 2.90 (m, 2 H) 1.60 (s, 2 H) 1.54 (t, *J*=7.03 Hz, 3 H) 0.70 - 0.94 (m, 8 H).

Example 15



***N,N*-dicyclopropyl-4-((4-((1,1-dioxidotetrahydro-3-thiophenyl)carbamoyl)-1,3-thiazol-2-yl)amino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

15


[00205] MS (ESI) *rt* = 1.88 min, *m/z* 583 (M+H).

[00206] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 9.31 (br. s., 1 H) 7.77 (s, 1 H) 7.69 - 7.75 (m, 1 H) 6.91 (s, 1 H) 4.85 - 4.99 (m, 1 H) 4.64 - 4.81 (m, 2 H) 4.07 (s, 3 H) 3.45 - 3.61 (m, 2 H) 3.35 (br. s., 1 H) 3.19 (dt, *J*=13.36, 8.12 Hz, 2 H) 2.75 - 2.91 (m, 2 H) 2.57 - 2.72 (m, 1 H) 2.42 (ddd, *J*=14.49, 7.53, 7.34 Hz, 1 H) 1.54 (t, *J*=6.90 Hz, 3 H) 0.69 - 0.94 (m, 8 H)

20

25

Example 16



***N,N*-dicyclopropyl-4-(4-(3,3-difluoropyrrolidine-1-carbonyl)thiazol-2-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**


5

[00207] MS (ESI) *rt* = 2.05 min, *m/z* 555 (*M*+*H*).

[00208] ¹H NMR (400 MHz, MeOD) δ ppm 8.00 (s, 1 H) 7.73 (s, 1 H) 7.70 (s, 1 H) 7.19 (s, 1 H) 4.68 (q, *J*=7.03 Hz, 2 H) 4.51 (t, *J*=12.80 Hz, 1 H) 4.34 (t, *J*=7.40 Hz, 1 H) 4.06 (s, 3 H) 3.96 (t, *J*=13.18 Hz, 1 H) 3.86 (t, *J*=7.65 Hz, 1 H) 2.87 - 3.02 (m, 2 H) 2.34 - 2.59 (m, 2 H) 1.45 (t, *J*=6.90 Hz, 3 H) 0.72 - 0.95 (m, 8 H).

10

Example 17



(4-(azetidine-1-carbonyl)thiazol-2-ylamino)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

15


[00209] MS (ESI) *rt* = 1.95 min, *m/z* 505 (*M*+*H*).

[00210] ¹H NMR (400 MHz, MeOD) δ ppm 8.03 (s, 1 H) 7.70 (s, 1 H) 7.21 (s, 1 H) 4.78 (t, *J*=7.53 Hz, 2 H) 4.69 (q, *J*=7.03 Hz, 2 H) 4.19 (t, *J*=7.78 Hz, 2 H) 4.08 (s, 3 H) 2.95 (ddd, *J*=7.22, 3.45, 3.14 Hz, 2 H) 2.27 - 2.48 (m, 2 H) 1.47 (t, *J*=7.03 Hz, 3 H) 0.69 - 0.93 (m, 8 H).

20

25

Example 18



Ethyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-4-carboxylate


5 **[00211]** This compound was prepared according to example 2 using ethyl bromopyruvate and *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **2B**) to yield ethyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-4-carboxylate as yellow solid.

10 **[00212]** MS (ESI) *m/z* 494.1 (M+H)

[00213] ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 11.42 (br s, 1H), 8.14 (s, 1H), 8.00 (s, 1H), 7.25 (s, 1H), 4.62 (q, 2H), 4.30 (q, 2H, *J* = 7.15 Hz), 4.05 (s, 3H), 2.89 – 2.97 (m, 2H), 1.39 (t, 3H, *J* = 7.01 Hz), 1.32 (t, 3H, *J* = 7.15 Hz), 0.74 – 0.80 (m, 4H), 0.64 – 0.70 (m, 4H)

15

Example 19



Ethyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-5-carboxylate

20

[00214] This compound was prepared according to example 2 using ethyl 2-chloro-2-formylacetate and *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo [4,5-d]pyrrolo [2,3-b]pyridine-7-carboxamide (example **2B**) to provide ethyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-


25

dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-5-carboxylate (18.5 mg, 17 % yield) as a pale yellow solid.

[00215] MS (ESI) m/z 494.0 (M+H)


[00216] ^1H NMR (400 MHz, DMSO- d_6) δ ppm 11.95 (s, 1H), 8.20 (s, 1H), 8.14 (s, 1H), 7.28 (s, 1H), 4.63 (q, 2H, $J = 7.19$ Hz), 4.28 (q, 2H, $J = 7.19$ Hz), 4.06 (s, 3H), 2.88 – 3.00 (m, 2H), 1.44 (t, 3H, $J = 7.03$ Hz), 1.32 (t, 3H, $J = 7.15$ Hz), 0.73 – 0.81 (m, 4H), 0.63 – 0.70 (m, 4H)

Example 20



***N*-cyclopropyl-2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-5-carboxamide**

20A Preparation of 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo [4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-5-carboxylic acid



[00217] To a solution of ethyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-5-carboxylate (**example 19**, 14.6 mg, 0.030 mmol) in EtOH (296 μL) was added 1 N NaOH aqueous solution (296 μL). The reaction mixture was stirred at rt 3 h and stirred at 50 $^{\circ}\text{C}$ 18 h. After cooling to room temperature, ethanol was removed by concentration *in vacuo*. The residue was acidified with 1 N HCl and the solid product was collected by filtration. 2-(7-(Dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-5-carboxylic acid was isolated as a pale yellow solid which was used crude in the next reaction.

[00218] MS (ESI) m/z 465.9 (M+H)


20 Preparation of *N*-cyclopropyl-2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-5-carboxamide

[00219] To a mixture of 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-5-carboxylic acid (example **20A**, 13.97 mg, 0.030 mmol) in DMF (300 μ L) was added cyclopropylamine (4.16 μ L, 0.060 mmol), HATU (14.83 mg, 0.039 mmol), and Hünig'sBase (15.72 μ L, 0.090 mmol). The reaction mixture was stirred at rt 1 h and concentrated *in vacuo*. The residue was taken up in MeOH and purified by preparative HPLC. *N*-Cyclopropyl-2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-5-carboxamide (3 mg, 19.82 % yield) was isolated as a white solid.

[00220] MS (ESI) m/z 505.0 (M+H)

[00221] ^1H NMR (400 MHz, DMSO- d_6) δ ppm 8.36 (br s, 1H), 8.15 (s, 1H), 8.02 (s, 1H), 7.25 (s, 1H), 4.63 (q, 2H, $J = 7.11$ Hz), 4.05 (s, 3H), 2.89 – 3.00 (m, 2H), 2.76 (tq, 1H, $J = 7.25, 3.70$ Hz), 1.41 (t, 3H, $J = 7.03$ Hz), 0.74 – 0.81 (m, 4H), 0.64 – 0.72 (m, 6H), 0.51 – 0.59 (m, 2H)

Example 21



N,N-dicyclopropyl-6-ethyl-1-methyl-4-(5-(morpholine-4-carbonyl)thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00222] This compound was prepared according to Example **20** using morpholine and 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-5-carboxylic acid (example **20A**) to


provide *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-(morpholine-4-carbonyl)thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide as a white solid.

[00223] MS (ESI) *m/z* 535.1 (M+H)

5 **[00224]** ¹H NMR (400 MHz, MeOD) δ ppm 8.08 (s, 1H), 7.79 – 7.93 (m, 1H), 7.26 (s, 1H), 4.74 (q, 2H, *J* = 7.28 Hz), 4.12 (s, 3H), 3.81 – 3.88 (m, 2H), 3.72 – 3.80 (m, 2H), 2.91 – 3.02 (m, 2H), 1.52 (t, 3H, *J* = 7.15 Hz), 0.83 – 0.93 (m, 4H), 0.72 – 0.84 (m, 4H)

10

Example 22



N,N-dicyclopropyl-6-ethyl-1-methyl-4-(5-(pyrrolidine-1-carbonyl)thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

15

[00225] This compound was prepared according to Example 20 using pyrrolidine and 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)thiazole-5-carboxylic acid (example 20A) to provide *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-(pyrrolidine-1-carbonyl)thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

20


[00226] MS (ESI) *m/z* 519.1 (M+H).

[00227] ¹H NMR (400 MHz, METHANOL-*d*₄) δ 8.58 (s, 1H), 7.97 (s, 1H), 7.28 (s, 1H), 4.71 (q, *J* = 6.7 Hz, 2H), 4.20 (s, 3H), 3.87 (t, *J* = 6.5 Hz, 2H), 3.67 (t, *J* = 6.7 Hz, 2H), 2.97 – 3.05 (m, 2H), 2.10 – 2.19 (m, 2H), 1.98 – 2.07 (m, 2H), 1.51 (t, *J* = 7.2

25

Hz, 3H), 0.88 – 0.95 (m, 4H), 0.80 – 0.86 (m, 4H)

Example 23



2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-N-isopropyl-N-methylthiazole-5-carboxamide

5

[00228] This compound was prepared according to Example **20** using *N*-methylpropan-2-amine and 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo [2,3-b]pyridin-4-ylamino)thiazole-5-carboxylic acid (example **20A**) to provide 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-


10 dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-*N*-isopropyl-*N*-methylthiazole-5-carboxamide

[00229] MS (ESI) *m/z* 521.1 (M+H).

[00230] ¹H NMR (400 MHz, METHANOL-*d*₄) δ : 8.68 (s, 1H), 7.87 (s., 1H), 7.32 (s, 1H), 4.74 (q, *J* = 7.0 Hz, 2H), 4.24 (s, 3H), 3.18 (br. s., 3H), 2.99 - 3.06 (m, 2H),

15 1.53 (t, *J* = 7.0 Hz, 3H), 1.35 (d, *J* = 6.8 Hz, 6H), 0.89 - 0.96 (m, 4H), 0.81 - 0.88 (m, 4H)

Example 24



20 **2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-N,N-dimethylthiazole-5-carboxamide**

[00231] This compound was prepared according to Example **20** using *N,N*-dimethylamine and 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo [2,3-b]pyridin-4-ylamino)thiazole-5-carboxylic acid


25 (example **20A**) to give 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-

dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-*N,N*-dimethylthiazole-5-carboxamide.

[00232] MS (ESI) m/z 493.1 (M+H).

[00233] ^1H NMR (400 MHz, METHANOL- d_4) δ : 8.56 (s, 1H), 7.91 (s, 1H), 7.31 (s, 1H), 4.72 (q, J = 7.0 Hz, 1H), 4.22 (s, 3H), 3.15 – 3.45 (br. S, 6 H), 2.97 - 3.05 (m, 2H), 2.91 (s, 1H), 1.54 (t, J = 7.0 Hz, 3H), 0.88 - 0.95 (m, 4H), 0.80 - 0.86 (m, 4H)

Example 25



10

2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-*N*-(1,1-Dioxo-tetrahydrothiophen-3-yl)-*N*-methylthiazole-5-carboxamide


15 **[00234]** This compound was prepared according to example 20

[00235] MS (ESI) m/z 597.0 (M+H).

[00236] ^1H NMR (400 MHz, METHANOL- d_4) δ : 8.74 (s, 1H), 7.99 (s, 1H), 7.37 (s, 1H), 5.34 - 5.45 (m, 1H), 4.78 (q, J = 6.9 Hz, 2H), 4.28 (s, 3H), 4.05 (s, 3H), 3.49 - 3.59 (m, 1H), 3.22 – 3.33 (m, 1H), 3.00 - 3.10 (m, 2H), 2.94 (s, 1H), 2.47 - 2.66 (m, 2H), 1.57 (t, J = 6.9 Hz, 3H), 1.34 – 1.45 (M, 1H), 0.91 - 0.99 (m, J = 6.5 Hz, 4H), 0.83 - 0.90 (m, 4H)

25

Example 26




***N,N*-dicyclopropyl-4-(4,5-dimethylthiazol-2-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

- 5 **[00237]** This compound was prepared according to example 2 using 3-bromo-2-butanone (0.034 mL, 0.323 mmol) to provide *N,N*-dicyclopropyl-4-(4,5-dimethylthiazol-2-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (59.17 mg, 44.3 % yield) as a pale yellow solid.

[00238] MS (ESI) *m/z* 450.1 (M+H)

- 10 **[00239]** ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.15 (s, 1H), 7.23 (s, 1H), 4.61 (q, 2H, *J* = 7.03 Hz), 4.04 (s, 3H), 2.88 – 2.98 (m, 2H), 2.29 (s, 3H), 2.19 (s, 3H), 1.38 (t, 3H, *J* = 7.03 Hz), 0.73 – 0.80 (m, 4H), 0.62 – 0.71 (m, 4H)

Example 27



15


***N,N*-dicyclopropyl-4-(5,6-dihydro-4H-cyclopenta[d]thiazol-2-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

[00240] Prepared using similar procedure as that of example 2

[00241] MS (ESI) *rt* = 1.95 min, *m/z* 462 (M+H).


- 20 **[00242]** ¹H NMR (500 MHz, MeOD) δ ppm 8.12 (s, 1 H) 7.22 (s, 1 H) 4.68 (q, *J*=7.12 Hz, 2 H) 4.10 (s, 3 H) 2.86 - 3.02 (m, 4 H) 2.80 (t, *J*=7.07 Hz, 2 H) 2.43 - 2.61 (m, 2 H) 1.46 (t, *J*=7.07 Hz, 3 H) 0.70 - 0.90 (m, 8 H).

Example 28



***N,N*-dicyclopropyl-6-ethyl-4-(4-ethyl-5-methylthiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

5 **28A** Preparation of 2-bromopentan-3-one



[00243] Prepared according to the procedure described in *Chem. Comm.* **2004**, 470-471.


10 **28** Preparation of *N,N*-dicyclopropyl-6-ethyl-4-(4-ethyl-5-methylthiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00244] Prepared from example **26A** using similar procedure as that of example **2**

[00245] MS (ESI) *m/z* 464 (M+H).

[00246] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 8.81 – 9.19 (m, 1 H) 7.69 (s, 1 H) 6.89 (s, 1 H) 4.75 (q, *J*=7.12 Hz, 2 H) 4.03 (s, 3 H) 2.78 – 2.93 (m, 2 H) 2.63 (q, *J*=7.48 Hz, 2 H) 2.37 (s, 3 H) 1.53 (t, *J*=7.04 Hz, 3 H) 1.26 (t, *J*=7.48 Hz, 3 H) 0.73 – 0.95 (m, 8 H).

Example 29



20

Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-methyl-4-(trifluoromethyl)thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide


[00247] This compound was prepared according to example 2 using 3-bromo-1,1,1-trifluorobutan-2-one to provide *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-methyl-4-(trifluoromethyl)thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide as a white solid.

5 [00248] MS (ESI) m/z 504.0 (M+H)

[00249] ^1H NMR (500 MHz, DMSO- d_6) δ ppm 8.14 (s, 1H), 7.25 (s, 1H), 4.62 (q, 2H, $J = 6.97$ Hz), 4.04 (s, 3H), 2.89 – 2.98 (m, 2H), 1.40 (t, 3H, $J = 6.87$ Hz), 0.73 – 0.80 (m, 4H), 0.65 – 0.71 (m, 4H)

10


Example 30



Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-methyl-5-(methylsulfonyl)thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

15

30A Preparation of 1-bromo-1-(methylsulfonyl)propan-2-one



[00250] To a solution of methanesulphonylacetone (0.5 g, 3.67 mmol) in AcOH (7.34 mL) and dichloromethane (11.02 mL) in a round bottom flask was added
 20 bromine (0.189 mL, 3.67 mmol) dropwise. The reaction mixture was stirred at rt 90 min. Solvent was removed *in vacuo*. The residue was taken up in ether and washed with saturated aqueous sodium bicarbonate (2 x). The organics were concentrated *in vacuo*. 1-Bromo-1-(methylsulfonyl)propan-2-one (0.332 g, 42% yield) was isolated as an oil.

25 [00251] MS (CI) m/z 215.0 (M+H)


30 Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-methyl-5-(methylsulfonyl)thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00252] This compound was prepared according to example 1 using 1-bromo-1-(methylsulfonyl)propan-2-one (**28A**, 32.5 mg, 0.151 mmol) to provide *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-methyl-5-(methylsulfonyl)thiazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (12.5 mg, 0.023 mmol, 30.6 % yield) as an off-white solid.

[00253] MS (ESI) *m/z* 514.1 (M+H)

[00254] ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 12.00 (br s, 1H), 8.18 (s, 1H), 7.27 (s, 1H), 4.59 (q, 2H, *J* = 6.78 Hz), 4.06 (s, 3H), 3.29 (s, 3H), 2.88 – 2.98 (m, 2H), 2.53 (s, 3H), 1.41 (t, 3H, *J* = 6.90 Hz), 0.72 – 0.81 (m, 4H), 0.62 – 0.71 (m, 4H)

Example 31




Preparation of 4-(5-chloro-4-methylthiazol-2-ylamino)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00255] This compound was prepared according to example 2 using 1,1-dichloropropan-2-one (14.26 μL, 0.147 mmol) to provide 4-(5-chloro-4-methylthiazol-2-ylamino)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (4.35 mg, 18.9% yield) as a pale green solid.

[00256] MS (ESI) *m/z* 470.1 (M+H)

[00257] ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 11.34 (s, 1H), 8.14 (s, 1H), 7.25 (s, 1H), 4.58 (q, 2H, *J* = 7.28 Hz), 4.04 (s, 3H), 2.88 – 2.98 (m, 2H), 2.25 (s, 3H), 1.40 (t, 3H, *J* = 6.90 Hz), 0.73 – 0.81 (m, 4H), 0.63 – 0.70 (m, 4H)

Example 32




Preparation of *N,N*-dicyclopropyl-4-(5,5-dimethyl-7-oxo-4,5,6,7-tetrahydrobenzo[d]thiazol-2-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00258] This compound was prepared according to example 2 using 2-chloro-5,5-dimethylcyclohexane-1,3-dione (34.9 mg, 0.200 mmol) to provide *N,N*-dicyclopropyl-4-(5,5-dimethyl-7-oxo-4,5,6,7-tetrahydrobenzo[d]thiazol-2-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (11.24 mg, 21.7% yield) as a pale yellow solid.

[00259] MS (ESI) m/z 518.2 (M+H)


[00260] ^1H NMR (400 MHz, DMSO- d_6) δ ppm 11.93 (s, 1H), 8.19 (s, 1H), 7.28 (s, 1H), 4.62 (q, 2H), 4.06 (s, 3H), 2.89 – 2.98 (m, 2H), 2.80 (s, 2H), 2.40 (s, 2H), 1.41 (t, 3H, $J = 7.03$ Hz), 1.09 (s, 6H), 0.73 – 0.80 (m, 4H), 0.63 – 0.71 (m, 4H)

Example 33




Preparation of 4-(5,5-dioxo-4,5,6,7-tetrahydro-5 λ^6 -thiopyrano[4,3-d]thiazol-2-ylamino)-6-ethyl-1-methyl-1,6-dihydro-1,3,5,6-tetraaza-as-indacene-7-carboxylic acid dicyclopropylamide

33A Preparation of 3-bromodihydro-2*H*-thiopyran-4(3*H*)-one



[00261] To a mixture of tetrahydrothiopyran-4-one (0.5 g, 4.30 mmol) in dichloromethane (51.2 mL) and methanol (20.49 mL) was added tetra-n-butylammonium tribromide (2.283 g, 4.73 mmol). The reaction mixture was stirred at
 5 rt 2.5 h. Solvent was removed *in vacuo*. The crude solid was triturated with ether (3 x). The combined ether layers were dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo*. The product was isolated as a pale orange oil. Material was used immediately in the next reaction.

10 **33B** Preparation of 3-bromo-1,1-dioxo-tetrahydro-1λ⁶-thiopyran-4-one



[00262] To a cooled (0 °C) of 3-bromodihydro-2*H*-thiopyran-4(3*H*)-one (example 33A, 0.791 g, 4.05 mmol) in dichloromethane (20.27 mL) was added mCPBA (1.749 g, 10.14 mmol) slowly. The reaction mixture was allowed to warm to room
 15 temperature. The reaction mixture was quenched with saturated aqueous sodium bicarbonate. The aqueous layer was extracted with dichloromethane (2 x). The combined organics were dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo* to afford 3-bromo-1,1-dioxo-tetrahydro-1λ⁶-thiopyran-4-one as a white solid which was used immediately in the next reaction.

20

33 Preparation of 4-(5,5-dioxo-4,5,6,7-tetrahydro-5λ⁶-thiopyrano[4,3-d]thiazol-2-ylamino)-6-ethyl-1-methyl-1,6-dihydro-1,3,5,6-tetraaza-as-indacene-7-carboxylic acid dicyclopropylamide


[00263] This compound was prepared according to example 1 using 3-bromo-1,1-dioxo-tetrahydro-1λ⁶-thiopyran-4-one (example 33B, 32.5 mg, 0.151 mmol) to
 25 provide 4-(5,5-dioxo-4,5,6,7-tetrahydro-5λ⁶-thiopyrano[4,3-d]thiazol-2-ylamino)-6-

ethyl-1-methyl-1,6-dihydro-1,3,5,6-tetraaza-as-indacene-7-carboxylic acid dicyclopropylamide (9 mg, 17.56 % yield) as a pale tan solid.

[00264] MS (ESI) m/z 526.2 (M+H)

[00265] ^1H NMR (400 MHz, DMSO- d_6) δ ppm 8.16 (s, 1H), 7.25 (s, 1H), 4.61 (q, 2H, $J = 6.78$ Hz), 4.54 (s, 2H), 4.04 (s, 3H), 3.49 (t, 2H, $J = 6.02$ Hz), 3.16 (t, 2H, $J = 8.16$ Hz), 2.89 – 2.97 (m, 2H), 1.38 (t, 3H, $J = 7.03$ Hz), 0.73 – 0.79 (m, 4H), 0.61 – 0.70 (m, 4H)

Example 34



10

Preparation of 4-(5-acetyl-4-methylthiazol-2-ylamino)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide


[00266] This compound was prepared according to example 2 using 3-chloro-2,4-pentanedione (36.0 μL , 0.302 mmol) to provide 4-(5-acetyl-4-methylthiazol-2-ylamino)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (76 mg, 100 % yield) as a pale yellow solid.

[00267] MS (ESI) m/z 478.2 (M+H)

[00268] ^1H NMR (400 MHz, DMSO- d_6) δ ppm 8.31 (s, 1H), 7.28 (s, 1H), 4.62 (q, 2H, $J = 6.94$ Hz), 4.08 (s, 3H), 2.89 – 2.99 (m, 2H), 2.61 (s, 3H), 1.44 (t, 3H, $J = 7.03$ Hz), 0.72 – 0.81 (m, 4H), 0.64 – 0.72 (m, 4H)


20

Example 35



Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-oxo-4,5-dihydrothiazolo[5,4-c]pyridin-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

5 **35A** Preparation of 3-bromopiperidine-2,4-dione



To a mixture of piperidine-2,4-dione (0.072 g, 0.637 mmol) in acetic acid (1.591 mL) was added bromine (0.033 mL, 0.637 mmol) dropwise. The reaction mixture was stirred at rt 70 min. Dichloromethane was added and the cloudy mixture was concentrated *in vacuo*. The residual solution was treated with ethyl ether. A white solid crashed out and the AcOH/ether was decanted. The white solid, presumed to be product was carried forward. The white solid started liquifying and turning red rapidly upon exposure to light and standing.

15 **35** Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-oxo-4,5-dihydrothiazolo[5,4-c]pyridin-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide


[00269] This compound was prepared according to example 2 using 3-bromopiperidine-2,4-dione (example 35A) to provide *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-oxo-4,5-dihydrothiazolo[5,4-c]pyridin-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (4.51 mg, 12.3% yield) as a white solid containing 10% *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-oxo-4,5,6,7-tetrahydrothiazolo[5,4-c]pyridin-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide.

25 **[00270]** MS (ESI) *m/z* 489.1 (M+H)

[00271] ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.20 (s, 1H), 7.32 – 7.47 (m, 1H), 7.29 (s, 1H), 6.62 (d, 1H, *J* = 7.03 Hz), 4.66 (q, 2H, *J* = 7.19 Hz), 4.07 (s, 3H), 2.88 – 3.06 (m, 2H), 1.44 (t, 3H, *J* = 7.03 Hz), 0.72 – 0.85 (m, 4H), 0.60 – 0.72 (m, 4H)

30

Example 36



Preparation of ethyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-6,7-dihydrothiazolo[5,4-c]pyridine-5(4H)-carboxylate


5

[00272] This compound was prepared according to example 2 using 3-bromo-4-oxo-piperidine-1-carboxylic acid ethyl ester to provide ethyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-6,7-dihydrothiazolo[5,4-c]pyridine-5(4H)-carboxylate (5 mg, 10 15.36 % yield) as a white solid.

[00273] MS (ESI) m/z 549.2 (M+H)

[00274] ^1H NMR (400 MHz, DMSO- d_6) δ ppm 10.83 (br s, 1H), 8.12 (s, 1H), 7.23 (s, 1H), 4.56 – 4.67 (m, 4H), 4.10 (q, 2H, $J = 7.19$ Hz), 4.04 (s, 3H), 3.72 (t, 2H, $J = 5.52$ Hz), 2.87 – 2.98 (m, 2H), 2.64 – 2.73 (m, 2H), 1.38 (t, 3H, $J = 7.03$ Hz), 1.22 15 (t, 3H, $J = 7.15$ Hz), 0.71 – 0.81 (m, 4H), 0.60 – 0.71 (m, 4H)

Example 37



Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4,5,6,7-tetrahydrothiazolo[5,4-c]pyridin-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide


[00275] This compound was prepared according to example 2 using 3-bromopiperidin-4-one, hydrobromide (33.8 mg, 0.130 mmol) to provide *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4,5,6,7-tetrahydrothiazolo[5,4-c]pyridin-2- 25

ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (30 mg, 60 % yield) as a pale yellow solid.

[00276] MS (ESI) m/z 477.3 (M+H)

[00277] ^1H NMR (400 MHz, DMSO- d_6) δ ppm 9.08 (br s, 2H), 8.25 (s, 1H), 7.27 (s, 1H), 4.63 (q, 2H, $J = 6.78$ Hz), 4.38 (br s, 2H), 4.06 (s, 3H), 2.85 – 2.98 (m, 4H), 1.38 (t, 3H, $J = 7.03$ Hz), 0.73 – 0.81 (m, 4H), 0.62 – 0.71 (m, 4H)

Example 38



10


4-(5-cyanothiazol-2-ylamino)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00278] To a mixture of 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example 1J, 60 mg, 0.177 mmol), 2-chlorothiazole-5-carbonitrile (38.5 mg, 0.266 mmol), Pd₂(dba)₃ (24.4 mg, 0.027 mmol), BINAP (49.7 mg, 0.080 mmol) and sodium *tert*-butoxide (27.3 mg, 0.284 mmol) was added DME (1 ml). The mixture was purged with argon gas for 5 min and heated to 85 °C for 3 h. The reaction mixture showed ~60% conversion by HPLC and was filtered through a plug of Celite and concentrated. The crude product was purified by preparative HPLC to afford 4-(5-cyanothiazol-2-ylamino)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide as a off-white solid.

[00279] MS (ESI) m/z 447.2 (M+H).

[00280] ^1H NMR (500 MHz, CD₃OD-CDCl₃) δ : 8.60 (s, 1H), 8.04 (s, 1H), 7.56 (s, 4H), 7.13 (s, 1H), 4.71 (q, $J = 7.1$ Hz, 2H), 4.20 (s, 3H), 2.84 - 2.91 (m, 2H), 1.52 (t, $J = 7.1$ Hz, 3H), 0.83 - 0.90 (m, 5H), 0.74 - 0.80 (m, 4H).

Example 39




4-(benzo[d]thiazol-2-ylamino)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

- 5 **[00281]** To a solution of 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **1J**, 21.2 mg, 0.063 mmol) in DMF (626 μ L) was added sodium hydride (10.02 mg, 0.251 mmol) in one portion. After 20 min, 2-bromo-1,3-benzothiazole (53.6 mg, 0.251 mmol) was added. The reaction mixture was stirred at room temperature for 16h. The reaction
- 10 mixture was quenched with water and diluted with ethyl acetate. The aqueous layer was extracted with ethyl acetate. The combined organics were washed with 10% LiCl, dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The crude residue was purified by flash chromatography using an Isco 4 g column eluting with 0 - 5% MeOH/CH₂Cl₂ to yield 4-(benzo[d]thiazol-2-ylamino)-*N,N*-
- 15 dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (6.8 mg, 21.87 % yield) as an off-white solid.


[00282] MS (ESI) *m/z* 472.2 (M+H)

- [00283]** ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 8.17 (s, 1H), 7.95 (d, 1H, *J* = 7.97 Hz), 7.67 (d, 1H, *J* = 8.25 Hz), 7.39 (t, 1H, *J* = 7.97 Hz), 7.27 (s, 1H), 7.22 (t, 1H, *J* =
- 20 8.52 Hz), 4.70 (q, 2H, *J* = 7.06 Hz), 4.06 (s, 3H), 2.88 – 2.99 (m, 2H), 1.65 (br s, 1H), 1.45 (t, 3H, *J* = 7.01 Hz), 0.74 – 0.84 (m, 4H), 0.64 – 0.73 (m, 4H)

Example 40




- 25 **Preparation of *N,N*-dicyclopropyl-6-ethyl-4-(5-fluorobenzo[d]thiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

40A Preparation of 5-fluoro-2-(methylthio)benzo[d]thiazole

[00284] To a solution of 5-fluoro-2-mercaptobenzothiazole (0.15 g, 0.810 mmol) in THF (8.10 mL) cooled to 0 °C was added sodium hydride (0.036 g, 0.891 mmol). After stirring 10 min, iodomethane (0.076 mL, 1.215 mmol) was added and the reaction mixture was slowly warmed to room temperature over 2 h. The reaction was diluted with ethyl acetate and quenched with saturated aqueous sodium bicarbonate. The organic layer was washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. 5-fluoro-2-(methylthio)benzo[d]thiazole (0.178 g, 110 % yield) was isolated as a white solid. Material was used without any further purification.

[00285] MS (ESI) *m/z* 200.0 (M+H)

[00286] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 7.68 (dd, 1H, *J* = 8.81, 5.04 Hz), 7.56 (dd, 1H, *J* = 9.57, 2.52 Hz), 7.07 (td, 1H, *J* = 8.81, 2.52 Hz), 2.80 (s, 3H)

40B Preparation of 5-fluoro-2-(methylsulfinyl)benzo[d]thiazole

[00287] 5-Fluoro-2-(methylthio)benzo[d]thiazole (example **40A**, 0.178 g, 0.893 mmol) was dissolved in CH₂Cl₂ (5.96 mL). mCPBA (0.154 g, 0.893 mmol) was added in one portion and the reaction mixture was stirred at room temperature. The reaction was quenched with saturated aqueous sodium bicarbonate and diluted with dichloromethane. The aqueous layer was extracted with dichloromethane (2 x). The combined organic were washed with brine, dried over anhydrous magnesium sulfate, filtered, and concentrated *in vacuo*. 5-Fluoro-2-(methylsulfinyl)benzo[d]thiazole (0.180 g, 94 % yield) was isolated as a white solid and used without further purification.

[00288] MS (ESI) m/z 216.0 (M+H)

[00289] ^1H NMR (500 MHz, CHLOROFORM- d) δ ppm 7.97 (dd, 1H, $J = 9.07$, 4.95 Hz), 7.76 (dd, 1H, $J = 9.07$, 2.47 Hz), 7.27 – 7.34 (m, 1H), 3.10 (s, 3H)

5 **40** Preparation of *N,N*-dicyclopropyl-6-ethyl-4-(5-fluorobenzo[d]thiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00290] To a solution of 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **1J**, 50 mg, 0.148 mmol) in DMF (1.5mL) was added sodium hydride (17.73 mg, 0.443 mmol).

10 After 20 min, 5-fluoro-2-(methylsulfinyl)benzo[d]thiazole (example **40B**, 95 mg, 0.443 mmol) was added in one portion. The reaction mixture was stirred at room temperature. The reaction mixture was diluted with ethyl acetate and quenched with water. The aqueous layer was extracted with ethyl acetate (2 x). The combined organic layers were washed with 10% lithium chloride solution (2 x), dried over


15 anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The crude residue was purified by flash chromatography using an Isco 12 g column eluting with 0-5% MeOH/ CH_2Cl_2 to provide *N,N*-dicyclopropyl-6-ethyl-4-(5-fluorobenzo[d]thiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (13 mg, 17.07 % yield).

20 [00291] MS (CI) m/z 490.2 (M+H)

[00292] ^1H NMR (400 MHz, DMSO- d_6) δ ppm 11.53 (br s, 1H), 8.19 (s, 1H), 7.95 – 8.02 (m, 1H), 7.47 (d, 1H, $J = 9.82$ Hz), 7.28 (s, 1H), 7.09 (t, 1H, $J = 8.18$ Hz), 4.69 (q, 2H, $J = 7.13$ Hz), 4.06 (s, 3H), 2.89 – 3.02 (m, 2H), 1.45 (t, 3H, $J = 7.05$ Hz), 0.74 – 0.81 (m, 4H), 0.63 – 0.71 (m, 4H)


25

Example 41



Preparation of *N,N*-dicyclopropyl-4-(1,3-dimethyl-1*H*-pyrazolo[3,4-*d*]thiazol-5-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide


- 5 **41A** Preparation of *N,N*-dicyclopropyl-6-ethyl-4-isothiocyanato-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide



- [00293] To a mixture of 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (194.5 mg, 0.575 mmol) in dichloromethane (2874 μ L) in a round bottom flask under nitrogen was added 1,1'-thiocarbonyldi-2(1*H*)-pyridone (133 mg, 0.575 mmol). The reaction mixture was stirred at room temperature for 16h. The reaction mixture was concentrated under reduced pressure. *N,N*-Dicyclopropyl-6-ethyl-4-isothiocyanato-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide was isolated as a yellow sticky solid. Material was used as is in subsequent reactions.

[00294] MS (ESI) m/z 381.1 (M+H)

- 20 **41B** Preparation of *N,N*-dicyclopropyl-4-(3-(1,3-dimethyl-1*H*-pyrazol-5-yl)thioureido)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide



- [00295] To a flask charged with *N,N*-dicyclopropyl-6-ethyl-4-isothiocyanato-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (example 41A, 0.0342 g, 0.090 mmol) was added THF (0.449 mL). 5-Amino-1,3-dimethylpyrazole (0.012 g, 0.108 mmol) was added followed by sodium hydride

(3.60 mg, 0.090 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was diluted with ethyl acetate and quenched with saturated aqueous sodium bicarbonate. The aqueous layer was extracted with ethyl acetate. The combined organics were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The crude material was used without further purification.

[00296] MS (ESI) m/z 492.2 (M+H)


41 Preparation of *N,N*-dicyclopropyl-4-(1,3-dimethyl-1*H*-pyrazolo[3,4-
d]thiazol-5-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-
7-carboxamide

[00297] *N,N*-Dicyclopropyl-4-(3-(1,3-dimethyl-1*H*-pyrazol-5-yl)thioureido)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **41B**, 41 mg, 0.083 mmol) was dissolved in chloroform (2780 μ L) and cooled to 0 °C. NIS (18.76 mg, 0.083 mmol) was slowly added and the reaction was warmed to room temperature. The reaction mixture was concentrated. Methanol and ether were added and the mixture was filtered. After standing for a week, product precipitated out of the solution. The product was triturated with methanol. *N,N*-Dicyclopropyl-4-(1,3-dimethyl-1*H*-pyrazolo[3,4-d]thiazol-5-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide was isolated as a pale yellow solid (8.1 mg, 20% yield).

[00298] MS (ESI) m/z 490.3 (M+H)

[00299] ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 11.39 (s, 1H), 8.17 (s, 1H), 7.26 (s, 1H), 4.66 (q, 2H, $J = 7.03$ Hz), 4.05 (s, 3H), 3.85 (s, 3H) 2.88 – 3.00 (m, 2H), 2.33 (s, 3H), 1.43 (t, 3H, $J = 7.03$ Hz), 0.72 – 0.82 (m, 4H), 0.61 – 0.73 (m, 4H)

Example 42




Methyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-5-methylthiazole-4-carboxylate

[00300] Methyl 3-bromo-2-oxobutanoate (320 mg, 1.642 mmol) was added to a suspension of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **2B**, 435 mg, 1.094 mmol) in EtOH (10 ml) and the reaction mixture was heated at 65 °C for 2h. The solvent was removed *in vacuo* and the residue purified by silica gel chromatography (2-10% MeOH/dichloromethane) to give methyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-5-methylthiazole-4-carboxylate (380 mg, 70% yield) as yellow solid.

[00301] MS (ESI) *rt* = 0.87 min, *m/z* 494 (M+H)


[00302] ¹H NMR (CHLOROFORM-*d*) δ: 9.41 (br. s., 1H), 7.78 (s, 1H), 6.89 (s, 1H), 4.73 (q, *J* = 7.0 Hz, 2H), 4.03 (s, 3H), 3.95 (s, 3H), 2.80 - 2.87 (m, 2H), 2.77 (s, 3H), 1.54 (t, *J* = 7.0 Hz, 3H), 0.83 - 0.90 (m, 4H), 0.73 - 0.80 (m, 4H).

Example 43



2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-*N,N*,5-trimethylthiazole-4-carboxamide

43A Preparation of 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-5-methylthiazole-4-carboxylic acid



[00303] 1N Aqueous NaOH (0.962 mL, 0.962 mmol) was added to a suspension of methyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-5-methylthiazole-4-carboxylate (example 41, 0.19 g, 0.385 mmol) in ethanol (2 mL) and the mixture was heated at 60 °C for 2h. The reaction was cooled to room temperature and acidified with 1N HCl to pH 4. The reaction mixture was concentrated and the residue triturated 5x with methanol. The combined methanol washes were concentrated, and the remaining solid washed 3x with 2:1 ether/dichloromethane provided 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-5-methylthiazole-4-carboxylic acid (105 mg, 57% yield).

[00304] MS (ESI) rt = 0.79 min, m/z 480 (M+H)

43 Preparation of 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-N,N,5-trimethylthiazole-4-carboxamide


[00305] A mixture of 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-5-methylthiazole-4-carboxylic acid (43A, 30 mg, 0.063 mmol), dimethylamine, 40% aqueous (0.016 mL, 0.125 mmol) and HATU (35.7 mg, 0.094 mmol) in DMF (1 mL) was stirred at room temperature for 3h. The solvent was evaporated off and the residue purified by silica gel chromatography (2-10% MeOH/dichloromethane). The material was further purified by preparative HPLC to give of 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo [2,3-b]pyridin-4-ylamino)-N,N,5-trimethylthiazole-4-carboxamide (20 mg, 62.5% yield) as off-white solid.

[00306] MS (ESI) m/z 507 (M+H)

[00307] ¹H NMR (400 MHz, CHLOROFORM-d) δ: 7.79 (s, 1H), 6.90 (s, 1H), 4.73 (q, J = 7.0 Hz, 2H), 4.05 (s, 3H), 3.12 (s, 3H), 3.11 (s, 3H), 2.80 - 2.87 (m, 2H), 2.53 (s, 3H), 1.53 (t, J = 7.2 Hz, 3H), 0.83 - 0.91 (m, 4H), 0.74 - 0.80 (m, 4H)

Examples **44-49** were prepared using the similar protocol as described for example **42**

Example 44



5


***N*-(2-cyanoethyl)-2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-N,5-dimethylthiazole-4-carboxamide**

10 **[00308]** MS (ESI) m/z 546 (M+H).

[00309] ^1H NMR (400 MHz, CHLOROFORM- d) δ ppm 9.34 (br. s., 1 H) 7.75 (d, $J=5.94$ Hz, 1 H) 6.90 (s, 1 H) 4.74 (q, $J=7.04$ Hz, 2 H) 4.06 (s, 3 H) 3.80 (t, $J=6.71$ Hz, 2 H) 3.11 - 3.32 (m, 3 H) 2.95 (t, $J=6.71$ Hz, 2 H) 2.78 - 2.91 (m, 2 H) 2.58 (d, $J=15.85$ Hz, 3 H) 1.54 (t, $J=7.04$ Hz, 3 H) 0.70 - 0.94 (m, 8 H).

15

Example 45



2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-N,5-dimethylthiazole-4-carboxamide


20

[00310] MS (ESI) m/z 493 (M+H).

[00311] ^1H NMR (500 MHz, CHLOROFORM- d) δ ppm 7.84 (s, 1 H) 7.40 (d, $J=4.99$ Hz, 1 H) 6.90 (s, 1 H) 4.73 (q, $J=6.94$ Hz, 2 H) 4.08 (s, 3 H) 2.99 (d, $J=4.99$ Hz, 3 H) 2.73 - 2.91 (m, 5 H) 2.14 (s, 1 H) 1.53 (t, $J=7.07$ Hz, 3 H) 0.66 - 0.97 (m, 8 H).

25

Example 46



2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-N-(2-methoxyethyl)-N,5-dimethylthiazole-4-carboxamide

5


[00312] MS (ESI) m/z 551 (M+H).

[00313] ^1H NMR (500 MHz, CHLOROFORM- d) δ ppm 7.73 (d, $J=3.05$ Hz, 1 H)

6.90 (s, 1 H) 4.74 (q, $J=7.21$ Hz, 2 H) 4.05 (s, 3 H) 3.72 (dd, $J=13.73, 4.58$ Hz, 2 H)

10 3.61 - 3.68 (m, 1 H) 3.57 (t, $J=5.41$ Hz, 1 H) 3.27 - 3.45 (m, 3 H) 3.17 (d, $J=8.32$ Hz, 3 H) 2.79 - 2.88 (m, 2 H) 2.54 (d, $J=5.83$ Hz, 3 H) 1.54 (t, $J=7.07$ Hz, 3 H) 0.73 - 0.92 (m, 8 H).

Example 47



15


4-(4-(azetidine-1-carbonyl)-5-methylthiazol-2-ylamino)-N,N-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00314] MS (ESI) m/z 519 (M+H).

20 **[00315]** ^1H NMR (500 MHz, CHLOROFORM- d) δ ppm 8.50 - 9.65 (m, 1 H) 8.00 (s, 1 H) 6.88 (s, 1 H) 4.68 (q, $J=7.21$ Hz, 2 H) 4.59 (t, $J=7.63$ Hz, 2 H) 4.18 (t, $J=7.77$ Hz, 2 H) 4.09 (s, 3 H) 2.78 - 2.89 (m, 2 H) 2.72 (s, 3 H) 2.30 (quin, $J=7.70$ Hz, 2 H) 1.49 (t, $J=7.07$ Hz, 3 H) 0.72 - 0.92 (m, 8 H).

25

Example 48




2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-N-ethyl-N,5-dimethylthiazole-4-carboxamide

5 **[00316]** MS (ESI) m/z 521 (M+H).

[00317] ^1H NMR (400 MHz, CHLOROFORM- d) δ ppm 7.77 (s, 1 H) 6.89 (s, 1 H) 4.74 (q, $J=7.04$ Hz, 2 H) 4.04 (s, 3 H) 3.36 – 3.66 (m, 2 H) 3.07 (d, $J=14.09$ Hz, 3 H) 2.72 – 2.90 (m, 2 H) 2.51 (s, 3 H) 2.11 (s, 1 H) 1.54 (t, $J=7.04$ Hz, 3 H) 1.13 – 1.35 (m, 3 H) 0.66 – 0.90 (m, 8 H).

10

Example 49




15 **2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-4-ylamino)-N-ethyl-5-methylthiazole-4-carboxamide**

[00318] MS (ESI) m/z 508 (M+H).


20 **[00319]** ^1H NMR (400 MHz, CHLOROFORM- d) δ ppm 7.78 (s, 1 H) 6.89 (s, 1 H) 4.73 (q, $J=7.04$ Hz, 2 H) 4.42 (q, $J=7.12$ Hz, 2 H) 4.04 (s, 3 H) 2.78 – 2.91 (m, 2 H) 2.75 (s, 3 H) 2.15 (s, 1 H) 1.54 (t, $J=7.04$ Hz, 3 H) 1.44 (t, $J=7.04$ Hz, 3 H) 0.68 – 0.91 (m, 8 H).

Example 50



2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-N,N,4-trimethylthiazole-5-carboxamid

- 5 **50A** Preparation of methyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4-methylthiazole-5-carboxylate




10

- [00320]** Methyl 2-bromo-3-oxobutanoate (213 mg, 1.094 mmol, prepared following the procedure described by Abu T. Khan et al. in *Tetrahedron Letters* **2006**, 47, 2751-2754) was added to a suspension of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example 15 **2B**, 290 mg, 0.730 mmol) in ethanol (5 ml) and the reaction mixture was heated at 65 °C for 2h. LC/MS shows completion of reaction (reaction remained as suspension all the time). The reaction mixture was concentrated to remove ethanol, and taken up in dichloromethane. The solution was washed with saturated sodium bicarbonate and the aqueous layer was back extracted with dichloromethane. The combined organic 20 layers were dried over magnesium sulfate, filtered, and concentrated *in vacuo*. The crude was purified by flash chromatography using an automated ISCO system (40 g column, eluting with 0-5% methanol / dichloromethane) to afford methyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4-methylthiazole-5-carboxylate (301 mg, 0.610 mmol, 84 % 25 yield) as a white solid.

[00321] MS (ESI) *m/z* 494.4 (M+H).

[00322] ^1H NMR (400 MHz, CHLOROFORM- d) δ : 7.75 (s, 1H), 6.92 (s, 1H), 4.76 (q, J = 7.0 Hz, 2H), 4.08 (s, 3H), 3.92 (s, 3H), 2.82 - 2.89 (m, 2H), 2.71 (s, 3H), 1.59 (t, J = 7.0 Hz, 3H), 0.85 - 0.93 (m, 4H), 0.75 - 0.82 (m, 4H)

5 **50B** Preparation of 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo [4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4-methylthiazole-5-carboxylic acid



[00323] A mixture of methyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4-methylthiazole-5-carboxylate (example **50A**, 288 mg, 0.583 mmol) and 1N NaOH (5.83 mL, 5.830 mmol) in methanol (5 mL) was heated at 65 °C for 18h. Methanol was evaporated and the aqueous solution was acidified to pH 4 with 1N HCl, solids crashed out of solution and were retained after filtration to afford the desired acid in pure form. The aqueous mixture was then extracted with dichloromethane, The organic layer was dried over magnesium sulfate, filtered and concentrated *in vacuo*. The crude was purified by flash chromatography using an automated ISCO system (40 g column, eluting with 0-8% methanol / dichloromethane). A total of 260 mg of 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4-methylthiazole-5-carboxylic acid (93 % yield) was obtained as a yellow solid.

[00324] MS (ESI) m/z 480.2 ($M+H$).

[00325] ^1H NMR (500 MHz, DMSO- d_6) δ : 8.03 (br. s., 1H), 7.06 (s, 1H), 4.40 (q, J = 7.0 Hz, 2H), 3.86 (s, 3H), 2.96 (s, 1H), 2.69 - 2.76 (m, 2H), 2.36 (s, 3H), 1.20 (t, J = 6.9 Hz, 3H), 0.56 (d, J = 5.5 Hz, 4H), 0.45 - 0.50 (m, 4H).

50 Preparation of 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo [4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-*N,N*,4-trimethylthiazole-5-carboxamide

[00326] A mixture of 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4-methylthiazole-5-carboxylic acid (example 50B, 51 mg, 0.106 mmol), dimethylamine (40% in water, 0.027 mL, 0.213 mmol), HATU (52.6 mg, 0.138 mmol) and 2,6-lutidine (0.025 mL, 0.213 mmol) in DMF (1 mL) was stirred at room temperature for 2h. Solvent was evaporated and the residue was purified by flash chromatography on silica gel using an automated ISCO system (24 g column, eluting with 2-8% methanol / dichloromethane). The product was further purified by preparative HPLC. 2-(7-(Dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo [4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-*N,N*,4-trimethylthiazole-5-carboxamide (17 mg, 0.032 mmol, 30.0 % yield) was obtained as a off-white solid.


[00327] MS (ESI) m/z 507.1 (M+H).

[00328] ^1H NMR (400 MHz, METHANOL- d_4) δ : 8.75 (s, 1H), 7.27 (s, 1H), 4.66 (q, J = 7.0 Hz, 2H), 4.21 (s, 3H), 3.19 (s, 6H), 2.95 - 3.02 (m, 2H), 2.39 (s, 3H), 1.47 (t, J = 7.2 Hz, 3H), 0.86 - 0.93 (m, 4H), 0.78 - 0.84 (m, 4H)

Examples 51-55 were synthesized using the same method as used for 50.

20


Example 51



N-(2-cyanoethyl)-2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-*N*,4-dimethylthiazole-5-carboxamide

[00329] MS (ESI) m/z 546.1 (M+H).

[00330] ^1H NMR (CDCl_3) δ : 7.77 (s, 1H), 6.89 (s, 1H), 4.71 (q, J = 6.9 Hz, 2H), 4.05 (s, 3H), 3.81 (t, J = 6.6 Hz, 2H), 3.31 (s, 3H), 2.78 - 2.86 (m, 4H), 2.44 (s, 3H), 1.53 (t, J = 7.0 Hz, 3H), 0.83 - 0.90 (m, 4H), 0.73 - 0.79 (m, 4H).

Example 52


2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-N-ethyl-N,4-dimethylthiazole-5-carboxamide

5

[00331] MS (ESI) m/z 521.1 (M+H).

[00332] ^1H NMR (400 MHz, CDCl_3) δ : 7.74 (s, 1H), 6.88 (s, 1H), 4.72 (q, $J = 7.0$ Hz, 2H), 4.04 (s, 3H), 3.57 (q, $J = 7.2$ Hz, 2H), 3.11 (s, 3H), 2.79 - 2.86 (m, 2H), 2.40 (s, 3H), 1.51 (t, $J = 7.0$ Hz, 3H), 1.25 (t, $J = 7.2$ Hz, 3H), 0.82 - 0.90 (m, 4H), 0.73 -

10 0.79 (m, 4H).

Example 53

2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-N-(2-methoxyethyl)-4-methylthiazole-5-carboxamide


15

[00333] MS (ESI) m/z 537.1 (M+H).

[00334] ^1H NMR (400 MHz, CDCl_3) δ : 7.74 (s, 1H), 6.89 (s, 1H), 6.13 (t, $J = 4.7$ Hz, 1H), 4.74 (q, $J = 7.0$ Hz, 2H), 4.05 (s, 3H), 3.65 (q, $J = 5.1$ Hz, 2H), 3.58 (t, $J = 4.8$ Hz, 2H), 3.42 (s, 3H), 2.79 - 2.87 (m, 2H), 2.65 (s, 3H), 1.58 (t, $J = 7.0$ Hz, 3H), 0.83 - 0.90 (m, 4H), 0.73 - 0.80 (m, 4H).

20

Example 54




2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-N,4-dimethylthiazole-5-carboxamide

5 [00335] MS (ESI) m/z 493.1 (M+H).

[00336] ^1H NMR (400 MHz, CDCl_3) δ : 7.77 (s, 1H), 6.86 (s, 1H), 5.87 - 5.94 (m, J = 4.6 Hz, 1H), 4.66 (q, J = 7.1 Hz, 2H), 4.04 (s, 3H), 3.04 (d, J = 4.6 Hz, 3H), 2.79 - 2.87 (m, 2H), 2.62 (s, 3H), 1.51 (t, J = 7.0 Hz, 3H), 0.83 - 0.90 (m, 4H), 0.73 - 0.80 (m, 4H).

10

Example 55




4-(5-(azetidine-1-carbonyl)-4-methylthiazol-2-ylamino)-N,N-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

15

[00337] MS (ESI) m/z 519.1 (M+H).





[00338] ^1H NMR (400 MHz, CDCl_3) δ : 7.73 (s, 1H), 6.90 (s, 1H), 4.73 (q, J = 7.0 Hz, 2H), 4.33 (br. s., 4H), 4.05 (s, 3H), 2.80 - 2.87 (m, 2H), 2.66 (s, 3H), 2.33 - 2.43 (m, 2H), 1.55 (t, J = 7.2 Hz, 3H), 0.83 - 0.90 (m, 4H), 0.73 - 0.80 (m, 4H).





[00339] Examples **56-70** were synthesized using same procedure as used for **50**







5

Ex. No.	R	Name	Retention time ((min)	MS (ESI) m/z
56		<i>N,N</i> -dicyclopropyl-6-ethyl-1-methyl-4-((4-methyl-5-((4-methyl-1-piperazinyl)carbonyl)-1,3-thiazol-2-yl)amino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.43	562.31
57		<i>N,N</i> -dicyclopropyl-6-ethyl-1-methyl-4-((4-methyl-5-(4-morpholinylcarbonyl)-1,3-thiazol-2-yl)amino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.70	549.25
58		<i>N,N</i> -dicyclopropyl-6-ethyl-1-methyl-4-((4-methyl-5-((2-(4-morpholinyl)ethyl)carbonyl)-1,3-thiazol-2-yl)amino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.46	592.3


Ex. No.	R	Name	Retention time ((min)	MS (ESI) m/z
59		4-((5-((2-acetamidoethyl)carbamoyl)-4-methyl-1,3-thiazol-2-yl)amino)- <i>N,N</i> -dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.51	564.27
60		4-((5-((cyanomethyl)(methyl)carbamoyl)-4-methyl-1,3-thiazol-2-yl)amino)- <i>N,N</i> -dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.78	532.25
61		4-((5-((cyanomethyl)carbamoyl)-4-methyl-1,3-thiazol-2-yl)amino)- <i>N,N</i> -dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.72	518.24
62		<i>N,N</i> -dicyclopropyl-4-((5-((1,1-dioxidotetrahydro-3-thiophenyl)carbamoyl)-4-methyl-1,3-thiazol-2-yl)amino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.68	597.24

Ex. No.	R	Name	Retention time ((min))	MS (ESI) m/z
63		<i>N,N</i> -dicyclopropyl-6-ethyl-4-((5-((trans-4-hydroxycyclohexyl)carbamoyl)-4-methyl-1,3-thiazol-2-yl)amino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.60	577.3
64		<i>N,N</i> -dicyclopropyl-6-ethyl-4-((5-(((3 <i>S</i>)-3-hydroxy-1-pyrrolidinyl)carbonyl)-4-methyl-1,3-thiazol-2-yl)amino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.51	549.28
65		<i>N,N</i> -dicyclopropyl-6-ethyl-1-methyl-4-((4-methyl-5-(methyl(2-(methylsulfonyl)ethyl)carbamoyl)-1,3-thiazol-2-yl)amino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.59	599.26
66		<i>N,N</i> -dicyclopropyl-6-ethyl-4-((5-((3-methoxy-1-azetidyl)carbonyl)-4-methyl-1,3-thiazol-2-yl)amino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.80	549.28

Ex. No.	R	Name	Retention time ((min)	MS (ESI) m/z
67		<i>N,N</i> -dicyclopropyl-4-((5-((3,3-difluorocyclobutyl)carbamoyl)-4-methyl-1,3-thiazol-2-yl)amino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	2.06	569.27
68		<i>N,N</i> -dicyclopropyl-4-((5-((2,2-difluoroethyl)carbamoyl)-4-methyl-1,3-thiazol-2-yl)amino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.88	543.23
69		<i>N,N</i> -dicyclopropyl-6-ethyl-1-methyl-4-((4-methyl-5-(tetrahydro-2H-pyran-4-ylcarbamoyl)-1,3-thiazol-2-yl)amino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.87	563.26
70		<i>N,N</i> -dicyclopropyl-4-((5-((1,1-dioxido-4-thiomorpholinyl)carbamoyl)-4-methyl-1,3-thiazol-2-yl)amino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide	1.66	597.26

Analysis Conditions- Column- Mac-Mod Halo 4.6 x 50 mm 2.7 um; Gradient-4 min from 0%B to 100%B; Flow rate- 4 mL/min; Solvent- A = 5 :95 Acetonitrile : Water; B = 90 :5 Acetonitrile : Water; Modifier = 10 mM NH₄OAc

5


Example 71

***N,N*-dicyclopropyl-6-ethyl-4-(4-ethyl-5-(1,1-dioxothiomorpholine-4-carbonyl)thiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

10

71A Preparation of methyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4-ethylthiazole-5-carboxylate

15



[00340] To a stirred suspension of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (103 mg, 0.259 mmol) in EtOH (5 ml) was added freshly prepared methyl 2-bromo-3-oxopentanoate¹ (81 mg, 0.389 mmol) at room temperature. The slurry was then stirred at 65 °C for 2h. The reaction mixture was then cooled to room temperature and concentrated *in vacuo* to afford a yellow solid. The solid was then taken up in methylene chloride and washed with saturated aqueous sodium bicarbonate solution. The organic phase was dried over anhydrous magnesium sulfate and concentrated *in*


vacuo. The crude residue was then purified by silica-gel flash chromatography using an Isco 40 g column eluting from 0-5% methanol/methylene chloride to afford methyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4-ethylthiazole-5-carboxylate (37.5 mg, 28.5 % yield) as a white solid.

[00341] MS (ESI) m/z 508.4 (M+H)

[00342] ^1H NMR (CDCl_3) δ ppm 9.38 (br s, 1 H), 7.75 (s, 1 H), 6.92 (s, 1 H), 4.76 (q, 2 H, $J = 6.94$ Hz), 4.08 (s, 3H), 3.92 (s, 3H), 3.14 (q, 2H, $J = 7.68$ Hz), 2.85 (m, 2H), 1.59 (t, 3H, $J = 6.94$ Hz), 1.34 (t, 3H, $J = 7.49$ Hz), 0.78-0.91 (m, 8H)

10

71B Preparation of 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4-ethylthiazole-5-carboxylic acid



[00343] To a stirred solution of methyl 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4-ethylthiazole-5-carboxylate (example **71A**, 103 mg, 0.203 mmol) in MeOH (2mL) was added NaOH (1.420 mL, 1.420 mmol). The solution was then stirred at 65 °C for 18 hrs. The reaction mixture was then cooled to room temperature and concentrated *in vacuo* to afford yellow slurry. The mixture was then diluted with water, brought to pH 4 with 1N HCl, and the precipitate collected by vacuum filtration. The filtrate was then extracted with CH_2Cl_2 ; the organic layer was dried over anhydrous magnesium sulfate and concentrated *in vacuo*. The crude residue was then purified by silica-gel flash chromatography using an Isco 40 g column eluting from 0-8% MeOH/ CH_2Cl_2 . Desired fractions were combined with the collected precipitate and concentrated to afford 2-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4-ethylthiazole-5-carboxylic acid (126 mg, 126 % yield) as an orange/yellow solid.

[00344] MS (ESI) m/z 494.2 (M+H)

[00345] ¹H NMR (CDCl₃) δ ppm 8.23 (s, 1 H), 7.28 (s, 1H), 4.62 (q, 2 H, *J* = 6.94 Hz), 4.07 (s, 3H), 3.04 (q, 2H, *J* = 7.49 Hz), 3.00 (m, 2H), 1.42 (t, 3H, *J* = 7.07 Hz), 1.23 (t, 3H, *J* = 7.63 Hz), 0.67-0.79 (m, 8H)


5 **71** Preparation of *N,N*-dicyclopropyl-6-ethyl-4-(4-ethyl-5-(1,1-dioxothiomorpholine-4-carbonyl)thiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00346] To a stirred solution of 2-(7-(dicyclopropylcarbonyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4-ethylthiazole-5-carboxylic acid (example **69B**, 33 mg, 0.067 mmol), HATU (50.8 mg, 0.134 mmol),
10 2,6-lutidine (0.020 mL, 0.174 mmol), and DMF (1 mL) was added thiomorpholine 1,1-dioxide (27.1 mg, 0.201 mmol). The solution was then stirred at 50 °C for 1.5 hrs. The reaction was then cooled to room temperature and diluted with methylene chloride. The solution was then washed with saturated aqueous sodium bicarbonate,
15 10% LiCl in H₂O, and saturated aqueous sodium bicarbonate again. The organic phase was dried over anhydrous magnesium sulfate and concentrated *in vacuo*. The crude residue was then purified by silica-gel flash chromatography using an Isco 40 g column eluting from 0-6% MeOH/CH₂Cl₂ to give *N,N*-dicyclopropyl-6-ethyl-4-(4-ethyl-5-(1,1-dioxothiomorpholine -4-carbonyl)thiazol-2-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (21.7 mg, 52.6 % yield)
20 as a pale yellow solid.

[00347] MS (ESI) *m/z* 611.3 (M+H)


[00348] ¹H NMR (CDCl₃) δ ppm 7.68 (s, 1 H), 6.83 (s, 1 H), 4.65 (q, 2 H, *J* = 7.04), 4.13 (m, 4H), 3.99 (s, 3H), 3.05 (m, 4H), 2.76 (m, 2H), 2.68 (q, 2H, *J* = 7.56
25 Hz), 1.46 (t, 3H, *J* = 7.04 Hz), 1.25 (t, 3H, *J* = 7.59 Hz), 0.68-0.80 (m, 8H)

Example 72



N,N-dicyclopropyl-6-ethyl-1-methyl-4-(5-methylisoxazol-3-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

72A Preparation of 4,4-bis(methylthio)but-3-en-2-one



5


[00349] A solution of acetone (2.205 mL, 30.0 mmol) in carbon disulfide (1.81 mL, 30.0 mmol) was added to a suspension of sodium *tert*-butoxide (5.76 g, 60.0 mmol) in benzene (30 mL) while keeping the reaction temperature not exceeding 10 °C. The reaction mixture was stirred at room temperature for 3h and iodomethane (3.75 mL, 60.0 mmol) was added at 10 °C. The resulting reaction mixture was stirred at room temperature overnight. The reaction mixture was diluted with ethyl acetate and washed twice with water. The organic layer was dried over magnesium sulfate, filtered and concentrated *in vacuo*. The crude product was purified by flash chromatography on silica gel using an automated ISCO system (80 g column, eluting with 10-35% ethyl acetate / hexanes) to give 4,4-bis(methylthio)but-3-en-2-one (3.456 g, 71.0 % yield) as a yellow solid.

15

[00350] ¹H NMR (400 MHz, CDCl₃) δ: 6.04 (s, 1H), 2.47 (s, 3H), 2.45 (s, 3H), 2.20 (s, 3H).

20

72B Preparation of (*Z*)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(1-(methylthio)-3-oxobut-1-enylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide



25

[00351] Sodium hydride (2.84 g, 71.1 mmol) was added to a solution of 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **1J**, 8.02 g, 23.70 mmol) in DMF (60 mL) and the

reaction mixture stirred at room temperature for 1 h. 4,4-bis(methylthio)but-3-en-2-one (example **70A**, 5.77 g, 35.5 mmol) in DMF (10 ml) was added and stirring continued for three days. LC/MS showed about 75% completion. The reaction mixture was diluted with ethyl acetate and water was added. An off-white precipitate formed and was collected by filtration to give 5.1 g of product. The filtrate was transferred to a separatory funnel. The layers were separated and the aqueous layer was extracted with dichloromethane (3x50 ml), the combined organic layers were washed with saturated sodium bicarbonate and dried over magnesium sulfate, filtered and concentrated *in vacuo*. The crude product was purified by flash chromatography on silica gel using an ISCO system (220 g column, eluting with 2-8% methanol / dichloromethane) to give (*Z*)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(1-(methylthio)-3-oxobut-1-enylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (6.6 g, 61.5% yield) along with 2.2 g of recovered **1J**.

[00352] MS (ESI) *m/z* 453.3 (M+H).

15 [00353] ¹H NMR (400 MHz, DMSO-*d*₆) δ: 14.42 (s, 1H), 8.16 (s, 1H), 7.24 (s, 1H), 5.54 (s, 1H), 4.52 (q, *J* = 7.0 Hz, 2H), 4.05 (s, 3H), 2.89 - 2.97 (m, 2H), 2.43 (s, 3H), 2.14 (s, 3H), 1.34 (t, *J* = 7.0 Hz, 3H), 0.71 - 0.80 (m, 4H), 0.63 - 0.70 (m, 4H).


72 Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-methylisoxazol-3-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00354] A mixture of (*Z*)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(1-(methylthio)-3-oxobut-1-enylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (22 mg, 0.049 mmol) and hydroxylamine (50% aqueous solution, 3.13 μL, 0.051 mmol) in ethanol (1 mL) was heated at reflux for 3h and LC/MS showed completion of reaction. The reaction mixture was diluted with methanol and purified by preparative HPLC to furnish *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-methylisoxazol-3-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (9 mg, 43.3 % yield) as a white solid.

30 [00355] MS (ESI) *m/z* 420.1 (M+H).

[00356] ^1H NMR (400 MHz, METHANOL- d_4) δ : 8.45 (s, 1H), 7.26 (s, 1H), 6.58 (s, 1H), 4.63 (q, $J = 7.2$ Hz, 2H), 4.17 (s, 3H), 2.94 - 3.02 (m, 2H), 2.33 (s, 3H), 1.47 (t, $J = 7.2$ Hz, 3H), 0.85 - 0.92 (m, 4H), 0.77 - 0.84 (m, 4H).

5

Example 73

***N,N*-dicyclopropyl-4-(1,5-dimethyl-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

10


[00357] A mixture of (*Z*)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(1-(methylthio)-3-oxobut-1-enylamino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (example 72B, 70 mg, 0.155 mmol) and *tert*-butyl 1-methylhydrazinecarboxylate (0.046 mL, 0.309 mmol) in acetic acid (1 mL) was stirred at 35 °C for 4 h (monitored by LC/MS until no starting material left). Formic acid (0.5 mL) was added and stirred at 60 °C for 6 h. Solvent was evaporated and the crude product was purified by flash chromatography on silica gel using an automated ISCO system (12 g column, eluting with 2-10% methanol / dichloromethane). The impure product was further purified by preparative HPLC to give *N,N*-dicyclopropyl-4-(1,5-dimethyl-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (45 mg, 63.9 % yield) as an off-white solid.

20

[00358] MS (ESI) m/z 433.2 ($M+H$).

[00359] ^1H NMR (500 MHz, CDCl_3) δ : 8.19 (s, 1H), 7.63 (s, 1H), 6.94 (s, 1H), 6.85 (s, 1H), 4.65 (q, $J = 7.0$ Hz, 2H), 3.99 (s, 3H), 3.71 (s, 3H), 2.79 - 2.85 (m, 2H), 2.33 (s, 3H), 1.49 (t, $J = 7.2$ Hz, 3H), 0.82 - 0.87 (m, 4H), 0.72 - 0.78 (m, 4H)

25

Alternative synthesis of example 73**73A** Preparation of 4-iodo-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

- 5 **[00360]** To a stirred suspension of 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (34.4 mg, 0.102 mmol) in diiodomethane (102 μ L) at 60 °C was slowly added isoamyl nitrite (27.4 μ L, 0.203 mmol). The reaction mixture was stirred at 60 °C 5 h. After cooling to rt, the reaction mixture was diluted with dichloromethane. Saturated aqueous sodium
- 10 bicarbonate was added and the aqueous layer was extracted with dichloromethane (3x). The combined organics were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The crude residue was placed on the pump for 16 h to remove any additional diiodomethane. LCMS was deemed pure enough to be carried forward crude. The product was used immediately in the next reaction.
- 15 **[00361]** MS (ESI) m/z 450.1 (M+H)

73 Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(1,5-dimethyl-1*H*-pyrazol-3-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide


- [00362]** To a vial containing 1,5-dimethyl-1*H*-pyrazol-3-amine (13.60 mg, 0.122 mmol), Pd₂(dba)₃ (4.67 mg, 5.10 μ mol), Xantphos (5.90 mg, 10.20 μ mol), and cesium carbonate (100 mg, 0.306 mmol) was added *N,N*-dicyclopropyl-6-ethyl-4-iodo-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (45.8 mg, 0.102 mmol) in DME (1020 μ L). The reaction mixture was sparged with argon for 5 min, the vial was capped, and the reaction mixture was heated at 90 °C 1 h. The
- 25 reaction mixture was diluted with dichloromethane and vacuum filtered through Celite. The filtrate was concentrated *in vacuo*. The crude residue was purified via flash column chromatography using an ISCO 12 g column eluting with 1 - 10% MeOH/CH₂Cl₂. *N,N*-Dicyclopropyl-4-(1,5-dimethyl-1*H*-pyrazol-3-ylamino)-6-

ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (14 mg, 31.7 % yield) was isolated as an off-white solid.

[00363] MS (ESI) m/z 433.2 (M+H).


[00364] ^1H NMR (500 MHz, CDCl_3) δ : 8.19 (s, 1H), 7.63 (s, 1H), 6.94 (s, 1H),
 5 6.85 (s, 1H), 4.65 (q, $J = 7.0$ Hz, 2H), 3.99 (s, 3H), 3.71 (s, 3H), 2.79 - 2.85 (m, 2H),
 2.33 (s, 3H), 1.49 (t, $J = 7.2$ Hz, 3H), 0.82 - 0.87 (m, 4H), 0.72 - 0.78 (m, 4H)

Example 74



10 ***N,N*-dicyclopropyl-6-ethyl-4-(1-ethyl-5-methyl-1H-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**


74A Preparation of *tert*-butyl 1,3-dioxoisindolin-2-yl(ethyl)carbamate



15 **[00365]** Diisopropyl azodicarboxylate (2.92 mL, 15.00 mmol) was added in one portion to a solution of *tert*-butyl 1,3-dioxoisindolin-2-ylcarbamate (2.62 g, 10 mmol, prepared following the procedure described by Nicolas Brosse et al. in *Eur. J. Org. Chem.* 4757-4764, **2003**), triphenylphosphine (3.93 g, 15.00 mmol) and ethanol (0.691 g, 15.00 mmol) in THF (20 mL) at 0 °C and the reaction solution was stirred at
 20 room temperature for 1h (monitored by TLC until completion). Solvent was evaporated and the residue was purified by flash chromatography on silica gel using an automated ISCO system (80 g column, eluting with 5-35% ethyl acetate / hexanes) to provide *tert*-butyl 1,3-dioxoisindolin-2-yl(ethyl)carbamate (2.6 g, 90 % yield) as a white solid which was used as it in the next step

25

74B Preparation of *tert*-butyl 1-ethylhydrazinecarboxylate



[00366] Methylhydrazine (1.415 mL, 26.9 mmol) was added to a solution of *tert*-butyl 1,3-dioxoisindolin-2-yl(ethyl)carbamate (example **74A**, 5.2 g, 17.91 mmol) in THF (40 mL) at 0 °C and the reaction mixture was stirred at room temperature

5 overnight. A white precipitate formed and was filtered off through a pad of Celite, The filtrate was concentrated *in vacuo*. The residue was dissolved in ethyl acetate (50 ml) and extracted with 1N HCl (3x30 ml), the acid layer was washed with ethyl acetate (50 ml) and basified to pH 10 by addition of 20% NaOH. The basic solution was then extracted with ethyl acetate (3x50 ml) and the combined organic layers were
10 washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo* to give *tert*-butyl 1-ethylhydrazinecarboxylate (2.5 g, 87 % yield) as colorless oil.

[00367] ¹H NMR (400 MHz, CDCl₃) δ: 3.90 (br. s., 2H), 3.35 (q, J = 7.0 Hz, 2H), 1.42 (s, 9H), 1.07 (t, J = 7.0 Hz, 3H)

15 **74** Preparation of *N,N*-dicyclopropyl-6-ethyl-4-(1-ethyl-5-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

[00368] A mixture of (*Z*)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(1-(methylthio)-3-oxobut-1-enylamino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-


20 carboxamide (example **74B**, 70 mg, 0.155 mmol) and *tert*-butyl 1-ethylhydrazinecarboxylate (49.6 mg, 0.309 mmol) in acetic acid (1 mL) was stirred at 35 °C for 4 h (monitored by LC/MS until no starting material left). Formic acid (1 mL) was added and the reaction mixture stirred at 60 °C for 6 h. The solvent was evaporated and the crude product was purified by flash chromatography on silica gel
25 using an automated ISCO system (12 g column, eluting with 2-10% methanol / dichloromethane). The material was further purified by preparative HPLC to afford *N,N*-dicyclopropyl-6-ethyl-4-(1-ethyl-5-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (38 mg, 53.4 % yield) as an off-white solid.

30 [00369] MS (ESI) *m/z* 447.3 (M+H).

[00370] ^1H NMR (500 MHz, CDCl_3) δ : 8.08 (s, 1H), 7.61 (s, 1H), 6.93 (s, 1H), 6.84 (s, 1H), 4.66 (q, $J = 7.1$ Hz, 2H), 4.02 (q, $J = 7.2$ Hz, 2H), 3.98 (s, 3H), 2.79 - 2.85 (m, 2H), 2.34 (s, 3H), 1.49 (t, $J = 7.1$ Hz, 3H), 1.41 (t, $J = 7.2$ Hz, 3H), 0.82 - 0.87 (m, 4H), 0.72 - 0.78 (m, 4H).

5

Example 75



4-(1-(2-cyanoethyl)-5-methyl-1H-pyrazol-3-ylamino)-N,N-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

10


[00371] The title compound was prepared using a method analogous to that used to prepare example 74.

[00372] MS (ESI) m/z 472.2 ($M+H$).

[00373] ^1H NMR (400 MHz, CDCl_3) δ : 8.05 (s, 1H), 7.64 (s, 1H), 6.99 (s, 1H), 6.85 (s, 1H), 4.65 (q, $J = 7.0$ Hz, 2H), 4.25 (t, $J = 6.8$ Hz, 2H), 4.01 (s, 3H), 2.92 (t, $J = 6.7$ Hz, 2H), 2.79 - 2.86 (m, 2H), 2.41 (s, 3H), 1.49 (t, $J = 7.2$ Hz, 3H), 0.82 - 0.88 (m, 4H), 0.72 - 0.80 (m, 4H).

20

Example 76



N,N-dicyclopropyl-6-ethyl-4-(1-(2-methoxyethyl)-5-methyl-1H-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide


[00374] The title compound was prepared using a method analogous to that used to prepare example 74.

[00375] MS (ESI) m/z 477.2 (M+H).

5 [00376] ^1H NMR (500 MHz, CDCl_3) δ : 8.09 (s, 1H), 7.61 (s, 1H), 6.94 (s, 1H), 6.84 (s, 1H), 4.66 (q, J = 6.8 Hz, 2H), 4.12 (t, J = 5.5 Hz, 2H), 3.98 (s, 3H), 3.74 (t, J = 5.8 Hz, 2H), 3.32 (s, 3H), 2.78 - 2.86 (m, 2H), 2.36 (s, 3H), 1.49 (t, J = 7.1 Hz, 3H), 0.81 - 0.89 (m, 4H), 0.71 - 0.80 (m, 4H).

10


Example 77



N,N-dicyclopropyl-6-ethyl-4-(1-(2-isopropoxyethyl)-5-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

15

77A Preparation of 2-isopropoxyethyl methanesulfonate



[00377] A round bottom flask was charged with 2-isopropoxyethanol (0.5 mL, 4.34 mmol) and dichloromethane (21.68 mL) and was cooled to 0 °C. Triethylamine (1.208 mL, 8.67 mmol) was added along with a tip of DMAP. Methanesulfonyl chloride (0.338 mL, 4.34 mmol) was added dropwise. The reaction mixture was slowly warmed to rt over 2 h. The reaction mixture was quenched with saturated aqueous ammonium chloride and transferred to a separatory funnel. The aqueous layer was extracted with dichloromethane (2 x). The combined organics were washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*.

20


25

2-Isopropoxyethyl methanesulfonate (1 g) was isolated as a clear oil and was immediately used in the next reaction.

[00378] ^1H NMR (400 MHz, CHLOROFORM- d) δ ppm 4.34 - 4.40 (m, 2H), 3.67 - 3.71 (m, 2H), 3.60 - 3.67 (m, 1H), 3.07 (s, 3H), 1.18 (d, 6H, J = 8.0 Hz)

5

77B Preparation of *tert*-butyl 1,3-dioxoisindolin-2-yl(2-isopropoxyethyl)carbamate




[00379] A round bottom flask charged with 2-isopropoxyethyl methanesulfonate (example **77A**, 0.791 g, 4.34 mmol) in acetonitrile (21.70 mL) was stirred under an atmosphere of nitrogen. *tert*-Butyl 1,3-dioxoisindolin-2-ylcarbamate (1.138 g, 4.34 mmol) was added along with potassium carbonate (2.399 g, 17.36 mmol) and benzyltriethylammonium chloride (0.198 g, 0.868 mmol). The reaction mixture was heated at 50 °C 24 h and cooled to rt over the weekend. The reaction mixture was diluted with ethyl acetate and water. The aqueous layer was extracted with ethyl acetate. The combined organics were washed with brine, dried over anhydrous sodium sulfate, vacuum filtered through a plug of silica gel and concentrated *in vacuo*. The crude oil was used without additional purification (1.26 g, 83% yield).

[00380] MS (ESI) m/z 249.2 ($M - C_4H_9CO_2$)

[00381] ^1H NMR (400 MHz, CHLOROFORM- d) as a mixture of rotomers δ ppm 7.86 - 7.94 (m, 2 H), 7.73 - 7.82 (m, 2 H), 3.79 - 3.90 (m, 2 H), 3.58 - 3.66 (m, 2 H), 3.41 - 3.55 (m, 1 H), 1.31 - 1.56 (m, 9 H), 0.87 - 1.05 (m, 6 H)

77C Preparation of *tert*-butyl 1-(2-isopropoxyethyl)hydrazinecarboxylate




- [00382] To a solution of *tert*-butyl 1,3-dioxoisindolin-2-yl(2-isopropoxyethyl)carbamate (example **77B**, 1.262 g, 3.62 mmol) in tetrahydrofuran (24.15 mL) at 0 °C was slowly added methylhydrazine (0.289 mL, 5.43 mmol). The reaction mixture was stirred at 0 °C for 3 h. The reaction mixture was filtered to
5 remove the white precipitate and the filtrate was concentrated *in vacuo*. The semisolid was triturated with ethyl acetate (2 x) and the triturant was concentrated *in vacuo*. This process was repeated to ultimately provide *tert*-butyl 1-(2-isopropoxyethyl)hydrazinecarboxylate (0.665 g, 84% yield) as a yellow oil.
- [00383] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 3.52 - 3.65 (m, 5 H), 1.48
10 (s, 9 H), 1.15 (d, 6 H, *J* = 6.27 Hz)

77 Preparation of *N,N*-dicyclopropyl-6-ethyl-4-(1-(2-isopropoxyethyl)-5-methyl-1H-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide


- 15 [00384] This compound was prepared according to example **73** using *tert*-butyl 1-(2-isopropoxyethyl)hydrazinecarboxylate (example **77C**, 29.8 mg, 0.137 mmol) to provide *N,N*-dicyclopropyl-6-ethyl-4-(1-(2-isopropoxyethyl)-5-methyl-1H-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (16.72 mg, 48.7% yield) as a white solid.
- 20 [00385] MS (ESI) *m/z* 505.4 (M+H)
- [00386] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 7.89 (s, 1 H), 6.99 (s, 1 H), 6.88 (s, 1H), 4.65 (q, 2 H, *J* = 7.03 Hz), 4.22 (t, 2H, *J* = 5.27 Hz), 4.06 (s, 3 H), 3.78 (t, 2H, *J* = 5.27 Hz), 3.52 (dt, 1H, *J* = 12.11, 6.12 Hz), 2.79 - 2.88 (m, 2 H), 2.41 (s, 3H), 1.50 (t, 3 H, *J* = 7.03 Hz), 1.09 (d, 6H, *J* = 6.27 Hz), 0.82 - 0.91 (m, 4 H),
25 0.71 - 0.81 (m, 4 H)

Example 78



***N,N*-dicyclopropyl-6-ethyl-4-(1-(2-(2-methoxyethoxy)ethyl)-5-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

- 5 **78A** Preparation of *tert*-butyl 1,3-dioxoisindolin-2-yl(2-(2-methoxyethoxy)ethyl) carbamate




- [00387] The compound was prepared according to Example **77B** using 1-bromo-2-(2-methoxyethoxy)ethane (0.259 mL, 1.906 mmol) to provide *tert*-butyl 1,3-dioxoisindolin-2-yl(2-(2-methoxyethoxy)ethyl)carbamate (0.583 g, 84 % yield) as a pale yellow oil.

[00388] MS (ESI) *m/z* 265.2 (M-C₄H₉CO₂)

- [00389] ¹H NMR (400 MHz, CHLOROFORM-*d*) as a mixture of rotomers δ ppm
7.85 - 7.94 (m, 2 H), 7.73 - 7.82 (m, 2 H), 3.83 - 3.93 (m, 2 H), 3.67 - 3.74 (m, 2 H),
15 3.48 - 3.59 (m, 2 H), 3.30 - 3.43 (m, 2 H), 3.16 - 3.28 (m, 3 H), 1.29 - 1.55 (m, 9 H)

78B Preparation of *tert*-butyl 1-(2-(2-ethoxyethoxy)ethyl)hydrazinecarboxylate



- [00390] The compound was prepared according to example **77C** using *tert*-butyl
20 1,3-dioxoisindolin-2-yl(2-(2-methoxyethoxy)ethyl)carbamate (0.583 g, 1.600 mmol)

to provide *tert*-butyl 1-(2-(2-methoxyethoxy)ethyl)hydrazinecarboxylate (0.4 g, 107% yield) as a yellow oil.

[00391] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 3.51 - 3.70 (m, 8 H), 3.39 (s, 3 H), 1.48 (s, 9 H)

5

78 Preparation of *N,N*-dicyclopropyl-6-ethyl-4-(1-(2-(2-methoxyethoxy)ethyl)-5-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide


[00392] This compound was prepared according to example 74 using *tert*-butyl 1-(2-(2-methoxyethoxy)ethyl)hydrazinecarboxylate (example 78B, 43.2 mg, 0.184 mmol) to provide *N,N*-dicyclopropyl-6-ethyl-4-(1-(2-(2-methoxyethoxy)ethyl)-5-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (17.34 mg, 36% yield) as a white solid.

[00393] MS (ESI) *m/z* 521.4 (M+H)

[00394] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 7.85 (br s, 1 H), 6.91 (s, 1 H), 6.87 (s, 1H), 4.65 (q, 2 H, *J* = 7.03 Hz), 4.20 (t, 2H, *J* = 5.27 Hz), 4.05 (s, 3 H), 3.87 (t, 2H, *J* = 5.65 Hz), 3.53 - 3.59 (m, 2H), 3.50 (ddd, 2H, *J* = 5.40, 2.89, 2.76 Hz), 2.77 - 2.87 (m, 2 H), 2.38 (s, 3H), 1.50 (t, 3 H, *J* = 7.03 Hz), 0.82 - 0.91 (m, 4 H), 0.71 - 0.79 (m, 4 H)

20


Example 79



Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-methyl-1-propyl-1*H*-pyrazol-3-ylamino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

25

79A Preparation of *tert*-butyl 1-propylhydrazinecarboxylate



[00395] To a solution of propylhydrazine, oxalic acid salt (0.5 g, 3.05 mmol) in ethanol (3 mL) in a round bottom flask under nitrogen was added triethylamine (0.425 mL, 3.05 mmol). The reaction mixture was cooled to 0 °C and BOC₂O (0.707 mL, 3.05 mmol) dissolved in ethanol (1 mL) was added dropwise over 10 min. The reaction mixture was allowed to warm to room temperature for 16h. Ethanol was removed by concentration *in vacuo*. The residue was taken up in water and ethyl acetate. The aqueous layer was extracted with ethyl acetate (3x). The combined organics were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The crude product was used as such in the next reaction.

[00396] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 3.30 - 3.36 (m, 2 H), 1.59 (dq, 3 H, *J*=14.40, 7.33 Hz), 1.48 (s, 9 H), 0.88 (t, 3 H, *J*=7.40 Hz)


79 Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-methyl-1-propyl-1*H*-pyrazol-3-ylamino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

[00397] This compound was prepared according to example 74 using *tert*-butyl 1-propylhydrazinecarboxylate (example 79A, 19.87 mg, 0.114 mmol) to provide *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-methyl-1-propyl-1*H*-pyrazol-3-ylamino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (15.6 mg, 59% yield) as a white solid.

[00398] MS (ESI) *m/z* 461.4 (*M*+*H*)


[00399] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 7.74 (br s, 1 H), 6.92 (s, 1 H), 6.86 (s, 1 H), 4.66 (q, 2 H, *J* = 7.11 Hz), 4.03 (s, 3 H), 3.95 (t, 2 H, *J* = 7.15 Hz), 2.78 - 2.87 (m, 2 H), 2.34 (s, 3 H), 1.87 (dq, 3 H, *J* = 14.49, 7.30 Hz), 1.50 (t, 3 H, *J* = 7.15 Hz), 0.96 (t, 3 H, *J* = 7.40 Hz), 0.83 - 0.89 (m, 4 H), 0.72 - 0.79 (m, 4 H)

Example 80



N,N-dicyclopropyl-4-(1-(2,2-difluoroethyl)-5-methyl-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide


- 5 **80A** Preparation of *tert*-butyl 2,2-difluoroethyl(1,3-dioxoisindolin-2-yl)carbamate



[00400] The title compound was prepared using a method analogous to that used to prepare example 77A

10

- 80B** Preparation of *tert*-butyl 1-(2,2-difluoroethyl)hydrazinecarboxylate



[00401] The title compound was prepared using a method analogous to that used to prepare example 77B.

- 15 [00402] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ: 5.94 (tt, *J* = 56.3, 4.4 Hz, 1H), 3.74 (td, *J* = 13.8, 4.4 Hz, 2H), 1.48 (s, 9H)

- 20 **80** Preparation of *N,N*-dicyclopropyl-4-(1-(2,2-difluoroethyl)-5-methyl-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide


[00403] The title compound was prepared using a method analogous to that used to prepare example 73.

[00404] MS (ESI) *m/z* 483.3 (*M*+*H*).

[00405] ^1H NMR 500 MHz, (CHLOROFORM- d) δ : 8.17 (br. s., 1H), 7.66 (s, 1H), 7.01 (s, 1H), 6.85 (s, 1H), 6.05 (tt, J = 56.0, 4.4 Hz, 1H), 4.65 (q, J = 6.9 Hz, 2H), 4.30 (td, J = 13.2, 4.3 Hz, 2H), 3.99 (s, 3H), 2.79 - 2.86 (m, 2H), 2.36 (s, 3H), 1.49 (t, J = 7.1 Hz, 3H), 0.81 - 0.88 (m, 4H), 0.72 - 0.79 (m, 4H).


5

Example 81



10 **4-(1-cyclobutyl-5-methyl-1H-pyrazol-3-ylamino)-N,N-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

81A Preparation of *tert*-butyl 1-cyclobutylhydrazinecarboxylate



15 [00406] The title compound was prepared following the procedure described by So Ok Park et al. in *J. Comb. Chem.* **11**, 315-326, **2009**.

81 Preparation of 4-(1-cyclobutyl-5-methyl-1H-pyrazol-3-ylamino)-N,N-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide


20 [00407] The title compound was prepared using a method analogous to that used to prepare example 73.

[00408] MS (ESI) m/z 473.4(M+H).

[00409] ^1H NMR (500 MHz, CHLOROFORM- d) δ : 8.05 (s, 1H), 6.91 (s, 1H), 6.79 (s, 1H), 4.68 (quin, J = 8.4 Hz, 1H), 4.61 (q, J = 7.1 Hz, 2H), 4.07 (s, 3H), 2.80 - 2.87 (m, 2H), 2.69 - 2.79 (m, 2H), 2.37 - 2.47 (m, 2H), 2.32 (s, 3H), 1.78 - 1.98 (m, 2H), 1.47 (t, J = 7.1 Hz, 3H), 0.82 - 0.91 (m, 4H), 0.72 - 0.80 (m, 4H).

25

Example 82



***N,N*-dicyclopropyl-6-ethyl-4-(1-(1-methoxypropan-2-yl)-5-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

5


[00410] Prepared as described in example 77.

[00411] MS (ESI) *m/z* 491 (M+H).

[00412] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 8.18 (br. s., 1 H) 7.63 (s, 1 H) 6.93 (s, 1 H) 6.85 (s, 1 H) 4.66 (q, *J*=7.04 Hz, 2 H) 4.29 - 4.48 (m, *J*=13.31, 6.82, 6.82, 6.71 Hz, 1 H) 4.00 (s, 3 H) 3.55 - 3.79 (m, 2 H) 3.29 (s, 3 H) 2.72 - 2.92 (m, 2 H) 2.36 (s, 3 H) 1.49 (t, *J*=7.04 Hz, 3 H) 1.46 (d, *J*=6.82 Hz, 2 H) 0.68 - 0.93 (m, 8 H).

15

Example 83



***N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-methyl-1-(tetrahydro-2*H*-pyran-4-yl)-1*H*-pyrazol-3-ylamino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**


20

[00413] Prepared using similar procedure as for example 77.

[00414] MS (ESI) m/z 503 (M+H).

[00415] ^1H NMR (500 MHz, CHLOROFORM- d) δ ppm 8.12 (s, 1 H) 7.64 (s, 1 H) 6.95 (s, 1 H) 6.86 (s, 1 H) 4.66 (q, $J=7.21$ Hz, 2 H) 4.09 - 4.23 (m, 3 H) 4.01 (s, 3 H) 3.55 (td, $J=12.00$, 1.80 Hz, 2 H) 2.73 - 2.89 (m, 2 H) 2.37 (s, 3 H) 2.25 - 2.36 (m, 2 H) 1.84 (dd, $J=12.76$, 2.22 Hz, 2 H) 1.50 (t, $J=7.07$ Hz, 3 H) 0.71 - 0.91 (m, 8 H).

Example 84




***N,N*-dicyclopropyl-4-(1-(2-(dimethylamino)ethyl)-5-methyl-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

[00416] Prepared using similar procedure as for example 77.

[00417] MS (ESI) m/z 490 (M+H).

[00418] ^1H NMR (500 MHz, CHLOROFORM- d) δ ppm 8.08 (s, 1 H) 7.64 (s, 1 H) 6.96 (s, 1 H) 6.86 (s, 1 H) 4.65 (q, $J=7.21$ Hz, 2 H) 4.24 (br. s., 2 H) 4.01 (s, 3 H) 3.01 - 3.22 (m, 2 H) 2.70 - 2.89 (m, 2 H) 2.50 (br. s., 6 H) 2.37 (s, 3 H) 1.49 (t, $J=7.07$ Hz, 3 H) 0.71 - 0.91 (m, 8 H).

Example 85



***N,N*-dicyclopropyl-6-ethyl-4-(1-(2-hydroxyethyl)-5-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**


5 [00419] Prepared using similar procedure as for example 77.

[00420] MS (ESI) *m/z* 463 (M+H).

[00421] ¹H NMR (500 MHz, MeOD) δ ppm 7.93 (s, 1 H) 7.16 (s, 1 H) 6.88 (s, 1 H) 4.60 (q, *J*=7.21 Hz, 2 H) 4.09 (t, *J*=5.55 Hz, 2 H) 4.05 (s, 3 H) 3.88 (t, *J*=5.55 Hz, 2 H) 2.88 - 3.00 (m, 2 H) 2.37 (s, 3 H) 1.43 (t, *J*=7.07 Hz, 3 H) 0.62 - 0.98 (m, 8 H).

10

Example 86



***N,N*-dicyclopropyl-4-(1-(2-(dimethylamino)-2-oxoethyl)-5-methyl-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

15


[00422] Prepared using similar procedure as for example 77.

[00423] MS (ESI) *m/z* 504 (M+H).

[00424] ¹H NMR (500 MHz, MeOD) δ ppm 7.93 (s, 1 H) 7.16 (s, 1 H) 6.99 (s, 1 H) 4.97 (s, 2 H) 4.61 (q, *J*=6.94 Hz, 2 H) 4.05 (s, 3 H) 3.15 (s, 3 H) 2.98 (s, 3 H) 2.79 - 2.97 (m, 2 H) 2.29 (s, 3 H) 1.44 (t, *J*=7.07 Hz, 3 H) 0.72 - 0.91 (m, 8 H).

20

Example 87



25

4-(1-(cyanomethyl)-5-methyl-1*H*-pyrazol-3-ylamino)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide


[00425] Prepared using similar procedure as for example 77.

5 [00426] MS (ESI) *m/z* 458 (M+H).

[00427] ¹H NMR (400 MHz, CHLOROFORM-*d*) TM ppm 8.17 (s, 1 H) 7.67 (s, 1 H) 7.10 (s, 1 H) 6.87 (s, 1 H) 4.93 (s, 2 H) 4.65 (q, *J*=7.04 Hz, 2 H) 4.01 (s, 3 H) 2.76 - 2.92 (m, 2 H) 2.43 (s, 3 H) 1.50 (t, *J*=7.04 Hz, 3 H) 0.68 - 0.93 (m, 8 H).

10

Example 88



***N,N*-dicyclopropyl-6-ethyl-4-(1-(3-methoxypropyl)-5-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

15


[00428] Prepared using similar procedure as for example 77.

[00429] MS (ESI) *m/z* 491 (M+H).

[00430] ¹H NMR (400 MHz, CHLOROFORM-*d*) TM ppm 8.11 (s, 1 H) 7.62 (s, 1 H) 6.92 (s, 1 H) 6.78 - 6.89 (m, 1 H) 4.66 (q, *J*=7.04 Hz, 2 H) 4.06 (t, *J*=6.60 Hz, 2 H) 3.98 (s, 3 H) 3.29 - 3.39 (m, 4 H) 2.75 - 2.91 (m, 2 H) 2.34 (s, 3 H) 2.08 (dq, *J*=6.38, 6.16 Hz, 2 H) 1.50 (t, *J*=7.04 Hz, 3 H) 0.70 - 0.94 (m, 8 H).

20

Example 89



***N,N*-dicyclopropyl-6-ethyl-4-(1-isopropyl-5-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**


[00431] Prepared using similar procedure as for example 77.

5 [00432] MS (ESI) *m/z* 461 (M+H).

[00433] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 8.15 (s, 1 H) 7.62 (s, 1 H) 6.91 (s, 1 H) 6.85 (s, 1 H) 4.66 (q, *J*=7.19 Hz, 2 H) 4.25 - 4.48 (m, 1 H) 3.99 (s, 3 H) 2.66 - 2.90 (m, *J*=7.13, 7.13, 3.85, 3.47 Hz, 2 H) 2.34 (s, 3 H) 1.38 - 1.58 (m, 9 H) 0.68 - 0.91 (m, 8 H).

10

Example 90



***N,N*-dicyclopropyl-6-ethyl-4-(1-isobutyl-5-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

15


[00434] Prepared using similar procedure as for example 77.

[00435] MS (ESI) *m/z* 475 (M+H).

[00436] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 8.11 (s, 1 H) 7.62 (s, 1 H) 6.93 (s, 1 H) 6.85 (s, 1 H) 4.66 (q, *J*=7.04 Hz, 2 H) 3.99 (s, 3 H) 3.74 (d, *J*=7.48 Hz, 2 H) 2.74 - 2.89 (m, 2 H) 2.33 (s, 3 H) 2.18 - 2.29 (m, 1 H) 1.50 (t, *J*=7.04 Hz, 3 H) 0.94 (d, *J*=6.60 Hz, 6 H) 0.72 - 0.88 (m, 8 H).

20

Example 91




25 **Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-methyl-1*H*-pyrazol-3-ylamino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

[00437] Prepared using similar protocol as for example 72 from hydrazine.

[00438] MS (ESI) m/z 419.3 (M+H)

[00439] ^1H NMR (400 MHz, CHLOROFORM- d) δ ppm 8.70 (br s, 1 H), 7.91 (br s, 1 H), 6.87 (s, 1 H), 6.09 (br s, 1 H), 4.64 (q, 2 H, $J = 7.03$ Hz), 4.08 (s, 3 H), 2.74 - 2.95 (m, 2 H), 2.41 (s, 3 H), 1.51 (t, 3 H, $J = 7.15$ Hz), 0.81 - 0.95 (m, 4 H), 0.70 - 0.81 (m, 4 H)

Example 92




N,N-dicyclopropyl-6-ethyl-1-methyl-4-((5-methyl-1-(2-(methylsulfonyl)ethyl)-1*H*-pyrazol-3-yl)amino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

[00440] Prepared using the same procedure as used for example 77.

[00441] MS (ESI) m/z 525.3 (M+H)

[00442] ^1H NMR (CDCl₃) δ ppm 8.25 (br s, 1H), 7.75 (s, 1H), 6.86 (s, 1H), 6.77 (s, 1H), 4.54 (q, 2H, $J = 8$ Hz), 4.36 (t, 2H, $J = 8$ Hz), 3.97 (s, 3H), 3.58 (t, 2H, $J = 8$ Hz), 2.71-2.77 (m, 2H), 2.44 (s, 3H), 2.34 (s, 3H), 1.41 (t, 3H, $J = 8$ Hz), 0.75-0.78 (m, 4H), 0.67-0.71 (m, 4H).

Example 93




***N,N*-dicyclopropyl-6-ethyl-4-((1-(*cis*-4-hydroxycyclohexyl)-5-methyl-1*H*-pyrazol-3-yl)amino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

5 [00443] Prepared using the same procedure as used for example 77.

[00444] MS (ESI) *m/z* 517.4 (M+H)

[00445] ¹H NMR (CDCl₃) δ ppm 8.14 (br s, 1H), 7.56 (s, 1H), 6.83 (s, 1H), 6.76 (s, 1H), 4.59 (q, 2H, *J* = 8 Hz), 4.01-4.02 (m, 1H), 3.93 (s, 3H), 3.88-3.93 (m, 1H), 2.73-2.77 (m, 2H), 2.28 (s, 3H), 2.25-2.33 (m, 2H), 1.93-1.96 (m, 2H), 1.59-1.73 (m, 10 4H), 1.42 (t, 3H, *J* = 8 Hz), 0.73-0.78 (m, 4H), 0.66-0.70 (m, 4H).

Example 94



15 ***N,N*-dicyclopropyl-6-ethyl-1-methyl-4-((5-methyl-1-(methylsulfonyl)-1*H*-pyrazol-3-yl)amino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**


[00446] Prepared using the same procedure as used for example 77.

[00447] MS (ESI) *m/z* 497.3 (M+H)

20 [00448] ¹H NMR (CDCl₃) δ ppm 8.46 (br s, 1H), 7.75 (s, 1H), 7.16 (s, 1H), 6.78 (s, 1H), 4.54 (q, 2H, *J* = 8 Hz), 3.97 (s, 3H), 3.19 (s, 3H), 2.72-2.76 (m, 2H), 2.51 (s, 3H), 1.41 (t, 3H, *J* = 8 Hz), 0.72-0.78 (m, 4H), 0.66-0.69 (m, 4H).


Example 95

25



***N,N*-dicyclopropyl-6-ethyl-4-(5-isopropyl-1-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**


5 **95A** Preparation of 4-methyl-1,1-bis(methylthio)pent-1-en-3-one



[00449] The title compound was prepared using a method analogous to that used to prepare Example **70A**.

[00450] ¹H NMR (400 MHz, CDCl₃) δ: 6.08 (s, 1H), 2.47 (s, 3H), 2.46 (s, 3H),
10 1.14 (s, 3H), 1.12 (s, 3H).

95B Preparation of (*Z*)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(4-methyl-1-(methylthio)-3-oxopent-1-enylamino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide



15 [00451] The title compound was prepared using a method analogous to that used to prepare example **70B**.

[00452] MS (ESI) *m/z* 481.4 (M+H).


20 **95** Preparation of *N,N*-dicyclopropyl-6-ethyl-4-(5-isopropyl-1-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

[00453] The title compound was prepared using a method analogous to that used to prepare example **73**.

[00454] MS (ESI) m/z 461.4 (M+H).


[00455] ^1H NMR (500 MHz, CDCl_3) δ : 7.80 (s, 1H), 7.07 (s, 1H), 6.86 (s, 1H), 4.60 (q, $J = 6.9$ Hz, 2H), 4.02 (s, 3H), 3.79 (s, 3H), 2.93 – 3.01 (m, 1H), 2.79 - 2.86 (m, 2H), 1.52 (t, $J = 7.1$ Hz, 3H), 1.35 (d, $J = 6.9$ Hz, 6H), 0.82 - 0.89 (m, 4H), 0.72 - 0.78 (m, 4H).

Example 96



10 *N,N*-dicyclopropyl-4-(5-cyclopropyl-1-ethyl-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

96A Preparation of 1-cyclopropyl-3,3-bis(methylthio)prop-2-en-1-one




15

[00456] The title compound was prepared using a method analogous to that used to prepare example 72A.

[00457] ^1H NMR (400 MHz, CDCl_3) δ : 6.23 (s, 1H), 2.50 (s, 3H), 2.48 (s, 3H), 1.86 - 1.94 (m, 1H), 1.09 (quin, $J = 3.7$ Hz, 2H), 0.82 - 0.88 (m, 2H).

20

96B Preparation of (*Z*)-*N,N*-dicyclopropyl-4-(3-cyclopropyl-1-(methylthio)-3-oxoprop-1-enylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide



[00458] The title compound was prepared using a method analogous to that used to prepare example 72B.

[00459] MS (ESI) m/z 479.3 (M+H).


5 [00460] ^1H NMR (400 MHz, CDCl_3) δ : 14.81 (s, 1H), 7.75 (s, 1H), 6.85 (s, 1H), 5.58 (s, 1H), 4.62 (q, $J = 7.0$ Hz, 2H), 4.00 (s, 3H), 2.75 - 2.85 (m, 2H), 2.45 (s, 3H), 1.76 - 1.86 (m, 1H), 1.46 (t, $J = 7.0$ Hz, 3H), 1.17 - 1.28 (m, 2H), 0.69 - 0.91 (m, 10H).

10 **96** Preparation of *N,N*-dicyclopentyl-4-(5-cyclopropyl-1-ethyl-1H-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00461] The title compound was prepared using a method analogous to that used to prepare example 73.


15 [00462] MS (ESI) m/z 473.3 (M+H).

Example 97



20 ***N,N*-dicyclopentyl-6-ethyl-1-methyl-4-(1-methyl-1H-pyrazol-3-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

97A Preparation of 4-bromo-*N,N*-dicyclopentyl-6-ethyl-1-methyl-1,6-dihydroimidazo [4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide



[00463] A round bottom flask charged with 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (134.4 mg, 0.397 mmol) under an atmosphere of nitrogen was treated with dibromomethane (397
5 μ L). Isoamyl nitrate (107 μ L, 0.794 mmol) was added and the reaction mixture was heated at 60 °C 3.5 h and cooled to rt. The reaction mixture was diluted with ethyl acetate and washed with saturated aqueous sodium bicarbonate. The organics were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. 4-Bromo-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-
10 b]pyridine-7-carboxamide (0.156 g, 0.388 mmol, 98% yield) was isolated as a brown glassy solid. The product was used immediately in the next reaction.

[00464] MS (ESI) m/z 404.2 (M+H)

97 Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(1-methyl-1*H*-pyrazol-3-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide
15


[00465] To a vial containing 1-methyl-1*H*-pyrazol-3-amine (11.89 mg, 0.122 mmol), Pd₂(dba)₃ (4.67 mg, 5.10 μ mol), Xantphos (5.90 mg, 10.20 μ mol), and cesium carbonate (100 mg, 0.306 mmol) was added 4-bromo-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example
20 **97A**, 41.0 mg, 0.102 mmol) in DME (1020 μ L). The reaction mixture was sparged with argon for 5 min, the vial was capped, and the reaction mixture was heated at 60 °C for 4 h and 80 °C for 16h. The reaction mixture was concentrated *in vacuo*. The crude residue was purified by flash column chromatography using an Isco 40 g column eluting with 0 – 5% MeOH/ CH₂Cl₂ followed by additional purification with
25 preparative HPLC. *N,N*-Dicyclopropyl-6-ethyl-1-methyl-4-(1-methyl-1*H*-pyrazol-3-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (13 mg, 0.030 mmol, 29.5% yield) was isolated as a white solid.

[00466] MS (ESI) m/z 419.3 (M+H)

[00467] ^1H NMR (400 MHz, CHLOROFORM- d) δ ppm 8.17 (s, 1 H), 7.64 (s, 1 H), 7.31 (d, 1H, $J = 2.26$ Hz), 7.16 (d, 1 H, $J = 2.26$ Hz), 6.86 (s, 1 H), 4.65 (q, 2 H, $J = 7.03$ Hz), 4.01 (s, 3 H), 3.85 (s, 3H), 2.78 - 2.87 (m, 2 H), 1.49 (t, 3 H, $J = 7.15$ Hz), 0.81 - 0.89 (m, 4 H), 0.71 - 0.80 (m, 4 H)

5

Example 98




10 ***N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(1-methyl-1*H*-pyrazol-4-ylamino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

[00468] Prepared using similar procedure as used for example 97.

[00469] ^1H NMR (500 MHz, CHLOROFORM- d) δ : 8.15 (s, 1H), 7.72 (s, 1H), 7.67 (s, 1H), 7.66 (br. s., 1H), 6.85 (s, 1H), 4.65 (q, $J = 7.2$ Hz, 2H), 4.01 (s, 3H), 3.95 (s, 3H), 2.79 - 2.86 (m, 2H), 1.50 (t, $J = 7.1$ Hz, 3H), 0.81 - 0.89 (m, 4H), 0.73 - 0.79 (m, 4H).

15

Example 99




20 ***N,N*-dicyclopropyl-6-ethyl-4-((5-(2-furyl)-1-methyl-1*H*-pyrazol-3-yl)amino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

[00470] Prepared in a similar method used for example 73.

[00471] MS (ESI) m/z 485.4 ($M+H$)

[00472] ¹H NMR (CDCl₃) δ ppm 8.03 (br s, 1H), 7.58 (s, 1H), 7.48 (m, 1H), 7.35 (s, 1H), 6.79 (s, 1H), 6.54 (d, 1H, *J* = 3.3 Hz), 6.47 (m, 1H), 4.58 (q, 2H, *J* = 7.04 Hz), 3.94 (s, 3H), 3.93 (s, 3H), 2.74 (m, 2H), 1.44 (t, 3H), 0.69-0.79 (m, 8H)

5

Example 100

N,N-dicyclopropyl-6-ethyl-1-methyl-4-((1-methyl-5-(trifluoromethyl)-1*H*-pyrazol-3-yl)amino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide


10

[00473] Prepared in a similar method used for example 73.

[00474] MS (ESI) *m/z* 487.3 (*M*+*H*)

[00475] ¹H NMR (CDCl₃) δ ppm 7.41 (s, 1H), 7.12 (s, 1H), 6.78 (s, 1H), 4.52 (q, 2H, *J* = 7.04 Hz), 4.11 (s, 3H), 3.86 (s, 3H), 2.74 (m, 2H), 1.42 (t, 3H, *J* = 7.15 Hz), 0.67-0.80 (m, 8H)


15

Example 101

3-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridin-4-ylamino)-1-ethyl-1*H*-pyrazole-5-carboxylic acid


20

101A Preparation of (*Z*)-*N,N*-dicyclopropyl-4-(4,4-dimethoxy-1-(methylthio)-3-oxobut-1-enylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide



[00476] Sodium hydride (142 mg, 3.55 mmol) was added to 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **1J**, 400 mg, 1.182 mmol) in DMF (7 mL) at room temperature and the mixture stirred for 30 min, 1,1-dimethoxy-4,4-bis(methylthio)but-3-en-2-one (394 mg, 1.773 mmol) was added and stirring continued overnight. The reaction mixture was diluted with ethyl acetate and washed with water, 10% lithium chloride and water. The organic layer was dried over magnesium sulfate, filtered and concentrated *in vacuo*. The crude product was purified by flash chromatography on silica gel using an automated ISCO system (80 g column, eluting with 2-5% methanol / dichloromethane). (*Z*)-*N,N*-Dicyclopropyl-4-(4,4-dimethoxy-1-(methylthio)-3-oxobut-1-enylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (484 mg, 0.944 mmol, 80 % yield) was obtained as a yellow solid.

101B Preparation of (*Z*)-*tert*-butyl 2-(1-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4,4-dimethoxy-3-oxobut-1-enyl)-1-ethylhydrazinecarboxylate




[00477] A mixture of (*Z*)-*N,N*-dicyclopropyl-4-(4,4-dimethoxy-1-(methylthio)-3-oxobut-1-enylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **101A**, 160 mg, 0.312 mmol) and *tert*-butyl 1-ethylhydrazinecarboxylate (125 mg, 0.780 mmol) in acetic acid (2 mL) was stirred at 35 °C overnight. Solvent was evaporated and the crude product was purified by flash chromatography on silica gel using an automated ISCO system (40 g column, eluting

with 2-8% methanol/dichloromethane). (*Z*)-*tert*-Butyl 2-(1-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4,4-dimethoxy-3-oxobut-1-enyl)-1-ethylhydrazinecarboxylate (195 mg, 0.312 mmol, 100 % yield) was obtained as a yellow solid.

5 [00478] MS (ESI) m/z 625.5 (M+H).

101C Preparation of *N,N*-dicyclopropyl-6-ethyl-4-(1-ethyl-5-formyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide



10

[00479] A mixture of (*Z*)-*tert*-butyl 2-(1-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-4,4-dimethoxy-3-oxobut-1-enyl)-1-ethylhydrazinecarboxylate (187 mg, 0.299 mmol) and TFA (25% in dichloroethane, 1.0 mL, 2.99 mmol) was heated at 60 °C for 0.5h and LC/MS

15 showed completion of reaction. Solvent was evaporated and the residue was partitioned between saturated sodium bicarbonate and dichloromethane. The layers were separated and aqueous layer was extracted with dichloromethane two more times. The combined organic layer was dried over magnesium sulfate, filtered and concentrated *in vacuo*. The crude product was purified by flash chromatography on silica gel using an automated ISCO system (24 g column, eluting with 1-5% methanol / dichloromethane). *N,N*-dicyclopropyl-6-ethyl-4-(1-ethyl-5-formyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (80 mg, 0.174 mmol, 58.0 % yield) was obtained as a yellow solid.

20

[00480] MS (ESI) m/z 461.4 (M+H).

25 [00481] ¹H NMR (500 MHz, CDCl₃) δ: 9.88 (s, 1H), 8.25 (s, 1H), 7.76 (s, 1H), 7.67 (s, 1H), 6.87 (s, 1H), 4.67 (d, *J* = 7.2 Hz, 2H), 4.52 (q, *J* = 7.1 Hz, 2H), 4.02 (s, 3H), 2.80 - 2.86 (m, 2H), 1.52 (t, *J* = 7.1 Hz, 3H), 1.44 (t, *J* = 7.2 Hz, 3H), 0.83 - 0.89 (m, 4H), 0.73 - 0.79 (m, 4H)

101 Preparation of 3-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-1-ethyl-1*H*-pyrazole-5-carboxylic acid

[00482] A mixture of *N,N*-dicyclopropyl-6-ethyl-4-(1-ethyl-5-formyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **101C**, 27 mg, 0.059 mmol) and OXONE (79 mg, 0.129 mmol) in DMF was stirred at room temperature for 20 hrs. Water was added and the mixture was extracted with dichloromethane 3 times. The organic layer was dried over magnesium sulfate, filtered and concentrated *in vacuo*. The crude product was


10 purified with preparative HPLC to give 3-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-1-ethyl-1*H*-pyrazole-5-carboxylic acid (20 mg, 98% yield)

[00483] MS (ESI) *m/z* 477.4 (M+H).

[00484] ¹H NMR (500 MHz, METHANOL-*d*₄) δ : 7.96 (s, 1H), 7.48 (s, 1H), 7.18 (s, 1H), 4.70 (q, *J* = 6.9 Hz, 2H), 4.60 (q, *J* = 7.2 Hz, 2H), 4.09 (s, 3H), 2.94 – 3.00 (m, 2H), 1.45 (t, *J* = 6.94 Hz, 3H), 1.44 (t, *J* = 7.21 Hz, 3H), 0.85 - 0.92 (m, 4H), 0.77 - 0.82 (m, 4H)

15

Example 102



20

***N,N*-dicyclopropyl-6-ethyl-4-(1-ethyl-5-(hydroxymethyl)-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

[00485] Sodium boronhydride (10 wt% on alumina, 13.14 mg, 0.035 mmol) was added to *N,N*-dicyclopropyl-6-ethyl-4-(1-ethyl-5-formyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **101C**, 16 mg, 0.035 mmol) in methanol (1 mL) at room temperature and the reaction mixture stirred for 0.5h. The solid alumina was filtered off and the filtrate was concentrated *in vacuo*. The crude product was purified by flash chromatography on

25


silica gel using an automated ISCO system (4 g column, eluting with 2-10% methanol / dichloromethane). *N,N*-dicyclopropyl-6-ethyl-4-(1-ethyl-5-(hydroxymethyl)-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (14 mg, 86 % yield) was obtained as a white solid.

5 [00486] MS (ESI) *m/z* 563.4 (M+H).

[00487] ¹H NMR (500 MHz, CDCl₃) δ: 8.09 (s, 1H), 7.59 (s, 1H), 7.08 (s, 1H), 6.81 (s, 1H), 4.72 (d, *J* = 4.7 Hz, 2H), 4.57 (q, *J* = 7.2 Hz, 2H), 4.15 (q, *J* = 7.3 Hz, 2H), 3.97 (s, 3H), 2.76 – 2.84 (m, 3H), 1.46 (t, *J* = 7.2 Hz, 3H), 1.41 (t, *J* = 7.1 Hz, 3H), 0.81 - 0.87 (m, 4H), 0.71 - 0.76 (m, 4H)

10

Example 103



N,N-dicyclopropyl-4-(5-(dimethylcarbamoyl)-1-ethyl-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

15


[00488] A mixture of 3-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridin-4-ylamino)-1-ethyl-1*H*-pyrazole-5-carboxylic acid (example **101**, 28 mg, 0.059 mmol), HATU (26.8 mg, 0.071 mmol) and dimethylamine (2M in THF, 59 μL, 0.118 mmol) in DMF (1 mL) was stirred at room temperature for 2h. Water was added and the reaction mixture was extracted with ethyl acetate 3 times. The combined organic layers were washed twice with 10% lithium chloride, dried over magnesium sulfate, filtered and concentrated *in vacuo*. The crude was purified by preparative HPLC (Phenomenex Luna 5u C18 column, 21.2x250 mm column, retention time 15.317 min, 30-100% gradient aqueous methanol over 20 minutes containing 10 mM ammonium acetate, flow rate 20 ml/min and monitoring at 254 nm). *N,N*-dicyclopropyl-4-(5-(dimethylcarbamoyl)-1-ethyl-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (20 mg, 66.9 % yield) was obtained as an off-white solid.

[00489] MS (ESI) *m/z* 504.5 (M+H).

[00490] ^1H NMR (500 MHz, CDCl_3) δ : 8.38 (br. s, 1H), 7.75 (s, 1H), 7.28 (s, 1H), 6.86 (s, 1H), 4.61 (q, $J = 7.2$ Hz, 2H), 4.27 (q, $J = 7.0$ Hz, 5H), 4.03 (s, 3H), 3.27 (s, 3H), 3.17 (s, 3H), 2.79 - 2.86 (m, 2H), 1.47 (t, $J = 7.14$ Hz, 3H), 1.46 (t, $J = 7.14$ Hz, 3H), 0.83 - 0.89 (m, 4H), 0.73 - 0.78 (m, 4H)

5

Example 104



N,N-dicyclopropyl-6-ethyl-4-(1-ethyl-5-(morpholine-4-carbonyl)-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

10


[00491] The title compound was prepared using a method analogous to that used to prepare example 103.

[00492] MS (ESI) m/z 546.5 ($M+H$).

15 [00493] ^1H NMR (500 MHz, CDCl_3) δ : 7.95 (s, 1H), 7.19 (s, 1H), 6.87 (s, 1H), 4.60 (q, $J = 6.9$ Hz, 2H), 4.28 (q, $J = 7.2$ Hz, 5H), 4.07 (s, 3H), 3.82 (br. s., 5H), 3.76 (br. s., 3H), 2.74 - 2.93 (m, 4H), 1.48 (t, $J = 7.14$ Hz, 3H), 1.47 (t, $J = 7.14$ Hz, 3H), 0.84 - 0.90 (m, 4H), 0.74 - 0.78 (m, 4H).

20

Example 105




N,N-dicyclopropyl-6-ethyl-4-(1-ethyl-5-(thiomorpholine 1,1-dioxide-4-carbonyl)-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

[00494] The title compound was prepared using a method analogous to that used to prepare example 103.

[00495] MS (ESI) m/z 594.5 (M+H).

[00496] ¹H NMR (500 MHz, CDCl₃) δ: 7.95 (s, 1H), 7.19 (s, 1H), 6.83 (s, 1H),
 5 4.55 (q, J = 7.03 Hz, 2H), 4.20 - 4.34 (m, 6H), 4.06 (s, 3H), 3.13 (br. s., 4H), 2.77 -
 2.89 (m, 2H), 1.46 (t, J = 7.21 Hz, 3H), 1.43 (t, J = 7.21 Hz, 3H), 0.82 - 0.88 (m, 4H),
 0.72 - 0.77(m, 4H).

Example 106



10

N,N-dicyclopropyl-6-ethyl-4-(1-ethyl-5-(2-morpholinoethylcarbamoyl)-1*H*-
 pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-
 7-carboxamide


15 [00497] The title compound was prepared using a method analogous to that used to
 prepare example 103.

[00498] MS (ESI) m/z 589.6 (M+H).

[00499] ¹H NMR (500 MHz, CDCl₃) δ: 8.18 (s, 1H), 7.68 (s, 1H), 7.45 (s, 1H),
 6.87 (s, 1H), 4.64 (q, J = 6.9 Hz, 2H), 4.54 (q, J = 7.2 Hz, 2H), 4.03 (s, 3H), 3.77 (br.
 20 s., 4H), 3.61 (br. s., 2H), 2.80 - 2.87 (m, 2H), 2.67 (br. s., 2H), 2.57 (br. s., 3H), 1.71
 (br. s., 3H), 1.53 (t, J = 7.1 Hz, 3H), 1.46 (t, J = 7.1 Hz, 3H), 0.84 - 0.90 (m, 4H), 0.74
 - 0.79 (m, 4H).

Example 107

25



***N,N*-dicyclopropyl-6-ethyl-4-(1-ethyl-5-(4-methylpiperazine-1-carbonyl)-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

5


[00500] The title compound was prepared using a method analogous to that used to prepare example **103**.

[00501] MS (ESI) *m/z* 559.5 (*M*+*H*).

[00502] ¹H NMR (500 MHz, CDCl₃) δ: 8.19 (s, 1H), 7.66 (s, 1H), 7.26 (s, 1H), 6.87 (s, 1H), 4.63 (q, *J* = 6.9 Hz, 2H), 4.26 (q, *J* = 7.2 Hz, 2H), 4.02 (s, 3H), 3.88 (br. s., 4H), 2.80 - 2.86 (m, 2H), 2.54 (br. s., 4H), 2.40 (br. s., 3H), 1.48 (t, *J* = 7.1 Hz, 3H), 1.47 (t, *J* = 7.1 Hz, 3H), 0.82 - 0.90 (m, 4H), 0.73 - 0.79 (m, 4H).

10

Example 108



15

4-(5-((cyanomethyl)(methyl)carbamoyl)-1-ethyl-1*H*-pyrazol-3-ylamino)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

[00503] The title compound was prepared using a method analogous to that used to prepare example **103**.


[00504] MS (ESI) *m/z* 529.3 (*M*+*H*).

[00505] ¹H NMR (500 MHz, CDCl₃) δ: 8.37 (br. s., 1H), 7.77 (s, 1H), 7.39 (s, 1H), 6.86 (s, 1H), 4.62 (q, *J* = 7.2 Hz, 2H), 4.54 (br. s., 2H), 4.27 - 4.34 (m, 1H), 4.04 (s,

20

3H), 3.41 (br. s., 3H), 2.79 - 2.86 (m, 2H), 1.47 (t, J = 7.2 Hz, 6H), 0.83 - 0.89 (m, 4H), 0.73 - 0.78 (m, 4H).

Example 109



5

4-((5-((2-acetamidoethyl)carbamoyl)-1-ethyl-1H-pyrazol-3-yl)amino)-N,N-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide


[00506] The title compound was prepared using a method analogous to that used to prepare example **103**.

[00507] MS (ESI) m/z 561.5 (M+H).

[00508] ¹H NMR (500 MHz, CDCl₃) δ: 7.77 (br. s., 2H), 7.40 (s, 1H), 6.72 (s, 1H), 4.48 - 4.56 (m, 4H), 3.99 (s, 3H), 3.61 - 3.66 (m, 2H), 3.52 - 3.58 (m, 2H), 2.81 (m, 2H), 2.02 (s, 3H), 1.44 (t, J = 7.1 Hz, 3H), 1.34 (t, J = 6.9 Hz, 3H), 1.26 (s, 1H), 0.82 - 0.90 (m, 4H), 0.70 - 0.77 (m, 4H)

15

Example 110



N,N-dicyclopropyl-4-((5-(diethylcarbamoyl)-1-ethyl-1H-pyrazol-3-yl)amino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide


20

[00509] The title compound was prepared using a method analogous to that used to prepare example **103**.

[00510] MS (ESI) m/z 532.4 (M+H).

[00511] ^1H NMR (500 MHz, CDCl_3) δ : 8.29 (br. S., 1H), 7.70 (s, 1H), 7.23 (s, 1H), 6.85 (s, 1H), 4.62 (q, $J = 6.9$ Hz, 2H), 4.20 (q, $J = 7.2$ Hz, 2H), 4.00 (s, 3H), 3.51 - 3.62 (m, 4H), 2.77 - 2.85 (m, 2H), 1.40 - 1.47 (m, 6H), 1.27 (t, $J = 7.1$ Hz, 6H), 0.80 - 0.88 (m, 4H), 0.72 - 0.77 (m, 4H)

5

Example 111

N,N-dicyclopropyl-4-(((5-(((2*R*,6*S*)-2,6-dimethyl-4-morpholinyl)carbonyl)-1-ethyl-1*H*-pyrazol-3-yl)amino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-
d]pyrrolo[2,3-b]pyridine-7-carboxamide


10

[00512] The title compound was prepared using a method analogous to that used to prepare example 103.

[00513] MS (ESI) m/z 574.4 ($M+H$).


15 [00514] ^1H NMR (500 MHz, CDCl_3) δ : 8.30 (br. S., 1H), 7.72 (s, 1H), 7.21 (s, 1H), 6.86 (s, 1H), 4.62 (q, $J = 6.9$ Hz, 2H), 4.21 - 4.28 (m, 2H), 4.07 - 4.20 (m, 2H), 4.01 (s, 3H), 3.61 (br. s, 2H), 2.91 (br. s, 1H), 2.79 - 2.85 (m, 2H), 2.60 (br. S. 1H), 1.47 (t, $J = 7.2$ Hz, 3H), 1.45 (t, $J = 7.2$ Hz, 3H), 1.07 - 1.36 (m, 6H), 0.81 - 0.88 (m, 4H), 0.72 - 0.78 (m, 4H)

20

Example 112


***N,N*-dicyclopropyl-4-(5-(dimethylcarbamoyl)-1-methyl-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

- 5 **112A** Preparation of (*Z*)-*tert*-butyl 2-(1-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridin-4-ylamino)-4,4-dimethoxy-3-oxobut-1-enyl)-1-methylhydrazinecarboxylate




- [00515] The title compound was prepared using a method analogous to that used to
10 prepare example **101B**.
[00516] MS (ESI) *m/z* 611.7 (*M*+*H*).

- 112B** Preparation of *N,N*-dicyclopropyl-6-ethyl-4-(5-formyl-1-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide
- 15 b]pyridine-7-carboxamide



- [00517] The title compound was prepared using a method analogous to that used to
 prepare Example **101C**.
20 [00518] MS (ESI) *m/z* 447.3 (*M*+*H*).

- 112C** Preparation of 3-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridin-4-ylamino)-1-methyl-1*H*-pyrazole-5-carboxylic acid



[00519] The title compound was prepared using a method analogous to that used to prepare example **101**.

5 **[00520]** MS (ESI) m/z 463.3 (M+H).


112 Preparation of *N,N*-dicyclopropyl-4-(5-(dimethylcarbamoyl)-1-methyl-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

10 **[00521]** A mixture of 3-(7-(dicyclopropylcarbamoyl)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridin-4-ylamino)-1-methyl-1*H*-pyrazole-5-carboxylic acid (39.8 mg, 0.086 mmol), HATU (49.0 mg, 0.129 mmol) and dimethylamine, 2M in THF (0.086 mL, 0.172 mmol) in DMF (2 mL) was stirred at room temperature for 2h. Water was added and the reaction mixture was extracted
15 with ethyl acetate 3 times. The combined organic layers were washed twice with 10% lithium chloride, dried over magnesium sulfate, filtered and concentrated *in vacuo*. The crude was purified by preparative HPLC to give *N,N*-dicyclopropyl-4-(5-(dimethylcarbamoyl)-1-methyl-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (23 mg, 54.1 % yield)
20 as an off-white solid.

[00522] MS (ESI) m/z 490.2 (M+H).

[00523] ^1H NMR (400 MHz, CDCl_3) δ : 8.28 (s, 1H), 7.67 (s, 1H), 7.34 (s, 1H), 6.86 (s, 1H), 4.61 (q, $J = 7.0$ Hz, 2H), 4.01 (s, 3H), 3.95 (s, 3H), 3.29 (s, 3H), 3.16 (s, 3H), 2.79 - 2.86 (m, 2H), 1.47 (t, $J = 7.0$ Hz, 3H), 0.82 - 0.89 (m, 4H), 0.72 - 0.79 (m,
25 4H).

Example 113



***N,N*-dicyclopropyl-6-ethyl-4-(1-methyl-5-(thiomorpholine 1,1-dioxide-4-carbonyl)-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**


5

[00524] The title compound was prepared using a method analogous to that used to prepare example **112**.

[00525] MS (ESI) *m/z* 580.3 (M+H).

[00526] ¹H NMR (400 MHz, CDCl₃) δ: 8.29 (s, 1H), 7.69 (s, 1H), 7.30 (s, 1H),
 10 6.83 (s, 1H), 4.57 (q, *J* = 6.8 Hz, 2H), 4.31 (br. s., 4H), 4.00 (s, 3H), 3.93 (s, 3H), 3.15 (br. s., 4H), 2.77 - 2.88 (m, 2H), 1.45 (t, *J* = 7.0 Hz, 3H), 0.80 - 0.91 (m, 4H), 0.69 - 0.80 (m, 4H).

Example 114



15

***N,N*-dicyclopropyl-4-(1-(2,2-difluoroethyl)-5-(dimethylcarbamoyl)-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**


20 **[00527]** The title compound was prepared using a method analogous to that used to prepare Example **112**; except utilizing example **80A**.

[00528] MS (ESI) *m/z* 540.4(M+H).

[00529] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ: 8.41 (br. s., 1H), 7.73 (s, 1H),
 7.43 (s, 1H), 6.86 (s, 1H), 6.17 (tt, *J* = 55.9, 4.2 Hz, 1H), 4.57 - 4.72 (m, 4H), 4.01 (s,

3H), 3.30 (s, 3H), 3.16 (s, 3H), 2.78 - 2.86 (m, 2H), 1.47 (t, $J = 7.0$ Hz, 3H), 0.80 - 0.89 (m, 4H), 0.71 - 0.80 (m, 4H).

Example 115



5

N,N-dicyclopropyl-6-ethyl-4-(5-(hydroxymethyl)-1-methyl-1H-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

10

[00530] Title compound was prepared by a method similar to that used for example 102.


[00531] MS (ESI) m/z 449.3 (M+H)

[00532] ^1H NMR (CDCl_3) δ ppm 8.11 (br s, 1H), 7.67 (s, 1H), 7.16 (s, 1H), 6.88 (s, 1H), 4.75 (d, 2H, $J = 5.94$ Hz), 4.67 (q, 2H, $J = 7.26$ Hz), 4.03 (s, 3H), 3.89 (s, 3H), 2.84 (m, 2H), 1.68 (t, 3H, $J = 6.16$ Hz), 1.50 (t, 3H, $J = 7.15$ Hz), 0.75-0.89 (m, 8H)

15

20

Example 116



N,N-dicyclopropyl-6-ethyl-1-methyl-4-(1-methyl-5-(morpholinomethyl)-1H-pyrazol-3-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide


25

[00533] To a stirred solution of *N,N*-dicyclopropyl-6-ethyl-4-(5-formyl-1-methyl-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (example **112B**, 22 mg, 0.049 mmol) and molecular sieves 4Å (30 mg) in MeOH (2 mL) was added morpholine (0.017 mL, 0.197 mmol). After the solution was stirred at room temperature for 12 hrs, sodium borohydride (7.46 mg, 0.197 mmol) was then added and stirred for 3 hrs. The solution was brought to pH = 4 with 1N HCl, neutralized with saturated aqueous sodium bicarbonate, and then extracted with CH₂Cl₂. The organic layer was dried over anhydrous magnesium sulfate and concentrated *in vacuo*. The residue was purified by silica-gel flash chromatography using an Isco 40 g column eluting from 0-10% MeOH/CH₂Cl₂ to yield *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(1-methyl-5-(morpholinomethyl)-1*H*-pyrazol-3-ylamino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (14.3 mg, 0.027 mmol, 54.9 % yield) as a white solid.

[00534] MS (ESI) *m/z* 518.3 (M+H)

[00535] ¹H NMR (CDCl₃) δ ppm 7.99 (br s, 1 H), 7.57 (s, 1 H), 6.99 (s, 1 H), 6.78 (s, 1H), 4.56 (q, 2 H *J* = 7.04 Hz), 3.93 (s, 3H), 3.77 (s, 3H), 3.64 (m, 4H), 3.48 (s, 2H), 2.75 (m, 2H), 2.45 (m, 4H), 1.42 (t, 3H *J* = 7.02 Hz), 0.66-0.80 (m, 8H)

Example 117




25 *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(1-methyl-5-((4-methylpiperazin-1-yl)methyl)-1*H*-pyrazol-3-ylamino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

[00536] Synthesized using a method analogous to that used to prepare example 116.

[00537] MS (ESI) m/z 531.4 (M+H)

[00538] ^1H NMR (CDCl_3) δ ppm 8.05 (br s, 1 H), 7.60 (s, 1 H), 7.01 (s, 1H), 6.78 (s, 1 H), 4.56 (q, 2 H $J = 6.75$ Hz), 3.95 (s, 3H), 3.72 (s, 3H), 3.61 (s, 2H), 2.96 (m, 4H), 2.76 (m, 4H), 2.74 (m, 2H), 2.71 (s, 3H), 1.42 (t, 3H, $J = 7.04$ Hz), 0.67-0.80 (m, 8H)

Example 118




N,N-dicyclopropyl-4-(5-((dimethylamino)methyl)-1-methyl-1*H*-pyrazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00539] Synthesized using a method analogous to that used to prepare example 116.

[00540] MS (ESI) m/z 476.4 (M+H)

[00541] ^1H NMR (CDCl_3) δ ppm 7.45 (s, 1 H), 7.33 (s, 1H), 6.84 (s, 1 H), 4.52 (q, 2H, $J = 7.04$ Hz), 4.25 (s, 2H), 3.98 (s, 3H), 3.90 (s, 3H), 2.80 (s, 6H), 2.77 (m, 2H), 1.37 (t, 3H, $J = 6.82$ Hz), 0.64-0.87 (m, 8H)


Example 119



N,N-dicyclopropyl-6-ethyl-4-(1-ethyl-5-((4-methylpiperazin-1-yl)methyl)-1*H*-pyrazol-3-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide


- 5 [00542] Prepared using intermediate **101B** and procedure as used for example **116**.
 [00543] MS (ESI) *m/z* 545.4 (*M*+*H*)
 [00544] ¹H NMR (CDCl₃) δ ppm 8.03 (s, 1 H), 7.57 (s, 1 H), 7.00 (s, 1H), 6.78 (s, 1 H), 4.55 (q, 2 H, *J* = 7.04 Hz), 4.03 (q, 2H, *J* = 5.72 Hz), 3.94 (s, 3H), 3.60 (s, 2H),
 10 2.93 (br s, 3H), 2.63-2.84 (br m, 10H), 1.42 (t, 3H, *J* = 7.04 Hz), 1.35 (t, 3H, *J* = 7.26 Hz), 0.69-0.79 (m, 8H)

Example 120



- 15 *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(2-methylthiazol-4-ylamino)-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

120A Preparation of 4-(2-chloroacetamido)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide



- 20 [00545] To a round bottom flask charged with 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (example **1J**, 47.2 mg, 0.139 mmol) was added dichloromethane (697 μL) and triethylamine (35.0 μL, 0.251 mmol). Chloroacetyl chloride (12.20 μL, 0.153 mmol) was added
 25 dropwise. The reaction mixture was stirred at rt for 16h. Additional chloroacetyl chloride (12.20 μL, 0.153 mmol) was added. The reaction mixture was diluted with

dichloromethane. The organics were washed with water, dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. 4-(2-Chloroacetamido)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (58.5 mg, 101 % yield) was isolated as an orange paste. Material was used in subsequent reactions with no further purification.

[00546] MS (ESI) *m/z* 415.0 (M+H)

[00547] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 8.04 (br s, 1H), 6.90 (s, 1H), 5.31 (s, 2H), 4.62 (q, 2H, *J* = 7.11 Hz), 4.24 (s, 3H), 2.77 – 2.88 (m, 2H), 1.47 (t, 3H, *J* = 7.03 Hz), 0.83 – 0.92 (m, 4H), 0.71 – 0.80 (m, 4H)


120 Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(2-methylthiazol-4-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00548] To a solution of 4-(2-chloroacetamido)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (58.5 mg, 0.141 mmol) in DMF (176 μ L) was added thioacetamide (12.71 mg, 0.169 mmol). The vial was capped and the reaction mixture was heated at 80 °C 3 h. The reaction mixture was concentrated *in vacuo*. The crude residue was taken up in MeOH and purified by prep-HPLC. *N,N*-Dicyclopropyl-6-ethyl-1-methyl-4-(2-methylthiazol-4-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (4.05 mg, 6.6% yield) was isolated as an off-white solid.


[00549] MS (ESI) *m/z* 436.0 (M+H)

[00550] ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.78 (s, 1H), 8.08 (s, 1H), 7.74 (s, 1H), 7.20 (s, 1H), 4.55 (q, 2H, *J* = 7.19 Hz), 4.03 (s, 3H), 2.87 – 2.98 (m, 2H), 2.65 (s, 3H), 1.34 (t, 3H, *J* = 7.03 Hz), 0.73 – 0.81 (m, 4H), 0.62 – 0.69 (m, 4H)

Example 121




***N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-methyloxazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

121A Preparation of 1-azidopropan-2-one

[00551] To a solution of chloroacetone (0.680 mL, 8.53 mmol) in acetone (10 mL)
5 and water (5.00 mL) was added sodium azide (0.555 g, 8.53 mmol). The reaction
mixture was stirred at rt 3.5 h. The reaction mixture was concentrated under reduced
pressure. The remaining aqueous solution was extracted with dichloromethane (2 x).
The combined organics were washed with brine, dried over anhydrous sodium sulfate,
filtered and concentrated *in vacuo* to yield 1-azidopropan-2-one (0.641 g, 6.47 mmol,
10 76 % yield) as a brown oil. This material was used without any purification.

[00552] ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 3.96 (s, 2H), 2.21 (s, 3H)

121B Preparation of *N,N*-dicyclopropyl-6-ethyl-4-isothiocyanato-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide

[00553] To a mixture of 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-
dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (example **1J**, 194.5 mg,
0.575 mmol) in dichloromethane (2874 μL) in a round bottom flask under nitrogen
20 was added 1,1'-thiocarbonyldi-2(1*H*)-pyridone (133 mg, 0.575 mmol). The reaction
mixture was stirred at rt for 16 h. The reaction mixture was concentrated under
reduced pressure. *N,N*-Dicyclopropyl-6-ethyl-4-isothiocyanato-1-methyl-1,6-
dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (180 mg, 94% yield)
was isolated as a yellow sticky solid. The material was used as is in subsequent
25 reactions.

[00554] MS (ESI) *m/z* 381.1 (M+H)

121 Preparation of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(5-methyloxazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide


[00555] To a solution of *N,N*-dicyclopropyl-6-ethyl-4-isothiocyanato-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **121B**, 45.7 mg, 0.120 mmol) in dioxane (343 μ L) was added 1-azidopropan-2-one (14.28 mg, 0.144 mmol) and triphenylphosphine (37.8 mg, 0.144 mmol). The contents of the flask were immersed into a preheated oil bath for 35 min and cooled to room temperature. The residue was diluted with ethyl acetate and quenched with saturated aqueous sodium bicarbonate. The aqueous layer was extracted with ethyl acetate.

The combined organics were washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The crude residue was purified by flash chromatography on an Isco 12 g column eluting with 1 – 10% MeOH/CH₂Cl₂. The pale yellow solid was taken up in MeOH and further purified by preparative HPLC. *N,N*-Dicyclopropyl-6-ethyl-1-methyl-4-(5-methyloxazol-2-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide was isolated (10 mg, 25% yield) as a white solid.

[00556] MS (ESI) *m/z* 420.0 (M+H)

[00557] ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 9.86 (s, 1H), 8.05 (s, 1H), 7.18 (s, 1H), 6.66 (d, 1H, *J* = 1.51 Hz), 4.41 (q, 2H, *J* = 7.28 Hz), 4.02 (s, 3H), 2.85 – 2.98 (m, 2H), 2.21 – 2.29 (m, 3H), 1.27 (t, 3H, *J* = 7.15 Hz), 0.71 – 0.80 (m, 4H), 0.60 – 0.70 (m, 4H)

Example 122



25 *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-(3-methyl-1,2,4-thiadiazol-5-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide


[00558] A mixture of *N,N*-dicyclopropyl-6-ethyl-4-isothiocyanato-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **121B**, 0.042

g, 0.111 mmol), acetamidine hydrochloride (10.49 mg, 0.111 mmol), and Hünig'sBase (0.058 mL, 0.333 mmol) in DMF (0.617 mL) was stirred at rt ON. Diethyl azodicarboxylate (0.088 mL, 0.222 mmol) was added and stirring continued at room temperature for 16 h. The reaction mixture was concentrated. The residue was taken up in MeOH and purified by preparative HPLC. *N,N*-Dicyclopropyl-6-ethyl-1-methyl-4-(3-methyl-1,2,4-thiadiazol-5-ylamino)-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (4.32 mg, 9% yield) was isolated as a pale yellow solid.

[00559] MS (ESI) m/z 437.1 (M+H)

[00560] ^1H NMR (400 MHz, MeOD) δ ppm 8.10 (s, 1 H), 7.28 (s, 1 H), 4.73 (q, 2 H, $J = 7.19$ Hz), 4.13 (s, 3 H), 2.93 - 3.01 (m, 2 H), 2.53 (s, 3 H), 1.53 (t, 3 H, $J = 7.03$ Hz), 0.83 - 0.92 (m, 4 H), 0.76 - 0.83 (m, 4 H)

Example 123




Preparation of *N,N*-dicyclopropyl-6-ethyl-4-(3-isopropyl-1,2,4-thiadiazol-5-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00561] This compound was prepared according to example 122 using isobutyrimidamide hydrochloride (15.08 mg, 0.123 mmol) and diisopropyl azodicarboxylate (47.8 μL , 0.246 mmol). *N,N*-Dicyclopropyl-6-ethyl-4-(3-isopropyl-1,2,4-thiadiazol-5-ylamino)-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (21 mg, 34.9 % yield) was isolated as an off-white solid.

[00562] MS (ESI) m/z 465.3 (M+H)

[00563] ^1H NMR (400 MHz, DMSO- d_6) δ ppm 12.24 (s, 1H), 8.18 (s, 1H), 7.28 (s, 1H), 4.64 (q, 2H, $J = 6.94$ Hz), 4.06 (s, 3H), 3.10 (ddd, 1H, $J = 13.93, 6.90, 6.78$ Hz), 2.86 - 3.00 (m, 2H), 1.42 (t, 3H, $J = 7.03$ Hz), 1.32 (s, 3H), 1.31 (s, 3H), 0.72 - 0.81 (m, 4H), 0.63 - 0.71 (m, 4H)

Example 124




5 **4-(1,2,4-thiadiazol-5-ylamino)-N,N-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide**

[00564] A mixture of *N,N*-dicyclopropyl-6-ethyl-1-methyl-4-thioureido-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (**example 2B**, 30 mg, 0.075 mmol) and 1,1-dimethoxy-*N,N*-dimethylmethanamine (1 g, 8.39 mmol) was
 10 heated at 50 °C for 2h. Excess 1,1-dimethoxy-*N,N*-dimethylmethanamine was evaporated *in vacuo* and the crude example was dissolved in dichloromethane (1 mL) and *O*-(mesitylsulfonyl)hydroxylamine (24.4 mg, 0.113 mmol) was added. The reaction mixture was stirred at room temperature for 3h. Water was added and the layers were separated. The organic layer was dried over magnesium sulfate, filtered
 15 and concentrated *in vacuo*. The crude residue was purified by preparative HPLC to give 4-(1,2,4-thiadiazol-5-ylamino)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (7 mg, 19.20 % yield) as a yellow solid.

[00565] MS (ESI) *m/z* 423.0 (M+H).

20 [00566] ¹H NMR (500 MHz, CD₃OD) δ: 8.32 (s, 1H), 8.27 (s, 1H), 7.38 (d, *J* = 4.4 Hz, 1H), 7.28 (s, 1H), 4.74 (q, *J* = 7.0 Hz, 2H), 4.15 (s, 3H), 2.94 - 3.00 (m, 2H), 1.53 (t, *J* = 7.0 Hz, 3H), 0.82 - 0.89 (m, 4H), 0.75 - 0.82 (m, 4H)

Example 125




***N,N*-dicyclopropyl-4-(1,5-dimethyl-1*H*-1,2,4-triazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide**

125A

Preparation of 4-(3-acetylthioureido)-N,N-dicyclopropyl-6-

5 ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide



[00567] To a solution of 4-amino-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **1J**, 60 mg, 0.177 mmol) in acetone (1.182 mL) was added acetyl isothiocyanate (0.020 mL, 0.230 mmol). The reaction mixture was stirred at room temperature for 6h. LC/MS showed still half of the starting material left. Another 0.5 equivalent of acetyl isothiocyanate was added and the reaction mixture was heated at 35 °C overnight. Solvent was evaporated and the crude product was purified by flash chromatography on silica gel using an automated ISCO system (24 g column, eluting with 2-8% methanol / dichloromethane). 4-(3-Acetylthioureido)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (34 mg, 43.6 % yield) was obtained as a yellow solid.

[00568] MS (ESI) m/z 440.1 (M+H).

125

Preparation of *N,N*-dicyclopropyl-4-(1,5-dimethyl-1*H*-1,2,4-

triazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide

[00569] A mixture of 4-(3-acetylthioureido)-*N,N*-dicyclopropyl-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine-7-carboxamide (example **125A**, 17 mg, 0.039 mmol) and methylhydrazine (2.486 μ L, 0.046 mmol) in acetic acid (0.5 mL) was heated at 80 $^{\circ}$ C for 4h, LC/MS showed starting material still remaining. Additional methylhydrazine (2.486 μ L, 0.046 mmol) was added and heating continued for another 2h, reaction complete. Solvent was evaporated and the crude

was purified by preparative HPLC to furnish *N,N*-dicyclopropyl-4-(1,5-dimethyl-1*H*-1,2,4-triazol-3-ylamino)-6-ethyl-1-methyl-1,6-dihydroimidazo[4,5-*d*]pyrrolo[2,3-*b*]pyridine-7-carboxamide (5.2 mg, 30.7 % yield) as an off-white solid.


[00570] MS (ESI) *m/z* 434.3 (M+H).

- 5 **[00571]** ¹H NMR (CDCl₃) δ: 7.96 (s, 1H), 7.65 (s, 1H), 6.83 (s, 1H), 4.67 (q, *J* = 7.1 Hz, 2H), 4.01 (s, 3H), 3.82 (s, 3H), 2.77 - 2.84 (m, 2H), 2.43 (s, 3H), 1.48 (t, *J* = 7.2 Hz, 3H), 0.79 - 0.86 (m, 4H), 0.71 - 0.77 (m, 4H).

WHAT IS CLAIMED IS:

1. A compound of the formula

5



(I)


or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein

10 X is C₁₋₄ alkyl;

Y is C₁₋₄ alkyl;

R is

15



any of which are optionally fused with a 5 or 6 membered carbocycle or heterocycle,
said heterocycle having one heteroatom selected from NR³ or S, said fused
20 carbocycle or heterocycle being optionally substituted with 0-3 R¹.

R¹ is H, halo, CN, C₁₋₆ alkyl substituted with 0-3 R^c, CF₃, CONR^aR^a, NR^aR^a,
COOR^b, SO₂-C₁₋₄alkyl, C(O)R^d, cycloalkyl substituted with 0-3 R^c, furanyl,
tetrahydropyranyl or pyridinyl;

R^2 is H, C_{1-6} alkyl substituted with 0-3 R^c , $C(O)O-C_{1-4}$ alkyl, SO_2-C_{1-4} alkyl, cycloalkyl substituted with 0-3 R^c , tetrahydropyranyl or R^2 is absent;

R^3 is H, $C(O)O-C_{1-4}$ alkyl, or R^3 is absent;

R^a is H, C_{1-6} alkyl substituted with 0-3 R^c , C_{3-6} cycloalkyl substituted with 0-3 R^c , tetrahydropyranyl or dioxotetrahydrothiophenyl;

R^b is H or C_{1-6} alkyl;

R^c is H, halo, CN, OH, $O-C_{1-4}$ alkyl, $O-C_{1-4}$ alkyl- $O-C_{1-4}$ alkyl, NH_2 , $N(C_{1-4}$ alkyl) $_2$, $C(O)N(C_{1-4}$ alkyl) $_2$, SO_2-C_{1-4} alkyl, or morpholinyl or piperazinyl either of which are optionally substituted with 0-1 C_{1-4} alkyl;

R^d is C_{1-6} alkyl, or azeridinyl, azetidiny, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, dioxidothiomorpholinyl or tetrahydropyranyl, any of which are substituted with 0-2 R^c ; and

R^e is H, halo, CN, C_{1-4} alkyl, OH, $O-C_{1-4}$ alkyl, SO_2-C_{1-4} alkyl, $NHC(O)-C_{1-4}$ alkyl, morpholinyl, $OC(O)-C_{1-4}$ alkyl, $C(O)N(C_{1-4}$ alkyl) $_2$, or $O-C_{1-4}$ alkyl- $O-C_{1-4}$ alkyl.

15


2. The compound according to claim 1


wherein

X is ethyl;

20 Y is methyl; and

R is





any of which are optionally substituted with 0-3 R¹;

or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof.


5

3. The compound according to claim 2

wherein

R is


10



4. The compound according to claim 2

15 wherein

R is




any of which are optionally substituted with 0-2 R^1

5

5. The compound according to claim 2

wherein



R is

R^1 is H, halo, CN, C_{1-6} alkyl substituted with 0-3 R^c , CF_3 , $CONR^aR^a$, $COOR^b$,

10 SO_2-C_{1-4} alkyl, $C(O)R^d$, cycloalkyl substituted with 0-3 R^c , or pyridinyl;

R^a is H, C_{1-6} alkyl substituted with 0-3 R^c , C_{3-6} cycloalkyl substituted with 0-3 R^c , tetrahydropyranyl, or dioxotetrahydrothiophenyl;

R^b is H or C_{1-6} alkyl;

R^c is H, halo, OH, $O-C_{1-4}$ alkyl, SO_2-C_{1-4} alkyl or morpholinyl;

15 R^d is C_{1-6} alkyl, or azetidiny, pyrrolidinyl, morpholinyl, piperazinyl or dioxidothiomorpholinyl, any of which are substituted with 0-2 R^c ; and


R^e is H, halo, CN, OH, $O-C_{1-4}$ alkyl, SO_2-C_{1-4} alkyl, $NHC(O)-C_{1-4}$ alkyl or morpholinyl;

or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof.

20

6. The compound according to claim 2

wherein



R^1 is H, halo, C_{1-6} alkyl substituted with 0-3 R^c , CF_3 , $CONR^aR^a$, $COOR^b$, $C(O)R^d$, cycloalkyl substituted with 0-3 R^c , or furanyl;

R^2 is H, C_{1-6} alkyl substituted with 0-3 R^c , SO_2-C_{1-4} alkyl, cycloalkyl
5 substituted with 0-3 R^c , or tetrahydropyranyl;

R^a is H, or C_{1-6} alkyl substituted with 0-3 R^c ;

R^b is H or C_{1-6} alkyl;

R^c is H, halo, CN, OH, $O-C_{1-4}$ alkyl, $O-C_{1-4}$ alkyl- $O-C_{1-4}$ alkyl, NH_2 , $N(C_{1-4}$
alkyl) $_2$, $C(O)N(C_{1-4}$ alkyl) $_2$, SO_2-C_{1-4} alkyl, or morpholinyl or piperazinyl, either of
10 which are optionally substituted with 0-1 C_{1-4} alkyl;

R^d is C_{1-6} alkyl, or morpholinyl, piperazinyl or dioxidothiomorpholinyl, any of
which are substituted with 0-2 R^c ;

R^e is H, C_{1-4} alkyl, CN, OH, $NHC(O)-C_{1-4}$ alkyl or morpholinyl;

or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof.

15

7. The compound according to claim 6
wherein

R^1 is C_{1-6} alkyl substituted with 0-3 R^c ; and

R^2 is C_{1-6} alkyl.

20

8. A pharmaceutical composition comprising a pharmaceutically
acceptable carrier and one or more compounds of any of claims 1-7 or
pharmaceutically acceptable salts, tautomers or stereoisomers thereof.

25 9. A compound of any of claims 1-7 or pharmaceutically acceptable salts,
tautomers or stereoisomers thereof, for use in therapy.

10. Use of a compound of any of claims 1-7 or pharmaceutically
acceptable salts, tautomers or stereoisomers thereof, for the manufacture of a
30 medicament for the treatment of myeloproliferative diseases or cancer.

11. Use of a compound according to Claim 10, wherein the cancer is selected from solid tumors of the pancreas, prostate, lung, head and neck, breast, colon, ovary, gastric cancer, multiple myeloma, melanoma, neuroblastoma, glioblastoma, systemic mastocytosis, and hematological malignancies such as acute myelogenous leukemia (including refractory acute myeloid leukemia) and acute lymphoid leukemia

12. Use of a compound according to Claim 10, wherein the myeloproliferative disease is polycythemia vera, essential thrombocytopenia or primary myelofibrosis.

13. One or more compound(s) according to claim 1 as an active agent for treating myeloproliferative diseases or cancer in a mammal.

14. One or more compound(s) as an active agent for treating cancer according to claim 13, wherein the cancer is selected from solid tumors of the pancreas, prostate, lung, head and neck, breast, colon, ovary, gastric cancer, multiple myeloma, melanoma, neuroblastoma, glioblastoma, systemic mastocytosis, and hematological malignancies such as acute myelogenous leukemia (including refractory acute myeloid leukemia) and acute lymphoid leukemia

15. One or more compound(s) as an active agent for treating myeloproliferative disease according to claim 13, wherein the myeloproliferative disease is polycythemia vera, essential thrombocytopenia or primary myelofibrosis.