

-1-

Modulation of Calcium Channels

This invention relates to the discovery that a group of 2-aryl-3-aroylebenzo[b]thiophenes are effective in modulating calcium channels, increasing the density of calcium channels in vascular and cardiac tissue, with no changes in inotropic or pressor response.

Replacement therapy with estrogen is generally acknowledged to produce beneficial effects on the cardiovascular system in postmenopausal women. See Knopt, *Obstet. Gynecol.*, 72, 23s-30s (1988). In postmenopausal women who receive estrogens, the cardiovascular mortality rate is reduced by about 30% to about 50%, and the cerebrovascular mortality rate is reduced by about 50%. See Stampfer et al., *N. Engl. J. Med.*, 325, 756-762 (1991). Although these beneficial cardiovascular effects may involve alterations in lipid profile, recent data suggests that estrogen may also have beneficial effects on the vascular responses of atherosclerotic coronary arteries. See Gisclard et al., *J. Pharmacol. and Experimental Therapeutics*, 244, 19-22 (1988); Williams et al., *Circulation*, 81, 1680-1687 (1990); Gangar et al., *Lancet*, 388, 839-842 (1991); and Williams et al., *JACC*, 20, 452-457 (1992). Both endothelial-independent and endothelial-dependent effects of estrogen have been described in vascular tissue. See Jiang et al., *Br. J. Pharmacol.*, 104, 1033-1037 (1991); Jiang et al., *American Journal of Physiology*, 32, H271-H275 (1992); Cheng and Gruetter, *European Journal Of Pharmacol.*, 215, 171-176 (1992); Mügge et al., *Cardiovas. Res.*, 27, 1939-1942 (1993); Salas et al., *European Journal of Pharmacol.*, 258, 47-55 (1994); Williams et al., *Circulation*, 81, 1680-1687 (1990); Cheng et al., *Life Sciences*, 10, 187-191 (1994); Gilligan et al., *Circulation*, 89, 2545-2551 (1994); and Reis et al., *Circulation*, 89, 52-60 (1994). Several reports have also

-2-


suggested that the vasodilating effects of estradiol and/or its ability to attenuate contractile responses may be mediated by inhibition of calcium influx via voltage dependent calcium channels. See Jiang et al., *Br. J.*

- 5 *Pharmacol.*, 104, 1033-1037 (1991); Jiang et al., *American Journal of Physiology*, 32, H271-H275 (1992); Collins et al., *Lancet*, 341, 1264 (1993); Muck et al., *Med. Sci. Res.*, 22, 19 (1994); and Salas et al., *European Journal of Pharmacol.*, 258, 47-55 (1994). Others have postulated that estradiol may
10 enhance cyclic AMP and cyclic GMP content, or increase ATP-sensitive potassium channels. See Mügge et al., *Cardiovas. Res.*, 27, 1939-1942 (1993); Sudhir et al., *Am. Heart J.*, 129, 726-732.

- The 2-aryl-3-arylbenzo[b]thiophene compounds that
15 are used in the methods of this invention were first developed by Jones and Suarez as anti-fertility agents. See U.S. Patent No. 4,133,814 (issued January 9, 1979). These compounds are generally useful in suppressing the growth of mammary tumors. Jones later found that a group of these
20 compounds are particularly useful for antiestrogen and antiandrogen therapy, especially in the treatment of mammary and prostatic tumors. See U.S. Patent 4,418,068 (issued November 29, 1983). One of these compounds, 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidino-
25 ethoxy)benzoyl]benzo[b]thiophene was clinically studied for the treatment of breast cancer. This compound is called raloxifene, formerly keoxifene.

- This invention provides methods for modulating calcium channels, increasing the density of calcium channels
30 in vascular and cardiac tissue with no changes in inotropic or pressor response, comprising administering to a warm-blooded animal in need thereof an effective amount of a compound of the formula

-3-



wherein R^1 and R^3 are independently hydrogen, C_1 - C_4 alkyl, $-\text{CO}-(C_1-C_6 \text{ alkyl})$, $-\text{CH}_2\text{Ar}$, or $-\text{CO}-\text{Ar}$, wherein Ar is phenyl or substituted phenyl;

R^2 is selected from the group consisting of pyrrolidino, hexamethyleneimino, and piperidino; or a pharmaceutically-acceptable salt thereof. The present invention also provides the use of the formula I compounds, or pharmaceutically-acceptable salts thereof, for the manufacture of a medicament for modulating calcium channels in vascular and cardiac tissue.

The present invention concerns the discovery that a select group of 2-aryl-3-aryloxybenzo[b]thiophenes (benzo[b]thiophenes), the compounds of formula I, are effective in modulating calcium channels, increasing the density of calcium channels in vascular and cardiac tissue, with no changes in inotropic or pressor response. Therefore, the present invention provides methods for modulating calcium channels in vascular and cardiac tissue. One aspect of the invention is a method for treating cardiac disorders, including but not limited to variant angina, exertional angina, unstable angina, ischemia-reperfusion injury to the myocardium, and arrhythmias. Another aspect is a method for treating cerebral vascular disorders, including but not limited to cerebral vasospasm due to arterial rupture, stroke, and migraine headaches. Another aspect is a method for treating renal disorders by increasing renal clearance due to increases in renal blood flow, useful for slowing of renal failure. Another aspect is a method for treating

-4-

gastrointestinal disorders, including but not limited to diseases related to diarrhea, such as IBS and IBD, diarrhea predominant. The therapeutic treatments provided by this invention are practiced by administering to a warm-blooded
5 animal in need thereof a pharmaceutically-effective amount of a compound of formula I or a pharmaceutically-acceptable salt thereof.

In the above formula, the term "C₁-C₆ alkyl" represents a straight, cyclic, or branched alkyl chain having
10 from one to six carbon atoms. Typical C₁-C₆ alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl, isopentyl, n-hexyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the like. The term "C₁-C₄ alkyl" represents a straight or
15 branched alkyl chain having one to four carbon atoms. Typical C₁-C₄ alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, secbutyl, isobutyl, and t-butyl.

The term "Ar" represents groups such as phenyl and substituted phenyl. The term "substituted phenyl", as used
20 herein, represents a phenyl group substituted with one or more moieties chosen from the group consisting of halogen, hydroxy, cyano, nitro, C₁-C₄ alkyl, C₁-C₄ alkoxy, acetyl, formyl, trichloromethyl, or trifluoromethyl. Examples of a substituted phenyl group include 4-chlorophenyl, 2,6-
25 dichlorophenyl, 2,5-dichlorophenyl, 3,4-dichlorophenyl, 3-chlorophenyl, 3-bromophenyl, 4-bromophenyl, 3,4-dibromophenyl, 3-chloro-4-fluorophenyl, 2-fluorophenyl, 4-hydroxyphenyl, 3-hydroxyphenyl, 2,4-dihydroxyphenyl, 3-nitrophenyl, 4-nitrophenyl, 4-cyanophenyl, 4-methylphenyl, 4-
30 ethylphenyl, 4-methoxyphenyl, 4-propylphenyl, 4-n-butylphenyl, 4-t-butylphenyl, 3-fluoro-2-methylphenyl, 2,3-difluorophenyl, 2,6-difluorophenyl, 2,6-dimethylphenyl, 2-fluoro-5-methylphenyl, 2,4,6-trifluorophenyl, 2-trifluoromethylphenyl, 2-chloro-5-trifluoromethylphenyl, 3,5-bis(trifluoromethyl)phenyl, 2-methoxyphenyl, 3-methoxyphenyl,
35 3,5-dimethoxyphenyl, 4-hydroxy-3-methylphenyl, 3,5-dimethyl-4-hydroxyphenyl, 2-methyl-4-nitrophenyl, 4-methoxy-2-

-5-

nitrophenyl, 2,4-dinitrophenyl, and the like. The term "C₁-C₄ alkoxy" represents groups such as methoxy, ethoxy, *n*-propoxy, isopropoxy, *n*-butoxy, *t*-butoxy, and the like. The term "halogen" represents fluoro, chloro, bromo, and iodo.

5 The term "pharmaceutically-effective amount" is used herein to represent an amount of the formula I compound that is capable of increasing the density of calcium channels in vascular and cardiac tissue. The particular dose of the formula I compound will of course be determined by the
10 particular circumstances surrounding the case, including the compound administered, the route of administration, the particular condition treated, and similar considerations.

 The term "modulating", as used herein, represents an increase in the density of calcium channels in vascular
15 and cardiac tissue, with no changes in inotropic or pressor response.

 The term "warm-blooded animal", as used herein, includes humans; companion animals, such as dogs and cats; and domestic animals, such as horses, cattle, sheep, swine,
20 goats and chickens. Preferably, the warm-blooded animal is a human or companion animal. More preferably, the warm-blooded animal is a human.

 While all the formula I compounds are useful for modulating calcium channels in vascular and cardiac tissue,
25 certain compounds are preferred. Preferably, R¹ and R³ are independently hydrogen, C₁-C₄ alkyl, -CO-(C₁-C₆ alkyl), or benzyl, and R² is piperidino or pyrrolidino. Representative compounds from this preferred group include 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-pyrrolidinoethoxy)benzoyl]benzo[b]-
30 thiophene, 6-methoxy-2-(4-methoxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]benzo[b]thiophene, 6-acetoxy-2-(4-acetoxyphenyl)-3-[4-(2-pyrrolidinoethoxy)benzoyl]benzo[b]thiophene, and 6-benzyloxy-2-(4-benzyloxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]benzo[b]thiophene.

35 More preferably, R¹ and R³ are independently hydrogen or C₁-C₄ alkyl, and R² is piperidino or pyrrolidino. Representative compounds from this more preferred group

-6-

include 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-pyrrolidinoethoxy)benzoyl]benzo[b]thiophene, 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]benzo[b]thiophene, 6-methoxy-2-(4-methoxyphenyl)-3-[4-(2-pyrrolidinoethoxy)benzoyl]benzo[b]thiophene, and 6-methoxy-2-(4-methoxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]benzo[b]thiophene. Most preferably, R¹ and R³ are hydrogen and R² is piperidino. This most preferred compound is 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]benzo[b]thiophene.

10 The formula I compounds used in the methods of the present invention can be made according to established procedures, such as those described in U.S. Patent Nos. 4,133,814, 4,418,068, and 4,380,635, all of which are incorporated by reference herein. In general, the process
15 starts with 6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene. This starting compound is protected, acylated at C-3 with a 4-(2-aminoethoxy)benzoyl group, and optionally deprotected to form the formula I compounds. Examples of the preparation of such compounds are provided in the U.S. Patents discussed
20 above.

 The compounds used in the methods of this invention form pharmaceutically-acceptable acid and, wherein R¹ and/or R³ is hydrogen, base addition salts with a wide variety of organic and inorganic acids and bases, including the
25 physiologically-acceptable salts which are often used in pharmaceutical chemistry. Typical inorganic acids used to form such salts include hydrochloric, hydrobromic, hydriodic, nitric, sulfuric, phosphoric, hypophosphoric, and the like. Salts derived from organic acids, such as aliphatic mono- and
30 dicarboxylic acids, phenyl-substituted alkanolic acids, hydroxyalkanoic and hydroxyalkandioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, may also be used. Such pharmaceutically-acceptable salts thus include acetate, phenylacetate, trifluoroacetate, acrylate, ascorbate,
35 benzoate, chlorobenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, methylbenzoate, o-acetoxybenzoate, naphthalene-2-benzoate, bromide, isobutyrate, phenylbutyrate,

-7-

and β -hydroxybutyrate, butyne-1,4-dioate, hexyne-1,6-dioate, caprate, caprylate, chloride, cinnamate, citrate, formate, fumarate, glycolate, heptanoate, decanoate, hippurate, lactate, malate, maleate, hydroxymaleate, malonate, mandelate, mesylate, nicotinate, isonicotinate, nitrate, oxalate, phthalate, terephthalate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, propiolate, propionate, phenylpropionate, salicylate, sebacate, succinate, suberate, sulfate, bisulfate, pyrosulfate, sulfite, bisulfite, sulfonate, benzenesulfonate, p-bromophenylsulfonate, chlorobenzene-sulfonate, ethanesulfonate, 2-hydroxyethanesulfonate, methanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, p-toluenesulfonate, xylenesulfonate, tartrate, and the like. The most preferred salt is the hydrochloride salt.

The pharmaceutically-acceptable acid addition salts are typically formed by reacting a compound of formula I with an equimolar or excess amount of acid. The reactants are generally combined in an organic solvent such as methanol, diethyl ether, or benzene. The salt normally precipitates out of solution within about one hour to 10 days and can be isolated by filtration, or the solvent can be stripped off by conventional means.

Bases commonly used for formation of salts include ammonium hydroxide and alkali and alkaline earth metal hydroxides, carbonates, as well as aliphatic primary, secondary, and tertiary amines, and aliphatic diamines. Bases especially useful in the preparation of addition salts include ammonium hydroxide, potassium carbonate, methylamine, diethylamine, ethylenediamine, and cyclohexylamine. These salts are generally prepared by reacting a formula I compound, wherein R^1 and/or R^3 are hydrogen, with one of the above bases in an organic solvent, such as methanol, diethyl ether, or benzene. The salts are isolated as described in the preceding paragraph.

These pharmaceutically-acceptable salts generally have enhanced solubility characteristics compared to the

-8-

compound from which they are derived, and thus are often more amenable to formulation as liquids or emulsions.

The formula I compounds are preferably formulated prior to administration such as in a pharmaceutical formulation comprising a compound of formula I and a pharmaceutically-acceptable carrier, diluent, or excipient. These pharmaceutical formulations are prepared by known procedures using well-known and readily available ingredients. In making these compositions, the active ingredient will usually be mixed with a carrier, diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper, or other container. When the carrier serves as a diluent, it may be a solid, semi-solid, or liquid material which acts as a vehicle, excipient, or medium for the active ingredient. The compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols, ointments containing, for example up to 10% by weight of active compound, soft and hard gelatin capsules, dermal patches, suppositories, sterile injectable solutions, and sterile packaged powders.

Some examples of suitable carriers, excipients, and diluents include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum, acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, micro-crystalline cellulose, polyvinylpyrrolidone, cross-linked polyvinylpyrrolidone, cellulose or derivatives thereof, water syrup, methyl cellulose, methyl and propyl hydroxybenzoates, talc, magnesium stearate and mineral oil. The formulations can additionally include lubricating agents, wetting agents (e.g. surfactant), emulsifying and suspending agents, disintegrating agents, preserving agents, sweetening agents, or flavoring agents. Compositions of the inventions may be formulated so as to provide quick, sustained, or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art.

-9-

The particular dosage of a compound of formula I required for modulating calcium channels in vascular and cardiac tissue, according to this invention, will depend upon the severity of the condition, the route of administration, and related factors that will be decided by the attending physician. Generally, effective daily doses will be from about 0.1 to about 1000 mg/day, and more typically from about 10 to about 100 mg/day. Such dosages will be administered to a subject in need thereof from once to about three times each day, or more often as needed to effectively treat the condition or symptom.

It is usually preferred to administer a compound of formula I in the form of an acid addition salt, as is customary in the administration of pharmaceuticals bearing a basic group, such as the piperidino group. For such purposes the following oral dosage forms are available.

In the formulations which follow, "Active ingredient" means a compound of formula I.

Formulation 1: Gelatin Capsules

Hard gelatin capsules are prepared using the following:

Ingredient	Quantity (mg/capsule)
Active ingredient	0.1 - 1000
Starch, NF	0 - 650
Starch flowable powder	0 - 650
Silicone fluid 350 centistokes	0 - 15

The ingredients are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules.

Examples of specific capsule formulations of raloxifene that have been made include those shown below:

-10-

Formulation 2: Raloxifene capsule

<u>Ingredient</u>	<u>Quantity (mg/capsule)</u>
Raloxifene	1
Starch, NF	112
Starch flowable powder	225.3
Silicone fluid 350 centistokes	1.7

Formulation 3: Raloxifene capsule

5

<u>Ingredient</u>	<u>Quantity (mg/capsule)</u>
Raloxifene	5
Starch, NF	108
Starch flowable powder	225.3
Silicone fluid 350 centistokes	1.7

Formulation 4: Raloxifene capsule

<u>Ingredient</u>	<u>Quantity (mg/capsule)</u>
Raloxifene	10
Starch, NF	103
Starch flowable powder	225.3
Silicone fluid 350 centistokes	1.7

10 *Formulation 5: Raloxifene capsule*

<u>Ingredient</u>	<u>Quantity (mg/capsule)</u>
Raloxifene	50
Starch, NF	150
Starch flowable powder	397
Silicone fluid 350 centistokes	3.0

The specific formulations above may be changed in compliance with the reasonable variations provided.

15

A tablet formulation is prepared using the ingredients below:

-11-

Formulation 6: Tablets

<u>Ingredient</u>	<u>Quantity (mg/tablet)</u>
Active ingredient	0.1 - 1000
Cellulose, microcrystalline	0 - 650
Silicon dioxide, fumed	0 - 650
Stearate acid	0 - 15

The components are blended and compressed to form tablets.

- 5 Alternatively, tablets each containing 0.1 - 1000 mg of Active ingredient are made up as follows:

Formulation 7: Tablets

<u>Ingredient</u>	<u>Quantity (mg/tablet)</u>
Active ingredient	0.1 - 1000
Starch	45
Cellulose, microcrystalline	35
Polyvinylpyrrolidone (as 10% solution in water)	4
Sodium carboxymethyl cellulose	4.5
Magnesium stearate	0.5
Talc	1

10

- The Active ingredient, starch, and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders which are then passed through a No. 14 mesh
15 U.S. sieve. The granules so produced are dried at 50°-60° C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate, and talc, previously passed through a No. 60 U.S. sieve, are then added to the granules which, after mixing, are compressed on a
20 tablet machine to yield tablets.

Suspensions each containing 0.1 - 1000 mg of Active ingredient per 5 mL dose are made as follows:

-12-

Formulation 8: Suspensions

Ingredient	Quantity (mg/5 ml)
Active ingredient	0.1 - 1000 mg
Sodium carboxymethyl cellulose	50 mg
Syrup	1.25 mg
Benzoic acid solution	0.10 mL
Flavor	q.v.
Color	q.v.
Purified water to	5 mL

The Active ingredient is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor, and color are diluted with some of the water and added, with stirring. Sufficient water is then added to produce the required volume.

Illustrative compounds that can be used in the methods of the present invention are shown in Table 1.

Table 1

Compound			
No.	R ¹ and R ³	R ²	Form
1	-C(O)-  -F	piperidino	base
2	-C(O)-  -F	piperidino	HCl
3	-C(O)- 	piperidino	base
4	-C(O)- 	piperidino	HCl
5	-C(O)CH ₂ CH ₂ CH ₃	piperidino	base
6	-C(O)CH ₂ CH ₂ CH ₃	piperidino	HCl
7	-C(O)C(CH ₃) ₃	piperidino	base
8	-C(O)C(CH ₃) ₃	piperidino	HCl
9	-C(O)CH ₂ C(CH ₃) ₃	piperidino	base

-13-

Table 1 (Cont'd)

Compound			
No.	R ¹ and R ³	R ²	Form
10	-C(O)CH ₂ C(CH ₃) ₃	piperidino	HCl
11	-C(O)-  -CH ₃	piperidino	HCl
12	-C(O)- 	piperidino	base
13	H	piperidino	base
14	H	piperidino	HCl
15	H	pyrrolodino	base
16	H	pyrrolodino	HCl
17	H	hexamethyleneimino	HCl
18	CH ₃	piperidino	HCl

The utility of the compounds of formula I is
 5 illustrated by the positive impact they have in at least one
 of the experiments described below.

Materials and Methods

Selection and dosing of rats were essentially as
 10 described by Sato et al. Sato et al., *J. Bone and Mineral
 Research*, 9, 715-724 (1994). Briefly, ovariectomized (ovex)
 virgin female rats (6 months old) were divided into 3 groups
 of 6, designated as: ovex; ethinyl estradiol (EE2, 0.1
 mg/kg/day p.o.); and raloxifene (compound 14, 1.0 mg/kg/day
 15 p.o.). A fourth group of sham-operated females (sham) served
 as a second control. Doses of EE2 and 14 were selected for
 comparable effects on bone density parameters (Sato et al.);
 incidentally they produced similarly significant (P, 0.05 vs
 ovex) effects at lowering total cholesterol (36±2 and 38±4
 20 mg/dl, EE2 and 14, respectively, vs 85±7 and 87±7 mg/dl, ovex
 and sham, respectively). Sham and ovex rats were
 administered vehicle (100 µg/g body weight of 20%
 hydroxypropyl-S-cyclodextrin). Animals were dosed for 35
 days and sacrificed by excess CO₂.

-14-

Hearts and aortas were carefully dissected, quickly frozen and stored at -70°C if membranes were not prepared immediately. Microsomal membrane vesicles were isolated from 3-4 g minced hearts or aortas from each group as previously described. Jones et al., *J. Biol. Chem.*, **254**, 530-535 (1979). Preparations were stored in 0.25 M sucrose/30 mM histidine at -70°C. Binding studies using increasing concentrations of the calcium channel ligand [³H] PN200-110 (0.01-4.0 nM) were done in 12x75 mm glass tubes (total volume 500 µl) at 23°C for 2 h using 100 (heart) or 200 (aorta) µg protein per tube. Assays were terminated by rapid filtration onto Whatman GF/C filter paper. Assay (and wash) buffer was 50 mM Tris/HCl (pH7.3), 1 mM EDTA and 12 mM MgCl₂. Nonspecific binding was defined as binding remaining in the presence of 1 µM nifedipine.

Radioligand binding affinity and receptor density were determined from saturation isotherm data using the nonlinear regression analysis program LUNDON-1. Lundeen and Gordon, , in *Receptor Binding in Drug Research*, 31-49, 1986.

Cardiovascular hemodynamic parameters in response to BAY k 8644 were determined in pithed rats from each of the four groups (sham, ovex, EE2, and 14) as described by Hayes and Bowling with the following modifications: the drug was administered through the femoral vein; and direct measurement of left ventricular systolic blood pressure was obtained by inserting a small section of PE 90 tubing attached to a pressure transducer directly into the left ventricle. Hayes and Bowling, *J. Pharmacol. and Exp. Ther.*, **241**, 861-869 (1987). Measurements were obtained of mean, systolic and diastolic blood pressure, heart rate, left ventricular systolic pressure and left ventricular dP/dt.

Results

EE2 and 14 effects on Ca²⁺ channel binding ([³H] PN-200-110) in cardiac and aortic tissues, and on *in vivo* hemodynamic responses to BAY k 8644 were determined and compared to ovex and sham controls. Whereas high affinity

-15-

5 dihydropyridine binding sites (B_{max}) in cardiac and aortic tissues were significantly increased in EE2- and 14-treated rats compared to ovex rats, binding affinities (K_d) were not significantly different among groups in either cardiac or aortic tissues.

Treatment group		chol	cardiac	[³ H]PN200-100		*p<0.05 vs ovex	
		mg/dL	B_{max} (fmol/mg)	K_d (pM)	B_{max} (fmol/mg)	K_d (nM)	
ovex	(n=5-7)	85±7	296±51	200±19	61±15	1.0±0.3	
10 sham	(n=4-6)	87±7	385±76	188±32	46±14	2.5±0.6	
EE2	(n=5-7)	36±2*	525±65*	204±21	133±26*	2.5±0.6	
Ralox	(n=4-5)	38±4*	535±80*	171±18	124±18*	1.3±0.6	


15 Despite the increase in Ca^{2+} channel B_{max} , *in vivo* contractile, heart rate and pressor responses to BAY k 8644 in EE2-treated rats were not increased compared to those of ovex or sham controls. Thus increased Ca^{2+} channel density did not result in a more sensitized response to the calcium channel agonist.



-16-

I claim:

1. A method for modulating calcium channels in vascular and cardiac tissue comprising administering to a warm-blooded animal in need thereof a pharmaceutically-effective amount of a compound having the formula



(I)

wherein R^1 and R^3 are independently hydrogen, C_1 - C_4 alkyl, $-CO-(C_1-C_6 \text{ alkyl})$, or $-CH_2Ar$, $-CO-Ar$, wherein Ar is phenyl or substituted phenyl;

R^2 is selected from the group consisting of pyrrolidine, hexamethyleneimino, and piperidino; or a pharmaceutically-acceptable salt thereof.

2. The method of Claim 1 wherein R^1 and R^3 are independently hydrogen, C_1 - C_4 alkyl, $-CO-(C_1-C_6 \text{ alkyl})$, or benzyl; and R^2 is piperidino or pyrrolidino.

3. The method of Claim 2 wherein R^1 and R^3 are independently hydrogen or C_1 - C_4 alkyl, and R^2 is piperidino or pyrrolidino.

4. The method of Claim 3 wherein R^1 and R^3 are hydrogen and R^2 is piperidino or pyrrolidino.


5. The method of Claim 4 wherein R^2 is piperidino.

6. The method of Claim 5 wherein said pharmaceutically-acceptable salt is the hydrochloride salt.

7. The method of Claim 4 wherein R^2 is pyrrolidino.


8. The method of Claim 7 wherein said pharmaceutically-acceptable salt is the hydrochloride salt.





CLAIMS :

1. The use of a compound of formula I




wherein

R^1 and R^3 are independently hydrogen, C_1 - C_4 alkyl, $-CO-(C_1$ - C_6 alkyl), or $-CH_2Ar$, $-CO-Ar$, wherein Ar is phenyl or substituted phenyl;

R^2 is selected from the group consisting of pyrrolidine, hexamethylenemino, and piperidino; or a pharmaceutically-acceptable salt thereof; for use in modulating calcium channels in vascular and cardiac tissue which comprises administering to a warm-blooded animal in need thereof a pharmaceutically-effective amount of a compound of formula I or a pharmaceutically acceptable salt or solvate thereof.

2. The use of Claim 1 wherein R^1 and R^3 are independently hydrogen, C_1 - C_4 alkyl, $-CO-(C_1$ - C_6 alkyl), or benzyl; and R_2 is piperidino or pyrrolidino.
3. The use of Claim 2 wherein R^1 and R^3 are independently hydrogen or C_1 - C_4 alkyl, and R^2 is piperidino or pyrrolidino.
4. The use of Claim 3 wherein R^1 and R^3 are hydrogen and R^2 is piperidino or pyrrolidino.
5. The use of Claim 4 wherein R^2 is piperidino.





-17-

6. The use of Claim 5 wherein said pharmaceutically-acceptable salt is the hydrochloride salt.

7. The use of Claim 4 wherein R^2 is pyrrolidino.

8. The use of Claim 7 wherein said pharmaceutically-acceptable salt is the hydrochloride salt.



9705 890-3

JMP's copy



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 43/40, C07D 413/00, 251/58, 207/00, 405/00, 333/74, 333/64		A1	(11) International Publication Number: WO 96/39833
			(43) International Publication Date: 19 December 1996 (19.12.96)
(21) International Application Number: PCT/US96/09161		(81) Designated States: AL, AM, AT, AU, AZ, BE, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 4 June 1996 (04.06.96)			
(30) Priority Data: 08/485,317 7 June 1995 (07.06.95) US			
(60) Parent Application or Grant (63) Related by Continuation US 08/485,317 (CON) Filed on 7 June 1996 (07.06.96)		Published With international search report.	
(71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).			
(72) Inventor; and (75) Inventor/Applicant (for US only): BOWLING, Nancy, L. [US/US]; 101 White Oak Drive, Greenfield, IN 46140 (US).			
(74) Agents: STRODE, Janelle, D. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).			
(54) Title: MODULATION OF CALCIUM CHANNELS			
(57) Abstract			
A method for modulating calcium channels, increasing the density of calcium channels in vascular and cardiac tissue with no changes in inotropic or pressor response, comprising administering to a warm-blooded animal in need thereof a pharmaceutically-effective amount of a compound having formula (I) wherein R¹ and R³ are independently hydrogen, C₁-C₄ alkyl, -CO-(C₁-C₆ alkyl), or -CH₂Ar, -CO-Ar, wherein Ar is phenyl or substituted phenyl; R² is selected from the group consisting of pyrrolidine, hexamethyleneimino, and piperidino; or a pharmaceutically-acceptable salt thereof.			
(I)			