


[FIG. 1]



[FIG. 2]

SEQ ID NO: 1

```

MSALGAVIAL LLWGQLFAVD SGNDVTDIAD DGCPKPPEIA HGYVEHSVRY QCKNYYKLRT 60
EGDGVYTLND KKQWINKAVG DKLPECEADD GCPKPPEIAH GYVEHSVRYQ CKNYYKL RTE 120
GDGVYTLNNE KQWINKAVGD KLPECEAVCG KPKNPANPVQ RILGGHLD AK GSFPWQAKMV 180
SHHNLT TGAT LINEQWLLTT AKNLFLNHSE NATAKDIAPT LTLYVGKKQL VEIEKVVLHP 240
NYSQVDIGLI KLKQKVSUNE RVMPICLPSK DYAEVGRVGY VSGWGRNANF KFTDHLKYVM 300
LPVADQDQCI RHYEGSTVPE KKTPKSPGV QPILNEHTFC AGMSKYQEDT CYGDAGSAFA 360
VHDLEEDTWY ATGILSFDKS CAVA EYGVYV KVT SIQDWVQ KTIAEN 406

```

[FIG. 3]

SEQ ID NO: 2

```

MSALGAVIAL LLWGQLFAVD SGNDVTDIAD DGCPKPPEIA HGYVEHSVRY QCKNYYKLRT 60
EGDGVYTLNN EKQWINKAVG DKLPECEAVC GKPKNPANPV QRILGGHLDA KGSFPWQAKM 120
VSHHNLTTGA TLINEQWLLT TAKNLFLNHS ENATAKDIAP TLTLYVGKKQ LVEIEKVVLH 180
PNYSQVDIGL IKLKQKSVN ERVMPICLPS KDYAEVGRVG YVSGWGRNAN FKFTDHLKYV 240
MLPVADQDQC IRHYEGSTVP EKKTPKSPVG VQPILNEHTF CAGMSKYQED TCYGDAGSAF 300
AVHDLEEDTW YATGILSFDK SCAVAEYGVY VKVTSIQDWV QKTIAEN 347

```

[FIG. 4]

SEQ ID NO: 3

```

MSALGAVVAL LLWGQLFAVD TGNEATDFTD DSCPKEPEIP NGYVEHLVRY QCKNYYRLRS 60
EGDGVYALNS EKQWVNKASG TKLPECEAVC GKPKNPVDQV QRIIGGLLDA KGSFPWQAKL 120
VSRHNLTTGA TLISEQWLLT TAQNLFLNHT PDAKPKDIAP TLKLYVGRKQ PVEIEKVVFH 180
PDYQEVDIGL IKLKEKVPVG ERVMPICLPS KDYAQVGRVG YVSGWGRNAN FNTELLKYV 240
TLPVADQDTC VKHYEGSTVP EKKTNRSSVG VQPILNEHTF CAGLSKFQED TCYGDAGSAF 300
VIHDEEDDTW YAAGILSFDK SCAVAEYGVY VKVPSILDWV QKTIAEN 347

```

[FIG. 5]


SEQ ID NO: 4

```


MRALGAVVAL LLCGQLFAAE TGNEATDATD DSCPKEPEIP KGYVEHVMRY HCQTYYYKLRT 60
AGDGVYTLDS NKQWTNKVTG EKLPECEAVC GKPKNPVDQV QRIMGGSLDA KGSFPWQAKM 120
ISHHNLTSGA TLINEQWLLT TAKNLRLGHK NDTAKDIAP TLRLYVGKKQ EVEIEKVIFH 180
PDNSTVDIGL IKLKQKVPVN ERVMPICLPS KDYVNVGLVG YVSGWGRNAN LNFTEHLKYV 240
MLPVADQEK C VQYYEGSTVP EKKTPKSPVG VQPILNEHTF CAGLSKYQED TCYGDAGSAF 300
AVHDKDDDTW YAAGILSFDK SCRTAEYGVY VRVTSILDWI QTTIADN 347

```


[FIG. 6]




[FIG. 7]




[FIG. 8]



[FIG. 9]



[FIG. 10]



Relationship between free-Hb concentration and Hp addition amount in free-Hb specific measurement