5-SUBSTITUTED IMIDAZOLES

TECHNICAL FIELD


This invention relates to novel 5-substituted imidazoles or pharmaceutically-acceptable salts thereof, processes for preparing them, pharmaceutical compositions containing them and their use in therapy. This invention particularly relates to compounds that are ligands for alpha 7 nicotinic acetylcholine receptors (α7 nAChRs).

BACKGROUND OF THE INVENTION

The use of compounds which bind nicotinic acetylcholine receptors in the treatment of a range of disorders involving reduced cholinergic function, such as Alzheimer's disease, cognitive or attention disorders, anxiety, depression, smoking cessation, neuroprotection, schizophrenia, analgesia, Tourette's syndrome, and Parkinson's disease has been discussed in McDonald *et al.* (1995) "Nicotinic Acetylcholine Receptors: Molecular Biology, Chemistry and Pharmacology", Chapter 5 in Annual Reports in Medicinal Chemistry, vol. 30, pp. 41-50, Academic Press Inc., San Diego, CA; and in Williams *et al.* (1994) "Neuronal Nicotinic Acetylcholine Receptors," Drug News & Perspectives, vol. 7, pp. 205-223.

DESCRIPTION OF THE INVENTION


This invention encompasses compounds of formula I:



I

wherein:


A represents:



II III IV or V

where R represents hydrogen or methyl, and

R^1 represents a moiety of Formula VI



VI,

Ar^1 is selected from a 5- or 6-membered aromatic or heteroaromatic ring having 0, 1 or 2 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms, or selected from an 8-, 9-

or 10-membered fused aromatic or heteroaromatic ring system having 0, 1, 2 or 3 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

D is selected from hydrogen, NR^2R^3, or E-Ar^2;

wherein

E is a single bond, -O-, -S-, or -NR^3-;

Ar^2 is selected from a 5- or 6-membered aromatic or heteroaromatic ring having 0, 1 or 2 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

where each Ar^1 or Ar^2 moiety may be unsubstituted or bear 1, 2 or 3 substituents selected from -R^3, -C_1-C_6alkyl, -C_2-C_6alkenyl, -C_2-C_6alkynyl, halogen, -CN, -NO_2, -CF_3, -$S(O)_nR^3$, -NR^2R^3, -$CH_2NR^2R^3$, -OR^3, -CH_2OR^3 or -CO_2R^4;

R^2 and R^3 are independently selected at each occurrence from hydrogen, -C_1-C_4alkyl, aryl, heteroaryl, -$C(O)R^4$, -$C(O)NHR^4$, -CO_2R^4 or -SO_2R^4, or

R^2 and R^3 in combination is -$(CH_2)_jG(CH_2)_k$- wherein G is oxygen, sulfur, NR^4, or a bond;

j is 2, 3 or 4;


k is 0, 1 or 2;

n is 0, 1 or 2, and

R^4 is independently selected at each occurrence from hydrogen, -C_{1-4}alkyl, aryl, or heteroaryl.

The invention also encompasses stereoisomers, enantiomers, *in vivo*-hydrolysable precursors and pharmaceutically-acceptable salts of compounds of formula I, pharmaceutical compositions and formulations containing them, methods of using them to treat diseases and conditions either alone or in combination with other therapeutically-active compounds or substances, processes and intermediates used to prepare them, uses of them as medicaments, uses of them in the manufacture of medicaments and uses of them for diagnostic and analytic purposes.


Compounds of the invention are those according to formula I:



I

wherein:


A represents:



II III IV V

where R represents hydrogen or methyl, and

R^1 represents a moiety of Formula VI



VI,

Ar^1 is selected from a 5- or 6-membered aromatic or heteroaromatic ring having 0, 1 or 2 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms, or selected from an 8-, 9- or 10-membered fused aromatic or heteroaromatic ring system having 0, 1, 2 or 3 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

D is selected from hydrogen, NR^2R^3, or E-Ar^2;

wherein

E is a single bond, -O-, -S-, or -NR^3-;

Ar^2 is selected from a 5- or 6-membered aromatic or heteroaromatic ring having 0, 1 or 2 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

where each Ar^1 or Ar^2 moiety may be unsubstituted or bear 1, 2 or 3 substituents selected from -R^3, -C_1-C_6alkyl, -C_2-C_6alkenyl, -C_2-C_6alkynyl, halogen, -CN, -NO_2, -CF_3, -$S(O)_nR^3$, -NR^2R^3, -$CH_2NR^2R^3$, -OR^3, -CH_2OR^3 or -CO_2R^4;

R^2 and R^3 are independently selected at each occurrence from hydrogen, -C_1-C_4alkyl, aryl, heteroaryl, -$C(O)R^4$, -$C(O)NHR^4$, -CO_2R^4 or -SO_2R^4, or

R^2 and R^3 in combination is -$(CH_2)_jG(CH_2)_k$- wherein G is oxygen, sulfur, NR^4, or a bond;

j is 2, 3 or 4;

k is 0, 1 or 2;


n is 0, 1 or 2, and

R^4 is independently selected at each occurrence from hydrogen, -C_{1-4}alkyl, aryl, or heteroaryl;

and stereoisomers, enantiomers, *in vivo*-hydrolysable precursors and pharmaceutically-acceptable salts thereof.


More particular compounds are those of formula I wherein:

A represents:



II

R^1 represents hydrogen or a moiety of Formula VI



VI,

wherein:

Ar^1 is selected from a 5- or 6-membered aromatic or heteroaromatic ring having 0 or 1 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

D is selected from hydrogen, NR^2R^3, or E-Ar^2;

wherein:

E is a single bond, -O-, -S-, or -NR^3-;

Ar^2 is selected from a 5- or 6-membered aromatic or heteroaromatic ring having 0 or 1 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

where each Ar^1 or Ar^2 moiety may be unsubstituted or bear 1, 2 or 3 substituents selected from halogen, -CN, -NO_2, -CF_3, -CH_3 or -C_2H_5;

R^2 and R^3 are independently selected at each occurrence from hydrogen, -C_1.C_4alkyl, aryl, heteroaryl, or


R^2 and R^3 in combination is -$(CH_2)_jG(CH_2)_k$- wherein G is oxygen;

j is 2, 3 or 4;

k is 0, 1 or 2;

and stereoisomers, enantiomers, *in vivo*-hydrolysable precursors and pharmaceutically-acceptable salts thereof.


Particular compounds of the invention are compounds according to formula I wherein A represents:



II

and stereoisomers, enantiomers and pharmaceutically-acceptable salts thereof.

Particular compounds of the invention are R-isomers of compounds of formula I in accord with formula VII,



VII

wherein A and R^1 are as defined for compounds of formula I.

Other particular compounds of the invention are those of formula VII wherein A is of formula II and R^1 is as defined for compounds of formula I.

Most particular compounds of the invention are those of formula VII wherein A is of formula II and R^1 is as defined for compounds of formula I.

Other particular compounds of the invention include those of formula I wherein E represents a single bond; or an enantiomer thereof, and pharmaceutically-acceptable salts thereof.

Particular compounds of the invention are those of formula I wherein Ar^1 is thiophenyl having optional substituents as defined herein.

Other particular compounds of the invention encompass those of formula I wherein Ar^1 is thiophenyl and Ar^2 is selected from hydrogen, halogen, phenyl, furanyl or thiophenyl having optional substituents as defined herein.

Particular compounds of the invention are those described herein and pharmaceutically-acceptable salts thereof.

In a further aspect the invention relates to compounds according to formula I wherein one or more of the atoms is a radioisotope of the same element. In a particular form of this aspect of the invention the compound of formula I is labeled with tritium. Such radio-labeled compounds are synthesized either by incorporating radio-labeled starting materials or, in the case of tritium, exchange of hydrogen for tritium by known methods. Known methods include (1) electrophilic halogenation, followed by reduction of the halogen in the presence of a tritium source, for example, by hydrogenation with tritium gas in the presence of a palladium catalyst, or (2) exchange of hydrogen for tritium performed in the presence of tritium gas and a suitable organometallic (e.g. palladium) catalyst.

Compounds of the invention labeled with tritium are useful for the discovery of novel medicinal compounds which bind to and modulate the activity, by agonism, partial agonism, or antagonism, of the α7 nicotinic acetylcholine receptor. Such tritium-labeled compounds may be used in assays that measure the displacement of a such compounds to assess the binding of ligand that bind to α7 nicotinic acetylcholine receptors.

In another aspect the invention relates to compounds according to formula I and their use in therapy and to compositions containing them.

In another aspect the invention encompasses the use of compounds according to formula I for the therapy of diseases mediated through the action of nicotinic acetylcholine receptors. A more particular aspect of the invention relates to the use of compounds of formula I for the therapy of diseases mediated through the action of α7 nicotinic acetylcholine receptors.

Another aspect of the invention encompasses a method of treatment or prophylaxis of diseases or conditions in which activation of the α7 nicotinic receptor is beneficial which method comprises administering a therapeutically-effective amount of a compound of the invention to a subject suffering from said disease or condition.

One embodiment of this aspect of the invention is a method of treatment or prophylaxis, wherein the disorder is anxiety, schizophrenia, mania or manic depression.

Another embodiment of this aspect of the invention is a method of treatment or prophylaxis of neurological disorders, psychotic disorders or intellectual impairment disorders, which comprises administering a therapeutically effective amount of a compound of the invention.

Another embodiment of this aspect of the invention is a method of treatment or prophylaxis, wherein the disorder is Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, or Attention Deficit Hyperactivity Disorder.

Another embodiment of this aspect of the invention is a method of treatment or prophylaxis, wherein the disorder is Parkinson's disease, Huntington's disease, Tourette's syndrome, or neurodegenerative disorders in which there is loss of cholinergic synapses.

Another embodiment of this aspect of the invention is a method of treatment or prophylaxis of jetlag, nicotine addiction, craving, pain, and for ulcerative colitis, which comprises administering a therapeutically effective amount of a compound of the invention.

Yet another embodiment of this aspect of the invention is a method for inducing the cessation of smoking which comprises administering an effective amount of a compound of the invention.

Another embodiment of this aspect of the invention is a pharmaceutical composition comprising a compound of the invention and at least one pharmaceutically-acceptable excipient, diluent, lubricant or carrier.

A further aspect of the invention relates to a pharmaceutical composition useful for treating or preventing a condition or disorder mentioned herein arising from dysfunction of nicotinic acetylcholine receptor neurotransmission in a mammal, preferably a human, comprising an amount of a compound of formula I, an enantiomer thereof or a pharmaceutically-acceptable salt thereof, effective in treating or preventing such disorder or condition, and at least one pharmaceutically-acceptable excipient, diluent, lubricant or carrier.

Another embodiment of this aspect of the invention relates to use of a pharmaceutical composition of the invention for the treatment, amelioration or prophylaxis of human diseases or conditions in which activation of the α7 nicotinic receptor is beneficial.

Another embodiment of this aspect of the invention is the use of the pharmaceutical composition of the invention for the treatment or prophylaxis of neurological disorders, psychotic disorders or intellectual impairment disorders.

Another embodiment of this aspect of the invention is the use of the pharmaceutical composition of the invention for the treatment or prophylaxis of Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, Attention Deficit Hyperactivity Disorder, anxiety, schizophrenia, or mania or manic depression, Parkinson's disease, Huntington's disease, Tourette's syndrome, neurodegenerative disorders in which there is loss of cholinergic synapse, jetlag, cessation of smoking, nicotine addiction including that resulting from exposure to products containing nicotine, craving, pain, and for ulcerative colitis.

A further aspect of the invention is the use of a compound according to the invention, an enantiomer thereof or a pharmaceutically-acceptable salt thereof, in the manufacture of a medicament for the treatment or prophylaxis of the diseases or conditions mentioned herein.

Another embodiment of this aspect of the invention is the use of a compound of the invention in the manufacture of a medicament for the treatment or prophylaxis of human diseases or conditions in which activation of the α7 nicotinic receptor is beneficial.

Another embodiment of this aspect of the invention is the use of a compound of the invention in the manufacture of a medicament for the treatment or prophylaxis of neurological disorders, psychotic disorders or intellectual impairment disorders.

Another embodiment of this aspect of the invention is the use of a compound of the invention in the manufacture of a medicament for treatment or prophylaxis of Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss or Attention Deficit Hyperactivity Disorder.

Another embodiment of this aspect of the invention is the use of a compound of the invention in the manufacture of a medicament for treatment or prophylaxis of anxiety, schizophrenia, or mania or manic depression.

Another embodiment of this aspect of the invention is the use of a compound of the invention in the manufacture of a medicament for treatment or prophylaxis of Parkinson's disease, Huntington's disease, Tourette's syndrome, or neurodegenerative disorders in which there is loss of cholinergic synapses.

Another embodiment of this aspect of the invention is the use of a compound as described above in the manufacture of a medicament for the treatment or prophylaxis of jetlag, pain, or ulcerative colitis.

Another aspect of the invention relates to the use of a compound of the invention in the manufacture of a medicament for facilitating the cessation of smoking or the treatment of nicotine addiction or craving including that resulting from exposure to products containing nicotine.

For the uses, methods, medicaments and compositions mentioned herein the amount of compound used and the dosage administered will, of course, vary with the compound employed, the mode of administration and the treatment desired. However, in general, satisfactory results are obtained when the compounds of the invention are administered at a daily dosage of from about 0.1 mg to about 20 mg/kg of animal body weight. Such doses may be given in divided doses 1 to 4 times a day or in sustained release form. For man, the total daily dose is in the range of from 5 mg to 1,400 mg, more preferably from 10 mg to 100 mg, and unit dosage forms suitable for oral administration comprise from 2 mg to 1,400 mg of the compound admixed with a solid or liquid pharmaceutical carriers, lubricants and diluents.

The compounds of formula I, an enantiomer thereof, and pharmaceutically-acceptable salts thereof, may be used on their own or in the form of appropriate medicinal preparations for enteral or parenteral administration. According to a further aspect of the invention, there is provided a pharmaceutical composition including preferably less than 80% and more preferably less than 50% by weight of a compound of the invention in admixture with an inert pharmaceutically-acceptable diluent, lubricant or carrier.

Examples of diluents, lubricants and carriers are:

– for tablets and dragees: lactose, starch, talc, stearic acid;

– for capsules: tartaric acid or lactose;

– for injectable solutions: water, alcohols, glycerin, vegetable oils;

– for suppositories: natural or hardened oils or waxes.

There is also provided a process for the preparation of such a pharmaceutical composition which process comprises mixing the ingredients.

Compounds according to the invention are agonists of nicotinic acetylcholine receptors. While not being limited by theory, it is believed that agonists of the α7 nicotinic acetylcholine receptor (nAChR) subtype are useful in the treatment or prophylaxis of neurological disorders, psychotic disorders and intellectual impairment disorders, and to have advantages over compounds which are or are also agonists of the α4 nAChR subtype. Therefore, compounds which are selective for the α7 nAChR subtype are preferred. The compounds of the invention are indicated as pharmaceuticals, in particular in the treatment or prophylaxis of neurological disorders, psychotic disorders and intellectual impairment disorders. Examples of psychotic disorders include schizophrenia, mania and manic depression, and anxiety. Examples of intellectual impairment disorders include Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, and Attention Deficit Hyperactivity Disorder. The compounds of the invention may also be useful as analgesics in the treatment of pain, chronic pain, and in the treatment or prophylaxis of Parkinson's disease, Huntington's disease, Tourette's syndrome, and neurodegenerative disorders in which there is loss of cholinergic synapses.

Compounds of the invention may further be useful for the treatment or prophylaxis of jetlag, for use in inducing the cessation of smoking, craving, and for the treatment or prophylaxis of nicotine addiction including that resulting from exposure to products containing nicotine.

It is also believed that compounds according to the invention are useful in the treatment and prophylaxis of ulcerative colitis.

The compounds of the invention have the advantage that they may be less toxic, be more efficacious, be longer acting, have a broader range of activity, be more potent, produce fewer side effects, are more easily absorbed or have other useful pharmacological properties when compared to known therapeutic moieties.


The compounds of formula I exist in tautomeric or enantiomeric forms, all of which are included within the scope of the invention. The various optical isomers may be isolated by separation of a racemic mixture of the compounds using conventional techniques, e.g. fractional crystallization, or chiral HPLC. Alternatively the individual enantiomers may be

made by reaction of the appropriate optically active starting materials under reaction conditions which will not cause racemization.


Methods of Preparation

Compounds of the invention may be prepared according to the synthetic schemes that follow, wherein A, D, E, Ar^1, and Ar^2 unless otherwise indicated, are as defined herein.


Method A –




Method B -




Method C –






Method D –



Method E –



General Experimental Procedures and Definitions

Commercial reagents were used without further purification. Mass spectra were recorded using either a Hewlett Packard 5988A or a MicroMass Quattro-1 Mass Spectrometer and are reported as *m/z* for the parent molecular ion. Room temperature refers to 20–25 °C.

SiO_2 chromatography was performed with an Isco CombiFlash Sq 16x instrument and pre-packaged disposable *RediSep* SiO_2 stationary phase columns (4, 12, 40, 120 gram sizes) with gradient elution at 5-125 mL/min of selected bi-solvent mixture, UV detection (190-760 nm range) or timed collection, 0.1mm flow cell path length.

Microwave heating was achieved with a Personal Chemistry Smith Synthesizer or a Personal Chemistry Emrys Optimizer (monomodal, 2.45 GHz, 300W max).

Supercritical Fluid Chromatography (SFC) was performed as a means of purification for selected compounds and intermediates.

Reverse Phase High Pressure Liquid Chromatography (RP-HPLC) was employed as a method of purification for selected compounds.

LC/MS HPLC method was generally performed with a Agilent Zorbax 5μ SB-C8 column 2.1mm x 5 cm. Solvents: A = H_2O with 0.05% TFA, B =10% H_2O, 90% Acetonitrile, 0.05% TFA. Gradient: (10-90%B over 3 min., 90% B hold through 4 min., -10% B at 5 min. and hold at 10% B until 6 min).

Unless otherwise indicated, halo includes chloro, bromo, fluoro and iodo and halogen refers to fluorine, chlorine, bromine, or iodine; C_{1-6}alkyl includes methyl, ethyl and linear, cyclic or branched propyl, butyl, pentyl or hexyl; C_{2-6}alkenyl includes ethenyl, 1-propenyl, 2-propenyl or 3-propenyl and linear, branched or cyclic butenyl, pentenyl or hexenyl; C_{2-6}alkynyl includes ethynyl or propynyl; the C_{1-4}alkyl groups referred to herein, e.g., methyl, ethyl, n-propyl, n-butyl, i-propyl, i-butyl, t-butyl, s-butyl, whether alone or part of another group, may be straight-chained or branched, and the C_{3-4} alkyl groups may also be cyclic, e.g., cyclopropyl, cyclobutyl. Alkyl groups referred to herein may optionally have one, two or three halogen atoms substituted thereon.

Pharmaceutically-acceptable derivatives include solvates and salts. For example, the compounds of formula I can form acid addition salts with acids, such as the conventional pharmaceutically-acceptable acids, for example, maleic, hydrochloric, hydrobromic, phosphoric, acetic, fumaric, salicylic, citric, lactic, mandelic, tartaric and methanesulfonic acids.

PHARMACOLOGY

The pharmacological activity of the compounds of the invention may be measured in the tests set out below:

Test A - Assay for affinity at α_7 nAChR subtype

^{125}I-α -Bungarotoxin (BTX) binding to rat hippocampal membranes. Rat hippocampi are homogenized in 20 volumes of cold homogenization buffer (HB: concentrations of constituents (mM): tris(hydroxymethyl)aminomethane 50; $MgCl_2$ 1; NaCl 120; KCl 5: pH 7.4). The homogenate is centrifuged for 5 minutes at 1000 g, the supernatant is saved and the pellet re-extracted. The pooled supernatants are centrifuged for 20 minutes at 12,000 g, washed, and re-suspended in HB. Membranes (30–80 μg) are incubated with 5 nM [^{125}I] α -BTX, 1 mg/mL BSA (bovine serum albumin), test drug, and either 2 mM CaCl2 or 0.5 mM EGTA [ethylene glycol-bis(β -aminoethylether)] for 2 hours at 21 °C, and then filtered and washed 4 times over Whatman glass fibre filters (thickness C) using a Brandel cell harvester. Pre-treating the filters for 3 hours with 1% (BSA/0.01% PEI (polyethyleneimine) in water is critical for low filter blanks (0.07% of total counts per minute). Non-specific binding is described by 100 μM (–)-nicotine, and specific binding is typically 75%.

Test B - Assay for affinity to the α_4 nAChR subtype

[^{3}H] -(–)-nicotine binding. Using a procedure modified from Martino-Barrows and Kellar (Mol Pharm (1987) 31:169-174), rat brain (cortex and hippocampus) is homogenised as in the [^{125}I] α -BTX binding assay, centrifuged for 20 minutes at 12,000 x g, washed twice, and then re-suspended in HB containing 100 μ M diisopropyl fluorophosphate. After 20 minutes at 4 °C, membranes (approximately 0.5 mg) are incubated with 3 nM [^{3}H] -(–)-nicotine, test drug, 1 μ M atropine, and either 2 mM $CaCl_2$ or 0.5 mM EGTA for 1 hour at 4 °C, and then filtered over Whatman glass fibre filters (thickness C) (pre-treated for 1 hour with 0.5% PEI) using a Brandel cell harvester. Non-specific binding is described by 100 μ M carbachol, and specific binding is typically 84%.

Binding data analysis for Tests A and B

IC_{50} values and pseudo Hill coefficients (n_H) are calculated using the non-linear curve fitting program ALLFIT (DeLean A, Munson P J and Rodbard D (1977) Am. J. Physiol., 235:E97-E102). Saturation curves are fitted to a one site model, using the non-linear regression program ENZFITTER (Leatherbarrow, R.J. (1987)), yielding K_D values of 1.67 and 1.70 nM for the [^{125}I]-α -BTX and [^{3}H] -(–)-nicotine ligands respectively. K_i values are estimated using the general Cheng-Prusoff equation:

$$K_i=[\ IC_{50}]\ /((2+([\text{ligand}]\ /K_D]\ )^n)^{1/n}-1)$$


where a value of n=1 is used whenever $n_H< 1.5$ and a value of n=2 is used when $n_H\geq 1.5$. Samples are assayed in triplicate and are typically ± 5%. K_i values are determined using 6 or more drug concentrations.

Compounds of the invention generally have binding affinities (K_i) of less than 2 μM in either Test A and or Test B.

EXAMPLES

The following examples are non-limiting and embody particular aspects of the invention.

Example 1: 3-Imidazol-1-yl-1-azabicyclo[2.2.2]octane.



a) 3-Imidazol-1-yl-1-azabicyclo[2.2.2]oct-2-ene.


To a solution of imidazole (6.53 g, 96 mmol) in anhydrous CH_2Cl_2 (50 mL) at 0 °C under a nitrogen atmosphere was added thionyl chloride (2.1 mL, 28.8 mmol) dropwise, resulting in a fine white precipitate. After 15 minutes the mixture was added to 3-quinuclidone hydrochloride salt (2.59 g, 16 mmol) in anhydrous CH_2Cl_2 (50 mL) under an inert atmosphere at 0 °C. Allowed to warm to ambient temperature, stirred for 2 days, then neutralized with saturated $NaHCO_3$ (aq). Adjusted to pH ~ 12 by the addition of 10% NaOH (aq) and extracted into CH_2Cl_2 (3x 100 mL). The combined organic extracts were dried over Na_2SO_4, filtered and concentrated *in vacuo*. The residue was subjected to flash chromatography (SiO_2 - 40 grams; gradient elution: 1-10% 3N NH_3/MeOH - CH_2Cl_2 at 50 mL/min over 40min) to afford the title compound (1a) as an orange-yellow tacky solid (640 mg, 23%). Mass spectrum (ES+): M + 1: 176 (100%); ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.72 (m,

2H), 1.79 (m, 2H), 2.69 (m, 2H), 3.02 (m, 2H), 3.26 (m, 1H), 6.53 (d, J = 2Hz, 1H), 7.11 (s, 1H), 7.15 (s, 1H), 7.71 (s, 1H).

b) 3-Imidazol-1-yl-1-azabicyclo[2.2.2]octane.

To a mixture of 3-imidazol-1-yl-1-azabicyclo[2.2.2]oct-2-ene (1a) (356 mg, 2.03 mmol) and ammonium formate (2.6 g, 40.6 mmol) in absolute EtOH (6 mL) was added 10% Pd/C (178mg, 50% w/w). The sealed reaction vessel was heated to 110 °C for 13 hrs before cooling to ambient temperature. The mixture was filtered through diatomaceous earth (0.75" by 2" diameter), washed liberally with EtOH and concentrated to a white-yellow tacky residue which was subsequently purified by flash chromatography (SiO_2 - 40 grams; gradient elution: 5-30% 3N NH_3/MeOH - EtOAc at 60 mL/min over 25min) to yield the title compound (188 mg, 52%). Mass spectrum (ES+): M + 1: 178 (100%); ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.50 (m, 1H), 1.64 (m, 1H), 1.77 (m, 2H), 2.09 (app q, J = 3 Hz, 1H), 2.81-3.05 (m, 4H), 3.15 (dd, J = 14.5, 4.4 Hz, 1H), 3.47 (ddd, J = 14.5, 9.7, 1.8 Hz, 1H), 4.26 (m, 1H), 7.05 (s, 1H), 7.08 (s, 1H), 7.62 (s, 1H).

Example 2: (R)-3-[5-(5-bromo-thiophen-2-yl)-imidazol-1-yl]-1-azabicyclo[2.2.2]octane.




a) 1-(5-Bromo-thiophen-2-yl)-2,2-dihydroxyethanone.

A mixture of selenium dioxide (8.66 g, 78 mmol), water (2.8 mL) and 1,4-dioxane (75 mL) was heated to reflux to dissolve all particulates (~15 min) before the addition of 2-acetyl-5-bromothiophene (8 g, 39 mmol) in one portion. After refluxing for 15 hrs, the mixture was cooled to ambient temperature, filtered through diatomaceous earth (1.5" by 3" diameter), washed liberally with Et_2O, and concentrated *in vacuo* to a yellow residue. The residue was subjected to Kugelrohr distillation (150 °C / 1 Torr) to yield a glyoxal as a yellow oil, which was immediately added to boiling water. Recrystallization from water, filtration, and drying (37 °C and 5 Torr) yielded the glyoxal hydrate, 1-(5-bromo-thiophen-2-yl)-2,2-dihydroxyethanone, as white-pink needles (6.723 g, 73%). ^{1}H NMR (300 MHz, DMSO-d_6) δ 5.43 (t, J = 6.1 Hz, 1H), 6.93 (d, J = 6.1 Hz, 2H), 7.37 (d, J = 4.1 Hz, 1H), 7.81 (d, J = 4.1 Hz, 1H); ^{13}C NMR (300 MHz, DMSO-d_6) δ 90.28, 122.32, 132.23, 135.73, 141.15, 189.6.

b) (R)-3-[5-(5-bromo-thiophen-2-yl)-imidazol-1-yl]-1-azabicyclo[2.2.2]octane.

Five individual round-bottomed thick-walled glass vials were charged with stir bars, (R)-(+)-3-aminoquinuclidine dihydrochloride (440 mg, 2.21 mmol), ammonium acetate (195 mg, 2.53 mmol), 1-(5-bromo-thiophen-2-yl)-2,2-dihydroxyethanone (500 mg, 2.21 mmol), formaldehyde (37%, aqueous – 210 μL, 2.53 mmol) and glacial AcOH/water (1:1, 3.5 mL). All vials were crimp sealed and subjected to microwave radiation[2] for 20 minutes at 120 °C. The resultant mixtures were combined, diluted with water, basified to pH ~ 13 with 20% NaOH (aq), and extracted with $CHCl_3$ (5 x 60 mL) and EtOAc (2 x 40 mL). The combined organic extracts were dried over $MgSO_4$, filtered, and concentrated by rotary evaporation to a brown gummy residue. The material was subjected to flash chromatography (SiO_2 – 120 grams; gradient elution: 5-10% 4N NH_3/MeOH – EtOAc at 90 mL/min over 48min) to provide a ~2:1 regio-isomeric mixture of 5- and 4-bromothienyl-imidazolyl compounds as an orange-tan gum (1.186 g, 33%) as determined by ^1H NMR signal integration. The mixture was subsequently purified by SFC with 25% MeOH/CO_2 with 0.5% DMEA at 50 mL/min, monitoring at 280 nm, to afford the title compound as a pale orange solid (700 mg, 20%). Mass spectrum (API+) M + 1: 338 (100%); ^1H NMR (300 MHz, $CDCl_3$) δ 1.48-1.80 (m, 4H), 1.96 (app q, J = 2.9 Hz, 1H), 2.74-3.16 (m, 5H), 3.37 (ddd, J = 14.5, 10, 2.2 Hz, 1H), 4.36 (m, 1H), 6.80 (d, J = 4 Hz, 1H), 7.06 (d, J = 4 Hz, 1H), 7.15 (s, 1H), 7.84 (s, 1H).


Example 3: 3-[(R)-5-(5-Furan-3-yl-thiophen-2-yl)-imidazol-1-yl]-1-azabicyclo[2.2.2]octane.



A conical thick-walled glass vial was charged with a stir bar, a 7:1 mixture based on integration of NMR signals of (R)-3-[4-(5-bromo-thiophen-2-yl)-imidazol-1-yl]-1-azabicyclo[2.2.2]octane (Example 6) and (R)-3-[5-(5-bromo-thiophen-2-yl)-imidazol-1-yl]-1-azabicyclo[2.2.2]octane (Example 7) (65 mg, 0.19 mmol), 3-furanboronic acid (62 mg, 0.23 mmol), dichlorobis(triphenylphosphine)-palladium (II) (7 mg, 0.0095 mmol), Cs_2CO_3 (124 mg, 0.38 mmol) and DME/H_2O/EtOH (7:3:2 - 1 mL). The vial was sealed under an argon purge and subjected to microwave radiation for 18 minutes at 150 °C. The resultant black slurry was filtered through a 0.7 μm GMF filter, washing with MeOH (1 x 1 mL), dilute with DMSO (1 x 1 mL), subjected to RP-HPLC purification (gradient elution: 5-45% acetonitrile

over 45 minutes at 24 mL/min - desired compound eluted at 15.4 minutes). The product-containing fractions were concentrated via centrifugal evaporation to yield the title compound as the bis-trifluoroacetic acid salt (67 mg, 64%). Mass spectrum (API+) M + 1: 326 (100%); ^{1}H NMR (300 MHz, MeOH-d_4) δ 2.04-2.23 (m, 4H), 2.59 (app d, J = 2.5 Hz, 1H), 3.34-3.47 (m, 3H), 3.62 (m, 1H), 3.74-3.94 (m, 2H), 5.22 (app t, J = 7.5 Hz, 1H), 6.76 (s, 1H), 7.35 (d, J = 4 Hz, 1H), 7.39 (d, J = 4 Hz, 1H), 7.60 (t, J = 1.8 Hz, 1H), 7.84 (d, J = 1 Hz, 1H), 7.94 (s, 1H), 9.54 (s, 1H).

Example 4: (R)-3-(5-Thiophen-2-yl-imidazol-1-yl)-1-azabicyclo[2.2.2]octane.



From the process described in Example 3, (R)-3-(5-thiophen-2-yl-imidazol-1-yl)-1-azabicyclo[2.2.2]octane was obtained as a by-product. The title compound was purified by RP-HPLC (eluted at 7.6 minutes) to afford the bis-trifluoroacetic acid salt as a clear, yellow gum. Mass spectrum (API+) M + 1: 260 (100%); ^{1}H NMR (300 MHz, MeOH-d_4) δ 1.98-2.10 (m, 3H), 2.14 (m, 1H), 2.55 (app t, J = 3 Hz, 1 H), 3.35-3.46 (m, 3H), 3.59 (m, 1H), 3.70–3.87 (m, 2H), 5.15 (app t, J = 4.5 Hz, 1H), 7.29 (dd, J = 4.0, 3.5 Hz, 1H), 7.47 (dd, J = 3.5, 1.0 Hz, 1H), 7.82 (m, 2H), 9.50 (s, 1H).

Compounds of the following structure prepared by methods analogous to those used in Example 3 are illustrated in Table 1:










Table 1

Example #	R^1	LC/MS retention time (min)	MS (M + 1) ion
4		0.30	260
5		1.61	336
6		0.33	337
7		1.54	342
8		1.51	342

CLAIMS


1. A compound according to formula I:



I

wherein:


A represents:



II III IV or V

where R represents hydrogen or methyl, and

R^1 represents hydrogen or a moiety of Formula VI



VI,

wherein:

Ar^1 is selected from a 5- or 6-membered aromatic or heteroaromatic ring having 0, 1 or 2 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms, or selected from an 8-, 9- or 10-membered fused aromatic or heteroaromatic ring system having 0, 1, 2 or 3 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

D is selected from hydrogen, NR^2R^3, or E-Ar^2;

wherein

E is a single bond, -O-, -S-, or -NR^3-;

Ar^2 is selected from a 5- or 6-membered aromatic or heteroaromatic ring having 0, 1 or 2 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

where each Ar^1 or Ar^2 moiety may be unsubstituted or bear 1, 2 or 3 substituents selected from -R^3, -C_1-C_6alkyl, -C_2-C_6alkenyl, -C_2-C_6alkynyl, halogen, -CN, -NO_2, -CF_3, -$S(O)_nR^3$, -NR^2R^3, -$CH_2NR^2R^3$, -OR^3, -CH_2OR^3 or -CO_2R^4;

R^2 and R^3 are independently selected at each occurrence from hydrogen, -C_1-C_4alkyl, aryl, heteroaryl, -$C(O)R^4$, -$C(O)NHR^4$, -CO_2R^4 or -SO_2R^4, or

R^2 and R^3 in combination is -$(CH_2)_jG(CH_2)_k$- wherein G is oxygen, sulfur, NR^4, or a bond;

j is 2, 3 or 4;

k is 0, 1 or 2;


n is 0, 1 or 2, and

R^4 is independently selected at each occurrence from hydrogen, -C_{1-4}alkyl, aryl, or heteroaryl;

and stereoisomers, enantiomers, *in vivo*-hydrolysable precursors and pharmaceutically-acceptable salts thereof.


2. A compound according to Claim 1 wherein:

A represents:



II

R^1 represents hydrogen or a moiety of Formula VI



VI,

wherein:

Ar^1 is selected from a 5- or 6-membered aromatic or heteroaromatic ring having 0 or 1 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

D is selected from hydrogen, NR^2R^3, or E-Ar^2;

wherein:

E is a single bond, -O-, -S-, or -NR^3-;

Ar^2 is selected from a 5- or 6-membered aromatic or heteroaromatic ring having 0 or 1 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

where each Ar^1 or Ar^2 moiety may be unsubstituted or bear 1, 2 or 3 substituents selected from halogen, -CN, -NO_2, -CF_3, -CH_3 or -C_2H_5;

R^2 and R^3 are independently selected at each occurrence from hydrogen, -C_1-C_4alkyl, aryl, heteroaryl, or


R^2 and R^3 in combination is -$(CH_2)_jG(CH_2)_k$- wherein G is oxygen;

j is 2, 3 or 4;

k is 0, 1 or 2;

and stereoisomers, enantiomers, *in vivo*-hydrolysable precursors and pharmaceutically-acceptable salts thereof.


3. A compound according to Claim 1, wherein A represents:



II

or an enantiomer thereof, and pharmaceutically-acceptable salts thereof.


4. A compound according to Claim 1, wherein said compound is an R-isomer in accord with formula VII,



VII

wherein A and R^1 are as defined for compounds of formula I.

5. A compound according to Claim 4, wherein A is of formula II



II

and R^1 is as defined for compounds of formula I.

6. A compound according to Claim 1, wherein E represents a single bond; or an enantiomer thereof, and pharmaceutically-acceptable salts thereof.

7. A compound according to Claim 1, wherein Ar^1 is thiophenyl.

8. A compound according to Claim 1, wherein Ar^1 is thiophenyl and Ar^2 is selected from hydrogen, halogen, phenyl, furanyl or thiophenyl having optional substituents.

9. A compound according to Claim 1, wherein one or more of the atoms is a radioisotope of the same element.

10. A method of treatment or prophylaxis of a disease or condition in which activation of the α7 nicotinic receptor is beneficial which method comprises administering a therapeutically-effective amount of a compound according to Claim 1 to a subject suffering from said disease or condition.

11. The method of Claim 10, wherein said disease or condition is anxiety, schizophrenia, mania or manic depression.

12. A method of treatment or prophylaxis of neurological disorders, psychotic disorders or intellectual impairment disorders, which comprises administering a therapeutically effective amount of a compound according to Claim 1.

13. The method of Claim 12, wherein said disorder is Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, Attention Deficit Hyperactivity Disorder, Parkinson's disease, Huntington's disease, Tourette's syndrome, neurodegenerative disorders in which there is loss of cholinergic synapses, jetlag, nicotine addiction, craving, pain, and for ulcerative colitis, which comprises administering a therapeutically effective amount of a compound according to Claim 1.

14. A method for inducing the cessation of smoking comprising administering an effective amount of a compound according to Claim 1.

15. A pharmaceutical composition comprising a compound according to Claim 1 and a pharmaceutically-acceptable diluent, lubricant or carrier.

16. A method of treatment or prophylaxis of a disease or condition in which activation of the α7 nicotinic receptor is beneficial which method comprises administering a therapeutically-effective amount of a pharmaceutical composition according to Claim 15 to a subject suffering from said disease or condition.

17. The method of Claim 16, wherein said disease or condition is anxiety, schizophrenia, mania or manic depression.

18. A method of treatment or prophylaxis of neurological disorders, psychotic disorders or intellectual impairment disorders, which comprises administering a therapeutically effective amount of a pharmaceutical composition according to Claim 15.

19. The method of Claim 12, wherein said disorder is Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, Attention Deficit Hyperactivity Disorder, Parkinson's disease, Huntington's disease, Tourette's syndrome, neurodegenerative disorders in which there is loss of cholinergic synapses, jetlag, nicotine addiction, craving, pain, and for ulcerative colitis, which comprises administering a therapeutically effective amount of a pharmaceutical composition according to Claim 15.

20. A method for inducing the cessation of smoking comprising administering an effective amount of a pharmaceutical composition according to Claim 15.

21. The use of a compound according to Claim 1, an enantiomer thereof or a pharmaceutically-acceptable salt thereof, in the manufacture of a medicament for the treatment or prophylaxis of human diseases or conditions in which activation of the $\alpha 7$ nicotinic receptor is beneficial selected from neurological disorders, psychotic disorders, intellectual impairment disorders, Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, Attention Deficit Hyperactivity Disorder, anxiety, schizophrenia, mania or manic depression, Parkinson's disease, Huntington's disease, Tourette's syndrome, or neurodegenerative disorders in which there is loss of cholinergic synapses.

22. The use of a compound according to Claim 1, in the manufacture of a medicament for the treatment or prophylaxis of jetlag, pain, or ulcerative colitis or to facilitate the cessation of smoking or the treatment of nicotine addiction or craving including that resulting from exposure to products containing nicotine.