

SELECTIVE REDUCTION OF CYSTEINE RESIDUES IN IL-17 ANTIBODIES

RELATED APPLICATIONS

This application claims priority to U.S. Provisional Patent Application No. 62/095,361, filed on December 22, 2014, which is incorporated by reference herein in its entirety.

TECHNICAL FIELD

The present disclosure relates to methods for selectively reducing CysL97 in a preparation of IL-17 antibodies or antigen binding fragments thereof, e.g., a preparation of secukinumab, that have been recombinantly produced by mammalian cells.

BACKGROUND OF THE DISCLOSURE

Classical antibodies are composed of two light chains (L) with a molecular weight of about 25kD each and two heavy chains (H) with a molecular weight of about 50kD each. The light and heavy chains are connected by a disulfide bond (L-S-S-H) and the two LH units are further linked between the heavy chains by two disulfide bonds. The general formula of a classical antibody is L-SS-H(-SS-)₂H-SS-L or simply H₂L₂ (HHLL). Besides these conserved inter-chain disulfide bonds, there are also conserved intra-chain disulfide. Both types of disulfide bonds are important for the stability and behavior (e.g., affinity) of an antibody. Generally, a disulfide bond is produced by two cysteine residues (Cys-SH) found at conserved positions in the antibody chains, which spontaneously form the disulfide bond (Cys-S-S-Cys). Disulfide bonds formation is determined by the redox potential of the environment and by the presence of enzymes specialized in thiol-disulfide exchange. The internal disulfide bonds (Cys-S-S-Cys) stabilize the three-dimensional structure of an antibody.

There are unusual antibodies that contain an additional free cysteine(s) (i.e., unpaired cysteine) that is involved in antigen recognition and binding. For these antibodies, modification of a free cysteine can have a negative effect on the activity and stability of the molecule, and can

lead to increased immunogenicity. As a result, processing of these antibodies can be difficult, as the end product may contain a substantial amount of inactive, misfolded and useless antibody material. US20090280131, which is incorporated by reference herein in its entirety, provides IL-17 antibodies, e.g., secukinumab (i.e., AIN457) with a free cysteine residue after the cis-proline in the light chain complementarity determining region (CDR) 3 loop (L-CDR3) (i.e., amino acid eight of L-CDR3 as set forth as SEQ ID NO:6, which corresponds to amino acid 97 of the light chain variable region as set forth as SEQ ID NO:10, herein after referred to as "CysL97"). In order to maintain full activity, the unpaired cysteine residue of secukinuamb cannot be masked by oxidative disulfide pairing with other cysteine residues or by oxidation with exogenous compounds (e.g., formation of mixed disulfides with other proteins, derivatization with cell metabolites [e.g., cysteine or glutathione], and formation of sulfoxides by oxygen). Unfortunately, because secukinuamb is manufactured using mammalian cells, undesired cell-based modifications of CysL97 do occur.

The literature describes refolding of mammalian proteins expressed in bacterial cells, which produce mammalian proteins as unfolded, insoluble aggregates having mixed disulfides (inclusion bodies). To obtain mammalian proteins from bacteria, inclusion body proteins are isolated, solubilized, and denatured with strong chaotropic reagents and reducing agents. Complete denaturation and reduction of disulfide bonds using a denaturing agent, reducing agent, disulfide adduct forming agent, and a mild oxidizing/reducing environment (pH 7-9) has also been used to properly refold plant proteins obtained from commercial sources or recombinantly produced in yeast (US 4,766,205). These processes, which employ complete denaturation and refolding of proteins, are expensive, caustic, time-consuming, and unnecessary for a protein produced in mammalian cells.

The use of reduction/oxidation coupling reagents to correct misfolding of non-naturally occurring Fc fusion proteins is known (WO02/68455). The Fc fusion proteins of WO02/68455 presumably contain interchain disulfides in the Fc region that are reduced and reoxidized by the disclosed process, but there is no teaching therein of how to produce a molecule having a selectively reduced cysteine residue. Moreover, an Fc fusion protein is simply not an antibody, a highly complex immunoglobulin that relies on numerous properly linked inter-chain and intra-chain disulfides for structure and activity.


US20050123532 provides methods of producing an antibody having a free cysteine by activating the antibody with a reducing agent, or by culturing the antibody-producing cells in a serum-free medium supplemented with L-cysteine. When using this cell-culture method, later processing steps, e.g., filtration, viral inactivation and chromatography, could lead to oxidation of the free cysteine produced by cell-culture methods. In such cases, the free cysteine is ideally protected during later steps by modifying the free thiol group with an oxidizing agent, which is itself later removed using various techniques, e.g., filtration (US20060257393). For commercial production, such methods require large quantities of reducing agent in the original culture medium, large quantities of an oxidizing agent during later processing, and additional filtration methods to remove the oxidizing agent, adding time and expense to the cost of production.

U.S. Patent No. 7,928,205 teaches a preference for using redox pairs for refolding IgG₂ antibodies obtained from mammalian cell cultures, as well as methods for decysteinylation of a free cysteine in the variable region of the 146B7 (IgG₁) antibody. The corresponding research publication, Banks et al. (2008) J. Pharmaceutical Sci. 97:764-779, teaches decysteinylation of an unpaired sulfhydryl in the variable region of a recombinant monoclonal IgG₁ antibody (MAB007). Banks et al. studied whether decysteinylation of MAB007 required the use of a strong denaturant (GdnHCL) and a reducing agent (cysteine) or whether selective reduction could occur in the presence of cysteine alone. The authors determined that cysteinylation was effectively removed from MAB007 in the presence and absence of denaturant.

None of the above references teach whether selective reduction of CysL97 in secukinumab is possible. Nor do the above references teach the reagents and conditions necessary for selective reduction of CysL97 in secukinumab, which depend upon, *inter alia*, the primary, secondary and tertiary structure of secukinumab; the position and location of oxidized CysL97 in secukinumab (e.g., solvent-accessible or inaccessible); and the relative strength of the conserved disulfide bonds in the antibody (e.g., whether CysL97 reacts first with a given reducing agent, or only after conserved cysteines have been reduced). Moreover, none of these references describe whether selective reduction of oxidized CysL97 in secukinumab would result in changes to the antibody structure (e.g., folding), chemical composition (e.g., deamidation), or properties (binding activity, propensity to aggregate or degrade), all of which could make it technically unfeasible/impractical to selectively reduce secukinumab at commercial scale.

SUMMARY OF THE DISCLOSURE

In order to maintain maximal secukinumab antigen-binding activity, we have determined that it is necessary during processing of secukinumab to convert CysL97 from masked (1) to free (2) form (see I, below) without significant reduction of the conserved disulfide bonds; otherwise lower activity and inactive lower molecular weight variants will form by chain unlinking ($H_2L_2 \rightarrow H_2L, HL_2, HL, H$ and L).



Introducing reducing conditions during commercial scale antibody preparation is counterintuitive (see, e.g., Trexler-Schmidt et al. 2010 Biotech and Bioengineering 106:452-61, which employs various reagents and methods to prevent antibody disulfide bond reduction during cell culture manufacturing of antibodies). Nevertheless, we have determined that it is possible to selectively reduce CysL97 in secukinumab during large scale commercial production in mammalian cells without significant denaturation of the antibody. Disclosed herein are methods for selectively reducing CysL97 in the antigen binding sites of the IL-17 antibodies (and fragments thereof) disclosed in US20090280131, particularly secukinumab. These methods assist in restoring the binding activity of these antibodies, and thus increase the bioactivity of preparations thereof. Furthermore, these methods assist in increasing the level of intact antibody and enhancing the homogeneity of preparations of these antibodies. The disclosed processes rely on the combined effect of particular ratios of antibody:reductant and controlled oxygen transfer rates in the system during incubation.

Accordingly, disclosed herein are methods for selectively reducing CysL97 in a preparation of IL-17 antibodies that have been recombinantly produced by mammalian cells, comprising:

- a) contacting the preparation with at least one reducing agent in a system to form a reducing mixture; and
- b) incubating the reducing mixture while maintaining a volumetric oxygen mass-transfer coefficient (k_{La}^*) in the system of $\leq$ about 0.37 h^{-1} , said k_{La}^* being calculated by adapting a dissolved oxygen curve to a saturation curve;

wherein the IL-17 antibodies each comprise an immunoglobulin heavy chain variable domain (V_H) comprising the three complementarity determining regions (CDRs) of the V_H set forth as

SEQ ID NO:8 and an immunoglobulin light chain variable domain (V_L) comprising the three CDRs of the V_L set forth as SEQ ID NO:10, and further wherein prior to step a) the initial percent oxygen saturation in the preparation is at least about 60%, as measured using an oxygen probe calibrated at 25°C.

Also disclosed herein are methods for selectively reducing CysL97 in a preparation of IL-17 antibodies that have been recombinantly produced by mammalian cells, comprising:

- a) contacting the preparation with a set of oxidation/reduction reagents selected from cysteine/cystine and cysteine/cystamine to form a reducing mixture; and
- b) incubating the reducing mixture at a temperature of about 37 °C under anaerobic conditions for at least about 4 hours, or incubating the reducing mixture at a temperature of about 18-24 °C for about 16-24 hours;

wherein the IL-17 antibodies each comprise an immunoglobulin heavy chain variable domain (V_H) comprising the three complementarity determining regions (CDRs) of the V_H set forth as SEQ ID NO:8 and an immunoglobulin light chain variable domain (V_L) comprising the three CDRs of the V_L set forth as SEQ ID NO:10.

Also disclosed herein are also purified preparations of secukinumab, wherein the level of intact secukinumab in the preparation is at least about 90%, as measured by sodium dodecyl sulfate capillary electrophoresis (CE-SDS), and wherein the level of activity of secukinumab in the preparation is at least about 90%, as measured by cystamine-CEX.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows the percentage of intact antibody (LHHL) over time after subsection to different reducing agents.

Figure 2A shows the percentage of intact antibody (LHHL) over time at room temperature after subsection to selective reduction with 8 mM cysteine under anaerobic conditions. **Figure 2B** shows the percentage of intact antibody (LHHL) over time at room temperature after subsection to selective reduction with 8 mM cysteine under aerobic conditions. **Figure 2C** shows the percentage of intact antibody (LHHL) over time at 37°C after subsection to selective reduction with 8 mM cysteine under anaerobic conditions. **Figure 2D** shows the percentage of intact antibody (LHHL) over time at 37°C after subsection to selective reduction with 8 mM cysteine under aerobic conditions.

Figure 3 shows the percentage of intact antibody over time at 37 °C temperature after subjection to selective reduction with 8 mM cysteine at various dissolved oxygen concentrations.

Figure 4A shows a 4D contour plot for activity by CEX. **Figure 4B** shows a 4D contour plot for purity for CE-SDS. The plots of **Figure 4** analyze the impact of and interaction of cysteine concentration, protein content and ratio of cysteine/cystine on the output parameters activity by CEX and purity by CE-SDS.

Figure 5A shows a 4D contour plot for activity by CEX. **Figure 5B** shows a 4D contour plot for purity for CE-SDS. The plots of **Figure 5** analyze the impact of and interaction of cysteine concentration, protein content and cystine concentration on the output parameters activity by CEX and purity by CE-SDS.

Figure 6 shows a 4D contour plot for activity by CEX, looking at the impact and interaction of pH, time, temperature and cysteine concentration.

Figure 7 shows a 4D contour plot for purity by CE-SDS, looking at the impact and interaction of pH, time, temperature and cysteine concentration.

Figure 8 shows the dissolved oxygen chart of the cysteine treatment of the confirmation run 1.

Figure 9 shows the dissolved oxygen chart of the cysteine treatment of the confirmation run 2.

Figure 10 shows the dissolved oxygen chart of the cysteine treatment of the confirmation run 3.

Figure 11 shows the dissolved oxygen chart of the cysteine treatment of the confirmation run 4.

Figure 12 compares the reaction kinetic of confirmation run 3 and confirmation run 4 with respect to activity by CEX and purity by CE-SDS.

Figure 13A shows a scaled and centered coefficient plot for activity by CEX of REACT.P.

Figure 13B shows a 4D contour plot for activity by CEX of REACT.P. The plots of **Figure 13** analyze the impact of and interaction of cysteine concentration, protein content and stirrer speed on the output parameters activity by CEX and purity by CE-SDS.

Figure 14A shows the dissolved oxygen profiles of the process characterization runs at 0 rpm (the numbers in the legend indicate run number, stirrer speed and cysteine and antibody concentration). **Figure 14B** shows the dissolved oxygen profiles of the process characterization runs at 50 rpm (the numbers in the legend indicate run number, stirrer speed and cysteine and antibody concentration). **Figure 14C** shows the dissolved oxygen profiles of the process characterization runs at 100 rpm (the numbers in the legend indicate run number, stirrer speed and cysteine and antibody concentration). **Figure 14D** compares the dissolved oxygen profiles of

the two runs performed at 50 mL scale at 50 rpm and 6.0 mM cysteine, one without antibody and the other with 12.7 g/L antibody.

Figure 15 shows selective reduction kinetics of activity by CEX at different incubation temperatures.

Figure 16 shows an overlay of dissolved oxygen charts from scale-down model qualification runs, with an indication of the timing of various steps of the process (cysteine addition, heating, incubation, cooling, and pH adjustment).

Figure 17 shows the kinetic for activity by CEX of the manufacturing-scale runs and the scale-down model runs during selective reduction.

DETAILED DESCRIPTION OF THE DISCLOSURE

It is an object of the disclosure to provide methods for selectively reducing CysL97 in the antigen binding sites of certain IL-17 antibodies or antigen binding fragments thereof, such as secukinumab. By “selectively reducing” is meant that CysL97 in a disclosed IL-17 antibody or antigen binding fragment thereof is reduced to an oxidized form without reduction of the conserved cysteine residues of these antibodies. The conserved cysteine residues, in the case of a classical IgG₁ antibody, are: two disulfide bridges in the hinge region, two inter-chain disulfide bridges (one in each Fab), four intra-chain disulfide bridges in the Fc region, and eight intra-chain disulfide bridges in the Fab portion of the antibody. During the selective reduction process, transient reduction of the conserved cysteines of some antibodies in a particular preparation may occur. However, upon completion of the reaction, the vast majority of the conserved cysteines that were transiently reduced will have reoxidized to form the conserved disulfide bonds found in typical antibodies, resulting in high purity and activity in the selectively reduced preparation (i.e., purified preparation) of antibodies. It will be understood that upon completion of the selective reduction reaction, the selectively reduced preparation (i.e., purified preparation) is not expected to contain 100% intact antibodies; instead the selectively reduced preparation will ideally contain at least about 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or about 100 % (relative to theoretical maximum), intact antibodies as measured by CE-SDS.

The term “comprising” encompasses “including” as well as “consisting” e.g. a composition “comprising” X may consist exclusively of X or may include something additional

e.g. $X + Y$.

The term “about” in relation to a numerical value x means, for example, $\pm 10\%$. When used in front of a numerical range or list of numbers, the term “about” applies to each number in the series, e.g., the phrase “about 1-5” should be interpreted as “about 1 – about 5”, or, e.g., the phrase “about 1, 2, 3, 4” should be interpreted as “about 1, about 2, about 3, about 4, etc.”

The relative molecular mass of secukinumab, based on post-translational amino acid sequence, is 147,944 Daltons. This molecular weight (i.e., 147,944 Daltons) is used in the calculation of secukinumab molarity values and molar ratios throughout the instant disclosure. However, during production in CHO cells, a C-terminal lysine is commonly removed from each heavy chain. The relative molecular mass of secukinumab lacking a C-terminal lysine from each heavy chain is 147,688 Daltons. A preparation of secukinumab contains a mixture of molecules with and without C-terminal lysine residues on the heavy chain. The secukinumab molarity values (and ratios employing these molarity values) used in the instant disclosure are therefore estimates, and the term “about”, “approximate” and the like in reference to these numerical values encompasses at least this variation in relative molecular mass and the resulting calculations made therewith.

The word “substantially” does not exclude “completely” e.g. a composition which is “substantially free” from Y may be completely free from Y. Where necessary, the word “substantially” may be omitted from the definition of the disclosure.

The term "antibody" as referred to herein includes whole antibodies and any antigen-binding portion or single chains thereof. A naturally occurring "antibody" is a glycoprotein comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as V_H) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as V_L) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The V_H and V_L regions can be further subdivided into regions of hypervariability, termed hypervariable regions or complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each V_H and V_L is composed of three CDRs and four FRs arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the

heavy and light chains contain a binding domain that interacts with an antigen. The constant regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (e.g., effector cells) and the first component (C1q) of the classical complement system.

The term "antigen-binding fragment" of an antibody as used herein, refers to fragments of an antibody that retain the ability to specifically bind to an antigen (e.g., IL-17). It has been shown that the antigen-binding function of an antibody can be performed by fragments of a full-length antibody. Examples of binding fragments encompassed within the term "antigen-binding portion" of an antibody include a Fab fragment, a monovalent fragment consisting of the V_L , V_H , CL and CH1 domains; a F(ab)2 fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; a Fd fragment consisting of the V_H and CH1 domains; a Fv fragment consisting of the V_L and V_H domains of a single arm of an antibody; a dAb fragment (Ward et al., 1989 Nature 341:544-546), which consists of a V_H domain; and an isolated CDR. Exemplary antigen binding sites include the CDRs of secukinumab as set forth in SEQ ID NOs:1-6 and 11-13 (**Table 1**), preferably the heavy chain CDR3. Furthermore, although the two domains of the Fv fragment, V_L and V_H , are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the V_L and V_H regions pair to form monovalent molecules (known as single chain Fv (scFv); see, e.g., Bird et al., 1988 Science 242:423-426; and Huston et al., 1988 Proc. Natl. Acad. Sci. 85:5879-5883). Such single chain antibodies are also intended to be encompassed within the term "antibody". Single chain antibodies and antigen-binding portions are obtained using conventional techniques known to those of skill in the art.

An "isolated antibody", as used herein, refers to an antibody that is substantially free of other antibodies having different antigenic specificities (e.g., an isolated antibody that specifically binds IL-17 is substantially free of antibodies that specifically bind antigens other than IL-17). The term "monoclonal antibody" or "monoclonal antibody composition" as used herein refer to a preparation of antibody molecules of single molecular composition. The term "human antibody", as used herein, is intended to include antibodies having variable regions in which both the framework and CDR regions are derived from sequences of human origin. A "human antibody" need not be produced by a human, human tissue or human cell. The human antibodies of the disclosure may include amino acid residues not encoded by human sequences

(e.g., mutations introduced by random or site-specific mutagenesis *in vitro*, by N-nucleotide addition at junctions *in vivo* during recombination of antibody genes, or by somatic mutation *in vivo*). In some embodiments of the disclosed processes and compositions, the IL-17 antibody is a human antibody, an isolated antibody, and/or a monoclonal antibody.

The term "IL-17" refers to IL-17A, formerly known as CTLA8, and includes wild-type IL-17A from various species (e.g., human, mouse, and monkey), polymorphic variants of IL-17A, and functional equivalents of IL-17A. Functional equivalents of IL-17A according to the present disclosure preferably have at least about 65%, 75%, 85%, 95%, 96%, 97%, 98%, or even 99% overall sequence identity with a wild-type IL-17A (e.g., human IL-17A), and substantially retain the ability to induce IL-6 production by human dermal fibroblasts.

The term " K_D " is intended to refer to the dissociation rate of a particular antibody-antigen interaction. The term " K_D ", as used herein, is intended to refer to the dissociation constant, which is obtained from the ratio of K_d to K_a (i.e. K_d/K_a) and is expressed as a molar concentration (M). K_D values for antibodies can be determined using methods well established in the art. A method for determining the K_D of an antibody is by using surface plasmon resonance, or using a biosensor system such as a Biacore® system. In some embodiments, the IL-17 antibody or antigen binding fragment, e.g., secukinumab, has a K_D of about 100-250 pM for human IL-17.

The term "affinity" refers to the strength of interaction between antibody and antigen at single antigenic sites. Within each antigenic site, the variable region of the antibody "arm" interacts through weak non-covalent forces with antigen at numerous sites; the more interactions, the stronger the affinity. Standard assays to evaluate the binding affinity of the antibodies toward IL-17 of various species are known in the art, including for example, ELISAs, western blots and RIAs. The binding kinetics (e.g., binding affinity) of the antibodies also can be assessed by standard assays known in the art, such as by Biacore analysis.

An antibody that "inhibits" one or more of these IL-17 functional properties (e.g., biochemical, immunochemical, cellular, physiological or other biological activities, or the like) as determined according to methodologies known to the art and described herein, will be understood to relate to a statistically significant decrease in the particular activity relative to that seen in the absence of the antibody (or when a control antibody of irrelevant specificity is present). An antibody that inhibits IL-17 activity affects a statistically significant decrease, e.g., by at least about 10% of the measured parameter, by at least 50%, 80% or 90%, and in certain

embodiments of the disclosed methods and compositions, the IL-17 antibody used may inhibit greater than 95%, 98% or 99% of IL-17 functional activity.

“Inhibit IL-6” as used herein refers to the ability of an IL-17 antibody or antigen binding fragment thereof (e.g., secukinumab) to decrease IL-6 production from primary human dermal fibroblasts. The production of IL-6 in primary human (dermal) fibroblasts is dependent on IL-17 (Hwang et al., (2004) *Arthritis Res Ther*; 6:R120-128). In short, human dermal fibroblasts are stimulated with recombinant IL-17 in the presence of various concentrations of an IL-17 binding molecule or human IL-17 receptor with Fc part. The chimeric anti-CD25 antibody Simulect® (basiliximab) may be conveniently used as a negative control. Supernatant is taken after 16 h stimulation and assayed for IL-6 by ELISA. An IL-17 antibody or antigen binding fragment thereof, e.g., secukinumab, typically has an IC₅₀ for inhibition of IL-6 production (in the presence 1 nM human IL-17) of about 50 nM or less (e.g., from about 0.01 to about 50 nM) when tested as above, i.e., said inhibitory activity being measured on IL-6 production induced by hu-IL-17 in human dermal fibroblasts. In some embodiments of the disclosed methods and compositions, IL-17 antibodies or antigen binding fragments thereof, e.g., secukinumab, and functional derivatives thereof have an IC₅₀ for inhibition of IL-6 production as defined above of about 20 nM or less, more preferably of about 10 nM or less, more preferably of about 5 nM or less, more preferably of about 2 nM or less, more preferably of about 1 nM or less.

The term "derivative", unless otherwise indicated, is used to define amino acid sequence variants, and covalent modifications (e.g., pegylation, deamidation, hydroxylation, phosphorylation, methylation, etc.) of an IL-17 antibody or antigen binding fragment thereof, e.g., secukinumab, according to the present disclosure, e.g., of a specified sequence (e.g., a variable domain). A “functional derivative” includes a molecule having a qualitative biological activity in common with the disclosed IL-17 antibodies. A functional derivative includes fragments and peptide analogs of an IL-17 antibody as disclosed herein. Fragments comprise regions within the sequence of a polypeptide according to the present disclosure, e.g., of a specified sequence. Functional derivatives of the IL-17 antibodies disclosed herein (e.g., functional derivatives of secukinumab) preferably comprise V_H and/or V_L domains having at least about 65%, 75%, 85%, 95%, 96%, 97%, 98%, or 99% overall sequence identity with the V_H and/or V_L sequences of the IL-17 antibodies and antigen binding fragments thereof disclosed

herein (e.g., the V_H and/or V_L sequences of **Table 1**), and substantially retain the ability to bind human IL-17 or, e.g., inhibit IL-6 production of IL-17 induced human dermal fibroblasts.

The phrase “substantially identical” means that the relevant amino acid or nucleotide sequence (e.g., V_H or V_L domain) will be identical to or have insubstantial differences (e.g., through conserved amino acid substitutions) in comparison to a particular reference sequence. Insubstantial differences include minor amino acid changes, such as 1 or 2 substitutions in a 5 amino acid sequence of a specified region (e.g., V_H or V_L domain). In the case of antibodies, the second antibody has the same specificity and has at least 50% of the affinity of the same. Sequences substantially identical (e.g., at least about 85% sequence identity) to the sequences disclosed herein are also part of this application. In some embodiments, the sequence identity of a derivative IL-17 antibody (e.g., a derivative of secukinumab, e.g., a secukinumab biosimilar antibody) can be about 90% or greater, e.g., 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or higher relative to the disclosed sequences.

“Identity” with respect to a native polypeptide and its functional derivative is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the residues of a corresponding native polypeptide, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent identity, and not considering any conservative substitutions as part of the sequence identity. Neither N- or C-terminal extensions nor insertions shall be construed as reducing identity. Methods and computer programs for the alignment are well known. The percent identity can be determined by standard alignment algorithms, for example, the Basic Local Alignment Search Tool (BLAST) described by Altschul et al. ((1990) J. Mol. Biol., 215: 403 410); the algorithm of Needleman et al. ((1970) J. Mol. Biol., 48: 444 453); or the algorithm of Meyers et al. ((1988) Comput. Appl. Biosci., 4: 11 17). A set of parameters may be the Blosom 62 scoring matrix with a gap penalty of 12, a gap extend penalty of 4, and a frameshift gap penalty of 5. The percent identity between two amino acid or nucleotide sequences can also be determined using the algorithm of E. Meyers and W. Miller ((1989) CABIOS, 4:11-17) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

"Amino acid(s)" refer to all naturally occurring L- α -amino acids, e.g., and include D-amino acids. The phrase "amino acid sequence variant" refers to molecules with some differences in their amino acid sequences as compared to the sequences according to the present

disclosure. Amino acid sequence variants of an antibody according to the present disclosure, e.g., of a specified sequence, still have the ability to bind the human IL-17 or, e.g., inhibit IL-6 production of IL-17 induced human dermal fibroblasts. Amino acid sequence variants include substitutional variants (those that have at least one amino acid residue removed and a different amino acid inserted in its place at the same position in a polypeptide according to the present disclosure), insertional variants (those with one or more amino acids inserted immediately adjacent to an amino acid at a particular position in a polypeptide according to the present disclosure) and deletional variants (those with one or more amino acids removed in a polypeptide according to the present disclosure).

The phrases “free cysteine”, “non-traditional cysteine” and “unpaired cysteine” interchangeably refer to a cysteine that is not involved in conserved antibody disulfide bonding. The free cysteine may be present in an antibody framework region or a variable region (e.g., within a CDR). In secukinumab, amino acid eight of L-CDR3 as set forth as SEQ ID NO:6, which corresponds to amino acid 97 of the light chain variable region as set forth as SEQ ID NO:10 (herein after referred to as CysL97) is a free cysteine. Each molecule of secukinumab comprises two such free cysteine residues – one in each V_L domain. The disclosed processes are capable of selectively reducing both free cysteine residues in secukinumab. In some embodiments, e.g., due to deletions and/or substitutions in the light chain of a disclosed IL-17 antibody or antigen binding fragment thereof, the free cysteine will not be present at position CysL97. In such case, the corresponding free cysteine is the target of the selective reduction reaction and is included within the term “CysL97”.

IL-17 Antibodies and Antigen Binding Fragments Thereof

The various disclosed processes and relate to the selective reduction of certain IL-17 antibodies or antigen binding fragments thereof (e.g., secukinumab). In one embodiment, the IL-17 antibody or antigen binding fragment thereof comprises at least one immunoglobulin heavy chain variable domain (V_H) comprising hypervariable regions CDR1, CDR2 and CDR3, said CDR1 having the amino acid sequence SEQ ID NO:1, said CDR2 having the amino acid sequence SEQ ID NO:2, and said CDR3 having the amino acid sequence SEQ ID NO:3. In one embodiment, the IL-17 antibody or antigen binding fragment thereof comprises at least one immunoglobulin light chain variable domain (V_L) comprising hypervariable regions CDR1’,

CDR2' and CDR3', said CDR1' having the amino acid sequence SEQ ID NO:4, said CDR2' having the amino acid sequence SEQ ID NO:5 and said CDR3' having the amino acid sequence SEQ ID NO:6. In one embodiment, the IL-17 antibody or antigen binding fragment thereof comprises at least one immunoglobulin heavy chain variable domain (V_H) comprising hypervariable regions CDR1-x, CDR2-x and CDR3-x, said CDR1-x having the amino acid sequence SEQ ID NO:11, said CDR2-x having the amino acid sequence SEQ ID NO:12, and said CDR3-x having the amino acid sequence SEQ ID NO:13.

In one embodiment, the IL-17 antibody or antigen binding fragment thereof comprises at least one immunoglobulin V_H domain and at least one immunoglobulin V_L domain, wherein: a) the V_H domain comprises (e.g., in sequence): i) hypervariable regions CDR1, CDR2 and CDR3, said CDR1 having the amino acid sequence SEQ ID NO:1, said CDR2 having the amino acid sequence SEQ ID NO:2, and said CDR3 having the amino acid sequence SEQ ID NO:3; or ii) hypervariable regions CDR1-x, CDR2-x and CDR3-x, said CDR1-x having the amino acid sequence SEQ ID NO:11, said CDR2-x having the amino acid sequence SEQ ID NO:12, and said CDR3-x having the amino acid sequence SEQ ID NO:13; and b) the V_L domain comprises (e.g., in sequence) hypervariable regions CDR1', CDR2' and CDR3', said CDR1' having the amino acid sequence SEQ ID NO:4, said CDR2' having the amino acid sequence SEQ ID NO:5, and said CDR3' having the amino acid sequence SEQ ID NO:6.

In one embodiment, the IL-17 antibody or antigen binding fragment thereof comprises: a) an immunoglobulin heavy chain variable domain (V_H) comprising the amino acid sequence set forth as SEQ ID NO:8; b) an immunoglobulin light chain variable domain (V_L) comprising the amino acid sequence set forth as SEQ ID NO:10; c) an immunoglobulin V_H domain comprising the amino acid sequence set forth as SEQ ID NO:8 and an immunoglobulin V_L domain comprising the amino acid sequence set forth as SEQ ID NO:10; d) an immunoglobulin V_H domain comprising the hypervariable regions set forth as SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:3; e) an immunoglobulin V_L domain comprising the hypervariable regions set forth as SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6; f) an immunoglobulin V_H domain comprising the hypervariable regions set forth as SEQ ID NO:11, SEQ ID NO:12 and SEQ ID NO:13; g) an immunoglobulin V_H domain comprising the hypervariable regions set forth as SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:3 and an immunoglobulin V_L domain comprising the hypervariable regions set forth as SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6; or h) an

immunoglobulin V_H domain comprising the hypervariable regions set forth as SEQ ID NO:11, SEQ ID NO:12 and SEQ ID NO:13 and an immunoglobulin V_L domain comprising the hypervariable regions set forth as SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6.

For ease of reference the amino acid sequences of the hypervariable regions of the secukinumab monoclonal antibody, based on the Kabat definition and as determined by the X-ray analysis and using the approach of Chothia and coworkers, is provided in **Table 1**, below.

Light-Chain		
CDR1'	Kabat	R-A-S-Q-S-V-S-S-S-Y-L-A (SEQ ID NO:4)
	Chothia	R-A-S-Q-S-V-S-S-S-Y-L-A (SEQ ID NO:4)
CDR2'	Kabat	G-A-S-S-R-A-T (SEQ ID NO:5)
	Chothia	G-A-S-S-R-A-T (SEQ ID NO:5)
CDR2'	Kabat	Q-Q-Y-G-S-S-P-C-T (SEQ ID NO:6)
	Chothia	Q-Q-Y-G-S-S-P-C-T (SEQ ID NO:6)
Heavy-Chain		
CDR1	Kabat	N-Y-W-M-N (SEQ ID NO:1)
CDR1-x	Chothia	G-F-T-F-S-N-Y-W-M-N (SEQ ID NO:11)
CDR2	Kabat	A-I-N-Q-D-G-S-E-K-Y-Y-V-G-S-V-K-G (SEQ ID NO:2)
CDR2-x	Chothia	A-I-N-Q-D-G-S-E-K-Y-Y (SEQ ID NO:12)
CDR3	Kabat	D-Y-Y-D-I-L-T-D-Y-Y-I-H-Y-W-Y-F-D-L (SEQ ID NO:3)
CDR3-x	Chothia	C-V-R-D-Y-Y-D-I-L-T-D-Y-Y-I-H-Y-W-Y-F-D-L-W-G (SEQ ID NO:13)

Table 1: Amino acid sequences of the hypervariable regions of the secukinumab monoclonal antibodies.

In preferred embodiments, the constant region domains preferably also comprise suitable human constant region domains, for instance as described in "Sequences of Proteins of Immunological Interest", Kabat E.A. et al, US Department of Health and Human Services, Public Health Service, National Institute of Health. DNA encoding the VL of secukinumab is set forth in SEQ ID NO:9. DNA encoding the V_H of secukinumab is set forth in SEQ ID NO:7.

In some embodiments, the IL-17 antibody or antigen binding fragment thereof (e.g., secukinumab) comprises the three CDRs of SEQ ID NO:10. In other embodiments, the IL-17 antibody or antigen binding fragment thereof comprises the three CDRs of SEQ ID NO:8. In other embodiments, the IL-17 antibody or antigen binding fragment thereof comprises the three CDRs of SEQ ID NO:10 and the three CDRs of SEQ ID NO:8. CDRs of SEQ ID NO:8 and SEQ ID NO:10 may be found in **Table 1**. The free cysteine in the light chain (CysL97) may be seen in SEQ ID NO:6.

In some embodiments, IL-17 antibody or antigen binding fragment thereof comprises the light chain of SEQ ID NO:14. In other embodiments, the IL-17 antibody or antigen binding fragment thereof comprises the heavy chain of SEQ ID NO:15 (with or without the C-terminal lysine). In other embodiments, the IL-17 antibody or antigen binding fragment thereof comprises the light chain of SEQ ID NO:14 and the heavy chain of SEQ ID NO:15 (with or without the C-terminal lysine). In some embodiments, the IL-17 antibody or antigen binding fragment thereof comprises the three CDRs of SEQ ID NO:14. In other embodiments, IL-17 antibody or antigen binding fragment thereof comprises the three CDRs of SEQ ID NO:15. In other embodiments, the IL-17 antibody or antigen binding fragment thereof comprises the three CDRs of SEQ ID NO:14 and the three CDRs of SEQ ID NO:15. CDRs of SEQ ID NO:14 and SEQ ID NO:15 may be found in **Table 1**.

Hypervariable regions may be associated with any kind of framework regions, though preferably are of human origin. Suitable framework regions are described in Kabat E.A. et al, *ibid*. The preferred heavy chain framework is a human heavy chain framework, for instance that of the secukinumab antibody. It consists in sequence, e.g. of FR1 (amino acid 1 to 30 of SEQ ID NO:8), FR2 (amino acid 36 to 49 of SEQ ID NO:8), FR3 (amino acid 67 to 98 of SEQ ID NO:8) and FR4 (amino acid 117 to 127 of SEQ ID NO:8) regions. Taking into consideration the determined hypervariable regions of secukinumab by X-ray analysis, another preferred heavy chain framework consists in sequence of FR1-x (amino acid 1 to 25 of SEQ ID NO:8), FR2-x

(amino acid 36 to 49 of SEQ ID NO:8), FR3-x (amino acid 61 to 95 of SEQ ID NO:8) and FR4 (amino acid 119 to 127 of SEQ ID NO:8) regions. In a similar manner, the light chain framework consists, in sequence, of FR1' (amino acid 1 to 23 of SEQ ID NO:10), FR2' (amino acid 36 to 50 of SEQ ID NO:10), FR3' (amino acid 58 to 89 of SEQ ID NO:10) and FR4' (amino acid 99 to 109 of SEQ ID NO:10) regions.

In one embodiment, the IL-17 antibody or antigen binding fragment thereof (e.g., secukinumab) is selected from a human IL-17 antibody that comprises at least: a) an immunoglobulin heavy chain or fragment thereof comprising a variable domain comprising, in sequence, the hypervariable regions CDR1, CDR2 and CDR3 and the constant part or fragment thereof of a human heavy chain; said CDR1 having the amino acid sequence SEQ ID NO:1, said CDR2 having the amino acid sequence SEQ ID NO:2, and said CDR3 having the amino acid sequence SEQ ID NO:3; and b) an immunoglobulin light chain or fragment thereof comprising a variable domain comprising, in sequence, the hypervariable regions CDR1', CDR2', and CDR3' and the constant part or fragment thereof of a human light chain, said CDR1' having the amino acid sequence SEQ ID NO: 4, said CDR2' having the amino acid sequence SEQ ID NO:5, and said CDR3' having the amino acid sequence SEQ ID NO:6.

In one embodiment, the IL-17 antibody or antigen binding fragment thereof is selected from a single chain antibody or antigen binding fragment thereof that comprises an antigen binding site comprising: a) a first domain comprising, in sequence, the hypervariable regions CDR1, CDR2 and CDR3, said CDR1 having the amino acid sequence SEQ ID NO:1, said CDR2 having the amino acid sequence SEQ ID NO:2, and said CDR3 having the amino acid sequence SEQ ID NO:3; and b) a second domain comprising, in sequence, the hypervariable regions CDR1', CDR2' and CDR3', said CDR1' having the amino acid sequence SEQ ID NO:4, said CDR2' having the amino acid sequence SEQ ID NO:5, and said CDR3' having the amino acid sequence SEQ ID NO:6; and c) a peptide linker which is bound either to the N-terminal extremity of the first domain and to the C-terminal extremity of the second domain or to the C-terminal extremity of the first domain and to the N-terminal extremity of the second domain.

Alternatively, an IL-17 antibody or antigen binding fragment thereof as used in the disclosed methods may comprise a derivative of the IL-17 antibodies set forth herein by sequence (e.g., a pegylated version of secukinumab). Alternatively, the V_H or V_L domain of an IL-17 antibody or antigen binding fragment thereof used in the disclosed methods may have V_H

or V_L domains that are substantially identical to the V_H or V_L domains set forth herein (e.g., those set forth in SEQ ID NO:8 and 10). A human IL-17 antibody disclosed herein may comprise a heavy chain that is substantially identical to that set forth as SEQ ID NO:15 (with or without the C-terminal lysine) and/or a light chain that is substantially identical to that set forth as SEQ ID NO:14. A human IL-17 antibody disclosed herein may comprise a heavy chain that comprises SEQ ID NO:15 (with or without the C-terminal lysine) and a light chain that comprises SEQ ID NO:14. A human IL-17 antibody disclosed herein may comprise: a) one heavy chain which comprises a variable domain having an amino acid sequence substantially identical to that shown in SEQ ID NO:8 and the constant part of a human heavy chain; and b) one light chain which comprises a variable domain having an amino acid sequence substantially identical to that shown in SEQ ID NO:10 and the constant part of a human light chain.

Alternatively, an IL-17 antibody or antigen binding fragment thereof used in the disclosed methods may be an amino acid sequence variant of the reference IL-17 antibodies set forth herein, as long as it contains CysL97. The disclosure also includes IL-17 antibodies or antigen binding fragments thereof (e.g., secukinumab) in which one or more of the amino acid residues of the V_H or V_L domain of secukinumab (but not CysL97), typically only a few (e.g., 1-10), are changed; for instance by mutation, e.g., site directed mutagenesis of the corresponding DNA sequences. In all such cases of derivative and variants, the IL-17 antibody or antigen binding fragment thereof is capable of inhibiting the activity of about 1 nM (= 30 ng/ml) human IL-17 at a concentration of about 50 nM or less, about 20 nM or less, about 10 nM or less, about 5 nM or less, about 2 nM or less, or more preferably of about 1 nM or less of said molecule by 50%, said inhibitory activity being measured on IL-6 production induced by hu-IL-17 in human dermal fibroblasts as described in Example 1 of WO 2006/013107.

In some embodiments, the IL-17 antibodies or antigen binding fragments thereof, e.g., secukinumab, bind to an epitope of mature human IL-17 comprising Leu74, Tyr85, His86, Met87, Asn88, Val124, Thr125, Pro126, Ile127, Val128, His129. In some embodiments, the IL-17 antibody, e.g., secukinumab, binds to an epitope of mature human IL-17 comprising Tyr43, Tyr44, Arg46, Ala79, Asp80. In some embodiments, the IL-17 antibody, e.g., secukinumab, binds to an epitope of an IL-17 homodimer having two mature human IL-17 chains, said epitope comprising Leu74, Tyr85, His86, Met87, Asn88, Val124, Thr125, Pro126, Ile127, Val128, His129 on one chain and Tyr43, Tyr44, Arg46, Ala79, Asp80 on the other chain. The residue

numbering scheme used to define these epitopes is based on residue one being the first amino acid of the mature protein (ie., IL-17A lacking the 23 amino acid N-terminal signal peptide and beginning with Glycine). The sequence for immature IL-17A is set forth in the Swiss-Prot entry Q16552. In some embodiments, the IL-17 antibody has a K_D of about 100-200 pM. In some embodiments, the IL-17 antibody has an IC_{50} of about 0.4 nM for *in vitro* neutralization of the biological activity of about 0.67 nM human IL-17A. In some embodiments, the absolute bioavailability of subcutaneously (s.c.) administered IL-17 antibody has a range of about 60 – about 80%, e.g., about 76%. In some embodiments, the IL-17 antibody, such as secukinumab, has an elimination half-life of about 4 weeks (e.g., about 23 to about 35 days, about 23 to about 30 days, e.g., about 30 days). In some embodiments, the IL-17 antibody (such as secukinumab) has a T_{max} of about 7-8 days.

Particularly preferred IL-17 antibodies or antigen binding fragments thereof used in the disclosed methods are human antibodies, especially secukinumab as described in Examples 1 and 2 of WO 2006/013107. Secukinumab is a recombinant high-affinity, fully human monoclonal anti-human interleukin-17A (IL-17A, IL-17) antibody of the IgG1/kappa isotype that is currently in clinical trials for the treatment of immune-mediated inflammatory conditions. Secukinumab (see, e.g., WO2006/013107 and WO2007/117749) has a very high affinity for IL-17, i.e., a K_D of about 100-200 pM and an IC_{50} for *in vitro* neutralization of the biological activity of about 0.67 nM human IL-17A of about 0.4 nM. Thus, secukinumab inhibits antigen at a molar ratio of about 1:1. This high binding affinity makes the secukinumab antibody particularly suitable for therapeutic applications. Furthermore, it has been determined that secukinumab has a very long half-life, i.e., about 4 weeks, which allows for prolonged periods between administration, an exceptional property when treating chronic life-long disorders, such as rheumatoid arthritis.

Disclosed herein are processes for selectively reducing CysL97 in preparations of the above-mentioned IL-17 antibodies and antigen binding fragments thereof (e.g., secukinumab). The disclosed methods conveniently may be performed on preparations of antibodies (e.g., IL-17 antibodies, e.g., secukinumab) to reduce cost. A “preparation” of antibodies refers to a composition (e.g., solution) having a plurality of an antibody molecule. A “preparation” includes any liquid composition comprising the IL-17 antibody or antigen binding fragment thereof. As such, a preparation may comprise, e.g., IL-17 antibody or antigen binding fragment thereof, e.g.,

secukinumab, in water or a buffer, in a column eluate, in a dialysis buffer, etc. In some embodiments, the initial preparation of antibodies comprises a pool of the IL-17 antibodies or antigen binding fragments thereof, e.g., secukinumab, in a buffer (e.g., a Tris, e.g., 1 mM – 1 M Tris, pH 6.0-8.0) or WFI. Prior to addition of the reducing agent to the antibody, the preparation may be adjusted by modifying dissolved oxygen levels, solution pH, antibody concentration, etc. In some embodiments, prior to addition of a reducing agent, the concentration of antibody (e.g., secukinumab) in the preparation is adjusted to between about 4 mg/ml – about 19.4 mg/ml, e.g., about 10 mg/ml - about 19.4 mg/ml, about 10 mg/ml - about 15.4 mg/ml, about 12 mg/ml - about 15 mg/ml, or about 13.5 mg/ml. In some embodiments, prior to addition of the reducing agent, the percent oxygen saturation in the preparation is adjusted to at least about 60% (as measured using an oxygen probe calibrated at 25°C), e.g., at least about 80%. In some embodiments, prior to addition of the reducing agent, the pH of the preparation is adjusted to about 7.3 - about 8.5, e.g., about 7.8 - about 8.2, e.g., about 7.9 - about 8.1, e.g., about 8.0. The concentration of antibody, pH and level of oxygen may also be adjusted immediately after (or even during) addition of the reductant, and thus should be interpreted as equivalent.

The preparations of IL-17 antibodies or antigen binding fragments thereof for use in the disclosed processes may be recombinantly produced by any mammalian cells using any mammalian cell line, e.g., Chinese hamster ovary cells (CHO) cells, mouse myeloma NS0 cells, baby hamster kidney (BHK) cells, human embryonic kidney cell line HEK-293, the human retinal cell line Per.C6 (Crucell, NL), HKB11 cell clone (derived from a hybrid cell fusion of HEK 293S with the Burkitt's lymphoma line 2B8), etc. By “recombinantly produced by mammalian cells” is meant that production of the antibody in the mammalian cells has been achieved using recombinant DNA technology. The IL-17 antibody preparation subjected to selective reduction may be a pool of antibodies harvested from the mammalian cells by centrifugation (with or without subsequent clarification). Alternatively, the IL-17 antibody preparation subjected to selective reduction may be a pool of antibodies from a further downstream chromatography step, e.g., an eluate from an affinity column (e.g., a protein A column), a cation exchange column, an anion exchange column, etc. Alternatively, the IL-17 antibody preparation subjected to selective reduction may be a pool of antibodies from a downstream filtration step, e.g., depth filtration, nanofiltration, ultrafiltration, etc. Alternatively, the IL-17 antibody preparation subjected to selective reduction may be a pool of

antibodies from a downstream step in which the pool has been treated to remove host cell proteins and/or to inactivate viri. In a one embodiment, the preparation of IL-17 antibodies subjected to selective reduction is a protein A eluate pool of antibodies.

Depending on the process conditions chosen, e.g., temperature, length of reaction time, pH, etc.) the concentration of antibody in the original preparation will vary. In some embodiments, the concentration of the IL-17 antibody used in the original preparation is between about 2 mg/ml to about 20 mg/ml, about 3.8 mg/ml to about 19.5 mg/ml, about 4 mg/ml to about 19.5 mg/ml, about 10 mg/ml to about 19.4 mg/ml, e.g., about 10 mg/ml to about 15.4 mg/ml, e.g., about 12 mg/ml to about 15 mg/ml, e.g., about 13.5 mg/ml of the IL-17 antibodies or antigen binding fragments thereof. Prior to selective reduction, the antibody concentration in the initial antibody preparation may be adjusted as desired using water for injection (WFI) or a buffer of choice.

The selective reduction processes described herein may be performed in any size vessel. In some embodiments, the vessel is lab-scale (e.g., 1L-2L). In other embodiments, the vessel is pilot-scale (e.g., 12 L-20 L). In further embodiments, the vessel is commercial-scale (e.g., greater than 10,000 L, e.g., 14,000 L, 15,000 L, 16,000 L, etc.).

Reducing Agents

Reducing agents are substances capable of electron donation in a redox (reduction-oxidation) reaction. Specifically, such agents are useful to deliver hydrogen to a masked (or blocked) cysteine present in the IL-17 antibody or antigen binding fragment thereof (e.g., secukinumab antibody). The process disclosed herein uses reducing agents for the selective reduction of the IL-17 antibody. Each reducing agent referred to herein include derivatives thereof (e.g., salts, esters and amides). Thus, e.g., reference to "cysteine" includes cysteine and cysteine-HCL, reference to "TCEP" includes TCEP and TCEP-HCL, reference to thioglycolic acid includes sodium thioglycolate, etc. Reducing agents for use in the disclosed methods include sodium bisulfate, ammonia, triethylsilane, glycylcysteine, sodium cyanoborohydride, ammonium thioglycolate, calcium thioglycolate, sodium thioglycolate, ascorbic acid, hydroquinone, aminomethanesulphonic acid, cysteic acid, cysteinesulphinic acid, ethanedisulphonic acid, ethanesulphonic acid, homotaurine, hypotaurine, isethionic acid, mercaptoethanesulphonic acid, N-methyltaurine (MTAU), TCEP (Tris(2-

carboxyethyl)phosphine hydrochloride), N-N-dimethyl-N-N bis(mercaptoacetyl)hydrazine (DMH), dithiothreitol (DTT), 2-mercaptoethanol (beta-mercaptoethanol), 2-mercaptoacetic acid (thioglycolic acid, TGA), cysteine (L-cysteine), cysteamine (beta-mercaptoethylamine, or MEA), glutathione, and combinations thereof. In some embodiments, the reducing agent for use in the disclosed process is a thiol-containing reducing agent (i.e., a compound having an R-SH group), e.g., an organosulfur compound. In some embodiments, the reducing agent for use in the disclosed process is, e.g., dithiothreitol (DTT), 2-mercaptoethanol, 2-mercaptoacetic acid, cysteine, cysteamine, glutathione and combinations.

The strength of a reducing agent is indicated by its oxidation-reduction potential (redox potential), E° , which is given in Volts (V) and traditionally determined at pH 7, 25°C. For example, the standard oxidation-reduction potential, E° , for CSH/CSSC is given as about -0.20 V to about -0.23 V (pH 7, 25°C) (P. C. Jocelyn (1967) *Eu. J. Biochem* 2:327-31; Liu “The role of Sulfur in Proteins,” in *The Proteins*, 3rd Ed. (ed. Neurath) p. 250-252 Academic Press 1977). The standard oxidation-reduction potential, E° , of DTT is given as about -0.33 V (pH 7, 25°C) (M.J.O'Neil, ed. by (2001). *Merck Index: an encyclopedia of chemicals, drugs, & biologicals*: 13th ed. (13. ed. ed.) United States: MERCK & CO INC.; Liu, *supra*). The standard oxidation-reduction potential, E° , of glutathione is given as about about -0.24 V or about -0.26 V (pH 7, 25°C) (Rost and Rapoport (1964) *Nature* 201:185; Gilbert (1990) *Adv. Enzymol. Relat. Areas Mol. Biol.* 63:69-172; Giles (2002) *Gen. Physiol Biophys* 21:65-72; Liu, *supra*). The standard oxidation-reduction potential, E° , of 2-mercaptoethanol is given as about -0.26 V (Lee and Whitesides (1990) *J. Org. Chem* 58:642-647). In some embodiments, the reducing agent has a standard oxidation-reduction potential, E° , similar to cysteine (e.g., about -0.20 V to about -0.23 V, about -0.20 V to about -0.22 V, about -0.20 V to about -0.21 V, about -0.21 V to about -0.23 V, about -0.21 V to -0.22 V, about -0.22 V to about -0.23 V, about -0.20 V, about -0.21 V, about -0.22 V, about -0.23 V).

The standard oxidation-reduction potential E° of thiol-containing compounds may be measured by thermal analysis, reduction of NAD⁺, polarography, reaction with Fe⁺⁺, or thiol-disulfide exchange studies (Jocelyn, *supra*; Borsook et al. (1937) *J. Biol. Chem* 117:281; Ghosh et al. (1932) *J. Indian Chem. Soc.* 9:43; Kolthoff et al. (1955) *J. Am. Chem. Soc.* 77:4739; Tanaka et al. (1955) *J. Am. Chem. Soc.* 77:2004; Kolthoff et al. (1955) *J. Am. Chem. Soc.* 77:4733; Eldjarn (1957) *J. Am. Chem. Soc.* 79:4589). In some embodiments, the standard

oxidation-reduction potential E° is determined by thermal analysis, polarography, reaction with Fe^{++} , or thiol-disulfide exchange studies, e.g., preferably by thiol-disulfide exchange studies. In some embodiments, the standard oxidation-reduction potential E° is determined at pH 7, 25°C .

The reducing agent, when combined with the antibody preparation, forms a “reducing mixture.” The reducing mixture may comprise excipients in addition to the reducing agent and the IL-17 antibody. For example, in certain embodiments, a small molar ratio of the oxidized form (e.g., cystine, cystamine) of the reducing agent may be added to the reducing mixture either simultaneously with the reducing agent or sequentially, e.g., 10-30 minutes or more after the start of incubation. For example, if cysteine is the reducing agent, then a small amount of cystine may be added to the reducing mixture, e.g., concurrently with the cysteine or, e.g., 15, 20, 30 minutes after cysteine is combined with the IL-17 antibody or antigen binding fragment thereof. Thus, in some embodiments the reducing mixture comprises a set of oxidation/reduction reagents. By “set of oxidation/reduction reagents” is meant a redox pair or redox couple, i.e., an oxidizing and reducing agent that appear on opposite sides of a half-equation (e.g., a reducing species and its corresponding oxidized form, e.g., $\text{Fe}^{2+}/\text{Fe}^{3+}$, cysteine/cystine, cysteamine/cystamine).

Depending on the reaction conditions (temperature, length of reaction time, quantity of IL-17 antibody or antigen binding fragment thereof, pH, etc.) the concentration of reducing agent used in a particular reducing mixture and selective reduction reaction will vary. In some embodiments, the amount of reducing agent used in the reducing mixture will vary from about 1 to about 20 mM. In some embodiments, the concentration of reducing agent employed in the reducing mixture is about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16 mM. In one embodiment, the amount of reducing agent (e.g., cysteine) is between about 4 mM and about 8 mM, e.g., 5.9 mM, 6 mM, 7.7 mM, 7.9 mM, 8 mM.

In one embodiment, the reducing agent is beta-mercaptoethanol. In certain embodiments, beta-mercaptoethanol is employed at a concentration of about 2.0 mM to about 8.0 mM.

In one embodiment, the reducing agent is glutathione. In certain embodiments, glutathione is employed at a concentration of about 2.0 mM to about 5.0 mM.

In one embodiment, the reducing agent is cysteamine. In certain embodiments, cysteamine is employed at a concentration of about 1.0 mM to 20 mM, about 4.0 mM to about 19 mM, about 2.0 mM to about 8.0 mM, about 4.0 mM to about 8.0 mM, about 4.8 mM to about 8.0 mM, about 5.5 mM to about 6.7 mM, or about 6.0 mM.

In one embodiment, the reducing agent is cysteine. In certain embodiments, the concentration of cysteine in the reducing mixture is about 1.0 mM to 20 mM, about 4.0 mM to about 19 mM, about 2.0 mM to about 8.0 mM, about 4.0 mM to about 8.0 mM, about 4.8 mM to about 8.0 mM, about 5.5 mM to about 6.7 mM, or about 6.0 mM. Cysteine concentration may be adjusted using a stock solution of, e.g., 120 mM cysteine-HCL.

Each IL-17 antibody has two CysL97 residues in need of selective reduction. The amount of reducing agent employed should therefore be sufficient to selectively reduce both CysL97 residues on a substantial portion of IL-17 antibodies in a preparation of antibodies, without concomitantly over-reducing the antibody by irreversibly reducing the traditional disulfide bonds. Depending on the reaction conditions (presence of oxidizing agent, temperature, length of reaction time, pH, etc.) the molar ratio of reducing agent:IL-17 antibody used in a particular reducing mixture and selective reduction reaction will vary. We have found that the molar ratio of reducing agent (e.g., cysteine):antibody (e.g., secukinumab) can range from about 11:1 (**Example 5.2**) to as high as about 546:1 (**Example 6.2**). In some embodiments of the disclosed methods, the molar ratio of reducing agent (e.g., cysteine):antibody (e.g., secukinumab) is between about 11:1 to about 462:1 (e.g., about 21:1), about 31:1 to about 545:1 (e.g., about 31:1 to about 156:1), about 21:1 to about 296:1, or about 46:1 to about 91:1. In other embodiments, the molar ratio of reducing agent (e.g., cysteine):antibody (e.g., secukinumab) is between about 23:1 to about 91:1 (e.g., about 23:1 to about 57:1), about 44:1 to about 275:1 (e.g., about 44:1), about 44:1 to about 66:1 (e.g., about 44:1 to about 66:1), preferably about 46:1 to about 118:1 (e.g., about 56:1 to about 118:1), more preferably about 54:1 to about 82:1. In one embodiment, the molar ratio of reducing agent (e.g., cysteine):antibody (e.g., secukinumab) is about 66:1.

If a higher molar ratio of reducing agent (e.g., cysteine):antibody (e.g., secukinumab) is used (representing excess reducing agent c.f. to antibody), then addition of a small amount of the corresponding oxidizing agent (e.g., cystine or cystamine) may be useful to mitigate the reductive power of the reducing agent (e.g., cysteine or cysteamine). This is particularly beneficial in an anaerobic environment. Thus, in some embodiments, selective reduction is carried out using a set of oxidation/reduction reagents (e.g., in an aerobic or anaerobic environment, preferably an anaerobic environment). In some embodiments, selective reduction is carried out using a molar ratio of reducing agent (e.g., cysteine):oxidizing agent (e.g., cystine)

of about 2:1 to about 80:1, about 4:1 to about 80:1, about 26:1 to about 80:1, about 2:1 to about 10:1 (e.g., about 6:1 to about 10:1), about 4:1 to about 28:1 (about 4:1 to about 18:1), about 27:1 to about 53:1 (e.g., about 27:1) in the reducing mixture. In certain embodiments, cysteine is used in the reducing mixture in combination with the oxidizing agent cystine or cystamine (preferably cystine). In some embodiments, selective reduction is carried out in conditions using about 4 mM -14 mM cysteine (e.g., about 7.7 mM to about 8.0 mM cysteine) and about 0.1 to about 1 mM cystine (e.g., about 0.1 to about 0.3 mM cystine) in the reducing mixture. In certain embodiments, the reducing mixture contains about 8.0 mM cysteine and about 0.1 mM cystine, about 7.9 mM cysteine and about 0.1 mM cystine, or about 7.7 mM cysteine and about 0.3 mM cystine. It will be understood that if an oxidizing agent, e.g., cystine, is employed in combination with the reducing agent, e.g., cysteine, in the disclosed process, the oxidizing agent, e.g., cystine, may be added at a point after the reducing agent, e.g., cysteine, is combined with the IL-17 antibody or antigen binding fragment thereof. For example, the IL-17 antibody or antigen binding fragment thereof may be combined with cysteine to form a reducing mixture, which is then incubated for, e.g., 15-30 minutes; thereafter, cystine may be added to the reaction.

Dissolved Oxygen

As used herein, “dissolved oxygen”, “dO₂” and “DO” refer to the amount of oxygen that is dissolved or carried in a given medium. It can be measured with an oxygen probe, such as an oxygen sensor or an optode in liquid media. DO is reported as either as a concentration (milligrams per liter (mg/L)) or as “percent saturation.” Milligrams per liter is the amount of oxygen in a liter of solvent and is also equivalent to parts per million = ppm. Percent oxygen saturation is the amount of oxygen in a solution relative to the total amount of oxygen that the solution can hold at a particular temperature.

As used herein, “initial percent oxygen saturation” refers to the amount of dissolved oxygen in the preparation of IL-17 antibodies (e.g., secukinumab) prior to contacting the preparation with the reducing agent in the vessel to form the reducing mixture. The initial percent oxygen saturation can be adjusted directly (e.g., by sparging) or indirectly (e.g., by stirring) to achieve a desired level of oxygen prior to the beginning of the selective reduction process. For example, in some embodiments the initial percent oxygen saturation in the IL-17 antibody preparation is adjusted to at least 40%, 50%, 60%, 70%, 80%, 90%, or even as high as

100% prior to contact with the reducing agent. This may be done in order to initially mitigate the power of the reducing agent once that agent is added to the IL-17 antibody preparation to form the reducing mixture, which avoids partial or complete reduction of the traditional disulfides of the antibody that otherwise would lead to loss of activity and purity. In preferred embodiments, the initial percent oxygen saturation in the IL-17 antibody preparation is adjusted to at least 60% (as measured using an oxygen probe calibrated at 25°C). In preferred embodiments, the initial percent oxygen saturation in the IL-17 antibody preparation is adjusted to at least 80% (as measured using an oxygen probe calibrated at 25°C).

We have determined that elevated oxygen levels during the cysteine treatment step can have a deleterious effect on antibody activity, which is likely due to the oxygen abrogating the reductive power of the cysteine, leading to insufficient reduction of C97 of secukinumab. This issue of oxygen uptake from the atmosphere can be managed by varying the amount of reducing agent, varying the molar ratio of reducing agent:antibody, using defined stirring speeds, or even employing stirring interruptions, especially when working at production scale. It will be understood that higher amounts of reducing agent (e.g., cysteine) are capable of handling higher oxygen levels in the reducing mixture. In some embodiments, during the incubation step of the selective reduction process, the oxygen saturation in the reducing mixture is generally maintained at a low percentage (e.g., less than about 40%, less than about 30%, less than about 20%, less than about 15%, less than about 10%, less than about 5%).

A loss of reductive power of the reducing agent (e.g., cysteine) likely leads to incomplete deblocking of CysL97-SH at earlier time points during the incubation step, and if there is no residual reducing agent (e.g., cysteine) available to protect deblocked Cys97L at later portions of the incubation step (or during the cooling step), then reoxidation of deblocked Cys97L-SH can occur. Therefore, ideally, a low percentage oxygen saturation will be maintained for at least about 60 minutes to about 330 minutes, e.g., at least about 60 minutes, at least about 90 minutes, at least about 120 minutes, at least about 150 minutes, at least about 180 minutes, at least about 210 minutes, at least about 240 minutes, at least about 270 minutes, at least about 300 minutes, or at least about 330 minutes. In some embodiments, this low percentage oxygen saturation will be maintained for the full incubation step, as well as part of the cooling step.

Percent oxygen saturation can be adjusted directly (e.g., by sparging) or indirectly (e.g., by stirring) to achieve a desired level of oxygen during incubation of the reducing mixture. In an

aerobic environment, a low percent oxygen saturation may be achieved by using intermittent (rather than continuous) mixing of the reducing solution, e.g., < 15 min/hr, e.g., <2 min/hr, or by using continuous stirring with a low spin speed. In an anaerobic environment, no (or little) oxygen is present that would lead to consumption of the reducing agent.

Volumetric oxygen mass-transfer (k_{La}^*)

When selective reduction is performed under aerobic conditions, the level of oxygen in the reaction is not controlled directly, but via other process parameters, e.g., stir speed. The physical setup of each reaction also influences the level of oxygen present in the reaction mixture. Therefore, it is important to identify a variable that can be used to compare the oxygen transferred into a solution between physical setups and during particular antibody processing steps – that variable is “ k_{La}^* ” (see, e.g., Garcia-Ochoa and Gomez (2009) *Biotechnology Advances* 27:153-176; Bandino et al. (2001) *Biochem. Engineering J.* 8:111-119; Juarez and Orejas (2001) *Latin Am. Appl. Res.* 31:433-439; Yange and Wang (1992) *Biotechnol. Prog.* 8:244-61). The k_{La}^* represents the amount of oxygen transferred into a solution over time via the headspace without sparging. This value is specific for each setup and scale, and depends on stirrer type, stirrer speed, filling volume and surface area of the solution in contact with the headspace, which is influenced by the individual geometry of each vessel. While the k_{La}^* of each physical setup differs, because the level of oxygen in the solution during the selective reduction process affects the activity and integrity of secukinumab, we expect that the selective reduction process, when performed in systems displaying similar k_{La}^* ranges, will lead to preparations of secukinumab having similar quality.

As used herein the term “system” encompasses both the physical setup (vessel, stir type, etc.) and the process conditions (fill volume, spin speed, etc.) that influence the oxygen transfer into a solution over time via the headspace without sparging, i.e., scale, stirrer speed, filling volume, surface area of the solution in contact with the headspace, which is influenced by the individual geometry of each vessel, etc.

As used herein the term “vessel” means any container in which the selective reduction reaction takes place. Vessels include, without limitation, bioreactors (e.g., steel, stirred tank, disposable or non-disposable, etc.) used for pilot and commercial scale antibody production, as well as common laboratory containers, such as flasks, tubes, etc. In some embodiments, the

vessel is a bioreactor capable of holding a volume of at least about 2 liters, at least about 100 liters, at least about 500 liters, at least about 1000 liters, at least about 2000 liters, at least about 5000 liters, at least about 10,000 liters, at least about 15,000 liters or greater.

The k_{La}^* cannot be directly determined in the oxygen transfer experiments. Instead, the dO_2 in a test solution is replaced by nitrogen and the increase of dO_2 over time is monitored using a calibrated dO_2 probe, which allows creation of an experimental dO_2 curve. Thereafter, the k_{La}^* values used herein are calculated for the particular systems by adapting the experimental dO_2 curve to a saturation curve (e.g., using Mathcad®) according the equation shown below:

$DO = C \times (1 - e^{-k_{La}^* \times (t - t_0)})$, where DO = the measured value of dissolved oxygen, C is the saturation value of oxygen (meaning 100 % when stirred infinitely and saturation is achieved), Euler's number $e = 2.718281\dots$, t = time point corresponding to the DO value, and t_0 = starting time point.

The equation represents the integrated form of an empirical formula established for determination of the oxygen transfer into solutions (k_{La}^* value). The formula was confirmed by different authors in various experiments (Doran, P. M. 1995. Bioprocess Engineering Principles, Academic Press, San Diego, California).

We have determined that the heating and cooling steps of the selective reduction process are generally more tolerable of a higher k_{La}^* than the incubation step. This is because elevated oxygen levels during the incubation step can have a deleterious effect on antibody activity, which is likely due to the increased oxygen transfer abrogating the reductive power of the reducing agent (e.g., cysteine), leading to insufficient reduction of C97 of secukinumab. In an aerobic environment, increasing the stir speed (or stir time) in the reducing mixture during the selective reduction process increases the k_{La}^* . Continuous stirring of the reducing mixture, which leads to higher k_{La}^* values, can be tolerated during the heating and cooling steps. In some embodiments, the k_{La}^* during the heating and cooling steps (separately) can be from about 0.12 h^{-1} to about 1.69 h^{-1} , about 0.08 h^{-1} to about 0.69 h^{-1} , about 0.24 h^{-1} to about 0.44 h^{-1} , about 0.39 h^{-1} to about 0.69 h^{-1} . In some embodiments, the k_{La}^* in the system during the heating or cooling step is $\leq$ about 0.69 h^{-1} , said k_{La}^* being calculated by adapting a saturation curve to a dissolved oxygen curve.

It is preferable to keep the k_{La}^* lower during the incubation step using, e.g., continuous stirring with low spin speeds, or, preferably, intermittent stirring. In some embodiments, the

reducing mixture is incubated while maintaining a volumetric oxygen mass-transfer coefficient (k_{La}^*) in the system of $\leq$ about 0.37 h^{-1} , $\leq$ about 0.27 h^{-1} , or $\leq$ about 0.18 h^{-1} , said k_{La}^* being calculated by adapting a dissolved oxygen curve to a saturation curve. If the k_{La}^* in the system during the incubation step of the selective reduction reaction is less than ($<$) 0.37 h^{-1} , then the molar ratio of reducing agent (e.g., cysteine):antibody can vary between 46.11:1 (about 46:1) to 118.36:1 (about 118:1) (for both shorter and longer incubation times, e.g., about 210 to about 330 minutes). The k_{La}^* in the system during the incubation step of the selective reduction reaction can be as high as ($\leq$) 0.37 h^{-1} if the molar ratio of reducing agent (e.g., cysteine):protein is between 69.89:1 (about 70:1) to 118.36:1 (about 118:1) (for shorter incubation times, e.g., up to about 240 minute incubation) or between 76.85 (about 77:1) to 118.36:1 (about 118:1) (for longer incubation times, e.g., up to about 300 minute incubation).

The entire selective reduction process (i.e., heating step, incubation step, and cooling step) can be generally performed using a k_{La}^* of $< 0.37\text{ h}^{-1}$, which includes about 210 - about 330 minute incubation time (e.g., about 240 minute - 300 minute incubation time), if the molar ratio of reducing agent (e.g., cysteine):antibody is between about 46:1 - about 118:1. The entire selective reduction process (i.e., heating step, incubation step, and cooling step) can be generally performed using a $k_{La}^* \leq 0.37\text{ h}^{-1}$ if the molar ratio of reducing agent (e.g., cysteine):antibody is between about 70:1 - about 118:1, and incubation is up to about 240 minutes; or if the molar ratio of reducing agent (e.g., cysteine):antibody is between about 77:1 - about 118:1, and incubation is up to about 300 minutes.

Further Process Components

The present disclosure provides a method for selective reduction of CysL97 in certain IL-17 antibodies, such as secukinumab, comprising contacting the antibody with a reducing agent to form a reducing mixture. It will be understood that the reducing mixture may comprise components in addition to the IL-17 antibody or antigen binding fragment thereof and the reducing agent. The reducing mixture may contain an aqueous component (e.g., a buffer, such as a Tris buffer), as well as reagents used to increase or decrease the pH of the reducing mixture, salt, EDTA, etc. Thus, while the reducing mixture will necessarily contain the IL-17 antibody (e.g., secukinumab) and the reducing agent, the reducing mixture may or may not contain additional components. In some embodiments, the reducing mixture contains a metal chelator

(e.g., EDTA, DMSA, DMPS). In some embodiments, the reducing mixture contains 1.3 mM to about 0.8 mM, about 1.1 to about 0.9 mM, or about 1.0 mM EDTA (e.g., di-Na-EDTA).

Depending on the reaction conditions (temperature, length of reaction time, quantity of IL-17 antibody, concentration of reducing agent, ratio reducing agent:antibody, etc.) the pH of a particular antibody preparation may vary. However, it is recognized that, based upon the reaction conditions described herein, such conditions can be varied in order to achieve the desired selective reduction. In some embodiments, the pH of the antibody preparation prior to contact with the reducing agent will vary from about 6.5 to about 9.5, e.g., about 7 to about 9. In other embodiments, the pH of the antibody preparation will be about 7.4 to about 8.5, about 7.8 to about 8.2, about 7.9 to about 8.1, or about 8.0. In some embodiments, the pH of the antibody preparation may be adjusted (e.g., following gassing to adjust the initial percent oxygen saturation) using a buffer, e.g., a 1M Tris buffer (e.g., 1M Tris buffer pH 10.8).

Following contacting of the preparation containing the IL-17 antibody or antigen binding fragment thereof (e.g., secukinumab) with the reducing agent, there may be an initial reduction in the level of intact IL-17 antibody or antigen binding fragment thereof (HLLH). The term "intact" refers to an antibody having all conserved disulfide bridges (e.g., 14 conserved disulfide bridges in the case of a classical IgG₁ antibody). The formation of various IL-17 antibody fragments, i.e., H₂L, HL₂, HL, H and L bands, as well as the intact H₂L₂ (HHLL) band can be determined using different analytical tools known in the art (such as SDS-PAGE, Cation-exchange HPLC). Preferably, the level of intact IL-17 antibody or antigen binding fragment thereof in the mixture is measured by sodium dodecyl sulfate capillary electrophoresis (CE-SDS). CE-SDS separates proteins according to their molecular size in an electric field. Non-reducing CE-SDS can be used to assess size variants in a preparation of antibodies. In some embodiments, the level of intact IL-17 antibody or antigen binding fragment thereof in the mixture decreases to at least about 80%, as measured by CE-SDS, within about 1-30 minutes (e.g., about 15 minutes) of addition of the reducing agent to the antibody preparation. In some embodiments, the level of intact IL-17 antibody or antigen binding fragment thereof in the mixture decreases to at least about 83%, as measured by CE-SDS. In some embodiments, the level of intact IL-17 antibody or antigen binding fragment thereof in the mixture decreases to between about 75% to about 87%. In some embodiments, the level of intact IL-17 antibody or antigen binding fragment thereof in the mixture decreases to at least about 38, 39, 40, 41, 45, 50,

55, 60, 65, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, or 87%, as measured by CE-SDS.

CE-SDS analyses may be performed using a Beckman Coulter PA-800 capillary electrophoresis system. Uncoated fused-silica capillaries with an inner diameter of 50 μm and a length of 30 cm (with 20 cm and 10 cm separation ranges for reducing CE-SDS and non-reducing CE-SDS analyses, respectively) are used for the analyses. The separation is monitored with a UV detector at 214 nm. For non-reducing CE-SDS analyses, antibody samples are diluted to 6.0 mg/mL with water, mixed thoroughly with non-reducing CE-SDS sample buffer (0.1 M Tris/1.0% SDS, pH 7.0) and 250 mM iodoacetamide, at a ratio of 20/75/5 (v/v/v), and heated at 70°C for 10 minutes to prevent disulfide bridge shuffling. The capillary temperature is set at 25°C for the separation. The electrophoresis is carried out at a constant voltage of 15 kV in the normal polarity mode for 20 minutes.

In some embodiments, the reducing mixture is heated. In some embodiments, heating occurs prior to the step of incubating. In some embodiments, the reducing mixture is heated to a temperature between about 32°C to about 42°C, to between about 35°C to about 39°C, to about 37°C. In some embodiments, the heating occurs for about 30 to about 120 minutes, about 45 to about 90 minutes, about 45 to about 75 minutes, about 60 minutes. During heating, the reducing mixture may be stirred, e.g., constantly or intermittently, using any means for stirring. Stirring may be axial (e.g., using a pitched blade impeller) or radial (e.g., using a rushton turbine). In some embodiments, during heating the reducing mixture is constantly stirred at 65-200 rpm (e.g., 50 rpm, 65 rpm, 75 rpm, 85 rpm, 100 rpm or 200 rpm).

During the incubating step, the reducing mixture will typically be incubated for a predetermined time to allow selective reduction of the free cysteine (e.g., the free cysteine of secukinumab). In some embodiments, incubation occurs following heating of the reducing mixture. Depending on the reaction conditions (reducing agent, temperature, quantity of IL-17 antibody or antigen binding fragment thereof, pH, etc.) the predetermined time for incubation of the reducing mixture will vary. In some embodiments, the time will vary between about 1 and 24 hours (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 16, 18, 20 or 24 hours). In some embodiments, incubating is performed for about 200 to about 500 minutes, about 210 to about 420 min., about 210 to about 330 minutes, about 240 to about 300 minutes, about 250 minutes.

During the incubating step, incubation will be performed at a predetermined temperature

to allow selective reduction of the target free cysteine (e.g., CysL97 in the antigen binding sites of secukinumab). Depending on the reaction conditions (reducing agent, time, quantity of IL-17 antibody or antigen binding fragment thereof, pH, etc.) the incubation temperature will vary. In some embodiments, the predetermined temperature will vary between about 20 to about 42°C. In some embodiments, the predetermined temperature will be about 32°C - about 42°C, between about 35°C - about 39°C, or about 37°C.

During the incubating step, the reducing mixture may be stirred to ensure product homogeneity while the reductant is incubated with the IL-17 antibody preparation, e.g., secukinumab. However, the level of oxygen in the vessel should be kept low during this portion of the selective reduction reaction in order to allow the reducing agent to effectively selectively reduce CysL97 in the IL-17 antibodies. Under aerobic conditions, low oxygen may be achieved by avoiding continuous stirring, e.g., by using intermittent stirring, e.g., ≤ 15 min/hr, e.g., ≤ 2 min/ hr. Under anaerobic conditions, transfer of oxygen is limited and therefore stirring may proceed for longer periods of time or may be continuous. Moreover, under anaerobic conditions, strict control of oxygen transfer (e.g., by regulated sparging) would also allow application of longer periods of stir time (including continuous stirring).

In some embodiments, the mixture is cooled following the incubating step. In some embodiments, the mixture is cooled to room temperature (e.g., a temperature between about 16°C to about 28°C). In some embodiments, cooling occurs for about 30 to about 120 minutes, about 45 to about 90 minutes, about 45 to about 75 minutes, about 60 minutes. During cooling, the reducing mixture may be stirred, e.g., constantly or intermittently using any means for stirring. Stirring may be axial (e.g., using a pitched blade impeller) or radial (e.g., using a rushton turbine). In some embodiments, during cooling the reducing mixture is constantly stirred at 65-200 rpm (e.g., 50 rpm, 65 rpm, 75 rpm, 85 rpm, 100 rpm or 200 rpm).

The selective reduction reaction may be quenched, e.g., using iodoacetamid or o-phosphoric acid (e.g., using a stock solution of 0.3 M o-phosphoric acid). In some embodiments, quenching occurs following the cooling step. In some embodiments, the selective reduction reaction is quenched by adjusting the pH of the mixture to between about 5.0 to about 5.5, about 5.1 to about 5.3, about 5.2. pH adjustment may be achieved using o-phosphoric acid.

The phrase “purified preparation” refers to a mixture of IL-17 antibodies or antigen binding fragments thereof that have been subjected to selective reduction. After completion of

selective reduction, there will be an increase in the level of intact IL-17 antibody in the purified preparation. The level of intact antibodies in the purified preparation after selective reduction may be measured via various well known techniques (e.g., non-reducing SDS PAGE, CE-SDS PAGE, size exclusion chromatography (SEC), HPLC). In some embodiments, the level of intact antibody is measured by CE-SDS. In some embodiments, after completion of selective reduction (e.g., after the cooling step), the level of intact IL-17 antibody or antigen binding fragment thereof in the purified preparation is at least about 80%, as measured by CE-SDS. In some embodiments the level of intact IL-17 antibody or antigen binding fragment thereof in the purified preparation is at least about 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or about 100 %, as measured by CE-SDS, after selective reduction. In some embodiments, the level of intact IL-17 antibody or antigen binding fragment thereof in the purified preparation is at least about 90%, as measured by CE-SDS, after selective reduction.

The activity (e.g., affinity, biological activity, etc.) of the antibodies in a preparation prior to selective reduction, during selective reduction or after selective reduction (i.e., in a purified preparation) may be measured via various well known techniques (see, e.g., WO2006/013107; WO2007/117749; Shen and Gaffen (2008) Cytokine. 41(2): 92-104). In certain embodiments, the activity is measured using an ELISA based assay or a cell-based assay (e.g., inhibition of IL-17 dependent release of IL-6 or GRO α from, e.g., C-20/A4 chondrocytes or BJ cell line). In some embodiments, activity is measured by a cystamine-CEX (cation exchange chromatography) method. The cystamine-CEX method includes derivatization of the antibody with cystamine (2,2'-dithiobis(ethylamine)), followed by analytical separation using cation exchange chromatography (CEX). Because the activity of the antibodies disclosed herein (e.g., secukinumab) is decreased if CysL97 is in oxidized form, derivatization of CysL97 with cystamine serves as a proxy to measure antibody activity, i.e., if selective reduction succeeds then reduced CysL97 can be derivitized with cystamine, whereas if selective reduction fails then oxidized CysL97 cannot be derivitized with cystamine. Derivatization by cystamine leads to an addition of one positive charge per free Cys97 residue. The resulting derivitized forms of secukinumab (e.g., +2, +1 charges) can then be separated from the non-derivitized form and quantified by CEX. A cystamine-derivitized secukinumab molecule with two cystamine bound to unpaired Cys97 on both light chains may be considered 100% biological active in theory. A cystamine-derivitized secukinumab molecule with addition of one cystamine bound to unpaired

Cys97 on one of the light chains may be considered 50% biological active. A cystamine-derivatized secukinumab molecule without any cystamine bound to the molecule may be considered biological inactive. The level of cystamine derivitization in a preparation of antibodies (e.g., a preparation of secukinumab antibodies), in comparison to the theoretical maximum level of cystamine derivitization in that preparation (e.g., expressed as a percentage of theoretical maximum) may then be used as a measure of the activity of the preparation.

In brief cystamine-CEX may be performed as follows. Antibody samples (50 µg) are first treated with carboxypeptidase B (1:40, w:w) to remove the C-terminal lysine in the heavy chain and then derivatized with 4 mM cystamine in 5 mM sodium acetate, 0.5 mM EDTA, pH4.7 at room temperature for 2 hours. The derivatization is stopped by addition of 2 µL of 1M phosphoric acid. CEX is performed on the cystamine-derivatized antibody samples using a ProPac™ WCX-10 analytical column (4 mm x 250 mm, Dionex). A gradient from 12.5 mM to 92.5 mM sodium chloride in 25 mM sodium phosphate, pH 6.0 at a flow rate of 1.0 ml/min is used for separation. Absorption at 220 nm is recorded by a UV detector (Agilent HPLC 1200).

Some initial preparations of IL-17 antibody with an oxidized free cysteine have activity levels as low as 45%. In some embodiments, prior to initiating the selective reduction process, the level of activity of the IL-17 antibodies or antigen binding fragments thereof in the preparation is less than about 80%, less than about 75%, less than about 70%, less than about 65%, less than about 60%, less than about 55%, less than about 50%, or less than about 45% (e.g., as measured by the cystamine-CEX method). During selective reduction, there will be an increase in the level of activity of the IL-17 antibodies in the purified preparation. In some embodiments, the level of activity of the IL-17 antibodies or antigen binding fragments thereof in the antibody preparation increases by at least about 15 percentage points (e.g., from about 60% to at least about 75%), at least about 20 percentage points (e.g., from about 60% to at least about 80%), at least about 25 percentage points (e.g., from about 60% to at least about 85%) or at least about 30 percentage points (e.g., from about 60% to at least about 90%) within about 60 minutes following contacting the antibody preparation with the reducing agent to form the reducing mixture (e.g., as measured by the cystamine-CEX method).

After selective reduction, the purified preparation will be enriched for IL-17 antibodies having the reduced form of CysL97 and will display an increased level of activity relative to the initial preparation. In some embodiments, after completion of selective reduction, the level of

activity of the IL-17 antibodies or antigen binding fragments thereof in the purified preparation is at least about 80% (relative to a theoretical maximum), as measured by cystamine-CEX, an ELISA, or cell-based binding assay (e.g., a cystamine-CEX assay). In some embodiments, after completion of selective reduction, the level of activity of the IL-17 antibodies or antigen binding fragments thereof in the purified preparation is at least about 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or about 100 %, as measured by CEX, an ELISA, or cell-based binding assay (e.g., a cystamine-CEX assay). In some embodiments, after completion of selective reduction, the level of activity of the IL-17 antibodies or antigen binding fragments thereof in the purified preparation is at least about 93%, as measured by cystamine-CEX, an ELISA, or cell-based binding assay (e.g., a cystamine-CEX assay).

Accordingly, disclosed herein are methods for selectively reducing CysL97 in a preparation of IL-17 antibodies that have been recombinantly produced by mammalian cells, comprising:

- a) contacting the preparation with at least one reducing agent in a system to form a reducing mixture; and
- b) incubating the reducing mixture while maintaining a volumetric oxygen mass-transfer coefficient (k_{La}^*) in the system of $\leq$ about 0.37 h^{-1} , said k_{La}^* being calculated by adapting a dissolved oxygen curve to a saturation curve;

wherein the IL-17 antibodies each comprise an immunoglobulin heavy chain variable domain (V_H) comprising the three complementarity determining regions (CDRs) of the V_H set forth as SEQ ID NO:8 and an immunoglobulin light chain variable domain (V_L) comprising the three CDRs of the V_L set forth as SEQ ID NO:10, and further wherein prior to step a) the initial percent oxygen saturation in the preparation is at least about 60%, as measured using an oxygen probe calibrated at 25°C.

Also disclosed herein are methods for selectively reducing CysL97 in a preparation of IL-17 antibodies that have been recombinantly produced by mammalian cells, comprising:

- a) contacting the preparation with a set of oxidation/reduction reagents selected from cysteine/cystine and cysteine/cystamine to form a reducing mixture; and
- b) incubating the reducing mixture at a temperature of about 37 °C under anaerobic conditions for at least about 4 hours, or incubating the reducing mixture at a temperature of about 18-24 °C for about 16-24 hours;

wherein the IL-17 antibodies each comprise an immunoglobulin heavy chain variable domain (V_H) comprising the three complementarity determining regions (CDRs) of the V_H set forth as SEQ ID NO:8 and an immunoglobulin light chain variable domain (V_L) comprising the three CDRs of the V_L set forth as SEQ ID NO:10.

Also disclosed herein are methods for selectively reducing CysL97 in a preparation of secukinumab antibodies that have been recombinantly produced by mammalian cells, comprising:

- a) adjusting the concentration of secukinumab in the preparation to between about 4 mg/ml – about 19.4 mg/ml, e.g., about 10 mg/ml - about 19.4 mg/ml, e.g., about 10 - about 15.4, e.g., about 12 mg/ml - about 15 mg/ml, e.g., about 13.5 mg/ml;
- b) adjusting the percent oxygen saturation in the preparation to at least about 60%, e.g., at least about 80%;
- c) adjusting the pH of the preparation to about 7.4 - about 8.5, e.g., about 7.8 - about 8.2, e.g., about 7.9 - about 8.1, e.g., about 8.0;
- d) contacting the preparation with cysteine in a vessel to form a reducing mixture, wherein the concentration of cysteine in the reducing mixture is about 4.0 mM - about 8.0 mM, e.g., about 4.8 mM - about 8.0 mM, e.g., about 5.5 mM - about 6.7 mM, e.g., about 6.0 mM;
- e) heating the reducing mixture to a temperature between about 32°C - about 42°C, e.g., to between about 35°C - about 39°C, e.g., to about 37°C, said heating occurring for about 45 - about 90 minutes, e.g., about 45 - about 75 minutes, e.g., about 60 minutes;
- f) incubating the reducing mixture from step e) at a temperature between about 20°C - about 42°C, e.g., 32°C - about 42°C, e.g., to between about 35°C - about 39°C, e.g., to about 37°C, said incubating occurring for about 210 - about 420 minutes, e.g., about 210 - about 330 minutes, e.g., about 240 - about 300 minutes, e.g., about 250 minutes while maintaining a volumetric oxygen mass-transfer coefficient (k_{La}^*) in the vessel of $\leq 0.37 \text{ h}^{-1}$, said k_{La}^* being calculated by adapting a saturation curve to a dissolved oxygen curve,
- g) cooling the mixture resultant from step f) to a temperature between about 16°C - about 28°C, said cooling occurring for about 45 - about 90 minutes, e.g., about 45 - about 75 minutes, e.g., about 60 minutes; and
- h) adjusting the pH of the mixture resultant from step g) to between about 5.1 - about 5.3, e.g., about 5.2.

Also disclosed herein are also purified preparations of secukinumab, wherein the level of intact secukinumab in the preparation is at least about 90%, as measured by sodium dodecyl sulfate capillary electrophoresis (CE-SDS), and wherein the level of activity of secukinumab in the preparation is at least about 90%, as measured by cystamine-CEX.

General

In some embodiments of the above methods, the IL-17 antibody or antigen binding fragment thereof comprises: i) an immunoglobulin heavy chain variable domain (VH) comprising the amino acid sequence set forth as SEQ ID NO:8; ii) an immunoglobulin light chain variable domain (VL) comprising the amino acid sequence set forth as SEQ ID NO:10; iii) an immunoglobulin VH domain comprising the amino acid sequence set forth as SEQ ID NO:8 and an immunoglobulin VL domain comprising the amino acid sequence set forth as SEQ ID NO:10; iv) an immunoglobulin VH domain comprising, in sequence, the hypervariable regions set forth as SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:3; v) an immunoglobulin VL domain comprising, in sequence, the hypervariable regions set forth as SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6; vi) an immunoglobulin VH domain comprising, in sequence, the hypervariable regions set forth as SEQ ID NO:11, SEQ ID NO:12 and SEQ ID NO:13; vii) an immunoglobulin VH domain comprising, in sequence, the hypervariable regions set forth as SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:3 and an immunoglobulin VL domain comprising, in sequence, the hypervariable regions set forth as SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6; and viii) an immunoglobulin VH domain comprising, in sequence, the hypervariable regions set forth as SEQ ID NO:11, SEQ ID NO:12 and SEQ ID NO:13 and an immunoglobulin VL domain comprising, in sequence, the hypervariable regions set forth as SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6. In some embodiments of the disclosed methods, the IL-17 antibody or antigen binding fragment thereof is a human antibody of the IgG₁ isotype. In some embodiments of the disclosed methods, the antibody is secukinumab.

The details of one or more embodiments of the disclosure are set forth in the accompanying description above. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, the preferred methods and materials are now described. Other features, objects, and advantages of the disclosure will be apparent from the description and from the claims. In the specification and

the appended claims, the singular forms include plural referents unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. All patents and publications cited in this specification are incorporated by reference. The following Examples are presented in order to more fully illustrate the preferred embodiments of the disclosure. These examples should in no way be construed as limiting the scope of the disclosed patient matter, as defined by the appended claims.

EXAMPLES

Example 1: Reduction of Secukinumab Protein A intermediate by different sulfhydryl agents

Example 1.1

A variety of sulfhydryl group-containing reducing agents (e.g., dithiothreitol (DTT), 2-mercaptoethanol, 2-mercaptoacetic acid, cysteine, cysteamine, glutathione) were screened for their use in selectively reducing the oxidized Cys97 in the light chain of secukinumab. In a first set of experiments, 1-mL portions of inactive starting material obtained from an early fermentation process after the Protein A capture step were incubated at 37°C at various pH with various concentrations of β -mercaptoethanol, cysteine and glutathione. After certain time-points, the samples were desalted by gel filtration into 20mM acetate pH 6 buffer and restoration of activity was determined by an ELISA. Also, the content of free sulfhydryl groups was determined by Ellman test (deblocking of the sulfhydryl group should result in a value of 2 Mol free-SH per mol antibody). The results are listed in the **Table 2**, below.

2-ME°	Activity % *	Free SH Mol/Mol	Cysteine	Activity % *	Free SH Mol/Mol	Glutathione	Activity % *	Free SH Mol/Mol
pH 7								
37°C 5mM/1h	59	1.3						
	/4h 85	1.6						
	/8h 85	1.6						
pH8								
37°C			1mM/2h	61	1.4			
			/8h	90	1.8			
2mM/1h	64	1.4	2mM/2h	80	1.8	2mM/2h	45	1.4
/4h	82	1.9	/4h	98	1.9	/8h	65	2.0
/8h	88	n.p.						

4mM/1h	88	1.3	4mM/1h	77	0.9	4mM/2h	59	1.6
/4h	91	1.8	/4h	105	1.9	/4h	94	1.8
/8h	106	2.0						
pH 9 37°C						8mM/1h	58	1.4
						/4h	75	1.7
			1mM/2h	48	1.2			
			/8h	71	1.5			
2mM/1h	82	n.p.	2mM/2h	78	1.6	2mM/2h	73	1.6
/4h	97	0.9	/4h	89	1.8	/8h	81	2.1
/8h	93	1.1						
4mM/1h	85	0.7	4mM/1h	74	1.5	4mM/2h	54	2.0
/4h	102	2.0	/4h	100	2.1	/4h	73	2.1
						8mM/1h	68	1.7
						/4h	81	2.1


Table 2: Activity and free-SH of samples following reaction at given conditions.

°: 2-Mercaptoethanol (2-ME)

*: % relative to reference

As a result of these studies, mercaptoethanol and cysteine in the pH range 7-9 at concentration of 1 to 20 mM and temperatures of 20-40°C were found to be most suitable.

In the course of further investigations, it was realized that antibody can be overreduced during exposure to the reducing agents with formation of antibody having partially reduced interchain disulfide bridges. This overreduction is reversible at atmospheric conditions when the antibody is isolated from the reaction mixture, e.g., by diafiltration as described above or by chromatography, because dissolved oxygen present in the buffers leads to spontaneous re-oxidation with re-formation of intact antibody.



Example 1.2 (anaerobic conditions)

In a second set of experiments, the treatments of secukinumab antibody obtained after the Protein A capture step were performed under anaerobic conditions by using de-aerated solutions and argon or nitrogen atmosphere to exclude effects of dissolved oxygen on the reduction outcome. In brief, a secukinumab solution from a Protein A capture step was adjusted to pH 8.0 by addition of a Tris base stock solution. The solution was then adjusted to a concentration of 8

mM cysteine and 1 mM EDTA by addition of a cysteine/EDTA stock solution (e.g., 200mM/12.5 mM EDTA at pH 8). The concentration of secukinumab in the reducing solution was 4 mg/mL (molar ratio of cysteine:antibody about 296:1). The mixture was incubated at room temperature for a period of 24 hours. At different time-points, samples were drawn and spiked with an excess of iodoacetamid, which stops the reaction by quenching sulhydryl groups of the reducing agent and antibody.

The same set-up was used for the experiments with 2-mercaptoethanol, 2-mercaptoacetic acid, cysteine, cysteamine and glutathione. For DTT, a concentration of 1mM was used.

The quenched samples were analyzed by capillary electrophoreses with SDS in the non-reduced mode (CE-SDS). This analytical method separates the different reduction products of antibody (HHL, HH, HL, H and L species) from intact antibody (LHHL) and quantifies them by on-line UV detection and area-integration of the obtained signals.

In **Figure 1**, the data for intact antibody (LHHL) from 0, 3, 15 min, 1h, 2h, 4h and 20-24h reduction treatment shows that DTT and β -mercapto acetic acid reduce the antibody completely without re-equilibration to intact antibody. Glutathione and β -mercaptoethanol also show pronounced reduction, but were able to induce a re-equilibration to intact antibody (LHHL). Cysteine reduces the antibody to about 50%, which is followed by a straight-forward re-equilibration to intact antibody. The data for cysteamine shows that this reagent either exerts little reduction or leads very quickly (within a time of less than 3min) to re-equilibration.

The reduction order found (i.e., DTT > β -mercapto acetic acid > β -mercaptoethanol > glutathione) correlates to published data for redox potentials or disulfide interchange. see *T. Liu in "The Proteins, 3rd Edition, Volume 3 (1977) p. 239* which gives following comments and data:

$$\text{Redox potential } E = E_0 + 0.059 \times \log [\text{R-SS-R}]/[\text{R-SH}]$$

Standard redox potential (E_0 in Volt) not directly measurable as electrode is poisoned by the sulfur. Therefore, redox potential had to be deduced from indirect measurements:

Dithiothreitol DTT/DTTox :	$E_0 \sim -0.33 \text{ V (pH 7, 25}^\circ\text{C)}$
Glutathione GSH/GSSG :	$E_0 \sim -0.24 \text{ V (pH 7, 25}^\circ\text{C)}$
Cysteine CSH/CSSC :	$E_0 \sim -0.22 \text{ V (pH 7, 25}^\circ\text{C)}$

Example 2: Cysteine Selectively Reduces Secukinumab Under Aerobic and Anaerobic Conditions.

Additional experiments were performed at room temperature (RT) (about 25°C) or 37°C using 8 mM cysteine, 4 mg/ml secukinumab, pH 8 (molar ratio of cysteine:antibody about 295.88:1). Reactions were performed under either aerobic (i.e., preparation of the solutions and the treatment was carried out under normal air) or anaerobic (i.e., preparation of the solutions and the treatment was carried out under exclusion or reduction of oxygen by de-aeration and subsequent working under argon or nitrogen atmosphere) conditions.

Under anaerobic conditions (e.g., nitrogen or argon atmosphere with 0% dissolved oxygen), in the early phase of the treatment there is a substantial degradation of antibody with formation of antibody fragments - a sign of over-reduction. The maximal decrease of intact antibody (LHHL) to about 40% occurs at around 15 minutes for experiments performed at either RT (**Figure 2A**) or 37°C (**Figure 2C**). Re-equilibration to intact antibody at 20 hours shows about 17.1% over-reduced variants remaining in the RT samples and 12.1% over-reduced variants remaining in the 37°C samples.

Under aerobic conditions, i.e., in presence of dissolved oxygen, there is a more moderate reaction. The maximal decrease of intact antibody (LHHL) to about 80% residual level occurs at around 15 minutes for experiments performed at RT (**Figure 2B**) and to about 73% for experiments performed at 37°C (**Figure 2D**). Re-equilibration to intact antibody shows only 5.8% over-reduced variants remaining in the RT samples after 20 hours and only 9.3% over-reduced variants remaining in the 37°C samples after 8 hours.

Thus, under anaerobic conditions the initial reduction is larger and re-equilibration to intact antibody is not as complete as under aerobic conditions.

Example 3: Influence of Dissolved Oxygen and Cystine on Selective Reduction

Because cysteine (Cys-SH) oxidizes in presence of air to cystine (Cys-SS-Cys), the difference observed under aerobic and anaerobic conditions could be caused by small amounts of cystine formed during preparation of cysteine solutions under aerobic conditions and also during the actual treatment of secukinumab with cysteine. In order to assess the influence of the level of

dO₂ and cystine (or cystamine) on selective reduction of secukinumab, we performed several additional preparative experiments with isolation of the antibody.

For some samples, selective reduction of 4 mg/ml secukinumab was performed with 8 mM cysteine (molar ratio of cysteine:antibody 295.88:1) in a solution having a pH 8.0 at 37°C under aerobic conditions at 100% dO₂ (no fumigation by nitrogen), 50%, and 20% dO₂ (stirring at ambient atmosphere with fumigation by nitrogen such that dissolved oxygen stayed at target level throughout the experiment), and anaerobic (0% dO₂) conditions (degassed solutions, full nitrogen atmosphere) without cystine, as well as under 100% dO₂ in presence of about 0.1 mM cystine [ratio cysteine:cystine = 80:1]. In one experiment, 0.3 mM cystine was added following an initial 30 minute incubation with 8 mM cysteine under anaerobic conditions. In another experiment, selective reduction was performed under anaerobic conditions using 7.7 mM cysteine and 0.3 mM cystine [ratio cysteine:cystine = 25.66:1]. In another sample, 0.1 mM cystamine was added instead of cystine.

In all these examples, samples were drawn at different time points. At the end of the treatment (240 min at 37°C), the antibody was isolated by adjusting the pH of the bulk solution to 5.0 and loading onto a cation-exchange column that binds the antibody. After a wash to remove the reducing agent, antibody was eluted by a salt and/or pH-gradient and analyzed by CE-SDS, SE-HPLC and CEX.

Figure 3 shows CE-SDS analysis of iodacetamid quenched samples drawn at different time of the treatment. Less dO₂ leads to greater reduction in the early phase and slower equilibration in the later phase of selective reduction. Level of intact antibody was substantially lower (70%) at the end of treatment in total anaerobic conditions (0% dO₂). Level of intact antibody was improved to above 90% when the reaction was performed under conditions of 50% or more dissolved oxygen. Addition of a small amount of cystine (or cysteamine) decreased the initial reduction of antibody.

Table 3 shows activity and purity of antibody obtained after the samples from different reactivation treatments (anaerobic and aerobic conditions) were purified using subsequent chromatography on SP-Sepharose FF.

Treatment conditions (37°C/pH 8.0/4h)	after SP- step			
	CE-SDS purity	SEC purity	CEX activity	Bioactivity

	%	%	%	%
Starting material	96.4	98.4	61.2	45
8mM cysteine/ 0% oxygene (anaerob)	82.3	98.1	92.8	92
8mM cysteine/0% oxygene /30min: + 0.3mM cystine	92.6	98.4	92.3	95
8mM cysteine/ 20% oxygene	96.9	98.0	88.6	99
8mM cysteine/ 50% oxygene	96.1	98.2	87.0	90
8mM cysteine/100% oxygene (aerob)	95.8	98.1	90.6	96
7.7mM cysteine/0.3mM cystine/anaerob	97.1	98.9	91.8	104
8mM cysteine/0.1mM cystine 100% oxygene (aerob)	94.5	98.0	82.4	108
8mM cysteine/0.1mM cystamine 100% oxygene (aerob)	94.8	98.6	86.1	86

Table 3: Activity and purity of antibody following different reactivation treatments.

With respect to bioactivity, in all cases fully active material was obtained (86-108 % of theoretical maximum) versus the non-selectively reduced secukinumab starting material, which had only 45% activity. With respect to CE-SDS purity, impaired purity (82.3%) was found in antibody obtained under completely anaerobic treatment lacking cystine, which was probably due to over-reduction under these conditions (note: addition of cystine to anaerobic reactions increased the purity of the samples as measured by CE-SDS, because reduction potential is smaller and by this over-reduction less likely). However, all other selective reduction treatments led to similar, and high, antibody quality and activity. We also noted that some CE-SDS purity values prior to the SP-Sepharose chromatography step were slightly lower than the values obtained following the chromatography step (e.g., 65% before vs 82% after) (data not shown), which suggests that additional re-oxidation may occur during the chromatography step. CEX activity was generally lower under the various aerobic conditions because there was more oxygen to mitigate the reductive power of the cysteine. Addition of the oxidative agents cystine and cystamine under aerobic conditions further decreased CEX activity.

Summary and Conclusions Drawn from Examples 1-3

We have determined that CysL97 of secukinumab is available for reduction in solution, without requiring partial unfolding of the full antibody structure, e.g., using guanidine HCl. Moreover, secukinumab overall is amenable to selective reduction, which, under controlled conditions, should allow activation of the antibody without substantial degradation.

Cysteine was found to be particularly ideal for selective reduction of secukinumab, as it

displayed only moderate over-reduction, coupled with fast equilibrium. In the first hour of selective reduction using cysteine, the antibody is partially reduced, e.g., up to 60% under anaerobic conditions and up to about 30% under fully-aerobic conditions. Subsequently, secukinumab slowly re-oxidizes to intact antibody. Re-oxidation of samples subjected to selective reduction with cysteine proceeded much more slowly at room temperature than reactions performed at 37°C (c.f. 21 hours for re-equilibrium at room temperature versus 8 hrs for equilibrium at 37°C). Room temperature samples also generally displayed smaller maximal decreases in intact antibody compared to selective reduction reactions performed at 37°C.

Under anaerobic conditions, we noted that the initial reduction of the antibody is larger and re-equilibration to intact secukinumab is not as complete as under aerobic conditions. However, addition of a small amount of cystine to anaerobic reactions resulted in improved purity and activity relative to anaerobic reactions performed in the absence of cystine. The aerobic reaction course can thus be simulated under anaerobic conditions when a small molar ratio of the oxidized form of the reducing reagent (e.g., cystine in the case of cysteine as the reducing agent) is present. Moreover, we noted that addition of cystine accelerated equilibrium of intact antibody – even when cystine was not present during the initial 30 minutes of incubation. Thus, selective reduction may be performed in the presence of air, as well as in the absence of dissolved oxygen by using an inert gas atmosphere (e.g., nitrogen or argon). If performed under fully anaerobic conditions, addition of a small molar ratio of the oxidized form of the reducing reagent is useful.

Example 4: Cysteine Selective Reduction Step Development Study: Proof of Concept (DoE1)

Example 4.1 - Study Design and Methods

The main purpose of the cysteine treatment step is to regain the full biological activity of the secukinumab antibody by the masking of the –SH group of the cysteine in position 97 of the light chain, which can occur during cell cultivation, harvesting and / or Protein A chromatography. In the following Example, the purity of the antibody is analyzed by a non-reducing CE-SDS method that monitors antibody integrity (called “purity by CE-SDS”) and the biologic activity of the antibody is analyzed by a cystamine-CEX chromatography method (called “activity by CEX”). In cystamine-CEX chromatography, cystamine is added to the

antibody sample and allowed to derivatize any free cysteine in secukinumab. The sample is then subjected to analytical separation by cation exchange chromatography, whereby cystamine-derivatized antibody species elute after non-derivatized antibody species because they carry an additional positive charge. The chromatogram is then compared to the chromatogram of the non-treated sample. The portion of species in the chromatogram of the treated sample having a shifted elution position is a direct measure of the abundance of free CysL97-SH in the original sample and can be used as surrogate marker for activity.

The main purpose of the proof of concept study (DoE1) was to check the applicability of design of experiments studies for improving of the cysteine treatment step to test a first set of parameter ranges to identify main influencing factors and support definition of operating ranges. Additionally, DoE1 was performed to evaluate if the experimental setup is applicable to detect the influence of input parameters on output parameters. The concept study was analyzed using Modde 9.0 (Umetrics) Software.

The input parameters are shown in **Table 4**. The addition of cystine (oxidized form) was tested as an additional input parameter, as the redox potential is influenced by the ratio reduced form/oxidized form, in this case the ratio cysteine/cystine. The ratio cysteine/cystine and cysteine concentration were investigated on 3 levels each and the antibody concentration ("content by ALC") on 2 levels. For product quality output parameters, activity by CEC and purity by CE-SDS were determined.

Name	Unit	Lower level	Medium level	Upper level
ratio cysteine/cystine	[-]	2.00	6.00	10.0
cysteine concentration	[mM]	0.50	5.25	10.0
content by ALC	[mg/mL]	3.25	n.a.	6.5

Table 4: Input parameters.

The design is a $3^2 \times 2$ Multilevel Factorial Design without center point runs. The design was chosen to gain data and process understanding for improving the cysteine treatment step. This type of design supports calculation of mathematical models with linear, interaction and quadratic terms for the input parameters ratio cysteine/cystine and cysteine concentration. The experimental design plan is given in **Table 5**.

RUN	Ratio cysteine to cystine [M/M]	Cysteine concentration [mM]	Protein Content by ALC [mg/mL]	Approx. molar ratio cysteine to protein [M/M]
React007_1	2	0.50	6.50	11.38
React007_2	6	10.00	6.50	227.61
React007_3	2	10.00	6.50	227.61
React007_4	10	10.00	6.50	227.61
React007_6	6	0.50	6.50	11.38
React007_7	10	10.00	3.25	455.21
React007_8	10	5.25	6.50	119.49
React007_9	6	10.00	3.25	455.21
React007_10	6	5.25	6.50	119.49
React007_11	2	10.00	3.25	455.21
React007_12	10	0.50	3.25	22.76
React007_13	6	5.25	3.25	238.99
React007_14	6	0.50	3.25	22.76
React007_15	10	5.25	3.25	238.99
React007_16	2	5.25	3.25	238.99
React007_17	2	0.50	3.25	22.76
React007_18	10	0.50	6.50	11.38

Table 5: Experimental design plan. The relative molecular mass of secukinumab, based on amino acid sequence, is 147,944 Daltons, which is used to calculate the molar ratio of cysteine to protein.

The selective reduction reactions were performed in 15 mL closed polypropylene tubes, which were placed in a water bath and warmed to 37 °C. The protein solution was diluted to the target concentration with WFI and pH was adjusted to pH 8.0 with 1 M Tris. After addition of the cysteine and cystine solution the tubes were tempered and incubated in a water bath for 4h. Then samples were withdrawn for determination of content by ALC. CE-SDS was performed with samples, which were withdrawn directly after incubation and the reaction was stopped by addition of a 10-fold excess (related to cysteine concentration) of iodoacetamide. The remaining solution was cooled to ambient temperature, the pH was adjusted to pH 5.0 - 5.5 and samples were withdrawn for CEX and SEC analytics.

The design was carried out according to the actual experimental design in **Table 6**.

Run	Molar ratio cysteine to cystine [M/M]	Cysteine concentration [mM]	Protein content by ALC [mg/mL]	Approx. molar ratio cysteine to protein [M/M]
React007_1	2	0.50	6.9	10.72
React007_2	6	10.00	6.3	234.83
React007_3	2	10.00	6.3	234.83
React007_4	10	10.00	6.3	234.83
React007_5	2	5.25	6.5	119.49
React007_6	6	0.50	6.8	10.88
React007_7	10	10.00	3.2	462.33
React007_8	10	5.25	6.5	119.49
React007_9	6	10.00	3.2	462.33
React007_10	6	5.25	6.6	117.68
React007_11	2	10.00	3.2	462.33
React007_12	10	0.50	3.5	21.13
React007_13	6	5.25	3.4	228.44
React007_14	6	0.50	3.5	21.13
React007_15	10	5.25	3.3	235.37
React007_16	2	5.25	3.4	228.44
React007_17	2	0.50	3.5	21.13
React007_18	10	0.50	6.6	11.21

Table 6: Actual experimental design. The relative molecular mass of secukinumab, based on amino acid sequence, is 147,944 Da, which is used to calculate the molar ratio of cysteine to protein.

Example 4.2 - Results of proof of concept study

The values for the output parameters are listed in **Table 7** together with the input parameters. The runs are listed with ascending values for activity by CEX. It can be seen that all runs with 0.5 mM concentration of cysteine have lower activities – probably because this cysteine concentration is too low to achieve complete unmasking of Cys97. In this set, a slight improvement in activity by CEX was noted when the ratio of cysteine:antibody was increased from about 11:1 to 21:1 (REACT00_17, _14 and _12). Of these three runs, those with a slightly higher ratio of cysteine:cystine (REACT007_14 and _12) (probably representing slight mitigation in the reductive power of cysteine) also had good purity by CE-SDS parameter. Then, there is a series of runs where the difference in activity output parameter is small (92.4 to 93.1%) and probably within the analytical accuracy, making interpretation among this group difficult. Finally, there is a set of 4 runs where the activity was the highest (93.7 to 95.1 %). Notably,

these runs did not contain the highest cysteine/protein ratio (462.33).

Run	Molar ratio cysteine to cystine [M/M]	Cysteine concentration [mM]	Protein content by ALC [mg/mL]	Approx. molar ratio cysteine to protein [M/M]	activity by CEX [%]	purity by CE-SDS [%]
React007_1	2	0.50	6.9	10.72	87.2	97.0
React007_6	6	0.50	6.8	10.88	88.8	95.0
React007_18	10	0.50	6.6	11.21	88.8	94.0
React007_17	2	0.50	3.5	21.13	89.3	78.0
React007_14	6	0.50	3.5	21.13	90.4	93.0
React007_12	10	0.50	3.5	21.13	90.9	95.0
React007_16	2	5.25	3.4	228.44	92.4	85.0
React007_3	2	10.00	6.3	234.83	92.8	83.0
React007_13	6	5.25	3.4	228.44	92.8	85.0
React007_9	6	10.00	3.2	462.33	92.9	68.0
React007_7	10	10.00	3.2	462.33	93.0	56.0
React007_8	10	5.25	6.5	119.49	93.0	88.0
React007_11	2	10.00	3.2	462.33	93.0	70.0
React007_4	10	10.00	6.3	234.83	93.1	77.0
React007_2	6	10.00	6.3	234.83	93.7	76.0
React007_15	10	5.25	3.3	235.37	94.0	78.0
React007_10	6	5.25	6.6	117.68	94.2	84.0
React007_5	2	5.25	6.5	119.49	95.2	87.0

Table 7: Input and output parameter values listed with ascending values for activity by CEX

Because of a good model fit it was possible to use contour plots to understand the main influencing parameters of the system. The quality of the models is represented by 4 tools, namely the R^2 , Q^2 , Model validity and the Standard Deviation (SD) of replicates. A model that is in good agreement with the data will have a R^2 and Q^2 close to 1.0 and Model validity above 0.25. Models with lower statistical significance have lower R^2 and Q^2 values. The model diagnostics for the quality output parameters indicate that the models for activity by CEX and purity by CE-SDS have high statistical significance ($R^2 = 0.88$ and $Q^2 = 0.73$ for activity by CEX and $R^2 = 0.84$ and $Q^2 = 0.63$ for purity by CE-SDS). The contour plot for activity by CEX (**Figure 4A**) teaches that activity by CEX values equal or higher than 93.0 % are achieved if the input parameter cysteine concentration is between 4.0 mM and 9.0 mM. Furthermore, the input factor content by ALC can vary from the low level (3.25 mg/mL) to the high level (6.5 mg/mL) when the input factor ratio cysteine / cystine is set to its center point of 6. To improve activity by CEX,

the model indicates that the input parameter content by ALC should be higher than the investigated upper level of 6.5 mg/mL.

In **Table 8**, the runs are listed according to increasing CE-SDS purity. Three groups can be seen: runs with lower purity ($\leq 70\%$), to which the runs with the high molar ratio of 462 Mol/Mol belong; runs with medium purity (70-83 %); and runs with higher purity ($>83\%$). Due to a good model fit, it is possible to also use a contour plot to understand how the input parameter influence the purity by CE-SDS values shown in **Table 8**. According to the contour plot for CE-SDS in **Figure 4B**, the area that comes closest to the theoretical optimum lies at the right lower side. This implies that the ratio cysteine/cystine should be set at a higher level and the cysteine concentration should be set to a lower level. Additionally, a higher content by ALC could be used to increase the purity by CE-SDS.

Run	Molar ratio cysteine to cystine [M/M]	Cysteine concentration [mM]	Protein content by ALC [mg/mL]	Approx. molar ratio cysteine to protein [M/M]	Activity by CEX [%]	Purity by CE- SDS [%]
React007_7	10	10.00	3.2	462.33	93.0	56
React007_9	6	10.00	3.2	462.33	92.9	68
React007_11	2	10.00	3.2	462.33	93.0	70
React007_2	6	10.00	6.3	234.83	93.7	76
React007_4	10	10.00	6.3	234.83	93.1	77
React007_15	10	5.25	3.3	235.37	94.0	78
React007_3	2	10.00	6.3	234.83	92.8	83
React007_10	6	5.25	6.6	117.68	94.2	84
React007_16	2	5.25	3.4	228.44	92.4	85
React007_13	6	5.25	3.4	228.44	92.8	85
React007_5	2	5.25	6.5	119.49	95.2	87
React007_8	10	5.25	6.5	119.49	93.0	88
React007_14	6	0.50	3.5	21.13	90.4	93
React007_18	10	0.50	6.6	11.21	88.8	94
React007_6	6	0.50	6.8	10.88	88.8	95
React007_12	10	0.50	3.5	21.13	90.9	95
React007_1	2	0.50	6.9	10.72	87.2	97

Table 8: Input and output parameters listed according to increasing CE-SDS purity.

Example 4.3 - Summary of Proof of Concept Study

The proof of concept study confirmed the applicability of the experimental setup to

evaluate and refine the input parameters of the cysteine treatment step. We noted that a high molar ratio of cysteine to protein can lead to low values of purity by CE-SDS. The models also suggested that increasing protein and cysteine concentration might improve purity by CE-SDS and activity by CEX. Thus, cysteine concentration and the influence of an increased parameter range for content by ALC on the product quality attributes activity by CEX and purity by CE-SDS was evaluated in more detail in the following study (Response Surface Design 1).

Example 5: Cysteine Selective Reduction Step Development Study: Response Surface Design 1

Example 5.1 - Study Design and Methods

Based on the data from **Example 4**, the input parameters of the selective reduction step were adjusted. The development study was analyzed using Modde 9.0 (Umetrics) Software. The input parameters are listed in **Table 9**. The factors are investigated on three levels each. The design was an adjusted version of the proof of concept study with adjusted levels of the input parameters content by ALC and cysteine concentration. The input parameter “ratio cysteine/cystine” was replaced by the parameter “concentration of cystine” to simplify the experimental setup and further the data evaluation.

Name	Unit	Lower level	Medium level	Upper level
Cystine concentration	[mM]	0.0	0.5	1.0
cysteine concentration	[mM]	4.0	9.0	14.0
content by ALC	[mg/mL]	3.8	11.6	19.5

Table 9: Input parameters.

The design is a 2^3 Center Composite Face (CCF) design with 4 center runs at center point conditions. This type of design supports calculation of mathematical models with linear, interaction and quadratic terms. The experimental design plan is given in **Table 10**.

Run	Protein content by ALC (mg/mL)	Cysteine concentration (mM)	Cystine concentration (mM)	Molar ratio cysteine to cystine (M/M)	Approx. molar ratio cysteine to protein (M/M)
React016_1	19.5	14	0.0	NA	106.22
React016_2 ¹	11.6	9	0.5	18	114.78
React016_3	3.8	9	0.5	18	350.39
React016_4	11.6	9	1.0	9	114.78

React016_5	19.5	9	0.5	18	68.28
React016_6	19.5	4	0.0	NA	30.35
React016_7	3.8	4	1.0	4	155.73
React016_8 ¹	11.6	9	0.5	18	114.78
React016_9	11.6	14	0.5	28	178.55
React016_10	3.8	4	0.0	NA	155.73
React016_11	11.6	4	0.5	8	51.02
React016_12	3.8	14	0.0	NA	545.06
React016_13	19.5	4	1.0	4	30.35
React016_14 ¹	11.6	9	0.5	18	114.78
React016_15 ¹	11.6	9	0.5	18	114.78
React016_16	19.5	14	1.0	14	106.22
React016_17	11.6	9	0.0	NA	114.78
React016_18	3.8	14	1.0	14	545.06

Table 10: Experimental design plan. 1) Center points. The relative molecular mass of secukinumab, based on amino acid sequence, is 147,944 Da, which is used to calculate the molar ratio of cysteine to protein.

The selective reduction reactions were performed in 15 mL closed polypropylene tubes, which were placed in a water bath and warmed to 37 °C. The design was carried out according to the actual experimental design in **Table 11**.

Run	Protein content by ALC (mg/mL)	Cysteine concentration (mM)	Cystine concentration (mM)	Molar ratio cysteine to cystine (M/M)	Approx. molar ratio cysteine to protein (M/M)
React016_1	19.5	14	0.0	NA	106.22
React016_2 ¹	11.9	9	0.5	18	111.89
React016_3	4.0	9	0.5	18	332.87
React016_4	11.5	9	1.0	9	115.78
React016_5	19.4	9	0.5	18	68.63
React016_6	19.4	4	0.0	NA	30.50
React016_7	3.8	4	1.0	4	155.73
React016_8 ¹	11.6	9	0.5	18	114.78
React016_9	11.5	14	0.5	28	180.11
React016_10	3.7	4	0.0	NA	159.94
React016_11	11.6	4	0.5	8	51.02
React016_12	3.8	14	0.0	NA	545.06
React016_13	19.3	4	1.0	4	30.66
React016_14 ¹	11.7	9	0.5	18	113.80
React016_15 ^{1, 2}	11.1	9	0.5	18	119.95

React016_16	17.8	14	1.0	14	116.36
React016_17	10.9	9	0.0	NA	122.16
React016_18	3.8	14	1.0	14	545.06

Table 11: Actual experimental design. 1) Center point; 2) This experiment is not regarded as a center point of the used DoE software, due to a deviation of more than 10% protein concentration from the originally planned center point. The relative molecular mass of secukinumab, based on amino acid sequence, is 147,944 Da, which is used to calculate the molar ratio of cysteine to protein.

Example 5.2 -Results of Response Surface Design 1

The input parameters and values for the output parameters are listed in **Table 12**. The runs are listed according increasing activity by CEX. It can be seen that runs with high CEX activity used higher cysteine concentration. The single run with lower activity (React016_13) had high protein concentration, low cysteine concentration and high cystine concentration.

The model diagnostics for the product quality output parameter were $R^2 = 0.95$, $Q^2 = 0.76$, $SD=0.20$, and Model validity = 0.78 for the activity by CEX and $R^2=0.79$, $Q^2=0.53$, $SD=2.91$, and Model validity = 0.83 for purity by CE-SDS, indicating significant models for both output parameters. Contour plots were calculated (**Figure 5**), to gain more insight into the impact of the input parameter settings and the impact on activity by CEX.

Run	Protein content by ALC (mg/mL)	Cysteine concentration (mM)	Cystine concentration (mM)	Molar ratio cysteine to cystine (M/M)	Approx. molar ratio cysteine to protein (M/M)	Activity by CEX [%]	Purity by CE-SDS [%]
React016_13	19.3	4	1	4	30.66	91.6	93
React016_7	3.8	4	1.0	4	155.73	93.3	92
React016_6	19.4	4	0.0	NA	30.50	93.5	94
React016_11	11.6	4	0.5	8	51.02	93.5	93
React016_5	19.4	9	0.5	18	68.63	94.1	85
React016_10	3.7	4	0.0	NA	159.94	94.2	79
React016_2 ¹	11.9	9	0.5	18	111.89	94.3	86
React016_3	4.0	9	0.5	18	332.87	94.3	85
React016_4	11.5	9	1.0	9	115.78	94.5	87
React016_8 ¹	11.6	9	0.5	18	114.78	94.5	91
React016_15 ^{1,2}	11.1	9	0.5	18	119.95	94.5	86
React016_14 ¹	11.7	9	0.5	18	113.80	94.7	86
React016_17	10.9	9	0.0	NA	122.16	94.7	85
React016_1	19.5	14	0.0	NA	106.22	94.9	76

React016_9	11.5	14	0.5	28	180.11	95.1	85
React016_12	3.8	14	0.0	NA	545.06	95.1	74
React016_16	17.8	14	1.0	14	116.36	95.1	78
React016_18	3.8	14	1.0	14	545.06	95.1	77

Table 12: Input and output parameters listed according increasing activity by CEX. 1) Center points; 2) This experiment is not regarded as a center point of the used DoE software, due to a deviation of more than 10% protein concentration from the originally planned center point.

As shown in the contour plot in **Figure 5**, to achieve a high activity by CEX, the most influential input parameter is the cysteine concentration, which should be set to its high level. But, even at the low level of cysteine concentration high CEX activity (> 93 %) is met if the content by ALC and the cystine concentration are not at the upper limit. The input parameters content by ALC and cystine concentration also have influence on the output parameter activity by CEX, but to a smaller extent.

In **Table 13** the runs are listed according to ascending CE-SDS purity. The runs can be divided into three groups. There was a series of six runs (mainly the 14 mM runs) with low purity (<80%), a series of runs with medium purity (85-91 %), and a series of runs with higher purity (>91%), the later all being 4mM cysteine runs.

Run	Protein content by ALC (mg/mL)	Cysteine concentration (mM)	Cystine concentration (mM)	Molar ratio cysteine to cystine (M/M)	Approx. molar ratio cysteine to protein (M/M)	Activity by CEX [%]	Purity by CE-SDS [%]
React016_12	3.8	14	0.0	NA	545.06	95.1	74
React016_1	19.5	14	0.0	NA	106.22	94.9	76
React016_18	3.8	14	1.0	14	545.06	95.1	77
React016_16	17.8	14	1.0	14	116.36	95.1	78
React016_10	3.7	4	0.0	NA	159.94	94.2	79
React016_5	19.4	9	0.5	18	68.63	94.1	85
React016_3	4.0	9	0.5	18	332.87	94.3	85
React016_17	10.9	9	0.0	NA	122.16	94.7	85
React016_9	11.5	14	0.5	28	180.11	95.1	85
React016_2 ¹	11.9	9	0.5	18	111.89	94.3	86
React016_15 ^{1, 2}	11.1	9	0.5	18	119.95	94.5	86
React016_14 ¹	11.7	9	0.5	18	113.80	94.7	86
React016_4	11.5	9	1.0	9	115.78	94.5	87
React016_8 ¹	11.6	9	0.5	18	114.78	94.5	91
React016_7	3.8	4	1.0	4	155.73	93.3	92

React016_13	19.3	4	1.0	4	30.66	91.6	93
React016_11	11.6	4	0.5	8	51.02	93.5	93
React016_6	19.4	4	0.0	NA	30.50	93.5	94

Table 13: Input and output parameters listed according to ascending CE-SDS purity. 1) Center points; 2) This experiment is not regarded as a center point of the used DoE software, due to a deviation of more than 10% protein concentration from the originally planned center point.

Based on the results the cysteine concentration should be low and the content by ALC and cystine concentration set to high to achieve higher purity by CE-SDS values.

The experiments React016_6, React016_7, React016_8 and React016_11 have high values for both two quality output parameters. To estimate the risk of failure of the cysteine treatment step based on the model established by this DoE, a design space estimator using Monte Carlo simulations was performed. The design space estimator provides a range where the factors can vary within, while the output parameters do not exceed the target range for product quality and process performance output parameter. It is suggested that content by ALC should be set to 12 mg/mL as ideal. The cysteine concentration has an ideal at 4.0 mM in this particular experiment. For the cystine concentration, the ideal is 0.32 mM.

According to the results of this study, the cysteine treatment step is mainly influenced by the cysteine concentration. High cysteine concentration promotes high activity by CEX, but leads simultaneously to low purity by CE-SDS. According to the design space estimator, the cysteine concentration should be kept at 4 mM and the content by ALC should be set to 12 mg/mL (ratio cysteine to protein of about 49).

For verification of the results of the statistical designs, target runs were performed in stirred 2L-bioreactors having an air overlay and open headspace. No stirring was applied during incubation. The process parameters are listed in **Table 14** and the analytical results are shown in **Table 15**.

Process parameter	Unit	Target run 1 (REACT019)	Target run 2 (REACT020)	Target run 3 (REACT021)
Protein concentration	[mg/mL]	12.90	13.00	12.60
Cysteine concentration	[mM]	4.00	8.00	4.00
Cystine concentration	[mM]	0.15	0.15	0.00
Temperature	[°C]	37.00	37.00	37.00
pH	[-]	8.00	8.00	8.00

Incubation time	[h]	4.00	4.00	4.00
Molar ratio cysteine/cystine	M/M	26.67	53.33	n.a.
Approx. molar ratio cysteine to protein	M/M	45.87	91.04	46.97

Table 14: Process parameters of target runs. The relative molecular mass of secukinumab, based on amino acid sequence, is 147,944 Da, which is used to calculate the molar ratio of cysteine to protein.

Process parameter	Unit	Target run 1 (REACT019)	Target run 2 (REACT020)	Target run 3 (REACT021)
Activity by CEX	[%]	94.8	94.3	95.4
Purity by CE-SDS	[%]	94.0	89.0	95.0

Table 15: Analytical Results of target runs.

All three runs lead to similar results regarding activity by CEX. This corresponds to the results of the previous DoE study indicating that 4.0 mM cysteine is sufficient to regain the activity of the antibody. In the 3rd target run (REACT021) the cystine was omitted, leading to comparable results for purity by CE-SDS with run 1 (REACT019) and demonstrating that, under the conditions tested, cystine has only minor influence on performance of the cysteine treatment step. An increase of the cysteine concentration to 8.0 mM (ratio of cysteine to protein of about 91) led, not to higher values of activity by CEX, but to lower purity by CE-SDS in the pool, falling below the development target of 90 %. Although cystine showed significance in the statistical design, a positive effect was not detectable in the target runs. Additionally, the influence in the statistical design on activity by CEX is minor (see **Figure 5**) and preparation of a cystine containing buffer is difficult due to the low solubility of cystine in non-basic buffers. Hence, cystine can be omitted in the cysteine treatment buffer.

Example 6: Cysteine Selective Reduction Step Development Study: Response Surface Design 2

Example 6.1 -Study Design and Methods

The main purpose of the response surface design 2 (DoE3) was the investigation of the process parameters temperature, time and pH on the cysteine treatment step in addition to the cysteine concentration. The data are also used to support definition of operating ranges and identify main influencing factors. The development study was analyzed using Modde 9.0 (Umetrics) Software. The input parameters are found in **Table 16**.

Name	Unit	Lower level	Medium level	Upper level
------	------	-------------	--------------	-------------

temperature	[°C]	32.0	37.0	42.0
time	[h]	1.0	4.0	7.0
pH	[-]	7.5	8.0	8.5
cysteine concentration	[mM]	2.0	5.0	8.0

Table 16: Input parameters.

The design is a Central Composite Face (CCF) design with 3 center point runs. The experimental design plan is given in **Table 17**. The design was chosen to evaluate the influence of additional process parameters, as the cysteine treatment solution was improved in the former experiments. The cysteine concentration was analyzed again as it was the most influencing input parameter on the cysteine treatment step and interactions with the input parameters were expected. This type of design supports calculation of mathematical models with linear, interaction and quadratic terms.

Run	Temperature [°C]	Time (h)	pH [-]	Cysteine concentration [mM]	Protein content by ALC [mg/mL]	Approx. molar ratio of cysteine to protein (M/M)
React027_1	42	7	8.5	8	13	91.04
React027_2	37	4	8.0	8	13	91.04
React027_3	32	1	7.5	8	13	91.04
React027_4	32	1	8.5	2	13	22.76
React027_5 ¹	37	4	8.0	5	13	56.90
React027_6	42	1	8.5	8	13	91.04
React027_7	32	1	8.5	8	13	91.04
React027_8	32	1	7.5	2	13	22.76
React027_9 ¹	37	4	8.0	5	13	56.90
React027_10	37	7	8.0	5	13	56.90
React027_11	42	7	7.5	2	13	22.76
React027_12	42	4	8.0	5	13	56.90
React027_13	32	7	7.5	8	13	91.04
React027_14	37	4	8.0	2	13	22.76
React027_15	37	4	7.5	5	13	56.90
React027_16	32	4	8.0	5	13	56.90
React027_17	42	1	8.5	2	13	22.76
React027_18	32	7	7.5	2	13	22.76
React027_19	37	1	8.0	5	13	56.90
React027_20	42	7	7.5	8	13	91.04
React027_21	42	1	7.5	8	13	91.04

React027_22 ¹	37	4	8.0	5	13	56.90
React027_23	42	7	8.5	2	13	22.76
React027_24	32	7	8.5	2	13	22.76
React027_25	37	4	8.5	5	13	56.90
React027_26	32	7	8.5	8	13	91.04
React027_27	42	1	7.5	2	13	22.76

Table 17: Experimental design plan. 1) Center points. The relative molecular mass of secukinumab, based on amino acid sequence, is 147,944 Da, which is used to calculate the molar ratio of cysteine to protein.

The selective reduction reactions were performed in 15 mL closed polypropylene tubes, which were placed in a water bath and warmed to 37 °C.

Example 6.2 - Results of Response Surface Design 2

The study was carried out according to the actual experimental design in **Table 18**.

Run	Temperature [°C]	Time (h)	pH [-]	Cysteine concentration [mM]	Protein content by ALC [mg/mL]	Approx. molar ratio of cysteine to protein [M/M]
React027_1	42	7	8.5	8	13	91.04
React027_2	37	4	8.0	8	13	91.04
React027_3	32	1	7.5	8	13	91.04
React027_4	32	1	8.4	2	13	22.76
React027_5 ^{1,2}	37	4	7.9	5	13	56.90
React027_6	42	1	8.5	8	13	91.04
React027_7	32	1	8.5	8	13	91.04
React027_8	32	1	7.3	2	13	22.76
React027_9 ¹	37	4	8.0	5	13	56.90
React027_10	37	7	8.0	5	13	56.90
React027_11	42	7	7.4	2	13	22.76
React027_12	42	4	8.0	5	13	56.90
React027_13	32	7	7.6	8	13	91.04
React027_14	37	4	7.9	2	13	22.76
React027_15	37	4	7.5	5	13	56.90
React027_16	32	4	8.0	5	13	56.90
React027_17	42	1	8.5	2	13	22.76
React027_18	32	7	7.4	2	13	22.76
React027_19	37	1	8.0	5	13	56.90
React027_20	42	7	7.6	8	13	91.04
React027_21	42	1	7.6	8	13	91.04

React027_22 ¹	37	4	8.0	5	13	56.90
React027_23	42	7	8.5	2	13	22.76
React027_24	32	7	8.5	2	13	22.76
React027_25	37	4	8.5	5	13	56.90
React027_26	32	7	8.5	8	13	91.04
React027_27	42	1	7.5	2	13	22.76

Table 18: Actual experimental design plan. 1) Center points. 2) Is not considered a center point of the used software. The relative molecular mass of secukinumab, based on amino acid sequence, is 147,944 Da, which is used to calculate the molar ratio of cysteine to protein.

The values for the product quality output parameters are listed in **Table 19** in ascending order with respect to the CEX activity results. The model diagnostics for the product quality output parameter are $R^2 = 0.93$, $Q^2 = 0.80$, Model Validity = 0.79 and SD = 0.42 for activity by CEX; $R^2 = 0.96$, $Q^2 = 0.89$, Model Validity = 0.67 and SD = 0.00 for purity by CE-SDS.

Run	Temperature [°C]	Time (h)	pH [-]	Cysteine concentration [mM]	Protein Content by ALC (mg/mL)	Approx. molar ratio of cysteine to protein (M/M)	activity by CEX [%]	purity by CE-SDS [%]
React027_8	32	1	7.3	2	13	22.76	89.1	91
React027_27	42	1	7.5	2	13	22.76	90.1	91
React027_3	32	1	7.5	8	13	91.04	90.3	61
React027_4	32	1	8.4	2	13	22.76	91.1	92
React027_7	32	1	8.5	8	13	91.04	91.7	65
React027_17	42	1	8.5	2	13	22.76	91.8	89
React027_18	32	7	7.4	2	13	22.76	92.5	93
React027_14	37	4	7.9	2	13	22.76	93.3	94
React027_21	42	1	7.6	8	13	91.04	93.3	77
React027_19	37	1	8.0	5	13	56.90	93.4	79
React027_23	42	7	8.5	2	13	22.76	93.6	93
React027_24	32	7	8.5	2	13	22.76	94.1	93
React027_11	42	7	7.4	2	13	22.76	94.2	95
React027_6	42	1	8.5	8	13	91.04	94.3	74
React027_16	32	4	8.0	5	13	56.90	94.4	86
React027_13	32	7	7.6	8	13	91.04	94.5	82
React027_9 ¹	37	4	8.0	5	13	56.90	95.0	89
React027_5 ^{1,2}	37	4	7.9	5	13	56.90	95.2	89
React027_10	37	7	8.0	5	13	56.90	95.3	90
React027_22 ¹	37	4	8.0	5	13	56.90	95.6	88
React027_25	37	4	8.5	5	13	56.90	95.7	87
React027_20	42	7	7.6	8	13	91.04	95.8	85

React027_12	42	4	8.0	5	13	56.90	95.9	90
React027_1	42	7	8.5	8	13	91.04	96.0	85
React027_15	37	4	7.5	5	13	56.90	96.1	88
React027_2	37	4	8.0	8	13	91.04	97.1	82
React027_26	32	7	8.5	8	13	91.04	97.1	82

Table 19: Input and output parameter values listed according to ascending activity by CEX. 1) Center point. 2) Is not considered a center point of the used software.

Table 19 shows that the runs can be divided mainly in three groups: lower activity (< 93 %), medium activity (93-95%), and high activity (>95%). Due to a good model fit, it is possible to use the contour plot (**Figure 6**) to identify the main influencing parameters of the system.

Figure 6 shows that medium to high incubation (e.g., 4 to 7 hours) time and high cysteine concentration is beneficial to increase activity by CEX. Incubation temperature and process pH have less influence on the output parameter activity by CEX. Nevertheless, the areas for purity of 95 % and above are the greatest in the plots for pH 8.0 and 8.5 at 37°C, suggesting that this is an optimal operating range.

In **Table 20**, the runs are listed by increasing purity by CE-SDS. The runs can be divided in three groups: runs with low purity (< 82%), runs with medium purity and runs with high purity (>90 %). All runs with high purity by CD-SDS employed 2 mM cysteine.

Run	Temperature [°C]	Time (h)	pH [-]	Cysteine concentration [mM]	Protein Content by ALC (mg/mL)	Approx. molar ratio of cysteine to protein (M/M)	activity by CEX [%]	purity by CE- SDS [%]
React027_3	32	1	7.5	8	13	91.04	90.3	61
React027_7	32	1	8.5	8	13	91.04	91.7	65
React027_6	42	1	8.5	8	13	91.04	94.3	74
React027_21	42	1	7.6	8	13	91.04	93.3	77
React027_19	37	1	8.0	5	13	56.90	93.4	79
React027_13	32	7	7.6	8	13	91.04	94.5	82
React027_2	37	4	8.0	8	13	91.04	97.1	82
React027_26	32	7	8.5	8	13	91.04	97.1	82
React027_20	42	7	7.6	8	13	91.04	95.8	85
React027_1	42	7	8.5	8	13	91.04	96.0	85
React027_16	32	4	8.0	5	13	56.90	94.4	86
React027_25	37	4	8.5	5	13	56.90	95.7	87
React027_22 ¹	37	4	8.0	5	13	56.90	95.6	88
React027_15	37	4	7.5	5	13	56.90	96.1	88
React027_17	42	1	8.5	2	13	22.76	91.8	89

React027_9 ¹	37	4	8.0	5	13	56.90	95.0	89
React027_5 ^{1,2}	37	4	7.9	5	13	56.90	95.2	89
React027_10	37	7	8.0	5	13	56.90	95.3	90
React027_12	42	4	8.0	5	13	56.90	95.9	90
React027_8	32	1	7.3	2	13	22.76	89.1	91
React027_27	42	1	7.5	2	13	22.76	90.1	91
React027_4	32	1	8.4	2	13	22.76	91.1	92
React027_18	32	7	7.4	2	13	22.76	92.5	93
React027_23	42	7	8.5	2	13	22.76	93.6	93
React027_24	32	7	8.5	2	13	22.76	94.1	93
React027_14	37	4	7.9	2	13	22.76	93.3	94
React027_11	42	7	7.4	2	13	22.76	94.2	95

Table 20: Input and output parameters listed with increasing purity by CE-SDS. 1) Center point. 2) Is not considered a center point of the used software.

Because of the good model fit, a contour plot (**Figure 7**) can be used to understand the process and the main influencing input parameters. **Figure 7** shows that medium to high incubation time and low cysteine concentration increase purity by CE-SDS. The input parameters incubation temperature and process pH have a minor influence on purity by CE-SDS.

Summary and Conclusions Drawn from Example 6

The development targets are fulfilled when the cysteine concentration is held at a low level, while the temperature can vary from center to high, the pH can vary from center to high, and the time can vary from center to high. To help estimate the input parameter ranges that lead to high product quality, a design space estimator using Monte Carlo simulations was run. For this simulation, a target of >93% for CEX activity and >90% for CE-SDS purity was set. The Monte Carlo simulations predicted that these quality targets are fulfilled when temperature is ideally at 39.1°C, but temperature can vary between 38.6 °C to 39.7 °C. The incubation time has an ideal at 5.0 h, but can vary between 4.0 h and 6.0 h. For the pH, the ideal is 8.2 and it can vary from 8.15 to 8.25. For the last input parameter, cysteine concentration, the ideal predicted is 2.4 mM and can vary between 2.2 mM to 2.5 mM. Within these ranges, the design space estimator has found that for the product quality output parameter activity by CEX the estimated DPMO (Defect Per Million Operations) value will be 110 and the DPMO value for Purity by CE-SDS will be 20. This means that 110 times out of one million the target limit for activity by CEX will not be met and 20 times out of a million the target limit for the purity by CE-SDS.

According to the results of the contour plots, and taking into account the results of the

previous development studies (DoE1 and DoE2), the target for incubation temperature was set to 37 °C. The incubation time target was set to 4 hours (240 minutes), as with the present experimental setup no heating and cooling time was included, which will be considered for manufacturing scale applications. Combined with heating and cooling time, an overall process time of approximately 6 hours will be applied. The influence of pH on purity by CE-SDS is not detectable and on activity by CEX the influence is minor. Therefore, the target was set to 8.0, as it is well within the tested range, the development target for activity by CEX can be safely met, and the risk of formation of acidic variants by CEX (enhanced by basic conditions) will be decreased. The results for the input parameter cysteine concentration indicates a low optimum level of 2.4 mM; however, most experiments employing 2 mM cysteine resulted in lower activity by CEX than was desired. This, combined with the results of the target runs, suggests that the target for cysteine concentration should be higher, i.e., 4 mM, during confirmation runs.

Example 7: Process Comparisons and Analysis of Continuous Stirring

Example 7.1 - Comparison of Process B2 and Process C

The considerations described in **Example 6** suggested that the selective reduction could preferably be done at higher protein concentration and lower cysteine concentration (4-6 mM) than so far carried out at manufacturing scale (Process B2 herein). **Table 21** compares the input parameters for earlier Process B2 and Proposed process C. Comparison runs were performed in duplicate at 50 mL scale in polypropylene tubes, which were placed in a water bath and warmed to 37°C. The results are shown in **Table 22**.

	Unit	Process B2	Proposed process C
Cysteine concentration	[mM]	8.00	4.00
Protein concentration ¹⁾	[mg/mL]	4.30	13.50
Approx. molar ratio cysteine to protein	[M/M]	275.24	43.84
pH	[-]	8.00	8.00
Incubation time	[h]	4.00	4.00
temperature	[°C]	37.00	37.00

Table 21: Input parameters of process B2 and proposed process C. 1) Before pH adjustment and cysteine addition.

The relative molecular mass of secukinumab, based on amino acid sequence, is 147,944 Da, which is used to calculate the molar ratio of cysteine to protein.

Run	Step performance	Unit	Activity by CEX	Purity by CE-SDS
REACT030	Proposed process C	[%]	95.5	95

REACT031	Proposed process C	[%]	95.0	95
REACT032	Process B2	[%]	96.7	79
REACT033	Process B2	[%]	97.0	79

Table 22: Output parameters of process B2 and proposed process C.

Table 22 shows that the modified cysteine treatment step of Process C leads to a significant increase in CE-SDS purity with only a very slight decrease in activity by CEX. The higher values of purity by CE-SDS are achieved by increased protein concentration and decreased cysteine concentration during the selective reductive step (i.e., a smaller molar ratio of cysteine to protein). This leads to a lower reductive power that is nevertheless still sufficient to ensure adequate activity of secukinumab.

Example 7.2 -Testing of Continuous Stirring During Incubation

During clinical manufacturing according to process B2, an intermittent mixing was applied (2 minutes every hour) to avoid inhomogeneities during the incubation phase at 37 °C. The feasibility of using continuous stirring to assure homogeneity throughout incubation was tested in a 2L bioreactor (run REACT029). The experimental setup and process conditions are shown in **Table 23**.

	Unit	REACT029
Cysteine concentration	[mM]	4.00
Protein concentration ¹⁾	[mg/mL]	13.50
Approx. molar ratio cysteine to protein	[M/M]	43.84
Stirrer speed during incubation	[rpm]	200.00
Filling volume	[L]	1.20
Incubation time	[h]	4.00

Table 23: Input parameters for testing continuous stirring during reaction. 1) Before pH adjustment and cysteine addition. The relative molecular mass of secukinumab, based on amino acid sequence, is 147,944 Da, which is used to calculate the molar ratio of cysteine to protein.

For REACT029, purity by CE-SDS was 96% and activity by CEX was 92.7 %, which is a lower CEX activity than shown by runs REACT030 and REACT 031 (proposed Process C) performed at small scale without stirring. Thus, increased oxygen transfer into the cysteine treatment solution caused by continuous mixing could abrogate the reductive power of the cysteine. Nevertheless, the activity by CEX in the cysteine treatment pool was close to the quality target. Since oxygen transfer and the ratio surface/volume decreases with scale-up to

manufacturing scale, this experiment was considered the worst case regarding high oxygen transfer. Continuous mixing was therefore used for the pilot scale runs ("Process C"), in order to ensure the homogeneity of the process solution.

Summary and Conclusions Drawn from Examples 5-7

We determined that the protein concentration during cysteine treatment can be increased threefold, which decreases process volumes significantly. We also demonstrated that lower cysteine concentrations provide similar product activity and increased product purity by CE-SDS after cysteine treatment. Thus, lower ratios of cysteine:protein are able to ensure quality product.

The data also shows that temperature and pH have minor influence on the cysteine treatment step and that addition of the oxidized form of cysteine – cystine – is not necessary to obtain quality product. Although the incubation time showed an ideal at 5 hours, we set the target to 4 hours, as with the present experimental setup no heating and cooling time was included, which must be accounted for during manufacturing-scale applications. Continuous stirring under open conditions without active aeration appeared to be feasible for use in the selective reduction method; however, as we will show in later Examples, the oxygen transfer into the solution should be limited (e.g., using low/slow stirring).

Example 8: Process Confirmation at Pilot Scale

The proposed purification conditions after process improvement were tested by process confirmation at 7 L pilot-scale in order to perform a first scale-up and obtain a first impression on the reproducibility and robustness of the process. The selective reduction step according to Process C (**Table 21**) was used in the pilot scale experiments (REACT035). Continuous stirring was applied. The dissolved oxygen concentration was measured by a pO_2 probe.

The activity by CEX of REACT035 following the selective reduction step according to confirmation run 1 was only 90.4%. Analysis of the dissolved oxygen curve (see **Figure 8**) revealed that there is a steady decrease of pO_2 in the first three hours of the treatment, indicating that oxygen consumption takes place in the solution faster than oxygen transfer into the solution triggered by stirring. We assume that oxygen consumption is caused by oxidation of the cysteine reagent (Cys-SH) to cystine or either cysteic acid (Cys-SO₃). The increase of the oxygen level in the last hour of the reaction would then indicate that the cysteine reagent had

been consumed and oxygen take up by the solution caused by stirring becomes visible. This implies that the reductive power of cysteine is not effective in the last period of the treatment, which could explain the impaired activity obtained from this run.

Example 8.1 - Investigation of the Cysteine Step During Confirmation Run 1

To investigate the root cause for the low activity (90.4%) determined by CEX in the first pilot scale run REACT035, the possible influencing factors were assessed. As mentioned above, a low cysteine concentration, introduction of oxygen by extended stirring during the reaction, and the scale change (increased reaction volume) could result in insufficient reductive power. Hence, the stirrer speed during incubation and cysteine concentration were evaluated.

The experimental plan and output parameter activity by CEX is shown in **Table 24**. In this experiment, the dissolved oxygen concentration was measured by a pO₂ probe.

Run	Cysteine concentration [mM]	Approx. molar ratio cysteine to protein [M/M]	Stirring during incubation [rpm]	Activity by CEX [%]
REACT035 ¹⁾	4	43.84	100	90.4
REACT038 ²⁾	4	43.84	0 ³⁾	92.5
REACT039 ²⁾	4	43.84	50	93.7
REACT040 ²⁾	6	65.75	50	91.9

Table 24: Investigation of stirring and cysteine concentration on the cysteine treatment step. 1) confirmation run 1, performed in a 20L bioreactor with 7 L filling volume. 2) experiment performed in a 2 L bioreactor with 1.7 L filling volume. 3) mixing every hour for 2 minutes with 50 rpm. The relative molecular mass of secukinumab, based on amino acid sequence, is 147,944 Da, which is used to calculate the molar ratio of cysteine to protein.

All three smaller scale reactions (REACT038, REACT039, and REACT040) showed improved activity by CEX in comparison to confirmation run 1 (REACT035) (**Table 24**). However, the corresponding activity by CEX was not increased in the experiment with elevated cysteine concentration (REACT040) or in the experiment without continuous mixing during incubation time (REACT038) when compared to REACT039. Hence, neither the cysteine concentration nor the stirring during incubation could be identified at this point as the main influencing factor. Nevertheless, the dissolved oxygen (dO₂) profile of the pilot scale confirmation run 1 (REACT035) (**Figure 8**) showed an increase in dO₂ after approximately 3 hours incubation, indicating that the reductive power of cysteine has been exhausted. The dO₂ chart of the two smaller scale experiments REACT039 and REACT040, in which continuous

stirring was performed, show no increase in dO_2 towards the end of the treatment (data not shown), indicating that the reductive power of the cysteine had not been exhausted in these smaller scale experiments.

Example 8.2 - Investigation of the Cysteine Step During Confirmation Run 2

The filling volume was 14 L instead of 7 L as in confirmation run 1 and two stirrers were used for stirring (radial stirrer at the bottom and in addition axial on the top). Cysteine concentration (4mM) and stirrer speed (100 rpm) were the same as in confirmation run 1, REACT 035. Following selective reduction, the activity of the antibody solution was only 85.5% measured by CEX. According to the dO_2 chart shown in **Figure 9**, the reductive power of the cysteine was exhausted even earlier than during confirmation run 1 (**Figure 8**), suggesting that the oxygen transfer into the solution by continuous stirring was more pronounced in this run.

The dO_2 curve trends in (**Figure 8 – 9**) suggest that the oxygen transfer over the headspace by continuous stirring is the main root cause for lower activity after the selective reduction step during confirmation run 1 and 2. With 7 L filling volume at confirmation run 1, only one stirrer (radial stirrer at the bottom) was submerged in the process solution of the cysteine treatment step. With 14 L filling volume at confirmation run 2, two stirrers (radial stirrer at the bottom and axial on the top) were submerged in the process solution, leading to even further increased oxygen transfer into the solution. Hence, the even poorer result of confirmation run 2 is most probably due to the elevated oxygen transfer into the process solution, visible by the dO_2 curve (**Figure 9**). Due to the higher oxygen transfer into the solution for confirmation run 2, the reductive power of the cysteine is exhausted earlier, resulting in less activity by CEX. Therefore, the oxygen transfer into the process intermediate should be restricted, e.g., by adjustment of mixing (speed, duration, frequency) during incubation.

Example 8.3 - Investigation of the Cysteine Step During Confirmation Run 3

The filling volume was 16 L instead of 14 L as in confirmation run 2 and the stirring profile was adapted. Instead of continuous stirring during the cysteine treatment, only mixing for 2 minutes every hour was applied, corresponding to earlier process B2. The activity of the antibody solution measured by CEX following the selective reduction step was 92.5%.

Due to omission of continuous mixing in the cysteine treatment step, oxygen transfer was

restricted, ensuring sufficient reductive power and leading to high activity by CEX. The dO₂ chart shown in **Figure 10** shows a steep decrease of the oxygen level - almost to zero % in the beginning of the treatment. This experiment demonstrates that regulating the amount of oxygen introduced during the cysteine treatment ensures proper reaction conditions and consequently a high activity by CEX.

Example 8.4 - Investigation of the Cysteine Step During Confirmation Run 4

Based on the results of confirmation run 3, the 4th pilot-scale confirmation run (REACT040) was performed with the identical setup and process conditions of confirmation run 3. However, the cysteine treatment in the pool was increased from 4 mM to 6 mM such that the process of confirmation run 4 is capable of handling higher oxygen transfer into the solution. This increased reductive power must be balanced against possible antibody over reduction. Nevertheless, based on the results of REACT040 (see **Table 24**), this adaptation represents a change with a low risk of failure. The activity by CEX of REACT040 following the selective reduction step was 91.9%.

In **Figure 11**, the dO₂ chart of confirmation run 4 is shown. Similarly to confirmation run 3, the dO₂ decreased after cysteine addition and remained at low levels, indicating that the reductive power of the cysteine in this experiment is not exhausted, while the product purity requirement determined by CE-SDS is fulfilled

In **Figure 12**, the reaction kinetic of confirmation run 3 and confirmation run 4 are compared with respect to activity by CEX and purity by CE-SDS. The kinetic of activity by CEX is comparable for both experiments; after approximately 2 h a plateau is reached, and thereafter the increase of activity is small in the following 2 hours treatment. The purity by CE-SDS curves are slightly different in the early phase of the reaction, in that the purity drop is more pronounced in the run at the higher cysteine concentration (6 mM). However, as both procedures lead to comparable product quality at the end of the cysteine treatment, both are appropriate to ensure adequate product quality. Nevertheless, considering the effect of elevated oxygen levels in the process solution, the process with the increased cysteine concentration applied in confirmation run 4 is expected to be more robust in respect to variations of the oxygen transfer into the process solution during cysteine treatment. Conclusively, a cysteine concentration of 6 mM (with 13.5 mg antibody = 65.75:1 molar ratio of cysteine:antibody [i.e.,

about 66:1]) was selected for the cysteine treatment step at manufacturing-scale, identical to the conditions of confirmation run 4. We also noted faster kinetics for CEX activity (plateau in 1 hour) in a run performed at a molar ratio of about 275:1 (cysteine:protein) (data not shown).

Summary and Conclusions Drawn from Example 8

We determined that elevated oxygen levels during the cysteine treatment step can have a deleterious effect on activity by CEX, which is likely due to the oxygen abrogating the reductive power of the cysteine, leading to insufficient reduction of C97 of secukinumab. Oxygen uptake from the atmosphere can be managed by varying the cysteine/protein ratio, using defined stirring speeds, or even employing stirring interruptions, when working at production scale.

Based on the results of the confirmation runs, the cysteine concentration was increased from 4 mM (molar ratio of cysteine to protein of 43.84) to 6 mM (molar ratio of cysteine to protein of 65.75), which should counterbalance the oxygen present in reaction solution. Furthermore, to reduce the level of dissolved oxygen present during the incubation phase of the selective reduction reaction, continuous mixing was replaced by mixing 2 minutes every hour during incubation. Although the selective reduction procedure was changed during process confirmation, a clear root cause was identified (level of oxygen transferred into solution during selective reduction), ensuring adequate activity by CEX in the drug substance by the adjusted cysteine treatment step.

Example 9: Characterization of the Selective Reduction Step

The influence of the input parameters protein concentration, cysteine concentration and stirrer speed on the selective reduction step were evaluated in a statistical design. The output parameters activity by CEX and purity by CE-SDS were used to assess product quality after selective reduction step and to define the proven acceptable ranges for the input parameters. Output parameters ranges were defined according to earlier design processes. These limits were used to define the input parameter ranges.

Additionally, the results of the three scale-down model qualification runs and seven selective reduction cycles performed at manufacturing-scale were used to assess product quality after selective reduction for the worst/best case studies regarding reductive power and the dedicated runs for incubation time and incubation temperature.

Example 9.1 – Process Characterization: Response Surface Design

Example 9.1.1 – Experimental Design and Methods.

Protein content, cysteine concentration and stirring mode during incubation (no stirring, continuous stirring at 50 or 100 rpm) were selected as input parameters for the response surface design, see **Table 25**.

Name	Unit	Lower level	Medium level	Upper level
Protein content by ALC	[mg/mL]	10	12.7	15.4
Dilution factor by TITR3 addition ¹⁾	[-]	15 (4.8 mM cysteine, 0.8 mM EDTA)	20 (6.0 M cysteine)	25 (8 mM cysteine; 1.3 mM EDTA)
Stirrer speed during incubation	[rpm]	0	50	100

Table 25: Input parameters. 1) corresponds to a defined cysteine concentration during the selective reduction. “Content by ALC” refers to the concentration of the antibody.

Various cysteine concentrations, expressed as “dilution factor by TITR3 addition,” were employed that reflect process conditions and potential variations, e.g., induced by weighing inaccuracy, inaccuracy of solution addition, etc. The antibody concentration contributes to the ratio of cysteine to protein and was also tested within the present design in order to detect potential interactions. Additionally, the stirrer speed during incubation was tested as an input parameter because, as shown in **Examples 8**, oxygen transferred from the headspace into the solution influences the selective reduction step.

A Central Composite Face Design with 3 center point runs was used. This type of design supports calculation of mathematical models with linear terms, interaction terms and quadratic terms. Secukinumab, INAKT.F (stored below -60°C) originating from manufacturing-scale run B012307 was used. Buffers used for process characterization were Titration Buffer AIN457-TITR1 (1M Tris base, pH 10.8 [pH range ≥ 10.0 , conductivity 0.10 – 0.30 mS/cm]), Titration buffer AIN457-TITR2 (0.3 M o-phosphoric acid, pH 1.4 [pH range ≤ 2.0 ; conductivity range 19.5-22.7 mS/cm]); Titration buffer AIN457-TITR3 (120 mM cysteine-HCL + 20 mM Di-Na-EDTA, pH 8.0 [pH range 7.8-8.2; conductivity range 14.7-18.3 mS/cm]).

The experiments were performed on the qualified scale-down model (described in detail in **Example 10**) in a stirred bioreactor (maximum 2 L volume) at ambient atmosphere with free

air exchange and monitoring of pH, dissolved oxygen, stirrer speed and temperature. The experimental design plan is shown in **Table 26**. All input parameters that were not part of the individual studies were held constant at the respective target according to **Table 27**.

Run	Protein content by ALC	Dilution factor by TITR3 addition	Stirrer speed during incubation	Cysteine	Approx. molar ratio cysteine to protein
	[mg/mL]	[-]	[rpm]	[mM]	[M/M]
REACT069 ¹	12.7	20	50	6.0	69.89
REACT070	12.7	20	0	6.0	69.89
REACT072	12.7	25	50	4.8	55.92
REACT073 ¹	12.7	20	50	6.0	69.89
REACT075	12.7	20	100	6.0	69.89
REACT077	15.4	15	0	8.0	76.85
REACT079	15.4	20	50	6.0	57.64
REACT080	10.0	15	0	8.0	118.36
REACT082	15.4	25	0	8.0	76.85
REACT084	10.0	25	100	4.8	71.01
REACT085	15.4	25	100	4.8	46.11
REACT086	10.0	20	50	6.0	88.77
REACT087	12.7	15	50	8.0	93.19
REACT088 ¹	12.7	20	50	6.0	69.89
REACT089	15.4	15	100	8.0	76.85
REACT090	10.0	25	0	4.8	71.01
REACT091	10.0	15	100	8.0	118.36

Table 26: Experimental design plan. 1) Center Point.

Process parameter	Unit	Target
Load pH	-	8
Load temperature	°C	as is
Protein concentration before pH adjustment and TITR3 addition	mg/mL	13.5
Incubation temperature	°C	37
Temperature after cooling	°C	22
Stirrer speed during heating and cooling	rpm	50
Stirrer speed during incubation ¹⁾	rpm	0

Heating time	min	60
Incubation time	min	240 REACTPT.300 and 300 for REACTP
Cooling time	min	60
pH of REACT.P	-	5.2
Dissolved oxygen	%	as is
Dilution factor with TITR3	-	1:20 (6 mM)

Table 27: Process parameters of the selective reduction step. 1) stirring for every hour for 2 minutes at a speed of 50 rpm.

INAKT.F was thawed in a hand warm water bath and diluted with WFI to target concentration. After pH adjustment, 1 L of the solution was transferred into the qualified 2 L bioreactor equipped with a dO₂ probe and stirrer. The selective reduction was started by adding the calculated volume of cysteine stock solution (TITR3 buffer) to achieve the cysteine target concentration listed in **Table 28**. The solution was heated to incubation temperature of 37 °C within 60 minutes under stirring of 50 rpm. The solution was then incubated at 37°C for 300 min either with continuous stirring at 50 or 100 rpm or with no stirring (0 rpm) according to the experimental design plan. After 300 minutes reaction time (60 minutes heating and 240 minutes incubation) a sample was withdrawn (REACT.PT300). After an additional 60 minutes incubation at 37°C (60 minutes heating and 300 minutes incubation), the solution was cooled to ambient temperature within 60 minutes at a stirrer speed of 50 rpm and another sample was withdrawn (REACT.P, total reaction time 420 min). This procedure makes it possible to evaluate the design at two time points (referred to as “REACT.PT300” and “REACT.P”) and to evaluate potential impact of incubation time.

Example 9.1.2 – Output Parameters of Response Surface Design

The values for the product quality output parameters for samples REACT.PT300 and REACT.P are listed in **Table 28** (ordered by ascending CEX activity in REACT PT300 samples) and **Table 29** (ordered by ascending CEX activity in REACT P samples). In general, the data shows that purity by CE-SDS was high (> 90 %) in all runs, however some runs had a lower activity by CEX (< 90 %). With respect to CEX activity in REACT.PT300 samples (**Table 28**), there was one run with low activity (89.4 %). In this run (REACT085) cysteine concentration was at the low limit, antibody concentration high (hence molar ratio cysteine to protein was the lowest), and stirring speed was the highest (highest oxygen transfer rate). Also, it can be seen

that there is a group of eight runs with high activity (more than 96 %) which contain all runs where no stirring took place.

	Antibody content by ALC	Stirrer speed during incubation	Cysteine conc.	Approx. molar ratio cysteine to protein	React. PT300 activity by CEX	React.P activity by CEX	React. PT300 purity by CE-SDS	React.P purity by CE-SDS
Run	[mg/mL]	[rpm]	[mM]	[M/M]	[%]	[%]	[%]	[%]
REACT085	15.4	100	4.8	46.11	89.4	84.0	95	94
REACT091	10.0	100	8.0	118.36	92.4	91.3	93	94
REACT084	10.0	100	4.8	71.01	94.3	88.9	95	95
REACT072	12.7	50	4.8	55.92	94.5	90.2	96	95
REACT089	15.4	100	8.0	76.85	95.2	93.7	95	94
REACT069 ¹	12.7	50	6.0	69.89	95.3	94.1	95	95
REACT075	12.7	100	6.0	69.89	95.7	88.9	95	95
REACT088 ¹	12.7	50	6.0	69.89	95.7	95.1	95	95
REACT079	15.4	50	6.0	57.64	96.0	94.4	95	95
REACT073 ¹	12.7	50	6.0	69.89	96.1	94.7	95	95
REACT082	15.4	0	8.0	76.85	96.1	96.2	95	96
REACT070	12.7	0	6.0	69.89	96.1	96.5	94	92
REACT087	12.7	50	8.0	93.19	96.5	96.0	94	94
REACT090	10.0	0	4.8	71.01	96.6	97.3	94	95
REACT086	10.0	50	6.0	88.77	96.7	95.4	94	94
REACT077	15.4	0	8.0	76.85	97.0	97.4	92	92
REACT080	10.0	0	8.0	118.36	97.8	97.5	90	91

Table 28: Input parameters and quality output parameter values REACT.PT300 and REACT.P sorted by ascending CEX for REACT.PT300 samples. 1) Center Point.

As shown in **Table 29**, with respect to CEX activity in REACT.P samples, there was a group of six runs resulting in lower activity (< 94 %), which group contains all runs having high stirrer speed. Notably, all runs with no stirring resulted in highest CEX activity (more than 96

%). Also it can be seen from **Table 29**, that CEX activity declined in REACT.P samples (compared to the CEX activity in REACT.PT300 samples) in runs carried out at 100 rpm, and (less pronounced) in runs carried out at 50 rpm. The highest decline in these runs is seen when the molar ratio of cysteine was at the lower end (e.g. decline from 89.4 to 84.0 % in REACT085 where molar ratio was about 46:1 and similar in REACT075, REACT084 and REACT072).

	Antibody content by ALC	Stirrer speed during incubation	Cysteine concen.	Approx. molar ratio cysteine to protein	React. PT300 activity by CEX	React. P activity by CEX	React. PT300 purity by CE-SDS	React. P purity by CE-SDS
Run	[mg/mL]	[rpm]	[mM]	[M/M]	[%]	[%]	[%]	[%]
REACT085	15.4	100	4.8	46.11	89.4	84.0	95	94
REACT075	12.7	100	6.0	69.89	95.7	88.9	95	95
REACT084	10.0	100	4.8	71.01	94.3	88.9	95	95
REACT072	12.7	50	4.8	55.92	94.5	90.2	96	95
REACT091	10.0	100	8.0	118.36	92.4	91.3	93	94
REACT089	15.4	100	8.0	76.85	95.2	93.7	95	94
REACT069 ¹	12.7	50	6.0	69.89	95.3	94.1	95	95
REACT079	15.4	50	6.0	57.64	96.0	94.4	95	95
REACT073 ¹	12.7	50	6.0	69.89	96.1	94.7	95	95
REACT088 ¹	12.7	50	6.0	69.89	95.7	95.1	95	95
REACT086	10.0	50	6.0	88.77	96.7	95.4	94	94
REACT087	12.7	50	8.0	93.19	96.5	96.0	94	94
REACT082	15.4	0	8.0	76.85	96.1	96.2	95	96
REACT070	12.7	0	6.0	69.89	96.1	96.5	94	92
REACT090	10.0	0	4.8	71.01	96.6	97.3	94	95
REACT077	15.4	0	8.0	76.85	97.0	97.4	92	92
REACT080	10.0	0	8.0	118.36	97.8	97.5	90	91

Table 29: Input parameters and quality output parameter values REACT.PT300 and REACT.P sorted by ascending CEX for REACT.P samples. 1) Center Point.

The statistical diagnosis (**Table 30**) indicates statistical significant models for purity by

CE-SDS for both time points and for activity by CEX for REACT.P at the end of the selective reduction. The model for activity by CEX for REACT.PT300 shows lower statistical significance (Q^2 of 0.15).

Output	R^2	Q^2	Model validity	SD of replicates
REACT.PT300				
activity by CEX	0.75	0.15	0.4	0.4
purity by CE-SDS	0.98	0.89	-0.2	0.0
REACT.P				
activity by CEX	0.92	0.75	0.48	0.5
purity by CE-SDS	0.87	0.59	-0.2	0.0

Table 30: Model Diagnosis for the product quality output parameters of REACT.PT300 and REACT.P. A model that is in good agreement with the data will have a R^2 and Q^2 close to 1.0 and Model validity above 0.25. Models with low statistical significance have low R^2 and Q^2 values.

Example 9.1.2.1 – Modelling Evaluation of Product Quality Output Parameters at Time Point 1 (REACT.PT300)

The coefficient plot and the contour plot for CEX activity at time point REACT.PT300 demonstrate that the input parameter stirrer speed (stir) during incubation affects activity by CEX (data not shown). The parameters protein concentration (contA) and cysteine concentration (dil-c) show no statistically significant effect on activity by CEX within the investigated range. In order to achieve a high activity by CEX the stirrer speed during incubation should be low.

As shown in **Table 29**, all the experiments at time point REACT.PT300 had at least 90% purity by CE-SDS. The coefficient plot (data not shown) demonstrates that all three input parameters affect purity by CE-SDS. There are also quadratic terms for content by ALC (contA) and stirrer speed (stir). There are interaction terms of dilution factor by TITR3 addition (dil-c representing the cysteine concentration) and stirrer speed, as well as an interaction term of the dilution factor by TITR3 addition and content by ALC. In the contour plot (data not shown), the model coefficients as visualized indicate that high stirrer speed, high dilution factor by TITR3 addition (respectively a low cysteine concentration), and medium to high content by ALC positively influence purity by CE-SDS. All these settings reduce the reductive power of cysteine and positively influence antibody integrity. Nevertheless, as all runs have high purity by CE-SDS, and the whole investigated input parameter range is appropriate to ensure adequate purity by CE-SDS after selective reduction.

Example 9.1.2.2 – Modelling of Product Quality Output Parameters at Time Point 2 (REACT.P)

The coefficient plot in **Figure 13A** and contour plot in **Figure 13B** for CEX activity at time point REACT.P confirms that especially the input parameter stirrer speed (stir) affects activity by CEX. Thus, in order to achieve a high activity by CEX, minimal stirring should be performed during incubation to limit oxygen transfer into the solution. Also, low dilution factor by TITR3 addition (dil-c, respectively a high cysteine concentration) is beneficial for activity by CEX after selective reduction. In addition, the interaction of stirrer speed and dilution by TITR3 addition has a significant – although small - impact on activity by CEX, whereas the interaction of content by ALC and dilution factor by TITR3 addition is borderline. The influence of content by ALC is not significant. The results clearly indicate that conditions which increase the reductive power, e.g., limited oxygen transfer by minimal stirring speed and higher cysteine concentration by lower dilution, result in higher activity by CEX.

As shown in **Table 29**, all the experiments at time point REACT.P had at least 90% purity by CE-SDS. The coefficient and contour plots (data not shown) for CE-SDS purity demonstrate that the input parameters content by ALC (representing protein concentration), dilution factor by TITR3 addition (dil-c, representing the cysteine concentration), and stirrer speed (stir, representing the oxygen transfer) have only a small effect on the purity by CE-SDS. The stirrer speed exhibits a quadratic effect and the interaction of dilution factor by TITR3 with stirrer speed is statistically significant. Purity by CE-SDS is improved by medium to high stirrer speed and high dilution factor by TITR3 addition. These settings reduce the reductive power of cysteine and positively influence antibody integrity. Nevertheless, as all runs meet the specified range for purity by CE-SDS, the whole investigated input parameter range is appropriate to ensure adequate purity by CE-SDS after selective reduction step.

Example 9.1.2.1 – Relationship of Dissolved Oxygen Levels to Output Parameters

The oxygen profiles of the reactions shown in **Table 26** were analyzed to determine the influence of the input parameters stirring speed, antibody content and cysteine concentration on levels of oxygen. Graphs of these oxygen profiles are provided in **Figure 14** (note: 0 to 1.00 on the x-axis corresponds to the 1 h heating phase, where stirring was at 50 rpm in all runs; 1.00 to

6.00 corresponds to the 5 h incubation phase, where stirring was either at 0 rpm [i.e., no stirring] 50, or 100 rpm; and 6.00 to 7.00 corresponds to the 1 h cooling phase, where stirring was at 50 rpm in all runs). All runs show a decrease of dissolved oxygen to low levels in the early phase. Notably, this decrease was less pronounced in runs with low antibody content (see runs REACT090 and REACT080 in the 0 rpm series (**Figure 14A**), REACT086 in the 50 rpm series (**Figure 14B**) and REACT084 and REACT091 in the 100 rpm series (**Figure 14C**)) and most pronounced in runs having high antibody content (see profiles of the runs with 15.4 mg/mL graphs of **Figure 14**).

In all runs without stirring during incubation (**Figure 14A**), the dO₂ level remained below about 20% during both the incubation and cooling phase. Also, except for REACT086, in the runs using 50 rpm stirring during incubation (**Figure 14B**), the dO₂ level stayed below about 20% during both the incubation and cooling step despite the oxygen transfer enabled in the incubation phase by 50 rpm stirring. However, in run REACT072, in which the cysteine concentration was low (4.8mM), a slight increase of the oxygen level can be seen in the final cooling phase. The runs with 100 rpm stirring during incubation (**Figure 14C**) showed very different profiles. First, as mentioned above, runs having low antibody content (10 mg/mL) (REACT084 and REACT091) showed a slower dO₂ decrease, which continued during the incubation phase. However, in run REACT084, in which the cysteine concentration was low (4.8 mM), the dO₂ level then increased during hours 5.00 and 6.00, the last hour of the incubation, to almost saturated level. Such an increase was observed also in run REACT085 even at an earlier time (hour 4.00). REACT085 also used a low level of cysteine, but high antibody content (15.4 mg/mL). An increase in the dO₂ level was also observed in run REACT075 during the cooling phase (hours 6.00 to 7.00). In run REACT075, the cysteine concentration was at the medium level (6.0 mM).

These profiles clearly indicate that oxygen is consumed in the reaction mixture at a rate higher than it is transferred into the mixture from the headspace, and that oxygen consumption is faster in runs having higher antibody concentration. This in turn suggests that the antibody itself may trigger dO₂ consumption. The increase of oxygen in later phases of selective reduction in runs having low cysteine levels indicates that the oxygen consumption is linked to a reaction with cysteine (Cys-SH), probably as follows: $4 \text{ Cys-SH} + \text{O}_2 \rightarrow 2 \text{ Cys-SS-Cys} + 2 \text{ H}_2\text{O}$ and/or

$2 \text{ Cys-SH} + 3 \text{ O}_2 \rightarrow 2 \text{ Cys-SO}_3\text{H}$. Indeed, the rate was fast enough to consume all cysteine after approximately 4 h in run REACT085, and after 5 h in run REACT084 (**Figure 14C**). After consumption of the cysteine, dO₂ levels then recovered to saturated levels (100% dO₂).

To study the impact of antibody on oxygen consumption, we performed two additional runs at 50 mL scale using 6 mM cysteine and a stirrer speed of 50 rpm. In one run, the antibody concentration was 12.7 mg/ml (molar ratio cysteine:antibody = 69.89:1, i.e., about 70:1); in the other run the antibody concentration was zero. As shown in **Figure 14D**, in the run without antibody there is a decrease of dO₂ to roughly 75% in the first hour, during which time the temperature of the reaction was raised from 20°C to 37°C. This 75% decrease of dO₂ fits well to the reported decrease of dO₂ saturation concentration from 8.9 mg/L at 20°C to 6.6 mg/L at 37°C (U.S. Geological Survey TWRI Book 9, April, 98). The dO₂ level remains constant during the remainder of the incubation phase. Once the cooling phase begins (after 6 hours), there is an increase in the dO₂ level due to the lowering of temperature from 37°C to 20°C. Thus, in the absence of antibody there is no consumption of oxygen in the solution, or consumption of oxygen by cysteine occurs at a low rate, such that the oxygen transfer from the headspace caused by stirring immediately compensates for that consumption. In contrast, in the run with antibody, there is a steady decrease in the dO₂ level up to approximately 5.5 hours. In the final 0.5 hour of the incubation phase there is a slight increase in the dO₂ level, followed by a stronger increase in the dO₂ level during the cooling phase (i.e., from hours 6 to 7). As such, in this antibody-containing run, probably most, if not all, cysteine was consumed after 5.5 hours. In fact, amino acid analysis of this run (data not shown) showed an elevated level of cystine, corresponding to almost complete consumption of the cysteine.

With this information, it is possible to correlate the low activity by CEX found in some runs with their dO₂ profiles. Run REACT085 (**Figure 14C**) (molar ratio cysteine: antibody about 46:1) had 89.4 % activity by CEX at REACT.PT300 (sample from 5.00 hour time-point) (see **Table 28**) because cysteine was consumed after the 4.00 time-point, leading to incomplete reduction of the antibody. REACT085 (**Figure 14C**) at the later time point REACT.P (sample from endpoint 7.00 hour) had even less activity by CEX (84.0 %, see **Table 29**) because the high level of oxygen in the last phase of the reaction, in the absence of cysteine, led to oxidative degradation of the antibody. Runs REACT084 (**Figure 14C**) (molar ratio cysteine: antibody

about 71:1) and REACT075 (**Figure 14C**) (molar ratio cysteine: antibody about 71:1) were the two other runs with relatively low activity at the later time point REACT.P (both 88.9%), although the activity was high at the earlier REACT.PT300 time point. In these runs, the reduction of the antibody seemed to be complete at time-point 5.00 hours (REACT.PT300 time point); however, degradation occurred by the later time point REACT.P due to the increased dO_2 level and absence of cysteine. Similarly, the relatively low activity found at later time point REACT.P for run REACT072 (**Figure 14B**) (90.2%) (molar ratio cysteine: antibody about 56:1) is explained by the increase in the oxygen level during the cooling phase of the reaction. The high activities obtained in runs at 0 rpm (even in run REACT082, in which antibody content was high and cysteine concentration was low [molar ratio cysteine: antibody about 46:1]) occurred because the oxygen transfer rate was low and consumption of cysteine was therefore nominal, ensuring complete reduction (and also protection) against oxidative degradation during the later cooling phase of the reduction.

In conclusion, oxygen transfer should be kept low in the reduction system because oxygen consumes the reductant (cysteine), which consumption is mediated (or accelerated) by the antibody itself. There is therefore a two-fold impact of this cysteine consumption: 1) a loss of reductive power of the cysteine leads to incomplete deblocking of CysL97-SH at earlier time point REACT.PT 300; and 2) if there is no residual cysteine available to protect deblocked Cys97L-SH during the time between REACT.P300 and REACT.P, then reoxidation of deblocked Cys97L-SH can occur.

Example 9.2 – Process Characterization: Worst/Best Case Scenarios

The main purpose of the worst / best case studies was the characterization of the input factors of the selective reduction step that were not tested during a design of experiments (DOE) study, but were assessed as possibly important in the process risk analysis. The data are also used to define the proven acceptable ranges and to support classification of the input parameters based on impact of their effect on product attributes.

Example 9.2.1 – Experimental Design and Methods.

Increased heating time was assessed as possibly significant, as it increases the oxygen transfer into the selective reduction solution due to extended stirring duration, which can lead to reduced reductive power by conversion of cysteine to cystine and hence less selective reduction

of the antibody. For the same reason, enhanced initial dO_2 level was assessed as possibly significant input parameter.

The oxidation of the sulfhydryl group of cysteine is pH dependent. Consequently, the influence of pH was assessed as possibly significant input parameter.

High reductive power can lead to over-reduction of the antibody and an extended cooling time could promote antibody reassembly. Thus, extended cooling time was also assessed as possibly significant input parameter

Although the temperature was already tested during earlier process development (32 – 42 °C), it was assessed as possibly significant. Typically, the chemical reaction rate decreases with a lower temperature. Therefore, the kinetic was determined at the upper and lower limit in dedicated runs.

The incubation time was also investigated. A low incubation time could lead to higher levels of over-reduced product in the selective reduction pool. Conversely, the extended incubation time could lead to lower activity by CEX as shown in the response surface design results (see **Table 29**, c.f. CEX activity of REACT.PT300 and REACT.P samples).

Three worst/best case sequences were defined to assess the influence of the parameters described above on the selective reduction step. The input parameters to assess the reductive power in a worst/best case sequence are shown in **Table 30**. The second worst/best case sequence was created to evaluate the influence of temperature on the selective reduction step (**Table 31**). The influence of the incubation time was evaluated in the third worst/best case study according to the input parameter listed in **Table 32**. The product quality output parameters used to assess the results of the worst/best case studies are activity by CEX and purity by CE-SDS.

Name	Unit	Lower limit	Upper limit
Heating time including stirring	[min]	45	90
Cooling time including stirring	[min]	45	90
Dissolved oxygen at start	[%]	60	100
Process pH	[-]	7.8	8.2

Table 30: Input parameters to assess influence of reductive power.

Name	Unit	Lower limit 1	Lower limit 2 ¹⁾	Upper limit
------	------	---------------	-----------------------------	-------------

Incubation temperature	[°C]	18	32	42
------------------------	------	----	----	----

Table 31: Input parameters to assess influence of incubation temperature. 1) 32°C was tested, as the reaction rate of the selective reduction at 18°C was not sufficient, resulting in incomplete reactivation.

Name	Unit	Lower limit	Upper limit
Incubation time	[min]	210	330

Table 32: Input parameters to assess influence of incubation time.

Dedicated runs were performed to evaluate the influence of additional possibly important parameters. Secukinumab, INAKT.F (stored below -60°C) originating from manufacturing-scale run B008530 was used. Buffers AIN457-TITR1, AIN457-TITR2 and AIN457-TITR3 are as described in **Example 10**. The experiments were performed on the qualified scale-down model (described in detail in **Example 10**) in an open stirred bioreactor (maximum 2 L volume) with pH, dissolved oxygen, stirrer speed and temperature monitoring. For all runs, the process parameters from **Table 27** were applied. All input parameters that were not part of the individual studies were held constant at the target in **Table 27** within the operating range of the standard sequence.

Example 9.2.2 – Results of Worst / Best Scenarios

The experimental design plan and the output parameters (purity by CE-SDS and activity by CEX) for investigation of the influence of heating time, cooling time, dO₂ at start and process pH, are shown in **Table 33**. The experimental design plan and the output parameters (purity by CE-SDS and activity by CEX) for investigation of the influence of incubation temperature are shown in **Table 34**. The experimental design plan and the output parameters (purity by CE-SDS and activity by CEX) for investigation of the incubation time are displayed in **Table 35**.

As shown in **Table 33**, duplicate runs with low reductive power, as well as high reductive power yielded high activity by CEX as well as purity by CE-SDS. Hence, the process is able to cover variations of the input parameters heating time and cooling time (including stirring), dO₂ level at start, and process pH within the tested ranges.

Run	Heating time [min]	Cooling time [min]	Dissolved oxygen at start [%]	Process pH [-]	Activity by CEX [%]	Purity by CE-SDS [%]
REACT076 ¹⁾	90	90	100	7.8	96.4	94
REACT078 ²⁾	45	45	60	8.2	97.4	93

REACT098 ¹⁾	90	90	100	7.8	95.8	95
REACT100 ²⁾	45	45	60	8.2	94.1	94

Table 33: Experimental design plan and output parameters for investigation of the influence of heating time, cooling time, dO₂ at start and process pH. 1) duplicate runs run with lower reductive power because of lower pH, higher dO₂ starting level and longer heating and cooling time. 2) duplicate runs with higher reductive power because of higher pH, lower dO₂ starting level and shorter heating and cooling time.

The results of the experiments to evaluate the influence of the incubation temperature are shown in **Table 34**. It can be seen that purity by CE-SDS was always high, whereas high activity by CEX was obtained only in the runs of 32°C and above.

Run	Incubation temperature [°C]	Activity by CEX [%]	Purity by CE-SDS [%]
REACT093	42	94.9	94
REACT095	42	95.2	94
REACT107	32	95.9	95
REACT109	32	97.3	95
REACT092	18	91.2	93
REACT099	18	90.4	95
REACT106 ¹⁾	37	95.6	96

Table 34: Experimental design plan and output parameters for investigation of the influence of incubation temperature. 1) Additional center point run.

Additionally the kinetic of the activity by CEX was determined at the different incubation temperatures. These results are shown in **Figure 15**. It can be seen that kinetics at 42°C is faster than at 32°C, especially in the later phase (60 min and more). However, at both temperatures the same plateau is reached at 240 min and REACT.P. At 18°C, the reaction rate is slower, leading to lower increase of activity by CEX within the tested step duration.

The results of the experiments to evaluate the influence of the incubation time are shown in **Table 35**. In all cases high activity and purity was obtained.

Run	Incubation time [min]	Activity by CEX [%]	Purity by CE-SDS [%]
REACT081 ¹⁾	210	96.7	95
REACT083 ²⁾	330	96.3	95
REACT094 ¹⁾	210	95.2	93
REACT096 ²⁾	330	96.2	94

Table 35: Experimental design plan and output parameters for investigation of the influence of incubation time. 1)

low incubation time. 2) high incubation time.

Example 9.3 – Classification and Justification of Process Parameters

Example 9.3.1 – Response Surface Design Study: Acceptable Ranges

Purity by CE-SDS was at least 90% in all experiments at time point 1 (REACT.PT300) and at the end of the selective reduction (REACT.P). To achieve high levels of purity by CE-SDS, the input parameter dilution factor by TITR3 addition should be set to high and stirrer speed should be set center to high. These conditions correspond to a low reductive power due to lower cysteine concentration and increased oxygen transfer over the headspace. The input parameter content by ALC, i.e., the antibody concentration, has significant influence at time point 1 (REACT.PT300) during the selective reduction step, and the ideal is around the center point. The influence of antibody concentration (content by ALC) on the absolute value of purity by CE-SDS is minor within the investigated ranges. Nevertheless, the specified purity value was met in all experiments and the whole range investigated for the parameters content by ALC (antibody concentration), dilution factor by TITR3 addition (cysteine concentration) and stirrer speed (oxygen transfer) is appropriate to ensure adequate purity by CE-SDS after selective reduction.

Activity by CEX was at least 90% for all but REACT085 (see **Table 28**) at time point 1 (REACT.PT300). This experiment was performed with stirrer speed at its high level, representing a higher oxygen transfer. Despite expectations, a longer incubation time (i.e., 60 additional minutes, time point REACT.P) was not particularly beneficial for the output parameter activity by CEX, as an additional two experiments (REACT075 and REACT084, see **Table 29**) displayed activity by CEX below 93%. These experiments have either high stirrer speed or high level of dilution factor by TITR3 addition, representing a low cysteine concentration and increased oxygen transfer over the headspace. These results suggest achieve highest activity by CEX, the input parameter stirrer speed should be set to a low level, implying a low oxygen transfer, and dilution factor by TITR3 addition should be set to center point or low level, implying a higher cysteine concentration. The input parameter content by ALC, representing the antibody concentration, had no significant influence on activity by CEX in these experiments.

The input parameter stirrer speed which mediates (oxygen transfer from headspace into solution is important for the selective reduction step and should be set to its lower level to ensure

adequate activity by CEX. According to an additional manufacturing-scale run (B018838, data not shown) a maximum of 15 minutes stirring per hour was tested, resulting in an acceptable product quality after the selective reduction step. However, the results of **Example 8** reveal that the oxygen transfer into solution by stirring is unique for each scale and setup and is controlled indirectly. Therefore, stirring conditions during incubation should be assessed in case of, e.g., process changes or scale-up. Stirring during incubation, which indirectly represents the level of dO_2 transfer into the reaction solution, is classified as a significant input parameter from a process perspective.

The input parameters content by ALC, namely the antibody concentration, and dilution factor by TITR3 addition, namely the cysteine concentration, show statistically significant effects, but are well-controlled and have minor influence on the selective reduction step. Hence, they are classified as non-key from a process point of view. As long as the mixing time during incubation is within the acceptable range, variations within the investigated ranges of cysteine concentration and antibody concentration will lead to adequate process performance and product quality after the selective reduction step. A list with the proven acceptable ranges for these input parameters is shown in **Table 36**.

Name	Unit	Lower limit	Upper limit
Mixing time during incubation	[min/h]	0	15 ¹⁾
content by ALC	[mg/mL]	10	15.4
dilution factor by TITR3 addition	[-]	15 (8 mM cysteine, 1.3 mM EDTA)	25 (4.8 mM cysteine, 0.8 mM EDTA)

Table 36: Acceptable ranges for stirrer speed during incubation, content by ALC and dilution factor by TITR3 addition (cysteine concentration). 1) according to characterization run at manufacturing-scale, batch B018838, SITE A, at 4h incubation time.

Example 9.3.2 – Worst/Best Case Studies: Acceptable Ranges

The input parameters heating time and cooling time (including stirring), dO_2 level at start and process pH have no significant influence regarding the product quality attributes activity by CEX and purity by CE-SDS within the investigated range displayed in **Table 37**. However, due to the experience of the response surface design, too high oxygen transfer into the selective reduction solution should be avoided, as it reduces reductive power. The oxygen transfer is

specific for each scale and setup, but in an aerobic process is controlled indirectly by other process conditions, such as stirring and airflow into the headspace. Therefore, stirring during heating and cooling, which affects the oxygen transfer, must be carefully evaluated, especially in case of, e.g., a process changes, process transfer, or scale-up. Accordingly, the heating and cooling time (including stirring) are classified as significant parameters from a process point of view. Although a range for heating and cooling time (including stirring) of 45 - 90 minutes was tested, the acceptable range was set to ≤ 90 minutes, as 90 minutes heating and cooling time (including stirring) reflects the worst case with respect to oxygen transfer leading to appropriate activity by CEX and purity by CE-SDS.

Name	Unit	Lower limit	Upper limit
Heating time including stirring	[min]	45	90
Cooling time including stirring	[min]	45	90
Dissolved oxygen at start	[%]	60	100
Process pH	[-]	7.8	8.2

Table 37: Tested ranges of heating time, cooling time, dissolved oxygen at start and process pH.

The results of the experiments investigating the input parameter incubation temperature demonstrate that an incubation temperature of 18°C leads to slow reaction kinetics resulting in insufficient reactivation. The low temperature is not sufficient to increase activity by CEX within an incubation time of 240 min to levels higher than 93.0 %. Therefore, the tested range of 18 – 42°C is narrowed to 32 – 42°C to ensure adequate activity by CEX after the selective reduction step, although the product quality attribute purity by CE-SDS can be achieved within an incubation temperature range of 18 – 42°C. The acceptable range for incubation temperature is therefore a lower limit of 32°C and an upper limit of 42°C.

Finally, the tested range of 210 – 330 min incubation time ensures adequate activity by CEX and purity by CE-SDS after selective reduction step.

Summary and Conclusions Drawn from Example 9

Based on the results of the process characterization study, the acceptable ranges for the investigated process input parameters were defined. The selective reduction step is characterized by the two product quality output parameters purity by CE-SDS and activity by CEX. The parameter classification and acceptable ranges are summarized in **Table 38**. The ratio of

cysteine:antibody for the PAR is about 46:1 to about 118:1. The ratio of cysteine:antibody for the SOR is about 54:1 to about 83:1.

Input Parameter	Unit	Suggested Operating Range (SOR)	Proven Acceptable Range (PAR)
Content by ALC ¹⁾	[mg/mL]	12.0 – 15.0	10.0 – 15.4
Dilution factor by TITR3 addition (cysteine concentration) ⁵⁾	[-]	1:18 – 1:22 (5.5 – 6.7 mM cysteine, 1.1 – 0.9 mM EDTA)	1:15 – 1:25 (4.8 – 8.0 mM cysteine, 1.3 – 0.8 mM EDTA)
Dissolved oxygen at start	[%]	≥ 80	≥ 60
Heating time including stirring	[min]	≤ 75	≤ 90
Mixing time during incubation	[min / hour]	0 – 5	0 – 15 ⁴⁾
Incubation time	[min]	240 – 300	210 – 330
Cooling time including stirring	[min]	≤ 75	≤ 90
Process pH ²⁾	[pH]	7.9 – 8.1	7.8 – 8.2
Process temperature ³⁾	[°C]	35 – 39	32 – 42

Table 38: Input parameters of selective reduction step including classification. 1) after dilution with WFI before pH adjustment and TITR3 addition, 2) before heating. 3) during incubation. 4) according to characterization run at manufacturing-scale, batch B018838, at 4 h incubation time. 5) related to intermediate volume after addition of WFI and pH adjustment with TITR1.

At the beginning of the selective reduction step, oxygen moderates the reductive power of the cysteine, while continuous transfer of oxygen into the solution results in an increased oxygen level and less antibody activity by CEX. However, at the end of the selective reduction step, the introduction of oxygen could enhance the formation of the disulfide bonds in the antibody that were disassociated at the beginning of the reaction when the reductive power was high, and therefore oxygen is important to ensure adequate purity by non-reducing CE-SDS. The importance of the oxygen transfer for antibody activity is demonstrated by the results of the response surface design study, in which a significant effect of the stirrer speed on oxygen transfer, and hence product activity by CEX, was observed (see **Table 28** and **29**). However, stirrer speed has a different effect on the level of dO₂ depending on the size of the vessel, the size of the stirrer, the type of stirrer, etc. Thus, it is important to identify a variable that can be used to compare the oxygen transferred into a solution between physical setups.

Example 10: Oxygen Transfer Rate (k_{La}^*) of the Scale-Down Model

In the previous example, we showed that, *inter alia*, the amount of dO_2 present during selective reduction, particularly during the incubation step has a strong influence on the quality and activity of secukinumab. When selective reduction is performed under aerobic conditions, the level of oxygen in the reaction is not controlled directly, but via other operating conditions, e.g., stir speed and airflow to the headspace. The physical setup of each reaction also influences the level of oxygen present in the reaction mixture (the equipment and operating conditions together form a “system”). “ k_{La}^* ” can be used to compare the oxygen transferred into a solution between physical setups and during particular antibody processing steps (see, e.g., Garcia-Ochoa and Gomez (2009) *Biotechnology Advances* 27:153-176; Bandino et al. (2001) *Biochem. Engineering J.* 8:111-119; Juarez and Orejas (2001) *Latin Am. Appl. Res.* 31:433-439; Yange and Wang (1992) *Biotechnol. Prog.* 8:244-61). The k_{La}^* represents the amount of oxygen transferred into a solution over time via the headspace without sparging. This value is specific for each setup and scale, and depends on stirrer type, stirrer speed, filling volume and surface area of the solution in contact with the headspace, which is influenced by the individual geometry of each vessel. While the k_{La}^* of each physical setup differs, because the level of oxygen in the solution during the selective reduction step significantly effects the activity and integrity of secukinumab, we expect that the selective reduction step, when performed in systems displaying similar k_{La}^* ranges, will lead to preparations of secukinumab having similar quality.

The k_{La}^* cannot be directly determined in the oxygen transfer experiments. Instead, the dO_2 in a test solution is replaced by nitrogen and the increase of dO_2 over time is monitored using a calibrated dO_2 probe, which allows creation of an experimental dO_2 curve. Thereafter, the k_{La}^* value is calculated for the particular system by adapting the experimental dO_2 curve to a saturation curve (e.g., using Mathcad®) according the equation shown below:

$DO = C \times (1 - e^{-k_{La}^* \times (t - t_0)})$, where DO = the measured value of dissolved oxygen, C is the saturation value of oxygen (meaning 100 % when stirred infinitely and saturation is achieved), $e = 2.718281\dots$, t = time point corresponding to the DO value, and t_0 = starting time point.

The equation represents the integrated form of an empirical formula established for determination of the oxygen transfer into solutions (k_{La}^* value). The formula was confirmed by

different authors in various experiments (Doran, P. M. 1995. Bioprocess Engineering Principles, Academic Press, San Diego, California, chapter 9.10.2, p. 210-213).

Example 10.1 – Experimental and Statistical Design Methods

Because the dissolved oxygen level is an important influencing factor for redox reactions, the oxygen transfer was assessed for qualification of a scale-down model and for comparison with manufacturing-scale.

A statistical design was used to determine the dependency of the k_{La}^* value with the input parameters of volume, air-flow into headspace, stirrer-speed and stirrer type. The input parameters are listed in **Table 39**. The factors were investigated on 3 levels, each according to a statistical design plan. The output parameter is the k_{La}^* value.

Name	Abbreviation	Unit	Lower limit	Upper limit
Volume	Vol	[L]	0.7	1.7
Airflow (headspace)	Air	[L/min]	0	0.5
Stirrer speed	Stir	[rpm]	50	200

Table 39 - Input parameters for k_{La}^* value determination of the scale-down model.

To determine the k_{La}^* value for each sample, a 2 L bioreactor was filled with water and heated to 37 °C. Airflow was controlled by a mass-flow meter and applied to the headspace. Thereafter, the solution was sparged with nitrogen gas to remove dO_2 from the water. Then, constant stirring was applied using a rushton turbine (radial). Over the course of time, the level of dO_2 in the water was recorded using an oxygen probe (calibrated at room temperature, about 18-25°C) until the dO_2 reached 90%. Thereafter, the k_{La}^* value was calculated by adapting the experimental dO_2 curve to a saturation curve as described above.

A Central Composite Face Centered design (CCF) with 3 center point runs was used. The design was chosen to determine the correlation between the most important process parameters and their influence on k_{La}^* . This type of design supports calculation of mathematical models with linear, interaction and quadratic terms.

Example 10.3 – Output Parameter (k_{La}^*) Values

The values for the experimental conditions and the process performance output parameter k_{La}^* are listed in **Table 40**.

Run	Airflow	Volume	Stirrer speed	Measured k_{La}^*
[-]	[L/min]	[L]	[rpm]	[1/h]
1	0.50	0.7	50	0.83
2	0.10	0.7	200	0.70
3	0.50	0.7	200	3.58
4	0.10	0.7	50	0.41
5	0.25	0.7	125	1.97
6	0.25	1.2	125	0.97
7	0.25	1.2	125	0.86
8	0.25	1.2	125	0.98
9	0.25	1.2	200	1.36
10	0.25	1.2	125	0.77
11	0.50	1.2	125	1.07
12	0.25	1.2	125	1.05
13	0.10	1.2	125	0.73
14	0.25	1.2	50	0.54
15	0.25	1.2	125	0.86
16	0.50	1.7	200	0.73
17	0.50	1.7	50	0.22
18	0.00	1.7	200	0.48
19	0.25	1.7	125	0.62
20	0.00	1.7	50	0.24

Table 40 - Statistical design – performed sequence and experimental conditions.

Example 10.4 – Statistical Diagnostic for Output parameter k_{La}^*

The quality of the models is represented by 4 tools; namely the R^2 , Q^2 , Model validity and the Standard Deviation (SD) of replicates. A model that explains the data well will have a R^2 and Q^2 close to 1.0 and Model validity above 0.25. In a model analysis, it was recognized that evaluation of the k_{La}^* results without transformation leads to a model in which the residuals do not meet the requirements of a normal distribution. This indicates that a transformation of the k_{La}^* value is necessary to describe the results in an adequate manner, which is sometimes observed for common k_{La}^* measurements in bioreactors. The model diagnostics for the process performance output parameter are $R^2 = 0.98$; $Q^2 = 0.91$ (following transformation by the power of -0.5); model validity = 0.56 and standard deviation of replicates = 0.06, indicating a model with high statistical significance.

Example 10.5 – Description of Mathematical Model

In the N-plot of residuals the standardized residuals (residuals of the responses divided by the standard deviation) are plotted on the horizontal axis and the normal probability on the vertical axis. Outliers can be graphically identified when they lay outside of ± 4 standardized residuals. Also, non-linear plots can indicate a model which needs an output parameter transformation (“Design of Experiments – Principles and Applications,” (1999-2008) MKS Umetrics AB, ed. Eriksson et al.). The N-plot of residuals for the process output parameter k_{La}^* (data not shown) indicates a good fit of the experimental result with the model. The values fit close to the regression line, which means that the model transformation is adequate (the standardized residuals meet the requirements of a normal distribution). Additionally, all experimental values are inside the range of ± 3 standardized residuals, indicating that the measured values do not include any outliers.

The coefficient plot for the process output parameter (data not shown), indicates that the input parameters, airflow, volume and stirrer speed have significant influence on the process parameter k_{La}^* . The contour plot (data not shown) illustrates the effect of airflow, volume and stirrer speed on k_{La}^* - the higher the airflow, the lower the volume (i.e. larger the headspace) and the higher the stirrer speed, the higher is the oxygen transfer over the head-space expressed as k_{La}^* .

Using the contour and coefficient plots, the corresponding k_{La}^* values of the experimental conditions were predicted with MODDE 8.02 by use of the statistical model. The results are shown in **Table 41**, which demonstrates the k_{La}^* range that was tested during process characterization (see **Example 9**) using the scale-down model.

	Airflow [L/min]	Volume [L]	Stirrer speed [rpm]	Predicted k_{La}^* [1/h]
Low stirrer speed	0	1.2	0	0.18
Center point	0	1.2	50	0.27
High stirrer speed	0	1.2	100	0.37
Worst/best case experiments	0	1.0	50	0.28

Table 41 - Parameter settings and corresponding k_{La}^* values used during process characterization studies.

Example 11: Oxygen Transfer Rate (k_{La}^*) at Manufacturing Scale

The k_{La}^* during constant stirring (i.e., the type of stirring used during the heating and cooling phase of the selective reduction process) at manufacturing-scale (1800 L vessel) was determined.

Example 11.1 - Determination of the k_{La}^* value at Manufacturing-Scale at SITE A

Example 11.1.1 – Experimental and Statistical Design Methods

A statistical design was used to determine the dependency of the k_{La}^* within the tested input parameter ranges. The input parameters are listed in **Table 42** below. The factors are investigated on 3 levels each according to the statistical design plan. The output parameter is the k_{La}^* value.

Name	Abbreviation	Unit	Lower limit	Upper limit
Volume	Vol	[L]	400	800
Stirrer speed	Stir	[rpm]	100	300
Temperature	Temp	[°C]	16	48

Table 42: Input parameters for k_{La}^* value determination at manufacturing-scale SITE A.

These experiments at SITE A were performed in a stainless steel vessel with a maximum working volume of 1800 L, a height of 2.7 m and a diameter of 1.0 m. To determine the k_{La}^* value for each sample, the vessel was filled with water and heated to the indicated temperature. Thereafter, nitrogen gas was applied to the headspace to remove dO_2 from the water. Then, constant stirring was applied (using a propeller stirrer from the bottom). Over the course of time, the level of dO_2 in the water was recorded using an oxygen probe (calibrated at room temperature, e.g., 18-25 °C) until the dO_2 reached 90%. Thereafter, the k_{La}^* value was calculated by adapting the experimental dO_2 curve to a saturation curve as described above. The input parameters and the corresponding results are shown in **Table 43**.

A Central Composite Face Centered (CCF) design with 4 center points was used. The design was chosen to determine the correlation between the most important input parameters and their influence on the oxygen transfer. This type of design supports calculation of mathematical models with linear, interaction and quadratic terms.

Example 11.1.2 – Output Parameter (k_{La}^*) Values

The values for the experimental conditions and process performance output parameter k_{La}^* are listed in **Table 43**.

Run [-]	Volume [L]	Stirrer speed [rpm]	Temperature [°C]	Measured k_{La}^* [1/h]
1	602	99	33	0.20
2	410	99	16	0.47
3	605	200	33	0.29
4	596	201	16	0.22
5	419	300	49	1.69
6	804	100	49	0.12
7	388	201	32	1.01
8	808	300	16	0.22
9	637	201	33	0.27
10	803	201	33	0.20
11	600	300	33	0.37
12	808	100	16	0.09
13	400	300	16	1.24
14	815	300	49	0.39
15	612	201	49	0.29
16	616	201	33	0.30
17	608	200	33	0.23
18	396	100	49	0.97

Table 43: Statistical design – performed sequence and experimental conditions.

Example 11.1.3 – Statistical Diagnostic for Output parameter k_{La}^*

The quality of the models is represented by 4 tools, namely the R^2 , Q^2 , Model validity and the Standard Deviation (SD) of replicates. A model that explains the data well will have a R^2 and Q^2 close to 1.0 and Model validity above 0.25. Models with low statistical significance have low R^2 and Q^2 values. It was recognized that evaluation of the k_{La}^* results without transformation leads to a model in which the residuals do not meet the requirements of a normal distribution. This indicates, that a transformation of the k_{La}^* is necessary to describe the results in an adequate manner. To gain a robust model, the output parameter was transformed by the power of -0.5. The model diagnostics for the process performance were $R^2=0.95$, $Q^2 = 0.88$, Model validity = 0.82, SD = 0.15, which indicates a model with high statistical significance.

Example 11.1.4 – Description of Mathematical Model

The N-plot of residuals for the process output parameter k_{La}^* value (data not shown) indicate a good fit of the experimental result with the model. The values fit close to the regression line, which means that the model transformation is adequate (the standardized residuals meet the requirements of a normal distribution). Additionally, all experimental values are inside the range ± 3 standardized residuals, indicating that the measured values do not include outliers.

A coefficient plot (data not shown) indicates that all input parameters, namely stirring speed, volume and temperature, have significant influence on the process parameter k_{La}^* value as the error bars do not cut the x-axis. Additionally, the interaction of stirrer speed and volume is significant. A contour plot of the k_{La}^* value (data not shown) illustrates the effect of temperature, volume and stirrer speed on k_{La}^* , i.e., the higher the temperature, the lower the volume and the higher the stirrer speed, the higher the oxygen transfer over the headspace expressed as k_{La}^* .

Example 11.1.5 – Results of Dedicated Run at Manufacturing Scale

The k_{La}^* value was determined in a dedicated experiment and compared with the prediction of the model calculated by MODDE 8.02. The results of the prediction and the dedicated run are shown in **Table 44**. According to these results, the mathematical model was able to predict the k_{La}^* value in an adequate manner.

Batch [-]	Volume [L]	Stirrer speed [rpm]	Temperature [°C]	k_{La}^* [1/h]
Model prediction	606	200	38	0.32
Result of the dedicated run	606	200	38	0.25

Table 44: k_{La}^* values (predicted and measured) of standard process conditions at manufacturing-scale (SITE A).

Example 11.1.6 – Predicted k_{La}^* for the Process conditions at SITE A

The k_{La}^* for the process conditions used during campaign AT493021 at SITE A were predicted with the model calculated by MODDE 8.02. The results are shown in **Table 45** with predicted k_{La}^* values of 0.05 h^{-1} to 0.69 h^{-1} when constant stirring is applied. The values indicate that the process step is robust over a broad range of k_{La}^* values resulting from different process volumes. The parameters having an influence on k_{La}^* (and hence the oxygen transfer) should be

considered for final assessment of process robustness, e.g., overall stirring time has a significant influence on the oxygen transfer.

Batch [-]	Cycle [-]	Volume [L]	Stirrer speed [rpm]	Temperature [°C]	Predicted k_{La}^* [h ⁻¹]
B010231	1	443	200	37	0.69
B010231	2	466	200	37	0.60
B010655	1	542	200	37	0.41
B010655	2	569	200	37	0.37
B010655	3	557	200	37	0.39
B012307	1	1348 ¹⁾	200	37	0.05
B013981	1	1142 ¹⁾	200	37	0.08

Table 45: k_{La}^* values of process conditions at manufacturing-scale (SITE A). 1) Volume exceeds the evaluated range. Extrapolation can lead to imprecise values.

Example 12: Qualification Runs and Comparison of Scale-Down Model to Manufacturing-Scale at Site A

Example 12.1 - Methods

Three scale-down model qualification runs (REACT065, REACT066, and REACT067) performed under standard conditions were compared with seven representative runs at manufacturing-scale. For the scale-down model qualification runs, secukinumab, INAKT.F (stored below -60°C) originating from manufacturing-scale run B008530 (campaign AT493021, SITE A), was used. The load (INAKT.F) was thawed in a hand warm water bath (15-35°C) before use. The steps of the full selective reduction procedure are displayed in **Table 46**.

Step	Buffer	Target pH	Comment
Concentration adjustment	WFI	NA	Adjust protein concentration to 12.0 – 15.0 (13.5) mg/mL*
Check and adjust of dissolved oxygen (DO) level	NA	NA	DO > 80%, start pH-adjustment 1
pH-adjustment 1	AIN457-TITR1 (1 M Tris)	7.9 – 8.1	Volume increase of approx. 5%
Addition of cysteine containing solution with stirring	AIN457-TITR3 (120 mM Cysteine HCl, 20 mM Na-EDTA, pH 8.0±0.2)	NA	DO > 80%, start cysteine treatment; addition of cysteine solution **under stirring

Heating time to reaction temperature with stirring	NA	NA	Heating from RT (18-25°C) to 37°C under stirring in max. 1 h (75 minutes). Stirrer speed
Incubation	NA	NA	Incubation time: 250 min with 2 min. stirring per hour
Cooling time after reaction to room temperature with stirring	NA	NA	Cooling from 37°C to RT (18-25°C) under stirring in max. 1h (75 minutes)
pH-adjustment 2	AIN457-TITR2 (0.3 M H ₃ PO ₄)	5.1 – 5.3	Volume increase of approx. 15%

Table 46: Description of selective reduction procedure. *calculation for addition of WFI: Vol (WFI) = (Vol

(INAKT.F)*c (INAKT.F) / 13.5 mg/mL) – Vol(INAKT.F); **calculation for addition of TITR3:

Vol(TITR3)=(Vol(INAKT.F)+Vol(WFI)+Vol(TITR1)) * factor (TITR3), wherein factor (TITR3) = 0.05 / 0.95 = 0.05263 (6 mM Cysteine in reactivation solution).

The process parameters listed in **Table 47** were applied for scale-down model qualification runs. The parameters correspond to the process conditions at manufacturing-scale (SITE A) and were scaled down appropriately.

Process parameter	Unit	Target	Operating Range
Load pH	-	8.0	7.9 – 8.1
Load temperature	°C	as is	18 – 25
Protein concentration before pH adjustment and TITR3 addition	mg/mL	13.5	12 - 15
Incubation temperature	°C	37	35 – 39
Temperature after cooling	°C	22	18 – 25
Stirrer speed during heating and cooling	rpm	50	40 – 60
Stirrer speed during incubation ¹	rpm	0	N/A
Heating time	min	60	≤ 60
Incubation time	min	250	240 - 300
Cooling time	min	60	≤ 60
pH of REACT.P	-	5.2	5.1 – 5.3
Dissolved oxygen	%	as is	> 80
Dilution factor with TITR3	-	1:20 (6 mM cysteine)	1:20 (6 mM cysteine)

Table 47 – Process parameters of the selective reduction step. ¹ stirring for 2 minutes each hour at 50 rpm.

Samples were withdrawn, pH was adjusted with 0.3 M phosphoric acid, pH 1.4 (AIN457-

TITR2) to 5.2 (acceptable range 5.0-5.4) within 5 minutes, and then the samples were transferred to the analytical lab. All retained samples were frozen and stored at $\leq -60^{\circ}\text{C}$. The qualification of the scale-down model was assessed by comparison of the product quality attributes activity by CEX and purity by CE-SDS, as previously described. The following time intervals were tested: 0, 10, 20, 30, 45, 60, 120, 240 min.

Example 12.2 - Results

Dissolved oxygen charts from three representative manufacturing-scale runs (data not shown) and the scale-down model qualification runs (**Figure 16**) show the characteristic shape of the selective reduction step. The curves show that the dissolved oxygen is removed from the system by oxidation of cysteine. According to development experience, extended stirring results in higher oxygen transfer and leads to insufficient activity by CEX. Hence, the selective reduction solution is only stirred for 2 - 15 minutes per hour during incubation, thereby limiting the oxygen transfer into the solution and keeping the dissolved oxygen level low until the selective reduction is stopped. During the cooling phase, the oxygen level increases slightly due to the oxygen transfer into the solution by continuous mixing, while a certain amount of cysteine is oxidized, rendering it unavailable as reducing agent for the dissolved oxygen.

For the manufacturing-scale runs 3-6 of campaign AT493021 at SITE A, the activity by CEX was on average 96.3 % with a standard deviation of 0.8 %, while the purity by CE-SDS was on average of 92 % with a standard deviation of 1 %. The results of each run are displayed in **Table 48**, along with the predicted k_{La}^* values derived from **Table 45**.

Run	Activity by CEX [%]	Purity by CE-SDS [%]	Predicted k_{La}^* [1/h]
B010231; Cycle#1	94.5	92	0.69
B010231; Cycle#2	96.7	93	0.60
B010655; Cycle#1	96.3	90	0.41
B010655; Cycle#2	96.4	90	0.37
B010655; Cycle#3	96.3	93	0.39
B012307; Cycle#1	96.8	92	0.05
B013981; Cycle#1	96.9	93	0.08
Standard deviation	0.8	1	
Average $\pm 3 \times \text{Stdev}$	96.3 ± 2.5	92 ± 4	

Table 48: Experimental results of the manufacturing-scale runs.

For the scale-down model qualification runs, the activity by CEX was on average value

95.0 % with a standard deviation of 1.6 %, while the purity by CE-SDS was on average 94 % with a standard deviation of 1 % (**Table 49**).

Run	Activity by CEX [%]	Purity by CE-SDS [%]
Qualification run#1; REACT065	95.9	93.0
Qualification run#2; REACT066	96.0	94.0
Qualification run#3; REACT067	93.1	95.0
Standard deviation	1.6	1.0
Average $\pm 3 \times \text{Stdev}$	95.0 $\pm$ 4.9	94 $\pm$ 3

Table 49: Experimental results of the scale-down model qualification runs.

In addition to the assessment of product quality at the end of the selective reduction step, the kinetic for activity by CEX was also monitored and the results of the scale-down model runs (average of REACT065, 66 and 67) and manufacturing-scale at SITE A were compared. The results are shown in **Figure 17**. The data obtained from the scale-down model are within the variation of the manufacturing scale, demonstrating that the kinetics at large-scale (Site A) and small-scale are comparable.

Example 13 – Combined analysis of k_{La}^* values from Examples 9-12

Due to the importance of the level of dissolved oxygen in the solution during the selective reduction reaction, controlled oxygen transfer from the head-space of a vessel is important for the selective reduction step. For each system, these variables can be captured as k_{La}^* values. The k_{La}^* is dependent on the stirrer speed, stirrer type, filling volume, temperature and surface area of the solution in contact with the headspace influenced by the individual geometry of each vessel. While k_{La}^* will change depending on each setup, from the above experiments, it can be seen that a range of k_{La}^* values are acceptable to ensure the quality and activity of secukinumab during the selective reduction process.

A predicted k_{La}^* range of $0.18 \text{ h}^{-1} - 0.37 \text{ h}^{-1}$ for the experimental conditions of the scale down studies was determined by use of the statistical model (**Table 41**). The k_{La}^* value decreased as the stir speed decreased, given a set volume. Because the response surface design experimental conditions and set up in **Example 9** are the same as those used with the scale-down model in **Example 10**, the predicted k_{La}^* values of **Table 41** can be correlated to the conditions and results in **Table 28**. This analysis is presented in **Table 50**.

56418 FF

Run	Protein (mg/ml)	Cysteine (mM)	Approx. molar ratio cysteine to protein (M/M)	kLa* (h ⁻¹)	Spin (RPM)	Activity by CEX (%) REACT.PT300 (240 min inc.)	Purity by CE-SDS (%) REACT.PT300 (240 min inc.)	Activity by CEX (%) REACT.P (300 min inc.)	Purity by CE-SDS (%) REACT.P (300 min inc.)
REACT082	15.4	4.8	46.11	0.18	0	96.1	95	96.2	96
REACT070	12.7	6.0	69.89	0.18	0	96.1	94	96.5	92
REACT090	10.0	4.8	71.01	0.18	0	96.6	94	97.3	95
REACT077	15.4	8.0	76.85	0.18	0	97.0	92	97.4	92
REACT080	10.0	8.0	118.36	0.18	0	97.8	90	97.5	91
REACT072	12.7	4.8	55.92	0.27	50	94.5	96	90.2	95
REACT079	15.4	6.0	57.64	0.27	50	96.0	95	94.4	95
REACT069	12.7	6.0	69.89	0.27	50	95.3	95	94.1	95
REACT073	12.7	6.0	69.89	0.27	50	96.1	95	94.7	95
REACT088	12.7	6.0	69.89	0.27	50	95.7	95	95.1	95
REACT086	10.0	6.0	88.77	0.27	50	96.7	94	95.4	94
REACT087	12.7	8.0	93.19	0.27	50	96.5	94	96.0	94
REACT085	15.4	4.8	46.11	0.37	100	89.4	95	84.0	94
REACT075	12.7	6.0	69.89	0.37	100	95.7	95	88.9	95
REACT084	10.0	4.8	71.01	0.37	100	94.3	95	88.9	95
REACT089	15.4	8.0	76.85	0.37	100	95.2	95	93.7	94
REACT091	10.0	8.0	118.36	0.37	100	92.4	93	91.3	94

Table 50 – kLa* analysis for output parameter values REACT.PT300 and REACT.P. Molar ratios are rounded to the nearest whole number. The relative

molecular mass of secukinumab, based on amino acid sequence, is 147,944 Daltons, which is used to calculate the molar ratio of cysteine to protein in column 4.

For experiments having a 240 minute incubation period (REACT.PT300) all reactions, except REACT085, resulted in very high selective reduction performance with adequate product quality as measured by CEX and CE-SDS (**Table 50**). REACT085 had a stir speed of 100 rpm, a k_{La}^* value of 0.37 h^{-1} during incubation, and a very low molar ratio of cysteine to antibody (about 46:1). The low molar ratio of cysteine to antibody, combined with high oxygen transfer into the vessel (represented by the high k_{La}^* value), likely resulted in insufficient reactivation.

For experiments having a 300 minute incubation period (REACT.P) all reactions, except REACT085, REACT075, and REACT084 resulted in very high selective reduction performance with adequate product quality as measured by CEX and CE-SDS (**Table 50**). REACT085, REACT075, and REACT084 had a stir speed of 100 rpm, a k_{La}^* value of 0.37 h^{-1} during incubation, and a low-medium molar ratio of cysteine to antibody (about 46:1 and about 71:1). The low-medium molar ratio of cysteine to antibody, combined with high oxygen transfer into the vessel (represented by the high k_{La}^* value), likely resulted in insufficient reactivation. The increased incubation time (relative to REACT.PT300 samples) appeared to contribute to poorer CEX values in some REACT.P runs.

As can be seen from **Table 50** reactions with CEX activity greater than 90% have the following characteristics regarding the k_{La}^* in the system during the incubation portion of the selective reduction reaction:

- 1) if the k_{La}^* in the system during the incubation step of the selective reduction reaction is $< 0.37 \text{ h}^{-1}$, then the molar ratio of cysteine:antibody can vary between about 46:1 to about 118:1 (for both shorter and longer incubation times, e.g., about 210 to about 330 minutes, e.g., 240-300 minutes); and
- 2) the k_{La}^* in the system during the incubation step of the selective reduction reaction can be as high as 0.37 h^{-1} if the molar ratio of cysteine:protein is between about 56:1 to about 118:1 (for shorter incubation times, e.g., up to about 240 minute incubation) or between about 77:1 to about 118:1 (for longer incubation times, e.g., up to about 300 minute incubation).

Of course, the k_{La}^* during the heating and cooling phase can be much higher than the k_{La}^* during the incubation phase, e.g., **Table 54** shows a k_{La}^* as high as 0.69 h^{-1} can be applied

during the heating/cooling phase and still produce high quality product.

Example 9 (process characterization studies) examined the output parameters activity by CEX and purity by CE-SDS using a scale-down model that applied constant stirring during the incubation phase. **Example 10** examined k_{La}^* values of the scale-down model that also applied constant stirring. **Example 11** examined the k_{La}^* values when using continuous stirring at manufacturing-scale. During actual manufacturing, continuous stirring is only applied during the heating and cooling phases of the selective reduction reaction, while intermittent stirring (e.g., 2 – 15 minutes per hour) is applied during the incubation phases of the selective reduction reaction. Therefore, during manufacturing, the k_{La}^* of the system during the incubation phases of the selective reduction reaction step will be much lower than the k_{La}^* values observed in **Examples 10 and 11**. Nevertheless, from the combined results of **Examples 9-11**, it can be seen that the entire selective reduction step (i.e., heating phase, incubation phase, and cooling phase) can be generally performed using a k_{La}^* of $< 0.37 \text{ h}^{-1}$, including about 240 to about 300 minute incubation phase, if the molar ratio of cysteine:antibody is between about 46:1 to about 118:1.

The Worst/Best case experiments of **Example 9** were performed using the same qualified scale-down model, using a molar ratio of cysteine: antibody of about 66:1 with negligible stirring (i.e., 2 minutes per hour) during the incubation phase (k_{La}^* of $< 0.18 \text{ h}^{-1}$). Since reactions incubated for 210 to 330 minutes in the Worst/Best case experiments of **Example 9** all produced high quality product, it is expected that the incubation time can be expanded to about 210- about 330 minutes if the k_{La}^* is low and the molar ratio of cysteine:antibody is medium (e.g., about 66:1).

Example 14– Preferred Selective Reduction Process Parameters and Procedures

The preferred process parameters of the selective reduction step are listed in **Table 51**. The molar ratio of cysteine:antibody for the set point is about 66:1, with a suggested operating range of about 54:1 to about 83:1, and a proven acceptable range of about 46:1 to about 118:1.

Parameter	Set-point / Target	Suggested Operating Range	Proven Acceptable Range (PAR)
Incubation temperature [°C]	37	35 – 39	32 – 42
Protein concentration before pH adjustment and addition of TITR3 buffer [g/L]	13.5	12 – 15	10.0 – 15.4
Process pH [-]	8	7.9 – 8.1	7.8 – 8.2

Dissolved oxygen (DO) at reactivation start [%]	as is	≥ 80	≥ 60
Dilution factor by AIN457-TITR3 addition	1:20 (6.0 mM cysteine, 1.0 mM EDTA)	1:18 – 1:22 (5.5 – 6.7 mM cysteine, 0.9 – 1.1 mM EDTA)	1:15 – 1:25 (4.8 – 8.0 mM cysteine, 1.3 – 0.8 mM EDTA)
Approx. molar ratio cysteine:antibody [M/M]	65.75:1 (about 66:1)	54.24:1 – 82.60:1 (about 54:1 to about 83:1)	46.11:1 – 118.36:1 (about 46:1 to about 118:1)
Incubation time for reactivation including stirring [min]	250	240 – 300	210 – 330
pH adjustment after temperature decrease [-]	5.2	5.1 – 5.3	
Heating time including stirring [min]	60	≤ 75	≤ 90
Stirring time during incubation [min / hour]	2	≤ 5	≤ 15
Stir speed during incubation [rpm]	75	65-85	65-85
Cooling time including stirring [min]	60	≤ 75	≤ 90

Table LLLLL: Selective reduction – list of process parameters.

The individual steps of the selective reduction procedure are displayed in **Table 52**. First, the protein concentration of the starting material is adjusted by addition of WFI to a target concentration of 13.5 mg/mL. Next, the pH is adjusted to 8.0 by addition of 1 M Tris. The dO₂ is checked and adjusted (e.g., to $\geq 60\%$), e.g., by further mixing, submerge aeration, etc., if the operating range is not met. The reaction is started by adding cysteine to 6 mM and thereafter the solution is heated (over about an hour) to incubation temperature (about 37°C). The solution is incubated for about 250 minutes with about 2 minutes stirring per hour to limit oxygen transfer from headspace into solution during incubation time. Finally, the solution is cooled to room temperature (over about an hour) and the pH is adjusted to about 5.2, e.g., with 0.3 M ortho-phosphoric acid to stop the reaction.

Step	Buffer	Target pH	Comment
Concentration adjustment	WFI	NA	Adjust protein concentration to 12.0 – 15.0 (13.5) mg/mL*
Check and adjust of dissolved oxygen (DO) level	NA	NA	DO > 80%, start pH-adjustment 1
pH-adjustment 1	AIN457-TITR1 (1 M Tris)	7.9 – 8.1	Volume increase of approx. 5%
Addition of cysteine	AIN457-TITR3	NA	DO > 80%, start cysteine

containing solution with stirring	(120 mM Cysteine HCl, 20 mM Na-EDTA, pH 8.0±0.2)		treatment; addition of cysteine solution **under stirring
Heating time to reaction temperature with stirring	NA	NA	Heating from RT (18-25°C) to 37°C under stirring in max. 1 h (75 minutes). Stirrer speed
Incubation	NA	NA	Incubation time: 250 min with 2 min. stirring per hour
Cooling time after reaction to room temperature with stirring	NA	NA	Cooling from 37°C to RT (18-25°C) under stirring in max. 1h (75 minutes)
pH-adjustment 2	AIN457-TITR2 (0.3 M H ₃ PO ₄)	5.1 – 5.3	Volume increase of approx.15%

Table 52: Description of selective reduction procedure. *calculation for addition of WFI: Vol (WFI) = (Vol

(INAKT.F)*c (INAKT.F) / 13.5 mg/mL) – Vol(INAKT.F); **calculation for addition of TITR3:

Vol(TITR3)=(Vol(INAKT.F)+Vol(WFI)+Vol(TITR1)) * factor (TITR3), wherein factor (TITR3) = 0.05 / 0.95 = 0.05263 (6 mM Cysteine in reactivation solution).

WHAT IS CLAIMED IS:

1. A method for selectively reducing CysL97 in a preparation of IL-17 antibodies that have been recombinantly produced by mammalian cells, comprising:
 - c) contacting the preparation with at least one reducing agent in a system to form a reducing mixture; and
 - d) incubating the reducing mixture while maintaining a volumetric oxygen mass-transfer coefficient (k_{La}^*) in the system of $\leq$ about 0.37 h^{-1} , said k_{La}^* being calculated by adapting a dissolved oxygen curve to a saturation curve;wherein the IL-17 antibodies each comprise an immunoglobulin heavy chain variable domain (V_H) comprising the three complementarity determining regions (CDRs) of the V_H set forth as SEQ ID NO:8 and an immunoglobulin light chain variable domain (V_L) comprising the three CDRs of the V_L set forth as SEQ ID NO:10; and wherein prior to step a) the initial percent oxygen saturation in the preparation is at least about 60%, as measured using an oxygen probe calibrated at 25°C.
2. The method according to claim 1, wherein the at least one reducing agent is selected from the group consisting of cysteamine, cysteine, and combinations thereof.
3. The method according to claim 1, wherein the at least one reducing agent is a thiol-containing reducing agent having a standard oxidation-reduction potential, E° , of about -0.20 V to about -0.23 V at pH 7.0, as measured by thiol-disulfide exchange.
4. The method according to claim 2, wherein the at least one reducing agent is cysteine.
5. The method according to claim 4, wherein the k_{La}^* in the system during step b) is $\leq$ about 0.37 h^{-1} , wherein the molar ratio of cysteine:IL-17 antibodies in the reducing mixture is between about 56:1 to about 118:1, and wherein the reducing mixture is incubated according to step b) for up to about 240 minutes.
6. The method according to claim 4, wherein the k_{La}^* in the system during step b) is $\leq$

about 0.37 h^{-1} , wherein the molar ratio of cysteine:IL-17 antibodies in the reducing mixture is between about 77:1 to about 118:1, and wherein the reducing mixture is incubated according to step b) for up to about 300 minutes.

7. The method according to claim 4, wherein the k_{LA}^* in the system during step b) is $<$ about 0.37 h^{-1} , and wherein the molar ratio of cysteine:IL-17 antibodies in the reducing mixture is between about 46:1 to about 118:1.

8. The method according to claim 4, wherein the k_{LA}^* in the system during step b) is $<$ about 0.37 h^{-1} , and wherein the molar ratio of cysteine:IL-17 antibodies in the reducing mixture is between about 54:1 to about 82:1.

9. The method according to claim 4, wherein the molar ratio of cysteine:IL-17 antibodies or in the reducing mixture is about 66:1.

10. The method according to any one of the preceding claims, wherein the k_{LA}^* in the system during step b) is $\leq$ about 0.27 h^{-1} , preferably $\leq$ about 0.18 h^{-1} .

11. The method according to any one of the preceding claims, wherein prior to step a) the preparation comprises about 4 mg/ml to about 19.4 mg/ml, e.g., about 10 mg/ml to about 19.4 mg/ml, e.g., about 10 mg/ml to about 15.4 mg/ml, e.g., about 12 mg/ml to about 15 mg/ml, e.g., about 13.5 mg/ml of the IL-17 antibodies.

12. The method according to claim 4, wherein the concentration of cysteine in the reducing mixture is about 4.0 mM to about 8.0 mM, e.g., about 4.8 mM to about 8.0 mM, e.g., about 5.5 mM to about 6.7 mM, e.g., about 6.0 mM.

13. The method according to any one of the preceding claims, wherein prior to step a) the pH of the preparation is about 7.3 to about 8.5, e.g., about 7.8 to about 8.2, e.g., about 7.9 to about 8.1, e.g., about 8.0.

14. The method according to any one of the preceding claims, wherein prior to step a) the initial percent oxygen saturation in the preparation is at least about 80% or at least about 60%, as measured using an oxygen probe calibrated at 25°C.
15. The method according to any one of the preceding claims, wherein the reducing mixture further comprises EDTA, and wherein the concentration of EDTA in the reducing mixture is about 1.3 mM to about 0.8 mM, e.g., about 1.1 to about 0.9 mM, e.g., about 1.0 mM.
16. The method according to any one of the preceding claims, wherein between step a) and step b), the reducing mixture is heated to a temperature between about 32°C to about 42°C, e.g., to between about 35°C to about 39°C, e.g., to about 37°C.
17. The method according to claim 16, wherein the reducing mixture is heated for about 45 to about 90 minutes, e.g., about 45 to about 75 minutes, e.g., about 60 minutes.
18. The method according to claim 16, wherein the k_{La}^* in the system during heating is $\leq$ about 0.69 h^{-1} , said k_{La}^* being calculated by adapting a saturation curve to a dissolved oxygen curve.
19. The method according to any one of the preceding claims, wherein the reducing mixture is incubated according to step b) at a temperature between about 20°C to about 42°C, e.g., about 32°C to about 42°C, e.g., about 35°C to about 39°C, e.g., about 37°C.
20. The method according to claim 7, wherein the reducing mixture is incubated according to step b) for about 210 to about 420 minutes, e.g., about 210 to about 330 minutes, e.g., about 240 to about 300 minutes, e.g., about 250 minutes.
21. The method according to any one of the preceding claims, wherein during step b) the reducing mixture is stirred for $\leq$ about 15 minutes per hour, e.g., $\leq$ about 5 minutes per hour, e.g., $\leq$ about 2 minutes per hour.

22. The method according to any one of the preceding claims, wherein the level of intact IL-17 antibodies in the reducing mixture following step a) decreases to at least about 80% as measured by sodium dodecyl sulfate capillary electrophoresis (CE-SDS).
23. The method according to any one of the preceding claims, further comprising:
c) cooling the mixture resultant from step b) to room temperature, e.g., between about 16 °C to about 28 °C.
24. The method according to claim 23, wherein the mixture is cooled according to step c) for about 45 to about 90 minutes, e.g., about 45 to about 75 minutes, e.g., about 60 minutes.
25. The method according to claim 23, wherein the k_{La}^* in the system during step c) is $\leq 0.69 \text{ h}^{-1}$, said k_{La}^* being calculated by adapting a dissolved oxygen curve to a saturation curve.
26. The method according to claim 23, further comprising:
d) adjusting the pH of the mixture resultant from step c) to between about 5.1 to about 5.3, e.g., about 5.2.
27. The method according to claim 26, wherein adjusting step d) comprises adding o-phosphoric acid to the mixture resultant from step c).
28. The method according to claim 26, wherein the level of intact IL-17 antibodies in the mixture resultant from step d) is at least about 90%, as measured by CE-SDS.
29. The method according to claim 26, wherein the level of activity of the IL-17 antibodies in the mixture resultant from step d) is at least about 90%, as measured by cystamine-cation exchange chromatography (cystamine-CEX).
30. The method according to any one of the above claims, wherein an oxidized form of the reducing reagent, e.g., cystine or cystamine, is not added to the reducing mixture.

31. The method according to any one of the above claims, wherein a denaturant, e.g., guanidine hydrochloride, is not added to the reducing mixture.
32. A method for selectively reducing CysL97 in a preparation of IL-17 antibodies that have been recombinantly produced by mammalian cells, comprising:
- a) contacting the preparation with a set of oxidation/reduction reagents selected from cysteine/cystine and cysteine/cystamine to form a reducing mixture; and
 - b) incubating the reducing mixture at a temperature of about 37 °C under anaerobic conditions for at least about 4 hours, or incubating the reducing mixture at a temperature of about 18-24 °C for about 16-24 hours;
- wherein the IL-17 antibodies each comprise an immunoglobulin heavy chain variable domain (V_H) comprising the three complementarity determining regions (CDRs) of the V_H set forth as SEQ ID NO:8 and an immunoglobulin light chain variable domain (V_L) comprising the three CDRs of the V_L set forth as SEQ ID NO:10.
33. The method according to claim 32, wherein the molar ratio of oxidation/reduction reagents in the reducing mixture is between about 4:1 to about 80:1.
34. The method according to claim 33, wherein the molar ratio of oxidation/reduction reagents in the reducing mixture is between about 26:1 to about 80:1.
35. The method according to claim 33, wherein the set of oxidation/reduction reagents is cysteine/cystine.
36. The method according to claim 33, wherein the set of oxidation/reduction reagents is cysteine/cystamine.
37. The method according to claim 32, wherein the molar ratio of cysteine:IL-17 antibodies in the reducing mixture is between about 21:1 to about 296:1 .
38. The method according to any one of the above claims, wherein the three CDRs of the V_H

set forth as SEQ ID NO:8 are, in sequence, the amino acids sequences set forth as SEQ ID NOs: 1, 2, and 3 or the amino acids sequences set forth as SEQ ID NOs: 11, 12, and 13 and wherein the three CDRs of the V_L set forth as SEQ ID NO:10 are, in sequence, the amino acids sequences set forth as SEQ ID NOs: 4, 5 and 6.

39. The method according to any one of the above claims, wherein the IL-17 antibodies each comprise:

- i) the V_L set forth as SEQ ID NO:10 and the V_H set forth as SEQ ID NO:8; or
- ii) the full length light chain set forth as SEQ ID NO:14 and the full length heavy chain set forth as SEQ ID NO:15, with or without the C-terminal lysine.

40. The method according to any one of the above claims, wherein the IL-17 antibodies are human antibodies of the IgG₁ isotype.

41. The method according to any one of the above claims, wherein the IL-17 antibodies are secukinumab antibodies.

42. The method according to any one of the above claims, wherein the level of activity of the IL-17 antibodies in the preparation increases by at least about 10 percentage points, at least about 15 percentage points, at least about 20 percentage points, at least about 25 percentage points, or at least about 30 percentage points within about 60 minutes following step b), as measured by cystamine-CEX.

43. A method for selectively reducing CysL97 in a preparation of secukinumab antibodies that have been recombinantly produced by mammalian cells, comprising:

- g) adjusting the concentration of secukinumab in the preparation to between about 4 mg/ml – about 19.4 mg/ml, e.g., about 10 mg/ml - about 19.4 mg/ml, e.g., about 10 - about 15.4, e.g., about 12 mg/ml - about 15 mg/ml, e.g., about 13.5 mg/ml;
- h) adjusting the percent oxygen saturation in the preparation to at least about 60%, e.g., at least about 80%;
- i) adjusting the pH of the preparation to about 7.4 - about 8.5, e.g., about

7.8 - about 8.2, e.g., about 7.9 - about 8.1, e.g., about 8.0;

j) contacting the preparation with cysteine in a vessel to form a reducing mixture, wherein the concentration of cysteine in the reducing mixture is about 4.0 mM - about 8.0 mM, e.g., about 4.8 mM - about 8.0 mM, e.g., about 5.5 mM - about 6.7 mM, e.g., about 6.0 mM;

k) heating the reducing mixture to a temperature between about 32°C - about 42°C, e.g., to between about 35°C - about 39°C, e.g., to about 37°C, said heating occurring for about 45 - about 90 minutes, e.g., about 45 - about 75 minutes, e.g., about 60 minutes;

l) incubating the reducing mixture from step e) at a temperature between about 20°C - about 42°C, e.g., 32°C - about 42°C, e.g., to between about 35°C - about 39°C, e.g., to about 37°C, said incubating occurring for about 210 - about 420 minutes, e.g., about 210 - about 330 minutes, e.g., about 240 - about 300 minutes, e.g., about 250 minutes while maintaining a volumetric oxygen mass-transfer coefficient (k_{La}^*) in the vessel of $\leq 0.37 \text{ h}^{-1}$, said k_{La}^* being calculated by adapting a saturation curve to a dissolved oxygen curve,

g) cooling the mixture resultant from step f) to a temperature between about 16°C - about 28°C, said cooling occurring for about 45 - about 90 minutes, e.g., about 45 - about 75 minutes, e.g., about 60 minutes; and

h) adjusting the pH of the mixture resultant from step g) to between about 5.1 - about 5.3, e.g., about 5.2.

44. A purified preparation of IL-17 antibodies made by any one of the preceding claims.

45. The purified preparation according to claim 44, wherein the IL-17 antibodies are secukinumab antibodies.

46. A purified preparation of secukinumab, wherein the level of intact secukinumab in the preparation is at least about 90%, as measured by sodium dodecyl sulfate capillary electrophoresis (CE-SDS), and wherein the level of activity of secukinumab in the preparation is at least about 90%, as measured by cystamine-CEX.