

DESCRIPTION

Polypeptide having D-lactate Dehydrogenase Activity, Polynucleotide Encoding the
Polypeptide, and Process for Production of D-lactic acid

TECHNICAL FIELD

5 [0001]

The present invention relates to a novel polypeptide for production of D-lactic acid, which polypeptide has a D-lactate dehydrogenase activity; a polynucleotide encoding the polypeptide; and a transformant into which the polynucleotide is introduced. The present invention also relates to a method of producing D-lactic
10 acid which comprises culturing the transformant.

BACKGROUND ART

[0002]

From the idea of carbon neutrality, strong attention has been given to polylactic acids (PLAs) which utilize a renewable resource, biomass, as starting
15 material. PLAs are relatively low cost and heat resistant having a melting point of about 170°C; therefore, they are expected as a biodegradable polymer which can be melt-molded. In addition, it is known that a stereocomplex polylactic acids can be obtained by mixing poly-L-lactic acid and poly-D-lactic acid (Patent Documents 1 to
3). Stereocomplex polylactic acids are known to exhibit a higher melting point and
20 higher crystallizability as compared to single polymers and provide useful molded articles as fibers, films and resin molded articles. For both of the starting materials thereof, L-lactic acid and D-lactic acid, there is a demand for a method of producing them with high purity and high efficiency.

[0003]

25 In nature, there exist bacteria which efficiently produce lactic acids, such as lactic acid bacteria, and some of those lactic acid production methods utilizing such bacteria have already been put to practical use. Examples of bacteria which

efficiently produce L-lactic acid include *Lactobacillus delbrueckii*. In addition, as bacteria which efficiently produce D-lactic acid, microbes such as *Sporolactobacillus laevolacticus* are known (Patent Document 4). In any of these cases, the amount of the accumulated lactic acid reached a high level in anaerobic culture; however, since

5 D-lactic acid and L-lactic acid are produced as by-product in L-lactic acid fermentation and D-lactic acid fermentation, respectively, the optical purity is lowered. In addition, it is extremely difficult to separate these lactic acids.

[0004]

Therefore, as a production method of lactic acid having a high optical purity,

10 it has been examined to introduce a gene encoding L-lactate dehydrogenase or D-lactate dehydrogenase into a yeast which does not intrinsically have lactic acid-producing ability and subject the yeast to L-lactic acid and D-lactic acid fermentation (Patent Documents 4 to 6 and Non-patent Document 1). With regard to L-lactic acid fermentation by genetically engineered yeast, by introducing a highly active gene

15 originated from *Xenopus laevis* which encodes the L-lactate dehydrogenase, lactic acid fermentation can be efficiently performed with high optical purity (Patent Document 4). On the other hand, with regard to D-lactic acid fermentation by genetically engineered yeast, although D-lactic acid having high optical purity can be obtained in the same manner as in the case of L-lactic acid fermentation, the yield

20 thereof is a problem (Patent Documents 5 and 6 and Non-patent Document 1).

PRIOR ART DOCUMENTS

PATENT DOCUMENTS

[0005]

Patent Document 1: JP S61-36321A

25 Patent Document 2: JP S63-241024A

Patent Document 3: JP 2000-17163A

Patent Document 4: WO2007/043253

Patent Document 5: JP 2007-074939A

Patent Document 6: WO2004/104202

NON-PATENT DOCUMENTS

[0006]

- 5 Non-patent Document 1: Ishida N, et al., Journal of Bioscience and Bioengineering (2006), 101, 172-7

SUMMARY OF THE INVENTION

PROBLEMS TO BE SOLVED BY THE INVENTION

[0007]

- 10 An object of the present invention is to realize highly productive D-lactic acid fermentation using a polypeptide having a D-lactate dehydrogenase activity higher than that of conventional polypeptides and a polynucleotide encoding the polypeptide.

MEANS FOR SOLVING THE PROBLEMS

[0008]

- 15 The present inventors intensively studied a variety of D-lactate dehydrogenases to discover a polypeptide which increases the production of D-lactic acid and also has a D-lactate dehydrogenase activity to yield only a small amount of by-product; and a polynucleotide encoding the polypeptide, thereby completing the present invention.

20 [0009]

That is, the present invention is constituted by the following modes.

[0010]

(1) A polypeptide which is any one of the following (A) to (C):

(A) a polypeptide having the amino acid sequence shown in SEQ ID NO:1 or 2;

- 25 (B) a polypeptide having the same amino acid sequence as shown in SEQ ID NO:1 or 2 except that one or several amino acids are substituted, deleted, inserted and/or added, which polypeptide has a D-lactate dehydrogenase activity; and

(C) a polypeptide having an amino acid sequence which has a sequence identity of not less than 80% to the amino acid sequence shown in SEQ ID NO:1 or 2, which polypeptide has a D-lactate dehydrogenase activity.

[0011]

5 (2) A polynucleotide which is any one of the following (a) to (e):

(a) a polynucleotide having the nucleotide sequence shown in SEQ ID NO:3 or 4;

(b) a polynucleotide having the same nucleotide sequence as shown in SEQ ID NO:3 or 4 except that one or several nucleotides are substituted, deleted, inserted and/or added, which polynucleotide encodes a polypeptide having a D-lactate

10 dehydrogenase activity;

(c) a polynucleotide which hybridizes under stringent conditions with the entirety or a part of the polynucleotide having the nucleotide sequence shown in SEQ ID NO:3 or 4 or a complementary strand thereof, which polynucleotide encodes a polypeptide having a D-LDH activity;

15 (d) a polynucleotide having a nucleotide sequence which has a sequence identity of not less than 80% to the nucleotide sequence shown in SEQ ID NO:3 or 4, which polynucleotide encodes a polypeptide having a D-lactate dehydrogenase activity; and

(e) a polynucleotide encoding the polypeptide according to (1).

[0012]

20 (3) A DNA construct in which the polynucleotide according to (2) and a promoter capable of expressing the polynucleotide are linked.

[0013]

(4) The DNA construct according to (3), characterized in that the above-described promoter is a promoter of pyruvate decarboxylase 1 gene (PDC1 gene),
25 suppression-of-exponential-defect 1 gene (SED1 gene) or glyceraldehyde-3-phosphate dehydrogenase 3 gene (TDH3 gene).

[0014]

(5) The DNA construct according to (3) or (4), wherein the above-described promoter is selected from any one of the following (I) to (III):

(I) a promoter having the nucleotide sequence shown in any one of SEQ ID NOs:5 to 7;

5 (II) a promoter having a nucleotide sequence which hybridizes under stringent conditions with the nucleotide sequence shown in any one of SEQ ID NOs:5 to 7 or a nucleotide sequence comprising a part thereof; and

(III) a promoter having the same nucleotide sequence as shown in any one of SEQ ID NOs:5 to 7 except that one or several nucleotides are deleted, substituted and/or
10 added.

[0015]

(6) A transformant in which the polynucleotide according to (2) or the DNA construct according to any one of (3) to (5) is introduced.

[0016]

15 (7) A transformed yeast in which the polynucleotide according to (2) or the DNA construct according to any one of (3) to (5) is introduced.

[0017]

(8) The transformed yeast according to (7), wherein at least one of pyruvate decarboxylase 1 gene (PDC1 gene), suppression-of-exponential-defect 1 gene (SED1
20 gene) and glyceraldehyde-3-phosphate dehydrogenase 3 gene (TDH3 gene) of the above-described transformed yeast is substituted with the polynucleotide according to (2) or the DNA construct according to any one of (3) to (5).

[0018]

(9) The transformed yeast according to (8), wherein at least one of the above-
25 described genes is PDC1 gene.

[0019]

(10) A method of producing D-lactic acid, which comprises the step of

culturing the transformant according to (6) or the transformed yeast according to any one of (7) to (9).

[0020]

(11) A transformant in which a polynucleotide encoding D-lactate
5 dehydrogenase originated from the family Limulidae or DNA construct in which the polynucleotide and a promoter capable of expressing the polynucleotide are linked is introduced.

[0021]

(12) A transformed yeast in which a polynucleotide encoding D-lactate
10 dehydrogenase originated from the genus *Limulus* or DNA construct in which the polynucleotide and a promoter capable of expressing the polynucleotide are linked is introduced.

[0022]

(13) The transformed yeast according to (12), wherein at least one of pyruvate
15 decarboxylase 1 gene (PDC1 gene), suppression-of-exponential-defect 1 gene (SED1 gene) and glyceraldehyde-3-phosphate dehydrogenase 3 gene (TDH3 gene) of the above-described transformed yeast is substituted with the above-described polynucleotide encoding D-lactate dehydrogenase originated from the family Limulidae or DNA construct in which the polynucleotide and a promoter capable of
20 expressing the polynucleotide are linked is introduced.

[0023]

(14) A method of producing D-lactic acid, which comprises the step of
culturing the transformant according to (11) or the transformed yeast according to
(12) or (13).

25 EFFECTS OF THE INVENTION

[0024]

According to the present invention, a polypeptide having a D-lactate

dehydrogenase activity suitable for fermentative production of D-lactic acid and a polynucleotide encoding the polypeptide are provided. In addition, by using the polynucleotide of the present invention, a transformant capable of highly producing D-lactic acid can be easily obtained, and in turn, D-lactic acid can be efficiently produced with high purity by culturing the transformant.

BRIEF DESCRIPTION OF THE DRAWINGS

[0025]

[Fig. 1] Fig. 1 shows the procedures for preparing one of the yeast SU042 strains according to the present invention.

[Fig. 2] Fig. 2 shows the results of membrane-integrated continuous culture of yeast SU042 strain according to the present invention.

MODE FOR CARRYING OUT THE INVENTION

[0026]

The modes for carrying out the present invention will now be explained in detail.

[0027]

The term "D-lactate dehydrogenase (hereinafter, also referred to as "D-LDH") activity" is an activity to convert reduced nicotinamide adenine dinucleotide (NADH) and pyruvic acid into D-lactic acid and oxidized nicotinamide adenine dinucleotide (NAD⁺).

[0028]

The present invention is characterized by including a polypeptide having the amino acid sequence shown in SEQ ID NO:1 or 2 or a homolog thereof, which polypeptide having a D-LDH activity.

[0029]

The polypeptide having the amino acid sequence shown in SEQ ID NO:1 or 2 is a polypeptide originated from *Limulus polyphemus*, which belongs to the family

Limulidae and genus *Limulus*.

[0030]

Examples of the homolog of the polypeptide having the amino acid sequence shown in SEQ ID NO:1 or 2 include a polypeptide having the same amino acid sequence as shown in SEQ ID NO:1 or 2 except that one or several, preferably 1 to 10, more preferably 1 to 5, still more preferably 1 or 2 amino acids are substituted, deleted, inserted and/or added, which polypeptide also has a D-LDH activity.

[0031]

Further, a homolog of the polypeptide shown in SEQ ID NO:3 or 4 may also be a polypeptide having an amino acid sequence which has a sequence identity of not less than 80%, preferably 90%, more preferably not less than 95% to the amino acid sequence shown in SEQ ID NO:1 or 2, which polypeptide also has a D-LDH activity. Here, the sequence identity of an amino acid sequence can be easily checked using the BLAST, which is a software widely used in the art. Anyone can use the BLAST through the NCBI (National Center for Biotechnology Information) homepage, and the identity can be easily checked by using default parameters.

[0032]

In addition, the homolog of the polypeptide having the amino acid sequence shown in SEQ ID NO:1 or 2 may also be a polypeptide originated from a living organism belonging to the family Limulidae, preferably a living organism belonging to the genus *Limulus* or *Tachypleus*, more preferably *Limulus polyphemus* belonging to the genus *Limulus*, or *Tachypleus tridentatus*, *Tachypleus gigas* or *Tachypleus rotundicauda* belonging to the genus *Tachypleus*, which polypeptide has a D-LDH activity.

[0033]

The polypeptide having the amino acid sequence shown in SEQ ID NO:1 or 2 can be extracted from *Limulus polyphemus* by a known method or prepared using a

known peptide synthesis method. Further, it can be also prepared by gene recombination technique using a polynucleotide encoding the amino acid sequence shown in SEQ ID NO:1 or 2. The polypeptide having the amino acid sequence shown in SEQ ID NO:1 or 2 can be also extracted from a living organism belonging to the genus *Tachypleus* by a known method or prepared using a known peptide synthesis method. Further, it can be also prepared by gene recombination technique using a polynucleotide encoding the amino acid sequence of the polypeptide.

[0034]

The homolog of the polypeptide having the amino acid sequence shown in SEQ ID NO:1 or 2 may have an improved D-LDH activity and/or an improved thermal stability as compared to the polypeptide having the amino acid sequence shown in SEQ ID NO:1 or 2. Here, the term "an improved D-LDH activity" refers to an improvement in the substrate affinity and molecular activity (Kcat) for pyruvic acid and NADH, as well as a shift in the optimum pH for the enzyme activity of the polypeptide closer to a pH suitable for growth of cells expressing the polypeptide. Such polypeptide may be artificially designed based on the polypeptide having the amino acid sequence shown in SEQ ID NO:1 or 2, or may be separated from nature. Further, a mutation may be randomly introduced to the D-lactate dehydrogenase gene by an evolutionary molecular engineering method to screen a preferred polypeptide therefrom.

[0035]

The present invention is characterized by including a polynucleotide having the nucleotide sequence shown in SEQ ID NO:3 or 4 or a homolog thereof, which polynucleotide encodes a polynucleotide having a D-LDH activity. Here, the origin of the "polynucleotide" does not matter and may be a cDNA, genomic DNA, synthetic DNA, mRNA, synthetic RNA, replicon RNA; however, it is preferably a DNA or RNA, more preferably DNA. Also, the "polynucleotide" may be either

single stranded or double stranded with a complementary strand thereof. Further, the "polynucleotide" may also contain a natural or artificial nucleotide derivative.

[0036]

5 The polynucleotide having the nucleotide sequence shown in SEQ ID NO:3 or 4 is a polynucleotide originated from *Limulus polyphemus* and characterized by encoding a polypeptide having the above-described D-LDH activity.

[0037]

10 Examples of the homolog of the polynucleotide shown in SEQ ID NO:3 or 4 include a polynucleotide having the same nucleotide sequence as shown in SEQ ID NO:3 or 4 except that one or several, preferably 1 to 40, more preferably 1 to 30, still more preferably 1 or 20, especially preferably 1 to 10, most preferably 1 to 5 nucleotides are substituted, deleted, inserted and/or added, which polynucleotide also encodes a polypeptide having a D-LDH activity.

[0038]

15 Further, examples of the homolog of the polynucleotide shown in SEQ ID NO:3 or 4 also include a polynucleotide which hybridizes under stringent conditions with the entirety or a part of the polynucleotide having the nucleotide sequence shown in SEQ ID NO:3 or 4 or a complementary strand thereof, which polynucleotide also encodes a polypeptide having a D-LDH activity. Here, the term
20 "a polynucleotide which hybridizes under stringent conditions" refers to, for example, a polynucleotide which is hybridized by a known hybridization technique (Current Protocols I Molecular Biology edit. Ausubel et al., (1987) Publish. John Wiley & Sons, Section 6.3-6.4) or the like using, as probe(s), one or a plurality of selected polynucleotides having at least 20, preferably 25, more preferably at least 30 arbitrary
25 consecutive nucleotides of the original nucleotide sequence. Here, stringent conditions can be achieved, for example, by performing hybridization in the presence of 50% formamide at a hybridization temperature of 37°C, 42°C for more stringent

condition, 65°C for still more stringent condition, and washing the resultant with a 0.1× to 2× SSC (saline-sodium citrate) solution (composition of 1× SSC solution: 150 mM sodium chloride and 15 mM sodium citrate).

[0039]

5 Further, the homolog of the polynucleotide shown in SEQ ID NO:3 or 4 may also be a polynucleotide having a nucleotide sequence which has a sequence identity of not less than 80%, more preferably not less than 90%, still more preferably not less than 95% to the nucleotide sequence shown in SEQ ID NO:3 or 4, which polynucleotide also encodes a polypeptide having a D-LDH activity. The sequence
10 identity of the nucleotide sequence of polynucleotide used herein can be determined by using the above-described gene analysis program BLAST or the like.

[0040]

Further, the homolog of the polynucleotide shown in SEQ ID NO:3 or 4 may also be a polynucleotide originated from a living organism belonging to the family
15 Limulidae, preferably a living organism belonging to the genus *Limulus* or *Tachypleus*, more preferably *Limulus polyphemus*, *Tachypleus tridentatus*, *Tachypleus gigas* or *Tachypleus rotundicauda*, which polynucleotide encodes a polypeptide having a D-LDH activity.

[0041]

20 Although the above-described polynucleotide can be prepared by cloning from a living organism belonging to the family Limulidae, preferably a living organism belonging to the family Limulidae and genus *Limulus* (hereinafter, these are also collectively referred to as "Limulus"), it can also be synthesized chemically or by employing the technique of Fujimoto et al. (Hideya Fujimoto, Production Method of
25 Synthetic Gene, Plant Cell Engineering Series 7, Plant PCR Experimental Protocol, 1997, Shujunsha Co., Ltd., p.95-100), which is known as a synthesis method of long-chain DNA. Normally, the polynucleotide cloned from *Limulus* can be isolated by

a known method. Examples of such method include one in which, using a primer synthesized based on a cDNA sequence which is highly conserved in the D-lactate dehydrogenase, the cDNA being prepared from poly A (+) RNA isolated from Limulus hemocytes, a partial fragment is amplified and sequenced, and the entire
5 ORF sequence thereof is then determined by 5' RACE and 3' RACE methods, followed by amplification by PCR to obtain a polynucleotide. Further, the cloning from Limulus can also be performed by complementing the D-lactate dehydrogenase function. The principle thereof is described in detail in DOMINIQUE, G., Appl Environ Microbiol, United States (1995), 61, 266-272. In this manner, the
10 polynucleotide of the present invention having a nucleotide sequence once determined can be once again obtained from Limulus hemocytes; however, it can be also directly synthesized chemically using a DNA synthesis apparatus.

[0042]

Here, the above-described polypeptide or polynucleotide can be modified as
15 appropriate by appropriately introducing a substitution(s), deletion(s), insertion(s) and/or addition mutation(s) to the amino acid sequence or nucleotide sequence thereof using a site-specific mutation introduction method (Current Protocols I Molecular Biology edit. Ausubel et al., (1987) Publish. John Wiley & Sons Section 8.1-8.5) or the like. Further, such modification of an amino acid sequence of
20 polypeptide or nucleotide sequence of polynucleotide is not restricted to those performed by artificial introduction or synthesis of a mutation, but encompasses not only those performed based on an artificial mutation treatment, but also those naturally occurred by a mutation in an amino acid.

[0043]

25 The above-described polynucleotide can be used for preparation of a D-lactic acid-producing cell by gene recombination. In order for a host cell in which the polynucleotide of the present invention is introduced by gene recombination

(hereinafter, also referred to as "transformant") to become able to produce D-lactic acid, it is required that the polypeptide having a D-LDH activity which is encoded by the polynucleotide be expressed in the transformant. In this case, as a method of allowing the polypeptide having a D-LDH activity to be expressed, it is useful to
5 utilize a DNA construct in which the polynucleotide and a promoter capable of expressing it are linked, that is, a DNA construct in which the polynucleotide is linked to the 3'-end of the promoter, and such DNA construct is also included in the present invention. The DNA construct of the present invention can be prepared by a method in which the polynucleotide (DNA) of the present invention and a promoter
10 capable of expressing the polynucleotide are each digested with an appropriate restriction enzyme(s) and then inserted to a restriction enzyme site or multicloning site of the below-described vector DNA to be linked or by linking the polynucleotide (DNA) of the present invention and the promoter by performing PCR.

[0044]

15 In a preferred mode of the above-described DNA construct, the DNA construct is carried by a vector such as a plasmid (DNA), bacteriophage (DNA), retrotransposon (DNA) or artificial chromosome (for example, YAC, PAC, BAC or MAC). Depending on the introduction mode of the DNA construct (inside or outside the host genome) and the type of the host cell, a prokaryotic vector,
20 eukaryotic vector, animal cell vector or plant cell vector, which is known in the art, is selected as appropriate.

[0045]

Here, examples of the plasmid include YCp-type *Escherichia coli*-yeast shuttle vectors such as pRS413, pRS415, pRS416, YCp50, pAUR112 and pAUR123;
25 YEp-type *Escherichia coli*-yeast shuttle vectors such as pYES32 and YEp13; YIp-type *Escherichia coli*-yeast shuttle vectors such as pRS403, pRS404, pRS405, pRS406, pAUR101 and pAUR135; plasmids originated from *Escherichia coli* (for

example, ColE-type plasmids such as pBR322, pBR325, pUC18, pUC19, pUC119, pTV118N, pTV119N, pBluescript, pHSG298, pHSG396 and pTrc99A; p1A-type plasmids such as pACYC177 and pACYC184; and pSC101-type plasmids such as pMW118, pMW119, pMW218 and pMW219); and plasmids originated from *Bacillus subtilis* (such as pUB110 and pTP5). Examples of the bacteriophage include λ -phages (such as Charon4A, Charon21A, EMBL3, EMBL4, λ gt100, gt11 and zap), ϕ X174, M13mp18 and M13mp19. An example of the retrotransposon is Ty factor and that of the YAC is pYACC2.

[0046]

The "promoter" in the above-described DNA construct means a nucleotide sequence involved in the initiation of transcription of mRNA from a gene and normally refers to an upstream sequence of the 5'-end of the gene existing on the chromosome. The length of the promoter nucleotide sequence is preferably 1 to 3,000 bp, more preferably 1 to 1,000 bp; however, it is not particularly restricted as long as the nucleotide sequence can initiate the transcription of mRNA of the gene which exists in downstream. Further, a mutation and operation for improving the transcription activity of a promoter are known and the "promoter" also includes those which are modified by a known method. The promoter in the above-described DNA construct is not particularly restricted as long as it has a promoter activity in the transformant introduced with the DNA construct; however, as described below, since it is preferred that the DNA construct of the present invention be introduced into a yeast, the promoter is preferably functional in the yeast.

[0047]

Preferred examples of the promoter functioning in yeast include those of acid phosphatase gene (PHO5), glyceraldehyde-3-phosphate dehydrogenase genes (TDH1, 2 and 3), alcohol dehydrogenase genes (ADH1, 2, 3, 4, 5, 6 and 7), galactose metabolism-related genes (GAL1, 7 and 10), cytochrome c gene (CYC1),

triosephosphate isomerase gene (TPI1), phosphoglycerate kinase gene (PGK1), phosphofructose kinase gene (PFK1) and pyruvate decarboxylase genes (PDC1, 5, 6), as well as those promoters of enolase-1 gene (ENO1), cell wall-related protein 2 gene (CWP2) and suppression-of-exponential-defect 1 gene (SED1) which are described and used in the international application PCT/JP 2008/072129 and exhibit a gene expression amount of not less than 5 times of the average relative expression amount of all genes after 50 hours or later from the start of yeast culture.

[0048]

More preferred examples of the promoter functioning in yeast include the PDC1 promoter and TDH3 promoter which are highly expressed in the ethanol fermentation pathway of yeast; and the SED1 promoter which is highly expressed in a yeast cultured for a prolonged period, and more specific examples include the PDC1 promoter shown in SEQ ID NO:5, the SED1 promoter shown in SEQ ID NO:6 and the TDH3 promoter shown in SEQ ID NO:7.

[0049]

Further, the PDC1 promoter, SED1 promoter or TDH3 promoter may also have, as long as its promoter activity is maintained, the same nucleotide sequence as shown in one of SEQ ID NOs:5 to 7, respectively, except that one or several, preferably 1 to 40, more preferably 1 to 30, still more preferably 1 or 20, especially preferably 1 to 10, most preferably 1 to 5 nucleotides are deleted, substituted and/or added.

[0050]

Further, the PDC1 promoter, SED1 promoter or TDH3 promoter may also have, as long as its promoter activity is maintained, a nucleotide sequence which hybridizes under stringent conditions with a nucleotide sequence containing the nucleotide sequence shown in one of SEQ ID NOs:5 to 7 or a part thereof, respectively. Here, a nucleotide sequence which "hybridizes under stringent

conditions" refers to, for example, a polynucleotide which is hybridized by a known hybridization technique (Current Protocols I Molecular Biology edit. Ausubel et al., (1987) Publish. John Wily & Sons, Section 6.3-6.4) or the like using, as probe(s), one or a plurality of selected polynucleotides having at least 20, preferably 25, more preferably at least 30 arbitrary consecutive nucleotides of the original nucleotide sequence. Here, stringent conditions can be achieved, for example, by performing hybridization in the presence of 50% formamide at a hybridization temperature of 37°C, 42°C for more stringent condition, 65°C for still more stringent condition, and washing the resultant with a 0.1× to 2× SSC (saline-sodium citrate) solution (composition of 1× SSC solution: 150 mM sodium chloride and 15 mM sodium citrate).

[0051]

The present invention also includes a transformant obtained by introducing the above-described polynucleotide or DNA construct into a host cell. The host cell is not particularly restricted as long as it stably retains the polynucleotide or DNA construct and examples thereof include bacteria such as *Escherichia coli*, *Bacillus subtilis* and lactic acid bacteria; yeast; insect cells; animal cells; and plant cells, and yeast is preferably used since it is acid-resistant and capable of growing even when D-lactic acid is highly produced by expressing a polypeptide having a D-LDH activity.

[0052]

Examples of the yeast include those belonging to the genera *Saccharomyces*, *Schizosaccharomyces*, *Kluyveromyces*, *Candida*, *Pichia*, *Hansenura*, *Yarrowia*, *Zygosaccharomyces*, *Torulopsis*, *Debaryomyces*, *Issachenkia* and *Fellomyces*; however, preferably, the yeast is one belonging to the genus *Saccharomyces*, *Candida* or *Kluyveromyces*, more preferably *Saccharomyces cerevisiae*, *Candida utilis*, *Candida glabrata*, *Candida albicans*, *Candida boidinii*, *Candida sonorensis*,

Kluyveromyces lactis or *Kluyveromyces marxianus*.

[0053]

Further, a polyploid yeast is preferably used in the present invention since it can stably maintain a high productivity of D-lactic acid over a prolonged period under simple operating conditions, so that D-lactic acid can be stably produced at a low cost. Here, the term "polyploid yeast" refers to yeast having 2 or more sets of chromosomes in the cell. The number of sets of chromosomes in the polyploid yeast is not particularly restricted; however, diploid yeast having 2 sets of chromosomes is preferred. Examples of the polyploid yeast include baker's yeasts, sake yeasts, wine yeasts and beer yeasts that are frequently used in the fermentation industry. The polyploid yeast to be used may be one isolated from a natural environment or one whose properties are partially modified by mutation or gene recombination.

[0054]

Further, in the present invention, a prototrophic yeast is preferably used since it not only allows the use of a culture medium containing less nutrients than conventional media, that is, a low-cost medium, but also enables stable maintenance of a high productivity of D-lactic acid over a prolonged period under simple operating conditions, so that lactic acid can be stably produced at a low cost. Here, the term "auxotrophic" means that a mutation occurred in a nutrient-synthesis gene of the wild-type yeast for some reason, resulting in a deficiency in the capacity to synthesize the nutrient. That is, "prototrophic yeast" is one which does not have genotype exhibiting auxotrophy or one in which such genotype is complemented. Examples of a method for preparing a prototrophic yeast by reverting the auxotrophy of an auxotrophic yeast include a method in which an auxotrophy is reverted by introducing a nutrient-synthesis gene by a gene recombination technique; and a method in which the processes of mating yeasts having different auxotrophies with

each other, allowing ascus formation and reverting a target auxotrophy are repeated until all of the auxotrophies are eventually reverted to obtain a prototrophic yeast. It is noted here that whether or not a yeast is prototrophic can be judged based on whether or not the yeast is capable of growing in SD medium, which is a minimal medium for yeasts.

[0055]

Further, yeast is a microorganism which vigorously performs ethanol fermentation, and in the metabolic pathway thereof, pyruvic acid which is a glycolytic product is converted to acetaldehyde by pyruvate decarboxylase, which is then converted to ethanol by alcohol dehydrogenase. Therefore, it is preferred that a yeast strain in which the pyruvate decarboxylase gene serving as the starting point of the ethanol metabolic pathway is destructed be used as host since this allows pyruvic acid, which should be naturally metabolized in the ethanol metabolic pathway, to be used in the D-lactic acid metabolic pathway.

[0056]

Particularly in the case of *Saccharomyces cerevisiae*, there are three types of genes encoding a pyruvate decarboxylase: pyruvate decarboxylase 1 (PDC1) gene, pyruvate decarboxylase 5 (PDC5) gene and pyruvate decarboxylase 6 (PDC6) gene. Among these genes, only the PDC1 and PDC5 are considered to have pyruvate dehydrogenase activity in the yeast cell; therefore, it is preferred that a strain having destructed PDC1 gene or PDC5 gene be used and it is more preferred that a strain having destructed PDC1 gene be used. In addition, as described in JP 2008-048726A, in the strain having destructed PDC1 gene, a strain having a mutant-type PDC5 gene which includes a mutation resulting in a decrease in the PDC5 activity is preferred, and a strain having a temperature-sensitive mutant-type PDC5 gene is more preferred. Here, the "temperature-sensitive mutant-type PDC5 gene" refers to a mutant-type PDC5 which has a property to exhibit a pyruvate decarboxylase

activity comparable to that of the wild-type PDC5 at a certain culturing temperature, but exhibit a deletion of or decrease in the PDC5 activity when the culturing temperature is changed to not lower than the certain culturing temperature. The normal culturing temperature of *Saccharomyces cerevisiae* is 28 to 30°C and it is preferred that the temperature at which such temperature sensitivity is exhibit be close to the normal culturing temperature since this makes the amount of heat required for changing the culturing temperature smaller, so that the culturing cost can be reduced. In particular, it is preferred that the temperature-sensitive mutant-type PDC5 exhibit the temperature sensitivity at a temperature of not lower than 34°C.

[0057]

Examples of a method of allowing a polypeptide having a D-LDH activity to be expressed in a transformed yeast by introducing the above-described polynucleotide or DNA construct to the yeast include, as already described, a method in which a vector containing the DNA construct is extrachromosomally introduced into a yeast and a method in which the polynucleotide or DNA construct is introduced into a yeast chromosome, and both of these method can be employed.

[0058]

In the method of introducing the above-described polynucleotide into a yeast chromosome, a construct which contains the polynucleotide of the present invention and a DNA segment for homologous recombination whose sequence is homologous to that of the yeast chromosome (hereinafter, also referred to as "DNA construct for homologous recombination") can be preferably used. The DNA segment for homologous recombination has a DNA sequence which is homologous to the one in the vicinity of the target site in the yeast chromosome to which the polynucleotide of the present invention is to be introduced. The construct contains at least one, preferably two such DNA segments for homologous recombination. In cases where the construct contains two DNA segments for homologous recombination, it is

preferred that the two DNA segments for homologous recombination each have a DNA sequence homologous to the DNA of the upstream or downstream of the target site located on the chromosome and that the polynucleotide of the present invention be linked between these DNA segments.

5 [0059]

The method of performing homologous recombination using the DNA construct for homologous recombination is not particularly restricted, and a method in which the DNA construct for homologous recombination is amplified by PCR and the resulting PCR fragment is inserted into a plasmid which is then introduced into a yeast or a method in which the PCR fragment is introduced into a yeast can be employed. Further, it is preferred that the DNA construct for homologous recombination contain a terminator for regulating the expression of the polynucleotide of the present invention. Here, the "terminator" means a sequence which terminates mRNA transcription from a gene and normally refers to a downstream sequence of the 3'-end of the gene existing on the chromosome.

15

[0060]

In addition, it is preferred that the DNA construct for homologous recombination contain, in addition to the polynucleotide or DNA construct of the present invention, a selective marker for making the selection of transformed yeast easy. Examples of the selective marker used herein include auxotrophic complementary genes such as URA3 and TRP1 and drug-resistant genes such as G418-resistant gene and neomycin-resistant gene.

20

[0061]

A DNA construct for homologous recombination containing the above-described polynucleotide, terminator and selective marker can be prepared by PCR in accordance with, for example, the following steps 1 to 3.

25

[0062]

Step 1: Using a plasmid in which a terminator is linked to downstream of the polynucleotide of the present invention as template and primers 1 and 2 as primer set, a fragment containing the polynucleotide of the present invention and the terminator is amplified by PCR. The primer 1 is designed in such a manner to add a sequence of not less than 40 bp complementary to the upstream of the introduction target site, and the primer 2 is designed based on a sequence originated from the plasmid which is located the downstream of the terminator.

[0063]

Step 2: Using a plasmid having a selective marker such as pRS400, pRS424 or pRS426 as template and primers 3 and 4 as primer set, a fragment containing the selective marker is amplified by PCR. The primer 3 is designed in such a manner to add a sequence of not less than 30 bp which has a homology to a sequence located downstream of the terminator of the PCR fragment obtained in the step 1, and the primer 4 is designed in such a manner to add a sequence of not less than 40 bp corresponding to the downstream of the introduction target site.

[0064]

Step 3: Using a mixture of the PCR fragments obtained in the steps 1 and 2 as template and primers 1 and 4 as primer set, PCR is performed to obtain a DNA construct for homologous recombination containing the polynucleotide of the present invention, terminator and yeast selective marker, in which sequences corresponding to the upstream and downstream of the introduction target site are added to both respective ends.

[0065]

In order to introduce the above-described DNA construct for homologous recombination into a yeast, a method such as transfection, cotransfection or electroporation can be employed. Specific examples thereof include a method using lithium acetate and protoplast method.

[0066]

When introducing the above-described polynucleotide into a yeast chromosome by homologous recombination, the introduction is carried out in such a manner that the polynucleotide can be regulated by a promoter of an endogenous gene located on the yeast chromosome. In this case, by the introduction of the above-described polynucleotide, the endogenous gene to be naturally regulated by the promoter may be simultaneously destructed and the exogenous polynucleotide of the present invention may be expressed in place of this endogenous gene. This method is particularly useful when the promoter is a high-expression promoter in yeast as described in the above and when an endogenous gene functioning to inhibit the D-lactic acid metabolic pathway of yeast is to be destructed.

[0067]

Preferred examples of the endogenous gene which serves as a target when introducing the above-described polynucleotide into a yeast chromosome include acid phosphatase gene (PHO5), glyceraldehyde-3-phosphate dehydrogenase genes (TDH1, 2 and 3), alcohol dehydrogenase genes (ADH1, 2, 3, 4, 5, 6 and 7), galactose metabolism-related genes (GAL1, 7 and 10), cytochrome c gene (CYC1), triosephosphate isomerase gene (TPI1), phosphoglycerate kinase gene (PGK1), phosphofructose kinase gene (PFK1) and pyruvate decarboxylase genes (PDC1, 5, 6), as well as promoters of enolase-1 gene (ENO1), cell wall-related protein 2 gene (CWP2) and suppression-of-exponential-defect 1 gene (SED1) which are described and used in the international application PCT/JP 2008/072129 and exhibit a gene expression amount of not less than 5 times of the average relative expression amount of all genes after 50 hours or later from the start of yeast culture. In particular, more preferred examples include the PDC1 gene and TDH3 gene which are highly expressed in the ethanol fermentation pathway of yeast; and the SED1 gene which is highly expressed in a yeast cultured for a prolonged period. As described in the

above, it is still more preferred that the above-described polynucleotide be introduced in such a manner to destruct at least the PDC1 gene serving as the starting point of the ethanol fermentation pathway or PDC5 gene, preferably the PDC1 gene, since this allows pyruvic acid metabolized in the ethanol metabolic pathway to be used in the D-lactic acid metabolic pathway.

[0068]

Further, as the method of introducing the above-described DNA construct into a yeast chromosome, in the same manner as in the case of introducing the above-described polynucleotide into a yeast chromosome, a DNA construct for homologous recombination can be prepared and introduced at an intended site located on the chromosome by homologous recombination. It is noted here that, since the above-described DNA construct already contains a promoter capable of expressing the above-described polynucleotide, it is not necessary to carry out the introduction of the DNA construct in such a manner that the above-described polynucleotide can be regulated by a promoter of an endogenous gene located on the yeast chromosome.

[0069]

Here, whether or not the above-described polynucleotide or DNA construct was introduced at the desired position on the yeast chromosome can be confirmed by PCR or Southern hybridization method. For example, the confirmation can be carried out by preparing a DNA from a transformed yeast and performing PCR with an introduction site-specific primer, followed by detection of the expected band by performing electrophoresis for the resulting PCR product. Alternatively, the confirmation can be carried out by performing PCR with a primer labeled with a fluorescent dye or the like. These methods are known to those skilled in the art.

[0070]

The present invention also includes a method of producing D-lactic acid which comprises the step of culturing the above-described transformant. When the

above-described transformant is cultured, expression of a polypeptide having a D-LDH activity is newly provided or enhanced in the transformant and, thus, D-lactic acid metabolic pathway is newly provided or enhanced. As a result, D-lactic acid can be obtained by carrying out the step of separating D-lactic acid from the culture.

5 The term "culture" used herein encompasses a culture supernatant, as well as transformant and homogenate thereof.

[0071]

The method of culturing the above-described transformant is not particularly restricted and a known method is employed. For example, as the method of
10 culturing the above-described transformed yeast, one described in "M.D. Rose et al., 'Methods In Yeast Genetics', Cold Spring Harbor Laboratory Press (1990)" or the like can be employed. Here, in cases where a plasmid containing the above-described DNA construct or DNA construct for homologous recombination contains a selective marker, by culturing the above-described transformant in a nutrient-free medium or
15 drug-supplemented medium depending on the selective marker, a desired transformant can be selected.

[0072]

As the medium in which the above-described transformant is cultured, as long as it contains a carbon source, nitrogen source, inorganic salts and the like that can be
20 utilized by the transformant and enables efficient culturing thereof, any natural medium or synthetic medium can be used. For example, in the case of a transformed yeast, as the carbon source, carbohydrates such as glucose, fructose, sucrose, galactose, maltose, raffinose, trehalose, sorbose, cellobiose, lactose, melibiose, melezitose, inulin, xylose, arabinose, ribose, rhamnose, glucosamine,
25 erythritol, ribitol, mannitol, glucitol, salicin and starch; organic acids such as acetic acid, propionic acid and citric acid; and alcohol such as ethanol and propanol can be used. As the nitrogen source, in addition to ammonium salts of an inorganic or

organic acid such as ammonia, ammonium chloride, ammonium sulfate, ammonium acetate and ammonium phosphate and other nitrogen-containing compounds, peptone, meat extract, corn steep liquor and the like can be used. As inorganic compound, monopotassium phosphate, magnesium phosphate, magnesium sulfate, sodium chloride, ferrous sulfate, manganese sulfate, copper sulfate, calcium carbonate and the like can be used.

[0073]

In the culturing of the above-described transformant, usually, the condition thereof is set to one in which good lactic acid productivity is attained, ranging from an aerobic condition, such as culture with shaking or culture with aeration and stirring, to an anaerobic condition where aeration is not performed, and the culturing is preferably carried out under a microaerobic condition or anaerobic condition.

The culturing can be carried out at a culturing temperature of 25 to 37°C; however, a temperature of 28 to 35°C is preferred. Since the pH of the culture decreases as D-

lactic acid is accumulated in the medium, it can be also neutralized by an alkaline substance such as calcium hydroxide, calcium carbonate, sodium hydroxide, potassium hydroxide, aqueous ammonia or ammonia gas. Further, in order to facilitate the subsequent purification of lactic acid, the culturing can be also carried out without such neutralization. Moreover, the culturing method is not particularly restricted as long as it can fermentatively produce D-lactic acid of interest, and batch culture, fed-batch culture, chemostat culture, continuous culture or the like can be employed. Preferably, the culturing is carried out by batch culture or the continuous culture using a membrane described in WO2007/097260, where fermentation culture is separated into a filtrate and non-filtrate using a separation membrane not liable to clogging and a desired fermentation product is collected from the filtrate, and at the same time, the non-filtrate is retained in or refluxed to the fermentation culture.

[0074]

The method of measuring D-lactic acid obtained in the above-described culture is not particularly restricted, and examples thereof include a method using HPLC and a method using the F-kit (manufactured by Roche).

[0075]

5 The D-lactic acid contained in the above-described fermentation culture can be separated and purified by a combination of those conventionally known methods of concentration, distillation, crystallization and the like, and examples thereof include a method in which the pH of filtered and separated fermentation liquid is adjusted to not higher than 1, followed by extraction with diethyl ether, ethyl acetate
10 or the like; a method in which lactic acid is allowed to adsorb to an ion-exchange resin followed by washing and elution; a method in which lactic acid is allowed to react with an alcohol in the presence of an acid catalyst to obtain an ester, which is then distilled; a method in which lactic acid is crystallized in the form of a calcium salt or lithium salt; and a separation and purification method in which the nanofilter
15 disclosed in WO2009/004922 and distillation are used in combination.

EXAMPLES

[0076]

The modes for carrying out the present invention will now be described by way of examples; however, these modes are merely examples of the present
20 invention and the present invention is not restricted thereto.

[0077]

(Example 1) Nucleotide sequencing of polynucleotides originated from *Limulus polyphemus* which encode a polypeptide having a D-LDH activity (hereinafter, referred to as "D-LDH genes originated from *Limulus polyphemus* ")

25 For preparation of cDNAs of *Limulus polyphemus*, polyA (+) RNA was purified. The polyA (+) RNA was isolated from the hemocytes of *Limulus polyphemus*. Total RNA was separated from the hemocytes of *Limulus polyphemus*

(purchased from Marine Biological Laboratory (USA)) in accordance with the AGPC method (see Experimental Medicine Vol. 9 (1991), p.1937-1940). The polyA (+) RNA was isolated from the thus separated total RNA using "Oligotex-dT30 Super Kit" (manufactured by Takara Bio Inc.). Next, using "SuperScript II Reverse Transcriptase" (Manufactured by Invitrogen) and oligo d (T) as primer, cDNAs were synthesized. Then, using the thus obtained reaction solution as template and the oligo DNAs shown in SEQ ID NOs:8 and 9 as primers, which oligo DNAs were designed with attention given to the DNA sequences highly conserved in the D-LDH genes, PCR was performed to amplify partial sequences of the D-LDH genes originated from *Limulus polyphemus*. The PCR was performed under the following conditions: 95°C (3 minutes); {95°C (30 seconds) - 45°C (30 seconds) - 72°C (30 seconds)} × 35 cycles; 72°C (5 minutes). Thereafter, the thus amplified DNA fragments were verified by electrophoresis using 1.0% agarose gel, and after purifying an amplified fragment of 600 bp using "QIA quick Gel extraction kit" (manufactured by Qiagen), this fragment was cloned into a plasmid vector "pGEMt-easy" (manufactured by Promega) and the DNA was sequenced. The nucleotide sequencing was performed using "Taq DyeDeoxy Terminator Cycle Sequencing Kit" (manufactured by Applied Biosystems) in accordance with Sanger's method. As a result, two kinds of partial sequences (D-LDH1 and D-LDH2) which are thought to correspond to the D-LDH genes originated from *Limulus polyphemus* were obtained.

[0078]

Next, using 5' RACE and 3' RACE methods, the entire ORF sequences of the D-LDH1 and D-LDH2 were clarified. The 5' RACE was performed by the following method. First, 50 µL of the oligo DNA shown in SEQ ID NO:10 (for D-LDH1) or SEQ ID NO:11 (for D-LDH2) which was adjusted to 100 µM, 50 units of T4 polynucleotide kinase (manufactured by Takara Bio Inc.), 10 µL of 10× kinase buffer (500 mM Tris-HCl pH7.6, 100 mM MgCl₂, 50 mM DTT, 1 mM spermidine, 1

mM EDTA), 1 μ L of 100 mM ATP and 34 μ L of distilled water were mixed and left to stand at 37°C for 1 hour to phosphorylate the 5'-end of the oligo DNA, and this oligo DNA was purified by performing a phenol-chloroform-isoamyl alcohol treatment and ethanol precipitation. Next, using the poly A (+) RNA of *Limulus polyphemus* as template, "SuperScript II Reverse Transcriptase" (manufactured by Invitrogen) and the above-described phosphorylated oligo DNA as primers, cDNA was synthesized. Thereafter, single-stranded cDNA was concatemerized by ligation reaction using "5'-Full RACE Core Set" (manufactured by Takara Bio Inc.) and the first PCR was performed using the thus produced concatemerized cDNA as template and the oligo DNAs shown in SEQ ID NOs:12 and 13 (for D-LDH1) or those shown in SEQ ID NOs:14 and 15 (for D-LDH2) as primers. Further, the second PCR was performed using the resulting reaction solution as template and the oligo DNAs shown in SEQ ID NOs:16 and 17 (for D-LDH1) or those shown in SEQ ID NOs:18 and 19 (for D-LDH2) as primers. The verification and purification of the amplified fragments and the nucleotide sequencing thereof were performed as described in the above.

[0079]

The 3' RACE was performed by the following method. First, using the polyA (+) RNA of *Limulus polyphemus* as template, "SuperScript II Reverse Transcriptase" (manufactured by Invitrogen) and oligo d (T) as primer, cDNA was synthesized. Next, using this reaction solution as template and the oligo d (T) and the oligo DNA shown in SEQ ID NO:13 (for D-LDH1) or SEQ ID NO:15 (for D-LDH2) as primers, the first PCR was performed. Further, the second PCR was performed using the resulting reaction solution as template and the oligo d (T) and the oligo DNA shown in SEQ ID NO:17 (for D-LDH1) or SEQ ID NO:19 (for D-LDH2) as primers. The verification and purification of the amplified fragments and the nucleotide sequencing thereof were performed as described in the above.

[0080]

The amino acid sequence (SEQ ID NO:1) of the D-LDH1 gene originated from *Limulus polyphemus* and the cDNA ORF sequence (SEQ ID NO:3), or the amino acid sequence (SEQ ID NO:2) of the D-LDH2 and the cDNA ORF sequence (SEQ ID NO:4) were determined by ligating the sequences obtained in the above-described method. It is noted here that the sequence identities between the amino acid sequences shown in SEQ ID NOs:1 and 2 and between the nucleotide sequences shown in SEQ ID NOs:3 and 4 were determined by BLAST to be 93% and 82%, respectively.

[0081]

(Example 2) Construction of a plasmid expressing the D-LDH genes originated from *Limulus polyphemus*

Using, as a primer set, the oligo DNAs shown in SEQ ID NOs:20 and 21 or those shown in SEQ ID NOs:22 and 23, which respectively correspond to the ORF 5'-side and ORF 3'-side of the D-LDH1 gene and D-LDH2 gene which were sequenced in Example 1 and are thought to be the D-LDH genes originated from *Limulus polyphemus* (hereinafter, referred to as "Lp. D-LDH1 gene" and "Lp. D-LDH2 gene", respectively) and, as template, the cDNA synthesized from the poly A (+) RNA of *Limulus polyphemus*, PCR was performed to clone the genes. The thus obtained DNA fragments were digested with restriction enzymes XhoI and NotI and introduced into the cleavage sites of yeast expression vector pTRS11 (carrying TDH3 promoter and selective marker URA3 gene; see JP 2006-280368A: it is noted here that, therein, the TDH3 promoter and terminator are indicated as GAPDH promoter and terminator), which was digested in the same manner with the restriction enzymes XhoI and NotI, to prepare plasmid vectors containing a DNA construct in which the TDH3 promoter and the Lp. D-LDH1 gene or Lp. D-LDH2 gene are linked. Hereinafter, these plasmid vectors are referred to as "pTRS205" (carrying the Lp. D-

LDH1) and "pTRS206" (carrying the Lp. D-LDH2). Here, the pTRS11 vector was prepared by digesting pNV11 vector (see Nature, vol. 357, 25 JUNE 1992, p.700) with the restriction enzyme XhoI and, after removing inserts, allowing it to self-ligate. [0082]

5 (Example 3) Introduction of the plasmid expressing the D-LDH genes originated from *Limulus polyphemus* into yeast

The pTRS205 and pTRS206 obtained as described in Example 2 were introduced into the yeast *Saccharomyces cerevisiae* SW029-1B strain (genotype: MATa ura TRP(Δ pdcl::TRP1) his ade lys leu) (hereinafter, referred to as "SW029-1B strain"). The introduction of the plasmids was carried out by a lithium acetate method using "YEASTMAKER Yeast Transformation System" (manufactured by Clontech) (for details, see the attached protocol). The SW029-1B strain used as the host is a strain which lacks uracil synthesis capacity and, by the action of the selective marker URA3 gene of the pTRS205 and 206, transformed yeast into which the pTRS205 and pTRS206 were introduced can be selected on a uracil-free medium. The introduction of the D-LDH gene-expressing vectors into the transformants obtained in this manner was confirmed by extracting genomic DNA containing plasmid DNA from the transformants cultured in a uracil-free liquid medium using a genomic DNA extraction kit "Dr. GenTLE" (manufactured by Takara Bio Inc.) and performing PCR using the thus extracted genomic DNA as template. As primers, those which were used for cloning the D-LDH genes originated from *Limulus polyphemus* were employed, respectively. As a result, it was confirmed that each of the D-LDH genes originated from *Limulus polyphemus* were respectively introduced into all of the transformed yeast cells.

25 [0083]

(Example 4) D-lactic acid productivity test 1

Using the *Saccharomyces cerevisiae* SW029-1B strains obtained as described

in Example 3 into which the pTRS205 and 206 were introduced (hereinafter, referred to as "SW029-1B/pTRS205 strain" and "SW029-1B/pTRS206 strain", respectively), D-lactic acid productivity test was carried out.

[0084]

5 SC3 medium having the composition shown in Table 1 from which uracil was removed (SC3-Ura medium) was added in test tubes in an amount of 10 mL. A small amount of each strain was inoculated thereto and cultured overnight at 30°C (preculture). Next, 10 mL of SC3-Ura medium was added in test tubes and 100 µL of each of the thus obtained preculture media was inoculated therein, followed by
10 culturing at 30°C with shaking (main culture). After 40 hours since the start of the main culture, the culture media were centrifuged and lactic acid contained in the resulting supernatants was measured by HPLC under the following conditions:
Column: "Shim-Pack SPR-H" (manufactured by Shimadzu Corporation)
Mobile phase: 5 mM p-toluenesulfonic acid (flow rate = 0.8 mL/min)
15 Reaction solution: 5 mM p-toluenesulfonic acid, 20 mM Bistris, 0.1 mM EDTA·2Na (flow rate = 0.8 mL/min)
Detection method: electroconductivity
Temperature: 45°C.

[0085]

20 In addition, based on the results of measuring the D-lactic acid and L-lactic acid concentrations by HPLC method under the following conditions, the optical purity of D-lactic acid was calculated using the following equations.

Column: "TSK-gel Enantio LI" (registered trademark: manufactured by Tosoh Corporation)

25 Mobile phase: 1 mM aqueous copper sulfate solution

Flow rate: 1.0 ml/min

Detection method: UV 254 nm

Temperature: 30°C

Optical purity (% e.e.) = $100 \times (D - L)/(D + L)$

Optical purity (%) = $100 \times D/(D + L)$

In the above equations, L represents the concentration of L-lactic acid and D
5 represents the concentration of D-lactic acid.

[0086]

[Table 1]

Glucose	100	g/L
Ammonium sulfate	1.5	g/L
Yeast Nitrogen base w/o amino acid and ammonium sulfate (manufactured by Difco)	1.7	g/L
19 standard amino acids other than leucine	152	mg/L
Leucine	760	mg/L
Inositol	152	mg/L
p-aminobenzoate	16	mg/L
Adenine	40	mg/L
Uracil	152	mg/L

[0087]

As a result, the concentration of the accumulated D-lactic acid was 13 g/L for
10 the SW029-1B/pTRS205 strain and 12 g/L for the SW029-1B/pTRS206 strain. In addition, only D-lactic acid was detected in the culture media and the level of L-lactic acid was not higher than the detection limit. It was confirmed that the transformed yeast was able to produce D-lactic acid by introducing therein the D-LDH genes cloned from *Limulus polyphemus*.

15 [0088]

(Example 5) Introduction of the D-LDH genes originated from *Limulus polyphemus* into yeast chromosome

By the following steps 1 to 3, a DNA construct for homologous recombination which contains the D-LDH genes originated from *Limulus*
20 *polyphemus* was prepared.

[0089]

[Step 1]

Using the pTRS205 carrying the Lp. D-LDH1 gene and the pTRS206 carrying the Lp. D-LDH2 gene that were obtained in Example 2 as templates and the oligo DNAs shown in SEQ ID NOs:24 and 25 or those shown in SEQ ID NOs:26 and 25 as primer set, PCR was performed to amplify DNA fragments of about 1.1 kb containing each of the D-LDH genes. Here, the oligo DNAs shown in SEQ ID NOs:24 and 26 were designed in such a manner that a sequence having a homology to the 65 bp region upstream of the PDC1 gene is added.

[0090]

10 [Step 2]

Next, using the plasmid pRS424 (GenBank Accession Number: U03453) as template and the oligo DNAs shown in SEQ ID NOs:27 and 28 as primer set, PCR was performed to amplify a DNA fragment of about 1.4 kb containing the TRP1 gene, which is a yeast selection marker. Here, the oligo DNA shown in SEQ ID NO:28 was designed in such a manner that a sequence having a homology to the 65 bp region downstream of the PDC1 gene is added.

[0091]

[Step 3]

After separating the DNA fragments by 1% agarose gel electrophoresis, they were purified using "QIA quick Gel extraction kit" (manufactured by Qiagen). Using a mixture of the thus obtained 1.1 kb fragments containing the respective D-LDH genes and 1.4 kb fragment containing the TRP1 gene as template and the oligo DNAs shown in SEQ ID NOs:24 and 28 or those shown in SEQ ID NOs:26 and 28 as primer set, PCR was performed to amplify a DNA construct of about 2.5 kb used for homologous recombination in which each of the D-LDH genes, TDH3 terminator and TRP 1 gene were linked.

[0092]

The DNA construct for homologous recombination prepared in the above-described manner was transformed into *Saccharomyces cerevisiae* SW092-2D strain (hereinafter, referred to as "SW092-2D strain"), which is a *Saccharomyces cerevisiae* NBRC10505 strain (hereinafter, referred to as "NBRC10505 strain") in which the lysine auxotrophy is reverted.

[0093]

The SW092-2D strain was prepared by the following method. Using the genomic DNA of the *Saccharomyces cerevisiae* BY4741 strain manufactured by Funakoshi Corporation as template and the oligonucleotides (SEQ ID NOs:29 and 30) as primer set, PCR was performed to amplify a PCR fragment of about 2 kb corresponding to the former half of the LYS2 gene. After separating the PCR fragment by 1% agarose gel electrophoresis and purifying it in accordance with a conventional method, the PCR fragment was transformed into the NBRC10505 strain to cancel the amber mutation in the LYS2 gene. By culturing the yeast strain in a lysine-free medium, a transformant in which the lysine synthesis capacity was reverted (NBRC10505(LYS2) strain) was selected. In order to confirm that the thus obtained transformant is a yeast in which the amber mutation in the LYS2 gene was canceled, the following steps were carried out. First, diploid cells were obtained by mating the thus obtained transformant with the *Saccharomyces cerevisiae* L0GY7 strain having the wild-type LYS2 gene and the diploid cells were cultured in an ascus formation medium to allow ascus formation. The resulting asci were anatomized using a micromanipulator to obtain respective haploid cells (tetrad) and the auxotrophy was examined for each of the haploid cells. All of the thus obtained haploid cells were confirmed to have lysine synthesis capacity. The thus obtained NBRC10505(LYS2) strain and the NBRC10506 strain were mated and the resulting asci were anatomized using a micromanipulator to obtain the SW092-2D strain (genotype: MATa ura3 leu2 trp1 his3 ade2 LYS2).

[0094]

The SW092-2D strain was transformed with the above-described DNA construct for homologous recombination and cultured in a tryptophan-free medium to select transformed yeasts. The transformed yeasts obtained in this manner are hereinafter referred to as "SW092-2D (Δ pdcl::Lp.D-LDH1-TRP1) strain" and "SW092-2D (Δ pdcl::Lp.D-DLH2-TRP1) strain".

[0095]

(Example 6) D-lactic acid productivity test 2

Using the SW092-2D (Δ pdcl::Lp.D-LDH1-TRP1) strain, SW092-2D (Δ pdcl::Lp.D-LDH2-TRP1) strain and a mini-jar fermentor (manufactured by B.E. Marubishi Co., Ltd., 5 L), their fermentation was evaluated.

[0096]

The SC3 medium shown in Table 1 was added to a test tube in an amount of 10 mL and a small amount of each strain was inoculated thereto and cultured overnight at 30°C (pre-preculture). Then, to 45 mL of SC3 medium placed in an Erlenmeyer flask, 5 mL of the thus obtained pre-preculture medium was added and cultured at 30°C for another 8 hours (preculture). The entire amount of the thus obtained preculture medium was inoculated to 1 L of SC3 medium placed in the mini-jar fermentor and cultured for 30 hours. The culturing conditions are shown below:

pH: pH5

Aeration: 100 mL/min

Stirring: 120 rpm

Neutralizer: 1N sodium hydroxide.

[0097]

The amount of the culture medium at the end of the culture period and the D-lactic acid and glucose concentrations in the culture medium were measured by

"Glucose Test Wako C" (registered trademark) (manufactured by Wako Pure Chemical Industries, Ltd.) to determine the lactic acid yield with respect to saccharide based on the added glucose calculated from the thus measured lactic acid and glucose concentrations. As a result, it was confirmed that the SW092-2D
 5 (Δ pdc1::Lp.D-LDH1-TRP1) and SW092-2D (Δ pdc1::Lp.D-LDH2-TRP1) produced D-lactic acid with a lactic acid yield with respect to saccharide of 45% and 42%, respectively.

[0098]

(Comparative Example 1) Cloning of the D-LDH gene originated from lactic acid
 10 bacterium and evaluation of fermentation

As a D-LDH gene capable of efficiently performing fermentative production of D-lactic acid in transformed yeast, the D-LDH gene originated from the *Leuconostoc mesenteroides* ATCC 9135 strain is known (see WO2004/104202). By cloning this D-LDH gene (hereinafter, referred to as "Lm. D-LDH gene") and
 15 introducing it into yeast to allow fermentation, the D-LDH activity of the Lm. D-LDH gene was examined by comparing it to that of the Lp. D-LDH1 gene or Lp. D-LDH2 gene originated from *Limulus polyphemus*.

[0099]

Using a primer set (SEQ ID NOs:31 and 32) which was designed by referring
 20 to the nucleotide sequences described in WO2004/104202 and the ATCC 9135 strain as template, the Lm. D-LDH gene was cloned by colony PCR (using "KOD-Plus-polymerase" manufactured by Toyobo Co., Ltd.). After purifying the thus obtained PCR-amplified fragment and phosphorylating the end thereof with "T4 Polynucleotide Kinase" (manufactured by Takara Bio Inc.), the fragment was ligated
 25 to the pUC118 vector (which had been digested with a restriction enzyme HincII and the cut surface had been dephosphorylated). The ligation was carried out using "DNA Ligation Kit Ver.2" (manufactured by Takara Bio Inc.). *Escherichia coli*

DH5 α was transformed with the ligation plasmid product and the plasmid DNA was collected to obtain a plasmid into which the Lm. D-LDH gene was subcloned. The thus obtained pUC118 plasmid into which the Lm. D-LDH gene was inserted was digested with restriction enzymes XhoI and NotI, and the resulting DNA fragments were each inserted to the XhoI/NotI cleavage site of the vector pTRS11 for yeast expression. In this manner, Lm. D-LDH gene-expressing plasmid pTRS207 was obtained.

[0100]

Next, the Lm. D-LDH gene was introduced into the PDC1 gene locus on the chromosome of the *Saccharomyces cerevisiae* SW092-2D strain in the same manner as in Example 5, except that the pTRS207 was used as template and the primer set shown in SEQ ID NO:33 was used in place of the primer shown in SEQ ID NO:24 or 26. The thus prepared transformed yeast is hereinafter referred to as "SW092-2D (Δ pdc1::Lm. D-LDH-TRP1)". Then, in the same manner as in Example 6, the D-lactate productivity was evaluated by batch culture. As a result, the yield with respect to saccharide by the SW092-2D (Δ pdc1::Lm. D-LDH-TRP1) was 38%. The result is shown in Table 2 along with the results of Example 6.

[0101]

[Table 2]

Strain name	SW092-2D (Δ pdc1::Lp. D-LDH1-TRP1)	SW092-2D (Δ pdc1::Lp. D-LDH2-TRP1)	SW092-2D (Δ pdc1::Lm. D-LDH-TRP1)
Lactic acid yield with respect to saccharide (%)	45	42	38

[0102]

(Reference Example 1) Preparation of a yeast introduced with the L-LDH gene originated from *Xenopus laevis*

In accordance with the method described in JP 2008-029329A, yeast in which the L-LDH gene originated from *Xenopus laevis* (X. L-LDH gene) was introduced

into the PDC1 gene locus was prepared. Here, as the yeast used for the introduction of the gene into the chromosome, a NBRC10506 strain in which the adenine auxotrophy is reverted (SU013-1D strain) was used. The SU013-1D strain was prepared by the following method. Using the plasmid pRS422 as template and the oligonucleotides (SEQ ID NOs:34 and 35) as primer set, PCR was performed to amplify a PCR fragment of about 2 kb corresponding to the ADE2 gene. After separating the PCR fragment by 1% agarose gel electrophoresis and purifying it in accordance with a conventional method, the PCR fragment was transformed to cancel the mutation in the ADE2 gene. By culturing the yeast strain in an adenine-free medium, a transformant in which the adenine synthesis capacity was reverted was selected.

[0103]

In order to confirm that the thus obtained transformant is a yeast in which the mutation in the AED2 gene was canceled, the following steps were carried out.

First, diploid cells were obtained by mating the thus obtained transformant with the *Saccharomyces cerevisiae* L0GY77 strain having the wild-type ADE2 gene and the diploid cells were cultured in an ascus formation medium to allow ascus formation. The resulting asci were anatomized using a micromanipulator to obtain respective haploid cells (tetrad) and the auxotrophy was examined for each of the haploid cells.

All of the thus obtained haploid cells were confirmed to have adenine synthesis capacity. The thus obtained NBRC10506(ADE2) strain and the NBRC10505 strain were mated and the resulting asci were anatomized using a micromanipulator to obtain the SU013-1D strain (genotype: MAT α ura3 leu2 trp1 his3 ADE2 lys2). The thus obtained transformed yeast is hereinafter referred to as "SU013-1D (Δ pdc1::X. L-LDH-TRP1) strain".

[0104]

(Example 7, Comparative Example 2) D-lactic acid productivity test 3

The SU013-1D (Δ pdc1::X. L-LDH-TRP1) strain prepared in Reference Example 1 was mated with each of the SW092-2D (Δ pdc1::Lp. D-LDH1-TRP1) strain, SW092-2D (Δ pdc1::Lp. D-LDH2-TRP1) strain and SW092-2D (Δ pdc1::Lm. D-LDH-TRP1) strain, which were prepared in Example 5 and Comparative Example 1, thereby preparing diploid yeasts having the L-LDH gene and D-LDH gene heterozygously at the PDC1 gene locus. The thus prepared diploid yeasts are hereinafter referred to as "Lp1-X strain, Lp2-X strain and Lm-X strain, respectively".

[0105]

Then, the Lp1-X strain, Lp2-X strain and Lm-X strain were batch-cultured under the same conditions as in Example 5, and the optical purity of the produced lactic acid was evaluated (Table 3).

[0106]

[Table 3]

Strain name	Lp1-X	Lp2-X	Lm-X
Yield of D-lactic acid (%)	48.3	47.5	43.8
Optical purity of D-lactic acid (%)	52.3	51.7	44.4

[0107]

As a result, the yeast having the D-LDH gene originated from *Limulus polyphemus* produced D-lactic acid with an optical purity of not less than 50%, while the optical purity of the D-lactic acid produced by the yeast having the D-LDH gene originated from *Leuconostoc mesenteroides* was 44.4%. Since the yield with respect to saccharide at which each yeast produces D-lactic acid and the optical purity of thereof are thought to be proportional to the D-LDH activity in the yeasts transformed with the D-LDH gene, from the results shown in Tables 2 and 3, it was confirmed that, when introduced into yeast, the D-LDH gene originated from *Limulus polyphemus* encodes a polypeptide having an activity higher than that of a polypeptide encoded by the D-LDH gene originated from *Leuconostoc mesenteroides*.

[0108]

(Example 8) Introduction of multiple copies of the D-LDH gene into the yeast chromosome and preparation of temperature-sensitive mutant-type PDC5 yeast

Next, preparation of a yeast introduced with the D-LDH gene originated from *Limulus polyphemus* also at gene loci other than the PDC1 gene locus was examined for the purpose of further improving the yield of D-lactic acid, and the D-LDH gene was introduced into the TDH3 gene whose introduction effect is described in JP 2008-029329A and the SED1 gene locus whose introduction effect is described in the international application PCT/JP 2008/072129. Further, the temperature-sensitive mutant-type PDC5 gene described in JP 2008-048726A was also introduced.

[0109]

[Introduction of the D-LDH gene into the TDH3 gene locus]

For introduction into the TDH3 gene locus, in the same manner as the preparation method of the pTRS150 described in JP 2008-029329A, a plasmid pTRS208 in which the TDH3 terminator of the pTRS205 carrying the Lp. D-LDH1 gene was substituted with the ADH1 terminator was prepared. Then, using the SEQ ID NO:36 in place of the SEQ ID NO:8 of JP 2008-029329A as primer and the pTRS208 as template, a DNA construct for homologous recombination to be used for introduction into the TDH3 gene locus was obtained.

[0110]

As the yeast into which the above-described DNA construct for homologous recombination is introduced, a leucine prototrophic strain of the SU013-1D strain (SW087-2C strain) was employed. The method of preparing the SW087-2C strain is described in the followings. Using the plasmid pRS425 as template and the oligonucleotides (SEQ ID NOs:37 and 38) as primer set, PCR was performed to amplify a PCR fragment of about 2 kb corresponding to the LEU2 gene. After separating the PCR fragment by 1% agarose gel electrophoresis and purifying it in accordance with a conventional method, the thus purified PCR fragment was

transformed into the SU013-1D strain to cancel the mutation in the LEU2 gene. By culturing the yeast strain in a leucine-free medium, a transformant in which the leucine synthesis capacity was reverted was selected. In order to confirm that the thus obtained transformant is a yeast in which the mutation in the LEU2 gene was canceled, the following steps were carried out. First, diploid cells were obtained by mating the thus obtained transformant with the *Saccharomyces cerevisiae* L0GY77 strain having the wild-type LEU2 gene and the diploid cells were cultured in an ascus formation medium to allow ascus formation. The resulting asci were anatomized using a micromanipulator to obtain respective haploid cells (tetrad) and the auxotrophy was examined for each of the haploid cells. All of the thus obtained haploid cells were confirmed to have leucine synthesis capacity. The thus obtained SU013-1D(LEU2) strain and the NBRC10505 strain were mated and the resulting asci were anatomized using a micromanipulator to obtain the SW087-2C strain (genotype: MAT α ura3 LEU2 trp1 his3 ADE2 lys2).

[0111]

The SW087-2C strain was transformed with the above-described DNA construct for homologous recombination and cultured in a uracil-free medium for selection to obtain a transformed yeast introduced with the Lp. D-LDH1 gene at the TDH3 gene locus. The thus obtained transformed yeast is hereinafter referred to as "SW087-2C (Δ TDH3::Lp. D-LDH-URA3) strain".

[0112]

[Introduction of the D-LDH gene into SED1 gene locus]

The introduction into the SED1 gene locus was carried out by modifying the method described in Example 2 of the international application PCT/JP 2008/072129.

That is, using the pTRS205 in place of the pTRS102 as PCR template and the SEQ ID NO:39 in place of the SEQ ID NO:14 of the above-described international application publication as primer, a DNA construct for homologous recombination to

be used for introduction into the SED1 gene locus was obtained.

[0113]

As the yeast into which the above-described DNA construct for homologous recombination is introduced, SW092-7D strain (genotype: MAT α ura3 leu2 trp1 his3 ADE2 LYS2), which was obtained by mating the NBRC10505(LYS2) strain
 5 ADE2 LYS2), which was obtained by mating the NBRC10505(LYS2) strain prepared in Example 5 with the NBRC10506(ADE2) strain prepared in Reference Example 1 and separating by anatomization of the resulting asci with a micromanipulator, was employed.

[0114]

10 The SW092-7D strain was transformed with the above-described DNA construct for homologous recombination and cultured in a histidine-free medium for selection to obtain a transformed yeast introduced with the Lp. D-LDH1 gene at the SED1 gene locus. The thus obtained transformed yeast is hereinafter referred to as "SW092-7D (Δ SED1::Lp. D-LDH-HIS3) strain".

15 [0115]

[Introduction of the temperature-sensitive mutant-type PDC5 gene (pdc5ts-9)]

As the yeast introduced with the temperature-sensitive mutant-type PDC5 gene, the yeast SW015 strain having the temperature-sensitive mutant-type PDC5 gene (pdc5ts-9), which is described in JP 2008-048726A, was employed. By
 20 mating the SW015 strain with the SW087-2C strain and separating by anatomization of the resulting asci with a micromanipulator, SW095-4B strain (genotype: MAT α ura3 LEU2 trp1 his3 ADE2 lys2 pdc5ts-9 Δ pdc1::TRP1) was obtained. Further, by mating the SW095-4B strain with the SW092-2D strain and separating by anatomization of the resulting asci with a micromanipulator, SW098-21B strain
 25 (genotype: MAT α ura3 LEU2 trp1 his3 ADE2 LYS2 pdc5ts-9) was obtained.

[0116]

[Introduction of the D-LDH gene into the PDC1, TDH3 and SED1 gene loci and

introduction of the temperature-sensitive mutant-type PDC5 gene (pdc5ts-9)]

By using the thus prepared SW092-2D (Δ PDC1::Lp. D-LDH1-TRP1) strain, SW087-2C (Δ TDH3::Lp. D-LDH1-URA3) strain, SW092-7D (Δ TDH3::Lp. D-LDH1-HIS3) and SW098-21B, diploid mating and tetrad production were repeated to obtain SU042 strain which contains the Lp. D-LDH1 gene at the PDC1, TDH3 and SED1 gene loci and the temperature-sensitive mutant-type PDC5 (pdc5ts-9), the SU042 strain being diploid and not auxotrophic for adenine, leucine and lysine. The procedures for preparing the SU042 strain are shown in Fig. 1.

[0118]

(Example 9) D-lactic acid production test 4

Using the SU042 strain prepared as described in Example 8, the D-lactic acid productivity was tested by batch culture. As the medium, the SC3 medium shown in Table 1 or a raw sugar medium (100 g/L "Yutosei" (manufactured by Muso Co., Ltd.), 1.5 g/L ammonium sulfate) was employed. First, the SU042 strain was cultured overnight with shaking in 5 ml of SC3 medium or raw sugar medium in a test tube (pre-preculture). In a 500 ml Sakaguchi flask, the resulting pre-preculture medium was inoculated to 50 ml of fresh SC3 medium or raw sugar medium and cultured for 24 hours with shaking (preculture). To a mini-jar fermentor (manufactured by Able Co., Ltd., 2 L), 1 L of SC3 medium or raw sugar medium was added, and the temperature was adjusted (32°C) and pH was controlled (pH 5, 5N calcium hydroxide). Culturing was carried out with aeration and stirring (at 200 mL/min and 400 rpm, respectively). As a result, the culture in the SC3 medium was finished in 30 hours and the yield with respect to saccharide was 63%. Further, the culture in the raw sugar medium was finished in 50 hours and the yield with respect to saccharide was 70%. From these results, it was confirmed that the performance of D-lactic acid fermentation was improved by the introduction of the Lp. D-LDH1 gene into the PDC1, TDH3 and SED1 gene loci and the introduction of the

temperature-sensitive mutation into the PDC5 gene.

[0118]

(Example 10) D-lactic acid productivity test by continuous culture

A continuous culture of the SU042 strain using the separation membrane
 5 described in WO2007/097260 was examined. As the medium, a raw sugar medium
 (75 g/L "Yutosei" (manufactured by Muso Co., Ltd.), 1.5 g/L ammonium sulfate) was
 employed.

The culturing conditions are shown below:

Fermentation vessel capacity: 2 (L)

10 Culture medium volume: 1.5 (L)

Separation membrane used: PVDF membrane filter (described in Reference Example
 2 of WO2007/097260)

Effective filtration area of the membrane separation element: 120 cm²

Temperature adjustment: 32 (°C)

15 Aeration volume in the fermentation reaction vessel: air 0.02 (L/min), nitrogen gas
 0.18 (L/min)

Stirring rate in the fermentation reaction vessel: 800 (rpm)

pH adjustment: adjusted to pH 5 with 5N calcium hydroxide

Sterilization: the culture vessel including the separation membrane element and
 20 medium used were all autoclaved at 121°C for 20 minutes.

[0119]

First, the SU042 strain was cultured at 30°C overnight with shaking in 10 ml
 of the raw sugar medium in a test tube (pre-pre-culture). The entire amount of
 the thus obtained culture medium was inoculated to 100 ml of fresh raw sugar
 25 medium and cultured in a 500 ml Sakaguchi flask for 24 hours at 30°C with shaking
 (pre-pre-culture). The resulting pre-pre-culture medium was inoculated to 1.5 L of
 the raw sugar medium loaded in a membrane integrated-type continuous culture

vessel (the apparatus shown in Fig. 2 of WO2007/097260). Removal of the culture medium by a peristaltic pump (200 mL/h) was started 50 hours after the initiation of the culture, and the culture was continued until 400 hours to measure the concentration of lactic acid, which is the substance produced, and the lactic acid production rate. The results are shown in Fig. 2. It is noted here that the lactic acid production rate during the continuous culture was calculated using the Equation 1 below.

[0120]

Lactic acid production rate (g/L·h) = product concentration in removed liquid (g/L) × removal rate of fermentation culture medium (L/hr) ÷ operational liquid volume of apparatus (L) ... (Equation 1)

[0121]

As a result of the continuous culture using the separation membrane, it was confirmed that the D-lactic acid yield with respect to saccharide was further improved to about 85% and that the D-lactic acid production rate was improved to 7.5 g/L·h.

[0122]

(Example 11) Introduction of the *Limulus polyphemus* D-LDH into *Candida utilis*

Next, using *Candida utilis*, which is a Crabtree-negative yeast, D-lactic acid was examined.

[0123]

(a) Preparation of a vector for introduction of the *Limulus polyphemus* D-LDH gene

First, a vector was constructed for introducing the D-LDH gene originated from *Limulus polyphemus* into the PDC1 gene locus on the *Candida utilis* chromosome. The *Candida utilis* NBRC0988 strain (hereinafter, referred to as "NBRC0988") was inoculated to YPD medium (1% Bacto Yeast Extract, 2% Bacto-peptone, 2% glucose) and cultured overnight at 30°C. From the thus obtained yeast

cells, genomic DNA was extracted in accordance with a conventional method and used as template in the subsequent PCR. First, using the oligo DNAs shown in SEQ ID NOs:40 and 41 as primer set, a fragment containing the terminator region of the PDC1 gene was amplified. It is noted here that, unless otherwise specified, KOD-plus- (manufactured by Toyobo Co., Ltd.) was used as DNA polymerase in accordance with the attached protocol. After digesting the thus obtained fragment of about 1kb with a restriction enzyme BssHI and purifying it, the resultant was ligated to pBluescriptIISK(+) which had been previously digested with restriction enzyme BssHI in the same manner. The thus obtained plasmid is hereinafter referred to as "pKS01".

[0124]

Next, in the same manner, using the genomic DNA of NBRC0988 as template and the oligo DNAs shown in SEQ ID NOs:42 and 43 as primer set, PCR was performed to amplify a fragment containing the PGK promoter. The thus obtained fragment of about 1 kb was purified and used in the subsequent experiments. Then, using the plasmid pLC1-hph having the hph gene capable of imparting resistance against hygromycin B which is a type of antibiotic substance as template and the oligo DNAs shown in SEQ ID NOs:44 and 45 as primer set, PCR was performed to amplify a fragment containing the hph gene, which was then purified to obtain a fragment of about 1.1 kb. Further, using the genomic DNA of NBRC0988 as template and the oligo DNAs shown in SEQ ID NOs:46 and 47 as primer set, PCR was performed to amplify a fragment containing the GAP terminator, which was then purified to obtain a fragment of about 500 bp.

[0125]

The fragment containing the PGK promoter, the fragment containing the hph gene and the fragment containing the GAP terminator that were obtained in the above-described manner were mixed, and using the oligo DNAs shown in SEQ ID

NOs:48 and 49 as primer set, PCR was performed to amplify a fragment in which the PGK promoter, hph gene and GAP terminator were linked. After phosphorylating the end of the thus obtained fragment of about 2.6 kb, the fragment was ligated to the pUC118 which had been digested with HincII and dephosphorylated. The thus
5 obtained plasmid is hereinafter referred to as "pKS02".

[0126]

Next, in the same manner, using the genomic DNA of NBRC0988 as template and the oligo DNAs shown in SEQ ID NOs:50 and 51 as primer set, PCR was performed to amplify a fragment containing the PGK terminator, which was then
10 purified to obtain a fragment of about 500 bp. Further, using the pKS02 obtained in the above-described manner as template and the oligo DNAs shown in SEQ ID NOs:52 and 53 as primer set, PCR was performed to amplify a fragment in which the PGK promoter, hph gene and GAP terminator were linked, which was then purified to obtain a fragment of about 2.6 kb.

15 [0127]

The fragment containing the PGK terminator and the fragment in which the PGK promoter, hph gene and GAP terminator were linked, which fragments were obtained in the above-described manner, were mixed, and using the oligo DNAs shown in SEQ ID NOs:50 and 53 as primer set, PCR was performed to amplify a
20 fragment in which the PGK terminator, PGK promoter, hph gene and GAP terminator were linked. After digesting the thus obtained fragment of about 3 kb with restriction enzymes BamHI and ClaI and purifying it, the resultant was ligated to the pKS02 which had been previously digested with restriction enzymes BamHI and ClaI in the same manner. The thus obtained plasmid is hereinafter referred to as
25 "pKS03".

[0128]

Next, in the same manner, using the genomic DNA of NBRC0988 as template

and the oligo DNAs shown in SEQ ID NOs:54 and 55 as primer set, PCR was performed to amplify a fragment containing the PDC1 promoter region, which was then purified to obtain a fragment of about 2.1 kb. Further, using the pTRS205 prepared in Example 2 as template and the oligo DNAs shown in SEQ ID NOs:56 and 57 as primer set, PCR was performed to amplify a fragment containing the D-LDH gene originated from *Limulus polyphemus*, which was then purified to obtain a fragment of about 1 kb.

[0129]

The fragment containing the PDC1 promoter region and the fragment containing the D-LDH gene originated from *Limulus polyphemus* that were obtained in the above-described manner were mixed, and using the oligo DNAs shown in SEQ ID NOs:54 and 57 as primer set, PCR was performed to amplify a fragment in which the fragment containing the PDC1 promoter region and the fragment containing the D-LDH gene originated from *Limulus polyphemus* were linked. After digesting the thus obtained fragment of about 3.1 kb with restriction enzymes NotI and BglII and purifying it, the resultant was ligated to the pKS03 which had been previously digested with restriction enzymes NotI and BamHI in the same manner. The thus obtained plasmid is hereinafter referred to as "pKS04".

[0130]

(b) Introduction of the *Limulus polyphemus* D-LDH gene into the PDC1 gene locus

The pKS04 obtained in the above-described manner was digested with restriction enzyme BglII and introduced into NBRC0988 by the electroporation method. A small amount of the cells was inoculated to YPD medium and cultured overnight at 30°C. After centrifuging the culture to discard the supernatant, the cells were washed with sterilized water and 1M sorbitol and eventually suspended in 1M sorbitol. Then, after mixing the cells and the pKS04 digested with restriction enzyme BglII and leaving the resulting mixture on ice for 5 minutes, the mixture was

transferred to an electroporation cuvette and subjected to electroporation at a capacitance of 25 μ F, voltage of 0.75 kV and resistance of 800 Ω . Thereafter, the mixture was transferred to YPD medium containing 1M sorbitol and cultured at 30°C for about 4 hours, and then applied to YPD medium supplemented with 600 μ g/L of hygromycin B. From the resulting hygromycin B-resistant strain, the genome thereof was extracted, and by verifying that fragments of about 3 kb and 2 kb were each amplified by PCR using the oligo DNAs shown in SEQ ID NOs:54 and 57 and SEQ ID NOs:56 and 51 as respective primer sets, it was confirmed that a transformed yeast in which the D-LDH gene originated from *Limulus polyphemus* was introduced into the PDC1 gene locus was obtained. The thus obtained transformed yeast is hereinafter referred to as "CuLpLDH strain".

[0131]

(Comparative Example 2) Introduction of D-LDH originated from lactic acid bacterium into *Candida utilis*

In the same manner as in Example 11, the D-LDH gene originated from a lactic acid bacterium, *Leuconostoc mesenteroides*, was introduced. It is noted here, however, that the oligo DNA shown in SEQ ID NO:58 was employed in place of the one shown in SEQ ID NO:55 used in Example 11. In addition, the pTRS207 obtained in Comparative Example 1 was used as template in place of the pTRS205 obtained in Example 2 and the oligo DNAs shown in SEQ ID NOs:59 and 60 were used in place of those shown in SEQ ID NOs:56 and 57. The thus obtained plasmid is hereinafter referred to as "pKS05".

[0132]

The plasmid obtained in the above-described manner was introduced into the NBRC0988 strain in the same manner as in Example 11. The thus obtained strain in which the D-LDH gene originated from a lactic acid bacterium, *Leuconostoc mesenteroides*, was introduced into the PDC1 gene locus is hereinafter referred to as

"CuLmLDH strain".

[0133]

(Example 12, Comparative Example 3) D-lactic productivity test in *Candida utilis*

Using the CuLpLDH strain and the CuLmLDH strain prepared in Example 11
5 and Comparative Example 2, respectively, their D-lactic acid productivity was
evaluated. YPD medium was added in an amount of 50 ml to a 500 ml Sakaguchi
flask, and the CuLpLDH and CuLmLDH strains were inoculated thereto in small
amounts and cultured at 30°C overnight with shaking (preculture). After collecting
the culture medium and washing it with fresh YPD medium, the culture medium was
10 placed and cultured in a mini-jar fermentor in which 1 L of YPD10 medium
(containing 10% glucose) was added. The culturing conditions are shown below:

Amount of initial inoculation: inoculated at $OD_{600} = 10$

pH: pH6

Aeration: 100 mL/min

15 Stirring: 120 rpm

Neutralizer: 1N calcium hydroxide

Culturing temperature: 35°C.

[0134]

The culture was carried out for 40 hours and the lactic acid concentration and
20 glucose concentration of the culture medium at 40 hours were analyzed. Glucose
had been completely consumed. In addition, as shown in Table 4, from the analysis
results of the productivity per glucose consumption (yield with respect to saccharide)
and the optical purity of D-lactic acid, the yeast (*Candida utilis*) introduced with the
D-LDH gene was found to be capable of fermentatively producing D-lactic acid and
25 the CuLpLDH strain introduced with the D-LDH gene originated from *Limulus*
polyphemus had a higher yield.

[0135]

(Example 13, Comparative Example 4) D-lactic productivity test 2 in *Candida utilis*

Next, using the CuLpLDH and CuLmLDH strains, their D-lactic acid productivity using xylose, which is a pentose, as sugar source was evaluated. YPD medium was added in an amount of 50 ml to a 500 ml Sakaguchi flask, and the CuLpLDH and CuLmLDH strains were inoculated thereto in small amounts and cultured at 30°C overnight with shaking (preculture). After collecting the culture medium and washing it with fresh YPD medium, the culture medium was placed and cultured in a mini-jar fermentor in which 1 L of YPX10 medium (1% yeast extract, 2% Bacto-peptone, 4% xylose) was added. The culturing conditions are shown below:

Amount of initial inoculation: inoculated at $OD_{600} = 10$

pH: pH6

Aeration: 100 mL/min

Stirring: 120 rpm

Neutralizer: 1N calcium hydroxide

Culturing temperature: 30°C.

[0136]

The culture was carried out for 60 hours and the lactic acid concentration and xylose concentration of the culture medium at 60 hours were analyzed. Xylose had been completely consumed. In addition, as shown in Table 4, from the analysis results of the productivity per xylose consumption (yield with respect to saccharide) and the optical purity of D-lactic acid, the yeast (*Candida utilis*) introduced with the D-LDH gene was found to be capable of fermentatively producing D-lactic acid using xylose as starting material and the CuLpLDH strain introduced with the D-LDH gene originated from *Limulus polyphemus* had a higher yield.

[0137]

[Table 4]

	Example 12	Example 13	Comparative Example 3	Comparative Example 4
Strain name	CuLpLDH	CuLpLDH	CuLmLDH	CuLmLDH
Sugar source	glucose	xylose	glucose	xylose
D-lactic acid yield with respect to saccharide (%)	29	23	22	17
Optical purity of D-lactic acid (% e.e.)	99.9	99.9	99.9	99.9

INDUSTRIAL APPLICABILITY

[0138]

By the present invention, it becomes able to stably produce D-lactic acid at a low cost, and since the produced D-lactic acid has a high optical purity, it is preferably used as a raw material for polymers.

CLAIMS

1. A polypeptide which is any one of the following (A) to (C):
 - (A) a polypeptide having the amino acid sequence shown in SEQ ID NO:1 or 2;
 - (B) a polypeptide having the same amino acid sequence as shown in SEQ ID NO:1 or
5 2 except that one or several amino acids are substituted, deleted, inserted and/or
added, which polypeptide has a D-lactate dehydrogenase activity; and
 - (C) a polypeptide having an amino acid sequence which has a sequence identity of
not less than 80% to the amino acid sequence shown in SEQ ID NO:1 or 2, which
polypeptide has a D-lactate dehydrogenase activity.
- 10 2. A polynucleotide which is any one of the following (a) to (e):
 - (a) a polynucleotide having the nucleotide sequence shown in SEQ ID NO:3 or 4;
 - (b) a polynucleotide having the same nucleotide sequence as shown in SEQ ID NO:3
or 4 except that one or several nucleotides are substituted, deleted, inserted and/or
added, which polynucleotide encodes a polypeptide having a D-lactate
15 dehydrogenase activity;
 - (c) a polynucleotide which hybridizes under stringent conditions with the entirety or a
part of the polynucleotide having the nucleotide sequence shown in SEQ ID NO:3 or
4 or a complementary strand thereof, which polynucleotide encodes a polypeptide
having a D-LDH activity;
 - 20 (d) a polynucleotide having a nucleotide sequence which has a sequence identity of
not less than 80% to the nucleotide sequence shown in SEQ ID NO:3 or 4, which
polynucleotide encodes a polypeptide having a D-lactate dehydrogenase activity; and
 - (e) a polynucleotide encoding the polypeptide according to claim 1.
3. A DNA construct in which the polynucleotide according to claim 2 and a
25 promoter capable of expressing said polynucleotide are linked.
4. The DNA construct according to claim 3, characterized in that said promoter
is a promoter of pyruvate decarboxylase 1 gene (PDC1 gene), suppression-of-

exponential-defect 1 gene (SED1 gene) or glyceraldehyde-3-phosphate dehydrogenase 3 gene (TDH3 gene).

5. The DNA construct according to claim 3 or 4, wherein said promoter is selected from any one of the following (I) to (III):

5 (I) a promoter having the nucleotide sequence shown in any one of SEQ ID NOs:5 to 7;

(II) a promoter having a nucleotide sequence which hybridizes under stringent conditions with the nucleotide sequence shown in any one of SEQ ID NOs:5 to 7 or a nucleotide sequence comprising a part thereof; and

10 (III) a promoter having the same nucleotide sequence as shown in any one of SEQ ID NOs:5 to 7 except that one or several nucleotides are deleted, substituted and/or added.

6. A transformant in which the polynucleotide according to claim 2 or the DNA construct according to any one of claims 3 to 5 is introduced.

15 7. A transformed yeast in which the polynucleotide according to claim 2 or the DNA construct according to any one of claims 3 to 5 is introduced.

8. The transformed yeast according to claim 7, wherein at least one of pyruvate decarboxylase 1 gene (PDC1 gene), suppression-of-exponential-defect 1 gene (SED1 gene) and glyceraldehyde-3-phosphate dehydrogenase 3 gene (TDH3 gene) of said
20 transformed yeast is substituted with the polynucleotide according to claim 2 or the DNA construct according to any one of claims 3 to 5.

9. The transformed yeast according to claim 8, wherein at least one of said genes is PDC1 gene.

10. A method of producing D-lactic acid, which comprises the step of culturing
25 the transformant according to claim 6 or the transformed yeast according to any one of claims 7 to 9.

11. A transformant in which a polynucleotide encoding D-lactate dehydrogenase

originated from the family Limulidae or a DNA construct in which said polynucleotide and a promoter capable of expressing said polynucleotide are linked is introduced.

12. A transformed yeast in which a polynucleotide encoding D-lactate dehydrogenase originated from the genus *Limulus* or a DNA construct in which said polynucleotide and a promoter capable of expressing said polynucleotide are linked is introduced.

13. The transformed yeast according to claim 12, wherein at least one of pyruvate decarboxylase 1 gene (PDC1 gene), suppression-of-exponential-defect 1 gene (SED1 gene) and glyceraldehyde-3-phosphate dehydrogenase 3 gene (TDH3 gene) of said transformed yeast is substituted with said polynucleotide encoding said D-lactate dehydrogenase originated from the family Limulidae or said DNA construct in which said polynucleotide and a promoter capable of expressing said polynucleotide are linked is introduced.

14. A method of producing D-lactic acid, which comprises the step of culturing the transformant according to claim 11 or the transformed yeast according to claim 12 or 13.

10029W001 sequence listing.txt
SEQUENCE LISTING

<110> TORAY INDUSTRIES, INC.

<120> Polypeptide having D-lactate dehydrogenase activity, polynucleotide encoding the same polypeptide and method for producing D-lactate

<130> 10029

<160> 60

<170> PatentIn version 3.1

<210> 1

<211> 326

<212> PRT

<213> Limulus polyphemus

<400> 1

Met Ser Lys Pro Lys Val Phe Val Thr Arg Pro Asp Val Pro Gln Ala
1 5 10 15

Gly Ile Asp Leu Leu Lys Glu Lys Cys Asp Val Glu Ile Tyr Asp Gln
20 25 30

Pro Met Pro Ile Pro Arg Asp Ala Leu Ile Lys Gly Val Gln Gly Lys
35 40 45

Asp Ala Leu Tyr Cys Leu Leu Thr Asp Lys Ile Asp Lys Asp Val Met
50 55 60

Asp Ala Ala Gly Pro Gln Leu Lys Val Ile Ala Thr Met Ser Val Gly
65 70 75 80

Phe Asp His Ile Asp Leu Asn Glu Cys Lys Ala Arg Asn Ile Ala Val
85 90 95

Ser Asn Thr Pro Asp Val Ser Thr Asp Ser Val Ala Glu Leu Thr Val
100 105 110

Thr Leu Leu Leu Val Cys Gly Arg Arg Ile Met Asp Ser Ala Asn Ala
115 120 125

Ile Lys Asn Gly Glu Trp Ile Tyr Ser Trp Ser Pro Leu Trp Leu Cys
130 135 140

Gly Arg Gly Leu Thr Asn Ser Thr Ile Gly Ile Val Gly Met Gly Arg
145 150 155 160

Ile Gly Gln Ala Val Met Lys Arg Leu Leu Pro Phe Gly Val Lys Lys
165 170 175

Val Leu Tyr Tyr Asp Leu Phe His Pro Ile Lys Pro Ala Glu Asp Met
180 185 190

10029w001 sequence listing.txt

Gly Ala Gln Tyr Val Glu Phe Glu Glu Leu Leu Lys Glu Ser Asp Phe
195 200 205

Val Val Ala Met Cys Asn Leu Ser Glu Gln Thr Lys Glu Leu Phe Asn
210 215 220

Ala Lys Ala Phe Ser Gln Met Lys Pro Thr Ala Val Phe Val Asn Thr
225 230 235 240

Ser Arg Gly Gly Val Val Gln Gln Asp Asp Leu Tyr Glu Ala Leu Lys
245 250 255

Asn Gly Lys Ile Arg Ala Ala Gly Leu Asp Val Met Ile Pro Glu Pro
260 265 270

Leu Pro Arg Asp His Lys Leu Thr Thr Leu Pro Asn Ile Val Leu Leu
275 280 285

Pro His Val Gly Ser Ala Glu Glu Ala Ala Arg Ile Glu Met Ala Thr
290 295 300

Leu Ala Ala Lys Asn Ile Leu Ser Val Leu Asp Gly Lys Pro Leu Val
305 310 315 320

Thr Pro Val Pro Met Pro
325

<210> 2
<211> 326
<212> PRT
<213> Limulus polyphemus

<400> 2

Met Ser Lys Pro Lys Val Phe Val Thr Arg Pro Asp Val Pro Gln Ala
1 5 10 15

Gly Ile Asp Leu Leu Lys Glu Lys Cys Asp Val Glu Ile Tyr Asp Gln
20 25 30

Pro Met Pro Ile Pro His Asp Ala Leu Ile Lys Gly Val Gln Gly Lys
35 40 45

Asp Ala Leu Tyr Cys Leu Leu Thr Asp Lys Ile Asp Lys Asp Val Met
50 55 60

Asp Ala Ala Gly Pro Gln Leu Lys Val Ile Ala Thr Met Ser Val Gly
65 70 75 80

Tyr Asp His Ile Asp Leu Asn Glu Cys Lys Ala Arg Asn Ile Val Val
85 90 95

Ser Asn Thr Pro Asp Val Ser Thr Asp Ser Val Ala Glu Leu Thr Val
Page 2

10029w001 sequence listing.txt

100

105

110

Thr Leu Leu Leu Val Val Gly Arg Arg Ile Phe Asp Ser Ala Cys Ala
115 120 125

Ile Lys Asn Gly Glu Trp Ile Tyr Ser Trp Ser Pro Leu Trp Leu Cys
130 135 140

Gly Arg Gly Leu Thr Asn Ser Thr Val Gly Ile Val Gly Met Gly Arg
145 150 155 160

Ile Gly Gln Ala Val Met Lys Arg Leu Leu Pro Phe Gly Val Lys Lys
165 170 175

Ile Leu Tyr Phe Asp Leu Phe His Pro Ile Lys Pro Ala Glu Asp Met
180 185 190

Gly Ala Gln Phe Val Glu Phe Glu Glu Leu Leu Lys Glu Ser Asp Phe
195 200 205

Val Val Ala Met Cys Asn Leu Ser Glu Glu Thr Lys Glu Ile Phe Asn
210 215 220

Ala Lys Ala Phe Ser Leu Met Lys Pro Thr Ala Val Phe Ile Asn Thr
225 230 235 240

Ser Arg Gly Gly Val Val Gln Gln Asp Asp Leu Tyr Glu Ala Leu Lys
245 250 255

Asn Gly Val Ile Arg Gly Ala Gly Leu Asp Val Met Val Pro Glu Pro
260 265 270

Leu Pro Arg Asp His Lys Leu Thr Thr Leu Pro Asn Ile Ile Leu Leu
275 280 285

Pro His Val Gly Ser Ala Glu Glu Ala Ala Arg Thr Glu Met Ala Thr
290 295 300

Leu Ala Ala Lys Asn Ile Leu Ser Val Leu Asp Gly Lys Pro Leu Leu
305 310 315 320

Thr Pro Val Gln Met Pro
325

<210> 3
<211> 981
<212> DNA
<213> Limulus polyphemus

<400> 3
atgagcaaac caaaggtatt tgtaaccaga ccagacgttc ctcaagcagg aattgacctt 60
ttaaaggaaa aatgtgatgt agaaatttac gaccagccga tgccgattcc acgagatgcc 120
Page 3

10029W001 sequence listing.txt

ttaattaaag gagttcaggg taaggatgct ctttattgtc ttctcacgga caaaattgac	180
aaggatgtga tggatgcagc aggacccag ctaaaggtaa tagcgaccat gtctgttgga	240
tttgaccata ttgaccttaa cgaatgtaag gcaagaaata ttgcggtcag caaactccc	300
gatgtttcca ctgactccgt agctgagctg actgtcacac tattgttggt ttgtggcaga	360
agaatcatgg attctgcaaa cgctattaaa aatggagagt ggatctactc ttggagtcca	420
ctgtggctat gtggtagagg attaaccaac agcacaatag gaattgtagg aatgggtaga	480
attggccaag ctgtcatgaa acgcctgttg cctttcggtg taaagaagggt tttgtattat	540
gatctatttc acccaatcaa acccgctgaa gacatgggag ctgagtatgt tgagtttgaa	600
gagttattga aagagtcaga ctttgtagtc gcaatgtgca atttatcaga acaaaccaag	660
gaattgttta atgccaaggc ttccagccaa atgaagccta cagcagtttt cgtaataact	720
agtcgcggag gagttgttca acaagatgac ctctatgagg ccctcaagaa tggcaagatc	780
cgtgctgcag ggcttgacgt aatgattcct gagcctctac cccgggatca caagcttact	840
actttacca acatagttct tcttcacat gttggcagtg ctgaggaagc agcaagaata	900
gaaatggcaa ccttggcagc taagaacatc ttgtccgttt tagatggaaa gcctcttgta	960
actcctgttc ccatgccata a	981

<210> 4
 <211> 981
 <212> DNA
 <213> *Limulus polyphemus*

<400> 4	
atgagcaagc ccaaagtatt tgctactaga cctgatgttc ctgaggcagg aattgatctt	60
ttgaaagaga aatgcgatgt cgaaatctat gatcagccta tgccaattcc acatgatgct	120
cttattaaag gagtgcaggg taaggacgct ctttattgtc tcctcaccga taagattgat	180
aaagatgtta tggatgcagc tgggcctcag ctaaaagtta tagctaccat gtctgttgga	240
tatgaccata ttgaccttaa tgaatgtaag gcaagaaata tagttgtaag taacactcct	300
gacgtatcca ctgactcagt agcagaactt actgttacac ttttattgggt tgttggtaga	360
agaatttttg attctgcctg tgccattaaa aatggagagt ggatttactc atggagtcct	420
ttgtggttgt gtggtcgagg attgaccaac agcaccgtag gaattgttggt tatgggaaga	480
attggtcaag cagtcatgaa gagacttcta ccgtttggag ttaagaaaat ttatacttt	540
gacctgtttc atccaatcaa gcctgcagaa gatatgggag ctgagtttgt tgaatttgaa	600
gagctgttga aagaatctga ttttgtagtt gccatgtgta atctgtctga ggaaacaaag	660
gagatattta acgccaaggc ttccagtcta atgaagccta ctgcagtttt tataaatacc	720
agccgtgggtg gagttgtgca gcaggatgac ttgtatgaag cccttaagaa tggcgtgatc	780
cgtggggccg gactggacgt gatggttcct gagcctttac cccgggacca caaactcaca	840
acactaccaa acataattct tctccgcgat gttggcagtg ctgaggaagc agcaagaaca	900

10029W001 sequence listing.txt

gaaatggcaa ctctggcagc caaaaacatc ctttccgttt tagatgggaa gccacttttg 960
actccagttc aaatgccata a 981

<210> 5
<211> 1000
<212> DNA
<213> *Saccharomyces cerevisiae*

<400> 5
aaaatgaagg ccaaatacaag gcgggaaggg acaaccagga cgtaaagggt agcctcccca 60
taacataaac tcaataaaat atatagtctt caacttgaaa aaggaacaag ctcatgcaaa 120
gaggtggtac ccgcacgccg aaatgcatgc aagtaaccta ttcaaagtaa tatctcatal 180
atgtttcatg agggtaacaa catgcgactg ggtgagcata tgttccgctg atgtgatgtg 240
caagataaac aagcaaggca gaaactaact tcttcttcat gtaataaaca cccccgcgt 300
ttatttacct atctctaaac ttcaacacct tatatcataa ctaatatattc ttgagataag 360
cacactgcac ccataccttc cttaaaaacg tagcttccag tttttggtgg ttccggcttc 420
cttcccgaatt ccgcccgcga aacgcatatt tttgttgcct ggtggcattt gcaaaatgca 480
taacctatgc atttaaaaga ttatgtatgc tcttctgact tttcgtgtga tgaggctcgt 540
ggaaaaaatg aataatttat gaatttgaga acaattttgt gttgttacgg tattttacta 600
tggaataatc aatcaattga ggattttatg caaatatcgt ttgaatattt ttccgaccct 660
ttgagtactt ttcttcataa ttgcataata ttgtccgctg cccctttttc tgttagacgg 720
tgtcttgatc tacttgctat cgttcaacac cactttattt tctaactatt ttttttttag 780
ctcatttgaa tcagcttatg gtgatggcac atttttgcat aaacctagct gtcctcgttg 840
aacataggaa aaaaaatat ataaacaagg ctctttcact ctccttgcaa tcagatttgg 900
gtttgttccc tttattttca tttttcttgt catattcctt tctcaattat tattttctac 960
tcataacctc acgcaaaata acacagtcaa atcaatcaaa 1000

<210> 6
<211> 1000
<212> DNA
<213> *Saccharomyces cerevisiae*

<400> 6
aggattttta tctgttggag ttaagggtgaa tacgtttttc catattgggg tatgcagctc 60
gaacctaaag tggatatgtac acatccccctc aagcacaccc attaccctta taggattaat 120
gtaagcaaca gcttacacgg aattggaaat actattcaac gatccatgca tctgccagat 180
tcggacatgc atattcccca attggatata gaaaattaac gtaaggcagt atcttttcac 240
aatgtacttg caacgcggcg acttaaagtt gaagtacaac ctgcagcagc ggctttttgt 300
acggtacgcc aaactgtcaa tggataatat tgcgtagacc gaaaaaggta atcctcaaca 360
ctacccgtgg tggatgacct aaagcagtaa tattggttgg aattatctcc cagacggcac 420
cgtctccccg agaaagctta gccccgaggt ctaccttcca tacaccactg attgctccac 480

10029w001 sequence listing.txt

gtcatgcggc cttctttcga ggacaaaaag gcatatatcg ctaaaattag ccatcagaac	540
cgttattgtt atttatatttt cattacgaaa gaggagaggg cccagcgcg cagagcacac	600
acggtcattg attactttat ttggctaaag atccatccct tctcgatgtc atctctttcc	660
attcttgtgt atttttgatt gaaaatgatt ttttgtccac taatttctaa aaataagaca	720
aaaagccttt aagcagtttt tcatccattt tactacggta aaatgaatta gtacggtatg	780
gctcccagtc gcattatttt tagattggcc gtaggggctg gggtagaact agagtaagga	840
acattgtctt gccctctttt gaactgtcat ataaatacct gacctatttt attctccatt	900
atcgtattat ctcacctctc tttttctatt ctcttgtaat tattgattta tagtcgtaac	960
tacaaagaca agcaaaataa aatacgttcg ctctattaag	1000

<210> 7
 <211> 1000
 <212> DNA
 <213> Saccharomyces cerevisiae

<400> 7 ctattttcga ggaccttgtc accttgagcc caagagagcc aagatttaaa ttttcctatg	60
acttgatgca aattcccaaa gctaataaca tgcaagacac gtacgggtcaa gaagacatat	120
ttgacctctt aacagggttca gacgcgactg cctcatcagt aagacccgtt gaaaagaact	180
tacctgaaaa aaacgaatat atactagcgt tgaatgtag cgtcaacaac aagaagtta	240
atgacgcgga ggccaaggca aaaagattcc ttgattacgt aaggaggatta gaatcatttt	300
gaataaaaaa cacgcttttt cagttcgagt ttatcattat caatactgcc atttcaaaga	360
atacgtaaat aattaatagt agtgattttc ctaactttat ttagtcaaaa aattagcctt	420
ttaattctgc tgtaacccgt acatgcccac aatagggggc gggttacaca gaatatataa	480
catcgtaggt gtctgggtga acagtttatt cctggcatcc actaaatata atggagcccg	540
ctttttaagc tggcatccag aaaaaaaaaa aatcccagca caaaatatt gttttcttca	600
ccaaccatca gttcataggt ccattctctt agcgcaacta cagagaacag gggcacaac	660
aggcaaaaaa cgggcacaac ctcaatggag tgatgcaacc tgcctggagt aaatgatgac	720
acaaggcaat tgaccacgc atgtatctat ctattttct tacaccttct attaccttct	780
gctctctctg atttggaaaa agctgaaaaa aaagggtgaa accagttccc tgaaattatt	840
cccctacttg actaataagt atataaagac ggtaggtatt gattgtaatt ctgtaaatct	900
atttcttaaa cttcttaaat tctactttta tagttagtct ttttttagt tttaaaacac	960
caagaactta gtttcgaata aacacacata aacaaacaaa	1000

<210> 8
 <211> 20
 <212> DNA
 <213> Artificial

<220>
 <223> primer

<400> 8		
atcdchacya tstcbgtggg		20
<210> 9		
<211> 17		
<212> DNA		
<213> Artificial		
<220>		
<223> primer		
<400> 9		
ggytcdggva ccatyac		17
<210> 10		
<211> 27		
<212> DNA		
<213> Artificial		
<220>		
<223> primer		
<400> 10		
ttgccattct tgagggcctc atagagg		27
<210> 11		
<211> 24		
<212> DNA		
<213> Artificial		
<220>		
<223> primer		
<400> 11		
gctgcacaac tccaccacgg ctgg		24
<210> 12		
<211> 24		
<212> DNA		
<213> Artificial		
<220>		
<223> primer		
<400> 12		
tcagctcagc tacggagtca gtgg		24
<210> 13		
<211> 24		
<212> DNA		
<213> Artificial		
<220>		
<223> primer		
<400> 13		
actattgttg gtttgtggca gaag		24
<210> 14		
<211> 24		
<212> DNA		
<213> Artificial		

10029w001 sequence listing.txt

```

<220>
<223> primer

<400> 14
taagttctgc tactgagtca gtgg
24

<210> 15
<211> 24
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 15
tagaagaatt ttgattctg cctg
24

<210> 16
<211> 24
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 16
aacatcgagg gtgttgctga ccgc
24

<210> 17
<211> 26
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 17
agagtggatc tactcttgga gtccac
26

<210> 18
<211> 24
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 18
tacgtcagga gtgttactta caac
24

<210> 19
<211> 26
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 19
agagtggatc tactcttgga gtccac
26

```

10029w001 sequence listing.txt

```

<210> 20
<211> 30
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 20
atgcctcgag atgagcaaac caaaggtatt 30

<210> 21
<211> 30
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 21
atgcgcggcc gcttatggca tgggaacagg 30

<210> 22
<211> 30
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 22
atgcctcgag atgagcaagc ccaaagtatt 30

<210> 23
<211> 30
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 23
atgcgcggcc gcttatggca tttgaactgg 30

<210> 24
<211> 75
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 24
attattattt tctactcata acctcacgca aaataacaca gtcaaatcaa tcaaaatgag 60
caaaccaaag gtatt 75

<210> 25
<211> 20
<212> DNA
<213> Artificial

<220>

```

10029w001 sequence listing.txt

<223> primer

<400> 25
aggcgtatca cgaggccctt 20

<210> 26
<211> 75
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 26
attattattt tctactcata acctcacgca aaataacaca gtcaaataca tcaaaatgag 60
caagcccaaa gtatt 75

<210> 27
<211> 60
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 27
gaattaattc ttgaagacga aagggcctcg tgatacgctt agattgtact gagagtgcac 60

<210> 28
<211> 80
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 28
tatttttctg tacataaaaa tgcttataaa actttaacta ataattagag attaaatcgc 60
ctgtgcggta tttcacaccg 80

<210> 29
<211> 24
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 29
gacaattctg gttagggtcca agag 24

<210> 30
<211> 24
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 30
ttaagctgct gcggagcttc cacg 24

10029w001 sequence listing.txt

<210>	31	
<211>	29	
<212>	DNA	
<213>	Artificial	
<220>		
<223>	primer	
<400>	31	
	ccctcgagat gaagattttt gcttacggc	29
<210>	32	
<211>	31	
<212>	DNA	
<213>	Artificial	
<220>		
<223>	primer	
<400>	32	
	ccgcggccgc ttaatattca acagcaatag c	31
<210>	33	
<211>	85	
<212>	DNA	
<213>	Artificial	
<220>		
<223>	primer	
<400>	33	
	tctcaattat tattttctac tcataacctc acgcaaaata acacagtcaa atcaatcaaa	60
	atgaagattt ttgcttacgg cattc	85
<210>	34	
<211>	22	
<212>	DNA	
<213>	Artificial	
<220>		
<223>	primer	
<400>	34	
	atggattcta gaacagttgg ta	22
<210>	35	
<211>	22	
<212>	DNA	
<213>	Artificial	
<220>		
<223>	primer	
<400>	35	
	ttacttgttt tctagataag ct	22
<210>	36	
<211>	75	
<212>	DNA	
<213>	Artificial	

10029W001 sequence listing.txt

```

<220>
<223> primer

<400> 36
ttagtttttaa aacaccaaga acttagtttc gaataaacac acataaacia acaaaatgag      60
caaaccaaag gtatt                                                              75

<210> 37
<211> 22
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 37
atgtctgccc ctaagaagat cg                                                  22

<210> 38
<211> 24
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 38
ttaagcaagg attttcttaa cttc                                              24

<210> 39
<211> 75
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 39
atttatagtc gtaactacaa agacaagcaa aataaaatac gttcgctcta ttaagatgag      60
caaaccaaag gtatt                                                              75

<210> 40
<211> 68
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 40
actcgcgcg c aagatctaag cggccgctaa tggatccaat aatcgatgct gtctttcttc      60
ttcatggg                                                                      68

<210> 41
<211> 37
<212> DNA
<213> Artificial

<220>

```

<223> primer

<400> 41
actcgcgcgc aagatctgaa cttctccaac aggtagc 37

<210> 42
<211> 99
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 42
cggccgccag ctgaagcttc gtacgtgca ggtcgacaac ccttaataata acttcgtata 60
atgtatgcta tacgaagtta tcctttgctg tgttctacc 99

<210> 43
<211> 68
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 43
tcgatcagaa acttctcgac agacgtcgcg gtgagttcag gctttttcat ctttatccgc 60
cagtatgt 68

<210> 44
<211> 50
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 44
atgaaaaagc ctgaactcac cgcgacgtct gtcgagaagt ttctgatcga 50

<210> 45
<211> 69
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 45
catgaggatc ataatttata acgtaatccc ataaataaaa gtcatacaat ctattccttt 60
gccctcgga 69

<210> 46
<211> 50
<212> DNA
<213> Artificial

<220>
<223> primer

10029w001 sequence listing.txt

<400> 46
attgtatgac ttttatttat gggattacgt tataaattat gatcctcatg 50

<210> 47
<211> 83
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 47
taggccacta gtggatctga tatcacctaa taacttcgta tagcatacat tatacgaagt 60
tattcattca tccctcacta tcg 83

<210> 48
<211> 20
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 48
ggccgccagc tgaagcttcg 20

<210> 49
<211> 20
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 49
aggccactag tggatctgat 20

<210> 50
<211> 44
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 50
actcgatcc ctgcaagcta ctttgtaatt aaacaaataa cggg 44

<210> 51
<211> 80
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 51
ggagactctt cacactgttg gcgtctatga ttcaagattg tcagtttcca tcgtggattg 60
gaatagatct ggtgaccttg 80

10029w001 sequence listing.txt

```

<210> 52
<211> 90
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 52
acaactattc caatccacga tggaaactga caatcttgaa tcatagacgc caacagtgtg      60
aagagtctcc agctgaagct tcgtacgctg                                         90

<210> 53
<211> 40
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 53
actcggccgg ccatcgatca ctagtggatc tgatatcacc                             40

<210> 54
<211> 38
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 54
actcgcggcc gctctagaca ccaactttga agataggg                               38

<210> 55
<211> 64
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 55
aatacctttg gtttgctcat ggtatcgatt gttttagttt tgtttgtttg ttgtgtataa      60
cggg                                         64

<210> 56
<211> 64
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 56
cccggttatac acaacaaaca aacaaaacta aaacaatcga taccatgagc aaaccaaagg      60
tatt                                         64

<210> 57
<211> 35

```

10029w001 sequence listing.txt

```

<212> DNA
<213> Artificial

<220>
<223> primer

<400> 57
actcagatct tcattatggc atgggaacag gagtt 35

<210> 58
<211> 64
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 58
ccgtaagcaa aaatcttcat ggtatcgatt gttttagttt tgtttgtttg ttgtgtataa 60
cggg 64

<210> 59
<211> 64
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 59
cccgttatac acaacaaaca aacaaaacta aaacaatcga taccatgaag atttttgctt 60
acgg 64

<210> 60
<211> 35
<212> DNA
<213> Artificial

<220>
<223> primer

<400> 60
actcagatct tcattaatat tcaacagcaa tagct 35

```