COMBINATION THERAPY INCLUDING AN MDM2 INHIBITOR AND ONE OR MORE ADDITIONAL PHARMACEUTICALLY ACTIVE AGENTS FOR THE TREATMENT OF CANCERS

Priority

This application claims the benefit of U.S. Provisional Patent Application No. 61/902,717, filed on November 11, 2013 which is hereby incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

The present invention provides combination therapy that includes an MDM2 inhibitor and one or more additional pharmaceutically active agents, particularly for the treatment of cancers. The invention also relates to pharmaceutical compositions that contain an MDM2 inhibitor and one or more additional pharmaceutically active agents for the treatment of cancers.

BACKGROUND OF THE INVENTION

p53 is a tumor suppressor and transcription factor that responds to cellular stress by activating the transcription of numerous genes involved in cell cycle arrest, apoptosis, senescence, and DNA repair. Unlike normal cells, which have infrequent cause for p53 activation, tumor cells are under constant cellular stress from various insults including hypoxia and pro-apoptotic oncogene activation. Thus, there is a strong selective advantage for inactivation of the p53 pathway in tumors, and it has been proposed that eliminating p53 function may be a prerequisite for tumor survival. In support of this notion, three groups of investigators have used mouse models to demonstrate that absence of p53 function is a continuous requirement for the maintenance of established tumors. When the investigators restored p53 function to tumors with inactivated p53, the tumors regressed.

p53 is inactivated by mutation and/or loss in 50% of solid tumors and 10% of liquid tumors. Other key members of the p53 pathway are also genetically or epigenetically altered in cancer. MDM2, an oncoprotein, inhibits p53 function, and it is activated by gene amplification at incidence rates that are reported to be as high as 10%. MDM2, in turn, is inhibited by another tumor suppressor, p14ARF. It has been suggested that alterations downstream of p53 may be responsible for at least partially inactivating the p53 pathway in $p53^{WT}$ tumors (p53 wild type). In support of this concept, some $p53^{WT}$ tumors appear to exhibit reduced apoptotic capacity, although their capacity to undergo cell cycle arrest remains intact. One cancer treatment strategy

involves the use of small molecules that bind MDM2 and neutralize its interaction with p53. MDM2 inhibits p53 activity by three mechanisms: 1) acting as an E3 ubiquitin ligase to promote p53 degradation; 2) binding to and blocking the p53 transcriptional activation domain; and 3) exporting p53 from the nucleus to the cytoplasm. All three of these mechanisms would be blocked by neutralizing the MDM2-p53 interaction. In particular, this therapeutic strategy could be applied to tumors that are $p53^{WT}$, and studies with small molecule MDM2 inhibitors have yielded promising reductions in tumor growth both *in vitro* and *in vivo*. Further, in patients with p53-inactivated tumors, stabilization of wild type p53 in normal tissues by MDM2 inhibition might allow selective protection of normal tissues from mitotic poisons. As used herein, MDM2 means a human MDM2 protein and p53 means a human p53 protein. It is noted that human MDM2 can also be referred to as HDM2 or hMDM2. Several MDM2 inhibitors are in human clinical trials for the treatment of various cancers.

The present invention relates to combination therapy with an MDM2 inhibitor and one or more additional pharmaceutically active agents, which particular combinations show enhanced anti-cancer activity in certain types of cancers over what is expected when the individual members of the combination therapy are used alone.

SUMMARY

In one aspect, the present disclosure refers to the use of a therapeutically effective amount of 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid and trametinib in the manufacture of a medicament for treating melanoma in a patient in need thereof. In still another aspect, the present disclosure refers to the use of a therapeutically effective amount of 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid and trametinib in the manufacture of a medicament for treating AML in a patient in need thereof. In example 1, the present disclosure provides a method of treating melanoma, the method comprising

administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a BRAF inhibitor.

In example 2, the present disclosure provides a method of example 1 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 3, the present disclosure provides a method of example 1 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 4, the present disclosure provides a method of any one of examples 1 to 3 wherein the BRAF inhibitor is dabrafenib.

In example 5, the present disclosure provides a method of any one of examples 1 to 3 wherein the BRAF inhibitor is AMG 2112819 or vemurafenib.

In example 6, the present disclosure provides a method of treating melanoma, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a pan-Raf inhibitor.

In example 7, the present disclosure provides a method of example 6 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 8, the present disclosure provides a method of example 6 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert -butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 9, the present disclosure provides a method of any one of examples 6 to 8 wherein the pan-RAF inhibitor is RAF265 or MLN-2480.

In example 10, the present disclosure provides a method of treating melanoma, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a MEK inhibitor.

In example 11, the present disclosure provides a method of example 10 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 12, the present disclosure provides a method of example 10 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3 – fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 13, the present disclosure provides a method of any one of examples 10 to 12 wherein the MEK inhibitor is trametinib.

In example 14, the present disclosure provides a method of any one of examples 10 to 12 wherein the MEK inhibitor is pimasertib, PD0325901, MEK162, TAK-733, GDC-0973 or AZD8330.

In example 15, the present disclosure provides a method of any one of examples 1 to 14 wherein the melanoma has a BRAF V600E or V600K mutation.

In example 16, the present disclosure provides a method of treating colon cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a BRAF inhibitor.

In example 17, the present disclosure provides a method of example 16 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 18, the present disclosure provides a method of example 16 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2- (tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 19, the present disclosure provides a method of any one of examples 16 to 18 wherein the BRAF inhibitor is dabrafenib.

In example 20, the present disclosure provides a method of any one of examples 16 to 18 wherein the BRAF inhibitor is AMG 2112819 or vemurafenib.

In example 21, the present disclosure provides a method of treating colon cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a pan-Raf inhibitor.

In example 22, the present disclosure provides a method of example 21 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 23, the present disclosure provides a method of example 21 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 24, the present disclosure provides a method of any one of examples 21 to 23 wherein the pan-RAF inhibitor is RAF265 or MLN-2480.

In example 25, the present disclosure provides a method of treating colon cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a MEK inhibitor.

In example 26, the present disclosure provides a method of example 25 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 27, the present disclosure provides a method of example 25 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-

chloro-3 –fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 28, the present disclosure provides a method of any one of examples 25 to 27 wherein the MEK inhibitor is trametinib.

In example 29, the present disclosure provides a method of any one of examples 25 to 27 wherein the MEK inhibitor is pimasertib, PD0325901, MEK162, TAK-733, GDC-0973 or AZD8330.

In example 30, the present disclosure provides a method of any one of examples 16 to 29 wherein the colon cancer has a BRAF V600E or V600K mutation.

In example 31, the present disclosure provides a method of treating liver cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a BRAF inhibitor.

In example 32, the present disclosure provides a method of example 31 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 33, the present disclosure provides a method of example 31 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 34, the present disclosure provides a method of any one of examples 31 to 33 wherein the BRAF inhibitor is dabrafenib, or a pharmaceutically acceptable salt thereof.

In example 35, the present disclosure provides a method of any one of examples 31 to 33 wherein the BRAF inhibitor is AMG 2112819 or vemurafenib.

In example 36, the present disclosure provides a method of treating liver cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a pan-Raf inhibitor.

In example 37, the present disclosure provides a method of example 36 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 38, the present disclosure provides a method of example 36 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-

chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 39, the present disclosure provides a method of any one of examples 36 to 38 wherein the pan-RAF inhibitor is RAF265 or MLN-2480.

In example 40, the present disclosure provides a method of treating liver cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a MEK inhibitor.

In example 41, the present disclosure provides a method of example 40 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 42, the present disclosure provides a method of example 40 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 43, the present disclosure provides a method of any one of examples 40 to 42 wherein the MEK inhibitor is trametinib.

In example 44, the present disclosure provides a method of any one of examples 40 to 42 wherein the MEK inhibitor is pimasertib, PD0325901, MEK162, TAK-733, GDC-0973 or AZD8330.

In example 45, the present disclosure provides a method of any one of examples 31 to 44 wherein the liver cancer has a BRAF V600E or V600K mutation.

In example 46, the present disclosure provides a method of treating bladder cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a MEK inhibitor.

In example 47, the present disclosure provides a method of example 46 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 48, the present disclosure provides a method of example 46 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 49, the present disclosure provides a method of any one of examples 46 to 48 wherein the MEK inhibitor is trametinib.

In example 50, the present disclosure provides a method of any one of examples 46 to 48 wherein the MEK inhibitor is pimasertib, PD0325901, MEK162, TAK-733, GDC-0973 or AZD8330.

In example 51, the present disclosure provides a method of treating AML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a MEK inhibitor.

In example 52, the present disclosure provides a method of example 51 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 53, the present disclosure provides a method of example 51 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 54, the present disclosure provides a method of any one of examples 51 to 53 wherein the MEK inhibitor is trametinib.

In example 55, the present disclosure provides a method of any one of examples 51 to 53 wherein the MEK inhibitor is pimasertib, PD0325901, MEK162, TAK-733, GDC-0973 or AZD8330.

In example 56, the present disclosure provides a method of any one of examples 51 to 55 wherein the AML has a FLT3-ITD mutation.

In example 57, the present disclosure provides a method of treating NSCLC, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a pan-Raf inhibitor.

In example 58, the present disclosure provides a method of example 57 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 59, the present disclosure provides a method of example 57 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 60, the present disclosure provides a method of any one of examples 57 to 59 wherein the pan-RAF inhibitor is RAF265 or MLN-2480.

In example 61, the present disclosure provides a method of treating NSCLC, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a MEK inhibitor.

In example 62, the present disclosure provides a method of example 61 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 63, the present disclosure provides a method of example 61 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3 – fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 64, the present disclosure provides a method of any one of examples 61 to 63 wherein the MEK inhibitor is trametinib.

In example 65, the present disclosure provides a method of any one of examples 61 to 63 wherein the MEK inhibitor is pimasertib, PD0325901, MEK162, TAK-733, GDC-0973 or AZD8330, or a pharmaceutically acceptable salt thereof.

In example 66, the present disclosure provides a method of any one of examples 57 to 65 wherein the NSCLC had a KRAS mutation.

In example 67, the present disclosure provides a method of treating kidney cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a MEK inhibitor.

In example 68, the present disclosure provides a method of example 67 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 69, the present disclosure provides a method of example 67 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 70, the present disclosure provides a method of any one of examples 67 to 69 wherein the MEK inhibitor is trametinib.

In example 71, the present disclosure provides a method of any one of examples 67 to 69 wherein the MEK inhibitor is pimasertib, PD0325901, MEK162, TAK-733, GDC-0973 or AZD8330, or a pharmaceutically acceptable salt thereof.

In example 72, the present disclosure provides a method of treating stomach cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a MEK inhibitor.

In example 73, the present disclosure provides a method of example 72 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 74, the present disclosure provides a method of example 72 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 75, the present disclosure provides a method of any one of examples 72 to 74 wherein the MEK inhibitor is trametinib.

In example 76, the present disclosure provides a method of any one of examples 72 to 74 wherein the MEK inhibitor is pimasertib, PD0325901, MEK162, TAK-733, GDC-0973 or AZD8330.

In example 77, the present disclosure provides a method of any one of examples 72 to 76 wherein the stomach cancer had a KRAS mutation.

In example 78, the present disclosure provides a method of treating prostate cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a PI3K pathway inhibitor.

In example 79, the present disclosure provides a method of example 78 wherein the PI3K pathway inhibitor is a PI3Kα selective inhibitor.

In example 80, the present disclosure provides a method of example 79 wherein the PI3Kα selective inhibitor is AMG 511, AMG2520765 or BYL719.

In example 81, the present disclosure provides a method of example 78 wherein the PI3K pathway inhibitor is a pan-PI3K inhibitor.

In example 82, the present disclosure provides a method of example 81 wherein the pan-PI3K inhibitor is BKM120 or GDC-0941.

In example 83, the present disclosure provides a method of example 78 wherein the PI3K pathway inhibitor is an AKT inhibitor.

In example 84, the present disclosure provides a method of example 83 wherein the AKT inhibitor is MK-2206, GDC-0068 or AZD5363.

In example 85, the present disclosure provides a method of example 78 wherein the PI3K pathway inhibitor is a dual PI3K/mTOR inhibitor.

In example 86, the present disclosure provides a method of example 85 wherein the dual PI3K/mTOR inhibitor is GDC-0980.

In example 87, the present disclosure provides a method of example 78 wherein the PI3K pathway inhibitor is an mTOR inhibitor.

In example 88, the present disclosure provides a method of example 87 wherein the mTOR inhibitor is AZD2014 or MLN0128.

In example 89, the present disclosure provides a method of any one of examples 78 to 88 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 90, the present disclosure provides a method of any one of examples 78 to 88 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 91, the present disclosure provides a method of treating breast cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a PI3K pathway inhibitor.

In example 92, the present disclosure provides a method of example 91 wherein the PI3K pathway inhibitor is a PI3Kα selective inhibitor.

In example 93, the present disclosure provides a method of example 92 wherein the PI3Kα selective inhibitor is AMG 511, AMG2520765 or BYL719.

In example 94, the present disclosure provides a method of claim 91 wherein the PI3K pathway inhibitor is a pan-PI3K inhibitor.

In example 95, the present disclosure provides a method of example 94 wherein the pan-PI3K inhibitor is BKM120 or GDC-0941.

In example 96, the present disclosure provides a method of example 91 wherein the PI3K pathway inhibitor is an AKT inhibitor.

In example 97, the present disclosure provides a method of example 96 wherein the AKT inhibitor is MK-2206, GDC-0068 or AZD5363.

In example 98, the present disclosure provides a method of example 91 wherein the PI3K pathway inhibitor is a dual PI3K/mTOR inhibitor.

In example 99, the present disclosure provides a method of example 98 wherein the dual PI3K/mTOR inhibitor is GDC-0980.

In example 100, the present disclosure provides a method of example 91 wherein the PI3K pathway inhibitor is an mTOR inhibitor.

In example 101, the present disclosure provides a method of example 100 wherein the mTOR inhibitor is AZD2014 or MLN0128.

In example 102, the present disclosure provides a method of any one of examples 91 to 101 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 103, the present disclosure provides a method of any one of examples 91 to 101 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 104, the present disclosure provides a method of any one of examples 91 to 103 wherein the breast cancer has a PI3K mutation.

In example 105, the present disclosure provides a method of treating endometrial cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a PI3K pathway inhibitor.

In example 106, the present disclosure provides a method of example 105 wherein the PI3K pathway inhibitor is a PI3Kα selective inhibitor.

In example 107, the present disclosure provides a method of example 106 wherein the PI3Kα selective inhibitor is AMG 511, AMG252076 or BYL719.

In example 108, the present disclosure provides a method of example 105 wherein the PI3K pathway inhibitor is a pan-PI3K inhibitor.

In example 109, the present disclosure provides a method of example 108 wherein the pan-PI3K inhibitor is BKM120 or GDC-0941.

In example 110, the present disclosure provides a method of example 105 wherein the PI3K pathway inhibitor is an AKT inhibitor.

In example 111, the present disclosure provides a method of example 110 wherein the AKT inhibitor is MK-2206, GDC-0068 or AZD5363.

In example 112, the present disclosure provides a method of example 105 wherein the PI3K pathway inhibitor is a dual PI3K/mTOR inhibitor.

In example 113, the present disclosure provides a method of example 112 wherein the dual PI3K/mTOR inhibitor is GDC-0980.

In example 114, the present disclosure provides a method of example 105 wherein the PI3K pathway inhibitor is an mTOR inhibitor.

In example 115, the present disclosure provides a method of example 114 wherein the mTOR inhibitor is AZD2014 or MLN0128.

In example 116, the present disclosure provides a method of any one of examples 105 to 115 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 117, the present disclosure provides a method of any one of examples 105 to 115 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 118, the present disclosure provides a method of treating NSCLC, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a PI3K pathway inhibitor.

In example 119, the present disclosure provides a method of example 118 wherein the PI3K pathway inhibitor is a PI3Kα selective inhibitor.

In example 120, the present disclosure provides a method of example 119 wherein the PI3Kα selective inhibitor is AMG 511, AMG2520765 or BYL719.

In example 121, the present disclosure provides a method of example 118 wherein the PI3K pathway inhibitor is a pan-PI3K inhibitor.

In example 122, the present disclosure provides a method of example 121 wherein the pan-PI3K inhibitor is BKM120 or GDC-0941.

In example 123, the present disclosure provides a method of example 118 wherein the PI3K pathway inhibitor is an AKT inhibitor.

In example 124, the present disclosure provides a method of example 123 wherein the AKT inhibitor is MK-2206, GDC-0068 or AZD5363.

In example 125, the present disclosure provides a method of example 118 wherein the PI3K pathway inhibitor is a dual PI3K/mTOR inhibitor.

In example 126, the present disclosure provides a method of example 125 wherein the dual PI3K/mTOR inhibitor is GDC-0980.

In example 127, the present disclosure provides a method of example 118 wherein the PI3K pathway inhibitor is an mTOR inhibitor.

In example 128, the present disclosure provides a method of example 127 wherein the mTOR inhibitor is AZD2014 or MLN0128.

In example 129, the present disclosure provides a method of any one of examples 118 to 128 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 130, the present disclosure provides a method of any one of examples 118 to 128 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 131, the present disclosure provides a method of treating head and neck cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a PI3K pathway inhibitor.

In example 132, the present disclosure provides a method of example 131 wherein the PI3K pathway inhibitor is a PI3Kα selective inhibitor.

In example 133, the present disclosure provides a method of example 132 wherein the PI3Kα selective inhibitor is AMG 511, AMG2520765 or BYL719.

In example 134, the present disclosure provides a method of example 131 wherein the PI3K pathway inhibitor is a pan-PI3K inhibitor.

In example 135, the present disclosure provides a method of example 134 wherein the pan-PI3K inhibitor is BKM120 or GDC-0941.

In example 136, the present disclosure provides a method of example 131 wherein the PI3K pathway inhibitor is an AKT inhibitor.

In example 137, the present disclosure provides a method of example 136 wherein the AKT inhibitor is MK-2206, GDC-0068 or AZD5363.

In example 138, the present disclosure provides a method of example 131 wherein the PI3K pathway inhibitor is a dual PI3K/mTOR inhibitor.

In example 139, the present disclosure provides a method of example 138 wherein the dual PI3K/mTOR inhibitor is GDC-0980.

In example 140, the present disclosure provides a method of example 131 wherein the PI3K pathway inhibitor is an mTOR inhibitor.

In example 141, the present disclosure provides a method of example 140 wherein the mTOR inhibitor is AZD2014 or MLN0128.

In example 142, the present disclosure provides a method of any one of examples 131 to 141 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 143, the present disclosure provides a method of any one of examples 131 to 141 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 144, the present disclosure provides a method of treating DLBCL, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a PI3K pathway inhibitor.

In example 145, the present disclosure provides a method of example 144 wherein the PI3K pathway inhibitor is a PI3Kα selective inhibitor.

In example 146, the present disclosure provides a method of example 145 wherein the PI3Kα selective inhibitor is AMG 511, AMG2520765 or BYL719.

In example 147, the present disclosure provides a method of example 144 wherein the PI3K pathway inhibitor is a pan-PI3K inhibitor.

In example 148, the present disclosure provides a method of example 147 wherein the pan-PI3K inhibitor is BKM120 or GDC-0941.

In example 149, the present disclosure provides a method of example 144 wherein the PI3K pathway inhibitor is an AKT inhibitor.

In example 150, the present disclosure provides a method of example 149 wherein the AKT inhibitor is MK-2206, GDC-0068 or AZD5363.

In example 151, the present disclosure provides a method of example 144 wherein the PI3K pathway inhibitor is a dual PI3K/mTOR inhibitor.

In example 152, the present disclosure provides a method of example 151 wherein the dual PI3K/mTOR inhibitor is GDC-0980.

In example 153, the present disclosure provides a method of example 144 wherein the PI3K pathway inhibitor is an mTOR inhibitor.

In example 154, the present disclosure provides a method of example 153 wherein the mTOR inhibitor is AZD2014 or MLN0128.

In example 155, the present disclosure provides a method of any one of examples 144 to 154 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 156, the present disclosure provides a method of any one of examples 144 to 154 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 157, the present disclosure provides a method of treating glioblastoma, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a PI3K pathway inhibitor.

In example 158, the present disclosure provides a method of example 157 wherein the PI3K pathway inhibitor is a PI3Kα selective inhibitor.

In example 159, the present disclosure provides a method of example 158 wherein the PI3Kα selective inhibitor is AMG 511, AMG2520765 or BYL719.

In example 160, the present disclosure provides a method of example 157 wherein the PI3K pathway inhibitor is a pan-PI3K inhibitor.

In example 161, the present disclosure provides a method of example 160 wherein the pan-PI3K inhibitor is BKM120 or GDC-0941.

In example 162, the present disclosure provides a method of example 157 wherein the PI3K pathway inhibitor is an AKT inhibitor.

In example 163, the present disclosure provides a method of example 162 wherein the AKT inhibitor is MK-2206, GDC-0068 or AZD5363.

In example 164, the present disclosure provides a method of example 157 wherein the PI3K pathway inhibitor is a dual PI3K/mTOR inhibitor.

In example 165, the present disclosure provides a method of example 164 wherein the dual PI3K/mTOR inhibitor is GDC-0980.

In example 166, the present disclosure provides a method of example 157 wherein the PI3K pathway inhibitor is an mTOR inhibitor.

In example 167, the present disclosure provides a method of example 166 wherein the mTOR inhibitor is AZD2014 or MLN0128.

In example 168, the present disclosure provides a method of any one of examples 157 to 167 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 169, the present disclosure provides a method of any one of examples 157 to 167 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 170, the present disclosure provides a method of treating bladder cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a PI3K pathway inhibitor.

In example 171, the present disclosure provides a method of example 170 wherein the PI3K pathway inhibitor is a PI3Kα selective inhibitor.

In example 172, the present disclosure provides a method of example 171 wherein the PI3Kα selective inhibitor is AMG 511, AMG2520765 or BYL719.

In example 173, the present disclosure provides a method of example 170 wherein the PI3K pathway inhibitor is a pan-PI3K inhibitor.

In example 174, the present disclosure provides a method of example 173 wherein the pan-PI3K inhibitor is BKM120 or GDC-0941.

In example 175, the present disclosure provides a method of example 170 wherein the PI3K pathway inhibitor is an AKT inhibitor.

In example 176, the present disclosure provides a method of example 175 wherein the AKT inhibitor is MK-2206, GDC-0068 or AZD5363.

In example 177, the present disclosure provides a method of example 170 wherein the PI3K pathway inhibitor is a dual PI3K/mTOR inhibitor.

In example 178, the present disclosure provides a method of example 177 wherein the dual PI3K/mTOR inhibitor is GDC-0980.

In example 179, the present disclosure provides a method of example 170 wherein the PI3K pathway inhibitor is an mTOR inhibitor.

In example 180, the present disclosure provides a method of example 179 wherein the mTOR inhibitor is AZD2014 or MLN0128.

In example 181, the present disclosure provides a method of any one of examples 170 to 180 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-

1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 182, the present disclosure provides a method of any one of examples 170 to 180 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 183, the present disclosure provides a method of treating AML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a PI3K pathway inhibitor.

In example 184, the present disclosure provides a method of example 183 wherein the PI3K pathway inhibitor is a PI3Kα selective inhibitor.

In example 185, the present disclosure provides a method of example 184 wherein the PI3Kα selective inhibitor is AMG 511, AMG2520765 or BYL719.

In example 186, the present disclosure provides a method of example 183 wherein the PI3K pathway inhibitor is a pan-PI3K inhibitor.

In example 187, the present disclosure provides a method of example 186 wherein the pan-PI3K inhibitor is BKM120 or GDC-0941.

In example 188, the present disclosure provides a method of example 183 wherein the PI3K pathway inhibitor is an AKT inhibitor.

In example 189, the present disclosure provides a method of example 188 wherein the AKT inhibitor is MK-2206, GDC-0068 or AZD5363.

In example 190, the present disclosure provides a method of example 183 wherein the PI3K pathway inhibitor is a dual PI3K/mTOR inhibitor.

In example 191, the present disclosure provides a method of example 190 wherein the dual PI3K/mTOR inhibitor is GDC-0980.

In example 192, the present disclosure provides a method of example 183 wherein the PI3K pathway inhibitor is an mTOR inhibitor.

In example 193, the present disclosure provides a method of example 192 wherein the mTOR inhibitor is AZD2014 or MLN0128.

In example 194, the present disclosure provides a method of any one of examples 183 to 193 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 195, the present disclosure provides a method of any one of examples 183 to 193 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 196, the present disclosure provides a method of any one of examples 183 to 195 wherein the AML has a FLT3 ITD mutation.

In example 197, the present disclosure provides a method of treating bladder cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a Bcl2/BclxL inhibitor.

In example 198, the present disclosure provides a method of example 197 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 199, the present disclosure provides a method of example 197 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 200, the present disclosure provides a method of any one of examples 197 to 199 wherein the Bcl2/BclxL inhibitor is navitoclax.

In example 201, the present disclosure provides a method of treating glioblastoma, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a Bcl2/BclxL inhibitor.

In example 202, the present disclosure provides a method of example 201 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 203, the present disclosure provides a method of example 201 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 204, the present disclosure provides a method of any one of examples 201 to 203 wherein the Bcl2/BclxL inhibitor is navitoclax.

In example 205, the present disclosure provides a method of treating head and neck cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a Bcl2/BclxL inhibitor.

In example 206, the present disclosure provides a method of example 205 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 207, the present disclosure provides a method of example 205 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 208, the present disclosure provides a method of any one of examples 205 to 207 wherein the Bcl2/BclxL inhibitor is navitoclax.

In example 209, the present disclosure provides a method of treating kidney cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a Bcl2/BclxL inhibitor.

In example 210, the present disclosure provides a method of example 209 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 211, the present disclosure provides a method of example 209 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 212, the present disclosure provides a method of any one of examples 209 to 211 wherein the Bcl2/BclxL inhibitor is navitoclax.

In example 213, the present disclosure provides a method of treating liver cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a Bcl2/BclxL inhibitor.

In example 214, the present disclosure provides a method of example 213 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 215, the present disclosure provides a method of example 213 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 216, the present disclosure provides a method of any one of examples 213 to 215 wherein the Bcl2/BclxL inhibitor is navitoclax.

In example 217, the present disclosure provides a method of treating sarcoma, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a Bcl2/BclxL inhibitor.

In example 218, the present disclosure provides a method of example 217 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 219, the present disclosure provides a method of example 217 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 220, the present disclosure provides a method of any one of examples 217 to 219 wherein the Bcl2/BclxL inhibitor is navitoclax.

In example 221, the present disclosure provides a method of treating AML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a Bcl2/BclxL inhibitor.

In example 222, the present disclosure provides a method of example 221 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 223, the present disclosure provides a method of example 221 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 224, the present disclosure provides a method of any one of examples 221 to 223 wherein the Bcl2/BclxL inhibitor is navitoclax.

In example 225, the present disclosure provides a method of any one of examples 221 to 224 wherein the AML has a FLT3 ITD mutation.

In example 226, the present disclosure provides a method of treating CML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a Bcl2/BclxL inhibitor.

In example 227, the present disclosure provides a method of example 226 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 228, the present disclosure provides a method of example 226 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 229, the present disclosure provides a method of any one of examples 226 to 228 wherein the Bcl2/BclxL inhibitor is navitoclax.

In example 230, the present disclosure provides a method of any one of examples 226 to 229 where the CML has a BCR-ABL mutation.

In example 231, the present disclosure provides a method of treating DLBCL, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a Bcl2/BclxL inhibitor.

In example 32, the present disclosure provides a method of example 231 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 233, the present disclosure provides a method of example 231 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 234, the present disclosure provides a method of any one of examples 231 to 233 wherein the Bcl2/BclxL inhibitor is navitoclax.

In example 235, the present disclosure provides a method of treating sarcoma, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a BCL2 inhibitor.

In example 236, the present disclosure provides a method of example 235 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-

(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 237, the present disclosure provides a method of example 235 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 238, the present disclosure provides a method of any one of examples 235 to 237 wherein the BCL2 inhibitor is ABT-199.

In example 239, the present disclosure provides a method of treating AML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a BCL2 inhibitor.

In example 240, the present disclosure provides a method of example 239 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 241, the present disclosure provides a method of example 239 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 242, the present disclosure provides a method of any one of examples 239 to 241 wherein the BCL2 inhibitor is ABT-199.

In example 243, the present disclosure provides a method of any one of examples 239 to 242 wherein the AML has a FLT3 ITD mutation.

In example 244, the present disclosure provides a method of treating CML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a BCL2 inhibitor.

In example 245, the present disclosure provides a method of example 244 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 246, the present disclosure provides a method of example 244 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 247, the present disclosure provides a method of any one of examples 244 to 246 wherein the BCL2 inhibitor is ABT-199.

In example 248, the present disclosure provides a method of any one of examples 244 to 247 where the CML has a BCR-ABL mutation.

In example 249, the present disclosure provides a method of treating DLBCL, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a BCL2 inhibitor.

In example 250, the present disclosure provides a method of example 249 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 251, the present disclosure provides a method of example 249 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 252, the present disclosure provides a method of any one of examples 249 to 251 wherein the BCL2 inhibitor is ABT-199.

In example 253, the present disclosure provides a method of treating endometrial cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a BCR-ABL inhibitor.

In example 254, the present disclosure provides a method of example 253 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 255, the present disclosure provides a method of example 253 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 256, the present disclosure provides a method of any one of examples 253 to 255 wherein the BCR-ABL inhibitor is dasatinib.

In example 257, the present disclosure provides a method of treating glioblastoma, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a BCR-ABL inhibitor.

In example 258, the present disclosure provides a method of example 257 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 259, the present disclosure provides a method of example 257 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 260, the present disclosure provides a method of any one of examples 257 to 259 wherein the BCR-ABL inhibitor is dasatinib.

In example 261, the present disclosure provides a method of treating CML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a BCR-ABL inhibitor.

In example 262, the present disclosure provides a method of example 261 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 263, the present disclosure provides a method of example 261 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 264, the present disclosure provides a method of any one of examples 261 to 263 wherein the BCR-ABL inhibitor is dasatinib.

In example 265, the present disclosure provides a method of any one of examples 261 to 264 where the CML has a BCR-ABL mutation.

In example 266, the present disclosure provides a method of treating endometrial cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a BCR-ABL inhibitor.

In example 267, the present disclosure provides a method of example 266 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 268, the present disclosure provides a method of example 266 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-

chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 269, the present disclosure provides a method of any one of examples 266 to 268 wherein the BCR-ABL inhibitor is dasatinib.

In example 270, the present disclosure provides a method of treating bladder cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a BCR-ABL inhibitor.

In example 271, the present disclosure provides a method of example 270 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 272, the present disclosure provides a method of example 270 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 273, the present disclosure provides a method of any one of examples 270 to 272 wherein the BCR-ABL inhibitor is dasatinib.

In example 274, the present disclosure provides a method of treating head and neck cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a BCR-ABL inhibitor.

In example 275, the present disclosure provides a method of example 274 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 276, the present disclosure provides a method of example 274 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 277, the present disclosure provides a method of any one of examples 274 to 276 wherein the BCR-ABL inhibitor is dasatinib.

In example 278, the present disclosure provides a method of treating kidney cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a HDAC inhibitor.

In example 279, the present disclosure provides a method of example 278 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 280, the present disclosure provides a method of example 278 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 281, the present disclosure provides a method of any one of examples 278 to 280 wherein the HDAC inhibitor is panobinostat.

In example 282, the present disclosure provides a method of treating liver cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a HDAC inhibitor.

In example 283, the present disclosure provides a method of example 282 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 284, the present disclosure provides a method of example 282 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 285, the present disclosure provides a method of any one of examples 282 to 284 wherein the HDAC inhibitor is panobinostat.

In example 286, the present disclosure provides a method of treating melanoma, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and a HDAC inhibitor.

In example 287, the present disclosure provides a method of example 286 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 288, the present disclosure provides a method of example 286 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 289, the present disclosure provides a method of any one of examples 286 to 288 wherein the HDAC inhibitor is panobinostat.

In example 290, the present disclosure provides a method of treating AML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and decitabine.

In example 291, the present disclosure provides a method of example 290 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 292, the present disclosure provides a method of example 290 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 293, the present disclosure provides a method of any one of examples 290 to 292 wherein the AML has a FLT3 ITD mutation.

In example 294, the present disclosure provides a method of treating AML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and cytarabine.

In example 295, the present disclosure provides a method of example 294 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 296, the present disclosure provides a method of example 294 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 297, the present disclosure provides a method of any one of examples 294 to 296 wherein the AML has a FLT3 ITD mutation.

In example 298, the present disclosure provides a method of treating AML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and doxorubicin.

In example 299, the present disclosure provides a method of example 298 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-

(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 300, the present disclosure provides a method of example 298 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 301, the present disclosure provides a method of any one of examples 298 to 300 wherein the AML has a FLT3 ITD mutation.

In example 302, the present disclosure provides a method of treating sarcoma, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and etoposide.

In example 303, the present disclosure provides a method of example 302 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 304, the present disclosure provides a method of example 302 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 305, the present disclosure provides a method of treating breast cancer, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and doxorubicin.

In example 306, the present disclosure provides a method of example 305 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 307, the present disclosure provides a method of example 305 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 308, the present disclosure provides a method of treating AML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and imatinib.

In example 309, the present disclosure provides a method of example 308 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 310, the present disclosure provides a method of example 308 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 311, the present disclosure provides a method of any one of examples 308 to 310 wherein the AML has a FLT3 ITD mutation.

In example 312, the present disclosure provides a method of treating AML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and ponatinib.

In example 313, the present disclosure provides a method of example 312 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 314, the present disclosure provides a method of example 312 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 315, the present disclosure provides a method of any one of examples 312 to 314 wherein the AML has a FLT3 ITD mutation.

In example 316, the present disclosure provides a method of treating AML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and bosutinib.

In example 317, the present disclosure provides a method of example 316 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 318, the present disclosure provides a method of example 316 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 319, the present disclosure provides a method of any one of examples 316 to 318 wherein the AML has a FLT3 ITD mutation

In example 320, the present disclosure provides a method of treating AML, the method comprising administering to a patient in need thereof a therapeutically effective amount of an MDM2 inhibitor and nilotinib.

In example 321, the present disclosure provides a method of example 320 wherein the MDM2 inhibitor is 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof.

In example 322, the present disclosure provides a method of example 320 wherein the MDM2 inhibitor is 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof.

In example 323, the present disclosure provides a method of any one of examples 320 to 322 wherein the AML has a FLT3 ITD mutation

In example 324, the present disclosure provides a method of treating melanoma, the method comprising administering to a patient in need thereof a therapeutically effective amount of 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, dabrafenib and trametinib.

In example 325, the present disclosure provides a method of treating melanoma, the method comprising administering to a patient in need thereof a therapeutically effective amount of 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, dabrafenib and trametinib.

In example 326, the present disclosure provides a method of any one of examples 324 to 325 wherein the melanoma has a BRAF V600E or V600K mutation.

In example 327, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; dabrafenib; and a pharmaceutically acceptable excipient.

In example 328, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic

acid, or a pharmaceutically acceptable salt thereof; dabrafenib; and a pharmaceutically acceptable excipient.

In example 329, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; AMG 2112819; and a pharmaceutically acceptable excipient.

In example 330, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; AMG 2112819; and a pharmaceutically acceptable excipient.

In example 331, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; RAF265; and a pharmaceutically acceptable excipient.

In example 332, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; RAF265; and a pharmaceutically acceptable excipient.

In example 333, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; MLN-2480; and a pharmaceutically acceptable excipient.

In example 334, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; MLN-2480; and a pharmaceutically acceptable excipient.

In example 335, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; trametinib; and a pharmaceutically acceptable excipient.

In example 336, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; trametinib; and a pharmaceutically acceptable excipient.

In example 337, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; nilotinib; and a pharmaceutically acceptable excipient.

In example 338, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; nilotinib; and a pharmaceutically acceptable excipient.

In example 339, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; pimasertinib; and a pharmaceutically acceptable excipient.

In example 340, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; pimasertinib; and a pharmaceutically acceptable excipient.

In example 341, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; PD0325901; and a pharmaceutically acceptable excipient.

In example 342, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; PD0325901; and a pharmaceutically acceptable excipient.

In example 343, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-

3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; MEK 162; and a pharmaceutically acceptable excipient.

In example 344, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; MEK 162; and a pharmaceutically acceptable excipient.

In example 345, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; TAK-733; and a pharmaceutically acceptable excipient.

In example 346, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; TAK-733; and a pharmaceutically acceptable excipient.

In example 347, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; GDC-0973; and a pharmaceutically acceptable excipient.

In example 348, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; GDC-0973; and a pharmaceutically acceptable excipient.

In example 349, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; AZD 8330; and a pharmaceutically acceptable excipient.

In example 350, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; AZD 8330; and a pharmaceutically acceptable excipient.

In example 351, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; vemurafenib; and a pharmaceutically acceptable excipient.

In example 352, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; vemurafenib; and a pharmaceutically acceptable excipient.

In example 353, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; AMG 511; and a pharmaceutically acceptable excipient.

In example 354, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; AMG 511; and a pharmaceutically acceptable excipient.

In example 355, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; AMG 2520765; and a pharmaceutically acceptable excipient.

In example 356, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; AMG 2520765; and a pharmaceutically acceptable excipient.

In example 357, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; BYL719; and a pharmaceutically acceptable excipient.

In example 358, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic

acid, or a pharmaceutically acceptable salt thereof; BYL719; and a pharmaceutically acceptable excipient.

In example 359, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; BKM 120; and a pharmaceutically acceptable excipient.

In example 360, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; BKM 120; and a pharmaceutically acceptable excipient.

In example 361, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; GDC-0941; and a pharmaceutically acceptable excipient.

In example 362, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; GDC-0941; and a pharmaceutically acceptable excipient.

In example 363, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; MK-2206; and a pharmaceutically acceptable excipient.

In example 364, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; MK-2206; and a pharmaceutically acceptable excipient.

In example 365, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; AZD 5363; and a pharmaceutically acceptable excipient.

In example 366, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; AZD 5363; and a pharmaceutically acceptable excipient.

In example 367, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; GDC-0068; and a pharmaceutically acceptable excipient.

In example 368, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; GDC-0068; and a pharmaceutically acceptable excipient.

In example 369, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; GDC-0980; and a pharmaceutically acceptable excipient.

In example 370, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; GDC-0980; and a pharmaceutically acceptable excipient.

In example 371, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; AZD2014; and a pharmaceutically acceptable excipient.

In example 372, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; AZD2014; and a pharmaceutically acceptable excipient.

In example 373, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-

3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; MLN0128; and a pharmaceutically acceptable excipient.

In example 374, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; MLN0128; and a pharmaceutically acceptable excipient.

In example 375, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; navitoclax; and a pharmaceutically acceptable excipient.

In example 376, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; navitoclax; and a pharmaceutically acceptable excipient.

In example 377, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; ABT-199; and a pharmaceutically acceptable excipient.

In example 378, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; ABT-199; and a pharmaceutically acceptable excipient.

In example 379, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; dasatinib; and a pharmaceutically acceptable excipient.

In example 380, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; dasatinib; and a pharmaceutically acceptable excipient.

In example 381, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; panobinostat; and a pharmaceutically acceptable excipient.

In example 382, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; panobinostat; and a pharmaceutically acceptable excipient.

In example 383, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; decitabine; and a pharmaceutically acceptable excipient.

In example 384, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; decitabine; and a pharmaceutically acceptable excipient.

In example 385, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; cytarabine; and a pharmaceutically acceptable excipient.

In example 386, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; cytarabine; and a pharmaceutically acceptable excipient.

In example 387, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; doxorubicin; and a pharmaceutically acceptable excipient.

In example 388, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic

acid, or a pharmaceutically acceptable salt thereof; doxorubicin; and a pharmaceutically acceptable excipient.

In example 389, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; etoposide; and a pharmaceutically acceptable excipient.

In example 390, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; etoposide; and a pharmaceutically acceptable excipient.

In example 391, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; imatinib; and a pharmaceutically acceptable excipient.

In example 392, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; imatinib; and a pharmaceutically acceptable excipient.

In example 393, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; ponatinib; and a pharmaceutically acceptable excipient.

In example 394, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; ponatinib; and a pharmaceutically acceptable excipient.

In example 395, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; bosutinib; and a pharmaceutically acceptable excipient.

In example 396, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; bosutinib; and a pharmaceutically acceptable excipient.

In example 397, the present disclosure provides a pharmaceutical composition comprising: 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, or a pharmaceutically acceptable salt thereof; dabrafenib; trametinib; and a pharmaceutically acceptable excipient.

In example 398, the present disclosure provides a pharmaceutical composition comprising: 4-(2-((3R,5R,6S)-1-((S)-2-(tert-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chloropheny1)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid, or a pharmaceutically acceptable salt thereof; dabrafenib; trametinib; and a pharmaceutically acceptable excipient.

In example 399, the present disclosure provides a method of treating melanoma comprising administering to a patient in need thereof a therapeutically effective amount of 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid, and dabrafenib.

In example 400, the present disclosure provides a combination of an MDM2 inhibitor medicament and a MEK inhibitor medicament for treating a solid tumor.

In example 401, the present disclosure provides a combination of an MDM2 inhibitor medicament and a MEK inhibitor medicament for treating AML.

In example 402, the present disclosure provides a combination of an MDM2 inhibitor medicament and a BRAF inhibitor medicament for treating a solid tumor.

In example 403, the present disclosure provides a combination of an MDM2 inhibitor medicament and a BRAF inhibitor medicament for treating AML.

In example 404, the present disclosure provides a use of an MDM2 inhibitor in combination with a BRAF inhibitor for manufacture of a medicament for the management or treatment of melanoma, liver cancer, AML or colon cancer in a subject.

In example 405, the present disclosure provides a use of an MDM2 inhibitor in combination with a MEK inhibitor for manufacture of a medicament for the management or treatment of melanoma, liver cancer, AML or colon cancer in a subject.

A further example of the disclosure includes the use of a combination comprising an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, a MEK inhibitor,

a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor. A further example of the disclosure includes the use of a combination for treatment of cancer, the combination comprising an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, an MEK inhibitor, a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor. A further example of the disclosure includes a method of using a combination comprising an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, a MEK inhibitor, a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor, for the treatment of cancer. A further example of the disclosure includes the use of a combination for treatment of cancer, the combination comprising an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, an MEK inhibitor, a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor, wherein the use comprises self-administering the combination.

A further example of the disclosure includes a method of treating cancer comprising prescribing a combination further comprising an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, a MEK inhibitor, a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor. A further example of the disclosure includes a method of treating cancer comprising prescribing to a subject in need thereof, a combination further comprising an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, a MEK inhibitor, a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor. A further example of the disclosure includes a method of treating cancer comprising prescribing a combination containing an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, a MEK inhibitor, a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor.

A further example of the disclosure includes a method of treating cancer using a combination comprising an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, a MEK inhibitor, a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor, wherein such method further comprises listing said combination in a formulary and directing a patient in need of such cancer treatment to administer the combination. A further example of the disclosure includes a method of treating cancer using a combination comprising an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, a MEK inhibitor, a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor, wherein such method further comprises listing said combination in a formulary and directing a patient in need of such cancer treatment to self-administer the combination.

A further example of the disclosure includes a method of treating cancer using a combination comprising an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, a MEK inhibitor, a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor, wherein such method further comprises selling said combination for self-administration to a patient in need of such cancer treatment.

A further example of the disclosure includes a method of using a combination comprising an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, a MEK inhibitor, a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor, for the treatment of cancer, wherein such method comprises purchasing said combination for self-administration by a patient in need of such cancer treatment. A further example of the disclosure includes a method of using a combination comprising an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, a MEK inhibitor, a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor, for the treatment of cancer, wherein such method comprises purchasing said combination for administration by a patient in need of such cancer treatment.

A further example of the disclosure includes a method of treating cancer comprising instructing a subject in need of such treatment to administer a combination comprising an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, a MEK inhibitor, a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor.

A further example of the disclosure includes a process of treating cancer comprising

A] prescribing

B] selling or advertising to sell,

C] purchasing,

D] instructing to self-administer, or

E] administering

of a combination described herein, wherein the combination has been approved by a regulatory agency for the treatment of cancer, to a subject in need of cancer treatment.

A further example of the disclosure includes a method of supplying a combination comprising an MDM2 inhibitor with another therapeutic agent selected from a RAF inhibitor, a MEK inhibitor, a Pi3K a selective inhibitor, an mTOR inhibitor, an AKT inhibitor or an Aurora kinase inhibitor for treating cancer, said method comprises reimbursing a physician, a formulary, a patient or an insurance company for the sale of said combination.

For clarity, the term “instructing” is meant to include information on a label approved by a regulatory agency, in addition to its commonly understood definition.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows data for the combination of compound 1 and compound A with A204 cells.

Figure 1a shows data for the combination of compound 1 and compound A with A204 cells.

Figure 2 shows data for the combination of compound 1 and compound A with A375sq2 cells.

Figure 2a shows data for the combination of compound 1 and compound A with A375sq2 cells.

Figure 3 shows data for the combination of compound 2 and compound B with A-427 cells.

Figure 3a shows data for the combination of compound 2 and compound B with A-427 cells.

Figure 4 shows data for the combination of compound 1 and compound B with C32 cells.

Figure 4a shows data for the combination of compound 1 and compound B with C32 cells.

Figure 5 shows data for the combination of compound 2 and compound B with C32 cells.

Figure 5a shows data for the combination of compound 2 and compound B with C32 cells.

Figure 6 shows data for the combination of compound 1 and compound A with G-361 cells.

Figure 6a shows data for the combination of compound 1 and compound A with G-361 cells.

Figure 7 shows data for the combination of compound 1 and compound A with LS 174T cells.

Figure 7a shows data for the combination of compound 1 and compound A with LS 174T cells.

Figure 8 shows data for the combination of compound 1 and compound A with MCF7 cells.

Figure 8a shows data for the combination of compound 1 and compound A with MCF7 cells.

Figure 9 shows data for the combination of compound 1 and compound B with NCI-H1666 cells.

Figure 9a shows data for the combination of compound 1 and compound B with NCI-H1666 cells.

Figure 10 shows data for the combination of compound 2 and compound B with NCI-H1666 cells.

Figure 10a shows data for the combination of compound 2 and compound B with NCI-H1666 cells.

Figure 11 shows data for the combination of compound 1 and compound A with RKO cells.

Figure 11a shows data for the combination of compound 1 and compound A with RKO cells.

Figure 12 shows data for the combination of compound 2 and compound B with RKO cells.

Figure 12a shows data for the combination of compound 2 and compound B with RKO cells.

Figure 13 shows data for the combination of compound 1 and compound B with RT4 cells.

Figure 13a shows data for the combination of compound 1 and compound B with RT4 cells.

Figure 14 shows the data for the combination of compound 2 and compound B with RT4 cells.

Figure 14a shows data for the combination of compound 2 and compound B with RT4 cells.

Figure 15 shows data for the combination of compound 1 and compound B with SH-4 cells.

Figure 15a shows data for the combination of compound 1 and compound B with SH-4 cells.

Figure 16 shows data for the combination of compound 2 and compound B with SH-4 cells.

Figure 16a shows data for the combination of compound 2 and compound B with SH-4 cells.

Figure 17 shows data for the combination of compound 1 and compound B with SK-HEP-1 cells.

Figure 17a shows data for the combination of compound 1 and compound B with SK-HEP-1 cells.

Figure 18 shows data for the combination of compound 2 and compound B with SK-HEP-1 cells.

Figure 18a shows data for the combination of compound 2 and compound B with SK-HEP-1 cells.

Figure 19 shows data for the combination of compound 3 and compound A with A204 cells.

Figure 19a shows data for the combination of compound 3 and compound A with A204 cells.

Figure 20 shows data for the combination of compound 3 and compound A with A375sq2 cells.

Figure 20a shows data for the combination of compound 3 and compound A with A375sq2 cells.

Figure 21 shows data for the combination of compound 3 and compound B with A-427 cells.

Figure 21a shows data for the combination of compound 3 and compound B with A-427 cells.

Figure 22 shows data for the combination of compound 4 and compound B with A-427 cells.

Figure 22a shows data for the combination of compound 4 and compound B with A-427 cells.

Figure 23 shows data for the combination of compound 3 and compound B with C32 cells.

Figure 23a shows data for the combination of compound 3 and compound B with C32 cells.

Figure 24 shows data for the combination of compound 4 and compound B with C32 cells.

Figure 24a shows data for the combination of compound 4 and compound B with C32 cells.

Figure 25 shows data for the combination of compound 3 and compound A with G-361cells.

Figure 25a shows data for the combination of compound 3 and compound A with G-361 cells. Figures 26 and 26a omitted.

Figure 27 shows data for the combination of compound 3 and compound A with LS 174T cells.

Figure 27a shows data for the combination of compound 3 and compound A with LS 174T cells.

Figure 28 shows data for the combination of compound 3 and compound A with MCF7 cells.

Figure 28a shows data for the combination of compound 3 and compound A with MCF7 cells.

Figure 29 shows data for the combination of compound 3 and compound B with NCI-H1666 cells.

Figure 29a shows data for the combination of compound 3 and compound B with NCI-H1666 cells.

Figure 30 shows data for the combination of compound 4 and compound B with NCI-H1666 cells.

Figure 30a shows data for the combination of compound 4 and compound B with NCI-H1666 cells.

Figure 31 shows data for the combination of compound 3 and compound A with RKO cells.

Figure 31a shows data for the combination of compound 3 and compound A with RKO cells.

Figure 32 shows data for the combination of compound 4 and compound B with RKO cells.

Figure 32a shows data for the combination of compound 4 and compound B with RKO cells.

Figure 33 shows data for the combination of compound 3 and compound B with RT4 cells.

Figure 33a shows data for the combination of compound 3 and compound B with RT4 cells.

Figure 34 shows data for the combination of compound 4 and compound B with RT4 cells.

Figure 34a shows data for the combination of compound 4 and compound B with RT4 cells.

Figure 35 shows data for the combination of compound 3 and compound B with SH-4 cells.

Figure 35a shows data for the combination of compound 3 and compound B with SH-4 cells.

Figure 36 shows data for the combination of compound 4 and compound B with SH-4 cells.

Figure 36a shows data for the combination of compound 4 and compound B with SH-4 cells.

Figure 37 shows data for the combination of compound 3 and compound B with SK-HEP-1 cells.

Figure 37a shows data for the combination of compound 3 and compound B with SK-HEP-1 cells.

Figure 38 shows data for the combination of compound 4 and compound B with SK-HEP-1 cells.

Figure 38a shows data for the combination of compound 4 and compound B with SK-HEP-1 cells.

Figure 39 shows data for the combination of compound 5 and compound A with A204 cells.

Figure 39a shows data for the combination of compound 5 and compound A with A204 cells.

Figure 40 shows data for the combination of compound 5 and compound A with A375sq2 cells.

Figure 40a shows data for the combination of compound 5 and compound A with A375sq2 cells.

Figure 41 shows data for the combination of compound 5 and compound B with CAL-51 cells.

Figure 41a shows data for the combination of compound 5 and compound B with CAL-51 cells.

Figure 42 shows data for the combination of compound 5 and compound A with G-361 cells.

Figure 42a shows data for the combination of compound 5 and compound A with G-361 cells.

Figure 43 shows data for the combination of compound 5 and compound B with HT-1197 cells.

Figure 43a shows data for the combination of compound 5 and compound B with HT-1197 cells.

Figure 44 shows data for the combination of compound 5 and compound A with LS 174T cells.

Figure 44a shows data for the combination of compound 5 and compound A with LS174T cells.

Figure 45 shows data for the combination of compound 5 and compound A with MCF7 cells.

Figure 45a shows data for the combination of compound 5 and compound A with MCF7 cells.

Figure 46 shows data for the combination of compound 5 and compound B with NCI-H460 cells.

Figure 46a shows data for the combination of compound 5 and compound B with NCI-H460 cells.

Figure 47 shows data for the combination of compound 5 and compound A with RKO cells.

Figure 47a shows data for the combination of compound 5 and compound A with RKO cells.

Figure 48 shows data for the combination of compound 6 and compound B with A204 cells.

Figure 48a shows data for the combination of compound 6 and compound B with A204 cells.

Figure 49 shows data for the combination of compound 6 and compound B with A2780 cells.

Figure 49a shows data for the combination of compound 6 and compound B with A2780 cells.

Figure 50 shows data for the combination of compound 6 and compound B with C32 cells.

Figure 50a shows data for the combination of compound 6 and compound B with C32 cells.

Figure 51 shows data for the combination of compound 6 and compound B with G-401cells.

Figure 51a shows data for the combination of compound 6 and compound B with G-401 cells.

Figure 52 shows data for the combination of compound 6 and compound B with SK-HEP-1 cells.

Figure 52a shows data for the combination of compound 6 and compound B with SK-HEP-1 cells.

Figures 53-56 omitted.

Figure 57 shows data for the combination of compound 8 and compound B with BV-173 cells.

Figure 57a shows data for the combination of compound 8 and compound B with BV-173 cells.

Figure 58 shows data for the combination of compound 8 and compound B with CML-T1 cells.

Figure 58a shows data for the combination of compound 8 and compound B with CML-T1 cells.

Figure 59 shows data for the combination of compound 9 and compound B with KNS-81-FD cells.

Figure 59a shows data for the combination of compound 9 and compound B with KNS-81-FD cells.

Figure 60 shows data for the combination of compound 9 and compound B with SW48 cells.

Figure 60a shows data for the combination of compound 9 and compound B with SW48 cells.

Figure 61 shows data for the combination of compound 10 and compound B with MDA-MB-175 VII cells.

Figure 61a shows data for the combination of compound 10 and compound B with MDA-MB-175 VII cells.

Figure 62 shows data for the combination of compound 10 and compound B with UACC-812 cells.

Figure 62a shows data for the combination of compound 10 and compound B UACC-812 cells.

Figure 63 shows data for the combination of compound 11 and compound A with HCT-116 cells.

Figure 63a shows data for the combination of compound 11 and compound A HCT-116 cells.

Figure 64 shows data for the combination of compound 13 and compound B with GDM-1 cells.

Figure 64a shows data for the combination of compound 13 and compound B with GDM-1 cells.

Figure 65 shows data for the combination of compound 13 and compound B with ML-2 cells.

Figure 65a shows data for the combination of compound 13 and compound B ML-2 cells.

Figure 66 shows data for the combination of compound 13 and compound B with MOLM-13 cells.

Figure 66a shows data for the combination of compound 13 and compound B with MOLM-13 cells.

Figure 67 shows data for the combination of compound 13 and compound B with OCI-AML3 cells.

Figure 67a shows data for the combination of compound 13 and compound B with OCI-AML3 cells.

Figure 68 shows data for the combination of compound 12 and compound B with GDM-1 cells.

Figure 68a shows data for the combination of compound 12 and compound B with GDM-1 cells.

Figure 69 shows data for the combination of compound 12 and compound B with ML-2 cells.

Figure 69a shows data for the combination of compound 12 and compound B with ML-2 cells.

Figure 70 shows data for the combination of compound 12 and compound B with MOLM-13 cells.

Figure 70a shows data for the combination of compound 12 and compound B with MOLM-13 cells.

Figure 71 shows data for the combination of compound 12 and compound B with OCI-AML3 cells.

Figure 71a shows data for the combination of compound 12 and compound B with OCI-AML3 cells.

Figure 72 shows data for combinations of AMG 232 and various MAP kinase pathway inhibitors.

Figure 73 shows data for combinations of AM-7209 and various MAP kinase pathway inhibitors.

Figure 74 shows data for combinations of RG7112 and various MAP kinase pathway inhibitors.

Figure 75 shows data for combinations of AMG 232 and various PI3 kinase pathway inhibitors.

Figure 76 shows data for combinations of AM-7209 and various PI3 kinase pathway inhibitors.

Figure 77 shows data for combinations of RG7112 and various PI3 kinase pathway inhibitors.

Figure 78 shows data for combinations of AMG 232 and various compounds active in the intrinsic apoptosis pathway.

Figure 79 shows data for combinations of AM-7209 and various compounds active in the intrinsic apoptosis pathway.

Figure 80 shows data for combinations of RG7112 and various compounds active in the intrinsic apoptosis pathway.

Figure 81 shows data for combinations of AMG 232 and various chemotherapeutic compounds.

Figure 82 shows data for combinations of AM-7209 and various chemotherapeutic compounds.

Figure 83 shows data for combinations of RG7112 and various chemotherapeutic compounds.

Figure 84 shows data for combinations of AMG 232 and various chemotherapeutic compounds.

Figure 85 shows data for combinations of AM-7209 and various chemotherapeutic compounds.

Figure 86 shows data for combinations of RG7112 and various chemotherapeutic compounds.

Figure 87 shows data for combinations of AMG 232 and various compounds in hematopoietic cell lines.

Figure 88 shows data for combinations of AM-7209 and various compounds in hematopoietic cell lines.

Figure 89 shows data for combinations of RG7112 and various compounds in hematopoietic cell lines.

Figure A shows tumor xenograft data for the combination of AMG 232 and cisplatin in an H460 tumor.

Figure B shows tumor xenograft data for the combination of AMG 232 and cisplatin in an HCT116 tumor.

Figure C shows tumor xenograft data for the combination of AMG 232 and CPT-11(irinotecan) in an HCT116 tumor.

Figure D shows tumor xenograft data for the combination of AMG 232 and doxorubicin in a SJSA-1 tumor.

Figure E shows tumor xenograft data for the combination of AMG 232 and BRAF inhibitor AMG 2112819 or MEK inhibitor 1009089 in a RKO tumor.

Figure F shows tumor xenograft data for the combination of RG7112 and PI3K inhibitor AMG 2520765 in a U87 tumor.

Figure G shows tumor xenograft data for the combination of AMG 232 and MEK inhibitor AMG 1009089 in an **A375** tumor.

Figure H shows tumor xenograft data for the combination of AMG 232 and BRAF inhibitor AMG 2112819 in an A375sq2 tumor.

Figure I shows tumor xenograft data for the combination of AMG 232, BRAF inhibitor AMG 2112819 and PI3K inhibitor 2539965 in a RKO tumor.

Figure J shows tumor xenograft data for various combinations of AMG 232, BRAF inhibitor AMG 2112819 and PI3k inhibitor AMG2539965 in a RKO tumor.

Figure K shows tumor xenograft data for the combination of AMG 232 and doxorubicin in a MOLM13 tumor.

Figure L shows tumor xenograft data for the combination of AMG 232 and MEK inhibitor AMG1009089 in a MOLM13 tumor.

Figure M shows tumor xenograft data for the combination of AMG 232 and cytarabine in a MOLM13 tumor.

Figure N shows tumor xenograft data for the combination of AMG 232 and decitabine in a MOLM13 tumor.

Figure O shows tumor xenograft data for the combination of AMG 232 and Sorafanib in a MOLM13 tumor.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides combination therapy that includes an MDM2 inhibitor and one or more additional pharmaceutically active agents, particularly for the treatment of cancers. The invention also relates to pharmaceutical compositions that contain an MDM2 inhibitor and one or more additional pharmaceutically active agents for the treatment of cancers.

The term "comprising" is meant to be open ended, including the indicated component but not excluding other elements.

The term "therapeutically effective amount" means an amount of a compound, or combination of compounds, that ameliorates, attenuates or eliminates one or more symptom of a particular disease or condition, or prevents or delays the onset of one of more symptom of a particular disease or condition.

The terms "patient" and “subject” may be used interchangeably and mean animals, such as dogs, cats, cows, horses, sheep and humans. Particular patients are mammals. The term patient includes males and females.

The term "pharmaceutically acceptable" means that the referenced substance, such as a compound, or a salt of the compound, or a formulation containing the compound, or a particular excipient, are suitable for administration to a patient.

The terms "treating", "treat" or "treatment" and the like include preventative (e.g., prophylactic) and palliative treatment. The term” treating” and the like, in accordance with the present invention, means reducing or eliminating cancers cells in a patient.

The term “excipient” means any pharmaceutically acceptable additive, carrier, diluent, adjuvant, or other ingredient, other than the active pharmaceutical ingredient (API), which is typically included for formulation and/or administration to a patient.

The phrase “compound(s) of the present invention” includes MDM2 inhibitors and/or the one or more additional pharmaceutically active agents according to the context of the use.

An “MDM2 inhibitor” is defined as a compound with a molecular weight less than about 1000 that binds MDM2 as shown with *in vitro* testing or by other means.

The compounds of the present invention are administered to a patient in a therapeutically effective amount. The compounds can be administered alone or as part of a pharmaceutically acceptable composition or formulation. In addition, the compounds or compositions can be administered all at once, as for example, by a bolus injection, multiple times, such as by a series of tablets, or delivered substantially uniformly over a period of time, as for example, using transdermal delivery. It is also noted that the dose of the compounds can be varied over time.

If the patient is to receive or is receiving multiple pharmaceutically active compounds, the compounds can be administered simultaneously or sequentially. For example, in the case of tablets, the active compounds may be found in one tablet or in separate tablets, which can be administered at once or sequentially in any order. In addition, it should be recognized that the compositions may be different forms. For example, one or more compounds may be delivered via a tablet, while another is administered via injection or orally as a syrup. All combinations, delivery methods and administration sequences are contemplated.

The term "cancer" means a physiological condition in mammals that is characterized by unregulated cell growth. General classes of cancers include carcinomas, lymphomas, sarcomas, and blastomas.

The compounds of the present invention can be used to treat cancer. The methods of treating a cancer comprise administering to a patient in need thereof a therapeutically effective amount of one or more compounds, or pharmaceutically acceptable salts of any of the compounds.

The compounds of the present invention can be used to treat tumors. The methods of treating a tumor comprise administering to a patient in need thereof a therapeutically effective amount of one or more compounds of the present invention, or pharmaceutically acceptable salts of any of the compounds.

The invention also concerns the use of the compounds in the manufacture of a medicament for the treatment of a condition such as a cancer.

Cancers which may be treated with compounds of the present invention include, without limitation, carcinomas such as cancer of the bladder, breast, colon, rectum, kidney, liver, lung (small cell lung cancer, and non-small-cell lung cancer), esophagus, gall-bladder, ovary, pancreas, stomach, cervix, thyroid, prostate, and skin (including squamous cell carcinoma); hematopoietic tumors of lymphoid lineage (including leukemia, acute lymphocytic leukemia, chronic myelogenous leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma and Burkett's lymphoma); hematopoietic tumors of myeloid lineage (including acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic leukemia); tumors of mesenchymal origin (including fibrosarcoma and rhabdomyosarcoma, and other sarcomas, e.g., soft tissue and bone); tumors of the central and peripheral nervous system (including astrocytoma, neuroblastoma, glioma and schwannomas); and other tumors (including melanoma, seminoma, teratocarcinoma, osteosarcoma, xenoderoma pigmentosum, keratoctanthoma, thyroid follicular cancer and Kaposi's sarcoma). Other cancers that can be treated with the compound of the present invention include endometrial cancer, head and neck cancer, glioblastoma, malignant ascites, and hematopoietic cancers.

Particular cancers that can be treated by the compounds of the present invention include soft tissue sarcomas, bone cancers such as osteosarcoma, breast tumors, bladder cancer, Li-Fraumeni syndrome, brain tumors, rhabdomyosarcoma, adrenocortical carcinoma, colorectal cancer, non-small cell lung cancer, and acute myelogenous leukemia (AML).

In a particular embodiment of the invention that relates to the treatment of cancers, the cancer is identified as p53wildtype ($p53^{WT}$). In another particular embodiment, the cancer is identified as $p53^{WT}$ and CDKN2A mutant. In another aspect, the present invention provides a diagnostic for determining which patients should be administered a compound of the present invention. For example, a sample of a patient's cancer cells may be taken and analyzed to determine the status of the cancer cells with respect to p53 and/or CDKN2A. In one aspect, a patient having a cancer that is $p53^{WT}$ will be selected for treatment over patients having a cancer that is mutated with respect to p53. In another aspect, a patient having a cancer that is both $p53^{WT}$ and has a mutant CDNK2A protein is selected over a patient that does not have these characteristics. In still another aspect, the patient has a cancer that is $p53^{WT}$ and exhibits MDM2 amplification. The taking of cancer cells for analyses is well known to those skilled in the art. The term "$p53^{WT}$" means a protein encoded by genomic DNA sequence no. NC_000017 version 9 (7512445..7531642)(GenBank); a protein encoded by cDNA sequence no. NM_000546 (GenBank); or a protein having the GenBank sequence no. NP_000537.3. The term "CDNK2A mutant" means a CDNK2A protein that in not wild type. The term "CDKN2A wild type" means a protein encoded by genomic DNA sequence no. 9:21957751-21984490 (Ensembl ID); a protein encoded by cDNA sequence no. NM_000077 (GenBank) or NM_058195 9GenBank) or; or a protein having the GenBank sequence no. NP_000068 or NP_478102.

The compounds of the present invention can also be used to treat hyperproliferative disorders such as thyroid hyperplasia (especially Grave's disease), and cysts (such as hypervascularity of ovarian stroma, characteristic of polycystic ovarian syndrome (Stein-Leventhal syndrome)).

The compounds of the present invention may be designated as follows in the application and figures.


Compound A	AMG 232
Compound B	AMG 2653149
Compound C	AM-7209
Compound 1	AMG 2112819
Compound 2	dabrafenib
Compound 3	PD0325901*
Compound 4	trametinib
Compound 5	AMG 511
Compound 6	panobinostat

Compound 7	not used
Compound 8	imatinib
Compound 9	erlotinib
Compound 10	lapatinib
Compound 11	cisplatin
Compound 12	cytarabine
Compound 13	AMG 900

*AMG 1009089 (also called herein 1009089 or Compound 3) is PD0325901.

The MDM2 inhibitors of the present invention include those disclosed in published PCT application WO2011/153,509. A particular compound disclosed in the application is AMG 232 (Example 362) having the structure and name shown below.


2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid




A particular synthesis of AMG 232 is set forth in U.S. provisional patent application number 61/833,196, filed June 10, 2013.

Procedures to Make Certain Intermediates and Starting Materials


Method for making



,




and




.

Step A. 2-(3-Chlorophenyl)-1-(4-chlorophenyl)ethanone




Sodium bis(trimethylsilyl)amide (1 M in tetrahydrofuran, 117 mL) was slowly added to a -78 °C solution of 2-(3-chlorophenyl) acetic acid (10 g, 58.6 mmol) in tetrahydrofuran (58 mL) over 1 hour. After stirring at -78 °C for 40 minutes, a solution of methyl 4-chlorobenzoate (10 g, 58.6 mmol) in tetrahydrofuran (35 mL) was added over a period of 10 minutes. The reaction was stirred at -78 °C for 3 hours then allowed to warm to 25 °C. After two hours at 25 °C, the reaction was quenched with saturated aqueous ammonium chloride solution, and most of the tetrahydrofuran was removed under reduced pressure. The residue was extracted with ethyl acetate (2 × 100 mL). The combined organic layers were washed with saturated sodium chloride solution, dried over sodium sulfate, filtered and the filtrate was concentrated. The product was recrystallized from ether/pentane to provide the title compound as a white solid. ^{1}H NMR (500 MHz, DMSO-d_6, δ ppm): 8.05 (m, 2H), 7.62 (m, 2H), 7.33 (m, 3H), 7.21 (br d, J = 7.3 Hz, 1H), 4.45 (s, 2H). MS (ESI) = 265.1 $[M + H]^+$.

Step B: Methyl 4-(3-chlorophenyl)-5-(4-chlorophenyl)-2-methyl-5-oxopentanoate



Methyl methacrylate (12.65 mL, 119 mmol) was added to a solution of 2-(3-chlorophenyl)-1-(4-chlorophenyl)ethanone (30 g, 113 mmol) in tetrahydrofuran (283 mL). Potassium *tert*-butoxide (1.27 g, 11.3 mmol) was then added and the reaction was stirred at room temperature for 2 days. The solvent was removed under a vacuum and replaced with 300 mL of ethyl acetate. The organic phase was washed with brine (50 mL), water (3 x 50 mL), and brine (50 mL). The organic phase was dried over magnesium sulfate, filtered and concentrated under a vacuum to afford methyl 4-(3-chlorophenyl)-5-(4-chlorophenyl)-2-methyl-5-oxopentanoate as an approximately 1:1 mixture of diastereomers. ^{1}H NMR (400 MHz, $CDCl_3$, δ ppm): 7.87 (m, 2H), 7.38 (m, 2H), 7.27-7.14 (series of m, 4H), 4.61 (m, 1H), 3.69 (s, 1.5H), 3.60 (s, 1.5 H), 2.45 (m, 1H), 2.34 (m, 1H), 2.10 (ddd, J = 13.9, 9.4, 5.5 Hz, 0.5H), 1.96 (ddd, J = 13.7, 9.0, 4.3 Hz, 0.5H), 1.22 (d, J = 7.0 Hz, 1.5H), 1.16 (d, J = 7.0, 1.5 H). MS (ESI) = 387.0 $[M + 23]^+$.

Step C: (3*S*, 5*R*,6*R*)-5-(3-Chlorophenyl)-6-(4-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one and (3*R*, 5*R*,6*R*)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one




Methyl 4-(3-chlorophenyl)-5-(4-chlorophenyl)-2-methyl-5-oxopentanoate (40 g, 104.0 mmol) was dissolved in 200 mL of anhydrous toluene and concentrated under a vacuum. The residue was placed under high vacuum for 2 hours before use. The compound was split into 2 x 20 g batches and processed as follows: methyl 4-(3-chlorophenyl)-5-(4-chlorophenyl)-2-methyl-5-oxopentanoate (20 g, 52.0 mmol) in anhydrous 2-propanol (104 mL) was treated with potassium *tert*-butoxide (2.33 g, 20.8 mmol) in a 250 mL glass hydrogenation vessel. $RuCl_2$(*S*-xylbinap)(*S*-DAIPEN) (0.191 g, 0.156 mmol, Strem Chemicals, Inc., Newburyport, MA) in 3.8 mL of toluene was added. After 1.5 hours, the vessel was pressurized to 50 psi (344.7 kPa) and purged with hydrogen five times and allowed to stir at room temperature. The reaction was recharged with additional hydrogen as needed. After 3 days, the reactions were combined and partitioned between 50% saturated ammonium chloride solution and ethyl acetate. The aqueous layer was extracted with ethyl acetate. The combined organic phases were washed with brine, dried over magnesium sulfate, filtered, and concentrated.

The crude product (predominantly, (4*R*,5*R*)-isopropyl 4-(3-chlorophenyl)-5-(4-chlorophenyl)-5-hydroxy-2-methylpentanoate) was dissolved in tetrahydrofuran (450 mL) and methanol (150 mL). Lithium hydroxide (1.4 M, 149 mL, 208 mmol) was added, and the solution was stirred at room temperature for 24 hours. The mixture was concentrated under a vacuum and the residue was redissolved in ethyl acetate. Aqueous 1N hydrochloric acid was added with stirring until the aqueous layer had a pH of about 1. The layers were separated and the organic phase was washed with brine, dried over magnesium sulfate, filtered and concentrated. The material was dissolved in 200 mL of anhydrous toluene and treated with pyridinium *p*-toluenesulfonate (PPTS, 0.784 g, 3.12 mmol). The reaction was heated to reflux under Dean-Stark conditions until the seco-acid was consumed (about 2 hours). The reaction was cooled to room temperature and washed with saturated sodium bicarbonate (50 mL) and brine (50 mL). The solution was dried over sodium sulfate, filtered and concentrated. The crude material was purified by flash chromatography on silica gel (120 g column; eluting with 100%

dichloromethane). The title compounds were obtained as a white solid with an approximate 94:6 enantiomeric ratio and a 7:3 mixture of methyl diastereomers. ^{1}H NMR (400 MHz, $CDCl_3$, δ ppm): 7.22-6.98 (series of m, 5H), 6.91 (dt, *J* = 7.4, 1.2 Hz, 0.3H), 6.81 (m, 2H), 6.73 (dt, *J* = 7.6, 1.4 Hz, 0.7H), 5.76 (d, *J* = 4.1 Hz, 0.3 H), 5.69 (d, *J* = 4.7 Hz, 0.7H), 3.67 (dt, *J* = 6.6, 4.3 Hz, 0.3H), 3.55 (td, *J* = 7.8, 4.7 Hz, 0.7 H), 2.96 (d of quintets, *J* = 13.5, 6.7 Hz, 0.7 H), 2.81 (m, 0.3 H), 2.56 (dt, *J* = 14.3, 8.0 Hz, 0.7 H), 2.32 (dt, *J* = 13.69, 7.0 Hz, 0.3 H), 2.06 (ddd, *J* = 13.7, 8.4, 4.1, 0.3 H), 1.85 (ddd, *J* = 14.1, 12.5, 7.4, 0.7 H), 1.42 (d, *J* = 7.0 Hz, 0.9 H), 1.41 (d, *J* = 6.7 Hz, 2.1H). MS (ESI) = 357.0 $[M + 23]^+$. $[\alpha]_D$ (22 °C, c = 1.0, CH_2Cl_2) = -31.9°; m.p. 98–99 °C.


Step D. (3*S*,5*R*,6*R*)-3-Allyl-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one



A solution of (3*S*, 5*R*,6*R*)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one and (3*R*,5*S*,6*S*)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one (4.5 g, 13.4 mmol) and allyl bromide (3.48 mL, 40.3 mmol) in tetrahydrofuran (22 mL) at -35 °C (acetonitrile/dry ice bath) was treated with a solution of lithium bis(trimethylsilyl)amide in tetrahydrofuran (1.0 M, 17.45 mL, 17.45 mmol). The reaction was allowed to warm to -5 °C over 1 hour and then was quenched with 50% saturated ammonium chloride. The reaction was diluted with 100 mL of ethyl acetate and the layers were separated. The organic phase was washed with brine, dried over magnesium sulfate, filtered and concentrated under a vacuum to afford the title compound as a white solid upon standing under a vacuum. Chiral SFC (92% CO_2, 8% methanol (20 mM ammonia), 5 mL/min, Phenomenex Lux-2 column (Phenomenex, Torrance, CA), 100 bar (10,000 kPa), 40 °C, 5 minute method) was used to determine that the compound had an enantiomeric ratio of 96:4. (Major enantiomer: title compound, retention time = 2.45 minutes, 96%; minor enantiomer (structure not shown, retention time = 2.12 min, 4%). The title compound was recrystallized by adding to heptane (4.7 g slurried in 40 mL) at reflux and 1.5 mL of toluene was added dropwise to solubilize. The solution was cooled to 0 °C. The white solid was filtered and rinsed with 20 mL of cold heptanes to afford a white powder. Chiral SFC (92% CO_2, 8% methanol, Phenomenex Lux-2 column, same method as above) indicated an enantiomeric ratio of 99.2:0.8. (major enantiomer, 2.45 min,


99.2%; minor enantiomer: 2.12 min, 0.8%) . ^{1}H NMR (400 MHz, $CDCl_3$, δ ppm): 7.24 (ddd, J = 8.0, 2.0, 1.2 Hz, 1H), 7.20–7.15 (series of m, 3H), 6.91 (t, J = 2.0 Hz, 1H), 6.78 (br d, J = 7.6 Hz, 1H), 6.60 (m, 2H), 5.84 (ddt, J = 17.6, 10.2, 7.4 Hz, 1H), 5.70 (d, J = 5.3 Hz, 1H), 5.21–5.13 (series of m, 2H), 3.82 (dt, J = 11.7, 4.5 Hz, 1H), 2.62 (ABX J_{AB} = 13.7 Hz, J_{AX} = 7.6 Hz, 1H), 2.53 (ABX, J_{AB} = 13.9 Hz, J_{BX} = 7.2 Hz, 1H). 1.99 (dd, J = 14.1, 11.9 Hz, 1H), 1.92 (ddd, J = 13.9, 3.9, 1.2 Hz, 1H). ^{13}C NMR ($CDCl_3$, 100 MHz, δ ppm): 175.9, 140.2, 134.5, 134.3, 134.0, 132.2, 129.8, 128.6, 128.0, 127.9, 127.8, 126.4, 119.9, 83.9, 44.5, 42.4, 40.7, 31.8, 26.1. MS (ESI) = 375.2 $[M + H]^+$. IR = 1730 cm^{-1}. $[\alpha]_D$ (24 °C, c = 1.0, CH_2Cl_2) = -191°. m.p. 111–114 °C.

Step E. (*S*)-2-((2*R*,3*R*)-2-(3-Chlorophenyl)-3-(4-chlorophenyl)-3-hydroxypropyl)-N-(*(S)*-1-hydroxy-3-methylbutan-2-yl)-2-methylpent-4-enamide




(3*S*,5*R*,6*R*)-3-Allyl-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one (113 g, 300.0 mmol) was combined with (*S*)-2-amino-3-methyllbutan-1-ol (93 g, 900.0 mmol) and the suspension was heated at 100 °C for 5 hours. The reaction mixture was cooled to room temperature, diluted with ethyl acetate (1000 mL) and washed with 1N hydrochloric acid (2X), water, and brine. The organic layer was dried over magnesium sulfate and concentrated under a vacuum to give the title compound as white solid which was used in next step without further purification.

Step F. (3*S*,5*S*,6*R*,8*S*)-8-allyl-6-(3-chlorophenyl)-5-(4-chlorophenyl)-3-isopropyl-8-methyl-2,3,5,6,7,8-hexahydrooxazolo[3,2-*a*]pyridin-4-ium trifluoromethanesulfonate



Trifluoromethanesulfonic anhydride (57 mL, 339 mmol) was added dropwise over 60 minutes via addition funnel to a solution of (*S*)-2-((2*R*,3*R*)-2-(3-chlorophenyl)-3-(4-chlorophenyl)-3-hydroxypropyl)-N-((*S*)-1-hydroxy-3-methylbutan-2-yl)-2-methylpent-4-enamide (73.7 g, 154 mmol) and 2,6-dimethylpyridine (78 mL, 678 mmol) in dichloromethane (700 mL) at -50 °C. The reaction mixture was stirred at -50 °C for one additional hour and concentrated under a vacuum to provide the title compound as a reddish solid which was used in next step without further purification.

Step G. (3*S*,5*R*,6*S*)-3-Allyl-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((*S*)-1-(isopropylthio)-3-methylbutan-2-yl)-3-methylpiperidin-2-one




(3*S*,5*S*,6*R*,8*S*)-8-Allyl-6-(3-chlorophenyl)-5-(4-chlorophenyl)-3-isopropyl-8-methyl-2,3,5,6,7,8-hexahydrooxazolo[3,2-*a*]pyridin-4-ium trifluoromethanesulfonate (736 mg, 1.242 mmol) was weighed into an oven dried 50 mL pear-bottom flask and dissolved in 20 mL dry toluene. The toluene was removed under a vacuum to remove trace water in the solid. The process was repeated twice, and the resulting residue was dried under a strong vacuum.

A solution of sodium isopropyl sulfide was prepared by adding potassium 2-methylpropan-2-olate (3.0 mL, 3.00 mmol, 1 M solution in tetrahydrofuran) to a solution of propane-2-thiol (331 mg, 4.35 mmol) in 8 mL dimethylformamide that had been prepared under nitrogen and cooled to 0 °C. The sulfide solution was allowed to stir at room temperature for five minutes and was cooled to 0°C. The dry (3*S*,5*S*,6*R*,8*S*)-8-allyl-6-(3-chlorophenyl)-5-(4-chlorophenyl)-3-isopropyl-8-methyl-2,3,5,6,7,8-hexahydrooxazolo[3,2-*a*]pyridin-4-ium trifluoromethanesulfonate (736 mg, 1.242 mmol) was dissolved in dimethylformamide (8 mL total) and transferred (3 transfers total) via syringe to the sulfide solution over the course of 5 minutes. After 5 minutes, the ice bath was removed and the pale orange solution was allowed to warm to room temperature.


After stirring overnight, the mixture was partitioned between ethyl acetate and saturated ammonium chloride solution. The aqueous phase was saturated in sodium chloride and back-

extracted three times. The combined organics were washed twice with saturated sodium bicarbonate, twice with brine, dried over sodium sulfate, filtered, and concentrated under a vacuum to provide a residue that was purified by silica gel column chromatography (80 g column, gradient elution of 0% to 50 % ethyl acetate in hexanes).

Method for making




Step A. (3*S*,5*R*,6*S*)-3-Allyl-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((*S*)-1-hydroxy-3-methylbutan-2-yl)-3-methylpiperidin-2-one




Lithium hydroxide hydrate (64.6 g, 1540 mmol) was added portionwise, over a 5 minute period, to a solution of (3*S*,5*S*,6*R*,8*S*)-8-allyl-6-(3-chlorophenyl)-5-(4-chlorophenyl)-3-isopropyl-8-methyl-2,3,5,6,7,8-hexahydrooxazolo[3,2-*a*]pyridin-4-ium trifluoromethanesulfonate (Step F above) dissolved in tetrahydrofuran (500 ml) and water (300 ml). The reaction mixture was stirred at room temperature for 1 hour and concentrated under a vacuum. The residue was dissolved in ethyl acetate (ca. 1.3 L) and the layers were separated. The organic layer was washed with 1N hydrochloric acid (ice cooled, with enough hydrochloric acid to protonate and remove any remaining 2,6-dimethylpyridine (300 mL x 2)), water and brine. The solvent was removed under a vacuum to give a residue which was purified by silica gel column chromatography (1500 g column, gradient elution of 0% to 50% ethyl acetate in hexanes. The product was also crystallized from cyclohexane.

Step B. (3*S*,5*S*,6*R*,8*S*)-8-Allyl-6-(3-chlorophenyl)-5-(4-chlorophenyl)-3-isopropyl-8-methyl-2,3,5,6,7,8-hexahydrooxazolo[3,2-*a*]pyridin-4-ium 4-methylbenzenesulfonate



(3*S*,5*R*,6*S*)-3-Allyl-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((*S*)-1-hydroxy-3-methylbutan-2-yl)-3-methylpiperidin-2-one (49.77 g, 98 mmol) was transferred to a 1000 mL flask containing 4-methylbenzenesulfonic acid hydrate (19.27 g, 101 mmol) and a stirring bar. The reactants were suspended in toluene (230 mL). The flask was equipped with a Dean Stark trap and reflux condenser, and the stirred mixture was heated at reflux in a preheated bath. After 1 hour, the solvent was carefully removed under a vacuum and the resulting residue was further dried under high vacuum. The title compound was taken to the next step without purification.

Step C. (3*S*,5*R*,6*S*)-3-Allyl-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((*S*)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methylpiperidin-2-one




(3*S*,5*S*,6*R*,8*S*)-8-Allyl-6-(3-chlorophenyl)-5-(4-chlorophenyl)-3-isopropyl-8-methyl-2,3,5,6,7,8-hexahydrooxazolo[3,2-*a*]pyridin-4-ium 4-methylbenzenesulfonate, dry, powdered potassium carbonate (26.9 g, 195 mmol) and propane-2-thiol (14 ml, 150 mmol) were added along with 200 mL freshly sparged dimethylformamide. The mixture was heated under argon at 50 °C. After about 21 hours, a solution of *meta*-chloroperbenzoic acid (68.2 g, 77% pure by weight, in 100 mL dimethylformamide) was transferred to a dropping funnel and rapidly added to the stirred reaction mixture while the flask was immersed in an ice bath. After 5 minutes, the resulting yellow solution was allowed to warm to room temperature. After 10 minutes, additional *meta*-chloroperbenzoic acid (12 g, 77% wt %) was added as a solid and the mixture was stirred at room temperature. Upon completion, the mixture was poured into ethyl acetate and washed with 1 M sodium hydroxide (500 mL) that had been poured into ice. The aqueous phase was back-extracted three times and washed with additional 1 M NaOH ((500 mL, also poured into ice). The aqueous layer was washed once with ethyl acetate and the organics were combined. Sodium

thiosulfate (1 M in water, 250 mL) was added to the organics in a large Erlenmeyer flask, and the mixture was stirred for twenty minutes. The organic phase was washed again with sodium thiosulfate (1 M in water, 250 mL) and the mixture was allowed to stand over the weekend. The organics were concentrated to ca. 500 mL, then sequentially washed with 10% aqueous citric acid, 1 M sodium hydroxide, and brine. The organics were dried over sodium sulfate, filtered, and concentrated to give the crude product. The residue was purified by flash column chromatography (1.5 kg silica gel column, gradient elution of 0% to 50% ethyl acetate in hexanes) to give the title compound as a white solid.

Synthesis of Compound AMG 232 (Alternative 1)

2-((3*R*,5*R*,6*S*)-5-(3-Chlorophenyl)-6-(4-chlorophenyl)-1-((*S*)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid




Ruthenium(III) chloride trihydrate (22 mg, 0.084 mmol) and sodium periodate (1.12 g, 5.24 mmol) were added to a mixture of (3*S*,5*R*,6*S*)-3-allyl-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((*S*)-1-(isopropylthio)-3-methylbutan-2-yl)-3-methylpiperidin-2-one (390 mg, 0.752 mmol) in acetonitrile (4.0 mL), carbon tetrachloride (4.0 mL),and water (6.0 mL). The resulting dark brown mixture was vigorously stirred at ambient temperature overnight. The mixture was filtered through a pad of diatomaceous earth, washing with ethyl acetate. The filtrate was partitioned between 2 M HCl and ethyl acetate. The aqueous phase was back-extracted twice with ethyl acetate, and the combined organics were washed with brine, dried over sodium sulfate, filtered, and concentrated under a vacuum to a residue that was purified by flash chromatography (40 g silica gel column, gradient elution of 0% to 15% isopropanol in hexanes). Fractions containing the desired product were combined, stripped of solvent, redissolved in minimal ACN/water, frozen, and lyophilized to give a white powder.

Subsequently, a mixture of (3*S*,5*R*,6*S*)-3-allyl-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((*S*)-1-(isopropylthio)-3-methylbutan-2-yl)-3-methylpiperidin-2-one (388 mg, 0.748 mmol), ruthenium(III) chloride trihydrate (19.56 mg, 0.075 mmol), and sodium periodate (1.15 g, 5.38

mmol) in acetonitrile (4 mL), carbon tetrachloride (4.00 mL), and water (4.00 mL) was vigorously stirred at ambient temperature. After four hours, the mixture was filtered through a pad of diatomaceous earth, and the filtrate was partitioned between ethyl acetate and 2 M HCl. The aqueous phase was back-extracted twice with ethyl acetate, and the combined organics were washed with brine, dried over sodium sulfate, filtered, and concentrated under a vacuum to a residue. The residue was purified by flash chromatography (40 g silica gel column, gradient elution of 0% to 15% isopropanol in hexanes). Fractions containing the product were concentrated and combined with the solid obtained in the prior experiment. The combined material was dissolved in minimal acetonitrile/water, frozen, and lyophilized overnight to give a white solid.

Synthesis of AMG 232 (Alternative 2)

2-((3*R*,5*R*,6*S*)-5-(3-Chlorophenyl)-6-(4-chlorophenyl)-1-((*S*)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid




Sodium periodate (2.85 g, 13.32 mmol) and ruthenium(III) chloride trihydrate (0.049 g, 0.189 mmol) were added to a mixture of (3*S*,5*R*,6*S*)-3-allyl-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((*S*)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methylpiperidin-2-one (1.73 g, 3.14 mmol) in acetonitrile (18 mL), carbon tetrachloride (18 mL), and water (27 mL). The mixture was stirred vigorously at room temperature for 25 hours. The mixture was diluted with 2M HCl and filtered through a pad of diatomaceous earth and rinsed with ethyl acetate. The organic layer was separated, washed with brine, dried over sodium sulfate, filtered, and concentrated under a vacuum. The material was purified twice by flash chromatography (120g silica gel, gradient elution of 0% to 20% isopropanol in hexanes; 120 g column, gradient elution of 0% to 15% gradient isopropanol in hexanes). It was purified once more by flash chromatography (220 g silica gel; gradient elution 0% to 20% isopropanol in hexanes, 45 minutes) using a method in which the purest fractions were concentrated and set aside and mixed fractions were pooled and resubjected to the chromatography.

Subsequently, a mixture of (3*S*,5*R*,6*S*)-3-allyl-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((*S*)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methylpiperidin-2-one (4.1 g, 7.45 mmol), ruthenium(III) chloride trihydrate (0.120 g, 0.459 mmol), and sodium periodate (6.73 g, 31.5 mmol) in acetonitrile (40 mL), carbon tetrachloride (40 mL), and water (60 mL) was vigorously stirred at ambient temperature for 23 hours. The reaction was diluted by addition of 2 M aqueous HCl and filtered through a diatomaceous earth pad, washing with copious ethyl acetate. Most of the organics were removed under a vacuum. The crude product was extracted into ethyl acetate, washed with brine, dried over sodium sulfate, filtered, and concentrated to a residue that was purified twice by flash chromatography (330 g silica gel column, gradient elution of 0% to 20% isopropanol in hexanes; 330 g silica gel column, gradient elution of 0% to 20% isopropanol in hexanes) to give an off-white foam. The material was purified by flash chromatography three additional times (220 g silica gel column; gradient elution 0% to 20% isopropanol in hexanes, 45 minutes) using a method in which the purest fractions were concentrated and set aside and mixed fractions were pooled and resubjected to the chromatography.

Mixed fractions from both experiments were combined and purified by flash chromatography twice more (220 g silica gel column; gradient elution 0% to 20% isopropanol in hexanes, 45 minutes), and again the pure fractions were set aside.
All of the pure fractions were combined, concentrated under a vacuum, dissolved in minimal acetonitrile/water and lyophilized.

Synthesis of AMG 232 (Alternative 3)

2-((3*R*,5*R*,6*S*)-5-(3-Chlorophenyl)-6-(4-chlorophenyl)-1-((*S*)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid




(3*S*,5*R*,6*S*)-3-Allyl-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((*S*)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methylpiperidin-2-one (5.05 g, 9.17 mmol) was weighed into a 500 mL round bottom flask containing a large stir bar and 2.04 g sodium periodate (2.04 g). The mixture was diluted with carbon tetrachloride (52 mL), acetonitrile, (52 mL) and water (78 mL). The

flask was immersed in a room temperature water bath and the internal temperature was monitored with a digital thermocouple.

Ruthenium chloride hydrate (approximately 50 mg) was added in a single portion. The internal temperature rose to 22 °C, then ice was added to the bath to cool the mixture. Additional ruthenium chloride hydrate (25 mg) was added 3 minutes later. After stirring for a total of thirty minutes, Three portions of sodium periodate (2.08 g, 2.07 g and 2.08 g) were slowly added on 15 minute intervals. The temperature was kept below 19 °C, and ice was quickly added to the bath if the internal temperature began to rise. The mixture was stirred at ambient temperature overnight. The mixture was filtered through a pad of diatomaceous earth and the filter cake was washed copiously with ethyl acetate. The filtrate was concentrated under a vacuum and partitioned between 2 M HCl (100 mL) and ethyl acetate (200 mL).

Two rounds of flash column chromatography (330 g silica gel, then 220 g silica gel, gradient elution of 0% to 20% isopropanol in hexanes) provided the title compound. A portion of this material was lyophilized from acetonitrile and water. The less pure fractions were repurified by two additional rounds of flash column chromatography (220 g then 330 g silica gel columns, gradient elution of 0% to 20% isopropanol in hexanes). The most pure fractions from both runs were combined, concentrated under a vacuum and lyophilized from acetonitrile and water to give the title compound.


Another particular MDM2 inhibitor is AM-7209 (Compound C herein), which is disclosed in U.S. provisional patent application number 61/770,901, filed February 28, 2013. (See Example No. 5 therein and below). AM-7209 has the following chemical name and structure: **4-(2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid**




.

EXAMPLE 1

2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-Butylsulfonyl)-1-cyclopropylethyl)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetic acid (Example 351 of WO2011/153509 (Amgen Inc.), published December 8, 2011.




Step A. 2-(3-Chlorophenyl)-1-(4-chlorophenyl)ethanone



Sodium bis(trimethylsilyl)amide (1 M in tetrahydrofuran, 117 mL) was slowly added to a -78 °C solution of 2-(3-chlorophenyl) acetic acid (10 g, 58.6 mmol) in tetrahydrofuran (58 mL) over 1 hour. After stirring at -78 °C for 40 minutes, a solution of methyl 4-chlorobenzoate (10 g, 58.6 mmol) in tetrahydrofuran (35 mL) was added over a period of 10 minutes. The reaction was stirred at -78 °C for 3 hours then allowed to warm to 25 °C. After two hours at 25 °C, the reaction was quenched with saturated aqueous ammonium chloride solution, and most of the tetrahydrofuran was removed under reduced pressure. The residue was extracted with ethyl acetate (2 × 100 mL). The combined organic layers were washed with saturated sodium chloride solution, dried over sodium sulfate, filtered and the filtrate was concentrated. The product was recrystallized from ether/pentane to provide the title compound as a white solid. ^{1}H NMR (500 MHz, DMSO-d_6, δ ppm): 8.05 (m, 2H), 7.62 (m, 2H), 7.33 (m, 3H), 7.21 (br d, J = 7.3 Hz, 1H), 4.45 (s, 2H). MS (ESI) = 265.1 $[M + H]^+$.


Step B: Methyl 4-(3-chlorophenyl)-5-(4-chlorophenyl)-2-methyl-5-oxopentanoate



Methyl methacrylate (12.65 mL, 119 mmol) was added to a solution of 2-(3-chlorophenyl)-1-(4-chlorophenyl)ethanone (30 g, 113 mmol, Example 1, Step A) in tetrahydrofuran (283 mL). Potassium *tert*-butoxide (1.27 g, 11.3 mmol) was then added and the reaction was stirred at room temperature for 2 days. The solvent was removed under a vacuum and replaced with 300 mL of ethyl acetate. The organic phase was washed with brine (50 mL), water (3 x 50 mL), and brine (50 mL). The organic phase was dried over magnesium sulfate, filtered and concentrated under a vacuum to afford methyl 4-(3-chlorophenyl)-5-(4-chlorophenyl)-2-methyl-5-oxopentanoate as an

approximately 1:1 mixture of diastereomers. ^{1}H NMR (400 MHz, $CDCl_3$, δ ppm): 7.87 (m, 2H), 7.38 (m, 2H), 7.27-7.14 (series of m, 4H), 4.61 (m, 1H), 3.69 (s, 1.5H), 3.60 (s, 1.5 H), 2.45 (m, 1H), 2.34 (m, 1H), 2.10 (ddd, J = 13.9, 9.4, 5.5 Hz, 0.5H), 1.96 (ddd, J = 13.7, 9.0, 4.3 Hz, 0.5H), 1.22 (d, J = 7.0 Hz, 1.5H), 1.16 (d, J = 7.0, 1.5 H). MS (ESI) = 387.0 $[M + 23]^+$.


Step C: (3*S*, 5*R*,6*R*)-5-(3-Chlorophenyl)-6-(4-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one and (3*R*, 5*R*,6*R*)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one



Methyl 4-(3-chlorophenyl)-5-(4-chlorophenyl)-2-methyl-5-oxopentanoate (40 g, 104.0 mmol, Example 1, Step B) was dissolved in 200 mL of anhydrous toluene and concentrated under a vacuum. The residue was placed under high vacuum for 2 hours before use. The compound was split into 2 x 20 g batches and processed as follows: methyl 4-(3-chlorophenyl)-5-(4-chlorophenyl)-2-methyl-5-oxopentanoate (20 g, 52.0 mmol) in anhydrous 2-propanol (104 mL) was treated with potassium *tert*-butoxide (2.33 g, 20.8 mmol) in a 250 mL glass hydrogenation vessel. $RuCl_2$(*S*-xylbinap)(*S*-DAIPEN) (0.191 g, 0.156 mmol, Strem Chemicals, Inc., Newburyport, MA) in 3.8 mL of toluene was added. After 1.5 hours, the vessel was pressurized to 50 psi (344.7 kPa) and purged with hydrogen five times and allowed to stir at room temperature. The reaction was recharged with additional hydrogen as needed. After 3 days, the reactions were combined and partitioned between 50% saturated ammonium chloride solution and ethyl acetate. The aqueous layer was extracted with ethyl acetate. The combined organic phases were washed with brine, dried over magnesium sulfate, filtered, and concentrated. The crude product (predominantly, (4*R*,5*R*)-isopropyl 4-(3-chlorophenyl)-5-(4-chlorophenyl)-5-hydroxy-2-methylpentanoate) was dissolved in tetrahydrofuran (450 mL) and methanol (150 mL). Lithium hydroxide (1.4 M, 149 mL, 208 mmol) was added, and the solution was stirred at room temperature for 24 hours. The mixture was concentrated under a vacuum and the residue was redissolved in ethyl acetate. Aqueous 1N hydrochloric acid was added with stirring until the aqueous layer had a pH of about 1. The layers were separated and the organic phase was washed with brine, dried over magnesium sulfate, filtered and concentrated. The material was dissolved in 200 mL of anhydrous toluene and treated with pyridinium *p*-toluenesulfonate (PPTS, 0.784 g, 3.12 mmol). The reaction was heated to reflux under Dean-Stark conditions until the seco-acid was consumed (about 2 hours). The reaction was cooled to room temperature and washed with

saturated sodium bicarbonate (50 mL) and brine (50 mL). The solution was dried over sodium sulfate, filtered and concentrated. The crude material was purified by flash chromatography on silica gel (120 g column; eluting with 100% dichloromethane). The title compounds were obtained as a white solid with an approximate 94:6 enantiomeric ratio and a 7:3 mixture of methyl diastereomers. ^{1}H NMR (400 MHz, $CDCl_3$, δ ppm): 7.22-6.98 (series of m, 5H), 6.91 (dt, J = 7.4, 1.2 Hz, 0.3H), 6.81 (m, 2H), 6.73 (dt, J = 7.6, 1.4 Hz, 0.7H), 5.76 (d, J = 4.1 Hz, 0.3 H), 5.69 (d, J = 4.7 Hz, 0.7H), 3.67 (dt, J = 6.6, 4.3 Hz, 0.3H), 3.55 (td, J = 7.8, 4.7 Hz, 0.7 H), 2.96 (d of quintets, J = 13.5, 6.7 Hz, 0.7 H), 2.81 (m, 0.3 H), 2.56 (dt, J = 14.3, 8.0 Hz, 0.7 H), 2.32 (dt, J = 13.69, 7.0 Hz, 0.3 H), 2.06 (ddd, J = 13.7, 8.4, 4.1, 0.3 H), 1.85 (ddd, J = 14.1, 12.5, 7.4, 0.7 H), 1.42 (d, J = 7.0 Hz, 0.9 H), 1.41 (d, J = 6.7 Hz, 2.1H). MS (ESI) = 357.0 $[M + 23]^+$. $[\alpha]_D$ (22 °C, c = 1.0, CH_2Cl_2) = -31.9°; m.p. 98–99 °C.


Step D. (3*S*,5*R*,6*R*)-3-Allyl-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one



A solution of (3*S*, 5*R*,6*R*)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one and (3*R*,5*S*,6*S*)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one (4.5 g, 13.4 mmol, Example 1, Step C) and allyl bromide (3.48 mL, 40.3 mmol) in tetrahydrofuran (22 mL) at -35 °C (acetonitrile/dry ice bath) was treated with a solution of lithium bis(trimethylsilyl)amide in tetrahydrofuran (1.0 M, 17.45 mL, 17.45 mmol). The reaction was allowed to warm to -5 °C over 1 hour and then was quenched with 50% saturated ammonium chloride. The reaction was diluted with 100 mL of ethyl acetate and the layers were separated. The organic phase was washed with brine, dried over magnesium sulfate, filtered and concentrated under a vacuum to afford the title compound as a white solid upon standing under a vacuum. Chiral SFC (92% CO_2, 8% methanol (20 mM ammonia), 5 mL/min, Phenomenex Lux-2 column (Phenomenex, Torrance. CA), 100 bar (10,000 kPa), 40 °C, 5 minute method) was used to determine that the compound had an enantiomeric ratio of 96:4. (Major enantiomer: title compound, retention time = 2.45 minutes, 96%; minor enantiomer (structure not shown, retention time = 2.12 min, 4%). The title compound was recrystallized by adding to heptane (4.7 g slurried in 40 mL) at reflux and 1.5 mL of toluene was added dropwise to solubilize. The solution was cooled to 0 °C. The white solid was filtered and rinsed with 20 mL of cold heptanes


to afford a white powder. Chiral SFC (92% CO_2, 8% methanol, Phenomenex Lux-2 column, same method as above) indicated an enantiomeric ratio of 99.2:0.8. (major enantiomer, 2.45 min, 99.2%; minor enantiomer: 2.12 min, 0.8%) . ^{1}H NMR (400 MHz, $CDCl_3$, δ ppm): 7.24 (ddd, J = 8.0, 2.0, 1.2 Hz, 1H), 7.20–7.15 (series of m, 3H), 6.91 (t, J = 2.0 Hz, 1H), 6.78 (br d, J = 7.6 Hz, 1H), 6.60 (m, 2H), 5.84 (ddt, J = 17.6, 10.2, 7.4 Hz, 1H), 5.70 (d, J = 5.3 Hz, 1H), 5.21–5.13 (series of m, 2H), 3.82 (dt, J = 11.7, 4.5 Hz, 1H), 2.62 (ABX J_{AB} = 13.7 Hz, J_{AX} = 7.6 Hz, 1H), 2.53 (ABX, J_{AB} = 13.9 Hz, J_{BX} = 7.2 Hz, 1H). 1.99 (dd, J = 14.1, 11.9 Hz, 1H), 1.92 (ddd, J = 13.9, 3.9, 1.2 Hz, 1H). ^{13}C NMR ($CDCl_3$, 100 MHz, δ ppm): 175.9, 140.2, 134.5, 134.3, 134.0, 132.2, 129.8, 128.6, 128.0, 127.9, 127.8, 126.4, 119.9, 83.9, 44.5, 42.4, 40.7, 31.8, 26.1. MS (ESI) = 375.2 $[M + H]^+$. IR = 1730 cm^{-1}. $[\alpha]_D$ (24 °C, c = 1.0, CH_2Cl_2) = -191°. m.p. 111–114 °C.

Step E. (2*S*)-2-((2*R*)-2-(3-Chlorophenyl)-3-(4-chlorophenyl)-3-hydroxypropyl)-*N*-((*S*)-1-cyclopropyl-2-hydroxyethyl)-2-methylpent-4-enamide



(3*S*,5*R*,6*R*)-3-Allyl-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one (125.0 g, 333 mmol, Example 1, Step D) was added to (*S*)-2-amino-2-cyclopropylethanol (101 g, 999 mmol) and the reaction mixture was heated at 110 °C under argon for 25 hours. The reaction mixture was diluted with isopropyl acetate, cooled to room temperature, and 3 M hydrochloric acid (400 mL) was added slowly. The mixture was stirred at room temperature for 20 minutes, and the layers were separated. The organic layer was washed with 1 M hydrochloric acid (200 mL) and brine, then dried over magnesium sulfate, filtered and concentrated under a vacuum to provide the desired product as a brown oil (159 g).


Step F. (3*S*,5*S*,6*R*,8*S*)-8-Allyl-6-(3-chlorophenyl)-5-(4-chlorophenyl)-3-cyclopropyl-8-methyl-2,3,5,6,7,8-hexahydrooxazolo[3,2-*a*]pyridin-4-ium 4-methylbenzenesulfonate



A 2 L 4-necked round-bottomed flask equipped with a magnetic stir bar, addition funnel, septa and internal temperature sensor was charged with *p*-toluenesulfonic anhydride (240 g, 734 mmol) and anhydrous dichloromethane (600 mL). The internal temperature was adjusted to 14 °C and the mixture was stirred for 10 minutes. A solution of (*S*)-2-((2*R*,3*R*)-2-(3-chlorophenyl)-3-(4-chlorophenyl)-3-hydroxypropyl)-*N*-((*S*)-1-cyclopropyl-2-hydroxyethyl)-2-methylpent-4-enamide (159.0 g, 334 mmol, Example 1, Step E) in anhydrous dichloromethane (400 mL) was added to the reaction mixture. The temperature increased to 17 °C before returning to 14 °C. The reaction mixture was cooled to 7 °C and 2,6-lutidine (160 mL, 1372 mmol) (dried over activated 4 Å molecular sieves) was added dropwise via addition funnel to the reaction mixture. The addition was complete after 1 hour. The reaction mixture was removed from the water bath and stirred at room temperature for 1 hour. The reaction mixture was heated at reflux for 16 hours. LCMS indicated that some intermediate remained. Additional *p*-toluenesulfonic anhydride (0.25 equiv) and lutidine (0.5 equiv) were added and the reaction mixture was heated at reflux for 8 hours. LCMS indicated that the reaction was complete. The reaction mixture was cooled to room temperature and added via addition funnel to 1 M aqueous sulfuric acid (764 mL, 764 mmol) with stirring. The addition took 30 minutes, and the solution was stirred at room temperature for 30 minutes thereafter. The layers were separated and the organic layer was dried over magnesium sulfate, filtered and concentrated under a vacuum to provide a brown syrup. To remove any dichloromethane from the syrup it was taken up in ethyl acetate and concentrated under a vacuum twice to provide a thick brown syrup. Ethyl acetate (2 L) was added and the mixture was heated at 60 °C until all of the syrup was dissolved (about 45 minutes). The solution was stirred while cooling to room temperature. Crystals had formed after 2 hours and the mixture was cooled to 10 °C for 1 hour before collecting the solid by vacuum filtration and washing with cold (10 °C) ethyl acetate. This provided 70 g of the desired product as an off-white crystalline solid. The filtrate was concentrated under a vacuum to 1.5 L and the mixture was stirred at 10 °C for 1.5 hours. The mixture was filtered under vacuum to provide a light brown crystalline solid that was shown to be lutidinium tosylate by NMR. The filtrate was concentrated under vacuum to provide a brown syrup (161g). Heptane was added to the syrup and the mixture was heated. A minimal amount of ethyl acetate was added until the material dissolved. The solution was cooled to room temperature and then placed in the freezer. The resulting solid was collected by vacuum filtration and washed with cold (0 °C) ethyl acetate to provide the desired product as an off-white crystalline solid (34g). The filtrate was concentrated to provide a dark brown oil and purified by flash chromatography on silica gel (1.5 kg SiO_2 column, gradient elution of 20% to 100% acetone in hexanes) to provide the desired product as a light brown syrup (73 g). ^{1}H NMR (500 MHz,


$CDCl_3$, δ ppm): -0.3 to -0.2 (m, 2H), 0.06–0.11 (m, 1H), 0.31–0.36 (m, 1H), 0.38–0.43 (m, 1H), 1.57 (s, 3H), 1.91 (dd, J = 3.7 and 13.9 Hz, 1H), 2.36 (s, 3H), 2.64 (dd, J = 7.3 and 13.7 Hz, 1H), 2.72 (dd, J = 7.6 and 13.7 Hz, 1H), 2.95 (t, J = 13.9 Hz, 1H), 3.32 (dt, J = 3.7 and 10.8 Hz, 1H), 4.47 (t, J = 8.6 Hz, 1H), 4.57–4.62 (m, 1H), 5.32 (d, J = 16.9 Hz, 1H), 5.35 (d, J = 10.3 Hz, 1H), 5.46 (t, J = 9.5 Hz, 1H), 5.82 (d, J = 10.5 Hz, 1H), 5.84–5.93 (m, 1H), 6.94 (br s, 1H), 7.04 (s, 1H), 7.14–7.20 (m, 5H), 7.28–7.40 (m, 3H), 7.88 (d, J = 8.1 Hz, 2H)). MS (ESI) 440.1 $[M+H]^+$.

Step G. (3*S*,5*R*,6*S*)-3-Allyl-1-((*S*)-2-(*tert*-butylthio)-1-cyclopropylethyl)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methylpiperidin-2-one



2-Methyl-2-propanethiol (0.195 mL, 1.796 mmol, dried over activated 4 Å molecular sieves) was added to a solution of lithium bis(trimethylsilyl)amide in tetrahydrofuran (1.0 M, 1.8 mL, 1.8 mmol) in anhydrous tetrahydrofuran (4 mL) at room temperature. The reaction mixture was heated at 60 °C. After 15 minutes at 60 °C, (3*S*,5*S*,6*R*,8*S*)-8-allyl-6-(3-chlorophenyl)-5-(4-chlorophenyl)-3-cyclopropyl-8-methyl-2,3,5,6,7,8-hexahydrooxazolo[3,2-*a*]pyridin-4-ium 4-methylbenzenesulfonate (1.00 g, 1.632 mmol, Example 1, Step F) was added as a solid. The reaction mixture was heated at 60 °C for 12 hours and then cooled to room temperature and diluted with water. The solution was extracted with ethyl acetate thrice and the organics were pooled, washed with brine, dried over sodium sulfate, decanted and concentrated under a vacuum to provide a brown oil. Purification by flash chromatography (80 g SiO_2 column, gradient elution of 10 to 60% ethyl acetate in hexanes provided the desired product as a colorless syrup. ^{1}H NMR (500 MHz, $CDCl_3$, δ ppm): -0.88 to -0.85 (m, 1H), -0.16 to -0.13 (m, 1H), 0.22–0.27 (m, 1H), 0.39–0.44 (m, 1H), 1.28 (s, 3H), 1.35 (s, 9H), 1.66–1.71 (m, 1H), 1.86 (dd, J = 3.2 and 13.5 Hz, 1H), 2.16 (t, J = 13.7, 1H), 2.21–2.27 (m, 1H), 2.60 (dd, J = 4.4 and 12.0 Hz, 1H), 2.65 (d, J = 7.6 Hz, 2H), 3.12 (dt, J = 3.2 and 10.3 Hz, 1H), 3.60 (t, J = 11.3 Hz, 1H), 4.68 (d, J = 10.3 Hz, 1H), 5.16–5.19 (m, 2H), 5.83–5.92 (m, 1H), 6.79 (d, J = 7.6 Hz, 1H), 6.93–7.04 (m, 3H), 7.09–7.16 (m, 2H), 7.19–7.24 (m, 2H). MS (ESI) 530.2 $[M+H]^+$.


Step H. 2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-Butylsulfonyl)-1-cyclopropylethyl)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetic acid



Ruthenium(III) chloride hydrate (30.0 mg, 0.135 mmol) was added to a solution of (3*S*,5*R*,6*S*)-3-allyl-1-((*S*)-2-(*tert*-butylthio)-1-cyclopropylethyl)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methylpiperidin-2-one (3.25 g, 6.13 mmol, Example 1, Step H) and sodium periodate (1.33 g) in ethyl acetate (12 mL), acetonitrile (12 mL) and water (18 mL) at 18 °C. The temperature rose to 25 °C upon addition. Additional sodium periodate was added in five 1.33 g portions over 30 minutes while maintaining the temperature below 22 °C. LCMS after 1.5 hours indicated that the reaction was incomplete, and sodium periodate (1 equivalent) was added. After 1.5 hours the reaction mixture was vacuum filtered, washed with ethyl acetate, and the layers were separated. The aqueous layer was extracted with ethyl acetate and the organics were combined, washed with brine, dried over magnesium sulfate, filtered and concentrated under a vacuum to provide a green oil. Purification by flash chromatography (330 g SiO_2 column, gradient elution of 0% to 20% isopropanol in hexanes provided the title compound as a white solid. ^{1}H NMR (500 MHz, $CDCl_3$, δ ppm): -1.15 to -1.05 (m, 1H), -0.35 to -0.25 (m, 1H), 0.18–0.28 (m, 1H), 0.33–0.40 (m, 1H), 1.45 (s, 9H), 1.51 (s, 3H), 1.86 (dd, J = 2.7 and 13.7 Hz, 1H), 1.87–1.93 (m, 1H), 2.47 (t, J = 13.9, 1H), 2.72–2.76 (m, 1H), 2.76 (d, J = 15.5 Hz, 1H), 2.93 (d, J = 13.7 Hz, 1H), 3.12 (d, J = 15.1 Hz, 1H), 3.12 (dt, J = 2.7 and 12.5 Hz, 1H), 4.29 (t, J = 11.5 Hz, 1H), 4.95 (d, J = 10.8 Hz, 1H), 6.86–6.89 (m, 1H), 6.96 (br s, 1H), 7.08–7.14 (m, 3H), 7.15–7.35 (m, 3H). MS (ESI) 580.2 $[M + H]^+$.

EXAMPLE 2


2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-Butylsulfonyl)-1-cyclopropylethyl)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamide



Oxalyl chloride (0.033 mL, 0.379 mmol) was added to a solution of 2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-butylsulfonyl)-1-cyclopropylethyl)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetic acid (0.200 g, 0.344 mmol, Example 1, Step H) in anhydrous dichloromethane (1.5 mL) at room temperature. The reaction mixture was stirred at room temperature for 1 hour and then concentrated under a vacuum to provide the acid chloride as a white foam (206 mg). Lithium bis(trimethylsilyl)amide (1.0 M in tetrahydrofuran, 0.516 mL, 0.516 mmol) and anhydrous tetrahydrofuran (0.5 mL) were added at room temperature. The reaction mixture was stirred at room temperature for 5.5 hours and was then diluted with 1 N hydrochloric acid and extracted with ethyl acetate thrice. The organics were pooled, washed with brine, dried over sodium sulfate, decanted and concentrated under a vacuum to provide a yellow foam. Purification by flash chromatography (12 g SiO_2 column; gradient elution of 35% to 100% ethyl acetate) provided the title compound as an off-white foam. ^{1}H NMR (500 MHz, $CDCl_3$, δ ppm): -1.10 to -1.00 (m, 1H), -0.38 to -0.325 (m, 1H), 0.17–0.26 (m, 1H), 0.30–0.38 (m, 1H), 1.43 (s, 3H), 1.44 (s, 9H), 1.85–1.92 (m, 1H), 2.00 (dd, J = 2.7 and 13.5 Hz, 1H), 2.39 (t, J = 13.7, 1H), 2.65–2.75 (m, 1H), 2.73–2.80 (m, 2H), 2.90–2.96 (m, 1H), 3.31 (dt, J = 2.9 and 10.8 Hz, 1H), 4.30–4.38 (m, 1H), 4.96 (d, J = 10.8 Hz, 1H), 5.63 (br s, 1H), 6.64 (br s, 1H), 6.90–6.91 (m, 1H), 7.00 (s, 2H), 7.06–7.11 (m, 3H), 7.12–7.29 (m, 2H). MS (ESI) 579.2 $[M + H]^+$.

EXAMPLE 3

2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-Butylsulfonyl)-1-cyclopropylethyl)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)-*N*-phenylacetamide




N-(3-Dimethylaminopropyl)-*N*′-ethylcarbodiimide hydrochloride (EDC, 0.117 g, 0.612 mmol) was added to a solution of 2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-butylsulfonyl)-1-cyclopropylethyl)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetic acid (0.118 g, 0.204 mmol, Example 1, Step H) and aniline (0.020 mL, 0.225 mmol) at 0 °C. After the addition was complete, the reaction mixture was removed from the ice bath and stirred at room temperature for 19 hours. The reaction mixture was diluted with ice-cold 1 M hydrochloric acid to adjust the pH to 1 and the solution was extracted twice with ether. The combined organic layer was washed with brine, dried over sodium sulfate, decanted and concentrated under a vacuum to provide an


orange oil. Purification by flash chromatography (12 g SiO_2 column, gradient elution of 15% to 100% ethyl acetate in hexanes provided the title compound as a white foam. ^{1}H NMR (500 MHz, $CDCl_3$, δ ppm): -1.32 to -1.20 (m, 1H), -0.40 to -0.28 (m, 1H), -0.28 to -0.10 (m, 1H), 0.30–0.40 (m, 1H), 1.45 (s, 9H), 1.47 (s, 3H), 1.94 (br s, 1H), 2.07 (dd, J = 2.7 and 13.7 Hz, 1H), 2.39 (t, J = 13.7, 1H), 2.67–2.73 (m, 2H), 2.95 (t, J = 13.5 Hz, 2H), 3.30 (dt, J = 2.7 and 11.0 Hz, 1H), 4.31 (br t, J = 11.7 Hz, 1H), 4.94 (d, J = 10.8 Hz, 1H), 6.86–6.89 (m, 1H), 6.99 (s, 1H), 7.02–7.09 (m, 6 H), 7.17 (t, J = 7.3 Hz, 1H), 7.38 (t, J = 8.3 Hz, 2H), 7.66 (d, J = 7.8 Hz, 2H). MS (ESI) 655.3 $[M + H]^+$.

EXAMPLE 4

2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-Butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetic acid




Step A. Methyl -4-chloro-3-fluorobenzoate




A solution of 4-chloro-3-fluoro benzoic acid (450.0 g, 2.586 mol, Fluorochem, Derbyshire, UK) in methanol (4.5 L) was cooled to 0 °C and thionyl chloride (450.0 mL) was added over 30 minutes. The reaction mixture was stirred for 12 hours at ambient temperature. The reaction was monitored by TLC. Upon completion, the solvent was removed under reduced pressure and the residue was quenched with 1.0 M sodium bicarbonate solution (500 mL). The aqueous layer was extracted with dichloromethane (2 x 5.0 L). The combined organic layer was washed with brine (2.5 L), dried over anhydrous sodium sulfate and concentrated under reduced pressure afforded the title compound as light brown solid. The crude compound was used in the next step without further purification. ^{1}H NMR (400 MHz, $CDCl_3$, δ ppm): 7.82–7.74 (m, 2H), 7.46 (dd, J = 8.2, 7.5 Hz, 1H), 3.92 (s, 3H).

Step B. 1-(4-chloro-3-fluorophenyl)-2-(3-chlorophenyl)ethanone



Sodium bis(trimethylsilyl)amide (1 M in tetrahydrofuran, 4 L, 4000 mmol) was added over 1 hour to a solution of 3-chlorophenyl acetic acid (250.0 g, 1465 mmol) in anhydrous tetrahydrofuran (1.75 L) at -78 °C under nitrogen. The resulting reaction mixture was stirred for an additional hour at -78 °C. Then, a solution of methyl-4-chloro-3-fluorobenzoate (221.0 g, 1175 mmol, Example 4, Step A) in tetrahydrofuran (500 mL) was added over 1 hour at -78 °C, and the resulting reaction mixture was stirred at the same temperature for 2 hours. The reaction was monitored by TLC. On completion, reaction mixture was quenched with 2 N hydrochloric acid (2.5 L) and aqueous phase was extracted with ethyl acetate (2 x 2.5 L). The combined organic layer was washed with brine (2.5 L), dried over anhydrous sodium sulfate and concentrated under reduced pressure to provide the crude material which was purified by flash column chromatography (silica gel: 100 to 200 mesh, product eluted in 2% ethyl acetate in hexane) to afford the title compound as a white solid. ^{1}H NMR (400 MHz, $CDCl_3$, δ ppm): 7.74 (ddd, J = 10.1, 8.9, 1.8 Hz, 2H), 7.56–7.48 (m, 1H), 7.26 (t, J = 6.4 Hz, 3H), 7.12 (d, J = 5.7 Hz, 1H), 4.22 (s, 2H). MS (ESI) 282.9 $[M + H]^+$.


Step C. Methyl 5-(4-chloro-3-fluorophenyl)-4-(3-chlorophenyl)-2-methyl-5-oxopentanoate




Methyl methacrylate (125.0 g, 1097 mmol) and potassium *tert*-butoxide (1 M in tetrahydrofuran, 115 mL, 115 mmol) were sequentially added to a solution of 1-(4-chloro-3-fluorophenyl)-2-(3-chlorophenyl)ethanone (327.0 g, 1160 mmol, Example 4, Step B) in anhydrous tetrahydrofuran (2.61 L), at 0 °C. The reaction mixture was stirred for 1 hour at 0 °C and then warmed to ambient temperature and stirred for 12 hours. On completion, the reaction was quenched with water (1.0 L) and extracted with ethyl acetate (2 x 2.5 L). The combined organic layer was washed with brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure to get the crude material which was purified by flash column chromatography (silica gel: 60 to 120 mesh, product eluted in 4% ethyl acetate in hexane) affording the title compound (mixture of diastereomers) as light yellow liquid. ^{1}H NMR (400 MHz, $CDCl_3$, δ ppm): 7.74–7.61 (m, 4H), 7.47–7.40 (m, 2H), 7.28–7.18 (m, 6H), 7.16–7.10 (m, 2H), 4.56 (m, 2H), 3.68 (s, 3H), 3.60 (s,

3H), 2.50–2.39 (m, 2H), 2.37–2.25 (m, 2H), 2.10–2.02 (m, 1H), 1.94 (ddd, J = 13.6, 9.1, 4.2 Hz, 1H), 1.21 (d, J = 7.0 Hz, 3H), 1.15 (d, J = 7.0 Hz, 3H). MS (ESI) 383.0 $[M + H]^+$.

Step D. (3*S*,5*R*,6*R*)-6-(4-Chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one and (3*R*,5*R*,6*R*)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one



and




In a 2000 mL reaction vessel charged with methyl 5-(4-chloro-3-fluorophenyl)-4-(3-chlorophenyl)-2-methyl-5-oxopentanoate (138.0 g, 360 mmol, Example 4, Step C) (which was cooled on ice for 10 minutes before transferring to a glove bag) anhydrous 2-propanol (500 mL), and potassium *tert*-butoxide (16.16 g, 144 mmol) were sequentially added while in a sealed glove bag under argon. This mixture was allowed to stir for 30 minutes. $RuCl_2$(*S*-xylbinap)(*S*-DAIPEN) (1.759 g, 1.440 mmol, Strem Chemicals, Inc., Newburyport, MA, weighed in the glove bag) in 30.0 mL toluene was added. The reaction was vigorously stirred at room temperature for 2 hours. The vessel was set on a hydrogenation apparatus, purged with hydrogen 3 times and pressurized to 50 psi (344.7 kPa). The reaction was allowed to stir overnight at room temperature. On completion, the reaction was quenched with water (1.5 L) and extracted with ethyl acetate (2 x 2.5 L). The organic layer was washed with brine (1.5 L), dried over anhydrous sodium sulfate and concentrated under reduced pressure to get crude material which was purified by flash column chromatography (silica gel; 60-120 mesh; product eluted in 12% ethyl acetate in hexane) to provide a dark colored liquid as a mixture of diastereomers.

The product was dissolved in (240.0 g, 581 mmol) in tetrahydrofuran (1.9 L) and methanol (480 mL), and lithium hydroxide monohydrate (2.5 M aqueous solution, 480.0 mL) was added. The reaction mixture was stirred at ambient temperature for 12 hours. On completion, the solvent was removed under reduced pressure and the residue was acidified with 2 N hydrochloric acid to a pH between 5 and 6. The aqueous phase was extracted with ethyl acetate (2 x 1.0 L). The combined organic layer was washed with brine (750 mL), dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure to provide a dark colored liquid, which was used without further purification.

A portion of the crude intermediate (25.4 g, predominantly seco acid) was added to a 500 mL round bottom flask, equipped with a Dean-Stark apparatus. Pyridinium *p*-toluenesulfonate (0.516

g, 2.053 mmol) and toluene (274 mL) were added, and the mixture was refluxed for 1 hour (oil bath temperature about 150 °C). The reaction was cooled to room temperature and concentrated under reduced pressure. The reaction was diluted with saturated aqueous sodium bicarbonate (150 mL), extracted with diethyl ether (2 × 150 mL), and washed with brine (150 mL). The combined organic layer was dried over magnesium sulfate, filtered and concentrated under reduced pressure. Purification by flash column chromatography (divided into 3 portions, 330 g SiO_2/each, gradient elution of 0% to 30% acetone in hexanes, 35 minutes) provided the title compounds as a pale yellow solid and a 1:1.6 mixture of diastereomers at C2. MS (ESI) 353.05 $[M + H]^+$.


Step E. (3*S*,5*R*,6*R*)-3-Allyl-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one



(3*S*,5*R*,6*R*)-6-(4-Chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one and (3*R*,5*R*,6*R*)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one (18 g, 51.0 mmol, Example 4, Step D) was added to an oven dried 500 mL round-bottom flask. The solid was dissolved in anhydrous toluene and concentrated to remove adventitious water. 3-Bromoprop-1-ene (11.02 mL, 127 mmol, passed neat through basic alumina prior to addition) in tetrahydrofuran (200 mL) was added and the reaction vessel was evacuated and refilled with argon three times. Lithium bis(trimethylsilyl)amide (1.0 M, 56.1 mL, 56.1 mmol) was added dropwise at -40 °C (dry ice/acetonitrile bath) and stirred under argon. The reaction was allowed to gradually warm to -10 °C and stirred at -10 °C for 3 hours. The reaction was quenched with saturated ammonium chloride (10 mL), concentrated, and the crude product was diluted in water (150 mL) and diethyl ether (200 mL). The layers were separated and the aqueous layer was washed twice more with diethyl ether (200 mL/each). The combined organic layer was washed with brine (100 mL), dried over magnesium sulfate, filtered, and concentrated under reduced pressure to a residue. The residue was purified by flash chromatography (2 x 330 g silica gel columns, gradient elution of 0% to 30% acetone in hexanes) to provide the title compound as a white solid. The product can alternatively be crystallized from a minimum of hexanes in dichloromethane. Enantiomeric excess was determined to be 87% by chiral SFC (90% CO_2, 10% methanol (20 mM ammonia), 5.0 mL/min, 100 bar (10,000 kPa), 40 °C, 5 minute method, Phenomenex Lux-2 (Phenomenex, Torrance, CA) (100 mm x 4.6 mm, 5 μm column), retention

times: 1.62 min. (minor) and 2.17 min. (major)). The purity could be upgraded to > 98% through recrystallization in hexanes and dichloromethane. ^{1}H NMR (400 MHz, $CDCl_3$, δ ppm): 7.24–7.17 (m, 3H), 6.94 (s, 1H), 6.80 (d, J = 7.5 Hz, 1H), 6.48 (dd, J = 10.0, 1.9 Hz, 1H), 6.40 (d, J = 8.3 Hz, 1H), 5.90–5.76 (m, 1H), 5.69 (d, J = 5.2 Hz, 1H), 5.20–5.13 (m, 2H), 3.81 (dd, J = 13.9, 6.9 Hz, 1H), 2.62 (dd, J = 13.8, 7.6 Hz, 1H), 2.50 (dd, J = 13.8, 7.3 Hz, 1H), 1.96 (d, J = 8.4 Hz, 2H), 1.40 (s, 3H). MS (ESI) 393.1 $[M + H]^+$.


Step F. (2*S*)-2-((2*R*)-3-(4-Chloro-3-fluorophenyl)-2-(3-chlorophenyl)-3-hydroxypropyl)-*N*-((*S*)-1-cyclopropyl-2-hydroxyethyl)-2-methylpent-4-enamide



Sodium methoxide (25% in methanol, 60.7 ml, 265 mmol) was added to a solution of (*S*)-2-amino-2-cyclopropylethanol hydrochloride (36.5 g, 265 mmol, NetChem Inc., Ontario, Canada) in methanol (177 mL) at 0 °C. A precipitate formed during the addition. After the addition was complete, the reaction mixture was removed from the ice bath and warmed to room temperature. The reaction mixture was filtered under a vacuum and the solid was washed with dichloromethane. The filtrate was concentrated under a vacuum to provide a cloudy brown oil. The oil was taken up in dichloromethane (150 mL), filtered under a vacuum and the solid phase washed with dichloromethane to provide the filtrate as a clear orange solution. The solution was concentrated under a vacuum to provide (*S*)-2-amino-2-cyclopropylethanol as a light brown liquid.


(3*S*,5*R*,6*R*)-3-Allyl-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyltetrahydro-2*H*-pyran-2-one (32 g, 81 mmol, Example 4, Step E) was combined with (*S*)-2-amino-2-cyclopropylethanol (26.7 g, 265 mmol) and the suspension was heated at 100 °C overnight. The reaction mixture was cooled to room temperature, diluted with ethyl acetate and washed with 1 N hydrochloric acid (2X), water, and brine. The organic layer was dried over magnesium sulfate and concentrated under vacuum to provide the title compound as a white solid. ^{1}H NMR (500 MHz, $CDCl_3$, δ ppm): 0.23–0.30 (m, 2H), 0.45–0.56 (m, 2H), 0.81 (m, 1H), 1.12 (s, 3H), 1.92–2.09 (m, 3H), 2.39 (dd, J = 13.6, 7.2 Hz, 1H), 2.86 (br s, 1H), 2.95 (dtd, J = 9.5, 6.3, 6.3, 2.9 Hz, 1H), 3.44 (dd, J = 11.0, 5.6 Hz, 1H), 3.49 (m, 1H), 3.61 (dd, J = 11.0, 2.9 Hz, 1H), 4.78 (d, J = 5.6 Hz, 1H), 4.95–5.13 (m, 2H), 5.63 (m, 1H), 5.99 (d, J = 6.4 Hz, 1H), 6.94–7.16 (m, 3H), 7.16–7.32 (m, 4H). MS (ESI) 494 $[M + H]^+$.

Step G. (3*S*,5*R*,6*S*)-3-Allyl-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-1-((*S*)-1-cyclopropyl-2-hydroxyethyl)-3-methylpiperidin-2-one




A solution of (2*S*)-2-((2*R*)-3-(4-chloro-3-fluorophenyl)-2-(3-chlorophenyl)-3-hydroxypropyl)-*N*-((*S*)-1-cyclopropyl-2-hydroxyethyl)-2-methylpent-4-enamide (40.2 g, 81 mmol, Example 4, Step F) in dichloromethane (80 mL) was added *p*-toluenesulfonic anhydride (66.3 g, 203 mmol) in dichloromethane (220 mL) at 0 °C ,and the reaction mixture was stirred for 10 minutes at same the temperature. 2,6-Lutidine (43.6 mL, 374 mmol, Aldrich, St. Louis, MO) was added dropwise via addition funnel at 0 °C. The reaction mixture was slowly warmed to room temperature, and then it was stirred at reflux. After 24 hours, sodium bicarbonate (68.3 g, 814 mmol) in water (600 mL) and 1,2-dichloroethane (300 mL) were added in succession. The reaction mixture was heated at reflux for an hour and then cooled to room temperature. The layers were separated and the aqueous layer was extracted with dichloromethane. The combined organic layer was washed with 1 N hydrochloric acid, water, and brine, then concentrated under reduced pressure. The residue was purified by flash chromatography (1.5 kg SiO_2 column, gradient elution of 10% to 50% ethyl acetate in hexanes) to provide the title compound as a white solid. ^{1}H NMR (500 MHz, $CDCl_3$, δ ppm): 0.06 (m, 1H), 0.26 (m, 1H), 0.57–0.67 (m, 2H), 0.85 (m, 1H), 1.25 (s, 3H), 1.85–2.20 (m, 2H), 2.57–2.65 (m, 2H), 3.09 (ddd, J = 11.8, 9.8, 4.8 Hz, 1H), 3.19 (t, J = 10.0 Hz, 1H), 3.36 (td, J = 10.3, 4.6 Hz, 1H), 3.63 (dd, J = 11.0, 4.6 Hz, 1H), 4.86 (d, J = 10.0 Hz, 1H), 5.16–5.19 (m, 2H), 5.87 (m, 1H), 6.77 (dd, J = 7.7, 1.6 Hz, 1H), 6.80–6.90 (m, 2H), 7.02 (t, J = 2.0 Hz, 1H), 7,16 (dd, J = 10.0, 7.7 Hz, 1H), 7.21 (dd, J = 10.0, 1.6 Hz, 1H), 7.29 (t, J = 10.0 Hz, 1H). MS (ESI) 476 $[M + H]^+$.

Step H. (3*S*,5*S*,6*R*,8*S*)-8-Allyl-5-(4-chloro-3-fluorophenyl)-6-(3-chlorophenyl)-3-cyclopropyl-8-methyl-2,3,5,6,7,8-hexahydrooxazolo[3,2-*a*]pyridin-4-ium 4-methylbenzenesulfonate



p-Toluenesulfonic acid monohydrate (30.3 g, 159 mmol, Aldrich, St. Louis, MO) was added to a solution of (3*S*,5*R*,6*S*)-3-allyl-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-1-((*S*)-1-cyclopropyl-2-hydroxyethyl)-3-methylpiperidin-2-one (73.6 g, 154 mmol) in toluene (386 mL). The reaction mixture was heated at reflux using a Dean-Stark apparatus. After 4 hours, the reaction was cooled and concentrated under reduced pressure to provide the title compound as a pale yellow syrup. The crude product was used in next step without further purification. ^{1}H NMR (500 MHz, $CDCl_3$, δ ppm): -0.25 to -0.10 (m, 2H), 0.08–0.18 (m, 1H), 0.33–0.50 (m, 2H), 1.57 (s, 3H), 1.92 (dd, *J* = 3.7 and 13.9 Hz, 1H), 2.37 (s, 3H), 2.63 (dd, *J* = 7.3 and 13.7 Hz, 1H), 2.72 (dd, *J* = 7.6 and 13.7 Hz, 1H), 2.93 (t, *J* = 13.7 Hz, 1H), 3.29 (m, 1H), 4.51 (t, *J* = 8.6 Hz, 1H), 4.57–4.63 (m, 1H), 5.33 (d, *J* = 17.1 Hz, 1H), 5.37 (d, *J* = 10.5 Hz, 1H), 5.47 (dd, *J* = 9.1 and 10.0 Hz, 1H), 5.75–5.93 (m, 2H), 6.80 (br s, 1H), 7.08 (s, 1H), 7.16–7.20 (m, 5H), 7.25–7.32 (m, 2H), 7.87 (d, *J* = 8.3 Hz, 2H). MS (ESI) 458 $[M + H]^+$.


Step I. (3*S*,5*R*,6*S*)-3-Allyl-1-((*S*)-2-(*tert*-butylthio)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methylpiperidin-2-one



2-Methyl-2-propanethiol (15.25 mL, 135 mmol, dried over activated 4 Å molecular sieves) was added to a solution of lithium bis(trimethylsilyl)amide in tetrahydrofuran (1.0 M, 135 mL, 135 mmol) at room temperature under argon in a 500 mL round-bottomed flask. The reaction mixture was heated to 60 °C. After 30 minutes, a solution of (3*S*,5*S*,6*R*,8*S*)-8-allyl-5-(4-chloro-3-fluorophenyl)-6-(3-chlorophenyl)-3-cyclopropyl-8-methyl-2,3,5,6,7,8-hexahydrooxazolo[3,2-*a*]pyridin-4-ium 4-methylbenzenesulfonate (78 g, 123 mmol, Example 4, Step H) in anhydrous tetrahydrofuran (100 mL) was added via cannula. The reaction mixture was heated at 60 °C for 3 hours and then cooled to room temperature. The reaction mixture was quenched with water and extracted thrice with ethyl acetate. The organics were pooled, washed with brine, dried over magnesium sulfate, filtered and concentrated under a vacuum to provide a yellow foam. Purification by flash column chromatography (1.5 kg SiO_2 column, gradient elution with 5% to 30% ethyl acetate in hexanes provided the title compound as an off-white foam. ^{1}H NMR (400 MHz, $CDCl_3$, δ ppm): -0.89 to -0.80 (m, 1H), -0.15 to -0.09 (m, 1H), 0.27–0.34 (m, 1H), 0.41–0.48 (m, 1H), 1.28 (s, 3H), 1.35 (s, 9H), 1.70–1.77 (m, 1H), 1.86 (dd, *J* = 3.1 and 13.5 Hz, 1H),

2.16 (t, J = 13.7, 1H), 2.17–2.23 (m, 1H), 2.60–2.63 (m, 3H), 3.09 (dt, J = 3.1 and 10.4 Hz, 1H), 3.62 (t, J = 11.1 Hz, 1H), 4.70 (d, J = 10.1 Hz, 1H), 5.16 (s, 1H), 5.19–5.21 (m, 1H), 5.82–5.93 (m, 1H), 6.65–6.80 (m, 1H), 6.80–6.83 (m, 1H), 6.84–6.98 (m, 1H), 7.05–7.07 (m, 1H), 7.12–7.18 (m, 2H), 7.19–7.26 (m, 1H). MS (ESI) 548.2 $[M + H]^+$.

Step J. 2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-Butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetic acid




Ruthenium(III) chloride hydrate (0.562 mg, 2.493 mmol) was added to a mixture of (3*S*,5*R*,6*S*)-3-allyl-1-((*S*)-2-(*tert*-butylthio)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methylpiperidin-2-one (62.17 g, 113 mmol, Example 4, Step I) and sodium periodate (24.67 g) in ethyl acetate (216 mL), acetonitrile (216 mL) and water (324 mL) at 20 °C. The temperature quickly rose to 29 °C. The reaction mixture was cooled to 20 °C and the remaining equivalents of sodium periodate were added in five 24.67 g portions over 2 hours, being careful to maintain an internal reaction temperature below 25 °C. The reaction was incomplete, so additional sodium periodate (13 g) was added. The temperature increased from 22 °C to 25 °C. After stirring for an additional 1.5 hours, the reaction mixture was filtered under a vacuum and washed with ethyl acetate. The layers were separated and the aqueous layer was extracted with ethyl acetate. The organics were pooled, washed with brine, dried over magnesium sulfate, filtered and concentrated under a vacuum to provide a dark green foam. Purification by flash column chromatography (1.5 kg SiO_2 column, gradient elution of 0% to 20% isopropanol in hexanes) provided an off-white foam. 15% Ethyl acetate in heptanes (970 mL) was added to the foam, and the mixture was heated at 80 °C until the foam dissolved. The solution was then cooled slowly, and at 60 °C the solution was seeded with previously obtained crystalline material. The mixture was cooled to room temperature and then allowed to stand at room temperature for 2 hours before collecting the solid by vacuum filtration to provide a white solid with a very pale pink hue (57.1 g). The mother liquor was concentrated under a vacuum to provide a pink foam (8.7 g). 15% ethyl acetate in heptanes (130 mL) was added to the foam, and it was heated at 80 °C to completely dissolve the material. The solution was cooled, and at 50 °C, it was seeded with crystalline material. After cooling to room temperature the solid was collected


by vacuum filtration to provide a white crystalline solid with a very pale pink hue. ^{1}H NMR (500 MHz, $CDCl_3$, δ ppm): -1.10 to -1.00 (m, 1H), -0.30 to -0.22 (m, 1H), 0.27–0.37 (m, 1H), 0.38–0.43 (m, 1H), 1.45 (s, 9H), 1.50 (s, 3H), 1.87 (dd, *J* = 2.7 and 13.7 Hz, 1H), 1.89–1.95 (m, 1H), 2.46 (t, *J* = 13.7, 1H), 2.69–2.73 (m, 1H), 2.78 (d, *J* = 14.9 Hz, 1H), 2.93 (dd, *J* = 2.0 and 13.7 Hz, 1H), 3.07 (d, *J* = 14.9 Hz, 1H), 3.11 (dt, *J* = 2.7 and 11.0 Hz, 1H), 4.30 (t, *J* = 13.5 Hz, 1H), 4.98 (d, *J* = 10.8 Hz, 1H), 6.75–6.87 (m, 1H), 6.88–6.90 (m, 1H), 6.98 (br s, 1H), 7.02–7.09 (m, 1H), 7.11–7.16 (m, 2H), 7.16–7.25 (m, 1H). MS (ESI) 598.1 $[M + H]^+$.

EXAMPLE 5

4-(2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-Butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid




Step A. Methyl 4-(2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoate



N-(3-Dimethylaminopropyl)-*N*′-ethylcarbodiimide hydrochloride (EDC, 76 g, 398 mmol) was added to a mixture of 2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetic acid (79.4 g, 133 mmol, Example 4, Step J) and methyl 4-amino-2-methoxybenzoate (26.4 g, 146 mmol) in pyridine (332 mL) at 3 °C. The mixture was allowed to warm to room temperature and was stirred at room temperature for 16 hours. The reaction mixture was cooled to 0 °C and added to an ice-cold solution of 1 M hydrochloric acid (1 L). Ether (1 L) was added and the layers were agitated and then separated. The organic layer was washed with 1 M hydrochloric acid (6 x 500 mL), saturated aqueous sodium bicarbonate (500 mL), brine (500 mL), dried over magnesium

sulfate, filtered and concentrated under a vacuum to provide an off-white foam. ^{1}H NMR (400 MHz, $CDCl_3$, δ ppm): -1.20 to -1.12 (m, 1H), -0.35 to -0.20 (m, 1H), 0.05–0.20 (m, 1H), 0.32–0.45 (m, 1H), 1.45 (s, 9H), 1.48 (s, 3H), 1.86–1.98 (m, 1H), 2.03 (dd, J = 2.7 and 13.7 Hz, 1H), 2.43 (t, J = 13.7, 1H), 2.64–2.75 (m, 1H), 2.80 (d, J = 14.3 Hz, 1H), 2.89–2.96 (m, 2H), 3.24 (dt, J = 2.5 and 10.8 Hz, 1H), 3.89 (s, 3H), 3.96 (s, 3H), 4.28–4.36 (m, 1H), 4.98 (d, J = 10.8 Hz, 1H), 6.85–6.93 (m, 3H), 6.99 (br s, 1H), 7.06–7.18 (m, 4 H), 7.82 (br s, 1H), 7.85 (d, J = 8.4 Hz, 1H), 8.81 (br s, 1H). MS (ESI) 761.2 $[M + H]^+$.

Step B. 4-(2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-Butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoic acid



A solution of lithium hydroxide monohydrate (18.2 g, 433 mmol) in water (295 mL) was added to a solution of methyl 4-(2-((3*R*,5*R*,6*S*)-1-((*S*)-2-(*tert*-butylsulfonyl)-1-cyclopropylethyl)-6-(4-chloro-3-fluorophenyl)-5-(3-chlorophenyl)-3-methyl-2-oxopiperidin-3-yl)acetamido)-2-methoxybenzoate (164.9 g, 217 mmol, Example 5, Step A) in tetrahydrofuran (591 mL) and methanol (197 mL) at room temperature. After stirring for 15 hours at room temperature, a trace amount of the ester remained, so the reaction mixture was heated at 50 °C for 1 hour. When the reaction was complete, the mixture was concentrated under a vacuum to remove the tetrahydrofuran and methanol. The thick mixture was diluted with water (1 L) and 1 M hydrochloric acid (1 L) was added. The resulting white solid was collected by vacuum filtration in a Büchner funnel. The vacuum was removed, and water (1 L) was added to the filter cake. The material was stirred with a spatula to suspend it evenly in the water. The liquid was then removed by vacuum filtration. This washing cycle was repeated three more times to provide a white solid. The solid was dried under vacuum at 45 °C for 3 days to provide the title compound as a white solid. ^{1}H NMR (500 MHz, DMSO-d_6) δ ppm -1.30 to -1.12 (m, 1H), -0.30 to -0.13 (m, 1H), 0.14–0.25 (m, 1H), 0.25–0.38 (m, 1H), 1.30 (s, 3H), 1.34 (s, 9H), 1.75–1.86 (m, 1H), 2.08–2.18 (m, 2H), 2.50–2.60 (m, 1H), 2.66 (d, J = 13.7, 1H), 3.02–3.16 (m, 2H), 3.40–3.50 (m, 1H), 3.77 (s, 3H), 4.05–4.20 (m, 1H), 4.89 (d, J = 10.5 Hz, 1H), 6.90–6.93 (m, 3H), 7.19 (d, J = 8.8

Hz, 1H), 7.22–7.26 (m, 3H), 7.40–7.50 (m, 1H), 7.54 (br s, 1H), 7.68 (d, J = 8.6 Hz, 1H) 10.44 (s, 1H), 12.29 (br s, 1H). MS (ESI) 747.2 $[M + H]^+$.

Another particular MDM2 inhibitor is Compound B (also known as AMG 2653149 or 2653149), which is Example 256 of published PCT application WO2011/153,509. Other MDM2 inhibitors that can be used in the combinations of the present invention include those disclosed in published PCT application WO2013/049250; U.S. provisional patent application number 61/766,635; and U.S. provisional patent application number 61/784,230. Still other MDM2 inhibitors that can be used in the combinations of the present invention include RG7112 (also known as RO504337), RG7388 (also known as idasanutin, and RO5503781), SAR405838 (also known as MI-773), SAR299155, MK-8242 (also known as SCH 900242), CGM097 and DS 3032. The structures of RG7112 and SAR299155 as well as other inhibitors of MDM2 that can be used in the present invention are shown in *Bioorganic & Medicinal Chemistry Letters* 23 (2013) 2480-2485, which summarizes pathways to the clinic for MDM2 inhibitors. Still other MDM2 inhibitors that can be used in the combinations of the present invention include RG7775 and Novartis CGM097.

The MDM2 inhibitors of the present invention can be used in combination with Aurora kinase inhibitors, such as those found in published PCT application WO2011/031842. A particular compound is AMG 900 (Example 1).

The MDM2 inhibitors of the present invention can be used in combination with MAP kinase pathway inhibitors. Examples of proteins in the MAP kinase pathway that can be inhibited and the inhibitors of such proteins used in combination with an MDM2 inhibitors are BRAF inhibitors, Pan-RAF inhibitors, and MEK inhibitors. There are three main RAF isoforms: ARAF, BRAF and CRAF. A pan-RAF inhibitor shows inhibitory activity on more than one RAF isoform. In contrast, a BRAF inhibitor exhibits more inhibitor activity (or selectivity) towards BRAF than the other RAF proteins.

The MDM2 inhibitors of the present invention can be used in combination with MEK inhibitors, such as those found in published PCT application WO2002/006213. A particular compound is N-(((2R)-2,3-dihydroxypropyl)oxy)-3,4-difluoro-2-((2-fluoro-4-iodophenyl)amino)benzamide, also known as AMG 1009089 or 1009089, (Example 39).

The MDM2 inhibitors of the present invention can be used in combination with BRAF inhibitors, such as those found in published PCT application WO2008/153,947. A particular compound is AMG 2112819 (also known as 2112819) (Example 56). Another particular BRAF inhibitor that can be used in the combinations of the present invention is dabrafenib. Another BRAF inhibitor that can be used in the combinations of the present invention is vemurafenib.

Pan-RAF inhibitors can also be used along with MDM2 inhibitors in the combinations of the present invention. Particular Pan–Raf inhibitor include RAF265 and MLN-2480.





The MDM2 inhibitors of the present invention can be used in combination with MEK inhibitors. Particular MEK inhibitors that can be used in the combinations of the present invention include PD0325901, trametinib, pimasertib, MEK162 [also known as binimetinib], TAK-733, GDC-0973 and AZD8330. A particular MEK inhibitor that can be used along with MDM2 inhibitor in the combinations of the present invention is trametinib (also called AMG 2712849 or 2712849). Another particular MEK inhibitor is N-(((2R)-2,3-dihydroxypropyl)oxy)-3,4-difluoro-2-((2-fluoro-4-iodophenyl)amino)benzamide, also known as AMG 1009089, 1009089 or PD0325901. Another particular MEK inhibitor that can be used in the combinations of the present invention includes cobimetinib.

In another aspect, the present invention relates to the use of the compound of the present invention in combination with one or more pharmaceutical agent that is an inhibitor of a protein in the phosphatidylinositol 3-kinase (PI3K) pathway. Examples of proteins in the PI3K pathway include PI3K, mTOR and PKB (also known as Akt or AKT). The PI3K protein exists in several isoforms including α, β, δ, or γ. It is contemplated that a PI3K inhibitor that can be used in the present invention can be selective for one or more isoform. By selective it is meant that the compounds inhibit one or more isoform more than other isoforms. Selectivity is a concept well known to those is the art and can be measured with well-known *in vitro* or cell-based activity assays. Preferred selectivity includes greater than 2-fold, preferably 10-fold, or more preferably 100-fold greater selectivity for one or more isoform over the other isoforms. In one aspect, the PI3K inhibitors that can be used in combination with compounds of the present invention are PI3K α selective inhibitors. In another aspect the compound is a PI3K δ selective inhibitor. In still another aspect the compound is a PI3K β selective inhibitor.

Examples of PI3K inhibitors that can be used in combination with one or more compounds of the present invention include those disclosed in the following: PCT published application no. WO2010/151791; PCT published application no. WO2010/151737; PCT published application no.WO2010/151735; PCT published application no. WO2010151740; PCT published application no. WO2008/118455; PCT published application no. WO2008/118454; PCT published application no. WO2008/118468; U.S. published application no. US20100331293; U.S. published application no. US20100331306; U.S. published application no. US20090023761; U.S. published application no. US20090030002; U.S. published application no. US20090137581;U.S. published application no. US2009/0054405; U.S. published application


no. U.S. 2009/0163489; U.S. published application no. US 2010/0273764; U.S. published application no. U.S. 2011/0092504; or PCT published application no. WO2010/108074.

Preferred PI3K inhibitors for use in combination with the compound of the present invention include:


;

;

; or

;

or a pharmaceutically acceptable salt thereof.

Also preferred is a compound of Formula IIa below, or a pharmaceutically acceptable salt thereof,



IIa

wherein X^1 is fluorine or hydrogen;Y^1 is hydrogen or methyl; and Z^1 is hydrogen or methyl. A particular PI3K inhibitor that can be used in the combinations of the present invention is AMG 511 (also known as AMG 2539965 or 2539965), which is Example 148 of published PCT application WO2010/126895.

Other PI3K inhibitors that can be used in combination with MDM2 inhibitors in the combinations of the present invention include Pan-PI3K inhibitors such as BKM120 and GDC-0941; PI3Kα selective inhibitors such as AMG 511 and BYL719; and PI3K β selective inhibitors such as GSK-2636771.

Compounds that inhibit both PI3K and mTOR (dual inhibitors) are known. In still another aspect, the present invention provides the use of dual PI3K and mTOR inhibitors for use in combination with MDM2 inhibitors. An example of a particular dual inhibitor is GDC-0980.

mTOR is a protein in the PI3K pathway. It is another aspect of the present invention to use an mTOR inhibitor in combination with MDM2 inhibitors. mTOR inhibitors that can be used

in combination with the compound of the present invention include those disclosed in the following documents: PCT published application no. WO2010/132598 and PCT published application no. WO2010/096314. mTOR inhibitors that can be used in combination with MDM2 inhibitors in the combinations of the present invention include AZD2014 and MLN0128.

PKB (AKT) is also a protein in the PI3K pathway. It is another aspect of the present invention to use an AKT inhibitor in combination with an MDM2 inhibitor. AKT inhibitors that can be used in combination with the compound of the present invention include those disclosed in the following documents: U.S. patent no. 7,354,944; U.S. patent no. 7,700,636; U.S. patent no. 7,919,514; U.S. patent no. 7,514,566; U.S. patent application publication no. US 2009/0270445 A1; U.S. patent no. 7,919,504; U.S. patent no. 7,897,619; or PCT published application no. WO 2010/083246 A1. Particular AKT inhibitors that can be used in combination with MDM2 inhibitors in the combinations of the present invention include MK-2206, GDC-0068 and AZD5363.

MDM2 inhibitors can also be used in combination with CDK4 and/or 6 inhibitors in the present invention CDK 4 and/or 6 inhibitors that can be used in the present combinations include those disclosed in the following documents: PCT published application no. WO 2009/085185 or U.S. patent application publication no. US2011/0097305.

Other compounds that can be used in combination with MDM2 inhibitors in the combinations of the present invention include compounds that inhibit proteins that are part of the intrinsic apoptosis pathway. Examples of such compounds include Bcl2/BclxL inhibitors such as navitoclax and Bcl2 inhibitors as such as ABT-199.

Other compounds that can be used in combination with MDM2 inhibitors in the combinations of the present invention include BCR-ABL inhibitors such as dasatinib and HDAC inhibitors such as panobinostat.

Other compounds that can be used in combination with MDM2 inhibitors in the combinations of the present invention include platinums, such as Cisplatin, Carboplatin and Oxaliplatin; Topoisomerase II inhibitors, typically of the anthracycline class, such as doxorubicin, daunorubicin, idarubicin, epirubicin, pegylated liposomal doxorubicin hydrochloride, myocet and etoposide; Topoisomerase I inhibitors such as irinotecan (CPT-11); DNA alkylation agents such as temozolomide; and nucleoside analogs such as cytarabine and decitabine.

Other compounds that can be used in combination with MDM2 inhibitors in the combinations of the present invention include receptor and non-receptor kinase inhibitors including tyrosine kinase inhibitors. Example of such compounds include imatinib, dasatinib, ponatinib, bosutinib, nilotininb, quizartinib, midostaurin, erlotinib and lapatinib.

The compound of the present invention can also be used in combination with pharmaceutically active agents that treat nausea. Examples of agents that can be used to treat nausea include: dronabinol; granisetron; metoclopramide; ondansetron; and prochlorperazine; or a pharmaceutically acceptable salt thereof.

The compound of the present invention may also be used in combination with radiation therapy, hormone therapy, surgery and immunotherapy, which therapies are well known to those skilled in the art.

Since one aspect of the present invention contemplates the treatment of the disease/conditions with a combination of pharmaceutically active compounds that may be administered separately, the invention further relates to combining separate pharmaceutical compositions in kit form. The kit comprises two separate pharmaceutical compositions: the compound of the present invention, and a second pharmaceutical compound. The kit comprises a container for containing the separate compositions such as a divided bottle or a divided foil packet. Additional examples of containers include syringes, boxes and bags. Typically, the kit comprises directions for the use of the separate components. The kit form is particularly advantageous when the separate components are preferably administered in different dosage forms (e.g., oral and parenteral), are administered at different dosage intervals, or when titration of the individual components of the combination is desired by the prescribing physician or veterinarian.

An example of such a kit is a so-called blister pack. Blister packs are well known in the packaging industry and are being widely used for the packaging of pharmaceutical unit dosage forms (tablets, capsules, and the like). Blister packs generally consist of a sheet of relatively stiff material covered with a foil of a preferably transparent plastic material. During the packaging process recesses are formed in the plastic foil. The recesses have the size and shape of the tablets or capsules to be packed. Next, the tablets or capsules are placed in the recesses and the sheet of relatively stiff material is sealed against the plastic foil at the face of the foil which is opposite from the direction in which the recesses were formed. As a result, the tablets or capsules are sealed in the recesses between the plastic foil and the sheet. Preferably the strength of the sheet is such that the tablets or capsules can be removed from the blister pack by manually applying pressure on the recesses whereby an opening is formed in the sheet at the place of the recess. The tablet or capsule can then be removed via said opening.

It may be desirable to provide a memory aid on the kit, e.g., in the form of numbers next to the tablets or capsules whereby the numbers correspond with the days of the regimen which the tablets or capsules so specified should be ingested. Another example of such a memory aid is a

calendar printed on the card, e.g., as follows "First Week, Monday, Tuesday, . . . etc . . . Second Week, Monday, Tuesday, . . . " etc. Other variations of memory aids will be readily apparent. A "daily dose" can be a single tablet or capsule or several pills or capsules to be taken on a given day. Also, a daily dose of a compound of the present invention can consist of one tablet or capsule, while a daily dose of the second compound can consist of several tablets or capsules and vice versa. The memory aid should reflect this and aid in correct administration of the active agents.

In another specific embodiment of the invention, a dispenser designed to dispense the daily doses one at a time in the order of their intended use is provided. Preferably, the dispenser is equipped with a memory-aid, so as to further facilitate compliance with the regimen. An example of such a memory-aid is a mechanical counter which indicates the number of daily doses that has been dispensed. Another example of such a memory-aid is a battery-powered micro-chip memory coupled with a liquid crystal readout, or audible reminder signal which, for example, reads out the date that the last daily dose has been taken and/or reminds one when the next dose is to be taken.

The compounds of the present invention and other pharmaceutically active compounds, if desired, can be administered to a patient either orally, rectally, parenterally, (for example, intravenously, intramuscularly, or subcutaneously) intracisternally, intravaginally, intraperitoneally, intravesically, locally (for example, powders, ointments or drops), or as a buccal or nasal spray. All methods that are used by those skilled in the art to administer a pharmaceutically active agent are contemplated.

Compositions suitable for parenteral injection may comprise physiologically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions, or emulsions, and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents, or vehicles include water, ethanol, polyols (propylene glycol, polyethylene glycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preserving, wetting, emulsifying, and dispersing agents. Microorganism contamination can be prevented by adding various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example, sugars, sodium chloride, and the like. Prolonged absorption of injectable pharmaceutical compositions can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

Solid dosage forms for oral administration include capsules, tablets, powders, and granules. In such solid dosage forms, the active compound is admixed with at least one inert customary excipient (or carrier) such as sodium citrate or dicalcium phosphate or (a) fillers or extenders, as for example, starches, lactose, sucrose, mannitol, and silicic acid; (b) binders, as for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia; (c) humectants, as for example, glycerol; (d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain complex silicates, and sodium carbonate; (a) solution retarders, as for example, paraffin; (f) absorption accelerators, as for example, quaternary ammonium compounds; (g) wetting agents, as for example, cetyl alcohol and glycerol monostearate; (h) adsorbents, as for example, kaolin and bentonite; and (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, or mixtures thereof. In the case of capsules, and tablets, the dosage forms may also comprise buffering agents.

Solid compositions of a similar type may also be used as fillers in soft and hard filled gelatin capsules using such excipients as lactose or milk sugar, as well as high molecular weight polyethylene glycols, and the like.

Solid dosage forms such as tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells, such as enteric coatings and others well known in the art. They may also contain opacifying agents, and can also be of such composition that they release the active compound or compounds in a certain part of the intestinal tract in a delayed manner. Examples of embedding compositions that can be used are polymeric substances and waxes. The active compound can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. In addition to the active compounds, the liquid dosage form may contain inert diluents commonly used in the art, such as water or other solvents, solubilizing agents and emulsifiers, as for example, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils, in particular, cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil, and sesame seed oil, glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, or mixtures of these substances, and the like.

Besides such inert diluents, the composition can also include adjuvants, such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents. Suspensions, in addition to the active compound, may contain suspending agents, as for example,

ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, or mixtures of these substances, and the like.

Compositions for rectal administration are preferable suppositories, which can be prepared by mixing the compounds of the present invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax, which are solid at ordinary room temperature, but liquid at body temperature, and therefore, melt in the rectum or vaginal cavity and release the active component.

Dosage forms for topical administration of the compound of the present invention include ointments, powders, sprays and inhalants. The active compound or compounds are admixed under sterile condition with a physiologically acceptable carrier, and any preservatives, buffers, or propellants that may be required. Ophthalmic formulations, eye ointments, powders, and solutions are also contemplated as being within the scope of this invention.

The compounds of the present invention can be administered to a patient at dosage levels in the range of about 0.1 to about 3,000 mg per day. For a normal adult human having a body weight of about 70 kg, a dosage in the range of about 0.01 to about 100 mg per kilogram body weight is typically sufficient. The specific dosage and dosage range that can be used depends on a number of factors, including the requirements of the patient, the severity of the condition or disease being treated, and the pharmacological activity of the compound being administered. A particular dosage of a compound of the present invention is the FDA approved dosage, if the compound has been approved.

The compounds of the present invention can be administered as pharmaceutically acceptable salts, esters, amides or prodrugs. The term "salts" refers to inorganic and organic salts of compounds of the present invention. The salts can be prepared in situ during the final isolation and purification of a compound, or by separately reacting a purified compound in its free base or acid form with a suitable organic or inorganic base or acid and isolating the salt thus formed. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, nitrate, acetate, oxalate, palmitiate, stearate, laurate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and laurylsulphonate salts, and the like. The salts may include cations based on the alkali and alkaline earth metals, such as sodium, lithium, potassium, calcium, magnesium, and the like, as well as non-toxic ammonium, quaternary ammonium, and amine cations including, but not limited to, ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine,

trimethylamine, triethylamine, ethylamine, and the like. See, for example, S. M. Berge, et al., "Pharmaceutical Salts," J Pharm Sci, 66: 1-19 (1977).

Examples of pharmaceutically acceptable esters of the compound of the present invention include C_1-C_8 alkyl esters. Acceptable esters also include C_5-C_7 cycloalkyl esters, as well as arylalkyl esters such as benzyl. C_1-C_4 alkyl esters are commonly used. Esters of compounds of the present invention may be prepared according to methods that are well known in the art.

Examples of pharmaceutically acceptable amides of the compound of the present invention include amides derived from ammonia, primary C_1-C_8 alkyl amines, and secondary C_1-C_8 dialkyl amines. In the case of secondary amines, the amine may also be in the form of a 5 or 6 membered heterocycloalkyl group containing at least one nitrogen atom. Amides derived from ammonia, C_1-C_3 primary alkyl amines and C_1-C_2 dialkyl secondary amines are commonly used. Amides of the compound of the present invention may be prepared according to methods well known to those skilled in the art.

The term "prodrug" means compounds that are transformed *in vivo* to yield a compound of the present invention. The transformation may occur by various mechanisms, such as through hydrolysis in blood. A discussion of the use of prodrugs is provided by T. Higuchi and W. Stella, "Prodrugs as Novel Delivery Systems," Vol. 14 of the A.C.S. Symposium Series, and in Bioreversible Carriers in Drug Design, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

To illustrate, because the compound of the invention contains a carboxylic acid functional group, a prodrug can comprise an ester formed by the replacement of the hydrogen atom of the acid group with a group such as (C_1-C_8 alkyl, (C_2-$C12_2$)alkanoyloxymethyl, 1-(alkanoyloxy)ethyl having from 4 to 9 carbon atoms, 1-methyl-1-(alkanoyloxy)ethyl having from 5 to 10 carbon atoms, alkoxycarbonyloxymethyl having from 3 to 6 carbon atoms, 1-(alkoxycarbonyloxy)ethyl having from 4 to 7 carbon atoms, 1-methyl-1-(alkoxycarbonyloxy)ethyl having from 5 to 8 carbon atoms, N-(alkoxycarbonyl)aminomethyl having from 3 to 9 carbon atoms, 1-(N-(alkoxycarbonyl)aminomethyl having from 4 to 10 carbon atoms, 3-phthalidyl, 4-crotonolactonyl, gamma-butyrolacton-4-yl, di-N,N-(C_1-C_2)alkylamino(C_2-C_3)alkyl (such as β-dimethylaminoethyl), carbamoyl-(C_1-C_2)alkyl, N,N-di(C_1-C_2)alkylcarbamoyl-(C_1-C_2)alkyl and piperidino-, pyrrolidino- or morpholino(C_2-$_3$)alkyl.

The compounds of the present invention may contain asymmetric or chiral centers, and therefore, exist in different stereoisomeric forms. It is contemplated that all stereoisomeric forms of the compound as well as mixtures thereof, including racemic mixtures, form part of the present invention. In addition, the present invention contemplates all geometric and positional isomers.

For example, if the compound contains a double bond, both the cis and trans forms (designated as Z and E, respectively), as well as mixtures, are contemplated.

Mixture of stereoisomers, such as diastereomeric mixtures, can be separated into their individual stereochemical components on the basis of their physical chemical differences by known methods such as chromatography and/or fractional crystallization. Enantiomers can also be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., an alcohol), separating the diastereomers and converting (e.g., hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. Also, some compounds may be atropisomers (e.g., substituted biaryls).

The compounds of the present invention may exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water (hydrate), ethanol, and the like. The present invention contemplates and encompasses both the solvated and unsolvated forms.

It is also possible that the compounds of the present invention may exist in different tautomeric forms. All tautomers of the compound of the present invention are contemplated. For example, all of the tautomeric forms of the tetrazole moiety are included in this invention. Also, for example, all keto-enol or imine-enamine forms of the compounds are included in this invention.

Those skilled in the art will recognize that the compound names and structures contained herein may be based on a particular tautomer of a compound. While the name or structure for only a particular tautomer may be used, it is intended that all tautomers are encompassed by the present invention, unless stated otherwise.

It is also intended that the present invention encompass compounds that are synthesized *in vitro* using laboratory techniques, such as those well known to synthetic chemists; or synthesized using *in vivo* techniques, such as through metabolism, fermentation, digestion, and the like. It is also contemplated that the compounds of the present invention may be synthesized using a combination of *in vitro* and *in vivo* techniques.

The present invention also includes isotopically-labelled compounds, which are identical to those recited herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as ^{2}H, ^{3}H, ^{13}C, ^{14}C, ^{15}N, ^{16}O, ^{17}O, ^{18}O, ^{31}P, ^{32}P, ^{35}S, ^{18}F, and ^{36}Cl. In one aspect, the present invention relates to compounds wherein one or more hydrogen atom is replaced with deuterium (^{2}H) atoms.

The compounds of the present invention that contains the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically-labelled compounds of the present invention, for example those into which radioactive isotopes such as ^{3}H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e., ^{3}H, and carbon-14, i.e., ^{14}C, isotopes are particularly preferred for their ease of preparation and detection. Further, substitution with heavier isotopes such as deuterium, i.e., ^{2}H, can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labelled compounds of this invention can generally be prepared by substituting a readily available isotopically labelled reagent for a non-isotopically labelled reagent.

The compounds of the present invention may exist in various solid states including crystalline states and as an amorphous state. The different crystalline states, also called polymorphs, and the amorphous states of the present compounds are contemplated as part of this invention.

In synthesizing the compounds of the present invention, it may be desirable to use certain leaving groups. The term "leaving groups" ("LG") generally refer to groups that are displaceable by a nucleophile. Such leaving groups are known in the art. Examples of leaving groups include, but are not limited to, halides (e.g., I, Br, F, Cl), sulfonates (e.g., mesylate, tosylate), sulfides (e.g., SCH_3), N-hydroxysuccinimide, N-hydroxybenzotriazole, and the like. Examples of nucleophiles include, but are not limited to, amines, thiols, alcohols, Grignard reagents, anionic species (e.g., alkoxides, amides, carbanions) and the like.

All patents, patent applications and other documents recited herein are hereby incorporated by reference in their entirety.

The examples presented below illustrate specific embodiments of the present invention. These examples are meant to be representative and are not intended to limit the scope of the claims in any manner

The following abbreviations may be used herein:

932 or 2705932	AMG 232
ADD	additivity
AML	acute myelogenous leukemia
ATP	adenosine triphosphate
Cispl	cisplatin

CML	chronic myelogenous leukemia
CPT-11	irinotecan
DIC	drug in capsules
DLBCL	diffuse large B-cell lymphoma
Dox	doxorubicin
GBM	glioblastoma
HPβCD	hydroxypropyl beta cyclodextrin
HPMC	hydroxypropyl methylcellulose
MDS	myelodysplastic syndrome
mpk	milligrams per kilogram
NHL	non-Hodgkin's lymphoma
NMR	nuclear magnetic resonance
NSCLC	non-small cell lung cancer
PBS	phosphate buffered saline
PCT	patent cooperation treaty
RTK	receptor tyrosine kinase
TGI	tumor growth inhibition
Tx begins	treatment begins

Cell Culture Reagents

Tween® 80	polyoxyethylene (20) sorbitan monooleate (Uniqema Americas, Inc., Wilmington, DE)
Pluronic® F68	polyoxyethylene-polyoxypropylene block co-polymer (BASF Corp., Mount Olive, NJ)

EXAMPLES

In Vitro Cell-Based Combination Studies

Cell lines were purchased from American Type Culture Collection (ATCC), German Collection of Microorganisms and Cell Cultures (DSMZ), and Japanese Collection of Research Bioresources (JCRB). Each line was cultured in its recommended growth medium. Cell line A375sq2 was made in accordance with the procedure set forth in J. Med. Chem. **2009**, 52, 6189-6192, footnote 13.

For Examples 1 to 71, cells were seeded into 384-well cell culture plates at initial densities ranging from 300 to 7500 cells per well in a 30 μL volume, depending on the growth rate of the cell line, so that adherent cells would remain at subconfluent densities by the end of the 72-hour treatment period. In order to determine the appropriate concentration range to test in subsequent combination experiments, 10 μL of a 19-point, two-fold serial titration of compound starting at a high final concentration of 20 μM, as well as a 0.25% dimethyl sulfoxide (DMSO) control were added to the cells 16 hours after seeding. CellTiter-Glo® Luminescent Cell Viability Assay (Promega; Madison, WI) was used to determine the number of viable cells based on quantitation of the amount of ATP present, an indicator of metabolically active cells. Luminescence was measured with an EnVision® Multilabel Reader (Perkin Elmer; Waltham, MA) for each cell line at time zero (V_0) before the addition of compounds, as well as after 72 hours of compound treatment (T_{72}). Growth inhibition (GI) was calculated according to the following equations, where V_{72} was luminescence of DMSO control at 72 hours and T_{72} was luminescence of the compound-treated sample: if $T_{72} \geq V_0$, then $GI = 100 \times (1 - ((T_{72}-V_0) / (V_{72} - V_0)))$; if $T_{72} < V_0$, then $GI = 100 \times (1 - ((T_{72}-V_0) / V_0))$. This formula is derived from the growth inhibition calculation used in the National Cancer Institute's NCI-60 high throughput screen. On a scale of 0 to 200 percent growth inhibition, a value of 0 represents uninhibited growth (i.e. DMSO control), 100 typically represents stasis (signal equivalent at time zero reading), and 200 represents complete cell killing. Sigmoidal dose response curves were plotted using a 4-parameter logistic model. For all combinations tested in any given cell line, the starting high concentration and dilution factor of each compound was chosen to well-define the curve maximum, curve minimum, and slope over a range of 9 doses.

Two-way combination experiments were conducted essentially as described above, with the following exceptions. To each well, 5 μL of a 9-point serial titration of the first compound (starting high final concentration and dilution factor determined as previously described) along with DMSO control was added to the cells in 10 identical rows (x-axis) of a 384-well plate. Then, 5 μL of a 9-point serial titration of the second compound (starting high final concentration and dilution factor determined as previously described) along with DMSO control were added to the cells in 10 identical columns (y-axis). The final concentration of DMSO in each well was 0.25%. Duplicate 100-well (10x10) matrices were run on each 384-well plate. Growth inhibition for each well of the matrix was calculated as previously described, and the data were analyzed for synergistic interactions using Chalice™ Analyzer software (Zalicus; Cambridge, MA) which generated synergy scores based on the Loewe Additivity model (Lehàr, J., et al. (2009). "Synergistic drug combinations tend to improve therapeutically relevant selectivity." Nat Biotech

27(7): 659-666) and Rickles, et al (2012) "Adenosine A2A and Beta-2 Adrenergic Receptor Agonists: Novel Selective and Synergistic Multiple Myeloma Targets Discovered through Systematic Combination Screening" Mol Cancer Therapeutics 11 (7): 1432.

The Loewe ADD (additivity) model quantifies combination effects. Combinations were ranked initially by Additivity Excess Volume, which is defined as ADD Volume = $\sum C_X,C_Y (I_{data} - I_{Loewe})$. where $I_{Loewe}(C_X,C_Y)$ is the inhibition that satisfies $(C_X/EC_X) + (C_Y/EC_Y) = 1$, and $EC_{X,Y}$ are the effective concentrations at I_{Loewe} for the single agent curves. A "Synergy Score" was also used, where the Synergy Score S = log f_X log $f_Y \sum I_{data} (I_{data}–I_{Loewe})$, summed over all non-single-agent concentration pairs, and where log $f_{X,Y}$ is the natural logarithm of the dilution factors used for each single agent. This effectively calculates a volume between the measured and Loewe additive response surfaces, weighted towards high inhibition and corrected for varying dilution factors. An uncertainty σ_S was calculated for each synergy score, based on the measured errors for the I_{data} values and standard error propagation.

In the examples shown, the Growth Inhibition (%) matrices contain the consensus growth inhibition values calculated from the luminescence data using the formulas described above; the ADD Model Growth Inhibition (%) matrices contain the predicted growth inhibition values based on the Loewe additivity model, which was derived from the modeled single agent growth inhibition curves; and the ADD Excess Growth Inhibition (%) matrices contain the values of growth inhibition in excess of the additivity model. The additivity model serves as a "null-hypothesis" and assumes no synergistic interaction between the two agents. Any activity observed after subtraction of the ADD model from the Growth Inhibition dose response matrix (= ADD Excess Growth Inhibition) is indicative of synergy.

For Examples 72-89, two-way combination experiments were carried out in a similar manner as described above, but using a high-throughput screening format. Cells were thawed from a liquid nitrogen preserved state. Screening began after the cells were expanded and were dividing at their expected doubling times. Cells were seeded in growth media in either black 1536-well or 384-well tissue culture treated plates at the cell densities as listed in the table below.

Cell Line	Plate Format	Cell Density
RT4	384	500
SJSA-1	1536	100
KS-1	1536	100
MCF7	384	500
RKO	1536	100
SNG-M	1536	100
RPMI-2650	1536	200

G-401	1536	100
CML-T1	1536	100
EOL-1	384	500
MOLM-13	384	500
SK-HEP-1	1536	100
A427	1536	100
DOHH-2	1536	100
22RV1	1536	100
A375	1536	100
C32	384	500
MKN45	1536	100
NOI-SNJ-1	1536	100
SW982	1536	100
HT-29	1536	100
PC-3	1536	100

Cells were equilibrated in assay plates via centrifugation and placed in incubators attached to the Dosing Modules at 37°C for 24 hours before treatment. At the time of treatment, a set of untreated assay plates were collected and ATP levels were measured by adding ATPLite 1step Luminescent Assay reagent (Perkin Elmer; Waltham, MA). These Tzero (T_0) plates were read using ultrasensitive luminescence on an EnVision® Multilabel Reader (Perkin Elmer; Waltham, MA). Treated assay plates were incubated with compound for 72 hours and were assayed for viable cell number at the endpoint. All data points were collected via automated processes; quality controlled; and analyzed using Chalice™ Analyzer software (Zalicus; Cambridge, MA) which generated synergy scores based on the Loewe Additivity model (Lehàr et al., supra). Assay plates were accepted if they passed the following quality control standards: relative luciferase values were consistent throughout the entire experiment, Z-factor scores were greater than 0.6, and untreated/vehicle controls behaved consistently on the plate. The synergistic interaction of each experimental combination was evaluated for statistical significance. The synergy scores calculated for individual replicates of heterologous combinations (AxB) were compared to the synergy scores of individual replicates of the component self-crosses (AxA and BxB) using two sample Student's t-test with unequal variance. Only those combinations in which the synergy score of AxB was statistically significant (p-value < 0.05) when compared to both AxA and BxB were considered synergistic.

Three-way combination experiments were conducted essentially as described above, with the following exceptions. Ten identical 384-well plates each containing duplicate 100-well two-way combinations were set-up as before, except that 3.3 μL of 3x final concentration of each compound were added. Then 3.3 μL of a single fixed concentration of the third compound were added to all of the wells of the matrix on a given plate. Thus the ten plates comprised the 9-point

serial titration of the third compound (z-axis; starting high final concentration and dilution factor determined as previously described), along with a DMSO control (i.e. no compound 3 added) plate. Growth inhibition for each well of the matrix was calculated as previously described. Examples of cell lines used in the above-identified experiments are set forth in the table below. The tissue column of the table indicate the type of tissue from which the cells were obtained, and the mutation column indicates certain mutations identified in the particular cell line.

Cell Line	**Tissue**	**Mutation**
HT-1197	Bladder	NRAS, PI3K
RT4	Bladder	TSC1
KNS-81-FD	Brain; CNS	EGFR
CAL-51	Breast	PI3K
MCF7	Breast	PI3K
MDA-MB-175-VII	Breast	HER2 autocrine loop
UACC-812	Breast	HER2 amplified
HCT116	Colon	KRAS, PI3K
LS 174T	Colon	KRAS, PI3K
RKO	Colon	BRAF, PI3K
SW48	Colon	EGFR
BV-173	Haematopoietic and lymphoid (CML)	BCR-ABL
CML-T1	Haematopoietic and lymphoid (CML)	BCR-ABL
GDM-1	Haematopoietic and lymphoid (AML)	None identified
ML-2	Haematopoietic and lymphoid (AML)	KRAS
MOLM-13	Haematopoietic and lymphoid (AML)	FLT3 ITD
OCI-AML3	Haematopoietic and lymphoid (AML)	Not known
CAL-54	Kidney	MET amplified
SK-HEP-1	Liver	BRAF
A427	Lung (NSCLC)	KRAS
A549	Lung	KRAS
NCI-H1666	Lung (NSCLC)	BRAF (not V600E)
NCI-H460	Lung (NSCLC)	KRAS, PI3K
A2780	Ovary	PTEN
A375sq2	Skin (Melanoma)	BRAF
A375	Skin (Melanoma)	BRAF

C32	Skin (Melanoma)	BRAF, PTEN
G-361	Skin (Melanoma)	BRAF
SH-4	Skin (Melanoma)	BRAF
A204	Soft tissue (rhabdomyosarcoma)	None identified
G-401	Soft tissue (kidney)	None identified
MKN45	Stomach	MET amplified

The results of the *in vitro* cell-based combination studies are shown in Figures 1 to 89.

In Figures 1 to71, the Growth Inhibition (%) matrices represent the consensus growth inhibition values from multiple experimental replicates for a given combination calculated according to formulas described above. For example, in Figure 1, the growth inhibition obtained when 2.5 μM Compound A and 3 μM Compound 1 were combined was 114%. The ADD Model Growth Inhibition (%) matrices represent the predicted growth inhibition values for an additive interaction between two compounds based on the Loewe additivity model and was derived from the experimental growth inhibition activities of each of the agents alone. The ADD Excess Growth Inhibition (%) matrices represent the growth inhibition values in excess of the additivity model. For example, in Figure 1a, when 2.5 μM Compound A was combined with 3 μM Compound 1, the growth inhibition in excess of the additivity model was 26% (114% experimental growth inhibition – 88% model = 26%). The shading of the matrices corresponds to the degree of growth inhibition, with darker shading/positive growth inhibition values representing larger effects; negative growth inhibition values were excluded from synergy score calculations. Synergy scores were calculated based on the sum of the excess growth inhibition values for a given combination, with normalization factors for the concentration ranges of the component agents tested and additional weighting given to synergistic interactions that occur at high effect levels (Lehàr et al., supra). For example, in Figure 1a, the synergy score for the combination of Compound A x Compound 1 was 0.532; the synergy score for the self-cross combination of Compound A x Compound A was 0.621; and the synergy score for the self-cross combination of Compound 1 x Compound 1 was 0.432.

In Figures 72 to 89, the first label on the Y axis indicates the target of the compound tested in combination with the particular MDM2 inhibitor. For example, in Figure 72, the first target is BRAF (the compound tested is vemurafenib) and the last is MEK (the compound tested is AZD8330) . The next section on the Y axis indicates the exact combination. For example, in Figure 72, AMG 232 x Vemurafenib means that the combination tested was AMG 232 and Vemurafenib. On the X axis, at the top of the grid, the cancer cell line that was tested is indicated. Above the cancer cell line, mutational status is indicated For example, in Figure 72,

KRAS and BRAF indicates that the particular cell lines below these designations contain a KRAS or BRAF mutation as indicated (mutation data obtained from the literature or the Sanger (Cosmic) or Broad Institute (Cancer Cell Line Encyclopedia) cancer genomics databases). It should be noted that the cell lines used in these experiments are known to those skilled in the art, and the various mutations associated with those cells lines may be readily determined by one skilled in the art. Also, on Figure 72, TP53 indicates that the designated cell lines contain a mutation in TP53. As mentioned above, MDM2 inhibitors show activity in cancer having wild type TP53. The shading in each box of the grid indicates the level of synergy identified, with darker indicating higher synergy. The number in the box is the synergy score, and if a number is underlined it indicates that the experiment showed statistical significance.

In Vivo Tumor Xenograft Combination Studies

In vivo tumor xenograft studies were conducted following these general procedures:

Tumor cells (Table 1) were cultured, harvested and implanted subcutaneously into the right flank of female athymic nude mice. When tumors reached about 200mm^3, mice were randomized into treatment groups (n=10/group) and treatment was initiated (on days indicated on graphs). Compound names, dosing frequency, and routes of administration are listed in Table 2. Tumor sizes and body weights were measured 2 to 3 times per week. Tumor volume was measured by digital calipers, calculated as L x W x H and expressed in mm^3. Statistical significance of observed differences between growth curves was evaluated by repeated measures analysis of covariance (RMANOVA) of the log transformed tumor volume data with Dunnett adjusted multiple comparisons comparing the control group to the treatment groups. For combination studies, RMANOVA was run with the combination group compared one to one with each single agent treatment group.

BD Matrigel™ Basement Membrane Matrix is a solubilized basement membrane preparation extracted from the Engelbreth-Holm-Swarm (EHS) mouse sarcoma (BD Biosciences, San Jose, CA)

All studies were measured in a blinded manner.

Table 1:

Cell Line	**Tumor Type**	**Cells/mouse**	**Source #**	**Matrigel (cells:matrigel)**
RKO	Colon	$5x10^6$	(ATCC) CRL-2577	1:1
SJSA-1	Osteosarcoma	$5x10^6$	(ATCC) CRL-2098	2:1
HCT116	Colorectal	$2x10^6$	(ATCC) CCL-247	2:1

A375sq2	Melanoma	$5x10^6$	See Reference above	2:1
NCI-H460	Non-small cell lung	$5x10^6$	(ATCC) HTB-177	no
U87	Glioblastoma	$5x10^6$	(ATCC) HTB-14	no
Molm13	Acute myelogenous leukemia	$2.5x10^6$	(DSMZ) AC-554	1:1

Table 2:

Treatment	**Route**	**Frequency**
AMG 232	PO	QD
1009089 (MEK)	PO	QD
cisplatin	IP	1x/wk
CPT-11	IP	1x/wk
doxorubicin	IV	1x/wk
2112819 (BRAF)	PO	QD
RG7112 (MDM2)	PO	QD
2520765 (PI3K)	PO	QD
cytarabine	IP	5 days on, 2 days off
decitabine	IP	3x/wk

Definition of abbreviations: PO: oral gavage IP: intraperitoneal IV: intravenous QD: once per day Wk: week

In vivo combination studies conducted:

1. AMG 232 + MEK (RKO),
2. AMG 232 + BRAF (RKO)
3. AMG 232 + cisplatin (H460)
4. AMG 232 + cisplatin (HCT-116)
5. AMG 232 + doxorubicin (SJSA-1)
6. AMG 232 + irinotecan (HCT116)
7. AMG 232 + MEK (A375sq2)
8. AMG 232 + BRAF (A375sq2)
9. AMG 232 + BRAF+ PI3K (RKO, triple combination)
10. AMG 232 + doxorubicin (Molm-13)
11. AMG 232 + MEK (Molm-13)

12. AMG 232 + cytarabine (Molm-13)
13. AMG 232 + decitabine (Molm-13)
14. AMG 232 + sorafenib (Molm-13)

The results of the *in vivo* tumor xenograft combination studies are shown in Figures A to O.

Table A below illustrates specific combinations of an MDM2 inhibitor with one or more additional pharmaceutically active agents for particular cancers types. The data obtained and summarized in the Figures indicates that the combinations set forth in Table A show enhanced anti-cancer activity over what is expected when the individual members of the combination therapy are used alone. It is noted that the magnitude of the therapeutic synergy that is seen can vary depending on the type of cancer treated and agent used.

Table A

MDM2 Inhibitor	**Additional Pharmaceutically Active Agent**	**Cancer Type**
AMG 232	vemurafenib	melanoma
AMG 232	vemurafenib	colon
AMG 232	vemurafenib	liver
AMG 232	vemurafenib	sarcoma
AMG 232	vemurafenib	AML
AMG 232	vemurafenib	CML
AMG 232	vemurafenib	DLBCL
AMG 232	vemurafenib	kidney
AMG 232	dabrafenib	melanoma
AMG 232	dabrafenib	colon
AMG 232	dabrafenib	liver
AMG 232	dabrafenib	sarcoma
AMG 232	dabrafenib	glioblastoma
AMG 232	dabrafenib	head and neck
AMG 232	dabrafenib	AML
AMG 232	dabrafenib	CML
AMG 232	dabrafenib	DLBCL
AMG 232	RAF265	melanoma

AMG 232	RAF265	colon
AMG 232	RAF265	liver
AMG 232	RAF265	sarcoma
AMG 232	RAF265	NSCLC
AMG 232	RAF265	stomach
AMG 232	RAF265	endometrium
AMG 232	RAF265	glioblastoma
AMG 232	RAF265	head and neck
AMG 232	RAF265	bladder
AMG 232	RAF265	AML
AMG 232	RAF265	CML
AMG 232	RAF265	DLBCL
AMG 232	MLN2480	melanoma
AMG 232	MLN2480	colon
AMG 232	MLN2480	liver
AMG 232	MLN2480	NSCLC
AMG 232	MLN2480	endometrium
AMG 232	MLN2480	bladder
AMG 232	MLN2480	AML
AMG 232	MLN2480	CML
AMG 232	MLN2480	DLBCL
AMG 232	trametinib	melanoma
AMG 232	trametinib	colon
AMG 232	trametinib	liver
AMG 232	trametinib	sarcoma
AMG 232	trametinib	NSCLC
AMG 232	trametinib	stomach
AMG 232	trametinib	prostate
AMG 232	trametinib	kidney
AMG 232	trametinib	glioblastoma
AMG 232	trametinib	breast
AMG 232	trametinib	head and neck
AMG 232	trametinib	bladder

AMG 232	trametinib	AML
AMG 232	trametinib	CML
AMG 232	trametinib	DLBCL
AMG 232	pimasertib	melanoma
AMG 232	pimasertib	colon
AMG 232	pimasertib	liver
AMG 232	pimasertib	NSCLC
AMG 232	pimasertib	stomach
AMG 232	pimasertib	prostate
AMG 232	pimasertib	kidney
AMG 232	pimasertib	glioblastoma
AMG 232	pimasertib	breast
AMG 232	pimasertib	head and neck
AMG 232	pimasertib	AML
AMG 232	pimasertib	CML
AMG 232	pimasertib	DLBCL
AMG 232	pimasertib	bladder
AMG 232	MEK162	melanoma
AMG 232	MEK162	colon
AMG 232	MEK162	liver
AMG 232	MEK162	NSCLC
AMG 232	MEK162	stomach
AMG 232	MEK162	prostate
AMG 232	MEK162	glioblastoma
AMG 232	MEK162	bladder
AMG 232	MEK162	AML
AMG 232	MEK162	CML
AMG 232	MEK162	DLBCL
AMG 232	TAK-733	melanoma
AMG 232	TAK-733	colon
AMG 232	TAK-733	liver
AMG 232	TAK-733	sarcoma
AMG 232	TAK-733	NSCLC

AMG 232	TAK-733	stomach
AMG 232	TAK-733	prostate
AMG 232	TAK-733	kidney
AMG 232	TAK-733	glioblastoma
AMG 232	TAK-733	breast
AMG 232	TAK-733	head and neck
AMG 232	TAK-733	bladder
AMG 232	TAK-733	AML
AMG 232	TAK-733	CML
AMG 232	TAK-733	DLBCL
AMG 232	GDC-0973	melanoma
AMG 232	GDC-0973	colon
AMG 232	GDC-0973	liver
AMG 232	GDC-0973	NSCLC
AMG 232	GDC-0973	stomach
AMG 232	GDC-0973	prostate
AMG 232	GDC-0973	kidney
AMG 232	GDC-0973	glioblastoma
AMG 232	GDC-0973	breast
AMG 232	GDC-0973	bladder
AMG 232	GDC-0973	head and neck
AMG 232	GDC-0973	sarcoma
AMG 232	GDC-0973	AML
AMG 232	GDC-0973	CML
AMG 232	GDC-0973	DLBCL
AMG 232	AZD8330	melanoma
AMG 232	AZD8330	colon
AMG 232	AZD8330	liver
AMG 232	AZD8330	sarcoma
AMG 232	AZD8330	NSCLC
AMG 232	AZD8330	stomach
AMG 232	AZD8330	prostate
AMG 232	AZD8330	kidney

AMG 232	AZD8330	glioblastoma
AMG 232	AZD8330	breast
AMG 232	AZD8330	head and neck
AMG 232	AZD8330	bladder
AMG 232	AZD8330	AML
AMG 232	AZD8330	CML
AMG 232	AZD8330	DLBCL
AMG 232	BKM120	prostate
AMG 232	BKM120	breast
AMG 232	BKM120	melanoma
AMG 232	BKM120	NSCLC
AMG 232	BKM120	kidney
AMG 232	BKM120	stomach
AMG 232	BKM120	head and neck
AMG 232	BKM120	bladder
AMG 232	BKM120	sarcoma
AMG 232	BKM120	AML
AMG 232	BKM120	CML
AMG 232	GDC-0941	prostate
AMG 232	GDC-0941	breast
AMG 232	GDC-0941	endometrium
AMG 232	GDC-0941	melanoma
AMG 232	GDC-0941	NSCLC
AMG 232	GDC-0941	kidney
AMG 232	GDC-0941	glioblastoma
AMG 232	GDC-0941	stomach
AMG 232	GDC-0941	head and neck
AMG 232	GDC-0941	bladder
AMG 232	GDC-0941	sarcoma
AMG 232	GDC-0941	AML
AMG 232	GDC-0941	CML
AMG 232	GDC-0941	DLBCL
AMG 232	BYL719	prostate

AMG 232	BYL719	breast
AMG 232	BYL719	endometrium
AMG 232	BYL719	melanoma
AMG 232	BYL719	NSCLC
AMG 232	BYL719	kidney
AMG 232	BYL719	glioblastoma
AMG 232	BYL719	stomach
AMG 232	BYL719	head and neck
AMG 232	BYL719	bladder
AMG 232	BYL719	sarcoma
AMG 232	BYL719	AML
AMG 232	BYL719	DLBCL
AMG 232	GSK-2636771	AML
AMG 232	GSK-2636771	DLBCL
AMG 232	MK-2206	prostate
AMG 232	MK-2206	breast
AMG 232	MK-2206	melanoma
AMG 232	MK-2206	endometrium
AMG 232	MK-2206	head and neck
AMG 232	MK-2206	sarcoma
AMG 232	MK-2206	AML
AMG 232	MK-2206	DLBCL
AMG 232	GDC-0068	prostate
AMG 232	GDC-0068	breast
AMG 232	GDC-0068	endometrium
AMG 232	GDC-0068	melanoma
AMG 232	GDC-0068	glioblastoma
AMG 232	GDC-0068	head and neck
AMG 232	GDC-0068	sarcoma
AMG 232	GDC-0068	AML
AMG 232	GDC-0068	DLBCL
AMG 232	AZD5363	prostate
AMG 232	AZD5363	breast

AMG 232	AZD5363	endometrium
AMG 232	AZD5363	melanoma
AMG 232	AZD5363	glioblastoma
AMG 232	AZD5363	head and neck
AMG 232	AZD5363	sarcoma
AMG 232	AZD5363	AML
AMG 232	AZD5363	CML
AMG 232	AZD5363	DLBCL
AMG 232	GDC-0980	prostate
AMG 232	GDC-0980	breast
AMG 232	GDC-0980	melanoma
AMG 232	GDC-0980	kidney
AMG 232	GDC-0980	glioblastoma
AMG 232	GDC-0980	stomach
AMG 232	GDC-0980	head and neck
AMG 232	GDC-0980	bladder
AMG 232	GDC-0980	liver
AMG 232	GDC-0980	sarcoma
AMG 232	GDC-0980	AML
AMG 232	GDC-0980	DLBCL
AMG 232	AZD2014	prostate
AMG 232	AZD2014	breast
AMG 232	AZD2014	endometrium
AMG 232	AZD2014	melanoma
AMG 232	AZD2014	NSCLC
AMG 232	AZD2014	kidney
AMG 232	AZD2014	glioblastoma
AMG 232	AZD2014	stomach
AMG 232	AZD2014	head and neck
AMG 232	AZD2014	bladder
AMG 232	AZD2014	liver
AMG 232	AZD2014	sarcoma
AMG 232	AZD2014	AML

AMG 232	AZD2014	DLBCL
AMG 232	MLN0128	prostate
AMG 232	MLN0128	breast
AMG 232	MLN0128	endometrium
AMG 232	MLN0128	melanoma
AMG 232	MLN0128	NSCLC
AMG 232	MLN0128	glioblastoma
AMG 232	MLN0128	stomach
AMG 232	MLN0128	head and neck
AMG 232	MLN0128	bladder
AMG 232	MLN0128	liver
AMG 232	MLN0128	sarcoma
AMG 232	MLN0128	AML
AMG 232	MLN0128	DLBCL
AMG 232	dasatinib	bladder
AMG 232	dasatinib	colon
AMG 232	dasatinib	endometium
AMG 232	dasatinib	glioblastoma
AMG 232	dasatinib	head and neck
AMG 232	dasatinib	kidney
AMG 232	dasatinib	NSCLC
AMG 232	dasatinib	melanoma
AMG 232	dasatinib	sarcoma
AMG 232	dasatinib	AML
AMG 232	dasatinib	CML
AMG 232	dasatinib	DLBCL
AMG 232	panobinostat	kidney
AMG 232	panobinostat	head and neck
AMG 232	panobinostat	melanoma
AMG 232	panobinostat	sarcoma
AMG 232	panobinostat	stomach
AMG 232	panobinostat	AML
AMG 232	panobinostat	DLBCL

AMG 232	panobinostat	liver
AMG 232	doxorubicin	breast
AMG 232	doxorubicin	AML
AMG 232	etoposide	sarcoma
AMG 232	cytarabine	AML
AMG 232	decitabine	AML
AMG 232	navitoclax	bladder
AMG 232	navitoclax	breast
AMG 232	navitoclax	endometrium
AMG 232	navitoclax	glioblastoma
AMG 232	navitoclax	head and neck
AMG 232	navitoclax	kidney
AMG 232	navitoclax	liver
AMG 232	navitoclax	melanoma
AMG 232	navitoclax	sarcoma
AMG 232	navitoclax	stomach
AMG 232	navitoclax	AML
AMG 232	navitoclax	CML
AMG 232	navitoclax	DLBCL
AMG 232	ABT-199	glioblastoma
AMG 232	ABT-199	head and neck
AMG 232	ABT-199	kidney
AMG 232	ABT-199	sarcoma
AMG 232	ABT-199	AML
AMG 232	ABT-199	CML
AMG 232	ABT-199	DLBCL
AMG 232	imatinib	CML
AMG 232	ponatinib	CML
AMG 232	bosutinb	CML
AMG 232	nilotinib	CML
AMG 232	quizartinib	AML
AMG 232	midostaurin	AML
AMG 232	cisplatin	Ovarian

AMG 232	cisplatin	Colon
AMG 232	cisplatin	NSCLC
AMG 232	cisplatin	esophageal/ stomach
AMG 232	cispaltin	Breast
AMG 232	doxorubicin	Breast
AMG 232	doxorubicin	stomach
AMG 232	doxorubicin	ovarian
AMG 232	doxorubicin	AML
AMG 232	doxorubicin	ALL
AMG 232	doxorubicin	MDS
AMG 232	doxorubicin	NHL
AMG 232	doxorubicin	Hodgkin's lymphoma
AMG 232	decitabine	MDS
AMG 232	sorafenib	kidney
AMG 232	sorafenib	liver
AMG 232	sorafenib	AML
AM-7209	vemurafenib	melanoma
AM-7209	vemurafenib	colon
AM-7209	vemurafenib	liver
AM-7209	vemurafenib	sarcoma
AM-7209	vemurafenib	AML
AM-7209	vemurafenib	CML
AM-7209	vemurafenib	DLBCL
AM-7209	dabrafenib	melanoma
AM-7209	dabrafenib	colon
AM-7209	dabrafenib	liver
AM-7209	dabrafenib	sarcoma
AM-7209	dabrafenib	glioblastoma
AM-7209	dabrafenib	stomach
AM-7209	dabrafenib	head and neck
AM-7209	dabrafenib	AML
AM-7209	dabrafenib	CML
AM-7209	dabrafenib	DLBCL

AM-7209	dabrafenib	prostate
AM-7209	dabrafenib	endometrium
AM-7209	RAF265	melanoma
AM-7209	RAF265	colon
AM-7209	RAF265	liver
AM-7209	RAF265	sarcoma
AM-7209	RAF265	NSCLC
AM-7209	RAF265	stomach
AM-7209	RAF265	endometrium
AM-7209	RAF265	kidney
AM-7209	RAF265	glioblastoma
AM-7209	RAF265	head and neck
AM-7209	RAF265	bladder
AM-7209	RAF265	AML
AM-7209	RAF265	CML
AM-7209	RAF265	DLBCL
AM-7209	MLN2480	melanoma
AM-7209	MLN2480	colon
AM-7209	MLN2480	liver
AM-7209	MLN2480	sarcoma
AM-7209	MLN2480	NSCLC
AM-7209	MLN2480	breast
AM-7209	MLN2480	AML
AM-7209	MLN2480	CML
AM-7209	MLN2480	DLBCL
AM-7209	trametinib	melanoma
AM-7209	trametinib	colon
AM-7209	trametinib	liver
AM-7209	trametinib	sarcoma
AM-7209	trametinib	NSCLC
AM-7209	trametinib	stomach
AM-7209	trametinib	endometrium
AM-7209	trametinib	prostate

AM-7209	trametinib	kidney
AM-7209	trametinib	glioblastoma
AM-7209	trametinib	breast
AM-7209	trametinib	head and neck
AM-7209	trametinib	bladder
AM-7209	trametinib	AML
AM-7209	trametinib	CML
AM-7209	trametinib	DLBCL
AM-7209	pimasertib	melanoma
AM-7209	pimasertib	colon
AM-7209	pimasertib	liver
AM-7209	pimasertib	NSCLC
AM-7209	pimasertib	stomach
AM-7209	pimasertib	endometrium
AM-7209	pimasertib	prostate
AM-7209	pimasertib	kidney
AM-7209	pimasertib	glioblastoma
AM-7209	pimasertib	breast
AM-7209	pimasertib	head and neck
AM-7209	pimasertib	bladder
AM-7209	pimasertib	AML
AM-7209	pimasertib	CML
AM-7209	pimasertib	sarcoma
AM-7209	MEK162	melanoma
AM-7209	MEK162	colon
AM-7209	MEK162	liver
AM-7209	MEK162	NSCLC
AM-7209	MEK162	stomach
AM-7209	MEK162	prostate
AM-7209	MEK162	kidney
AM-7209	MEK162	glioblastoma
AM-7209	MEK162	head and neck
AM-7209	MEK162	bladder

AM-7209	MEK162	AML
AM-7209	MEK162	CML
AM-7209	TAK-733	melanoma
AM-7209	TAK-733	colon
AM-7209	TAK-733	liver
AM-7209	TAK-733	sarcoma
AM-7209	TAK-733	NSCLC
AM-7209	TAK-733	stomach
AM-7209	TAK-733	endometrium
AM-7209	TAK-733	prostate
AM-7209	TAK-733	kidney
AM-7209	TAK-733	glioblastoma
AM-7209	TAK-733	breast
AM-7209	TAK-733	head and neck
AM-7209	TAK-733	bladder
AM-7209	TAK-733	AML
AM-7209	TAK-733	CML
AM-7209	TAK-733	DLBCL
AM-7209	GDC-0973	melanoma
AM-7209	GDC-0973	colon
AM-7209	GDC-0973	liver
AM-7209	GDC-0973	sarcoma
AM-7209	GDC-0973	NSCLC
AM-7209	GDC-0973	stomach
AM-7209	GDC-0973	endometrium
AM-7209	GDC-0973	prostate
AM-7209	GDC-0973	kidney
AM-7209	GDC-0973	glioblastoma
AM-7209	GDC-0973	breast
AM-7209	GDC-0973	head and neck
AM-7209	GDC-0973	bladder
AM-7209	GDC-0973	AML
AM-7209	GDC-0973	CML

AM-7209	GDC-0973	DLBCL
AM-7209	AZD8330	melanoma
AM-7209	AZD8330	colon
AM-7209	AZD8330	liver
AM-7209	AZD8330	sarcoma
AM-7209	AZD8330	NSCLC
AM-7209	AZD8330	stomach
AM-7209	AZD8330	prostate
AM-7209	AZD8330	kidney
AM-7209	AZD8330	glioblastoma
AM-7209	AZD8330	breast
AM-7209	AZD8330	head and neck
AM-7209	AZD8330	bladder
AM-7209	AZD8330	AML
AM-7209	AZD8330	CML
AM-7209	AZD8330	DLBCL
AM-7209	BKM120	prostate
AM-7209	BKM120	breast
AM-7209	BKM120	melanoma
AM-7209	BKM120	NSCLC
AM-7209	BKM120	kidney
AM-7209	BKM120	glioblastoma
AM-7209	BKM120	stomach
AM-7209	BKM120	head and neck
AM-7209	BKM120	bladder
AM-7209	BKM120	sarcoma
AM-7209	BKM120	AML
AM-7209	BKM120	CML
AM-7209	GDC-0941	prostate
AM-7209	GDC-0941	breast
AM-7209	GDC-0941	colon
AM-7209	GDC-0941	endometrium
AM-7209	GDC-0941	melanoma

AM-7209	GDC-0941	NSCLC
AM-7209	GDC-0941	kidney
AM-7209	GDC-0941	glioblastoma
AM-7209	GDC-0941	stomach
AM-7209	GDC-0941	head and neck
AM-7209	GDC-0941	bladder
AM-7209	GDC-0941	sarcoma
AM-7209	GDC-0941	AML
AM-7209	GDC-0941	DLBCL
AM-7209	GDC-0941	liver
AM-7209	BYL719	prostate
AM-7209	BYL719	breast
AM-7209	BYL719	colon
AM-7209	BYL719	melanoma
AM-7209	BYL719	NSCLC
AM-7209	BYL719	kidney
AM-7209	BYL719	glioblastoma
AM-7209	BYL719	stomach
AM-7209	BYL719	head and neck
AM-7209	BYL719	bladder
AM-7209	BYL719	liver
AM-7209	BYL719	sarcoma
AM-7209	BYL719	AML
AM-7209	BYL719	DLBCL
AM-7209	GSK-2636771	breast
AM-7209	GSK-2636771	NSCLC
AM-7209	GSK-2636771	liver
AM-7209	GSK-2636771	sarcoma
AM-7209	GSK-2636771	AML
AM-7209	GSK-2636771	DLBCL
AM-7209	MK-2206	prostate
AM-7209	MK-2206	breast
AM-7209	MK-2206	colon

AM-7209	MK-2206	endometrium
AM-7209	MK-2206	melanoma
AM-7209	MK-2206	NSCLC
AM-7209	MK-2206	glioblastoma
AM-7209	MK-2206	stomach
AM-7209	MK-2206	head and neck
AM-7209	MK-2206	sarcoma
AM-7209	MK-2206	liver
AM-7209	MK-2206	AML
AM-7209	MK-2206	CML
AM-7209	MK-2206	DLBCL
AM-7209	GDC-0068	prostate
AM-7209	GDC-0068	breast
AM-7209	GDC-0068	colon
AM-7209	GDC-0068	endometrium
AM-7209	GDC-0068	melanoma
AM-7209	GDC-0068	NSCLC
AM-7209	GDC-0068	glioblastoma
AM-7209	GDC-0068	stomach
AM-7209	GDC-0068	head and neck
AM-7209	GDC-0068	liver
AM-7209	GDC-0068	sarcoma
AM-7209	GDC-0068	AML
AM-7209	GDC-0068	DLBCL
AM-7209	AZD5363	prostate
AM-7209	AZD5363	breast
AM-7209	AZD5363	colon
AM-7209	AZD5363	endometrium
AM-7209	AZD5363	NSCLC
AM-7209	AZD5363	glioblastoma
AM-7209	AZD5363	stomach
AM-7209	AZD5363	head and neck
AM-7209	AZD5363	liver

AM-7209	AZD5363	sarcoma
AM-7209	AZD5363	AML
AM-7209	AZD5363	CML
AM-7209	AZD5363	DLBCL
AM-7209	GDC0980	prostate
AM-7209	GDC0980	breast
AM-7209	GDC0980	colon
AM-7209	GDC0980	endometrium
AM-7209	GDC0980	melanoma
AM-7209	GDC0980	NSCLC
AM-7209	GDC0980	kidney
AM-7209	GDC0980	glioblastoma
AM-7209	GDC0980	stomach
AM-7209	GDC0980	head and neck
AM-7209	GDC0980	bladder
AM-7209	GDC0980	sarcoma
AM-7209	GDC0980	AML
AM-7209	GDC0980	DLBCL
AM-7209	AZD2014	prostate
AM-7209	AZD2014	breast
AM-7209	AZD2014	colon
AM-7209	AZD2014	melanoma
AM-7209	AZD2014	NSCLC
AM-7209	AZD2014	kidney
AM-7209	AZD2014	stomach
AM-7209	AZD2014	head and neck
AM-7209	AZD2014	bladder
AM-7209	AZD2014	liver
AM-7209	AZD2014	sarcoma
AM-7209	AZD2014	AML
AM-7209	AZD2014	DLBCL
AM-7209	MLN0128	prostate
AM-7209	MLN0128	breast

AM-7209	MLN0128	colon
AM-7209	MLN0128	endometrium
AM-7209	MLN0128	melanoma
AM-7209	MLN0128	NSCLC
AM-7209	MLN0128	glioblastoma
AM-7209	MLN0128	stomach
AM-7209	MLN0128	head and neck
AM-7209	MLN0128	bladder
AM-7209	MLN0128	liver
AM-7209	MLN0128	sarcoma
AM-7209	MLN0128	AML
AM-7209	MLN0128	DLBCL
AM-7209	navitoclax	bladder
AM-7209	navitoclax	breast
AM-7209	navitoclax	colon
AM-7209	navitoclax	endometrium
AM-7209	navitoclax	glioblastoma
AM-7209	navitoclax	head and neck
AM-7209	navitoclax	kidney
AM-7209	navitoclax	liver
AM-7209	navitoclax	NSCLC
AM-7209	navitoclax	melanoma
AM-7209	navitoclax	sarcoma
AM-7209	navitoclax	stomach
AM-7209	navitoclax	AML
AM-7209	navitoclax	CML
AM-7209	navitoclax	DLBCL
AM-7209	ABT-199	glioblastoma
AM-7209	ABT-199	head and neck
AM-7209	ABT-199	liver
AM-7209	ABT-199	sarcoma
AM-7209	ABT-199	AML
AM-7209	ABT-199	CML

AM-7209	ABT-199	DLBCL
AM-7209	doxorubicin	AML
AM-7209	etoposide	sarcoma
AM-7209	etoposide	stomach
AM-7209	Irinotecan	colon
AM-7209	cytarabine	AML
AM-7209	decitabine	AML
AM-7209	dasatinib	bladder
AM-7209	dasatinib	colon
AM-7209	dasatinib	endometrium
AM-7209	dasatinib	glioblastoma
AM-7209	dasatinib	head and neck
AM-7209	dasatinib	kidney
AM-7209	dasatinib	liver
AM-7209	dasatinib	NSCLC
AM-7209	dasatinib	melanoma
AM-7209	dasatinib	prostate
AM-7209	dasatinib	sarcoma
AM-7209	dasatinib	AML
AM-7209	dasatinib	CML
AM-7209	dasatinib	DLBCL
AM-7209	panobinostat	endometrium
AM-7209	panobinostat	head and neck
AM-7209	panobinostat	kidney
AM-7209	panobinostat	liver
AM-7209	panobinostat	melanoma
AM-7209	panobinostat	sarcoma
AM-7209	panobinostat	stomach
AM-7209	panobinostat	AML
AM-7209	panobinostat	CML
AM-7209	panobinostat	DLBCL
AM-7209	ponatinib	CML
AM-7209	imatinib	CML

AM-7209	bosutinib	CML
AM-7209	nilotinib	CML
AM-7209	quizartinib	AML
AM-7209	midostaurin	AML
AM-7209	cisplatin	Ovarian
AM-7209	cisplatin	Colon
AM-7209	cisplatin	NSCLC
AM-7209	cisplatin	esophageal/ stomach
AM-7209	cispaltin	Breast
AM-7209	doxorubicin	Breast
AM-7209	doxorubicin	stomach
AM-7209	doxorubicin	ovarian
AM-7209	doxorubicin	AML
AM-7209	doxorubicin	ALL
AM-7209	doxorubicin	MDS
AM-7209	doxorubicin	NHL
AM-7209	doxorubicin	Hodgkin's lymphoma
AM-7209	decitabine	MDS
AM-7209	sorafenib	kidney
AM-7209	sorafenib	liver
AM-7209	sorafenib	AML

CLAIMS

1. Use of a therapeutically effective amount of 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid and trametinib in the manufacture of a medicament for treating melanoma in a patient in need thereof.

2. The use of claim 1, further comprising a therapeutically effective amount of dabrafenib.

3. The use of claim 1, wherein the melanoma has a BRAF V600E or V600K mutation.

4. The use of claim 2, wherein the melanoma has a BRAF V600E or V600K mutation.

5. Use of a therapeutically effective amount of 2-((3R,5R,6S)-5-(3-chlorophenyl)-6-(4-chlorophenyl)-1-((S)-1-(isopropylsulfonyl)-3-methylbutan-2-yl)-3-methyl-2-oxopiperidin-3-yl)acetic acid and trametinib in the manufacture of a medicament for treating AML in a patient in need thereof.

6. The use of claim 5, wherein the AML has a FLT3-ITD mutation.

Figure 1-Example 1; A204 cells





Figure 1a-Example 1; A204 cells



	Synergy Score
A x 1	0.532
A self-cross	0.621
1 self-cross	0.432

Figure 2-Example 2; A375sq2





Figure 2a-Example 2; A375sq2 cells



-	-	-	-	-	-	-	-	-	-
3	3	12	1	6	14	16	23	26	27
-2	-1	0	-4	5	7	11	22	19	18
-4	-3	3	-3	-5	2	15	15	18	13
0	1	2	0	4	-1	7	12	10	11
7	5	2	1	4	0	6	9	7	10
-5	10	3	1	6	-2	2	5	7	-3
-2	2	6	6	-7	0	-2	1	-1	2
0	-1	4	-1	7	-3	-2	1	-3	2
0	4	2	-5	2	1	-3	5	-4	1

	Synergy Score
A x 1	1.25
A self-cross	0.333
1 x self-cross	0.226

Figure 3-Example 3; A-427 cells





Figure 3a-Example 3; A-427 cells



	Synergy Score
B x 2	5.73
B self-cross	0.712
2 self-cross	0.169

Figure 4-Example 4; C32 cells





Figure 4a-Example 4; C32 cells



	Synergy Score
2653149 x 2112819	16.3
2653149 self	0.886
2112819 self	4.74

Figure 5-Example 5; C32 cells





Figure 5a-Example 5; C32 cells



	Synergy Score
B x 2	39.4
B self-cross	0.73
2 self-cross	3.27

Figure 6-Example 6; G-361 cells





Figure 6a-Example 6; G-361 cells



	Synergy Score
A x 1	5.97
A self-cross	0.0953
1 self-cross	2.26

Figure 7-Example 7; LS 174T cells





Figure 7a-Example 7; LS 174T cells



	Synergy Score
A x 1	1.76
A self-cross	0.578
1 self-cross	1.43

Figure 8-Example 8; MCF7 cells





Figure 8a-Example 8, MCF7 cells



	Synergy Score
A x 1	1.36
A self-cross	0.753
1 self-cross	0.955

Figure 9-Example 9; NCI-H1666 cells





Figure 9a-Example 9; NCI-H1666 cells



	Synergy Score
B x 1	4.3
B self-cross	0.117
1 self-cross	0.532

Figure 10-Example 10; NCI-H1666 cells





Figure 10a-Example 10; NCI-H1666 cells



B x 2	2.18
B self-cross	0.117
2 self-cross	0.233

Figure 11-Example 11; RKO cells







Figure 11a-Example 11; RKO cells




	Synergy Score
A x 1	8.21
A self-cross	0.235
1 self-cross	1.33

Figure 12-Example 12; RKO cells





Example 12a-Example 12; RKO cells



	Synergy Score
B x 2	7.6
B self-cross	0.154
2 self-cross	1.73

Figure 13-Example 13; RT4 cells





Figure 13a-Example 13; RT4 cells



	Synergy Score
B x 1	3.5
B self-cross	1.37
1 self-cross	0.835

Figure 14-Example 14; RT4 cells





Figure 14a-Example 14; RT4 cells



	Synergy Score
B x 2	2.05
B self-cross	1.33
2 self-cross	0.405

Figure 15-Example 15; SH-4 cells





Figure 15a-Example 15; SH-4 cells



	Synergy Score
B x 1	8.84
B self-cross	0.259
1 self-cross	2.21

Figure 16-Example 16; SH-4 cells





Figure 16a-Example 16; SH-4 cells



	Synergy Score
B x 2	12.1
B self-cross	0.29
2 self-cross	2.71

Figure 17-Example 17; SK-HEP-1





Figure 17a-Example 17; SK-HEP-1



	Synergy Score
B x 1	4.51
B self-cross	0.886
1 self-cross	0.923

Figure 18-Example 18; SK-HEP-1 cells





Figure 18a-Example 18; SK-HEP-1 cells



	Synergy Score
B x 2	4.95
B self-cross	0.827
2 self-cross	0.738

Figure 19-Example 19; A204 cells





Figure 19a-Example 19; A204 cells



	Synergy Score
A x 3	4.71
A self-cross	0.621
3 self-cross	0.278

Figure 20-Example 20; A375sq2 cells





Figure 20a-Example 20; A375sq2 cells



	Synergy Score
A x 3	1.3
A self-cross	0.333
3 self-cross	0.101

Figure 21-Example 21; A-427 cells





Figure 21a-Example 21; A-427 cells



	Synergy Score
B x 3	10.7
B self-cross	0.712
3 self-cross	1.34

Figure 22-Example 22; A-427 cells





Figure 22a-Example 22; A-427 cells



	Synergy Score
B x4	26
B self-cross	1.67
4 self-cross	3.27

Figure 23-Example 23; C32 cells





Figure 23a-Example 23; C32 cells



	Synergy Score
B x 3	22.9
B self-cross	0.886
3 self-cross	4.69

Figure 24-Example 24; C32 cells





Figure 24a-Example 24; c32 cells



	Synergy Score
B x 4	37.7
B self-cross	0.73
4 self-cross	4.59

Figure 25-Example 25; G-361 cells





Figure 25a-Example 25; G-361 cells



	Synergy Score
A x 3	4.36
A self-cross	0.144
3 self-cross	0.697

Figure 27-Example 27; LS 174T cells





Figure 27a-Example 27; LS 174T cells



	Synergy Score
A x 3	5.85
A self-cross	0.578
3 self-cross	0.257

Figure 28-Example 28; MCF7 cells





Figure 28a-Example 28; MCF7 cells



	Synergy Score
A x 3	1.31
A self-cross	0.753
3 self-cross	0.805

Figure 29-Example 29; NCI-H1666 cells





Figure 29a-Example 29; NCI-H1666 cells



	Synergy Score
B x 3	6.2
B self-cross	0.117
3 self-cross	2.22

Figure 30-Example 30; NCI-H1666 cells





Figure 30a-Example 30; NCI-H1666 cells



	Synergy Score
B x 4	8.97
B self-cross	0.117
4 self-cross	0.602

Figure 31-Example 31; RKO cells





Figure 31a-Example 31; RKO cells



	Synergy Score
A x 3	17.6
A self-cross	0.235
3 self-cross	0.368

Figure 32-Example 32; RKO cells





Figure 32a-Example 32; RKO cells



	Synergy Score
A x 4	29.3
B self-cross	0.154
4 self-cross	2.47

Figure 33-Example 33; RT4 cells





Figure 33a-Example 33; RT4 cells



	Synergy Score
B x 3	4.67
B self-cross	1.37
3 self-cross	0.666

Figure 34-Example 34; RT4 cells





Figure 34a-Figure 34; RT4 cells



	Synergy Score
B x 4	36.8
B self-cross	1.33
4 self-cross	1.23

Figure 35-Example 35; SH-4 cells





Figure 35a-Example 35; SH-4 cells



	Synergy Score
B x 3	9.82
B self-cross	0.259
3 self-cross	3.29

Figure 36-Example 36; SH-4 cells





Figure 36a-Example 36; SH-4 cells



	Synergy Score
B x 4	9.13
B self-cross	0.29
4 self-cross	1.93

Figure 37-Example 37; SK-HEP-1 cells





Figure 37a-Example 37; SK-HEP-1 cells



	Synergy Score
B x 3	9.09
B self-cross	0.886
3 self-cross	0.538

Figure 38-Example 38; SK-HEP-1 cells





Figure 38a-Example 38; SK-HEP-1 cells



	Synergy Score
B x 4	9.38
B self-cross	0.118
4 self-cross	0.901

Figure 39-Example 39; A204 cells





Figure 39a-Example 39; A204 cells



	Synergy Score
A x 5	4.62
A self-cross	0.621
5 self-cross	1.57

Figure 40-Example 40; A375sq2 cells





Figure 40a-Example 40; A375sq2 cells



	Synergy Score
A x 5	2.14
A self-cross	0.333
5 self-cross	0.294

Figure 41-Example 41; CAL-51 cells





Figure 41a-Example 41; CAL-51 cells



	Synergy Score
B x 5	26.4
B self-cross	0.386
5 self-cross	3.85

Figure 42-Example 42; G-361 cells





Figure 42a-Example 42; G-361 cells



	Synergy Score
A x 5	4.48
A self-cross	0.144
5 self-cross	1.93

Figure 43-Example 43; HT-1197 cells





Figure 43a-Example 43; HT1197 cells



	Synergy Score
B x 5	13.8
B self-cross	0.901
5 self-cross	0.799

Figure 44-Example 44; LS 174T cells





Figure 44a-Example 44; LS 174T cells



	Synergy Score
A x 5	4.56
A self-cross	0.578
5 self-cross	0.312

Figure 45-Example 45; MCF7 cells





Figure 45a-Example 45; MCF7 cells



	Synergy Score
A x 5	6.32
A self-cross	0.753
5 self-cross	2.48

Figure 46-Example 46; NCI-H460 cells





Figure 46a-Example 46; NCI-H460 cells



	Synergy Score
B x 5	10.1
B self-cross	0.496
5 self-cross	0.683

Figure 47-Example 47; RKO cells





Figure 47a-Example 47; RKO cells



	Synergy Score
A x 5	3.06
A self-cross	0.235
5 self-cross	0.485

Figure 48-Example 48; A204 cells





Figure 48a-Example 48; A204 cells



	Synergy Score
B x 6	19.4
B self-cross	0.792
6 self-cross	3.52

Figure 49-Example 49; A2780 cells





Figure 49a-Example 49; A2780 cells



	Synergy Score
B x 6	20
B cross-self	0.561
6 cross-self	4.38

Figure 50-Example 50; C32 cells





Figure 50a- Example 50; C32 cells



	Synergy Score
B x 6	15.7
B self-cross	1.09
6 self-cross	1.75

Figure 51-Example 51; G-401 cells





Figure 51a-Example 51; G-401 cells



	Synergy Score
B x 6	26.8
B self-cross	0.548
6 self-cross	4.14

Figure 52-Example 52; SK-HEP-1 cells





Figure 52a-Example 52; SK-HEP-1 cells



	Synergy Score
B x 6	10.3
B self-cross	0.827
6 self-cross	0.98

Figure 57-Example 57; BV-173 cells





Figure 57a-Example 57; BV-173 cells



	Synergy Score
B x 8	13.3
B self-cross	2.68
8 self-cross	4.72

Figure 58-Example 58; CML-T1 cells





Figure 58a-Example 58; CML-T1 cells



	Synergy Score
B x 8	32.6
B self-cross	3.43
8 self-cross	6.61

Figure 59-Example 59; KNS-81-FD cells





Figure 59a-Example 59; KNS-81-FD cells



	Synergy Score
B x 9	1.92
B self-cross	1.34
9 self-cross	0.103

Figure 60-Example 60; SW48 cells





Figure 60a-Example 60; SW48 cells



	Synergy Score
B x 9	5.06
B self-cross	2.23
9 self-cross	0.59

Figure 61-Example 61; MDA-MB-175-VII cells





Figure 61a-Example 61; MDA-MB-175-VII cells



	Synergy Score
B x 10	5.29
B self-cross	1.53
10 self-cross	3.11

Figure 62-Example 62; UACC-812 cells





Figure 62a-Example 62; UACC-812 cells



	Synergy Score
B x 10	9.93
B self-cross	2.74
10 self-cross	2.06

Figure 63-Example 63; HCT 116 cells





Figure 63a-Example 63; HCT 116 cells



	Synergy Score
A x 11	1.32
A self-cross	1.27
11 self-cross	0.244

Figure 64-Example 64; GDM-1 cells





Figure 64a-Example 64; GDM-1 cells



	Synergy Score
B x 13	9.25
B self-cross	9.93
13 self-cross	2.71

Figure 65-Example 65; ML-2 cells





Figure 65a-Example 65; ML-2 cells



	Synergy Score
B x 13	7.24
B self-cross	14.9
13 self-cross	1.06

Figure 66-Example 66; MOLM-13 cells





Figure 66a-Example 66; MOLM-13 cells



	Synergy Score
B x 13	5.66
B self-cross	.428
13 self-cross	.146

Figure 67-Example 67; OCI-AML3 cells






Figure 67a-Example 67; OCI-AML3 cells




	Synergy Score
B x 13	9.62
B self-cross	3.91
13 self-cross	1.75

Figure 68-Example 68; GDM-1 cells



Example 68a-Figure 68; GDM-1 cells



	Synergy Score
B x 12	12.5
B self-cross	9.93
12 self-cross	7.87

Figure 69-Example 69; ML-2 cells





Figure 69a-example 69; ML-2 cells



	Synergy Score
B x 12	11
B self-cross	14.9
12 self-cross	8.62

Figure 70-Example 70; MOLM-13 cells





Figure 70a-Example70; MOLM-13 cells



	Synergy Score
B x 12	15.4
B self-cross	0.428
12 self-cross	1.7

Figure 71-Example 71; OCI-AML3 cells





Figure 71a-Example 71; OCI-AML3 cells



	Synergy Score
B x 12	13.6
B self-cross	3.91
12 self-cross	5.81

Figure 72
AMG 232 – MAPKi Combinations




Figure 73
AM-7209 – MAPKi Combinations




Figure 74
RG7112 – MAPKi Combinations




Figure 75
AMG 232 – PI3Ki Combinations




Figure 76
AM-7209 – PI3Ki Combinations




Figure 77

RG7112 – PI3Ki Combinations




Figure 78

AMG 232 – Intrinsic Apoptosis Combinations




Figure 79

AM-7209 – Intrinsic Apoptosis Combinations




Figure 80
RG7112 – Intrinsic Apoptosis Combinations




Figure 81
AMG 232 – Miscellaneous Combinations




Figure 82
AM-7209 – Miscellaneous Combinations




Figure 83
RG7112 – Miscellaneous Combinations




Figure 84

AMG 232 – Chemotherapeutic Combinations

		MCF7 (Breast)	RKO (Colon)	KS-1 (GBM)	A427 (NSCLC)	SJSA-1 (Sarcoma)	SW982 (Sarcoma)	MKN45 (Stomach)	NCI-SNU-1 (Stomach)	EOL-1 (AML)	MOLM-13 (AML)	TP53 HT-29 (Colon)	TP53 PC-3 (Prostate)
Platinums	AMG 232 x Cisplatin				0.70							0.19	0.48
	AMG 232 x Oxaliplatin		8.85									0.08	0.18
Topoisomerase II	AMG 232 x Doxorubicin	4.63								4.37	8.63	1.14	2.14
	AMG 232 x Etoposide					2.16	9.05	2.86	1.88			1.92	0.52
Topoisomerase I	AMG 232 x Irinotecan		1.69									0.95	0.67
DNA Alkylation	AMG 232 x Temozolomide			2.61								0.04	0.18
Nucleoside Analogs	AMG 232 x Cytarabine									6.90	8.28	0.56	1.33
	AMG 232 x Decitabine									12.49	13.71	0.07	0.86

Figure 85

AM-7209 – Chemotherapeutic Combinations




Figure 86
RG7112 – Chemotherapeutic Combinations




Figure 87

AMG 232 – Haematopoietic Lines




Figure 88

AM-7209 – Haematopoietic Lines




Figure 89

RG7112 – Haematopoietic Lines




Figure A

MDM2 Inhibitor and Cisplatin in H460 Tumor




Figure B

MDM2 Inhibitor and Cisplatin in HCT116 Tumor





Figure C

MDM2 Inhibitor and CPT-11 in HCT116 Tumor




Vehicle (2705932): 15% HPßCD, 1% Pluronic F68, pH 8


Figure D

MDM2 inhibitor and Doxorubicin in SJSA-1 Tumors

MDM2 Inhibition + Doxorubicin (5mpk)




MDM2 Inhibition + Doxorubicin (10mpk)




Vehicle (2705932): 15% HPßCD, 1% Pluronic F68, pH 9.4

Figure E
MDM2 Inhibitor and BRAF or MEK Inhibition in RKO Tumors

MDM2 Inhibition + BRAF Inhibition




MDM2 Inhibition + MEK Inhibition



1 DIC in 1009089 group on day 11 (N=9)
1 DIC in '819/'932 combination group on day 17 (N=9)
Vehicle (2112819/1009089): 2% HPMC, 1% Tween 80, pH 2.2
Vehicle (2705932): 15% HPßCD, 1% Pluronic F68, pH 8


Blinded study (daily oral dosing)


Figure F
MDM2 Inhibitor and PI3K Inhibitor in U87 Tumor



All curves are derived from a single blinded study (dai)ly oral dosing

Figure G
MDM2 Inhibitor and MEK Inhibitor in A375 Tumors






Group	% TGI
Vehicle+ Vehicle	0.0
Vehicle + 2705932 100 mg/kg	90.5
1009089 5 mg/kg+ Vehicle	101.6
1009089 5 mg/kg+ 2705932 100 mg/kg	120.1

Group	% TGI
Vehicle+ Vehicle	0.0
Vehicle+2705932 5 mg/kg	29.1
1009089 0.5 mg/kg+ Vehicle	43.7
1009089 0.5 mg/kg+ 2705932 5 mg/kg	57.0

Vehicle (1009089): 2% HPMC, 1% Tween 80, pH 2.2

Vehicle (2705932): 15% HPßCD, 1% Pluronic F68, pH 8


Figure H
MDM2 Inhibitor and BRAF Inhibitor in A375sq2 Tumors



* p<0.0001 compare to 2705932 100mg/kg

Vehicle (2112819): 2% HPMC, 1% Tween 80, pH 2.2
Vehicle (2705932): 15% HPßCD, 1% Pluronic F68, pH 8

Figure I
Combination of MDM2, PI3K and BRAF Inhibition in RKO Tumors



Group	% TGI
Vehicle 2 (pH2.2) + Vehicle 1 (pH 9.4)	0.0
Vehicle 2 (pH2.2) + 2705932 100 mpk	21.7
2539965 10 mpk + Vehicle 1 (pH 9.4)	76.4
2112819 10 mpk + Vehicle 1 (pH 9.4)	19.5
2539965 10 mpk + 2705932 100 mpk	85.7
2112819 10 mpk + 2705932 100 mpk	96.6
2539965 +2112819+ Vehicle 1 (pH 9.4)	83.2
2539965 +2112819+2705932 100 mg/kg	R= 30.7

Blinded Study

RMANOVA: p values are comparisons with the triple combination group

* $p < 0.0001$


$p = 0.0090$


Vehicle 1 (2705932): 15% HPßCD, 1% Pluronic F68, pH 9.4


Vehicle 2 (2112819/2539965): 15% HPßCD, 1% Pluronic F68, pH 2.2

Figure J

Combination of MDM2, PI3K and BRAF Inhibition in the RKO Tumor Model (single agents vs. double combinations)








Blinded study:
RMANOVA p values are comparisons to the double combination groups

Vehicle 1 (2705932): 15% HPßCD, 1% Pluronic F68, pH 9.4
Vehicle 2 (2112819/2539965): 15% HPßCD, 1% Pluronic F68, pH 2.2


Figure K

MDM2 Inhibitor and Doxorubicin in Molm13 Tumors

MDM2 Inhibition + Doxorubicin (5mpk)




MDM2 Inhibition + Doxorubicin (10mpk)



Vehicle (2705932): 15% HPßCD, 1% Pluronic F68, pH 9.4


Figure L

MDM2 Inhibitor and MEK inhibitor in Molm13 Tumors



Vehicle (2705932): 15% HPßCD, 1% Pluronic F68, pH 9.4
Vehicle (1009089): 2% HPMC, 1% Tween 80, pH 2.2


Figure M
MDM2 Inhibitor and Cytarabine in Molm13 Tumors



Vehicle: 15% HPβCD, 1% Pluronic F68, pH 9.4 and Saline

Figure N


MDM2 Inhibitor and Decitabine in Molm13 Tumors



Vehicle (2705932): 15% HPßCD, 1% Pluronic F68, pH 9.4

Figure O

MDM2 Inhibition and Sorafenib inMolm13 Tumors



Vehicle (674064): 2% HPMC, 1% Tween 80, pH 2.2

Vehicle (2705932): 15% HPßCD, 1% Pluronic F68, pH 9.4