

BLOOD BRAIN BARRIER PENETRATING SELECTIVE INHIBITORS OF ROCK2 PROTEIN KINASES AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATION(S)

5 This application claims priority from U.S. Provisional Application No. 63/496,836 filed April 18, 2023, which is incorporated herein by reference in its entirety.

TECHNICAL FIELD

10 The present invention relates to the novel substituted bicyclic derivatives that can selectively inhibit ROCK2 and/or Rho-kinase mediated phosphorylation of myosin light chain phosphates, the process for their preparation, and formulation thereof and their use for preparing a medicament, as well as pharmacologically acceptable salts, diastereomers, enantiomers, racemates.

BACKGROUND

15 The Rho-associated coiled-coil containing protein kinases (ROCKs/Rho-kinase/Rho-associated kinase) are downstream effectors of the small GTPase Rho (Rho A, Rho B, Rho C, and Rho E) and belong to the family of serine/threonine kinases. The active GTP bound form of Rho mediates several biological functions through the action of ROCKs, including smooth muscle contractions, cell motility, and cytokinesis. The ROCK proteins were identified in 1996 as proteins that bind to Rho GTPase. Two proteins were independently isolated as p160 and
20 p164. Later, these were recognized as ROCK-1 and ROCK-2, respectively and as isoforms of the Rho-associated kinase. The two isoforms, ROCK-1 (ROCK-b/p160) and ROCK-2 (ROCK-a/p164), share 92% similarity in their amino acid sequence. Their structure comprises an N-terminally located catalytic kinase domain, followed by a coiled-coil containing region (600 amino acids) with a Rho-binding domain and a pleckstrin homology (PH) domain at the C
25 terminus. They have varying locations, and distinct physiological roles have been identified for each. The ROCK-1 transcript (gene located on chromosome 18) is ubiquitous, with more prominent expression in liver, kidney, spleen, testis, thymus, and blood corpuscles, whereas


ROCK-2 mRNA (chromosome 2) is expressed more abundantly in skeletal muscles and brain, suggesting that they have specialized roles in these locations.

ROCK inhibitors have been considered for use in numerous diseases, such as cerebral ischemia, hypertension, erectile dysfunction, glaucoma, osteoporosis, cardiac hypertrophy, diabetic cardiomyopathy, retinopathy, pulmonary hypertension, and atherosclerosis. However, their implementation is limited because of a lack of knowledge regarding the involvement of the particular ROCK isoform. Whether isoform-specific targeting or combined ROCK inhibition would provide a better therapeutic outcome is yet to be verified. Despite their incomplete specificity towards the ROCK isoforms as well as other serine/threonine kinases, such as PRK2, PKC, cAMP-dependent protein kinase, and citron kinase, some nonspecific ROCK2 inhibitors have shown promising results in certain pathological states, such as glaucoma and hypertension. However, further work is required for isoform-selective ROCK2 inhibitors to be of clinical use, although several research-based studies have helped resolve much ambiguity over ROCK-1 and ROCK-2-specific functions. Since ROCKs play a central role in the organization of the actin cytoskeleton, it might be anticipated that the complete inhibition of both isoforms could cause adverse events in patients. Therefore, selective ROCK2 inhibition could have a greater specific therapeutic index over the dual inhibition of ROCK1 and ROCK2. Thus, understanding the discreet functions of each isoform in a particular disorder could help improve safety and specificity and expand the therapeutic applications of ROCK inhibitors.

20

SUMMARY

In one aspect, the present invention provides novel chemical compounds represented by (1) below, which is capable of inhibiting Rho-associated coiled-coil forming protein serine/threonine kinases (ROCKs):



wherein X, Y, Z¹, Z², Z³, R¹, R², R³ and R⁴ are defined in the detailed description of the invention below.

In another aspect, the present invention provides pharmaceutically acceptable salts, diastereomers, enantiomers, racemates, hydrates, solvates, prodrugs of the novel compounds.

5 Also, the present invention provides pharmaceutically acceptable salts, hydrates, or solvates of the diastereomers, enantiomers, or racemates.

In still another aspect, the present invention provides pharmaceutical compositions each comprising the compound, salt, diastereomer, enantiomer, racemate, hydrate, solvate, prodrug, or a combination thereof.


10 In yet another aspect, the present invention provides methods of treating or alleviating certain ROCK-mediated diseases or disorders by using the compounds, salts, diastereomers, enantiomers, racemates, hydrates, solvates, prodrugs, or compositions. Non-limiting examples of the diseases or disorders include cardiovascular, pulmonary, inflammatory, neurological, or proliferative diseases or disorders.

15 In a further aspect, the present invention provides methods of preparing the compounds, salts, diastereomers, enantiomers, racemates, hydrates, solvates, and prodrugs.

DETAILED DESCRIPTION

1. Compounds

20 In one aspect, the present invention provides a compound represented by Formula (1), a pharmaceutically acceptable salt, diastereomer, enantiomer, racemate, hydrate, solvate, or prodrug thereof:



wherein:

X is CH₂ or O;

Y is N or CH;

Z^1 , Z^2 and Z^3 each are independently N or CH;

n is 0, 1, 2 or 3;

R^1 is H, heterocycloalkyl, C₃-C₇cycloalkyl, or C₁-C₆alkyl wherein a nitrogen on
5 heterocycloalkyl can be optionally substituted with C₁-C₃alkyl, and wherein the C₃-C₇cycloalkyl
or C₁-C₆alkyl can be optionally substituted with one or more suitable substituents, for example,
C₁-C₆alkyl or OH;

R^2 is H, amino, NR⁶R⁷, or NR⁸R⁹;

R^3 is 5-6 membered heteroaryl or bicyclic heteroaryl, wherein the heteroaryl or bicyclic
10 heteroaryl can be optionally substituted with one or more suitable substituents, for example,
halogen, CN, CHF₂, CF₃, C₁-C₃alkyl, or amino, wherein the 5-6 membered heteroaryl or bicyclic
heteroaryl has 1-3 heteroatoms selected from the group consisting of oxygen and nitrogen;


R^4 and R^5 each are independently H, Cl, F, OH, OCD³-C₁-C₃alkyl, or C₁-C₃alkoxy,
wherein the C₁-C₃alkyl or C₁-C₃alkoxy can be optionally substituted with one or more suitable
15 substituents, for example, amino, OH or OMe, OEt, or OPrⁱ;

R^6 and R^7 each are independently H, CD₃, C₁-C₆alkyl, or C₃-C₇cycloalkyl, wherein the
C₁-C₆alkyl or C₃-C₇cycloalkyl can be optionally substituted one or more suitable substituents, for
example, hydroxyl or C₁-C₃alkoxy; and

R^7 and R^8 , taken together with the nitrogen atom to which they are bonded a 4-6
20 membered saturated monocyclic group having no heteroatom other than the nitrogen atom to
which R^7 and R^8 are bonded, wherein the 4-6 membered saturated or partially saturated
monocyclic group is optionally substituted at one or two carbon atoms with, halogen, amino,
hydroxyl or C₁-C₃alkoxy.

In some embodiments, R^1 may be H, heterocycloalkyl C₃-C₇cycloalkyl, or C₁-C₆alkyl
25 wherein a nitrogen on heterocycloalkyl can be optionally substituted with methyl or ethyl, wherein
the C₃-C₇cycloalkyl or C₁-C₆alkyl can be optionally substituted with one or more suitable
substituents, for example, C₁-C₃alkyl or OH.

In some embodiments, R^3 may be one of the following group:



In some embodiments, R^4 and R^5 each may be independently H, Cl, F, OH, CH_3 ,
 5 methoxy, OCD_3 , ethoxy, or iso-propoxy.

In some embodiments, R^6 and R^7 each may be independently H, CD_3 , C_1 - C_6 alkyl, or C_3 - C_7 cycloalkyl, wherein the C_1 - C_6 alkyl or C_3 - C_7 cycloalkyl is optionally substituted with, hydroxyl methoxy or ethoxy.

In some embodiments R^7 and R^8 , taken together with the nitrogen atom to which they are
 10 bonded a 4-6 membered saturated monocyclic group having no heteroatom other than the nitrogen atom to which R^7 and R^8 are bonded, wherein the 4-6 membered saturated or partially saturated monocyclic group is optionally substituted at one or two carbon atoms with, halogen, amino, hydroxyl or methoxy.

The articles “a” and “an” are used in this disclosure to refer to one or more than one (i.e.,
 15 to at least one) of the grammatical objects of the article. By way of example, “an element” means one element or more than one element.

The term “and/or” is used in this disclosure to mean either “and” or “or” unless indicated
 otherwise. The use of the term “or” is used to mean “and/or” unless explicitly indicated to refer
 to alternatives only or the alternative are mutually exclusive, although the disclosure supports a
 20 definition that refers to only alternatives and “and/or.”

By "optional" or "optionally," it is meant that the subsequently described event or circumstance may or may not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "optionally substituted aryl" encompasses both "aryl" and "substituted aryl" as defined herein. It will be understood by those ordinarily skilled in the art, with respect to any group containing one or more substituents, that such groups are not intended to introduce any substitution or substitution patterns that are sterically impractical, synthetically non-feasible, and/or inherently unstable.

The term "optionally substituted" unless otherwise specified means that a group may be unsubstituted or substituted by one or more (e.g., 0, 1, 2, 3, 4, or 5 or more, or any range derivable therein) of the substituents listed for that group in which said substituents may be the same or different. In an embodiment, an optionally substituted group has 1 substituent. In another embodiment, an optionally substituted group has 2 substituents. In another embodiment, an optionally substituted group has 3 substituents. In another embodiment, an optionally substituted group has 4 substituents. In another embodiment, an optionally substituted group has 5 substituents. For instance, an alkyl group that is optionally substituted can be a fully saturated alkyl chain (i.e., a pure hydrocarbon). Alternatively, the same optionally substituted alkyl group can have substituents different from hydrogen. For instance, it can, at any point along the chain be bonded to a halogen atom, a hydroxyl group, or any other substituent described herein. Thus, the term "optionally substituted" means that a given chemical moiety has the potential to contain other functional groups, but does not necessarily have any further functional groups.

The term "alkyl," used alone or as part of a larger moiety such as "arylalkyl" or "cycloalkyl" refers to a straight or branched hydrocarbon radical having from 1 to 10 carbon atoms, or from 1-8 carbon atoms, or from 1-6 carbon atoms, or from 1-4 carbon atoms (unless stated otherwise) and includes, for example, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, isobutyl, *tert*-butyl, *n*-pentyl, *iso*-pentyl, *n*-hexyl and the like. An alkyl can be unsubstituted or substituted with one or more suitable substituents.

The term "cycloalkyl" refers to a monocyclic or polycyclic hydrocarbon ring group having from 3 to 10 carbon atoms, or from 3 to 7 carbon atoms, in the hydrocarbon ring (unless stated otherwise) and includes, for example, cyclopropyl, cycloheptyl, cyclooctyl, cyclodecyl,

cyclobutyl, adamantyl, norpinanyl, decalanyl, norbornyl, cyclohexyl, cyclopentyl, and the like. A cycloalkyl group can be unsubstituted or substituted with one or more suitable substituents.

The term "hetero" refers to the replacement of at least one carbon atom member in a ring system with at least one heteroatom such as nitrogen, sulfur, sulfoxide, sulfone, and oxygen.

5 The term "heterocycloalkyl" means a non-aromatic saturated or a non-aromatic unsaturated monocyclic or a non-aromatic saturated or non-aromatic unsaturated polycyclic ring having from 2 to 9 carbon atoms, or from 2 to 7 carbon atoms, in the ring (unless stated otherwise) and at least one heteroatom, preferably, 1 to 4 heteroatoms selected from nitrogen, sulfur (including oxidized sulfur such as sulfone or sulfoxide) and oxygen. The ring or ring
10 system of the heterocycloalkyl group can be linked to another moiety of the compound via a carbon atom or a nitrogen atom, if such an atom is present. A heterocycloalkyl group can have a total of 3-10, or 3-8, or 5-8, atoms in the ring system (unless otherwise stated). A heterocycloalkyl group can have one or more carbon-carbon double bonds or carbon-heteroatom double bonds in the ring group as long as the ring group is not rendered aromatic by their
15 presence.

Examples of heterocycloalkyl groups include azetidiny, aziridiny, pyrrolidiny, piperidiny, piperaziny, homopiperaziny, morpholino, thiomorpholino, tetrahydrofurany, tetrahydrothiofurany, tetrahydropyrany, pyranly, and the like. A heterocycloalkyl group can be unsubstituted or substituted with one or more suitable substituents.

20 As used herein, the term "halo" includes fluoro, chloro, bromo, and iodo.

As used herein, the term "alkoxy" refers to the alkyl groups above bound through oxygen, examples of which include methoxy, ethoxy, *iso*-propoxy, *tert*-butoxy, and the like. In addition, alkoxy also refers to polyethers such as -O-(CH₂)₂-O-CH₃, and the like. An alkoxy can be unsubstituted or substituted with one or more suitable substituents.

25 As used herein, the term "aryl" refers to unsubstituted or substituted aromatic monocyclic or polycyclic groups and includes, for example, carbocyclic aromatic groups such as phenyl, naphthyl and the like, as well as heteroaromatic groups such as pyridyl, furanyl, thiophenyl, and the like. The term "aryl" also includes an aromatic ring (such as a phenyl or pyridyl ring) fused to a non-aromatic carbocyclic or heterocyclic ring. The term "aryl" may be interchangeably used

with "aryl ring," aromatic group," and "aromatic ring." Heteroaryl groups have 4 to 14 atoms in the heteroaromatic ring(s), 1 to 9 of which are independently selected from the group consisting of oxygen, sulfur and nitrogen. Heteroaryl groups have 1-3 heteroatoms in a 5-8 membered aromatic group. An aryl or heteroaryl can be a mono- or bicyclic aromatic group. Typical aryl and heteroaryl groups include, for example, phenyl, quinolinyl, indazolyl, indolyl, dihydrobenzodioxynyl, 3-chlorophenyl, 2,6-dibromophenyl, pyridyl, pyrimidinyl, 3-methylpyridyl, benzothienyl, 2,4,6-tribromophenyl, 4-ethylbenzothienyl, furanyl, 3,4-diethylfuranyl, naphthyl, 4,7-dichloronaphthyl, pyrrole, pyrazole, imidazole, thiazole, and the like. An aryl or heteroaryl can be unsubstituted or substituted with one or more suitable substituents.

As used herein, the term "hydroxyl" or "hydroxy" refers to -OH.

As used herein, the term "amino" refers to -NH₂.

As used herein, the term "hydroxyalkyl" refers to any hydroxyl derivative of alkyl radical. The term "hydroxyalkyl" includes any alkyl radical having one or more hydrogen atoms replaced by a hydroxy group.

As used herein, the term "arylalkyl" includes any alkyl radical having one or more hydrogen atoms replaced by an aryl group, e.g., a benzyl group, a phenethyl group, and the like.

A "substituent," as used herein, refers to a molecular moiety that is covalently bonded to an atom within a molecule of interest. For example, a ring substituent may be a moiety such as a halogen, alkyl group, haloalkyl group or other group that is covalently bonded to an atom (preferably a carbon or nitrogen atom) that is a ring member. Substituents of aromatic groups are generally covalently bonded to a ring carbon atom. The term "substitution" refers to replacing a hydrogen atom in a molecular structure with a substituent, such that the valence on the designated atom is not exceeded, and such that a chemically stable compound (*i.e.*, a compound that can be isolated, characterized, and tested for biological activity) results from the substitution.

As described above, certain groups can be unsubstituted or substituted with one or more suitable substituents by other than hydrogen at one or more available positions, typically 1, 2, 3, 4 or 5 positions, by one or more suitable groups (which may be the same or different). Certain

groups, when substituted, are substituted with 1, 2, 3 or 4 independently selected substituents. Suitable substituents include halo, alkyl, haloalkyl, aryl, hydroxy, alkoxy, hydroxyalkyl, amino, and the like.

As used herein, the term “pharmaceutically acceptable” refers to a material, such as a carrier or diluent, which does not abrogate the biological activity or properties of the compounds described herein. Such materials are administered to an individual without causing undesirable biological effects or interacting in a deleterious manner with any of the components of the composition in which it is contained.

As used herein, the term “pharmaceutically acceptable salt” refers to a formulation of a compound that does not cause significant irritation to an organism to which it is administered and does not abrogate the biological activity and properties of the compounds described herein.

Pharmaceutically acceptable salt forms include pharmaceutically acceptable acidic/anionic or basic/cationic salts (UK Journal of Pharmaceutical and Biosciences Vol. 2(4), 01-04, 2014, which is incorporated herein by reference). Pharmaceutically acceptable acidic/anionic salts include acetate, benzenesulfonate, benzoate, bicarbonate, bitartrate, bromide, calcium edetate, camsylate, carbonate, chloride, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, glyceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isethionate, lactate, lactobionate, malate, maleate, malonate, mandelate, mesylate, methylsulfate, mucate, napsylate, nitrate, pamoate, pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, sulfate, hydrogensulfate, tannate, tartrate, teoclate, tosylate, and triethiodide salts. Pharmaceutically acceptable basic/cationic salts include, the sodium, potassium, calcium, magnesium, diethanolamine, *N*-methyl-*D*-glucamine, *L*-lysine, *L*-arginine, ammonium, ethanolamine, piperazine and triethanolamine salts.

A pharmaceutically acceptable acid salt is formed by reaction of the free base form of a compound of Formula 1 with a suitable inorganic or organic acid including, but not limited to, hydrobromic, hydrochloric, sulfuric, nitric, phosphoric, succinic, maleic, formic, acetic, propionic, fumaric, citric, tartaric, lactic, benzoic, salicylic, glutamic, aspartic, *p*-toluenesulfonic, benzenesulfonic, methanesulfonic, ethanesulfonic, naphthalenesulfonic such as 2-naphthalenesulfonic, or hexanoic acid. A pharmaceutically acceptable acid addition salt of a

compound of Formula 1 can comprise or be, for example, a hydrobromide, hydrochloride, sulfate, nitrate, phosphate, succinate, maleate, formate, acetate, propionate, fumarate, citrate, tartrate, lactate, benzoate, salicylate, glutamate, aspartate, *p*-toluenesulfonate, benzenesulfonate, methanesulfonate, ethanesulfonate, naphthalenesulfonate (e.g., 2-naphthalenesulfonate) or
5 hexanoate salt.

The free acid or free base forms of the compounds of the invention may be prepared by methods known to those of ordinary skill in the art (e.g., for further details see L.D. Bigley, S.M. Berg, D.C. Monkhouse, in “*Encyclopedia of Pharmaceutical Technology*”. Eds, J. Swarbrick and J.C. Boylam, Vol 13, Marcel Dekker, Inc., 1995, pp.453-499; the entire teachings of which are
10 incorporated herein by reference) from the corresponding base addition salt or acid addition salt form, respectively. For example, a compound of the invention in an acid addition salt form may be converted to the corresponding free base form by treating with a suitable base (e.g., ammonium hydroxide solution, sodium hydroxide, and the like). A compound of the invention in a base addition salt form may be converted to the corresponding free acid by treating with a
15 suitable acid (e.g., hydrochloric acid, etc.).

Prodrug derivatives of the compounds of the invention may be prepared by methods known to those of ordinary skill in the art (e.g., for further details see Saulnier *et al.*, *Bioorg. Med. Chem. Letters*, 1994, 4, 1985; Daniela Hartmann Jornada *et al.*, *Molecules* 2016, 21, 42; the entire teachings of which are incorporated herein by reference). Protected derivatives of the
20 compounds of the invention may be prepared by means known to those of ordinary skill in the art. A detailed description of techniques applicable to the creation of protecting groups and their removal can be found in T. W. Greene, “*Green’s Protective Groups in Organic Chemistry*,” 4th edition, John Wiley and Sons, Inc., 2006, the entire teachings of which are incorporated herein by reference.

Compounds of the invention may be prepared as their individual stereoisomers by
25 reaction of a racemic mixture of the compound with an optically active resolving agent to form a pair of diastereoisomeric compounds, separating the diastereomers and recovering the optically pure enantiomers. Resolution of enantiomers may be carried out using covalent diastereomeric derivatives of the compounds of the invention, or by using dissociable complexes (e.g.,
30 crystalline diastereomeric salts). Diastereomers have distinct physical properties (e.g., melting

points, boiling points, solubility, reactivity, etc.) and may be readily separated by taking advantage of these dissimilarities. The diastereomers may be separated by chromatography, or by separation/resolution techniques based upon differences in solubility. The optically pure enantiomer is then recovered, along with the resolving agent, by any practical means that would not result in racemization. A more detailed description of the techniques applicable to the resolution of stereoisomers of compounds from their racemic mixture can be found in Jean Jacques, Andre Collet and Samuel H. Wilen, "*Enantiomers, Racemates and Resolutions*," John Wiley And Sons, Inc., 1981, the entire teachings of which are incorporated herein by reference.

As used herein, the term "solvate" refers to a complex of variable stoichiometry formed by a solute (in this invention, a compound of Formula 1 or a pharmaceutically acceptable salt thereof) and a solvent. Such solvents for the purpose of the invention may not interfere with the biological activity of the solute. Non-limiting examples of suitable solvents include water, acetone, methanol, ethanol and acetic acid. Preferably the solvent used is a pharmaceutically acceptable solvent. Non-limiting examples of suitable pharmaceutically acceptable solvents include water, ethanol and acetic acid.

2. Compositions

In another aspect, the present invention provides a pharmaceutical composition comprising the compound, salt, diastereomer, enantiomer, racemate, hydrate, solvate, prodrug, or a pharmaceutical combination thereof. The composition may further comprise an additional component. A non-limiting example of the additional component includes a pharmaceutically acceptable carrier, diluent, excipient, and a combination thereof.

As used herein, the term "pharmaceutical composition" refers to a mixture of a compound described herein with other chemical components, such as carriers, stabilizers, diluents, dispersing agents, suspending agents, thickening agents, and/or excipients.

As used herein, the term "pharmaceutical combination" means a product that results from the mixing or combining of more than one active ingredient.

As used herein, the term "acceptable" with respect to a formulation, composition or ingredient, as used herein, means having no persistent detrimental effect on the general health of the subject being treated.

As used herein, the term “carrier” refers to chemical compounds or agents that facilitate the incorporation of a compound described herein into cells or tissues.

As used herein, the term “diluent” refers to chemical compounds that are used to dilute a compound described herein prior to delivery. Diluents can also be used to stabilize compounds described herein.

Suitable pharmaceutically acceptable carriers, diluents, adjuvants, or excipients for use in the pharmaceutical compositions of the invention include tablets (coated tablets) made of for example collidone or shellac, gum Arabic, talc, titanium dioxide or sugar, capsules (gelatin), solutions (aqueous or aqueous-ethanolic solution), syrups containing the active substances, emulsions or inhalable powders (of various saccharides such as lactose or glucose, salts and mixture of these excipients with one another) and aerosols (propellant-containing or –free inhale solutions).

Excipients which may be used include, for example, water, pharmaceutically acceptable organic solvents such as paraffins (e.g., petroleum fractions), vegetable oils (e.g. groundnut or sesame oil), mono- or polyfunctional alcohols (e.g., ethanol or glycerol), carriers such as natural mineral powders (e.g., kaoline, clays, talc, chalk), synthetic mineral powders (e.g., highly dispersed silicic acid and silicates), sugars (e.g., cane sugar, lactose and glucose), emulsifiers (e.g., lignin, spent sulphite liquors, methylcellulose, starch and polyvinylpyrrolidone) and lubricants (e.g., magnesium stearate, talc, stearic acid and sodium lauryl sulphate).

3. Methods of Using Compounds or Compositions

In another aspect, the present invention provides methods of treating or alleviating certain protein kinase-mediated diseases or conditions by administering a subject or patient a therapeutically effective amount of the compound, salt, diastereomer, enantiomer, racemate, hydrate, solvate, prodrug, or composition. In some embodiments, the present invention provides methods of treating or alleviating certain ROCK-mediated diseases or disorders by administering a subject or patient a therapeutically effective amount of the compound, salt, diastereomer, enantiomer, racemate, hydrate, solvate, prodrug, or composition. In some embodiments, the present invention provides methods of treating or alleviating a disease or disorder in which ROCK is known to play a role.

In a further aspect, the present invention provides methods of inhibiting enzyme activity, particularly ROCK1, ROCK2, PKC δ , PKC θ , PRK1, GSK3, PRK2, NEK1 and NEK4 kinase activity, by administering a subject or patient a therapeutically effective amount of the compound, salt, diastereomer, enantiomer, racemate, hydrate, solvate, prodrug, or composition.

5 In yet another aspect, the present invention provides methods of inhibiting protein kinase activity, such as, for example, ROCK kinase activity, in a biological sample by contacting the biological sample with the compound, salt, diastereomer, enantiomer, racemate, hydrate, solvate, prodrug, or composition.

As used herein, the term “inhibitor” refers to a compound which inhibits one or more
10 kinases described herein. For example, the term “ROCK inhibitor” refers to a compound which inhibits the ROCK receptor or reduces the signaling effect.

As used herein, the term “protein kinase-mediated disease” or a “disorder or disease or condition mediated by inappropriate protein kinase activity” refers to any disease state mediated or modulated by protein kinases described herein. Such disease states include, but are not
15 limited to, fibrotic disorders; pulmonary fibrosis including cystic and idiopathic pulmonary fibrosis, radiation induced lung injury, liver fibrosis including cirrhosis, cardiac fibrosis including arterial fibrosis, endomyocardial fibrosis, old myocardial infarction, arterial stiffness, atherosclerosis, restenosis, arthrofibrosis, Crohn’s disease, myelofibrosis, Peyronie’s diseases, nephrogenic systemic fibrosis, progressive massive fibrosis, retroperitoneal cavity fibrosis,
20 scleroderma/systemic sclerosis, mediastinal fibrosis, Keloids and hypertrophic scars, glial scarring, or renal fibrosis; cardiovascular disease or disorder, such as, for example, cerebral vasospasm, hypertension, atherosclerosis, angina, myocardial infarction, ischemic/reperfusion injury, stroke, bronchial asthma; glaucoma, pre-term labor, erectile dysfunction, or renal disease, such as, for example, chronic renal failure, chronic nephritis, diabetic nephropathy, and IgA
25 nephropathy; a neurological disease or disorder, such as for example, spinal-cord injury, Alzheimer’s disease, multiple sclerosis, or neuropathic pain; and proliferative disorders, such as, for example, retinopathy, fibrosis, or invasive/metastatic cancers. Such cancers include adenocarcinoma, adrenocortical cancer; bladder cancer; bone cancer; brain cancer; breast cancer; cancer of the buccal cavity; cervical cancer; colon cancer; colorectal cancer; endometrial or
30 uterine carcinoma; epidermoid carcinoma; esophageal cancer; eye cancer; follicular carcinoma;

gallbladder cancer; gastrointestinal cancer; cancer of the genitourinary tract; glioblastoma; hairy cell carcinoma; head and neck cancer; hepatic carcinoma; hepatocellular cancer; Hodgkin's disease; keratoacanthoma; kidney cancer; large cell carcinoma; cancer of the large intestine; laryngeal cancer; liver cancer; lung cancer, such as, for example, adenocarcinoma of the lung, small-cell lung cancer, squamous carcinoma of the lung, non-small cell lung cancer; melanoma; myeloproliferative disorders; neuroblastoma; ovarian cancer; papillary carcinoma; pancreatic cancer; cancer of the peritoneum; prostate cancer; rectal cancer; salivary gland carcinoma; sarcoma; squamous cell cancer; small cell carcinoma; cancer of the small intestine; stomach cancer; testicular cancer; thyroid cancer; and vulval cancer. In particular embodiments, the treated cancer is melanoma, breast cancer, colon cancer, or pancreatic cancer.

As used herein, the term "treat," "treating" or "treatment" refers to methods of alleviating, abating or ameliorating a disease or condition symptoms, preventing additional symptoms, ameliorating or preventing the underlying metabolic causes of symptoms, inhibiting the disease or condition, arresting the development of the disease or condition, relieving the disease or condition, causing regression of the disease or condition, relieving a condition caused by the disease or condition, or stopping the symptoms of the disease or condition either prophylactically and/or therapeutically.

As used herein, the term "subject" or "patient" encompasses mammals and non-mammals. Examples of mammals include, but are not limited to, humans, chimpanzees, apes, monkeys, cattle, horses, sheep, goats, swine; rabbits, dogs, cats, rats, mice, guinea pigs, and the like. Examples of non-mammals include, but are not limited to, birds, fishes and the like.

As used herein, the term "administration" or "administering" of the subject compound refers to providing a compound of the invention and/or prodrugs thereof to a subject in need of treatment.

As used herein, the term "effective amount" or "therapeutically effective amount" refer to a sufficient amount of a compound described herein being administered which will relieve to some extent one or more of the symptoms of the disease or condition being treated. The result can be reduction and/or alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. For example, an "effective amount" for therapeutic uses is the amount of the composition comprising a compound as disclosed herein required to

provide a clinically significant decrease in disease symptoms. An appropriate “effective” amount in any individual case may be determined using techniques, such as a dose escalation study. By way of example only, a therapeutically effective amount of a compound of the invention may be in the range of e.g., about 0.01 mg/kg/day to about 1000 mg/kg/day, from
5 about 0.1 mg/kg/day to about 500 mg/kg/day, from about 0.1 mg (x2)/kg/day to about 500 mg (x2)/kg/day.

The compounds of the present invention were screened against a kinase panel and inhibited the activity of at least one kinase on the panel. Examples of kinases include, but not limited to, ROCK1 and ROCK2.

10 The compounds described herein are inhibitors of ROCK kinase activity and have therapeutic benefit in the treatment of disorders associated with inappropriate kinase activity, in particular in the treatment and prevention of disease states mediated by kinases, including ROCK kinase. Therefore, the present invention provides methods of regulating and, in particular, inhibiting signal transduction cascades in which a kinase plays a role. The method generally
15 involves administering to a subject or contacting a cell expressing the kinase with an effective amount of a compound described herein, a salt, a diastereomer, an enantiomer, a racemate, a hydrate, a solvate, a prodrug, and/or a composition thereof, to regulate or inhibit the signal transduction cascade. The methods are also used to regulate and, in particular, inhibit downstream processes or cellular responses elicited by activation of the particular kinase signal
20 transduction cascade. The methods are also practiced in *in vitro* contexts or in *in vivo* contexts as a therapeutic approach towards the treatment or prevention of diseases characterized by, caused by, or associated with activation of the kinase-dependent signal transduction cascade.

For the therapeutic uses of compounds provided herein, including compounds of Formula 1, salts, diastereomers, enantiomers, racemates, hydrates, solvates, or prodrugs thereof, such
25 compounds are administered in therapeutically effective amounts either alone or as part of a pharmaceutical composition. Accordingly, provided herein are pharmaceutical compositions, which comprise at least one compound provided herein, including at least one compound of Formula 1, a pharmaceutically acceptable salt, a diastereomer, an enantiomer, a racemate, a hydrate, a solvate, or prodrug thereof, and one or more pharmaceutically acceptable carriers,
30 diluents, adjuvant or excipients.

In addition, such compounds and compositions are administered singly or in combination with one or more additional therapeutic agents. Non-limiting examples of the additional therapeutic agents may include immune checkpoint inhibitors and immunogenic cell death (ICD)-inducing chemotherapeutic agents. Non-limiting examples of the immune checkpoint inhibitor may include PD-1 inhibitors, PD-L1 inhibitors, and CTLA-4 inhibitors. Non-limiting examples of the immunogenic cell death (ICD)-inducing chemotherapeutic agent may include doxorubicin, idarubicin, mitoxantrone, tautomycin, calyculin A, salubrinal, oxaliplatin, bleomycin, and cyclophosphamide. The methods of administration of such compounds and compositions include, but are not limited to, intravenous administration, inhalation, oral administration, rectal administration, parenteral, intravitreal administration, subcutaneous administration, intramuscular administration, intranasal administration, dermal administration, topical administration, ophthalmic administration, buccal administration, tracheal administration, bronchial administration, sublingual administration or optic administration. Compounds provided herein are administered by way of known pharmaceutical formulations, including tablets, capsules or elixirs for oral administration, suppositories for rectal administration, sterile solutions or suspensions for parenteral or intramuscular administration, lotions, gels, ointments or creams for topical administration, and the like. In some embodiments, such pharmaceutical compositions are formulated as tablets, pills, capsules, a liquid, an inhalant, a nasal spray solution, a suppository, a solution, a gel, an emulsion, an ointment, eye drops or ear drops.

The therapeutically effective amount will vary depending on, among others, the disease indicated, the severity of the disease, the age and relative health of the subject, the potency of the compound administered, the mode of administration and the treatment desired. The required dosage will also vary depending on the mode of administration, the particular condition to be treated and the effect desired.

The compounds of Formula 1 are useful for inhibiting one or more protein kinases and for treating diseases and disorders that are mediated by the protein kinases, such as cancer, autoimmune diseases, fibrotic disorders, cardiovascular disease, and neurodegenerative diseases.

The term “biological sample,” as used herein, means a sample outside an animal and includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from an animal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or

extracts thereof Inhibition of kinase activity, particularly ROCK kinase activity, in a biological sample is useful for a variety of purposes known to one of skill in the art. Examples of such purposes include, but are not limited to, biological specimen storage and biological assays.

The term “ROCK-mediated disease” or “condition,” as used herein, means any disease or
5 other deleterious condition in which ROCK is known to play a role. ROCK participates in a variety of important physiological functions in the vasculature, including smooth muscle contraction; cell proliferation, such as, for example, vascular smooth muscle cell proliferation; and cell adhesion and migration (see Hu & Lee, *Expert Opin. Ther. Targets*, 9(4):715-36, 2005; Shimokawa & Takeshita, *Arterioscler. Thromb. Vase. Biol.* 25(9):1767- 75, 2005). ROCK
10 participates in inflammatory responses due to leukocyte migration, such as, for example, autoimmune disease and allergic reactions (see Wettschureck et al., *J. Mol. Med.* 80:629-38, 2002). Abnormal activation of the Rho/ROCK pathway has been observed in various disorders of the central nervous system (see Mueller et al, *Nature Rev.*, 4:387-98, 2005). In addition, ROCK has been implicated in tumor cell migration and invasion (Riento & Ridley, *Nature Rev.*
15 4:446-56, 2004) and in osteoporosis (Ohnaka et al., *Biochem. Biophys., Res. Commun.* 287(2):337-4, 2001).

Specifically, the present invention relates to a method of treating or lessening the severity of a cardiovascular disease or disorder, such as, for example, cerebral vasospasm, hypertension, atherosclerosis, angina, myocardial infarction, ischemic/reperfusion injury, stroke, bronchial
20 asthma; glaucoma, pre-term labor, erectile dysfunction, or renal disease, such as, for example, chronic renal failure, chronic nephritis, diabetic nephropathy, and IgA nephropathy; a neurological disease or disorder, such as for example, spinal-cord injury, Alzheimer’s disease, multiple sclerosis, or neuropathic pain; and proliferative disorders, such as, for example, retinopathy, fibrosis, or invasive/metastatic cancers. Such cancers include adenocarcinoma,
25 adrenocortical cancer; bladder cancer; bone cancer; brain cancer; breast cancer; cancer of the buccal cavity; cervical cancer; colon cancer; colorectal cancer; endometrial or uterine carcinoma; epidermoid carcinoma; esophageal cancer; eye cancer; follicular carcinoma; gallbladder cancer; gastrointestinal cancer; cancer of the genitourinary tract; glioblastoma; hairy cell carcinoma; head and neck cancer; hepatic carcinoma; hepatocellular cancer; Hodgkin’s disease;
30 keratoacanthoma; kidney cancer; large cell carcinoma; cancer of the large intestine; laryngeal cancer; liver cancer; lung cancer, such as, for example, adenocarcinoma of the lung, small-cell

lung cancer, squamous carcinoma of the lung, non-small cell lung cancer; melanoma; myeloproliferative disorders, neuroblastoma; ovarian cancer; papillary carcinoma; pancreatic cancer; cancer of the peritoneum; prostate cancer; rectal cancer; salivary gland carcinoma; sarcoma; squamous cell cancer; small cell carcinoma; cancer of the small intestine; stomach
5 cancer; testicular cancer; thyroid cancer; and vulval cancer. In particular embodiments, the treated cancer is melanoma, breast cancer, colon cancer, or pancreatic cancer.

In another aspect, the invention provides a method of treating a fibrotic disorder in a subject comprising administering to the subject a therapeutically effective amount of a compound of Formula I. Non-limiting examples of fibrotic disorders are pulmonary fibrosis
10 including cystic and idiopathic pulmonary fibrosis, radiation induced lung injury, liver fibrosis including cirrhosis, cardiac fibrosis including arterial fibrosis, endomyocardial fibrosis, old myocardial infarction, arterial stiffness, atherosclerosis, restenosis, arthrofibrosis, Crohn's disease, myelofibrosis, Peyronie's diseases, nephrogenic systemic fibrosis, progressive massive fibrosis, retroperitoneal cavity fibrosis, scleroderma/systemic sclerosis, mediastinal fibrosis,
15 Keloids and hypertrophic scars, glial scarring, or renal fibrosis.

In one aspect, the present invention provides methods for treating a cell-proliferative disease or condition, such as cancer, comprising administering to a subject in need of such treatment a therapeutically effective amount of the compound of Formula 1, a pharmaceutically acceptable salt, a diastereomer, an enantiomer, a racemate, a hydrate, a solvate, a prodrug
20 thereof, or a pharmaceutical composition or medicament thereof, wherein the cell proliferative disease or condition include, for example, lymphoma, osteosarcoma, melanoma, breast cancer, renal cancer, prostate cancer, colorectal cancer, thyroid cancer, ovarian cancer, pancreatic cancer, neuronal cancer, lung cancer, uterine cancer or gastrointestinal cancer. In one aspect, the present invention provides methods of inhibiting growth of cancer cells with the compound
25 described herein, salt, diastereomer, enantiomer, racemate, hydrate, solvate, prodrug, or any combination thereof, or the composition described herein.

In certain embodiments, protein kinase-mediated diseases or conditions are inflammatory diseases or conditions, respiratory diseases or autoimmune diseases or conditions, such as asthma, chronic obstructive pulmonary disease (COPD), adult respiratory distress syndrome
30 (ARDS), ulcerative colitis, Crohn's disease, bronchitis, dermatitis, allergic rhinitis, psoriasis,

scleroderma, urticaria, rheumatoid arthritis, multiple sclerosis, cancer, breast cancer, HIV associated diseases or lupus.

In another aspect, the present invention provides methods for treating a neurological/neurodegenerative disease or condition by administering to a subject a therapeutically effective amount of the compound described herein, salt, diastereomer, enantiomer, racemate, hydrate, solvate, prodrug, or any combination thereof, or the composition described herein. In certain embodiment, such neurological/neurodegenerative disease or condition includes, for example, Alzheimer's disease, cerebral edema, cerebral ischemia, multiple sclerosis, cerebral cavernous malformations (CCMs), neuropathies, Parkinson's disease, blunt or surgical trauma (including post-surgical cognitive dysfunction and spinal cord or brain stem injury), as well as the neurological aspects of disorders such as degenerative disc disease and sciatica.

In another aspect, the present invention provides methods for treating a cardiovascular disease by administering to a subject a therapeutically effective amount of the compound described herein, salt, diastereomer, enantiomer, racemate, hydrate, solvate, prodrug, or any combination thereof, or the composition described herein. Such a cardiovascular disease affects the heart or blood vessels and includes, for example, atherosclerosis, arrhythmia, angina, myocardial ischemia, myocardial infarction, cardiac or vascular aneurysm, vasculitis, stroke, peripheral obstructive arteriopathy of a limb, an organ, or a tissue, reperfusion injury following ischemia of an organ or a tissue, endotoxic, surgical, or traumatic shock, hypertension, valvular heart disease, heart failure, abnormal blood pressure, vasoconstriction, vascular abnormality, or inflammation.

In another aspect, the present invention provides methods of treating cancer comprising administering to a subject in need a composition comprising a therapeutically effective amount of at least one of the compounds described herein, salts, diastereomers, enantiomers, racemates, hydrates, solvates, or prodrugs thereof and a therapeutically effective amount of at least one immune checkpoint inhibitor, wherein the cancer is adenocarcinoma, adrenocortical cancer, bladder cancer, bone cancer, brain cancer, breast cancer, buccal cavity cancer, cervical cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, epidermoid carcinoma, esophageal cancer, eye cancer, follicular carcinoma, gallbladder cancer, gastrointestinal cancer,

genitourinary tract cancer, glioblastoma, hairy cell carcinoma, head and neck cancer, hepatic carcinoma, hepatocellular cancer, Hodgkin's disease, keratoacanthoma, kidney cancer, large cell carcinoma, large intestine cancer, laryngeal cancer, liver cancer, lung adenocarcinoma, small-cell lung cancer, lung squamous carcinoma, non-small cell lung cancer, melanoma, a
5 myeloproliferative disorder, neuroblastoma, ovarian cancer, papillary carcinoma, pancreatic cancer, peritoneal cancer, prostate cancer, rectal cancer, salivary gland carcinoma, sarcoma, squamous cell cancer, small cell carcinoma, small intestine cancer, stomach cancer, testicular cancer, thyroid cancer, vulvar cancer, or any combination thereof. Examples of the checkpoint inhibitor include, not being limited to, a PD-1 inhibitor, a PD-L1 inhibitor, and a CTLA-4
10 inhibitor.

In another aspect, the present invention provides methods of treating a cancer comprising administering to a subject in need a composition comprising a therapeutically effective amount of at least one of the compounds described herein, salts, diastereomers, enantiomers, racemates, hydrates, solvates, or prodrugs thereof and a therapeutically effective amount of at least one
15 immunogenic cell death (ICD)-inducing chemotherapeutic, wherein the cancer is adenocarcinoma, adrenocortical cancer, bladder cancer, bone cancer, brain cancer, breast cancer, buccal cavity cancer, cervical cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, epidermoid carcinoma, esophageal cancer, eye cancer, follicular carcinoma, gallbladder cancer, gastrointestinal cancer, genitourinary tract cancer, glioblastoma, hairy cell
20 carcinoma, head and neck cancer, hepatic carcinoma, hepatocellular cancer, Hodgkin's disease, keratoacanthoma, kidney cancer, large cell carcinoma, large intestine cancer, laryngeal cancer, liver cancer, lung adenocarcinoma, small-cell lung cancer, lung squamous carcinoma, non-small cell lung cancer, melanoma, a myeloproliferative disorder, neuroblastoma, ovarian cancer, papillary carcinoma, pancreatic cancer, peritoneal cancer, prostate cancer, rectal cancer, salivary
25 gland carcinoma, sarcoma, squamous cell cancer, small cell carcinoma, small intestine cancer, stomach cancer, testicular cancer, thyroid cancer, vulvar cancer, or any combination thereof. Examples of the immunogenic cell death (ICD)-inducing chemotherapeutic include, not being limited to, doxorubicin, idarubicin, mitoxantrone, tautomycin, calyculin A, salubrinal, oxaliplatin, bleomycin, and cyclophosphamide.

30 In the above methods for using the compound of the invention, the compound described herein, salt, diastereomer, enantiomer, racemate, hydrate, solvate, or prodrug thereof is

administered to a system comprising cells or tissues. In certain embodiments, the compound described herein, a salt, a diastereomer, an enantiomer, a racemate, a hydrate, a solvate, a prodrug, or any combination thereof is administered to a human or animal subject. In still certain embodiments, a pharmaceutical composition or a medicament comprising at least one of
5 the compound, the salt, the diastereomer, the enantiomer, the racemate, the hydrate, the solvate, the prodrug, or any combination thereof is administered to a human or animal subject.


4. Methods of Preparing Compounds

In a further aspect, the present invention provides methods of preparing the compounds, salts, diastereomers, enantiomers, racemates, hydrates, solvates, and prodrugs. In some
10 embodiments, the compounds, salts, diastereomers, enantiomers, racemates, hydrates, solvates, or prodrugs may be prepared by methods including, but not limited to, one or more of the following methods:

- (a) optionally converting a compound of the invention into a pharmaceutically acceptable salt;
- 15 (b) optionally converting a salt form of a compound of the invention to a non-salt form;
- (c) optionally converting an unoxidized form of a compound of the invention into a pharmaceutically acceptable *N*-oxide;
- (d) optionally resolving an individual isomer of a compound of the invention from a mixture of stereoisomers;
- 20 (e) optionally converting a non-derivatized compound of the invention into a pharmaceutically acceptable prodrug derivative; and
- (f) optionally converting a prodrug derivative of a compound of the invention to its non-derivatized form.

Exemplary methods for preparing the compounds of the invention are described herein,
25 including in the Examples, which will be described in detail below. Some embodiments of the invention provide processes for preparing the compounds of the present invention, as illustrated in Method 1 or Method 2 below.

Method 1




- A bicyclic acid (2) was reacted with thionyl chloride or oxalyl chloride in dichloromethane in the presence of catalytic amount of GMF to provide its acyl chloride (3). The resulting acyl chloride (3) was coupled with a bicyclic heteroaryl compound (4) in a Friedel-Craft acylation reaction to provide ketone (5) in an organic solvent (dichloromethane or dichloroethane) in the presence of a Lewis acid (aluminum chloride, titanium chloride or stannous chloride). The resulted ketone (5) was reacted with an alkyl bromide, alkyl chloride or alkyl epoxide to afford a *N*-alkylated compound (6) in an organic solvent (DMF, acetone, dioxane, acetonitrile, dichloromethane or dichloroethane) with an inorganic base (NaHCO₃, Na₂CO₃, K₂CO₃, Cs₂CO₃, K₂HPO₄ or K₃PO₄) at room temperature (RT)-150 °C for 5-24 hours.

The *N*-alkylated compound (6) also was alternatively prepared starting from a bicyclic heteroaryl compound (4) by *N*-alkylation at the above mentioned reaction condition to give (7), followed by Friedel-Craft acylation reaction with an acyl chloride (3).

- The *N*-alkylated compound (6) was reacted with a boronic acid or a boronic ester with Suzuki reaction condition using a palladium catalyst (Pd(OAc)₂, Pd₂(dba)₃, Pd(PPh₃)₄ or Pd(dppf)Cl₂·DCM) to provide the compound (1) in the presence of an inorganic base (NaHCO₃, Na₂CO₃, K₂CO₃, Cs₂CO₃, CsF, K₂HPO₄ or K₃PO₄) at room temperature (RT)-150 °C for 5-24 hours.

The *N*-alkylated compound (6) also was alternatively prepared starting from a bicyclic as shown below in Method 2.

Method 2



A bicyclic heteroaryl compound (4) was reacted with *N*-iodosuccinimide (NIS) in an organic solvent (DMF, acetone, dioxane, acetonitrile, dichloromethane or dichloroethane) at RT for 2-24h to provide an iodobicyclic heteroaryl compound (8). The iodobicyclic heteroaryl compound (8) was reacted with an alkyl bromide, alkyl chloride or alkyl epoxide to afford a *N*-alkylated compound (10) in an organic solvent (DMF, acetone, dioxane, acetonitrile, dichloromethane or dichloroethane) with an inorganic base (NaHCO₃, Na₂CO₃, K₂CO₃, Cs₂CO₃, K₂HPO₄ or K₃PO₄) at room temperature (RT)-150 °C for 5-24 hours. A *N*-alkylated compound (10) was alternatively prepared from bicyclic heteroaryl compound (4) by N-alkylation to (9), followed by iodination of (9) using *N*-iodosuccinimide (NIS) as aforementioned. A bicyclic acid (2) was coupled with *N*-methoxymethanamine hydrochloride by a coupling reagent (DCC, EDCI, HATU, HBTU, PyBop or PyBrop) in an organic solvent (DMF, tetrahydrofuran, dichloromethane or dichloroethane) at RT for 2-24h to provide a Weinreb amide (11). The Weinreb amide (11) was converted to a ketone (6) by reacting with iodobicyclic heteroaryl compound (10) in the presence of Grignard reagent at low temperature (-78-0 °C) in THF.

Non-limiting examples of the compounds described herein may include:

(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(3-methoxy-1H-pyrazol-4-yl)pyrrolo[3,2-*c*]pyridin-3-yl]methanone;

- (6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(3-methoxy-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]methanone dihydrochloride;
- (6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(3-methoxy-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]methanone dimethanesulfonic acid;
- 5 (6-(5-chloro-1H-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1H-pyrrolo[3,2-c]pyridin-3-yl)((S)-6-methoxychroman-3-yl)methanone;
- (6-(5-chloro-1H-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1H-pyrrolo[3,2-c]pyridin-3-yl)((S)-6-methoxychroman-3-yl)methanone dihydrochloride;
- (6-(5-chloro-1H-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1H-pyrrolo[3,2-c]pyridin-3-yl)((S)-6-methoxychroman-3-yl)methanone dihydrobromide;
- 10 (6-(5-chloro-1H-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1H-pyrrolo[3,2-c]pyridin-3-yl)((S)-6-methoxychroman-3-yl)methanone dimethanesulfonic acid;
- (6-Chlorochroman-3-yl)-[6-(3-chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]pyrrolo[2,3-b]pyridin-3-yl]methanone;
- 15 [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]pyrrolo[2,3-b]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone;
- (6-Chlorochroman-3-yl)-[6-(3-chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]pyrrolo[3,2-c]pyridin-3-yl]methanone;
- (6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]pyrrolo[3,2-c]pyridin-3-yl)-(6-fluorochroman-3-yl)methanone;
- 20 [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]pyrrolo[2,3-b]pyridin-3-yl]-(6-methoxychroman-3-yl)methanone;
- [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]pyrrolo[3,2-c]pyridin-3-yl]-(6-methoxychroman-3-yl)methanone;
- 25 [1-(Azetidin-3-yl)-6-(3-chloro-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone;

(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)pyrrolo[2,3-b]pyridin-3-yl]methanone;

(6-Chlorochroman-3-yl)-[6-(3-chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]methanone;

5 [1-(Azetidin-3-yl)-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

(6-Chlorochroman-3-yl)-[6-(5-methoxy-1H-pyrazol-4-yl)-1-[(1-methylpyrrolidin-2-yl)methyl]indol-3-yl]methanone;

10 (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[(1-methylpyrrolidin-2-yl)methyl]indol-3-yl]methanone;

[1-(Azetidin-3-ylmethyl)-6-(5-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

[1-(Azetidin-3-ylmethyl)-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

15 [1-[2-(Dimethylamino)ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;

(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]methanone;

20 (S)-(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]methanone;

(R)-(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]methanone;

(6-Chlorochroman-3-yl)-[6-(5-fluoro-1H-pyrazol-4-yl)-1-[(1-methylpyrrolidin-2-yl)methyl]indol-3-yl]methanone;

25 (6-Chlorochroman-3-yl)-[6-(5-methoxy-1H-pyrazol-4-yl)-1-[(1R)-1-methylpyrrolidin-2-yl]methyl]indol-3-yl]methanone;

- (6-Chlorochroman-3-yl)(6-(5-methoxy-1H-pyrazol-4-yl)-1-(((S)-1-methylpyrrolidin-2-yl)methyl)-1H-indol-3-yl)methanone;
- [1-(Azetidin-2-ylmethyl)-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- 5 [1-(Azetidin-2-ylmethyl)-6-(5-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- [1-(Azetidin-2-ylmethyl)-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- 10 (6-Chlorochroman-3-yl)-[2-(5-chloro-1H-pyrazol-4-yl)-7-[2-(dimethylamino)ethyl]pyrrolo[2,3-d]pyrimidin-5-yl]methanone;
- (6-Chlorochroman-3-yl)-[7-[2-(dimethylamino)ethyl]-2-(5-fluoro-1H-pyrazol-4-yl)pyrrolo[2,3-d]pyrimidin-5-yl]methanone;
- (1-(((R)-Azetidin-2-yl)methyl)-6-(5-chloro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;
- 15 (1-(((S)-Azetidin-2-yl)methyl)-6-(5-chloro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;
- [1-[[2S]-Azetidin-2-yl]methyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-methoxychroman-3-yl)methanone;
- [1-(Azetidin-2-ylmethyl)-6-(5-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- 20 [1-[[2R]-Azetidin-2-yl]methyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- (1-(((S)-Azetidin-2-yl)methyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-chlorochroman-3-yl)methanone;
- 25 (1-(((R)-Azetidin-2-yl)methyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;

[1-(2-Aminoethyl)-6-(3-chloro-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]-(6-chlorochroman-3-yl)methanone;

[1-(2-Aminopropyl)-6-(3-chloro-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone;

5 [1-(2-Aminoethyl)-6-(3-chloro-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone;

[1-(2-Aminoethyl)-6-(3-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;

10 [1-[(2R)-2-Aminopropyl]-6-(3-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

[1-[(2R)-2-Aminopropyl]-6-(3-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

[1-[(2S)-2-Aminopropyl]-6-(3-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

15 [1-[(2S)-2-Aminopropyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

[1-[(2S)-2-Aminopropyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

20 [1-[(2S)-2-Aminobutyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

(6-(5-chloro-1H-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1H-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone;

[1-[(2R)-2-Aminobutyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

25 (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[(2S)-2-(dimethylamino)propyl]indol-3-yl]methanone;

- (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[[1-(dimethylamino)cyclopropyl]methyl]indol-3-yl]methanone;
- (6-Chlorochroman-3-yl)-[1-[(2R)-2-(dimethylamino)propyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]methanone;
- 5 (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[(2R)-2-(dimethylamino)propyl]indol-3-yl]methanone;
- (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[(2R)-2-(dimethylamino)propyl]indol-3-yl]methanone;
- 10 (6-Chlorochroman-3-yl)-[6-(5-fluoro-1H-pyrazol-4-yl)-1-[(1-methylazetidin-3-yl)methyl]indol-3-yl]methanone;
- (6-Chlorochroman-3-yl)-[6-(5-methoxy-1H-pyrazol-4-yl)-1-[(1-methylazetidin-3-yl)methyl]indol-3-yl]methanone;
- [6-(5-Fluoro-1H-pyrazol-4-yl)-1-[[2R)-1-methylazetidin-2-yl]methyl]indol-3-yl)-(6-methoxychroman-3-yl)methanone;
- 15 [6-(5-Chloro-1H-pyrazol-4-yl)-1-[[1S)-1-methylazetidin-2-yl]methyl]indol-3-yl)-(6-methoxychroman-3-yl)methanone;
- (6-Methoxychroman-3-yl)-[6-(5-methoxy-1H-pyrazol-4-yl)-1-[[1S)-1-methylazetidin-2-yl]methyl]indol-3-yl]methanone;
- [6-(5-Fluoro-1H-pyrazol-4-yl)-1-[[2S)-1-methylazetidin-2-yl]methyl]indol-3-yl)-(6-methoxychroman-3-yl)methanone;
- 20 (6-Methoxychroman-3-yl)-[6-(5-methoxy-1H-pyrazol-4-yl)-1-[[1R)-1-methylazetidin-2-yl]methyl]indol-3-yl]methanone;
- [6-(5-Chloro-1H-pyrazol-4-yl)-1-[[1R)-1-methylazetidin-2-yl]methyl]indol-3-yl)-(6-methoxychroman-3-yl)methanone;
- 25 (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[[1R)-1-methylazetidin-2-yl]methyl]indol-3-yl]methanone;

- [1-[[[(1R)-1-Ethylazetidin-2-yl]methyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-methoxychroman-3-yl)methanone;
- (6-Chlorochroman-3-yl)-[6-(5-fluoro-1H-pyrazol-4-yl)-1-[[[(1R)-1-methylazetidin-2-yl]methyl]indol-3-yl]methanone;
- 5 [6-(5-Chloro-1H-pyrazol-4-yl)-1-[[[(1R)-1-ethylazetidin-2-yl]methyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;
- [1-[2-(Azetidin-1-yl)ethyl]-6-(3-chloro-1H-pyrazol-4-yl)pyrrolo[2,3-b]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone;
- [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(methylamino)ethyl]pyrrolo[3,2-c]pyridin-3-yl]-(6-
10 fluorochroman-3-yl)methanone;
- [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(methylamino)ethyl]pyrrolo[2,3-b]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone;
- [1-[2-(Azetidin-1-yl)ethyl]-6-(3-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- 15 [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(methylamino)ethyl]indol-3-yl]-(6-fluorochroman-3-yl)methanone;
- (6-Chlorochroman-3-yl)-[1-[3-(dimethylamino)propyl]-6-(3-methoxy-1H-pyrazol-4-yl)indol-3-yl]methanone;
- (S)-(6-Chlorochroman-3-yl)(1-(3-(dimethylamino)propyl)-6-(3-methoxy-1H-pyrazol-4-
20 yl)-1H-indol-3-yl)methanone;
- (R)-(6-Chlorochroman-3-yl)(1-(3-(dimethylamino)propyl)-6-(3-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone;
- (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[2-(methylamino)ethyl]indol-3-yl]methanone;
- 25 (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[2-[ethyl(methyl)amino]ethyl]indol-3-yl]methanone;

(6-Chlorochroman-3-yl)-[1-[2-[ethyl(methyl)amino]ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]methanone;

(6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[2-(diethylamino)ethyl]indol-3-yl]methanone;

5 (6-Chlorochroman-3-yl)-[1-[2-(diethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]methanone;

[6-(5-Chloro-1H-pyrazol-4-yl)-1-[2-(diethylamino)ethyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;

10 [1-[2-(Diethylamino)ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-methoxychroman-3-yl)methanone;

[6-(5-Chloro-1H-pyrazol-4-yl)-1-[2-(methylamino)ethyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;

[6-(5-Fluoro-1H-pyrazol-4-yl)-1-[2-(methylamino)ethyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;

15 [6-(5-Chloro-1H-pyrazol-4-yl)-1-[2-[ethyl(methyl)amino]ethyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;

[1-[2-[Ethyl(methyl)amino]ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-methoxychroman-3-yl)methanone;

20 [1-[(1-Aminocyclopropyl)methyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

[1-[(1-Aminocyclopropyl)methyl]-6-(5-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

(1-((1-Aminocyclopropyl)methyl)-6-(5-chloro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;

25 (1-((1-Aminocyclopropyl)methyl)-6-(5-fluoro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-chlorochroman-3-yl)methanone;

- (1-((1-Aminocyclopropyl)methyl)-6-(5-fluoro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-fluorochroman-3-yl)methanone;
- (1-((1-Aminocyclopropyl)methyl)-6-(5-fluoro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;
- 5 2,3-Dihydro-1,4-benzodioxin-3-yl-[1-[2-(dimethylamino)ethyl]-6-(1H-pyrazol-4-yl)indol-3-yl]methanone;
- Chroman-3-yl-[1-[2-(dimethylamino)ethyl]-6-(1H-pyrazol-4-yl)indol-3-yl]methanone
- [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;
- 10 (S)-(6-(3-Chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;
- (S)-(6-(3-Chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dihydrochloride;
- (R)-(6-(3-Chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-
- 15 methoxychroman-3-yl)methanone;
- [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(6-fluoro-2,3-dihydro-1,4-benzodioxin-3-yl)methanone;
- (6-Chloro-2,3-dihydro-1,4-benzodioxin-3-yl)-[6-(3-chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]methanone;
- 20 [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(2,3-dihydro-1,4-benzodioxin-3-yl)methanone;
- [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(6-fluorochroman-3-yl)methanone;
- (S)-[6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(6-
- 25 fluorochroman-3-yl)methanone;
- (R)-[6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(6-fluorochroman-3-yl)methanone dimethanesulfonic acid;

(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(3-methylisoxazol-4-yl)indol-3-yl]methanone;

(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(2-methylpyrazol-3-yl)indol-3-yl]methanone;

5 [6-(2-Amino-4-pyridyl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(6-chlorochroman-3-yl)methanone;

[1-[2-(Dimethylamino)ethyl]-6-(3-methylisoxazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;

10 [1-[2-(Dimethylamino)ethyl]-6-(2-methylpyrazol-3-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;

(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-methoxy-1*H*-pyrazol-4-yl)indol-3-yl]methanone;

(*R*)-(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-methoxy-1*H*-pyrazol-4-yl)indol-3-yl]methanone;

15 (*S*)-(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-methoxy-1*H*-pyrazol-4-yl)indol-3-yl]methanone;

(*R*)-(6-(5-Chloro-1*H*-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1*H*-indol-3-yl)(6-chlorochroman-3-yl)methanone;

20 (*S*)-(6-(5-Chloro-1*H*-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1*H*-indol-3-yl)(6-chlorochroman-3-yl)methanone;

[1-[2-(Dimethylamino)ethyl]-6-(5-methoxy-1*H*-pyrazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;

(*S*)-[1-[2-(Dimethylamino)ethyl]-6-(5-methoxy-1*H*-pyrazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;

25 (*S*)-[1-[2-(Dimethylamino)ethyl]-6-(5-methoxy-1*H*-pyrazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone dihydrochloride;

- (R)-[1-[2-(Dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;
- [1-[2-(Dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-methoxychroman-3-yl)methanone;
- 5 (6-Chloro-2,3-dihydro-1,4-benzodioxin-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]methanone;
- (2,3-Dihydrobenzo[b][1,4]dioxin-2-yl)(1-(2-(dimethylamino)ethyl)-6-(1H-pyrazol-4-yl)-1H-indazol-3-yl)methanone;
- (S)-(2,3-Dihydrobenzo[b][1,4]dioxin-2-yl)(1-(2-(dimethylamino)ethyl)-6-(1H-pyrazol-4-yl)-1H-indazol-3-yl)methanone;
- 10 (6-Chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-methoxy-1H-pyrazol-4-yl)-1H-indazol-3-yl)methanone;
- (S)-(6-Chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-methoxy-1H-pyrazol-4-yl)-1H-indazol-3-yl)methanone;
- 15 (R)-(6-Chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-methoxy-1H-pyrazol-4-yl)-1H-indazol-3-yl)methanone;
- (6-Chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-fluoro-1H-pyrazol-4-yl)-1H-indazol-3-yl)methanone;
- (1-(2-(Dimethylamino)ethyl)-6-(3-methoxy-1H-pyrazol-4-yl)-1H-indazol-3-yl)(6-methoxychroman-3-yl)methanone;
- 20 (6-(3-Chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indazol-3-yl)(6-methoxychroman-3-yl)methanone;
- (1-(2-(Dimethylamino)ethyl)-6-(3-methoxy-1H-pyrazol-4-yl)-1H-indazol-3-yl)(6-fluorochroman-3-yl)methanone;
- 25 (6-(3-Chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indazol-3-yl)(6-fluorochroman-3-yl)methanone;

- (S)-(6-(3-Chloro-1*H*-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1*H*-indazol-3-yl)(6-fluorochroman-3-yl)methanone;
- Chroman-3-yl(1-(2-(dimethylamino)ethyl)-6-(5-(trifluoromethyl)-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)methanone;
- 5 (6-(3-Fluoro-1*H*-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1*H*-indazol-3-yl)(6-fluorochroman-3-yl)methanone;
- (6-(3-Fluoro-1*H*-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1*H*-indazol-3-yl)((S)-6-fluorochroman-3-yl)methanone;
- (1-(2-(Dimethylamino)ethyl)-6-(3-fluoro-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)(6-fluorochroman-3-yl)methanone;
- 10 (7-Chloro-2,3-dihydrobenzo[*b*][1,4]dioxin-2-yl)(1-(2-(dimethylamino)ethyl)-6-(3-methoxy-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)methanone;
- (7-Chloro-2,3-dihydrobenzo[*b*][1,4]dioxin-2-yl)(1-(2-(dimethylamino)ethyl)-6-(3-fluoro-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)methanone;
- 15 (6-(3-Chloro-1*H*-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1*H*-indazol-3-yl)(7-chloro-2,3-dihydrobenzo[*b*][1,4]dioxin-2-yl)methanone;
- (1-(2-(Dimethylamino)ethyl)-6-(3-methoxy-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)(7-fluoro-2,3,4,5-tetrahydrobenzo[*b*]oxepin-4-yl)methanone;
- (6-(3-Chloro-1*H*-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1*H*-indazol-3-yl)(7-fluoro-2,3,4,5-tetrahydrobenzo[*b*]oxepin-4-yl)methanone;
- 20 (1-(2-(Dimethylamino)ethyl)-6-(3-methoxy-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)(7-methoxy-2,3,4,5-tetrahydrobenzo[*b*]oxepin-4-yl)methanone;
- (6-(3-Chloro-1*H*-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1*H*-indazol-3-yl)(7-methoxy-2,3,4,5-tetrahydrobenzo[*b*]oxepin-4-yl)methanone;
- 25 (6-(5-chloro-1*H*-pyrazol-4-yl)-1-(((R)-1-methylazetidin-2-yl)methyl)-1*H*-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone dihydrochloride;

(6-(5-chloro-1H-pyrazol-4-yl)-1-(((R)-1-methylazetidin-2-yl)methyl)-1H-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone dihydrobromide;

(6-(5-chloro-1H-pyrazol-4-yl)-1-(((R)-1-methylazetidin-2-yl)methyl)-1H-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone dimethanesulfonic acid;

5 (6-(5-chloro-1H-pyrazol-4-yl)-1-(((R)-1-ethylazetidin-2-yl)methyl)-1H-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone;

(6-(5-chloro-1H-pyrazol-4-yl)-1-(((R)-1-ethylazetidin-2-yl)methyl)-1H-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone dihydrobromide;

10 (6-(5-chloro-1H-pyrazol-4-yl)-1-(((R)-1-ethylazetidin-2-yl)methyl)-1H-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone dimethanesulfonic acid;

((S)-6-chlorochroman-3-yl)(1-((R)-2-(dimethylamino)propyl)-6-(5-fluoro-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone;

((S)-6-chlorochroman-3-yl)(1-((R)-2-(dimethylamino)propyl)-6-(5-fluoro-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone hydrobromide;

15 (S)-(6-chlorochroman-3-yl)(1-(2-(diethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone;

(S)-(6-chlorochroman-3-yl)(1-(2-(diethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone dihydrochloride;

20 (S)-(6-chlorochroman-3-yl)(1-(2-(diethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone dimethanesulfonic acid;

(S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(diethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;

(S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(diethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dihydrochloride;

(S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(diethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dihydrobromide

(S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(diethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dimethanesulfonic acid;

5 (S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(ethyl(methyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone

(S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(ethyl(methyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dihydrochloride;

10 (S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(ethyl(methyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dihydrobromide;

(S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(ethyl(methyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dimethanesulfonic acid;

(S)-(6-(3-chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dihydrobromide;

15 (S)-(6-(3-chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dimethanesulfonic acid;

(6-(3-chloro-1H-pyrazol-4-yl)-1-(((R)-1-(2-fluoroethyl)azetidin-2-yl)methyl)-1H-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone;

20 (6-(3-chloro-1H-pyrazol-4-yl)-1-(((R)-1-(2,2-difluoroethyl)azetidin-2-yl)methyl)-1H-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone;

(6-(3-chloro-1H-pyrazol-4-yl)-1-(((R)-1-(2,2,2-trifluoroethyl)azetidin-2-yl)methyl)-1H-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone;

(S)-(6-(3-chloro-1H-pyrazol-4-yl)-1-(2-(ethyl(2-fluoroethyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;

(S)-(6-(3-chloro-1H-pyrazol-4-yl)-1-(2-((2,2-difluoroethyl)(ethyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;

(S)-(6-(3-chloro-1H-pyrazol-4-yl)-1-(2-(ethyl(2,2,2-trifluoroethyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;

5 (S)-(1-(2-(bis(2-fluoroethyl)amino)ethyl)-6-(3-chloro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;

(S)-(6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone hydrochloride;

10 (S)-(6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone dihydrobromide;

(S)-(6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone dimethanesulfonic acid;

(6-(3-fluoro-1H-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1H-indazol-3-yl)((S)-6-fluorochroman-3-yl)methanone dihydrobromide.

15

EXAMPLES

The present invention is further exemplified by the following examples that illustrate the preparation of compounds of Figure 1 according to the invention. The examples are for illustrative purpose only and are not intended, nor should they be construed as limiting the invention in any manner. Those skilled in the art will appreciate that variations and modifications can be made without changing the scope of the invention.

20 Nuclear magnetic resonance (NMR) and mass spectrometry (MS) spectra obtained for compounds described in the examples below and those described herein were consistent with that of the compounds of formulae herein.

25 Liquid chromatography-mass spectrometry (LC-MS) Method:


1. Samples are run on Agilent Technologies 6120 MSD system with a Zorbax Eclipse XDB-C18 (3.5 μ) reverse phase column (4.6 x 50 mm) run at room temperature with flow rate of 1.5 mL/minute.
2. The mobile phase uses solvent A (water/0.1 % formic acid) and solvent B (acetonitrile/0.1 % formic acid): 95 %/5 % to 0 %/100 % (A/B) for 5 minute.
3. The mass spectra (m/z) were recorded using electrospray ionization (ESI).
4. Ionization data was rounded to the nearest integer.

Proton NMR Spectra:

- Unless otherwise indicated, all ^1H NMR spectra are run on a Varian series Mercury 400 or 500 MHz. All observed protons are reported as parts-per-million (ppm) downfield from tetramethylsilane using conventional abbreviations for designation of major peaks: e.g., s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet) and brs (broad singlet).

Example 1: Preparation of Compounds


General Schem 1



15

Compound 1-1: (6-chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(3-methoxy-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]methanone (followed step 5a)

Scheme 1



Step 1: A mixture of 6-chlorochromane-3-carboxylic acid (6.0 g, 28.2 mmol), thionyl chloride (21.0 mL, 282.2 mmol), and DMF (0.1 mL, catalyst) in toluene (60 mL) was heated to reflux with stirring for 3 hours. The mixture was cooled to room temperature and concentrated to remove volatile organic solvents. The crude was dissolved in acetonitrile (100 mL) and concentrated to provide **1a** as a pale blown oil. The **1a** was used in the next step without further purification.

Step 2: A mixture of **1a** (5.4 g, 23 mmol) and 6-bromo-1H-pyrrolo[3,2-c]pyridine (3.7 g, 19 mmol) in dichloromethane (60 mL) was cooled in an ice bath, and aluminum chloride (9.4 g, 70 mmol) was added. After being stirred for 1 hour at 0 °C, the reaction was quenched with a saturated aqueous NH₄Cl solution (100 mL) at 0 °C to form solid precipitation. The mixture was then warmed to room temperature and stirred for an additional 2 hours. The solids were collected by filtration and then rinsed with methanol. The resulting solids were dried under high vacuum to provide **1b** as a white solid in 80% yield (7.4 g). LC/MS found 391.0 and 393.0 [M+H]⁺.

Step 3: A mixture of **1b** (800 mg, 2.1 mmol), potassium carbonate (1.76 g, 1.3 mmol, 770 μL), and 2-chloro-N,N-dimethylethanamine hydrochloride (690 mg, 6.4 mmol) in DMF (20 mL) was heated to 70 °C for 16 hours. The mixture was poured over water (100 mL) and extracted with EtOAc (100 mL). The organic layer was separated, washed with brine (100 mL),

dried over Na₂SO₄, filtered, and concentrated. The residue was purified by flash column chromatography on silica gel column using a gradient of 20-100% EtOAc in hexane to give **1c** as a white solid in 20 % yield (202 mg). LC/MS found 463.1 and 465.0 [M+H]⁺.

Step 4: A mixture of Intermediate **1c** (202 mg, 43 μmol), [3-methoxy-1-(p-tolylsulfonyl)pyrazol-4-yl]boronic acid (190 mg, 65 μmol), potassium carbonate (2 M in water, 54 μL), and tetrakis(triphenylphosphine)palladium (25.0 mg, 22 μmol) in 1,4-dioxane (10 mL) was degassed via sparging with argon for 5 minutes, and the resulting mixture was heated to 70 °C and shaken overnight. The mixture was cooled to room temperature, concentrated, poured over water (50 mL), and extracted twice with EtOAc (50 mL). The combined organic layers were dried over Na₂SO₄, filtered, and concentrated to afford **1d**, and the residue was used for the next step without further purification.

Step 5a for a tosyl (Ts) protected pyrazole compound A solution of **1d** (crude, 230 mg) in THF (1 mL) was cooled on an ice bath and treated with 2N sodium hydroxide solution (2.0 mL). The solution was stirred at 0 °C for 30 minutes, and it was neutralized with 1N HCl solution (4.0 mL), diluted with EtOAc (30 mL) and saturated Na₂CO₃ solution (30 mL). The aqueous phase was separated and extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered, treated with silica gel, and evaporated under reduced pressure. The material was purified by silica gel column chromatography to provide a compound **1-1** (6-chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(3-methoxy-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]methanone (101 mg, 210 μmol, 58% yield for step 4 and 5a) as a white solid. LC/MS found 480.2 [M+H]⁺.

Compound 1-2: ((6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-pyrrolo[3,2-c]pyridin-3-yl)methanone dihydrochloride)













Step 6a: To a solution of **1-1** (50 mg, 104 μmol) in acetone (1 mL), was slowly added a 4M solution of HCl in dioxane (55 μL) at room temperature to form solid precipitation. After being stirred for additional 1 hour at room temperature, was added ter-butylethylether (2 mL) for a complete precipitation. The solids were collected by filtration under nitrogen atmosphere, rinsed with ter-butylethylether, and dried under vacuum to give **1-2** (47.5 mg, 82%) as white powder. LC/MS found 480.2 [M+H]⁺.






















Compound 1-3: ((6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-pyrrolo[3,2-c]pyridin-3-yl)methanone dimethanesulfonic acid)

Step 6a: To a solution of **1-1** (50 mg, 104 μmol) in acetone (1 mL), was slowly added a 1M solution of methane sulfonic acid in acetone (220 μL) at room temperature to form solid precipitation. After being stirred for additional 1 hour at room temperature, was added ter-butylethylether (2 mL) for a complete precipitation. The resulting solids were collected by filtration under nitrogen atmosphere, rinsed with ter-butylethylether and then dried under vacuum to give **1-3** (60.3 mg, 90%) as white powder. LC/MS found 480.2 $[\text{M}+\text{H}]^+$.

































The following compounds **1-4** to **1-51** shown in Table 1 were prepared by the method (General Scheme 1) similar to that described for the preparation of compound **1-1**, **1-2** or **1-3** using appropriate (4) with an appropriate boronic acid/ester (R^3) and appropriate R^1R^2 substituent.

Table 1





































Compd	R^1R^2 Substituent	R^3 boronic acid	(4)	Structure	$[\text{M}+\text{H}]^+$
1-2					480.2
1-3					480.2
1-4					492.3
1-5					492.3































1-6				 2HBr	492.3
1-7				 2MeSO ₃ H	492.3
1-8					480.2
1-9					484.2
1-10				 ¹ H NMR (500 MHz, CHLOROFORM- <i>d</i>) δ ppm 10.17 - 10.46 (m, 1 H) 8.42 (d, <i>J</i> =8.2 Hz, 1 H) 8.04 (s, 1 H) 7.77 (s, 1 H) 7.68 (s, 1 H) 7.41 - 7.49 (m, 1 H) 6.77 - 6.84 (m, 1 H) 6.69 - 6.75 (m, 1 H) 6.67 (d, <i>J</i> =2.9 Hz, 1 H) 4.48 - 4.54 (m, 1 H) 4.32 (br s, 1 H) 4.06 - 4.16 (m, 1 H) 3.69 - 3.81 (m, 4 H) 3.36 (dd, <i>J</i> =16.3, 11.3 Hz, 1 H) 2.91 (ddd, <i>J</i> =16.5, 4.8, 1.8 Hz, 2 H) 2.06 - 2.21 (m, 4 H)	468.2
1-11					484.2
1-12					468.2





























1-13					480.2
1-14					480.2
1-15					452.1
1-16					468.2
1-17					483.2
1-18					463.2
1-19					505.3
1-20					509.2

1-21					481.2
1-22					477.3
1-23					451.2
1-24					467.2
1-25					467.2
1-26					467.2
1-27					493.2
1-28					505.3


1-29					505.3
1-30					466.2
1-31					481.2
1-32					477.3
1-33					485.1
1-34					469.2
1-35					477.2
1-36					477.2
1-37					461.2

1-38					461.2
1-39					481.1
1-40					465.2
1-41					473.3
1-42					473.3
1-43					479.2
1-44					491.2
1-45					475.2
1-46				 ¹ H NMR (500 MHz, CHLOROFORM- <i>d</i>) δ ppm 8.42 (d, <i>J</i> =8.2 Hz, 1 H) 8.30 - 8.38 (m, 1 H) 8.02 (d, <i>J</i> =4.3 Hz, 1 H) 7.76 (s, 1 H) 7.68 (s, 1 H) 7.44 (dd, <i>J</i> =8.4, 1.4 Hz, 1 H) 6.81 (dd, <i>J</i> =8.9, 1.8 Hz, 1 H) 6.70 - 6.75 (m, 1 H) 6.67 (dd, <i>J</i> =7.2,	491.2

				2.9 Hz, 1 H) 4.48 - 4.56 (m, 1 H) 4.24 - 4.30 (m, 1 H) 4.05 - 4.16 (m, 1 H) 3.70 - 3.81 (m, 4 H) 3.29 - 3.48 (m, 3 H) 2.75 - 2.99 (m, 2 H) 1.90 - 2.24 (m, 6 H)	
1-47					487.2
1-48					475.2
1-49					487.3
1-50				 ¹ H NMR (500 MHz, CHLOROFORM- <i>d</i>) δ ppm 10.19 - 10.52 (m, 1 H) 8.42 (d, <i>J</i> =8.4 Hz, 1 H) 8.03 (br s, 1 H) 7.77 (s, 1 H) 7.68 (s, 1 H) 7.45 (br d, <i>J</i> =8.2 Hz, 1 H) 6.81 (d, <i>J</i> =8.9 Hz, 1 H) 6.72 (br dd, <i>J</i> =8.8, 2.7 Hz, 1 H) 6.67 (br dd, <i>J</i> =5.5, 2.7 Hz, 1 H) 4.47 - 4.55 (m, 1 H) 4.30 (br s, 1 H) 4.11 (q, <i>J</i> =10.4 Hz, 1 H) 3.70 - 3.80 (m, 4 H) 3.29 - 3.40 (m, 1 H) 2.91 (br dd, <i>J</i> =14.6, 2.0 Hz, 1 H) 2.05 - 2.15 (m, 3 H)	491.2
1-51					491.2
1-52				 2HCl	491.3
1-53				 2HBr	491.3
1-54				 2MeSO ₃ H	491.2


1-55					495.1
1-56					501.3
1-57					479.2
1-58					501.3
1-59					501.3
1-60					501.3
1-61					501.3

General scheme 2



Compound 2-1: [1-(2-aminoethyl)-6-(3-chloro-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]-(6-chlorochroman-3-yl)methanone

Scheme 2A



Step 1: A mixture of **1b** (80 mg, 213 μ mol), potassium carbonate (176.8 mg, 1.3 mmol, 77 μ L), and 2-bromoethan-1-ol (87 mg, 639 μ mol) in DMF (2 mL) was heated to 70 °C and stirred overnight. The mixture was poured over water (10 mL) and extracted with EtOAc (10 mL). The organic layer was separated, washed with brine (10 mL), dried over Na₂SO₄, filtered, and concentrated. The residue was purified by flash column chromatography on silica gel column using a gradient of 10-100% EtOAc in hexane to give ([6-bromo-1-(2-hydroxyethyl)pyrrolo[3,2-c]pyridin-3-yl]-(6-chlorochroman-3-yl)methanone) as a white solid (**2a** 68 mg, 68 % yield). LC/MS found 435.1 and 437.0 [M+H]⁺.

Step 2: A mixture of **2a** (56 mg, 129 μ mol) and triethylamine (27 μ L, 193 μ mol) in DCM (3 mL) was cooled in an ice bath, treated with MsCl (15 μ L, 193 μ mol), and stirred for 1 hour at 0 °C. The solution was diluted with DCM (10 mL), washed sequentially with water and brine (10 mL each), dried (Na₂SO₄), treated with silica gel, and then evaporated under reduced pressure. The material was purified by silica gel column chromatography to provide 2-(6-bromo-3-(6-chlorochromane-3-carbonyl)-1H-pyrrolo[3,2-c]pyridin-1-yl)ethyl methanesulfonate (**2b**, 59 mg, 115 μ mol, 89% yield) as a clear oil. LC/MS found 513.0, 515.1 [M+H]⁺.

Step 3: A mixture of **2b** (29 mg, 56 μ mol) and sodium azide (46 mg, 113 μ mol) in DMF (2 mL) was heated to 60 °C and shaken overnight. The reaction mixture was diluted with EtOAc (10 mL), washed sequentially with water and brine (10 mL each), dried (Na₂SO₄), treated with silica gel, and evaporated under reduced pressure. The material was purified by silica gel column chromatography to provide (1-(2-azidoethyl)-6-bromo-1H-pyrrolo[3,2-c]pyridin-3-yl)(6-chlorochroman-3-yl)methanone (**2c**, 19 mg, 41 μ mol, 73% yield) as a white solid. LC/MS found 460.2, 462.2 [M+H]⁺.


Step 4: A mixture of **2c** (19 mg, 41 μ mol), triphenylphosphane (14 mg, 54 μ mol), and water (2~3 drops) in THF (1.5 mL) was heated to 55 °C and shaken overnight. The mixture was concentrated, poured over water (10 mL), and extracted with EtOAc (10 mL). The aqueous layer was separated and extracted with EtOAc (5 mL). The organic layers were combined, dried over silica gel, and evaporated under reduced pressure. The residue was purified by silica gel column chromatography to provide (1-(2-aminoethyl)-6-bromo-1H-pyrrolo[3,2-c]pyridin-3-yl)(6-chlorochroman-3-yl)methanone (**2d**, 16 mg, 37 μ mol, 89% yield) as a white solid. LC/MS found 434.0, 436.0 [M+H]⁺.

Step 5: A mixture of **2d** (16 mg, 37 μmol), tert-butyl 3-chloro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-1-carboxylate (24 mg, 72 μmol), potassium carbonate (2 M in water, 53 μL), and tetrakis(triphenylphosphine)palladium (2.5 mg, 2.1 μmol) in 1,4-dioxane (2 mL) was degassed via sparging with argon or nitrogen for 5 minutes, heated to 70 $^{\circ}\text{C}$, and shaken overnight. The mixture was cooled to room temperature, concentrated, poured over water (5 mL), and extracted with EtOAc (5 mL) twice. The combined organic layer was dried over Na_2SO_4 , filtered, and concentrated to afford tert-butyl 4-(1-(2-aminoethyl)-3-(6-chlorochroman-3-carbonyl)-1H-pyrrolo[3,2-c]pyridin-6-yl)-5-chloro-1H-pyrazole-1-carboxylate **2e** and then it was used to next step without further purification.

Step 6: A solution of crude **2e** (21 mg) in DCM (1.0 mL) was treated with TFA (0.5 mL) and stirred at ambient temperature. After being stirred for 1.5 hours at rt, the mixture was concentrated under reduced pressure to provide a residue. The material was dissolved in EtOAc (10 mL) and washed with 2M sodium carbonate solution (10 mL). The aqueous layer was separated and extracted with EtOAc (5 mL), and the combined organics were dried over Na_2SO_4 , filtered, treated with silica gel, and evaporated under reduced pressure. The material was purified by silica gel column chromatography to provide (1-(2-aminoethyl)-6-(3-chloro-1H-pyrazol-4-yl)-1H-pyrrolo[3,2-c]pyridin-3-yl)(6-chloro-2H-chromen-3-yl)methanone (**2-1**, 8.3 mg, 18.2 μmol ; 25% yield for steps 5 and 6) as a white solid. LC/MS found 456.1 $[\text{M}+\text{H}]^+$.

Compound 2-12: (6-chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[(2S)-2-(dimethylamino)propyl]indol-3-yl]methanone

Scheme 2B



Step 1: A mixture of [1-[(1-aminocyclopropyl)methyl]-6-bromo-indol-3-yl]-(6-chlorochroman-3-yl)methanone (**2f**, 22 mg, 48 μmol) prepared with similar method to **2d**, formaldehyde (36 μL , 479 μmol), and acetic acid (12 μL , 210 μmol) in THF (1 mL) was stirred at room temperature. At about 30 minutes the reaction mixture was cooled in an ice bath and
5 treated with sodiumcyanoborohydride (6.6 mg, 105 μmol). The reaction mixture was stirred for additional 30min and quenched with aq. sat. NH_4Cl solution (1 mL). The mixture was concentrated, diluted with EtOAc (15 mL), washed sequentially with water and brine (15 mL each), dried (Na_2SO_4), treated with silica gel, and evaporated under reduced pressure. The material was purified by silica gel column chromatography to provide (6-bromo-1-((S)-2-(dimethylamino)propyl)-1H-indol-3-yl)(6-chlorochroman-3-yl)methanone (**2g**, 23 mg, 47 μmol ,
10 98% yield) as a pale brown solid. LC/MS found 475.1, 477.2 $[\text{M}+\text{H}]^+$.






















Step 2: A mixture of **2g** (23 mg, 48 μmol), tert-butyl 5-chloro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-1-carboxylate (25 mg, 73 μmol), potassium carbonate (2 M in water, 60 μL , 121 μmol), and [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (2.0 mg, 2.4 μmol) in 1,4-dioxane (2 mL) was degassed via sparging with argon or
15 nitrogen for 5 minutes, heated to 70 $^\circ\text{C}$, and shaken overnight. The mixture was cooled to room temperature, concentrated, poured over water (5 mL), and extracted with EtOAc (5 mL) twice. The combined organic layers were dried over Na_2SO_4 , filtered, and concentrated to afford tert-butyl 3-chloro-4-(3-(6-chlorochromane-3-carbonyl)-1-((S)-2-(dimethylamino)propyl)-1H-indol-6-yl)-1H-pyrazole-1-carboxylate **2h**, and then it was used to the next step without further
20 purification.

Step 3a for a Boc protected pyrazole compound A solution of crude **2h** (29 mg) in DCM (1.0 mL) was treated with TFA (0.5 mL) and stirred at room temperature. At about 1.5 hours the solvents were evaporated under reduced pressure, dissolved in EtOAc (10 mL), and
25 washed with 2M Na_2CO_3 solution (10 mL). The aqueous layer was separated and extracted with EtOAc (5 mL), and the combined organic layers were dried over Na_2SO_4 , filtered, treated with silica gel, and evaporated under reduced pressure. The residue was purified by silica gel column chromatography to provide tert-butyl 3-chloro-4-(3-(6-chlorochromane-3-carbonyl)-1-((S)-2-(dimethylamino)propyl)-1H-indol-6-yl)-1H-pyrazole-1-carboxylate (**2-12**, 21 mg, 42 μmol ; 87%
30 yield for steps 2 and 3a) as a pale brown solid. LC/MS found 497.3 $[\text{M}+\text{H}]^+$.

The following compounds **2-2** to **2-16** shown in Table 2 were prepared by the method (General Scheme 2) similar to that described for the preparation of compound **2-1** or **2-12** using appropriate **2a** (azaindolic/indolic core) and appropriate boronic acid/ester (R^3).


Table 2

Compd	R^3 boronic acid	(9)	Structure	$[M+H]^+$
2-2				454.2
2-3				440.1
2-4				439.2
2-5				469.2
2-6				465.2

2-7				469.2
2-8				453.1
2-9				465.2
2-10				479.3
2-11				479.2
2-12				497.3
2-13				509.2


2-14				493.3
2-15				497.2
2-16				481.2
2-17				481.2

General Scheme 3



Compound 3-1: [1-[2-(azetidin-1-yl)ethyl]-6-(3-chloro-1H-pyrazol-4-yl)pyrrolo[2,3-b]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone

Scheme 3



Step 1: A mixture of **3a** [6-bromo-1-(2-hydroxyethyl)pyrrolo[2,3-b]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone, 85.0 mg, 203 μmol) prepared by the similar method in Scheme2, and triethylamine (56.5 μL , 405 μmol) in DCM (1.5 mL) was cooled in an ice bath, treated with methanesulfonyl chloride (15 μL , 193 μmol), and stirred for 1 hour at 0 °C. The solution was diluted with DCM (10 mL), washed sequentially with water and brine (10 mL each), dried (Na_2SO_4), treated with silica gel, and evaporated under reduced pressure. The material was purified by silica gel column chromatography to provide 2-(6-bromo-3-(6-fluorochroman-3-carbonyl)-1H-pyrrolo[2,3-b]pyridin-1-yl)ethyl methanesulfonate (**3b**, 86 mg, 173 μmol , 85% yield) as a pale brown oil. LC/MS found 497.1, 499.2 $[\text{M}+\text{H}]^+$.

Step 2: A mixture of **3b** (20 mg, 40 μmol), potassium carbonate (25 mg, 181 μmol), and azetidine hydrochloride (11 mg, 121 μmol) in DMF (1 mL) was heated to 50 °C and shaken overnight. The reaction mixture was diluted with EtOAc (10 mL), washed sequentially with water and brine (10 mL each), dried (Na_2SO_4), treated with silica gel, and evaporated under reduced pressure. The residue was purified by silica gel column chromatography to provide (1-(2-(azetidin-1-yl)ethyl)-6-bromo-1H-pyrrolo[2,3-b]pyridin-3-yl)(6-fluorochroman-3-

yl)methanone (**3c**, 4.7 mg, 10 μ mol, 25% yield) as a pale brown solid. LC/MS found 458.2, 460.1 $[M+H]^+$.

Step 3: A mixture of **3c** (4.7 mg, 10 μ mol), tert-butyl 3-chloro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-1-carboxylate (6.1 mg, 18 μ mol) or 5-chloro-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-tosyl-1H-pyrazole, potassium carbonate (2 M in water, 13 μ L), and tetrakis(triphenylphosphine)palladium (3.5 mg, 2.1 μ mol) in 1,4-dioxane (2 mL) was degassed via sparging with argon or nitrogen for 5 minutes, heated to 70 °C, and shaken overnight. The mixture was cooled to room temperature, concentrated, poured over water (5 mL), and extracted with EtOAc (x2) twice. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated to afford intermediate **3d-Boc** (Boc (tert-butyl 4-(1-(2-(azetidin-1-yl)ethyl)-3-(6-fluorochroman-3-carbonyl)-1H-pyrrolo[2,3-b]pyridin-6-yl)-5-chloro-1H-pyrazole-1-carboxylate) or **3d-Ts** ([1-[2-(azetidin-1-yl)ethyl]-6-[3-methoxy-1-(p-tolylsulfonyl)pyrazol-4-yl]indol-3-yl]-(6-chlorochroman-3-yl)methanone). The Intermediate **3d-Boc** or **3d-Ts** was used to the next step without further purification.

Step 4a: A solution of crude **3d-Boc**, (6.5 mg) in DCM (1.0 mL) was treated with TFA (0.5 mL) and stirred at ambient temperature. At about 1.5 hours the solvents were evaporated under reduced pressure to provide a residue. The material was dissolved in EtOAc (10 mL) and washed with 2M Na₂CO₃ solution (10 mL). The aqueous layer was separated and extracted with EtOAc (5 mL), and the combined organics were dried over Na₂SO₄, filtered, treated with silica gel, and evaporated under reduced pressure. The material was purified by silica gel column chromatography to provide [1-[2-(azetidin-1-yl)ethyl]-6-(3-chloro-1H-pyrazol-4-yl)pyrrolo[2,3-b]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone (**3-1**, 2.1 mg, 4.4 μ mol; 42% yield for steps 3 and 4a) as a pale brown solid. LC/MS found 480.2 $[M+H]^+$.

Step 4b: A solution of **3d-Ts**, (crude, 30 mg) in THF (1 mL) was cooled on an ice bath, then treated with aq. 2N NaOH solution (0.20 mL). The solution was stirred at 0°C for 30 minutes, and it was neutralized with aqueous 1N HCl solution (0.40 mL), diluted with EtOAc (10 mL) and saturated aqueous Na₂CO₃ solution (10 mL). The aqueous phase was separated and extracted with EtOAc (x2). The combined organic layers were, dried over Na₂SO₄, filtered, treated with silica gel, and evaporated under reduced pressure. The material was purified by silica gel column chromatography to provide [1-[2-(azetidin-1-yl)ethyl]-6-(3-methoxy-1H-






















pyrazol-4-yl]indol-3-yl]-(6-chlorochroman-3-yl)methanone (**3-4**, 4.0 mg, 8.5 μ mol, 17% yield for step 3 and 4b) as a white solid. LC/MS found 491.2 $[M+H]^+$.

The following compounds **3-2** to **3-19** shown in Table 3 were prepared by the method (General Scheme 3) similar to that described for the preparation of compound **3-1** or **3-4** using appropriate (7) (azaindolic/indolic core), appropriate boronic acid/ester (R^3), and an appropriate R^1 substituent.

Table 3

Compd	R^1R^2 Substituent	R^3 boronic acid	(7)	Structure	$[M+H]^+$
3-2	$-NH_2$				454.2
3-3	$-NH_2$				454.2
3-4					491.2
3-5	$-NH_2$				453.2
3-6	$-NH-$				493.2
3-7	$-NH-$				493.2


3-8					493.2
3-9					469.2
3-10					497.2
3-11					493.3
3-12					511.2
3-13					507.3
3-14					507.2

3-15				 2HCl	507.2
3-16				 2MeSO ₃ H	507.2
3-17				 Et-N-Et	507.2
3-18				 Et-N-Et	507.2
3-19				 Et-N-Et 2HCl	507.2
3-20				 Et-N-Et 2HBr	507.2
3-21				 Et-N-Et 2MeSO ₃ H	507.2

3-22					491.3
3-23					465.2
3-24					449.2
3-25					493.2
3-26					493.2
3-27					493.2
3-28					493.2


3-29					493.2
3-30					477.3

General Scheme 4



Compound 4-1 [1-[(1-aminocyclopropyl)methyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone

Scheme 4



Step 1: A mixture of (6-bromo-1H-indol-3-yl)-(6-chlorochroman-3-yl)methanone **1b** (300 mg, 768 μmol), potassium carbonate (318 mg, 2.3 mmol), and methyl 1-(bromomethyl)cyclopropane carboxylate (445 mg, 2.3 mmol) in DMF (4 mL) was heated to 70 $^{\circ}\text{C}$ and shaken overnight. The reaction mixture was diluted with EtOAc (15 mL), washed sequentially with water and brine (15 mL each), dried (Na_2SO_4), treated with silica gel, and evaporated under reduced pressure. The material was purified by silica gel column chromatography to provide methyl 1-((6-bromo-3-(6-chlorochromane-3-carbonyl)-1H-indol-1-yl)methyl)cyclopropane-1-carboxylate (**4a**, 380 mg, 756 μmol , 98% yield) as a pale brown solid. LC/MS found 502.2, 504.3 $[\text{M}+\text{H}]^+$.

Step 2: A mixture of **4a** (380 mg, 756 μmol) and $\text{LiOH}\cdot\text{H}_2\text{O}$ (2M, 3.8 mL, 7.6 mmol) in THF (4 mL) was stirred at room temperature overnight. The mixture was acidified to pH 3 to 4 with aqueous 1N HCl solution, and the resulting precipitate was vacuum filtered, washed with water, and dried under a vacuum overnight to provide 1-((6-bromo-3-(6-chlorochromane-3-carbonyl)-1H-indol-1-yl)methyl)cyclopropane-1-carboxylic acid, (**4b** 290 mg, 593 μmol , 78% yield) as a pale brown solid. LC/MS found 487.1, 489.1 $[\text{M}+\text{H}]^+$.

Step 3: A mixture of **4b** (290 mg, 593 μmol), Triethylamine (165 μL , 1.19 mmol), and [azido(phenoxy)phosphoryl]oxybenzene (218 μL , 1.01 mmol) in THF (3 mL) was shaken

overnight at rt. The reaction mixture was diluted with EtOAc (10 mL), washed sequentially with water and brine (10 mL each), dried (Na₂SO₄), treated with silica gel, and evaporated under reduced pressure. The material was purified by silica gel column chromatography to provide 1-((6-bromo-3-(6-chlorochromane-3-carbonyl)-1H-indol-1-yl)methyl)cyclopropane-1-carbonyl azide (**4c**, 225 mg, 438 μ mol, 73% yield) as a pale brown solid. LC/MS found 513.2, 513.2 [M+H]⁺.

Step 4: A solution of **4c** (215 mg, 418 μ mol) in tert-butanol (1 mL) and toluene (2 mL) was heated to 100 °C and shaken overnight. The reaction mixture was cooled to room temperature, concentrated, diluted with EtOAc (10 mL), washed sequentially with water and brine (10 mL each), dried (Na₂SO₄), treated with silica gel, and evaporated under reduced pressure. The material was purified by silica gel column chromatography to provide tert-butyl 1-((6-bromo-3-(6-chlorochromane-3-carbonyl)-1H-indol-1-yl)methyl)cyclopropyl)carbamate (**4d**, 170 mg, 304 μ mol, 72% yield) as a pale brown solid. LC/MS found (not observed in LCMS) [M+H]⁺.

Step 5: A mixture of **4d** (45 mg, 80 μ mol), tert-butyl 5-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-1-carboxylate (42 mg, 129 μ mol), potassium carbonate (2 M in water, 100 μ L), and cyclopentyl(diphenyl)phosphane;dichloromethane;dichloropalladium;iron (3.3 mg, 4.0 μ mol) in 1,4-dioxane (2 mL) was degassed via sparging with argon or nitrogen for 5 minutes, heated to 70 °C, and shaken overnight. The mixture was cooled to room temperature, concentrated, poured over water (5 mL), and extracted with EtOAc (5 mL) twice. The combined organic layer was dried over Na₂SO₄, filtered, and concentrated to afford tert-butyl 4-(1-((tert-butoxycarbonyl)amino)cyclopropyl)methyl)-3-(6-chlorochromane-3-carbonyl)-1H-indol-6-yl)-3-methoxy-1H-pyrazole-1-carboxylate **4e** and then used to the next step without further purification.




Step 6 A solution of crude **4e** (54 mg) in DCM (1.0 mL) was treated with TFA (0.5 mL) and stirred at ambient temperature. At about 1.5 hours the solvents were evaporated under reduced pressure to provide a residue. The material was dissolved in EtOAc (10 mL) and washed with 2M Na₂CO₃ solution (10 mL). The aqueous layer was separated and extracted with EtOAc (5 mL), and the combined organics were dried over Na₂SO₄, filtered, treated with silica gel, and evaporated under reduced pressure. The material was purified by silica gel column

chromatography to provide [1-[(1-aminocyclopropyl)methyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone (**4-1**, 25 mg, 52 μ mol; 65% yield for steps 5 and 6) as a pale brown solid. LC/MS found 480.2 $[M+H]^+$.


The following compounds **4-2** to **4-6** shown Table 4 were prepared by the method (General Scheme 4) similar to that described for the preparation of compound **4-1** using appropriate **1b** (azaindolic/indolic core), appropriate boronic acid/ester (R^3), and an appropriate R^1R^2 substituent.

Table 4

Compd	R^3 boronic acid	1b	Structure	MS found $[M+H]^+$
4-2				481.2
4-3				477.2
4-4				465.3
4-5				465.2


4-6				461.2
-----	---	---	--	-------

General Scheme 5



Compound 5-1: (6-chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(3-methyl-1H-pyrazol-4-yl)indol-3-yl]methanone

Scheme 5A



Step 1: A solution of 6-bromo-1H-indole (5.00 g, 25.5 mmol) in DMF (150 mL) was treated with sodium hydride (60% dispersion in mineral oil; 3.08 g, 77.0 mmol) in portions. The solution was stirred at ambient temperature for 30 minutes, 2-dimethylaminoethyl chloride hydrochloride (4.83 g, 33.5 mmol) was added, and the mixture was stirred at 70 °C for five hours. The mixture was diluted with EtOAc (500 mL), washed sequentially with water (3x500 mL) and brine (300 mL), dried (Na₂SO₄), filtered, and evaporated under reduced pressure. The residue was purified by silica gel column chromatography to provide 2-(6-bromo-1H-indol-1-yl)-N,N-dimethylethan-1-amine (**5a**, 5.93 g, 22.2 mmol, 87% yield) as a yellow liquid. LC/MS found 267.1, 269.1 [M+H]⁺.

Step 2: A solution of **5a** (2.51 g, 9.39 mmol) in DMF (90 mL) was cooled on an ice bath, then treated with N-iodosuccinimide (2.22 g, 9.9 mmol), and the ice bath was removed. After stirring for 90 minutes, the reaction was diluted with EtOAc (500 mL) and washed with water (500 mL). The aqueous phase was extracted with EtOAc (150 mL), and the combined organic layers were washed with brine (2x450 mL), dried (Na₂SO₄), filtered, and evaporated under reduced pressure. The residue was purified by silica gel column chromatography to provide 2-(6-

bromo-3-iodo-indol-1-yl)-*N,N*-dimethyl-ethanamine (**5b**, 2.12 g, 5.38 mmol, 57% yield) as a red liquid that solidified on standing. LC/MS found 393.0, 395.0 [M+H]⁺.

Step 3: To **5b** (2.75 g, 7.00 mmol) in a 250 mL round bottom flask was added THF (100 mL) under nitrogen. The solution was cooled on a dry ice / acetonitrile bath (approximate
5 temperature -42 °C) and treated dropwise (3 minutes) with isopropylmagnesium chloride (2 M in THF, 4.90 mL). After three minutes the cooling bath was replaced with an ice bath, and the mixture was stirred for 60 minutes. The reaction was cooled back down to -42 °C, and a solution of 6-chloro-*N*-methoxy-*N*-methyl-chromane-3-carboxamide (**6-16**, 2.68 g, 10.49 mmol) in THF (27 mL) was added dropwise (3 minutes). The cold bath was removed, and the reaction was
10 stirred for 3 hours. The reaction was quenched by addition of 25% aqueous NH₄Cl (20 mL) and stirred for 10 minutes. The THF was removed under reduced pressure. The aqueous residue was diluted with water to 200 mL and extracted with DCM (200 mL). The organic layer was washed with brine (150 mL), dried over Na₂SO₄, treated with silica gel, and evaporated under reduced pressure. The residue was purified by silica gel column chromatography to provide [6-bromo-1-
15 [2-(dimethylamino)ethyl]indol-3-yl]-(6-chlorochroman-3-yl)methanone (**5c**, 1.02 g, 2.2 mmol, 32% yield) as a yellow solid. LC/MS 461.1, 463.1 [M+H]⁺.


Step 4: A suspension **5c** (143.5 mg, 310.8 μmol), *tert*-butyl 3-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-1-carboxylate (146.7 mg, 476.0 μmol), PdCl₂(dppf)-CH₂Cl₂ (39.1 mg, 47.88 μmol), and potassium phosphate tribasic (198.2 mg, 933.7
20 μmol) in dioxane (5 mL) and water (0.3 mL) in a 40 mL vial was bubbled with nitrogen for 12 minutes and shaken at 70 °C overnight. The solution was evaporated under reduced pressure, and the residue was dissolved in EtOAc (30 mL), washed sequentially with water (10 mL) and brine (10 mL), dried over Na₂SO₄, and concentrated under reduced pressure to provide impure *tert*-butyl 4-[3-(6-chlorochromane-3-carbonyl)-1-[2-(dimethylamino)ethyl]indol-6-yl]-3-methyl-
25 pyrazole-1-carboxylate (**5d**, 290.7 mg) as a brown solid. The resulting residue was used without further purification. LC/MS found 563.3 [M+H]⁺.

Step 5: A solution of impure **5d** (290.7 mg) in DCM (2 mL) and MeOH (0.1 mL) was treated with TFA (2 mL). The solution was stirred at ambient temperature for 25 minutes. The solvents were evaporated under reduced pressure. The residue was dissolved in MeOH (10 mL)
30 and evaporated, twice. The residue was dissolved in EtOAc (30 mL), washed sequentially with

aqueous NaOH (1M, 5 mL) and brine (5 mL), dried over Na₂SO₄, treated with silica gel, and then concentrated under reduced pressure. The residue was purified by silica gel column chromatography to provide (6-chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(3-methyl-1H-pyrazol-4-yl)indol-3-yl]methanone (**5-1**, 99.3 mg, 214.48 μ mol; 69% yield for steps 4 and 5) as a tan solid. LC/MS found 463.3 [M+H]⁺.

Compound 5-29: (6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-methoxy-1H-pyrazol-4-yl)-1H-indazol-3-yl)methanone

Scheme 5B



Step 1: To a mixture of 6-bromo-3-iodo-1H-indazole (973 mg, 3.01 mmol), potassium carbonate (1.25 g, 9.04 mmol), and sodium iodide (98 mg, 0.65 mmol) in acetone (15 mL) was added 2-chloro-N,N-dimethyl-ethanamine hydrochloride (479 mg, 3.33 mmol). The mixture was heated to reflux with stirring overnight, cooled to room temperature, poured over H₂O, and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue was purified by a silica gel column chromatography using a gradient of 10-40% EtOAc in heptanes to provide 2-(6-bromo-3-iodo-1H-indazol-1-yl)-N,N-dimethylethan-1-amine as an off-white solid in 81% yield (**5e**, 965 mg, 2.5 mmol). LC/MS found 394.0 [M+H]⁺.

Step 2: To a solution of **5e** (542 mg, 1.38 mmol) in dry THF (10 mL) cooled to -40 °C was added dropwise *iso*-propylmagnesium chloride (2.0 M in THF, 0.96 mL, 1.9 mL). The mixture was allowed to stir at this temperature for 1 hour before the dropwise addition of a


solution of 6-chloro-*N*-methoxy-*N*-methyl-chromane-3-carboxamide (**6-16**, 459 mg, 1.80 mmol) in dry THF (4 mL). The mixture was warmed up to room temperature and allowed to stir for 2 hours. The reaction was quenched by the addition of saturated aqueous NH₄Cl and extracted with EtOAc (x2). The combined organic layers were washed with brine, dried over Na₂SO₄, and then concentrated to dryness. The residue was purified by silica gel column chromatography using a gradient of 50-100% EtOAc in heptanes to provide (6-bromo-1-(2-(dimethylamino)ethyl)-1*H*-indazol-3-yl)(6-chlorochroman-3-yl)methanone as a white solid in 11% yield (**5f**, 72 mg, 0.16 mmol). LC/MS found 462.1 [M+H]⁺.

Step 3: Starting with **5f** and 3-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-tosyl-1*H*-pyrazole by using the Suzuki protocol Step 4 in Scheme 5, (6-Chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-methoxy-1-tosyl-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)methanone was obtained as a light brown solid (**5g**, 47 mg, 69 μmol). LC/MS found 635.3 [M+H]⁺.

Step 4: Starting with **5g** (6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-methoxy-1-tosyl-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl) by using the general deprotection Step5a in Scheme 1, (6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-methoxy-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)methanone was obtained as a yellow (**5-29**, 11 mg, 23 μmol). LC/MS found 480.2 [M+H]⁺.

Compound 5-34: (6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-fluoro-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)methanone

Scheme 5C





Step 1: To an oven dried 40 mL vial equipped with a stir bar, were charged with [6-bromo-1-[2-(dimethylamino)ethyl]indazol-3-yl]-(6-chlorochroman-3-yl)methanone (**5h**, 399 mg, 0.86 mmol), bis(pinacolato) diboron (329 mg, 1.30 mmol), potassium acetate (170 mg, 1.73

mmol), and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (35 mg, 42 μ mol). The vial was sealed with a septum cap. 1,4-dioxane (9 mL) was added via syringe. The mixture was purged with N₂ for 10 minutes and heated to 100 °C with stirring overnight. The mixture was cooled to room temperature and filtered through a plug of diatomaceous earth, and the filtrate was concentrated to dryness. The residue was purified by silica gel column chromatography using a gradient of 1-7% CH₃OH in CH₂Cl₂ to provide **5i** (6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-indazol-3-yl)methanone as a sticky brown solid in quantitative yield. LC/MS found 510.3 [M+H]⁺.

Step 2: Starting with *tert*-butyl 4-bromo-3-fluoro-1*H*-pyrazole-1-carboxylate and **5a** using general Suzuki reaction followed by deprotection of Step 4 and **5a** in Scheme 5, (6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-fluoro-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)methanone was obtained as a slightly brown solid (**5-34**, 127 mg, 31% yield). LC/MS found 468.3 [M+H]⁺.

Alternatively, racemic mixtures of compounds were separated by supercritical fluid chromatography as below.

Compound 5-19: [(3*S*)-6-Chlorochroman-3-yl]-[1-[2-(dimethylamino)ethyl]-6-(5-methoxy-1*H*-pyrazol-4-yl)indol-3-yl]methanone & Compound 5-20: [(3*R*)-6-chlorochroman-3-yl]-[1-[2-(dimethylamino)ethyl]-6-(5-methoxy-1*H*-pyrazol-4-yl)indol-3-yl]methanone





A sample of racemic Compounds **5-18** was separated by supercritical fluid chromatography under the following conditions: Column, NanoMicro UniChiral AS-5H 21 x 250 mm; Mobile phase, 45% 2-propanol and 0.25% diethylamine in CO₂; Flow rate, 70 mL/min;

Sample, 149.5 mg dissolved in 6 mL methanol and 6 mL dichloromethane; Injection, 1.0 mL; detection: 254 nm.

Compound 5-19: second-eluting peak, 64.7 mg, 100% purity, 97.4% ee. LC/MS found 479.2 $[M+H]^+$.

5 **Compound 5-20:** first-eluting peak, 66.0 mg, 100% purity, 100% ee. LC/MS found 479.2 $[M+H]^+$.

Compound 5-22: (S)-(6-(3-chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-chlorochroman-3-yl)methanone & **Compound 5-23: (R)-(6-(3-chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-chlorochroman-3-yl)methanone**
10 and





A sample of racemic Compound **5-21** was separated by supercritical fluid chromatography under the following conditions: Column, "ChiralPak IC-H 21 x 250 mm; Mobile phase, 40% methanol in CO₂; Flow rate, 70 mL/min; Sample, 19.4 mg dissolved in 2.0
15 mL methanol and 2.0 mL dichloromethane; Injection, 0.75 mL; Detection: 220 nm".

Compound 5-22: second-eluting peak, 80.2 mg, 100% purity, 96.4% ee. LC/MS found 452.2 $[M+H]^+$.

Compound 5-23: first-eluting peak, 77.5 mg, 100% purity, 100% ee. LC/MS found 452.2 $[M+H]^+$.

20 **Compound 5-25: [1-[2-(Dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-[(3R)-6-fluorochroman-3-yl]methanone** & **Compound 5-27: [1-[2-(Dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-[(3S)-6-fluorochroman-3-yl]methanone**



A sample of racemic Compound **5-24** was separated by supercritical fluid chromatography under the following conditions: Column, ChiralPak AD-H 21 x 250 mm; Mobile phase, 40% 2-propanol and 0.25% diethylamine in CO₂; Flow rate, 70 mL/min; Sample, 82.0 mg dissolved in 4.0 mL methanol and 4.0 mL dichloromethane; Injection, 2.0 mL; Detection, 254 nm.



















Compound 2-25: second-eluting peak, 31.2 mg, 100% purity, 100% ee. LC/MS found 463.3 [M+H]⁺.
















Compound 5-27: first-eluting peak, 29.2 mg, 100% purity, 100% ee. LC/MS found 463.3 [M+H]⁺.



















The following compounds **5-2** to **5-18** and **5-28** to **5-51** shown in Table 5 were prepared by methods similar to those described for the preparation of compound **5-1**, **5-29** or **5-34**, using the appropriate boronic acid/ester and Weinreb amide.

Table 5



















Compd	R ³ boronic acid or ester	Weinreb amide	Structure	[M+H] ⁺
5-2				417.2

5-3				415.3
5-3				479.3
5-4				479.3
5-5				479.3
5-6				479.3
5-7				479.3



















5-8				479.3
5-9				523.2
5-10				541.3
5-11				459.2
5-12				524.2

5-13				543.1
5-14				561.2
5-15				543.3
5-16				469.1
5-17				485.2
5-18				451.2



















5-19				467.2
5-20				467.2
5-21				467.2
5-22				467.2
5-23				464.2
5-24				463.3













5-25				475.2
5-26				448.2
5-27				447.3
5-28				479.2
5-29				479.2
5-30				479.3



















5-31				479.3
5-32				479.3
5-33				479.3
5-34				483.2
5-35				483.2
5-36				483.2




5-37				463.3
5-38				463.3
5-39				463.3
5-40				463.3
5-41				475.3
5-42				481.2

5-43				418.2
5-44				418.2
5-45				480.3
5-46				480.3
5-47				480.3
5-48			 ¹ H NMR (500 MHz, CHLOROFORM- <i>d</i>) δ ppm 8.42 (d, <i>J</i> =8.5 Hz, 1 H), 7.88 (d, <i>J</i> =1.8 Hz, 1 H), 7.80 (s, 1 H), 7.58 (dd, <i>J</i> =9.7, 1.1 Hz, 1 H), 7.22 (d, <i>J</i> =2.4 Hz, 1 H), 7.18 (dd, <i>J</i> =8.7, 2.4 Hz, 1 H), 6.90 (d, <i>J</i> =8.7 Hz, 1 H), 4.71 - 4.75 (m, 1 H), 4.67 (t, <i>J</i> =6.8 Hz, 2 H), 4.39 - 4.46 (m, 1 H), 4.33 (t, <i>J</i> =10.5 Hz, 1 H), 3.36	452.2


			(dd, $J=16.1, 10.5$ Hz, 1 H), 3.14 (dd, $J=16.9, 4.7$ Hz, 1 H), 3.01 (t, $J=6.8$ Hz, 2 H), 2.44 (s, 6 H)	
5-49				476.3
5-50				480.2
5-51				464.2
5-52				468.3
5-53				468.3
5-54				484.2

5-55			 ¹H NMR (500 MHz, CHLOROFORM-<i>d</i>) δ ppm 9.52 (br s, 1 H), 8.32 (dd, <i>J</i>=8.5, 0.5 Hz, 1 H), 7.76 (d, <i>J</i>=2.0 Hz, 1 H), 7.70 (s, 1 H), 7.47 (dd, <i>J</i>=8.4, 1.2 Hz, 1 H), 6.78 - 6.88 (m, 3 H), 4.60 - 4.64 (m, 1 H), 4.58 (t, <i>J</i>=6.9 Hz, 2 H), 4.30 - 4.38 (m, 1 H), 4.19 - 4.25 (m, 1 H), 3.28 (dd, <i>J</i>=16.6, 10.3 Hz, 1 H), 3.04 (dd, <i>J</i>=16.2, 4.6 Hz, 1 H), 2.92 (t, <i>J</i>=6.8 Hz, 2 H), 2.34 (s, 6 H)	452.2
5-56			 ¹H NMR (500 MHz, CHLOROFORM-<i>d</i>) δ ppm 11.06 (br s, 1 H), 8.27 (d, <i>J</i>=8.4 Hz, 1 H), 7.65 (s, 1 H), 7.52 - 7.59 (m, 1 H), 7.38 (br d, <i>J</i>=8.4 Hz, 1 H), 6.77 - 6.89 (m, 3 H), 4.72 (br d, <i>J</i>=6.6 Hz, 1 H), 4.61 (br d, <i>J</i>=10.5 Hz, 1 H), 4.49 (br dd, <i>J</i>=14.1, 4.2 Hz, 1 H), 4.29 - 4.38 (m, 1 H), 4.23 (t, <i>J</i>=9.8 Hz, 1 H), 3.84 (br d, <i>J</i>=2.1 Hz, 1 H), 3.55 (br s, 1 H), 3.28 (br dd, <i>J</i>=16.4, 10.1 Hz, 1 H), 3.00 - 3.09 (m, 1 H), 2.87 - 2.99 (m, 1 H), 2.22 (br s, 2 H), 2.09 (br s, 3 H)	465.2
5-57				465.3
5-58			 HBr	465.2

5-59				482.3
5-60				470.2
5-61				486.2
5-62				478.3
5-63				482.3
5-64				490.3


5-65				494.3
------	---	---	--	-------

General Scheme 6



Weinreb Intermediate 6-1: 6-Chloro-*N*-methoxy-*N*-methyl-chromane-3-carboxamide

5

































Weinreb Intermediate 6-1











A suspension of 6-chlorochromane-3-carboxylic acid (964.5 mg, 4.54 mmol), *N,O*-dimethyl hydroxylamine hydrochloride (665.3 mg, 6.82 mmol), and HBTU (2.15 g, 5.67 mmol) in DMF (33 mL) was treated with *N,N*-diisopropylethylamine (18.4 mmol, 3.2 mL). The solution was stirred at ambient temperature overnight. The solution was diluted with EtOAc (300 mL), washed successively with water (3x300 mL) and brine (300 mL), dried (Na₂SO₄), filtered, and evaporated under reduced pressure. The material was purified by silica gel column chromatography to provide 6-chloro-*N*-methoxy-*N*-methyl-chromane-3-carboxamide (Weinreb Intermediate 9-1; 1.06 g, 4.15 mmol, 91% yield) as a clear, colorless oil. LC/MS found 256.0 [M+H]⁺.

The following Weinreb amide Intermediates **6-2** through **6-21** shown in Table 6 were prepared by methods similar to those described for the preparation of Intermediate **6-1**, using the appropriate carboxylic acid.

Table 6

Intermediate	Carboxylic acid	Weinreb Intermediate	LCMS [M+H] ⁺
6-2			224.1
6-3			222.2
6-4			239.9
6-5			251.9
6-6			242.1
6-7			242.3
6-8			258.1

6-9			254.1
6-10			258.0
6-11			292.1
6-12			252.1
6-13			266.1
6-14			208.1
6-15			252.1
6-16			256.2

6-17			256.3
6-18			256.2
6-19			240.1
6-20			240.2
6-21			224.2

Example 2: ROCK Inhibition Assay

Kinase IC₅₀ was determined through an *in vitro* assay based on LANCE Ultra TR-FRET (Time-Resolved Fluorescence Resonance Energy Transfer) homogeneous technologies method (Perkin Elmer). Recombinant ROCK1 (amino acids 1-477) and ROCK2 (amino acids 5-554) proteins were purchased from Carna Biosciences and SignalChem. Compound activities were measured by Envision and IC₅₀s were calculated. The assays were performed in white LUMITRAC™ 200 96 well half-area microplates from Greiner Bio-One. The kinase reaction buffer consisted of 50 mM HEPES pH 7.5, 1 mM EGTA, 10 mM MgCl₂, 2 mM DTT and 0.01 % Tween-20. Kinases were incubated with 50 nM (ULight-CREBtide) substrate in the presence of 1 mM (ROCK1) or 1 mM (ROCK2) of ATP. The kinase reaction was carried out for 1 hour before addition of a stopping buffer to a final of 10 mM EDTA and 0.6 nM of LANCE Ultra Europium anti-phospho-CREB (Ser133) antibody (PerkinElmer TRF0200) in LANCE detection

buffer. All assay incubations were performed at room temperature, and the microplates were sealed with polyester film during that time. The reaction was incubated for 1 hour, and the signal was read in Envision in TR-FRET mode (excitation at 320 nm and emission at 615/ 665 nm).

The compounds of (1) exhibited useful pharmacological properties. As used herein, a way to describe potency of inhibitory activity (nM) is a value of inhibitory activity at 50 % (IC₅₀). The results are shown in Table 7 below, where an IC₅₀ of less than 20 nM is defined as “A,” an IC₅₀ of between 21 nM and 200 nM is defined as “B,” an IC₅₀ of between 201 nM and 500 nM is defined as “C,” and an IC₅₀ of greater than 501 nM is defined as “D.” Table 7 illustrates the inhibition of ROCK1 and ROCK2 by representative compounds of (1).

10

Table 7. Inhibition Activity of ROCK1 and ROCK2 at 1mM of ATP

Compound	ROCK1	ROCK2	Compound	ROCK1	ROCK2
1-1	D	B	3-10	D	B
1-2	D	B	3-11	D	B
1-3	D	B	3-12	D	B
1-4	D	A	3-13	D	A
1-8	D	C	3-14	D	A
1-9	C	B	3-17	D	B
1-10	D	B	3-20	C	A
1-11	D	D	3-21	C	B
1-12	D	C	3-22	C	A
1-13	D	A	3-23	D	A
1-14	D	C	3-24	D	B
1-15	D	B	3-25	C	A
1-16	D	D	3-26	D	A
1-17	D	B	3-28	D	B
1-18	D	B	3-30	C	B
1-19	D	B	4-1	A	A
1-20	D	D	4-2	A	A
1-21	A	A	4-3	B	A
1-22	A	A	4-4	B	A
1-23	D	B	4-5	A	A

1-24	D	B	4-6	B	A
1-25	C	B	5-1	D	C
1-26	C	A	5-2	D	C
1-27	C	C	5-3	D	C
1-28	D	D	5-3	D	A
1-29	C	B	5-4	D	A
1-30	D	A	5-8	D	A
1-31	D	B	5-9	C	A
1-32	D	A	5-10	C	A
1-33	D	D	5-11	B	B
1-34	D	C	5-12	D	B
1-35	B	A	5-13	C	A
1-36	C	A	5-14	D	B
1-37	D	A	5-15	D	B
1-38	C	A	5-16	D	C
1-39	D	B	5-17	D	D
1-40	D	A	5-18	C	C
1-41	C	A	5-19	D	B
1-42	C	A	5-20	C	A
1-43	B	A	5-21	C	A
1-44	D	A	5-23	D	C
1-45	D	B	5-24	D	D
1-46	D	B	5-25	D	C
1-47	D	A	5-26	D	D
1-48	D	B	5-27	D	D
1-49	D	B	5-28	D	A
1-50	D	A	5-29	D	C
1-51	D	A	5-30	C	A
1-55	D	A	5-34	C	A
1-56	D	B	5-35	D	C

1-57	D	C	5-36	C	A
1-58	D	B	5-37	D	A
1-59	D	A	5-38	D	A
2-1	B	A	5-39	D	A
2-2	B	A	5-40	D	C
2-3	B	A	5-41	C	B
2-4	A	A	5-42	D	C
2-5	B	A	5-43	D	B
2-6	B	A	5-44	C	B
2-7	A	A	5-45	D	A
2-8	D	D	5-46	C	A
2-9	C	A	5-47	D	B
2-10	A	A	5-48	D	A
2-11	B	A	5-49	B	A
2-12	D	A	5-50	B	A
2-13	B	B	5-51	D	B
2-14	C	A	5-52	C	A
2-15	C	A	5-53	C	A
2-16	D	A	5-54	D	C
2-17	C	A	5-55	C	A
3-1	D	A	5-56	D	B
3-2	D	B	5-57	C	A
3-3	D	B	5-59	D	B
3-4	D	C	5-60	D	B
3-5	C	A	5-61	D	B
3-6	D	B	5-62	D	D
3-7	C	B	5-63	D	D
3-8	D	C	5-64	D	D
3-9	D	C	5-65	D	D

Example 3: Blood–Brain Barrier (BBB) Penetrating Properties in Rat and Mouse

The compounds of (1) exhibited useful blood–brain barrier (BBB) penetrating properties. The concentrations of compounds of (1) in blood and brain were measured from plasma and brain samples taken 0.25 and 2 hours after intravenous (IV) administration of the compounds of (1) at 1 mg/kg in CD-1 mouse and Sprague-Dawley rat (n=3). After
5 euthanizing the mouse and rat, blood samples from cardiac puncture and brain tissue were collected. Plasma or brain homogenate was vortexed for 1 min. with internal standard in Methanol/Acetonitrile (1:1, v/v), and centrifuged at 4000–4500 rpm for 15 min. at room temperature to precipitate proteins. Subsequently, the supernatant was introduced into the LC/MS/MS system. An AB Sciex API 5500 mass spectrometer system (Sciex, USA) was used
10 for quantitative analysis.

The concentrations compounds of (1) in plasma and brain sample were determined using the standard calibration curve and brain-to-plasma concentration ratio was calculated. Table 8 illustrates the blood–brain barrier (BBB) ratio by representative compounds of (1).


Table 8. Blood–brain barrier (BBB) ratio

Compound	time (hour)	Brain/Plasma (mouse)		Brain/Plasma (rat)	
		Mean	SD	Mean	SD
1-55	0.25	0.67	0.07	0.4	0.02
	2	0.8	0.19	0.59	0.15
3-17	0.25	0.73	0.14	1.05	0.22
	2	0.92	0.43	0.98	0.12
3-24	0.25	0.96	0.33	1.37	0.27
	2	0.54	0.09	1.38	0.17
5-56	0.25	7.32	0.56	12.4	2.1
	2	10.2	1.6	20.2	6.6

15

CLAIMS

1. A compound of Formula (1), a pharmaceutically acceptable salt of the compound, or a prodrug of the compound or the salt:



5 wherein:

X is CH₂ or O;

Y is or CH;

Z¹, Z² and Z³ each are independently N or CH;

n is 0, 1, 2 or 3;

10 R¹ is H, heterocycloalkyl C₃-C₇cycloalkyl, or C₁-C₆alkyl wherein a nitrogen on the heterocycloalkyl is optionally substituted with C₁-C₃alkyl, wherein the C₃-C₇cycloalkyl or C₁-C₆alkyl is optionally substituted with one or more substituents selected from the group consisting of C₁-C₆alkyl and OH;

R² is H, amino, NR⁶R⁷, or NR⁸R⁹;

15 R³ is 5-6 membered heteroaryl or bicyclic heteroaryl, wherein the heteroaryl or bicyclic heteroaryl is optionally substituted with one or more substituents selected from the group consisting of halogen, CN, CHF₂, CF₃, C₁-C₃alkyl, and amino, wherein the 5-6 membered heteroaryl or bicyclic heteroaryl has 1-3 heteroatoms selected from the group consisting of oxygen and nitrogen;


20 R⁴ and R⁵ each are independently H, Cl, F, OH, CD₃, C₁-C₃alkyl, or C₁-C₃alkoxy, wherein the C₁-C₃alkyl or C₁-C₃alkoxy is optionally substituted with one or more substituents selected from the group consisting of amino, OH, and C₁-C₃alkoxy;

R⁶ and R⁷ each are independently H, CD₃, C₁-C₆alkyl, or C₃-C₇cycloalkyl, wherein the C₁-C₆alkyl or C₃-C₇cycloalkyl is optionally substituted with hydroxyl or C₁-C₃alkoxy; and

R^7 and R^8 , taken together with the nitrogen atom to which they are bonded a 4-6 membered saturated monocyclic group having no heteroatom other than the nitrogen atom to which R^7 and R^8 are bonded, wherein the 4-6 membered saturated or partially saturated monocyclic group is optionally substituted at one or two carbon atoms with halogen, amino, hydroxyl, or C_1 - C_3 alkoxy.

2. The compound, salt, or prodrug of claim 1, wherein R^1 is H, heterocycloalkyl C_3 - C_7 cycloalkyl, or C_1 - C_6 alkyl wherein a nitrogen on heterocycloalkyl is optionally substituted with methyl or ethyl, wherein the C_3 - C_7 cycloalkyl or C_1 - C_6 alkyl is optionally substituted with one or more suitable substituents selected from the group consisting of methyl, ethyl, and OH.

10 3. The compound, salt, or prodrug of claim 1, R^3 is selected from the following group:



4. The compound, salt, or prodrug of claim 1, wherein R^4 and R^5 each are independently H, Cl, F, OH, CH_3 , methoxy, OCD_3 , ethoxy, or iso-propoxy.

15 5. The compound, salt, or prodrug of claim 1, wherein R^6 and R^7 each are independently H, CD_3 , C_1 - C_6 alkyl, or C_3 - C_7 cycloalkyl, wherein the C_1 - C_6 alkyl or C_3 - C_7 cycloalkyl is optionally substituted with hydroxyl methoxy or ethoxy.

6. The compound, salt, or prodrug of claim 1, wherein R^8 and R^9 , taken together with the nitrogen atom to which they are bonded a 4-6 membered saturated monocyclic group having no
20 heteroatom other than the nitrogen atom to which R^9 and R^9 are bonded, wherein the 4-6

membered saturated or partially saturated monocyclic group is optionally substituted at one or two carbon atoms selected from the group consisting of halogen, hydroxyl, and methoxy.

7. A pharmaceutical composition comprising the compound, salt, or prodrug of claim 1 and an additional component selected from the group consisting of a pharmaceutically acceptable carrier, diluent, excipient, adjuvant, and a combination thereof.

8. A compound represented by the following list, a pharmaceutically acceptable salt of the compound, or a prodrug of the compound or the salt:

(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(3-methoxy-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]methanone;

10 (6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(3-methoxy-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]methanone;

(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(3-methoxy-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]methanone;

15 (6-(5-chloro-1H-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1H-pyrrolo[3,2-c]pyridin-3-yl)((S)-6-methoxychroman-3-yl)methanone

(6-(5-chloro-1H-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1H-pyrrolo[3,2-c]pyridin-3-yl)((S)-6-methoxychroman-3-yl)methanone dihydrochloride;

(6-(5-chloro-1H-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1H-pyrrolo[3,2-c]pyridin-3-yl)((S)-6-methoxychroman-3-yl)methanone dihydrobromide;

20 (6-(5-chloro-1H-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1H-pyrrolo[3,2-c]pyridin-3-yl)((S)-6-methoxychroman-3-yl)methanone dimethanesulfonic acid;

(6-Chlorochroman-3-yl)-[6-(3-chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]pyrrolo[2,3-b]pyridin-3-yl]methanone;

25 [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]pyrrolo[2,3-b]pyridin-3-yl]- (6-fluorochroman-3-yl)methanone;

(6-Chlorochroman-3-yl)-[6-(3-chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]pyrrolo[3,2-c]pyridin-3-yl]methanone;

- [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]pyrrolo[3,2-c]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone;
- [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]pyrrolo[2,3-b]pyridin-3-yl]-(6-methoxychroman-3-yl)methanone;
- 5 [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]pyrrolo[3,2-c]pyridin-3-yl]-(6-methoxychroman-3-yl)methanone;
- [1-(Azetidin-3-yl)-6-(3-chloro-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone;
- (6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)pyrrolo[2,3-b]pyridin-3-yl]methanone;
- 10 (6-Chlorochroman-3-yl)-[6-(3-chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]methanone;
- [1-(Azetidin-3-yl)-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- 15 (6-Chlorochroman-3-yl)-[6-(5-methoxy-1H-pyrazol-4-yl)-1-[(1-methylpyrrolidin-2-yl)methyl]indol-3-yl]methanone;
- (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[(1-methylpyrrolidin-2-yl)methyl]indol-3-yl]methanone;
- [1-(Azetidin-3-ylmethyl)-6-(5-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- 20 [1-(Azetidin-3-ylmethyl)-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- [1-[2-(Dimethylamino)ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;
- 25 (6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]methanone;

- (S)-(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]methanone;
- (R)-(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]methanone;
- 5 (6-Chlorochroman-3-yl)-[6-(5-fluoro-1H-pyrazol-4-yl)-1-[(1-methylpyrrolidin-2-yl)methyl]indol-3-yl]methanone;
- (6-Chlorochroman-3-yl)-[6-(5-methoxy-1H-pyrazol-4-yl)-1-[(1R)-1-methylpyrrolidin-2-yl)methyl]indol-3-yl]methanone;
- (6-Chlorochroman-3-yl)(6-(5-methoxy-1H-pyrazol-4-yl)-1-(((S)-1-methylpyrrolidin-2-yl)methyl)-1H-indol-3-yl)methanone;
- 10 [1-(Azetidin-2-ylmethyl)-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- [1-(Azetidin-2-ylmethyl)-6-(5-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- [1-(Azetidin-2-ylmethyl)-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- 15 (6-Chlorochroman-3-yl)-[2-(5-chloro-1H-pyrazol-4-yl)-7-[2-(dimethylamino)ethyl]pyrrolo[2,3-d]pyrimidin-5-yl]methanone;
- (6-Chlorochroman-3-yl)-[7-[2-(dimethylamino)ethyl]-2-(5-fluoro-1H-pyrazol-4-yl)pyrrolo[2,3-d]pyrimidin-5-yl]methanone;
- 20 (1-(((R)-Azetidin-2-yl)methyl)-6-(5-chloro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;
- (1-(((S)-Azetidin-2-yl)methyl)-6-(5-chloro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;
- 25 [1-(((2S)-Azetidin-2-yl)methyl)-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-methoxychroman-3-yl)methanone;

[1-(Azetidin-2-ylmethyl)-6-(5-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

[1-[(2R)-Azetidin-2-yl]methyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

5 (1-(((S)-Azetidin-2-yl)methyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-chlorochroman-3-yl)methanone;

(1-(((R)-Azetidin-2-yl)methyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;

10 [1-(2-Aminoethyl)-6-(3-chloro-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]-(6-chlorochroman-3-yl)methanone;

[1-(2-Aminopropyl)-6-(3-chloro-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone;

[1-(2-Aminoethyl)-6-(3-chloro-1H-pyrazol-4-yl)pyrrolo[3,2-c]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone;

15 [1-(2-Aminoethyl)-6-(3-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;

[1-[(2R)-2-Aminopropyl]-6-(3-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

20 [1-[(2R)-2-Aminopropyl]-6-(3-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

[1-[(2S)-2-Aminopropyl]-6-(3-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

[1-[(2S)-2-Aminopropyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

25 [1-[(2S)-2-Aminopropyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;

- [1-[(2S)-2-Aminobutyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- (6-(5-chloro-1H-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1H-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone;
- 5 [1-[(2R)-2-Aminobutyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[(2S)-2-(dimethylamino)propyl]indol-3-yl]methanone;
- (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[[1-(dimethylamino)cyclopropyl]methyl]indol-3-yl]methanone;
- 10 (6-Chlorochroman-3-yl)-[1-[(2R)-2-(dimethylamino)propyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]methanone;
- (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[(2R)-2-(dimethylamino)propyl]indol-3-yl]methanone;
- 15 (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[(2R)-2-(dimethylamino)propyl]indol-3-yl]methanone;
- (6-Chlorochroman-3-yl)-[6-(5-fluoro-1H-pyrazol-4-yl)-1-[(1-methylazetidin-3-yl)methyl]indol-3-yl]methanone;
- (6-Chlorochroman-3-yl)-[6-(5-methoxy-1H-pyrazol-4-yl)-1-[(1-methylazetidin-3-yl)methyl]indol-3-yl]methanone;
- 20 [6-(5-Fluoro-1H-pyrazol-4-yl)-1-[[2R)-1-methylazetidin-2-yl]methyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;
- [6-(5-Chloro-1H-pyrazol-4-yl)-1-[(1S)-1-methylazetidin-2-yl]methyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;
- 25 (6-Methoxychroman-3-yl)-[6-(5-methoxy-1H-pyrazol-4-yl)-1-[(1S)-1-methylazetidin-2-yl]methyl]indol-3-yl]methanone;

- [6-(5-Fluoro-1H-pyrazol-4-yl)-1-[[[(2S)-1-methylazetidin-2-yl]methyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;
- (6-Methoxychroman-3-yl)-[6-(5-methoxy-1H-pyrazol-4-yl)-1-[[[(1R)-1-methylazetidin-2-yl]methyl]indol-3-yl]methanone;
- 5 [6-(5-Chloro-1H-pyrazol-4-yl)-1-[[[(1R)-1-methylazetidin-2-yl]methyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;
- (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[[[(1R)-1-methylazetidin-2-yl]methyl]indol-3-yl]methanone;
- [1-[[[(1R)-1-Ethylazetidin-2-yl]methyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-methoxychroman-3-yl)methanone;
- 10 (6-Chlorochroman-3-yl)-[6-(5-fluoro-1H-pyrazol-4-yl)-1-[[[(1R)-1-methylazetidin-2-yl]methyl]indol-3-yl]methanone;
- [6-(5-Chloro-1H-pyrazol-4-yl)-1-[[[(1R)-1-ethylazetidin-2-yl]methyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;
- 15 [1-[2-(Azetidin-1-yl)ethyl]-6-(3-chloro-1H-pyrazol-4-yl)pyrrolo[2,3-b]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone;
- [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(methylamino)ethyl]pyrrolo[3,2-c]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone;
- [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(methylamino)ethyl]pyrrolo[2,3-b]pyridin-3-yl]-(6-fluorochroman-3-yl)methanone;
- 20 [1-[2-(Azetidin-1-yl)ethyl]-6-(3-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(methylamino)ethyl]indol-3-yl]-(6-fluorochroman-3-yl)methanone;
- 25 (6-Chlorochroman-3-yl)-[1-[3-(dimethylamino)propyl]-6-(3-methoxy-1H-pyrazol-4-yl)indol-3-yl]methanone;

(S)-(6-Chlorochroman-3-yl)(1-(3-(dimethylamino)propyl)-6-(3-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone;

(R)-(6-Chlorochroman-3-yl)(1-(3-(dimethylamino)propyl)-6-(3-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone;

5 (6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[2-(methylamino)ethyl]indol-3-yl]methanone;

(6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[2-ethyl(methyl)amino]ethyl]indol-3-yl]methanone;

10 (6-Chlorochroman-3-yl)-[1-[2-[ethyl(methyl)amino]ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]methanone;

(6-Chlorochroman-3-yl)-[6-(5-chloro-1H-pyrazol-4-yl)-1-[2-(diethylamino)ethyl]indol-3-yl]methanone;

(6-Chlorochroman-3-yl)-[1-[2-(diethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]methanone;

15 [6-(5-Chloro-1H-pyrazol-4-yl)-1-[2-(diethylamino)ethyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;

[1-[2-(Diethylamino)ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-methoxychroman-3-yl)methanone;

20 [6-(5-Chloro-1H-pyrazol-4-yl)-1-[2-(methylamino)ethyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;

[6-(5-Fluoro-1H-pyrazol-4-yl)-1-[2-(methylamino)ethyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;

[6-(5-Chloro-1H-pyrazol-4-yl)-1-[2-[ethyl(methyl)amino]ethyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;

25 [1-[2-[Ethyl(methyl)amino]ethyl]-6-(5-fluoro-1H-pyrazol-4-yl)indol-3-yl]-(6-methoxychroman-3-yl)methanone;

- [1-[(1-Aminocyclopropyl)methyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- [1-[(1-Aminocyclopropyl)methyl]-6-(5-chloro-1H-pyrazol-4-yl)indol-3-yl]-(6-chlorochroman-3-yl)methanone;
- 5 (1-[(1-Aminocyclopropyl)methyl]-6-(5-chloro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;
- (1-[(1-Aminocyclopropyl)methyl]-6-(5-fluoro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-chlorochroman-3-yl)methanone;
- 10 (1-[(1-Aminocyclopropyl)methyl]-6-(5-fluoro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-fluorochroman-3-yl)methanone;
- (1-[(1-Aminocyclopropyl)methyl]-6-(5-fluoro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;
- 2,3-Dihydro-1,4-benzodioxin-3-yl-[1-[2-(dimethylamino)ethyl]-6-(1H-pyrazol-4-yl)indol-3-yl]methanone;
- 15 Chroman-3-yl-[1-[2-(dimethylamino)ethyl]-6-(1H-pyrazol-4-yl)indol-3-yl]methanone;
- [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(6-methoxychroman-3-yl)methanone;
- (S)-(6-(3-Chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;
- 20 (S)-(6-(3-Chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dihydrochloride;
- (R)-(6-(3-Chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;
- 25 [6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(6-fluoro-2,3-dihydro-1,4-benzodioxin-3-yl)methanone;
- (6-Chloro-2,3-dihydro-1,4-benzodioxin-3-yl)-[6-(3-chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]methanone;

[6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(2,3-dihydro-1,4-benzodioxin-3-yl)methanone;

[6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(6-fluorochroman-3-yl)methanone;

5 (S)-[6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(6-fluorochroman-3-yl)methanone;

(S)-[6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(6-fluorochroman-3-yl)methanone hydrochloride;

10 (S)-[6-(3-Chloro-1H-pyrazol-4-yl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(6-fluorochroman-3-yl)methanone dimethanesulfonic acid;;

(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(3-methylisoxazol-4-yl)indol-3-yl]methanone;

(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(2-methylpyrazol-3-yl)indol-3-yl]methanone;

15 [6-(2-Amino-4-pyridyl)-1-[2-(dimethylamino)ethyl]indol-3-yl]-(6-chlorochroman-3-yl)methanone;

[1-[2-(Dimethylamino)ethyl]-6-(3-methylisoxazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;

20 [1-[2-(Dimethylamino)ethyl]-6-(2-methylpyrazol-3-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;

(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]methanone;

(R)-(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]methanone;

25 (S)-(6-Chlorochroman-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]methanone;

- (R)-(6-(5-Chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-chlorochroman-3-yl)methanone;
- (S)-(6-(5-Chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-chlorochroman-3-yl)methanone;
- 5 [1-[2-(Dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;
- (S)-[1-[2-(Dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;
- (S)-[1-[2-(Dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone dihydrochloride;
- 10 (R)-[1-[2-(Dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-fluorochroman-3-yl)methanone;
- [1-[2-(Dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]-(6-methoxychroman-3-yl)methanone;
- 15 (6-Chloro-2,3-dihydro-1,4-benzodioxin-3-yl)-[1-[2-(dimethylamino)ethyl]-6-(5-methoxy-1H-pyrazol-4-yl)indol-3-yl]methanone;
- (2,3-Dihydrobenzo[b][1,4]dioxin-2-yl)(1-(2-(dimethylamino)ethyl)-6-(1H-pyrazol-4-yl)-1H-indazol-3-yl)methanone
- (S)-(2,3-Dihydrobenzo[b][1,4]dioxin-2-yl)(1-(2-(dimethylamino)ethyl)-6-(1H-pyrazol-4-yl)-1H-indazol-3-yl)methanone;
- 20 (6-Chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-methoxy-1H-pyrazol-4-yl)-1H-indazol-3-yl)methanone;
- (S)-(6-Chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-methoxy-1H-pyrazol-4-yl)-1H-indazol-3-yl)methanone;
- 25 (R)-(6-Chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-methoxy-1H-pyrazol-4-yl)-1H-indazol-3-yl)methanone;

- (6-Chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(3-fluoro-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)methanone;
- (1-(2-(Dimethylamino)ethyl)-6-(3-methoxy-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)(6-methoxychroman-3-yl)methanone;
- 5 (6-(3-Chloro-1*H*-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1*H*-indazol-3-yl)(6-methoxychroman-3-yl)methanone;
- (1-(2-(Dimethylamino)ethyl)-6-(3-methoxy-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)(6-fluorochroman-3-yl)methanone;
- 10 (6-(3-Chloro-1*H*-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1*H*-indazol-3-yl)(6-fluorochroman-3-yl)methanone;
- (*S*)-(6-(3-Chloro-1*H*-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1*H*-indazol-3-yl)(6-fluorochroman-3-yl)methanone;
- Chroman-3-yl(1-(2-(dimethylamino)ethyl)-6-(5-(trifluoromethyl)-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)methanone;
- 15 (6-(3-Fluoro-1*H*-pyrazol-4-yl)-1-(((*S*)-1-methylazetidin-2-yl)methyl)-1*H*-indazol-3-yl)(6-fluorochroman-3-yl)methanone;
- (6-(3-Fluoro-1*H*-pyrazol-4-yl)-1-(((*S*)-1-methylazetidin-2-yl)methyl)-1*H*-indazol-3-yl)((*S*)-6-fluorochroman-3-yl)methanone;
- (1-(2-(Dimethylamino)ethyl)-6-(3-fluoro-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)(6-fluorochroman-3-yl)methanone;
- 20 (7-Chloro-2,3-dihydrobenzo[*b*][1,4]dioxin-2-yl)(1-(2-(dimethylamino)ethyl)-6-(3-methoxy-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)methanone;
- (7-Chloro-2,3-dihydrobenzo[*b*][1,4]dioxin-2-yl)(1-(2-(dimethylamino)ethyl)-6-(3-fluoro-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)methanone;
- 25 (6-(3-Chloro-1*H*-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1*H*-indazol-3-yl)(7-chloro-2,3-dihydrobenzo[*b*][1,4]dioxin-2-yl)methanone;

- (1-(2-(Dimethylamino)ethyl)-6-(3-methoxy-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)(7-fluoro-2,3,4,5-tetrahydrobenzo[*b*]oxepin-4-yl)methanone;
- (6-(3-Chloro-1*H*-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1*H*-indazol-3-yl)(7-fluoro-2,3,4,5-tetrahydrobenzo[*b*]oxepin-4-yl)methanone;
- 5 (1-(2-(Dimethylamino)ethyl)-6-(3-methoxy-1*H*-pyrazol-4-yl)-1*H*-indazol-3-yl)(7-methoxy-2,3,4,5-tetrahydrobenzo[*b*]oxepin-4-yl)methanone;
- (6-(3-Chloro-1*H*-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1*H*-indazol-3-yl)(7-methoxy-2,3,4,5-tetrahydrobenzo[*b*]oxepin-4-yl)methanone;
- 10 (6-(5-chloro-1*H*-pyrazol-4-yl)-1-(((*R*)-1-methylazetidin-2-yl)methyl)-1*H*-indol-3-yl)((*S*)-6-methoxychroman-3-yl)methanone dihydrochloride;
- (6-(5-chloro-1*H*-pyrazol-4-yl)-1-(((*R*)-1-methylazetidin-2-yl)methyl)-1*H*-indol-3-yl)((*S*)-6-methoxychroman-3-yl)methanone dihydrobromide;
- (6-(5-chloro-1*H*-pyrazol-4-yl)-1-(((*R*)-1-methylazetidin-2-yl)methyl)-1*H*-indol-3-yl)((*S*)-6-methoxychroman-3-yl)methanone dimethanesulfonic acid;
- 15 (6-(5-chloro-1*H*-pyrazol-4-yl)-1-(((*R*)-1-ethylazetidin-2-yl)methyl)-1*H*-indol-3-yl)((*S*)-6-methoxychroman-3-yl)methanone;
- (6-(5-chloro-1*H*-pyrazol-4-yl)-1-(((*R*)-1-ethylazetidin-2-yl)methyl)-1*H*-indol-3-yl)((*S*)-6-methoxychroman-3-yl)methanone dihydrobromide;
- (6-(5-chloro-1*H*-pyrazol-4-yl)-1-(((*R*)-1-ethylazetidin-2-yl)methyl)-1*H*-indol-3-yl)((*S*)-6-methoxychroman-3-yl)methanone dimethanesulfonic acid;
- 20 ((*S*)-6-chlorochroman-3-yl)(1-((*R*)-2-(dimethylamino)propyl)-6-(5-fluoro-1*H*-pyrazol-4-yl)-1*H*-indol-3-yl)methanone;
- ((*S*)-6-chlorochroman-3-yl)(1-((*R*)-2-(dimethylamino)propyl)-6-(5-fluoro-1*H*-pyrazol-4-yl)-1*H*-indol-3-yl)methanone hydrobromide;

(S)-(6-chlorochroman-3-yl)(1-(2-(diethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone;

(S)-(6-chlorochroman-3-yl)(1-(2-(diethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone dihydrochloride;

5 (S)-(6-chlorochroman-3-yl)(1-(2-(diethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone dimethanesulfonic acid;

(S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(diethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;

10 (S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(diethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dihydrochloride;

(S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(diethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dihydrobromide

(S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(diethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dimethanesulfonic acid;

15 (S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(ethyl(methyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone

(S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(ethyl(methyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dihydrochloride;

20 (S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(ethyl(methyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dihydrobromide;

(S)-(6-(5-chloro-1H-pyrazol-4-yl)-1-(2-(ethyl(methyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dimethanesulfonic acid;

(S)-(6-(3-chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dihydrobromide;

(S)-(6-(3-chloro-1H-pyrazol-4-yl)-1-(2-(dimethylamino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone dimethanesulfonic acid;

(6-(3-chloro-1H-pyrazol-4-yl)-1-(((R)-1-(2-fluoroethyl)azetidin-2-yl)methyl)-1H-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone;

5 (6-(3-chloro-1H-pyrazol-4-yl)-1-(((R)-1-(2,2-difluoroethyl)azetidin-2-yl)methyl)-1H-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone;

(6-(3-chloro-1H-pyrazol-4-yl)-1-(((R)-1-(2,2,2-trifluoroethyl)azetidin-2-yl)methyl)-1H-indol-3-yl)((S)-6-methoxychroman-3-yl)methanone;

10 (S)-(6-(3-chloro-1H-pyrazol-4-yl)-1-(2-(ethyl(2-fluoroethyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;

(S)-(6-(3-chloro-1H-pyrazol-4-yl)-1-(2-((2,2-difluoroethyl)(ethyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;

(S)-(6-(3-chloro-1H-pyrazol-4-yl)-1-(2-(ethyl(2,2,2-trifluoroethyl)amino)ethyl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;

15 (S)-(1-(2-(bis(2-fluoroethyl)amino)ethyl)-6-(3-chloro-1H-pyrazol-4-yl)-1H-indol-3-yl)(6-methoxychroman-3-yl)methanone;

(S)-(6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone hydrochloride;

20 (S)-(6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone dihydrobromide;

(S)-(6-chlorochroman-3-yl)(1-(2-(dimethylamino)ethyl)-6-(5-methoxy-1H-pyrazol-4-yl)-1H-indol-3-yl)methanone dimethanesulfonic acid;

(6-(3-fluoro-1H-pyrazol-4-yl)-1-(((S)-1-methylazetidin-2-yl)methyl)-1H-indazol-3-yl)((S)-6-fluorochroman-3-yl)methanone dihydrobromide.

9. Use of the compound, salt, or prodrug of claim 1 for preparing a medicament for treating a disease that can be treated by inhibition of a ROCK enzyme.
10. Use of the compound, salt, or prodrug of claim 1 for treating or alleviating a fibrotic disorder, a cardiovascular disease or disorder, an inflammatory disease or disorder, a
5 neurological disease or disorder, or a proliferative disease or disorder.
11. The use of claim 10, wherein the fibrotic disorder is pulmonary fibrosis selected from the group consisting of cystic and idiopathic pulmonary fibrosis, radiation induced lung injury, liver fibrosis including cirrhosis, cardiac fibrosis including arterial fibrosis, endomyocardial fibrosis,
10 old myocardial infarction, arterial stiffness, atherosclerosis, restenosis, arthrofibrosis, Crohn's disease, myelofibrosis, Peyronie's diseases, nephrogenic systemic fibrosis, progressive massive fibrosis, retroperitoneal cavity fibrosis, scleroderma/systemic sclerosis, mediastinal fibrosis, Keloids and hypertrophic scars, glial scarring, and renal fibrosis.
12. The use of claim 10, wherein the cardiovascular disease or disorder is selected from the
15 group consisting of angina, atherosclerosis, cerebral vasospasm, cerebrovascular contraction, coronary vasospasm, endothelial dysfunction, erectile dysfunction, glaucoma, hypertension, ischemic/reperfusion injury, myocardial hypertrophy, myocardial infarction, peripheral circulation disorder, pre-term labor, Raynaud's Disease, renal disease, and stroke.
13. The use of claim 10, wherein the proliferative disease or disorder is an invasive or
20 metastatic cancer selected from the group consisting of adenocarcinoma, adrenocortical cancer, bladder cancer, bone cancer, brain cancer, breast cancer, cancer of the buccal cavity, cervical cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, epidermoid carcinoma, esophageal cancer, eye cancer, follicular carcinoma, gallbladder cancer, gastrointestinal cancer, cancer of the genitourinary tract, glioblastoma, hairy cell carcinoma,
25 head and neck cancer, hepatic carcinoma, hepatocellular cancer, Hodgkin's disease, keratoacanthoma, kidney cancer, large cell carcinoma, cancer of the large intestine, laryngeal cancer, liver cancer, adenocarcinoma of the lung, small-cell lung cancer, squamous carcinoma of the lung, non-small cell lung cancer, melanoma, a myeloproliferative disorder, neuroblastoma,

ovarian cancer, papillary carcinoma, pancreatic cancer, cancer of the peritoneum, prostate cancer, rectal cancer, salivary gland carcinoma, sarcoma, squamous cell cancer, small cell carcinoma, cancer of the small intestine, stomach cancer, testicular cancer, thyroid cancer, and vulval cancer.

- 5 14. The use of claim 10, wherein the neurological disease or disorder is selected from the group consisting of Huntington's disease, Parkinson's Disease, Alzheimer's, Amyotrophic lateral sclerosis (ALS), Batten disease, dementia, spinal muscular atrophy, motor neuron diseases, cerebral cavernous malformations (CCMs), spinocerebellar ataxia, acute or chronic pain, dementia, neuronal degeneration, spinal cord injury, cerebral vasospasm, and multiple sclerosis.

10