

METHODS AND SYSTEMS FOR ASSESSMENT OF CLINICAL INFERTILITY

CROSS-REFERENCE

- [0001] This application claims the benefit of U.S. Provisional Application Nos. 61/077,439, filed July 1, 2008, and 61/081,596, filed July 17, 2008, which applications are incorporated by reference herein in their entirety.

GOVERNMENT RIGHTS

- [0002] This invention was made with government support under federal grant nos. R01 GM067250 and R01 HD057970 awarded by the National Institutes of Health. The United States Government has certain rights in this invention.

BACKGROUND OF THE INVENTION

- [0003] Reproductive failure is a serious problem that has been addressed clinically by various assisted reproductive technologies, including *in vitro* fertilization (IVF) and embryo transfer (ET). These procedures might be expected to yield exceptionally high conception rates as *in vitro* fertilization provides embryos that appear normal at a morphological level for transfer into a fully primed recipient. Despite these efforts the success rate of IVF/ET is less than ideal. In the published data for IVF/ET in the United States and Canada in 1994, there were 26,961 initiated cycles of standard IVF. Of these, 86.2% led to retrieval and of these 90.2% led to a transfer. However, the overall success rate in terms of clinical pregnancies was 22.7% per initiated cycle and a 29.1% pregnancy rate per transfer.
- [0004] Additionally, there appears to be a high incidence of early pregnancy loss after *in vitro* fertilization with a biochemical pregnancy rate of 18% and a spontaneous abortion rate of 27%. Thus, it appears that the IVF technique has been well optimized but implantation failure may be the cause for a large number of losses with ET and this peri-implantational loss is an area of potential improvement. A major factor in the success rate of various assisted reproductive technologies is endometrial receptivity, a transient state that must be coordinated with embryo development to implantation-competent blastocysts.

[0005] IVF is an expensive procedure and can be psychologically traumatic for a patient. Surgical procedures are required to collect eggs from a female for IVF and, following fertilization, further surgery is required to implant the fertilized eggs in the womb. The recipient must then wait for a period of time before it can be determined whether or not pregnancy has been established. In some cases, pregnancy may never be achieved despite repeated attempts, and these cases can represent a considerable expense to the patient and society, both in financial and human terms.

[0006] Therefore, until success rates of IVF can be improved, it would be desirable to be able to identify recipients for whom IVF is unlikely to be successful prior to treatment, so that such patients may avoid the above mentioned costs and trauma of the IVF procedure.

[0007] The present invention addresses these needs.

SUMMARY OF THE INVENTION

[0008] Methods and computer-based systems for facilitating assessment of clinical infertility are provided. The methods and systems can be implemented to, for example, facilitate assessment of a subject for an in vitro fertilization treatment cycle, including determining probability of a live birth event. The methods and systems can be implemented to, for example, facilitate a determination of success of implantation of embryos, selection of an optimal number of embryos to transfer, and determination of success in subsequent in vitro fertilization treatment cycles following an unsuccessful treatment cycle.

[0009] These and other objects, advantages, and features of the invention will become apparent to those persons skilled in the art upon reading the details of the invention as more fully described below.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010] The invention is best understood from the following detailed description when read in conjunction with the accompanying drawings. It is emphasized that, according to common practice, the various features of the drawings are not to-scale.

On the contrary, the dimensions of the various features are arbitrarily expanded or reduced for clarity. Included in the drawings are the following figures.

- [0011] **FIG 1** shows the source of the data. **Panel A** shows the IVF cycles performed in 2005. **Panel B** shows the utilization of oocytes and embryos in 665 fresh, non-donor IVF cycles. All numbers in Panel A indicate the number of cycles and numbers in Panel B indicate the number of oocytes or embryos. Fresh cycles are defined by ovarian stimulation of gonadotropins and embryo transfer performed within the same cycle; cryopreserved cycles utilize embryos that were obtained and cryopreserved from a previous cycle; “freeze-all” are cycles in which ovarian stimulation was performed, but embryos were cryopreserved instead of being transferred back within the same cycle for medical or non-medical reasons. 157 cycles were removed from analysis for a variety of medical and non-medical reasons that did not result in fresh embryo transfer.
- [0012] **FIG 2** shows the distribution of all embryos from 665 fresh, non-donor IVF cases according to their cell number on Day 3.
- [0013] **FIG 3** shows variables and their relative importance in determining A) number of 8-cell embryos (**Panel A**), day 3 FSH (**Panel B**), and the total number of embryos (**Panel C**).
- [0014] **FIG 4** shows that translational block of cyclin A2 by *Ccna2*-MO causes embryos to arrest at the 2-cell stage. **Panel A** shows *Ccna2* expression in 1- to 2-cell embryos by RT-PCR. **Panel B** shows nuclear cyclin A2 localization was absent in $83.4 \pm 6.0\%$ of *Ccna2*-MO-injected embryos but present in all uninjected embryos; $p < 0.01$). **Panel C** shows that ~50kD cyclin A2 protein was not detected in *Ccna2*-MO-injected embryos.
- [0015] **FIG 5, panel A** shows that *Ccna2*-MO induced higher rates of 2-cell stage arrest compared to controls ($p < 0.01$). **Panel B** shows that only $1.8 \pm 1.8\%$ of *Ccna2*-MO-injected embryos reached blastocyst stage ($p = 0.06$ compared to *Ccna2*-MM). **Panel C** shows that the rates of 2-cell stage arrest decreased with the concentration of *Ccna2*-MO ($p = 0.05$ for 0.5 mM; $p = 0.01$ for 0.25 mM). **Panel D** shows that the rate of blastocyst development at were higher at 0.25 mM than at 0.75 mM ($p < 0.001$). All columns and error bars represent mean $\pm$ s.e.m., respectively, from at least three independent sets of experiments. Scale bar 40 μ m.

See Table 1 for the targeted sequence of all MOs used, and Table 7 for the total number of embryos tested in each set of experiments.

[0016] **FIG 6** shows the mechanisms that regulate mammalian embryo development at the earliest stages. In wild type (+/-) embryos, maternal transcripts are present before embryonic genome activation (EGA), while maternal and embryonic transcripts are present at maternal-embryonic transition stage, both resulting in production of a normal gene product (**Panel A**). In homozygous null mutant (-/-) embryos generated from a mother that is heterozygous (+/-) for the null mutation, persistent maternal transcripts and/or proteins may “rescue” or delay the phenotype onset (**Panel B**). In contrast, homozygous null mutant embryos generated from a homozygous mutant female, or a female with oocyte-specific gene deletion, the observed defects may reflect oocyte defects, rather than specific gene requirement in the early embryo (**Panel C**). Therefore, these strategies do not address the precise roles of specific genes at the cusp of EGA or during EGA, when both maternal and early embryonic transcripts may be present simultaneously. Cytoplasmic microinjection of antisense morpholino oligonucleotides (MOs) into wild type embryo just at or before EGA results in specific translational block of both maternal and embryonic gene transcripts (**Panel D**). Since MOs persist for at least a few cycles of cell division, gene-specific translational block is presumably effective until the morula-blastocyst stages. The absence of gene product during these developmental stages would reveal critical gene function and unmask early phenotypes that may not be detectable in conventional gene-targeting strategies by homologous recombination and transgenesis. While this model is well established in other species, it shifts the paradigm from investigating function of embryonic genes to that of gene products regardless of their maternal or embryonic origin.

[0017] **FIG 7** shows the experimental strategy. Embryos at the 2 pronuclei (2PN) or 1-cell stage were collected from wild type matings, and injected with an antisense morpholino oligomer (MO) that was designed to target a specific gene. MO binds to 5' UTR or transcription start site and blocks translation by steric hindrance. Microinjected embryos and uninjected control embryos were cultured *in vitro* and observed for developmental phenotypes such as fragmentation, or arrest at the 2-cell, 4-cell, multicell, or morula stages. If a gene-specific MO produces the

same phenotype consistently, while the mismatch control MO allows normal development, then knockdown of the gene of interest was validated by immunocytochemistry and/or immunoblotting. Mechanism of gene function was further investigated by obtaining global gene expression profiles from injected and control embryos at the mid-2-cell stage (43 hours post-HCG). Candidate downstream genes were tested for differential expression, and gene function in the early embryo.

[0018] **FIG 8** shows Oct4 expression in mouse zygote by single embryo RT-PCR.

[0019] **FIG 9** shows that Oct4 is required for early embryo development prior to formation of the blastocyst. **Panel A** shows that *Oct4*-MO-injected embryos arrested at multicell stage, while controls reached blastocyst stage. **Panel B** shows that *Oct4* knockdown induced higher arrest rates at the 1- to multicell stages; $p < 0.01$. **Panel C** shows that None of the *Oct4*-MO-injected embryos developed to blastocysts. **Panel D** shows that the rates of arrest at the 1- to multicell stages decreased with concentration of *Oct4*-MO ($p < 0.01$). **Panel E** shows that there was a non-significant trend for higher rates of blastocyst development with decreasing concentrations of *Oct4*-MO.

[0020] **FIG 10** shows that decreased Oct4 expression was evident by the 4-cell stage in *Oct4*-MO-injected embryos. Oct4 signal was absent in embryos injected with *Oct4*-MO, but its nuclear localization was present in uninjected and mismatch controls. Scale bar 40 μm .

[0021] **FIG 11, Panel A**, shows that nuclear Oct4 expression is absent in *Oct4*-MO-injected embryos (top panel) but present in *Oct4*-MM-injected embryos and uninjected. **Panel B** shows that compared to no coinjection or mEYFP mRNA coinjection, coinjection of 36 or 3.6 ng/ μL of *Oct4* mRNA resulted in partial rescue of the *Oct4*-MO-induced phenotype by specifically decreasing the arrest rates at the 1- to multicell stages, which resulted in higher rates blastocyst development ($p < 0.01$). **Panel C** shows that overexpression of *Oct4* mRNA induced higher developmental arrest in a dose-dependent manner, such that blastocyst rates are significantly lower after injection of Oct4 mRNA at 36 or 90 ng/ μL , compared to overexpression of mEYFP ($p < 0.01$). Gene overexpression was validated by q-PCR or immunocytochemistry (data not shown). (Scale bar = 40 μm).

- [0022] **FIG 12** shows decreased Oct4 expression at the multicell stage in *Oct4*-MO-injected embryos. Only $6.4 \pm 3.2\%$ of *Oct4*-MO-injected embryos showed nuclear Oct4 signal, while $88.9 \pm 11.1\%$ of *Oct4*-MM-injected embryos and $82.7 \pm 10.9\%$ of uninjected control embryos showed unequivocal nuclear Oct4 expression at the multicell stage; $p < 0.05$.
- [0023] **FIG 13** shows confirmation of the requirement of Oct4 in early embryo development by *Oct4*E4-MO, an antisense morpholino that targets the splice site of exon 4 of *Oct4*. **Panel A** shows sites targeted by the two morpholinos, *Oct4*-MO and *Oct4*E4-MO. *Oct4*-MO targets the 25 nucleotides starting at the ATG start site, while *Oct4*E4-MO targets the splice site at the intron (I)-exon (E) boundary of the 4th exon (E4). Removal of E4 is expected to result in a protein product that lacks the DNA-binding and activation domains. **Panel B** shows that $64.6 \pm 19.9\%$ of embryos injected with *Oct4*E4-MO, while none that were injected with the mismatch control, *Oct4*E4-MM, arrested at the 2-cell stage. **Panel C** shows that blastocyst development is severely compromised after injection of *Oct4*E4-MO compared to the mismatch control, *Oct4*E4-MM.
- [0024] **FIGS 14 and 15** show gene regulation by Oct4. **FIG 14, Panel A** shows unsupervised clustering of 3 *Oct4* knockdown, 3 *Ccna2* knockdown, and 6 uninjected (NI) pooled embryo samples, and increased (red) and decreased (blue) gene expression. Scale is standard deviation **FIG 14, Panel B** shows intersection of differentially expressed genes in *Oct4* and *Ccna2* knockdown embryos. **FIG 14, Panel C** and **FIG 15** show relative expression of *Oct4* knockdown by single-embryo q-PCR for **FIG 14, Panel C** overexpressed genes ($p < 0.05$ for *Hes5*) and **FIG 15** downregulated genes ($*p < 0.001$, $**p < 0.05$, $***p < 0.1$). TR translational repression. Error bars indicate s.e.m.
- [0025] **FIG 16** shows models of Oct4 function at the 1- to 2-cell stage. **Panel A** shows genes whose Absolute (ΔC_T) (red) is greater than its s.e.m. (blue), are ordered clockwise based on increasing s.e.m. (Log scale.) **Panel B** shows proposed regulation of various modules by Oct4 via transcriptional and post-transcriptional mechanisms.
- [0026] **FIG 17** shows a summary of the data source for the live birth analysis. The numbers in the boxes indicate the number of fresh cycles performed in the Stanford

IVF center through 2003-06. The cycles that were not included were related to: gestational carriers, Cryo. ET or cryopreserved embryo transfer (cycles which utilized frozen embryos, irrespective of whether the cycle was performed at Stanford or outside), donor oocytes (non-self oocytes), recipient cycles and cycles in which no gonadotropin was injected. 9 cycles were excluded from analysis for duplicated entries and unknown outcomes. The cycles analyzed were further classified into Cycles 1, 2, 3 or beyond, depending on the number of times the patient returned for a fresh IVF cycle.

[0027] **FIG 18** shows a summary of live birth outcomes. Different boxes represented C1 (first-IVF cycle), C2 (second-IVF cycle), and C3 (third-IVF cycle). Each box shows the number of cycles with live birth (LB+) or no live birth (LB-) outcomes, the percentage of the total number (N) of cycles analyzed and the number of patients that dropped out and did not return.

[0028] **FIG 19** is a graph showing distribution of embryos with respect to developmental stage. The distribution of 2969 fresh, non-donor IVF cases from all three cycles according to their cell number on day 3 of *in vitro* culture. The number of cells per embryo and the maximum number of embryos (~40%) that had 8 cells were similar amongst cycles C1 (no fill pattern), C2 (horizontal fill pattern), and C3 (vertical fill pattern).

[0029] **FIG 20** shows prediction models and their temporal relationship with a typical IVF cycle. An IVF cycle comprises of down-regulation of endogenous gonadotropins and the menstrual cycle using oral contraceptive pill (OCP), periodic FSH injections and cycle monitoring by ultrasound with or without serum estradiol, followed by hCG injection to simulate endogenous LH surge, oocyte retrieval, fertilization by IVF or ICSI, *in vitro* embryo culture, embryo transfer, and cryopreservation of excess embryos. The three black stars represent the points at which patients can take an informed decision about further treatment options and whether to continue treatment. The Pre-IVF model extends until the treatment start date. The Pre-OR (oocyte retrieval) model extends until the hCG injection. The Post-IVF model includes all variables until embryo transfer.

[0030] **FIGS 21A, 21B, and 21C** show tree models for three analysis groups: Pre-IVF (21A), Pre-OR (21B) and Post-IVF (21C). The pie-charts show the percentage

of live births (LB) among the total number (N) of cycles analyzed. Terminal populations are labeled as population 1 (Pop 1), etc., with its live birth rate indicated.

- [0031] **FIG 22** shows inclusion and exclusion criteria for 2007 data set.
- [0032] **FIG 23** shows that Pre-IVF, Pre-OR and Post-IVF models identify distinct populations with differential live birth rates.
- [0033] **FIG 24A, 24B, and 24C** show comparison of live birth rates between Pre-IVF (Fig. 23A), Pre-OR- (Fig. 23 B), Post-IVF models and an age-based control model.
- [0034] **FIG 25** shows that Post-IVF model C1 population assignment predicts differential live birth rates in C2 and cumulative live birth rates in subsequent cycles (C2 and C3).
- [0035] **FIG 26** shows results of gene chip experiments performed at the 1-cell stage.
- [0036] **FIG 27** shows results of gene chip experiments performed at the 4-cell stage.
- [0037] **FIG 28** shows results of gene chip experiments performed at the 8-cell stage.
- [0038] **FIG 29** shows a correlation of two gene chip experiments showing expression of a panel of genes at various cell stages.
- [0039] **FIG 30** shows a correlation of two gene chip experiments showing expression of a panel of genes at various cell stages.
- [0040] Before the present embodiments are described, it is to be understood that this invention is not limited to particular embodiments described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.
- [0041] Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limits of that range is also specifically disclosed. Each smaller range between any stated value or intervening value in a

stated range and any other stated or intervening value in that stated range is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included or excluded in the range, and each range where either, neither or both limits are included in the smaller ranges is also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in the invention.

[0042] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, some potential and preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited. It is understood that the present disclosure supercedes any disclosure of an incorporated publication to the extent there is a contradiction.

[0043] It must be noted that as used herein and in the appended claims, the singular forms "a", "an", and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a cell" includes a plurality of such cells and reference to "the compound" includes reference to one or more compounds and equivalents thereof known to those skilled in the art, and so forth.

[0044] It is further noted that the claims may be drafted to exclude any optional element. As such, this statement is intended to serve as antecedent basis for use of such exclusive terminology as "solely", "only" and the like in connection with the recitation of claim elements, or the use of a "negative" limitation.

[0045] The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention. Further, the dates of publication provided may be different from the actual publication dates which may need to be independently confirmed.

DEFINITIONS

- [0046] The terms “subject,” “individual,” and “patient” are used interchangeably herein to refer to a mammal being assessed for treatment and/or being treated. In an embodiment, the mammal is a human, such as a female human. The terms “subject,” “individual,” and “patient” thus encompass individuals in need of assessment of clinical infertility, including those who have undergone or are candidates for an in vitro fertilization cycle.
- [0047] As used herein, the term “correlates,” or “correlates with,” and like terms, refers to a statistical association between instances of two events, where events include numbers, data sets, and the like. For example, when the events involve numbers, a positive correlation (also referred to herein as a “direct correlation”) means that as one increases, the other increases as well. A negative correlation (also referred to herein as an “inverse correlation”) means that as one increases, the other decreases.
- [0048] A “biological sample” encompasses a variety of sample types obtained from an individual. The definition encompasses blood and other liquid samples of biological origin, solid tissue samples such as a biopsy specimen or tissue cultures or cells derived therefrom and the progeny thereof. The definition also includes samples that have been manipulated in any way after their procurement, such as by treatment with reagents; washed; or enrichment for certain cell populations, such as cancer cells. The definition also includes sample that have been enriched for particular types of molecules, e.g., nucleic acids, polypeptides, etc. The term “biological sample” encompasses a clinical sample, and also includes tissue obtained by surgical resection, tissue obtained by biopsy, cells in culture, cell supernatants, cell lysates, tissue samples, organs, bone marrow, blood, plasma, serum, and the like. A “biological sample” includes a sample obtained from a patient’s uterus, including embryo cultures.
- [0049] The terms “gene product” and “expression product” are used interchangeably herein in reference to a gene, to refer to the RNA transcription products (transcripts) of the gene, including mRNA and the polypeptide translation products of such RNA transcripts, whether such product is modified post-translationally or not. The terms “gene product” and “expression product” are used

interchangeably herein, in reference to an RNA, particularly an mRNA, to refer to the polypeptide translation products of such RNA, whether such product is modified post-translationally or not. A gene product can be, for example, an unspliced RNA, an mRNA, a splice variant mRNA, a polypeptide, a post-translationally modified polypeptide, a splice variant polypeptide, etc.

- [0050] As used herein, the term “normalized expression level” refers to an expression level of a response indicator gene relative to the level of an expression product of a reference gene(s).
- [0051] As used herein, the terms “label” and “detectable label” refer to a molecule capable of being detected, where such molecules include, but are not limited to, radioactive isotopes, fluorescers (fluorophores), chemiluminescers, chromophores, enzymes, enzyme substrates, enzyme cofactors, enzyme inhibitors, chromophores, dyes, metal ions, metal sols, ligands (e.g., biotin, avidin, strepavidin or haptens), intercalating dyes and the like. The term “fluorescer” or “fluorophore” refers to a substance or a portion thereof which is capable of exhibiting fluorescence in a detectable range.
- [0052] As used herein, the term “target nucleic acid region” or “target nucleic acid” refers to a nucleic acid with a “target sequence” to be detected (e.g., in a method involving nucleic acid hybridization and/or amplification). The target nucleic acid may be either single-stranded or double-stranded and may or may not include other sequences besides the target sequence (e.g., the target nucleic acid may or may not include nucleic acid sequences upstream or 5' flanking sequence, and may or may not include downstream or 3' flanking sequence. Where detection is by amplification, these other sequences in addition to the target sequence may or may not be amplified with the target sequence.
- [0053] The term “primer” or “oligonucleotide primer” as used herein, refers to an oligonucleotide which acts to initiate synthesis of a complementary nucleic acid strand when placed under conditions in which synthesis of a primer extension product is induced, e.g., in the presence of nucleotides and a polymerization-inducing agent such as a DNA or RNA polymerase and at suitable temperature, pH, metal ion concentration, and salt concentration. Primers are generally of a length compatible with their use in synthesis of primer extension products, and can be in

the range of between about 8 nucleotides and about 100 nucleotides (nt) in length, such as about 10 nt to about 75 nt, about 15 nt to about 60 nt, about 15 nt to about 40 nt, about 18 nt to about 30 nt, about 20 nt to about 40 nt, about 21 nt to about 50 nt, about 22 nt to about 45 nt, about 25 nt to about 40 nt, and so on, e.g., in the range of between about 18 nt and about 40 nt, between about 20 nt and about 35 nt, between about 21 and about 30 nt in length, inclusive, and any length between the stated ranges. Primers can be in the range of between about 10-50 nucleotides long, such as about 15-45, about 18-40, about 20-30, about 21-25 nt and so on, and any length between the stated ranges. In some embodiments, the primers are not more than about 10, 12, 15, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 55, 60, 65, or 70 nucleotides in length. In this context, the term “about” may be construed to mean 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 more nucleotides either 5' or 3' from either termini or from both termini.

[0054] Primers are in many embodiments single-stranded for maximum efficiency in amplification, but may alternatively be double-stranded. If double-stranded, the primer is in many embodiments first treated to separate its strands before being used to prepare extension products. This denaturation step is typically effected by heat, but may alternatively be carried out using alkali, followed by neutralization. Thus, a “primer” is complementary to a template, and complexes by hydrogen bonding or hybridization with the template to give a primer/template complex for initiation of synthesis by a polymerase, which is extended by the covalent addition of bases at its 3' end.

[0055] A “primer pair” as used herein refers to first and second primers having nucleic acid sequence suitable for nucleic acid-based amplification of a target nucleic acid. Such primer pairs generally include a first primer having a sequence that is the same or similar to that of a first portion of a target nucleic acid, and a second primer having a sequence that is complementary to a second portion of a target nucleic acid to provide for amplification of the target nucleic acid or a fragment thereof. Reference to “first” and “second” primers herein is arbitrary, unless specifically indicated otherwise. For example, the first primer can be designed as a “forward primer” (which initiates nucleic acid synthesis from a 5' end of the target nucleic acid) or as a “reverse primer” (which initiates nucleic acid

synthesis from a 5' end of the extension product produced from synthesis initiated from the forward primer). Likewise, the second primer can be designed as a forward primer or a reverse primer.

[0056] As used herein, the term “probe” or “oligonucleotide probe”, used interchangeably herein, refers to a structure comprised of a polynucleotide, as defined above, which contains a nucleic acid sequence complementary to a nucleic acid sequence present in the target nucleic acid analyte (e.g., a nucleic acid amplification product). The polynucleotide regions of probes may be composed of DNA, and/or RNA, and/or synthetic nucleotide analogs. Probes are generally of a length compatible with their use in specific detection of all or a portion of a target sequence of a target nucleic acid, and are in many embodiments in the range of between about 8 nt and about 100 nt in length, such as about 8 to about 75 nt, about 10 to about 74 nt, about 12 to about 72 nt, about 15 to about 60 nt, about 15 to about 40 nt, about 18 to about 30 nt, about 20 to about 40 nt, about 21 to about 50 nt, about 22 to about 45 nt, about 25 to about 40 nt in length, and so on, e.g., in the range of between about 18-40 nt, about 20-35 nt, or about 21-30 nt in length, and any length between the stated ranges. In some embodiments, a probe is in the range of between about 10-50 nucleotides long, such as about 15-45, about 18-40, about 20-30, about 21-28, about 22-25 and so on, and any length between the stated ranges. In some embodiments, the primers are not more than about 10, 12, 15, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 55, 60, 65, or 70 nucleotides in length. In this context, the term “about” may be construed to mean 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 more nucleotides either 5' or 3' from either termini or from both termini.

[0057] Where a nucleic acid is said to hybridize to a recited nucleic acid sequence, hybridization is under stringent conditions. An example of stringent hybridization conditions is hybridization at 50°C or higher and 0.1×SSC (15 mM sodium chloride/1.5 mM sodium citrate). Another example of stringent hybridization conditions is overnight incubation at 42°C in a solution: 50 % formamide, 5 × SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5 × Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1 × SSC at about 65°C. Stringent

hybridization conditions are hybridization conditions that are at least as stringent as the above representative conditions, where conditions are considered to be at least as stringent if they are at least about 80% as stringent, e.g., at least about 90% as stringent as the above specific stringent conditions.

DETAILED DESCRIPTION OF THE INVENTION

[0058] As noted above, methods and computer-based systems for facilitating assessment of clinical infertility are provided. The methods and systems can be implemented to, for example, facilitate assessment of a subject for an in vitro fertilization treatment cycle. The methods and systems can be implemented to, for example, facilitate a determination of success for implantation of embryos, selection of an optimal number of embryos to transfer, and determination of success in subsequent in vitro fertilization treatment cycles following an unsuccessful treatment cycle.

[0059] In certain embodiments, the method includes obtaining items of information from a female subject to provide a profile for the female subject, wherein each item of information relate to preselected patient variables, comparing the profile for the female subject to a library of known profile patterns known to be indicative of responsiveness to an in vitro fertilization procedure using an algorithm based upon the preselected patient variables, wherein the comparing provides an evaluation of the female subject for an in vitro fertilization procedure. In certain embodiments, the in vitro fertilization procedure is at least a second vitro fertilization procedure for said female subject.

[0060] The items of information may be provided by the female subject based on a written or electronic questionnaire or may be requested, transcribed, or otherwise logged, by a health care practitioner, such as a doctor, nurse, technician, or the like, during or concurrent with a medical evaluation that may optionally be associated with a determination to undergo a first or subsequent in vitro fertilization cycle.

[0061] Exemplary items of information relating to preselected patient variables include, but are not limited to: patient characteristics, such as age, previous infertility history, clinical diagnosis; clinical treatment information, such as type of

medication, number of days of stimulation, number of oocytes, etc.; conventional embryo morphology data, such as number of embryos, developmental stage, grade, and the like. In some embodiments, the information includes age, total number of embryos, rate of cleavage arrest, number of 8-cell embryos, day 3 follicle stimulating hormone (FSH) level, and number of 8-cell embryos transferred.

[0062] In certain embodiments, the in vitro fertilization procedure provides for a live birth event following the in vitro fertilization procedure. In such embodiments, the method provides a probability of a live birth event occurring resulting from the first or subsequent in vitro fertilization cycle based at least in part on items of information from the female subjects.

[0063] In some embodiments, the female subject is a pre-in vitro fertilization (pre-IVF) procedure patient. In certain embodiments, the items of information relating to preselected patient variables for determining the probability of a live birth event for a pre-IVF procedure patient may include age, diminished ovarian reserve, 3 follicle stimulating hormone (FSH) level, body mass index, polycystic ovarian disease, season, unexplained female infertility, number of spontaneous miscarriages, year, other causes of female infertility, number of previous pregnancies, number of previous term deliveries, endometriosis, tubal disease, tubal ligation, male infertility, uterine fibroids, hydrosalpinx, and male infertility causes.

[0064] In some embodiments, the female subject is a pre-surgical (pre-OR) procedure patient (pre-OR is also referred to herein as pre-oocyte retrieval). In certain embodiments, the items of information relating to preselected patient variables for determining the probability of a live birth event for a pre-OR procedure patient may include age, endometrial thickness, total number of oocytes, total amount of gonatropins administered, number of total motile sperm after wash, number of total motile sperm before wash, day 3 follicle stimulating hormone (FSH) level, body mass index, sperm collection, age of spouse, season number of spontaneous miscarriages, unexplained female infertility, number of previous term deliveries, year, number of previous pregnancies, other causes of female infertility, endometriosis, male infertility, tubal ligation, polycystic ovarian disease, tubal disease, sperm from donor, hydrosalpinx, uterine fibroids, and male infertility causes.

[0065] In some embodiments, the female subject is a post-in vitro fertilization (post-IVF) procedure patient. In certain embodiments, the items of information relating to preselected patient variables for determining the probability of a live birth event for a post-IVF procedure patient may include blastocyst development rate, total number of embryos, total amount of gonatropins administered, endometrial thickness, flare protocol, average number of cells per embryo, type of catheter used, percentage of 8-cell embryos transferred, day 3 follicle stimulating hormone (FSH) level, body mass index, number of motile sperm before wash, number of motile sperm after wash, average grade of embryos, day of embryo transfer, season, number of spontaneous miscarriages, number of previous term deliveries, oral contraceptive pills, sperm collection, percent of unfertilized eggs, number of embryos arrested at 4-cell stage, compaction on day 3 after transfer, percent of normal fertilization, percent of abnormally fertilized eggs, percent of normal and mature oocytes, number of previous pregnancies, year, polycystic ovarian disease, unexplained female infertility, tubal disease, male infertility only, male infertility causes, endometriosis, other causes of female infertility, uterine fibroids, tubal ligation, sperm from donor, hydrosalpinx, performance of ICSI, or assisted hatching.

[0066] Additional examples of parameters are provided in the examples section, including, for example, Tables 13 and 15.

[0067] In certain embodiments, the method includes obtaining items of information relating to at least two preselected patient variables, or more. As such, in other embodiments, the method includes obtaining items of information relating to at 3 or more preselected patient variables, including 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 12 or more, 15 or more, 17 or more, 20 or more, and the like.

[0068] In certain embodiments, the method includes assigning a weighted relative importance to each preselected patient variable in relation to other preselected patient variables. For example, in an analysis of a Pre-IVF model subject, the preselected patient variables of age and diminished ovarian reserve are given a higher relative importance value over other preselected patient variables, such as, for example, body mass index, tubular disease, and endometriosis. In such

embodiments, the sum of the relative importance of each preselected patient variables will equal 100.

[0069] In another embodiment, in an analysis of a Pre-OR model subject, the preselected patient variables of total amount of gonatropins, endometrial thickness and age are given a higher relative importance value over other preselected patient variables, such as, for example, number of previous pregnancies, tubal ligation and number of spontaneous miscarriages. In such embodiments, the sum of the relative importance of each preselected patient variables will equal 100.

[0070] In yet another embodiment, in an analysis of a Post-IVF model subject, the preselected patient variables of blastocyst development rate, total number of embryos, total amount of gonatropins administered, and endometrial thickness are given a higher relative importance value over other preselected patient variables, such as, for example, body mass index, number of motile sperm before wash, number of motile sperm after wash, average grade of embryos, and day of embryo transfer. In such embodiments, the sum of the relative importance of each preselected patient variables will equal 100.

[0071] In some embodiments, the comparing includes applying a decision rule. In some embodiments, the data analysis algorithm comprises the use of a classification tree. In other embodiments, the data analysis algorithm is nonparametric, such as using a Wilcoxon Signed Rank Test. In certain embodiments, the data analysis algorithm detects differences in a distribution of feature values. In some embodiments, the data analysis algorithm includes using a multiple additive regression tree. In some embodiments, the data analysis algorithm is a logistic regression.

[0072] In further embodiments, the method includes assessment of a gene expression profile of an arrested embryo from the female subject. For example, embryos that have arrested, for example embryos that have fewer than about 5 cells on day 3 following in vitro fertilization, are tested for a relative expression level of a panel of genes that are critical to embryo development. The gene expression profile is then compared to a control gene expression profile.

[0073] Any gene that for which normalized expression level is correlated (either positively or negatively) with infertility or likelihood of success or failure of an in

in vitro fertilization cycle is suitable for use with the methods of the invention. Exemplary genes include, but are not limited to, Oct4, Eif3c, Papola, Piwil2, Eif3b, Eif4b, Rbm3, Cpsf4 and other genes found to be down- or upregulated upon Oct4 knockdown, or the knockdown of another gene encoding a pluripotency regulator (e.g. Sall4). Other exemplary genes include those listed in Tables 8A, 8B, 9A, and 9B in attached Appendixes A-D, as well as Table 12. These gene products are referred to herein as “infertility indicator gene products”; and genes encoding the response indicator gene products are referred to as “infertility indicator genes.” Normalized expression levels of one or more of these infertility indicator genes can be determined to assess a female patient for an in vitro fertilization treatment cycle. Infertility indicator genes were identified as described in detail below in the Examples. Other genes that are suitable for use in the analysis can be identified using the methods described here in the Examples section.

[0074] In carrying out a subject assessment, a sample comprising an infertility indicator gene is assayed for a level of an infertility indicator gene product(s). Where the gene product being assayed is a nucleic acid, a nucleic acid sample (e.g., a sample comprising nucleic acid) is obtained from an embryo cell. Where the gene product being assayed is a protein, a polypeptide sample (e.g., a sample comprising polypeptides) is obtained from an embryo.

[0075] Nucleic acid (including mRNA) and/or polypeptide infertility indicator gene products can be obtained from an embryo, including an oocyte, such as an arrested embryo or oocyte (e.g., an embryo or oocyte having less than about 8 cells on about day 3 following fertilization, including about 7 cells, about 6 cells, about 5 cells, about 4 cells, about 3 cells, etc.), using standard methods. Levels of nucleic acid and/or polypeptide gene products can be measured using any of a variety of well-known methods, including those described in the Examples below.

[0076] An expression level of a response indicator gene is normalized relative to the level of an expression product of a reference gene(s). Assessing the infertility likelihood is conducted by comparing the normalized expression level to a range of values of normalized expression levels of the gene product in an embryo cell.

[0077] Normalized expression level of one or more infertility indicator genes can be carried out to assess the likelihood that a patient will respond positively or

negatively to an in vitro fertilization treatment cycle. Normalized expression level of a single infertility indicator gene can be carried out to assess the likelihood that a patient will respond positively or negatively to an in vitro fertilization treatment cycle. In addition, normalized expression level of two or more infertility indicator genes can be carried out to assess the likelihood that a patient will respond positively or negatively to an in vitro fertilization treatment cycle. The analysis can be more stringent, e.g., the optimal number of embryos to transfer to the female patient to maximize the likelihood of a live birth outcome while minimizing the likelihood of multiple gestations. The analysis can be less stringent, e.g., the likelihood that a patient will exhibit a beneficial response to an in vitro fertilization treatment cycle.

[0078] In some embodiments, the analysis includes determining the optimal number of embryos to transfer in order to minimize the probability of multiple gestation events in a subject. In such embodiments, the subject is first identified as a subject having a high probability of having multiple gestation events. The subject is then analyzed to determine the optimal number of embryos to transfer in order to provide for a single live birth event following the in vitro fertilization cycle.

[0079] It will be appreciated that assessment of likelihood that a patient will respond positively or negatively in vitro fertilization treatment cycle can be conducted by determining normalized expression levels of two or more infertility indicator genes (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more infertility indicator genes), or any combination of one or more sets of infertility indicator genes. The assessment can involve analyzing expression levels of a combination of infertility indicator genes, and determining normalized expression levels of the combination of infertility indicator genes, where the infertility indicator gene products can include gene products that are positively correlated with an in vitro fertilization treatment cycle and gene products that are negatively correlated with an in vitro fertilization treatment cycle. For example, a normalized level of a first gene that positively correlates with an in vitro fertilization treatment cycle, and a normalized level of a second gene that negatively correlates with an in vitro fertilization treatment cycle, can be determined.

Determining a normalized level of a gene product

- [0080] As discussed above, the expression level of an infertility indicator gene is normalized, thereby providing a normalized value. The expression level of an infertility indicator gene is normalized relative to the level of an expression product of a reference gene(s).
- [0081] For example, the expression level of an infertility indicator gene can be normalized relative to the mean level of gene products of two or more reference genes. As an example, the expression level of an infertility indicator gene can be normalized relative to the mean level of gene products of all assayed genes, or a subset of the assayed genes, where a subset of the assayed genes can include 3, 4, 5, 6, 7, 8, 9, or more assayed genes.
- [0082] Suitable reference genes include, but are not limited to, glyceraldehyde-3-phosphate dehydrogenase (GAPDH) (see, e.g., GenBank Accession No. NM_002046; phosphoglycerate kinase 1 (see, e.g., GenBank Accession No. NM_000291); lactate dehydrogenase A (see, e.g., GenBank Accession No. NM_005566); ribosomal protein L32 (see, e.g., GenBank Accession No. NM_000994); ribosomal protein S18 (see, e.g., GenBank Accession No. NM_022551); tubulin, beta polypeptide (TUBB) (see, e.g., GenBank Accession No. NM_001069); and beta actin (see, e.g., GenBank Accession No. NM_001101). See, e.g., Eisenberg and Levanon (2003) *Trends in Genetics* 19:362, for a list of additional suitable reference genes.
- [0083] The level of an RNA transcript as measured by TaqMan® RT-PCR refers to the cycle threshold (Ct) value. The lower the Ct, the greater the amount of mRNA present in the sample. The expression value of a RNA transcript in a sample is normalized, e.g., by first determining the mean expression value in Ct of designated reference genes in a sample (Ct_{Ref}). The normalized expression value for a gene (Ct_{Gene}) is then calculated as $Ct_{Gene} - Ct_{Ref}$. Optionally, the normalized expression values for all genes can be adjusted, e.g., so that all adjusted normalized Ct have a value >0.

Determining a probability of beneficial response

[0084] A normalized level of an infertility indicator gene product determined for an individual patient can be compared to normalized expression level values for the infertility indication gene determined in a population of patients for which the clinical outcome is already known in order to determine an individual patient's probability of beneficial response to an in vitro fertilization treatment cycle. Normalized expression level values (e.g., expressed as Ct) correlated with a probability can be used. For example, a normalized level of a response indicator gene product can be compared graphically, to determine the probability of beneficial response to an in vitro fertilization treatment cycle.

[0085] The analyses and determinations described herein in connection with a subject method for assessing likelihood of response can be made without the need for assessing any change in the level of a response indicator gene over time.

Classification Tree Analysis

[0086] One approach to analyze this data is to use a classification tree algorithm that searches for patterns and relationships in large datasets. A "classification tree" is a recursive partition to evaluate a female subject for an in vitro fertilization procedure using a series of questions that are designed to accurately place the patient into one of the classes. Each question asks whether a patient's condition satisfies a given predictor, with each answer being used to guide the user down the classification tree until a class into which the patient falls can be determined. As used herein, a "predictor" is the range of values of the features—such as, for example, age, total number of embryos, rate of cleavage arrest, number of 8-cell embryos, day 3 follicle stimulating hormone (FSH) level, and number of 8-cell embryos transferred.

Multiple Additive Regression Trees

[0087] An automated, flexible modeling technique that uses multiple additive regression trees (MART) may also be used to classify sets of features as belonging to one of two populations. A MART model uses an initial offset, which specifies a constant that applies to all predictions, followed by a series of regression trees. Its

fitting is specified by the number of decision points in each tree, the number of trees to fit, and a "granularity constant" that specifies how radically a particular tree can influence the MART model. For each iteration, a regression tree is fitted to estimate the direction of steepest descent of the fitting criterion. A step having a length specified by the granularity constant is taken in that direction. The MART model then consists of the initial offset plus the step provided by the regression tree. The differences between the observed and predicted values are recalculated, and the cycle proceeds again, leading to a progressive refinement of the prediction. The process continues either for a predetermined number of cycles or until some stopping rule is triggered.

[0088] The number of splits in each tree is a particularly meaningful fitting parameter. If each tree has only one split, the model looks only at one feature and has no capability for combining two predictors. If each tree has two splits, the model can accommodate two-way interactions among features. With three trees, the model can accommodate three-way interactions, and so forth.

[0089] The value of sets of features in predicting class status was determined for data sets with features and known class status. MART provides a measure of the contribution or importance of individual features to the classification decision rule. Specifically, the degree to which a single feature contributes to the decision rule upon its selection at a given tree split can be measured to provide a ranking of features by their importance in determining the final decision rule. Repeating the MART analysis on the same data set may yield a slightly different ranking of features, especially with respect to those features that are less important in establishing the decision rule. Sets of predictive features and their corresponding biomarkers that are useful for the present invention, therefore, may vary slightly from those set forth herein.

[0090] One exemplary implementation of the MART technology is found in a module, or "package," for the R statistical programming environment (see Venables et al., in *Modern Applied Statistics with S*, 4th ed. (Springer, 2002); www.r-project.org). Results reported in this document were calculated using R versions 1.7.0 and 1.7.1. The module implementing MART, written by Dr. Greg Ridgeway, is called "gbm" and is also available for download (see www.r-project.org). The

MART algorithm is amenable to ten-fold cross-validation. The granularity parameter was set to 0.05, and the gbm package's internal stopping rule was based on leaving out 20% of the data cases at each marked iteration. The degree of interaction was set to one, so no interactions among features were considered. The gbm package estimates the relative importance of each feature on a percentage basis, which cumulatively equals 100% for all the features of the biomarker profile. The features with highest importance, which together account for at least 90% of total importance, are reported as potentially having predictive value. Note that the stopping rule in the fitting of every MART model contributes a stochastic component to model fitting and feature selection. Consequently, multiple MART modeling runs based on the same data may choose slightly, or possibly even completely, different sets of features. Such different sets convey the same predictive information; therefore, all the sets are useful in the present invention. Fitting MART models a sufficient number of times is expected to produce all the possible sets of predictive features within a profile. Accordingly, the disclosed sets of predictors are merely representative of those sets of features that can be used to classify individuals into populations.

Wilcoxon Signed Rank Test Analysis

[0091] In yet another method, a nonparametric test such as a Wilcoxon Signed Rank Test can be used to identify individual biomarkers of interest. The features in a biomarker profile are assigned a "p-value," which indicates the degree of certainty with which the biomarker can be used to classify individuals as belonging to a particular reference population. Generally, a p-value having predictive value is lower than about 0.05. Biomarkers having a low p-value can be used by themselves to classify individuals. Alternatively, combinations of two or more biomarkers can be used to classify individuals, where the combinations are chosen on the basis of the relative p-value of a biomarker. In general, those biomarkers with lower p-values are preferred for a given combination of items of information. Combinations of at least three, four, five, six, 10, 20 or 30 or more biomarkers also can be used to classify individuals in this manner. The artisan will understand that the relative p-

value of any given biomarker may vary, depending on the size of the reference population.

Analysis Results Reporting

- [0092] As discussed above, evaluation of a female subject for an in vitro fertilization procedure, including determining probability of a live birth event, is done by obtaining and comparing items of information from the female subject to a library of known profile patterns known to be indicative of responsiveness to an in vitro fertilization procedure using an algorithm based upon said preselected patient variables and optionally evaluating the normalized expression level of one or more fertility response genes. In some embodiments, a patient's evaluation is provided in a report. Thus, in some embodiments, the method further includes a step of preparing or generating a report that includes information regarding the patient's likelihood of successes for an in vitro fertilization procedure. For example, a subject method can further include a step of generating or outputting a report providing the results of a patient's evaluation, which report can be provided in the form of an electronic medium (e.g., an electronic display on a computer monitor), or in the form of a tangible medium (e.g., a report printed on paper or other tangible medium).
- [0093] A report that includes information regarding the patient's evaluation is provided to a user. The evaluation may include, for example, a determination of success for implantation of embryos, selection of an optimal number of embryos to transfer, and determination of success in subsequent in vitro fertilization treatment cycles following an unsuccessful treatment cycle.
- [0094] The report generator may also perform one or more of sample gathering, sample processing, and data generation, e.g., the report generator may also perform one or more of: a) sample gathering; b) sample processing; c) measuring a level of an infertility indicator gene product(s); d) measuring a level of a reference gene product(s); and e) determining a normalized level of a infertility indicator gene product(s). Alternatively, an entity other than the report generator can perform one or more sample gathering, sample processing, and data generation.

[0095] For clarity, it should be noted that the term “user,” which is used interchangeably with “client,” is meant to refer to a person or entity to whom a report is transmitted, and may be the same person or entity who does one or more of the following: a) collects a sample; b) processes a sample; c) provides a sample or a processed sample; and d) generates data (e.g., level of a response indicator gene product(s); level of a reference gene product(s); normalized level of a response indicator gene product(s)) for use in the likelihood assessment. In some cases, the person(s) or entity(ies) who provides sample collection and/or sample processing and/or data generation, and the person who receives the results and/or report may be different persons, but are both referred to as “users” or “clients” herein to avoid confusion. In certain embodiments, e.g., where the methods are completely executed on a single computer, the user or client provides for data input and review of data output. A “user” can be a health professional (e.g., a clinician, a laboratory technician, a physician (e.g., a Reproductive Endocrinologist), etc.).

[0096] In embodiments where the user only executes a portion of the method, the individual who, after computerized data processing according to the methods of the invention, reviews data output (e.g., results prior to release to provide a complete report, a complete, or reviews an “incomplete” report and provides for manual intervention and completion of an interpretive report) is referred to herein as a “reviewer.” The reviewer may be located at a location remote to the user (e.g., at a service provided separate from a healthcare facility where a user may be located).

Report

[0097] A “report,” as described herein, is an electronic or tangible document which includes report elements that provide information of interest relating to a subject likelihood assessment and its results. A subject report includes at least an evaluation for an in vitro fertilization treatment cycle, e.g., facilitate a determination of success for implantation of embryos, selection of an optimal number of embryos to transfer, and determination of success in subsequent in vitro fertilization treatment cycles following an unsuccessful treatment cycle. A subject report can be completely or partially electronically generated. A subject report can further include one or more of: 1) information regarding the testing facility; 2) service provider information; 3)

patient data; 4) sample data; 5) an interpretive report, which can include various information including: a) indication; b) test data, where test data can optionally include normalized level of one or more infertility indicator gene products; and 6) other features.

- [0098] Where government regulations or other restrictions apply (e.g., requirements by health, malpractice, or liability insurance), all results, whether generated wholly or partially electronically, are subjected to a quality control routine prior to release to the user.

Testing facility information

- [0099] The report can include information about the testing facility, which information is relevant to the hospital, clinic, or laboratory in which sample gathering and/or data generation was conducted. Data generation can include one or more of: a) measuring a level of a gene product(s) (e.g., an infertility indicator gene product(s), a reference gene product(s)); b) determination of a normalized level of an infertility indicator gene product. This information can include one or more details relating to, for example, the name and location of the testing facility, the identity of the lab technician who conducted the assay and/or who entered the input data, the date and time the assay was conducted and/or analyzed, the location where the sample and/or result data is stored, the lot number of the reagents (e.g., kit, etc.) used in the assay, and the like. Report fields with this information can generally be populated using information provided by the user.

Service provider information

- [00100] The report can include information about the service provider, which may be located outside the healthcare facility at which the user is located, or within the healthcare facility. Examples of such information can include the name and location of the service provider, the name of the reviewer, and where necessary or desired the name of the individual who conducted sample gathering and/or data generation. Report fields with this information can generally be populated using data entered by the user, which can be selected from among pre-scripted selections (e.g., using a drop-down menu). Other service provider information in the report can include

contact information for technical information about the result and/or about the interpretive report.

Patient data

[00101] The patient data can include patient medical history (which can include, e.g., data about prior or current in vitro fertilization treatment cycles), personal history; administrative patient data (that is, data that are not essential to the likelihood assessment), such as information to identify the patient (e.g., name, patient date of birth (DOB), gender, mailing and/or residence address, medical record number (MRN), room and/or bed number in a healthcare facility), insurance information, and the like), the name of the patient's physician or other health professional who ordered the response likelihood assessment and, if different from the ordering physician, the name of a staff physician who is responsible for the patient's care (e.g., primary care physician). Report fields with this information can generally be populated using data entered by the user.

[00102] Exemplary items of information include, but are not limited to: patient characteristics, such as age, previous infertility history, clinical diagnosis; clinical treatment information, such as type of medication, number of days of stimulation, number of oocytes, etc.; conventional embryo morphology data, such as number of embryos, developmental stage, grade, and the like. In some embodiments, the information includes age, total number of embryos, rate of cleavage arrest, number of 8-cell embryos, day 3 follicle stimulating hormone (FSH) level, number of 8-cell embryos transferred, age, diminished ovarian reserve, endometrial thickness, blastocyst rate, total number of embryos, total number of oocytes, total amount of gonatropins administered, and number of total motile sperm.

Sample data

[00103] The sample data can provide information about the embryo analyzed in the likelihood assessment, such as the number of days following fertilization resulting in arrest, and the date and time collected. Report fields with this information can generally be populated using data entered by the user, some of which may be provided as pre-scripted selections (e.g., using a drop-down menu).

Interpretive report

- [00104] The interpretive report portion of the report includes information generated after processing of the data as described herein. The interpretive report can include an evaluation of a female subject for an in vitro fertilization procedure. The interpretive report can include, for example, Result of analysis comparing the profile for the female subject to a library of known profile patterns known to be indicative of responsiveness to an in vitro fertilization procedure using an algorithm based upon said preselected patient variables, and optionally Result of normalized level of infertility indicator gene(s) (e.g., “normalized level of infertility indicator gene(s)”; Interpretation; and, optionally, Recommendation(s).
- [00105] The Interpretation portion of the report can include a Recommendation(s). Where the results indicate a determination of success for implantation of embryos, selection of an optimal number of embryos to transfer, and determination of success in subsequent in vitro fertilization treatment cycles following an unsuccessful treatment cycle.
- [00106] It will be readily appreciated that the report can include all or some of the elements above, with the proviso that the report generally includes at least the elements sufficient to provide the analysis requested by the user (e.g., likelihood assessment).

Additional features

- [00107] It will also be readily appreciated that the reports can include additional elements or modified elements. For example, where electronic, the report can contain hyperlinks which point to internal or external databases which provide more detailed information about selected elements of the report. For example, the patient data element of the report can include a hyperlink to an electronic patient record, or a site for accessing such a patient record, which patient record is maintained in a confidential database. This latter embodiment may be of interest in an in-hospital system or in-clinic setting.

Computer-Based Systems and Methods

[00108] The methods and systems described herein can be implemented in numerous ways. In one embodiment of particular interest, the methods involve use of a communications infrastructure, for example the internet. Several embodiments of the invention are discussed below. It is also to be understood that the present invention may be implemented in various forms of hardware, software, firmware, processors, or a combination thereof. The methods and systems described herein can be implemented as a combination of hardware and software. The software can be implemented as an application program tangibly embodied on a program storage device, or different portions of the software implemented in the user's computing environment (e.g., as an applet) and on the reviewer's computing environment, where the reviewer may be located at a remote site associated (e.g., at a service provider's facility).

[00109] The various elements of the computing device, such as a the input device, may be associated with other elements of the system via a wired connection or a wireless connection, including, for example, a wireless LAN connection, Bluetooth connection protocol, ZigBee connection protocol, radio-frequency connection protocol, or a cellular phone connection protocol, including code derived multiple access (CDMA) or via a global system for mobile communication (GSM).

[00110] For example, during or after data input by the user, portions of the data processing can be performed in the user-side computing environment. For example, the user-side computing environment can be programmed to provide for defined test codes to denote a likelihood "score," where the score is transmitted as processed or partially processed responses to the reviewer's computing environment in the form of test code for subsequent execution of one or more algorithms to provide a results and/or generate a report in the reviewer's computing environment.

[00111] The application program for executing the algorithms described herein may be uploaded to, and executed by, a machine comprising any suitable architecture. In general, the machine involves a computer platform having hardware such as one or more central processing units (CPU), a random access memory (RAM), and input/output (I/O) interface(s). The computer platform also includes an operating system and microinstruction code. The various processes and functions described

herein may either be part of the microinstruction code or part of the application program (or a combination thereof) which is executed via the operating system. In addition, various other peripheral devices may be connected to the computer platform such as an additional data storage device and a printing device.

[00112] As a computer system, the system generally includes a processor unit. The processor unit operates to receive information, which generally includes subject information data, such as age, total number of embryos, rate of cleavage arrest, number of 8-cell embryos, day 3 follicle stimulating hormone (FSH) level, number of 8-cell embryos transferred, age, diminished ovarian reserve, endometrial thickness, blastocyst rate, total number of embryos, total number of oocytes, total amount of gonatropins administered, and number of total motile sperm. This information received can be stored at least temporarily in a database, and data analyzed to generate a report as described above.

[00113] Part or all of the input and output data can also be sent electronically; certain output data (e.g., reports) can be sent electronically or telephonically (e.g., by facsimile, e.g., using devices such as fax back). Exemplary output receiving devices can include a display element, a printer, a facsimile device and the like. Electronic forms of transmission and/or display can include email, interactive television, and the like. In an embodiment of particular interest, all or a portion of the input data and/or all or a portion of the output data (e.g., usually at least the final report) are maintained on a web server for access, preferably confidential access, with typical browsers. The data may be accessed or sent to health professionals as desired. The input and output data, including all or a portion of the final report, can be used to populate a patient's medical record which may exist in a confidential database at the healthcare facility.

[00114] A system for use in the methods described herein generally includes at least one computer processor (e.g., where the method is carried out in its entirety at a single site) or at least two networked computer processors (e.g., where data is to be input by a user (also referred to herein as a "client") and transmitted to a remote site to a second computer processor for analysis, where the first and second computer processors are connected by a network, e.g., via an intranet or internet). The system can also include a user component(s) for input; and a reviewer component(s) for

review of data, generated reports, and manual intervention. Additional components of the system can include a server component(s); and a database(s) for storing data (e.g., as in a database of report elements, e.g., interpretive report elements, or a relational database (RDB) which can include data input by the user and data output. The computer processors can be processors that are typically found in personal desktop computers (e.g., IBM, Dell, Macintosh), portable computers, mainframes, minicomputers, or other computing devices, such as Smartphone devices, including, for example, an Apple® iPhone® device.

[00115] The networked client/server architecture can be selected as desired, and can be, for example, a classic two or three tier client server model. A relational database management system (RDMS), either as part of an application server component or as a separate component (RDB machine) provides the interface to the database.

[00116] In one embodiment, the architecture is provided as a database-centric client/server architecture, in which the client application generally requests services from the application server which makes requests to the database (or the database server) to populate the report with the various report elements as required, particularly the interpretive report elements, especially the interpretation text and alerts. The server(s) (e.g., either as part of the application server machine or a separate RDB/relational database machine) responds to the client's requests.

[00117] The input client components can be complete, stand-alone personal computers offering a full range of power and features to run applications. The client component usually operates under any desired operating system and includes a communication element (e.g., a modem or other hardware for connecting to a network), one or more input devices (e.g., a keyboard, mouse, keypad, or other device used to transfer information or commands), a storage element (e.g., a hard drive or other computer-readable, computer-writable storage medium), and a display element (e.g., a monitor, television, LCD, LED, or other display device that conveys information to the user). The user enters input commands into the computer processor through an input device. Generally, the user interface is a graphical user interface (GUI) written for web browser applications.

[00118] The server component(s) can be a personal computer, a minicomputer, or a mainframe and offers data management, information sharing between clients,

network administration and security. The application and any databases used can be on the same or different servers.

[00119] Other computing arrangements for the client and server(s), including processing on a single machine such as a mainframe, a collection of machines, or other suitable configuration are contemplated. In general, the client and server machines work together to accomplish the processing of the present invention.

[00120] Where used, the database(s) is usually connected to the database server component and can be any device which will hold data. For example, the database can be a any magnetic or optical storing device for a computer (e.g., CDROM, internal hard drive, tape drive). The database can be located remote to the server component (with access via a network, modem, etc.) or locally to the server component.

[00121] Where used in the system and methods, the database can be a relational database that is organized and accessed according to relationships between data items. The relational database is generally composed of a plurality of tables (entities). The rows of a table represent records (collections of information about separate items) and the columns represent fields (particular attributes of a record). In its simplest conception, the relational database is a collection of data entries that "relate" to each other through at least one common field.

[00122] Additional workstations equipped with computers and printers may be used at point of service to enter data and, in some embodiments, generate appropriate reports, if desired. The computer(s) can have a shortcut (e.g., on the desktop) to launch the application to facilitate initiation of data entry, transmission, analysis, report receipt, etc. as desired.

Computer-Readable Storage Media

[00123] The invention also contemplates a computer-readable storage medium (e.g. CD-ROM, memory key, flash memory card, diskette, etc.) having stored thereon a program which, when executed in a computing environment, provides for implementation of algorithms to carry out all or a portion of the methods of analysis of evaluating a subject for an in vitro fertilization procedure as described herein. Where the computer-readable medium contains a complete program for carrying

out the methods described herein, the program includes program instructions for collecting, analyzing and generating output, and generally includes computer readable code devices for interacting with a user as described herein, processing that data in conjunction with analytical information, and generating unique printed or electronic media for that user.

- [00124] Where the storage medium provides a program which provides for implementation of a portion of the methods described herein (e.g., the user-side aspect of the methods (e.g., data input, report receipt capabilities, etc.)), the program provides for transmission of data input by the user (e.g., via the internet, via an intranet, etc.) to a computing environment at a remote site. Processing or completion of processing of the data is carried out at the remote site to generate a report. After review of the report, and completion of any needed manual intervention, to provide a complete report, the complete report is then transmitted back to the user as an electronic document or printed document (e.g., fax or mailed paper report). The storage medium containing a program according to the invention can be packaged with instructions (e.g., for program installation, use, etc.) recorded on a suitable substrate or a web address where such instructions may be obtained. The computer-readable storage medium can also be provided in combination with one or more reagents for carrying determining subject information data, e.g., material for determining rate of cleavage arrest, number of 8-cell embryos, day 3 follicle stimulating hormone (FSH) level, and number of 8-cell embryos transferred, and the like.

Kits

- [00125] The materials for use in the methods of the present invention are suited for preparation of kits produced in accordance with well known procedures. The invention thus provides kits comprising reagents, which may include gene-specific or gene-selective probes and/or primers useful for assaying the expression of genes disclosed herein and for assessing the likelihood of response to an in vitro treatment cycle.
- [00126] For example, a subject kit can include one or more nucleic acid probes that hybridize specifically to nucleic acid infertility indicator gene products. A subject

kit can include, e.g., one or more nucleic acid probes, where each of the one or more probes hybridizes specifically to a different response indicator gene product. For example, a subject kit can include probes that hybridize specifically to nucleic acid products of infertility indicator genes including, but are not limited to, Oct4, Eif3c, Papola, Piwil2, Eif3b, Eif4b, Rbm3, and Cpsf4. As another example, a subject kit can include a set of two or more nucleic acid probes, where each probe of the set hybridizes to a nucleic acid product of a different infertility indicator gene. For example, a subject kit can include a set of two, three, four, five, six, seven, or more, nucleic acid probes, where each probe of the set hybridizes to a nucleic acid product of a different member of a set of infertility indicator genes.

[00127] In some cases, a subject kit will include, in addition to a probe that hybridizes specifically to a nucleic acid product of an infertility indicator gene, one or more probes that hybridize specifically to a reference gene product. Such probes can be used in determining normalized expression levels of an infertility indicator gene.

[00128] A subject kit can include one or more nucleic acid primer pairs, where the primer pairs, when used as primers in a polymerase chain reaction, amplify a target nucleic acid response indicator gene product, or a target region of a nucleic acid response indicator gene product. A subject kit can include primer pairs for multiple infertility indicator genes.

[00129] Exemplary sequences of nucleic acid primers and probes are provided in the Examples described herein. Those skilled in the art will readily appreciate that other probe and primer sequences are also possible, and are readily obtained based on known nucleotide sequences of infertility indicator genes, and/or based on known nucleotide sequences of reference genes.

[00130] In addition to the above-mentioned probes and primers, a subject kit can comprise reagents for the extraction and/or isolation of RNA from single cell embryos, in particular fixed paraffin-embedded tissue samples and/or reagents for preparing a cDNA copy of an mRNA, and/or reagents for nucleic acid amplification. Exemplary reagents include those required for use of a FLUIDIGM® BIOMARK® 48.48 Dynamic array system, comparable single-cell

gene expression analysis platform (RT-PCR), or emerging technology such as next-generation, whole-transcriptome sequencing at the single-cell level.

[00131] Primers and probes can be designed based on known sequences of infertility indicator genes, and are readily synthesized by standard techniques, e.g., solid phase synthesis via phosphoramidite chemistry, as disclosed in U.S. Pat. Nos. 4,458,066 and 4,415,732, incorporated herein by reference; Beaucage et al. (1992) Tetrahedron 48:2223-2311; and Applied Biosystems User Bulletin No. 13 (1 Apr. 1987). Other chemical synthesis methods include, for example, the phosphotriester method described by Narang et al., Meth. Enzymol. (1979) 68:90 and the phosphodiester method disclosed by Brown et al., Meth. Enzymol. (1979) 68:109. Poly(A) or poly(C), or other non-complementary nucleotide extensions may be incorporated into probes using these same methods. Hexaethylene oxide extensions may be coupled to probes by methods known in the art. Cload et al. (1991) J. Am. Chem. Soc. 113:6324-6326; U.S. Pat. No. 4,914,210 to Levenson et al.; Durand et al. (1990) Nucleic Acids Res. 18:6353-6359; and Horn et al. (1986) Tet. Lett. 27:4705-4708.

[00132] A probe or a primer can include a detectable label. Exemplary labels include fluorochromes, e.g. fluorescein isothiocyanate (FITC), rhodamine, Texas Red, phycoerythrin, allophycocyanin, 6-carboxyfluorescein (6-FAM), 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein (JOE), 6-carboxy-X-rhodamine (ROX), 6-carboxy-2',4',7',4,7-hexachlorofluorescein (HEX), 5-carboxyfluorescein (5-FAM), N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA), Cy5, Cy3, and the like; and radioactive labels (e.g., ³²P, etc.).

[00133] Probes and primers for inclusion in a subject kit include those useful in various amplification and/or detection systems. Exemplary amplification and/or detection systems include Sunrise™ primer-based systems, Molecular Beacons, the Taqman™ system, an Amplifluor™ hairpin primer-based system, a Scorpions technology (e.g., bi-functional molecules containing a PCR primer element covalently linked to a probe element), a Light Upon Extension or LUX™-based system, and a FLUIDIGM® BIOMARK® 48.48 Dynamic array system. Further exemplary detection systems include those based on a melt-curve analysis, and using intercalating dyes such as the fluorescent dye SYBR Green.

[00134] The kits may optionally comprise reagent(s) with an identifying description or label or instructions relating to their use in the methods of the present invention. The kits may comprise containers (including microtiter plates suitable for use in an automated implementation of the method), each with one or more of the various reagents (typically in concentrated form) utilized in the methods of the invention, including, for example, pre-fabricated microarrays, buffers, the appropriate nucleotide triphosphates (e.g., dATP, dCTP, dGTP and dTTP; or rATP, rCTP, rGTP and UTP), reverse transcriptase, DNA polymerase, RNA polymerase, and one or more probes and primers of the present invention (e.g., appropriate length poly(T) or random primers linked to a promoter reactive with the RNA polymerase).

[00135] Instructions for the use of mathematical algorithms used to evaluate a female subject for an in vitro fertilization treatment cycle, including determining a probability of a live birth event, can also be included in a subject kit. In such embodiments, the kits will further include a written or electronic medium, or instructions to access a remote database, as described above, to provide and/or receive information, which generally includes subject information data, such as age, total number of embryos, rate of cleavage arrest, number of 8-cell embryos, day 3 follicle stimulating hormone (FSH) level, number of 8-cell embryos transferred, age, diminished ovarian reserve, endometrial thickness, blastocyst rate, total number of embryos, total number of oocytes, total amount of gonatropins administered, and number of total motile sperm, in order to carry out the methods as described above.

EXAMPLES

[00136] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the present invention, and are not intended to limit the scope of what the inventors regard as their invention nor are they intended to represent that the experiments below are all or the only experiments performed. Efforts have been made to ensure accuracy with respect to numbers used (e.g. amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated

otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

Materials and Methods

[00137] The following methods and materials were used in the examples below.

Data Collection, Inclusion and Exclusion Criteria

[00138] Data related to clinical diagnoses, IVF treatment protocol and monitoring, embryology data and treatment outcomes for all IVF cycles performed between January 1, 2003 and December 31, 2006 at Stanford University Medical Center were retrieved from BabySentryPro (BabySentry Ltd, Limassol, Cyprus), a widely used fertility database management system, or obtained from medical and embryology records as necessary. Retrospective data collection, de-identification, and analysis were performed according to a Stanford University Institutional Review Board-approved protocol. The inclusion criteria for data analysis were fresh, stimulated, non-donor oocyte IVF cycles.

[00139] In certain experiments, cycles that did not result in embryo transfer for any reason, cycles performed for women aged over 45, and those performed for preimplantation genetic screening were excluded.

[00140] In other experiments, we did not exclude cycles that were canceled at any point due to medical or non-medical reasons after the IVF treatment had started. We excluded cryopreserved embryo transfer cycles which used embryos that had been conceived in a different IVF clinic.

Defining Cycle Numbers

[00141] We renumbered all IVF cycles according to the following method. For each patient, the first fresh IVF cycle appearing in this four-year data set is called "Cycle 1". The final outcome of all embryo transfer treatments that utilize fresh or cryopreserved embryos conceived from Cycle 1 is defined as the outcome for Cycle 1. The second fresh IVF cycle that appears is defined "Cycle 2", and so on. Our

database could not accommodate data on IVF cycles that were performed elsewhere.

Assessment of Embryo Development

[00142] The standard clinical protocols for IVF treatment, fertilization, embryo culture, embryo assessment, cryopreservation criteria, and clinical outcomes are described below. The normal progression of human embryo development *in vitro* is characterized by the appearance of 2 pronuclei at 16-20 hours after insemination as evidence of fertilization on Day 1, with Day 0 as the day of oocyte retrieval. By late Day 1, embryo development has reached the 2-cell stage, followed by the 4-cell and 8-cell stages on Days 2 and 3, respectively. On Days 4 and 5, embryo development is characterized by the establishment of the morula and blastocyst stages, respectively. All embryos were available for evaluation on Day 3. The day of embryo transfer was determined by the number of blastomeres on Day 3. In general, if 4 or more 8-cell embryos were present, we would recommend extended embryo culture until Day 5, when blastocyst transfer, which has been associated with higher pregnancy rates, would be performed. If fewer than four 8-cell embryos were present, embryo transfer would be performed on Day 3.

Patient, IVF Cycle, and Embryo Parameters

[00143] 30 variables for association with IVF treatment outcomes were analyzed, as listed in Table 1, under four main categories: patient characteristics and clinical diagnoses, IVF cycle characteristics, embryo cohort parameters, and parameters of transferred embryos. The cleavage arrest rate was defined as the percentage of embryos within a cohort with 4 or fewer cells on Day 3 of *in vitro* culture. All other variables were self-explanatory.

Definition of Outcomes

[00144] Statistical analyses were performed based on the dichotomous outcomes of live birth versus no live birth. No live birth encompassed all other outcomes such as negative serum β -hCG, or positive serum β -hCG followed by biochemical pregnancy, spontaneous abortion, and ectopic pregnancy.

Statistical Analysis

- [00145] Since some patients underwent more than one IVF cycle during the study period, the analyses were performed based on treatment cycles rather than patients. Statistical analyses were performed based on the dichotomous outcomes of no pregnancy, as defined by negative serum β -hCG, and pregnancy, as defined by positive serum β -hCG, and included biochemical pregnancy, clinical pregnancy, spontaneous abortion, and ectopic pregnancy. We performed pair-wise logistic regression of each variable to the outcome and determined the Pearson correlation coefficient between each pair of continuous variables.
- [00146] For the main analyses, boosted classification trees were constructed by MART[®] to identify non-redundant prognostic variables, which were then further analyzed by CART to identify thresholds that would define them as categorical variables. MART[®] is a robust method used to identify interactive structure of variables that are predictive of outcomes. The use of cross-validation and boosting in parameter selection and model assessment in MART[®] also preserve parsimony and prevent over-fitting. In the MART[®] tree constructions, the whole data set is divided into 10 subsets to achieve 10 fold cross validation for model assessment. The same 10 fold cross validation was repeated 1000 times to perform a robust prediction rate estimation and identify tree models with the highest prediction rates in the CART. While MART[®] is powerful in selecting non-redundant prognostic variables from a large set of highly interactive variables, CART analysis results in simple algorithms, and more easily understood “decision trees”, that are used in the medical literature (Guznick et al., N Engl J Med 345: 1388-1393. (2001)). Thus non-redundant, prognostic variables identified by MART[®] to confer prediction were analyzed by CART to further define prognostic thresholds.
- [00147] In some experiments, if patients underwent multiple fresh IVF cycles during the study period, only the first three cycles were analyzed. We generated the models based on cycle 1 (C1) data only. Data from C2 and C3 were used to test inter-cycle validation of the pre-transfer model only.
- [00148] In such experiments, we performed pair-wise logistic regression of each variable to the outcome and determined the Pearson correlation coefficient between

each pair of continuous variables. We generated three models based on data that were available at three different time points prior to IVF treatment (pre-IVF), oocyte retrieval (pre-OR), and after embryo transfer (post-IVF). We generated each model by ranking variables according to their relative influence using gradient boosting machine (GBM), the R-software implementation of MART[®], followed by construction of regression tree models using top-ranked variables with Rpart software. MART[®] is a robust method used to identify interactive structure of variables that are predictive of outcomes. The use of cross-validation and boosting in parameter selection and model assessment in MART[®] also preserve parsimony and prevent over-fitting. In the MART[®] tree constructions, the whole data set is divided into 10 subsets to achieve 10 fold cross validation for model assessment. The same 10 fold cross validation was repeated 1000 times to perform a robust prediction rate estimation and identify tree models with the highest prediction rates. While MART[®] is powerful in selecting non-redundant prognostic variables from a large set of highly interactive variables.

Treatment Protocols used for Assisted Reproductive technologies (ART)

[00149] The majority of our analyzed IVF cycles were performed in patients with poor ovarian reserve, severe male factor infertility, tubal infertility, anovulatory disorders, or unexplained etiology. In general, one of three stimulation protocols was used in each treatment cycle: luteal downregulation (long) was used for most patients, and microdose lupron (flare) and antagonist protocols were used primarily for patients with presumed diminished ovarian reserve or with a history of previously failed IVF cycles. The long protocol consisted of luteal downregulation using 0.5 mg leuprolide acetate, which was decreased to 0.25 mg with stimulation. In the flare protocol, microdose lupron (0.04 mg s.c. bid) was started after 2-4 weeks of oral contraceptive pills. In the antagonist protocol, GnRH antagonist was initiated when the lead follicle reached 14mm in size. In all three protocols, baseline ultrasound testing was performed to document that no cysts > 1.5 cm were present in the ovaries. When baseline criteria were met, gonadotropin therapy using recombinant FSH with human menopausal gonadotropin was begun. Stimulation was generally achieved using daily dosing of a total of 150-600 IU per day in order

to maximize follicular recruitment. Ultrasound monitoring of follicular growth was performed starting on cycle day 7 and then every 1-3 days as indicated. Serum estradiol levels were monitored as necessary.

- [00150] A dose of 10,000 IU of human chorionic gonadotropin was administered when at least two follicles reached an average diameter of >17mm. Transvaginal ultrasound-guided oocyte retrieval was performed 34-36 hours after hCG administration in the standard fashion with monitored anesthetic care.

Oocyte Fertilization and Embryo Culture

- [00151] Oocytes were cultured in groups before fertilization under mineral oil in approximately 125 µl droplets of Sage Cleavage Medium (Cooper Surgical, Inc, Trumbull, CT) with 10% Serum Protein Substitute (SPS, Irvine Scientific, Santa Ana, CA). Oocytes destined for conventional IVF were cultured in groups of 5 and oocytes destined for intracytoplasmic sperm injection (ICSI) were cultured in groups of up to 20 after stripping the cumulus cells. Oocytes were inseminated with sperm if the semen analysis was normal and fertilization was expected to be normal. Oocytes were inseminated conventionally between 4-6 hours after retrieval. If the semen analysis was abnormal or poor fertilization was expected, then the oocytes were injected with sperm using ICSI. The fertilized oocytes were cultured in groups of up to 20 under mineral oil in approximately 125 µl droplets of Sage Cleavage Medium (Cooper Surgical, Inc, Trumbull, CT) with 10% SPS at 37°C in a humidified atmosphere of 5% O₂, 5% CO₂ and 90% N₂. Fertilization check was performed 16~18 hours after insemination or ICSI. The zygotes with clear two pronuclei were cultured for another 48 hours in Sage Cleavage Medium with 10% SPS. Oocytes with single pronucleus (1PN) or three or more pronuclei were considered abnormally fertilized. If Day 5 blastocyst transfer was indicated, extended embryo culture was performed in Quinn's Advantage Blastocyst medium (Cooper Surgical) with 10% SPS for 48 hours before transfer. Of note, the same culture media was used during the study time period.

Cleavage and Grading of Embryos

[00152] A single team of experienced embryologists evaluated the embryos on post-retrieval day 3, 68 to 72 hours after oocyte harvest. Embryos were examined for cleavage (cell number) and grade, which includes cytoplasmic fragmentation. Embryos were graded as follows on Day 3: Grade 1, blastomeres have equal size and no cytoplasmic fragmentation; Grade 2, blastomeres have equal size and minor cytoplasmic fragmentation involving < 10% of the embryo; Grade 3, blastomeres have unequal size and fragmentation involving 10-20% of the embryo; Grade 4, blastomeres have equal or unequal size, and moderate to significant cytoplasmic fragmentation covering 20-50% of the embryo; and Grade 5, few blastomeres and severe fragmentation covering $\geq 50\%$ of the embryo (Veeck et al., (1999) *An Atlas of Human Gametes and Conceptuses*. New York: Parthenon Publishing. 47-50). Of note, cytoplasmic fragmentation in embryos was easily differentiated from cleavage based on the size of the fragments, their location within the embryo, and the absence of a nucleus. In contrast, cytoplasmic fragmentation of oocytes was not included as a variable because it was extremely rare. The presence or absence of compaction was routinely noted by our embryologists on Day 3. As compaction was observed in <10% of embryos, we did not include this variable in the analysis.

Assisted Hatching

[00153] Indications for assisted hatching (AH) in our center were advanced maternal age, elevated FSH level, and/or a history of multiple failed assisted reproduction cycles. On the day of embryo transfer, embryos for hatching were placed in phosphate-buffered saline (PBS) with 10% SPS. AH was accomplished by using the ZILOS-tk laser (Hamilton Thorne Biosciences, Beverly, MA) to make a hole in an area of the zona pellucida that was between blastomeres. Embryos were then rinsed and returned to the media until transfer.

Embryo Transfer and Cryopreservation

[00154] Ultrasound-guided embryo transfer was performed using a Tefcat or Echotip Softpass catheter (Cook Ob/Gyn, Spencer, IN). Progesterone supplementation with

vaginal suppositories was performed in all patients. For patients having embryo transfer on Day 3, any remaining embryos with more than five blastomeres were placed in extended culture for another 2 or 3 days. Any expanding, expanded and hatching blastocysts with good inner cell mass and trophectoderm were frozen on day 5 or day 6. In addition, cryopreservation was performed on excess embryos, severe ovarian hyperstimulation syndrome, and fertility preservation due to medical or social reasons. Excess embryos that were not transferred were commonly discarded if they were not of sufficient quality for cryopreservation or if patients did not opt for embryo cryopreservation due to non-medical reasons. In cases of pre-implantation genetic diagnosis, embryos that tested positive for genetic diseases or aneuploidy were also discarded.

Clinical Outcomes

[00155] Serum quantitative β -hCG levels were obtained at 8-10 days after embryo transfer, and followed serially until the diagnosis of clinical pregnancy was made by the presence of a gestational sac on transvaginal ultrasound. Outcomes other than clinical pregnancy included: 1) no pregnancy if serum quantitative β -hCG was negative; 2) biochemical pregnancy as defined by decreasing serum quantitative β -hCG levels before a gestational sac could be visualized by transvaginal ultrasound; 3) spontaneous abortion as defined by pregnancy loss after a gestational sac was visualized by transvaginal ultrasound; and 4) ectopic pregnancy; and 5) other abnormal gestations such as gestational trophoblastic disease. Live birth outcomes were obtained by follow-up contact with patients as part of routine clinical care, but they were not used in this study.

Regression Trees

[00156] Clinical IVF data, especially when considering oocyte and embryo parameters, often do not lend themselves to meaningful analysis by multivariate logistic regression. The high degree of interaction and multicollinearity of many relevant variables interfere with conventional multivariate regression. In these situations, the regression and classification tree models (CART), has been widely

used in clinical research (Fonarow GC, *Jama* 293: 572-580 (2005); Friedman J (1999) Greedy function approximation: A stochastic boosting machine. Technical Report, Department of Statistics, Stanford University; Friedman J (1999) Stochastic gradient boosting. Technical Report, Department of Statistics, Stanford University; Friedman J (2002) Tutorial: Getting started with MART in R. Department of Statistics, Stanford University; Friedman et al., *Stat Med* 22: 1365-1381 (2003); Guzick et al., *N Engl J Med* 345: 1388-1393 (2001); Pilote et al., *N Engl J Med* 335: 1198-1205 (1996)). Here, we used Multiple Additive Regression Tree (MART®), a more powerful statistical method that combines “boosting” with CART to “boost” or increase accuracy in the CART method, to identify non-redundant prognostic variables.

[00157] In general, regression trees have several key advantages: 1) ability to consider all types of clinical IVF and embryology data, including numeric, ordinal, binary, and categorical variables; 2) ability to handle missing values well based on a “surrogate” splitting technique without the need for imputation; 3) ability to generate results that are invariant to monotonic data transformation and thus eliminate the need to test different methods of data transformation or metrics; 4) ability to generate trees that are immune to the effects of extreme outliers; 5) ability to generate trees that inherently explore and identify interactions of variables that would otherwise need to be explicitly stated in a multiple logistic regression. Most importantly, regression trees can consider a large number of variables, including ones that may turn out to be irrelevant, even if only a small number of variables have significant statistical impact on outcomes. This ability to consider many variables is critical for analysis of IVF outcomes, as many variables, such as percentage of 8-cell embryos, number of 8-cell embryos, percentage of 8-cell embryos transferred, and number of 8-cell embryos transferred, may be highly interactive; thus, arbitrarily selecting one of them may compromise completeness of data and introduce bias, while including all of them may cause the conventional multivariate regression to breakdown.

[00158] The results from MART® may help to identify variables that could be re-analyzed by multivariate logistic regression. Often, they may identify thresholds, or “cut-offs”, that will be used to create categorical variables to segregate cases into

subgroups for further inter-group comparison of characteristics by conventional methods such as t-tests, chi-square analysis, or Wilcoxon rank sum test. For example, CART analysis was used by Guzick et al. to classify men as subfertile, of indeterminate fertile status, or fertile based on threshold values for sperm concentration, motility, and morphology, exemplifies the power of this strategy in infertility research (Guzick et al., N Engl J Med 345: 1388-1393 (2001)).

Embryo Culture

[00159] 3-5 week old wild type F1 (C57BL6xDBA/2) females (Charles River) were superovulated by intraperitoneal injections of 5 IU of pregnant mare's serum gonadotropin (Sigma) followed by 5 IU of human chorionic gonadotropin (Sigma) 48 hours later, and mated overnight with wild type males. Mice were sacrificed by cervical dislocation 17 hours after hCG injection, and 1-cell embryos were released from oviducts. Cumulus cells were removed by hyaluronidase (Sigma) treatment and pipetting. Pre-implantation embryos at the two pronuclei stage were recovered, pooled from 3-6 females in M2 media (Chemicon International), followed by immediate cytoplasmic microinjection and culture in Human Tubal Fluid with 10% serum supplement (In-Vitro Fertilization, Inc.) microdrops under mineral oil (Sigma) in mixed gas (90% nitrogen, 5% oxygen, 5% carbon dioxide; Praxair) at 37°C, and cultured at ten embryos per 20 µL drop.

Microinjection of Antisense Morpholino Oligonucleotides

[00160] 25-nt, antisense morpholino oligonucleotides (MOs) that specifically target the 5'UTR or translational start site, or controls mismatched at 5 nts were purchased from Gene Tools, LLC. (See Table 1 for sequence details). We had determined 0.6-0.75 mM to be the maximal concentration that would allow normal rates of blastocyst development (data not shown). Hence, unless otherwise specified, 5-10 pL of 0.75 mM *Ccna2*-MO (0.60 mM for *Oct4*-MO) was injected into the cytoplasm of each embryo on an inverted microscope (Olympus IX70) equipped with hydraulic micromanipulation system (IM300 Microinjector, Narishige, Japan). 10 uninjected control embryos were used in each experiment, which was performed at least three times. The mean percentage and standard error

of the mean (mean $\pm$ s.e.m.) of embryos progressing to, or arresting at, each developmental stage were calculated, and statistical significance was determined by calculating the p-value using 2-tailed Student's t-test.

Table 1: Antisense Morpholino Oligonucleotides Target Gene-Specific Sequence in the 5' UTR and/or Start Site

Name	Sequence	GC content, %
<i>Ccna2</i> -MO-1	5'-TCGAGGTGCCCCGGCATCGCGGCTCC-3' (SEQ ID NO:01)	76
<i>Ccna2</i> -MO-2	5'-CTGTCCGGCGGCAGAGCGTTCACAGC-3' (SEQ ID NO:02)	68
<i>Ccna2</i> -MM-1	5'-TCCAGGTCCCCCGCATCCCGGATCC-3' (SEQ ID NO:03)	72
<i>Oct4</i> -MO	5'-AGTCTGAAGCCAGGTGTCCAGCCAT-3' (SEQ ID NO:04)	56
<i>Oct4</i> -MM	5'-ACTCTCAAGCCACGTGTGCAGCGAT-3' (SEQ ID NO:05)	56
<i>Oct4E4</i> -MO	5'-CTCCGATTTCATATCTGGGCAGGG-3' (SEQ ID NO:06)	56
<i>Oct4E4</i> -MM	5'-CTGCGATTTCATATGTGCGCACGG-3' (SEQ ID NO:07)	56
Standard control*	5'-TCCAGGTCCCCCGCATCCCGGATCC-3' (SEQ ID NO:08)	72

* splice site of mutated human β -globin gene
Mismatched nucleotides are underlined

Immunoblot and immunocytochemistry

[00161] Injected and control embryos were collected at 2-cell stage (43 hours post human chorionic gonadotropin (hCG) administration) and washed in PBS containing 3mg/mL polyvinylpyrrolidone (PVP). For immunoblot, injected and control embryos were lysed in RIPA buffer (50 mM Tris-HCl, pH7.5, 150 mM NaCl, 1 mM EDTA, 1% Nonident P-40, 2 mg/ml aprotinin, 2 mg/ml leupeptin, 1 mg/ml pepstatin, and 20 mg/ml phenylmethylsulfonyl) containing a phosphatase inhibitor cocktail (Roche), boiled in Laemli buffer, and stored at -80°C until lysates from 75 embryos were collected for each condition. Samples were loaded and analyzed by electrophoresis on 10% Tris-HCl polyacrylamide gel, semi-dry transferred onto nitrocellulose membrane (Bio-Rad), blocked in 0.1% Tween-20-1% Casein TBS Blocking Solution (Bio-Rad), incubated overnight in 1:250 diluted primary rabbit polyclonal anti-cyclin A2 antibody (Santa Cruz Biotechnologies, sc-751) at 4°C and in 1:2000 diluted secondary donkey anti-rabbit horseradish

peroxidase-linked antibody (Amersham, NA934V) for one hour at room temperature, and visualized using ECL Blotting Detection Reagent (Amersham). Immunocytochemistry was performed according to standard protocol. Briefly, embryos were fixed in 4% paraformaldehyde-PBS solution, permeabilized in 0.1% Triton X-PBS and treated with ImageIT FX Signal Enhancer solution (Invitrogen) at RT, incubated in 1:100 diluted primary antibody overnight at 4°C, in 1:10,000 diluted secondary antibody for one hour, followed by 3µM DAPI for 10 min., and were mounted in VectaShield Mounting Medium (Vector Laboratories, H-1000). Controls were performed in parallel with normal rabbit or mouse serum control. All antibodies were diluted in 1% BSA. Embryos were imaged by confocal microscopy using LSM 510 Confocal Laser Scanning Microscope or epifluorescence microscopy or the Axiovert 200 microscope equipped with an AxioCam digital camera (Zeiss) using fixed parameters and exposure times. Primary antibodies were purchased from Santa Cruz Biotechnologies (anti-cyclin A2 rabbit polyclonal (sc-751), normal rabbit IgG (sc-2027), mouse monoclonal anti-Oct4 (sc-5279), and normal mouse IgG (sc-2025). Secondary antibodies were purchased from Molecular Probes (Alexa Fluor 594 goat anti-rabbit IgG (A-11012) and goat-anti mouse IgG (A-11001).

mRNA synthesis by *in vitro* transcription.

- [00162] Full-length mouse cDNA clone, Oct4-pSPORT (clone ID 30019896) (Open Biosystems), and a plasmid encoding the fluorescence mitotic biosensor, a modified enhanced yellow fluorescence protein (mEYFP), were sequence-verified, linearized by restriction enzyme digest, and used as templates. 5' capped and polyadenylated mRNA transcripts were transcribed *in vitro* (mMessage and PolyA-Tail kits, Ambion), which were then quantitated by UV spectroscopy, and analyzed by electrophoresis to confirm size.

RNA sample preparation for gene chip experiments

- [00163] Samples containing 20 pooled injected or control mouse embryos were washed through 3 drops of PBS/PVP and collected for total RNA extraction and isolation (Picupure Total RNA Isolation Kit, Molecular Devices Corp.), to yield 10

μL of total RNA. 5 μL of total RNA, or the equivalent of 10 mouse embryos, was subjected to two rounds of amplification (WT-Ovation Pico system, Nugen) according to manufacturer's instructions. The quality of ssDNA resulting from the second round amplification was tested on the Bioanalyzer 2100 (Agilent), and a typical yield of 5-8 μg per sample was quantitated by the ND-1000 UV spectrophotometer (Nanodrop Technologies). Direct biotin labeling and fragmentation were performed (FL-Ovation cDNA Biotin Module (Nugen). Fragmented, labeled, ssDNA samples were submitted to Stanford University PAN Core Facility for hybridization to the GeneChip® Mouse Genome 430 2.0 Array (Affymetrix), and laser scanning.

Statistical analysis for gene chip experiments

[00164] Raw data from a total of 12 gene chips (3 *Ccna2*-MO and their uninjected controls, and 3 *Oct4*-MO and their uninjected controls) and were normalized by dChip (Li C & Wong WH (2003) in *The analysis of gene expression data: methods and software*. (Springer, New York), pp. 120-141). Unsupervised clustering analysis was performed for genes that have: 1) expression level greater than 500 Signal Intensity (SI) in at least 10 percent of the samples; 2) standard deviation to mean ratio > 0.4 and < 1000 across the samples. Subsequent analyses were not restricted to these criteria. The lists of differentially expressed genes and their ranking were generated by the method proposed by Johnson and Wong, which is based on fold change, logged fold change and unpaired t-statistic (Nicholas Johnson and W.H.W. (2007) Combining scientific and statistical significance in gene ranking. Unpublished.). Differential expression was defined by a threshold of 5 percent median false discovery rate (FDR) estimated from 300 random permutations across the samples. To find significantly enriched gene ontology (GO) terms (Ashburner M, *et al.* (2000) Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* 25, 25-29), each gene list was mapped to Entrez gene identification numbers (NCBI) and tested by GOSTAT PACKAGE (Beissbarth T & Speed TP (2004) Gostat: find statistically overrepresented Gene Ontologies within a group of genes. *Bioinformatics* 20, 1464-1465) against a set of "universe" genes defined by "P" call in at least 2 out of 3

treated samples or 4 out of 6 control samples by (MAS, Affymetrix). To estimate the FDR, 50 tests were performed with randomly generated lists of genes from the “universe” gene set, and the average number of enriched GO terms were calculated (Ashburner et al.; Beissbarth et al.). The cut off for p-value was chosen to reflect FDR of ~10 percent. Differentially expressed genes with possible Oct4-binding sites were identified by comparison with putative Oct4-regulated genes previously identified by Zhou *et. al.* (Zhou Q, Chipperfield H, Melton DA, & Wong WH (2007) A gene regulatory network in mouse embryonic stem cells. *Proc Natl Acad Sci U S A* 104, 16438-16443), followed by mapping of probe sets to Refseq using the BIOCONDUCTOR ANNOT PACKAGE (R.C. Gentleman VJC, D.M. Bates, B. Bolstad, M. Dettling, S. Dudoit, B. Ellis, L. Gautier, Y. Ge, J. Gentry, K. Hornik, T. Hothorn, W. Huber, S. Iacus, R. Irizarry, F. Leisch, C. Li, M. Maechler, A.J. Rossini, G. Sawitzki, C. Smith, G. Smyth, L. Tierney, J.Y.H. Yang, J. Zhang. (2004) Bioconductor: Open software development for computational biology and bioinformatics. *Genome Biology* 5, R80).

RT-PCR, q-PCR, and single embryo q-PCR analysis

[00165] Lysis buffer was added to PBS/PVP washed embryos (Cells-to-cDNA kit, Ambion) and samples were treated with 1 µL of DNase I. Reverse transcription (RT) was performed by using 1.0 µL of SuperScript III RT enzyme (200 U/µL) as per protocol (Invitrogen). Amplification of gene-specific product by TaqPolymerase High Fidelity Kit (Invitrogen) was performed on the thermocycler (Mastercycler gradient 5331, Eppendorf) as follows: 94.0°C for 2 min., 94.0°C for 15 sec., 60.0°C for 30 sec., 68.0°C for 45 sec, 68.0°C for 7 minutes for 50 cycles. (See Table 2 for all primer sequence and TaqMan probes). Single embryo qRT-PCR was performed using the Biomark 48.48 Dynamic Array system (Fluidigm, South San Francisco, CA). Single embryos were treated with acid tyrode, and collected in 10 µl reaction buffer, followed by preamplification as per manufacturer’s instructions (TaqMan Gene Expression Assay, Applied Biosystems; Table S8). Amplified cDNA was loaded onto a 48.48 Dynamic Array using the NanoFlex IFC controller (Fluidigm). Threshold cycle (C_T) as a measurement of relative fluorescence intensity was extracted from the BioMark PCR analysis

software (Fluidigm). All reactions were performed in duplicates or triplicates along with negative RT, PBS, and positive controls in at least three to five independent experiments. Data for each gene assayed were tested with a linear model (ANCOVA) in which $C_T \sim \beta_0 + \beta_1 * \text{Condition} + \beta_2 * C_T[\text{Gapdh}] + \beta_3 * C_T[\text{beta-actin}]$, where “Condition” referred to no injection or *Oct4* knockdown. C_T values were directly used in data analysis, as gene expression at the single-cell level has been shown to follow a lognormal distribution.

Table 2. Taqman® Probes and Gene-Specific Primers Used in q-PCR

Probe Name	Applied Biosystems Assay ID/ Catalog No.
Bmpr1a	Mm01208758_m1
Pou5f1	Mm00658129_gH
Klf9	Mm00495172_m1
Fgf4	Mm00438917_m1
Sall4	Mm01240680_m1
Bclaf1	Mm00464127_m1
Yy1	Mm00456392_m1
Pknox1	Mm00479320_m1
Gata4	Mm00484689_m1
Mta2	Mm00488671_m1
Rest1	Mm00803268_m1
Tcf15	Mm00626495_m1
Dppa5	Mm01171664_g1
Bmpr1a	Mm00477650_m1
Fgfr1l	Mm00475318_g1
Il17rd	Mm00460340_m1
Ubtf	Mm00456972_m1
Gtf3c4	Mm00557022_m1
Gtf3c2	Mm00510828_m1
Polr3e	Mm00491765_m1
Polr3a	Mm00805896_m1
Eif3b	Mm00659801_m1
Eif3s10	Mm00468721_m1
Piwil2	Mm00502383_m1
Eif4e	Mm00725633_s1
Polr2h	Mm01344328_g1
Eif2c5	Mm01305462_m1
Eif3c	Mm01278697_m1
Eif5b	Mm01227234_m1
Papola	Mm01334253_m1
Eif3e	Mm01700222_g1
Sox2	Mm03053810_s1
Mouse β -actin	4352341E
Mouse β -actin	4352933E
Mouse GAPDH	4352932E
Gene Specific Primers for Oct4 (302 bp product)	
Forward	GGCGTTCTCTTTGGAAAGGTGTT (SEQ NO:09)
Reverse	CTCGAACCACATCCTTCTCT (SEQ NO:10)
Gene Specific Primers for Ccna2 (212 bp product)	
Forward	GATAGATTCTCTCCTCCATG (SEQ NO:11)
Reverse	TCACACACTTAGTGTCTCTGG (SEQ NO:12)

EXAMPLE 1

CLINICAL AND EMBRYOLOGY DATA

[00166] Of all 1117 IVF treatments performed at Stanford University in 2005, 822 were fresh IVF cycles that used the patients' own oocytes (**FIG 1, panel A**). Based on our exclusion criteria, 157 cycles were excluded for a variety of medical and non-medical reasons.

[00167] The 157 cycles that were excluded consisted of: cancelled oocyte retrieval due to poor ovarian stimulation (63 cycles), cancelled embryo transfer due to complete lack of embryo development (8 cycles), cancelled embryo transfer due to unexpected medical or non-medical reasons (35 cycles), cycles that were not treated with gonadotropins (3 cycles), missing outcomes (8 cycles), and women being ≥ 45 years of age based on age alone (29 cycles). In our study, 160 patients underwent subsequent repeat cycles after a previously failed attempt for a total of 368 cycles. Of these, 126 patients had 2 IVF cycles performed, 25 patients had 3 cycles, 6 patients had 4 cycles, 1 patient had 5 cycles and 2 patients had 6 cycles performed in the same year at the same institution. 511 cycles (76.8%) with Day 3 embryo transfer and 154 cycles (23.2%) with Day 5 transfer, or a total of 665 IVF cycles, fulfilled the inclusion and exclusion criteria for analyses.

[00168] Clinical and embryology data on the remaining 665 cycles that satisfied inclusion and exclusion criteria, and their 4144 embryos, respectively, were analyzed to test the hypothesis that cohort-specific variables predict IVF cycle outcomes (**FIG 1**). Of those 4144 embryos, the number of blastomeres or cells on day 3 was recorded for 4002 embryos (96.6%). Overall, 38.8% had 8 cells, the developmentally appropriate cell number, while 18.2% of embryos had ≤ 4 cells, and 33.6% had 5-7 cells (**FIG 2**).

EXAMPLE 2

PROGNOSTIC SIGNIFICANCE AND CORRELATION OF VARIABLES

[00169] We systematically examined the association of each variable with IVF outcomes, and the correlation of each pair of variables. Pair-wise logistic regression tests confirmed many known prognostic variables, including female age, day 3

FSH, and the number of 8-cell embryos. However, in addition to these known prognostic variables, we observed that cohort-specific variables such as fertilization rate and the rate of cleavage arrest were also significantly associated with IVF cycle outcome ($p < 0.001$; Table 3). In contrast, except for male factor infertility ($p < 0.05$), none of the conventional clinical infertility diagnoses were significantly associated with IVF outcomes. Notably, despite a high degree of correlation between many variables and age or day 3 FSH level, which estimates ovarian aging, neither age nor day 3 FSH level was correlated to cohort-specific embryo parameters (Table 4). Collectively, these results suggest that determinants other than age-related mechanisms and clinical diagnoses impact cohort-specific embryo developmental competence.

Table 4: Correlation between each pair of variables

	Age	Gravidity	Maximum Day 3 FSH level	No. of oocytes	Average grade	Average grade of embryos transferred	No. of embryos	Percentage of 8-cell stage embryos	Percentage of embryos with ≤ 4 cells	Average cell no. of embryos	No. of embryos transferred	Percentage of transferred embryos at the 8-cell stage	Percentage of transferred embryos at the ≤ 4 -cell stage	Average cell no. of embryos transferred	Fertilization rate	No. of 8-cell embryos	No. of 8-cell embryos transferred
Age	1.00	0.19	0.13	-0.30	-0.01	0.12	-0.25	0.03	0.03	0.03	0.25	-0.20	0.15	-0.13	0.03	-0.17	-0.02
Gravidity	0.19	1.00	0.01	-0.06	-0.09	-0.03	-0.03	0.08	-0.03	0.11	0.06	0.03	0.15	0.02	0.03	0.06	0.09
Maximum Day 3 FSH level	0.13	0.01	1.00	-0.24	0.02	0.05	-0.20	-0.02	-0.05	0.02	0.00	-0.20	0.35	-0.24	0.03	-0.20	-0.10
No. of oocytes	-0.30	-0.06	-0.24	1.00	-0.02	-0.25	0.86	0.02	0.06	-0.39	0.13	0.04	-0.21	0.20	-0.27	0.68	0.38
Average grade	-0.01	-0.09	0.02	-0.02	1.00	0.84	0.02	0.35	0.29	-0.36	0.09	-0.42	0.41	0.77	-0.41	-0.32	-0.17
Average grade of embryos transferred	0.12	-0.03	0.05	-0.25	0.84	1.00	-0.28	-0.34	0.28	-0.36	0.05	-0.41	0.43	0.77	-0.41	-0.32	-0.17
No. of embryos	-0.25	-0.03	-0.20	0.86	0.02	-0.28	1.00	0.05	-0.51	0.05	-0.15	0.77	0.41	0.77	-0.41	-0.32	-0.17
Percentage of 8-cell stage embryos	0.03	0.08	-0.02	0.06	-0.35	-0.34	0.05	1.00	0.61	-0.15	0.77	0.41	0.77	0.41	0.77	-0.41	-0.32
Percentage of embryos with ≤ 4 cells	0.03	-0.03	-0.03	-0.05	0.29	0.28	-0.08	-0.51	1.00	-0.83	0.04	-0.41	0.87	-0.70	-0.12	-0.32	-0.35
Average cell no. of embryos	0.03	0.11	0.02	0.06	-0.39	-0.36	0.05	0.61	-0.83	1.00	-0.03	-0.22	0.43	0.77	-0.41	-0.32	-0.35
No. of embryos transferred	0.25	0.06	-0.03	0.13	0.04	0.09	0.14	-0.15	0.04	-0.03	1.00	-0.22	0.43	0.77	-0.41	-0.32	-0.35
Percentage of transferred embryos at the 8-cell stage	-0.20	0.03	-0.20	0.35	-0.24	-0.42	0.41	0.77	0.41	-0.22	0.77	0.41	0.77	0.41	0.77	-0.41	-0.32
Percentage of transferred embryos at the ≤ 4 -cell stage	0.15	0.03	0.00	-0.21	0.20	0.31	-0.28	0.41	0.87	-0.67	0.02	-0.48	0.58	0.14	0.64	0.70	0.41
Average cell no. of embryos transferred	-0.13	0.02	-0.05	0.27	-0.28	-0.41	0.35	-0.41	0.43	0.77	-0.41	0.77	0.41	0.77	-0.41	-0.32	-0.35
Fertilization rate	0.03	0.03	0.04	-0.06	0.04	0.06	0.04	0.04	0.04	-0.06	0.02	-0.48	0.58	0.14	0.64	0.70	0.41
No. of 8-cell embryos	-0.17	0.06	-0.20	0.68	-0.32	-0.41	0.77	-0.41	0.43	0.77	-0.41	0.77	0.41	0.77	-0.41	-0.32	-0.35
No. of 8-cell embryos transferred	-0.02	0.09	-0.10	0.38	-0.17	-0.32	-0.35	-0.41	0.43	0.77	-0.41	0.77	0.41	0.77	-0.41	-0.32	-0.35

EXAMPLE 3**THRESHOLDS OF NON-REDUNDANT, PROGNOSTIC VARIABLES DEFINING
HUMAN EMBRYO COHORT PHENOTYPES**

[00170] Sequential Multiple Additive Regression Tree (MART[®]) and Classification and Regression Tree (CART) analyses of all 30 variables (listed in Table 3 and its legend) determined that IVF cycle outcomes were most accurately predicted at ~70% by using only four non-redundant variables: total number of embryos, rate of cleavage arrest in an embryo cohort, the number of 8-cell embryos in a cohort, and day 3 FSH level. Remarkably, these four variables all describe the embryo cohort rather than individual embryos, and were more informative than age, clinical diagnoses, or any measures of the transferred embryos. Interestingly, the total number of embryos, day 3 FSH, and the number of 8-cell embryos depended on and thus captured the effects of many other variables. In contrast, the rate of cleavage arrest was independent of any of those known variables. (Details on MART[®] and CART analyses are reported in Table 5, and **FIG 3**).

Table 3: Association of each variable with pregnancy outcome

Variables	Estimate*	S.E.	P-Value
Patient Characteristics and Clinical Diagnoses[†]			
Age	-0.10	0.02	2.16E-007
Maximum Day 3 FSH level	-0.08	0.03	1.70E-003
Gravidity	0.036	0.066	5.86E-001
Male Factor (infertility diagnosis)	0.50	0.24	3.71E-002
IVF Cycle Characteristics			
Microdose lupron (flare) protocol	-1.14	0.24	2.53E-006
Antagonist protocol	-0.74	0.19	9.98E-005
Performance of ICSI	-0.15	0.16	3.47E-001
No. of oocytes	0.08	0.01	1.58E-009
Embryo Cohort Parameters			
Fertilization rate	1.24	0.36	5.37E-004
No. of embryos	0.14	0.02	2.67E-012
Average cell no. of embryos	0.29	0.06	6.34E-006
No. of 8-cell embryos	0.26	0.04	2.88E-012
Percentage of 8-cell embryos	0.76	0.28	5.75E-003
Cleavage arrest rate [‡]	-1.28	0.35	2.76E-004
Average grade of embryos	-0.091	0.17	5.88E-001
Parameters of Transferred Embryos			
Day 5 embryo transfer [§]	1.40	0.19	7.51E-013
No. of embryos transferred	0.0058	0.053	9.12E-001
Average cell no. of embryos transferred	0.47	0.07	2.19E-010
Percentage of transferred embryos at the 8-cell stage	1.33	0.21	5.35E-010
No. of 8-cell embryos transferred	0.41	0.08	4.40E-008
No. of embryos with ≤4 cells transferred	-2.14	0.49	1.06E-005
Average grade of embryos transferred	-0.52	0.17	2.61E-003
* Positive and negative estimates indicate association with positive and negative pregnancy outcomes, respectively.			
†Clinical infertility diagnoses that were not significantly associated with pregnancy outcome (p-value > 0.05) were not listed: uterine factor, polycystic ovarian syndrome, endometriosis, tubal ligation, tubal disease, hydrosalpinges, unexplained infertility, and "other diagnoses". Each IVF case may have more than one clinical infertility diagnosis.			
‡Cleavage arrest rate is defined as the percentage of embryos with 4 or fewer cells on Day 3 of <i>in vitro</i> culture.			
§Day 5 embryo transfer is arbitrarily listed under Parameters of Transferred Embryos. It can also be considered an Embryo Cohort Parameter, as it depends on the total number of embryos and the number of 8-cell embryos.			

Table 5: Models generated by MART[®] to identify non-redundant, prognostic variables

Model	Degree of Interaction	Learning Rate*	Tree Number [†]	Cross-validated (CV) Prediction Error Rate ^{††}
1	2	0.0100	2	0.300800
2	3	0.0010	3	0.300800
8	20	0.0010	4	0.308300
6	10	0.0001	4	0.315800
3	4	0.1000	30	0.323300
4	5	0.0020	16	0.323300
5	6	0.0100	16	0.330800
7	15	0.0001	3	0.330800

* One of the "boosting" tuning parameters in MART[®] to prevent "over-fitting" of the data²
[†] Number of trees constructed by MART[®]
^{††} 10-fold cross validation error

[00171] Of the prognostic thresholds identified, the most robust phenotypes are A1 and A2, and B1 and B2 (Table 6). Number of embryos < 6 or ≥ 6 is used by all 5 top CART models, defines all other phenotypes (B to F), and can be applied to all cases. Specifically, the phenotype defined by having fewer than 6 embryos, has an odds ratio of 3.9 for no pregnancy compared to cases with ≥ 6 embryos (95% Confidence Interval [CI], 2.8 to 5.5). Similarly, the next most robust phenotypes are defined by the number of embryos and cleavage arrest rate, such that for cases with ≥ 6 embryos, those with cleavage arrest rate > 14.6% are 3.0 times more likely to result in no pregnancy than those with cleavage arrest rate ≤ 14.6% (95% CI, 1.9 to 4.9).

Table 6: Prognostic thresholds defining cohort-specific phenotypes

	Embryos (No.)*	Cleavage Arrest (%)*	8-cell mbryo (No.)*	FSH (mIU/mL)*	Pregnancy - No. (%)†	No Pregnancy No. (%)‡	Applicable Cases - No. (%) §	No. Trees¶	Reference Condition	Odds Ratio	95% Confidence Interval (C.I.)
A1**	≥6				177 (57.7)	130 (42.3)	307 (46.2)	5			
A2	< 6				92 (25.7)	266 (74.3)	358 (53.8)	5	A1	3.9	(2.8, 5.5)
B1	≥6	≤ 14.6			112 (70.4)	47 (29.6)	159 (23.9)	4			
B2	≥6	> 14.6			65 (43.9)	83 (56.1)	148 (22.3)	4	B1	3.0	(1.9, 4.9)
B3	≥6	14.6 – 52.8			62 (47.3)	69 (52.7)	131 (19.7)	1	B1	2.6	(1.6, 4.3)
B4	≥6	≤ 52.8			174 (60.0)	116 (40.0)	290 (43.6)	n/a			
B5	≥6	> 52.8			3 (17.6)	14 (82.3)	17 (2.6)	1	B1 B4	10.6 6.7	(3.2, 49.6) (2.1, 30.9)
C1	≥6		≥ 2		157 (63.6)	90 (36.4)	247 (37.1)	1			
C2	≥6		< 2		20 (33.3)	40 (66.7)	60 (9.2)	1	C1	3.5	(1.9, 6.4)
D1	≥6	> 14.6	≥ 2		51 (53.1)	45 (46.9)	96 (14.4)	1			
D2	≥6	> 14.6	< 2		14 (26.9)	38 (73.1)	52 (7.8)	1	D1	3.0	(1.5, 6.5)
E1	≥6	> 14.6	≥ 2	≤ 4.6	14 (82.4)	3 (17.6)	17 (2.6)	1			
E2	≥6	> 14.6	≥ 2	> 4.6	34 (46.9)	37 (53.1)	71 (12.2)	1	E1	4.8	(1.4, 23.4)

* Cohort phenotypes defined by thresholds of non-redundant prognostic variables. Each set of conditions (A-E) use "AND" as the operator where more than one condition is listed.

† No. of cases that satisfy the threshold conditions and have pregnancy outcome. This percentage is calculated by using the No. Applicable Cases as denominator. In general, conditions that discriminate between pregnancy and no pregnancy outcomes more highly are more robust and are expected to be more useful in both clinical management and translational research.

‡ No. of cases that satisfy the threshold conditions and have no pregnancy outcome. This percentage is calculated by using the No. Applicable Cases as denominator.

§ The No. Applicable Cases is the total number of cases that satisfy the threshold conditions. This percentage is calculated by using the total number of cycles (665) as the denominator. In general, the larger the number of applicable cases, the more useful the set of conditions are for clinical management and counseling. However, for the purpose of translational research, conditions that define a smaller number of cases may have more specific correlates on a molecular level.

¶ No. Trees shows the number of CART trees that utilize each set of conditions. There are a total of 5 trees. (See Supplemental Results.) Increased utilization indicates "usefulness" or "robustness" of that particular set of conditions.

|| Reference condition against which the Odds Ratio and 95% C.I. for having no pregnancy is calculated.

** Conditions A-E are listed from most robust and "useful" to least "useful" based on: the number of trees that utilize each set of conditions, the number of applicable cases, and the odds ratio and 95% C.I.

[00172] In contrast, the rest of the thresholds listed in Table 6 are used by only 1 CART model each, and is applicable to fewer cases. However, as some of those phenotypes describe very specific subset of cases and have odds ratios that are highly discriminatory, they may be extremely useful depending on the clinical or

translational research context. For example, for cases with ≥ 6 embryos, having cleavage arrest rates of 14.6-52.8% and $> 52.8\%$ increase the odds of no pregnancy by 2.6 (95% CI 1.6 to 4.3) and 10.6 (95% CI 3.2 to 49.6), respectively, when compared to cases with cleavage rates of $\leq 14.6\%$.

EXAMPLE 3

ANALYSES BY MART[®] AND CART MODELS

[00173] To overcome the challenges presented by the highly interactive nature and potentially non-linear dependence amongst variables, and their multicollinearity, all 30 variables listed in Table 5 and its legend were analyzed by MART[®] to generate models that utilize non-redundant prognostic variables. Eight models were generated by MART[®] with increasing complexity of individual trees for boosting, while the learning rate was chosen to produce a minimum test error for the given complexity. Among them, the 3 top-ranking models had cross-validated (CV) prediction error rates of 0.301 to 0.308, which separated them well from the other five models, whose CV error rates ranged from 0.315 to 0.331 (Table 5). The third model, however, was based on very complex individual trees yet showed a larger CV error than the two preceding models; thus it was excluded from the analysis. The resulting two top models, consistent with characteristics of meaningful regression tree models, used very few trees and shared common features, such that the two models collectively used only 5 trees containing 5 variables, while some of the other models used up to 16 to 30 trees each.

[00174] These 5 non-redundant prognostic variables were: total number of embryos, the rate of cleavage arrest, the number of 8-cell embryos, day 3 FSH level, and the number of 8-cell embryos transferred. In each model, removal of each of the first four variables while keeping all other parameters constant, increased the error rate of the model, thus confirming their significant contribution. However, removal of the number of 8-cell embryos transferred, did not alter the error rate (data not shown), which suggested that this variable was less relevant than others.

[00175] 24 models representing all possible combinations of these 5 variables were analyzed by CART to further define thresholds that have prognostic significance.

The top 5 trees generated by CART had superior prediction scores (0.6828 to 0.6950) compared to the rest of the models (0.565 to 0.6700). As there were shared features, these top 5 tree models utilize mostly the same threshold conditions for 4 variables, while the variable, number of 8-cell embryos transferred, was not utilized by any of these top models. Therefore, IVF cycle outcomes could be most accurately predicted at ~70% by using only four non-redundant variables that are more informative than age, clinical diagnoses, or any measures of the transferred embryos.

[00176] In order to understand which factors amongst patient characteristics, diagnoses, and IVF treatment characteristics, in turn determined these four non-redundant, cohort-specific prognostic variables, tree models were constructed by MART to represent the dependence of each of these four prognostic variables. The total number of embryos, day 3 FSH, and the number of 8-cell embryos depended on and thus captured the effects of many other variables (**FIG 3**). In contrast, the rate of cleavage arrest was independent of any of those known variables.

EXAMPLE 4

MORPHOLINO MEDIATED GENE KNOCKOUT STUDY

[00177] Here, we provide proof-of-concept of the efficiency and specificity of MO-mediated gene knockdown by testing the procedure on the *Ccna2* gene. We then report the novel role of Oct4 that was revealed by MO-mediated gene knockdown. *Ccna2*, the gene encoding cell cycle regulator cyclin A2, has been suggested as an important transcriptional regulator in embryonic genome activation (Hara KT, *et al.* (2005) *Dev Biol* 286, 102-113), a critical developmental milestone at the 1- to 2-cell stages for which few clear mechanisms or regulators have emerged. Consistent with the literature, MO-mediated *Ccna2* knockdown decreased cyclin A2 protein expression. In addition, our results showed that cyclin A2 is required for development beyond the 2-cell stage (**FIG 4**, panels A-C, **FIG 5**, panels A-D, Tables 1 and 7). MOs block translation of transcripts by steric hindrance in an efficient and gene-specific manner, which has been well established in zebrafish and other model organisms (Gore AV, *et al.* (2005) *Nature* 438, 1030-1035; Imai *et al.*, (2006). *Science* 312, 1183-1187; Sumanas S & Larson JD (2002) *Brief Funct*

Genomic Proteomic 1, 239-256; Yamada L, *et al.* (2003) *Development* 130, 6485-6495). Most importantly, MOs mediate rapid knockdown of transcripts regardless of their maternal or embryonic origin, before activation of downstream genes can provide partial “rescue” of the phenotype. Unlike siRNAs, MOs function independently of endogenous pathways, so the MO-mediated translational block should not limit knockdown efficiency or interfere with the endogenous mRNA degradation machinery.

[00178] By combining MO-mediated gene knockdown with global gene expression profiling and single-embryo level quantitative RT-PCR (q-PCR), we determined the influence of Oct4 on gene expression, and analyzed the Oct4-regulated gene network in the early embryo (**FIG 6** and **FIG 7**). Consistent with the literature, we confirmed *Oct4* gene expression at the 1-cell stage (**FIG 8**). After 1-cell embryos were microinjected with 0.6 mM *Oct4*-MO, the rate of developmental arrest at the 1- to multicell stages was dramatically higher than that observed for uninjected and mismatch (*Oct4*-MM) controls (**FIG 9**, panels A and B). Of embryos injected with *Oct4*-MO that reached the multicell stage, $86.8 \pm 8.3\%$ arrested and did not form morulae, compared to $10.5 \pm 10.5\%$ embryos injected with *Oct4*-MM ($p < 0.01$; data not shown). Most remarkably, none of the *Oct4*-MO-injected embryos developed to blastocysts, compared to relatively high blastocyst rates of *Oct4*-MM-injected and uninjected embryos ($p < 0.01$; **FIG 9**, panel C). Further, the specificity of *Oct4*-MO is supported by the direct relationship between the phenotype severity and presumed “gene-dosage” as titrated by *Oct4*-MO concentration (**FIG 9**, panels D and E). Oct4 protein expression was indeed reduced in *Oct4*-MO-injected embryos at the 4-cell (**FIG 10**) and multi-cell stages (**FIG 11**, panel A, and **FIG 12**). Injection of another MO, targeting an intron-exon boundary in *Oct4*, confirmed that disruption of Oct4 function is detrimental to development before the blastocyst stage (**FIG 13**, panels A-C).

[00179] The critical function of Oct4 at the 1- to 2-cell transition was embryo-autonomous. The effects of Oct4 knockdown could not be rescued by media conditioned by uninjected embryos or the *in vivo* environment provided by transferring injected embryos to oviducts of appropriately timed surrogate mothers (data not shown). Co-injection of low (3.6 ng/ μ L) or high (3.6 ng/ μ L)

concentrations of unaltered, full-length *Oct4* mRNA with 0.6 mM *Oct4*-MO resulted in a decrease in the percentage of embryos arresting at the 1- to multicell stages, compared to co-injection of control mRNA encoding a modified enhanced yellow fluorescent protein (mEYFP) (**FIG 11**, panel *B*). No embryos co-injected with *Oct4* mRNA arrested at the multicell stage, while $80.0 \pm 5.2\%$ arrested after co-injection of mEYFP mRNA ($p < 0.01$; data not shown). Thus, co-injection of *Oct4* mRNA, but not control mRNA, partially rescued the *Oct4*-MO-induced multicell stage arrest phenotype. The failure of achieving full rescue of the phenotype may be due to insufficient *Oct4* expression after co-injection of *Oct4* mRNA at the low concentration, inappropriately high *Oct4* expression at the high concentration, or relative instability of the *in vitro* transcribed *Oct4* mRNA compared to *Oct4*-MO; further, these possibilities are not mutually exclusive. Hence, we next tested whether *Oct4* over-expression itself would interfere with development.

[00180] Injection of *Oct4* mRNA resulted in over-expression as quantified by q-PCR, but did not cause ectopic expression of Oct4 protein (data not shown). Oct4 over-expression indeed induced developmental arrest in a dosage-dependent manner, while injection of comparable or greater amounts of mEYFP mRNA interfered minimally with blastocyst development (**FIG 11**, panel *C*). The “gene dosage” effect of *Oct4* RNA injection may be due to enhanced Oct4 functions in transcriptional regulation, or Oct4 over-production may allow non-specific promoter-binding, or extra Oct4 causes inappropriate sequestration and subsequent inactivation of co-factors. Collectively, these data definitively showed *Oct4* expression was required, and that its correct level was critical to early embryo development, just as pluripotency of ESCs depends on a precise range of *Oct4* expression levels (Stefanovic S & Puceat M (2007) *Cell Cycle* 6, 8-10).

Table 7. Summary of the Number of Embryos Tested and the Number of Experiments Performed for Each Condition

Experiment	Condition	No. Injected Embryos	No. Uninjected Embryos	No. experiments
<i>Ccna2</i> knockdown	<i>Ccna2</i> -MO 0.75 mM	94	65	8
	<i>Ccna2</i> -MM 0.75 mM	75	101	9
<i>Ccna2</i> -MO Immunocytochemistry	<i>Ccna2</i> -MO 0.75 mM	26	28	3
<i>Ccna2</i> "gene dosage"	<i>Ccna2</i> -MO, 0.5 mM	39	30	3
	<i>Ccna2</i> -MO, 0.25 mM	50	29	3
<i>Oct4</i> knockdown	<i>Oct4</i> -MO, 0.60 mM	41	37	4
	<i>Oct4</i> -MM, 0.60 mM	57	30	3
	<i>Oct4</i> -MO, 0.40 mM	79	25	3
	<i>Oct4</i> -MO, 0.20 mM	63	30	3
	<i>Oct4</i> E4-MO, 0.60 mM	35	30	3
	<i>Oct4</i> E4-MM, 0.60 mM	32	30	3
<i>Oct4</i> -MO Immunocytochemistry	<i>Oct4</i> -MO, 0.60 mM	32	22	3
	<i>Oct4</i> -MM, 0.60 mM	20	20	3
Rescue	0.036µg/µl <i>Oct4</i> mRNA+ 0.60mM <i>Oct4</i> -MO	32	30	3
	0.036µg/µl <i>EYFP</i> mRNA + 0.60mM <i>Oct4</i> -MO	72	30	3
Overexpression	0.09µg/µl <i>Oct4</i> mRNA	24	20	3
	0.036µg/µl <i>Oct4</i> mRNA	33	25	3
	3.6ng/µl <i>Oct4</i> mRNA	35	30	3
	0.09µg/µl <i>EYFP</i> mRNA	41	20	2
<i>Oct4</i> -MO/Conditioned Media	<i>Oct4</i> -MO, 0.60 mM, conditioned media	20	20	2
	<i>Oct4</i> -MO, 0.60 mM, unconditioned media	20	20	2
<i>Ccna2</i> -MO gene chip	<i>Ccna2</i> -MO, 0.75 mM	60	60	3
<i>Oct4</i> -MO gene chip	<i>Oct4</i> -MO, 0.6 mM	60	60	3
<i>Oct4</i> -MO single embryo QPCR	<i>Oct4</i> -MO, 0.60 mM, 2-cell stage	14-20	14-20	3-5
	Control-MO, 0.60 mM, 2-cell stage	5-10	n/a	2

EXAMPLE 5**GENE REGULATION BY OCT4**

[00181] To dissect the mechanisms of Oct4 function, we compared the global gene expression profile of *Oct4* knockdown embryos to the effects of *Ccna2* knockdown and to uninjected controls at the mid-2-cell stage. The goal was to identify differential gene expression that coincides with the first major wave of embryonic genome activation at the mid-2-cell stage (Hamatani T, Carter MG, Sharov AA, & Ko MS (2004) *Dev Cell* 6, 117-1317; Wang QT, *et al.* (2004) *Dev Cell* 6, 133-

1448) (Tables 8, A and B, 9, A and B). Analysis by an unsupervised algorithm showed that the embryo samples clustered according to the experimental conditions, which further supported the specific and non-random effects of gene knockdown (FIG 14, panel A). At an arbitrary threshold false detection rate (FDR) of 0.05, the Oct4-regulated gene set is five times larger compared to the number of genes changing expression in response to cyclin A2 knockdown (FIG 14, panel B). The different sizes of the under-expressed versus over-expressed *Oct4*-knockdown gene sets suggest that *Oct4* may be predominantly activating rather than repressing transcription (FIG 14, panel B). Some of the Oct4 candidate target genes have previously been identified as putative Oct4 targets based on mESC chromatin immunoprecipitation (ChIP) data or genomic sequence analysis of Oct4-binding sites (Zhou Q, Chipperfield H, Melton DA, & Wong WH (2007) *Proc Natl Acad Sci U S A* 104, 16438-16443) (Table 10).

[00182] The list of cyclin A2-regulated genes was rich in genes encoding factors for chromatin modification and remodelling ($p=0.005$), nucleotide metabolism ($p=0.01$), and chromosome organization ($p=0.01$; Tables 11 A and B). Oct4-regulated genes were significantly enriched for translation ($p=1.1 \times 10^{-4}$) and RNA processing functions ($p=3.0 \times 10^{-5}$) (Tables 11, C and D). Comparison of our data with published Oct4-regulated networks in mouse ESCs indicates that *Oct4* has distinct and specific post-transcriptional and translational regulatory functions mediated by its control of genes encoding subunits in eukaryotic translation initiation factors (Eif), including *Eif3c*, and *Eif3b*. Interestingly, these two Eif subunits are evolutionarily conserved from yeast to human, and are amongst the six subunits comprising the functional core of mammalian Eif3, the largest of the Eif complexes (Masutani M, Sonenberg N, Yokoyama S, & Imataka H (2007) *Embo J* 26, 3373-3383). In addition to its embryo-specific function, Oct4 also controls the expression of genes encoding post-transcriptional regulators Dppa5 and Piwil2 (also known as Mili), as it does in ESCs (data not shown). Dppa5 is an embryo-, germ cell- and ESC-specific RNA-binding protein whose role in maternal-embryonic transition is not known. However, Piwil2 and its bound pi-RNAs, are known for their role in regulating retrotransposons in the fully-grown mouse oocyte (Watanabe T, *et al.* (2008) *Nature* 453, 539-543.26). Collectively, our data indicate

that the Oct4 has a distinct, developmental stage-specific role in controlling genes encoding post-transcriptional regulators, in addition to its conserved functions shared amongst pluripotent cell types.

Table 8A: Genes That Have Higher Expression Levels In *Oct4*-MO-Injected Compared To Uninjected Embryos (Following the Examples Section)

Table 8B: Genes That Have Lower Expression Levels In *Oct4*-MO-Injected Compared To Uninjected Embryos (Following the Examples Section)

Table 9A: Genes That Have Higher Expression Levels In *Ccna2*-MO-Injected Compared To Uninjected Embryos (Following the Examples Section)

Table 9B: Genes That Have Lower Expression Levels In *Ccna2*-MO-Injected Compared To Uninjected Embryos (Following the Examples Section)

Table 10: Oct4 Candidate Target Genes That Have Putative Oct4 Binding Sites Based on Genomic Sequence Analysis or Mouse ESC Chromatin Precipitation (ChIP) Data

Downregulated in *Oct4* knockdown embryos:

1700021F05Rik: RIKEN cDNA 1700021F05 gene
 2900073H19Rik: RIKEN cDNA 2900073H19 gene
 Arhgap8: Rho GTPase activating protein 8
 Cbfa2t2h: core-binding factor, runt domain, alpha subunit 2, translocated to, 2 homolog (human)
 Cdt1: chromatin licensing and DNA replication factor 1
 Didol1: death inducer-obliterators 1
 Dpm1: dolichol-phosphate (beta-D) mannosyltransferase 1
 Dppa5: developmental pluripotency associated 5
 Eif4e2: eukaryotic translation initiation factor 4E member 2
 Elovl6: ELOVL family member 6, elongation of long chain fatty acids (yeast)
 Ets5: ets variant gene 5
 Fkbp4: FK506 binding protein 4
 Gnb2l1: guanine nucleotide binding protein (G protein), beta polypeptide 2 like 1
 Hexb: hexosaminidase B
 Igf2bp1: insulin-like growth factor 2 mRNA binding protein 1
 Klf9: Kruppel-like factor 9
 Mkrn1: makorin, ring finger protein, 1
 Mtf2: metal response element binding transcription factor 2
 Mygl1: melanocyte proliferating gene 1
 Pa2g4: proliferation-associated 2G4
 Pitpnc1: phosphatidylinositol transfer protein, cytoplasmic 1
 Ppm1a: protein phosphatase 1A, magnesium dependent, alpha isoform
 Rest: RE1-silencing transcription factor

Rif1 /// LOC671598: Rap1 interacting factor 1 homolog (yeast) /// similar to Telomere-associated protein RIF1 (Rap1-interacting factor 1 homolog) (mRif1)
 Slc19a3: solute carrier family 19 (sodium/hydrogen exchanger), member 3
 Slc22a12: solute carrier family 22 (organic anion/cation transporter), member 12
 Slc25a36: solute carrier family 25, member 36
 Tfrc: transferrin receptor
 Tnpo3: transportin 3
 Ube2h: ubiquitin-conjugating enzyme E2H
 Ube2o: ubiquitin-conjugating enzyme E2O
 Zfp297b: zinc finger protein 297B

Upregulated in *Oct4* knockdown embryos

BC022623: cDNA sequence BC022623
 2810429O05Rik: RIKEN cDNA 2810429O05 gene
 Blcap: bladder cancer associated protein homolog (human)
 Icosl: icos ligand
 Sox2: SRY-box containing gene 2
 Bcas2: breast carcinoma amplified sequence 2
 Ldlr: low density lipoprotein receptor
 Zfp219: zinc finger protein 219
 Arl4c /// LOC632433: ADP-ribosylation factor-like 4C /// similar to ADP-ribosylation factor-like protein 7
 Tcl1: T-cell lymphoma breakpoint 1
 Dcpl1a: decapping enzyme
 Nes: nestin
 Rbpsiuh: Recombining binding protein suppressor of hairless (*Drosophila*)

Table 11A. Functional Categories That Were Enriched In Downregulated Genes In The *Ccna2* Knockdown Model

GOBPID	P-value	Term
GO:0050875	1.1E-3	cellular physiological process**
GO:0016568	4.0E-3	chromatin modification*
GO:0042254	4.0E-3	ribosome biogenesis and assembly**
GO:0006338	4.8E-3	chromatin remodeling*
GO:0007028	0.01	cytoplasm organization and biogenesis**
GO:0044237	0.01	cellular metabolism**
GO:0007582	0.01	physiological process
GO:0006139	0.01	nucleobase, nucleoside, nucleotide and nucleic acid metabolism*
GO:0008152	0.01	metabolism
GO:0006376	0.01	mRNA splice site selection*
GO:0006413	0.01	translational initiation**
GO:0007001	0.01	chromosome organization and biogenesis (sensu Eukaryota)*
GO:0007001		
GO:0006996	0.01	organelle organization and biogenesis*
GO:0007046	0.01	ribosome biogenesis**
GO:0043170	0.02	macromolecule metabolism**
GO:0051276	0.02	chromosome organization and biogenesis
GO:0006325	0.02	establishment and/or maintenance of chromatin architecture
GO:0006323	0.02	DNA packaging*
GO:0016043	0.02	cell organization and biogenesis

GO:0044238	0.03	primary metabolism**
GO:0000245	0.03	spliceosome assembly
GO:0006486	0.03	protein amino acid glycosylation
GO:0009059	0.03	macromolecule biosynthesis**
GO:0043413	0.03	biopolymer glycosylation
GO:0009101	0.04	glycoprotein biosynthesis
GO:0006364	0.04	rRNA processing
GO:0006412	0.04	protein biosynthesis**
GO:0006259	0.04	DNA metabolism
GO:0016072	0.04	rRNA metabolism**
GO:0009100	0.05	glycoprotein metabolism

Notes:

**Indicates the categories that were enriched in the *Oct4* data set as well.

* indicates functional categories that were specifically enriched in *Ccna2* knockdown.

Table 11B. Functional Categories That Were Enriched In Upregulated Genes In The *Ccna2* Knockdown Model

GOBPID	P-value	Term
GO:0044262	7.1E-4	cellular carbohydrate metabolism

Table 11C. Functional Categories That Were Enriched In Downregulated Genes In The *Oct4* Knockdown Model

GOBPID	P-value	Term
GO:0006412	1.8E-7	protein biosynthesis**
GO:0009059	1.3E-6	macromolecule biosynthesis**
GO:0006413	3.6E-6	translational initiation**
GO:0009058	1.8E-5	biosynthesis
GO:0044249	2.1E-5	cellular biosynthesis
GO:0042254	2.3E-5	ribosome biogenesis and assembly**
GO:0006396	3.0E-5	RNA processing*
GO:0007046	3.3E-5	ribosome biogenesis**
GO:0016070	3.9E-5	RNA metabolism**
GO:0007028	7.0E-5	cytoplasm organization and biogenesis**
GO:0043037	1.1E-4	Translation*
GO:0043170	1.2E-4	macromolecule metabolism**
GO:0019538	1.6E-4	protein metabolism
GO:0044238	3.1E-4	primary metabolism**
GO:0044267	3.1E-4	cellular protein metabolism
GO:0050875	3.3E-4	cellular physiological process**
GO:0044237	5.3E-4	cellular metabolism**
GO:0044260	5.7E-4	cellular macromolecule metabolism

Note:

**Indicates categories that were enriched in the *Ccna2* data set as well.

* indicates functional categories that were specifically enriched in *Oct4* knockdown.

Table 11D. Functional Categories That Were Enriched In Upregulated Genes In The *Oct4* Knockdown Model

GOBPID	P-value	Term
GO:0044262	2.0E-4	Alcohol metabolism

EXAMPLE 6

OCT4 FUNCTION AT THE 1- TO 2-CELL STAGE

- [00183] In order to understand the role of Oct4 in reprogramming the early embryo, we examined its role in embryonic genome activation and maternal transcript degradation. Overall, Oct4 regulates gene expression pertinent to basic machinery required for the entire spectrum of gene regulation, including transcription involving all three RNA polymerases, translation, RNA processing such as regulation of polyadenylation, and mRNA degradation proteins (Table 12). High levels of mRNA from developmental genes, such as *Six1*, *Nestin*, and *Hoxa3*, indicated that Oct4 was required for their repression, while excessive levels of maternal transcripts that would normally be rapidly degraded, such as *Zar1* and *Nobox1*, indicated that Oct4 knockdown interfered with the mRNA degradation machinery. Thus, Oct4 has developmental stage- and cell-specific functions, and has an important role in the processes that mark maternal-embryonic transition.
- [00184] To further define the Oct4-regulated gene network, we selected 42 genes representing transcriptional, post-transcriptional and signaling functions for q-PCR assays. We analyzed RNA from single *Oct4*-MO-injected and control embryos and focused on genes that were under-expressed in *Oct4* knockdown. After removing data related to 3 genes for which there were technical difficulties, expression changes of 39 genes were appropriately measured based on analysis using a linear model (See Methods and Materials). Of those, 34 or ~ 87% showed altered expression levels in *Oct4* knockdown in the expected directions (**FIG 14**, panel C and **FIG 15**), while 5 genes, including *Sox2*, did not change (data not shown). 21 of the 34 genes, or ~ 62%, showed statistically significant differential expression by q-PCR at $p < 0.05$ or less, while injection of a control MO did not alter expression of any of the genes assayed (data not shown). Thus, we have proven that Oct4 directly or indirectly regulates genes encoding the entire spectrum of transcriptional and post-transcriptional regulators at the 1- to 2-cell stages.
- [00185] Our single-embryo data allowed us to go beyond simply validating our gene chip data. Methods using samples comprised of pooled cells or embryos, generate

relative gene expression that represents an average of all cells assayed, but they cannot discern between genes that are consistently differentially regulated versus those with a tendency towards stochastic changes; similarly, rare outlier embryos expressing unique transcriptomes are not recognized (Bengtsson M, Stahlberg A, Rorsman P, & Kubista M (2005) *Genome Res* 15, 1388-1392; Chang HH, *et al.* (2008) *Nature* 453, 544-547; Warren L, Bryder D, Weissman IL, & Quake SR (2006) *Proc Natl Acad Sci U S A* 103, 17807-17812). By analyzing quantitative expression data at the single-embryo level, we were able to make this discrimination. We presume genes whose relative expression is consistent amongst single embryos have a higher likelihood to be essential nodes in a gene regulatory network, which is expected to respond to perturbations in a consistent and predictable manner. The gene set was restricted to genes whose differential expression (represented by the difference in threshold cycles, ΔC_T) ΔC_T is greater than expression differences amongst single embryos (represented by standard error of the mean, s.e.m.). We propose a hierarchy in the Oct4-regulated gene network in which 29 genes are ordered based on their increasing s.e.m., or inter-embryo variation and presumed decreasing biological significance in this network (**FIG 16**, panel A). Taken together, we have identified and ranked potential key nodes of this network in a quantitative fashion.

[00186] The data shows that in the unique developmental context of maternal-embryonic transition, concomitant with massive mRNA degradation and dramatic reprogramming, Oct4 controls the expression of many transcriptional regulators. Oct4 also maintains the expression of many genes, such as *Eif3c*, *Papola*, *Piwil2*, *Eif3b*, *Eif4e*, *Rbm3* and *Cpsf4*, that are involved in the post-transcriptional control. Through its influence on both the transcriptional and post-transcriptional regulators, Oct4 can directly or indirectly affect many essential processes, such as chromatin remodelling, epigenetic regulation, apoptosis, cell cycle regulation, and signalling, during early developmental program (**FIG 16**, panel B).

Table 12: Candidate Oct4-Regulated Genes That Function In Transcription, Translation, RNA Processing, Chromatin Remodeling, Signaling, Apoptosis And The Cell Cycle.

	Gene Symbol	Fold change	False detection rate (FDR)	T. P-value
POL I transcription				
upstream binding transcription factor, RNA polymerase I	Ubtf	0.49	0.02667	0.00424
arginyl-tRNA synthetase	Rars	0.47	0.03200	0.00131
POL II transcription				
BTAF1 RNA polymerase II, B-TFIID transcription factor-associated, (Mot1 homolog, <i>S. cerevisiae</i>)	Btaf1	0.12	0.03834	0.00833
mediator of RNA polymerase II transcription, subunit 8 homolog (yeast)	Med8	0.25	0.03405	0.00142
transcription elongation factor B (SIII), polypeptide 3	Tceb3	0.29	0.01471	0.00001
TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor	Taf9	0.30	0.02151	0.00027
polymerase (RNA) II (DNA directed) polypeptide H	Polr2h	0.42	0.02586	0.00859
elongation factor RNA polymerase II 2	Eli2	0.49	0.03696	0.00098
cofactor required for Sp1 transcriptional activation, subunit 2	Med14	0.50	0.03794	0.00156
POL III transcription				
general transcription factor IIIC, polypeptide 4	Gtf3c4	0.07	0.00000	0.00002
general transcription factor IIIC, polypeptide 5	Gtf3c5	0.34	0.03209	0.00212
BRF2, subunit of RNA polymerase III transcription initiation factor, BRF1-like	Brf2	0.34	0.04385	0.05104
general transcription factor IIIC, polypeptide 2, beta	Gtf3c2	0.35	0.03314	0.00047
general transcription factor III C 1	Gtf3c1	0.47	0.03303	0.00236
polymerase (RNA) III (DNA directed) polypeptide A	Polr3a	0.55	0.04421	0.01694
polymerase (RNA) III (DNA directed) polypeptide E	Polr3e	0.56	0.03366	0.00134
Translational control, RNA processing, and posttranscriptional regulation				
Translational initiation				
eukaryotic translation initiation factor 2C, 5	Eif2c5	0.09	0.01370	0.00022
eukaryotic translation initiation factor 5B	Eif5b	0.14	0.03802	0.00002
eukaryotic translation initiation factor 3, subunit 9 (eta)	Eif3b	0.29	0.03803	0.00358
eukaryotic translation initiation factor 3, subunit 8	Eif3c	0.36	0.00000	0.00016
eukaryotic translation initiation factor 3, subunit 3 (gamma)	Eif3h	0.42	0.03226	0.00485
eukaryotic translation initiation factor 3, subunit 4 (delta)	Eif3g	0.47	0.04924	0.01507
integrin beta 4 binding protein	Eif6	0.53	0.02756	0.00236
eukaryotic translation initiation factor 4, gamma 1	Eif4g1	0.59	0.04065	0.00041
eukaryotic translation initiation factor 2, subunit 1 alpha	Eif2s1	0.60	0.04879	0.00311
eukaryotic translation initiation factor 4E /// hypothetical LOC630527	Eif4e	0.65	0.03742	0.00177
eukaryotic translation initiation factor 3, subunit 10 (theta)	Eif3s10	0.66	0.04070	0.01798
Translational repression				
piwi-like homolog 2 (<i>Drosophila</i>)	Piwi2	0.34	0.02778	0.00051
eukaryotic translation initiation factor 4E member 2	Eif4e2	0.46	0.02553	0.01084

3' end processing				
poly (A) polymerase alpha	Papola	0.10	0.02475	0.00014
cleavage and polyadenylation specific factor 6	Cpsf6	0.20	0.03256	0.00009
cleavage and polyadenylation specific factor 4	Cpsf4	0.48	0.03542	0.00233
Translational termination				
eukaryotic translation termination factor 1	Etf1	0.36	0.02717	0.00207
Nonsense mediated decay (NMD)				
eukaryotic translation initiation factor 3, subunit 6	Eif3e	0.21	0.00000	0.00000
Other RNA binding and processing				
integrator complex subunit 7	Ints7	0.10	0.02757	0.00004
RNA binding motif protein 3	Rbm3	0.20	0.03791	0.00037
RNA binding motif protein 4	Rbm4	0.42	0.04967	0.00236
developmental pluripotency associated 5	Dppa5	0.46	0.03672	0.00786
RNA binding motif protein 5	Rbm5	0.52	0.03415	0.00245
proliferation-associated 2G4	Pa2g4	0.70	0.04956	0.01729
Pre-mRNA processing and transport				
Heterogeneous nuclear ribonucleoprotein U		0.18	0.03688	0.00020
PRP4 pre-mRNA processing factor 4 homolog (yeast)	Prpf4	0.34	0.03828	0.00128
RNA, U transporter 1	Snupn	0.35	0.01681	0.00227
PRP38 pre-mRNA processing factor 38 (yeast) domain containing B	Prpf38b	0.48	0.04103	0.00548
protein phosphatase 1G (formerly 2C), magnesium-dependent, gamma isoform	Ppm1g	0.48	0.02463	0.00012
heterogeneous nuclear ribonucleoprotein A/B	Hnrpab	0.58	0.03233	0.00100
heterogeneous nuclear ribonucleoprotein A1	Hnrpa1	0.63	0.04981	0.00961
heterogeneous nuclear ribonucleoprotein K	Hnrpk	0.64	0.04473	0.01239
heterogeneous nuclear ribonucleoprotein M	Hnrpm	0.63	0.04884	0.00724
Transcriptional regulation				
homeodomain interacting protein kinase 3	Hipk3	0.17	0.04003	0.00031
Kruppel-like factor 9	Klf9	0.17	0.00000	0.00020
nuclear respiratory factor 1	Nrf1	0.20	0.01923	0.00073
integrator complex subunit 4	Ints4	0.34	0.03520	0.00080
homeodomain interacting protein kinase 1	Hipk1	0.35	0.01786	0.00073
Pbx/knotted 1 homeobox	Pknox1	0.37	0.03796	0.00484
proline, glutamic acid and leucine rich protein 1	Pelp1	0.40	0.02415	0.00573
core-binding factor, runt domain, alpha subunit 2, translocated to, 2 homolog (human)		0.41	0.04076	0.02903
YY1 transcription factor	Yy1	0.43	0.03853	0.00074
GATA binding protein 4	Gata4	0.45	0.03963	0.00325
metastasis-associated gene family, member 2	Mta2	0.51	0.03395	0.00326
BCL2-associated transcription factor 1	Bclaf1	0.54	0.02664	0.00242
RE1-silencing transcription factor	Rest	0.62	0.03827	0.00983
ets variant gene 5	Etv5	0.50	0.04900	0.00576
makorin, ring finger protein, 1	Mkrn1	0.58	0.03793	0.00310
Chromatin remodeling, epigenetic regulation				
jumonji domain containing 1B	Kdm3b	0.14	0.03770	0.00132

jumonji, AT rich interactive domain 1C (Rbp2 like)	Kdm5c	0.23	0.01408	0.00075
sirtuin 1 ((silent mating type information regulation 2, homolog) 1 (S. cerevisiae)	Sirt1	0.28	0.01136	0.00002
methyltransferase like 2	Mettl2	0.37	0.03333	0.00010
jumonji domain containing 1A	Kdm3a	0.38	0.04874	0.00764
SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily c, member 1	Rsc8	0.41	0.03665	0.01049
jumonji, AT rich interactive domain 1B (Rbp2 like)	Kdm5b	0.57	0.03738	0.00388
eukaryotic translation initiation factor 2, subunit 3, structural gene X-linked /// similar to eukaryotic translation initiation factor 2, subunit 3, structural gene X-linked		0.57	0.04905	0.00183
Histone H3-specific HAT activity				
general transcription factor IIIC, polypeptide 4	Gtf3c4	0.07	0.00000	0.00002
general transcription factor IIIC, polypeptide 5	Gtf3c5	0.34	0.03209	0.00212
GCN5 general control of amino acid synthesis-like 2 (yeast)	Gcn5	0.34	0.02525	0.00159
general transcription factor IIIC, polypeptide 2, beta	Gtf3c2	0.35	0.03314	0.00047
general transcription factor III C 1	Gtf3c1	0.47	0.03303	0.00236
H3 histone, family 3A	H3f3a	0.53	0.04006	0.00865
Signaling pathways (Fgf4, Bmp, Toll, Egf, Mapk, Erk, Igf2, Amp)				
fibroblast growth factor 4	Fgf4	0.09	0.02591	0.00088
signaling intermediate in Toll pathway-evolutionarily conserved	Ecsit	0.17	0.01935	0.00000
protein kinase, AMP-activated, gamma 1 non-catalytic subunit	Prkag1	0.19	0.03786	0.00010
insulin-like growth factor 2 mRNA binding protein 1	Imp1	0.26	0.04930	0.00047
bone morphogenetic protein receptor, type 1A	Bmpr1a	0.35	0.02347	0.00005
fibroblast growth factor receptor-like 1	Fgfr1l	0.34	0.03349	0.01815
interleukin 17 receptor D	Il17rd	0.35	0.01961	0.00052
mitogen activated protein kinase 1	Mapk1	0.39	0.04889	0.00710
splicing factor, arginine/serine-rich 15 /// similar to splicing factor, arginine/serine-rich 15	Sfrs15	0.47	0.02113	0.00047
protein phosphatase 1A, magnesium dependent, alpha isoform	Ppm1a	0.46	0.02525	0.00280
protein kinase C, delta	Prkcd	0.45	0.03465	0.02184
insulin-like growth factor 2 receptor	Igf2r	0.53	0.04738	0.00600
Apoptosis				
junction-mediating and regulatory protein	Jmy	0.21	0.00000	0.00036
sphingosine-1-phosphate phosphatase 1	Sgpp1	0.23	0.01538	0.00160
cathepsin B	Ctsb	0.28	0.00000	0.00024
death inducer-obliterators 1	Dido1	0.46	0.03440	0.00196
Cell cycle, DNA replication, DNA repair, cell division, chromosome segregation, cell growth, cell size				
origin recognition complex, subunit 4-like (S. cerevisiae)	Orc4l	0.17	0.01852	0.00383
purine rich element binding protein B	Purb	0.29	0.02669	0.00019
chromosome segregation 1-like (S. cerevisiae)	Cse1l	0.34	0.03254	0.00002
minichromosome maintenance deficient 5, cell division cycle 46 (S. cerevisiae)	Mcm5	0.34	0.01852	0.00004
kinesin family member 11	Kif11	0.34	0.01938	0.00170
replication factor C (activator 1) 2	Rfc3	0.40	0.04420	0.00919
kinesin family member 22	Kif22	0.42	0.02667	0.00033

protein phosphatase 2 (formerly 2A), catalytic subunit, beta isoform	Ppp2cb	0.43	0.03243	0.00658
centromere protein E	Cenpe	0.54	0.03801	0.00299
Rap1 interacting factor 1 homolog (yeast) /// similar to Telomere-associated protein RIF1 (Rap1-interacting factor 1 homolog) (mRif1)	mRif1	0.63	0.04515	0.00228

EXAMPLE 6

DETEERMINATION OF PROBABILITY OF LIVE BIRTH EVENT

[00187] In the US, 7.3 million couples suffer from clinical infertility, for which more than ~ 120,000 IVF treatment cycles are performed per year. IVF is the most effective treatment for many infertility couples, but it is largely empiric, and may not result in a live birth for some couples despite multiple attempts. IVF treatment has revolutionized how physicians can help subfertile patients, but success rates with the use of autologous oocytes appear to have reached a plateau in the past decade. At an average cost of \$12,000 per cycle, IVF treatment alone costs health care payers, most of whom are the couples themselves, more than \$1 billion per year. However, the decision-making process for IVF can be daunting because IVF counseling and management decisions are often made based on the woman's chronological age, with non-standardized ways to adjust for various estimates of ovarian reserve and embryo quality. Further, quality of care is thought to vary amongst IVF clinics, but head-to-head comparisons are not possible in the absence of scientifically and rigorously defined prognostic stratification of patients. For a treatment that is physically, emotionally, and financially demanding, patients may feel that they are rolling the dice when deciding on whether to start or repeat IVF treatment. Consequently, there may be incongruence in patient- or self-selection for IVF, such that some patients with truly poor prognosis may develop unrealistic expectations, while others with truly good prognosis may miss opportunities for effective treatment. Although numerous publications have reported variables, including chronological age of the female patient, that are significantly associated with IVF outcomes, their contribution relative to outcomes is not clear, and we are even less certain how to directly apply those findings to counsel patients.

[00188] Accordingly, in the example we applied MART to analyze dichotomous live birth outcomes of a larger, four-year data set on 3338 fresh, non-oocyte donor IVF cycles and their associated embryos, including subsequent transfer of cryopreserved

embryos from 2003 to 2006. Importantly, 57 variables pertaining to clinical diagnoses, treatment response, and embryo developmental parameters from 1,879 fresh, first-cycles were used to generate prediction models in an unbiased manner, with no pre-selection of variables based on previous literature. We identified four prognostic factors -- the percentage of blastocysts (Blast Rate), total number of embryos (Embryo Number), total amount of gonadotropins required (TG), and endometrial thickness (EndoTh) -- that determine live birth outcomes. The prognostic contribution of other variables, such as age and the number of 8-cell stage embryos on day 3, were optional once these four prognostic variables were known. Finally, the results show how our approach and findings can be applied to immediately improve patient counseling and management protocols, and how these rigorously defined prognostic criteria can fuel our concerted effort to improve quality assurance and decrease rates of multiple gestation.

[00189] Between January 1, 2003 to December 31, 2006, 5037 IVF treatments were performed in total, with annual volumes that were comparable across years. Of those, 3347 were fresh IVF cycles that used the patients' own oocytes and that fulfilled inclusion criteria. After applying exclusion criteria, 3338 cycles remained, of which 1879 were first-IVF cycles (C1), 778 were second-IVF cycles performed for patients who did not have live birth outcomes in C1 and who returned for their second treatment (C2); 312 were third-IVF cycles (C3); 369 cycles were fourth, fifth and sixth cycles, which were not analyzed (**FIG. 17** and **FIG. 18**).

[00190] Data from C1, C2, and C3 were subsequently analyzed separately. Overall, 41% % of embryos had 8 blastomeres, the developmentally appropriate cell number, while 17% % of embryos had ≤ 4 cells, and 33% % had 5-7 cells, and 9% % had more than 8 cells (**FIG. 19**). This profile of *in vitro* human embryo development in IVF was consistent with previous reports. Further, the number of blastomeres formed by 10,687, 3,932 and 1,573 embryos by Day 3 of *in vitro* culture in C1, C2, and C3, were not significantly different, which suggested that embryo development was comparable amongst the first three cycles ($p=0.6$; **FIG 19**).

[00191] We used C1 data only to generate our prediction models, because patients dropped out for a variety of reasons for which there were no controls (**FIG. 18**), and

we did not want significant prognostic factors to be masked by cycle number; we did not know whether the drop out rates were comparable amongst patient subsets that varied in their prognoses. Consistent with other reports, a significant number of patients who had no live births dropped out after each unsuccessful cycle, and did not return for subsequent treatments. Nonetheless, C2 and C3 data were used to test whether the prediction models held true for subsequent treatments.

Variables Associated with Live Birth Outcomes

[00192] We first focused on 1879 C1 cycles and their associated 10,687 embryos that satisfied all inclusion and exclusion criteria. We systematically examined the association of each of 57 variables with live birth outcomes. These variables were chosen based on the quality and completeness of data entry, rather than prior scientific or clinical knowledge. Pair-wise logistic regression confirmed significant association of many variables with live birth outcomes, including patient's age, male partner's age, Day 3 FSH, and the number of prior clinical pregnancy losses ($p < 0.05$) (Table 13). As expected, diminished ovarian reserve (DOR), a clinical diagnosis based on poor ovarian response in previous non-IVF infertility treatments or clomid challenge test, was highly negatively associated with live birth outcomes ($p < 0.0001$), while the diagnosis of polycystic ovarian syndrome, which tends to increase ovarian response, was positively associated with live births ($p < 0.0001$).

Table 13. Variables and their association with IVF live birth outcomes.

Table 13A. Continuous Variables					
Variables	Estimate ¹	S.E.	p-value	Mean ⁵	StDev
Pre-IVF					
Age of patient	-1.27E-01	1.24E-02	4.12E-24	40.99	4.56
Age of male partner	-6.31E-02	1.02E-02	2.28E-09	42.95	5.82
Body mass index	-4.55E-03	6.17E-03	5.55E-01	24.90	9.75
No. of previous pregnancies ²	-1.16E-01	4.29E-02	1.07E-02	1.04	1.33
No. of previous term deliveries ³	-3.05E-02	9.05E-02	7.49E-01	0.27	0.60
Spontaneous miscarriages ⁴	-2.01E-01	7.36E-02	1.04E-02	0.41	0.82
Serum d.3 FSH (IU/L)	-6.93E-02	1.60E-02	2.89E-05	8.05	4.92
Year	2.24E-02	4.73E-02	6.83E-01	4.32	1.12
Pre-OR					
Total amount of recombinant FSH (IU/L)	-4.46E-04	3.94E-05	1.15E-28	3,512.03	1,587.94
Total amount of human menopausal gonadotropins (IU/L)	-1.01E-03	1.18E-04	4.69E-17	1,257.11	544.41
Total amount of gonadotropin	-3.55E-04	3.14E-05	1.36E-28	4,769.14	1,990.39
Endometrial Thickness (mm)	2.22E-01	2.66E-02	2.94E-16	10.00	2.10
No. of sperm motile after wash (million/mL)	-3.92E-04	2.92E-04	2.53E-01	128.41	244.60
No. of sperm motile before wash (million/mL)	-2.71E-04	3.39E-04	5.22E-01	106.46	198.93
Post-IVF					
Total no. of oocytes	1.06E-01	8.66E-03	2.53E-33	10.24	6.63
Percentage of normal and mature oocytes	3.36E+00	5.86E-01	2.69E-08	0.80	0.38
Normal fertilization (%)	1.32E+00	2.40E-01	9.74E-08	0.65	0.24
Unfertilized eggs (%)	-1.08E+00	2.48E-01	2.79E-05	0.27	0.23
Abnormally fertilized eggs (%)	-9.87E-01	4.34E-01	3.56E-02	0.08	0.14
Total number of embryos	1.78E-01	1.21E-02	2.43E-47	5.92	4.94
Blastocyst development (%)	3.05E+00	2.41E-01	3.03E-35	0.15	0.23
Compaction on day 3	2.66E-01	2.76E-01	4.51E-01	0.09	0.19
Average no. of cells per embryos	2.90E-01	4.57E-02	7.05E-10	6.76	1.32
No. embryos arrested at? 4 cells	-1.32E-02	2.74E-03	3.00E-06	16.69	23.44
No. of 8-cell embryos	2.59E-01	2.25E-02	1.71E-29	2.73	2.74
Percentage of 8 cell embryos (%)	9.40E-03	1.89E-03	1.50E-06	40.49	29.09
No. of embryos cryopreserved	3.23E+00	2.88E-01	2.33E-28	0.11	0.19
Average grade of embryos	-1.49E-01	1.11E-01	2.53E-01	1.83	0.61
Total no. of transferred embryos	9.59E-02	3.51E-02	1.04E-02	2.28	1.49
Average no. of cells per transferred embryos	5.37E-01	6.02E-02	2.07E-18	7.41	1.24
Average grade of transferred embryos	-6.33E-01	1.24E-01	8.24E-07	1.62	0.62
No. of transferred embryos arrested with ?4 cells	-2.36E-02	3.29E-03	7.12E-13	8.56	24.50
No. of 8 cell embryos transferred	3.26E-01	5.43E-02	5.59E-09	1.47	1.07
Percentage of 8 cell embryos transferred	1.67E-02	1.62E-03	3.65E-24	59.61	39.12

Table 13B. Categorical Variables					
Variables	Estimate	S.E.	p-value	Mean ⁶	StDev
Pre-IVF					
<i>Diminished ovarian reserve</i>	-1.07E+00	1.23E-01	1.40E-17	0.39	0.49
<i>Polycystic ovarian syndrome</i>	7.58E-01	1.66E-01	1.01E-05	0.09	0.29
<i>Unexplained female infertility</i>	5.74E-01	1.70E-01	1.37E-03	0.09	0.28
<i>Other causes for infertility</i>	-2.52E-01	1.14E-01	4.19E-02	0.34	0.47
<i>Tubal disease</i>	-9.40E-02	1.66E-01	6.24E-01	0.12	0.32
<i>Uterine Fibroids</i>	-1.87E-01	1.95E-01	4.51E-01	0.09	0.28
<i>Endometriosis</i>	-9.18E-02	1.53E-01	6.23E-01	0.14	0.35
<i>Male infertility</i>	4.51E-02	1.07E-01	7.09E-01	0.42	0.49
<i>Tubal Ligation</i>	4.09E-02	3.72E-01	9.12E-01	0.02	0.14
<i>Hydrosalpinx</i>	-7.22E-02	2.97E-01	8.08E-01	NA	NA
<i>Season: summer</i>	9.88E-02	1.54E-01	6.15E-01	0.22	0.42
<i>Season: winter</i>	1.32E-01	1.42E-01	4.63E-01	0.30	0.46
<i>Season: fall</i>	3.27E-01	1.54E-01	5.05E-02	0.20	0.40
Pre-OR					
<i>Oral contraception</i>	7.33E-01	1.85E-01	1.39E-04	0.87	0.34
<i>Sperm collected surgically</i>	1.57E-01	2.57E-01	6.23E-01	0.04	0.20
<i>Sperm collection from frozen</i>	2.61E-01	3.21E-01	5.22E-01	0.03	0.16
<i>Sperm from donor</i>	-4.03E-01	4.56E-01	4.84E-01	0.02	0.13
Post-IVF					
<i>Assisted Hatching</i>	-8.58E-01	1.26E-01	4.03E-11	0.33	0.47
<i>Day 5 Embryo transfer</i>	1.42E+00	1.20E-01	6.89E-31	0.27	0.44
<i>Antagonist protocol</i>	-1.35E+00	1.21E-01	6.30E-28	0.47	0.50
<i>Flare protocol</i>	-1.70E+00	1.84E-01	8.98E-20	0.18	0.39
<i>Echoip catheter</i>	6.83E-01	1.47E-01	7.64E-06	0.14	0.35
<i>Other catheter</i>	-2.69E-01	4.67E-01	6.24E-01	0.02	0.13
<i>Performance of ICSI</i>	3.34E-01	1.07E-01	3.07E-03	0.41	0.49

FOOTNOTE:

Each variable was tested for its association with live birth outcomes in IVF by logistic regression, which gave estimate, standard error of the estimate (S.E.), p-value, mean and standard deviation (S.D.). Results for continuous variables and categorical variables were listed in Table 1A and Table 1B, respectively.

¹Positive and negative estimates indicate association with positive and negative pregnancy outcomes, respectively.

²Number of previous clinical pregnancies as defined by positive serum human chorionic gonadotropin (hCG) or pregnancy test.

³Number of previous deliveries carried to term, 37wks.

⁴Miscarriages refer to developmental arrest or clinical pregnancy loss at or after 5wks gestation. Season, seemingly irrelevant, was included as a negative control.

[00193] We noted that many pairs of variables, such as age and total amount of gonadotropins used, were highly correlated with one another (Table 14). Although these findings are consistent with the presumed complex and poorly understood biological mechanisms amongst ovarian aging, ovarian hormone production and egg quality, those interactions could not be studied well by ANOVA using Chi-

square statistics tests (data not shown), presumably because they do not interact in a linear fashion, or multiple conditions may affect the nature of the interactions.

[00194] We chose to analyze data and generate prediction models by constructing boosted classification trees by MART[®] to identify non-redundant prognostic variables. MART[®] is a robust method used to identify interactive structure of variables that are predictive of outcomes. The use of cross-validation and boosting in parameter selection and model assessment in MART[®] also preserve parsimony and prevent over-fitting. Finally, MART does not presume the absence or presence of interactions, or the nature of the interactions.

Table 14A and 14B. Correlation Between Every Pair of Variables.

Table 14A

	Age of patient	Age of spouse	Body Mass Index	No. of previous pregnancies	No. of previous term deliveries	Spontaneous miscarriages	Serum d.2 FSH	Year	Total amount of FSH (IU/ml)	Total amount of HMG (IU/L)	Total amount of gonadotropin (IU/L)	Endometrial thickness (mm)	Total no. of sperm motile after wash (mill/mL)	Total no. of sperm motile before wash (mill/mL)	Total no. of oocytes	Percentage of normal and mature oocytes	Normal fertilization (%)
Pre-IVF Variables																	
Age of patient	1.00	0.56	0.04	0.22	0.13	0.14	0.17	-0.27	0.43	0.33	0.44	-0.16	0.04	0.05	-0.37	-0.20	-0.05
Age of spouse	0.56	1.00	0.01	0.10	0.09	0.06	0.08	-0.19	0.23	0.19	0.24	-0.12	-0.06	-0.05	-0.20	-0.08	-0.05
Body Mass Index	0.04	0.01	1.00	0.08	0.05	0.05	-0.04	0.07	0.00	0.00	0.00	-0.05	-0.01	0.00	0.06	-0.05	-0.04
No. of previous pregnancies	0.22	0.10	0.08	1.00	0.58	0.70	0.01	-0.01	0.08	0.06	0.08	-0.07	0.07	0.05	-0.04	-0.16	0.00
No. of previous term deliveries	0.13	0.09	0.05	0.58	1.00	0.14	0.05	-0.01	0.02	0.00	0.02	0.05	0.01	0.02	0.00	-0.09	0.01
Spontaneous miscarriages	0.14	0.06	0.05	0.70	0.14	1.00	-0.02	0.01	0.06	0.05	0.06	-0.07	0.05	0.03	-0.02	-0.13	0.00
Serum d.2 FSH	0.17	0.08	-0.04	0.01	0.05	-0.02	1.00	0.03	0.35	0.28	0.36	-0.11	0.00	0.00	-0.30	-0.13	-0.06
Year	-0.27	-0.19	0.07	-0.01	-0.01	0.01	0.03	1.00	-0.05	-0.06	-0.06	-0.04	-0.11	-0.08	0.11	-0.01	0.04
Pre-OR Variables																	
Total amount of FSH (IU/ml)	0.43	0.23	0.00	0.08	0.02	0.06	0.35	-0.05	1.00	0.66	0.98	-0.16	0.01	0.03	-0.55	-0.06	-0.05
Total amount of HMG (IU/L)	0.33	0.19	0.00	0.06	0.00	0.05	0.28	-0.06	0.66	1.00	0.80	-0.11	0.01	0.03	-0.44	-0.09	-0.06
Total amount of gonadotropin (IU/L)	0.44	0.24	0.00	0.08	0.02	0.06	0.36	-0.06	0.98	0.80	1.00	-0.16	0.02	0.03	-0.56	-0.07	-0.05
Endometrial thickness (mm)	-0.16	-0.12	-0.05	-0.07	0.05	-0.07	-0.11	-0.04	-0.16	-0.11	-0.16	1.00	0.01	0.01	0.14	0.37	0.01
Total no. of sperm motile after wash (mill/mL)	0.04	-0.06	-0.01	0.07	0.01	0.05	0.00	-0.11	0.01	0.01	0.02	0.01	1.00	0.62	-0.03	-0.19	0.01
Total no. of sperm motile before wash (mill/mL)	0.05	-0.05	0.00	0.05	0.02	0.03	0.00	-0.08	0.03	0.03	0.03	0.01	0.62	1.00	0.01	-0.08	-0.01
Post-IVF Variables																	
Total no. of oocytes	-0.37	-0.20	0.06	-0.04	0.00	-0.02	-0.30	0.11	-0.55	-0.44	-0.56	0.14	-0.03	0.01	1.00	0.13	0.03
Percentage of normal and mature oocytes	-0.20	-0.08	-0.05	-0.16	-0.09	-0.13	-0.13	-0.01	-0.06	-0.09	-0.07	0.37	-0.19	-0.08	0.13	1.00	-0.04
Normal fertilization (%)	-0.05	-0.05	-0.04	0.00	0.01	0.00	-0.06	0.04	-0.05	-0.06	-0.05	0.01	0.01	-0.01	0.03	-0.04	1.00
Unfertilized eggs (%)	0.01	0.03	0.03	-0.02	-0.01	-0.02	0.06	-0.08	0.04	0.03	0.04	0.01	-0.04	-0.04	-0.01	0.05	-0.83
Abnormally fertilized eggs (%)	0.06	0.03	0.01	0.04	0.00	-0.02	-0.01	0.07	0.02	0.05	0.03	-0.02	0.05	0.09	-0.03	-0.01	-0.35
Total no. of embryos	-0.34	-0.20	0.02	-0.03	0.00	-0.01	-0.27	0.10	-0.45	-0.36	-0.46	0.21	-0.01	-0.02	0.87	0.48	0.39
Blastocyst development (%)	-0.35	-0.21	0.06	0.01	0.03	0.02	-0.20	0.19	-0.41	-0.32	-0.42	0.12	0.00	-0.01	0.47	0.03	0.15
Compaction on d.3 (%)	0.09	0.05	-0.04	0.04	0.01	0.02	-0.02	-0.24	-0.05	-0.04	-0.05	-0.01	0.01	-0.01	-0.04	0.02	0.05
Average no. of cells per embryo	0.00	-0.01	0.04	0.07	0.02	0.05	-0.01	-0.04	-0.06	-0.07	-0.06	0.05	0.02	0.02	0.03	0.02	-0.01
No. of embryos arrested at =4 cells	-0.03	0.00	-0.01	-0.04	0.00	-0.04	0.00	0.08	0.02	0.04	0.03	-0.04	-0.03	0.00	0.00	0.01	-0.03
No. of 8 cell embryos	-0.23	-0.14	0.06	0.02	0.02	0.01	-0.22	0.02	-0.41	-0.32	-0.42	0.13	0.00	-0.01	0.65	0.03	0.26
Percentage of 8 cell embryos (%)	0.03	0.01	0.03	0.05	0.01	0.02	-0.06	-0.10	-0.08	-0.06	-0.08	0.06	0.01	0.04	0.06	0.01	0.05
Cryopreserved embryos (%)	-0.37	-0.24	0.03	-0.04	-0.02	-0.04	-0.19	0.11	-0.37	-0.31	-0.38	0.10	0.03	0.01	0.43	0.00	0.17
Average grade of embryos	-0.05	-0.03	-0.08	0.04	0.03	0.04	-0.01	0.16	0.04	0.02	0.04	0.01	-0.04	0.01	-0.02	0.01	0.01
Total no. of embryos transferred	0.13	0.08	-0.03	-0.02	-0.05	-0.03	-0.13	-0.07	0.02	0.04	0.03	0.11	0.02	0.00	0.08	0.59	0.23
Average no. of cells per transferred embryo	-0.16	-0.10	0.04	0.03	0.01	0.04	-0.12	-0.01	-0.21	-0.18	-0.22	0.10	0.01	0.00	0.28	0.02	0.11
Average grade of embryos transferred	0.11	0.06	-0.08	0.04	0.03	0.04	0.08	0.12	0.20	0.14	0.20	-0.05	-0.05	0.02	-0.23	-0.03	-0.06
No. of transferred embryos with arrest at = 4 cells	0.09	0.06	-0.01	-0.01	0.01	-0.03	0.08	0.04	0.15	0.13	0.15	-0.07	-0.02	0.02	-0.19	0.01	-0.12
No. of 8 cell embryos transferred	0.03	0.04	0.04	0.00	0.00	-0.02	-0.14	-0.10	-0.18	-0.11	-0.17	0.08	0.02	-0.01	0.29	-0.01	0.18
Percentage of 8 cell embryos transferred (%)	-0.19	-0.10	0.04	0.00	0.01	-0.01	-0.16	-0.02	-0.30	-0.21	-0.30	0.13	0.01	0.01	0.36	0.03	0.15

Table 14B

	Unfertilized eggs (%)	Abnormally fertilized eggs (%)	Total no. of embryos	Blastocyst development (%)	Compaction on d.3 (%)	Average no. of cells per embryo	No. of embryos arrested at =4 cells	No. of 8 cell embryos	Percentage of 8 cell embryos (%)	Cryopreserved embryos (%)	Average grade of embryos	Total no. of embryos transferred	Average no. of cells per transferred embryo	Average grade of embryos transferred	No. of transferred embryos with arrest at = 4 cells	No. of 8 cell embryos transferred	Percentage of 8 cell embryos transferred (%)
Pre-IVF Variables																	
Age of patient	0.01	0.06	-0.34	-0.35	0.09	0.00	-0.03	-0.23	0.03	-0.37	-0.05	0.13	-0.16	0.11	0.09	0.03	-0.19
Age of spouse	0.03	0.03	-0.20	-0.21	0.05	-0.01	0.00	-0.14	0.01	-0.24	-0.03	0.08	-0.10	0.06	0.06	0.04	-0.10
Body Mass Index	0.03	0.01	0.02	0.06	-0.04	0.04	-0.01	0.05	0.03	0.03	-0.08	-0.03	0.04	-0.08	-0.01	0.04	0.04
No. of previous pregnancies	-0.02	0.04	-0.03	0.01	0.04	0.07	-0.04	0.02	0.05	-0.04	0.04	-0.02	0.03	0.04	-0.01	0.00	0.00
No. of previous term deliveries	-0.01	0.00	0.00	0.03	0.01	0.02	0.00	0.02	0.01	-0.02	0.03	-0.05	0.01	0.03	0.01	0.00	0.01
Spontaneous miscarriages	-0.02	0.02	-0.01	0.02	0.02	0.05	-0.04	0.01	0.02	-0.04	0.04	-0.03	0.04	0.04	-0.03	-0.02	-0.01
Serum d.2 FSH	0.06	-0.01	-0.27	-0.20	-0.02	-0.01	0.00	-0.22	-0.06	-0.19	-0.01	-0.13	-0.12	0.08	0.08	-0.14	-0.16
Year	-0.08	0.07	0.10	0.19	-0.24	-0.04	0.08	0.02	-0.10	0.11	0.16	-0.07	-0.01	0.12	0.04	-0.10	-0.02
Pre-OR Variables																	
Total amount of FSH (IU/ml)	0.04	0.02	-0.45	-0.41	-0.05	-0.06	0.02	-0.41	-0.08	-0.37	0.04	0.02	-0.21	0.20	0.15	-0.18	-0.30
Total amount of HMq (IU/L)	0.03	0.05	-0.36	-0.32	-0.04	-0.07	0.04	-0.32	-0.06	-0.31	0.02	0.04	-0.18	0.14	0.13	-0.11	-0.21
Total amount of gonadotropin (IU/L)	0.04	0.03	-0.46	-0.42	-0.05	-0.06	0.03	-0.42	-0.08	-0.38	0.04	0.03	-0.22	0.20	0.15	-0.17	-0.30
Endometrial thickness (mm)	0.01	-0.02	0.21	0.12	-0.01	0.05	-0.04	0.13	0.06	0.10	0.01	0.11	0.10	-0.05	-0.07	0.08	0.13
Total no. of sperm motile after wash (mil/mL)	-0.04	0.05	-0.01	0.00	0.01	0.02	-0.03	0.00	0.01	0.03	-0.04	0.02	0.01	-0.05	-0.02	0.02	0.01
Total no. of sperm motile before wash (mil/mL)	-0.04	0.09	-0.02	-0.01	-0.01	0.02	0.00	-0.01	0.04	0.01	0.01	0.00	0.00	0.02	0.02	-0.01	0.01
Post-IVF Variables																	
Total no. of oocytes	-0.01	-0.03	0.87	0.47	-0.04	0.03	0.00	0.65	0.06	0.43	-0.02	0.08	0.28	-0.23	-0.19	0.29	0.36
Percentage of normal and mature oocytes	0.05	-0.01	0.48	0.03	0.02	0.02	0.01	0.03	0.01	0.00	0.01	0.59	0.02	-0.03	0.01	-0.01	0.03
Normal fertilization (%)	-0.83	-0.35	0.39	0.15	0.05	-0.01	-0.03	0.26	0.05	0.17	0.01	0.23	0.11	-0.06	-0.12	0.18	-0.15
Unfertilized eggs (%)	1.00	-0.24	-0.34	-0.17	-0.04	-0.02	0.05	-0.24	-0.05	-0.18	-0.01	-0.18	-0.12	0.06	0.12	-0.16	-0.14
Abnormally fertilized eggs (%)	-0.24	1.00	-0.11	0.03	-0.01	0.06	-0.03	-0.04	0.00	0.00	-0.01	-0.09	0.02	0.00	0.01	-0.04	-0.01
Total no. of embryos	-0.34	-0.11	1.00	0.52	-0.02	0.02	-0.01	0.77	0.07	0.49	0.00	0.24	0.31	-0.24	-0.22	0.33	0.40
Blastocyst development (%)	-0.17	0.03	0.52	1.00	0.00	0.25	-0.20	0.61	0.26	0.69	-0.18	-0.26	0.34	-0.35	-0.22	0.18	0.44
Compaction on d.3 (%)	-0.04	-0.01	-0.02	0.00	1.00	0.13	-0.09	0.08	0.19	0.03	-0.30	0.00	0.10	-0.27	-0.06	0.10	0.13
Average no. of cells per embryo	-0.02	0.06	0.02	0.25	0.13	1.00	-0.82	0.33	0.57	0.22	-0.38	-0.09	0.77	-0.37	-0.59	0.33	0.42
No. of embryos arrested at =4 cells	0.05	-0.03	-0.01	-0.20	-0.09	-0.82	1.00	-0.27	-0.47	-0.18	0.29	0.06	-0.62	0.27	0.78	-0.27	-0.34
No. of 8 cell embryos	-0.24	-0.04	0.77	0.61	0.08	0.33	-0.27	1.00	0.53	0.60	-0.21	-0.09	0.43	-0.42	-0.30	0.53	0.64
Percentage of 8 cell embryos (%)	-0.05	0.00	0.07	0.26	0.19	0.57	-0.47	0.53	1.00	0.23	-0.34	-0.11	0.46	-0.37	-0.36	0.60	0.77
Cryopreserved embryos (%)	-0.18	0.00	0.49	0.69	0.03	0.22	-0.18	0.60	0.23	1.00	-0.17	-0.32	0.32	-0.33	-0.20	0.17	0.42
Average grade of embryos	-0.01	-0.01	0.00	-0.18	-0.30	-0.38	0.29	-0.21	-0.34	-0.17	1.00	0.09	-0.26	0.86	0.16	-0.18	-0.23
Total no. of embryos transferred	-0.18	-0.09	0.24	-0.26	0.00	-0.09	0.06	-0.09	-0.11	-0.32	0.09	1.00	-0.14	0.16	0.06	0.39	-0.21
Average no. of cells per transferred embryo	-0.12	0.02	0.31	0.34	0.10	0.77	-0.62	0.43	0.46	0.32	-0.26	-0.14	1.00	-0.39	-0.76	0.42	0.55
Average grade of embryos transferred	0.06	0.00	-0.24	-0.35	-0.27	-0.37	0.27	-0.42	-0.37	-0.33	0.86	0.16	-0.39	1.00	0.27	-0.30	-0.43
No. of transferred embryos with arrest at = 4 cells	0.12	0.01	-0.22	-0.22	-0.06	-0.59	0.78	-0.30	-0.36	-0.20	0.16	0.06	-0.76	0.27	1.00	-0.35	-0.43
No. of 8 cell embryos transferred	-0.16	-0.04	0.33	0.18	0.10	0.33	-0.27	0.53	0.60	0.17	-0.18	0.39	0.42	-0.30	-0.35	1.00	0.73
Percentage of 8 cell embryos transferred (%)	-0.14	-0.01	0.40	0.44	0.13	0.42	-0.34	0.64	0.77	0.42	-0.23	-0.21	0.55	-0.43	-0.43	0.73	1.00

IVF Treatment Time Point-Specific Prognostic Factors and Models

[00195] Consistent with our goal to facilitate real life decision-making, we used boosted tree from MART to generate three prediction models, each of which utilized variables that would be available at a specific time point during IVF planning and treatment. As shown by the schematics in FIG. 20, the Pre-IVF model was limited to 21 variables primarily pertaining to patients' baseline characteristics, clinical diagnoses; the Pre-oocyte retrieval (Pre-OR) model utilized the same 21 variables pertaining to patient's response to ovarian stimulation in addition to 9 variables used in the Pre-IVF model, for a total of 30 variables; the Post-IVF model used all variables in Pre-IVF and Pre-OR in addition to data

pertaining to embryo development, and embryo parameters related to transfer or cryopreservation for a total of 57 variables.

[00196] **Pre-IVF Model.** MART analysis of 1879 C1 patients with respect to 21 continuous and categorical variables that are known prior to starting IVF treatment (see Table 13), showed that each of patient's age and the diagnosis of diminished ovarian reserve (DOR) alone, as well as their interaction, predicted live birth outcomes. At our center, DOR has been routinely used to describe patients who showed a history of poor response to ovarian stimulation by gonadotropins in controlled ovarian hyperstimulation/intrauterine insemination (COH/TUI) treatment. While the presence of DOR is a negative predictor for LB, it only pertained to ~12% of the population. The rest of the patients were stratified by age. However, the model identified age thresholds (i.e. 40.5, 42.5, and 46.5) that were vastly different from thresholds (i.e. <35, 36-37, 38-40, 41-42, >42) that are arbitrarily and commonly used in the literature (ref). Overall, 5 patient subsets, hereafter populations, were discerned by this model with LB rates of 21%, 39%, 27%, 17%, and 4% for 222 (12% of all C1 patients), 690 (37%), 176 (9%), 581 (31%), and 199 (11%) patients, respectively (**FIG. 21A**).

[00197] **Pre-OR Model.** In this model, 30 variables known by the time of oocyte retrieval, from the same population comprising 1879 C1 patients, were analyzed by MART. Three independent prognostic factors were identified – total amount of gonadotropins required (TG), age, and endometrial thickness (EndoTh). Again, although each of these prognostic factors had been reported to be associated with LB outcomes, for the first time, we are objectively defining their thresholds and the nature of their potential interaction, which are critical for clinical application. Overall, this model identified 7 distinct populations with respect to their LB rates spanning 2%, 51%, 28%, 32%, 18%, 37%, and 8% for 26 (1.4% of all C1 patients), 387 (21%), 58 (3%), 200 (11%), 662 (35%), 141 (8%), and 405 (22%) patients in Populations (Pops) 1 to 7 (**FIG. 21B**). In addition to precisely parsed populations, the striking findings were the miniscule chance of LB outcomes for patients who had EndoTh < 7.05 mm (Pop 1), those who fared poorly according to all 3 prognostic factors (Pop 5); and Age > 44.5 (Pop 7). A mere combination of Pops 1,

5, and 7 showed that 1119 or ~60% of all C1 patients had a very low LB rate of 13.5%.

[00198] **Post-IVF Model.** The Post-IVF model was generated by MART based on all 57 variables listed in Table 1. Of note, this analysis was restricted to 1664 of the 1879 C1 patients because 215 patients did not complete their IVF treatment and had “cycle cancellation” primarily due to suboptimal ovarian response. Four significant prognostic factors – Blast Rate, Embryo Number, TG, and EndoTh sufficed to differentiate 6 populations with LB rates ranging from 7%, 39%, 17%, 54%, 36%, and 72% for 361 (19% of patients analyzed), 96 (5.5%), 643 (40%), 316 (19%), 198 (12%), and 95 (6%) patients in Populations 1 to 6 (**FIG. 21C**).

Model Validation in Returning Patients and an Independent Data Set.

[00199] We validated our models by testing whether live birth rates amongst populations are significantly different in related data sets comprised of patients whose C1 did not result in a live birth and who returned for repeat treatment in 778 2nd-cycles (C2) and 312 3rd-cycles (C3). To further validate our results, we also tested an independent data set comprised of C1 data from 343 non-redundant patients who had IVF treatment in 2007. (See **FIG. 22**) Interestingly, we noted that the composition of the overall IVF patient population altered in between cycles, and between data sets.

[00200] We tested differential live birth rates amongst populations by ANOVA Chi-square test, which revealed that differential LB outcomes amongst populations remained significantly different in C2 and C3 from the 2003-2006 data set as well as C1 from the independent data set from 2007, for each of the Pre-IVF, Pre-OR, and Post-IVF models with p-values at 1.17E-50, 8.69E-71, and 1.64E-96, respectively (**FIG. 23**).

[00201] Then we validated our models in a different way, by testing whether the live birth rates of each prognostically-defined population was stable and unaltered amongst C1, C2, C3 of 2003-2006 data (inter-cycle, hereafter), and between C1 of 2007 and C1 of 2003-2006 data sets. The inter-cycle comparison showed that all populations remained similar for the Pre-OR and Post-IVF models; in the Pre-IVF model, only Pop 2 and Pop 4 showed that their live birth rates altered amongst

cycles ($p=0.02$). (See Table 15 for p-values of inter-cycle comparisons.) Further, comparison of populations between the 2003-2006 and 2007 data sets indicated that the Pre-IVF and Post-IVF models, and most of the Pre-OR models have remained constant and highly reproducible ($p \geq 0.5$); Pops 1 and 6 of the Pre-OR model had differential live birth outcomes between the two data sets.

Table 15. Live birth rates were compared between age-based control model and each of Pre-IVF, Pre-OR and Post-IVF models across populations for each age group.

	Age groups				
	<35	35-37	38-40	41-42	≥ 43
<i>Pre-IVF</i>	0.6	0.1	0.01	NA	NA
<i>Pre-OR</i>	0.04	1.8E-04	3.2E-07	4.1E-05	2.1E-06
<i>Post-IVF</i>	3.2E-07	3.7E-05	9.5E-16	3.7E-05	1.6E-09

[00202] Collectively, our results showed that all three models were effective in stratifying patients with differential live birth rates, and this ability to stratify by prognosis is reproducible in subsequent cycles and in an independent data set, despite changes in the composition of the general IVF patient population. Further, we confirmed that the prognostic stratification and live birth prediction used in each model were reproducible and validated in an independent data set, as well as in repeat cycles of returning patients.

Models predicted live birth outcomes more specifically than age.

[00203] We tested whether these prediction models defined patient populations better and more specifically than an age-based control model (hereafter, control). (See Methods for age-based control model details.) We show the population-specific, observed live birth outcomes compared to live birth rates predicted by the control for each five age categories, <35, 35-37, 38-40, 41-42, and ≥ 43 (**FIG. 24A-C**). Here, age categories were defined by conventional standards used by the Society for Assisted Reproductive Technologies (SART) national registry and most of the literature (ref). These comparisons highlight the tendency for age to overestimate or underestimate live birth rates. For example, the control model

tends to overestimate live birth rates in Pop 1 of the Pre-IVF model, Pops 5 and 7 of the Pre-OR model, and Pops 1 and 3 of the Post-IVF model (**FIG 24A-C**). In the comparison with the Pre-OR model, control model also tends to underestimate live birth rates in Pops 2 and 3, and in age groups 41-42 and ≥ 43 of Pop 6, while comparison with the Post-IVF model, age tends to underestimate Pops 4 and 6.

[00204] We compared each of Pre-IVF, Pre-OR and Post-IVF models to the control with respect to the live birth rates in each of five age categories. Both the Pre-OR and Post-IVF models predicted live birth rates significantly better than the control model across populations in all five age groups (p-values range from 0.04 to 0.5E-16, see **Table 15** for all p-values of these comparisons). In the Pre-IVF model, in which age is a key prognostic factor, the Pre-IVF model predicted outcomes better than the control for the age group 38-40, presumably because it also considers the diagnosis of diminished ovarian reserve ($p=0.01$). In summary, amongst patients who are undergoing their first fresh IVF cycle 17.7% of patients would receive more accurate and personalized live birth outcomes prediction from using the Pre-IVF model and 76% of patients would benefit from using the Pre-OR model. The utility of the Post-IVF model is to facilitate decision-making regarding subsequent IVF treatment in the event that live birth does not result from the first treatment, its use as a prediction tool for subsequent cycles is analyzed more in-depth later.

Prognostic Stratification Predicts Subsequent and Cumulative Live Birth Rates

[00205] We asked whether population assignment in C1 predicts live birth rates in C2 and the cumulative live birth rates in C2 and C3, for patients who do not have a live birth from C1. These questions precisely address the frequent scenario in which a couple needs to decide whether to repeat IVF treatment after an unsuccessful cycle. We addressed these questions by assigning patients to populations based on the Post-IVF model, followed by tracking patients' outcomes in C2 and C3. We considered both observed and hypothetical cumulative live birth rates. The observed cumulative rates were conservatively calculated based on the actual patients who returned for C2 and C3; the nominator was number of live births in C2 and C3, and the denominator was the total number of C2 cycles. The hypothetical

cumulative rates were calculated based on the probability of having live birth outcomes in C2 or C3, and assumed that the patients who dropped out were no different from the ones who returned; therefore, they are not affected by the limited number of patients who chose to have C3. Specifically, C1 assignment to Pops 1 and 3 resulted in observed cumulative live birth rates of 15.1% and 18.4%, respectively, and 27.5% for Pop 5. In contrast, Pops 2, 4 and 6 had much higher cumulative live birth rates of 56.7%, 42.9%, and 75%, respectively; the hypothetical cumulative live birth rates had a trend towards being slightly higher (FIG. 25). Therefore, population assignment based on prognostic factors in C1 re-defines the IVF population into subsets that have distinct and predictable cumulative live birth rates in C2 and C3.

Gradient Boost Analysis to Identify Relative Importance of Preselected Patient Variables

[00206] We used gradient boosted machine (GBM) to perform analyses to determine the "relative influence" of each variable on live birth outcomes for all three models. Briefly, we performed 10-fold cross validation for 1879, 1879, and 1664 fresh cycles for Pre-IVF, Pre-OR, and Post-IVF models, respectively. The optimal numbers of trees were 4677, 8680, and 10,745 for Pre-IVF, Pre-OR, and Post-IVF models, respectively; 25,000 trees (i.e. more than the minimum required) were analyzed for each model. The results are shown in Tables 16A, 16B, and 16C. The variables are ranked according to decreasing relative influence/importance such that the most important variables are ranked at the top. The numbers for relative influence add up to 100 in each model.

[00207] These GBM outputs contain thousands of trees which cannot be visualized and are conceptually too abstract for most scientific and clinical discussions. Therefore, we chose the top variables for further analysis to construct simpler tree models that could be visualized with Rpart. The advantages of simpler tree models were that they were conceptually more easily understood by scientists, clinicians, and lay people; distinct sub-populations can be characterized and subjected to further analyses to address specific questions, and to explore the utility of the models.

Tables 16A, 16B, and 16C. List of variables and their relative importance in determining live birth outcomes in the Pre-IVF, Pre-OR and Post-IVF model respectively.

Table 16A: Pre-IVF Model

Pre-IVF model	
Variables	Relative Importance
<i>Age of patient</i>	45.38
<i>Diminished ovarian reserve</i>	14.82
<i>Age of spouse</i>	11.55
<i>Serum d. 3 FSH (IU/L)</i>	10.71
<i>Body Mass Index</i>	6.37
<i>Polycystic ovarian syndrome</i>	2.88
<i>Season</i>	1.75
<i>Unexplained female infertility</i>	1.35
<i>Spontaneous miscarriages</i>	0.97
<i>Year</i>	0.89
<i>Other causes of female infertility</i>	0.76
<i>No. of previous pregnancies</i>	0.74
<i>No. of previous term deliveries</i>	0.64
<i>Endometriosis</i>	0.51
<i>Tubal disease</i>	0.48
<i>Tubal ligation</i>	0.09
<i>Male infertility only</i>	0.07
<i>Uterine fibroids</i>	0.02
<i>Hydrosalpinx</i>	0.01
<i>Male infertility causes</i>	0.01

Table 16B: Pre-OR Model

Pre-OR model	
Variables	Relative Importance
<i>Total amount of gonadotropin</i>	35.22
<i>Endometrial Thickness</i>	17.58
<i>Age of patient</i>	14.16
<i>No. of sperm motile after wash (million/mL)</i>	8.05
<i>No. of sperm motile before wash (million/mL)</i>	6.87
<i>Sperm collection</i>	6.03
<i>Age of spouse</i>	4.27
<i>Body Mass Index</i>	2.91
<i>Serum d. 3 FSH (IU/L)</i>	2.41
<i>Season</i>	0.75
<i>Spontaneous miscarriages</i>	0.39
<i>Unexplained female infertility</i>	0.39
<i>No. of previous term deliveries</i>	0.26
<i>Year</i>	0.26
<i>No. of previous pregnancies</i>	0.21
<i>Other causes of female infertility</i>	0.10
<i>Endometriosis</i>	0.04
<i>Male infertility only</i>	0.02
<i>Tubal ligation</i>	0.02
<i>Polycystic ovarian syndrome</i>	0.02
<i>Tubal disease</i>	0.02
<i>Sperm from donor</i>	0.00
<i>Hydrosalpinx</i>	0.00
<i>Uterine fibroids</i>	0.00
<i>Male infertility causes</i>	0.00

Table 16C: Post-IVF Model

Post-IVF model	
Variables	Relative Importance
<i>Blastocyst development (%)</i>	27.30
<i>Total amount of gonadotropin</i>	10.58
<i>Total no. of embryos</i>	8.95
<i>Endometrial Thickness</i>	7.24
<i>Flare protocol</i>	6.13
<i>Average no. of cells per embryo</i>	4.97
<i>Catheter used</i>	4.04
<i>Percentage of 8 cell embryos</i>	3.54
<i>Serum d. 3 FSH (IU/L)</i>	3.17
<i>Body Mass Index</i>	3.16
<i>No. of sperm motile before wash (million/mL)</i>	2.99
<i>No. of sperm motile after wash (million/mL)</i>	2.88
<i>Age of patient</i>	2.56
<i>Average grade of embryos</i>	2.49
<i>Day of embryo transfer</i>	1.88
<i>Season</i>	1.32
<i>Spontaneous miscarriages</i>	0.93
<i>No. of previous term deliveries</i>	0.74
<i>Oral contraceptive pills</i>	0.68
<i>Sperm collection</i>	0.58
<i>Unfertilized eggs (%)</i>	0.56
<i>No. embryos arrested at= 4 cells</i>	0.54
<i>Compaction on day 3</i>	0.51
<i>Normal fertilization (%)</i>	0.48
<i>Abnormally fertilized eggs (%)</i>	0.48
<i>Percentage of normal and mature oocytes</i>	0.48
<i>No. of previous pregnancies</i>	0.26
<i>Year</i>	0.17
<i>Polycystic ovarian syndrome</i>	0.12
<i>Unexplained female infertility</i>	0.11
<i>Tubal disease</i>	0.05
<i>Male infertility only</i>	0.05
<i>Male infertility causes</i>	0.03
<i>Endometriosis</i>	0.02
<i>Other causes of female infertility</i>	0.01
<i>Uterine fibroids</i>	0.01
<i>Tubal ligation</i>	0.00
<i>Sperm from donor</i>	0.00
<i>Hydrosalpinx</i>	0.00
<i>Performance of ICSI</i>	0.00
<i>Assisted Hatching</i>	0.00

EXAMPLE 7**DETERMINATION OF MOLECULAR FINGERPRINT FOR NORMAL VS.
ABNORMAL HUMAN EMBRYOS**

[00208] We successfully tested the expression of a panel of genes in dissociated blastomeres of control cryopreserved-thawed human embryos that are at the 1-cell, 4-cell, and 8-cell stages. (Patients donated those control embryos to research because they had completed their families.) We analyzed the data by considering both standard deviation and mean. The results of gene chip experiments on the 1-cell, 4-cell, and 8-cell stages are provided in FIGS 25 – 30. These results show the utility of gene expression analysis at the single-cell level as well examining expression of many genes simultaneously. As such, the results show that the fingerprinting analysis for the embryos could be used to provide information that may improve prediction of live birth outcomes for subsequent IVF cycles.

[00209] The preceding merely illustrates the principles of the invention. It will be appreciated that those skilled in the art will be able to devise various arrangements which, although not explicitly described or shown herein, embody the principles of the invention and are included within its spirit and scope. Furthermore, all examples and conditional language recited herein are principally intended to aid the reader in understanding the principles of the invention and the concepts contributed by the inventors to furthering the art, and are to be construed as being without limitation to such specifically recited examples and conditions. Moreover, all statements herein reciting principles, aspects, and embodiments of the invention as well as specific examples thereof, are intended to encompass both structural and functional equivalents thereof. Additionally, it is intended that such equivalents include both currently known equivalents and equivalents developed in the future, i.e., any elements developed that perform the same function, regardless of structure. The scope of the present invention, therefore, is not intended to be limited to the exemplary embodiments shown and described herein. Rather, the scope and spirit of present invention is embodied by the appended claims.

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
12561	0610039J04Rik: RIKEN cDNA 0610039J04 gene	726.94	952.88	925.99	702.03	806.57	331	405.8	335.4	516.81	1.682481163	0.04796748	0.09424119	11.4121
17702	0610040J01Rik: RIKEN cDNA 0610040J01 gene	408.14	292.64	232.51	164.99	152.49	79.1	50.46	95.62	57.89	3.108117559	0.027310924	0.06003501	19.3713
14297	1110007M04Rik: RIKEN cDNA 1110007M04 gene	1261.71	969.97	1148.54	891.24	883.27	742.9	514.8	528.3	552.68	1.643608215	0.047535211	0.09174589	11.9566
23774	1300003B13Rik: RIKEN cDNA 1300003B13 gene	366.9	654.15	629.58	353.58	521.61	292.6	233.5	199.1	273.86	1.761395346	0.047778709	0.09288628	11.6452
2769	1500005A01Rik: RIKEN cDNA 1500005A01 gene	190.67	162.34	88.43	60.38	43.44	74.49	60.43	32.43	37.65	2.858882197	0.048085485	0.09187593	12.0161
22816	1500011B03Rik: RIKEN cDNA 1500011B03 gene	322.26	247.98	673.66	122.44	129.61	103.8	91.83	188.2	110.61	3.332395687	0.028446389	0.06021882	19.6218
22897	1500041J02Rik: RIKEN cDNA 1500041J02 gene	1893.92	1849.2	1693.29	1451	1424	1116	1066	1017	1101.9	1.515255995	0.046834345	0.09180399	11.8767
39273	1700013H16Rik: RIKEN cDNA 1700013H16 gene	758.43	675.7	737.99	758.17	463.94	313.6	215	215.7	339.42	1.884057091	0.045634921	0.08490741	12.9579
39378	1700080O16Rik /// LOC665895 /// LOC671223 /// LOC676479: RIKEN cDNA 1700080O16 gene /// similar to melanoma antigen family A, 5 /// similar to melanoma antigen family A, 5 /// similar to melanoma antigen family A, 5	200.58	99.12	179.18	2.68	13.11	53.79	4.95	4.98	14.24	10.21610667	0.010416667	0.0375	38.9031
25993	1700129I15Rik: RIKEN cDNA 1700129I15 gene	669.05	306.9	110.37	38.64	337.75	62.42	8.11	130.2	9.06	3.706501527	0.032414911	0.06916261	16.8302
1469	1810021J13Rik: RIKEN cDNA 1810021J13 gene	197.73	84.01	193.51	54.69	91.43	27.17	26.09	36.55	61.85	3.191953791	0.03257329	0.06910423	16.8574
38853	1810044A24Rik: RIKEN cDNA 1810044A24 gene	327.76	445.4	180.89	194.87	99.06	230.3	90.63	104.2	122.74	2.266582724	0.044401544	0.08662162	12.7189
14361	2010109N14Rik: RIKEN cDNA 2010109N14 gene	2153.09	1615.8	2175.97	1431.2	1545.7	1498	990.7	1212	998.06	1.548963257	0.036834925	0.07317417	15.5482
12883	2010301N04Rik: RIKEN cDNA 2010301N04 gene	137.6	388.98	247.94	70.02	143.76	72.7	44.83	53.42	47.03	3.587733926	0.023560209	0.0575829	21.1602
4342	2410022L05Rik: RIKEN cDNA 2410022L05 gene	1336.22	1283.4	1702.28	919.13	704.51	611.2	198	869.9	354.78	2.363351515	0.019047619	0.04355556	28.4416

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
4343	2410022L05Rik: RIKEN cDNA 2410022L05 gene	1587.18	1243	1840.86	1006.1	799.09	735.3	264.5	1067	383.1	2.195358325	0.020833333	0.04227431	29.5735
7831	2510042P03Rik: RIKEN cDNA 2510042P03 gene	609.91	476.61	483.28	205.72	292.53	253.9	143.3	227.1	130.33	2.506046408	0.026923077	0.06301282	18.4494
38903	2600010E01Rik: RIKEN cDNA 2600010E01 gene	303.78	312.45	368.93	57.76	100.13	37.76	33.32	102.2	59.89	5.037893122	0.01010101	0.03750842	38.2228
12473	2610001J05Rik: RIKEN cDNA 2610001J05 gene	296.77	409.38	503.49	245.92	295.87	217.2	127.3	127.2	112.09	2.149210241	0.042988741	0.08313886	13.1993
17629	2610200G18Rik: RIKEN cDNA 2610200G18 gene	507.41	353.97	562.56	226.7	318.05	160	138.3	150	181.72	2.424181549	0.02284264	0.05854484	20.816
8796	2610204L23Rik: RIKEN cDNA 2610204L23 gene	2035.34	2238.2	2213.21	1158	1919.6	996.2	1310	1294	1439.4	1.598253609	0.035814607	0.07222846	15.755
13166	2610206B13Rik: RIKEN cDNA 2610206B13 gene	225.38	66.31	200.47	99.87	45.82	54.05	28.4	45.37	49.13	3.050830647	0.03313253	0.07128514	16.2151
15069	2700060E02Rik: RIKEN cDNA 2700060E02 gene	1283.21	1570.9	1406.02	866.04	1267.3	698.8	686.8	853.4	876.63	1.623218225	0.047315742	0.09020625	12.2015
12465	2810026P18Rik: RIKEN cDNA 2810026P18 gene	2572.92	1625.6	2626.66	1339.2	1910.7	1650	1370	1348	1217.5	1.544918909	0.02722323	0.06469449	17.9141
1357	2810037C14Rik: RIKEN cDNA 2810037C14 gene	1234.26	1441.4	1159.06	686.13	824.88	637.2	505.8	807.8	513.31	1.929321084	0.027184466	0.06282848	18.5328
7425	2810405K02Rik: RIKEN cDNA 2810405K02 gene	178.7	133.85	23.59	37.21	19.15	47.42	21.42	5.11	43.31	3.872134547	0.047822374	0.09246513	11.7588
2836	2810429O05Rik: RIKEN cDNA 2810429O05 gene	545.26	616.74	406.07	256.07	314.66	153.4	93.44	189.3	192.29	2.615411559	0.024390244	0.05095238	24.3184
8173	2900092E17Rik: RIKEN cDNA 2900092E17 gene	364.43	535.89	536.98	201.3	208.16	223.1	249.4	258.8	279.55	2.023952855	0.034950071	0.07232525	15.8264
31900	3 days neonate thymus cDNA, RIKEN full-length enriched library, clone:A630060L04 product:unclassifiable, full insert sequence	217.23	198.01	208.5	35.11	62.24	25.96	128.6	54.01	42.21	3.58368285	0.026156942	0.06102616	18.9222

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
23783	3110040M04Rik /// LOC668050 /// LOC671025 /// LOC671035 /// LOC671043 /// LOC671049 /// LOC676755 /// LOC676758: RIKEN cDNA 3110040M04 gene /// similar to U2-associated SR140 protein /// hypothetical protein LOC668055 /// similar to U2-associated SR140 protein /// similar to U2-associated SR140 protein /// similar to U2-associated SR140 protein /// similar to U2-associated SR140 protein /// similar to U2-associated SR140 protein	262.9	66.64	204.12	24.31	69.95	13.99	53.82	27.08	11.49	5.319577352	0.019607843	0.04245098	28.9087
24812	3110045A19Rik: RIKEN cDNA 3110045A19 gene	133.79	102.29	11.36	16.34	11.54	14.64	9.74	11.08	9.95	6.752353664	0.035664336	0.07221445	15.7371
41988	4930406H16Rik /// Nalp4e: RIKEN cDNA 4930406H16 gene /// NACHT, leucine rich repeat and PYD containing 4E	504.28	1414.9	933.88	507.4	694.22	548.5	512.1	494.6	292.39	1.871318333	0.045098039	0.08532353	12.8793
25457	4930452B06Rik: RIKEN cDNA 4930452B06 gene	159.43	56.45	170.72	54.77	36.42	26.26	16.28	39.2	26.39	3.879189243	0.029668412	0.06595695	17.569
27714	4930455F23Rik: RIKEN cDNA 4930455F23 gene	534.1	677.74	522.3	346.74	399.19	274.9	281.7	256.8	176.61	1.997868651	0.047457627	0.09223446	11.7332
12139	4933407C03Rik: RIKEN cDNA 4933407C03 gene	132.88	24.46	352.91	23.21	29.24	24.42	23.67	16.78	20.79	7.389037724	0.023746702	0.05663149	21.4433
5148	4933411K20Rik: RIKEN cDNA 4933411K20 gene	738.65	1039.5	815.98	385.61	591.61	566	375.2	430.3	545.57	1.792608768	0.048440066	0.09399288	11.4661

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
39323	4933416M07Rik: RIKEN cDNA 4933416M07 gene	131.45	240.96	376.74	164.45	78.57	97.69	70.38	90	47.91	2.729143898	0.042071197	0.08097807	13.6443
29217	4933427D06Rik: RIKEN cDNA 4933427D06 gene	908.64	946.8	1089.77	626.62	734.75	426.3	393.3	601.6	587.3	1.747950978	0.047619048	0.09166961	11.9754
32880	5330427D05Rik: RIKEN cDNA 5330427D05 gene	288.66	223.45	219.68	127.67	89.86	117.7	46.92	105.9	66.56	2.639125809	0.047120419	0.09182373	11.9053
12662	5730507H05Rik: RIKEN cDNA 5730507H05 gene	812.73	648.02	809.21	424.8	297.08	456.7	492.6	353.6	405.37	1.868164517	0.035161744	0.07214721	15.7684
10438	5830415L20Rik: RIKEN cDNA 5830415L20 gene	1798.9	1735.1	1623.57	447.1	794.05	612.1	547.7	578.6	472.45	2.988146002	0	0.02944444	58.7902
10605	6430548M08Rik: RIKEN cDNA 6430548M08 gene	172.77	77.66	129.96	52.6	22.23	16.06	21.88	26.12	38.85	4.280297063	0.038167939	0.07466497	15.0198
26997	6720463M24Rik: RIKEN cDNA 6720463M24 gene	217.15	218.01	225.79	73.98	49.43	68.05	56.28	107.7	70.8	3.10166827	0.032786885	0.07150025	16.1248
31488	7420416P09Rik: RIKEN cDNA 7420416P09 gene	891.83	777.85	767.35	491.65	642.89	411.6	237.4	519.4	353.02	1.835126771	0.048171276	0.09178709	12.027
29615	A230048G03Rik /// LOC670614: RIKEN cDNA A230048G03 gene /// similar to CG2747-PB, isoform B	171.76	66.93	163.5	48.01	41.96	51.96	61.61	18.34	49.85	2.960217863	0.043032787	0.08315232	13.2034
22603	A230062G08Rik: RIKEN cDNA A230062G08 gene	322.83	261.74	354.1	323.26	201.66	68.32	42.84	53.83	127.44	2.29686181	0.047858942	0.09292751	11.655
41840	A430033K04Rik: RIKEN cDNA A430033K04 gene	1229.01	998.13	1215.48	461.08	557.03	342.2	761	837.6	205.38	2.175946907	0.016877637	0.04720113	26.7452
15795	A830039H10Rik: RIKEN cDNA A830039H10 gene	285.01	249.94	242.98	101.83	100.81	86.69	105.9	99.37	94.16	2.642649682	0.042253521	0.0812026	13.6501
14842	Adcy3: adenylate cyclase 3	758.96	864.83	804.58	572.29	686.9	438.5	319.3	409.5	337.76	1.757014688	0.048333333	0.09301667	11.6011
27502	Adult male medulla oblongata cDNA, RIKEN full-length enriched library, clone:6330441D15 product:hypothetical protein, full insert sequence	149.12	176.59	33.92	30.73	59.6	50.95	19.27	31.34	26.32	3.296182576	0.042900919	0.0831018	13.1896
42166	Agpr: Agouti related protein	231.44	178.62	94.37	33.38	179.24	13.68	1	3.54	25.21	3.940089826	0.033182504	0.07107089	16.2397
30358	AI449441: expressed sequence AI449441	151.46	251.91	122.96	52.19	97.84	34.47	27.9	63.13	28.11	3.466802793	0.044847328	0.08782443	12.5897

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
27056	Akap7: A kinase (PRKA) anchor protein 7	474.68	818.9	809.2	330.68	516.16	412.1	350.4	355.2	259.52	1.891006214	0.028520499	0.0653773	17.7396
40856	Arhgef15: Rho guanine nucleotide exchange factor (GEF) 15	75.81	156.72	162.47	102.39	39.12	15.89	18.53	57.97	26.4	3.034959662	0.04506232	0.08746245	12.6344
14676	Arl4c /// LOC632433: ADP-ribosylation factor-like 4C /// similar to ADP-ribosylation factor-like protein 7	311.79	210.2	41.38	104.69	73.13	31.74	50.31	49.49	54.66	3.095269491	0.042752868	0.08291971	13.3239
14653	Arpc5l: actin related protein 2/3 complex, subunit 5-like	2473.77	2293.9	2924.51	1813.4	2433.6	1537	1353	1754	1246.7	1.517524443	0.048559671	0.0937668	11.4908
40302	Arrdc3: arrestin domain containing 3	183.69	264.91	361.58	62.76	138.75	89.69	46.24	105.3	72.66	3.144010245	0.026209677	0.06090726	18.9442
18069	Arx: aristaless related homeobox gene (Drosophila)	297.32	299.7	123.78	72.45	57.19	29.97	24.05	38.12	24.48	5.853975473	0.016129032	0.03553763	46.1316
29021	Ascl1: achaete-scute complex homolog-like 1 (Drosophila)	78.29	281.43	176.69	53.61	123.08	40.97	87.55	31.01	24.18	2.976748058	0.046793761	0.09179376	11.8701
7414	Atp6v1g1: ATPase, H+ transporting, lysosomal V1 subunit G1	814.9	317.26	545.09	311.05	237.03	227	233.4	269.9	361.16	2.046137987	0.042708333	0.08292361	13.3161
27011	Atp8a1: ATPase, aminophospholipid transporter (APLT), class I, type 8A, member 1	113.19	302.99	336.88	111.03	98.91	55.24	45.16	134.9	41.74	3.092585368	0.024615385	0.05401026	22.9511
10707	Atpbd4: ATP binding domain 4	644.58	851.86	1031.97	378.68	582.62	175.6	246.6	575.3	407.51	2.137051719	0.026785714	0.05929315	19.8453
26830	AU022751: expressed sequence AU022751	1575.02	1805.1	1927.95	1006.1	1485.3	930.7	1148	1148	924.02	1.598423276	0.037974684	0.07783659	14.1867
28921	AU041783: expressed sequence AU041783	728.29	1031.6	715.32	268.18	470.17	621.1	289.7	297.9	348.87	2.1562451	0.03485064	0.07223803	15.8165
17440	Aurkc: aurora kinase C	1462.84	1182.8	1533.48	1312.5	1219.2	927.9	377.4	555.4	561.63	1.687182324	0.047372954	0.09207293	11.8231
15571	Auts2: autism susceptibility candidate 2	892.1	1352.7	1127.84	816.47	892.95	829.9	561.9	507.8	366.68	1.696647131	0.047493404	0.09168279	11.9562
8718	AW060207: expressed sequence AW060207	432.59	609.26	574.19	250.29	379.24	218.9	259.2	166.6	184.98	2.214784968	0.026946108	0.06165669	18.8051

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
29405	AW146299: expressed sequence AW146299	284.07	216.2	332.39	121.93	161.67	105.2	66.41	71.03	112.37	2.607562828	0.036290323	0.07364247	15.4019
29099	B230312I18Rik: RIKEN cDNA B230312I18 gene	420.5	374.15	459.33	169.13	94.51	275.9	97.89	101.3	61.44	3.134479828	0.016949153	0.04735876	26.7539
11389	B930041F14Rik: RIKEN cDNA B930041F14 gene	26.1	145.4	149.78	15.65	34.72	7.96	30.02	31.98	43.39	3.924749572	0.036096257	0.07358289	15.378
35357	B930096F20Rik: RIKEN cDNA B930096F20 gene	232.53	102.36	229.72	30.99	16.44	13.59	17.81	40.92	54.55	6.478600115	0.010309278	0.03721649	38.8416
33323	B930096F20Rik: RIKEN cDNA B930096F20 gene	162.51	79.13	176.86	11.2	5.38	6.12	14.66	19.82	13.65	11.81702668	0.018691589	0.03785047	36.9537
30001	BB176347: Expressed sequence BB176347	141.82	106.11	79.54	18.32	14.54	23.04	8.15	16.92	11.44	7.087328211	0.029616725	0.06619048	17.5516
27229	BC010981: cDNA sequence BC010981	388.39	149.2	438	121.27	178.69	160.1	120.7	85.02	190.36	2.279043147	0.03271028	0.07022845	16.5397
10371	BC021367: cDNA sequence BC021367	255.84	599.36	133.32	56.72	58.42	9.17	12.43	144.6	45.71	6.045808997	0.015151515	0.03744949	34.4898
10893	BC022623: cDNA sequence BC022623	784.51	724.29	716.83	301.14	357.98	339.1	388.4	240.7	107.29	2.566144551	0.021912351	0.04861886	26.0618
28308	BC047219: cDNA sequence BC047219	402.73	298.41	481.25	99.48	304.76	138.3	227.8	47.95	55.07	2.707649679	0.02690583	0.05922272	19.8718
31719	BC052883: cDNA sequence BC052883	3547.69	3957.6	3518.81	3150.9	2830	3065	843.8	1089	690.96	1.889427057	0.024390244	0.05562782	21.7701
14991	Bcas2: breast carcinoma amplified sequence 2	933.79	898.82	1023.13	656.47	683.52	725.1	238.6	384.1	299.55	1.911914063	0.037546934	0.07496454	14.8965
4790	Blcap: bladder cancer associated protein homolog (human)	1031.43	970.72	1005.09	398.98	455.59	428.7	593.8	433.6	467.82	2.1646422	0.025236593	0.0543428	23.1594
7998	Btf3: basic transcription factor 3	1578.04	1824.5	1702.45	1348.1	1356.3	1018	743.3	874.9	818.67	1.657623607	0.036809816	0.07552147	14.7389
27083	C030048B08Rik: RIKEN cDNA C030048B08 gene	408.49	460.05	549.24	205.03	223.41	242.7	206.1	243.9	160.9	2.211756263	0.036991369	0.07569667	14.7567
31101	C330046E03: hypothetical protein C330046E03	4819.33	3225.1	3559.24	755.8	537.22	920.6	495.6	490.6	1006.3	5.517476831	0	0.005	199.636
43660	C80008: expressed sequence C80008	387.88	488.55	927.38	95.78	305.61	74.02	92.37	277.2	122.15	3.730232751	0.018867925	0.03821803	32.1199
41465	C87977: expressed sequence C87977	1813.37	1754	1731.6	1381	1479.5	1226	879	1054	758.51	1.563607178	0.0472103	0.09198283	11.8126

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
22912	Calcoccl: calcium binding and coiled coil domain 1	551.7	236.85	422.22	259.49	280.12	113.5	190.8	154.3	163.4	2.084497585	0.044790652	0.08589094	12.8067
8101	Camk2g: calcium/calmodulin-dependent protein kinase II gamma	336.31	254.03	256.42	116.66	94.65	136.1	97.12	109.2	105.92	2.567339761	0.04613936	0.08864721	12.4665
11172	Car9: carbonic anhydrase 9	471.81	315.93	321.74	92.43	324.34	121.6	91.77	148.2	106.79	2.507101133	0.045	0.08457333	13.0189
270	Carhsp1: calcium regulated heat stable protein 1	783.29	492.05	805.9	323.58	378.15	367.8	428.5	461.3	410.2	1.756676458	0.038106236	0.077602	14.2229
15424	Ccdc102a /// LOC672218: coiled-coil domain containing 102A /// similar to CG31638-PA	88.88	90.52	37.42	3.07	8.66	4.42	4.14	2.85	2.42	16.96557121	0.027027027	0.06286358	18.4783
4096	Ccdc56: coiled-coil domain containing 56	392.74	395.1	319.83	208.46	301.88	148.4	177.1	110	108.37	2.101282392	0.044310171	0.08402148	13.0746
22624	Ccdc69: Coiled-coil domain containing 69	962.65	1366.1	1349.11	634.59	688.1	744	606.2	369.6	463.47	2.098090669	0.019230769	0.0436859	28.4695
22741	Ccdc69: coiled-coil domain containing 69	1132.93	1623.1	1851.99	805.4	1046.3	856	984.6	482	775.42	1.861933951	0.022535211	0.05503286	22.1289
22623	Ccdc69: coiled-coil domain containing 69	1050.67	1566.1	1589.84	762.24	1001.9	968.9	918.1	524.8	763.31	1.703329676	0.027079304	0.06286912	18.4874
16411	Ccni: cyclin I	493.47	644.26	702.9	315.9	404.45	474.9	206.1	318.3	253.34	1.865932038	0.042944785	0.08317314	13.192
16410	Ccni: cyclin I	995.58	940.06	798.15	555.77	798.27	475.2	498.2	513.8	309.27	1.735447687	0.045499505	0.08530828	12.9222
44690	Ccnjl: cyclin J-like	710.33	626.07	346.33	315.27	210.57	294	79.62	182.2	113.19	2.816638072	0.017647059	0.03870588	31.472
329	Cd24a: CD24a antigen	1205	1433.8	1104.89	719.21	978.74	647.6	776.5	731.8	580.28	1.688576461	0.047178538	0.08963306	12.3197
6122	Cdc25b: cell division cycle 25 homolog B (S. cerevisiae)	1318	1113.9	1472.91	1078.3	923.92	886.1	576.7	563.7	362.23	1.778541969	0.036144578	0.07362784	15.3818
15802	Cdc9111: CDC91 cell division cycle 91-like 1 (S. cerevisiae)	49.58	345.26	607.45	52.03	65.68	20.93	52.8	45.12	27.03	7.604916727	0	0.03	60.8874
12970	Cdca1: cell division cycle associated 1	674.36	667	543.88	296.75	413.81	246.4	141.6	321.8	221.52	2.296412693	0.022038567	0.05520661	21.9807
42049	Cdk9: cyclin-dependent kinase 9 (CDC2-related kinase)	139.64	199.95	196.21	14.31	19.14	29.4	35.55	8.3	51.82	6.76003028	0.014285714	0.03807143	33.6461
8112	Cdkal1: CDK5 regulatory subunit associated protein 1-like 1	879.67	767.38	813.54	253.82	513.83	385.5	494.5	208.3	137.41	2.468897841	0.022177419	0.04872312	26.1509

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
3196	Cebpb: CCAAT/enhancer binding protein (C/EBP), beta	132	304.11	204.27	35.5	57.66	60.4	8.94	15.35	29.75	6.169364162	0.018404908	0.03862986	31.7317
19532	Chchd5: coiled-coil-helix-coiled-coil-helix domain containing 5	220.46	43.46	183.56	95.26	39.57	75.13	40.76	8.84	10.24	3.317123795	0.042857143	0.0835	13.1659
21181	Chchd7: coiled-coil-helix-coiled-coil-helix domain containing 7	143.11	45.85	147.48	54.41	37.61	40.61	18.63	32.67	27.57	3.181465721	0.045320197	0.08514614	12.9126
14624	Chd4: chromodomain helicase DNA binding protein 4	618.5	949.18	1142.49	391	589.16	319.2	415.8	423.6	527.37	2.033097654	0.0243309	0.05852393	20.5004
15480	Chd4: chromodomain helicase DNA binding protein 4	1585.68	1818.1	1935.42	1226.8	1598.3	1198	946.6	909.1	980.36	1.556702038	0.044834308	0.0858642	12.8114
15075	Clu: clusterin	196.31	133.06	97.6	58.66	66.23	26.33	41.53	27.45	12.04	3.676972098	0.04567079	0.08815414	12.5518
2308	Cml1: camello-like 1	47.86	117.93	133.93	4.92	25.59	4.56	62.29	5.09	4.57	5.601196038	0.032520325	0.06910569	16.8467
14516	Cnot3: CCR4-NOT transcription complex, subunit 3	796.98	668.81	778.76	366.48	721.23	392.7	402.2	266.4	437.76	1.735481277	0.048182587	0.09296422	11.6855
3016	Cops7b: COP9 (constitutive photomorphogenic) homolog, subunit 7b (Arabidopsis thaliana)	630.4	592.56	784.56	381.57	313.14	269.3	386	255.5	166.93	2.265261447	0.027422303	0.06459476	17.9807
9970	Csrp1: cysteine and glycine-rich protein 1	592.11	830.02	1000.51	402.09	254.56	515.3	117.8	186.2	282.36	2.755708736	0.014388489	0.03784173	33.681
42439	Cxcl1: chemokine (C-X-C motif) ligand 1	22.86	80.3	90.94	1.97	4.77	2.02	4.71	2.48	4.31	19.16090819	0.033639144	0.07090724	16.3534
3504	Cxcl1: chemokine (C-X-C motif) ligand 1	47.52	119.73	89.85	17.25	11.6	15.44	9.7	10.43	6.75	7.224954335	0.036904762	0.07630159	14.5178
16786	Cxcr4: chemokine (C-X-C motif) receptor 4	358.53	513.32	197.69	121.26	183.91	87.45	18.1	32.91	17.61	4.637672361	0.010526316	0.03778947	38.9113
30909	Cyp11a1: cytochrome P450, family 11, subfamily a, polypeptide 1	258.01	292.08	442.1	107.55	178.36	106.1	178.5	12.51	54.95	3.11031348	0.032051282	0.06915064	16.7746
41238	Cyp13: cysteine-rich perinuclear theca 3	423.83	155.74	395.11	142.52	146.78	84.29	153.7	281.9	54.27	2.257667006	0.036855037	0.07558149	14.7418

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
17076	D10Jhu81e: DNA segment, Chr 10, Johns Hopkins University 81 expressed	1166.3	278.18	584.98	41.06	273.73	194.1	342.2	66.45	240.74	3.504143933	0.01744186	0.0394186	31.3457
21185	D10Wsu102e: DNA segment, Chr 10, Wayne State University 102, expressed	2792.95	2498.7	2814.16	1716.1	1847	1338	1962	1712	1576.6	1.596805904	0.026666667	0.06302222	18.3654
31660	D11Ert497e: DNA segment, Chr 11, ERATO Doi 497, expressed	1363.15	1027.4	1189.28	742.66	758.15	478.8	680.6	595	536.27	1.888339413	0.027726433	0.06448552	18.0557
32944	D130073L02Rik: RIKEN cDNA D130073L02 gene	1475.86	1476.2	1683.68	818.11	1061.5	984.4	665.5	689	969.73	1.787029488	0.03422619	0.0715377	16.1158
19130	D14Ert500e: DNA segment, Chr 14, ERATO Doi 500, expressed	681.58	601.92	716.43	390.44	459.16	256.8	360.4	265.9	425.26	1.853614907	0.045226131	0.08424791	13.0523
29688	D2Ert612e: DNA segment, Chr 2, ERATO Doi 612, expressed	319.43	176.06	370.14	220.95	120.36	106	116.2	108.1	103.63	2.233105886	0.043294615	0.08247448	13.4481
7623	D2Ert750e: DNA segment, Chr 2, ERATO Doi 750, expressed	394.9	506.08	652.28	243.42	291.26	438	219.8	255.7	110.89	1.9925724	0.047619048	0.09278752	11.6257
17383	Dbndd1: dysbindin (dystrobrein binding protein 1) domain containing 1	464.15	301.44	304.82	104.24	166.01	137.6	190.5	70.39	112.63	2.739969027	0.037174721	0.07516316	14.8291
41824	Dbx1: developing brain homeobox 1	276.19	187.2	124.35	38.52	58.77	7.62	29.55	11.63	10.84	7.49047346	0.01459854	0.03824818	33.796
6290	Dcp1a: decapping enzyme	577.73	499.84	520.9	251.84	361.9	184	289.1	278.2	245.94	1.984530675	0.04600939	0.08865102	12.4452
542	Dctn3: dynactin 3	540.25	352.08	532.14	252.9	202.65	271.3	222	154	123.8	2.322461258	0.02244389	0.05806318	20.7509
1811	Ddit3: DNA-damage inducible transcript 3	499.34	650.01	844	254.47	351.31	188.1	199	274.8	371.53	2.432175213	0.022727273	0.05471591	22.2271
9363	Ddx19a /// Ddx19b: DEAD (Asp-Glu-Ala-Asp) box polypeptide 19a /// DEAD (Asp-Glu-Ala-Asp) box polypeptide 19b	1578.61	1941.1	1533.78	1167.2	1152.7	885.9	897.4	1007	1064.7	1.636896915	0.048	0.09173926	12.0151

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
2222	Ddx19a: DEAD (Asp-Glu-Ala-Asp) box polypeptide 19a	1009.15	1114.6	1292.36	497.39	638.59	587.9	581.9	501.6	556.5	2.031071632	0.024154589	0.0584219	20.4566
36061	Dgat1: diacylglycerol O-acyltransferase 1	289.38	324.34	39.41	98.71	107.58	55.58	34.36	61.27	22.93	3.43364088	0.027600849	0.059908	19.4349
24651	Dhfr: dihydrofolate reductase	426.78	75.47	340.75	63.57	73.72	47.34	58.38	22.96	36.07	5.582042114	0.018867925	0.03471698	49.1336
16466	Dhrs3: dehydrogenase/reductase (SDR family) member 3	175.77	358.46	397.37	163.62	207.39	44.67	59.21	66.67	115.31	2.836482105	0.02247191	0.05490637	22.1263
36303	Diras2: DIRAS family, GTP-binding RAS-like 2	83.58	244.42	311.89	57.09	96.41	29.57	84.61	77.33	98.72	2.884141257	0.03164557	0.06949895	16.6642
14159	Dmpk: dystrophia myotonic-protein kinase	130.39	64.65	251.65	43.94	24.65	45.01	36.88	47.29	31.12	3.903097558	0.044573643	0.08648256	12.7504
18463	Dnab5: DnaJ (Hsp40) homolog, subfamily B, member 5	173.41	174.8	38.49	23.48	43.07	40.6	23.62	34.57	27	4.021004471	0.024943311	0.05890401	20.0001
1165	Dnalc4: dynein, axonemal, light chain 4	627.08	485.5	527.62	152.51	208.44	124.3	259.5	384	378.63	2.176154117	0.037783375	0.07515533	14.9173
4618	Dscr6: Down syndrome critical region homolog 6 (human)	364.43	202.42	53.05	100.55	71.25	41.67	57.79	52.26	26.72	3.539858383	0.033383915	0.07102175	16.2903
14348	Dsp: desmoplakin	348.74	368.07	172.59	17.12	59.21	37.98	120.1	62.81	35.6	5.344951923	0.018181818	0.03509091	48.1504
44606	Dynlt3: dynein light chain Tctex-type 3	1184.16	850.7	1225.34	804.47	862.49	821.8	513.8	484.1	546.26	1.616793787	0.044487427	0.0865087	12.7286
30501	E130311K13Rik: RIKEN cDNA E130311K13 gene	292.14	72.05	149.12	73.07	44.21	47.25	43.95	46.76	53.87	3.321212513	0.048373645	0.09281346	11.6139
1360	Egr1: early growth response 1	863.82	594.38	874.96	651.4	743.85	349.3	362.2	301.7	296.64	1.725097598	0.047790803	0.0904749	12.1522
28198	Ehd2 /// LOC673251: EH-domain containing 2 /// similar to EH-domain containing 2	552.22	499.34	187.9	118.31	170.8	197.3	120	329.7	107.65	2.375217983	0.024691358	0.0582963	20.6613
3769	Ehf: ets homologous factor	422.95	299.88	362.77	181.23	308.34	201	118.5	82.28	115.52	2.156471301	0.048128342	0.09172906	12.0251
6059	Eif2ak1: eukaryotic translation initiation factor 2 alpha kinase 1	601.03	422.26	534.76	313.33	270.08	133.6	249.2	221.6	232.52	2.194034895	0.032019704	0.06823755	16.968

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
22574	Eif2b1: eukaryotic translation initiation factor 2B, subunit 1 (alpha)	639.36	591.13	452.83	426.81	356.95	299.3	292.9	218.6	271.63	1.804037146	0.047939445	0.09294365	11.6628
41276	Emx2: empty spiracles homolog 2 (Drosophila)	234.91	298.01	341.22	97.13	165.98	119.9	85.52	182	66.68	2.437578428	0.037585421	0.078265	14.0886
27167	Faah: fatty acid amide hydrolase	147.03	240.15	277.1	108.99	47.52	58.21	61.45	124.1	50.02	2.950192081	0.030252101	0.06778151	17.1504

WO 2010/003020

PCT/US2009/049447

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
38293	Fank1: fibronectin type 3 and ankyrin repeat domains 1	2043.72	1333.2	2424.93	1297.9	895.96	1029	224.5	906.6	897.4	2.20974613	0.014705882	0.03808824	33.9309
38294	Fank1: fibronectin type 3 and ankyrin repeat domains 1	924.97	617.49	1115.41	530.43	486.47	334.5	325.3	374.2	450.52	2.125114436	0.022792023	0.0548433	22.228
28859	Fbxw14 /// LOC668758 /// LOC673189: F-box and WD-40 domain protein 14 /// similar to F-box and WD-40 domain protein 14 /// similar to F-box and WD-40 domain protein 14	1689.47	1449.1	1379.4	771.7	977.11	774	924.7	979.8	930.52	1.686457855	0.04250797	0.08190932	13.5148
17631	Fetub: fetuin beta	339.98	98.9	176.78	28.16	22.02	28.89	47.19	149.6	27.08	4.064701416	0.030050083	0.06791319	17.1106
18896	Fgfl: fibroblast growth factor 1	681.86	369.41	720.7	327.77	349.5	445.6	311.8	285.3	249.7	1.799219174	0.038548753	0.07857143	14.042
16430	G6pdx: glucose-6-phosphate dehydrogenase X-linked	1968.98	1787.5	2156.26	787.91	1751.9	1060	912.4	963.3	1065.4	1.807953743	0.022900763	0.05866836	20.8236
20610	Galm: galactose mutarotase	325.39	308.96	339.4	107.5	239.54	178.8	127.7	148.9	72.69	2.225459947	0.047488584	0.09021005	12.2255
32346	Gas2l1: growth arrest-specific 2 like 1	161.84	235.35	19.91	17.07	61.3	43.92	20.12	11.33	52.21	4.050497694	0.045454545	0.0852668	12.9177
12777	Gch1: GTP cyclohydrolase 1	398.77	412.01	533.59	236.21	254.72	224.9	133.7	148.3	134.69	2.37426818	0.0373599	0.07496056	14.8721
38302	Ghitm: Growth hormone inducible transmembrane protein	167.4	64.72	61.47	5.07	13.4	2.06	3.34	4.13	8.35	16.15350757	0.037714286	0.07841524	14.1066
27801	Gm129: gene model 129, (NCBI)	962.68	506.38	860.85	158.2	317.4	272.4	320.2	434.4	279.11	2.615437289	0.014925373	0.03823383	34.0765
31138	Gm1568: gene model 1568, (NCBI)	543.23	1089.3	807.32	324.93	568.66	388.6	233.9	198.5	219.96	2.522390451	0.01754386	0.04565789	27.2403

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
16058	Gm428 /// LOC623180 /// LOC623197 /// LOC623210 /// LOC623219: gene model 428, (NCBI) /// hypothetical LOC623180 /// hypothetical LOC623197 /// hypothetical LOC623210 /// hypothetical LOC623219	1914.89	1546.2	1987.58	718.98	1301.8	938	1086	1344	986.94	1.709161859	0.026694045	0.06009582	19.1631
17875	Gna14: guanine nucleotide binding protein, alpha 14	300.19	342.97	390.91	109.58	273.46	141.3	122.7	96.2	128.54	2.372264281	0.037558685	0.07721831	14.351
27857	Gnaz: guanine nucleotide binding protein, alpha z subunit	291.26	148.83	411.78	101.45	157.63	114.7	126.2	93.34	143.95	2.311190092	0.045837231	0.08869036	12.4247
3764	Gnb4: guanine nucleotide binding protein, beta 4	800.06	500.22	743.39	483.85	537.75	253.1	157.9	393.7	258.05	1.960947433	0.033674963	0.07097609	16.075
1723	Gng3: guanine nucleotide binding protein (G protein), gamma 3 subunit	1141.11	1006.1	938.92	376.96	611.22	390.9	606.5	482.2	588.54	2.019579614	0.026369168	0.06075727	19.0026
38463	Gng4: guanine nucleotide binding protein (G protein), gamma 4 subunit	29.31	68.46	318.22	20.93	10.51	15.87	14.53	12.26	10.91	9.786848606	0.04808476	0.09437924	11.4171
14851	Gng5: guanine nucleotide binding protein (G protein), gamma 5 subunit	751.62	1021.2	1032.02	380.3	617.1	541.3	307.9	450.5	359.29	2.111806714	0.021428571	0.05125	24.5286
19438	Gprc5b: G protein-coupled receptor, family C, group 5, member B	306.88	118.93	6.69	4.77	9.28	5.32	8.02	2.74	32.26	13.86440135	0.023346304	0.05102464	25.4672
14609	Grina: glutamate receptor, ionotropic, N-methyl D-aspartate-associated protein 1 (glutamate binding)	1034.42	812.15	1228.38	477.25	540.8	546.5	285.4	670.3	531.6	2.015164771	0.027088036	0.0592927	19.9298
15539	Grn: granulin	1352.59	514.79	1057.39	324.87	349.47	237.6	326.6	429.6	434.48	2.782090489	0.012820513	0.03683761	41.8478
21945	Grn: granulin	1042.32	422.35	831.7	282.08	293.46	176.8	252.5	333.1	327.99	2.756746699	0.017094017	0.03763533	35.8799
44280	Gyg: glycogenin	1636.36	1537.6	1912.79	587.38	955.09	642.4	640.7	735.8	665.37	2.406956662	0.01754386	0.03795322	36.2379
16505	Gyg: glycogenin	1843.18	1330	1474.57	451.38	788.27	545	633.2	808.2	736.02	2.346160253	0.01863354	0.03867495	31.8723
21695	Hadhscl: L-3-hydroxyacyl-Coenzyme A dehydrogenase, short chain	176.86	172.31	47.15	7.97	25.17	8.67	37.31	48.56	24.68	5.202415332	0.024475524	0.05073427	24.4128

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
20748	Hba-a1: hemoglobin alpha, adult chain 1	145.01	91.26	185.79	31.03	31.41	48.27	14.05	70.23	28.93	3.769739193	0.037667072	0.07575132	14.6733
34409	Hdlbp: High density lipoprotein (HDL) binding protein	65.43	20.02	222.9	4.86	15.41	2.9	23.01	5.26	6.4	10.66217151	0.048414023	0.09280467	11.6202
21722	Hes5: hairy and enhancer of split 5 (Drosophila)	77.81	670.65	255.74	37.89	6.51	4.11	5.4	49.42	10.01	17.72013411	0	0.03571429	76.3251
14548	Hes6: hairy and enhancer of split 6 (Drosophila)	874.59	449.21	52.72	14.43	115.08	53.85	47.95	155.1	45.4	6.375581853	0	0.03063063	57.3556
294	Hey1: hairy/enhancer-of-split related with YRPW motif 1	782.14	951.79	1330.97	370.68	579.38	458.1	340.9	320.2	328.5	2.556469371	0.019900498	0.04260365	29.0333
40573	Hist2h2be: histone 2, H2be	45.26	112.92	234.65	9.51	47.79	32.3	12.23	14.75	13.01	6.062659156	0.032357473	0.07042116	16.4512
15290	Hk1: hexokinase 1	97.53	40.72	292.16	23.48	13.09	10.02	17.7	8.41	70.37	6.016775005	0.045279383	0.08702954	12.6818
1997	Hnmt: histamine N-methyltransferase	291.41	218.29	287.13	133.75	161.6	140.3	63.03	104.4	82.11	2.325899764	0.042619543	0.08295565	13.3018
20448	Hoxa3: homeo box A3	67.37	168.06	195.07	30.43	54.67	3.04	114.4	21.42	18.67	3.549198236	0.036643026	0.07695823	14.4359
38029	Hspa12a: heat shock protein 12A	99.91	395.12	430.36	63.95	34.34	2.89	11.27	190.3	15.72	5.810743776	0	0.02886179	56.2068
20415	Hspa1a: heat shock protein 1A	665.51	521.22	682.91	384.53	418.07	300.9	177.6	256.6	240.21	2.103200405	0.026717557	0.06280534	18.4144
20345	Hspa1b: heat shock protein 1B	1356.32	1044	604.27	612.34	519.36	458.8	326.2	890.6	200.5	1.997912074	0.030201342	0.06785794	17.1351
11286	Hspa1b: heat shock protein 1B	603.12	550.63	273.69	292.64	253.66	243.1	178.7	397.6	109.8	1.934869094	0.04518664	0.08519974	12.8968
1396	Hspa2: heat shock protein 2	1671.68	1450.9	1466.81	768.33	1325.6	888	653	1010	758.87	1.698705433	0.041845494	0.08110515	13.6108
3507	Icosl: icos ligand	698.31	470.26	803.79	259.27	478.54	329.9	299.9	350.5	227.76	2.027216478	0.030508475	0.06688701	17.2922
14244	Id2: inhibitor of DNA binding 2	159.2	175.61	225	87.31	76.02	44.55	20.82	47.44	37.67	3.567827666	0.036262204	0.0725709	15.7007
3942	Ier3: immediate early response 3	135.39	282.8	200.44	43.45	88.87	51.5	102.8	52.75	30	3.349649403	0.032407407	0.07047325	16.4563

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
5812	Igh-VJ558 /// LOC238447 /// LOC544903 /// LOC544907: immunoglobulin heavy chain (J558 family) /// similar to immunoglobulin heavy chain variable region /// similar to immunoglobulin mu-chain /// similar to anti-poly(dC) monoclonal antibody heavy chain	796.82	447.15	483.63	357.22	539.3	266.5	77.64	114.2	116.29	2.348686715	0.047531993	0.09001828	12.2414
18787	Ing3: inhibitor of growth family, member 3	797.55	862.17	1017.87	636.48	680.07	589	200.3	400.2	302.52	1.906748702	0.041980624	0.08096161	13.6303
40325	Insig1: insulin induced gene 1	2692.54	2321.2	2684.68	1690.3	1802.6	1778	1643	1631	1533.5	1.527618507	0.035150646	0.07234338	15.8472
16512	Irf1: interferon regulatory factor 1	789.3	575.41	867.38	269.32	522.82	348.8	278.6	292.6	194.46	2.341447296	0.022900763	0.05101781	25.2531
2812	Irx3: Iroquois related homeobox 3 (Drosophila)	348.55	332.29	138.19	5.68	121.77	63.47	127.6	69.42	60.1	3.656465546	0.017857143	0.04482143	27.5757
18056	Itga9: integrin alpha 9	602.24	988.31	1038.91	552.93	662.74	418.5	555.2	457.9	546.34	1.646726537	0.042268041	0.08298625	13.2595
1704	Jun: Jun oncogene	824.08	580.75	436.92	74.84	119.42	26.57	109.4	37.89	28.35	9.290272138	0	0.02583333	113.684
38760	Kalrn: kalirin, RhoGEF kinase	455.34	326.78	526.39	276.27	277.62	210.1	85.33	54.31	126.1	2.541487006	0.028697572	0.05972774	19.7266
17540	Kcnq1: potassium voltage-gated channel, subfamily Q, member 1	253.97	94.35	476.89	62.84	42.18	56.37	133.6	55.74	24.75	4.395844986	0.020661157	0.04779614	26.4773
38646	Kif13a: Kinesin family member 13A	341.25	421.32	771.62	218.81	258.08	529	77.71	47.73	84.33	2.524106842	0.047169811	0.09193539	11.8117
25517	Kif24: kinesin family member 24	338.53	277.95	398.92	132.82	85.27	159.4	147.6	148.5	106.19	2.604290899	0.029801325	0.06786424	17.0703
16345	Klhl13: kelch-like 13 (Drosophila)	1515.8	1527.5	1239.11	1454.6	785.49	962.1	538.2	329.8	445.64	1.896626556	0.026966292	0.05925094	19.8985
13958	Ldhb: lactate dehydrogenase B	2740.69	2312.3	3295.47	1762	2507	1685	1417	2027	1678.1	1.507554068	0.045105566	0.087508	12.6371
5980	Ldlr: low density lipoprotein receptor	335.33	308.94	424.04	117.76	245.41	172	99.41	87.75	147.65	2.456054441	0.038283063	0.07753674	14.2499

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
29964	Ldlrad3: low density lipoprotein receptor class A domain containing 3	349.24	268.52	370.69	153.24	103.08	147.7	93.05	223.7	96.65	2.418344628	0.033873343	0.07123711	16.0846
916	Llgl1: lethal giant larvae homolog 1 (Drosophila)	1297.45	1170.9	1328.72	919.12	927.65	845	526.8	632.3	506.16	1.742963703	0.036451169	0.07264099	15.6391
21355	Lmo7: LIM domain only 7	2299.11	2394	2530.26	1703.4	2212.2	1824	1271	1373	1270.7	1.496467727	0.04749788	0.09230421	11.7338
32615	LOC240444: Similar to Potassium voltage-gated channel subfamily G member 2 (Voltage-gated potassium channel subunit Kv6.2) (Cardiac potassium channel subunit)	218.5	275.91	425.72	86.21	127.38	61.45	46.82	105.4	143.32	3.22507492	0.026315789	0.06088394	18.9799
14891	LOC545161 /// LOC637273: similar to Peroxiredoxin 1 (Thioredoxin peroxidase 2) (Thioredoxin-dependent peroxide reductase 2) (Osteoblast specific factor 3) (OSF-3) (Macrophage 23 kDa stress protein) /// similar to Peroxiredoxin-1 (Thioredoxin peroxidase 2) (Thioredoxin-dependent peroxide reductase 2) (Osteoblast-specific factor 3) (OSF-3) (Macrophage 23 kDa stress protein)	1517.86	1912.8	1961.88	1213.5	1373.7	1085	657.6	832.6	661.9	1.851793474	0.023890785	0.05119454	24.0588
19912	LOC574530: metallothionein 1K	226.98	259.54	118.5	143.89	63.21	9.37	10.47	15.52	12.43	4.747302758	0.018957346	0.04349131	28.4084

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
34813	LOC627488 /// LOC627520 /// LOC667692 /// LOC667695 /// LOC673990: similar to THO complex subunit 4 (Tho4) (RNA and export factor binding protein 1) (REF1-I) (Ally of AML-1 and LEF-1) (Aly/REF) /// similar to THO complex subunit 4 (Tho4) (RNA and export factor binding protein 1) (REF1-I) (Ally of AML-1 and LEF-1) (Aly/REF) /// similar to THO complex subunit 4 (Tho4) (RNA and export factor binding protein 1) (REF1-I) (Ally of AML-1 and LEF-1) (Aly/REF) /// similar to THO complex subunit 4 (Tho4) (RNA and export factor binding protein 1) (REF1-I) (Ally of AML-1 and LEF-1) (Aly/REF) /// similar to THO complex subunit 4 (Tho4) (RNA and export factor binding protein 1) (REF1-I) (Ally of AML-1 and LEF-1) (Aly/REF)	3712.88	3330.1	4059.81	2547.1	3079.4	2572	1852	2591	2371.8	1.479044054	0.036719706	0.07589147	14.7015

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
36676	LOC665521 /// LOC665546 /// LOC665755 /// LOC666203: similar to PRAME family member 8 /// similar to PRAME family member 8 /// similar to PRAME family member 8 /// similar to PRAME family member 8	1690.74	1078.8	1867.35	435.93	1040.6	413.3	298	288.9	409.86	3.212720848	0	0.02811594	69.6907
35372	LOC666806: similar to X- linked eukaryotic translation initiation factor 1A	1437.33	1258.1	1431.69	830.47	904.29	911.7	639.8	803.3	394.95	1.840602429	0.028508772	0.06025585	19.6427
23907	LOC677447: similar to RIKEN cDNA 5730590G19- like	169.23	96.13	289.52	47.67	49.17	46.11	50.1	30.45	38.54	4.235078614	0.026415094	0.06295597	18.313
2237	Lypd3: Ly6/Plaur domain containing 3	398.36	354	401.18	228.49	210.36	118.7	246.3	99.17	235.65	2.026189368	0.047863248	0.09232479	11.7737
22323	Lysmd3: LysM, putative peptidoglycan-binding, domain containing 3	453.85	368.82	661.12	92.84	321.95	197.1	125	290.1	91.81	2.6523721	0.026748971	0.0601166	19.1722
14392	Marcks11 /// LOC673071: MARCKS-like 1 /// similar to MARCKS-related protein (MARCKS-like protein 1) (Macrophage myristoylated alanine-rich C kinase substrate) (Mac-MARCKS) (MacMARCKS) (Brain protein F52)	1185.33	1143.2	1024.39	984.94	699.06	786.1	242.8	549.7	257.94	1.904808706	0.034833091	0.07184809	15.9639
14326	Marcks11: MARCKS-like 1	1090.15	1065.5	915.39	902.84	654.38	711.6	198.7	505.8	271.67	1.892871561	0.036890646	0.07380764	15.2728
9962	Mbd2: methyl-CpG binding domain protein 2	705.4	147.62	1019.73	112.36	370.48	107.5	65.92	14.24	271.83	3.974764411	0.014285714	0.036	43.8036
1105	Meal: male enhanced antigen 1	542.92	668.6	745.81	373.15	443.27	291.3	373	331.2	364.96	1.798256252	0.047787611	0.09168732	11.9939
14305	Mm.10009.3	245.87	217.16	188.38	53.3	153.42	55.43	78.1	79.43	58.1	2.726819875	0.04253112	0.08289765	13.2956
43719	Mm.150985.1	393	518.17	908.05	363.36	370.12	406.2	195.3	152.2	111.3	2.276315543	0.045053869	0.08550767	12.8666

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
4333	Mm.25004.1	145.84	123.93	76.52	26.6	44.95	15.88	28.28	3.92	34.91	4.481558173	0.037313433	0.07497512	14.8637
9278	Mm.37131.1	290.79	409.6	147.87	153.64	67.38	113.6	173	47.35	77.8	2.681312429	0.035100287	0.07225406	15.847
38596	Mm.45490.1	458.52	300.11	583.52	157.64	158.29	215.6	52.94	334	76.83	2.696759027	0.024193548	0.05605735	21.6519
22096	Mm.69144.1	564.45	289.9	595.48	130.38	120.9	231.2	100.7	88.48	177.73	3.41381462	0.012345679	0.0373251	41.2274
21487	Mm.9772.3	207.88	220.36	148.09	29.05	85.16	103.9	122.9	53.97	65.17	2.504911335	0.04757859	0.0923676	11.7365
8468	Mocs2: molybdenum cofactor synthesis 2	451.72	641.1	548.64	204.64	269.88	172.8	296.6	378.2	132.88	2.256395453	0.02972028	0.06585082	17.5962
29403	Mrpl9: mitochondrial ribosomal protein L9	687.37	522.93	342.88	378.66	145.59	253.6	220.5	250.8	282.44	2.028206166	0.043112513	0.08283912	13.3899
6034	Mrps23: mitochondrial ribosomal protein S23	189.41	237.99	295.52	122.32	119.68	82.57	55.82	57.01	56.99	2.924492809	0.035310734	0.07228343	15.7753
44561	Mrps23: mitochondrial ribosomal protein S23	133.78	165.37	174.49	82.76	82.15	57.97	48.91	9.82	35.54	2.986851647	0.048430493	0.09099851	12.1046
18672	Mt4: metallothionein 4	471.49	401.25	465.5	283.03	160.15	374.6	77.11	182.5	150.05	2.180449535	0.037926675	0.07498946	14.9389
8584	Mtap: methylthioadenosine phosphorylase	301.33	399.52	615.48	133.87	304.21	110.8	88.98	14.43	153.22	3.268395635	0.025559105	0.05434505	23.2205
3549	Mthfd2: methylenetetrahydrofolate dehydrogenase (NAD+ dependent), methenyltetrahydrofolate cyclohydrolase	983.54	1203.2	1182.24	590.25	760.27	536.8	554.4	527.2	690.45	1.841304716	0.02661597	0.06293409	18.3588
27588	Mxd1: MAX dimerization protein 1	993.73	911.74	644.1	466.93	617.29	512.7	418	374.6	279.54	1.910455702	0.029255319	0.06550827	17.6984
6161	Mxd1: MAX dimerization protein 1	181.67	333.48	504.31	168.5	123.99	172.7	110.9	62.78	111.84	2.716133587	0.029462738	0.06636049	17.5011
20325	Myo5b: myosin Vb	768.64	656.09	488.09	409.12	389.77	359	270.7	380	263.4	1.84641373	0.044616877	0.08649855	12.7532
29664	Myom2: myomesin 2	1192.6	1602.7	1540.02	1131.7	1059.7	972	571.8	941.8	785.78	1.587230631	0.043340381	0.082537	13.4485
28881	Nalp4f: NACHT, leucine rich repeat and PYD containing 4F	2150.11	2492.3	2718.12	1515.2	1789.6	1160	566.2	1035	625.39	2.199965628	0.010638298	0.03808511	38.9565
11528	Nalp6: NACHT, leucine rich repeat and PYD containing 6	317.78	138.89	39.96	17.77	77.33	22.29	55.67	9.16	5.59	5.288642777	0.031695721	0.06955098	16.6649
41896	Nalp9b: NACHT, LRR and PYD containing protein 9b	743.42	818.13	815.08	320.66	556.55	474.3	285.5	410.1	285.42	2.037874176	0.026974952	0.06290944	18.4675

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
18033	Ncoa4 /// LOC627557: nuclear receptor coactivator 4 /// similar to nuclear receptor coactivator 4	552.62	540.61	580.1	250.08	280.29	212.8	299.8	200.4	190.95	2.333403057	0.027108434	0.06128514	18.8762
21797	Ndr1: N-myc downstream regulated gene 1	760.12	747.8	336.89	280.11	487.1	357.4	374.4	244.3	197.19	1.901454324	0.031796502	0.0690938	16.7306
4919	Ndr1: N-myc downstream regulated-like	442.34	476.68	210.38	300.65	162.3	171.9	239.9	189.1	144.37	1.869588969	0.048053024	0.0935156	11.5462
10414	Nefl: neurofilament, light polypeptide	1726.22	1655.4	1459.28	902.95	1121	600.9	759.2	965.7	548.27	1.976692432	0.022813688	0.05089987	25.2396
2584	Nes: nestin	259.68	196.54	282.1	75.72	199.89	71.08	119.4	87.37	86.2	2.308404202	0.046753247	0.09178066	11.8681
6998	Neurog2: neurogenin 2	231.05	418.13	218.55	123.46	78.36	80.47	63.67	209.6	71.81	2.766114122	0.047928513	0.09439751	11.403
9918	Nobox: NOBOX oogenesis homeobox	396.92	408.09	475.01	129.03	371.41	87.59	172.8	98.82	245.05	2.317449397	0.038327526	0.07741773	14.2596
17216	Nudcd2: NudC domain containing 2	1083.58	1300.2	1922.19	656.24	404.23	762.1	293	236.5	633.56	2.884501117	0.012987013	0.03718615	41.8566
2861	Nudcd2: NudC domain containing 2	653.94	528.12	803.02	415.16	310.05	389.2	346.4	283.8	405.47	1.846517339	0.047745358	0.09168288	11.986
3960	Nupr1: nuclear protein 1	312.62	225.23	176.51	47.1	36.87	54.82	8.86	46.3	100.04	4.859757135	0.018099548	0.04529412	27.5871
3961	Nupr1: nuclear protein 1	208.1	133.13	133.4	11.64	27.13	47.78	1.98	17.52	68.53	5.437392599	0.029982363	0.06595532	17.6434
2981	Oas1c: 2'-5' oligoadenylate synthetase 1C	1384.52	1483.8	1747.71	600.25	1009	720.6	552.8	343.8	414.35	2.535773144	0.00990099	0.03722772	37.9655
1142	Oas1d /// Oas1e: 2'-5' oligoadenylate synthetase 1D /// 2'-5' oligoadenylate synthetase 1E	2035.81	1982.1	2708.21	777.35	1074.7	732.5	695	757.1	651.09	2.869641642	0	0.02992754	53.6191
19318	Obfc2b: oligonucleotide/oligosaccharid e-binding fold containing 2B	433.1	115.04	679.94	161.64	146	98.16	107.4	78.98	143.91	3.336675225	0.020512821	0.04188034	29.5008
10199	Odf2: outer dense fiber of sperm tails 2	1037.05	1156.3	1111.85	662.64	858.1	614	392.7	651.3	460.77	1.816309428	0.035	0.0724	15.828
22813	Osbp2: oxysterol binding protein 2	269.05	335.57	568.18	144.89	117.87	130.8	68.72	106.4	53.36	3.771121722	0.016666667	0.04731944	26.6072
43535	Pak7: P21 (CDKN1A)- activated kinase 7	394.46	1077	817.61	505.83	477.53	544	153	241.8	217.14	2.13996971	0.027322404	0.06455373	17.9674

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
41333	Paqr5: progesterin and adipoQ receptor family member V	724.43	758.22	613.93	502.88	471.52	563.1	113.6	175.4	120.97	2.153154125	0.032102729	0.06917603	16.7783
849	Pdlim1: PDZ and LIM domain 1 (elfin)	300.99	412.46	570	141.35	241.6	138.6	163.9	202.1	181.18	2.402069959	0.033923304	0.07121436	16.0941
41503	Pex10 /// LOC668173 /// LOC671348: peroxisome biogenesis factor 10 /// similar to peroxisome biogenesis factor 10 isoform 1 /// similar to Peroxisome assembly protein 10 (Peroxin-10) (Peroxisome biogenesis factor 10) (RING finger protein 69)	280.61	206.54	96.17	72.22	80.02	54.72	83.5	24.39	29.12	3.391691136	0.028776978	0.06503597	17.8229
12239	Pgam5: phosphoglycerate mutase family member 5	110.6	127.3	151.21	91.12	15.55	11.93	56.81	11.64	29.05	3.601203147	0.046956522	0.09189855	11.8815
6474	Phkg1: phosphorylase kinase gamma 1	178.41	106.98	171.74	40.05	80.47	88.56	39.25	46.03	40.22	2.732560225	0.04787234	0.09175532	11.9989
4052	Pitpm2: phosphatidylinositol transfer protein, membrane-associated 2	84.85	63.94	331.06	11.48	54.66	9.66	23.36	16.57	16.8	7.24137931	0.046918123	0.08945109	12.2952
19362	Plac8: placenta-specific 8	662.3	661.38	795.38	468	575.56	338.1	183	196.8	196.89	2.164105864	0.033823529	0.07121569	16.08
20205	Plec1 /// LOC671535: plectin 1 /// similar to poly (ADP-ribose) polymerase family, member 10	121.35	70.86	157.14	22.86	52.58	29.9	14.04	42.18	35.14	3.552109812	0.047738693	0.09283082	11.6437
27915	Plekhl1: pleckstrin homology domain containing, family G (with RhoGef domain) member 1	1853.45	1666.1	1893.1	1079.7	1334.2	984	491.4	722.9	597.2	2.078008684	0.015873016	0.0412522	29.9928
11164	Plk2: polo-like kinase 2 (Drosophila)	498.11	383.42	423.89	177.35	110.99	143.6	134.9	38.56	65.38	3.89207078	0.016949153	0.03774011	35.7785
15242	Pofut2: protein O-fucosyltransferase 2	1778.06	1154.3	2374.19	882.1	1201.9	822.3	491.1	875.2	1214.3	1.934223068	0.02484472	0.05419255	23.0226
14202	Pofut2: protein O-fucosyltransferase 2	841.36	629.59	1545.03	432.55	580.35	395.1	328.1	451.2	735.19	2.064043252	0.042842215	0.08301289	13.3317

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
10997	Pold3: polymerase (DNA-directed), delta 3, accessory subunit	1231.65	1183.2	1391.33	824.72	900.21	640.4	329.6	569.7	526.55	2.007913041	0.023980815	0.05843325	20.4291
34590	Pold3: polymerase (DNA-directed), delta 3, accessory subunit	423.34	669.16	843.98	290.44	334.78	246.5	303.9	237.9	260.63	2.313402702	0.024523161	0.05544959	21.8711
38961	Ppp1r3d: protein phosphatase 1, regulatory subunit 3D	575.14	503.37	526.53	166.64	221.68	213.9	94.06	104	89.52	3.607723257	0.009615385	0.03676282	37.8147
36182	Ppp4c: protein phosphatase 4, catalytic subunit	571.66	453.52	340.78	131.19	406.4	205.3	175.5	111.2	158.93	2.298802602	0.035050072	0.07244158	15.8315
14726	Prdx1: peroxiredoxin 1	2781.18	3417.9	3506.42	2190.9	2626.8	2285	1226	1574	1195.6	1.749054555	0.017094017	0.04625356	26.9776
14063	Prdx1: peroxiredoxin 1	2078.16	2456.5	2879.78	1833.6	1734.2	1642	870.6	902.3	801.35	1.904999994	0.022088353	0.04886212	26.0918
13692	Prdx1: peroxiredoxin 1	1982.74	2435.2	2494.71	1597.2	1890.1	1553	859.4	1144	886.29	1.743495	0.023460411	0.05467253	22.4906
295	Prdx1: peroxiredoxin 1	687.71	1005.3	1004.39	538.28	643.24	491.2	434.4	689.3	439.9	1.666929311	0.041019956	0.08053954	13.8032
12996	Prdx2: peroxiredoxin 2	2137.92	1606.1	1921.64	1079.9	1587.9	1273	690.7	1588	625.66	1.655379356	0.037457435	0.07846008	14.0576
2801	Prdx2: peroxiredoxin 2	3503.34	3323.1	3762.91	2292.1	2644.3	1942	2586	2530	2441.4	1.467059107	0.041800643	0.08119328	13.6017
42136	Prmt3: protein arginine N-methyltransferase 3	165	52.27	96.5	11.63	3.78	14.62	15.11	34.44	6.54	7.286809104	0.035302594	0.07236792	15.8727
2615	Prss8: protease, serine, 8 (prostasin)	471.01	140.12	249.79	75.51	130.38	26.33	79.43	130.7	70.09	3.36008118	0.029151943	0.06553592	17.6734
8719	Pstpip1: proline-serine-threonine phosphatase-interacting protein 1	44.1	110.11	91.14	10.97	19.4	25.08	18.91	12.49	10.27	5.052512356	0.047161572	0.09181659	11.9108
43158	Ptdss2: phosphatidylserine synthase 2	1203.34	418.9	557.48	508.13	304.61	307.9	188.3	147	386.5	2.366084658	0.046685341	0.08895425	12.3985
41102	Rab3gap1: RAB3 GTPase activating protein subunit 1	460.17	347.13	470.57	199.81	233.06	254.1	132.1	145	138.52	2.317921277	0.026266417	0.06385866	18.2065
16961	Rap2b: RAP2B, member of RAS oncogene family	808.09	565.99	369.3	354.3	287.58	166.3	57.48	218.1	87	2.978227632	0.018604651	0.04384496	28.1739
8013	Ras11b: RAS-like, family 11, member B	426.89	253.01	172	67.51	155.36	83.97	28.63	67.32	76.13	3.557587906	0.022988506	0.0580613	20.1914
16931	Rassf1: Ras association (RalGDS/AF-6) domain family 1	551.69	251.68	515.62	171.98	181.71	219.6	213.6	114.1	183.5	2.432506201	0.024767802	0.05414861	23.0186

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
6797	Rassf5: Ras association (RalGDS/AF-6) domain family 5	720.77	665.14	560.86	336.47	286.71	262.6	254.6	257.9	267.99	2.336764273	0.022099448	0.05517495	22.0115
19264	Rbm7: RNA binding motif protein 7	1311.84	1104.7	1399.09	852.61	1194.9	761.2	564.4	793.3	454.85	1.651371739	0.044444444	0.08668921	12.7196
30077	Rbpsuh: Recombining binding protein suppressor of hairless (Drosophila)	229.78	181.62	164.15	47.11	63.14	54.63	33.26	159.1	37.35	2.917131272	0.047963801	0.09049472	12.1646
17285	Rdh11: retinol dehydrogenase 11	266.28	352.96	301.12	131.17	93.27	93.1	130.5	206.4	92.85	2.463127752	0.039370079	0.07937008	13.9553
3831	Rela: v-rel reticuloendotheliosis viral oncogene homolog A (avian)	1165.05	845.26	817.64	573.79	604.68	707	323.3	379.4	435.95	1.870281638	0.047038328	0.09183798	11.8894
19286	Rexo2: REX2, RNA exonuclease 2 homolog (S. cerevisiae)	496.13	531.04	541.91	329.27	450.19	238	210.9	214.8	230.19	1.875420574	0.048006509	0.09429889	11.4137
3543	Rgs2: regulator of G-protein signaling 2	982.72	955.41	1090.21	552.53	601.28	456.3	215.4	392.9	290.79	2.413808495	0.018181818	0.04524242	27.6143
3542	Rgs2: regulator of G-protein signaling 2	755.61	753.62	945.56	289.77	380.15	277.6	309.3	389.5	287.36	2.539035188	0.022140221	0.05134071	24.8558
38623	Rgs2: regulator of G-protein signaling 2	439.17	401.76	570.74	252.29	232.03	224.9	47.94	265.8	134.38	2.439549995	0.035714286	0.07222222	15.7414
40641	Rnd1: Rho family GTPase 1	341.36	140.63	50.42	22.78	40.06	50.85	49.94	22.23	23.68	5.081702778	0.031862745	0.06855664	16.9255
996	Rnd3: Rho family GTPase 3	236.49	269.23	240.29	57.64	90.84	104.7	34.95	86.4	66.49	3.383342026	0.027196653	0.06002789	19.3316
995	Rnd3: Rho family GTPase 3	346.48	273.15	396.01	104.43	242.51	112.8	134.1	41.05	157.66	2.563194024	0.031897927	0.06903775	16.7571
44629	Rnf185: Ring finger protein 185	3391.02	3265.6	3308.22	2163.5	2326.9	1837	1543	1821	2169	1.680373212	0.022222222	0.05532407	22.0213
19135	Rnf185: ring finger protein 185	3100.72	2881.4	3164.78	1942	2217.8	1890	2233	2234	2296.7	1.427717853	0.047833935	0.09054753	12.1523
19484	Sc5d: sterol-C5-desaturase (fungal ERG3, delta-5-desaturase) homolog (S. cerevisiae)	156.19	125.87	470.87	12.69	92.23	44.12	37.46	10.13	75.32	5.53726788	0.026768642	0.06290631	18.417
21214	Scarb2: scavenger receptor class B, member 2	412.98	136.41	11.58	17.91	8.27	79.58	23.24	12.15	32.61	6.456837017	0.048995984	0.09481928	11.3271

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
14524	Sdhc: succinate dehydrogenase complex, subunit C, integral membrane protein	655.46	629.8	569.58	133.4	248.7	128	367.2	95.09	439.58	2.627289338	0.023715415	0.05048748	25.7165
16404	Sh3bp2: SH3-domain binding protein 2	939.44	869.33	963.81	543.92	577.31	563.7	456.3	460.2	392.82	1.851886733	0.036885246	0.0731102	15.5576
11436	Six1: sine oculis-related homeobox 1 homolog (Drosophila)	890.55	896.49	591.29	257.89	404.29	126.8	116.8	139.6	145.28	3.995178901	0	0.02714286	66.8915
16644	Slc20a1: solute carrier family 20, member 1	755.61	816.13	1040.94	591.52	587.71	537.5	338.8	299.8	249.93	2.00568847	0.036745407	0.07408136	15.2304
5126	Slc25a15: solute carrier family 25 (mitochondrial carrier ornithine transporter), member 15	2097.82	2285.9	2519.54	1552.6	1649.2	1444	801.2	1031	1205.9	1.796727072	0.024657534	0.05515982	21.9246
42635	Slc7a14: Solute carrier family 7 (cationic amino acid transporter, y+ system), member 14	850.08	861.15	971.47	367.62	635.72	604.9	171.7	384	263.26	2.210530653	0.024213075	0.05830508	20.4881
15341	Slc9a3r1: solute carrier family 9 (sodium/hydrogen exchanger), isoform 3 regulator 1	1945.59	1874.9	1042.14	1213.4	1279.4	924.9	732.2	1247	905.41	1.543013175	0.04248366	0.08125999	13.6676
15340	Slc9a3r1: solute carrier family 9 (sodium/hydrogen exchanger), isoform 3 regulator 1	1511.18	1791	882.39	908.38	1070.8	773.7	603.5	1141	742.49	1.597206796	0.04744186	0.08977984	12.3407
29662	Slc4a1: solute carrier organic anion transporter family, member 4a1	1518.9	1099.7	758.84	781.47	590.77	472.9	410.2	560.3	482.97	2.047771782	0.023752969	0.05806809	20.3739

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
22832	Smurf1 /// LOC640390: SMAD specific E3 ubiquitin protein ligase 1 /// similar to Smad ubiquitination regulatory factor 1 (Ubiquitin- protein ligase SMURF1) (Smad-specific E3 ubiquitin ligase 1) (hSMURF1)	353.27	405.72	431.46	185.4	183.4	173.3	130.3	182.9	293.86	2.071951336	0.04549763	0.08836651	12.5257
7515	Snap29: synaptosomal- associated protein	1533.29	1209.3	1300.59	1097.4	921.79	856.9	676.1	678.9	687.34	1.644127897	0.047818792	0.09289989	11.6499
15156	Snord22: small nucleolar RNA, C/D box 22	244.55	88.91	292.33	138.36	78.77	71.21	96.16	68.67	49.02	2.492243971	0.041755889	0.08124197	13.5908
10772	Snrbp2: U2 small nuclear ribonucleoprotein B	1280	1540.4	1365.87	675.42	748.01	668.4	399.1	607.5	397.8	2.394785166	0.013513514	0.03754505	33.014
20449	Snrbp2: U2 small nuclear ribonucleoprotein B	1101.45	1113.1	1328.87	444.18	372.6	458.6	607.3	424.6	345.21	2.67191229	0.016260163	0.03837398	35.0616
1262	Sox2: SRY-box containing gene 2	258.38	397.22	227.54	100.5	112.85	106.8	94.1	81.95	80.38	3.06347995	0.032307692	0.07045128	16.4379
7733	Srd5a2l: steroid 5 alpha- reductase 2-like	951.51	803.16	917.18	539.46	538.83	573.3	345.4	553.4	479.19	1.763841853	0.042576419	0.08121543	13.679
37863	Ssr3: signal sequence receptor, gamma	878.88	551.79	1339.32	496.85	533.72	503.9	432	471.5	368.38	1.974066235	0.036986301	0.07271233	15.6044
17274	St3gal5: ST3 beta-galactoside alpha-2,3-sialyltransferase 5	585.5	294.61	591.89	101.96	252.48	156	129.7	214.2	153.36	2.921765366	0.015037594	0.03827068	34.2305
10512	Stat6: signal transducer and activator of transcription 6	280.93	369.43	537.91	177.71	247.51	140.3	74.96	158.2	163.26	2.470544207	0.038560411	0.07460154	15.0996
5687	Surf5: surfeit gene 5	808.83	625.57	901.86	425.58	600.45	367.5	196.1	219.1	363.82	2.150727493	0.024630542	0.05834154	20.6282
16783	Taf13: TAF13 RNA polymerase II, TATA box binding protein (TBP)- associated factor	2786.83	2273	2785.38	1860.7	2037.4	1751	1700	1367	1418.5	1.548259965	0.030769231	0.06631909	17.421
14515	Taok3: TAO kinase 3	840.36	751.13	664	644.14	406.28	520.9	288.5	201	341.2	1.877923992	0.042726348	0.08343845	13.1523

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
21642	Tax1bp3 /// Rpl13: Tax1 (human T-cell leukemia virus type I) binding protein 3 /// ribosomal protein L13	871.73	903.94	1048.44	382.12	394.43	282.3	507.1	145.2	438.71	2.627237926	0.016666667	0.04101852	30.5599
502	Taz: tafazzin	510.91	478.79	738.95	285.8	194.27	193.5	116.3	261.3	97.64	3.009750152	0.023529412	0.05086275	25.5673
6617	Tcl1: T-cell lymphoma breakpoint 1	4221.74	3745.3	4210.23	2520.7	3350.8	3381	2300	2780	3125.2	1.394976158	0.044399596	0.08380424	13.0961
19537	Thecd1: thioesterase domain containing 1	749.49	760.8	818.91	422.7	601.57	400.7	231.5	336.5	417.73	1.932376769	0.036599764	0.07688705	14.4348
20152	Thrap3: thyroid hormone receptor associated protein 3	308.4	324.33	307.94	159.33	140.04	211.4	90.03	100.2	151.4	2.207264707	0.048242028	0.09426547	11.4399
16803	Tle6: transducin-like enhancer of split 6, homolog of Drosophila E(spl)	1270.14	833.99	1229.7	374.24	654.97	691.3	407.2	562.4	495.75	2.092904854	0.017241379	0.04622126	27.0809
15195	Tm2d2: TM2 domain containing 2	934.68	572.01	641.41	527.63	320.67	388.1	290.4	286.3	252.38	2.079960494	0.041896362	0.08087468	13.7569
21204	Tmed10 /// LOC634748: transmembrane emp24-like trafficking protein 10 (yeast) /// similar to transmembrane trafficking protein	371.69	689.99	932.01	227.36	344.45	312	356.4	335.3	331.6	2.090851887	0.029307282	0.06547661	17.7043
15880	Tmed10 /// LOC634748: transmembrane emp24-like trafficking protein 10 (yeast) /// similar to transmembrane trafficking protein	293.24	526.59	744.83	183.86	260.8	259	306.9	282.8	262.97	2.01075635	0.042769857	0.08350984	13.1531
15864	Tmed10 /// LOC634748: transmembrane emp24-like trafficking protein 10 (yeast) /// similar to transmembrane trafficking protein	325.23	522.46	807.94	214.86	292.41	263.9	328.4	287.9	276.07	1.990478194	0.048141892	0.09307995	11.6733
21746	Tmem109: transmembrane protein 109	658.76	639.21	690.17	367.58	528.21	245.5	273.8	272.9	273.33	2.027348928	0.036	0.07345778	15.3724

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
14887	Tmem109: transmembrane protein 109	827.27	787.55	910.96	487.81	701.24	347.9	350.2	365	381.93	1.917784713	0.037974684	0.07493671	14.9506
8336	Tmem38a: transmembrane protein 38a	965.95	199.02	508.25	119.7	369.44	96.15	251.5	101.5	150.48	3.073850902	0.024561404	0.05080702	24.4359
10674	Tor1a: torsin family 1, member A (torsin A)	1760.16	1607.9	1566.85	1294.5	1207	1120	851.9	880.3	816.44	1.599693348	0.047144152	0.09024781	12.187
7470	Tpbp: trophoblast glycoprotein	135.67	173.87	65.62	38.44	9.01	28.02	6.79	31.8	103.96	3.441519127	0.047454702	0.0919931	11.8366
20812	Tpi1: triosephosphate isomerase 1	723.43	510.71	278.29	243.4	202.66	199.6	498.9	79.81	92.94	2.29629236	0.03686088	0.07673801	14.4886
22085	Transcribed locus	244.72	168.35	553.92	235.2	92.48	72.5	7.42	50.24	1	4.21493331	0.035410765	0.07242682	15.7775
43618	Transcribed locus	1331.08	1132.8	759.97	700.03	776.17	653.8	600.2	448.6	142.5	1.941371472	0.0390625	0.07442274	15.175
30279	Transcribed locus	115.59	77.5	3.01	1	6.1	4.39	6.91	7.21	3.07	13.67503487	0.042162162	0.08108468	13.6466
31948	Transcribed locus	948.04	695.42	1011.76	481.07	569.52	551.4	635.4	513.4	462.89	1.652432858	0.046961326	0.08948742	12.2985
36521	Transcribed locus	163.5	288.96	275.95	153.58	132.29	129	127.8	62.86	77.98	2.131349485	0.049114332	0.094635	11.3498
14557	Transcribed locus, moderately similar to XP_574723.1 PREDICTED: similar to LRRGT00097 [Rattus norvegicus]	389.93	446.02	354.24	146.9	73.38	178.6	150.5	243.5	285.69	2.207018682	0.045135406	0.0843096	13.0436
31203	Transcribed locus, weakly similar to XP_417295.1 PREDICTED: similar to TAF3 protein [Gallus gallus]	833.28	557.28	868.7	344.25	284.45	400.2	464.4	244.1	342.7	2.172271392	0.022727273	0.05269481	23.5482
16554	Trfp: Trf (TATA binding protein-related factor)-proximal protein homolog (Drosophila)	1346.13	1510.8	1419.34	707.69	420.35	502.5	668.7	1297	686.12	1.997305029	0.023323615	0.0545967	22.442
11566	Trp11: thyroid hormone receptor interactor 11	259.32	592.74	496.79	341.11	199.04	280	193.4	155.1	127.48	2.081494398	0.037878788	0.07494108	14.9376
39305	Troap: trophinin associated protein	208.07	262.5	390.33	82.47	98.83	71.69	101.3	44.16	107.74	3.401287978	0.025114155	0.0585312	20.0849
8651	Trpc2: transient receptor potential cation channel, subfamily C, member 2	1159.04	683.82	924.49	504.31	574.63	394.4	398.4	456.4	182.58	2.204436195	0.02359882	0.05485742	22.5014
17247	Ttk: Ttk protein kinase	442.51	328.09	595.94	156.99	233.89	156.6	291.1	218.8	142.18	2.278383087	0.037859008	0.07429939	15.191

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
20704	Tubb2b: tubulin, beta 2b	2089.18	1881.4	2187.75	1223.2	1598.2	1370	901.7	1304	685.75	1.739163956	0.02832244	0.06042847	19.5561
1904	Ube2a: ubiquitin-conjugating enzyme E2A, RAD6 homolog (S. cerevisiae)	668.24	817.55	665.53	322.75	424.91	331.9	208	371.1	379.5	2.110968831	0.036401099	0.07271978	15.6237
6873	Ube2i /// LOC546265 /// LOC669417: ubiquitin-conjugating enzyme E2I /// similar to Chain A, Human Ubiquitin-Conjugating Enzyme Ubc9 /// similar to Ubiquitin-conjugating enzyme E2 I (Ubiquitin-protein ligase I) (Ubiquitin carrier protein I) (SUMO-1-protein ligase) (SUMO-1-conjugating enzyme) (Ubiquitin-conjugating enzyme UbcE2A)	881.06	714.36	770.75	667.98	572.39	299.1	178.7	412.2	456.5	1.829369083	0.049193548	0.09460215	11.3561
34066	Ube2w: ubiquitin-conjugating enzyme E2W (putative)	257.73	223.92	304.28	83.65	208.87	56.27	64.14	66.41	62.1	2.903110225	0.032258065	0.07035842	16.4371
30305	Umodl1: uromodulin-like 1	918.71	1173.2	1195.72	468.99	479.34	394.6	140.4	315.4	169.66	3.340415155	0	0.02952381	59.0622
27170	Unc5a: unc-5 homolog A (C. elegans)	251.93	17.43	149.64	24.49	31.01	22.93	71.25	19.8	14.94	4.543975708	0.036948749	0.07631307	14.5235
18877	Vil2: villin 2	241.97	205.21	109.2	76.45	75.17	88.63	56.94	39.13	59.75	2.809503371	0.038596491	0.07755945	14.307
36640	Wdr27: WD repeat domain 27	258.28	184.34	219.19	117.86	87.15	48.94	74.51	75.45	48.47	2.925903002	0.0375	0.07488333	14.8953
40438	Wdr40b: WD repeat domain 40B	412.95	601.38	440.4	258.01	180.34	165.1	177.8	333.6	44.28	2.509864477	0.034023669	0.07133136	16.1014
20216	Wdr82: WD repeat domain containing 82	930.14	666.97	795.89	478.36	548.79	548.8	439.9	334.8	377.74	1.754141621	0.047417443	0.09220999	11.7317
44710	Wfdc3: WAP four-disulfide core domain 3	840.41	352.61	513.04	317.4	228.96	204.5	83.53	153.4	181.23	2.918761708	0.024647887	0.0509507	24.4457

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
4516	Xlr3a /// Xlr3b /// MGC76689: X-linked lymphocyte-regulated 3A /// X- linked lymphocyte-regulated 3B /// hypothetical protein LOC574437	360.83	588.71	622.92	230.28	410.11	221.9	206	332.5	245.03	1.910945836	0.036817102	0.07669438	14.4846
6776	Xmr /// LOC546272 /// LOC619991 /// LOC664810 /// LOC664861 /// LOC664877 /// LOC664890 /// LOC664906 /// LOC664923 /// LOC664944 /// LOC664989: Xmr protein /// similar to Xlr-related, meiosis regulated /// similar to Xlr-related, meiosis regulated /// similar to Synaptonemal complex protein 3 (SCP-3) /// similar to Xlr-related, meiosis regulated /// similar to Xlr- related, meiosis regulated /// similar to Xlr-related, meiosis regulated /// similar to Xlr- related, meiosis regulated /// similar to Xlr-related, meiosis regulated /// similar to Xlr- related, meiosis regulated /// regulated	402.45	431.9	419.31	132.31	243.77	319	93.22	113	120.66	2.453418398	0.029411765	0.06629181	17.4967

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
6777	Xmr /// LOC546272 /// LOC619991 /// LOC664810 /// LOC664861 /// LOC664877 /// LOC664890 /// LOC664906 /// LOC664923 /// LOC664944 /// LOC664989: Xmr protein /// similar to Xlr-related, meiosis regulated /// similar to Xlr-related, meiosis regulated /// similar to Synaptonemal complex protein 3 (SCP-3) /// similar to Xlr-related, meiosis regulated /// similar to Xlr- related, meiosis regulated /// similar to Xlr-related, meiosis regulated /// similar to Xlr- related, meiosis regulated /// similar to Xlr-related, meiosis regulated /// similar to Xlr- related, meiosis regulated /// similar to Xlr-related, meiosis regulated	416.14	408.25	408.94	135.94	270.28	290.5	105.7	134.5	133.3	2.304945055	0.038860104	0.0744905	15.1475
15133	Ywhaq: tyrosine 3- monooxygenase/tryptophan 5- monooxygenase activation protein, theta polypeptide	823.51	1230.4	1166.2	689.41	787.92	682.6	668.7	504.4	743.91	1.579678045	0.048561151	0.09080635	12.127
14866	Ywhaz: tyrosine 3- monooxygenase/tryptophan 5- monooxygenase activation protein, zeta polypeptide	1349.62	789.35	427.74	385.13	670.02	394.3	284.5	439.5	466.58	1.944403621	0.048956661	0.09480203	11.319
27398	Zarl: zygote arrest 1	308.67	302.35	296.84	96.91	62.32	61.6	36.14	73.34	105	4.17109646	0.016853933	0.04059925	30.8299
8460	Zfp219: zinc finger protein 219	362.85	474.33	421.09	139.56	154.27	248.9	222.6	276.2	128.61	2.150613169	0.041622199	0.08153682	13.5586
15729	Zfp3612: zinc finger protein 36, C3H type-like 2	1493.92	1568.6	1661.69	1083.2	1187.2	1122	833.5	1041	686.1	1.587178186	0.047877145	0.09053899	12.154

Table 8A

Row	Gene	Oct4.1	Oct4.2	Oct4.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
208791	Zfp422-rs1: zinc finger protein 422, related sequence 1	807.14	637.1	963.93	639.73	531.32	551.3	241.2	450.1	320.33	1.761665271	0.049153908	0.09463336	11.3527
24731	Zfp444: zinc finger protein 444	153.25	73.77	591	9.86	2.23	35.58	45.91	75.68	121.03	5.63588136	0.041202673	0.08044915	13.8348
7994	Zfp503: zinc finger protein 503	499.68	1344.6	631.84	248.83	28.76	97.85	46.17	549	78.64	4.71964318	0.02	0.036	49.7065
7995	Zfp503: zinc finger protein 503	201.34	138.04	36.43	20	36.65	32.65	17.34	48.18	62.9	3.452232225	0.04757548	0.08988411	12.2507
44691	Zfp68: zinc finger protein 68	836.97	770.67	353.93	367.82	511.34	379.8	260.4	345.8	315.87	1.798780376	0.037125749	0.07644311	14.5373
8911	Zfp71-rs1: zinc finger protein 71, related sequence 1	118.31	304.51	288.84	100.79	115.62	155.5	55.72	138.6	55.13	2.290505311	0.043478261	0.08252033	13.4658
17202	Zfp90: zinc finger protein 90	148.51	168.8	284.96	41.22	44.14	22.58	38.95	17.6	56.83	5.442526658	0.024539877	0.05421268	22.8885
1832	Zmat2: zinc finger, matrin type 2	271.97	245.47	570.3	68.8	63.18	133.6	110.3	99.83	40.13	4.218008376	0.023972603	0.05133562	24.0851

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
52	transferrin receptor	624.69	241.34	260.79	1658.5	1221.5	1475.4	2092.35	1394.63	2043.46	0.227967388	0	0.00015951
154	eukaryotic translation initiation factor 3, subunit 8	1574.05	898.06	884.18	3064.3	2464.01	3091	3612.17	2968.05	3465.95	0.359626325	0	0.00015921
785	transmembrane emp24 protein transport domain containing 6	158.97	64.18	182.56	508.83	418.62	577.06	745.64	731.85	729.56	0.218619664	0	0.00022179
1785	cathepsin B	725.96	401.03	413.68	1480.9	1297.15	1599.4	2070.39	1870.47	2628.6	0.28148163	0	0.00024093
4369	ATP-binding cassette, sub-family F (GCN20), member 1	415.1	244.32	320.89	937.07	895.73	1089.7	1088.85	1163.3	2060.89	0.270972664	0	0.0003167
4798	junction-mediating and regulatory protein	143.62	100.64	83.34	547.49	231.34	463.45	532.88	603.44	674.49	0.214602255	0	0.00036372
4799	junction-mediating and regulatory protein	290.52	233.15	287.38	1194	1060.93	1111.5	1676.86	1290.77	1588.17	0.204752955	0	6.13E-006
5181	acylphosphatase 1, erythrocyte (common) type	16.91	108.49	111.19	532.02	571.18	559.42	528.01	477.38	466.65	0.15095098	0	0.00038783
7126	transferrin receptor	689.39	491.5	251.38	1428.2	1227.04	1221.1	2091.98	1833.64	2227.85	0.285602903	0	0.00068689
10805	RIKEN cDNA 9130011J15 gene	285.53	256.11	271.55	782.98	735.67	749.47	1214.59	1679.66	1121.66	0.258811622	0	0.0002899
10887	phosphatidylserine synthase 2	708.39	150.51	190.22	1424.2	1092.46	1221.8	1035.72	1127.98	1999.15	0.265555293	0	0.00127543
12367	Kruppel-like factor 9	42.96	50.16	156.96	413.31	485.3	353.82	523.55	415.15	759.55	0.16950669	0	0.00019734
13878	general transcription factor IIIC, polypeptide 4	27.87	16.57	35.71	474.48	220.1	217.44	516.42	384.03	566.54	0.067380969	0	2.35E-005
13972	eukaryotic translation initiation factor 3, subunit 6	203.48	187.05	266.56	1013.4	919.72	998.66	1287.5	1098.19	1049.17	0.206416571	0	1.86E-006
14002	phosphatidylserine synthase 2	602.84	152.26	203.21	1152.7	870.06	1008.7	1052.55	1114.61	1781.74	0.274571264	0	0.00102264
14683	diacylglycerol kinase, epsilon	190.03	434.95	115.62	1081.8	777.24	733.1	1097.92	905.2	997.74	0.264831986	0	0.00044972
15504	torsin family 2, member A	231.7	99.13	132.44	803.51	625.23	692.06	515.53	1579.92	860.9	0.182492146	0	0.00031663
15713	acylphosphatase 1, erythrocyte (common) type	9.09	70.55	99.27	431.09	432.79	469.83	452.15	386.11	368.38	0.140854607	0	0.00068273
15791	eukaryotic translation initiation factor 3, subunit 6	117.88	107.21	148.87	614.35	581.81	595.12	849.71	634.84	809.58	0.183070977	0	2.91E-006
18122	acylphosphatase 1, erythrocyte (common) type	24.1	104.37	109.38	657.52	691.59	706.49	594.96	581.28	463.27	0.12873771	0	0.00010853
18375	solute carrier family 19 (sodium/hydrogen exchanger), member 3	227.44	127.66	156.95	731.75	478.33	711.17	1052.97	795.48	1291.81	0.202330925	0	0.0001809
19172	transmembrane protein 111	330.63	240.26	230.31	1271.5	949.26	808.68	1152.76	998.66	1763.99	0.230733122	0	7.44E-005

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
21863	Kruppel-like factor 9	126.03	324.94	264.78	1413.6	1073.94	1024.2	1693.05	992.37	1695.4	0.181373344	0	8.56E-005
21873	eukaryotic translation initiation factor 3, subunit 8	688.45	338.22	292.12	1367.9	940	1082	1811.77	1859.33	2228.32	0.283934988	0	0.00112513
22149	phosphatidylserine synthase 2	1200.27	227.6	292.55	2416.8	1921.95	2063.5	1639.91	1905.6	3229.72	0.261113788	0	0.00137219
22419	eukaryotic translation initiation factor 3, subunit 6	253.42	148.89	203.42	867.23	627.04	653.69	1252.51	944.86	908.52	0.23058519	0	7.39E-005
27213	DNA segment, Chr 19, Brigham & Women's Genetics 1357 expressed	566.7	267.24	1036.42	1513.8	1212.97	2166.7	2823.83	2513.75	3203.16	0.278448166	0	0.00268403
28204	galactosidase, beta 1	426.39	169.6	51.27	1713.4	1707.87	1919	2551.5	2521.75	2747.95	0.098357318	0	0.00014849
28632	ankyrin repeat domain 38	29.64	6.72	16.18	346.3	291.12	247.57	166	148.99	316.27	0.069302556	0	4.02E-005
29867	RIKEN cDNA 4933433K01 gene	145.78	41.38	31.11	448.82	321.98	385.46	480.01	555.13	902.54	0.141095173	0	0.00031856
33396	Transcribed locus	352.3	82.9	105.87	670.9	481.47	592.69	1401.57	1113.63	1247.74	0.196466957	0	0.00143556
41050	RIKEN cDNA 4930427A07 gene	225.06	114.11	217.2	462.21	493.17	756.34	822.19	1212.27	977.74	0.23555437	0	0.00068706
29064	ubiquitin specific peptidase 22	470.08	134.25	91.36	877.44	673.24	719.13	843.91	842.44	1309.02	0.264260671	0.01064	0.00127129
21759	eukaryotic translation initiation factor 3, subunit 8	873.58	406.65	383.89	1537.6	1184.08	1444.6	1689.92	1790.34	2140.93	0.340049716	0.01111	0.00060111
22191	ribosomal protein L24 /// similar to ribosomal protein L24 /// similar to ribosomal protein L24	1172.49	672.1	952.29	1781.4	1430.15	1946.1	2400.65	2576.5	2545.94	0.441122173	0.01124	0.0018963
2935	sirtuin 1 ((silent mating type information regulation 2, homolog) 1 (S. cerevisiae)	159.78	122.53	182.71	551.66	431.36	568.18	612.19	521.34	692.06	0.275421332	0.01136	2.03E-005
4370	ATP-binding cassette, sub-family F (GCN20), member 1	427.95	279.01	441.26	1100.4	946.89	1266	1325.41	1128.42	2317.25	0.284061348	0.01136	0.00045761
16122	Transcribed locus	93.98	30.44	15.4	215.29	193.76	209.2	360.32	327.29	349.09	0.168971872	0.01149	0.00047997
153	eukaryotic translation initiation factor 3, subunit 8	475.14	206.19	253.03	1108.7	888.34	784.61	1267.74	1026.05	1313.35	0.292501663	0.01163	0.00024627
8022	ATP-binding cassette, sub-family F (GCN20), member 2	152.19	156.91	162.97	363.57	305.96	451.65	884.29	887.81	683.59	0.263957035	0.01163	0.00200161
28457	solute carrier family 30 (zinc transporter), member 1	233.42	27.36	61.68	315.4	343.75	619.79	393.91	491.5	503.8	0.241710549	0.01176	0.00194384
21824	thioredoxin-like 2	294.52	175.68	266.05	909.58	693.73	804.9	643.13	850.93	762.17	0.315686342	0.0119	4.42E-005

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	tpvals
7389	inositol polyphosphate-5-phosphatase E	25.77	21.31	39.99	241.41	177.04	187.24	342.97	181.11	290.47	0.122613079	0.01205	1.28E-005
16808	cathepsin B	130.23	101.77	67.38	288.01	181.13	261.76	690.28	580.37	717.78	0.22018659	0.0122	0.00427226
39214	1-acylglycerol-3-phosphate O-acyltransferase 5 (lysophosphatidic acid acyltransferase, epsilon)	499.54	525.47	363.14	1210.6	1108.45	1066.1	1281.2	1561.27	1458.73	0.361200628	0.0125	7.86E-005
33996	abhydrolase domain containing 6	156.99	100	43.52	553.9	326.89	296.3	593.6	167.18	618.84	0.235075546	0.01266	0.00413915
53	transferrin receptor	409.32	206.62	95.99	595.48	591.14	503.44	983.73	948.43	1145.38	0.298653411	0.01316	0.00301392
9791	PQ loop repeat containing 2	13.76	7.56	6.29	292.48	156.56	55.45	60.34	219.02	172.47	0.057742178	0.01333	0.00024888
7125	transferrin receptor	361.01	329.39	377.07	974.9	842.63	895.37	1430.85	1069.76	1417.72	0.321952338	0.01351	0.00011751
14950	eukaryotic translation initiation factor 2C, 5	44.09	7.7	7.32	207.56	115.15	151.62	314.47	175.57	291.88	0.094105473	0.0137	0.00022399
18250	apoptotic peptidase activating factor 1	36.46	51.69	50.02	277.53	197.48	388.51	297.58	228.07	287.72	0.164793159	0.01389	8.99E-006
10657	jumonji, AT rich interactive domain 1C (Rbp2 like)	55.24	122.77	112.59	314.35	218.11	459.64	670.73	493.09	411.48	0.226376879	0.01408	0.00075321
43911	gb:BM122336 /DB_XREF=gi:17106104 /DB_XREF=L0508F03-3 /CLONE=L0508F03 /FEA=EST /CNT=3 /TID=Mm.218397.1 /TIER=ConsEnd /STK=2 /UG=Mm.218397 /UG_TITLE=ESTs	75.29	12.69	9.34	166.18	32.25	202.06	433.2	516.27	301.93	0.117828669	0.01449	0.00986852
18703	transcription elongation factor B (SIII), polypeptide 3	231.44	240.04	198.89	638.35	629.15	741.99	859.61	921.16	783.83	0.293116226	0.01471	1.39E-005
16204	protective protein for beta-galactosidase	474.09	446.21	267.96	1117.8	569.65	1350.3	1365.1	642.54	1356.92	0.371197851	0.01493	0.00629467
15395	Vacuolar protein sorting 54 (yeast)	714.94	245.49	144.2	1535.2	886.77	1048.1	1134.23	921.8	1216.19	0.327673024	0.01515	0.00227705
4980	sphingosine-1-phosphate phosphatase 1	262.03	173.69	51.88	1215.3	589.36	577.14	509.63	536.21	763.86	0.232661338	0.01538	0.0016041
54	transferrin receptor	434.35	354.1	332.55	1138.5	818.28	858.07	1238.66	1119.89	1324.05	0.345056322	0.01563	0.00010839
27902	potassium inwardly-rectifying channel, subfamily K, member 6	44.82	139.14	63.27	323.91	232.15	319.5	516.23	473.27	565.17	0.203462224	0.01587	0.00046105

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
39802	neuropilin (NRP) and tolloid (TLL)-like 2	87	64.53	146.38	522.51	475.14	750.45	427.95	275.96	347.24	0.212849871	0.01639	0.00032483
384	ribosomal protein S15	256.87	118.98	155.05	330.72	201.23	1135.8	727.25	514.63	1106.92	0.264353593	0.01667	0.01955601
6466	go:0010514 /DB_XREF=gi:14010854 /GEN=Usmg3 /FEA=FLmRNA /CNT=2 /TID=Mm.218588.1 /TIER=FL /STK=1 /UG=Mm.218588 /LL=83678 /DEF=Mus musculus upregulated during skeletal muscle growth 3 (Usmg3), mRNA. /PROD=upregulated	228.12	205.44	238.32	906	787.64	785.29	730.81	656.42	846.69	0.285126834	0.01667	3.78E-006
34766	Vacuolar protein sorting 33A (yeast)	315.2	115.51	221.53	607.77	456.63	589.67	807.65	809.12	987.11	0.306363391	0.01667	0.00105604
38885	RNA, U transporter 1	515.79	234.67	214.8	722.54	562.88	810.9	985.1	1105.78	1283.74	0.352868063	0.01681	0.00227145
8821	peptidase (mitochondrial processing) alpha	73.11	56.51	41.28	381.32	318.1	307.73	329.1	251.1	518.94	0.16227585	0.01695	1.65E-005
10448	RIKEN cDNA 2610028H07 gene	638.95	517.06	757.25	1387	1282.58	1540.6	1558.37	2078.88	1484.02	0.410069668	0.01724	0.00022595
14001	phosphatidylserine synthase 2	252.19	121.81	137.37	847.59	360.31	528.7	764.33	553.32	801.99	0.265216895	0.01724	0.0006063
21527	DEAD (Asp-Glu-Ala-Asp) box polypeptide 49	173.52	35.84	59.84	496.4	167.03	182.83	391.02	405.54	415.71	0.26154586	0.01739	0.00468851
29344	Adult male cecum cDNA, RIKEN full-length enriched library, clone:9130409M07 product: hypothetical protein, full insert sequence	90.7	19.1	227.83	613.93	432.25	327.99	562.83	671.26	386.9	0.225450393	0.01754	0.00292411
13579	RIKEN cDNA 9430010O03 gene	1276.81	930.39	852.58	2128.7	1524.54	2270.8	2042.66	2601.49	3336.82	0.440098756	0.0177	0.00221166
8699	homeodomain interacting protein kinase 1	703.08	358.75	437.43	1281.1	954.65	1185.9	1575.69	1648.73	1861.89	0.352435769	0.01786	0.00072822
16053	expressed sequence C77815	226.98	68.5	120.21	355.27	288.1	364.69	689.78	629.82	713.08	0.273413709	0.01786	0.00247561
33270	gene model 104, (NCBI)	167.88	96.62	189.61	623.96	423.95	354.11	625.11	416.17	774.69	0.282232077	0.01802	0.00055679
3951	solute carrier family 25, member 36	196.31	138.07	287.87	552.92	426.31	612.23	716.16	716.8	838.71	0.322148103	0.01818	0.00045152
8247	melanocyte proliferating gene 1	38.3	8.3	8.89	124.7	125.7	87.63	225.51	241.69	255.09	0.104666516	0.01835	0.00028537

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
24977	origin recognition complex, subunit 4-like (S. cerevisiae)	64.74	3.28	40.64	285.69	259.43	282.71	113.94	217.49	143.37	0.166831717	0.01852	0.00382844
28887	deficient 5, cell division cycle 46 (S. cerevisiae)	521.19	337.32	431.77	1405.9	1199.34	1296.8	1001.57	1386.43	1286.19	0.34061178	0.01852	3.87E-005
44778	gb:BB473929 /DB_XREF=gi:16439785 /DB_XREF=BB473929 /CLONE=D330003M06 /FEA=EST /CNT=5 /TID=Mm.214471.1 /TIER=ConsEnd /STK=3 /UG=Mm.214471 /UG_TITLE=ESTs, Weakly similar to ENV1_HUMAN RETROVIRUS-RELATED ENV POLYPROTEIN (H.sapiens)	186.55	85.65	81.04	314.37	247.68	303.97	629.46	581.85	788.79	0.246493517	0.01887	0.00263775
9574	solute carrier family 1 (neuronal/epithelial high affinity glutamate transporter, system Xag), member 1	86.76	56.92	100.42	341.84	267.96	357.7	469.01	318.11	475.25	0.21893653	0.01905	4.04E-005
8320	DEAD (Asp-Glu-Ala-Asp) box polypeptide 27	112.56	127.64	180.35	447.67	410.04	494.24	453.03	489.33	871.51	0.265681561	0.01908	0.00016822
5137	nuclear respiratory factor 1	96.06	22.21	58.59	206.74	207.72	216.54	387.53	439.3	340.7	0.196671726	0.01923	0.00072954
12240	kelch-like 21 (Drosophila)	945.8	323.32	551.74	1286.9	1071.87	1407.3	2344.72	2073.51	1947.4	0.359439618	0.01923	0.00324608
1375	signaling intermediate in Toll pathway-evolutionarily conserved	36.01	35.11	41.82	287.35	199.74	197.73	218.85	209.08	241.65	0.166774956	0.01935	7.77E-007
20342	kinesin family member 11	285.49	271.4	175.23	802.55	400.74	666.11	896.31	538.17	967.58	0.34279614	0.01938	0.00170065
32215	RIKEN cDNA 4833426J09 gene	123.19	52.54	98.29	403.04	266.83	290.65	664.64	325.8	820.34	0.197755566	0.01942	0.00102827
40316	(human)	76.89	53.93	138.9	311.28	272.58	273.85	386.44	406.16	550.1	0.245154312	0.01948	0.00031454
22448	ribosomal protein L37a	1002.33	507.96	700.13	1369.4	1086.69	1508.1	1821.16	2432.09	2770.43	0.402340412	0.01953	0.00609027
14883	similar to tripartite motif- containing 43	2079.61	1375.77	1494.31	3821.7	3156.2	3686.6	3576.25	3726.61	3888.5	0.452940224	0.01961	0.00012531
17514	dolichol-phosphate (beta-D) mannosyltransferase 1	258.66	211.61	223.09	752.95	658.07	688.27	654.69	634.23	705.73	0.338725042	0.01961	4.11E-006

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
20498	interleukin 17 receptor D	209.33	454.26	402.87	1218.7	961.8	1018.6	741.16	890.6	1180.07	0.354838495	0.01961	0.00052241
8691	YLP motif containing 1	212.15	172.37	259.8	629.55	420.34	439.07	700.52	1020	768.8	0.32391888	0.01969	0.00118415
1363	protein tyrosine phosphatase, non-receptor type 1	179.44	81.35	70.37	316.61	354.23	389.06	459.26	401.39	646.69	0.257989125	0.01974	0.00033685
4707	RIKEN cDNA 2310008H09 gene	197.94	200.34	46.69	511.26	431.89	592.74	490.78	359.97	815.86	0.277889149	0.01984	0.00233408
29582	HECT domain containing 1	631.08	441.62	450.4	1198	860.21	1101.7	1757.53	1238.84	1402.21	0.403019138	0.01987	0.00082061
12090	pyridoxal (pyridoxine, vitamin B6) kinase	799	486.45	774.66	1353.4	1207.1	1465.8	1752.21	1584.83	2835.54	0.403985912	0.02	0.00279501
19076	B-cell receptor-associated protein 31	467.79	406.25	272.93	836.98	845.26	926.32	1177.38	1170.25	1387.16	0.361629108	0.02	0.00035378
22059	DNA segment, Chr 9, ERATO Doi 720, expressed	78.25	101.45	110.24	395.68	368.76	211.62	565.85	296.89	522.68	0.24555787	0.02	0.00042704
14968	Casitas B-lineage lymphoma- like 1	59.21	39.31	73.98	227.96	165.93	243.84	387.77	297.25	353.48	0.205819011	0.02013	0.00013549
3646	RIKEN cDNA 0610007P06 gene	95.48	69.48	144.37	251.3	275.76	304.24	515.25	489.89	705.11	0.243418386	0.02016	0.00113655
7606	caseinolytic peptidase X (E.coli)	305.08	204.23	250.93	827.98	642.13	720.65	860.05	674.03	951.76	0.325125091	0.02041	4.40E-005
14623	DNA segment, Chr 19, ERATO Doi 721, expressed	1485.68	1003.23	618.92	2282.7	1674.85	1684	2958.22	2051.06	2943.84	0.457211329	0.02041	0.00546228
21740	ubiquitin specific peptidase 22	61.29	89.92	72.62	339.83	313.96	288.37	464.56	442.81	598.41	0.182872129	0.02041	3.09E-005
40844	RIKEN cDNA A530082C11 gene	103.02	148.62	109.7	459.34	222.66	476.49	456.67	365.07	541.49	0.286582174	0.02049	0.0004431
41846	DNA segment, Chr 14, Abbott 1 expressed	186.91	203.53	137.66	493.13	316.45	653.17	637.61	639.35	503.72	0.325642915	0.02055	0.00045287
12366	Kruppel-like factor 9	16.35	11.35	21.12	152.11	133.07	165.7	191.43	117.35	292.03	0.092841046	0.02066	1.05E-005
16275	ankyrin repeat domain 10	957.77	535.96	734	1791.6	1402.79	1508.8	1900.92	1549.28	1875.29	0.444274929	0.02069	0.00043489
15513	solute carrier family 44, member 2	306.38	221.45	247.39	842.17	546.42	703.47	972.75	1046.06	1193.47	0.292296497	0.02083	0.00026173
30136	DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 31	298.29	107.94	733.56	541.89	894.27	2088.9	557.59	1736.89	1228.52	0.323432306	0.02083	0.02150807
5443	phosphatidylinositol glycan, class N	16.63	32.81	16.53	182.07	166.32	169.44	257.91	95.85	192.79	0.123959488	0.02098	3.41E-005
11492	splicing factor, arginine/serine- rich 15 /// similar to splicing factor, arginine/serine-rich 15	1024.33	1160.82	1136.04	2097	1886.86	1968.6	2810.75	2721.99	2515.66	0.474425893	0.02113	0.00046687

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
512	ribosomal protein L37a	554.13	423.04	428.11	1126.8	809.26	769.11	1242.39	1292.95	1650.52	0.407856568	0.02128	0.00194373
19536	TATA box binding protein (TBP)-associated factor	311.53	245.56	144.13	955.42	630.33	651.71	788.7	630.81	1019.52	0.299891585	0.02151	0.00027298
13981	bolA-like 2 (E. coli)	1123.84	508.83	707.52	1607	1111.28	1182.5	2058.81	2007.06	2217.36	0.459584404	0.02174	0.00773939
16339	CNDP dipeptidase 2 (metallopeptidase M20 family)	29.26	96.28	61.29	542.76	229.08	165.59	150.97	419.23	414.14	0.194435338	0.02174	0.00246061
40577	RIKEN cDNA D330037H05 gene /// similar to La-related protein 4 (La ribonucleoprotein domain family member 4) /// similar to La-related protein 4 (La ribonucleoprotein domain family member 4)	2109.33	1591.5	1770.31	3282.8	2998.53	3238.5	4490.22	4321.16	4046.76	0.488976197	0.02198	0.00074647
23071	RIKEN cDNA 3300001M20 gene	252.81	164.8	211.21	708.72	562.73	499.17	858.23	902.18	429.21	0.317566612	0.02222	0.0005967
5178	RIKEN cDNA 1700021F05 gene	1666.18	665.55	1064.07	2327.8	1899.07	2094.8	2300.95	2524.08	2681.17	0.491153722	0.02339	0.00283379
9650	bone morphogenetic protein receptor, type 1A	216.85	211.07	155.24	631.62	531	550.21	566.68	440.17	655.89	0.345517942	0.02347	4.52E-005
20927	ribosomal protein S15a	1300.6	959.53	1013.49	2281.4	1908	2020.4	1644.46	2363.4	2951.07	0.497181577	0.02358	0.00167999
9017	L-2-hydroxyglutarate dehydrogenase	651.86	266.18	527.12	1034.2	590.33	965.17	1621.21	1345.63	1417.59	0.414435677	0.02367	0.00986099
19095	ribosomal protein S25	1492.69	819.24	1298.17	1951	1634.91	2069.3	2782.56	3268.32	3608.72	0.471453683	0.02381	0.00935608
23146	RIKEN cDNA 4933417E01 gene	1387.41	981.03	1339	2342.5	1725.81	1965.4	2559.65	2606.75	2682.38	0.534117054	0.02392	0.00225043
14122	ribosomal protein L37 /// similar to ribosomal protein L37	2088.46	1026.24	1539.86	2325.4	1980.64	2858.3	3213.87	3926.6	5118.59	0.47927296	0.02395	0.01585526
19182	solute carrier family 12, member 2	967.39	592.93	609.84	1657.9	1103.01	1407.2	2393.42	1532.65	1828.34	0.437422461	0.0241	0.00251573
9201	proline, glutamic acid and leucine rich protein 1	510.07	270.17	230.83	665.9	602.67	521.24	1107.52	1067.04	1083.99	0.400553843	0.02415	0.00573436
7982	RIKEN cDNA 2610012O22 gene	269.15	194.81	225.6	623.68	421.41	548.5	719.07	776.11	829.08	0.352009393	0.02424	0.0003918
6045	Son of sevenless homolog 1 (Drosophila)	324.06	248.6	173.51	684.56	368.73	779.87	937.33	493.55	698.66	0.376596765	0.02439	0.00317609

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
8238	RIKEN cDNA 2900073H19 gene	238.99	108.15	83.52	574.46	220.64	424.2	345.06	612.16	561.13	0.314620203	0.02439	0.00372284
1087	(formerly 2C), magnesium-dependent, gamma isoform	787.65	715.87	989.87	1836.5	1670.65	1777.8	1497.19	1797.05	1727.82	0.483825524	0.02463	0.00012463
12224	solute carrier family 44, member 2	389.99	279.49	239.58	762.05	891.14	732.46	894.68	755.73	873.96	0.370287697	0.02469	5.28E-005
8376	poly (A) polymerase alpha	27.35	12.11	9.5	148.02	91.72	167.2	349.39	111.73	134.94	0.097627119	0.02475	0.00014113
15635	DEAD (Asp-Glu-Ala-Asp) box polypeptide 54	1421.61	650.28	836.92	1354.5	1685.81	1727.6	2190.64	2797.01	3685.12	0.432837698	0.025	0.01163473
22224	WD repeat domain 81	30.24	51.94	30.12	149.76	94.94	145.32	314.53	347.69	678.34	0.129783079	0.025	0.00279921
6961	defensin related cryptdin 3	151.47	120.24	161.96	432.42	333.51	435.02	555.4	446.67	466.28	0.32493163	0.0251	4.13E-005
2811	metal response element binding transcription factor 2	1221.23	1231.8	963.63	2148.1	1573.31	2029	2913.63	2368.89	2633.34	0.500013171	0.02513	0.00216856
38660	thioredoxin-like 2 /// similar to thioredoxin-like 2	256.9	121.83	146.19	592.86	461.89	532.76	354.26	505.22	432.44	0.364599938	0.02517	0.00046608
19232	proteasome (prosome, macropain) subunit, beta type 4	308.03	183.74	228.7	497.19	390.48	588.01	679.25	828.99	1020.79	0.359811322	0.02521	0.0025591
7130	GCN5 general control of amino acid synthesis-like 2 (yeast)	166.77	199.37	129.42	236.69	512.49	594.15	527.41	562.67	457.51	0.342838958	0.02525	0.00158602
39148	protein phosphatase 1A, magnesium dependent, alpha isoform	928.91	352.17	441.26	1276.6	1065.68	1257	1597.52	1131.12	1120.34	0.462479307	0.02525	0.0028015
20760	RIKEN cDNA 6330412F12 gene	1221.12	833.76	1060.24	2026.1	1856.48	2309.3	2009.25	2276.23	2705.78	0.472593299	0.02532	0.00036731
30288	RIKEN cDNA D630023B12 gene	231.8	124.88	215.74	650.6	584.87	451.15	354.09	367.11	666.9	0.372339595	0.02534	0.00162651
6136	matrix metalloproteinase 19	680.52	322.99	248.89	1034.6	954	994.51	1010.69	1245.39	922.69	0.406499315	0.02538	0.00106165
504	glutamate dehydrogenase 1	522.62	453.19	388.25	953.35	635.51	724.4	1144.23	1119.64	1119.85	0.478871262	0.02542	0.00299594
19855	acyl-CoA synthetase long-chain family member 4	127.18	124.68	134.3	543.08	402.43	284.25	500.48	291.57	601.29	0.294430254	0.02548	0.000421
10996	methionyl aminopeptidase 1	241.66	153.56	132.59	424.1	450.43	589.31	730.27	394.02	744.54	0.316749033	0.02551	0.00058301
12146	scaffold attachment factor B2	210.74	107.75	267.22	331.79	464.58	1217.3	391.5	663.29	603.9	0.31898202	0.02551	0.00753413
14456	eukaryotic translation initiation factor 4E member 2	608.3	392.9	580.62	1063.5	641.07	821.13	1540.18	1462.72	1376.89	0.458135605	0.02553	0.01083693
24272	ribosomal protein S21	2211.44	1389.97	1767.7	3057.7	2659.66	3238.2	3772.83	3453.75	5642.06	0.492033374	0.02564	0.00500878
20122	histocompatibility 47	1061.18	593.13	563.78	1601.4	1273.75	1337.6	1610.56	1602.25	1546.14	0.494465907	0.02575	0.00117621

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
1829	integrin beta 5	20.55	31.81	35.62	138.64	106.21	113.56	245.56	233.83	262.94	0.159856097	0.02577	0.0001947
8820	peptidase (mitochondrial processing) alpha	103.32	47.96	17.42	213.06	180.94	210.53	289.87	282.06	241.71	0.237912239	0.02577	0.00089259
8632	polymerase (RNA) II (DNA directed) polypeptide H	546.89	407.89	194.5	750.55	562.77	678.85	1016.14	1159.35	1244.52	0.424701322	0.02586	0.00859166
18309	fibroblast growth factor 4	35.44	14.3	20.86	432.43	172.1	100.94	143.65	149.66	594.64	0.088614427	0.02591	0.00088338
29855	ankyrin repeat domain 35	194.21	126.07	115.62	458.93	307.25	358.15	396.59	1145.29	528.2	0.272914247	0.02595	0.00318398
23808	RIKEN cDNA 4632434I11 gene	448.86	431.73	363.62	671.77	596.73	936.9	1130.57	1078.94	1163.22	0.446102905	0.02597	0.0029039
16734	squamous cell carcinoma antigen recognized by T-cells 1	321.84	208.57	257.48	621.17	329.5	867.92	525.03	704.23	1019.17	0.387453221	0.02604	0.00692468
20423	splicing factor, arginine/serine-rich 15 /// similar to splicing factor, arginine/serine-rich 15	1714.9	1723.34	1637.59	2614.4	2538.82	2537.6	4326.68	3486.43	3541.53	0.533024807	0.02609	0.00359332
21533	ribosomal protein L41	2356.65	1636.04	2315.68	3506.7	3203.28	3590.5	4290.25	3926.82	4970.51	0.537155245	0.02618	0.00181749
11378	gb:AV003927 /DB_XREF=gi:4780777 /DB_XREF=AV003927 /CLONE=0610038F02 /FEA=mRNA /CNT=8 /TID=Mm.89136.2 /TIER=Stack /STK=8 /UG=Mm.89136 /LL=15078 /UG_GENE=H3f3a /UG_TITLE=H3 histone, family 3A	1053.61	887.11	842.46	1543.8	1584.51	1875.9	1827.18	1989.27	2602.7	0.487282112	0.0262	0.00095764
10575	Yip1 domain family, member 4	1301.23	782.33	716.95	1652.4	1414.23	1672.3	2397.45	2218.58	2274.82	0.481613455	0.02632	0.00312624
27086	RIKEN cDNA 6720458F09 gene	514.62	69.3	305.05	616.13	470.79	506.67	1261.3	1536.3	928.03	0.334248254	0.02643	0.01918525
7894	WD repeat domain 36	254.38	272.47	200.29	610.23	547.12	595.16	573.36	653.39	981.04	0.367214605	0.0265	0.00024551
20279	UTP20, small subunit (SSU) processome component, homolog (yeast)	265.64	117.47	77.53	395.94	294.27	367.33	559.06	746.95	535	0.317841679	0.02655	0.0029134
1828	integrin beta 5	24.85	44.95	37.18	101.84	100.6	157.78	300.13	259.71	279.75	0.178328235	0.0266	0.00087414
11029	F-box only protein 33	352.6	363.65	331.07	680.26	745.22	715.11	875.23	916.58	1126.49	0.414051304	0.0266	0.00023586
29619	BCL2-associated transcription factor 1	2093.59	1414.28	1845.33	3045.3	2613.48	2893.6	4169.09	3260.8	3703.5	0.543864934	0.02664	0.00241845

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
7972	kinesin family member 22	361.65	356.42	276.4	873.55	734.19	599.86	722.03	806.28	1047.44	0.415804823	0.02667	0.00033015
20855	upstream binding transcription factor, RNA polymerase I	926.95	606.37	732.79	1499.9	977.24	1296.7	1732.93	1838.77	1899.81	0.490214572	0.02667	0.00423551
3937	purine rich element binding protein B	92.81	87.2	83.07	292.59	182.12	245.32	380.5	365.18	322.38	0.294258119	0.02669	0.00018866
15556	ataxin 10	17.33	32.34	12.45	228.41	198.1	179.58	181.72	19.95	77.06	0.140412739	0.02674	0.01259115
32318	Guanine nucleotide binding protein (G protein), beta polypeptide 1-like	38.16	26.14	37.97	167.8	192.62	140.2	315.19	211.09	159.05	0.172469328	0.02675	3.09E-005
20865	transportin 3	498.71	536.39	322.85	1055.4	938.14	988.25	892.24	975.15	1289.64	0.442415456	0.02676	0.00038598
4981	sphingosine-1-phosphate phosphatase 1	1139.7	516.96	256.51	1556.5	1386.43	1522.7	1138.21	1603.75	1205.39	0.454813871	0.02688	0.00679675
19730	gb:M11310.1 /DB_XREF=gi:192009 /FEA=FLmRNA /CNT=3 /TID=Mm.1786.2 /TIER=FL /STK=2 /UG=Mm.1786 /LL=11821 /UG_GENE=Aprt /UG_TITLE=adenine phosphoribosyl transferase /DEF=Mouse adenine phosphoribosyltransferase (APRT), complete cds. /FL=gb:M11310.1	324.96	291.64	351.88	787.01	359.57	481.37	843.3	1266.51	1056.9	0.403982764	0.02688	0.01927971
22636	glutamine fructose-6-phosphate transaminase 1	91.19	137.72	9.09	335.73	165.1	314.89	566.93	599.99	90.41	0.229613372	0.02691	0.02247234
2926	ubiquitin-conjugating enzyme E2H	294.34	174.38	235.24	787.87	546.3	553.52	647.84	590.18	714.62	0.366614327	0.02697	0.00012805
27358	RIKEN cDNA 9630055N22 gene /// similar to Feline leukemia virus subgroup C receptor-related protein 1 (Feline leukemia virus subgroup C receptor) (hFLVCR)	51.52	68.1	41.88	364.66	211.27	222.89	271.25	214.45	185.62	0.219706967	0.02698	4.17E-005
9307	sorting nexin 6	301.12	372.14	208.12	830.58	453.64	637.73	836.86	766.12	1038.77	0.386256765	0.02703	0.00175737
16924	torsin family 1, member B	614.56	254.05	272.1	1028.7	649.17	887.96	740.56	964.62	1195.38	0.417351759	0.02703	0.00265516

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
13406	RIKEN cDNA 1700081L11 gene	279.8	322.06	289.78	766.08	332.23	382.41	846.57	836.8	863.11	0.442808899	0.02708	0.02078566
4267	eukaryotic translation termination factor 1	631.46	328.73	441.06	971.21	1389.44	1088.4	1011.03	1046.31	2206.07	0.363373978	0.02717	0.0020725
10910	nucleoporin 107	200.91	280.21	189.53	608.9	413.29	503.35	667.94	577.6	680.86	0.388564112	0.02717	0.00027743
3500	G patch domain containing 4	532.74	280.11	245.54	950.82	529.03	789.31	962.71	725.77	902.34	0.435553233	0.02727	0.00265492
16856	solute carrier family 12, member 2	171.34	87.64	209.04	527.45	337.21	487.48	447.9	572.27	515.56	0.324128164	0.02732	0.00041255
23353	GIN5 complex subunit 3 (Psf3 homolog)	212.75	16.22	83.74	345.6	181.35	376.3	448.35	272.34	362.05	0.314915986	0.02737	0.00945197
11567	thyroid hormone receptor associated protein 3	409.89	226.84	478.01	731.74	781.37	771.17	1116.95	1028.78	1020.12	0.409069141	0.0274	0.00102386
20123	RIKEN cDNA D230025D16 gene	566.99	244.72	230.85	773.96	545.87	752.66	969.51	843.84	1512.66	0.386240622	0.02747	0.00551593
23022	gene model 608, (NCBI)	19.79	34.73	13.36	118.5	152.45	170.4	284.71	124.46	269.07	0.121258675	0.02747	8.52E-005
957	sarcosine dehydrogenase	102.85	51.26	39.67	276.87	330.62	409.42	265.49	87.25	247.39	0.239672488	0.02752	0.00405166
11737	integrin beta 4 binding protein	1478.64	784.12	1168.22	2140.2	1696.43	2027.5	2235.45	2368.91	2496.33	0.529274561	0.02756	0.00235946
12467	integrator complex subunit 7	15.64	11.95	9.45	107.56	81.04	69.17	167.85	135.22	200.6	0.097289347	0.02757	3.84E-005
28696	cDNA sequence BC031781	206.5	200.11	102.8	539.87	415.28	426.91	561.46	529.25	585.07	0.3331829	0.02762	0.0002035
2616	dodecenoyl-Coenzyme A delta isomerase (3,2 trans-enoyl-Coenzyme A isomerase)	213.16	144.86	245.49	656.83	765.69	443.88	467.21	472	709.62	0.343368713	0.02765	0.00043803
16782	Traf and Tnf receptor associated protein	86.91	90.66	92.25	320.57	262.64	310.65	271.8	329.04	442.44	0.278575632	0.02778	1.87E-005
17246	piwi-like homolog 2 (Drosophila)	194.88	162.18	83.38	485.71	471.02	370.39	490.29	309.46	467.42	0.339545695	0.02778	0.00050921
28263	zinc finger protein 297B	346.22	252.82	249.98	689.62	506.85	491.93	824.26	739.66	843.71	0.414557511	0.02778	0.00089669
13852	CCR4-NOT transcription complex, subunit 1	624.54	555.94	435.99	1003.5	854.27	1131.5	1220.49	1252.71	1422.68	0.46955327	0.02788	0.00078009
23277	biogenesis of lysosome-related organelles complex-1, subunit 2	79.81	76.06	64.03	325.63	170.46	148.71	292.97	413.85	462.88	0.242380821	0.02793	0.00148942
42753	RIKEN cDNA 2010002M12 gene	174.49	77.58	21.59	574.43	290.32	438.63	251.82	312.33	262.98	0.256896236	0.02799	0.00346194
2019	cDNA sequence BC038286	125.97	73.92	68.54	284.36	290.06	209.54	343.44	337.31	491.7	0.274410783	0.028	0.00030981

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
39706	/DB_XREF=gi:12854789 /GEN=Hpk1 /FEA=mRNA /CNT=3 /TID=Mm.160710.1 /TIER=ConsEnd /STK=2 /UG=Mm.160710 /UG_TITLE=Mus musculus adult male testis cDNA, RIKEN full-length enriched library, clone:4930557G23:homeodomain interacting protein kinase 1, full insert sequence /DEF=Mus musculus adult male testis cDNA, RIKEN full-length enriched library, clone:4930557G23:homeodomain interacting protein kinase 1,	615.34	546.59	659.88	1182.5	1131.4	1245.5	1548.48	1390.42	1818.55	0.438098857	0.02804	0.00032472
27310	G patch domain and KOW motifs	61.52	159.03	86.57	532.82	393.07	275.19	336.31	260.97	331.72	0.288364756	0.02809	0.00056609
15631	Max dimerization protein 3	231.87	142.18	388.83	680.62	398.74	1726.9	635.95	544.19	1682.23	0.269158034	0.0282	0.01298519
1794	translocase of inner mitochondrial membrane 10 homolog (yeast)	445.43	218.47	297.38	1237.7	550.92	497.86	560.37	1173.81	1446.18	0.351674081	0.02825	0.01197186
40394	RIKEN cDNA A230097K15 gene	542.56	392.94	536.72	1135.4	943.8	714.8	1147.59	1095.12	1136.24	0.476993833	0.0283	0.00099635
13932	cytochrome c oxidase, subunit XVII assembly protein homolog (yeast)	374.36	512.87	771.27	1560.4	816.01	1231.3	757.93	1492.77	1259.95	0.465978119	0.02834	0.00846149
12583	nucleolar and coiled-body phosphoprotein 1	1396.29	962.28	1269.5	2433.1	1712.66	2129	2601.27	2478.57	2899.66	0.509051699	0.02841	0.00166888
44733	RIKEN cDNA 2610510H03 gene	1694.62	1302.55	1307.24	2154.8	2197.38	2580.1	2853.86	2696.93	3033.96	0.554799252	0.02841	0.00127502
24057	RIKEN cDNA 6330548G22 gene	1941.19	1205.89	1387.66	2922.2	2488.55	2691	2668.23	2814.66	3260.93	0.53838802	0.02846	0.00088299
9979	G patch domain containing 4	1180.84	937.75	328.1	2341.4	1228.15	1953.3	2071.81	1558.28	1584.77	0.455720416	0.02857	0.00998652
15379	sorbitol dehydrogenase	27.38	63.24	77.06	217.25	225.56	241.16	303.1	194.63	333.28	0.221362658	0.02857	0.00015891
1395	CD320 antigen	376.3	186.45	150.95	519.75	347.27	458.89	885.75	732.03	898.14	0.371539753	0.02874	0.00681172

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	tpvals
10147	homeodomain interacting protein kinase 1 /// similar to homeodomain-interacting protein kinase 1	162.22	95.7	238.88	242.84	377.63	282.15	900.14	726.92	970.38	0.283880848	0.02874	0.01486244
40408	zinc finger and BTB domain containing 11	154.16	137.47	107.76	393.41	317.18	313.89	408.57	545.45	573.96	0.312945159	0.02885	0.00024451
12344	La ribonucleoprotein domain family, member 1 /// similar to la related protein isoform 1	383.9	74.36	193.38	510.75	581.65	447.49	684.6	681.6	1010.2	0.332784344	0.0289	0.00428131
801	proteasome (prosome, macropain) subunit, alpha type 6	317.41	182.75	245.98	577.9	392.79	559.5	873.25	678.68	1083.53	0.358234609	0.02896	0.00296154
12365	gb:AW545056 /DB_XREF=gi:7187569 /DB_XREF=C0190C04-3 /CLONE=C0190C04 /FEA=mRNA /CNT=182 /TID=Mm.29909.1 /TIER=Stack /STK=60 /UG=Mm.29909 /LL=69023 /UG_GENE=1810005K14Rik /UG_TITLE=RIKEN cDNA 1810005K14 gene	575.81	256.8	387.06	829.02	524.34	892.94	1101.05	1108.62	1025.51	0.44501485	0.02907	0.00545304
13580	RIKEN cDNA 9430010O03 gene	252.11	205.75	110.44	548.01	407.24	529.86	651.64	535.7	411.67	0.368533001	0.0293	0.00067634
29034	jumonji, AT rich interactive domain 1A (Rbp2 like)	37.2	22.61	18.65	222.58	113.9	160.57	93.92	144	233.42	0.162042152	0.02951	0.00011963
8637	Bardet-Biedl syndrome 2 homolog (human)	145.67	71.74	140.97	337.79	236.48	247.56	500.41	586.34	389.87	0.311844939	0.02961	0.00216673
21932	Transcribed locus	449.07	163.81	87.36	583.75	309.89	572.13	653.87	636.22	766.49	0.397598194	0.0297	0.00950051
24471	TSC22 domain family 2	1111.19	838.9	701.35	1530.6	1318.86	1453.7	2101.28	2010.43	1899.33	0.514136405	0.0298	0.00250066
12713	transportin 3	236.77	170.07	152.07	455.61	333.47	459.35	782.62	552.05	536.83	0.358283679	0.0299	0.00088863
4235	zinc finger protein 644	25.19	29.71	183.22	387.55	173.75	307.41	242.2	191.55	329.75	0.291776181	0.03064	0.00413242
27956	RIKEN cDNA 4933439C20 gene /// phosphatidylserine decarboxylase	214.02	187.62	218.88	489.82	385.39	354.78	580.99	667.79	1214.05	0.336068371	0.03073	0.00567336
14945	proteasome (prosome, macropain) subunit, alpha type 6	673.17	339.26	437.73	823.99	814.83	842.51	1149.91	1450.18	1653.82	0.430618656	0.03081	0.00552425

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
8311	CDNA clone IMAGE:30031514	1455.82	885.16	844.81	1802.7	1683.15	1977.4	2026.15	1949.49	2208.37	0.547043051	0.03107	0.00167117
18423	stromal antigen 2	448.64	362.26	221.01	1147.6	777.91	1089.8	665.09	575.32	749.19	0.412363183	0.03116	0.00282021
7826	methionine adenosyltransferase II, alpha	1253.06	613.26	919.4	1482.9	1371.95	1652.5	2216.98	1838.93	2120.48	0.521490234	0.03143	0.00499621
8396	zinc finger protein 639	651.33	592	510.89	1190.3	1159.1	1416.3	1011.3	1284.01	1188.3	0.483970154	0.03152	0.00017759
19114	WD repeat domain 36	205.63	144.93	118.6	423.92	347.67	404.86	409.04	556.77	672.92	0.333307284	0.03161	0.00038691
9191	ubiquitin A-52 residue ribosomal protein fusion product 1 /// similar to ubiquitin A-52 residue ribosomal protein fusion product 1 /// similar to ubiquitin A-52 residue ribosomal protein fusion product 1 /// similar to ubiquitin A-52 residue ribosomal protein fusion product	1179.37	800.7	703.02	1290.2	1670.19	1344.1	2225.44	1773.66	3566.28	0.452084904	0.0317	0.01417489
6104	brix domain containing 1	189.12	152.83	356.96	579.34	457.58	505.53	797.43	691.14	566.73	0.388526162	0.03179	0.00117216
14528	gene model 288, (NCBI)	611.12	289.52	223.86	897.94	493.54	694.95	841.18	913.47	964.44	0.468003463	0.03188	0.00748606
14234	ribosomal protein S3	1603.17	992.86	1195.88	2140	1511	1797.1	2583.61	2433.02	3136.49	0.557583805	0.03191	0.01375262
1610	GRIP1 associated protein 1	156.21	52.05	43.1	267.72	230.72	274.17	284.24	349.37	343.89	0.287250516	0.03198	0.00064653
607	arginyl-tRNA synthetase	475.28	273.88	474.37	856.71	656.69	777.19	931.87	958.54	1034	0.469234899	0.032	0.00130575
8157	general transcription factor IIIC, polypeptide 5	212.08	99.85	122.9	316.44	273.49	369.01	425.12	499.92	698.25	0.336786421	0.03209	0.00211539
10540	peroxisome proliferative activated receptor, gamma, coactivator-related 1	902.33	445.31	392.62	1139.4	999.53	1033.5	1236.07	1335.21	2000.98	0.449403209	0.03215	0.00487984
17591	zinc finger protein 292	33.35	44.66	45.16	200.8	146.72	175.4	256.79	209.89	116.94	0.222621866	0.03216	6.65E-005
16780	histocompatibility 47	1837.26	1122.62	1324.02	2610	1889.17	2344.4	2583.68	2481.42	2674.07	0.587533816	0.03217	0.00342811
257	eukaryotic translation initiation factor 3, subunit 3 (gamma)	627.16	426.41	780.66	1198.3	1019.38	1204.3	1897.89	1173.9	2308.62	0.41675632	0.03226	0.00484988
13101	heterogeneous nuclear ribonucleoprotein A/B	1618.69	1405.47	1477.8	2463.2	2093.97	2498.9	2936.45	2830.56	2597.97	0.583869773	0.03233	0.00099967
33525	Transcribed locus	83.97	35.52	295.51	432.87	220.51	436.57	389.83	242.22	680.37	0.34549216	0.03235	0.01240034

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
2568	ribosomal protein L30 /// similar to ribosomal protein L30 /// similar to ribosomal protein L30 /// similar to ribosomal protein L30 /// similar to ribosomal protein L30	1254.95	735.95	965.54	1689.5	1596.35	1943.3	1668.9	1938.22	2788.5	0.508643152	0.03235	0.00325586
7517	RIKEN cDNA 1810009K13 gene	143.62	57.05	68.62	322.65	136.04	252.87	427.73	276.99	553.43	0.273431114	0.03236	0.00435699
19614	TIP41, TOR signalling pathway regulator-like (S. cerevisiae)	1754.37	1366.54	1274.92	2383.4	2108.26	2070.9	2846.61	2746.03	2667.1	0.593140169	0.03241	0.0026507
5982	protein phosphatase 2 (formerly 2A), catalytic subunit, beta isoform	551.48	339.48	280.15	775	546.79	692.17	928.41	1201.26	1257.08	0.433687423	0.03243	0.00657847
9877	influenza virus NS1A binding protein	975.18	492.65	387.23	858.37	991.15	1163.2	1663.53	2058.91	1252.97	0.464454715	0.03248	0.01279151
16885	chromosome segregation 1-like (S. cerevisiae)	121.07	97.77	145.66	350.27	345.42	378.07	298.67	387.07	410.09	0.336008186	0.03254	2.39E-005
29192	cleavage and polyadenylation specific factor 6	17.6	44.66	34.12	174.3	133	221.06	127.6	192.74	129.81	0.196993388	0.03256	8.84E-005
8733	transmembrane emp24 protein transport domain containing 5	207.72	120.4	223.01	611.94	436.79	466.68	476.75	506.84	551.44	0.361344593	0.03257	0.00015655
8642	/DB_XREF=gi:14318706 /FEA=FLmRNA /CNT=134 /TID=Mm.29500.1 /TIER=FL+Stack /STK=29 /UG=Mm.29500 /DEF=Mus musculus, hypothetical protein FLJ10788, clone MGC:6884 IMAGE:2651599, mRNA, complete cds.	180.91	186.12	138.75	443.86	379.97	493.4	573.62	451.43	424.69	0.365584014	0.03261	5.25E-005
27736	RIKEN cDNA C730049O14 gene	468.55	201.7	280.69	854.88	650.64	580.29	733.69	724.56	697.53	0.448388458	0.03263	0.00126179
15165	glutaredoxin 2 (thioltransferase)	187.31	23.63	26.17	284.27	207.99	195.31	286.09	281.75	295.36	0.305796475	0.03264	0.00335924

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
34773	coiled-coil domain containing 66	327.65	244.56	99.43	681.69	370.27	538.09	732.89	447.88	631.69	0.394790904	0.03268	0.00477611
22170	hexosaminidase B	466.76	254.86	157.95	680.55	374.26	443.5	767.08	800.87	731.5	0.463204626	0.03271	0.01339699
41702	PREDICTED: Mus musculus similar to aspartoacylase-3 (LOC629627), mRNA	52.96	7.24	13.47	124.13	107.95	76.94	201.8	231.95	168.21	0.161737909	0.03274	0.00107041
8599	mitochondrial ribosomal protein S6	84.99	90.76	80.52	172.16	157.7	188.73	395.52	436.4	635.42	0.258085632	0.03279	0.00751919
12768	zinc finger protein 644	15.61	32.03	46.1	366.23	40.06	93.41	192.4	133.37	321.07	0.163518063	0.03279	0.01131652
19905	junction-mediating and regulatory protein	169.13	233.3	148.29	543.3	293	427.21	657.56	428.21	656.69	0.366417496	0.03284	0.00158297
13983	bolA-like 2 (E. coli)	640.93	283.51	474.89	891.08	567.86	705.83	1189.82	1104.23	1324	0.483961112	0.03286	0.01507785
12572	BCL2-associated transcription factor 1	2243.95	1513.18	2041.07	3292.2	2693.98	3229.5	3913.71	3100.47	3629.4	0.583930579	0.03293	0.00247576
13491	phosphatidylinositol 4-kinase type 2 alpha	106.26	77.85	116.88	271.2	295.15	204.29	626.53	277.93	551.44	0.270365679	0.03294	0.00205473
8482	nucleolar complex associated 2 homolog (S. cerevisiae)	984.9	484.36	571.5	1289.3	1202.67	1169.9	1470.81	1389.35	1671.63	0.498135738	0.03297	0.00193977
18925	hippocampus abundant gene transcript 1	1626.37	984.96	926.25	1800.2	1384	1711.6	2541.97	2468.73	3077.91	0.544894435	0.03302	0.02020823
29821	general transcription factor III C 1	649.54	697.58	431.58	1080.6	901.53	1212.7	1451.96	1192.11	1707.24	0.471422324	0.03303	0.00235796
28901	RIKEN cDNA 4932408B21 gene	57.42	28.78	34.91	186.88	112.52	187.44	217.22	178.97	178.07	0.228272547	0.03307	6.72E-005
27240	nucleoporin 98	708.33	558.18	484.95	1086.1	858.65	840.28	1314.93	1255.99	1590.24	0.504291543	0.0331	0.00479921
16347	DEAD (Asp-Glu-Ala-Asp) box polypeptide 21	691.31	341.98	479.23	933.21	720.44	1161.8	1068.9	1079.18	1463.87	0.470644611	0.03313	0.00388951
12433	general transcription factor IIIC, polypeptide 2, beta	134.75	114	117.11	326.86	320.97	220.76	337.18	352.3	512.23	0.3534367	0.03314	0.00046942
16878	nuclear fragile X mental retardation protein interacting protein 1	735.99	520.27	616.98	1152	1106.58	1108.4	1669.92	1264.41	1709.52	0.467676882	0.03316	0.00109882
20819	ubiquitin-conjugating enzyme E2C	1431.97	1109.57	1108.32	1340.1	1457.9	1689.9	3079.28	3096.31	3132.89	0.529102477	0.03318	0.04620446
15367	adhesion regulating molecule 1	2376.07	1243.06	1825.49	2794.9	2622.99	3207.4	2488.6	3090.31	3801.22	0.604778024	0.0332	0.00947177
35487	TatD DNase domain containing 1	49.58	54.22	10.94	234.77	114.06	144.99	202.21	154.66	288.54	0.201434302	0.03323	0.00129083

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
26869	guanine monphosphate synthetase	122.92	142.78	107.6	432.72	251.71	382.77	333.62	321.76	588.17	0.323098561	0.03324	0.0004596
17750	Zinc finger protein 644	18.41	38	141.75	287.13	126.78	223.87	231.07	167.08	286.89	0.299602365	0.03327	0.00457717
10563	ring finger protein 11	81.84	78.97	111.45	413.57	203.45	266.73	291.63	298.49	401.2	0.290399825	0.03333	0.00019063
22525	ribosomal protein L23	1694.63	1408.99	1322.27	2543.8	2122.05	2440.6	2267.39	2629.52	3731.63	0.562553901	0.03333	0.00413512
41190	methyltransferase like 2	201.66	120.68	113.21	444.44	370.04	370.14	371.11	445.47	374.11	0.366731079	0.03333	9.87E-005
22167	CNDP dipeptidase 2 (metallopeptidase M20 family)	22.88	61.95	32.07	330.23	79.86	180.33	84.76	218.18	158.8	0.22220955	0.0334	0.00380262
1037	craniofacial development protein 1	632.43	487.38	759.93	1272.7	874.72	1051.5	1332.88	1198.09	1629.04	0.510874572	0.03341	0.00315993
19490	Rho GTPase activating protein 8	561.58	232.84	159.57	755.26	534.11	863.06	783.69	493.49	842.26	0.446638123	0.03343	0.00744318
23936	RIKEN cDNA 2810004A10 gene	611.4	319.12	196.57	795.57	922.98	799.63	775.73	795.6	585.57	0.482169289	0.03346	0.00506537
38670	fibroblast growth factor receptor-like 1	170.84	122.93	50.89	527.58	138.93	388.29	327.68	457.46	183.49	0.340669062	0.03349	0.01815324
13982	bolA-like 2 (E. coli)	667.26	327.26	632.69	997.91	533.2	734.58	1519.41	1322.18	1597.39	0.485396	0.03353	0.03999237
21923	gene /// RIKEN cDNA 5830484A20 gene /// similar to Sp110 nuclear body protein ///	936.03	563.58	807.33	1249.3	1254.37	1612.3	1349.57	1971.41	2092.11	0.48419199	0.03354	0.00309151
7522	sortilin 1	47.52	42.59	83.69	467.9	79.92	299.98	346.19	142.97	127.38	0.237376566	0.0336	0.01339749
20481	SDA1 domain containing 1	1362.71	859.8	1118.87	2204.8	1724.2	2050.9	2051.86	2375.74	1968.44	0.539978684	0.03364	0.00101078
677	cathepsin C	620.48	261.03	388.32	1142.3	623.95	597.3	1072.15	767.24	1113.22	0.47772632	0.03365	0.01077421
12728	polymerase (RNA) III (DNA directed) polypeptide E	1173.78	856.85	818.59	1893.6	1486.88	1477.6	1672.05	1702.36	2013.3	0.556175975	0.03366	0.00133563
171	ribosomal protein S26	1005.17	525.6	798.19	1159.7	1132.3	1291	1650.32	1768.46	3066.94	0.462613383	0.03373	0.01726663
21453	ribosomal protein S8	2653.11	1674.67	2415.56	3453.2	3323.49	3368.9	3676.28	3927.51	5180.08	0.588181318	0.03373	0.00586965
18058	AF4/FMR2 family, member 4	66.6	79.82	102.25	260.49	269.44	166.59	260.12	196.38	405.8	0.319049024	0.0338	0.00066396
2114	torsin family 1, member B	611.51	202.39	243.41	852.18	486.97	714.88	856.26	824.23	655.93	0.481640834	0.03386	0.00820538
7324	metastasis-associated gene family, member 2	808.65	595.65	569.46	1393.2	875.73	1334.4	1064.04	1538.48	1479.47	0.513642858	0.03395	0.00325528
20207	PHD finger protein 17	45.11	97.31	48.99	150.19	219.06	156.4	230.13	395.67	311.67	0.261646345	0.03398	0.00115284
22719	myelin protein zero-like 1	26.63	58.06	86.57	211.23	241.63	192.89	146.38	553.53	185.92	0.223638334	0.034	0.00221299
13129	mediator of RNA polymerase II transcription, subunit 8 homolog (yeast)	88.18	61.52	35.11	199.79	152.12	177.8	282.24	471.76	209.96	0.247457604	0.03405	0.00141734

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
33684	activating transcription factor 7 interacting protein 2	465.25	439.02	701.39	1507.1	999.44	1317.4	817.23	803.3	1090.48	0.491410733	0.03407	0.00480905
9652	bone morphogenetic protein receptor, type 1A	30.66	33.23	24.13	164.86	83.03	98.67	231.61	148.75	169.39	0.196405262	0.03412	0.00020267
20214	RNA binding motif protein 5	571.27	767.82	724.89	970.89	1114.44	1258.3	1664.76	1551.93	1412.32	0.517765759	0.03415	0.00244522
16715	spermatogenesis associated 5	137.92	102.31	61.45	459.08	274.64	228.49	327.14	221.78	356.12	0.323127594	0.03417	0.00093717
24191	cancer susceptibility candidate 5 /// similar to cancer susceptibility candidate 5 isoform 2	152.1	164.87	125.69	368.06	261.62	337.15	517.57	426.13	722.68	0.336213215	0.03421	0.00163987
27633	vitamin K epoxide reductase complex, subunit 1-like 1	158.7	49.79	76.93	398.67	216.04	284.84	293.31	284.28	381.74	0.307088139	0.03423	0.00085041
7003	WD repeat domain 77	1094.94	580.22	464.08	1369.7	1345.02	1526.4	1197.88	1576.35	1489.59	0.503058814	0.03427	0.00254454
344	glycine decarboxylase	449.54	174.14	151.67	613.06	279.64	804.84	631.24	715.88	615.34	0.423688525	0.03431	0.01299645
1917	solute carrier family 12, member 2	500.3	207.9	501	1012.6	574.08	711.55	831.2	898.12	1082.66	0.473252358	0.03432	0.00723914
15722	DEAD (Asp-Glu-Ala-Asp) box polypeptide 17	385.08	297.48	284.8	475.57	561.19	524.9	865.28	1016.7	851.49	0.450445039	0.03434	0.00548161
3811	DNA segment, human DXS9928E	157.43	142.21	194.44	493.98	482.27	475.73	458.09	304.57	395.93	0.378522698	0.03438	0.00016711
14157	RIKEN cDNA 2610101J03 gene	1078.89	693.47	683.97	1413	1116.89	1409.9	2146.69	1725.33	1648.32	0.519305374	0.0344	0.00470655
21888	death inducer-obliterator 1	489.88	352.87	242.07	885.74	746.18	817.23	629.22	636.1	1048.16	0.455555019	0.0344	0.00196002
6127	Willi/Angelman syndrome 2 homolog (human)	146.49	115.62	62.97	277.82	180.98	205.37	423.86	504.78	426.8	0.32192354	0.03448	0.00497845
10418	pantothenate kinase 3	571.77	464.21	419.81	1151.2	770.81	1060.5	873.11	919.21	1010	0.503316456	0.03455	0.00060585
7942	torsin family 2, member A	56.41	57.84	15.89	182.4	117.93	153.01	167.89	264.45	201.53	0.239401772	0.03455	0.00088711
6639	sorting nexin 3	120.78	103.65	110.21	326.23	216.07	202.87	341.2	336.98	604.44	0.330053901	0.03456	0.00250308
3398	abhydrolase domain containing 6	330.84	210.9	246.57	422.75	380.61	691.33	684.84	889.36	650.71	0.423868158	0.03459	0.00458787
640	translocase of inner mitochondrial membrane 8 homolog a1 (yeast)	878.17	670.48	908.3	1804.6	1319.02	1419.3	1722.45	1433.31	1537.61	0.532026734	0.03462	0.00067228
7006	protein kinase C, delta	411.25	383.64	115.26	712.82	315.54	733.67	910.26	661.36	733.23	0.447591274	0.03465	0.02184294
15510	Williams-Beuren syndrome chromosome region 1 homolog	387.38	65.5	224.42	655.34	402.2	299.57	683.74	452.6	930.59	0.395614537	0.03469	0.0206498
1970	midasin homolog (yeast)	72.03	19.88	33.69	203.55	127.77	190.18	139.01	179.55	201.68	0.241135024	0.03474	0.0003368
3936	purine rich element binding protein B	1414.47	939.03	992.32	1815.5	1442.75	1699.3	2127.53	2327.95	2198.24	0.576307096	0.03476	0.00541495

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
11214	thioesterase superfamily member 4	209.2	140.63	221.22	557.73	459.97	598.38	413.32	339.06	663.38	0.376701937	0.03481	0.00078993
20608	RIKEN cDNA 2610304G08 gene	85.77	64.41	24.73	216	142.41	211.13	393.72	237.84	189.99	0.251471867	0.03483	0.00126552
401	KT112 homolog, chromatin associated (S. cerevisiae)	204.06	74	121.41	306.83	197.69	220.06	546.92	436.97	511.52	0.359884504	0.03484	0.00978944
4187	gb:AV006589 /DB_XREF=gi:4783576 /DB_XREF=AV006589 /CLONE=1100006A23 /FEA=EST /CNT=1 /TID=Mm.198379.1 /TIER=ConsEnd /STK=0 /UG=Mm.198379 /LL=99251 /UG_GENE=AV006589 /UG_TITLE=expressed sequence AV006589	750.81	258.88	477.14	770.5	790.87	855.25	1131.06	1138.45	1223.11	0.503222073	0.03488	0.00891259
20263	ATP-binding cassette, sub-family F (GCN20), member 1	1974.08	1512.28	1591.36	2799.7	2379.24	2881.9	3476.53	3087.76	3006.5	0.575978511	0.03492	0.00134869
6063	nuclear receptor coactivator 6 interacting protein	96.49	66.05	71.48	230.69	170.41	262.11	380.42	250.08	337.13	0.286993206	0.035	0.00026083
7582	associated 3	772.56	580.96	729.21	1398.6	1166.76	1297.1	1691.5	1392.13	1417.23	0.498064757	0.03503	0.00033862
7829	signal recognition particle receptor ('docking protein')	1178.28	396.35	506.11	1274.3	1169.1	1329.2	1193.62	1444.27	1684.77	0.51406442	0.03505	0.00684649
1149	solute carrier family 34 (sodium phosphate), member 2	1671.27	1451.17	1598.15	2771.9	2571.18	2756.1	2423.89	2605.98	2900.33	0.588995153	0.03509	0.00034828
23975	RIKEN cDNA D530033C11 gene	15.02	9.78	12.59	135.75	30.07	112.3	106.24	70.57	148.16	0.12399476	0.03512	0.00060996
3060	T-cell immunoglobulin and mucin domain containing 2	695.08	248.62	328	826.75	592.21	729.65	1034.03	1032.95	1807.45	0.422278451	0.03518	0.01412352
7965	integrator complex subunit 4	108.79	143.67	117.13	263.16	266.45	284.38	553.99	382.53	444.37	0.336774676	0.0352	0.00079899
15134	ribosomal protein S8	2879.67	1822.12	2555.47	3715	3487.55	3736.8	3939.03	4199.88	5464.29	0.591403484	0.03525	0.0054164
10616	sarcolemma associated protein	302.65	177.5	294.36	828.01	459.32	599.96	670.2	546.64	765.82	0.400268737	0.03526	0.00086754
19022	folate receptor 1 (adult)	295.13	157.89	124.56	512.13	492.87	451.56	490.43	421.28	529.28	0.39866784	0.03526	0.00048712
39934	RIKEN cDNA 2410019A14 gene	948.04	617.96	722.82	1276.2	1461.42	1309	1701.46	1341.04	1647.3	0.523969663	0.03527	0.00085077

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
27541	hypothetical protein 5430421B17	29.79	29.13	17.85	111.26	148.96	160.01	179.04	101.84	124.88	0.185886028	0.03534	1.86E-005
13353	cyclin L2	366.32	286.12	242.58	661.8	568.59	569.17	785.19	691.43	896.88	0.428951417	0.03535	0.00044813
8628	cleavage and polyadenylation specific factor 4	437.6	464.66	259.65	801.65	552.59	757.67	924.15	849.4	912.93	0.484291606	0.03542	0.00233476
15335	RIKEN cDNA 2810409H07 gene	74.5	32.14	51.2	212.66	134.11	162.73	340.99	117.8	348.25	0.239780029	0.03544	0.00207122
21143	RIKEN cDNA 2900097C17 gene	55.86	42.84	49.42	191.63	97.34	143.81	238.34	247.68	216.44	0.260949227	0.03544	0.00048204
27810	mannosidase 2, alpha 2	187.42	198.71	152.25	448.07	341.04	401.19	578.87	431.83	634.48	0.379745228	0.03549	0.00040675
21839	B-cell receptor-associated protein 31	682.51	376.13	741.32	1355.9	913.47	1360.2	1021.68	1106.77	1232.25	0.514990859	0.03564	0.00340949
27178	spermatogenesis associated 2	1161.91	964.7	1257.96	2142.9	1761.61	1953.7	1664.48	2350.84	2119.38	0.564429895	0.03571	0.00117285
27596	Expressed sequence AI195381	164.06	50.24	97.33	314.61	235.05	369.8	208.02	349.59	306.67	0.34941191	0.03579	0.00171003
8359	SEH1-like (S. cerevisiae)	34.37	35.38	21.55	153.44	52.09	164.71	185.96	151.36	190.7	0.203281901	0.03581	0.0010864
21158	phosphoserine aminotransferase 1	762.08	527.82	423.16	1268.1	1111.16	1110.2	1163.28	1112.98	1052.34	0.502508789	0.03586	0.00058213
20989	ribosomal protein L37	265.68	159.08	539.01	502.7	517.5	685.71	1247.62	1948.95	3246.7	0.236531774	0.0359	0.02475044
20676	reticulin 4	1574.4	894.92	1056.14	1903.9	1775.22	1903.2	2525.25	2161.78	2275.69	0.562051115	0.03594	0.0030746
11523	ornithine decarboxylase, structural 1 /// similar to Ornithine decarboxylase (ODC)	1456.62	734.62	1199.01	1588.9	1504	1639.4	2484.21	2389.48	2308.26	0.569109376	0.03598	0.01491284
6500	lysophospholipase 3	47.88	151.19	39.47	198.56	143.95	151.5	374.75	347.89	480.81	0.281055224	0.03599	0.00760032
788	transmembrane protein 49	214.62	75.2	94.84	335.84	369.33	256.77	372.81	272.48	580.32	0.351681104	0.03602	0.00252792
10737	SNAP-associated protein	273.69	269.17	162.25	547.62	550.93	515.68	525.29	571.28	535.32	0.434432492	0.03605	0.00015943
9707	leukocyte specific transcript 1	150.44	125.82	60.94	231.95	196.75	216.83	442.18	529.47	633.7	0.29961748	0.03608	0.00854339
10862	cyclin G associated kinase	189.88	169.84	165.37	414.6	338.87	451.82	691.18	343.8	583.16	0.371951846	0.03617	0.00088473
12376	p53 and DNA damage regulated 1	548.67	181.79	376.6	789.75	643.57	739.9	846.2	866	924.2	0.460350463	0.03618	0.00344332
12925	enhancer of rudimentary homolog (Drosophila)	971.59	772.44	792.75	1277.8	1052.43	1432.2	1720.87	1915.14	1795	0.551868619	0.03625	0.0060476
980	fibrillarlin	684.3	450.34	486.4	856.32	653.88	1067.7	1220.0	1189.37	1652.28	0.488299694	0.03627	0.01051729
29430	trans-acting transcription factor 6	683.7	635.1	523.7	922.49	838	1035.4	1440.35	1366.46	1588.4	0.512799038	0.03632	0.0059478
22752	phosphoglucomutase 3	117.47	70.55	76.49	280.19	226.01	244.57	283.48	365.08	342.8	0.303680189	0.03636	8.87E-005

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
14032	seizure related 6 homolog like 2	23.56	3.91	6.09	51.61	168.7	132.62	164.68	29.4	81.1	0.106860263	0.0364	0.0018215
21869	similar to ubiquitin A-52 residue ribosomal protein fusion product	1092.71	549.03	901.08	1326.5	1044.1	1421.4	1855.2	1998.4	2994.86	0.477956614	0.03646	0.02013608
15947	expressed sequence AI593864	59.53	130.96	145.41	238.39	245.29	390.12	422.46	217.98	589.3	0.319366401	0.03648	0.00409729
360	ankyrin repeat domain 10	1631.2	1255.22	1597.8	2366.3	1976.89	2233.4	2873.13	2724.96	2719.92	0.602127754	0.03656	0.00313083
13232	brix domain containing 5	249.7	149.34	101.69	500.78	396.36	411.4	449.86	312.53	480.19	0.392556995	0.03664	0.00094472
7575	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily c, member 1	169.06	151.5	175.74	345.56	264.16	208.33	557.93	543.22	522.86	0.40646012	0.03665	0.01048791
847	developmental pluripotency associated 5	637.03	556.11	625.16	1200.8	846.84	1332.8	1135.98	1117.74	2353.11	0.455300635	0.03672	0.00785991
19522	WW domain containing adaptor with coiled-coil	1020.92	699.47	883.58	1264.2	1408.61	1531	1740.17	1592.41	1789.74	0.558426352	0.03672	0.0014871
21628	uridine monophosphate synthetase	98.18	94.72	68.1	288.06	248.98	175.65	306.21	226.92	452.59	0.307346283	0.0368	0.00065722
15391	ras responsive element binding protein 1	141.53	165.74	217.71	421.99	422.79	503.87	510.83	456.58	227.76	0.41274933	0.03686	0.00261093
13913	Heterogeneous nuclear ribonucleoprotein U	22.26	32.92	25.16	157.59	122.26	145.62	79.73	130.3	273.63	0.1767404	0.03688	0.00020361
28627	solute carrier family 19 (sodium/hydrogen exchanger), member 3	485.43	151.49	233.17	737.26	511.1	592.5	613.8	620.42	958.53	0.431419994	0.03694	0.00454474
18771	elongation factor RNA polymerase II 2	604.78	436.5	353.74	1119.5	733.54	995.87	939.53	968.34	900.94	0.493137707	0.03696	0.00098
10803	TBC1 domain family, member 20	126.56	81.16	223.11	336.16	248.27	209.16	442.89	430.88	702.71	0.363558882	0.037	0.01061646
13848	expressed sequence AA408556	215.08	107.35	165.18	295.85	310.39	421.99	454.92	383.23	796.93	0.366168414	0.03707	0.00397733
1683	XPA binding protein 1	410.06	452.6	766.71	1401.7	1053.22	878.62	1090.9	995.33	1162.3	0.495088976	0.03712	0.00193188
19044	ERGIC and golgi 3	44.81	65.7	48.44	93.9	72.33	245.35	177.49	438.27	277.99	0.243539948	0.03714	0.01416792
27096	like) 10	20.8	3.27	12.57	89.8	103.08	86.03	136.13	47.95	210.38	0.108825757	0.03717	0.00063987
727	6-phosphofructo-2- kinase/fructose-2,6- biphosphatase 3	63.36	20.84	112.36	213.6	178.27	137.99	367.86	210.92	349.81	0.269546436	0.03724	0.00380938

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
2739	RGS16	351.74	117.12	76.23	503.15	162.89	413.91	695.01	773.54	1692.2	0.257075483	0.03727	0.02968997
24940	DNA segment, Chr 8, Brigham & Women's Genetics 1414 expressed	118.78	53.72	55.79	186.53	261.76	155.04	261.02	198.73	393.02	0.313563629	0.0373	0.00157746
10152	heat shock protein 110	1398.12	830.88	1073.63	1668.8	1565.54	2160	2008.28	1789.22	2316.3	0.573965664	0.03731	0.0040989
19155	kinesin family member 22	398.22	450.27	354.46	841.6	654.47	559	989.59	886.55	947.43	0.49314973	0.03736	0.00219119
7723	phosphoribosylaminoimidazole carboxylase, phosphoribosylaminoribosylami	391.13	442.83	339.16	827.89	516.64	792.43	952.32	677.57	929.92	0.499543303	0.03737	0.002641
11302	jumonji, AT rich interactive domain 1B (Rbp2 like)	1673.42	942.45	1166.08	2053.4	1794.29	2099.4	2527.42	2125.87	2560.26	0.574734303	0.03738	0.00388238
13496	ribosomal protein L38	1996.91	1375.66	2224.12	2929.6	2359.26	2980.6	2797.37	3517.85	4415.85	0.58910883	0.03741	0.01204374
18935	eukaryotic translation initiation factor 4E /// hypothetical LOC630527	2312.87	1828.13	2131.71	3207.1	2800.93	3286.2	3402.34	3230.56	3391.04	0.649411718	0.03742	0.00176804
10415	translocase of inner mitochondrial membrane 17a	1200.54	871.03	958.28	1480.2	1104.86	1390.9	2541.11	2448.39	2310.83	0.537385623	0.03744	0.02830428
11848	TNFAIP3 interacting protein 1	321.33	225.29	228.01	806.72	511.77	512.88	594.28	486.89	1024.8	0.393478846	0.03745	0.00199172
29413	DCN1, defective in cullin neddylation 1, domain containing 3 (S. cerevisiae)	208.2	181.55	129.44	483.09	350.59	483.79	304.94	421.64	437.94	0.418365908	0.0375	0.0005084
19069	myc induced nuclear antigen	198.52	93.12	89.6	515.36	223.46	260.41	415.61	176.74	570.73	0.352622889	0.03754	0.01215379
22250	karyopherin (importin) alpha 1	487.51	288.58	322.38	647.95	625.2	643.16	953.35	982.64	857.55	0.466456469	0.03755	0.00191203
20912	ribosomal protein S24	1503.87	981.91	1382.51	1903.5	1484.96	1825.7	2662.36	2502.71	2852.52	0.584700379	0.03757	0.01797688
40811	gem (nuclear organelle) associated protein 5	66.66	15.96	108.86	341.52	213.42	149.85	199.83	127.31	288.52	0.290022341	0.0376	0.00628726
19408	RIKEN cDNA 1810020D17 gene	70.67	116.83	133.45	300.46	275.02	220.85	330.43	391.71	361.99	0.341352648	0.03761	0.00033814
16317	GM2 ganglioside activator protein	1460.32	965.39	1158.55	1585.9	1681.36	1877.4	2813.6	2472.1	2183.15	0.568322595	0.03761	0.00878651
15388	RIKEN cDNA 2610042O14 gene	385.77	161.98	351.7	704.56	394.83	574.8	563.17	1018.26	1190.97	0.4045572	0.03763	0.01369527
18450	brix domain containing 1	368.46	490.29	890	1496.2	1126.8	1048.1	945.53	776.39	1481.18	0.508783516	0.03764	0.01101669
72	oncogene family	79.2	22.75	80.1	172.14	152.77	222.72	326.45	170.57	344.7	0.262064994	0.03765	0.00211936
22384	DEAD (Asp-Glu-Ala-Asp) box polypeptide 54	319.22	262.39	315.65	522.53	433.98	484.6	736.35	779.63	1015.23	0.451756153	0.03768	0.00631429

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
32708	cDNA sequence BC068171	1424.91	1570.38	1471.39	2308.1	2391.78	2968.3	2287.82	2852.25	2449.75	0.585484655	0.03768	0.00081003
13654	ubiquitin-conjugating enzyme E2O	656.86	318.47	383.2	924.15	728.47	782.62	1092.38	932.79	1270.08	0.474140955	0.03769	0.00312346
15701	proteasome (prosome, macropain) subunit, beta type 4	711.54	339.67	377.73	882.09	954.28	936.36	747.49	930.36	1311.81	0.495953936	0.0377	0.00367955
43187	jumonji domain containing 1B	13.53	5.03	18.14	102.01	28.28	55.5	131.11	118.55	104.24	0.136004002	0.0377	0.00131957
44702	zinc finger protein 407	25.36	30.71	38.29	227.66	134.41	183.53	100.04	82.93	152.65	0.214157645	0.03772	0.00029598
14551	ribosomal protein S24	1773.15	1158.27	1610.42	2156.3	1692.91	2124.5	2913.41	2832.19	3114.02	0.61238471	0.03774	0.0206946
379	zinc finger, A20 domain containing 2	1713.11	1445.5	1218.63	2269.6	2150.04	2173.5	2659.81	2431.14	3458.7	0.578127838	0.03774	0.00444974
20984	splicing factor, arginine/serine-rich 1 (ASF/SF2)	2351.06	1787.06	1796.58	3052.8	2627.08	2864.6	3491.69	3251.35	3629.06	0.627461802	0.03775	0.00349604
17472	amine oxidase, copper containing 3	69.46	19.83	7.36	85.43	273.68	193.82	107.32	107.65	335.47	0.175190553	0.03778	0.00363781
29573	RIKEN cDNA 4930535B03 gene	86.18	17.31	12.72	377.01	80.79	145.39	180.67	103.3	181.19	0.217550428	0.03778	0.00481092
738	ubiquitin-like 1 (sentrin) activating enzyme E1A	239.75	189.44	212.92	490.87	339.53	437.9	482.8	633.06	755.85	0.408985959	0.03779	0.00153851
8834	solute carrier family 39 (metal ion transporter), member 6	926.32	580.02	618.52	917.82	1152.33	1077.5	1430.48	1479.23	1557.51	0.558085418	0.0378	0.00688556
3709	guanine nucleotide binding protein 13, gamma	60.13	20.72	5.19	102.97	138.52	82.41	150.51	146.09	148.15	0.223873024	0.0378	0.00231997
36724	interferon-induced protein 35	1116.96	271.9	305	985.49	1595.47	746.92	620.37	1621.46	1471.57	0.48112275	0.03782	0.03491781
240	minichromosome maintenance deficient 5, cell division cycle	3.51	116.93	27.29	275.96	137.29	122.81	381.1	145.29	202.31	0.233609539	0.03783	0.00879308
12377	p53 and DNA damage regulated 1	725.38	231.39	442.6	715.57	694.98	822.93	1022.53	1047.48	1095.64	0.5183687	0.03784	0.01170324
12147	scaffold attachment factor B2	121.67	55.05	141.82	202.46	205.91	445.2	256.47	382.54	338.31	0.34796192	0.03785	0.00328657
11762	activating transcription factor 7 interacting protein 2	225.98	243.54	179.98	629.85	458.47	542.77	416.46	320.25	530.93	0.448127283	0.03785	0.00146075
1985	protein kinase, AMP-activated, gamma 1 non-catalytic subunit	41.91	31.4	14.25	149.26	132.45	140.8	117.68	165.53	217.77	0.189628475	0.03786	9.79E-005
9245	solute carrier family 1 (neutral amino acid transporter), member 5	10.9	13.02	15.54	68.86	61.47	39.47	135.13	217.53	154.55	0.116571395	0.03787	0.00095526
12998	splicing factor, arginine/serine-rich 1 (ASF/SF2)	665.47	668.28	445.98	808.48	849.06	987.04	1495.44	1250.6	1317.15	0.530647294	0.03789	0.00796067

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
22322	drebrin-like	16.85	15.82	9.22	61.28	99.35	52.2	236.1	101.96	127.64	0.123472801	0.0379	0.00042543
23370	RNA binding motif protein 3	39.43	20.38	13.54	161.7	136.81	113.63	64.65	102.43	161.88	0.197948995	0.03791	0.00037432
15912	MBD2-interacting zinc finger	391.59	235.52	209.12	595.63	489.39	563.7	600.44	667.79	618.68	0.473030266	0.03793	0.0007328
21512	makorin, ring finger protein, 1	1072.98	807.82	1014.98	1919.8	1380.03	1377.3	1555.19	1824.23	2003.22	0.575718958	0.03793	0.00309783
18256	cofactor required for Spl transcriptional activation, subunit 2	478.92	496.91	461.96	789.66	829.56	748.74	1285.92	1020.77	1061.21	0.501333715	0.03794	0.0015605
18900	GTP binding protein 4	1225.17	664.56	760.91	1492.5	1158.36	1447.9	2173.69	1808.18	1582.59	0.548608423	0.03795	0.00801715
29040	expressed sequence AA673488	484.11	218.8	155.21	630.18	425.47	677.11	680.05	395.77	933.82	0.458593416	0.03796	0.01368159
5392	Pbx/knotted 1 homeobox	136.46	110.17	38.04	161.58	245.89	197.24	309.66	326.55	315.1	0.36589504	0.03796	0.00483993
21315	importin 7	1897.13	1382.87	1447.66	2615.2	2037.98	2474.9	2454.84	2754.77	2633.6	0.631563479	0.03798	0.00311793
29563	cDNA sequence BC068171	663.48	885.66	893.34	1434	1361.4	1572.1	1455.55	1492.34	1452.45	0.557143297	0.038	0.00034246
15723	centromere protein E	638.18	667.92	635.85	1062.6	840.39	1209.8	1570.6	1201.87	1244.4	0.544756305	0.03801	0.00298774
14383	eukaryotic translation initiation factor 5B	20.96	15.05	9.7	81.27	106.34	102.48	124.36	76.5	147.21	0.14325561	0.03802	1.95E-005
10833	eukaryotic translation initiation factor 3, subunit 9 (eta)	88.55	23.79	36.88	226.1	76.88	140.2	208.42	168.48	203.88	0.291456698	0.03803	0.00358346
657	FK506 binding protein 4	880.01	451.63	640.66	1073.2	846.29	1012.7	1672.69	1558.58	1505.66	0.514345769	0.03803	0.01132742
30138	anti-Mullerian hormone type 2 receptor	144.16	58.14	204.59	277.7	226.3	364.45	492.22	340.87	636.14	0.34811437	0.03803	0.00675705
20815	phosphatidylinositol transfer protein, cytoplasmic 1	80.4	58.16	113.91	393.06	254.86	712.53	266.43	91.96	256.22	0.255658056	0.03804	0.01559053
6603	integrin alpha 6	593.83	573.74	427.75	925.3	1185.7	1056.8	971.42	1090.26	915.58	0.519219472	0.03807	0.0004221
12382	EBNA1 binding protein 2	939.3	650.84	646.44	1300.7	865.58	1123.9	1609.43	1404.18	1650.05	0.562392505	0.038	0.01050595
5650	transmembrane protein 49	178.72	95.35	104.83	304.43	188.62	201.94	473.19	397.83	447.36	0.376383874	0.038	0.00693338
16338	proteasome (prosome, macropain) subunit, beta type 2	230.23	131.35	156.94	402.72	255.26	414.95	581.61	681.86	363.66	0.384080354	0.03814	0.00393859
9656	PRP4 pre-mRNA processing factor 4 homolog B (yeast)	299.22	298.4	149.09	561.48	546.09	598.99	669.75	360.25	512.38	0.459669923	0.03815	0.00313005
21930	transcription factor-like 5 (basic helix-loop-helix)	21.09	38.28	22.9	250.9	115.79	62.22	136.62	119.48	296.43	0.167649906	0.03819	0.00144596
21395	guanine nucleotide binding protein (G protein), beta polypeptide 2 like 1	226.37	103.26	158.36	279.05	215.92	341.37	489.8	485.86	609.01	0.403129272	0.0382	0.01001628
484	Sec61 alpha 1 subunit (S. cerevisiae)	2767.22	1784.91	2139.48	3540.8	2633.86	3422.7	3549.86	3286.01	3718.68	0.664131518	0.03822	0.00962031

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
41081	RIKEN cDNA 2700023E23 gene	383.97	263.74	193.02	576.37	466.4	578.03	626.96	633.72	924.84	0.441754766	0.03826	0.00216237
22751	RE1-silencing transcription factor	2083.8	1792.43	1660.4	2301.8	2518.88	2548.3	3491.19	3525.95	3353.07	0.624228963	0.03827	0.00982946
6782	Ribosomal protein L41	2170.73	1455.5	2121.76	2890.9	2673.78	2954.3	3287.73	3064.33	3459.93	0.627133956	0.03827	0.00351568
23799	PRP4 pre-mRNA processing factor 4 homolog (yeast)	139.76	105.45	114.11	272.96	233.13	303.29	299.62	458.97	563.81	0.337107957	0.03828	0.00127622
5220	guanylate cyclase activator 1a (retina)	1137.82	874.49	1002.71	1303.4	1212.1	1554.2	2394.16	2176.13	2314.5	0.550467392	0.03829	0.01816163
22682	gb:BB204543 /DB_XREF=gi:8869496 /DB_XREF=BB204543 /CLONE=A430058H10 /FEA=EST /CNT=1 /TID=Mm.215042.2 /TIER=ConsEnd /STK=0 /UG=Mm.215042 /UG_TITLE=ESTs	101.6	39.07	67.86	262.21	64.78	123.41	308.26	237.22	438.68	0.290723288	0.03833	0.02815887
7211	ADP-ribosylation factor 4	228.83	119.48	108.04	327.09	279.11	307.49	457.63	425.84	551.85	0.388546664	0.03833	0.00201733
2543	galactosidase, alpha	273.94	147.47	118.3	418.22	369.57	248.33	523.01	438.32	654.08	0.407093263	0.03834	0.00570311
19658	cDNA sequence BC021395	585.54	403.84	321.3	781.58	616.26	679.87	1012.69	951.38	1013.92	0.518495955	0.03834	0.00493793
27841	BTAF1 RNA polymerase II, B- TFIID transcription factor- associated, (Mot1 homolog, S. cerevisiae)	28.85	6.86	11.93	42.58	39.42	232.36	46.01	239.9	187.23	0.120990476	0.03834	0.00833485
30576	Transcribed locus	510.18	377.24	438.04	946.62	823.44	1198.9	651.44	766.79	926.47	0.498886833	0.0384	0.00188992
29457	WD repeat domain 46	218.17	112.46	94.73	408.68	196.3	309.16	631.99	252.43	611.74	0.352948986	0.0384	0.01177698
19166	solute carrier family 39 (zinc transporter), member 4	46.7	42.22	31.16	170.36	69.43	62.77	295.72	198.68	333.58	0.212429458	0.03846	0.01086785
43448	TatD DNase domain containing 1	77.05	76.25	15.42	213.53	120.79	170.2	248.59	156.85	449.66	0.248186993	0.03846	0.00550496
6729	YY1 transcription factor	283.2	139.65	276.18	630.7	488.3	503.98	593.02	531.08	482.62	0.432880971	0.03853	0.00073673
3998	procollagen, type V, alpha 3	33.3	2.46	6.94	108.03	51.77	59.8	243.66	175.36	95.74	0.11628857	0.03853	0.00187396
14394	acetyltransferase 2	315.87	400.24	148.72	745.68	497.68	1107.4	690.17	601.4	547.17	0.412858906	0.03864	0.00534374
4785	RIKEN cDNA 2410016F19 gene	2626.86	1700.7	1801.16	2574.1	2660.57	2995.8	4101.89	3403.46	3437.85	0.639284686	0.03864	0.01416184
19007	GTP binding protein 3	315.42	126.56	117.97	465.37	336.42	416.21	408.64	480.29	597.69	0.414069259	0.03872	0.00246244

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	tpvals
7933	protein regulator of cytokinesis 1	717.27	432.74	439.28	852.9	910.14	799.31	1102.98	1075.94	1372.07	0.519941636	0.03905	0.0038711
18762	RIKEN cDNA 1810003N24 gene	844.39	363.6	396.78	978.22	773.69	906.85	985.65	1181.43	1314.69	0.522681267	0.0394	0.00727294
10372	IMP4, U3 small nucleolar ribonucleoprotein, homolog (yeast)	1079.23	720.29	726.16	1229.9	1160.97	1507.8	1381.36	1559.21	2228.6	0.557065661	0.03946	0.00884479
41173	gb:BB209183 /DB_XREF=gi:8874136 /DB_XREF=BB209183 /CLONE=A430091G17 /FEA=EST /CNT=18 /TID=Mm.129698.1 /TIER=Stack /STK=17 /UG=Mm.129698 /UG_TITLE=ESTs	496.81	534.73	672.17	964.31	1013.34	1068.5	823.07	1042.47	1568.62	0.525809642	0.03951	0.00333501
15338	RAB20, member RAS oncogene family	486.8	419.24	302.02	593.19	602.78	611.39	1207.04	970.25	1217.61	0.46443661	0.03951	0.01195215
15902	Crx opposite strand transcript 1	216.2	196.78	180.31	438.71	365.54	284.88	517.72	489.36	491.37	0.458567465	0.03955	0.00125293
13638	ribosomal protein S21	2584.81	1818.73	2032.86	3125.7	2859.55	3137.7	3666.04	3359.24	5278.12	0.600791827	0.03957	0.0123343
7634	cyclin M2	191.94	160.39	185.32	529.15	281.25	348.56	414.53	418.85	581.74	0.417741484	0.03961	0.00131644
3158	GATA binding protein 4	243.29	314.72	216.04	548.17	372.29	460.72	660.84	563.02	829.97	0.450682822	0.03963	0.00324729
14053	denticless homolog (Drosophila)	117.57	78.14	95.78	349.89	205.74	183.08	336.21	206.1	338.68	0.359930851	0.03967	0.0011816
21575	tubulin, beta 5	38.96	56.74	38.01	155.85	185.38	153.18	142.88	165.48	195.42	0.267904908	0.03967	1.01E-005
1352	peptidylprolyl isomerase D (cyclophilin D) /// lysosomal-associated membrane protein 3 /// similar to peptidylprolyl isomerase D	707.11	447.21	263.78	1140.3	699.01	928.31	1030.22	754.1	902.76	0.519960034	0.03969	0.00795448
17069	autophagy-related 3 (yeast)	314.5	241.22	433.46	642.61	635.28	677.38	1423.01	439.15	1211.18	0.393420846	0.03973	0.01356134
20883	RIKEN cDNA 2900053A13 gene	46.21	35.31	27.79	124.96	115.95	155.72	172.07	129.07	221.68	0.237772581	0.03978	6.40E-005
12626	dipeptidase 3	414.84	426.47	469.53	773.35	660.61	710.24	1012.53	968.29	930.73	0.518554122	0.03979	0.00111782
2421	zinc finger protein 292	14.9	34.16	12.24	99.79	76.57	80.72	137	132.44	131.78	0.186237278	0.03985	0.00012921
10640	kelch-like 22 (Drosophila)	63.35	36.5	154.28	373.75	196.24	394.95	84.02	140.12	513.75	0.29847959	0.03988	0.03136841
5230	von Hippel-Lindau syndrome homolog	242.32	87.46	145.37	415.71	238.33	239.9	369.93	417.03	500.62	0.435613701	0.03994	0.00838939

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
12058	ring finger protein (C3H2C3 type) 6	554.18	420.14	401.07	655.12	583.25	868.77	1096.24	908.12	1006.93	0.537426516	0.03997	0.0070928
10617	sarcolemma associated protein	102.17	105.27	51.27	194.14	150.33	224.98	300.15	333.64	268.15	0.351653878	0.04	0.00175258
39100	nucleolar protein 9	340.36	166.21	323.61	458.72	342.33	403.84	935.22	768.76	875.51	0.438740296	0.04	0.02347893
10978	homeodomain interacting protein kinase 3	11.81	29.62	9.18	120.44	57.72	112.28	96.61	68.27	152.13	0.166630998	0.04003	0.00030594
13562	cDNA sequence BC010304	1318.59	952.75	1281.75	1970.4	1624.18	2021.9	1486.9	2242.25	2206.77	0.615126394	0.04003	0.00682695
7422	H3 histone, family 3A	878.81	631.98	853.13	1174.7	1128.36	1357.7	1349.45	1593.59	2334.29	0.528957723	0.04006	0.00865031
16942	SUMO/sentrin specific peptidase 3	2271.54	1602.51	1721.79	2780.1	2659.11	2911.5	2929.1	2911.41	3250.48	0.641664431	0.04006	0.00238324
21171	ribosomal protein L29 /// hypothetical protein LOC669999 /// similar to 60S ribosomal protein L29	1696.19	634.99	1377.16	1987.6	2266.44	1883.6	1592.12	2337.53	4074.9	0.524436456	0.04009	0.02917422
18445	Yip1 domain family, member 4	782.63	357.97	422.77	943.49	888.23	831.02	1009.32	1081.25	1031.26	0.540531103	0.04011	0.00347794
23583	CXXC finger 6	166.09	59.38	58.14	251.54	278.86	292.88	369.76	167.47	233.69	0.355802283	0.04012	0.00239894
11104	RAN binding protein 5	1214.6	825.31	930.44	1439.4	1397.73	1332.6	1951.59	1977.64	2086.45	0.583253563	0.04015	0.00776708
20617	placenta specific 9	196.43	91.79	72.6	352.41	222.51	218.3	278.58	511.3	340.48	0.37515466	0.04017	0.00430094
2867	tumor necrosis factor receptor superfamily, member 12a	301.25	159.52	151.05	479.97	246.27	564	471.44	369.03	575.57	0.452148336	0.0402	0.00844923
16007	hypothetical LOC435970 /// similar to gonadotropin inducible ovarian transcription factor 2	678.45	317.43	184.75	688.24	630.48	750.43	1021.21	760.72	1173.03	0.469985729	0.04022	0.01029519
38800	Glutamine fructose-6-phosphate transaminase 1	154.6	86.66	72.39	201.4	144.57	299	425.52	262.9	388.13	0.364385169	0.04027	0.00670534
13746	deoxyhypusine synthase	40.12	11.88	30.86	40.89	74.65	90.18	175.94	258.3	230.11	0.190465245	0.04028	0.00951161
12807	peptidyl prolyl isomerase H /// hypothetical protein LOC624822 /// similar to Peptidyl-prolyl cis-trans isomerase H (PPIase H) (Rotamase H)	48.4	49.08	49.99	166.67	194.54	121.42	198.1	204.78	97.2	0.300129234	0.04032	0.00035827
13130	protein	102.04	67.67	139.68	333.55	151.31	287.34	230.45	274.04	374.05	0.374850067	0.04033	0.00283292
2362	cofilin 2, muscle	978.51	785.68	989.09	1233.6	1142.57	1301.9	2052.78	1861.5	1865.78	0.582208806	0.04038	0.01428345

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
6144	eukaryotic translation initiation factor 4E member 2	106.88	47.69	39.81	160	100.21	135.61	343.08	263.59	456.25	0.266503969	0.04038	0.00981135
14010	eukaryotic translation initiation factor 5B	630.83	605.07	661.68	984.87	922.28	1001.4	1288.76	1369.41	1283.35	0.554029868	0.04039	0.00170399
17195	ariadne homolog 2 (Drosophila)	1041.91	675.07	711.19	1533	934.55	1369.8	1554.44	1033.03	1685.49	0.598785251	0.04042	0.01816852
38383	mitochondrial ribosomal protein S6	1289.27	719.43	948.29	1789.4	1403.2	1559.5	1291.59	2038.19	1795.1	0.598760357	0.04043	0.00896259
16985	mitochondrial ribosomal protein L39	192.78	74.48	65.23	382.08	203.89	182.76	355.09	320.19	281.17	0.385455431	0.04046	0.00469527
2261	mitochondrial ribosomal protein L39	11.28	52.99	34.16	169.89	124.95	91.91	119.25	152.47	153.1	0.242566877	0.04049	0.00084464
39073	phosphatidylinositol 3-kinase, catalytic, beta polypeptide	76.55	96.39	113.73	271.48	148.58	263.12	289.7	255.76	314.46	0.371550774	0.0405	0.00077467
5910	proteasome (prosome, macropain) 26S subunit, non-ATPase, 14	774.34	476.98	629.32	1287.5	896.48	1249	1056.77	933.88	1219.72	0.566167074	0.0405	0.00346014
38875	RIKEN cDNA 1110008B24 gene	723.82	473.87	306.58	908.37	546.03	872.14	1023.54	1117.82	1343	0.517740797	0.04052	0.01875828
44707	expressed sequence AF013969 /// similar to CG5514-PB, isoform B /// region containing expressed sequence AF013969; RIKEN cDNA A230054D04 gene	143.57	109.96	108.78	314.91	203.78	285.29	503.61	242.43	417.29	0.36833036	0.04052	0.00212647
8041	ring finger and WD repeat domain 3	1896.6	1829.31	1380.91	2772	2658.21	3056.3	2072.36	2195.33	2804.08	0.656477432	0.04054	0.01044829
28154	methyltransferase like 2	51.31	48.28	45.52	144.77	183.34	169.06	101.31	165.93	286.78	0.276081849	0.04056	0.00042431
30269	nucleolar protein 11	29.76	6.59	11.32	94.82	98.86	59.84	176.47	125.5	88.54	0.148036582	0.04058	0.00033177
7189	valosin containing protein /// similar to Transitional endoplasmic reticulum ATPase (TER ATPase) (15S Mg(2+)-ATPase p97 subunit) (Valosin-containing protein) (VCP)	441.58	368.14	517.63	943.18	758.44	743.25	937.83	818.5	891.57	0.521267363	0.04058	0.0004249
16306	ubiquitin specific peptidase 10	257.9	263.88	466.98	838.32	498.36	592.08	738.05	449.15	994.08	0.481143736	0.04059	0.0102128

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
24333	activating transcription factor 7 interacting protein 2	998.36	673.13	860.25	1461.8	1354.64	1323.4	1579.88	1425.89	1629.02	0.577057117	0.0406	0.00102143
18466	host cell factor C1	217.35	291.13	173.2	369.82	317.79	306.44	711.47	663.73	675.05	0.447840226	0.04061	0.01723647
7354	hippocampus abundant gene transcript 1	392.76	231.65	210.52	490.55	398.57	461.1	662.98	750.7	838.3	0.463566709	0.04063	0.00703408
11196	eukaryotic translation initiation factor 4, gamma 1	795.82	790.01	913.29	1492.1	1297.24	1320.6	1521.77	1501.26	1366.38	0.588076665	0.04065	0.00040865
12088	pyridoxal (pyridoxine, vitamin B6) kinase	341.59	275.6	123.17	435.93	526.52	455.27	643.14	513.95	833.89	0.434394344	0.04067	0.0051941
955	eukaryotic translation initiation factor 3, subunit 10 (theta)	3497.53	2343.31	2333.08	3666.5	3233.57	3908.4	5268.73	4157.74	4381.74	0.664095243	0.0407	0.01798116
21792	Zinc finger protein 444	311.39	109.74	33.14	452.64	198.57	309.81	294.53	537.31	689.29	0.36602945	0.0407	0.01990646
2633	WD repeat domain 33	130.08	90.77	188.21	291.14	355.77	593.08	234.78	271.81	596.3	0.349194154	0.04072	0.00518948
16350	complement component 1, q subcomponent binding protein	1157.96	830.97	657.36	1662.1	1312.42	1366.6	1453.75	1506.64	2023.78	0.567554294	0.04074	0.00450461
16278	RIKEN cDNA 2610524G07 gene	445.79	163.3	359.19	701.15	580.47	1519.9	587.27	478.76	1412.36	0.366776895	0.04075	0.0187662
41375	core-binding factor, runt domain, alpha subunit 2, translocated to, 2 homolog (human)	362.76	159.53	39.4	549.54	295.88	488.83	219.51	745.72	448.33	0.408827393	0.04076	0.02903456
9888	vitelliform macular dystrophy 2-like protein 1	26.77	68.92	92.66	159.89	205.9	333.32	240.53	128.6	182.61	0.301155214	0.04079	0.0025478
12466	ribosomal protein S24	1501.15	1020.57	1349	1942.8	1498.52	1804.5	2346.82	2369.96	2625.8	0.614966159	0.04082	0.01377549
6758	zinc finger protein 143	70.96	68.45	69.67	183.06	92.13	284.13	176.81	250.89	303.16	0.324109814	0.04085	0.0039866
23587	GTPase activating protein and VPS9 domains 1	333.58	155.43	192.66	424.38	502.98	467.39	685.61	645.29	353.59	0.442752108	0.04086	0.00362749
12738	type 2B	125.53	76.26	48.2	253.24	181.84	235.81	167.98	240.57	313.04	0.359057222	0.04087	0.00119569
14480	Son cell proliferation protein	352.01	285.74	318.7	532.78	480.48	448.55	821.48	910.68	792.96	0.479792723	0.04088	0.00694346
11012	SET translocation	915.35	704.29	595.07	1259.9	1330.58	1160.7	1068.77	1370.79	1550.02	0.572220299	0.04088	0.00220896
2866	tumor necrosis factor receptor superfamily, member 12a	320.47	172.16	158.2	480.91	257.13	582.58	497.84	384.37	599.54	0.464485418	0.0409	0.00962175
38041	sorting nexin 6	279.12	83.74	111.89	319.18	284.78	562.5	346.75	388.99	594.39	0.380318755	0.0409	0.00500672
13631	protein tyrosine phosphatase-like A domain containing 1	221.52	140.45	439.19	494.52	549.29	538.97	573.99	612.57	1361.41	0.387900502	0.04091	0.00955865

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
40666	bromodomain and PHD finger containing, 3	123.25	66.62	84.15	270.72	241.5	263.39	137.62	214.11	397.32	0.359450632	0.04092	0.00275523
25023	cyclin M4	81.3	61.69	90.14	148.79	264.46	223.53	165.53	233.02	314.14	0.345513424	0.04095	0.00066368
15839	peroxisome proliferator activated receptor binding protein	3.63	2.38	10.41	65.58	38.61	88.28	38.4	118.58	104.52	0.072339582	0.04096	0.00013507
14494	DnaJ (Hsp40) homolog, subfamily B, member 12	183.48	163.99	62.66	294.14	274.02	311.49	605.82	320.93	536.74	0.350068711	0.04097	0.00463062
10865	xylulokinase homolog (H. influenzae)	570.91	332.39	251.87	818.91	532.81	562.41	921.59	711.51	946.94	0.514074901	0.04101	0.00971793
16446	Unc-51 like kinase 1 (C. elegans)	688.42	447.89	387.79	998.96	689.04	942.76	783.34	1059.4	1087.82	0.548107284	0.04102	0.00483918
12517	PRP38 pre-mRNA processing factor 38 (yeast) domain containing B	453.67	395.38	239.57	654.65	564.35	746.42	1167.03	624.92	806.21	0.477090355	0.04103	0.0054774
12358	DEAH (Asp-Glu-Ala-His) box polypeptide 40	72.27	61.52	54.09	167.12	149.31	199	299.91	175.4	477.91	0.255854016	0.04106	0.00156953
13777	general transcription factor III C 1	161.54	193.59	135.49	406.5	272.1	321.91	350.92	440.43	553.06	0.418453508	0.04108	0.00119698
1549	spermatogenesis associated 5	119.35	125.19	136.28	318.86	254.39	257.63	387.86	321.28	353.57	0.402220122	0.04111	0.000123
7228	activity-dependent neuroprotective protein	454.34	277.92	262.38	806.44	487.17	695.53	665.1	564.28	782.06	0.497247899	0.04113	0.00268655
17344	glutamine fructose-6-phosphate transaminase 1	374.81	257.95	316.1	699.57	464.92	769.86	656.51	363.82	746.9	0.51267837	0.04113	0.0111343
22765	purine rich element binding protein B	155.98	78.28	45.27	264.07	195.74	234.5	347.63	237.43	304.74	0.352917411	0.04118	0.00165563
2822	FUS interacting protein (serine-arginine rich) 1	2250.39	1979.57	1833.46	2877.6	2467.65	3105.4	3470.04	3149.84	3432.49	0.65539575	0.04119	0.00493355
29854	ADP-ribosylation factor-like 15	102.07	159.07	161.91	336.3	397.43	326.19	521.65	339.7	306.09	0.379866748	0.04131	0.00035486
14013	DEAD (Asp-Glu-Ala-Asp) box polypeptide 52	388.08	161.92	274.42	654.08	454.68	714	468.82	637.43	504.22	0.480259115	0.0414	0.00404719
44759	expressed sequence AA763515	8.75	107.57	93.59	266.69	476.12	367.11	115.8	208.23	76.04	0.278028331	0.04145	0.03222834
25723	nucleolar protein 9	814.25	472.11	250.05	1027.7	836.53	843.91	990.85	1023.2	913.87	0.545203243	0.04161	0.00953711
19626	sestrin 2	492.49	377.96	378.76	576.84	439.85	945.27	773.15	957.31	978.1	0.534934012	0.04167	0.02023325
17889	phosphatidylinositol 4-kinase type 2 beta	199.31	230.48	234.25	466.28	327.46	508.52	589.11	402.49	715.19	0.441361892	0.04173	0.0022702

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
24760	RIKEN cDNA 4933402C05 gene	526.44	324.33	277.42	582.5	656.43	516.22	821.01	823.62	1015	0.511096816	0.04177	0.00786959
40677	RIKEN cDNA D530033C11 gene	62.8	31.87	92.29	229.92	169.71	276.49	176.84	152.21	205.02	0.308976276	0.04182	0.00070361
40756	cDNA sequence BC057371	266.3	343.51	142.19	489.41	461.09	497.84	765.56	698.43	524.25	0.437644402	0.04183	0.00285669
15702	ribosomal protein S17	3413.68	2189.66	2823.31	4101	3483.88	3592.5	4611.75	4168.53	4913.62	0.677620117	0.04188	0.01435464
16556	nuclear import 7 homolog (S. cerevisiae)	1117.61	701.46	924.84	1468.6	1454.75	1208.1	1615.11	1894.95	1641.93	0.591138237	0.04193	0.00457958
11703	poly (A) polymerase alpha	15.3	45.46	34.26	129.02	144.67	119.25	158.08	76.86	140.95	0.247180781	0.04198	0.0004212
5457	homeodomain interacting protein kinase 1	982.38	849.34	1301	1736.4	1266.8	1727.8	1609.86	1888.44	2141.79	0.604126672	0.04203	0.00752055
21337	carboxypeptidase D	52.84	88.41	40.22	258.05	184.47	187.52	155.3	168.61	253.58	0.300563961	0.04209	0.00026393
21183	ubiquitin associated protein 2-like	704.73	268.76	278.2	1175.1	736.68	781.12	765.49	711.76	1019.45	0.48238121	0.04214	0.00655066
16501	eukaryotic translation initiation factor 3, subunit 10 (theta)	1962.76	1345.41	1507.34	2379.6	1854.13	2427.1	3453.2	2549.07	3057.45	0.612638481	0.04218	0.01453233
791	Morf4 family associated protein 1	1393.96	988.45	1124.79	1873	1359.87	1748.1	1989.43	2140.18	2023.47	0.629994243	0.04224	0.00786029
9218	protein arginine N-methyltransferase 6	1192.36	520.85	617.17	1347.3	1072.8	1130.9	1415.39	1423.29	1354.94	0.601806934	0.04277	0.01108551
39489	RIKEN cDNA 2610206G21 gene	385.68	426.03	475.42	734.42	556.38	887.02	879.79	778.82	850.91	0.549194212	0.04282	0.00202463
14334	WD repeats and SOF domain containing 1	431.5	295.6	235.2	806.65	558.06	634.33	784.95	603.44	597.88	0.482923537	0.04287	0.00128243
24209	RIKEN cDNA 2610020H08 gene	32.11	23.32	52.07	145.28	75.19	165.2	243.85	127.48	115.24	0.24649179	0.04302	0.00099377
20784	DEAD (Asp-Glu-Ala-Asp) box polypeptide 10	70.39	42.43	116.81	234.66	140.44	219.65	270.65	147.92	253.11	0.36264144	0.04305	0.00237097
7934	protein regulator of cytokinesis 1	882.29	561.67	562.14	1130.7	990.99	915.08	1248.9	1220.25	1392.55	0.581610598	0.04307	0.00476743
8707	zinc finger, CCCH-type with G patch domain	51.69	25.13	56.62	118.43	68.27	140.57	216.62	178.1	256.	0.272855536	0.043	0.00395268
21424	ubiquitin specific peptidase 36	186.19	106.7	198.3	424.4	291.16	431.35	322.19	484.24	311.35	0.433790055	0.04312	0.001588
7817	signal peptide peptidase 3	87.29	42.07	37.9	188.55	124.22	96.45	188.49	216.63	182.0	0.33574547	0.04317	0.00179556
2217	stathmin-like 3	280.8	139.85	201.65	475.25	288.43	374.17	564.1	475.36	454.23	0.472953202	0.0432	0.00392245

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
20309	cofactor required for Sp1 transcriptional activation, subunit 7	401.97	260.33	241.18	724.75	562.78	486.97	470.51	624.15	970.19	0.470642166	0.04322	0.00490214
1349	RIKEN cDNA 0610009D07 gene	161.97	142.69	83.83	332.41	184.17	281.59	388.71	302.32	482.73	0.394020072	0.04326	0.00393886
10566	CUG triplet repeat, RNA binding protein 1	368.59	320.29	329.52	818.02	671.83	693.42	734.46	490.43	505.7	0.520406964	0.04332	0.00216732
8579	cell cycle progression 1	220.05	202.83	244.27	519.44	428.94	471.56	477.1	413	524.9	0.470662518	0.04333	8.49E-005
15916	BTB and CNC homology 1	299.52	142.08	280.99	509.03	500.69	370.86	552.06	427.69	703.89	0.471630627	0.04338	0.00429197
4645	nucleolar protein 7	551.77	243.19	360.68	894.12	728.94	555.6	772.7	534.61	874.95	0.529998257	0.04348	0.00928765
29317	NIMA (never in mitosis gene a)-related expressed kinase 2	692.79	642.8	418.9	917.15	746.99	927.21	1171.25	1253.89	1243.99	0.560496959	0.04353	0.00770205
11419	tropomyosin 3, gamma	574.31	530.72	522.22	1153.2	848.47	1058.4	816.32	879.3	1247.36	0.542146497	0.04358	0.0016093
1594	NIMA (never in mitosis gene a)-related expressed kinase 2	750.07	650.55	453.52	1026.9	842.77	1001	1176.28	1304.61	1168.12	0.568777283	0.04363	0.00385126
23285	RIKEN cDNA 1110004B13 gene	32.86	16.77	16.39	47.07	45.2	114.19	120.64	162.14	140.35	0.209723788	0.04369	0.00294454
8141	FUS interacting protein (serine-arginine rich) 1	657.7	606.69	401.25	1289.5	963.26	940.8	1005.18	1015.79	962.73	0.539283789	0.04379	0.0014819
694	heme oxygenase (decycling) 2	228.15	127.34	175.71	429.17	306.91	318.12	522.16	364.93	537.19	0.428649818	0.0438	0.0018308
21921	WD repeat domain 33	423.94	236.64	225.43	650.55	442.71	521.71	588.06	600.77	812.98	0.489944094	0.04384	0.0034789
23670	BRF2, subunit of RNA polymerase III transcription initiation factor, BRF1-like	101.79	131.02	195.54	374.57	304.79	715.04	381.23	100.11	630.97	0.341762709	0.04385	0.05104204
29502	transmembrane protein 39b	176.13	81.2	214.01	384.29	398.39	485.44	203.27	287.14	367.76	0.443344981	0.04389	0.00918906
17866	caspase 3	628.36	670.27	458.61	1113.9	883.27	1268.1	856.19	1204.37	987.32	0.556689339	0.0439	0.00269251
21277	ribosomal protein L23	4511.5	3680.37	3884.36	5456.7	5071.7	5720.7	6020.5	5596.08	6154.6	0.709944642	0.04396	0.00538158
6830	N-acetylated alpha-linked acidic dipeptidase 2	977.43	735.87	1119.53	1576.5	1513.42	1599.2	1649.89	1277.79	1688.28	0.608881433	0.0440	0.0026034
14469	growth arrest specific 2	296.03	215.2	25.66	413.0	244.06	357.68	862.42	415.16	513.27	0.382727402	0.044	0.02873344
12046	RIKEN cDNA 2610304G08 gene	258.49	114.73	87.48	371.57	264.27	368.45	247.09	401.4	444.6	0.439307902	0.04412	0.00518255
688	EMG1 nucleolar protein homolog (S. cerevisiae)	747.94	398.7	373.77	892.83	714.84	908.6	952.79	986.65	1139.24	0.543492715	0.04414	0.00487152
12472	phosphoribosyl pyrophosphate amidotransferase	1491.3	1292.29	865.49	1886.3	1875.57	1977.5	2339.73	2006.58	1731.57	0.617588445	0.04415	0.005014
6656	solute carrier family 30 (zinc transporter), member 5	297.02	228.45	223.91	509.29	464.77	462.7	542.34	536.0	601.37	0.480914365	0.04415	0.00017236

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
20828	survival motor neuron domain containing 1	1251.34	889.9	1044.49	1444.9	1548.98	1671.9	1831.75	1993.53	1836.28	0.616951332	0.04419	0.00303399
1798	replication factor C (activator 1) 2	112.63	81.23	219.08	447.59	301.17	360.79	232.07	199.2	517.32	0.401274938	0.0442	0.00918665
15103	polymerase (RNA) III (DNA directed) polypeptide A	722.68	439.95	880.64	968.12	883.9	1401.1	997.47	1601.38	1547.52	0.552273944	0.04421	0.01694181
25102	Protein phosphatase 2 (formerly 2A), catalytic subunit, beta isoform	174.77	85.22	66.46	204.94	186.65	182.56	357.01	457.2	468.36	0.351641605	0.04424	0.00971819
11003	programmed cell death 2-like	554.47	517.02	316.42	881.18	628.34	852.95	854.94	1019.81	858.95	0.544687481	0.04425	0.00361332
1176	myeloid cell leukemia sequence 1	967.2	721.92	533.68	1440.5	1391.54	1486.5	898.34	715.94	1658.09	0.585653235	0.04426	0.04611204
519	zinc finger and BTB domain containing 17	53.35	146.21	217.39	394.13	287.82	382.18	413.55	194.42	1009.27	0.310997736	0.04435	0.01661823
7589	MYB binding protein (P160) 1a	627.85	220.85	196.3	619.81	559.46	543.97	759.99	756.52	1180.53	0.472820726	0.04436	0.01351934
15255	hexosaminidase B	452.07	322.98	329.02	829.4	691.76	686.04	625.54	661.85	990.61	0.492316954	0.04441	0.00102269
13783	5,10-methylenetetrahydrofolate reductase	92.68	23.2	61.28	119.07	236.68	128.56	224.28	167.49	212.1	0.325607896	0.04442	0.0031695
22165	ribosomal protein S23	1550.31	917.07	1521.18	2053.3	1593.83	1798.2	2341.16	2253.28	2552.84	0.633474792	0.04447	0.01650885
1699	ELOVL family member 6, elongation of long chain fatty acids (yeast)	123.87	55.45	27.87	238.32	107.52	110.47	373.69	190.73	287.91	0.316649346	0.04452	0.01238244
30087	RIKEN cDNA 4930515G01 gene	30.95	26.16	15.9	110.2	64.69	112.74	138.13	116.5	168.78	0.205361161	0.04458	0.00016379
964	ankyrin repeat and MYND domain containing 2	67.44	57.45	75.7	180.49	149.85	218.49	301.23	86.12	363.81	0.308602374	0.04462	0.007918
12034	RIKEN cDNA 1100001I22 gene	632	348.59	642.76	973.51	834.59	1122.8	900.12	946.7	1813.9	0.49254739	0.04463	0.00948749
6044	Son of sevenless homolog 1 (Drosophila)	230.34	128.95	112.0	438.22	273.17	373.78	444.95	294.97	346.44	0.43407183	0.04468	0.00166919
3604	podoplanin	129.35	118.81	88.89	294.23	216.59	222.67	344.13	264.7	548.92	0.356432817	0.04469	0.00216879
2245	heterogeneous nuclear ribonucleoprotein K	1641.05	1528.27	1311.2	2707.7	2502.94	2859.9	1831.48	2175.84	1879.69	0.642018685	0.04473	0.01239415
8159	ribosomal protein S11	1905.1	755.25	1126.75	2116.3	1573.32	1611.4	2104.2	2067.46	2512.79	0.631950714	0.04489	0.02807283
19749	SET and MYND domain containing 5	83.2	60.13	96.49	253.0	152.74	155.94	279.3	195.6	293.15	0.36069938	0.04494	0.00109287

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
18114	nucleolar and coiled-body phosphoprotein 1	447.22	412.6	479.77	809.13	599.42	798.12	1127.99	790.66	1288.01	0.494922719	0.04504	0.00517173
21966	ring finger protein 7	978.6	679.37	530.73	1504.8	910.25	1298.1	1164.47	1209.11	1470.39	0.579244127	0.04509	0.00830564
33824	Rap1 interacting factor 1 homolog (yeast) /// similar to Telomere-associated protein RIF1 (Rap1-interacting factor 1 homolog) (mRif1)	1336.02	1290.49	1278.51	1985.5	1725.98	2113.1	2351.85	1874.39	2319.62	0.631344433	0.04515	0.00227917
14479	solute carrier family 5 (sodium-dependent vitamin transporter), member 6	1025.49	706.6	869.51	1379.5	1274.12	1219.6	1518.78	1593.95	1642.94	0.602996207	0.0452	0.00271465
19667	RIKEN cDNA 1500011J06 gene	2082.92	1586.47	1424.05	2592.1	2138.83	2584.9	3671.38	2264.15	2862.89	0.632167509	0.04561	0.01397112
21211	protein tyrosine phosphatase-like A domain containing 1	488.96	206.07	471.03	602.68	571.37	650.06	818.59	878.21	1244.5	0.489384964	0.04566	0.01423314
20126	expressed sequence AA408296	1123.36	793.87	596.32	1599.9	1397.86	1490.4	981.23	1241.52	1475.94	0.614046445	0.04591	0.0132658
20457	splicing factor, arginine/serine-rich 1 (ASF/SF2)	2421.27	1796.6	1659.9	2947.2	2521.55	2808.5	3148.99	2942.53	3230.58	0.667956494	0.04596	0.00568803
5056	synaptosomal-associated protein 23	749.26	643.08	613.45	1161.8	1012.37	1087.2	1424.69	1123.3	1276.51	0.566141158	0.04602	0.00087345
23156	peptidylprolyl isomerase (cyclophilin)-like 1	304.9	357.27	348.55	845.83	673.8	668.59	562.18	556.37	635.42	0.51277082	0.04607	0.00062062
14444	AT rich interactive domain 4B (Rbp1 like)	54.11	86.93	51.53	199.78	154.66	258.1	146.18	272.78	173.72	0.319559914	0.04675	0.00042303
979	fibrillarin	527.33	324.24	338.18	693.11	444.03	799.63	1106.82	681.18	990.6	0.50462636	0.04681	0.01468203
14590	RIKEN cDNA 4732471D19 gene	51.05	36.62	123.36	196.47	132.63	198.82	195.08	238.56	382.09	0.314114539	0.04688	0.0029813
2168	PWP1 homolog (S. cerevisiae)	253.54	123.82	124.06	398.21	231.41	365.55	410.58	364.75	416.54	0.458537567	0.04691	0.00338099
11080	ATP-binding cassette, sub-family F (GCN20), member 1	147.69	158.88	142.97	345.18	164.53	215.87	707.42	447.31	892.23	0.324280263	0.04693	0.03515947
19751	ankyrin repeat and MYND domain containing 2	3.72	28.84	6.61	112.14	69.74	61.26	111.73	38.77	117.79	0.153178343	0.04696	0.00113502
14756	Kruppel-like factor 9	210.02	29.37	90.94	354.82	272.53	212.2	254.78	274.03	358.42	0.382596509	0.04698	0.00747019
15073	YTH domain family 1	215.69	353.03	190.75	634.54	438.28	329.92	608.02	420.79	788.44	0.471721962	0.04701	0.01026759
27459	RNA pseudouridylate synthase domain containing 3	35.48	20.56	13.51	94.07	113.99	134.1	103.42	130.55	442.34	0.136577415	0.04707	0.00101169
182	Trk-fused gene	784.41	340.53	295.58	928.5	564.98	679.38	871.81	1022.58	1079.02	0.552058093	0.04717	0.02052607

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
16390	cell division cycle 2 homolog A (S. pombe)	1520.87	991.88	1873.5	1922.5	2470.74	2908.9	2019.67	2007.39	3556.92	0.589307355	0.0472	0.02144756
2140	claudin 6	377.34	362.31	248.66	836.91	293.71	421.11	822.38	709.62	736.86	0.517359884	0.04722	0.04780535
23988	RIKEN cDNA 2210011C24 gene	72.37	81.79	46.68	176.2	192.27	212.67	138.15	189.52	394.55	0.308188068	0.04725	0.00136374
14006	La ribonucleoprotein domain family, member 5	423.27	344.14	388.76	594.63	701.46	735.35	726.67	737.66	830.55	0.534481962	0.04727	0.00036524
8270	insulin-like growth factor 2 receptor	434.23	266.75	433.37	781.98	500.48	705.3	745.85	596	939.69	0.53139859	0.04738	0.0059995
15878	zinc finger protein 289	81.46	122.04	76.07	252.08	206.46	182.53	595.85	231.71	743.7	0.252738063	0.04741	0.00831163
27919	casein kinase 1, epsilon	10.44	8.6	20.89	54.93	27.16	86.67	149.6	94.28	120.88	0.14968511	0.04752	0.00167154
3281	ubiquitously expressed transcript	138.54	90.92	30.95	203.48	182.02	211.77	247.35	247.04	374.41	0.355249067	0.04754	0.0043143
16269	RIKEN cDNA 5730403B10 gene	77.47	94.8	164.04	275.16	166.89	142.51	330.15	361.39	315.31	0.422656638	0.04759	0.01185988
13417	IBR domain containing 3	1392.62	1504.44	1813.39	2414	2004.88	2264.8	2704.73	2941.42	2367	0.641018401	0.04762	0.00504366
23624	protein phosphatase 1A, magnesium dependent, alpha isoform	690.04	292.96	486.52	859.53	850.37	1201.6	707.21	1015.61	891.22	0.531900954	0.04765	0.00698808
10324	caspase 3	74.65	40.81	90.44	109.36	153.83	190.24	284.59	203.65	344.59	0.320153002	0.0477	0.00417898
17368	RB1-inducible coiled-coil 1	901.84	849.94	864.96	1457.9	1313.88	1382.7	1641.67	1296.45	1849.88	0.585240717	0.04772	0.00145884
40892	Smg-7 homolog, nonsense mediated mRNA decay factor (C. elegans)	670.82	509.13	594.16	909.72	935.57	910.25	1666.94	1373.11	1068.38	0.516934078	0.04777	0.00541515
14699	low density lipoprotein receptor-related protein associated protein 1	75.98	47.15	136.94	150.37	243.04	316.66	208.78	258.74	221.87	0.37167193	0.0478	0.00218354
9696	protein phosphatase 1A, magnesium dependent, alpha isoform	281.46	111.9	158.2	565.15	304.18	544.53	369.01	388.93	396.41	0.429528738	0.04783	0.00343167
12475	DEAD (Asp-Glu-Ala-Asp) box polypeptide 10	278.5	119.19	117.37	431.43	222.53	267.29	496.73	357.89	488.83	0.454868194	0.04785	0.0119792
12733	UTP11-like, U3 small nucleolar ribonucleoprotein, (yeast)	1652.37	1247.73	876.1	2092.8	2024.87	1852.2	1938.24	1738.63	2374.9	0.62823646	0.04788	0.00801569
16773	ribosomal protein L36a-like	2127.47	1381.86	1791.8	2318	2118.29	2209.7	3314.32	3112.02	3538.96	0.638255937	0.04798	0.02608004
2899	phospholipase C, delta 4	138.83	132.49	147.63	335.85	224.94	209.08	437.46	324.88	403.75	0.432808529	0.04808	0.0029441

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
8894	solute carrier family 25 (mitochondrial carrier, phosphate carrier), member 25	207.47	125.99	152.31	472.41	322.97	339.68	379.28	255.45	490.21	0.429884956	0.04813	0.00168341
12544	small nuclear RNA activating complex, polypeptide 3	209.56	210.89	181.97	483.56	393.07	376.22	534.26	381.24	455.1	0.459257847	0.0483	0.00019626
21985	ribosomal protein S24	1695.18	1059.62	1483.47	2021.2	1554.34	1896.5	2472.65	2477.22	2756.87	0.643194644	0.0484	0.02471089
7941	MOB1, Mps One Binder kinase activator-like 1B (yeast)	559.19	330.53	241.51	736.04	605.17	626.58	639.84	1041.82	1058.44	0.480567728	0.04846	0.00718575
7576	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily c, member 1	431.94	141.49	156.2	467.97	379.07	430.91	503.05	853.05	827.37	0.421578427	0.04851	0.01096394
3178	poly A binding protein, cytoplasmic 1	1067.51	748.21	815.73	1481.8	1078.86	1271.7	1876.97	1398.42	2496.99	0.547952194	0.04868	0.01791445
535	proteasome (prosome, macropain) subunit, beta type 7	683.88	672	512.62	1128.4	1099.82	764.78	900.56	1120.79	1147.58	0.606465831	0.04873	0.00532769
10969	jumonji domain containing 1A	120.68	130.02	130.27	359	253.12	231.81	183.78	364.53	594.57	0.383499177	0.04874	0.00763501
23568	BAT2 domain containing 1	1319.8	1222.06	1121.76	2051.9	1915.46	1889.3	2016.68	1664.87	1912.55	0.639891151	0.04878	0.00118966
20689	eukaryotic translation initiation factor 2, subunit 1 alpha	873.2	655.24	823.56	1212.8	1067.32	1162.7	1408.93	1473.86	1531.6	0.598688079	0.04879	0.00311329
13607	expressed sequence AU014645	1511.34	1084.22	1023.43	1777.7	1571.13	1910.8	1920.8	1977.06	2014.19	0.647889341	0.04882	0.00448388
10857	heterogeneous nuclear ribonucleoprotein M	1632.02	1042.99	1052.83	1963.8	1558.88	1934.4	1928.28	2260.45	2173.42	0.630807682	0.04884	0.00723852
16722	WD repeat domain 12	55.39	143.68	73.22	227.44	175.28	219.03	230.08	395.68	302.9	0.351249024	0.04887	0.0019824
38631	mitogen activated protein kinase 1	192.12	80.19	241.96	690.93	358.51	359.34	325.02	315.25	558.26	0.394483203	0.04889	0.00710119
29078	RIKEN cDNA D330037H05 gene /// similar to La-related protein 4 (La ribonucleoprotein domain family member 4)	1816.97	1506.16	1606.86	2248.8	2319.28	2211	2969.76	2795.29	2623.9	0.65004971	0.04892	0.00408689
10770	proteasome (prosome, macropain) 26S subunit, ATPase 2	97.46	73.58	87.82	238.69	150.72	194.9	269.55	209.28	386.58	0.357117236	0.04893	0.00134467

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
22147	ERO1-like (S. cerevisiae) /// ribosomal protein L31	1617.14	815.67	1036.5	1391.6	1541.45	2086.7	1682.06	2068.83	2407.9	0.62070841	0.04895	0.02270587
14175	Sec61 alpha 1 subunit (S. cerevisiae)	1726.8	1072.28	1292.76	2085.2	1791.82	2048.5	2261.11	2116.35	2459.79	0.641215536	0.04899	0.00585437
22705	ets variant gene 5	505.73	222.21	367.1	690.41	565.33	752.06	668.44	1034.4	639.83	0.503412275	0.049	0.00575794
16198	t-complex protein 1	1790.6	1179.11	1413.64	2167.2	1848.4	2128.3	2997.31	2246.75	2828.77	0.61664846	0.04902	0.01018077
20029	cDNA sequence BC024969	39.51	47.41	37.05	94.02	167.9	143.54	230.28	67.15	149.97	0.290715944	0.04904	0.00244613
6054	eukaryotic translation initiation factor 2, subunit 3, structural gene X-linked /// similar to eukaryotic translation initiation factor 2, subunit 3, structural gene X-linked	752.35	559.3	604.82	1247.6	942.02	997.47	1342.47	1152.54	1009.05	0.572833831	0.04905	0.00183039
21965	similar to PRAME family member 8 /// expressed sequence AU018829	2409.02	1514.52	1777.52	2919.6	2279.5	3009.7	2492.54	2865.6	3135.81	0.682651308	0.04905	0.01414262
7639	nucleolar protein 11	103.2	75.19	118.35	278.21	178.97	208.17	352.34	374.83	213.44	0.369548432	0.04907	0.0014996
14311	expressed sequence C78212	421.11	232.74	227.15	553.16	581.34	405.68	580.26	619.3	637.51	0.521726257	0.04909	0.003325
7266	ubiquitin-conjugating enzyme E2B, RAD6 homology (S. cerevisiae)	527.65	343.53	429.59	890.61	526.22	809.78	736.6	884.92	789.17	0.56100317	0.0491	0.00503997
11354	Mus musculus, clone IMAGE:3983419, mRNA	72	27.43	24.72	143.74	119.31	60.58	222.57	201.82	145.95	0.277749813	0.04912	0.00430038
10003	RNA guanylyltransferase and 5'-phosphatase	106.94	84.63	87.19	301.26	182.18	150.86	353.53	154.09	362.86	0.370499342	0.04914	0.00611854
1462	exosome component 5	40.73	61.74	17.43	87.19	68.73	139.55	145.69	246.81	177.61	0.277039673	0.04916	0.00522601
8293	ring finger and SPRY domain containing 1	633.65	573.63	579.76	1137.8	1373.91	1161.8	843.63	923.02	883.55	0.565188996	0.04916	0.00324511
8271	insulin-like growth factor 2 receptor	74.29	195.52	490.69	528.54	492.61	528.94	459.73	649.21	498.88	0.481647672	0.04917	0.01382869
8020	pleckstrin homology domain containing, family F (with FYVE domain) member 2	625.38	324.51	341.8	1052.9	850.6	998.02	505.77	626.89	737.95	0.54134623	0.04918	0.01741884
39491	zinc finger protein 710	727.55	440.46	524.24	900.66	784.47	1074.7	975.36	1333.25	979.43	0.559617585	0.04919	0.00456617
1486	DnaJ (Hsp40) homolog, subfamily B, member 9	2390.47	1648.72	1910.19	2935.8	2556.3	2981.4	2867.17	2942.9	3211.21	0.68013085	0.0492	0.00549139
15093	ribosomal protein S17	2459.15	1443.18	2004.58	2845.5	2446.97	2579.6	3506.76	2970.35	3555.66	0.659810063	0.0492	0.0169152
2115	torsin family 1, member B	256.19	89.29	93.92	278.8	251.82	363.38	456.6	304.99	408.7	0.425715379	0.04921	0.00453646

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
27418	cDNA sequence BC035295	443.69	439.27	554.8	570.76	834.16	708.51	1041.56	1097.16	911.05	0.556925937	0.04922	0.00861834
17466	G patch domain containing 4	560.23	265.59	219.67	798.37	518.64	726.56	747.62	528.15	616.45	0.531273264	0.04923	0.00884928
2013	eukaryotic translation initiation factor 3, subunit 4 (delta)	271.74	95.04	221.24	323.99	248.6	395.58	465.04	463.27	611.39	0.468939778	0.04924	0.01506929
27189	RIKEN cDNA E430025E21 gene	155.47	124.82	115.94	304.08	355.62	222.62	391.26	261.61	311.02	0.429236111	0.04925	0.00055666
7846	mannosidase, alpha, class 2C, member 1	6.08	8.03	3.61	38.99	14.56	85.09	59.15	145.44	156.62	0.07090127	0.04928	0.00203805
15948	ATPase type 13A3	1601.83	1562.47	1794.92	2141.2	2094.83	2038.2	3190.74	2860.13	3084.85	0.643639106	0.0493	0.01688862
3056	insulin-like growth factor 2 mRNA binding protein 1	23.6	23.13	59.05	86.87	175.53	124.87	113.81	134.37	168.17	0.263258754	0.0493	0.00046644
31866	M-phase phosphoprotein 1	584.18	393.46	472.99	807.12	805.27	884.71	916.93	822.02	1167.48	0.536919384	0.04933	0.00130358
11154	growth factor, erv1 (S. cerevisiae)-like (augmenter of liver regeneration)	390.69	114.24	604.44	777.64	571.65	1161.3	360.32	727.15	1012.77	0.481198735	0.04935	0.04517723
4640	cyclin M3	240.03	342.07	62.47	256.37	380.42	457.8	470.6	681.96	476.52	0.473309909	0.04937	0.02795838
12184	phosphatidylinositol transfer protein, cytoplasmic 1	70.98	70.85	77.64	242.87	150.65	236.21	204.73	89.26	219.98	0.383789459	0.0494	0.00477932
16753	COX4 neighbor	595.77	603.5	354.09	995.56	953.48	706.37	894.23	725.76	1113.09	0.576547419	0.04941	0.00769649
8770	TruB pseudouridine (psi) synthase homolog 2 (E. coli)	1598.35	788.01	714.92	2065.6	1575.69	1737.5	1380.01	1601.91	1777.98	0.611769533	0.04943	0.01407303
35301	gb:AV268519 /DB_XREF=gi:16389684 /DB_XREF=AV268519 /CLONE=4930535J16 /FEA=EST /CNT=4 /TID=Mm.38413.1 /TIER=ConsEnd /STK=2 /UG=Mm.38413 /UG_TITLE=ESTs	446.43	227.18	224.84	668.78	557.23	627.97	642.03	432.62	473.25	0.528207932	0.04945	0.00529613
11453	cDNA sequence BC010304	237.84	308.43	345.14	374.3	548.32	431.79	1196.49	751.72	656.51	0.450306002	0.04951	0.02034917
14840	ERO1-like (S. cerevisiae) ///	1914.61	1048.97	1332.05	1815.1	1860.89	2459.7	2005.25	2401.75	2725.53	0.647509527	0.04952	0.01837773
16309	ribosomal protein L31	5.67	27.92	7.3	114.53	67.49	114.67	39.64	77.22	93.19	0.162167495	0.04953	0.00068104
7219	prion protein	3806.66	2690.44	2523.5	4149.4	3627.73	3863	4934.86	4376.47	4773.38	0.701316785	0.04956	0.01729256
22327	proteasome (prosome, macropain) subunit, alpha type 4	310.97	225.94	272.82	597.48	410.26	388.07	523.64	561.2	597.35	0.52614035	0.04956	0.00218884

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
17151	exonuclease 3"-5" domain-like 2	69.67	29.49	42.91	151.69	68.29	135.72	157.58	233.27	169.66	0.310125408	0.04957	0.00330747
10743	sorbitol dehydrogenase	167.8	89.01	200.21	398.11	251.14	352.86	403.04	323.86	301.79	0.450088635	0.04957	0.00219161
6064	nuclear receptor coactivator 6 interacting protein	1342.43	996.25	1101.97	1935.4	1486.31	1789.3	1893.95	2042.34	1527.18	0.64464665	0.04958	0.00539514
14546	exocyst complex component 8	38.13	48.15	41.87	130.42	147.06	97.95	144.96	178.44	148.81	0.30236893	0.04961	4.94E-005
14589	RIKEN cDNA 4732471D19 gene	50.72	36.59	43.82	139.69	116.02	153.6	151.77	176.54	113.67	0.308073629	0.04963	3.73E-005
17331	ribosomal protein L15	1103.58	552.38	1197.41	1368.2	1136.31	1371.7	1675.87	2054.56	2318.95	0.574953378	0.04963	0.02851657
29240	catenin (cadherin associated protein), delta 1	55.32	151.05	51.37	125.57	252.74	142.05	216.39	309.83	274.46	0.390207715	0.04966	0.00867468
8570	expressed sequence AI840826	48.97	43.58	48.65	146.36	99.71	108.05	225.48	172.99	158.23	0.310050284	0.04967	0.00041891
29169	RNA binding motif protein 4	177.47	100.66	79.8	250.81	194.96	248.25	310.54	382.4	312.05	0.421339486	0.04967	0.00235887
40440	RIKEN cDNA C130032F08 gene	1581.36	805.74	827.62	1302.8	1329.25	1510.7	2220.74	2011.87	2336.34	0.600228723	0.0497	0.03109115
18873	basic leucine zipper and W2 domains 1	1856.21	1429.11	1724.42	2392.9	2189.52	2377.5	2784.24	2571.65	2854.64	0.660457255	0.04971	0.00401185
13899	karyopherin (importin) beta 1	30.07	36.79	45.53	113.82	67.63	72.13	158.24	180.43	197.68	0.284556859	0.04972	0.00325133
18692	GA repeat binding protein, alpha	1521.21	896.36	993.48	1692.9	1670.86	1806.1	2042.84	1734.73	2022.7	0.621877894	0.04976	0.00459458
42681	activating signal cointegrator 1 complex subunit 3	300.81	273.13	256.77	436.29	426.49	352.72	661.15	697.71	694.37	0.508276915	0.04976	0.00901142
7056	solute carrier family 22 (organic anion/cation transporter), member 12	78	127.98	90.24	109.02	174.7	301.99	259.4	365.4	248.19	0.406139671	0.04977	0.01328366
2136	peroxisomal membrane protein 2	28.37	71.63	15.47	123.16	112.33	103.5	137.14	245.59	211.62	0.247431296	0.0498	0.00176646
12220	histone aminotransferase 1	95.59	63.13	63.73	206.98	126.57	201.1	258.49	221.04	202.13	0.365775454	0.0498	0.00050866
12258	splicing factor, arginine/serine-rich 1 (ASF/SF2)	1549.68	1176.89	1231.71	2058.8	1861.02	1951.5	1990.7	2140.9	2193.12	0.649106926	0.0498	0.00186597
12983	PAK1 interacting protein 1	75.66	34.9	47.76	190.93	119.37	100.33	294.26	127.72	233.4	0.297030074	0.0498	0.00310024
12327	ribosomal protein L31	1611.89	822.91	1110.43	1525.	1520.0	1916.3	1780.17	2157.72	2472.13	0.623534481	0.0498	0.01946329
14684	heterogeneous nuclear ribonucleoprotein A1	1389.69	888.32	1178.26	1690.6	1425.06	1871.4	1956.12	2275.22	1706.98	0.632700133	0.0498	0.00961426
21465	translocase of outer mitochondrial membrane 20 homolog (yeast)	1215	813.02	1005.8	1760.6	1492.5	1269.6	1678.3	1498.07	2176.74	0.614391051	0.04982	0.00941987

Table 8B

Row	Gene	10/04/01	10/04/02	10/04/03	NI.1.1	NI.1.2	NI.1.3	NI.2.1	NI.2.2	NI.2.3	foldchange	fdr	t.pvals
13601	guanine nucleotide binding protein-like 3 (nucleolar)	627.77	328.73	461.31	838.59	772.45	739.06	836.66	835.1	1505.97	0.512971636	0.04985	0.00920993
25426	DNA segment, Chr 5, Wayne State University 178, expressed	31.53	65.26	42.32	176	76.21	139.61	139.72	309.45	160.76	0.277733966	0.04985	0.00304201
22109	RIKEN cDNA 2700007P21 gene	563.57	337.22	396.2	764.62	661.9	636.46	1052.76	777.27	896.81	0.541561061	0.04986	0.00415005
34705	uroporphyrinogen decarboxylase	312.94	195.7	307.86	802.17	395.61	379.25	967.5	396.14	683.09	0.450636908	0.04986	0.02054484
3486	homeodomain interacting protein kinase 3	514.46	276.53	265.41	530.11	301.61	663.23	1201.72	793.58	994.76	0.471080332	0.04986	0.05047199
23671	polycomb group ring finger 5	728.53	398.49	398.87	916.23	850.34	1205.5	787.59	737.65	933.74	0.561910338	0.04986	0.00662276
3870	zinc finger protein 292	61.98	31.99	50.41	171.7	60.09	122.44	175.23	148.97	187.23	0.333572072	0.0499	0.00486358
1959	N-myc downstream regulated gene 3	19.32	37.3	32.34	163.21	83.33	51.59	232.87	162	63.09	0.235315902	0.04991	0.00686013
21127	coiled-coil domain containing 86	132.83	55.7	44.7	210.48	136.27	157.44	194.19	308.63	357.29	0.341904273	0.04991	0.00496129
7468	trans-golgi network protein	504.95	301.85	270.73	671.72	450.68	487.94	700.65	843.07	702.27	0.558837029	0.04991	0.01315788
4093	component of oligomeric golgi complex 4	747.08	426.75	620.57	928.72	864.53	976.28	1190.49	1078.38	2087.76	0.503609237	0.04995	0.01527683
19091	phosphoserine aminotransferase 1	962.82	668.17	307.26	1569	751.88	888.84	1338.76	947.09	1617.53	0.544979622	0.04995	0.03781093
7812	ATPase, Na+/K+ transporting, alpha 1 polypeptide	47.24	70.25	39.58	362.38	201.58	163.49	145.12	138.2	113.32	0.279461609	0.04995	0.0017652

Table 9A

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	R	MeanFDR	T
8013	Ras11b: RAS-like, family 11, member B	1175.28	1162.12	866.08	67.51	155.4	83.97	28.63	67.32	76.13	13.37793368	0	0.013333333	281.069
16505	Gyg: glycogenin	2774.07	3647.22	2911.43	451.38	788.3	545	633.2	808.2	736.02	4.71113938	0	0.036666667	146.474
44280	Gyg: glycogenin	2716.28	3189.75	2680.95	587.38	955.1	642.4	640.7	735.8	665.37	4.063169251	0	0.046666667	115.459
19189	Hsbp1: heat shock factor binding protein 1	2054.28	1787.07	1312.02	326.46	423.8	376.3	415.9	610.1	588.56	3.760019554	0	0.056666667	88.0219
38879	0610008C08Rik /// LOC621156 /// LOC631337: RIKEN cDNA 0610008C08 gene /// hypothetical LOC621156 /// hypothetical protein LOC631337	574.74	794.44	810.38	108.25	279.7	167.9	96.73	84.66	122.89	5.068037018	0	0.044285714	82.2657
21617	Ywhab: tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, beta polypeptide	1175.16	1329.81	1339.93	250.85	326.1	341.3	285.4	335.3	415.3	3.935031906	0	0.045	79.1967
12776	Gpsn2: glycoprotein, synaptic 2	960.08	781.16	213.63	107.93	146.7	123.1	160.8	174.9	137.46	4.595531107	0	0.043333333	75.6419
38623	Rgs2: regulator of G-protein signaling 2	963.98	951.57	378.04	252.29	232	224.9	47.94	265.8	134.38	3.963622853	0	0.041666667	72.6641
14769	Ywhab: tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, beta polypeptide	984.18	1329.73	1254.81	241.01	312	348.2	323.2	330.1	398.25	3.654921319	0	0.043333333	69.7329
21642	Tax1bp3 /// Rpl13: Tax1 (human T-cell leukemia virus type I) binding protein 3 /// ribosomal protein L13	1245.68	2205.53	913.44	382.12	394.4	282.3	507.1	145.2	438.71	4.060385047	0	0.041666667	68.3463
22387	BC018601: cDNA sequence	871.11	1011.02	988.69	262.99	239.9	197.1	251.4	209.6	269.45	4.013785582	0	0.041555556	67.9884
28288	Zfp291: zinc finger protein 291	987.39	413.21	165.77	31.66	208.2	60.95	9.02	50.48	51.43	7.609463431	0	0.038333333	65.2239
15663	Ninj1: ninjurin 1	768.76	910.41	1337.77	236.5	145	273.4	260.1	211.1	352.74	4.080392223	0	0.037121212	60.0212
27398	Zar1: zygote arrest 1	196.34	550.87	430.24	96.91	62.32	61.6	36.14	73.34	105	5.409708024	0	0.0376	56.505
8065	Hsbp1: heat shock factor binding protein 1	1479.03	1221.16	1148.4	399.02	455.2	495.6	385.2	442.7	494.47	2.880433196	0.0222222	0.053111111	42.8537
20436	Gltcr2: glioma tumor suppressor candidate region gene 2	229.4	345.29	561.03	117.87	44.11	14.4	87.12	56.32	105.97	5.334648536	0.0227273	0.054166667	42.8606

Table 9A

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	R	MeanFDR	T
10123	Hspb1: heat shock protein 1	451.26	233.48	504.65	36.78	60.45	103.9	181.6	70.72	133.05	4.055475996	0.0232558	0.054263566	43.1189
32039	Myo5b: myosin Vb	932.01	832.86	419.75	166.98	224	201.3	246.4	398.8	209.84	3.018911206	0.025641	0.057435897	43.5021
19373	Mtap: methylthioadenosine phosphorylase	430.05	544.69	232.61	76.62	179.7	135.6	24.53	69.42	68.87	4.352771519	0.0263158	0.05877193	43.6173
15861	Ptk9l: protein tyrosine kinase 9-like (A6-related protein)	3879.84	515.28	668.16	278.34	548	167.5	39.25	201.2	238.1	6.877821117	0.027027	0.053873874	45.4725
5039	Ywhab: tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, beta polypeptide	786.43	1401.89	1268.98	550.3	550.1	542.5	548.9	344.8	534.84	2.251344699	0.0275229	0.057889908	31.9962
7118	Zfp313: zinc finger protein 313	664.63	959.1	606.33	102.21	233.1	60.95	260.6	238.4	243.17	3.917884751	0.0277778	0.052314815	45.8903
27301	Chsyl: carbohydrate (chondroitin) synthase 1	3606.66	3856.21	3309.81	1920.1	1952	1650	1778	2224	1984.7	1.872169743	0.0283019	0.058207547	32.1594
23482	Wdr51b: WD repeat domain 51B	549.71	635.29	186.24	78.52	121.3	128.3	70.8	228.5	105.47	3.74236511	0.0285714	0.052857143	46.074
22475	Sept11: septin 11	379.67	680.28	606.49	144.25	278.7	239.7	234.4	123.4	131.01	2.894758327	0.0288462	0.05875	32.2155
16314	2700060E02Rik: RIKEN cDNA 2700060E02 gene	1657.64	2133.91	1718.85	759.77	1151	666.9	740.4	552.1	918.56	2.301557302	0.0291262	0.05776699	32.463
40450	Pfdn4: prefoldin 4	835.74	1351.12	1275.03	465.73	482.3	379.6	367.1	416.9	416.99	2.738241285	0.0294118	0.052941176	46.429
3542	Rgs2: regulator of G-protein signaling 2	910.45	1375.3	511.1	289.77	380.2	277.6	309.3	389.5	287.36	2.89283424	0.0294118	0.05622549	37.2147
38514	Prkaca: protein kinase, cAMP dependent, catalytic, alpha	1061.04	1151.27	898.98	699.13	230.1	414.1	446.6	245.9	459.58	2.493540321	0.0294118	0.058267974	32.4698
9746	Ptpnj: protein tyrosine phosphatase, receptor type, J	313.63	348.74	149.56	38.69	84.77	85.95	59.83	36.02	100.14	4.005574741	0.029703	0.058778878	32.499
21697	Zfp313: zinc finger protein 313	3162.11	3960.29	3102.1	1059.1	1648	1223	1678	2090	1792.6	2.154513004	0.0298507	0.054328358	37.9163
19162	Gtf2b: general transcription factor IIB	952.55	1437.02	1358.53	532.39	552.3	466.5	425.1	572.2	544.43	2.423695714	0.030303	0.05479798	37.9814
5038	Ywhab: tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, beta polypeptide	842.81	1251.66	740.5	218.36	255.6	347.1	475.5	409.7	250.61	2.897586852	0.030303	0.058754209	32.6829

Table 9A

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	R	MeanFDR	T
16058	Gm428 /// LOC623180 /// LOC623197 /// LOC623210 /// LOC623219: gene model 428, (NCBI) /// hypothetical LOC623180 /// hypothetical LOC623197 /// hypothetical LOC623210 /// hypothetical LOC623219	1730.14	2814.61	2209.92	718.98	1302	938	1086	1344	986.94	2.118852909	0.0309278	0.059450172	32.7088
1904	Ube2a: ubiquitin-conjugating enzyme E2A, RAD6 homolog (S. cerevisiae)	835.58	1090.31	1082.16	322.75	424.9	331.9	208	371.1	379.5	2.9516296	0.03125	0.053854167	46.9047
1431	SrpK2: serine/arginine-rich protein specific kinase 2	1149.91	1761.01	1427.58	473.26	698.9	399.7	966.1	423.3	728.78	2.351516006	0.03125	0.06	32.7117
11264	2610510J17Rik: RIKEN cDNA 2610510J17 gene	957.03	1024.23	729.2	175.1	254.7	235.5	303	296	462.73	3.138886637	0.0322581	0.054086022	47.3645
16336	Uchl1: ubiquitin carboxy-terminal hydrolase L1	1255.01	1792.27	1364.58	298.99	536.1	340.2	818.6	772.9	910.66	2.399366965	0.0325203	0.061382114	30.5168
16195	Bpgm: 2,3-bisphosphoglycerate mutase	1084.63	1316.44	933.03	223.6	542.2	298.2	441	450.4	441.04	2.78256733	0.0327869	0.055846995	38.5724
15069	2700060E02Rik: RIKEN cDNA 2700060E02 gene	1911.39	1978.28	1849.47	866.04	1267	698.8	686.8	853.4	876.63	2.186763905	0.032967	0.058608059	33.4602
1378	Sec61b: Sec61 beta subunit	629.4	1333.69	789.37	274.35	516.3	341.2	201.2	346.8	199.6	2.929224024	0.0330579	0.061019284	30.716
277	Herpud2: HERPUD family member 2	1579.89	1939.26	1921.14	868.46	883.8	596.7	827.7	1099	877.44	2.11163511	0.0333333	0.059185185	33.4759
22096	Mm.69144.1	453.06	395.64	858.44	130.38	120.9	231.2	100.7	88.48	177.73	4.019684715	0.0337079	0.057453184	33.9669
28335	Usp27x: ubiquitin specific peptidase 27, X chromosome	1037.34	637.29	144.44	150.48	158.3	218	119.3	163.4	242.26	3.459491841	0.0338983	0.056384181	39.0302
1111	Nek7: NIMA (never in mitosis gene a)-related expressed kinase 7	717.12	574.95	336.93	210.68	267.8	172	134	130	199.51	2.924911121	0.0338983	0.060988701	30.8699
12221	Cpa1: carboxypeptidase A1	1233.98	1187.56	1071.87	304.36	410.4	369.7	630.8	505.1	743.84	2.357067674	0.0344828	0.061005747	30.9745
10438	5830415L20Rik: RIKEN cDNA 5830415L20 gene	1215.2	1852.35	1334.99	447.1	794.1	612.1	547.7	578.6	472.45	2.550718424	0.0348837	0.059224806	33.9864
20740	Mm.27444.1	337.59	481.47	424.2	81.2	144.9	122.1	82.14	158	135.08	3.437268455	0.0350877	0.06	31.3576
6654	Hmgn1: high mobility group nucleosomal binding domain 1	2802.24	3952.07	3639.58	1578	1726	1428	2308	2200	2525.7	1.766840536	0.0357143	0.061142857	29.2973
2773	Lmo1: LIM domain only 1	728.82	1098.32	1034.76	305.5	364.2	304	486.9	565.8	414.43	2.345012373	0.036	0.060986667	30.4439

Table 9A

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	R	MeanFDR	T
16848	Ube2a: ubiquitin-conjugating enzyme E2A, RAD6 homolog (S. cerevisiae)	986.42	1081.65	857.17	294.92	401.6	308.1	491.7	435.7	385.52	2.524566112	0.0363636	0.058787879	31.7403
28139	Mall: mal, T-cell differentiation protein-like	140.43	181.82	65.34	2.1	15.41	3.36	15.11	1	10.85	16.20698307	0.037037	0.058436214	34.7455
23084	2410127E18Rik: RIKEN cDNA 2410127E18 gene	526.43	815.91	633.99	308.65	278.7	283.9	152.4	171.6	168.65	2.897993299	0.037037	0.060518519	29.7435
42427	Mrg1: myeloid ecotropic viral integration site-related gene 1	1187.24	1519.64	1212.95	887.49	524.6	553.7	452.6	589.9	467.49	2.255569398	0.037037	0.063101852	25.4067
34813	LOC627488 /// LOC627520 /// LOC667692 /// LOC667695 /// LOC673990: similar to THO complex subunit 4 (Tho4) (RNA and export factor binding protein 1) (REF1-I) (Ally of AML-1 and LEF-1) (Aly/REF) /// similar to THO complex subunit 4 (Tho4) (RNA and export factor binding protein 1) (REF1-I) (Ally of AML-1 and LEF-1) (Aly/REF) /// similar to THO complex subunit 4 (Tho4) (RNA and export factor binding protein 1) (REF1-I) (Ally of AML-1 and LEF-1) (Aly/REF) /// similar to THO complex subunit 4 (Tho4) (RNA and export factor binding protein 1) (REF1-I) (Ally of AML-1 and LEF-1) (Aly/REF) /// similar to THO complex subunit 4 (Tho4) (RNA and export factor binding protein 1) (REF1-I) (Ally of AML-1 and LEF-1) (Aly/REF)	2997.33	4389.29	4558.04	2547.1	3079	2572	1852	2591	2371.8	1.591199861	0.0372093	0.063224806	25.4182

Table 9A

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	R	MeanFDR	T
14324	2700060E02Rik: RIKEN cDNA 2700060E02 gene	1736.22	1743.31	1787.5	771.81	1146	670	697.7	850.4	807.84	2.130990215	0.0373134	0.060746269	29.8051
5988	Ak311: adenylate kinase 3 alpha-like 1	923.36	955.8	444.05	293.2	479	230.4	457.2	322.9	400.3	2.128475753	0.0373832	0.063317757	25.4548
1245	Tnfaip8: tumor necrosis factor, alpha-induced protein 8	360.72	565.75	517.2	173.14	264.6	167.2	150.3	188	131.01	2.688048113	0.0375587	0.063067293	25.4969
29349	Srrm2: serine/arginine repetitive matrix 2	422.76	520.49	235.84	276.19	142	111.1	23.71	54.04	95.17	3.358560971	0.0378788	0.060505051	30.0013
18213	Gnas: GNAS (guanine nucleotide binding protein, alpha stimulating) complex locus	749.56	1054.77	1007.27	303.09	467.2	274.3	329.4	595	414.54	2.359288924	0.0381679	0.060916031	30.0133
12033	Zfp313: zinc finger protein 313	2750.76	3476.54	2573.42	687.45	1148	821.6	1426	1577	1289.5	2.532599515	0.0384615	0.048589744	51.0983
27637	Nr2e1: nuclear receptor subfamily 2, group E, member 1	1154.42	1358.16	898.95	372.14	477.2	439.9	220.9	478.6	491.58	2.750834556	0.0384615	0.055705128	40.7351
2211	Fxc1: fractured callus expressed transcript 1	319.96	265.72	286.07	4.63	47.28	3.92	13.61	90.27	179.09	5.146103896	0.0384615	0.058162393	35.2462
23218	5730405I09Rik: RIKEN cDNA 5730405I09 gene	1466.98	1315.46	1000.54	584.51	638.8	679.3	643.1	491.1	601.7	2.079343049	0.038835	0.062718447	25.8477
9918	Nobox: NOBOX oogenesis homeobox	557.22	579.54	288.37	129.03	371.4	87.59	172.8	98.82	245.05	2.580168012	0.0390244	0.062520325	25.9252
27981	1110003E01Rik: RIKEN cDNA 1110003E01 gene	935.07	637	979.71	438.48	639.8	244.7	270.6	373.4	222.36	2.331073921	0.0392157	0.060915033	28.4683
23766	4930431B09Rik: RIKEN cDNA 4930431B09 gene	288.09	929.85	626.58	234.92	253.4	170	197.8	182.5	246.51	2.870446163	0.0396825	0.061349206	30.3043
2222	Ddx19a: DEAD (Asp-Glu-Ala-Asp) box polypeptide 19a	1171.65	1506.5	1210.32	497.39	638.6	587.9	581.9	501.6	556.5	2.311923278	0.0397727	0.062424242	27.0664
9267	BC004728: cDNA sequence BC004728	2914.05	3310.43	4063.56	1784.9	1884	1762	2063	1805	2090.6	1.80671016	0.039801	0.063034826	25.9867
11008	Sec24b: SEC24 related gene family, member B (S. cerevisiae)	491.17	371.38	563.69	38.39	140.5	93.28	159.3	103.5	197.21	3.896085448	0.04	0.056866667	41.2419
20332	2610510J17Rik: RIKEN cDNA 2610510J17 gene	920.34	904.56	875.79	268.6	445	387.9	292.4	305.2	492.61	2.464504236	0.04	0.061555556	28.6135
1259	Eefsec: eukaryotic elongation factor, selenocysteine-tRNA-specific	420.39	175.37	131.6	27.72	33.52	41.27	19.45	30.53	65.01	6.688367816	0.04	0.062038095	27.1963

Table 9A

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	R	MeanFDR	T
17879	Ehd4: EH-domain containing 4	1018.34	1327.03	1393.04	514.73	741	621.1	418.3	856.2	476.56	2.06100183	0.04	0.062966667	26.0088
7236	Snx9: sorting nexin 9	2034.25	2223.65	2179.13	937.58	906.1	769.4	1117	1452	1343	1.972975541	0.0402685	0.060939597	28.7304
19870	Ggtal1: glycoprotein galactosyltransferase alpha 1, 3	2076.84	1911.12	1747.57	1033.9	1405	1131	712.7	1003	500.21	1.982372856	0.040404	0.063013468	26.0912
4975	Ywhag: 3-monooxygenase/tryptophan 5-monooxygenase activation protein, gamma polypeptide	1701.12	2425.78	1962.92	1046	935.4	694.1	728.4	780.6	1139.8	2.287531271	0.0405405	0.057837838	35.922
3840	Atp6v1c1: ATPase, H ⁺ transporting, lysosomal V1 subunit C1	1562.18	1993.36	1324.28	641.08	676.2	434.4	835.1	1096	751.79	2.200907908	0.0405405	0.061306306	28.7411
1499	Kdelr2: KDEL (Lys-Asp-Glu-Leu) endoplasmic reticulum protein retention receptor 2	1134.97	1840.76	1296.63	454.68	740.1	649.3	505.6	898.4	565.51	2.240609193	0.0405405	0.063108108	25.1968
19039	Csda: cold shock domain protein A	1886.38	2541.08	2229.73	996.1	1299	937.6	1283	1407	1167.3	1.877663438	0.040724	0.06321267	25.2231
9095	Dnahc8: dynein, axonemal, heavy chain 8	275.09	424.82	466.62	57.34	144.7	48.48	81.76	152.8	74.6	4.169007541	0.0408163	0.054013605	41.8892
4403	Gadd45b: Growth arrest and DNA damage-inducible 45 beta	325.8	1088.25	1693.72	59.09	210.2	129.8	401.4	453.8	881.65	2.91008774	0.0408163	0.061360544	28.7836
23361	Fbxo28: F-box protein 28	728.16	665.63	382.02	171.44	132.3	122.3	209.7	213.5	344.89	2.974406646	0.0410959	0.056164384	36.4751
6486	G6pd2 /// G6pdx: glucose-6-phosphate dehydrogenase 2 /// glucose-6-phosphate dehydrogenase X-linked	919.61	1851.91	1150.83	309.7	590.7	402	725.6	646.1	587.89	2.404866938	0.0412371	0.063487973	26.2283
22429	Gorasp2: golgi reassembly stacking protein 2	1053.97	1211.58	759.04	341.93	459	419.6	583.3	420.2	553.71	2.177773618	0.0414747	0.063195084	25.3481
6921	Kif17: kinesin family member 17	1050.79	1040.1	651.51	576.54	561.7	248.7	435.1	356.1	394	2.132420979	0.0416667	0.063368056	26.3333
13584	Dcp1a: decapping enzyme	1250.95	1233.21	734.43	514.44	536.9	508.7	712.4	438.6	473.45	2.021454389	0.0418848	0.063560209	26.3363
4967	Ncoa4 /// LOC627557: nuclear receptor coactivator 4 /// similar to nuclear receptor coactivator 4	819.67	1349.74	597.1	225.35	376.4	254.5	372.8	517.1	318.51	2.679817891	0.0419162	0.06257485	27.4886
14382	Pcid2: PCI domain containing 2	208.21	208.84	103.64	13.83	77.97	8.25	39.19	18.14	57.28	4.85129973	0.0424242	0.063030303	27.5603
9970	Csrp1: cysteine and glycine-rich protein 1	707.37	1081.76	451.86	402.09	254.6	515.3	117.8	186.2	282.36	2.54908518	0.042471	0.065933076	23.6583

Table 9A

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	R	MeanFDR	T
9611	Ptprrj /// AW125753: protein tyrosine phosphatase, receptor type, J /// expressed sequence AW125753	276.18	258.57	380.78	49.33	87.25	38.19	50.96	146.2	102.3	3.861366512	0.0425532	0.064255319	24.6082
12280	2610528K11Rik: RIKEN cDNA 2610528K11 gene	847.62	350.15	887.05	191.27	286.5	257.4	379.5	362.4	372.64	2.254139704	0.0426829	0.063130081	27.6
14887	Tmem109: transmembrane protein 109	1049.13	1100.4	693.15	487.81	701.2	347.9	350.2	365	381.93	2.158401859	0.0427807	0.064171123	26.423
8101	Camk2g: calcium/calmodulin-dependent protein kinase II gamma	283.06	354.25	367.2	116.66	94.65	136.1	97.12	109.2	105.92	3.04563095	0.0428016	0.065823606	23.7561
22457	Sepw1: selenoprotein W, muscle 1	2758.54	3164.45	4104.83	1517.9	1644	1398	1870	1618	3061.9	1.805050019	0.0429688	0.065950521	23.7661
15377	Tm2d2: TM2 domain containing 2	761.31	348.95	1367.84	465.42	283.5	356.7	295.1	314.2	225.89	2.553662885	0.0431373	0.066183007	23.7726
13610	Pnrc1: proline-rich nuclear receptor coactivator 1	1875.98	2497.69	2218.37	912.68	1173	1023	1143	1267	1271.5	1.9415163	0.0432099	0.063333333	27.6831
22912	Calcocol1: calcium binding and coiled coil domain 1	645.5	564.97	318.61	259.49	280.1	113.5	190.8	154.3	163.4	2.632509534	0.0432432	0.06372973	26.5901
7409	Tgfb2: transforming growth factor, beta 2	793.78	1590.37	1213.53	412.2	635.7	521.1	714.9	491.8	663.43	2.09223397	0.0433071	0.065721785	23.8689
38991	Ttll11: tubulin tyrosine ligase-like family, member 11	629.88	997.15	450.48	230.82	216.8	257.8	240	375.5	152.52	2.819887748	0.0434783	0.063967391	26.5994
21361	Slc25a4: solute carrier family 25 (mitochondrial carrier, adenine nucleotide translocator), member 4	609.28	302.35	741.44	237.03	343.3	259.4	201.2	188.6	100.24	2.486193413	0.0434783	0.065144928	24.6327
21528	Cnn3: calponin 3, acidic	842.6	930.24	788.12	354.82	542.4	337.9	180.8	351.3	266.95	2.518163806	0.04375	0.062583333	27.9931
15242	Pofut2: protein O-fucosyltransferase 2	1778.35	1399.41	2187.94	882.1	1202	822.3	491.1	875.2	1214.3	1.955775711	0.0438247	0.064940239	24.0113
23699	Cd164l2: D164 sialomucin-like 2	294.25	488.64	324.23	53.73	65.52	83.92	138.4	143.4	129.99	3.600624431	0.0438596	0.064809942	24.7437
37480	4930562C15Rik: RIKEN cDNA 4930562C15 gene	266.99	324.53	138.98	37.48	48.81	45.36	87.04	111.8	47.11	3.868968805	0.043956	0.063791209	26.6715
16487	Wfs1: Wolfram syndrome 1 homolog (human)	539	852.8	431.85	108.02	256.6	262.6	278.2	201	172.7	2.851658705	0.044	0.065093333	24.0215
20014	Arrb2: arrestin, beta 2	261.8	442.51	421.66	79.45	122.9	130.7	191.7	126.2	149.35	2.813975283	0.0440529	0.064375918	24.8594

Table 9A

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	R	MeanFDR	T
33055	2700050L05Rik: RIKEN cDNA 2700050L05 gene	504.87	984.48	426.53	189.02	221.1	184	321.7	177.9	178.35	3.012176811	0.0443686	0.067963595	22.5402
408	Fkbp8: FK506 binding protein 8	590.14	427.04	302.19	164.62	241.9	117.1	99.71	136.4	185.58	2.791638014	0.0444444	0.069534392	21.8165
39254	8430429K09Rik: RIKEN cDNA 8430429K09 gene	483.61	603.3	382.14	140.52	172.4	143.3	254.2	103.1	239.14	2.791278738	0.0445344	0.06560054	24.0684
159	Bpgm: 2,3-bisphosphoglycerate mutase	4125.5	4888.39	3943.66	2336.5	3162	2111	2186	2053	2562.9	1.798272305	0.044586	0.063078556	28.0958
1105	Meal1: male enhanced antigen 1	641.79	1052.56	800.16	373.15	443.3	291.3	373	331.2	364.96	2.291779211	0.044586	0.069522293	21.8514
38903	2600010E01Rik: RIKEN cDNA 2600010E01 gene	247.28	200.25	369.05	57.76	100.1	37.76	33.32	102.2	59.89	4.175811813	0.0446429	0.064449405	24.9266
15195	Tm2d2: TM2 domain containing 2	807.85	315.89	1436.51	527.63	320.7	388.1	290.4	286.3	252.38	2.479036756	0.0446735	0.068373425	22.5493
21185	D10Wsu102e: DNA segment, Chr 10, Wayne State University 102, expressed	2625.46	3167.45	2893.72	1716.1	1847	1338	1962	1712	1576.6	1.711233117	0.0447284	0.069158679	21.9226
4343	2410022L05Rik: RIKEN cDNA 2410022L05 gene	1605.23	1541.87	1099.08	1006.1	799.1	735.3	264.5	1067	383.1	1.995685441	0.044898	0.06537415	24.1286
12245	Dnm1l: dynamin 1-like	714.07	890.45	642.35	203.05	404.7	362.8	383.8	214.8	288.04	2.419788054	0.0449827	0.068373702	22.606
109	Atp6v1b2: ATPase, H+ transporting, lysosomal V1 subunit B2	1155.57	1164.53	728.39	468.28	602.1	390.5	479.6	654.4	600.14	1.908347679	0.0450161	0.069464094	21.9365
41382	9230115E21Rik: RIKEN cDNA 9230115E21 gene	1049.24	1705.49	1742	671.13	1012	681.8	714.7	866.2	989.16	1.822353437	0.045082	0.065560109	24.1432
1430	Srpk2: serine/arginine-rich protein specific kinase 2	1999.89	2326.25	2092.34	1043.1	1218	1110	1149	875.2	960.85	2.019612501	0.0451613	0.06172043	28.271
16430	G6pdx: glucose-6-phosphate dehydrogenase X-linked	1912.61	2849.68	1863.38	787.91	1752	1060	912.4	963.3	1065.4	2.025944759	0.0451613	0.068913978	22.0228
2300	Sdhb: succinate dehydrogenase complex, subunit B, iron sulfur (lp)	538.02	229.97	832.03	138.81	236.8	218.7	38.77	316.8	160.01	2.883334535	0.0452675	0.065775034	24.1473
6223	Zbtb20: zinc finger and BTB domain containing 20	134.97	146.13	16	4.9	11.08	4.9	9.14	2.83	47.12	7.430286357	0.0452962	0.068222997	22.6625
13279	Dcp1a: decapping enzyme	370.06	844.47	493.12	102.24	141.4	115.2	321.2	207.5	303.81	2.866964391	0.0453074	0.068727077	22.0548

Table 9A

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	R	MeanFDR	T
9363	Ddx19a /// Ddx19b: DEAD (Asp-Glu-Ala-Asp) box polypeptide 19a /// DEAD (Asp-Glu-Ala-Asp) box polypeptide 19b	1900.93	2603.49	1537.03	1167.2	1153	885.9	897.4	1007	1064.7	1.956903393	0.0454545	0.06739899	23.3704
10998	Pold3: polymerase (DNA-directed), delta 3, accessory subunit	1214.68	1819.91	1832.16	850.28	830.5	871	1004	1059	901.76	1.764581113	0.045614	0.068175439	22.7446
18210	Lnk: linker of T-cell receptor pathways	473.25	331.54	156.1	95.8	81.69	130.4	71.84	51.82	156.54	3.267666463	0.0456432	0.065172891	24.3452
21746	Tmem109: transmembrane protein 109	786.47	847.01	508.13	367.58	528.2	245.5	273.8	272.9	273.33	2.183845573	0.0457746	0.068004695	22.7842
16900	Ube2j1: ubiquitin-conjugating enzyme E2, J1	1048.14	731.11	425.79	189.15	297.8	190.2	412.6	359	345.7	2.457827565	0.0458333	0.065013889	24.4002
14774	Surf4: surfeit gene 4	2026.91	2349.99	1707.51	1143.2	1323	1046	1020	1066	1054.3	1.82922382	0.0462046	0.069229923	22.1641
6591	Dbi: diazepam binding inhibitor	748.16	1411.18	1571.45	551.45	796.2	566.5	644.3	687.8	749.36	1.867435173	0.0462633	0.067793594	22.8636
9161	Llg1: lethal giant larvae homolog 1 (Drosophila)	1430.08	1103.73	1765.18	919.12	927.7	845	526.8	632.3	506.16	1.97334894	0.0463576	0.069392936	22.1803
14725	Aldh9a1: aldehyde dehydrogenase 9, subfamily A1	2464.65	2087.14	1545.48	936.87	1225	1018	1310	1014	1062.2	1.857250556	0.0464135	0.065316456	24.4636
22776	Abhd13: abhydrolase domain containing 13	1448.41	1446.62	1230.77	644.65	823	682.5	642.8	658.9	939.64	1.879049135	0.0464396	0.070846233	21.5641
8799	Arl8a: ADP-ribosylation factor-like 8A	450.63	484.52	240.82	185.21	144.7	165.6	126.1	202.3	138.16	2.444844075	0.0465116	0.071792636	20.9911
4729	Tcl1b3: T-cell leukemia/lymphoma 1B, 3	3067.43	4225.86	3944.03	1750.2	2137	2048	2719	3084	2461.6	1.582715554	0.0465839	0.070662526	21.5931
8719	Pstpip1: proline-serine-threonine phosphatase-interacting protein 1	133	150.18	47.71	10.97	19.4	25.08	18.91	12.49	10.27	6.814044481	0.0466667	0.069311111	22.2252
5059	Rab18: RAB18, member RAS oncogene family	1682.03	1655.73	1019.13	929.17	817.7	742.1	723.3	1044	764.41	1.735622591	0.0467836	0.072017544	21.015
11799	Runx1t1: runt-related transcription factor 1; translocated to, 1 (cyclin D-related)	720.19	672.38	443.2	238.13	239.4	221.5	336.5	333.9	281.49	2.224016573	0.046875	0.0708125	21.6373
3543	Rgs2: regulator of G-protein signaling 2	871.75	1343.48	839.92	552.53	601.3	456.3	215.4	392.9	290.79	2.435178026	0.0469799	0.069519016	22.2513
11937	Defb8: defensin beta 8	322.3	225.13	147.33	19.16	51.47	86.7	48.97	79.85	82	3.774331115	0.0470588	0.072215686	21.0551

Table 9A

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	R	MeanFDR	T
22002	Anapc5: anaphase-promoting complex subunit 5	2197.5	1065.37	1661.25	652.28	960.9	647.2	1135	645.4	1044.3	1.936841774	0.0471014	0.067777778	22.9611
270	Carhsp1: calcium regulated heat stable protein 1	695.96	1334.14	836.29	323.58	378.2	367.8	428.5	461.3	410.2	2.419384517	0.047138	0.06959596	22.2624
11459	Lhx8: LIM homeobox protein 8	915.65	1329.12	940.38	474.05	516.8	519	579.1	373.5	302.8	2.303731001	0.0471698	0.070597484	21.6799
34771	2700050L05Rik: RIKEN cDNA 2700050L05 gene	83.83	110.47	358.76	26.54	30.56	15.02	19.41	28.26	5.59	8.822140692	0.0472441	0.074041995	20.0479
412	Orc3l: origin recognition complex, subunit 3-like (S. cerevisiae)	512.79	479.04	423.31	81.65	207.6	115.3	178	144.3	323.47	2.694683525	0.0472727	0.067612121	22.9795
8298	Rap1a: RAS-related protein-1a	1038.15	1345.27	1417.09	535.11	752.6	615	727.4	726.5	676.38	1.884757581	0.0473373	0.071577909	21.187
18549	H1f0: H1 histone family, member 0	1417.73	1021.56	619.53	522.35	740.5	456.5	358.3	405.5	425.71	2.103018928	0.0479042	0.070588822	21.3322
29960	LOC433810: similar to transmembrane protein SHREW1	689.22	717.53	441.92	260.52	343.3	371.8	226.1	240.7	301.53	2.120070873	0.048	0.072835556	20.2788
2778	Ggtal: glycoprotein galactosyltransferase alpha 1, 3	615.11	636.67	665.46	268.83	335	231.9	263.9	175.6	342.73	2.369976637	0.048048	0.070640641	21.3411
23726	1700010D01Rik /// MGC118250: RIKEN cDNA 1700010D01 gene /// similar to hypothetical protein LOC76386	514.54	738.88	626.14	342.68	278.7	196.3	235.7	321	246.24	2.319556713	0.0482574	0.073038427	20.3038
12425	Zcchc3: zinc finger, CCHC domain containing 3	601.6	494.73	290.75	165.02	163.9	195.7	202.5	148.4	244.54	2.476795886	0.0483384	0.070956692	21.3504
10133	Trim25: tripartite motif protein 25	616.96	638.71	337.03	143.65	248.1	271	339.9	229.3	207.55	2.21282094	0.0484848	0.071151515	21.3508
8706	Car10: carbonic anhydrase 10	126.12	168.15	16.02	9.61	16.81	12.12	1.76	21.12	7.68	8.98089725	0.0485075	0.067524876	23.2315
7747	Arpc4: actin related protein 2/3 complex, subunit 4	1172.92	1818.6	1389.88	803.24	994.5	747.6	842.5	574.3	530.53	1.950496372	0.0485934	0.075302643	19.7699
3909	Ches1: checkpoint suppressor 1	691.62	717.97	330.8	186.5	461.5	200.6	209.6	288.8	218.2	2.223671686	0.0486891	0.067752809	23.2316
1396	Hspa2: heat shock protein 2	1165.3	1847.2	1706.76	768.33	1326	888	653	1010	758.87	1.746794118	0.0487179	0.0752564	19.7913

Table 9A

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	R	MeanFDR	T
43852	LOC545637 /// LOC623272 /// LOC623281 /// LOC636750 /// LOC667780: hypothetical LOC545637 /// hypothetical LOC623272 /// hypothetical LOC623281 /// hypothetical protein LOC636750 /// hypothetical protein LOC667780	3154.18	5046.7	4038.1	1861.5	3004	2393	2912	2908	2107.5	1.611845948	0.0487805	0.073486902	20.3393
1723	Gng3: guanine nucleotide binding protein (G protein), gamma 3 subunit	939.26	1673.4	1032.44	376.96	611.2	390.9	606.5	482.2	588.54	2.385349318	0.0488722	0.067531328	23.2874
85452	Reep2: receptor accessory protein 2	1276.5	951.38	773.57	228.14	348.3	268	801	510.5	544.43	2.223139853	0.0489297	0.07108053	21.4329
26935	Bahd1: bromo adjacent homology domain containing 1	496.5	645.98	493.5	200.26	217.8	255.2	202.4	244.1	139.48	2.598567276	0.0489691	0.075197595	19.8532
927	Mod1 /// LOC624892 /// LOC677317: malic enzyme, supernatant /// similar to NADP- dependent malic enzyme (NADP- ME) (Malic enzyme 1) /// similar to NADP-dependent malic enzyme (NADP-ME) (Malic enzyme 1)	998.06	922.67	686.51	467.78	580.6	523.9	272.3	372.6	349.48	2.031565455	0.0491803	0.072905282	20.4366
38626	Mfap2: microfibillar-associated protein 2	509.37	115.67	349.24	27.37	198.3	120.4	76.61	70.93	137.03	3.089568568	0.0492308	0.071117949	21.4898
33836	Transcribed locus	22.68	284.45	137.88	67.18	11.32	10.03	13.47	8.36	30.26	6.329256151	0.0494792	0.0740625	20.0011
13825	Plekhc1: pleckstrin homology domain containing, family C (with FERM domain) member 1	433.88	1351.76	1329.49	686.75	581.8	481.9	519	457	620.73	1.861357091	0.0495868	0.072901745	20.4943
20764	Txn2: thioredoxin 2	417.2	626.98	484.3	144.42	244.2	113.1	228.5	145.6	295.77	2.609329521	0.0498615	0.072816251	20.5416

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
3646	0610007P06Rik: RIKEN cDNA 0610007P06 gene	53.21	84.8	297.2	251.3	275.76	304.24	515.25	489.89	705.11	0.342476048	0.04973822	0.0741274	20.0279
4498	0610016J10Rik: RIKEN cDNA 0610016J10 gene	4.86	1.17	7.33	143.31	63.22	200.98	78.49	177.41	3.9	0.04004136	0.042105263	0.06375439	26.3724
21983	1110005A23Rik: RIKEN LOC625193 /// LOC636633: RIKEN cDNA 1110005A23 gene /// similar to cytokine induced protein 29 kDa /// similar to cytokine induced	278.64	96.47	238.75	422.57	521.22	543.58	778.62	488.14	494.19	0.377955374	0.048632219	0.07113475	21.3759
38875	1110008B24Rik: RIKEN cDNA 1110008B24 gene	377.94	192.85	622.98	908.37	546.03	872.14	1023.5	1117.8	1343	0.41087267	0.044303797	0.06272152	28.0809
14747	1110014K08Rik: RIKEN LOC664786 /// LOC664849 /// LOC669054 /// LOC672175: RIKEN cDNA 1110014K08 gene /// hypothetical protein LOC664786 /// hypothetical protein LOC664849 /// hypothetical protein LOC669054 /// hypothetical protein LOC672175	181.23	248.59	247.67	383.41	393.19	537.89	753.56	769.32	870.69	0.365414799	0.041666667	0.06175926	28.8862
40498	1110034A24Rik: RIKEN cDNA 1110034A24 gene	43.95	64.41	50.95	319.84	238.19	282.21	220.54	188.83	248.29	0.212711129	0.034965035	0.06109557	29.101
23309	1600002K03Rik: RIKEN cDNA 1600002K03 gene	107	59.9	245.28	683.77	429.59	346.82	750.02	281.7	595.32	0.267023406	0.033333333	0.05611111	38.745
41245	2010107H07Rik: RIKEN cDNA 2010107H07 gene	194.12	113.48	300.45	660.66	354.66	360.51	650.65	483.92	1044.3	0.342111408	0.040816327	0.06326531	26.1485
12325	2310036O22Rik: RIKEN cDNA 2310036O22 gene	355.44	333.42	817.17	1346.9	1033.9	1084.7	1228.3	1278.2	1731	0.391021738	0.039473684	0.05811404	35.6782
16489	2410016O06Rik: RIKEN cDNA 2410016O06 gene	569.29	93.03	367.09	890.36	502.25	860.54	1000.8	906.94	1398.2	0.370354647	0.035714286	0.0597619	31.4657

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
15550	2510039O18Rik: RIKEN cDNA 2510039O18 gene	530.8	99.11	420.85	1242.7	731.24	1077.5	660.72	895.16	784.09	0.389791149	0.044198895	0.06355433	26.7591
19105	2510039O18Rik: RIKEN cDNA 2510039O18 gene	89.8	83.01	199.83	321.54	214.67	216.65	401.71	481.55	470.2	0.353830377	0.049450549	0.07271978	20.4938
35340	2600011C06Rik: RIKEN cDNA 2600011C06 gene	1515.45	1353.66	1480.64	2833.1	2791.7	2941	2356.4	2682.9	2361.5	0.544860232	0.041176471	0.06201961	27.4584
2216	2610029G23Rik: RIKEN cDNA 2610029G23 gene	572.06	520.64	750.01	1471	1141.3	1096.6	1730.3	1327.7	1092	0.46894679	0.046931408	0.06761733	22.9544
12846	2810409H07Rik: RIKEN cDNA 2810409H07 gene	86.52	29.18	100.39	480.67	269.97	281.71	245.18	142.85	326.22	0.247440742	0.042553191	0.0643617	26.3759
8238	2900073H19Rik: RIKEN cDNA 2900073H19 gene	211.83	95.11	192.63	574.46	220.64	424.2	345.06	612.16	561.13	0.36496265	0.044444444	0.06426667	24.9235
13588	cDNA, RIKEN full-length enriched library, clone:A630012F14 product:unclassifiable, full insert sequence	573.65	184.37	288.58	996.46	770.78	953.56	685.85	734.41	780.12	0.425345141	0.045751634	0.06929194	22.0658
36328	3110050K21Rik: RIKEN cDNA 3110050K21 gene	311.28	401.01	530.93	1030.8	761.38	1078.3	1332.5	1029.5	1510.5	0.368745523	0.036144578	0.05738956	34.6953
24498	3110050K21Rik: RIKEN cDNA 3110050K21 gene	451.12	293.97	202.68	849.56	599.42	547.68	1108.3	684.07	839.07	0.409568423	0.043103448	0.06469828	24.629
23071	3300001M20Rik: RIKEN cDNA 3300001M20 gene	108.79	169.76	262.21	708.72	562.73	499.17	858.23	902.18	429.21	0.27309456	0.02173913	0.05210145	42.8378
32215	4833426J09Rik: RIKEN cDNA 4833426J09 gene	101.24	83.15	123.05	403.04	266.83	290.65	664.64	325.8	820.34	0.221874211	0.03	0.05903333	32.5436
41050	4930427A07Rik: RIKEN cDNA 4930427A07 gene	154.04	258.9	438	462.21	493.17	756.34	822.19	1212.3	977.74	0.36026859	0.036036036	0.05921922	31.6007
24760	4933402C05Rik: RIKEN cDNA 4933402C05 gene	184.74	248.77	343.55	582.5	656.43	516.22	821.01	823.62	1015	0.352026602	0.031578947	0.06042105	32.7419
23146	4933417E01Rik: RIKEN cDNA 4933417E01 gene	1233.06	595.65	1580.61	2342.5	1725.8	1965.4	2559.7	2606.8	2682.4	0.491168017	0.035714286	0.05678571	34.675
23294	4933440H19Rik: RIKEN cDNA 4933440H19 gene	222.08	110.71	164.48	490.16	293.12	477.16	332.22	512.35	433.13	0.391838118	0.049079755	0.07127812	21.4347
23653	5730406M06Rik: RIKEN cDNA 5730406M06 gene	175.07	173.9	279.19	681.23	606.53	632.71	617.02	514.97	512.55	0.352402939	0.037593985	0.06057644	29.9153

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
39158	5730406M06Rik: RIKEN cDNA 5730406M06 gene	475.1	352.64	504.28	956.32	865.18	1125.5	938.04	1078.7	754.09	0.46591231	0.047318612	0.07053628	21.7093
27731	5730406M06Rik: RIKEN cDNA 5730406M06 gene	984.39	305.06	745.9	1521.6	1377	1419.4	1034.7	1357.1	1056.2	0.524172156	0.048192771	0.07079317	21.3436
27732	5730406M06Rik: RIKEN cDNA 5730406M06 gene	964.44	283.46	700.66	1444	1271	1384.7	996.72	1324	1014.8	0.5241432	0.049315068	0.07291324	20.452
40394	A230097K15Rik: RIKEN cDNA A230097K15 gene	739.64	322.54	497.24	1135.4	943.8	714.8	1147.6	1095.1	1136.2	0.50524631	0.044871795	0.06926282	21.9347
40844	A530082C11Rik: RIKEN cDNA A530082C11 gene	132.33	27.77	268.5	459.34	222.66	476.49	456.67	365.07	541.49	0.339926717	0.044444444	0.06285185	26.878
13848	AA408556: expressed sequence AA408556	171.39	60.02	172.24	295.85	310.39	421.99	454.92	383.23	796.93	0.303119051	0.044843049	0.06391629	25.0467
44961	AA591059: Expressed sequence AA591059	56.29	33.55	46.52	130.59	70.28	207.6	293.95	193.18	298.12	0.228462286	0.044067797	0.06924294	22.3522
20263	Abcf1: ATP-binding cassette, sub-family F (GCN20), member 1	1565.6	1512.55	1825.57	2799.7	2379.2	2881.9	3476.5	3087.8	3006.5	0.556241255	0.038461538	0.06267628	25.7414
3399	Abhd6: abhydrolase domain containing 6	74	36.87	87.35	553.9	326.89	296.3	593.6	167.18	618.84	0.15505865	0	0.04020833	66.0252
3398	Abhd6: abhydrolase domain containing 6	209.56	278.43	237.36	422.75	380.61	691.33	684.84	889.36	650.71	0.390015055	0.040229885	0.06212644	27.2306
7228	Adnp: activity-dependent neuroprotective protein	290.38	98.62	340.91	806.44	487.17	695.53	665.1	564.28	782.06	0.364902089	0.036764706	0.06019608	29.7163
28853	AI449441: expressed sequence AI449441	152.43	274.6	83.3	549.74	575.03	368.81	540.27	415.64	365.12	0.362629281	0.044176707	0.06532798	24.0235
43126	AI553587: expressed sequence AI553587	45.94	59.16	164.25	440.23	390.53	659.33	420.52	199.69	203.51	0.232819462	0.035087719	0.05421053	39.7968
27478	Alkbh1: alkB, alkylation repair homolog 1 (E. coli)	367.13	252.58	364.88	684.73	472.21	782.39	1187.3	1539.4	783.2	0.361366497	0.039408867	0.06277504	25.948
41185	Ankrd11: ankyrin repeat domain 11	194.44	140.3	145.55	605.72	556.3	490.08	635.77	558.13	541.28	0.283567726	0.03076923	0.05533333	38.1328

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
17472	Aoc3: amine oxidase, copper containing 3	14.01	4.63	37.83	85.43	273.68	193.82	107.32	107.65	335.47	0.102359136	0.0390625	0.06083333	30.258
20486	Aspm: asp (abnormal spindle)-like, microcephaly associated (Drosophila)	199.52	137.22	144.24	521.87	545.05	411.51	527.32	557.97	436.97	0.3205796	0.027777778	0.0583642	32.0107
43330	Aspm: asp (abnormal spindle)-like, microcephaly associated (Drosophila)	158.72	62.35	100.75	406.86	421	369.44	451.38	451.4	443.97	0.252998172	0.031746032	0.0562963	38.2554
32443	Aspm: asp (abnormal spindle)-like, microcephaly associated (Drosophila)	292.51	187.37	247.14	625.22	519.02	571.3	712.94	561.32	557.81	0.409864669	0.04787234	0.07400709	20.1586
30016	Atrx: Alpha thalassemia/mental retardation syndrome X-linked homolog (human)	120.99	183.34	102.72	558.59	325.41	576.41	338.6	384.09	383.86	0.317145573	0.048148148	0.06746914	23.1632
15284	AU040096: expressed sequence AU040096	1327.17	938.18	1971.18	2586.3	2194.2	3019.8	2743.2	2441.9	2607.7	0.543384867	0.037914692	0.06336493	25.5363
8675	B230354K17Rik: RIKEN cDNA B230354K17 gene	814.67	430.9	611.56	2576.3	1355.6	1878.6	1425.7	1638.7	1475.3	0.358860153	0.025	0.05741667	43.3599
15916	Bach1: BTB and CNC homology 1	214.4	51.49	267.98	509.03	500.69	370.86	552.06	427.69	703.89	0.348454093	0.038647343	0.06276973	25.7553
8637	Bbs2: Bardet-Biedl syndrome 2 homolog (human)	118.85	61.19	192.91	337.79	236.48	247.56	500.41	586.34	389.87	0.324523048	0.045602606	0.06908795	22.0603
8704	BC003965: cDNA sequence BC003965	318.96	314.61	589.74	1242.2	913.8	906.78	1067.1	675.98	1003.8	0.421130361	0.044692737	0.06277467	26.9326
31179	BF642829: expressed sequence BF642829	55.1	17.77	90.7	564.5	291.99	569.64	719.96	231.69	348.62	0.11998973	0	0.04888889	85.4286
1110	Bub3: budding uninhibited by benzimidazoles 3 homolog (S. cerevisiae)	163.69	146.39	195.19	499.57	444.31	468.08	309.06	361.39	554.54	0.383223042	0.046728972	0.07061267	21.6291
18450	Bxdc1: brix domain containing 1	521.77	121.43	552.63	1496.2	1126.8	1048.1	945.53	776.39	1481.2	0.347916279	0.034482759	0.05551724	39.3557

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
10968	C430004E15Rik: RIKEN cDNA C430004E15 gene	125.61	72.26	122.92	346.64	170.85	293.11	500.16	350.92	598.27	0.283891237	0.038277512	0.06291866	25.6253
27736	C730049O14Rik: RIKEN cDNA C730049O14 gene	279.36	161.54	258.16	854.88	650.64	580.29	733.69	724.56	697.53	0.329621675	0.04	0.05737778	35.8955
34773	Ccdc66: coiled-coil domain containing 66	152.66	109.96	220.57	681.69	370.27	538.09	732.89	447.88	631.69	0.284019738	0.036363636	0.05381818	40.4713
29123	Cdca2: cell division cycle associated 2	209.02	187.77	269.92	719	518.46	652.95	696.51	623.21	617.99	0.348322414	0.032258065	0.06094086	30.5018
41131	Cdca2: cell division cycle associated 2	160.58	179.33	233.01	408.1	505.8	456.62	590.37	765.51	518.81	0.353086549	0.042918455	0.06473534	24.6229
8311	CDNA clone IMAGE:30031514	1064.61	532.91	1115.12	1802.7	1683.2	1977.4	2026.2	1949.5	2208.4	0.465796823	0.031914894	0.05836879	33.0714
4656	Cebpz: CCAAT/enhancer binding protein zeta	801.7	466.96	1043.21	2020.2	1604.2	1858.7	1691.2	1501.9	1690.1	0.446033561	0.028571429	0.05825397	32.2117
15723	Cenpe: centromere protein E	595.49	593.11	496.28	1062.6	840.39	1209.8	1570.6	1201.9	1244.4	0.472642964	0.043290043	0.06494949	24.6295
39617	Ckap2l: cytoskeleton associated protein 2-like	98.96	178.8	150.55	296.31	387.77	321.94	430.71	480.09	416.15	0.367180032	0.047761194	0.07039801	21.3319
16339	Cndp2: CNDP dipeptidase 2 (metallopeptidase M20 family)	95.72	12.15	118.99	542.76	229.08	165.59	150.97	419.23	414.14	0.23609485	0.032786885	0.06057377	30.7138
13932	Cox17: cytochrome c oxidase, subunit XVII assembly protein homolog (yeast)	609.8	99.15	883.83	1560.4	816.01	1231.3	757.93	1492.8	1260	0.447513191	0.04494382	0.06275281	26.9721
1878	Crlz1: charged amino acid rich leucine zipper 1	547.14	492.43	937.54	1867.1	1324.2	1689.6	1664.4	1073.1	1696.2	0.424524879	0.035714286	0.05494048	39.9162
677	Ctsc: cathepsin C	560.59	170.75	535.58	1142.3	623.95	597.3	1072.2	767.24	1113.2	0.476631541	0.047619048	0.07378307	20.1335
10567	Cugbp1: CUG triplet repeat, RNA binding protein 1	144.44	139.03	182.01	475.93	188.63	348.36	686.59	538.56	396.31	0.353388653	0.046822742	0.0693534	22.2407
10566	Cugbp1: CUG triplet repeat, RNA binding protein 1	454.48	148.16	264.74	818.02	671.83	693.42	734.46	490.43	505.7	0.443235067	0.04719764	0.07229105	21.0766

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
41846	D14Abb1e: DNA segment, Chr 14, Abbott 1 expressed	166.7	221.03	266.86	493.13	316.45	653.17	637.61	639.35	503.72	0.40364059	0.047445255	0.06781022	22.983
18143	D3Erd300e: DNA segment, Chr 3, ERATO Doi 300, expressed	651.01	741.63	789.6	1222.2	863.98	1291	1855.7	1684.9	1719.9	0.505282071	0.042635659	0.06616279	23.6601
38498	D530033C11Rik: RIKEN cDNA D530033C11 gene	890.62	74.97	669.69	1354.2	1034.7	1547.4	1172.7	959.68	1175.8	0.45145483	0.043650794	0.06470899	24.0102
29413	Dcun1d3: DCN1, defective in cullin neddylation 1, domain containing 3 (S. cerevisiae)	112.69	125.06	198.24	483.09	350.59	483.79	304.94	421.64	437.94	0.35132293	0.044715447	0.06539295	24.0964
15722	Ddx17: DEAD (Asp-Glu-Ala-Asp) box polypeptide 17	190.78	456.61	194.03	475.57	561.19	524.9	865.28	1016.7	851.49	0.391801878	0.04109589	0.06312024	25.2918
16347	Ddx21: DEAD (Asp-Glu-Ala-Asp) box polypeptide 21	496.86	124.45	757.01	933.21	720.44	1161.8	1068.9	1079.2	1463.9	0.428886151	0.040462428	0.06244701	27.2505
27562	Depdc1b: DEP domain containing 1B	18.58	56.46	148.7	299.93	171.91	324.22	316.42	295.91	377.86	0.250513646	0.038461538	0.06087179	30.1319
14683	Dgke: diacylglycerol kinase, epsilon	343.02	114.18	420.65	1081.8	777.24	733.1	1097.9	905.2	997.74	0.313911367	0.03030303	0.0530303	46.7677
7310	Dnajb11: DnaJ (Hsp40) homolog, subfamily B, member 11	556.18	649.73	756.84	1341.8	1134.7	1077.5	1277.8	1479.7	1531.5	0.5005132	0.046511628	0.06949059	22.185
14950	Eif2c5: eukaryotic translation initiation factor 2C, 5	73.43	7.99	58.25	207.56	115.15	151.62	314.47	175.57	291.88	0.222360199	0.046296296	0.07078189	21.539
22419	Eif3s6: eukaryotic translation initiation factor 3, subunit 6	381.28	298.29	530.64	867.23	627.04	653.69	1252.5	944.86	908.52	0.460694538	0.047021944	0.07048067	21.6691
395	Eif3s7: eukaryotic translation initiation factor 3, subunit 7 (zeta)	29.43	61.47	88.84	294.07	351.69	339.68	230.39	167.87	157.32	0.233274065	0.038961039	0.06056277	28.4566

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
153	Eif3s8: eukaryotic translation initiation factor 3, subunit 8	647.41	374.43	471.72	1108.7	888.34	784.61	1267.7	1026.1	1313.4	0.467559382	0.048327138	0.06748451	23.1855
18771	Eli2: elongation factor RNA polymerase II 2	298.51	184.57	392.63	1119.5	733.54	995.87	939.53	968.34	900.94	0.309562316	0.033333333	0.05511111	47.6055
17713	Etnk1: Ethanolamine kinase 1	159.03	73.22	93.44	636.72	598.27	683.95	479.48	375.04	558.65	0.195485743	0	0.03666667	65.0229
42522	Fancm: Fanconi anemia, complementation group M	235.24	234.96	201.67	561.14	449.79	487.17	710.1	690.61	645.65	0.379109935	0.039215686	0.0625	25.9442
33019	Fancm: Fanconi anemia, complementation group M	324.67	275.04	276.12	624.04	629.44	728.5	955.82	882.68	792.98	0.379684662	0.04137931	0.06165517	28.8393
16073	Fancm: Fanconi anemia, complementation group M	372.48	232.77	365.4	583.78	592.92	710.05	789.6	732.26	922.82	0.448189166	0.048648649	0.07340541	20.3188
14660	Fen1: flap structure specific endonuclease 1	461.57	282.56	398.88	1199.4	1125.7	829.3	821.82	831.08	814.02	0.406671055	0.049056604	0.0674717	23.3419
6438	Fv1: Friend virus susceptibility 1	38.2	81.88	76.52	241.25	95.92	244.58	302.26	159.26	316.04	0.289264406	0.048991354	0.07193084	20.9108
18071	Fzd7: frizzled homolog 7 (Drosophila)	340.48	178.18	288.66	840.79	815.36	819.63	1126.2	951.13	832.39	0.299813573	0.023809524	0.05539683	43.1524
23587	Gapvd1: GTPase activating protein and VPS9 domains 1	129.82	259.4	193.47	424.38	502.98	467.39	685.61	645.29	353.59	0.378463517	0.048710602	0.07160458	20.8995
22636	Gfpt1: glutamine fructose-6-phosphate transaminase 1	38.52	89.06	120.23	335.73	165.1	314.89	566.93	599.99	90.41	0.239077687	0.042328042	0.06405644	26.3748
28204	Glb1: galactosidase, beta 1	1082.26	154.29	824.12	1713.4	1707.9	1919	2551.5	2521.8	2748	0.31313842	0	0.041	74.6783
33270	Gm104: gene model 104, (NCBI)	270.98	111.96	218.82	623.96	423.95	354.11	625.11	416.17	774.69	0.373997433	0.04676259	0.06773381	22.9284
29375	Gm1564: gene model 1564, (NCBI)	200.31	157.46	220.69	547.93	543.11	730.67	543.28	424.1	561.78	0.345259589	0.041420118	0.06234714	27.4591
3500	Gpatc4: G patch domain containing 4	440.58	205.69	292.42	950.82	529.03	789.31	962.71	725.77	902.34	0.386293771	0.034782609	0.05953623	31.3345
9979	Gpatc4: G patch domain containing 4	1300.71	154.78	656.61	2341.4	1228.2	1953.3	2071.8	1558.3	1584.8	0.393399692	0.042553191	0.05496454	42.1856

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
29821	Gtf3c1: general transcription factor III C 1	1082.02	314.5	452.3	1080.6	901.53	1212.7	1452	1192.1	1707.2	0.490006758	0.049608355	0.07422106	20.0083
13878	Gtf3c4: general transcription factor IIIC, polypeptide 4	249.77	12.55	161.14	474.48	220.1	217.44	516.42	384.03	566.54	0.355996822	0.045454545	0.06888528	22.0575
7302	Gtpbp4: GTP binding protein 4	505.74	834.95	1152	1902.8	1140.8	1610.7	1950.4	1522.4	1943.6	0.495037098	0.044025157	0.06255765	28.0427
20122	H47: histocompatibility 47	624.17	411.69	853.39	1601.4	1273.8	1337.6	1610.6	1602.3	1546.1	0.421159518	0.03125	0.05572917	38.1717
16780	H47: histocompatibility 47	1553.67	678.07	1382.64	2610	1889.2	2344.4	2583.7	2481.4	2674.1	0.495709628	0.034482759	0.05862069	33.9839
14429	H47: histocompatibility 47	1937.91	764.71	1187.44	2611	1869.7	2265.5	2327.5	2522.8	2346.6	0.557987889	0.044217687	0.06821995	22.4819
29582	Hectd1: HECT domain containing 1	886.78	469.71	529.57	1198	860.21	1101.7	1757.5	1238.8	1402.2	0.499059992	0.049350649	0.07448485	19.9476
8699	Hipk1: homeodomain interacting protein kinase 1	992.48	402.72	711.08	1281.1	954.65	1185.9	1575.7	1648.7	1861.9	0.495129872	0.045627376	0.06700887	23.4149
8270	Igf2r: insulin-like growth factor 2 receptor	327.8	119.68	339.68	781.98	500.48	705.3	745.85	596	939.69	0.36875366	0.039370079	0.06091864	30.3013
8271	Igf2r: insulin-like growth factor 2 receptor	169.96	242.38	158.23	528.54	492.61	528.94	459.73	649.21	498.88	0.361359253	0.046218487	0.06540616	24.4395
22514	Ilf3: interleukin enhancer binding factor 3	170.07	35.32	121.1	560.62	295.08	376.34	300.48	289.94	360.17	0.299171183	0.046594982	0.06761051	22.9134
7251	Incenp: inner centromere protein	393.46	237.19	218.41	663.52	579.94	476.28	892.65	816.19	920.58	0.390447811	0.042944785	0.06310838	27.6464
21315	Ipo7: importin 7	1777.24	743.47	1888.27	2615.2	2038	2474.9	2454.8	2754.8	2633.6	0.588991329	0.048128342	0.07298574	20.2873
27085	Isg20l2: interferon stimulated exonuclease gene 20-like 2	416.94	227.53	402	1147.7	772.85	990.8	770.46	676.05	841.48	0.402541876	0.044247788	0.06415929	24.9122
4798	Jmy: junction-mediating and regulatory protein	165.05	95.76	208.4	547.49	231.34	463.45	532.88	603.44	674.49	0.30736729	0.036585366	0.05780488	34.7455
4799	Jmy: junction-mediating and regulatory protein	886.39	251.37	566.34	1194	1060.9	1111.5	1676.9	1290.8	1588.2	0.43020715	0.04347826	0.06289855	27.7616
19905	Jmy: junction-mediating and regulatory protein	187.78	141.18	219.35	543.3	293	427.2	657.56	428.2	656.69	0.36481402	0.04609929	0.06806147	22.8305

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
2694	Kctd9: potassium channel tetramerisation domain containing 9	171.19	85.46	175.44	439.95	379.75	353.67	485.57	295.15	503.25	0.351672947	0.044520548	0.0681621	22.5441
20342	Kif11: kinesin family member 11	351.66	218.63	237.15	802.55	400.74	666.11	896.31	538.17	967.58	0.37806277	0.035460993	0.06096927	29.2285
19155	Kif22: kinesin family member 22	291.5	250.11	329.46	841.6	654.47	559	989.59	886.55	947.43	0.35709542	0.028985507	0.05560386	37.1006
7972	Kif22: kinesin family member 22	202.55	190.1	260.35	873.55	734.19	599.86	722.03	806.28	1047.4	0.273030408	0.034482759	0.05172414	48.7204
15135	Kif2c: kinesin family member 2C	167.17	158.49	331.1	532.54	543.91	578.45	476.24	472.46	1022	0.362287382	0.047297297	0.06951577	22.3005
21863	Klf9: Kruppel-like factor 9	1075.11	184.91	618.97	1413.6	1073.9	1024.2	1693.1	992.37	1695.4	0.476142088	0.046242775	0.07190751	20.9288
9017	L2hgdh: L-2-hydroxyglutarate dehydrogenase	356.73	197.34	475.3	1034.2	590.33	965.17	1621.2	1345.6	1417.6	0.295197523	0	0.03680556	58.3346
12344	Larp1 /// LOC631268: La ribonucleoprotein domain family, member 1 /// similar to la related protein isoform 1	347.33	50.62	240.65	510.75	581.65	447.49	684.6	681.6	1010.2	0.326124981	0.028037383	0.05772586	32.144
27871	Lemd3: LEM domain containing 3	355.22	46.19	248.38	639.18	502.36	726.84	660.72	542.9	577.78	0.356070777	0.044871795	0.06168803	28.2329

WO 2010/003020

PCT/US2009/049447

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
20064	Mgat2: mannoside acetylglucosaminyltransferase 2	469.73	304.14	823.62	1323.3	860.58	1140.1	1038.6	1107.9	1146.1	0.482879947	0.045138889	0.06842593	22.6204
19069	Mina: myc induced nuclear antigen	78.98	36.19	157.22	515.36	223.46	260.41	415.61	176.74	570.73	0.251943523	0.035211268	0.06105634	29.1747
21512	Mkrl1: makorin, ring finger protein, 1	648.36	512.8	1294.71	1919.8	1380	1377.3	1555.2	1824.2	2003.2	0.488259093	0.04519774	0.06293785	26.9961
6466	Mm.218588.1	254.43	33.9	245.82	906	787.64	785.29	730.81	656.42	846.69	0.226678125	0	0.04019608	65.2427
12365	Mm.29909.1	281.86	28.14	399.88	829.02	524.34	892.94	1101.1	1108.6	1025.5	0.25901034	0	0.03666667	63.2122
29718	Mnab: membrane associated DNA binding protein	118.68	46.27	176.41	283.79	408.91	313.29	413.11	248.67	694.29	0.289035842	0.040909091	0.06340909	25.2435
8643	Mobk1b: MOB1, Mps One Binder kinase activator-like 1B (yeast)	331.14	291.72	770.54	1112.7	863.99	1204.8	1306.9	1049	1225.8	0.412053466	0.030612245	0.05911565	32.694
7941	Mobk1b: MOB1, Mps One Binder kinase activator-like 1B (yeast)	254.91	242.34	466.56	736.04	605.17	626.58	639.84	1041.8	1058.4	0.409444571	0.047745358	0.07390805	20.1378
791	Mrfap1: Morf4 family associated protein 1	768.71	852.78	1066.33	1873	1359.9	1748.1	1989.4	2140.2	2023.5	0.48280997	0.03539823	0.0600295	31.3818
2811	Mtf2: metal response element binding transcription factor 2	1035.21	1134.68	1619.57	2148.1	1573.3	2029	2913.6	2368.9	2633.3	0.554570812	0.044303797	0.06973629	21.7746
16360	Ndufc1: NADH dehydrogenase (ubiquinone) 1, subcomplex unknown, 1	305.64	283.11	391.61	1442.1	822.36	1025.9	1048.3	788.41	1348.7	0.302783972	0.035714286	0.0497619	49.4498
14278	Nme1: expressed in non-metastatic cells 1, protein	58.62	18.15	143.92	450.69	102.76	119.29	425.45	295.56	236.64	0.270720502	0.048387097	0.07321685	20.3045
23157	Osbpl7: oxysterol binding protein-like 7	208.36	123.05	167.34	291.45	385.78	284.54	615.7	485.5	504.16	0.38856622	0.048843188	0.07529563	19.8199
10534	Oxnad1: oxidoreductase NAD-binding domain containing 1	230.13	169.82	311.6	693.65	782.99	681.59	465.88	438.48	324.58	0.42014425	0.047368421	0.0742193	20.0481
10418	Pank3: pantothenate kinase 3	425.65	241.73	280.88	1151.2	770.81	1060.5	873.11	919.21	1010	0.327845955	0.024390244	0.05658537	43.2753
3006	Pdgfa: platelet derived growth factor, alpha	301.5	109.27	223.39	769.44	1041	424.85	654.37	303.32	583.26	0.335864925	0.03649635	0.06034063	29.6355

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
12377	Pdrg1: p53 and DNA damage regulated 1	372.77	252.95	540.97	715.57	694.98	822.93	1022.5	1047.5	1095.6	0.432177036	0.040201005	0.06324958	26.0102
12376	Pdrg1: p53 and DNA damage regulated 1	285.31	282.96	545.59	789.75	643.57	739.91	846.21	866	924.2	0.463178117	0.048850575	0.0717433	20.9062
727	Pfkfb3: 6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 3	39.57	6.24	92.23	213.6	178.27	137.99	367.86	210.92	349.81	0.189296856	0.040935673	0.06185185	27.429
14762	Pgd: phosphogluconate dehydrogenase	929.25	508.03	877.51	1851.3	1253.4	1745.2	1403.8	1529.3	1992.3	0.473598818	0.039473684	0.06089912	28.5711
15044	Pgd: phosphogluconate dehydrogenase	864.91	479.01	841.06	1775	1198.1	1646	1302.6	1437.5	1852.4	0.474396469	0.042168675	0.06273092	27.5296
15538	Pgd: phosphogluconate dehydrogenase	641.54	351.22	569.81	1194	750.15	1094.5	1037.2	1017.5	1353.6	0.48475081	0.046376812	0.07173913	20.9781
28991	Plcd4: phospholipase C, delta 4	85.67	50.61	173.34	335.85	224.94	209.08	437.46	324.88	403.75	0.319861981	0.045454545	0.06839161	22.7015
16204	Ppqb: protective protein for beta-galactosidase	440.09	225.35	350.75	1117.8	569.65	1350.3	1365.1	642.54	1356.9	0.317445293	0	0.03811594	58.4023
1352	Ppid /// Lamp3 /// LOC671770: peptidylprolyl isomerase D (cyclophilin D) /// lysosomal-associated membrane protein 3 /// similar to peptidylprolyl isomerase D	177.09	265.27	578.82	1140.3	699.01	928.31	1030.2	754.1	902.76	0.37442549	0.032608696	0.05804348	33.4589
39148	Ppmla: protein phosphatase 1A, magnesium dependent, alpha isoform	678.05	348.48	381.66	1276.6	1065.7	1257	1597.5	1131.1	1120.3	0.378124375	0.027777778	0.05555556	36.5516
9696	Ppmla: protein phosphatase 1A, magnesium dependent, alpha isoform	132.05	88.38	124.3	565.15	304.18	544.53	369.01	388.93	396.41	0.268459355	0.033333333	0.06136111	30.7357
32940	Ppp1r3e: Protein phosphatase 1, regulatory (inhibitor) subunit 3E	5.81	13.38	13.14	172.3	123.7	36.66	111.76	102.91	75.79	0.103768135	0.049723757	0.07308471	20.499

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
15897	Pramel7: preferentially expressed antigen in melanoma like 7	228.49	144.58	867.63	970.28	608.23	1186	963.95	982.9	977.84	0.436162808	0.038095238	0.06279365	25.6083
15898	Pramel7: preferentially expressed antigen in melanoma like 7	883.77	273.04	670.2	1610.9	1098.7	1406.8	1266	1347.1	1586.8	0.439380494	0.03875969	0.06098191	30.2002
7006	Prkcd: protein kinase C, delta	235.8	93.35	81.71	712.82	315.54	733.67	910.26	661.36	733.23	0.202051696	0	0.03909091	73.8504
9218	Prmt6: protein arginine N-methyltransferase 6	738.14	448.83	800.51	1347.3	1072.8	1130.9	1415.4	1423.3	1354.9	0.513255025	0.044827586	0.06847126	22.56
41444	Prpf19: PRP19/PSO4 pre-mRNA processing factor 19 homolog (S. cerevisiae)	3993.02	2736.93	2277.87	4887.6	4659.6	4958.5	4053.2	4990.6	4177.1	0.64976133	0.04664723	0.07188533	21.0008
33047	Prr11: proline rich 11	401.64	173.26	234.39	568.15	688.18	606.56	817.5	1134	740.86	0.355318761	0.036231884	0.06014493	29.5659
35103	Prr11: proline rich 11	265.25	149.08	442.11	729.69	571.34	733.59	873.03	564.31	1014.1	0.381823653	0.040697674	0.06170543	27.3805
21158	Psat1: phosphoserine aminotransferase 1	608.23	383.49	505.63	1268.1	1111.2	1110.2	1163.3	1113	1052.3	0.439232447	0.043010753	0.06390681	26.5312
14945	Psma6: proteasome (prosome, macropain) subunit, alpha type 6	297.18	193.17	579.11	823.99	814.83	842.51	1149.9	1450.2	1653.8	0.31757146	0.039215686	0.05627451	41.0024
14001	Ptdss2: phosphatidylserine synthase 2	286.3	79.67	434.04	847.59	360.31	528.7	764.33	553.32	801.99	0.414917121	0.045901639	0.06926776	22.0925
3935	Purb: purine rich element binding protein B	255.68	124.63	215.26	618.23	457.69	488.12	438.8	529.05	494.18	0.393626056	0.04851752	0.07339623	20.3063
16286	Rab1: RAB1, member RAS oncogene family	24.78	29.32	63.21	200.41	213.31	158.37	260.27	168.49	186.44	0.197609683	0.037735849	0.06314465	25.5279
15338	Rab20: RAB20, member RAS oncogene family	448.47	171.13	416.25	593.19	602.78	611.39	1207	970.25	1217.6	0.398230769	0.045977011	0.06689655	23.5179
4228	Rbm35b: RNA binding motif protein 35b	187.48	63.54	221.76	590.29	628.16	453.65	363.16	321	412.3	0.34153495	0.047619048	0.06794872	22.9952
12058	Rnf6: ring finger protein (C3H2C3 type) 6	508.38	239.28	418.61	655.12	583.25	868.77	1096.2	908.12	1006.9	0.455713959	0.047794118	0.06800245	23.0191
12057	Rnf6: ring finger protein (C3H2C3 type) 6	861.12	396.56	614.69	1246.8	778.27	939.73	1510	1251.2	1413.2	0.524537304	0.049095607	0.07514212	19.8721

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
1293	Rrs1: RRS1 ribosome biogenesis regulator homolog (S. cerevisiae)	618.8	397.87	439.42	1053.2	649.43	1070.2	1408.1	1440.1	939.09	0.443918374	0.042372881	0.06426554	24.5877
8403	Rwdd4a: RWD domain containing 4A	120.1	127.06	198.13	517.17	347.78	401.78	607.87	348.3	679.49	0.30684367	0.04109589	0.06171233	28.8054
20481	Sdad1: SDA1 domain containing 1	850.27	363.33	872.76	2204.8	1724.2	2050.9	2051.9	2375.7	1968.4	0.337163067	0	0.03539683	62.6786
15545	Sf3b2: splicing factor 3b, subunit 2	805.59	473.95	826.78	1604.5	2014.8	1868.8	1091.8	1101.2	1830	0.442914549	0.028169014	0.05497653	36.8781
12258	Sfrs1: splicing factor, arginine/serine-rich 1 (ASF/SF2)	1313.28	772.48	1279.84	2058.8	1861	1951.5	1990.7	2140.9	2193.1	0.551915042	0.048780488	0.07106707	21.4237
4980	Sgpp1: sphingosine-1-phosphate phosphatase 1	171.97	94.46	279.34	1215.3	589.36	577.14	509.63	536.21	763.86	0.260417512	0.032258065	0.05677419	38.3871
4981	Sgpp1: sphingosine-1-phosphate phosphatase 1	1314.88	188.41	379.99	1556.5	1386.4	1522.7	1138.2	1603.8	1205.4	0.447708184	0.033613445	0.06098039	30.8513
18042	Sgpp1: sphingosine-1-phosphate phosphatase 1	31.01	17.03	39.22	176.04	188.98	215.43	109.27	132.56	153.26	0.178895791	0.044354839	0.06548387	24.0389
1917	Slc12a2: solute carrier family 12, member 2	437.13	331.4	303.71	1012.6	574.08	711.55	831.2	898.12	1082.7	0.419649444	0.041450777	0.06310881	26.2992
16856	Slc12a2: solute carrier family 12, member 2	153.39	179.89	175.06	527.45	337.21	487.48	447.9	572.27	515.56	0.352051858	0.043715847	0.06347905	26.6694
21716	Slc1a4: solute carrier family 1 (glutamate/neutral amino acid transporter), member 4	60.95	35.65	114.42	292.45	192.95	238.64	177.44	338.73	362.16	0.263384861	0.043668122	0.06537118	24.6402
28457	Slc30a1: solute carrier family 30 (zinc transporter), member 1	317.74	103.14	73.18	315.4	343.75	619.79	393.91	491.5	503.8	0.370338999	0.047619048	0.07036706	21.3192
23716	Sltn: SAFB-like, transcription modulator	370.13	284.47	502.78	1011.1	760.09	1052	1179.5	795.6	946.04	0.402964314	0.034188034	0.06111111	30.8935

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
7575	Smarca1: SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily c, member 1	97.38	219.78	155.82	345.56	264.16	208.33	557.93	543.22	522.86	0.38736149	0.049222798	0.0747323	19.9146
6834	Smarca1: SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily e, member 1	873.62	915.47	718.44	1533.2	1416.9	1757.6	1787.9	1926.5	1710.1	0.49496406	0.041666667	0.06248016	27.4725
6044	Sos1: Son of sevenless homolog 1 (Drosophila)	134.91	61.07	109.98	438.22	273.17	373.78	444.95	294.97	346.44	0.281792101	0.039735099	0.06119205	28.5896
6045	Sos1: Son of sevenless homolog 1 (Drosophila)	391.93	223.27	270.95	684.56	368.73	779.87	937.33	493.55	698.66	0.447245565	0.048913043	0.07344203	20.356
27178	Spata2: spermatogenesis associated 2	1669.65	719.38	963.1	2142.9	1761.6	1953.7	1664.5	2350.8	2119.4	0.559020019	0.047477745	0.07050445	21.2734
15690	Spint2: serine protease inhibitor, Kunitz type 2	120.15	171.41	256.46	612.14	318.08	490.9	552.94	367.02	613.78	0.370927895	0.041284404	0.06328746	25.2973
21622	Stt3a: STT3, subunit of the oligosaccharyltransferase complex, homolog A (S. cerevisiae)	295.43	563.39	363.82	1099.3	829.32	1154.4	987.79	752.3	707.18	0.442161261	0.045801527	0.06703562	23.437
6577	suppressor of Ty 4 homolog 1 (S. cerevisiae) /// suppressor of Ty 4 homolog 2 (S. cerevisiae)	782.52	691.08	832.99	1708.4	1408.4	1683.3	1689.4	2017.7	2324.6	0.425897785	0.034090909	0.0580303	33.971
1745	Tacc3: transforming, acidic coiled-coil containing protein 3	738.85	414.57	618.79	1661.2	1187.9	1268.8	1516.9	1282.6	1702	0.411215669	0.035294118	0.05729412	34.3614
13841	Tbl2: transducin (beta)-like 2	293.54	36.54	247.36	684.47	780.75	476.82	685.89	393.26	569.1	0.321667609	0.0375	0.057375	35.1428
41362	Terf2ip: telomeric repeat binding factor 2, interacting protein	127.85	108.83	96.14	286.88	330.88	351.36	345.81	519.43	330.16	0.307523146	0.04797048	0.06747847	23.1137
182	Tfg: Trk-fused gene	388.74	156.37	629.82	928.5	564.98	679.38	871.81	1022.6	1079	0.456614208	0.046428571	0.0677619	22.8893

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
7468	Tgoln1: trans-golgi network protein	305.53	236.42	345.51	671.72	450.68	487.94	700.65	843.07	702.27	0.46026144	0.047493404	0.0739314	20.0806
3061	Timd2: T-cell immunoglobulin and mucin domain containing 2	608.39	151.76	231.44	1866.9	1351.1	1401.1	1475.4	1242.5	1993.4	0.212552423	0	0.06166667	97.4881
1794	Timm10: translocase of inner mitochondrial membrane 10 homolog (yeast)	207.76	55.6	617.54	1237.7	550.92	497.86	560.37	1173.8	1446.2	0.322267911	0.038961039	0.05835498	35.3576
785	Tmed6: transmembrane emp24 protein transport domain containing 6	254.44	20.44	279.48	508.83	418.62	577.06	745.64	731.85	729.56	0.298720754	0.037037037	0.05450617	40.4979
23223	Tmpo: thymopoietin	372.55	239.05	785	1127.4	1035.5	974.18	1028.2	951.54	1084.8	0.450399171	0.042735043	0.06450142	24.6087
13926	Tomm22: translocase of outer mitochondrial membrane 22 homolog (yeast)	1013.62	632.86	904.75	2273.2	1748.6	1962.2	1195.2	1371.2	1705.4	0.497524304	0.040609137	0.06323181	26.1029
16924	Tor1b: torsin family 1, member B	272.73	156.58	572.98	1028.7	649.17	887.96	740.56	964.62	1195.4	0.36670801	0.037974684	0.05776371	35.2
21932	Transcribed locus	332.77	106.56	140.03	583.75	309.89	572.13	653.87	636.22	766.49	0.328962199	0.032258065	0.0578853	33.3722
43944	Transcribed locus	53.89	21.65	39.31	224.22	170.32	140.35	49.01	235.17	219.21	0.221231267	0.046153846	0.06666667	23.5864
24471	Tsc22d2: TSC22 domain family 2	936.98	613.69	879.83	1530.6	1318.9	1453.7	2101.3	2010.4	1899.3	0.47129429	0.035971223	0.06131894	29.3405
10769	Ttfl: transcription termination factor 1	42.37	81.67	164.99	294.59	226.16	368.82	336.41	367.68	251.53	0.313279391	0.045936396	0.06791519	22.8092
14944	Txndc1: thioredoxin domain containing 1	424.55	133.69	462.39	960.94	645.2	984.25	612.43	1057.4	631.05	0.417324638	0.045454545	0.06567493	24.2087
22950	Unkl: unkempt-like (Drosophila)	270.84	139.44	154.65	747.39	286.66	573.67	677.07	831.48	565.56	0.306874571	0.028571429	0.05490476	37.0961
16306	Usp10: ubiquitin specific peptidase 10	173.21	255.12	344.42	838.32	498.36	592.08	738.05	449.15	994.08	0.376030404	0.041025641	0.06345299	26.1651
12733	Utp11: UTP11-like, U3 small nucleolar ribonucleoprotein, (yeast)	955.84	514.61	1811.24	2092.8	2024.9	1852.2	1938.2	1738.6	2374.9	0.545964687	0.046920821	0.07209189	21.0368

Table 9B

Row	Gene	CycA2.1	CycA2.2	CycA2.3	NI1.1	NI1.2	NI1.3	NI2.1	NI2.2	NI2.3	FoldChange	MedianFDR	MeanFDR	T
27185	Utp15: UTP15, U3 small nucleolar ribonucleoprotein, homolog (yeast)	222.46	131.63	302.63	1588.6	802.68	700.69	623.36	1475.8	537.97	0.22925845	0.037037037	0.0491358	50.5511
15510	Wbscr1: Williams-Beuren syndrome chromosome region 1 homolog (human)	117.48	136.2	356.85	655.34	402.2	299.57	683.74	452.6	930.59	0.356613825	0.046025105	0.06517434	24.4325
26959	Wdr3: WD repeat domain 3	772.19	388.92	483.21	1122.9	1121.4	1247.2	1065.7	962.84	1081.4	0.498168587	0.049046322	0.07336966	20.3716
21792	Zfp444: Zinc finger protein 444	75.01	31.45	145.74	452.64	198.57	309.81	294.53	537.31	689.29	0.203210926	0.041666667	0.055	41.8997
39491	Zfp710: zinc finger protein 710	371.26	333.7	313.27	900.66	784.47	1074.7	975.36	1333.3	979.43	0.336722951	0.037735849	0.05540881	40.5004

CLAIMS

That which is claimed is:

1. A method of evaluating a female subject for an in vitro fertilization procedure, comprising:
 - obtaining items of information from a female subject to provide a profile for said female subject, wherein each item of information relate to preselected patient variables;
 - comparing said profile for said female subject to a library of known profile patterns known to be indicative of responsiveness to an in vitro fertilization procedure using an algorithm based upon said preselected patient variables,
 - wherein said comparing provides an evaluation of said female subject for an in vitro fertilization procedure.
2. The method of claim 1, wherein said in vitro fertilization procedure is at least a second vitro fertilization procedure for said female subject.
3. The method of claim 1, wherein said in vitro fertilization procedure comprises determining a number of embryos for implantation.
4. The method of claim 1, wherein said in vitro fertilization procedure provides for a live birth event.
5. The method of claim 1, wherein said female subject is a pre-in vitro fertilization (pre-IVF) procedure patient.
6. The method of claim 5, wherein said preselected patient variables comprises age, diminished ovarian reserve, 3 follicle stimulating hormone (FSH) level, body mass index, polycystic ovarian disease, season, unexplained female infertility, number of spontaneous miscarriages, year, other causes of female infertility, number of previous pregnancies, number of previous term deliveries, endometriosis, tubal disease, tubal ligation, male infertility, uterine fibroids, hydrosalpinx, or male infertility causes.
7. The method of claim 1, wherein said female subject is a pre-surgical (pre-OR) procedure patient.

8. The method of claim 7, wherein said preselected patient variables comprises age, endometrial thickness, total number of oocytes, total amount of gonatropins administered, number of total motile sperm after wash, number of total motile sperm before wash, day 3 follicle stimulating hormone (FSH) level, body mass index, sperm collection, age of spouse, season number of spontaneous miscarriages, unexplained female infertility, number of previous term deliveries, year, number of previous pregnancies, other causes of female infertility, endometriosis, male infertility, tubal ligation, polycystic ovarian disease, tubal disease, sperm from donor, hydrosalpinx, uterine fibroids, or male infertility causes.

9. The method of claim 1, wherein said female subject is a post-in vitro fertilization (post-IVF) procedure patient.

10. The method of claim 9, wherein said preselected patient variables comprises blastocyst development rate, total number of embryos, total amount of gonatropins administered, endometrial thickness, flare protocol, average number of cells per embryo, type of catheter used, percentage of 8-cell embryos transferred, day 3 follicle stimulating hormone (FSH) level, body mass index, number of motile sperm before wash, number of motile sperm after wash, average grade of embryos, day of embryo transfer, season, number of spontaneous miscarriages, number of previous term deliveries, oral contraceptive pills, sperm collection, percent of unfertilized eggs, number of embryos arrested at 4-cell stage, compaction on day 3 after transfer, percent of normal fertilization, percent of abnormally fertilized eggs, percent of normal and mature oocytes, number of previous pregnancies, year, polycystic ovarian disease, unexplained female infertility, tubal disease, male infertility only, male infertility causes, endometriosis, other causes of female infertility, uterine fibroids, tubal ligation, sperm from donor, hydrosalpinx, performance of ICSI, or assisted hatching.

11. The method of claim 1, wherein said method comprises obtaining items of information relating to at least two preselected patient variables.

12. The method of claim 1, wherein said method comprises obtaining items of information relating to at least five preselected patient variables.

13. The method of claim 12, wherein preselected patient variables is given a weighted relative importance in relation to other preselected patient variables.

14. The method of claim 1, wherein said preselected patient variables comprises age, total number of embryos, rate of cleavage arrest, number of 8-cell embryos, day 3 follicle stimulating hormone (FSH) level, number of 8-cell embryos transferred.

15. The method of claim 1, further comprising determining an expression level of one or more infertility indicator genes and comparing said expression level to a control gene expression profile known to be indicative of responsiveness to an in vitro fertilization procedure.

16. The method of claim 15, wherein said infertility indicator genes comprise Oct4, Eif3c, Papola, Piwil2, Eif3b, Eif4b, Rbm3, and Cpsf4.

17. The method of claim 15, wherein said expression level of one or more infertility indicator genes is done by using a BioMark® Dynamic Array, RT-PCR, or whole whole-transcriptome sequencing.

18. The method of claim 15, wherein said expression level of one or more infertility indicator genes is determined for an embryo.

19. The method of claim 18, wherein said embryo is in arrested development following fertilization.

20. The method of claim 15, wherein said expression level of one or more infertility indicator genes is determined for an oocyte.

21. The method of claim 20, wherein said oocyte is in arrested development following fertilization.

22. The method of claim 1, wherein said comparing said profile for said female subject comprises applying a decision rule.

23. The method of claim 1, wherein said data analysis algorithm comprises use of a classification tree.

24. The method of claim 1, wherein said data analysis algorithm is nonparametric.

25. The method of claim 24, wherein said data analysis algorithm detects differences in a distribution of feature values.

26. The method of claim 25, wherein said nonparametric algorithm comprises using a Wilcoxon Signed Rank Test.

27. The method of claim 1, wherein said data analysis algorithm comprises using a multiple additive regression tree or by gradient boosted machine (GBM).

28. The method of claim 1, wherein said data analysis algorithm is a logistic regression.

29. A method of evaluating a female subject for an in vitro fertilization procedure, comprising:

obtaining items of information from a female subject to provide a profile for said female subject, wherein each item of information relate to preselected patient variables;

determining an expression level of one or more infertility indicator genes;

comparing said profile for said female subject to a library of known profile patterns known to be indicative of responsiveness to an in vitro fertilization procedure using an algorithm based upon said preselected patient variables; and

comparing said expression level of said one or more infertility indicator genes to a control gene expression profile known to be indicative of responsiveness to an in vitro fertilization procedure

wherein said comparing said profile for said female subject and said expression level of said one or more infertility indicator genes provides an evaluation of said female subject for an in vitro fertilization procedure.

30. The method of claim 29, wherein said in vitro fertilization procedure is at least a second vitro fertilization procedure for said female subject.

31. The method of claim 29, wherein said in vitro fertilization procedure comprises a determining a number of embryos for implantation.

32. The method of claim 29, wherein said in vitro fertilization procedure provides for a live birth event.

33. The method of claim 29, wherein said in vitro fertilization procedure provides for a live birth event.

34. The method of claim 29, wherein said female subject is a pre-in vitro fertilization (pre-IVF) procedure patient.

35. The method of claim 34, wherein said preselected patient variables comprises age, diminished ovarian reserve, 3 follicle stimulating hormone (FSH) level, body mass index, polycystic ovarian disease, season, unexplained female infertility, number of spontaneous miscarriages, year, other causes of female infertility, number of previous pregnancies, number of previous term deliveries, endometriosis, tubal disease, tubal ligation, male infertility, uterine fibroids, hydrosalpinx, or male infertility causes.

36. The method of claim 29, wherein said female subject is a pre-surgical (pre-OR) procedure patient.

37. The method of claim 36, wherein said preselected patient variables comprises age, endometrial thickness, total number of oocytes, total amount of gonatropins administered, number of total motile sperm after wash, number of total motile sperm before wash, day 3 follicle stimulating hormone (FSH) level, body mass index, sperm collection, age of spouse, season number of spontaneous miscarriages, unexplained female infertility, number of previous term deliveries, year, number of previous pregnancies, other causes of female infertility, endometriosis, male infertility, tubal ligation, polycystic ovarian disease, tubal disease, sperm from donor, hydrosalpinx, uterine fibroids, or male infertility causes.

38. The method of claim 29, wherein said female subject is a post-in vitro fertilization (post-IVF) procedure patient.

39. The method of claim 38, wherein said preselected patient variables comprises blastocyst development rate, total number of embryos, total amount of gonatropins administered, endometrial thickness, flare protocol, average number of cells per embryo, type of catheter used, percentage of 8-cell embryos transferred, day 3 follicle stimulating hormone (FSH) level, body mass index, number of motile sperm before wash, number of motile sperm after wash, average grade of embryos, day of embryo transfer, season, number of spontaneous miscarriages, number of previous term deliveries, oral contraceptive pills, sperm collection, percent of unfertilized eggs, number of embryos arrested at 4-cell stage, compaction on day 3 after transfer, percent of normal fertilization,

percent of abnormally fertilized eggs, percent of normal and mature oocytes, number of previous pregnancies, year, polycystic ovarian disease, unexplained female infertility, tubal disease, male infertility only, male infertility causes, endometriosis, other causes of female infertility, uterine fibroids, tubal ligation, sperm from donor, hydrosalpinx, performance of ICSI, or assisted hatching.

40. The method of claim 29, wherein said method comprises obtaining items of information relating to at least two preselected patient variables.

41. The method of claim 29, wherein said method comprises obtaining items of information relating to at least five preselected patient variables.

42. The method of claim 41, wherein preselected patient variables is given a weighted relative importance in relation to other preselected patient variables.

43. The method of claim 29, wherein said preselected patient variables comprises age, total number of embryos, rate of cleavage arrest, number of 8-cell embryos, day 3 follicle stimulating hormone (FSH) level, and number of 8-cell embryos transferred.

44. The method of claim 29, wherein said infertility indicator genes comprise Oct4, Eif3c, Papola, Piwi12, Eif3b, Eif4b, Rbm3, and Cpsf4.

45. The method of claim 29, wherein said expression level of one or more infertility indicator genes is done by using a BioMark® Dynamic Array, RT-PCR, or whole whole-transcriptome sequencing.

46. The method of claim 29, wherein said expression level of one or more infertility indicator genes is determined for an embryo.

47. The method of claim 46, wherein said embryo is in arrested development following fertilization.

48. The method of claim 29, wherein said expression level of one or more infertility indicator genes is determined for an oocyte.

49. The method of claim 48, wherein said embryo is in arrested development following fertilization.

50. The method of claim 29, wherein said comparing said profile for said female subject comprises applying a decision rule.

51. The method of claim 29, wherein said data analysis algorithm comprises use of a classification tree.

52. The method of claim 29, wherein said data analysis algorithm is nonparametric.

53. The method of claim 52, wherein said data analysis algorithm detects differences in a distribution of feature values.

54. The method of claim 52, wherein said nonparametric algorithm comprises using a Wilcoxon Signed Rank Test.

55. The method of claim 29, wherein said data analysis algorithm comprises using a multiple additive regression tree or by gradient boosted machine (GBM).

56. The method of claim 29, wherein said data analysis algorithm is a logistic regression.

57. A system for analysis of data in evaluating a female subject for an in vitro fertilization procedure, comprising:

a computing environment;

an input device, connected to the computing environment, to receive data from a user, wherein the data received comprises items of information from a female subject to provide a profile for said female subject, wherein each item of information relate to preselected patient variables,

an output device, connected to the computing environment, to provide information to the user; and

a computer readable storage medium having stored thereon at least one algorithm to provide for comparing the profile for the female subject to a library of known profile patterns known to be indicative of responsiveness to an in vitro fertilization procedure,

wherein the system provides results that can be used for generation of a report providing an analysis of data in evaluating a female subject for an in vitro fertilization procedure.

58. The system of claim 57, wherein the computing environment comprises a local computer local to the user and a remote computer at a site remote to the user, wherein the local computer and the remote computer are connected through a network, and wherein the computer readable storage medium is provided on the remote computer.

59. The system of claim 57, wherein said input device is a personal computing device or a Smartphone device.

60. The system of claim 57, wherein said connection of the input device to the computing device is via a wired connection or a wireless connection.

61. The system of claim 57, wherein said wireless connection is via a wireless LAN connection, Bluetooth connection protocol, ZigBee connection protocol, radio-frequency connection protocol, or a cellular phone connection protocol.

62. The system of claim 61, wherein said cellular phone connection is via code derived multiple access (CDMA) or via a global system for mobile communication (GSM).

63. The system of claim 57, wherein said in vitro fertilization procedure is at least a second vitro fertilization procedure for said female subject.

64. The system of claim 57, wherein said in vitro fertilization procedure comprises a determining a number of embryos for implantation.

65. The system of claim 57, wherein said in vitro fertilization procedure provides for a live birth event.

66. The system of claim 57, wherein said in vitro fertilization procedure provides for a live birth event.

67. The system of claim 57, wherein said female subject is a pre-in vitro fertilization (pre-IVF) procedure patient.

68. The system of claim 57, wherein said preselected patient variables comprises age, diminished ovarian reserve, 3 follicle stimulating hormone (FSH) level, body mass index, polycystic ovarian disease, season, unexplained female infertility, number of spontaneous miscarriages, year, other causes of female infertility, number of previous pregnancies, number of previous term deliveries, endometriosis, tubal disease, tubal ligation, male infertility, uterine fibroids, hydrosalpinx, or male infertility causes.

69. The system of claim 57, wherein said female subject is a pre-surgical (pre-OR) procedure patient.

70. The system of claim 69, wherein said preselected patient variables comprises age, endometrial thickness, total number of oocytes, total amount of gonatropins administered, number of total motile sperm after wash, number of total motile sperm before wash, day 3 follicle stimulating hormone (FSH) level, body mass index, sperm collection, age of spouse, season number of spontaneous miscarriages, unexplained female infertility, number of previous term deliveries, year, number of previous pregnancies, other causes of female infertility, endometriosis, male infertility, tubal ligation, polycystic ovarian disease, tubal disease, sperm from donor, hydrosalpinx, uterine fibroids, or male infertility causes.

71. The system of claim 57, wherein said female subject is a post-in vitro fertilization (post-IVF) procedure patient.

72. The system of claim 71, wherein said preselected patient variables comprises blastocyst development rate, total number of embryos, total amount of gonatropins administered, endometrial thickness, flare protocol, average number of cells per embryo, type of catheter used, percentage of 8-cell embryos transferred, day 3 follicle stimulating hormone (FSH) level, body mass index, number of motile sperm before wash, number of motile sperm after wash, average grade of embryos, day of embryo transfer, season, number of spontaneous miscarriages, number of previous term deliveries, oral contraceptive pills, sperm collection, percent of unfertilized eggs, number of embryos arrested at 4-cell stage, compaction on day 3 after transfer, percent of normal fertilization, percent of abnormally fertilized eggs, percent of normal and mature oocytes, number of previous pregnancies, year, polycystic ovarian disease, unexplained female infertility, tubal disease, male infertility only, male infertility causes, endometriosis, other causes of female

infertility, uterine fibroids, tubal ligation, sperm from donor, hydrosalpinx, performance of ICSI, or assisted hatching.

73. The system of claim 57, wherein said method comprises obtaining items of information relating to at least two preselected patient variables.

74. The system of claim 57, wherein said method comprises obtaining items of information relating to at least five preselected patient variables.

75. The system of claim 74, wherein preselected patient variables is given a weighted relative importance in relation to other preselected patient variables.

76. The system of claim 57, wherein said preselected patient variables comprises age, total number of embryos, rate of cleavage arrest, number of 8-cell embryos, day 3 follicle stimulating hormone (FSH) level, and number of 8-cell embryos transferred.

77. The system of claim 57, wherein the data received further comprises an expression level of one or more infertility indicator genes.

78. The system of claim 77, wherein said infertility indicator genes comprise Oct4, Eif3c, Papola, Piwil2, Eif3b, Eif4b, Rbm3, and Cpsf4.

79. The system of claim 77, wherein said expression level of one or more infertility indicator genes is done by using a BioMark® Dynamic Array, RT-PCR, or whole whole-transcriptome sequencing.

80. The system of claim 77, wherein said expression level of one or more infertility indicator genes is determined for an embryo.

81. The system of claim 80, wherein said embryo is in arrested development following fertilization.

82. The system of claim 77, wherein said expression level of one or more infertility indicator genes is determined for an oocyte.

83. The system of claim 82, wherein said embryo is in arrested development following fertilization.

84. The system of claim 57, wherein said comparing comprises applying a decision rule.

85. The system of claim 57, wherein said data analysis algorithm comprises use of a classification tree.

86. The system of claim 57, wherein said data analysis algorithm is nonparametric.

87. The system of claim 86, wherein said data analysis algorithm detects differences in a distribution of feature values.

88. The system of claim 86, wherein said nonparametric algorithm comprises using a Wilcoxon Signed Rank Test.

89. The system of claim 57, wherein said data analysis algorithm comprises using a multiple additive regression tree or by gradient boosted machine (GBM).

90. The system of claim 57, wherein said data analysis algorithm is a logistic regression.