

SYNTHESIS OF TETRACYCLINES AND ANALOGUES THEREOF

Related Applications

[0001] The present application claims priority under 35 U.S.C. § 119(e) to U.S. provisional patent applications, USSN 60/660,947, filed March 11, 2005, and USSN 60/573,623, filed May 21, 2004, each of which is incorporated herein by reference.


Government Support

[0002] The work described herein was supported, in part, by grants from the National Institutes of Health (R01 AI48825) and the National Science Foundation (predoctoral fellowship R10964). The United States government may have certain rights in the invention.

Background of the Invention

[0001] The tetracyclines are broad spectrum anti-microbial agents that are widely used in human and veterinary medicine (Schappinger *et al.*, "Tetracyclines: Antibiotic Action, Uptake, and Resistance Mechanisms" *Arch. Microbiol.* 165:359-69, 1996; Mitscher, *Medicinal Research Series*, Vol. 9, The Chemistry of the Tetracycline Antibiotics, Marcel Dekker Inc. New York, 1978). The total production of tetracyclines by fermentation or semi-synthesis is measured in the thousands of metric tons per year. The first tetracycline, chlorotetracycline (1) (Aureomycin™), was isolated from the soil bacterium *Streptomyces aureofaciens* by Lederle Laboratories (Wyeth-Ayerst Research) in the 1945 (Duggar, *Ann. N.Y. Acad. Sci.* 51:177-181, 1948; Duggar, Aureomycin and Preparation of Some, U.S. Patent 2,482,055, 1949; incorporated herein by reference). Oxytetracycline (2) was isolated soon after from *S. rimosus* by scientists at Pfizer Laboratories (Finlay *et al. Science* 111:85, 1950). The structures of chlorotetracycline and oxytetracycline were elucidated by scientists at Pfizer in collaboration with R. B. Woodward and co-workers at Harvard University (Hochstein *et al. J. Am. Chem. Soc.* 74:3708-3709, 1952; Hochstein *et al. J. Am. Chem. Soc.* 75:5455-75, 1953; Stephens *et al. J. Am. Chem. Soc.* 74:4976-77, 1952; Stephens *et al. J. Am. Chem. Soc.* 76:3568-75, 1954). Tetracycline (3) was later prepared by the


hydrogenolysis of chlorotetracycline and was found to retain the anti-microbial activity of chlorotetracycline and oxytetracycline and had increased stability (Boothe *et al. J. Am. Chem. Soc.* 75:4621, 1953; Conover *et al. J. Am. Chem. Soc.* 75:4622-23, 1953). Tetracycline was later found to be a natural product of *S. aureofaciens*, *S. viridofaciens*, and *S. rimosus*.



[0002] The primary tetracyclines of clinical importance today include tetracycline (3) (Boothe *et al. J. Am. Chem. Soc.* 75:4621, 1953), oxytetracycline (2, Terramycin™) (Finlay *et al. Science* 111:85, 1950), doxycycline (Stephens *et al. J. Am. Chem. Soc.* 85:2643, 1963), and minocycline (Martell *et al. J. Med. Chem.* 10:44, 1967; Martell *et al. J. Med. Chem.* 10:359, 1967). The tetracyclines exert their anti-microbial activity by inhibition of bacterial protein synthesis (Bentley and O'Hanlon, Eds., *Anti-Infectives: Recent Advances in Chemistry and Structure-Activity Relationships* The Royal Society of Chemistry: Cambridge, UK, 1997). Most tetracyclines are bacteriostatic rather than bactericidal (Rasmussen *et al. Antimicrob. Agents Chemother.* 35:2306-11, 1991; Primrose and Wardlaw, Ed. "The Bacteriostatic and Bacteriocidal Action of Antibiotics" *Sourcebook of Experiments for the Teaching of Microbiology* Society for General Microbiology, Academic Press Ltd., London, 1982). It has been proposed that after tetracycline passes through the cytoplasmic membrane of a bacterium it chelates Mg^{+2} , and this tetracycline- Mg^{+2} complex binds the 30S subunit

of the bacterial ribosome (Goldman *et al. Biochemistry* 22:359-368, 1983). Binding of the complex to the ribosome inhibits the binding of aminoacyl-tRNAs, resulting in inhibition of protein synthesis (Wissmann *et al. Forum Mikrobiol.* 292-99, 1998; Epe *et al. EMBO J.* 3:121-26, 1984). Tetracyclines have also been found to bind to the 40S subunit of eukaryotic ribosome; however, they do not achieve sufficient concentrations in eukaryotic cells to affect protein synthesis because they are not actively transported in eukaryotic cells (Epe *et al. FEBS Lett.* 213:443-47, 1987).


[0003] Structure-activity relationships for the tetracycline antibiotics have been determined empirically from 50 years of semi-synthetic modification of the parent structure (Sum *et al. Curr. Pharm. Design* 4:119-32, 1998). Permutations with the upper left-hand portion of the natural product, also known as the hydrophobic domain, have provided new therapeutically active agents, while modifications of the polar hydrophobic domain result in a loss of activity. However, semi-synthesis by its very nature has limited the number of tetracycline analogs that can be prepared and studied.



Tetracycline(3)

[0004] The tetracyclines are composed of four linearly fused six-membered rings with a high density of polar functionality and stereochemical complexity. In 1962, Woodward and co-workers reported the first total synthesis of racemic 6-desmethyl-6-deoxytetracycline (sancycline, 4), the simplest biologically active tetracycline (Conover *et al. J. Am. Chem. Soc.* 84:3222-24, 1962). The synthetic route was a remarkable achievement for the time and proceeded by the stepwise construction of the rings in a linear sequence of 22 steps (overall yield ~0.003%). The first enantioselective synthesis of (-)-tetracycline (3) from the A-ring precursor D-glucosamine (34 steps, 0.002% overall yield) was reported by Tatsuda and co-workers in 2000 (Tatsuta *et al. Chem. Lett.* 646-47, 2000). Other approaches to the synthesis of tetracycline antibiotics, which have also proceeded by the stepwise assembly of the




ABCD ring system beginning with D or CD precursors, include the Shemyakin synthesis of ($\pm$)-12a-deoxy-5a,6-anhydrotetracycline (Gurevich *et al. Tetrahedron Lett.* 8:131, 1967; incorporated herein by reference) and the Muxfeldt synthesis of ($\pm$)-5-oxytetracycline (terramycin, 22 steps, 0.06% yield) (Muxfeldt *et al. J. Am. Chem. Soc.* 101:689, 1979; incorporated herein by reference). Due to the length and poor efficiency of the few existing routes to tetracyclines, which were never designed for synthetic variability, synthesis of tetracycline analogs is still limited.



[0005] There remains a need for a practical and efficient synthetic route to tetracycline analogs, which is amenable to the rapid preparation of specific analogs that can be tested for improved antibacterial and potentially antitumor activity. Such a route would allow the preparation of tetracycline analogs which have not been prepared before.

Summary of the Invention


[0006] The present invention centers around novel synthetic approaches for preparing tetracycline analogs. These synthetic approaches are particularly useful in preparing 6-deoxytetracyclines, which are more stable towards acid and base than 6-hydroxytetracyclines. Doxycycline and minocycline, the two most clinically important tetracyclines, as well as tigecycline, an advanced clinical candidate, are members of the 6-deoxytetracycline class.




The approaches are also useful in preparing 6-hydroxytetracyclines, pentacyclines, hexacyclines, C5-substituted tetracyclines, C5-unsubstituted tetracyclines, tetracyclines with heterocyclic D-rings, and other tetracycline analogs.

[0007] These novel synthetic approaches to tetracycline analogs involve a convergent synthesis of the tetracycline ring system using a highly functionalized chiral enone (**5**) as a key intermediate. The first approach involves the reaction of the enone with an anion formed by the deprotonation of a toluate (**6**) or metallation of a benzylic halide as shown below. The deprotonation of a toluate is particularly useful in preparing 6-deoxytetracyclines with or without a C5-substituent. The metallation (*e.g.*, metal-halogen exchange (*e.g.*, lithium-halogen exchange), metal-metalloid exchange (*e.g.*, lithium-metalloid exchange)) is particularly useful in preparing 6-deoxytetracyclines with or without a C5-substituent as well as pentacyclines. Any organometallic reagent may be used in the cyclization process. Particularly useful reagents may include lithium reagents, Grignard reagents, zero-valent metal reagents,

and ate complexes. In certain embodiments, milder conditions for the cyclization reaction may be preferred.




[0008] The second approach involves reacting the enone (5) in a Diels-Alder-type reaction with a diene (7) or a benzocyclobutenol (8).



In both these approaches, the chiral enone provides the functionalized A and B rings of

the tetracycline core, and the D-ring is derived from the toluate (6), benzylic halide, or benzocyclobutenol (8). In bringing these two portions of the molecule together in a stereoselective manner the C-ring is formed. These approaches not only allow for the stereoselective and efficient synthesis of a wide variety of tetracycline analogs never before prepared, but they also allow for preparation of tetracycline analogs in which the D-ring is replaced with a heterocycle, 5-membered ring, or other ring system. They also allow the preparation of various pentacyclines or higher cyclines containing aromatic and non-aromatic carbocycles and heterocycles.

[0009] Through the oxidation at C6 of 6-deoxytetracycline analogs, 6-oxytetracycline analogs may be prepared as shown in the scheme below:





The 6-deoxytetracycline is transformed into an aromatic naphthol intermediate which undergoes spontaneous autoxidation to form the hydroperoxide. Hydrogenolysis of the hydroperoxide results in the 6-oxytetracycline. This oxidation of 6-deoxytetracycline analogs can be used to prepare tetracyclines in which the D-ring is replaced with a heterocycle, 5-membered ring, or other ring system as well as pentacyclines and other polycyclines containing aromatic and non-aromatic carbocycles and heterocycles.

[0010] The present invention not only provides synthetic methods for preparing these tetracycline analogs but also the intermediates, including chiral enones (5), toluates (6), dienes (7), benzylic halides, and benzocyclobutenol (8), used in these syntheses, and novel derivatives accessed by them.

[0011] Some of the broad classes of compounds available through these new approaches and considered to be a part of the present invention include tetracyclines and various analogs. Important subclasses of tetracyclines include 6-deoxytetracyclines with or without a C5-hydroxyl group, and 6-hydroxytetracyclines with or without a C5-hydroxyl group. Many of the analogs available through these new approaches have never been synthesized before given the limitations of semi-synthetic approaches and

earlier total syntheses. For example, certain substitutions about the D-ring become accessible using the present invention's novel methodologies. In certain classes of compounds of the invention, the D-ring of the tetracyclines analog, which is usually a phenyl ring, is replaced with a heterocyclic moiety, which may be bicyclic or tricyclic. In other classes, the D-ring is replaced with a non-aromatic ring. The size of the D-ring is also not limited to six-membered rings, but instead it may be three-membered, four-membered, five-membered, seven-membered, or larger. In the case of pentacyclines, the five rings may or may not be linear in arrangement. Each of the D- and E-rings may be heterocyclic or carbocyclic, may be aromatic or non-aromatic, and may contain any number of atoms ranging from three to ten atoms. In addition, higher cyclines such as hexacyclines may be prepared. In certain classes, the C-ring may not be fully formed, leading to dicyclines with the A-B fused ring system intact. The compounds of the invention include isomers, stereoisomers, enantiomers, diastereomers, tautomers, protected forms, pro-drugs, salts, and derivatives of any particular compound.





[0012] The present invention also includes intermediates useful in the synthesis of compounds of the present invention. These intermediates include chiral enones, toluates, benzylic halides, and benzocyclobutenol. The intermediates includes various substituted forms, isomers, tautomers, stereoisomers, salts, and derivatives thereof.

[0013] In another aspect, the present invention provides methods of treatment and pharmaceutical composition including the novel compounds of the present invention. The pharmaceutical compositions may also include a pharmaceutically acceptable excipient. The methods and pharmaceutical compositions may be used to treat any infection including cholera, influenza, bronchitis, acne, malaria, urinary tract infections, sexually transmitted diseases including syphilis and gonorrhea, Legionnaires' disease, Lyme disease, Rocky Mountain spotted fever, Q fever, typhus, bubonic plague, gas gangrene, leptospirosis, whooping cough, and anthrax. In certain embodiments, the infections are caused by tetracycline-resistant organisms. In certain instances, the compounds of the invention exhibit anti-neoplastic or anti-proliferative activity, in which case the compounds may be useful in the treatment of diseases such as cancer, autoimmune diseases, inflammatory diseases, and diabetic retinopathy. The methods and compositions may be used to treat disease in humans and other animals including domesticated animals. Any mode of administration including oral and parenteral administration of the pharmaceutical composition may be used.

[0014] Given past work in the synthesis of tetracyclines, the present inventive strategies represent a breakthrough, providing new synthetic routes to tetracyclines and various analogs. The ability to prepare a wide variety of tetracycline analogs and the

use of some of these compounds in the treatment of diseases such as cancer and infectious diseases marks an advance not only in synthetic organic chemistry but also in medicine. The tetracycline class of antibiotics has played a major role in the treatment of infectious diseases in human and veterinary medicine for the past 50 years; however, with the high use of these antibiotics over many years resistance has become a major problem. The present invention fortunately allows for the development of tetracycline analogs with activity against tetracycline-resistant organisms. Therefore, the developments described herein will allow the tetracycline class of antibiotics to remain part of a physician's armamentarium against infection diseases.

Definitions

[0015] Definitions of specific functional groups and chemical terms are described in more detail below. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed., inside cover, and specific functional groups are generally defined as described therein. Additionally, general principles of organic chemistry, as well as specific functional moieties and reactivity, are described in "Organic Chemistry", Thomas Sorrell, University Science Books, Sausalito: 1999, the entire contents of which are incorporated herein by reference.

[0016] Certain compounds of the present invention may exist in particular geometric or stereoisomeric forms. The present invention contemplates all such compounds, including *cis*- and *trans*-isomers, *R*- and *S*-enantiomers, diastereomers, (D)-isomers, (L)-isomers, the racemic mixtures thereof, and other mixtures thereof, as falling within the scope of the invention. Additional asymmetric carbon atoms may be present in a substituent such as an alkyl group. All such isomers, as well as mixtures thereof, are intended to be included in this invention.

[0017] Isomeric mixtures containing any of a variety of isomer ratios may be utilized in accordance with the present invention. For example, where only two isomers are combined, mixtures containing 50:50, 60:40, 70:30, 80:20, 90:10, 95:5, 96:4, 97:3, 98:2, 99:1, or 100:0 isomer ratios are all contemplated by the present invention. Those of ordinary skill in the art will readily appreciate that analogous ratios

are contemplated for more complex isomer mixtures.

[0018] If, for instance, a particular enantiomer of a compound of the present invention is desired, it may be prepared by asymmetric synthesis, or by derivation with a chiral auxiliary, where the resulting diastereomeric mixture is separated and the auxiliary group cleaved to provide the pure desired enantiomers. Alternatively, where the molecule contains a basic functional group, such as amino, or an acidic functional group, such as carboxyl, diastereomeric salts are formed with an appropriate optically-active acid or base, followed by resolution of the diastereomers thus formed by fractional crystallization or chromatographic means well known in the art, and subsequent recovery of the pure enantiomers.

[0019] One of ordinary skill in the art will appreciate that the synthetic methods, as described herein, utilize a variety of protecting groups. By the term “protecting group”, as used herein, it is meant that a particular functional moiety, *e.g.*, O, S, or N, is temporarily blocked so that a reaction can be carried out selectively at another reactive site in a multifunctional compound. In preferred embodiments, a protecting group reacts selectively in good yield to give a protected substrate that is stable to the projected reactions; the protecting group should be selectively removable in good yield by readily available, preferably non-toxic reagents that do not attack the other functional groups; the protecting group forms an easily separable derivative (more preferably without the generation of new stereogenic centers); and the protecting group has a minimum of additional functionality to avoid further sites of reaction. As detailed herein, oxygen, sulfur, nitrogen, and carbon protecting groups may be utilized. Hydroxyl protecting groups include methyl, methoxymethyl (MOM), methylthiomethyl (MTM), *t*-butylthiomethyl, (phenyldimethylsilyl)methoxymethyl (SMOM), benzyloxymethyl (BOM), *p*-methoxybenzyloxymethyl (PMBM), (4-methoxyphenoxy)methyl (*p*-AOM), guaiacolmethyl (GUM), *t*-butoxymethyl, 4-pentenylloxymethyl (POM), siloxymethyl, 2-methoxyethoxymethyl (MEM), 2,2,2-trichloroethoxymethyl, bis(2-chloroethoxy)methyl, 2-(trimethylsilyl)ethoxymethyl (SEMOR), tetrahydropyranyl (THP), 3-bromotetrahydropyranyl, tetrahydrothiopyranyl, 1-methoxycyclohexyl, 4-methoxytetrahydropyranyl (MTHP), 4-methoxytetrahydrothiopyranyl, 4-methoxytetrahydrothiopyranyl S,S-dioxide, 1-[(2-

chloro-4-methyl)phenyl]-4-methoxypiperidin-4-yl (CTMP), 1,4-dioxan-2-yl, tetrahydrofuranyl, tetrahydrothiofuranyl, 2,3,3a,4,5,6,7,7a-octahydro-7,8,8-trimethyl-4,7-methanobenzofuran-2-yl, 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 1-methyl-1-methoxyethyl, 1-methyl-1-benzyloxyethyl, 1-methyl-1-benzyloxy-2-fluoroethyl, 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 2-(phenylselenyl)ethyl, *t*-butyl, allyl, *p*-chlorophenyl, *p*-methoxyphenyl, 2,4-dinitrophenyl, benzyl, *p*-methoxybenzyl, 3,4-dimethoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-halobenzyl, 2,6-dichlorobenzyl, *p*-cyanobenzyl, *p*-phenylbenzyl, 2-picolyl, 4-picolyl, 3-methyl-2-picolyl *N*-oxido, diphenylmethyl, *p,p'*-dinitrobenzhydryl, 5-dibenzosuberyl, triphenylmethyl, α -naphthyl diphenylmethyl, *p*-methoxyphenyl diphenylmethyl, di(*p*-methoxyphenyl)phenylmethyl, tri(*p*-methoxyphenyl)methyl, 4-(4'-bromophenacyloxyphenyl)diphenylmethyl, 4,4',4''-tris(4,5-dichlorophthalimidophenyl)methyl, 4,4',4''-tris(levulinoyloxyphenyl)methyl, 4,4',4''-tris(benzoyloxyphenyl)methyl, 3-(imidazol-1-yl)bis(4',4''-dimethoxyphenyl)methyl, 1,1-bis(4-methoxyphenyl)-1'-pyrenylmethyl, 9-anthryl, 9-(9-phenyl)xantheryl, 9-(9-phenyl-10-oxo)anthryl, 1,3-benzodithiolan-2-yl, benzisothiazolyl S,S-dioxido, trimethylsilyl (TMS), triethylsilyl (TES), triisopropylsilyl (TIPS), dimethylisopropylsilyl (IPDMS), diethylisopropylsilyl (DEIPS), dimethylhexylsilyl, *t*-butyldimethylsilyl (TBDMS), *t*-butyldiphenylsilyl (TBDPS), tribenzylsilyl, tri-*p*-xylylsilyl, triphenylsilyl, diphenylmethylsilyl (DPMS), *t*-butylmethoxyphenylsilyl (TBMPS), formate, benzoylformate, acetate, chloroacetate, dichloroacetate, trichloroacetate, trifluoroacetate, methoxyacetate, triphenylmethoxyacetate, phenoxyacetate, *p*-chlorophenoxyacetate, 3-phenylpropionate, 4-oxopentanoate (levulinate), 4,4-(ethylenedithio)pentanoate (levulinoyldithioacetal), pivaloate, adamantate, crotonate, 4-methoxycrotonate, benzoate, *p*-phenylbenzoate, 2,4,6-trimethylbenzoate (mesitoate), alkyl methyl carbonate, 9-fluorenylmethyl carbonate (Fmoc), alkyl ethyl carbonate, alkyl 2,2,2-trichloroethyl carbonate (Troc), 2-(trimethylsilyl)ethyl carbonate (TMSEC), 2-(phenylsulfonyl) ethyl carbonate (Psec), 2-(triphenylphosphonio) ethyl carbonate (Peoc), alkyl isobutyl carbonate, alkyl vinyl carbonate alkyl allyl carbonate, alkyl *p*-nitrophenyl carbonate, alkyl benzyl carbonate, alkyl *p*-methoxybenzyl carbonate, alkyl 3,4-dimethoxybenzyl carbonate, alkyl *o*-

nitrobenzyl carbonate, alkyl *p*-nitrobenzyl carbonate, alkyl *S*-benzyl thiocarbonate, 4-ethoxy-1-naphthyl carbonate, methyl dithiocarbonate, 2-iodobenzoate, 4-azidobutyrate, 4-nitro-4-methylpentanoate, *o*-(dibromomethyl)benzoate, 2-formylbenzenesulfonate, 2-(methylthiomethoxy)ethyl, 4-(methylthiomethoxy)butyrate, 2-(methylthiomethoxymethyl)benzoate, 2,6-dichloro-4-methylphenoxyacetate, 2,6-dichloro-4-(1,1,3,3-tetramethylbutyl)phenoxyacetate, 2,4-bis(1,1-dimethylpropyl)phenoxyacetate, chlorodiphenylacetate, isobutyrate, monosuccinoate, (*E*)-2-methyl-2-butenate, *o*-(methoxycarbonyl)benzoate, α -naphthoate, nitrate, alkyl *N,N,N',N'*-tetramethylphosphorodiamidate, alkyl *N*-phenylcarbamate, borate, dimethylphosphinothioyl, alkyl 2,4-dinitrophenylsulfenate, sulfate, methanesulfonate (mesylate), benzylation, and tosylate (Ts). For protecting 1,2- or 1,3-diols, the protecting groups include methylene acetal, ethylidene acetal, 1-*t*-butylethylidene ketal, 1-phenylethylidene ketal, (4-methoxyphenyl)ethylidene acetal, 2,2,2-trichloroethylidene acetal, acetonide, cyclopentylidene ketal, cyclohexylidene ketal, cycloheptylidene ketal, benzylidene acetal, *p*-methoxybenzylidene acetal, 2,4-dimethoxybenzylidene ketal, 3,4-dimethoxybenzylidene acetal, 2-nitrobenzylidene acetal, methoxymethylene acetal, ethoxymethylene acetal, dimethoxymethylene ortho ester, 1-methoxyethylidene ortho ester, 1-ethoxyethylidene ortho ester, 1,2-dimethoxyethylidene ortho ester, α -methoxybenzylidene ortho ester, 1-(*N,N*-dimethylamino)ethylidene derivative, α -(*N,N'*-dimethylamino)benzylidene derivative, 2-oxacyclopentylidene ortho ester, di-*t*-butylsilylene group (DTBS), 1,3-(1,1,3,3-tetraisopropylidisiloxanylidene) derivative (TIPDS), tetra-*t*-butoxydisiloxane-1,3-diylidene derivative (TBDS), cyclic carbonates, cyclic boronates, ethyl boronate, and phenyl boronate. Amino-protecting groups include methyl carbamate, ethyl carbamate, 9-fluorenylmethyl carbamate (Fmoc), 9-(2-sulfo)fluorenylmethyl carbamate, 9-(2,7-dibromo)fluorenylmethyl carbamate, 2,7-di-*t*-butyl-[9-(10,10-dioxo-10,10,10,10-tetrahydrothioxanthyl)]methyl carbamate (DBD-Tmoc), 4-methoxyphenacyl carbamate (Phenoc), 2,2,2-trichloroethyl carbamate (Troc), 2-trimethylsilyl ethyl carbamate (Teoc), 2-phenylethyl carbamate (hZ), 1-(1-adamantyl)-1-methylethyl carbamate (Adpoc), 1,1-dimethyl-2-haloethyl carbamate, 1,1-dimethyl-2,2-dibromoethyl carbamate (DB-*t*-BOC), 1,1-dimethyl-2,2,2-trichloroethyl carbamate

(TCBOC), 1-methyl-1-(4-biphenyl)ethyl carbamate (Bpoc), 1-(3,5-di-*t*-butylphenyl)-1-methylethyl carbamate (*t*-Bumeoc), 2-(2'- and 4'-pyridyl)ethyl carbamate (Pyoc), 2-(*N,N*-dicyclohexylcarboxamido)ethyl carbamate, *t*-butyl carbamate (BOC), 1-adamantyl carbamate (Adoc), vinyl carbamate (Voc), allyl carbamate (Alloc), 1-isopropylallyl carbamate (Ipaoc), cinnamyl carbamate (Coc), 4-nitrocinnamyl carbamate (Noc), 8-quinolyl carbamate, *N*-hydroxypiperidinyl carbamate, alkylthio carbamate, benzyl carbamate (Cbz), *p*-methoxybenzyl carbamate (Moz), *p*-nitobenzyl carbamate, *p*-bromobenzyl carbamate, *p*-chlorobenzyl carbamate, 2,4-dichlorobenzyl carbamate, 4-methylsulfinylbenzyl carbamate (MsZ), 9-anthrylmethyl carbamate, diphenylmethyl carbamate, 2-methylthioethyl carbamate, 2-methylsulfonyl ethyl carbamate, 2-(*p*-toluenesulfonyl)ethyl carbamate, [2-(1,3-dithianyl)]methyl carbamate (Dmoc), 4-methylthiophenyl carbamate (Mtpc), 2,4-dimethylthiophenyl carbamate (Bmpc), 2-phosphonioethyl carbamate (Peoc), 2-triphenylphosphonioisopropyl carbamate (Ppoc), 1,1-dimethyl-2-cyanoethyl carbamate, *m*-chloro-*p*-acyloxybenzyl carbamate, *p*-(dihydroxyboryl)benzyl carbamate, 5-benzisoxazolylmethyl carbamate, 2-(trifluoromethyl)-6-chromonylmethyl carbamate (Tcroc), *m*-nitrophenyl carbamate, 3,5-dimethoxybenzyl carbamate, *o*-nitrobenzyl carbamate, 3,4-dimethoxy-6-nitrobenzyl carbamate, phenyl(*o*-nitrophenyl)methyl carbamate, phenothiazinyl-(10)-carbonyl derivative, *N'*-*p*-toluenesulfonylaminocarbonyl derivative, *N'*-phenylaminothiocarbonyl derivative, *t*-amyl carbamate, *S*-benzyl thiocarbamate, *p*-cyanobenzyl carbamate, cyclobutyl carbamate, cyclohexyl carbamate, cyclopentyl carbamate, cyclopropylmethyl carbamate, *p*-decyloxybenzyl carbamate, 2,2-dimethoxycarbonylvinyl carbamate, *o*-(*N,N*-dimethylcarboxamido)benzyl carbamate, 1,1-dimethyl-3-(*N,N*-dimethylcarboxamido)propyl carbamate, 1,1-dimethylpropynyl carbamate, di(2-pyridyl)methyl carbamate, 2-furanylmethyl carbamate, 2-iodoethyl carbamate, isoborynl carbamate, isobutyl carbamate, isonicotinyl carbamate, *p*-(*p'*-methoxyphenylazo)benzyl carbamate, 1-methylcyclobutyl carbamate, 1-methylcyclohexyl carbamate, 1-methyl-1-cyclopropylmethyl carbamate, 1-methyl-1-(3,5-dimethoxyphenyl)ethyl carbamate, 1-methyl-1-(*p*-phenylazophenyl)ethyl carbamate, 1-methyl-1-phenylethyl carbamate, 1-methyl-1-(4-pyridyl)ethyl carbamate, phenyl carbamate, *p*-(phenylazo)benzyl carbamate, 2,4,6-tri-*t*-butylphenyl carbamate,

4-(trimethylammonium)benzyl carbamate, 2,4,6-trimethylbenzyl carbamate, formamide, acetamide, chloroacetamide, trichloroacetamide, trifluoroacetamide, phenylacetamide, 3-phenylpropanamide, picolinamide, 3-pyridylcarboxamide, *N*-benzoylphenylalanyl derivative, benzamide, *p*-phenylbenzamide, *o*-nitrophenylacetamide, *o*-nitrophenoxyacetamide, acetoacetamide, (*N*'-dithiobenzoyloxycarbonylamino)acetamide, 3-(*p*-hydroxyphenyl)propanamide, 3-(*o*-nitrophenyl)propanamide, 2-methyl-2-(*o*-nitrophenoxy)propanamide, 2-methyl-2-(*o*-phenylazophenoxy)propanamide, 4-chlorobutanamide, 3-methyl-3-nitrobutanamide, *o*-nitrocinnamide, *N*-acetylmethionine derivative, *o*-nitrobenzamide, *o*-(benzoyloxymethyl)benzamide, 4,5-diphenyl-3-oxazolin-2-one, *N*-phthalimide, *N*-dithiasuccinimide (Dts), *N*-2,3-diphenylmaleimide, *N*-2,5-dimethylpyrrole, *N*-1,1,4,4-tetramethyldisilylazacyclopentane adduct (STABASE), 5-substituted 1,3-dimethyl-1,3,5-triazacyclohexan-2-one, 5-substituted 1,3-dibenzyl-1,3,5-triazacyclohexan-2-one, 1-substituted 3,5-dinitro-4-pyridone, *N*-methylamine, *N*-allylamine, *N*-[2-(trimethylsilyl)ethoxy]methylamine (SEM), *N*-3-acetoxypyrrolamine, *N*-(1-isopropyl-4-nitro-2-oxo-3-pyrroline-3-yl)amine, quaternary ammonium salts, *N*-benzylamine, *N*-di(4-methoxyphenyl)methylamine, *N*-5-dibenzosuberylamine, *N*-triphenylmethylamine (Tr), *N*-[(4-methoxyphenyl)diphenylmethyl]amine (MMTr), *N*-9-phenylfluorenylamine (PhF), *N*-2,7-dichloro-9-fluorenylmethyleneamine, *N*-ferrocenylmethylamino (Fcm), *N*-2-picolylamino *N*'-oxide, *N*-1,1-dimethylthiomethyleneamine, *N*-benzylideneamine, *N*-*p*-methoxybenzylideneamine, *N*-diphenylmethyleneamine, *N*-[(2-pyridyl)mesityl]methyleneamine, *N*-(*N*',*N*'-dimethylaminomethylene)amine, *N*,*N*'-isopropylidenediamine, *N*-*p*-nitrobenzylideneamine, *N*-salicylideneamine, *N*-5-chlorosalicylideneamine, *N*-(5-chloro-2-hydroxyphenyl)phenylmethyleneamine, *N*-cyclohexylideneamine, *N*-(5,5-dimethyl-3-oxo-1-cyclohexenyl)amine, *N*-borane derivative, *N*-diphenylborinic acid derivative, *N*-[phenyl(pentacarbonylchromium- or tungsten)carbonyl]amine, *N*-copper chelate, *N*-zinc chelate, *N*-nitroamine, *N*-nitrosoamine, amine *N*-oxide, diphenylphosphinamide (Dpp), dimethylthiophosphinamide (Mpt), diphenylthiophosphinamide (Ppt), dialkyl phosphoramidates, dibenzyl phosphoramidate, diphenyl phosphoramidate, benzenesulfenamide, *o*-nitrobenzenesulfenamide (Nps), 2,4-

dinitrobenzenesulfenamide, pentachlorobenzenesulfenamide, 2-nitro-4-methoxybenzenesulfenamide, triphenylmethylsulfenamide, 3-nitropyridinesulfenamide (Npys), *p*-toluenesulfonamide (Ts), benzenesulfonamide, 2,3,6-trimethyl-4-methoxybenzenesulfonamide (Mtr), 2,4,6-trimethoxybenzenesulfonamide (Mtb), 2,6-dimethyl-4-methoxybenzenesulfonamide (Pme), 2,3,5,6-tetramethyl-4-methoxybenzenesulfonamide (Mte), 4-methoxybenzenesulfonamide (Mbs), 2,4,6-trimethylbenzenesulfonamide (Mts), 2,6-dimethoxy-4-methylbenzenesulfonamide (iMds), 2,2,5,7,8-pentamethylchroman-6-sulfonamide (Pmc), methanesulfonamide (Ms), β -trimethylsilylethanesulfonamide (SES), 9-anthracenesulfonamide, 4-(4',8'-dimethoxynaphthylmethyl)benzenesulfonamide (DNMBS), benzylsulfonamide, trifluoromethylsulfonamide, and phenacylsulfonamide.. Exemplary protecting groups are detailed herein, however, it will be appreciated that the present invention is not intended to be limited to these protecting groups; rather, a variety of additional equivalent protecting groups can be readily identified using the above criteria and utilized in the method of the present invention. Additionally, a variety of protecting groups are described in *Protective Groups in Organic Synthesis*, Third Ed. Greene, T.W. and Wuts, P.G., Eds., John Wiley & Sons, New York: 1999, the entire contents of which are hereby incorporated by reference.

[0020] It will be appreciated that the compounds, as described herein, may be substituted with any number of substituents or functional moieties. In general, the term “substituted” whether preceded by the term “optionally” or not, and substituents contained in formulas of this invention, refer to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. When more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. As used herein, the term “substituted” is contemplated to include all permissible substituents of organic compounds. In a broad aspect, the permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and nonaromatic substituents of organic compounds. For purposes of this invention, heteroatoms such as nitrogen may have hydrogen substituents and/or any permissible substituents of organic compounds described herein

which satisfy the valencies of the heteroatoms. Furthermore, this invention is not intended to be limited in any manner by the permissible substituents of organic compounds. Combinations of substituents and variables envisioned by this invention are preferably those that result in the formation of stable compounds useful in the treatment, for example, of infectious diseases or proliferative disorders. The term "stable", as used herein, preferably refers to compounds which possess stability sufficient to allow manufacture and which maintain the integrity of the compound for a sufficient period of time to be detected and preferably for a sufficient period of time to be useful for the purposes detailed herein.

[0021] The term "aliphatic", as used herein, includes both saturated and unsaturated, straight chain (*i.e.*, unbranched), branched, acyclic, cyclic, or polycyclic aliphatic hydrocarbons, which are optionally substituted with one or more functional groups. As will be appreciated by one of ordinary skill in the art, "aliphatic" is intended herein to include, but is not limited to, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, and cycloalkynyl moieties. Thus, as used herein, the term "alkyl" includes straight, branched and cyclic alkyl groups. An analogous convention applies to other generic terms such as "alkenyl", "alkynyl", and the like. Furthermore, as used herein, the terms "alkyl", "alkenyl", "alkynyl", and the like encompass both substituted and unsubstituted groups. In certain embodiments, as used herein, "lower alkyl" is used to indicate those alkyl groups (cyclic, acyclic, substituted, unsubstituted, branched or unbranched) having 1-6 carbon atoms.

[0022] In certain embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-20 aliphatic carbon atoms. In certain other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-10 aliphatic carbon atoms. In yet other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-8 aliphatic carbon atoms. In still other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-6 aliphatic carbon atoms. In yet other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-4 carbon atoms. Illustrative aliphatic groups thus include, but are not limited to, for example, methyl, ethyl, n-propyl, isopropyl, cyclopropyl, -CH₂-cyclopropyl, vinyl, allyl, n-butyl, sec-butyl, isobutyl, tert-

butyl, cyclobutyl, -CH₂-cyclobutyl, n-pentyl, sec-pentyl, isopentyl, tert-pentyl, cyclopentyl, -CH₂-cyclopentyl, n-hexyl, sec-hexyl, cyclohexyl, -CH₂-cyclohexyl moieties and the like, which again, may bear one or more substituents. Alkenyl groups include, but are not limited to, for example, ethenyl, propenyl, butenyl, 1-methyl-2-buten-1-yl, and the like. Representative alkynyl groups include, but are not limited to, ethynyl, 2-propynyl (propargyl), 1-propynyl, and the like.

[0023] The term “alkoxy”, or “thioalkyl” as used herein refers to an alkyl group, as previously defined, attached to the parent molecule through an oxygen atom or through a sulfur atom. In certain embodiments, the alkyl, alkenyl, and alkynyl groups contain 1-20 aliphatic carbon atoms. In certain other embodiments, the alkyl, alkenyl, and alkynyl groups contain 1-10 aliphatic carbon atoms. In yet other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-8 aliphatic carbon atoms. In still other embodiments, the alkyl, alkenyl, and alkynyl groups contain 1-6 aliphatic carbon atoms. In yet other embodiments, the alkyl, alkenyl, and alkynyl groups contain 1-4 aliphatic carbon atoms. Examples of alkoxy, include but are not limited to, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, tert-butoxy, neopentoxy, and n-hexoxy. Examples of thioalkyl include, but are not limited to, methylthio, ethylthio, propylthio, isopropylthio, n-butylthio, and the like.

[0024] The term “alkylamino” refers to a group having the structure -NHR', wherein R' is aliphatic, as defined herein. In certain embodiments, the aliphatic group contains 1-20 aliphatic carbon atoms. In certain other embodiments, the aliphatic group contains 1-10 aliphatic carbon atoms. In yet other embodiments, the aliphatic group employed in the invention contain 1-8 aliphatic carbon atoms. In still other embodiments, the aliphatic group contains 1-6 aliphatic carbon atoms. In yet other embodiments, the aliphatic group contains 1-4 aliphatic carbon atoms. Examples of alkylamino groups include, but are not limited to, methylamino, ethylamino, n-propylamino, iso-propylamino, cyclopropylamino, n-butylamino, tert-butylamino, neopentylamino, n-pentylamino, hexylamino, cyclohexylamino, and the like.

[0025] The term “dialkylamino” refers to a group having the structure -NRR', wherein R and R' are each an aliphatic group, as defined herein. R and R' may be the same or different in an dialkylamino moiety. In certain embodiments, the aliphatic

groups contains 1-20 aliphatic carbon atoms. In certain other embodiments, the aliphatic groups contains 1-10 aliphatic carbon atoms. In yet other embodiments, the aliphatic groups employed in the invention contain 1-8 aliphatic carbon atoms. In still other embodiments, the aliphatic groups contains 1-6 aliphatic carbon atoms. In yet other embodiments, the aliphatic groups contains 1-4 aliphatic carbon atoms. Examples of dialkylamino groups include, but are not limited to, dimethylamino, methyl ethylamino, diethylamino, methylpropylamino, di(n-propyl)amino, di(iso-propyl)amino, di(cyclopropyl)amino, di(n-butyl)amino, di(tert-butyl)amino, di(neopentyl)amino, di(n-pentyl)amino, di(hexyl)amino, di(cyclohexyl)amino, and the like. In certain embodiments, R and R' are linked to form a cyclic structure. The resulting cyclic structure may be aromatic or non-aromatic. Examples of cyclic diaminoalkyl groups include, but are not limited to, aziridinyl, pyrrolidinyl, piperidinyl, morpholinyl, pyrrolyl, imidazolyl, 1,3,4-trianolyl, and tetrazolyl.

[0026] Some examples of substituents of the above-described aliphatic (and other) moieties of compounds of the invention include, but are not limited to aliphatic; heteroaliphatic; aryl; heteroaryl; arylalkyl; heteroarylalkyl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; F; Cl; Br; I; -OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)₂R_x; -NR_x(CO)R_x wherein each occurrence of R_x independently includes, but is not limited to, aliphatic, heteroaliphatic, aryl, heteroaryl, arylalkyl, or heteroarylalkyl, wherein any of the aliphatic, heteroaliphatic, arylalkyl, or heteroarylalkyl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic, and wherein any of the aryl or heteroaryl substituents described above and herein may be substituted or unsubstituted. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples that are described herein.

[0027] In general, the terms “aryl” and “heteroaryl”, as used herein, refer to stable mono- or polycyclic, heterocyclic, polycyclic, and polyheterocyclic unsaturated moieties having preferably 3-14 carbon atoms, each of which may be substituted or unsubstituted. Substituents include, but are not limited to, any of the previously

mentioned substituents, *i.e.*, the substituents recited for aliphatic moieties, or for other moieties as disclosed herein, resulting in the formation of a stable compound. In certain embodiments of the present invention, “aryl” refers to a mono- or bicyclic carbocyclic ring system having one or two aromatic rings including, but not limited to, phenyl, naphthyl, tetrahydronaphthyl, indanyl, indenyl, and the like. In certain embodiments of the present invention, the term “heteroaryl”, as used herein, refers to a cyclic aromatic radical having from five to ten ring atoms of which one ring atom is selected from S, O, and N; zero, one, or two ring atoms are additional heteroatoms independently selected from S, O, and N; and the remaining ring atoms are carbon, the radical being joined to the rest of the molecule via any of the ring atoms, such as, for example, pyridyl, pyrazinyl, pyrimidinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, quinolinyl, isoquinolinyl, and the like.

[0028] It will be appreciated that aryl and heteroaryl groups can be unsubstituted or substituted, wherein substitution includes replacement of one, two, three, or more of the hydrogen atoms thereon independently with any one or more of the following moieties including, but not limited to: aliphatic; heteroaliphatic; aryl; heteroaryl; arylalkyl; heteroarylalkyl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; -F; -Cl; -Br; -I; -OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)₂R_x; -NR_x(CO)R_x, wherein each occurrence of R_x independently includes, but is not limited to, aliphatic, heteroaliphatic, aryl, heteroaryl, arylalkyl, or heteroarylalkyl, wherein any of the aliphatic, heteroaliphatic, arylalkyl, or heteroarylalkyl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic, and wherein any of the aryl or heteroaryl substituents described above and herein may be substituted or unsubstituted. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples that are described herein.

[0029] The term “cycloalkyl”, as used herein, refers specifically to groups having three to seven, preferably three to ten carbon atoms. Suitable cycloalkyls

include, but are not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like, which, as in the case of other aliphatic, heteroaliphatic, or heterocyclic moieties, may optionally be substituted with substituents including, but not limited to aliphatic; heteroaliphatic; aryl; heteroaryl; arylalkyl; heteroarylalkyl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; -F; -Cl; -Br; -I; -OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)₂R_x; -NR_x(CO)R_x, wherein each occurrence of R_x independently includes, but is not limited to, aliphatic, heteroaliphatic, aryl, heteroaryl, arylalkyl, or heteroarylalkyl, wherein any of the aliphatic, heteroaliphatic, arylalkyl, or heteroarylalkyl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic, and wherein any of the aryl or heteroaryl substituents described above and herein may be substituted or unsubstituted. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples that are described herein.

[0030] The term “heteroaliphatic”, as used herein, refers to aliphatic moieties that contain one or more oxygen, sulfur, nitrogen, phosphorus, or silicon atoms, *e.g.*, in place of carbon atoms. Heteroaliphatic moieties may be branched, unbranched, cyclic or acyclic and include saturated and unsaturated heterocycles such as morpholino, pyrrolidinyl, *etc.* In certain embodiments, heteroaliphatic moieties are substituted by independent replacement of one or more of the hydrogen atoms thereon with one or more moieties including, but not limited to aliphatic; heteroaliphatic; aryl; heteroaryl; arylalkyl; heteroarylalkyl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; -F; -Cl; -Br; -I; -OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)₂R_x; -NR_x(CO)R_x, wherein each occurrence of R_x independently includes, but is not limited to, aliphatic, heteroaliphatic, aryl, heteroaryl, arylalkyl, or heteroarylalkyl, wherein any of the aliphatic, heteroaliphatic, arylalkyl, or heteroarylalkyl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic, and wherein any of the aryl or heteroaryl substituents described above and

herein may be substituted or unsubstituted. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples that are described herein.

[0031] The terms “halo” and “halogen” as used herein refer to an atom selected from fluorine, chlorine, bromine, and iodine.

[0032] The term “haloalkyl” denotes an alkyl group, as defined above, having one, two, or three halogen atoms attached thereto and is exemplified by such groups as chloromethyl, bromoethyl, trifluoromethyl, and the like.

[0033] The term “heterocycloalkyl” or “heterocycle”, as used herein, refers to a non-aromatic 5-, 6-, or 7- membered ring or a polycyclic group, including, but not limited to a bi- or tri-cyclic group comprising fused six-membered rings having between one and three heteroatoms independently selected from oxygen, sulfur and nitrogen, wherein (i) each 5-membered ring has 0 to 1 double bonds and each 6-membered ring has 0 to 2 double bonds, (ii) the nitrogen and sulfur heteroatoms may be optionally be oxidized, (iii) the nitrogen heteroatom may optionally be quaternized, and (iv) any of the above heterocyclic rings may be fused to a benzene ring. Representative heterocycles include, but are not limited to, pyrrolidinyl, pyrazolinyl, pyrazolidinyl, imidazolinyl, imidazolidinyl, piperidinyl, piperazinyl, oxazolidinyl, isoxazolidinyl, morpholinyl, thiazolidinyl, isothiazolidinyl, and tetrahydrofuryl. In certain embodiments, a “substituted heterocycloalkyl or heterocycle” group is utilized and as used herein, refers to a heterocycloalkyl or heterocycle group, as defined above, substituted by the independent replacement of one, two or three of the hydrogen atoms thereon with but are not limited to aliphatic; heteroaliphatic; aryl; heteroaryl; arylalkyl; heteroarylalkyl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; -F; -Cl; -Br; -I; -OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)₂R_x; -NR_x(CO)R_x, wherein each occurrence of R_x independently includes, but is not limited to, aliphatic, heteroaliphatic, aryl, heteroaryl, arylalkyl, or heteroarylalkyl, wherein any of the aliphatic, heteroaliphatic, arylalkyl, or heteroarylalkyl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic,

and wherein any of the aryl or heteroaryl substituents described above and herein may be substituted or unsubstituted. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples which are described herein.

[0034] “Carbocycle”: The term "carbocycle", as used herein, refers to an aromatic or non-aromatic ring in which each atom of the ring is a carbon atom.

[0035] “Independently selected”: The term “independently selected” is used herein to indicate that the R groups can be identical or different.

[0036] “Labeled”: As used herein, the term “labeled” is intended to mean that a compound has at least one element, isotope, or chemical compound attached to enable the detection of the compound. In general, labels typically fall into three classes: a) isotopic labels, which may be radioactive or heavy isotopes, including, but not limited to, ^2H , ^3H , ^{32}P , ^{35}S , ^{67}Ga , $^{99\text{m}}\text{Tc}$ (Tc-99m), ^{111}In , ^{123}I , ^{125}I , ^{169}Yb and ^{186}Re ; b) immune labels, which may be antibodies or antigens, which may be bound to enzymes (such as horseradish peroxidase) that produce detectable agents; and c) colored, luminescent, phosphorescent, or fluorescent dyes. It will be appreciated that the labels may be incorporated into the compound at any position that does not interfere with the biological activity or characteristic of the compound that is being detected. In certain embodiments, hydrogen atoms in the compound are replaced with deuterium atoms (^2H) to slow the degradation of compound *in vivo*. Due to isotope effects, enzymatic degradation of the deuterated tetracyclines may be slowed thereby increasing the half-life of the compound *in vivo*. In certain embodiments of the invention, photoaffinity labeling is utilized for the direct elucidation of intermolecular interactions in biological systems. A variety of known photophores can be employed, most relying on photoconversion of diazo compounds, azides, or diazirines to nitrenes or carbenes (See, Bayley, H., Photogenerated Reagents in Biochemistry and Molecular Biology (1983), Elsevier, Amsterdam.), the entire contents of which are hereby incorporated by reference. In certain embodiments of the invention, the photoaffinity labels employed are o-, m- and p-azidobenzoyls, substituted with one or more halogen moieties, including, but not limited to 4-azido-2,3,5,6-tetrafluorobenzoic acid.

[0037] “Tautomers”: As used herein, the term “tautomers” are particular

isomers of a compound in which a hydrogen and double bond have changed position with respect to the other atoms of the molecule. For a pair of tautomers to exist there must be a mechanism for interconversion. Examples of tautomers include keto-enol forms, imine-enamine forms, amide-imino alcohol forms, amidine-aminidine forms, nitroso-oxime forms, thio ketone-enethiol forms, *N*-nitroso-hydroxyazo forms, nitro-*aci*-nitro forms, and pyridione-hydroxypyridine forms.

[0038] Definitions of non-chemical terms used throughout the specification include:

[0039] “Animal”: The term animal, as used herein, refers to humans as well as non-human animals, including, for example, mammals, birds, reptiles, amphibians, and fish. Preferably, the non-human animal is a mammal (*e.g.*, a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, a primate, or a pig). A non-human animal may be a transgenic animal.

[0040] “Associated with”: When two entities are “associated with” one another as described herein, they are linked by a direct or indirect covalent or non-covalent interaction. Preferably, the association is covalent. Desirable non-covalent interactions include hydrogen bonding, van der Waals interactions, hydrophobic interactions, magnetic interactions, electrostatic interactions, *etc.*

[0041] “Effective amount”: In general, the “effective amount” of an active agent or the microparticles refers to an amount sufficient to elicit the desired biological response. As will be appreciated by those of ordinary skill in this art, the effective amount of a compound of the invention may vary depending on such factors as the desired biological endpoint, the pharmacokinetics of the compound, the disease being treated, the mode of administration, and the patient. For example, the effective amount of a tetracycline analog antibiotic is the amount that results in a sufficient concentration at the site of the infection to kill the microorganism causing the infection (bacteriocidal) or to inhibit the reproduction of such microorganisms (bacteriostatic). In another example, the effective amount of tetracycline analog antibiotic is the amount sufficient to reverse clinical signs and symptoms of the infection, including fever, redness, warmth, pain, chills, cultures, and pus production.

Brief Description of the Drawing

- [0042] *Figure 1* shows the modular synthesis of tetracycline and tetracycline analogs starting from benzoic acid.
- [0043] *Figure 2* depicts the total synthesis of (-)-tetracycline starting from benzoic acid and involving an *o*-quinone dimethide Diels-Alder reaction between the chiral enone **10** and the benzocyclobutenol **11**. The overall yield for the 17 step synthesis was 1.1%.
- [0044] *Figure 3* is the total synthesis of (-)-doxycycline in 18 steps (overall yield 8.2%). The synthesis includes the reaction of the chiral enone **23** with the anion **24** to yield the tetracycline core. The first seven steps are identical to the first seven steps in the synthesis of (-)-tetracycline shown in *Figure 2*.
- [0045] *Figure 4* shows a first and second generation synthesis of isoxazole **4** used in the synthesis of (-)-tetracycline and (-)-doxycycline as shown in *Figure 2*.
- [0046] *Figure 5* shows the synthesis of benzocyclobutenol **11** used in the synthesis of (-)-tetracycline as shown in *Figure 2*.
- [0047] *Figure 6* shows the synthesis of dicyclines. Dicyclines preserve the hydrophilic region thought to be important for the antimicrobial activity of tetracyclines.
- [0048] *Figure 7* depicts the synthesis of tricyclines via a Diels-Alder reaction with the chiral enone **10** and a diene (**41**). Tricyclines preserve the hydrophobic region thought to be important for antimicrobial activity.
- [0049] *Figure 8* shows the synthesis of pentacyclines.
- [0050] *Figure 9* shows the synthesis of bridge pentacyclines by reacting anion **47** with a chiral enone.
- [0051] *Figure 10* shows five compounds that may be used as analog platforms for the synthesis of tetracycline analogs.
- [0052] *Figure 11* is a scheme showing the synthesis of a pyridone/hydroxypyridine analog of sancycline.

[0053] *Figure 12* shows the total synthesis of 6-deoxytetracycline from benzoic acid in 14 steps (overall yield 8%). The first ten steps are identical to the first 10 steps in the synthesis of (-)-tetracycline shown in *Figure 2*.

[0054] *Figure 13A* shows the synthesis of a pyridine analog of sancycline, 7-aza-10-deoxysancycline. *Figure 13B* shows the synthesis of 10-deoxysancycline.

[0055] *Figure 14A* and *14B* show a number of examples of heterocyclines, tetracycline analogs, pentacyclines, and polycyclines potentially accessible via the inventive method.

[0056] *Figure 15* shows the chemical structures of various tetracycline antibiotics. (-)-Tetracycline (**1**) was first produced semi-synthetically, by hydrogenolysis of the fermentation product aureomycin (7-chlorotetracycline), but later was discovered to be a natural product and is now produced by fermentation (M. Nelson, W. Hillen, R. A. Greenwald, Eds., *Tetracyclines in Biology, Chemistry and Medicine* (Birkhauser Verlag, Boston, 2001); incorporated herein by reference). (-)-Doxycycline (**2**) and minocycline (**3**) are clinically important non-natural antibiotics and are both manufactured by multi-step chemical transformations of fermentation products (semi-synthesis) (M. Nelson, W. Hillen, R. A. Greenwald, Eds., *Tetracyclines in Biology, Chemistry and Medicine* (Birkhauser Verlag, Boston, 2001); incorporated herein by reference). Structures **4-6** are representative of tetracycline-like molecules that cannot be prepared by any known semi-synthetic pathway, but which are now accessible by the convergent assembly depicted in *Figure 15B*. *Figure 15B* depicts a generalized Michael-Dieckmann reaction sequence that forms the C-ring of tetracyclines from the coupling of structurally varied carbanionic D-ring precursors with either of the AB precursors **7** or **8**.

[0057] *Figure 16* shows the transformation of benzoic acid in 7 steps to the key bicyclic intermediate **14**. This product is then used to prepare the AB precursor enone **7** by the 4-step sequence shown, or to enone **8**, AB precursor to 6-deoxy-5-hydroxytetracycline derivatives, by the 8-step sequence shown.

[0058] *Figure 17* shows the synthesis of the clinically important antibiotic (-)-doxycycline (**2**) by the convergent coupling of the *o*-toluate anion derived from **18** and the AB precursor enone **8**.

[0059] *Figure 18* shows the synthesis of structurally diverse 6-deoxytetracyclines by coupling of structurally diverse D-ring precursors and AB precursors **7** or **8**. The number of steps and overall yields from benzoic acid are shown in parentheses below each structure synthesized. MIC values ($\mu\text{g/mL}$) are also shown for whole-cell antibacterial testing of each analog against 5 Gram-positive and 5-Gram-negative microorganisms. Corresponding MICs for tetracycline (**1**), a testing control, appear at bottom.

[0060] *Figure 19* shows a crystalline Michael adduct as the product of a lithium anion and a chiral enone.

[0061] *Figure 20* shows the synthesis of a pentacycline via a Michael-Dieckman reaction sequence.


[0062] *Figure 21* shows the synthesis of various novel tetracycline analogs and their corresponding D-ring precursor. These compounds represent significant gaps in the tetracycline fields, likely missing from the literature for lack of a viable synthesis.

[0063] *Figure 22* shows alternative sequences to AB enone precursors from 1*S*,2*R*-*cis*-dihydroxybenzoic acid.

[0064] *Figure 23* shows novel routes to AB precursors. These routes do not involve the microbial dihydroxylation of benzoic acid.

Detailed Description of Certain Preferred Embodiments of the Invention

[0065] The present invention provides a strategy for the synthesis of tetracycline analogs via a convergent synthesis using as an intermediate, the highly functionalized chiral enone **9** as shown below:



wherein R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or

unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_C$; $=O$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_D$; $=O$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_5 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_E$; $-CN$; $-SCN$; $-SR_E$; or $-N(R_E)_2$; wherein each occurrence of R_E is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_6 is selected from the group consisting of hydrogen, halogen, substituted or unsubstituted aliphatic, substituted or unsubstituted heteroaliphatic, substituted or unsubstituted alkoxy, $-OH$, $-CN$, $-SCN$, $-SH$, alkylthio, arylthio, $-NO_2$, amino, alkyl amino, and dialkyl amino groups;


P is independently selected from the group consisting of hydrogen or a protecting group. The chiral enone 9 can be reacted with anions of phthalides, anions of toluates, benzocyclobutenole, or dienes to yield tetracycline analogs including heterocyclic tetracyclines, dicyclines, tricyclines, pentacyclines, heterocyclic pentacyclines, polycyclines, and heterocyclic polycyclines. These new compounds are tested for anti-microbial activity against microbes including traditionally tetracycline-sensitive organisms as well as organisms known to be tetracycline-resistant. Compounds found to be bacteriocidal or bacteriostatic are used in formulating pharmaceutical for the treatment of infections in human and veterinary medicine. The compounds are also tested for anti-proliferative activity. Such compounds are useful in the treatment of antiproliferative diseases including cancer, anti-inflammatory diseases, autoimmune diseases, benign neoplasms, and diabetic retinopathy. The inventive approach to the synthesis of tetracycline analogs allows for the efficient synthesis of many compounds never before prepared or available using earlier routes and semi-synthetic techniques.

Compounds

[0066] Compounds of the present invention include tetracycline analogs, heterocyclic tetracycline analogs, dicyclines, tricyclines, pentacyclines, heterocyclic pentacyclines, bridged pentacyclines, heterocyclic polycyclines, bridged polycyclines, and other polycyclines. Particularly useful compounds of the present invention include those with biological activity. In certain embodiments, the compounds of the invention exhibit antimicrobial activity. For example, the compound may have a mean inhibitory concentration, with respect to a particular bacteria, of less than 50 $\mu\text{g/mL}$, preferably less than 25 $\mu\text{g/mL}$, more preferably less than 5 $\mu\text{g/mL}$, and most preferably less than 1 $\mu\text{g/mL}$. For example, infection caused by the following organisms may be treated with antimicrobial compounds of the invention: Gram-positives—*Staphylococcus aureus*, *Streptococcus* Group A, *Streptococcus viridans*, *Streptococcus pneumoniae*; Gram-negatives—*Neisseria meningitidis*, *Neisseria gonorrhoeae*, *Haemophilus influenzae*, *Escherichia coli*, *Bacteroides fragilis*, other *Bacteroides*; and Others—

Mycoplasma pneumoniae, *Treponema pallidum*, *Rickettsia*, and *Chlamydia*. In other embodiments, the compounds of the invention exhibit antiproliferative activity.

[0067] In certain embodiments, the tetracycline analogs of the present invention are represented by the formula:



The D-ring of 10 may include one, two, or three double bonds. In certain embodiments, the D-ring is aromatic. In other embodiments, the D-ring includes only one double bond, and in yet other embodiments, the D-ring includes two double bonds which may or may not be in conjugation. The D-ring may be substituted with various groups R_7 , R_6 , and R_8 as defined below.

In 10, R_1 can be hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_A$; $=O$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. In certain embodiments, R_1 is hydrogen. In other embodiments, R_1 is lower alkyl, alkenyl, or alkynyl. In yet other embodiments, R_1 is methyl, ethyl, n-propyl, cyclopropyl, or isopropyl. In still other embodiments R_1 is methyl.

R_2 may be hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted

or unsubstituted, branched or unbranched heteroaryl; $-OR_B$; $=O$; $-C(=O)R_B$; $-CO_2R_B$; $-CN$; $-SCN$; $-SR_B$; $-SOR_B$; $-SO_2R_B$; $-NO_2$; $-N(R_B)_2$; $-NHC(O)R_B$; or $-C(R_B)_3$; wherein each occurrence of R_B is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. In certain embodiments, R_2 is hydrogen. In other embodiments, R_2 is hydroxyl or a protected hydroxyl group. In certain embodiments, R_2 is alkoxy. In yet other embodiments, R_2 is a lower alkyl, alkenyl, or alkynyl group. In certain embodiments, R_1 is methyl, and R_2 is hydroxyl. In other embodiments, R_1 is methyl, and R_2 is hydrogen. In certain embodiments, R_1 and R_2 are taken together to form a carbocyclic or heterocyclic ring system spiro-linked to **10**.

R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_C$; $=O$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. In certain embodiments, R_3 is hydrogen. In other embodiments, R_3 is a hydroxyl group or a protected hydroxyl group. In yet other embodiments, R_3 is alkoxy. In still further embodiments, R_3 is lower alkyl, alkenyl, or alkynyl.

R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_D$; $=O$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently a hydrogen, a protecting group, an aliphatic

moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. In certain embodiments, R_4 is hydrogen. In other embodiments, R_4 is a hydroxyl group or a protected hydroxyl group. In yet other embodiments, R_4 is alkoxy. In still further embodiments, R_4 is lower alkyl, alkenyl, or alkynyl. In certain embodiments, both R_3 and R_4 are hydrogen. In other embodiments, R_3 and R_4 are taken together to form a carbocyclic or heterocyclic ring system spiro-linked to the B-ring of **10**.

R_5 may be hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_E$; $-CN$; $-SCN$; $-SR_E$; or $-N(R_E)_2$; wherein each occurrence of R_E is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. In certain embodiments, R_5 is amino, alkylamino, or dialkylamino; preferably dimethylamino, diethylamino, methyl(ethyl)amino, dipropylamino, methyl(propyl)amino, or ethyl(propyl)amino. In other embodiments, R_5 is hydroxyl, protected hydroxyl, or alkoxy. In yet other embodiments, R_5 is sulphydryl, protected sulphydryl, or alkylthioxy.

R_7 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_G$; $=O$; $-C(=O)R_G$; $-CO_2R_G$; $-CN$; $-SCN$; $-SR_G$; $-SOR_G$; $-SO_2R_G$; $-NO_2$; $-N(R_G)_2$; $-NHC(O)R_G$; or $-C(R_G)_3$; wherein each occurrence of R_G is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. In certain embodiments, R_7 is hydroxyl, protected hydroxyl,


alkoxy, lower alkyl, lower alkenyl, lower alkynyl, or halogen.

R_6 and R_8 are absent if the dashed line between the carbon atoms which R_6 and R_8 are attached to represents a bond, or are each selected independently from the group consisting of hydrogen, halogen, substituted or unsubstituted aliphatic, substituted or unsubstituted heteroaliphatic, substituted or unsubstituted alkoxy, -OH, -CN, -SCN, -SH, alkylthio, -NO₂, amino, alkyl amino, and dialkyl amino groups. In certain embodiments, R_6 and R_8 are absent. In other embodiments, R_6 or R_8 is absent.

[0068] The variable n is an integer in the range of 0 to 8, inclusive. As will be appreciated by one of skill in the art, when the D-ring is aromatic n is an integer between 0 and 4, preferably between 1 and 3, more preferable between 1 and 2. In certain embodiments, when n is 2, the substituents R_7 are in the *ortho* configuration. In other embodiments, when n is 2, the substituents R_7 are in the *para* configuration. And in yet other embodiments, when n is 2, the substituents R_7 are in the *meta* configuration.



[0069] A dashed line in formula 10 may represent a bond or the absence of a bond.

[0070] As will be appreciated by one of skill in this art, compounds of formula 10 include derivatives, labeled forms, salts, pro-drugs, isomers, and tautomers thereof. Derivatives include protected forms. Salts include any pharmaceutically acceptable salts including HCl, HBr, HI, acetate, and fatty acid (*e.g.*, lactate, citrate, myristoleate, oleate, valerate) salts. In certain embodiments, the inventive compound exists in zwitterionic form at neutral pH with the R_5 being a protonated amino group and the C-3 hydroxyl group deprotonated as shown in formula 10a.




Isomers include geometric isomers, diastereomers, and enantiomers. Tautomers include both keto and enol forms of carbonyl moieties as well as various tautomeric forms of substituted and unsubstituted heterocycles. For example, the B-ring as shown


in formula **10** includes an enol moiety as drawn, but the enol may exist as the keto form in certain compounds as shown below in formula **10b** and **10c**:



Other tautomeric forms will be appreciated by one of skill in the art and will depend on the substitution pattern of the core ring structure. The formulae drawn are only given as examples and do not in any way represent the full range of tautomers that may exist for a particular compound.


[0071] Various subclasses of compounds of the formula **10** which include a substituted or unsubstituted aromatic D-ring are shown below. These subclasses include unsubstituted, monosubstituted, disubstituted, and trisubstituted D-ring.












wherein the definitions of R_1 , R_2 , R_3 , R_4 , and R_5 are as described above, and R_7 is halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_G$; $=O$; $-C(=O)R_G$; $-CO_2R_G$; $-CN$; $-SCN$; $-SR_G$; $-SOR_G$; $-SO_2R_G$; $-NO_2$; $-N(R_G)_2$; $-NHC(O)R_G$; or $-C(R_G)_3$; wherein each occurrence of R_G is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. In certain embodiments, R_7 is hydroxyl, protected hydroxyl, alkoxy, lower alkyl, lower alkenyl, lower alkynyl, or halogen. In other embodiments, R_7 is cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; or cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic. In yet other embodiments, R_7 is amino, alkylamino, or dialkylamino. In other embodiments, R_7 is substituted or unsubstituted cyclic, heterocyclic, aryl, or heteroaryl. In certain embodiments, R_7 is branched or unbranched acyl.


[0072] Various subclasses of compounds of the formula **10** which include a hydroxyl group at C10 are shown:









wherein the definitions of R_1 , R_2 , R_3 , R_4 , R_5 , R_E , and R_7 are as described above. In certain embodiments, the compounds are 6-deoxytetracyclines as shown in the




formulae below:





wherein R_2 is hydrogen, and the definitions of R_1 , R_3 , R_4 , R_5 , R_E , and R_7 are as described above.

[0073] In another aspect of the invention, the carbocyclic D-ring of tetracycline is replaced with a heterocyclic or carbocyclic moiety as shown in formula (11):



The definitions of R_1 , R_2 , R_3 , R_4 , and R_5 are as described above for formula 10. The D-

ring represented by  can be a substituted or unsubstituted aryl, heteroaryl, carbocyclic, or heterocyclic moiety, in which each occurrence of X is selected from the group consisting of -O-, -S-, -NR₇-, -C(R₇)₂-; n is an integer in the range of 1 to 5, inclusive; and the bonds between adjacent X moieties are either single or double bonds.


In certain embodiments,  is a polycyclic ring system such as a bicyclic or

tricyclic moiety. In other embodiments,  is a monocyclic moiety. In yet other


embodiments,  is a substituted or unsubstituted heterocyclic moiety. In certain

embodiments,  is not a substituted or unsubstituted phenyl ring. In other


embodiments,  is a pyridinyl moiety as shown:



In another embodiment,  is selected from the group consisting of




In yet another embodiment, $(X)_n$ is a five-membered heterocyclic ring selected from the group consisting of:



Various tetracyclines (heterocycles) of the invention are also shown in *Figure 14*.




[0074] Other compounds of the invention include pentacyclines of the formula:



wherein R_1 , R_2 , R_3 , R_4 , R_5 , and $(X)_n$ are as defined above. In certain embodiments, the rings of the compound are linear. In other embodiments, the ring system is not

linear. Each occurrence of the ring $(X)_n$, in certain embodiments, is a monocyclic

ring system. Each occurrence of $(X)_n$ is heterocyclic or carbocyclic. $(X)_n$ is three-membered, four-membered, five-membered, six-membered, or seven-membered; preferably, five-membered or six-membered. Other classes of pentacyclines include compounds of the formulae (12), (13), and (14):



wherein R_1 , R_2 , R_3 , R_4 , R_5 , and R_7 are as defined above. In formulae **12**, **13**, and **14**,

 represents a substituted or unsubstituted aryl, heteroaryl, carbocyclic, or heterocyclic moiety, in which each occurrence of X is selected from the group consisting of -O-, -S-, -NR₈-, -C(R₈)₂-; n is an integer in the range of 1 to 5, inclusive; and the bonds between adjacent X moieties are either single or double bonds. In certain


embodiments,  is a polycyclic ring system such as a bicyclic or tricyclic moiety.

In other embodiments,  is a monocyclic moiety. In other embodiments,  is a substituted or unsubstituted, aromatic or nonaromatic carbocyclic moiety, for


example a phenyl ring. In yet other embodiments,  is a substituted or

unsubstituted heterocyclic moiety. In certain embodiments,  is not a substituted


or unsubstituted phenyl ring. In other embodiments,  is a pyridinyl moiety as shown:




In another embodiment,  is selected from the group consisting of








In yet another embodiment,  is a five-membered heterocyclic ring selected from the group consisting of:




[0075] Various subclasses of the formula (12) include:



[0076] Various subclasses of the formula (13) include:




and




[0077]

Various subclasses of the formula (14) include:




Various pentacyclines of the invention are also shown in *Figure 14*.

[0078] In certain embodiments, the tetracycline analogs of the present invention are represented by the formula:



wherein X is nitrogen, sulfur, and oxygen, and R_1 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , and n are defined as above with the caveat that when X is S or O, R_1 is absent.

[0079] Other classes of compounds of the invention include dicyclines of the formula (15).



wherein R_3 , R_4 , and R_5 are as defined above. P is hydrogen or a protecting group. R_9 is hydrogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_1$; $-CN$; $-SCN$; $-SR_1$; or $-N(R_1)_2$; wherein each occurrence of R_1 is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. In certain embodiments, R_9 is hydrogen or lower (C_1 - C_6) alkyl, alkenyl, or alkynyl. In other embodiments, R_9 is a vinyl group. In yet other embodiments, R_9 is a substituted or unsubstituted aryl group. In still other embodiments, R_9 is a substituted or unsubstituted heterocyclic group.

R₁₀ is cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched aryl; or substituted or unsubstituted, branched or unbranched heteroaryl moiety. In certain embodiments, R₁₀ is a substituted or unsubstituted phenyl ring. In certain embodiments, R₁₀ is a substituted or unsubstituted heterocyclic ring. In certain embodiments, R₁₀ is a substituted or unsubstituted aryl ring. In other embodiments, R₁₀ is a lower (C₁-C₆) alkyl, alkenyl, or alkynyl group.


Methods of Synthesis

[0080] The present invention also includes all steps and methodologies used in preparing the compounds of the invention as well as intermediates along the synthetic route. The present invention provides for the modular synthesis of tetracyclines and its various analogs by joining a highly functionalized chiral enone, which will become the A- and B-rings of the tetracycline core, with a molecule which will become the D-ring of the tetracycline core. The joining of these two intermediates results in the formation of the C-ring, preferably in an enantioselective manner. This methodology also allows for the synthesis of pentacyclines, hexacyclines, or higher ring systems as well as the incorporation of heterocycles into the ring system. In particular, the joining of these two fragments includes various nucleophilic addition reactions and cycloaddition reactions with enone (9) as described above.

[0081] The synthesis begins with the preparation of the enone (9) starting from benzoic acid. As shown in *Figure 2*, the first step of the synthesis involves the microbial dihydroxylation of benzoic acid using *Alcaligenes eutrophus*. The diol (1 in *Figure 2*), which is preferably optically pure, then undergoes hydroxyl-directed epoxidation to yield the allylic epoxide (2 in *Figure 2*). Protection and rearrangement of allylic epoxide 2 yielded the isomeric allylic epoxide (3 in *Figure 2*). The metalated isoxazole (4 in *Figure 2*) was added to the isomeric allylic epoxide to yield 5 (*Figure 2*), which was subsequently metalated to close the six-membered ring by nucleophilic attack of the epoxide. The intermediate 6 (*Figure 2*) was then rearranged, deprotected, and oxidized to yield the chiral enone 9 (*Figure 2*). As will be appreciated by one of

skill in this art, functionalization and rearrangement of intermediates 6, 7, 8, and 9 in Figure 2 will allow for the preparation of different class of compounds of the invention.

[0082] In one embodiment, enone (9) is reacted with an anion resulting from the deprotonation of toluate (6). The toluate of formula:



wherein R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_A$; $=O$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_7 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_G$; $=O$; $-C(=O)R_G$; $-CO_2R_G$; $-CN$; $-SCN$; $-SR_G$; $-SOR_G$; $-SO_2R_G$; $-NO_2$; $-N(R_G)_2$; $-NHC(O)R_G$; or $-C(R_G)_3$; wherein each occurrence of R_G is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety; and


n is an integer in the range of 0 to 3, inclusive;

R_9 is $-OR_I$; $-CN$; $-SCN$; $-SR_I$; or $-N(R_I)_2$; wherein each occurrence of R_I is

independently a hydrogen, a protecting group; a cyclic or acyclic, substituted or unsubstituted aliphatic moiety; a cyclic or acyclic, substituted or unsubstituted aliphatic heteroaliphatic moiety; a substituted or unsubstituted aryl moiety; or a substituted or unsubstituted heteroaryl moiety; and

P is selected from the group consisting of hydrogen, lower (C_1 - C_6) alkyl group, an acyl group, and a protecting group;

is deprotonated under basic conditions (*e.g.*, LDA, HMDS), and the resulting anion is reacted with an enone of formula:



wherein R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_C$; $=O$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;


R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_D$; $=O$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety;

alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_5 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_E$; $-CN$; $-SCN$; $-SR_E$; or $-N(R_E)_2$; wherein each occurrence of R_E is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_6 is selected from the group consisting of hydrogen, halogen, substituted or unsubstituted aliphatic, substituted or unsubstituted heteroaliphatic, substituted or unsubstituted alkoxy, $-OH$, $-CN$, $-SCN$, $-SH$, alkylthio, arylthio, $-NO_2$, amino, alkyl amino, and dialkyl amino groups; and


P is independently selected from the group consisting of hydrogen or a protecting group;
to form the product:



wherein R_1 , R_3 , R_4 , R_5 , R_7 , P , and n are as defined above;


R_2 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_B$; $=O$; $-C(=O)R_B$; $-CO_2R_B$; $-CN$; $-SCN$; $-SR_B$; $-SOR_B$; $-SO_2R_B$; $-NO_2$; $-N(R_B)_2$; $-NHC(O)R_B$; or $-C(R_B)_3$; wherein each occurrence of R_B is independently a hydrogen, a protecting group, an aliphatic

moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. As will be appreciated by one of skill in this art, the toluate may be further substituted in certain embodiments. In addition, the phenyl ring of the toluate may be substituted for an aromatic heterocyclic ring such as a pyridine ring as shown in *Figures 11 and 13*. Other examples of carbocyclic and heterocyclic analogs of toluate (6) include:



Other toluates are shown in *Figure 21*. In certain embodiments, polycyclic toluates are used in the Michael-Dieckmann reaction sequence to form pentacyclines, hexacyclines, or higher cyclines. Toluates useful in preparing pentacyclines are exemplified by the

formula:



wherein R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_A$; $=O$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

each R_7 is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_G$; $=O$; $-C(=O)R_G$; $-CO_2R_G$; $-CN$; $-SCN$; $-SR_G$; $-SOR_G$; $-SO_2R_G$; $-NO_2$; $-N(R_G)_2$; $-NHC(O)R_G$; or $-C(R_G)_3$; wherein each occurrence of R_G is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;



represents a substituted or unsubstituted aryl, heteroaryl, carbocyclic, or heterocyclic moiety, in which each occurrence of X is selected from the group consisting of $-O-$, $-S-$, $-NR_8-$, $-C(R_8)_2-$;

R_8 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted,


branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_H$; $=O$; $-C(=O)R_H$; $-CO_2R_H$; $-CN$; $-SCN$; $-SR_H$; $-SOR_H$; $-SO_2R_H$; $-NO_2$; $-N(R_H)_2$; $-NHC(O)R_H$; or $-C(R_H)_3$; wherein each occurrence of R_H is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

n is an integer in the range of 1 to 5, inclusive; and

the bonds between adjacent X moieties are either single or double bonds; and

R_9 is selected from the group consisting of substituted or unsubstituted aryl or heteroaryl groups.

[0083] In another embodiment, enone (9) is reacted with an anion, which is generated through metallation (*e.g.*, metal-halogen exchange, metal-metalloid exchange, lithium-halogen exchange, lithium-tin exchange, *etc.* by reacting the toluate with the appropriate metal reagent) of a toluate of the the following formula:



wherein R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_A$; $=O$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety;


alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_7 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_G$; $=O$; $-C(=O)R_G$; $-CO_2R_G$; $-CN$; $-SCN$; $-SR_G$; $-SOR_G$; $-SO_2R_G$; $-NO_2$; $-N(R_G)_2$; $-NHC(O)R_G$; or $-C(R_G)_3$; wherein each occurrence of R_G is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

n is an integer in the range of 0 to 3, inclusive;

R_9 is selected from the group consisting of substituted or unsubstituted aryl or heteroaryl groups; and

Y is a halogen or $Sn(R_Y)_3$, wherein R_Y is alkyl. The anion generated is reacted with an enone of formula:



wherein R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_C$; $=O$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy;


or heteroarylthio moiety;

R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_D$; $=O$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_5 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_E$; $-CN$; $-SCN$; $-SR_E$; or $-N(R_E)_2$; wherein each occurrence of R_E is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_6 is selected from the group consisting of hydrogen, halogen, substituted or unsubstituted aliphatic, substituted or unsubstituted heteroaliphatic, substituted or unsubstituted alkoxy, $-OH$, $-CN$, $-SCN$, $-SH$, alkylthio, arylthio, $-NO_2$, amino, alkyl amino, and dialkyl amino groups; and

P is independently selected from the group consisting of hydrogen or a protecting group; to generate the product of formula:




wherein R_1 , R_3 , R_4 , R_5 , R_7 , P , and n are as defined above; and

R_2 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_B$; $=O$; $-C(=O)R_B$; $-CO_2R_B$; $-CN$; $-SCN$; $-SR_B$; $-SOR_B$; $-SO_2R_B$; $-NO_2$; $-N(R_B)_2$; $-NHC(O)R_B$; or $-C(R_B)_3$; wherein each occurrence of R_B is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety.

[0084] Any metal may be used in the metallation reaction to generate the metal anionic reagent to be reacted with the enone. In certain embodiments, the metal is a Group I element on the periodic chart. In other embodiments, the metal is a Group II element on the periodic chart. In other embodiments, the metal is a transition metal. Exemplary metals useful in the metallation reaction include sodium, lithium, calcium, aluminium, cadmium, copper, beryllium, arsenic, antimony, tin, magnesium, titanium, zinc, manganese, iron, cobalt, nickel, zinc, platinum, palladium, mercury, and ruthenium. In certain preferred embodiments, the metal is chosen from lithium, magnesium, titanium, zinc, and copper. In yet other embodiments, the metal is magnesium, lithium, sodium, beryllium, zinc, mercury, arsenic, antimony, or tin. In certain particular embodiments, a lithium-halogen exchange is used. The lithium-halogen exchange may be performed *in situ* in the presence of the enone. The lithium-halogen exchange may be performed using any lithium reagent including, for example, alkylolithium reagents, *n*-butyllithium, *t*-butyllithium, phenyl lithium, mesityl lithium, and methyllithium. In certain embodiments, other organometallics reagents are generated and reacted with the enone. Examples include Grignard reagents, zero-valent metal complexes, ate complexes, *etc.* In certain embodiments, the metal reagent is a magnesium reagent including, but not limited to, magnesium metal, magnesium anthracene, activated magnesium turnings, *etc.* In certain embodiments, the reagent is zinc-based. The reagent may be generated *in situ* in the presence of the enone, or the

reagent may be generated separately and later contacted with the enone. In certain embodiments, milder conditions for the cyclization are used (*e.g.*, a zinc reagent).


[0085] As will be appreciated by one of skill in this art, the toluate may be further substituted in certain embodiments. In addition, the phenyl ring of the toluate may be substituted for an aromatic heterocyclic ring or ring system such as a pyridine ring. Examples of carbocyclic and heterocyclic analogs of toluate include:



In certain embodiments, the halogen Y is bromine. In other embodiments, Y is iodine. In yet other embodiments, Y is chloride. In certain embodiments, Y is a metalloid (*e.g.*, tin, selenium, tellurium, *etc.*). In certain embodiments, Y is $-\text{SnR}_3$, wherein each occurrence of R is independently alkyl (*e.g.*, $-\text{Sn}(\text{CH}_3)_3$). After the metallation

reaction, Y is a metal such as lithium, magnesium, zinc, copper, antimony, sodium, *etc.* In certain embodiments, R₁ is hydrogen or lower alkyl (C₁-C₆). In certain particular embodiments, R₁ is hydrogen. Other toluates are shown in *Figure 21*.

[0086] In other embodiments, polycyclic toluates may be used to prepare pentacyclines, hexacyclines, or higher cyclines. Toluates useful in the preparation of such cyclines are of the formula:



wherein R₁ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; -OR_A; =O; -C(=O)R_A; -CO₂R_A; -CN; -SCN; -SR_A; -SOR_A; -SO₂R_A; -NO₂; -N(R_A)₂; -NHC(O)R_A; or -C(R_A)₃; wherein each occurrence of R_A is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

each R₇ is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; -OR_G; =O; -C(=O)R_G; -CO₂R_G; -CN; -SCN; -SR_G; -SOR_G; -SO₂R_G; -NO₂; -N(R_G)₂; -NHC(O)R_G; or -C(R_G)₃; wherein each occurrence of R_G is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;



represents a substituted or unsubstituted aryl, heteroaryl, carbocyclic, or heterocyclic moiety, in which each occurrence of X is selected from the group consisting of -O-, -S-, -NR₈-, -C(R₈)₂-;


R₈ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; -OR_H; =O; -C(=O)R_H; -CO₂R_H; -CN; -SCN; -SR_H; -SOR_H; -SO₂R_H; -NO₂; -N(R_H)₂; -NHC(O)R_H; or -C(R_H)₃; wherein each occurrence of R_H is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

n is an integer in the range of 1 to 5, inclusive; and

the bonds between adjacent X moieties are either single or double bonds;


R₉ is selected from the group consisting of substituted or unsubstituted aryl or heteroaryl groups; and

Y is a halogen or Sn(R_Y)₃, wherein R_Y is alkyl. In certain embodiments, the halogen Y is bromine. In certain embodiments, the halogen Y is bromine. In other embodiments, Y is iodine. In yet other embodiments, Y is chloride. In certain embodiments, Y is a metalloid (*e.g.*, tin, selenium, tellurium, *etc.*). In certain embodiments, Y is -SnR₃, wherein each occurrence of R is independently alkyl (*e.g.*, -Sn(CH₃)₃). After the metallation reaction, Y is a metal such as lithium, magnesium, zinc, copper, sodium, mercury, antimony, *etc.* In certain embodiments, R₁ is hydrogen or lower alkyl (C₁-C₆). In certain particular embodiments, R₁ is hydrogen. In certain embodiments, R₉ is phenyl or substituted phenyl. In certain embodiments, *ortho*-R₇ is alkoxy such as methoxy. In other embodiments, R₇ is hydrogen. Exemplary polycyclic toluates include:




[0087]


Compounds of the formula below with a heterocyclic C-ring:



may be prepared by Michael-Dieckmann closure of a D-ring precursor derived from the corresponding anilide, phenol, or thiophenol. A representative example using anthranilic acid (*i.e.*, anilide as the nucleophile in the Michael addition reaction) is shown below:



[0088] In another embodiment, the enone (9) is reacted with a benzocyclobutenol in an *o*-quinone dimethide Diels-Alder reaction. The enone of formula:



wherein R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_C$; $=O$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;


R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_D$; $=O$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently a hydrogen, a protecting group, an aliphatic

moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_5 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_E$; $-CN$; $-SCN$; $-SR_E$; or $-N(R_E)_2$; wherein each occurrence of R_E is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_6 is selected from the group consisting of hydrogen, halogen, substituted or unsubstituted aliphatic, substituted or unsubstituted heteroaliphatic, substituted or unsubstituted alkoxy, $-OH$, $-CN$, $-SCN$, $-SH$, alkylthio, arylthio, $-NO_2$, amino, alkyl amino, and dialkyl amino groups;

P is independently selected from the group consisting of hydrogen or a protecting group; is reacted under suitable conditions (*e.g.*, heat) with a benzocyclobutenol of formula:



wherein R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_A$; $=O$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety;


alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_7 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_G$; $=O$; $-C(=O)R_G$; $-CO_2R_G$; $-CN$; $-SCN$; $-SR_G$; $-SOR_G$; $-SO_2R_G$; $-NO_2$; $-N(R_G)_2$; $-NHC(O)R_G$; or $-C(R_G)_3$; wherein each occurrence of R_G is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

P are each selected independently from the group consisting of hydrogen or a protecting group; and

n is an integer in the range of 0 to 3, inclusive;

to form the product of formula:




wherein R_1 , R_3 , R_4 , R_5 , R_6 , R_7 , and P are defined as above; and

R_2 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_B$; $=O$; $-C(=O)R_B$; $-CO_2R_B$; $-CN$; $-SCN$; $-SR_B$; $-SOR_B$; $-SO_2R_B$; $-NO_2$; $-N(R_B)_2$; $-NHC(O)R_B$; or $-C(R_B)_3$; wherein each occurrence of R_B is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy;

or heteroarylthio moiety. As will be appreciated by one of skill in this art, the reactants may be substituted further and still fall within the claimed invention. For example, the phenyl ring of the benzocyclobutenol ring may be further substituted.

[0089] In another embodiment, the enone is reacted with a diene in a Diels-Alder reaction to yield a tricyclic. The enone of formula:



wherein R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_C$; $=O$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;


R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_D$; $=O$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_5 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted,

branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; -OR_E; -CN; -SCN; -SR_E; or -N(R_E)₂; wherein each occurrence of R_E is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R₆ is selected from the group consisting of hydrogen, halogen, substituted or unsubstituted aliphatic, substituted or unsubstituted heteroaliphatic, substituted or unsubstituted alkoxy, -OH, -CN, -SCN, -SH, alkylthio, arylthioxy, -NO₂, amino, alkyl amino, and dialkyl amino groups; are as defined above; and

P is independently selected from the group consisting of hydrogen or a protecting group; is reacted under suitable conditions (*e.g.*, heat) with a diene of formula:




wherein R₁ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; -OR_A; =O; -C(=O)R_A; -CO₂R_A; -CN; -SCN; -SR_A; -SOR_A; -SO₂R_A; -NO₂; -N(R_A)₂; -NHC(O)R_A; or -C(R_A)₃; wherein each occurrence of R_A is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety; and

P are each selected independently from the group consisting of hydrogen and


protecting groups;

to yield a protected tricycline of formula:



wherein R_2 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_B$; $=O$; $-C(=O)R_B$; $-CO_2R_B$; $-CN$; $-SCN$; $-SR_B$; $-SOR_B$; $-SO_2R_B$; $-NO_2$; $-N(R_B)_2$; $-NHC(O)R_B$; or $-C(R_B)_3$; wherein each occurrence of R_B is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. As will be appreciated by one of skill in this art, the enone and diene may be further substituted and still be encompassed within the present invention.

[0090] In yet another embodiment, the enone is reacted with an anion of a phthalide or cyano-phthalide. The enone of formula:



wherein R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_C$; $=O$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein

each occurrence of R_C is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;


R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_D$; $=O$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_5 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_E$; $-CN$; $-SCN$; $-SR_E$; or $-N(R_E)_2$; wherein each occurrence of R_E is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_6 is selected from the group consisting of hydrogen, halogen, substituted or unsubstituted aliphatic, substituted or unsubstituted heteroaliphatic, substituted or unsubstituted alkoxy, $-OH$, $-CN$, $-SCN$, $-SH$, alkylthio, arylthio, $-NO_2$, amino, alkyl amino, and dialkyl amino groups; and

P is independently selected from the group consisting of hydrogen or a protecting group;

is reacted under basic conditions (*e.g.*, LDA, Ph_3CLi) with the anion of the phthalide of formula:




wherein R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_A$; $=O$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

R_7 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_G$; $=O$; $-C(=O)R_G$; $-CO_2R_G$; $-CN$; $-SCN$; $-SR_G$; $-SOR_G$; $-SO_2R_G$; $-NO_2$; $-N(R_G)_2$; $-NHC(O)R_G$; or $-C(R_G)_3$; wherein each occurrence of R_G is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

P are each selected independently from the group consisting of hydrogen, lower alkyl group, acyl group, or a protecting group; and

n is an integer in the range of 0 to 3, inclusive;
to yield a product of formula:



wherein R_2 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_B$; $=O$; $-C(=O)R_B$; $-CO_2R_B$; $-CN$; $-SCN$; $-SR_B$; $-SOR_B$; $-SO_2R_B$; $-NO_2$; $-N(R_B)_2$; $-NHC(O)R_B$; or $-C(R_B)_3$; wherein each occurrence of R_B is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety.

[0091] The products of the above reactions are then further functionalized, reduced, oxidized, rearranged, protected, and deprotected to yield the final desired product. Various exemplary reactions used in the final syntheses of the compounds of the invention are shown in *Figure 2, 3, 11, 12, and 13*. As will be appreciated by one of skill in the art, various isolation and purification techniques including flash chromatography, crystallization, distillation, HPLC, thin layer chromatography, extraction, filtration, *etc.* may be used in the course of synthesizing compounds of the invention. These techniques may be used in the preparation or purification of intermediates, reagents, products, starting materials, or solvents.

Pharmaceutical Compositions

[0092] This invention also provides a pharmaceutical preparation comprising at least one of the compounds as described above and herein, or a pharmaceutically acceptable derivative thereof, which compounds inhibit the growth of or kill microorganisms, and, in certain embodiments of special interest are inhibit the growth of or kill tetracycline-resistant organisms including chlortetracycline-resistant

organisms, oxytetracycline-resistant organisms, demeclocycline-resistant organisms, doxycycline-resistant organisms, minocycline-resistant organisms, or any organisms resistant to antibiotics of the tetracycline class used in human or veterinary medicine. In other embodiments, the compounds show cytostatic or cytotoxic activity against neoplastic cells such as cancer cells. In yet other embodiments, the compounds inhibit the growth of or kill rapidly dividing cells such as stimulated inflammatory cells.

[0093] As discussed above, the present invention provides novel compounds having antimicrobial and antiproliferative activity, and thus the inventive compounds are useful for the treatment of a variety of medical conditions including infectious diseases, cancer, autoimmune diseases, inflammatory diseases, and diabetic retinopathy. Accordingly, in another aspect of the present invention, pharmaceutical compositions are provided, wherein these compositions comprise any one of the compounds as described herein, and optionally comprise a pharmaceutically acceptable carrier. In certain embodiments, these compositions optionally further comprise one or more additional therapeutic agents, *e.g.*, another anti-microbial agent or another anti-proliferative agent. In other embodiments, these compositions further comprise an anti-inflammatory agent such as aspirin, ibuprofen, acetaminophen, *etc.*, pain reliever, or anti-pyretic.

[0094] It will also be appreciated that certain of the compounds of the present invention can exist in free form for treatment, or where appropriate, as a pharmaceutically acceptable derivative thereof. According to the present invention, a pharmaceutically acceptable derivative includes, but is not limited to, pharmaceutically acceptable salts, esters, salts of such esters, or any other adduct or derivative which upon administration to a patient in need is capable of providing, directly or indirectly, a compound as otherwise described herein, or a metabolite or residue thereof, *e.g.*, a prodrug.

[0095] As used herein, the term "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M.

Berge, *et al.* describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 66: 1-19, 1977; incorporated herein by reference. The salts can be prepared *in situ* during the final isolation and purification of the compounds of the invention, or separately by reacting the free base functionality with a suitable organic or inorganic acid. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid, or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, harnisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate, and aryl sulfonate.

[0096] Additionally, as used herein, the term "pharmaceutically acceptable ester" refers to esters which hydrolyze *in vivo* and include those that break down readily in the human body to leave the parent compound or a salt thereof. Suitable ester groups include, for example, those derived from pharmaceutically acceptable aliphatic carboxylic acids, particularly alkanoic, alkenoic, cycloalkanoic and alkanedioic acids, in which each alkyl or alkenyl moiety advantageously has not more than 6 carbon atoms. Examples of particular esters include formates, acetates,

propionates, butyrates, acrylates and ethylsuccinates. In certain embodiments, the esters are cleaved by enzymes such as esterases.

[0097] Furthermore, the term "pharmaceutically acceptable prodrugs" as used herein refers to those prodrugs of the compounds of the present invention which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals with undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention. The term "prodrug" refers to compounds that are rapidly transformed *in vivo* to yield the parent compound of the above formula, for example by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, Vol. 14 of the A.C.S. Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference.

[0098] As described above, the pharmaceutical compositions of the present invention additionally comprise a pharmaceutically acceptable carrier, which, as used herein, includes any and all solvents, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. Remington's *Pharmaceutical Sciences*, Fifteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1975) discloses various carriers used in formulating pharmaceutical compositions and known techniques for the preparation thereof. Except insofar as any conventional carrier medium is incompatible with the anti-cancer compounds of the invention, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutical composition, its use is contemplated to be within the scope of this invention. Some examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; Cremophor; Solutol;

excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil; sesame oil; olive oil; corn oil and soybean oil; glycols; such a propylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

Uses of Compounds and Pharmaceutical Compositions

[0099] The invention further provides a method of treating infections and inhibiting tumor growth. The method involves the administration of a therapeutically effective amount of the compound or a pharmaceutically acceptable derivative thereof to a subject (including, but not limited to a human or animal) in need of it.

[00100] The compounds and pharmaceutical compositions of the present invention may be used in treating or preventing any disease or conditions including infections (*e.g.*, skin infections, GI infection, urinary tract infections, genito-urinary infections, systemic infections), proliferative diseases (*e.g.*, cancer), and autoimmune diseases (*e.g.*, rheumatoid arthritis, lupus). The compounds and pharmaceutical compositions may be administered to animals, preferably mammals (*e.g.*, domesticated animals, cats, dogs, mice, rats), and more preferably humans. Any method of administration may be used to deliver the compound of pharmaceutical compositions to the animal. In certain embodiments, the compound or pharmaceutical composition is administered orally. In other embodiments, the compound or pharmaceutical composition is administered parenterally.

[00101] In yet another aspect, according to the methods of treatment of the present invention, bacteria are killed, or their growth is inhibited by contacting the bacteria with an inventive compound or composition, as described herein. Thus, in still another aspect of the invention, a method for the treatment of infection is provided comprising administering a therapeutically effective amount of an inventive compound,

or a pharmaceutical composition comprising an inventive compound to a subject in need thereof, in such amounts and for such time as is necessary to achieve the desired result. In certain embodiments of the present invention a "therapeutically effective amount" of the inventive compound or pharmaceutical composition is that amount effective for killing or inhibiting the growth of bacteria. The compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for killing or inhibiting the growth of bacteria. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular compound, its mode of administration, its mode of activity, and the like. The compounds of the invention are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts.

[00102] Furthermore, after formulation with an appropriate pharmaceutically acceptable carrier in a desired dosage, the pharmaceutical compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, as an oral or nasal spray, or the like, depending on the severity of the infection being treated. In certain embodiments, the compounds of the invention may be administered orally or parenterally at dosage levels sufficient to deliver from about 0.001 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 50 mg/kg, preferably from about 0.1 mg/kg to about 40 mg/kg, preferably from about

0.5 mg/kg to about 30 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 0.1 mg/kg to about 10 mg/kg, and more preferably from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect. The desired dosage may be delivered three times a day, two times a day, once a day, every other day, every third day, every week, every two weeks, every three weeks, or every four weeks. In certain embodiments, the desired dosage may be delivered using multiple administrations (*e.g.*, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, or more administrations).

[00103] Liquid dosage forms for oral and parenteral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents. In certain embodiments for parenteral administration, the compounds of the invention are mixed with solubilizing agents such as Cremophor, alcohols, oils, modified oils, glycols, polysorbates, cyclodextrins, polymers, and combinations thereof.

[00104] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any

bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[00105] The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[00106] In order to prolong the effect of a drug, it is often desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

[00107] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

[00108] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia,

c) humectants such as glycerol, d) disintegrating agents such as agar--agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

[00109] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[00110] The active compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, *e.g.*, tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active

ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

[00111] Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, ear drops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispensing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

[00112] It will also be appreciated that the compounds and pharmaceutical compositions of the present invention can be employed in combination therapies, that is, the compounds and pharmaceutical compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies employed may achieve a desired effect for the same disorder (for example, an inventive compound may be administered concurrently with another anticancer agent), or they may achieve different effects (*e.g.*, control of any adverse effects).

[00113] In still another aspect, the present invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention, and in certain embodiments, includes an additional approved therapeutic agent for use as a combination therapy. Optionally associated with such container(s) can be a notice in

the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceutical products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

[00114] These and other aspects of the present invention will be further appreciated upon consideration of the following Examples, which are intended to illustrate certain particular embodiments of the invention but are not intended to limit its scope, as defined by the claims.

Examples

Example 1-Synthesis of (-)-Tetracycline


[00115] **General Procedures.** All reactions were performed in flame-dried round bottomed or modified Schlenk (Kjeldahl shape) flasks fitted with rubber septa under a positive pressure of argon, unless otherwise noted. Air- and moisture-sensitive liquids and solutions were transferred via syringe or stainless steel cannula. Where necessary (so noted), solutions were deoxygenated by alternative freeze (liquid nitrogen)/evacuation/ thaw cycles ($\geq$ three iterations). Organic solutions were concentrated by rotary evaporation at ~ 25 Torr (house vacuum). Flash column chromatography was performed on silica gel (60 Å, standard grade) as described by Still *et al.* (Still, W. C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, *43*, 2923-2925; incorporated herein by reference). Analytical thin-layer chromatography was performed using glass plates pre-coated with 0.25 mm 230-400 mesh silica gel impregnated with a fluorescent indicator (254 nm). Thin layer chromatography plates were visualized by exposure to ultraviolet light and/or exposure to ceric ammonium molybdate or an acidic solution of *p*-anisaldehyde followed by heating on a hot plate.

[00116] **Materials.** Commercial reagents and solvents were used as received with the following exceptions. Chlorotrimethylsilane, triethylamine, diisopropylamine, 2,2,6,6-tetramethylpiperidine, *N,N,N',N'*-tetramethylethylenediamine, DMPU, HMPA, and *N,N*-diisopropylethylamine were distilled from calcium hydride under dinitrogen atmosphere. Benzene, dichloromethane, ethyl ether, methanol, pyridine,

tetrahydrofuran, hexane, acetonitrile, *N,N*-dimethylformamide, and toluene were purified by the method of Pangborn *et al.* (Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. *Organometallics* **1996**, *15*, 1518-1520; incorporated herein by reference). The molarity of *n*-butyllithium, *s*-butyllithium, and *t*-butyllithium were determined by titration with a tetrahydrofuran solution of 2-butanol using triphenylmethane as an indicator (Duhamel, L.; Palquevent, J.-C. *J. Org. Chem.* **1979**, *44*, 3404-3405; incorporated herein by reference).

[00117] Instrumentation. Proton nuclear magnetic resonance (^1H NMR) spectra and carbon nuclear magnetic resonance (^{13}C NMR) were recorded with Varian Unity/Inova 600 (600 MHz), Varian Unity/Inova 500 (500 MHz/125 MHz), or Varian Mercury 400 (400 MHz/100 MHz) NMR spectrometers. Chemical shifts for protons are reported in parts per million scale (δ scale) downfield from tetramethylsilane and are referenced to residual protium in the NMR solvents (CHCl_3 : δ 7.26, $\text{C}_6\text{D}_5\text{H}$: δ 7.15, D_2HCO_2 : δ 3.31, CDHCl_2 : δ 5.32, $(\text{CD}_2\text{H})\text{CD}_3\text{SO}$: δ 2.49). Chemical shifts for carbon are reported in parts per million (δ scale) downfield from tetramethylsilane and are referenced to the carbon resonances of the solvent (CDCl_3 : δ 77.0, C_6D_6 : δ 128.0, D_3CO_2 : δ 44.9, CD_2Cl_2 : δ 53.8, $(\text{CD}_3)_2\text{SO}$: δ 39.5). Data are represented as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br = broad), integration, coupling constant in Hz, and assignment. Infrared (IR) spectra were obtained using a Perkin-Elmer 1600 FT-IR spectrophotometer referenced to a polystyrene standard. Data are represented as follows: frequency of the absorption (cm^{-1}), intensity of absorption (s = strong, sb = strong broad, m = medium, w = weak, br = broad), and assignment (where appropriate). Optical rotations were determined on a JASCO DIP-370 digital polarimeter equipped with a sodium lamp source using a 200- μL or 2-mL solution cell. High resolution mass spectra were obtained at the Harvard University Mass Spectrometry Facilities.

Microbial Dihydroxylation Product DRS1:



Preparation of Glycerol Stock Solutions

[00118] *Alcaligenes eutrophus* B9 cells (lyophilized powder, 20 mg, generously supplied by Prof. George D. Hegeman (Indiana University); Reiner, A. M.; Hegeman, G. D. *Biochemistry* **1971**, *10*, 2530.) were suspended in nutrient broth (5 mL, prepared by dissolving 8 g of Difco Bacto® Nutrient Broth in 1 L of nanopure water followed by sterilization in an autoclave at 125 °C) in a 20-mL sterile culture tube. Aqueous sodium succinate solution (16.7 µL of a 2.5 M aqueous solution, 5 mM final concentration) was added, and the culture tube was shaken at 250 rpm at 30 °C until cell growth became apparent (3 d). An aliquot (250 µL) of the cellular suspension was then transferred to 5 mL of Hutner's mineral base medium (HMB, see paragraph below) containing sodium succinate (16.7 µL of a 2.5 M aqueous solution, 5 mM final concentration) in a 20-mL sterile culture tube. The culture tube was shaken at 250 rpm for 2 d at 30 °C, whereupon an aliquot (250 µL) of the fermentation solution was subcultured in a sterile Erlenmeyer flask containing 50 mL of HMB and aqueous sodium succinate solution (167 µL of a 2.5 M solution, 5 mM final concentration). The flask was shaken at 250 rpm for 24 h at 30 °C. The resulting solution was used directly for the preparation of glycerol stock solutions. Thus, a portion of the subcultured cellular suspension (5 mL) was diluted with an equal volume of sterile glycerol, and the resulting solution was divided equally into ten 2-mL sterile Eppendorf tubes. The individual stock solutions were then stored at –80 °C.

Hutner's Mineral Base Medium

[00119] Hutner's mineral base medium (HMB) was prepared as follows. Solid potassium hydroxide (400 mg) was dissolved in 500 mL of nanopure water in a 2-L Erlenmeyer flask. Nitrilotriacetic acid (200 mg), magnesium sulfate (283 mg), calcium chloride dihydrate (67 mg), ammonium molybdate (0.2 mg), iron (II) sulfate (2.0 mg), Hutner's Metals 44 solution (1 mL, see paragraph below), ammonium sulfate (1.0 g), potassium dihydrogen phosphate (2.72 g) and sodium monohydrogen phosphate heptahydrate (5.36 g) were added sequentially. The solution was diluted to a total volume of 1 L and the pH was adjusted to 6.8 with concentrated hydrochloric acid. The medium was sterilized by filtration or by heating in an autoclave.


[00120] Hutner's Metals 44 solution was prepared as follows. Concentrated sulfuric acid (100 μ L) was added to nanopure water (50 mL) in a 250-mL Erlenmeyer flask. Solid EDTA (0.50 g), zinc sulfate heptahydrate (2.20 g), iron (II) sulfate heptahydrate (1.0 g), copper (I) sulfate (0.39 g), cobalt (II) nitrate hexahydrate (50 mg) and sodium tetraborate decahydrate (36 mg) were then added in sequence, followed by 50 mL of nanopure water.

Cellular Dihydroxylation of Sodium Benzoate

[00121] A sterile pipette tip was streaked across the surface of a frozen glycerol stock solution to produce small shards (ca. 10 mg). The frozen shards were added to a sterile 125 mL Erlenmeyer flask containing HMB (25 mL) and aqueous sodium succinate solution (140 μ L of a 1.5 M solution, 5 mM final concentration). The flask was shaken at 250 rpm for 2 days at 30 °C. An aliquot (10 mL) of the white, heterogeneous solution was transferred using a sterile pipette to a mammalian cell growth jar containing HMB (6 L) and aqueous sodium succinate solution (20 mL of a 1.5 M solution, 5 mM final concentration). The jar was warmed on a hot plate to an internal temperature of 30 °C; cotton-filtered air was sparged through the medium. After 2 days, the white, heterogeneous solution was treated with aqueous sodium benzoate solution (18 mL of a 1.0 M solution) and aqueous sodium succinate solution (10 mL of a 1.5 M solution), inducing dihydroxylation. The resulting mixture was aerated vigorously for 6 hours at an internal temperature of 30 °C. After induction, sufficient aqueous sodium benzoate solution (24 to 48 mL of a 1.0 M solution, depending on the rate of consumption) was added hourly to maintain a concentration of 10-20 mM (determined by UV absorbance at 225 nm). Aqueous sodium succinate solution (10 mL of a 1.5 M solution) was added every fourth hour. These additions proceeded over 18 hours, then the solution was aerated overnight at an internal temperature of 30 °C, to ensure complete conversion. The fermentation broth was centrifuged, in portions, at 6000 rpm (Sorvall GS-3 rotor, model SLA-3000) to remove cellular material. The supernatant was concentrated to a volume of 400 mL using a rotary evaporator (bath temperature <45 °C). The concentrate was cooled to 0 °C and then acidified to pH 3.0 using concentrated aqueous hydrochloric acid. The acidified

aqueous solution was extracted repeatedly with ethyl acetate (8 × 500 mL, 4 × 800 mL, 8 × 1 L). The ethyl acetate extracts were dried over sodium sulfate before concentration, using a rotary evaporator (bath temperature <45 °C), providing a pale yellow solid residue. Trituration of the residue with dichloromethane (2 × 200 mL) followed by drying in vacuo afforded pure (1*S*,2*R*)-1,2-dihydroxycyclohexa-3,5-diene-1-carboxylic acid (**DRS1**) as a white powder mp 95-96 °C dec (38 g, 74%, $[\alpha]_D -114.8$ (*c* 0.5 in EtOH), lit., $[\alpha]_D -106$ (*c* 0.5 in EtOH) Jenkins, G. N.; Ribbons, D. W.; Widdowson, D. A.; Slawin, A. M. Z.; Williams, D. J. *J. Chem. Soc. Perkin Trans. 1* **1995**, 2647.).


Epoxide **DRS2**:



[00122] *m*-Chloroperoxybenzoic acid (*m*CPBA was purified as follows: 50 g of 77% *m*CPBA (Alrich) was dissolved in benzene (1 L), the benzene solution was then washed with pH 7.4 phosphate buffer (3 × 1 L) and dried over Na₂SO₄ for 3 hours and concentrated (<40 °C, thermal detonation hazard) to provide pure *m*CPBA as a white solid; 10.7 g, 62.3 mmol, 1.2 equiv) was added in three equal portions over 30 min. to a suspension of the microbial dihydroxylation product **DRS1** (8.10 g, 51.9 mmol, 1.0 equiv) in ethyl acetate (400 mL) at 23 °C. The heterogeneous solution was stirred for 10 h, then was diluted with benzene (80 mL) and stirred for 1 h. The supernatant was decanted and the solid residue was triturated with benzene (2 × 15 mL). The resulting pasty solid was dried in vacuo to provide the epoxide **DRS2** as an amorphous white powder (7.36 g, 83%).

[00123] mp 87-91 °C; ¹H NMR (400 MHz, CD₃OD) δ 6.23 (dd, 1H, *J* = 9.6, 3.9 Hz, =CHC(OCH)), 5.92 (dd, 1H, *J* = 9.6, 1.9 Hz, =CHC(CO₂H)), 4.40 (d, 1H, *J* = 1.3 Hz, CHOH), 3.58 (dd, 1H, *J* = 4.4, 1.3 Hz, CHCHOH), 3.49 (m, 1H, =CCHO); ¹³C NMR (100 MHz, CD₃OD) δ 175.8, 135.1, 128.8, 75.4, 70.9, 57.5, 50.3; FTIR (neat), cm⁻¹ 3381 (s, OH), 1738 (s, C=O), 1608 (m), 1255 (m), 1230 (m), 1084 (m, C-O); HRMS (CI) *m/z* calcd for (C₇H₈O₅+NH₄)⁺ 190.0715, found 190.0707.


Epoxide **DJB1**:



[00124] A solution of trimethylsilyldiazomethane in hexanes (2.0 M, 25.5 mL, 51.0 mmol, 1.2 equiv) was added to a solution of the epoxide **DRS2** (7.36 g, 42.8 mmol, 1.0 equiv) in methanol-benzene (1:3, 160 mL) at 23 °C. Extensive gas evolution was observed upon addition. The yellow solution was stirred for 5 min, then was concentrated, affording a light yellow solid. The solid was dried by azeotropic distillation from benzene (2 × 25 mL), and the dried solid was suspended in dichloromethane (200 mL). Triethylamine (20.8 mL, 149 mmol, 3.5 equiv) and *tert*-butyldimethylsilyl trifluoromethanesulfonate (29.4 mL, 128 mmol, 3.0 equiv) were then added in sequence, providing a homogeneous solution. The reaction solution was stirred at 23 °C for 30 min. An aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 300 mL) was added followed by dichloromethane (100 mL). The organic phase was separated and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a brown oil. The product was purified by flash column chromatography (5:95 ethyl acetate-hexanes), affording the epoxide **DJB1** as a light yellow oil (12.4 g, 70% over 2 steps).

R_f 0.50 (1:4 ethyl acetate-hexanes); ¹H NMR (400 MHz, CDCl₃) δ 5.95 (dd, 1H, *J* = 9.8, 3.4 Hz, =CHCOTBS), 5.89 (ddd, 1H, *J* = 9.8, 2.9, 1.5 Hz, =CHCHOCCO₂), 4.63 (d, 1H, *J* = 3.9 Hz, O₂CCCHOTBS), 4.42 (m, 1H, =CCHOTBS), 3.78 (s, 3H, OCH₃), 3.31 (d, 1H, *J* = 2.0 Hz, CHOCCO₂), 0.90 (s, 9H, C(CH₃)₃), 0.89 (s, 9H, C(CH₃)₃), 0.09 (s, 3H, SiCH₃), 0.08 (s, 6H, SiCH₃), 0.07 (s, 3H, SiCH₃); ¹³C NMR (100 MHz, CDCl₃) δ 170.2, 138.7, 122.6, 69.3, 68.4, 59.7, 52.5, 52.0, 25.9, 25.7, 18.3, 18.2, -4.18, -4.27, -4.45, -5.21; FTIR (neat), cm⁻¹ 1759 (m, C=O), 1736 (s, C=O), 1473 (m), 1256 (w), 1253 (s), 1150 (s, C-O), 1111 (m, C-O), 1057 (s, C-O), 940 (m); HRMS (ES) *m/z* calcd for (C₂₀H₃₈O₅Si₂)⁺ 414.2258, found 414.2239.


Isoxazole **MGC2** (Method A):



[00125] Triethylamine (37.5 mL, 0.269 mol, 1.15 equiv), 4-(dimethylamino)pyridine (289 mg, 2.34 mmol, 0.01 equiv), and methanesulfonyl chloride (20.8 mL, 0.269 mol, 1.15 equiv) were added in sequence to a solution of the alcohol **MGC1** (prepared in two steps from commercially available methyl 3-hydroxy-5-isoxazolecarboxylate as previously reported by: Reiss, R.; Schön, M.; Laschat, S.; Jäger, V. *Eur. J. Org. Chem.* **1998**, 473-479.) (48.0 g, 0.234 mol, 1.0 equiv) in dichloromethane (450 mL) at 0 °C. The reaction mixture was stirred at 0 °C for 2.5 h, then was concentrated, affording an orange oil. Chilled dimethylamine (condensed using a cold finger with dry ice/acetone, 26.2 mL, 0.480 mol, 2.0 equiv) was added to a mixture of the orange oil prepared above and *N,N*-dimethylformamide (150 mL) at 0 °C, providing a homogenous solution. The solution was stirred at 0 °C for 2 h, then was allowed to warm to 23 °C; stirring was continued at that temperature for 24 h. The solution was partitioned between saturated aqueous sodium bicarbonate solution-brine (2:1, 300 mL) and ethyl acetate-hexanes (1:1, 500 mL). The organic phase was separated and washed with brine (2 × 200 mL), and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, furnishing a brown residue. The product was purified by flash column chromatography (1:4 to 1:1 ethyl acetate-hexanes), affording the isoxazole **MGC2** as a light yellow oil (40.1 g, 74%).

R_f 0.34 (1:1 ethyl acetate-hexanes); 1H NMR (500 MHz, $CDCl_3$) δ 7.43-7.31 (m, 5H, ArH), 5.82 (s, 1H, =CH), 5.23 (s, 2H, OCH_2Ar), 3.48 (s, 2H, $CH_2N(CH_3)_2$), 2.27 (s, 6H, $N(CH_3)_2$); ^{13}C NMR (125 MHz, $CDCl_3$) δ 171.9, 171.2, 136.1, 128.8, 128.5, 128.7, 94.8, 71.7, 55.1, 45.3; FTIR (neat), cm^{-1} 2950 (s, CH), 1615 (s), 1494 (s), 1452 (s), 1136 (m); HRMS (ES) m/z calcd for $(C_{13}H_{16}N_2O_2)^+$ 232.1212, found 232.1220.


Isoxazole **MGC4**:



[00126] Chilled dimethylamine (condensed into a reaction vessel submerged in a 0 °C bath using a cold finger with dry ice/acetone, 106 mL, 1.94 mol, 2.2 equiv) was added dropwise via cannula to a solution of the isoxazole **MGC3** (prepared in two steps from glyoxylic acid as reported by: Pevarello, P.; Varasi, M. *Synth. Commun.* **1992**, 22, 1939.) (174 g, 0.884 mol, 1.0 equiv) in acetonitrile (2 L) at 0 °C. The reaction mixture was stirred at 0 °C for 2 h, then the cooling bath was removed. The reaction mixture was allowed to warm to 23 °C; stirring was continued at that temperature for 8 h. The mixture was partitioned between brine-saturated aqueous sodium bicarbonate solution (1:1, 1.5 L) and ethyl acetate (1.5 L). The organic phase was separated and the aqueous phase was further extracted with ethyl acetate (3 × 400 mL). The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated to a volume of 500 mL, resulting in the formation of a white precipitate. The concentrate was filtered and the filtrate was concentrated, providing the isoxazole **MGC4** as an orange oil (143 g, 79%). An analytical sample was prepared by flash column chromatography (1:9 to 2:8 ethyl acetate-hexanes), affording the isoxazole **MGC4** as a light yellow oil.

[00127] R_f 0.30 (1:4 ethyl acetate-hexanes); ^1H NMR (300 MHz, CDCl_3) δ 6.26 (s, 1H, vinyl), 3.63 (s, 2H, $\text{CH}_2\text{N}(\text{CH}_3)_2$), 2.30 (s, 6H, $\text{N}(\text{CH}_3)_2$); ^{13}C NMR (100 MHz, CDCl_3) δ 172.1, 140.5, 106.8, 54.5, 45.3; FTIR (neat), cm^{-1} 3137 (w), 2945 (m), 2825 (m), 2778 (m), 1590 (s), 1455 (m), 1361 (m), 1338 (s), 1281 (s), 1041 (m); HRMS (ES) m/z calcd for $(\text{C}_6\text{H}_9\text{BrN}_2\text{O}+\text{H})^+$ 204.9976, found 204.9969.


Isoxazole **MGC2** (Method B):



[00128] Sodium metal (32.63 g, 1.42 mol, 2.03 equiv) was added portionwise

over 8 h to benzyl alcohol (1 L) at 23 °C. The resulting mixture was stirred vigorously for 24 h, then was transferred via large bore cannula to the neat isoxazole **MGC4** (143 g, 0.700 mol, 1.0 equiv) at 23 °C. The resulting light brown mixture was placed in an oil bath preheated to 120 °C and was stirred for 20 h at that temperature. Ethyl acetate (2 L) was added to the cooled reaction mixture and stirring was continued for 15 min. Aqueous hydrochloric acid (1.0 M, 2 L) was added and the aqueous phase was separated. The organic phase was further extracted with two 300-mL portions of 1.0 M aqueous hydrochloric acid. The aqueous phases were combined and the pH adjusted to 9 by slow addition of aqueous sodium hydroxide (6.0 M, approx. 350 mL). The resulting mixture was extracted with dichloromethane (3 × 500 mL). The organic extracts were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, yielding the isoxazole **MGC2** as a yellow oil (102 g, 63%). An analytical sample was prepared by flash column chromatography (3:7 ethyl acetate-hexanes, then 5:95 methanol in ethyl acetate), affording the isoxazole **MGC2** as a light yellow oil (spectroscopic data was identical to that obtained for material prepared by Method A).

Ketone **MGC5**:




[00129] A solution of *n*-butyllithium in hexanes (2.47 M, 16.0 mL, 39.5 mmol, 1.0 equiv) was added to a solution of the isoxazole **MGC2** (9.16 g, 39.5 mmol, 1.0 equiv) in tetrahydrofuran (150 mL) at -78 °C. The resulting rust-colored solution was stirred at -78 °C for 1 h whereupon a solution of the methyl ester **DJB1** (9.82 g, 23.7 mmol, 0.6 equiv) in tetrahydrofuran (6 mL) was added dropwise via cannula. The transfer was quantitated with two 1-mL portions of tetrahydrofuran. The resulting brown solution was stirred at -78 °C for 1 h, then an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 250 mL) was added. The biphasic mixture was allowed to warm to 23 °C, then was extracted with dichloromethane (2 × 300 mL). The organic

extracts were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a yellow oil. The product was purified by flash column chromatography (1:9 to 1:3 ethyl acetate-hexanes), affording the ketone **MGC5** as a light yellow solid (10.6 g, 73%).

[00130] R_f 0.59 (1:3 ethyl acetate-hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.44-7.35 (m, 5H, ArH), 5.90 (ddd, 1H, $J = 9.8, 5.9, 2.0$ Hz, =CHCHOSi), 5.82 (dd, 1H, $J = 9.8, 3.4$ Hz, =CHCHOCC), 5.31 (m, 2H, OCH_2Ar), 4.58 (d, 1H, $J = 4.2$ Hz, (O)CCCHOSi), 4.27 (m, 1H, =CHCHOSi), 3.94 (d, 1H, $J = 15.6$ Hz, CHH'N), 3.77 (d, 1H, $J = 15.6$ Hz, CHH'N), 3.17 (dd, 1H, $J = 3.4, 1.5$ Hz, HCOCC(O)), 2.35 (s, 6H, $\text{N}(\text{CH}_3)_2$), 0.89 (s, 9H, $\text{C}(\text{CH}_3)_3$), 0.83 (s, 9H, $\text{C}(\text{CH}_3)_3$), 0.06 (s, 3H, SiCH₃), 0.05 (s, 3H, SiCH₃), 0.04 (s, 3H, SiCH₃), -0.07 (s, 3H, SiCH₃); ^{13}C NMR (125 MHz, CDCl_3) δ 191.8, 176.3, 168.9, 136.5, 135.5, 128.8, 128.7, 125.0, 106.9, 72.4, 69.6, 67.8, 67.4, 55.3, 52.6, 45.9, 26.2, 26.0, 18.5, 18.3, -3.1, -3.8, -3.8, -5.1; FTIR (neat), cm^{-1} 2952 (s, CH), 1682 (s, C=O), 1594 (s), 1502 (s), 1456 (m), 1097 (s, C-O), 774 (s); HRMS (FAB) m/z calcd for $(\text{C}_{32}\text{H}_{50}\text{N}_2\text{O}_6\text{Si}_2+\text{Na})^+$ 637.3105, found 637.3097.

Ketones **MGC6** and **MGC7**:



[00131] Solid lithium trifluoromethanesulfonate (76.0 mg, 0.490 mmol, 0.05 equiv) was added to a solution of the ketone **MGC5** (6.02 g, 9.80 mmol, 1.0 equiv) in toluene (500 mL) at 23 °C. The resulting heterogeneous light yellow mixture was placed in an oil bath preheated to 65 °C and was stirred at that temperature for 3 h. The reaction mixture was cooled to 23 °C and was filtered. The solids were washed with toluene (50 mL) and the filtrate was concentrated, providing a yellow oil. The oil was covered with dichloromethane-trifluoroacetic acid (10:1, 165 mL) and the resulting mixture was stirred at 23 °C for 18 h. Aqueous sodium bicarbonate solution (300 mL) was added and extensive gas evolution was observed upon addition. The biphasic mixture was extracted with diethyl ether (4 × 300 mL) and the organic extracts were

combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a brown oil. The product was purified by flash column chromatography (1:9 to 1:5 ethyl acetate-hexanes), affording the ketone **MGC6** as a white foam (3.20 g, 62%) and the ketone **MGC7** as a viscous yellow oil (1.68 g, 28%).


Ketone MGC6:

[00132] R_f 0.52 (1:3 ethyl acetate-hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.45 (m, 2H, ArH), 7.36-7.30 (m, 3H, ArH), 5.96 (bs, 1H, =CH), 5.45 (bs, 1H, =CH), 5.32 (m, 2H, OCHH'Ar), 5.33 (bs, 1H, CHOSi), 4.15 (d, 1H, $J = 8.8$ Hz, CHOSi), 3.59 (d, 1H, $J = 3.9$ Hz, CHN(CH₃)₂), 3.34 (bs, 1H, C₃CH), 2.57 (bs, 1H, OH), 2.39 (s, 6H, N(CH₃)₂), 0.90 (s, 9H, C(CH₃)₃), 0.16 (s, 3H, SiCH₃), 0.11 (s, 3H, SiCH₃); ^{13}C NMR (100 MHz, C_6D_6) δ 189.2, 178.3, 168.6, 135.3, 128.5, 128.4, 128.3, 125.4, 106.4, 79.8, 72.3, 72.2, 67.1, 63.6, 42.9, 26.1, 18.5, -4.0, -4.8; FTIR (neat), cm^{-1} 3549 (bs, OH), 3455 (bs, OH), 2942 (s, CH), 1698 (s, C=O), 1617 (m), 1508 (s), 1032 (s, C-O), 906 (s); HRMS (ES) m/z calcd for $(\text{C}_{26}\text{H}_{36}\text{N}_2\text{O}_6\text{Si}+\text{H})^+$ 501.2421, found 501.2422.

Ketone MGC7:

[00133] R_f 0.64 (1:5 ethyl acetate-hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.50 (d, 2H, $J = 1.5$ Hz, ArH), 7.40-7.32 (m, 3H, ArH), 5.94 (dd, 1H, $J = 9.7, 6.4$ Hz, =CHCHCHOSi), 5.76 (d, 1H, $J = 9.7$ Hz, =CHCOH), 5.37 (d, 1H, $J = 12.2$ Hz, OCHH'Ph), 5.32 (d, 1H, $J = 12.2$ Hz, OCHH'Ph), 4.09 (d, 1H, $J = 2.9$ Hz, HOCCHOSi), 4.03 (s, 1H, OH), 3.88 (m, 1H, NCHCHCHOSi), 3.74 (d, 1H, $J = 3.9$ Hz, (CH₃)₂NCH), 2.46 (s, 6H, N(CH₃)₂), 0.91 (s, 9H, C(CH₃)₃), 0.87 (s, 9H, C(CH₃)₃), 0.06 (s, 3H, SiCH₃), 0.05 (s, 3H, SiCH₃), 0.04 (s, 3H, SiCH₃), 0.03 (s, 3H, SiCH₃); ^{13}C NMR (125 MHz, CDCl_3) δ 194.9, 173.9, 170.5, 135.8, 132.6, 128.8, 128.5, 128.3, 127.9, 106.2, 81.6, 74.8, 72.0, 71.7, 69.5, 44.6, 43.2, 26.1, 25.9, 18.7, 18.2, -3.6, -4.1, -4.3, -4.3; FTIR (neat), cm^{-1} 3461 (bs, OH), 2940 (s, CH), 1693 (s, C=O), 1663 (s), 1647 (m), 1503 (m), 1080 (s, C-O), 774 (s); HRMS (ES) m/z calcd for $(\text{C}_{32}\text{H}_{50}\text{N}_2\text{O}_6\text{Si}_2+\text{H})^+$ 615.3285, found 615.3282.


Alkene DRS3:



[00134] Diethyl azodicarboxylate (472 μL , 3.00 mmol, 3.0 equiv) was added to a solution of the ketone **MGC6** (500 mg, 1.00 mmol, 1.0 equiv) and triphenylphosphine (789 mg, 3.00 mmol, 3.0 equiv) in toluene (6.0 mL) at 0 $^{\circ}\text{C}$. The mixture was stirred at 0 $^{\circ}\text{C}$ for 90 min whereupon a solution of 2-nitrobenzenesulfonyl hydrazine (651 mg, 3.00 mmol, 3.0 equiv) in tetrahydrofuran (3 mL) was added dropwise via cannula. The resulting mixture was stirred at 0 $^{\circ}\text{C}$ for 10 min, then was allowed to warm to 23 $^{\circ}\text{C}$; stirring was continued at that temperature for 23 h. An aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 30 mL) was added and the resulting biphasic mixture was extracted with dichloromethane ($2 \times 50 \text{ mL}$). The organic extracts were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a yellow sludge. The product was purified by flash column chromatography (95:5 to 1:9 ethyl acetate-hexanes), affording the alkene **DRS3** as a white solid (356 mg, 74%).

[00135] R_f 0.65 (1:3 ethyl acetate-hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.46 (d, 2H, $J = 6.8 \text{ Hz}$, ArH), 7.39-7.34 (m, 3H, ArH), 5.81 (m, 1H, $=\text{CHCH}_2$), 5.55 (dd, 1H, $J = 10.3, 2.0 \text{ Hz}$, $=\text{CHCOSi}$), 5.39 (d, 1H, $J = 12.2 \text{ Hz}$, OCHH'Ph), 5.35 (d, 1H, $J = 12.2 \text{ Hz}$, OCHH'Ph), 4.15 (s, 1H, CHOSi), 4.04 (bs, 1H, OH), 3.76 (d, 1H, $J = 10.7 \text{ Hz}$, $\text{CHN}(\text{CH}_3)_2$), 2.58 (dd, 1H, $J = 10.7, 3.9 \text{ Hz}$, C_3CH), 2.47 (m, 8H, $\text{N}(\text{CH}_3)_2$, $=\text{CCH}_2$), 0.86 (s, 9H, $\text{C}(\text{CH}_3)_3$), -0.05 (s, 3H, SiCH_3), -0.13 (s, 3H, SiCH_3); ^{13}C NMR (125 MHz, CDCl_3) δ 191.5, 183.3, 167.9, 135.3, 128.8, 128.7, 128.5, 127.4, 106.8, 78.3, 72.6, 72.0, 67.9, 60.7, 43.0, 42.1, 26.0, 25.8, 23.6, 18.2, -4.6 , -5.0 ; FTIR (neat), cm^{-1} 3528 (w, OH), 2933 (s, CH), 1702 (s, $\text{C}=\text{O}$), 1600 (m), 1507 (s), 1092 (s, C-O), 1061 (s, C-O); HRMS (ES) m/z calcd for $(\text{C}_{26}\text{H}_{36}\text{N}_2\text{O}_5\text{Si}+\text{H})^+$ 485.2472, found 485.2457.

Diol **DRS4**:



[00136] Acetic acid (83.0 μL , 1.44 mmol, 2.0 equiv) and a solution of tetrabutylammonium fluoride in tetrahydrofuran (1.0 M, 1.44 mL, 1.44 mmol, 2.0 equiv) were added in sequence to a solution of the olefin **DRS3** (350 mg, 0.723 mmol, 1.0 equiv) in tetrahydrofuran (7.0 mL) at 0 °C. The resulting light gray solution was stirred at 0 °C for 30 min, then was allowed to warm to 23 °C; stirring was continued at that temperature for 5 h. The reaction mixture was concentrated, providing a brown oil. The product was purified by flash column chromatography (1:4 to 1:1 ethyl acetate-hexanes), affording the diol **DRS4** as a waxy white solid (202 mg, 76%).

[00137] R_f 0.38 (1:1 ethyl acetate-hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.51-7.48 (m, 2H, ArH), 7.42-7.36 (m, 3H, ArH), 5.84 (m, 1H, =CHCH₂), 5.55 (m, 1H, =CHCOH), 5.36 (m, 2H, OCH₂Ph), 4.15 (d, 1H, J = 8.1 Hz, CHOH), 3.69 (d, 1H, J = 8.8 Hz, CHN(CH₃)₂), 2.67 (m, 1H, C₃CH), 2.47 (s, 6H, N(CH₃)₂), 2.43 (dd, 1H, J = 7.7, 1.5 Hz, =CCHH'), 2.36 (m, 1H, =CCHH'); FTIR (neat), cm^{-1} 3492 (w, OH), 3272 (s, OH), 1703 (s, C=O), 1606 (m), 1509 (s), 1008 (s, C-O), 732 (s); HRMS (ES) m/z calcd for $(\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_5+\text{H})^+$ 371.1607, found 371.1601.

Cyclohexenone **DRS5**:




[00138] Solid *o*-iodoxybenzoic acid (558 mg, 1.99 mmol, 3.0 equiv) was added to a solution of the diol **DRS4** (246 mg, 0.665 mmol, 1.0 equiv) in dimethylsulfoxide (5.0 mL) at 23 °C. The resulting heterogeneous mixture was stirred for 5 min whereupon it became homogeneous. The brown reaction mixture was stirred at 23 °C for 36 h. Water (10 mL) was added resulting in the precipitation of excess *o*-iodoxybenzoic acid. The mixture was filtered and the filtrate was partitioned between saturated aqueous sodium bicarbonate solution-brine (1:1, 20 mL) and ethyl acetate-

hexanes (2:1, 45 mL). The organic phase was separated and the aqueous phase was further extracted with a 45-mL portion of ethyl acetate-hexanes (2:1). The organic extracts were combined and washed with aqueous sodium sulfite solution (2.0 M, 50 mL), brine (50 mL), and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing the cyclohexenone **DRS5** as a light brown foam (206 mg, 84%).

[00139] R_f 0.15 (1:3 ethyl acetate-hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.48 (d, 2H, $J = 7.3$ Hz, ArH), 7.40-7.34 (m, 3H, ArH), 6.98 (m, 1H, =CHCH₂), 6.12 (ddd, 1H, $J = 12.2, 2.0, 2.0$ Hz, =CHC(O)), 5.35 (m, 2H, OCH₂Ar), 4.75 (bs, 1H, OH), 3.85 (d, 1H, $J = 9.8$ Hz, CHN(CH₃)₂), 2.82 (m, 3H, C₃CH, =CCH₂), 2.48 (s, 6H, N(CH₃)₂); ^{13}C NMR (125 MHz, CDCl_3) δ 192.8, 188.2, 182.8, 167.6, 149.7, 135.0, 128.9, 128.8, 128.6, 128.3, 107.9, 79.7, 72.8, 60.4, 45.5, 42.4, 25.4; FTIR (neat), cm^{-1} 3447 (w, OH), 1707 (s, C=O), 1673 (s, C=O), 1600 (m), 1512 (s), 1018 (s, C-O), 730 (s); HRMS (ES) m/z calcd for $(\text{C}_{20}\text{H}_{20}\text{N}_2\text{O}_5+\text{H})^+$ 369.1450, found 369.1454.

Silyl-Cyclohexenone **DRS6**:




[00140] 2,6-Lutidine (75.0 μL , 0.640 mmol, 5.0 equiv) and *tert*-butyldimethylsilyl trifluoromethanesulfonate (88.0 μL , 0.380 mmol, 3.0 equiv) were added in sequence to a solution of the cyclohexenone **DRS5** (47.0 mg, 0.130 mmol, 1.0 equiv) in dichloromethane (3 mL) at 23 °C. The mixture was stirred at 23 °C for 3 h, then an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 15 mL) was added. The biphasic mixture was extracted with dichloromethane (2 $\times$ 20 mL) and the organic extracts were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, affording the silyl-cyclohexenone **DRS6** as a white crystalline solid (56.0 mg, 91%).

[00141] Mp 157-158 °C (dec); R_f 0.54 (1:3 ethyl acetate-hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.51 (d, 2H, $J = 1.5$ Hz, ArH), 7.50-7.34 (m, 3H, ArH), 6.94 (m, 1H,

=CHCH₂), 6.10 (ddd, 1H, J = 10.3, 1.5, 1.5 Hz, =CHC(O)), 5.36 (m, 2H, OCH₂Ar), 3.79 (d, 1H, J = 10.7 Hz, CHN(CH₃)₂), 2.83 (m, 2H, =CCH₂), 2.78 (m, 1H, C₃CH), 2.46 (s, 6H, N(CH₃)₂), 0.84 (s, 9H, C(CH₃)₃), 0.27 (s, 3H, SiCH₃), 0.06 (s, 3H, SiCH₃); ¹³C NMR (125 MHz, CDCl₃) δ 193.4, 187.9, 181.6, 167.7, 149.5, 135.2, 128.8, 128.8, 128.8, 128.6, 108.6, 83.5, 72.8, 59.8, 48.1, 42.2, 26.3, 25.8, 19.3, -2.2, -3.8; FTIR (neat), cm⁻¹ 2942 (s), 1719 (s, C=O), 1678 (s, C=O), 1602 (m), 1510 (s), 1053 (s, C-O), 733 (s); HRMS (ES) m/z calcd for (C₂₆H₃₄N₂O₅Si+H)⁺ 483.2315, found 483.2321.

Ketone **MGC9**:




[00142] A solution of methylmagnesium bromide in ether (3.15 M, 11.6 mL, 36.7 mmol, 1.07 equiv) was added to a solution of the aldehyde **MGC8** (synthesized in 2 steps from commercially available 3-benzyloxy benzyl alcohol as reported by: Hollinshead, S. P.; Nichols, J. B.; Wilson, J. W. *J. Org. Chem.* **1994**, 59, 6703.) (10.0 g, 34.3 mmol, 1.0 equiv) in tetrahydrofuran (90 mL) at -5 °C (NaCl/ice bath). The light brown solution was stirred at -5 °C for 60 min, then was partitioned between saturated aqueous ammonium chloride solution (400 mL) and ethyl acetate (400 mL). The organic phase was separated and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a light yellow oil (10.1 g, 95% crude). The product was used without further purification.

[00143] Sodium bromide (846 mg, 8.22 mmol, 0.25 equiv) and 2,2,6,6-tetramethyl-1-piperidinyloxy (51.0 mg, 0.329 mmol, 0.01 equiv) were added in sequence to a solution of the light yellow oil prepared above (10.1 g, 32.8 mmol, 1.0 equiv) in tetrahydrofuran (30 mL) at 0 °C. A freshly prepared solution of sodium bicarbonate (690 mg, 8.22 mmol, 0.25 equiv) in commercial Clorox bleach (90 mL) was cooled to 0 °C and was added in one portion to the mixture prepared above at 0 °C. The resulting bright yellow mixture was stirred vigorously at 0 °C for 1.5 h whereupon sodium sulfite (1.0 g) was added. The resulting mixture was stirred for 15 min at 23

°C, then was partitioned between water (400 mL) and ethyl acetate (400 mL). The organic phase was separated and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a light brown oil. The product was crystallized from ethanol, furnishing the ketone **MGC9** as a white solid (8.08 g, 80% over 2 steps).

[00144] R_f 0.80 (3:7 ethyl acetate-hexanes); ^1H NMR (400 MHz, CDCl_3) δ 7.26-7.48 (m, 6H, ArH), 6.98 (m, 2H, ArH), 5.19 (s, 2H, OCH_2Ph), 2.62 (s, 3H, C(=O)CH_3); ^{13}C NMR (100 MHz, CDCl_3) δ 202.4, 155.5, 144.4, 136.3, 128.9, 128.7, 128.3, 127.2, 120.3, 115.2, 109.1, 71.3, 30.9; FTIR (neat), cm^{-1} 3065 (w), 3032 (w), 2918 (m), 1701 (s, C=O), 1565 (m), 1426 (m), 1300 (s), 1271 (s), 1028 (m); HRMS (ES) m/z calcd for $(\text{C}_{15}\text{H}_{13}\text{O}_2\text{Br}+\text{H})^+$ 304.0099, found 304.0105.


Epoxide **MGC10**:



[00145] Dimethylsulfoxide (90 mL) was added dropwise via syringe to a mixture of solid trimethylsulfoxonium iodide (694 mg, 3.15 mmol, 1.3 equiv) and solid sodium hydride (60% in oil, 126 mg, 3.15 mmol, 1.3 equiv, washed with three 2-mL portions of *n*-hexane) at 23 °C. Vigorous gas evolution was observed upon addition. The resulting cloudy gray mixture was stirred at 23 °C for 40 min, then a solution of the ketone **MGC9** (8.08 g, 26.5 mmol, 1.0 equiv) in dimethylsulfoxide (30 mL) was added dropwise via cannula. The transfer was quantitated with a 2-mL portion of dimethylsulfoxide. The resulting orange mixture was stirred at 23 °C for 35 h, then was partitioned between brine (1 L) and ether (500 mL). The organic phase was separated and the aqueous phase was further extracted with one 500-mL portion of ether. The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a yellow oil. The product was purified by flash column chromatography (5:95 ethyl acetate-hexanes), affording the epoxide **MGC10** as a clear oil (7.94 g, 94%).

[00146] R_f 0.90 (3:7 ethyl acetate-hexanes); ^1H NMR (300 MHz, CDCl_3) δ 7.20-7.52 (m, 6H, ArH), 7.10 (dd, 1H, $J = 7.5, 1.2$ Hz, *o*-ArH), 6.88 (dd, 1H, $J = 8.1, 1.2$ Hz, *o*-ArH), 5.16 (s, 2H, OCH_2Ph), 3.03 (d, 1H, $J = 4.8$ Hz, $\text{CHH}'\text{OCCH}_3$), 2.87 (d, 1H, $J = 4.8$ Hz, $\text{CHH}'\text{OCCH}_3$), 1.67 (s, 3H, COCH_3); ^{13}C NMR (100 MHz, CDCl_3) δ 155.0, 143.4, 136.7, 128.8, 128.4, 128.2, 127.2, 121.2, 112.8, 112.3, 71.2, 59.7, 55.9, 22.9; FTIR (neat), cm^{-1} 3034 (w), 2981 (w), 2925 (w), 1595 (w), 1567 (s), 1469 (s), 1423 (s), 1354 (s), 1300 (s), 1266 (s), 1028 (s); HRMS (ES) m/z calcd for $(\text{C}_{16}\text{H}_{15}\text{O}_2\text{Br}+\text{H})^+$ 318.0255, found 318.0254.

Benzocyclobutenol **MGC11**:



[00147] A solution of *n*-butyllithium in hexanes (1.60 M, 8.25 mL, 13.6 mmol, 1.4 equiv) was added dropwise via syringe down the side of a reaction vessel containing a solution of the epoxide **MGC10** (3.11 g, 9.74 mmol, 1.0 equiv) in tetrahydrofuran (90 mL) at -78°C . The resulting yellow solution was stirred at -78°C for 20 min whereupon a suspension of magnesium bromide (3.95 g, 21.4 mmol, 2.2 equiv) in tetrahydrofuran (25 mL) was added dropwise via cannula. The transfer was quantitated with two 2.5-mL portions of tetrahydrofuran. The resulting cloudy mixture was stirred at -78°C for 60 min, then the cooling bath was removed and the reaction mixture was allowed to warm to 23°C . The mixture became clear upon warming and was stirred at 23°C for 1 h. The reaction mixture was poured into aqueous Rochelle's salt solution (10% wt/wt, 1 L) and the resulting mixture was extracted with ethyl acetate (2×400 mL). The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing an off-white solid. The product was purified by flash column chromatography (1:9 to 2:9 ethyl acetate-hexanes), affording the *trans*-benzocyclobutenol **MGC11** as a white solid (1.57 g, 67%).

[00148] R_f 0.50 (3:7 ethyl acetate-hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.44

(br d, 2H, $J = 7.5$ Hz, ArH), 7.38 (br t, 2H, $J = 7.5$ Hz, ArH), 7.22-7.34 (m, 2H, ArH), 6.82 (d, 1H, $J = 8.5$ Hz, *o*-ArH), 6.75 (d, 1H, $J = 7.5$ Hz, *o*-ArH), 5.35 (d, 1H, $J = 12.0$ Hz, OCHH'Ph), 5.25 (d, 1H, $J = 12.0$ Hz, OCHH'Ph), 4.71 (br d, 1H, $J = 5.5$ Hz, CHOH), 3.31 (br q, 1H, $J = 7.0$ Hz, CHCH₃), 2.21 (br d, 1H, $J = 7.0$ Hz, OH), 1.38 (d, 3H, $J = 7.0$ Hz, CHCH₃); ¹³C NMR (100 MHz, CDCl₃) δ 154.0, 148.9, 137.4, 131.5, 128.5, 128.4, 127.8, 127.3, 115.2, 114.6, 77.6, 71.2, 50.6, 16.5; FTIR (neat), cm⁻¹ 3249 (m, OH), 2958 (w), 1602 (m), 1580 (s), 1453 (s), 1261 (s), 1039 (s); HRMS (ES) m/z calcd for (C₁₆H₁₆O₂+H)⁺ 240.1150, found 240.1154.


Benzocyclobutenol MGC12:



[00149] Triethylamine (336 μ L, 2.41 mmol, 1.4 equiv) and triethylsilyl trifluoromethanesulfonate (468 μ L, 2.07 mmol, 1.2 equiv) were added in sequence to a solution of the benzocyclobutenol **MGC11** (500 mg, 1.72 mmol, 1.0 equiv) in dichloromethane (10 mL) at 23 °C. The light yellow solution was stirred at 23 °C for 15 min, then was partitioned between water (30 mL) and dichloromethane (30 mL). The organic phase was separated and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a yellow oil. The product was purified by flash column chromatography (5:95 ethyl acetate-hexanes), affording the benzocyclobutenol **MGC12** (609 mg, 99%) as a clear oil.

[00150] R_f 0.85 (1:4 ethyl acetate-hexanes); ¹H NMR (400 MHz, CDCl₃) δ 7.48-7.32 (m, 5H, ArH), 7.24 (m, 2H, ArH), 6.82 (d, 1H, $J = 8.4$ Hz, *o*-ArH), 6.74 (d, 1H, $J = 7.2$ Hz, *o*-ArH), 5.37 (d, 1H, $J = 11.2$ Hz, CHH'Ph), 5.20 (d, 1H, $J = 11.2$ Hz, CHH'Ph), 4.87 (d, 1H, $J = 1.6$ Hz, CHOTES), 3.45 (dq, 1H, $J = 7.2, 1.6$ Hz, CHCH₃), 1.42 (d, 3H, $J = 7.2$ Hz, CHCH₃), 0.98 (t, 9H, $J = 7.6$ Hz, TES), 0.56 (q, 6H, $J = 7.6$ Hz, TES); ¹³C NMR (100 MHz, CDCl₃) δ 154.2, 148.8, 137.6, 131.3, 129.0, 128.7, 128.1, 127.8, 115.1, 114.7, 71.7, 49.9, 16.9, 7.1, 5.2, 5.1; FTIR (neat), cm⁻¹ 2952 (w), 2923 (w), 2854 (w), 1606 (w), 1469 (w), 1371 (m), 1265 (s), 1086 (s), 1057 (s), 1048 (s); HRMS (ES) m/z calcd for (C₂₂H₃₀O₂Si+H)⁺ 354.2015, found 354.2006.

Vinyl Sulfide MGC13:



[00151] Solid pyridinium hydrobromide perbromide (293 mg, 0.917 mmol, 2.5 equiv) was added to a solution of the cyclohexenone **DRS5** (135 mg, 0.367 mmol, 1.0 equiv) in dichloromethane (4 mL) at 23 °C. The brown solution was stirred vigorously at 23 °C for 17 h whereupon sodium sulfite (150 mg, 1.19 mmol, 3.25 equiv) was added. The resulting mixture was partitioned between an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 30 mL) and dichloromethane (30 mL). The organic phase was separated and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a light brown foamy solid. The product was used immediately without further purification.


R_f 0.45 (2:3 ethyl acetate-hexanes); ^1H NMR (500 MHz, C_6D_6) δ 7.24 (d, 2H, $J = 7.0$ Hz, *o*-ArH), 7.02 (t, 2H, $J = 7.0$ Hz, *m*-ArH), 6.99 (d, 1H, $J = 7.0$ Hz, *p*-ArH), 6.42 (ddd, 1H, $J = 6.0, 3.5, 2.0$ Hz, CH=CB_r), 5.12 (d, 1H, $J = 12.5$ Hz, CHH'Ph), 5.03 (d, 1H, $J = 12.5$ Hz, CHH'Ph), 4.00 (br s, 1H, OH), 3.25 (d, 1H, $J = 11.0$ Hz, CHN(CH₃)₂), 2.28-2.22 (m, 2H, CH₂CH, CH₂CH), 2.16 (dd, 1H, $J = 18.0, 6.0$ Hz, CH₂CH), 1.83 (s, 6H, N(CH₃)₂); FTIR (neat), cm^{-1} 3397 (m, OH), 3063 (m), 2943 (m), 1714 (s, C=O), 1606 (s), 1514 (s), 1477 (s), 1371 (m), 1022 (m); HRMS (ES) m/z calcd for $(\text{C}_{20}\text{H}_{19}\text{O}_5\text{BrN}_2)^+$ 447.0555, found 447.0545.

[00152] Benzenethiol (39.0 μL , 0.378 mmol, 1.03 equiv) and 1,8-diazabicyclo[5,4,0]undec-7-ene (56.0 μL , 0.378 mmol, 1.03 equiv) were added in sequence to a solution of the product prepared above (164 mg, 0.367 mmol, 1.0 equiv) in *N,N*-dimethylformamide (4 mL) at 0 °C. The resulting dark brown mixture was stirred vigorously at 0 °C for 25 min, then was partitioned between ethyl acetate-hexanes (1:1, 30 mL) and an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 30 mL). The organic phase was separated and the aqueous phase was further extracted with two 15-mL portions of ethyl acetate-hexanes (1:1). The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was

filtered and the filtrate was concentrated, providing a brown oil. The product was purified by flash column chromatography (15:85 to 1:4 ethyl acetate-hexanes), furnishing the vinyl sulfide **MGC13** as a white foam (116 mg, 66% over two steps).

[00153] R_f 0.47 (2:3 ethyl acetate-hexanes); ^1H NMR (500 MHz, C_6D_6) δ 7.34 (dd, 2H, $J = 7.0, 1.0$ Hz, *o*-ArH), 7.23 (d, 2H, $J = 6.5$ Hz, *o*-ArH), 6.85-7.04 (m, 6H, ArH), 6.27 (ddd, 1H, $J = 6.0, 3.0, 1.0$ Hz, CH=CSPH), 5.11 (d, 1H, $J = 12.0$ Hz, OCHH'Ph), 5.02 (d, 1H, $J = 12.0$ Hz, OCHH'Ph), 4.62 (br s, 1H, OH), 3.42 (d, 1H, $J = 10.5$ Hz, CHN(CH₃)₂), 2.44 (ddd, 1H, $J = 20.0, 5.5, 3.0$ Hz, CH₂CH), 2.27-2.34 (m, 2H, CH₂CH, CH₂CH), 1.87 (s, 6H, N(CH₃)₂); ^{13}C NMR (100 MHz, CDCl_3) δ 188.9, 187.4, 182.5, 167.6, 145.4, 135.3, 135.2, 132.8, 132.6, 129.5, 128.6, 128.4, 128.3, 128.0, 127.8, 108.1, 80.3, 72.5, 59.8, 45.7, 41.4, 25.9; FTIR (neat), cm^{-1} 3445 (w, OH), 3056 (w), 2943 (m), 2800 (w), 1711 (s, C=O), 1682 (s), 1600 (m), 1507 (s), 1471 (s), 1451 (m), 1333 (m), 1020 (m); HRMS (ES) m/z calcd for $(\text{C}_{26}\text{H}_{24}\text{O}_5\text{N}_2\text{S}+\text{H})^+$ 477.1484, found 447.1465.

Diel-Alder Addition Product **MGC14** and Lactone **MGC15**:



[00154] A reaction vessel containing a mixture of the vinylsulfide **MGC13** (131 mg, 0.275 mmol, 1.0 equiv) and the benzocyclobutenol **MGC12** (750 mg, 2.11 mmol, 7.7 equiv) was placed in an oil bath preheated to 85 °C. The light yellow solution was stirred at 85 °C for 48 h, then was allowed to cool to 23 °C. The cooled mixture was purified by flash column chromatography (1:19 to 1:4 ethyl acetate-hexanes), affording the Diels-Alder addition product **MGC14** as an off-white foamy solid (145 mg, 64%), the lactone **MGC15** as a clear oil (20.0 mg, 9%), and the recovered benzocyclobutenol **MGC12** as a clear oil (650 mg).

Diels-Alder Addition Product **MGC14**:

[00155] mp 178-179 °C; R_f 0.55 (2:3 ethyl acetate-hexanes); ^1H NMR (600 MHz, C_6D_6) δ 7.27 (d, 2H, $J = 7.2$ Hz, *o*-ArH), 7.06-7.22 (m, 8H, ArH), 6.92-6.96 (m,


3H, ArH), 6.85 (d, 1H, $J = 7.2$ Hz, ArH), 6.70-6.75 (m, 3H, ArH), 6.55 (d, 1H, $J = 8.4$ Hz, *o*-ArH), 5.75 (s, 1H, CHOTES), 5.29 (br s, 1H, OH), 5.16 (d, 1H, $J = 12.0$ Hz, OCHH'Ph), 5.10 (d, 1H, $J = 12.0$ Hz, OCHH'Ph), 4.66 (d, 1H, $J = 10.8$ Hz, OCHH'Ph'), 4.63 (d, 1H, $J = 10.8$ Hz, OCHH'Ph'), 4.36 (d, 1H, $J = 6.6$ Hz, CHN(CH₃)₂), 3.02 (dq, 1H, $J = 7.8, 6.0$ Hz, CH₃CH), 2.77 (ddd, 1H, $J = 6.6, 6.0, 4.2$ Hz, CHCHN(CH₃)₂), 2.41-2.52 (m, 2H, CHCHH'CH, CH₃CHCHCH₂), 2.08 (s, 6H, N(CH₃)₂), 1.83 (ddd, 1H, $J = 13.2, 4.2, 4.2$ Hz, CHCHH'CH), 1.34 (d, 3H, $J = 7.8$ Hz, CH₃CH), 0.70 (t, 9H, $J = 7.8$ Hz, Si(CH₂CH₃)₃), 0.48 (d, 6H, $J = 7.8$ Hz, Si(CH₂CH₃)₃); ¹³C NMR (100 MHz, CDCl₃) δ 196.3, 186.1, 181.4, 168.3, 156.3, 143.9, 137.6, 136.6, 135.4, 130.6, 129.8, 129.3, 128.6, 128.5, 128.4, 128.2, 128.0, 127.8, 125.4, 121.1, 109.3, 108.4, 80.6, 72.4, 70.2, 66.0, 62.5, 61.7, 43.2, 42.0, 38.1, 37.2, 27.4, 20.5, 6.9, 4.9; FTIR (neat), cm⁻¹ 3490 (w, OH), 3063 (w), 3023 (w), 2951 (m), 2871 (m), 1715 (s, C=O), 1602 (m), 1589 (m), 1513 (s), 1457 (s), 1366 (m), 1260 (s), 1065 (s), 1012 (s); HRMS (FAB) m/z calcd for (C₄₈H₅₄O₇N₂SSi+Na)⁺ 853.3318, found 853.3314.

Lactone MGC15:

[00156] R_f 0.55 (3:7 ethyl acetate-hexanes); ¹H NMR (600 MHz, C₆D₆) δ 7.34 (d, 2H, $J = 7.2$ Hz, *o*-ArH), 7.02-7.18 (m, 11H, ArH), 6.72-6.84 (m, 4H, ArH), 6.54 (d, 1H, $J = 7.8$ Hz, *o*-ArH), 5.73 (s, 1H, CHOTES), 5.49 (d, 1H, $J = 6.6$ Hz, (C=O)OCHC=O), 5.20 (s, 2H, OCH₂Ph), 4.60 (d, 1H, $J = 11.4$ Hz, OCHH'Ph'), 4.57 (d, 1H, $J = 11.4$ Hz, OCHH'Ph'), 3.49 (d, 1H, $J = 11.4$ Hz, CHN(CH₃)₂), 3.23 (dq, 1H, $J = 9.0, 7.2$ Hz, CH₃CH), 2.49 (m, 1H, CH₃CHCHCHH'), 2.30-2.40 (m, 2H, CHCHN(CH₃)₂, CH₃CHCHCH₂), 2.16 (dd, 1H, $J = 12.0, 0.6$ Hz, CH₃CHCHCHH'), 1.96 (s, 6H, N(CH₃)₂), 1.33 (d, 3H, $J = 7.2$ Hz, CH₃CH), 0.73 (t, 9H, $J = 7.8$ Hz, Si(CH₂CH₃)₃), 0.46-0.62 (m, 6H, Si(CH₂CH₃)₃); ¹³C NMR (100 MHz, CDCl₃) δ 196.4, 176.0, 170.0, 157.9, 156.0, 144.0, 136.6, 136.5, 135.6, 129.8, 129.7, 129.4, 128.9, 128.6, 128.4, 128.3, 128.2, 128.1, 127.8, 125.1, 121.2, 108.8, 101.9, 75.9, 72.1, 70.1, 64.7, 64.6, 62.9, 41.4, 36.7, 35.6, 27.7, 21.7, 6.9, 4.9; FTIR (neat), cm⁻¹ 3062 (w), 3033 (w), 2950 (m), 2874 (m), 1731 (s, C=O), 1599 (m), 1590 (m), 1514 (s), 1453 (s), 1365 (m), 1259 (s), 1120 (s), 1059 (s), 1010 (s); HRMS (ES) m/z calcd for

$$(\text{C}_{48}\text{H}_{54}\text{O}_7\text{N}_2\text{SSi}+\text{H})^+ \text{ 831.3499, found 831.3509.}$$


Alcohol MGC16:



[00157] Triethylamine trihydrofluoride (200 μ L, 1.23 mmol, 8.5 equiv) was added to a solution of the Diels-Alder addition product **MGC14** (120 mg, 0.144 mmol, 1.0 equiv) in tetrahydrofuran (6 mL) at 23 $^{\circ}$ C. The mixture was stirred vigorously at 23 $^{\circ}$ C for 12 h, then was partitioned between an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 30 mL) and ethyl acetate (30 mL). The organic phase was separated and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a light brown solid. The product was purified by flash column chromatography (1:4 to 1:1 ethyl acetate-hexanes), affording the alcohol **MGC16** as a colorless oil (78.3 mg, 76%).

[00158] R_f 0.20 (2:3 ethyl acetate-hexanes); ^1H NMR (600 MHz, C_6D_6) δ 7.69 (dd, 2H, $J = 7.2, 0.6$ Hz, *o*-ArH), 7.24 (d, 2H, $J = 7.2$ Hz, ArH), 6.92-7.06 (m, 12H, ArH), 6.76 (d, 1H, $J = 7.8$ Hz, ArH), 6.47 (d, 1H, $J = 8.4$ Hz, *o*-ArH), 5.44 (br s, 1H, CHOH), 5.18 (d, 1H, $J = 12.0$ Hz, OCHH'Ph), 5.16 (d, 1H, $J = 12.0$ Hz, OCHH'Ph), 4.57 (d, 1H, $J = 12.6$ Hz, OCHH'Ph'), 4.52 (d, 1H, $J = 12.6$ Hz, OCHH'Ph'), 3.44 (dq, 1H, $J = 6.6, 5.4$ Hz, CH_3CH), 2.98 (d, 1H, $J = 3.0$ Hz, $\text{CHN}(\text{CH}_3)_2$), 2.90 (m, 1H, $\text{CHCHN}(\text{CH}_3)_2$), 2.76 (br s, 1H, OH), 2.32 (m, 1H, $\text{CH}_3\text{CHCHCH}_2$), 1.94 (m, 1H, $\text{CH}_3\text{CHCHCH}_2$), 1.79 (s, 6H, $\text{N}(\text{CH}_3)_2$), 1.07 (m, 1H, $\text{CH}_3\text{CHCHCH}_2$), 0.84 (d, 3H, $J = 6.6$ Hz, CH_3CH); ^{13}C NMR (100 MHz, CDCl_3) δ 202.5, 185.6, 179.2, 168.9, 156.9, 139.4, 139.1, 137.1, 136.5, 135.3, 130.5, 129.6, 128.8, 128.7, 128.6, 128.5, 128.4, 128.3, 127.8, 126.9, 124.7, 119.3, 110.0, 106.8, 82.3, 72.5, 69.9, 66.4, 64.2, 59.3, 43.0, 39.1, 37.8, 32.6, 25.3, 16.8; FTIR (neat), cm^{-1} 3435 (w, OH), 3066 (w), 2964 (w), 2933 (w), 2871 (w), 1738 (s, C=O), 1698 (s, C=O), 1614 (m), 1583 (m), 1513 (s), 1471 (s), 1453 (s), 1369 (m), 1263 (m), 1035 (m), 1014 (m); HRMS (ES) m/z calcd for $(\text{C}_{42}\text{H}_{40}\text{O}_7\text{N}_2\text{S}+\text{H})^+$ 717.2634, found 717.2631.

Triketone MGC17:




[00159] Solid *o*-iodoxybenzoic acid (459 mg, 1.64 mmol, 15.0 equiv) was added in one portion to a solution of the alcohol **MGC16** (78.3 mg, 0.109 mmol, 1.0 equiv) in dimethylsulfoxide (3.0 mL) at 23 °C. The resulting heterogeneous mixture was stirred for 5 min whereupon it became homogeneous. The reaction vessel was protected from light and was placed in an oil bath preheated to 35 °C. The brown solution was stirred vigorously at 35 °C for 18 h, then was partitioned between saturated aqueous sodium bicarbonate solution-brine-water (2:1:1, 75 mL) and ethyl acetate-ether (1:2, 35 mL). The organic phase was separated and the aqueous phase was further extracted with two 25-mL portions ethyl acetate-ether (1:2). The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a yellow oil. The product was purified by flash column chromatography (1:2 ethyl acetate-hexanes), affording the ketone **MGC17** as a yellow oil (61.7 mg, 79%).

[00160] R_f 0.45 (2:3 ethyl acetate-hexanes); ^1H NMR (600 MHz, C_6D_6) δ 7.57 (d, 2H, $J = 7.2$ Hz, *o*-ArH), 7.40 (d, 2H, $J = 7.2$ Hz, ArH), 7.18-7.23 (m, 3H, ArH), 6.94-7.06 (m, 6H, ArH), 6.76-6.84 (m, 3H, ArH), 6.59 (d, 1H, $J = 7.8$ Hz, ArH), 6.53 (d, 1H, $J = 8.4$ Hz, *o*-ArH), 5.09 (d, 1H, $J = 12.6$ Hz, OCHH'Ph), 4.96 (d, 1H, $J = 12.6$ Hz, OCHH'Ph), 4.77 (d, 1H, $J = 12.0$ Hz, OCHH'Ph'), 4.72 (d, 1H, $J = 12.0$ Hz, OCHH'Ph'), 4.48 (br s, 1H, OH), 4.06 (dq, 1H, $J = 7.2, 3.0$ Hz, CH₃CH), 3.15 (d, 1H, $J = 12.0$ Hz, CHN(CH₃)₂), 2.20 (ddd, 1H, $J = 12.6, 5.4, 3.0$ Hz, CH₃CHCHCH₂), 2.13 (ddd, 1H, $J = 12.0, 3.0, 0.6$ Hz, CHCHN(CH₃)₂), 1.81-1.88 (m, 7H, N(CH₃)₂, CH₃CHCHCHH'), 1.78 (ddd, 1H, $J = 13.8, 5.4, 0.6$ Hz, CH₃CHCHCHH'), 1.01 (d, 3H, $J = 7.2$ Hz, CH₃CH); ^{13}C NMR (100 MHz, CDCl_3) δ 200.3, 187.5, 183.1, 167.8, 160.6, 146.4, 138.2, 137.1, 135.3, 134.3, 131.7, 129.6, 128.9, 128.6, 128.5, 128.4, 128.3, 127.7, 126.7, 121.3, 118.0, 112.8, 108.3, 82.9, 77.5, 72.4, 70.3, 58.1, 47.0, 44.1, 32.4,

18.7, 18.0, 16.3; FTIR (neat), cm^{-1} 3457 (w, OH), 3063 (w), 2939 (w), 2878 (w), 2795 (w), 1727 (s, C=O), 1704 (s, C=O), 1667 (m, C=O), 1593 (s), 1513 (s), 1471 (s), 1453 (s), 1371 (m), 1276 (m), 1044 (m); HRMS (ES) m/z calcd for $(\text{C}_{42}\text{H}_{38}\text{O}_7\text{N}_2\text{S}+\text{H})^+$ 715.2478, found 715.2483.

Peroxide MGC18:



[00161] A solution of trifluoroacetic acid in dichloromethane (1.0 M, 0.189 mL, 0.189 mmol, 2.5 equiv) and a solution of *m*-chloroperoxybenzoic acid in dichloromethane (0.5 M, 0.228 mL, 0.114 mmol, 1.5 equiv) were added in sequence to a solution of the sulfide **MGC17** (54.2 mg, 0.0758 mmol, 1.0 equiv) in dichloromethane (4.0 mL) at $-78\text{ }^\circ\text{C}$. The resulting cloudy mixture was stirred at $-78\text{ }^\circ\text{C}$ for 10 min, then the $-78\text{ }^\circ\text{C}$ bath was replaced with a $0\text{ }^\circ\text{C}$ bath. The mixture became homogeneous upon warming. The solution was stirred at $0\text{ }^\circ\text{C}$ for 30 min, then was partitioned between an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 10 mL) and dichloromethane (10 mL). The organic phase was separated and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a bright yellow oil. The oil was taken up in toluene (1 mL) and dried by azeotropic distillation at $40\text{ }^\circ\text{C}$ under high vacuum. The resulting yellow oil was dissolved in chloroform (2 mL) and the reaction vessel was exposed to atmospheric oxygen. The mixture was allowed to stand until oxidation was complete as evidenced by ^1H NMR spectroscopy. The mixture was filtered and the filtrate was concentrated, providing the peroxide **MGC18** as a brown oil. The product was reduced immediately to tetracycline.

[00162] The peroxide **MGC18** can also be prepared by following the procedure reported by Wasserman (*J. Am. Chem. Soc.* **1986**, *108*, 4237-4238.):


[00163] A solution of trifluoroacetic acid in dichloromethane (1.0 M, 24.5 μL ,

0.0245 mmol, 2.5 equiv) and a solution of *m*-chloroperoxybenzoic acid in dichloromethane (0.5 M, 29.4 μ L, 0.0147 mmol, 1.5 equiv) were added in sequence to a solution of the sulfide **MGC17** (7.00 mg, 0.00979 mmol, 1.0 equiv) in dichloromethane (0.5 mL) at -78 $^{\circ}$ C. The resulting cloudy mixture was stirred at -78 $^{\circ}$ C for 10 min, then the -78 $^{\circ}$ C bath was replaced with a 0 $^{\circ}$ C bath. The mixture became homogeneous upon warming. The solution was stirred at 0 $^{\circ}$ C for 30 min, then was partitioned between an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 8 mL) and dichloromethane (8 mL). The organic phase was separated and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a bright yellow oil. The oil was taken up in toluene (1 mL) and dried by azeotropic distillation at 40 $^{\circ}$ C under high vacuum. The resulting yellow oil was dissolved in chloroform (2 mL) and *meso*-tetraphenylporphine (0.6 mg, 0.979 μ mol, 0.10 equiv) was added in one portion. Oxygen gas was bubbled through the resulting mixture under UV irradiation (200 W Hg lamp) for 10 min. The mixture was concentrated to 0.5 mL and was diluted with methanol (5 mL) resulting in precipitation of *meso*-tetraphenylporphine. The resulting mixture was filtered and the filtrate was concentrated, providing the peroxide **MGC18** a light yellow solid.

[00164] R_f 0.10 (2:3 ethyl acetate-hexanes); ^1H NMR (500 MHz, C_6D_6 , keto tautomer reported) δ 8.95 (br s, 1H, OOH), 7.48 (d, 2H, $J = 7.0$ Hz, *o*-ArH), 7.28 (d, 2H, $J = 7.0$ Hz, ArH), 6.96-7.16 (m, 8H, ArH), 6.53 (d, 1H, $J = 8.0$ Hz, ArH), 5.14 (d, 1H, $J = 12.0$ Hz, OCHH'Ph), 5.03 (d, 1H, $J = 12.0$ Hz, OCHH'Ph), 4.83 (d, 1H, $J = 12.5$ Hz, OCHH'Ph'), 4.74 (d, 1H, $J = 12.5$ Hz, OCHH'Ph'), 4.60 (br s, 1H, OH), 3.54 (d, 1H, $J = 11.0$ Hz, CHCHN(CH₃)₂), 3.12 (dd, 1H, $J = 18.0, 0.5$ Hz, CHCHH'CH), 2.82 (dd, 1H, $J = 18.0, 4.5$ Hz, CHCHH'CH), 2.44 (ddd, 1H, $J = 11.0, 4.5, 0.5$ Hz, CHCHN(CH₃)₂), 1.86 (s, 6H, N(CH₃)₂), 1.01 (s, 3H, CH₃); ^{13}C NMR (100 MHz, C_6D_6 , enol and keto tautomers reported) δ 194.4, 188.6, 187.8, 187.2, 182.3, 178.4, 171.9, 167.7, 165.6, 159.5, 158.4, 147.9, 145.9, 137.0, 136.8, 136.6, 135.4, 135.3, 134.5, 134.3, 133.5, 133.4, 133.1, 132.9, 131.0, 130.8, 130.2, 129.9, 129.7, 129.2, 128.9, 126.8, 126.7, 124.5, 124.3, 122.2, 118.6, 116.9, 116.5, 113.4, 113.3, 113.2, 108.2, 107.9, 103.3, 83.7, 81.7, 80.1, 79.1, 72.4, 70.7, 70.4, 63.9, 59.1, 46.1, 44.9, 41.4, 40.8, 31.5, 30.0, 26.8, 22.9, 21.4; FTIR (neat film), cm^{-1} 3035 (w), 2946 (w), 1907 (w), 1731

(s, C=O), 1410 (s), 1379 (m), 1235 (m), 1170 (m), 1136 (m); HRMS (ES) m/z calcd for $(C_{36}H_{32}O_9N_2+H)^+$ 637.2186, found 637.2190.

(–)-Tetracycline (MGC29):



[00165] Pd black (14.1 mg, 0.133 mmol, 1.75 equiv) was added in one portion to a solution of the peroxide **MGC18** (48.2 mg, 0.0758 mmol, 1.0 equiv) in dioxane (3 mL) at 23 °C. An atmosphere of hydrogen was introduced by briefly evacuating the flask, then flushing with pure hydrogen (1 atm). The Pd catalyst was initially present as a fine dispersion, but aggregated into clumps within 5 min. The yellow heterogeneous mixture was stirred at 23 °C for 2 h, then was filtered through a plug of cotton. The filtrate was concentrated, affording a yellow solid. The product was purified by preparatory HPLC on a Phenomenex Polymerx DVB column (10 μ M, 250 x 10 mm, flow rate 4.0 mL/min, Solvent A: methanol-0.005 N aq. HCl (1:4), Solvent B: acetonitrile) using an injection volume of solvent A (500 μ L) containing oxalic acid (10 mg) and an isocratic elution of 5% B for 2 min, then a gradient elution of 5-50% B for 20 min. The peak eluting at 11-16 min was collected and concentrated, affording (–)-tetracycline hydrochloride as a yellow powder (16.0 mg, 44% from triketone **MGC17**), which was identical with natural (–)-tetracycline hydrochloride in all respects.

[00166] ^1H NMR (600 MHz, CD_3OD , hydrochloride) δ 7.50 (dd, 1H, J = 8.4, 7.8 Hz, ArH), 7.13 (d, 1H, J = 7.8 Hz, ArH), 6.91 (d, 1H, J = 8.4 Hz, ArH), 4.03 (s, 1H, $\text{CHN}(\text{CH}_3)_2$), 2.96-3.04 (m, 7H, $\text{HOC}(\text{CH}_3)\text{CH}$, $\text{N}(\text{CH}_3)_2$), 2.91 (br dd, 1H, J = 12.6, 2.4 Hz, $(\text{CH}_3)_2\text{NCHCH}$), 2.18 (ddd, 1H, J = 12.6, 6.0, 2.4 Hz, $\text{CHCHH}'\text{CH}$), 1.90 (ddd, 1H, J = 12.6, 12.6, 12.0 Hz, $\text{CHCHH}'\text{CH}$), 1.60 (s, 3H, CH_3); ^{13}C NMR (100 MHz, CD_3OD) δ 195.4, 174.5, 163.8, 148.3, 137.8, 118.7, 116.4, 116.0, 107.5, 96.5, 74.7, 71.2, 70.1, 43.5, 43.0, 35.9, 27.8, 22.9; UV max (0.1 N HCl), nm 217, 269, 356; $[\alpha]_D = -251^\circ$ (c = 0.12 in 0.1 M HCl); lit. (*The Merck Index: An Encyclopedia of Chemicals*,

127.6, 107.3, 80.8, 76.9, 72.4, 64.8, 54.6, 46.3, 41.5, 26.2, 18.4, -2.9, -4.2; FTIR (neat), cm^{-1} 3499 (m, OH), 2930 (m), 2856 (m), 2799 (w), 1704 (s, C=O), 1605 (s), 1514 (s), 1471 (s), 1362 (s), 1255 (s), 1144 (s), 1053 (s); HRMS (ES) m/z calcd for $(\text{C}_{26}\text{H}_{35}\text{BrN}_2\text{O}_5\text{Si}+\text{H})^+$ 563.1577, found 563.1575.

Allylic Sulfide **MGC20**:




[00169] Triethylamine (0.229 mL, 1.64 mmol, 1.3 equiv) and benzenethiol (0.150 mL, 1.45 mmol, 1.15 equiv) were added in sequence to a solution of the allylic bromide **MGC19** (712 mg, 1.26 mmol, 1.0 equiv) in acetonitrile (17 mL) at 0 °C. The mixture was stirred at 0 °C for 20 min, then the cooling bath was removed. The reaction mixture was allowed to warm to 23 °C and stirring was continued at that temperature for 10 min. The reaction mixture was partitioned between ethyl acetate (100 mL) and an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 100 mL). The organic phase was separated and the aqueous phase was further extracted with an additional 30-mL portion of ethyl acetate. The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, furnishing a clear oil. The product was purified by flash column chromatography (0.01:2:8 to 0.013:7 triethylamine-ethyl acetate-hexanes), affording the allylic sulfide **MGC20** as a white foamy solid (728 mg, 97%).

[00170] R_f 0.65 (3:7 ethyl acetate-hexanes); ^1H NMR (400 MHz, C_6D_6) δ 7.35 (d, 2H, $J = 7.2$ Hz, *o*-ArH), 7.19 (m, 2H, *o*-ArH), 6.95 (m, 3H, *p,m*-ArH), 6.89 (m, 2H, *p,m*-ArH), 6.83 (d, 1H, $J = 7.2$ Hz, *p*-ArH), 5.51 (m, 1H, CH=CHCHSPh), 5.12 (m, 2H, CHOTBS, OCHH'Ph), 5.05 (d, 1H, $J = 12.4$ Hz, OCHH'Ph), 4.73 (dt, 1H, $J = 10.0, 2.0$ Hz, CH=CHCHSPh), 4.38 (m, 1H, CH=CHCHSPh), 3.47 (m, 1H, CHCHN(CH₃)₂), 2.92 (d, 1H, $J = 2.0$ Hz, CHCHN(CH₃)₂), 1.75 (s, 6H, N(CH₃)₂), 1.14 (s, 9H, SiC(CH₃)₃), 0.35 (s, 3H, SiCH₃), 0.31 (s, 3H, SiCH₃); ^{13}C NMR (125 MHz, C_6D_6) δ 189.9, 177.0, 168.9, 136.7, 135.2, 131.3, 130.3, 129.2, 128.5, 128.4, 128.3, 126.2, 124.0, 106.2, 79.2, 72.4, 71.7, 63.2, 49.8, 43.4, 39.0, 26.6, 19.1, -2.9, -4.5; FTIR

(neat), cm^{-1} 3310 (m, OH), 2927 (m), 2854 (m), 2792 (w), 1697 (s, C=O), 1621 (s), 1505 (s), 1470 (s), 1365 (s), 1254 (s), 1145 (s), 1089 (s); HRMS (ES) m/z calcd for $(\text{C}_{32}\text{H}_{40}\text{N}_2\text{O}_5\text{SSi}+\text{H})^+$ 593.2505, found 593.2509.

Lower R_f Sulfoxide **MGC21**:



[00171] (–)-[(8,8)-(Dichlorocamphoryl)sunfonyl]oxaziridine (118 mg, 0.395 mmol, 1.5 equiv) was added to a solution of the allylic sulfide **MGC20** (156 mg, 0.263 mmol, 1.0 equiv) in dichloromethane (2 mL) at 23 °C. The mixture was stirred at 23 °C for 20 h, then was concentrated, providing a light brown solid. The product was purified by flash column chromatography (0.001:2:8 to 0.001:3:7 triethylamine-ethyl acetate-hexanes), affording the lower R_f allylic sulfoxide **MGC21** as a white solid (165 mg, 99%).

[00172] R_f 0.18 (3:7 ethyl acetate-hexanes); ^1H NMR (400 MHz, C_6D_6) δ 7.43 (dd, 2H, $J = 8.0, 1.5$ Hz, o -ArH), 7.16 (m, 2H, o -ArH), 6.92 (m, 6H, p,m -ArH), 5.43 (m, 1H, $\text{CH}=\text{CHCHS}(\text{O})\text{Ph}$), 5.33 (d, 1H, $J = 5.0$ Hz, CHOTBS), 5.09 (d, 1H, $J = 11.5$ Hz, $\text{OCHH}'\text{Ph}$), 5.02 (m, 2H, $\text{CH}=\text{CHCHS}(\text{O})\text{Ph}$, $\text{OCHH}'\text{Ph}$), 3.73 (m, 1H, $\text{CH}=\text{CHCHS}(\text{O})\text{Ph}$), 3.41 (m, 1H, $\text{CHCHN}(\text{CH}_3)_2$), 2.85 (d, 1H, $J = 2.5$ Hz, $\text{CHCHN}(\text{CH}_3)_2$), 1.70 (s, 6H, $\text{N}(\text{CH}_3)_2$), 1.12 (s, 9H, $\text{SiC}(\text{CH}_3)_3$), 0.39 (s, 3H, SiCH_3), 0.36 (s, 3H, SiCH_3); ^{13}C NMR (125 MHz, C_6D_6) δ 189.5, 176.9, 168.8, 145.5, 135.2, 130.2, 129.9, 129.0, 128.5, 128.4, 128.3, 127.8, 124.3, 122.9, 106.1, 79.3, 72.4, 70.6, 67.8, 63.1, 43.4, 38.5, 26.6, 19.2, –2.6, –4.7; FTIR (neat), cm^{-1} 3310 (m, OH), 2927 (m), 2854 (m), 2792 (w), 1697 (s, C=O), 1621 (s), 1505 (s), 1470 (s), 1365 (s), 1254 (s), 1145 (s), 1089 (s); HRMS (ES) m/z calcd for $(\text{C}_{32}\text{H}_{40}\text{N}_2\text{O}_6\text{SSi}+\text{H})^+$ 609.2455, found 609.2452.

Rearranged Allylic Alcohol **MGC22**:




R_f 0.40 (3:7 ethyl acetate-hexanes); ^1H NMR (500 MHz, C_6D_6) δ 7.30 (d, 2H, $J = 7.0$ Hz, *o*-ArH), 7.06 (dd, 2H, $J = 7.5, 7.0$ Hz, *m*-ArH), 7.00 (d, 1H, $J = 7.5$ Hz, *p*-ArH), 5.85 (m, 1H, =CHCHOH), 5.42 (br d, 1H, $J = 10.5$ Hz, =CHCHOTBS), 5.16 (d, 1H, $J = 12.5$ Hz, OCHH'Ph), 5.06 (d, 1H, $J = 12.5$ Hz, OCHH'Ph), 4.44 (m, 1H, =CHCHOH), 4.31 (br s, 1H, OH), 4.07 (br s, 1H, =CHCHOTBS), 3.34 (br s, 1H, OH), 3.33 (d, 1H, $J = 11.5$ Hz, CHCHN(CH₃)₂), 2.75 (br d, 1H, $J = 11.5$ Hz, CHCHN(CH₃)₂), 2.03 (s, 6H, N(CH₃)₂), 0.89 (s, 9H, Si(CH₃)₃), -0.11 (s, 3H, SiCH₃), -0.13 (s, 3H, SiCH₃); ^{13}C NMR (100 MHz, C_6D_6) δ 189.7, 182.2, 167.7, 135.2, 129.2, 128.8, 128.3, 128.2, 106.6, 78.6, 71.9, 68.1, 64.1, 59.6, 48.8, 41.2, 25.5, 17.8, -5.2, -5.6; FTIR (neat), cm^{-1} 3515 (m, OH), 2917 (m), 2852 (m), 1708 (s, C=O), 1601 (s), 1511 (s), 1471 (m), 1369 (m), 1254 (m), 1100 (m), 1022 (m); HRMS (ES) m/z calcd for $(\text{C}_{26}\text{H}_{36}\text{N}_2\text{O}_6\text{Si}+\text{H})^+$ 501.2421, found 501.2424.

110

dichloromethane (3 mL) at 23 °C. The reaction mixture was stirred at 23 °C for 2 h, then was partitioned between ethyl acetate (50 mL) and saturated aqueous sodium bicarbonate solution (50 mL). The organic phase was separated and the aqueous phase was further extracted with an additional 30-mL portion of ethyl acetate. The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a clear oil (180 mg, 99%). The product was used in the next step without further purification. An analytical sample was prepared by purification of the crude reaction mixture by flash column chromatography (0.001:2:8 to 0.001:3:7 triethylamine-ethyl acetate-hexanes), affording the benzyl carbonate **MGC23** as a white solid.

[00175] R_f 0.60 (3:7 ethyl acetate-hexanes); ^1H NMR (500 MHz, C_6D_6) δ 7.26 (d, 2H, $J = 7.0$ Hz, *o*-ArH), 7.02 (m, 8H, ArH), 5.75 (br dd, 1H, $J = 10.5, 3.0$ Hz, =CHCHOCO₂Bn), 5.70 (br dd, 1H, $J = 10.5, 2.5$ Hz, =CHCHOTBS), 5.37 (m, 1H, =CHCHOCO₂Bn), 5.10 (d, 1H, $J = 12.5$ Hz, OCHH'Ph), 5.06 (d, 1H, $J = 12.5$ Hz, OCHH'Ph), 4.91 (d, 1H, $J = 12.0$ Hz, OCHH'Ph'), 4.88 (d, 1H, $J = 12.0$ Hz, OCHH'Ph'), 4.41 (m, 1H, =CHCHOTBS), 3.38 (d, 1H, $J = 7.5$ Hz, CHCHN(CH₃)₂), 3.11 (m, 1H, CHCHN(CH₃)₂), 1.92 (s, 6H, N(CH₃)₂), 0.92 (s, 9H, Si(CH₃)₃), 0.02 (s, 3H, SiCH₃), -0.02 (s, 3H, SiCH₃); ^{13}C NMR (100 MHz, C_6D_6) δ 188.9, 179.9, 168.3, 155.2, 135.6, 135.4, 133.2, 128.6, 128.5, 128.4, 128.3, 127.7, 124.9, 107.0, 77.3, 72.2, 71.6, 69.6, 66.6, 60.3, 44.4, 42.2, 25.9, 18.2, -4.8, -4.8; FTIR (neat), cm^{-1} 3532 (w, OH), 2948 (m), 2842 (m), 1738 (s, C=O), 1708 (s, C=O), 1608 (s), 1512 (s), 1471 (m), 1383 (m), 1258 (s), 1101 (m); HRMS (ES) m/z calcd for $(\text{C}_{34}\text{H}_{42}\text{N}_2\text{O}_8\text{Si}+\text{H})^+$ 635.2789, found 635.2786.

Diol **MGC24**:



[00176] Acetic acid (40.0 μL , 0.709 mmol, 2.5 equiv) and a solution of tetrabutylammonium fluoride in tetrahydrofuran (1.0 M, 0.709 mL, 0.709 mmol, 2.5 equiv) were added in sequence to a solution of the benzyl carbonate **MGC23** (180 mg,

0.284 mmol, 1.0 equiv) in tetrahydrofuran (3 mL) at 23 °C. The resulting yellow solution was stirred at 23 °C for 4 h, then was partitioned between ethyl acetate (50 mL) and an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 50 mL). The organic phase was separated and the aqueous phase was further extracted with two 20-mL portions of ethyl acetate. The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a brown oil. The product was purified by flash column chromatography (2:8 to 1:1 ethyl acetate-hexanes), affording the diol **MGC24** as a white solid (135 mg, 92% over 2 steps).

[00177] R_f 0.15 (3:7 ethyl acetate-hexanes); ^1H NMR (500 MHz, C_6D_6) δ 7.24 (d, 2H, $J = 7.0$ Hz, *o*-ArH), 7.02 (m, 8H, ArH), 5.68 (br dd, 1H, $J = 10.5, 2.5$ Hz, =CHCHOCO₂Bn), 5.63 (br dd, 1H, $J = 10.5, 3.0$ Hz, =CHCHOH), 5.26 (m, 1H, =CHCHOCO₂Bn), 5.09 (d, 1H, $J = 12.0$ Hz, OCHH'Ph), 5.05 (d, 1H, $J = 12.0$ Hz, OCHH'Ph), 4.89 (d, 1H, $J = 12.0$ Hz, OCHH'Ph'), 4.86 (d, 1H, $J = 12.0$ Hz, OCHH'Ph'), 4.16 (m, 1H, =CHCHOH), 3.24 (d, 1H, $J = 6.5$ Hz, CHCHN(CH₃)₂), 2.94 (m, 1H, CHCHN(CH₃)₂), 2.25 (br s, 1H, OH), 1.82 (s, 6H, N(CH₃)₂); ^{13}C NMR (100 MHz, CDCl_3) δ 168.1, 154.8, 135.1, 134.9, 132.2, 128.9, 128.9, 128.8, 128.7, 128.6, 126.4, 106.7, 76.6, 72.9, 71.3, 70.3, 64.9, 60.3, 44.4, 43.3; FTIR (neat), cm^{-1} 3468 (m, OH), 3034 (w), 2949 (m), 2798 (m), 1738 (s, C=O), 1705 (s, C=O), 1606 (s), 1513 (s), 1475 (m), 1379 (m), 1261 (s), 1022 (m); HRMS (ES) m/z calcd for $(\text{C}_{28}\text{H}_{28}\text{N}_2\text{O}_8 + \text{H})^+$ 521.1929, found 521.1926.

Cyclohexenone **MGC25**:




[00178] Solid *o*-iodoxybenzoic acid (79.0 mg, 0.281 mmol, 6.5 equiv) was added in one portion to a solution of the diol **MGC24** (22.5 mg, 0.0433 mmol, 1.0 equiv) in dimethylsulfoxide (0.7 mL) at 23 °C. The reaction mixture was initially heterogeneous, but became homogeneous within 5 min. The brown reaction mixture was protected from light and was stirred vigorously at 23 °C for 12 h. The resulting orange reaction

mixture was partitioned between ether (20 mL) and water (20 mL). The organic phase was separated and the aqueous phase was further extracted with two 10 mL-portions ether. The organic phases were combined and washed with saturated aqueous sodium bicarbonate solution (8 mL, containing 30 mg of sodium bisulfite) and brine (10 mL). The washed solution was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated, yielding the cyclohexenone **MGC25** as a white oily solid (22.2 mg, 99%).

[00179] R_f 0.33 (2:3 ethyl acetate-hexanes); ^1H NMR (400 MHz, C_6D_6) δ 7.22 (d, 2H, $J = 6.8$ Hz, $o\text{-ArH}$), 6.99 (m, 8H, ArH), 6.12 (ddd, 1H, $J = 10.4, 4.0, 1.2$ Hz, $\text{CH}=\text{CHCHOCO}_2\text{Bn}$), 5.74 (dd, 1H, $J = 10.4, 1.2$ Hz, $\text{CH}=\text{CHCHOCO}_2\text{Bn}$), 5.41 (ddd, 1H, $J = 4.0, 1.2, 1.2$ Hz, $\text{CH}=\text{CHCHOCO}_2\text{Bn}$), 5.18 (br s, 1H, OH), 5.08 (d, 1H, $J = 12.0$ Hz, $\text{OCHH}'\text{Ph}$), 5.01 (d, 1H, $J = 12.0$ Hz, $\text{OCHH}'\text{Ph}$), 4.89 (d, 1H, $J = 12.4$ Hz, $\text{OCHH}'\text{Ph}'$), 4.83 (d, 1H, $J = 12.4$ Hz, $\text{OCHH}'\text{Ph}'$), 3.28 (d, 1H, $J = 8.4$ Hz, $\text{CHCHN}(\text{CH}_3)_2$), 2.85 (ddd, 1H, $J = 8.4, 4.0, 1.2$ Hz, $\text{CHCHN}(\text{CH}_3)_2$), 1.92 (s, 6H, $\text{N}(\text{CH}_3)_2$); ^{13}C NMR (100 MHz, C_6D_6) δ 192.3, 186.2, 180.5, 167.8, 154.8, 141.8, 135.3, 135.2, 129.9, 128.6, 128.6, 128.5, 128.4, 127.8, 107.7, 78.9, 72.5, 69.9, 59.9, 48.4, 41.9; FTIR (neat), cm^{-1} 3442 (m, OH), 3030 (w), 2948 (m), 2793 (m), 1742 (s, C=O), 1711 (s, C=O), 1608 (s), 1510 (s), 1448 (m), 1376 (m), 1258 (s), 1056 (m); HRMS (ES) m/z calcd for $(\text{C}_{28}\text{H}_{26}\text{N}_2\text{O}_8+\text{H})^+$ 519.1767, found 519.1773.

Silyl-Cyclohexenone **MGC26**:




[00180] Triethylamine (172 μL , 1.24 mmol, 3.5 equiv) and *tert*-butyldimethylsilyl trifluoromethanesulfonate (243 μL , 1.06 mmol, 3.0 equiv) were added in sequence to a solution of the cyclohexenone **MGC25** (183 mg, 0.353 mmol, 1.0 equiv) in tetrahydrofuran (8 mL) at 0 $^\circ\text{C}$. The reaction mixture was stirred at 0 $^\circ\text{C}$ for 40 min, then was partitioned between ethyl acetate (50 mL) and an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 50 mL). The organic phase was

separated and the aqueous phase was further extracted with a 25-mL portion of ethyl acetate. The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a yellow oily solid. The product was purified by flash column chromatography (1:9 to 2:8 ethyl acetate-hexanes), affording the silyl-cyclohexenone **MGC26** as a clear oil (207 mg, 93%).

[00181] R_f 0.50 (3:7 ethyl acetate-hexanes); ^1H NMR (400 MHz, C_6D_6) δ 7.21 (dd, 2H, $J = 7.5, 1.0$ Hz, *o*-ArH), 7.15 (d, 2H, $J = 8.0$ Hz, *o*-ArH), 7.05 (t, 2H, $J = 8.0$ Hz, *m*-ArH), 6.98 (m, 4H, *m,p*-ArH), 6.30 (ddd, 1H, $J = 10.5, 5.0, 2.0$ Hz, CH=CHCHOCO₂Bn), 5.68 (dd, 1H, $J = 10.5, 1.0$ Hz, CH=CHCHOCO₂Bn), 5.65 (br d, 1H, $J = 5.0$ Hz, CH=CHCHOCO₂Bn), 5.10 (d, 1H, $J = 12.5$ Hz, OCHH'Ph), 5.01 (d, 1H, $J = 12.5$ Hz, OCHH'Ph), 4.95 (d, 1H, $J = 12.5$ Hz, OCHH'Ph'), 4.82 (d, 1H, $J = 12.5$ Hz, OCHH'Ph'), 3.11 (d, 1H, $J = 11.0$ Hz, CHCHN(CH₃)₂), 2.94 (br d, 1H, $J = 11.0$ Hz, CHCHN(CH₃)₂), 1.96 (s, 6H, N(CH₃)₂), 1.08 (s, 9H, SiC(CH₃)₃), 0.59 (s, 3H, SiCH₃), 0.29 (s, 3H, SiCH₃); ^{13}C NMR (100 MHz, C_6D_6) δ 193.3, 186.7, 180.3, 167.8, 154.9, 140.9, 135.6, 135.3, 129.9, 128.6, 128.5, 128.5, 128.4, 128.0, 127.8, 108.6, 82.4, 72.4, 69.6, 69.3, 59.7, 50.2, 41.4, 26.5, 19.6, -1.9, -3.4; FTIR (neat), cm^{-1} 2930 (m), 2855 (m), 1745 (s, C=O), 1722 (s, C=O), 1691 (m), 1613 (m), 1513 (s), 1473 (m), 1455 (m), 1378 (m), 1264 (s), 1231 (s), 1046 (m); HRMS (ES) m/z calcd for $(\text{C}_{34}\text{H}_{40}\text{N}_2\text{O}_8+\text{H})^+$ 633.2632, found 633.2620.

Michael-Dieckmann Addition Product **MGC27**:




[00182] A solution of *n*-butyllithium in hexanes (1.55 M, 155 μL , 0.241 mmol, 5.1 equiv) was added to a solution of *N,N,N',N'*-tetramethylethylenediamine (39.0 μL ,

0.261 mmol, 5.5 equiv) and diisopropyl amine (34.0 μ L, 0.249 mmol, 5.25 equiv) in tetrahydrofuran (1 mL) at -78 $^{\circ}$ C. The resulting mixture was stirred vigorously at -78 $^{\circ}$ C for 30 min whereupon a solution of the ester **CDL-I-280** (73.0 mg, 0.213 mmol, 4.5 equiv) in tetrahydrofuran (1 mL) was added dropwise via cannula. The resulting deep red mixture was stirred vigorously at -78 $^{\circ}$ C for 75 min, then a solution of the silyl-cyclohexenone **MGC26** (30.0 mg, 0.0474 mmol, 1.0 equiv) in tetrahydrofuran (1 mL) was added dropwise via cannula. The resulting light red mixture was allowed to warm slowly to 0 $^{\circ}$ C over 2 h, then was partitioned between an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 10 mL) and dichloromethane (10 mL). The organic phase was separated and the aqueous phase was further extracted with two 10-mL portions of dichloromethane. The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a yellow oil. The product was purified by preparatory HPLC on a Coulter Ultrasphere ODS column (10 μ M, 250 x 10 mm, flow rate 3.5 mL/min, Solvent A: methanol, Solvent B: water) using an injection volume of 400 μ L (methanol) and an isocratic elution of 10% B for 75 min. The peak eluting at 36-42 min was collected and concentrated, affording the Michael-Dieckmann addition product **MGC27** (33.0 mg, 80%) as a light yellow solid.

[00183] R_f 0.35 (1:4 ethyl acetate-hexanes); ^1H NMR (500 MHz, C_6D_6) δ 16.55 (br s, 1H, enol), 7.26 (d, 2H, $J = 7.0$ Hz, o -ArH), 7.14 (d, 2H, $J = 7.5$ Hz, ArH), 6.85-7.05 (m, 6H, ArH), 6.66-6.74 (m, 2H, ArH), 6.51 (dd, 1H, $J = 9.0, 1.5$ Hz, ArH), 5.73 (br d, 1H, $J = 4.0$ Hz, BnOCO_2CH), 5.17 (d, 1H, $J = 12.5$ Hz, $\text{OCHH}'\text{Ph}$), 5.03 (d, 1H, $J = 12.5$ Hz, $\text{OCHH}'\text{Ph}$), 4.99 (d, 1H, $J = 12.5$ Hz, $\text{OCHH}'\text{Ph}'$), 4.93 (d, 1H, $J = 12.5$ Hz, $\text{OCHH}'\text{Ph}'$), 3.58 (d, 1H, $J = 11.5$ Hz, $\text{CHCHN}(\text{CH}_3)_2$), 3.35 (dd, 1H, $J = 12.5, 4.0$ Hz, CH_3CHCH), 2.99 (d, 1H, $J = 11.5$ Hz, $\text{CHCHN}(\text{CH}_3)_2$), 2.56 (dq, 1H, $J = 12.5, 7.0$ Hz, CH_3CH), 2.18 (s, 6H, $\text{N}(\text{CH}_3)_2$), 1.33 (s, 9H, $\text{C}(\text{CH}_3)_3$), 1.16 (d, 3H, $J = 7.0$ Hz, CH_3CH), 1.11 (s, 9H, $\text{C}(\text{CH}_3)_3$), 0.61 (s, 3H, CH_3), 0.36 (s, 3H, CH_3); ^{13}C NMR (100 MHz, CDCl_3) δ 189.7, 186.3, 180.9, 178.4, 167.9, 154.7, 152.1, 150.8, 145.9, 136.1, 135.5, 133.9, 128.7, 128.6, 128.5, 127.3, 123.8, 122.7, 122.6, 108.9, 105.5, 83.0, 82.9, 74.8, 72.4, 69.2, 60.8, 52.7, 43.2, 38.4, 27.5, 26.6, 19.5, 16.3, -1.8 , -2.7 ; FTIR (neat film), cm^{-1} 2974 (w), 2933 (w), 2851 (w), 1760 (s, $\text{C}=\text{O}$), 1748 (s, $\text{C}=\text{O}$), 1723 (s,

C=O), 1606 (m), 1513 (m), 1471 (m), 1370 (m). 1260 (s), 1232 (s), 1148 (s); HRMS (ES) m/z calcd for $(C_{48}H_{56}O_{12}N_2Si)^+$ 881.3681, found 881.3684.


Initial Deprotection of Michael-Dieckmann Addition Product **MGC28**:



[00184] Hydrofluoric acid (1.2 mL, 48% aqueous) was added to a polypropylene reaction vessel containing a solution of the Michael-Dieckmann addition product **MGC27** (33.0 mg, 0.0375 mmol, 1.0 equiv) in acetonitrile (7.0 mL) at 23 °C. The resulting mixture was stirred vigorously at 23 °C for 60 h, then was poured into water (50 mL) containing K_2HPO_4 (7.0 g). The resulting mixture was extracted with ethyl acetate (3×20 mL). The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, furnishing the pentacyclic phenol **MGC28** as a yellow oil (25.0 mg, 99%). The product was used in the next step without further purification.

[00185] R_f 0.05 (1:4 ethyl acetate-hexanes); 1H NMR (600 MHz, C_6D_6 , crude) δ 14.86 (br s, 1H, enol), 11.95 (s, 1H, phenol), 7.23 (d, 2H, $J = 7.8$ Hz, *o*-ArH), 7.14 (d, 2H, $J = 7.2$ Hz, *o*-ArH), 6.94-7.02 (m, 6H, ArH), 6.86 (t, 1H, $J = 8.4$ Hz, ArH), 6.76 (d, 1H, $J = 8.4$ Hz, ArH), 6.28 (d, 1H, $J = 7.8$ Hz, ArH), 5.46 (dd, 1H, $J = 3.6, 3.0$ Hz, $BnOCO_2CH$), 5.12 (d, 1H, $J = 12.0$ Hz, $OCHH'Ph$), 5.04 (d, 1H, $J = 12.0$ Hz, $OCHH'Ph$), 4.92 (s, 2H, OCH_2Ph), 3.41 (d, 1H, $J = 9.6$ Hz, $CHCHN(CH_3)_2$), 2.82 (dd, 1H, $J = 9.6, 3.0$ Hz, $CHCHN(CH_3)_2$), 2.65 (dd, 1H, $J = 13.2, 3.6$ Hz, CH_3CHCH), 2.78 (dq, 1H, $J = 13.2, 7.2$ Hz, CH_3CH), 2.05 (s, 6H, $N(CH_3)_2$), 1.04 (d, 3H, $J = 7.2$ Hz, CH_3CH); ^{13}C NMR (100 MHz, C_6D_6 , crude) δ 193.4, 186.2, 181.3, 172.3, 167.9, 163.3, 154.6, 145.8, 136.6, 135.8, 128.6, 128.4, 127.2, 116.8, 116.0, 115.6, 107.6, 104.7, 76.8, 73.9, 72.5, 69.5, 60.3, 48.7, 43.0, 41.8, 37.5, 15.3; FTIR (neat film), cm^{-1} 3424 (m, OH), 3059, 3030, 2925, 2857, 1744 (s, C=O), 1713 (s, C=O), 1614 (s), 1582 (s), 1455 (s), 1252 (s); HRMS (ES) m/z calcd for $(C_{37}H_{34}O_{10}N_2+H)^+$ 667.2292, found 667.2300.

(-)-Doxycycline (MGC30):




[00186] Pd black (7.00 mg, 0.0657 mmol, 1.75 equiv) was added in one portion to a solution of the pentacyclic phenol **MGC28** (25.0 mg, 0.0375 mmol, 1.0 equiv) in tetrahydrofuran-methanol (1:1, 2.0 mL) at 23 °C. An atmosphere of hydrogen was introduced by briefly evacuating the flask, then flushing with pure hydrogen (1 atm). The Pd catalyst was initially present as a fine dispersion, but aggregated into clumps within 5 min. The yellow heterogeneous mixture was stirred at 23 °C for 2 h, then was filtered through a plug of cotton. The filtrate was concentrated, affording a yellow oil (>95% doxycycline based on ¹H NMR analysis). The product was purified by preparatory HPLC on a Phenomenex Polymerx DVB column (10 μM, 250 x 10 mm, flow rate 4.0 mL/min, Solvent A: methanol-0.005 N aq. HCl (1:4), Solvent B: acetonitrile) using an injection volume of solvent A (400 μL) containing oxalic acid (10 mg) and an isocratic elution of 5% B for 2 min, then a gradient elution of 5-50% B for 20 min. The peak eluting at 12-17 min was collected and concentrated, affording (-)-doxycycline hydrochloride as a yellow powder (16.2 mg, 90%), which was identical with natural (-)-doxycycline hydrochloride in all respects.

[00187] ¹H NMR (600 MHz, CD₃OD, hydrochloride) δ 7.47 (t, 1H, *J* = 8.4 Hz, ArH), 6.93 (d, 1H, *J* = 8.4 Hz, ArH), 6.83 (d, 1H, *J* = 8.4 Hz, ArH), 4.40 (s, 1H, (CH₃)₂NCH), 3.53 (dd, 1H, *J* = 12.0, 8.4 Hz, CHOH), 2.95 (s, 3H, N(CH₃)CH₃'), 2.88 (s, 3H, N(CH₃)CH₃'), 2.80 (d, 1H, *J* = 12.0 Hz, CHCHN(CH₃)₂), 2.74 (dq, 1H, *J* = 12.6, 6.6 Hz, CH₃CH), 2.58 (dd, 1H, *J* = 12.6, 8.4 Hz, CH₃CHCH), 1.55 (d, 3H, *J* = 6.6 Hz, CH₃CHCH); ¹³C NMR (100 MHz, CD₃OD) δ 195.3, 188.2, 173.8, 172.1, 163.2, 149.0, 137.7, 117.1, 116.9, 116.6, 108.4, 96.0, 74.5, 69.8, 66.9, 47.5, 43.4, 43.0, 41.9, 40.0, 16.3; UV max (0.01 N methanolic HCl), nm 218, 267, 350; [α]_D = -109° (c = 0.16 in 0.01 M methanolic HCl); lit. (*The Merck Index: An Encyclopedia of Chemicals*,

Drugs, and Biologicals, 12th ed. Budavari, S.; O'Neal, M. J.; Smith, A.; Heckelman, P. E.; Kinneary, J. F., Eds.; Merck & Co.: Whitehouse Station, NJ, 1996; entry 3496.) UV max (0.01 N methanolic HCl), nm 267, 351; $[\alpha]_D = -110^\circ$ ($c = 1$ in 0.01 M methanolic HCl); HRMS (ES) m/z calcd for $(C_{22}H_{24}O_8N_2+H)^+$ 445.1611, found 445.1603.

Example 3-Synthesis of 6-Deoxytetracycline

Ester CDL-I-280:




[00188] A solution of *sec*-butyllithium in cyclohexane (1.40 M, 24.0 mL, 33.6 mmol, 2.6 equiv) was added to a solution of *N,N,N',N'*-tetramethylethylenediamine (4.9 mL, 33 mmol, 2.5 equiv) in tetrahydrofuran (25 mL) at -78°C . The resulting yellow solution was cooled to -90°C (internal temperature) in a liquid nitrogen-ethanol bath. A solution of *o*-anisic acid (2.00 g, 13.1 mmol, 1.0 equiv) in tetrahydrofuran (10 mL) was added dropwise via cannula over a period of 30 min to the yellow solution. The resulting orange suspension was stirred for an additional 30 min at -90°C , then was allowed to warm to -78°C over 15 min, whereupon iodoethane (4.2 mL, 52 mmol, 4.0 equiv) was added. The mixture was allowed to warm to 23°C over 15 min, then was partitioned between water (50 mL) and ether (50 mL). The aqueous layer was separated and diluted with aqueous hydrochloric acid (1.0 M, 100 mL). The resulting mixture was extracted with ethyl acetate (4×80 mL). The organic layers were combined and then dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a brown oil (1.8 g). ^1H NMR (500 MHz, CDCl_3) analysis of the crude product showed an 8:2 ratio of the carboxylic acid **CDL-I-279** (δ 3.89, OCH_3) and unreacted anisic acid (δ 4.07, OCH_3). Oxalyl chloride (1.0 mL, 11 mmol, 0.8 equiv) and *N,N*-dimethylformamide (100 μL) were added in sequence to a solution of the residue in dichloromethane (20 mL) at 23°C . Vigorous gas evolution was observed upon addition of *N,N*-dimethylformamide. The reaction

mixture was stirred for 2 h at 23 °C, whereupon phenol (1.4 g, 15 mmol, 1.1 equiv), pyridine (2.4 mL, 30 mmol, 2.3 equiv), and 4-(dimethylamino)pyridine (10 mg, 0.081 mmol, 0.006 equiv) were added in sequence at 23 °C. The resulting brown reaction mixture was then stirred for 2 h at 23 °C. Aqueous hydrochloric acid (1 M, 50 mL) was added and the resulting mixture was extracted with ethyl acetate (2 × 50 mL). The organic layers were combined, then washed with an aqueous sodium hydroxide solution (0.1 M, 50 mL), followed by brine (50 mL), and were then dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a clear oil. The product was purified by flash column chromatography (5:95 ethyl acetate-hexanes), affording the ester **CDL-I-280** as a colorless oil (1.7 g, 50%).

[00189] R_f 0.28 (5:95 ethyl acetate-hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.56 (t, 2H, $J = 7.8$ Hz, ArH), 7.37 (t, 1H, $J = 7.8$ Hz, ArH), 7.31-7.26 (m, 3H, ArH), 6.93 (d, 1H, $J = 7.8$ Hz, ArH), 6.85 (d, 1H, $J = 8.3$ Hz, ArH), 3.91 (s, 3H, OCH_3), 2.79 (q, 2H, $J = 7.8$ Hz, CH_2CH_3), 1.33 (t, 3H, $J = 7.8$ Hz, CH_2CH_3); ^{13}C NMR (125 MHz, CDCl_3) δ 166.9, 156.5, 150.8, 142.8, 130.9, 129.5, 125.9, 122.5, 121.6, 120.9, 108.5, 55.9, 26.6, 15.6; FTIR (neat film), cm^{-1} 2970 (m), 1740 (s, C=O), 1583 (s), 1488 (s), 1471 (s), 1438 (m), 1298 (w), 1270 (s), 1236 (s), 1186 (s), 1158 (m), 1091 (m), 1046 (s), 1001 (w); HRMS (ES) m/z calcd for $(\text{C}_{16}\text{H}_{16}\text{O}_3+\text{H})^+$ 257.1178, found 257.1183.

Phenol **CDL-I-298**:




[00190] A solution of boron tribromide in dichloromethane (1.0 M, 5.2 mL, 5.2 mmol, 2.0 equiv) was added to a solution of the ester **CDL-I-280** (662 mg, 2.58 mmol, 1.0 equiv) in dichloromethane (10 mL) at 0 °C. The resulting yellow solution was stirred for 70 min at 0 °C, whereupon saturated aqueous sodium bicarbonate solution (50 mL) was added. The resulting biphasic mixture was stirred for 20 min at 0 °C, dichloromethane (50 mL) was added, the layers were separated, and the aqueous phase

was further extracted with dichloromethane (50 mL). The organic layers were combined and then dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing the phenol **CDL-I-298** as a colorless oil (605 mg, 97%).

[00191] R_f 0.47 (5:95 ethyl acetate-hexanes); ^1H NMR (500 MHz, CDCl_3) δ 10.94 (s, 1H, OH), 7.49 (t, 2H, $J = 7.8$ Hz, ArH), 7.41 (t, 1H, $J = 7.8$ Hz, ArH), 7.35 (t, 1H, $J = 7.3$ Hz, ArH), 7.24 (d, 2H, $J = 7.8$ Hz, ArH), 6.93 (d, 1H, $J = 8.3$ Hz, ArH), 6.85 (d, 1H, $J = 8.3$ Hz, ArH), 3.13 (q, 2H, $J = 7.8$ Hz, CH_2CH_3), 1.34 (t, 3H, $J = 7.8$ Hz, CH_2CH_3); ^{13}C NMR (125 MHz, CDCl_3) δ 170.3, 163.2, 149.8, 147.8, 135.1, 129.7, 126.4, 122.0, 121.6, 115.9, 111.1, 29.8, 16.4; FTIR (neat film), cm^{-1} 2973 (w), 1670 (s, C=O), 1609 (m), 1588 (m), 1490 (w), 1444 (m), 1311 (m), 1295 (m), 1234 (m), 1187 (s), 1162 (s), 1105 (m); HRMS (ES) m/z calcd for $(\text{C}_{15}\text{H}_{14}\text{O}_3 + \text{H})^+$ 243.1021, found 243.1014.

Ester **CDL-I-299**:




[00192] *N,N*-diisopropylethylamine (520 μL , 2.99 mmol, 1.2 equiv), di-*t*-butyl dicarbonate (645 mg, 2.96 mmol, 1.2 equiv), and 4-(dimethylamino)pyridine (31 mg, 0.25 mmol, 1.5 equiv) were added in sequence to a solution of the phenol **CDL-I-298** (605 mg, 2.50 mmol, 0.1 equiv) in dichloromethane (10 mL) at 23 $^\circ\text{C}$. The reaction mixture was stirred for 1 h at 23 $^\circ\text{C}$, whereupon saturated aqueous ammonium chloride solution (50 mL) was added. Dichloromethane (50 mL) was added, the layers were separated, and the aqueous phase was extracted with dichloromethane (50 mL). The organic layers were combined and then dried over sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a brown oil. The product was purified by flash column chromatography (1:9 ether-hexanes), affording the ester **CDL-I-299** as a colorless oil, which crystallized upon standing overnight at -14 $^\circ\text{C}$ (733 mg,

86%), mp 58 °C.

[00193] R_f 0.23 (1:9 ether-hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.46-7.42 (m, 3H, ArH), 7.31-7.26 (m, 3H, ArH), 7.22 (d, 1H, $J = 7.3$ Hz, ArH), 7.15 (d, 1H, $J = 7.3$ Hz, ArH), 2.86 (q, 2H, $J = 7.3$ Hz, CH_2CH_3), 1.46 (s, 9H, Boc), 1.31 (t, 3H, $J = 7.3$ Hz, CH_2CH_3); ^{13}C NMR (125 MHz, CDCl_3) δ 165.1, 151.6, 150.6, 148.7, 144.5, 131.3, 129.4, 126.8, 126.1, 125.4, 121.7, 120.5, 83.8, 27.5, 26.8, 15.6; FTIR (neat film), cm^{-1} 2964 (w), 1754 (s, C=O), 1586 (w), 1491 (w), 1467 (w), 1457 (w), 1368 (w), 1278 (s), 1234 (s), 1190 (s), 1145 (s), 1051 (m); HRMS (ES) m/z calcd for $(\text{C}_{20}\text{H}_{22}\text{O}_5 + \text{NH}_4)^+$ 360.1811, found 360.1808.

Michael-Dieckmann Addition Product **CDL-I-287**:




[00194] A solution of *n*-butyllithium in hexanes (1.45 M, 47 μL , 0.068 mmol, 6.8 equiv) was added to a solution of diisopropylamine (10 μL , 0.071 mmol, 7.1 equiv) and *N,N,N',N'*-tetramethylethylenediamine (10 μL , 0.066 mmol, 6.6 equiv) in tetrahydrofuran (300 μL) at -78 °C. The resulting solution was stirred at -78 °C for 30 min whereupon a solution of the ester **CDL-I-299** (17 mg, 0.050 mmol, 5.0 equiv) in tetrahydrofuran (200 μL) was added, forming a deep red solution. The solution was stirred at -78 °C for 75 min, then a solution of the enone **DRS6** (5.0 mg, 0.010 mmol, 1.0 equiv) in tetrahydrofuran (100 μL) was added at -78 °C. The color of the reaction mixture remained deep red following the addition. The mixture was allowed to warm to 0 °C over 150 min. Upon reaching 0 °C, an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 15 mL) was added. The resulting yellow mixture was extracted with dichloromethane (3 $\times$ 15 mL). The organic layers were combined and

then dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a yellow oil. The product was purified by preparatory HPLC on a Coulter Ultrasphere ODS column (5 μ m, 250 $\times$ 10 mm, flow rate 3.5 mL/min, Solvent A: water, Solvent B: methanol, UV detection at 350 nm) using an injection volume of 500 μ L methanol with an isocratic elution of 89.5% B. The peak eluting at 31–40 min was collected and concentrated affording the Michael-Dieckmann product **CDL-I-287** as a light yellow solid (6.1 mg, 83%), mp 114 $^{\circ}$ C.

[00195] R_f 0.37 (2:8 tetrahydrofuran-hexanes); ^1H NMR (500 MHz, CDCl_3) δ (s, 1H, 16.24, enol-OH), 7.55–7.50 (m, 3H, ArH), 7.40–7.35 (m, 4H, ArH), 7.10 (d, 1H, J = 7.8 Hz, ArH), 5.39–5.34 (m, 2H, OCH_2Ph), 3.92 (d, 1H, J = 10.7 Hz, $\text{CHN}(\text{CH}_3)_2$), 2.81–2.71 (m, 2H, CH_3CH , CH_3CHCH), 2.55 (dd, 1H, J = 10.7, 5.7 Hz, $\text{CHCHN}(\text{CH}_3)_2$), 2.48 (s, 6H, $\text{N}(\text{CH}_3)_2$), 2.40 (d, 1H, J = 14.7 Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 2.31 (ddd, 1H, J = 14.7, 9.3, 5.7, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 1.56 (s, 3H, CH_3), 1.55 (s, 9H, Boc), 0.84 (s, 9H, TBS), 0.27 (s, 3H, TBS), 0.13 (s, 3H, TBS); ^{13}C NMR (125 MHz, CDCl_3) δ 187.4, 183.1, 182.8, 181.6, 167.6, 151.7, 150.2, 147.4, 135.0, 134.0, 128.5, 128.5, 123.4, 123.0, 122.4, 108.3, 107.4, 94.8, 83.9, 81.5, 72.5, 61.5, 46.4, 41.9, 39.5, 34.9, 27.7, 26.0, 20.7, 19.0, 16.0, -2.6, -3.7; FTIR (neat film), cm^{-1} 2923 (m), 2841 (m), 1759 (s, C=O), 1718 (s, C=O), 1605 (s), 1508 (s), 1467 (m), 1456 (m), 1369 (m), 1277 (s), 1262 (m), 1231 (s), 1144 (s), 1005 (w); HRMS (ES) m/z calcd for $(\text{C}_{40}\text{H}_{50}\text{N}_2\text{O}_9\text{Si}+\text{H})^+$ 731.3364, found 731.3370.

6-Deoxytetracycline **CDL-I-322**




[00196] Hydrofluoric acid (0.6 mL, 48% aqueous) was added to a polypropylene reaction vessel containing a solution of the Michael-Dieckmann addition product **CDL-I-287** (15 mg, 0.021 mmol, 1.0 equiv) in acetonitrile (3.5 mL) at 23 $^{\circ}$ C. The reaction mixture was stirred at 23 $^{\circ}$ C for 55 h, then was poured into water (20 mL) containing

K₂HPO₄ (4.0 g). The resulting mixture was extracted with ethyl acetate (4 × 20 mL). The organic phases were combined and then dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a light yellow oil. Pd black (7.6 mg, 0.071 mmol, 3.4 equiv) was added in one portion to a solution of the residue in methanol-tetrahydrofuran (1:1, 2 mL). An atmosphere of hydrogen gas was introduced by briefly evacuating the flask, then flushing with pure hydrogen (1 atm). The mixture was stirred at 23 °C for 2 h. Within 5 min, the color changed from light yellow to dark yellow. The reaction mixture was filtered through a plug of cotton. The filtrate was concentrated, affording a yellow oil (10 mg). The product was purified by preparatory HPLC on a Phenomenex Polymerx DVB column (10 μm, 250 × 10 mm, flow rate 5 mL/min, Solvent A: methanol-0.02 N HCl (1:4), Solvent B: acetonitrile, UV detection at 365 nm) using an injection volume of 400 μL methanol containing oxalic acid monohydrate (10 mg) and an isocratic elution of 18% B for 15 min, then a linear gradient elution of 18-60% B in 15 min. The peak eluting at 17.5-22.5 min was collected and concentrated to give 6-deoxytetracycline hydrochloride (**CDL-I-322**·HCl) as a yellow powder (8.1 mg, 81%).

[00197] ¹H NMR (500 MHz, CD₃OD, hydrochloride) δ 7.49 (t, 1H, *J* = 7.8 Hz, ArH), 6.95 (d, 1H, *J* = 7.8 Hz, ArH), 6.84 (d, 1H, *J* = 7.8 Hz, ArH), 4.09 (s, 1H, CHN(CH₃)₂), 3.03 (br s, 3H, N(CH₃)), 2.97 (br s, 3H, N(CH₃)), 2.90 (br d, 1H, *J* = 12.7 Hz, CHCHN(CH₃)₂), 2.67 (ddd, 1H, *J* = 12.7, 12.7, 5.2 Hz, CH₃CHCH), 2.61-2.56 (m, 1H, CH₃CH), 2.30 (ddd, *J* = 13.7, 5.2, 2.9 Hz, CHH'CHCHN(CH₃)₂), 1.54 (ddd, *J* = 13.7, 12.7, 12.7 Hz, CHH'CHCHN(CH₃)₂), 1.38 (d, 3H, *J* = 6.8 Hz, CH₃CH). HRMS (ES) *m/z* calcd for (C₂₂H₂₄N₂O₇+H)⁺ 429.1662, found 429.1660.

Example 4-Synthesis of a Pyridone Sancycline Analog


Phenyl Ester CDL-II-464:



[00198] 2,4,6-Trichlorobenzoyl chloride (356 μL , 2.28 mmol, 1.1 equiv) was added to a solution of the carboxylic acid **CDL-II-417** (reported by A.N. Osman, M.M. Ismail, M.A. Barakat, *Revue Roumaine de Chimie* **1986**, 31, 615-624) (534 mg, 2.08 mmol, 1.0 equiv) and triethylamine (320 μL , 2.28 mmol, 1.1 equiv) in tetrahydrofuran (25 mL) at 23 $^{\circ}\text{C}$. A white precipitate was formed upon addition. The reaction mixture was stirred for 30 min at 23 $^{\circ}\text{C}$. A solution of phenol (489 mg, 5.20 mmol, 2.5 equiv) and 4-(dimethylamino)pyridine (583 mg, 5.20 mmol, 2.5 equiv) in tetrahydrofuran (10 mL) was added via cannula to the reaction mixture prepared above at 0 $^{\circ}\text{C}$. The resulting mixture was allowed to warm to 23 $^{\circ}\text{C}$ over 10 min, and was stirred for 90 min at that temperature. An aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 30 mL) was then added and the resulting mixture was extracted with dichloromethane ($3 \times 30 \text{ mL}$). The organic extracts were combined and then dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate concentrated, providing a colorless oil. The product was purified by flash column chromatography (6:94 ethyl acetate-hexanes), affording the phenyl ester **CDL-II-464** as a white solid (590 mg, 85%), mp 65 $^{\circ}\text{C}$.

[00199] R_f 0.33 (1:9 ethyl acetate-hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.49 (d, 2H, $J = 7.3 \text{ Hz}$, ArH), 7.40-7.24 (m, 6H, ArH), 7.14 (d, 2H, $J = 7.3 \text{ Hz}$, ArH), 6.69 (s, 1H, pyr-H), 5.49 (s, 2H, CH_2Ph), 2.47 (s, 3H, CH_3), 2.43 (s, 3H, CH_3); ^{13}C NMR (125 MHz, CDCl_3) δ 165.9, 160.1, 157.8, 150.7, 148.5, 137.3, 129.4, 128.3, 127.7, 127.6, 125.9, 121.7, 118.1, 113.4, 67.8, 24.1, 19.2; FTIR (neat film), cm^{-1} 1738 (s, C=O), 1600 (s), 1569 (s), 1492 (m), 1441 (m), 1400 (m), 1333 (s), 1272 (s), 1185 (s), 1159 (m), 1097 (m), 1051 (s); HRMS (ES) m/z calcd for $(\text{C}_{21}\text{H}_{19}\text{NO}_3 + \text{H})^+$ 334.1443, found 334.1442.

Michael-Dieckmann Addition Product **CDL-II-466**:




[00200] A solution of *n*-butyllithium in hexanes (1.67 M, 80 μL , 0.13 mmol, 4.2 equiv) was added to a solution of diisopropylamine (20 μL , 0.14 mmol, 4.5 equiv) in tetrahydrofuran (2.5 mL) at $-78\text{ }^{\circ}\text{C}$. The resulting solution was allowed to warm to $0\text{ }^{\circ}\text{C}$ over 15 min. *N,N'*-dimethylpropyleneurea (17 μL , 0.14 mmol, 4.5 equiv) was added to the mixture prepared above at $0\text{ }^{\circ}\text{C}$, whereupon the mixture was cooled to $-78\text{ }^{\circ}\text{C}$. A solution of the ester **CDL-II-464** (31 mg, 0.093 mmol, 3.0 equiv) in tetrahydrofuran (250 μL) was then added at $-78\text{ }^{\circ}\text{C}$. The resulting yellow solution was stirred for 5 min at $-78\text{ }^{\circ}\text{C}$, then a solution of the enone **DRS6** (15 mg, 0.031 mmol, 1.0 equiv) in tetrahydrofuran (250 μL) was added at $-78\text{ }^{\circ}\text{C}$. The resulting deep red mixture was allowed to warm to $0\text{ }^{\circ}\text{C}$ over 4 h. Acetic acid (40 μL) was added at to the deep red mixture at $0\text{ }^{\circ}\text{C}$, followed by an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 15 mL). The resulting yellow mixture was extracted with dichloromethane ($3 \times 15\text{ mL}$). The organic extracts were combined and then dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a yellow oil. The product was purified by preparatory HPLC on a Coulter Ultrasphere ODS column (5 μm , $250 \times 10\text{ mm}$, flow rate 3.5 mL/min, Solvent A: water, Solvent B: methanol, UV detection at 350 nm) using an injection volume of 500 μL DMSO and a gradient elution of 92-100% B over 30 min. The peak eluting at 21-29 min was collected and concentrated to give enol **CDL-II-466** as a light yellow solid (15.0 mg, 67%).

[00201] R_f 0.55 (3:7 ethyl acetate-hexanes); ^1H NMR (600 MHz, CD_2Cl_2) δ 16.05 (s, 1H, enol-OH), 7.52-7.26 (m, 10H, ArH), 6.66 (s, 1H, pyr-H), 5.57 (d, 1H, $J =$

min. The peak at 7.0 min indicated the desired product. After stirring for 3 h at 23 °C the deprotection was complete and the mixture was concentrated to a yellow oil. The crude mixture was purified by preparatory HPLC on a Phenomenex Polymerx DVB column (10 μ m, 250 $\times$ 10 mm, flow rate 4 ml/min, Solvent A: 0.01 N aqueous hydrochloric acid, Solvent B: acetonitrile, UV detection at 365 nm) using an injection volume of 500 μ L methanol containing oxalic acid monohydrate (30 mg) and a linear gradient of 0-20% B over 40 min. The peak eluting at 20-29 min was collected and concentrated to give the hydrochloride of **CDL-II-460** as a yellow powder (4.8 mg, 74%).

[00203] ^1H NMR (500 MHz, CD_3OD , hydrochloride) δ 6.37 (s, 1H, ArH), 4.06 (s, 1H, $\text{CHN}(\text{CH}_3)_2$), 3.05-2.95 (m, 8H, $\text{N}(\text{CH}_3)_2$, $\text{CHCHN}(\text{CH}_3)_2$, $\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.79 (dd, 1H, $J = 16.1, 3.9$ Hz, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.55 (dd, 1H, $J = 16.1, 16.1$ Hz, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.40 (s, 3H, Ar- CH_3), 2.18 (br. d, 1H, $J = 12.7$ Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 1.59 (ddd, 1H, $J = 12.7, 12.7, 12.7$ Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$); ^{13}C NMR (100 MHz, $(\text{CD}_3)_2\text{SO}$) δ 187.3, 183.5, 177.8, 172.1, 160.6, 159.8, 153.3, 115.3, 107.2, 106.9, 95.6, 74.2, 68.4, 41.5, 35.7, 34.5, 33.9, 31.0, 19.2; HRMS (ES) m/z calcd for $(\text{C}_{21}\text{H}_{23}\text{N}_3\text{O}_7+\text{H})^+$ 430.1614, found 430.1607.


Example 5-Synthesis of Pyridine Sancycline Analog (7-Aza-10-Deoxysancycline)



[00204] A solution of 2-methyl-nicotinic acid ethyl ester **JDB1-67-SM** (0.589 g, 3.56 mmol, 1.0 equiv), aqueous sodium hydroxide (1.0 M, 3.9 mL, 3.9 mmol, 1.1 equiv), and ethanol (5 mL) was heated at reflux for 18 h. The reaction mixture was allowed to cool to 23 °C, and was concentrated, affording the carboxylate salt (710 mg) as a white solid. Oxalyl chloride (357 μ L, 4.09 mmol, 1.15 equiv) was added to a mixture of the carboxylate salt in dichloromethane (20 mL) at 23 °C. Vigorous gas evolution was observed upon addition. The reaction mixture was stirred at 23 °C for 30 min, then *N,N*-dimethylformamide (20 μ L) was added. After stirring for an additional

30 min at 23 °C, phenol (837 mg, 8.90 mmol, 2.5 equiv), pyridine (864 μ L, 10.7 mmol, 3.0 equiv), and dimethylaminopyridine (3 mg) were added in sequence. The resulting solution was stirred for 90 min at 23 °C, whereupon an aqueous potassium phosphate buffer solution (pH 7.05, 0.2 M, 5.0 mL) was added. The resulting mixture was partitioned between water (30 mL) and ethyl acetate (50 mL). The aqueous phase was extracted with an additional 50-mL portion of ethyl acetate. The organic layers were combined and washed with an aqueous sodium hydroxide solution (50 mL, 1M), brine (50 mL), and then dried over anhydrous sodium sulfate. The dried solution was decanted and concentrated, affording a colorless oil (900 mg). The product was purified by flash column chromatography (25:75 ethyl acetate-hexanes), providing the ester **JDB1-67** as a colorless oil (500 mg, 66%).


[00205] R_f 0.15 (3:7 ethyl acetate-hexanes); ^1H NMR (300 MHz, CDCl_3) δ 8.70 (dd, 1H, $J = 1.7, 4.95$ Hz, pyr-H), 8.44 (dd, 1H, $J = 1.7, 7.8$ Hz, pyr-H), 7.48-7.43 (m, 2H, ArH), 7.33-7.20 (m, 4H, ArH, pyr-H), 2.93 (s, 1H, CH_3); ^{13}C NMR (100 MHz, CDCl_3) δ 164.8, 160.8, 152.4, 150.5, 138.9, 129.5, 126.1, 124.5, 121.6, 121.0, 25.0; FTIR (neat film), cm^{-1} 3406 (m), 1948 (w), 1747 (s), 1578 (s), 1487 (s), 1435 (s), 1273 (s), 1237 (s), 1191 (s), 1046 (s), 915 (m), 822 (m), 749 (s), 689 (s); HRMS (ES) m/z calcd for $(\text{C}_{13}\text{H}_{11}\text{NO}_2 + \text{H})^+$ 214.0868, found 214.0866.



[00206] A solution of *n*-butyllithium in hexanes (1.47 M, 136 μ L, 0.200 mmol, 8.03 equiv) was added to a solution of diisopropylamine (26.5 μ L, 0.202 mmol, 8.05 equiv) in tetrahydrofuran (0.750 mL) at -78 °C. The reaction mixture was briefly (10 min) transferred to an ice bath, with stirring, then was cooled to -78 °C. Hexamethylphosphoramide (49.0 μ L, 0.399 mmol, 16.0 equiv) was added to the mixture prepared above at -78 °C. The resulting mixture was stirred for 5 minutes


whereupon a colorless solution was formed. The resulting solution was added dropwise via cannula to a solution of the ester **JDB1-67** (36.0 mg, 0.169 mmol, 6.79 equiv), and the enone **DRS6** (12.2 mg, 0.0249 mmol, 1.00 equiv) in tetrahydrofuran (1 mL) at $-95\text{ }^{\circ}\text{C}$ dropwise via cannula. The light red mixture was allowed to warm to $-50\text{ }^{\circ}\text{C}$ over 50 min and was then partitioned between an aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 5.0 mL) and dichloromethane (25 mL). The organic phase was separated and the aqueous phase was further extracted with dichloromethane ($3 \times 15\text{ mL}$). The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was decanted and concentrated, affording a yellow solid. The product was purified by preparatory HPLC on a Coulter Ultrasphere ODS column ($10\text{ }\mu\text{m}$, $250 \times 10\text{ mm}$, 3.5 mL/min , Solvent A: water, Solvent B: methanol, UV detection at 350 nm) using an injection volume of $500\text{ }\mu\text{L}$ methanol and a linear gradient elution of 85-100% B over 30 min. The peak at 21-27 min was collected and concentrated to give enol **JDB1-87** as a white solid (11.0 mg, 72%).

[00207] R_f 0.07 (3:7 ethyl acetate-hexanes); ^1H NMR (500 MHz, CD_2Cl_2) δ 15.21 (s, 1H, enol), 8.63 (d, 1H, $J = 4.5\text{ Hz}$, pyr-H), 8.19 (d, 1H, $J = 7.5\text{ Hz}$, pyr-H), 7.54-7.43 (m, 5H, ArH), 7.34 (d, 1H, $J = 4.5, 7.5\text{ Hz}$, pyr-H), 5.36 (d, 1H, $J = 12.0\text{ Hz}$, OCHH'Ph), 5.33 (d, 1H, $J = 12.0\text{ Hz}$, OCHH'Ph), 4.03 (d, 1H, $J = 10.7\text{ Hz}$, CHN(CH₃)₂), 3.36-3.31 (m, 1H, CHCH₂CHCHN(CH₃)₂), 3.23 (dd, 1H, $J = 16.3, 5.6\text{ Hz}$, CHH'CHCH₂CHCHN(CH₃)₂), 2.99 (dd, 1H, $J = 16.3, 16.3\text{ Hz}$, CHH'CHCH₂CHCHN(CH₃)₂), 2.63 (ddd, 1H, $J = 1.6, 4.4, 10.7\text{ Hz}$, CHCHN(CH₃)₂), 2.54-2.48 (m, 7H, N(CH₃)₂, CHH'CHCHN(CH₃)₂), 2.19 (dd, 1H, $J = 1.6, 14.5\text{ Hz}$, CHH'CHCHN(CH₃)₂), 0.87 (s, 9H, TBS), 0.26 (s, 3H, TBS), 0.13 (s, 3H, TBS); ^{13}C NMR (100 MHz, CD_2Cl_2) δ 187.7, 183.5, 182.6, 182.2, 167.9, 161.2, 153.4, 137.6, 134.1, 129.2, 129.1, 129.1, 126.8, 123.0, 108.7, 106.9, 82.2, 73.0, 61.8, 47.0, 42.1, 41.4, 30.1, 28.4, 26.1, 23.2, 19.3, $-2.4, -3.5$; HRMS (ES) m/z calcd for $(\text{C}_{33}\text{H}_{39}\text{N}_3\text{O}_6\text{Si}+\text{H})^+$ 602.2686, found 602.2686.



[00208] Pd black (3.0 mg, 0.028 mmol, 2.6 equiv) was added in one portion to a solution of the enol **JDB1-87** (6.5 mg, 0.011 mmol, 1.0 equiv) in dioxane-methanol (7:2, 9.0 mL) at 23 °C. An atmosphere of hydrogen was introduced by briefly evacuating the flask, then flushing with pure hydrogen (1 atm). The green mixture was stirred for 7 hr, and then filtered through a plug of cotton. The filtrate was concentrated, providing the carboxamide as a yellow oil (7.0 mg). Aqueous hydrofluoric acid (48%, 0.5 mL) was added to a polypropylene reaction vessel containing a solution of the carboxamide in acetonitrile (4.5 mL) at 23 °C. The reaction mixture was heated to 35 °C and was stirred at that temperature for 27 hr. The excess hydrofluoric acid was quenched with methoxytrimethylsilane (3.5 mL, 25 mmol). The reaction mixture was concentrated, affording a yellow solid. The product was purified by preparatory HPLC on a Phenomenex Polymerx DVB column (10 μ m, 250 $\times$ 10 mm, 4 mL/min, Solvent A: 0.5% trifluoroacetic acid in water, Solvent B: 0.5% trifluoroacetic acid in methanol-acetonitrile (1:1), UV detection at 350 nm) using an injection volume of 500 μ L methanol and a linear gradient of 0-20% B over 40 min. The peak at 35-45 min was collected and concentrated to give a yellow oil. The oil was dissolved in 1 mL methanol, treated with concentrated hydrochloric acid (20 μ L), and then concentrated to give the hydrochloride of **JDB1-109** as a yellow powder (3.7 mg, 86%).

[00209] ^1H NMR (500 MHz, CD_3OD , hydrochloride) δ 8.79-8.77 (m, 2H, pyr-H) 7.91 (dd, 1H, J = 6.8, 6.8 Hz, pyr-H), 4.12 (s, 1H, $\text{CHN}(\text{CH}_3)_2$), 3.41-3.22 (m, 2H, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$, $\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 3.11-3.00 (m, 8H, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$, $\text{CHCHN}(\text{CH}_3)_2$, $\text{N}(\text{CH}_3)_2$), 2.34 (ddd, 1H, J = 12.9, 4.4, 2.4 Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 1.77 (ddd, 1H, J = 12.9, 12.9, 12.9 Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$); HRMS (ES) m/z calcd for $(\text{C}_{20}\text{H}_{21}\text{N}_3\text{O}_6+\text{H})^+$ 400.1508, found

Example 6-Synthesis of 10-Deoxysancycline

[00210] *N,N*-dimethylformamide (20 μL) was added to a solution of the carboxylic acid **JDB1-113-SM** (500 mg, 3.67 mmol, 1.0 equiv) and oxalyl chloride (367 μL , 4.22 mmol, 1.15 equiv) in dichloromethane (20 mL) at 23 $^{\circ}\text{C}$. Vigorous gas evolution was observed. After stirring for 80 min at 23 $^{\circ}\text{C}$, phenol (863 mg, 9.18 mmol, 2.5 equiv), pyridine (890 μL , 11.0 mmol, 3.0 equiv), and dimethylaminopyridine (3 mg) were added in sequence. The resulting solution was stirred for 90 min at 23 $^{\circ}\text{C}$, whereupon an aqueous potassium phosphate buffer solution (pH 7.05, 0.2 M, 5.0 mL) was added. The resulting mixture was partitioned between water (30 mL) and ethyl acetate (50 mL). The aqueous phase was extracted with an additional 50-mL portion of ethyl acetate. The organic layers were combined and washed with an aqueous sodium hydroxide solution (50 mL, 1M), brine (50 mL), and then dried over anhydrous sodium sulfate. The dried solution was decanted and concentrated, affording a colorless oil (850 mg). The product was purified by flash column chromatography (25:75 ethyl acetate-hexanes), providing the ester **JDB1-113** as a colorless oil (774 mg, 99%).

[00211] R_f 0.43 (3:7 ethyl acetate-hexanes); ^1H NMR (300 MHz, CDCl_3) δ 8.18 (d, 1H, $J = 8.1$ Hz, ArH), 7.49-7.20 (m, 8H, ArH, OArH), 2.69 (s, 3H, ArCH₃); ^{13}C NMR (100 MHz, CDCl_3) δ 165.8, 150.9, 141.3, 132.7, 132.0, 131.2, 129.5, 128.5, 125.9, 125.8, 121.8, 22.0; FTIR (neat film), cm^{-1} 3046 (w), 2923 (w), 1739 (s), 1594 (m), 1487 (m), 1287 (m), 1241 (s), 1189 (s), 1159 (m), 1041 (s), 733 (s); HRMS (ES) m/z calcd for $(\text{C}_{14}\text{H}_{12}\text{O}_2 + \text{NH}_4)^+$ 230.1181, found 230.1187.

deoxysancycline hydrochloride (**JDB1-130**·HCl) as a white powder (9.1 mg, 91%).

[00215] ^1H NMR (500 MHz, CD_3OD , hydrochloride) δ 7.96 (d, 1H, $J = 7.3$ Hz, ArH) 7.51 (dd, 1H, $J = 7.3, 7.3$ Hz, ArH), 7.39 (dd, 1H, $J = 7.3, 7.3$ Hz, ArH), 7.30 (d, 1H, $J = 7.3$ Hz, ArH), 4.04 (s, 1H, $\text{CHN}(\text{CH}_3)_2$), 3.31-2.99 (m, 8H, $\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$, $\text{CHCHN}(\text{CH}_3)_2$, $\text{N}(\text{CH}_3)_2$), 2.87 (dd, 1H, $J = 15.4, 4.3$ Hz, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.61 (dd, 1H, $J = 15.4, 15.4$ Hz, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.21 (ddd, $J = 12.8, 5.0, 2.5$ Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 1.66 (ddd, 1H, $J = 12.8, 12.8, 12.8$ Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$).

Example 7-A Convergent, Enantioselective Synthetic Route to Structurally Diverse 6-Deoxytetracycline Antibiotics

[00216] Among tetracyclines, semi-synthetic approaches have led to the discovery of the 6-deoxytetracyclines doxycycline (**2** in Figure 15A) and minocycline (**3** in Figure 15A), clinically the most important agents in the class. 6-Deoxytetracyclines exhibit considerably improved chemical stability as compared to their 6-hydroxy counterparts and show equal or greater potencies in antibacterial assays (Stephens *et al.*, *J. Am. Chem. Soc.* **85**, 2643 (1963); M. Nelson, W. Hillen, R. A. Greenwald, Eds., *Tetracyclines in Biology, Chemistry and Medicine* (Birkhauser Verlag, Boston, 2001); each of which is incorporated herein by reference). It is evident that at present neither semi-synthesis nor modified biosynthesis is capable of addressing the great majority of novel structures that a chemist might wish to explore in pursuit of a lead structure like tetracycline; structures such as the D-ring heterocyclic analogs **4** and **5** in Figure 15A, or new ring systems such as the pentacycline **6** (Figure 15A) are exemplary. Absent a viable laboratory synthetic pathway, these structures and the regions of complex chemical space they represent must be ceded in the search for new antibiotics. Here, we report a short and efficient route for the synthesis of enantiomerically pure members of the 6-deoxytetracyclines from benzoic acid. The route we describe allows for the synthesis of 6-deoxytetracyclines (both with or without an hydroxyl group at C5) by a notably late-stage coupling reaction of the AB precursors **7** or **8** (Figure 15B) with a variety of different D-ring precursors, and has provided

compounds such as doxycycline (**2** in Figure 15A), the heterocyclic analogs **4** and **5** (Figure 15A), the pentacycline **6** (Figure 15A), as well as other 6-deoxytetracycline analogs.

[00217] The strategic advantage of a synthetic approach involving a late-stage C-ring construction ($AB + D \rightarrow ABCD$, Figure 15B) is that much of the polar functionality known to play a role in the binding of tetracyclines to the bacterial ribosome lies within the AB fragment (D. E. Brodersen *et al.*, *Cell* **103**, 1143 (2000); M. Pioletti *et al.*, *EMBO J.* **20**, 1829 (2001); each of which is incorporated herein by reference), while enormous structural variation on or near the D-ring is not only permissible, but has been cited as a means to overcome bacterial resistance. The advanced clinical candidate tigecycline (P.-E. Sum, P. Petersen, *Bioorg. Med. Chem. Lett.* **9**, 1459 (1999); incorporated herein by reference), a minocycline derivative with a D-ring substituent, is exemplary, and is reported to be one of the most promising new antibiotics under evaluation by the FDA (K. Bush, M. Macielag, M. Weidner-Wells, *Curr. Opin. Microbiol.* **7**, 466 (2004); incorporated herein by reference). Classically, approaches to the synthesis of the tetracycline antibiotics have proceeded by stepwise assembly of the ABCD ring system and begin with D or CD precursors, as exemplified by the Woodward synthesis of ($\pm$)-6-deoxy-6-demethyltetracycline (sancycline, 25 steps, ~0.002% yield) (J. J. Korst *et al.*, *J. Am. Chem. Soc.* **90**, 439 (1968); incorporated herein by reference), the Shemyakin synthesis of ($\pm$)-12a-deoxy-5a,6-anhydrotetracycline (A. I. Gurevich *et al.*, *Tetrahedron Lett.* **8**, 131 (1967); incorporated herein by reference), and the Muxfeldt synthesis of ($\pm$)-5-oxytetracycline (terramycin, 22 steps, 0.06% yield) (H. Muxfeldt *et al.*, *J. Am. Chem. Soc.* **101**, 689 (1979); incorporated herein by reference). Only one published synthesis of (–)-tetracycline itself has appeared, this from D-glucosamine (an A-ring precursor, 34 steps, 0.002% yield) (K. Tatsuta *et al.*, *Chem. Lett.* 646 (2000); incorporated herein by reference), while the most efficient construction of the tetracycline ring system thus far is undoubtedly the synthesis of ($\pm$)-12a-deoxytetracycline by the Stork laboratory (16 steps, 18-25% yield) (G. Stork *et al.*, *J. Am. Chem. Soc.* **118**, 5304 (1996); incorporated herein by reference). The latter research served to identify C12a oxygenation as perhaps the greatest challenge in tetracycline synthesis (it could not be achieved with

12a-deoxytetracycline as substrate), a conclusion supported by the results of prior synthetic efforts (J. J. Korst *et al.*, *J. Am. Chem. Soc.* **90**, 439 (1968); A. I. Gurevich *et al.*, *Tetrahedron Lett.* **8**, 131 (1967); H. Muxfeldt *et al.*, *J. Am. Chem. Soc.* **101**, 689 (1979); each of which is incorporated herein by reference). The problem is significant, for C12a oxygenation appears to greatly enhance antimicrobial activity (W. Rogalski, in *Handbook of Experimental Pharmacology*, J. J. Hlavka, J. H. Boothe, Eds. (Springer-Verlag, New York, 1985), vol. 78, chap. 5; incorporated herein by reference). A key feature of the synthetic approach to 6-deoxytetracyclines that we have developed is that it introduces the C12a hydroxyl group in the first step of the sequence (Figure 16) and uses the stereogenic center produced in that step to elaborate all others in the target molecule. To protect the vinylogous carbamic acid function of the A-ring we used the 5-benzyloxyisoxazole group developed by Stork and Hagedorn for that purpose (G. Stork, A. A. Hagedorn, III, *J. Am. Chem. Soc.* **100**, 3609 (1978); incorporated herein by reference), an innovation that proved critically enabling in the present work, while the dimethylamino group of the A-ring was incorporated without modification.

[00218] Our synthesis of 6-deoxytetracyclines was initiated by whole-cell, microbial dihydroxylation of benzoic acid with a mutant strain of *Alcaligenes eutrophus* (A. M. Reiner, G. D. Hegeman, *Biochemistry* **10**, 2530 (1971); A. G. Myers *et al.*, *Org. Lett.* **3**, 2923 (2001); each of which is incorporated herein by reference), producing the diol **9** (Figure 16) with >95% ee in 79% yield (90-g batch, ~13 g/L, Figure 16). Hydroxyl-directed epoxidation of the microcrystalline product (**9**, *m*-CPBA, EtOAc) provided the α -oriented epoxide **10** (Figure 16) in 83% yield; esterification of this product (trimethylsilyldiazomethane) followed by bis-silylation and concomitant epoxide isomerization in the presence of *tert*-butyldimethylsilyl triflate (3 equiv.), afforded the epoxy ester **11** (Figure 16) in 70% yield (A. G. Myers *et al.*, *Org. Lett.* **3**, 2923 (2001); incorporated herein by reference). Separately, 3-benzyloxy-5-dimethylaminomethylisoxazole, prepared on the mole-scale by a simple four-step sequence from glyoxylic acid (D. M. Vyas, Y. Chiang, T. W. Doyle, *Tetrahedron Lett.* **25**, 487 (1984); P. Pevarello, M. Varasi, *Synth. Commun.* **22**, 1939 (1992); each of which is incorporated herein by reference), was deprotonated at C4

with *n*-butyllithium, and the resulting organolithium reagent (**12** in Figure 16) was then added to the epoxy ester **11** (Figure 16), forming the ketone **13** (73%) (Figure 16). In a noteworthy transformation, and a key step of the synthesis, exposure of the ketone **13** (Figure 16) to lithium triflate (5 mol %) at 60 °C, followed by selective removal of the allylic silyl ether of the rearranged product (TFA), afforded the tricyclic AB precursor **14** (Figure 16) in 62% yield after purification by flash column chromatography. The transformation of **13** to **14** (Figure 16) is believed to involve initial S_N-prime opening of the allylic epoxide by the *N,N*-dimethylamino group followed by ylide formation and [2,3]-sigmatropic rearrangement, a process that is reminiscent of the Sommelet-Hauser rearrangement (S. H. Pine, *Organic Reactions*, **18**, 403 (1970); incorporated herein by reference). Compound **14** (Figure 16) possesses the requisite *cis* stereochemistry of the AB fusion as well as an α -oriented *N,N*-dimethylamino substituent (confirmed by X-ray crystallographic analysis of a derivative), and serves as a common intermediate for the synthesis of both the AB precursor enone **7** (4 steps, 49% yield, Figure 16) and the AB precursor to 5- α -hydroxy-6-deoxytetracyclines, enone **8** (8 steps, 56% yield, Figure 16), as detailed in sequence below.

[00219] To synthesize the AB precursor enone **7** (Figure 15), intermediate **14** was subjected to reductive transposition (A. G. Myers, B. Zheng, *Tetrahedron Lett.* **37**, 4841 (1996); incorporated herein by reference) in the presence of triphenylphosphine, diethyl azodicarboxylate, and *o*-nitrobenzenesulfonyl hydrazide (added last, a procedural variant), affording the transposed cycloalkene **15** in 74% yield. Hydrolysis of the silyl ether group within **15** (HCl, methanol), oxidation of the resulting allylic alcohol (IBX, DMSO) (M. Frigerio, M. Santagostino, *Tetrahedron Lett.* **35**, 8019 (1994); incorporated herein by reference), and protection of the remaining (tertiary) carbinol (TBSOTf, 2,6-lutidine) (E. J. Corey *et al.*, *Tetrahedron Lett.* **22**, 3455 (1981); incorporated herein by reference) then provided the enone **7** (Figure 15) in 66% yield (3 steps) after flash column chromatography. By a somewhat longer but slightly more efficient sequence the intermediate **14** (Figure 15) could also be transformed into the enone **8** (Figure 15), the AB precursor to 5- α -hydroxy-6-deoxytetracyclines. This sequence involved the transformation of **14** (Figure 15) into the phenylthio ether **16** (with net retention), diastereoselective sulfoxidation using a chiral oxidant (F. A. Davis

et al., *J. Org. Chem.* **57**, 7274 (1992); incorporated herein by reference) (99:1 selectivity), and Mislow-Evans rearrangement (E. N. Prilezhaeva, *Russ. Chem. Rev.* **70**, 897 (2001); incorporated herein by reference), producing the allylic alcohol **17** in 66% yield (4 steps). High diastereoselectivity in the sulfoxidation step was essential, for only one diastereomer (the major isomer under the conditions specified) underwent efficient thermal rearrangement. After protection of the allylic alcohol **17** (Figure 15) using benzyl chloroformate, a sequence nearly identical to the final three steps of the synthesis of **7** (Figure 15) was employed to transform the resulting benzyl carbonate into the enone **8** (Figure 15) in 85% yield (56% yield and 8 steps from **14**).

[00220] 6-Deoxytetracyclines were assembled with all requisite functionality and stereochemistry in a single operation. In this process the AB precursors **7** or **8** (Figure 15) are coupled with a range of different carbanionic D-ring precursors in a Michael-Dieckmann reaction sequence (T.-L. Ho, *Tandem Organic Reactions* (Wiley, New York, 1992); incorporated herein by reference) that forms two carbon-carbon bonds and the C-ring of the 6-deoxytetracyclines (Figures 15B, 17, and 18). The process is perhaps best illustrated in detail by the 3-step synthesis of (–)-doxycycline from the AB precursor **8** (Figure 17). Deprotonation of the D-ring precursor **18** (4.5 equiv, LDA, TMEDA, THF, –78 °C), synthesized in 5 steps (42% yield) from anisic acid, followed by addition of the enone **8** (1 equiv, –78 → 0 °C), provided the tetracyclic coupling product **19** (Figure 17) in diastereomerically pure form in 79% yield after purification by rp-HPLC. Removal of the protective groups (2 steps, 90% yield) and purification (rp-HPLC) afforded (–)-doxycycline hydrochloride (18 steps, 8.3% yield from benzoic acid). A remarkable feature of the convergent coupling reaction that produces the tetracyclic product **19** (Figure 17) is its stereoselectivity. Although in theory four diastereomeric products can be formed, largely one was produced, corresponding in configuration (5a*R*, 6*R*) to that of known biologically active 6-deoxytetracyclines. A minor diastereomeric impurity, believed to be 6-epi-**19** (Figure 17), was also isolated in separate rp-HPLC fractions (<7% yield). Michael-Dieckmann cyclization sequences (T.-L. Ho, *Tandem Organic Reactions* (Wiley, New York, 1992); incorporated herein reference) and condensations of *o*-toluate anions in particular (F. J. Leeper, J. Staunton, *J.C.S. Chem. Comm.*, 406 (1978); F. M. Hauser, R. P. Rhee, *J. Org. Chem.* **43**, 178,

(1978); J. H. Dodd, S. M. Weinreb, *Tetrahedron Lett.* **20**, 3593 (1979); each of which is incorporated herein by reference) are extensively precedented in synthesis, but we are unaware of any example exhibiting the high degree of diastereoselectivity of the present case. Phenyl ester activation in toluate condensations is also precedented, though in a system that forms a fully aromatized cyclization product (White *et al.*, *J. Org. Chem.* **51**, 1150 (1986); incorporated herein by reference). We observed that the presence of the phenyl ester group of the D-ring precursor **18** (Figure 17) was essential for successful cyclization to occur; anions derived from simple alkyl esters and phthalide-derived anions underwent Michael addition, but the resulting adducts did not cyclize. Perhaps even more remarkable than the condensation that produces **19** (Figure 17) is the parallel transformation of **18** with the enone **7** (Figure 18, entry 1), which forms (–)-6-deoxytetracycline in protected form with >20:1 diastereoselectivity, in 81% yield after purification by rp-HPLC (diastereomerically pure; a minor diastereomer, epimeric at C6, was also isolated separately). It appears that additions to **7** and **8** proceed almost exclusively by addition to the “top” face of each enone (as drawn), producing C5a-stereochemistry corresponding to natural tetracyclines, though why this should be the case is not obvious.

[00221] As the examples of entries 2-5 (Figure 18) show, efficient and stereoselective condensations are not restricted to the *o*-toluate anion derived from the D-ring substrate **18** (Figure 17); the novel D-ring heterocyclic analogs **4** and **5** (Figure 18) were synthesized by a related sequence from *o*-toluate anions of very different structures, as was the pentacycline derivative **6** (Figure 18). In each case it was necessary to optimize the specific conditions for *o*-toluate anion generation and trapping. For entries 3-5 (Figure 18) anion generation was best conducted *in situ*, in the presence of the enone **7**, either by selective deprotonation (entry 3) or by lithium-halogen exchange (entries 4 and 5). A number of potentially competing non-productive reaction sequences (*e.g.*, enolization of **7**) might have occurred during *in situ* anion generation; the observed efficiencies of the transformations are surprising in light of this. It is also noteworthy that *in situ* anion generation permits the use of *o*-toluates lacking an *o*-alkoxy substituent (entries 3 and 4), substrates known to be problematic from prior studies (F. M. Hauser *et al.*, *Synthesis* 72 (1980); incorporated herein by

reference). Finally, *o*-toluate anion formation by *in situ* or stepwise halogen-metal exchange (entries 4 and 5) is unprecedented.

[00222] The efficiencies of the synthetic sequences have allowed for the preparation of sufficient quantities of each tetracycline analog for antibacterial testing using standard serial-dilution techniques (5-20 mg amounts). Minimum inhibitory concentrations (MICs) are reported for each analog in whole-cell antimicrobial assays using five Gram-positive and five Gram-negative organisms (Figure 18). Thus far, the pentacycline derivative **6** (Figure 18) has shown the most promising antibacterial properties, with activity equal to or greater than tetracycline in each of the Gram-positive strains examined, including strains with resistance to tetracycline, methicillin, and vancomycin.

Experimentals

[00223] **General Procedures.** All reactions were performed in flame-dried round bottomed or modified Schlenk (Kjeldahl shape) flasks fitted with rubber septa under a positive pressure of argon, unless otherwise noted. Air- and moisture-sensitive liquids and solutions were transferred via syringe or stainless steel cannula. Organic solutions were concentrated by rotary evaporation at ~25 Torr (house vacuum). Flash column chromatography was performed on silica gel (60 Å, standard grade) as described by Still *et al.* (Still, W. C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, *43*, 2923-2925; incorporated herein by reference). Analytical thin-layer chromatography was performed using glass plates pre-coated with 0.25 mm 230-400 mesh silica gel impregnated with a fluorescent indicator (254 nm). Thin-layer chromatography plates were visualized by exposure to ultraviolet light and/or exposure to ceric ammonium molybdate or an acidic solution of *p*-anisaldehyde followed by heating on a hot plate.


[00224] **Materials.** Commercial reagents and solvents were used as received with the following exceptions. Triethylamine, diisopropylamine, *N,N,N',N'*-tetramethylethylene-diamine, DMPU, HMPA, and *N,N*-diisopropylethylamine were distilled from calcium hydride under an atmosphere of dinitrogen. Dichloromethane, methanol, tetrahydrofuran, acetonitrile, and toluene were purified by the method of Pangborn *et al.* (Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.;

Timmers, F. J. *Organometallics* **1996**, *15*, 1518-1520; incorporated herein by reference).

[00225] Instrumentation. Proton nuclear magnetic resonance (^1H NMR) spectra and carbon nuclear magnetic resonance (^{13}C NMR) spectra were recorded with Varian Unity/Inova 600 (600 MHz), Varian Unity/Inova 500 (500 MHz/125 MHz), or Varian Mercury 400 (400 MHz/100 MHz) NMR spectrometers. Chemical shifts for protons are reported in parts per million (δ scale) and are referenced to residual protium in the NMR solvents (CHCl_3 : δ 7.26, $\text{C}_6\text{D}_5\text{H}$: δ 7.15, D_2HCO : δ 3.31, CDHCl_2 : δ 5.32, $(\text{CD}_2\text{H})\text{CD}_3\text{SO}$: δ 2.49). Chemical shifts for carbon are reported in parts per million (δ scale) and are referenced to the carbon resonances of the solvent (CDCl_3 : δ 77.0, C_6D_6 : δ 128.0, CD_3OD : δ 44.9, CD_2Cl_2 : δ 53.8, $(\text{CD}_3)_2\text{SO}$: δ 39.5). Data are represented as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br = broad), integration, coupling constant in Hz, and assignment. Infrared (IR) absorbance spectra were obtained using a Perkin-Elmer 1600 FT-IR spectrophotometer referenced to a polystyrene standard. Data are represented as follows: frequency of the absorption (cm^{-1}), intensity of the absorption (s = strong, m = medium, w = weak, br = broad), and assignment (where appropriate). Optical rotations were determined using a JASCO DIP-370 digital polarimeter equipped with a sodium lamp source. High-resolution mass spectra were obtained at the Harvard University Mass Spectrometry Facilities.

Synthesis of (–)-Doxycycline

Cyclization Step:




[00226] A solution of *n*-butyllithium in hexanes (1.55 M, 155 μ L, 0.240 mmol, 5.1 equiv) was added to a solution of *N,N,N',N'*-tetramethylethylenediamine (39 μ L, 0.26 mmol, 5.5 equiv) and diisopropylamine (34 μ L, 0.25 mmol, 5.1 equiv) in tetrahydrofuran (1 mL) at -78 $^{\circ}$ C. The resulting mixture was stirred vigorously at -78 $^{\circ}$ C for 30 min whereupon a solution of 2-(phenoxycarbonyl)-3-ethylphenyl *t*-butyl carbonate (73.0 mg, 0.213 mmol, 4.5 equiv) in tetrahydrofuran (1 mL) was added dropwise via cannula. The resulting deep-red mixture was stirred vigorously at -78 $^{\circ}$ C for 75 min, then a solution of enone **8** (30.0 mg, 0.0474 mmol, 1 equiv) in tetrahydrofuran (1 mL) was added dropwise via cannula. The resulting light-red mixture was allowed to warm slowly to 0 $^{\circ}$ C over 2 h. The ice-cold product solution was then partitioned between aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 10 mL) and dichloromethane (10 mL). The organic phase was separated and the aqueous phase was further extracted with two 10-mL portions of dichloromethane. The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a yellow oil. The product was purified by preparatory HPLC on a Coulter Ultrasphere ODS column [10 μ m, 250 x 10 mm, UV detection at 350 nm, injection volume: 400 μ L (methanol), isocratic elution with methanol-water (9:1), flow rate: 3.5 mL/min]. Fractions eluting at 36-42 min were collected and concentrated, affording the pentacyclic addition product depicted in diastereomerically pure form (33.0 mg, 79%, a light-yellow solid).

[00227] R_f 0.35 (1:4 ethyl acetate-hexanes); ^1H NMR (500 MHz, C_6D_6) δ 16.55 (br s, 1H, enol), 7.26 (d, 2H, $J = 7.0$ Hz, *o*-ArH), 7.14 (d, 2H, $J = 7.5$ Hz, ArH), 6.85-

7.05 (m, 6H, ArH), 6.66-6.74 (m, 2H, ArH), 6.51 (dd, 1H, $J = 9.0, 1.5$ Hz, ArH), 5.73 (br d, 1H, $J = 4.0$ Hz, BnOCO₂CH), 5.17 (d, 1H, $J = 12.5$ Hz, OCHH'Ph), 5.03 (d, 1H, $J = 12.5$ Hz, OCHH'Ph), 4.99 (d, 1H, $J = 12.5$ Hz, OCHH'Ph'), 4.93 (d, 1H, $J = 12.5$ Hz, OCHH'Ph'), 3.58 (d, 1H, $J = 11.5$ Hz, CHCHN(CH₃)₂), 3.35 (dd, 1H, $J = 12.5, 4.0$ Hz, CH₃CHCH), 2.99 (d, 1H, $J = 11.5$ Hz, CHCHN(CH₃)₂), 2.56 (dq, 1H, $J = 12.5, 7.0$ Hz, CH₃CH), 2.18 (s, 6H, N(CH₃)₂), 1.33 (s, 9H, C(CH₃)₃), 1.16 (d, 3H, $J = 7.0$ Hz, CH₃CH), 1.11 (s, 9H, C(CH₃)₃), 0.61 (s, 3H, CH₃), 0.36 (s, 3H, CH₃); ¹³C NMR (100 MHz, CDCl₃) δ 189.7, 186.3, 180.9, 178.4, 167.9, 154.7, 152.1, 150.8, 145.9, 136.1, 135.5, 133.9, 128.7, 128.6, 128.5, 127.3, 123.8, 122.7, 122.6, 108.9, 105.5, 83.0, 82.9, 74.8, 72.4, 69.2, 60.8, 52.7, 43.2, 38.4, 27.5, 26.6, 19.5, 16.3, -1.8, -2.7; FTIR (neat film), cm⁻¹ 2974 (w), 2933 (w), 2851 (w), 1760 (s, C=O), 1748 (s, C=O), 1723 (s, C=O), 1606 (m), 1513 (m), 1471 (m), 1370 (m), 1260 (s), 1232 (s), 1148 (s); HRMS (ES) m/z calcd for (C₄₈H₅₆ N₂O₁₂Si)⁺ 881.3681, found 881.3684.


Deprotection Step 1:



[00228] Concentrated aqueous hydrofluoric acid (48 wt %, 1.2 mL) was added to a polypropylene reaction vessel containing a solution of the purified pentacyclic addition product from the experiment above (33.0 mg, 0.0375 mmol, 1 equiv) in acetonitrile (7.0 mL) at 23 °C. The resulting mixture was stirred vigorously at 23 °C for 60 h, then was poured into water (50 mL) containing dipotassium hydrogenphosphate (7.0 g). The resulting mixture was extracted with ethyl acetate (3 × 20 mL). The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, affording the product depicted as a yellow oil (25.0 mg, 100%). This product was used in the next step without further purification.

[00229] *R_f* 0.05 (1:4 ethyl acetate-hexanes); ¹H NMR (600 MHz, C₆D₆, crude) δ 14.86 (br s, 1H, enol), 11.95 (s, 1H, phenol), 7.23 (d, 2H, *J* = 7.8 Hz, *o*-ArH), 7.14 (d, 2H, *J* = 7.2 Hz, *o*-ArH), 6.94-7.02 (m, 6H, ArH), 6.86 (t, 1H, *J* = 8.4 Hz, ArH), 6.76 (d, 1H, *J* = 8.4 Hz, ArH), 6.28 (d, 1H, *J* = 7.8 Hz, ArH), 5.46 (dd, 1H, *J* = 3.6, 3.0 Hz, BnOCO₂CH), 5.12 (d, 1H, *J* = 12.0 Hz, OCHH'Ph), 5.04 (d, 1H, *J* = 12.0 Hz, OCHH'Ph), 4.92 (s, 2H, OCH₂Ph), 3.41 (d, 1H, *J* = 9.6 Hz, CHCHN(CH₃)₂), 2.82 (dd, 1H, *J* = 9.6, 3.0 Hz, CHCHN(CH₃)₂), 2.65 (dd, 1H, *J* = 13.2, 3.6 Hz, CH₃CHCH), 2.78 (dq, 1H, *J* = 13.2, 7.2 Hz, CH₃CH), 2.05 (s, 6H, N(CH₃)₂), 1.04 (d, 3H, *J* = 7.2 Hz, CH₃CH); ¹³C NMR (100 MHz, C₆D₆, crude) δ 193.4, 186.2, 181.3, 172.3, 167.9, 163.3, 154.6, 145.8, 136.6, 135.8, 128.6, 128.4, 127.2, 116.8, 116.0, 115.6, 107.6, 104.7, 76.8, 73.9, 72.5, 69.5, 60.3, 48.7, 43.0, 41.8, 37.5, 15.3; FTIR (neat film), cm⁻¹ 3424 (m, OH), 3059, 3030, 2925, 2857, 1744 (s, C=O), 1713 (s, C=O), 1614 (s), 1582 (s), 1455 (s), 1252 (s); HRMS (ES) *m/z* calcd for (C₃₇H₃₄N₂O₁₀+H)⁺ 667.2292, found 667.2300.

Deprotection Step 2:




[00230] Palladium black (7.00 mg, 0.0657 mmol, 1.75 equiv) was added in one portion to a solution of the product from the procedure above (25.0 mg, 0.0375 mmol, 1 equiv) in tetrahydrofuran-methanol (1:1, 2.0 mL) at 23 °C. An atmosphere of hydrogen was introduced by briefly evacuating the flask, then flushing with pure hydrogen (1 atm). The palladium catalyst was initially observed to be a fine dispersion, but aggregated into clumps within 5 min. The yellow heterogeneous mixture was stirred at 23 °C for 2 h, then was filtered through a plug of cotton. The filtrate was concentrated, affording a yellow oil. The product was purified by preparatory HPLC on a Phenomenex Polymerx DVB column (10 μ m, 250 x 10 mm, UV detection at 350 nm, Solvent A: methanol-0.005 N aq. HCl (1:4), Solvent B: acetonitrile, injection volume: 400 μ L (solvent A containing 10 mg oxalic acid), isochratic elution with 5% B for 2 min, then gradient elution with 5 $\rightarrow$ 50% B for 20 min, flow rate: 4.0 mL/min]. Fractions eluting at 12-17 min were collected and concentrated, affording (-)-doxycycline hydrochloride as a yellow powder (16.2 mg, 90%), which was identical with natural (-)-doxycycline hydrochloride [reverse-phase HPLC (co-injection), ^1H NMR (including measurement of an admixture of synthetic and natural doxycycline), ^{13}C NMR, $[\alpha]_{\text{D}}$, UV).

[00231] ^1H NMR (600 MHz, CD_3OD , hydrochloride) δ 7.47 (t, 1H, J = 8.4 Hz, ArH), 6.93 (d, 1H, J = 8.4 Hz, ArH), 6.83 (d, 1H, J = 8.4 Hz, ArH), 4.40 (s, 1H, $(\text{CH}_3)_2\text{NCH}$), 3.53 (dd, 1H, J = 12.0, 8.4 Hz, CHOH), 2.95 (s, 3H, $\text{N}(\text{CH}_3)\text{CH}_3'$), 2.88 (s, 3H, $\text{N}(\text{CH}_3)\text{CH}_3'$), 2.80 (d, 1H, J = 12.0 Hz, CHCHN $(\text{CH}_3)_2$), 2.74 (dq, 1H, J = 12.6, 6.6 Hz, CH_3CH), 2.58 (dd, 1H, J = 12.6, 8.4 Hz, CH_3CHCH), 1.55 (d, 3H, J = 6.6 Hz, CH_3CHCH); ^{13}C NMR (100 MHz, CD_3OD) δ 195.3, 188.2, 173.8, 172.1, 163.2, 149.0, 137.7, 117.1, 116.9, 116.6, 108.4, 96.0, 74.5, 69.8, 66.9, 47.5, 43.4, 43.0, 41.9, 40.0, 16.3; UV max (0.01 M methanolic HCl), nm 218, 267, 350; $[\alpha]_{\text{D}} = -109^\circ$ (c = 0.16 in 0.01 M methanolic HCl); HRMS (ES) m/z calcd for $(\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_8 + \text{H})^+$ 445.1611, found 445.1603.

[00232] Literature values (*The Merck Index: An Encyclopedia of Chemicals, Drugs, and Biologicals*, 12th ed. Budavari, S.; O'Neal, M. J.; Smith, A.; Heckelman, P. E.; Kinneary, J. F., Eds.; Merck & Co.: Whitehouse Station, NJ, 1996; entry 3496.): UV max (0.01 M methanolic HCl), nm 267, 351; $[\alpha]_D = -110^\circ$ (c = 1 in 0.01 M methanolic HCl).

Synthesis of (–)-6-Deoxytetracycline

Cyclization Step:




[00233] A solution of *n*-butyllithium in hexanes (1.65 M, 75 μ L, 0.12 mmol, 3.9 equiv) was added to a solution of diisopropylamine (17 μ L, 0.12 mmol, 3.9 equiv) and *N,N,N',N'*-tetramethylethylenediamine (19 μ L, 0.13 mmol, 4.1 equiv) in tetrahydrofuran (1 mL) at -78°C . The resulting solution was stirred at -78°C for 30 min whereupon a solution of 2-(phenoxycarbonyl)-3-ethylphenyl *t*-butyl carbonate (31.8 mg, 0.093 mmol, 3.0 equiv) in tetrahydrofuran (250 μ L) was added dropwise via syringe. The resulting deep-red mixture was stirred at -78°C for 90 min, then a solution of enone 7 (15.0 mg, 0.031 mmol, 1 equiv) in tetrahydrofuran (250 μ L) was added dropwise via syringe. The resulting deep-red mixture was allowed to warm slowly to 0°C over 3 h. The ice-cold product solution was then partitioned between aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 15 mL) and dichloromethane (15 mL). The organic phase was separated and the aqueous phase was further extracted with two 15-mL portions of dichloromethane. The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a yellow oil. The product was purified by preparatory HPLC on a Coulter Ultrasphere ODS column [5 μ m, 250 $\times$ 10

mm, UV detection at 350 nm, injection volume: 500 μ L (methanol), isocratic elution with methanol-water (89:11), flow rate: 3.5 mL/min]. Fractions eluting at 39-60 min were collected and concentrated, affording the pentacyclic addition product depicted in diastereomerically pure form (18.5 mg, 81%, a light-yellow foam).

[00234] R_f 0.37 (2:8 tetrahydrofuran-hexanes); ^1H NMR (500 MHz, CDCl_3) δ (s, 1H, 16.24, enol-OH), 7.55-7.50 (m, 3H, ArH), 7.40-7.35 (m, 4H, ArH), 7.10 (d, 1H, J = 7.8 Hz, ArH), 5.39-5.34 (m, 2H, OCH_2Ph), 3.92 (d, 1H, J = 10.7 Hz, $\text{CHN}(\text{CH}_3)_2$), 2.81-2.71 (m, 2H, CH_3CH , CH_3CHCH), 2.55 (dd, 1H, J = 10.7, 5.7 Hz, $\text{CHCHN}(\text{CH}_3)_2$), 2.48 (s, 6H, $\text{N}(\text{CH}_3)_2$), 2.40 (d, 1H, J = 14.7 Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 2.31 (ddd, 1H, J = 14.7, 9.3, 5.7, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 1.56 (s, 3H, CH_3), 1.55 (s, 9H, Boc), 0.84 (s, 9H, TBS), 0.27 (s, 3H, TBS), 0.13 (s, 3H, TBS); ^{13}C NMR (125 MHz, CDCl_3) δ 187.4, 183.1, 182.8, 181.6, 167.6, 151.7, 150.2, 147.4, 135.0, 134.0, 128.5, 128.5, 123.4, 123.0, 122.4, 108.3, 107.4, 94.8, 83.9, 81.5, 72.5, 61.5, 46.4, 41.9, 39.5, 34.9, 27.7, 26.0, 20.7, 19.0, 16.0, -2.6, -3.7; FTIR (neat film), cm^{-1} 2923 (m), 2841 (m), 1759 (s, C=O), 1718 (s, C=O), 1605 (s), 1508 (s), 1467 (m), 1456 (m), 1369 (m), 1277 (s), 1262 (m), 1231 (s), 1144 (s), 1005 (w); HRMS (ES) m/z calcd for $(\text{C}_{40}\text{H}_{50}\text{N}_2\text{O}_9\text{Si}+\text{H})^+$ 731.3364, found 731.3370.

Deprotection:




[00235] Concentrated aqueous hydrofluoric acid solution (48 wt %, 0.6 mL) was added to a polypropylene reaction vessel containing a solution of the purified pentacyclic addition product from the experiment above (15.0 mg, 0.0205 mmol, 1 equiv) in acetonitrile (3.5 mL) at 23 $^{\circ}\text{C}$. The reaction mixture was stirred at 23 $^{\circ}\text{C}$ for 55 h, then was poured into water (20 mL) containing dipotassium hydrogenphosphate (4.0 g). The resulting mixture was extracted with ethyl acetate (4×20 mL). The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, affording a light-yellow oil. The residue was dissolved in methanol-tetrahydrofuran (1:1, 2 mL) and to the resulting

solution was added palladium black (7.6 mg, 0.071 mmol, 3.5 equiv) in one portion. An atmosphere of hydrogen gas was introduced by briefly evacuating the flask, then flushing with pure hydrogen (1 atm). The yellow mixture was stirred at 23 °C for 2 h, then was filtered through a plug of cotton. The filtrate was concentrated, affording a yellow oil (10 mg). The product was purified by preparatory HPLC on a Phenomenex Polymerx DVB column [10 μ m, 250 $\times$ 10 mm, UV detection at 365 nm, Solvent A: methanol-0.02 N HCl (1:4), Solvent B: acetonitrile, injection volume: 400 μ L (methanol containing 10 mg oxalic acid), isocratic elution with 18% B for 15 min, then gradient elution with 18 $\rightarrow$ 60% B over 15 min, flow rate: 5 mL/min]. Fractions eluting at 17.5-22.5 min were collected and concentrated, affording 6-deoxytetracycline hydrochloride as a yellow powder (8.1 mg, 85%).

[00236] ^1H NMR (500 MHz, CD_3OD , hydrochloride) δ 7.49 (t, 1H, $J = 7.8$ Hz, ArH), 6.95 (d, 1H, $J = 7.8$ Hz, ArH), 6.84 (d, 1H, $J = 7.8$ Hz, ArH), 4.09 (s, 1H, $\text{CHN}(\text{CH}_3)_2$), 3.03 (br s, 3H, $\text{N}(\text{CH}_3)_2$), 2.97 (br s, 3H, $\text{N}(\text{CH}_3)_2$), 2.90 (br d, 1H, $J = 12.7$ Hz, $\text{CHCHN}(\text{CH}_3)_2$), 2.67 (ddd, 1H, $J = 12.7, 12.7, 5.2$ Hz, CH_3CHCH), 2.61-2.56 (m, 1H, CH_3CH), 2.30 (ddd, 1H, $J = 13.7, 5.2, 2.9$ Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 1.54 (ddd, 1H, $J = 13.7, 12.7, 12.7$ Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 1.38 (d, 3H, $J = 6.8$ Hz, CH_3CH); UV max (0.01 M methanolic HCl), nm 269, 353; $[\alpha]_{\text{D}} = -142^\circ$ ($c = 0.20$ in 0.01 M methanolic HCl); HRMS (ES) m/z calcd for $(\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_7 + \text{H})^+$ 429.1662, found 429.1660.

Synthesis of a (–)-D-ring Pyridone Analog of Tetracycline


Cyclization Step:



[00237] A solution of *n*-butyllithium in hexanes (1.67 M, 80 μ L, 0.13 mmol, 4.3 equiv) was added to a solution of diisopropylamine (20 μ L, 0.14 mmol, 4.6 equiv) in tetrahydrofuran (2.5 mL) at -78 $^{\circ}$ C. The resulting solution was allowed to warm to 0 $^{\circ}$ C over 15 min. *N,N*-dimethylpropyleneurea (17 μ L, 0.14 mmol, 4.5 equiv) was added and the resulting solution was cooled to -78 $^{\circ}$ C. A solution of phenyl 2-(benzyloxy)-4,6-dimethylpyridine-3-carboxylate (31.0 mg, 0.0930 mmol, 2.99 equiv) in tetrahydrofuran (250 μ L) was then added via syringe to the cooled reaction solution. The resulting yellow solution was stirred for 5 min at -78 $^{\circ}$ C, then a solution of enone 7 (15.0 mg, 0.0311 mmol, 1 equiv) in tetrahydrofuran (250 μ L) was added via syringe. The resulting deep-red mixture was allowed to warm to 0 $^{\circ}$ C over 4 h. Acetic acid (40 μ L) was added to the deep-red mixture at 0 $^{\circ}$ C. The ice-cold product solution was then partitioned between aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 15 mL) and dichloromethane (15 mL). The organic phase was separated and the aqueous phase was further extracted with two 15-mL portions of dichloromethane. The organic extracts were combined and then dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, providing a yellow oil. The product was purified by preparatory HPLC on a Coulter Ultrasphere ODS column [5 μ m, 250 $\times$ 10 mm, UV detection at 350 nm, Solvent A: water, Solvent B: methanol, injection volume: 500 μ L DMSO, gradient elution with 92 $\rightarrow$ 100% B over 30 min, flow rate: 3.5 mL/min]. Fractions eluting at 21–29 min were collected and concentrated, affording the pentacyclic addition product depicted in diastomerically pure form (15.0 mg, 67%, a light-yellow solid).

[00238] R_f 0.55 (3:7 ethyl acetate-hexanes); ^1H NMR (600 MHz, CD_2Cl_2) δ 16.05 (s, 1H, enol-OH), 7.52-7.26 (m, 10H, ArH), 6.66 (s, 1H, pyr-H), 5.57 (d, 1H, J = 12.7 Hz, OCHH'Ph), 5.43 (d, J = 12.7 Hz, 1H, OCHH'Ph), 5.33-5.28 (m, 2H, OCH₂Ph), 3.99 (d, 2H, J = 10.5 Hz, CHN(CH₃)₂), 3.04-3.00 (m, 1H, CHCH₂CHCHN(CH₃)₂), 2.84 (dd, 1H, J = 16.1, 4.9 Hz, CHH'CHCH₂CHCHN(CH₃)₂), 2.74 (dd, 1H, J = 16.1, 16.1 Hz, CHH'CHCH₂CHCHN(CH₃)₂), 2.53 (dd, 1H, J = 10.5, 3.9 Hz, CHCHN(CH₃)₂), 2.51-2.43 (m, 10H, N(CH₃)₂, Ar-CH₃, CHH'CHCHN(CH₃)₂), 2.07 (d, 1H, J = 14.2 Hz, CHH'CHCHN(CH₃)₂), 0.82 (s, 9H, TBS), 0.22 (s, 3H, TBS), 0.10 (s, 3H, TBS); ^{13}C NMR (100 MHz, CD_2Cl_2) δ 187.9, 185.2, 182.5, 178.8, 167.9, 161.9, 161.8, 154.8, 137.9, 135.6, 129.1, 129.0, 129.0, 128.7, 127.9, 127.9, 116.4, 111.6, 108.6, 107.5, 82.0, 73.0, 68.1, 61.7, 46.9, 42.0, 39.2, 28.6, 26.1, 24.6, 23.0, 19.3, -2.4, -3.5; FTIR (neat film), cm^{-1} 2939 (m), 2857 (w), 1720 (s, C=O), 1593 (s), 1510 (s), 1469 (m), 1449 (m), 1326 (s), 1254 (m), 1187 (w), 1157 (m), 1090 (m), 1064 (m), 1007 (m); HRMS (ES) m/z calcd for $(\text{C}_{41}\text{H}_{47}\text{N}_3\text{O}_7\text{Si}+\text{H})^+$ 722.3262, found 722.3261.

Deprotection:




[00239] Pearlman's catalyst (10 mg, 0.0094 mmol, 0.68 equiv) was added to a solution of the purified pentacyclic addition product from the experiment above (10 mg, 0.014 mmol, 1 equiv) in dioxane-methanol (1:1, 10 mL) at 23 °C. An atmosphere of hydrogen gas was introduced by briefly evacuating the flask, then flushing with pure hydrogen (1 atm). The reaction mixture was observed to form a green color within 10 min. After stirring at 23 °C for 2 h, the reaction mixture was filtered through a plug of cotton and the filtrate was concentrated. The oily yellow residue was dissolved in methanol (10 mL) and to the resulting solution was added concentrated aqueous hydrochloric acid solution (37 wt %, 100 μL) at 23 °C. The reaction mixture was stirred at 23 °C for 3 h, then was concentrated. The product was purified by preparatory HPLC on a Phenomenex Polymerx DVB column [10 μm , 250 $\times$ 10 mm, UV detection at 365 nm, Solvent A: 0.01 N aqueous hydrochloric acid, Solvent B: acetonitrile, injection volume: 500 μL (methanol containing 30 mg oxalic acid), linear gradient with 0 $\rightarrow$ 20% B over 40 min, flow rate: 4 ml/min]. Fractions eluting at 20-29 min were collected and concentrated, affording the D-ring pyridone hydrochloride as a yellow powder (4.8 mg, 74%).

[00240] ^1H NMR (500 MHz, CD_3OD , hydrochloride) δ 6.37 (s, 1H, ArH), 4.06 (s, 1H, $\text{CHN}(\text{CH}_3)_2$), 3.05-2.95 (m, 8H, $\text{N}(\text{CH}_3)_2$, $\text{CHCHN}(\text{CH}_3)_2$, $\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.79 (dd, 1H, $J = 16.1, 3.9$ Hz, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.55 (dd, 1H, $J = 16.1, 16.1$ Hz, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.40 (s, 3H, Ar- CH_3), 2.18 (br. D, 1H, $J = 12.7$ Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 1.59 (ddd, 1H, $J = 12.7, 12.7, 12.7$ Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$); ^{13}C NMR (100 MHz, $(\text{CD}_3)_2\text{SO}$) δ 187.3, 183.5, 177.8, 172.1, 160.6, 159.8, 153.3, 115.3, 107.2, 106.9, 95.6, 74.2, 68.4, 41.5, 35.7, 34.5, 33.9, 31.0, 19.2; UV max (0.01 M methanolic HCl), nm 267, 370; $[\alpha]_D = -146^\circ$ ($c = 0.43$ in 0.01 M methanolic HCl); HRMS (ES) m/z calcd for $(\text{C}_{21}\text{H}_{23}\text{N}_3\text{O}_7 + \text{H})^+$ 430.1614, found 430.1607.

Synthesis of a (–)-Pentacycline

Cyclization Step:




[00241] A solution of *n*-butyllithium in hexanes (2.65 M, 107 μ L, 0.284 mmol, 4.03 equiv) was added to a solution of phenyl 3-(bromomethyl)-1-methoxynaphthalene-2-carboxylate (105 mg, 0.283 mmol, 4.02 equiv) and enone **7** (34.0 mg, 0.0705 mmol, 1 equiv) in tetrahydrofuran (2.80 mL) at $-100\text{ }^{\circ}\text{C}$. The resulting light-red reaction mixture was allowed to warm to $0\text{ }^{\circ}\text{C}$ over 70 min. The ice-cold product solution was then partitioned between aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 15 mL) and dichloromethane (15 mL). The organic phase was separated and the aqueous phase was further extracted with two 15-mL portions of dichloromethane. The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered, and the filtrate was concentrated, affording a yellow solid. The product was purified by preparatory HPLC on a Coulter Ultrasphere ODS column [10 μ m, 250×10 mm, UV detection at 350 nm, Solvent A: water, Solvent B: methanol, two separate injections (750 μ L each, acetonitrile), isocratic elution with 94% B for 20 min followed by a linear gradient elution with 94 $\rightarrow$ 100% B over 20 min, flow rate: 3.5 mL/min]. Fractions eluting at 24–38 min were collected and concentrated, affording the hexacyclic addition product in diastereomerically pure form (36.1 mg, 75%, a white solid).

[00242] R_f 0.37 (3:7 ethyl acetate-hexanes); ^1H NMR (500 MHz, CDCl_3) δ 16.25 (s, 1H, enol-OH), 8.30 (d, 1H, $J = 8.3$ Hz, ArH), 7.75 (d, 1H, $J = 7.8$ Hz, ArH), 7.59–7.34 (m, 7H, ArH), 7.26 (s, 1H, ArH), 5.38 (s, 2H, OCH_2Ph), 4.02 (s, 3H, OCH_3), 3.99 (d, 1H, $J = 10.7$ Hz, $\text{CHN}(\text{CH}_3)_2$), 3.08–3.05 (m, 2H, $\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$, 2.95–2.90 (m, 1H, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.58 (dd, 1H, $J = 10.7, 5.9$ Hz, $\text{CHCHN}(\text{CH}_3)_2$), 2.51 (s, 6H, $\text{N}(\text{CH}_3)_2$), 2.50–2.48 (m, 1H, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 2.20–2.14 (m, 1H, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 0.82 (s, 9H, TBS), 0.29 (s, 3H, TBS), 0.13 (s, 3H, TBS); ^{13}C NMR (125 MHz, CDCl_3) δ 187.9, 184.1,

183.0, 182.0, 167.8, 159.2, 137.5, 136.7, 135.3, 129.5, 128.8, 128.7, 128.5, 127.5, 126.4, 124.2, 121.8, 119.5, 108.7, 108.7, 82.4, 72.8, 63.8, 61.6, 46.8, 42.1, 40.7, 29.3, 26.2, 23.1, 19.3, -2.2, -3.5; FTIR (neat film), cm^{-1} 2934 (m), 2852 (m), 1718 (s, C=O), 1610 (s), 1513 (s), 1472 (m), 1452 (m), 1369 (m), 1339 (w), 1293 (m), 1252 (m), 1190 (w), 1159 (m), 1067 (m), 1026 (w), 1011 (w); HRMS (ES) m/z calcd for $(\text{C}_{39}\text{H}_{44}\text{N}_2\text{O}_7\text{Si}+\text{H})^+$ 681.2996, found 681.2985.

Deprotection:




[00243] Concentrated aqueous hydrofluoric acid solution (48 wt %, 1.0 mL) was added to a polypropylene reaction vessel containing a solution of the purified hexacyclic addition product from the experiment above (24.0 mg, 0.035, 1 equiv) in acetonitrile (9.0 mL) at 23 °C. The reaction mixture was stirred at 23 °C for 22 h, then was poured into water (50 mL) containing dipotassium hydrogenphosphate (12.0 g). The resulting mixture was extracted with ethyl acetate (3 × 50 mL). The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, affording a yellow oil. The residue was dissolved in methanol-dioxane (1:1, 5 mL) and to the resulting solution was added palladium black (10.0 mg, 0.0940 mmol, 2.67 equiv) in one portion. An atmosphere of hydrogen gas was introduced by briefly evacuating the flask, then flushing with pure hydrogen (1 atm). The yellow mixture was stirred at 23 °C for 4 h, then was filtered through a plug of cotton. The filtrate was concentrated, affording a yellow oil. The residue was dissolved in dichloromethane (4.5 mL) and to the resulting solution was added a solution of boron tribromide (1.0 M in dichloromethane, 0.5 mL, 14 equiv) at -78 °C. The dark-red mixture was stirred at -78 °C for 15 min, then at 23 °C for 3.5 h. Methanol (20 mL) was added and the resulting yellow solution was stirred at 23 °C for 1 h. The solution was concentrated, affording a yellow oil. The product was purified by preparatory HPLC on a Phenomenex Polymerx DVB column [7 μm , 150 × 21.2

mm, UV detection at 350 nm, Solvent A: 0.01 N HCl, Solvent B: acetonitrile, injection volume: 500 μ L (methanol containing 10 mg oxalic acid), gradient elution with 25 $\rightarrow$ 50% B over 60 min, flow rate: 6 mL/min]. Fractions eluting at 30-35 min were collected and concentrated, affording the pentacycline hydrochloride as a yellow powder (13.1 mg, 74%).

[00244] ^1H NMR (600 MHz, CD_3OD , hydrochloride) δ 8.36 (d, 1H, $J = 7.7$ Hz, ArH), 7.74 (d, 1H, $J = 7.7$ Hz, ArH), 7.64 (dd, 1H, $J = 7.7, 7.7$ Hz, ArH), 7.50 (dd, 1H, $J = 7.7, 7.7$ Hz, ArH), 7.1 (s, 1H, ArH), 4.10 (s, 1H, $\text{CHN}(\text{CH}_3)_2$), 3.13-2.97 (m, 9H, $\text{N}(\text{CH}_3)_2$, $\text{CHCHN}(\text{CH}_3)_2$, $\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.67 (dd, 1H, $J = 14.3, 14.3$ Hz, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.22 (ddd, 1H, $J = 13.6, 4.9, 2.9$ Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 1.64 (ddd, 1H, $J = 13.6, 13.6, 13.6$ Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$); UV max (0.01 M methanolic HCl), nm 268, 345, 402; $[\alpha]_{\text{D}} = -113^\circ$ ($c = 0.18$ in 0.01 M methanolic HCl); HRMS (ES) m/z calcd for $(\text{C}_{25}\text{H}_{24}\text{N}_2\text{O}_7 + \text{H})^+$ 465.1662, found 465.1656.

Synthesis of (–)-7-Aza-10-Deoxysancycline

Cyclization Step:




[00245] A solution of *n*-butyllithium in hexanes (2.65 M, 33.0 μ L, 0.0945 mmol, 5.00 equiv) was added to a solution of diisopropylamine (13.2 μ L, 0.0945 mmol, 5.00 equiv) in tetrahydrofuran (0.750 mL) at -78°C . The resulting solution was briefly warmed in an ice bath (10 min), then was cooled to -78°C .

Hexamethylphosphoramide (33.0 μ L, 0.189 mmol, 10.0 equiv) was added, producing a colorless solution, and this solution was then transferred (cold) dropwise via cannula to a solution containing phenyl 2-methylpyridine-3-carboxylate (16.0 mg, 0.0755 mmol, 4.00 equiv) and enone 7 (9.1 mg, 0.019 mmol, 1 equiv) in tetrahydrofuran (0.750 mL) at -95°C , forming a light-red mixture. The reaction solution was allowed to warm to -50°C over 50 min. The product solution was then partitioned between aqueous

potassium phosphate buffer solution (pH 7.0, 0.2 M, 10 mL) and dichloromethane (25 mL). The organic phase was separated and the aqueous phase was further extracted with three 15-mL portions of dichloromethane. The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, affording a yellow solid. The product was purified by preparatory HPLC on a Coulter Ultrasphere ODS column [10 μ m, 250 $\times$ 10 mm, UV detection at 350 nm, Solvent A: water, Solvent B: methanol, injection volume: 500 μ L (methanol), gradient elution of 85 $\rightarrow$ 100% B over 30 min, flow rate: 3.5 mL/min]. Fractions eluting at 21-27 min were collected and concentrated, affording the pentacyclic addition product in diastereomerically pure form (8.6 mg, 76%, a white solid).

[00246] R_f 0.07 (3:7 ethyl acetate-hexanes); ^1H NMR (500 MHz, CD_2Cl_2) δ 15.21 (s, 1H, enol), 8.63 (d, 1H, J = 4.5 Hz, pyr-**H**), 8.19 (d, 1H, J = 7.5 Hz, pyr-**H**), 7.54-7.43 (m, 5H, Ar**H**), 7.34 (d, 1H, J = 4.5, 7.5 Hz, pyr-**H**), 5.36 (d, 1H, J = 12.0 Hz, OCH**H**'Ph), 5.33 (d, 1H, J = 12.0 Hz, OCH**H**'Ph), 4.03 (d, 1H, J = 10.7 Hz, CHN(CH₃)₂), 3.36-3.31 (m, 1H, CHCH₂CHCHN(CH₃)₂), 3.23 (dd, 1H, J = 16.3, 5.6 Hz, CH**H**'CHCH₂CHCHN(CH₃)₂), 2.99 (dd, 1H, J = 16.3, 16.3 Hz, CH**H**'CHCH₂CHCHN(CH₃)₂), 2.63 (ddd, 1H, J = 1.6, 4.4, 10.7 Hz, CHCHN(CH₃)₂), 2.54-2.48 (m, 7H, N(CH₃)₂, CH**H**'CHCHN(CH₃)₂), 2.19 (dd, 1H, J = 1.6, 14.5 Hz, CH**H**'CHCHN(CH₃)₂), 0.87 (s, 9H, TBS), 0.26 (s, 3H, TBS), 0.13 (s, 3H, TBS); ^{13}C NMR (100 MHz, CD_2Cl_2) δ 187.7, 183.5, 182.6, 182.2, 167.9, 161.2, 153.4, 137.6, 134.1, 129.2, 129.1, 129.1, 126.8, 123.0, 108.7, 106.9, 82.2, 73.0, 61.8, 47.0, 42.1, 41.4, 30.1, 28.4, 26.1, 23.2, 19.3, -2.4, -3.5; HRMS (ES) m/z calcd for (C₃₃H₃₉N₃O₆Si+H)⁺ 602.2686, found 602.2686.


Deprotection:



[00247] Palladium black (3.0 mg, 0.028 mmol, 2.6 equiv) was added in one portion to a solution of the purified pentacyclic addition product from the experiment above (6.5 mg, 0.011 mmol, 1 equiv) in dioxane-methanol (7:2, 9.0 mL) at 23 °C. An atmosphere of hydrogen was introduced by briefly evacuating the flask, then flushing with pure hydrogen (1 atm). The resulting green mixture was stirred at 23 °C for 7 hr, then was filtered through a plug of cotton. The filtrate was concentrated, affording a yellow oil (7.0 mg). The residue was dissolved in acetonitrile (4.5 mL), transferred to a polypropylene reaction vessel, and concentrated aqueous hydrofluoric acid solution (48 wt %, 0.5 mL) was added to the resulting solution at 23 °C. The reaction mixture was heated to 35 °C for 27 hr. Excess hydrofluoric acid was quenched by the addition of methoxytrimethylsilane (3.5 mL, 25 mmol). The reaction mixture was concentrated, affording a yellow solid. The product was purified by preparatory HPLC on a Phenomenex Polymerx DVB column [10 μ m, 250 $\times$ 10 mm, UV detection at 350 nm, Solvent A: 0.5% trifluoroacetic acid in water, Solvent B: 0.5% trifluoroacetic acid in methanol-acetonitrile (1:1), injection volume: 500 μ L (methanol), gradient elution with 0 $\rightarrow$ 20% B over 40 min, flow rate: 4 mL/min]. Fractions eluting at 35-45 min were collected and concentrated to give a yellow oil. The oil was dissolved in methanolic HCl (1.0 mL, 0.10 M) and concentrated, affording 7-aza-10-deoxysancycline hydrochloride as a yellow powder (3.7 mg, 79%). ^1H NMR (500 MHz, CD_3OD , hydrochloride) δ 8.79-8.77 (m, 2H, pyr-**H**) 7.91 (dd, 1H, J = 6.8, 6.8 Hz, pyr-**H**), 4.12 (s, 1H, $\text{CHN}(\text{CH}_3)_2$), 3.41-3.22 (m, 2H, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), $\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$, 3.11-3.00 (m, 8H, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$, $\text{CHCHN}(\text{CH}_3)_2$, $\text{N}(\text{CH}_3)_2$), 2.34 (ddd, 1H, J = 12.9, 4.4, 2.4 Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 1.77 (ddd, 1H, J = 12.9, 12.9, 12.9 Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$); UV max (0.01 M methanolic HCl), nm 264, 345; $[\alpha]_D = -154^\circ$ (c = 0.15 in 0.01 M methanolic HCl); HRMS (ES) m/z calcd for $(\text{C}_{20}\text{H}_{21}\text{N}_3\text{O}_6 + \text{H})^+$ 400.1508, found 400.1504.

Synthesis of (–)-10-Deoxysancycline

Cyclization Step:




[00248] A solution of *n*-butyllithium in hexanes (2.65 M, 59 μ L, 0.16 mmol, 4.0 equiv) was added to a solution of phenyl 2-(bromomethyl)benzoate (45.6 mg, 0.157 mmol, 3.97 equiv) and enone **7** (19.0 mg, 0.0394 mmol, 1 equiv) in tetrahydrofuran (1.57 mL) at $-100\text{ }^{\circ}\text{C}$. The resulting light-red solution was allowed to warm to $0\text{ }^{\circ}\text{C}$ over 30 min. The ice-cold product solution was then partitioned between aqueous potassium phosphate buffer solution (pH 7.0, 0.2 M, 5 mL) and dichloromethane (25 mL). The organic phase was separated and the aqueous phase was further extracted with an additional 15-mL portion of dichloromethane. The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, affording a yellow solid. The product was purified by preparatory HPLC on a Coulter Ultrasphere ODS column [10 μ m, 250×10 mm, Solvent A: water, Solvent B: methanol, injection volume: 1.0 mL (methanol), gradient elution with 85 $\rightarrow$ 100% B over 30 min, UV detection at 350 nm, flow rate: 3.5 mL/min]. Fractions eluting at 25-30 min were collected and concentrated, affording the pentacyclic addition product in diastereomerically pure form (19.2 mg, 81%, a white solid).

[00249] R_f 0.46 (3:7 ethyl acetate-hexanes); ^1H NMR (500 MHz, CD_2Cl_2) δ 15.53 (s, 1H, enol), 7.94 (d, 1H, $J = 7.9$ Hz, ArH), 7.54 - 7.28 (m, 8H, ArH, OCH_2ArH), 5.37-5.34 (m, 2H, OCH_2Ph), 4.05 (d, 1H, $J = 10.7$ Hz, $\text{CHN}(\text{CH}_3)_2$), 3.24-3.18 (m, 1H, $\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.99 (dd, 1H, $J = 15.5, 5.6$ Hz, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.88 (dd, 1H, $J = 15.5, 15.5$ Hz, $\text{CHH}'\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$), 2.61 (dd, 1H, $J = 4.4, 10.7$ Hz, $\text{CHCHN}(\text{CH}_3)_2$), 2.54-2.44 (m, 7H, $\text{N}(\text{CH}_3)_2$, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 2.14 (d, 1H, $J = 14.3$ Hz, $\text{CHH}'\text{CHCHN}(\text{CH}_3)_2$), 0.86 (s, 9H, TBS), 0.25 (s, 3H, TBS), 0.12 (s, 3H, TBS); ^{13}C NMR (100 MHz, CD_2Cl_2) δ 187.8, 183.0, 182.8, 182.4, 167.7, 141.7, 135.4, 133.4,

130.9, 129.0, 128.9, 128.9, 128.1, 127.5, 126.5, 108.5, 106.8, 82.1, 72.8, 61.5, 58.5, 46.9, 41.9, 38.6, 29.0, 25.9, 23.1, 19.1, -2.6, -3.7; HRMS (ES) m/z calcd for $(C_{34}H_{40}N_3O_6Si+H)^+$ 601.2734, found 601.2730.

Deprotection:



[00250] Concentrated aqueous hydrofluoric acid solution (48 wt %, 1.1 mL) was added to a polypropylene reaction vessel containing a solution of the pentacyclic addition product from the experiment above (15.1 mg, 0.0251 mmol, 1 equiv) in acetonitrile (10 mL) at 23 °C. The resulting solution was stirred vigorously at 23 °C for 12 h, then was poured into water (50 mL) containing dipotassium hydrogenphosphate (4.7 g) and the product was extracted with ethyl acetate (3 × 25 mL). The organic phases were combined and dried over anhydrous sodium sulfate. The dried solution was filtered and the filtrate was concentrated, affording a yellow solid (12.2 mg, 99%). The residue was dissolved in methanol-dioxane (1:1, 3.0 mL) and palladium black (6.5 mg, 0.061 mmol, 2.4 equiv) was added to the resulting solution in one portion. An atmosphere of hydrogen was introduced by briefly evacuating the flask, then flushing with pure hydrogen (1 atm). The resulting light-yellow mixture was stirred at 23 °C for 20 min, then was filtered through a plug of cotton. The filtrate was concentrated, affording a yellow solid. The product was purified by preparatory HPLC on a Phenomenex Polymerx DVB column [10 μm, 250 × 10 mm, UV detection at 350 nm, Solvent A: 0.01 N HCl, Solvent B: acetonitrile, injection volume: 1.0 mL (methanol containing 10 mg oxalic acid), gradient elution with 5→50% B over 30 min, flow rate: 5 mL/min]. Fractions eluting at 16-22 min were collected and concentrated, affording 10-deoxysancycline hydrochloride as a white powder (9.1 mg, 83%).

[00251] ^1H NMR (500 MHz, CD_3OD , hydrochloride) δ 7.96 (d, 1H, $J = 7.3$ Hz, ArH) 7.51 (dd, 1H, $J = 7.3, 7.3$ Hz, ArH), 7.39 (dd, 1H, $J = 7.3, 7.3$ Hz, ArH), 7.30 (d, 1H, $J = 7.3$ Hz, ArH), 4.04 (s, 1H, $\text{CHN}(\text{CH}_3)_2$), 3.31-2.99 (m, 8H, $\text{CHCH}_2\text{CHCHN}(\text{CH}_3)_2$, $\text{CHCHN}(\text{CH}_3)_2$, $\text{N}(\text{CH}_3)_2$), 2.87 (dd, 1H, $J = 15.4, 4.3$ Hz,

CHH'CHCH₂CHCHN(CH₃)₂), 2.61 (dd, 1H, $J = 15.4, 15.4$ Hz, CHH'CHCH₂CHCHN(CH₃)₂), 2.21 (ddd, $J = 12.8, 5.0, 2.5$ Hz, CHH'CHCHN(CH₃)₂), 1.66 (ddd, 1H, $J = 12.8, 12.8, 12.8$ Hz, CHH'CHCHN(CH₃)₂); UV max (0.01 M methanolic HCl), nm 264, 348; $[\alpha]_D = -147^\circ$ ($c = 0.15$ in 0.01 M methanolic HCl); HRMS (ES) m/z calcd for (C₂₁H₂₂N₂O₆+H)⁺ 399.1556, found 399.1554.

Biological testing.

[00252] Whole-cell antibacterial activity was determined according to methods recommended by the NCCLS (National Committee for Clinical Laboratory Standards. 2002. *Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically: approved standard-fifth edition*. NCCLS document M100-S12. National Committee for Clinical Laboratory Standards. Wayne, PA.; incorporated herein by reference). Test compounds were dissolved in dimethyl sulfoxide (DMSO) and the resulting solutions were diluted in water (1:10) to produce stock solutions with a final concentration of 256 µg tetracycline analog per mL. In a 96-well microtiter plate, 50-µL aliquots of stock solutions were diluted serially into cation-adjusted Mueller-Hinton broth (MHB; Becton-Dickinson, Cockeysville, MD). Test organisms (50 µL aliquots of solutions $\sim 5 \times 10^5$ CFU/mL) were then added to the appropriate wells of the microtiter plate. Inoculated plates were incubated aerobically at 35 °C for 18-24 h. The MIC was the lowest concentration of compound determined to inhibit visible growth. Five Gram-positive and five Gram-negative bacterial strains were examined in minimum inhibitory concentration (MIC) assays. The Gram-positive strains were *Staphylococcus aureus* ATCC 29213, *Staphylococcus epidermidis* ACH-0016, *Staphylococcus haemolyticus* ACH-0013, *Enterococcus faecalis* ATCC 700802 (a VRE or vancomycin-resistant enterococcus strain), and *Staphylococcus aureus* ATCC 700699 (carrying the *tetM* resistance gene). The Gram-negative strains were *Pseudomonas aeruginosa* ATCC 27853, *Klebsiella pneumoniae* ATCC 13883, *E. coli* ATCC 25922, *E. coli* ACH-0095 (multiply antibiotic-resistant), and *E. coli* ATCC 53868::pBR322 (containing a plasmid encoding tetracycline-resistance). These strains are listed again below, along with certain other details of their origins and known resistance to antibiotics.

Bacterial strains

Gram-Positive Organisms:

<i>Staphylococcus aureus</i> ATCC 29213	QC strain for MIC testing
<i>Staphylococcus aureus</i> ATCC 700699	Methicillin- and tetracycline-resistant clinical isolate with intermediate resistance to vancomycin
<i>Staphylococcus epidermidis</i> ACH-0018	Clinical isolate (Achillion strain collection)
<i>Staphylococcus haemolyticus</i> ACH-0013	Clinical isolate (Achillion strain collection)
<i>Enterococcus faecalis</i> ATCC 700802	Vancomycin-resistant clinical isolate

Gram-Negative Organisms:

<i>E. coli</i> ATCC 25922	QC strain for MIC testing
<i>E. coli</i> ATCC 53868::pBR322	Laboratory strain carrying a plasmid with a tetracycline-resistance marker
<i>E. coli</i> ACH-0095	Multiply-resistant clinical isolate (Achillion strain collection)

Klebsiella pneumoniae ATCC 13883

QC strain for MIC testing


Pseudomonas aeruginosa ATCC 27853

QC strain for MIC testing

ATCC = American Type Culture Collection, Manassas, VA

Example 8-Alternative Routes to Tetracycline Analogs

[00253] Many of the studies described above show the generation of the carbanionic D-ring precursor by metalization of phenyl esters of *o*-toluate derivatives. These self-condensation reactions at times required to use of up to 4-5 equivalents of a given D-ring precursor. The presence of an electron-withdrawing substituent on the α -carbon greatly improves the efficiency of metalation and coupling as described in Example 7 and elsewhere herein. Lithium-halogen exchange of benzylic bromides conducted *in situ* in the presence of the AB electrophile has been found to provide coupling products where benzylic metalation fails (see Example 7). These benzylic bromides can be prepared with surprising efficiencies (near quantitative yields) and are surprisingly stable. The developments may lead to a coupling reaction that could be conductable on a multi-kilo scale. Many different phenyl ester substituents (see below) may be used to optimize a coupling reaction.



The optimal group for benzylic metalation, however, may not be the same as the optimal group for lithium-halogen exchange. In addition, for the lithium-halogen exchange process, besides ester modification, other metal reagents may be used including, but not limited to, other alkyllithium reagents (*e.g.*, phenyllithium, mesityllithium), Grignard reagents (*e.g.*, *iso*-propylmagnesium chloride) and zinc-based systems. Barbier-type couplings will be explored using a variety of zero-valent metals for coupling.


The AB-ring precursors may also be prepared by alternative routes. The step-count for the synthesis of most 6-deoxytetracycline analogs is 14 from benzoic acid. Eleven of these 14 steps are dedicated to the synthesis of the AB-ring precursor. Any improvements in the length or efficiency of the route to these AB-ring precursors will have a substantial impact on the synthesis overall. Alternative syntheses of the AB-ring precursor are shown in Figures 22 and 23. Among the strategies for alternative A-ring closure sequences are intramolecular Michael additions, palladium-mediated processes, and iminium ion induce closures. Hypervalent iodine reagents may also be used instead of microbial dihydroxylation in the synthesis of the AB-ring precursors as shown in Figure 23.

Other Embodiments

[00254] The foregoing has been a description of certain non-limiting preferred embodiments of the invention. Those of ordinary skill in the art will appreciate that various changes and modifications to this description may be made without departing from the spirit or scope of the present invention, as defined in the following claims.

Claims

1. A method of preparing a compound of formula:



or a salt thereof,

wherein:

R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_A$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_2 is hydrogen;

R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_C$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_D$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio,

arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_5 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_E$; $-CN$; $-SCN$; $-SR_E$; or $-N(R_E)_2$; wherein each occurrence of R_E is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;


each occurrence of R_7 is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_G$; $-C(=O)R_G$; $-CO_2R_G$; $-CN$; $-SCN$; $-SR_G$; $-SOR_G$; $-SO_2R_G$; $-NO_2$; $-N(R_G)_2$; $-NHC(O)R_G$; or $-C(R_G)_3$; wherein each occurrence of R_G is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

each P' is independently hydrogen or a protecting group;

P is hydrogen, lower (C_1 - C_6) alkyl, acyl, or a protecting group; and

n is an integer in the range of 0 to 3, inclusive;

by reacting an anion of a toluate, wherein the toluate has the formula:




wherein R_9 is $-OR_I$, $-CN$, $-SCN$, $-SR_I$, or $-N(R_I)_2$; and each R_I is independently hydrogen; a protecting group; a cyclic or acyclic, substituted or unsubstituted aliphatic; a cyclic or acyclic, substituted or unsubstituted heteroaliphatic; a substituted or unsubstituted aryl; or a substituted or unsubstituted heteroaryl;

and wherein the anion is formed by deprotonation of the toluate under basic


conditions;

with an enone of formula:



wherein R₆ is hydrogen.

2. The method of claim 1, wherein R₁ is methyl.
3. The method of claim 1, wherein R₉ is -OR₁; and R₁ is a substituted or unsubstituted aryl; or a substituted or unsubstituted heteroaryl.
4. The method of claim 3, wherein R₁ is substituted or unsubstituted phenyl.
5. The method of claim 1, wherein P is a BOC protecting group.
6. The method of claim 1, wherein P is benzyl.
7. The method of claim 1, wherein P is C₁₋₆ alkyl.
8. A method of preparing a tetracycline of formula:



or a salt thereof,

wherein:

R₁ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched

heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_A$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_2 is hydrogen;

R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_C$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_D$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;


R_5 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_E$; $-CN$; $-SCN$; $-SR_E$; or $-N(R_E)_2$; wherein each occurrence of R_E is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

each occurrence of R_7 is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted

aryl; substituted or unsubstituted heteroaryl; $-OR_G$; $-C(=O)R_G$; $-CO_2R_G$; $-CN$; $-SCN$; $-SR_G$; $-SOR_G$; $-SO_2R_G$; $-NO_2$; $-N(R_G)_2$; $-NHC(O)R_G$; or $-C(R_G)_3$; wherein each occurrence of R_G is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio; and

n is an integer in the range of 0 to 3, inclusive;

by reacting an anion of a toluate, wherein the toluate has the formula:




wherein:

R_9 is $-OR_I$, $-CN$, $-SCN$, $-SR_I$, or $-N(R_I)_2$; and each R_I is independently hydrogen; a protecting group; a cyclic or acyclic, substituted or unsubstituted aliphatic; a cyclic or acyclic, substituted or unsubstituted heteroaliphatic; a substituted or unsubstituted aryl; or a substituted or unsubstituted heteroaryl; and

P is hydrogen, lower (C_1 - C_6) alkyl, acyl, or a protecting group;

wherein the anion is formed by deprotonation of the toluate under basic conditions;


with an enone of formula:



wherein:


R_6 is hydrogen and each P' is independently hydrogen or a protecting group;

to form a compound of formula:




or salt thereof; and

deprotecting the compound to provide a tetracycline of formula:



or salt thereof.

9. The method of claim 8, wherein R₁ is methyl.
10. The method of claim 8, wherein R₉ is -OR_I; and R_I is a substituted or unsubstituted aryl; or a substituted or unsubstituted heteroaryl.
11. The method of claim 10, wherein R_I is substituted or unsubstituted phenyl.
12. The method of claim 8, wherein P is a BOC protecting group.
13. The method of claim 8, wherein P is benzyl.
14. The method of claim 8, wherein P is C₁₋₆ alkyl.
15. A method of preparing a compound of formula:



or a salt thereof;

wherein:

R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_A$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_2 is hydrogen;

R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_C$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_D$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_5 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or

unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_E; -CN; -SCN; -SR_E; or -N(R_E)₂; wherein each occurrence of R_E is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;


R₆ is selected from the group consisting of hydrogen, halogen, substituted or unsubstituted aliphatic, substituted or unsubstituted heteroaliphatic, substituted or unsubstituted alkoxy, -OH, -CN, -SCN, -SH, alkylthio, -NO₂, amino, alkyl amino, dialkyl amino, and -SAr, wherein Ar is substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl;

each occurrence of R₇ is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_G; -C(=O)R_G; -CO₂R_G; -CN; -SCN; -SR_G; -SOR_G; -SO₂R_G; -NO₂; -N(R_G)₂; -NHC(O)R_G; or -C(R_G)₃; wherein each occurrence of R_G is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

each P' is independently hydrogen or a protecting group;


n is an integer in the range of 0 to 3, inclusive;

by reacting an enone of formula:




or salt thereof;

with a benzocyclobutenol of formula:




16. The method of claim 15, wherein the enone is of the formula:




or a salt thereof.

17. The method of claim 15, wherein the benzocyclobutenol is of the formula:



18. A method of preparing a tetracycline of formula:



or a salt thereof;

wherein:

R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_A$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; -

NHC(O)R_A; or -C(R_A)₃; wherein each occurrence of R_A is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R₂ is hydrogen;

R₃ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_C; -C(=O)R_C; -CO₂R_C; -CN; -SCN; -SR_C; -SOR_C; -SO₂R_C; -NO₂; -N(R_C)₂; -NHC(O)R_C; or -C(R_C)₃; wherein each occurrence of R_C is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R₄ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_D; -C(=O)R_D; -CO₂R_D; -CN; -SCN; -SR_D; -SOR_D; -SO₂R_D; -NO₂; -N(R_D)₂; -NHC(O)R_D; or -C(R_D)₃; wherein each occurrence of R_D is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;


R₅ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_E; -CN; -SCN; -SR_E; or -N(R_E)₂; wherein each occurrence of R_E is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

each occurrence of R₇ is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_G; -C(=O)R_G; -CO₂R_G; -CN; -SCN; -SR_G; -SOR_G; -SO₂R_G; -NO₂; -N(R_G)₂; -NHC(O)R_G; or -C(R_G)₃; wherein each occurrence of R_G is

independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio; and


n is an integer in the range of 0 to 3, inclusive;

by reacting an enone of formula:




or salt thereof,

with a benzocyclobutenol of formula:




wherein:

R₆ is hydrogen, and each P' is independently hydrogen or a protecting group;
to form a compound of formula:




or salt thereof; and

oxidizing and deprotecting the compound to provide a tetracycline of formula:




or salt thereof.

19. The method of claim 20, wherein the enone is of the formula:




or a salt thereof.

20. The method of claim 20, wherein the benzocyclobutenol is of the formula:



21. A method of preparing a compound of formula:



or a salt thereof;

wherein:

R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted

heteroaryl; $-OR_A$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_2 is hydrogen;

R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_C$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_D$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;


R_5 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_E$; $-CN$; $-SCN$; $-SR_E$; or $-N(R_E)_2$; wherein each occurrence of R_E is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_6 is selected from the group consisting of hydrogen, halogen, substituted or unsubstituted aliphatic, substituted or unsubstituted heteroaliphatic, substituted or unsubstituted alkoxy, $-OH$, $-CN$, $-SCN$, $-SH$, alkylthio, $-NO_2$, amino, alkyl amino, dialkyl amino, and $-SAr$, wherein Ar is substituted or unsubstituted aryl, or substituted or


unsubstituted heteroaryl;

each P' is independently hydrogen or a protecting group; and


n is an integer in the range of 0 to 3, inclusive;



or salt thereof; with a diene of formula:




22. The method of claim 21, wherein the enone is of the formula:



or salt thereof.

23. The method of claim 21, wherein R₁ is C₁-C₆ alkyl.

24. A method of preparing a compound of formula:



or a salt thereof;

wherein:

R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_A$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_2 is hydrogen;

R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_C$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_D$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_5 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_E$; $-CN$; $-SCN$; $-SR_E$; or $-N(R_E)_2$; wherein each occurrence of R_E is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;


each occurrence of R_7 is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_G$; $-C(=O)R_G$; $-CO_2R_G$; $-CN$; $-SCN$; $-SR_G$; $-SOR_G$; $-SO_2R_G$; $-NO_2$; $-N(R_G)_2$; $-NHC(O)R_G$; or $-C(R_G)_3$; wherein each occurrence of R_G is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

P is hydrogen, lower (C_1 - C_6) alkyl, acyl, or a protecting group;


each P' is independently hydrogen or a protecting group; and

n is an integer in the range of 0 to 3, inclusive;

by reacting under basic conditions an anion of a phthalide, wherein the phthalide has the formula:




with an enone having the formula:



or salt thereof; wherein R_6 is hydrogen.

25. A method of preparing a tetracycline of formula:



or salt thereof;

wherein:

R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_A$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_2 is hydrogen;

R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_C$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_D$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;


R_5 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_E$; $-CN$; $-SCN$; $-SR_E$; or $-N(R_E)_2$; wherein each occurrence of R_E is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

each occurrence of R_7 is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_G$; $-C(=O)R_G$; $-CO_2R_G$; $-CN$; $-SCN$; $-SR_G$; $-SOR_G$; $-SO_2R_G$; $-NO_2$; $-N(R_G)_2$; $-NHC(O)R_G$; or $-C(R_G)_3$; wherein each occurrence of R_G is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio; and


n is 0 or an integer in the range of 1 to 3, inclusive;

the method comprising steps of:

reacting under basic conditions an anion of a phthalide, wherein the phthalide has the formula:




wherein P is hydrogen, lower (C_1 - C_6) alkyl, acyl, or a protecting group, with an enone having the formula:




or salt thereof, wherein R_6 is hydrogen, and each P' is independently hydrogen or a protecting

to form a compound having the formula:




and

deprotecting the compound to form a tetracycline of formula:



or salt thereof.

26. A method of preparing a compound of formula:



wherein:

R₁ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_A; -C(=O)R_A; -CO₂R_A; -CN; -SCN; -SR_A; -SOR_A; -SO₂R_A; -NO₂; -N(R_A)₂; -NHC(O)R_A; or -C(R_A)₃; wherein each occurrence of R_A is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio,

arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R₂ is hydrogen;

R₃ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_C; -C(=O)R_C; -CO₂R_C; -CN; -SCN; -SR_C; -SOR_C; -SO₂R_C; -NO₂; -N(R_C)₂; -NHC(O)R_C; or -C(R_C)₃; wherein each occurrence of R_C is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R₄ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_D; -C(=O)R_D; -CO₂R_D; -CN; -SCN; -SR_D; -SOR_D; -SO₂R_D; -NO₂; -N(R_D)₂; -NHC(O)R_D; or -C(R_D)₃; wherein each occurrence of R_D is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R₅ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_E; -CN; -SCN; -SR_E; or -N(R_E)₂; wherein each occurrence of R_E is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;


each occurrence of R₇ is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_G; -C(=O)R_G; -CO₂R_G; -CN; -SCN; -SR_G; -SOR_G; -SO₂R_G; -NO₂; -N(R_G)₂; -NHC(O)R_G; or -C(R_G)₃; wherein each occurrence of R_G is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or

heteroarylthio;

each P' is independently hydrogen or a protecting group; and


n is an integer in the range of 0 to 3, inclusive;

by reacting an anion of a toluate, wherein the toluate has the formula:



wherein R₉ is substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl; and Y is a halogen or Sn(alkyl)₃, and wherein the anion is produced by metallation of the toluate;

with an enone of formula:



or salt thereof; wherein R₆ is hydrogen.

27. The method of claim 26, wherein R₁ is hydrogen.

28. The method of claim 26, wherein Y is Br.


29. The method of claim 26, wherein Y is I..

30. The method of claim 26, wherein Y is -Sn(Me)₃.

31. The method of claim 26, wherein R₉ is methyl.

32. The method of claim 26, wherein R₉ is substituted or unsubstituted phenyl.

33. The method of claim 26, wherein the metal reagent is a lithium reagent.
34. The method of claim 33, wherein the lithium reagent is selected from the group consisting of *n*-butyl lithium, phenyllithium, *t*-butyl lithium, methyl lithium, and mesityl lithium.
35. The method of claim 26, wherein the metal reagent is a zinc reagent.
36. The method of claim 26, wherein the metal reagent is a magnesium reagent.
37. The method of claim 26, wherein the metallated toluate is a Grignard reagent.
38. The method of claim 26, wherein the metal reagent is a zero valent metal or an ate complex.
39. A method of preparing a tetracycline of formula:



or salt thereof;

wherein:

R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_A$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroarylthio, or heteroarylthio;

R₂ is hydrogen;

R₃ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_C; -C(=O)R_C; -CO₂R_C; -CN; -SCN; -SR_C; -SOR_C; -SO₂R_C; -NO₂; -N(R_C)₂; -NHC(O)R_C; or -C(R_C)₃; wherein each occurrence of R_C is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R₄ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_D; -C(=O)R_D; -CO₂R_D; -CN; -SCN; -SR_D; -SOR_D; -SO₂R_D; -NO₂; -N(R_D)₂; -NHC(O)R_D; or -C(R_D)₃; wherein each occurrence of R_D is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R₅ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_E; -CN; -SCN; -SR_E; or -N(R_E)₂; wherein each occurrence of R_E is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;


each occurrence of R₇ is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_G; -C(=O)R_G; -CO₂R_G; -CN; -SCN; -SR_G; -SOR_G; -SO₂R_G; -NO₂; -N(R_G)₂; -NHC(O)R_G; or -C(R_G)₃; wherein each occurrence of R_G is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

and


n is 0 or an integer in the range of 1 to 3, inclusive;

the method comprising steps of:

reacting an anion of a toluate, wherein the toluate has the formula:




wherein Y is halogen or Sn(alkyl)₃; and R₉ is substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl, and wherein the anion is produced by metallation of the toluate, with an enone of the formula:




or salt thereof;

wherein R₆ is hydrogen and each P' is independently hydrogen or a protecting group; and to produce a compound of the formula:



or salt thereof; and


deprotecting the compound to provide a tetracycline having the formula:



or salt thereof.

40. The method of claim 39, wherein R_1 is hydrogen.
41. The method of claim 39, wherein Y is Br.
42. The method of claim 39, wherein Y is I.
43. The method of claim 39, wherein Y is $-\text{Sn}(\text{Me})_3$.
44. The method of claim 39, wherein R_9 is phenyl.
45. The method of claim 39, wherein R_9 is substituted phenyl.
46. The method of claim 39, wherein the metallation of the toluate is performed with a lithium reagent.
47. The method of claim 46, wherein the lithium reagent is selected from the group consisting of *n*-butyl lithium, phenyllithium, *t*-butyl lithium, methyl lithium, and mesityl lithium.
48. The method of claim 39, wherein the metal reagent is a zinc reagent.
49. The method of claim 39, wherein the metal reagent is a magnesium reagent.

50. The method of claim 39, wherein the metallated toluate is a Grignard reagent.
51. The method of claim 39, wherein the metal reagent is a zero valent metal or ate complex.
52. A method of preparing compound of the formula:



or salt thereof;

wherein:

R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_A$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_2 is hydrogen;

R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_C$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched

heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_D$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_5 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_E$; $-CN$; $-SCN$; $-SR_E$; or $-N(R_E)_2$; wherein each occurrence of R_E is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

each occurrence of R_7 is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_G$; $-C(=O)R_G$; $-CO_2R_G$; $-CN$; $-SCN$; $-SR_G$; $-SOR_G$; $-SO_2R_G$; $-NO_2$; $-N(R_G)_2$; $-NHC(O)R_G$; or $-C(R_G)_3$; wherein each occurrence of R_G is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

each occurrence of P' is independently hydrogen or a protecting group;



represents a substituted or unsubstituted aryl; a substituted or unsubstituted heteroaryl; a substituted or unsubstituted carbocyclic; or a substituted or unsubstituted heterocyclic in which each occurrence of X is selected from the group consisting of $-O-$, $-S-$, $-NR_8-$, $-C(R_8)_2-$;


R_8 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_H$; $=O$, $-C(=O)R_H$; $-CO_2R_H$; $-CN$; $-SCN$; $-SR_H$; $-SOR_H$; $-SO_2R_H$; $-NO_2$; -

$N(R_H)_2$; $-NHC(O)R_H$; or $-C(R_H)_3$; wherein each occurrence of R_H is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

m is an integer in the range of 1 to 5, inclusive; and


the bonds between adjacent X moieties are either single or double bonds;

by reacting an anion of a toluate, wherein the toluate has the formula:



wherein R_9 is selected from the group consisting of substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl; and Y is a halogen or $Sn(alkyl)_3$, and wherein the anion is formed by metalation of the toluate,

with an enone of formula:



or salt thereof; wherein R_6 is hydrogen.


53. The method of claim 52, wherein R_1 is hydrogen.

54. The method of claim 52, wherein Y is Br.

55. The method of claim 52, wherein Y is I.

56. The method of claim 52, wherein Y is $-Sn(Me)_3$.

57. The method of claim 52, wherein R₉ is phenyl.
58. The method of claim 52, wherein R₉ is substituted phenyl.
59. The method of claim 52, wherein the metallation of the toluate is performed with a lithium reagent.
60. The method of claim 59, wherein the lithium reagent is selected from the group consisting of n-butyl lithium, phenyllithium, *t*-butyl lithium, methyl lithium, and mesityl lithium.
61. The method of claim 52, wherein the metal reagent is a zinc reagent.
62. The method of claim 52, wherein the metal reagent is a magnesium reagent.
63. The method of claim 52, wherein the metallated toluate is a Grignard reagent.
64. The method of claim 52, wherein the metal reagent is a zero valent metal or an ate complex.
65. A method of preparing a pentacycline of formula:



or salt thereof;

wherein:

R₁ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched

heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_A$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

R_2 is hydrogen;

R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_C$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;


R_4 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_D$; $-C(=O)R_D$; $-CO_2R_D$; $-CN$; $-SCN$; $-SR_D$; $-SOR_D$; $-SO_2R_D$; $-NO_2$; $-N(R_D)_2$; $-NHC(O)R_D$; or $-C(R_D)_3$; wherein each occurrence of R_D is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

each occurrence of R_E is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;

each occurrence of R_7 is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_G$; $-C(=O)R_G$; $-CO_2R_G$; $-CN$; $-SCN$; $-SR_G$; $-SOR_G$; $-SO_2R_G$; $-NO_2$; $-N(R_G)_2$; $-NHC(O)R_G$; or $-C(R_G)_3$; wherein each occurrence of R_G is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or

heteroarylthio;

each occurrence of P' is independently hydrogen or a protecting group;




represents a substituted or unsubstituted aryl; a substituted or unsubstituted heteroaryl; a substituted or unsubstituted carbocyclic; or a substituted or unsubstituted heterocyclic in which each occurrence of X is selected from the group consisting of -O-, -S-, -NR₈-, -C(R₈)₂-;

R₈ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; -OR_H; =O, -C(=O)R_H; -CO₂R_H; -CN; -SCN; -SR_H; -SOR_H; -SO₂R_H; -NO₂; -N(R_H)₂; -NHC(O)R_H; or -C(R_H)₃; wherein each occurrence of R_H is independently hydrogen, a protecting group, aliphatic, heteroaliphatic, acyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, amino, alkylamino, dialkylamino, heteroaryloxy, or heteroarylthio;


m is an integer in the range of 1 to 5, inclusive; and

the bonds between adjacent X moieties are either single or double bonds;


by reacting an anion of a toluate, wherein the toluate has the formula:



wherein R₉ is selected from the group consisting of substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl; and Y is a halogen or Sn(alkyl)₃, and wherein the anion is formed by metalation of the toluate, with an enone of formula:




or salt thereof; wherein R_6 is hydrogen and wherein R_5 is $-N(R_E)_2$,
to provide a compound of formula:



or salt thereof, wherein R_5 is $-N(R_E)_2$; and

deprotecting the compound to provide a pentacycline having the formula:



or salt thereof.

66. The method of claim 65, wherein R_1 is hydrogen.

67. The method of claim 65, wherein Y is Br.

68. The method of claim 65, wherein Y is I.

69. The method of claim 65, wherein Y is $-Sn(Me)_3$.

70. The method of claim 65, wherein R₉ is phenyl.
71. The method of claim 65, wherein R₉ is substituted phenyl.
72. The method of claim 65, wherein the metallation of the toluate is performed with a lithium reagent.
73. The method of claim 72, wherein the lithium reagent is selected from the group consisting of n-butyl lithium, phenyllithium, *t*-butyl lithium, methyl lithium, and mesityl lithium.
74. The method of claim 65, wherein the metal reagent is a zinc reagent.
75. The method of claim 65, wherein the metal reagent is a magnesium reagent.
76. The method of claim 65, wherein the metallated toluate is a Grignard reagent.
77. The method of claim 65, wherein the metal reagent is a zero valent metal or an ate complex.