

- 1 -

PROCESS FOR THE PREPARATION OF 1-BENZYL-3-HYDROXYMETHYL- 1H- INDAZOLE
AND ITS DERIVATIVES AND REQUIRED MAGNESIUM INTERMEDIATES


SCOPE OF THE INVENTION

5 This invention relates to a process for the preparation of 1-benzyl-3-hydroxymethyl-1H-indazole.

In particular, the present invention relates to the process for the conversion of 1-benzyl-3-hydroxymethyl-1H-indazole according to formula (II) below, to the 1-benzyl-3-hydroxymethyl-1H-indazole
10 according to formula (I) below.

STATE OF THE ART

European patent EP-B-0 382 276 describes some 1-benzyl-3-hydroxymethyl-1H-indazole derivatives of formula (A) having analgesic
15 activity:



in which

R and R', which may be the same or different, are H or C₁₋₅ alkyl, and
R'' is H or C₁₋₄ alkyl,

20 possibly in the form of its salt with a pharmaceutically-acceptable organic or inorganic base when R'' is H.

- 2 -

In turn, European patent EP-B-0 510 748 on the other hand describes the use of the same derivatives for the preparation of a pharmaceutical composition active in the treatment of auto-immune diseases.

5 In addition to this, document EP-B1-0 858 337 describes a pharmaceutical composition comprising a compound of formula (A) in which $R = R' = CH_3$ and $R'' = H$, and an immunosuppressant.

European patent EP-B-1 005 332 describes the use of the same derivatives to prepare a pharmaceutical composition active in the
10 treatment of conditions deriving from the production of MCP-1.

Finally international patent application WO 2008/061671 describes the use of a compound of formula (A) to reduce blood levels of triglycerides, cholesterol and glucose.

Various processes for the preparation of compounds of formula (A)
15 are described in the abovementioned patent EP-B-0 382 276.

The processes described in patent EP-B-0 382 276 have as their key point the preparation of 1-benzyl-3-hydroxymethyl-1H-indazole, from which the compounds of formula (A) can be obtained via three different reaction routes.

20 The first reaction route provides for the conversion of 1-benzyl-3-hydroxymethyl-1H-indazole into the corresponding alcoholate which is then caused to react with $X-CRR'-COOR''$, where X is a leaving group selected from the group comprising halogens, arylene-SO₂-O-, or alkylene-SO₂-O-, to yield the compounds of formula (A).

25 The second reaction route provides for the conversion of 1-benzyl-3-hydroxymethyl-1H-indazole into the corresponding 3-halogenomethyl derivative which is subsequently caused to react with an alcoholate of formula $MeO-CRR'-COOR''$, where Me is an alkali metal, to yield the compounds of formula (A).

- 3 -

The third reaction route provides for the reaction of 1-benzyl-3-hydroxymethyl-1H-indazole with chloroform and a ketone of formula $O=CRR'$ in the presence of an alkaline base such as sodium hydroxide to yield the compounds of formula (A) where R'' is hydrogen.

5 Preparation of the key intermediate 1-benzyl-3-hydroxymethyl-1H-indazole by the processes described in patent EP-B-0 382 276 is carried out by reducing the corresponding 3-carboxylic acid with a suitable reducing agent, such as for example aluminium lithium hydride ($LiAlH_4$).

10

BRIEF DESCRIPTION OF THE INVENTION

The Applicant has become aware that the processes of synthesis known in the art and described in the abovementioned patent EP-B-0 382 276 have a number of disadvantages.

15 Firstly, 1-benzyl-1(H)-indazol-3-carboxylic acid is not a product which can be easily obtained on the market, and is rather expensive. In particular there are few suppliers and the synthesis routes described in the literature provide for benzylation of the corresponding 1(H)-indazol-3-carboxylic acid, which is also expensive and not easy to obtain. In the
20 second place, reduction of the 1-benzyl-1(H)-indazol-3-carboxylic acid to obtain 1-benzyl-3-hydroxymethyl-1H-indazole proceeds with high dilution factors.

Furthermore the second reaction route provides for the use of thionyl chloride to convert the 1-benzyl-3-hydroxymethyl-1H-indazole into the
25 corresponding 3-chloromethyl derivative. The use of thionyl chloride, a highly toxic substance, gives rise to considerable safety and management problems in industrial processes.

Finally, the third reaction route (the Bargellini reaction) has shown industrial disadvantages in low yields (less than 50%), the production of
30 carbon monoxide, a toxic and flammable gas, and the generation of


- 4 -

significant exothermic phenomena which are difficult to manage industrially (Davis *et al. Synthesis*, 12, (2004), 1959-1962). Apart from this the Bargellini reaction finds better application in the synthesis of ethers from phenols and not from aliphatic alcohols (US 3,262,850;


5 Cvetovich *et al., J. Org. Chem.*, (2005), 70, 8560-8563).

The Applicant has therefore considered the problem of developing a new process for the preparation of 1-benzyl-3-hydroxymethyl-1H-indazole in order to obtain compounds of formula (A) which is capable of overcoming the abovementioned disadvantages. In particular the

10 Applicant has extended the problem to the preparation of 1-benzyl-3-hydroxymethyl-1H-indazoles having the following formula (II).



in order to obtain compounds having the following formula (I)



15 in which the substituents from R₁ to R₁₂ have the meanings indicated below in the detailed description and in the claims.

- 5 -

The Applicant has found a new process for the preparation of 1-benzyl-3-hydroxymethyl-1H-indazoles of formula (II) to obtain compounds of formula (I) which considerably improves on one hand industrial applicability, yields and the costs of the new process in
5 comparison with the processes known hitherto, and on the other hand the quality of the compounds obtained using it.

The Applicant has surprisingly found that 1-benzyl-3-hydroxymethyl-1H-indazole, or its derivatives of formula (II), can easily be obtained by reacting a Grignard reagent having the formula (IV) described below
10 with a suitable electrophilic compound such as, for example, aldehydes, ketones or amides, with subsequent reduction, if necessary, of the intermediate carbonyl compound.

In particular the Applicant has surprisingly found that the Grignard reagent of formula (IV) is easily obtained from 1-benzyl-3-halogeno-1H-indazole or its derivatives of formula (III) described below through a
15 halogen/magnesium exchange reaction with Grignard reagents of the alkyl magnesium halide type, at low temperature.

The Applicant also believes that the Grignard reagents of formula (IV) are not known in the art.

20 In fact the only indazole organometallic derivatives known in the art, with the metal in the 3 position, are those with metals such as zinc (Knochel *et al.*, *Synlett* 2005, 267) or copper (Knochel *et al.*, *Synthesis* 2006, 15, 2618 and Knochel *et al.*, *Synlett* 2004, 13, 2303-2306), while reactions degrading the indazole ring in an attempt to prepare the
25 corresponding 3-organolithium compounds (Welch *et al.*, *Synthesis*, 1992, 937) and 3-organosodium derivatives (Tertov *et al.*, *Zhurnal Organicheskoi Khimii* 1970, 6; 2140) are known.

The said 1-benzyl-3-halogeno-1H-indazole derivatives of formula (III), which are widely known in the literature, can easily be obtained by
30 halogenation of 1H-indazole in the 3 position with subsequent


- 6 -

benzylation in the 1 position (Collot *et al.*, *Tetrahedron*, 1999, 55, 6917; Coller *et al.*, *Aust. J. Chem.* 1974, 27, 2343).

The Applicant has also surprisingly found that 1-benzyl-3-hydroxymethyl-1H-indazole, or its derivatives of formula (II), can be readily converted into the corresponding 3-halogenomethyl derivatives by mere treatment with hydrohalogen acids, and then to compounds of formula (I) by etherification with the appropriate hydroxycarboxylic acid or ester of formula (VI).

Alternatively the Applicant has also surprisingly found that 1-benzyl-3-hydroxymethyl-1H-indazole, or its derivatives of formula (II), can easily be converted into compounds of formula (I) by etherification with the appropriate α -halogenocarboxylic acid or ester of formula (VII) described below.

Accordingly, this invention relates to a process for the preparation of 1-benzyl-3-hydroxymethyl-1H-indazole derivatives represented by the following formula (I):




in which the substituents from R₁ to R₁₂ have the meanings indicated below in the detailed description and in the claims,

20 in which,


a) the 1-benzyl-3-hydroxymethyl-1H-indazole or a derivative thereof represented by the following formula (II):

- 7 -




is caused to react with the hydrohalogen acid of formula HX'' , where X'' is a halogen atom selected from the group consisting in chlorine, bromine and iodine, preferably chlorine, to form 1-benzyl-3-

- 5 halogenomethyl-1H-indazole or a derivative thereof represented by the following formula (V):



- b) the 1-benzyl-3-halogenomethyl-1H-indazole or a derivative thereof represented by the abovementioned formula (V) is caused to react in the presence of a strong base with a compound represented by the
- 10 following formula (VI):




- 8 -

to form the 1-benzyl-3-hydroxymethyl-1H-indazole derivatives represented by the abovementioned formula (I)..

5

DETAILED DESCRIPTION OF THE INVENTION

Advantageously, the process for the preparation of 1-benzyl-3-hydroxymethyl-1H-indazole and its derivatives represented by the following formula (II):



10

in which

R₁ and R₂, which may be the same or different, are hydrogen or an alkyl group having from 1 to 6 carbon atoms,

R₃, R₄ and R₈, which may be the same or different, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, and a halogen atom,

15

R₅ may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, a halogen atom, or together with one of R₆ and R₇ may form a ring having 5 or 6 carbon atoms, and


20

R₆ and R₇, which may be the same or different, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, or one of R₆ and R₇ together with R₅ may form a ring having 5 or 6 carbon atoms,.

provides that


- 9 -

a) a 1-benzyl-3-halogeno-1H-indazole of formula (III):



in which X is a halogen atom selected from iodine and bromine, preferably iodine, and R₃-R₈ have the abovementioned meanings,


5 is caused to react with an alkyl magnesium halide of formula RMgX' where R is an alkyl group from 1 to 6 carbon atoms and X' is a halogen atom selected from bromine and chlorine, preferably chlorine, to form intermediate compound (IV):



10 b) the said intermediate compound (IV) is caused to react with a carbonyl compound of formula R₁-CO-R₂, where R₁ and R₂ have the abovementioned meanings, to form a compound of formula (II), or alternatively to b)

b') the said intermediate compound (IV) is caused to react with an
15 amide of formula R'R''N-CO-R₁, where R' and R'', which may be the same or different, are an alkyl group having 1 to 3 carbon atoms and R₁ has the abovementioned meanings, to form an intermediate compound (VIII):

- 10 -



which is caused to react with a carbonyl group reducing agent to form a compound of formula (II).

Advantageously, stage a) is performed in the presence of a suitable
 5 solvent, such as for example tetrahydrofuran, 2-methyl-tetrahydrofuran, diethyl ether, dioxane, t-butyl-methyl ether, dibutyl ether, xylene, toluene, dichloromethane, chloroform, n-hexane, n-heptane and their mixtures and so on, preferably 2-methyl-tetrahydrofuran, tetrahydrofuran, toluene, xylene and their mixtures, and even more
 10 preferably 2-methyl-tetrahydrofuran.

The alkyl magnesium halide of formula RMgX' used in stage a) may be methylMgCl, ethylMgCl, n-propylMgCl, i-propylMgCl, n-butylMgCl, i-butylMgCl, sec-butylMgCl, t-butylMgCl, n-pentylMgCl, n-hexylMgCl, allylMgCl, cyclohexylMgCl, methylMgBr, ethylMgBr, n-propylMgBr,
 15 i-propylMgBr, n-butylMgBr, i-butylMgBr, sec-butylMgBr, t-butylMgBr, n-pentylMgBr, n-hexylMgBr, allylMgBr, cyclohexylMgBr, and preferably i-propylMgCl. These reagents may be obtained commercially or prepared according to methods extensively described in the literature (Silverman *et al.*, *Handbook of Grignard reagents*, Chapter 2, CRC
 20 Press).

Advantageously, the exchange reaction in stage a) may be catalysed by the addition of lithium salts, for example LiCl, as described in the literature (Knochel *et al.*, *Chem. Commun.*, 2005, 543).

- 11 -

Advantageously, stage a) is carried out at a temperature of between -30°C and +30°C, preferably at a temperature of between -20°C and -10°C.

Advantageously, stage a) is carried out using a molar ratio between
5 the alkyl magnesium halide of formula RMgX' and the 1-benzyl-3-halogeno-1H-indazole of formula (III) of between 1 and 4, preferably between 1.5 and 4.

Advantageously, stage b) is carried out in the presence of a suitable solvent, such as for example tetrahydrofuran, 2-methyl-tetrahydrofuran,
10 diethyl ether, dioxane, t-butylmethyl ether, dibutyl ether, xylene, toluene, dichloromethane, chloroform, n-hexane, n-heptane and their mixtures, and so on, preferably 2-methyl-tetrahydrofuran, tetrahydrofuran, toluene, xylene and their mixtures, but preferably 2-methyl-tetrahydrofuran.

Advantageously, stage b) is carried out using a carbonyl compound
15 selected from the group of aldehydes, such as for example formaldehyde, acetaldehyde, propanal, butanal, pentanal, hexanal, and the like, and ketones such as for example acetone, ethylmethyl ketone, isobutylmethyl ketone, and so on. Formaldehyde is preferably used, and
20 in particular polymers such as suitably depolymerised paraformaldehyde or trioxane are used as a source of formaldehyde.

Advantageously, stage b) is carried out using a molar ratio between the 1-benzyl-3-halogeno-1H-indazole of formula (III) and the carbonyl compound of formula $\text{R}_1\text{-CO-R}_2$ of between 1 and 6.

Advantageously, stage b) is carried out at a temperature of between
25 -30°C and +30°C, preferably at a temperature of between -10°C and 0°C.

Advantageously, stage b') is carried out using an alkyl amide selected from the group comprising N,N-dimethylformamide,
30 N,N-diethylformamide, N,N-di-n-propylformamide, N,N-

- 12 -

dimethylacetamide, N,N-diethylacetamide, N,N-di-n-propylacetamide, N,N-dimethylpropionamide, N,N-diethylpropionamide, N,N-di-n-propylpropionamide, preferably N,N-dimethylformamide.

Advantageously, stage b') is carried out at a temperature of between
5 -30°C and +30°C, preferably at a temperature of between -10°C and 0°C.

In particular, stage b') is carried out using a molar ratio between the 1-benzyl-3-halogeno-1H-indazole of formula (III) and the amide of formula R'R"N-CO-R₁ of between 1 and 4.

10 Advantageously, the carbonyl group reducing agent used in stage b') is selected from the group comprising hydrides, such as for example NaBH₄, KBH₄, LiBH₄, Zn(BH₄)₂, Ca(BH₄)₂, NaAlH₄, LiAlH₄, Et₃SiH, Bu₃SnH, i-Bu₂AlH, 70% NaAlH₂(OCH₂CH₂OCH₃)₂ in toluene, and derivatives. The carbonyl group reducing agent is preferably 70%
15 NaAlH₂(OCH₂CH₂OCH₃)₂ in toluene. Carbonyl group reducing agents are extensively reported in the literature (described, for example, in Smith, March, *March's Advanced Organic Chemistry*, 5th ed., pages 1197-1205, John Wiley & Sons, Inc. and Carey, Sundberg, *Advanced Organic Chemistry*, 4th ed., pages 262-290).

20 Advantageously, the reduction in stage b') is carried out in the presence of a suitable solvent such as for example tetrahydrofuran, 2-methyl-tetrahydrofuran, diethyl ether, dioxane, t-butylmethyl ether, dibutyl ether, xylene, toluene, dichloromethane, chloroform, n-hexane, n-heptane, methanol, ethanol, n-propanol, i-propanol, diglyme (bis-(2-methoxyethyl)ether), pyridine, dimethylsulphoxide (DMSO), acetic acid,
25 their mixtures and so on, preferably toluene, xylene, tetrahydrofuran, 2-methyl-tetrahydrofuran and their mixtures.

Advantageously the reduction in stage b') is carried out at a temperature of between 10° and 100°C, preferably at a temperature of
30 between 20°C and 60°C.

- 13 -

In particular, reduction of the carbonyl group in stage b') is carried out using a number of equivalents of hydride of between 1 and 3, preferably 2.

Advantageously, the R₁-R₈ groups in formulae (II), (III), (IV) and (VIII) described above may have the following meanings.


Preferably, R₁ and R₂, which may be the same or different, are represented by a hydrogen atom, or an alkyl group having from 1 to 3 carbon atoms.

Preferably, R₃, R₄ and R₈, which may be the same or different, may be hydrogen, a methyl group, an ethyl group, a methoxy group, an ethoxy group, a chlorine atom and a fluorine atom.

Advantageously, R₅ may be hydrogen, a methyl group, an ethyl group, a methoxy group, an ethoxy group, a chlorine atom and a fluorine atom, or together with one of R₆ and R₇ may form a ring having 6 carbon atoms.

Preferably, R₆ and R₇, which may be the same or different, may be hydrogen, a methyl group, an ethyl group, or one of R₆ and R₇ together with R₅ may form a ring having 6 carbon atoms.

The present invention relates to a process for the preparation of 1-benzyl-3-hydroxymethyl-1H-indazole derivatives represented by the following formula (I):



in which

- 14 -


R_1 - R_8 have the meanings in formula (II) above,

R_{10} and R_{11} , which may be the same or different, are hydrogen or an alkyl group having from 1 to 5 carbon atoms, and

R_{12} is hydrogen or an alkyl group having from 1 to 4 carbon atoms.

5 The process for preparation of the 1-benzyl-3-hydroxymethyl-1H-indazole derivatives shown in the abovementioned formula (I) according to the invention provides that

a) the 1-benzyl-3-hydroxymethyl-1H-indazole or a derivative thereof represented by the following formula (II):




10

in which R_1 - R_8 have the abovementioned meanings,

is caused to react with a hydrohalogen acid of formula HX'' , where X'' is a halogen atom selected from the group consisting in chlorine, bromine and iodine, preferably chlorine, to form 1-benzyl-3-


15 halogenomethyl-1H-indazole or a derivative thereof represented by the following formula (V):

- 15 -



in which R_1 - R_8 and X'' have the abovementioned meanings,

- b) the 1-benzyl-3-halogenomethyl-1H-indazole or a derivative thereof represented by the abovementioned formula (V) are caused to react in the presence of a strong base with a compound represented by the following formula (VI):



- in which R_{10} , R_{11} , which may be the same or different, and R_{12} have the abovementioned meanings, to form the 1-benzyl-3-hydroxymethyl-1H-indazole derivatives represented by the above formula (I).

- Preferably the abovementioned process for the preparation of the 1-benzyl-3-hydroxymethyl-1H-indazole derivatives represented by formula (I) may comprise formation of the salt of the carboxyl group represented by $-COOR_{12}$ by treatment with a pharmaceutically acceptable organic or inorganic base. This treatment may be carried out directly on the corresponding acid when R_{12} is hydrogen, or following the reaction of hydrolysing the ester when R_{12} is an alkyl group from 1 to 4 carbon atoms.

- 16 -

Advantageously, stage a) is carried out in aqueous solution or in organic solvent. The hydrohalogen acid of formula HX" used is concentrated or dilute hydrochloric acid, hydrobromic acid or hydroiodic acid, preferably hydrochloric acid in a concentration such as to have a
5 molar ratio between the acid and the compound of formula (II) of between 1 and 20, preferably between 1 and 5, and even more preferably approximately 3.

Advantageously, stage a) is carried out at a temperature of between 25°C and 100°C, preferably at a temperature of between 60°C and
10 90°C.

The organic solvent preferably used in stage a) is selected from the group comprising toluene, xylene, acetic acid, dioxane, dibutylether, 2-methyl-tetrahydrofuran.

Advantageously, stage b) is carried out in aprotic solvents, such as
15 for example tetrahydrofuran, dioxane, N,N-dimethylformamide, toluene, N-methylpyrrolidone, dimethylsulphoxide, hexamethylphosphoramide, acetone, isobutylmethyl ketone, methylethyl ketone or their mixtures, preferably toluene or N,N-dimethylformamide and their mixtures.

The strong base used in stage b) is preferably selected from the
20 group comprising sodium hydride, metallic sodium, metallic potassium, butyl lithium, lithium-diisopropyl amide, sodium amide, potassium hydride, preferably sodium hydride.

Advantageously, stage b) is carried out using an α -hydroxy acid selected from the group comprising hydroxyacetic acid, lactic acid,
25 α -hydroxyisobutyric acid, α -hydroxybutyric acid, 2-ethyl-2-hydroxybutyric acid, 2-hydroxyisovaleric acid, 2-hydroxy-3,3-dimethylbutyric acid, 2-hydroxyisocaproic acid, preferably α -hydroxyisobutyric acid.

Advantageously, stage b) is carried out using an α -hydroxyester
30 selected from the group comprising methyl glycolate, ethyl glycolate,

butyl glycolate, methyl lactate, ethyl lactate, butyl lactate, t-butyl lactate, isopropyl lactate, isobutyl lactate, methyl-2-hydroxyisobutyrate, ethyl-2-hydroxyisobutyrate, ethyl-2-hydroxyvalerate, t-butyl-2-hydroxybutyrate, preferably ethyl-2-hydroxyisobutyrate.

- 5 Preferably, the molar ratio between the 1-benzyl-3-halomethyl-1H-indazole of formula (V) and the α -hydroxy acid or ester of formula (VI) is between 1 and 2, preferably approximately 1.2.

In particular, the molar ratio between the α -hydroxy acid of formula (VI) and the strong base is between 1 and 3, preferably approximately
10 2. Similarly the molar ratio between the α -hydroxy ester of formula (VI) and the strong base is between 1 and 1.5, preferably approximately 1.

Advantageously, the groups R_1 - R_{12} in formulae (I), (II), (V), (VI) and (VII) described above may have the following meanings.

Preferably R_1 and R_2 , which may be the same or different, are
15 represented by a hydrogen atom, or an alkyl group having from 1 to 3 carbon atoms.

Preferably, R_3 , R_4 and R_8 , which may be the same or different, may be hydrogen, a methyl group, an ethyl group, a methoxy group, an ethoxy group, a chlorine atom and a fluorine atom.

20 Advantageously, R_5 may be hydrogen, a methyl group, an ethyl group, a methoxy group, an ethoxy group, a chlorine atom and a fluorine atom, or together with one of R_6 and R_7 may form a ring having 6 carbon atoms.

Preferably, R_6 and R_7 , which may be the same or different, may be
25 hydrogen, a methyl group, an ethyl group, or one of R_6 and R_7 together with R_5 may form a ring having 6 carbon atoms.

Preferably, R_{10} and R_{11} , which may be the same or different, are hydrogen or an alkyl group having 1 to 3 carbon atoms, and R_{12} is hydrogen or an alkyl group having 1 to 3 carbon atoms.

- 18 -

The following examples are intended to illustrate this invention without however restricting it in any way.

EXPERIMENTAL PART

5 The compounds 3-iodo-1H-indazole and 1-benzyl-3-iodo-1H-indazole were prepared according to the procedure reported by Collot *et al.* (*Tetrahedron*, 55, 6917, 1999). The compound 3-bromo-1H-indazole was prepared using the procedure reported by Collier *et al.* (*Aust. J. Chem.* 1974, 27, 2343).

10 **Example 1**

Preparation of 1-benzyl-3-bromoindazole

3-bromo-1H-indazole (90.4 g, 0.459 mol, 1.0 eq.) and toluene (450 mL) were placed in a 1 litre flask fitted with a mechanical stirrer under an atmosphere of nitrogen. Then potassium t-butoxide (t-BuOK, 15 54.2 g, 0.483 mol, 1.05 eq.) was added at room temperature over about half an hour and benzyl bromide (86.3 g, 0.505 mol, 1.1 eq.) was added over approximately 1.5 hours. The mixture was left stirred at the same temperature until the reaction was complete (checked by TLC, approximately 3 hours). Then 0.1 M HCl (45 mL) and water (90 mL) 20 were added and the resulting phases were separated. The organic phase was washed with water, and the solvent was evaporated off at reduced pressure in order to obtain a red oily residue. The product was then precipitated through the addition of n-heptane, filtered and dried under vacuum at room temperature. Yield: 65.9 g of beige solid (yield 25 50%).

¹H NMR (300 MHz, DMSO-d₆) δ (ppm) 5.67 (s, 2H), 7.29 (m, 6H), 7.50 (ddd, 1H, J = 8.6 Hz, 6.9 Hz, 1.0 Hz), 7.60 (dd, 1H, J = 8.2 Hz, 0.7 Hz), 7.80 (dd, 1H, J = 8.6 Hz, 0.7 Hz).

¹³C NMR (300 MHz, DMSO-d₆) δ (ppm) 52.2, 110.4, 119.5, 121.7, 30 122.9, 127.4, 127.4, 127.6, 127.7, 128.6, 128.6, 129.6, 136.9, 140.5.

Example 2*Preparation of 1-benzyl-3-hydroxymethyl-1H-indazole*

- A solution of i-propylmagnesium chloride (i-PrMgCl) in
- 5 2-methyltetrahydrofuran (Me-THF) was prepared in a suitably thoroughly dried flask maintained under an atmosphere of nitrogen from magnesium metal (Mg, 10.91 g, 0.4489 mol, 1.5 eq.) activated with iodine crystals and a solution of isopropyl chloride (i-PrCl, 41.0 mL, 0.4489 mol, 1.5 eq.) in anhydrous Me-THF (185 mL). After cooling to
- 10 approximately -10°C a solution of 1-benzyl-3-iodo-1H-indazole (100 g, 0.2993 mol, 1.0 eq.) in anhydrous Me-THF (120 mL) was added over 1 hour keeping the temperature constant. The reaction mixture was kept stirred for a further hour to complete the halogen/magnesium exchange, and a yellow suspension was obtained. Gaseous formaldehyde
- 15 (generated by heating a suspension of 54 g of paraformaldehyde in 150 mL of xylene at approximately 115°C) was passed over this for approximately two hours at a temperature of below 0°C. When the reaction was complete, dilute H₃PO₄ was added and the excess of re-polymerised paraformaldehyde was removed by filtration. The phases
- 20 were separated and the organic phase was then washed with a dilute solution of NaHCO₃ and then concentrated. The product, precipitated out by the addition of n-hexane, was collected by filtration and dried. Yield: 56.8 g of white solid (79.6%).
- mp: 85-86°C
- 25 ¹H NMR (300 MHz, DMSO-d₆) δ (ppm) 4.79 (d, 2H, J = 5.8 Hz), 5.27 (t, 1H, J = 5.8 Hz), 5.6 (s, 2H), 7.12 (t, 1H, J = 7.5 Hz), 7.28 (m, 5H), 7.36 (t, 1H, J = 7.2 Hz), 7.64 (d, 1H, J = 8.5 Hz), 7.86 (d, 1H, J = 8.2 Hz).

- 20 -

^{13}C NMR (300 MHz, DMSO- d_6) δ (ppm) 51.6, 56.6, 109.6, 120.0, 120.9, 122.2, 126.2, 127.3, 127.3, 127.4, 128.5, 128.5, 137.7, 140.3, 145.2.

5 **Example 3**

Preparation of 1-benzyl-3-hydroxymethyl-1H-indazole

A solution of 2M i-PrMgCl in THF (69 mL, 138 mmol, 4.0 eq.) was added to a suitably thoroughly dried flask maintained under a nitrogen atmosphere. The solution was cooled to approximately -10°C . A
10 solution of 1-benzyl-3-bromo-1H-indazole (10 g, 34.8 mmol, 1.0 eq.) in anhydrous THF (40 mL) was added over approximately one hour keeping the temperature constant. The reaction mixture was kept stirred for at least 6 hours, and a yellow suspension was obtained. Gaseous formaldehyde (generated by heating a suspension of 16.7 g of
15 paraformaldehyde in 60 mL of xylene at approximately 115°C) was passed over this for approximately two hours at a temperature of below 0°C . When the reaction was complete, dilute H_3PO_4 was added and the excess of re-polymerised paraformaldehyde was removed by filtration. Me-THF (60 mL) was added to the mixture and the phases were
20 separated. The organic phase was washed with a dilute solution of NaHCO_3 . After concentration of the organic phase an oily residue containing the product was obtained. Subsequent purification of the crude product by silica gel chromatography yielded 2.8 g of white solid (yield: 34%).
25 mp: $85-86^\circ\text{C}$
 ^1H NMR (300 MHz, DMSO- d_6) δ (ppm) 4.79 (d, 2H, $J = 5.8$ Hz), 5.27 (t, 1H, $J = 5.8$ Hz), 5.6 (s, 2H), 7.12 (t, 1H, $J = 7.5$ Hz), 7.28 (m, 5H), 7.36 (t, 1H, $J = 7.2$ Hz), 7.64 (d, 1H, $J = 8.5$ Hz), 7.86 (d, 1H, $J = 8.2$ Hz).

- 21 -

^{13}C NMR (300 MHz, DMSO- d_6) δ (ppm) 51.6, 56.6, 109.6, 120.0, 120.9, 122.2, 126.2, 127.3, 127.3, 127.4, 128.5, 128.5, 137.7, 140.3, 145.2.

5 **Example 4**

Preparation of 1-benzyl-3-chloromethyl-1H-indazole

1-benzyl-3-hydroxymethyl-1H-indazole (400 g, 1.7 mol, 1 eq.), toluene (1.6 L) and concentrated HCl (422 mL, 5.1 mol, 3.0 eq.) were added to a three-necked flask fitted with a mechanical stirrer and reflux
10 condenser.

The reaction mixture was heated to approximately 90°C and kept stirred until the reaction was complete (checked by TLC, approximately two hours). After cooling to room temperature NaCl was added (approximately 10 g), the phases were separated, and the aqueous
15 phase was discharged. The organic phase was washed with a saturated solution of NaHCO_3 (approximately 100 mL) and then concentrated. The product, precipitated out by the addition of n-hexane (approximately 500 mL), was filtered and dried. Yield: 398.2 g of white solid (91%).

mp: 89-91°C

20 ^1H NMR (300 MHz, DMSO- d_6) δ (ppm) 5.14 (s, 2H), 5.65 (s, 2H), 7.27 (m, 6H), 7.43 (m, 1H), 7.12 (d, 1H, $J = 8.5$ Hz), 7.88 (d, 1H, $J = 8.2$ Hz)

^{13}C NMR (300 MHz, DMSO- d_6) δ (ppm) 38.2, 51.8, 110.2, 120.1, 120.9, 121.7, 126.7, 127.3, 127.3, 127.5, 128.5, 128.5, 137.2, 140.4,
25 140.6.

Example 5

Preparation of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid

- 22 -

Ethyl-2-hydroxyisobutyrate (18.5 g, 140 mmol, 1.2 eq.), toluene (100 mL) and DMF (20 mL) were placed in a three-necked flask fitted with a mechanical stirrer and a reflux condenser under an inert atmosphere. A dispersion of 60% NaH (5.6 g, 140 mmol, 1.2 eq.) was added to the mixture in portions over a period of approximately 1.5 hours. A solution of 1-benzyl-3-chloromethyl-1H-indazole (30 g, 117 mmol, 1 eq.) in toluene (90 mL) and DMF (60 mL) was then added dropwise. The reaction mixture was heated to approximately 90°C and kept at that temperature until the reaction was complete (checked by TLC, approximately 10 hours). After cooling to room temperature the mixture was washed with acidified water and water. The organic phase was concentrated under reduced pressure and the oily residue obtained was treated with 10 M NaOH (36 mL) at reflux temperature for at least 3 hours. The product, which was precipitated out by the addition of concentrated HCl, was filtered and dried. Yield: 32.3 g of white solid (85%).

mp: 133-134°C.

Elemental analysis:

Calculated: C (70.35), H (6.21), N (8.64),

Found: C (70.15), H (6.17), N (8.63).

¹H NMR (300 MHz, DMSO-d₆) δ (ppm) 1.44 (s, 6H), 4.76 (s, 2H), 5.60 (s, 2H), 7.14 (t, 1H, J = 7.6 Hz), 7.20-7.34 (m, 5H), 7.37 (ddd, 1H, J = 8.3 Hz, 7.0 Hz, 1.1 Hz), 7.66 (d, 1H, J = 8.4 Hz), 7.94 (d, 1H, J = 8.1 Hz), 12.77 (s, 1H).

¹³C NMR (300 MHz, DMSO-d₆) δ (ppm) 24.48, 24.48, 51.63, 59.65, 76.93, 109.69, 120.22, 121.06, 122.62, 126.28, 127.36, 127.36, 127.44, 128.46, 128.46, 137.49, 140.31, 141.97, 175.46.

Example 6

Preparation of 1-benzyl-1H-indazol-3-carbaldehyde

- 23 -

A solution of i-PrMgCl in THF was prepared in a suitably thoroughly dried flask maintained under a nitrogen atmosphere from magnesium metal (Mg, 164 mg, 6.75 mmol, 1.5 eq.) activated with iodine crystals and a solution of i-PrCl (0.62 mL, 6.75 mmol, 1.5 eq.) in anhydrous THF
5 (2.8 mL).

After cooling to approximately -10°C a solution of 1-benzyl-3-iodo-1H-indazole (1.5 g, 4.5 mmol, 1.0 eq.) in anhydrous THF (5 mL) was added to the reaction mixture over one hour keeping the temperature constant. The reaction mixture was kept stirred for a further hour to
10 complete the halogen/magnesium exchange, yielding a yellow suspension. While stirring, dimethylformamide (DMF) (1.4 mL, 18 mmol, 4 eq.) was added to the suspension over one hour at a temperature below 0°C and the reaction mixture was kept stirred at the same temperature until the reaction was complete (checked by TLC).

15 Dilute H₃PO₄ and toluene were added to the reaction mixture and the phases were separated. The organic phase was washed with a solution of dilute NaHCO₃. After the organic phase had been concentrated, the product, precipitated out by the addition of n-hexane, was filtered and dried. Yield: 1.0 g of yellowish solid (94%).

20 ¹H NMR (300 MHz, DMSO-d₆) δ (ppm) 5.84 (s, 2H), 7.32 (m, 5H), 7.39 (ddd, 1H, J = 8.1 Hz, 7.0 Hz, 1.0 Hz), 7.53 (ddd, 1H, J = 8.4 Hz, 7.0 Hz, 1.2 Hz) 7.90 (dt, 1H, J = 8.5 Hz, 1.0 Hz), 8.16 (dt, 1H, J = 8.1 Hz, 1.2 Hz), 10.19 (s, 1H).

¹³C NMR (300 MHz, DMSO-d₆) δ (ppm) 52.9, 111.0, 121.0, 121.2,
25 124.2, 127.5, 127.6, 127.6, 127.9, 128.6, 128.6, 136.2, 140.7, 142.4, 186.8.

Example 7

Preparation of 1-benzyl-3-hydroxymethyl-1H-indazole

- 24 -

1-benzyl-1H-indazol-3-carbaldehyde (2.36 g, 10 mmol, 1 eq.) and toluene (12 mL) were placed in a thoroughly dried 100 mL flask fitted with a magnetic stirrer and inerted with nitrogen. A 70% solution of sodium dihydro-bis(2-methoxyethoxy)aluminate in toluene (2.8 mL,
5 10 mmol, 2 eq.) was then slowly added to the solution at room temperature. Once the reaction was complete (after approximately 15 minutes), 2 M HCl (10 mL), H₂O (10 mL) and toluene (15 mL) were added. The phases were separated and the aqueous phase was extracted twice with toluene. The pooled organic phases were washed
10 with water and concentrated. The product was then precipitated out by adding n-hexane, filtered and dried. Yield: 1.95 g of white solid (82.0%).
mp: 85-86°C.

¹H NMR (300 MHz, DMSO-d₆) δ (ppm) 4.79 (d, 2H, J = 5.8 Hz), 5.27 (t, 1H, J = 5.8 Hz), 5.6 (s, 2H), 7.12 (t, 1H, J = 7.5 Hz), 7.28 (m, 5H)
15 7.36 (t, 1H, J = 7.2 Hz), 7.64 (d, 1H, J = 8.5 Hz), 7.86 (d, 1H, J = 8.2 Hz).


¹³C NMR (300 MHz, DMSO-d₆) δ (ppm) 51.6, 56.6, 109.6, 120.0, 120.9, 122.2, 126.2, 127.3, 127.3, 127.4, 128.5, 128.5, 137.7, 140.3, 145.2.

20

- 25 -

CLAIMS

1. A process for the preparation of 1-benzyl-3-hydroxymethyl-1H-indazole represented by the following formula (I):
- 5 indazole represented by the following formula (I):




in which


- R_1 and R_2 , which may be the same or different, are hydrogen or an alkyl group having 1 to 6 carbon atoms,
- 10 R_3 , R_4 and R_8 , which may be the same or different, may be hydrogen, an alkyl group having 1 to 5 carbon atoms, an alkoxy group having 1 to 3 carbon atoms and a halogen atom,
- R_5 may be hydrogen, an alkyl group having 1 to 5 carbon atoms, an alkoxy group having 1 to 3 carbon atoms, a halogen atom, or together
- 15 with one of R_6 and R_7 may form a ring having 5 or 6 carbon atoms,
- R_6 and R_7 , which may be the same or different, may be hydrogen, an alkyl group having 1 to 5 carbon atoms, or one of R_6 and R_7 together with R_5 may form a ring having 5 or 6 carbon atoms,
- R_{10} and R_{11} , which may be the same or different, are hydrogen or an
- 20 alkyl group having 1 to 5 carbon atoms, and
- R_{12} is hydrogen or an alkyl group having 1 to 4 carbon atoms,
- in which

- 26 -

a) 1-benzyl-3-hydroxymethyl-1H-indazole or a derivative thereof represented by the following formula (II):




is caused to react with a hydrohalogen acid of formula HX'' , where X'' is a halogen atom selected from the group comprising chlorine, bromine and iodine, preferably chlorine, to form 1-benzyl-3-halogenomethyl-1H-indazole or a derivative thereof represented by the following formula (V):



b) 1-benzyl-3-halogenomethyl-1H-indazole or a derivative thereof represented by the abovementioned formula (V) is caused to react in the presence of a strong base with a compound represented by the following formula (VI):

- 27 -



to form the 1-benzyl-3-hydroxymethyl-1H-indazole represented by the abovementioned formula (I).

- 5 2. The process of preparation according to Claim 1, in which the said 1-benzyl-3-hydroxymethyl-1H-indazole represented by formula (I) are caused to form salts with the carboxyl group -COOR_{12} by treatment with a pharmaceutically acceptable organic or inorganic base.
- 10 3. The process of preparation according to Claim 1 wherein R_1 and R_2 , which may be the same or different, are represented by a hydrogen atom, or an alkyl group having 1 to 3 carbon atoms.
- 15 4. The process of preparation according to Claim 1 wherein R_3 , R_4 and R_8 , which may be the same or different, may be hydrogen, a methyl group, an ethyl group, a methoxy group, an ethoxy group, a chlorine atom and a fluorine atom,
- 20 5. The process of preparation according to Claim 1 wherein R_5 may be hydrogen, a methyl group, an ethyl group, a methoxy group, an ethoxy group, a chlorine atom and a fluorine atom, or together with R_6 and R_7 may form a ring having 6 carbon atoms, and
- 25 6. The process of preparation according to Claim 1 wherein R_6 and R_7 , which may be the same or different, may be hydrogen, a methyl group, an ethyl group, or one of R_6 and R_7 together with R_5 may form a ring having 6 carbon atoms.