


least a fluorine atom, a chlorine atom or a bromine atom as the halogen atom; R³ is a fluorine atom or a chlorine atom; R⁴ is a trifluoromethyl group.

Further, particularly more preferred is one wherein R¹ is a hydrogen atom or a methyl group; R² is a haloalkyl group represented by the formula, -CF₂CF₂X, in which X is a hydrogen atom, a fluorine atom, a chlorine atom or a bromine atom; R³ is a chlorine atom; R⁴ is a trifluoromethyl group; the most preferred being one wherein R¹ is a methyl group; R² is a haloalkyl group represented by the formula, -CF₂CF₂X, in which X is a hydrogen atom, a fluorine atom, a chlorine atom or a bromine atom; R³ is a chlorine atom; R⁴ is a trifluoromethyl group.

The compounds of the present invention can be produced according to the following reaction scheme.



wherein R¹, R², R³ and R⁴ are each as defined above and A is a halogen atom.

The compound of the present invention can be produced by reacting halide compounds having the formula [III] with imidazole derivatives having the formula [II]

The reaction may be carried out at about -5°C to about 150°C for about 1 to 24 hours in a solvent in the presence of a reagent for removing a hydrogen halide.

The amounts of the reagents used in the reaction may be 1-2 equivalents of the halide compounds having the formula [III] and 1-4 equivalents of the reagent for removing a hydrogen halide to one equivalent of the imidazole derivatives having the formula [II].

Examples of the solvent which is used for the both reactions described above include aliphatic hydrocarbons such as hexane, heptane, ligroin, petroleum ether, etc.; aromatic hydrocarbons such as benzene, toluene, xylene, etc.; halogenated hydrocarbons such as chloroform, carbon tetrachloride, dichloroethane, chlorobenzene, dichlorobenzene, etc.; ethers such as diethyl ether, diisopropyl ether, dioxan, tetrahydrofuran, ethylene glycol dimethyl ether, etc.; ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone, isophorone, cyclohexanone, etc.; esters such as ethyl acetate, butyl acetate, etc.; nitro compounds such as nitroethane, nitrobenzene, etc.; nitriles such as acetonitrile, isobutyronitrile, etc.; tertiary amines such as pyridine, triethylamine, N,N-diethylaniline, tributylamine, N-methylmorpholine, etc.; acid amides such as formamide, N,N-dimethylformamide, N,N-dimethylacetamide, etc.; sulfur compounds such as dimethylsulfoxide, sulfolane, etc.; or mixtures thereof.

Examples of the reagent of removing hydrogen halide include organic bases such as pyridine, triethylamine, N,N-diethylaniline, etc.; inorganic bases such as sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium hydrogen carbonate, calcium carbonate, sodium hydride, etc.; alkali metal alkoxides such as sodium methoxide, sodium ethoxide, etc.

After completion of the reaction, post-treatment follows in a conventional manner. If necessary and desired, the product may further be purified by chromatography, distillation, recrystallization, etc.

The imidazole derivatives having the formula [II] and the halide compounds having the formula [III] which

are used as raw materials for the compounds of the present invention are prepared by the methods described in U.S. Patent 3,868,458, U.S. Patent 3,940,484, U.S. Patent 3,996,366, J. Org. Chem., 47, 2867 (1982), Japan Patent (laid open) 86-286,370 and U.S. Patent 3,888,932, U.S. Patent 3,928,416, European Patent 23,100, European Patent 34,402, West German Patent 2,606,393; West German Patent 3,545,570, U.S. Patent 4,184,041, British Patent 2,002,368, British Patent 1,121,211, Japan Patent (laid open) 84-20,269 respectively, or in a manner similar to the methods.

Examples of the compounds of the present invention are shown in Table 1 below.

10

15

20

25

30

35

40

45

50

55

R1	R2	R3	R4
H	CF ₃	F	CF ₃
H	CF ₃	Cl	CF ₃
H	CF ₃	Br	CF ₃
H	CF ₃	NO ₂	CF ₃
H	CF ₃	CF ₃	CF ₃
H	CF ₂ CF ₂ H	F	CF ₃
H	CF ₂ CF ₂ H	Cl	CF ₃
H	CF ₂ CF ₂ H	Br	CF ₃
H	CF ₂ CF ₂ H	NO ₂	CF ₃
H	CF ₂ CF ₂ H	CF ₃	CF ₃
H	CF ₂ CF ₃	F	CF ₃
H	CF ₂ CF ₃	Cl	CF ₃
H	CF ₂ CF ₃	Br	CF ₃
H	CF ₂ CF ₃	NO ₂	CF ₃
H	CF ₂ CF ₃	CF ₃	CF ₃
H	CF ₂ CF ₂ Cl	F	CF ₃
H	CF ₂ CF ₂ Cl	Cl	CF ₃

- Cont'd -

(Cont'd)

5	H	CF ₂ CF ₂ Cl	Br	CF ₃
	H	CF ₂ CF ₂ Cl	NO ₂	CF ₃
10	H	CF ₂ CF ₂ Cl	CF ₃	CF ₃
	H	CF ₂ CF ₂ Br	F	CF ₃
15	H	CF ₂ CF ₂ Br	Cl	CF ₃
	H	CF ₂ CF ₂ Br	Br	CF ₃
20	H	CF ₂ CF ₂ Br	NO ₂	CF ₃
	H	CF ₂ CF ₂ Br	CF ₃	CF ₃
25	H	CF ₂ CF ₂ CF ₃	F	CF ₃
	H	CF ₂ CF ₂ CF ₃	Cl	CF ₃
30	H	CF ₂ CF ₂ CF ₃	Br	CF ₃
	H	CF ₂ CF ₂ CF ₃	NO ₂	CF ₃
35	H	CF ₂ CF ₂ CF ₃	CF ₃	CF ₃
	H	CF ₂ CF ₂ CF ₂ CF ₃	Cl	CF ₃
40	H	CF ₂ CF ₂ CF ₂ CF ₃	Br	CF ₃
	CH ₃	CF ₃	F	CF ₃
50	CH ₃	CF ₃	Cl	CF ₃

- Cont'd -

(Cont'd)

5	CH ₃	CF ₃	Br	CF ₃
	CH ₃	CF ₃	NO ₂	CF ₃
10	CH ₃	CF ₃	CF ₃	CF ₃
	CH ₃	CF ₂ CF ₂ H	F	CF ₃
15	CH ₃	CF ₂ CF ₂ H	Cl	CF ₃
	CH ₃	CF ₂ CF ₂ H	Br	CF ₃
20	CH ₃	CF ₂ CF ₂ H	NO ₂	CF ₃
	CH ₃	CF ₂ CF ₂ H	CF ₃	CF ₃
25	CH ₃	CF ₂ CF ₃	F	CF ₃
	CH ₃	CF ₂ CF ₃	Cl	CF ₃
30	CH ₃	CF ₂ CF ₃	Br	CF ₃
	CH ₃	CF ₂ CF ₃	NO ₂	CF ₃
35	CH ₃	CF ₂ CF ₃	CF ₃	CF ₃
	CH ₃	CF ₂ CF ₂ Cl	F	CF ₃
40	CH ₃	CF ₂ CF ₂ Cl	Cl	CF ₃
	CH ₃	CF ₂ CF ₂ Cl	Br	CF ₃
45	CH ₃	CF ₂ CF ₂ Cl	NO ₂	CF ₃
50	CH ₃	CF ₂ CF ₂ Cl		

- Cont'd -

(Cont'd)

5	CH ₃	CF ₂ CF ₂ Cl	CF ₃	CF ₃
	CH ₃	CF ₂ CF ₂ Br	F	CF ₃
10	CH ₃	CF ₂ CF ₂ Br	Cl	CF ₃
	CH ₃	CF ₂ CF ₂ Br	Br	CF ₃
15	CH ₃	CF ₂ CF ₂ Br	NO ₂	CF ₃
	CH ₃	CF ₂ CF ₂ Br	CF ₃	CF ₃
20	CH ₃	CF ₂ CF ₂ CF ₃	F	CF ₃
	CH ₃	CF ₂ CF ₂ CF ₃	Cl	CF ₃
25	CH ₃	CF ₂ CF ₂ CF ₃	Br	CF ₃
	CH ₃	CF ₂ CF ₂ CF ₃	NO ₂	CF ₃
30	CH ₃	CF ₂ CF ₂ CF ₃	CF ₃	CF ₃
	CH ₃	CF ₂ CF ₂ CF ₃	Cl	CF ₃
35	CH ₃	CF ₂ CF ₂ CF ₃	Br	CF ₃
	CH ₃	CF ₂ CF ₂ CF ₃	F	CF ₃
40	CH ₃	CF ₂ CF ₂ CF ₃	Cl	CF ₃
	C ₂ H ₅	CF ₃	Br	CF ₃
45	C ₂ H ₅	CF ₃	F	CF ₃
	C ₂ H ₅	CF ₃	Cl	CF ₃
50	C ₂ H ₅	CF ₃	Br	CF ₃
	C ₂ H ₅	CF ₃	CF ₃	CF ₃

- Cont'd -

(Cont'd)

5	C ₂ H ₅	CF ₂ CF ₂ H	F	CF ₃
	C ₂ H ₅	CF ₂ CF ₂ H	Cl	CF ₃
10	C ₂ H ₅	CF ₂ CF ₂ H	Br	CF ₃
	C ₂ H ₅	CF ₂ CF ₂ H	CF ₃	CF ₃
15	C ₂ H ₅	CF ₂ CF ₃	F	CF ₃
	C ₂ H ₅	CF ₂ CF ₃	Cl	CF ₃
20	C ₂ H ₅	CF ₂ CF ₃	Br	CF ₃
	C ₂ H ₅	CF ₂ CF ₃	CF ₃	CF ₃
25	C ₂ H ₅	CF ₂ CF ₂ Cl	F	CF ₃
	C ₂ H ₅	CF ₂ CF ₂ Cl	Cl	CF ₃
30	C ₂ H ₅	CF ₂ CF ₂ Cl	Br	CF ₃
35	C ₂ H ₅	CF ₂ CF ₂ Cl	CF ₃	CF ₃
	C ₂ H ₅	CF ₂ CF ₂ Br	F	CF ₃
40	C ₂ H ₅	CF ₂ CF ₂ Br	Cl	CF ₃
45	C ₂ H ₅	CF ₂ CF ₂ Br	Br	CF ₃
	C ₂ H ₅	CF ₂ CF ₂ Br	CF ₃	CF ₃
50	C ₂ H ₅	CF ₂ CF ₂ CF ₃	F	CF ₃

- Cont'd -

(Cont'd)

5	C ₂ H ₅	CF ₂ CF ₂ CF ₃	Cl	CF ₃
	C ₂ H ₅	CF ₂ CF ₂ CF ₃	Br	CF ₃
10	C ₂ H ₅	CF ₂ CF ₂ CF ₃	CF ₃	CF ₃
	C ₃ H ₇ -n	CF ₃	F	CF ₃
15	C ₃ H ₇ -n	CF ₃	Cl	CF ₃
	C ₃ H ₇ -n	CF ₃	Br	CF ₃
20	C ₃ H ₇ -n	CF ₃	CF ₃	CF ₃
	C ₃ H ₇ -n	CF ₂ CF ₂ H	F	CF ₃
25	C ₃ H ₇ -n	CF ₂ CF ₂ H	Cl	CF ₃
30	C ₃ H ₇ -n	CF ₂ CF ₂ H	Br	CF ₃
	C ₃ H ₇ -n	CF ₂ CF ₂ H	CF ₃	CF ₃
35	C ₃ H ₇ -n	CF ₂ CF ₃	F	CF ₃
	C ₃ H ₇ -n	CF ₂ CF ₃	Cl	CF ₃
40	C ₃ H ₇ -n	CF ₂ CF ₃	Br	CF ₃
	C ₃ H ₇ -n	CF ₂ CF ₃	CF ₃	CF ₃
45	C ₃ H ₇ -n	CF ₂ CF ₂ Cl	F	CF ₃
50	C ₃ H ₇ -n	CF ₂ CF ₂ Cl	Cl	CF ₃

55

- Cont'd -

(Cont'd)

5	C ₃ H ₇ -n	CF ₂ CF ₂ Cl	Br	CF ₃
	C ₃ H ₇ -n	CF ₂ CF ₂ Cl	CF ₃	CF ₃
10	C ₃ H ₇ -n	CF ₂ CF ₂ Br	F	CF ₃
	C ₃ H ₇ -n	CF ₂ CF ₂ Br	Cl	CF ₃
15	C ₃ H ₇ -n	CF ₂ CF ₂ Br	Br	CF ₃
	C ₃ H ₇ -n	CF ₂ CF ₂ Br	CF ₃	CF ₃
20	C ₃ H ₇ -n	CF ₂ CF ₂ CF ₃	F	CF ₃
	C ₃ H ₇ -n	CF ₂ CF ₂ CF ₃	Cl	CF ₃
25	C ₃ H ₇ -n	CF ₂ CF ₂ CF ₃	Br	CF ₃
	C ₃ H ₇ -n	CF ₂ CF ₂ CF ₃	CF ₃	CF ₃
30	C ₃ H ₇ -iso	CF ₃	F	CF ₃
35	C ₃ H ₇ -iso	CF ₃	Cl	CF ₃
	C ₃ H ₇ -iso	CF ₃	Br	CF ₃
40	C ₃ H ₇ -iso	CF ₃	CF ₃	CF ₃
	C ₃ H ₇ -iso	CF ₂ CF ₂ H	F	CF ₃
45	C ₃ H ₇ -iso	CF ₂ CF ₂ H	Cl	CF ₃
50	C ₃ H ₇ -iso	CF ₂ CF ₂ H	Br	CF ₃

- Cont'd -

(Cont'd)

5	C ₃ H ₇ -iso	CF ₂ CF ₂ H	CF ₃	CF ₃
	C ₃ H ₇ -iso	CF ₂ CF ₃	F	CF ₃
10	C ₃ H ₇ -iso	CF ₂ CF ₃	Cl	CF ₃
	C ₃ H ₇ -iso	CF ₂ CF ₃	Br	CF ₃
15	C ₃ H ₇ -iso	CF ₂ CF ₃	CF ₃	CF ₃
	C ₃ H ₇ -iso	CF ₂ CF ₂ Cl	F	CF ₃
20	C ₃ H ₇ -iso	CF ₂ CF ₂ Cl	Cl	CF ₃
	C ₃ H ₇ -iso	CF ₂ CF ₂ Cl	Br	CF ₃
25	C ₃ H ₇ -iso	CF ₂ CF ₂ Cl	CF ₃	CF ₃
30	C ₃ H ₇ -iso	CF ₂ CF ₂ Br	F	CF ₃
	C ₃ H ₇ -iso	CF ₂ CF ₂ Br	Cl	CF ₃
35	C ₃ H ₇ -iso	CF ₂ CF ₂ Br	Br	CF ₃
40	C ₃ H ₇ -iso	CF ₂ CF ₂ Br	CF ₃	CF ₃
	C ₃ H ₇ -iso	CF ₂ CF ₂ CF ₃	F	CF ₃
45	C ₃ H ₇ -iso	CF ₂ CF ₂ CF ₃	Cl	CF ₃
	C ₃ H ₇ -iso	CF ₂ CF ₂ CF ₃	Br	CF ₃
50	C ₃ H ₇ -iso	CF ₂ CF ₂ CF ₃	CF ₃	CF ₃

55

- Cont'd -

(Cont'd)

5	H	CF ₃	Cl	OCHF ₂
	H	CF ₂ CF ₂ H	Cl	OCHF ₂
10	H	CF ₂ CF ₃	Cl	OCHF ₂
	H	CF ₂ CF ₂ Cl	Cl	OCHF ₂
15	H	CF ₂ CF ₂ Br	Cl	OCHF ₂
	H	CF ₂ CF ₂ CF ₃	Cl	OCHF ₂
20	CH ₃	CF ₃	Cl	OCHF ₂
	CH ₃	CF ₂ CF ₂ H	Cl	OCHF ₂
25	CH ₃	CF ₂ CF ₃	Cl	OCHF ₂
	CH ₃	CF ₂ CF ₂ Cl	Cl	OCHF ₂
30	CH ₃	CF ₂ CF ₂ Br	Cl	OCHF ₂
	CH ₃	CF ₂ CF ₂ CF ₃	Cl	OCHF ₂
35	H	CF ₃	Br	OCHF ₂
	H	CF ₂ CF ₂ H	Br	OCHF ₂
40	H	CF ₂ CF ₃	Br	OCHF ₂
	H	CF ₂ CF ₂ Cl	Br	OCHF ₂
45	H	CF ₂ CF ₂ Br	Br	OCHF ₂
	H	CF ₂ CF ₂ CF ₃	Br	OCHF ₂
50				
55				

- Cont'd -

(Cont'd)

5	CH ₃	CF ₃	Br	OCHF ₂
	CH ₃	CF ₂ CF ₂ H	Br	OCHF ₂
10	CH ₃	CF ₂ CF ₃	Br	OCHF ₂
	CH ₃	CF ₂ CF ₂ Cl	Br	OCHF ₂
15	CH ₃	CF ₂ CF ₂ Br	Br	OCHF ₂
	CH ₃	CF ₂ CF ₂ CF ₃	Br	OCHF ₂
20	H	CF ₃	Cl	OCClF ₂
	H	CF ₂ CF ₂ H	Cl	OCClF ₂
25	H	CF ₂ CF ₃	Cl	OCClF ₂
	H	CF ₂ CF ₂ Cl	Cl	OCClF ₂
30	H	CF ₂ CF ₂ Br	Cl	OCClF ₂
	H	CF ₂ CF ₂ CF ₃	Cl	OCClF ₂
35				
	CH ₃	CF ₃	Cl	OCClF ₂
40	CH ₃	CF ₂ CF ₂ H	Cl	OCClF ₂
	CH ₃	CF ₂ CF ₃	Cl	OCClF ₂
45	CH ₃	CF ₂ CF ₂ Cl	Cl	OCClF ₂
	CH ₃	CF ₂ CF ₂ Br	Cl	OCClF ₂
50				
	CH ₃	CF ₂ CF ₂ CF ₃	Cl	OCClF ₂
55				

Examples of harmful insects against which the compounds of the present invention exhibit remarkable

effects include the following:

Harmful insects belonging to Hemiptera:

- 5 Planthoppers such as small brown planthopper (Laodelphax striatellus), brown planthopper (Nilaparvata lugens), white-backed rice planthopper (Sogatella furcifera), etc.; leafhoppers such as green rice leafhopper (Nephotettix cincticeps), (Nephotettix virescens), etc.; aphids, bugs, whiteflies, scales, lace bugs, psyllids, etc.

Lepidoptera:

- 10 Pyralid moths such as rice stem borer (Chilo suppressalis), rice leafroller (Cnaphalocrocis medinalis), Indian meal moth (Plodia interpunctella), etc.; moths such as tobacco cutworm (Spodoptera litura), rice armyworm (Pseudaletia separata), cabbage armyworm (Mamestra brassicae), etc.; Pieridae such as common cabbageworm (Pieris rapae crucivora), etc.; Tortricidae or tortricid moths such as Adoxophyes spp., Grapholita spp., etc.; Carposinidae, Lyonetiid moths (Lyonetiidae), tussock moths (Lymantriidae), beet semi-looper (Autographa nigrisigna); harmful insects belonging to Agrothis spp. such as turnip cutworm (Agrothis segetum), black cutworm (Agrothis ipsilon); harmful insects belonging to Hilliothis spp.; diamondback moth (Plutella xylostella), clothes moths (Tineidae), casemaking clothes moth (Tinea translucens), webbing clothes moth (Tineola bisselliella); etc.

20 Harmful insects belonging to Diptera:

- Mosquitos such as common mosquito (Culex pipiens pallens), Culex tritaeniorhynchus, etc.; Aedes spp. such as Aedes aegypti, Aedes albopictus, etc.; Anopheles spp. such as Anopheles sinensis, etc.; midges (Chironomidae); Muscidae such as housefly (Musca domestica), false stablefly (Muscina stabulans), etc.; Calliphoridae; Sarcophagidae; lesser housefly (Fannia canicularis), Anthomyiidae or anthomyiid flies such as seedcorn maggot (Delia platura), onion maggot (Della antiqua), etc.; fruit flies (Tephritidae); small fruit flies (Drosophilidae); moth flies (Psychodidae); black flies (Simuliidae); Tabanidae; stable flies (Stomoxys); etc.

30 Harmful insects belonging to Coleoptera:

- Corn root worms such as western corn rootworm (Diabrotica virgifera), southern corn root worm (Diabrotica undecimpunctata), etc.; scarabs (Scarabaeidae) such as cupreous chafer (Anomala cuprea), soybean beetle (Anomala rufocuprea), etc.; weevils such as maize weevil (Sitophilus zeamais), rice water weevil (Lissorhoptrus oryzophilus), adzuki bean weevil (Callosobruchys chinensis), etc.; darkling beetles (Tenebrionidae) such as yellow mealworm (Tenebrio molitor), red flour beetle (Tribolium castaneum), etc.; leaf beetles (Chrysomelidae) such as cucurbit leaf beetle (Aulacophora femoralis), striped flea beetles (Phyllotreta striolata), etc.; Anobiidae; Epilachna spp. such as twenty-eight-spotted ladybirds (Epilachna vigintioctopunctata), etc.; powderpost beetles (Lyctidae); false powderpost beetles (Bostrychidae), Cerambycidae; robe beetle (Paderus fusipes), etc.

Harmful insects belonging to Dictyoptera:

- 45 German cockroach (Blattella germanica), smokybrown cockroach (Periplaneta fuliginosa), American cockroach (Periplaneta americana), brown cockroach (Periplaneta brunnea), oriental cockroach (Blatta orientalis), etc.

Harmful insects belonging to Thysanoptera:

- 50 Thrips palmi, flower thrips (Thrips hawaiiensis), etc.

Harmful insects belonging to Hymenoptera:

- 55 ants (Formicidae); hornets (Vespididae); bethylid wasps (Bethylidae); sawflies (Tenthredinidae) such as cabbage sawfly (Athalia rosae ruficornis), etc.

Harmful insects belonging to Orthoptera:

mole crickets (Gryllotalpidae); grasshoppers (Acrididae), etc.;

5 Harmful insects belonging to Aphaniptera:

Purex irritans, etc.

Harmful insects belonging to Anoplura:

10

Pediculus humanus capitis, Phthirus pubis, etc.

Harmful insects belonging to Isoptera:

15

Reticulitermes speratus, Formosan subterranean termite (Coptotermes formosanus), etc.

Moreover, the compounds of the present invention are very effective to the insects which develop the resistance against conventional insecticides.

20

In the case that the compounds of the present invention are used as the active ingredient of insecticidal compositions, the compounds may be used as they are, without adding any other components but in general, the compounds are mixed with a solid carrier, a liquid carrier, a gaseous carrier, a feed, etc. and, if necessary and desired, the mixture is further supplemented with a surfactant and other adjuvants used to prepare insecticidal preparations and prepared into forms such as oil sprays, emulsifiable concentrates, wettable powders, flowable concentrated, granules, dusts, aerosol, fumigants (fogging, etc.), poison bait, etc.

25

These formulations may contain any effective amount of insecticide, generally 0.01 to 95% by weight of the compounds of the present invention as the active ingredient, together with an inert carrier.

30

Examples of the solid carrier used for making formulations include fine powders or granulates, etc. of clays (kaolin clay, diatomaceous earth, synthetic hydrated silicon dioxide, bentonite, Fubasami clay, terra alba, etc.), talc, ceramics, other inorganic minerals (sericite, quartz, sulfur, activated carbon, calcium carbonate, hydrated silica, etc.), chemical fertilizers (ammonium sulfate, ammonium phosphate, ammonium nitrate, urea, ammonium chloride, etc.), etc. Examples of the liquid carrier include water, alcohols (methanol, ethanol, etc.), ketones (acetone, methyl ethyl ketone, etc.), aromatic hydrocarbons (benzene, toluene, xylene, ethylbenzene, methyl-naphthalene, etc.), aliphatic hydrocarbons (hexane, cyclohexane, kerosene, gas oil, etc.), esters (ethyl acetate, butyl acetate, etc.), nitriles (acetonitrile, isobutyronitrile, etc.), ethers (diisopropyl ether, dioxan, etc.), acid amides (N,N-dimethylformamide, N,N-dimethylacetamide, etc.), halogenated hydrocarbons (dichloromethane, trichloroethane, carbon tetrachloride, etc.), dimethylsulfoxide; vegetable oils such as soybean oil, cotton seed oil, etc. Examples of the gaseous carrier, i.e., propellant, include freon gas, butane gas, LPG (liquefied petroleum gas), dimethyl ether, carbon dioxide, etc.

35

Examples of the surfactant include alkyl sulfates, alkyl sulfonic acid salts, alkylaryl sulfonic acid salts, alkyl aryl ethers and polyoxyethylene derivatives thereof, polyethylene glycol ether, polyvalent alcohol esters, sugar alcohol derivatives, etc.

40

Examples of the adjuvants such as binders, dispersing agents, etc. for formulations include casein, gelatin, polysaccharides (starch powders, gum arabic, cellulose derivatives, alginic acid, etc.), lignin derivatives, bentonite, sugars, synthetic water-soluble high molecular substances (polyvinyl alcohol, polyvinylpyrrolidone, polyacrylic acid, etc.). Examples of the stabilizer include PAP (acidic isopropyl phosphate), BHT (2,6-di-tert-butyl-4-methylphenol), BHA (mixture of 2-tert-butyl-4-methoxyphenol and 3-tert-butyl-4-methoxyphenol), vegetable oils, mineral oils, surfactants, fatty acids or esters thereof, and the like.

45

As a base material for the poison baits, there are, for example, feed components such as crop powders, essential vegetable oil, sugars, crystalline cellulose etc.; antioxidants such as dibutylhydroxytoluene, nordihydroguaiaretic acid, etc.; preservatives such as dehydroacetic acid, etc.; feeding error preventing agents such as red pepper powders, etc.; incentive flavor such as cheese flavor, onion flavor, etc.

50

The thus obtained formulations may be used as they are or after diluting with water, etc. Alternatively, the formulations may be used as admixture with other insecticides, nematocides, acaricides, bacteriocides, herbicides, plant growth regulators, synergistic agents, fertilizers, soil conditioners, animal feed, etc., or may also be used simultaneously with them, without mixing therewith.

55

Where the compounds of the present invention are used as insecticides for agricultural use, the dose is generally 0.1 g to 100 g per 10 ares; when emulsifiable concentrates, wettable powders, flowable concentrates, etc. are used after diluting them with water, the concentration is 0.1 ppm to 500 ppm. Granules, dusts, etc. may be used as they are, without diluting them. For purposes of household and public hygiene, emulsifi-

able concentrates, wettable powders, flowable concentrates, etc. are diluted with water in a concentration of 0.1 ppm to 500 ppm; oils, aerosol, fumigants, poison baits, etc. may be used as they are.

These doses and concentrations may vary depending upon kind of formulations, timing for application, place applied, method for application, kinds of insect, condition of damages, etc. and may be increased or decreased, irrespective of the ranges set forth above.

Hereafter the present invention is described in more detail, by referring to synthesis examples, formulation examples and test examples.

Synthesis Example 1 (Synthesis of Compound No. (3))


To a solution of 0.37 g (2 m mol) of 4(5)-pentafluoroethylimidazole in 5 ml of N,N-dimethyl formamide was added 80 mg (2 m mol) of an oily sodium hydride (60%) while cooling with ice, followed by stirring at the same temperature for 10 minutes. After the reaction was completed, to the reaction mixture was added dropwise 0.43 g (2 m mol) of 2,3-dichloro-5-trifluoromethylpyridine, followed by stirring at room temperature for 8 hours. After the reaction was completed, the reaction mixture was poured into water, and extracted with ethyl acetate. Further, then the residue was washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The obtained product was subjected to silica gel chromatography to give 0.24 g of 1-(3-chloro-5-trifluoromethylpyridin-2-yl)-4-pentafluoroethylimidazole.
m.p. 52.0°C

Synthesis Example 2 (Synthesis of Compound No. (11))

To a solution of 0.22 g (1 m mol) of 2-methyl-4(5)-(2-chloro-1,1,2,2-tetrafluoroethyl) imidazole in 5 ml of N,N-dimethylformamide was added both of 0.21 g (1.5 m mol) of anhydrous potassium carbonate and 0.22 g (1 m mol) of 2,3-dichloro-5-trifluoromethylpyridine, followed by stirring at 80-85°C for 7 hours. After the reaction was completed, the reaction mixture was poured into water, and extracted with ethyl acetate. Further, then the residue was washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The obtained product was subjected to silica gel chromatography to give 0.27 g of 1-(3-chloro-5-trifluoromethylpyridin-2-yl)-2-methyl-4-(2-chloro-1,1,2,2-tetrafluoroethyl) imidazole.
m.p. 137.5°C

Examples of the present invention prepared in the same manner as above are shown in Table 2.

Table 2



Compound No.	R ¹	R ²	R ³	R ⁴	Physical constant
(1)	H	CF ₃	Cl	CF ₃	m.p. 73.3°C
(2)	H	CF ₂ CF ₂ H	Cl	CF ₃	25.4 n _D 1.4813
(3)	H	CF ₂ CF ₃	Cl	CF ₃	m.p. 52.0°C
(4)	H	CF ₂ CF ₃	CF ₃	CF ₃	23.9 n _D 1.4220
(5)	H	CF ₂ CF ₂ Br	Cl	CF ₃	24.0 n _D 1.4999
(6)	H	CF ₂ CF ₂ Br	CF ₃	CF ₃	24.5 n _D 1.4550
(7)	CH ₃	CF ₂ CF ₂ CF ₃	Cl	CF ₃	m.p. 77.9°C
(8)	CH ₃	CF ₃	Cl	CF ₃	m.p. 73.5°C
(9)	CH ₃	CF ₂ CF ₂ H	Cl	CF ₃	m.p. 147.9°C
(10)	CH ₃	CF ₂ CF ₃	Cl	CF ₃	m.p. 85.3°C
(11)	CH ₃	CF ₂ CF ₂ Cl	Cl	CF ₃	m.p. 137.5°C
(12)	CH ₃	CF ₂ CF ₂ Br	Cl	CF ₃	m.p. 63-65°C
(13)	CH ₃	CF ₂ CF ₂ Br	CF ₃	CF ₃	m.p. 60.3°C

Next, Formulation examples are shown, wherein parts are all by weight and the compounds of the present invention are designated by the compound numbers shown in Table 2.

Formulation Example 1. Emulsifiable concentrate

After 10 parts each of Compounds (1) through (13) of the present invention are dissolved in 35 parts of xylene and 35 parts of dimethylformamide, 14 parts of polyoxyethylene styrylphenyl ether and 6 parts of cal-

cium dodecylbenzenesulfonate are added to the solutions. The resulting mixtures are thoroughly mixed stirred to give 10% emulsifiable concentrate, respectively.

Formulation Example 2. Wettable powder

After 20 parts of Compound (1) through (13) of the present invention are added to a mixture of 4 parts of sodium laurylsulfate, 2 parts of calcium ligninsulfonate, 20 parts of synthetic hydrated silicon dioxide fine powders and 54 parts of diatomaceous earth, the mixture is mixed and stirred with a juice mixer to give 20% wettable powder.

Formulation Example 3. Granule

After 5 parts of synthetic hydrated silicon dioxide fine powders, 5 parts of sodium dodecylbenzenesulfonate, 30 parts of bentonite and 55 parts of clay are added to 5 parts of Compound (1) through (13) of the present invention, the mixture is thoroughly mixed and stirred. A suitable amount of water is further added to the mixture followed by stirring. The mixture is granulated with a granulator and air-dried to give 5% granule.

Formulation Example 4. Dust

After 1 part of Compound (7) of the present invention is dissolved in a appropriate amount of acetone, 5 parts of synthetic hydrated silicon dioxide fine powders, 0.3 part of PAP and 93.7 parts of clay are added to the solution. The mixture is mixed and stirred with a juice mixer and acetone is evaporated off to give 1% dust.

Formulation Example 5. Flowable concentrate

After 20 parts of Compound (12) of the present invention and 1.5 part of sorbitan trioleate are mixed with 28.5 parts of an aqueous solution containing 2 parts of polyvinyl alcohol, the mixture is finely divided (less than 3 μ in particle diameter) with a sand grinder. Then, 40 parts of aqueous solution containing 0.05 part of xanthane gum and 0.1 part of aluminum magnesium silicate are added to the powders and 10 parts of propylene glycol are further added thereto. The mixture is thoroughly mixed and stirred to give 20% flowable concentrate for aqueous suspension.

Formulation Example 6. Oil spray

After 0.1 part of Compound (1) through (13) of the present invention is dissolved in 5 parts of xylene and 5 parts of trichloroethane, the solution is mixed with 89.9 parts of deodorized kerosene to give 0.1% oil spray.

Formulation Example 7. Oil-based aerosol

After 0.1 part of Compound (1) through (13) of the present invention, 0.2 part of tetramethrin, 0.1 part of d-phenothrin, 10 parts of trichloroethane and 59.6 parts of deodorized kerosene are mixed with each other and dissolved. The solution is filled in an aerosol container. After a valve is mounted to the container, 30 parts of propellant (liquefied petroleum gas) are filled under pressure through the valve to give oil-based aerosol.

Formulation Example 8. Water-based aerosol

After 0.2 part of Compound (11) of the present invention, 0.2 part of d-allethrin, 0.2 part of d-phenothrin, 5 parts of xylene, 3.4 parts of deodorized kerosene and 1 part of emulsifier [ATMOS 300 (registered trademark, Atlas Chemical Co., Ltd.)] are mixed with each other and dissolved. The solution and 50 parts of distilled water are filled in an aerosol container. After a valve is mounted to the container, 40 parts of propellant (liquefied petroleum gas) are filled under pressure through the valve to give water-based aerosol.

Formulation Example 9. Mosquito coil

After 0.3 g of d-allethrin is added to 0.3 g of Compound (12) of the present invention, the mixture is dissolved in 20 ml of acetone. The solution is then uniformly mixed with 99.4 g of carrier for mosquito-coil (tapa powder : sake lees powder : wood powder of 4 : 3 : 3) with stirring and 120 ml of water is then added to the mixture. The mixture is thoroughly kneaded, molded and dried to give mosquito-coil.

Formulation Example 10. Electric mosquito mat

Acetone is added to 0.4 g of Compound (12) of the present invention, 0.4 g of d-allethrin and 0.4 g of piperonyl butoxide to dissolve and make the whole volume 10 ml. This solution, 0.5 ml, is uniformly impregnated with a base material for electric mat (a mixture of cotton linter and pulp solidified in a plate-like form.) having 2.5 cm × 1.5 cm and a thickness of 0.3 cm to give an electric mosquito mat.

Formulation Example 11. Fumigant

After 100 mg of Compound (12) of the present invention is dissolved in a appropriate amount of acetone, the solution is impregnated with a porous ceramic plate having 4.0 cm × 4.0 cm and a thickness of 1.2 cm to give a fumigant.

Formulation Example 12. Poison bait

After 10 mg of Compound (1) through (13) of the present invention is dissolved in a 0.5 ml acetone, the solution is applied to 5 g of the powder of dry animal food. The powder is dried to give a 0.5% poison bait.

Next, effectiveness of the compounds of the present invention as the active ingredient of insecticidal compositions is described below, with reference to test examples, wherein the compounds of the present invention are designated by the compound numbers shown in Table 2 and compounds used for comparison and control are designated by the compound numbers shown in Table 3.

Table 3

Compound Symbol	Chemical Structure	Note
(A)		Compound described in U.S. Patent 3,868,458 and 3,940,484
(B)		Compound described in U.S. Patent 3,996,366

Test Example 1 (Insecticidal test on nymphs of brown planthopper)

The emulsifiable concentrate of the test compound prepared according to Formulation Example 1 was diluted with water (corresponding to 500, 5, 0.5 ppm) and a rice plant seedling (length of about 12 cm) was immersed in the dilution for a minute. After air-drying, the rice plant seedling was put in a test tube and about 30 nymphs of brown planthopper (*Nilaparvata lugens*) were released. Six days after, the nymphs were observed if they were alive or dead. Criterion for the judgment is as follows.

- a: no insect was alive.
- b: alive insects were 5 or less.
- c: alive insects were 6 or more.

The results are shown in Table 4.

Table 4

	<u>Test Compound</u>	<u>Concentration (ppm)</u>	<u>Efficacy</u>
5	(1)	500	a
	(2)	500	a
		5	a
10	(3)	500	a
		5	a
	(4)	500	a
15	(5)	500	a
		5	a
	(6)	500	a
20	(7)	500	a
	(8)	500	a
25	(9)	500	a
		5	a
		0.5	a
	(10)	500	a
30		5	a
		0.5	a
	(11)	500	a
35		5	a
		0.5	a
	(12)	500	a
		5	a
40		0.5	a
	(13)	500	a
45	Untreated	—	c

Test Example 2 (Insecticidal test on southern corn rootworm)

On the bottom of a polyethylene cup having a diameter of 5.5 cm, a filter paper which is of the same size was laid down and 1 ml of an aqueous dilution (500 or 50 ppm) of the emulsifiable concentrate of the test compound prepared according to Formulation Example 1 was dropped onto the filter paper and one corn sprout was put as feed. About 30 eggs of southern corn rootworm (*Diabrotica undecimpunctata*) were put in the cup. Eight days after the cup was covered, dead or alive larvae hatched were examined. Criterion for the judgment is as follows.

- a: no insect was alive.
 b: alive insects were 5 or less.
 c: alive insects were 6 or more.

The results are shown in Table 5.

Table 5

	<u>Test Compound</u>	<u>Concentration (ppm)</u>	<u>Efficacy</u>
5	(1)	500	a
		50	a
10	(2)	500	a
		50	a
	(3)	500	a
		50	a
15	(4)	500	a
		50	a
	(5)	500	a
20		50	a
	(6)	500	a
		50	a
25	(7)	500	a
		50	a
	(8)	500	a
30		50	a
	(9)	500	a
		50	a
35	(10)	500	a
		50	a
	(11)	500	a
		50	a
40	(12)	500	a
		50	a
45	(13)	500	a
		50	a
	Untreated		c

50 Test Example 3 (Insecticidal test on common mosquito)

The emulsifiable concentrate of the test compound prepared according to Formulation Example 1 was diluted with water and 0.7 ml of the dilution was added to 100 ml of ion exchange water (concentration of the effective ingredient was 3.5 ppm). In the mixture were released 20 last instar larvae of common mosquito (Culex pipiens pallens). One day after the release, mortality was examined.

55 Criterion for the judgment is as follows.

- a: 90% or more
b: not less than 10% but less than 90%

c: less than 10%

The results are shown in Table 6.

Table 6

	<u>Test Compound</u>	<u>Efficacy</u>
5		
10	(1)	a
	(2)	a
	(3)	a
15	(4)	a
	(5)	a
20	(6)	a
	(7)	a
	(8)	a
25	(9)	a
	(10)	a
	(11)	a
30	(12)	a
	(13)	a
35	Untreated	c

Test Example 4 (Insecticidal test on German cockroach)

On the bottom of a polyethylene cup having a diameter of 5.5 cm, a filter paper which is of the same size was laid down and 0.7 ml of an aqueous dilution (500 ppm) of the emulsifiable concentrate of the test compound prepared according to Formulation Example 1 was dropped onto the filter paper. As feed, 30 mg of sucrose was uniformly spread thereon. In the cup, 10 adult males of German cockroach (*Blattella germanica*) were released. Six days after the cup was covered, dead or alive insects were examined to determine mortality.

The results are shown in Table 7.

Table 7

5	<u>Test Compound</u>	<u>Mortality (%)</u> <u>500 ppm</u>
	(1)	100
10	(2)	100
	(3)	100
	(4)	100
15	(5)	100
	(6)	100
20	(7)	100
	(8)	100
	(9)	100
25	(10)	100
	(11)	100
	(12)	100
30	(13)	100
	(A)	0
35	(B)	0
	Untreated	0

Test Example 5 (Insecticidal test on housefly)


On the bottom of a polyethylene cup having a diameter of 5.5 cm, a filter paper which is of the same size was laid down and 0.7 ml of an aqueous dilution (500 ppm) of the emulsifiable concentrate of the test compound prepared according to Formulation Example 1 was dropped onto the filter paper. As feed, 30 mg of sucrose was uniformly spread thereon. In the cup, 10 adult females of housefly (*Musca domestica*) were released. Forty eight hours after the cup was covered, dead or alive insects were examined to determine mortality (2 replications). The results are shown in Table 8.

Table 8

5	Test Compound	Mortality (%)
		500 ppm
	(1)	100
10	(2)	100
	(3)	100
	(4)	100
15	(5)	100
	(6)	100
	(7)	100
20	(8)	100
	(9)	100
25	(10)	100
	(11)	100
	(12)	100
30	(13)	100
	Untreated	0

Claims

1. A pyridylimidazole derivative having the formula:



wherein R¹ is a hydrogen atom, a C₁-C₃ alkyl group, a C₁-C₃ alkylthio group or a C₂-C₃ alkoxyalkyl group; R² is a hydrogen atom or a C₁-C₄ haloalkyl group; R³ is a halogen atom, a nitro group or a trifluoromethyl group; R⁴ is a C₁-C₃ haloalkyl group or a C₁-C₃ haloalkoxy group.


2. A pyridylimidazole derivative according to claim 1, wherein R¹ is a hydrogen atom or a C₁-C₃ alkyl group; R² is a C₁-C₃ haloalkyl group which comprises at least a fluorine atom, a chlorine atom or a bromine atom as the halogen atom; R³ is a fluorine atom, a chlorine atom or a trifluoromethyl group; R⁴ is a C₁-C₃ ha-

Table 8

<u>Test Compound</u>	<u>Mortality (%)</u> <u>500 ppm</u>
(1)	100
(2)	100
(3)	100
(4)	100
(5)	100
(6)	100
(7)	100
(8)	100
(9)	100
(10)	100
(11)	100
(12)	100
(13)	100
Untreated	0

Claims

1. A pyridylimidazole derivative having the formula:




wherein R¹ is a hydrogen atom, a C₁-C₃ alkyl group, a C₁-C₃ alkylthio group or a C₂-C₃ alkoxyalkyl group; R² is a hydrogen atom or a C₁-C₄ haloalkyl group; R³ is a halogen atom, a nitro group or a trifluoromethyl group; R⁴ is a C₁-C₃ haloalkyl group or a C₁-C₃ haloalkoxy group.

2. A pyridylimidazole derivative according to claim 1, wherein R¹ is a hydrogen atom or a C₁-C₃ alkyl group; R² is a C₁-C₃ haloalkyl group which comprises at least a fluorine atom, a chlorine atom or a bromine atom as the halogen atom; R³ is a fluorine atom, a chlorine atom or a trifluoromethyl group; R⁴ is a C₁-C₃ ha-




loalkyl group which comprises at least a fluorine atom as the halogen atom.

3. A pyridylimidazole derivative according to claim 1, wherein R¹ is a hydrogen atom or a methyl group; R² is a C₂ haloalkyl group which comprises at least a fluorine atom, a chlorine atom or a bromine atom as the halogen atom; R³ is a fluorine atom or a chlorine atom; R⁴ is a trifluoromethyl group.
4. A pyridylimidazole derivative according to claim 1, wherein R¹ is a hydrogen atom or a methyl group; R² is a haloalkyl group represented by the formula, -CF₂CF₂X, in which X is a hydrogen atom, a fluorine atom, a chlorine atom or a bromine atom; R³ is a chlorine atom; R⁴ is a trifluoromethyl group.
5. A pyridylimidazole derivative according to claim 1, wherein R¹ is a methyl group; R² is a haloalkyl group represented by the formula, -CF₂CF₂X, in which X is a hydrogen atom, a fluorine atom, a chlorine atom or a bromine atom; R³ is a chlorine atom; R⁴ is a trifluoromethyl group.
6. A pyridylimidazole derivative according to claim 1, which is 1-(3-chloro-5-trifluoromethylpyridin-2-yl)-4-(1,1,2,2-tetrafluoroethyl)imidazole,
1-(3-chloro-5-trifluoromethylpyridin-2-yl)-4-pentafluoroethylimidazole,
1-(3-chloro-5-trifluoromethylpyridin-2-yl)-4-(2-bromo-1,1,2,2-tetrafluoroethyl)imidazole,
1-(3-chloro-5-trifluoromethylpyridin-2-yl)-2-methyl-4-(1,1,2,2-tetrafluoroethyl)imidazole,
1-(3-chloro-5-trifluoromethylpyridin-2-yl)-2-methyl-4-pentafluoroethylimidazole,
1-(3-chloro-5-trifluoromethylpyridin-2-yl)-2-methyl-4-(2-chloro-1,1,2,2-tetrafluoroethyl)imidazole, or
1-(3-chloro-5-trifluoromethylpyridin-2-yl)-2-methyl-4-(2-bromo-1,1,2,2-tetrafluoroethyl)imidazole.
7. A process for producing a pyridylimidazole derivative according to claim 1, which comprises reacting a halide compound having the formula;



wherein R³ is a halogen atom, a nitro group or a trifluoromethyl group; R⁴ is a C₁-C₃ haloalkyl group or a C₁-C₃ haloalkoxy group and A is a halogen atom, with an imidazole derivative having the formula;




wherein R¹ is a hydrogen atom, a C₁-C₃ alkyl group, a C₁-C₃ alkylthio group or a C₂-C₃ alkoxyalkyl group and R² is a hydrogen atom or a C₁-C₄ haloalkyl group.

8. An insecticidal composition which comprises a pyridylimidazole derivative according to claim 1 and an inert carrier.
9. A method for controlling insect pests which comprises applying a pyridylimidazole derivative according to claim 1 to the insect pests or to a locus where insect pests may propagate.
10. Use of a pyridylimidazole derivative according to claim 1 as an insecticide.

Patentansprüche


1. Pyridylimidazolderivat der Formel






in der R¹ ein Wasserstoffatom, einen C₁-C₃-Alkyl-, C₁-C₃-Alkylthio- oder C₂-C₃-Alkoxyalkylrest bedeutet, R² ein Wasserstoffatom oder einen C₁-C₄-Haloalkylrest bedeutet, R³ ein Halogenatom, eine Nitro- oder Trifluormethylgruppe bedeutet, R⁴ einen C₁-C₃-Haloalkyl- oder C₁-C₃-Haloalkoxyrest bedeutet.

2. Pyridylimidazolderivat nach Anspruch 1, wobei R¹ ein Wasserstoffatom oder einen C₁-C₃-Alkylrest bedeutet, R² einen C₁-C₃-Haloalkylrest bedeutet, der als Halogenatom mindestens ein Fluor-, Chlor- oder Bromatom umfaßt, R³ ein Fluor- oder Chloratom oder eine Trifluormethylgruppe bedeutet, R⁴ einen C₁-C₃-Haloalkylrest bedeutet, der als Halogenatom mindestens ein Fluoratom umfaßt.
3. Pyridylimidazolderivat nach Anspruch 1, wobei R¹ ein Wasserstoffatom oder eine Methylgruppe bedeutet, R² einen C₂-Haloalkylrest bedeutet, der als Halogenatom mindestens ein Fluor-, Chlor- oder Bromatom umfaßt, R³ ein Fluor- oder Chloratom bedeutet, R⁴ eine Trifluormethylgruppe bedeutet.
4. Pyridylimidazolderivat nach Anspruch 1, wobei R¹ ein Wasserstoffatom oder eine Methylgruppe bedeutet, R² einen Haloalkylrest der Formel -CF₂CF₂X bedeutet, in der X ein Wasserstoff-, Fluor-, Chlor- oder Bromatom bedeutet, R³ ein Chloratom bedeutet, R⁴ eine Trifluormethylgruppe bedeutet.
5. Pyridylimidazolderivat nach Anspruch 1, wobei R¹ eine Methylgruppe bedeutet, R² einen Haloalkylrest der Formel -CF₂CF₂X bedeutet, in dem X ein Wasserstoff-, Fluor-, Chlor- oder Bromatom bedeutet, R³ ein Chloratom bedeutet, R⁴ eine Trifluormethylgruppe bedeutet.
6. Pyridylimidazolderivat nach Anspruch 1, nämlich
 - 1-(3-Chlor-5-trifluormethylpyridin-2-yl)-4-(1,1,2,2-tetrafluorethyl)imidazol,
 - 1-(3-Chlor-5-trifluormethylpyridin-2-yl)-4-pentafluorethylimidazol,
 - 1-(3-Chlor-5-trifluormethylpyridin-2-yl)-4-(2-brom-1,1,2,2-tetrafluorethyl)imidazol,
 - 1-(3-Chlor-5-trifluormethylpyridin-2-yl)-2-methyl-4-(1,1,2,2-tetrafluorethyl)imidazol,
 - 1-(3-Chlor-5-trifluormethylpyridin-2-yl)-2-methyl-4-pentafluorethylimidazol,
 - 1-(3-Chlor-5-trifluormethylpyridin-2-yl)-2-methyl-4-(2-chlor-1,1,2,2-tetrafluorethyl)imidazol oder
 - 1-(3-Chlor-5-trifluormethylpyridin-2-yl)-2-methyl-4-(2-brom-1,1,2,2-tetrafluorethyl)imidazol.
7. Verfahren zur Herstellung eines Pyridylimidazolderivats nach Anspruch 1, umfassend die Umsetzung einer Halogenverbindung der Formel



in der R³ ein Halogenatom, eine Nitro- oder Trifluormethylgruppe bedeutet, R⁴ einen C₁-C₃-Haloalkyl- oder C₁-C₃-Haloalkoxyrest bedeutet und A ein Halogenatom bedeutet, mit einem Imidazolderivat der Formel






in der R¹ ein Wasserstoffatom, einen C₁-C₃-Alkyl-, C₁-C₃-Alkylthio- oder C₂-C₃-Alkoxyalkylrest bedeutet und R² ein Wasserstoffatom oder einen C₁-C₄-Haloalkylrest bedeutet.

8. Insektizide Zusammensetzung, umfassend ein Pyridylimidazolderivat nach Anspruch 1 und einen inerten Träger.
9. Verfahren zur Bekämpfung von schädlichen Insekten, umfassend die Anwendung eines Pyridylimidazolderivats nach Anspruch 1 gegen die Insekten oder an Orten, an denen sich die Insekten vermehren können.
10. Verwendung eines Pyridylimidazolderivats nach Anspruch 1 als Insektizid.

Revendications

1. Dérivé de pyridylimidazole de formule:




dans laquelle R¹ est un atome d'hydrogène, un groupement alkyle en C₁-C₃, un groupement alkylthio en C₁-C₃ ou un groupement alkoxyalkyle en C₂-C₃; R² est un atome d'hydrogène ou un groupement halogénoalkyle en C₁-C₄; R³ est un atome d'halogène, un groupement nitro ou un groupement trifluorométhyle; R⁴ est un groupement halogénoalkyle en C₁-C₃ ou un groupement halogénoalcoxy en C₁-C₃.

2. Dérivé de pyridylimidazole selon la revendication 1, dans lequel R¹ est un atome d'hydrogène ou un groupement alkyle en C₁-C₃; R² est un groupement halogénoalkyle en C₁-C₃ qui comporte au moins un atome de fluor, un atome de chlore ou un atome de brome en tant qu'atome d'halogène; R³ est un atome de fluor, un atome de chlore ou un groupement trifluorométhyle; R⁴ est un groupement halogénoalkyle en C₁-C₃ qui comporte au moins un atome de fluor en tant qu'atome d'halogène.
3. Dérivé de pyridylimidazole selon la revendication 1, dans lequel R¹ est un atome d'hydrogène ou un groupement méthyle; R² est un groupement halogénoalkyle en C₂ qui contient au moins un atome de fluor, un atome de chlore ou un atome de brome en tant qu'atome d'halogène; R³ est un atome de fluor ou un atome de chlore; R⁴ est un groupement trifluorométhyle.
4. Dérivé de pyridylimidazole selon la revendication 1, dans lequel R¹ est un atome d'hydrogène ou un groupement méthyle; R² est un groupement halogénoalkyle représenté par la formule -CF₂CF₂X dans laquelle X est un atome d'hydrogène, un atome de fluor, un atome de chlore ou un atome de brome; R³ est un atome de chlore; R⁴ est un groupement trifluorométhyle.
5. Dérivé de pyridylimidazole selon la revendication 1, dans lequel R¹ est un groupement méthyle; R² est un groupement halogénoalkyle représenté par la formule -CF₂CF₂X dans laquelle X est un atome d'hydrogène, un atome de fluor, un atome de chlore ou un atome de brome; R³ est un atome de chlore; R⁴




est un groupement trifluorométhyle.

6. Dérivé de pyridylimidazole selon la revendication 1, qui est le 1-(3-chloro-5-trifluorométhylpyridin-2-yl)-4-(1,1,2,2-tétrafluoroéthyl)imidazole,
 le 1-(3-chloro-5-trifluorométhylpyridin-2-yl)-4-pentafluoroéthylimidazole,
 le 1-(3-chloro-5-trifluorométhylpyridin-2-yl)-4-(2-bromo-1,1,2,2-tétrafluoroéthyl)imidazole,
 le 1-(3-chloro-5-trifluorométhylpyridin-2-yl)-2-méthyl-4-(1,1,2,2-tétrafluoroéthyl)imidazole,
 le 1-(3-chloro-5-trifluorométhylpyridin-2-yl)-2-méthyl-4-pentafluoroéthylimidazole,
 le 1-(3-chloro-5-trifluorométhylpyridin-2-yl)-2-méthyl-4-(2-chloro-1,1,2,2-tétrafluoroéthyl)imidazole, ou
 le 1-(3-chloro-5-trifluorométhylpyridin-2-yl)-2-méthyl-4-(2-bromo-1,1,2,2-tétrafluoroéthyl)imidazole.
7. Procédé de production d'un dérivé de pyridylimidazole selon la revendication 1, qui consiste à faire réagir un halogénure de formule



dans laquelle R³ est un atome d'halogène, un groupement nitro ou un groupement trifluorométhyle; R⁴ est un groupement halogénoalkyle en C₁-C₃ ou un groupement halogénoalcoxy en C₁-C₃ et A est un atome d'halogène, avec un dérivé d'imidazole de formule



dans laquelle R¹ est un atome d'hydrogène, un groupement alkyle en C₁-C₃, un groupement alkylthio en C₁-C₃ ou un groupement alcoxyalkyle en C₂-C₃, et R² est un atome d'hydrogène ou un groupement halogénoalkyle en C₁-C₄.

8. Composition insecticide qui contient un dérivé de pyridylimidazole selon la revendication 1 et un support inerte.
9. Procédé de lutte contre des insectes parasites qui consiste à appliquer un dérivé de pyridylimidazole selon la revendication 1 aux insectes parasites ou à un lieu où peuvent se propager les insectes parasites.
10. Utilisation d'un dérivé de pyridylimidazole selon la revendication 1 en tant qu'insecticide.

