

CLAIMS:

1. A recombinant adeno-associated virus (rAAV) having an AAV capsid and a vector genome comprising a frataxin (FXN) gene having the sequence of SEQ ID NO: 3 or a sequence at least 95% identical thereto that encodes human frataxin protein having a sequence of SEQ ID NO: 2, and regulatory sequences operably linked to the FXN gene which direct expression of the FXN gene in targeted human cells.
2. The rAAV according to claim 1, wherein
 - (i) the vector genome comprises an AAV2 5' inverted terminal repeat (ITR), cytomegalovirus immediate early (CMV IE) enhancer, a chicken beta actin promoter, a chimeric intron comprising a chicken beta actin splice donor and a rabbit beta-globin acceptor element, the FXN gene, a polyadenylation signal sequence (polyA), and an AAV2 3' ITR, optionally comprising the sequence of nucleotides 198 to 2737 of SEQ ID NO: 8 or nucleotides 198 to 2736 of SEQ ID NO: 12; and/or
 - (ii) the vector genome further comprises at least one, at least two, or at least three tandem repeats of dorsal root ganglion (DRG)-specific miRNA targeted sequences.
3. The rAAV according to claim 2, wherein the at least two or at least three miRNA target sequences are the same.
4. The rAAV according to claim 2, wherein the polyA is rabbit beta globin polyA.
5. The rAAV according to any one of claims 1 to 4, wherein the AAV capsid is an AAVrh91 capsid, or an AAV clade F capsid.
6. The rAAV according to claim 5, wherein the clade F capsid is an AAVhu68 capsid.
7. An aqueous pharmaceutical composition comprising a formulation buffer and a stock of the rAAV according to any one of claims 1 to 6.

8. The pharmaceutical composition according to claim 7, wherein the formulation buffer comprises:

an artificial cerebrospinal fluid comprising buffered saline and one or more of sodium, calcium, magnesium, potassium, or mixtures thereof; and
a surfactant.

9. The pharmaceutical composition according to claim 8, wherein the surfactant is present at 0.0005 % w/w to 0.001% w/w of the pharmaceutical composition, and/or wherein the composition is at a pH in the range of 7.5 to 7.8, or 6.2 to 7.7, or 7.

10. The rAAV according to any one of claims 1 to 6 or the pharmaceutical composition according to any one of claims 7 to 9, wherein the rAAV or pharmaceutical composition is suitable for intravenous administration, intrathecal administration, and/or intraparenchymal (dentate nucleus) administration.

11. A first rAAV according to any one of claims 1 to 6 and a second rAAV according to any one of claims 1 to 6 for use in a treatment regimen for Friedreich's ataxia (FRDA) comprising dual-route of administration of rAAV, the regimen comprising of intravenous administration of said first rAAV for intravenous administration and said second rAAV for intraparenchymal (dentate nucleus (and intraparenchymal (dentate nucleus (IDN))) or intrathecal administration .

12. Use of a first rAAV according to any one of claims 1 to 6 and of a second rAAV according to any one of claims 1 to 6 in the preparation of a medicament for a treatment regimen for Friedreich's ataxia (FRDA) comprising dual-route of administration of rAAV, the regimen comprising said first rAAV for intravenous administration and said second rAAV for intraparenchymal (dentate nucleus) or intrathecal administration.

13. The first rAAV and second rAAV according to claim 11 or the use according to claims 12, wherein the intraparenchymal (dentate nucleus (IDN)) administration is to be performed unilaterally or bilaterally.

14. The first rAAV and second rAAV according to claim 11 or the use according to claim 12, wherein the intravenous and intraparenchymal (dentate nucleus (IDN)) administrations of the rAAV are to be performed sequentially and within a 24-hour period.

15. An rAAV according to any one of claims 1 to 6 or a pharmaceutical composition according to any one of claims 7 to 9 for use in treatment of Friedreich's ataxia (FRDA), the treatment comprising a dual-route administration according to any one of claims 11, 13 to 14, wherein an amelioration of FRDA includes reduction of cardiac and/or neurological symptoms.

16. The use according to any one of claims 12 to 14, wherein an amelioration of FRDA includes reduction of cardiac and/or neurological symptoms.

17. The rAAV according to any one of claims 1 to 6 or the pharmaceutical composition according to any one of claims 7 to 9 for use in treating FRDA in a subject.

18. Use of the rAAV according to any one of claims 1 to 6 or the pharmaceutical composition according to any one of claims 7 to 9 in the preparation of a medicament for treating FRDA in a subject.

19. The rAAV or the pharmaceutical composition according to claim 17 or the use according to claim 18, wherein the rAAV or the pharmaceutical composition is to be administered intravenously, intrathecally, or via intraparenchymal injection.

20. The rAAV or the pharmaceutical composition according to claim 17 and 19, or the use according to claims 18 to 19, wherein the rAAV is to be administered by a first route of administration and by a second route of administration.

21. The rAAV, pharmaceutical composition or use of claim 20, wherein the first route is intravenous and the second route is intrathecal, or wherein the first route is intravenous and the second route is intraparenchymal.

22. The rAAV, pharmaceutical composition or use according to claim 21, wherein the intraparenchymal (dentate nucleus (IDN)) administration is to be performed unilaterally or bilaterally, and/or wherein the amount of dose of rAAV to be administered intravenously and the amount of rAAV to be administered IDN is a ratio of 10 to 1.

23. The rAAV, pharmaceutical composition or use according to any one of claims 17 to 22, wherein the first route and second route are to be administered simultaneously or sequentially and/or wherein the first route and second route are administered within 24 hours of one another.

24. The rAAV, pharmaceutical composition or use according to any one of claims 17 to 23, wherein treating FRDA comprises ameliorating one or more symptoms of FRDA in the subject.

25. The rAAV, pharmaceutical composition or use of claim 24, wherein the amelioration of one or more symptoms of FRDA includes increased average life span, reduction in progression towards neuromuscular decline, improvement in neuromuscular development, reduction in progression towards cardiomyopathy, or improvement in cardiac symptoms, or any combination thereof.

26. The rAAV, pharmaceutical composition or use of claim 25, wherein the improvement in cardiac symptoms is a reduction or elimination of an arrhythmia in the subject.

27. A plasmid comprising

(i) an expression cassette comprising a frataxin (FXN) gene having the sequence of SEQ ID NO: 3 or a sequence 95% identical thereto that encodes human frataxin protein having a sequence of SEQ ID NO: 2, or

(ii) the vector genome of SEQ ID NO: 8 or 12.

28. The plasmid according to claim 27, the expression cassette further comprising a promoter comprising a cytomegalovirus immediate early (CMV IE) enhancer, a chicken beta-actin promoter, a chimeric intron comprising a chicken beta actin splice donor and a rabbit beta-globin splice acceptor element, and a polyA and/or wherein the expression cassette is flanked by an AAV2 5' ITR and an AAV2 3' ITR.

29. A host cell comprising the plasmid according to any one of claims 27 to 28.