

Claims

1. A compound of formula (I):



(I)

and/or a pharmaceutically acceptable salt thereof, and/or solvates, racemic mixtures, enantiomers, diasteromers, and tautomers thereof, wherein

A is  ;

R_1 is chosen from H, -OH, halo, C_{1-6} alkyl, C_{1-6} alkoxyl, $-NH_2$, $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl$)_2$, oxo, or C_{3-8} cycloalkyl;

each of R_2 is independently chosen from H, deuterium, halo, -OH, $-NH_2$, -CN, -SH, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, C_{3-8} cycloalkyl, oxo, $-OR_5$, $-OCOR_5$, $-NHR_5$, $-N(R_5)(C_{1-4}$ alkyl), $-COR_5$, $-NHCOR_5$, or 3-8 membered heterocyclyl containing one or more heteroatoms independently chosen from N, O, and S; in which each of said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-8} cycloalkyl or 3-8 membered heterocyclyl containing one or more heteroatoms independently chosen from N, O, and S is optionally substituted with one or more groups chosen from deuterium, halo, -CN, -OH, -SH, $-NH_2$, $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl$)_2$, or C_{1-6} alkoxyl; or two R_2, which attach to the same carbon atom, together with the carbon atom they are attached to form a 3-5 membered cycloalkyl which is optionally substituted with one or more halo or deuterium;

R_3' and R_4' are both H;

R_3 and R_4 are independently chosen from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-12} cycloalkyl, 3-12 membered heterocyclyl containing one or more heteroatoms independently chosen from N, O, and S, phenyl, 5-12 membered heteroaryl containing one or more heteroatoms independently chosen from N, O, and S, $-C(O)R_5$, $-OR_5$, or $-NHR_5$, in which each of said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-12} cycloalkyl,

3-12 membered heterocyclyl containing one or more heteroatoms independently chosen from N, O, and S, phenyl, or 5-12 membered heteroaryl containing one or more heteroatoms independently chosen from N, O, and S is optionally substituted with one or more R_6; provided that R_3 and R_4 are not H simultaneously; provided that when one of R_3 and R_4 is optionally substituted phenyl or optionally substituted 5-6 membered heteroaryl containing one or more heteroatoms independently chosen from N, O, and S, the other one is $-OR_5$ or $-NHR_5$;

or R_3 and R_3' are independently chosen from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-12} cycloalkyl, 3-12 membered heterocyclyl containing one or more heteroatoms independently chosen from N, O, and S, phenyl, 5-12 membered heteroaryl containing one or more heteroatoms independently chosen from N, O, and S, $-C(O)R_5$, $-OR_5$, or $-NHR_5$, in which each of said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-12} cycloalkyl, 3-12 membered heterocyclyl containing one or more heteroatoms independently chosen from N, O, and S, phenyl, or 5-12 membered heteroaryl containing one or more heteroatoms independently chosen from N, O, and S is optionally substituted with one or more R_6; R_4 and R_4' together with the N atom they are attached to form a 3-8 membered heterocyclic ring containing one or more heteroatoms independently chosen from N, O, and S optionally substituted by one or more R_6;

R_5 is chosen from C_{1-6} alkyl or C_{3-8} cycloalkyl, each of which is optionally substituted with one or more groups independently chosen from halo, -CN, -OH, -SH, $-NH_2$, or C_{1-6} alkoxyl;

each of R_6 is independently chosen from deuterium, halo, -CN, -OH, -SH, $-NH_2$, C_{1-6} alkoxyl, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-8} cycloalkyl, 3-8 membered heterocyclyl containing one or more heteroatoms independently chosen from N, O, and S, phenyl, or 5-6 membered heteroaryl containing one or more heteroatoms independently chosen from N, O, and S, in which each of said C_{1-6} alkoxyl, C_{1-6} alkyl, C_{3-8} cycloalkyl, 3-8 membered heterocyclyl containing one or more heteroatoms independently chosen from N, O, and S, phenyl, or 5-6 membered heteroaryl containing one or more heteroatoms independently chosen from N, O, and S is optionally substituted with one or more groups independently chosen from halo, -CN, -OH, -SH, $-NH_2$, C_{1-6} alkoxyl, C_{2-6} alkynyl, or C_{1-6} alkyl;

m is 0, 1, 2, 3, 4, 5, or 6;

n is 0, 1, or 2.

2. The compound of formula (I) according to claim 1, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, R_1 is chosen from H, -OH or halo.

3. The compound of formula (I) according to claim 2, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, R_1 is -OH.

4. The compound of formula (I) according to claim 1, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, each of R_2 is independently chosen from H, deuterium, halo, -OH, $-NH_2$, -CN, -SH, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, C_{3-8} cycloalkyl, oxo, $-OR_5$, $-OCOR_5$, $-NHR_5$, $-N(R_5)(C_{1-4}$ alkyl), $-NHCOR_5$, or 3-8 membered heterocyclyl containing one or more heteroatoms independently chosen from N, O, and S.

5. The compound of formula (I) according to claim 4, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, each of R_2 is independently chosen from H, deuterium, halo, C_{1-6} alkyl, or C_{1-6} haloalkyl.

6. The compound of formula (I) according to claim 1, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, R_3 and R_4 are independently chosen from C_{1-6} alkyl, C_{3-12} cycloalkyl, 3-12 membered heterocyclyl containing one or more heteroatoms independently chosen from N, O, and S, phenyl, 5-12 membered heteroaryl containing one or more heteroatoms independently chosen from N, O, and S, $-C(O)R_5$, $-OR_5$, or $-NHR_5$, in which each of said C_{1-6} alkyl, C_{3-12} cycloalkyl, 3-12 membered heterocyclyl

containing one or more heteroatoms independently chosen from N, O, and S, phenyl, or 5-12 membered heteroaryl containing one or more heteroatoms independently chosen from N, O, and S is optionally substituted with one or more R_6.

7. The compound of formula (I) according to claim 6, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, R_3 and R_4 are independently chosen from C_{1-6} alkyl substituted with one or more halo, 5-12 membered heteroaryl containing one or more heteroatoms independently chosen from N, O, and S substituted with C_{1-6} haloalkyl, or $-OR_5$.

8. The compound of formula (I) according to claim 1, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, R_3 and R_4 are independently chosen from C_{1-6} alkyl substituted with one or more halo.

9. The compound of formula (I) according to claim 1, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, R_5 is C_{1-6} alkyl optionally substituted with one or more halo.


10. The compound of formula (I) according to claim 1, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, each of R_6 is independently chosen from deuterium, halo, -CN, -OH, $-NH_2$, C_{1-6} alkoxyl, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-8} cycloalkyl, 3-8 membered heterocyclyl containing one or more heteroatoms independently chosen from N, O, and S, phenyl, or 5-6 membered heteroaryl containing one or more heteroatoms independently chosen from N, O, and S, in which each of said C_{1-6} alkoxyl, C_{1-6} alkyl, C_{3-8} cycloalkyl, 3-8 membered heterocyclyl containing one or more heteroatoms independently chosen from N, O, and S, phenyl, or 5-6 membered heteroaryl containing

one or more heteroatoms independently chosen from N, O, and S is optionally substituted with one or more halo.

11. The compound of formula (I) according to claim 1, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, n is 1.


12. The compound of formula (I) according to claim 1, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, R_3 is chosen from H, C_{1-6} alkyl optionally substituted by C_{1-6} haloalkyl, or 5-12 membered heteroaryl containing one or more heteroatoms independently chosen from N, O, and S optionally substituted by C_{1-6} haloalkyl; R_3'is H; R_4 and R_4' together with the N atom they are attached to form a 3-8 membered heterocyclic ring containing one or more heteroatoms independently chosen from N, O, and S optionally substituted by one or more groups chosen from halo, -OH, or C_{1-6} haloalkyl.

13. The compound of formula (I) according to any one of claims 1-12, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, the compound of formula (I) has the structure of formula (I-1)




(I-1) .

14. The compound of formula (I) according to any one of claims 1-12, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, the compound of formula (I) has the structure of formula (II); wherein X is halo; p is 0, 1, or 2; m is 0, 1, or 2;



(II).



15. The compound of formula (I) according to claim 14, or a pharmaceutically acceptable salts thereof, and/or solvates, racemic mixture, enantiomers, diasteromers, and tautomers thereof, wherein, the compound of formula (I) has the structure of formula (II-1); wherein X is halo; p is 0, 1, or 2; m is 0, 1, or 2;



































(II-1)

















.

















16. A compound selected from:

















Compound	Structure	Compound	Structure
1		152	















2		153	
3		154	
4		155	
5		156	
6		157	
7		158	
8		159	
9		160	




10		161	
11		162	
12		163	
13		164	
14		165	
15		166	
16		167	
17		168	














18		169	
19		170	
20		171	
21		172	
22		173	
23		174	
24		175	
25		176	















26		177	
27		178	
28		179	
29		180	
30		181	
31		182	
32		183	
33		184	














34		186	
35		187	
36		188	
37		189	
38		190	
39		191	
40		192	
41		193	














42		194	
43		195	
44		196	
45		197 & 198	
46			
47		199	
48		200	














49		201 & 202	
50			
51		203	
52		204	
53		205	
		206	
55		207	






56		208	
57		209	
		210	
59		211	
60		212	
61		213	
62		214	














63		215	
64		216	
65		217	
66		218	
67		219	
68		220	
69		221	

















		222 & 223	
71			
72		224	
73		225	
74		226	
75		227	
76		228	



77		229	
78		230 & 231	
79			
80		232	
81		233	
82			
83		235	















84		236 & 237	
85			
		238	
87		239	
89		240	
90		241	
91		242	















92		243	
93		244	
94			















95 & 96		246 & 247	
			
97& 98			
		249 & 250	
99			
100 & 101		251 & 252	
			















102 & 103		253 & 254	
			
104 & 105		255 & 256	
			
106 & 107		257 & 258	
			
108 & 109		259	
		260	












110 & 111		261	
		262 & 263	
112 & 113			
		264 & 265	
114 & 115			
		266 267 268 269	
116 & 117			

118 & 119			
		270	
120 & 121		271	
		272 & 273	
122			
123		274	
124		275	

125		276	
126		277	
127		278	
128		279	
129		280	
130		281	
131		282	

132		283	
133		284	
134		285 & 286	
135			
136		287 & 288	
137			
138		289 & 290	

139			
140		291 & 292	
141			
142		293	
143		294	
144		295	
145		296	

146 & 147		297	
		298	
148 & 149		299	
		300	
150 & 151		301	
			

or a pharmaceutically acceptable salt thereof.

17. A pharmaceutical composition comprising the compound of any one of claims 1-16, or a pharmaceutically acceptable salt thereof, and optionally comprising at least one pharmaceutically acceptable excipient.

18. A use of a compound of any one of claims 1-16 or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating a disease induced by IDH mutation in a subject wherein, the disease induced by IDH mutation is cancer, which is chosen from solid tumors, neurogliocytoma, or hematological malignant tumor.

19. The use of claim 18, wherein the cancer is chosen from leukemia, lymphoma, or myeloma.

20. The use of claim 18, wherein, the cancer is chosen from acute myeloid leukemia (AML), acute promyelocytic leukemia (APL), glioblastoma (GBM), myelodysplastic syndrome (MDS), myeloproliterative neoplasms (MPN), cholangiocarcinoma, chondrosarcoma, giant cell tumor, intestinal cancer, melanoma, lung cancer, or non-Hodgkin's lymphoma (NHL).

21. The use of claim 20, wherein the cancer is intrahepatic cholangiocarcinoma (IHCC).