


CLAIMS

[Claim 1]


A pharmaceutical composition for treating colorectal cancer and/or lung cancer,

the pharmaceutical composition comprising a compound of the following formula (I) or a salt thereof and one or more pharmaceutically acceptable excipients:



wherein

R^1 is the following formula (II) or (III):



R^{1a} and R^{1b} , which are the same as or different from each other, are H or F,


R^2 is halogen, C_{1-3} alkyl, cyclopropyl, or vinyl,

R^3 is the following formula (IV):




R^4 is C_{1-3} alkyl, oxetanyl, tetrahydrofuranyl,


tetrahydropyranyl, optionally substituted pyrazolyl, optionally substituted pyridyl, optionally substituted pyrimidinyl, optionally substituted pyrrolidinyl, or optionally substituted piperidinyl,
 R^5 is ethyl, isopropyl, tert-butyl, or C_{3-6} cycloalkyl,
 R^{6a} and R^{6b} , which are the same as or different from each other, are H or C_{1-3} alkyl optionally substituted with a group selected from the group consisting of F, OH, and $N(CH_3)_2$, or R^{6a} and R^{6b} form cyclopropyl together with the carbon to which they are attached,
 R^7 is H, halogen, or a group selected from the group consisting of the following formulae (VI), (VII), (VIII), and (IX):



R^{7a} is H or C_{1-3} alkyl optionally substituted with OH,
 X is O,
 Y is phenylene or pyridinediyl,
 L is a bond, C_{1-3} alkylene, or $C=O$,
 Z is NH or a group selected from the group consisting of the following formulae (X), (XI), and (XII):



or, $Y-L-Z$ is the following formula (XIII):



[Claim 2]

The pharmaceutical composition according to claim 1, wherein the compound of the formula (I) is selected from the group consisting of

(4R)-1-[(2S)-2-(4-{4-[({6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-{4-[4-(hydroxymethyl)-1,3-thiazol-5-yl]phenyl}ethyl]-L-prolinamide,


(4R)-1-[(2S)-2-(4-{4-[({6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(2-oxo-1,3-oxazolidin-3-yl)phenyl]ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-2-{[1-(2,2-difluoroethyl)piperidin-4-yl]oxy}-7-(6-fluoro-5-methyl-1H-indazol-4-yl)quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(1-methyl-1H-pyrazol-5-yl)phenyl]ethyl]-L-prolinamide, and
 (4R)-1-[(2S)-2-(4-{4-[({6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-[(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl]-4-hydroxy-L-prolinamide.


[Claim 3]

Use of a compound of the following formula (I) or a salt thereof in the manufacture of a pharmaceutical composition for treating colorectal cancer and/or lung cancer:



wherein

R¹ is the following formula (II) or (III):



R^{1a} and R^{1b}, which are the same as or different from each

other, are H or F,

R² is halogen, C₁₋₃ alkyl, cyclopropyl, or vinyl,

R³ is the following formula (IV):







R⁴ is C₁₋₃ alkyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, optionally substituted pyrazolyl, optionally substituted pyridyl, optionally substituted pyrimidinyl, optionally substituted pyrrolidinyl, or optionally substituted piperidinyl,

R⁵ is ethyl, isopropyl, tert-butyl, or C₃₋₆ cycloalkyl,

R^{6a} and R^{6b}, which are the same as or different from each other, are H or C₁₋₃ alkyl optionally substituted with a group selected from the group consisting of F, OH, and N(CH₃)₂, or R^{6a} and R^{6b} form cyclopropyl together with the carbon to which they are attached,

R⁷ is H, halogen, or a group selected from the group consisting of the following formulae (VI), (VII), (VIII), and (IX):




R^{7a} is H or C₁₋₃ alkyl optionally substituted with OH,

X is O,


Y is phenylene or pyridinediyl,

L is a bond, C₁₋₃ alkylene, or C=O,

Z is NH or a group selected from the group consisting of the following formulae (X), (XI), and (XII):




or, Y-L-Z is the following formula (XIII):




[Claim 4]

A compound of the following formula (I) or a salt thereof for use in the treatment of colorectal cancer and/or lung cancer:



wherein

R¹ is the following formula (II) or (III):




R^{1a} and R^{1b}, which are the same as or different from each other, are H or F,

R² is halogen, C₁₋₃ alkyl, cyclopropyl, or vinyl,


R³ is the following formula (IV):




R^4 is C_{1-3} alkyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, optionally substituted pyrazolyl, optionally substituted pyridyl, optionally substituted pyrimidinyl, optionally substituted pyrrolidinyl, or optionally substituted piperidinyl,
 R^5 is ethyl, isopropyl, tert-butyl, or C_{3-6} cycloalkyl,
 R^{6a} and R^{6b} , which are the same as or different from each other, are H or C_{1-3} alkyl optionally substituted with a group selected from the group consisting of F, OH, and $N(CH_3)_2$, or R^{6a} and R^{6b} form cyclopropyl together with the carbon to which they are attached,
 R^7 is H, halogen, or a group selected from the group consisting of the following formulae (VI), (VII), (VIII), and (IX):



R^{7a} is H or C_{1-3} alkyl optionally substituted with OH,
X is O,
Y is phenylene or pyridinediyl,
L is a bond, C_{1-3} alkylene, or $C=O$,
Z is NH or a group selected from the group consisting of the following formulae (X), (XI), and (XII):



or, Y-L-Z is the following formula (XIII):



[Claim 5]

A pharmaceutical composition for treating colorectal cancer and/or lung cancer,

the pharmaceutical composition comprising a compound of the formula (I) or a salt thereof and one or more pharmaceutically acceptable excipients,

wherein the compound of the formula (I) is selected from the group consisting of

(4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-{4-[4-(hydroxymethyl)-1,3-thiazol-5-yl]phenyl}ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-

indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(2-oxo-1,3-oxazolidin-3-yl)phenyl]ethyl}-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-2-{[1-(2,2-difluoroethyl)piperidin-4-yl]oxy}-7-(6-fluoro-5-methyl-1H-indazol-4-yl)quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(1-methyl-1H-pyrazol-5-yl)phenyl]ethyl}-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-{(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl}-4-hydroxy-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide, and

(4R)-1-[(2S)-2-(4-{4-[({6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-[(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl]-4-hydroxy-L-prolinamide.

[Claim 6]

The pharmaceutical composition according to claim 5, wherein the compound of the formula (I) is selected from the group consisting of

(4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-{4-[4-(hydroxymethyl)-1,3-thiazol-5-yl]phenyl}ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-

yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(2-oxo-1,3-oxazolidin-3-yl)phenyl]ethyl}-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-2-{[1-(2,2-difluoroethyl)piperidin-4-yl]oxy}-7-(6-fluoro-5-methyl-1H-indazol-4-yl)quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(1-methyl-1H-pyrazol-5-yl)phenyl]ethyl}-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-{(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl}-4-hydroxy-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-

yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide, and
 (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-{(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl}-4-hydroxy-L-prolinamide.

[Claim 7]

The pharmaceutical composition according to claim 5, wherein the compound of the formula (I) is selected from the group consisting of

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide,
 (4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-{4-[4-(hydroxymethyl)-1,3-thiazol-5-yl]phenyl}ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(2-oxo-1,3-oxazolidin-3-yl)phenyl]ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-2-[[1-(2,2-difluoroethyl)piperidin-4-yl]oxy]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(1-methyl-1H-pyrazol-5-yl)phenyl]ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-[(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl]-4-hydroxy-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide, and

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-[(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl]-4-hydroxy-L-prolinamide.

[Claim 8]

Use of a compound of the following formula (I) or a salt thereof in the manufacture of a pharmaceutical composition for treating colorectal cancer and/or lung cancer: wherein the compound of the formula (I) is selected from the group consisting of

(4R)-1-[(2S)-2-(4-{4-[({6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-

indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-(hydroxymethyl)-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(2-oxo-1,3-oxazolidin-3-yl)phenyl]ethyl]-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-2-{[1-(2,2-difluoroethyl)piperidin-4-yl]oxy}-7-(6-fluoro-5-methyl-1H-indazol-4-yl)quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(1-methyl-1H-pyrazol-5-yl)phenyl]ethyl]-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-[(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl]-4-hydroxy-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide, and
 (4R)-1-[(2S)-2-(4-{4-[({6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-[(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl]-4-hydroxy-L-prolinamide.

[Claim 9]

The use of Claim 8, the compound of the formula (I) is selected from the group consisting of

(4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide,
 (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-{4-[4-

(hydroxymethyl)-1,3-thiazol-5-yl]phenyl]ethyl}-L-prolinamide,
 (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(2-oxo-1,3-oxazolidin-3-yl)phenyl]ethyl}-L-prolinamide,
 (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-2-[[1-(2,2-difluoroethyl)piperidin-4-yl]oxy}-7-(6-fluoro-5-methyl-1H-indazol-4-yl)quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide,
 (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(1-methyl-1H-pyrazol-5-yl)phenyl]ethyl}-L-prolinamide,
 (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-[(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-

2-hydroxyethyl}-4-hydroxy-L-prolinamide,
 (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide, and
 (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-[(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl]-4-hydroxy-L-prolinamide.

[Claim 10]

The use of Claim 8, the compound of the formula (I) is selected from the group consisting of

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide,
 (4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-

yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-{4-[4-(hydroxymethyl)-1,3-thiazol-5-yl]phenyl}ethyl]-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(2-oxo-1,3-oxazolidin-3-yl)phenyl]ethyl}-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-2-{[1-(2,2-difluoroethyl)piperidin-4-yl]oxy}-7-(6-fluoro-5-methyl-1H-indazol-4-yl)quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(1-methyl-1H-pyrazol-5-yl)phenyl]ethyl}-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-

yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-{(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl}-4-hydroxy-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide, and (4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-{(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl}-4-hydroxy-L-prolinamide.

[Claim 11]

A compound of the following formula (I) or a salt thereof for use in therapy:

wherein the compound of the formula (I) is selected from the group consisting of

(4R)-1-[(2S)-2-(4-{4-[({(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(4-methyl-

1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide,
 (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-{4-[4-(hydroxymethyl)-1,3-thiazol-5-yl]phenyl}ethyl}-L-prolinamide,
 (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(2-oxo-1,3-oxazolidin-3-yl)phenyl]ethyl}-L-prolinamide,
 (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-2-[[1-(2,2-difluoroethyl)piperidin-4-yl]oxy]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide,
 (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(1-methyl-1H-pyrazol-5-yl)phenyl]ethyl}-L-prolinamide,
 (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-

yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-{(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl}-4-hydroxy-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide, and (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-{(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl}-4-hydroxy-L-prolinamide.

[Claim 12]

The compound for use of Claim 11, the compound of the formula (I) is selected from the group consisting of (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[(6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-

indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-{4-[4-(hydroxymethyl)-1,3-thiazol-5-yl]phenyl}ethyl]-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(2-oxo-1,3-oxazolidin-3-yl)phenyl]ethyl}-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-2-{[1-(2,2-difluoroethyl)piperidin-4-yl]oxy}-7-(6-fluoro-5-methyl-1H-indazol-4-yl)quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(1-methyl-1H-pyrazol-5-yl)phenyl]ethyl}-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-

indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-{(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl}-4-hydroxy-L-prolinamide, (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide, and (4R)-1-[(2S)-2-(4-{4-[({(7M)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-{(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl}-4-hydroxy-L-prolinamide.

[Claim 13]

The compound for use of Claim 11, the compound of the formula (I) is selected from the group consisting of (4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-{(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl}-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-{4-[4-(hydroxymethyl)-1,3-thiazol-5-yl]phenyl}ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(2-oxo-1,3-oxazolidin-3-yl)phenyl]ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-2-{[1-(2,2-difluoroethyl)piperidin-4-yl]oxy}-7-(6-fluoro-5-methyl-1H-indazol-4-yl)quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(1-methyl-1H-pyrazol-5-yl)phenyl]ethyl]-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(oxan-4-yl)oxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-[(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl]-4-hydroxy-L-prolinamide,

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-4-hydroxy-N-[(1R)-2-hydroxy-1-[4-(4-methyl-1,3-thiazol-5-yl)phenyl]ethyl]-L-prolinamide, and

(4R)-1-[(2S)-2-(4-{4-[({(7P)-6-cyclopropyl-4-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl]-7-(6-fluoro-5-methyl-1H-indazol-4-yl)-2-[(2S)-2-methoxypropoxy]quinazolin-8-yl}oxy)methyl]phenyl}-1H-1,2,3-triazol-1-yl)-3-methylbutanoyl]-N-[(1R)-1-[4-(1-ethyl-1H-pyrazol-5-yl)phenyl]-2-hydroxyethyl]-4-hydroxy-L-prolinamide.