**WHAT IS CLAIMED IS:**

1. A method of treating hidradenitis suppurativa in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a compound which inhibits JAK1 and/or JAK2, or a pharmaceutically acceptable salt thereof, wherein the compound is:

ruxolitinib;

ruxolitinib, wherein one or more hydrogen atoms are replaced by deuterium atoms;

{1-{1-[3-Fluoro-2-(trifluoromethyl)isonicotinoyl]piperidin-4-yl}-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile;

4-{3-(Cyanomethyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-1-yl}-N-[4-fluoro-2-(trifluoromethyl)phenyl]piperidine-1-carboxamide;

[3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]-1-(1-{[2-(trifluoromethyl)pyrimidin-4-yl]carbonyl}piperidin-4-yl)azetidin-3-yl]acetonitrile;

4-[3-(cyanomethyl)-3-(3',5'-dimethyl-1H,1'H-4,4'-bipyrazol-1-yl)azetidin-1-yl]-2,5-difluoro-N-[(1S)-2,2,2-trifluoro-1-methylethyl]benzamide;

((2R,5S)-5-{2-[(1R)-1-hydroxyethyl]-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl}tetrahydro-2H-pyran-2-yl)acetonitrile;

3-[1-(6-chloropyridin-2-yl)pyrrolidin-3-yl]-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]propanenitrile;

3-(1-[1,3]oxazolo[5,4-b]pyridin-2-ylpyrrolidin-3-yl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]propanenitrile;

4-[(4-{3-cyano-2-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]propyl}piperazin-1-yl)carbonyl]-3-fluorobenzonitrile;

4-[(4-{3-cyano-2-[3-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrrol-1-yl]propyl}piperazin-1-yl)carbonyl]-3-fluorobenzonitrile;

[trans-1-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]-3-(4-{[2-(trifluoromethyl)pyrimidin-4-yl]carbonyl}piperazin-1-yl)cyclobutyl]acetonitrile;

{trans-3-(4-{[4-[(3-hydroxyazetidin-1-yl)methyl]-6-(trifluoromethyl)pyridin-2-yl]oxy}piperidin-1-yl)-1-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]cyclobutyl}acetonitrile;

{trans-3-(4-{[4-{[(2S)-2-(hydroxymethyl)pyrrolidin-1-yl]methyl}-6-(trifluoromethyl)pyridin-2-yl]oxy}piperidin-1-yl)-1-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]cyclobutyl}acetonitrile;

{trans-3-(4-{[4-{[(2R)-2-(hydroxymethyl)pyrrolidin-1-yl]methyl}-6-(trifluoromethyl)pyridin-2-yl]oxy}piperidin-1-yl)-1-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]cyclobutyl}acetonitrile;

4-(4-{3-[(dimethylamino)methyl]-5-fluorophenoxy}piperidin-1-yl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]butanenitrile;

5-{3-(cyanomethyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-1-yl}-N-isopropylpyrazine-2-carboxamide;

4-{3-(cyanomethyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-1-yl}-2,5-difluoro-N-[(1S)-2,2,2-trifluoro-1-methylethyl]benzamide;

5-{3-(cyanomethyl)-3-[4-(1H-pyrrolo[2,3-b]pyridin-4-yl)-1H-pyrazol-1-yl]azetidin-1-yl}-N-isopropylpyrazine-2-carboxamide;

{1-(cis-4-{[6-(2-hydroxyethyl)-2-(trifluoromethyl)pyrimidin-4-yl]oxy}cyclohexyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile;

{1-(cis-4-{[4-[(ethylamino)methyl]-6-(trifluoromethyl)pyridin-2-yl]oxy}cyclohexyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile;

{1-(cis-4-{[4-(1-hydroxy-1-methylethyl)-6-(trifluoromethyl)pyridin-2-yl]oxy}cyclohexyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile;

{1-(cis-4-{[4-{[(3R)-3-hydroxypyrrolidin-1-yl]methyl}-6-(trifluoromethyl)pyridin-2-yl]oxy}cyclohexyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile;

{1-(cis-4-{[4-{[(3S)-3-hydroxypyrrolidin-1-yl]methyl}-6-(trifluoromethyl)pyridin-2-yl]oxy}cyclohexyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile;

{trans-3-(4-{[4-({[(1S)-2-hydroxy-1-methylethyl]amino}methyl)-6-(trifluoromethyl)pyridin-2-yl]oxy}piperidin-1-yl)-1-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]cyclobutyl}acetonitrile;

{trans-3-(4-{[4-({[(2R)-2-hydroxypropyl]amino}methyl)-6-(trifluoromethyl)pyridin-2-yl]oxy}piperidin-1-yl)-1-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]cyclobutyl}acetonitrile{trans-3-(4-{[4-({[(2S)-2-hydrox;

ypropyl]amino}methyl)-6-(trifluoromethyl)pyridin-2-yl]oxy}piperidin-1-yl)-1-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]cyclobutyl}acetonitrile;

{trans-3-(4-{[4-(2-hydroxyethyl)-6-(trifluoromethyl)pyridin-2-yl]oxy}piperidin-1-yl)-1-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]cyclobutyl}acetonitrile;

or a pharmaceutically acceptable salt of any of the aforementioned.

2. The method of claim 1, wherein the compound or salt is selective for JAK1 and JAK2 over JAK3 and TYK2.

3. The method of claim 2, wherein the compound is ruxolitinib, or a pharmaceutically acceptable salt thereof.

4. The method of claim 3, wherein the compound is ruxolitinib, or a pharmaceutically acceptable salt thereof, wherein one or more hydrogen atoms are replaced by deuterium atoms.

5. The method of claim 3, wherein the salt is ruxolitinib phosphate.

6. The method of claim 1, wherein the compound or salt is selective for JAK1 over JAK2, JAK3, and TYK2.

7. The method of claim 6, wherein the compound is {1-{1-[3-fluoro-2-(trifluoromethyl)isonicotinoyl]piperidin-4-yl}-3[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile, or a pharmaceutically acceptable salt thereof.

8. The method of claim 7, wherein the salt is {1-{1-[3-fluoro-2-(trifluoromethyl)isonicotinoyl]piperidin-4-yl}-3[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile adipic acid salt.

9. The method of claim 6, wherein the compound is 4-[3-(cyanomethyl)-3-(3',5'-dimethyl-1H,1'H-4,4'-bipyrazol-1-yl)azetidin-1-yl]-2,5-difluoro-N-[(1S)-2,2,2-trifluoro-1-methylethyl]benzamide, or a pharmaceutically acceptable salt thereof.

10. The method of claim 9, wherein the salt is 4-[3-(cyanomethyl)-3-(3',5'-dimethyl-1H,1'H-4,4'-bipyrazol-1-yl)azetidin-1-yl]-2,5-difluoro-N-[(1S)-2,2,2-trifluoro-1-methylethyl]benzamide phosphoric acid salt.

11. The method of claim 6, wherein the compound is ((2R,5S)-5-{2-[(1R)-1-hydroxyethyl]-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl}tetrahydro-2H-pyran-2-yl)acetonitrile, or a pharmaceutically acceptable salt thereof.

12. The method of claim 6, wherein the compound is ((2R,5S)-5-{2-[(1R)-1-hydroxyethyl]-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl}tetrahydro-2H-pyran-2-yl)acetonitrile monohydrate.

13. The method of any one of claims 7-12, wherein the compound or salt is administered at a dosage of 15, 30, 60 or 90 mg on a free base basis.

14. The method of any one of claims 1-13, further comprising administering an additional therapeutic agent.

15. The method of claim 14, wherein the additional therapeutic agent is an antibiotic, a retinoid, a corticosteroid, an anti-TNF-alpha agent, or an immunosuppressant.

16. The method of claim 15, wherein the antibiotic is clindamycin, doxycycline, minocycline, trimethoprim-sulfamethoxazole, erythromycin, metronidazole, rifampin, moxifloxacin, dapsone, or a combination thereof.

17. The method of claim 15, wherein the retinoid is etretinate, acitretin, or isotretinoin.

18. The method of claim 15, wherein the corticosteroid is triamcinolone, dexamethasone, fluocinolone, cortisone, prednisone, prednisolone or flumetholone.

19. The method of claim 15, wherein the anti–TNF-alpha agent is infliximab, etanercept, or adalimumab.

20. The method of claim 15, wherein the immunosuppressant is methotrexate, cyclosporin A, mycophenolate mofetil, or mycophenolate sodium.

21. The method of claim 14, wherein the additional therapeutic agent is finasteride, metformin, adapalene, or azelaic acid.

22. The method of any one of claims 1-21, wherein the administrating of the compound or salt is topical.

23. The method of any one of claims 1-21, wherein the administering of the compound or salt is oral.

24. The method of any one of claims 1-23, wherein the method results in a 10%, 20%, 30%, 40%, or 50% improvement in HiSCR (Hidradenitis Suppurativa Clinical Response).