

INTERLEUKIN-2/INTERLEUKIN-2 RECEPTOR ALPHA FUSION PROTEINS
AND METHODS OF USE

FIELD OF THE INVENTION

The presently disclosed subject matter generally relates to methods and compositions for modulating the immune response employing an Interleukin-2/Interleukin-2 Receptor alpha fusion protein.

REFERENCE TO A SEQUENCE LISTING SUBMITTED AS
A TEXT FILE VIA EFS-WEB

The official copy of the sequence listing is submitted concurrently with the specification as a text file via EFS-Web, in compliance with the American Standard Code for Information Interchange (ASCII), with a file name of 464173seqlist.txt, a creation date of July 30, 2015 and a size of 139 KB. The sequence listing filed via EFS-Web is part of the specification and is hereby incorporated in its entirety by reference herein.

STATEMENT OF FEDERAL FUNDING

This invention was made with government support under grant number R01 DK093866, awarded by the National Institute of Health (NIH), National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). The government has certain rights in the invention.

BACKGROUND OF THE INVENTION

Interleukin-2 (IL-2) is a biologic that has been used in attempts to boost immune responses in cancer and HIV/AIDS patients. More recently lower doses of IL-2 have been used to selectively boost tolerance to suppress unwanted immune responses associated with autoimmune-like attack of self tissues. Importantly, these low doses of IL-2 have not shown any signs of enhancing or re-activation of autoreactive T cells. Nevertheless, IL-2 has important drawbacks as a therapeutic, including a very short-half life *in vivo*, which limits its efficacy, and toxicity at high doses. For these reasons

new IL-2 biologics are needed having improved pharmacokinetics and durability of responses for use.

SUMMARY OF THE INVENTION

5 Various methods and compositions are provided which can be employed to modulate the immune system. Compositions include a fusion protein comprising: (a) a first polypeptide comprising Interleukin-2 (IL-2) or a functional variant or fragment thereof; and (b) a second polypeptide, fused in frame to the first polypeptide, wherein the second polypeptide comprises an extracellular domain of Interleukin-2 Receptor
10 alpha (IL-2R α) or a functional variant or fragment thereof, and wherein the fusion protein has IL-2 activity.

Various methods are provided for decreasing the immune response in a subject comprising administering to a subject in need of a decrease in the immune response a therapeutically effective amount of the IL-2/IL-2R α fusion protein disclosed herein.

15 Further provided are methods for increasing the immune response in a subject comprising administering to a subject in need of an increase in the immune response a therapeutically effective amount of the IL-2/IL-2R α fusion protein disclosed herein. Further provided are methods for increasing T regulatory cell activity.

Additional methods including enhancing the immunogenicity of a vaccine or
20 overcoming a suppressed immune response to a vaccine in a subject, comprising: (a) administering to the subject a therapeutically effective amount of the IL-2/IL-2R α fusion protein disclosed herein; and, (b) administering to the subject a vaccine, wherein the fusion protein enhances the immunogenicity of the vaccine or overcomes the suppressed immune response to the vaccine.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 provides a schematic of an IL-2/IL-2R α fusion protein, where L=leader peptide, LK=linker region, G=glycine, H=histidine, and T= termination codon.

FIG. 2A and FIG. 2B provide the deduced protein sequences of non-limiting
30 examples of IL-2/IL-2R α fusion proteins. FIG. 2A provides the deduced protein sequences of non-limiting examples of mouse IL-2/IL-2R α fusion proteins. The sequences of mouse IL-2 and IL-2R α are shown above and below, respectively, of the fusion proteins. The sequence denoted as IL-2 is set forth in SEQ ID NO: 3; the sequence denoted as IL-2-(G4S)4-IL-2R α is set forth in SEQ ID NO: 54; the sequence

denoted as IL-2-(G4S)5-IL-2R α is set forth in SEQ ID NO:55; the sequence denoted as IL-2-(G3S)4-IL-2R α is set forth in SEQ ID NO:56; the sequence denoted as IL-2-(G3S)3-IL-2R α is set forth in SEQ ID NO: 57; and the extracellular domain of IL-2R α is set forth in SEQ ID NO: 10. FIG. 2B provides the deduced protein sequences of non-limiting examples of human IL-2/IL-2R α fusion proteins. The sequences of human IL-2 and IL-2R α are shown above and below, respectively, of the fusion proteins. The sequence denoted as IL-2 is set forth in SEQ ID NO: 1; the sequence denoted as IL-2-(G3S)2-IL-2R α is set forth in SEQ ID NO: 58; the sequence denoted as IL-2-(G3S)3-IL-2R α is set forth in SEQ ID NO:59; the sequence denoted as IL-2-(G3S)4-IL-2R α is set forth in SEQ ID NO:60; the sequence denoted as IL-2-(G4S)4-IL-2R α is set forth in SEQ ID NO: 61; and the extracellular domain of IL-2R α is set forth in SEQ ID NO: 7.

FIG. 3 shows the bioactivity of IL-2/IL-2R α fusion proteins. COS-7 cells were transfected with the IL-2/IL-2R α fusion cDNAs with the indicated linkers. Supernatants from these cells were cultured with anti-CD3 activated T cell blasts to assess IL-2 activity. (A) Proliferative responses by the T blasts after dilutions of the indicated fusion proteins. (B) Effect of anti-IL-2 on proliferation stimulated by a 1:2 dilution of the culture supernatant containing the indicated fusion proteins.

FIG. 4 shows the activity of purified IL-2/IL-2R α fusion proteins. Supernatants of transfected CHO cells were used to purify IL-2/(G₃S)₃/IL-2R α and IL-2/(Gly₄Ser)₄/IL-2R α by Nickel-based affinity chromatography to the 6x-His tag. (A) IL-2 bioactivity measure by proliferation of anti-CD3 T cell blast to the indicated purified fusion protein. (B) The effect of each purified fusion protein to inhibit the binding of PC61 and 7D4 anti-IL-2R α monoclonal antibodies, directed to non-ligand binding site, to anti-CD3 activated T cell blasts.

FIG. 5 shows that a monoclonal anti-IL-2R α antibody that is directed to the IL-2 binding site of IL-2R α cannot bind to the IL-2/IL-2R α fusion protein. Purified fusion proteins with variable linkers, as indicated, were first incubated with the 3C7 anti-IL-2R α monoclonal antibody, directed to the ligand binding site of IL-2R α or the 7D4 monoclonal antibody, directed to a non-ligand binding site of IL-2R α . The capacity of 3C7 or 7D4 to then bind to cell surface IL-2R α was assessed using IL-2R α -transfected EL4 cells.

FIG. 6 shows the biochemical properties of purified IL-2/IL-2R α . (A) Purified IL-2/IL-2R α was subjected to SDS-PAGE under reducing and non-reducing conditions;

IL-2/IL-2R α was visualized by Western blot analysis by probing with an antibody directed to 6x-His tag of the fusion protein. **(B)** The indicated amount of purified IL-2/(Gly₃Ser)₃/IL-2R α was subjected to SDS-PAGE under reducing conditions followed by Coomassie Blue staining.

FIG. 7 shows the effect of IL-2/IL-2R α fusion protein on IL-2-dependent signal transduction *in vivo*. C57BL/6 mice received a single injection i.p. of IL-2/(G₃S)₃/IL-2R α (4000 units of IL-2 activity) and pSTAT5 levels in the indicated spleen cell populations were immediately assessed. pSTAT5 levels were determined 0.5 hr after injection of the IL-2/IL-2R α fusion protein. For CD4⁺ T cells, the cells were gated to exclude Foxp3⁺ Treg cells

FIG. 8 shows the effect of IL-2/IL-2R α fusion protein on Tregs cells *in vivo*. NOD mice were injected i.p. 3 times (day 1, 3, 5) with the indicated amount of IL-2 activity associated with IL-2/(G₃S)₃/IL-2R α . The effect on Tregs was assessed for the spleen, pancreatic lymph nodes (PLN) and pancreas 24 hr after the last injection. Evaluated were the proportion of Tregs in CD4⁺ T cells; the mean fluorescent intensity (MFI) for CD25 expression by Tregs after normalization to CD25 expression by Tregs from control treated mice; the proliferative status of Tregs as assessed by expression of the proliferative marker Ki67; and the % of Tregs that expressed Klrp1, which marks an IL-2-dependent terminally differentiated subpopulation.

FIG. 9 shows the comparison of IL-2/IL-2R α fusion protein and recombinant IL-2 to induce changes in Treg cells *in vivo*. C57BL/6 mice were injected i.p. 3 times (day 1, 3, 5) with IL-2/(G₃S)₃/IL-2R α (2000 Units), recombinant human IL-2 (25,000 Units) or preformed complexes of anti-IL-2 (Jes-6.1; 5 μ g) and mouse IL-2 (10,000 Units) (IL2/IC). The effect on Tregs was assessed for the spleen 24, 72 hr and 1 week after the last injection. Treg were evaluated as described in FIG. 8.

FIG. 10 shows Limited application of low-dose IL-2 delays diabetes in NOD mice. NOD mice (8 mice/group) received IL-2/IL-2R α , soluble IL-2R α , or PBS according to the schedule in (A). Urine and blood glucose levels were monitored until mice reached 40 weeks of age. Mice were considered diabetic after 2 consecutive readings of glucose levels >250 mg/dl.

FIG. 11 demonstrates high-dose IL-2/IL-2R α enhances the development of CD8⁺ T cell memory. C57BL/6 mice received congenic class I-restricted ovalbumin

(OVA)-specific OT-I T cell receptor transgenic T cells. These mice were immunized and treated with a single application of IL-2/(G₃S)₃/IL-2R α fusion protein, IL2/IC containing 15,000 units of IL-2, or recombinant IL-2 (25,000 Units). At the indicated times, the relative proportion of OT-I T cells within the total CD8⁺ T cell compartment in peripheral blood was assessed.

FIG. 12 shows the type of persistent OT-I memory cells supported by high-dose IL-2/IL-2R α fusion protein: (A) Gating strategy to identify effector-memory (EM) and central memory (CM) cells. (B) Distribution of OT-I memory cells 28 and 202 day post immunization for mice that also received IL-2/IL-2R α (12,000 units).

FIG. 13 shows characterization of human IL-2/IL-2R α fusion proteins containing glycine/serine linkers of variable length, as shown. (A) IL-2-bioactivity of purified human IL-2//IL-2R α using the CTLL bioassay. (B) Western blot analysis of human IL-2/IL-2R α fusion proteins after SDS-PAGE under reducing conditions.

FIG. 14 shows human IL-2/IL-2R α fusion protein bind monoclonal anti-IL-2R α antibodies. Purified fusion proteins with the indicated linkers were first incubated with the BC96 anti-IL-2R α monoclonal antibody, directed to the ligand binding region of human IL-2R α or the M-A257 monoclonal antibody, directed to a non-ligand binding region of human IL-2R α . The capacity of BC96 or M-A257 to then bind to cell surface IL-2R α was assessed using IL-2R α -transfected CHO cells.

FIG. 15 shows IL-2 interacts with the IL-2 binding site of IL-2R α in the context of human IL-2/IL-2R α fusion proteins. IL-2-bioactivity of the indicated fusion proteins with variable glycine/serine linkers was assessed using CTLL cells. Mut refers to fusion proteins where IL-2R α contained Arg³⁵→Thr, Arg³⁶→Ser mutations. Western blot analysis confirmed similar amounts of all fusion proteins (not shown).

DETAILED DESCRIPTION OF THE INVENTION

The present inventions now will be described more fully hereinafter with reference to the accompanying drawings, in which some, but not all embodiments of the invention are shown. Indeed, these inventions may be embodied in many different forms and should not be construed as limited to the embodiments set forth herein; rather, these embodiments are provided so that this disclosure will satisfy applicable legal requirements. Like numbers refer to like elements throughout.

Many modifications and other embodiments of the inventions set forth herein will come to mind to one skilled in the art to which these inventions pertain having the benefit of the teachings presented in the foregoing descriptions and the associated drawings. Therefore, it is to be understood that the inventions are not to be limited to the specific embodiments disclosed and that modifications and other embodiments are intended to be included within the scope of the appended claims. Although specific terms are employed herein, they are used in a generic and descriptive sense only and not for purposes of limitation.

I. Overview

Current technology relies on the use of recombinant Interleukin-2 (IL-2), which has poor pharmacological properties, especially a short half-life that limits its usefulness. Provided herein are Interleukin-2/Interleukin-2 receptor alpha (IL-2/IL-2R α) fusion proteins have intrinsic properties that separate them from recombinant IL-2 and other IL-2 fusion proteins. First, the size of the IL-2/IL-2R α fusion protein will increase its half-life *in vivo*. Second, the weak interaction between IL-2 and IL-2R α (one subunit of the IL-2R) in the context of the IL-2/IL-2R α fusion protein provides another mechanism to prolong the availability of the IL-2. While not being limited to the specific mechanism of action, the prolonged availability of the IL-2 activity might occur through a competitive interaction between the IL-2 moiety with IL-2R α of the IL-2/IL-2R α fusion and with cells that express the IL-2R.

II. Interleukin-2/Interleukin-2 Receptor Alpha Fusion Proteins And Polynucleotides Encoding the Same

A fusion protein is provided which comprises a first polypeptide comprising interleukin-2 (IL-2) or a functional variant or fragment thereof fused in frame to a second polypeptide comprising or consisting of the extracellular domain of the Interleukin-2 Receptor Alpha (IL-2R α) polypeptide or a functional variant or fragment thereof.

As used herein, “fusion protein” refers to the in frame genetic linkage of at least two heterologous polypeptides. Upon transcription/translation, a single protein is made. In this way, multiple proteins, or fragments thereof can

be incorporated into a single polypeptide. "Operably linked" is intended to mean a functional linkage between two or more elements. For example, an operable linkage between two polypeptides fuses both polypeptides together in frame to produce a single polypeptide fusion protein. In a particular aspect,
5 the fusion protein further comprises a third polypeptide which, as discussed in further detail below, can comprise a linker sequence.

The IL-2/IL-2R α fusion protein or the active variant or fragment thereof can have one or more the following properties/activities: (1) increasing activity of regulatory T cells (Tregs) and/or increasing immune
10 tolerance in low dose IL-2 based therapies; (2) increasing immune response and memory in higher dose therapies; (3) increasing IL-2 availability when compared to recombinant IL-2; and/or (4) increasing persistent IL-2 stimulation of IL-2R bearing lymphocytes *in vivo*. Such activity and methods of assaying are disclosed in further detail elsewhere herein. See, for example,
15 Example 1 provided herein.

In one non-limiting embodiment, an increased activity of Tregs that results from the IL-2/IL-2R α fusion protein or the active variant or fragment thereof can be assayed in a variety of ways including, for example, (1) an increased representation and number of Tregs in the CD4⁺ T cell compartment;
20 (2) upregulation of IL-2-dependent CD25; (3) increased proliferation as assessed by expression of the proliferative marker Ki67; and (4) an increased fraction of IL-2-dependent terminally differentiated KIrg1⁺ Treg subset. Such effects on Tregs can be seen in, for example, in the spleen and the inflamed pancreas.

In one non-limiting embodiment, the IL-2/IL-2R α fusion protein or the active variant or fragment thereof increases tolerogenic and immune suppressive Tregs and immunity through increasing T effector/memory responses and, in further embodiments, it exhibits improved pharmacokinetics by delivering such responses at (1) lower effective levels of IL-2 activity
25 compared to native or recombinant IL-2; (2) displays more persistent biological responses than native or recombinant IL-2; and/or (3) retains the hierarchy with Tregs responsive at lower level doses than T effector/memory cells.
30

In specific embodiments, the fusion protein has an improved activity over the native or recombinant IL-2. For example, the effect of the IL-2/IL-2R α fusion protein can increase tolerogenic Tregs at about 2 fold, 5 fold, 10 fold, 20 fold, 30 fold, 40 fold, 50 fold, 60 fold, 70 fold, 80 fold, 90 fold, 100 fold 150 fold, 200 fold or lower level IL-2 activity in comparison to native or recombinant IL-2. In other embodiments, the IL-2/IL-2R α fusion protein is more effective than native or recombinant IL-2 in inducing persistent augmentation of Tregs and related properties.

Various IL-2 and IL-2R α fragments and variants from a variety of organism can be used to generate the IL-2/IL-2R α extracellular domain fusion proteins provided herein. Such components are discussed in further detailed elsewhere herein. Examples of non-limiting unprocessed IL-2/IL-2R α extracellular domain fusion proteins are set forth in SEQ ID NO: 17, 19, 21, 23, 25, 27, 36, 38, 44, 46, 54, 55, 56, 57, 58, 59, 60, and 61, while non-limiting examples of mature forms of the IL-2/IL-R α extracellular domain fusion proteins are set forth in SEQ ID NOS: 16, 18, 20, 22, 24, 26, 37, 39, 43, 45, 62 and 64. Non-limiting examples of polynucleotides encoding such fusion proteins are set forth in SEQ ID NO: 29, 30, 31, 32, 33, 34, 42, 47, 48, 49, 63, and 65.

The term "secretory signal sequence" denotes a polynucleotide sequence that encodes a polypeptide (a "secretory peptide") that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of the cell in which it is synthesized. The larger polypeptide is commonly cleaved to remove the secretory peptide during the transit through the secretory pathway. As used herein, a "mature" form of a fusion protein or polypeptide comprises the processed form of the polypeptide that has had the secretory peptide removed. As used herein, the "unprocessed" form of the fusion protein retains the secretory peptide sequence.

Biologically active fragments and variants of the mature and unprocessed form of the IL-2/IL-R α extracellular domain fusion proteins, and the polynucleotide encoding the same, are also provided. Such a functional polypeptide fragment can comprise at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500 or more continuous amino acids of any one of SEQ ID NO: 16, 17, 18, 19, 20,

21, 22, 23, 24, 25, 26, 27, 36, 37, 38, 39, 43, 44, 45, 46, 54, 55, 56, 57, 58, 59, 60, 61, 62, or 64. Alternatively, a functional polypeptide variant can comprise at least 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to the sequence set forth in SEQ ID NO: 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 36, 37, 38, 39, 43, 44, 45, 46, 54, 55, 56, 57, 58, 59, 60, 61, 62, or 64.

Active variants and fragments of polynucleotides encoding the IL-2/IL-R α extracellular domain fusion proteins are further provided. Such polynucleotide can comprise at least 100, 200, 300, 400, 500, 600, 700, 800, 1000, 1100, 1200, 1300, 1500, 1800, 2000 continuous nucleotides of SEQ ID NO: 29, 30, 31, 32, 33, 34, 42, 47, 48, 49, 63 or 65 or the polynucleotide encoding the polypeptides set forth in SEQ ID NO: 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 36, 37, 38, 39, 43, 44, 45, 46, 54, 55, 56, 57, 58, 59, 60, 61, 62, or 64 and continue to encode a functional IL-2/IL-R α extracellular domain fusion protein. Alternatively, a functional polynucleotide can comprise at least 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to the sequence set forth in SEQ ID NO: 29, 30, 31, 32, 33, 34, 42, 47, 48, 49, 63 or 65 or the polynucleotide encoding the polypeptides set forth in SEQ ID NO: 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 36, 37, 38, 39, 43, 44, 45, 46, 54, 55, 56, 57, 58, 59, 60, 61, 62, or 64 and continue to encode a functional IL-2/IL-R α extracellular domain fusion proteins.

It is further recognized that the components of the IL-2/IL-2R α fusion protein can be found any order. In one embodiment, the IL-2 polypeptide is at the N-terminus and the extracellular domain of IL-2R α is at the C-terminus of the fusion protein.

i. Interleukin-2

As used herein, "Interleukin-2" or "IL-2" refers to any native or recombinant IL-2 from any vertebrate source, including mammals such as primates (e.g. humans) and rodents (e.g. mice and rats), and domesticated or agricultural mammals unless otherwise indicated. The term encompasses unprocessed IL-2, as well as, any form of IL-2 that results from processing in the cell (i.e, the mature form of IL-2). The term also encompasses naturally occurring variants and fragments of IL-2, e.g. splice variants or allelic variants, and non-naturally occurring variants. The amino acid

sequence of an exemplary mature form of human IL-2 (having the 20 amino acid signal sequence) is shown in SEQ ID NO: 2. Unprocessed human IL-2 additionally comprises an N-terminal 20 amino acid signal peptide (SEQ ID NO: 1), which is absent in the mature IL-2 molecule. The amino acid sequence of an exemplary mature form of mouse IL-2 (having the 20 amino acid signal sequence) is shown in SEQ ID NO: 4. Unprocessed mouse IL-2 additionally comprises an N-terminal 20 amino acid signal peptide (SEQ ID NO: 3), which is absent in the mature IL-2 molecule. See also FIG. 2A and FIG. 2B. By a "native IL-2", also termed "wild-type IL-2", is meant a naturally occurring or recombinant IL-2.

Additional nucleic acid and amino acid sequences for IL-2 are known. See, for example, GenBank Accession Nos: Q7JFM2 (*Aotus lemurinus* (Gray-bellied night monkey)); Q7JFM5 (*Aotus nancymae* (Ma's night monkey)); P05016 (*Bos taurus* (Bovine)); Q29416 (*Canis familiaris* (Dog) (*Canis lupus familiaris*)); P36835 (*Capra hircus* (Goat)); and, P37997 (*Equus caballus* (Horse)).

Biologically active fragments and variants of IL-2 are also provided. Such IL-2 active variants or fragments will retain IL-2 activity. The phrase "biological activity of IL-2" refers to one or more of the biological activities of IL-2, including but not limited to, the ability to stimulate IL-2 receptor bearing lymphocytes. Such activity can be measured both *in vitro* and *in vivo*. IL-2 is a global regulator of immune activity and the effects seen here are the sum of such activities. For example, it is regulates survival activity (Bcl-2), induces T effector activity (IFN-gamma, Granzyme B, and Perforin), and promotes T regulatory activity (FoxP3). See, for example, Malek *et al.* (2010) *Immunity* 33(2):153-65, herein incorporated by reference in its entirety.

Biologically active variants of IL-2 are known. See, for example, US Application Publications 20060269515 and 20060160187 and WO 99/60128, each of which is herein incorporated by reference.

Biologically active fragments and variants of IL-2 can be employed in the fusion proteins disclosed herein. Such a functional fragment can comprise at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 75, 100, 125, 150 or more continuous amino acids of SEQ ID NO: 1, 2, 3, or 4. Alternatively, a functional variant can comprise at least 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to the sequence set forth in SEQ ID NO: 1, 2, 3, or 4.

Active variants and fragments of polynucleotides encoding the IL-2 proteins are further provided. Such polynucleotide can comprise at least 100, 200, 300, 400, 500, 600, 700 continuous nucleotides of polypeptide encoding SEQ ID NO: 1, 2, 3, or 4, and continue to encode a protein having IL-2 activity. Alternatively, a functional polynucleotide can comprise at least 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to the polypeptide encoding the amino sequence set forth in SEQ ID NO: 1, 2, 3, or 4 and continue to encode a functional IL-2 polypeptide.

10 *ii. Interleukin-2 Receptor Alpha*

The term "CD25" or "IL-2 receptor α " or "IL-2R α " as used herein, refers to any native or recombinant IL-2R α from any vertebrate source, including mammals such as primates (e.g. humans) and rodents (e.g., mice and rats) and domesticated or agricultural mammals unless otherwise indicated. The term also encompasses naturally occurring variants of IL-2R α , e.g. splice variants or allelic variants, or non-naturally occurring variants. Human IL-2 exerts its biological effects via signaling through its receptor system, IL-2R. IL-2 and its receptor (IL-2R) are required for T-cell proliferation and other fundamental functions which are crucial of the immune response. IL-2R consists of 3 noncovalently linked type I transmembrane proteins which are the alpha (p55), beta (p75), and gamma (p65) chains. The human IL-2R alpha chain contains an extracellular domain of 219 amino acids, a transmembrane domain of 19 amino acids, and an intracellular domain of 13 amino acids. The secreted extracellular domain of IL-2R alpha (IL-2R-a) can be employed in the fusion proteins describe herein.

The amino acid sequence of an exemplary mature form of human IL-2R α is shown in SEQ ID NO: 6. Unprocessed human IL-2R α is shown in SEQ ID NO: 5. The extracellular domain of SEQ ID NO: 6 is set forth in SEQ ID NO: 7. The amino acid sequence of an exemplary mature form of mouse IL-2R α is shown in SEQ ID NO: 9. Unprocessed mouse IL-2R α is shown in SEQ ID NO: 8. The extracellular domain of SEQ ID NO: 9 is set forth in SEQ ID NO: 10. By a "native IL-2R α ", also termed "wild-type IL-2R α ", is meant a naturally occurring or recombinant IL-2R α . The sequence of a native human IL-2R α molecule is shown in SEQ ID NO: 5 and 6.

Nucleic acid and amino acid sequences for IL-2R α are known. See, for example, GenBank Accession Nos: NP_001030597.1 (*P.troglodytes*); NP_001028089.1 (*M.mulatta*); NM_001003211.1 (*C.lupus*); NP_776783.1 (*B.taurus*); NP_032393.3 (*M.musculus*); and, NP_037295.1 (*R.norvegicus*), each of which is herein incorporated by reference.

Biologically active fragments and variants of the extracellular domain of IL-2R α are also provided. Such IL-2R α extracellular domain active variants or fragments will retain the IL-2R α extracellular domain activity. The phrase “biological activity of the IL-2R α extracellular domain” refers to one or more of the biological activities of extracellular domain of IL-2R α , including but not limited to, the ability to enhance intracellular signaling in IL-2 receptor responsive cells. Non-limiting examples of biologically active fragments and variants of the IL-2R α are disclosed, for example, in Robb et al., *Proc. Natl. Acad. Sci. USA*, 85:5654-5658, 1988, which is herein incorporated by reference.

Biologically active fragments and variants of the extracellular domain of IL-2R α can be employed in the fusion proteins disclosed herein. Such a functional fragment can comprise at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 75, 100, 125, 150, 175, 200, 215 or greater continuous amino acids of the extracellular domain of any one of SEQ ID NO: 6, 9, 7, 10, 5, or 8. Alternatively, a functional variant can comprise at least 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to the sequence set forth in SEQ ID NO: 6, 9, 7, 10, 5, or 8.

In one embodiment, the fusion proteins provided herein can comprise at least one mutation within the extracellular domain of IL-2R α . In a specific embodiment, the Arginine at position 35 of IL-2R α can be mutated to a Threonine and/or the Arginine at position 36 of IL-2R α can be mutated to a Serine. Such a fusion protein can have increased IL-2 activity compared to a fusion protein not comprising these mutations in the extracellular domain of IL-2R α and/or compared to native or recombinant IL-2. The amino acid sequences of exemplary fusion proteins comprising IL-2R α with mutations within the extracellular domain of IL-2R α are set forth in SEQ ID NOS: 62 and 64. In one embodiment, the fusion protein comprises the amino acid sequence of any one of SEQ ID NO: 62 or 64; or a sequence having at least 80%, 85%, 90%, or 95% to any one of SEQ ID NO: 62 or 64.

Active variants and fragments of polynucleotides encoding the extracellular domain of IL-2R α are further provided. Such polynucleotide can comprise at least 100, 200, 300, 400, 500, 600 or greater continuous nucleotides of polypeptide encoding SEQ ID NO: 6, 9, 7, 10, 5, or 8 and
5 continue to encode a protein having the extracellular domain activity of IL-2R α . Alternatively, a functional polynucleotide can comprise at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to the polypeptide encoding the amino sequence set forth in SEQ ID NO: 6, 9, 7, 10, 5, or 8 and continue to encode a protein having the
10 extracellular domain activity of IL-2R α .

iii. Additional Components

The IL-2/IL-2R α fusion proteins can further comprise additional elements. Such elements can aid in the expression of the fusion protein, aid in the secretion of
15 the fusion protein, improve the stability of the fusion protein, allow for more efficient purification of the protein, and/or modulate the activity of the fusion protein.

“Heterologous” in reference to a polypeptide or polynucleotide is a polypeptide or polynucleotide that originates from a different protein or polynucleotide. The additional components of the fusion protein can originate from
20 the same organism as the other polypeptide components of the fusion protein, or the additional components can be from a different organism than the other polypeptide components of the fusion protein.

In one embodiment, the IL-2/IL-2R α fusion protein comprises a linker sequence located between the IL-2 polypeptide and the IL-2R α polypeptide.
25 The linker can be of any length and can comprise at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 50 or 60 or more amino acids. In one embodiment, the linker sequence comprises glycine amino acid residues. In other instances, the linker sequence comprises a combination of glycine and serine amino acid residues.
30 Such glycine/serine linkers can comprises any combination of the amino acid residues, including, but not limited to, the peptide GGGs or GGGGS or repeats of the same, including 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more repeats of these given peptides. For example, linker sequences can comprise GGGSGGGSGGGs (SEQ ID NO: 13) (also noted as (Gly₃Ser)₃);

GGGSGGGSGGGSGGG (SEQ ID NO: 11) (also noted as (Gly₃Ser)₄); or (Gly₃Ser)₅; (Gly₃Ser)₆; (Gly₃Ser)₇, etc. Linker sequences can further comprise (Gly₄Ser)₃ as set forth in SEQ ID NO: 50; GGGSGGGSGGGSGGGSGGG (SEQ ID NO: 40) (also noted as (Gly₄Ser)₄);

- 5 GGGSGGGSGGGSGGGSGGGSGGG (SEQ ID NO: 41) (also noted as (Gly₄Ser)₅); (Gly₄Ser)₂, (Gly₄Ser)₁, (Gly₄Ser)₆; (Gly₄Ser)₇; (Gly₄Ser)₈, etc. In addition, active variants and fragments of any linker can further be employed in the fusion protein disclosed herein.

- It is further recognized that the polynucleotide encoding the IL-2/IL-2R α fusion protein can comprise additional elements that aid in the translation of the fusion protein. Such sequences include, for example, Kozak sequences attached to the 5' end of the polynucleotide encoding the fusion protein. The Kozak consensus sequence is a sequence which occurs on eukaryotic mRNA that plays a role in the initiation of the translation process and has the consensus (gcc)gccRccAUGG (SEQ ID NO: 35); wherein (1) a lower case letter denotes the most common base at a position where the base can nevertheless vary; (2) upper case letters indicate highly-conserved bases, *i.e.* the 'AUGG' sequence is constant or rarely, if ever, changes, with the exception being the IUPAC ambiguity code 'R' which indicates that a purine (adenine or guanine) is normally observed at this position; and (3) the sequence in brackets ((gcc)) is of uncertain significance. In one embodiment, the Kozak sequence comprises the sequence set forth in SEQ ID NO: 53.
- 10
15
20

- In one non-limiting embodiment, the IL-2/IL-2R α fusion protein comprises an IL-2 leader optimized Kozak sequence as set forth in SEQ ID NO: 28 or a functional variant or fragment thereof. A functional variant or fragment of a Kozak sequence will retain the ability to increase translation of the protein when compared to the level of translation from a sequence lacking the leader. Such a functional fragment can comprise at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40 continuous nucleotides of a kozak sequence or the sequence set forth in SEQ ID NO: 28 or 53. Alternatively, a functional variant can comprise at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to the kozak sequence or the sequence set forth in SEQ ID NO: 28 or 53.
- 25
30

In still further embodiments, the IL-2/IL-2R α fusion protein comprises one or more tags at the C-terminus to aid in the purification of the polypeptide.

Such tags are known and include, for example, a Histidine tag. In specific embodiments a 6X His tag is employed. It is further recognized that an additional linker sequence can be employed between the fusion protein and the His tag.

5 Non-limiting embodiment of an IL-2/IL-2R α fusion protein is set forth in FIG. 1, FIG. 2A, and FIG. 2B. Such a fusion protein comprises a leader peptide, IL-2 or a functional variant or fragment thereof, a variable linker, IL-2R α , a glycine linker, 6x his tag, and two termination codons.

10 *iv. Variants and Fragments*

a. Polynucleotides

Fragments and variants of the polynucleotides encoding the IL-2/IL-2R α extracellular domain fusion protein or the various components contained therein (i.e., the IL-2R α extracellular domain, the IL-2R α polypeptides, the linker sequences
15 and/or Kozak sequences) can be employed in the various methods and compositions of the invention. By "fragment" is intended a portion of the polynucleotide and hence the protein encoded thereby or a portion of the polypeptide. Fragments of a polynucleotide may encode protein fragments that retain the biological activity of the native protein and hence have IL-2 activity, IL-2R α extracellular domain activity, IL-
20 2/IL-2R α fusion protein activity, or if encoding a linker sequence, provide for the desired activity of the IL-2/IL-2R α fusion protein.

A biologically active portion of a IL-2R α extracellular domain, IL-2 polypeptide, IL-2/IL-2R α fusion protein, Kozak sequence, or linker sequence can be prepared by isolating a portion of one of the polynucleotides encoding the portion of
25 the IL-2R α extracellular domain or IL-2 polypeptide and expressing the encoded portion of the polypeptide (e.g., by recombinant expression *in vitro*), and assessing the activity of the portion of the IL-2R α extracellular domain or/and IL-2 polypeptide or the activity of the IL-2/ILR α fusion protein.

"Variant" sequences have a high degree of sequence similarity. For
30 polynucleotides, conservative variants include those sequences that, because of the degeneracy of the genetic code, encode the amino acid sequence of one of the IL-2R α extracellular domain polypeptides, IL-2 polypeptides, IL-2/IL-2R α fusion proteins, or linker sequences. Variants such as these can be identified with the use of well-known molecular biology techniques, as, for example, polymerase chain reaction (PCR) and

hybridization techniques. Variant polynucleotides also include synthetically derived nucleotide sequences, such as those generated, for example, by using site-directed mutagenesis but which still encode an IL-2R α extracellular domain, IL-2 polypeptide, IL-2/IL-2R α fusion protein, a Kozak sequence, or the linker sequence.

5

b. Polypeptides

By "variant" protein is intended a protein derived from the native protein by deletion (so-called truncation) or addition of one or more amino acids to the N-terminal and/or C-terminal end of the native protein; deletion or addition of one or
10 more amino acids at one or more sites in the native protein; or substitution of one or more amino acids at one or more sites in the native protein. Variant proteins are biologically active, that is they continue to possess the desired biological activity, that is, IL-2/IL-2R α fusion protein activity, IL-2 activity or IL-2R α extracellular domain activity. Such variants may result from, for example, genetic polymorphism or from
15 human manipulation. Biologically active variants of a IL-2/IL-2R α fusion protein or any one of its components (i.e., an IL-2R α extracellular domain polypeptide, a IL-2 polypeptide, or a linker sequence) will have at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity to the amino acid sequence for the native protein
20 as determined by sequence alignment programs and parameters described elsewhere herein. A biologically active variant of a protein may differ from that protein by as few as 1-15 amino acid residues, as few as 1-10, such as 6-10, as few as 5, as few as 4, 3, 2, or even 1 amino acid residue.

Proteins may be altered in various ways including amino acid substitutions,
25 deletions, truncations, and insertions. Methods for such manipulations are generally known in the art. For example, amino acid sequence variants of the IL-2R α extracellular domain, IL-2 polypeptide, IL-2/IL-2R α fusion protein, or linker sequences can be prepared by mutations in the DNA. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Kunkel
30 (1985) *Proc. Natl. Acad. Sci. USA* 82:488-492; Kunkel *et al.* (1987) *Methods in Enzymol.* 154:367-382; U.S. Patent No. 4,873,192; Walker and Gastra, eds. (1983) *Techniques in Molecular Biology* (MacMillan Publishing Company, New York) and the references cited therein. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the protein of interest may be found in the model of

Dayhoff *et al.* (1978) *Atlas of Protein Sequence and Structure* (Natl. Biomed. Res. Found., Washington, D.C.), herein incorporated by reference. Conservative substitutions, such as exchanging one amino acid with another having similar properties, may be preferable.

5 Thus, the polynucleotides disclosed herein can include the naturally occurring sequences, the "native" sequences, as well as mutant forms. Likewise, the proteins used in the methods of the invention encompass naturally occurring proteins as well as variations and modified forms thereof. Such variants will continue to possess the ability to implement a recombination event. Generally, the mutations made in the
10 polynucleotide encoding the variant polypeptide should not place the sequence out of reading frame, and/or create complementary regions that could produce secondary mRNA structure. See, EP Patent Application Publication No. 75,444.

Variant polynucleotides and proteins also encompass sequences and proteins derived from a mutagenic and recombinogenic procedure such as DNA shuffling.

15 With such a procedure, one or more different IL-2R α extracellular domain or IL-2 coding sequences can be manipulated to create a new IL-2R α extracellular domain or IL-2 polypeptides possessing the desired properties. In this manner, libraries of recombinant polynucleotides are generated from a population of related sequence polynucleotides comprising sequence regions that have substantial sequence identity
20 and can be homologously recombined *in vitro* or *in vivo*. Strategies for such DNA shuffling are known in the art. See, for example, Stemmer (1994) *Proc. Natl. Acad. Sci. USA* 91:10747-10751; Stemmer (1994) *Nature* 370:389-391; Crameri *et al.* (1997) *Nature Biotech.* 15:436-438; Moore *et al.* (1997) *J. Mol. Biol.* 272:336-347; Zhang *et al.* (1997) *Proc. Natl. Acad. Sci. USA* 94:4504-4509; Crameri *et al.* (1998)
25 *Nature* 391:288-291; and U.S. Patent Nos. 5,605,793 and 5,837,458.

III. Polynucleotides Encoding the IL-2/IL-2R α Fusion Proteins and Methods of Making

30 Compositions further include isolated polynucleotides that encode the various fusion proteins described herein above, and variants and fragments thereof. Vectors and expression cassettes comprising the polynucleotides described herein are further disclosed. Expression cassettes will generally include a promoter operably linked to a polynucleotide and a transcriptional and translational termination region.

The use of the term "polynucleotide" is not intended to limit the present invention to polynucleotides comprising DNA. Those of ordinary skill in the art will recognize that polynucleotides, can comprise ribonucleotides and combinations of ribonucleotides and deoxyribonucleotides. Such deoxyribonucleotides and
5 ribonucleotides include both naturally occurring molecules and synthetic analogues.

An "isolated" or "purified" polynucleotide or protein, or biologically active portion thereof, is substantially or essentially free from components that normally accompany or interact with the polynucleotide or protein as found in its naturally occurring environment. Thus, an isolated or purified polynucleotide or protein is
10 substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. Optimally, an "isolated" polynucleotide is free of sequences (optimally protein encoding sequences) that naturally flank the polynucleotide (i.e., sequences located at the 5' and 3' ends of the polynucleotide) in
15 the genomic DNA of the organism from which the polynucleotide is derived. For example, in various embodiments, the isolated polynucleotide can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb, or 0.1 kb of nucleotide sequence that naturally flank the polynucleotide in genomic DNA of the cell from which the polynucleotide is derived. A protein that is substantially free of cellular material
20 includes preparations of protein having less than about 30%, 20%, 10%, 5%, or 1% (by dry weight) of contaminating protein. When the protein of the invention or biologically active portion thereof is recombinantly produced, optimally culture medium represents less than about 30%, 20%, 10%, 5%, or 1% (by dry weight) of chemical precursors or non-protein-of-interest chemicals.

25 Conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art may be employed herein. Such techniques are explained fully in the literature. See, e.g., Sambrook *et al.*, "Molecular Cloning: A Laboratory Manual" (1989); "Current Protocols in Molecular Biology" Volumes I-III [Ausubel, R. M., ed. (1994)]; "Cell Biology: A Laboratory Handbook" Volumes I-III [J. E. Celis, ed. (1994)]; "Current Protocols in Immunology" Volumes I-III [Coligan,
30 J. E., ed. (1994)]; "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid Hybridization" [B.D. Hames & S.J. Higgins eds. (1985)]; "Transcription And Translation" [B.D. Hames & S.J. Higgins, eds. (1984)]; "Animal Cell Culture" [R.I.

Freshney, ed. (1986)]; "Immobilized Cells And Enzymes" [IRL Press, (1986)];
B. Perbal, "A Practical Guide To Molecular Cloning" (1984).

A vector which comprises the above-described polynucleotides operably
linked to a promoter is also provided herein. A nucleotide sequence is "operably
5 linked" to an expression control sequence (e.g., a promoter) when the expression
control sequence controls and regulates the transcription and translation of that
sequence. The term "operably linked" when referring to a nucleotide sequence
includes having an appropriate start signal (e.g., ATG) in front of the nucleotide
sequence to be expressed and maintaining the correct reading frame to permit
10 expression of the sequence under the control of the expression control sequence and
production of the desired product encoded by the sequence. If a gene that one desires
to insert into a recombinant nucleic acid molecule does not contain an appropriate
start signal, such a start signal can be inserted in front of the gene. A "vector" is a
replicon, such as plasmid, phage or cosmid, to which another nucleic acid segment
15 may be attached so as to bring about the replication of the attached segment. The
promoter may be, or is identical to, a bacterial, yeast, insect or mammalian promoter.
Further, the vector may be a plasmid, cosmid, yeast artificial chromosome (YAC),
bacteriophage or eukaryotic viral DNA.

Other numerous vector backbones known in the art as useful for expressing
20 protein may be employed. Such vectors include, but are not limited to: adenovirus,
simian virus 40 (SV40), cytomegalovirus (CMV), mouse mammary tumor virus
(MMTV), Moloney murine leukemia virus, DNA delivery systems, i.e. liposomes,
and expression plasmid delivery systems. Further, one class of vectors comprises
DNA elements derived from viruses such as bovine papilloma virus, polyoma virus,
25 baculovirus, retroviruses or Semliki Forest virus. Such vectors may be obtained
commercially or assembled from the sequences described by methods well-known in
the art.

A host vector system for the production of a polypeptide which comprises the
vector of a suitable host cell is provided herein. Suitable host cells include, but are
30 not limited to, prokaryotic or eukaryotic cells, e.g. bacterial cells (including gram
positive cells), yeast cells, fungal cells, insect cells, and animal cells. Numerous
mammalian cells may be used as hosts, including, but not limited to, the mouse
fibroblast cell NIH 3T3, CHO cells, HeLa cells, Ltk⁻ cells, etc. Additional animal
cells, such as R1.1, B-W and L-M cells, African Green Monkey kidney cells (e.g.,

COS 1, COS 7, BSC1, BSC40, and BMT10), insect cells (e.g., Sf9), and human cells and plant cells in tissue culture can also be used.

A wide variety of host/expression vector combinations may be employed in expressing the polynucleotide sequences presented herein. Useful expression vectors, for example, may consist of segments of chromosomal, non-chromosomal and synthetic DNA sequences. Suitable vectors include derivatives of SV40 and known bacterial plasmids, e.g., *E. coli* plasmids col El, pCR1, pBR322, pMB9 and their derivatives, plasmids such as RP4; phage DNAs, e.g., the numerous derivatives of phage λ , e.g., NM989, and other phage DNA, e.g., M13 and filamentous single stranded phage DNA; yeast plasmids such as the 2 μ plasmid or derivatives thereof; vectors useful in eukaryotic cells, such as vectors useful in insect or mammalian cells; vectors derived from combinations of plasmids and phage DNAs, such as plasmids that have been modified to employ phage DNA or other expression control sequences; and the like.

Any of a wide variety of expression control sequences (sequences that control the expression of a nucleotide sequence operably linked to it) may be used in these vectors to express the polynucleotide sequences provided herein. Such useful expression control sequences include, for example, the early or late promoters of SV40, CMV, vaccinia, polyoma or adenovirus, the *lac* system, the *trp* system, the *TAC* system, the *TRC* system, the *LTR* system, the major operator and promoter regions of phage λ , the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the promoters of acid phosphatase (e.g., Pho5), the promoters of the yeast α -mating factors, and other sequences known to control the expression of genes of prokaryotic or eukaryotic cells or their viruses, and various combinations thereof.

It will be understood that not all vectors, expression control sequences and hosts will function equally well to express the polynucleotide sequences provided herein. Neither will all hosts function equally well with the same expression system. However, one skilled in the art will be able to select the proper vectors, expression control sequences, and hosts without undue experimentation to accomplish the desired expression without departing from the scope of this invention. For example, in selecting a vector, the host must be considered because the vector must function in it. The vector's copy number, the ability to control that copy number, and the expression

of any other proteins encoded by the vector, such as antibiotic markers, will also be considered.

In selecting an expression control sequence, a variety of factors will normally be considered. These include, for example, the relative strength of the system, its
5 controllability, and its compatibility with the particular nucleotide sequence or gene to be expressed, particularly as regards potential secondary structures. Suitable unicellular hosts will be selected by consideration of, e.g., their compatibility with the chosen vector, their secretion characteristics, their ability to fold proteins correctly, and their fermentation requirements, as well as the toxicity to the host of the product
10 encoded by the nucleotide sequences to be expressed, and the ease of purification of the expression products.

In preparing the expression cassette, the various polynucleotides may be manipulated, so as to provide for the polynucleotide sequences in the proper orientation and, as appropriate, in the proper reading frame. Toward this end,
15 adapters or linkers may be employed to join the polynucleotides or other manipulations may be involved to provide for convenient restriction sites, removal of superfluous DNA, removal of restriction sites, or the like. For example, linkers such as two glycines may be added between polypeptides. Methionine residues encoded by atg nucleotide sequences may be added to allow initiation of gene transcription. For
20 this purpose, *in vitro* mutagenesis, primer repair, restriction, annealing, resubstitutions, e.g., transitions and transversions, may be involved.

Further provided is a method of producing a polypeptide which comprises expressing a polynucleotide encoding a fusion protein disclosed herein in a host cell under suitable conditions permitting the production of the polypeptide and recovering
25 the polypeptide so produced.

IV. Methods of Use

Various methods are provided for modulating an immune response. As used herein, the term “modulating” includes inducing, inhibiting,
30 potentiating, elevating, increasing, or decreasing a given activity or response.

By “subject” is intended mammals, e.g., primates, humans, agricultural and domesticated animals such as, but not limited to, dogs, cats, cattle, horses, pigs, sheep, and the like. In one embodiment, the subject undergoing treatment with the pharmaceutical formulations provided herein is a human.

A “therapeutically effective amount” of an IL-2/IL-2R α fusion protein refers to the amount of the IL-2/IL-2R α fusion protein sufficient to elicit a desired biological response. As will be appreciated by one of ordinary skill in the art, the absolute amount of a particular IL-2/IL-2R α fusion protein that is effective can vary depending on such factors as the desired biological endpoint, the IL-2/IL-2R α fusion protein to be delivered, the target cell or tissue, and the like. One of ordinary skill in the art will further understand that an effective amount can be administered in a single dose, or can be achieved by administration of multiple doses (i.e., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more doses).

i. Methods for Increasing an Immune Response

Various methods are provided for increasing the immune response in a subject. Such methods comprise administering to a subject in need of an increase in the immune response a therapeutically effective amount of an IL-2/IL-2R α fusion protein. As such, in specific embodiments, transient application of higher doses of IL-2 are employed to boosted immune effector and memory responses.

It is further recognized that the various IL-2/IL-2R α fusion protein can be used in combination with an antigen to enhance the immune response to the antigen. Thus, the IL-2/IL-2R α fusion protein can also be used as a vaccine adjuvant especially to boost cell-mediated immune memory.

For example, the IL-2/IL-2R α fusion protein can be used to enhance a vaccine preparation. Thus, the various IL-2/IL-2R α fusion proteins are useful for increasing the efficacy of anti-cancer vaccines or for vaccines that are poorly immunogenic. Further provided are methods for enhancing the efficacy or immunogenicity of a vaccine in a subject, or overcoming a suppressed immune response to a vaccine in a subject, including (i) administering to the subject a therapeutically effective amount of an IL-2/IL-2R α fusion protein and (ii) administering to the subject a vaccine.

By “vaccine” is intended a composition useful for stimulating a specific immune response (or immunogenic response) in a subject. In some embodiments, the immunogenic response is protective or provides protective immunity. For example, in the case of a disease-causing organism the vaccine enables the subject to better resist infection with or disease progression from the organism against which the vaccine is directed. Alternatively, in the case of a cancer, the vaccine strengthens the subject’s natural defenses against cancers that have already developed. These types

of vaccines may also prevent the further growth of existing cancers, prevent the recurrence of treated cancers, and/or eliminate cancer cells not killed by prior treatments.

Representative vaccines include, but are not limited to, vaccines against
5 diphtheria, tetanus, pertussis, polio, measles, mumps, rubella, hepatitis B,
Haemophilus influenzae type b, varicella, meningitis, human immunodeficiency virus,
tuberculosis, Epstein Barr virus, malaria, hepatitis E, dengue, rotavirus, herpes,
human papillomavirus, and cancers. Vaccines of interest include the two vaccines
that have been licensed by the U.S. Food and Drug Administration to prevent virus
10 infections that can lead to cancer: the hepatitis B vaccine, which prevents infection
with the hepatitis B virus, an infectious agent associated with liver cancer (*MMWR
Morb. Mortal. Wkly. Rep.* 46:107-09, 1997); and Gardasil™, which prevents infection
with the two types of human papillomavirus that together cause 70 percent of cervical
cancer cases worldwide (Speck and Tyring, *Skin Therapy Lett.* 11:1-3, 2006). Other
15 treatment vaccines of interest include therapeutic vaccines for the treatment of cancer,
cervical cancer, follicular B cell non-Hodgkin's lymphoma, kidney cancer, cutaneous
melanoma, ocular melanoma, prostate cancer, and multiple myeloma.

By “enhancing the efficacy” or “enhancing the immunogenicity” with regard
to a vaccine is intended improving an outcome, for example, as measured by a change
20 in a specific value, such as an increase or a decrease in a particular parameter of an
activity of a vaccine associated with protective immunity. In one embodiment,
enhancement refers to at least a 5%, 10%, 25%, 50%, 100% or greater than 100%
increase in a particular parameter. In another embodiment, enhancement refers to at
least a 5%, 10%, 25%, 50%, 100% or greater than 100% decrease in a particular
25 parameter. In one example, enhancement of the efficacy/immunogenicity of a
vaccine refers to an increase in the ability of the vaccine to inhibit or treat disease
progression, such as at least a 5%, 10%, 25%, 50%, 100%, or greater than 100%
increase in the effectiveness of the vaccine for that purpose. In a further example,
enhancement of the efficacy/immunogenicity of a vaccine refers to an increase in the
30 ability of the vaccine to recruit the subject's natural defenses against cancers that have
already developed, such as at least a 5%, 10%, 25%, 50%, 100%, or greater than
100% increase in the effectiveness of the vaccine for that purpose.

Similarly, by “overcoming a suppressed immune response” with regard to a
vaccine is intended improving an outcome, for example, as measured by a change in a

specific value, such as a return to a formerly positive value in a particular parameter of an activity of a vaccine associated with protective immunity. In one embodiment, overcoming refers to at least a 5%, 10%, 25%, 50%, 100% or greater than 100% increase in a particular parameter. In one example, overcoming a suppressed immune response to a vaccine refers to a renewed ability of the vaccine to inhibit or treat disease progression, such as at least a 5%, 10%, 25%, 50%, 100%, or greater than 100% renewal in the effectiveness of the vaccine for that purpose. In a further example, overcoming a suppressed immune response to a vaccine refers to a renewed ability of the vaccine to recruit the subject's natural defenses against cancers that have already developed, such as at least a 25%, 50%, 100%, or greater than 100% renewal in the effectiveness of the vaccine for that purpose.

By "therapeutically effective amount" is intended an amount that is useful in the treatment, prevention or diagnosis of a disease or condition. As used herein, a therapeutically effective amount of an IL-2/IL-2R α fusion protein is an amount which, when administered to a subject, is sufficient to achieve a desired effect, such as modulating an immune response in a subject without causing a substantial cytotoxic effect in the subject. As outlined above, a therapeutically effective amount of an IL-2/IL-2R α fusion protein can be administered to a subject to increase an immune response, enhance the immune response to an antigen, enhance the efficacy or immunogenicity of a vaccine in a subject, or to overcome a suppressed immune response to a vaccine. The effective amount of an IL-2/IL-2R α fusion protein useful for modulating such functions will depend on the subject being treated, the severity of the affliction, and the manner of administration of the IL-2/IL-2R α fusion protein. Exemplary doses include about 10^4 to about 10^7 IU of IL-2 activity per adult, about 10^4 to 10^5 IU of IL-2 activity per adult, about 10^5 to about 10^6 IU of IL-2 activity per adult, about 10^6 to about 10^7 IU of IL-2 activity per adult. In other instances, the therapeutically effective dose of the IL-2/IL-2R α fusion protein is about 10^5 IU of IL-2 activity $\pm$ 100-fold, is about 10^5 IU of IL-2 activity $\pm$ 10-fold, about 10^5 IU of IL-2 activity $\pm$ 2-fold, about 10^5 IU of IL-2 activity $\pm$ 20-fold, about 10^5 IU of IL-2 activity $\pm$ 30-fold, about 10^5 IU of IL-2 activity $\pm$ 40-fold, about 10^5 IU of IL-2 activity $\pm$ 50-fold, about 10^5 IU of IL-2 activity $\pm$ 60-fold, about 10^5 IU of IL-2 activity $\pm$ 70-fold, about 10^5 IU of IL-2 activity $\pm$ 80-fold, or about 10^5 IU of IL-2 activity $\pm$ 90-fold. In a specific non-limiting embodiment, a human IL-2 fusion protein is administered at this dosage.

In one embodiment, the reference standard for the mouse IL-2 fusion protein is the mouse IL-2 is from eBiosciences (Catalog Number: 14-8021). Briefly, the bioactivity of mouse IL-2 from eBioscience is as follows: The ED50 of this protein, as measured by CTLL-2 cell proliferation assay, is less than or equal to 175 pg/mL.

5 This corresponds to a specific activity of greater than or equal to 5.7×10^6 Units/mg.

In another embodiment, the reference standard for the human IL-2 fusion protein is the human IL-2 drug Aldesleukin (Proleukin). Thus, the IL-2 fusion proteins disclosed herein are directly compared to the fusion protein to the IL-2 drug that is used in low dose or high dose IL-2 therapy. IL-2 activity for mouse and human
10 IL-2 use the same assay and their activity in units/mg are similar. With respect to the human IL-2 drug, i.e. aldesleukin (Proleukin), the standard measure of an amount IL-2 is the International Unit (IU) which technically is not a fixed amount but the amount that produces a fixed effect in a specific assay of biological activity, i.e. CTLL proliferation assay. In practice, the manufacture of IL-2 is standardized and there is a
15 conversion between drug weight and International Units. It is $1.1 \text{ mg IL-2} = 18$ million IU (abbreviated 18 MIU).

It is furthermore understood that appropriate doses of a functional agent depend upon the potency of the active agent with respect to the activity to be modulated. Such appropriate doses may be determined using the assays described
20 herein. In addition, it is understood that the specific dose level for any particular animal subject will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, gender, and diet of the subject, the time of administration, the route of administration, the rate of excretion, and/or any drug combination.

25 When administration is for the purpose of treatment, administration may be for either a prophylactic or therapeutic purpose. When provided prophylactically, the substance is provided in advance of any symptom. The prophylactic administration of the substance serves to prevent or attenuate any subsequent symptom. When provided therapeutically, the substance is provided at (or shortly after) the onset of a
30 symptom. The therapeutic administration of the substance serves to attenuate any actual symptom.

The skilled artisan will appreciate that certain factors may influence the dosage required to effectively treat a subject, including but not limited to, the severity of the disease or disorder, previous treatments, the general health and/or age of the

subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of an IL-2/IL-2R α fusion protein can include a single treatment or, preferably, can include a series of treatments. It will also be appreciated that the effective dosage of an IL-2/IL-2R α fusion protein used for treatment may
5 increase or decrease over the course of a particular treatment. Changes in dosage may result and become apparent from the results of diagnostic assays as described herein.

Therapeutically effective amounts of an IL-2/IL-2R α fusion protein can be determined by animal studies. When animal assays are used, a dosage is administered to provide a target *in vivo* concentration similar to that which has been shown to be
10 effective in the animal assays.

ii. Methods for Decreasing an Immune Response

Various methods are provided for decreasing the immune response in a subject. Such methods comprise administering to a subject in need of a decrease in
15 the immune response a therapeutically effective amount of an IL-2/IL-2R α fusion protein.

There is much interest to harness the suppressive power of Tregs to inhibit unwanted immune responses. Data in mouse and man shows that enhancing IL-2R signaling with a low dose of IL-2 selectively boosts Tregs and enhances immune
20 tolerogenic mechanisms. IL-2/IL-2R α fusion proteins provided herein represent a new and improved form of IL-2 that more potentially enhances Tregs. Thus, the IL-2/IL-2R α fusion proteins can be administered to patients with autoimmune diseases, chronic graft versus host disease, transplant rejection reactions, and other conditions where the goal is to suppress self-reactivity.

For example, a therapeutically effective amount of an IL-2/IL-2R α fusion protein that promotes immune tolerance can find use, for example, in treating a subject having an autoimmune or an inflammatory disorder, including but not limited to, graft rejections and allergies. Thus, in one embodiment, a method of treating a subject having an autoimmune or inflammatory disorder is provided. Such a method
25 comprises administering to the subject a therapeutically effective amount of an IL-2/IL-2R α fusion protein.
30

Non-limiting examples of autoimmune disorders that can be treated or prevented include type1 diabetes, multiple sclerosis, rheumatoid arthritis, celiac disease, systemic lupus erythematosus, juvenile idiopathic arthritis,

Crohn's disease, ulcerative colitis or systemic sclerosis, graft versus host disease, HCV-induced vasculitis, alopecia areata or psoriasis.

Additional autoimmune diseases include those where there is already an indication that Tregs may be impaired and would benefit from IL-2-dependent boosting of Tregs. In this regard, single nucleotide polymorphisms (SNPs) in IL-2, IL-2R α , or IL-2R13 have been associated as a genetic risk for type 1 diabetes, multiple sclerosis, rheumatoid arthritis, celiac disease, systemic lupus erythematosus, juvenile idiopathic arthritis, Crohn's disease, ulcerative colitis, and systemic sclerosis. Studies suggest that the genetic risk is related to impaired Treg numbers and/or activity. In addition, low dose IL-2 therapy has shown to benefit patients with chronic GvHD and HCV-induced vasculitis. Thus, such patients populations can also be administered a therapeutically effective amount of an IL-2/IL-2R α fusion protein.

In other embodiments, the IL-2/IL-2R α fusion protein can be used in combination with a therapeutic agent to reduce the immune response to the agent (i.e. protein). For example, the IL-2/IL-2R α fusion protein can be used in combination with a therapeutic protein which must be chronically administered to a subject. Thus, in a specific embodiment, the method comprises includes administering to the subject at least one additional therapeutic agent in combination with an IL-2/IL-2R α fusion protein. Such therapeutic agents, include but are not limited to, a cytokine, a glucocorticoid, an anthracycline (*e.g.*, doxorubicin or epirubicin), a fluoroquinolone (*e.g.*, ciprofloxacin), an antifolate (*e.g.*, methotrexate), an antimetabolite (*e.g.*, fluorouracil), a topoisomerase inhibitor (*e.g.*, camptothecin, irinotecan or etoposide), an alkylating agent (*e.g.*, cyclophosphamide, ifosfamide, mitolactol, or melphalan), an antiandrogen (*e.g.*, flutamide), an antiestrogen (*e.g.*, tamoxifen), a platinum compound (*e.g.*, cisplatin), a vinca alkaloid (*e.g.*, vinorelbine, vinblastine or vindesine), or mitotic inhibitor (*e.g.*, paclitaxel or docetaxel).

Moreover, the therapeutically effective amount of the IL-2/IL-2R α fusion protein can further be administered in combination therapies to increase Tregs and tolerance. Such combination therapies can comprises the therapeutically effective amount of the IL-2/IL-2R α fusion protein in combination with anti-TNF α or other agents to inhibit inflammatory responses.

The therapeutically effective amount of an IL-2/IL-2R α fusion protein useful for decreasing an immune response will depend on the subject being treated, the

severity of the affliction, and the manner of administration of the IL-2/IL-2R α fusion protein. Exemplary doses include about 10^3 IU to about 10^6 IU of IL-2 activity per adult or about 10^4 IU to about 10^6 IU of IL-2 activity per adult. Exemplary doses include about 10^3 to about 10^6 IU of IL-2 activity per adult, about 10^3 to about 10^4 IU of IL-2 activity per adult, about 10^4 to about 10^6 IU of IL-2 activity per adult, about 10^4 to 10^5 IU of IL-2 activity per adult, or about 10^5 to about 10^6 IU of IL-2 activity per adult. In other instances, the therapeutically effective dose of the IL-2/IL-2R α fusion protein is about 10^4 IU of IL-2 activity $\pm$ 100-fold, is about 10^4 IU of IL-2 activity $\pm$ 10-fold, about 10^4 IU of IL-2 activity $\pm$ 2-fold, about 10^4 IU of IL-2 activity $\pm$ 20-fold, about 10^4 IU of IL-2 activity $\pm$ 30-fold, about 10^4 IU of IL-2 activity $\pm$ 40-fold, about 10^4 IU of IL-2 activity $\pm$ 50-fold, about 10^4 IU of IL-2 activity $\pm$ 60-fold, about 10^4 IU of IL-2 activity $\pm$ 70-fold, about 10^4 IU of IL-2 activity $\pm$ 80-fold, or about 10^4 IU of IL-2 activity $\pm$ 90-fold. In a specific non-limiting embodiment, a human IL-2 fusion protein is administered at this dosage.

In one embodiment, the reference standard for the mouse IL-2 fusion protein is the mouse IL-2 is from eBiosciences (Catalog Number: 14-8021). Briefly, the bioactivity of mouse IL-2 from eBioscience is as follows: The ED50 of this protein, as measured by CTLL-2 cell proliferation assay, is less than or equal to 175 pg/mL. This corresponds to a specific activity of greater than or equal to 5.7×10^6 Units/mg.

In another embodiment, the reference standard for the human IL-2 fusion protein is the human IL-2 drug Aldesleukin (Proleukin). Thus, the IL-2 fusion proteins disclosed herein are directly compared to the fusion protein to the IL-2 drug that is used in low dose or high dose IL-2 therapy. IL-2 activity for mouse and human IL-2 use the same assay and their activity in units/mg are similar. With respect to the human IL-2 drug, i.e. aldesleukin (Proleukin), the standard measure of an amount IL-2 is the International Unit (IU) which technically is not a fixed amount but the amount that produces a fixed effect in a specific assay of biological activity, i.e. CTLL proliferation assay. In practice, the manufacture of IL-2 is standardized and there is a conversion between drug weight and International Units. It is $1.1 \text{ mg IL-2} = 18 \text{ million IU}$ (abbreviated 18 MIU).

It is furthermore understood that appropriate doses of a functional agent depend upon the potency of the active agent with respect to the expression or activity to be modulated. Such appropriate doses may be determined using the assays described herein. In addition, it is understood that the specific dose level for any

particular animal subject will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, gender, and diet of the subject, the time of administration, the route of administration, the rate of excretion, and/or any drug combination.

5 When administration is for the purpose of treatment, administration may be for either a prophylactic or therapeutic purpose. When provided prophylactically, the substance is provided in advance of any symptom. The prophylactic administration of the substance serves to prevent or attenuate any subsequent symptom. When provided therapeutically, the substance is provided at (or shortly after) the onset of a
10 symptom. The therapeutic administration of the substance serves to attenuate any actual symptom.

 The skilled artisan will appreciate that certain factors may influence the dosage required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the
15 subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of an IL-2/IL-2R α fusion protein can include a single treatment or, preferably, can include a series of treatments. It will also be appreciated that the effective dosage of an IL-2/IL-2R α fusion protein used for treatment may increase or decrease over the course of a particular treatment. Changes in dosage may
20 result and become apparent from the results of diagnostic assays as described herein.

 Therapeutically effective amounts of an IL-2/IL-2R α fusion protein can be determined by animal studies. When animal assays are used, a dosage is administered to provide a target tissue concentration similar to that which has been shown to be effective in the animal assays.

25

iii. Pharmaceutical Composition

 The various IL-2/IL-2R α fusion proteins disclosed herein (also referred to herein as "active compounds") can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the fusion protein
30 and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active

substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

5 A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), and transmucosal. In addition, it may be desirable to administer a therapeutically effective amount of the pharmaceutical composition locally to an
10 area in need of treatment. This can be achieved by, for example, local or regional infusion or perfusion during surgery, topical application, injection, catheter, suppository, or implant (for example, implants formed from porous, non-porous, or gelatinous materials, including membranes, such as sialastic membranes or fibers), and the like. In another embodiment, the therapeutically effective amount of the
15 pharmaceutical composition is delivered in a vesicle, such as liposomes (see, e.g., Langer, *Science* 249:1527-33, 1990 and Treat *et al.*, in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez Berestein and Fidler (eds.), Liss, N.Y., pp. 353-65, 1989).

 In yet another embodiment, the therapeutically effective amount of the
20 pharmaceutical composition can be delivered in a controlled release system. In one example, a pump can be used (see, e.g., Langer, *Science* 249:1527-33, 1990; Sefton, *Crit. Rev. Biomed. Eng.* 14:201-40, 1987; Buchwald *et al.*, *Surgery* 88:507-16, 1980; Saudek *et al.*, *N. Engl. J. Med.* 321:574-79, 1989). In another example, polymeric materials can be used (see, e.g., Levy *et al.*, *Science* 228:190-92, 1985; During *et al.*,
25 *Ann. Neurol.* 25:351-56, 1989; Howard *et al.*, *J. Neurosurg.* 71:105-12, 1989). Other controlled release systems, such as those discussed by Langer (*Science* 249:1527-33, 1990), can also be used.

 Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for
30 injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be

adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes, or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile
5 aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL9 (BASF; Parsippany, NJ), or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the
10 extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures
15 thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion, and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will
20 be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride, in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active
25 compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the
30 preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying, which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from a pressurized container or dispenser that contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means.

5 For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the
10 active compounds are formulated into ointments, salves, gels, or creams as generally known in the art. The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will
15 protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the
20 art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No.
25 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated with each unit containing a predetermined quantity of active
30 compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the

limitations inherent in the art of compounding such a functional compound for the treatment of individuals.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

5

iv. Kits

As used herein, a “kit” comprises an IL-2/IL-2R α fusion protein for use in modulating the immune response, as described elsewhere herein. The terms “kit” and “system,” as used herein are intended to refer to at least one or more IL-2/IL-2R α fusion protein which, in specific embodiments, are in combination with one or more other types of elements or components (*e.g.*, other types of biochemical reagents, containers, packages, such as packaging intended for commercial sale, instructions of use, and the like).

15 *V. Sequence Identity*

As described above, active variants and fragments of the IL-2/IL-2R α fusion proteins or the polynucleotide encoding the same, including the various components of the IL-2/IL-2R α fusion protein are provided. Such components include, IL-2, the extracellular domain of IL-2R α , the linker sequences or the Kozak sequence. The activity retained by the active variant or fragment of the fusion protein or a given component of the fusion protein is discussed in further detail elsewhere herein.

Such variants can have at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to a given reference polypeptide or polynucleotide. A fragment can comprise at least 10, 20, 30, 50, 75, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1500, 2000 contiguous nucleotides of a given reference nucleotide sequence or up to the full length of a given nucleotide reference sequence; or a fragment can comprise at least 10, 20, 30, 40, 50, 60, 70, 75, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200 contiguous amino acids or up to the full length of a given reference polypeptide sequence.

As used herein, “sequence identity” or “identity” in the context of two polynucleotides or polypeptide sequences makes reference to the residues in the two sequences that are the same when aligned for maximum correspondence over a specified comparison window. When percentage of sequence identity is used in reference to proteins it is recognized that residue positions which are not identical

often differ by conservative amino acid substitutions, where amino acid residues are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. When sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the substitution. Sequences that differ by such conservative substitutions are said to have "sequence similarity" or "similarity". Means for making this adjustment are well known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score between zero and 1. The scoring of conservative substitutions is calculated, e.g., as implemented in the program PC/GENE (Intelligenetics, Mountain View, California).

As used herein, "percentage of sequence identity" means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison, and multiplying the result by 100 to yield the percentage of sequence identity.

Unless otherwise stated, sequence identity/similarity values provided herein refer to the value obtained using GAP Version 10 using the following parameters: % identity and % similarity for a nucleotide sequence using GAP Weight of 50 and Length Weight of 3, and the nwsgapdna.cmp scoring matrix; % identity and % similarity for an amino acid sequence using GAP Weight of 8 and Length Weight of 2, and the BLOSUM62 scoring matrix; or any equivalent program thereof. By "equivalent program" is intended any sequence comparison program that, for any two sequences in question, generates an alignment having identical nucleotide or amino acid residue matches and an identical percent sequence identity when compared to the corresponding alignment generated by GAP Version 10.

As used herein, the singular terms “a,” “an,” and “the” include plural referents unless context clearly indicates otherwise. Similarly, the word “or” is intended to include “and” unless the context clearly indicates otherwise. It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or
5 molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description.

The subject matter of the present disclosure is further illustrated by the following non-limiting examples.

10 EXPERIMENTAL

IL-2 is a biologic that has been used in attempts to boost immune responses in cancer and HIV/AIDS patients. More recently much lower doses of IL-2 have been used to selectively boost tolerance to suppress unwanted immune responses associated with autoimmune-like attack of self-tissues. Importantly, these low doses of IL-2 have not
15 shown signs of enhancing or re-activation of autoreactive T cells. Nevertheless, IL-2 has important drawbacks as a therapeutic, including a very short-half life *in vivo*, which limits its efficacy, and toxicity at high doses. For these reasons a new IL-2 biologic has been produced with the goals of improving its pharmacokinetics and durability of responses for use 1) in low dose IL-2-based therapy to boost regulatory T
20 cells (Tregs) and immune tolerance and 2) in adjuvant therapy with higher doses to boost immune responses and memory. To achieve these goals, IL-2/IL-2R α fusion proteins have been developed, where these fusions were designed to increase IL-2 availability by increasing persistent IL-2 stimulation of IL-2R-bearing lymphocytes *in vivo*. These fusions consist of engineered proteins as follows (FIG. 1): 1) a leader
25 sequence of IL-2 that contains an optimized Kozak sequence for efficient translation; 2) the full length sequence of IL-2; 3) a glycine or glycine/serine linker sequence of variable length; 4) the coding sequence of the expressed extracellular domain of IL-2R α ; 5) a 2 amino acid glycine spacer; 6) a six amino acid poly-histidine region for purification; and 7) two termination codons. The predicted protein sequences from
30 these mouse and human cDNAs are shown for IL-2/(GlySer)/IL-2R α fusion proteins in FIG. 2A and FIG. 2B, respectively. These cDNAs were cloned into the pCIneo expression vector and used for expression of these fusion proteins in COS7 cells. Analysis of the culture supernatants indicated that each mouse fusion protein exhibited IL-2 bioactivity *in vitro*, with optimal activity associated with the IL-2/(Gly₃Ser)₃/IL-

2R α fusion protein (FIG. 3A). Accordingly, inclusion of anti-IL-2 in this bioassay completely inhibited proliferation (FIG. 3B). Larger amounts of IL-2/(Gly₃Ser)₃/IL-2R α and IL-2/(Gly₄Ser)₄/IL-2R α were prepared after expression in CHO cells and purified by affinity chromatography through binding of the 6x His tag of the fusion protein to immobilized nickel. The mouse IL-2/(Gly₃Ser)₃/IL-2R α fusion protein showed greater IL-2 bioactivity than IL-2/(Gly₄Ser)₄/IL-2R α (FIG. 4A) even though both fusion proteins similarly inhibited the binding of two anti-IL2R α antibodies (PC61 and 7D4) (FIG. 4B) to cells expressing IL-2R α , confirming greater IL-2 activity is associated with the former fusion protein. The inhibition of binding of PC61 and 7D4 also indicates that IL-2R α portion of the fusion protein retained sufficient tertiary structure to bind these antibodies. However, these fusion proteins did not inhibit the binding of a monoclonal antibody (3C7) directed to the IL-2 binding site of IL-2R α , to cells expressing IL-2R α . This result implies that IL-2 within the IL-2/IL-2R α fusion protein is spatially near the binding site of IL-2R α (FIG. 5). Western blot analysis of these fusion proteins showed that IL-2/IL-2R α was 55-65 kDa, with somewhat faster mobility under non-reducing condition, and that it was approximately 15 kDa larger than that observed for soluble IL-2R α (FIG. 6A). Correspondingly, direct analysis of the purified mouse IL-2/(Gly₃Ser)₃/IL-2R α by SDS-PAGE was consistent with a heterogeneous 55-65 kDa monomer protein (FIG. 6B), which is the expected size for an IL-2 (15 kDa) and IL-2R α (40-50kDa) fusion molecule (FIG. 6), where IL-2R α shows size heterogeneity due to extensive variable glycosylation (Malek and Korthy, *J. Immunol.* 136:4092-4098, 1986). An immediate consequence of IL-2-dependent signal transduction is tyrosine phosphorylation of STAT5 (pSTAT5). Treatment of mice with mouse IL-2/(Gly₃Ser)₃/IL-2R α resulted in extensive and selective activation of pSTAT5 in Tregs 30 min post-treatment (FIG. 7). Dose-response studies showed that mouse IL-2/(Gly₃Ser)₃/IL-2R α affected a number of key activities of Tregs *in vivo* (FIG. 8). These effects on Tregs included: increased representation (FIG. 8A) and number (not shown) of Tregs in the CD4⁺ T cell compartment; upregulation of IL-2-dependent CD25 (FIG.8B); increased proliferation as assessed by expression of the proliferative marker Ki67 (FIG. 8C); and increased fraction of IL-2-dependent terminally differentiated Klrp1⁺ Treg subset (FIG. 8D). These effects were most striking for Tregs in the spleen and the inflamed pancreas of non-obese diabetic (NOD) mice. 1000 units of IL-2 activity, as measured in the standard CTLL IL-2 bioassay,

associated with IL-2/(Gly₃Ser)₃/IL-2R α showed lower, but readily measurable effects on Tregs (FIG. 8). C57/BL6 mice treated with IL-2/(Gly₃Ser)₃/IL-2R α (2000 units of IL-2 activity) were compared to mice that received recombinant IL-2 (25,000 units) or agonist complexes of IL-2/anti-IL-2 (IL2/IC) (10,000 units of IL-2 activity) (FIG. 9).

5 IL-2/(Gly₃Ser)₃/IL-2R α was much more effective than recombinant IL-2 and slightly more effective than IL2/IC in inducing persistent augmentation of Tregs and related properties (FIG. 9). These increases in tolerogenic Tregs occurred at 5- and 12.5-fold lower levels of IL-2 activity in comparison to IL2/IC and recombinant IL-2, respectively. When considering IL-2-dependent activation of pSTAT5 in Tregs
10 directly ex vivo (FIG. 9), these data suggest a biological half-life of approximately 72 hours for IL-2/IL-2R α . Pre-diabetic NOD mice underwent a short course of treatment with low amounts of IL-2/IL-2R α (FIG. 10). A delay in the onset of diabetes was observed in those mice that were treated with 800 U of IL-2 activity associated with IL-2/IL-2R α . With respect to immunity, application of a single high dose of IL-
15 2/(Gly₃Ser)₃/IL-2R α (12,000 U of IL-2 activity) also substantially boosted CD8⁺ T cell responses, especially long-lived memory cells (FIG. 11). Early after immunization (day 28), CD44^{hi} CD62L^{lo} CD127^{hi} effector-memory (EM) cells dominated the memory pool; however, with increasing time CD44^{hi} CD62L^{hi} CD127^{hi} central memory (CM) cells increased, and CM cells dominated the memory pool 202 days
20 post-immunization (FIG. 12). Thus, IL-2/(Gly₃Ser)₃/IL-2R α functions in an analogous manner to recombinant IL-2 to boost tolerogenic and immune suppressive Tregs and immunity through increasing T effector/memory responses, but it exhibits improved pharmacokinetics by delivering such responses: 1) at a lower effective levels of IL-2 activity; 2) with more persistent biological responses; and 3) retaining the hierarchy
25 with Tregs responsive at lower doses than T effector/memory cells. These findings support the notion that IL-2/IL-2R α fusion proteins represent an improved and new class of drugs to deliver IL-2 activity to selectively boost immune tolerance or immune memory when administered at the proper dose and regimen.

IL-2/IL-2R α fusion proteins were also produced that comprise human IL-2
30 and human IL-2R α (FIG. 1, FIG. 2B). These cDNAs were expressed in CHO cells and the secreted fusion proteins were purified on Nickel affinity chromatography based on the 6x-His tag. Fusion proteins varied in the length of the glycine/serine linkers in an analogous manner to those used for mouse IL-2/IL-2R α . All 4 of the

resulting human IL-2/IL-2R α fusion proteins exhibited IL-2 bioactivity using the mouse CTLL assay (FIG. 13A). Western blot analysis confirmed that human IL-2/IL-2R α also showed a heterogeneous band between 55-60 kDa (FIG. 13B), consistent with highly glycosylated molecules expected for IL-2 linked to IL-2R α .

5 The IL-2/IL-2R α fusion proteins with the (G₃S)₃ and especially the (G₄S)₄ linkers may have greater activity because less fusion protein was seen even though equivalent amount of IL-2 activity was loaded on each lane (FIG. 13B). The capacity of the fusion protein to inhibit the binding of anti-IL-2R α monoclonal antibodies, M-A257 and BC96, to cells bearing human IL-2R α indicates that IL-2R α of the fusion protein retained sufficient tertiary structure to bind these antibodies (FIG. 14). However, these fusion proteins only partially inhibited the binding of a monoclonal antibody (BC96) directed to the IL-2 binding site of IL-2R α , implying that IL-2 within the IL-2/IL-2R α fusion protein is spatially near the binding site of IL-2R α . Moreover, we estimated the specific activity of the mouse and human IL-2/IL-2R α fusion proteins containing the (G₃S)₃ linker to be 80 and 2000 pM, respectively, for 1 unit/ml of IL-2 bioactivity activity. These values are much higher than the activity of recombinant IL-2, which is 10 pM at 1 unit/ml. The distinct activities between human and mouse IL-2/IL-2R α is at least partially accounted for by a relative ineffectiveness of the human fusion protein to support the proliferation of mouse CTLL cells in the bioassay compared to the mouse fusion proteins or mouse and human recombinant IL-2 (not shown). These relatively low specific activities and the antibody blocking results (FIG. 5 and FIG. 12) raised the possibility that there is a specific intramolecular interaction between IL-2 and IL-2R α in the context of the fusion protein that limits the amount of IL-2 in the fusion protein to stimulate cells bearing the IL-2R. To directly test this notion, two arginine residues within the IL-2 binding site of human IL-2R α (see Robb *et al.*, *Proc. Natl. Acad. Sci. USA*, 85:5654-5658, 1988) were mutated to threonine and serine. We detected much greater bioactivity associated with these mutant IL-2R fusion proteins (FIG. 15); the specific activity of the mutated IL-2/IL-2R α fusion proteins was estimated to be approximately 5 pM for 1 unit/ml of IL-2 activity, a value very similar to recombinant IL-2. Thus, these data indicate that human IL-2/IL-2R α is biologically active and one specific mechanism of action that accounts for the prolonged IL-2 activity in these fusion proteins is through a competitive interaction

between the IL-2 moiety with the IL-2 binding region of IL-2R α of the fusion protein and with cells that express the IL-2R.

Table 1. Summary of Sequences

SEQ ID NO	AA/NT	Source	Description	
1	AA	human	IL-2-unprocessed	GenBank Acc. No. AAB46883 IL-2 myrmqlisci alsalvtns aptssstkt qqlqlehlld lqmilnginn yknpltrmltlfkfympkka telkhlqcle eelkpleevl nlaqsknfhl rprdlisin vivlelkgsettfmceyade tativeflnr witfcqsiis tlt
2	AA	Human	IL-2 – mature form	GenBank AAB46883 with first 20 aa removed aptssstkt qqlqlehlld lqmilnginn yknpltrmltlfkfympkka telkhlqcle eelkpleevl nlaqsknfhl rprdlisin vivlelkgse ttfmceyade tativeflnr witfcqsiis tlt
3	AA	Mouse	IL-2 unprocessed	Acc No. P04351 MYSMQLASCV TLTLVLLVNS APTSSSTSSS TAEAQQQQQQ QQQQQQHLEQ LLMDLQELLS RMENYRNKL PRMLTFKFYL PKQATELKDL QCLEDELGPL RHVLDLTQSK SFQLEDAENF ISNIRVTVVK LKGSNTFEC QFDESATVV DFLRRWIAFC QSIISTSPQ
4	AA	Mouse	IL-2 mature form	Mature form of Acc No. P04351 APTSSSTSSS TAEAQQQQQQ QQQQQQHLEQ LLMDLQELLS RMENYRNKL PRMLTFKFYL PKQATELKDL QCLEDELGPL RHVLDLTQSK SFQLEDAENF ISNIRVTVVK LKGSNTFEC QFDESATVV DFLRRWIAFC QSIISTSPQ
5	AA	Human	IL-2R α unprocessed form	Genebank Acc No. NP_000408.1 mdsyllmwgl ltfimvpgcq aelcdddppe iphatfkama ykegtmlnce ckrgfrrikslymlctgn sshsswdnqc qctssatnt tkqvtpppee qkerkttemq spmqpvdqaslpghcreppp weneateriy hfvgqmvvy qcvqgyralh rgpaesvckm thgktrwtqpqlctgemet sqfpgeekpq aspegrpese tscvttdf qiqtemaatm etsiftteyqvavagcvfl isvllsglt wqrrqrksrr ti
6	AA	Human	IL-2R α mature form	First 1-21 AA removed from NP_000408.1 elcdddppe iphatfkama ykegtmlnce ckrgfrrikslymlctgn sshsswdnqc qctssatnt tkqvtpppee qkerkttemq spmqpvdqas lpghcreppp weneateriy hfvgqmvvy qcvqgyralh rgpaesvckm thgktrwtqpqlctgemet sqfpgeekpq aspegrpese tscvttdf qiqtemaatm etsiftteyqvavagcvfl isvllsglt wqrrqrksrr ti
7	AA	Human	Mature form of IL-2R α extracellular domain	ELCDDDPPEIPHATFKAMAYKEGTMLNCECKRGFRRIKSGSLYMLCTGN SSHSSWDNQCCQCTSSATRNNTKQVTPQPEEQKERKTTEMQSPMQPVD QASLPGHCREPPPWENEATERIYHFVVGQMVVYQCVQGYRALHRGPA ESVCKMTHGKTRWTQPQLICTGEMETSQFPGEEKPQASPEGRPESETS CLVTTDFQIQTEMAATMETSIFTTEYQ
8	AA	Mouse	IL-2R α unprocessed form	Acc No. NP_032393.3 meprllmlgf lsitvpscr aelclydppe vpatfkals ykngtilnce ckrgfrrikelvymrlgns wssncqctsn shdksrkvt aqlehqkeqq tttdmqkptq smhqenltghcrepppwkhe dskriyhve gqsvhyecip gykalrgpa isickmkcgk tgwtqpqlctvderehrfl aseesqgsrn sspesetscp ittdfpqpt ettamtetfv ltmeykvavascflililsgltwqhr wrksrti
9	AA	Mouse	IL-2R α mature form	aa 1-21 removed from Acc No. NP_032393.3 elclydppe vpatfkals ykngtilnce ckrgfrlike lymrlgns wssncqctsn shdksrkvt aqlehqkeqq tttdmqkptq smhqenltgh crepppwkhe dskriyhve gqsvhyecip gykalrgpa isickmkcgk tgwtqpqlct vderehrfl aseesqgsrn sspesetscp ittdfpqpt ettamtetfv ltmeykva scflilil ilsgltwqhr wrksrti

				ACAACAGACATGCAGAAGCCAACACAGTCTATGCACCAAGAGAACCT TACAGGTCAGTGCAGGGAGCCACCTCCTTGGAAACATGAAGATTCCA AGAGAATCTATCATTTCTGGAAGGACAGAGTGTTCACTACGAGTGT ATTCCGGGATACAAGGCTCTACAGAGAGGTCTGCTATTAGCATCTG CAAGATGAAGTGTGGGAAAACGGGGTGGACTCAGCCCCAGCTCACA TGTGTAGATGAAAGAGAACACCACCGATTCTGGCTAGTGAGGAATC TCAAGGAAGCAGAAATTTCTCTCCGAGAGTGAGACTTCTGCCCA TAACCACCACAGACTTCCACAACCCACAGAAACAAGTGAATGACG GAGACATTTGTGCTCACAAATGGAGTATAAGGGTGGACATCACCATCA CCATCACTAATAA
32	NT	Human	Unprocessed form IL-2 (Gly4Ser)4- extracellular domain of IL-2 R α	ATGGACAGGATGCAACTCCTGTCTTGCAATTGCACTAAGTCTTGCACTT GTCACAAACAGTGCACCTACTTCAAGTTCTACAAAGAAAACACAGCT ACAACTGGAGCATTTACTGCTGGATTACAGATGATTTGAATGGAAT TAATAATTACAAGAATCCAAACTCACCAGGATGCTCACATTTAAGTT TTACATGCCCAAGAAGGCCACAGAACTGAAACATCTTCAGTGTCTAG AAGAAGAACTCAAACCTCTGGAGGAAGTGCTAAATTTAGCTCAAAGC AAAACTTTCACTTAAGACCCAGGGACTTAATCAGCAATATCAACGTA ATAGTTCTGGAATAAAGGGATCTGAAACAACATTCATGTGTGAATA TGCTGATGAGACAGCAACCATTGTAGAATTTCTGAACAGATGGATTA CCTTTTGTCAAAGCATCATCTCAACACTGACTggtggaggatctggtgga ggtggatcagggtgggtggatccggtggagggtggtatct GAGCTCTGTGACGATGACCCGCCAGAGATCCCACACGCCACATTCAA AGCCATGGCCTACAAGGAAGGAACCATGTTGAACTGTGAATGCAAG AGAGGTTTCCGCAGAATAAAAAGCGGGTCACTCTATATGCTCTGTAC AGGAACTCTAGCCACTCGTCTGGGACAACCAATGTCAATGCACAA GCTCTGCCACTCGGAACACAACGAAACAAGTGACACCTCAACCTGAA GAACAGAAAGAAAGGAAAACACAGAAATGCAAAGTCCAATGCAGC CAGTGGACCAAGCGAGCCTTCCAGGTCACTGCAGGGAACCTCCACCA TGGGAAAATGAAGCCACAGAGAGAAATTTATCATTTCTGTTGGGGC AGATGGTTTATTATCAGTGCGTCCAGGGATACAGGGCTCTACACAGA GGTCTGCTGAGAGCGTCTGCAAAATGACCCACGGGAAGACAAGGT GGACCCAGCCCCAGCTCATATGCACAGGTGAAATGGAGACCATCA GTTTCCAGGTGAAGAGAAGCCTCAGGCAAGCCCCGAAGGCCGCTCT GAGAGTGAGACTTCTGCCTCGTCACAACAACAGATTTCAAATACA GACAGAAATGGCTGCAACCATGGAGACGTCCATATTTACAACAGAGT ACCAGGGTGGACATCACCATCACCATCACTAATAA
33	NT	Human	Unprocessed form IL-2 (Gly3Ser)4- extracellular domain of IL-2 R α	ATGGACAGGATGCAACTCCTGTCTTGCAATTGCACTAAGTCTTGCACTT GTCACAAACAGTGCACCTACTTCAAGTTCTACAAAGAAAACACAGCT ACAACTGGAGCATTTACTGCTGGATTACAGATGATTTGAATGGAAT TAATAATTACAAGAATCCAAACTCACCAGGATGCTCACATTTAAGTT TTACATGCCCAAGAAGGCCACAGAACTGAAACATCTTCAGTGTCTAG AAGAAGAACTCAAACCTCTGGAGGAAGTGCTAAATTTAGCTCAAAGC AAAACTTTCACTTAAGACCCAGGGACTTAATCAGCAATATCAACGTA ATAGTTCTGGAATAAAGGGATCTGAAACAACATTCATGTGTGAATA TGCTGATGAGACAGCAACCATTGTAGAATTTCTGAACAGATGGATTA CCTTTTGTCAAAGCATCATCTCAACACTGACTggtggagggttctggtggagg tcagggtggagggttcgggtggagggttctGAGCTCTGTGACGATGACCCGCCAGA GATCCACACGCCACATTCAAAGCCATGGCCTACAAGGAAGGAACCA TGTTGAACTGTGAATGCAAGAGAGGTTTCCGCAGAATAAAAAGCGG GTCACTCTATATGCTCTGTACAGGAACTCTAGCCACTCGTCTGGGA CAACCAATGTCAATGCACAAGCTCTGCCACTCGGAACACAACGAAAC AAGTGACACCTCAACCTGAAGAACAGAAAGAAAGGAAAACACAGA AATGCAAAGTCCAATGCAGCCAGTGGACCAAGCGAGCCTTCAGGTC ACTGCAGGGAACCTCCACATGGGAAAATGAAGCCACAGAGAGAAT TTATCATTTCTGTTGGTGGGACAGATGGTTTATTATCAGTGCGTCCAGG GATACAGGGCTCTACACAGAGGTCTGCTGAGAGCGTCTGCAAAATG ACCCACGGGAAGACAAGGTGGACCCAGCCCCAGCTCATATGCACAG GTGAAATGGAGACCAGTCAGTTTCCAGGTGAAGAGAAGCCTCAGGC AAGCCCCGAAGGCCGCTCTGAGAGTGAGACTTCTGCCTCGTCACAA CAACAGATTTCAAATACAGACAGAAATGGCTGCAACCATGGAGACG TCCATATTTACAACAGAGTACCAGGGTGGACATCACCATCACCATCAC TAATAA

34	NT	Human	Unprocessed form of IL-2 (Gly3Ser)3-extracellular domain of IL-2 R α	ATGGACAGGATGCAACTCCTGTCTTGCATTGCACTAAGTCTTGCACTT GTCACAAACAGTGCACCTACTTCAAGTTCTACAAAGAAAACACAGCT ACAACTGGAGCATTACTGCTGGATTACAGATGATTTTGAATGGAAT TAATAATTACAAGAATCCCAAACCTACCAGGATGCTCACATTTAAGTT TTACATGCCCAAGAAGGCCACAGAAGTAAACATCTTCAGTGTCTAG AAGAAGAACTCAAACCTCTGGAGGAAGTCTAAATTTAGCTCAAAGC AAAACTTTCACTTAAGACCCAGGGACTTAATCAGCAATATCAACGTA ATAGTTCTGGAATAAGGGATCTGAAACAACATTCATGTGTGAATA TGCTGATGAGACAGCAACCATTGTAGAATTTCTGAACAGATGGATTA CCTTTTGTCAAAGCATCATCTCAACACTGACTgggtggaggttctgggtggaggt tcaggtggaggttcgGAGCTCTGTGACGATGACCCGCCAGAGATCCCACA CGCCACATTCAAAGCCATGGCCTACAAGGAAGGAACCATGTTGAACT GTGAATGCAAGAGAGGTTTCCGCAGAATAAAAAGCGGGTCACTCTAT ATGCTCTGTACAGGAACTCTAGCCACTCGTCTGGGACAACCAATG TCAATGCACAAGCTCTGCCACTCGGAACACAACGAAACAGTGACAC CTCAACCTGAAGAACAGAAAGAAAGGAAAACACAGAAATGCAAAG TCCAATGCAGCCAGTGGACCAAGCGAGCCTTCCAGGTCACTGCAGGG AACCTCCACCATGGGAAAATGAAGCCACAGAGAGAATTTATCATTTT GTGGTGGGGCAGATGGTTTATTATCAGTGCCTCCAGGGATACAGGG CTCTACACAGAGGTCTGCTGAGAGCGTCTGCAAAATGACCCACGGG AAGACAAGGTGGACCCAGCCCCAGCTCATATGCACAGGTGAAATGG AGACCAGTCAGTTTCCAGGTGAAGAGAAGCCTCAGGCAAGCCCCGA AGGCCGTCTGAGAGTGAGACTTCTGCCTCGTCACAACAACAGATT TTCAAATACAGACAGAAATGGCTGCAACCATGGAGACGTCCATATTT ACAACAGAGTACCAGGGTGGACATCACCATCACCATCACTAATAA (gcc)gccRccAUGG
35	NT		Kozak consensus	(gcc)gccRccAUGG
36	AA	Mouse	unprocessed form of IL-2 (Gly3Ser)3-extracellular domain of IL-2 R α	MYSMQLASCVTLTLLVNSAPTSSSTSSSTAEAQQQQQQQQQQQH LEQLLMDLQELLSRMENYRNLKLPRLMTFKFYLPKQATELKDLQCLEDEL GPLRHVLDLTQSKSFQLEDAENFISNIRVTVVKLGSDNTFECQFDDESA TVVDFLRRWIAFCQSIISTSPQGGGSGGGSGGGSELCLYDPPEVPNATFK ALSYKNGTILNCECKRGFRRLKELVYMRLGNSWSSNCQCTSNSHDKSR KQVTAQLEHQKEQQTTTDMQKPTQSMHQENLTGHCREPPPWKHEDS KRIYHFVEGQSVHYECIPGYKALQRGPAISIC27KMKCGKTGWTPQLTC VDEREHHRFLASEESQGSRNSSPESETSCPIITTTDFPQPTETTAMTETFVL TMEYK
37	AA	Mouse	Mature form of IL-2 (Gly3Ser)3-extracellular domain of IL-2 R α	MDSMQLASCVTLTLLVNSAPTSSSTSSSTAEAQQQQQQQQQQQH LEQLLMDLQELLSRMENYRNLKLPRLMTFKFYLPKQATELKDLQCLEDEL GPLRHVLDLTQSKSFQLEDAENFISNIRVTVVKLGSDNTFECQFDDESA TVVDFLRRWIAFCQSIISTSPQGGGSGGGSGGGSELCLYDPPEVPNATFK ALSYKNGTILNCECKRGFRRLKELVYMRLGNSWSSNCQCTSNSHDKSR KQVTAQLEHQKEQQTTTDMQKPTQSMHQENLTGHCREPPPWKHEDS KRIYHFVEGQSVHYECIPGYKALQRGPAISICKMKCGKTGWTPQLTCV DEREHRFLASEESQGSRNSSPESETSCPIITTTDFPQPTETTAMTETFVL TMEYK
38	AA	Human	unprocessed form of IL-2 (Gly4Ser)5-extracellular domain of IL-2 R α	MDRMQLLSCIALSLALVTNSAPTSSSTKKTLQLEHLLDLQMILNGINN YKNPKLRLMTFKFYMPKKATELKHLCLEELKPLEEVLNLAQSKNFHL RPRDLISINIVILEKGSETTFMCEYADETATVEFLNRWITFCQSIISTLT GGGGSGGGSGGGSGGGSGGGSGGGSELDDDPPEIPHATFKAMAYK EGTMLNCECKRGFRRIKSGSLYMLCTGNSSHSSWDNQCCQCTSSATRNT TKQVTPQPEEQKERKTTEMQSPMQPVDQASLPGHCREPPPWENEATE RIYHFVVGQMVVYQCVQGYRALHRGPAESVCKMTHGKTRWTPQLIC TGEMETSQFPGEKPQASPEGRPESETSCLVTTTDFQIQTEMAATMETS IFTTEYQ
39	AA	human	Mature form of human IL-2 (Gly4Ser)5-extracellular domain of IL-2 R α	APTSSSTKKTLQLEHLLDLQMILNGINNYPKLPRLMTFKFYMPKKA TELKHLQCLEELKPLEEVLNLAQSKNFHLRPRDLISINIVILEKGSETTF MCEYADETATVEFLNRWITFCQSIISTLTGGGGSGGGSGGGSGGGG SGGGSELDDDPPEIPHATFKAMAYKEGTMNLNCECKRGFRRIKSGSLY MLCTGNSSHSSWDNQCCQCTSSATRNTTKQVTPQPEEQKERKTTEMQS PMQPVQDQASLPGHCREPPPWENEATERIYHFVVGQMVVYQCVQGYR ALHRGPAESVCKMTHGKTRWTPQLICTGEMETSQFPGEKPQASPEG RPESETSCLVTTTDFQIQTEMAATMETSIFTTEYQ

				GPAESVCKMTHGKTRWTQPOLICTGEMETSQFPGEKPQASPEGRPES ETSCLVTITDFQIQTEMAATMETSIFTTEYQ
47	NT	Human	Unprocessed form IL-2 (Gly3Ser)2- extracellular domain of IL-2 R α	ATGGACAGGATGCAACTCCTGTCTTGCAATTGCACTAAGTCTTGCACTT GTCACAAACAGTGACCTACTTCAAGTTCTACAAAGAAAACACAGCT ACAACTGGAGCATTTACTGCTGGATTACAGATGATTTTGAATGGAAT TAATAATTACAAGAATCCCAAACCTACCAGGATGCTCACATTTAAGTT TTACATGCCCAAGAAGGCCACAGAAGTGAACATCTTCAGTGTCTAG AAGAAGAACTCAAACCTCTGGAGGAAGTGCTAAATTTAGCTCAAAGC AAAAACTTTCACTTAAGACCCAGGGACTTAATCAGCAATATCAACGTA ATAGTTCTGGAAGTAAAGGGATCTGAAACAACATTCATGTGTGAATA TGCTGATGAGACAGCAACCATTGTAGAATTTCTGAACAGATGGATTA CCTTTTGTCAAAGCATCATCTCAACACTGACTggtggagggttctggtggagg tcaGAGCTCTGTGACGATGACCCGCCAGAGATCCCACACGCCACATTC AAAGCCATGGCCTACAAGGAAGGAACCATGTTGAAGTGTGAATGCA AGAGAGGTTTCCGCAGAATAAAAAGCGGGTCACTCTATATGCTCTGT ACAGGAACTCTAGCCACTCGTCTGGGACAACCAATGTCAATGCAC AAGCTCTGCCACTCGGAACACAACGAAACAAGTGACACCTCAACCTG AAGAACAGAAAGAAAGGAAAACACAGAAATGCAAGTCCAATGCA GCCAGTGGACCAAGCGAGCCTTCCAGGTCACTGCAGGGAACCTCCAC CATGGGAAAATGAAGCCACAGAGAGAATTTATCATTTCGTGGTG GCAGATGGTTTATTATCAGTGCGTCCAGGGATACAGGGCTCTACACA GAGGTCTGCTGAGAGCGTCTGCAAAATGACCCACGGGAAGACAAG GTGGACCCAGCCCCAGCTCATATGCACAGGTGAAATGGAGACCAGTC AGTTTCCAGGTGAAGAGAAGCCTCAGGCAAGCCCCGAAGGCCGTCC TGAGAGTGAGACTTCCTGCCTCGTCACAACAACAGATTTTCAAATACA GACAGAAATGGCTGCAACCATGGAGACGTCCATATTTACAACAGAGT ACCAGGGTGGACATCACCATCACCATCACTAATAA
48	NT	Human	Unprocessed form IL-2 (Gly3)- extracellular domain of IL-2 R α	ATGGACAGGATGCAACTCCTGTCTTGCAATTGCACTAAGTCTTGCACTT GTCACAAACAGTGACCTACTTCAAGTTCTACAAAGAAAACACAGCT ACAACTGGAGCATTTACTGCTGGATTACAGATGATTTTGAATGGAAT TAATAATTACAAGAATCCCAAACCTACCAGGATGCTCACATTTAAGTT TTACATGCCCAAGAAGGCCACAGAAGTGAACATCTTCAGTGTCTAG AAGAAGAACTCAAACCTCTGGAGGAAGTGCTAAATTTAGCTCAAAGC AAAAACTTTCACTTAAGACCCAGGGACTTAATCAGCAATATCAACGTA ATAGTTCTGGAAGTAAAGGGATCTGAAACAACATTCATGTGTGAATA TGCTGATGAGACAGCAACCATTGTAGAATTTCTGAACAGATGGATTA CCTTTTGTCAAAGCATCATCTCAACACTGACTggtggagggtGAGCTCTGT GACGATGACCCGCCAGAGATCCCACACGCCACATTCAAAGCCATGGC CTACAAGGAAGGAACCATGTTGAAGTGTGAATGCAAGAGAGGTTTCC GCAGAATAAAAAGCGGGTCACTCTATATGCTCTGTACAGGAACTCT AGCCACTCGTCTGGGACAACCAATGTCAATGCACAAGCTCTGCCAC TCGGAACACAACGAAACAAGTGACACCTCAACCTGAAGAACAGAAA GAAAGGAAAACACAGAAATGCAAGTCCAATGCAGCCAGTGGACC AAGCGAGCCTTCCAGGTCACTGCAGGGAACCTCCACATGGGAAAAT GAAGCCACAGAGAGAATTTATCATTTCGTGGTGAGGATGGTTTA TTATCAGTGCGTCCAGGGATACAGGGCTCTACACAGAGGTCTGCTG AGAGCGTCTGCAAAATGACCCACGGGAAGACAAGGTGGACCCAGCC CCAGCTCATATGCACAGGTGAAATGGAGACCAGTCAGTTTCCAGGTG AAGAGAAGCCTCAGGCAAGCCCCGAAGGCCGTCTGAGAGTGAGAC TTCCTGCCTCGTCACAACAACAGATTTTCAAATACAGACAGAAATGGC TGCAACCATGGAGACGTCCATATTTACAACAGAGTACCAGGGTGGAC ATCACCATCACCATCACTAATAA
49	NT	Human	Unprocessed form IL-2 (Gly4Ser)5- extracellular domain of IL-2 R α	ATGGACAGGATGCAACTCCTGTCTTGCAATTGCACTAAGTCTTGCACTT GTCACAAACAGTGACCTACTTCAAGTTCTACAAAGAAAACACAGCT ACAACTGGAGCATTTACTGCTGGATTACAGATGATTTTGAATGGAAT TAATAATTACAAGAATCCCAAACCTACCAGGATGCTCACATTTAAGTT TTACATGCCCAAGAAGGCCACAGAAGTGAACATCTTCAGTGTCTAG AAGAAGAACTCAAACCTCTGGAGGAAGTGCTAAATTTAGCTCAAAGC AAAAACTTTCACTTAAGACCCAGGGACTTAATCAGCAATATCAACGTA ATAGTTCTGGAAGTAAAGGGATCTGAAACAACATTCATGTGTGAATA TGCTGATGAGACAGCAACCATTGTAGAATTTCTGAACAGATGGATTA CCTTTTGTCAAAGCATCATCTCAACACTGACTggtggagggtgagtcagggtga ggtgagtcggtggagggtgagtcagggtggagggtgagtcagggtgagtcgagct

				CTGTGACGATGACCCGCCAGAGATCCCACACGCCACATTCAAAGCCA TGGCCTACAAGGAAGGAACCATGTTGAAGTGTGAATGCAAGAGAGG TTTCCGCAGAATAAAAAGCGGGTCACTCTATATGCTCTGTACAGGAA ACTCTAGCCACTCGTCTGGGACAACCAATGTCAATGCACAAGCTCTG CCACTCGGAACACAACGAAACAAGTGACACCTCAACCTGAAGAACAG AAAGAAAGGAAAACACAGAAATGCAAAAGTCCAATGCAGCCAGTGG ACCAAGCGAGCCTTCCAGGTCACTGCAGGGAACCTCCACCATGGGAA AATGAAGCCACAGAGAGAATTTATCATTTCTGGTGGGGCAGATGGT TTATTATCAGTGCCTCCAGGGATACAGGGCTCTACACAGAGGTCTGT CTGAGAGCGTCTGCAAAATGACCCACGGGAAGACAAGGTGGACCCA GCCCCAGCTCATATGCACAGGTGAAATGGAGACCAAGTTCAG GTGAAGAGAAGCCTCAGGCAAGCCCCGAAGGCCGTCTGAGAGTGA GACTTCTGCTCGTCACAACAACAGATTTTCAAATACAGACAGAAAT GGCTGCAACCATGGAGACGTCCATTTTACAACAGAGTACCAGGGTG GACATCACCATCACCATCAATAA
50	AA		(G1y4Ser)3 linker	GGGGSGGGSGGGGS
51	AA		(G1y4Ser)2 linker	GGGGSGGGGS
52	AA		(G1y4Ser)1 linker	GGGGS
53	NT		Kozak sequence	gccaccATGG
54	AA	Mouse	Unprocessed form of IL-2 (G1y4Ser)4- extracellular domain of IL-2 R α + glycine spacer and poly-histidine region	MDSMQLASCVTLTLVLLVNSAPTSSSTSSSTAEAQQQQQQQQQQH LEQLLMDLQELLSRMENYRNKLPRMLTFKFYLPKQATELKDLCLEDEL GPLRHVLDLTQSKSFQLEDAENFISNIRVTVVKLKGSNTFECQFDDESA TVVDFLRRWIAFCQSIISTSPQGGGSGGGSGGGSGGGSGGGSELCLYDP PEVPNATFKALSYKNGTILNCECKRGFRRLKELVYMRLGNSWSSNCQC TSNSHDKSRKQVTAQLEHQKEQQTDDMQKPTQSMHQENLTGHCRE PPPWKHEDSKRIYHFVEGQSVHYECIPGYKALQRGPAISICKMKCGKTG WTQPQLTCVDEREHRFLASEESQGSRNSSPESETSCPIITDDFPQPTET TAMTETFVLTMEYKGGHHHHHH
55	AA	Mouse	Unprocessed form of IL-2 (G1y4Ser)5- extracellular domain of IL-2 R α + glycine spacer and poly-histidine region	MDSMQLASCVTLTLVLLVNSAPTSSSTSSSTAEAQQQQQQQQQQH LEQLLMDLQELLSRMENYRNKLPRMLTFKFYLPKQATELKDLCLEDEL GPLRHVLDLTQSKSFQLEDAENFISNIRVTVVKLICGSNTFECQFDDESA TVVDFLRRWIAFCQSIISTSPQGGGSGGGSGGGSGGGSGGGSGGGSE LCLYDPPEVPNATFKALSYKNGTILNCECKRGFRRLKELVYMRLGNSWS SNCQCTSNHDKSRKQVTAQLEHQKEQQTDDMQKPTQSMHQENLT GHCREPPPWKHEDSKRIYHFVEGQSVHYECIPGYKALQRGPAISICKMKC GKTGWTQPQLTCVDEREHRFLASEESQGSRNSSPESETSCPIITDDFPQ PTETTAMTETFVLTMEYKGGHHHHHH
56	AA	Mouse	Unprocessed form of IL-2 (Gly3Ser)4- extracellular domain of IL-2 R α + glycine spacer and poly-histidine region	MDSMQLASCVTLTLVLLVNSAPTSSSTSSSTAEAQQQQQQQQQQH LEQLLMDLQELLSRMENYRNKLPRMLTFKFYLPKQATELKDLCLEDEL GPLRHVLDLTQSKSFQLEDAENFISNIRVTVVKLKGSNTFECQFDDESA TVVDFLRRWIAFCQSIISTSPQGGGSGGGSGGGSGGGSELCLYDPPEVP NATFKALSYKNGTILNCECKRGFRRLKELVYMRLGNSWSSNCQCTSNS HDKSRKQVTAQLEHQKEQQTDDMQKPTQSMHQENLTGHCREPPPW KHEDSKRIYHFVEGQSVHYECIPGYKALQRGPAISICKMKCGKTGWTQP QLTCVDEREHRFLASEESQGSRNSSPESETSCPIITDDFPQPTETTAMTE TFVLTMEYKGGHHHHHH
57	AA	Mouse	Unprocessed form of IL-2 (G1y3Ser)3- extracellular domain of IL-2 R α + glycine spacer and poly-histidine region	MDSMQLASCVTLTLVLLVNSAPTSSSTSSSTAEAQQQQQQQQQQH LEQLLMDLQELLSRMENYRNKLPRMLTFKFYLPKQATELKDLCLEDEL GPLRHVLDLTQSKSFQLEDAENFISNIRVTVVKLKGSNTFECQFDDESA TVVDFLRRWIAFCQSIISTSPQGGGSGGGSGGGSELCLYDPPEVPNATFK ALSYKNGTILNCECKRGFRRLKELVYMRLGNSWSSNCQCTSNHDKSR KQVTAQLEHQKEQQTDDMQKPTQSMHQENLTGHCREPPPWKHEDS KRIYHFVEGQSVHYECIPGYKALQRGPAISICKMKCGKTGWTQPQLTCV DEREHRFLASEESQGSRNSSPESETSCPIITDDFPQPTETTAMTETFVLT MEYKGGHHHHHH

58	AA	Human	Unprocessed form IL-2 (Gly3Ser)2-extracellular domain of IL-2 R α + glycine spacer and poly-histidine region	MDRMQLLSICIALSLALVTNSAPTSSSTKKTKLQLEHLLDLQMILNGINN YKNPKLTRMLTFKFYMPKKATELKHLCLEELKPLEEVLNLAQSKNFHL RPRDLISININIVLELKGSETTFMCEYADETATIVEFLNRWITFCQSIISTLT GGGSGGGSELCDDDPPEIPHATFKAMAYKEGTMLNCECKRGFRRIKSGS LYMLCTGNSSHSSWDNQCQCTSSATRNNTTKQVTPQPEEQKERKTTEM QSPMQPVDQASLPGHCREPPPWENEATERIYHFVVGQMVYYQCVQG YRALHRGPAESVCKMTHGKTRWTQPQLICTGEMETSQFPGECKPQASP EGRPESETSLVTTTDFQIQTEMAATMETSIFTTEYQGGHHHHHH
59	AA	Human	Unprocessed form of IL-2 (Gly3Ser)3-extracellular domain of IL-2 R α + glycine spacer and poly-histidine region	MDRMQLLSICIALSLALVTNSAPTSSSTKKTKLQLEHLLDLQMILNGINN YKNPKLTRMLTFKFYMPKKATELKHLCLEELKPLEEVLNLAQSKNFHL RPRDLISININIVLELKGSETTFMCEYADETATIVEFLNRWITFCQSIISTLT GGGSGGGSGGGSELCDDDPPEIPHATFKAMAYKEGTMLNCECKRGFRRIKSGSLYMLCTGNSSHSSWDNQCQCTSSATRNNTTKQVTPQPEEQKERKT TEMQSPMQPVDQASLPGHCREPPPWENEATERIYHFVVGQMVYYQC VQGYRALHRGPAESVCKMTHGKTRWTQPQLICTGEMETSQFPGECKP QASPEGRPESETSLVTTTDFQIQTEMAATMETSIFTTEYQGGHHHHHH
60	AA	Human	Unprocessed form IL-2 (Gly3Ser)4-extracellular domain of IL-2 R α + glycine spacer and poly-histidine region	MDRMQLLSICIALSLALVTNSAPTSSSTKKTKLQLEHLLDLQMILNGINN YKNPKLTRMLTFKFYMPKKATELKHLCLEELKPLEEVLNLAQSKNFHL RPRDLISININIVLELKGSETTFMCEYADETATIVEFLNRWITFCQSIISTLT GGGSGGGSGGGSGGGSELCDDDPPEIPHATFKAMAYKEGTMLNCECK RGFRRIKSGSLYMLCTGNSSHSSWDNQCQCTSSATRNNTTKQVTPQPEE QKERKTTEMQSPMQPVDQASLPGHCREPPPWENEATERIYHFVVGQ MVYYQCVQGYRALHRGPAESVCKMTHGKTRWTQPQLICTGEMETSQF PGECKPQASPEGRPESETSLVTTTDFQIQTEMAATMETSIFTTEYQGGH HHHHH
61	AA	Human	Unprocessed form IL-2 (Gly4Ser)4-extracellular domain of IL-2 R α + glycine spacer and poly-histidine region	MDRMQLLSICIALSLALVTNSAPTSSSTKKTKLQLEHLLDLQMILNGINN YKNPKLTRMLTFKFYMPKKATELKHLCLEELKPLEEVLNLAQSKNFHL RPRDLISININIVLELKGSETTFMCEYADETATIVEFLNRWITFCQSIISTLT GGGSGGGSGGGSGGGSGGGSELCDDDPPEIPHATFKAMAYKEGTML NCECKRGFRRIKSGSLYMLCTGNSSHSSWDNQCQCTSSATRNNTTKQVT PQPEEQKERKTTEMQSPMQPVDQASLPGHCREPPPWENEATERIYHFV VGQMVYYQCVQGYRALHRGPAESVCKMTHGKTRWTQPQLICTGEME TSQFPGECKPQASPEGRPESETSLVTTTDFQIQTEMAATMETSIFTTEY QGGHHHHHHH
62	AA	Human	Mature form IL-2 (Gly3Ser)3-extracellular domain of mutIL-2 R α	APTSSSTKKTKLQLEHLLDLQMILNGINNYKNPKLTRMLTFKFYMPKKA TELKHLCLEELKPLEEVLNLAQSKNFHLRPRDLISININIVLELKGSETTF MCEYADETATIVEFLNRWITFCQSIISTLTGGGSGGGSGGGSELCDDDPP EIPHATFKAMAYKEGTMLNCECKRGFTSIKSGSLYMLCTGNSSHSSWDN QCQCTSSATRNNTTKQVTPQPEEQKERKTTEMQSPMQPVDQASLPGHC REPPPWENEATERIYHFVVGQMVYYQCVQGYRALHRGPAESVCKMTH GKTRWTQPQLICTGEMETSQFPGECKPQASPEGRPESETSLVTTTDFQI QTEMAATMETSIFTTEYQ
63	NT	Human	Unprocessed form IL-2 (Gly3Ser)3-extracellular domain of mutIL-2 R α Mut	ATGGACAGGATGCAACTCCTGTCTTGCACTGCACTAAGTCTTGCACTT GTCACAAACAGTGACCTACTTCAAGTTCTACAAAGAAAACACAGCT ACAACTGGAGCATTACTGCTGGATTACAGATGATTTGAATGGAAT TAATAATTACAAGAATCCCAAACCTCACCAGGATGCTCACATTTAAGTT TTACATGCCCAAGAAGGCCACAGAACTGAAACATCTTCAGTGTCTAG AAGAAGAACTCAAACCTCTGGAGGAAGTCTAAATTTAGCTCAAAGC AAAACTTTCACTTAAGACCCAGGGACTTAATCAGCAATATCAACGTA ATAGTTCTGGAACATAAAGGGATCTGAAACAACATTCTATGTGTGAATA TGCTGATGAGACAGCAACCATTGTAGAATTTCTGAACAGATGGATTA CCTTTTGTCAAAGCATCATCTCAACACTGACTggagggttctggagggt tcagggtggagggtcgGAGCTCTGTGACGATGACCCGCCAGAGATCCCACA CGCCACATTCAAAGCCATGGCCTACAAGGAAGGAACCATGTTGAAGT GTGAATGCAAGAGAGGTTTCACTCAATAAAAAGCGGGTCACTCTAT ATGCTCTGTACAGGAACTCTAGCCACTCGTCTGGGACAACCAATG TCAATGCACAAGCTCTGCCACTCGGAACACAAACGAAACAAGTGACAC CTCAACCTGAAGAACAGAAAGAAAGGAAAACACAGAAATGCAAG TCCAATGCAGCCAGTGGACCAAGCGAGCCTTCCAGGTCACTGCAGGG

				AACCTCCACCATGGGAAAAATGAAGCCACAGAGAGAATTTATCATTTCTGTGGTGGGGCAGATGGTTTATTATCAGTGCGTCCAGGGATACAGGGCTCTACACAGAGGTCTGCTGAGAGCGTCTGCAAAATGACCCACGGGAAGACAAGGTGGACCCAGCCCCAGCTCATATGCACAGGTGAAATGGAGACCAGTCAGTTTCCAGGTGAAGAGAAAGCCTCAGGCAAGCCCCGAGAGGCCGTCTGAGAGTGAGACTTCTGCCTCGTCACAACAACAGATTCAAATACAGACAGAAATGGCTGCAACCATGGAGACGTCCATATTTACAACAGAGTACCAGGGTGGACATCACCATCACCATCACTAATAA
64	AA	Human	Mature form IL-2 (Gly4Ser)4-extracellular domain of mutIL-2 R α	APTSSSTKKTQLQLEHLLDLQMILNGINNYKNPKLTRMLTFKFMPPKKA TELKHLQCLEEEELKPLEEVNLQAQSKNFHLRPDLISNINVIVLELKGSETTF MCEYADETATIVEFLNRWITFCQSIISTLTGGGGSGGGSGGGSGGGG SELCDDDPPEIPHATFKAMAYKEGTMNLNCEKRGFTSIKSGSLYMLCTG NSSHSSWDNQCQCTSSATRNNTTKQVTPQPEEQKERKTTEMQSPMQPV DQASLPGHCREPPPWENEATERIYHFVVGQMVVYQCVQGYRALHRGP AESVCKMTHGKTRWTQPQLICTGEMETSQFPGEKPKQASPEGRPESET SCLVTTTDFQIQTEMAATMETSIFTTEYQ
65	NT	Human	Unprocessed form IL-2 (Gly4Ser)4-extracellular domain of mutIL-2 R α	ATGGACAGGATGCAACTCCTGTCTTGCATTGCACTAAGTCTTGCACTT GTCACAAACAGTGACCTACTTCAAGTTCTACAAAGAAAACACAGCT ACAACTGGAGCATTTACTGCTGGATTACAGATGATTTTGAATGGAAT TAATAATTACAAGAATCCAAACTCACCAGGATGCTCACATTTAAGTT TTACATGCCCAAGAAGGCCACAGAAGTCAACATCTTCAGTCTAG AAGAAGAAGTCAAACCTCTGGAGGAAGTCTAAATTTAGCTCAAAGC AAAAACTTCACTTAAGACCCAGGGACTTAATCAGCAATATCAACGTA ATAGTTCTGGAAGTAAAGGGATCTGAAACAACATTCATGTGTGAATA TGCTGATGAGACAGCAACCATTGTAGAATTTCTGAACAGATGGATTA CCTTTTGTCAAAGCATCATCTCAACACTGACTggtggaggatcggtgga ggtggatcaggaggaggatccggaggaggatct GAGCTCTGTGACGATGACCCGCCAGAGATCCACACGCCACATTCAA AGCCATGGCCTACAAGGAAGGAACCATGTTGAAGTGTGAATGCAAG AGAGGTTTCACTCAATAAAAAGCGGGTCACTCTATATGCTCTGTACA GGAACTCTAGCCACTCGTCTGGGACAACCAATGTCAATGCACAAG CTCTGCCACTCGGAACACAACGAAACAAGTGACACCTCAACCTGAAG AACAGAAAAGAAAGGAAAACACAGAAATGCAAAGTCCAATGCAGCC AGTGGAACCAAGCGAGCCTTCCAGGTCACTGCAGGGAACTCCACCAT GGGAAAATGAAGCCACAGAGAGAATTTATCATTTCTGTTGGTGGGCA GATGGTTTATTATCAGTGCGTCCAGGGATACAGGGCTCTACACAGAG GTCCTGCTGAGAGCGTCTGCAAAATGACCCACGGGAAGACAAGGTG GACCCAGCCCCAGCTCATATGCACAGGTGAAATGGAGACCAGTCAGT TTCCAGGTGAAGAGAAGCCTCAGGCAAGCCCCGAAGCCGCTCTGA GAGTGAGACTTCTGCCTCGTCACAACAACAGATTTTCAAATACAGAC AGAAATGGCTGCAACCATGGAGACGTCCATATTACAACAGAGTACC AGGGTGGACATCACCATCACCATCACTAATAA

All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

Claims

1. A fusion protein comprising:
 - (a) a first polypeptide comprising Interleukin-2 (IL-2); and
 - (b) a second polypeptide, fused in frame to the first polypeptide by a linker, wherein the linker is the sequence set forth in SEQ ID NO: 13, and wherein the second polypeptide comprises an extracellular domain of Interleukin-2 Receptor alpha (IL-2R α), which is capable of binding to IL-2;
wherein the fusion protein has IL-2 activity.
2. The fusion protein of claim 1, wherein the fusion protein has an increased IL-2 activity when compared to native IL-2.
3. The fusion protein of any one of claims 1 or 2, wherein the fusion protein has an increased persistent IL-2 stimulation of IL-2R bearing lymphocytes *in vivo* when compared to native IL-2.
4. The fusion protein of any one of claims 1-3, wherein the first polypeptide comprising IL-2 has at least 75% sequence identity to SEQ ID NO: 2.
5. The fusion protein of any one of claims 1-4, wherein the second polypeptide comprising the extracellular domain of IL-2R α has at least 75% sequence identity to SEQ ID NO: 7.

6. The fusion protein of any one of claims 1-5, wherein the first polypeptide comprising IL-2 has at least 75%, 80%, 85%, 90%, 95%, or 99% sequence identity to SEQ ID NO: 2.

7. The fusion protein of any one of claims 1-6, wherein the second polypeptide comprising the extracellular domain of IL-2R α has at least 75%, 80%, 85%, 90%, 95%, or 99% sequence identity to SEQ ID NO: 7.

8. The fusion protein of any one of claims 1-7, comprising

- (a) the amino acid sequence of any one of SEQ ID NO: 26, 27, 36, 37, 57, or 59; or,
- (b) a sequence having at least 75%, 80%, 85%, 90%, or 95% sequence identity to any one of SEQ ID NO: 26, 27, 36, 37, 57, or 59.

9. The fusion protein of any one of claims 1-7, comprising the amino acid sequence of any one of SEQ ID NO: 26, 27, 36, 37, 57, or 59.

10. The fusion protein of any one of claims 1-7, comprising the amino acid sequence of SEQ ID NO: 26.

11. The fusion protein of any one of claims 1-10, wherein the fusion protein comprises one or more substitutions, deletions, truncations, or insertions in the extracellular domain of IL-2R α , as compared to the polypeptide comprising the extracellular domain of IL-2R α , as set forth in SEQ ID NO: 7.

12. The fusion protein of claim 9, wherein the fusion protein comprises:
 - (a) the amino acid sequence of SEQ ID NO: 62; or
 - (b) a sequence having at least 75%, 80%, 85%, 90%, or 95% sequence identity to SEQ ID NO: 62.
13. The fusion protein of claim 9, comprising the amino acid sequence of SEQ ID NO: 62.
14. A polynucleotide comprising a nucleotide sequence encoding the fusion protein of any one of claims 1-13.
15. A host cell comprising the polynucleotide of claim 14.
16. The host cell of claim 15, wherein the host cell comprises a CHO cell or a COS cell.
17. A method for producing a fusion protein of any one of claims 1-13, the method comprising (i) introducing into a host cell a polynucleotide encoding the fusion protein of any one of claims 1-13 and (ii) expressing the fusion protein in the host cell.
18. The method of claim 17, wherein the polynucleotide encoding the fusion protein is operably linked to a promoter active in the host cell.

19. Use of the fusion protein of any one of claims 1 and 4-13 in the manufacture of a medicament for the treatment of an autoimmune disease or inflammatory disorder in a subject in need thereof, wherein:

- (a) the fusion protein has an increased IL-2 activity when compared to native IL-2, or
- (b) the fusion protein has an increased persistent IL-2 stimulation of IL-2R bearing lymphocytes *in vivo* when compared to native IL-2.

20. The use of claim 19, wherein the subject has an autoimmune disease.

21. The use of claim 20, wherein the autoimmune disease comprises type 1 diabetes, multiple sclerosis, rheumatoid arthritis, celiac disease, systemic lupus erythematosus, juvenile idiopathic arthritis, Crohn's disease, ulcerative colitis or systemic sclerosis, graft versus host disease, psoriasis, alopecia areata, or HCV-induced vasculitis.

22. The use of any one of claims 19-21, wherein a therapeutically effective amount of the fusion protein comprises 10^3 to 10^6 IU of IL-2 activity per subject or $10^4 \pm 100$ fold of IL-2 activity per subject.

23. Use of the fusion protein of any one of claims 1 and 4-13 for the manufacture of a medicament in enhancing the immunogenicity of an anti-cancer vaccine or a vaccine that is poorly immunogenic in a subject in need thereof, wherein:

- (a) the fusion protein has an increased IL-2 activity when compared to native IL-2, or

(b) the fusion protein has an increased persistent IL-2 stimulation of IL-2R bearing lymphocytes *in vivo* when compared to native IL-2.

24. Use of the fusion protein of any one of claims 1 and 4-13 for the manufacture of a medicament in overcoming a suppressed immune response to an anti-cancer vaccine or a vaccine that is poorly immunogenic in a subject in need thereof, wherein:

(a) the fusion protein has an increased IL-2 activity when compared to native IL-2, or

(b) the fusion protein has an increased persistent IL-2 stimulation of IL-2R bearing lymphocytes *in vivo* when compared to native IL-2.

25. The use of claim 23 or 24, wherein the fusion protein and the vaccine are to be administered sequentially, in any order.

26. The use of claim 23 or 24, wherein the fusion protein and the vaccine are to be administered simultaneously.

27. The use of any one of claims 23-26, wherein the vaccine is a cancer vaccine.

28. The use of any one of claims 23-27, wherein a therapeutically effective amount of the fusion protein comprises at least 10^4 to 10^7 IU of IL-2 activity per subject or at least 10^5+100 of IL-2 activity per subject.

29. The use of any one of claims 19-28, wherein the subject is a human.