

DESCRIPTION

METHOD OF PRODUCING SUBSTANCE

5 Technical Field

[0001]

The present invention relates to a method of producing a substance by using cells.

10 Background Art

[0002]

Cell culturing and substance production

Cells generally exist as three-dimensional aggregates in the body. However, when cells are cultured in an artificial environment, it is common to use the classical plate culture method in which the cells are cultured two-dimensionally in a manner plated as a monolayer on the bottom of the culturing vessel, or a suspension culture method in which cells are cultured while dispersed in a liquid culture solution. Cells most suited for the plate culture method are cells having relatively high adhesion, but even when such suitable cells are used, differences in the culturing environment can often result in significant changes in the properties of the cells. With suspension culture methods as well, certain cells are suitable while others are not.

[0003]

With increasing demand for *in vivo* proteins to be used for medical purposes, such as vaccines, enzymes, hormones, antibodies, cytokines and the like, interest is becoming increasingly focused on mass production of such *in vivo* proteins by cell culturing. For suspended cells of *E. coli* and the like, research is being conducted on techniques for mass culturing in large-scale culturing tanks. Mass culturing of suspended cells using large-scale culturing tanks requires large volumes of culture solution and an agitating apparatus. Increasing focus is

also being directed toward research in which substances are produced using adherent cells, as research on such cells continues to progress. In the case of adherent cells, the cells will only expand two-dimensionally when the classical plate culture method is employed, and therefore a large culturing area is necessary. In order to perform mass production of *in vivo* proteins, etc., a lot of researches are also being conducted on cell culture carriers and bioreactors for three-dimensional and mass cell culture.

[0004]

Microcarriers, which are microparticles on which cells can adhere and grow, are being widely studied as typical cell culturing supports (PTL 3). Different types of microcarriers have been studied and developed, and many are available on the market. They are often used for production of vaccines and proteins, and widely employed as methodologies in upscalable systems. In microcarrier culturing, however, the microcarriers must be adequately stirred and diffused to avoid their aggregation, and this places a limit on the cell culture volume. Moreover, for production of a substance, for example, the procedure is methodologically complex as well, since fine particles must be separated with a fractionating filter or the like in order to separate the carrier itself. In addition, since the form of the particulate matter of a microcarrier used in microcarrier culturing is limited, it is impossible to avoid the properties that arise from its form.

[0005]

Alternative methods to microcarrier culturing have been discovered, such as methods of continuous mass culturing of spheroid cells by three-dimensional culturing using methyl cellulose or gellan gum. It is indeed possible to achieve mass culturing of cells in a bioreactor using a cellulose sponge. Being a large

closed system, however, there are many restrictions on its operation, such as the inability to easily contact the culturing environment.

[0006]

5 It has been a goal to establish a novel system that is suited for convenience and automation and that allows efficient production of numerous substances in large quantities in cells.

[0007]

10 Porous polyimide film

 The term "polyimide" is a general term for polymers including imide bonds in the repeating unit. An "aromatic polyimide" is a polymer in which aromatic compounds are directly linked by imide bonds. An aromatic polyimide has an aromatic-aromatic conjugated structure via an imide bond, and therefore has a strong rigid molecular structure, and since the imide bonds provide powerful intermolecular force, it has very high levels of thermal, mechanical and chemical properties.

20 **[0008]**

 Porous polyimide films have been utilized in the prior art for filters and low permittivity films, and especially for battery-related purposes, such as fuel cell electrolyte membranes and the like. PTLs 6 to 8 describe porous polyimide films with numerous macrovoids, having excellent permeability for gases and the like, high porosity, excellent smoothness on both surfaces, relatively high strength and, despite high porosity, also excellent resistance against compression stress in the film thickness direction. All of these are porous polyimide films formed via amic acid.

Citation List

[Patent Literature]

35 **[0009]**

[PTL 1] Japanese Examined Patent Publication No. HEI 6-30570

[PTL 2] Japanese Patent No. 2683946

[PTL 3] WO2003/054174

[PTL 4] Japanese Unexamined Patent Publication No. 2009-261410

5 [PTL 5] WO2000/068371

[PTL 6] WO2010/038873

[PTL 7] Japanese Unexamined Patent Publication No. 2011-219585

10 [PTL 8] Japanese Unexamined Patent Publication No. 2011-219586

[Non-patent literature]

[0010]

15 [NPL 1] Hogwood et al., Current Opinion in Biotechnology 2014, 30:153-160

[NPL 2] Kurokawa et al., Journal of Bioscience and Bioengineering VOL. 111 No. 5, 600-604, 2011

DISCLOSURE OF THE INVENTION

20 Problems to be Solved by the Invention

[0011]

25 It is an object of the present invention to provide a method for producing a substance using cells, a substance-producing apparatus and kit, and use of the same.

Means for Solving the Problems

[0012]

30 As a result of diligent research conducted with the aim of providing a method and system that allow convenient and efficient production of substances in cells, the present inventors have completed this invention upon finding that by conducting culturing using a porous polyimide film, it is possible to culture large quantities of cells in a limited space without any
35 special pretreatment, taking advantage of high-density culture characteristics to accomplish efficient substance

production. Thus, one feature of the invention is culturing of cells using a porous polyimide film.

[0013]

Cells grow stably and independently by utilizing the large diameter communicating pores in the porous polyimide film, through which the cells are able to pass, so that even when large quantities of cells are present in comparison to plate culturing, it is possible to ensure contact of the communicating pores with the medium and thereby allow stable growth to be continued. The same structural features also contribute in the production of substances such as proteins and viruses, allowing the produced substances to be efficiently supplied to the medium. Moreover, due to the thin-film properties, flexible properties, form stability and free shapeability of porous polyimide films, it is possible to insert large amounts of sheets in small unit spaces and to layer multiple films, while the high heat resistance and solvent resistance of porous polyimide films allow convenient and rapid sterilization of the sheets to be accomplished by various means. In addition, the three-dimensional scaffold structure of a porous polyimide film offers the possibility of increasing substance productivity per cell, compared to flat culturing.

[0014]

The present invention preferably includes, but is not limited to, the following modes.

[Mode 1]

A method of producing a substance using cells, the method including applying the cells to a porous polyimide film and culturing the cells to produce the substance by the cells.

[Mode 2]

The method according to mode 1, wherein the substance is selected from the group consisting of proteins, glycoproteins and viruses.

[Mode 3]

The method according to mode 1 or 2, wherein the cells are cultured under stationary culture conditions.

[Mode 4]

5 The method according to mode 1 or 2, wherein the cells are cultured under rotating culture or stirred conditions.

[Mode 5]

The method according to mode 1 or 2, wherein the cells are cultured in a continuous manner.

10 [Mode 6]

The method according to mode 5, including:

setting a cell culturing apparatus in an incubator and culturing the cells, wherein

the cell culturing apparatus includes:

15 a culturing unit that houses one or more porous polyimide films to support cells, and that comprises a culture medium supply port and a culture medium discharge port, and

20 a culture medium-supply unit comprising a culture medium housing vessel, a culture medium supply line, and a liquid conveyance pump that conveys the medium through the culture medium supply line, the first end of the culture medium supply line contacting the medium in the culture medium housing vessel, and the second end of the culture medium supply line being in communication with
25 the culturing unit interior via the culture medium supply port of the culturing unit.

[Mode 7]

30 The method according to mode 6, wherein the cell culturing apparatus does not have an air supply port, an air discharge port and an oxygen permeation film.

[Mode 8]

35 The method according to any one of modes 1 to 7, wherein the porous polyimide film is a porous polyimide film including a polyimide obtained from a tetracarboxylic dianhydride and a diamine.

[Mode 9]

The method according to mode 8, wherein the porous polyimide film is a colored porous polyimide film obtained by forming a polyamic acid solution composition including a polyamic acid solution obtained from a tetracarboxylic dianhydride and a diamine, and a coloring precursor, and then heat treating it at 250°C or higher.
[Mode 10]

The method according to mode 8 or 9, wherein the porous polyimide film is a porous polyimide film with a multilayer structure, having two different surface layers and a macro-void layer.
[Mode 11]

The method according to mode 10, wherein the film thickness of the porous polyimide film is no greater than 75 μm .
[Mode 12]

The method according to any one of modes 1 to 11, wherein the porous polyimide film is:
i) folded,
ii) wound into a roll,
iii) connected as sheets or fragments by a filamentous structure, or
iv) bound into a rope,
and suspended or anchored in the cell culture medium in the cell culturing vessel.
[Mode 13]

The method according to any one of modes 1 to 12, using two or more porous polyimide films layered either above and below or left and right in the cell culture medium.
[Mode 14]

The method according to any one of modes 1 to 13, wherein the cells are transformed by genetic engineering technology so as to express a substance.
[Mode 15]

The method according to any one of modes 1 to 14, wherein the cells are selected from the group consisting

of animal cells, insect cells, plant cells, yeast cells
and bacteria.

[Mode 16]

5 The method according to mode 15, wherein the animal
cells are cells derived from an animal belonging to the
subphylum Vertebrata.

[Mode 17]

10 The method according to mode 15 or 16, wherein the
animal cells are selected from the group consisting of
Chinese hamster ovary tissue-derived cells (CHO cells),
African green monkey kidney-derived established cell
lines (Vero cells), human hepatic cancer-derived cells
(HepG2 cells), canine kidney epithelial cell-derived cell
lines (MDCK cells) and human hepatic cancer tissue-
15 derived established cell lines (huGK-14).

[Mode 18]

20 The method according to mode 15, wherein the
bacteria are selected from the group consisting of lactic
acid bacteria, *E. coli*, *Bacillus subtilis* and
cyanobacteria.

[Mode 19]

25 A substance-producing apparatus for use in the
method according to any one of modes 1 to 18, including a
porous polyimide film.

[Mode 20]

The substance-producing apparatus according to mode
19, wherein two or more porous polyimide films are
layered either above and below or left and right.

[Mode 21]

30 A kit for use in the method according to any one of
modes 1 to 18, including a porous polyimide film.

[Mode 22]

35 Use of a porous polyimide film in the method
according to any one of modes 1 to 18.

Effect of the Invention

[0015]

The method of the invention, when used to carry out culturing using a porous polyimide film with an apparatus and kit, allows a large volume of cells to be efficiently cultured by placing one or more sheets together in a limited space in various forms. With the method of the invention it has become possible to produce substances by efficient and convenient cell culturing.

Brief Description of the Drawings

[0016]

Fig. 1 shows changes in cell counts during gas phase subculturing after long term culturing of human skin fibroblasts.

Mode for Carrying Out the Invention

[0017]

I. Method of producing substance

The present invention relates to a method of producing a substance using cells. The method of the invention includes applying cells to a porous polyimide film and culturing the cells to produce a substance by the cells.

[0018]

1. Cells

There are no particular restrictions on the type of cells that can be utilized for the method of the invention, and it may be used for growth of any type of cells.

[0019]

For example, the cells may be selected from the group consisting of animal cells, insect cells, plant cells, yeast cells and bacteria. Animal cells are largely divided into cells from animals belonging to the subphylum Vertebrata, and cells from non-vertebrates (animals other than animals belonging to the subphylum Vertebrata). There are no particular restrictions on the source of the animal cells, for the purpose of the

present specification. Preferably, they are cells from an animal belonging to the subphylum Vertebrata. The subphylum Vertebrata includes the superclass Agnatha and the superclass Gnathostomata, the superclass

5 Gnathostomata including the class Mammalia, the class Aves, the class Amphibia and the class Reptilia. Preferably, they are cells from an animal belonging to the class Mammalia, generally known as mammals. Mammals are not particularly restricted but include, preferably,
10 mice, rats, humans, monkeys, pigs, dogs, sheep and goats.

[0020]

While not a restriction, the animal cells used are preferably cells selected from the group consisting of Chinese hamster ovary-derived cells (CHO cells), African
15 green monkey kidney-derived established cell lines (Vero cells), human hepatic cancer-derived cells (HepG2 cells), canine kidney epithelial cell-derived cell lines (MDCK cells), human hepatic cancer tissue-derived established cell lines (huGK-14), normal human fibroblast-like
20 synovial membrane cells (HFLS cells) and chronic rheumatoid arthritis patient-derived fibroblast-like synovial membrane cells (HFLS-RA cells).

[0021]

There are also no particular restrictions on sources
25 of plant cells, for the purpose of the present specification. Suitable cells are from plants including bryophytes, pteridophytes and spermatophytes.

Plants from which spermatophyte cells are derived include both monocotyledons and dicotyledons. While not
30 restrictive, monocotyledons include Orchidaceae plants, Poaceae plants (rice, corn, barley, wheat, sorghum and the like) and Cyperaceae plants. Dicotyledons include plants belonging to many subclasses including the subclass Chrysanthemum, the subclass Magnoliidae and the
35 subclass Rosidae.

[0022]

Algae may be considered cell-derived organisms.

These include different groups, from the eubacteria
Cyanobacteria (blue-green algae), to eukaryotic
monocellular organisms (diatoms, yellow-green algae,
dinoflagellates and the like) and multicellular marine
5 algae (red algae, brown algae and green algae).

[0023]

There are no particular limitations on the types of
archaebacteria or bacteria for the purpose of the present
specification. Archaebacteria are composed of groups
10 comprising methanogenic bacteria, extreme halophilic
bacteria, thermophilic acidophilic bacteria,
hyperthermophilic bacteria and the like. Bacteria are
selected from the group consisting of, for example,
lactic acid bacteria, *E. coli*, *Bacillus subtilis* and
15 cyanobacteria.

[0024]

The types of animal cells or plant cells that may be
used for the method of the invention are not particularly
restricted, but are preferably selected from the group
20 consisting of pluripotent stem cells, tissue stem cells,
somatic cells and germ cells.

[0025]

The term "pluripotent stem cells", for the purpose
of the invention, is intended as a comprehensive term for
25 stem cells having the ability to differentiate into cells
of a variety of tissues (pluripotent differentiating
power). While not restrictive, pluripotent stem cells
include embryonic stem cells (ES cells), induced
pluripotent stem cells (iPS cells), embryonic germ cells
30 (EG cells) and germ stem cells (GS cells). They are
preferably ES cells or iPS cells. Particularly preferred
are iPS cells, which are free of ethical problems, for
example. The pluripotent stem cells used may be any
publicly known ones, and for example, the pluripotent
35 stem cells described in International Patent Publication
No. WO2009/123349 (PCT/JP2009/057041) may be used.

[0026]

The term "tissue stem cells" refers to stem cells that are cell lines capable of differentiation but only to limited specific tissues, though having the ability to differentiate into a variety of cell types (pluripotent
5 differentiating power). For example, hematopoietic stem cells in the bone marrow are the source of blood cells, while neural stem cells differentiate into neurons. Additional types include hepatic stem cells from which the liver is formed and skin stem cells that form skin
10 tissue. Preferably, the tissue stem cells are selected from among mesenchymal stem cells, hepatic stem cells, pancreatic stem cells, neural stem cells, skin stem cells and hematopoietic stem cells.

[0027]

15 The term "somatic cells" refers to cells other than germ cells, among the cells composing a multicellular organism. In sexual reproduction these are not passed on to the next generation. Preferably, the somatic cells are selected from among hepatocytes, pancreatic cells,
20 muscle cells, bone cells, osteoblasts, osteoclasts, chondrocytes, adipocytes, skin cells, fibroblasts, pancreatic cells, renal cells and lung cells, or blood cells such as lymphocytes, erythrocytes, leukocytes, monocytes, macrophages or megakaryocytes.

25 **[0028]**

The term "germ cells" refers to cells having the role of passing on genetic information to the succeeding generation in reproduction. These include, for example, gametes for sexual reproduction, i.e. the ova, egg cells,
30 sperm, sperm cells, and spores for asexual reproduction.

[0029]

The cells may also be selected from the group consisting of sarcoma cells, established cell lines and transformants. The term "sarcoma" refers to cancer
35 occurring in non-epithelial cell-derived connective tissue cells, such as the bone, cartilage, fat, muscle or blood, and includes soft tissue sarcomas, malignant bone

tumors and the like. Sarcoma cells are cells derived from sarcoma. The term "established cell line" refers to cultured cells that are maintained *in vitro* for long periods and reach a stabilized character and can be semi-permanently subcultured. Cell lines derived from various tissues of various species including humans exist, such as PC12 cells (from rat adrenal medulla), CHO cells (from Chinese hamster ovary), HEK293 cells (from human embryonic kidney), HL-60 cells from (human leukocytes) and HeLa cells (from human cervical cancer), Vero cells (from African green monkey kidney epithelial cells), MDCK cells (from canine renal tubular epithelial cells) and HepG2 cells (from human hepatic cancer). The term "transformants" refers to cells with an altered genetic nature by extracellularly introduced nucleic acid (DNA and the like).

[0030]

The cells are not particularly restricted so long as they are cells that can exhibit a desired substance. The cells may naturally express the substance, or they may be transformed by genetic engineering technology so as to produce the substance. The cells are preferably cells that have been transformed by genetic engineering technology so as to produce the substance. Suitable methods are known for transformation of animal cells, plant cells and bacteria. (For example, MOLECULAR CLONING: A Laboratory Manual (Fourth Edition), Michael R Green and Joseph Sambrook, 2012, (Cold Spring Harbor Laboratory Press), Mutation Research 760 (2014) 36-45, Reviews in Mutation Research).

[0031]

2. Substance

The type of substance that can be produced by cells using the method of the invention is not particularly restricted, so long as the substance can be produced in the cells naturally or by genetic engineering technology. Preferably, the substance is one selected from the group

consisting of proteins (including polypeptides),
glycoproteins and viruses.

[0032]

5 Examples of proteins include physiologically active
proteins such as erythropoietin, insulin and albumin,
cytokines such as tumor necrosis factor α , interleukin-6
(IL-6), interleukin-8 (IL-8), granulocyte colony
stimulating factor (G-CSF) and interferon, enzymes such
10 as thrombin and trypsin, and monoclonal antibodies
containing antibody drugs. Physiologically active
proteins and monoclonal antibodies are preferred.

[0033]

15 Examples of glycoproteins include collagen,
fibronectin and hyaluronic acid. Fibronectin is
preferred.

[0034]

Examples of viruses include influenza virus and
adenovirus. Influenza virus is preferred.

[0035]

20 Examples of substances naturally expressed in
specific cells according to the invention include
antibody drugs expressed by transformants, and human
collagen or fibronectin expressed by human skin
fibroblasts.

25 **[0036]**

3. Porous polyimide film

Polyimide is a general term for polymers containing
imide bonds in the repeating unit, and usually it refers
to an aromatic polyimide in which aromatic compounds are
30 directly linked by imide bonds. An aromatic polyimide
has an aromatic-aromatic conjugated structure via an
imide bond, and therefore has a strong rigid molecular
structure, and since imide bonds have powerful
intermolecular force, it has very high levels of thermal,
35 mechanical and chemical properties.

[0037]

The porous polyimide film used for the invention is

preferably a porous polyimide film including (as the main component) a polyimide obtained from a tetracarboxylic dianhydride and a diamine, and more preferably it is a porous polyimide film comprising a polyimide obtained from a tetracarboxylic dianhydride and a diamine. The phrase "including as the main component" means that it essentially contains no components other than the polyimide obtained from a tetracarboxylic dianhydride and a diamine, as constituent components of the porous polyimide film, or that it may contain them but they are additional components that do not affect the properties of the polyimide obtained from the tetracarboxylic dianhydride and diamine.

[0038]

The porous polyimide film used for the invention also includes colored porous polyimide films obtained by forming a polyamic acid solution composition containing a polyamic acid solution obtained from a tetracarboxylic acid component and a diamine component, and a coloring precursor, and then heat treating it at 250°C or higher.

[0039]

Polyamic acid

A polyamic acid is obtained by polymerization of a tetracarboxylic acid component and a diamine component. A polyamic acid is a polyimide precursor that can be cyclized to a polyimide by thermal imidization or chemical imidization.

[0040]

The polyamic acid used may be any one that does not have an effect on the invention, even if a portion of the amic acid is imidized. Specifically, the polyamic acid may be partially thermally imidized or chemically imidized.

[0041]

When the polyamic acid is to be thermally imidized, there may be added to the polyamic acid solution, if necessary, an imidization catalyst, an organic

phosphorus-containing compound, or fine particles such as inorganic fine particles or organic fine particles.

Also, when the polyamic acid is to be chemically imidized, there may be added to the polyamic acid solution, if necessary, a chemical imidization agent, a dehydrating agent, or fine particles such as inorganic fine particles or organic fine particles. Even if such components are added to the polyamic acid solution, they are preferably added under conditions that do not cause precipitation of the coloring precursor.

[0042]

Coloring precursor

A coloring precursor to be used for the invention is a precursor that generates a colored substance by partial or total carbonization under heat treatment at 250°C or higher.

[0043]

Coloring precursors to be used for the invention are preferably uniformly dissolved or dispersed in a polyamic acid solution or polyimide solution and subjected to thermal decomposition by heat treatment at 250°C or higher, preferably 260°C or higher, even more preferably 280°C or higher and more preferably 300°C or higher, and preferably heat treatment in the presence of oxygen such as air, at 250°C, preferably 260°C or higher, even more preferably 280°C or higher and more preferably 300°C or higher, for carbonization to produce a colored substance, more preferably producing a black colored substance, with carbon-based coloring precursors being most preferred.

[0044]

The coloring precursor, when being heated, first appears as a carbonized compound, but compositionally it contains other elements in addition to carbon, and also includes layered structures, aromatic crosslinked structures and tetrahedron carbon-containing disordered structures.

[0045]

Carbon-based coloring precursors are not particularly restricted, and for example, they include tar or pitch such as petroleum tar, petroleum pitch, coal
5 tar and coal pitch, coke, polymers obtained from acrylonitrile-containing monomers, ferrocene compounds (ferrocene and ferrocene derivatives), and the like. Of these, polymers obtained from acrylonitrile-containing monomers and/or ferrocene compounds are preferred, with
10 polyacrylnitrile being preferred as a polymer obtained from an acrylonitrile-containing monomer.

[0046]

The tetracarboxylic dianhydride used may be any tetracarboxylic dianhydride, selected as appropriate
15 according to the properties desired. Specific examples of tetracarboxylic dianhydrides include biphenyltetracarboxylic dianhydrides such as pyromellitic dianhydride, 3,3',4,4'-biphenyltetracarboxylic dianhydride (s-BPDA) and 2,3,3',4'-
20 biphenyltetracarboxylic dianhydride (a-BPDA), oxydiphthalic dianhydride, diphenylsulfone-3,4,3',4'-tetracarboxylic dianhydride, bis(3,4-dicarboxyphenyl)sulfide dianhydride, 2,2-bis(3,4-dicarboxyphenyl)-1,1,1,3,3,3-hexafluoropropane
25 dianhydride, 2,3,3',4'-benzophenonetetracarboxylic dianhydride, 3,3',4,4'-benzophenonetetracarboxylic dianhydride, bis(3,4-dicarboxyphenyl)methane dianhydride, 2,2-bis(3,4-dicarboxyphenyl)propane dianhydride, p-phenylenebis(trimellitic acid monoester acid anhydride),
30 p-biphenylenebis(trimellitic acid monoester acid anhydride), m-terphenyl-3,4,3',4'-tetracarboxylic dianhydride, p-terphenyl-3,4,3',4'-tetracarboxylic dianhydride, 1,3-bis(3,4-dicarboxyphenoxy)benzene dianhydride, 1,4-bis(3,4-dicarboxyphenoxy)benzene
35 dianhydride, 1,4-bis(3,4-dicarboxyphenoxy)biphenyl dianhydride, 2,2-bis[(3,4-dicarboxyphenoxy)phenyl]propane dianhydride, 2,3,6,7-naphthalenetetracarboxylic

dianhydride, 1,4,5,8-naphthalenetetracarboxylic dianhydride, 4,4'-(2,2-hexafluoroisopropylidene)diphthalic dianhydride, and the like. Also preferably used is an aromatic
5 tetracarboxylic acid such as 2,3,3',4'-diphenylsulfonetetracarboxylic acid. These may be used alone or in appropriate combinations of two or more.

[0047]

Particularly preferred among these are at least one
10 type of aromatic tetracarboxylic dianhydride selected from the group consisting of biphenyltetracarboxylic dianhydride and pyromellitic dianhydride. As a biphenyltetracarboxylic dianhydride there may be suitably used 3,3',4,4'-biphenyltetracarboxylic dianhydride.

[0048]

Any desired diamine may be used as a diamine. Specific examples of diamines include the following.

1) Benzenediamines with one benzene nucleus, such as 1,4-diaminobenzene (paraphenylenediamine), 1,3-
20 diaminobenzene, 2,4-diaminotoluene and 2,6-diaminotoluene;

2) diamines with two benzene nuclei, including diaminodiphenyl ethers such as 4,4'-diaminodiphenyl ether and 3,4'-diaminodiphenyl ether, and 4,4'-
25 diaminodiphenylmethane, 3,3'-dimethyl-4,4'-diaminobiphenyl, 2,2'-dimethyl-4,4'-diaminobiphenyl, 2,2'-bis(trifluoromethyl)-4,4'-diaminobiphenyl, 3,3'-dimethyl-4,4'-diaminodiphenylmethane, 3,3'-dicarboxy-4,4'-diaminodiphenylmethane, 3,3',5,5'-tetramethyl-4,4'-
30 diaminodiphenylmethane, bis(4-aminophenyl)sulfide, 4,4'-diaminobenzanilide, 3,3'-dichlorobenzidine, 3,3'-dimethylbenzidine, 2,2'-dimethylbenzidine, 3,3'-dimethoxybenzidine, 2,2'-dimethoxybenzidine, 3,3'-diaminodiphenyl ether, 3,4'-diaminodiphenyl ether, 4,4'-
35 diaminodiphenyl ether, 3,3'-diaminodiphenyl sulfide, 3,4'-diaminodiphenyl sulfide, 4,4'-diaminodiphenyl sulfide, 3,3'-diaminodiphenylsulfone, 3,4'-

diaminodiphenylsulfone, 4,4'-diaminodiphenylsulfone,
3,3'-diaminobenzophenone, 3,3'-diamino-4,4'-
dichlorobenzophenone, 3,3'-diamino-4,4'-
dimethoxybenzophenone, 3,3'-diaminodiphenylmethane, 3,4'-
5 diaminodiphenylmethane, 4,4'-diaminodiphenylmethane, 2,2-
bis(3-aminophenyl)propane, 2,2-bis(4-aminophenyl)propane,
2,2-bis(3-aminophenyl)-1,1,1,3,3,3-hexafluoropropane,
2,2-bis(4-aminophenyl)-1,1,1,3,3,3-hexafluoropropane,
3,3'-diaminodiphenyl sulfoxide, 3,4'-diaminodiphenyl
10 sulfoxide and 4,4'-diaminodiphenyl sulfoxide;

3) diamines with three benzene nuclei, including
1,3-bis(3-aminophenyl)benzene, 1,3-bis(4-
aminophenyl)benzene, 1,4-bis(3-aminophenyl)benzene, 1,4-
bis(4-aminophenyl)benzene, 1,3-bis(4-
15 aminophenoxy)benzene, 1,4-bis(3-aminophenoxy)benzene,
1,4-bis(4-aminophenoxy)benzene, 1,3-bis(3-aminophenoxy)-
4-trifluoromethylbenzene, 3,3'-diamino-4-(4-
phenyl)phenoxybenzophenone, 3,3'-diamino-4,4'-di(4-
phenylphenoxy)benzophenone, 1,3-bis(3-aminophenyl
20 sulfide)benzene, 1,3-bis(4-aminophenyl sulfide)benzene,
1,4-bis(4-aminophenyl sulfide)benzene, 1,3-bis(3-
aminophenylsulfone)benzene, 1,3-bis(4-
aminophenylsulfone)benzene, 1,4-bis(4-
aminophenylsulfone)benzene, 1,3-bis[2-(4-
25 aminophenyl)isopropyl]benzene, 1,4-bis[2-(3-
aminophenyl)isopropyl]benzene and 1,4-bis[2-(4-
aminophenyl)isopropyl]benzene;

4) diamines with four benzene nuclei, including
3,3'-bis(3-aminophenoxy)biphenyl, 3,3'-bis(4-
30 aminophenoxy)biphenyl, 4,4'-bis(3-aminophenoxy)biphenyl,
4,4'-bis(4-aminophenoxy)biphenyl, bis[3-(3-
aminophenoxy)phenyl]ether, bis[3-(4-
aminophenoxy)phenyl]ether, bis[4-(3-
aminophenoxy)phenyl]ether, bis[4-(4-
35 aminophenoxy)phenyl]ether, bis[3-(3-
aminophenoxy)phenyl]ketone, bis[3-(4-
aminophenoxy)phenyl]ketone, bis[4-(3-

aminophenoxy)phenyl]ketone, bis[4-(4-aminophenoxy)phenyl]ketone, bis[3-(3-aminophenoxy)phenyl]sulfide, bis[3-(4-aminophenoxy)phenyl] sulfide, bis[4-(3-aminophenoxy)phenyl] sulfide, bis[4-(4-aminophenoxy)phenyl] sulfide, bis[3-(3-aminophenoxy)phenyl]sulfone, bis[3-(4-aminophenoxy)phenyl]sulfone, bis[4-(3-aminophenoxy)phenyl]sulfone, bis[4-(4-aminophenoxy)phenyl]sulfone, bis[3-(3-aminophenoxy)phenyl]methane, bis[3-(4-aminophenoxy)phenyl]methane, bis[4-(3-aminophenoxy)phenyl]methane, bis[4-(4-aminophenoxy)phenyl]methane, 2,2-bis[3-(3-aminophenoxy)phenyl]propane, 2,2-bis[3-(4-aminophenoxy)phenyl]propane, 2,2-bis[4-(3-aminophenoxy)phenyl]propane, 2,2-bis[4-(4-aminophenoxy)phenyl]propane, 2,2-bis[3-(3-aminophenoxy)phenyl]-1,1,1,3,3,3-hexafluoropropane, 2,2-bis[3-(4-aminophenoxy)phenyl]-1,1,1,3,3,3-hexafluoropropane, 2,2-bis[4-(3-aminophenoxy)phenyl]-1,1,1,3,3,3-hexafluoropropane and 2,2-bis[4-(4-aminophenoxy)phenyl]-1,1,1,3,3,3-hexafluoropropane.

[0049]

These may be used alone or in mixtures of two or more. The diamine used may be appropriately selected according to the properties desired.

[0050]

Preferred among these are aromatic diamine compounds, with 3,3'-diaminodiphenyl ether, 3,4'-diaminodiphenyl ether, 4,4'-diaminodiphenyl ether, paraphenylenediamine, 1,3-bis(3-aminophenyl)benzene, 1,3-bis(4-aminophenyl)benzene, 1,4-bis(3-aminophenyl)benzene, 1,4-bis(4-aminophenyl)benzene, 1,3-bis(4-aminophenoxy)benzene and 1,4-bis(3-aminophenoxy)benzene being preferred for use. Particularly preferred is at least one type of diamine selected from the group consisting of benzenediamines, diaminodiphenyl ethers and

bis(aminophenoxy)phenyl.

[0051]

From the viewpoint of heat resistance and dimensional stability under high temperature, the porous polyimide film to be used for the invention is preferably formed from a polyimide obtained by combination of a tetracarboxylic dianhydride and a diamine, having a glass transition temperature of 240°C or higher, or without a distinct transition point at 300°C or higher.

[0052]

From the viewpoint of heat resistance and dimensional stability under high temperature, the porous polyimide film to be used for the invention is preferably a porous polyimide film comprising one of the following aromatic polyimides.

(i) An aromatic polyimide comprising at least one tetracarboxylic acid unit selected from the group consisting of biphenyltetracarboxylic acid units and pyromellitic acid units, and an aromatic diamine unit,

(ii) an aromatic polyimide comprising a tetracarboxylic acid unit and at least one type of aromatic diamine unit selected from the group consisting of benzenediamine units, diaminodiphenyl ether units and bis(aminophenoxy)phenyl units, and/or,

(iii) an aromatic polyimide comprising at least one type of tetracarboxylic acid unit selected from the group consisting of biphenyltetracarboxylic acid units and pyromellitic acid units, and at least one type of aromatic diamine unit selected from the group consisting of benzenediamine units, diaminodiphenyl ether units and bis(aminophenoxy)phenyl units.

[0053]

While not restrictive, the porous polyimide film for use in the invention may be a porous polyimide film with a multilayer structure, having at least two surface layers (A-surface and B-surface), and a macro-void layer

sandwiched between the two surface layers. Preferably, the porous polyimide film is a porous polyimide film wherein the macro-void layer has a partition bonded to the surface layers (A-surface and B-surface) and a plurality of macro-voids with mean pore sizes of 10 to 500 μm in the planar direction of the film, surrounded by the partition and the surface layers (A-surface and B-surface), wherein the macro-void layer partition and the surface layers (A-surface and B-surface) each have thicknesses of 0.01 to 20 μm , with a plurality of pores with mean pore sizes of 0.01 to 100 μm , the pores being optionally communicating with each other, and also having a partial or total multilayer structure in communication with the macro-voids, where the total film thickness is 5 to 500 μm and the porosity is 40% or greater and less than 95%.

[0054]

The total film thickness of the porous polyimide film used for the invention is not limited, but may be 20 to 75 μm according to one mode. Differences in the film thickness may be observed as differences in cell growth rate, cell morphology, cell saturation within the plate, and the like.

[0055]

According to the invention, when the porous polyimide film used has two different surface layers (A-surface and B-surface), and a macro-void layer sandwiched between the two surface layers, the mean pore size of the holes in the A-surface may differ from the mean pore size of the holes in the B-surface. Preferably, the mean pore size of the holes in the A-surface is smaller than the mean pore size of the holes in the B-surface. More preferably, the mean pore size of the holes in the A-surface is smaller than the mean pore size of the holes in the B-surface, with the mean pore size of the holes in the A-surface being 0.01 to 50 μm , 0.01 μm to 40 μm , 0.01

μm to 30 μm, 0.01 μm to 20 μm or 0.01 μm to 15 μm, and the mean pore size of the holes in the B-surface being 20 μm to 100 μm, 30 μm to 100 μm, 40 μm to 100 μm, 50 μm to 100 μm or 60 μm to 100 μm. Most preferably, the A-surface is a mesh structure having small holes with a mean pore size of no greater than 15 μm, such as 0.01 μm to 15 μm, and the B-surface is a large-hole structure with a mean pore size of 20 μm or greater, such as 20 μm to 100 μm. According to the invention, when the porous polyimide film is used in which the mean pore size of the holes in the A-surface is smaller than the mean pore size of the holes in the B-surface, cells may be seeded on the A-surface, or on the B-surface. Preferably, cells are seeded on the A-surface.

[0056]

The total film thickness of the porous polyimide film used for the invention can be measured using a contact thickness gauge.

The mean pore size of the surface of the porous polyimide film can be determined by measuring the pore area of 200 or more open holes from a scanning electron micrograph of the porous film surface, and calculating the mean diameter from the average value for the pore areas according to the following formula (1), assuming the pore shapes to be circular.

$$2 \times \sqrt{(S a / \pi)} \quad (1)$$

Mean pore size =

(wherein Sa represents the average value for the pore areas)

The porosity of the porous polyimide film used for the invention can be determined by measuring the film thickness and mass of the porous film cut out to a prescribed size, and performing calculation from the

basis weight according to the following formula (2).

$$\text{Porosity (\%)} = \frac{(1 - w / (S \times d \times D)) \times 100}{(2)}$$

5 (wherein S represents the area of the porous film, d represents the total film thickness, w represents the measured mass, and D represents the polyimide density, the polyimide density being defined as 1.34 g/cm³.)

[0057]

10 For example, the porous polyimide films described in International Patent Publication No. WO2010/038873, Japanese Unexamined Patent Publication No. 2011-219585 and Japanese Unexamined Patent Publication No. 2011-219586 may also be used in the invention.

15

[0058]

Naturally, the porous polyimide film to which cells are applied in the method of the invention is preferred to be in a state including no biological components other than those of the loaded cells, i.e. in a sterilized state. The method of the invention preferably includes a step of pre-sterilizing the porous polyimide film. A porous polyimide film has very excellent heat resistance and is lightweight, allows free selection of the shape and size, and is easy to treat for sterilization. Any desired sterilization treatment may be conducted, such as dry heat sterilization, steam sterilization, sterilization with a microbicide such as ethanol, or electromagnetic wave sterilization using UV or gamma rays.

30

[0059]

The cells that have been seeded on the surface of the porous polyimide film can stably grow and proliferate on the surface and/or in the interior of the film. The cells may be in a variety of different forms, depending on the location of growth and proliferation in the film.

35

According to one mode of the invention, growth may be carried out while moving the surface and interior of the porous polyimide film and changing the form, depending on the type of cell.

5

[0060]

4. Application of cells to porous polyimide film

There are no particular restrictions on the specific steps for application of the cells to the porous
10 polyimide film. It is possible to carry out the steps described throughout the present specification, or to employ any desired method suited for applying cells to a film-like support. Application of cells to the porous polyimide film in the method of the invention includes,
15 but is not limited to, the following modes.

[0061]

(A) A mode including a step of seeding cells on the surface of a porous polyimide film;

(B) A mode including a step of:

20 placing a cell suspension on the dried surface of a porous polyimide film,

allowing it to stand, or moving the porous polyimide film to promote efflux of the liquid, or stimulating part of the surface to cause absorption of the cell suspension
25 into the film, and

retaining the cells in the cell suspension inside the film and allowing the water to flow out; and

(C) A mode including a step of:

30 wetting one or both sides of a porous polyimide film with a cell culture solution or a sterilized liquid,

loading a cell suspension into the wetted porous polyimide film, and

retaining the cells in the cell suspension inside the film and allowing the water to flow out.

35

[0062]

Mode (A) includes a step of directly seeding cells or a cell mass on the surface of a porous polyimide film.

Alternatively, it includes a mode of placing a porous polyimide film in a cell suspension and wetting the cell culture solution from the surface of the film.

[0063]

5 Cells seeded on the surface of a porous polyimide film adhere to the porous polyimide film and infiltrate into the interiors of the pores. Preferably, the cells adhere spontaneously to the porous polyimide film without applying any particular exterior physical or chemical
10 force. The cells that have been seeded on the surface of the porous polyimide film can stably grow and proliferate on the surface and/or in the interior of the film. The cells may be in a variety of different forms, depending on the location of the film used for growth and
15 proliferation.

[0064]

 For mode (B), a cell suspension is placed on the dried surface of a porous polyimide film. The porous polyimide film is allowed to stand, or the porous
20 polyimide film is moved to promote efflux of the liquid, or part of the surface is stimulated to cause absorption of the cell suspension into the film, so that the cell suspension permeates into the film. While it is not our intention to be constrained by theory, this is believed
25 to be due to the properties of each of the surface forms of the porous polyimide film. According to this mode, the cells are absorbed and seeded in the locations of the film where the cell suspension has been loaded.

[0065]

30 Alternatively, as according to mode (C), after all or a portion of one or both sides of the porous polyimide film has been wetted with the cell culture solution or sterilized liquid, the cell suspension may be loaded into the wetted porous polyimide film. This will
35 significantly increase the transit rate of the cell suspension.

[0066]

For example, a method of wetting a portion of the film edges, for the main purpose of preventing fly loss of the film, may be used (hereunder referred to as "single-point wetting method"). The single-point wetting method is nearly the same as the dry method (mode (B)) in which the film essentially is not wetted. However, it is possible that cell solution permeation through the film is more rapid at the small wetted portions. There may also be used a method in which all of one or both sides of the porous polyimide film that have been thoroughly wetted (hereunder this will also be referred to as "wet film") is loaded with a cell suspension (this will hereunder be referred to as "wet film method"). In this case, the entire porous polyimide film has a greatly increased transit rate for the cell suspension.

[0067]

According to modes (B) and (C), the cells in the cell suspension are retained in the film, while the water flows out. This allows treatment such as increasing the concentration of cells in the cell suspension and flowing out of unwanted non-cellular components together with the water.

[0068]

Mode (A) will also be referred to as "natural seeding", and modes (B) and (C) as "suction seeding".

[0069]

Preferably, but not restrictively, the viable cells are selectively retained in the porous polyimide film. Thus, according to a preferred mode of the invention, the viable cells are retained in the porous polyimide film, and the dead cells preferentially flow out together with the water.

[0070]

The sterilized liquid used for mode (C) is not particularly restricted, and may be a sterilized buffering solution or sterilized water. A buffering solution may be, for example, (+) or (-) Dulbecco's PBS,

or (+) or (-) Hank's Balanced Salt Solution. Examples of buffering solutions are listed in Table 1 below.

[0071]

[Table 1]

Component	Concentration (mmol/L)	Concentration (g/L)
NaCl	137	8.00
KCl	2.7	0.20
Na ₂ HPO ₄	10	1.44
KH ₂ PO ₄	1.76	0.24
pH (-)	7.4	7.4

5

[0072]

In the method of the invention, application of cells to the porous polyimide film further includes a mode of adding adherent cells in a floating state as a suspension together with the porous polyimide film, to adhere the cells with the film (entangling). For example, for application of the cells to the porous polyimide film in the method of the invention, the cell culture medium, the cells and one or more of the porous polyimide films may be placed in the cell culturing vessel. When the cell culture medium is a liquid, the porous polyimide film is in a floating state in the cell culture medium. The cells can adhere to the porous polyimide film due to the properties of the porous polyimide film. Thus, even with cells that are not suited for natural suspension culture, the porous polyimide film allows culturing in a floating state in the cell culture medium. The cells preferably spontaneously adhere to the porous polyimide film. Here, "adhere spontaneously" means that the cells are retained on the surface or in the interior of the porous polyimide film without applying any particular exterior physical or chemical force.

[0073]

5. Cell culturing

The invention includes culturing of cells that have

30

been applied to a porous polyimide film and production of a substance by the cells.

[0074]

Application of cells to a porous polyimide film and
5 their culturing is described as follows in
PCT/JP2014/070407.

[0075]

Cell culturing can be classified into culturing
where the cultured cells are adhesion culture-type cells
10 or suspension culture-type cells, depending on the state
in the cell culture. Adhesion culture-type cells are
cultured cells that adhere and grow on a culturing
vessel, with the medium being exchanged at the time of
subculture. Suspension culture-type cells are cultured
15 cells that grow in a suspended state in a medium, and
generally the medium is not exchanged with each
subculture but dilution culture is carried out. Because
suspension culture allows culturing in a suspended state,
i.e. in a liquid, mass culturing becomes possible, and
20 because it is three-dimensional culturing, unlike with
adherent cells that grow only on the culturing vessel
surface, the advantage of increased culturable cell count
per unit space is afforded.

[0076]

25 In the method of the invention, when the porous
polyimide film is used in a state suspended in the cell
culture medium, two or more fragments of the porous
polyimide film may be used. Since the porous polyimide
film is a flexible thin-film, using such fragments that
30 are suspended in the culture solution, for example,
allows a porous polyimide film with a large surface area
to be added into a fixed volume of cell culture medium.
In the case of normal culturing, the container base area
constitutes the area limit in which cell culture can be
35 accomplished, but with cell culturing using the porous
polyimide film of the invention, all of the large surface
area of the previously added porous polyimide film

constitutes area in which cell culturing can be accomplished. The porous polyimide film allows the cell culture solution to pass through, allowing supply of nutrients, oxygen and the like even into the folded film, for example.

[0077]

The sizes and shapes of the porous polyimide film fragments are not particularly restricted. The shapes may be as desired, such as circular, elliptical, quadrilateral, triangular, polygonal or string-like.

[0078]

Because the porous polyimide film of the invention is flexible, it can be used with varying shapes. Instead of a flat form, the porous polyimide film can also be used by working into a three-dimensional shape. For example, porous polyimide films may be: i) folded, ii) wound into a roll, iii) connected as sheets or fragments by a filamentous structure, or iv) bound into a rope, for suspension or fixing in the cell culture medium in the cell culturing vessel. By forming into shapes such as i) to iv), it is possible to place a large amount of porous polyimide films into a fixed volume of cell culture medium, similar to using fragments. Furthermore, since each fragment can be treated as an aggregate, it is possible to aggregate and move the cell masses together, for overall high applicability.

[0079]

With the same concept as fragment aggregates, two or more porous polyimide films may be used in a layered form either above and below or left and right in the cell culture medium. Layering includes a mode in which portions of the porous polyimide films overlap. Layered culturing allows culturing of cells at high density in a narrow space. It is also possible to further layer a film on a film on which cells are already growing, setting it to create a multilayer of different cell types. The number of layered porous polyimide films is

not particularly restricted.

[0080]

Two or even more forms of the cell culturing method described above may be used in combination. For example, 5 using any of the methods of modes (A) to (C), first the cells may be applied to the porous polyimide film and then the cell-adhered porous polyimide film may be used for suspension culture. Alternatively, the step of application to the porous polyimide film may be a 10 combination of two or more of the methods of any of modes (A) to (C).

[0081]

In the method of the invention, preferably the cells grow and proliferate on the surface or in the interior of 15 the porous polyimide film. By the method of the invention, it is possible to carry out continuous growth of cells for 2 days or longer, more preferably 4 days or longer and even more preferably 6 days or longer.

[0082]

20 In the method of the invention, application of cells and culturing are carried out on a porous polyimide film, thereby allowing culturing of large volumes of cells to be accomplished since large numbers of cells grow on the multisided connected pore sections on the inside, and the 25 surfaces on the porous polyimide film.

[0083]

Moreover, in the method of the invention, it is possible to culture large volumes of cells efficiently while drastically reducing the amount of medium used for 30 cell culturing compared to the prior art. For example, large volumes of cells can be cultured even when all or a portion of the porous polyimide film is not in contact with the liquid phase of the cell culture medium. Also, the total volume of the cell culture medium in the cell 35 culturing vessel, with respect to the total porous polyimide film volume including the cell survival zone, can be significantly reduced below that of methods of the

prior art.

[0084]

Using the method of the invention, cells can be satisfactorily cultured for a long time even under
5 conditions in which the total volume of the cell culture medium in the cell culturing vessel is 10,000 times or less of the total sum of the porous polyimide film volume including the cell survival zone. Moreover, cells can be satisfactorily cultured for a long time even under
10 conditions in which the total volume of the cell culture medium in the cell culturing vessel is 1000 times or less of the total sum of the porous polyimide film volume including the cell survival zone. In addition, cells can be satisfactorily cultured for a long time even under
15 conditions in which the total volume of the cell culture medium in the cell culturing vessel is 100 times or less of the total sum of the porous polyimide film volume including the cell survival zone. Cells can also be satisfactorily cultured for a long time even under
20 conditions in which the total volume of the cell culture medium in the cell culturing vessel is 10 times or less of the total sum of the porous polyimide film volume including the cell survival zone.

25 **[0085]**

Conventionally known three-dimensional culturing base materials for cells include the porous polystyrene base material Alvetex[®](ReproCell), the polycaprolactone base material 3D Insert-PCL (3D Biotech), and the
30 polystyrene base material 3D Insert-PS (3D Biotech). When cells have been cultured using a porous polyimide film according to the method of the invention, the substance productivity per unit volume is much higher compared to when cells have been cultured using a
35 conventionally known three-dimensional culture base material.

[0086]

Moreover, in the method of the invention, the cells are cultured in the porous polyimide film without forming cell masses (spheroids). Since the cells under such culturing conditions can stably grow for prolonged periods without detaching, it is possible to obtain the produced substance as a clarified liquid.

[0087]

In addition, the porous polyimide film can be frozen and stored while supporting large volumes of cells. The porous polyimide film can also be thawed when desired, and used for production of protein, without passing through a preculturing step. The ability of the cells to produce the substance does not decrease even after the freezing and thawing steps. For use of thawed cells in the prior art, it has been necessary to preculture the thawed cells and then culture the surviving cells after supporting them on a desired base material or support. By using the porous polyimide film, however, it is possible to carry out culturing, freezing, storage, thawing and re-culturing of cells while they are adhering to the porous polyimide film, without a preculturing step after thawing as has been required in the prior art. Therefore, by preparing large amounts of a frozen cell-supporting porous polyimide films, large volumes of cells can be used at any desired time without an amplifying culturing step. Such steps of freezing, storage, thawing and culturing may be repeated several times.

[0088]

6. Cell culturing and substance production system and culturing conditions

In the method of the invention, the cell culturing and substance production system and culturing conditions may be set as appropriate according to the type of cells used. Culturing methods suited for various cells including animal cells, plant cells and bacteria are publicly known, and a person skilled in the art may carry out culturing of cells suited for the porous polyimide

film, using any publicly known method. The cell culture medium may also be prepared as appropriate for the type of cells.

[0089]

5 Cell culture methods and cell culture media for animal cells may be found in the Cell Culture Media Catalog of Lonza Group, Ltd., for example. Cell culture methods and cell culture media for plant cells may also be found in the Plant Tissue Culturing Media Series by
10 Wako Corp. Japan, for example. Cell culture methods and cell culture media for bacteria may also be found in the General Bacterial Media Catalog of BD Corp., for example. The cell culture medium to be used in the method of the invention may be in any form such as a liquid medium,
15 semi-solid medium or solid medium. Also, a liquid medium in mist form may be sprayed into the cell culturing vessel to contact the medium with the cell-supporting porous polyimide film.

[0090]

20 The cell culture using a porous polyimide film may also be combined with another suspension culture support such as a microcarrier, cellulose sponge or the like.

[0091]

The method of the invention is not particularly
25 restricted in terms of the form and scale of the system used for the culturing, and any scale from cell culturing dish to a flask, plastic bag, test tube or large tank may be used, as appropriate. These include, for example, Cell Culture Dish by BD Falcon, and Nunc Cell Factory by
30 Thermo Scientific. By using a porous polyimide film according to the invention, it has become possible to carry out culturing even of cells that have not been capable of natural suspension culture, using an apparatus intended for suspension culture, in a state similar to
35 suspension culturing. The apparatus for suspension culture that is used may be, for example, a spinner flask or rotating culturing flask by Corning, Inc. As an

environment allowing a similar function to be obtained, there may be used a hollow fiber culturing system such as the FiberCell[®] System by Veritas.

5 **[0092]**

According to the invention, the cells may also be cultured under stationary culture conditions. By intermittently replacing the medium, it is possible to isolate the useful substance that is produced.

10 Application of the porous polyimide film can also drastically increase efficiency in stationary culturing using a single use culturing bag.

[0093]

15 According to the invention, the cells may also be cultured under rotating culture or stirred conditions. Continuously swinging the single use culturing bag will allow its use on a very large scale. The same also applies for rotating culture or stirred culture with a spinner flask. Moreover, in each of these methods, a
20 continuous or intermittent medium exchange system may be mounted to tailor it for long term culturing.

[0094]

25 According to the invention, the cells can be cultured in a continuous manner. For example, the culturing in the method of the invention may be carried out in a manner with continuous circulation such as continuous addition and recovery of the medium on the porous polyimide film, or exposure of the porous polyimide film sheet to air using an open apparatus.

30 **[0095]**

Cell culturing according to the invention may be carried out in a system in which a cell culture medium is continuously or intermittently supplied to a cell culturing vessel from cell culture medium supply means
35 installed outside of the cell culturing vessel. The system may be such that the cell culture medium is circulated between the cell culture medium supply means

and the cell culturing vessel.

[0096]

The invention includes a mode in which a cell
culturing apparatus is set in an incubator and the cells
5 are cultured. When it is to be used in a system in which
the cell culture medium is continuously or intermittently
supplied to the cell culturing vessel from cell culture
medium supply means installed outside of the cell
culturing vessel, the system may be a cell culturing
10 apparatus including a culturing unit which is the cell
culturing vessel, and a culture medium-supply unit which
is the cell culture medium supply means, wherein the
culturing unit is a culturing unit that houses one or
more porous polyimide films to support cells, and that
15 comprises a culture medium supply port and a culture
medium discharge port, and the culture medium-supply unit
is a culture medium-supply unit comprising a culture
medium housing vessel, a culture medium supply line, and
a liquid conveyance pump that conveys the medium through
20 the culture medium supply line, the first end of the
culture medium supply line contacting the medium in the
culture medium housing vessel, and the second end of the
culture medium supply line being in communication with
the culturing unit interior via the culture medium supply
25 port of the culturing unit.

[0097]

In the cell culturing apparatus, the culturing unit
may be a culturing unit that does not comprise an air
supply port and an air discharge port, or it may be a
30 culturing unit that comprises an air supply port and an
air discharge port. Even if the culturing unit does not
comprise an air supply port and air discharge port, the
oxygen, etc. necessary for cell culturing is adequately
supplied to the cells through the medium. Furthermore,
35 in the cell culturing apparatus described above, the
culturing unit may further comprise a culture medium
discharge line, the first end of the culture medium

discharge line being connected to the culture medium housing vessel, the second end of the culture medium discharge line being in communication with the culturing unit interior via the culture medium discharge port of the culturing unit, and the medium being able to circulate through the culture medium-supply unit and the culturing unit.

[0098]

7. Production of substance by cells

The invention produces a desired substance by cells, by culturing the cells as described above. The produced substance may be a substance accumulating inside the cells, or a substance secreted from the cells. The produced substance may be collected by a known method, according to the type and properties of the substance. When the substance is secreted from the cells, the substance can be recovered from the cell culture medium. When the produced substance is a substance that accumulates inside the cells, the cells may be disrupted by a known method such as chemical treatment using a cytolytic agent or the like, ultrasonic treatment or physical treatment using a homogenizer or a shredding DispoTube, to move the substance out of the cells and recover it. The method of disrupting the cells may be any one applied as appropriate by a person skilled in the art depending on the type of cells and the type of substance.

[0099]

II. Substance-producing apparatus

The invention also relates to an apparatus for production of a substance by cell culturing, to be used in the method of the invention, the apparatus including a porous polyimide film. In the substance-producing apparatus of the invention, the porous polyimide film may be used in a fixed state, or it may be used in a floating state in the cell culture medium, and it may be either placed in the medium or exposed from the medium. In the

substance-producing apparatus, two or more porous polyimide films may be layered either above and below or left and right. The layered aggregates or cluster may be either placed in the medium or exposed from the medium.

5 **[0100]**

 The substance-producing apparatus for cell culturing of the invention may be a known cell culturing apparatus, in any desired form so long as it includes a porous polyimide film. The shape and scale of the culturing apparatus is not particularly restricted, and any scale from a dish or test tube to a large tank may be used, as appropriate. These include, for example, Cell Culture Dish by BD Falcon, and Nunc Cell Factory by Thermo Scientific. By using a porous polyimide film according to the invention, it has become possible to carry out culturing even of cells that have not been suitable for natural suspension culture, by using an apparatus intended for suspension culture, in a state similar to suspension culture. The apparatus for suspension culture that is used may be, for example, a spinner flask or rotating culture flask by Corning, Inc. As an environment allowing a similar function to be obtained, there may be used a hollow fiber culturing system such as the FiberCell[®] System by Veritas.

25 **[0101]**

 The substance-producing apparatus for cultured cells according to the invention may be a continuous circulating or open apparatus, wherein medium is continuously added to and recovered from sheets on the mesh, and the method may also be carried out with a type that exposes the porous polyimide film sheets to air.

[0102]

 Means may also be provided for recovering the substance that has been produced by the cells according to the invention. For example, by directly connecting a semipermeable membrane or the like to circulating added medium, it is possible to construct an efficient and

prolonged culturing/removal system in which sugars or amino acids are added while removing unwanted substances such as lactic acid.

[0103]

5 III. Kit

 The present invention also relates to a kit for use in the method of the invention, the kit including a porous polyimide film.

[0104]

10 The kit of the invention may include constituent elements necessary for cell culturing and substance production and recovery, in addition to the porous polyimide film, as appropriate. For example, it may include the cells to be applied to the porous polyimide
15 film, the cell culture medium, a continuous medium-supply apparatus, a continuous medium circulating apparatus, a scaffold or module to support the porous polyimide film, a cell culturing apparatus, ELISA to confirm the substance production, cell disruption means (for example,
20 a cytolytic agent and a shredding Dispo Homogenizer), substance recovering means (for example, an ultrafiltration centrifugation tube, coprecipitation reagent set and tube, and reagents such as antibodies), and the instruction manual for the kit.

25 **[0105]**

 While not restrictive, one mode includes a package containing either one or a plurality of sterilized porous polyimide films stored in a transparent pouch, in a form allowing their use for cell culturing, or a kit having a
30 sterile liquid encapsulated together with a porous polyimide film in the same pouch, in the form of an integrated film/liquid allowing efficient suction seeding.

35 **[0106]**

IV. Use

 The invention further includes use of a porous

polyimide film for the method of the invention described above.

Examples

5 [0107]

The present invention will now be explained in greater detail by examples. It is to be understood, however, that the invention is not limited to these examples. A person skilled in the art may easily
10 implement modifications and changes to the invention based on the description in the present specification, and these are also encompassed within the technical scope of the invention. Unless otherwise specified, the term "porous polyimide film" refers to a porous polyimide film
15 with a total film thickness of 25 μm and a porosity of 73%. Each porous polyimide film had at least two different surface layers (A-surface and B-surface), and a macro-void layer sandwiched between the two surface layers. The mean pore size of the holes in the A-surface
20 was 6 μm , and the mean pore size of the holes in the B-surface was 46 μm .

[0108]

The porous polyimide films used in the following examples were prepared by forming a polyamic acid
25 solution composition including a polyamic acid solution obtained from 3,3',4,4'-biphenyltetracarboxylic dianhydride (s-BPDA) as a tetracarboxylic acid component and 4,4'-diaminodiphenyl ether (ODA) as a diamine component, and polyacrylamide as a coloring precursor,
30 and performing heat treatment at 250°C or higher.

[0109]

<Cells and materials used>

- Normal human skin fibroblasts (Lonza, product code CC-
35 2511)
- Normal human fibroblast-like synovial membrane cells

HFLS (Cell Applications, Inc.)

- Chronic rheumatoid arthritis patient-derived fibroblast-like synovial membrane cells HFLS-RA (Cell Applications, Inc.)

5 • HepG2 (HEPG2-500 by CET (Cellular Engineering Technologies, Inc.))

- CHO-K1 (cat. 85051005 by Public Health England)

- CHO DP-12 (ATCC CRL-12445)

10 • CHO-K1 medium (Ham's F-12 087-08335 by Wako Pure Chemical Industries, Ltd.)

- CHO DP-12 medium (IMDM 098-06465 by Wako Pure Chemical Industries, Ltd.)

- Cell Counting Kit 8 (CK04, Dojindo Laboratories)

- Stainless steel mesh (60 mesh E9117 by Qholia)

15 • 2 cm × 2 cm sterilized square vessel (cat. 103 by Thermo Fisher Scientific)

- Penicillin-Streptomycin-Amphotericin B Suspension (X100) (161-23181 by Wako Pure Chemical Industries, Ltd.)

- Microscope, image software

20 LSM 700 by Carl Zeiss, software: ZEN

[0110]

Example 1

Spontaneous substance production by normal human skin fibroblasts

25 For this example, normal human skin fibroblasts were used for seeding and cell culturing on a porous polyimide film, and the amount of fibronectin produced on the petri dish was confirmed.

 After adding 1 ml of cell culture medium (2% FBS, 30 Fibroblast Media, Lonza) to a 2 cm × 2 cm square sterilized vessel, a 1.4 cm-square porous polyimide film was immersed in the medium with the A-surface of the mesh structure facing upward. Separately, there was prepared a normal human skin fibroblast suspension with normal 35 human skin fibroblasts suspended at 4.2×10^6 cells per 1 ml of medium (of which viable cells were 4.2×10^6 and

dead cells were 4.0×10^4 , for a viable cell rate of 99%). The cell suspension was added to the cell culture medium in the square vessel at $4 \times 10^4/\text{cm}^2$ per sheet, and culturing was carried out for 14 days in the square vessel. There was also prepared a normal human skin fibroblast suspension with normal human skin fibroblasts suspended at 3.0×10^6 cells per 1 ml of medium (of which viable cells were 2.9×10^6 and dead cells were 1.6×10^5 , for a viable cell rate of 95%). After adding 1 ml of cell culture medium to a 5 cm^2 petri dish, adding the cell suspension at $1.0 \times 10^4/\text{cm}^2$ and culturing for 43 days, cell culturing was carried out in a petri dish.

[0111]

There were prepared a 5 cm^2 petri dish containing a single cell-cultured sheet and a 5 cm^2 petri dish in which cells had been cultured without the sheet.

The medium supernatant was discarded, 1 ml of fresh cell culture medium was added to each, incubation was carried out for 2 days at 37°C with 5% CO_2 , and the medium supernatant was recovered. The amount of fibronectin in the recovered supernatant was quantified by ELISA (Table 2). Table 2 shows the amounts of fibronectin spontaneously produced using human skin fibroblasts, using the porous polyimide film sheet and petri dish as different culturing scaffolds. As shown in Table 2, using the porous polyimide film sheet resulted in more than twice the amount of spontaneous production of fibronectin than when using the petri dish.

[0112]

[Table 2]

Comparison of fibronectin production per unit area
(Example 1)

Culturing scaffold	Fibronectin (ng/cm^2)
Porous polyimide film	3.2×10^3
Petri dish	1.4×10^3

[0113]

Example 2

Spontaneous substance production by normal human
fibroblast-like synovial membrane cells HFLS and chronic
5 rheumatoid arthritis patient-derived fibroblast-like
synovial membrane cells HFLS-RA

For this example, normal human fibroblast-like
synovial membrane cells HFLS and chronic rheumatoid
arthritis patient-derived fibroblast-like synovial
10 membrane cells HFLS-RA were used for seeding and cell
culturing on porous polyimide films, after which the
amounts of IL-6 production in the petri dishes were
confirmed.

[0114]

15 After adding 1 ml of cell culture medium (10% FBS,
Synoviocyte Growth Medium, Cell Applications, Inc.) to a
2 cm × 2 cm square sterilized vessel, a 1.4 cm-square
porous polyimide film was immersed in the medium with the
A-surface of the mesh structure facing upward.
20 Separately, an HFLS cell suspension was prepared with the
HFLS cells suspended at 1.5×10^6 cells per 1 ml of medium
(of which 1.4×10^6 were viable cells and 2.0×10^4 were
dead cells, for a viable cell rate of 99%). The cell
suspension was added to the cell culture medium in the
25 square vessel at $4.0 \times 10^4/\text{cm}^2$ per sheet, and culturing
was carried out for 13 days in the square vessel. Also,
1 ml of cell culture medium was added to a 5 cm² petri
dish, and an HFLS cell suspension was separately prepared
with the HFLS cells suspended at 5.9×10^5 cells per 1 ml
30 of medium (of which 5.7×10^5 were viable cells and $2.0 \times$
 10^4 were dead cells, for a viable cell rate of 97%). The
cell suspension was added at $7 \times 10^3/\text{cm}^2$ and cultured for
8 days, and cell culturing was carried out in a petri
dish.

35 **[0115]**

Similarly, an HFLS-RA cell suspension was prepared

with the HFLS-RA cells suspended at 2.2×10^6 cells per 1 ml of medium, of which 2.0×10^6 were viable cells and 1.6×10^5 were dead cells, for a viable cell rate of 93%. The cell suspension was added to the cell culture medium in the square vessel at $4.0 \times 10^4/\text{cm}^2$ per sheet, and culturing was carried out for 13 days in the square vessel. Also, 1 ml of cell culture medium was added to a 5 cm^2 petri dish, and an HFLS-RA cell suspension was separately prepared with the HFLS-RA cells suspended at 5.8×10^5 cells per 1 ml of medium (of which 5.6×10^5 were viable cells and 2.0×10^4 were dead cells, for a viable cell rate of 97%). The cell suspension was added at $7.0 \times 10^3/\text{cm}^2$ and cultured for 8 days, and cell culturing was carried out in a petri dish.

15 **[0116]**

Two 5 cm^2 petri dishes were prepared, and one cell cultured HFLS sheet and one HFLS-RA sheet was placed in them, respectively. Also prepared were 5 cm^2 petri dishes of cell cultured HFLS and HFLS-RA without the sheets.

20 **[0117]**

The medium supernatant was discarded, 1 ml of fresh cell culture medium was added to each, incubation was carried out for 3 days at 37°C 5% CO_2 , and the medium supernatant was recovered. The amount of IL-6 in the recovered supernatant was quantified by ELISA (Table 3). Table 3 shows the amounts of IL-6 spontaneously produced using the normal human fibroblast-like synovial membrane cells, with the porous polyimide film sheets and petri dishes used as different culturing scaffolds. As shown in Table 3, using the porous polyimide film sheets resulted in more than about 25 times the amount of spontaneous production of IL-6 with normal human fibroblast-like synovial membrane cells HFLS and more than about 5 times the amount with chronic rheumatoid arthritis patient-derived fibroblast-like synovial membrane cells HFLS-RA, than when using the petri dishes.

[0118]

[Table 3]

IL-6 production by human fibroblast-like synovial
membrane cells (Example 2)

Cells and culturing scaffold	IL-6 (pg/ml)
Human fibroblast-like synovial membrane cells/Porous polyimide film	2.7×10^2
RA human fibroblast-like synovial membrane cells/Porous polyimide film	3.5×10^3
Human fibroblast-like synovial membrane cells/Petri dish	1.1×10^1
RA human fibroblast-like synovial membrane cells/Petri dish	7.1×10^2

5

[0119]

Example 3

Spontaneous albumin production by human hepatic cancer-
derived cell line, HepG2 cells

10 For this example, the human hepatic cancer-derived
cell line HepG2 was used for seeding and cell culturing
on a porous polyimide film, after which the amount of
albumin production spontaneously released into the medium
during a fixed period was confirmed.

15 **[0120]**

 After adding 1 ml of cell culture medium (10% FBS,
HEPG2.E.MEDIA-450, Cellular Engineering Technologies
Inc.) into a 2 cm × 2 cm square sterilized vessel, a 1.4
cm-square porous polyimide film was immersed in the
20 medium with the A-surface of the mesh structure facing
upward. Separately, a HepG2 cell suspension was prepared
with the HepG2 cells suspended at 1.6×10^6 cells per 1 ml
of medium (of which 1.4×10^6 were viable cells and $1.3 \times$
 10^5 were dead cells, for a viable cell rate of 92%). The
25 cell suspension was added to the cell culture medium in
the square vessel at $2 \times 10^4/\text{cm}^2$ per sheet, and culturing
was carried out for 21 days in the square vessel. Also,
2 ml of cell culture medium was added to a 10 cm² petri
dish and the cell suspension was added at $2 \times 10^4/\text{cm}^2$ for

cell culturing for 21 days, and cell culturing was carried out in a petri dish.

[0121]

There were prepared a 10 cm² petri dish containing
5 three cell-cultured sheets or a single sheet, and a 10 cm²
petri dish in which cells had been cultured. The medium
supernatant was discarded, 2 ml of fresh cell culture
medium was added to each, incubation was carried out for
3 hours at 37°C with 5% CO₂, and the medium supernatant
10 was recovered. The amount of albumin in the recovered
supernatant was quantified by ELISA (Tables 4 and 5).
Tables 4 and 5 show the amounts of albumin spontaneously
produced using the HepG2 cells, with the porous polyimide
film sheets and petri dish used as different culturing
15 scaffolds. Table 4 shows the amount of albumin
production per unit area of the film, and Table 5 shows
the amount of albumin production per cell. Since there
was no correlation between the number of sheets present
and the albumin production amount per cell, the
20 production amount per porous polyimide film being
constant, it is therefore expected that substance
production proceeds in a proportional manner with
increasing cell count.

[0122]

25 [Table 4]

Albumin production per unit area by HepG2 cells
(Example 3)

Culturing scaffold and cell amount	Albumin (ng/cm ²)
Porous polyimide films: 3 sheets	2.8×10^2
Porous polyimide films: 1 sheet	3.2×10^2
Petri dish	2.5×10^1

[0123]

[Table 5]

Albumin production per cell by HepG2 cells (Example 3)

Culturing scaffold and cell amount	Albumin (pg/cell)
Porous polyimide films: 3 sheets	2.8×10^{-1}
Porous polyimide films: 1 sheet	2.7×10^{-1}
Petri dish	9.7×10^{-2}

[0124]

5 Example 4

Substance production by G-CSF-producing CHO-K1 cell line

For this example, G-CSF was produced by cell culturing using G-CSF-producing CHO-K1 cells, and the amount of G-CSF released into the medium was measured.

10 **[0125]**

A suspension was prepared with the G-CSF-producing CHO-K1 cells suspended at 4.1×10^6 cells per 1 ml of medium (of which 3.6×10^6 were viable cells and 4.1×10^5 were dead cells, for a viable cell rate of 90%). The cell suspension was seeded on the A-surface of a 8×12.5 cm rectangular porous polyimide film at $4.1 \times 10^4/\text{cm}^2$ per sheet, and 20 ml of cell culture medium (1% FBS, Ham's F-12, product of Wako Pure Chemical Industries, Ltd.) was added to a 10×14 cm rectangular sterilized vessel, which was then immersed into medium. The medium was exchanged twice a week in the rectangular vessel, and culturing was carried out for 7 days. The medium supernatant of the cultured sheet was removed, 20 ml of fresh cell culture medium was added, incubation was performed for 24 hours at 37°C with 5% CO_2 , and the medium supernatant was recovered. The amount of G-CSF in the recovered supernatant was quantified by ELISA (Table 6).

[Table 6]

Culturing scaffold	G-CSF (pg/cell/day)
Porous polyimide film	3.4×10^{-2}

[0126]

Example 5

Substance production by anti-human IL-8-producing CHO DP-12 cells

5 For this example, the amount of anti-human IL-8 antibody produced and released in the medium by cell culturing using human anti-IL-8 antibody-producing CHO DP-12 cells was measured, to determine the efficiency of a cell culture system using a porous polyimide film. As
10 a comparative example, the amount of antibody production obtained by culturing in a common petri dish was also examined.

[0127]

15 After adding 0.5 ml of cell culture medium (2% FBS, IMDM, product of Wako Pure Chemical Industries, Ltd.) to a 2 cm × 2 cm square sterilized vessel, each sterilized 1.4 cm-square porous polyimide film was immersed therein with the A-surface of the mesh structure facing upward. A human anti-IL-8-producing CHO DP-12 cell suspension was
20 added to the sheets in each medium at 4×10^4 cells per sheet, and continuous cell culturing was carried out, with medium exchange at a frequency of twice a week. After cell culturing for 85 days, a CCK8 was used to measure the cell count. Two cell-cultured sheets were
25 each placed in a 10 cm² petri dish, 2 ml of cell culture medium was added to each, and incubation was performed for 24 hours at 37°C with 5% CO₂, after which the medium supernatants were recovered. The amount of anti-human IL-8 antibody in the recovered supernatant was quantified
30 by ELISA (Porous polyimide film 1 and Porous polyimide film 2 in Table 7).

[0128]

35 After adding 12 ml of cell culture medium to a 60 cm² petri dish, there was prepared a suspension of 5.3×10^6 human anti-IL-8-producing CHO DP-12 cells per 1 ml of medium (of which 5.3×10^6 were viable cells and 2.8×10^5

were dead cells, for a viable cell rate of 95%), seeding the cells at $2.0 \times 10^4/\text{cm}^2$. After 24 hours of culturing, the medium supernatant was recovered. The amount of human anti-IL-8 antibody in the recovered supernatant was quantified by ELISA (Petri dish 1 and Petri dish 2 in Table 7).

[0129]

[Table 7]

Culturing scaffold	Human anti-IL-8 antibody (pg/cell/day)
Porous polyimide film 1	16.2
Porous polyimide film 2	19.3
Petri dish 1	5.4
Petri dish 2	4.6

[0130]

Surprisingly, when the porous polyimide films were used as culturing scaffolds, the amount of substance production in each cell of the CHO-K1 cells increased by more than three times compared to using the petri dishes.

[0131]

Example 6

Substance production by anti-human IL-8-producing CHO DP-12 cells

For this example, in cell culturing using human anti-IL-8 antibody-producing CHO DP-12 cells, where the cultured cells were frozen while on the porous polyimide film sheet, rethawed and then cultured, it was examined whether or not this produced any change in anti-human IL-8 antibody produced and released into the medium when ordinary culturing was continued.

[0132]

After adding 0.5 ml of cell culture medium (2% FBS, IMDM, product of Wako Pure Chemical Industries, Ltd.) to a 2 cm $\times$ 2 cm square sterilized vessel, six sterilized 1.4 cm-square porous polyimide films were immersed therein with the A-surfaces of the mesh structures facing

upward. A human anti-IL-8-producing CHO DP-12 cell suspension was added to the sheets in each medium at 4×10^4 cells per sheet, and continuous cell culturing was carried out, with medium exchange at a frequency of twice a week. After cell culturing for 78 days, a CCK8 was used to measure the cell count. Next, two of the six porous polyimide film sheets supporting cells by cell culturing were each placed in a 10 cm^2 petri dish, 2 ml of cell culture medium was added to each, and incubation was performed for 24 hours at 37°C with 5% CO_2 , after which the medium supernatants were recovered. The amounts of anti-human IL-8 antibody in the recovered supernatants were quantified by ELISA (Non-frozen sheet 1 and Non-frozen sheet 2 in Table 8).

[0133]

Also, four of the six porous polyimide film sheets supporting cells by cell culturing were transferred to cryopreservation bags under sterile conditions, and 3 ml of CELLBANKER was added as a cell cryopreservation liquid. After freezing with a programmed freezer at -80°C under two different conditions (1°C per minute or 1°C every 10 minutes), it was stored at -80°C for 24 hours and transferred into liquid nitrogen. After 3 days, the contents were thawed by heating the bag at 37°C , 2 ml of medium was added and the mixture was allowed to stand in an incubator for 24 hours, after which each sheet was transferred to a $2 \text{ cm} \times 2 \text{ cm}$ square sterilized vessel, 1 ml of cell culture medium was added and culturing was continued for 3 days. Next, the sheets were each placed in a 10 cm^2 petri dish, 2 ml of cell culture medium was added to each, and incubation was performed for 24 hours at 37°C with 5% CO_2 , after which the medium supernatants were recovered. The amounts of anti-human IL-8 antibody in the recovered supernatants were quantified by ELISA (Frozen sheets 1 to 4 in Table 8). No change in anti-IL-8 production was seen due to the freezing.

[0134]

[Table 8]

	Human anti-IL-8 antibody (pg/cell/day)
Frozen sheet 1	26.4
Frozen sheet 2	18.0
Frozen sheet 3	24.0
Frozen sheet 4	28.7
Non-frozen sheet 1	16.2
Non-frozen sheet 2	19.3

5 **[0135]**

Example 7

Confirming growth by gas phase subculturing during long
term culturing of human skin fibroblasts

After adding 2 ml of medium to a 6 cm-diameter petri
10 dish, human skin fibroblasts were seeded on the A-
surfaces of the mesh structures of 1.4 cm-square
sterilized porous polyimide films, at 4×10^4 cells per
sheet, and culturing was carried out for 1 month. The
sheets were then cut into quarter portions and culturing
15 was continued for a total of 230 days of culturing.
Next, three 1.4 cm-square stainless steel meshes were
stacked and set at the center of a 3.5 cm dish, and the
porous polyimide film was placed thereover and sandwiched
with two empty 1.4 cm-square sterilized porous polyimide
20 films. When 1 ml of medium was added in this state, the
medium reached approximately the height of the sheets.
They were then directly transferred into a CO₂ incubator,
the medium was exchanged at a rate of twice per week, and
cell culturing was subsequently continued.

25 **[0136]**

After 7 days of culturing, each sheet was separately
isolated and culturing was continued in each sheet.
After 7, 10, 16, 21, 28, 42 and 56 days, the cell counts
were measured using a CCK8, and the cell growth behaviors
30 on the original sheets and the subsequently set empty
porous polyimide films were observed with a CCK8, based

on staining. The behavior was observed whereby the cells efficiently migrated from the porous polyimide films in which long term culturing of human skin fibroblasts had taken place, to the empty porous polyimide films, and continuously proliferated. The results are shown in Fig. 1.

[0137]

ELISA measurement was performed, using a human skin fibroblast-cultured sheet wherein continuous long term culturing had been carried out for 294 days on a porous polyimide film without gas phase subculturing, and a base sheet that had been gas-phase subcultured for the same period up to the 230th day and two sheets (upper and lower) that had been cultured by gas phase culturing for 56 days after subculturing, and the fibronectin produced by the living human skin fibroblasts was compared with the amount of fibronectin released in 24 hours into the medium in which the sheets had been cultured. Stable production of fibronectin was confirmed without any effect of the culturing period or gas phase subculturing. The results are shown in Table 9. For comparison, the fibronectin amount produced from two sheets cultured for 13 days with a porous polyimide film was also recorded.

[0138]

[Table 9]

Entry (days cultured and condition)	Fibronectin production per area (ng/cm ² /day)
Porous polyimide film, normal culturing, day 13 (Run 1)	480
Porous polyimide film, normal culturing, day 13 (Run 2)	376
Sheet cultured for 294 days (without gas phase subculturing)	760
Sheet cultured for 294 days (gas phase subcultured base sheet)	709
Sheet cultured for 56 days after gas phase subculturing (Upper)	338
Sheet cultured for 56 days after gas phase subculturing (Lower)	266

[0139]

5 Example 8

Freezing of human skin fibroblasts and substance
production

After adding 0.5 ml of cell culture medium to a 2 cm
× 2 cm square sterilized vessel, a 1.4 cm-square
10 sterilized porous polyimide film was immersed therein
with the A-surface of the mesh structure facing upward.
After adding a suspension of 4×10^4 human skin
fibroblasts per sheet onto the sheet in the medium,
culturing was initiated in a CO₂ incubator. Cell
15 culturing was continued with exchange of medium at a
frequency of twice per week, and after cell culturing for
49 days, a CCK8 was used to measure the cell count, which
was 9.1×10^4 .

[0140]

20 The cell-growing sheet was placed in a
cryopreservation bag containing 3 ml of CELLBANKER, and
after freezing to -80°C in a programmed freezer, by
lowering the temperature 1°C every 10 minutes, it was
stored at -80°C for 24 hours and then transferred to
25 liquid nitrogen. After 5 days, the bag was warmed to 37°C

to thaw the contents, 2 ml of medium was added, and the mixture was allowed to stand for 24 hours in an incubator. After 24 hours, 5 days, 8 days, 13 days, 21 days, 29 days and 35 days, a CCK8 was used to measure the cell count. After 24 hours, 8 days, 21 days and 35 days, the specific activity was found to be 34%, 91%, 105% and 152%, respectively. After culturing for 35 days, the amount of fibronectin production by the human skin fibroblasts growing on the same sheet was measured by ELISA. The results are shown in Table 10. For comparison, the fibronectin amount produced from two sheets cultured for 13 days with a porous polyimide film without freezing was also recorded. There was no damage by the freezing, and continuous substance production was confirmed.

[0141]

[Table 10]

Entry (days cultured and condition)	Fibronectin production per area (ng/cm ² /day)
Porous polyimide film, normal culturing, day 13 (Run 1)	480
Porous polyimide film, normal culturing, day 13 (Run 2)	376
Sheet cultured for 35 days after freezing and thawing	685

CLAIMS

1. A method of producing a substance using cells, the method including applying the cells to a porous polyimide film and culturing the cells to produce the substance by the cells.

2. The method according to claim 1, wherein the substance is selected from the group consisting of proteins, glycoproteins and viruses.

3. The method according to claim 1 or 2, wherein the cells are cultured under stationary culture conditions.

4. The method according to claim 1 or 2, wherein the cells are cultured under rotating culture or stirred conditions.

5. The method according to claim 1 or 2, wherein the cells are cultured in a continuous manner.

6. The method according to claim 5, including:
 setting a cell culturing apparatus in an incubator and culturing the cells, wherein
 the cell culturing apparatus includes:
 a culturing unit that houses one or more porous polyimide films to support cells, and that comprises a culture medium supply port and a culture medium discharge port, and
 a culture medium-supply unit comprising a culture medium housing vessel, a culture medium supply line, and a liquid conveyance pump that conveys the medium through the culture medium supply line, the first end of the culture medium supply line contacting the medium in the culture medium housing vessel, and the second end of the culture medium supply line being in communication with the culturing unit interior via the culture medium supply port of the culturing unit.

7. The method according to claim 6, wherein the cell culturing apparatus does not have an air supply port, an air discharge port and an oxygen permeation film.

8. The method according to any one of claims 1 to 7, wherein the porous polyimide film is a porous polyimide film comprising a polyimide obtained from a tetracarboxylic dianhydride and a diamine.

5 9. The method according to claim 8, wherein the porous polyimide film is a colored porous polyimide film obtained by forming a polyamic acid solution composition including a polyamic acid solution obtained from a tetracarboxylic dianhydride and a diamine, and a coloring precursor, and then heat treating it at 250°C or higher.

10 10. The method according to claim 8 or 9, wherein the porous polyimide film is a porous polyimide film with a multilayer structure, having two different surface layers and a macro-void layer.

15 11. The method according to claim 10, wherein the film thickness of the porous polyimide film is no greater than 75 μm .

 12. The method according to any one of claims 1 to 11, wherein the porous polyimide film is:

- 20 i) folded,
 ii) wound into a roll,
 iii) connected as sheets or fragments by a filamentous structure, or
 iv) bound into a rope,

25 and suspended or anchored in the cell culture medium in the cell culturing vessel.

 13. The method according to any one of claims 1 to 12, using two or more porous polyimide films layered either above and below or left and right in the cell culture medium.

30 14. The method according to any one of claims 1 to 13, wherein the cells are transformed by genetic engineering technology so as to express a substance.

35 15. The method according to any one of claims 1 to 14, wherein the cells are selected from the group consisting of animal cells, insect cells, plant cells, yeast cells and bacteria.

16. The method according to claim 15, wherein the animal cells are cells derived from an animal belonging to the subphylum Vertebrata.

5 17. The method according to claim 15 or 16, wherein the animal cells are selected from the group consisting of Chinese hamster ovary tissue-derived cells (CHO cells), African green monkey kidney-derived established cell lines (Vero cells), human hepatic cancer-derived cells (HepG2 cells), canine kidney epithelial cell-
10 derived cell lines (MDCK cells) and human hepatic cancer tissue-derived established cell lines (huGK-14).

18. The method according to claim 15, wherein the bacteria are selected from the group consisting of lactic acid bacteria, *E. coli*, *Bacillus subtilis* and
15 cyanobacteria.

19. A substance-producing apparatus for use in the method according to any one of claims 1 to 18, including a porous polyimide film.

20 20. The substance-producing apparatus according to claim 19, wherein two or more porous polyimide films are layered either above and below or left and right.

21. A kit for use in the method according to any one of claims 1 to 18, including a porous polyimide film.

25 22. Use of a porous polyimide film for the method according to any one of claims 1 to 18.