5

CIRCULATING RNA SIGNATURES SPECIFIC TO PREECLAMPSIA

CONTINUING APPLICATION DATA

This application claims the benefit of U.S. Provisional Application Serial No. 62/676,436 filed May 25, 2018, and U.S. Provisional Application Serial No. 62/848,219 filed May 15, 2019, each of which is incorporated by reference herein in its entirety.

FIELD OF INVENTION

The present invention relates generally to methods and materials for use in the detection and early risk assessment for the pregnancy complication preeclampsia.

BACKGROUND

Preeclampsia is a condition that occurs only during pregnancy, affecting 5% to 8% of all pregnancies. It is the direct cause of 10%-15% of maternal deaths and 40% of fetal deaths. The three main symptoms of preeclampsia may include high blood pressure, swelling of hands and feet, and excess protein in the urine (proteinuria), occurring after week 20 of pregnancy. Other signs and symptoms of preeclampsia may include severe headaches, changes in vision (including temporary loss of vision, blurred vision or light sensitivity), nausea or vomiting, decreased urine output, decreased platelets levels (thrombocytopenia), impaired liver function, and shortness of breath, caused by fluid in the lung.

The more severe the preeclampsia and the earlier it occurs in pregnancy, the greater the risks for mother and baby. Preeclampsia may require induced labor and delivery or delivery by cesarean delivery. Left untreated, preeclampsia can lead to serious, even fatal, complications for both the mother and baby. Complications of preeclampsia include fetal growth restriction, low birth weight, preterm birth, placental abruption, HELLP syndrome (hemolysis, elevated liver enzymes, and low platelet count syndrome), eclampsia (a severe form of preeclampsia that leads to seizures), organ damage, including kidney, liver, lung, heart, or eye damage, stroke or other brain injury. See, for example, “Preeclampsia - Symptoms and causes - Mayo Clinic,” April 3,

2018, available at on the worldwide web at mayoclinic.org/diseases-conditions/preeclampsia/symptoms-causes/syc-20355745.

With early detection and treatment, most women can deliver a healthy baby if preeclampsia is detected early and treated with regular prenatal care. Although various protein biomarkers display changed levels in maternal serum at presymptomatic stages, these biomarkers lack discriminative and predictive power in individual patients (Karumanchi and Granger, 2016, *Hypertension*; 67(2): 238–242). Thus, the identification of biomarkers for the early detection of preeclampsia is critical for the early diagnosis and treatment of preeclampsia.

SUMMARY OF THE INVENTION

The present invention includes a method of detecting preeclampsia and/or determining an increased risk for preeclampsia in a pregnant female, the method including:

identifying in a biosample obtained from the pregnant women a plurality of circulating RNA (C-RNA) molecules;

wherein a plurality of C-RNA molecules is selected from:

(a) a plurality of C-RNA molecules encoding at least a portion of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, up to all seventy-five of ARRDC2, JUN, SKIL, ATP13A3, PDE8B, GSTA3, PAPPA2, TIPARP, LEP, RGP1, USP54, CLEC4C, MRPS35, ARHGEF25, CUX2, HEATR9, FSTL3, DDI2, ZMYM6, ST6GALNAC3, GBP2, NES, ETV3, ADAM17, ATOH8, SLC4A3, TRAF3IP1, TTC21A, HEG1, ASTE1, TMEM108, ENC1, SCAMP1, ARRDC3, SLC26A2, SLIT3, CLIC5, TNFRSF21, PPP1R17, TPST1, GATSL2, SPDYE5, HIPK2, MTRNR2L6, CLCN1, GINS4, CRH, C10orf2, TRUB1, PRG2, ACY3, FAR2, CD63, CKAP4, TPCN1, RNF6, THTPA, FOS, PARN, ORAI3,

ELMO3, SMPD3, SERPINF1, TMEM11, PSMD11, EBI3, CLEC4M, CCDC151, CPAMD8, CNFN, LILRA4, ADA, C22orf39, PI4KAP1, and ARFGAP3; or

(b) a plurality of C-RNA molecules encoding at least a portion of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, any twenty six of more, or all twenty-seven of TIMP4, FLG, HTRA4, AMPH, LCN6, CRH, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, CLEC4C, KRT5, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, LEP, NES, and VSIG4; or

(c) a plurality of C-RNA molecules encoding at least a portion of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, up to all one hundred twenty-two of CYP26B1, IRF6, MYH14, PODXL, PPP1R3C, SH3RF2, TMC7, ZNF366, ADCY1, C6, FAM219A, HAO2, IGIP, IL1R2, NTRK2, SH3PXD2A, SSUH2, SULT2A1, FMO3, FSTL3, GATA5, HTRA1, C8B, H19, MN1, NFE2L1, PRDM16, AP3B2, EMP1, FLNC, STAG3, CPB2, TENC1, RP1L1, A1CF, NPR1, TEK, ERRFI1, ARHGEF15, CD34, RSPO3, ALPK3, SAMD4A, ZCCHC24, LEAP2, MYL2, NRG3, ZBTB16, SERPINA3, AQP7, SRPX, UACA, ANO1, FKBP5, SCN5A, PTPN21, CACNA1C, ERG, SOX17, WWTR1, AIF1L, CA3, HRG, TAT, AQP7P1, ADRA2C, SYNPO, FN1, GPR116, KRT17, AZGP1, BCL6B, KIF1C, CLIC5, GPR4, GJA5, OLAH, C14orf37, ZEB1, JAG2, KIF26A,

APOLD1, PNMT, MYOM3, PITPNM3, TIMP4, HTRA4, AMPH, LCN6, CRH, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, LEP, NES, VSIG4, HBG2, CADM2, LAMP5, PTGDR2, NOMO1, NXF3, PLD4, BPIFB3, PACSIN1, CUX2, FLG, CLEC4C, and KRT5; or

(d) a plurality of C-RNA molecules encoding at least a portion of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, any twenty-four or more, any twenty-five or more, any twenty-six or more, any twenty-seven or more, any twenty-eight or more, any twenty-nine or more, or all thirty of VSIG4, ADAMTS2, NES, FAM107A, LEP, DAAM2, ARHGEF25, TIMP3, PRX, ALOX15B, HSPA12B, IGFBP5, CLEC4C, SLC9A3R2, ADAMTS1, SEMA3G, KRT5, AMPH, PRG2, PAPPA2, TEAD4, CRH, PITPNM3, TIMP4, PNMT, ZEB1, APOLD1, PLD4, CUX2, and HTRA4; or

(e) a plurality of C-RNA molecules encoding at least a portion of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, any twenty-four or more, any twenty-five or more, or all twenty-six of ADAMTS1, ADAMTS2, ALOX15B, AMPH, ARHGEF25, CELF4, DAAM2, FAM107A, HSPA12B, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LEP, LRRC26, NES, OLAH, PACSIN1, PAPPA2, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, and VSIG4; or

(f) a plurality of C-RNA molecules encoding at least a portion of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, or all twenty-two of ADAMTS1, ADAMTS2, ALOX15B, ARHGEF25, CELF4, DAAM2, FAM107A, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LEP, LRRC26, NES, OLAH, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, and VSIG4; or

(g) a plurality of C-RNA molecules encoding at least a portion of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, or all eleven of CLEC4C, ARHGEF25, ADAMTS2, LEP, ARRDC2, SKIL, PAPPA2, VSIG4, ARRDC4, CRH, and NES (including in some embodiments, the seven of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, and VSIG4; the eight of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, SKIL, and VSIG4; the eight of ADAMTS2, ARHGEF25, ARRDC4, CLEC4C, LEP, NES, SKIL, and VSIG4; the ten of ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, CRH, LEP, PAPPA2, SKIL, and VSIG4; the of six of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, and SKIL; or the eight of ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, LEP, PAPPA2, and SKIL); or

(h) a plurality of C-RNA molecules encoding at least a portion of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, or all twenty-four of LEP, PAPPA2, KCNA5, ADAMTS2, MYOM3,

ATP13A3, ARHGEF25, ADA, HTRA4, NES, CRH, ACY3, PLD4, SCT, NOX4, PACSIN1, SERPINF1, SKIL, SEMA3G, TIPARP, LRRC26, PHEX, LILRA4, and PER1

is indicative of preeclampsia and/or an increased risk for preeclampsia in the pregnant women.

The present invention includes a method of detecting preeclampsia and/or determining an increased risk for preeclampsia in a pregnant female, the method including:

obtaining a biosample from the pregnant female;

purifying a population of circulating RNA (C-RNA) molecules from the biosample;

identifying protein coding sequences encoded by the C-RNA molecules within the purified population of C-RNA molecules;

wherein protein coding sequences encoded by the C-RNA molecules encoding at least a portion of a protein is selected from:

(a) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, any fifty or more, any seventy or more, or all seventy-five of ARRDC2, JUN, SKIL, ATP13A3, PDE8B, GSTA3, PAPPA2, TIPARP, LEP, RGP1, USP54, CLEC4C, MRPS35, ARHGEF25, CUX2, HEATR9, FSTL3, DDI2, ZMYM6, ST6GALNAC3, GBP2, NES, ETV3, ADAM17, ATOH8, SLC4A3, TRAF3IP1, TTC21A, HEG1, ASTE1, TMEM108, ENC1, SCAMP1, ARRDC3, SLC26A2, SLIT3, CLIC5, TNFRSF21, PPP1R17, TPST1, GATSL2, SPDYE5, HIPK2, MTRNR2L6, CLCN1, GINS4, CRH, C10orf2, TRUB1, PRG2, ACY3, FAR2, CD63, CKAP4, TPCN1, RNF6, THTPA, FOS, PARN, ORAI3, ELMO3, SMPD3, SERPINF1, TMEM11, PSMD11, EBI3, CLEC4M, CCDC151, CPAMD8, CNFN, LILRA4, ADA, C22orf39, PI4KAP1, and ARFGAP3; or

(b) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine

or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, any twenty six of more, or all twenty-seven of TIMP4, FLG, HTRA4, AMPH, LCN6, CRH, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, CLEC4C, KRT5, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, LEP, NES, and VSIG4; or

(c) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, any fifty or more, any seventy-five or more, any one hundred or more, or all one hundred twenty-two of CYP26B1, IRF6, MYH14, PODXL, PPP1R3C, SH3RF2, TMC7, ZNF366, ADCY1, C6, FAM219A, HAO2, IGIP, IL1R2, NTRK2, SH3PXD2A, SSUH2, SULT2A1, FMO3, FSTL3, GATA5, HTRA1, C8B, H19, MN1, NFE2L1, PRDM16, AP3B2, EMP1, FLNC, STAG3, CPB2, TENC1, RP1L1, A1CF, NPR1, TEK, ERRFI1, ARHGEF15, CD34, RSPO3, ALPK3, SAMD4A, ZCCHC24, LEAP2, MYL2, NRG3, ZBTB16, SERPINA3, AQP7, SRPX, UACA, ANO1, FKBP5, SCN5A, PTPN21, CACNA1C, ERG, SOX17, WWTR1, AIF1L, CA3, HRG, TAT, AQP7P1, ADRA2C, SYNPO, FN1, GPR116, KRT17, AZGP1, BCL6B, KIF1C, CLIC5, GPR4, GJA5, OLAH, C14orf37, ZEB1, JAG2, KIF26A, APOLD1, PNMT, MYOM3, PITPNM3, TIMP4, HTRA4, AMPH, LCN6, CRH, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, LEP, NES, VSIG4, HBG2, CADM2, LAMP5, PTGDR2, NOMO1, NXF3, PLD4, BPIFB3, PACSIN1, CUX2, FLG, CLEC4C, and KRT5; or

(d) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, any twenty-four or more, any twenty-five or more, any twenty-six or more, any twenty-seven or more, any twenty-eight or more, any twenty-nine or more, or all thirty of VSIG4, ADAMTS2, NES, FAM107A, LEP, DAAM2, ARHGEF25, TIMP3, PRX, ALOX15B, HSPA12B, IGFBP5, CLEC4C, SLC9A3R2, ADAMTS1, SEMA3G, KRT5, AMPH, PRG2, PAPPA2, TEAD4, CRH, PITPNM3, TIMP4, PNMT, ZEB1, APOLD1, PLD4, CUX2, and HTRA4; or

(e) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, any twenty-four or more, any twenty-five or more, or all twenty-six of ADAMTS1, ADAMTS2, ALOX15B, AMPH, ARHGEF25, CELF4, DAAM2, FAM107A, HSPA12B, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LEP, LRRC26, NES, OLAH, PACSIN1, PAPPA2, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, and VSIG4; or

(f) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, or all twenty-two of ADAMTS1, ADAMTS2, ALOX15B, ARHGEF25, CELF4, DAAM2, FAM107A, HTRA4, IGFBP5,

KCNA5, KRT5, LCN6, LEP, LRRC26, NES, OLAH, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, and VSIG4; or

(g) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, or all eleven of CLEC4C, ARHGEF25, ADAMTS2, LEP, ARRDC2, SKIL, PAPPA2, VSIG4, ARRDC4, CRH, and NES (including in some embodiments, the seven of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, and VSIG4; the eight of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, SKIL, and VSIG4; the eight of ADAMTS2, ARHGEF25, ARRDC4, CLEC4C, LEP, NES, SKIL, and VSIG4; the ten of ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, CRH, LEP, PAPPA2, SKIL, and VSIG4; the of six of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, and SKIL; or the eight of ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, LEP, PAPPA2, and SKIL); or

(h) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, or all twenty-four of LEP, PAPPA2, KCNA5, ADAMTS2, MYOM3, ATP13A3, ARHGEF25, ADA, HTRA4, NES, CRH, ACY3, PLD4, SCT, NOX4, PACSIN1, SERPINF1, SKIL, SEMA3G, TIPARP, LRRC26, PHEX, LILRA4, and PER1

is indicative of preeclampsia and/or an increased risk for preeclampsia in the pregnant women.

In some aspects, identifying protein coding sequences encoded by C-RNA molecules within the biosample includes hybridization, reverse transcriptase PCR, microarray chip analysis, or sequencing.

In some aspects, identifying protein coding sequences encoded by the C-RNA molecules within the biosample includes sequencing, including, for example, massively parallel sequencing of clonally amplified molecules and/or RNA sequencing.

In some aspects, the method further includes removing intact cells from the biosample; treating the biosample with a deoxynuclease (DNase) to remove cell free DNA (cfDNA); synthesizing complementary DNA (cDNA) from C-RNA molecules in the biosample; and/or enriching the cDNA sequences for DNA sequences that encode proteins by exome enrichment prior to identifying protein coding sequence encoded by the circulating RNA (C-RNA) molecules.

The present invention includes a method of detecting preeclampsia and/or determining an increased risk for preeclampsia in a pregnant female, the method including:

obtaining a biological sample from the pregnant female;

removing intact cells from the biosample;

treating the biosample with a deoxynuclease (DNase) to remove cell free DNA (cfDNA);

synthesizing complementary DNA (cDNA) from RNA molecules in the biosample;

enriching the cDNA sequences for DNA sequences that encode proteins (exome enrichment);

sequencing the resulting enriched cDNA sequences; and

identifying protein coding sequences encoded by enriched C-RNA molecules;

wherein protein coding sequences encoded by the C-RNA molecules encoding at least a portion of a protein selected from:

(a) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, up to all seventy-five of ARRDC2, JUN, SKIL, ATP13A3, PDE8B, GSTA3, PAPPA2, TIPARP, LEP, RGP1, USP54, CLEC4C, MRPS35, ARHGEF25, CUX2, HEATR9, FSTL3, DDI2, ZMYM6, ST6GALNAC3, GBP2, NES, ETV3, ADAM17, ATOH8, SLC4A3, TRAF3IP1, TTC21A, HEG1, ASTE1, TMEM108, ENC1, SCAMP1, ARRDC3, SLC26A2, SLIT3, CLIC5, TNFRSF21, PPP1R17, TPST1, GATSL2, SPDYE5, HIPK2, MTRNR2L6, CLCN1, GINS4, CRH, C10orf2, TRUB1, PRG2, ACY3,

FAR2, CD63, CKAP4, TPCN1, RNF6, THTPA, FOS, PARN, ORAI3, ELMO3, SMPD3, SERPINF1, TMEM11, PSMD11, EBI3, CLEC4M, CCDC151, CPAMD8, CNFN, LILRA4, ADA, C22orf39, PI4KAP1, and ARFGAP3; or

(b) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, any twenty six of more, or all twenty-seven of TIMP4, FLG, HTRA4, AMPH, LCN6, CRH, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, CLEC4C, KRT5, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, LEP, NES, and VSIG4; or

(c) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, up to all one hundred twenty-two of CYP26B1, IRF6, MYH14, PODXL, PPP1R3C, SH3RF2, TMC7, ZNF366, ADCY1, C6, FAM219A, HAO2, IGIP, IL1R2, NTRK2, SH3PXD2A, SSUH2, SULT2A1, FMO3, FSTL3, GATA5, HTRA1, C8B, H19, MN1, NFE2L1, PRDM16, AP3B2, EMP1, FLNC, STAG3, CPB2, TENC1, RP1L1, A1CF, NPR1, TEK, ERRFI1, ARHGEF15, CD34, RSPO3, ALPK3, SAMD4A, ZCCHC24, LEAP2, MYL2, NRG3, ZBTB16, SERPINA3, AQP7, SRPX, UACA, ANO1, FKBP5, SCN5A, PTPN21, CACNA1C, ERG, SOX17, WWTR1, AIF1L, CA3, HRG, TAT, AQP7P1, ADRA2C, SYNPO, FN1, GPR116, KRT17, AZGP1, BCL6B, KIF1C, CLIC5, GPR4, GJA5, OLAH, C14orf37, ZEB1, JAG2, KIF26A, APOLD1, PNMT, MYOM3, PITPNM3, TIMP4, HTRA4, AMPH, LCN6, CRH,

TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, LEP, NES, VSIG4, HBG2, CADM2, LAMP5, PTGDR2, NOMO1, NXF3, PLD4, BPIFB3, PACSIN1, CUX2, FLG, CLEC4C, and KRT5; or

(d) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, any twenty-four or more, any twenty-five or more, any twenty-six or more, any twenty-seven or more, any twenty-eight or more, any twenty-nine or more, or all thirty of VSIG4, ADAMTS2, NES, FAM107A, LEP, DAAM2, ARHGEF25, TIMP3, PRX, ALOX15B, HSPA12B, IGFBP5, CLEC4C, SLC9A3R2, ADAMTS1, SEMA3G, KRT5, AMPH, PRG2, PAPPA2, TEAD4, CRH, PITPNM3, TIMP4, PNMT, ZEB1, APOLD1, PLD4, CUX2, and HTRA4; or

(e) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, any twenty-four or more, any twenty-five or more, or all twenty-six of ADAMTS1, ADAMTS2, ALOX15B, AMPH, ARHGEF25, CELF4, DAAM2, FAM107A, HSPA12B, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LEP, LRRC26, NES, OLAH, PACSIN1, PAPPA2, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, and VSIG4; or

(f) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any

seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, or all twenty-two of ADAMTS1, ADAMTS2, ALOX15B, ARHGEF25, CELF4, DAAM2, FAM107A, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LEP, LRRC26, NES, OLAH, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, and VSIG4; or

(g) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, or all eleven of CLEC4C, ARHGEF25, ADAMTS2, LEP, ARRDC2, SKIL, PAPPA2, VSIG4, ARRDC4, CRH, and NES (including in some embodiments, the seven of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, and VSIG4; the eight of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, SKIL, and VSIG4; the eight of ADAMTS2, ARHGEF25, ARRDC4, CLEC4C, LEP, NES, SKIL, and VSIG4; the ten of ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, CRH, LEP, PAPPA2, SKIL, and VSIG4; the of six of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, and SKIL; or the eight of ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, LEP, PAPPA2, and SKIL); or

(h) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, or all twenty-four of LEP, PAPPA2, KCNA5, ADAMTS2, MYOM3, ATP13A3, ARHGEF25, ADA, HTRA4, NES, CRH, ACY3, PLD4, SCT, NOX4, PACSIN1, SERPINF1, SKIL, SEMA3G, TIPARP, LRRC26, PHEX, LILRA4, and PER1

is indicative of preeclampsia and/or an increased risk for preeclampsia in the pregnant women.

The present invention includes a method of identifying a circulating RNA signature associated with an increased risk of preeclampsia, the method including obtaining a biological sample from the pregnant female; removing intact cells from the biosample; treating the

biosample with a deoxynuclease (DNase) to remove cell free DNA (cfDNA); synthesizing complementary DNA (cDNA) from RNA molecules in the biosample; enriching the cDNA sequences for DNA sequences that encode proteins (exome enrichment); sequencing the resulting enriched cDNA sequences; and identifying protein coding sequences encoded by enriched C-RNA molecules.

The present invention includes a method including:

obtaining a biological sample from the pregnant female;

removing intact cells from the biosample;

treating the biosample with a deoxynuclease (DNase) to remove cell free DNA (cfDNA);

synthesizing complementary DNA (cDNA) from RNA molecules in the biosample;

enriching the cDNA sequences for DNA sequences that encode proteins (exome enrichment);

sequencing the resulting enriched cDNA sequences; and

identifying protein coding sequences encoded by the enriched C-RNA molecules;

wherein the protein coding sequences include at least a portion of a protein selected from:

(a) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five up to all seventy-five ARRDC2, JUN, SKIL, ATP13A3, PDE8B, GSTA3, PAPPA2, TIPARP, LEP, RGP1, USP54, CLEC4C, MRPS35, ARHGEF25, CUX2, HEATR9, FSTL3, DDI2, ZMYM6, ST6GALNAC3, GBP2, NES, ETV3, ADAM17, ATOH8, SLC4A3, TRAF3IP1, TTC21A, HEG1, ASTE1, TMEM108, ENC1, SCAMP1, ARRDC3, SLC26A2, SLIT3, CLIC5, TNFRSF21, PPP1R17, TPST1, GATSL2, SPDYE5, HIPK2, MTRNR2L6, CLCN1, GINS4, CRH, C10orf2, TRUB1, PRG2, ACY3, FAR2, CD63, CKAP4, TPCN1, RNF6, THTPA, FOS, PARN, ORAI3, ELMO3, SMPD3, SERPINF1, TMEM11, PSMD11, EBI3, CLEC4M, CCDC151, CPAMD8, CNFN, LILRA4, ADA, C22orf39, PI4KAP1, and ARFGAP3; or

(b) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, any twenty six of more, or all twenty-seven of TIMP4, FLG, HTRA4, AMPH, LCN6, CRH, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, CLEC4C, KRT5, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, LEP, NES, and VSIG4; or

(c) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, up to all one hundred twenty-two of CYP26B1, IRF6, MYH14, PODXL, PPP1R3C, SH3RF2, TMC7, ZNF366, ADCY1, C6, FAM219A, HAO2, IGIP, IL1R2, NTRK2, SH3PXD2A, SSUH2, SULT2A1, FMO3, FSTL3, GATA5, HTRA1, C8B, H19, MN1, NFE2L1, PRDM16, AP3B2, EMP1, FLNC, STAG3, CPB2, TENC1, RP1L1, A1CF, NPR1, TEK, ERRFI1, ARHGEF15, CD34, RSPO3, ALPK3, SAMD4A, ZCCHC24, LEAP2, MYL2, NRG3, ZBTB16, SERPINA3, AQP7, SRPX, UACA, ANO1, FKBP5, SCN5A, PTPN21, CACNA1C, ERG, SOX17, WWTR1, AIF1L, CA3, HRG, TAT, AQP7P1, ADRA2C, SYNPO, FN1, GPR116, KRT17, AZGP1, BCL6B, KIF1C, CLIC5, GPR4, GJA5, OLAH, C14orf37, ZEB1, JAG2, KIF26A, APOLD1, PNMT, MYOM3, PITPNM3, TIMP4, HTRA4, AMPH, LCN6, CRH, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2,

FAM107A, LEP, NES, VSIG4, HBG2, CADM2, LAMP5, PTGDR2, NOMO1, NXF3, PLD4, BPIFB3, PACSIN1, CUX2, FLG, CLEC4C, and KRT5; or

(d) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, any twenty-four or more, any twenty-five or more, any twenty-six or more, any twenty-seven or more, any twenty-eight or more, any twenty-nine or more, or all thirty of VSIG4, ADAMTS2, NES, FAM107A, LEP, DAAM2, ARHGEF25, TIMP3, PRX, ALOX15B, HSPA12B, IGFBP5, CLEC4C, SLC9A3R2, ADAMTS1, SEMA3G, KRT5, AMPH, PRG2, PAPPA2, TEAD4, CRH, PITPNM3, TIMP4, PNMT, ZEB1, APOLD1, PLD4, CUX2, and HTRA4; or

(e) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, any twenty-four or more, any twenty-five or more, or all twenty-six of ADAMTS1, ADAMTS2, ALOX15B, AMPH, ARHGEF25, CELF4, DAAM2, FAM107A, HSPA12B, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LEP, LRRC26, NES, OLAH, PACSIN1, PAPPA2, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, and VSIG4; or

(f) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, or all twenty-two of ADAMTS1, ADAMTS2,

ALOX15B, ARHGEF25, CELF4, DAAM2, FAM107A, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LEP, LRRC26, NES, OLAH, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, and VSIG4; or

(g) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, or all eleven of CLEC4C, ARHGEF25, ADAMTS2, LEP, ARRDC2, SKIL, PAPPA2, VSIG4, ARRDC4, CRH, and NES (including in some embodiments, the seven of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, and VSIG4; the eight of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, SKIL, and VSIG4; the eight of ADAMTS2, ARHGEF25, ARRDC4, CLEC4C, LEP, NES, SKIL, and VSIG4; the ten of ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, CRH, LEP, PAPPA2, SKIL, and VSIG4; the of six of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, and SKIL; or the eight of ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, LEP, PAPPA2, and SKIL); or

(h) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, or all twenty-four of LEP, PAPPA2, KCNA5, ADAMTS2, MYOM3, ATP13A3, ARHGEF25, ADA, HTRA4, NES, CRH, ACY3, PLD4, SCT, NOX4, PACSIN1, SERPINF1, SKIL, SEMA3G, TIPARP, LRRC26, PHEX, LILRA4, and PER1.

In some aspects, the biosample includes plasma.

In some aspects, the biosample is obtained from a pregnant female at less than 16 weeks gestation or at less than 20 weeks gestation.

In some aspects, the biosample is obtained from a pregnant female at greater than 20 weeks gestation.

The present invention includes a circulating RNA (C-RNA) signature for an elevated risk of preeclampsia, the C-RNA signature encoding at least a portion of any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, any fifty or more, any seventy or more, up to all seventy-five ARRDC2, JUN, SKIL, ATP13A3, PDE8B, GSTA3, PAPPA2, TIPARP, LEP, RGP1, USP54, CLEC4C, MRPS35, ARHGEF25, CUX2, HEATR9, FSTL3, DDI2, ZMYM6, ST6GALNAC3, GBP2, NES, ETV3, ADAM17, ATOH8, SLC4A3, TRAF3IP1, TTC21A, HEG1, ASTE1, TMEM108, ENC1, SCAMP1, ARRDC3, SLC26A2, SLIT3, CLIC5, TNFRSF21, PPP1R17, TPST1, GATSL2, SPDYE5, HIPK2, MTRNR2L6, CLCN1, GINS4, CRH, C10orf2, TRUB1, PRG2, ACY3, FAR2, CD63, CKAP4, TPCN1, RNF6, THTPA, FOS, PARN, ORAI3, ELMO3, SMPD3, SERPINF1, TMEM11, PSMD11, EBI3, CLEC4M, CCDC151, CPAMD8, CNFN, LILRA4, ADA, C22orf39, PI4KAP1, and ARFGAP3.

The present invention includes a circulating RNA (C-RNA) signature for an elevated risk of preeclampsia, the C-RNA signature encoding at least a portion of any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, any twenty six of more, or all twenty-seven of TIMP4, FLG, HTRA4, AMPH, LCN6, CRH, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, CLEC4C, KRT5, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, LEP, NES, and VSIG4.

The present invention includes a circulating RNA (C-RNA) signature for an elevated risk of preeclampsia, the C-RNA signature encoding a least a portion of a plurality of CYP26B1, IRF6, MYH14, PODXL, PPP1R3C, SH3RF2, TMC7, ZNF366, ADCY1, C6, FAM219A, HAO2, IGIP, IL1R2, NTRK2, SH3PXD2A, SSUH2, SULT2A1, FMO3, FSTL3, GATA5,

HTRA1, C8B, H19, MN1, NFE2L1, PRDM16, AP3B2, EMP1, FLNC, STAG3, CPB2, TENC1, RP1L1, A1CF, NPR1, TEK, ERRFI1, ARHGEF15, CD34, RSPO3, ALPK3, SAMD4A, ZCCHC24, LEAP2, MYL2, NRG3, ZBTB16, SERPINA3, AQP7, SRPX, UACA, ANO1, FKBP5, SCN5A, PTPN21, CACNA1C, ERG, SOX17, WWTR1, AIF1L, CA3, HRG, TAT, AQP7P1, ADRA2C, SYNPO, FN1, GPR116, KRT17, AZGP1, BCL6B, KIF1C, CLIC5, GPR4, GJA5, OLAH, C14orf37, ZEB1, JAG2, KIF26A, APOLD1, PNMT, MYOM3, PITPNM3, TIMP4, HTRA4, AMPH, LCN6, CRH, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, LEP, NES, VSIG4, HBG2, CADM2, LAMP5, PTGDR2, NOMO1, NXF3, PLD4, BPIFB3, PACSIN1, CUX2, FLG, CLEC4C, and KRT5.

The present invention includes a circulating RNA (C-RNA) signature for an elevated risk of preeclampsia, the C-RNA signature encoding at least a portion of any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, any twenty-four or more, any twenty-five or more, any twenty-six or more, any twenty-seven or more, any twenty-eight or more, any twenty-nine or more, or all thirty of VSIG4, ADAMTS2, NES, FAM107A, LEP, DAAM2, ARHGEF25, TIMP3, PRX, ALOX15B, HSPA12B, IGFBP5, CLEC4C, SLC9A3R2, ADAMTS1, SEMA3G, KRT5, AMPH, PRG2, PAPPA2, TEAD4, CRH, PITPNM3, TIMP4, PNMT, ZEB1, APOLD1, PLD4, CUX2, and HTRA4.

The present invention includes a circulating RNA (C-RNA) signature for an elevated risk of preeclampsia, the C-RNA signature encoding at least a portion of any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, any twenty-four or more, any twenty-five or more, or all twenty-six of ADAMTS1, ADAMTS2, ALOX15B, AMPH, ARHGEF25, CELF4, DAAM2, FAM107A, HSPA12B, HTRA4, IGFBP5, KCNA5, KRT5,

LCN6, LEP, LRRC26, NES, OLAH, PACSIN1, PAPPA2, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, and VSIG4.

The present invention includes a circulating RNA (C-RNA) signature for an elevated risk of preeclampsia, the C-RNA signature encoding at least a portion of any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, or all twenty-two of ADAMTS1, ADAMTS2, ALOX15B, ARHGEF25, CELF4, DAAM2, FAM107A, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LEP, LRRC26, NES, OLAH, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, and VSIG4.

The present invention includes a circulating RNA (C-RNA) signature for an elevated risk of preeclampsia, the C-RNA signature encoding at least a portion of any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, or all eleven of CLEC4C, ARHGEF25, ADAMTS2, LEP, ARRDC2, SKIL, PAPPA2, VSIG4, ARRDC4, CRH, and NES, including in some embodiments, the seven of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, and VSIG4; the eight of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, SKIL, and VSIG4; the eight of ADAMTS2, ARHGEF25, ARRDC4, CLEC4C, LEP, NES, SKIL, and VSIG4; the ten of ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, CRH, LEP, PAPPA2, SKIL, and VSIG4; the of six of ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, and SKIL; or the eight of ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, LEP, PAPPA2, and SKIL.

The present invention includes a circulating RNA (C-RNA) signature for an elevated risk of preeclampsia, the C-RNA signature encoding at least a portion of any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, any twenty-three or more, or all twenty-four of LEP, PAPPA2, KCNA5, ADAMTS2, MYOM3, ATP13A3, ARHGEF25, ADA, HTRA4, NES, CRH,

ACY3, PLD4, SCT, NOX4, PACSIN1, SERPINF1, SKIL, SEMA3G, TIPARP, LRRC26, PHEX, LILRA4, and PER1.

The present invention includes a solid support array comprising a plurality of agents capable of binding and/or identifying a C-RNA signature as described herein.

The present invention includes a kit comprising a plurality of probes capable of binding and/or identifying a C-RNA signature as described herein.

The present invention includes a kit comprising a plurality of primers for selectively amplifying a C-RNA signature as described herein.

As used herein, the term “nucleic acid” is intended to be consistent with its use in the art and includes naturally occurring nucleic acids or functional analogs thereof. Particularly useful functional analogs are capable of hybridizing to a nucleic acid in a sequence specific fashion or capable of being used as a template for replication of a particular nucleotide sequence. Naturally occurring nucleic acids generally have a backbone containing phosphodiester bonds. An analog structure can have an alternate backbone linkage including any of a variety of those known in the art. Naturally occurring nucleic acids generally have a deoxyribose sugar (e.g. found in deoxyribonucleic acid (DNA)) or a ribose sugar (e.g. found in ribonucleic acid (RNA)). A nucleic acid can contain any of a variety of analogs of these sugar moieties that are known in the art. A nucleic acid can include native or non-native bases. In this regard, a native deoxyribonucleic acid can have one or more bases selected from the group consisting of adenine, thymine, cytosine or guanine and a ribonucleic acid can have one or more bases selected from the group consisting of uracil, adenine, cytosine or guanine. Useful non-native bases that can be included in a nucleic acid are known in the art. The term "template" and “target," when used in reference to a nucleic acid, is intended as a semantic identifier for the nucleic acid in the context of a method or composition set forth herein and does not necessarily limit the structure or function of the nucleic acid beyond what is otherwise explicitly indicated.

As used herein, “amplify,” “amplifying” or “amplification reaction” and their derivatives, refer generally to any action or process whereby at least a portion of a nucleic acid molecule is replicated or copied into at least one additional nucleic acid molecule. The additional nucleic acid molecule optionally includes sequence that is substantially identical or substantially complementary to at least some portion of the target nucleic acid molecule. The target nucleic acid molecule can be single-stranded or double-stranded and the additional nucleic acid molecule

can independently be single-stranded or double-stranded. Amplification optionally includes linear or exponential replication of a nucleic acid molecule. In some embodiments, such amplification can be performed using isothermal conditions; in other embodiments, such amplification can include thermocycling. In some embodiments, the amplification is a multiplex amplification that includes the simultaneous amplification of a plurality of target sequences in a single amplification reaction. In some embodiments, "amplification" includes amplification of at least some portion of DNA and RNA based nucleic acids alone, or in combination. The amplification reaction can include any of the amplification processes known to one of ordinary skill in the art. In some embodiments, the amplification reaction includes polymerase chain reaction (PCR).

As used herein, "amplification conditions" and its derivatives, generally refers to conditions suitable for amplifying one or more nucleic acid sequences. Such amplification can be linear or exponential. In some embodiments, the amplification conditions can include isothermal conditions or alternatively can include thermocyling conditions, or a combination of isothermal and thermocycling conditions. In some embodiments, the conditions suitable for amplifying one or more nucleic acid sequences include polymerase chain reaction (PCR) conditions. Typically, the amplification conditions refer to a reaction mixture that is sufficient to amplify nucleic acids such as one or more target sequences, or to amplify an amplified target sequence ligated to one or more adapters, e.g., an adapter-ligated amplified target sequence. Generally, the amplification conditions include a catalyst for amplification or for nucleic acid synthesis, for example a polymerase; a primer that possesses some degree of complementarity to the nucleic acid to be amplified; and nucleotides, such as deoxyribonucleotide triphosphates (dNTPs) to promote extension of the primer once hybridized to the nucleic acid. The amplification conditions can require hybridization or annealing of a primer to a nucleic acid, extension of the primer and a denaturing step in which the extended primer is separated from the nucleic acid sequence undergoing amplification. Typically, but not necessarily, amplification conditions can include thermocycling; in some embodiments, amplification conditions include a plurality of cycles where the steps of annealing, extending and separating are repeated. Typically, the amplification conditions include cations such as Mg^{++} or Mn^{++} and can also include various modifiers of ionic strength.

As used herein, the term "polymerase chain reaction" (PCR) refers to the method of K. B. Mullis U.S. Pat. Nos. 4,683,195 and 4,683,202, which describes a method for increasing the concentration of a segment of a polynucleotide of interest in a mixture of genomic DNA without cloning or purification. This process for amplifying the polynucleotide of interest consists of introducing a large excess of two oligonucleotide primers to the DNA mixture containing the desired polynucleotide of interest, followed by a series of thermal cycling in the presence of a DNA polymerase. The two primers are complementary to their respective strands of the double-stranded polynucleotide of interest. The mixture is denatured at a higher temperature first and the primers are then annealed to complementary sequences within the polynucleotide of interest molecule. Following annealing, the primers are extended with a polymerase to form a new pair of complementary strands. The steps of denaturation, primer annealing and polymerase extension can be repeated many times (referred to as thermocycling) to obtain a high concentration of an amplified segment of the desired polynucleotide of interest. The length of the amplified segment of the desired polynucleotide of interest (amplicon) is determined by the relative positions of the primers with respect to each other, and therefore, this length is a controllable parameter. By virtue of repeating the process, the method is referred to as the "polymerase chain reaction" (hereinafter "PCR"). Because the desired amplified segments of the polynucleotide of interest become the predominant nucleic acid sequences (in terms of concentration) in the mixture, they are said to be "PCR amplified." In a modification to the method discussed above, the target nucleic acid molecules can be PCR amplified using a plurality of different primer pairs, in some cases, one or more primer pairs per target nucleic acid molecule of interest, thereby forming a multiplex PCR reaction.

As used herein, the term "primer" and its derivatives refer generally to any polynucleotide that can hybridize to a target sequence of interest. Typically, the primer functions as a substrate onto which nucleotides can be polymerized by a polymerase; in some embodiments, however, the primer can become incorporated into the synthesized nucleic acid strand and provide a site to which another primer can hybridize to prime synthesis of a new strand that is complementary to the synthesized nucleic acid molecule. The primer can include any combination of nucleotides or analogs thereof. In some embodiments, the primer is a single-stranded oligonucleotide or polynucleotide. The terms "polynucleotide" and "oligonucleotide" are used interchangeably herein to refer to a polymeric form of nucleotides of any length, and

may comprise ribonucleotides, deoxyribonucleotides, analogs thereof, or mixtures thereof. The terms should be understood to include, as equivalents, analogs of either DNA or RNA made from nucleotide analogs and to be applicable to single stranded (such as sense or antisense) and double-stranded polynucleotides. The term as used herein also encompasses cDNA, that is complementary or copy DNA produced from an RNA template, for example by the action of reverse transcriptase. This term refers only to the primary structure of the molecule. Thus, the term includes triple-, double- and single-stranded deoxyribonucleic acid ("DNA"), as well as triple-, double- and single-stranded ribonucleic acid ("RNA").

As used herein, the terms "library" and "sequencing library" refer to a collection or plurality of template molecules which share common sequences at their 5' ends and common sequences at their 3' ends. The collection of template molecules containing known common sequences at their 3' and 5' ends may also be referred to as a 3' and 5' modified library.

The term "flowcell" as used herein refers to a chamber comprising a solid surface across which one or more fluid reagents can be flowed. Examples of flowcells and related fluidic systems and detection platforms that can be readily used in the methods of the present disclosure are described, for example, in Bentley et al., *Nature* 456:53-59 (2008), WO 04/018497; US 7,057,026; WO 91/06678; WO 07/123744; US 7,329,492; US 7,211,414; US 7,315,019; US 7,405,281, and US 2008/0108082.

As used herein, the term "amplicon," when used in reference to a nucleic acid, means the product of copying the nucleic acid, wherein the product has a nucleotide sequence that is the same as or complementary to at least a portion of the nucleotide sequence of the nucleic acid. An amplicon can be produced by any of a variety of amplification methods that use the nucleic acid, or an amplicon thereof, as a template including, for example, PCR, rolling circle amplification (RCA), ligation extension, or ligation chain reaction. An amplicon can be a nucleic acid molecule having a single copy of a particular nucleotide sequence (e.g. a PCR product) or multiple copies of the nucleotide sequence (e.g. a concatameric product of RCA). A first amplicon of a target nucleic acid is typically a complimentary copy. Subsequent amplicons are copies that are created, after generation of the first amplicon, from the target nucleic acid or from the first amplicon. A subsequent amplicon can have a sequence that is substantially complementary to the target nucleic acid or substantially identical to the target nucleic acid.

As used herein, the term “array” refers to a population of sites that can be differentiated from each other according to relative location. Different molecules that are at different sites of an array can be differentiated from each other according to the locations of the sites in the array. An individual site of an array can include one or more molecules of a particular type. For example, a site can include a single target nucleic acid molecule having a particular sequence or a site can include several nucleic acid molecules having the same sequence (and/or complementary sequence, thereof). The sites of an array can be different features located on the same substrate. Exemplary features include without limitation, wells in a substrate, beads (or other particles) in or on a substrate, projections from a substrate, ridges on a substrate or channels in a substrate. The sites of an array can be separate substrates each bearing a different molecule. Different molecules attached to separate substrates can be identified according to the locations of the substrates on a surface to which the substrates are associated or according to the locations of the substrates in a liquid or gel. Exemplary arrays in which separate substrates are located on a surface include, without limitation, those having beads in wells.

The term “Next Generation Sequencing (NGS)” herein refers to sequencing methods that allow for massively parallel sequencing of clonally amplified molecules and of single nucleic acid molecules. Non-limiting examples of NGS include sequencing-by-synthesis using reversible dye terminators, and sequencing-by-ligation.

The term “sensitivity” as used herein is equal to the number of true positives divided by the sum of true positives and false negatives.

The term “specificity” as used herein is equal to the number of true negatives divided by the sum of true negatives and false positives.

The term “enrich” herein refers to the process of amplifying nucleic acids contained in a portion of a sample. Enrichment includes specific enrichment that targets specific sequences, e.g., polymorphic sequences, and non-specific enrichment that amplifies the whole genome of the DNA fragments of the sample.

As used herein, the term “each,” when used in reference to a collection of items, is intended to identify an individual item in the collection but does not necessarily refer to every item in the collection unless the context clearly dictates otherwise.

As used herein, “providing” in the context of a composition, an article, a nucleic acid, or a nucleus means making the composition, article, nucleic acid, or nucleus, purchasing the

composition, article, nucleic acid, or nucleus, or otherwise obtaining the compound, composition, article, or nucleus.

The term “and/or” means one or all of the listed elements or a combination of any two or more of the listed elements.

The words “preferred” and “preferably” refer to embodiments of the disclosure that may afford certain benefits, under certain circumstances. However, other embodiments may also be preferred, under the same or other circumstances. Furthermore, the recitation of one or more preferred embodiments does not imply that other embodiments are not useful, and is not intended to exclude other embodiments from the scope of the disclosure.

The terms “comprises” and variations thereof do not have a limiting meaning where these terms appear in the description and claims.

It is understood that wherever embodiments are described herein with the language “include,” “includes,” or “including,” and the like, otherwise analogous embodiments described in terms of “consisting of” and/or “consisting essentially of” are also provided.

Unless otherwise specified, “a,” “an,” “the,” and “at least one” are used interchangeably and mean one or more than one.

Also herein, the recitations of numerical ranges by endpoints include all numbers subsumed within that range (e.g., 1 to 5 includes 1, 1.5, 2, 2.75, 3, 3.80, 4, 5, etc.).

Reference throughout this specification to “one embodiment,” “an embodiment,” “certain embodiments,” or “some embodiments,” etc., means that a particular feature, configuration, composition, or characteristic described in connection with the embodiment is included in at least one embodiment of the disclosure. Thus, the appearances of such phrases in various places throughout this specification are not necessarily referring to the same embodiment of the disclosure. Furthermore, the particular features, configurations, compositions, or characteristics may be combined in any suitable manner in one or more embodiments.

For any method disclosed herein that includes discrete steps, the steps may be conducted in any feasible order. And, as appropriate, any combination of two or more steps may be conducted simultaneously.

The above summary of the present disclosure is not intended to describe each disclosed embodiment or every implementation of the present disclosure. The description that follows more particularly exemplifies illustrative embodiments. In several places throughout the

application, guidance is provided through lists of examples, which examples can be used in various combinations. In each instance, the recited list serves only as a representative group and should not be interpreted as an exclusive list.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1. A schematic of the relationships between placental health, maternal response, and fetal response.

Figure 2. Origins of circulating RNA (C-RNA).

Figure 3. Library prep workflow for C-RNA.

Figure 4. Validation of C-RNA approach comparing 3rd trimester pregnant and non-pregnant samples.

Figure 5. Validation of C-RNA approach using longitudinal pregnancy samples.

Figure 6. Description of clinical studies.

Figure 7. Sequencing data characteristics.

Figure 8. Classification of PE without any selection of genes, relying of entire data set.

Figure 9. Description of bootstrapping method.

Figure 10. Classification of preeclampsia samples with bootstrapping approach.

Figure 11. Examination of over-abundant preeclampsia genes.

Figure 12. Standard Adaboost Model.

Figure 13. Independent cohort allows further validation of preeclampsia signature.

Figure 14. Performance of standard adaboost model in classification of preeclampsia.

Figure 15. Classification of preeclampsia with standard DEX TREAT analysis.

Figure 16. Selection of genes and classification of preeclampsia with jackknifing approach.

Figure 17. Validation of TREAT, Bootstrapping, and Jackknifing approaches in independent PEARL biobank cohort.

Figure 18. A diagram of the bioinformatic approach to build AdaBoost Refined models.

Figure 19. Relative abundance of genes utilized by AdaBoost Refined model and their predictive capability on independent datasets.

Figure 20. Identifying C-RNA signatures specific to preeclampsia in Nextera Flex generated libraries using standard TREAT analysis and jackknifing approach.

Figure 21. Relative abundance of genes utilized by AdaBoost Refined model on Nextera Flex generated libraries and their predictive power in RGH14 dataset.

Figures 22A-22D. Validation of a clinic-friendly, whole-exome C-RNA analysis method. Fig. 22A is a schematic of the sequencing library preparation method; all steps after blood collection can be performed in a centralized processing lab. Temporal changes of transcripts altered throughout the course of pregnancy (Fig. 22B). Overlap of genes identified in C-RNA pregnancy progression studies (Fig. 22C). Tissues expressing the 91 genes unique to the pregnancy time course study (Fig. 22D).

Figures 23A-23C. Sample collection for PE clinical studies. Panels illustrate the time of blood collection (triangles) and gestational age at birth (squares) for each individual in the iPC study (Fig. 23A) and the PEARL study (Fig. 23B). The red line indicates the threshold for term birth. Preterm birth rates are significantly elevated in early-onset PE cohorts (Fig. 23C). *** $p < 0.001$ by Fisher's exact test.

Figures 24A-24G. Differential analysis of C-RNA identifies preeclampsia biomarkers. Fold change and abundance of transcripts altered in PE (Fig. 24A). One-sided confidence p-value intervals were calculated after jackknifing for each gene detected by standard analysis methods (Fig. 24B). Transcript abundance fold-change determined by whole exome sequencing and by qPCR for (21) genes (Fig. 24C). * $p < 0.05$ by Student's T-test. Tissue distribution of affected genes (Fig. 24D). Hierarchical clustering of iPC samples (average linkage, squared Euclidean distance) (Fig. 24E). Clustering of early-onset PE (Fig 24F) and late-onset PE (Fig. 24G) samples from the PEARL study.

Figures 25A-25E. AdaBoost classifies preeclampsia samples across cohorts. Heatmap illustrating the relative abundance of the transcripts used by machine learning in each cohort (Fig. 25A). The height of each block reflects each gene's importance. ROC curves for each dataset (Fig. 25B). Distributions (KDE) of AdaBoost Scores. The orange line indicates the optimal boundary to discriminate PE and control samples (Fig. 25C). Concordance of genes identified by differential analysis and those used in AdaBoost (Fig. 25D). Tissue distribution of AdaBoost genes (Fig 25E).

Figures 26A-26C. C-RNA data integrity when blood is stored in different collection tubes. Comparing the abundance of previously detected C-RNA pregnancy markers from blood stored overnight in different tube types to immediate processing after collection in EDTA tubes

(Fig. 26A). Scatterplots comparing transcript FPKM values for C-RNA prepared from the same individual after different blood storage durations (Fig. 26B). Pearson's correlation coefficient, R, is more variable when using EDTA tubes (cf, Cell-Free) (Fig. 26C).

Figures 27A and 27B. The effect of plasma volume on C-RNA data quality. A meta-analysis was performed with data from nine independent studies to determine the appropriate plasma input for the protocol. Noise (biological coefficient of variation, EdgeR) was calculated from biological replicates within each study (Fig. 27A). Library complexity (bound population, Preseq) was calculated for each sample (Fig. 27B). ** $p < 0.01$, *** $p < 0.001$ by ANOVA with Tukey's HSD correction, with study as a blocking variable.

Figures 28A-28C. Pregnancy marker tissue specificity. Pie charts showing tissue specificity of the genes detected in pregnancy by three independent studies, using either the full set of altered genes (Fig. 28A), the transcripts unique to each study (Fig. 28B), or intersecting gene sets (Fig. 28C).

Figures 29A-29E. Jackknifing excludes genes that are not universally altered in preeclampsia. Schematic of the jackknifing approach used to determine how consistently transcripts were altered across PE samples (Fig. 29A). Average abundance and noise for each differentially abundant gene (Fig. 29B). ROC area under the curve values for each affected transcript provide a measure of how separated C-RNA transcript abundance distributions are for control and PE samples (Fig. 29C). * $p < 0.05$ by Mann-Whitney U test. Hierarchical clustering of iPC samples using the genes excluded after jackknifing (Fig. 29D). Tissue distribution of excluded transcripts (Fig. 29E). The decreased contribution of the fetus and placenta may suggest the maternal component of PE is most variable between individuals.

Figures 30A-30D. AdaBoost model development strategy. The RGH014 dataset was divided into 6 pieces (Fig. 30A). The "Holdout Subset" contained 10% of the samples (randomly selected) as well as the 3 samples which were incorrectly clustered when using differentially abundant genes (as with Fig. 24C) and was fully excluded from model building. The remaining samples were divided at random into 5 evenly sized "Test Subsets." For each test subset, training data was composed of all non-holdout and non-testing samples. Gene counts for training and testing data were TMM-normalized in edgeR, and then standardized to mean 0 and standard deviation 1 for each gene. For each train/test sample set, AdaBoost models (90 estimators, 1.6 learning rate) were built 10 times from the training data (Fig. 30B). Feature

pruning was performed, removing genes below an incrementally increasing importance threshold and assessing performance by Matthew's correlation coefficient when predicting testing data. The model with the best performance – and fewest genes, in the case of a tie – was retained. Estimators from all 50 independent models were combined into a single AdaBoost model (Fig. 30C). Feature pruning was performed on the resulting ensemble, this time using the percent of models which used a gene to set threshold values and performance measured by average log loss value across test subsets. ROC curve after applying the final AdaBoost model to the holdout data (Fig. 30D). All samples segregated correctly, except for two of the three samples which also misclustered by HCA.

Figures 31A-31E. The effect of hyperparameter selection and feature pruning on machine learning performance. Heatmap of a grid search to identify the optimal hyperparameters for AdaBoost (Fig. 31A). Matthew's correlation coefficient was used as a measure of performance. Flattened views of performance for each hyperparameter (Fig. 31B). Arrows indicate the values selected for model construction. Fig. 31C shows the impact of pruning individual AdaBoost models on performance (as in Fig. 30B). Solid lines are the average for all 10 models, and the shaded region shows the standard deviation. The number of AdaBoost models using each gene observed in the pre-pruned ensemble (Fig. 31D). Model performance when pruning the combined AdaBoost ensemble (Fig. 31E). The orange lines in Fig. 31D and Fig. 31E show the threshold applied to generate the final AdaBoost model.

The schematic drawings are not necessarily to scale. Like numbers used in the figures may refer to like components. However, it will be understood that the use of a number to refer to a component in a given figure is not intended to limit the component in another figure labeled with the same number. In addition, the use of different numbers to refer to components is not intended to indicate that the different numbered components cannot be the same or similar to other numbered components.

DETAILED DESCRIPTION

Provided herein are signatures of circulating RNA found in the maternal circulation that are specific to preeclampsia and the use of such signatures in noninvasive methods for the diagnosis of preeclampsia and the identification of pregnant women at risk for developing preeclampsia.

While most of the DNA and RNA in the body is located within cells, extracellular nucleic acids can also be found circulating freely in the blood. Circulating RNA, also referred to herein as "C-RNA," refers to extracellular segments of RNA found in the bloodstream. C-RNA molecules originate predominately from two sources: one, released into the circulation from dying cells undergoing apoptosis, and two, contained within exosomes shed by living cells into the circulation. Exosomes are small membranous vesicles about 30–150 nm of diameter released from many cell types into the extracellular space and are found in a wide variety of body fluids, including serum, urine, and breast milk and carrying protein, mRNA, and microRNA. The lipid bilayer structure of exosomes protects the RNAs contained within from degradation by RNases, providing for stability in blood. See, for example, Huang et al., 2013, *BMC Genomics*; 14:319; And Li et al., 2017, *Mol Cancer*; 16:145). Evidence is accumulating that exosomes have specialized functions and play a role in such processes as coagulation, intercellular signaling, and waste management (van der Pol et al., 2012, *Pharmacol Rev*; 64(3):676–705). See, also, Samos et al., 2006, *Ann N Y Acad Sci*; 1075:165–173; Zernecke et al., 2009, *Sci Signal*; 2:ra81; Ma et al., 2012, *J Exp Clin Cancer Res*; 31:38; and Sato-Kuwabara et al., 2015, *Int J Oncol*; 46:17–27.

With the methods described herein, the C-RNA molecules found in maternal circulation function as biomarkers of fetal, placental, and maternal health and provide a window into the progression of pregnancy. Described herein are C-RNA signatures within the maternal circulation that are indicative of pregnancy, C-RNA signatures within the maternal circulation that are temporally associated with the gestational stage of pregnancy, and C-RNA signatures within the maternal circulation that are indicative of the pregnancy complication preeclampsia.

A C-RNA signature within the maternal circulation indicative of preeclampsia includes a plurality of C-RNA molecules encoding at least a portion of a plurality of proteins selected from ARRDC2, JUN, SKIL, ATP13A3, PDE8B, GSTA3, PAPPA2, TIPARP, LEP, RGP1, USP54, CLEC4C, MRPS35, ARHGEF25, CUX2, HEATR9, FSTL3, DDI2, ZMYM6, ST6GALNAC3,

GBP2, NES, ETV3, ADAM17, ATOH8, SLC4A3, TRAF3IP1, TTC21A, HEG1, ASTE1, TMEM108, ENC1, SCAMP1, ARRDC3, SLC26A2, SLIT3, CLIC5, TNFRSF21, PPP1R17, TPST1, GATSL2, SPDYE5, HIPK2, MTRNR2L6, CLCN1, GINS4, CRH, C10orf2, TRUB1, PRG2, ACY3, FAR2, CD63, CKAP4, TPCN1, RNF6, THTPA, FOS, PARN, ORAI3, ELMO3, SMPD3, SERPINF1, TMEM11, PSMD11, EBI3, CLEC4M, CCDC151, CPAMD8, CNFN, LILRA4, ADA, C22orf39, PI4KAP1, and ARFGAP3. This C-RNA signature is the Adaboost General signature obtained with the TruSeq library prep method shown in Table 1 below, also referred to herein as "list (a)" or "(a)."

A C-RNA signature within the maternal circulation indicative of preeclampsia includes a plurality of C-RNA molecules encoding at least a portion of a plurality of proteins selected from TIMP4, FLG, HTRA4, AMPH, LCN6, CRH, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, CLEC4C, KRT5, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, LEP, NES, and VSIG4. This C-RNA signature is the Bootstrapping signature obtained with the TruSeq library prep method shown in Table 1 below, also referred to herein as "list (b)" or "(b)."

A C-RNA signature within the maternal circulation indicative of preeclampsia includes a plurality of C-RNA molecules encoding at least a portion of a protein selected from CYP26B1, IRF6, MYH14, PODXL, PPP1R3C, SH3RF2, TMC7, ZNF366, ADCY1, C6, FAM219A, HAO2, IGIP, IL1R2, NTRK2, SH3PXD2A, SSUH2, SULT2A1, FMO3, FSTL3, GATA5, HTRA1, C8B, H19, MN1, NFE2L1, PRDM16, AP3B2, EMP1, FLNC, STAG3, CPB2, TENC1, RP1L1, A1CF, NPR1, TEK, ERRFI1, ARHGEF15, CD34, RSPO3, ALPK3, SAMD4A, ZCCHC24, LEAP2, MYL2, NRG3, ZBTB16, SERPINA3, AQP7, SRPX, UACA, ANO1, FKBP5, SCN5A, PTPN21, CACNA1C, ERG, SOX17, WWTR1, AIF1L, CA3, HRG, TAT, AQP7P1, ADRA2C, SYNPO, FN1, GPR116, KRT17, AZGP1, BCL6B, KIF1C, CLIC5, GPR4, GJA5, OLAH, C14orf37, ZEB1, JAG2, KIF26A, APOLD1, PNMT, MYOM3, PITPNM3, TIMP4, HTRA4, AMPH, LCN6, CRH, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, LEP, NES, VSIG4, HBG2, CADM2, LAMP5, PTGDR2, NOMO1, NXF3, PLD4, BPIFB3, PACSIN1, CUX2, FLG, CLEC4C, and KRT5. This C-RNA signature is the Standard DEX Treat signature obtained with the TruSeq library prep method shown in Table 1 below, also referred to herein as "list (c)" or "(c)."

A C-RNA signature within the maternal circulation indicative of preeclampsia includes a plurality of C-RNA molecules encoding at least a portion of a protein selected from VSIG4, ADAMTS2, NES, FAM107A, LEP, DAAM2, ARHGEF25, TIMP3, PRX, ALOX15B, HSPA12B, IGFBP5, CLEC4C, SLC9A3R2, ADAMTS1, SEMA3G, KRT5, AMPH, PRG2, PAPPA2, TEAD4, CRH, PITPNM3, TIMP4, PNMT, ZEB1, APOLD1, PLD4, CUX2, and HTRA4. This C-RNA signature is the Jacknifing signature obtained with the TruSeq library prep method shown in Table 1 below, also referred to herein as "list (d)" or "(d)."

A C-RNA signature within the maternal circulation indicative of preeclampsia includes a plurality of C-RNA molecules encoding at least a portion of a protein selected from ADAMTS1, ADAMTS2, ALOX15B, AMPH, ARHGEF25, CELF4, DAAM2, FAM107A, HSPA12B, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LEP, LRRC26, NES, OLAH, PACSIN1, PAPPA2, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, and VSIG4. This C-RNA signature is the Standard DEX Treat signature obtained with the Nextera Flex for Enrichment library prep method shown in Table 1 below, also referred to herein as "list (e)" or "(e)."

A C-RNA signature within the maternal circulation indicative of preeclampsia includes a plurality of C-RNA molecules encoding at least a portion of a protein selected from ADAMTS1, ADAMTS2, ALOX15B, ARHGEF25, CELF4, DAAM2, FAM107A, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LEP, LRRC26, NES, OLAH, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, and VSIG4. This C-RNA signature is the Jacknifing signature obtained with the Nextera Flex for Enrichment library prep method shown in Table 1 below, also referred to herein as "list (f)" or "(f)."

A C-RNA signature within the maternal circulation indicative of preeclampsia includes a plurality of C-RNA molecules encoding at least a portion of a protein selected from CLEC4C, ARHGEF25, ADAMTS2, LEP, ARRDC2, SKIL, PAPPA2, VSIG4, ARRDC4, CRH, and NES. This C-RNA signature is the Adaboost Refined TruSeq signature obtained with the TruSeq library prep method shown in Table 1 below, also referred to herein as "AdaBoost Refined 1," "list (g)," or "(g)."

In some embodiments, a C-RNA signature within the maternal circulation indicative of preeclampsia includes C-RNA molecules encoding at least a portion of a protein selected from ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, and VSIG4 (also referred to herein as "AdaBoost Refined 2"), ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP,

PAPPA2, SKIL, and VSIG4 (also referred to herein as "AdaBoost Refined 3"), ADAMTS2, ARHGEF25, ARRDC4, CLEC4C, LEP, NES, SKIL, and VSIG4 (also referred to herein as "AdaBoost Refined 4"), ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, CRH, LEP, PAPPA2, SKIL, and VSIG4 (also referred to herein as "AdaBoost Refined 5"), ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, and SKIL (also referred to herein as "AdaBoost Refined 6"), or ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, LEP, PAPPA2, and SKIL (also referred to herein as "AdaBoost Refined 7").

A C-RNA signature within the maternal circulation indicative of preeclampsia includes a plurality of C-RNA molecules encoding at least a portion of a protein selected from LEP, PAPPA2, KCNA5, ADAMTS2, MYOM3, ATP13A3, ARHGEF25, ADA, HTRA4, NES, CRH, ACY3, PLD4, SCT, NOX4, PACSIN1, SERPINF1, SKIL, SEMA3G, TIPARP, LRRC26, PHEX, LILRA4, and PER1. This C-RNA signature is the Adaboost Refined Nextera Flex signature obtained with the Nextera Flex for Enrichment library prep method shown in Table 1 below, also referred to herein as "list (h)" or "(h)."

In some embodiments, a C-RNA signature within the maternal circulation indicative of preeclampsia includes a plurality of C-RNA molecules encoding at least a portion of a protein selected from of any one or more of any of (a), (b), (c), (d), (e), (f), (g), and/or (h) in combination with any one or more of any of (a), (b), (c), (d), (e), (f), (g), and/or (h).

The Examples provided herewith describe the eight gene lists summarized above that distinguish preeclampsia and control pregnancies. Each was identified by using different analysis methods and/or distinct datasets. However, there is a high degree of concordance between many of these gene sets. Identifying a transcript as altered in preeclampsia C-RNA with multiple approaches indicates that said transcript has higher predictive value for classification of this disease. Thus, the importance of the transcripts identified by all differential expression analyses and by all AdaBoost models was combined and ranked. Genes assigned lower ranks are not unimportant or uninformative, but they may be less robust for classification of preeclampsia across cohorts and sample preparations.

First, the transcripts identified when using all differential expression analyses (Standard DEX Treat, bootstrapping and the jackknifing) for both library preparation methods (TruSeq and Nextera Flex for Enrichment) were combined. Table 2 below shows the relative importance for all of the 125 transcripts identified by the different analysis methods. Transcripts identified

across every analysis method and both library preparations are the strongest classifiers and assigned an importance ranking of 1. Transcripts that were identified by three or more analysis methods and were detected with both library preparations were given an importance ranking of 2. Transcripts identified by the most stringent analysis method, jackknifing but only one of the library preparations were assigned an importance ranking of 3. Transcripts identified in two of the five analysis methods were given an importance ranking of 4. Transcripts that were only identified in the Standard DEX Treat method, the most broad and inclusive analysis, were given the lowest importance ranking of 5.

Then, the 91 transcripts identified across all AdaBoost models (AdaBoost General and AdaBoost Refined) and both library preparations (Table 3 below) were combined. When generating the refined AdaBoost models for each library preparation, observed slight variations had been observed in the gene set obtained each time a model was built from the same data. This is a natural result of randomness used by AdaBoost to search through the large whole-exome C-RNA data. To obtain a representative list of genes, model building for refined AdaBoost was run a minimum of nine separate times and all genes used by one or more models reported. The percent of models that included each transcript are reported in Table 3 (Frequency Used By AdaBoost). AdaBoost assigns its own "importance" value to each transcript, which reflects how much the abundance of that transcript influences determining whether a sample is from a preeclampsia patient. These AdaBoost importance values were averaged across each refined AdaBoost model a given transcript was used by (Table 3, Average AdaBoost Model Importance).

Transcripts identified across all AdaBoost analyses and library preparations were assigned the highest importance ranking of 1. Transcripts identified in the refined AdaBoost model for a single library preparation method with over 90% frequency used by AdaBoost were assigned an importance ranking of 2. Generally, these transcripts also have higher AdaBoost model importance, consistent with increased predictive capabilities. Transcripts identified in the refined AdaBoost model for a single library preparation method but used by less than 90% of AdaBoost models were assigned an importance ranking of 3. Transcripts identified only in the general AdaBoost model for a single library preparation were given the lowest importance ranking of 4.

Table 2 lists every gene identified by DEX analysis across all analysis approaches and library preps. Rank 1 = Transcript identified across every analysis method and library prep method. Rank 2 = Transcript identified both library preps, and 3/5 analysis methods. Rank 3= Identified in one library prep method, in jacknifing, are most stringent analysis. Rank 4 = identified in 2/5 analyses. And, Rank 5 = Only identified in Standard DEX Treat method, our most relaxed analysis method.

Table 3 lists every gene identified by Adaboost analysis across both library preps. Rank 1 = identified in both library prep methods and the refined adaboost models. Rank 2 = Identified in one library prep method, present in refined adaboost model with high model importance and frequency. Rank 3 = identified in one library prep method, present in refined adaboost model with medium model importance and frequency. And, Rank 4 = Identified in one library prep, not present in the refined adaboost model.

Table 4 below is a glossary of all of the various genes recited herein. The information was obtained from the HUGO Gene Nomenclature Committee at the European Bioinformatics Institute.

Table 1: Composite Gene Listing

DIFFERENTIAL EXPRESSION APPROACHES					Machine Learning Approaches		
Standard DEX TREAT	Bootstrapping	Jackknifing	Standard DEX TREAT	Jackknifing	Adaboost General	Adaboost Refined TruSeq	Adaboost Refined Nextera Flex
Library Prep: TruSeq	Library Prep: TruSeq	Library Prep: TruSeq	Library Prep: Nextera Flex for Enrichment	Library Prep: Nextera Flex for Enrichment	Library Prep: TruSeq	Library Prep: TruSeq	Library Prep: Nextera Flex for Enrichment
Broadest DEX list for TST170	Refined subset of Standard DEX TREAT	Refined subset of Standard DEX TREAT	Broadest DEX list for Nextera	Refined subset of Standard DEX TREAT for Nextera	Broadest adaboost	Imrpoved model building for Adaboost, to improve unverisal signal	Imrpoved model building for Adaboost, to improve unverisal signal
122 genes	27 genes	30 genes	26 genes	22 genes	75 genes	11 genes	24 genes
FLG	FLG	VSIG4	ADAMTS1	ADAMTS1	ARHGEF25	CLEC4C	LEP
KRT5	KRT5	ADAMTS2	ADAMTS2	ADAMTS2	CLEC4C	ARHGEF25	PAPPA2
HBG2	CLEC4C	NES	ALOX15B	ALOX15B	CRH	ADAMTS2	KCNA5
NXF3	TEAD4	FAM107A	AMPH	ARHGEF25	CUX2	LEP	ADAMTS2
CLEC4C	SEMA3G	LEP	ARHGEF25	CELF4	LEP	ARRDC2	MYOM3
BPIFB3	ADAMTS1	DAAM2	CELF4	DAAM2	NES	SKIL	ATP13A3
LAMP5	IGFBP5	ARHGEF25	DAAM2	FAM107A	PAPPA2	PAPPA2	ARHGEF25
CADM2	HSPA12B	TIMP3	FAM107A	HTRA4	PRG2	VSIG4	ADA
CUX2	SLC9A3R2	PRX	HSPA12B	IGFBP5	ACY3	ARRDC4	HTRA4
PACSIN1	PRX	ALOX15B	HTRA4	KCNA5	ADA	CRH	NES
PTGDR2	TIMP3	HSPA12B	IGFBP5	KRT5	ADAM17	NES	CRH
PLD4	ARHGEF25	IGFBP5	KCNA5	LCN6	ARFGAP3		ACY3
NOMO1	HTRA4	CLEC4C	KRT5	LEP	ARRDC2		PLD4
SH3RF2	NES	SLC9A3R2	LCN6	LRRC26	ARRDC3		SCT
ZNF366	TIMP4	ADAMTS1	LEP	NES	ASTE1		NOX4
SH3PXD2A	PAPPA2	SEMA3G	LRRC26	OLAH	ATOH8		PACSIN1
SULT2A1	FAM107A	KRT5	NES	PRX	ATP13A3		SERPINF1
FAM219A	PRG2	AMPH	OLAH	PTGDR2	C10orf2		SKIL
PPP1R3C	AMPH	PRG2	PACSIN1	SEMA3G	C22orf39		SEMA3G
NFE2L1	DAAM2	PAPPA2	PAPPA2	SLC9A3R2	CCDC151		TIPARP
PODXL	LCN6	TEAD4	PRX	TIMP3	CD63		LRRC26
HTRA1	ALOX15B	**CRH**	PTGDR2	VSIG4	CKAP4		PHEX
EMP1	CRH	PITPNM3	SEMA3G		CLCN1		LILRA4
H19	VSIG4	TIMP4	SLC9A3R2		CLEC4M		PER1
IGIP	LEP	PNMT	TIMP3		CLIC5		
SSUH2	ADAMTS2	ZEB1	VSIG4		CNFN		
C6	ARMS2	APOLD1			CPAMD8		
ARHGEF15		PLD4			DDI2		
IRF6		CUX2			EBI3		
NPR1		HTRA4			ELMO3		
ALPK3					ENC1		
ZCCHC24					ETV3		
SAMD4A					FAR2		
STAG3					FOS		
RP1L1					FSTL3		
A1CF					GATSL2		
MN1					GBP2		
CD34					GINS4		
MYH14					GSTA3		
TENC1					HEATR9		
FSTL3					HEG1		
PRDM16					HIPK2		
FMO3					JUN		
UACA					LILRA4		

TEK					MRPS35		
SOX17					MTRNR2L6		
FLNC					ORAI3		
TMC7					PARN		
KIF1C					PDE8B		
CLIC5					PI4KAP1		
SYNPO					PPP1R17		
CACNA1C					PSMD11		
ERG					RGP1		
PTPN21					RNF6		
NTRK2					SCAMP1		
WWTR1					SERPINF1		
CYP26B1					SKIL		
ZEB1					SLC26A2		
AIF1L					SLC4A3		
C8B					SLIT3		
KIF26A					SMPD3		
ZBTB16					SPDYE5		
BCL6B					ST6GALNAC3		
FKBP5					THTPA		
FN1					TIPARP		
AQP7					TMEM108		
IL1R2					TMEM11		
ERRFI1					TNFRSF21		
SRPX					TPCN1		
GJA5					TPST1		
GPR116					TRAF3IP1		
JAG2					TRUB1		
MYL2					TTC21A		
ADCY1					USP54		
NRG3					ZMYM6		
GPR4							
PITPNM3							
SERPINA3							
CPB2							
ADRA2C							
ANO1							
CA3							
C14orf37							
TEAD4							
TAT							
LEAP2							
HAO2							
SEMA3G							
ADAMTS1							
APOLD1							
IGFBP5							
HSPA12B							
GATA5							
SLC9A3R2							
RSPO3							
AQP7P1							
PRX							
PNMT							
MYOM3							
HRG							
TIMP3							
ARHGEF25							
HTRA4							
SCN5A							
OLAH							

NES							
TIMP4							
PAPPA2							
AZGP1							
FAM107A							
PRG2							
AMPH							
AP3B2							
KRT17							
DAAM2							
LCN6							
ALOX15B							
CRH							
VSIG4							
LEP							
ADAMTS2							
ARMS2							

Table 2: DEX Analysis

		TruSeq				Nextera Flex for enrichment		
Gene	Importance Ranking	Standard DEX TREAT	Fold Change	Bootstrapping	Jackknifing	Standard DEX	Fold Change	Jackknifing
ADAMTS1	1	Y	1.79	Y	Y	Y	+3.2	Y
ADAMTS2	1	Y	3.61	Y	Y	Y	+12.2	Y
ALOX15B	1	Y	2.51	Y	Y	Y	+5.3	Y
ARHGEF25	1	Y	2.02	Y	Y	Y	+3.8	Y
DAAM2	1	Y	2.48	Y	Y	Y	+5.4	Y
FAM107A	1	Y	2.31	Y	Y	Y	+4.3	Y
HTRA4	1	Y	2.03	Y	Y	Y	+4.0	Y
IGFBP5	1	Y	1.81	Y	Y	Y	+3.4	Y
KRT5	1	Y	-2.52	Y	Y	Y	-4.8	Y
LEP	1	Y	3.48	Y	Y	Y	+8.1	Y
NES	1	Y	2.15	Y	Y	Y	+4.2	Y
PRX	1	Y	1.93	Y	Y	Y	+3.3	Y
SEMA3G	1	Y	1.78	Y	Y	Y	+3.5	Y
SLC9A3R2	1	Y	1.85	Y	Y	Y	+3.4	Y
TIMP3	1	Y	2.01	Y	Y	Y	+3.7	Y
VSIG4	1	Y	3.03	Y	Y	Y	+8.2	Y
PAPPA2	2	Y	2.20	Y	Y	Y	+4.2	N
AMPH	2	Y	2.37	Y	Y	Y	+4.1	N
HSPA12B	2	Y	1.82	Y	Y	Y	+3.2	N
PTGDR2	2	Y	-1.67	N	N	Y	-3.5	Y
LCN6	2	Y	2.49	Y	N	Y	+4.4	Y
OLAH	2	Y	2.07	N	N	Y	+5.2	Y
APOLD1	3	Y	1.80	N	Y	N	NA	N
CUX2	3	Y	-1.73	N	Y	N	NA	N
PITPNM3	3	Y	1.66	N	Y	N	NA	N
PLD4	3	Y	-1.59	N	Y	N	NA	N
PNMT	3	Y	1.98	N	Y	N	NA	N
CLEC4C	3	Y	-1.86	Y	Y	N	NA	N
CRH	3	Y	2.54	Y	Y	N	NA	N
PRG2	3	Y	2.36	Y	Y	N	NA	N
TEAD4	3	Y	1.74	Y	Y	N	NA	N
TIMP4	3	Y	2.17	Y	Y	N	NA	N
CELF4	3	N	NA	N	N	Y	+5.3	Y
KCNA5	3	N	NA	N	N	Y	-4.0	Y
LRRC26	3	N	NA	N	N	Y	-4.4	Y
ARMS2	4	Y	4.43	Y	N	N	NA	N
FLG	4	Y	-3.05	Y	N	N	NA	N
PACSIN1	4	Y	-1.70	N	N	Y	-3.4	N
A1CF	5	Y	1.37	N	N	N	NA	N
ADCY1	5	Y	1.62	N	N	N	NA	N
ADRA2C	5	Y	1.69	N	N	N	NA	N
AIF1L	5	Y	1.48	N	N	N	NA	N
ALPK3	5	Y	1.35	N	N	N	NA	N
ANO1	5	Y	1.69	N	N	N	NA	N
AP3B2	5	Y	2.40	N	N	N	NA	N
AQP7	5	Y	1.51	N	N	N	NA	N
AQP7P1	5	Y	1.88	N	N	N	NA	N
ARHGEF15	5	Y	1.33	N	N	N	NA	N
AZGP1	5	Y	2.27	N	N	N	NA	N
BCL6B	5	Y	1.50	N	N	N	NA	N
BPIFB3	5	Y	-1.80	N	N	N	NA	N
C14orf37	5	Y	1.73	N	N	N	NA	N
C6	5	Y	1.33	N	N	N	NA	N
C8B	5	Y	1.49	N	N	N	NA	N
CA3	5	Y	1.72	N	N	N	NA	N

CACNA1C	5	Y	1.42	N	N	N	NA	N
CADM2	5	Y	-1.76	N	N	N	NA	N
CD34	5	Y	1.37	N	N	N	NA	N
CLIC5	5	Y	1.41	N	N	N	NA	N
CPB2	5	Y	1.69	N	N	N	NA	N
CYP26B1	5	Y	1.48	N	N	N	NA	N
EMP1	5	Y	1.30	N	N	N	NA	N
ERG	5	Y	1.43	N	N	N	NA	N
ERRFI1	5	Y	1.54	N	N	N	NA	N
FAM219A	5	Y	1.24	N	N	N	NA	N
FKBP5	5	Y	1.50	N	N	N	NA	N
FLNC	5	Y	1.40	N	N	N	NA	N
FMO3	5	Y	1.39	N	N	N	NA	N
FN1	5	Y	1.51	N	N	N	NA	N
FSTL3	5	Y	1.38	N	N	N	NA	N
GATA5	5	Y	1.82	N	N	N	NA	N
GJA5	5	Y	1.55	N	N	N	NA	N
GPR116	5	Y	1.56	N	N	N	NA	N
GPR4	5	Y	1.65	N	N	N	NA	N
H19	5	Y	1.32	N	N	N	NA	N
HAO2	5	Y	1.75	N	N	N	NA	N
HBG2	5	Y	-2.15	N	N	N	NA	N
HRG	5	Y	1.99	N	N	N	NA	N
HTRA1	5	Y	1.29	N	N	N	NA	N
IGIP	5	Y	1.32	N	N	N	NA	N
IL1R2	5	Y	1.53	N	N	N	NA	N
IRF6	5	Y	1.34	N	N	N	NA	N
JAG2	5	Y	1.57	N	N	N	NA	N
KIF1C	5	Y	1.41	N	N	N	NA	N
KIF26A	5	Y	1.49	N	N	N	NA	N
KRT17	5	Y	2.47	N	N	N	NA	N
LAMP5	5	Y	-1.77	N	N	N	NA	N
LEAP2	5	Y	1.74	N	N	N	NA	N
MN1	5	Y	1.37	N	N	N	NA	N
MYH14	5	Y	1.38	N	N	N	NA	N
MYL2	5	Y	1.60	N	N	N	NA	N
MYOM3	5	Y	1.99	N	N	N	NA	N
NFE2L1	5	Y	1.26	N	N	N	NA	N
NOMO1	5	Y	-1.49	N	N	N	NA	N
NPR1	5	Y	1.34	N	N	N	NA	N
NRG3	5	Y	1.62	N	N	N	NA	N
NTRK2	5	Y	1.45	N	N	N	NA	N
NXF3	5	Y	-1.96	N	N	N	NA	N
PODXL	5	Y	1.27	N	N	N	NA	N
PPP1R3C	5	Y	1.25	N	N	N	NA	N
PRDM16	5	Y	1.38	N	N	N	NA	N
PTPN21	5	Y	1.44	N	N	N	NA	N
RP1L1	5	Y	1.36	N	N	N	NA	N
RSPO3	5	Y	1.87	N	N	N	NA	N
SAMD4A	5	Y	1.35	N	N	N	NA	N
SCN5A	5	Y	2.03	N	N	N	NA	N
SERPINA3	5	Y	1.67	N	N	N	NA	N
SH3PXD2A	5	Y	1.23	N	N	N	NA	N
SH3RF2	5	Y	1.18	N	N	N	NA	N
SOX17	5	Y	1.40	N	N	N	NA	N
SRPX	5	Y	1.55	N	N	N	NA	N
SSUH2	5	Y	1.33	N	N	N	NA	N
STAG3	5	Y	1.36	N	N	N	NA	N
SULT2A1	5	Y	1.23	N	N	N	NA	N

SYNPO	5	Y	1.42	N	N	N	NA	N
TAT	5	Y	1.74	N	N	N	NA	N
TEK	5	Y	1.39	N	N	N	NA	N
TENC1	5	Y	1.38	N	N	N	NA	N
TMC7	5	Y	1.40	N	N	N	NA	N
UACA	5	Y	1.39	N	N	N	NA	N
WWTR1	5	Y	1.47	N	N	N	NA	N
ZBTB16	5	Y	1.50	N	N	N	NA	N
ZCCHC24	5	Y	1.35	N	N	N	NA	N
ZEB1	5	Y	1.48	N	Y	N	NA	N
ZNF366	5	Y	1.23	N	N	N	NA	N

Table 3: Adaboost Analysis

	Importance Ranking	Fold Change in Preeclampsia		Frequency Used By Adaboost		Average AdaBoost Model Importance	
		TruSeq	Nextera Flex	TruSeq	Nextera Flex	TruSeq	Nextera Flex
ADAMTS2	1	+11.6	+12.2	100%	100%	9%	8%
ARHGEF25	1	+4.1	+3.8	100%	100%	11%	5%
CRH	1	+5.7	+3.9	14%	100%	2%	4%
LEP	1	+10.7	+8.1	100%	100%	8%	17%
NES	1	+4.5	+4.2	7%	100%	4%	4%
PAPPA2	1	+4.9	+4.2	64%	100%	3%	8%
SKIL	1	+1.5	+1.4	86%	78%	3%	3%
ACY3	2	ND	-2.3	ND	100%	ND	3%
ADA	2	ND	-1.6	ND	100%	ND	5%
ARRDC2	2	+1.8	ND	93%	ND	3%	ND
ATP13A3	2	ND	+1.5	ND	100%	ND	5%
CLEC4C	2	-3.6	ND	100%	ND	18%	ND
HTRA4	2	ND	+4.0	ND	100%	ND	5%
KCNA5	2	ND	-4.0	ND	100%	ND	8%
MYOM3	2	ND	+4.2	ND	100%	ND	7%
NOX4	2	ND	-1.8	ND	100%	ND	2%
PACSIN1	2	ND	-3.4	ND	100%	ND	2%
PLD4	2	ND	-2.7	ND	100%	ND	3%
SCT	2	ND	-3.3	ND	100%	ND	3%
SERPINF1	2	ND	-1.6	ND	100%	ND	2%
VSIG4	3	+8.1	ND	43%	ND	3%	ND
ARRDC4	3	+2.0	ND	36%	ND	4%	ND
LILRA4	3	ND	-2.7	ND	33%	ND	1%
LRRC26	3	ND	-4.4	ND	56%	ND	2%
PER1	3	ND	+2.2	ND	33%	ND	1%
PHEX	3	ND	-2.2	ND	56%	ND	2%
SEMA3G	3	ND	+3.5	ND	67%	ND	5%
TIPARP	3	ND	+1.2	ND	67%	ND	2%
ADAM17	4	ND	ND	ND	ND	ND	ND
ARFGAP3	4	ND	ND	ND	ND	ND	ND
ARRDC3	4	ND	ND	ND	ND	ND	ND
ASTE1	4	ND	ND	ND	ND	ND	ND
ATOH8	4	ND	ND	ND	ND	ND	ND
C10orf2	4	ND	ND	ND	ND	ND	ND
C22orf39	4	ND	ND	ND	ND	ND	ND
CCDC151	4	ND	ND	ND	ND	ND	ND
CD63	4	ND	ND	ND	ND	ND	ND
CKAP4	4	ND	ND	ND	ND	ND	ND
CLCN1	4	ND	ND	ND	ND	ND	ND
CLEC4M	4	ND	ND	ND	ND	ND	ND
CLIC5	4	ND	ND	ND	ND	ND	ND
CNFN	4	ND	ND	ND	ND	ND	ND
CPAMD8	4	ND	ND	ND	ND	ND	ND
CUX2	4	ND	ND	ND	ND	ND	ND

DDI2	4	ND	ND	ND	ND	ND	ND
EBI3	4	ND	ND	ND	ND	ND	ND
ELMO3	4	ND	ND	ND	ND	ND	ND
ENC1	4	ND	ND	ND	ND	ND	ND
ETV3	4	ND	ND	ND	ND	ND	ND
FAR2	4	ND	ND	ND	ND	ND	ND
FOS	4	ND	ND	ND	ND	ND	ND
FSTL3	4	ND	ND	ND	ND	ND	ND
GATSL2	4	ND	ND	ND	ND	ND	ND
GBP2	4	ND	ND	ND	ND	ND	ND
GINS4	4	ND	ND	ND	ND	ND	ND
GSTA3	4	ND	ND	ND	ND	ND	ND
HEATR9	4	ND	ND	ND	ND	ND	ND
HEG1	4	ND	ND	ND	ND	ND	ND
HIPK2	4	ND	ND	ND	ND	ND	ND
JUN	4	ND	ND	ND	ND	ND	ND
MRPS35	4	ND	ND	ND	ND	ND	ND
MTRNR2L6	4	ND	ND	ND	ND	ND	ND
ORAI3	4	ND	ND	ND	ND	ND	ND
PARN	4	ND	ND	ND	ND	ND	ND
PDE8B	4	ND	ND	ND	ND	ND	ND
PI4KAP1	4	ND	ND	ND	ND	ND	ND
PPP1R17	4	ND	ND	ND	ND	ND	ND
PRG2	4	ND	ND	ND	ND	ND	ND
PSMD11	4	ND	ND	ND	ND	ND	ND
RGP1	4	ND	ND	ND	ND	ND	ND
RNF6	4	ND	ND	ND	ND	ND	ND
SCAMP1	4	ND	ND	ND	ND	ND	ND
SLC26A2	4	ND	ND	ND	ND	ND	ND
SLC4A3	4	ND	ND	ND	ND	ND	ND
SLIT3	4	ND	ND	ND	ND	ND	ND
SMPD3	4	ND	ND	ND	ND	ND	ND
SPDYE5	4	ND	ND	ND	ND	ND	ND
ST6GALNAC3	4	ND	ND	ND	ND	ND	ND
THTPA	4	ND	ND	ND	ND	ND	ND
TMEM108	4	ND	ND	ND	ND	ND	ND
TMEM11	4	ND	ND	ND	ND	ND	ND
TNFRSF21	4	ND	ND	ND	ND	ND	ND
TPCN1	4	ND	ND	ND	ND	ND	ND
TPST1	4	ND	ND	ND	ND	ND	ND
TRAF3IP1	4	ND	ND	ND	ND	ND	ND
TRUB1	4	ND	ND	ND	ND	ND	ND
TTC21A	4	ND	ND	ND	ND	ND	ND
USP54	4	ND	ND	ND	ND	ND	ND
ZMYM6	4	ND	ND	ND	ND	ND	ND

Table 4: Gene Glossary

Gene Symbol Used in Patent	Official Gene Symbol	Approved Name	HGNC ID	Location
KRT5	KRT5	keratin 5	HGNC:6442	12q13.13
CUX2	CUX2	cut like homeobox 2	HGNC:19347	12q24.11-q24.12
CLEC4C	CLEC4C	C-type lectin domain family 4 member C	HGNC:13258	12p13.31
PLD4	PLD4	phospholipase D family member 4	HGNC:23792	14q32.33
ALOX15B	ALOX15B	arachidonate 15-lipoxygenase type B	HGNC:434	17p13.1
PRG2	PRG2	proteoglycan 2, pro eosinophil major basic protein	HGNC:9362	11q12.1
HTRA4	HTRA4	HtrA serine peptidase 4	HGNC:26909	8p11.22
AMPH	AMPH	amphiphysin	HGNC:471	7p14.1
PNMT	PNMT	phenylethanolamine N-methyltransferase	HGNC:9160	17q12
LEP	LEP	leptin	HGNC:6553	7q32.1
PAPPA2	PAPPA2	pappalysin 2	HGNC:14615	1q25.2
CRH	CRH	corticotropin releasing hormone	HGNC:2355	8q13.1
TIMP4	TIMP4	TIMP metallopeptidase inhibitor 4	HGNC:11823	3p25.2
APOLD1	APOLD1	apolipoprotein L domain containing 1	HGNC:25268	12p13.1
ARHGEF25	ARHGEF25	Rho guanine nucleotide exchange factor 25	HGNC:30275	12q13.3
TIMP3	TIMP3	TIMP metallopeptidase inhibitor 3	HGNC:11822	22q12.3
SEMA3G	SEMA3G	semaphorin 3G	HGNC:30400	3p21.1
IGFBP5	IGFBP5	insulin like growth factor binding protein 5	HGNC:5474	2q35
PRX	PRX	periaxin	HGNC:13797	19q13.2
PITPNM3	PITPNM3	PITPNM family member 3	HGNC:21043	17p13.2-p13.1
FAM107A	FAM107A	family with sequence similarity 107 member A	HGNC:30827	3p14.3-p14.2
TEAD4	TEAD4	TEA domain transcription factor 4	HGNC:11717	12p13.33
HSPA12B	HSPA12B	heat shock protein family A (Hsp70) member 12B	HGNC:16193	20p13
NES	NES	nestin	HGNC:7756	1q23.1
SLC9A3R2	SLC9A3R2	SLC9A3 regulator 2	HGNC:11076	16p13.3
ZEB1	ZEB1	zinc finger E-box binding homeobox 1	HGNC:11642	10p11.22
ADAMTS1	ADAMTS1	ADAM metallopeptidase with thrombospondin type 1 motif 1	HGNC:217	21q21.3
DAAM2	DAAM2	dishevelled associated activator of morphogenesis 2	HGNC:18143	6p21.2
ADAMTS2	ADAMTS2	ADAM metallopeptidase with thrombospondin type 1 motif 2	HGNC:218	5q35.3
VSIG4	VSIG4	V-set and immunoglobulin domain containing 4	HGNC:17032	Xq12
ARRDC2	ARRDC2	arrestin domain containing 2	HGNC:25225	19p13.11
SKIL	SKIL	SKI like proto-oncogene	HGNC:10897	3q26.2
ARRDC4	ARRDC4	arrestin domain containing 4	HGNC:28087	15q26.2
KCNA5	KCNA5	potassium voltage-gated channel subfamily A member 5	HGNC:6224	12p13.32
MYOM3	MYOM3	myomesin 3	HGNC:26679	1p36.11
ATP13A3	ATP13A3	ATPase 13A3	HGNC:24113	3q29
ADA	ADA	adenosine deaminase	HGNC:186	20q13.12
ACY3	ACY3	aminoacylase 3	HGNC:24104	11q13.2
SCT	SCT	secretin	HGNC:10607	11p15.5
NOX4	NOX4	NADPH oxidase 4	HGNC:7891	11q14.3
PACSIN1	PACSIN1	protein kinase C and casein kinase substrate in neurons 1	HGNC:8570	6p21.3
SERPINF1	SERPINF1	serpin family F member 1	HGNC:8824	17p13.3
TIPARP	TIPARP	TCDD inducible poly(ADP-ribose) polymerase	HGNC:23696	3q25.31
LRRC26	LRRC26	leucine rich repeat containing 26	HGNC:31409	9q34.3
PHEX	PHEX	phosphate regulating endopeptidase homolog X-linked	HGNC:8918	Xp22.11
LILRA4	LILRA4	leukocyte immunoglobulin like receptor A4	HGNC:15503	19q13.42
PER1	PER1	period circadian regulator 1	HGNC:8845	17p13.1
CELF4	CELF4	CUGBP Elav-like family member 4	HGNC:14015	18q12.2
LCN6	LCN6	lipocalin 6	HGNC:17337	9q34.3
OLAH	OLAH	oleoyl-ACP hydrolase	HGNC:25625	10p13
PTGDR2	PTGDR2	prostaglandin D2 receptor 2	HGNC:4502	11q12.2
JUN	JUN	Jun proto-oncogene, AP-1 transcription factor subunit	HGNC:6204	1p32.1
PDE8B	PDE8B	phosphodiesterase 8B	HGNC:8794	5q13.3
GSTA3	GSTA3	glutathione S-transferase alpha 3	HGNC:4628	6p12.2
RGP1	RGP1	RGP1 homolog, RAB6A GEF complex partner 1	HGNC:21965	9p13.3
USP54	USP54	ubiquitin specific peptidase 54	HGNC:23513	10q22.2
MRPS35	MRPS35	mitochondrial ribosomal protein S35	HGNC:16635	12p11.22

HEATR9	HEATR9	HEAT repeat containing 9	HGNC:26548	17q12
FSTL3	FSTL3	follistatin like 3	HGNC:3973	19p13.3
DDI2	DDI2	DNA damage inducible 1 homolog 2	HGNC:24578	1p36.21
ZMYM6	ZMYM6	zinc finger MYM-type containing 6	HGNC:13050	1p34.3
ST6GALNAC3	ST6GALNAC3	ST6 N-acetylgalactosaminide alpha-2,6-sialyltransferase 3	HGNC:19343	1p31.1
GBP2	GBP2	guanylate binding protein 2	HGNC:4183	1p22.2
ETV3	ETV3	ETS variant 3	HGNC:3492	1q23.1
ADAM17	ADAM17	ADAM metallopeptidase domain 17	HGNC:195	2p25.1
ATOH8	ATOH8	atonal bHLH transcription factor 8	HGNC:24126	2p11.2
SLC4A3	SLC4A3	solute carrier family 4 member 3	HGNC:11029	2q35
TRAF3IP1	TRAF3IP1	TRAF3 interacting protein 1	HGNC:17861	2q37.3
TTC21A	TTC21A	tetratricopeptide repeat domain 21A	HGNC:30761	3p22.2
HEG1	HEG1	heart development protein with EGF like domains 1	HGNC:29227	3q21.2
ASTE1	ASTE1	asteroid homolog 1	HGNC:25021	3q22.1
TMEM108	TMEM108	transmembrane protein 108	HGNC:28451	3q22.1
ENC1	ENC1	ectodermal-neural cortex 1	HGNC:3345	5q13.3
SCAMP1	SCAMP1	secretory carrier membrane protein 1	HGNC:10563	5q14.1
ARRDC3	ARRDC3	arrestin domain containing 3	HGNC:29263	5q14.3
SLC26A2	SLC26A2	solute carrier family 26 member 2	HGNC:10994	5q32
SLIT3	SLIT3	slit guidance ligand 3	HGNC:11087	5q34-q35.1
CLIC5	CLIC5	chloride intracellular channel 5	HGNC:13517	6p21.1
TNFRSF21	TNFRSF21	TNF receptor superfamily member 21	HGNC:13469	6p12.3
PPP1R17	PPP1R17	protein phosphatase 1 regulatory subunit 17	HGNC:16973	7p14.3
TPST1	TPST1	tyrosylprotein sulfotransferase 1	HGNC:12020	7q11.21
GATSL2	**CASTOR2**	**cytosolic arginine sensor for mTORC1 subunit 2**	**HGNC:37073**	**7q11.23**
SPDYE5	SPDYE5	speedy/RINGO cell cycle regulator family member E5	HGNC:35464	7q11.23
HIPK2	HIPK2	homeodomain interacting protein kinase 2	HGNC:14402	7q34
MTRNR2L6	MTRNR2L6	MT-RNR2 like 6	HGNC:37163	7q34
CLCN1	CLCN1	chloride voltage-gated channel 1	HGNC:2019	7q34
GINS4	GINS4	GINS complex subunit 4	HGNC:28226	8p11.21
C10orf2	**TWNK**	**twinkle mtDNA helicase**	**HGNC:1160**	**10q24.31**
TRUB1	TRUB1	TruB pseudouridine synthase family member 1	HGNC:16060	10q25.3
FAR2	FAR2	fatty acyl-CoA reductase 2	HGNC:25531	12p11.22
CD63	CD63	CD63 molecule	HGNC:1692	12q13.2
CKAP4	CKAP4	cytoskeleton associated protein 4	HGNC:16991	12q23.3
TPCN1	TPCN1	two pore segment channel 1	HGNC:18182	12q24.13
RNF6	RNF6	ring finger protein 6	HGNC:10069	13q12.13
THTPA	THTPA	thiamine triphosphatase	HGNC:18987	14q11.2
FOS	FOS	Fos proto-oncogene, AP-1 transcription factor subunit	HGNC:3796	14q24.3
PARN	PARN	poly(A)-specific ribonuclease	HGNC:8609	16p13.12
ORAI3	ORAI3	ORAI calcium release-activated calcium modulator 3	HGNC:28185	16p11.2
ELMO3	ELMO3	engulfment and cell motility 3	HGNC:17289	16q22.1
SMPD3	SMPD3	sphingomyelin phosphodiesterase 3	HGNC:14240	16q22.1
TMEM11	TMEM11	transmembrane protein 11	HGNC:16823	17p11.1
PSMD11	PSMD11	proteasome 26S subunit, non-ATPase 11	HGNC:9556	17q11.2
EBI3	EBI3	Epstein-Barr virus induced 3	HGNC:3129	19p13.3
CLEC4M	CLEC4M	C-type lectin domain family 4 member M	HGNC:13523	19p13.2
CCDC151	CCDC151	coiled-coil domain containing 151	HGNC:28303	19p13.2
CPAMD8	CPAMD8	C3 and PZP like alpha-2-macroglobulin domain containing 8	HGNC:23228	19p13.11
CNFN	CNFN	cornifelin	HGNC:30183	19q13.2
C22orf39	C22orf39	chromosome 22 open reading frame 39	HGNC:27012	22q11.21
PI4KAP1	PI4KAP1	phosphatidylinositol 4-kinase alpha pseudogene 1	HGNC:33576	22q11.21
ARFGAP3	ARFGAP3	ADP ribosylation factor GTPase activating protein 3	HGNC:661	22q13.2
FLG	FLG	filaggrin	HGNC:3748	1q21.3
ARMS2	ARMS2	age-related maculopathy susceptibility 2	HGNC:32685	10q26.13
CYP26B1	CYP26B1	cytochrome P450 family 26 subfamily B member 1	HGNC:20581	2p13.2
IRF6	IRF6	interferon regulatory factor 6	HGNC:6121	1q32.2
MYH14	MYH14	myosin heavy chain 14	HGNC:23212	19q13.33
PODXL	PODXL	podocalyxin like	HGNC:9171	7q32.3
PPP1R3C	PPP1R3C	protein phosphatase 1 regulatory subunit 3C	HGNC:9293	10q23.32
SH3RF2	SH3RF2	SH3 domain containing ring finger 2	HGNC:26299	5q32

TMC7	TMC7	transmembrane channel like 7	HGNC:23000	16p12.3
ZNF366	ZNF366	zinc finger protein 366	HGNC:18316	5q13.1
ADCY1	ADCY1	adenylate cyclase 1	HGNC:232	7p12.3
C6	C6	complement C6	HGNC:1339	5p13.1
FAM219A	FAM219A	family with sequence similarity 219 member A	HGNC:19920	9p13.3
HAO2	HAO2	hydroxyacid oxidase 2	HGNC:4810	1p12
IGIP	IGIP	IgA inducing protein	HGNC:33847	5q31.3
IL1R2	IL1R2	interleukin 1 receptor type 2	HGNC:5994	2q11.2
NTRK2	NTRK2	neurotrophic receptor tyrosine kinase 2	HGNC:8032	9q21.33
SH3PXD2A	SH3PXD2A	SH3 and PX domains 2A	HGNC:23664	10q24.33
SSUH2	SSUH2	ssu-2 homolog	HGNC:24809	3p25.3
SULT2A1	SULT2A1	sulfotransferase family 2A member 1	HGNC:11458	19q13.33
FMO3	FMO3	flavin containing dimethylaniline monoxygenase 3	HGNC:3771	1q24.3
GATA5	GATA5	GATA binding protein 5	HGNC:15802	20q13.33
HTRA1	HTRA1	HtrA serine peptidase 1	HGNC:9476	10q26.13
C8B	C8B	complement C8 beta chain	HGNC:1353	1p32.2
H19	H19	H19 imprinted maternally expressed transcript	HGNC:4713	11p15.5
MN1	MN1	MN1 proto-oncogene, transcriptional regulator	HGNC:7180	22q12.1
NFE2L1	NFE2L1	nuclear factor, erythroid 2 like 1	HGNC:7781	17q21.3
PRDM16	PRDM16	PR/SET domain 16	HGNC:14000	1p36.32
AP3B2	AP3B2	adaptor related protein complex 3 subunit beta 2	HGNC:567	15q25.2
EMP1	EMP1	epithelial membrane protein 1	HGNC:3333	12p13.1
FLNC	FLNC	filamin C	HGNC:3756	7q32.1
STAG3	STAG3	stromal antigen 3	HGNC:11356	7q22.1
CPB2	CPB2	carboxypeptidase B2	HGNC:2300	13q14.13
TENC1	**TNS2**	**tensin 2**	**HGNC:19737**	**12q13.13**
RP1L1	RP1L1	RP1 like 1	HGNC:15946	8p23.1
A1CF	A1CF	APOBEC1 complementation factor	HGNC:24086	10q11.23
NPR1	NPR1	natriuretic peptide receptor 1	HGNC:7943	1q21.3
TEK	TEK	TEK receptor tyrosine kinase	HGNC:11724	9p21.2
ERRFI1	ERRFI1	ERBB receptor feedback inhibitor 1	HGNC:18185	1p36.23
ARHGEF15	ARHGEF15	Rho guanine nucleotide exchange factor 15	HGNC:15590	17p13.1
CD34	CD34	CD34 molecule	HGNC:1662	1q32.2
RSPO3	RSPO3	R-spondin 3	HGNC:20866	6q22.33
ALPK3	ALPK3	alpha kinase 3	HGNC:17574	15q25.3
SAMD4A	SAMD4A	sterile alpha motif domain containing 4A	HGNC:23023	14q22.2
ZCCHC24	ZCCHC24	zinc finger CCHC-type containing 24	HGNC:26911	10q22.3
LEAP2	LEAP2	liver enriched antimicrobial peptide 2	HGNC:29571	5q31.1
MYL2	MYL2	myosin light chain 2	HGNC:7583	12q24.11
NRG3	NRG3	neuregulin 3	HGNC:7999	10q23.1
ZBTB16	ZBTB16	zinc finger and BTB domain containing 16	HGNC:12930	11q23.2
SERPINA3	SERPINA3	serpin family A member 3	HGNC:16	14q32.13
AQP7	AQP7	aquaporin 7	HGNC:640	9p13.3
SRPX	SRPX	sushi repeat containing protein X-linked	HGNC:11309	Xp11.4
UACA	UACA	uveal autoantigen with coiled-coil domains and ankyrin repeats	HGNC:15947	15q23
ANO1	ANO1	anoctamin 1	HGNC:21625	11q13.3
FKBP5	FKBP5	FKBP prolyl isomerase 5	HGNC:3721	6p21.31
SCN5A	SCN5A	sodium voltage-gated channel alpha subunit 5	HGNC:10593	3p22.2
PTPN21	PTPN21	protein tyrosine phosphatase non-receptor type 21	HGNC:9651	14q31.3
CACNA1C	CACNA1C	calcium voltage-gated channel subunit alpha1 C	HGNC:1390	12p13.33
ERG	ERG	ETS transcription factor ERG	HGNC:3446	21q22.2
SOX17	SOX17	SRY-box 17	HGNC:18122	8q11.23
WWTR1	WWTR1	WW domain containing transcription regulator 1	HGNC:24042	3q25.1
AIF1L	AIF1L	allograft inflammatory factor 1 like	HGNC:28904	9q34.12-q34.13
CA3	CA3	carbonic anhydrase 3	HGNC:1374	8q21.2
HRG	HRG	histidine rich glycoprotein	HGNC:5181	3q27.3
TAT	TAT	tyrosine aminotransferase	HGNC:11573	16q22.2
AQP7P1	AQP7P1	aquaporin 7 pseudogene 1	HGNC:32048	9q13
ADRA2C	ADRA2C	adrenoceptor alpha 2C	HGNC:283	4p16.3
SYNPO	SYNPO	synaptopodin	HGNC:30672	5q33.1
FN1	FN1	fibronectin 1	HGNC:3778	2q35

GPR116	**ADGRF5**	**adhesion G protein-coupled receptor F5**	**HGNC:19030**	**6p12.3**
KRT17	KRT17	keratin 17	HGNC:6427	17q21.2
AZGP1	AZGP1	alpha-2-glycoprotein 1, zinc-binding	HGNC:910	7q22.1
BCL6B	BCL6B	BCL6B transcription repressor	HGNC:1002	17p13.1
KIF1C	KIF1C	kinesin family member 1C	HGNC:6317	17p13.2
GPR4	GPR4	G protein-coupled receptor 4	HGNC:4497	19q13.32
GJA5	GJA5	gap junction protein alpha 5	HGNC:4279	1q21.2
C14orf37	**ARMH4**	**armadillo like helical domain containing 4**	**HGNC:19846**	**14q23.1**
JAG2	JAG2	jagged canonical Notch ligand 2	HGNC:6189	14q32.33
KIF26A	KIF26A	kinesin family member 26A	HGNC:20226	14q32.33
HBG2	HBG2	hemoglobin subunit gamma 2	HGNC:4832	11p15.4
CADM2	CADM2	cell adhesion molecule 2	HGNC:29849	3p12.1
LAMP5	LAMP5	lysosomal associated membrane protein family member 5	HGNC:16097	20p12.2
NOMO1	NOMO1	NODAL modulator 1	HGNC:30060	16p13.11
NXF3	NXF3	nuclear RNA export factor 3	HGNC:8073	Xq22.1
BPIFB3	BPIFB3	BPI fold containing family B member 3	HGNC:16178	20q11.21
CGB8	CGB8	chorionic gonadotropin subunit beta 8	HGNC:16453	19q13.33
CGB5	CGB5	chorionic gonadotropin subunit beta 5	HGNC:16452	19q13.33
ZSCAN23	ZSCAN23	zinc finger and SCAN domain containing 23	HGNC:21193	6p22.1
HSPA1A	HSPA1A	heat shock protein family A (Hsp70) member 1A	HGNC:5232	6p21.33
PMAIP1	PMAIP1	phorbol-12-myristate-13-acetate-induced protein 1	HGNC:9108	18q21.32
C8orf4	**TCIM**	**transcriptional and immune response regulator**	**HGNC:1357**	**8p11.21**
ITM2B	ITM2B	integral membrane protein 2B	HGNC:6174	13q14.2
IFIT2	IFIT2	interferon induced protein with tetratricopeptide repeats 2	HGNC:5409	10q23.31
CD74	CD74	CD74 molecule	HGNC:1697	5q33.1
HSPA6	HSPA6	heat shock protein family A (Hsp70) member 6	HGNC:5239	1q23.3
TFAP2A	TFAP2A	transcription factor AP-2 alpha	HGNC:11742	6p24.3
TRPV6	TRPV6	transient receptor potential cation channel subfamily V member 6	HGNC:14006	7q34
EXPH5	EXPH5	exophilin 5	HGNC:30578	11q22.3
CAPN6	CAPN6	calpain 6	HGNC:1483	Xq23
ALDH3B2	ALDH3B2	aldehyde dehydrogenase 3 family member B2	HGNC:411	11q13.2
RAB3B	RAB3B	RAB3B, member RAS oncogene family	HGNC:9778	1p32.3
MUC15	MUC15	mucin 15, cell surface associated	HGNC:14956	11p14.3
GRHL2	GRHL2	grainyhead like transcription factor 2	HGNC:2799	8q22.3
CSHL1	CSHL1	chorionic somatomammotropin hormone like 1	HGNC:2442	17q23.3
CSH2	CSH2	chorionic somatomammotropin hormone 2	HGNC:2441	17q23.3
KISS1	KISS1	KiSS-1 metastasis suppressor	HGNC:6341	1q32.1
CGA	CGA	glycoprotein hormones, alpha polypeptide	HGNC:1885	6q14.3
PLAC4	PLAC4	placenta enriched 4	HGNC:14616	21q22.2
PSG1	PSG1	pregnancy specific beta-1-glycoprotein 1	HGNC:9514	19q13.2
GH2	GH2	growth hormone 2	HGNC:4262	17q23.3
PSG3	PSG3	pregnancy specific beta-1-glycoprotein 3	HGNC:9520	19q13.2
PSG4	PSG4	pregnancy specific beta-1-glycoprotein 4	HGNC:9521	19q13.31
PSG7	PSG7	pregnancy specific beta-1-glycoprotein 7 (gene/pseudogene)	HGNC:9524	19q13.31
PSG11	PSG11	pregnancy specific beta-1-glycoprotein 11	HGNC:9516	19q13.31
CSH1	CSH1	chorionic somatomammotropin hormone 1	HGNC:2440	17q23.3
PSG2	PSG2	pregnancy specific beta-1-glycoprotein 2	HGNC:9519	19q13.31
HSD3B1	HSD3B1	hydroxy-delta-5-steroid dehydrogenase, 3 beta- and steroid delta-isomerase 1	HGNC:5217	1p12
LGALS14	LGALS14	galectin 14	HGNC:30054	19q13.2
FCGR1C	FCGR1CP	Fc fragment of IgG receptor Ic, pseudogene	HGNC:3615	1q21.1
PSG5	PSG5	pregnancy specific beta-1-glycoprotein 5	HGNC:9522	19q13.31
LAGALS13	LGALS13	galectin 13	HGNC:15449	19q13.2
GCM1	GCM1	glial cells missing transcription factor 1	HGNC:4197	6p12.1

The term "plurality" refers to more than one element. For example, the term is used herein in reference to a number of C-RNA molecules that serve as a signature indicative of preeclampsia.

A plurality may include any two, any three, any four, any five, any six, any seven, any eight, any nine, any ten, any eleven, any twelve, any thirteen, any fourteen, any fifteen, any sixteen, any seventeen, any eighteen, any nineteen, any twenty, any twenty-one, any twenty-two, any twenty-three, any twenty-four, any twenty-five, any twenty-six, any twenty-seven, any twenty-eight, any twenty-nine, any thirty, any thirty-one, any thirty-two, any thirty-three, any thirty-four, any thirty-five, any thirty-six, any thirty-seven, any thirty-eight, any thirty- nine, any forty, any forty-one, any forty- two, any forty-three, any forty-four, any forty-five, any forty-six, any forty-seven, any forty-eight, any forty-nine, any fifty, any fifty-one, any fifty- two, any fifty-three, any fifty-four, any fifty-five, any fifty-six, any fifty-seven, any fifty-eight, any fifty-nine, any sixty, any sixty-one, any sixty-two, any sixty-three, any sixty-four, any sixty-five, any sixty-six, any sixty-seven, any sixty-eight, any sixty-nine, any seventy, any seventy-one, any seventy-two, any seventy-three, any seventy-four, any seventy-five, any seventy-six, any seventy-seven, any seventy-eight, any seventy-nine, any eighty, any eighty-one, any eighty-two, any eighty-three, any eighty-four, any eighty-five, any eighty-six, any eighty-seven, any eighty-eight, any eighty-nine, any ninety, any ninety-one, any ninety- two, any ninety-three, any ninety-four, any ninety-five, any ninety-six, any ninety-seven, any ninety-eight, any ninety-nine, any one hundred, any one hundred and one, any one hundred and two, any one hundred and three, any one hundred and four, any one hundred and five, any one hundred and six, any one hundred and seven, any one hundred and eight, any one hundred and nine, any one hundred ten, any one hundred eleven, any one hundred twelve, any one hundred thirteen, any one hundred fourteen, any one hundred fifteen, any one hundred sixteen, any one hundred seventeen, any one hundred eighteen, any one hundred nineteen, any one hundred twenty, any one hundred twenty-one, or any one hundred twenty-two of the molecules recited in a list described herein. A plurality may include a least any of the numbers recited above. A plurality may include more than any of the numbers recited above. A plurality may include a range of any of those recited above. In some embodiments, a C-RNA signature indicative of preeclampsia includes just one of the biomarkers recited above.

The identification and/or quantification of one of these C-RNA signatures within a sample obtained from a subject can be used to determine that the subject suffers from preeclampsia or is at a risk of developing preeclampsia.

A sample may be a biological sample or biosample, including but not limited to blood, serum, plasma, sweat, tears, urine, sputum, lymph, saliva, amniotic fluid, a tissue biopsy, swab, or smear, including for example, but not limited to, a placental tissue sample. In some preferred embodiments, a biological sample is a cell free plasma sample. A biological sample may be a maternal sample obtained from a pregnant female subject.

As used herein, the term "subject" refers to a human subject as well as a non-human mammalian subject. Although the examples herein concern humans and the language is primarily directed to human concerns, the concept of this disclosure is applicable to any mammal, and is useful in the fields of veterinary medicine, animal sciences, research laboratories and such.

A subject may be a pregnant female, including a pregnant female in any gestational stages of pregnancy. The gestational stage of pregnancy may be, for example, the first trimester, the second trimester, including late second trimester, or the third trimester, including early third trimester. The gestational stage of pregnancy may be, for example, before 16 weeks of pregnancy, before 20 weeks of pregnancy, or after 20 weeks of pregnancy. The gestational stage of pregnancy may be, for example, 8-18 weeks of pregnancy, 10-14 weeks of pregnancy, 11-14 weeks of pregnancy, 11-13 weeks, or 12-13 weeks of pregnancy.

The discovery of cell-free fetal nucleic acids in maternal plasma has opened up new possibilities for noninvasive prenatal diagnosis. Over the last few years, a number of approaches have been demonstrated to allow such circulating fetal nucleic acids to be used for the prenatal detection of chromosomal aneuploidies. Any of the methods described for example in Poon et al., 2000, *Clin Chem*; 1832-4; Poon et al., 2001, *Ann N Y Acad Sci*; 945:207-10; Ng et al., 2003, *Clin Chem*; 49(5):727-31; Ng et al., 2003, *Proc Natl Acad Sci U S A.*;100(8):4748-53; Tsui et al., 2004, *J Med Genet*; 41(6):461-7; Go et al., 2004, *Clin Chem*; 50(8):1413-4; Smets et al., 2006, *Clin Chim Acta*; 364(1-2):22-32; Tsui et al., 2006, *Methods Mol Biol*; 336:123-34; Purwosunu et al., 2007, *Clin Chem*; 53(3):399-404; Chim et al., 2008, *Clin Chem*; 54(3):482-90; Tsui and Lo, 2008, M*ethods Mol Biol*; 444:275-89; Lo, 2008, *Ann N Y Acad Sci*; 1137:140-143; Miura et al., 2010, *Prenat Diagn*; 30(9):849-61; Li et al., 2012, *Clin Chim Acta*; 413(5-6):568-76; Williams et

al., 2013, *Proc Natl Acad Sci U S A*; 110(11):4255-60; Tsui et al., 2014, *Clin Chem*; 60(7):954-62; Tsang et al., 2017, *Proc Natl Acad Sci U S A*; 114(37):E7786-E7795, and US Patent Publication US 2014/0243212 may be used in the methods described herein.

The detection and identification of biomarkers of a C-RNA signature within the maternal circulation indicative of preeclampsia or a risk for developing preeclampsia may involve any of a variety of technologies. For example, biomarkers may be detected in serum by radioimmunoassay or the polymerase chain reaction (PCR) technique may be used.

In various embodiments, the identification of the biomarkers of a C-RNA signature within the maternal circulation indicative of preeclampsia or a risk for developing preeclampsia may involve sequencing the C-RNA molecules. Any of a number of sequencing technologies can be utilized, including, but not limited to, any of a variety of high-throughput sequencing techniques.

In some embodiments, the C-RNA population within a maternal biosample may be subject to enrichment of RNA sequences the include protein-coding sequences prior to sequencing. Any of a variety of platforms available for whole-exome enrichment and sequencing may be used, including but not limited to the Agilent SureSelect Human All Exon platform (Chen et al., 2015a, *Cold Spring Harb Protoc;* 2015(7):626-33. doi: 10.1101/pdb.prot083659); the Roche NimbleGen SeqCap EZ Exome Library SR platform (Chen et al., 2015b, *Cold Spring Harb Protoc*; 2015(7):634-41. doi: 10.1101/pdb.prot084855); or the Illumina TruSeq Exome Enrichment platform (Chen et al., 2015c, *Cold Spring Harb Protoc*; 2015(7):642-8. doi:10.1101/pdb.prot084863). See also "TruSeq™ Exome Enrichment Guide," Catalog # FC-930-1012 Part # 15013230 Rev. B November 2010 and Illumina's "TruSeq™ RNA Sample Preparation Guide," Catalog #RS-122-9001DOC Part # 15026495 Rev. F March 2014.

In particular embodiments, biomarkers of a C-RNA signature within the maternal circulation indicative of preeclampsia or a risk for developing preeclampsia may be detected and identified using microarray techniques. In this method, polynucleotide sequences of interest are plated, or arrayed, on a microchip substrate. The arrayed sequences are then hybridized with a maternal biosample, or a purified and/or enriched portion thereof. Microarrays may include a variety of solid supports including, but not limited to, beads, glass microscope slides, glass wafers, gold, silicon, microchips, and other plastic, metal, ceramic, or biological surfaces.

Microarray analysis can be performed by commercially available equipment, following manufacturer's protocols, such as by using Illumina's technology.

With obtaining, shipping, storing, and/or processing blood samples for the preparation of circulating RNA, steps may be taken to stabilize the sample and/or prevent the disruption of cell membranes resulting in the release of cellular RNAs into the sample. For example, in some embodiments, blood samples may be collected, shipped, and/or stored in tubes that have cell- and DNA-stabilizing properties, such as Streck Cell-Free DNA BCT® blood collection tubes, prior to processing into plasma. In some embodiments, blood samples are not exposed to EDTA. See, for example, Qin et al., 2013, *BMC Research Notes*; 6:380 and Medina Diaz et al., 2016, *PLoS ONE*; 11(11):e0166354.

In some embodiments, blood samples are processed into plasma within about 24 to about 72 hours of the blood draw, and in some embodiments, within about 24 hours of the blood draw. In some embodiments, blood samples are maintained, stored, and/or shipped at room temperature prior to processing into plasma.

In some embodiments, blood samples are maintained, stored, and/or shipped without exposure to chilling (for example, on ice) or freezing prior to processing into plasma.

The disclosure includes kits for use in the diagnosis of preeclampsia and the identification of pregnant women at risk for developing preeclampsia. A kit is any manufacture (e.g. a package or container) including at least one reagent, e.g. a probe, for specifically detecting a C-RNA signature within the maternal circulation as described herein that is indicative of preeclampsia or a risk for developing preeclampsia. The kit may be promoted, distributed, or sold as a unit for performing the methods of the present disclosure.

The use of signatures of circulating RNA found in the maternal circulation specific to preeclampsia in noninvasive methods for the diagnosis of preeclampsia and the identification of pregnant women at risk for developing preeclampsia may be combined with appropriate monitoring and medical management. For example, further tests may be ordered. Such test may include, for example, blood tests to measure liver function, kidney function, and/or platelet and various clotting proteins, urine analysis to measure protein or creatinine levels, fetal ultrasound to measure monitor fetal growth, weight, and amniotic fluid, a nonstress test to measure how fetal heart rate with fetal movement, and/or a biophysical profile using ultrasound to measure your fetal breathing, muscle tone, and movement and the volume of amniotic fluid may be

ordered. Therapeutic interventions may include, for example, increasing the frequency of prenatal visits, antihypertensive medications to lower blood pressure, corticosteroid medications, anticonvulsant medications, bed rest, hospitalization, and/or early delivery. See, for example, Townsend et al., 2016 “Current best practice in the management of hypertensive disorders in pregnancy,” *Integr Blood Press Control*; 9: 79–94.

Therapeutic interventions may include the administration of low dose aspirin to pregnant women identified at risk of for developing preeclampsia. A recent multicenter, double-blind, placebo-controlled trial demonstrated that treatment of women at high risk for preterm preeclampsia with low-dose aspirin resulted in a lower incidence of this diagnosis compared to placebo (Rolnik et al., 2017, "Aspirin versus Placebo in Pregnancies at High Risk for Preterm Preeclampsia," *N Engl J Med*; 377(7):613-622). Dosages of low dose aspirin include, but are not limited to, about 50 to about 150 mg per day, about 60 to about 80 mg per day, about 100 or more mg per day, or about 150 mg per day. Administration may begin, for example, at or before 16 weeks of gestation or from 11 to 14 weeks of gestation. Administration may continue thru 36 weeks of gestation.

The present invention is illustrated by the following examples. It is to be understood that the particular examples, materials, amounts, and procedures are to be interpreted broadly in accordance with the scope and spirit of the invention as set forth herein.

EXAMPLES

Example 1

C-RNA Signatures Unique to Pregnancy

The presence of circulating nucleic acid in maternal plasma provides a window into the progression and health of the fetus and the placenta (Fig. 1). Circulating RNA (C-RNA) is detected in maternal circulation and originates from two predominant sources. A significant fraction of C-RNA originates from apoptotic cells, which release vesicles containing C-RNA into the blood stream. C-RNA also enters maternal circulation through the shedding of active signaling vesicles such as exosomes and microvesicles from a variety of cell types. As shown in Fig. 2, C-RNA is therefore comprised of the byproducts of cell death as well as active signaling products. Characteristics of C-RNA include generation through common processes, release from cells throughout the body, and stable and contained in vesicles. It represents a circulating transcriptome that reflects tissue-specific changes in gene expression, signaling, and cell death.

C-RNA has the potential to be an excellent biomarker for at least the following reasons:

1) All C-RNA is contained within membrane bound vesicles, which protects the C-RNA from degradation, making it quite stable in the blood.

2) C-RNA originates from all cell types. For example, C-RNA has been shown to contain transcripts from both the placenta and the developing fetus. The diverse origins of C-RNA give it the potential to be a rich repository for accessing information on both fetal and overall maternal health.

C-RNA libraries were prepared from plasma samples using standard Illumina library prep and whole exome enrichment technology. This is shown in Fig. 3. Specifically, Illumina TruSeq™ library prep and RNA Access Enrichment were used. Using this approach, libraries were generated that have 90% of the reads aligning to the human coding region (Fig. 3 and Fig. 7). Samples were downsampled to 50 M reads and ≥40 M mapped reads were used for downstream analysis. Samples were processed using the C-RNA workflow shown in Fig. 3. Dual Indexed libraries. Sequenced 50X50 on Hiseq2000

As shown in Fig. 4, comparing results from plasma samples from third trimester pregnant women to plasma samples from non-pregnant women provides a clear signature unique to

pregnancy. The top twenty differential abundance genes of this signature are CSHL1, CSH2, KISS1, CGA, PLAC4, PSG1, GH2, PSG3, PSG4, PSG7, PSG11, CSH1, PSG2, HSD3B1, GRHL2, LGALS14, FCGR1C, PSG5, LGALS13, and GCM1. The majority of the genes identified in the pregnancy signature are placentally expressed and also correlate with published data. These results also confirm that placental RNA can be accessed in in the maternal circulation.

Example 2

C-RNA Signatures Across Gestational Age

This example characterized C-RNA signatures across different gestational ages throughout pregnancy. It is expected that the changes in C-RNA signatures at different time points longitudinally across pregnancy will be more subtle than the differences between C-RNA signatures of pregnant and non-pregnant samples noted in Example 1. As shown in Fig. 5, clear temporal changes in C-RNA profiles of the signature genes were observed as pregnancy progressed, with a clear group of genes upregulated in the first trimester and clear group of genes that increase in the third trimester.

These genes included CGB8, CGB5, ZSCAN23, HSPA1A, PMAIP1, C8orf4, ITM2B, IFIT2, CD74, HSPA6, TFAP2A,TRPV6, EXPH5, CAPN6, ALDH3B2, RAB3B, MUC15, GSTA3, GRHL2, and CSHL1, as listed in Fig. 5.

These genes may also include CSHL1, CSH2, KISS1, CGA, PLAC4, PSG1, GH2, PSG3, PSG4, PSG7, PSG11, CSH1, PSG2, HSD3B1, GRHL2, LGALS14, FCGR1C, PSG5, LGALS13, and GCM1.

These changes throughout the course of pregnancy correlate with published data from both Steve Quake and Dennis Lo. See, for example, Maron et al., 2007, "Gene expression analysis in pregnant women and their infants identifies unique fetal biomarkers that circulate in maternal blood," *J Clin Invest*; 117(10):3007–3019; Koh et al., 2014, "Noninvasive in vivo monitoring of tissue-specific global gene expression in humans," *Proc Natl Acad Sci U S A*; 111(20):7361-6; and Ngo et al., 2018, "Noninvasive blood tests for fetal development predict gestational age and preterm delivery," *Science*; 360(6393):1133-1136. C-RNA signatures correlating with patterns of gene expression of the placenta were found. Thus, this approach is

able to detect subtle changes within pregnancy and provides non-invasive means to monitor placental health.

Example 3

C-RNA Signatures of Preeclampsia

With this example, C-RNA signatures unique to preeclampsia were identified. C-RNA signatures were determined in samples collected from pregnant women diagnosed with preeclampsia from two studies, the RGH14 Study (registered with clinical trials.gov as NCT0208494) and the Pearl Study (also referred to herein as the Pearl Biobank; registered with clinical trials.gov as NCT02379832)), were assayed (Fig. 6). Two tubes of blood were collected at the time of diagnosis for preeclampsia. Eighty controls samples matched for gestational age were collected to minimize transcriptional variability not related to the preeclampsia disease state and to control for gestational age differences in C-RNA signatures. Samples from the RGH14 study were used to identify a set of biologically relevant genes, and the predictive value of these biomarkers was validated in an independent cohort of samples from the Pearl Biobank.

In the analysis of the RGH14 data, C-RNA signatures unique to preeclampsia (PE) were identified using four different methods, the TREAT method, a Bootstrap method, a jackknifing method, and the Adaboost method. Example 3 focuses on the first 3 analysis methods and Example four focuses on the Adaboost method.

The t-test relative to threshold (TREAT) statistical method utilizing the EDGR program allows researchers to formally test (with associated p-values) whether the differential expression in a microarray experiment is greater than a given (biologically meaningful) threshold. See McCarthy and Smyth, 2009 "Testing significance relative to a fold-change threshold is a TREAT," *Bioinformatics*; 25(6):765-71 for a more detailed description of the TREAT statistical method and Robinson et al., 2010, "edgeR: a Bioconductor package for differential expression analysis of digital gene expression data," *Bioinformatics*; 26:139-140 for a more detailed description of the EDGR program. See Freund and Schapire, 1997, "A Decision-Theoretic Generalization of On-Line Learning and an Application to Boosting," *Journal of Computer and Systems Sciences*; 55(1):119-139 and Pedregosa et al., 2011, "Scikit-learn: Machine Learning in

Python," *JMLR*; 12:2825-2830 for a more detailed description of the Adaboost method. The Adaboost method will be discussed in Example 4.

In the first method, standard statistical testing (TREAT method) was used to identify genes that are statistically different in the RGH14 preeclampsia cohort of 40 patients as compared to a subset of matched controls (40 patients). 122 genes were identified as statistically different in the preeclampsia cohort (40 patients) as compared to a subset of matched controls (40 patients) (Fig. 8, right panel). These genes include CYP26B1, IRF6, MYH14, PODXL, PPP1R3C, SH3RF2, TMC7, ZNF366, ADCY1, C6, FAM219A, HAO2, IGIP, IL1R2, NTRK2, SH3PXD2A, SSUH2, SULT2A1, FMO3, FSTL3, GATA5, HTRA1, C8B, H19, MN1, NFE2L1, PRDM16, AP3B2, EMP1, FLNC, STAG3, CPB2, TENC1, RP1L1, A1CF, NPR1, TEK, ERRFI1, ARHGEF15, CD34, RSPO3, ALPK3, SAMD4A, ZCCHC24, LEAP2, MYL2, NRG3, ZBTB16, SERPINA3, AQP7, SRPX, UACA, ANO1, FKBP5, SCN5A, PTPN21, CACNA1C, ERG, SOX17, WWTR1, AIF1L, CA3, HRG, TAT, AQP7P1, ADRA2C, SYNPO, FN1, GPR116, KRT17, AZGP1, BCL6B, KIF1C, CLIC5, GPR4, GJA5, OLAH, C14orf37, ZEB1, JAG2, KIF26A, APOLD1, PNMT, MYOM3, PITPNM3, TIMP4, HTRA4, AMPH, LCN6, CRH, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, LEP, NES, VSIG4, HBG2, CADM2, LAMP5, PTGDR2, NOMO1, NXF3, PLD4, BPIFB3, PACSIN1, CUX2, FLG, CLEC4C, and KRT5.

The TREAT method did not identify a set of genes that 100% accurately classifies the preeclampsia patients into a separate group (Fig. 15). However, focusing in on these identified genes did improve classification compared to using the entire data set of all measured genes (Fig. 8, left panel). This highlights the value of focusing in on a subset of genes for prediction. However, with the TREAT method, a significant amount of variability was observed in the genes identified depending on which controls were selected. To deal with this biological variability and further improve the predictive value of our gene list, a second bootstrapping approach was developed.

In the RGH14 study more control samples (80) are available than preeclampsia patient samples (40). Thus, the RGH14 cohort of 40 preeclampsia patient samples was compared to a random selection of 40 controls samples (still matched for gestational age) and a gene list that is statistically different in the preeclampsia cohort was identified. As shown in Fig. 9, this was

then repeated 1,000 times, to identify how often a set of genes was identified. A significant subset of genes only show up less than 10 times out of the 1,000 iterations (less than 1% of the 1,000 iterations). These low frequency genes most likely are due to biological noise and may not reflect a gene that is universally specific to preeclampsia. So, the gene list was further downselected by requiring a gene to be considered as statistically different in the preeclampsia cohort only if identified in 50% of the 1,000 iterations performed (Fig. 9, right panel). As shown in Fig. 10, differential transcript abundance with the additional bootstrapping selection distinguishes preeclampsia samples from healthy controls. Using this additional requirement helped address biological variability and further improved the ability to classify preeclampsia samples correctly.

Using this Bootstrap method, 27 genes were identified as statistically associated with preeclampsia. These genes include TIMP4, FLG, HTRA4, AMPH, LCN6, CRH, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, CLEC4C, KRT5, PRG2, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, LEP, NES, and VSIG4. The genes identified with this bootstrapping method had excellent concordance with published data. Approximately 75% of these genes are expressed by the placenta. As shown in Fig. 11, there is overlap with known markers of preeclampsia, including PAPPA and CRH. And, a significant number of these genes are involved in embryo development, extracellular matrix remodeling, immune regulation, and cardiovascular function, all pathways known to be dysregulated in preeclampsia.

A third jackknifing approach was also developed to capture the subset of genes with the highest predictive value. This approach is similar to the bootstrapping method. Patients from both preeclampsia and control groups were randomly subsampled and differentially abundant genes identified 1,000 times. Instead of using the frequency with which a gene is identified as statistically different, the jackknifing approach calculated confidence intervals (95%, one-sided) for the p-value of each transcript. Genes where this confidence interval exceeded 0.05 were excluded. (Fig. 16, left panel).

Using the jackknifing approach, 30 genes were identified as predictive of preeclampsia: VSIG4, ADAMTS2, NES, FAM107A, LEP, DAAM2, ARHGEF25, TIMP3, PRX, ALOX15B, HSPA12B, IGFBP5, CLEC4C, SLC9A3R2, ADAMTS1, SEMA3G, KRT5, AMPH, PRG2, PAPPA2, TEAD4, CRH, PITPNM3, TIMP4, PNMT, ZEB1, APOLD1, PLD4, CUX2, HTRA4.

As shown in Fig. 16 right panel, this approach gave good classification of preeclampsia patients in the RGH14 data set (compare Fig. 15 (TREAT), Fig. 10 (bootstrapping) and Fig. 16 (jackknifing)). Each identified gene list was also used to classify preeclampsia samples in the independent Pearl Biobank dataset. As shown in Fig. 17, each gene list was able to classify preeclampsia samples.

All genes identified by the bootstrapping and jackknifing methods are represented in the 122 TREAT method genes (Table 2, DEX analysis, TruSeq library prep method). The bootstrapping and jackknifing approach gene lists are highly concordant, with over 70% of genes in common. Nearly 90% of transcripts identified by any approach exhibit increased transcript abundance in preeclampsia patients, consistent with elevated signaling and/or cell death in this disease.

Example 4

Identification of C-RNA Signatures with Adaboost

With this example an alternative approach, a publicly available machine learning algorithm called adaboost, was used to identify a specific C-RNA signature associated with preeclampsia. As shown in Fig. 12, this approach identifies a set of genes that has the most predictive power to classify a sample as preeclampsia (PE) or normal. Using this gene list, the clearest separation of a preeclampsia cohort from healthy controls was observed. However, this approach can also be very susceptible to overtraining to the samples used to build the model. Thus, the predictive model was validated using a completely independent data set from the PEARL study (Fig. 13). Using this Adaboost gene list, 85% of the preeclampsia samples were accurately classify with 85% specificity (Fig. 14). Overall, the Adaboost machine learning approach built the most accurate predictive model for preeclampsia.

Using the Adaboost method, 75 genes were identified as statistically associated with preeclampsia (Table 3, AdaBoost Analysis, TruSeq library prep method). These genes include ARRDC2, JUN, SKIL, ATP13A3, PDE8B, GSTA3, PAPPA2, TIPARP, LEP, RGP1, USP54, CLEC4C, MRPS35, ARHGEF25, CUX2, HEATR9, FSTL3, DDI2, ZMYM6, ST6GALNAC3, GBP2, NES, ETV3, ADAM17,ATOH8, SLC4A3, TRAF3IP1, TTC21A, HEG1, ASTE1, TMEM108, ENC1, SCAMP1, ARRDC3, SLC26A2, SLIT3, CLIC5, TNFRSF21, PPP1R17,

TPST1, GATSL2, SPDYE5, HIPK2, MTRNR2L6, CLCN1, GINS4, CRH, C10orf2, TRUB1, PRG2, ACY3, FAR2, CD63, CKAP4, TPCN1, RNF6, THTPA, FOS, PARN, ORAI3, ELMO3, SMPD3, SERPINF1, TMEM11, PSMD11, EBI3, CLEC4M, CCDC151, CPAMD8, CNFN, LILRA4, ADA, C22orf39, PI4KAP1, and ARFGAP3.

A refined AdaBoost model was also developed for robust classification of PE samples. In order to create a generalized machine learning model that could accurately predict new samples, we used a rigorous approach that avoided overfitting to a single dataset and validated the final classifier with samples not used for model building. As illustrated in Fig. 18, the RGH14 dataset was divided into 6 pieces by random selection: a holdout subset with 12% of samples which was excluded from model building, and 5 evenly sized test subsets. For each iteration subsets were designated as training data or test samples. This process, starting at building the AdaBoost model was repeated for a minimum of 10 times on this data subset. After 50 high performing models were built for the 5 test-train subsets, the estimators from all models were merged into a single AdaBoost model.

Using the refined AdaBoost model, 11 genes were identified as statically associated with preeclampsia. These genes include CLEC4C, ARHGEF25, ADAMTS2, LEP, ARRDC2, SKIL, PAPPA2, VSIG4, ARRDC4, CRH and NES. The performance of this predictive model was validated using the hold out data set from RGH14 as well as in the completely independent Pearl Biobank cohort (Fig. 19).

AdaBoost Model Creation Description. The AdaBoost classification approach was refined in order to obtain more specific gene sets (AdaBoost Refined 1-7) by the following approach, also illustrated in Fig. 18. The RGH14 dataset was divided into 6 pieces by random selection: a holdout subset with 12% of samples which was excluded from model building, and 5 evenly sized test subsets.

For each of the test subsets, training data was assigned as all samples in neither the holdout or test samples. Gene counts for the test and training samples were TMM-normalized in edgeR, then standardized such that the training data has mean of 0 and standard deviation of 1 for each gene. An AdaBoost model with 90 estimators and 1.6 learning rate was then fit to the training data. Feature pruning was then performed by determining the feature importance of each gene in the model and testing the impact of eliminating estimators using genes with importance below a threshold value. The threshold resulting in the best performance (as

measured by Matthew's correlation coefficient on test data classification) with the fewest genes was selected, and that model retained. This process, starting at building the AdaBoost model was repeated for a minimum of 10 times on this data subset.

After all 50 plus models were built for the 5 test-train subsets, the estimators from all models were merged into a single AdaBoost model. Feature pruning was performed again, this time using the percent of models incorporating a gene to for threshold values and assessing performance with the average negative log loss value for the classification of each test subset. The model which obtained the maximal negative log loss value with the fewest genes was selected as the final AdaBoost model.

AdaBoost Gene Lists. Upon repetition of this process, slight variations were observed in the genes selected for the final model, due to innate randomization in the AdaBoost algorithm implementation, however performance remained high for predicting the test data, holdout data, and independent (Pearl) datasets.

Eleven total genes were observed in at least one of 14 AdaBoost Refined models generated: ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, CRH, LEP, NES, PAPPA2, SKIL, VSIG4 (AdaBoost Refined 1), although no models were generated that included all simultaneously.

Two observed gene sets offered the highest performance on classification of independent data. These are AdaBoost Refined 2: ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, VSIG4 and AdaBoost Refined 3: ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, SKIL, VSIG4.

Four additional gene sets performed almost as highly as AdaBoost Refined 2-3. These are AdaBoost Refined 4: ADAMTS2, ARHGEF25, ARRDC4, CLEC4C, LEP, NES, SKIL, VSIG4; AdaBoost Refined 5: ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, CRH, LEP, PAPPA2, SKIL, VSIG4; AdaBoost Refined 6: ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, SKIL; and AdaBoost Refined 7: ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, LEP, PAPPA2, SKIL.

Example 5

Identification of C-RNA signature with transposome based library prep

The RGH14 samples were also processed through the Illumina Nextera Flex for Enrichment protocol, enriched for whole exome and sequenced to >40 million reads. This approach is more sensitive and robust for low inputs, thus likely to identify additional genes predictive of preeclampsia. This dataset was run through three analysis methods, standard differential expression analysis (TREAT), jackknifing, and the refined Adaboost model. See Example 3 and Example 4 for detailed description of these analysis methods.

Changing the method for generating libraries altered the genes detected in all three analysis methods. For the TREAT method, 26 genes were identified as differentially abundant in preeclampsia, with the majority again showing elevated abundance in preeclampsia (See Table 2, DEX Analysis, Nextera Flex for Enrichment library prep method). These genes include ADAMTS1, ADAMTS2, ALOX15B, AMPH, ARHGEF25, CELF4, DAAM2, FAM107A, HSPA12B, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LEP, LRRC26, NES, OLAH, PACSIN1, PAPPA2, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, VSIG4. Fig. 20 shows classification of the RGH14 samples with this gene list.

Applying the jackknifing analysis method downselected the TREAT list to 22 genes identified as differentially abundant in preeclampsia. These genes included ADAMTS1, ADAMTS2, ALOX15B, ARHGEF25, CELF4, DAAM2, FAM107A, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LEP, LRRC26, NES, OLAH, PRX, PTGDR2, SEMA3G, SLC9A3R2, TIMP3, VSIG4. The improved performance of this list is shown in Fig. 20.

The refined AdaBoost model approach was applied to this data, as described in Example 4. Using this method, 24 genes are identified as statically associated with preeclampsia (Table 3, AdaBoost Analysis, Nextera Flex for Enrichment library prep method). These genes include LEP, PAPPA2, KCNA5, ADAMTS2, MYOM3, ATP13A3, ARHGEF25, ADA, HTRA4, NES, CRH, ACY3, PLD4, SCT, NOX4, PACSIN1, SERPINF1, SKIL, SEMAG3, TIPARP, LRRC26, PHEX, LILRA4, and PER1. The performance of this predictive model is indicated in Fig. 21.

Example 6

Circulating transcriptome measurements from maternal blood detects early-onset preeclampsia signature

5 Molecular tools to non-invasively monitor pregnancy health from conception to birth would enable accurate detection of pregnancies at risk for adverse outcomes. Circulating RNA (C-RNA) is released by all tissues into the bloodstream, offering an accessible, comprehensive measurement of placental, fetal and maternal health (Koh et al., 2014, *Proceedings of the National Academy of Sciences*; 111:7361-7366; and Tsui et al., 2014, *Clinical Chemistry*;
10 60:954-962). Preeclampsia (PE), a prevalent and potentially fatal pregnancy complication, is placental in origin but gains a substantial maternal component as the disease progresses (Staff et al., 2013, *Hypertension*; 61:932-942; and Chaiworapongsa et al., 2014, *Nature Reviews Nephrology*; 10, 466-480). Yet purported biomarkers have shown limited clinical utility (Poon and Nicolaides, 2014, *Obstetrics and Gynecology International*; 2014:1-11; Zeisler et al., 2016,
15 *N Engl J Med*; 374:13-22; and Duhig etal., 2018, *F1000Research*; 7:242). Hypothesizing that characterization of the circulating transcriptome may identify better biomarkers, C-RNA was analyzed from 113 pregnancies, 40 at the time of early-onset PE diagnosis. Using a novel workflow, differences were identified in the abundance of 30 transcripts which are consistent with the biology of PE and represent placental, fetal, and maternal contributions. Further, a
20 machine learning model was developed, demonstrating that only seven C-RNA transcripts are required to classify PE in two independents cohorts (92-98% accuracy). The global measurements of C-RNA disclosed in this example highlight the utility in monitoring both maternal and fetal health and hold great promise for the diagnosis and prediction of at-risk pregnancies.

25 Several studies have begun to investigate and identify potential biomarkers in C-RNA for a range of pregnancy complications (Pan et al., 2017, *Clinical Chemistry*; 63:1695-1704; Whitehead et al., 2016, *Prenatal Diagnosis*; 36:997-1008; Tsang et al., 2017, *Proc Natl Acad Sci USA*; 114: E7786-E7795; and Ngo et al., 2018, *Science*; 360:1133-1136). However, these studies have involved few patients and have been limited to monitoring small numbers of genes
30 – almost exclusively placental and fetal derived transcripts. Measurements of the entire circulating transcriptome are difficult to perform because they require specific upfront sample

collection and processing to minimize variability and contamination from cell lysis (Chiu et al., 2001, *Clinical Chemistry*; 47:1607-1613; and Page et al., 2013, *PLoS ONE*; 8: e77963). This complex workflow makes large clinical sample collections difficult to achieve because the labor required for immediate processing of blood samples is infeasible for many clinics (Marton and Weiner, 2013, *BioMed Research International*; 2013:891391). Therefore, with this example, a method was established that allows overnight shipment of blood to a processing lab where every step of sample preparation is performed in a controlled environment, providing a scalable platform for clinical trial level assessments (Fig 22A).

The lynchpin of this method is the ability to ship blood overnight to a processing lab. The C-RNA pregnancy signal was assessed after overnight, room-temperature shipping in several tube types (Figs. 26A-26C). Blood stored in EDTA tubes, the gold standard used by prior C-RNA studies, exhibited a reduction in the abundance of pregnancy-associated transcripts and overall instability of the transcriptomic profile (Qin et al., 2013, *BMC Research Notes*; 6:380). In contrast, the predominant tube type used for Non-Invasive Prenatal Testing (NIPT), Cell-Free DNA BCT (Streck), retained the signal from placental transcripts and had improved technical reproducibility (Fig. 26B) (Medina Diaz et al., 2016, *PLoS ONE*; 11:e0166354).

Shipment of blood allowed us to easily obtain an average of 5 mL plasma per patient from a single tube of blood. The difference in C-RNA data quality was assessed when using varying plasma volumes and determined that using < 2 mL plasma significantly increased noise and decreased library complexity (Figs. 27A and 27B). Thus 4 mL of plasma was used for the studies of this example to maximize confidence in data quality.

This novel workflow was validated by recapitulating previous work monitoring C-RNA dynamics of >10,000 transcripts per healthy pregnancy from first to third trimester. Using 152 samples collected serially from 45 healthy pregnancies (Pre-Eclampsia and Growth Restriction Longitudinal Study Control Cohort - PEARL; NCT02379832; Table 5), 156 significantly altered transcripts were identified, with the majority increasing in abundance as pregnancy progresses (Fig. 22B). 42% of the altered genes were identified in prior C-RNA studies (Fig. 22C) (Koh et al., 2014, *Proceedings of the National Academy of Sciences*; 111:7361-7366; and Tsui et al., 2014, *Clinical Chemistry*; 60:954-962). Of the 91 transcripts identified only in this study, 64% are expressed by placental and/or fetal tissues (Figs. 22D and 28A-28C). Presumably, the remaining genes reflect the maternal response to pregnancy.

Study Design

For the next phase of investigation, the workflow was applied on clinical samples to measure C-RNA changes in PE (iPC, Illumina Preeclampsia Cohort). PE is a heterogeneous disorder and associated with different severity and patient outcomes based on whether it manifests before (early-onset) or after (late-onset) 34 gestational weeks (Staff et al., 2013, *Hypertension*; 61:932-942; Chaiworapongsa et al., 2014, *Nature Reviews Nephrology*; 10, 466-4803; and Dadelszen et al., 2003, *Hypertension in Pregnancy*; 22:143-148). This study to focused on the more severe early-onset form of PE and defined strict diagnostic criteria with clear inclusion and exclusion requirements – most critically excluding any individuals with a history of chronic hypertension – in order to obtain a clean cohort (Table 6) (Nakanishi et al., 2017, *Pregnancy Hypertension*; 7:39-43; and Hiltunen et al., 2017, *PLoS ONE*; 12:e0187729). Maternal characteristics, pregnancy outcomes, and medications in use were recorded throughout the study (Table 7). 113 samples were collected across 8 sites (Table 8), 40 at the time of PE diagnosis, and 73 controls gestationally-age matched within 1 week (Fig. 23A). All but one woman with PE gave birth prematurely, in contrast to 9.5% of controls, confirming these diagnostic criteria as identifying individuals severely impacted by this disease (Fig. 23C).

All samples were randomly distributed across multiple processing batches, then sequenced to ≥40 M reads. Standard differential expression analysis using the full cohort identified 42 altered transcripts, with 37 increased in PE (Fig. 24A, blue and orange). However, of concern was the high variability observed in the genes detected as altered when different subsets of controls were selected for analysis.

To address this discrepancy, a jackknifing approach was incorporated which allowed the identification of the genes that are most consistently altered (Figs. 24A and 24B, orange). 1,000 iterations of differential analysis with randomly selected sample subsets were performed, which allowed the construction of confidence intervals for the p-values associated with each putatively altered transcript (Fig. 29A). 12 genes whose confidence interval exceeded 0.05 were excluded (Fig. 24B). These genes would not have been excluded by simply setting a threshold for baseline abundance or biological variance (Fig. 29B), however it was observed that these transcripts have lower predictive value (Fig. 29C). Hierarchical clustering indicates these genes are not altered

universally in the PE cohort, and thus lack sensitivity (73%) for accurate classification of this condition (Fig. 29D).

The analysis then focused on the refined 30 gene set, 60% of which have previously been associated with PE (Namli et al., 2018, *Hypertension in Pregnancy*; 37: 9-17; Than et al., 2018, *Frontiers in Immunology*; 9:1661; Kramer et al., 2016, *Placenta*; 37:19-25; Winn et al., 2008, *Endocrinology*; 150:452-462; and Liu et al., 2018, *Molecular Medicine Reports*; 18:2937-2944). qPCR analysis confirmed 19 of 20 genes as significantly altered in PE (Fig. 24C, Table 9). Strikingly, 40% of these genes encode for extracellular or secreted protein products. Additionally, nearly all genes are involved in PE relevant processes, including extracellular matrix (ECM) remodeling, pregnancy duration, placental/fetal development, angiogenesis, and hypoxia response (Table 10). 67% of these transcripts were expressed by the placenta and/or fetus (Fig. 24D). In the remaining maternally expressed transcripts, cardiovascular and immune functions were well represented (Table 10). Hierarchical clustering of these genes effectively segregated PE and control samples with 98% sensitivity and 97% specificity (Fig. 24E). Intriguingly, clinical data for the two misidentified controls indicated potentially confounding health problems, as suggested by their use of hypertensive medication (Table 7).

Using the genes identified in iPC, the ability to cluster a cohort of samples obtained from an independent biobank was assessed - the Pre-Eclampsia and Growth Restriction Longitudinal Study (PEARL; NCT02379832; Figs. 23B and 23C, Table 11). This cohort consisted of both early- (diagnosed at < 34 weeks); and late-onset PE with gestationally age-matched controls. Early-onset PE samples clustered separately from matched controls with 83% sensitivity and 92% specificity, further validating the relevance of these transcripts (Fig. 24F). In contrast, no clustering was observed for the late-onset PE and matched control samples (Fig. 24G).

The iPC data was then used to build an AdaBoost model for robust classification of PE samples. In order to create a generalized machine learning model that could accurately predict new samples, a rigorous approach was used that avoided overfitting to a single dataset and validated the final classifier with samples not used for model building (Figs. 30A-30D and Figs. 31A-31E). Surprisingly, the final model only utilized 7 genes, 3 of which have not been previously reported (Fig. 25A). For the entire iPC cohort, this model classified samples with extremely high accuracy (AUC=0.99, sensitivity=98%, specificity=99%; Figs. 25B and 25C, blue). Early-onset PE PEARL samples were also accurately classified (AUC=0.88,

sensitivity=100%, specificity=83%; Figs. 25B and 25C, pink). Unexpectedly, late-onset PE PEARL samples were also classified with reasonable accuracy (AUC=0.74, sensitivity=75%, specificity=67%; Figs. 25B and 25C, green).

This gene set was highly concordant with transcripts identified by differential abundance analysis (Fig. 25D; Table 10). The classifier relied on both placentally and maternally expressed transcripts (Fig. 25E). All genes used by the model form protein products that are either extracellular or membrane bound. Despite the small number of genes selected by AdaBoost, a diversity of PE-relevant functions was observed, specifically cardiovascular function and angiogenesis, immune regulation, fetal development, and ECM remodeling.

Methods

Prospective Clinical Sample Collection. Pregnant patients were recruited in an Illumina sponsored clinical study protocol in compliance with the International Conference on Harmonization for Good Clinical Practice. Following informed consent, 20 mL whole blood samples were collected from 40 pregnant women with a diagnosis of preeclampsia before 34 weeks gestation with severe features defined per ACOG guidelines (Table 6). Samples from 76 healthy pregnancies were also collected and were matched for gestational age to the preeclampsia group. Three control samples developed term preeclampsia after blood collection and were excluded from data analysis. For detailed inclusion and exclusion criteria, see Table 6. Patient clinical history, treatment and birth outcome information were also recorded (Table 7).

Patients were recruited across 8 different clinical sites, including University of Texas Medical Branch (Galveston, Texas), Tufts Medical Center (Boston, MA), Columbia University Irving Medical Center (New York, NY), Winthrop University Hospital (Mineola, NY), St. Peter's University Hospital (New Brunswick, NJ), Christiana Care (Newark, DE), Rutgers University Robert Wood Johnson Medical School (New Brunswick, NJ) and New York Presbyterian/Queens (New York, NY). The clinical protocol and informed consent were approved by each clinical site's Institutional Review Board. See Table 8 for patient distribution across clinical sites.

PEARL Validation Cohort Study Design. Illumina obtained plasma samples from the Preeclampsia and Growth Restriction Longitudinal study (PEARL; NCT02379832) to be used as an independent validation cohort. Plasma samples were obtained after the study had reached

completion. PEARL samples were collected at the Centre hospitalier universitaire de Québec (CHU de Québec) with principal investigator Emmanual Bujold, MD, MSc. A group of 45 control pregnancies and 45 case pregnancies were recruited in this study and written informed consent was obtained for all patients. Only participants above 18 years of age were eligible, and all pregnancies were singleton.

Preeclampsia Group. The criteria for preeclampsia was defined based on the Society of Obstetricians and Gynecologists of Canada (SOGC) June 2014 criteria for preeclampsia, with a gestational age requirement between 20 and 41 weeks. A blood sample was taken once at the time of diagnosis.

Control Group. 45 pregnant women who were expected to have a normal pregnancy were recruited between 11 and 13 weeks gestational age. Each enrolled patient was followed longitudinally with blood drawn at 4 timepoints throughout pregnancy until birth. The control women were divided into three subgroups and subsequent follow up blood draws were staggered to cover the entire range of gestational ages throughout pregnancy (Table 5).

The PEARL control samples were used for two purposes. 153 longitudinal samples from 45 individual women were used to monitor placental dynamics throughout pregnancy. Additionally, control samples were selected for comparison to the preeclampsia cohort, which were matched for gestational age and used to validate the model.

Study Sample Processing. All samples from the Illumina prospective collection and the PEARL samples were processed identically by investigators blinded to disease status. Two tubes of blood were collected per patient in Cell-Free DNA BCT tubes (Streck) following the manufacturer instructions. Blood samples were stored and shipped at room temperature overnight and processed within 72 hours. Blood was centrifuged at 1,600 x g for 20 minutes at room temperature, plasma transferred to a new tube and centrifuged additional 10 minutes at 16,000 x g to remove residual cells. Plasma was stored at -80°C until use. Circulating RNA was extracted from 4.5 mL of plasma using the Circulating Nucleic Acid Kit (Qiagen) followed by DNAse I digestion (Thermofisher) according to manufacturer's instructions.

cDNA Synthesis and Library Prep. Circulating RNA was fragmented at 94°C for 8 minutes followed by random hexamer primed cDNA synthesis using the Illumina TruSight Tumor 170 Library Prep kit (Illumina). Illumina sequencing library prep was carried out according to TST170 Tumor Library Prep Kit for RNA, with the following modifications to

accommodate low RNA inputs. All reactions were reduced to 25% of original volume and the ligation adaptor was used at 1 in 10 dilution. Library quality was assessed using High Sensitivity DNA Analysis kits on the Agilent Bioanalzyer 2100 (Agilent).

Whole Exome Enrichment. Sequencing libraries were quantified using Quant-iT PicoGreen dsDNA Kit (ThermoFisher Scientific), normalized to 200 ng input and pooled to 4 samples per enrichment reaction. Whole exome enrichment was carried out according to the TruSeq RNA Access Library Prep guide (Illumina). Additional blocking oligos lacking the 5' biotin designed against hemoglobin genes HBA1, HBA2, and HBB were included in the enrichment reaction to reduce enrichment of these genes in the sequencing libraries. Final enrichment libraries were quantified using Quant-IT Picogreen dsDNA Kit (ThermoFisher Scientific), normalized and pooled for paired end 50 by 50 sequencing on Illumina HiSeq 2000 platforms to a minimum depth of 40 million reads per sample.

Data Analysis. Unless otherwise noted, all statistical testing was two-sided. Non-parametric testing was used when data were not normally distributed. Sequencing reads were mapped to human reference genome (hg19) with tophat (v2.0.13), and transcript abundance quantified with featureCounts (subread-1.4.6) against RefGene coordinates (obtained 10/27/2014). Tissue expression data were obtained from Body Atlas (CorrelationEngine, BaseSpace, Illumina, Inc) (Kupershmidt, et al. , 2010, *PLoS ONE* **5**; 10.1371/ journal.pone.0013066). vGenes with expression ≥ 2-fold higher than the median expression across all tissues in the placenta or any of the fetal tissues (brain, liver, lung, and thyroid) were assigned to that group. Subcellular localization was obtained from UniProt.

Differential expression analysis was performed in R (v3.4.2) with edgeR (v3.20.9), after exclusion of genes with a CPM ≤ 0.5 in < 25% of samples. The dataset was normalized by the TMM method, and differentially abundant genes identified by the glmTreat test for a log fold change ≥ 1 followed by Bonferroni-Holm p-value correction. The same process was used for each jackknifing iteration, using 90% of samples in each group selected by random sampling without replacement. After 1,000 jackknifing iterations, the one-sided 95% confidence interval for gene-wise p-values was calculated with statsmodels (v0.8.0). Hierarchical clustering analysis was performed with squared Euclidean distance and average linkage.

AdaBoost was performed in python with scikit-learn (v0.19.1, sklearn.ensemble. AdaBoostClassifier). Optimal hyperparameter values (90 estimators, 1.6 learning rate) were

determined by grid search, using Matthew's correlation coefficient to quantify performance. The overall AdaBoost model development strategy is illustrated in Figs. 31A-31E. Datasets (TMM-normalized log CPM values of genes with a CPM ≤ 0.5 in < 25% of samples) were standardized (sklearn.preprocessing. StandardScaler) prior to fitting classifiers. The same scaler fit on training data was applied to the corresponding testing dataset; all 5 scalers for the 5 training datasets were averaged for use with the final model. The decision_function score was used to construct ROC curves and determine sample classification.

RT-qPCR validation assay and analysis. C-RNA was isolated and converted to cDNA from 2mls of plasma from 19 Preeclampsia (PE) and 19 matched control samples, which were selected randomly. The cDNA was pre-amplified using the TaqMan Preamp master Mix (cat: 4488593) for 16 cycles and diluted 10-fold to a final volume of 500µL. For qPCR, the reaction mixture contained 5µL of diluted pre-amplified cDNA, 10µL of TaqMan gene expression master mix (cat: 4369542), 1µL of TaqMan Probe, and 4µL of water using the manufacturer's instructions. For each TaqMan probe (Table 9), three qPCR reactions were carried out per diluted cDNA sample and the *Cq* values were determined using Bio-Rad CFX manager software. To determine gene abundance for each target gene, the $\Delta\Delta Cq = 2^{-(\text{target } Cq - \text{ref } Cq_{avg})}$ was calculated using the mean Cq values between five reference gene probes (ref Cq_{avg}). To determine the fold change (PE/CTRL) for each probe, the ΔΔCq values for each sample was divided by the average ΔΔCq value for the matched control group.

Tube type study. To assess the effects of tube type and overnight shipping on circulating RNA quality, blood was drawn from pregnant and non-pregnant females in the following tube types: K2 EDTA (Beckton Dickinson), ACD (Beckton Dickinson), Cell Free RNA BCT tube (Streck), and 1 Cell Free DNA BCT tube (Streck). 8 mL of blood was drawn into each tube and shipped overnight either on ice packs (EDTA and ACD) or shipped at room temperature (Cell Free RNA and DNA BCT tubes). All shipped blood tubes were processed into plasma within 24 hours of the blood draw. As a reference, 8 mL of blood was also drawn into K2 EDTA tubes and processed within 4 hours into plasma on site and shipped as plasma on dry ice. All plasma processing and circulating RNA extraction was carried out as described in the methods section. 3 mL of plasma was used per condition to generate sequencing libraries for enrichment using Illumina protocols as described.

Reproducibility Study. Plasma was obtained from 10 individuals and split into 4 mL, 1 mL, and 0.5 mL volumes, with 3 replicates for each volume. Circulating RNA extraction (Qiagen Circulating Nucleic Acid Kit) and random primed cDNA synthesis were carried out on all samples as previously described. For libraries using 4.5 mL plasma inputs, sequencing libraries were generated using the TST170 Tumor Library Prep Kit as described above. For 1 mL and 0.5 mL inputs, the Accel-NGS 1S Plus DNA Library Kit (Swift Biosciences) was used to generate libraries. Whole exome enrichment and sequencing was carried out on all samples using as described above.

Discussion

This study focused on identifying differences that are universal to early-onset PE, supporting the ultimate goal of clinically actionable biomarker discovery. This required tailoring the analysis methods to account for the variability observed in the data. This variance stems from both the substantial biological noise in C-RNA measurements as well as the phenotypic diversity of PE. C-RNA is inherently more variable than single tissue transcriptomics because it represents a combination of cell death, signaling, and gene expression across all organs. Furthermore, PE exhibits a wide range of maternal and fetal outcomes which may be associated with different underlying molecular causes. While the genes that were eliminated may be biologically relevant in PE, they were not universal in the cohort. Interestingly, the excluded transcripts were elevated in specific women, who may represent a molecular subset of PE. Larger cohorts will help elucidate if C-RNA can delineate PE subtypes, which is crucial to understanding the diverse pathophysiology of this condition.

The most universal set of transcripts was identified by AdaBoost. The success of this method was underscored by highly accurate classification of an independent early-onset PE cohort (PEARL). These samples were collected from a different population with significantly relaxed inclusion and exclusion criteria, for instance including women in the control group who had chronic hypertension, gestational diabetes, or Alport syndrome – none of which were misidentified as having PE. In contrast to hierarchical clustering, 17 of 24 individuals from the late-onset PE cohort were correctly classified by machine learning model of this example, surprising given the suggestion that early- and late-onset PE are distinct conditions. The findings of this example suggest there may be some pathways universally altered in all PE.

In every assessment, C-RNA revealed changes in placental, fetal, and maternally expressed transcripts. One of the most striking trends observed in PE samples was the increased abundance of myriad ECM remodeling and cell migration/invasion proteins (FAM107A, SLC9A3R2, TIMP4, ADAMTS1, PRG2, TIMP3, LEP, ADAMTS2, ZEB1, HSPA12B), tracking with dysfunctional extravillous trophoblast invasion and remodeling of maternal vessels characteristic in this disease. The maternal side of early-onset PE manifests as cardiovascular dysfunction, inflammation, and preterm birth (PNMT, ZEB1, CRH), all of which show molecular signs of aberrant behavior in the data of this example.

Table 5: PEARL Control Cohort Gestational Age Distribution for 45 healthy pregnancies

Control Groups	**Recruitment**	**Follow up visit #1**	**Follow up visit #2**	**Follow up visit #3**
Group 1 (n=15)	$11^{0/7}$-$13^{6/7}$ weeks	$14^{0/7}$-$17^{6/7}$ weeks	$26^{0/7}$-$28^{6/7}$ weeks	$35^{0/7}$-$37^{6/7}$ weeks
Group 2 (n=15)	$11^{0/7}$-$13^{6/7}$ weeks	$18^{0/7}$-$21^{6/7}$ weeks	$29^{0/7}$-$31^{6/7}$ weeks	$35^{0/7}$-$37^{6/7}$ weeks
Group 3 (n=15)	$11^{0/7}$-$13^{6/7}$ weeks	$22^{0/7}$-$25^{6/7}$ weeks	$32^{0/7}$-$34^{6/7}$ weeks	$35^{0/7}$-$37^{6/7}$ weeks

Table 6: Diagnostic Criteria for Preeclampsia with Severe Features and Inclusion/Exclusion Criteria

Blood Pressure	1) Systolic BP ≥160mmHG or diastolic BP≥110mmHg measured on at least 2 occasions 4 hours apart while on bedrest but before the onset of labor or measured on 1 occasion only, if antihypertensive therapy is initiated due to severe hypertension
Proteinuria	Measured by one of the following: 1) Excretion of ≥300 mg of protein in a 24hr period 2) Protein/creatinine value of at least 0.3 3) qualitative determination with urine dipstick of ≥1+
OR	
Blood Pressure	1) Systolic BP ≥140 mmHg or diastolic ≥ 90 mmHG
With one of the following features	1) Thombocytopenia (<100,000 platelets/mL) 2) Impaired liver function 3) Newly developed renal insufficiency 4) Pulmonary edema 5) New onset cerebral disturbances or scotomata

Preeclampsia Cohort

Inclusion Criteria	Exclusion Criteria
1. Women 18 years of age or older	1. Known Malignancy
2. Pregnant women with a viable singleton	2. History of maternal organ or bone marrow

gestation
3. Gestational age between 20 0/7 and 33 6/7 weeks determined by ultrasound and/or LMP per ACOG guidelines.
4. Preeclampsia diagnosed with severe features per ACOG guidelines

transplant
3. Maternal blood transfusion in the last 8 weeks
4. Chronic Hypertension diagnosed prior to current pregnancy
5. Type I, II or gestational diabetes
6. Fetal anomaly or known chromosome abnormality
7. Active Labor

Control Cohort

Inclusion Criteria
1. Women 18 years of age or older
2. Pregnant women with a viable singleton gestation
3. Gestational age between 20 0/7 and 33 6/7 weeks determined by ultrasound and/or LMP per ACOG guidelines.

Exclusion Criteria
1. Known Malignancy
2. History of maternal organ or bone marrow transplant
3. Maternal blood transfusion in the last 8 weeks
4. Chronic Hypertension diagnosed prior to current pregnancy
5. Type I, II or gestational diabetes
6. Fetal anomaly or known chromosome abnormality
7. Active Labor
8. Thrombocytopenia (<100,000 plts/mL)
9. Impaired liver function
10. Newly developed renal insufficiency (serum creatine > 1.1 mg/dL)
11. Pulmonary edema
12. New Onset cerebral disturbances or scotomata
13. Preeclampsia in prior or current pregnancy
14. Fetal growth restriction

Table 7: Study characteristics for Illumina Preeclampsia Cohort

		Early Onset PE Cohort	Control Cohort
	Sample Size	n=40	n=73
Gestational Age at Sample Collection (weeks.days)		30.5 (+/- 2.6)	30.5 (+/- 2.6)
Maternal Characteristics			
Ethnicity	(% Hispanic)	35% (n =14)	41.1% (n=30)
Race	% Caucasian	35% (n =14)	46.6% (n=34)
	% African American	27.5% (n =11)	17.8% (n=13)
	% Asian	7.5% (n=3)	13.7% (n=10)
	% Unknown	30% (n=12)	20.5% (n=15)
	% Other	0.0%	1.4% (n=1)
Maternal Age (years, mean +/- SD)		30.4 (+/-5.7)	29.7 (+/-5.3)
Maternal BMI (kg/m2, mean+/-SD)		34.2 (+/-5.8)	30.1 (+/-5.6)
Gravida (% Nulliparous)		32.5% (n=13)	38.4% (n=28)
Chronic Hypertension		0% (n=0)	0% (n=0)
Type I, II Diabetes		0% (n=0)	0% (n=0)
Gestational Diabetes		0% (n=0)	0% (n=0)
Birth Outcomes			
Gestational Age at Birth (weeks.days)		31.5 (+/- 3.1)	38.9(+/-1.8)
Full Term		2.5% (n=1)	90.4% (n=66)
Preterm (<37 weeks)		97.5% (n=39)	9.6% (n=7)
Sex (% male)		37.5% (n=15)	42.5% (n=31)
Birth Weight (kg)		1.4 (+/-0.52)	3.2 (+/- 0.55)
Small for Gestational Age*		45% (n=18)	9.6%(n=7)
Stillbirth		2.5% (n=1)	0% (n=0)
Medications for treatment of:			
PE/Hypertension	MgSO4	82.5% (n=33)	4.1% (n=3)
	Antenatal Steroids	95.0% (n=38)	6.8% (n=5)
	Anti-Hypertensive	75.0% (n=30)	5.3% (n=4)
	Aspirin	20.0% (n=8)	0% (n=0)
Other Conditions	Analgesics	60.0% (n=24)	11.8% (n=9)
	Antimicrobials	12.5% (n=5)	5.5% (n=4)
	Antihistamines	32.5% (n=13)	13.7% (n=10)
	Asthma	10.0% (n=4)	2.7% (n=2)
	Psychoactive	15.0% (n=6)	5.5% (n=4)
	Hypothyroidism	7.5% (n=3)	2.7% (n=2)
Pregnancy Symptoms	Antiemetics	25.0% (n=10)	5.5% (n=4)
	Antacids	27.5% (n=11)	8.2% (n=6)
	Anti-constipation	15.0% (n=6)	11.8% (n=9)
	Prenatal Vitamins	17.5% (n=7)	31.5% (n=23)
	Iron Supplement	10% (n=4)	12.3% (n=9)

*Defined as birthweight <10% of population for male or female fetus

Table 8: Medical Center Collection Site Patient Distribution

Clinical Site	**Location**	**Number PE patients**	**Number of controls**
University of Texas Medical Branch	Galveston, Texas	4	11
Tufts Medical Center	Boston, MA	10	17
Columbia University Irving Medical Center	New York, NY	4	9
Winthrop University Hospital	Mineola, NY	5	9
St. Peter's University Hospital	New Brunswick, NJ	3	6
Christiana Care	Newark, DE	7	13
Rutgers University Robert Wood Johnson Medical School	New Brunswick, NJ	5	8
New York Presbyterian/Queens	New York, NY	2	3
	Total Samples collected	**40**	**76**

Table 9: Genes validated by TaqMan qPCR

Gene Name	Assay ID	Type	RefSeq
ABHD12	Hs01018050_m1	Reference	NM_001042472.2
ABHD12	Hs01018050_m1	Target	NM_001042472.2
ADAMTS2	Hs01029111_m1	Target	NM_014244.4
ALOX15B	Hs00153988_m1	Target	NM_001039130.1
ARHGEF25	Hs00384780_g1	Target	NM_001111270.2
ARRDC2	Hs01006434_g1	Target	NM_001286826.1
CLEC4C	Hs01092460_m1	Target	NM_130441.2
DAAM2	Hs00322497_m1	Target	NM_001201427.1
FAM107A	Hs00200376_m1	Target	NM_001076778.2
HSPA12B	Hs00369554_m1	Target	NM_001197327.1
HTRA4	Hs00538137_m1	Target	NM_153692.3
IGFBP5	Hs00181213_m1	Target	NM_000599.3
KRBOX4	Hs01063506_gH	Reference	NM_001129898.1
KRT5	Hs00361185_m1	Target	NM_000424.3
LEP	Hs00174877_m1	Target	NM_000230.2
NES	Hs00707120_s1	Target	NM_006617.1
NME3	Hs01573872_g1	Reference	NM_002513.2
PAPPA2	Hs01060983_m1	Target	NM_020318.2
PITPNM3	Hs01107787_m1	Target	NM_001165966.1
PLD4	Hs00975488_m1	Target	NM_001308174.1
PRG2	Hs00794928_m1	Target	NM_001243245.2
TIMP3	Hs00165949_m1	Target	NM_000362.4
TIMP4	Hs00162784_m1	Target	NM_003256.3
VSIG4	Hs00907325_m1	Target	NM_001184830.1
WNT7A	Hs00171699_m1	Reference	NM_004625.3
ZEB1	Hs01566408_m1	Target	NM_001128128.2
ZNF138	Hs00864088_gH	Reference	NM_001271638.1

Table 10

Gene Symbol	Analysis	Previous Literature Reports	Change in PE	Tissue Expression Category	Sub-Cellular Location	Function(s)
ARRDC2	AdaBoost	No	Increase*	Other (Skeletal Muscle; Globus Pallidus; Lung)	Membrane	Protein Trafficking
ALOX15B	DEX	Yes	Increase	Fetal	Nucleus; Cytoskeleton; Cytosol; Membrane	Cell Cycle; Immune Function; Cardiovascular Function
AMPH	DEX	No	Increase	Fetal	Cytoskeleton; Membrane	Synaptic Vesicle Endocytosis
CUX2	DEX	No	Decrease	Fetal	Nucleus	Cell Cycle; Fetal Development; DNA Damage Response
FAM107A	DEX	No	Increase	Fetal	Cytoskeleton; Membrane; Nucleus	Cell Migration/Invasion; Cell Cycle; ECM Regulation
IGFBP5	DEX	Yes	Increase	Fetal	Extracellular or Secreted	Fetal Development; IGF Signaling
NES	DEX	Yes	Increase	Fetal	Cytoskeleton	Fetal Development; Cell Cycle
PITPNM3	DEX	No	Increase	Fetal	Membrane	Phosphatidylinositol Regulation
PRX	DEX	Yes	Increase	Fetal	Membrane	Cell Structure/Composition
TEAD4	DEX	Yes	Increase	Fetal	Nucleus	Placental Development
PNMT	DEX	Yes	Increase	Other (Adrenal Gland Cortex; Adrenal Gland; Skeletal Muscle Psoas)	Cytosol	Epinephrine Synthesis; Cardiovascular Function; Pregnancy Duration
DAAM2	DEX	Yes	Increase	Other (Corpus Callosum; Globus Pallidum External; Nodose Nucleus)	Extracellular or Secreted	Fetal Development
SLC9A3R2	DEX	No	Increase	Other (Heart Ventricle; Liver; Parotid Gland)	Membrane; Nucleus	ECM Regulation; Cell Structure/Composition
HSPA12B	DEX	No	Increase	Other (Heart Ventricle; Lung; Spleen)	unknown	Angiogenesis; Cardiovascular Function; Cell Migration/Invasion; Hypoxia Response
PLD4	DEX	No	Decrease	Other (Nodose Nucleus; Subthalamic Nucleus; Corpus Callosum)	Membrane	Phosphatidylinositol Regulation; Immune Function
TIMP4	DEX	No	Increase	Other (Omental Adipose Tissue; Subcutaneous Adipose Tissue; Joint Synovium)	Extracellular or Secreted	ECM Regulation; Immune Function
KRT5	DEX	Yes	Decrease	Other (Oral Mucosa; Pharyngeal Mucosa; Esophagus)	Cytoskeleton	Cell Structure/Composition
ZEB1	DEX	No	Increase	Other (Synovial Membrane; Aorta; Myometrium)	Nucleus	Immune Function; Cell Migration/Invasion; Fetal Development; Pregnancy Duration
APOLD1	DEX	Yes	Increase	Placental	Plasma Membrane	Angiogenesis; Cardiovascular Function; Hypoxia Response; Fetal Development
HTRA4	DEX	Yes	Increase	Placental	Extracellular or Secreted	IGF Signaling; Placental Development
SEMA3G	DEX	No	Increase	Placental	Extracellular or Secreted	Cell Migration/Invasion
ADAMTS1	DEX	Yes	Increase	Placental/Fetal	Extracellular or Secreted	ECM Regulation; Fetal Development; Angiogenesis
CRH	DEX	Yes	Increase	Placental/Fetal	Extracellular or Secreted	Pregnancy Duration; Fetal Development; Cardiovascular Function
PRG2	DEX	Yes	Increase	Placental/Fetal	Extracellular or Secreted	Immune Function; ECM Regulation; IGF Signaling
TIMP3	DEX	Yes	Increase	Placental/Fetal	Extracellular or Secreted	ECM Regulation; Immune Function; Angiogenesis
ARHGEF25	DEX & AdaBoost	No	Increase	Other (Hippocampus; Myometrium; Cerebellum)	Membrane; Sarcomere	
CLEC4C	DEX & AdaBoost	Yes	Decrease	Other (Rectum Colon; Ascending Colon; Substantia Nigra Reticulata)	Membrane	Immune Function
LEP	DEX & AdaBoost	Yes	Increase	Placental	Extracellular or Secreted	Energy Homeostasis; Immune Function; Angiogenesis; Fetal Development; ECM Regulation
PAPPA2	DEX & AdaBoost	Yes	Increase	Placental	Extracellular or Secreted	Fetal Development; IGF Signaling
VSIG4	DEX & AdaBoost	Yes	Increase	Placental	Membrane	Immune Function
ADAMTS2	DEX & AdaBoost	No	Increase	Placental/Fetal	Extracellular or Secreted	ECM Regulation; Angiogenesis; Fetal Development

Key

Increase* indicates the change was not statistically different

CorrelationEngine Body Atlas was used to find the 3 top tissues expressing genes in the "Other" category

UniProt was used to determine sub-cellular localization...as a note; I merged all "membrane" classifications to one category (so Plasma Membrane; ER Membrane; etc are not distinct)

Table 11: Study characteristics for Illumina Preeclampsia Cohort

		Early Onset PE	Early Onset Control	Late Onset PE	Late Onset Control
Sample Size		n=12	n=12	n=12	n=12
Gestational Age at Sample Collection (weeks.days)		29.2 (+/- 2.3)	29.3 (+/-2.3)	35.6 (+/-1.3)	35.9 (+/-0.8)
Maternal Characteristics					
Ethnicity	(% Hispanic)	0% (n=0)	0% (n=0)	0% (n=0)	0% (n=0)
Race	% Caucasian	91.7% (n=11)	100% (n=12)	100% (n=12)	100% (n=12)
	% African	8.3% (n=1)	0% (n=0)	0% (n=0)	0% (n=0)
Maternal Age (years, mean +/- SD)		29.3 (+/- 3.5)	30.1 (+/- 3.8)	30.2 (+/- 4.8)	29.4 (+/- 3.2)
Maternal BMI (kg/m2, mean+/-SD)		33.6 (+/- 9.0)	28.5 (+/- 7.0)	32.2 (+/- 4.9)	27.9 (+/- 4.5)
Gravida (% Nulliparous)		60% (n=6)	58.3% (n=7)	75% (n=9)	75% (n=9)
Chronic Hypertension		13.3% (n=2)	8.3% (n=1)	8.3% (n=1)	0% (n=0)
Type I, II Diabetes		13.3% (n=2)	0% (n=0)	25.0% (n=3)	0% (n=0)
Gestational Diabetes		13.3% (n=2)	33.3% (n=4)	8.3% (n=1)	16.7% (n=2)
Other Health Condition		0% (n=0)	8.3% (n=1)	0% (n=0)	0% (n=0)
Birth Outcomes					
Gestational Age at Birth (weeks.days)		30.3 (+/- 3.4)	39.0 (+/-1.5)	37.0 (+/- 1.4)	39.7 (+/- 1.6)
Full Term		0% (n=0)	91.7% (n=11)	75.0% (n=9)	91.7% (n=11)
Preterm (<37 weeks)		100% (n=12)	8.3% (n=1)	25.0% (n=3)	8.3% (n=1)
Sex (% male)		75% (n=9)	58.3% (n=7)	66.7% (n=8)	58.3% (n=7)
Birth Weight (kg)		1.3 (+/- 0.54)	3.2 (+/-0.40)	2.7 (+/- 0.55)	3.4 (+/- 0.54)
Fetal Growth Restriction		50.0% (n=6)	0% (n=0)	8.3% (n=1)	0% (n=0)
Small for Gestational Age*		25% (n=3)	0% (n=0)	33.3% (n=4)	25% (n=3)
Stillbirth		0% (n=0)	0% (n=0)	0% (n=0)	0% (n=0)
HELLP		25.0% (n=3)	0% (n=0)	0% (n=0)	0% (n=0)
Medications for treatment of:					
PE/Hypertension	MgSO4	83.3% (n=10)	0% (n=0)	33.3% (n=4)	0% (n=0)
	Antenatal Steroids	100% (n=12)	0% (n=0)	25.0% (n=3)	0% (n=0)
	Anti-Hypertensive	100% (n=12)	8.3% (n=1)	75% (n=9)	0% (n=0)
	Aspirin	8.3% (n=1)	25.0% (n=3)	25.0% (n=3)	8.3% (n=1)

*Defined as birthweight <10% of population for male or female fetus

The complete disclosure of all patents, patent applications, and publications, and electronically available material (including, for instance, nucleotide sequence submissions in, e.g., GenBank and RefSeq, and amino acid sequence submissions in, e.g., SwissProt, PIR, PRF, PDB, and translations from annotated coding regions in GenBank and RefSeq) cited herein are incorporated by reference in their entirety. Supplementary materials referenced in publications (such as supplementary tables, supplementary figures, supplementary materials and methods, and/or supplementary experimental data) are likewise incorporated by reference in their entirety. In the event that any inconsistency exists between the disclosure of the present application and the disclosure(s) of any document incorporated herein by reference, the disclosure of the present application shall govern. The foregoing detailed description and examples have been given for clarity of understanding only. No unnecessary limitations are to be understood therefrom. The disclosure is not limited to the exact details shown and described, for variations obvious to one skilled in the art will be included within the disclosure defined by the claims.

Unless otherwise indicated, all numbers expressing quantities of components, molecular weights, and so forth used in the specification and claims are to be understood as being modified in all instances by the term “about.” Accordingly, unless otherwise indicated to the contrary, the numerical parameters set forth in the specification and claims are approximations that may vary depending upon the desired properties sought to be obtained by the present disclosure. At the very least, and not as an attempt to limit the doctrine of equivalents to the scope of the claims, each numerical parameter should at least be construed in light of the number of reported significant digits and by applying ordinary rounding techniques.

Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the disclosure are approximations, the numerical values set forth in the specific examples are reported as precisely as possible. All numerical values, however, inherently contain a range necessarily resulting from the standard deviation found in their respective testing measurements.

All headings are for the convenience of the reader and should not be used to limit the meaning of the text that follows the heading, unless so specified.

What is claimed is:

Claim 1. A method of detecting preeclampsia in a pregnant female, the method comprising:

identifying in a biosample obtained from the pregnant women a plurality of circulating RNA (C-RNA) molecules;

wherein the biosample comprises blood, plasma, or serum;

wherein the identification of a plurality of C-RNA molecules is selected from:

(a) a plurality of C-RNA molecules encoding a coding region of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, any fifty or more, or all of arrestin domain containing 2 (ARRDC2), Jun proto-oncogene, AP-1 transcription factor subunit (JUN), ski like proto-oncogene (SKIL), ATPase 13A3 (ATP13A3), phosphodiesterase 8B (PDE8B), glutathione S-transferase alpha 3 (GSTA3), TCDD inducible poly (ADP-ribose) polymerase (TIPARP), RGP1 homolog, RAB6A GEF complex partner 1 (RGP1), ubiquitin specific peptidase 54 (USP54), C-type lectin domain family 4 member C (CLEC4C), mitochondrial ribosomal protein S35 (MRPS35), Rho guanine nucleotide exchange factor 25 (ARHGEF25), cut like homeobox 2 (CUX2), HEAT repeat containing 9 (HEATR9), follistatin like 3 (FSTL3), DNA damage inducible 1 homolog 2 (DDI2), zinc finger MYM-type containing 6 (ZMYM6), ST6 N-acetylgalactosaminide alpha-2,6-sialyltransferase 3 (ST6GALNAC3), guanylate binding protein 2 (GBP2), ETS variant 3 (ETV3), ADAM metallopeptidase domain 17 (ADAM17), atonal bHLH transcription factor 8 (ATOH8), solute carrier family 4 member 3 (SLC4A3), TRAF3 interacting protein 1 (TRAF3IP1),

tetratricopeptide repeat domain 21A (TTC21A), heart development protein with EGF like domains 1 (HEG1), asteroid homolog 1 (ASTE1), transmembrane protein 108 (TMEM108), ectodermal-neural cortex 1 (ENC1), secretory carrier membrane protein 1 (SCAMP1), arrestin domain containing 3 (ARRDC3), solute carrier family 26 member 2 (SLC26A2), slit guidance ligand 3 (SLIT3), chloride intracellular channel 5 (CLIC5), TNF receptor superfamily member 21 (TNFRSF21), protein phosphatase 1 regulatory subunit 17 (PPP1R17), tyrosylprotein sulfotransferase 1 (TPST1), cytosolic arginine sensor for mTORC1 subunit 2 (GATSL2), speedy/RINGO cell cycle regulator family member E5 (SPDYE5), homeodomain interacting protein kinase 2 (HIPK2), MT-RNR2 like 6 (MTRNR2L6), chloride voltage-gated channel 1 (CLCN1), GINS complex subunit 4 (GINS4), twinkle mtDNA helicase (C10orf2), TruB pseudouridine synthase family member 1 (TRUB1), aminoacylase 3 (ACY3), fatty acyl-CoA reductase 2 (FAR2), CD63 molecule (CD63), cytoskeleton associated protein 4 (CKAP4), two pore segment channel 1 (TPCN1), ring finger protein 6 (RNF6), thiamine triphosphatase (THTPA), Fos proto-oncogene, AP-1 transcription factor subunit (FOS), poly(A)-specific ribonuclease (PARN), ORAI calcium release-activated calcium modulator 3 (ORAI3), engulfment and cell motility 3 (ELMO3), sphingomyelin phosphodiesterase 3 (SMPD3), serpin family F member 1 (SERPINF1), transmembrane protein 11 (TMEM11), proteasome 26S subunit, non-ATPase 11 (PSMD11), C-type lectin domain family 4 member M (CLEC4M), coiled-coil domain containing 151 (CCDC151), C3 and PZP like alpha-2-macroglobulin domain containing 8 (CPAMD8), cornifelin (CNFN), eukocyte immunoglobulin like receptor A4 (LILRA4), adenosine deaminase (ADA), chromosome 22 open reading frame 39 (C22orf39), phosphatidylinositol 4-kinase alpha pseudogene 1 (PI4KAP1), and ADP ribosylation factor GTPase activating protein 3 (ARFGAP3); or

(b) a plurality of C-RNA molecules encoding a coding region of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, or all twenty of TIMP metallopeptidase inhibitor 4 (TIMP4), filaggrin (FLG), HtrA serine peptidase 4 (HTRA4), amphiphysin (AMPH), lipocalin 6 (LCN6), TEA domain transcription factor 4 (TEAD4), age-related maculopathy susceptibility 2 (ARMS2), semaphorin 3G (SEMA3G), ADAM metallopeptidase with thrombospondin type 1 motif 1 (ADAMTS1), arachidonate 15-lipoxygenase type B (ALOX15B), SLC9A3 regulator 2 (SLC9A3R2), insulin like growth factor binding protein 5 (IGFBP5), heat shock protein family A (Hsp70) member 12B (HSPA12B), C-type lectin domain family 4 member C (CLEC4C), keratin 5 (KRT5), periaxin (PRX), Rho guanine nucleotide exchange factor 25 (ARHGEF25), α disintegrin and metallopeptidase with thrombospondin type 1 motif 2 (ADAMTS2), dishevelled associated activator of morphogenesis 2 (DAAM2), and family with sequence similarity 107 member A (FAM107A); or

(c) a plurality of C-RNA molecules encoding a coding region of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, any fifty or more, any seventy-five or more, any one hundred or more, or all of cytochrome P450 family 26 subfamily B member 1 (CYP26B1), interferon regulatory factor 6 (IRF6), myosin heavy chain 14 (MYH14), podocalyxin like (PODXL), protein phosphatase 1 regulatory subunit 3C (PPP1R3C), SH3 domain containing ring finger 2 (SH3RF2), transmembrane channel like 7 (TMC7), zinc finger protein 366

(ZNF366), adenylate cyclase 1 (ADCY1), complement C6 (C6), family with sequence similarity 219 member A (FAM219A), hydroxyacid oxidase 2 (HAO2), IgA inducing protein (IGIP), interleukin 1 receptor type 2 (IL1R2), neurotrophic receptor tyrosine kinase 2 (NTRK2), SH3 and PX domains 2A (SH3PXD2A), ssu-2 homolog (SSUH2), sulfotransferase family 2A member 1 (SULT2A1), flavin containing dimethylaniline monoxygenase 3 (FMO3), follistatin like 3 (FSTL3), GATA binding protein 5 (GATA5), HtrA serine peptidase 1 (HTRA1), complement C8 beta chain (C8B), MN1 proto-oncogene, transcriptional regulator (MN1), nuclear factor, erythroid 2 like 1 (NFE2L1), PR/SET domain 16 (PRDM16), adaptor related protein complex 3 subunit beta 2 (AP3B2), epithelial membrane protein 1 (EMP1), filamin C (FLNC), stromal antigen 3 (STAG3), carboxypeptidase B2 (CPB2), tensin 2 (TENC1), RP1 like 1 (RP1L1), APOBEC1 complementation factor (A1CF), natriuretic peptide receptor 1 (NPR1), TEK receptor tyrosine kinase (TEK), ERBB receptor feedback inhibitor 1 (ERRFI1), Rho guanine nucleotide exchange factor 15 (ARHGEF15), CD34 molecule (CD34), R-spondin 3 (RSPO3), alpha kinase 3 (ALPK3), sterile alpha motif domain containing 4A (SAMD4A), zinc finger CCHC-type containing 24 (ZCCHC24), liver enriched antimicrobial peptide 2 (LEAP2), myosin light chain 2 (MYL2), neuregulin 3 (NRG3), zinc finger and BTB domain containing 16 (ZBTB16), serpin family A member 3 (SERPINA3), aquaporin 7 (AQP7), sushi repeat containing protein X-linked (SRPX), uveal autoantigen with coiled-coil domains and ankyrin repeats (UACA), anoctamin 1 (ANO1), FKBP prolyl isomerase 5 (FKBP5), sodium voltage-gated channel alpha subunit 5 (SCN5A), protein tyrosine phosphatase non-receptor type 21 (PTPN21), calcium voltage-gated channel subunit alpha1 C (CACNA1C), ETS transcription factor ERG (ERG), SRY-box 17 (SOX17), WW domain containing transcription regulator 1 (WWTR1), allograft inflammatory factor 1 like (AIF1L), carbonic anhydrase 3 (CA3), histidine rich glycoprotein (HRG), tyrosine aminotransferase (TAT), aquaporin 7 pseudogene 1 (AQP7P1), adrenoceptor alpha 2C (ADRA2C), synaptopodin (SYNPO), adhesion G protein-coupled receptor F5 (GPR116), keratin 17 (KRT17), alpha-2-glycoprotein 1, zinc-binding (AZGP1), BCL6B

transcription repressor (BCL6B), kinesin family member 1C (KIF1C), chloride intracellular channel 5 (CLIC5), G protein-coupled receptor 4 (GPR4), gap junction protein alpha 5 (GJA5), oleoyl-ACP hydrolase (OLAH), armadillo like helical domain containing 4 (C14orf37), zinc finger E-box binding homeobox 1 (ZEB1), jagged canonical Notch ligand 2 (JAG2), kinesin family member 26A (KIF26A), apolipoprotein L domain containing 1 (APOLD1), phenylethanolamine N-methyltransferase (PNMT), myomesin 3 (MYOM3), PITPNM family member 3 (PITPNM3), TIMP4, HTRA4, AMPH, LCN6, TEAD4, ARMS2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, IGFBP5, HSPA12B, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, hemoglobin subunit gamma 2 (HBG2), cell adhesion molecule 2 (CADM2), lysosomal associated membrane protein family member 5 (LAMP5), prostaglandin D2 receptor 2 (PTGDR2), NODAL modulator 1 (NOMO1), nuclear RNA export factor 3 (NXF3), phospholipase D family member 4 (PLD4), BPIFB3, protein kinase C and casein kinase substrate in neurons 1 (PACSIN1), CUX2, FLG, CLEC4C, and KRT5; or

(d) a plurality of C-RNA molecules encoding a coding region of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or all twenty-three of ADAMTS2, FAM107A, DAAM2, ARHGEF25, , PRX, ALOX15B, HSPA12B, IGFBP5, CLEC4C, SLC9A3R2, ADAMTS1, SEMA3G, KRT5, AMPH, TEAD4, PITPNM3, TIMP4, PNMT, ZEB1, APOLD1, PLD4, CUX2, and HTRA4; or

(e) a plurality of C-RNA molecules encoding a coding region of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more,

any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, or all twenty-one of ADAMTS1, ADAMTS2, ALOX15B, AMPH, ARHGEF25, CUGBP Elav-like family member 4 (CELF4), DAAM2, FAM107A, HSPA12B, HTRA4, IGFBP5, potassium voltage-gated channel subfamily A member 5 (KCNA5), KRT5, LCN6, leucine rich repeat containing 26 (LRRC26), OLAH, PACSIN1, PRX, PTGDR2, SEMA3G, and SLC9A3R2; or

(f) a plurality of C-RNA molecules encoding a coding region of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or all of ADAMTS1, ADAMTS2, ALOX15B, ARHGEF25, CELF4, DAAM2, FAM107A, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LRRC26, OLAH, PRX, PTGDR2, SEMA3G, and SLC9A3R2; or

(g) a plurality of C-RNA molecules encoding a coding region of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, or all six of CLEC4C, ARHGEF25, ADAMTS2, ARRDC2, SKIL, and arrestin domain containing 4 (ARRDC4); or

(h) a plurality of C-RNA molecules encoding a coding region of a protein selected from any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or all twenty of KCNA5, ADAMTS2, MYOM3, ATP13A3, ARHGEF25, ADA, HTRA4, ACY3, PLD4, secretin (SCT), NADPH oxidase 4 (NOX4), PACSIN1, SERPINF1, SKIL, SEMA3G, TIPARP, LRRC26, phosphate regulating endopeptidase homolog X-linked (PHEX), LILRA4, and period circadian regulator 1 (PER1), is indicative of preeclampsia in the pregnant women.

Claim 2. A method of detecting preeclampsia in a pregnant female, the method comprising:

purifying a population of circulating RNA (C-RNA) molecules from a biosample obtained from the pregnant female;

wherein the biosample comprises blood, plasma, or serum;

identifying protein coding sequences encoded by the C-RNA molecules within the purified population of C-RNA molecules;

wherein the identification of protein coding sequences encoded by the C-RNA molecules encoding a coding region of a protein selected from:

(a) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, any fifty or more, or all of ARRDC2, JUN, SKIL, ATP13A3, PDE8B, GSTA3, TIPARP, RGP1, USP54, CLEC4C, MRPS35, ARHGEF25, CUX2, HEATR9, FSTL3, DDI2, ZMYM6, ST6GALNAC3, GBP2, ETV3, ADAM17, ATOH8, SLC4A3, TRAF3IP1, TTC21A, HEG1, ASTE1, TMEM108, ENC1, SCAMP1, ARRDC3, SLC26A2, SLIT3, CLIC5, TNFRSF21, PPP1R17, TPST1, GATSL2, SPDYE5, HIPK2, MTRNR2L6, CLCN1, GINS4, C10orf2, TRUB1, ACY3, FAR2, CD63, CKAP4, TPCN1, RNF6, THTPA, FOS, PARN, ORAI3, ELMO3, SMPD3, SERPINF1, TMEM11, PSMD11, CLEC4M, CCDC151, CPAMD8, CNFN, LILRA4, ADA, C22orf39, PI4KAP1, and ARFGAP3; or

(b) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, or all twenty of TIMP4, FLG, HTRA4, AMPH, LCN6, TEAD4, ARMS2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, IGFBP5, HSPA12B, CLEC4C, KRT5, PRX, ARHGEF25, ADAMTS2, DAAM2, and FAM107A; or

(c) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, any fifty or more, any seventy-five or more, any one hundred or more, or all of CYP26B1, IRF6, MYH14, PODXL, PPP1R3C, SH3RF2, TMC7, ZNF366, ADCY1, C6, FAM219A, HAO2, IGIP, IL1R2, NTRK2, SH3PXD2A, SSUH2, SULT2A1, FMO3, FSTL3, GATA5, HTRA1, C8B, MN1, NFE2L1, PRDM16, AP3B2, EMP1, FLNC, STAG3, CPB2, TENC1, RP1L1, A1CF, NPR1, TEK, ERRFI1, ARHGEF15, CD34, RSPO3, ALPK3, SAMD4A, ZCCHC24, LEAP2, MYL2, NRG3, ZBTB16, SERPINA3, AQP7, SRPX, UACA, ANO1, FKBP5, SCN5A, PTPN21, CACNA1C, ERG, SOX17, WWTR1, AIF1L, CA3, HRG, TAT, AQP7P1, ADRA2C, SYNPO, GPR116, KRT17, AZGP1, BCL6B, KIF1C, CLIC5, GPR4, GJA5, OLAH, C14orf37, ZEB1, JAG2, KIF26A, APOLD1, PNMT, MYOM3, PITPNM3, TIMP4, HTRA4, AMPH, LCN6, TEAD4, ARMS2, PAPPA2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, TIMP3, IGFBP5, HSPA12B, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, HBG2, CADM2, LAMP5, PTGDR2, NOMO1, NXF3, PLD4, BPIFB3, PACSIN1, CUX2, FLG, CLEC4C, and KRT5; or

(d) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, or all twenty-three of ADAMTS2, FAM107A, DAAM2, ARHGEF25, PRX, ALOX15B, HSPA12B, IGFBP5, CLEC4C, SLC9A3R2, ADAMTS1, SEMA3G, KRT5, AMPH, TEAD4, PITPNM3, TIMP4, PNMT, ZEB1, APOLD1, PLD4, CUX2, and HTRA4; or

(e) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, or all twenty-one of ADAMTS1, ADAMTS2, ALOX15B, AMPH, ARHGEF25, CELF4, DAAM2, FAM107A, HSPA12B, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LRRC26, OLAH, PACSIN1, PRX, PTGDR2, SEMA3G, and SLC9A3R2; or

(f) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, or all seventeen of ADAMTS1, ADAMTS2, ALOX15B, ARHGEF25, CELF4, DAAM2, FAM107A, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LRRC26, OLAH, PRX, PTGDR2, SEMA3G, and SLC9A3R2; or

(g) any one or more, any two or more, any three or more, any four or more, any five or more, or all six of CLEC4C, ARHGEF25, ADAMTS2, ARRDC2, SKIL, and ARRDC4; or

(h) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, or all twenty of KCNA5, ADAMTS2, MYOM3, ATP13A3, ARHGEF25, ADA, HTRA4, ACY3, PLD4, SCT, NOX4, PACSIN1, SERPINF1, SKIL, SEMA3G, TIPARP, LRRC26, PHEX, LILRA4, and PER1,

is indicative of preeclampsia in the pregnant women.

Claim 3. The method of claim 2, wherein identifying the plurality of C-RNA molecules or identifying protein coding sequences encoded by C-RNA molecules within the biosample comprises hybridization, reverse transcriptase PCR, microarray chip analysis, or sequencing.

Claim 4. The method of claim 3, wherein sequencing comprises:

a) massively parallel sequencing of clonally amplified molecules; or

b) RNA sequencing.

Claim 5. The method of claim any one of claims 1 to 4, further comprising:

removing intact cells from the biosample;

treating the biosample with a deoxynuclease (DNase) to remove cell free DNA (cfDNA);

synthesizing complementary DNA (cDNA) from C-RNA molecules in the biosample; and/or

enriching the cDNA sequences for DNA sequences that encode proteins by exome enrichment;

prior to identifying protein coding sequence encoded by the circulating RNA (C-RNA) molecules.

Claim 6. A method of detecting preeclampsia in a pregnant female, the method comprising:

removing intact cells from a biosample obtained from the pregnant female;

wherein the biosample comprises blood, plasma, or serum;

treating the biosample with a deoxynuclease (DNase) to remove cell free DNA (cfDNA);

synthesizing complementary DNA (cDNA) from RNA molecules in the biosample;

enriching the cDNA sequences for DNA sequences that encode proteins (exome enrichment);

sequencing the resulting enriched cDNA sequences; and

identifying protein coding sequences encoded by enriched C-RNA molecules;

wherein the identification of protein coding sequences encoded by the C-RNA molecules encoding a coding region of a protein selected from**:**

(a) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any twenty four or more, any twenty five or more, any fifty or more, or all of ARRDC2, JUN, SKIL, ATP13A3, PDE8B, GSTA3, TIPARP, RGP1, USP54, CLEC4C, MRPS35, ARHGEF25, CUX2, HEATR9, FSTL3, DDI2, ZMYM6, ST6GALNAC3, GBP2, ETV3, ADAM17, ATOH8, SLC4A3, TRAF3IP1, TTC21A, HEG1, ASTE1, TMEM108, ENC1, SCAMP1, ARRDC3, SLC26A2, SLIT3, CLIC5, TNFRSF21, PPP1R17, TPST1, GATSL2, SPDYE5, HIPK2, MTRNR2L6, CLCN1, GINS4, C10orf2, TRUB1, ACY3, FAR2, CD63, CKAP4, TPCN1, RNF6, THTPA, FOS, PARN, ORAI3, ELMO3, SMPD3, SERPINF1, TMEM11, PSMD11, CLEC4M, CCDC151, CPAMD8, CNFN, LILRA4, ADA, C22orf39, PI4KAP1, and ARFGAP3; or

(b) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, or all twenty of TIMP4, FLG, HTRA4, AMPH, LCN6, TEAD4, ARMS2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, IGFBP5, HSPA12B, CLEC4C, KRT5, PRX, ARHGEF25, ADAMTS2, DAAM2, and FAM107A; or

(c) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty one or more, any twenty two or more, any twenty three or more, any

twenty four or more, any twenty five or more, any fifty or more, any seventy-five or more, any one hundred or more, or all of CYP26B1, IRF6, MYH14, PODXL, PPP1R3C, SH3RF2, TMC7, ZNF366, ADCY1, C6, FAM219A, HAO2, IGIP, IL1R2, NTRK2, SH3PXD2A, SSUH2, SULT2A1, FMO3, FSTL3, GATA5, HTRA1, C8B, MN1, NFE2L1, PRDM16, AP3B2, EMP1, FLNC, STAG3, CPB2, TENC1, RP1L1, A1CF, NPR1, TEK, ERRFI1, ARHGEF15, CD34, RSPO3, ALPK3, SAMD4A, ZCCHC24, LEAP2, MYL2, NRG3, ZBTB16, SERPINA3, AQP7, SRPX, UACA, ANO1, FKBP5, SCN5A, PTPN21, CACNA1C, ERG, SOX17, WWTR1, AIF1L, CA3, HRG, TAT, AQP7P1, ADRA2C, SYNPO, GPR116, KRT17, AZGP1, BCL6B, KIF1C, CLIC5, GPR4, GJA5, OLAH, C14orf37, ZEB1, JAG2, KIF26A, APOLD1, PNMT, MYOM3, PITPNM3, TIMP4, HTRA4, AMPH, LCN6, TEAD4, ARMS2, SEMA3G, ADAMTS1, ALOX15B, SLC9A3R2, IGFBP5, HSPA12B, PRX, ARHGEF25, ADAMTS2, DAAM2, FAM107A, HBG2, CADM2, LAMP5, PTGDR2, NOMO1, NXF3, PLD4, BPIFB3, PACSIN1, CUX2, FLG, CLEC4C, and KRT5; or

(d) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, any twenty-one or more, any twenty-two or more, or all twenty-three of ADAMTS2, FAM107A, DAAM2, ARHGEF25, PRX, ALOX15B, HSPA12B, IGFBP5, CLEC4C, SLC9A3R2, ADAMTS1, SEMA3G, KRT5, AMPH, TEAD4, PITPNM3, TIMP4, PNMT, ZEB1, APOLD1, PLD4, CUX2, and HTRA4; or

(e) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, any twenty or more, or all twenty-one of ADAMTS1, ADAMTS2, ALOX15B, AMPH, ARHGEF25, CELF4, DAAM2, FAM107A, HSPA12B, HTRA4, IGFBP5,

KCNA5, KRT5, LCN6, LRRC26, OLAH, PACSIN1, PRX, PTGDR2, SEMA3G, and SLC9A3R2; or

(f) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, or all eighteen of ADAMTS1, ADAMTS2, ALOX15B, ARHGEF25, CELF4, DAAM2, FAM107A, HTRA4, IGFBP5, KCNA5, KRT5, LCN6, LRRC26, OLAH, PRX, PTGDR2, SEMA3G, and SLC9A3R2; or

(g) any one or more, any two or more, any three or more, any four or more, any five or more, or all six of CLEC4C, ARHGEF25, ADAMTS2, ARRDC2, SKIL, and ARRDC4; or

(h) any one or more, any two or more, any three or more, any four or more, any five or more, any six or more, any seven or more, any eight or more, any nine or more, any ten or more, any eleven or more, any twelve or more, any thirteen or more, any fourteen or more, any fifteen or more, any sixteen or more, any seventeen or more, any eighteen or more, any nineteen or more, or all twenty of KCNA5, ADAMTS2, MYOM3, ATP13A3, ARHGEF25, ADA, HTRA4, ACY3, PLD4, SCT, NOX4, PACSIN1, SERPINF1, SKIL, SEMA3G, TIPARP, LRRC26, PHEX, LILRA4, and PER1,

is indicative of preeclampsia in the pregnant women.

Claim 7. The method of any one of claims 1 to 6, wherein the biosample is obtained from a pregnant female: at less than 16 weeks gestation; at less than 20 weeks gestation; or at greater than 20 weeks gestation.

Claim 8. The method of any one of claims 1 to 7, wherein the biosample is a blood sample and the blood sample is collected, shipped, and/or stored in a tube that has cell- and DNA-stabilizing properties prior to processing the blood sample into plasma.

Claim 9. The method of claim 1 to 7, wherein the biosample is a blood sample and the blood sample is:

not exposed to EDTA prior to processing the blood sample into plasma;

processed into plasma within 24 to 72 hours of the blood draw;

maintained, stored, and/or shipped at room temperature prior to processing into plasma; and/or

maintained, stored, and/or shipped without exposure to chilling or freezing prior to processing into plasma.

Claim 10. The method of any one of claims 1 to 7, wherein the plurality of C-RNA molecules further comprises corticotropin releasing hormone (CRH), fibronectin 1 (FN1), pappalysin 2 (PAPPA2), TIMP metallopeptidase inhibitor 3 (TIMP3), leptin (LEP), H19 imprinted maternally expressed transcript (H19), proteoglycan 2, pro eosinophil major basic protein (PRG2), nestin (NES), V-set and immunoglobulin domain containing 4 (VSIG4), and/or Epstein-Barr virus induced 3 (EB13).

Claim 11. The method of claim 10, wherein the identification of a plurality of C-RNA molecules consisting of:

ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, leptin (LEP), pappalysin 2 (PAPPA2), and V-set and immunoglobulin domain containing 4 (VSIG4);

ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, PAPPA2, SKIL, and VSIG4;

ADAMTS2, ARHGEF25, arrestin domain containing 4 (ARRDC4), CLEC4C, LEP, nestin (NES), SKIL, and VSIG4;

ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, corticotropin releasing hormone (CRH), LEP, PAPPA2, SKIL, and VSIG4;

ADAMTS2, ARHGEF25, ARRDC2, CLEC4C, LEP, and SKIL; or

ADAMTS2, ARHGEF25, ARRDC2, ARRDC4, CLEC4C, LEP, PAPPA2, and SKIL

is indicative of preeclampsia in the pregnant women.