

COSMETIC OR PHARMACEUTICAL COMPOSITIONS COMPRISING METALLOPROTEINASE INHIBITORS

5 **FIELD OF THE INVENTION**

The present invention refers to peptides capable of inhibiting the activity of matrix metalloproteinases (MMP) and to cosmetic or pharmaceutical compositions containing such peptides which can be used in the treatment and/or care of skin, mucosae and/or scalp, preferably for the treatment and/or care of those conditions, disorders and/or
10 pathologies of skin, mucosae and/or scalp resulting from MMP overexpression or an increase in the MMP activity.

BACKGROUND OF THE INVENTION

The skin is made up of two layers: epidermis and dermis. The outer layer is the
15 epidermis which is made up mainly of keratinocytes, melanocytes and Langerhans cells and its basic function is to retain body water, act as a barrier against harmful chemical agents as well as against pathogen agents, and perform cellular renovation processes. The inner layer, dermis, formed by fibroblasts, adipocytes and macrophages is tightly connected to the epidermis through the basal membrane and it
20 contains numerous nerve endings which provide tactile and temperature sensations. It also houses hair follicles, sweat glands, sebaceous glands, Apocrine Glands and blood vessels, and one of its main functions is to keep the skin elasticity and appearance.

The dermis also includes the extracellular matrix, formed by a group of extracellular
25 proteins (fibrous proteins, glycoproteins and proteoglycans) whose principal function is to keep skin structure. Correct tissue functioning and development depend on the right formation of the extracellular matrix and on the right regulation of its components [Wiberg C., Klatt A.R., Wagener R., Paulsson M., Bateman J.F., Heinegard D. and Morgelin M. (2003) "Complexes of matrilin-1 and biglycan or decorin connect collagen
30 VI microfibrils to both collagen II and aggrecan" *J. Biol. Chem.* 278:37698-37704]. The two most important fibrous proteins in the extracellular matrix are collagen and elastin, which are responsible for the mechanical properties of the tissues such as the ability to resist tension, compression, extensibility and torsion. Proteoglycans have a structural and metabolic function, while glycoproteins, together with proteoglycans, work as a
35 union bridge between matrix components and cells [Aumailley M. and Gayraud B.

- 2 -

(1998) "Structure and biological activity of the extracellular matrix" *J. Mol. Med.* 76:253-265; Culav E.M., Clark C.H. and Merrilees M.J. (1999) "Connective tissues: matrix composition and its relevance to physical therapy" *Phys. Ther.* 79:308-319; Scott J.E. (2003) "Elasticity in extracellular matrix 'shape modules' of tendon, cartilage, etc. A sliding proteoglycan-filament model" *J. Physiol.* 553:335-343].

Collagens are a family of fibrous proteins of the extracellular matrix that constitute a 25% of the total proteic mass in mammals. They have been classified in more than 20 families, all of them having individual characteristics which fulfill specific functions in different tissues.

The main characteristic of collagen is its helicoidal structure formed by the association of three polypeptide chains rich in glycine and proline. Alterations in its aminoacid composition cause dysfunction and loss of its mechanical properties [Culav E.M., Clark C.H. and Merrilees M.J. (1999) "Connective tissues: matrix composition and its relevance to physical therapy" *Phys. Ther.* 79:308-319]. These polypeptide chains can associate one to the other and form fibrils, which have a diameter of 10-300nm and a length of up to hundreds of micrometers in mature tissues. These fibrils are often added into major structures, such as cable bunching, which can be seen through electronic microscopy as collagen fibers of many micrometers in diameter. This process is known as fibrillogenesis [Aumailley M. and Gayraud B. (1998) "Structure and biological activity of the extracellular matrix" *J. Mol. Med.* 76:253-265]. Not all collagens have the ability to form fibrils; only I, II, III, V and XI type collagens, which are known as fibrillar collagens.

An adult dermis is basically formed by fibrillar collagens type I, III and V. Type I collagens represent 80-90% of the total collagen of the dermis. Generally, type I collagen fibers feature a bigger diameter, which correlates with its ability to withstand a bigger mechanical load. Type III collagen intervenes in tissue extensibility, and as years go by, it is replaced by type I collagen molecules, process which is partly responsible for mature skins being less extensible than young skins. Type V collagen associates with types I and III regulating the diameter of fibrils [*The Biology of the Skin*, Freinkel R.K. and Woodley D.T., eds. The Parthenon Publishing Group, 2001; Culav E.M., Clark C.H. and Merrilees M.J. (1999) "Connective tissues: matrix composition and its relevance to physical therapy" *Phys. Ther.* 79:308-319].

- 3 -

Collagen fibers are in constant renewal process, but such renewal decreases with age, causing the thinning of dermis. Besides, even though collagen fibers organization provides collagen network with great resistance, collagen fibers are sensitive to certain enzymes known as matrix metalloproteases (MMP). MMPs belong to a family of proteolytic enzymes (endoproteases) which contain a zinc atom coordinated with three cysteine residues and one residue of methionine in its active center and which can, collectively, degrade macromolecular components from the extracellular matrix and from the basal laminas into a neutral pH (collagen, elastin, etc.).

The family of matrix metalloproteases is classified according to its structural similarity and its substrate specificity [Woessner J.F. (1991) "Matrix metalloproteinases and their inhibitors in connective tissue remodeling" *Faseb J.* 5:2145-2154; Miyazaki K. and Higashi S. (1996) "Matrix metalloproteinases: their structures and functions, with special reference to their roles in tumor invasion and metastasis" *Seikagaku* 68:1791-1807]. Within the family of MMPs there are collagenases which degrade fibrillar collagen (MMP-1 or interstitial collagenase, MMP-8 or neutrophil collagenase, MMP-3 or collagenase 3), gellatinases which degrade type IV collagen or any other form of denaturalized collagen (MMP-2 or gellatinase A 72kDa and MMP-9 or gellatinase B 92kDa), stromelysins whose wide spectrum of activity is directed to the extracellular matrix proteins such as glycoproteins like fibronectin or laminin and proteoglycans, among others (MMP-3 or stromelysin 1, MMP-10 or stromelysin 2 and MMP-11 or stromelysin 3), matrilysin (MMP-7) metalloelastase (MMP-12) or the membrane metalloproteases (MMP-14, MMP-15, MMP-16 and MMP-17).

Metalloproteases are produced and secreted in an inactive way (proenzyme), which is later activated in the extracellular environment by the loss of the propeptide region of its sequence. The members of this protein family can activate one another. The MMP activity regulation can take place in different ways: regulating gens expression (transcription and transfer), regulating inactive process activation or acting locally on the active process.

MMPs play an important role in different skin, mucosae and/or scalp conditions and disorders in which there is a degradation and destruction of extracellular proteins [Kahari V.M. and Saarialho-Kere U. (1997) "Matrix metalloproteinases in skin" *Exp.*

- 4 -

Dermatol. 6:199-213]. Among the different pathologies described in which there is a MMP overexpression or an increase of MMP activity in connective tissue cells, we find chronic ulcer [Miyoshi H., Kanekura T., Aoki T. and Kanzaki T. (2005) "Beneficial effects of tissue inhibitor of metalloproteinases-2 (TIMP-2) on chronic dermatitis" *J. Dermatol.* 32:346-353], psoriasis [Flisiak I., Mysliwiec H. and Chodyncka B. (2005) "Effect of psoriasis treatment on plasma concentrations of metalloproteinase-1 and tissue inhibitor of metalloproteinase-1" *J. Eur. Acad. Dermatol. Venereol.* 9:418-421; Suomela S., Kariniemi A.L., Impola U., Karvonen S.L., Snellman E., Uurasmaa T., Peltonen J., Saarialho-Kere U. (2003) "Matrix metalloproteinase-19 is expressed by keratinocytes in psoriasis" *Acta Derm. Venereol.* 83:108-114], oral pathologies such as gingivitis and periodontitis [Reynolds J.J. and Meikle M.C. (1997) "The functional balance of metalloproteinases and inhibitors in tissue degradation: relevance to oral pathologies" *J. R. Coll. Surg. Edinb.* 42:154-160], skin cancer [Ntayi C., Hornebeck W. and Bernard P. (2004) "Involvement of matrix metalloproteinases (MMPs) in cutaneous melanoma progression" *Pathol. Biol. (Paris)* 52:154-159; Kerkela E. and Saarialho-Kere U. (2003) "Matrix metalloproteinases in tumor progression: focus on basal and squamous cell skin cancer" *Exp. Dermatol.* 12:109-125] and tumor invasion and metastasis [Sato H., Takino T. and Miyamori H. (2005) "Roles of membrane-type matrix metalloproteinase-1 in tumor invasion and metastasis" *Cancer Sci.* 96:212-217].

MMPs also play a key role in different physiological situations in which the extracellular matrix is degraded or reconstructed, such as the extracellular matrix proteolytic remodeling, including tissue morphogenesis during development, tissue repair and angiogenesis [Kahari V.M. and Saarialho-Kere U. (1997) "Matrix metalloproteinases in skin" *Exp. Dermatol.* 6:199-213]. In a particular way, MMPs have a crucial role in connective tissue remodeling [Abraham D., Ponticos M. and Nagase H. (2005) "Connective tissue remodeling: cross-talk between endothelins and matrix metalloproteinases" *Curr. Vasc. Pharmacol.* 3:369-379], for example collagen degradation by MMPs makes the skin look wrinkled and flaccid.

Likewise, MMPs participate in skin aging. Different factors, including exposure to ultraviolet (UV) radiation, produce collagen degradation, with all the consequences it entails on skin structure and/or firmness, particularly on those skin areas exposed to the solar light such as the face, ears, neck, scalp, arms and hands.

- 5 -

Skin damage associated to chronic exposition (repetitive irradiation) or high exposition (strong irradiation) to UVA and/or UVB rays has been studied; particularly it is known that

- UVB rays (290-300nm; 5% of total UV rays) with more energetic wavelength, especially affect epidermic cells (keratinocytes) acting over its DNA.
- UVA rays (320-400nm; 95% of total UV rays) have a stronger penetration grade and also act over dermic cells such as fibroblasts and they act indirectly generating free radicals.

Moreover, prolonged exposure to UV radiation, particularly to UVA and/or UVB radiation stimulates MMP expression [Fisher G.J., Datta S.C., Talwar H.S., Wang Z.Q., Varani J., Kang S. and Voorhees J.J. (1996) "Molecular basis of sun-induced premature skin ageing and retinoid antagonism" *Nature* 379:335-339; Fisher G.J., Wang Z.Q., Datta S.C., Varani J., Kang S. and Voorhees J.J. (1997) "Pathophysiology of Prematur Skin Aging Induced by Ultraviolet Light" *New Eng. J. Med.* 337:1419-1429; Fisher G.J., Choi H.C., Bata-Csorgo Z., Shao Y., Datta S., Wang Z.Q., Kang S. and Voorhees J.J. (2001) "Ultraviolet irradiation increases matrix metalloproteinase-8 protein in human skin in vivo" *J. Invest. Dermatol.* 117:219-226], especially matrix metalloelastase type 1 (MMP-1). This is one of the components of photoinduced skin aging (or photoaging) [Rittie L. and Fisher G.J. (2002) "UV-light-induced signal cascades and skin aging" *Ageing Res. Rev.* 1:705-720]. Besides, it is known that MMP-1, MMP-2 and MMP-9 activity increases with age and that this increase, together with cell growth deceleration, contributes to chronologic skin aging [EP 1 005 333 B1]. Similarly, smokers' skin also has a premature aging aspect in which MMPs are overexpressed [Lahmann C., Bergemann J., Harrison G. and Young A.R. (2001) "Matrix metalloproteinase-1 and skin aging in smokers" *Lancet* 357:935-936].

Another skin and/or scalp pathologies or disorders, associated to MMP overexpression or to an increase of MMP activity in the connective tissue is acne [Papakonstantinou E., Aletras A.J., Glass E., Tsogas P., Dionyssopoulos A., Adjaye J., Fimmel S., Gouvousis P., Herwig R., Lehrach H., Zouboulis C.C. and Karakiulakis G. (2005) "Matrix metalloproteinases of epithelial origin in facial sebum of patients with acne and their regulation by isotretinoin" *J. Invest. Dermatol.* 125:673-684]. It is described that skins affected by acne have high levels of MMP-1.

Likewise, rosacea is a skin and/or scalp pathology or disorder in which MMPs are also

- 6 -

involved. Rosacea is characterized by an increase of angiogenesis and inflammation. Angiogenesis refers to the process of new blood vessels formation and it includes benign conditions such as rosacea and malignant processes such as cancer. Matrix degrading enzymes, present in tissue extracellular matrix facilitate angiogenesis since they allow new blood vessels to penetrate the matrix. MMPs represent a kind of enzymes involved in such processes [Sapadin A.N., Fleischmajer R. (2006) "Tetracyclines: Nonantibiotic properties and their clinical implications" *J. Am. Acad. Derm.* 54:258-265].

People with dermatitis, including contact dermatitis and atopic dermatitis, also have high levels of some MMPs [Herouy Y., Mellios P., Bandemir E., Dichmann S., Nockowski P., Schöpf E. and Norgauer J. (2001) "Inflammation in stasis dermatitis upregulates MMP-1, MMP-2 and MMP-13 expression" *J. Dermatol. Sci.* 25:198-205; Devillers A.C., van Toorenenbergen A.W., Klein Heerenbrink G.J., Muldert P.G. and Oranje A.P. (2007) "Elevated levels of plasma matrix metalloproteinase-9 in patients with atopic dermatitis: a pilot study" *Clin. Exp. Dermatol.* 32:311-313; Miyoshi H., Kanekura T., Aoki T. and Kanzaki T. (2005) "Beneficial effects of tissue inhibitor of metalloproteinases-2 (TIMP-2) on chronic dermatitis" *J. Dermatol.* 32:346-353]. "Dermatitis" is defined as those skin conditions, disorders or pathologies that cause inflammation, including contact dermatitis, atopic dermatitis, sensitive skin and eczema. It is also known that MMPs are involved in perifollicular matrix degradation, and thus, in hair loss. Specifically, cytokines and the epidermal growth factor stimulate MMP-9 production in the lower epithelial compartment of hair root, such mechanism controls capillary follicle involution observed in alopecia [Jarrousse F., Boisnic S., Branchet M.C., Beranger J.Y., Godeau G., Breton L., Bernard B.A. and Mahé Y.F. (2001) "Identification of clustered cells in human hair follicle responsible for MMP-9 gelatinolytic activity: consequences for the regulation of hair growth" *Int. J. Dermatol.* 40:385-392]. Thus, overexpressed MMP inhibition during alopecic processes could be effective in delaying, and even preventing, hair loss [EP 1 076 549 B1].

Also, MMP activity is related to scar formation in tissues containing collagen. "Scar formation is defined as the formation of an abnormal morphological collagen structure due to previous injuries or due to the healing process of tissue containing collagen on the skin.

- 7 -

Healing processes consist of three stages: (1) inflammation, (2) tissue formation and (3) tissue remodeling. A necessary stage in the healing process is extracellular matrix degradation: in order for the cells to proliferate in the wounded area and regenerate it, it is necessary that the extracellular matrix be degraded. Such degradation is made
5 through MMPs. Healing process stages are regulated by a balance between the different MMPs and it has been described that an excess of MMP activity causes chronic ulcers. For example, an overexpression of MMP-8 can be associated to the pathogenesis of leg chronic ulcers. Likewise, diabetic ulcers are characterized by a prolonged inflammation, decrease collagen synthesis and high MMP levels.

10

Most scars consist of collagen fibers irregularly organized as well as an excess of collagen. Scars have different causes (accidents, surgery, skin diseases, burns, acne, infections and accidents in general), but not all scars are the same. Different kinds of scars can be grouped in

15

- Flat and pale scars: formed as a result of the body's natural healing process.

- Sunken scars: formed by skin attached to deeper structures, such as muscles, or due to loss of fat in internal tissues. These scars are recessed into the skin and are usually the result of an injury.

20

- Hypertrophic scars: appear when the body produces an excess of collagen during the healing process. These scars elevate over the skin surface and contain irregularly organized collagen.

25

- Keloid scars: formed as a result of an imbalance in the production of collagen during the healing process. These scars not only elevate over skin surface, but also they extend beyond the boundary of the original wound and can continue to grow indefinitely.

30

- Acne scars: formed in skin affected by acne. The scar can be sunken or become a keloid. People who have had chicken pox can have similar scars.

35

- Stretched scars: occur when the skin around a healing wound is put under tension during the healing process. Initially, the scar may appear normal but can widen and thin over a period of weeks or months. This can occur when the

- 8 -

wound is close to a joint and is stretched during movement or it may be due to poor healing because of general ill health or malnutrition.

- Stretch marks: develop when the skin is stretched rapidly, for example during pregnancy or the adolescent growth spurt.

Therefore, skin scar reduction is desirable both from the pathological point of view, as healing during fibrotic processes, and from the cosmetic point of view, as in the case of softening the aspect of scars caused by acne or stretch marks.

It has also been described that during adipocytes proliferation and differentiation, MMPs are overexpressed [Traurig M.T., Permana P.A., Nair S., Kobes S., Bogardus C. and Baier L.J. (2006) "Differential expression of matrix metalloproteinase 3 (MMP3) in preadipocytes/stromal vascular cells from nonobese nondiabetic versus obese nondiabetic Pima Indians" *Diabetes* 55:3160-3165]. MMP activity inhibition with several specific inhibitors prevents adipocytes differentiation. An especially interesting fact is that MMP inhibitors are able to reduce the accumulation of lipogenic markers (triglycerides) in adipocyte cultures [Demeulemeester D., Collen D. and Lijnen H.R. (2005) "Effect of matrix metalloproteinase inhibition on adipose tissue development" *Biochem. Biophys. Res. Commun.* 329:105-110]. Thus, MMP inhibitors can be developed as anti cellulite agents and help reduce orange peel skin aspect.

MMP activity is also responsible for the extracellular matrix disorganization that surrounds lymphatic and blood vessels. Matrix deterioration around blood vessels allows for a passive vasodilatation which gives place to capillary visibility or telangiectasia, or couperosis. Besides, this microcapillary passive dilatation can cause local blood vessel bursts which can give place to bags under the eyes or dark circles in the periorbital area. Furthermore, MMPs have an influence over vein wall mechanical properties, which can make veins fragile and consequently lead to the development of varicose veins.

Apart from the relation of MMPs to tissue matrix degradation, it has been suggested that MMPs are also involved in different pathologies that concur with an abnormal metabolism of the connective tissue or basal membrane matrix such as arthritis (rheumatoid arthritis, osteoarthritis, etc.), bone diseases (osteoporosis, etc.), ectopic

- 9 -

angiogenesis, multiple sclerosis, tumors metastasis and tissue ulcers (cornea, stomach, epidermis, etc.) [EP 0 927 161 B1]. Therefore, an MMP inhibitor could be effective in treating and preventing those pathologies caused by an abnormal metabolism of the tissular matrix.

5

Then, it is widely accepted that MMP activity regulation is highly important for the basal membrane and extracellular matrix protection, as well as for preventing and improving signs of aging. In the context of the present invention, the term "aging" refers to changes experienced by the skin with the passing of years (chrono-aging), or due to sun exposition (photoaging) or due to environmental agents like tobacco smoke, extreme cold or wind weather conditions, chemical pollutants or pollution and it includes all visible external changes as well as those perceptible by touch, such as for example and in a non-limiting sense, development of skin discontinuities such as wrinkles, thin lines, cracks, irregularities or roughness, increase of pore size, loss of elasticity, loss of firmness, loss of smoothness, loss of the capacity to recuperate after deformation, skin hanging such as cheek hanging, appearance of eye pouches or double chin, among others, changes of the skin color, such as marks, reddening, bags under the eyes or the appearance of hyperpigmented areas such as age marks or freckles among others, anomalous differentiation, hyperkeratinization, elastosis, keratosis, orange-peel skin, loss of collagen structuring and other histological changes of the stratum corneum, of the dermis, epidermis, vascular system (for example the appearance of spider veins or telangiectasias) or of those tissues close to the skin, among others.

25 The cosmetic and pharmaceutical sectors have identified numerous compounds and plant extracts which are effective as MMP inhibitors and there are different bibliographical revisions in literature about MMPs, pathologies associated to their overexpression or their activity increase and the different families of compounds and plant extracts useful to their inhibition. In the state of the art there have been described different approximations to control MPPs activity, including small molecules [Levy D.E., Lapierre E., Liang W., Ye W., Lange C.W., Li X., Grobelny D., Casabonne M., Tyrrell D., Holme K., Nadzan A. and Galaray R.E. (1998) "Matrix metalloproteinase inhibitors: A structure activity study" *J. Med. Chem.* 41:199-223; Wojtowicz-Praga S.M., Dickson R.B. and Hawkins M.J. (1997) "Matrix metalloproteinase inhibitors" *Investigational new*
30 *Drugs* 15:61-75; Duivenvuurden W.C.M., Hirte H.W. and Singh G. (1997) "Use of

- 10 -

tetracycline as an inhibitor of matrix metalloproteinase activity secreted by human bone metastasizing cancer cells" *Invasion and Metas.* 17:312-322] peptidic inhibitors [Otake S., Monta Y. and Morikawa T. (1994) "Inhibition of matrix metalloproteinases by peptidyl hydroxamic acids" *Biochem. Biophys. Res. Comm.* 199:1442-1446] or antibodies against MMPs [Su J-L., Becherer D., Edwards C., Bukhart W., McMgeehan G.M. and Champion B.R. (1995) "Monoclonal antibodies against human collagenase and stromelysin" *Hybridoma* 14:383-390]. Cosmetic industry has made important efforts to offset MMPs activity and the age-related loss of functionality of extracellular matrix components caused by MMPs. Balance between production and degradation of skin essential biomolecules such as collagen evolves with aging towards degradation processes, which leads to, for example, a progressive thinning and disorganization of the dermis which produces dermis flaccidity and a subsequent formation of wrinkles. Therefore, those methods which allow to delay or prevent extracellular matrix degradation will have a potential beneficial effect on mature skins or on aged and/or photo-aged skins; allowing them to partially recover the mechanical properties (elasticity, flexibility and firmness) which they have lost due to age or sun exposure and/or environmental pollutants and thus show a better appearance with fewer wrinkles and a smoother skin. Likewise, MMP inhibition is also an important aspect for the cosmetic sector for applications other than delaying the aging and/or photo-aging, such as for example hair growth modulation [EP 1 076 549 B1] or wound treatments [US 2004/0127420 A1; US 2003/0166567 A1].

Despite the great number of existing compounds and/or extracts, there is still a need to identify new more effective and selective MMP inhibitors.

In the present invention there are described peptides which are effective in MMP inhibition, imitating on this way the function of endogenous MMP inhibitors (TIMP, matrix metalloproteinase tissue inhibitor). The peptide sequence of the invention is not contained in proenzymatic MMP sequences, such as the peptide sequences described in US 2004/0127420 A1 and US 2003/0166567 A1. Sequences similar to the peptides of the invention, without the citrulline residue on the carboxy-terminal (C-terminal), are found in sequences of different enzymes or have enzymatic activity [WO 2004/033668 A2; WO 99/00489 A1]; there is not any clue in the state of the art that suggests the effectiveness of the peptides of the invention as MMP inhibitors, so a person skilled in the art could not deduce the nature of the peptides which inhibit


MMPs.

DESCRIPTION OF THE INVENTION

The present invention provides a solution to the above mentioned problem. Surprisingly, the applicant of the present invention has found that certain peptides, whose amino acid sequence does not derive from natural products, are capable of inhibiting MMPs, mainly human MMP-1, MMP-2, MMP-3 and/or MMP-9.

Therefore, peptides in the present invention provide a simple and effective and risk-free solution for the treatment and/or care of skin, mucosae and/or scalp which comprises the application on the skin, mucosae and/or scalp or the oral or parenteral administration of a peptide of general formula (I) to a mammal, as it is described below.

In a first aspect, the invention refers to a peptide according to the general formula (I)



(I)

stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof, characterized in that:

AA₁ is -Arg-;

AA₂ is selected from the group consisting of -His- and -Asn-;

AA₃ AA₂ is selected from the group consisting of -His- and -Arg-;

AA₄ is -Cit-;

R₁ is selected from the group consisting of H, substituted or non-substituted non-cyclic aliphatic group, substituted or non-substituted alicyclyl, substituted or non-substituted heterocyclyl, substituted or non-substituted heteroarylalkyl, substituted or non-substituted aryl, substituted or non-substituted aralkyl, and R₅-CO-; and

R₂ is selected from the group consisting of -NR₃R₄, -OR₃ and -SR₃; wherein R₃ and R₄ are independently selected from the group consisting of H, substituted or non-substituted non-cyclic aliphatic group, substituted or non-substituted alicyclyl, substituted or non-substituted heterocyclyl, substituted or non-substituted heteroarylalkyl, substituted or non-substituted aryl and substituted or non-substituted aralkyl;

wherein R₅ is selected from the group consisting of H, substituted or non-

- 12 -

substituted non-cyclic aliphatic group, substituted or non-substituted alicyclyl, substituted or non-substituted aryl, substituted or non-substituted aralkyl, substituted or non-substituted heterocyclyl and substituted or non-substituted heteroarylalkyl.

5

Another aspect of this invention is a process to obtain these peptides of general formula (I).

10

Another aspect of this invention is aimed at a cosmetic or pharmaceutical composition comprising a cosmetic or pharmaceutical effective amount of at least one peptide of general formula (I), stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof, and at least one cosmetically or pharmaceutically acceptable excipient or adjuvant.

15

In another aspect, the invention is aimed at the use of a peptide of general formula (I), stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof, in the preparation of a cosmetic or pharmaceutical composition for the treatment and/or care of skin, mucosae and/or scalp.

20

DETAILED DESCRIPTION OF THE INVENTION

The peptides of the invention are peptides not derived from natural products; which have an important MMP inhibiting activity and thus, they are useful for the treatment and/or care of those conditions, disorders and/or pathologies of the skin, mucosae and/or scalp resulting from an MMP overexpression or from an increased MMP activity.

25

Definitions

For a better understanding of the present invention, there are hereby included the meanings of some terms and expressions, as used in the context of the invention.

30

In the present description, the abbreviations used for amino acids follow the rules of IUPAC-IUB Commission on Biochemical Nomenclature specified in *Eur. J. Biochem.* (1984) 138:9-37 and in *J. Biol. Chem.* (1989) 264:633-673.

35

So, for example Gly represents $\text{NH}_2\text{-CH}_2\text{-COOH}$, Gly- represents $\text{NH}_2\text{-CH}_2\text{-CO-}$, -Gly represents $\text{-NH-CH}_2\text{-COOH}$ and -Gly- represents $\text{-NH-CH}_2\text{-CO-}$. Therefore, the dash

- 13 -

representing the peptide bond, eliminates the OH from the 1-carboxyl group of the amino acid (herein represented in the conventional non-ionized form) when it is placed to the right of the symbol, and it eliminates the H from the 2-amino group of the amino acid when it is placed to the left of the symbol; both modifications can be applied to the

5





Symbol	Residue	Symbol	Residue
-Arg-		-Cit-	
-His-		-Asn-	

Table 1

10

In this description, the abbreviation "Ac-" is used to designate the acetyl group (CH₃-CO-) and the abbreviation "Palm-" is used to designate palmitoyl group (CH₃-(CH₂)₁₄-CO-).

15 The term "non-cyclic aliphatic group" is used in the present invention to cover, for example and in a non-limiting sense, linear or branch alkyl, alkenyl and alkynyl groups.

The term "alkyl group" refers to a linear or branched saturated group, which has between 1 and 24, preferably between 1 and 16, more preferably between 1 and 14, even more preferably between 1 and 12, even more preferably 1, 2, 3, 4, 5 or 6 carbon, atoms and which is bound to the rest of the molecule by a single bond, including, for example and in a non-limiting sense, methyl, ethyl, isopropyl, isobutyl, *tert*-butyl, heptyl, octyl, decyl, dodecyl, lauryl, hexadecyl, octadecyl, amyl, 2-ethylhexyl, 2-methylbutyl, 5-methylhexyl and the like.

20

The term "alkenyl group" refers to a group having between 2 and 24, preferably between 2 and 16, more preferably between 2 and 14, even more preferably between 2 and 12, even more preferably 2, 3, 4, 5 or 6 carbon atoms, with one or more carbon-carbon double bonds, preferably with 1, 2 or 3 conjugated or non-conjugated carbon-carbon double bonds, and which is bound to the rest of the molecule by a single bond, including, for example and in a non-limiting sense, vinyl, oleyl, linoleyl group and the like.

10 The term "alkynyl group" refers to a group having between 2 and 24, preferably between 2 and 16, more preferably between 2 and 14, even more preferably between 2 and 12, even more preferably 2, 3, 4, 5 or 6 carbon atoms, with one or more conjugated or non-conjugated carbon-carbon triple bonds, preferably with 1, 2 or 3 conjugated or non-conjugated carbon-carbon triple bonds, and which is bound to the
15 rest of the molecule by a single bond, including, for example and in a non-limiting sense, ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, pentynyl group, such as for example 1-pentylnyl and the like.

The term "alicyclic group" is used in the present invention to cover, for example and in
20 a non-limiting sense, cycloalkyl, cycloalkenyl or cycloalkylnyl groups.

The term "cycloalkyl" refers to a saturated mono- or polycyclic aliphatic group which has between 3 and 24, preferably between 3 and 16, more preferably between 3 and 14, even more preferably between 3 and 12, even more preferably 3, 4, 5 or 6 carbon
25 atoms, and which is bound to the rest of the molecule by a single bond, including, for example and in a non-limiting sense, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, methyl cyclohexyl, dimethyl cyclohexyl, octahydroindene, decahydronaphthalene, dodecahydrophenalene and the like.

30 The term "cycloalkenyl group" refers to a non-aromatic mono- or polycyclic aliphatic group which has between 5 and 24, preferably between 5 and 16, more preferably between 5 and 14, even more preferably between 5 and 12, even more preferably 5 or 6 carbon atoms, with one or more carbon-carbon double bonds, preferably with 1, 2 or 3 conjugated or non-conjugated carbon-carbon double bonds, and which is bound to
35 the rest of the molecule by a single bond, including, for example and in a non-limiting

sense, cyclopent-1-en-1-yl group and the like.

The term "cycloalkynyl group" refers to a non-aromatic mono- or polycyclic aliphatic group which has between 5 and 24, preferably between 5 and 16, more preferably
5 between 5 and 14, even more preferably between 5 and 12, even more preferably 5 or 6 carbon atoms, with one or more carbon-carbon triple bonds, preferably 1, 2 or 3 conjugated or non-conjugated triple bonds carbon-carbon, and which is bound to the rest of the molecule by a single bond, including, for example and in a non-limiting sense, cyclohex-1-yn-1-yl group and the like.

10

The term "aryl group" refers to an aromatic group which has between 6 and 30, preferably between 6 and 18, more preferably between 6 and 10, even more preferably between 6 or 10 carbon atoms, comprising 1, 2, 3 or 4 aromatic nuclei, bound by a carbon-carbon bond, or fused, including, for example and in a non-limiting sense,
15 phenyl, naphthyl, diphenyl, indenyl, phenanthryl or anthranil among others; or an aralkyl group.

15

The term "aralkyl group" refers to an alkyl group substituted with an aromatic group, having between 7 and 24 carbon atoms and including, for example and in a non-
20 limiting sense, $-(CH_2)_{1-6}$ -phenyl, $-(CH_2)_{1-6}$ -(1-naphthyl), $-(CH_2)_{1-6}$ -(2-naphthyl), $-(CH_2)_{1-6}$ -CH(phenyl)₂ and the like.

20

The term "heterocyclyl group" refers to a 3-10 member hydrocarbon ring, in which one or more of the atoms of the ring, preferably 1, 2 or 3 atoms of the ring, are elements
25 other than carbon, such as nitrogen, oxygen or sulfur and which can be saturated or unsaturated. For the purposes of this invention, the heterocycle can be a monocyclic, bicyclic or tricyclic cyclic system, which can include systems of fused rings; and the nitrogen, oxygen or sulfur atoms can be optionally oxidized in the heterocyclic radical; the nitrogen atom can be optionally quaternized, and the heterocyclyl radical can be
30 partially or completely saturated or be aromatic. The term heterocyclyl more preferably refers to a 5 or 6 member ring.

30

The term "heteroarylalkyl group" refers to an alkyl group substituted with a substituted or non-substituted aromatic heterocyclyl group, wherein the alkyl group has 1 to 6
35 carbon atoms and the aromatic heterocyclyl group has between 2 to 24 carbon atoms

35

- 16 -

and from 1 to 3 atoms different from carbon, and including, for example and in a non-limiting sense, $-(CH_2)_{1-6}$ -imidazolyl, $-(CH_2)_{1-6}$ -triazolyl, $-(CH_2)_{1-6}$ -thienyl, $-(CH_2)_{1-6}$ -furyl, $-(CH_2)_{1-6}$ -pyrrolidinyl and the like.

As it is understood in this technical area, there can be a certain level of substitution on the above defined radicals. Thus, there can be substitution in any of the groups in the present invention. This document references to substituted groups in the groups of the present invention indicate that the specified radical can be substituted in one or more available positions by one or more substituents, preferably in 1, 2 or 3 positions, more preferably in 1 or 2 positions, and even more preferably in 1 position. Said substituents include, for example and in a non-limiting sense, C_1 - C_4 alkyl; hydroxyl; C_1 - C_4 alkoxy; amino; C_1 - C_4 aminoalkyl; C_1 - C_4 carbonyloxy; C_1 - C_4 oxycarbonyl; halogen such as fluorine, chlorine, bromine and iodine; cyano; nitro; azido; C_1 - C_4 alkylsulfonyl; thiol; C_1 - C_4 alkylthio; aryloxy such as phenoxy; $-NR_b(C=NR_b)NR_bR_c$; wherein R_b and R_c are independently selected from the group consisting of H, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_3 - C_{10} cycloalkyl, C_6 - C_{18} aryl, C_7 - C_{17} aralkyl, 3-10 member heterocyclyl or protective group of the amino group.

Compounds of the invention

The compounds of the invention are defined by the general formula (I)



wherein R_1 , AA_1 , AA_2 , AA_3 , AA_4 and R_2 have the previously defined meaning.

The R_1 and R_2 groups are bound to amino-terminal (*N*-terminal) and carboxy-terminal (*C*-terminal) ends of the peptide sequence.

According to an embodiment of the present invention, R_1 is selected from the group consisting of H or R_5 -CO-, wherein R_5 is selected from the group consisting of the substituted or non-substituted C_1 - C_{24} alkyl radical, substituted or non-substituted C_2 - C_{24} alkenyl, substituted or non-substituted C_2 - C_{24} alkynyl, substituted or non-substituted C_3 - C_{24} cycloalkyl, substituted or non-substituted C_5 - C_{24} cycloalkenyl, substituted or non-substituted C_5 - C_{24} cycloalkynyl, substituted or non-substituted C_6 - C_{30} aryl, substituted or non-substituted C_7 - C_{24} aralkyl, substituted or non-substituted 3-10 member heterocyclyl, and substituted or non-substituted heteroarylalkyl with from 2 to 24 carbon atoms and from 1 to 3 atoms different from carbon and an alkyl chain of 1 to

- 17 -

6 carbon atoms. More preferably, R_1 is selected from H, acetyl, *tert*-butanoyl, hexanoyl, 2-methylhexanoyl, cyclohexanecarboxyl, octanoyl, decanoyl, lauroyl, myristoyl, palmitoyl, stearoyl, oleoyl and linoleoyl. Even more preferably, R_1 is H, acetyl, lauroyl, myristoyl or palmitoyl. In an even more preferred embodiment, radicals R_1 are H, acetyl or palmitoyl.

According to another preferred embodiment, R_2 is $-NR_3R_4$, $-OR_3$ or $-SR_3$ wherein R_3 and R_4 are independently selected from the group consisting of H, substituted or non-substituted C_1 - C_{24} -alkyl, substituted or non-substituted C_2 - C_{24} alkenyl, substituted or non-substituted C_2 - C_{24} alkynyl, substituted or non-substituted C_3 - C_{24} cycloalkyl, substituted or non-substituted C_5 - C_{24} cycloalkenyl, substituted or non-substituted C_5 - C_{24} cycloalkynyl, substituted or non-substituted C_6 - C_{30} aryl, substituted or non-substituted C_7 - C_{24} aralkyl, substituted or non-substituted 3-10 member heterocyclyl, and substituted or non-substituted heteroarylalkyl with from 2 to 24 carbon atoms and from 1 to 3 atoms different from carbon and an alkyl chain of 1 to 6 carbon atoms. Optionally, R_3 and R_4 can be bound by a saturated or unsaturated carbon-carbon bond, forming a cycle with the nitrogen atom. More preferably R_2 is $-NR_3R_4$ or $-OR_3$, wherein R_3 and R_4 are independently selected from the group consisting of H, substituted or non-substituted C_1 - C_{24} alkyl, substituted or non-substituted C_2 - C_{24} alkenyl, substituted or non-substituted C_2 - C_{24} alkynyl, substituted or non-substituted C_3 - C_{10} cycloalkyl, substituted or non-substituted C_6 - C_{15} aryl and substituted or non-substituted 3-10 member heterocyclyl, and substituted or non-substituted heteroarylalkyl with a 3 to 10 members ring and an alkyl chain of 1 to 6 carbon atoms. More preferably, R_3 and R_4 are selected from the group consisting of H, methyl, ethyl, hexyl, dodecyl or hexadecyl. More preferably, R_3 is H and R_4 is selected from the group consisting of H, methyl, ethyl, hexyl, dodecyl or hexadecyl.

According to an even more preferred embodiment, R_2 is selected from $-OH$ and $-NH_2$.

Even more preferably, R_1 is acetyl and R_2 is $-OH$.

According to an embodiment of the present invention AA_1 is $-Arg-$, AA_2 is $-His-$, AA_3 is $-His-$ and AA_4 is $-Cit-$.

According to an embodiment of the present invention, AA_1 is $-Arg-$, AA_2 is $-Asn-$, AA_3 is

- 18 -

-Arg- and AA₄ is -Cit-.

According to another embodiment of the present invention, R₁ is selected from the group consisting of H, acetyl, lauroyl, myristoyl or palmitoyl, AA₁ is -L-Arg-, AA₂ is
 5 -L-His-, AA₃ is -L-His-, AA₄ is -L-Cit- and R₂ is -NR₃R₄ or -OR₃ wherein R₃ and R₄ are independently selected from the group consisting of H, methyl, ethyl, hexyl, docetyl and hexadecyl groups, preferably R₂ is -OH or -NH₂. Even more preferably, R₁ is acetyl and R₂ is -OH.

10 According to another embodiment of the present invention R₁ is selected from the group consisting of H, acetyl, lauroyl, myristoyl or palmitoyl, AA₁ is -L-Arg-, AA₂ is -L-Asn-, AA₃ is -L-Arg-, AA₄ is -L-Cit- and R₂ is -NR₃R₄ or -OR₃ wherein R₃ and R₄ are independently selected from the group consisting of H, methyl, ethyl, hexyl, docetyl and hexadecyl groups, preferably R₂ is -OH or -NH₂. Even more preferably, R₁ is acetyl
 15 and R₂ is -OH.

In a preferred way, the compounds of formula (I) are selected from the group consisting of:

Ac-Arg-His-His-Cit-OH,
 Ac-Arg-Asn-Arg-Cit-OH,
 20 Ac-Arg-Asn-His-Cit-OH,
 Ac-Arg-His-Arg-Cit-OH,
 Palm-Arg-His-His-Cit-OH,
 Palm-Arg-Asn-Arg-Cit-OH,
 Palm-Arg-Asn-His-Cit-OH,
 25 Palm-Arg-His-Arg-Cit-OH,
 Ac-Arg-His-His-Cit-NH-(CH₂)₁₅-CH₃,
 Ac-Arg-Asn-Arg-Cit-NH-(CH₂)₁₅-CH₃,
 H-Arg-His-His-Cit-NH₂,

and

30 H-Arg-Asn-Arg-Cit-NH₂,

mixtures thereof or cosmetically or pharmaceutically acceptable salts thereof.

The peptides of the present invention can exist as stereoisomers or mixtures of stereoisomers; for example, the amino acids that form them can have a L-, D-configuration or they can be racemic independently from one another. Therefore, it is
 35 possible to obtain isomeric mixtures as well as racemic or diastereomeric mixtures or

- 19 -

pure diastereoisomers or enantiomers, depending on the number of asymmetric carbons and on which isomers or isomeric mixtures are present. Preferred structures for the peptides of the invention are pure isomers, that is to say, enantiomers or diastereoisomers.

5

For example when it is stated that AA₁ can be -Arg-, it is understood that AA₁ is selected from -L-Arg-, -D-Arg- or racemic or non-racemic mixtures of both of them. Likewise, when it is said that AA₂ can be -His-, it is understood that it can be -L-His-, -D-His- or racemic or non-racemic mixtures of both of them. The methods described in the present document allow the person skilled in the art to obtain each of the stereoisomers of the peptide of the invention by choosing the amino acid with the suitable configuration.

10

Within the scope of the present invention, there are also included cosmetically or pharmaceutically acceptable salts of peptides provided by this invention. The term "cosmetically or pharmaceutically acceptable salts" means a salt generally recognized for its use in animals, and more particularly in human beings, and it includes the salts used to form base addition salts, either inorganic, such as for example and in a non-limiting sense, lithium, sodium, potassium, calcium, magnesium or aluminium, among others, or organic such as for example and in a non-limiting sense, ethylamine, diethylamine, ethylenediamine, ethanolamine, diethanolamine, arginine, lysine, histidine, or piperazine among others; or acid addition salts, either organic, such as for example and in a non-limiting sense, acetate, citrate, lactate, malonate, maleate, tartrate, fumarate, benzoate, aspartate, glutamate, succinate, oleate, trifluoroacetate, oxalate, pamoate or gluconate among others, or inorganic, such as for example and in a non-limiting sense, chloride, sulfate, borate, or carbonate among others. The nature of the salt is not critical, as long as it is cosmetically or pharmaceutically acceptable. The cosmetically or pharmaceutically acceptable salts of the peptides the invention can be obtained by conventional methods well known in the state of the art [*Berge S.M., Bighley L.D. and Monkhouse D.C. (1977) "Pharmaceutical Salts" J. Pharm. Sci. 66:1-19*].

20

25

30

Preparation methods

Invention peptides synthesis, its stereoisomers or its cosmetically or pharmaceutically acceptable salts can be made by means of conventional methods, known in the state

35

- 20 -

of the art, such as by peptide synthesis methods in solid phase [Stewart J.M. and Young J.D. (1984) "Solid Phase Peptide Synthesis, 2nd edition" Pierce Chemical Company, Rockford, Illinois; Bodanzsky M. and Bodanzsky A. (1984) "The practice of Peptide Synthesis" Springer Verlag, New York; Lloyd-Williams P., Albericio F. and Giralt E. (1997) "Chemical Approaches to the Synthesis of Peptides and Proteins" CRC, Boca Raton, FL, USA], solution synthesis, a combination synthesis in solid phase methods and in solution or enzymatic synthesis [Kullmann W. (1980) "Proteases as catalysts for enzymic syntheses of opioid peptides" J.Biol.Chem. 255:8234-8238]. Peptides can also be obtained by bacterial strain fermentation modified or not, by genetic engineering so as to obtain the desired sequences or by animal or vegetal protein controlled hydrolysis, preferably vegetal, releasing peptide fragments containing at least the desired sequence.

For example, a method for obtaining the peptides of the invention having the (I) formula, includes the following stages:

- amino acid coupling, having the N-terminal end protected and C-terminal free, on an amino acid with its N-terminal end free and the C-terminal end protected or joined to a solid support;
- protecting group elimination from the N-terminal end;
- coupling sequence repetition and N-terminal end elimination until the desired peptide sequence is obtained;
- protecting group elimination from the C-terminal end or solid support excision.

Preferably, C-terminal end is joined to a solid support and the process takes place in solid phase, thus, it comprises the amino acid coupling with the N-terminal protected end and the C-terminal free end on an amino acid with its N-terminal free and the C-terminal joined to a polymeric support; protecting group elimination from the N-terminal end; and repetition of this sequence as many times as needed until a tetrapeptide is obtained, and finally the original polymeric support synthesized peptide is removed through excision.

Functional groups in amino acid side chains are conveniently protected by temporal or permanent protecting groups throughout the synthesis, and can simultaneously or orthogonally be unprotected to the polymeric support peptide excision.

- 21 -

Alternatively, solid phase synthesis can take place via a convergent strategy coupling a dipeptide or a tripeptide over the polymeric support or on a dipetide or amino acid previously joined to the polymeric support. Convergent synthesis strategies are widely known by subject experts and are described in Lloyd-Williams P., Albericio F. and
5 Giralt E. en "*Convergent solid-phase peptide synthesis*" (1993) *Tetrahedron* 49:11065-11133.

The process can comprise additional stages such as *N*-terminal and *C*-terminal unprotection and/or random order polymeric support peptide excision, using standard
10 processes and conditions known in the field, after which the functional groups of said ends can be modified. Optional *N*-terminal and *C*-terminal ends modification can be done with a formula (I) peptide anchored to the polymeric support or once the peptide has been removed from the polymeric support.

15 Optionally, R_1 can be introduced through the *N*-terminal end reaction of the peptide of the invention with a R_1-X wherein R_1 has the above mentioned meaning and X is a salient group, such as for example and in a non-limiting sense, tosyl group, mesyl group and halogen groups among others, by a nucleophilic substitution reaction, in presence of the right bases and solvents and wherein such fragments have those
20 functional groups which do not participate in N-C bond formation, conveniently protected by temporal or permanent protected groups.

Optionally and/or additionally, R_2 radicals can be introduced by a HR_2 compound reaction wherein R_2 is $-OR_3$, $-NR_3R_4$ or $-SR_3$, with a complementary fragment
25 corresponding to the formula (I) peptide wherein R_2 is $-OH$ in the presence of the right solvent and base such as for example, *N,N* diisopropylethylamine or triethylamine or an additive such as 1-hydroxybenzotriazole (HOBt) or hydroxybenzotriazole (HOAt) and a dehydrant agent such as, carbodiimide, an uranium salt, a phosphonium salt or an amidinium salt, among others or by previously forming an acyl halide with, for example,
30 thionyl chloride, so as to obtain a peptide of general formula (I) according to the invention, wherein such fragments present those functional groups that do not participate in N-C bond formation, conveniently protected by temporal or permanent protected groups or alternatively, other R_2 radicals can be introduced by simultaneous incorporation to the process of polymeric support peptide excision.

35

- 22 -

A subject expert will easily understand that C-terminal and N-terminal ends unprotection/excision stages and its later derivatization can take place in any order, according to processes known in the field [Smith, M. B. and March, J. (1999) "March's Advanced Organic Chemistry Reactions, Mechanisms and Structure", 5th Edition, John Wiley & Sons, 2001].

The term "protecting group" refers to a group which blocks an organic functional group and can be eliminated under controlled conditions. Protecting groups, their relative reactivities and the conditions under which they stay inert are known by the subject expert.

Examples of representative protecting groups for the amino group are amides such as, amide acetate, benzoate amide, pivalate amide; carbamates such as, benzyloxycarbonyl (Cbz), *para*-nitrobenzyloxycarbonyl (pNZ), *tert*-butyloxycarbonyl (Boc), 2,2,2-trichloroethoxycarbonyl (Troc), 2-(trimethylsilyl)ethoxycarbonyl (Teoc), 9-fluorenylmethoxycarbonyl (Fmoc), allyloxycarbonyl (Alloc), among others; preferably, Boc or Fmoc.

Examples of representative groups for the carboxyl group are esters such as, *tert*-butyl ester (tBu), allyl ester (All), triphenylmethyl ester (trityl ester, Trt), cyclohexyl ester (cHex), benzyl ester (Bzl), *o*-nitrobenzyl ester, *p*-nitrobenzyl ester, *p*-methoxybenzyl ester, trimethylsilyl ester, among others; invention preferred protecting groups are All, tBu, cHex, Bzl and Trt esters.

Trifunctional amino acids can be protected during synthetic process with temporal or permanent orthogonal protecting groups to N-terminal and C-terminal end protecting groups. Arginine guanidine group can be protected with 2,2,5,7,8-pentamethylchroman-6-sulfonyl (Pmc) group, 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl (Pbf), *para*-toluenesulfonyl (tosyl, Tos) or 4-methoxy-2,3,6-trimethylbenzenesulfonyl (Mtr), among others; the histidine imidazole group can be protected with tosyl group (Tos), *tert*-butyloxycarbonyl group (Boc), trityl group (Trt), methyltrityl group (Mtt) or the 2,4-dinitrophenyl group (Dnp) among others; and the asparagine amide group can be protected with trityl group (Trt) or the xanthyl group (Xan) or being used without protection from the amide group.

35

- 23 -

In a preferred embodiment, the group protection strategy used is the strategy in which amino groups are protected through Boc, carboxyl groups are protected through Bzl, cHex or All, the arginine side chain is protected with Mtr or Tos, the asparagine chain is used without protection and the histidine chain is protected with Tos or Dnp.

5

In another preferred embodiment, the group protection strategy used is the strategy in which amino groups are protected through Fmoc, carboxyl groups are protected with tBu, All or Trt, the arginine side chain is protected with Pmc or Pbf, the asparagine chain is protected with Trt and histidine chain with Trt or Mtt.

10

Examples of these and other additional protecting groups, its introduction and elimination can be found described in [Greene T.W. and Wuts P.G.M., (1999) "Protective groups in organic synthesis" John Wiley & Sons, New York; Atherton B. and Sheppard R.C. (1989) "Solid Phase Peptide Synthesis: A practical approach" IRL Oxford University Press]. The term "protecting groups" includes also polymeric supports employed in solid phase synthesis.

15

When the synthesis is totally or partially done in solid phase, some of the solid supports to be used in the invention method, are polystyrene supports, polyethyleneglycol engrafted in polystyrene and similar ones, such as for example and in a non-limiting sense, *p*-methylbenzhydrylamine resins (MBHA) [Matsueda G.R. and Stewart J.M. (1981) "A *p*-methylbenzhydrylamine resin for improved solid-phase synthesis of peptide amides" *Peptides* 2:45-50], 2-chlorotrityl resins [Barlos K., Gatos D., Kallitsis J., Papaphotiu G., Sotiriu P., Wenqing Y. and Schäfer W. (1989) "Darstellung geschützter Peptid-fragmente unter Einsatz substituierter Triphenylmethyl-harze" *Tetrahedron Lett.* 30:3943-3946; Barlos K., Gatos D., Kapos S., Papaphotiu G., Schäfer W. and Wenqing Y. (1989) "Veresterung von partiell geschützten Peptid-fragmenten mit Harzen. Einsatz von 2-Chlorotritylchlorid zur Synthese von Leu15 -gastrin I" *Tetrahedron Lett.* 30:3947-3951], TentaGel® resins (Rapp Polymere GmbH), ChemMatrix® resins (Matrix Innovation, Inc) and similar ones, which may or may not include a labile spacer such as, 5-(4-aminomethyl-3,5-dimethoxy-phenoxy) valeric acid (PAL) [Albericio F., Kneib-Cordonier N., Biancalana S., Gera L., Masada R.I., Hudson D. and Barany G. (1990) "Preparation and application of the 5-(4-(9-fluorenylmethyloxycarbonyl)aminomethyl-3,5-dimethoxy-phenoxy)-valeric acid (PAL) handle for the solid-phase synthesis of C-terminal peptide amides under mild

35

- 24 -

conditions" *J. Org. Chem.* 55:3730-3743], the 2-[4-aminomethyl-(2,4-dimethoxyphenyl)] phenoxyacetic acid (AM) [Rink H. (1987) "Solid-phase synthesis of protected peptide fragments using a trialkoxy-diphenyl-methylester resin" *Tetrahedron Lett.* 28:3787-3790], Wang [Wang S.S. (1973) "p-Alkoxybenzyl Alcohol Resin and p-Alkoxybenzyloxycarbonylhydrazide Resin for Solid Phase Synthesis of Protected Peptide Fragments" *J.Am.Chem.Soc.* 95:1328-1333] and similar ones which enable simultaneous unprotection and excision of the polymeric support peptide.

Cosmetic or pharmaceutical compositions

10 The peptides of the invention can be administered to inhibit MMPs by any means making contact of the peptides with the site of action thereof in the body of a mammal, preferably human beings, and in the form of a composition containing them.

In this sense, another aspect of the invention is a cosmetic or pharmaceutical composition comprising at least one peptide of general formula (I), stereoisomers thereof, mixtures thereof or its cosmetically or pharmaceutically acceptable salts with at least one cosmetically or pharmaceutically acceptable adjuvant. These compositions can be prepared by conventional methods, known persons skilled in the art [*"Harry's Cosmeticology"*, Eight edition (2000) Rieger M.M., ed., New York Chemical Pub., NY, US; *"Remington: The Science and Practice of Pharmacy"*, Twentieth edition (2003) Genaro A.R., ed., Lippincott Williams & Wilkins, Philadelphia, US].

The peptides of the present invention have a variable water-solubility, according to the nature of their sequence or the possible modifications they have at the N-terminal and/or C-terminal ends. Therefore, the peptides of the present invention can be incorporated into the compositions by means of aqueous solution, and those which are not water-soluble can be solubilized in conventional cosmetically or pharmaceutically acceptable solvents, such as for example and in a non-limiting sense, ethanol, propanol, isopropanol, propylene glycol, glycerin, butylene glycol or polyethylene glycol or any combination thereof.

The cosmetically or pharmaceutically effective amount of the peptides of the invention to be administered to treat a condition, disorder and/or pathology, as well as their dosage, will depend on diverse factors, including the age, condition of the patient, the severity of the disorder or pathology, the route and frequency of administration and on

- 25 -

the particular nature of the peptides to be used.

“Cosmetically or pharmaceutically effective amount” means a non-toxic amount of peptide(s) which is enough to provide the desired effect. Peptides of the invention are
5 used in the cosmetic or pharmaceutical composition of the present invention at concentrations than are cosmetically or pharmaceutically effective to obtain the desired effect; preferably, regarding to the total weight of the composition, between 0.00000001% (by weight) and 20% (by weight); preferably between 0.000001% (by weight) and 20% (by weight), more preferably between 0.0001% (by weight) and 10%
10 (by weight) and even more preferably between 0.0001% (by weight) and 5% (by weight).

The peptides of the invention can also be incorporated into cosmetic or pharmaceutical delivery systems and/or sustained release systems.

15 The term “delivery systems” refers to a diluent, adjuvant, excipient or carrier with which the peptide of the invention is administered. Such cosmetic or pharmaceutical carriers can be liquids; such as water, oils or surfactants, including the ones with petroliferous, animal, plant or synthetic origin; such as for example and in a non-limiting sense peanut oil, soybean oil, mineral oil, sesame oil, castor oils, polysorbates, sorbitan
20 esters, ether sulfate, sulfates, betaines, glucosides, maltosides, fatty alcohols, nonoxynols, poloxamers, polyoxyethylenes, polyethylene glycols, dextrose, glycerol, digitonin and the like.

In “*Remington's Pharmaceutical Sciences*” by E.W. Martin, there are solvents,
25 adjuvants or excipients described as suitable carriers.

The term “sustained release” is used in a conventional sense referring to a delivery system for a compound providing the gradual release of said compound during a period of time and preferably, although not necessarily, with constant levels of
30 compound release throughout a period of time.

Examples of delivery systems or sustained release systems are liposomes, mixed liposomes, milliparticles, microparticles, nanoparticles, solid lipidic nanoparticles, sponges, cyclodextrins, vesicles, micelles, surfactant mixed micelles, phospholipid-
35 surfactant mixed micelles, millispheres, microspheres, nanospheres, lipospheres,

- 26 -

millicapsules, microcapsules, nanocapsules, as well as microemulsions and nanoemulsions, which can be added in order to obtain a greater penetration of the active ingredient and/or to improve its pharmacokinetic and pharmacodynamic properties.

5

The sustained release formulations can be prepared by means of methods known in the state of the art, and the compositions containing them can be administered, for example, by topical administration, including the adhesive patches and non-adhesive patches and microelectric patches, or by systemic administration, such as for example
10 and in a non-limiting sense, by oral, nasal, rectal route, subcutaneous implantation or injection, or direct implantation or injection in a specific part of the body, and, preferably, they have to release a relatively constant amount of the peptides of the invention. The amount of peptide contained in the sustained release formulation will depend, for example, on the site of administration, the kinetics and the duration of the
15 release of the peptide of the invention, as well as the nature of the condition, disorder and/or pathology to be treated or prevented.

The peptides of the present invention can also be absorbed on solid organic polymers or solid mineral supports, such as for example and in a non-limiting sense, talcum
20 powder, bentonite, silica, starch and maltodextrin among others.

The peptides of the invention can also be incorporated into fabrics, non-woven fabrics or medical devices which are in direct contact with the skin, mucosae and/or the scalp, so that they release the peptides of the invention either by biodegradation of the
25 anchorage system to the fabric, non-woven fabric or medical devices or by the friction of the latter with the body, by body moisture, by pH of the skin or by body temperature. Likewise, the fabrics and non-woven fabrics can be used to make garments which are in direct contact with the body. Preferably, the fabrics, non-woven fabrics and medical devices containing the peptides of the invention are used for the treatment and/or care
30 of those conditions, disorders and/or pathologies of the skin, mucosae and/or the scalp, which result from MMP overexpression or an increase in the MMP activity.

Examples of fabrics, non-woven fabrics, garments, medical devices and means for immobilizing the peptides to them, including the delivery systems and/or sustained
35 release systems described above, are described in the literature and are known in the

- 27 -

state of the art [Schaab C.K. (1986) "Impregnating Fabrics With Microcapsules", HAPPI May 1986; Nelson G. (2002) "Application of microencapsulation in textiles" *Int. J. Pharm.* 242:55-62; "Biofunctional Textiles and the Skin" (2006) *Curr. Probl. Dermatol.* v.33, Hipler U.C. and Elsner P., eds. S. Karger AG, Basel, Switzerland; Malcom R.K., McCullagh S.D., Woolfson A.D., Gorman S.P., Jones D.S. and Cuddy J. (2004) "Controlled release of a model antibacterial drug from a novel self-lubricating silicone biomaterial" *J. Cont. Release* 97:313-320]. Preferred fabrics, non-woven fabrics, garments, medical devices are bandages, gauzes, T-shirts, socks, pantyhose, underwear, girdles, gloves, diapers, sanitary napkins, dressings, bedcovers, wipes, hydrogels, adhesive patches, non-adhesive patches, microelectric patches and/or face masks.

Cosmetic or pharmaceutical preparations containing the peptides of the present invention, stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof, can be used in different types of formulations of topical or transdermal application which will include, optionally, the cosmetically or pharmaceutically acceptable excipients necessary for the formulation of the desired dosage form [Faulí i Trillo C. (1993) in "Tratado de Farmacia Galénica", Luzán 5, S.A. Ediciones, Madrid].

The topical or transdermal application formulations can be presented in any solid, liquid or semi-solid dosage form, such as for example and in a non-limiting sense, creams, multiple emulsions such as for example and in a non-limiting sense emulsions of oil and/or silicone in water, emulsions of water in oil and/or silicone, emulsions of the water/oil/water or water/silicone/water type and emulsion of the oil/water/oil or silicon/water/silicon type, anhydrous compositions, aqueous dispersions, oils, milks, balsams, foams, lotions, gels, cream gels, hydroalcoholic solutions, hydroglycolic solutions, liniments, saline solutions, soaps, shampoos, conditioners, sera, polysaccharide films, unguents, mousses, ointments, powders, bars, pencils and aerosols or sprays, including leave-on formulations and rinse-off formulations. These topical or transdermal application formulations can be incorporated by techniques known by the person skilled in the art to different types of solid accessories, such as for example and in a non-limiting sense, wipes, hydrogels, adhesive patches, non-adhesive patches, microelectric patches or face masks; or they can be incorporated to different makeup line products such as makeup foundations; such as for example fluid

- 28 -

or solid makeup foundations, makeup removal lotions, makeup removal milks, under eye concealers, eye shadows, lipsticks, lip protectors, lip glosses and powders, among others.

- 5 The cosmetic or pharmaceutical compositions of the invention can include agents which increase the percutaneous absorption of the peptides of the present invention, such as for example and in a non-limiting sense, dimethylsulfoxide, dimethylacetamide, dimethylformamide, surfactants, azone (1-dodecylazacycloheptan-2-one), alcohol, urea, ethoxydiglycol, acetone, propylenglycol or polyethyleneglycol among others.
- 10 Likewise, the cosmetic or pharmaceutical compositions object of the present invention can be applied to local areas to be treated by iontophoresis, sonophoresis, electroporation, microelectric patches, mechanic pressure, osmotic pressure gradient, occlusive treatment, microinjections, or pressure needle-free injections by means of pressure, such as for example, oxygen pressure injections, or any combination thereof,
- 15 for the purpose of achieving greater penetration of the peptide of the invention. The application area will be determined by the nature of the condition, disorder and/or pathology to be prevented or treated.

- Likewise, the cosmetic compositions containing the peptides of the present invention,
- 20 stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof can be used in different kinds of formulations for their oral administration, preferably in the form of oral cosmetics, such as for example and in a non-limiting sense, capsules, including gelatin capsules, tablets, including sugar coated tablets, powders, granulated forms, chewing gums, solutions, suspensions, emulsions,
- 25 syrups, polysaccharides films, jellies or gelatins as well as any other presentation known by a person skilled in the art. Particularly, the peptides of the invention can be incorporated into any form of functional or fortified food such as for example and in a non-limiting sense, diet bars or compact or non-compact powders. Such powders can be solubilized in water, soda, dairy products, soy derivatives, or they can be
- 30 incorporated into diet bars. The peptides of the present invention can be formulated with excipients and adjuvants usual for oral compositions or food supplements such as for example and in a non-limiting sense, fat components, aqueous components, humectants, preservatives, texturizing agents, flavors, aromas, antioxidants and colorants commonly found in the food industry.

35

- 29 -

Cosmetic or pharmaceutical compositions containing the peptides of the present invention, stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof can be administrated, not only by transdermal or topical routes, but also by any other type of suitable route, for example by oral or parenteral route, for which they will include the pharmaceutically acceptable excipients necessary for the formulation of the desired dosage form. In the context of the present invention, the term "parenteral" includes nasal route; rectal route; subcutaneous, intradermal or intravascular injections; such as intravenous, intramuscular, intravitreal, spinal, intracranial, intraarticular, intrathecal, and intraperitoneal injections; as well as any other similar injection or infusion technique. A revision of the different pharmaceutical dosage form of active ingredients and of the necessary excipients for obtaining said dosage form can be found in *"Tratado de Farmacia Galénica"*, C. Faulí i Trillo, 1993, Luzán 5, S.A. Ediciones, Madrid.

Among the cosmetically or pharmaceutically acceptable adjuvants described in the present invention, there are included additional ingredients commonly used in compositions for the treatment and/or care of skin, mucosae and/or scalp such as for example and in a non-limiting sense, other agents inhibiting MMP, agents stimulating or inhibiting melanine synthesis, whitening or depigmenting agents, propigmenting agents, self-tanning agents, anti-age agents, NO-synthase inhibiting agents, antioxidant agents, free radicals scavengers and/or anti-atmospheric pollution agents, anti-glycation agents, emulsifying agent, emollients, organic solvents, liquid propellants, skin conditioners such as humectants, substances retaining moisture, alphahydroxyacids, betahydroxyacids, moisturizers, epidermic hydrolytic enzymes, vitamins, pigments or colorants, dyes, gelling polymers, thickeners, surfactants, softeners, anti-wrinkle agents, agents capable of reducing or treating bags under the eyes, exfoliating agents, antimicrobial agents, fungicide agents, fungistatic agents, bactericide agents, bacteriostatic agents, agents stimulating the synthesis of dermal or epidermal macromolecules and/or agents capable of preventing or inhibiting their degradation, agents stimulating collagen synthesis, agents stimulating elastin synthesis, agents stimulating decorine synthesis, agents stimulating laminin synthesis, agents stimulating defensin synthesis, agents stimulating chaperone synthesis, agents stimulating aquaporin synthesis, agents stimulating hyaluronic acid synthesis, agents stimulating the synthesis of lipids and components of the stratum corneum (ceramides, fatty acids, etc.), other agents inhibiting collagen degradation, agents inhibiting elastin

- 30 -

degradation, agents stimulating fibroblast proliferation, agents stimulating keratinocyte proliferation, agents stimulating adipocyte proliferation, agents stimulating melanocyte proliferation, agents stimulating keratinocyte differentiation, agents stimulating adipocyte differentiation, agents inhibiting acetylcholinesterase, dermorelaxing agents, agents stimulating glycosaminoglycan synthesis, DNA repairing agents, DNA protecting agents, anti-itching agents, agents for the treatment and/or care of sensitive skin, firming agents, anti-stretch mark agents, astringent agents, agents regulating sebum production, agents stimulating lipolysis, anticellulitic agents, agents stimulating healing, coadjutant healing agents, agents stimulating reepithelizing, coadjutant reepithelizing agents, cytokine growth factors, calming agents, anti-inflammatory agents, agents acting on capillary circulation and/or microcirculation, agents stimulating angiogenesis, agents inhibiting vascular permeability, agents acting on cell metabolism, agents intended to improve the dermal-epidermal junction, agents inducing hair growth, agents inhibiting or delaying hair growth, preservatives, perfumes, chelating agent, plant extracts, essential oils, marine extracts, agents coming from a bio-fermentation process, mineral salts, cell extracts and sunscreens (organic or mineral photoprotecting agents that are active against ultraviolet A and/or B rays), among others, as long as they are physically and chemically compatible with the remaining components of the composition and specially with the peptides of general formula (I) contained in the composition of the present invention. Likewise, the nature of these additional ingredients should not unacceptably alter the benefits of the peptides of the present invention. Said additional ingredients can be synthetic or natural such as for example plant extracts, or come from a bio-fermentation process. Additional examples can be found in *CTFA Cosmetic Ingredient Handbook, Eleventh Edition (2006)*.

An additional aspect of the present invention refers to a cosmetic or pharmaceutical composition containing a cosmetically or pharmaceutically effective amount of at least one peptide of the invention, stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof and also a cosmetically or pharmaceutically effective amount of at least one synthetic component, natural extract or product from biofermentation process with activity inhibiting MMP such as for example and in a non-limiting sense, natural extracts containing ursolic acid, isoflavones like genistein, quercetin, carotenoid, lycopene, soybean extract, blueberry extract, rosemary extract, *Trifolium pratense* (red clover) extract, *Phormium tenax*

- 31 -

(formio) extract, kakkon-tō extract, salvia extract, retinol and its derivatives, retinoic acid and its derivatives, sapogenins such as for example and in a non-limiting sense, diosgenin, hecogenin, smilagenin, sarsapogenin, tigogenin, yamogenin and yucagenin among others, Collalift® [INCI: Hydrolyzed Malt Extract], Juvenesce [INCI: Ethoxydiglicol and Caprylic Triglyceride, Retinol, Ursolic Acid, Phytonadione, Ilomastat] or EquiStat [INCI: Pyrus Malus Fruit Extract, Glycine Soja Seed Extract] marketed by Coletica/Engelhard, Pepha®-Timp [INCI: Human Oligopeptide-20], Regu-Age [INCI: Hydrolyzed Rice Bran Protein, Glycine Soja Protein, Oxido Reductases] or Colhibin [INCI: Hydrolyzed Rice Protein] marketed by Pentapharm, Lipeptide [INCI: Hydrolyzed vegetable protein] marketed by Lipotec, Litchiderm™ [INCI: Litchi Chinensis pericarp extract] or Arganyl™ [INCI: Argania Spinosa Leaf Extract] marketed by Laboratories Sérobiologiques/Cognis, MDI Complex® [INCI: glycosaminoglycans] or ECM-Protect® [INCI: Water (Aqua), Dextran, Tripeptide-2] marketed by Atrium Innovations, Dakaline [INCI: Prunus amygdalus dulcis, Anogeissus leiocarpus bark extract] marketed by Soliance, Homeostatine [INCI: Enteromorpha compressa, Caesalpinia Spinosa] marketed by Provital, Timp-Peptide [proposed INCI: Acetyl Hexapeptide] or ECM Moduline [proposed INCI: Palmitoyl tripeptide] marketed by Infinitec Activos, IP2000 [INCI: Dextran, Trifluoroacetyl tripeptide-2] marketed by Institut Européen de Biologie Cellulaire, Actimp 1.9.3® [INCI: Hydrolyzed Lupine Protein] marketed by Expanscience Laboratories, Vitaderm® [INCI: Alcohol, Water (Aqua), Glycerin, Hydrolyzed Rice Protein, Ilex Aquifolium Extract, Sodium Ursolate, Sodium Oleanolate] marketed by Rahn, adapalene, tetracycline and its derivatives such as, minocycline, rolitetracycline, chlortetracycline, metacycline, oxytetracycline, doxycycline, demeclocycline and its salts, Batimastat [BB94; 4-(*N*-hydroxyamine) -2*R*-isobutyl -3*S*-(thiophen-2-ylthiomethyl)succinyl]-L-phenylalanine-*N*-methanamide], Marimastat [BB2516; 2*S*-[*N*4(*R*^{*}),2*R*^{*},3*S*]]-*N*4[2,2-dimethyl-1-[methylaninocarbonyl]propyl]-*N*1,2-dihydroxy-3-(2-methylpropyl) butanediamine], among others.

Likewise, the cosmetic or pharmaceutical compositions of the present invention can additionally contain a cosmetically or pharmaceutically effective amount of at least one analgesic and/or anti-inflammatory compound aiming at reducing swelling and irritation associated to inflammatory processes wherein there is MMP overexpression and/or overactivity. Among these compounds we can highlight synthetic compounds such as hydrocortisone, clobetasol, dexamethasone, prednisone, paracetamol, acetylsalicylic acid, amoxiprin, benorilate, choline salicylate, diflunisal, faislamine, methyl salicylate,

- 32 -

magnesium salicylate, salsalate, diclofenac, aceclofenac, acemetacin, bromfenac, etodolac, indomethacin, sulindac, tolmetin, ibuprofen, carprofen, fenbufen, fenoprofen, flurbiprofen, ketoprofen, ketorolac, loxoprofen, naproxen, oxaprozine, tiaprofenic acid, suprofen, mefenamic acid, meclofenamate, meclofenamic acid, tolfenamic acid, nabumetone, phenylbutazone, azapropazone, metamizole, oxyphenbutazone, sulfinpyrazone, piroxicam, lornoxicam, meloxicam, tenoxicam, celecoxib, etoricoxib, lumiracoxib, parecoxib, rofecoxib, valdecoxib, nimesulide, licofelone, omega-3 fatty acid and its biometabolites, morphine, codeine, oxycodone, hydrocodone, diamorphine, pethidine, tramadol, brupenorphine, benzocaine, lidocaine, chloroprocaine, tetracaine, procaine, tricyclic antidepressants, amitriptyline, carbamazepine, gabapentin, pregabalin, sabolol, panthenol, biotin, tocopheryl lauriminodipropionate phosphate, disodium, ciclopirox olamine, nordihydroguaiaretic acid, NeutrazenTM [INCI: Water (Aqua), Butylenen Glycol, Dextran, Palmitoyl Tetrapeptide-8] marketed by Atrium Innovations, Meliprene[®] [INCI: Dextran, Acetyl Heptapeptide-1] marketed by Institut Europeen de Biologie Cellulaire, coenzyme Q10 or alkylglycerol ethers, or natural extracts or essential oils with analgesic and/or anti-inflammatory activity such as for example and in a non-limiting sense, madecassoside, echinacine, amaranth seed oil, sandalwood oil, placenta extract, peach tree leaf extract, *Aloe vera*, *Arnica montana*, *Artemisia vulgaris*, *Asarum maximum*, *Calendula officinalis*, *Capsicum*, *Centipeda cunninghamii*, *Chamomilla recutita*, *Crinum asiaticum*, *Hamamelis virginiana*, *Harpagophytum procumbens*, *Hypericum perforatum*, *Lilium candidum*, *Malva sylvestris*, *Melaleuca alternifolia*, *Origanum majorana*, *Salix alba*, *Silybum marianum*, *Tanacetum parthenium* or *Uncaria guianensis*, among others.

Additionally, the present invention refers to a cosmetic or pharmaceutical composition comprising a cosmetically or pharmaceutically effective amount of at least one peptide of general formula (I), stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof and also a cosmetically or pharmaceutically effective amount of at least one extract or extract combination with healing and/or reepithelizing activity or effective as coadjuvants in healing and/or reepithelizing processes such as extracts of *Centella asiatica*, *Rosa moschata*, *Echinacea angustifolia*, *Symphytum officinal*, *Equisetum arvense*, *Hypericum perforatum*, *Mimosa tenuiflora*, *Aloe vera*, Polyplant[®] Epithelizing [INCI: Calendula Officinalis, Hypericum Perforatum, Chamomilla Recutita, Rosmarinus Officinalis] marketed by Provital, Cytokinol[®] LS 9028 [INCI: Hydrolyzed Casein, Hydrolyzed Yeast Protein, Lysine HCl]

- 33 -

marketed by Laboratories Sérobiologiques/Cognis or Deliner® [INCI: Zea May (Corn) Kernel Extract] marketed by Coletica/Engelhard among others, and/or a cosmetically or pharmaceutically effective amount of at least one synthetic compound, extract or product coming from a biofermentation process with healing and/or reepithelizing activity or effective as coadjuvants in healing and/or reepithelizing processes such as for example and in a non-limiting sense, cadherins, integrins, selectins, hyaluronic acid receptors, immunoglobulins, fibroblast growth factors, connective tissue growth factor, platelet growth factor, endothelial vascular growth factor, epidermal growth factor, insulin-like growth factor, keratinocyte growth factor, colony-stimulating factors, transforming growth factor-beta, tumor necrosis factor-alpha, interferons, interleukins, matrix metalloproteases, protein tyrosine phosphatase receptors, Antarcticine® [INCI: Pseudoalteromonas Ferment Extract] or Decorinyl™ [INCI: Tripeptide-10 Citrulline], marketed by Lipotec, among others.

An additional aspect of the present invention refers to a cosmetic or pharmaceutical composition comprising a cosmetically or pharmaceutically effective amount of at least one peptide of general formula (I), stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof and also a cosmetically or pharmaceutically effective amount of at least one extract with anti-wrinkle and/or antiaging activity such as for example and in a non-limiting sense, extracts of *Vitis vinifera*, *Rosa canina*, *Curcuma longa*, *Iris pallida*, *Theobroma cacao*, *Ginkgo biloba*, or *Dunaliella salina*, among others, and/or at least one synthetic compound, extract or product coming from a biofermentation process with anti-wrinkle and/or antiaging activity such as for example and in a non-limiting sense, Matrixyl® [INCI: Palmitoyl Pentapeptide-3] or Matrixyl 3000® [INCI: Palmitoyl Tetrapeptide-3, Palmitoyl Oligopeptide] marketed by Sederma, Vialox® [INCI: Pentapeptide-3] or Syn-ake® [INCI: Dipeptide Diaminobutyryl Benzylamide Diacetate] marketed by Pentapharm, Myoxinol™ [INCI: Hydrolyzed Hibiscus Esculentus Extract] marketed by Laboratoires Sérobiologiques/Cognis, Algisum C® [INCI: Methylsilanol Mannuronate] or Hydroxyprolisilane CN® [INCI: Methylsilanol Hydroxyproline Aspartate] marketed by Exsymol, Argireline® [INCI: Acetyl Hexapeptide-8] (Acetyl hexapeptide-8), Leuphasyl® [INCI: Pentapeptide-18] (Pentapéptido-18), Aldenine® [INCI: Hydrolyzed wheat protein, hydrolyzed soy protein, tripeptide-1], Trylagen™ [INCI: Pseudoalteromonas Ferment Extract, Hydrolyzed Wheat Protein, Hydrolyzed Soy Protein, Tripeptide-10 Citrulline, Tripeptide-1]; Eyeseryl® [INCI: Acetyl Tetrapeptide-5] or Lipochroman-6 [INCI:

- 34 -

- Dimethylmethoxy Chromanol] marketed by Lipotec, Kollaren® [INCI: Tripeptide-1, Dextran] marketed by Institut Europeen de Biologie Cellulaire, Collaxyl® [INCI: Hexapeptide-9] or Quintescine® [INCI: Water, Butylene Glycol, Dipeptide-4] marketed by Vincience, BONT-L-Peptide [proposed INCI: Palmitoyl Hexapeptide] marketed by
- 5 Infinitec Activos, Ca²⁺ channel antagonists such as alverine, manganese or magnesium salts, certain secondary or tertiary amines, retinol and its derivatives, idebenone and its derivatives, coenzyme Q10 and derivatives, boswellic acid and its derivatives or agonist of chloride channel, among others.
- 10 An additional aspect of present invention refers to a cosmetic or pharmaceutical composition comprising a cosmetically or pharmaceutically effective amount of at least one peptide of general formula (I), and also a cosmetically or pharmaceutically effective amount of at least one extract or combination of extracts with refirming, rethickening and/or restructuring activity, such as for example and in a non-limiting sense, *Malpighia*
- 15 *punicitolia*, *Cynara scolymus*, *Gossypium herbaceum*, *Aloe Barbadensis*, *Panicum miliaceum*, *Morus nigra*, *Sesamum indicum*, *Glycine soy*, *Triticum vulgare*, Pronalen® Refirming HSC [INCI: Triticum vulgare, Silybum Marianum, Glycine Soy, Equisetum Arvense, Alchemilla Vulgaris, Medicago Sativa, Raphanus Sativus] or Polyplant® Refirming [INCI: Coneflower, Asiatic Centella, Fucus, Fenugreek] marketed
- 20 by Provital, Lanablue® [INCI: Sorbitol, Algae Extract] marketed by Atrium Innovations, Pepha®-Nutrix [INCI: Natural Nutrition Factor] marketed by Pentapharm, or plant extracts containing isoflavones or else at least one synthetic compound, extract or product coming from a biofermentation process with refirming, rethickening and/or restructuring activity, such as for example and in a non-limiting sense, Biopeptide EL™
- 25 [INCI: Palmitoyl Oligopeptide], Biopeptide CL™ [INCI: Palmitoyl Oligopeptide], Vexel® [INCI: Water (Aqua), Propylene Glycol, Lecithin, Caffeine, Palmitoyl Carnitine], Matrixyl® [INCI: Palmitoyl Pentapeptide-3], Matrixyl 3000® [INCI: Palmitoyl Tetrapeptide-3, Palmitoyl Oligopeptide] or Bio-Bustyl™ [INCI: Glyceryl Polymethacrylate, Rahnella Soy Protein Ferment, Water (Aqua), Propylene Glycol,
- 30 Glycerin, PEG-8, Palmitoyl Oligopeptide] marketed by Sederma, Dermosaccharides® HC [INCI: Glycerin, Water (Aqua), Glycosaminoglycans, Glycogen], Aglycal® [INCI: Mannitol, Cyclodextrin, Glycogen, Aratostaphylos Uva Ursi Leaf Extract], Cytokinol® LS [INCI: Hydrolyzed Casein, Hydrolyzed Yeast Protein, Lysine HCL] or Firmiderm® LS9120 [INCI: Terminalia Catappa Leaf extract, Sambucus
- 35 Negra Flower Extract, PVP, Tannic Acid] marketed by Laboratoires

- 35 -

Sérobiologiques/Cognis, Liftline® [INCI: Hydrolyzed wheat protein], Raffermin® [INCI: Hydrolyzed Soy Flour] or Ridulisse C® [Hydrolyzed Soy Protein] marketed by Silab, Serilesine® [INCI: hexapeptide-10] or Decorinyl™ [INCI: Tripeptide-10 Citrulline] marketed by Lipotec, Ursolisome® [INCI: Lecithin, Ursolic Acid, Atelocollagen, Xanthan Gum, Sodium Chondroitin Sulfate] or Collalift® [INCI: Hydrolyzed Malt Extract] marketed by Coletica/Engelhard, Syn®-Coll [INCI: Palmitoyl Tripeptide-5] marketed by Pentapharm, Hydriame® [INCI : Water (Aqua), Glycosaminoglycans, Sclerotium Gum] marketed by Atrium Innovations or IP2000 [INCI: Dextran, Trifluoroacetyl tripeptide-2] marketed by Institut Europeen de Biologie Cellulaire, among others.

10

Applications

Another aspect of the present invention refers to a cosmetic or pharmaceutical method for the treatment and/or care of those conditions, disorders and/or pathologies of the skin, mucosae and/or scalp in mammals, preferably in human beings, who benefit from MMP inhibition; comprising the administration of an effective amount of at least one peptide of general formula (I), stereoisomers thereof, mixtures thereof or cosmetically or pharmaceutically acceptable salts thereof, preferably in a cosmetic or pharmaceutical composition containing them. The present invention also provides a cosmetic or pharmaceutical method to inhibit MMPs, preferably skin, mucosae and/or scalp MMPs.

20

Likewise, the present invention provides a cosmetic or pharmaceutical method for treatment and/or care of those conditions, disorders and/or pathologies of the skin, mucosae and/or scalp which are caused by MMP overexpression or by an increase in the MMP activity, comprising the application on skin, mucosae and/or scalp or the oral or parenteral administration of a cosmetic or pharmaceutical composition containing at least one peptide of the invention, stereoisomers thereof, mixtures thereof or cosmetically or pharmaceutically acceptable salts thereof.

25

Preferably, among the conditions, disorders and/or pathologies of the skin, mucosae and/or scalp to be treated and/or cared caused by MMP overexpression or by an increase of MMP activity, there are included acne, rosacea, psoriasis, dermatitis, atopic dermatitis, eczema, sensitive skins, gingivitis, periodontitis, skin cancer, tumor invasions, aging skin, photoaging skin, wrinkles, expression wrinkles, stretch marks, keloids, hypertrophic scars, cellulitis, orange peel skin, tumor metastasis, ulcers,

35

- 36 -

diabetic ulcers, telangiectasia, cuperosis, varicose veins, eye dark circles, bags under the eye, alopecia and hair loss.

5 Compositions containing the peptides of the present invention, stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof can be applied on skin, mucosae and/or scalp or be orally or parenterally administrated, as required to treat and/or care for a condition, disorder and/or pathology.

10 Application or administration frequency can vary widely, depending on each individual's needs; there being suggested an application or administration range from once a month up to ten times a day, preferably from once a week up to four times a day, more preferably from three times a week up to three times a day, even more preferably, once or twice a day.

15 An additional aspect of the present invention refers to the use of at least one peptide of general formula (I), stereoisomers thereof, mixtures thereof or cosmetically or pharmaceutically acceptable salts thereof, in the preparation of a cosmetic or pharmaceutical composition for the treatment and/or care of skin, mucosae and/or scalp.

20 Additionally, the present invention refers to the use of at least one peptide of general formula (I), stereoisomers thereof, mixtures thereof or cosmetically or pharmaceutically acceptable salts thereof, in the preparation of a cosmetic or pharmaceutical composition for MMP inhibition, preferably skin, mucosae and/or scalp MMPs.

25 Likewise, another aspect of the present invention refers to the use of at least one peptide of general formula (I), stereoisomers thereof, mixtures thereof or cosmetically or pharmaceutically acceptable salts thereof, in the preparation of a cosmetic or pharmaceutical composition for the treatment and/or care of skin, mucosae and/or
30 those conditions, disorders and/or pathologies of the scalp caused by a MMP overexpression or by an increased MMP activity. Preferably, cosmetic or pharmaceutical compositions are created to treat and/or care those skin, mucosae and/or scalp areas affected by acne, rosacea, psoriasis, dermatitis, atopic dermatitis, eczema, sensitive skins, gingivitis, periodontitis, skin cancer, tumor invasions, aging
35 skin, photoaging skin, wrinkles, expression wrinkles, stretch marks, keloids,

- 37 -

hypertrophic scars, cellulitis, orange peel skin, tumor metastasis, ulcers, diabetic ulcers, telangiectasia, cuperosis, varicose veins, eye dark circles, bags under the eyes, alopecia and hair loss.

5 According to another important aspect, the present invention refers to the use of a peptide of general formula (I) in the preparation of a cosmetic or pharmaceutical composition for skin treatment aiming at reducing, delaying and/or preventing aging and photoaging signs.

10 In an additional embodiment, the present invention refers to the use of at least one of the peptides of general formula (I), stereoisomers thereof, mixtures thereof or cosmetically or pharmaceutically acceptable salts thereof, in the preparation of a cosmetic or pharmaceutical composition for oral treatment or hygiene. Preferably, the cosmetic or pharmaceutical composition is used for gingivitis and periodontitis
15 treatment or prevention. Oral hygiene cosmetic or pharmaceutical composition examples include toothpaste, oral elixirs for mouth rinsing or chewing gum, among others.

An additional aspect of the present invention refers to the use of at least one of the
20 peptides of general formula (I), stereoisomers thereof, mixtures thereof or cosmetically or pharmaceutically acceptable salts thereof, in the preparation of a cosmetic or pharmaceutical composition for hair treatment or hygiene. Preferably, the cosmetic or pharmaceutical composition is employed for alopecia and hair loss treatment or prevention. Hair hygiene cosmetic or pharmaceutical composition examples include
25 shampoo, hair conditioners, hair lotions, hair tonics or scalp masks, among others.

Another aspect of the present invention refers to the use of at least one of the peptides of general formula (I), stereoisomers thereof, mixtures thereof or cosmetically or pharmaceutically acceptable salts thereof, in the preparation of a cosmetic or
30 pharmaceutical composition for body treatment or hygiene.

The following specific examples are useful in illustrating the nature of the present invention. These examples are solely included for illustrative purposes and are not to be interpreted as limitations to the herein claimed invention.

EXAMPLES

General Methodology

All reagents and solvents are synthetic quality and are used without any additional treatment.

Abbreviations

The abbreviations used for amino acids follow the IUPAC-IUB Commission on Biochemical Nomenclature specified in *Eur. J. Biochem.* (1984) 138:9-37 and in *J. Biol. Chem.* (1989) 264:633-673.

10

Ac, acetyl; All, alyl; Alloc, allyloxycarbonyl; AM, 2-[4-aminomethyl (2,4 dimethoxyphenyl)] acid; Arg, arginine; Asn, asparagine; Boc, *tert*-butoxycarbonyl; Bzl, benzyl; Cbz, benzyloxycarbonyl; cHex, cyclohexyl; Cit, citrulline; CITrt-®, 2-chlorotrityl resin; cps, centipoise; C-terminal, carboxy-terminal; DCM, dichloromethane; DIEA, *N,N*-diisopropylamine; DIPCDI, *N,N'*-diisopropylcarbodiimide; DMF, *N,N*-dimethylformamide; Dnp, 2,4-dinitrophenyl; DPPC, dipalmitoylphosphatidylcholine; equiv, equivalent; ESI-MS, electrospray ionization mass spectrometry; Fmoc, 9-fluorenylmethoxycarbonyl; His, histidine; HOAt, 1-hydroxybenzotriazole; HOBt, 1-hydroxybenzotriazole; HPLC, high-performance liquid chromatography; INCI, International Nomenclature of Cosmetic Ingredients; MBHA, resin *p*-metalbenzhydrylamine resin; MeCN, acetonitrile; MeOH, methanol; mLV, multilaminar vesicles; MMP, matrix metalloproteases; Mtr, 4-methoxy-2,3,6 trimethylbenzenesulfonyl; Mtt, methyltrityl; *N*-terminal, amino-terminal; PAL, 5-(4-aminomethyl-3,5-dimethoxyphenoxy) valeric acid; Palm, palmitoyl; Pbf, 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl ; Pmc, 2,2,5,7,8- pentamethyl-croman-6-sulfonic; pNZ, *p*-nitrobenzyloxycarbonyl; ®, resin; tBu, *tert*-butyl; Teoc, 2-(trimethylsilyl)ethoxycarbonyl; TFA, trifluoroacetic acid; THF, tetrahydrofuran; TIMP, tissue inhibitor of metalloproteases matrix; TIS, triisopropylsilane; Tos, *para*-toluenesulfonyl or tosyl; Troc, 2,2,2- trichloroethyl-oxycarbonyl; Trt, triphenylmethyl or trityl; ULV, unilaminar vesicles; UV, ultraviolet; Xan, xanthyl.

30

Chemical synthesis

All synthetic processes are performed with polypropylene syringes equipped with porous polyethylene discs or in Pyrex® reactors equipped with a porous plate. Soluble solvents and reagents are eliminated by suction. Fmoc group elimination is carried out

35

- 39 -

with piperidine-DMF (2:8, v/v) (1 x 1min, 1 x 5min; 5mL/g resin) [Lloyd-Williams P., Albericio F. and Giralt, E. (1997) "Chemical Approaches to the Synthesis of Peptides and Proteins" CRC, Boca Raton, FL, USA].

- 5 Washings between unprotection, coupling and again unprotection stages have been carried out with DMF (3 x 1min) using 10mL solvent/g resin each time. Coupling reactions have been carried out with 3mL solvent/g resin. Coupling control is carried out by ninhydrin test [Kaiser E., Colescott R.L., Bossinger C.D. and Cook P.I. (1970) "Color test for detection of free terminal amino groups in the solid-phase synthesis of peptides" *Anal. Biochem.* 34:595-598]. All synthetic transformations and washing have
10 been carried out at room temperature.

EXAMPLE 1

Obtaining Fmoc-AA₁-AA₂-AA₃-AA₄-O-2-CITrt-®

15

- 3.5g Fmoc-L-Cit-OH were added (8.8mmol, 1equiv), dissolved in 55mL DCM, then 1.3mL DIEA (7.6mmol, 0.86equiv) were added on the dried 2-chlorotrityl resin (5.5g, 8.8mmol). That solution was left in agitation during 5min, after which 2.5mL DIEA were added (14.6mmol, 1.66equiv). Then, it reacted during 40min. Remaining chloride
20 groups were blocked by treating them with 4.4mL of MeOH.

- Fmoc N-terminal group was unprotected as described in the general methods and 13.63g Fmoc-L-His(Trt)-OH or 14.27g Fmoc-L-Arg(Pbf)-OH (22mmol, 2.5equiv) were incorporated the peptidyl-resin in presence of DIPCDI (3.39mL, 22mmol, 2.5equiv) and
25 HOBt (3.37g, 22mmol, 2.5equiv) using DMF as solvent during one hour. Afterwards, the resin was washed as described in the general methods and the Fmoc group unprotection treatment was repeated so as to incorporate the subsequent amino acid. Following the described protocols, 13.63g Fmoc-L-His(Trt)-OH or 13.13g Fmoc-L-Asn(Trt)-OH (22mmol, 2.5equiv) and 14.27g of Fmoc-L-Arg(Pbf)-OH (22mmol,
30 2.5equiv) were sequentially coupled in presence of each 37g HOBt (22mmol, 2.5equiv) and 3.39mL DIPCDI (22mmol, 2.5equiv) coupling.

Once the synthesis was over, peptidyl-resins were washed with DCM (5 x 3min) and dried by nitrogen flow.

35

- 40 -

EXAMPLE 2*Obtaining Fmoc-AA₁-AA₂-AA₃-AA₄-AM-MBHA-®*

6.85g Fmoc-AM-MBHA resin of functionalization 0.73mmol/g (5mmol) were treated
5 with piperidine-DMF, according to the described protocol so as to eliminate the Fmoc
group. 9.9g Fmoc-L-Cit-OH (25mmol, 5equiv) were added on the unprotected resin in
presence of DIPCDI (3.85mL, 25mmol, 5equiv) and HOBt (3.85g, 25mmol, 5equiv)
using DMF as solvent during 1 hour.

10 The resin was later washed as described in the general methods and the Fmoc group
unprotection treatment was repeated so as to incorporate the subsequent amino acid.
According to the described protocols, 15.49g Fmoc-L-His(Trt)-OH or 16.22g
Fmoc-L-Arg(Pbf)-OH (25mmol, 5equiv), 15.49g Fmoc-L-His(Trt)-OH or 14.92g
Fmoc-L-Asn(Trt)-OH (25mmol, 5equiv) and 16.22g Fmoc-L-Arg(Pbf)-OH (25mmol,
15 5equiv) were sequentially coupled in presence of each 3.85g HOBt (25mmol, 5equiv)
and 3.85mL DIPCDI (25mmol, 5equiv) coupling.

Once the synthesis was over, peptidyl-resins were washed with DCM (5 x 3min) and
dried by nitrogen flow.

20

EXAMPLE 3*Fmoc N-terminal protecting group general excision process*

Fmoc N-terminal Group was unprotected from peptidyl-resins obtained in examples 1
25 and 2 as it is described in the general methods (20% piperidine in DMF, 1 x 5min + 1 x
20min). Peptidyl-resins were washed with DMF (5 x 1min), DCM (4 x 1min), Diethyl
ether (4 x 1min) and vacuum dried.

EXAMPLE 4

30

*R₁ palmitoyl group introduction process: Obtaining Palm-AA₁-AA₂-AA₃-AA₄-O-2-CITrt-®
and Palm-AA₁-AA₂-AA₃-AA₄-AM-MBHA-®.*

2.56g of palmitic acid (10mmol, 10equiv) pre-dissolved in DMF (1mL) were added on
35 1mmol of the peptidyl-resins obtained in example 3, in presence of 1.53g HOBt

- 41 -

(10mmol, 10equiv) and 1.54mL DIPCDI (10mmol, 10equiv). They were left to react during 15 hours, after which resins were washed with THF (5 x 1min), DCM (5 x 1min), DMF (5 x 1min), MeOH (5 x 1min), DMF (5 x 1min), THF (5 x 1min), DMF (5 x 1min), DCM (4 x 1min), ether (3 x 1min), and vacuum dried.

5

EXAMPLE 5

R₁ acetyl group introduction process: Obtaining Ac-AA₁-AA₂-AA₃-AA₄-O-2-CITrt-® and Ac-AA₁-AA₂-AA₃-AA₄-AM-MBHA-®.

10 1mmol of the peptidyl-resins obtained in example 3 were treated with 25equiv Acetic anhydride in presence of 25equiv DIEA using 5mL DMF as solvent. They were left to react during 30min, after which peptidyl-resins were washed with DMF (5 x 1min), DCM (4 x 1min), diethyl ether (4 x 1min) and vacuum dried.

15 **EXAMPLE 6**

Polymeric support excision process: Obtaining H-AA₁-AA₂-AA₃-AA₄-OH, Ac-AA₁-AA₂-AA₃-AA₄-OH, Palm-AA₁-AA₂-AA₃-AA₄-OH, H-AA₁-AA₂-AA₃-AA₄-NH₂, Ac-AA₁-AA₂-AA₃-AA₄-NH₂ and Palm-AA₁-AA₂-AA₃-AA₄-NH₂.

20 200mg of dried peptidyl-resins obtained in examples 3, 4 and 5 were treated with 5mL of TFA-TIS-H₂O (90:5:5) during 2 hours at room temperature with agitation. Filtrations were collected on 50mL cold diethyl ether, filtered through polypropylene syringes equipped with porous polyethylene discs and washed 5 times with 50mL diethyl ether. Final-precipitates were vacuum dried.

25

HPLC analysis of the peptides obtained in gradients of MeCN (+0.07% TFA) in H₂O (+0.1% TFA) showed purity higher than 85% in all cases. The identity of the peptides obtained was confirmed by ES-MS.

30 **EXAMPLE 7**

Polymeric support excision process and functionalization with substituted R₂ amine: Obtaining Ac-AA₁-AA₂-AA₃-AA₄-NH-(CH₂)₁₅-CH₃.

35 Ac-AA₁-AA₂-AA₃-AA₄-OH peptides with completely protected side chains were obtained by treating 150mg of Ac-AA₁-AA₂-AA₃-AA₄-O-2-CITrt-® peptidyl-resins from example 6,

- 42 -

previously vacuum dried in presence of KOH, with 3mL of a 3% solution of TFA in DCM during 5min. Filtrations were collected on 50mL of cold diethyl ether and the treatment was repeated three times. Ethereal solutions were submitted to rotavaporation at room dryness and temperature, precipitates were resuspended in 50% MeCN in H₂O and lyophilised. 10mg of the obtained raw peptides were weighted, and 3equiv hexadecylamine and 25mL DMF anhydride were added. 2equiv DIPCDI were added and left to react with magnetic agitation at 47°C. Reactions were controlled by HPLC through disappearance of the starting material, which were complete after 24-48 hours. Solvents to dryness were evaporated and coevaporated twice with DCM. The residues obtained [Ac-AA₁-AA₂-AA₃-AA₄-NH-(CH₂)₁₅-CH₃ with completely protected side chains] were resuspended in 25mL of a mixture of TFA-DCM-anisole (49:49:2) and were left to react during 30min at room temperature. 250mL of cold diethyl ether were added, solvents were evaporated under reduced pressure and two additional coevaporations were done with ether. Residues were dissolved in a mixture of 50% MeCN in H₂O and lyophilised.

HPLC analysis of the peptides obtained in gradients of MeCN (+0.07% TFA) in H₂O (+0.1% TFA) showed a purity higher than 70% in all cases. The identity of the peptides obtained was confirmed by ES-MS.

EXAMPLE 8

Collagenase inhibition trial.

Peptides were resuspended in water in the presence of 0.5% DMSO. The trial was carried out in black microplates having 96 well and the EnzChek® Gelatinase/Collagenase Assay Kit (Molecular Probes) was used. To that end, peptides were preincubated at 2mg/mL during 1hour with 0.1unit/mL of type IV collagenase at room temperature with moderate agitation, after said time fluorescein conjugated substrate (DQ™ Gelatin) was added to a final concentration of 25µg/mL and reactions were incubated for 2 hours at room temperature with agitation and protected from light. The substrate, whose fluorescence is inhibited, is directed into collagenase releasing fluorescent fragments, monitored by fluorescence with a FLUOstar galaxy reader (BMG LabTechnologies), using 485nm filters for excitation and 520nm for emission.

Table 2 details those peptides whose collagenase inhibition values are higher than

- 43 -

25%. Inhibition values were normalized with respect to medium inhibition basal values.

Table 2. Collagenase activity inhibition percentage

	Compound	Inhibition %
5	Ac-L-Arg-L-His-L-Arg-L-Cit-OH	60.5
	Ac-L-Arg-L-His-L-His-L-Cit-OH	60.6
	Ac-L-Arg-L-Asn-L-Arg-L-Cit-OH	58.9
	Ac-L-Arg-L-His-L-His-L-Cit-CONH-(CH ₂) ₁₅ -CH ₃	55.4
	Palm-L-Arg-L-His-L-His-L-Cit-NH ₂	36.7
10	Palm-L-Arg-L-Asn-L-Arg-L-Cit-NH ₂	31.3
	H-L-Arg-L-Asn-L-His-L-Cit-OH	26.8
	Ac-L-Arg-L-Asn-L-His-L-Cit-CONH-(CH ₂) ₁₅ -CH ₃	26.8

EXAMPLE 9

15 *MMP-1, MMP-2, MMP-3 and MMP-9 inhibition.*

Human MMPs were reconstructed in 50mM Tris-HCl, 150mM NaCl, 5mM CaCl₂, 0.2mM NaN₃ at a 7.6 pH and activated by reaction with 10mM of 4-aminophenylmercuric acetate (dissolved in 0.01M NaOH) at a 10:1 to 37°C ratio for 4-6h. Activated proteases (0.35µg/mL) were preincubated with peptides to a final 0.5mM concentration in a 96-well black microplate for 1 hour at room temperature. After preincubation, 25µg/mL substrate (DQTM Gelatin) were added to the wells and samples were incubated for 16 hours at room temperature and protected against the light. Released fluorescence by marked gelatin digestion was measured with an automated multiplate fluorescence reader, exciting at 485nm and reading at 520nm.

The results were corrected from basal fluorescent value in MMP and product absence and normalized with respect to control fluorescence. Table 3 details the best inhibition values for peptides.

Table 3. Human MMPs inhibited by the peptides of the invention

	MMP-1	MMP-2	MMP-3	MMP-9
Ac-L-Arg-L-His-L-His-L-Cit-OH	34.7	101.4	39.1	63.1
Ac-L-Arg-L-Asn-L-Arg-L-Cit-OH	-37.0	78.7	21.9	69.5

- 44 -

Ac-L-Arg-L-Asn-L-Arg-L-Cit-OH	-25.2	95.3	88.1	89.6
-------------------------------	-------	------	------	------

EXAMPLE 10

Cosmetic composition containing Palm-L-Arg-L-Asn-L-His-L-Cit-NH₂ preparation.

5

The following formulation was prepared as described in the present invention:

INGREDIENT (INCI nomenclature)		WEIGHT %
A	MINERAL OIL	8.0
10 A	STEARIC ACID	2.4
A	CETEARYL ALCOHOL	1.6
A	BEESWAX	0.8
B	GLYCERINE	2.4
B	AQUA (WATER)	63.4
15 C	CARBOMER	0.3
C	TRIETHANOLAMINE	0.9
D	AQUA (WATER)	15.0
D	<i>Palm-L-Arg-L-Asn-L-His-L-Cit-NH₂</i> (0.01%)	5.0
D	LECITHIN	0.4

20

Phase A components were weighed in a big enough reactor and the mixture was heated at 80°C so as to melt the waxes. Phase B components were weighed in an adequate recipient for the whole content and heated at 70°C. Phase A was slowly added to phase B under intense agitation, and phase C was later added to that mixture under agitation. After the addition, it was left to cool with soft agitation and when the mixture was at room temperature an aqueous solution of *Palm-L-Arg-L-Asn-L-His-L-Cit-NH₂* and lecithin was added, homogenized and pH was corrected with triethanolamine.

25

30 The pH obtained from the cream was 6-7 and viscosity 10.000-15.000cps (6/50).

EXAMPLE 11

Liposomes containing Ac-L-Arg-L-His-L-His-L-Cit-OH preparation.

- 45 -

Dipalmitoylphosphatidylcholine (DPPC) was weighed and dissolved in chloroform. The solvent was vacuum evaporated until obtaining a fine phospholipid layer and said layer was hydrated by treatment at 55°C with a peptide aqueous solution to the desired concentration (containing Phenonip®), and MLV liposomes were obtained. ULV liposomes were obtained by submerging MLV liposomes in an ultrasound bath at 55°C during 2-minute 8 cycles in 5 minutes intervals. ULV liposomes size was reduced by passing them through a high pressure extrusion system.

INGREDIENT (INCI nomenclature)	WEIGHT %
PHOSPHATIDYLCHOLINE	4.0
<i>Ac-L-Arg-L-His-L-His-L-Cit-OH</i>	0.2
PHENOXYETHANOL, METHYLPARABEN, ETHYLPARABEN, BUTYLPARABEN, PROPYLPARABEN, ISOBUTYLPARABEN	0.50
AQUA (WATER)	c.s.p. 100

EXAMPLE 12

Facial cream composition containing H-L-Arg-L-His-L-Arg-L-Cit-NH₂.

Preparation

- Mix Phase A components and heat at 70°C.
- Mix Phase B components and heat at 70°C.
- Add Phase C on Phase B agitating with homogenizer (Silverson) during 5 minutes.
- Add Phase A little by little into the mixture of phases B and C with homogenizer keeping homogenization for 5 minutes.
- Initiate cooling up to 30-35°C with soft agitation. Add phase D at 50°C. Keep agitation. Add previously solubilised Phases E and F at 35-38°C.

INGREDIENT (INCI Nomenclature)	WEIGHT %
A BUTYROSPERMUM PARKII	3.5-4.5
A CETEARYL ETHYLHEXANOATE	3-5
A GLYCERYL STEARATE S.E.	1.5-2.5
A SQUALANE	0.5-1
A PEG-100 STEARATE	1
A POLYSORBATE 60	0.30

- 46 -

	A	CETYL PALMITATE	1.5-2.5
	A	DIMETHICONE	2.5-3.5
	A	CETEARYL ALCOHOL	1.5-2.5
	A	PALMITIC ACID	0.5
5	B	AQUA (WATER)	2
	B	GLYCERIN	1.5-2.5
	B	BUTYLENE GLYCOL	1-3
	B	MANNITOL	0.5-1.5
	B	HYDROGENATED LECITHIN	0.5-1.5
10	B	PROPYLENE GLYCOL	0.5-1.5
	C	CARBOMER	0.4
	C	ETHYLHEXYL PALMITATE	1.5-2.5
	D	TROMETHAMINE	0.4
	D	AQUA (WATER)	1
15	E	PRESERVATIVES	q.s.
	F	<i>H-L-Arg-L-His-L-Arg-L-Cit-NH₂</i>	0.10
	F	AQUA (WATER)	c.s.p.100

EXAMPLE 13

20 *Liposome gel composition preparation containing Ac-L-Arg-L-His-L-His-L-Cit-OH.*

The liposomes of example 11 were spread in water with preservatives (EDTA, imidazolidinyl urea and Phenonip®) under soft agitation. Hispagel® 200 [INCI: Aqua, glycerin, glyceryl polyacrylate] was added and softly agitated until a homogenous mixture was obtained.

	INGREDIENT (INCI nomenclature)	WEIGHT %
	LIPOSOMES CONTAINING <i>Ac-L-Arg-L-His-L-His-L-Cit-OH</i> (1%)	10.00
	DISODIUM EDTA	0.15
30	IMIDAZOLIDINYL UREA	0.10
	PHENOXYETHANOL, METHYLPARABEN, ETHYLPARABEN, BUTYLPARABEN, PROPYLPARABEN, ISOBUTYLPARABEN	0.50
	AQUA (WATER)	29.25
	AQUA (WATER), GLYCERIN, GLYCERYL POLYACRYLATE	60.00

35

EXAMPLE 14

Mixed micelles composition containing Ac-L-Arg-L-His-L-His-L-Cit-OH preparation.

	INGREDIENT (INCI nomenclature)	WEIGHT %
5	A AQUA (WATER)	c.s.p.100
	A PHENOXYETHANOL	0.5
	A CAPRILYL GLYCOL	0.5
	A POTASIUM SORBATE	0.3
	B AQUA (WATER), PSEUDOALTEROMONAS FERMENT EXTRACT	7.5
10	B Ac-L-Arg-L-His-L-His-L-Cit-OH	0.025
	B AQUA (WATER), PENTAPEPTIDE-18	20
	B LECITHIN	4.0
	C XANTHAN GUM	0.4
	D AQUA (WATER), CAPRILYL/CAPRYL GLUCOSIDE	30

15

In the right container for the complete sample, phase A ingredients were weighed and lightly heated at 30°C to help dissolve part of the preservatives. After that, phase B components were added and homogenized under moderate agitation.

20 Then, phase C was added under continuous agitation, after which phase D was added with slow agitation so as not to produce foam.

pH was adjusted to 5.5-6.5.

25 **EXAMPLE 15**

Composition containing Ac-L-Arg-L-His-L-His-L-Cit-OH for the treatment and/or prevention of stretch marks

30 In the right container for the complete sample, phase A components were weighed and lightly heated at 30°C to help dissolve part of the preservatives. After that, phase B components were mixed and added onto phase A and the composition was homogenized under moderate agitation.

INGREDIENT (INCI nomenclature)	WEIGHT %
--------------------------------	----------

- 48 -

	A	GLYCERIN	50
	A	PHENOXYETHANOL	0.50
	A	CAPRILYL GLYCOL	0.50
	B	AQUA (WATER), PSEUDOALTEROMONAS FERMENT EXTRACT	7.50
5	B	<i>Ac-L-Arg-L-His-L-His-L-Cit-OH</i>	0.025
	B	AQUA (WATER)	c.s.p.100

EXAMPLE 16*Hair lotion composition containing Ac-L-Arg-L-Asn-L-His-L-Cit-NH₂.*

10

INGREDIENT (INCI nomenclature)		WEIGHT %
A	ALCOHOL DENAT.	50-60
A	PANTHENOL	0.05-0.15
A	ZINC RICINOLEATE	0.05-0.10
A	FRAGRANCE	0.02
B	AQUA (WATER)	c.s.p.100
B	<i>Ac-L-Arg-L-Asn-L-His-L-Cit-NH₂</i>	0.01

Preparation:

20

- Mix Phase A components.
- Mix Phase B components.
- Slowly add Phase B onto Phase A with agitation until complete homogenization.


EXAMPLE 1725 *Collutary composition containing Ac-L-Arg-L-His-L-His-L-Cit-OH.*

Components were mixed until complete homogenization.

INGREDIENT (INCI nomenclature)		WEIGHT %
	<i>Ac-L-Arg-L-His-L-His-L-Cit-OH</i>	0.10
30	SODIUM SACCARIN	0.01-0.03
	SORBITOL	4-6
	PROPYLENE GLYCOL	8-12
	PEG-60 HYDROGENATED CASTOR OIL	1-3
	AQUA (WATER)	c.s.p.100

CLAIMS

1. A peptide of general formula (I)



(I)

- 5 stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof, characterized in that:

AA₁ is -Arg-;

AA₂ is selected from the group consisting of -His- and -Asn-;

AA₃ is selected from the group consisting of -His- and -Arg-;

10 AA₄ is -Cit-;

R₁ is selected from the group consisting of H, substituted or non-substituted non-cyclic aliphatic group, substituted or non-substituted alicyclyl, substituted or non-substituted heterocyclyl, substituted or non-substituted heteroarylalkyl, substituted or non-substituted aryl, substituted or non-substituted aralkyl, and

15 R₅-CO-; and

R₂ is selected from the group consisting of -NR₃R₄, -OR₃ and -SR₃; wherein R₃ and R₄ are independently selected from the group consisting of H, substituted or non-substituted non-cyclic aliphatic group, substituted or non-substituted alicyclyl, substituted or non-substituted heterocyclyl, substituted or non-

20 substituted heteroarylalkyl, substituted or non-substituted aryl and substituted or non-substituted aralkyl

wherein R₅ is selected from the group consisting of H, substituted or non-substituted non-cyclic aliphatic group, substituted or non-substituted alicyclyl, substituted or non-substituted aryl, substituted or non-substituted aralkyl, substituted or non-substituted heterocyclyl and substituted or non-substituted heteroarylalkyl.

2. Peptide according to claim 1, **characterized in that** R₁ is H, or a R₅-CO- group wherein R₅ is selected from the group consisting of substituted or non substituted C₁-C₂₄ alkyl, substituted or non substituted C₂-C₂₄ alkenyl, substituted or non substituted C₂-C₂₄ alkynyl, substituted or non substituted C₃-C₂₄ cycloalkyl, substituted or non substituted C₅-C₂₄ cycloalkenyl, substituted or non substituted C₅-C₂₄ cycloalkynyl, substituted or non substituted C₆-C₃₀ aryl, substituted or non substituted C₇-C₂₄ aralkyl, substituted or non substituted 3 to 10 member heterocyclyl and substituted or non substituted heteroarylalkyl with from 2 to 24 carbon atoms and from
- 35 1 to 3 atoms different from carbon and an alkyl chain of 1 to 6 carbon atoms.

3. Peptide according to claim 2, **characterized in that** R_1 is selected from the group consisting of H, acetyl, *tert*-butanoyl, hexanoyl, 2-methylhexanoyl, cyclohexanecarboxyl, octanoyl, decanoyl, lauroyl, myristoyl, palmitoyl, stearyl, oleoyl and linoleoyl.
4. Peptide according to any of the preceding claims, **characterized in that** R_2 is $-NR_3R_4$ or $-OR_3$, wherein R_3 and R_4 are independently selected from the group consisting of H, substituted or non substituted C_1 - C_{24} alkyl, substituted or non substituted C_2 - C_{24} alkenyl, substituted or non substituted C_2 - C_{24} alkynyl, substituted or non substituted C_3 - C_{24} cycloalkyl, substituted or non substituted C_5 - C_{24} cycloalkenyl, substituted or non substituted C_5 - C_{24} cycloalkynyl, substituted or non substituted C_6 - C_{30} aryl, substituted or non substituted C_7 - C_{24} aralkyl, substituted or non substituted 3 to 10 member heterocyclyl and substituted or non substituted heteroarylalkyl with from 2 to 24 carbon atoms and from 1 to 3 atoms different from carbon and an alkyl chain of 1 to 6 carbon atoms.
5. Peptide according to claim 4, **characterized in that** R_3 and R_4 are selected from the group consisting of H, methyl, ethyl, hexyl, dodecyl and hexadecyl.
6. Peptide according to any of the preceding claims, **characterized in that** AA_2 is -His- and AA_3 is -His-.
7. Peptide according to any of the claims 1 to 5, **characterized in that** AA_2 is -Asn- and AA_3 is -Arg-.
8. Peptide according to any of the claims 1 to 6, **characterized in that** R_1 is H, acetyl, lauroyl, myristoyl or palmitoyl, AA_1 is -L-Arg-, AA_2 is -L-His-, AA_3 is -L-His-, AA_4 is -L-Cit, and R_2 is $-NR_3R_4$ or $-OR_3$ wherein R_3 and R_4 are independently selected from H, methyl, ethyl, hexyl, dodecyl and hexadecyl groups.
9. Peptide according to any of the claims 1 to 5 and 7, **characterized in that** R_1 is H, acetyl, lauroyl, myristoyl or palmitoyl, AA_1 is -L-Arg-, AA_2 is -L-Asn-, AA_3 is -L-Arg-, AA_4 is -L-Cit, and R_2 is $-NR_3R_4$ or $-OR_3$ wherein R_3 and R_4 are independently selected from H, methyl, ethyl, hexyl, dodecyl and hexadecyl groups.

10. Peptide according to any of the claims 1 to 9, **characterized in that** R₁ is selected from the group consisting of H, acetyl and palmitoyl and R₂ is selected from the group consisting of -OH and -NH₂.

5

11. Process for obtaining a peptide of general formula (I), stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof as defined in any of the claims 1 to 10, **characterized in that** said process is carried out on solid phase or in solution.

10

12. Process according to claim 11, **characterized in that** the free amino groups are protected by Boc or Fmoc, the free carboxyl groups are protected with tBu, All or Trt, the arginine side chain is protected with Pmc, Mtr, Tos or Pbf, the histidine side chain is protected with Trt, Dnp, Boc, Mtt or Tos and the asparagine side chain is free or

15

13. Cosmetic or pharmaceutical composition comprising a cosmetically or pharmaceutically effective amount of at least one peptide of general formula (I), stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof, as defined in any of the claims 1 to 10, and at least one cosmetically or pharmaceutically acceptable excipient or adjuvant.

20

14. Cosmetic or pharmaceutical composition, according to claim 13, **characterized in that** the peptide of general formula (I) is at a concentration comprised between 0.000001% and 20% by weight, with regard to total weight of the composition.

25

15. Cosmetic or pharmaceutical composition, according to claim 14, **characterized in that** said concentration is comprised between 0.0001% and 5% by weight, with regard to total weight of the composition.

30

16. Cosmetic or pharmaceutical composition, according to any of the claims 13 to 15, **characterized in that** the peptide of general formula (I), stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof are incorporated into cosmetic or pharmaceutical delivery system and/or sustained release system, selected from group consisting of liposomes, mixed liposomes, millicapsules,

35

- 52 -

microcapsules, nanocapsules, sponges, cyclodextrins, vesicles, micelles, surfactant mixed micelles, phospholipid-surfactant mixed micelles, millispheres, microspheres, nanospheres, lipospheres, microemulsions, nanoemulsions, milliparticles, microparticles, nanoparticles and solid lipidic nanoparticles.

5

17. Cosmetic or pharmaceutical composition, according to any of the claims 13 to 16, **characterized in that** the peptide of general formula (I), stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof is absorbed on an solid organic polymer or cosmetically or pharmaceutically acceptable solid support, selected from the group consisting of talcum powder, bentonite, silica, starch and maltodextrin.

10

18. Cosmetic or pharmaceutical composition, according to any of the claims 13 to 17, **characterized in that** said composition is selected from the group consisting of creams, multiple emulsions, anhydrous compositions, aqueous dispersions, oils, milks, balsams, foams, lotions, gels, cream gels, hydroalcoholic solutions, hydroglycolic solutions, liniments, saline solutions, soaps, shampoos, conditioners, sera, unguents, mousses, ointments, powders, bars, pencils, sprays, aerosols, capsules, gelatin capsules, tablets, sugar coated tablets, powders, granulated forms, chewing gum, solutions, suspensions, emulsions, syrups, polysaccharides films, jellies and gelatins.

15

20

19. Cosmetic or pharmaceutical composition, according to any of the claims 13 to 17, **characterized in that** said composition is a product selected from the group consisting of under eye removers, makeup foundations, makeup remover lotions, milk make up remover, eye shadows, lipsticks, lip glosses and powders.

25

20. Cosmetic or pharmaceutical composition, according to any of the claims 13 to 17, **characterized in that** the peptide of general formula (I), stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof are incorporated into a fabric, a non-woven fabric or a medical device.

30

21. Cosmetic or pharmaceutical composition, according to claim 20, **characterized in that** said fabric, non-woven fabric or medical device is selected from the group consisting of bandages, gauzes, T-shirts, socks, pantyhose, underwear, girdles, gloves, diapers, sanitary napkins, dressings, bedcovers, wipes, hydrogels, adhesive

35

- 53 -

patches, non-adhesive patches, microelectric patches and face masks.

22. Cosmetic or pharmaceutical composition, according to any of the claims 13 to 20,
characterized in that said composition additionally comprises a cosmetically or
5 pharmaceutically effective amount of at least one selected active agent consisting of
agents inhibiting matrix metalloelastases, agents stimulating or inhibiting melanine
synthesis, whitening or depigmenting agents, propigmenting agents, self-tanning
agents, anti-age agents, NO-synthase inhibiting agents, antioxidant agents, free
radicals scavengers and/or anti-atmospheric pollution agents, anti-glycation agents,
10 emulsifying agent, emollients, organic solvents, liquid propellants, skin conditioners
such as humectants, substances retaining moisture, alphahydroxyacids,
betahydroxyacids, moisturizers, epidermic hydrolytic enzymes, vitamins, pigments or
colorants, dyes, gelling polymers, thickeners, surfactants, softeners, anti-wrinkle
agents, agents capable of reducing or treating bags under the eyes, exfoliating agents,
15 antimicrobial agents, fungicide agents, fungistatic agents, bactericide agents,
bacteriostatic agents, agents stimulating the synthesis of dermal or epidermal
macromolecules and/or agents capable of preventing or inhibiting their degradation,
agents stimulating collagen synthesis, agents stimulating elastin synthesis, agents
stimulating decorine synthesis, agents stimulating laminin synthesis, agents stimulating
20 defensin synthesis, agents stimulating chaperone synthesis, agents stimulating
aquaporin synthesis, agents stimulating hyaluronic acid synthesis, agents stimulating
the synthesis of lipids and components of the stratum corneum, agents stimulating the
synthesis of ceramides, agents inhibiting collagen degradation, agents inhibiting elastin
degradation, agents stimulating fibroblast proliferation, agents stimulating keratinocyte
25 proliferation, agents stimulating adipocyte proliferation, agents stimulating melanocyte
proliferation, agents stimulating keratinocyte differentiation, agents stimulating
adipocyte differentiation, agents inhibiting acetylcholinesterase, dermorelaxing agents,
agents stimulating glycosaminoglycan synthesis, DNA repairing agents, DNA
protecting agents, anti-itching agents, agents for the treatment and/or care of sensitive
30 skin, firming agents, anti-stretch mark agents, astringent agents, agents regulating
sebum production, agents stimulating lipolysis, anticellulitic agents, agents stimulating
healing, coadjuvant healing agents, agents stimulating reepithelizing, coadjuvant
reepithelizing agents, cytokine growth factors, calming agents, anti-inflammatory
agents, agents acting on capillary circulation and/or microcirculation, agents
35 stimulating angiogenesis, agents inhibiting vascular permeability, agents acting on cell

- 54 -

metabolism, agents intended to improve the dermal-epidermal junction, agents inducing hair growth, agents inhibiting or delaying hair growth, preservatives, perfumes, chelating agent, plant extracts, essential oils, marine extracts, agents coming from a bio-fermentation process, mineral salts, cell extracts and sunscreens (organic or mineral photoprotecting agents that are active against ultraviolet A and/or B rays), or mixtures thereof.

23. Cosmetic or pharmaceutical composition, according to claim 22, **characterized in that** said active agent has a synthetic origin or is a plant extract or comes from a biofermentation process.

24. Cosmetic or pharmaceutical composition, according to any of the claims 22 to 23, **characterized in that** said active agent is an agent with healing and/or reepithelizing activity or a natural extract or essential oil with intrinsic healing and/or reepithelizing activity.

25. Cosmetic or pharmaceutical composition, according to claim 24, **characterized in that** agent with healing and/or reepithelizing activity is *Pseudoalteromonas* ferment extract.

26. Cosmetic or pharmaceutical composition, according to any of the claims 22 to 23, **characterized in that** said active agent is an agent with anti-wrinkle and/or anti-aging activity or a natural extract or essential oil with intrinsic anti-wrinkle and/or anti-aging activity.

27. Cosmetic or pharmaceutical composition, according to claim 26, **characterized in that** said active agent with anti-wrinkle and/or anti-aging activity is Pentapeptide-18 and/or Acetyl hexapeptide-8.

28. Cosmetic or pharmaceutical composition, according to any of the claims 22 to 23, **characterized in that** said composition comprises at least one agent selected from the group consisting of an anti-wrinkle and/or anti-aging activity agent, or a natural extract or essential oil of anti-wrinkle and/or anti-aging intrinsic activity and at least one agent selected from the group an agent with healing and/or reepithelizing activity or a natural extract or essential oil with intrinsic healing and/or reepithelizing activity.

- 55 -

29. Cosmetic or pharmaceutical composition, according to claim 28, **characterized in that** said anti-wrinkle and/or anti-aging intrinsic activity agent is Pentapeptide-18 and/or Acetyl hexapeptide-8 and the healing and/or reepithelizing activity agent is *Pseudoalteromonas* ferment extract.

30. Cosmetic or pharmaceutical composition, according to claims 16 and 29, **characterized in that** said cosmetically or pharmaceutically acceptable delivery system or sustained release system is a phospholipid-surfactant mixed micelle.

31. Cosmetic or pharmaceutical composition, according to any of the claims 22 to 23, **characterized in that** said active agent is an agent with refirming, rethickening and/or restructuring activity or a natural extract or essential oil with intrinsic refirming, rethickening and/or restructuring activity.

32. Cosmetic or pharmaceutical composition, according to any of the claims 22 to 23, **characterized in that** said active agent is an anti-inflammatory and/or analgesic agent or a natural extract or essential oil with intrinsic anti-inflammatory and/or analgesic activity.

33. Cosmetic or pharmaceutical composition, according to any of the claims 22 to 23, **characterized in that** said active agent is a matrix metalloprotease inhibitor agent or a natural extract or essential oil with intrinsic activity inhibiting matrix metalloproteases.

34. Use of a peptide of general formula (I), stereoisomers thereof, mixtures thereof or the cosmetically or pharmaceutically acceptable salts thereof, according to any of the claims 1 to 10, in the preparation of a cosmetic or pharmaceutical composition for the treatment and/or care of skin, mucosae and/or scalp.

35. Use, according to claim 34, wherein said treatment and/or care consists of inhibiting at least one skin, mucosae and/or scalp matrix metalloprotease.

36. Use, according to claim 35, wherein said matrix metalloproteases are selected from the group consisting of human MMP-1, MMP-2, MMP-3 and MMP-9.

- 56 -

37. Use, according to any of the claims 35 to 36, wherein said treatment and/or care is provided by topical or transdermal administration of the cosmetic or pharmaceutical composition.

5 38. Use, according to claim 37, wherein said topical or transdermal administration by performed by means of iontophoresis, sonophoresis, electroporation, mechanic pressure, osmotic pressure gradient, occlusive treatment, microinjections, pressure needle-free injections by means of pressure,, microelectric patches or any combination thereof.

10 39. Use, according to any of the claims 35 to 36, wherein said treatment and/or care is provided by means of oral administration of the cosmetic or pharmaceutical composition.

15 40. Use, according to any of the claims 34 to 39, for the treatment and/or care of skin, mucosae and/or scalp conditions, disorders and/or pathologies caused by matrix metalloproteases overexpression or by an increased matrix metalloproteases activity.

20 41. Use, according to claim 40, wherein said conditions, disorders and/or pathologies are selected from the group consisting of acne, rosacea, psoriasis, dermatitis, atopic dermatitis, eczema, sensitive skins, gingivitis, periodontitis, skin cancer, tumor invasions, aging skin, photoaging skin, wrinkles, expression wrinkles, stretch marks, keloids, hypertrophic scars, cellulitis, orange peel skin, tumour metastasis, ulcers, diabetic ulcers, telangiectasia, cuperosis, varicose veins, eye dark circles, bags under
25 the eyes, alopecia and hair loss.

42. Use, according to any of the claims 34 to 41, wherein said skin treatment is performed for reducing, delaying and/or preventing aging and/or photoaging signs.

30 43. Use, according to any of the claims 34 to 41, for the treatment of oral cavity mucosae or for oral hygiene.

44. Use, according to any of the claims 34 to 41, for the treatment of hair and/or scalp and/or hair hygiene.

35

- 57 -

45. Use, according to any of the claims 32 to 39, for the treatment of the body skin or body hygiene.