




FIG. 1A






FIG. 1B



EPSP, excitatory postsynaptic potential; LTD, long-term depression; ns, not significant; STP, short-term potentiation.


* $p < 0.05$

FIG. 1C



ANOVA, analysis of variance; fEPSP, field excitatory postsynaptic potentials; HFS, high-frequency stimulation; ns, not significant versus vehicle.

FIG. 2



* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ versus vehicle.
ANOVA, analysis of variance; fEPSP, field excitatory postsynaptic potentials; HFS, high-frequency stimulation.

FIG. 3




FIG. 4B




FIG. 4A






FIG. 5




LS, least-squares; PANSS NSFS, Positive and Negative Syndrome Scale – Negative Symptom Factor Score; qd, once daily; SE, standard error.

FIG. 6




* $p < 0.05$.
BACS, Brief Assessment of Cognition in Schizophrenia; LS, least-squares; qd, once daily; SE, standard error.

FIG. 7



* $p < 0.05$.
LS, least-squares; ns, not significant; qd, once daily; SCoRS, Schizophrenia Cognition Rating Scale;
SE, standard error.

FIG. 8



^aAssessments on day 7 and day 8 may be performed 1 day before or after (i.e., day 6 or day 8, and day 7 or day 9, respectively)

FIG. 9




FIG. 10




FIG. 11B




FIG. 11A




FIG. 12A

FIG. 12B

15/19




FIG. 13B




FIG. 13A




FIG. 14B




FIG. 14A






FIG. 15



D=Day; No.=number; OS=onsite; QD=once daily; V=virtual; W=Week

FIG. 16A



D=Day; F/U=follow-up; No.=number; OS=onsite; QD=once daily; V=virtual; W=Week
^a Early Termination Visit

FIG. 16B