

METHOD FOR TRANSFERRING ONE OR MORE ACTIVE INGREDIENTS BETWEEN DIFFERENT PHASE CARRIERS

BACKGROUND OF THE INVENTION

5 1. Field of the Invention

The present invention relates to a method for controlling administration, particularly to a method for transferring one or more active ingredients between different phase carriers.

10 2. Description of Related Art

US patent No. 5,689,533 discloses "Ophthalmic Pharmaceutical Composition", which provides a method of administering a drug to an eye including the steps of: (a) admixing a pharmaceutically acceptable hydrocarbonaceous
15 semi-solid or oil which contains the drug with water at a temperature above the melting point of the semi-solid or oil; and (b) nebulizing the admixture to form liquid drops; and (c) applying the liquid drops to the eye.

20 SUMMARY OF THE PRESENT INVENTION

It is an object of the present invention to provide a method, which enables one or more active ingredients to be conveniently transferred between different phase carriers, so that the active ingredient can be stored and used optionally.

It is another object of the present invention to provide a method, which enables one or more active ingredients to be transferred between different phase carriers. For example, the active ingredient unstable in water can be stored in an oil phase carrier (oil phase) or non-aqueous liquid phase carrier to keep the active ingredient stable. When desired to use, then the active ingredient can be transferred to water or water solution (aqueous phase carrier), and then be transferred to liquid fine drops (vapor phase).

It is the other object of the present invention to provide a method, which enables one or more active ingredients stored in a solid phase carrier (solid phase) to be transferred to an aqueous phase carrier, and then be transferred to liquid fine drops (vapor phase).

One aspect of this invention relates to a method for transferring one or more active ingredients between different phase carriers comprising the steps of: (a) providing a solid, semi-solid, or non-aqueous liquid drug which contains at least one active ingredient; (b) admixing said drug with water or water solution to form an admixture; and (c) nebulizing said admixture to form liquid fine drops containing said active ingredients.

The aforesaid solid drug can be a solid active ingredient (or active ingredients) without a carrier, or an active ingredient (or active ingredients) mixed with a solid

carrier. The solid drug can be in the form of powder, granule, tablet or capsule, wherein the powder is preferred. In order to control the amount of the active ingredient in the powder, a solid powdered diluent can be added. The solid powdered
5 diluent can be selected from the group consisting of dicalcium phosphate, calcium sulfate, lactose, kaolin, sodium chloride, dry starch, and powdered sugar.

The aforesaid semi-solid drug can be an active ingredient (or active ingredients) mixed with a grease, fat or
10 wax carrier.

The aforesaid non-aqueous liquid drug can be an active ingredient (or active ingredients) stored in a non-aqueous liquid carrier such as oil phase carrier, alcohol carrier, or other organic carriers. The oil phase carrier is not limited here, but
15 must enable the active ingredient to remain stable therein. The oil phase carrier can be animal oil (for example: lanolin), plant oil (for example: olive oil, corn oil, soybean oil), or mineral oil.

There is no special limitation on the water solution of
20 step (b). Example of the water solution is an aqueous solution containing active ingredients.

The drug of the present invention can be contained one or more active ingredients. The active ingredient is selected
* from the group consisting of ophthalmic agents, dermal agents
25 (including cosmetic agents), ENT (ear, nose and throat) agents,

and respiratory agents.

The ophthalmic agents can be selected from the group consisting of vitamins, steroids (e.g. betamethasone, or dexamethasone), cephalosporins (e.g. cephadrine, or cephalexin), penicillins (e.g. amoxicillin, or ampicillin),
5 tetracyclines (e.g. chlorotetracycline, or demeclocycline), macrolides (e.g. erythromycin base, erythromycin propionate), aminoglycosides (e.g. gentamicin, tobramycin or neomycin), sulfonamides (e.g. sulfamethomidin, sulfamethoxypyridazine),
10 polypeptides (e.g. bacitracin) and naphazoline hydrochloride.

The dermal agents can be selected from the group consisting of vitamins, steroids (e.g. betamethasone, or dexamethasone), cephalosporins (e.g. cephadrine, or cephalexin), penicillins (e.g. amoxicillin, or ampicillin),
15 tetracyclines (e.g. chlorotetracycline, or demeclocycline), macrolides (e.g. erythromycin base, erythromycin propionate), aminoglycosides (e.g. gentamicin, tobramycin or neomycin), sulfonamides (e.g. sulfamethomidin, sulfamethoxypyridazine), polypeptides (e.g. bacitracin) hydroquinone, piroxicam, and
20 indomethacin.

The ENT (ear, nose and throat) agents can be selected from the group consisting of vitamins, steroids (e.g. betamethasone, or dexamethasone), cephalosporins (e.g. cephadrine, or cephalexin), penicillins (e.g. amoxicillin, or
25 ampicillin), tetracyclines (e.g. chlorotetracycline, or

demeclocycline), macrolides (e.g. erythromycin base, erythromycin propionate), aminoglycosides (e.g. gentamicin, tobramycin or neomycin), sulfonamides (e.g. sulfamethomidin, sulfamethoxypyridazine), polypeptides (e.g. bacitracin),
5 naphazoline hydrochloride, and acetylcysteine.

The respiratory agents can be selected from the group consisting of vitamins, steroids (e.g. betamethasone, or dexamethasone), cephalosporins (e.g. cephadrine, or cephalexin), penicillins (e.g. amoxicillin, or ampicillin),
10 tetracyclines (e.g. chlorotetracycline, or demeclocycline), macrolides (e.g. erythromycin base, erythromycin propionate), aminoglycosides (e.g. gentamicin, tobramycin or neomycin), sulfonamides (e.g. sulfamethomidin, sulfamethoxypyridazine), polypeptides (e.g. bacitracin), ephedrine sulfate,
15 methylephedrine, acetylcysteine, isoproterenol sulfate, and aminophylline.

In order to transfer the active ingredient into water after the drug is mixed with water or water solution to form the admixture in the step (b), it may be necessary to stir, oscillate,
20 or heat the admixture.

The nebulizer is a well-known device and is easily obtained and operated by one skilled in the art. In the present invention, the nebulizer is used for providing energy to oscillate water and form a liquid fine drops escaping from the
25 surface of the admixture (active ingredient with water or water

solution). The temperature of the nebulized fine drops of liquid was about 25°C . The active ingredient of the admixture can be brought out with liquid fine drops to achieve the object of controlling drug delivery.

5

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The characteristic of the present invention is the transfer of active ingredients. The method provided in the present invention enables one or more active ingredients stored in a carrier in a certain phase (such as an oil phase carrier, a grease phase carrier, or a solid phase carrier) to transfer into an admixture (water or water solution phase). Then, the active ingredient can be brought out with liquid fine drops (vapor phase) from the admixture by means of ultrasonic vibration.

The present invention can be applied in many situations. For example, the active ingredient (e.g. cephalosporin, vitamin C, or penicillin, etc) unstable in water can be stored in an oil phase carrier, a grease phase carrier, or a solid phase carrier to form a non-aqueous drug to keep the active ingredient stable. When desired to use, then the drug can be mixed with water or water solution to form an admixture. The admixture was then placed in a nebulizer to form fine drops of liquid. The nebulized fine drops of liquid can then be easily and smoothly delivered to

the patient's body by gentle currents of air (e.g. provided by a fan installed in the nebulizer).

In general, the oil phase carrier or the grease phase carrier in the admixture is not brought out with liquid fine drops.

- 5 Only the active ingredient is brought out with liquid fine drops. The method of the present invention enables an active ingredient (or active ingredients) to be conveniently transferred between different phase carriers.

- The present invention is demonstrated in more detail
10 with reference to the following examples, which are only illustrative and are not intended to limit the scope of the present invention.

Example 1

- 15 (The preparation of a drug containing cephalexin monohydrate):

10mg cephalexin monohydrate (an active ingredient) is added into a 10ml flask, and then mineral oil is added to obtain a 10ml drug containing 0.1%(W/V) cephalexin monohydrate.

- 20 Examples 2 to 5

(The preparation of drugs containing cephalexin monohydrate):

Repeat the steps in example 1, but 50mg, 100mg, 250mg, and 500mg cephalexin monohydrate are added instead to obtain pharmaceuticals containing 0.5%(W/V), 1.0%(W/V),

2.5%(W/V), and 5.0%(W/V) cephalixin monohydrate respectively.

Examples 6 and 7

5 (The preparation of drugs containing cephalixin sodium):

Repeat the steps in example 1, but 100mg and 500mg cephalixin sodium are respectively added instead to obtain the pharmaceuticals containing 1.0%(W/V) and 5.0%(W/V) cephalixin sodium.

10

Test Example 1

(nebulization and HPLC analysis):

200 μ l(V1) drug obtained in example 1 is added into a cup containing 5ml(V2) water to form an admixture, and then
15 the cup is placed in a nebulizer (Model 8050, Formosa-CJ Business Corp., Taipei, Taiwan) which nebulized the admixture in the cup into liquid fine drops. After nebulized for a period of time, there are some oil phase liquid still remained in the cup, and the liquid fine drops is collected and cooled
20 down to become liquid which is a water solution containing the active ingredient. The concentration of the active ingredient in the collected liquid (liquid fine drops) is then determined by HPLC analysis, and the results are listed in table 1, wherein the term "theoretical concentration" is the concentration of the

active ingredient in the admixture before nebulization, and the term " liquid fine drops concentration " is the concentration of the active ingredient in the collected liquid (liquid fine drops).

5 Test Examples 2 to 7

(nebulization and HPLC analysis):

Repeat the steps in test example 1, but the drugs obtained in examples 2 to 7 are respectively added instead, and the results of HPLC analysis are listed in table 1.

10

Table 1:

test example	active ingredient	theoretical concentration (mg/ml)	liquid fine drops concentration (mg/ml)	recovery ratio(%)
1	cephalexin monohydrate	0.04	0.0302	75.38
2	cephalexin monohydrate	0.20	0.1635	81.74
3	cephalexin monohydrate	0.40	0.3889	97.23
4	cephalexin monohydrate	1.00	0.9614	96.14
5	cephalexin monohydrate	2.00	1.3847	68.74
6	cephalexin sodium	0.40	0.3659	91.49
7	cephalexin sodium	2.00	1.6590	82.95

theoretical concentration = (concentration of drug(%W/V)) ×
(amount of drug(=V1)) / (amount of water in the cup(=V2))
recovery ratio (%) = (liquid fine drops concentration) /
(theoretical concentration) × 100%

5

The active ingredient contained in the drug obtained in examples 1 to 7 are all stored in the oil phase carriers. The drugs obtained in examples 1 to 7 are respectively added into water to form 2-phase (oil/aqueous) admixtures. The admixtures can
10 then be conveniently nebulized with a nebulizer by means of ultrasonic vibration. According to results listed in table 1, the active ingredients stored in the drugs (oil phase) can be transferred to the admixtures (oil/aqueous phase), and can then be transferred to liquid fine drops (vapor phase) by nebulizing
15 the admixture with a nebulizer. Additionally, the high recovery ratio (for example in test example 3 the recovery ratio is 97.23%) in the method of the present invention indicate the active ingredients (cephalexin monohydrate and cephalexin sodium) can be easily and effectively transferred between different
20 phases.

Example 8

(The preparation of a 1-phase (aqueous) admixture containing

tobramycin):

15mg solid tobramycin is added into 5ml water to obtain an admixture containing 3 mg/ml tobramycin.

5 Example 9

(The preparation of a 2-phase (oil/aqueous) admixture containing tobramycin and cephalexin monohydrate):

5ml water solution containing 3 mg/ml tobramycin is added into a cup for nebulization, and then a 200 μ l mixture
10 consisting of 1%(W/V) cephalexin monohydrate in mineral oil is added to obtain a 2-phase (oil/aqueous) admixture containing 3mg/ml tobramycin and 0.4mg/ml cephalexin monohydrate.

15 Example 10

(The preparation of a 2-phase (oil/aqueous) admixture containing cephradine monohydrate and cephalexin monohydrate):

5mg cephradine monohydrate and 5mg cephalexin
20 monohydrate are added into a 10ml flask, and then mineral oil is added to the volume obtain a 10ml oil phase drug containing 0.5%(W/V) cephradine monohydrate and 0.5%(W/V) cephalexin monohydrate in mineral oil. Then 200 μ l(=V1)

above oil phase drug is added into a cup containing 5ml(=V2) water and stirred to form a 2-phase (oil/aqueous) admixture containing 0.2mg/ml cephradine monohydrate and 0.2mg/ml cephalixin.

5

Test Examples 8 to 10

(nebulization and HPLC analysis):

Repeat the steps in test example 1, but the admixtures obtained in examples 8 to 10 are respectively added instead,
10 and the results of HPLC analysis are listed in table 2.

Table 2:

test example	active ingredient	theoretical concentration (mg/ml)	Liquid fine drops concentration (mg/ml)	recovery ratio (%)
8	tobramycin	3	2.7668	92.23
9	tobramycin	3	2.8916	96.39
	cephalexin monohydrate	0.4	0.3323	83.07
10	cephradine monohydrate	0.2	0.1535	76.74
	cephalexin monohydrate	0.2	0.1948	97.40

In example 8, the active ingredient (tobramycin) stored
15 in the solid phase carrier transfers into the aqueous phase carrier

after adding water for preparing the admixture. The result of test example 8, as listed in table 2, indicates the drug obtained in example 8 can be nebulized to form liquid fine drops by the nebulizer. According to results listed in table 2, the collected liquid (collected from liquid fine drops) containing the active ingredient (tobramycin). Additionally, the high recovery of 92.23% indicates the active ingredient (tobramycin) stored in a solid phase carrier can be successfully transferred to vapor phase (liquid fine drops).

10 In example 9, another active ingredient (cephalexin monohydrate) stored in an oil phase carrier is added into the admixture of example 8 to form a 2-phase (oil/aqueous) admixture. The result of test example 9, as listed in table 2, indicates that the active ingredient (cephalexin monohydrate) stored in the oil phase carrier and the active ingredient (tobramycin) stored in the aqueous phase carrier can be mixed to obtain the 2-phase (oil/aqueous) admixture, and then be nebulized into liquid fine drops. The collected liquid (liquid fine drops) is water solution primarily containing the active ingredients (tobramycin and cephalexin monohydrate). Both recovery ratios over 80% indicate that the mixture of the active ingredients stored in the aqueous phase carrier and in the oil phase carrier can be transferred to vapor phase (liquid fine drops).

In example 10, the admixture is prepared by adding the drug (cephradine monohydrate and cephalixin monohydrate stored in the oil phase carrier) into water. The result of test example 10, as listed in table 2, indicates that the admixture can
5 form liquid fine drops by nebulization, and the collected liquid (liquid fine drops) is water solution primarily containing the active ingredients (cephradine monohydrate and cephalixin monohydrate), so that two active ingredients stored in oil phase carrier can be transferred to vapor phase (liquid fine drops).

10 From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention, and, without departing from the scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. Thus, other embodiments are also within
15 the claims.



What is Claimed is:

1. A method for transferring one or more active ingredients between different phase carriers comprising:
 - 5 (a) providing a solid, semi-solid, or non-aqueous liquid drug which contains at least one active ingredient;
 - (b) admixing said drug with water or water solution to form an admixture; and
 - (c) nebulizing said admixture to form liquid fine drops
- 10 containing said active ingredients.
2. The method of claim 1, wherein said solid drug is selected from the group consisting of powder, granule, tablet and capsule.
3. The method of claim 2, wherein said powder contains
- 15 at least one active ingredient with or without a solid carrier.
4. The method of claim 3, wherein said solid carrier is a powder diluent selected from the group consisting of dicalcium phosphate, calcium sulfate, lactose, kaolin, sodium chloride, dry starch, and powdered sugar.
- 20 5. The method of claim 1, wherein said semi-solid drug contains at least one active ingredient and a carrier selected from the group consisting of grease carrier, fat carrier and wax carrier.
6. The method of claim 1, wherein said non-aqueous
- 25 liquid drug contains at least one active ingredient and a carrier



selected from the group consisting of oil phase carrier, alcoholic carrier and organic solution carrier.

7. The method of claim 6, wherein said oil phased carrier is selected from the group consisting of lanolin, olive
5 oil, corn oil, soybean oil, and mineral oil.

8. The method of claim 1, wherein said drug contains two or more active ingredients.

9. The method of claim 1, wherein said admixture contains two or more active ingredients.

10 10. The method of claim 1, wherein said admixture is nebulized by a nebulizer.

11. The method of claim 1, wherein said active ingredient is selected from ophthalmic agents, dermal agents, ENT agents, and respiratory agents.

15 12. The method of claim 11, wherein said ophthalmic agent is selected from the group consisting of vitamins, steroids, cephalosporins, penicillins, tetracyclines, macrolides, aminoglycosides, sulfonamides, polypeptides, and naphazoline hydrochloride.

20 13. The method of claim 11, wherein said dermal agent selected from the group consisting of vitamins, steroids, cephalosporins, penicillins, tetracyclines, macrolides, aminoglycosides, sulfonamides, polypeptides, hydroquinone, piroxicam, and indomethacin.

25 14. The method of claim 11, wherein said ENT agent is

selected from the group consisting of vitamins, steroids, cephalosporins, penicillins, tetracyclines, macrolides, aminoglycosides, sulfonamides, polypeptides, naphazoline hydrochloride, and acetylcysteine.

- 5 15. The method of claim 1, wherein said respiratory agent is selected from the group consisting of vitamins, steroids, cephalosporins, penicillins, tetracyclines, macrolides, aminoglycosides, sulfonamides, polypeptides, ephedrine sulfate, methylephedrine, acetylcyseine, isoproterenol sulfate,
10 and aminophylline.

