(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
3 May 2001 (03.05.2001)

PCT

(10) International Publication Number
WO 01/30353 A1

(51) International Patent Classification[7]: A61K 31/505, 31/616, A61P 7/02 // (A61K 31/616, 31:505)

(21) International Application Number: PCT/EP00/10123

(22) International Filing Date: 14 October 2000 (14.10.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
99121121.0 22 October 1999 (22.10.1999) EP

(71) Applicant *(for all designated States except US)*: **BOEHRINGER INGELHEIM PHARMA KG** [DE/DE]; 55218 Ingelheim/Rhein (DE).

(72) Inventor; and
(75) Inventor/Applicant *(for US only)*: **EISERT, Wolfgang** [DE/DE]; Nettelbeckstrasse 19, 30175 Hannover (DE).

(74) Agent: **LAUDIEN, Dieter**; Boehringer Ingelheim GmbH, B Patente, 55216 Ingelheim/Rhein (DE).

(81) Designated States *(national)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States *(regional)*: ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:
— *With international search report.*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/30353 A1

(54) **Title:** USE OF DIPYRIDAMOLE OR MOPIDAMOL IN THE MANUFACTURE OF A MEDICAMENT FOR THE TREATMENT AND PREVENTION OF FIBRIN-DEPENDENT MICROCIRCULATION DISORDERS

(57) **Abstract:** A method of treatment of the human or non-human animal body for treating fibrin-dependent microcirculation disorders is disclosed, for example microcirculation disorders caused by metabolic diseases, inflammatory reactions or autoimmune diseases, furthermore peripheral microcirculation disorders or microcirculation disorders associated with increased cell fragmentation, which method comprises administering to a human or non-human animal body in need of such treatment an effective amount of a pharmaceutical composition containing a pyrimido-pyrimidine selected from dipyridamole, mopidamol and the pharmaceutically acceptable salts thereof, and the use of said pyrimido-pyrimidine for the manufacture of a corresponding pharmaceutical composition.

USE OF DIPYRIDAMOLE OR MOPIDAMOL IN THE MANUFACTURE OF A MEDICAMENT FOR THE TREATMENT AND PREVENTION OF FIBRIN-DEPENDENT MICROCIRCULATION DISORDERS

Field of the Invention

This invention relates to a method of treating fibrin-dependent microcirculation disorders using dipyridamole or mopidamol as active principle, providing a lasting improvement of microcirculation under treatment, and the use of dipyridamole or mopidamol for the manufacture of a corresponding pharmaceutical composition.

Background of the Invention

Dipyridamole {2,6-bis(diethanolamino)-4,8-dipiperidino-pyrimido[5,4-d]pyrimidine}, closely related substituted pyrimidopyrimidines and their preparation have been described in e.g. U.S.Patent 3,031,450. Further related substituted pyrimidopyrimidines and their preparation have been described in e.g. GB 1,051,218, inter alia the compound mopidamol {2,6-bis-(diethanolamino)-4-piperidinopyrimido[5,4-d]pyrimidine}. Dipyridamole was introduced as a *coronary vasodilator* in the early 1960s. It is also well known having *platelet aggregation inhibitor properties* due to the inhibition of adenosine uptake. Subsequently, dipyridamole was shown to reduce thrombus formation in a study of arterial circulation of the brain in a rabbit model. These investigations led to its use as an *antithrombotic agent*; it soon became the therapy of choice for such applications as stroke prevention, maintaining the patency of coronary bypass and valve-replacement, as well as for treatment prior to coronary angioplasty.

Furthermore, the European Stroke Prevention Study 2 (ESPS-2; J Neurol Sci. 1996; 143: 1-13; Neurology 1998; 51: 17-19) proved

that treatment by dipyridamole alone was as effective as low-dose aspirin in the reduction of stroke risk, and combination therapy with dipyridamole and aspirin was more than twice as effective as aspirin alone.

Dipyridamole appears to inhibit thrombosis through multiple mechanisms. Early studies showed that it inhibits the uptake of adenosine, which was found to be a potent endogenous anti-thrombotic compound. Dipyridamole was also shown to inhibit cyclic AMP phosphodiesterase, thereby increasing intracellular c-AMP.

By laboratory models reflecting the complex physiology of the blood vessel it could be shown that the vasculature is not a passive conduit, but interacts profoundly with the blood through an intricate system of checks and balances to protect its integrity after vascular accident. Therefore the endothelium produces prostacyclin, a potent inhibitor of aggregation. The normal endothelium is not thrombogenic and prevents the attachment of platelets. Various stimulants precipitate the release of endothelium-derived relaxing factor (EDRF), which inhibits platelet adhesion and aggregation. At the same time, intracellular increase in cGMP was shown to be responsible for relaxation of smooth muscle cells following administration of nitro compounds. Thus the endothelium can inhibit thrombus formation by two separate mechanisms, one mediated by prostacyclin and c-AMP, and the other by EDRF and c-GMP. Dipyridamole appears to enhance both of these anti-thrombotic mechanisms of the vessel wall, in addition to its adenosine-sparing effects. It stimulates prostacyclin production by increasing intracellular levels of cAMP, and it enhances the strongly anti-thrombotic nitric oxide system by increasing cGMP.

Dipyridamole also has *antioxidant properties* (Free Radic. Biol. Med. 1995; 18: 239-247) that may contribute to its antithrombotic effect. When oxidized, low density lipoproteins

become recognized by the scavenger receptor on macrophages, which is assumed to be the necessary step in the development of atherosclerosis (Ann. Rev. Med. 1992; 43: 219-25).

The inhibition of free radical formation by dipyridamole has been found to inhibit fibrinogenesis in experimental liver fibrosis (Hepatology 1996; 24: 855-864) and to suppress oxygen radicals and proteinuria in experimental animals with amino-nucleoside nephropathy (Eur. J. Clin. Invest. 1998; 28: 877-883; Renal Physiol. 1984; 7: 218-226). Inhibition of lipid peroxidation also has been observed in human nonneoplastic lung tissue (Gen. Pharmacol. 1996; 27: 855-859).

Mopidamol is known to possess antithrombotic and additionally antimetastatic properties.

Summary of the Invention

It has now surprisingly been found that dipyridamole and mopidamol exert a protective effect on the vessel wall thereby strongly influencing the interaction of the vessel wall with the fibrin pathway of the coagulation system resulting in an essential reduction of fibrin accumulation after stimulating clot formation.

It is known that vascular damages accelerate fibrin accumulation since the prothrombinase complex becomes significantly more potent when settled on negatively charged phospholipids of the cellular membrane. By stabilizing the cellular membrane, less negatively charged phosphadidylserines may become exposed on the outer cell membrane, offering fewer opportunities for the prothrombinase complex to bind to the phospholipids, and thereby preventing the prothrombinase complex from operating at its full conversion rate to turn prothrombin into thrombin which is responsible for the conversion of fibrinogen into fibrin.

Platelet accretion and fibrin accumulation are the basic pathways involved in clot formation. It has been shown that the time course of these two pathways differs essentially (Thromb. Haemost. 1993; 69 (Abstr.): 569) proving that both mechanisms are not as stringently coupled as it was anticipated. Whereas the activity of dipyridamole and mopidamol as platelet aggregation inhibitor is well known it is a new finding that these agents additionally are *inhibitors of fibrin accumulation* mediated by their capacity to stabilize cellular membranes of the vessel wall. This effect is especially important in small vessels or capillary vessels where the ratio of vessel wall surface area to blood volume is high, and provides a new approach for treatment and prevention of fibrin-dependent microcirculation disorders. Therefore dipyridamole and mopidamol may have therapeutic potential in a variety of diseases involving progressive dysfunction of medium and small-sized vessels.

The known vasodilating activity of dipyridamole was generally considered to be more important in the bigger vessels and in the context of short-term treatment or prevention of acute conditions. In using dipyridamole as a stress test agent it is known that by short-term high-dose infusion of dipyridamole the vascular autoregulation lags behind thereby showing disproportional perfusion. This is used to differentiate lesser increase in blood flow in post-stenotic areas compared with bigger increase in healthy segments of the circulation by nuclear imaging or echocardiography. In case of long-term oral treatment plasma dipyridamole as well as correlated adenosine levels increase over a period of several hours allowing the autoregulatory systems to compensate whereby under "stress test" conditions dipyridamole plasma levels as well as adenosine levels reach their peaks within four minutes. It has been found that treatment according to the present invention provides a lasting effect on small or capillary vessels and thereby a lasting improvement of microcirculation.

The finding that dipyridamole and mopidamol have significant inhibitory effects on fibrin accumulation via the vessel wall and a stabilizing effect on cell membranes provides a rationale also for combination treatment together with other antithrombotic agents, such as platelet aggregation inhibitors, e.g. acetylsalicalic acid (ASA), clopidogrel or ticlopidine or the pharmaceutically acceptable salts thereof, fibrinogen receptor antagonists (Abciximab, RDGS-peptides, synthetic i.v. or oral fibrinogen antagonists, e.g. fradafiban, lefradafiban or pharmaceutically acceptable salts thereof), heparin and heparinoids or antithrombins, or for combination treatment using additional cardiovascular therapies such as treatment with ACE inhibitors, Angiotensin II antagonists, Ca-antagonists or lipid-lowering agents such as the statins.

ASA inhibits aggregation through direct effects on the platelet, in more detail, by irreversibly acetylating platelet cyclooxygenase, thus inhibiting the production of thromboxane, which is strongly thrombotic. In high doses, however, aspirin crosses over into endothelial cells (N. Eng. J. Med. 1984; 311: 1206-1211), where it interrupts the production of prostacyclin, a potent natural inhibitor of platelet aggregation and by-product of the "arachidonic cascade" (N. Engl. J. Med. 1979; 300: 1142-1147). These observations led to the concept of low-dose antiplatelet therapy with ASA to maximize inhibition of thromboxane while minimizing the loss of prostacyclin (Lancet 1981; 1: 969-971). In combination with dipyridamole or mopidamol according to the invention also the low-dose ASA concept is preferred.

Viewed from one aspect the present invention provides a method of treatment of the human or non-human animal body, preferably mammalian body, for treating or preventing fibrin-dependent microcirculation disorders or of disease states where such microcirculation disorders are involved, said method comprising administering to said body an effective amount of a pharmaceutical composition comprising a pyrimido-pyrimidine selec-

ted from dipyridamole, mopidamol and the pharmaceutically acceptable sals thereof, dipyridamole being preferred, optionally in combination with one or more other antithrombotic agents.

Viewed from a different aspect the present invention provides the use of a pyrimido-pyrimidine selected from dipyridamole, mopidamol and the pharmaceutically acceptable salts thereof

for the manufacture of an oral pharmaceutical composition comprising the pyrimido-pyrimidine as the only active compound adapted to provide a daily dosage of 25 to 450 mg of the pyrimido-pyrimidine for the treatment of the human or non-human animal body for treating or preventing fibrin-dependent microcirculation disorders or of disease states where such microcirculation disorders are involved selected from the group consisting of

microcirculation disorders caused by metabolic diseases where vascular damages are involved,

such as diabetic angiopathy, especially diabetic microangiopathy, e.g. diabetic gangrene, diabetic retinopathy, diabetic neuropathy or ulcus cruris,

microcirculation disorders caused by inflammatory reactions,

such as morbus crohn,

microcirculation disorders caused by autoimmune diseases,

such as autoimmune chronic-active hepatitis (idiopathic hepatitis), primary-biliary cirrhosis or (autoimmune associated) multiple sclerosis,

peripheral microcirculation disorders,

such as Raynaud's disease, tinnitus or
sudden loss of hearing,

microcirculation disorders associated with increased cell fragmentation,

such as tumor diseases or thrombotic-thrombocytopenic purpura (TTP),

and, as further indications,
nephrosclerosis,
prerenal hypertension,
haemolytic-uremic syndrome (HUS),
arterial hypertension,
vascular dementia,
Sudeck's disease,
central-veneous thrombosis of the eye,
homocystine-induced vasculopathy,
ischemic or coronary heart diseases,
prevention of myocardial infarction or reinfarction and
treatment or prevention of atherosclerosis.

In another aspect, the present invention provides the use of a pyrimido-pyrimidine selected from dipyridamole, mopidamol and the pharmaceutically acceptable salts thereof, in combination with

one or more other antithrombotic agents selected from acetylsalicalic acid (ASA), clopidogrel, ticlopidine and the pharmaceutically acceptable salts thereof, fibrinogen receptor antagonists, heparin, heparinoids and antithrombins,

or in combination with an ACE inhibitor, an Angiotensin II antagonist, a Ca-antagonist or a lipidlowering agent,

for the manufacture of a pharmaceutical composition for the treatment of the human or non-human animal body for treating or preventing disorders or disease states as defined above,

wherein said composition is adapted to provide the pyrimido-pyrimidine in an oral daily dosage of 25 to 450 mg when combined with clopidogrel, ticlopidine and the pharmaceutically acceptable salts thereof, fibrinogen receptor antagonists, heparin, heparinoids or antithrombins, or with an ACE inhibitor, Angiotensin II antagonist, Ca-antagonist or lipidlowering agent, and,

when combined with ASA, said composition is adapted to provide the pyrimido-pyrimidine in an oral daily dosage of 50 to 300 mg and the ASA in an oral daily dosage of 10 to 30 mg.

In yet another aspect, the present invention provides a pharmaceutical composition comprising a pyrimido-pyrimidine selected from dipyridamole, mopidamol and the pharmaceutically acceptable salts thereof adapted to provide the pyrimido-pyrimidine in an oral daily dosage of 25 to 450 mg, in combination with

one or more other antithrombotic agents selected from clopidogrel, ticlopidine and the pharmaceutically acceptable salts thereof, fibrinogen receptor antagonists, heparin, heparinoids and antithrombins,

or in combination with an ACE inhibitor, an Angiotensin II antagonist, a Ca-antagonist or a lipidlowering agent.

Detailed Description of the Invention

The invention provides a new approach for the treatment and prevention of fibrin-dependent microcirculation disorders associated with progressive dysfunction of medium and small-sized vessels comprising administering to a person in need of such treatment an effective amount of a pharmaceutical composition containing a pyrimido-pyrimidine selected from dipyridamole, mopidamol and the pharmaceutically acceptable salts thereof.

Fibrin-dependent microcirculation disorders are meant to be such disorders where fibrin deposition is involved in pathogenesis or progression of dysfunction of medium or small-sized vessels leading to a variety of clinical pictures. Metabolic diseases such as diabetes mellitus are known to cause said microcirculation disorders, however, also inflammatory reactions may cause microcirculation disorders due to local fibrinogen release from the tissue site of inflammation. Furthermore it is assumed that microcirculation disorders also can be caused by autoimmune reactions.

The indication "fibrin-dependent microcirculation disorders" should be understood in a non-limiting manner to comprise

microcirculation disorders caused by metabolic diseases where vascular damages are involved,

such as diabetic angiopathy, especially diabetic micro-angiopathy, e.g. diabetic gangrene, diabetic retinopathy, diabetic neuropathy or ulcus cruris,

microcirculation disorders caused by inflammatory reactions,

such as morbus crohn,

microcirculation disorders caused by autoimmune diseases,

such as autoimmune chronic-active hepatitis (idiopathic hepatitis), primary-biliary cirrhosis or (autoimmune associated) multiple sclerosis,

peripheral microcirculation disorders,

such as Raynaud's disease, tinnitus or sudden loss of hearing,

microcirculation disorders associated with increased cell fragmentation,

such as tumor diseases or thrombotic-thrombocytopenic purpura (TTP),

or, as further indications,
nephrosclerosis,
prerenal hypertension,
haemolytic-uremic syndrome (HUS),
arterial hypertension,
vascular dementia,

Alzheimer's disease,
Sudeck's disease,
central-veneous thrombosis of the eye,
ischemic optic neuropathy,
homocystine-induced vasculopathy,
ischemic or coronary heart diseases,
prevention of myocardial infarction or reinfarction and
treatment or prevention of atherosclerosis.

The indication "fibrin-dependent microcirculation disorders" also includes corresponding disorders of the myocardium. Thus the present invention provides a method for improving the blood supply of the myocardium in a person in need of such treatment, for example in a person suffering from ischemic or coronary heart disease, as well as a method for prevention of myocardial infarction or reinfarction.

Furthermore, the present invention also provides a method of treatment or prevention of atherosclerosis since administration of dipyridamole or mopidamol supports or helps to improve or restore the microcirculation supplying the vessels.

As already mentioned hereinbefore dipyridamole, mopidamol or a pharmaceutically acceptable salt thereof can be used alone in a monopreparation or in combination with other antithrombotic agents for the treatment of fibrin-dependent microcirculation disorders.

It is of advantage to maintain a plasma level of dipyridamole or mopidamol of about 0.2 to 5 μmol/L, preferably of about 0.4 to 5 μmol/L, especially of about 0.5 to 2 μmol/L or particularly of about 0.8 to 1.5 μmol/L. This can be achieved using any of the oral dipyridamole retard, instant or the parenteral formulations on the market, the retard formulations being preferred, for instance those available under the trademark Persantin®, or, for the combination therapy with low-dose ASA, using those formulations available under the

trademark Asasantin® or Aggrenox®. Dipyridamol retard formulations are also disclosed in EP-A-0032562, instant formulations are disclosed in EP-A-0068191 and combinations of ASA with dipyridamole are disclosed in EP-A-0257344 which are incorporated by reference. In case of mopidamol also oral retard, instant or a parenteral formulations can be used, e.g. those disclosed in GB 1,051,218 or EP-A-0,108,898 which are incorporated by reference, retard formulations being preferred.

Dipyridamole or mopidamol can be administered orally in a daily dosage of 25 to 450 mg, preferably 50 to 240 mg, most preferred 75 to 200 mg. For long-term treatment it is of advantage to administer repeated doses such as a dose of 25 mg dipyridamole retard or any other instant release formulation three or four times a day. For parenteral administration dipyridamole could be given in a dosage of 0.5 to 5 mg/kg body weight, preferably 1 to 3.5 mg/kg body weight, during 24 hours as slow i.v. infusion (not faster than 0.2 mg/min).

Dipyridamole or mopidamol in combination with low-dose ASA may be administered orally in a daily dosage of 10 to 30 mg of ASA together with 50 to 300 mg of dipyridamole or mopidamol, preferably 80 to 240 mg of dipyridamole or mopidamol, for instance in a weight ratio between 1 to 5 and 1 to 12, most preferred a weight ratio of 1 to 8, for instance 25 mg of ASA together with 200 mg of dipyridamole or mopidamol.

Other antithrombotic compounds would be given at 0.1 to 10 times, preferably at 0.3 to 5.0 times, most preferred at 0.3 to 2.0 times the clinically described dose (e.g. Rote Liste® 1999; fradafiban, lefradafiban: EP-A-0483667), together with a daily dosage of 25 to 450 mg, preferably 50 to 240 mg, most preferred 75 to 200 mg of dipyridamole or mopidamol.

For combination treatment using dipyridamole or mopidamol together with ACE inhibitors any ACE inhibitor known in the art would be suitable, e.g. benazepril, captopril, ceronapril, enalapril, fosinopril, imidapril, lisinopril, moexipril, quinapril, ramipril, trandolapril or perindopril, using the dosages known in the art, for instance as described in Rote Liste® 1999, Editio Cantor Verlag Aulendorf.

For combination treatment using dipyridamole or mopidamol together with Angiotensin II antagonists any Angiotensin II antagonist known in the art would be suitable, e.g. the sartans such as candesartan, eprosartan, irbesartan, losartan, telmisartan, valsartan, olmesartan or tasosartan, using the dosages known in the art, for instance as described in Rote Liste® 1999, Editio Cantor Verlag Aulendorf.

For combination treatment using dipyridamole or mopidamol together with Ca-antagonists any Ca-antagonist known in the art would be suitable, e.g. nifedipine, nitrendipine, nisoldipine, nilvadipine, isradipine, felodipine or lacidipine, using the dosages known in the art, for instance as described in Rote Liste® 1999, Editio Cantor Verlag Aulendorf.

For combination treatment using dipyridamole or mopidamol together with statins any statin known in the art would be suitable, e.g. lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin or cerivastatin, using the dosages known in the art, for instance as described in Rote Liste® 1999, Editio Cantor Verlag Aulendorf.

It should be noted that such microcirculation disorders associated with increased cell fragmentation, as mentioned hereinbefore, especially accelerate fibrin accumulation due to the potentiated free cellular membrane areas activating the prothrombinase complex. These microcirculation disorders, for

example tumor diseases or thrombotic-thrombocytopenic purpura, are preferably treated using high doses of dipyridamole or mopidamol. This means that a plasma level of dipyridamole or mopidamol of about 0.2 to 50 μmol/L, preferably of about 0.5 to 20 μmol/L, particularly of about 0.5 to 10 μmol/L should be maintained preferably by slow i.v. infusion. For oral treatment of these indications dipyridamole or mopidamol should be administered in a daily dosage of about 150 to 1000 mg, preferably 200 to 800 mg, most preferred 200 to 600 mg.

Microcirculation disorders associated with increased cell fragmentation also can be treated by a combination of dipyridamole or mopidamol with low-dose ASA using the high doses of dipyridamole or mopidamol mentioned above, together with an oral daily dosage of about 10 to 30 mg of ASA, preferably of about 25 mg of ASA.

In order to study the inhibition of fibrin accumulation by dipyridamole the following experiment was carried out:

Study Using Radio Labeled Platelets and Fibrinogen In Vivo

The effects of dipyridamole and heparin were investigated using platelets radiolabeled with ^{99}Tc and fibrinogen labeled with ^{123}J. The method is described in Nuclear Instruments and Methods in Physics Research A. 1994, 353: 448-452. An energy-sensitive solid-state radiation detector was placed to surround each carotid artery in rabbits. After inducing injury through balloon angioplasty, the accretion of platelets and fibrinogen was monitored for four hours. It was found that platelet and fibrinogen accretion had dissimilar time courses. Accretion was not detected in radiolabeled platelets injected 30 minutes after injury; apparently the angioplasty site had been passivated by endogenous platelet adherence. In contrast, fibrinogen accretion did not reach a plateau in control or treated animals even after four hours, indicating that different stimuli or triggers may be regulating fibrinogen at va-

rious time points after injury (Fig. 1). Treatment with heparin reduced accumulation of both platelets and fibrinogen. Treatment with dipyridamole also produced a reduction in platelet aggregation, which was similar to that seen with heparin; the reduction in accretion of fibrinogen was far greater with dipyridamole than with heparin, however (Fig. 2).

FIGURE LEGENDS:

Fig. 1: Simultaneous detection of ^{99}Tc labeled platelets and ^{123}I-labeled fibrinogen at one minute interval after angioplasty of the common carotid artery of rabbits. Control (no treatment) group (N=6) showed after injury a rapid increase of platelets and a gradual build up of fibrinogen. Treatment with heparin (100U /kg bolus followed by 25U/ kg/h infusion) showed reduction of platelet as well as fibrinogen accretion. No injury measurements showing constant radioactivity.

Fig. 2: Deposition of radioactive labeled platelets (^{99}Tc) and fibrinogen (^{123}I) at angioplasty site after treatment with dipyridamole (0.25 mg/kg followed by 0.45 mg/kg/hr). Platelet deposition is reduced, but fibrinogen accretion is almost entirely blocked during the first four hours after angioplasty.

CLAIMS

1. The use of a pyrimido-pyrimidine selected from dipyridamole, mopidamol and the pharmaceutically acceptable salts thereof for the manufacture of an oral pharmaceutical composition comprising the pyrimido-pyrimidine as the only active compound adapted to provide a daily dosage of 25 to 450 mg of the pyrimido-pyrimidine for the treatment of the human or non-human animal body for treating or preventing fibrin-dependent microcirculation disorders or of disease states where such microcirculation disorders are involved selected from from the group consisting of

microcirculation disorders caused by metabolic diseases where vascular damages are involved,

such as diabetic angiopathy, especially diabetic microangiopathy, e.g. diabetic gangrene, diabetic retinopathy, diabetic neuropathy or ulcus cruris,

microcirculation disorders caused by inflammatory reactions,

such as morbus crohn,

microcirculation disorders caused by autoimmune diseases,

such as autoimmune chronic-active hepatitis (idiopathic hepatitis), primary-biliary cirrhosis or (autoimmune associated) multiple sclerosis,

peripheral microcirculation disorders,

such as Raynaud's disease, tinnitus or
sudden loss of hearing,

microcirculation disorders associated with increased cell fragmentation,

such as tumor diseases or thrombotic-thrombocytopenic purpura (TTP),

and, as further indications,
nephrosclerosis,
prerenal hypertension,
haemolytic-uremic syndrome (HUS),
arterial hypertension,
vascular dementia,
Sudeck's disease,
central-veneous thrombosis of the eye,
homocystine-induced vasculopathy,
ischemic or coronary heart diseases,
prevention of myocardial infarction or reinfarction and
treatment or prevention of atherosclerosis.

2. The use of claim 1, wherein the pharmaceutical composition comprises the pyrimido-pyrimidine in an oral retard or instant formulation.

3. The use of a pyrimido-pyrimidine selected from dipyridamole, mopidamol and the pharmaceutically acceptable salts thereof, in combination with

one or more other antithrombotic agents selected from acetylsalicalic acid (ASA), clopidogrel, ticlopidine and the pharmaceutically acceptable salts thereof, fibrinogen receptor antagonists, heparin, heparinoids and antithrombins,

or in combination with an ACE inhibitor, an Angiotensin II antagonist, a Ca-antagonist or a lipidlowering agent,

for the manufacture of a pharmaceutical composition for the treatment of the human or non-human animal body for treating or preventing disorders or disease states as defined in claim 1,

wherein said composition is adapted to provide the pyrimido-pyrimidine in an oral daily dosage of 25 to 450 mg when combined with clopidogrel, ticlopidine and the pharmaceutically acceptable salts thereof, fibrinogen receptor antagonists, heparin, heparinoids or antithrombins, or with an ACE inhibitor, Angiotensin II antagonist, Ca-antagonist or lipidlowering agent, and,

when combined with ASA, said composition is adapted to provide the pyrimido-pyrimidine in an oral daily dosage of 50 to 300 mg and the ASA in an oral daily dosage of 10 to 30 mg.

4. The use of any of claims 1 to 3, characterized in that the pyrimido-pyrimidine is dipyridamole or a pharmaceutically acceptable salt thereof.

5. A pharmaceutical composition comprising a pyrimido-pyrimidine selected from dipyridamole, mopidamol and the pharmaceutically acceptable salts thereof adapted to provide the pyrimido-pyrimidine in an oral daily dosage of 25 to 450 mg, in combination with

one or more other antithrombotic agents selected from clopidogrel, ticlopidine and the pharmaceutically acceptable salts thereof, fibrinogen receptor antagonists, heparin, heparinoids and antithrombins,

or in combination with an ACE inhibitor, an Angiotensin II antagonist, a Ca-antagonist or a lipidlowering agent.

6. The pharmaceutical composition of claim 5 comprising the pyrimido-pyrimidine in combination with clopidogrel or ticlopidine.

7. The pharmaceutical composition of claim 5 comprising the pyrimido-pyrimidine in combination with an ACE inhibitor selected from benazepril, captopril, ceronapril, enalapril, fosinopril, imidapril, lisinopril, moexipril, quinapril, ramipril, trandolapril or perindopril.

8. The pharmaceutical composition of claim 5 comprising the pyrimido-pyrimidine in combination with an Angiotensin II antagonist selected from the sartans.

9. The pharmaceutical composition of claim 8 wherein the sartan is selected from candesartan, eprosartan, irbesartan, losartan, telmisartan, valsartan, olmesartan and tasosartan.

10. The pharmaceutical composition of claim 5 comprising the pyrimido-pyrimidine in combination with a Ca-antagonist selected from nifedipine, nitrendipine, nisoldipine, nilvadipine, isradipine, felodipine and lacidipine.

11. The pharmaceutical composition of claim 5 comprising the pyrimido-pyrimidine in combination with a lipidlowering agent selected from the statins.

12. The pharmaceutical composition of claim 11 wherein the statin is selected from lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin and cerivastatin.

13. The pharmaceutical composition of any of claims 5 to 12 wherein the pyrimido-pyrimidine is dipyridamole or a pharmaceutically acceptable salt thereof.