

DESCRIPTION

CELL CULTURE METHOD AND SCREENING METHOD

Technical Field

5 [0001]

The present invention relates to a cell culture method and a screening method.

Background Art

[0002]

10 A technique of using cells isolated from tissue for tests and examinations is essential in biotechnology-related fields. Such a technique is widely used for a diagnosis of diseases and pathoses, a search for a new drug and an assessment of its effect, an animal inspection, a plant inspection, a test of an environment pollutant and
15 so on. Therefore, cells used in the biotechnology field are becoming increasingly diversified.

[0003]

Although some isolated cells are immediately used for tests in a floating state, others are cultured in a state of being adherent to a
20 culture dish and used for various tests and examinations in most cases. Primary cells and cell lines used for cell culture are required to exhibit drug sensitivity, toxicity reaction or the like of a similar level to a test in vivo, so-called an in vivo test. In other words, an in vivo-like cell function is required in a cell culture container. Further, because
25 isolation for obtaining primary cells is complicated and cell culture lines used for a cell culture test are expensive, a test method with a small number of cells is desired.

[0004]

30 Recently, discontinuation of development in the clinical testing phase has been an issue. This is due to an animal species difference in the pharmacokinetics study phase. Heretofore, in pharmacokinetics studies in the preclinical phase, in vivo pharmacokinetics has been predicted by using an animal such as a rat, a dog or a monkey. However, it has becoming evident that the prediction is virtually

invalid in a clinical testing using a human. Therefore, in the prediction of pharmacokinetics or the like for humans, using a human sample is the most effective and convenient way, and it is important for conducting efficient drug development and safe clinical testing.

5 [0005]

In the pharmacokinetics study that examines in vivo pharmacokinetics, absorption, metabolism and excretion in the liver are mainly examined, and a human sample to be used is liver slices, liver cells, liver microsomes or the like. Among those, the liver slices
10 are not easily obtainable, and the liver microsomes can be used only for a metabolism test with limited metabolic enzymes. Thus, use of the liver cells is considered to be the most effective.

[0006]

In screening, a culture dish to be used is a petri dish made of
15 resin or a 6-well, 12-well, 48-well or 96-well plate. The size of the entire plate is substantially the same, and as the number of wells increases, the size of one well decreases. One well corresponds to one culture dish. Further, with the recent trend toward micronization, a 384-well plate made up of a larger number of culture dishes with a
20 smaller diameter has been started to be used, and the one adaptable for a desired screening method is used. The bottom of such a culture dish is a flat-shaped, and the bottom surface is used as a culture surface.

[0007]

However, if a hitherto-used culture dish is used for culture of
25 tissue cells, there are cases where the original function disappears and dedifferentiation occurs and where undifferentiated cells do not differentiate, which raises an issue that a target cell function is not expressed. For example, if fresh human liver cells are cultured on a normal flat plate, the function of metabolic enzymes when isolated is
30 significantly lowered in one day or so, and therefore a drug metabolism test is conducted in four hours from seeding the cells onto the plate in some cases. There is thus a problem that it is impossible to make use for a test with long-hours culture and a problem that it is impossible to investigate long-hours metabolic stability.

[0008]

To overcome the above problems, an attempt to coat a culture container surface with a biological material (glycoprotein, protein etc.) of human or animal origin (cf. Patent Document 1), an attempt to
5 culture in polymer gel (cf. Patent Document 2), and an attempt to form a liver cell mass in a micro-container (cf. Patent Document 3) have been made.

[Patent Document 1]

Japanese Unexamined Patent Application Publication No. 8-319317

10 [Patent Document 2]

Japanese Unexamined Patent Application Publication No.8-308562

[Patent Document 3]

International Patent Publication No. WO2008/130025

Disclosure of Invention

15 **Technical Problem**

[0009]

However, in the method disclosed in Patent Document 1, there are problems such that the biological material as coating is special and high cost, it is difficult to form a uniform cell assembly in a culture
20 container, and the in vivo function cannot be sustained over a long period of time. In the method disclosed in Patent Document 2 also, there are problems such that the size of a cell assembly cannot be controlled, microscopic observation is not easy, and operability is complicated as a screening substrate. Further, because a commercially
25 available dish or plate is used as a supporting container in both of the above methods, efficient screening with the minimum number of cells required is difficult. In the method disclosed in Patent Document 3 also, although improvement of liver cell metabolic activity in the early stage of culture is possible, maintaining the metabolic activity for two
30 weeks or longer is difficult.

[0010]

An object of the present invention is to provide a cell culture method whereby an in vivo function can be sustained over a long period of time and culture can be conducted using the minimum number

of cells required.

Technical Solution

[0011]

A cell culture method according to the present invention
5 includes culturing undifferentiated cells in a layered state in a
partitioned micro-space and obtaining differentiated cells. The layered
state means that cells are laminated two or more layers. When
screening a pharmaceutical agent, undifferentiated cells capable of
differentiating into liver cells, intestinal epithelial cells, nerve cells,
10 myocardial cells and vascular endothelial cells are preferred.
Particularly, in the prediction of pharmacokinetics or the like for
humans, undifferentiated cells are preferably human cells. The
undifferentiated cells are preferably stem cells, precursor cells or the
like.

15 [0012]

Further, it is particularly preferred that the partitioned micro-
space is a micro-container in a cell culture container having a plurality
of micro-containers on a surface. The micro-container preferably has a
bottom surface area of $9 \times 10^{-4} \text{ mm}^2$ to $9 \times 10^{-2} \text{ mm}^2$, and a side wall
20 with a height of 15 μm to 300 μm and a width of 3 μm to 15 μm .
Further, in order to facilitate microscopic observation, it is preferred
that a region where the micro-container is formed in the cell culture
container has transparency.

[0013]

25 A screening method according to the present invention includes
placing a plurality of cells cultured by the above-described culture
method and screening a compound. Further, in order to conduct culture
using the minimum number of cells required, it is preferred that the
cell culture container includes a plurality of partitioned spots each
30 made up of a plurality of micro-containers.

Advantageous Effects

[0014]

According to the present invention, it is possible to provide a
cell culture method whereby an in vivo function can be sustained

uniformly over a long period of time and culture can be conducted using the minimum number of cells required.

Brief Description of Drawings

[0015]

5 Fig. 1 is a plan view showing a structure of a cell culture container used for a cell culture method according to an example 1;

Fig. 2 is a cross-sectional view showing a structure of a cell culture container used for the cell culture method according to the example 1;

10 Fig. 3 is a plan view showing a structure of a cell culture container used for a screening method according to an embodiment; and

Fig. 4 is a cross-sectional view showing a structure of a cell culture container used for the screening method according to the embodiment.

15 Explanation of Reference

[0016]

10 CELL CULTURE UNIT

11 MICRO-CONTAINER

12 SIDE WALL OF MICRO-CONTAINER

20 13 SPOT

14 SIDE WALL OF SPOT

Best Mode for Carrying Out the Invention

[0017]

In the present invention, undifferentiated cells such as liver
 25 stem cells are cultured in a layered state with a uniform size in a micro-container, which is a partitioned micro-space. An especially important point in the invention is to use undifferentiated cells having proliferation potency, to culture in a partitioned micro-space, and to culture in a layered fashion. It is thereby possible to form a cell mass
 30 containing differentiated liver parenchymal cells, for example, and improve the function. Further, because liver stem cells undergo undifferentiated proliferation, it is possible to sustain the cell function. The present invention provides such a culture method and a screening method using the same.

[0018]

On the surface of a culture container used in the culture method and the screening method according to the present invention, a pit and projection pattern or a plurality of micro-containers or a culture space is formed. By optimizing the width and the height of a side wall (projection) that partitions the micro-containers, it is possible to culture cells only in the micro-containers and maintain a uniform differentiated state. Note that a partitioned culture space made of gel may be formed instead of the micro-containers.

[0019]

The dimensions of a micro-container surrounded by side walls should be in an optimum range for culturing cells. If the bottom surface area of the micro-container is too large, a cell partially extends thinly, and a uniform layered state is not formed as in the case of culture on a flat plate. On the other hand, if the bottom surface area of the micro-container is too small, a cell cannot be contained. Therefore, the dimensions of the space are preferably in the range to contain several to several tens of cells according to cell species to be cultured.

[0020]

Further, the side wall of the micro-container also should be in an optimum range for culturing cells. If the width of the side wall is too wide, a cell adheres to the top surface of the side wall, which is unsuited to culture. If the width of the side wall is too narrow, preparation is difficult. If the height of the side wall is too low, a cell goes over the side wall, which is unsuited to culture. If the height of the side wall is too high, preparation is difficult and further the material is hard to diffuse, which degrades the culture environment.

[0021]

Hereinafter, an embodiment of the present invention is described. Note that, however, the present invention is not limited to the following embodiment. Further, to clarify the explanation, the following description and the drawings are appropriately simplified.

[0022]

Embodiment

A structure of a micro-container of a cell culture unit used for cell culture according to an embodiment is described hereinafter with reference to Figs. 1 and 2. Fig. 1 is a plan view showing a structure of a micro-container according to the embodiment, and Fig. 2 is a cross-sectional view along line II-II in Fig. 1. As shown in Fig. 1, a cell culture unit 10 includes a micro-container 11 and a side wall 12. On a culture surface of the cell culture unit 10, a plurality of side walls 12 are formed in a mesh pattern, and a space surrounded on all four sides by the side walls 12 is the micro-container 11.

[0023]

In Fig. 1, the width a of the bottom surface of the micro-container 11, and the width b and the height c of the side wall 12 for partitioning the micro-containers 11 are shown. It is necessary to satisfy $3\mu\text{m} \leq b \leq 15\mu\text{m}$ and $c/b \geq 2$. If the width b of the side wall 12 is more than $15\mu\text{m}$, a cell adheres to the top surface of the side wall, which is unsuited to culture. On the other hand, if the width b of the side wall 12 is less than $3\mu\text{m}$, preparation is difficult. If the height of the side wall is too low, a cell goes over the side wall, which is unsuited to culture. If the height c of the side wall 12 is less than two times the width b of the side wall 12, a cell cultured in the micro-container 11 goes over it and moves to the adjacent micro-container 11. Further, specifically, when human fetal liver cells are layered in a square micro-container with one side of $100\mu\text{m}$, the height c of the side wall 12 is preferably $15\mu\text{m}$ to $300\mu\text{m}$, and more preferably $50\mu\text{m}$ to $150\mu\text{m}$. If the height c of the side wall is too high, preparation is difficult and further the material is hard to diffuse, which degrades the culture environment. The side wall 12 may have a multi-step shape.

[0024]

The bottom surface shape of the micro-container 11 is not particularly limited, and various shapes may be employed other than a square, circle, or polygonal shape. The bottom surface area is preferably $9 \times 10^{-4} \text{ mm}^2$ to $9 \times 10^{-2} \text{ mm}^2$. Further, an isotropic shape is preferable, and when the bottom surface has a rectangular shape, it is

preferred that the long side is 1 to 1.5 times the short side.

[0025]

The cell culture unit used in the present invention may have partitioned spots each made up of a plurality of micro-containers required for one screening as shown in Figs. 3 and 4 in order to minimize the number of cells required. For example, in the case of using a micro-container in a square shape with one side of 200 μm and a height of 50 μm which provides a high differentiation efficiency when the minimum number of cells required for screening is about 1000, nine micro-containers are required; therefore, by preparing a spot in which the space is partitioned into nine micro-containers and providing a plurality of spots, it is possible to perform high-throughput screening that allows simultaneous examination of a plurality of reagents or pharmaceutical agents.

[0026]

Fig. 3 is a plan view showing another structure of a cell culture unit according to the embodiment, and Fig. 4 is a cross-sectional view along line IV-IV in Fig. 3. Fig. 3 shows a side wall 14 that partitions a plurality of micro-containers and a partitioned spot 13. The height d of the side wall 14 may be set so that the capacity can keep a supernatant fluid such as a culture solution or a reaction solution without drying, and it can be defined appropriately.

[0027]

Although a method of creating the pit and projection pattern of the cell culture unit according to the present invention is not particularly limited, it may be a method such as transcription molding using a mold, 3D optical modeling, precision machine cutting, wet etching, dry etching, laser processing or electrical discharge machining, for example. It is preferred to appropriately select such a manufacturing method in consideration of the purpose of the cell culture container, a required processing accuracy, a cost or the like.

[0028]

As a specific example of the transcription molding method using a mold, a method of creating a pit and projection pattern by resin

molding with use of a metal structure as a mold may be used. This method is preferable because it enables reproduction of the shape of the metal structure onto resin as a pit and projection pattern with a high transcription rate, and use of a general-purpose resin material can reduce a material cost. Such a method of using the metal structure is advantageous in that it is low cost and satisfies a high dimensional accuracy.

[0029]

A method of manufacturing the above-described metal structure may be plating onto a resist pattern formed by photolithography or a resin pattern formed by 3D optical modeling, precision machine cutting, wet etching, dry etching, laser processing, electrical discharge machining or the like, for example. It is preferred to appropriately select the method in consideration of a purpose, a required processing accuracy, a cost or the like.

[0030]

A method of forming a pit and projection pattern on resin by using the metal structure obtained above as a mold may be injection molding, press molding, monomer-cast molding, solvent cast molding, hot embossing molding, roll transcription by extrusion molding or the like, for example. It is preferred to use the injection molding in terms of productivity and mold transcription ability.

[0031]

A material of a screening chip according to the present invention is not particularly limited as long as it has self-bearing properties, and synthetic resin, silicon, glass or the like may be used, for example. In terms of cost and cell visibility by microscopic observation, it is preferred to use transparent synthetic resin as a material.

[0032]

The transparent synthetic resin may be acrylic resin such as polymethyl methacrylate and methyl methacrylate-styrene copolymer, styrene resin such as polystyrene, olefin resin such as cycloolefin, ester resin such as polyethylene terephthalate and polylactic acid, silicone resin such as polydimethylsiloxane, polycarbonate resin or the

like, for example. Such resin may contain various kinds of additives such as coloring agent, dispersing agent and thickening agent within the range that does not lose transparency.

[0033]

5 In the screening chip according to the present invention, surface treatment may be performed on the surface side of the pit and projection pattern to form a modified layer and/or a coating layer for the purpose of improving surface hydrophilicity, biocompatibility, cellular affinity or the like.

10 [0034]

A method of forming the modified layer is not particularly limited as long as it is not a method that loses self-bearing properties or a method that causes extreme surface roughness of 100 μm or more, and it may be a method like chemical treatment such as drug treatment, solvent treatment and introduction of graft polymer by surface graft polymerization, or physical treatment such as corona discharge, ozone treatment and plasma treatment, for example.

[0035]

20 Further, a method of forming the coating layer is not particularly limited, and it may be a method like dry coating such as sputtering and vapor deposition, wet coating such as inorganic material coating and polymer coating or the like, for example.

[0036]

25 It is preferred to provide hydrophilicity onto the pit and projection pattern in order to inject a culture solution without mixing of air bubbles, and inorganic vapor deposition is preferable as a method to form a uniform hydrophilic film.

[0037]

30 Further, in the case of taking cellular affinity into consideration, it is preferred to make coating of cytophilic protein such as collagen or fibronectin, for example. In order to uniformly coat a collagen solution or the like, it is preferred to make coating after forming the above-described hydrophilic film. Because culture on an extracellular matrix surface is generally desirable in cell culture

in imitation of the in vivo environment, it is particularly preferred to provide an organic film made of extracellular matrix suitable for cultured cells after providing a uniform hydrophilic inorganic film as described above.

5 [0038]

Cells to be cultured by the culture method and the screening method according to the present invention are preferably undifferentiated cells, and they may be liver stem cells, liver precursor cells, bowel stem cells, intestinal precursor cells, mesenchymal stem
10 cells, cardiac muscle precursor cells, embryonic stem cells or the like, for example, and cell species may be appropriately selected according to the purpose of screening. For example, when the purpose is screening in assumption of metabolic response of a pharmaceutical agent in a liver, liver stem cells capable of differentiating into liver
15 cells or the like are used. Particularly, human liver stem cells are used when the purpose is metabolic response of a pharmaceutical agent in a human.

[0039]

The screening method according to the present invention places
20 cells only in a micro-container for culturing cells and makes the function similar to that in vivo expressed in the space, and it is thus necessary to seed an appropriate number of cells. The cell seeding density is preferably 1.0×10^4 - 5.0×10^6 cells/cm². For example, when the micro-container has a square shape with one side of 100 μ m, it is
25 preferably 1.0×10^4 - 1.0×10^5 cells/cm².

[0040]

When cells to be layered are liver cells or intestinal epithelium cells, it is preferred to measure the gene expression level, metabolic enzyme activity, transporter activity or the like and conduct screening.

30 Further, when cells to be layered are nerve cells or myocardial cells, it is preferred to measure the gene expression level, enzyme activity action potential or the like and conduct screening.

When cells to be layered are vascular endothelial cells, it is preferred to visually check vascularization and conduct screening.

[0041]

[Example]

Examples of the cell culture method according to the present invention are described hereinafter; however, the present invention is not limited to those examples.

<Culture Method of Fetal Liver Cells>

A frozen stock of human fetal liver cells was seeded onto type IV collagen coat dish (available from BD), cultured for about ten days, and proliferated. After that, on a pit and projection pattern substrate coated with 0.03% type IV collagen (available from Nitta Gelatin Inc.), liver cells were seeded at a rate of five cells in one micro-container (3.8×10^4 cells/cm²) and cultured for three weeks with 5% CO₂ and 37°C. The composition of a culture solution used was such that 10% fetal bovine serum, 1μg/ml insulin, 1×10^{-7} M dexamethasone, 10mM nicotinamide, 2mM L-glutamine, 50 μm β-mercaptoethanol, 5mM HEPES, 59μg/ml penicillin, 100μg/ml streptomycin were added to a DMEM/F12 culture medium, and, from the first day after seeding, 25ng/ml HGF, 20ng/ml EGF, 10ng/ml oncostatin M were further added to the culture solution. With use of a fresh culture medium of 0.5 mL having the same composition, a culture medium is replaced every several days.

[0042]

<Gene Expression Analysis>

Gene expression of cytochrome P450 (CYP), which is a drug metabolizing enzyme of a liver, was evaluated by retrieving RNA from the cells that has been cultured for a predetermined number of days and conducting real-time PCR after cDNA synthesis.

[0043]

[Example 1]

The pit and projection pattern shape shown in Fig. 1 where a=100 μm, b=10 μm and c=50 μm was created by photolithography, Ni electrolytic plating was performed, and a mold having the corresponding pit and projection shape was obtained. Using the mold, pattern transcription was made onto polystyrene by hot embossing

molding, and a resin substrate with the above dimensions was produced. A silicon dioxide film of 100 nm was deposited on the resin substrate surface by vacuum deposition, γ -ray sterilization was performed, and thereby a pit and projection pattern substrate was obtained. After coating the pit and projection substrate with IV collagen, human fetal liver cells were cultured.

[0044]

[Comparative Example 1]

Using a commercially available (Falcon(registered trademark) available from Becton, Dickinson and Company) γ -ray sterilized flat 24-well culture plate, after coating the plate with IV collagen, human fetal liver cells were cultured.

[0045]

Table 1 shows the gene expression levels of CYP3A4, CYP2D6 and CYP2C9 after three weeks of culture in the example 1 and the comparative example 1. The table shows values when each CYP expression level after three weeks in the comparative example 1 is 1. In the example 1, the expression levels were higher than in the comparative example 1 for any CYP, and the function was sustained for three weeks of culture.

[Table 1]

	Example 1	Comparative Example 1
CYP3A4	4.5	1.0
CYP2D6	8.5	1.0
CYP2C9	2.3	1.0

Industrial Applicability

[0046]

The present invention is applicable to a cell culture method that cultures cells isolated from tissue.

14
CLAIMS

1. A cell culture method comprising:

5 culturing undifferentiated cells in a layered state in a
partitioned micro-space and obtaining differentiated cells.

2. The cell culture method according to Claim 1, wherein the
undifferentiated cells are undifferentiated cells capable of
differentiating into any of liver parenchymal cells, intestinal
10 epithelium cells, nerve cells, myocardial cells, and vascular
endothelial cells.

3. The cell culture method according to Claim 1 or 2, wherein
the undifferentiated cells are human cells.
15

4. The cell culture method according to one of Claims 1 to 3,
wherein the partitioned micro-space is a micro-container in a cell
culture container having a plurality of micro-containers on a surface.

20 5. The cell culture method according to Claim 4, wherein the
micro-container has a bottom surface area of 9×10^{-4} mm² to 9×10^{-2}
mm², and a side wall with a height of 15 μ m to 300 μ m and a width of
3 μ m to 15 μ m.

25 6. The cell culture method according to Claim 4 or 5, wherein a
region where the micro-container is formed in the cell culture
container has transparency.

7. A screening method comprising:

30 placing a plurality of cells cultured by the culture method
according to one of Claims 1 to 6 and screening a compound.

8. A screening method comprising:

placing a plurality of cells cultured by the culture method

according to one of Claims 4 to 6 and screening a compound,

wherein the cell culture container includes a plurality of partitioned spots each made up of the plurality of micro-containers.