

THE USE OF SPIROCYCLIC COMPOUNDS AS TRYPTOPHAN HYDROXYLASE INHIBITORS

5

FIELD OF THE INVENTION

The present invention is directed to spirocyclic compounds which are inhibitors of tryptophan hydroxylase (TPH), particularly isoform 1 (TPH1), that are useful in the treatment of diseases or disorders associated with peripheral serotonin including, for example,
10 gastrointestinal, cardiovascular, pulmonary, inflammatory, metabolic, and low bone mass diseases, as well as serotonin syndrome, and cancer.

BACKGROUND OF THE INVENTION

Serotonin (5-hydroxytryptamine, 5-HT) is a neurotransmitter that modulates central and
15 peripheral functions by acting on neurons, smooth muscle, and other cell types. 5-HT is involved in the control and modulation of multiple physiological and psychological processes. In the central nervous system (CNS), 5-HT regulates mood, appetite, and other behavioral functions. In the GI system, 5-HT plays a general prokinetic role and is an important mediator of sensation (e.g., nausea and satiety) between the GI tract and the brain. Dysregulation of the
20 peripheral 5-HT signaling system has been reported to be involved in the etiology of several conditions (see for example: Mawe, G. M. & Hoffman, J. M. Serotonin Signalling In The Gut- functions, Dysfunctions And Therapeutic Targets. *Nature Reviews. Gastroenterology & Hepatology* **10**, 473–486 (2013); Gershon, M. D. 5-hydroxytryptamine (serotonin) In The Gastrointestinal Tract. *Current Opinion in Endocrinology, Diabetes, and Obesity* **20**, 14–21
25 (2013); Lesurtel, M., Soll, C., Graf, R. & Clavien, P.-A. Role of Serotonin In The Hepato- gastrointestinal Tract: An Old Molecule For New Perspectives. *Cellular And Molecular Life Sciences : CMLS* **65**, 940–52 (2008)). These include osteoporosis (e.g. Kode, A. *et al.* FOXO1 Orchestrates The Bone-suppressing Function Of Gut-derived Serotonin. *The Journal of Clinical Investigation* **122**, 3490–503 (2012); Yadav, V. K. *et al.* Pharmacological Inhibition Of Gut-
30 derived Serotonin Synthesis Is A Potential Bone Anabolic Treatment For Osteoporosis. *Nature Medicine* **16**, 308–12 (2010); Yadav, V. K. *et al.* Lrp5 Controls Bone Formation By Inhibiting

Serotonin Synthesis In The Duodenum. *Cell* **135**, 825–37 (2008)), cancer (e.g. Liang, C. *et al.* Serotonin Promotes The Proliferation Of Serum-deprived Hepatocellular Carcinoma Cells Via Upregulation Of FOXO3a. *Molecular Cancer* **12**, 14 (2013); Soll, C. *et al.* Serotonin Promotes Tumor Growth In Human Hepatocellular Cancer. *Hepatology* **51**, 1244–1254 (2010); Pai, V. P *et al.* Altered Serotonin Physiology In Human Breast Cancers Favors Paradoxical Growth And Cell Survival. *Breast Cancer Research : BCR* **11**, R81 (2009); Engelman, K., Lovenberg, W. & Sjoerdsma, A. Inhibition Of Serotonin Synthesis By Para-chlorophenylalanine In Patients With The Carcinoid Syndrome. *The New England Journal of Medicine* **277**, 1103–8 (1967)), cardiovascular (e.g. Robiolio, P. A. *et al.* Carcinoid Heart Disease : Correlation of High Serotonin Levels With Valvular Abnormalities Detected by Cardiac Catheterization and Echocardiography. *Circulation* **92**, 790–795 (1995).), diabetes (e.g. Sumara, G., Sumara, O., Kim, J. K. & Karsenty, G. Gut-derived Serotonin Is A Multifunctional Determinant To Fasting Adaptation. *Cell Metabolism* **16**, 588–600 (2012)), atherosclerosis (e.g. Ban, Y. *et al.* Impact Of Increased Plasma Serotonin Levels And Carotid Atherosclerosis On Vascular Dementia. *Atherosclerosis* **195**, 153–9 (2007)), as well as gastrointestinal (e.g. Manocha, M. & Khan, W. I. Serotonin and GI Disorders: An Update on Clinical and Experimental Studies. *Clinical and Translational Gastroenterology* **3**, e13 (2012); Ghia, J.-E. *et al.* Serotonin Has A Key Role In Pathogenesis Of Experimental Colitis. *Gastroenterology* **137**, 1649–60 (2009); Sikander, A., Rana, S. V. & Prasad, K. K. Role Of Serotonin In Gastrointestinal Motility And Irritable Bowel Syndrome. *Clinica Chimica Acta; International Journal of Clinical Chemistry* **403**, 47–55 (2009); Spiller, R. Recent Advances In Understanding The Role Of Serotonin In Gastrointestinal Motility In Functional Bowel Disorders: Alterations In 5-HT Signalling And Metabolism In Human Disease. *Neurogastroenterology and Motility: The Official Journal of The European Gastrointestinal Motility Society* **19 Suppl 2**, 25–31 (2007); Costedio, M. M., Hyman, N. & Mawe, G. M. Serotonin And Its Role In Colonic Function And In Gastrointestinal Disorders. *Diseases of the Colon and Rectum* **50**, 376–88 (2007); Gershon, M. D. & Tack, J. The Serotonin Signaling System: From Basic Understanding To Drug Development For Functional GI Disorders. *Gastroenterology* **132**, 397–414 (2007); Mawe, G. M., Coates, M. D. & Moses, P. L. Review Article: Intestinal Serotonin Signalling In Irritable Bowel Syndrome. *Alimentary Pharmacology & Therapeutics* **23**, 1067–76 (2006); Crowell, M. D. Role Of Serotonin In The Pathophysiology Of The Irritable Bowel Syndrome. *British Journal of Pharmacology* **141**,

1285–93 (2004)), pulmonary (e.g. Lau, W. K. W. *et al.* The Role Of Circulating Serotonin In The Development Of Chronic Obstructive Pulmonary Disease. *PloS One* **7**, e31617 (2012); Eggermayer, P., Town, G. I. & Peacock, A. J. Role Of Serotonin In The Pathogenesis Of Acute And Chronic Pulmonary Hypertension. *Thorax* **54**, 161–168 (1999)), inflammatory (e.g. 5 Margolis, K. G. *et al.* Pharmacological Reduction of Mucosal but Not Neuronal Serotonin Opposes Inflammation In Mouse Intestine. *Gut* doi:10.1136/gutjnl-2013-304901 (2013); Duerschmied, D. *et al.* Platelet Serotonin Promotes The Recruitment Of Neutrophils To Sites Of Acute Inflammation In Mice. *Blood* **121**, 1008–15 (2013); Li, N. *et al.* Serotonin Activates Dendritic Cell Function In The Context Of Gut Inflammation. *The American Journal of* 10 *Pathology* **178**, 662–71 (2011)), or liver diseases or disorders (e.g. Ebrahimkhani, M. R. *et al.* Stimulating Healthy Tissue Regeneration By Targeting The 5-HT_{2B} Receptor In Chronic Liver Disease. *Nature Medicine* **17**, 1668–73 (2011)). The large number of pharmaceutical agents that block or stimulate the various 5-HT receptors is also indicative of the wide range of medical disorders that have been associated with 5-HT dysregulation (see for example: Wacker, D. *et al.* 15 Structural Features For Functional Selectivity At Serotonin Receptors. *Science (New York, N.Y.)* **340**, 615–9 (2013)).

The rate-limiting step in 5-HT biosynthesis is the hydroxylation of tryptophan by dioxygen, which is catalyzed by tryptophan hydroxylase (TPH; EC 1.14.16.4) in the presence of the cofactor (6R)-L-erythro-5,6,7,8-tetrahydrobiopterin (BH₄). The resulting oxidized product, 20 5-hydroxytryptophan (5-HTT) is subsequently decarboxylated by an aromatic amino acid decarboxylase (AADC; EC 4.1.1.28) to produce 5-HT. Together with phenylalanine hydroxylase (PheOH) and tyrosine hydroxylase (TH), TPH belongs to the pterin-dependent aromatic amino acid hydroxylase family.

Two vertebrate isoforms of TPH, namely TPH1 and TPH2, have been identified. TPH1 25 is primarily expressed in the pineal gland and non-neuronal tissues, such as enterochromaffin (EC) cells located in the gastrointestinal (GI) tract. TPH2 (the dominant form in the brain) is expressed exclusively in neuronal cells, such as dorsal raphe or myenteric plexus cells. The peripheral and central systems involved in 5-HT biosynthesis are isolated, with 5-HT being unable to cross the blood-brain barrier. Therefore, the pharmacological effects of 5-HT can be 30 modulated by agents affecting TPH in the periphery, mainly TPH1 in the gut.


A small number of phenylalanine-derived TPH1 inhibitors are known. One example, p-chlorophenylalanine (pCPA), a very weak and unselective irreversible inhibitor of TPH, has proven effective in treating chemotherapy-induced emesis, as well as diarrhea, in carcinoid tumor patients. However, pCPA is distributed centrally and, as a result, its administration has been linked to the onset of depression and other alterations of CNS functions in patients and animals. p-Ethynyl phenylalanine is a more selective and more potent TPH inhibitor than pCPA (Stokes, A. H. *et al.* p-Ethynylphenylalanine: A Potent Inhibitor Of Tryptophan Hydroxylase. *Journal of Neurochemistry* **74**, 2067–73 (2000), but also affects central 5-HT production and, like pCPA, is believed to irreversibly interfere with the synthesis of TPH (and possibly other proteins).

More recently, bulkier phenylalanine-derived TPH inhibitors have been reported to reduce intestinal 5-HT concentration without affecting brain 5-HT levels (Zhong, H. *et al.* Molecular dynamics simulation of tryptophan hydroxylase-1: binding modes and free energy analysis to phenylalanine derivative inhibitors. *International Journal of Molecular Sciences* **14**, 9947–62 (2013); Ouyang, L. *et al.* Combined Structure-Based Pharmacophore and 3D-QSAR Studies on Phenylalanine Series Compounds as TPH1 Inhibitors. *International Journal of Molecular Sciences* **13**, 5348–63 (2012); Camilleri, M. LX-1031, A Tryptophan 5-hydroxylase Inhibitor, And Its Potential In Chronic Diarrhea Associated With Increased Serotonin. *Neurogastroenterology and Motility: The Official Journal of The European Gastrointestinal Motility Society* **23**, 193–200 (2011); Cianchetta, G. *et al.* Mechanism of Inhibition of Novel Tryptophan Hydroxylase Inhibitors Revealed by Co-crystal Structures and Kinetic Analysis. *Current chemical genomics* **4**, 19–26 (2010); Jin, H. *et al.* Substituted 3-(4-(1,3,5-triazin-2-yl)-phenyl)-2-aminopropanoic Acids As Novel Tryptophan Hydroxylase Inhibitors. *Bioorganic & Medicinal Chemistry Letters* **19**, 5229–32 (2009); Shi, Z.-C. *et al.* Modulation Of Peripheral Serotonin Levels By Novel Tryptophan Hydroxylase Inhibitors For The Potential Treatment Of Functional Gastrointestinal Disorders. *Journal of medicinal chemistry* **51**, 3684–7 (2008); Liu, Q. *et al.* Discovery And Characterization of Novel Tryptophan Hydroxylase Inhibitors That Selectively Inhibit Serotonin Synthesis In The Gastrointestinal Tract. *The Journal of Pharmacology and Experimental Therapeutics* **325**, 47–55 (2008)).

There is a current need to selectively reduce intestinal 5-HT levels as a means for treating and preventing 5-HT-associated diseases. The TPH1 inhibitors described herein are intended to address this need.

5 SUMMARY OF THE INVENTION

The present invention relates to a TPH-inhibiting compound of Formula I:



or a pharmaceutically acceptable salt thereof, wherein constituent variables are defined herein.

10 The present invention further relates to a pharmaceutical composition comprising a TPH-inhibiting compound of the invention, or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier.

The present invention further relates to a method of inhibiting TPH, such as TPH1, by contacting the TPH enzyme with a compound of Formula I, or a pharmaceutically acceptable salt thereof.

15 The present invention further relates to a method of lowering peripheral serotonin in a patient comprising administering to the patient an effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof.

20 The present invention further relates to a method of treating or preventing a disease in a patient comprising administering to the patient a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof.


The present invention further relates to a compound of Formula I, or a pharmaceutically acceptable salt thereof, for use in the treatment or prevention of disease in a patient.

The present invention further relates to use of a compound of Formula I or a pharmaceutically acceptable salt thereof, for the preparation of a medicament for the treatment or prevention of disease in a patient.

5 DETAILED DESCRIPTION

Compounds

The present invention relates to a TPH-inhibiting compound of Formula I:



10 or a pharmaceutically acceptable salt thereof, wherein:

Ring A is C₃₋₁₀ cycloalkyl, C₆₋₁₀ aryl, 4 to 10-membered heterocycloalkyl, or 5 to 10-membered heteroaryl;

L is O or NR⁴;

W is N or CR⁵;

15 X is N or CR⁶;

Y is N or CR⁷;

wherein only one of X and Y is N;

R¹ is H, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, 4-10 membered heterocycloalkyl, 5-10 membered heteroaryl, phenyl, -(CR⁸R⁹)_pOC(O)R¹⁰, -(CR⁸R⁹)_pNR¹¹R¹², or -(CR⁸R⁹)_pC(O)NR¹¹R¹², wherein
20 said C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, 4-10 membered heterocycloalkyl, 5-10 membered heteroaryl, and phenyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from F, Cl, Br, CN, C₁₋₄ alkyl, and C₁₋₄ haloalkyl;

R² and R³ are each independently selected from H, C₁₋₄ alkyl, and C₁₋₄ haloalkyl;

R⁴ is H or C₁₋₄ alkyl;

R⁵ and R⁶ are each independently selected from H, halo, and C₁₋₄ alkyl;

R⁷ is H, C₁₋₄ alkyl, C₂₋₆ alkenyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, (5-10 membered heteroaryl)-C₁₋₄ alkyl, NR¹³R¹⁴, OR¹⁵, C(O)R¹⁶,
5 S(O)_qR¹⁷, wherein said C₁₋₄ alkyl, C₂₋₆ alkenyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, and (5-10 membered heteroaryl)-C₁₋₄ alkyl are each optionally substituted by 1, 2, or 3 substituents selected from halo, C₁₋₄ alkyl, C₂₋₆ alkenyl, amino, C₁₋₄ alkylamino, C₂₋₈ dialkylamino, hydroxy, and C₁₋₄ alkoxy;

10 R⁸ and R⁹ are each independently selected from H and C₁₋₄ alkyl;

R¹⁰ is C₁₋₆ alkyl optionally substituted by 1, 2 or 3 substituents independently selected from C₁₋₆ haloalkyl, C₃₋₁₀ cycloalkyl, OR^a, and NR^cR^d;

R¹¹ and R¹² are each independently selected from H and C₁₋₆ alkyl;

R¹³ is H or C₁₋₄ alkyl;

15 R¹⁴ is H, C₁₋₄ alkyl, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, or (5-10 membered heteroaryl)-C₁₋₄ alkyl, C(O)R^{b1}, C(O)OR^{a1}, C(O)NR^{c1}R^{d1}, S(O)R^{b1}, S(O)₂R^{b1}, or S(O)₂NR^{c1}R^{d1}, wherein said C₁₋₄ alkyl, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, and (5-10 membered heteroaryl)-C₁₋₄ alkyl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, C₁₋₄ alkyl, C₁₋₄ haloalkyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1},
20 S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};

25 or R¹³ and R¹⁴ together with the N atom to which they are attached form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, halo, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1},
30 OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}C(O)OR^{a1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, and S(O)₂NR^{c1}R^{d1},

wherein said C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, and 5-6 membered heteroaryl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}C(O)OR^{a1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, and S(O)₂NR^{c1}R^{d1};

R¹⁵ is H, C₁₋₄ alkyl, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, or (5-10 membered heteroaryl)-C₁₋₄ alkyl, wherein said C₁₋₄ alkyl, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, and (5-10 membered heteroaryl)-C₁₋₄ alkyl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}C(O)OR^{a1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, and S(O)₂NR^{c1}R^{d1};

R¹⁶ is C₁₋₄ alkyl or NR^{18a}R^{18b} wherein said C₁₋₄ alkyl is optionally substituted by 1, 2, or 3 substituents independently selected from halo, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}C(O)OR^{a1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, and S(O)₂NR^{c1}R^{d1};

R¹⁷ is C₁₋₄ alkyl, NR^{18a}R^{18b}, or OR^{18c}, wherein said C₁₋₄ alkyl is optionally substituted by 1, 2, or 3 substituents independently selected from halo, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}C(O)OR^{a1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, and S(O)₂NR^{c1}R^{d1};

R^{18a} and R^{18b} are each independently selected from H and C₁₋₄ alkyl wherein said C₁₋₄ alkyl is optionally substituted by 1, 2, or 3 substituents independently selected from halo, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1},

R^B is H, Cy^2 , halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$, $C(O)OR^{a3}$, $OC(O)R^{b3}$, $OC(O)NR^{c3}R^{d3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)OR^{a3}$, $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}S(O)R^{b3}$, $NR^{c3}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, $S(O)R^{b3}$, $S(O)NR^{c3}R^{d3}$, $S(O)_2R^{b3}$, or $S(O)_2NR^{c3}R^{d3}$, wherein said C_{1-6} alkyl and C_{2-6} alkenyl are each optionally

5 substituted with 1, 2, 3, 4, or 5 substituents independently selected from Cy^2 , halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$, $C(O)OR^{a3}$, $OC(O)R^{b3}$, $OC(O)NR^{c3}R^{d3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)OR^{a3}$, $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}S(O)R^{b3}$, $NR^{c3}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, $S(O)R^{b3}$, $S(O)NR^{c3}R^{d3}$, $S(O)_2R^{b3}$, and $S(O)_2NR^{c3}R^{d3}$;

R^C and R^D are each independently selected from H, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, NO_2 , OR^{a4} , SR^{a4} , $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$, $NR^{c4}S(O)R^{b4}$, $NR^{c4}S(O)_2R^{b4}$, $NR^{c4}S(O)_2NR^{c4}R^{d4}$, $S(O)R^{b4}$, $S(O)NR^{c4}R^{d4}$, $S(O)_2R^{b4}$, and $S(O)_2NR^{c4}R^{d4}$; wherein said C_{1-6} alkyl and C_{2-6} alkenyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered

10 heterocycloalkyl, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, NO_2 , OR^{a4} , SR^{a4} , $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$, $NR^{c4}S(O)R^{b4}$, $NR^{c4}S(O)_2R^{b4}$, $NR^{c4}S(O)_2NR^{c4}R^{d4}$, $S(O)R^{b4}$, $S(O)NR^{c4}R^{d4}$, $S(O)_2R^{b4}$, and $S(O)_2NR^{c4}R^{d4}$;

15

Cy^1 and Cy^2 are each independently selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, each of which is optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy} ;

20

each R^{Cy} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)OR^{a5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}S(O)R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, and $S(O)_2NR^{c5}R^{d5}$, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C_{1-6} alkyl, CN, NO_2 , OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)OR^{a5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$,

25

30

substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₄ alkyl, halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

5 or any R^c and R^d together with the N atom to which they are attached form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6},
10 wherein said C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, and 5-6 membered heteroaryl are optionally substituted by 1, 2, or 3 substituents independently selected from halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};
15 NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

or any R^{c1} and R^{d1} together with the N atom to which they are attached form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6},
20 wherein said C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, and 5-6 membered heteroaryl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};
25 NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

or any R^{c2} and R^{d2} together with the N atom to which they are attached form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, and 5-6 membered heteroaryl, C₁₋₆ haloalkyl, halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6},
30 C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6},

aryl, 5-6 membered heteroaryl, C₁₋₆ haloalkyl, halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6}, wherein said C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀

aryl, and 5-6 membered heteroaryl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

each R^{a6}, R^{b6}, R^{c6}, and R^{d6} is independently selected from H, C₁₋₄ alkyl, C₂₋₄ alkenyl, C₃₋₇ cycloalkyl, phenyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl, wherein said C₁₋₄ alkyl, C₂₋₄ alkenyl, C₃₋₇ cycloalkyl, phenyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl are each optionally substituted by 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ alkylamino, and di(C₁₋₄ alkyl)amino;


n is 1 or 2;

p is 1, 2, or 3; and

q is 1 or 2;

wherein any aforementioned 4-10 or 4-7 membered heterocycloalkyl group optionally comprises 1, 2, or 3 oxo substituents, wherein each oxo substituent that is present is substituted on a ring-forming carbon, nitrogen, or sulfur atom of the 4-10 or 4-7 membered heterocycloalkyl group.

In some embodiments, the present invention relates to a TPH-inhibiting compound of Formula I:



or a pharmaceutically acceptable salt thereof, wherein:

Ring A is C₃₋₁₀ cycloalkyl, C₆₋₁₀ aryl, 4 to 10-membered heterocycloalkyl, or 5 to 10-membered heteroaryl;

L is O or NR⁴;

5 W is N or CR⁵;

X is N or CR⁶;

Y is N or CR⁷;

wherein only one of X and Y is N;

10 R¹ is H, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, phenyl, -(CR⁸R⁹)_pOC(O)R¹⁰, -(CR⁸R⁹)_pNR¹¹R¹², or -(CR⁸R⁹)_pC(O)NR¹¹R¹², wherein said C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, and phenyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from F, Cl, Br, CN, C₁₋₄ alkyl, and C₁₋₄ haloalkyl;

R² and R³ are each independently selected from H, C₁₋₄ alkyl, and C₁₋₄ haloalkyl;

R⁴ is H or C₁₋₄ alkyl;

15 R⁵ and R⁶ are each independently selected from H, halo, and C₁₋₄ alkyl;

R⁷ is H, C₁₋₄ alkyl, C₂₋₆ alkenyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, (5-10 membered heteroaryl)-C₁₋₄ alkyl, NR¹³R¹⁴, OR¹⁵, C(O)R¹⁶, S(O)_qR¹⁷, wherein said C₁₋₄ alkyl, C₂₋₆ alkenyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, and (5-10 membered heteroaryl)-C₁₋₄ alkyl are each optionally substituted by 1, 2, or 3 substituents selected from halo, C₁₋₄ alkyl, C₂₋₆ alkenyl, amino, C₁₋₄ alkylamino, C₂₋₈ dialkylamino, hydroxy, and C₁₋₄ alkoxy;

R⁸ and R⁹ are each independently selected from H and C₁₋₄ alkyl;

25 R¹⁰ is C₁₋₆ alkyl optionally substituted by 1, 2 or 3 substituents independently selected from C₁₋₆ haloalkyl, C₃₋₁₀ cycloalkyl, OR^a, and NR^cR^d;

R¹¹ and R¹² are each independently selected from H and C₁₋₆ alkyl;

R¹³ is H or C₁₋₄ alkyl;

30 R¹⁴ is H, C₁₋₄ alkyl, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, or (5-10 membered heteroaryl)-C₁₋₄ alkyl, C(O)R^{b1}, C(O)OR^{a1},

C(O)NR^{c1}R^{d1}, S(O)R^{b1}, S(O)₂R^{b1}, or S(O)₂NR^{c1}R^{d1}, wherein said C₁₋₄ alkyl, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, and (5-10 membered heteroaryl)-C₁₋₄ alkyl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, C₁₋₄ alkyl, C₁₋₄ haloalkyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};

or R¹³ and R¹⁴ together with the N atom to which they are attached form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, halo, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}C(O)OR^{a1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, and S(O)₂NR^{c1}R^{d1}, wherein said C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, and 5-6 membered heteroaryl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}C(O)OR^{a1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, and S(O)₂NR^{c1}R^{d1};

R¹⁵ is H, C₁₋₄ alkyl, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, or (5-10 membered heteroaryl)-C₁₋₄ alkyl, wherein said C₁₋₄ alkyl, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, and (5-10 membered heteroaryl)-C₁₋₄ alkyl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}C(O)OR^{a1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, and S(O)₂NR^{c1}R^{d1};

R¹⁶ is C₁₋₄ alkyl or NR^{18a}R^{18b} wherein said C₁₋₄ alkyl is optionally substituted by 1, 2, or 3 substituents independently selected from halo, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl,

C₆₋₁₀ aryl, 5-6 membered heteroaryl, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}C(O)OR^{a1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, and S(O)₂NR^{c1}R^{d1};

R¹⁷ is C₁₋₄ alkyl, NR^{18a}R^{18b}, or OR^{18c}, wherein said C₁₋₄ alkyl is optionally substituted by

1, 2, or 3 substituents independently selected from halo, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}C(O)OR^{a1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, and S(O)₂NR^{c1}R^{d1};

R^{18a} and R^{18b} are each independently selected from H and C₁₋₄ alkyl wherein said C₁₋₄ alkyl is optionally substituted by 1, 2, or 3 substituents independently selected from halo, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}C(O)OR^{a1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, and S(O)₂NR^{c1}R^{d1};

or R^{18a} and R^{18b} together with the N atom to which they are attached form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, halo, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}C(O)OR^{a1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, and S(O)₂NR^{c1}R^{d1}, wherein said C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, and 5-6 membered heteroaryl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, CN, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}C(O)OR^{a1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, and S(O)₂NR^{c1}R^{d1};

R^{18c} is H, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₇ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, or (5-10 membered heteroaryl)-C₁₋₄ alkyl, wherein said C₁₋₆ alkyl, C₃₋₇ cycloalkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, C₆₋₁₀ aryl, C₆₋₁₀ aryl-C₁₋₄ alkyl, 4-10 membered heterocycloalkyl, (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, 5-10 membered heteroaryl, and

(5-10 membered heteroaryl)-C₁₋₄ alkyl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, C₁₋₄ alkyl, C₁₋₄ haloalkyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};

R^A is H, Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, or S(O)₂NR^{c2}R^{d2}, wherein said C₁₋₆ alkyl and C₂₋₆ alkenyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, and S(O)₂NR^{c2}R^{d2};

R^B is H, Cy², halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)OR^{a3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}S(O)R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, or S(O)₂NR^{c3}R^{d3}, wherein said C₁₋₆ alkyl and C₂₋₆ alkenyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from Cy², halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)OR^{a3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}S(O)R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, and S(O)₂NR^{c3}R^{d3};

R^C and R^D are each independently selected from H, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, NO₂, OR^{a4}, SR^{a4}, C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)OR^{a4}, OC(O)R^{b4}, OC(O)NR^{c4}R^{d4}, NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4}, NR^{c4}C(O)NR^{c4}R^{d4}, NR^{c4}S(O)R^{b4}, NR^{c4}S(O)₂R^{b4}, NR^{c4}S(O)₂NR^{c4}R^{d4}, S(O)R^{b4}, S(O)NR^{c4}R^{d4}, S(O)₂R^{b4}, and S(O)₂NR^{c4}R^{d4}; wherein said C₁₋₆ alkyl and C₂₋₆ alkenyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, NO₂, OR^{a4}, SR^{a4}, C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)OR^{a4}, OC(O)R^{b4}, OC(O)NR^{c4}R^{d4}, NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4}, NR^{c4}C(O)NR^{c4}R^{d4}, NR^{c4}S(O)R^{b4}, NR^{c4}S(O)₂R^{b4}, NR^{c4}S(O)₂NR^{c4}R^{d4}, S(O)R^{b4}, S(O)NR^{c4}R^{d4}, S(O)₂R^{b4}, and S(O)₂NR^{c4}R^{d4};

Cy¹ and Cy² are each independently selected from C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, each of which is optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from RCy;

each R^{Cy} is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₆₋₁₀

5 aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO₂, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)OR^{a5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}S(O)R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, and S(O)₂NR^{c5}R^{d5}, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, CN, NO₂, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)OR^{a5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}S(O)R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, and S(O)₂NR^{c5}R^{d5};

15 each R^a, R^{a1}, R^{a2}, R^{a3}, R^{a4}, and R^{a5} is independently selected from H, C₁₋₆ alkyl, C₁₋₄ haloalkyl, C₂₋₆ alkenyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₄ alkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, (5-10 membered heteroaryl)-C₁₋₄ alkyl, or (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₄ alkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, (5-10 membered heteroaryl)-C₁₋₄ alkyl, and (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₄ alkyl, halo, CN, OR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

25 each R^{b1}, R^{b2}, R^{b3}, R^{b4}, and R^{b5} is independently selected from H, C₁₋₆ alkyl, C₁₋₄ haloalkyl, C₂₋₆ alkenyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₄ alkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, (5-10 membered heteroaryl)-C₁₋₄ alkyl, or (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₄ alkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, (5-10 membered heteroaryl)-C₁₋₄ alkyl, and (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₄ alkyl, halo, CN, OR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

membered heterocycloalkyl)-C₁₋₄ alkyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₄ alkyl, halo, CN, OR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

each R^c, R^d, R^{c1}, R^{d1}, R^{c2}, R^{d2}, R^{c3}, R^{d3}, R^{c4}, R^{d4}, R^{c5}, and R^{d5} is independently selected from H, C₁₋₆ alkyl, C₁₋₄ haloalkyl, C₂₋₆ alkenyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₄ alkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, (5-10 membered heteroaryl)-C₁₋₄ alkyl, or (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₄ alkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, (5-10 membered heteroaryl)-C₁₋₄ alkyl, and (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₄ alkyl, halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

or any R^c and R^d together with the N atom to which they are attached form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6}, wherein said C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, and 5-6 membered heteroaryl are optionally substituted by 1, 2, or 3 substituents independently selected from halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

or any R^{c1} and R^{d1} together with the N atom to which they are attached form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6},

OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6}, wherein said C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, and 5-6 membered heteroaryl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

or any R^{c2} and R^{d2} together with the N atom to which they are attached form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group optionally substituted with 1, 2, or 3 substituents

independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, and 5-6 membered heteroaryl, C₁₋₆ haloalkyl, halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6}, wherein said C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, and 5-6 membered heteroaryl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

or any R^{c3} and R^{d3} together with the N atom to which they are attached form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group optionally substituted with 1, 2, or 3 substituents

independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, C₁₋₆ haloalkyl, halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6}, wherein said C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, and 5-6 membered heteroaryl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

or any R^{c4} and R^{d4} together with the N atom to which they are attached form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group optionally substituted with 1, 2, or 3 substituents

independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, C₁₋₆ haloalkyl, halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6}, wherein said C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, and 5-6 membered heteroaryl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

or any R^{c5} and R^{d5} together with the N atom to which they are attached form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, 5-6 membered heteroaryl, C₁₋₆ haloalkyl, halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6}, wherein said C₁₋₆ alkyl, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl, C₆₋₁₀ aryl, and 5-6 membered heteroaryl are each optionally substituted by 1, 2, or 3 substituents independently selected from halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

each R^{a6}, R^{b6}, R^{c6}, and R^{d6} is independently selected from H, C₁₋₄ alkyl, C₂₋₄ alkenyl, C₃₋₇ cycloalkyl, phenyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl, wherein said C₁₋₄ alkyl, C₂₋₄ alkenyl, C₃₋₇ cycloalkyl, phenyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl are each optionally substituted by 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ alkylamino, and di(C₁₋₄ alkyl)amino;

n is 1 or 2;

p is 1, 2, or 3; and

q is 1 or 2;

wherein any aforementioned 4-10 or 4-7 membered heterocycloalkyl group optionally comprises 1, 2, or 3 oxo substituents, wherein each oxo substituent that is present is substituted

on a ring-forming carbon, nitrogen, or sulfur atom of the 4-10 or 4-7 membered heterocycloalkyl group.

In some embodiments, L is O.

In some embodiments, L is NR^4 .

5 In some embodiments, W is CR^5 ; X is N; and Y is CR^7 .

In some embodiments, W is N; X is N; and Y is CR^7 .

In some embodiments, W is CR^5 ; X is CR^6 ; and Y is N.

In some embodiments, W is CR^5 ; X is CR^6 ; and Y is CR^7 .

In some embodiments, W is N; X is CR^6 ; and Y is CR^7 .

10 In some embodiments, R^2 is H and R^3 is H.

In some embodiments, R^2 is H and R^3 is C_{1-4} alkyl.

In some embodiments, R^2 is H and R^3 is methyl.

In some embodiments, R^2 is H and R^3 is C_{1-4} haloalkyl.

In some embodiments, R^2 is H and R^3 is trifluoromethyl.

15 In some embodiments, n is 1.

In some embodiments, n is 2.

In some embodiments, R^1 is H.

In some embodiments, R^1 is C_{1-10} alkyl, C_{3-10} cycloalkyl, phenyl, $-(\text{CR}^8\text{R}^9)_p\text{OC}(\text{O})\text{R}^{10}$, $-(\text{CR}^8\text{R}^9)_p\text{NR}^{11}\text{R}^{12}$, or $-(\text{CR}^8\text{R}^9)_p\text{C}(\text{O})\text{NR}^{11}\text{R}^{12}$, wherein said C_{1-10} alkyl, C_{3-10} cycloalkyl, and
20 phenyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from F, Cl, Br, CN, C_{1-4} alkyl, and C_{1-4} haloalkyl.

In some embodiments, R^1 is C_{1-10} alkyl.

In some embodiments, R^1 is ethyl.

In some embodiments, R^4 is H.

25 In some embodiments, R^5 is H.

In some embodiments, R^6 is H.

In some embodiments, R^7 is other than H.

In some embodiments, R^7 is C_{1-4} alkyl, $\text{NR}^{13}\text{R}^{14}$, or OR^{15} .

In some embodiments, R^7 is $\text{NR}^{13}\text{R}^{14}$.

30 In some embodiments, R^7 is NH_2 .

In some embodiments, R^7 is C_{1-4} alkyl.

- In some embodiments, R^7 is OR^{15} .
- In some embodiments, Ring A is C_{3-10} cycloalkyl.
- In some embodiments, Ring A is C_{6-10} aryl.
- In some embodiments, Ring A is phenyl.
- 5 In some embodiments, Ring A is 4 to 10-membered heterocycloalkyl.
- In some embodiments, Ring A is phenyl, adamantanyl, naphthyl, 1,2,3,4-tetrahydroquinoxaliny, 3,4-dihydroquinazoliny, 1,2,3,4-tetrahydroquinazoliny, or pyridyl.
- In some embodiments, Ring A is 5 to 10-membered heteroaryl.
- In some embodiments, at least one of R^A , R^B , R^C , and R^D is other than hydrogen.
- 10 In some embodiments, at least two of R^A , R^B , R^C , and R^D are other than hydrogen.
- In some embodiments, R^A is Cy^1 .
- In some embodiments, R^A is C_{6-10} aryl or 5-10 membered heteroaryl, each of which is optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy} .
- In some embodiments, R^A is 5-10 membered heteroaryl optionally substituted by 1, 2, 3,
- 15 4, or 5 substituents independently selected from R^{Cy} .
- In some embodiments, R^A is 5 to 6-membered heteroaryl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy} .
- In some embodiments, R^A is pyrazolyl which is optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy} .
- 20 In some embodiments, R^A is 3-methyl-1H-pyrazol-1-yl.
- In some embodiments, R^A is C_{6-10} aryl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy} .
- In some embodiments, R^A is phenyl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy} .
- 25 In some embodiments, R^B is H.
- In some embodiments, R^B is Cy^2 , halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$, $C(O)OR^{a3}$, $OC(O)R^{b3}$, $OC(O)NR^{c3}R^{d3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)OR^{a3}$, $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}S(O)R^{b3}$, $NR^{c3}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, $S(O)R^{b3}$, $S(O)NR^{c3}R^{d3}$, $S(O)_2R^{b3}$, and $S(O)_2NR^{c3}R^{d3}$, wherein said C_{1-6} alkyl
- 30 and C_{2-6} alkenyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from Cy^2 , halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b3}$,

C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)OR^{a3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}S(O)R^{b3}, NR^{c1}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, and S(O)₂NR^{c3}R^{d3}.

In some embodiments, R^B is Cy².

5 In some embodiments, R^B is C₆₋₁₀ aryl or 5-10 membered heteroaryl, each of which is optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy}.

In some embodiments, R^B is halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)OR^{a3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}S(O)R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, and S(O)₂NR^{c3}R^{d3}, wherein said C₁₋₆ alkyl and C₂₋₆ alkenyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from Cy², halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)OR^{a3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}S(O)R^{b3}, NR^{c1}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, and S(O)₂NR^{c3}R^{d3}.

In some embodiments, R^B is halo.

In some embodiments, R^C is H.


In some embodiments, R^C is halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, NO₂, OR^{a4}, SR^{a4}, C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)OR^{a4}, OC(O)R^{b4}, OC(O)NR^{c4}R^{d4}, NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4}, NR^{c4}C(O)NR^{c4}R^{d4}, NR^{c4}S(O)R^{b4}, NR^{c4}S(O)₂R^{b4}, NR^{c4}S(O)₂NR^{c4}R^{d4}, S(O)R^{b4}, S(O)NR^{c4}R^{d4}, S(O)₂R^{b4}, and S(O)₂NR^{c4}R^{d4}; wherein said C₁₋₆ alkyl and C₂₋₆ alkenyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, NO₂, OR^{a4}, SR^{a4}, C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)OR^{a4}, OC(O)R^{b4}, OC(O)NR^{c4}R^{d4}, NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4}, NR^{c4}C(O)NR^{c4}R^{d4}, NR^{c4}S(O)R^{b4}, NR^{c4}S(O)₂R^{b4}, NR^{c4}S(O)₂NR^{c4}R^{d4}, S(O)R^{b4}, S(O)NR^{c4}R^{d4}, S(O)₂R^{b4}, and S(O)₂NR^{c4}R^{d4}.

In some embodiments, R^D is H.

In some embodiments, R^D is halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, NO₂, OR^{a4}, SR^{a4}, C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)OR^{a4}, OC(O)R^{b4}, OC(O)NR^{c4}R^{d4}, NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4}, NR^{c4}C(O)NR^{c4}R^{d4}, NR^{c4}S(O)R^{b4}, NR^{c4}S(O)₂R^{b4}, NR^{c4}S(O)₂NR^{c4}R^{d4}, S(O)R^{b4}, S(O)NR^{c4}R^{d4}, S(O)₂R^{b4}, and S(O)₂NR^{c4}R^{d4}; wherein said C₁₋₆ alkyl and C₂₋₆ alkenyl are each


optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, NO₂, OR^{a4}, SR^{a4}, C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)OR^{a4}, OC(O)R^{b4}, OC(O)NR^{c4}R^{d4}, NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4}, NR^{c4}C(O)NR^{c4}R^{d4}, NR^{c4}S(O)R^{b4},
5 NR^{c4}S(O)₂R^{b4}, NR^{c4}S(O)₂NR^{c4}R^{d4}, S(O)R^{b4}, S(O)NR^{c4}R^{d4}, S(O)₂R^{b4}, and S(O)₂NR^{c4}R^{d4}.

In some embodiments, the compounds of the invention have Formula IIa:




IIa.

In some embodiments, the compounds of the invention have Formula IIb:




IIb.

In some embodiments, the compounds of the invention have Formula IIc:




IIc.

In some embodiments, the compounds of the invention have Formula IId:



IId.

In some embodiments, the compounds of the invention have Formula IIe:



Ile.

In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or Ile, L is O.

5 In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or Ile, L is NR₄.

In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or Ile, R³ is H.

10 In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or Ile, R² is CF₃ and R³ is H.

In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or Ile, R¹ is H or C₁₋₁₀ alkyl.

In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or Ile, R^A is Cy¹.

15 In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or Ile, R^A is C₆₋₁₀ aryl or 5-10 membered heteroaryl, each of which is optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy}.

20 In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or Ile, R^A is 5-10 membered heteroaryl optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy}.

In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or Ile, R^A is 5 to 6-membered heteroaryl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.

25 In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or Ile, R^A is C₆₋₁₀ aryl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.

In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or Ile, R^A is phenyl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.

30 In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or Ile, R^B is Cy².

In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or IIe, R^B is H, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, OR^{a3} , $C(O)NR^{c3}R^{d3}$, or $C(O)OR^{a3}$, wherein said C_{1-6} alkyl and C_{2-6} alkenyl are each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} haloalkyl, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$, $C(O)OR^{a3}$, $OC(O)R^{b3}$, $OC(O)NR^{c3}R^{d3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)OR^{a3}$, $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}S(O)R^{b3}$, $NR^{c1}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, $S(O)R^{b3}$, $S(O)NR^{c3}R^{d3}$, $S(O)_2R^{b3}$, and $S(O)_2NR^{c3}R^{d3}$.


In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or IIe, R^C is H.


10 In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or IIe, R^D is H.

In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or IIe, R^5 is H.

15 In some embodiments, where the compounds of the invention have Formula IIa, IIb, IIc, IId, or IIe, R^6 is H.

In some embodiments, the compounds of the invention have Formula IIIa or IIIb:





IIIb.

In some embodiments, where the compounds of the invention have Formula IIIa or IIIb, R² is CF₃.

5 In some embodiments, where the compounds of the invention have Formula IIIa or IIIb, R¹ is H or C₁₋₁₀ alkyl.

In some embodiments, where the compounds of the invention have Formula IIIa or IIIb, R^A is Cy¹.

10 In some embodiments, where the compounds of the invention have Formula IIIa or IIIb, R^A is C₆₋₁₀ aryl or 5-10 membered heteroaryl, each of which is optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy}.

In some embodiments, where the compounds of the invention have Formula IIIa or IIIb, R^A is 5-10 membered heteroaryl optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy}.

15 In some embodiments, where the compounds of the invention have Formula IIIa or IIIb, R^A is 5 to 6-membered heteroaryl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.

In some embodiments, where the compounds of the invention have Formula IIIa or IIIb, R^A is C₆₋₁₀ aryl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.

20 In some embodiments, where the compounds of the invention have Formula IIIa or IIIb, R^A is phenyl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.


In some embodiments, where the compounds of the invention have Formula IIIa or IIIb, R^B is Cy².

In some embodiments, where the compounds of the invention have Formula IIIa or IIIb, R^B is H, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, OR^{a3} , $C(O)NR^{c3}R^{d3}$, or $C(O)OR^{a3}$, wherein said C_{1-6} alkyl and C_{2-6} alkenyl are each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} haloalkyl, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$, $C(O)OR^{a3}$, $OC(O)R^{b3}$, $OC(O)NR^{c3}R^{d3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)OR^{a3}$, $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}S(O)R^{b3}$, $NR^{c1}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, $S(O)R^{b3}$, $S(O)NR^{c3}R^{d3}$, $S(O)_2R^{b3}$, and $S(O)_2NR^{c3}R^{d3}$.

In some embodiments, where the compounds of the invention have Formula IIIa or IIIb, R^C is H.

In some embodiments, where the compounds of the invention have Formula IIIa or IIIb, R^D is H.

In some embodiments, the compounds of the invention have Formula IV:



In some embodiments, where the compounds of the invention have Formula IV, R^2 is CF_3 .

In some embodiments, where the compounds of the invention have Formula IV, R^1 is H or C_{1-10} alkyl.

In some embodiments, where the compounds of the invention have Formula IV, R^A is Cy^1 .

In some embodiments, where the compounds of the invention have Formula IV, R^A is C_{6-10} aryl or 5-10 membered heteroaryl, each of which is optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy} .

In some embodiments, where the compounds of the invention have Formula IV, R^A is 5-10 membered heteroaryl optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy}.

In some embodiments, where the compounds of the invention have Formula IV, R^A is 5 to 6-membered heteroaryl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.

In some embodiments, where the compounds of the invention have Formula IV, R^A is C₆₋₁₀ aryl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.

In some embodiments, where the compounds of the invention have Formula IV, R^A is phenyl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.


In some embodiments, where the compounds of the invention have Formula IV, R^B is Cy².

In some embodiments, where the compounds of the invention have Formula IV, R^B is H, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, OR^{a3}, C(O)NR^{c3}R^{d3}, or C(O)OR^{a3}, wherein said C₁₋₆ alkyl and C₂₋₆ alkenyl are each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ haloalkyl, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)OR^{a3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}S(O)R^{b3}, NR^{c1}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, and S(O)₂NR^{c3}R^{d3}.

In some embodiments, where the compounds of the invention have Formula IV, R^C is H.

In some embodiments, where the compounds of the invention have Formula IV, R^D is H.

In some embodiments, the compounds of the invention have Formula Va:



Va.

In some embodiments, where the compounds of the invention have Formula Va, R² is CF₃.

5 In some embodiments, where the compounds of the invention have Formula Va, R¹ is H or C₁₋₁₀ alkyl.

In some embodiments, where the compounds of the invention have Formula Va, R^A is Cy¹.

10 In some embodiments, where the compounds of the invention have Formula Va, R^A is C₆₋₁₀ aryl or 5-10 membered heteroaryl, each of which is optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy}.

In some embodiments, where the compounds of the invention have Formula Va, R^A is 5-10 membered heteroaryl optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy}.

15 In some embodiments, where the compounds of the invention have Formula Va, R^A is 5 to 6-membered heteroaryl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.


In some embodiments, where the compounds of the invention have Formula Va, R^A is C₆₋₁₀ aryl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.

20 In some embodiments, where the compounds of the invention have Formula Va, R^A is phenyl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.

In some embodiments, where the compounds of the invention have Formula Va, R^B is Cy².

25 In some embodiments, where the compounds of the invention have Formula Va, R^B is H, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, OR^{a3}, C(O)NR^{c3}R^{d3}, or C(O)OR^{a3}, wherein said C₁₋₆ alkyl and C₂₋₆ alkenyl are each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ haloalkyl, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)OR^{a3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}S(O)R^{b3}, NR^{c1}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, and S(O)₂NR^{c3}R^{d3}.

30 In some embodiments, the compounds of the invention have Formula Vb:



Vb.

In some embodiments, where the compounds of the invention have Formula Vb, R² is CF₃.

5 In some embodiments, where the compounds of the invention have Formula Vb, R¹ is H or C₁₋₁₀ alkyl.

In some embodiments, where the compounds of the invention have Formula Vb, R^A is Cy¹.

10 In some embodiments, where the compounds of the invention have Formula Vb, R^A is C₆₋₁₀ aryl or 5-10 membered heteroaryl, each of which is optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy}.

In some embodiments, where the compounds of the invention have Formula Vb, R^A is 5-10 membered heteroaryl optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy}.

15 In some embodiments, where the compounds of the invention have Formula Vb, R^A is 5 to 6-membered heteroaryl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.


In some embodiments, where the compounds of the invention have Formula Vb, R^A is C₆₋₁₀ aryl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.

20 In some embodiments, where the compounds of the invention have Formula Vb, R^A is phenyl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.

In some embodiments, where the compounds of the invention have Formula Vb, R^B is Cy².

In some embodiments, where the compounds of the invention have Formula Vb, R^B is H, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, OR^{a3} , $C(O)NR^{c3}R^{d3}$, or $C(O)OR^{a3}$, wherein said C_{1-6} alkyl and C_{2-6} alkenyl are each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} haloalkyl, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$,
5 $C(O)OR^{a3}$, $OC(O)R^{b3}$, $OC(O)NR^{c3}R^{d3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)OR^{a3}$, $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}S(O)R^{b3}$, $NR^{c1}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, $S(O)R^{b3}$, $S(O)NR^{c3}R^{d3}$, $S(O)_2R^{b3}$, and $S(O)_2NR^{c3}R^{d3}$.

In some embodiments, the compounds of the invention have Formula VI:



VI.

In some embodiments, where the compounds of the invention have Formula VI, R^2 is CF_3 .

In some embodiments, where the compounds of the invention have Formula VI, R^1 is H
15 or C_{1-10} alkyl.


In some embodiments, where the compounds of the invention have Formula VI, R^B is Cy^2 .

In some embodiments, where the compounds of the invention have Formula VI, Cy^2 is phenyl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy} .

In some embodiments, where the compounds of the invention have Formula VI, R^B is H,
20 halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, OR^{a3} , $C(O)NR^{c3}R^{d3}$, or $C(O)OR^{a3}$, wherein said C_{1-6} alkyl and C_{2-6} alkenyl are each optionally substituted with 1, 2, or 3 substituents

independently selected from halo, C₁₋₆ haloalkyl, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)OR^{a3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}S(O)R^{b3}, NR^{c1}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, and S(O)₂NR^{c3}R^{d3}.

- 5 In some embodiments, where the compounds of the invention have Formula VI, R^C is H.
In some embodiments, where the compounds of the invention have Formula VI, R^D is H.
In some embodiments, the compounds of the invention have Formula VIA:



- 10 VIA.

In some embodiments, where the compounds of the invention have Formula VIA, R² is CF₃.

In some embodiments, where the compounds of the invention have Formula VIA, R¹ is H or C₁₋₁₀ alkyl.


- 15 In some embodiments, where the compounds of the invention have Formula VIA, R^B is Cy².

In some embodiments, where the compounds of the invention have Formula VIA, Cy² is phenyl optionally substituted by 1, 2, or 3 substituents independently selected from R^{Cy}.

- 20 In some embodiments, where the compounds of the invention have Formula VIA, R^B is H, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, CN, OR^{a3}, C(O)NR^{c3}R^{d3}, or C(O)OR^{a3}, wherein said C₁₋₆ alkyl and C₂₋₆ alkenyl are each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ haloalkyl, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3},

$C(O)OR^{a3}$, $OC(O)R^{b3}$, $OC(O)NR^{c3}R^{d3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)OR^{a3}$,
 $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}S(O)R^{b3}$, $NR^{c1}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, $S(O)R^{b3}$, $S(O)NR^{c3}R^{d3}$,
 $S(O)_2R^{b3}$, and $S(O)_2NR^{c3}R^{d3}$.

In some embodiments, the compounds of the invention have Formula VII:



VII

wherein a is 0, 1, 2, or 3.

In some embodiments, where the compounds of the invention have Formula VII, R^2 is CF_3 .

In some embodiments, where the compounds of the invention have Formula VII, R^1 is H or C_{1-10} alkyl.

In some embodiments, where the compounds of the invention have Formula VII, R^B is Cy^2 .

In some embodiments, where the compounds of the invention have Formula VII, R^B is H, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, OR^{a3} , $C(O)NR^{c3}R^{d3}$, or $C(O)OR^{a3}$, wherein said C_{1-6} alkyl and C_{2-6} alkenyl are each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} haloalkyl, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$, $C(O)OR^{a3}$, $OC(O)R^{b3}$, $OC(O)NR^{c3}R^{d3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)OR^{a3}$, $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}S(O)R^{b3}$, $NR^{c1}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, $S(O)R^{b3}$, $S(O)NR^{c3}R^{d3}$, $S(O)_2R^{b3}$, and $S(O)_2NR^{c3}R^{d3}$.

In some embodiments, where the compounds of the invention have Formula VII, R^B is H or halo.


In some embodiments, where the compounds of the invention have Formula VII, R^B is halo.

In some embodiments, where the compounds of the invention have Formula VII, R^C is H.

In some embodiments, where the compounds of the invention have Formula VII, R^D is H.

5 In some embodiments, where the compounds of the invention have Formula VII, R^{Cy} is halo, C_{1-6} alkyl, C_{1-6} haloalkyl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, and $S(O)_2NR^{c5}R^{d5}$, wherein said C_{1-6} alkyl and 4-10 membered heterocycloalkyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C_{1-6} alkyl, CN, NO_2 , OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$,
10 $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)OR^{a5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}S(O)R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, and $S(O)_2NR^{c5}R^{d5}$.

In some embodiments, the compounds of the invention have Formula VIII:



15 VIII

wherein a is 0, 1, 2, or 3.

In some embodiments, where the compounds of the invention have Formula VIII, R^2 is CF_3 .

20 In some embodiments, where the compounds of the invention have Formula VIII, R^1 is H or C_{1-10} alkyl.

In some embodiments, where the compounds of the invention have Formula VIII, R^B is Cy^2 .

In some embodiments, where the compounds of the invention have Formula VIII, R^B is H, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, OR^{a3} , $C(O)NR^{c3}R^{d3}$, or $C(O)OR^{a3}$, wherein said C_{1-6} alkyl and C_{2-6} alkenyl are each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} haloalkyl, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$,
5 $C(O)OR^{a3}$, $OC(O)R^{b3}$, $OC(O)NR^{c3}R^{d3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)OR^{a3}$,
 $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}S(O)R^{b3}$, $NR^{c1}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, $S(O)R^{b3}$, $S(O)NR^{c3}R^{d3}$,
 $S(O)_2R^{b3}$, and $S(O)_2NR^{c3}R^{d3}$.

In some embodiments, where the compounds of the invention have Formula VIII, R^B is H or halo.

10 In some embodiments, where the compounds of the invention have Formula VIII, R^B is halo.

In some embodiments, where the compounds of the invention have Formula VIII, R^C is H.

15 In some embodiments, where the compounds of the invention have Formula VIII, R^D is H.

In some embodiments, where the compounds of the invention some embodiments have Formula VIII, R^{Cy} is halo, C_{1-6} alkyl, C_{1-6} haloalkyl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, and $S(O)_2NR^{c5}R^{d5}$, wherein said C_{1-6} alkyl and 4-10 membered heterocycloalkyl are each optionally substituted with 1, 2, 3,
20 4, or 5 substituents independently selected from halo, C_{1-6} alkyl, CN, NO_2 , OR^{a5} , SR^{a5} , $C(O)R^{b5}$,
 $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)OR^{a5}$,
 $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}S(O)R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$,
 $S(O)_2R^{b5}$, and $S(O)_2NR^{c5}R^{d5}$.

In some embodiments, where the compounds of the invention have Formula VIII, a is 0.

25 In some embodiments, the chiral carbon to which $-C(O)OR^1$ is attached has an *S* configuration.

In some embodiments, the carbon to which $-R^2$ is attached is chiral and has an *R* configuration.

30 It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, can also be provided in combination in a single embodiment. Conversely, various features of the invention which are, for brevity, described in

the context of a single embodiment, can also be provided separately or in any suitable subcombination.

The term "substituted" means that an atom or group of atoms formally replaces hydrogen as a "substituent" attached to another group. The hydrogen atom is formally removed and replaced by a substituent. A single divalent substituent, *e.g.*, oxo, can replace two hydrogen atoms. The term "optionally substituted" means unsubstituted or substituted. The substituents are independently selected, and substitution may be at any chemically accessible position. It is to be understood that substitution at a given atom is limited by valency. Throughout the definitions, the term " C_{i-j} " indicates a range which includes the endpoints, wherein *i* and *j* are integers and indicate the number of carbons. Examples include C_{1-4} , C_{1-6} , and the like.

The term "n-membered" where *n* is an integer typically describes the number of ring-forming atoms in a moiety where the number of ring-forming atoms is *n*. For example, piperidinyl is an example of a 6-membered heterocycloalkyl ring, pyrazolyl is an example of a 5-membered heteroaryl ring, pyridyl is an example of a 6-membered heteroaryl ring, and 1, 2, 3, 4-tetrahydro-naphthalene is an example of a 10-membered cycloalkyl group.

At various places in the present specification various aryl, heteroaryl, cycloalkyl, and heterocycloalkyl rings are described. Unless otherwise specified, these rings can be attached to the rest of the molecule at any ring member as permitted by valency. For example, the term "a pyridine ring" or "pyridinyl" may refer to a pyridin-2-yl, pyridin-3-yl, or pyridin-4-yl ring.

For compounds of the invention in which a variable appears more than once, each variable can be a different moiety independently selected from the group defining the variable. For example, where a structure is described having two *R* groups that are simultaneously present on the same compound, the two *R* groups can represent different moieties independently selected from the group defined for *R*.

As used herein, the term " C_{i-j} alkyl," employed alone or in combination with other terms, refers to a saturated hydrocarbon group that may be straight-chain or branched, having *i* to *j* carbon atoms. In some embodiments, the alkyl group contains from 1 to 10, 1 to 6, 1 to 4, or from 1 to 3 carbon atoms. Examples of alkyl moieties include, but are not limited to, chemical groups such as methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *s*-butyl, and *t*-butyl.

As used herein, the term " C_{i-j} alkoxy," employed alone or in combination with other terms, refers to a group of formula -O-alkyl, wherein the alkyl group has *i* to *j* carbon atoms.

Example alkoxy groups include methoxy, ethoxy, and propoxy (*e.g.*, *n*-propoxy and isopropoxy). In some embodiments, the alkyl group has 1 to 3 carbon atoms or 1 to 4 carbon atoms.

As used herein, " C_{i-j} alkenyl" refers to an alkyl group having one or more double carbon-carbon bonds and having *i* to *j* carbon atoms. In some embodiments, the alkenyl moiety contains
5 2 to 6 or to 2 to 4 carbon atoms. Example alkenyl groups include, but are not limited to, ethenyl, *n*-propenyl, isopropenyl, *n*-butenyl, *sec*-butenyl, and the like.

As used herein, the term " C_{i-j} alkylamino" refers to a group of formula -NH(alkyl), wherein the alkyl group has *i* to *j* carbon atoms. In some embodiments, the alkyl group has 1 to 6 or 1 to 4 carbon atoms.

10 As used herein, the term "di- C_{i-j} -alkylamino" refers to a group of formula -N(alkyl)₂, wherein the two alkyl groups each has, independently, *i* to *j* carbon atoms. In some embodiments, each alkyl group independently has 1 to 6 or 1 to 4 carbon atoms.

As used herein, the term "thio" refers to a group of formula -SH.

As used herein, the term " C_{i-j} alkylthio" refers to a group of formula -S-alkyl, wherein the
15 alkyl group has *i* to *j* carbon atoms. In some embodiments, the alkyl group has 1 to 6 or 1 to 4 carbon atoms.

As used herein, the term "amino" refers to a group of formula -NH₂.

As used herein, the term " C_{i-j} aryl," employed alone or in combination with other terms, refers to a monocyclic or polycyclic (*e.g.*, having 2, 3 or 4 fused rings) aromatic hydrocarbon
20 having *i* to *j* ring-forming carbon atoms, such as, but not limited to, phenyl, 1-naphthyl, 2-naphthyl, anthracenyl, phenanthrenyl, and the like. In some embodiments, aryl is C_{6-10} aryl. In some embodiments, the aryl group is a naphthalene ring or phenyl ring. In some embodiments, the aryl group is phenyl.

As used herein, the term "arylalkyl" refers to a group of formula - C_{i-j} alkyl-(C_{i-j} aryl). In
25 some embodiments, arylalkyl is C_{6-10} aryl- C_{1-3} alkyl. In some embodiments, arylalkyl is C_{6-10} aryl- C_{1-4} alkyl. In some embodiments, arylalkyl is benzyl.

As used herein, the term "carbonyl," employed alone or in combination with other terms, refers to a -C(=O)- group.

As used herein, the term "carboxy" refers to a group of formula -C(=O)OH.

30 As used herein, the term " C_{i-j} cycloalkyl," employed alone or in combination with other terms, refers to a non-aromatic cyclic hydrocarbon moiety having *i* to *j* ring-forming carbon

atoms, which may optionally contain one or more alkenylene groups as part of the ring structure. Cycloalkyl groups can include mono- or polycyclic (*e.g.*, having 2, 3 or 4 fused rings) ring systems. Also included in the definition of cycloalkyl are moieties that have one or more aromatic rings (aryl or heteroaryl) fused to the cycloalkyl ring, for example, benzo or pyrido derivatives of cyclopentane, cyclopentene, cyclohexane, and the like. Where the cycloalkyl group includes a fused aromatic ring, the cycloalkyl group can be attached at either an atom in the aromatic or non-aromatic portion. One or more ring-forming carbon atoms of a cycloalkyl group can be oxidized to form carbonyl linkages. In some embodiments, cycloalkyl is C₃₋₁₀ or C₃₋₇ cycloalkyl, which can be monocyclic or polycyclic. Exemplary cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptatrienyl, norbornyl, norpinyl, norcarnyl, adamantanyl and the like. In some embodiments, the cycloalkyl group is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

As used herein, the term "cycloalkylalkyl" refers to a group of formula –C_{i-j} alkyl–(C_{i-j} cycloalkyl). In some embodiments, cycloalkylalkyl is C₃₋₇ cycloalkyl–C₁₋₃ alkyl, wherein the cycloalkyl portion is monocyclic. In some embodiments, cycloalkylalkyl is C₃₋₇ cycloalkyl–C₁₋₄ alkyl.

As used herein, "C_{i-j} haloalkoxy" refers to a group of formula –O-haloalkyl having i to j carbon atoms. An example haloalkoxy group is OCF₃. An additional example haloalkoxy group is OCHF₂. In some embodiments, the alkyl group has 1 to 6 or 1 to 4 carbon atoms.

As used herein, the term "halo" refers to a halogen atom selected from F, Cl, I or Br. In some embodiments, "halo" refers to a halogen atom selected from F, Cl, or Br. In some embodiments, the halo group is F.

As used herein, the term "C_{i-j} haloalkyl," employed alone or in combination with other terms, refers to an alkyl group having from one halogen atom to 2s+1 halogen atoms which may be the same or different, where "s" is the number of carbon atoms in the alkyl group, wherein the alkyl group has i to j carbon atoms. In some embodiments, the haloalkyl group is fluoromethyl, difluoromethyl, or trifluoromethyl. In some embodiments, the haloalkyl group is trifluoromethyl. In some embodiments, the haloalkyl group has 1 to 6 or 1 to 4 carbon atoms.

As used herein, the term "heteroaryl," employed alone or in combination with other terms, refers to a monocyclic or polycyclic (*e.g.*, having 2, 3 or 4 fused rings) aromatic moiety, having one or more heteroatom ring members selected from nitrogen, sulfur and oxygen. In some

embodiments, the heteroaryl group is a 5- to 10-membered heteroaryl ring, which is monocyclic or bicyclic and which has 1, 2, 3, or 4 heteroatom ring members independently selected from nitrogen, sulfur and oxygen. In some embodiments, the heteroaryl group is a 5- to 6-membered heteroaryl ring, which is monocyclic and which has 1, 2, 3, or 4 heteroatom ring members independently selected from nitrogen, sulfur and oxygen. When the heteroaryl group contains more than one heteroatom ring member, the heteroatoms may be the same or different. The nitrogen atoms in the ring(s) of the heteroaryl group can be oxidized to form N-oxides. Example heteroaryl groups include, but are not limited to, pyridine, pyrimidine, pyrazine, pyridazine, pyrrole, pyrazole, azolyl, oxazole, thiazole, imidazole, furan, thiophene, quinoline, isoquinoline, indole, benzothiophene, benzofuran, benzisoxazole, imidazo[1,2-*b*]thiazole, purine, and the like.

A 5-membered heteroaryl is a heteroaryl group having five ring-forming atoms comprising carbon and one or more (*e.g.*, 1, 2, or 3) ring atoms independently selected from N, O, and S. Example five-membered heteroaryls include thienyl, furyl, pyrrolyl, imidazolyl, thiazolyl, oxazolyl, pyrazolyl, isothiazolyl, isoxazolyl, 1,2,3-triazolyl, tetrazolyl, 1,2,3-thiadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-triazolyl, 1,2,4-thiadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-triazolyl, 1,3,4-thiadiazolyl, and 1,3,4-oxadiazolyl.

A six-membered heteroaryl is a heteroaryl group having six ring-forming atoms wherein one or more (*e.g.*, 1, 2, or 3) ring atoms are independently selected from N, O, and S. Example six-membered heteroaryls include pyridyl, pyrazinyl, pyrimidinyl, triazinyl and pyridazinyl.

As used herein, the term "heteroarylalkyl" refers to a group of formula $-C_{i-j}$ alkyl- (heteroaryl). In some embodiments, heteroarylalkyl 5-10 membered heteteroaryl- C_{1-4} alkyl, wherein the heteroaryl portion is monocyclic or bicyclic and has 1, 2, 3, or 4 heteroatom ring members independently selected from nitrogen, sulfur and oxygen. In some embodiments, the heteroarylalkyl is 5-6 membered heteteroaryl- C_{1-3} alkyl or 5-6 membered heteteroaryl- C_{1-4} alkyl, wherein the heteroaryl portion is monocyclic and has 1, 2, 3, or 4 heteroatom ring members independently selected from nitrogen, sulfur and oxygen.

As used herein, the term "heterocycloalkyl," employed alone or in combination with other terms, refers to a non-aromatic ring or ring system, which optionally contains one or more alkenylene groups as part of the ring structure, and which has at least one heteroatom ring member independently selected from nitrogen, sulfur and oxygen. When the heterocycloalkyl groups contains more than one heteroatom, the heteroatoms may be the same or different.

Heterocycloalkyl groups can include mono- or polycyclic (*e.g.*, having 2, 3 or 4 fused rings) ring systems, including spiro systems. Also included in the definition of heterocycloalkyl are moieties that have one or more aromatic rings (aryl or heteroaryl) fused to the non-aromatic ring, for example, 1,2,3,4-tetrahydro-quinoline, dihydrobenzofuran and the like. Where the

5 heterocycloalkyl group includes a fused aromatic ring, the heterocycloalkyl group can be attached at either an atom in the aromatic or non-aromatic portion. The carbon atoms or heteroatoms in the ring(s) of the heterocycloalkyl group can be oxidized (*e.g.* have one or two oxo substituents) to form a carbonyl, or sulfonyl group (or other oxidized linkage) or a nitrogen atom can be quaternized. In some embodiments, the heterocycloalkyl group is 5- to 10-
10 membered, which can be monocyclic or bicyclic and which has 1, 2, 3, or 4 heteroatom ring members independently selected from nitrogen, sulfur and oxygen. In some embodiments, the heterocycloalkyl group is 5- to 6-membered or 5- to 7-membered. Examples of heterocycloalkyl groups include 1, 2, 3, 4-tetrahydroquinoline, dihydrobenzofuran, azetidine, azepane, pyrrolidine, piperidine, piperazine, morpholine, thiomorpholine, and pyran. Further examples of
15 heterocycloalkyl groups include 2-oxotetrahydrofuranyl, 2-oxopyrrolidinyl, 2-oxoimidazolidinyl, 1-oxo-1,2,3,4-tetrahydroisoquinolin-6-yl, and 2-oxo-1,3-dioxolan-4-yl.

As used herein, the term "heterocycloalkylalkyl" refers to a group of formula $-C_{i-j}$ alkyl- (heterocycloalkyl). In some embodiments, heterocycloalkylalkyl is 5-10 membered heterocycloalkyl- C_{1-3} alkyl or 5-10 membered heterocycloalkyl- C_{1-4} alkyl, wherein the
20 heterocycloalkyl portion is monocyclic or bicyclic and has 1, 2, 3, or 4 heteroatom ring members independently selected from nitrogen, sulfur and oxygen. In some embodiments, heterocycloalkylalkyl is 5-6 membered heterocycloalkyl- C_{1-4} alkyl wherein the heterocycloalkyl portion is monocyclic and has 1, 2, 3, or 4 heteroatom ring members independently selected from nitrogen, sulfur and oxygen.

25 The compounds described herein can be asymmetric (*e.g.*, having one or more stereocenters). All stereoisomers, such as enantiomers and diastereoisomers, are intended unless otherwise indicated. Compounds of the present invention that contain asymmetrically substituted carbon atoms can be isolated in optically active or racemic forms. Methods on how to prepare optically active forms from optically inactive starting materials are known in the art, such as by
30 resolution of racemic mixtures or by stereoselective synthesis. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein,

and all such stable isomers are contemplated in the present invention. *Cis* and *trans* geometric isomers of the compounds of the present invention may be isolated as a mixture of isomers or as separated isomeric forms.

Resolution of racemic mixtures of compounds can be carried out by any of numerous methods known in the art. An example method includes fractional recrystallization using a chiral resolving acid which is an optically active, salt-forming organic acid. Suitable resolving agents for fractional recrystallization methods are, for example, optically active acids, such as the D and L forms of tartaric acid, diacetyltartaric acid, dibenzoyltartaric acid, mandelic acid, malic acid, lactic acid or the various optically active camphorsulfonic acids such as β -camphorsulfonic acid. Other resolving agents suitable for fractional crystallization methods include stereoisomerically pure forms of α -methylbenzylamine (*e.g.*, *S* and *R* forms, or diastereoisomerically pure forms), 2-phenylglycinol, norephedrine, ephedrine, N-methylephedrine, cyclohexylethylamine, 1,2-diaminocyclohexane, and the like.

Resolution of racemic mixtures can also be carried out by elution on a column packed with an optically active resolving agent (*e.g.*, dinitrobenzoylphenylglycine). Suitable elution solvent composition can be determined by one skilled in the art.

Compounds of the invention can also include tautomeric forms. Tautomeric forms result from the swapping of a single bond with an adjacent double bond together with the concomitant migration of a proton. Tautomeric forms include prototropic tautomers which are isomeric protonation states having the same empirical formula and total charge. Example prototropic tautomers include ketone – enol pairs, amide - imidic acid pairs, lactam – lactim pairs, amide - imidic acid pairs, enamine – imine pairs, and annular forms where a proton can occupy two or more positions of a heterocyclic system, for example, *1H*- and *3H*-imidazole, *1H*-, *2H*- and *4H*-1, 2, 4-triazole, *1H*- and *2H*- isoindole, and *1H*- and *2H*-pyrazole.

Compounds of the invention can also include all isotopes of atoms occurring in the intermediates or final compounds. Isotopes include those atoms having the same atomic number but different mass numbers. For example, isotopes of hydrogen include tritium and deuterium.

The term "compound," as used herein, is meant to include all stereoisomers, geometric isomers, tautomers, and isotopes of the structures depicted. Compounds herein identified by name or structure as one particular tautomeric form are intended to include other tautomeric forms unless otherwise specified. Compounds herein identified by name or structure without

specifying the particular configuration of a stereocenter are meant to encompass all the possible configurations at the stereocenter. For example, if a particular stereocenter in a compound of the invention could be R or S, but the name or structure of the compound does not designate which it is, then the stereocenter can be either R or S.

5 All compounds, and pharmaceutically acceptable salts thereof, can be found together with other substances such as water and solvents (*e.g.*, hydrates and solvates) or can be isolated.

In some embodiments, the compounds of the invention, or salts thereof, are substantially isolated. By "substantially isolated" is meant that the compound is at least partially or substantially separated from the environment in which it was formed or detected. Partial
10 separation can include, for example, a composition enriched in the compounds of the invention. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the compounds of the invention, or salt thereof. Methods for isolating compounds and their salts are routine in the art.

15 The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

20 The expressions, "ambient temperature" and "room temperature," as used herein, are understood in the art, and refer generally to a temperature, *e.g.*, a reaction temperature, that is about the temperature of the room in which the reaction is carried out, for example, a temperature from about 20 °C to about 30 °C.

The present invention also includes pharmaceutically acceptable salts of the compounds
25 described herein. As used herein, "pharmaceutically acceptable salts" refers to derivatives of the disclosed compounds wherein the parent compound is modified by converting an existing acid or base moiety to its salt form. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts of
30 the present invention include the conventional non-toxic salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically acceptable salts of

the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, non-aqueous media like ether, EtOAc, alcohols (*e.g.*, methanol, ethanol, iso-propanol, or butanol) or acetonitrile (CH₃CN) are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th Ed., (Mack Publishing Company, Easton, 1985), p. 1418, Berge *et al.*, *J. Pharm. Sci.*, **1977**, 66(1), 1-19, and in Stahl *et al.*, *Handbook of Pharmaceutical Salts: Properties, Selection, and Use*, (Wiley, 2002).

The below Table is a key to some abbreviations used throughout.

Abbreviations

atm	atmosphere
BOC	tert-butyl-oxy-carbonyl
CAS#	Chemical Abstract Service registry number
CBS	Corey-Bakshi-Shibata (catalyst)
CH ₃ CN	Acetonitrile
CBZ	Carbobenzyloxy
DIPEA	N,N-diisopropylethylamine
DMAP	4-dimethylaminopyridine
DME	dimethylether
DMF	dimethyl formamide
dppf	1,1'-bis(diphenylphosphino)ferrocene
EDCI	1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride
ee	enantiomeric excess
EtOAc	ethyl acetate
h	hour(s)
min	minute(s)
HOAT	1-hydroxy-7-azabenzotriazole
HOAc	acetic acid
HPLC	high-performance liquid chromatography

KOAc	potassium acetate
LAH	lithium aluminum hydride
LDA	lithium diisopropylamide
mCPBA	3-meta-chloroperoxybenzoic acid
MeOH	Methanol
MS	mass spectrometry
MTBE	methyl t-butyl ether
NH ₄ OH	ammonium hydroxide
NMP	1-methyl-2-pyrrolidone
PAH	pulmonary arterial hypertension
PE	petroleum ether
PheOH	phenylalanine hydroxylase
Prep-TLC	preparative thin-layer chromatography
p-TSA	para-toluene sulfonic acid
RT	room temperature
SNAr	nucleophilic aromatic substitution
TBAF	tetrabutylammonium fluoride
tBuOH	tert-butanol
TBTU	O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate
TEA	Triethylamine
TFA	trifluoroacetic acid
TH	tyrosine hydroxylase
THF	Tetrahydrofuran
TLC	thin-layer chromatography
TMS	Trimethylsilyl
TMSI	Trimethylsilyl iodide
TPH	tryptophan hydroxylase


Synthesis

Procedures for making compounds described herein are provided below with reference to Schemes 1-10. Optimum reaction conditions and reaction times may vary depending on the

particular reactants used. Unless otherwise specified, solvents, temperatures, pressures and other reaction conditions are readily selected by one of ordinary skill in the art. Specific procedures are provided in the Examples section. Compounds are named using the “structure to name” function included in ChemDraw[®] v.12 (Perkin-Elmer).

Typically, reaction progress may be monitored by thin layer chromatography (TLC) or HPLC-MS if desired. Intermediates and products may be purified by chromatography on silica gel, recrystallization, HPLC and/or reverse phase HPLC. In the reactions described below, it may be necessary to protect reactive functional groups (such as hydroxy, amino, thio, or carboxy groups) to avoid their unwanted participation in the reactions. The incorporation of such groups, and the methods required to introduce and remove them are known to those skilled in the art (for example, see Greene, Wuts, *Protective Groups in Organic Synthesis*, 2nd Ed. (1999)). One or more deprotection steps in the synthetic schemes may be required to ultimately afford compounds of Formula I. The protecting groups depicted in the schemes are used as examples, and may be replaced by other compatible alternative groups. Starting materials used in the following schemes can be purchased or prepared by methods described in the chemical literature, or by adaptations thereof, using methods known by those skilled in the art. The order in which the steps are performed can vary depending on the protecting or functional groups introduced and the reagents and reaction conditions used, but would be apparent to those skilled in the art.

Compounds of Formula I can be prepared as shown in general in Scheme 1. Briefly, in step 1, an alcohol (where the ring substituted by R^A, R^B, R^C, R^D corresponds to ring A of Formula I) (see, e.g., Intermediate 1) in dioxane is treated with a dichloro heterocycle (e.g., 2-amino-4,6-dichloropyrimidine) in the presence of a base (e.g., Cs₂CO₃), and heated for several hours (e.g. 12-24 h) at reflux. In step 2, a spirocycle of formula **B** (e.g., (S)-2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate) is added to a solution of compound **A** in a solvent (e.g., dioxane) in the presence of a base (e.g. Na₂CO₃), and heated to reflux to provide a compound of formula **C**. In step 3, the amino protecting group (P) (e.g. CBZ or BOC) of a compound of formula **C** is removed (e.g. with TMSI, transition metal-catalyzed hydrogenation, or strong acid depending on the nature of the protecting group). In step 4, a compound of formula **D** is obtained by ester hydrolysis (e.g. with LiOH in aqueous THF). In some instances, the sequence of steps 3 and 4 can be reversed.



Scheme 1


5 Alcohols (e.g., Intermediate 1) used in Scheme 1 can be prepared as shown in Scheme 2. Briefly, in step 1, to a solution of base (e.g. potassium *t*-butoxide) in a solvent (e.g. DMSO) is added 3-methyl pyrazole and an aryl bromide **E** (e.g., 1,4-dibromo-2-fluorobenzene), and the mixture is heated for several hours (e.g. 12-24 h) to provide a compound of formula **F**. In step 2, a compound of formula **F** is treated with a Grignard reagent (e.g., *i*-PrMgCl) in a solvent (e.g.,

10 THF), then reacted with ethyl trifluoroacetate in a solvent (e.g., THF) to provide a ketone of formula **G**.

 Alternatively, a ketone of formula **G** can be obtained by treating first a fluoro aromatic compound of formula **E1** with a strong base (e.g., LDA), then trapping the intermediate aryl lithium with trifluoroacetic acid ethyl ester to give a compound of formula **F1** (Step 1a). In a

15 subsequent step 2a, 3-methyl pyrazole can be introduced onto a ketone of formula **F1** via an $\text{S}_{\text{N}}\text{Ar}$ reaction in the presence of base (e.g., K_2CO_3) under solvent reflux (e.g., toluene). In step 3, a ketone of formula **G** is converted stereospecifically into a chiral alcohol of formula **H** via either chiral transfer hydrogenation (e.g., with potassium formate) in the presence of a transition


metal catalyst (e.g., pentamethyl cyclopentadienyl iridium (III) chloride dimer) and a chiral ligand (e.g., (1R,2R)-(-)-N-(4-toluene sulfonyl)-1,2-diphenyl ethylene diamine) in a solvent (e.g., acetonitrile), or alternatively with a borane reagent (e.g. catechol borane) and a chiral catalyst (e.g. (S)-2-methyl-CBS oxazaborolidine) in a solvent (e.g., THF). Alternatively, an alcohol of formula **K** can be made in a similar fashion starting from a ketone of formula **J** (step 2c). A ketone of formula **J** can be prepared in one step (step 2c) by reacting the aryl ester of formula **E2** with a nucleophilic silylating agent (e.g., trimethyl(trifluoromethyl)silane) in the presence of a fluoride source (e.g., TBAF) in an inert solvent (e.g., THF).



Scheme 2

Other types of oxygen or nitrogen linker groups (L-groups) can be installed as shown in Scheme 3. Briefly, in step 1, to a spirocyclic compound of **B** (e.g., (S)-2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate) in dioxane is added a di-halo heterocycle (e.g., 2-amino-4,6-dichloropyrimidine) in the presence of a base (e.g., Cs_2CO_3) under solvent reflux (e.g., dioxane) to provide a compound of formula **M**. In step 2, to a compound of formula **M** in a solvent (e.g., dioxane) is added an alcohol or an amine of formula **O** (e.g., Intermediate 7 or


16) in the presence of a base (e.g., Cs_2CO_3). After heating at reflux for several hours (e.g., 12-24 h), a compound of formula **P** is obtained. In step 3, the amino protecting group (P) (e.g., CBZ or BOC) of a compound of formula **P** is removed (e.g., with TMSI, transition metal-catalyzed hydrogenation, or acid). Then, in step 4, a compound of formula **Q** is obtained by ester hydrolysis (e.g., with LiOH in aqueous THF). In some instances, the sequence of steps 3 and 4 can be reversed.



Scheme 3


For certain substituents and substitution patterns, palladium-mediated coupling reactions (e.g., Suzuki or Stille reactions) can be used, as shown in Schemes 4a, 4b, and 4c. Briefly, in step 1, to a compound of formula **R** in a solvent (e.g., aqueous dioxane) is added a boronic acid or boronate (e.g., phenyl boronic acid) in the presence of a palladium catalyst (e.g., PdCl₂(dppf)-CH₂Cl₂) and a base (e.g., KHCO₃), and the mixture heated to reflux for several hours (e.g., 12-24) to provide a compound of formula **S**. In step 3, the amino protecting group (P) (e.g., CBZ or BOC) of a compound of formula **S** is removed (e.g., with TMSI, transition metal-catalyzed hydrogenation, or acid). Then, in step 4, a compound of formula **T** is obtained by ester hydrolysis (e.g., with LiOH in aqueous THF). In some instances, the sequence of steps 2 and 3 can be reversed. A similar set of conditions can be used when starting with a compound of formula **U** or **X**, to obtain a compound of formula **W** or **AA**, respectively (Schemes 4b and 4c).

BOC) of a compound of formula **S** is removed (e.g., with TMSI, transition metal-catalyzed




Scheme 4a

5




Scheme 4b




Scheme 4c

Various substitutions of the central 6-membered ring (e.g., the ring containing W, X, and Y) can be accomplished as shown in Scheme 5. Briefly, in step 1, to a solution of a methyl sulfide of formula **AB** in an inert solvent (e.g., CH_2Cl_2) is added an oxidant (e.g., m-CPBA). The solution is stirred at RT for several hours (e.g., 12-24 h) to provide a sulfone of formula **AC**. In step 2, to a solution of a compound of formula **AC** in a solvent (e.g., dioxane) is added a spirocyclic compound of formula **B** (e.g., (S)-2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate) in the presence of a base (e.g., Cs_2CO_3), and the mixture is heated for several hours (e.g., 12-24 h) to provide a sulfone of formula **AD**. In step 3, the ester group is saponified (e.g., with LiOH) in an aqueous or alcoholic solvent (e.g., aqueous THF) to provide an acid of formula **AE**. In step 4, heating an acid of formula **AE** in the presence of an alcohol or an amine (e.g., phenol) and a base (e.g., Cs_2CO_3) for several hours (e.g., 16-24 h) in a solvent (e.g., dioxane), followed in step 5 by deprotection of the amine (e.g. with TMSI, transition metal-catalyzed hydrogenation, or acid) provides a compound of formula **AF**.




Scheme 5

Ester group substituents can be introduced by the general method of Scheme 6. Briefly, in step 1, to a solution of an acid of formula **AG** in an inert solvent (e.g., CH_2Cl_2) is added a coupling reagent (e.g., EDCI and DMAP), followed by an alcohol (e.g., propanol) to provide a compound of formula **AH**. In step 2, the benzyl groups of the benzyl ester and of the N-CBZ group can be removed with reagents such as TMSI or by transition metal-catalyzed hydrogenation (e.g., H_2 with Pd/C), affording a compound of formula **AI**. In case the amino protecting group is a BOC, an additional step 3, involving treatment with a strong acid (e.g., TFA), can be used for the final deprotection.




Scheme 6

5 Ethyl esters can be generally prepared according to Scheme 7. Briefly, deprotection of the amino group in a compound of formula **AJ**, can be accomplished either with the use of a dealkylating agent (e.g., TMSI) or via transition metal-catalyzed hydrogenation (e.g., H₂ with Pd/C) if the protecting group is CBZ, or with a strong acid (e.g., TFA or HCl), if the protecting group is BOC, to provide **AK**. It will be recognized by those skilled in the art that many other protecting groups can be used alternatively (for example, see Greene, Wuts, Protective Groups in Organic Synthesis, 2nd Ed. (1999)).




Scheme 7

Various esters can be made via direct alcohol coupling to the acid, as shown in Scheme 8, or via alkylation of the acid, as shown in Scheme 9. Briefly, an amino acid of formula **AL** is dissolved in an alcoholic solvent (e.g., n-octanol), optionally in the presence of a co-solvent (e.g., toluene), and heated in the presence of acid (e.g., p-TSA) for several hours (e.g., 12-24 h), optionally in the presence of a water trapping material (e.g., molecular sieve) or apparatus (e.g., Dean-Stark trap) to produce an ester of formula **AM**. Alternatively, in step 1, an acid of formula **AN** is dissolved in a solvent (e.g., DMF) in the presence of a base (e.g., K₂CO₃) and treated with an alkyl halide (e.g., 2-chloro-ethyl-dimethyl-amine). After heating the solution for several hours (e.g., 12-24 h), an ester of formula **AO** is obtained. In step 2, removal of the amino protecting group (e.g., with an acid like TFA in an inert solvent such as CH₂Cl₂ in case of a BOC protecting group) provides an ester of formula **AP**. Other compatible deprotection methods apparent to those skilled in the art can be applied for other types of amino protecting groups.




Scheme 8



Scheme 9

t-Butyl esters can be made via direct alcohol coupling to the acid, as shown in Scheme 10. Briefly, in step 1, an acid of formula **AQ** is dissolved in a solvent (e.g., DMF) in the presence of *t*-butanol, and treated with a coupling agent (e.g., EDCI and DMAP) to provide a compound of formula **AR**. In step 2, removal of the amino protecting group is achieved as described earlier to afford a compound of formula **AS**.



Scheme 10

Methods of Use

The compounds of the invention can be used to inhibit the activity of the TPH1 enzyme in a cell by contacting the cell with an inhibiting amount of a compound of the invention. The cell can be part of the tissue of a living organism, or can be in culture, or isolated from a living organism. Additionally, the compounds of the invention can be used to inhibit the activity of the TPH1 enzyme in an animal, individual, or patient, by administering an inhibiting amount of a compound of the invention to the cell, animal, individual, or patient.

Compounds of the invention can also lower peripheral serotonin levels in an animal, individual, or patient, by administering an effective amount of a compound of the invention to the animal, individual, or patient. In some embodiments, the compounds of the invention can lower levels of peripheral serotonin (e.g., 5-HT in the GI tract) selectively over non-peripheral serotonin (e.g., 5-HT in the CNS). In some embodiments, the selectivity is 2-fold or more, 3-fold or more, 5-fold or more, 10-fold or more, 50-fold or more, or 100-fold or more.

As TPH1 inhibitors that can lower peripheral serotonin levels, the compounds of the invention are useful in the treatment and prevention of various diseases associated with abnormal expression or activity of the TPH1 enzyme, or diseases associated with elevated or abnormal peripheral serotonin levels. In some embodiments, the treatment or prevention includes

administering to a patient in need thereof a therapeutically effective amount of a TPH1 inhibitor of the invention.

Biological assays, some of which are described herein, can be used to determine the inhibitory effect of compounds against TPH (such as TPH1) *in vitro* and/or *in vivo*. *In vitro* biochemical assays for human, mouse, and rat TPH1 and human TPH2, PheOH, and TH may be used to measure inhibition of enzyme activity and the selectivity among TPH1, TPH2, PheOH, and TH. In addition, the efficacy of these compounds can be determined, for example, by measuring their effect on intestinal 5-HT levels in rodents after oral administration.

Diseases treatable or preventable by administering a TPH1 inhibitor of the invention include bone disease such as, for example, osteoporosis, osteoporosis pseudoglioma syndrome (OPPG), osteopenia, osteomalacia, renal osteodystrophy, Paget's disease, fractures, and bone metastasis. In some embodiments, the disease is osteoporosis, such as primary type 1 (e.g., postmenopausal osteoporosis), primary type 2 (e.g., senile osteoporosis), and secondary (e.g., steroid- or glucocorticoid-induced osteoporosis).

The present invention further includes methods of treating or preventing bone fracture such as, for example, osteoporotic or traumatic fracture, or surgical fractures associated with an orthopedic procedure (e.g., limb lengthening, bunion removal, an increase in bone formation associated with a prosthesis, bone metastasis, or spinal fusion).

Further diseases treatable or preventable by the methods of the invention include cardiovascular diseases such as atherosclerosis and pulmonary hypertension (PH), including idiopathic or familial PH, and also including PH associated with or brought on by other diseases or conditions. In some embodiments, the PH disease is pulmonary arterial hypertension (PAH).

The types of PAH treatable according to the methods of the invention include (1) idiopathic (IPAH), (2) familial (FPAH), and (3) associated (APAH) which is the most common type of PAH. The latter is PAH which is associated with other medical conditions including, for example, (1) collagen vascular disease (or connective tissue disease) which include autoimmune diseases such as scleroderma or lupus; (2) congenital heart and lung disease; (3) portal hypertension (e.g., resulting from liver disease); (4) HIV infection; (5) drugs (e.g., appetite suppressants, cocaine, and amphetamines); (6) other conditions including thyroid disorders, glycogen storage disease, Gaucher disease, hereditary hemorrhagic telangiectasia, hemoglobinopathies, myeloproliferative disorders, and splenectomy. APAH can also be PAH

associated with abnormal narrowing in the pulmonary veins and/or capillaries such as in pulmonary veno-occlusive disease (PVOD) and pulmonary capillary hemangiomatosis. Another type of PAH is associated with persistent pulmonary hypertension of the newborn (PPHN).

Further diseases treatable or preventable by the methods of the invention include

- 5 metabolic diseases such as diabetes and hyperlipidemia; pulmonary diseases such as chronic obstructive pulmonary disease (COPD), and pulmonary embolism; gastrointestinal diseases such as IBD, colitis, chemotherapy-induced emesis, diarrhea, carcinoid syndrome, celiac disease, Crohn's disease, abdominal pain, dyspepsia, constipation, lactose intolerance, MEN types I and II, Ogilvie's syndrome, pancreatic cholera syndrome, pancreatic insufficiency,
- 10 pheochromocytoma, scleroderma, somatization disorder, Zollinger-Ellison Syndrome, or other gastrointestinal inflammatory conditions; liver diseases such as chronic liver disease; cancers such as liver cancer, breast cancer, cholangiocarcinoma, colon cancer, colorectal cancer, neuroendocrine tumors, pancreatic cancer, prostate cancer, and bone cancer (e.g., osteosarcoma, chondrosarcoma, Ewing's sarcoma, osteblastoma, osteoid osteoma, osteochondroma,
- 15 enchondroma, chondromyxoid fibroma, aneurysmal bone cyst, unicameral bone cyst, giant cell tumor, and bone tumors); blood diseases (e.g., myeloproliferative syndrome, myelodysplastic syndrome, Hodgkin's lymphoma, non-Hodgkin's lymphoma, myeloma, and anemia such as aplastic anemia and anemia associated with kidney disease; and blood cancers (e.g., leukemias such as acute lymphocytic leukemia (ALL), chronic lymphocytic leukemia (CLL), acute
- 20 myeloid leukemia (AML), and chronic myeloid leukemia (CML)).

The compounds of the invention are also useful in the treatment and prevention of serotonin syndrome.

- In some embodiments, the present invention includes methods of lowering plasma cholesterol, lowering plasma triglycerides, lowering plasma glycerol, lowering plasma free fatty
- 25 acids in a patient by administering to said patient a therapeutically effective amount of a compound of the invention.

The compounds of the invention are also useful in the treatment and prevention of inflammatory disease, such as allergic airway inflammation (e.g., asthma).

As used herein, the term "cell" is meant to refer to a cell that is *in vitro*, *ex vivo* or *in vivo*.

- 30 In some embodiments, an *ex vivo* cell can be part of a tissue sample excised from an organism

such as a mammal. In some embodiments, an *in vitro* cell can be a cell in a cell culture. In some embodiments, an *in vivo* cell is a cell living in an organism such as a mammal.

As used herein, the term “contacting” refers to the bringing together of indicated moieties in an *in vitro* system or an *in vivo* system. For example, “contacting” the enzyme with a compound of the invention includes the administration of a compound of the present invention to an individual or patient, such as a human, having the TPH1 enzyme, as well as, for example, introducing a compound of the invention into a sample containing a cellular or purified preparation containing the TPH1 enzyme.

As used herein, the term “individual” or “patient,” used interchangeably, refers to any animal, including mammals, preferably mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, or primates, and most preferably humans.

As used herein, the phrase “therapeutically effective amount” refers to the amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal, individual or human that is being sought by a researcher, veterinarian, medical doctor or other clinician.

As used herein the term “treating” or “treatment” refers to 1) inhibiting the disease; for example, inhibiting a disease, condition or disorder in an individual who is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (*i.e.*, arresting further development of the pathology and/or symptomatology), or 2) ameliorating the disease; for example, ameliorating a disease, condition or disorder in an individual who is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (*i.e.*, reversing the pathology and/or symptomatology).

As used herein the term “preventing” or “prevention” refers to inhibiting onset or worsening of the disease; for example, in an individual who may be predisposed to the disease, condition or disorder but does not yet experience or display the pathology or symptomatology of the disease.

Combination Therapy

One or more additional pharmaceutical agents or treatment methods can be used in combination with the compounds of the present invention for treatment or prevention of various diseases, disorders or conditions disclosed herein. The agents can be combined with the present

compounds in a single dosage form, or the agents can be administered simultaneously or sequentially in separate dosage forms.

Example pharmaceutical agents that may be useful in a combination therapy for blood disorders like blood cancers include parathyroid hormone, anti-sclerostin antibodies, cathepsin K inhibitors, and anti-Dickopf 1.

Example pharmaceutical agents that may be useful in a combination therapy for cancer include leuprolide, goserelin, buserelin, flutamide, nilutamide, ketoconazole, aminoglutethimide, mitoxantrone, estramustine, doxorubicin, etoposide, vinblastine, paclitaxel, carboplatin, and vinorelbine. Therapies that can be combined with TPH inhibition include radiation therapy, high-intensity focused ultrasound, or surgery (e.g., removal of diseased tissues). Other drugs for use in treating cancer include testolactone, anastrozole, letrozole, exemestane, vorozole, formestane, fadrozole, GnRH-analogues, temozolomide, bavituximab, cyclophosphamide, fluorouracil, fulvestrant, gefitinib, trastuzumab, IGF-1 antibodies, lapatinib, methotrexate, olaparib, BSI-201, pazopanib, rapamycin, ribavirin, sorafenib, sunitinib, tamoxifen, docetaxel, vatalinib, bevacizumab, and octreotide.

Example pharmaceutical agents that may be useful in combination therapy for cardiovascular or pulmonary diseases include endothelin receptor antagonists such as ambrisentan, BMS-193884, bosentan, darusentan, SB-234551, sitaxsentan, tezosentan and macitentan. Anticoagulants such as warfarin, acenocoumarol, phenprocoumon, phenindione, heparin, fondaparinux, argatroban, bivalirudin, lepirudin, and ximelagatran may also be useful in combination therapy. Pharmaceutical agents for combination therapy further include calcium channel blockers like amlodipine, felodipine, nicardipine, nifedipine, nimodipine, nisoldipine, nitrendipine, lacidipine, lercanidipine, phenylalkylamines, verapamil, gallopamil, diltiazem, and menthol. Prostacyclins like epoprostenol, iloprost and treprostinil may also be combined with the TPH inhibitors of the invention. Further pharmaceutical agents for combination therapy in cardiovascular or pulmonary diseases include PDE5 inhibitors like sildenafil, tadalafil, and vardenafil; diuretics like furosemide, ethacrynic acid, torasemide, bumetanide, hydrochlorothiazide, spironolactone, mannitol, nitric oxide or nitric oxide releasers, and soluble guanylate cyclase stimulators, such as riociguat. Yet further pharmaceutical agents for combination therapy include APJ receptor agonists (WO 2013/111110); IP receptor agonists

(WO 2013/105057; WO 2013/105066; WO 2013/105061; WO 2013/105063; WO 2013/105065; WO 2013/105058); and PDGF receptor inhibitors (WO 2013/030802).

Example pharmaceutical agents that may be useful in combination therapy for metabolic disorders include HSL inhibitors such as those disclosed in International Patent Publications
5 WO2006/074957; WO2005/073199; WO2004/11 1031; WO2004/111004; WO2004/035550; WO2003/051841; WO2003/051842; and WO2001/066531.

Example pharmaceutical agents that may be useful in combination therapy for bone disorders and diseases include bisphosphonates such as etidronate, clodronate, tiludronate, pamidronate, neridronate, olpadronate, alendronate, ibandronate, risedronate, cimadronate,
10 zoledronate, and the like. Serotonin receptor modulators, such as 5-HT_{1B}, 5-HT_{2A}, and 5-HT_{2B} agonists or antagonists, may also be useful in combination therapy for bone disease. Other useful agents for combination therapy include selective serotonin reuptake inhibitors (SSRI), anti-serotonin antibodies, and beta blockers such as IPS339, ICII 18,551, butaxamine, metipranolol, nadol, oxprenolol, penbutolol, pindolol, propranolol, timolol, and sotalol. Further
15 useful agents for combination therapy for the treatment of bone disorders, such as osteoporosis, include teriparatide, strontium ranelate, raloxifene, and denosumab.

Administration, Pharmaceutical Formulations, Dosage Forms

The compounds of the invention can be administered to patients (animals and humans) in
20 need of such treatment in appropriate dosages that will provide prophylactic and/or therapeutic efficacy. The dose required for use in the treatment or prevention of any particular disease or disorder will typically vary from patient to patient depending on, for example, particular compound or composition selected, the route of administration, the nature of the condition being treated, the age and condition of the patient, concurrent medication or special diets then being
25 followed by the patient, and other factors. The appropriate dosage can be determined by the treating physician.

A compound of this invention can be administered orally, subcutaneously, topically, parenterally, by inhalation spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. Parenteral
30 administration can involve subcutaneous injections, intravenous or intramuscular injections or infusion techniques.

Treatment duration can be as long as deemed necessary by a treating physician. The compositions can be administered one to four or more times per day. A treatment period can terminate when a desired result, for example a particular therapeutic effect, is achieved. Or a treatment period can be continued indefinitely.

5 In some embodiments, the pharmaceutical compositions can be prepared as solid dosage forms for oral administration (e.g., capsules, tablets, pills, dragees, powders, granules and the like). A tablet can be prepared by compression or molding. Compressed tablets can include one or more binders, lubricants, glidants, inert diluents, preservatives, disintegrants, or dispersing agents. Tablets and other solid dosage forms, such as capsules, pills and granules, can include
10 coatings, such as enteric coatings.

 Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable aqueous or organic solvents, or mixtures thereof, and powders. Liquid dosage forms for oral administration can include, for example, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. Suspensions
15 can include one or more suspending agents

 Dosage forms for transdermal administration of a subject composition include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants.

 Compositions and compounds of the present invention can be administered by aerosol which can be administered, for example, by a sonic nebulizer.

20 Pharmaceutical compositions of this invention suitable for parenteral administration include a compound of the invention together with one or more pharmaceutically acceptable sterile isotonic aqueous or non-aqueous solutions, dispersions, suspensions or emulsions. Alternatively, the composition can be in the form of a sterile powder which can be reconstituted into a sterile injectable solutions or dispersion just prior to use.

25 The invention will be described in greater detail by way of specific examples. The following examples are offered for illustrative purposes, and are not intended to limit the invention in any manner. Those of skill in the art will readily recognize a variety of non-critical parameters which can be changed or modified to yield essentially the same results. The compounds of the Examples were found to be inhibitors of TPH1 as described below.

EXAMPLES

The compounds described herein can be prepared in a number of ways based on the teachings contained herein and synthetic procedures known in the art. In the description of the synthetic methods described below, it is to be understood that all proposed reaction conditions, including choice of solvent, reaction atmosphere, reaction temperature, duration of the experiment and workup procedures, can be chosen to be the conditions standard for that reaction, unless otherwise indicated. It is understood by one skilled in the art of organic synthesis that the functionality present on various portions of the molecule should be compatible with the reagents and reactions proposed. Substituents not compatible with the reaction conditions will be apparent to one skilled in the art, and alternate methods are therefore indicated. The starting materials for the examples are either commercially available or are readily prepared by standard methods from known materials.

¹H NMR Spectra were acquired on one or more of three instruments: (1) Agilent UnityInova 400 MHz spectrometer equipped with a 5 mm Automation Triple Broadband (ATB) probe (the ATB probe was simultaneously tuned to ¹H, ¹⁹F and ¹³C); (2) Agilent UnityInova 500 MHz spectrometer; or (3) Varian Mercury Plus 400 MHz spectrometer. Several NMR probes were used with the 500 MHz NMR spectrometer, including both 3 mm and 5 mm ¹H, ¹⁹F and ¹³C probes and a 3 mm X¹H¹⁹F NMR probe (usually X is tuned to ¹³C). For typical ¹H NMR spectra, the pulse angle was 45 degrees, 8 scans were summed and the spectral width was 16 ppm (-2 ppm to 14 ppm). Typically, a total of about 32768 complex points were collected during the 5.1 second acquisition time, and the recycle delay was set to 1 second. Spectra were collected at 25 °C. ¹H NMR Spectra were typically processed with 0.3 Hz line broadening and zero-filling to about 131072 points prior to Fourier transformation. Chemical shifts were expressed in ppm relative to tetramethylsilane. The following abbreviations are used herein: br = broad signal, s = singlet, d = doublet, dd = double doublet, ddd = double double doublet, dt = double triplet, t = triplet, td = triple doublet, tt = triple triplet q = quartet, m = multiplet.

Liquid chromatography - mass spectrometry (LCMS) experiments to determine retention times and associated mass ions were performed using one or more of the following Methods A, B, and C:

Method A: Waters BEH C18, 3.0 x 30 mm, 1.7 μm, was used at a temperature of 50 °C and at a flow rate of 1.5 mL/min, 2 μL injection, mobile phase: (A) water with 0.1% formic acid

and 1% acetonitrile, mobile phase (B) MeOH with 0.1% formic acid; retention time given in minutes. Method A details: (I) ran on a Binary Pump G1312B with UV/Vis diode array detector G1315C and Agilent 6130 mass spectrometer in positive and negative ion electrospray mode with UV PDA detection with a gradient of 15-95% (B) in a 2.2 min linear gradient (II) hold for 0.8 min at 95% (B) (III) decrease from 95-15% (B) in a 0.1 min linear gradient (IV) hold for 0.29 min at 15% (B);

Method B: An Agilent Zorbax Bonus RP, 2.1 x 50 mm, 3.5 μ m, was used at a temperature of 50 °C and at a flow rate of 0.8 mL/min, 2 μ L injection, mobile phase: (A) water with 0.1% formic acid and 1% acetonitrile, mobile phase (B) MeOH with 0.1% formic acid; retention time given in minutes. Method details: (I) ran on a Binary Pump G1312B with UV/Vis diode array detector G1315C and Agilent 6130 mass spectrometer in positive and negative ion electrospray mode with UV-detection at 220 and 254 nm with a gradient of 5-95% (B) in a 2.5 min linear gradient (II) hold for 0.5 min at 95% (B) (III) decrease from 95-5% (B) in a 0.1 min linear gradient (IV) hold for 0.29 min at 5% (B).

Method C: An API 150EX mass spectrometer linked to a Shimadzu LC-10AT LC system with a diode array detector was used. The spectrometer had an electrospray source operating in positive and negative ion mode. LC was carried out using an Agilent ZORBAX XDB 50 x 2.1 mm C18 column and a 0.5 mL/minute flow rate. Solvent A: 95% water, 5% acetonitrile containing 0.01% formic acid; Solvent B: acetonitrile. The gradient was shown as below. 0-0.5 min: 2% solvent (B); 0.5-2.5 min: 2% solvent B to 95% solvent (B); 2.5-4.0 min: 95% solvent (B); 4.0-4.2 min: 95% solvent (B) to 2% solvent B; 4.2-6.0 min: 2% solvent (B).

Microwave experiments were carried out using a Biotage Initiator™, which uses a single-mode resonator and dynamic field tuning. Temperatures from 40-250 °C were achieved, and pressures of up to 20 bars were reached.

Preparative HPLC purification was carried out using either a C18-reverse-phase column from Genesis (C18) or a C6-phenyl column from Phenomenex (C6 Ph) (100 × 22.5 mm i.d. with 7 micron particle size, UV detection at 230 or 254 nm, flow 5-15mL/min), eluting with gradients from 100-0 to 0-100 % water/acetonitrile or water/MeOH containing 0.1% formic acid. Fractions containing the required product (identified by LCMS analysis) were pooled, the organic fraction removed by evaporation, and the remaining aqueous fraction lyophilised, to give the product.

Chiral HPLC was carried out using a Chiralpak AD column, 4.4 mm x 250 mm, particle size 5 micron

Compounds which required column chromatography were purified manually or fully automatically using either a Biotage SP1TM Flash Purification system with Touch Logic

- 5 ControlTM or a Combiflash Companion[®] with pre-packed silica gel Isolute[®] SPE cartridge, Biotage SNAP cartridge or Redisep[®] Rf cartridge respectively.

Preparation of alcohols and amines


- 10 The chiral alcohols drawn below are shown in their absolute configuration (unless otherwise shown). Their enantiopurity (% ee) can be determined via Mosher ester analysis and analyzed as described in the literature (Dale, J. A. & Mosher, H. S. Nuclear Magnetic Resonance Enantiomer Regents. Configurational Correlations Via Nuclear Magnetic Resonance Chemical Shifts Of Diastereomeric Mandelate, O-Methylmandelate, and alpha-Methoxy alpha-Trifluoromethylphenylacetate (MTPA) Esters. *J. Am. Chem. Soc.* **95**, 512–519 (1973)).
- 15 The chiral alcohols of the invention are preferably enantiomerically enriched, for example, to $\geq 95\%$ ee.

Representative Mosher ester preparation

- To a solution of (R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol (46 mg, 0.20 mmol, Intermediate 3) was added pyridine (138 mg, 1.7 mmol)
- 20 followed by the addition of either (S or R)- α -methoxy- α -trifluoromethyl-phenylacetyl chloride (10 mg, 0.40 mmol). The reaction was stirred for 12 h, then the material was purified directly on silica gel chromatography (EtOAc/heptane) to provide the “Mosher ester” which was analyzed by ¹H NMR for enantiomeric purity.

25

Intermediate 1: (R)-1-(4-Bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol



Step 1: Potassium *t*-butoxide (16.3 g, 145 mmol) was dissolved in DMSO (100 mL). To this solution was added 3-methyl pyrazole (10.4 g, 120 mmol) and the reaction was heated at 50 °C for 30 min. 1,4-Dibromo-2-fluorobenzene (31 g, 120 mmol) was then added and the reaction stirred at 50 °C for 16 h. The reaction was cooled to RT and extracted with water and EtOAc, washed with brine, dried over Na₂SO₄, and then filtered and concentrated *in vacuo*. Purification by normal phase silica gel column chromatography (EtOAc/heptane) provided 1-(2,5-dibromophenyl)-3-methyl-1H-pyrazole.


Step 2: 1-(2,5-dibromophenyl)-3-methyl-1H-pyrazole (23.0 g, 73 mmol) from Step 1 was dissolved in 200 mL of THF and cooled to 0 °C. *i*-Propyl magnesium chloride (2.0 M in THF, 40 mL) was added dropwise and the reaction was stirred for 45 min, then ethyl trifluoroacetate (10.5 mL) was added. The reaction was stirred for 30 min at 0 °C, then 10% HCl is added dropwise (400 mL). The reaction was extracted with water and EtOAc, washed with brine, dried over Na₂SO₄, filtered, and then concentrated *in vacuo*. Purification by normal phase silica gel column chromatography (EtOAc/heptane) provided 1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanone.

Step 3: METHOD A: Pentamethylcyclopentadienyl iridium (III) chloride dimer (CAS# 12354-84-6) (10.4 mg) and (1R,2R)-(-)-N-(4-toluene sulfonyl)-1,2-diphenyl ethylene diamine (CAS# 144222-34-4) (9.2 mg) were combined in water (120 mL), then heated to 50 °C for 5 h to provide the “Iridium complex.” 1-[4-Bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl]-2,2,2-trifluoroethanone (16 g, 48 mmol) was dissolved in acetonitrile (120 mL) to which the Iridium complex and potassium formate (3.1 g, 3.7 mmol) were added. The reaction mixture was heated to 50 °C for 8 h. The reaction mixture was then cooled to RT, partitioned between water and EtOAc, and extracted. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Recrystallization from hot heptane (200 mL) provided the title compound.

METHOD B: Alternatively, the trifluoromethyl (or other prochiral) ketones of formula **G** or **J** (scheme 2) were asymmetrically reduced as follows (see for example: Corey, E. J. & Link, J. O. A General, Catalytic, and Enantioselective Synthesis of Alpha-amino Acids. *J. Am. Chem. Soc.*

114, 1906–1908 (1992)): Catechol borane (95 mL, 1 M in THF) and (S)-2-methyl-CBS oxazaborolidine (2.6 g, 9.6 mmol) were mixed in a jacketed glass reactor. The mixture was stirred at RT for 20 min, then the jacket was cooled to -78 °C. At a reaction temperature of -65 °C, 1-[4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl]-2,2,2-trifluoroethanone (16 g, 48 mmol) in THF (150 mL) was added dropwise over 2 h. The reaction was then warmed to -36 °C and held at this temperature for 22 h. Then the reaction was quenched with 3 N NaOH (100 mL) while maintaining a reaction temperature of <-25 °C. The reaction was then warmed to 0 °C and H₂O₂ (30%, 100 mL) was added over 30 min, then warmed to RT for 4 h. The reaction mixture was quenched with 1 N NaOH, extracted with ether, washed with brine, dried over Na₂SO₄, and concentrated *in vacuo*. Purification on normal phase silica gel chromatography (EtOAc/heptane) provided the product as a viscous oil.

Intermediate 2: (R)-1-(5-Bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol




Step 1: Diisopropylamine (4.40 mL, 31.4 mmol) was dissolved in THF (28 mL) and cooled to -40 °C. Then n-butyllithium (12.6 mL, 2.5 M in hexanes, 31.4 mmol) was added dropwise, and the reaction was stirred at -40 °C for 1 h, then cooled to -78 °C. A solution of 1-bromo-4-fluorobenzene (5 g, 28.6 mmol) in THF (6.0 mL) was added, and the reaction was stirred at -78 °C for 1 h. Trifluoroacetic acid ethyl ester (3.73 mL, 31.4 mmol) in THF (6.0 mL) was then added, and the reaction was slowly warmed to 0 °C over an hour. The reaction was quenched with NH₄Cl (aq. sat), and extracted with EtOAc, washed with brine, and dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column chromatography (EtOAc/heptane) provided 1-(5-bromo-2-fluorophenyl)-2,2,2-trifluoroethanone.

Step 2: 1-(5-bromo-2-fluorophenyl)-2,2,2-trifluoroethanone (2.20 g, 8.12 mmol) from Step 1, K₂CO₃ (1.68 g, 12.2 mmol), and 3-methyl-1H-pyrazole (1.33 g, 16.2 mmol) were stirred in

toluene (10 mL). The reaction was then heated to 110 °C for 16 h. The reaction was cooled, and water and EtOAc were added. The toluene-EtOAc layer is removed *in vacuo*, and then the reaction is extracted with water and EtOAc, washed with brine, and dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column chromatography (EtOAc/heptane) provided 1-[5-bromo-2-(3-methyl-pyrazol-1-yl)-phenyl]-2,2,2-trifluoroethanone.

Step 3: The title compound was prepared using the Iridium complex-catalyzed hydrogenation as described for Intermediate 1, (R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol.

Intermediate 3: (R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol




Step 1: Potassium *t*-butoxide (3.9 g, 0.33 mmol) was dissolved in DMSO (25 mL). To this solution was added 3-methyl pyrazole (2.7 g, 0.33 mmol) and the reaction was heated at 50 °C for 30 min. 1-Bromo-4-chloro-2-fluorobenzene (4.6 g, 0.22 mmol) was then added and the reaction was stirred at 50 °C for 16 h. The reaction was cooled to RT and extracted with water and EtOAc, washed with brine, and dried over Na₂SO₄, filtered and concentrated *in vacuo*. Purification by normal phase silica gel column chromatography (EtOAc/heptane) provided 1-(2-bromo-5-chlorophenyl)-3-methyl-1H-pyrazole and 1-(2-bromo-5-chlorophenyl)-5-methyl-1H-pyrazole as a 4:1 mixture that was used in the next step directly.

Step 2: The mixture from Step 1 (8 g, 0.39 mmol) was dissolved in 160 mL of THF and cooled to 0 °C. *i*-Propyl magnesium chloride (2.0 M in THF, 23 mL) was added dropwise and the reaction stirred for 45 min, then ethyl trifluoroacetate (6 mL) was added. The reaction was stirred for 30 min at 0 °C, then 10% HCl was added dropwise (40 mL). The reaction was extracted with water and EtOAc, washed with brine, and dried over Na₂SO₄, filtered, and concentrated *in vacuo*.

Purification by normal phase silica gel column chromatography (EtOAc/heptane) provided 1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanone as a white solid.


Step 3: The title compound was prepared using the Iridium complex-catalyzed hydrogenation, as described for Intermediate 1 (R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol.

Intermediate 4: (R)-1-(5-chloro-2-(2,2,2-trifluoro-1-hydroxyethyl)phenyl)pyrrolidin-2-one



To a solution of (R)-1-(4)-2,2,2-trifluoroethanol (300 mg, 1.04 mmol) in toluene (7 mL) was added pyrrolidin-2-one (89 mg, 1.04 mmol), (1S,2S)-N1,N2-dimethylcyclohexane-1,2-diamine (74 mg, 0.52 mmol), CuI (50 mg, 0.26 mmol) and K₂CO₃ (360 mg, 2.6 mmol). The reaction was heated in a sealed tube to 130 °C for 12 h and then cooled to RT. The solids were filtered and the product was purified by normal phase silica gel chromatography (EtOAc:petroleum ether) to provide the title compound as a white solid.

Intermediate 5: (R)-2,2,2-Trifluoro-1-(2-methyl-1H-benzo[d]imidazol-4-yl)ethanol




Step 1: 4-Bromo-2-methyl-1H-benzimidazole (500 mg, 2.37 mmol) was dissolved in THF (8 mL) and cooled to -78 °C. *n*-Butyllithium (2.3 mL, 2.5 M in hexanes, 5.7 mmol) was added dropwise and the reaction was stirred at -78 °C for 30 min. Trifluoroacetic acid ethyl ester (339 μL, 2.8 mmol) was added and the reaction was stirred at 0 °C for 1 h. The reaction was quenched with HCl (2 N, 4 mL), then extracted with water and EtOAc, washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column chromatography (CH₂Cl₂/MeOH/NH₄OH) provided 2,2,2-trifluoro-1-(2-methyl-1H-benzoimidazol-4-yl)-ethanone.

Step 2: The title compound was prepared using the Iridium complex-catalyzed hydrogenation, as described for Intermediate 1 (R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol.

5

Intermediate 6: 1-(4-Chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethanol




Step 1: 1-(2-bromo-5-chlorophenyl)-3-methyl-1H-pyrazole/1-(2-bromo-5-chlorophenyl)-5-methyl-1H-pyrazole mixture (Intermediate 3, step 1) (1.00 g, 3.68 mmol) was dissolved in THF (6 mL) and cooled to 0 °C. *i*-Propyl magnesium chloride (2.76 mL, 2.0 M in THF, 5.52 mmol) was added dropwise and allowed to warm to RT over 30 min. The reaction was then cooled to -15 °C. Acetyl chloride (481 μ L, 5.5 mmol) was added and the reaction was warmed to RT for 3 h. The reaction was quenched with HCl (2 N, 4 mL), then extracted with water and EtOAc, washed with brine, and dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column chromatography (EtOAc/heptane) provided 1-[4-chloro-2-(3-methyl-pyrazol-1-yl)-phenyl]-ethanone.

Step 2: 1-[4-Chloro-2-(3-methyl-pyrazol-1-yl)-phenyl]-ethanone (400 mg, 1.70 mmol) from Step 1 was dissolved in MeOH (10 mL) and cooled to 0 °C. NaBH₄ (129 mg, 3.41 mmol) was added portionwise, then the reaction was warmed to RT, stirred for 30 min, then quenched with acetone. The MeOH was removed *in vacuo* then the residue was partitioned between water and EtOAc and extracted several times. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column chromatography (CH₂Cl₂/MeOH/NH₄OH) provided the title compound.


25

Intermediate 7: 1-(2,6-dibromophenyl)ethanol




To a solution of 1-(2,6-dibromophenyl)-2,2,2-trifluoroethanone (CAS# 1208078-23-2) (3 g, 9 mmol) in EtOH (50 mL) was added NaBH₄ (340 mg, 9 mmol) at 5 °C. The reaction was warmed to RT for 1 h, then extracted with EtOAc NaHCO₃, brine, and dried over Na₂SO₄ filtered and concentrated *in vacuo* to provide 1-(2,6-dibromophenyl)-2,2,2-trifluoroethanol as a light yellow oil.

Intermediate 8: 1-(2,5-dibromophenyl)ethanol



This compound was made as described above for Intermediate 7, 1-(2,6-dibromophenyl)-2,2,2-trifluoroethanol, starting with 1-(2,5-dibromophenyl)-2,2,2-trifluoroethanone to provide a light yellow oil.

Intermediate 9: (4-Chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)methanol







Step 1: 1-(2-bromo-5-chlorophenyl)-3-methyl-1H-pyrazole/1-(2-bromo-5-chlorophenyl)-5-methyl-1H-pyrazole mixture (Intermediate 3, step 1) (1.00 g, 3.68 mmol) was dissolved in THF (6 mL) then cooled to 0 °C. *i*-Propyl magnesium chloride (2.76 mL, 2.0 M in THF, 5.52 mmol) was added dropwise and the reaction was warmed to RT for 30 min. The reaction was then cooled to -15 °C and paraformaldehyde (166 mg, 5.5 mmol) was added. The reaction mixture was allowed to warm to RT and stirred for 1 h. DMF (500 mL) was added and the reaction was stirred for an additional 1 h. The reaction was quenched with HCl (2 N, 4 mL), diluted with

water, extracted with EtOAc, washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column chromatography (EtOAc/heptane) provided 4-chloro-2-(3-methyl-pyrazol-1-yl)-benzaldehyde.


- 5 *Step 2:* 4-Chloro-2-(3-methyl-pyrazol-1-yl)-benzaldehyde (446 mg, 2.03 mmol) from Step 1 was dissolved in MeOH (14 mL) and cooled to 0 °C. NaBH₄ (175 mg, 4.61 mmol)) was added portionwise. The reaction mixture was allowed to warm to RT, and after 90 min was quenched with acetone. The MeOH was removed *in vacuo*. The residue was partitioned between water and EtOAc and then extracted. The combined organic layers were washed with brine, dried over
- 10 Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column chromatography (EtOAc/heptane) provided the title compound.

- Using the procedure described for Intermediate 3, (R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol, the following alcohols (Intermediates 10-15) shown
- 15 in the Table below were prepared starting with the appropriately substituted 1-bromo-2-fluorobenzene.

No.	Name	Structure	LCMS (MH ⁺)
Intermediate 10	(R)-2,2,2-trifluoro-1-(4-methyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethanol		271
Intermediate 11	(R)-2,2,2-trifluoro-1-(4-methoxy-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethanol		287

Intermediate 12	(R)-1-(3-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol		291
Intermediate 13	(R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(trifluoromethyl)phenyl)ethanol		325
Intermediate 14	(R)-2,2,2-trifluoro-1-(4-fluoro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethanol		274
Intermediate 15	(R)-2,2,2-trifluoro-1-(6-methyl-2-(3-methyl-1H-pyrazol-1-yl)pyridin-3-yl)ethanol		272

Intermediate 16: (2-Phenoxy-6-(piperidin-1-yl)phenyl)methanamine




Step 1: To a solution of phenol (415 mg, 4.5 mmol) in 60 mL of DMF was added NaH (60%, 6.0 mmol) at 0 °C. The reaction was stirred for 1 h, then 2-fluoro-6-(piperidin-1-yl)benzonitrile (CAS# 646989-68-6) (612 mg, 3.0 mmol) was added and the reaction stirred for 48 h at RT. The reaction mixture was then diluted with water and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, then concentrated *in vacuo*.

Purification by normal phase silica gel column chromatography (EtOAc/heptane) provided 2-phenoxy-6-(piperidin-1-yl)benzonitrile as an off-white solid.

Step 2: To 2-phenoxy-6-(piperidin-1-yl)benzonitrile (250 mg, 0.9 mmol) from Step 1 in 20 mL of MeOH was added Raney Nickel (5%) and NH₄OH (2 mL). The reaction was stirred under 1 atm of H₂ at RT for 2 h. The solid was filtered away and the filtrate was concentrated *in vacuo* to provide the title compound as a viscous oil.


Intermediate 17: (R)-1-(4-Chloro-2-(2-methoxyethoxy)phenyl)-2,2,2-trifluoroethanol



Step 1: 1-Bromo-4-chloro-2-(2-methoxy-ethoxy)-benzene (CAS# 1245563-20-5) (5.00 g, 18.8 mmol) was dissolved in THF (30 mL) and cooled to 0 °C. *i*-Propylmagnesium bromide (11.3 mL, 2.0 M in THF, 22.6 mmol) was added dropwise, and the reaction was stirred at 10 °C for 30 min, then warmed to RT for 16 h. The reaction was then cooled to -15 °C and trifluoroacetic acid ethyl ester (3.37 mL, 28.2 mmol) was added. The reaction was stirred at 10 °C for 1 h. The reaction was quenched with HCl (2 N, 38 mL) at 0 °C. The reaction mixture was diluted with water and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, then concentrated *in vacuo*. Purification by normal phase silica gel column chromatography (EtOAc/heptane) provided 1-(4-chloro-2-(2-methoxyethoxy)phenyl)-2,2,2-trifluoroethanone.


Step 2: The title compound was prepared using the Iridium complex-catalyzed hydrogenation as described for Intermediate 1 (R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol.

Intermediate 18: (R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethanol




To a solution of (R)-1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethanol (300 mg, 1.1 mmol) in dioxane (12 mL) was added phenyl boronic acid (185 mg, 1.5 mmol), Pd₂(dppf)Cl₂ (35 mg, 0.07 mmol) and Na₂CO₃ (3 mL, 2.0 M, aq). The reaction was heated to 90 °C for 2 h, then cooled to RT, and concentrated *in vacuo*. The residue was taken up in CH₂Cl₂, washed with brine, and extracted with CH₂Cl₂. The combined organic layers were dried over Na₂SO₄. Purification by normal phase silica gel column (EtOAc/hexanes) to provide a white solid.

Intermediate 19: (R)-1-(4-chloro-2-(5-chlorothiophen-2-yl)phenyl)-2,2,2-trifluoroethanol



This compound was made in the same way as described for Intermediate 18 (R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethanol to provide a white solid.

Intermediate 20: (R)-2,2,2-trifluoro-1-(6-methyl-2-(3-methyl-1H-pyrazol-1-yl)pyridin-3-yl)ethanol



Step 1: To the solution of 2-chloro-6-methylnicotinic acid (5 g, 29.1 mmol) in CH₂Cl₂ (40 mL) was added oxalyl dichloride (8.1 g, 63.8 mmol) at 0 °C and the reaction mixture was stirred for 2 h. The mixture was concentrated and 40 mL of methanol was then added at 0 °C and the reaction mixture was stirred at RT for 12 h. The mixture was then concentrated *in vacuo* and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered,

and concentrated *in vacuo* to provide methyl 2-chloro-6-methylnicotinate that is used without further purification as a light yellow solid.


Step 2: To a solution of 3-methyl-1H-pyrazole (1.1 g, 13.4 mmol) in DMF (5 ml) was added sodium hydride (1.0 g, 60% in oil) at 0 °C. The reaction mixture was stirred for 1 h at 0 °C and then . A solution of methyl 2-chloro-6-methylnicotinate (4.3 g, 23.16 mmol) in DMF (5 mL) was added dropwise to the reaction mixture at 0 °C. After addition, the mixture was heated to 80 °C and stirred for 12 h. After this time, the mixture was poured into ice-water and extracted and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated in vacuo and then purified by normal phase silica gel column (EtOAc/hepate) to provide methyl 6-methyl-2-(3-methyl-1H-pyrazol-1-yl)nicotinate as a brown semi-solid.

Step 3: To a solution of methyl 6-methyl-2-(3-methyl-1H-pyrazol-1-yl)nicotinate (3.7 g, 16 mmol) and trimethyl(trifluoromethyl)silane (11.4 g, 80.2 mmol) in toluene (60 ml), was added dropwise at -78°C and then the solution of tetrabutyl ammonium fluoride (1.6 mL, 1.0 M in THF) was added dropwise to the reaction mixture at -78 °C. After addition, the mixture was warmed slowly up to RT and stirred for 12 h. The reaction mixture was concentrated and the resulting residue was dissolved in methanol (30 mL). 6 N HCl (30 mL) was added to the reaction mixture and the resulting mixture was stirred for 2 h. The reaction mixture was concentrated, adjusted to pH 6 with sat. NaHCO₃ and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated in vacuo and purified by normal phase silica gel column (EtOAc/hepate) to provide 2,2,2-trifluoro-1-(6-methyl-2-(3-methyl-1H-pyrazol-1-yl)pyridin-3-yl)ethanone as a brown semi-solid.

Step 4: A solution of (S)-(-)-2-Butyl-CBS-oxazaborolidine solution (3.0 ml 1.0 M in toluene) and catecholborane (30 ml 1.0 M in THF) was stirred at RT for 30 min. The mixture was then cooled to -70 °C and 2,2,2-trifluoro-1-(6-methyl-2-(3-methyl-1H-pyrazol-1-yl)pyridin-3-yl)ethanone (1 g, 2.9 mmol) in THF (16 mL) was added dropwise. After addition, the reaction mixture was warmed up to -32 °C and stirred for 12 h. After this time, 3N NaOH (18 mL) was added followed by H₂O₂ (18 mL) and the temperature of the reaction mixture was increased to

RT for 30 min and then extracted with ethyl. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated in vacuo and purified by normal phase silica gel column (EtOAc/hepate) to provide the title compound as a yellow solid.

5 **Intermediate 38: (R)-1-(5-bromo-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethanol**



- 10 *Step 1:* A solution of 2,4-dibromo-benzoic acid, (2.3 g, 18.8 mmol), phenyl boronic acid (5 g, 17.9 mmol), Pd₂(dba)₃ (818 mg, 8.9 mmol) and LiOH (1.65 g, 39.3 mmol) in a 1:1 mixture of NMP/water (100 mL) was heated to 70 °C for 2 d. After this time, the reaction mixture was cooled to RT, and the reaction mixture was adjusted to pH = 4-5 with 3 N HCl. The mixture was then extracted with ethyl acetate and the combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo* and purified by normal phase silica gel column (EtOAc/PE 10:1 to 1:1) to afford 5-bromo-[1,1'-biphenyl]-2-carboxylic acid as a colorless oil.
- 15 *Step 2:* To a solution of 5-bromo-[1,1'-biphenyl]-2-carboxylic acid (5 g, 18.2 mmol) in MeOH (30 mL) was added SOCl₂ (10 mL) dropwise. The reaction mixture was heated to 70 °C for 2 h, then cooled to RT. The mixture was concentrated, adjusted to pH= 7-8 with saturated aqueous NaHCO₃ and extracted with ethyl acetate. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo* and purified by normal phase silica gel column (EtOAc/PE 50:1) to afford methyl 5-bromo-[1,1'-biphenyl]-2-carboxylate as a colorless oil.
- 20

- 25 *Step 3:* A solution of methyl 5-bromo-[1,1'-biphenyl]-2-carboxylate (2.2 g, 6.9 mmol) in THF (50 mL) was cooled to 0 °C. LiAlH₄ (380 mg, 10 mmol) was added slowly. The reaction mixture was stirred at RT for 2 h, after which water (1 mL) was added slowly to quench the reaction. The solid was removed by filtration and the filtrate was concentrated *in vacuo* to provide (5-bromo-[1,1'-biphenyl]-2-yl)methanol as a white solid that was used directly without further purification.





Step 4: To a solution of (5-bromo-[1,1'-biphenyl]-2-yl)methanol (2.0 g, 8.4 mmol) in CH₂Cl₂ (30 mL) was added Dess-Martin Periodinane (4.3 g, 10 mmol). The reaction mixture was stirred at RT for 2 h and then the solids were filtered and the resultant filtrate was concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc:PE = 1:50) afforded 5-bromo-[1,1'-biphenyl]-2-carbaldehyde as a colorless oil.

Step 5: To a solution of 5-bromo-[1,1'-biphenyl]-2-carbaldehyde (1.9 g, 7.3 mmol) and was added TMSCF₃ (1.2g, 8.7 mmol) in THF (20 mL) and cooled to 0 °C. To this solution was added TBAF (1.46 mL, 1M in THF) and the reaction mixture was warmed to RT for 3 h. After this time, the mixture was treated with 3 N HCl (5 mL) and stirred for 12 h. Then the reaction mixture was diluted with water and extracted with ethyl acetate. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, concentrated *in vacuo* and purified by normal phase silica gel column (EtOAc:PE = 1:10) to afford 1-(5-bromo-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethanol as a colorless oil.

Step 6: To a solution of 1-(5-bromo-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethanol (1.8 g, 5.5 mmol) in CH₂Cl₂ (30 mL) was added Dess-Martin Periodinane (3g, 7.1 mmol). The reaction mixture was stirred at RT for 2 h and then the solids were filtered. The resultant filtrate was concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc:PE = 1:50) afforded 1-(5-bromo-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethanone as a colorless oil.


Step 7: 1-(5-Bromo-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethanone (1.3 g, 3.9 mmol) in CH₃CN (10 mL) was reduced to the chiral alcohol using the chiral iridium catalyst (METHOD A) at RT. The reaction mixture was then charged with potassium formate (725 mg, 8.6 mmol) and the mixture was stirred at 40 °C for 12 h. Then the reaction was diluted with water and extracted with ethyl acetate. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, concentrated *in vacuo* and purified by normal phase silica gel column (EtOAc:PE = 1:10) to afford the title compound as a colorless oil.

Using the procedure described for Intermediate 3, (R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol, the following alcohols (Intermediates 39-42) shown in the Table below were prepared starting with the appropriately substituted pyrazole.

No.	Name	Structure	LCMS (MH ⁺)
Intermediate 39	(R)-1-(4-chloro-2-(3-(trifluoromethyl)-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol		345
Intermediate 40	(R)-1-(2-(3-(tert-butyl)-1H-pyrazol-1-yl)-4-chlorophenyl)-2,2,2-trifluoroethanol		334
Intermediate 41	(R)-1-(4-chloro-2-(3-isopropyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol		319
Intermediate 42	(R)-1-(4-chloro-2-(3-cyclopropyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol		317

5











Intermediate 43: (R)-1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethanol



A solution of dichloro(pentamethylcyclopentadienyl)iridium (III) dimer ($[\text{Cp}^*\text{IrCl}_2]_2$, 14 mg, 0.02 mmol) and (1*R*,2*R*)-(-)-(4-toluenesulfonyl)-1,2-diphenylethylenediamine (14 mg, 0.04 mmol) in water (7 mL) was prepared at RT. The resulting mixture was heated to 40 °C for 3 h to provide a homogeneous orange solution. To this active catalyst solution at 40 °C was added potassium formate (143 mg, 171 mmol), and a solution of 1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethanol (CAS# 1033805-23-0, 98 mg, 0.34 mmol) in CH_3CN (70 mL). The reaction mixture was then stirred at 40 °C for 2 h and then cooled to RT and the layers were separated. The aqueous layer was extracted with MTBE and the combined organic layers were dried over Na_2SO_4 , filtered, and concentrated *in vacuo* to provide the title compound that was used without further purification.

The following alcohols and amines in the table below are useful in preparing compounds of the invention. They are either commercially available or can be prepared by known synthetic procedures. CAS registry numbers are provided for each.

No.	Name	CAS Registry #	Structure	Ex#
21	(R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethanol	1033805-15-0		10a & 10e
22	(R)-1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethanol	1033805-25-2		34a-34ae
23	(R)-1-(5-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol	1033805-72-9		10k


24	1-(adamantan-1-yl)ethanamine	13392-28-4		38
25	(adamantan-1-yl)methanamine	17768-41-1		41c
26	[1,1'-biphenyl]-3-ylmethanamine	177976-49-7		39d
27	naphthalen-2-ylmethanamine	2018-90-8		39a
28	1-(adamantan-1-yl)ethanol	26750-08-3		41c
29	(R)-1-(naphthalen-2-yl)ethanamine	3906-16-9		39e
30	(R)-2,2,2-trifluoro-1-(naphthalen-2-yl)ethanol	68200-42-0		59b
31	[1,1'-biphenyl]-4-ylmethanamine	712-76-5		39b
32	(2-(piperidin-1-yl)phenyl)methanamine	72752-54-6		41a
33	(R)-1-(4-bromophenyl)-2,2,2-trifluoroethanol	80418-12-8		55a- 55db

No. = Intermediate number; Ex# = Used in the preparation of the following example(s)

Preparation of boronic acids and esters

The boronic acids and esters used in biaryl couplings are either commercially available or
5 can be readily synthesized from the corresponding bromide using routine synthetic methods.
The following Intermediate 34 is a representative example.


Intermediate 34: 6-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-3,4-dihydroquinolin-2(1H)-one



To a solution of 6-bromo-3,4-dihydroquinolin-2(1H)-one (200 mg, 0.89 mmol) in 5 mL of acetonitrile was added pinacoldiboron (300 mg, 1.2 mmol), Pd(dppf)₂Cl (30 mg, 0.09mmol), KOAc (250 mg, 2.1 mmol) and triethyl amine (1mL). The reaction was heated to 87 °C for 24 h, then cooled to RT. The solids were filtered away, and the solvent was removed *in vacuo*, then extracted with EtOAc, water, brine and dried over Na₂SO₄. The solvent was removed *in vacuo* to provide an off-white solid which was used without further purification.

Spirocyclic amino esters preparation

Intermediate 35: (S)-2-Benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate




Step 1: (3S)-8-Tert-butyl 3-ethyl 2,8-diazaspiro[4.5]decane-3,8-dicarboxylate [Example 24 in US Pat. Pub. No. 2012/0101280] (50 g, 160 mmol) in CH₂Cl₂ (500 mL), and Et₃N (51.7 g, 512 mmol) was cooled to 0 °C. Benzyl chloroformate (34.1 g, 205 mmol) was added dropwise and the mixture was stirred at 0 °C for 3 h. The reaction mixture was washed with water, extracted with CH₂Cl₂, dried over Na₂SO₄, and concentrated *in vacuo* to provide (S)-2-benzyl 8-tert-butyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3,8-tricarboxylate as a light yellow oil which was used directly without further purification.

Step 2: To a solution of (S)-2-benzyl 8-tert-butyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3,8-tricarboxylate (79 g, 160 mmol, Step 1) in CH₂Cl₂ (400 mL) was added TFA (182 g, 1600 mmol) dropwise at RT. The reaction mixture was stirred for 3 h then concentrated *in vacuo*. The residue was quenched with saturated NaHCO₃ and solid NaHCO₃ was added until no further gas

evolution was noted. The mixture was extracted with EtOAc and the combined organic layers were concentrated *in vacuo*. Purification by normal phase silica gel column chromatography (CH₂Cl₂/MeOH/NH₄OH) provided the title compound as a light yellow solid.


5 **Intermediate 36: (S)-2-Tert-butyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate**



Step 1: (S)-2-Benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (2.4 g, 6.9 mmol) in HCl/dioxane (50 mL, 3.3 N) was stirred for 2 h at RT. The solvent was then removed *in vacuo* to provide (S)-2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate hydrochloride
10 which was used directly without further purification.

Step 2: To a solution of (S)-2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate hydrochloride and BOC₂O (1.5 g, 6.9 mmol) in EtOH (50 mL) was added Pd/C (10%, 2.4 g) and HOAc (cat.). The mixture was degassed and blanked under H₂ then stirred at 45 °C at 50 psi of
15 H₂ for 12 h. The solid was filtered away and the filtrate concentrated *in vacuo* to provide the title compound as a viscous solid.

Intermediate 37: Methyl 3,9-diazaspiro[5.5]undecane-2-carboxylate



20 To a solution of 3,9-diazaspiro[5.5]undecane-2-carboxylic acid, 3-[(4-methoxyphenyl)methyl]-9-(phenylmethyl)-methyl ester [CAS# 1314388-32-3] (50 mg, 0.12 mmol) in MeOH (2 mL) and water (2 mL) was added a catalytic amount of TFA. The mixture was hydrogenated using a H-cube apparatus under 80 °C / 80 bar 2 cycles. The reaction mixture was cooled to RT then concentrated *in vacuo* to provide the title compound as a white solid which is used directly.

25

General synthetic methods

Methods for removal of N-Carbobenzyloxy (N-CBZ) protecting group


Method A - Hydrogenation over Pd/C: To a solution of N-CBZ protected compound (1 eq.) in EtOAc was added HOAc (100 μ L) and 5% (w/w) Pd/C (5 mol%). The reaction mixture was degassed, blanketed under H₂ (balloon) 3 times, then stirred at RT for 2 h. The reaction was then filtered through a pad of celite that was rinsed with 1:9 MeOH:EtOAc. The filtrate was concentrated *in vacuo*. The product was purified by column chromatography using an Isco Gold reversed phase silica cartridge (H₂O:HOAc: 99:1 to MeOH:AcOH 99:1).

Method B - Dealkylation with TMSI: To a solution of N-CBZ protected compound (1 eq.) in CH₃CN was added a solution of TMSI (2.2 eq.) in CH₃CN (0.2 M). The reaction mixture was stirred at RT for 2 h then quenched with 1 N HCl to pH 1. The product was purified by column chromatography using an Isco Gold reversed phase silica cartridge (H₂O:HOAc: 99:1 to MeOH:AcOH 99:1).

General ester hydrolysis with lithium hydroxide: To a solution of an ethyl ester compound (1 eq) in THF (0.18 M) and water (1.4 M) was added LiOH-H₂O (10 eq). The mixture was stirred at RT for 1 h. Water was added and the pH was adjusted to 6.5 with 1 N HCl. THF was removed *in vacuo*, then the solid was precipitated, washed with water, and dried *in vacuo* to yield the corresponding carboxylic acid.

The compounds of the examples were isolated either in the neutral zwitterionic form or as a TFA or HCl salt.

Example 1u: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a solution of (R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol (160 mg, 0.2 mmol, Intermediate 1) in dioxane (2 mL) was added 2-amino-4,6-dichloropyrimidine (100 mg, 0.16 mmol) and Cs₂CO₃ (48 g, 0.16 mmol). The reaction was heated to 80 °C for 16 h, cooled to RT, and filtered. The solvent was removed *in vacuo* and the residue was dissolved in a mixture of CH₂Cl₂ and heptane, concentrated to half the volume, filtered, and concentrated again *in vacuo*. Purification via normal phase silica gel chromatography (CH₂Cl₂/Heptane) provided 4-[(1R)-1-[4-bromo-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoro-ethoxy]-6-chloro-pyrimidin-2-amine as an off-white solid.

Step 2: To a solution of 4-[(1R)-1-[4-bromo-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoro-ethoxy]-6-chloro-pyrimidin-2-amine (125 mg, 0.3 mmol, Step 1) in dioxane (3 mL) was added (S)-2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (95 mg, 0.3 mmol) and Na₂CO₃ (182 mg, 0.35 mmol). The reaction was heated to 90 °C for 130 h, then cooled to RT, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

Step 3: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (300 mg, 0.4 mmol, Step 2) in ethanol (2 mL) and water (0.5 mL) was added phenylboronic acid (143 mg, 0.8 mmol), PdCl₂(PPh₃)₂ (41 mg, 0.058 mmol), and Cs₂CO₃ (390 mg, 1.2 mmol). The reaction was heated to 60 °C for 16 h, then cooled to RT, filtered through celite and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

Step 4: A solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (240 mg, 0.4 mmol, Step 3) in EtOAc (5 mL) was hydrogenated using an H-Cube

apparatus and a 10% (w/w) Pd/C cartridge with a flow rate of 1.0 mL/min at RT. Purification on normal phase silica gel (EtOAc/heptane) provided (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate.


5

Step 5: To a solution of (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (50 mg, 0.08 mmol) from Step 4 in THF (2.0 mL) and water (0.2 mL), was added lithium hydroxide monohydrate (58 mg, 0.05 mmol). The reaction mixture was stirred at RT for 2 h, then the solution was neutralized with 1 N HCl, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided the title compound as an off-white solid as the zwitterionic form.

10

Example 1m: (S)-8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid

15



Step 1: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (Step 2, Example 1u) (300 mg, 0.4 mmol, Step 2) in ethanol (2 mL) and water (0.5 mL) was added (3,4-dimethylphenyl)boronic acid (120 mg, 0.8 mmol), PdCl₂(PPh₃)₂ (41 mg, 0.058 mmol), and Cs₂CO₃ (390 mg, 1.2 mmol). The reaction was heated to 60 °C for 16 h, then cooled to RT, filtered through celite and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

20


25

Step 2: A solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (220 mg, 0.3 mmol) in EtOAc (5 mL) was

hydrogenated using an H-Cube apparatus and a 10% (w/w) Pd/C cartridge with a flow rate of 1.0 mL/min at RT. Purification on normal phase silica gel (EtOAc/heptane) provided (S)-ethyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate.

Step 3: To a solution of (S)-ethyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (50 mg, 0.08 mmol) from Step 2 in THF (2.0 mL) and water (0.2 mL), was added lithium hydroxide monohydrate (58 mg, 0.05 mmol). The reaction mixture was stirred at RT for 2 h, then the solution was neutralized with 1 N HCl and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided the title compound as an off-white solid as the zwitterionic form.

Example 1cq: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(hydroxymethyl)-4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




Step 1: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (Step 2, Example 1u) (300 mg, 0.4 mmol, Step 2) in ethanol (2 mL) and water (0.5 mL) was added (3-(hydroxymethyl)-4-methylphenyl)boronic acid (CAS# 1451391-54-0; 120 mg, 0.7 mmol), PdCl₂(PPh₃)₂ (41 mg, 0.058 mmol), and Cs₂CO₃ (390 mg, 1.2 mmol). The

reaction was heated to 60 °C for 16 h, then cooled to RT, filtered through celite and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(hydroxymethyl)-4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate a white solid.

Step 2: A solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(hydroxymethyl)-4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (200 mg, 0.24 mmol,) in EtOAc (5 mL) was hydrogenated using an H-Cube apparatus and a 10% (w/w) Pd/C cartridge with a flow rate of 1.0 mL/min at RT. Purification on normal phase silica gel (EtOAc/heptane) provided (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(hydroxymethyl)-4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate.

Step 3: To a solution of (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(hydroxymethyl)-4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (50 mg, 0.08 mmol) from Step 2 in THF (2.0 mL) and water (0.2 mL), was added lithium hydroxide monohydrate (58 mg, 0.05 mmol). The reaction mixture was stirred at RT for 2 h, then the solution was neutralized with 1 N HCl, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided the title compound as an off-white solid as the zwitterionic form.

Example 1cr: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(hydroxymethyl)-3'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was made as described for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(hydroxymethyl)-4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 1cq) using (4-
5 (hydroxymethyl)-3-methylphenyl)boronic acid (CAS# 1218790-88-5).

Using the generic scheme below, the following examples of Table 1a were prepared as described above for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 1u). The boronic acid was generally used to make the analogues below, however, where it was
10 not available, the corresponding boronate was used.

























Table 1a.













Ex. No.	Cy	CAS Name	LCMS (MH ⁺)
1a		(3S)-8-(2-amino-6-((1R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-(methylsulfinyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	671
1b		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-(methylthio)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro [4.5]decane-3-carboxylic acid	655
1c		(S)-8-(2-amino-6-((R)-1-(3'-carboxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	652
1d		(S)-8-(2-amino-6-((R)-1-(3'-carboxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	652.5
1e		(S)-8-(2-amino-6-((R)-1-(4'-carboxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	652.6
1f		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(1,2,3,6-tetrahydropyridin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	613.5
1g		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(pyridin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	609.6
1h		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(1-methyl-1H-pyrazol-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-	612.6

		2,8-diazaspiro[4.5]decane-3-carboxylic acid	
1i		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(isoxazol-4-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	599.6
1j		(S)-8-(2-amino-6-((R)-1-(4-(3,6-dihydro-2H-pyran-4-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	614.6
1k		(S)-8-(6-((R)-1-(4-(1-acetyl-1,2,3,6-tetrahydropyridin-4-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	655.7
1l		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	666.7
1m		(S)-8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	636.7
1n		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-methoxypyridin-4-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	639.6
1o		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(3-methyl-1H-indazol-6-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	662.3
1p		(S)-8-(2-amino-6-((R)-1-(4'-(tert-butyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	664.8
1q		(S)-8-(2-amino-6-((R)-1-(4'-ethoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	652.7
1r		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-methoxypyrimidin-5-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	639.6
1s		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(6-methoxypyridin-3-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	639.6











1u		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	608.6
1v		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-2',3',4',5'-tetrahydro-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	636
1w		(S)-8-(2-amino-6-((R)-1-(3'-cyano-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	633
1x		(S)-8-(6-((R)-1-(4'-(acetamidomethyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	679
1y		(S)-8-(6-((R)-1-(4'-(2-acetamidoethyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	693
1z		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(quinolin-7-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	659
1aa		(S)-8-(6-((R)-1-(4-(1H-indol-6-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	567
1ab		(S)-8-(2-amino-6-((R)-1-(4'-(aminomethyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	637
1ac		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	626
1ad		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(quinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	659
1ae		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	622











1af		(S)-8-(2-amino-6-((R)-1-(3',4'-dichloro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	677
1ag		(S)-8-(2-amino-6-((R)-1-(3',4'-difluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	644
1ah		(S)-8-(2-amino-6-((R)-1-(4'-chloro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	643
1ai		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(pyrimidin-5-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	610
1ak		(S)-8-(2-amino-6-((R)-1-(3'-chloro-3-(3-methyl-1H-pyrazol-1-yl)-5'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	711
1al		(S)-8-(2-amino-6-((R)-1-(3'-chloro-4'-ethoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	687
1am		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	676
1an		(S)-8-(2-amino-6-((R)-1-(3'-chloro-5'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	657
1ao		(S)-8-(2-amino-6-((R)-1-(4'-chloro-3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	661
1ap		(S)-8-(2-amino-6-((R)-1-(3'-ethoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	652
1aq		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	640

1ar		(S)-8-(2-amino-6-((R)-1-(3'-chloro-4'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	661
1as		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-3'-(trifluoromethoxy)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	692
1at		(S)-8-(2-amino-6-((R)-1-(3',5'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	637
1au		(S)-8-(2-amino-6-((R)-1-(3',4'-difluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	644
1av		(S)-8-(2-amino-6-((R)-1-(3',5'-difluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	644
1aw		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	694
1ax		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	684
1ay		(S)-8-(2-amino-6-((R)-1-(3'-ethoxy-5'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	670
1az		(S)-8-(2-amino-6-((R)-1-(3'-(tert-butyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	664
1ba		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-fluoro-3'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	640
1bb		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-isopropyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	650

1bc		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	666
1bd		(S)-8-(2-amino-6-((R)-1-(4'-chloro-3'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	656
1be		(S)-8-(2-amino-6-((R)-1-(3'-carbamoyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	651
1bf		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-3',5'-bis(trifluoromethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	744
1bg		(S)-8-(2-amino-6-((R)-1-(3'-ethoxy-4'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	670
1bh		(S)-8-(2-amino-6-((R)-1-(4'-chloro-3',5'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	671
1bi		(S)-8-(2-amino-6-((R)-1-(3',5'-dichloro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	677
1bj		(S)-8-(2-amino-6-((R)-1-(3'-(tert-butyl)-5'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	778
1bk		(S)-8-(2-amino-6-((R)-1-(3'-chloro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	642
1bl		(S)-8-(2-amino-6-((R)-1-(3'-chloro-3-(3-methyl-1H-pyrazol-1-yl)-4'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	711

1bm		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-methoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	638
1bn		(S)-8-(2-amino-6-((R)-1-(4'-ethoxy-3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	670
1bo		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3',4',5'-trifluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	662
1bp		(S)-8-(2-amino-6-((R)-1-(3'-chloro-4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	657
1bq		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-4'-(trifluoromethoxy)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	706
1br		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-5'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	684
1bs		(S)-8-(2-amino-6-((R)-1-(3'-chloro-5'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	661
1bt		(S)-8-(2-amino-6-((R)-1-(4'-chloro-3-(3-methyl-1H-pyrazol-1-yl)-3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	710
1bu		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-5'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	694
1bv		(S)-8-(2-amino-6-((R)-1-(3'-chloro-4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	701

1bw		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(naphthalen-2-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	659
1bx		(S)-8-(2-amino-6-((R)-1-(4'-(benzyloxy)-3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	733
1by		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	681
1bz		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-4'-propoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	685
1ca		(S)-8-(2-amino-6-((R)-1-(4'-butoxy-3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	698
1cb		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-(5-methyl-1,3,4-oxadiazol-2-yl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	709
1cc		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-(methanesulfonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	687
1cd		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-propoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	668
1ce		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-((2-morpholinoethyl)carbamoyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	764
1cf		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-sulfamoyl-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	689

1cg		(S)-8-(2-amino-6-((R)-1-(4'-carbamoyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	652
1ch		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-(methylcarbamoyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	666
1ci		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-methoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	657
1cj		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-(piperazine-1-carbonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	721
1ck		(S)-8-(2-amino-6-((R)-1-(4'-(dimethylcarbamoyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	680
1cl		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isobutoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	681
1cm		(S)-8-(2-amino-6-((R)-1-(4'-(diethylcarbamoyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	707
1cn		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-(neopentyloxy)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	695
1co		(S)-8-(2-amino-6-((R)-1-(4-(chroman-6-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	665
1cp		(S)-8-(2-amino-6-((R)-1-(4-(cinnolin-6-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	661





1cq		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(hydroxymethyl)-4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	652
1cr		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(hydroxymethyl)-3'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	653
1cs		(S)-8-(2-amino-6-((R)-1-(4-(6-ethoxypyridin-3-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	654
1ct		(S)-8-(2-amino-6-((S)-1-(3',4'-bis(hydroxymethyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	669

Table 1b.

¹H NMR Data for Compounds of Table 1a

Ex. No.	¹ H NMR
1a	¹ H NMR (MeOH-d ₄): δ ppm 1.29 (m, 3H), 1.68 (q, J = 6.3 Hz, 4H), 2.10 (dd, J = 13.6, 8.1 Hz, 1H), 2.41 (s, 4H), 2.85 (s, 3H), 3.24 (m, 2H), 3.62 (m, 1H), 3.71 (s, 2H), 3.79 (dd, J = 13.7, 5.8 Hz, 2H), 4.44 (t, J = 8.5 Hz, 1H), 4.83 (s, 2H), 6.44 (d, J = 2.4 Hz, 1H), 6.92 (q, J = 6.2 Hz, 1H), 7.88 (m, 8H)
1b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.31 (m, 3H), 1.60 (t, J = 5.6 Hz, 4H), 2.06 (dd, J = 13.4, 7.0 Hz, 1H), 2.40 (s, 4H), 2.51 (s, 3H), 2.80 (s, 1H), 3.13 (d, J = 11.5 Hz, 1H), 3.25 (d, J = 11.0 Hz, 1H), 3.53 (dd, J = 22.1, 9.7 Hz, 2H), 3.69 (d, J = 14.9 Hz, 2H), 4.14 (t, J = 8.1 Hz, 1H), 4.93 (s, 1H), 6.42 (d, J = 2.3 Hz, 1H), 6.79 (q, J = 6.5 Hz, 1H), 7.34 (m, 2H), 7.63 (dd, J = 8.9, 2.1 Hz, 3H), 7.77 (m, 2H), 7.97 (d, J = 2.3 Hz, 1H)
1c	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.4 (d, J = 18.0 Hz, 1H), 1.68 (q, J = 6.8, 5.6 Hz, 4H), 2.4 (dd, J = 13.6, 8.1 Hz, 1H), 2.41 (s, 4H), 2.85 (s, 1H), 3.25 (m, 2H), 3.64 (m, 1H), 3.72 (s, 1H), 3.79 (d, J = 13.8 Hz, 2H), 4.44 (q, J = 8.6 Hz, 1H), 6.5 (d, J = 2.4 Hz, 1H), 6.92 (dd, J = 10.5, 4.4 Hz, 1H), 7.95 (m, 5H), 8.1 (d, J = 2.5 Hz, 1H), 8.4(m, 2H)
1d	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.68 (dt, J = 9.0, 5.8 Hz, 4H), 2.09 (dd, J = 13.6, 8.1 Hz, 1H), 2.41 (s, 4H), 3.24 (m, 2H), 3.72 (m, 4H), 4.45 (t, J = 8.5 Hz, 1H), 6.44 (d, J = 2.3 Hz, 1H), 6.90 (q, J = 6.2 Hz, 1H), 7.61 (t, J = 7.8 Hz, 1H), 7.75 (d, J = 1.7 Hz, 1H), 7.85 (m, 2H), 7.98 (m, 2H), 8.08 (dt, J = 7.8, 1.3 Hz, 1H), 8.34 (t, J = 1.8 Hz, 1H)
1e	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.68 (dt, J = 9.0, 5.8 Hz, 4H), 2.09 (dd, J = 13.6, 8.1 Hz, 1H), 2.41 (s, 4H), 3.24 (m, 2H), 3.72 (m, 4H), 4.45 (t, J = 8.5 Hz, 1H),

	6.44 (d, J = 2.3 Hz, 1H), 6.90 (q, J = 6.2 Hz, 1H), 7.61 (t, J = 7.8 Hz, 1H), 7.75 (d, J = 1.7 Hz, 1H), 7.85 (m, 2H), 7.98 (m, 2H), 8.08 (dt, J = 7.8, 1.3 Hz, 1H), 8.34 (t, J = 1.8 Hz, 1H)
1f	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.47 - 1.69 (m, 4 H) 1.97 - 2.13 (m, 1 H) 2.19 - 2.35 (m, 1 H) 2.37 (d, J=0.34 Hz, 0 H) 2.66 - 2.81 (m, 2 H) 3.05 - 3.17 (m, 1 H) 3.18 - 3.28 (m, 1 H) 3.33 - 3.40 (m, 2 H) 3.41 - 3.72 (m, 4 H) 3.73 - 3.83 (m, 2 H) 3.99 - 4.13 (m, 1 H) 5.71 (s, 1 H) 6.32 (d, J=0.39 Hz, 1 H) 6.41 (d, J=2.29 Hz, 2 H) 6.67 - 6.79 (m, 1 H) 7.49 (d, J=1.81 Hz, 1 H) 7.55 - 7.64 (m, 1 H) 7.72 (d, J=8.40 Hz, 2 H) 7.92 (d, J=2.29 Hz, 1 H)
1g	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.49 - 1.69 (m, 4 H) 2.06 (dd, J=13.47, 7.03 Hz, 1 H) 2.31 (dd, J=13.42, 9.32 Hz, 1 H) 2.41 (s, 3 H) 3.12 (d, J=12.00 Hz, 1 H) 3.25 (d, J=11.76 Hz, 1 H) 3.38 - 3.57 (m, 2 H) 3.58 - 3.76 (m, 2 H) 4.08 (dd, J=9.13, 7.17 Hz, 1 H) 5.74 (s, 1 H) 6.44 (d, J=2.34 Hz, 1 H) 6.87 (q, J=6.62 Hz, 1 H) 7.75 - 7.80 (m, 2 H) 7.82 (s, 1 H) 7.89 (s, 2 H) 8.02 (d, J=2.34 Hz, 1 H), 8.57 - 8.69 (m, 2 H)
1h	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.61 - 1.84 (m, 5 H) 2.10 (dd, J=13.62, 8.49 Hz, 1 H) 2.40 (s, 3 H) 2.47 (dd, J=13.76, 8.88 Hz, 1 H) 3.25 - 3.29 (m, 2 H) 3.73 - 3.91 (m, 4 H) 3.94 (s, 3 H) 4.53 (t, J=8.64 Hz, 1 H) 6.42 (d, J=2.39 Hz, 1 H) 6.81 (q, J=5.94 Hz, 1 H) 7.61 - 7.70 (m, 2 H) 7.71 - 7.78 (m, 1 H) 7.90 - 7.96 (m, 2 H) 8.12 (s, 1 H)
1i	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 - 1.40 (m, 5 H), 1.60-1.65 (m, 8 H), 2.40 (m, 5 H) 2.47 (dd, J=13.76, 8.88 Hz, 1 H) 3.25 - 3.29 (m, 2 H) 3.73 - 3.91 (m, 4 H) 3.94 (s, 3 H) 4.53 (t, J=8.64 Hz, 1 H), 5.6 (s, 1H), 6.42 (d, J=2.39 Hz, 2 H) 6.70 (m, 1 H) 7.61 - 7.70 (m, 2 H) 7.71 - 7.78 (m, 1 H) 7.90 - 7.96 (m, 2 H) 8.12 (s, 1 H)
1j	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.58 - 1.86 (m, 4 H) 2.01 - 2.20 (m, 1 H) 2.40 (s, 3 H) 2.43 - 2.61 (m, 3 H) 3.61 - 3.75 (m, 2 H) 3.86 (s, 4 H) 3.93 (t, J=5.44 Hz, 2 H) 4.25 - 4.37 (m, 2 H) 4.49 - 4.69 (m, 1 H) 6.42 (d, J=2.24 Hz, 2 H) 6.52 (br. s., 1 H) 6.77 - 6.88 (m, 1 H) 7.51 (d, J=1.32 Hz, 1 H) 7.60 - 7.73 (m, 2 H) 7.91 (d, J=2.34 Hz, 1 H)
1k	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.72 (d, J=18.21 Hz, 4 H) 2.09 (dd, J=13.62, 8.49 Hz, 1 H) 2.18 (d, J=14.50 Hz, 3 H) 2.39 (s, 3 H) 2.48 (dd, J=13.64, 8.91 Hz, 1 H) 2.58 (br. s., 1 H) 2.66 (br. s., 1 H) 3.60 - 3.95 (m, 6 H) 4.24 (br. s., 2 H) 4.55 (t, J=8.71 Hz, 1 H) 6.33 (br. s., 1 H) 6.42 (d, J=2.34 Hz, 1 H) 6.46 (br. s., 1 H) 6.75 - 6.87 (m, 1 H) 7.52 (s, 1 H) 7.62 - 7.74 (m, 2 H) 7.92 (d, J=2.34 Hz, 1 H)
1l	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J=6.05 Hz, 6 H) 1.49 (d, J=5.47 Hz, 4 H) 1.70 - 1.86 (m, 1 H) 1.98 - 2.15 (m, 1 H) 2.37 (s, 3 H) 2.69 (d, J=11.13 Hz, 1 H) 2.93 (s, 1 H) 3.35 - 3.52 (m, 2H) 3.53 - 3.64 (m, 2 H) 3.64 - 3.73 (m, 1 H) 4.59 (s, 1 H) 5.71 (s, 1 H) 6.38 (d, J=2.15 Hz, 1 H) 6.68 - 6.82 (m, 1 H) 6.93 (d, J=8.79 Hz, 2 H) 7.44 - 7.58 (m, 3 H) 7.64 (d, J=1.37 Hz, 1 H) 7.67 - 7.78 (m, 1 H) 7.93 (d, J=2.15 Hz, 1 H)
1m	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.51 (d, J=5.47 Hz, 4 H) 1.71 - 1.86 (m, 1 H) 2.01 - 2.17 (m, 1 H) 2.28 (s, 3 H) 2.31 (s, 3 H) 2.39 (s, 3 H) 2.64 - 2.78 (m, 1 H) 2.90 - 3.05 (m, 1 H) 3.36 - 3.54 (m, 2 H) 3.55 - 3.79 (m, 3 H) 5.73 (s, 1 H) 6.41 (d, J=2.15 Hz, 1 H) 6.69 - 6.87 (m, 1 H) 7.20 (s, 1 H) 7.33 - 7.40 (m, 1 H) 7.43 (s, 1 H) 7.59 (d, J=1.37 Hz, 1 H) 7.70 (s, 1 H) 7.75 (s, 1 H) 7.96 (d, J=2.15 Hz, 1 H)
1n	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.53 (d, J=5.86 Hz, 4 H) 1.75 - 1.87 (m, 1 H) 2.04 - 2.17 (m, 1 H) 2.41 (s, 3 H) 2.64 - 2.76 (m, 1 H) 2.91 - 3.04 (m, 1 H) 3.38 - 3.54 (m, 2 H) 3.55 - 3.74 (m, 3 H), 3.95 (s, 3 H) 5.72 (s, 1 H) 6.44 (d, J=2.34 Hz, 1 H) 6.79 - 6.92 (m, 1 H) 7.12 (s, 1 H) 7.23 - 7.31 (m, 1 H) 7.74 (d, J=1.17 Hz, 1 H) 7.79 - 7.89 (m, 2 H) 8.02 (d, J=2.15 Hz, 1 H) 8.15 - 8.25 (m, 1 H)

1o	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.60 (br. s., 4 H) 2.02 - 2.17 (m, 1 H) 2.24 - 2.39 (m, 1 H) 2.43 (s, 3 H) 2.53 - 2.66 (m, 4 H) 3.07 - 3.17 (m, 1 H) 3.21 - 3.29 (m, 1 H) 3.40 - 3.59 (m, 2 H) 3.61 - 3.80 (m, 2 H) 4.00 - 4.18 (m, 1 H) 5.77 (s, 1 H) 6.45 (d, J=2.15 Hz, 1 H) 6.75 - 6.90 (m, 1 H) 7.36 - 7.56 (m, 1 H) 7.73 (d, J=4.10 Hz, 2 H) 7.77 - 7.89 (m, 4 H) 8.01 (d, J=2.15 Hz, 1 H)
1p	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.24 - 1.42 (m, 9 H) 1.60 (br. s., 4 H) 2.02 - 2.12 (m, 1 H) 2.42 (s, 4 H) 3.05 - 3.17 (m, 1 H) 3.20 - 3.29 (m, 1 H) 3.41 - 3.79 (m, 4 H) 4.02 - 4.17 (m, 1 H) 5.78 (s, 1 H) 6.43 (d, J=2.15 Hz, 1 H) 6.73 - 6.88 (m, 1 H) 7.51 (d, J=8.40 Hz, 2 H) 7.57 - 7.69 (m, 3 H) 7.78 (s, 2 H) 7.98 (d, J=2.15 Hz, 1 H)
1q	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.41 (t, J=7.03 Hz, 3 H) 1.60 (br. s., 4 H) 1.95 - 2.14 (m, 1 H) 2.27 - 2.38 (m, 1 H) 2.41 (s, 3 H) 3.14 (s, 1 H) 3.20 - 3.29 (m, 1 H) 3.41 - 3.59 (m, 2 H) 3.60 - 3.83 (m, 2 H) 3.99 - 4.20 (m, 3 H) 5.77 (s, 1 H) 6.43 (d, J=2.34 Hz, 1 H) 6.71 - 6.85 (m, 1 H) 7.00 (d, J=8.79 Hz, 2 H) 7.52 - 7.65 (m, 3 H) 7.72 (d, J=1.56 Hz, 1 H) 7.76 (s, 1 H) 7.97 (d, J=2.34 Hz, 1 H)
1r	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.62 (d, J=4.88 Hz, 4 H) 2.02 - 2.13 (m, 1 H) 2.27 - 2.38 (m, 1 H) 2.42 (s, 3 H) 3.16 (s, 1 H) 3.26 (s, 1 H) 3.41 - 3.59 (m, 2 H) 3.60 - 3.78 (m, 2 H) 3.99 - 4.19 (m, 4 H) 5.76 (s, 1 H) 6.45 (d, J=2.34 Hz, 1 H) 6.82 - 6.96 (m, 1 H) 7.74 (d, J=1.56 Hz, 1 H) 7.81 (d, J=1.56 Hz, 1 H) 7.86 (s, 1 H) 8.03 (d, J=2.15 Hz, 1 H) 8.92 (s, 2 H)
1s	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.59 (br. s., 4 H) 2.01 - 2.14 (m, 1 H) 2.31 (br. s., 1 H) 2.41 (s, 3 H) 3.07 - 3.17 (m, 1 H) 3.21 - 3.29 (m, 1 H) 3.40 - 3.59 (m, 2 H) 3.67 (d, J=5.47 Hz, 2 H) 3.95 (s, 3 H) 4.09 (d, J=1.17 Hz, 1 H) 5.75 (s, 1 H) 6.43 (d, J=2.15 Hz, 1 H) 6.82 (d, J=6.44 Hz, 1 H) 6.88 (d, J=8.79 Hz, 1 H) 7.64 (d, J=1.56 Hz, 1 H) 7.68 - 7.76 (m, 1 H) 7.77 - 7.87 (m, 1 H) 7.93 - 8.07 (m, 2 H) 8.45 (d, J=2.34 Hz, 1 H)
1u	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.42 - 1.74 (m, 4 H) 2.05 (dd, J=13.50, 7.20 Hz, 1 H) 2.40 (s, 4 H) 3.07 - 3.16 (m, 1 H) 3.17 - 3.29 (m, 1 H) 3.38 - 3.59 (m, 2 H) 3.59 - 3.78 (m, 2 H) 4.11 (dd, J=9.10, 7.25 Hz, 1 H) 6.42 (d, J=2.34 Hz, 1 H) 6.74 - 6.86 (m, 1 H) 7.34 - 7.41 (m, 1 H) 7.42 - 7.50 (m, 2 H) 7.60 - 7.69 (m, 3 H) 7.71 - 7.77 (m, 1 H) 7.77 - 7.83 (m, 1 H) 7.97 (d, J=2.00 Hz, 1 H)
1v	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.24 (t, J=7.59 Hz, 3 H) 1.50 - 1.69 (m, 4 H) 2.06 (dd, J=13.42, 7.13 Hz, 1 H) 2.32 (dd, J=13.45, 9.20 Hz, 1 H) 2.37 (s, 3 H) 2.72 (q, J=7.61 Hz, 2 H) 3.08 - 3.27 (m, 2 H) 3.39 - 3.78 (m, 4 H) 4.08 (dd, J=9.13, 7.17 Hz, 1 H) 5.74 (s, 1 H) 6.36 (d, J=2.34 Hz, 1 H) 6.71 (q, J=6.65 Hz, 1 H) 7.26 - 7.34 (m, 1 H) 7.35 - 7.44 (m, 1 H) 7.56 (s, 1 H) 7.82 (d, J=2.29 Hz, 1 H)
1w	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.51 - 1.64 (m, 4 H) 1.96 (s, 2 H) 2.04 (dd, J=13.23, 7.32 Hz, 1 H) 2.25 - 2.34 (m, 1 H) 2.38 (s, 3 H) 3.09 (d, J=11.67 Hz, 1 H) 3.23 (d, J=11.81 Hz, 1 H) 3.38 - 3.56 (m, 2 H) 3.59 - 3.73 (m, 2 H) 4.00 - 4.10 (m, 1 H) 5.73 (s, 1 H) 6.41 (d, J=2.34 Hz, 1 H) 6.79 - 6.89 (m, 1 H) 7.61 - 7.67 (m, 1 H) 7.69 - 7.88 (m, 4 H) 7.96 - 8.02 (m, 2 H) 8.08 (t, J=1.59 Hz, 1 H)
1x	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.49 - 1.64 (m, 4 H) 1.97 (s, 3 H) 1.98 (s, 3 H) 2.04 (dd, J=13.35, 7.15 Hz, 1 H) 2.29 (dd, J=13.47, 9.18 Hz, 1 H) 2.38 (s, 3 H) 3.05 - 3.26 (m, 2 H) 3.39 - 3.73 (m, 4 H) 4.05 (dd, J=9.18, 7.22 Hz, 1 H) 4.38 (s, 2 H) 5.73 (s, 1 H) 6.40 (d, J=2.29 Hz, 1 H) 6.77 (q, J=6.67 Hz, 1 H) 7.37 (d, J=8.40 Hz, 2 H) 7.59 - 7.66 (m, 3 H) 7.71 - 7.81 (m, 2 H) 7.95 (d, J=2.34 Hz, 1 H)
1y	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.57 (t, J=4.44 Hz, 4 H) 1.89 (s, 3 H) 1.97 (s, 4 H) 2.04 (dd, J=13.37, 7.13 Hz, 1 H) 2.29 (dd, J=13.28, 9.18 Hz, 1 H) 2.38 (s, 3 H) 2.82

	(t, J=7.32 Hz, 2 H) 3.06 - 3.25 (m, 2 H) 3.40 (t, J=7.32 Hz, 2 H) 3.43 - 3.73 (m, 4 H) 4.05 (dd, J=9.18, 7.27 Hz, 1 H) 5.74 (s, 1 H) 6.40 (d, J=2.25 Hz, 1 H) 6.77 (q, J=6.80 Hz, 1 H) 7.31 (d, J=8.30 Hz, 2 H) 7.57 - 7.63 (m, 3H) 7.70 - 7.80 (m, 2 H) 7.95 (d, J=2.34 Hz, 1 H)
1z	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.53 - 1.61 (m, 4 H) 1.97 (s, 3 H) 2.04 (dd, J=13.32, 7.03 Hz, 1 H) 2.30 (dd, J=13.59, 9.30 Hz, 1 H) 2.40 (s, 3 H) 3.06 - 3.26 (m, 2 H) 3.40 - 3.72 (m, 4 H) 4.06 (dd, J=8.91, 7.25 Hz, 1 H) 5.76 (s, 1 H) 6.42 (d, J=2.29 Hz, 1 H) 6.85 (q, J=6.51 Hz, 1 H) 7.55 (dd, J=8.27, 4.32 Hz, 1 H) 7.82 (d, J=1.71 Hz, 1 H) 7.85 - 7.99 (m, 3 H) 8.00 - 8.07 (m, 2 H) 8.29 (s, 1 H) 8.39 (d, J=7.61 Hz, 1 H) 8.88 (dd, J=4.30, 1.66 Hz, 1 H)
1aa	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.59 (d, J=4.69 Hz, 4 H) 1.99 - 2.13 (m, 1 H) 2.24 - 2.39 (m, 1 H) 3.03 - 3.14 (m, 1 H) 3.17 - 3.27 (m, 1 H) 3.38 - 3.54 (m, 2 H) 3.55 - 3.75 (m, 2 H) 3.99 - 4.14 (m, 1 H) 5.56 (s, 1 H) 6.46 (d, J=2.93 Hz, 1 H) 6.58 - 6.71 (m, 1 H) 7.27 (d, J=3.12 Hz, 1 H) 7.29 - 7.36 (m, 1 H) 7.54 - 7.66 (m, 4 H) 7.70 (d, J=8.20 Hz, 2 H)
1ab	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.52 - 1.62 (m, 4 H) 1.92 (s, 5 H) 1.99 - 2.09 (m, 1 H) 2.27 (dd, J=13.42, 9.22 Hz, 1 H) 2.38 (s, 3 H) 3.06 - 3.26 (m, 2 H) 3.39 - 3.73 (m, 4 H) 4.00 - 4.08 (m, 1 H) 4.13 (s, 2 H) 5.72 (s, 1 H) 6.41 (d, J=2.29 Hz, 1 H) 6.78 (q, J=6.49 Hz, 1 H) 7.53 (d, J=8.30 Hz, 2 H) 7.66 (d, J=1.66 Hz, 1 H) 7.73 - 7.84 (m, 4 H) 7.97 (d, J=2.25 Hz, 1 H)
1ac	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.51 - 1.70 (m, 4 H) 2.06 (dd, J=13.37, 7.13 Hz, 1 H) 2.31 (dd, J=13.28, 9.37 Hz, 1 H) 2.41 (s, 3 H) 3.08 - 3.19 (m, 1 H) 3.20 - 3.29 (m, 1 H) 3.39 - 3.78 (m, 4 H), 4.01 - 4.19 (m, 1 H) 5.75 (s, 1 H) 6.43 (d, J=2.15 Hz, 1 H) 6.84 (d, J=6.64 Hz, 1 H) 7.06 - 7.19 (m, 1 H) 7.37 - 7.52 (m, 3 H) 7.65 (d, J=1.56 Hz, 1 H) 7.70 - 7.77 (m, 1 H) 7.78 - 7.86 (m, 1 H) 8.00 (d, J=2.15 Hz, 1 H)
1ad	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.53 - 1.68 (m, 4 H) 2.00 - 2.11 (m, 1 H) 2.25 - 2.36 (m, 1 H) 2.44 (s, 3 H) 3.03 - 3.13 (m, 1 H) 3.18 - 3.26 (m, 1 H) 3.42 - 3.60 (m, 2 H) 3.62 - 3.80 (m, 2 H) 3.98 - 4.12 (m, 1 H) 5.73 - 5.86 (m, 1 H) 6.38 - 6.53 (m, 1 H) 6.80 - 6.96 (m, 1 H) 7.56 - 7.64 (m, 1 H) 7.82 - 7.93 (m, 2 H) 7.93 - 8.00 (m, 1 H) 8.03 - 8.09 (m, 1 H) 8.16 (s, 2 H) 8.28 - 8.38 (m, 1 H), 8.42 - 8.55 (m, 1 H) 8.80 - 8.97 (m, 1 H)
1ae	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (t, J = 6.7 Hz, 1H), 1.30 (d, J = 16.8 Hz, 3H), 1.57 (q, J = 8.1, 5.5 Hz, 4H), 1.98 (m, 1H), 2.24 (m, 1H), 2.38 (d, J = 10.5 Hz, 6H), 2.99 (d, J = 11.6 Hz, 1H), 3.16 (d, J = 11.5 Hz, 1H), 3.48 (ddt, J = 20.2, 13.1, 6.1 Hz, 3H), 3.65 (dd, J = 13.9, 6.2 Hz, 2H), 3.96 (t, J = 8.0 Hz, 1H), 5.75 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.77 (q, J = 6.6 Hz, 1H), 7.26 (d, J = 7.8 Hz, 2H), 7.58 (m, 3H), 7.74 (m, 2H), 7.96 (d, J = 2.4 Hz, 1H)
1af	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.90 (m, 1H), 1.27 (m, 5H), 1.51 (dt, J = 10.5, 5.6 Hz, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.09 (dd, J = 13.1, 8.7 Hz, 1H), 2.40 (s, 3H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.54 (m, 4H), 3.84 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.1, 1.7 Hz, 2H), 5.73 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.84 (q, J = 6.5 Hz, 1H), 7.64 (m, 3H), 7.80 (m, 3H), 8.01 (d, J = 2.4 Hz, 1H)
1ag	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 14.8 Hz, 1H), 1.58 (s, 8H), 2.05 (m, 2H), 2.39 (s, 8H), 3.12 (d, J = 11.1 Hz, 2H), 3.24 (d, J = 11.2 Hz, 3H), 3.47 (s, 3H), 3.53 (s, 1H), 3.64 (s, 4H), 4.11 (s, 2H), 4.96 (s, 1H), 6.42 (d, J = 2.0 Hz, 2H), 6.83 (q, J = 6.6 Hz, 2H), 7.33 (q, J = 9.0 Hz, 2H), 7.47 (t, J = 6.1 Hz, 2H), 7.65 (m, 6H), 7.79 (d, J = 8.1 Hz, 2H), 7.99 (d, J = 2.0 Hz, 2H)

1ah	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 10.2 Hz, 1H), 1.57 (t, J = 5.3 Hz, 5H), 2.04 (dd, J = 13.2, 6.9 Hz, 1H), 2.34 (m, 5H), 3.09 (d, J = 11.8 Hz, 1H), 3.22 (m, 2H), 3.48 (dd, J = 25.5, 12.5 Hz, 3H), 3.64 (s, 3H), 4.06 (t, J = 7.9 Hz, 1H), 4.83 (s, 2H), 4.93 (s, 1H), 5.74 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.81 (q, J = 6.7 Hz, 1H), 7.45 (d, J = 8.2 Hz, 2H), 7.65 (m, 3H), 7.76 (m, 3H), 7.98 (d, J = 2.3 Hz, 1H)
1ai	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.91 (tt, J = 8.8, 4.7 Hz, 1H), 1.32 (m, 3H), 1.58 (h, J = 5.2, 4.4 Hz, 8H), 1.99 (m, 2H), 2.25 (dd, J = 13.4, 9.0 Hz, 2H), 2.40 (s, 6H), 3.03 (d, J = 11.5 Hz, 2H), 3.18 (d, J = 11.5 Hz, 2H), 3.48 (ddt, J = 21.1, 13.0, 5.9 Hz, 5H), 3.62 (dt, J = 11.3, 6.3 Hz, 4H), 3.98 (t, J = 7.9 Hz, 2H), 5.73 (s, 2H), 6.44 (d, J = 2.4 Hz, 2H), 6.90 (q, J = 6.6 Hz, 2H), 7.86 (m, 6H), 8.04 (d, J = 2.4 Hz, 2H), 9.15 (d, J = 11.9 Hz, 6H)
1ak	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.67 (dt, J = 11.3, 5.3 Hz, 4H), 2.08 (dd, J = 13.6, 8.1 Hz, 1H), 2.40 (m, 4H), 3.25 (q, J = 11.8, 9.3 Hz, 3H), 3.67 (m, 4H), 4.43 (t, J = 8.5 Hz, 1H), 6.44 (d, J = 2.4 Hz, 1H), 6.95 (q, J = 6.3 Hz, 1H), 7.77 (dt, J = 5.4, 1.8 Hz, 2H), 7.86 (d, J = 1.4 Hz, 2H), 7.96 (m, 1H), 8.05 (d, J = 2.4 Hz, 2H)
1al	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.44 (t, J = 7.0 Hz, 3H), 1.57 (t, J = 5.6 Hz, 4H), 2.02 (dd, J = 13.4, 7.0 Hz, 1H), 2.28 (dd, J = 13.3, 9.1 Hz, 1H), 2.39 (s, 3H), 3.06 (d, J = 11.6 Hz, 1H), 3.21 (d, J = 11.6 Hz, 1H), 3.47 (dd, J = 22.4, 13.7 Hz, 3H), 3.65 (dd, J = 13.8, 6.9 Hz, 2H), 4.03 (t, J = 8.1 Hz, 1H), 4.14 (q, J = 7.0 Hz, 2H), 4.93 (s, 2H), 5.75 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.6 Hz, 1H), 7.12 (d, J = 8.7 Hz, 1H), 7.57 (m, 2H), 7.73 (m, 3H), 7.98 (d, J = 2.4 Hz, 1H)
1am	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 3.7 Hz, 2H), 1.55 (m, 4H), 1.92 (dd, J = 13.4, 7.2 Hz, 1H), 2.19 (t, J = 10.6 Hz, 1H), 2.40 (s, 3H), 2.88 (d, J = 11.4 Hz, 1H), 3.10 (d, J = 11.5 Hz, 1H), 3.47 (dd, J = 22.2, 15.6 Hz, 3H), 3.64 (s, 3H), 3.85 (t, J = 8.1 Hz, 1H), 5.74 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.83 (q, J = 6.6 Hz, 1H), 7.69 (m, 3H), 7.82 (m, 2H), 7.99 (m, 3H)
1an	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.17 (t, J = 7.0 Hz, 1H), 1.29 (m, 1H), 1.57 (d, J = 5.9 Hz, 4H), 2.00 (dd, J = 13.4, 7.0 Hz, 1H), 2.26 (dd, J = 13.1, 9.4 Hz, 1H), 2.38 (d, J = 9.1 Hz, 6H), 3.03 (d, J = 11.6 Hz, 1H), 3.18 (d, J = 11.5 Hz, 1H), 3.47 (ddt, J = 20.8, 13.2, 6.0 Hz, 2H), 3.62 (h, J = 7.1, 6.4 Hz, 3H), 4.00 (t, J = 7.9 Hz, 1H), 4.89 (s, 9H), 5.74 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.82 (q, J = 6.9, 6.4 Hz, 1H), 7.21 (s, 1H), 7.40 (s, 1H), 7.45 (s, 1H), 7.60 (d, J = 1.9 Hz, 1H), 7.68 (dd, J = 8.0, 1.8 Hz, 1H), 7.78 (d, J = 8.2 Hz, 1H), 7.99 (d, J = 2.3 Hz, 1H)
1ao	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (m, 2H), 1.55 (q, J = 7.5, 5.1 Hz, 4H), 1.94 (dd, J = 13.2, 6.9 Hz, 1H), 2.21 (dd, J = 13.1, 8.9 Hz, 1H), 2.39 (s, 3H), 2.94 (d, J = 11.5 Hz, 1H), 3.12 (d, J = 11.2 Hz, 1H), 3.46 (ddt, J = 20.2, 13.0, 5.9 Hz, 2H), 3.63 (dd, J =

	13.7, 6.1 Hz, 2H), 3.90 (t, J = 8.0 Hz, 1H), 4.85 (d, J = 9.0 Hz, 1H), 5.73 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.83 (q, J = 6.6 Hz, 1H), 7.57 (m, 4H), 7.77 (m, 2H), 8.00 (d, J = 2.4 Hz, 1H)
1ap	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 11.9 Hz, 2H), 1.39 (t, J = 7.0 Hz, 3H), 1.59 (t, J = 5.7 Hz, 4H), 2.05 (dd, J = 13.4, 7.2 Hz, 1H), 2.39 (s, 4H), 3.12 (d, J = 11.6 Hz, 1H), 3.24 (d, J = 11.7 Hz, 1H), 3.51 (ddt, J = 25.1, 13.2, 5.8 Hz, 2H), 3.67 (dd, J = 13.8, 5.7 Hz, 2H), 4.10 (dq, J = 14.0, 7.7, 7.0 Hz, 3H), 6.41 (d, J = 2.3 Hz, 1H), 6.79 (q, J = 6.6 Hz, 1H), 6.93 (dd, J = 8.2, 2.5 Hz, 1H), 7.19 (m, 2H), 7.34 (t, J = 7.9 Hz, 1H), 7.62 (d, J = 1.8 Hz, 1H), 7.75 (m, 2H), 7.97 (d, J = 2.4 Hz, 1H)
1aq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (t, J = 7.2 Hz, 1H), 1.28 (s, 4H), 1.58 (t, J = 5.6 Hz, 4H), 2.05 (dd, J = 13.5, 7.1 Hz, 1H), 2.29 (d, J = 1.8 Hz, 4H), 2.39 (s, 3H), 3.10 (d, J = 11.8 Hz, 1H), 3.24 (d, J = 11.5 Hz, 1H), 3.49 (ddt, J = 21.1, 12.9, 5.8 Hz, 3H), 3.68 (ddt, J = 18.9, 12.4, 6.2 Hz, 3H), 4.07 (m, 1H), 5.75 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.80 (q, J = 6.5 Hz, 1H), 7.37 (m, 3H), 7.63 (d, J = 1.8 Hz, 1H), 7.76 (m, 2H), 7.98 (d, J = 2.3 Hz, 1H)
1ar	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.57 (t, J = 5.8 Hz, 4H), 1.99 (dd, J = 13.3, 7.0 Hz, 1H), 2.25 (dd, J = 13.3, 9.1 Hz, 1H), 2.39 (s, 3H), 3.00 (d, J = 11.3 Hz, 1H), 3.17 (d, J = 11.5 Hz, 1H), 3.31 (d, J = 2.4 Hz, 4H), 3.49 (m, 2H), 3.65 (dd, J = 13.8, 6.6 Hz, 2H), 3.97 (t, J = 8.1 Hz, 1H), 5.74 (s, 1H), 6.42 (d, J = 2.3 Hz, 1H), 6.83 (q, J = 6.6 Hz, 1H), 7.33 (t, J = 8.8 Hz, 1H), 7.65 (m, 2H), 7.73 (dd, J = 8.2, 1.9 Hz, 1H), 7.82 (m, 2H), 8.00 (d, J = 2.3 Hz, 1H)
1as	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.60 (m, 4H), 2.05 (dd, J = 13.4, 7.1 Hz, 1H), 2.32 (dd, J = 13.4, 9.1 Hz, 1H), 2.40 (s, 3H), 3.12 (d, J = 11.7 Hz, 1H), 3.24 (d, J = 11.7 Hz, 1H), 3.51 (dq, J = 24.6, 7.3, 6.5 Hz, 2H), 3.66 (dt, J = 11.7, 6.2 Hz, 2H), 4.11 (dd, J = 9.1, 7.1 Hz, 1H), 6.42 (d, J = 2.3 Hz, 1H), 6.83 (q, J = 6.3 Hz, 1H), 7.32 (m, 1H), 7.58 (dd, J = 15.5, 7.5 Hz, 2H), 7.69 (m, 2H), 7.80 (m, 2H), 8.01 (d, J = 2.3 Hz, 1H)
1at	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.57 (m, 4H), 1.97 (dd, J = 13.1, 6.8 Hz, 1H), 2.23 (m, 2H), 2.37 (d, J = 18.2 Hz, 9H), 2.98 (d, J = 11.6 Hz, 1H), 3.15 (d, J = 11.5 Hz, 1H), 3.48 (ddt, J = 20.5, 13.0, 5.8 Hz, 2H), 3.65 (dd, J = 13.4, 5.8 Hz, 2H), 3.94 (dd, J = 9.1, 7.1 Hz, 1H), 5.75 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.6 Hz, 1H), 7.03 (s, 1H), 7.27 (d, J = 1.6 Hz, 2H), 7.59 (d, J = 1.8 Hz, 1H), 7.73 (m, 2H), 7.97 (d, J = 2.4 Hz, 1H)
1au	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 3.4 Hz, 2H), 1.60 (t, J = 5.7 Hz, 4H), 2.06 (dd, J = 13.5, 7.2 Hz, 1H), 2.39 (s, 4H), 3.13 (d, J = 11.7 Hz, 1H), 3.25 (d, J = 11.7 Hz, 1H), 3.52 (m, 2H), 3.67 (dt, J = 12.0, 6.4 Hz, 2H), 4.14 (dd, J = 9.1, 7.2 Hz,

	¹ H, 4.91 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.83 (q, J = 6.6 Hz, 1H), 7.36 (dt, J = 10.4, 8.4 Hz, 1H), 7.51 (ddt, J = 8.1, 3.9, 1.6 Hz, 1H), 7.65 (m, 2H), 7.77 (m, 2H), 7.99 (d, J = 2.3 Hz, 1H)
1av	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.57 (dt, J = 6.7, 3.0 Hz, 4H), 1.98 (dd, J = 13.3, 7.0 Hz, 1H), 2.24 (dd, J = 13.4, 9.2 Hz, 1H), 2.40 (s, 3H), 2.99 (d, J = 11.5 Hz, 1H), 3.16 (d, J = 11.5 Hz, 1H), 3.48 (ddt, J = 20.5, 13.2, 5.9 Hz, 2H), 3.65 (dq, J = 11.1, 5.0 Hz, 2H), 3.95 (t, J = 8.2 Hz, 1H), 5.74 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.84 (q, J = 6.6 Hz, 1H), 6.99 (tt, J = 9.0, 2.3 Hz, 1H), 7.35 (m, 2H), 7.68 (d, J = 1.9 Hz, 1H), 7.79 (m, 2H), 8.02 (d, J = 2.4 Hz, 1H)
1aw	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (m, 1H), 1.30 (d, J = 13.3 Hz, 2H), 1.51 (q, J = 6.5, 5.7 Hz, 5H), 1.77 (dd, J = 13.0, 6.9 Hz, 1H), 2.05 (dd, J = 13.0, 8.9 Hz, 1H), 2.40 (s, 3H), 2.62 (d, J = 11.0 Hz, 1H), 2.93 (d, J = 11.0 Hz, 1H), 3.42 (d, J = 14.2 Hz, 2H), 3.49 (s, 1H), 3.62 (dt, J = 16.7, 6.6 Hz, 3H), 5.72 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.83 (q, J = 6.6 Hz, 1H), 7.44 (m, 1H), 7.69 (d, J = 1.9 Hz, 1H), 7.81 (m, 2H), 8.01 (m, 3H)
1ax	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (m, 2H), 1.32 (m, 13H), 1.57 (t, J = 5.4 Hz, 4H), 2.01 (m, 1H), 2.29 (dd, J = 13.4, 9.2 Hz, 1H), 2.39 (s, 3H), 2.80 (s, 1H), 3.07 (d, J = 11.6 Hz, 1H), 3.17 (s, 1H), 3.50 (m, 2H), 3.66 (d, J = 13.8 Hz, 2H), 4.03 (t, J = 8.1 Hz, 1H), 4.65 (p, J = 6.1 Hz, 1H), 5.75 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.78 (q, J = 6.6 Hz, 1H), 7.16 (t, J = 8.6 Hz, 1H), 7.45 (m, 2H), 7.61 (d, J = 1.8 Hz, 1H), 7.74 (m, 2H), 7.98 (d, J = 2.4 Hz, 1H)
1ay	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.88 (m, 2H), 1.28 (s, 2H), 1.39 (t, J = 7.0 Hz, 3H), 1.59 (d, J = 5.6 Hz, 4H), 2.04 (dd, J = 13.5, 7.1 Hz, 1H), 2.39 (s, 4H), 3.10 (d, J = 11.7 Hz, 1H), 3.24 (m, 1H), 3.49 (ddt, J = 25.3, 13.1, 5.9 Hz, 2H), 3.66 (dq, J = 12.1, 5.6 Hz, 2H), 4.07 (dq, J = 12.7, 7.2 Hz, 3H), 5.75 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.71 (dt, J = 10.8, 2.2 Hz, 1H), 6.80 (p, J = 6.6 Hz, 1H), 7.01 (m, 2H), 7.63 (d, J = 1.8 Hz, 1H), 7.76 (m, 2H), 8.00 (d, J = 2.4 Hz, 1H)
1az	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.90 (m, 1H), 1.36 (s, 13H), 1.59 (m, 4H), 2.05 (dd, J = 13.4, 6.9 Hz, 1H), 2.33 (dt, J = 13.7, 6.0 Hz, 1H), 2.40 (s, 3H), 3.12 (d, J = 11.4 Hz, 1H), 3.24 (d, J = 12.1 Hz, 1H), 3.40 (s, 1H), 3.53 (d, J = 14.7 Hz, 1H), 3.68 (d, J = 13.4 Hz, 2H), 4.10 (s, 1H), 4.76 (m, 5H), 4.83 (s, 2H), 5.01 (s, 1H), 6.42 (d, J = 2.2 Hz, 1H), 6.77 (q, J = 6.6 Hz, 1H), 7.42 (m, 3H), 7.65 (m, 4H), 7.76 (m, 2H), 7.98 (d, J = 2.3 Hz, 1H)
1ba	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (d, J = 12.1 Hz, 3H), 1.59 (t, J = 5.2 Hz, 4H), 2.05 (dd, J = 13.4, 7.0 Hz, 1H), 2.34 (m, 7H), 3.11 (d, J = 11.7 Hz, 1H), 3.24 (d, J = 11.8 Hz, 1H), 3.51 (m, 2H), 3.68 (dt, J = 13.1, 6.3 Hz, 2H), 4.09 (dd, J = 9.2, 6.9 Hz, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.79 (q, J = 6.7 Hz, 1H), 7.11 (t, J = 9.1 Hz, 1H), 7.55 (m,

	3H), 7.74 (m, 2H), 7.98 (d, J = 2.3 Hz, 1H)
1bb	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (d, J = 6.9 Hz, 6H), 1.59 (t, J = 5.6 Hz, 4H), 2.05 (dd, J = 13.4, 6.9 Hz, 1H), 2.32 (dd, J = 13.4, 8.9 Hz, 1H), 2.40 (s, 3H), 2.96 (h, J = 6.8 Hz, 1H), 3.12 (d, J = 11.8 Hz, 1H), 3.24 (d, J = 11.7 Hz, 1H), 3.51 (m, 2H), 3.67 (m, 2H), 4.10 (t, J = 8.3 Hz, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.5 Hz, 1H), 7.27 (m, 1H), 7.37 (t, J = 7.7 Hz, 1H), 7.48 (m, 2H), 7.62 (d, J = 1.8 Hz, 1H), 7.76 (m, 2H), 7.98 (d, J = 2.4 Hz, 1H)
1bc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (m, 1H), 1.29 (m, 9H), 1.55 (d, J = 5.8 Hz, 4H), 1.99 (m, 1H), 2.26 (dd, J = 13.4, 9.1 Hz, 1H), 2.39 (s, 3H), 3.04 (d, J = 11.6 Hz, 1H), 3.17 (d, J = 11.6 Hz, 1H), 3.47 (ddt, J = 20.9, 13.0, 5.9 Hz, 2H), 3.62 (dq, J = 11.5, 5.7 Hz, 2H), 4.01 (m, 1H), 4.64 (hept, J = 5.9 Hz, 1H), 5.74 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.79 (q, J = 6.6 Hz, 1H), 6.91 (dd, J = 8.1, 2.4 Hz, 1H), 7.16 (ddd, J = 8.5, 5.3, 1.7 Hz, 2H), 7.32 (t, J = 7.9 Hz, 1H), 7.64 (m, 3H), 7.77 (d, J = 8.3 Hz, 1H), 7.97 (d, J = 2.4 Hz, 1H)
1bd	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.59 (d, J = 5.6 Hz, 4H), 2.05 (dd, J = 13.4, 7.0 Hz, 1H), 2.31 (dd, J = 13.4, 9.3 Hz, 1H), 2.41 (d, J = 11.1 Hz, 6H), 3.11 (d, J = 11.7 Hz, 1H), 3.24 (d, J = 11.7 Hz, 1H), 3.49 (ddd, J = 27.1, 12.8, 5.9 Hz, 2H), 3.66 (dq, J = 13.8, 6.0 Hz, 2H), 4.07 (dd, J = 9.2, 7.0 Hz, 1H), 5.75 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.80 (q, J = 6.6 Hz, 1H), 7.45 (m, 2H), 7.63 (dd, J = 7.2, 2.0 Hz, 2H), 7.76 (m, 2H), 7.98 (d, J = 2.3 Hz, 1H)
1be	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 3.9 Hz, 4H), 1.59 (t, J = 5.8 Hz, 7H), 2.05 (dd, J = 13.3, 7.0 Hz, 2H), 2.31 (dd, J = 13.4, 9.2 Hz, 2H), 2.40 (s, 5H), 3.11 (d, J = 11.7 Hz, 2H), 3.24 (m, 4H), 3.50 (dq, J = 23.7, 7.4, 6.5 Hz, 4H), 3.67 (dq, J = 13.1, 6.4 Hz, 4H), 4.09 (t, J = 8.1 Hz, 2H), 4.84 (s, 1H), 4.93 (s, 1H), 6.42 (d, J = 2.3 Hz, 2H), 6.84 (q, J = 6.5 Hz, 2H), 7.57 (t, J = 7.8 Hz, 2H), 7.74 (s, 2H), 7.88 (m, 7H), 7.99 (d, J = 2.4 Hz, 2H), 8.19 (t, J = 1.9 Hz, 2H)
1bf	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.57 (t, J = 5.7 Hz, 4H), 1.99 (dd, J = 13.3, 7.0 Hz, 1H), 2.25 (dd, J = 13.3, 9.1 Hz, 1H), 2.41 (s, 3H), 3.01 (d, J = 11.6 Hz, 1H), 3.17 (d, J = 11.5 Hz, 1H), 3.48 (dq, J = 23.5, 7.0, 6.5 Hz, 2H), 3.64 (dd, J = 13.0, 5.8 Hz, 2H), 3.97 (dd, J = 9.0, 7.1 Hz, 1H), 5.73 (s, 1H), 6.44 (d, J = 2.4 Hz, 1H), 6.87 (q, J = 6.6 Hz, 1H), 7.79 (d, J = 1.8 Hz, 1H), 7.88 (m, 2H), 8.00 (s, 1H), 8.07 (d, J = 2.3 Hz, 1H), 8.29 (s, 2H)
1bg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.43 (t, J = 7.0 Hz, 3H), 1.64 (q, J = 5.8 Hz, 4H), 2.08 (dd, J = 13.5, 7.7 Hz, 1H), 2.40 (s, 4H), 3.23 (m, 2H), 3.64 (m, 4H), 4.18 (q, J = 7.0 Hz, 2H), 4.32 (t, J = 8.4 Hz, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.83 (q, J = 6.4 Hz, 1H), 7.20 (m, 2H), 7.36 (dd, J = 8.0, 2.1 Hz, 1H), 7.65 (d, J = 1.6 Hz, 1H), 7.77 (m, 2H), 7.99 (d, J = 2.4 Hz, 1H)

1bh	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.59 (d, J = 5.4 Hz, 4H), 2.04 (dd, J = 13.4, 7.0 Hz, 1H), 2.29 (m, 2H), 2.41 (d, J = 9.6 Hz, 9H), 3.10 (d, J = 11.7 Hz, 1H), 3.24 (m, 1H), 3.49 (ddt, J = 20.7, 13.0, 6.0 Hz, 2H), 3.65 (dt, J = 13.2, 7.2 Hz, 2H), 4.06 (dd, J = 9.2, 7.1 Hz, 1H), 5.75 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.79 (q, J = 6.6 Hz, 1H), 7.45 (s, 2H), 7.62 (d, J = 1.8 Hz, 1H), 7.75 (m, 2H), 7.98 (d, J = 2.4 Hz, 1H)
1bi	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.57 (m, 4H), 1.99 (dd, J = 13.4, 7.1 Hz, 1H), 2.26 (dd, J = 13.3, 9.1 Hz, 1H), 2.40 (s, 3H), 3.02 (d, J = 11.6 Hz, 1H), 3.18 (d, J = 11.6 Hz, 1H), 3.48 (dq, J = 23.5, 7.1, 6.6 Hz, 2H), 3.64 (dq, J = 11.4, 5.5 Hz, 2H), 3.99 (dd, J = 9.2, 7.0 Hz, 1H), 5.73 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.85 (q, J = 6.6 Hz, 1H), 7.46 (t, J = 1.9 Hz, 1H), 7.66 (t, J = 1.8 Hz, 3H), 7.73 (dd, J = 8.3, 2.0 Hz, 1H), 7.82 (d, J = 8.1 Hz, 1H), 8.02 (d, J = 2.4 Hz, 1H)
1bj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (m, 1H), 1.34 (s, 10H), 1.61 (t, J = 5.9 Hz, 4H), 2.06 (dd, J = 13.5, 7.3 Hz, 1H), 2.39 (m, 7H), 3.14 (d, J = 11.6 Hz, 1H), 3.25 (m, 1H), 3.55 (m, 2H), 3.68 (s, 2H), 4.18 (dd, J = 9.0, 7.3 Hz, 1H), 6.42 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.5 Hz, 1H), 7.28 (m, 2H), 7.45 (t, J = 1.7 Hz, 1H), 7.60 (d, J = 1.8 Hz, 1H), 7.75 (m, 2H), 7.98 (d, J = 2.3 Hz, 1H)
1bk	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 2H), 1.58 (t, J = 5.5 Hz, 4H), 2.04 (dd, J = 13.3, 6.9 Hz, 1H), 2.30 (dd, J = 13.3, 9.1 Hz, 1H), 2.40 (s, 3H), 3.09 (d, J = 11.7 Hz, 1H), 3.23 (d, J = 12.2 Hz, 1H), 3.51 (m, 2H), 3.66 (dd, J = 13.9, 6.6 Hz, 2H), 4.05 (t, J = 8.2 Hz, 1H), 4.62 (s, 1H), 5.75 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.82 (q, J = 6.5 Hz, 1H), 7.43 (m, 2H), 7.63 (m, 2H), 7.75 (m, 3H), 8.00 (d, J = 2.3 Hz, 1H)
1bl	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.57 (t, J = 5.6 Hz, 4H), 2.00 (dd, J = 13.5, 7.0 Hz, 1H), 2.26 (dd, J = 13.3, 9.1 Hz, 1H), 2.40 (s, 3H), 3.03 (d, J = 11.6 Hz, 1H), 3.18 (d, J = 11.5 Hz, 1H), 3.47 (ddd, J = 20.9, 14.0, 6.5 Hz, 2H), 3.64 (t, J = 7.4 Hz, 2H), 3.99 (dd, J = 9.1, 7.1 Hz, 1H), 5.74 (s, 1H), 6.43 (d, J = 2.3 Hz, 1H), 6.86 (q, J = 6.6 Hz, 1H), 7.81 (m, 5H), 7.98 (d, J = 1.6 Hz, 1H), 8.03 (d, J = 2.4 Hz, 1H)
1bm	¹ H NMR (400 MHz, MeOH-d ₄): δ 1.54 (d, J=2.93 Hz, 4 H), 1.82 - 1.99 (m, 1 H), 2.09 - 2.24 (m, 1 H), 2.40 (s, 3 H), 2.79 - 2.93 (m, 1 H), 2.99 - 3.14 (m, 1 H), 3.37 - 3.55 (m, 2 H), 3.56 - 3.72 (m, 2 H), 3.82 (s, 4 H), 5.74 (s, 1 H), 6.41 (d, J=2.15 Hz, 1 H), 6.70 - 6.84 (m, 1 H), 6.99 (d, J=8.79 Hz, 2 H), 7.50 - 7.63 (m, 3 H), 7.64 - 7.71 (m, 1 H), 7.71 - 7.80 (m, 1 H), 7.95 (d, J=2.15 Hz, 1 H)
1bn	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (m, 1H), 1.28 (s, 2H), 1.42 (t, J = 7.0 Hz, 3H), 1.58 (t, J = 5.2 Hz, 4H), 2.05 (dd, J = 13.4, 7.0 Hz, 1H), 2.31 (dd, J = 13.4, 9.0 Hz, 1H), 2.39 (s, 3H), 3.11 (d, J = 11.8 Hz, 1H), 3.24 (d, J = 11.7 Hz, 1H), 3.51 (m, 2H), 3.67 (m, 2H), 4.13 (m, 3H), 4.63 (s, 1H), 5.75 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.6 Hz, 1H), 7.15 (t, J = 8.6 Hz, 1H), 7.45 (m, 2H), 7.60 (d, J = 1.8 Hz, 1H), 7.73 (m, 2H), 7.97 (d, J = 2.4 Hz, 1H)


1bo	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 8.3 Hz, 1H), 1.59 (t, J = 5.3 Hz, 4H), 2.05 (dd, J = 13.5, 7.0 Hz, 1H), 2.32 (dd, J = 13.6, 9.2 Hz, 1H), 2.39 (s, 3H), 3.12 (d, J = 11.6 Hz, 1H), 3.25 (m, 1H), 3.49 (ddd, J = 24.6, 12.8, 5.8 Hz, 2H), 3.66 (dq, J = 12.7, 6.0 Hz, 2H), 4.09 (t, J = 8.1 Hz, 1H), 5.76 (s, 1H), 6.42 (d, J = 2.3 Hz, 1H), 6.85 (q, J = 6.6 Hz, 1H), 7.54 (m, 2H), 7.68 (d, J = 1.8 Hz, 1H), 7.78 (m, 2H), 8.01 (d, J = 2.3 Hz, 1H)
1bp	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 8.6 Hz, 1H), 1.59 (d, J = 5.9 Hz, 4H), 2.05 (dd, J = 13.5, 7.3 Hz, 1H), 2.39 (m, 7H), 3.13 (d, J = 11.6 Hz, 1H), 3.24 (m, 2H), 3.52 (dq, J = 25.3, 6.3 Hz, 2H), 3.68 (dd, J = 13.7, 5.9 Hz, 2H), 4.16 (m, 1H), 6.42 (d, J = 2.3 Hz, 1H), 6.82 (q, J = 6.5 Hz, 1H), 7.37 (d, J = 7.9 Hz, 1H), 7.51 (dd, J = 7.9, 2.0 Hz, 1H), 7.63 (d, J = 1.9 Hz, 1H), 7.73 (m, 3H), 7.98 (d, J = 2.4 Hz, 1H)
1bq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 8.0 Hz, 1H), 1.59 (m, 4H), 2.05 (dd, J = 13.4, 7.0 Hz, 1H), 2.38 (d, J = 13.7 Hz, 7H), 3.13 (d, J = 11.7 Hz, 1H), 3.25 (d, J = 11.6 Hz, 1H), 3.53 (dt, J = 31.7, 10.3 Hz, 2H), 3.67 (dd, J = 13.5, 7.0 Hz, 2H), 4.15 (m, 1H), 4.89 (s, 17H), 6.42 (d, J = 2.4 Hz, 1H), 6.82 (q, J = 6.6 Hz, 1H), 7.32 (m, 1H), 7.61 (m, 4H), 7.76 (m, 2H), 7.99 (d, J = 2.3 Hz, 1H)
1br	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.90 (s, 1H), 1.29 (m, 8H), 1.60 (d, J = 7.7 Hz, 4H), 2.02 (m, 2H), 2.39 (s, 4H), 3.14 (d, J = 11.6 Hz, 1H), 3.25 (d, J = 11.4 Hz, 1H), 3.51 (m, 2H), 3.66 (dd, J = 13.6, 6.6 Hz, 2H), 4.11 (dt, J = 21.0, 7.6 Hz, 1H), 4.65 (h, J = 6.0 Hz, 1H), 6.42 (d, J = 2.3 Hz, 1H), 6.69 (dt, J = 10.9, 2.2 Hz, 1H), 6.82 (q, J = 6.3 Hz, 1H), 6.99 (m, 2H), 7.61 (d, J = 1.8 Hz, 1H), 7.71 (m, 1H), 7.78 (d, J = 8.1 Hz, 1H), 7.99 (d, J = 2.3 Hz, 1H)
1bs	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.09 (s, 1H), 0.90 (q, J = 8.4, 7.7 Hz, 1H), 1.29 (d, J = 8.6 Hz, 3H), 1.59 (d, J = 5.7 Hz, 5H), 2.05 (dd, J = 13.5, 6.9 Hz, 1H), 2.31 (dd, J = 13.6, 9.4 Hz, 1H), 2.40 (s, 3H), 3.13 (d, J = 11.6 Hz, 1H), 3.25 (m, 1H), 3.61 (m, 5H), 4.08 (m, 1H), 5.74 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.85 (q, J = 6.7 Hz, 1H), 7.24 (dt, J = 8.5, 2.1 Hz, 1H), 7.45 (dt, J = 9.6, 2.0 Hz, 1H), 7.58 (t, J = 1.7 Hz, 1H), 7.68 (d, J = 1.9 Hz, 1H), 7.79 (m, 2H), 8.02 (d, J = 2.3 Hz, 1H)
1bt	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (d, J = 1.7 Hz, 1H), 1.58 (t, J = 5.7 Hz, 4H), 2.03 (dd, J = 13.4, 7.0 Hz, 1H), 2.29 (dd, J = 13.4, 9.2 Hz, 1H), 2.40 (s, 3H), 3.07 (d, J = 11.6 Hz, 1H), 3.22 (d, J = 11.6 Hz, 1H), 3.49 (m, 2H), 3.65 (dd, J = 13.0, 6.7 Hz, 2H), 4.04 (dd, J = 9.2, 7.0 Hz, 1H), 5.73 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.84 (q, J = 6.6 Hz, 1H), 7.77 (m, 5H), 7.92 (dd, J = 8.4, 2.2 Hz, 1H), 8.04 (dd, J = 14.2, 2.3 Hz, 2H)
1bu	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.60 (d, J = 5.4 Hz, 4H), 2.05 (dd, J = 13.5, 7.1 Hz, 1H), 2.40 (s, 4H), 3.11 (d, J = 11.7 Hz, 1H), 3.24 (d, J = 11.7 Hz, 1H), 3.49 (ddt, J = 21.1, 13.3, 6.0 Hz, 2H), 3.67 (dt, J = 12.9, 6.1 Hz, 2H), 4.07 (dd, J = 9.2, 7.1 Hz, 1H), 5.74 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.86 (q, J = 6.6 Hz, 1H), 7.50 (m, 1H), 7.80 (m, 5H), 8.04 (d, J = 2.4 Hz, 1H)
1bv	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (d, J = 6.0 Hz, 7H), 1.55 (d, J = 5.9 Hz, 4H), 1.98 (dd, J = 13.3, 7.0 Hz, 1H), 2.25 (dd, J = 13.3, 9.1 Hz, 1H), 2.39 (s, 2H), 3.01 (d, J = 11.5 Hz, 1H), 3.17 (d, J = 11.6 Hz, 1H), 3.47 (ddt, J = 20.8, 13.0, 6.0 Hz, 2H), 3.64 (dd, J = 13.9, 6.3 Hz, 2H), 3.98 (dd, J = 9.2, 7.1 Hz, 1H), 4.67 (p, J = 6.1 Hz, 1H), 4.89 (s, 11H), 5.74 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.79 (q, J = 6.6 Hz, 1H), 7.13 (d, J = 8.7 Hz, 1H), 7.55 (m, 2H), 7.71 (m, 4H), 7.98 (d, J = 2.4 Hz, 1H)
1bw	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.54 (t, J = 5.7 Hz, 4H), 1.95 (dd, J = 13.3, 7.0 Hz, 1H), 2.22 (dd, J = 13.3, 9.0 Hz, 1H), 2.41 (s, 3H), 2.95 (d, J = 11.5 Hz, 1H), 3.13 (d,

	J = 11.4 Hz, 1H), 3.47 (ddt, J = 19.9, 12.9, 5.9 Hz, 2H), 3.64 (dd, J = 13.3, 6.4 Hz, 2H), 3.94 (t, J = 8.1 Hz, 1H), 4.91 (s, 10H), 5.77 (s, 1H), 6.43 (d, J = 2.3 Hz, 1H), 6.83 (q, J = 6.6 Hz, 1H), 7.49 (m, 2H), 7.83 (m, 7H), 8.01 (d, J = 2.4 Hz, 1H), 8.13 (d, J = 1.9 Hz, 1H)
1bx	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.59 (m, 4H), 2.05 (dd, J = 13.5, 7.2 Hz, 1H), 2.39 (s, 4H), 3.12 (d, J = 11.7 Hz, 1H), 3.24 (d, J = 11.8 Hz, 1H), 3.51 (m, 2H), 3.67 (dd, J = 13.7, 6.2 Hz, 2H), 4.13 (dd, J = 9.1, 7.2 Hz, 1H), 5.19 (s, 2H), 6.41 (d, J = 2.3 Hz, 1H), 6.79 (q, J = 6.5 Hz, 1H), 7.21 (t, J = 8.6 Hz, 1H), 7.42 (m, 7H), 7.60 (d, J = 1.8 Hz, 1H), 7.72 (m, 2H), 7.97 (d, J = 2.3 Hz, 1H)
1by	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.33 (dd, J = 6.1, 1.6 Hz, 6H), 1.58 (t, J = 5.2 Hz, 4H), 2.04 (dd, J = 13.5, 7.1 Hz, 1H), 2.22 (s, 3H), 2.35 (m, 4H), 3.10 (d, J = 11.8 Hz, 1H), 3.23 (d, J = 11.8 Hz, 1H), 3.49 (ddt, J = 20.8, 13.6, 5.9 Hz, 2H), 3.66 (dd, J = 13.6, 6.9 Hz, 2H), 4.07 (dd, J = 9.2, 7.1 Hz, 1H), 4.63 (p, J = 6.1 Hz, 1H), 5.76 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.75 (q, J = 6.7 Hz, 1H), 6.97 (d, J = 8.2 Hz, 1H), 7.45 (d, J = 8.2 Hz, 2H), 7.57 (d, J = 1.8 Hz, 1H), 7.71 (m, 2H), 7.96 (d, J = 2.3 Hz, 1H)
1bz	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.06 (t, J = 7.4 Hz, 3H), 1.29 (m, 1H), 1.58 (d, J = 5.9 Hz, 4H), 1.83 (h, J = 7.1 Hz, 2H), 2.02 (dd, J = 13.4, 6.9 Hz, 1H), 2.29 (dd, J = 13.3, 9.1 Hz, 1H), 2.39 (s, 3H), 3.08 (d, J = 11.6 Hz, 1H), 3.21 (d, J = 11.5 Hz, 1H), 3.48 (ddd, J = 21.8, 12.6, 5.8 Hz, 2H), 3.65 (dd, J = 13.6, 7.3 Hz, 2H), 4.04 (q, J = 7.1, 6.4 Hz, 3H), 4.97 (s, 1H), 5.75 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.5 Hz, 1H), 7.14 (t, J = 8.6 Hz, 1H), 7.44 (m, 2H), 7.59 (d, J = 1.9 Hz, 1H), 7.72 (m, 2H), 7.98 (d, J = 2.3 Hz, 1H)
1ca	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.99 (t, J = 7.4 Hz, 3H), 1.53 (m, 6H), 1.79 (dq, J = 8.6, 6.5 Hz, 2H), 1.93 (dd, J = 13.2, 7.0 Hz, 1H), 2.20 (dd, J = 13.3, 9.1 Hz, 1H), 2.39 (s, 3H), 2.91 (d, J = 11.4 Hz, 1H), 3.11 (d, J = 11.4 Hz, 1H), 3.47 (ddt, J = 20.0, 13.0, 5.9 Hz, 2H), 3.64 (dd, J = 13.8, 5.7 Hz, 2H), 3.88 (t, J = 8.0 Hz, 1H), 4.07 (t, J = 6.4 Hz, 2H), 5.75 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.78 (q, J = 6.7 Hz, 1H), 7.14 (t, J = 8.6 Hz, 1H), 7.43 (m, 2H), 7.59 (d, J = 1.9 Hz, 1H), 7.71 (m, 2H), 7.98 (d, J = 2.4 Hz, 1H)
1cb	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (s, 1H), 1.28 (s, 1H), 1.62 (d, J = 6.2 Hz, 8H), 2.06 (dd, J = 13.3, 7.3 Hz, 2H), 2.41 (s, 8H), 2.65 (s, 5H), 3.15 (d, J = 12.0 Hz, 2H), 3.25 (m, 2H), 3.52 (s, 3H), 3.58 (s, 2H), 3.70 (d, J = 13.3 Hz, 4H), 4.21 (t, J = 8.4 Hz, 2H), 5.89 (s, 1H), 6.00 (m, 1H), 6.44 (d, J = 2.3 Hz, 2H), 6.89 (q, J = 6.4 Hz, 2H), 7.80 (m, 10H), 8.04 (d, J = 2.3 Hz, 2H), 8.12 (t, J = 7.9 Hz, 2H)
1cc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 6.2 Hz, 1H), 1.57 (m, 4H), 1.99 (dd, J = 13.4, 7.0 Hz, 1H), 2.25 (dd, J = 13.4, 9.1 Hz, 1H), 2.40 (s, 3H), 3.01 (d, J = 11.6 Hz, 1H), 3.15 (m, 4H), 3.32 (s, 1H), 3.48 (ddt, J = 20.3, 12.8, 5.9 Hz, 2H), 3.65 (dd, J = 13.9, 7.0 Hz, 2H), 3.97 (dd, J = 9.1, 7.0 Hz, 1H), 4.89 (m, 2H), 5.75 (s, 1H), 6.44 (d, J = 2.4 Hz, 1H), 6.85 (q, J = 6.6 Hz, 1H), 7.75 (d, J = 1.7 Hz, 1H), 7.84 (m, 2H), 7.94 (d, J = 8.5 Hz, 2H), 8.03 (m, 3H)
1cd	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.05 (t, J = 7.4 Hz, 3H), 1.51 (q, J = 6.5, 5.7 Hz, 4H), 1.80 (h, J = 6.7, 6.1 Hz, 3H), 1.89 (d, J = 1.5 Hz, 1H), 2.07 (dd, J = 12.7, 9.2 Hz, 1H), 2.39 (s, 3H), 2.67 (d, J = 11.2 Hz, 1H), 2.94 (t, J = 11.8 Hz, 1H), 3.40 (m, 3H), 3.64 (m, 3H), 3.96 (t, J = 6.4 Hz, 2H), 4.89 (m, 1H), 5.75 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.75 (m, 1H), 6.99 (m, 2H), 7.59 (dd, J = 9.0, 2.0 Hz, 3H), 7.71 (m, 2H), 7.97 (d, J = 2.4 Hz, 1H)

1ce	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (d, J = 16.9 Hz, 4H), 1.57 (s, 5H), 2.02 (m, 1H), 2.27 (dd, J = 13.3, 8.8 Hz, 1H), 2.40 (s, 3H), 2.60 (m, 6H), 3.06 (d, J = 11.4 Hz, 1H), 3.20 (d, J = 11.3 Hz, 1H), 3.34 (s, 1H), 3.48 (s, 3H), 3.56 (t, J = 6.7 Hz, 2H), 3.70 (m, 7H), 4.02 (s, 1H), 4.98 (d, J = 6.2 Hz, 1H), 5.76 (s, 1H), 6.43 (d, J = 2.3 Hz, 1H), 6.82 (q, J = 6.5 Hz, 1H), 7.71 (s, 1H), 7.80 (m, 4H), 7.93 (d, J = 8.0 Hz, 2H), 8.01 (d, J = 2.3 Hz, 1H)
1cf	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.90 (t, J = 6.4 Hz, 1H), 1.30 (dd, J = 12.6, 4.9 Hz, 6H), 1.55 (m, 5H), 1.93 (dd, J = 13.2, 7.0 Hz, 1H), 2.19 (qd, J = 9.4, 3.3 Hz, 1H), 2.40 (s, 3H), 2.92 (d, J = 11.4 Hz, 1H), 3.11 (d, J = 11.3 Hz, 1H), 3.48 (m, 2H), 3.65 (dd, J = 13.7, 6.3 Hz, 2H), 3.88 (dd, J = 8.9, 7.2 Hz, 1H), 4.87 (d, J = 12.3 Hz, 1H), 4.97 (d, J = 12.9 Hz, 2H), 5.75 (s, 1H), 6.43 (d, J = 2.3 Hz, 1H), 6.83 (q, J = 6.7 Hz, 1H), 7.73 (s, 1H), 7.84 (m, 4H), 8.00 (m, 3H)
1cg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (s, 1H), 1.58 (d, J = 5.9 Hz, 4H), 2.03 (dd, J = 13.4, 6.9 Hz, 1H), 2.30 (dd, J = 13.3, 9.2 Hz, 1H), 2.40 (s, 3H), 3.11 (d, J = 11.7 Hz, 1H), 3.23 (d, J = 11.5 Hz, 1H), 3.48 (ddd, J = 28.3, 12.4, 5.7 Hz, 2H), 3.65 (dd, J = 13.7, 7.2 Hz, 2H), 4.07 (m, 1H), 5.76 (s, 1H), 6.43 (d, J = 2.3 Hz, 1H), 6.82 (q, J = 6.5 Hz, 1H), 7.70 (d, J = 1.7 Hz, 1H), 7.79 (dt, J = 13.1, 8.1 Hz, 4H), 7.99 (m, 3H)
1ch	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 3.6 Hz, 1H), 1.63 (q, J = 5.8 Hz, 5H), 2.07 (dd, J = 13.5, 7.5 Hz, 1H), 2.37 (dd, J = 13.5, 9.0 Hz, 1H), 3.04 (s, 3H), 3.15 (d, J = 24.6 Hz, 6H), 3.27 (m, 1H), 3.52 (dt, J = 24.6, 8.3 Hz, 2H), 3.65 (m, 2H), 4.23 (t, J = 8.1 Hz, 1H), 6.66 (q, J = 7.0 Hz, 1H), 7.51 (d, J = 8.1 Hz, 2H), 7.68 (m, 6H)
1ci	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.58 (d, J = 5.6 Hz, 4H), 2.03 (dd, J = 13.3, 7.1 Hz, 1H), 2.30 (dd, J = 13.4, 9.2 Hz, 1H), 2.39 (s, 3H), 3.09 (d, J = 11.7 Hz, 1H), 3.22 (d, J = 11.7 Hz, 1H), 3.49 (ddd, J = 21.1, 12.5, 5.6 Hz, 2H), 3.66 (dd, J = 13.7, 6.7 Hz, 2H), 3.90 (s, 3H), 4.06 (dd, J = 9.2, 7.1 Hz, 1H), 5.76 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.6 Hz, 1H), 7.17 (t, J = 8.9 Hz, 1H), 7.46 (m, 2H), 7.60 (d, J = 1.8 Hz, 1H), 7.73 (m, 2H), 7.98 (d, J = 2.3 Hz, 1H)
1cj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (d, J = 17.0 Hz, 1H), 1.57 (d, J = 5.5 Hz, 4H), 2.01 (dd, J = 13.2, 7.0 Hz, 1H), 2.28 (dd, J = 13.3, 9.0 Hz, 1H), 2.40 (s, 3H), 2.79 (s, 2H), 2.91 (s, 2H), 3.07 (d, J = 12.1 Hz, 1H), 3.20 (d, J = 11.4 Hz, 1H), 3.49 (m, 4H), 3.65 (dd, J = 13.5, 6.9 Hz, 2H), 3.74 (s, 2H), 4.02 (t, J = 8.1 Hz, 1H), 4.99 (s, 1H), 5.76 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.82 (q, J = 6.6 Hz, 1H), 7.52 (d, J = 7.9 Hz, 2H), 7.76 (m, 5H), 8.01 (d, J = 2.4 Hz, 1H)
1ck	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (d, J = 11.1 Hz, 1H), 1.51 (q, J = 6.8, 6.0 Hz, 4H), 1.78 (dd, J = 13.0, 7.0 Hz, 1H), 1.89 (s, 2H), 2.07 (dd, J = 13.1, 9.1 Hz, 1H), 2.40 (s, 3H), 2.68 (d, J = 11.1 Hz, 1H), 2.95 (d, J = 11.1 Hz, 1H), 3.03 (s, 3H), 3.11 (s, 3H), 3.22 (s, 2H), 3.45 (m, 3H), 3.63 (q, J = 7.9, 7.5 Hz, 3H), 5.75 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.82 (q, J = 6.6 Hz, 1H), 7.53 (d, J = 7.9 Hz, 2H), 7.70 (m, 1H), 7.80 (m, 4H), 8.01 (d, J = 2.5 Hz, 1H)
1cl	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.04 (d, J = 6.7 Hz, 6H), 1.29 (m, 1H), 1.60 (d, J = 6.0 Hz, 5H), 2.05 (ddd, J = 13.7, 7.0, 3.9 Hz, 2H), 2.36 (m, 4H), 3.13 (d, J = 11.9 Hz, 1H), 3.24 (d, J = 11.6 Hz, 1H), 3.51 (ddd, J = 25.5, 14.3, 7.1 Hz, 2H), 3.69 (d, J = 13.8 Hz, 3H), 3.77 (d, J = 6.4 Hz, 2H), 4.14 (t, J = 8.3 Hz, 1H), 4.93 (s, 8H), 6.41 (d, J = 2.3 Hz, 1H), 6.77 (q, J = 6.6 Hz, 1H), 6.99 (m, 2H), 7.60 (m, 3H), 7.72 (m, 2H), 7.96 (d, J = 2.3 Hz, 1H)
1cm	¹ H NMR (400 MHz, MeOH-d ₄): δ 1.14 (t, J = 7.1 Hz, 3H), 1.26 (t, J = 7.3 Hz, 3H),

	1.55 (q, J = 4.8 Hz, 4H), 1.90 (m, 1H), 2.18 (dd, J = 13.2, 9.0 Hz, 1H), 2.40 (s, 3H), 2.87 (d, J = 11.4 Hz, 1H), 3.09 (d, J = 11.3 Hz, 1H), 3.30 (m, 4H), 3.54 (dddd, J = 37.2, 30.6, 15.1, 5.9 Hz, 6H), 3.84 (dd, J = 9.0, 6.9 Hz, 1H), 5.76 (s, 1H), 6.42 (d, J = 2.3 Hz, 1H), 6.83 (q, J = 6.6 Hz, 1H), 7.48 (m, 2H), 7.70 (d, J = 1.6 Hz, 1H), 7.80 (m, 4H), 8.01 (d, J = 2.3 Hz, 1H)
1cn	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.84 (s, 3H), 1.06 (s, 10H), 1.30 (m, 1H), 1.56 (t, J = 5.4 Hz, 4H), 2.00 (dd, J = 13.4, 6.8 Hz, 1H), 2.27 (dd, J = 13.2, 9.0 Hz, 1H), 2.41 (s, 3H), 3.04 (d, J = 11.5 Hz, 1H), 3.19 (d, J = 11.4 Hz, 1H), 3.48 (ddt, J = 20.4, 13.0, 5.9 Hz, 2H), 3.65 (s, 4H), 4.03 (t, J = 8.1 Hz, 1H), 5.77 (s, 1H), 6.43 (d, J = 2.2 Hz, 1H), 6.79 (q, J = 6.6 Hz, 1H), 6.98 (d, J = 8.4 Hz, 2H), 7.60 (m, 4H), 7.75 (d, J = 8.2 Hz, 1H), 7.97 (d, J = 2.3 Hz, 1H)
1co	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (d, J = 5.8 Hz, 1H), 1.55 (t, J = 5.7 Hz, 4H), 1.98 (m, 3H), 2.25 (dd, J = 13.3, 9.2 Hz, 1H), 2.39 (s, 3H), 2.80 (t, J = 6.5 Hz, 2H), 3.02 (d, J = 11.6 Hz, 1H), 3.17 (d, J = 11.6 Hz, 1H), 3.47 (ddt, J = 20.5, 13.0, 5.9 Hz, 2H), 3.64 (dt, J = 13.8, 5.9 Hz, 2H), 4.00 (dd, J = 9.2, 7.1 Hz, 1H), 4.16 (m, 2H), 5.75 (s, 1H), 6.40 (d, J = 2.3 Hz, 1H), 6.76 (m, 2H), 7.33 (d, J = 6.5 Hz, 2H), 7.54 (d, J = 1.8 Hz, 1H), 7.63 (dd, J = 8.3, 1.9 Hz, 1H), 7.71 (d, J = 8.3 Hz, 1H), 7.95 (d, J = 2.4 Hz, 1H)
1cp	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.59 (t, J = 5.3 Hz, 4H), 2.05 (dd, J = 13.5, 7.1 Hz, 1H), 2.31 (dd, J = 13.4, 9.3 Hz, 1H), 2.42 (s, 3H), 3.11 (d, J = 11.7 Hz, 1H), 3.24 (d, J = 11.7 Hz, 1H), 3.51 (m, 2H), 3.67 (s, 2H), 4.07 (dd, J = 9.3, 7.1 Hz, 1H), 4.89 (s, 1H), 5.78 (s, 1H), 5.99 (m, 1H), 6.46 (d, J = 2.3 Hz, 1H), 6.89 (q, J = 6.5 Hz, 1H), 7.91 (m, 2H), 8.00 (dd, J = 8.3, 2.0 Hz, 1H), 8.07 (d, J = 2.4 Hz, 1H), 8.34 (m, 3H), 8.55 (d, J = 8.9 Hz, 1H), 9.33 (d, J = 5.9 Hz, 1H)
1cq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.97 (s, 1H), 7.77 (s, 2H), 7.67 (d, J = 15.8 Hz, 2H), 7.51 (s, 1H), 7.26 (d, J = 7.8 Hz, 1H), 6.42 (s, 1H), 4.69 (s, 2H), 4.14 (s, 1H), 3.68 (s, 2H), 3.51 (s, 3H), 3.23 (s, 1H), 3.13 (d, J = 11.6 Hz, 1H), 2.38 (d, J = 14.0 Hz, 6H), 2.05 (s, 1H), 1.60 (s, 4H), 1.29 (s, 3H)
1cr	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.97 (s, 2H), 7.74 (q, J = 8.3 Hz, 4H), 7.62 (s, 2H), 7.46 (d, J = 7.4 Hz, 5H), 6.78 (q, J = 6.7 Hz, 2H), 6.41 (s, 2H), 5.75 (s, 2H), 4.65 (s, 3H), 4.07 (t, J = 8.1 Hz, 2H), 3.64 (s, 4H), 3.52 (d, J = 6.9 Hz, 1H), 3.46 (d, J = 16.2 Hz, 4H), 3.22 (d, J = 11.8 Hz, 2H), 3.10 (d, J = 11.8 Hz, 2H), 2.38 (d, J = 7.9 Hz, 10H), 2.29 (s, 1H), 2.03 (dd, J = 13.4, 7.0 Hz, 2H), 1.58 (d, J = 5.6 Hz, 7H), 1.30 (d, J = 13.4 Hz, 5H)
1cs	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.43 (d, J = 2.5 Hz, 1H), 7.99 (q, J = 3.3 Hz, 2H), 7.79 (d, J = 8.3 Hz, 1H), 7.72 (d, J = 8.1 Hz, 1H), 7.63 (s, 1H), 6.83 (dd, J = 20.4, 7.6 Hz, 2H), 6.42 (d, J = 2.3 Hz, 1H), 5.74 (s, 1H), 4.35 (q, J = 7.0 Hz, 2H), 3.96 (d, J = 8.9 Hz, 1H), 3.64 (s, 3H), 3.48 (dd, J = 25.9, 11.9 Hz, 2H), 3.15 (d, J = 11.7 Hz, 1H), 2.99 (d, J = 11.4 Hz, 1H), 2.39 (s, 3H), 2.24 (s, 1H), 2.02 – 1.94 (m, 1H), 1.56 (s, 4H), 1.38 (t, J = 7.1 Hz, 3H), 1.28 (s, 1H).


Example 1cp: (S)-8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-(trifluoromethyl)-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 1u) starting with (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 10d).

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.57 (br. s., 4 H) 1.91 - 2.01 (m, 1 H) 2.18 - 2.27 (m, 1 H) 2.33 (d, J=11.71 Hz, 6 H) 2.88 - 3.00 (m, 1 H) 3.08 - 3.19 (m, 1 H) 3.38 - 3.56 (m, 2 H) 3.58 - 3.75 (m, 2 H) 3.85 - 3.98 (m, 1 H) 5.65 (s, 1 H) 6.55 - 6.70 (m, 1 H) 6.92 - 7.04 (m, 1 H) 7.19 - 7.28 (m, 1 H) 7.38 - 7.46 (m, 1 H) 7.46 - 7.53 (m, 1 H) 7.72 (s, 1 H) 7.83 (s, 2 H) 8.22 - 8.35 (m, 1 H) LCMS (MH⁺): 690.

Example 2: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(piperidin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: A solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(1,2,3,6-tetrahydropyridin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (Example 1f) (150 mg, 0.15 mmol) in MeOH (5 mL) was hydrogenated in an H-Cube apparatus using a 10% (w/w) Pd/C cartridge with a flow rate of 1.0 mL/min at RT. The resulting eluent was concentrated in vacuo and The product was

purified by column chromatography using an Isco Gold reversed phase silica cartridge (100% CH₂Cl₂ to 90:9:1 CH₂Cl₂:MeOH:conc. NH₄OH) to provide (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(1,2,3,6-tetrahydropyridin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate.

5


Step 2: Hydrolysis of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(1,2,3,6-tetrahydropyridin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate using the LiOH general method provided the title compound as an off-white solid.

10

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.49 - 1.69 (m, 4 H) 2.00 - 2.18 (m, 3 H) 2.21 - 2.35 (m, 1 H) 2.38 (s, 3 H) 2.92 - 3.05 (m, 1 H) 3.08 (d, J=0.44 Hz, 2 H) 3.10 - 3.18 (m, 2 H) 3.25 (d, J=11.71 Hz, 1 H) 3.38 - 3.72 (m, 7 H) 4.09 (t, J=7.88 Hz, 1 H) 5.69 (s, 1 H) 6.41 (d, J=2.29 Hz, 1 H) 6.74 (q, J=6.80 Hz, 1 H) 7.34 (d, J=1.12 Hz, 1 H) 7.43 (d, J=8.15 Hz, 1 H) 7.71 (d, J=8.44 Hz, 1 H) 7.91 (d, J=2.20 Hz, 1 H). LCMS (MH⁺): 613.

15

Example 3a: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(1-(methanesulfonyl)piperidin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



20

Step 1: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(1,2,3,6-tetrahydropyridin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (320 mg, 0.413 mmol) in CH₂Cl₂ (5.0 mL) was added methanesulfonyl chloride (47 mg, 0.41 mmol) and triethylamine (94 mg, 0.83 mmol), and the reaction was stirred for 1.5 h at RT and then concentrated *in vacuo*. The product was purified by column chromatography using an Isco Gold reversed phase silica cartridge (100% CH₂Cl₂ to

25

90:9:1 CH₂Cl₂:MeOH:conc. NH₄OH) to provide (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(1-(methylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as an off-white solid.

5

Step 2: A solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(1-(methylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (290 mg, 0.340 mmol, Step 1) in MeOH (10 mL) was hydrogenated in an H-Cube apparatus using a 10% (w/w) Pd/C cartridge with a flow rate of 1.0 mL/min at RT. The resulting eluent was concentrated in vacuo and The product was purified by column chromatography using an Isco Gold reversed phase silica cartridge (100% CH₂Cl₂ to 90:9:1 CH₂Cl₂:MeOH:conc. NH₄OH) to provide (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(1-(methylsulfonyl)piperidin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate.

15

Step 3: Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(1-(methylsulfonyl)piperidin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provided the title compound as an off-white solid.

20

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.53 - 1.65 (m, 4 H) 1.80 (qd, J=12.57, 3.98 Hz, 2 H) 1.94 - 2.02 (m, 2 H) 2.02 - 2.12 (m, 1 H) 2.31 (dd, J=13.42, 9.27 Hz, 1 H) 2.38 (s, 3 H) 2.67 - 2.94 (m, 3 H) 2.86 (s, 3 H) 3.07 - 3.28 (m, 2 H) 3.37 - 3.74 (m, 4 H) 3.78 - 3.92 (m, 2 H) 4.08 (dd, J=9.15, 7.20 Hz, 1 H) 5.71 (s, 1 H) 6.39 (d, J=2.29 Hz, 1 H) 6.64 - 6.82 (m, 1 H) 7.31 (d, J=1.71 Hz, 1 H) 7.42 (dd, J=8.25, 1.76 Hz, 1 H) 7.67 (d, J=8.10 Hz, 1 H) 7.89 (d, J=2.29 Hz, 1 H). LCMS (MH⁺): 693.

25

Using the generic scheme below, the following examples of Table 2a can be prepared as described above for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(1-(methylsulfonyl)piperidin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 3a).

30





Table 2a.




Ex. No.	Cy	CAS Name	LCMS (MH ⁺)
3b		(S)-8-(6-((R)-1-(4-(1-acetylpiperidin-4-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	656.7
3c		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(tetrahydro-2H-pyran-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	615.6

Table 2b.

NMR Data for Compounds of Table 2a

Ex. No.	¹ H NMR
3b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.54 - 1.82 (m, 6 H) 1.86 - 1.99 (m, 2 H) 2.05 - 2.18 (m, 4 H) 2.36 - 2.38 (m, 3 H) 2.48 (dd, J=13.69, 8.86 Hz, 1 H) 2.66 - 2.81 (m, 1 H) 2.88 - 3.03 (m, 1 H) 3.19 - 3.27 (m, 1 H) 3.31 - 3.40 (m, 1 H) 3.60 - 3.95 (m, 4 H) 4.05 (d, J=13.08 Hz, 1 H) 4.55 (t, J=8.66 Hz, 1 H) 4.67 (d, J=13.13 Hz, 1 H) 6.39 (d, J=2.39 Hz, 1 H) 6.50 (br. s., 1 H) 6.79 - 6.87 (m, 1 H), 7.36 (s, 1 H) 7.47 (dd, J=8.22, 1.64 Hz, 1 H) 7.64 (d, J=8.30 Hz, 1 H) 7.86 (d, J=2.39 Hz, 1 H)
3c	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.59 (d, J=5.08 Hz, 4 H) 1.72 - 1.89 (m, 4 H) 2.06 (dd, J=13.45, 7.15 Hz, 1 H) 2.32 (dd, J=13.45, 9.25 Hz, 1 H) 2.38 (s, 3 H) 2.82 - 2.95 (m, 1 H) 3.07 - 3.16 (m, 1 H) 3.25 (d, J=11.76 Hz, 1 H) 3.36 - 3.74 (m, 6 H) 4.03 (dt, J=11.16, 2.96 Hz, 2 H) 4.08 (dd, J=9.15, 7.20 Hz, 1 H) 5.71 (s, 1 H) 6.39 (d, J=2.29 Hz, 1 H) 6.72 (q, J=6.75 Hz, 1 H) 7.29 (d, J=1.71 Hz, 1 H) 7.41 (dd, J=8.20, 1.76 Hz, 1 H) 7.67 (d, J=8.10 Hz, 1 H) 7.88 (d, J=2.34 Hz, 1 H)

Example 4: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-4'-(methoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



5
Step 1: To a solution of (S)-8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (product of Step 3, Example 10m) (135 mg, 0.18 mmol) in dioxane (2 mL) was added (3-methoxy-4-(methoxycarbonyl)phenyl)boronic acid (84 mg, 0.4
10 mmol) and Cs₂CO₃ (48 mg, 0.16 mmol). The reaction was heated to 80 °C for 16 h, cooled to RT, and filtered. The solvent was removed *in vacuo*. Purification via normal phase silica gel chromatography (CH₂Cl₂/Heptane) provided (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-4'-(methoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid
15 as an off-white solid.


Step 2: N-CBZ Deprotection was accomplished via method B to yield (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-4'-(methoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid as an off-white solid.

5

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.66 (d, J=5.47 Hz, 4 H) 2.03 - 2.17 (m, 1 H) 2.42 (s, 4 H) 3.16 - 3.30 (m, 2 H) 3.47 - 3.81 (m, 4 H) 3.89 (s, 3 H) 3.97 (s, 3 H) 4.26 - 4.45 (m, 1 H) 6.40 - 6.52 (m, 1 H) 6.82 - 6.96 (m, 1 H) 7.30 - 7.37 (m, 1 H) 7.40 (s, 1 H) 7.76 (s, 1 H) 7.80 - 7.93 (m, 4 H) 7.99 - 8.09 (m, 1 H). LCMS: 696.7.

10

Example 5a: (S)-8-(2-amino-6-((R)-1-(3'-(ethoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




15 The title compound was made according to the procedures described for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-4'-(methoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 4).

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.42 (t, J=7.13 Hz, 3 H) 1.61 (br. s., 4 H) 2.02 - 2.14 (m, 1 H) 2.28 - 2.40 (m, 1 H) 2.42 (s, 3 H) 3.06 - 3.19 (m, 1 H) 3.21 - 3.30 (m, 1 H) 3.40 - 3.60 (m, 2 H) 3.62 - 3.80 (m, 2 H) 4.01 - 4.19 (m, 1 H) 4.41 (d, J=7.22 Hz, 2 H) 5.76 (s, 1 H) 6.45 (d, J=2.34 Hz, 1 H) 6.79 - 6.92 (m, 1 H) 7.60 (s, 1 H) 7.70 (d, J=1.56 Hz, 1 H) 7.80 (d, J=1.56 Hz, 1 H) 7.84 (s, 1 H) 7.90 - 7.97 (m, 1 H) 8.02 (d, J=2.15 Hz, 1 H) 8.05 (s, 1 H) 8.31 (s, 1 H) 680.7. LCMS (MH⁺): 578.7.

25

Example 5b: (S)-8-(2-amino-6-((R)-1-(4'-(ethoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




- 5 The title compound was made according to the procedures described for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-4'-(methoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 4).

¹H NMR (400 MHz, MeOH-d₄): δ ppm 0.88 (m, 4H), 1.30 (d, J = 17.4 Hz, 10H), 1.40 (t, J = 7.1 Hz, 4H), 1.59 (d, J = 5.8 Hz, 5H), 2.05 (dd, J = 13.5, 7.2 Hz, 1H), 2.35 (m, 5H), 3.11 (d, J = 11.7 Hz, 1H), 3.24 (d, J = 11.7 Hz, 1H), 3.49 (ddd, J = 28.1, 12.7, 5.7 Hz, 2H), 3.66 (dd, J = 13.2, 7.3 Hz, 3H), 4.07 (t, J = 8.1 Hz, 1H), 4.38 (q, J = 7.1 Hz, 2H), 4.82 (d, J = 9.7 Hz, 1H), 4.91 (s, 2H), 5.75 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.83 (q, J = 6.5 Hz, 1H), 7.72 (d, J = 1.6 Hz, 1H), 7.81 (m, 4H), 8.00 (d, J = 2.4 Hz, 1H), 8.10 (m, 2H). LCMS (MH⁺): 681.

15

Example 6: (S)-8-(2-amino-6-((R)-1-(4-(3-carboxypropyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a solution of 9-borabicyclo[3.3.1]nonane (2.0 mL, 0.5 M in THF, 1.0 mmol) was added methyl but-3-enoate (100 μ L, 1.0 mmol) and stirred at RT for 2 h to prepare the 9-BBN/butane solution.


5 *Step 2:* To a solution of (S)-8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (product of Step 3, Example 10m) (250 mg, 0.32 mmol) in THF (2 mL) was added sequentially PdCl₂(dppf)CH₂Cl₂ (8 mg, 0.01 mmol), NaOEt (66 mg, 1 mmol) and the prepared 9-BBN/butene solution from Step 1. The reaction was heated to 65 °C for 2 h,
10 then cooled to RT. The reaction was extracted with EtOAc, brine and dried over Na₂SO₄ and concentrated *in vacuo*. The product was purified by column chromatography using an Isco Gold reversed phase silica cartridge (100% CH₂Cl₂ to 90:9:1 CH₂Cl₂:MeOH:conc. NH₄OH) to provide (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(4-methoxy-4-oxobutyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-
15 dicarboxylate as an off-white solid.

Step 3: N-CBZ Deprotection was accomplished via method B to provide (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(4-methoxy-4-oxobutyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as an off-white solid.

20 *Step 4:* Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(4-methoxy-4-oxobutyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate was carried out using the LiOH general method providing the title compound as an off-white solid.

25 ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.40 - 1.61 (m, 4 H) 1.76 - 1.93 (m, 3 H) 2.24 (t, J=7.35 Hz, 2 H) 2.27 - 2.37 (m, 4 H) 2.58 - 2.74 (m, 2 H) 3.10 (br. s., 2 H) 3.53 (br. s., 4 H) 4.42 (br. s., 1 H) 5.71 (br. s., 1 H) 6.00 (br. s., 2 H) 6.38 (d, J=2.20 Hz, 1 H) 7.00 (q, J=6.87 Hz, 1 H) 7.29 (d, J=1.51 Hz, 1 H) 7.32 - 7.41 (m, 1 H) 7.60 (s, 1 H) 8.05 (d, J=2.29 Hz, 1 H) 8.94 (br. s., 1 H)
30 10.20 (br. s, 1 H) 12.14 (br. s., 1 H). LCMS (MH⁺): 618.6.

Example 7: (S)-8-(2-amino-6-((R)-1-(4-(2-carboxyethyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




5 The title compound was made as described for (S)-8-(2-amino-6-((R)-1-(4-(3-carboxypropyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 6).

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.56 (t, J=5.54 Hz, 4 H) 1.97 (s, 2 H) 2.04 (dd, J=13.30, 7.10 Hz, 1 H) 2.29 (dd, J=13.67, 9.18 Hz, 1 H) 2.35 (s, 3 H) 2.59 - 2.68 (m, 2 H) 2.97 (t, J=7.49 Hz, 2 H) 3.06 - 3.13 (m, 1 H) 3.23 (d, J=11.86 Hz, 1 H) 3.39 - 3.55 (m, 2 H) 3.57 - 3.75 (m, 2 H) 4.06 (dd, J=9.05, 7.30 Hz, 1 H) 5.72 (s, 1 H) 6.36 (d, J=2.29 Hz, 1 H) 6.71 (q, J=6.61 Hz, 1 H) 7.28 (d, J=1.61 Hz, 1 H) 7.37 (dd, J=8.20, 1.46 Hz, 1 H) 7.62 (d, J=8.10 Hz, 1 H) 7.83 (d, J=2.25 Hz, 1 H). LCMS (MH⁺): 604.

15

Example 9: (S)-8-(2-amino-6-((R)-1-(4-(3-ethoxy-3-oxopropyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




20 *Step 1:* To a solution of (S)-8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-

diazaspiro[4.5]decane-3-carboxylic acid (product of Step 3, Example 10m) (240 mg, 0.33 mmol) in ethanol (8 mL) was added (E)-ethyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)acrylate (110 mg, 0.49 mmol), PdCl₂(PPh₃)₂ (20 mg, 0.049 mmol) and KHCO₃ (170 mg, 0.05 mmol). The reaction was heated to 80 °C for 2 h, cooled to RT, and filtered. The solvent was removed
5 *in vacuo*. Purification via normal phase silica gel chromatography (CH₂Cl₂/heptane) provided ((S)-8-(2-amino-6-((R)-1-(4-((E)-3-ethoxy-3-oxoprop-1-en-1-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-(((2E,4Z)-2-vinylhexa-2,4-dien-1-yl)oxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid as a white solid.

10 *Step 2:* To a solution of ((S)-8-(2-amino-6-((R)-1-(4-((E)-3-ethoxy-3-oxoprop-1-en-1-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-(((2E,4Z)-2-vinylhexa-2,4-dien-1-yl)oxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (180 mg, 0.15 mmol) in MeOH (5 mL) was hydrogenated in an H-Cube apparatus using a 10% (w/w) Pd/C cartridge with a flow rate of 1.0 mL/min at RT. The resulting eluent was concentrated in
15 *vacuo* and the product was purified by column chromatography using an Isco Gold reversed phase silica cartridge (100% CH₂Cl₂ to 90:9:1 CH₂Cl₂:MeOH:conc. NH₄OH) to provide the title compound as a white solid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.14 (t, J=7.15 Hz, 3 H) 1.50 - 1.68 (m, 4 H) 1.94 - 1.99
20 (m, 2 H) 2.04 (dd, J=13.45, 7.20 Hz, 1 H) 2.30 (dd, J=13.47, 9.27 Hz, 1 H) 2.35 (s, 3 H) 2.66 (t, J=7.54 Hz, 2 H) 2.97 (t, J=7.52 Hz, 2 H) 3.07 - 3.14 (m, 1 H) 3.23 (d, J=11.76 Hz, 1 H) 3.39 - 3.72 (m, 4 H) 4.01 - 4.11 (m, 3 H) 5.70 (s, 1 H) 6.36 (d, J=2.34 Hz, 1 H) 6.72 (q, J=6.72 Hz, 1 H) 7.27 (d, J=1.61 Hz, 1 H) 7.35 (dd, J=8.15, 1.61 Hz, 1 H) 7.62 (d, J=8.05 Hz, 1 H) 7.83 (d, J=2.34 Hz, 1 H). LCMS (MH⁺): 632.1

25 **Example 10d: (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid**




Step 1: To a solution of (1R)-1-[4-chloro-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoroethanol (40 g, 138 mmol) in dioxane (400 mL) was added 4,6-dichloropyrimidin-2-amine (113 g, 690 mmol) and Cs₂CO₃ (132 g, 405 mmol). The mixture was heated for 24 h at 80 °C. The reaction was then cooled to RT and filtered. The solvent was removed *in vacuo*, then CH₂Cl₂ and heptane was added. The solvent volume was reduced until a solid precipitated out. The solid was filtered and the procedure repeated several times to provide 4-chloro-6-[(1R)-1-[4-chloro-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoroethoxy]pyrimidin-2-amine as a white solid.

Step 2: To a solution of 4-chloro-6-[(1R)-1-[4-chloro-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoroethoxy]pyrimidin-2-amine (57.3 g, 137 mmol, Step 1) in dioxane (500 mL) was added (S)-2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (48 g, 124.9 mmol), and NaHCO₃ (31.5 g, 375 mmol). After 5 h, an additional amount of NaHCO₃ (31.5 g, 375 mmol) was added and the reaction mixture was heated to 90 °C for 36 h. The reaction was then cooled to RT and filtered. Purification by normal phase silica gel column (EtOAc/heptane) provided (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

Step 3: N-CBZ Deprotection was accomplished via method B to provide (S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate an off-white solid.


Step 4: Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provided the title compound as an off-white solid.

Using the generic scheme below, the following examples of Table 3a were prepared as described above for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 10d).



5

Table 3a.



* Stereochemistry defined in name in table below

Ex. No.	R'	R''	R'''	CAS Name	LCMS (MH ⁺)
10a	H	H	H	8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-	532

				diazaspiro[4.5]decane-3-carboxylic acid	
10b	H	Cl	H	8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	566
10c	H	Cl	H	(R)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid R-Spiro	566
10d	H	Cl	H	(S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	566
10e	H	H	H	(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	532
10f	H	H	Cl	(S)-8-(2-amino-6-((R)-1-(3-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	566.9
10g	H	CF ₃	H	(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(trifluoromethyl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	600.6
10h	H	CH ₃	H	(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-methyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	546.6
10i	H	F	H	8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-fluoro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	550.5
10j	H		H	8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-methoxy-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	564.6
10k	Cl	H	H	8-(2-amino-6-((R)-1-(5-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	566.9
10l	H		H	(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-methoxy-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	564.6

10m	H	Br	H	(S)-8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	611
10n	Br	H	H	(S)-8-(2-amino-6-((R)-1-(5-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	611.5


Table 3b.

NMR Data for Compounds of Table 3a

Ex. No.	NMR
10a	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.59 (br. s., 4 H) 1.97 - 2.12 (m, 1 H) 2.24 - 2.35 (m, 1 H) 2.39 (s, 3 H) 3.11 (s, 1 H) 3.22 (s, 1 H) 3.40 - 3.58 (m, 2 H) 3.66 (br. s., 2 H) 3.95 - 4.17 (m, 1 H), 5.73 (s, 1 H) 6.39 (s, 1 H) 6.70 - 6.88 (m, 1 H) 7.42 (d, J=7.52 Hz, 1 H) 7.53 (dd, J=12.93, 7.57 Hz, 2 H) 7.75 (d, J=7.52 Hz, 1 H) 7.87 (s, 1 H)
10b	¹ H NMR (400 MHz, MeOH-d ₄) δ ppm 1.44 - 1.74 (m, 4 H) 1.88 - 2.06 (m, 1 H) 2.17 - 2.31 (m, 1 H) 2.39 (s, 3 H) 2.86 - 3.04 (m, 1 H) 3.09 3.21 (m, 1 H) 3.41-3.57 (m, 2 H) 3.58-3.77 (m, 2H) 3.85-4.05 (m, 1H) 5.63-5.76 (m, 1 H) 6.36-6.48 (m, 1 H) 6.76 6.91 (m, 1 H) 7.46-7.60 (m, 2 H) 7.67 - 7.79 (m, 1 H) 7.90 - 8.03 (m, 1 H)
10c	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.62 (br. s., 4 H) 2.04 - 2.17 (m, 1 H) 2.41 (s, 4 H) 3.10 - 3.21 (m, 1 H) 3.27 (s, 1 H) 3.44 - 3.58 (m, 2 H) 3.60 - 3.79 (m, 2 H) 4.05 - 4.18 (m, 1 H) 5.71 (s, 1 H) 6.44 (d, J=2.15 Hz, 1 H) 6.75 - 6.91 (m, 1 H) 7.52 (s, 2 H) 7.66 - 7.80 (m, 1 H) 7.96 (d, J=2.15 Hz, 1 H)
10d	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.59 (t, J=5.30 Hz, 4 H) 1.97 - 2.12 (m, 1 H) 2.31 (dd, J=13.45, 9.25 Hz, 1 H) 2.38 (s, 3 H) 3.11 (d, J=11.76 Hz, 1 H) 3.25 (d, J=11.71 Hz, 1 H) 3.38 - 3.57 (m, 2 H), 3.58 - 3.74 (m, 2 H) 4.08 (dd, J=9.15, 7.15 Hz, 1 H) 5.69 (s, 1 H) 6.41 (d, J=2.39 Hz, 1 H) 6.82 (q, J=6.61 Hz, 1 H) 7.44 - 7.57 (m, 2 H) 7.71 (d, J=8.35 Hz, 1 H) 7.93 (d, J=2.34 Hz, 1 H)
10e	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.71 (dt, J=18.13, 5.48 Hz, 4 H) 2.08 (dd, J=13.62, 8.49 Hz, 1 H) 2.37 (s, 3 H) 2.47 (dd, J=13.59, 8.96 Hz, 1 H) 3.62 - 3.90 (m, 4 H) 4.54 (t, J=8.71 Hz, 1 H) 6.38 (d, J=2.34 Hz, 1 H) 6.48 (br. s., 1 H) 6.85 (q, J=6.04 Hz, 1 H) 7.46 (dd, J=7.86, 1.07 Hz, 1 H) 7.52 - 7.59 (m, 1 H) 7.61 - 7.67 (m, 1 H) 7.70 (d, J=7.76 Hz, 1 H) 7.84 (d, J=2.39 Hz, 1 H)
10f	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.49 - 1.68 (m, 4 H) 2.04 (dd, J=13.45, 7.15 Hz, 1 H) 2.30 (dd, J=13.45, 9.25 Hz, 1 H) 2.35 (s, 3 H) 3.05 - 3.26 (m, 2 H) 3.38 - 3.77 (m, 5 H) 4.06 (dd, J=9.10, 7.15 Hz, 1 H) 5.60 (s, 1 H) 6.18 (q, J=6.56 Hz, 1 H) 6.39 (d, J=2.34 Hz, 1 H) 7.49 - 7.59 (m, 1 H) 7.60 - 7.74 (m, 3 H)
10g	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.51 - 1.64 (m, 4 H) 2.03 (dd, J=13.45, 7.15 Hz, 1 H) 2.29 (dd, J=13.47, 9.27 Hz, 1 H) 2.37 (s, 3 H) 3.03 - 3.25 (m, 2 H) 3.37 - 3.54 (m, 2 H) 3.56 - 3.75 (m, 2 H), 4.04 (dd, J=9.08, 7.22 Hz, 1 H) 5.66 (s, 1 H) 6.42 (d, J=2.34 Hz, 1 H) 6.90 (q, J=6.54 Hz, 1 H) 7.73 (s, 1 H) 7.78 (d, J=8.25 Hz, 1 H) 7.91 (d, J=8.35 Hz, 1 H) 7.98 (d, J=2.34 Hz, 1 H)

10h	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.58 (t, J=5.25 Hz, 4 H) 2.05 (dd, J=13.45, 7.15 Hz, 1 H) 2.30 (dd, J=1.00 Hz, 1 H) 2.37 (s, 3 H) 2.40 (s, 3 H) 3.05 - 3.17 (m, 1 H) 3.21 - 3.29 (m, 1 H) 3.36 - 3.75 (m, 4 H) 4.09 (dd, J=9.10, 7.25 Hz, 1 H) 5.73 (s, 1 H) 6.37 (d, J=2.25 Hz, 1 H) 6.71 (d, J=6.69 Hz, 1 H) 7.23 (d, J=0.68 Hz, 1 H) 7.31 (d, J=8.10 Hz, 1 H) 7.60 (d, J=8.05 Hz, 1 H) 7.84 (d, J=2.29 Hz, 1 H)
10i	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.52 - 1.91 (m, 4 H) 2.05 - 2.16 (m, 1 H) 2.40 (s, 3 H) 2.45 - 2.69 (m, 1 H) 3.52 - 4.13 (m, 4 H) 4.57 (d, J=17.28 Hz, 1 H) 6.43 (d, J=2.25 Hz, 1 H) 6.88 - 7.09 (m, 1 H) 7.23 - 7.51 (m, 2 H) 7.68 - 7.83 (m, 1 H) 7.92 (d, J=2.29 Hz, 1 H)
10j	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.59 (d, J=4.54 Hz, 4 H) 2.00 - 2.12 (m, 1 H) 2.27 - 2.35 (m, 1 H) 2.38 (s, 3 H) 3.05 - 3.17 (m, 1 H) 3.25 (d, J=11.71 Hz, 1 H) 3.48 (dd, J=1.17, 0.20 Hz, 2 H) 3.66 (d, J=5.52 Hz, 2 H) 3.85 (s, 3 H) 4.08 (dd, J=9.08, 7.27 Hz, 1 H) 5.72 (s, 1 H) 6.38 (d, J=2.29 Hz, 1 H) 6.67 (d, J=6.69 Hz, 1 H) 6.94 (d, J=2.64 Hz, 1 H) 7.06 (dd, J=8.83, 2.59 Hz, 1 H) 7.63 (d, J=8.83 Hz, 1 H) 7.87 (d, J=2.29 Hz, 1 H)
10k	¹ H NMR (400 MHz, CHLOROFORM-d): δ ppm 1.18-1.36 (m, 3 H) 1.43 (t, J=6.74 Hz, 3 H) 1.54-2.29 (m, 6 H) 2.39 (br. s., 3 H) 3.78 (br. s., 4 H) 4.26 (br. s., 2 H) 4.42 (d, J=6.15 Hz, 2 H) 5.53 (br. s., 1 H), 6.36 (s, 1 H) 6.59 (br. s., 1 H) 7.48 (d, J=7.96 Hz, 1 H), 7.61 (br. s. 1 H) 8.16 (d, J=8.05 Hz, 1 H) 8.34 (br. s., 1 H)
10l	¹ H NMR (400 MHz, DICHLOROMETHANE-d ₂): δ ppm 1.40 - 1.61 (m, 4 H) 1.95 (dd, J=12.89, 5.86 Hz, 1 H) 2.14 - 2.28 (m, 1 H) 2.36 (s, 3 H) 3.07 (d, J=1.00 Hz, 1 H) 3.16 (d, J=1.00 Hz, 1 H) 3.36 (br. s., 2 H), 3.54 (br. s., 2 H) 3.79 (s, 3 H) 4.08 (t, J=7.71 Hz, 1 H) 4.71 - 5.04 (m, 2 H) 5.47 (s, 1 H) 6.30 (d, J=2.10 Hz, 1 H) 6.65 (q, J=7.11 Hz, 1 H) 6.87 (d, J=2.64 Hz, 1 H) 6.95 (dd, J=8.86, 2.61 Hz, 1 H), 7.61 (d, J=8.74 Hz, 1 H) 7.65 (d, J=2.20 Hz, 1 H)
10m	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.64 (d, J=4.69 Hz, 4 H) 2.03 - 2.15 (m, 1 H) 2.40 (s, 4 H) 3.12 - 3.31 (m, 2 H) 3.43 - 3.63 (m, 2 H) 3.64 - 3.78 (m, 2 H) 4.16 - 4.34 (m, 1 H) 6.43 (d, J=2.34 Hz, 1 H) 6.76 - 6.91 (m, 1 H) 7.67 (dd, J=5.76, 4.20 Hz, 3 H) 7.94 (d, J=2.15 Hz, 1 H)
10n	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.47 - 1.71 (m, 4 H) 1.90 (dd, J=13.15, 9.15 Hz, 1 H) 2.24 - 2.39 (m, 4 H) 3.13 (t, J=5.25 Hz, 2 H) 3.66 (br. s., 4 H) 4.39 - 4.51 (m, 2 H) 6.05 (s, 1 H) 6.42 (d, J=2.34 Hz, 1 H) 7.25 (d, J=5.27 Hz, 1 H) 7.51 (d, J=8.59 Hz, 1 H) 7.78 (s, 1 H) 7.85 (dd, J=8.54, 2.29 Hz, 1 H) 8.11 (d, J=2.34 Hz, 1 H) 8.95 (d, J=6.69 Hz, 1 H) 10.20 (br. s., 1 H)

Example 10o: (S)-8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




The title compound was prepared as described for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 10d) starting with (R)-1-(5-bromo-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethanol (Intermediate 38).

5

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.29 (d, J = 7.7 Hz, 2H), 1.61 (q, J = 6.5, 5.3 Hz, 4H), 2.06 (dd, J = 13.5, 7.4 Hz, 1H), 2.36 (dd, J = 13.5, 9.1 Hz, 1H), 3.15 (d, J = 11.9 Hz, 1H), 3.26 (d, J = 11.7 Hz, 1H), 3.47 (ddt, J = 21.7, 13.4, 5.8 Hz, 2H), 3.63 (m, 2H), 4.18 (t, J = 8.2 Hz, 1H), 6.63 (q, J = 6.8 Hz, 1H), 7.50 (m, 7H). LCMS (MH⁺): 607.

10

Example 10p: (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-(trifluoromethyl)-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




15 The title compound was prepared as described for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 10d) starting with (R)-1-(4-chloro-2-(3-(trifluoromethyl)-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol (Intermediate 39).

20 ¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.53 (d, J=5.08 Hz, 4 H) 1.77 - 1.87 (m, 1 H) 2.03 - 2.20 (m, 1 H) 2.75 (s, 1 H) 2.99 (s, 1 H) 3.37 -3.53 (m, 2 H) 3.54 - 3.66 (m, 2 H) 3.66 - 3.77 (m, 1 H) 5.56 (s, 1 H) 6.53 - 6.70 (m, 1 H) 6.96 (d, J=2.34 Hz, 1 H) 7.62 (dd, J=4.30, 2.34 Hz, 2 H), 7.76 (s, 1 H) 8.25 (d, J=1.37 Hz, 1 H). LCMS (MH⁺): 620.

25


Example 10pa: (S)-8-(2-amino-6-((R)-1-(2-(3-(tert-butyl)-1H-pyrazol-1-yl)-4-chlorophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 10d) starting with (R)-1-(2-(3-(tert-butyl)-1H-pyrazol-1-yl)-4-chlorophenyl)-2,2,2-trifluoroethanol (Intermediate 40).

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.40 (s, 9 H) 1.51 - 1.68 (m, 4 H) 1.99 - 2.12 (m, 1 H) 2.25 - 2.41 (m, 1 H) 3.05 - 3.16 (m, 1 H) 3.20 - 3.28 (m, 1 H) 3.38 - 3.55 (m, 2 H) 3.56 - 3.73 (m, 2 H) 4.00 - 4.16 (m, 1 H) 5.57 (s, 1 H) 6.52 (d, J=2.34 Hz, 1 H) 7.15 - 7.28 (m, 1 H) 7.44 - 7.53 (m, 1 H) 7.56 (d, J=1.95 Hz, 1 H) 7.68 - 7.79 (m, 1 H) 7.95 (d, J=2.34 Hz, 1 H). LCMS (MH⁺): 609.

Example 10q: (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-isopropyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




The title compound was prepared as described for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]

decane-3-carboxylic acid (Example 10d) starting with (R)-1-(4-chloro-2-(3-isopropyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol (Intermediate 41).

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.36 (dd, J=6.93, 1.07 Hz, 6 H) 1.57 (br. s., 4 H) 1.86 - 2.03 (m, 1 H) 2.15 - 2.30 (m, 1 H) 2.86 - 3.00 (m, 1 H) 3.02 - 3.19 (m, 2 H) 3.39 - 3.55 (m, 2 H) 3.57 - 3.73 (m, 2 H) 3.82 - 3.98 (m, 1 H) 5.63 (s, 1 H) 6.40 - 6.56 (m, 1 H) 6.93 - 7.10 (m, 1 H) 7.54 (s, 2 H) 7.67 - 7.78 (m, 1 H) 7.91 - 8.02 (m, 1 H). LCMS (MH⁺): 595.


Example 10r: (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-cyclopropyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 10d) starting with (R)-1-(4-chloro-2-(3-cyclopropyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol (Intermediate 42).


¹H NMR (400 MHz, MeOH-d₄): δ ppm 0.77 - 0.90 (m, 2 H) 0.95 - 1.08 (m, 2 H) 1.49 - 1.65 (m, 4 H) 1.80 - 1.95 (m, 1 H) 1.99 - 2.10 (m, 1 H) 2.10 - 2.24 (m, 1 H) 2.74 - 2.85 (m, 1 H) 3.00 - 3.11 (m, 1 H) 3.38 - 3.69 (m, 4 H) 3.72 - 3.84 (m, 1 H) 5.56 - 5.70 (m, 1 H) 6.29 - 6.38 (m, 1 H) 6.89 - 7.05 (m, 1 H) 7.52 (s, 2 H) 7.67 - 7.77 (m, 1 H) 7.86 - 7.98 (m, 1 H). LCMS (MH⁺): 593.

Example 11: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(6-methyl-2-(3-methyl-1H-pyrazol-1-yl)pyridin-3-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 10d) starting with (S)-2,2,2-trifluoro-1-(6-methyl-2-(3-methyl-1H-pyrazol-1-yl)pyridin-3-yl)ethanol (Intermediate 20)

Example 12a: (S)-8-(2-amino-6-((R)-1-(4-ethyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




Step 1: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (300 mg, 0.388 mmol, see Example 1u) in EtOH:H₂O (15 mL) was added 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (90 mg, 0.58 mmol), KHCO₃ (389 mg, 3.88 mmol), and PdCl₂(PPh₃)₂ (41 mg, 0.058 mmol). The reaction mixture was heated to 80 °C for 1 h, then cooled to RT. The reaction was diluted with water, extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification with a 40 g Isco RediSep silica cartridge (EtOAc:heptane) provided (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-vinylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

Step 2: N-CBZ Deprotection was accomplished via method A, which also reduced the olefin, to provide (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-ethyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

5

Step 3: Hydrolysis of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-ethyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate using the LiOH general method provided the title compound as a white solid.


10 Using the same scheme below, the following examples of Table 4a were prepared as described above for (S)-8-(2-amino-6-((R)-1-(4-ethyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 12a).



15

20

Table 4a.




Ex. No.	R	CAS Name	LCMS (MH+)
12a		(S)-8-(2-amino-6-((R)-1-(4-ethyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	560
12b		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-propylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	575
12c		(S)-8-(2-amino-6-((R)-1-(4-butyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	588

Table 4b.

NMR Data for Compounds of Table 4a

Ex. No.	NMR
12a	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.59 Hz, 3 H), 1.50 - 1.69 (m, 4 H), 2.01 - 2.35 (m, 2 H), 2.37 (s, 3 H), 2.72 (q, J = 7.57 Hz, 2 H), 3.05 - 3.28 (m, 2 H), 3.40 - 3.76 (m, 4 H), 4.08 (dd, J = 8.88, 7.32 Hz, 1 H), 5.72 (s, 1 H), 6.38 (d, J = 2.25 Hz, 1 H), 6.71 (q, J = 6.70 Hz, 1 H), 7.25 (d, J = 1.56 Hz, 1 H), 7.35 (dd, J = 8.18, 1.59 Hz, 1 H), 7.63 (d, J = 8.15 Hz, 1 H), 7.85 (d, J = 2.29 Hz, 1 H)
12b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.96 (t, J = 7.35 Hz, 3 H), 1.49 - 1.62 (m, 4 H), 1.62 - 1.77 (m, 2 H), 2.01 - 2.35 (m, 2 H), 2.37 (s, 3 H), 2.59 - 2.74 (m, 2 H), 3.06 - 3.29 (m, 2 H), 3.39 - 3.77 (m, 4 H), 4.08 (dd, J = 9.05, 7.30 Hz, 1 H), 5.72 (s, 1 H), 6.37 (d, J = 2.29 Hz, 1 H), 6.71 (q, J = 6.72 Hz, 1 H), 7.23 (d, J = 1.56 Hz, 1 H), 7.33 (dd, J = 8.15, 1.56 Hz, 1 H), 7.63 (d, J = 8.05 Hz, 1 H), 7.85 (d, J = 2.29 Hz, 1 H)
12c	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.94 (t, J = 7.35 Hz, 3 H), 1.38 (dq, J = 14.92, 7.39 Hz, 2 H), 1.49 - 1.72 (m, 6 H), 2.01 - 2.35 (m, 2 H), 2.37 (s, 3 H), 2.60 - 2.74 (m, 2 H), 3.07 - 3.28 (m, 2 H), 3.40 - 3.74 (m, 4 H), 4.08 (dd, J = 9.15, 7.20 Hz, 1 H), 5.71 (s, 1 H), 6.38 (d, J = 2.15 Hz, 1 H), 6.63 - 6.77 (m, 1 H), 7.23 (d, J = 1.61 Hz, 1 H), 7.33 (dd, J = 8.10, 1.66 Hz, 1 H), 7.63 (d, J = 8.05 Hz, 1 H), 7.85 (d, J = 2.29 Hz, 1 H)

Example 13: (3S)-8-(2-amino-6-((1R)-1-(4-(1,2-dihydroxyethyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



- 5 *Step 1:* To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-vinylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (product of Step 1, Example 12b) (373 mg, 0.518 mmol) in 4:1 acetone:H₂O (20 mL) was added OsO₄ (313 µL of a 4% (w/w) aqueous solution, 325 mg, 0.0518 mmol) and N-methylmorpholine-N-oxide (214 µL of a 50% (w/w) aqueous solution, 242 mg, 1.04 mmol).
- 10 The reaction was stirred at RT for 24 h, concentrated *in vacuo*, and the residue was purified by chromatography on a 50 g Isco Gold RediSep reversed phase silica cartridge (H₂O:HOAc : 99:1 to EtOH:HOAc 99:1). A second purification on a 40 g Isco RediSep silica cartridge eluting (CH₂Cl₂ 100% to 90:9:1 CH₂Cl₂:EtOH:NH₄OH) provided (3S)-2-benzyl 3-ethyl 8-(2-amino-6-((1R)-1-(4-(1,2-dihydroxyethyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-
- 15 trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.


Step 2: N-CBZ deprotection was accomplished via method A to provide (3S)-ethyl 8-(2-amino-6-((1R)-1-(4-(1,2-dihydroxyethyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid.

- 20 *Step 3:* Hydrolysis of (3S)-ethyl 8-(2-amino-6-((1R)-1-(4-(1,2-dihydroxyethyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provides the title compound as a white solid.

- 25 ¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.49 - 1.66 (m, 4 H) 2.05 (dd, J=13.50, 7.20 Hz, 1 H) 2.31 (dd, J=13.45, 9.20 Hz, 1 H) 2.38 (s, 3 H) 3.04 - 3.28 (m, 2 H) 3.38 - 3.76 (m, 6 H) 4.08 (dd,

J=8.98, 7.27 Hz, 1 H) 4.67 - 4.79 (m, 1 H) 5.72 (d, J=2.15 Hz, 1 H) 6.39 (d, J=2.29 Hz, 1 H) 6.77 (q, J=6.65 Hz, 1 H) 7.45 (s, 1 H) 7.52 (d, J=8.20 Hz, 1 H) 7.71 (d, J=8.15 Hz, 1 H) 7.88 (dd, J=4.20, 2.34 Hz, 1 H). LCMS (MH⁺): 592.

5 **Example 14: (S)-8-(2-amino-6-((R)-1-(4-cyano-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid**



10 *Step 1:* To a solution of (3S)-2-benzyl 3-ethyl 8-(2-amino-6-(1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (730 mg, 1.0 mmol), was added ZnCN₂ (280 mg, 2.4 mmol), Zn (64 mg, 1.0 mmol), DMA (10 mL), and Pd(P-t-Bu₃)₂ (78 mg, 0.15 mmol). The reaction mixture was heated in a sealed vial at 115 °C for 2 h, then cooled to RT, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptate) provided (3S)-2-benzyl 3-ethyl
15 8-(2-amino-6-(1-(4-cyano-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a viscous oil.

Step 2: N-CBZ Deprotection was accomplished via Method A to provide (3S)-ethyl 8-(2-amino-6-(1-(4-cyano-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-
20 diazaspiro[4.5]decane-3-carboxylate as an off-white solid.


Step 3: Hydrolysis of (3S)-ethyl 8-(2-amino-6-(1-(4-cyano-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provides the title compound as an off-white solid.

25

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.47 - 1.71 (m, 4 H) 1.95 - 2.10 (m, 1 H) 2.20 - 2.33 (m, 1 H) 2.36 (s, 3 H) 2.96 - 3.24 (m, 2 H) 3.35 - 3.54 (m, 2 H) 3.55 - 3.79 (m, 2 H) 3.92 - 4.13 (m, 1 H) 5.65 (s, 1 H) 6.42 (d, J=2.15 Hz, 1 H) 6.95 (q, J=6.72 Hz, 1 H) 7.70 - 7.91 (m, 3 H) 7.97 (d, J=2.25 Hz, 1 H). LCMS (MH⁺): 556.

5

Example 15: (S)-8-(2-amino-6-((R)-1-(4-carbamoyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



10 *Step 1:* To a solution of (3S)-2-benzyl 3-ethyl 8-(2-amino-6-(1-(4-cyano-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (150 mg, 0.2 mmol, see Ex. 14) in toluene (10 mL) was added acetaldehyde oxime (240 mg, 4 mmol) and InCl₃ (44 mg, 0.2 mmol). The reaction was heated to 110 °C for 3 h, then cooled to RT, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/hepate) provided (3S)-2-benzyl 3-ethyl 8-(2-amino-6-(1-(4-carbamoyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

20 *Step 2:* N-CBZ Deprotection was accomplished via Method A to provide (3S)-ethyl 8-(2-amino-6-(1-(4-carbamoyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid.


Step 3: Hydrolysis of (3S)-ethyl 8-(2-amino-6-(1-(4-carbamoyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provides the title compound as a white solid.

25

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.56 (t, J=4.98 Hz, 5 H) 2.03 (dd, J=13.47, 7.03 Hz, 1 H) 2.23 - 2.33 (m, 2 H) 2.35 - 2.39 (m, 3 H) 3.04 - 3.12 (m, 1 H) 3.22 (d, J=11.71 Hz, 1 H) 3.37 - 3.72 (m, 5 H) 4.05 (dd, J=9.20, 7.05 Hz, 1 H) 5.70 (s, 1 H) 6.40 (d, J=2.39 Hz, 1 H) 6.82 - 6.92 (m, 1 H) 7.80 (d, J=8.10 Hz, 1 H) 7.87 - 7.97 (m, 4 H). LCMS (MH⁺): 575.

5

Example 16: (S)-8-(2-amino-6-((R)-1-(4-carboxy-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a solution of (3S)-2-benzyl 3-ethyl 8-(2-amino-6-(1-(4-cyano-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (0.35 g, 0.50 mmol, see Ex. 14) in MeOH (5 mL) and water (1 mL) was added LiOH-H₂O (0.20 g, 5 mmol). The mixture was heated to 50 °C overnight. The reaction was then cooled to RT, and the reaction was acidified with 6N HCl to pH=1. Concentration *in vacuo* followed by reverse phase HPLC purification (MeOH/water/HOAc) provided (3S)-8-(2-amino-6-(1-(4-carboxy-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid as a white solid.


Step 2: N-CBZ Deprotection was accomplished via Method A to provide the title compound as a white solid.

20

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.57 (t, J=5.42 Hz, 4 H) 2.03 (dd, J=13.42, 7.42 Hz, 1 H) 2.25 - 2.35 (m, 2 H) 2.37 (s, 2 H) 3.04 - 3.13 (m, 1 H) 3.16 - 3.25 (m, 1 H) 3.38 - 3.75 (m, 5 H) 4.06 (dd, J=9.03, 7.32 Hz, 1 H) 5.72 (s, 1 H) 6.39 (d, J=2.29 Hz, 1 H) 6.78 - 6.89 (m, 1 H) 7.76 (d, J=8.15 Hz, 1 H) 7.90 (d, J=2.34 Hz, 1 H) 7.95 (d, J=1.42 Hz, 1 H) 8.04 (dd, J=8.13, 1.59 Hz, 1 H). LCMS (MH⁺): 576.

25

Example 17: (S)-8-(2-amino-6-((R)-1-(4-(ethoxycarbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



5 *Step 1:* To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (1.50 g, 1.94 mmol, See Ex. 1u) in THF (20 mL), MeOH (10 mL) and water (10 mL) was added LiOH-H₂O (0.80 g, 19.4 mmol), and the reaction was stirred at RT for 4 h. The pH of the reaction mixture was adjusted to 6.5 with 6 N HCl, and the organic solvents were removed *in vacuo* to provide a white solid that is filtered away. The reaction mixture was then partitioned between water and EtOAc, and extracted. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, then concentrated *in vacuo* to provide (2S)-8-[2-amino-6-[(1R)-1-[4-bromo-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoro-ethoxy]pyrimidin-4-yl]-3-benzyloxycarbonyl-3,8-diazaspiro[4.5]decane-2-carboxylic acid as a white solid that is used directly without further purification.


Step 2: To a solution of (2S)-8-[2-amino-6-[(1R)-1-[4-bromo-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoro-ethoxy]pyrimidin-4-yl]-3-benzyloxycarbonyl-3,8-diazaspiro[4.5]decane-2-carboxylic acid (74 mg, 0.10 mmol, Step 2) in EtOH (4 mL) was added KHCO₃ (84 mg, 1.0 mmol). The reaction mixture was degassed, fitted with a 1 atm CO balloon, then treated with PdCl₂(PPh₃)₂ (14 mg, 0.02 mmol). The reaction was degassed once more with 1 atm CO and then heated to 80 °C for 12 h. The reaction was cooled to RT, concentrated *in vacuo* and the residue was partitioned between water and EtOAc, and extracted. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (CH₂Cl₂/AcOH/EtOH) provided (2S)-8-[2-amino-6-[(1R)-1-[4-

ethoxycarbonyl-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoro-ethoxy] pyrimidin-4-yl]-3-benzyloxycarbonyl-3,8-diazaspiro[4.5]decane-2-carboxylic acid as a white solid.

Step 3: N-CBZ Deprotection was accomplished via Method A to provide the title compound as a white solid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.37 (t, J=7.13 Hz, 3 H) 1.58 (d, J=4.30 Hz, 4 H) 1.97 (s, 2 H) 2.04 (dd, J=13.47, 7.27 Hz, 1 H) 2.30 (dd, J=13.59, 9.25 Hz, 1 H) 2.38 (s, 3 H) 3.05 - 3.27 (m, 2 H) 3.39 - 3.76 (m, 4 H) 3.99 - 4.10 (m, 1 H) 4.37 (q, J=7.13 Hz, 2 H) 5.68 (s, 1 H) 6.41 (d, J=2.34 Hz, 1 H) 6.84 (q, J=6.67 Hz, 1 H) 7.83 (d, J=8.10 Hz, 1 H) 7.94 (d, J=2.34 Hz, 1 H) 7.99 (d, J=1.61 Hz, 1 H) 8.09 (dd, J=8.27, 1.68 Hz, 1 H). LCMS (MH⁺): 604.

Example 18a: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(((1,1,1,3,3,3-hexafluoro-2-methylpropan-2-yl)oxy)carbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a solution of (S)-8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (product of Step 3, Example 10m) (1.2 g, 1.6 mmol) in DMF (16 mL) was added benzyl bromide (0.27 g, 1.6 mmol) and NaHCO₃ (0.67 g, 8.0 mmol). The reaction was then heated to 60 °C for 2 h, cooled to RT, and stirred for 12 h. The precipitate was filtered, washed with EtOAc and the filtrate concentrated *in vacuo*. The residue was partitioned between water and EtOAc, and extracted. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc /heptane) provided (S)-dibenzyl 8-(2-amino-6-((R)-1-(4-bromo-2-(3-

methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as the white solid.

Step 2: To a solution of (S)-dibenzyl 8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate from Step 1 (415 mg, 0.50 mmol) in 1,4-dioxane (8 mL) and water (4 mL) was added KHCO₃ (420 mg, 5.0 mmol), and the reaction was degassed with 1 atm CO. Then PdCl₂(PPh₃)₂ (140 mg, 0.10 mmol) was added and the reaction mixture was treated with 1 atm CO (balloon). The reaction mixture was heated to 80 °C for 12 h, then cooled to RT, and concentrated *in vacuo*.

The residue was partitioned between water and EtOAc, and extracted. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*.


Purification by normal phase silica gel column (CH₂Cl₂/MeOH/NH₄OH) provided

4-[(1R)-1-[2-amino-6-[(2S)-2,3-bis(benzyloxycarbonyl)-3,8-diazaspiro[4.5]decan-8-yl]pyrimidin-4-yl]oxy-2,2,2-trifluoro-ethyl]-3-(3-methylpyrazol-1-yl)benzoic acid as a white solid.

Step 3: To a solution of 4-[(1R)-1-[2-amino-6-[(2S)-2,3-bis(benzyloxycarbonyl)-3,8-diazaspiro[4.5]decan-8-yl]pyrimidin-4-yl]oxy-2,2,2-trifluoro-ethyl]-3-(3-methylpyrazol-1-yl)benzoic acid (80 mg, 0.1 mmol) in CH₂Cl₂ (4 mL) was added DMAP (73 mg, 0.6 mmol), (CF₃)₂MeCOH (108 mg, 0.6 mmol), followed by EDCI (114 mg, 0.6 mmol). The reaction mixture was stirred at RT for 12 h, diluted with CH₂Cl₂ and washed with water. The aqueous solution was extracted with CH₂Cl₂. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc / heptane) provided dibenzyl (2S)-8-[2-amino-6-[(1R)-2,2,2-trifluoro-1-[2-(3-methylpyrazol-1-yl)-4-[2,2,2-trifluoro-1-methyl-1-(trifluoromethyl)ethoxy]carbonyl-phenyl]ethoxy]pyrimidin-4-yl]-3,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.


Step 4: N-CBZ Deprotection was accomplished via Method A to provide the title compound as a white solid.

Using the generic scheme below, the following examples of Table 5a were prepared as described above for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(((1,1,1,3,3,3-hexafluoro-2-methylpropan-2-yl)oxy)carbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 18a).



5

Table 5a.









Ex. No.	R	CAS Name	LCMS (MH+)
18a		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(((1,1,1,3,3,3-hexafluoro-2-methylpropan-2-yl)oxy)carbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	740
18b		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(propoxycarbonyl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	618
18c		(S)-8-(2-amino-6-((R)-1-(4-(butoxycarbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	632
18d		(S)-8-(2-amino-6-((R)-1-(4-(tert-butoxycarbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	632
18e		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(isobutoxycarbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	632
18f		(S)-8-(2-amino-6-((R)-1-(4-((cyclopentyl)oxy)carbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	644


Table 5b.

NMR Data for Compounds of Table 5a

Ex. No.	NMR
18a	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.58 (br. s., 4 H) 1.97 (s, 1 H) 2.04 (dd, J=13.50, 7.20 Hz, 1 H) 2.12 (s, 3 H) 2.31 (dd, J=13.45, 9.30 Hz, 1 H) 2.38 (s, 3 H) 3.04 - 3.27 (m, 2 H) 3.38 - 3.55 (m, 2 H) 3.64 (dd, J=13.23, 5.56 Hz, 2 H) 4.07 (t, J=8.08 Hz, 1 H) 5.67 (s, 1 H) 6.43 (d, J=2.34 Hz, 1 H) 6.85 (q, J=6.69 Hz, 1 H) 7.90 (d, J=8.20 Hz, 1 H) 7.96 (dd, J=8.20, 2.00 Hz, 2 H) 8.06 (dd, J=8.27, 1.73 Hz, 1 H)
18b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.01 (t, J=7.44 Hz, 3 H) 1.58 (d, J=4.49 Hz, 4 H) 1.72 - 1.85 (m, 2 H) 1.97 (s, 1 H) 2.04 (dd, J=13.35, 7.25 Hz, 1 H) 2.30 (dd, J=13.52, 9.13 Hz, 1 H) 2.38 (s, 3 H), 3.06 - 3.26 (m, 2 H) 3.38 - 3.72 (m, 4 H) 4.00 - 4.12 (m, 1 H) 4.29 (t, J=6.64 Hz, 2 H) 5.68 (s, 1 H) 6.42 (d, J=2.44 Hz, 1 H) 6.84 (q, J=6.57 Hz, 1 H) 7.84 (d, J=8.20 Hz, 1 H) 7.95 (d, J=2.34 Hz, 1H) 7.98 (d, J=1.61 Hz, 1 H) 8.09 (dd, J=8.22, 1.64 Hz, 1 H)
18c	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.97 (t, J=7.42 Hz, 3 H) 1.46 (dq, J=15.01, 7.48

	Hz, 2 H) 1.58 (d, J=4.83 Hz, 4 H) 1.68 - 1.82 (m, 2 H) 1.97 (s, 1 H) 2.04 (dd, J=13.52, 7.03 Hz, 1 H) 2.30 (dd, J=13.42, 9.18 Hz, 1 H) 2.38 (s, 3 H) 3.07 - 3.25 (m, 2 H) 3.38 - 3.71 (m, 4 H) 4.06 (dd, J=9.15, 7.00 Hz, 1 H) 4.33 (t, J=6.61 Hz, 2 H) 5.68 (s, 1 H) 6.42 (d, J=2.39 Hz, 1 H) 6.84 (q, J=6.44 Hz, 1 H) 7.84 (d, J=8.30 Hz, 1 H) 7.95 (d, J=2.29 Hz, 1 H) 7.98 (d, J=1.61 Hz, 1 H) 8.08 (dd, J=8.25, 1.71 Hz, 1 H)
18d	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.57 (s, 13 H) 1.97 (s, 2 H) 2.04 (dd, J=13.50, 7.15 Hz, 1 H) 2.30 (dd, J=14.06, 9.96 Hz, 1 H) 2.38 (s, 3 H) 3.08 - 3.26 (m, 2 H) 3.38 - 3.74 (m, 4 H) 4.01 - 4.14 (m, 1 H) 5.68 (s, 1 H) 6.41 (d, J=2.34 Hz, 1 H) 6.80 (q, J=6.64 Hz, 1 H) 7.80 (d, J=8.15 Hz, 1 H) 7.92 (dd, J=7.88, 1.93 Hz, 2 H) 8.02 (dd, J=8.27, 1.59 Hz, 1 H)
18e	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.00 (d, J=6.74 Hz, 6 H) 1.52 - 1.64 (m, 4 H) 1.97 (s, 2 H) 2.00 - 2.12 (m, 2 H) 2.30 (dd, J=13.45, 9.35 Hz, 1 H) 2.38 (s, 3 H) 3.07 - 3.26 (m, 2 H) 3.37 - 3.55 (m, 2 H) 3.58 - 3.70 (m, 2 H) 4.06 (dd, J=9.03, 7.17 Hz, 1 H) 4.12 (d, J=6.59 Hz, 2 H) 5.68 (s, 1 H) 6.42 (d, J=2.39 Hz, 1 H) 6.84 (q, J=6.51 Hz, 1 H) 7.84 (d, J=8.35 Hz, 1 H) 7.95 (d, J=2.34 Hz, 1 H) 7.98 (d, J=1.61 Hz, 1 H) 8.09 (dd, J=8.27, 1.68 Hz, 1 H)
18f	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.54 - 1.94 (m, 11 H) 1.97 (s, 3 H) 2.04 (dd, J=13.35, 7.15 Hz, 1 H) 2.24 - 2.35 (m, 1 H) 2.38 (s, 3 H) 3.02 - 3.27 (m, 2 H) 3.37 - 3.81 (m, 4 H) 3.95 - 4.22 (m, 1H) 5.32 - 5.44 (m, 1 H) 5.67 (s, 1 H) 6.41 (d, J=2.39 Hz, 1 H) 6.82 (d, J=6.39 Hz, 1 H) 7.82 (d, J=8.30 Hz, 1 H) 7.94 (d, J=1.85 Hz, 2 H) 8.06 (dd, J=8.15, 1.71 Hz, 1 H)

Example 19a: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-5-vinylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



5

Step 1: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(5-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (500 mg, 0.65 mmol) in 4:1 EtOH:H₂O (25 mL) was added 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (150 mg, 0.971 mmol), KHCO₃ (648 mg, 6.47 mmol), and PdCl₂(PPh₃)₂ (68 mg, 0.097 mmol). The reaction mixture was heated to 80 °C for 1.75 h, then cooled to RT, and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification via a 40 g Isco RediSep

10

silica cartridge eluting (EtOAc/hepate) provides (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-5-vinylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as an off-white solid.

- 5 *Step 2:* N-CBZ Deprotection was accomplished via Method B to provide (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-5-vinylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as an off-white solid.

- Step 3:* Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-5-vinylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provided the title compound as a white solid.
- 10

- Using the generic scheme below, the following examples of Table 6a were prepared as described above for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-5-vinylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 19a).
- 15





Table 6a.

Ex. No.	R	CAS Name	LCMS (MH+)
19a		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-5-vinylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	558.6
19b		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-5-((E)-prop-1-en-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	572.6
19c		(S)-8-(2-amino-6-((R)-1-(5-((E)-but-1-en-1-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	585.5
19d		(S)-8-(2-amino-6-((R)-1-(5-((E)-2-carboxyvinyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	602.6

Table 6b.**NMR Data for Compounds of Table 6a**

Ex. No.	NMR
19a	¹ H NMR (400 MHz, MeOH- d ₄): δ ppm 1.59 (m, 4 H) 2.06 (dd, J=13.42, 7.17 Hz, 1 H) 2.31 (dd, J=13.42, 9.18 Hz, 1 H) 2.38 (s, 3 H) 3.18 (m, 2 H) 3.59 (m, 4 H) 4.07 (dd, J=9.20, 7.20 Hz, 1 H) 5.36 (d, J=10.98 Hz, 1 H) 5.75 (s, 1 H) 5.85 (d, J=17.62 Hz, 1 H) 6.39 (d, J=2.34 Hz, 1 H) 6.80 (m, 2 H) 7.38 (d, J=8.30 Hz, 1 H) 7.63 (dd, J=8.25, 2.00 Hz, 1 H) 7.74 (s, 1 H) 7.87 (d, J=2.29 Hz, 1 H)
19b	¹ H NMR (400 MHz, MeOH- d ₄): δ ppm 1.59 (m, 3 H) 1.90 (dd, J=6.32, 1.20 Hz, 3 H) 2.06 (dd, J=13.47, 7.13 Hz, 1 H) 2.31 (dd, J=13.45, 9.25 Hz, 1 H) 2.37 (s, 3 H) 3.18 (m, 2 H) 3.57 (m, 4 H) 4.08 (dd, J=9.18, 7.17 Hz, 1 H) 5.75 (s, 1 H) 6.39 (m, 3 H) 6.75 (q, J=6.67 Hz, 1 H) 7.32 (d, J=8.25 Hz, 1 H) 7.52 (dd, J=8.30, 2.00 Hz, 1 H) 7.65 (s, 1 H) 7.84 (d, J=2.34 Hz, 1 H)
19c	¹ H NMR (400 MHz, MeOH- d ₄): δ ppm 1.11 (t, J=7.47 Hz, 3 H) 1.59 (d, J=4.59 Hz, 4 H) 2.06 (dd, J=13.37, 7.22 Hz, 1 H) 2.28 (m, 3 H) 2.37 (s, 3 H) 3.18 (m, 2 H) 3.59 (m, 4 H)

	4.07 (dd, J=9.10, 7.20 Hz, 1 H) 5.76 (s, 1 H) 6.40 (m, 3 H) 6.76 (m, 1 H) 7.33 (d, J=8.25 Hz, 1 H) 7.54 (dd, J=8.30, 2.05 Hz, 1 H) 7.66 (s, 1 H) 7.84 (d, J=2.29 Hz, 1 H)
19d	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.57 (t, J=5.44 Hz, 4 H) 1.97 (s, 3 H) 2.04 (dd, J=13.72, 7.27 Hz, 1 H) 2.30 (dd, J=13.32, 9.18 Hz, 1H) 2.37 (s, 3 H) 3.07 - 3.25 (m, 2 H) 3.40 - 3.55 (m, 2 H) 3.65 (dd, J=9.27, 4.73 Hz, 2 H) 4.07 (t, J=7.98 Hz, 1 H) 5.75 (s, 1 H) 6.40 (d, J=2.34 Hz, 1 H) 6.51 (d, J=16.20 Hz, 1 H) 6.94 (q, J=6.52 Hz, 1 H) 7.46 (d, J=8.30 Hz, 1 H) 7.66 (d, J=15.86 Hz, 1 H) 7.78 (dd, J=8.32, 1.88 Hz, 1 H) 7.87 (s, 1 H) 7.92 (d, J=2.34 Hz, 1 H)

Using the generic scheme below, the following examples of Table 7a can be prepared as described above for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-5-vinylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 19a),

5 by substituting the alkylidene borolane with a boronic acid or ester.













Table 7a.



Ex. No.	Cy	CAS Name	LCMS (MH ⁺)
---------	----	----------	-------------------------

19e		(S)-8-(2-amino-6-((R)-1-(3',4'-dimethyl-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	536.7
19f		(S)-8-(2-amino-6-((R)-1-(3'-carboxy-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	652
19g		(S)-8-(2-amino-6-((R)-1-(4'-carboxy-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	652
19h		(S)-8-(2-amino-6-((R)-1-(3'-((E)-2-carboxyvinyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	678
19i		(S)-8-(2-amino-6-((R)-1-(4'-((E)-2-carboxyvinyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	678
19j		(S)-8-(2-amino-6-((R)-1-(3'-(2-carboxyethyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	680
19k		(S)-8-(2-amino-6-((R)-1-(4'-(2-carboxyethyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	680
19l		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(hydroxymethyl)-3'-methyl-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	652







19m		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(hydroxymethyl)-4'-methyl-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	652
19n		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	608
19o		(S)-8-(2-amino-6-((R)-1-(3',4'-difluoro-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	644
19p		(S)-8-(2-amino-6-((R)-1-(3',4'-dichloro-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	677
19q		(S)-8-(2-amino-6-((R)-1-(4'-chloro-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	643
19r		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(hydroxymethyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	639

Table 7b.


NMR Data for Compounds of Table 7a

Ex. No.	NMR
19e	¹ H NMR (400 MHz, MeOH- d ₄): δ ppm 1.57 (m, 4 H) 2.04 (dd, J=13.62, 6.98 Hz, 1 H) 2.32 (d, J=11.96 Hz, 6 H) 2.40 (s, 3 H) 3.16 (m, 2 H) 3.55 (m, 4 H) 4.07 (dd, J=9.18, 7.22 Hz, 1 H) 5.79 (s, 1 H) 6.40 (d, J=2.29 Hz, 1 H) 6.85 (m, 1 H) 7.21 (d, J=7.76 Hz, 1 H) 7.31 (m, 1 H) 7.36 (s, 1 H) 7.45 (d, J=8.25 Hz, 1 H) 7.75 (dd, J=8.27, 2.12 Hz, 1 H) 7.90 (d, J=2.20 Hz, 2 H)
19f	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.53 - 1.67 (m, 4 H) 2.05 (dd, J=13.42, 7.17 Hz, 1 H) 2.30 (dd, J=13.42, 9.22 Hz, 1 H) 2.40 (s, 3 H) 3.06 - 3.27 (m, 2 H) 3.39 - 3.74 (m, 4 H) 4.08 (dd, J=9.13, 7.27 Hz, 1 H) 5.79 (s, 1 H) 6.42 (d, J=2.29 Hz, 1 H) 6.92 (q, J=6.62 Hz, 1 H) 7.53 (d, J=8.25 Hz, 1 H) 7.57 (t, J=7.76 Hz, 1 H) 7.77 - 7.87 (m, 2 H) 7.94 (d, J=2.34 Hz, 1 H) 7.97 (d, J=1.42 Hz, 1 H) 8.04 (dt, J=7.79, 1.23 Hz, 1 H) 8.24 (t, J=1.61 Hz, 1 H)
19g	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.47 - 1.67 (m, 4 H) 2.05 (dd, J=13.45, 7.20 Hz, 1 H) 2.31 (dd, J=13.37, 9.27 Hz, 1 H) 2.40 (s, 3 H) 2.99 - 3.28 (m, 2 H) 3.39 - 3.78 (m, 4 H)

	4.08 (dd, J=9.08, 7.27 Hz, 1 H) 5.79 (s, 1 H) 6.42 (d, J=2.29 Hz, 1 H) 6.86 - 7.01 (m, 1 H) 7.53 (d, J=8.30 Hz, 1 H) 7.64 - 7.77 (m, 2 H) 7.85 (dd, J=8.30, 2.15 Hz, 1 H) 7.94 (d, J=2.34 Hz, 1 H) 7.99 (d, J=1.32 Hz, 1 H) 8.08 - 8.18 (m, 2 H)
19h	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.59 (t, J=5.54 Hz, 4 H) 2.04 (dd, J=13.45, 7.39 Hz, 1 H) 2.32 (dd, J=13.50, 9.25 Hz, 1 H) 2.41 (s, 3 H) 3.07 - 3.26 (m, 2 H) 3.41 - 3.76 (m, 4 H) 4.08 (dd, J=9.01, 7.30 Hz, 1 H) 5.81 (s, 1 H) 6.42 (d, J=2.29 Hz, 1 H) 6.57 (d, J=16.01 Hz, 1 H) 6.86 - 6.97 (m, 1 H) 7.48 - 7.57 (m, 2 H) 7.60 - 7.68 (m, 2 H) 7.73 (d, J=16.01 Hz, 1 H) 7.77 (bs, 1 H) 7.83 (dd, J=8.25, 2.10 Hz, 1 H) 7.93 - 7.96 (m, 2 H)
19i	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.50 - 1.65 (m, 4 H) 2.05 (dd, J=13.45, 7.20 Hz, 1 H) 2.31 (dd, J=13.40, 9.30 Hz, 1 H) 2.40 (s, 3 H) 3.05 - 3.28 (m, 2 H) 3.40 - 3.74 (m, 4 H) 4.07 (dd, J=9.10, 7.25 Hz, 1 H) 5.79 (s, 1 H) 6.42 (d, J=2.29 Hz, 1 H) 6.54 (d, J=16.01 Hz, 1 H) 6.91 (q, J=6.72 Hz, 1 H) 7.51 (d, J=8.25 Hz, 1 H) 7.61 - 7.75 (m, 5 H) 7.82 (dd, J=8.30, 2.15 Hz, 1 H) 7.93 (d, J=2.34 Hz, 1 H) 7.97 (s, 1 H)
19j	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.51 - 1.66 (m, 4 H) 2.04 (dd, J=13.50, 7.15 Hz, 1 H) 2.31 (dd, J=13.37, 9.18 Hz, 1 H) 2.40 (s, 3 H) 2.65 (t, J=7.61 Hz, 2 H) 2.99 (t, J=7.59 Hz, 2 H) 3.06 - 3.27 (m, 2 H) 3.40 - 3.78 (m, 4 H) 4.08 (dd, J=8.98, 7.42 Hz, 1 H) 5.80 (s, 1 H) 6.41 (d, J=2.34 Hz, 1 H) 6.88 (q, J=6.61 Hz, 1 H) 7.27 (d, J=7.32 Hz, 1 H) 7.35 - 7.41 (m, 1 H) 7.41 - 7.51 (m, 3 H) 7.77 (dd, J=8.27, 2.12 Hz, 1 H) 7.88 - 7.97 (m, 2 H)
19k	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.57 (d, J=3.37 Hz, 4 H) 2.04 (dd, J=13.40, 7.20 Hz, 1 H) 2.30 (dd, J=13.35, 9.20 Hz, 1 H) 2.40 (s, 3 H) 2.63 (t, J=7.61 Hz, 2 H) 2.96 (t, J=7.57 Hz, 2 H) 3.03 - 3.26 (m, 2 H) 3.39 - 3.76 (m, 4 H) 4.07 (dd, J=9.03, 7.32 Hz, 1 H) 5.78 (s, 1 H) 6.41 (d, J=2.29 Hz, 1 H) 6.86 (q, J=6.54 Hz, 1 H) 7.34 (d, J=8.25 Hz, 2 H) 7.46 (d, J=8.30 Hz, 1 H) 7.52 (d, J=8.25 Hz, 2 H) 7.76 (dd, J=8.27, 2.12 Hz, 1 H) 7.89 - 7.92 (m, 2 H)
19l	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.45 - 1.65 (m, 4 H) 2.00 - 2.09 (m, 1 H) 2.30 (dd, J=13.40, 9.25 Hz, 1 H) 2.40 (s, 6 H) 3.03 - 3.27 (m, 2 H) 3.39 - 3.76 (m, 4 H) 4.07 (dd, J=9.10, 7.25 Hz, 1 H), 4.67 (s, 2 H) 5.79 (s, 1 H) 6.41 (d, J=2.25 Hz, 1 H) 6.86 (q, J=6.64 Hz, 1 H) 7.36 - 7.53 (m, 4 H) 7.77 (dd, J=8.30, 2.15 Hz, 1 H) 7.91 (d, J=2.44 Hz, 2 H)
19m	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.46 - 1.69 (m, 4 H) 2.00 - 2.10 (m, 1 H) 2.30 (dd, J=13.45, 9.25 Hz, 1 H) 2.37 (s, 3 H) 2.40 (s, 3 H) 3.03 - 3.27 (m, 2 H) 3.39 - 3.76 (m, 4 H) 4.07 (dd, J=9.13, 7.22 Hz, 1 H) 4.70 (s, 2 H) 5.78 (s, 1 H) 6.41 (d, J=2.25 Hz, 1 H) 6.85 (q, J=6.57 Hz, 1 H) 7.26 (d, J=7.91 Hz, 1 H) 7.43 (dd, J=7.81, 1.95 Hz, 1 H) 7.47 (d, J=8.30 Hz, 1 H) 7.64 (d, J=1.81 Hz, 1 H) 7.79 (dd, J=8.27, 2.12 Hz, 1 H) 7.91 (d, J=2.29 Hz, 1 H) 7.94 (s, 1 H)
19n	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 7.2 Hz, 1H), 1.56 (d, J = 6.3 Hz, 4H), 2.03 (d, J = 12.8 Hz, 1H), 2.30 (d, J = 12.4 Hz, 1H), 2.39 (s, 3H), 3.09 (d, J = 11.5 Hz, 1H), 3.22 (d, J = 11.7 Hz, 1H), 3.47 (t, J = 18.6 Hz, 2H), 3.63 (s, 2H), 4.07 (s, 1H), 4.64 (s, 1H), 5.78 (s, 1H), 6.41 (d, J = 2.1 Hz, 1H), 6.87 (q, J = 6.5 Hz, 1H), 7.44 (m, 4H), 7.59 (d, J = 7.4 Hz, 2H), 7.64 (s, 1H), 7.77 (m, 1H), 7.91 (m, 2H)
19o	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (d, J = 18.0 Hz, 1H), 1.57 (d, J = 6.1 Hz, 4H), 2.04 (dd, J = 13.9, 6.4 Hz, 1H), 2.30 (dd, J = 13.5, 8.4 Hz, 1H), 2.39 (s, 3H), 3.11 (d, J = 11.6 Hz, 1H), 3.23 (d, J = 11.4 Hz, 1H), 3.48 (dq, J = 21.6, 7.6, 6.8 Hz, 2H), 3.64 (dd, J = 13.8, 6.9 Hz, 2H), 4.08 (m, 1H), 4.87 (s, 12H), 5.78 (s, 1H), 6.41 (d, J = 2.0 Hz, 1H), 6.91 (q, J = 6.6 Hz, 1H), 7.36 (m, 2H), 7.50 (t, J = 9.3 Hz, 2H), 7.74 (dd, J = 8.3, 2.2 Hz,

	1H), 7.90 (dd, J = 7.9, 2.1 Hz, 2H)
19p	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (s, 1H), 1.44 (s, 1H), 1.52 (q, J = 5.9 Hz, 4H), 1.85 (m, 1H), 2.11 (dd, J = 13.2, 8.8 Hz, 1H), 2.39 (s, 3H), 2.77 (d, J = 11.3 Hz, 1H), 3.01 (d, J = 11.3 Hz, 1H), 3.45 (ddt, J = 19.8, 12.8, 5.8 Hz, 2H), 3.61 (m, 2H), 3.74 (t, J = 8.0 Hz, 1H), 5.78 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.93 (q, J = 6.6 Hz, 1H), 7.54 (m, 3H), 7.75 (m, 2H), 7.92 (dd, J = 11.1, 2.0 Hz, 2H)
19q	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.57 (t, J = 5.0 Hz, 4H), 2.03 (dd, J = 13.3, 6.9 Hz, 1H), 2.29 (dd, J = 13.4, 9.0 Hz, 1H), 2.39 (s, 3H), 3.08 (d, J = 11.6 Hz, 1H), 3.22 (d, J = 11.6 Hz, 1H), 3.48 (ddt, J = 20.4, 13.2, 5.9 Hz, 2H), 3.65 (dd, J = 13.7, 6.5 Hz, 2H), 4.05 (t, J = 8.0 Hz, 1H), 5.77 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.89 (q, J = 6.6 Hz, 1H), 7.47 (m, 3H), 7.58 (m, 2H), 7.77 (dd, J = 8.3, 2.2 Hz, 1H), 7.91 (t, J = 2.4 Hz, 2H)
19r	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.94 – 7.80 (m, 9H), 7.60 (d, J = 8.1 Hz, 6H), 7.50 (dd, J = 20.7, 8.1 Hz, 9H), 6.94 (q, J = 6.2 Hz, 3H), 6.42 (d, J = 2.3 Hz, 3H), 4.66 (s, 5H), 4.38 (t, J = 8.4 Hz, 3H), 3.73 (s, 6H), 3.63 – 3.55 (m, 1H), 3.29 – 3.18 (m, 5H), 2.40 (s, 9H), 2.07 (dd, J = 13.5, 7.8 Hz, 3H), 1.70 – 1.61 (m, 10H), 1.28 (s, 1H).

Example 20: (S)-8-(2-amino-6-((R)-1-(2'-(ethoxycarbonyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




5 The title compound was made using the procedure described for (S)-8-(2-amino-6-((R)-1-(3'-(ethoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 5a) starting with (S)-8-(2-amino-6-((R)-1-(5-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid.

10

¹H NMR (400 MHz, MeOH-d₄): δ ppm 0.89 (t, J=7.15 Hz, 3 H) 1.60 (t, J=5.54 Hz, 4 H) 2.06 (dd, J=13.50, 7.25 Hz, 1 H) 2.33 (dd, J=13.42, 9.27 Hz, 1 H) 2.40 (s, 3 H) 3.08 - 3.28 (m, 2 H) 3.39 - 3.73 (m, 4 H) 3.74 - 3.98 (m, 2 H) 4.08 (dd, J=9.08, 7.32 Hz, 1 H) 5.74 (s, 1 H) 6.42 (d, J=2.34 Hz, 1 H) 6.88 (q, J=6.75 Hz, 1 H) 7.38 (dd, J=7.71, 0.93 Hz, 1 H) 7.45 - 7.56 (m, 4 H) 7.58 - 7.65 (m, 1 H) 7.82 (dd, J=7.69, 1.20 Hz, 1 H) 7.95 (d, J=2.34 Hz, 1 H). LCMS (MH⁺): 680.


Example 21: (S)-8-(2-amino-6-((R)-1-(4'-(ethoxycarbonyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was made using the procedure described for (S)-8-(2-amino-6-((R)-1-(3'-(ethoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 5) starting with (S)-8-(2-amino-6-((R)-1-(5-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.41 (t, J=7.15 Hz, 3 H) 1.58 (br. s., 4 H) 2.05 (dd, J=13.50, 7.15 Hz, 1 H) 2.30 (dd, J=13.42, 9.18 Hz, 1 H) 2.40 (s, 3 H) 3.03 - 3.28 (m, 2 H) 3.37 - 3.76 (m, 4 H) 4.07 (dd, J=9.13, 7.22 Hz, 1 H) 4.39 (q, J=7.13 Hz, 2 H) 5.78 (s, 1 H) 6.42 (d, J=2.25 Hz, 1 H) 6.86 - 7.01 (m, 1 H) 7.53 (d, J=8.30 Hz, 1 H) 7.66 - 7.77 (m, 2 H) 7.84 (dd, J=8.30, 2.20 Hz, 1 H) 7.94 (d, J=2.29 Hz, 1 H) 7.99 (d, J=1.51 Hz, 1 H) 8.06 - 8.17 (m, 2 H). LCMS (MH⁺): 680.

Example 22a: (S)-8-(2-amino-6-((R)-1-(5-ethyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



5 *Step 1:* (S)-Ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-5-vinylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (100 mg, 0.171 mmol) in MeOH (2 mL) was hydrogenated via an H-Cube apparatus using a 10% (w/w) Pd/C cartridge with a flow rate of 1.0 mL/min at RT. The catalyst was filtered and the filtrate was concentrated *in vacuo*. The residue was lyophilized from 1:1 H₂O:CH₃CN to provide (S)-ethyl
10 8-(2-amino-6-((R)-1-(5-ethyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid which was used directly in the next step.

Step 2: Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-1-(5-ethyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using
15 the LiOH general method provided the title compound as a white solid.

Using the same generic scheme below, the following examples of Table 8a can be prepared as described above for (S)-8-(2-amino-6-((R)-1-(5-ethyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid
20 (Example 22a).





Table 8a.



Ex. No.	R	CAS Name	LCMS (MH ⁺)
22a		(S)-8-(2-amino-6-((R)-1-(5-ethyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	561
22b		S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-5-propylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	575
22c		(S)-8-(2-amino-6-((R)-1-(5-butyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	589

5


Table 8b.

NMR Data for Compounds of Table 8a

Ex. No.	NMR
---------	-----

22a	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.24 (t, J=7.59 Hz, 3 H) 1.57 (m, 4 H) 2.06 (dd, J=13.42, 7.13 Hz, 1 H) 2.32 (dd, J=13.45, 9.20 Hz, 1 H) 2.37 (s, 3 H) 2.72 (q, J=7.61 Hz, 2 H) 3.18 (m, 2 H) 3.57 (m, 4 H) 4.08 (dd, J=9.13, 7.17 Hz, 1 H) 5.74 (s, 1 H) 6.36 (d, J=2.34 Hz, 1 H) 6.71 (q, J=6.65 Hz, 1 H) 7.31 (m, 1 H) 7.39 (m, 1 H) 7.56 (s, 1 H) 7.82 (d, J=2.29 Hz, 1 H)
22b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.91 (t, J=7.35 Hz, 2 H) 1.62 (m, 6 H) 2.06 (dd, J=13.52, 7.17 Hz, 1 H) 2.31 (dd, J=13.45, 9.25 Hz, 1 H) 2.37 (s, 3 H) 2.66 (t, J=7.52 Hz, 2 H) 3.18 (m, 2 H) 3.56 (m, 4 H) 4.08 (dd, J=9.13, 7.17 Hz, 1 H) 5.74 (s, 1 H) 6.36 (d, J=2.29 Hz, 1 H) 6.70 (q, J=6.70 Hz, 1 H) 7.31 (m, 1 H) 7.37 (m, 1 H) 7.53 (s, 1 H) 7.82 (d, J=2.29 Hz, 1 H)
22c	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.92 (t, J=7.37 Hz, 2 H) 1.32 (dq, J=14.94, 7.38 Hz, 2 H) 1.60 (m, 6 H) 2.06 (dd, J=13.37, 7.22 Hz, 1 H) 2.31 (dd, J=13.45, 9.25 Hz, 1 H) 2.37 (s, 3 H) 2.69 (t, J=7.59 Hz, 2 H) 3.18 (m, 2 H) 3.58 (m, 4 H) 4.08 (dd, J=9.20, 7.25 Hz, 1 H) 5.75 (s, 1 H) 6.36 (d, J=2.15 Hz, 1 H) 6.69 (q, J=6.62 Hz, 1 H) 7.30 (m, 1 H) 7.37 (m, 1 H) 7.53 (s, 1 H) 7.82 (d, J=2.29 Hz, 1 H)

Example 23: (S)-8-(2-Amino-6-((R)-1-(5-(ethoxycarbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



5

Step 1: To a solution of (S)-8-(2-amino-6-((R)-1-(5-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (product of Step 3, Example 10m) (180 mg, 0.24 mmol) in ethanol (2 mL) was added Pd(PPh₃)₂Cl₂ (34 mg, 0.048 mmol), KHCO₃ (242 mg, 2.4 mmol).

10 A balloon of CO was fitted and the reaction mixture was heated to 80 °C for 20 h, then cooled to RT. The reaction was quenched with water, and extracted with EtOAc. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated *in vacuo*.


Purification by normal phase silica gel column (CH₂Cl₂/MeOH/AcOH) provided (S)-8-(2-amino-6-((R)-1-(5-(ethoxycarbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)

15 pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid as an off-white solid.

Step 2: N-CBZ Deprotection of (S)-8-(2-amino-6-((R)-1-(5-(ethoxycarbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy) pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid was accomplished via Method A to provide the title compound as an off-white solid.

¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.34 (t, J=7.10 Hz, 3 H) 1.51 - 1.71 (m, 4 H) 1.90 (dd, J=13.28, 9.18 Hz, 1 H) 2.26 - 2.40 (m, 4 H) 3.13 (br. s., 2 H) 3.66 (br. s., 4 H) 4.29 - 4.52 (m, 4 H) 6.07 (s, 1 H) 6.47 (d, J=2.39 Hz, 1 H) 7.48 (d, J=6.05 Hz, 1 H) 7.72 (d, J=8.40 Hz, 1 H) 8.15 (dd, J=8.40, 1.95 Hz, 1 H) 8.19 - 8.29 (m, 2 H) 8.96 (d, J=5.56 Hz, 1 H) 10.36 (d, J=4.49 Hz, 1 H). LCMS (MH⁺): 604.


Example 24: (S)-8-(2-Amino-6-((R)-1-(5-carboxy-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy) pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Hydrolysis of (S)-8-(2-amino-6-((R)-1-(5-(ethoxycarbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 23) using the LiOH general method provides the title compound as a white solid.

¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.45 - 1.65 (m, 4 H) 1.83 - 1.95 (m, 1 H) 2.26 - 2.38 (m, 4 H) 3.12 (br. s., 2 H) 3.61 (br. s., 4 H) 4.36 - 4.51 (m, 1 H) 5.93 (br. s., 1 H) 6.46 (d, J=2.39 Hz, 1 H) 7.40 (m, J=5.80 Hz, 1 H) 7.67 (d, J=8.35 Hz, 1 H) 8.11 (dd, J=8.35, 1.95 Hz, 1 H) 8.21 (d, J=2.39 Hz, 1 H) 8.25 (s, 1 H) 8.93 (m, J=4.40 Hz, 1 H) 10.09 (br. s., 1 H). LCMS (MH⁺): 576.

Example 25: (S)-8-(2-Amino-6-((R)-2,2,2-trifluoro-1-(4-(hydroxymethyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy) pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (386 mg, 0.50 mmol) in DMF (10 mL) and Et₃N (0.35 mL, 2.5 mmol) was added
 5 (n-octyl)₃SiH (368 mg, 1.0 mmol). The mixture was degassed under 1 atm of CO balloon and PdCl₂(PPh₃)₂ (72 mg, 0.10 mmol) was added, then degassed again with 1 atm of CO, and heated to 80 °C for 12 h. The reaction was cooled to RT and concentrated *in vacuo*. The residue was diluted with water then extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Normal phase column chromatography on silica gel
 10 (EtOAc / heptane) provided (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-formyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a light yellow solid contaminated with about 25% of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as by-product. The mixture was used directly in the
 15 next step.


Step 2: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-formyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (36 mg, 0.05 mmol) in dichloroethane (2 mL) was added NaCNBH₃ (1M in THF,
 20 1 mL, 0.5 mmol), followed by a few drops of HOAc. The mixture was stirred at RT for 3 h then concentrated *in vacuo*. The residue was dissolved in MeOH and purified on reverse phase HPLC (MeOH/H₂O/HOAc) to provide (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(hydroxymethyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a sticky solid that was used without further
 25 purification.

Step 3: N-CBZ Deprotection was accomplished via Method B to provide (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(hydroxymethyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid.

- 5 Step 4: Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(hydroxymethyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provided the title compound as a white solid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.57 (t, J=5.15 Hz, 4 H) 1.91 - 2.12 (m, 7 H) 2.30 (dd, J=13.23, 9.42 Hz, 1 H) 2.36 (s, 3 H) 3.07 - 3.26 (m, 2 H) 3.39 - 3.54 (m, 2 H) 3.58 - 3.70 (m, 2 H) 3.99 - 4.13 (m, 1 H) 4.65 (s, 2 H) 5.71 (s, 1 H) 6.37 (d, J=2.34 Hz, 1 H) 6.74 (q, J=6.65 Hz, 1 H) 7.39 (s, 1 H) 7.45 (d, J=8.20 Hz, 1 H) 7.68 (d, J=8.10 Hz, 1 H) 7.84 (d, J=2.34 Hz, 1 H).
LCMS (MH⁺): 562.

- 15 **Example 26: (S)-8-(2-amino-6-((R)-1-(4-((dimethylamino)methyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid**



- Step 1: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-formyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (166 mg, 0.23 mmol, see Ex. 25) in dichloroethane (4 mL) and HOAc (10 mg) was added NaBH(OAc)₃ (242 mg, 1.15 mmol) and Me₂NH (2M in THF, 0.58 mL, 1.15 mmol). The reaction mixture was stirred at RT for 20 h then concentrated *in vacuo*. The residue was dissolved in MeOH (1 mL) and purified by reverse phase HPLC (MeOH/H₂O/HOAc) to provide
25 (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-((dimethylamino) methyl)-2-(3-methyl-1H-pyrazol-


1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

Step 2: N-CBZ Deprotection was accomplished via Method A to provide (S)-ethyl 8-(2-amino-6-
5 ((R)-1-(4-((dimethylamino)methyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid.

Step 3: Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-1-(4-((dimethylamino)methyl)-2-(3-methyl-
1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-
10 carboxylate using the LiOH general method provided the title compound as a white solid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.66 - 1.81 (m, 4 H) 2.10 (dd, J=13.62, 8.54 Hz, 1 H)
2.38 (s, 3 H) 2.49 (dd, J=13.62, 8.88 Hz, 1 H) 2.88 (s, 3 H) 2.90 (s, 3 H) 3.58 - 3.90 (m, 4 H)
4.37 - 4.49 (m, 2 H) 4.56 (t, J=8.69 Hz, 1 H) 6.37 (br. s., 1 H) 6.43 (d, J=2.34 Hz, 1 H) 7.06 -
15 7.12 (m, 1 H) 7.71 - 7.78 (m, 2 H) 7.85 (d, J=8.10 Hz, 1 H) 7.98 (d, J=2.39 Hz, 1 H). LCMS
(MH⁺): 589.

Example 27: (S)-8-(6-((R)-1-(4-Bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a solution of 1(R)-1-[4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl]-2,2,2-trifluoroethanol (15.7 g, 46.3 mmol, Intermediate 1) in dioxane (200 mL) was added 4,6-dichloro-2-methylpyrimidine (30.6 g, 51 mmol) and Cs₂CO₃ (61.2 g, 187 mmol). The reaction mixture was heated to 80 °C for 30 h, then cooled to RT, and filtered. The residue was
25 concentrated *in vacuo* and purified by normal phase column chromatography on silica gel

(CH₂Cl₂ /heptane) to provide (R)-4-(1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-6-chloro-2-methylpyrimidine as a white solid.


Step 2: To a solution of (R)-4-(1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-6-chloro-2-methylpyrimidine (21 g) in dioxane (200 ml) was added (S)-2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (15 g) and Na₂CO₃ (14 g). The reaction was heated to 90 °C for 48 h, then cooled to RT, filtered, and concentrated *in vacuo*. Purification of the residue on normal phase column chromatography on silica gel (EtOAc /heptane) provided (S)-2-benzyl 3-ethyl 8-(6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as an off-white solid.

Step 4: N-CBZ Deprotection was accomplished via Method A to provide (S)-ethyl 8-(6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate an off-white solid.

Step 5: Hydrolysis of (S)-ethyl 8-(6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provided the title compound as an off-white solid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.64 (br. s., 4 H) 2.10 (d, J=7.03 Hz, 1 H) 2.28 (s, 3 H) 2.35 (dd, J=13.37, 9.27 Hz, 1 H) 2.39 (s, 3 H) 3.10 - 3.20 (m, 1 H) 3.28 (d, J=11.91 Hz, 1 H) 3.45 - 3.67 (m, 2 H) 3.75 (br. s., 2 H) 4.10 (dd, J=8.98, 7.22 Hz, 1 H) 6.17 (s, 1 H) 6.43 (d, J=2.15 Hz, 1 H) 7.01 (d, J=6.44 Hz, 1 H) 7.58 - 7.75 (m, 3 H) 8.03 (d, J=2.15 Hz, 1 H). LCMS (MH⁺): 609.

Example 28: (S)-8-(6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-methyl pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described above for (S)-8-(6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (by replacing 1(R)-1-[4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl]-2,2,2-trifluoroethanol with 1(R)-1-[4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl]-2,2,2-trifluoroethanol, Intermediate 3) and obtained as an off-white solid.

¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.34 - 1.54 (m, 4 H) 1.82 (dd, J=13.01, 6.76 Hz, 1 H) 1.99 - 2.08 (m, 1 H) 2.11 (s, 3 H) 2.30 (s, 3 H) 2.92 (d, J=11.52 Hz, 1 H) 3.06 (d, J=11.52 Hz, 1 H) 3.42 - 3.65 (m, 4 H) 3.70 (dd, J=8.91, 7.00 Hz, 1 H) 6.15 (s, 1 H) 6.42 (s, 1 H) 7.43 (q, J=6.93 Hz, 1 H) 7.54 - 7.61 (m, 1 H) 7.64 (d, J=2.10 Hz, 1 H) 7.70 (d, J=8.44 Hz, 1 H) 8.19 (d, J=2.39 Hz, 1 H) 8.70 (br. s., 1 H). LCMS (MH⁺): 565.

General biaryl coupling (Suzuki) procedures

Biaryl coupling method A

Step 1: To a mixture of (S)-2-((benzyloxy)carbonyl)-8-(6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (product of Step 3, Example 10m) (150 mg, 0.2 mmol), an arylboronic acid (0.4 mmol), Pd(N,N-dimethyl μ-alaninate)₂ (3.42 mg, 0.01 mmol), and K₃PO₄ (128 mg, 0.6 mmol) were added water (3.0 mL) and EtOH (3.0 mL). The mixture was stirred at 50 °C for 12 h. The reaction was then cooled to RT, diluted with water, and extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The target biaryl compounds were purified by normal phase silica gel column (CH₂Cl₂:MeOH).

Step 2: Subsequent N-CBZ deprotection via method A afforded the final target spirocyclic amino acids.

Biaryl coupling method B

Step 1: To a mixture of (S)-2-((benzyloxy)carbonyl)-8-(6-((R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-

5 diazaspiro[4.5]decane-3-carboxylic acid (product of Step 3, Example 10m) (150 mg, 0.2 mmol), an arylboronic acid (0.4 mmol), Pd(OAc)₂ • (1,1,3,3-tetramethyl-2-N-butylguanidine)₂ (5.7 mg, 0.01 mmol), and K₂CO₃ (83.5 mg, 0.61 mmol) was added water (1.0 mL) and dioxane (3.0 mL). The reaction mixture was stirred at 44°C for 24 h. The reaction mixture was then cooled to RT, diluted with water, and extracted with EtOAc. The combined organic layers were dried over
10 Na₂SO₄, filtered, and concentrated *in vacuo*. The target biaryl compounds were purified by normal phase silica gel column (CH₂Cl₂:MeOH).

Step 2: Subsequent N-CBZ deprotection via method A afforded the final target spirocyclic amino acids.

15 Using the generic scheme below and employing the biaryl coupling method A, the following examples of Table 9 were prepared.










Table 9.




Ex. No.	Cy	CAS Name	LCMS (MH+)
29a		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(4-(2-methoxypyridin-4-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	638
29b		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-(methylsulfonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	686
29c		(S)-8-(6-((R)-1-(3',4'-difluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	645
29d		(S)-8-(6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	635
29e		(S)-8-(6-((R)-1-(3'-(ethoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	680
29f		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(4-(6-methoxypyridin-3-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	639
29g		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(4-(2-methoxypyrimidin-5-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	640
29h		(S)-8-(6-((R)-1-(2',4'-dimethoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	668



29i		(S)-8-(6-((R)-1-(4'-(ethoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	679
29j		(S)-8-(6-((R)-1-(4'-(dimethylcarbamoyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	678
29k		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(4-(2-methoxypyridin-3-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	639
29l		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-methoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	655.6
29m		(S)-8-(6-((R)-1-(3'-(dimethylcarbamoyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	679












Using the generic scheme above with the biaryl coupling method B, the following examples of Table 10 were prepared.



Table 10.




5

Ex. No.	Cy	CAS Name	LCMS (MH+)
29n		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(2',4',6'-trimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	650
29o		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	666

29p		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(2'-methoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	638
29q		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-4'-(methoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	695
29r		(S)-8-(6-((R)-1-(4'-(tert-butyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	663
29s		(S)-8-(6-((R)-1-(4'-ethoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-methylpyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	652
29t		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-(trifluoromethoxy)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	692
29u		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(3'-(methoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	666
29v		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(pyrimidin-5-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	609
29w		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	637
29x		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(3'-isopropyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	650
29y		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	626
29z		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(pyridin-3-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	609

29aa		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	638
29ab		(S)-8-(2-methyl-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(pyridin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	608

Example 30a: 8-(6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-phenoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



- 5 *Step 1:* To a solution of (R)-1-[4-chloro-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoroethanol (5.00 g, 17.2 mmol) and 4,6-dichloro-2-(methylthio)pyrimidine (3.36 g, 17.2 mmol) in dioxane (250 mL) was added Cs₂CO₃ (16.8 g, 51.6 mmol). The reaction mixture was then heated to 70 °C for 90 h, then cooled to RT. The reaction mixture was quenched with water and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered,
- 10 and concentrated *in vacuo*. Purification on a 120 g Isco RediSep silica cartridge (EtOAc:heptane) provided 4-chloro-6-[(R)-1-[4-chloro-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoroethoxy]-2-methylsulfanylpurimidine as a white solid.

- Step 2:* To a solution of 4-chloro-6-[(R)-1-[4-chloro-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-
- 15 trifluoroethoxy]-2-methylsulfanylpurimidine (4 g, 8.95 mmol) in CH₂Cl₂ (200 mL) was added *m*-CPBA (4.2 g of a 77% (w/w) source, 18.8 mmol) and the reaction was stirred at RT for 15 h. The reaction was then diluted with saturated NaHCO₃, and extracted with CH₂Cl₂. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification on a 120 g Isco RediSep silica cartridge (EtOAc:heptane) provided 4-

chloro-6-[(1*R*)-1-[4-chloro-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoroethoxy]-2-methylsulfonylpyrimidine as an off-white solid.

Step 3: To a solution of 4-chloro-6-[(1*R*)-1-[4-chloro-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoroethoxy]-2-methylsulfonylpyrimidine (2.49 g, 5.17 mmol) in dioxane (100 mL) was added 2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (1.8 g, 5.2 mmol), Cs₂CO₃ (5.06 g, 15.5 mmol), and the reaction mixture was heated to 100 °C for 1.5 h. The reaction mixture was cooled to RT, quenched with brine, and extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification on a 120 g Isco RediSep silica cartridge (EtOAc:heptane) provided (S)-2-benzyl 3-ethyl 8-(6-((*R*)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-(methylsulfonyl)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid (1.3 g) in addition to (S)-2-benzyl 3-ethyl 8-(4-chloro-6-((*R*)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-2-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate.

Step 4: To a solution of 2-benzyl 3-ethyl 8-(6-((*R*)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-(methylsulfonyl)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (2.10 g, 2.65 mmol) in 2:1 THF:H₂O (90 mL) was added LiOH (127 mg, 5.3 mmol), and the reaction was stirred at RT for 21 h, after which additional LiOH (65 mg, 2.6 mmol) was added, and the reaction was stirred for 8 h longer. The reaction was then quenched with 1 N HCl to pH<1, and extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated *in vacuo* to provide 2-((benzyloxy)carbonyl)-8-(6-((*R*)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-(methylsulfonyl)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid as an off-white solid which was used directly without further purification.

Step 5: To a solution of 2-((benzyloxy)carbonyl)-8-(6-((*R*)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-(methylsulfonyl)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (300 mg, 0.393 mmol) in 1,4-dioxane (10 mL) was added phenol (74 mg, 0.79 mmol), Cs₂CO₃ (512 mg, 1.5 mmol), and the reaction was heated to 70 °C for 21 h. The reaction was then cooled to RT, diluted with water, acidified to pH<1 with 1

N HCl, and extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification on a 50 g Isco Gold RediSep reverse phase silica cartridge (H₂O:HOAc : 99:1 MeOH:HOAc 99:1) provided 2-((benzyloxy)carbonyl)-8-(6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-

5 phenoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid as an off-white solid.

Step 6: N-CBZ Deprotection was accomplished via Method B to provide the title compound as an off-white solid.

10 Using the generic scheme below, the following examples of Table 11a were prepared as described above for 8-(6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-phenoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 30a).





Table 11a.



Ex. No.	R	CAS Name	LCMS (MH+)
30a		8-(6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-phenoxypyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	644
30b		8-(6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-(cyclohexyloxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	649


Table 11b.

NMR Data for Compounds of Table 11a

Ex. No.	NMR
30a	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.57 (br. s., 4 H), 2.00 - 2.31 (m, 2 H), 2.32 (s, 3 H), 3.06 - 3.28 (m, 2 H), 3.36 - 3.71 (m, 4 H), 4.07 (dd, J = 8.83, 7.37 Hz, 1 H), 6.11 (s, 1 H), 6.30 (d, J = 2.34 Hz, 1 H), 6.70 (q, J = 6.43 Hz, 1 H), 6.97 - 7.06 (m, 2 H), 7.10 - 7.20 (m, 1 H), 7.26 - 7.36 (m, 2 H), 7.47 (d, J = 2.15 Hz, 1 H), 7.54 (dd, J = 8.54, 2.15 Hz, 1 H), 7.71 (d, J = 8.54 Hz, 1 H), 7.86 (d, J = 2.39 Hz, 1 H).
30b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.16 - 1.95 (m, 14 H), 2.04 - 2.35 (m, 2 H), 2.36 (s, 3 H), 3.07 - 3.30 (m, 2 H), 3.43 - 3.82 (m, 4 H), 4.09 (dd, J = 8.86, 7.39 Hz, 1 H), 4.80 - 4.95 (m, 1 H), 5.98 (s, 1 H), 6.37 (d, J = 2.39 Hz, 1 H), 7.01 - 7.13 (m, 1 H), 7.45 - 7.55 (m, 2 H), 7.70 (d, J = 9.08 Hz, 1 H), 8.12 (d, J = 2.34 Hz, 1 H).

5


Example 31: 8-(6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-(cyclohexylamino)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described above by replacing the alcohol in Step 5 of Example 30a with cyclohexyl amine.

- 5 ¹H NMR (400 MHz, MeOH-d₄): δ ppm 0.99 - 1.95 (m, 14 H), 2.02 - 2.37 (m, 2 H), 2.38 (s, 3 H), 3.07 - 3.29 (m, 2 H), 3.41 - 3.77 (m, 5 H), 4.09 (dd, *J* = 9.10, 7.15 Hz, 1 H), 5.60 (s, 1 H), 6.39 (d, *J* = 2.39 Hz, 1 H), 6.87 - 7.21 (m, 1 H), 7.49 (dtd, *J* = 4.48, 2.26, 2.26, 2.12 Hz, 2 H), 7.70 (d, *J* = 9.03 Hz, 1 H), 7.87 (d, *J* = 2.34 Hz, 1 H). LCMS (MH⁺): 650.

10 **Example 32: (S)-8-(6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-(cyclobutanecarboxamido)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid**




- Step 1:* To a solution of (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (product of Step 3, Example 10m) (300 mg, 0.412 mmol) in pyridine (1.0 mL) was added cyclobutanecarbonyl chloride (54 mg, 0.045 mmol). The reaction mixture was stirred at RT for 3 h, then diluted with EtOAc, and washed with 0.5 N HCl. The organic layer was dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification on a 40 g Isco RediSep silica cartridge (EtOAc/heptane) provides (S)-2-benzyl 3-ethyl 8-(6-((R)-1-(4-
- 15
- 20

chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-(cyclobutanecarboxamido) pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as an off-white solid.

Step 2: The title compound was prepared by the N-CBZ removal using the general method B to provide a white solid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.66 (d, J = 4.30 Hz, 4 H), 1.78 - 1.99 (m, 2 H), 2.03 - 2.38 (m, 6 H), 2.39 (s, 3 H), 3.12 - 3.32 (m, 2 H), 3.47 - 3.90 (m, 5 H), 4.10 (dd, J = 9.10, 7.20 Hz, 1 H), 6.03 (s, 1 H), 6.41 (d, J = 2.34 Hz, 1 H), 6.82 - 6.98 (m, 1 H), 7.45 - 7.57 (m, 2 H), 7.73 (d, J = 8.49 Hz, 1 H), 7.97 (d, J = 2.34 Hz, 1 H). LCMS (MH⁺): 649.


Example 33: (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(2-oxopyrrolidin-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 10d) starting with (R)-1-(5-chloro-2-(2,2,2-trifluoro-1-hydroxyethyl)phenyl)pyrrolidin-2-one.

¹H NMR (DMSO-d₆): δ ppm 1.23 (m, 1H), 1.40 (m, 4H), 1.81 (dd, J = 13.2, 6.9 Hz, 1H), 2.07 (m, 4H), 2.45 (d, J = 8.1 Hz, 2H), 2.91 (d, J = 11.5 Hz, 3H), 3.06 (d, J = 11.6 Hz, 1H), 3.47 (d, J = 6.9 Hz, 3H), 3.66 (m, 3H), 5.54 (s, 1H), 6.09 (s, 2H), 6.74 (q, J = 6.9 Hz, 1H), 7.55 (m, 3H). LCMS (MH⁺): 570.

Example 34c: (S)-8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a solution of (R)-1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethanol (Intermediate 43) (400 mg, 1.4 mmol) in dioxane (25 mL) was added 4,6-dichloropyrimidin-2-amine (1.1 g, 7 mmol) and Cs₂CO₃ (1.3 g, 4 mmol). The mixture was heated for 24 h at 80 °C. The reaction was then cooled to RT and filtered. The solvent was removed *in vacuo*, then CH₂Cl₂ and heptane was added. The solvent volume was reduced until a solid precipitated out. The solid was filtered and the procedure repeated several times to provide (R)-4-(1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethoxy)-6-chloropyrimidin-2-amine as a white solid.


Step 2: To a solution of (R)-4-(1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethoxy)-6-chloropyrimidin-2-amine (100 mg, 0.24 mmol, Step 1) in dioxane (5 mL) was added (S)-2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (100 mg, 0.29 mmol), and NaHCO₃ (300 mg, 3.5 mmol). After 5 h, an additional amount of NaHCO₃ (300 mg, 3.5 mmol) was added and the reaction mixture was heated to 90 °C for 36 h. The reaction was then cooled to RT and filtered. Purification by normal phase silica gel column (EtOAc/heptane) provided (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

Step 3: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (100 mg, 0.13 mmol) in 10:1 dioxane:water (5 mL) was phenyl boronic acid (33 mg, 0.27 mmol), KHCO₃ (27 mg, 0.3 mmol), and PdCl₂(dppf)-CH₂Cl₂ (6 mg, 0.007 mmol). The reaction was heated to 100 °C for 15 h, cooled to RT, and concentrated *in vacuo*. The residue was diluted with water, and extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-

Step 2: N-CBZ Deprotection was accomplished via method B to provide (S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-sulfamoyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid.


- 5 Step 3: Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-sulfamoyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provided the title compound as an off-white solid.

10 Using the generic scheme below, the following examples of Table 12a can be prepared as described above for (S)-8-(2-amino-6-((R)-1-(5-chloro-3'-sulfamoyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 34u).












15










Table 12a.





















* Stereochemistry defined in name in table below










Ex. No.	Cy	CAS Name	LCMS (MH+)
34a		(S)-8-(2-amino-6-((R)-1-(3',5-dichloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	597
34b		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	577
34c		8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	563
34d		8-(2-amino-6-((R)-1-(2'-amino-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	577
34e		8-(2-amino-6-((R)-1-(5-chloro-3'-nitro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	606
34f		8-(2-amino-6-((R)-1-(3'-amino-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	577
34g		8-(2-amino-6-((R)-1-(5-chloro-4'-nitro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	607
34h		8-(2-amino-6-((R)-1-(4'-amino-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	577
34i		(S)-8-(2-amino-6-((R)-1-(4-chloro-2-(6-methylpyridin-2-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	578
34j		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(ethylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	655











34k		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(propylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	669
34l		(S)-8-(2-amino-6-((R)-1-(3'-(butylsulfonyl)-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	682
34m		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(hydroxymethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	592
34n		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(methylsulfonamido)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	656
34o		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(2-oxopyrrolidin-1-yl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	646
34p		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(3-methyl-2-oxoimidazolidin-1-yl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	660.5
34q		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	630
34r		(S)-8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	563
34s		(S)-8-(2-amino-6-((R)-1-(4-chloro-2-(5-chlorothiophen-2-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	604
34t		(S)-8-(2-amino-6-((R)-1-(4-chloro-2-(1-methyl-1H-pyrazol-3-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	566












34u		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-sulfamoyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	641
34v		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-hydroxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	578
34w		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	640
34x		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-cyano-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	587
34y		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-methoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	592
34z		(S)-8-(2-amino-6-((R)-1-(3'-(aminomethyl)-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	591
34aa		(S)-8-(6-((R)-1-(3'-(acrylamidomethyl)-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	645
34ab		(S)-8-(2-amino-6-((R)-1-(3'-carboxy-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	606
34ac		(S)-8-(2-amino-6-((R)-1-(3'-carbamoyl-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	605

34ad		(S)-8-(2-amino-6-((R)-1-(5-chloro-4'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	640
34ae		(S)-8-(2-amino-6-((R)-1-(5-chloro-4'-sulfamoyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	641
34af		(S)-8-(2-amino-6-((R)-1-(4',5-dichloro-3'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	615
34ag		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-isopropoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	621
34ah		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-ethoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	607
34ai		(S)-8-(2-amino-6-((R)-1-(3',5-dichloro-4'-ethoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	642
34aj		(S)-8-(2-amino-6-((R)-1-(3',5-dichloro-4'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	611
34ak		(S)-8-(2-amino-6-((R)-1-(3',5-dichloro-4'-isopropoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	655

34al		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-fluoro-4'-isopropoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	639
34am		(S)-8-(2-amino-6-((R)-1-(4',5-dichloro-3'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	665
34an		(S)-8-(2-amino-6-((R)-1-(3',5-dichloro-5'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	615
34ao		(S)-8-(2-amino-6-((R)-1-(3'-(tert-butyl)-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	619
34ap		(S)-8-(2-amino-6-((R)-1-(3',5-dichloro-5'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	665
34aq		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-fluoro-5'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	648
34ar		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-methoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	593
34as		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	580
34at		(S)-8-(2-amino-6-((R)-1-(4',5-dichloro-3'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	611
34au		(S)-8-(2-amino-6-((R)-1-(5-chloro-3',5'-difluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	598

34av		(S)-8-(2-amino-6-((R)-1-(3',5-dichloro-4'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	615
34aw		(S)-8-(2-amino-6-((R)-1-(5-chloro-3',4'-difluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	598
34ax		(S)-8-(2-amino-6-((R)-1-(3',5-dichloro-4'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	665
34ay		(S)-8-(2-amino-6-((R)-1-(5-chloro-3',4'-dimethyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	591
34az		(S)-8-(2-amino-6-((R)-1-(5-chloro-4'-ethoxy-3'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	625
34ba		(S)-8-(2-amino-6-((R)-1-(5-chloro-3',5'-dimethyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	591
34bb		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-methyl-4'-(trifluoromethoxy)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	661
34bc		(S)-8-(2-amino-6-((R)-1-(4',5-dichloro-3',5'-dimethyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	625
34bd		(S)-8-(2-amino-6-((R)-1-(5-chloro-4'-fluoro-3'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	595

34be		(S)-8-(2-amino-6-((R)-1-(3',5-dichloro-5'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	611
34bf		(S)-8-(2-amino-6-((R)-1-(5-chloro-3',4',5'-trifluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	616
34bg		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(trifluoromethoxy)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	696
34bh		(S)-8-(2-amino-6-((R)-1-(5-chloro-3',5'-bis(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	698
34bi		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-isopropyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	605
34bj		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3',5,5'-trichloro-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	631
34bk		(S)-8-(2-amino-6-((R)-1-(5-chloro-4'-fluoro-3'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	648
34bl		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-fluoro-5'-isopropoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	639
34bm		(S)-8-(2-amino-6-((R)-1-(3'-(tert-butyl)-5-chloro-5'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	633
34bn		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-fluoro-4'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	595

34bo		(S)-8-(2-amino-6-((R)-1-(4-chloro-2-(pyridin-3-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	563
34bp		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-ethoxy-4'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	625
34bq		(S)-8-(2-amino-6-((R)-1-(3'-(tert-butyl)-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	619
34br		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(prop-1-en-2-yl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	603
34bs		(S)-8-(2-amino-6-((R)-1-(4-chloro-2-(2-(dimethylamino)pyridin-4-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	603
34bt		(S)-8-(2-amino-6-((R)-1-(4-chloro-2-(naphthalen-2-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	613
34bu		(S)-8-(2-amino-6-((R)-1-(4-chloro-2-(2-isopropylpyridin-4-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	606
34bv		(S)-8-(2-amino-6-((R)-1-(5-chloro-4'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	525
34bw		(S)-8-(2-amino-6-((R)-1-(4',5-dichloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	597
34bx		(S)-8-(2-amino-6-((R)-1-(5-chloro-4'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	577
34by		(S)-8-(2-amino-6-((R)-1-(5-chloro-2',3',4',5'-tetrahydro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	567

34bz		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-isobutoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	635
34ca		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(pyrrolidine-1-carbonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	660
34cb		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(cyclopentyloxy)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	647
34cc		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(((1R,4R)-4-hydroxycyclohexyl)carbamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	740
34cd		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-ethyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	591
34ce		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-isopropyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	633
34cf		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	703

34cg		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(morpholine-4-carbonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	676
34ch		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(4-methylpiperazine-1-carbonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	689
34ci		(S)-8-(2-amino-6-((R)-1-(4-chloro-2-(2-methylthiazol-5-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	584
34cj		(S)-8-(2-amino-6-((R)-1-(4-chloro-2-(1-methyl-2-oxo-1,2-dihydropyridin-3-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	594
34ck		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(N-methylsulfamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	656
34cl		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(N,N-dimethylsulfamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	670
34cm		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(methylcarbamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	620
34cn		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(dimethylcarbamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	634
34co		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(diethylcarbamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	662








34cp		(S)-8-(6-((R)-1-(2-(1H-benzo[d]imidazol-4-yl)-4-chlorophenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	603
34cq		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(piperazine-1-carbonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	675
34cr		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(4-cyclopropylpiperazine-1-carbonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	716
34cs		(S)-8-(2-amino-6-((R)-1-(4-chloro-2-(pyridin-2-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	564
34ct		(S)-8-(2-amino-6-((R)-1-(4-chloro-2-(pyrimidin-2-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	564
34cu		(S)-8-(2-amino-6-((R)-1-(4-chloro-2-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	565
34cv		(S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(2-methoxyethoxy)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	637

Table 12b.

NMR Data for Compounds of Table 12a

Ex. No.	NMR
34a	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.31 (d, J = 15.3 Hz, 1H), 1.67 (d, J = 7.3 Hz, 4H), 2.10 (dd, J = 13.6, 8.1 Hz, 1H), 2.46 (m, 1H), 3.25 (t, J = 12.0 Hz, 2H), 3.52 (s, 2H), 3.63 (m, 3H), 4.45 (t, J = 8.6 Hz, 1H), 4.83 (d, J = 3.0 Hz, 1H), 6.59 (q, J = 6.5 Hz, 1H), 7.32 (q, J = 1.8 Hz, 1H), 7.39 (m, 1H), 7.52 (m, 4H), 7.70 (d, J = 8.4 Hz, 1H)
34b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.60 (q, J = 5.6 Hz, 4H), 2.06 (dd, J = 13.4, 7.2 Hz, 1H), 2.33 (dd, J = 13.5, 9.2 Hz, 1H), 2.43 (s, 3H), 3.13 (d, J = 11.8 Hz, 1H), 3.26 (d, J = 11.7 Hz, 1H), 3.47 (m, 2H), 3.62 (tt, J = 9.2, 4.9 Hz, 2H), 4.10 (dd, J = 9.1, 7.1 Hz, 1H), 4.61 (s, 1H), 5.48 (s, 1H), 6.66 (q, J = 6.9 Hz, 1H), 7.27 (m, 4H), 7.42 (m, 2H), 7.67 (d, J = 8.5 Hz, 1H)
34c	¹ H NMR (400 MHz, MeOH -d ₄): δ ppm 1.62 (d, J=4.88 Hz, 4 H) 2.08 (dd, J=13.47,

	7.22 Hz, 1 H) 2.34 (dd, J=13.37, 9.27 Hz, 1 H) 3.08 - 3.19 (m, 1H) 3.28 (d, J=11.71 Hz, 1 H) 3.38 - 3.56 (m, 2 H) 3.63 (d, J=5.66 Hz, 2 H) 4.11 (dd, J=8.98, 7.22 Hz, 1 H) 5.51 (s, 1 H) 6.66 (d, J=6.83 Hz, 1 H) 7.30 (d, J=2.15 Hz, 1 H) 7.41 - 7.52 (m, 4 H) 7.52 - 7.61 (m, 2 H) 7.69 (d, J=8.59 Hz, 1 H)
34d	¹ H-NMR (400 MHz, MeOH -d4): δ ppm 1.9 (m,4H), 1.98 (m,1H), 2.26 (m,1H), 3.01 (m,1H), 3.17 (m,1H), 3.48 (m,2H), 3.60 (m,2H), 3.95 (m,1H), 5.53-5.52 (d,1H), 6.26-6.22 (q,1H), 6.97-6.69 (m, 3H), 7.31-7.17(m,2H), 7.47-7.44 (m,1H), 7.74-7.63 (m,1H)
34e	¹ H NMR (400 MHz, DMSO-d6): δ ppm 1.61 (m, 4 H), 2.07-2.04 (m, 1 H), 2.37-2.33 (m, 1 H), 3.15-3.12 (d, 1 H, J=11.8 Hz), 3.25 (d, 1 H, J=11.8 Hz), 3.50-3.47 (m, 2 H), 3.67-3.66 (m, 2 H), 4.11-4.07 (t, 1H), 5.58 (s, 1 H), 6.58-6.53 (q, 1 H, J=6.8 Hz), 7.36 (s, 1 H), 7.53-7.51 (d, 1 H, J=8.4 Hz), 7.70-7.67 (d, 1 H, J=8.0 Hz), 7.82-7.78 (m, 2 H), 8.38-8.36 (d, 1 H, J=8.0 Hz), 8.58 (s, 1 H)
34f	¹ H NMR (400 MHz, MeOH -d4): δ ppm 1.58 (m, 4 H), 2.05-2.02 (m, 1 H), 2.31-2.30 (m, 1 H), 3.28-3.21 (d, 1 H, J =11.8 Hz), 3.48-3.46 (m, 2 H, J =11.8 Hz), 3.68-3.51 (m, 2 H), 4.08-4.01 (q, 1 H, J =7.0 Hz), 5.44 (s, 1 H), 6.76-6.69 (m, 4 H), 7.26-7.21 (m, 2 H), 7.41-7.40 (d, 1 H, J =8.4 Hz), 7.66-7.64 (d, 1 H, J =8.4 Hz)
34g	¹ H-NMR 400 MHz, MeOH -d4): δ ppm 1.28 (m,2H), 1.63 (m 4H), 2.10-2.04 (m,1H), 2.42-2.36 (m,1H), 3.19-3.16 (d, J =6.0,2H), 3.26 (s,1H), 3.65 (m,2H), 4.28-4.24 (t, J=16.0,1H), 5.58 (s 1H), 6.62-6.57 (m, 1H), 7.37-7.36 (d, J =4.0,1H), 7.54-7.51 (dd, J=12.0, 4.0,1H), 7.72-7.70 (d, J =8.0,1H), 7.78-7.76 (d, J =8.0,2H), 8.43-8.41 (d, J=8.0,2H)
34h	¹ H-NMR (400 MHz, MeOH -d4): δ ppm 1.29 (m,2H), 1.58 (m,4H), 2.07-2.02 (m,1H), 2.33-2.28(m,1H), 3.11-3.08 (d, J=12.0,1H), 3.24-3.21 (d, J=12.0, 1H), 3.48-3.41(m,2H), 3.60-3.55(m,2H), 4.08-4.04 (t, J=16.0, 1H), 5.39 (d, J =2.0,1H), 6.66-6.63 (m,1H),), 6.86-6.84 (d, J =8.02H), 7.19-7.17 (d, J =8.0,2H), 7.25-7.24 (d, J =4.0,1H), 7.37-7.35 (dd, J =8.0, 6.0,1H), 7.63-7.61(d, J=8.0,1H)
34i	¹ H NMR (400 MHz, MeOH-d4): δ 7.88 (t, J=7.68 Hz,1 H), 7.70 (d, J=8.52 Hz,1 H), 7.50 (m, 3 H), 7.35 (d, J=7.76 Hz,1 H), 6.99 (q, J=6.96 Hz, 1H), 5.69 (s, 1 H), 4.06 (t, J=7.48 Hz,2 H), 3.62 (m, 2 H), 3.48 (m, 2 H), 3.22 (d,J=11.64 Hz, 1H),3.09 (d, J=11.44 Hz,1 H), 2.61 (s, 3 H), 2.30 (m, 1 H), 2.03 (m,1H), 1.57 (m, 4 H).
34j	¹ H NMR (400 MHz, MeOH-d4): δ ppm 1.25 (t, J = 7.4 Hz, 3H), 1.63 (m, 4H), 2.11-2.06 (m, 1H), 2.38-2.31 (m, 1H), 3.16-3.13 (m, 1H), 3.26 (m, 1H), 3.31 (m, 2H), 3.55-3.50 (m, 2H), 3.71-3.64 (m, 2H), 4.11 (M, 1H), 5.61 (s, 1H), 6.63 (m, 1H) , 7.36 (s, 1H), 7.52-7.50 (m, 1H), 7.71-7.68 (m, 1H), 7.76-7.75 (m, 1H), 7.83 (t, J = 7.8 Hz, 1H), 8.04 (d, J = 7.2 Hz, 1H) 8.43 (s, 1H)
34k	¹ H NMR (400 MHz, MeOH -d4): δ ppm 0.96 (t, J=12.0, 4H), 1.70-1.62 (m, 8H), 2.06 (s, 1H), 2.32 (s, 1H),3.24 (d, J = 12.0, 1H), 3.50(s, 2H), 3.67(s, 2H), 4.07 (s, 1H), 4.63 (s, 1H), 5.61 (s, 1H), 6.62 (q, J=8.0, 1H), 7.37(s,1H), 7.50 (d,1H,J=8.0),7.79-7.69 (m,2H), 7.83(t,1H,J=8.0), 8.03(d,1H,J=8.0), 8.45 (s, 1H)
34l	¹ H NMR (400 MHz, MeOH-d4): δ ppm 8.47 (s, 1 H), 8.03 (d, 1 H), 7.74 (t, 1 H), 7.67 (m, 2 H), 7.51-7.49 (d, 1 H), 7.37 (s, 1 H), 6.64-6.59 (q, 1 H), 5.62 (s, 1 H), 4.12-4.08 (t, 1 H), 3.67 (m, 2 H), 3.50 (m, 2 H), 3.26 (d, 1 H), 3.13 (d, 1 H), 2.35-2.32 (m, 1 H), 2.05 (m, 1 H), 1.63 (m, 6 H), 1.34 (q, 2 H), 0.84-0.80 (t, 3 H)
34m	¹ H NMR (400 MHz, MeOH -d4): δ ppm 7.74-7.66 (m, 2H), 7.48-7.39 (m, 3H), 7.27 (m, 2H), 6.73-6.71 (m, 1H), 5.53 (s, 1H), 4.73 (s, 2H), 4.08 (t, J = 7.1 Hz, 1H), 3.63 (m, 2H), 3.47 (m, 2H), 3.27-3.24 (m, 1H), 3.14-3.11 (m, 1H), 2.36-2.30 (m, 1H), 2.08-2.03 (m,

	1H), 1.60 (m, 4H)
34n	¹ H NMR (400 MHz, MeOH -d ₄) δ ppm 7.66 (d, 1 H, J=8.6 Hz), 7.50 (m, 3 H), 7.31 (m, 2 H), 7.24 (d, 1 H, J=8.2 Hz), 6.61 (m, 1 H), 4.21 (m, 1 H), 3.63 (m, 2 H), 3.48 (m, 2 H), 3.21 (m, 1 H), 3.18 (m, 1 H), 3.01 (s, 3 H), 2.37 (m, 1 H), 2.07 (m, 1 H), 1.62 (m, 4 H)
34o	¹ H-NMR (400 MHz, MeOH-d ₄): δ ppm 7.97 (s, 1H), 7.60-7.67 (m, 2H), 7.52-7.54 (m, 1H), 7.40 -7.46 (m, 1H), 7.31 - 7.31 (m, 1H), 7.22 - 7.24 (m, 1H), 6.61 -6.66 (m, 1H), 5.51 (s, 1H), 4.39 -4.04 (m, 4H), 3.52 -3.60 (m, 2H), 3.42 -3.50 (m, 2H), 3.15 -3.18 (d, 1H), 2.96 -2.99 (d, 1H), 2.60 -2.64 (m, 2H), 2.18 -2.28 (m, 3H), 1.96 -2.00 (m, 1H), 1.58 -1.59 (m, 4H)
34p	¹ H-NMR (400 MHz, DMSO-d ₆): δ ppm 8.00 (s, 1H), 7.56 -7.58 (m, 1H), 7.43 -7.49 (m, 2H), 7.26 -7.27 (m, 2H), 6.97 -7.97 (m, 1H), 6.59 -6.55 (m, 1H), 5.48 (s, 1H), 3.78 -3.82 (m, 1H), 3.70 -3.74 (m, 2H), 3.39-3.44 (m, 6H), 2.98 -3.02 (d, 1H), 2.82 -2.85 (d, 1H), 2.70 (s, 3H), 2.04 -2.11 (m, 1H), 1.69 -1.75 (m, 1H), 1.36 -1.40 (m, 4H)
34q	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.64 (t, J = 5.8 Hz, 4H), 2.08 (dd, J = 13.5, 7.6 Hz, 1H), 2.40 (dd, J = 13.5, 9.0 Hz, 1H), 3.19 (d, J = 11.8 Hz, 1H), 3.28 (d, J = 12.2 Hz, 1H), 3.51 (m, 2H), 3.66 (m, 2H), 4.28 (t, J = 8.4 Hz, 1H), 4.87 (s, 16H), 6.53 (q, J = 6.7 Hz, 1H), 7.34 (d, J = 2.3 Hz, 1H), 7.51 (dd, J = 8.6, 2.3 Hz, 1H), 7.75 (m, 5H)
34r	¹ H NMR (400 MHz, MeOH-d) δ ppm 1.62 (d, J=4.88 Hz, 4 H) 2.08 (dd, J=13.47, 7.22 Hz, 1 H) 2.34 (dd, J=13.37, 9.27 Hz, 1 H) 3.08 - 3.19 (m, 1H) 3.28 (d, J=11.71 Hz, 1 H) 3.38 - 3.56 (m, 2 H) 3.63 (d, J=5.66 Hz, 2 H) 4.11 (dd, J=8.98, 7.22 Hz, 1 H) 5.51 (s, 1 H) 6.66 (d, J=6.83 Hz, 1 H) 7.30 (d, J=2.15 Hz, 1 H) 7.41 - 7.52 (m, 4 H) 7.52 - 7.61 (m, 2 H) 7.69 (d, J=8.59 Hz, 1 H)
34s	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.60 (d, J=5.47 Hz, 4 H) 1.98 - 2.12 (m, 1 H) 2.26 - 2.39 (m, 1 H) 3.07 - 3.17 (m, 1 H) 3.20 - 3.29 (m, 1 H) 3.38 - 3.55 (m, 2 H) 3.56 - 3.71 (m, 2 H) 4.01 - 4.15 (m, 1 H) 5.51 (s, 1 H) 6.74 - 6.89 (m, 1 H) 7.11 (s, 1 H) 7.14 (s, 1 H) 7.42 (d, J=2.15 Hz, 11 H) 7.44 - 7.53 (m, 1 H) 7.61 - 7.73 (m, 1 H)
34t	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.51 (d, J=4.69 Hz, 4 H) 1.84 - 2.00 (m, 1 H) 2.09 - 2.31 (m, 1 H) 2.82 - 3.00 (m, 1 H) 3.02 - 3.20 (m, 1 H) 3.32 - 3.64 (m, 4 H) 3.84 - 3.94 (m, 1 H) 3.98 (s, 3 H), 5.50 (s, 1 H) 6.63 (d, J=1.95 Hz, 1 H) 7.13 - 7.27 (m, 1 H) 7.39 (d, J=1.56 Hz, 1 H) 7.55 (d, J=1.76 Hz, 1 H) 7.64 (d, J=8.59 Hz, 1 H) 7.71 (d, J=1.76 Hz, 1 H)
34u	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.52 - 1.75 (m, 4 H) 2.07 (dd, J=13.40, 7.30 Hz, 1 H) 2.34 (dd, J=13.42, 9.18 Hz, 1 H) 3.07 - 3.29 (m, 2 H) 3.40 - 3.78 (m, 4 H) 4.10 (dd, J=9.10, 7.25 Hz, 1 H) 5.59 (s, 1 H) 6.61 (q, J=6.59 Hz, 1 H) 7.31 (d, J=2.20 Hz, 1 H) 7.49 (dd, J=8.52, 2.22 Hz, 1 H) 7.61 (d, J=7.03 Hz, 1 H) 7.65 - 7.80 (m, 2 H) 7.97 - 8.10 (m, 1 H) 8.32 (br. s., 1 H)
34v	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.52 - 1.71 (m, 4 H) 2.07 (dd, J=13.42, 7.27 Hz, 1 H) 2.33 (dd, J=13.47, 9.27 Hz, 1 H) 3.08 - 3.29 (m, 2 H) 3.36 - 3.76 (m, 4 H) 4.09 (dd, J=9.15, 7.20 Hz, 1 H) 5.48 (s, 1 H) 6.74 (q, J=7.00 Hz, 1 H) 6.87 (d, J=7.47 Hz, 1 H) 6.91 (ddd, J=8.19, 2.48, 0.85 Hz, 1 H) 7.05 (d, J=0.73 Hz, 1 H) 7.28 (d, J=2.20 Hz, 1 H) 7.32 (t, J=7.88 Hz, 1 H) 7.43 (dd, J=8.49, 2.25 Hz, 1 H) 7.67 (d, J=8.49 Hz, 1 H)
34w	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.43 - 1.76 (m, 4 H) 2.08 (dd, J=13.45, 7.15 Hz, 1 H) 2.34 (dd, J=13.42, 9.22 Hz, 1 H) 3.13 - 3.29 (m, 5 H) 3.41 - 3.77 (m, 4 H) 4.10 (dd, J=9.18, 7.17 Hz, 1 H) 5.60 (s, 1 H) 6.57 (q, J=6.57 Hz, 1 H) 7.35 (d, J=2.15 Hz, 1 H) 7.51 (dd, J=8.52, 2.17 Hz, 1 H) 7.70 (d, J=8.54 Hz, 1 H) 7.72 - 7.78 (m, 1 H) 7.78 - 7.87 (m, 1 H) 8.04 - 8.15 (m, 1 H) 8.41 (d, J=0.39 Hz, 1 H)

34x	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.46 - 1.69 (m, 4 H) 1.90 (dd, J=13.25, 9.20 Hz, 1 H) 2.35 (dd, J=13.35, 8.66 Hz, 1 H) 3.14 (br. s., 2 H) 3.64 (br. s., 4 H) 4.45 (t, J=6.49 Hz, 1 H) 5.84 (br. s., 1 H), 6.56 (q, J=6.77 Hz, 1 H) 7.48 (d, J=1.07 Hz, 1 H) 7.62 - 7.69 (m, 2 H) 7.75 - 7.82 (m, 1 H) 7.83 - 7.91 (m, 1 H) 7.92 - 8.00 (m, 2 H) 8.97 (br. s., 1 H) 10.23 (br. s., 1 H)
34y	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.46 - 1.71 (m, 4 H) 1.91 (dd, J=13.32, 9.27 Hz, 1 H) 2.27 - 2.40 (m, 1 H) 3.14 (br. s., 2 H) 3.63 (d, J=5.37 Hz, 4 H) 3.81 (s, 3 H) 4.36 - 4.53 (m, 1 H) 5.85 (br. s., 1 H) 6.72 (q, J=6.62 Hz, 1 H) 6.94 - 7.10 (m, 3 H) 7.40 (d, J=2.05 Hz, 1 H) 7.49 (t, J=7.96 Hz, 1 H) 7.57 - 7.70 (m, 2 H) 8.96 (d, J=5.71 Hz, 1 H) 10.27 (br. s., 1 H)
34z	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.43 - 1.76 (m, 4 H) 1.92 (dd, J=13.18, 9.32 Hz, 1 H) 2.35 (dd, J=13.30, 8.57 Hz, 1 H) 3.14 (br. s., 2 H) 3.67 (br. s., 4 H) 3.97 - 4.18 (m, 2 H) 4.44 (t, J=6.88 Hz, 1 H) 5.93 (br. s., 1 H) 6.75 (q, J=6.57 Hz, 1 H) 7.39 (d, J=1.66 Hz, 1 H) 7.53 (br. s., 1 H) 7.57 - 7.71 (m, 5 H) 8.58 (br. s., 3 H) 9.01 (br. s., 1 H) 10.55 (br. s., 1 H)
34aa	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.68 - 1.86 (m, 5 H) 2.13 (dd, J=13.69, 8.66 Hz, 1 H) 2.54 (dd, J=13.72, 8.98 Hz, 1 H) 3.56 (br. s., 1 H) 3.67 (br. s., 3 H) 4.44 - 4.63 (m, 3 H) 5.70 (dd, J=9.42, 2.54 Hz, 1 H) 6.19 - 6.35 (m, 2 H) 6.58 (br. s., 1 H) 7.28 (d, J=7.57 Hz, 1 H) 7.34 - 7.40 (m, 2 H) 7.43 (d, J=7.86 Hz, 1 H) 7.47 - 7.56 (m, 2 H) 7.71 (d, J=8.64 Hz, 1 H)
34ab	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.44 - 1.69 (m, 4 H) 1.91 (dd, J=13.28, 9.18 Hz, 1 H) 2.35 (dd, J=13.15, 8.61 Hz, 1 H) 3.14 (br. s., 2 H) 3.64 (br. s., 4 H) 4.37 - 4.53 (m, 1 H) 5.87 (br. s., 1 H) 6.62 (q, J=6.78 Hz, 1 H) 7.43 (t, J=1.22 Hz, 1 H) 7.65 (s, 2 H) 7.70 (d, J=4.78 Hz, 2 H) 7.99 - 8.12 (m, 1 H) 8.26 (br. s., 1 H) 8.96 (d, J=5.03 Hz, 1 H) 10.25 (br. s., 1 H)
34ac	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.62 (d, J=4.15 Hz, 4 H) 2.02 - 2.14 (m, 1 H) 2.26 - 2.43 (m, 1 H) 3.08 - 3.29 (m, 2 H) 3.40 - 3.77 (m, 4 H) 4.09 (dd, J=8.98, 7.27 Hz, 1 H) 5.55 (s, 1 H) 6.55 - 6.70 (m, 1 H) 7.30 (d, J=2.05 Hz, 1 H) 7.47 (dd, J=8.47, 2.07 Hz, 1 H) 7.51 - 7.59 (m, 1 H) 7.59 - 7.65 (m, 1 H) 7.67 (d, J=8.30 Hz, 1 H) 7.96 (dd, J=8.52, 1.00 Hz, 1 H) 8.32 - 8.50 (m, 1 H)
34ad	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.50 - 1.76 (m, 4 H) 2.07 (dd, J=13.20, 7.15 Hz, 1 H) 2.34 (dd, J=13.47, 9.27 Hz, 1 H) 3.14 (d, J=11.71 Hz, 1 H) 3.23 (s, 3 H) 3.27 (d, J=11.86 Hz, 1 H) 3.40 - 3.76 (m, 4 H) 4.09 (dd, J=9.03, 7.27 Hz, 1 H) 5.54 (s, 1 H) 6.60 (q, J=6.64 Hz, 1 H) 7.34 (d, J=2.15 Hz, 1 H) 7.52 (dd, J=8.49, 2.20 Hz, 1 H) 7.72 (d, J=8.54 Hz, 1 H) 7.78 (d, J=7.76 Hz, 2 H) 8.14 (d, J=8.64 Hz, 2 H)
34ae	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.61 (q, J=4.88 Hz, 4 H) 2.07 (dd, J=13.37, 7.13 Hz, 1 H) 2.33 (dd, J=13.42, 9.22 Hz, 1 H) 3.08 - 3.30 (m, 2 H) 3.38 - 3.74 (m, 4 H) 4.10 (dd, J=8.91, 7.35 Hz, 1 H), 5.52 (s, 1 H) 6.53 - 6.69 (m, 1 H) 7.33 (d, J=2.20 Hz, 1 H) 7.50 (dd, J=8.52, 2.22 Hz, 1 H) 7.67 (d, J=7.96 Hz, 2 H) 7.71 (d, J=8.54 Hz, 1 H) 8.08 (d, J=8.64 Hz, 2 H)
34af	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.62 (q, J = 6.0, 5.0 Hz, 17H), 2.07 (dd, J = 13.4, 7.2 Hz, 5H), 2.33 (dd, J = 13.4, 9.2 Hz, 5H), 3.15 (d, J = 11.8 Hz, 5H), 3.27 (d, J = 11.8 Hz, 13H), 3.49 (m, 9H), 3.64 (ddt, J = 15.7, 10.7, 5.2 Hz, 9H), 4.11 (dd, J = 9.2, 7.1 Hz, 5H), 6.61 (q, J = 6.7 Hz, 4H), 7.33 (m, 7H), 7.47 (m, 8H), 7.66 (m, 8H)
34ag	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.86 (m, 1H), 1.34 (dd, J = 9.4, 6.0 Hz, 7H), 1.58 (t, J = 5.7 Hz, 4H), 1.98 (dd, J = 13.3, 7.0 Hz, 1H), 2.26 (dd, J = 13.3, 9.0 Hz, 1H), 3.00

	(d, J = 11.5 Hz, 1H), 3.16 (d, J = 11.5 Hz, 1H), 3.46 (ddt, J = 19.2, 12.6, 5.9 Hz, 2H), 3.62 (dt, J = 12.8, 7.1 Hz, 2H), 3.96 (t, J = 8.1 Hz, 1H), 4.69 (p, J = 6.0 Hz, 1H), 5.50 (s, 1H), 6.73 (q, J = 6.9 Hz, 1H), 6.95 (m, 1H), 7.04 (dd, J = 8.5, 2.4 Hz, 1H), 7.20 (s, 1H), 7.28 (d, J = 2.3 Hz, 1H), 7.42 (m, 2H), 7.66 (d, J = 8.5 Hz, 1H)
34ah	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 7.3 Hz, 1H), 1.41 (t, J = 7.0 Hz, 3H), 1.61 (q, J = 6.2, 5.4 Hz, 4H), 2.07 (dd, J = 13.5, 7.3 Hz, 1H), 2.35 (dd, J = 13.5, 9.1 Hz, 1H), 3.14 (d, J = 11.8 Hz, 1H), 3.26 (d, J = 11.7 Hz, 1H), 3.33 (s, 1H), 3.48 (m, 2H), 3.66 (dd, J = 14.5, 6.2 Hz, 2H), 4.13 (tt, J = 9.7, 7.2 Hz, 3H), 4.87 (s, 17H), 6.74 (q, J = 6.9 Hz, 1H), 6.97 (d, J = 7.6 Hz, 1H), 7.04 (dd, J = 8.3, 2.5 Hz, 1H), 7.19 (s, 1H), 7.29 (d, J = 2.2 Hz, 1H), 7.44 (m, 2H), 7.67 (d, J = 8.5 Hz, 1H)
34ai	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.48 (t, J = 7.0 Hz, 3H), 1.60 (m, 4H), 2.06 (dd, J = 13.4, 6.9 Hz, 1H), 2.33 (dd, J = 13.4, 8.9 Hz, 1H), 3.13 (d, J = 11.6 Hz, 1H), 3.25 (d, J = 11.4 Hz, 1H), 3.48 (ddd, J = 21.0, 14.2, 7.2 Hz, 2H), 3.64 (q, J = 8.9, 8.1 Hz, 2H), 4.09 (t, J = 8.3 Hz, 1H), 4.20 (q, J = 6.9 Hz, 2H), 4.88 (s, 15H), 5.52 (s, 1H), 6.63 (q, J = 6.7 Hz, 1H), 7.24 (m, 2H), 7.34 (d, J = 8.3 Hz, 1H), 7.43 (dd, J = 8.5, 2.3 Hz, 2H), 7.57 (s, 1H), 7.65 (d, J = 8.4 Hz, 1H)
34aj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.58 (dd, J = 7.1, 4.1 Hz, 4H), 1.99 (dd, J = 13.4, 7.1 Hz, 1H), 2.29 (m, 1H), 2.46 (s, 3H), 3.02 (d, J = 11.6 Hz, 1H), 3.18 (d, J = 11.6 Hz, 1H), 3.46 (ddt, J = 21.0, 13.5, 6.0 Hz, 2H), 3.63 (m, 2H), 3.98 (dd, J = 9.2, 7.1 Hz, 1H), 5.52 (s, 1H), 6.60 (q, J = 6.6 Hz, 1H), 7.28 (m, 2H), 7.45 (dd, J = 8.3, 2.4 Hz, 2H), 7.56 (m, 1H), 7.66 (d, J = 8.5 Hz, 1H)
34ak	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 6.2 Hz, 1H), 1.40 (d, J = 6.0 Hz, 6H), 1.60 (d, J = 5.2 Hz, 4H), 2.03 (dd, J = 13.4, 6.8 Hz, 1H), 2.30 (dd, J = 13.3, 8.9 Hz, 1H), 2.81 (s, 1H), 3.07 (d, J = 11.6 Hz, 1H), 3.22 (d, J = 11.8 Hz, 1H), 3.48 (m, 2H), 3.64 (d, J = 9.7 Hz, 2H), 4.03 (t, J = 7.9 Hz, 1H), 4.75 (m, 1H), 5.53 (s, 1H), 6.63 (q, J = 6.8 Hz, 1H), 7.29 (m, 3H), 7.43 (dd, J = 8.6, 2.3 Hz, 1H), 7.58 (s, 1H), 7.65 (d, J = 8.5 Hz, 1H)
34al	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.39 (d, J = 6.0 Hz, 6H), 1.60 (m, 4H), 2.06 (dd, J = 13.5, 7.2 Hz, 1H), 2.33 (dd, J = 13.4, 9.2 Hz, 1H), 3.13 (d, J = 11.7 Hz, 1H), 3.26 (d, J = 11.7 Hz, 1H), 3.48 (m, 2H), 3.64 (q, J = 12.4, 10.4 Hz, 2H), 4.10 (dd, J = 9.2, 7.2 Hz, 1H), 4.70 (hept, J = 5.9 Hz, 1H), 5.53 (s, 1H), 6.67 (q, J = 6.8 Hz, 1H), 7.25 (m, 4H), 7.43 (dd, J = 8.5, 2.3 Hz, 1H), 7.65 (d, J = 8.4 Hz, 1H)
34am	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.61 (q, J = 5.7 Hz, 4H), 2.06 (dd, J = 13.4, 7.1 Hz, 1H), 2.33 (dd, J = 13.4, 9.2 Hz, 1H), 3.12 (d, J = 11.7 Hz, 1H), 3.26 (d, J = 11.6 Hz, 1H), 3.48 (ddt, J = 18.1, 13.6, 6.0 Hz, 2H), 3.65 (tt, J = 11.7, 4.8 Hz, 2H), 4.08 (dd, J = 9.2, 7.1 Hz, 1H), 5.54 (s, 1H), 6.49 (q, J = 6.6 Hz, 1H), 7.35 (d, J = 2.2 Hz, 1H), 7.51 (dd, J = 8.5, 2.3 Hz, 1H), 7.69 (d, J = 8.5 Hz, 1H), 7.82 (td, J = 17.6, 16.6, 4.9 Hz, 3H)
34an	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.88 (d, J = 7.8 Hz, 1H), 1.29 (d, J = 6.7 Hz, 2H), 1.62 (q, J = 5.9, 5.4 Hz, 4H), 2.07 (dd, J = 13.4, 7.2 Hz, 1H), 2.33 (dd, J = 13.4, 9.2 Hz, 1H), 3.13 (d, J = 11.7 Hz, 1H), 3.26 (d, J = 11.7 Hz, 1H), 3.49 (ddd, J = 21.4, 12.5, 5.9 Hz, 2H), 3.65 (td, J = 12.3, 10.3, 5.5 Hz, 2H), 4.10 (dd, J = 9.2, 7.1 Hz, 1H), 5.55 (s, 1H), 6.59 (q, J = 6.7 Hz, 1H), 7.25 (d, J = 14.9 Hz, 1H), 7.35 (m, 2H), 7.44 (s, 1H), 7.51 (dd, J = 8.5, 2.2 Hz, 1H), 7.69 (d, J = 8.5 Hz, 1H)
34ao	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (m, 2H), 1.30 (d, J = 11.8 Hz, 3H), 1.36 (s, 10H), 1.60 (m, 5H), 2.34 (s, 1H), 2.81 (s, 1H), 3.14 (d, J = 11.6 Hz, 1H), 3.28 (m, 4H),

	3.45 (s, 2H), 3.62 (s, 3H), 4.15 (t, J = 8.1 Hz, 1H), 4.85 (s, 38H), 6.62 (t, J = 6.7 Hz, 1H), 7.28 (m, 2H), 7.46 (m, 5H), 7.66 (d, J = 8.5 Hz, 1H)
34ap	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 7.5 Hz, 2H), 1.65 (s, 3H), 2.08 (dd, J = 13.5, 7.9 Hz, 1H), 2.44 (t, J = 11.2 Hz, 1H), 3.24 (dd, J = 14.0, 11.6 Hz, 1H), 3.61 (m, 4H), 4.41 (t, J = 8.4 Hz, 1H), 6.48 (q, J = 6.6 Hz, 1H), 7.38 (d, J = 2.2 Hz, 1H), 7.54 (dd, J = 8.5, 2.2 Hz, 1H), 7.71 (d, J = 8.6 Hz, 1H), 7.86 (t, J = 1.5 Hz, 2H)
34aq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (d, J = 2.2 Hz, 1H), 1.60 (s, 4H), 2.03 (d, J = 13.1 Hz, 1H), 2.30 (t, J = 11.1 Hz, 1H), 3.08 (d, J = 11.6 Hz, 1H), 3.23 (d, J = 12.3 Hz, 1H), 3.50 (dt, J = 24.2, 8.0 Hz, 2H), 3.64 (d, J = 10.2 Hz, 2H), 4.03 (t, J = 7.8 Hz, 1H), 4.58 (s, 1H), 5.55 (s, 1H), 6.52 (q, J = 6.7 Hz, 1H), 7.37 (d, J = 2.2 Hz, 1H), 7.53 (dd, J = 8.5, 2.3 Hz, 1H), 7.66 (m, 4H)
34ar	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.57 (q, J = 7.9, 6.6 Hz, 4H), 1.92 (dd, J = 13.2, 7.0 Hz, 1H), 2.19 (dd, J = 13.2, 9.0 Hz, 1H), 2.88 (d, J = 11.4 Hz, 1H), 3.10 (d, J = 11.4 Hz, 1H), 3.45 (ddt, J = 20.3, 13.1, 6.1 Hz, 2H), 3.61 (m, 2H), 3.87 (s, 4H), 5.48 (s, 1H), 6.70 (q, J = 6.9 Hz, 1H), 7.04 (m, 2H), 7.16 (s, 1H), 7.29 (d, J = 2.2 Hz, 1H), 7.45 (m, 2H), 7.67 (d, J = 8.5 Hz, 1H)
34as	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 10.0 Hz, 2H), 1.61 (d, J = 5.6 Hz, 5H), 2.06 (dd, J = 13.4, 7.2 Hz, 1H), 2.33 (dd, J = 13.4, 9.2 Hz, 1H), 3.13 (d, J = 11.7 Hz, 1H), 3.26 (d, J = 11.7 Hz, 1H), 3.49 (dt, J = 21.9, 7.2 Hz, 3H), 3.65 (ddt, J = 15.1, 10.1, 5.2 Hz, 3H), 4.09 (dd, J = 9.2, 7.1 Hz, 1H), 4.86 (s, 26H), 5.53 (s, 1H), 6.64 (q, J = 6.8 Hz, 1H), 7.28 (m, 5H), 7.47 (dd, J = 8.5, 2.3 Hz, 1H), 7.55 (m, 1H), 7.68 (d, J = 8.5 Hz, 1H)
34at	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.35 (s, 1H), 1.60 (q, J = 5.1 Hz, 4H), 2.03 (dd, J = 13.3, 7.1 Hz, 1H), 2.30 (dd, J = 13.4, 9.1 Hz, 1H), 2.45 (s, 3H), 3.08 (d, J = 11.6 Hz, 1H), 3.23 (d, J = 11.7 Hz, 1H), 3.31 (s, 3H), 3.47 (ddt, J = 19.8, 12.7, 5.7 Hz, 2H), 3.63 (dt, J = 12.9, 7.5 Hz, 2H), 4.04 (dd, J = 9.0, 7.3 Hz, 1H), 5.51 (s, 1H), 6.62 (q, J = 6.8 Hz, 1H), 7.30 (m, 3H), 7.45 (dd, J = 8.5, 2.3 Hz, 1H), 7.52 (d, J = 8.1 Hz, 1H), 7.66 (d, J = 8.5 Hz, 1H)
34au	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 6.2 Hz, 1H), 1.62 (q, J = 5.5 Hz, 4H), 2.07 (dd, J = 13.5, 7.3 Hz, 1H), 2.35 (dd, J = 13.5, 9.2 Hz, 1H), 3.15 (d, J = 11.8 Hz, 1H), 3.27 (d, J = 11.7 Hz, 1H), 3.49 (ddt, J = 21.6, 13.6, 6.3 Hz, 2H), 3.66 (ddt, J = 15.6, 10.1, 4.9 Hz, 2H), 4.14 (dd, J = 9.1, 7.3 Hz, 1H), 4.85 (d, J = 3.1 Hz, 16H), 6.63 (q, J = 6.8 Hz, 1H), 7.12 (m, 3H), 7.34 (d, J = 2.2 Hz, 1H), 7.50 (dd, J = 8.5, 2.2 Hz, 1H), 7.69 (d, J = 8.5 Hz, 1H)
34av	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.88 (m, 1H), 1.28 (s, 3H), 1.61 (q, J = 6.1 Hz, 4H), 2.07 (dd, J = 13.4, 7.1 Hz, 1H), 2.33 (dd, J = 13.5, 9.0 Hz, 1H), 3.13 (d, J = 11.6 Hz, 1H), 3.26 (d, J = 11.8 Hz, 2H), 3.48 (ddt, J = 20.7, 12.7, 5.7 Hz, 2H), 3.65 (q, J = 8.9, 6.2 Hz, 2H), 4.10 (m, 1H), 4.90 (s, 1H), 5.55 (s, 1H), 6.57 (q, J = 6.8 Hz, 1H), 7.42 (m, 5H), 7.67 (d, J = 8.3 Hz, 2H)
34aw	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (t, J = 7.7 Hz, 2H), 1.60 (q, J = 6.1, 4.9 Hz, 4H), 2.03 (dd, J = 13.3, 7.1 Hz, 1H), 2.30 (dd, J = 13.4, 9.0 Hz, 1H), 3.07 (d, J = 11.6 Hz, 1H), 3.22 (d, J = 11.6 Hz, 1H), 3.48 (ddt, J = 20.9, 13.3, 5.7 Hz, 2H), 3.64 (tt, J = 10.8, 5.3 Hz, 2H), 4.03 (t, J = 8.1 Hz, 1H), 4.87 (s, 17H), 5.54 (s, 1H), 6.61 (q, J = 6.7 Hz, 1H), 7.32 (t, J = 5.0 Hz, 2H), 7.46 (m, 3H), 7.54 (s, 1H), 7.67 (d, J = 8.5 Hz, 1H)
34ax	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (dd, J = 17.9, 6.7 Hz, 2H), 1.54 (m, 4H), 1.79 (dd, J = 12.9, 7.0 Hz, 1H), 2.06 (td, J = 16.5, 14.8, 7.8 Hz, 1H), 2.66 (d, J = 11.0 Hz,

	1H), 2.97 (d, J = 11.1 Hz, 1H), 3.45 (ddt, J = 20.1, 13.2, 6.0 Hz, 2H), 3.62 (m, 3H), 5.50 (d, J = 16.5 Hz, 1H), 6.54 (q, J = 6.7 Hz, 1H), 7.34 (d, J = 2.3 Hz, 1H), 7.52 (dd, J = 8.5, 2.3 Hz, 1H), 7.68 (dd, J = 24.2, 8.1 Hz, 2H), 7.84 (m, 1H), 7.97 (d, J = 8.1 Hz, 1H)
34ay	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (m, 2H), 1.28 (s, 2H), 1.40 (s, 10H), 1.60 (q, J = 5.5 Hz, 4H), 2.06 (dd, J = 13.5, 7.1 Hz, 1H), 2.35 (d, J = 3.7 Hz, 7H), 3.13 (d, J = 11.6 Hz, 1H), 3.25 (d, J = 11.6 Hz, 1H), 3.47 (dq, J = 22.4, 7.8, 6.8 Hz, 2H), 3.63 (dd, J = 13.9, 7.3 Hz, 2H), 4.11 (t, J = 8.3 Hz, 1H), 6.66 (q, J = 6.8 Hz, 1H), 7.17 (d, J = 7.1 Hz, 2H), 7.26 (m, 2H), 7.41 (dd, J = 8.5, 2.2 Hz, 1H), 7.65 (d, J = 8.5 Hz, 1H)
34az	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.88 (q, J = 10.5, 8.0 Hz, 1H), 1.30 (d, J = 12.4 Hz, 4H), 1.46 (t, J = 7.0 Hz, 5H), 1.61 (d, J = 5.7 Hz, 8H), 2.06 (dd, J = 13.4, 7.1 Hz, 2H), 2.33 (dd, J = 13.3, 8.8 Hz, 2H), 3.13 (d, J = 11.5 Hz, 2H), 3.26 (d, J = 11.8 Hz, 2H), 3.48 (m, 4H), 3.62 (d, J = 12.9 Hz, 3H), 4.19 (m, 5H), 4.84 (s, 2H), 6.67 (q, J = 6.7 Hz, 2H), 7.27 (m, 7H), 7.43 (dd, J = 8.6, 2.2 Hz, 2H), 7.65 (d, J = 8.5 Hz, 2H)
34ba	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.59 (m, 4H), 2.04 (dd, J = 13.4, 7.2 Hz, 1H), 2.38 (s, 7H), 3.08 (d, J = 11.7 Hz, 1H), 3.23 (d, J = 11.7 Hz, 1H), 3.46 (m, 2H), 3.64 (dt, J = 14.7, 5.8 Hz, 2H), 4.04 (dd, J = 9.2, 7.1 Hz, 1H), 5.48 (s, 1H), 6.67 (q, J = 6.9 Hz, 1H), 7.03 (s, 2H), 7.12 (s, 1H), 7.25 (d, J = 2.3 Hz, 1H), 7.42 (dd, J = 8.5, 2.3 Hz, 1H), 7.66 (d, J = 8.5 Hz, 1H)
34bb	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 3H), 1.58 (m, 4H), 1.98 (dd, J = 13.2, 7.1 Hz, 1H), 2.25 (dd, J = 13.3, 9.1 Hz, 1H), 2.40 (s, 3H), 2.98 (d, J = 11.5 Hz, 1H), 3.17 (d, J = 11.5 Hz, 1H), 3.46 (ddt, J = 20.0, 13.0, 6.1 Hz, 2H), 3.63 (dq, J = 12.7, 6.3 Hz, 2H), 3.95 (dd, J = 9.1, 7.0 Hz, 1H), 4.89 (s, 17H), 5.52 (s, 1H), 6.61 (q, J = 6.7 Hz, 1H), 7.30 (d, J = 2.3 Hz, 1H), 7.46 (m, 4H), 7.67 (d, J = 8.4 Hz, 1H)
34bc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.88 (d, J = 7.5 Hz, 1H), 1.28 (s, 3H), 1.60 (q, J = 5.5 Hz, 4H), 2.05 (dd, J = 13.6, 7.3 Hz, 1H), 2.30 (m, 1H), 2.44 (d, J = 2.6 Hz, 6H), 3.10 (d, J = 11.7 Hz, 1H), 3.26 (m, 2H), 3.47 (ddd, J = 16.0, 12.4, 6.6 Hz, 2H), 3.62 (d, J = 12.7 Hz, 2H), 4.06 (dd, J = 9.2, 7.2 Hz, 1H), 5.50 (d, J = 2.5 Hz, 1H), 6.64 (q, J = 6.7 Hz, 1H), 7.21 (s, 2H), 7.27 (d, J = 2.3 Hz, 1H), 7.44 (dd, J = 8.5, 2.3 Hz, 1H), 7.66 (d, J = 8.5 Hz, 1H)
34bd	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (t, J = 7.7 Hz, 1H), 1.29 (d, J = 5.9 Hz, 5H), 1.61 (q, J = 5.6 Hz, 4H), 2.06 (dd, J = 13.5, 7.2 Hz, 1H), 2.33 (m, 4H), 2.84 (s, 1H), 3.12 (d, J = 11.7 Hz, 1H), 3.25 (d, J = 11.7 Hz, 1H), 3.48 (m, 2H), 3.64 (ddt, J = 15.0, 10.2, 5.1 Hz, 2H), 4.08 (dd, J = 9.2, 7.1 Hz, 1H), 5.52 (s, 1H), 6.63 (q, J = 6.8 Hz, 1H), 7.26 (m, 4H), 7.43 (dd, J = 8.5, 2.3 Hz, 1H), 7.66 (d, J = 8.5 Hz, 1H)
34be	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.58 (m, 4H), 1.99 (dd, J = 13.3, 7.1 Hz, 1H), 2.27 (dd, J = 13.3, 9.1 Hz, 1H), 2.42 (s, 3H), 3.01 (d, J = 11.6 Hz, 1H), 3.19 (d, J = 11.5 Hz, 1H), 3.47 (ddt, J = 21.2, 13.6, 6.9 Hz, 2H), 3.64 (dq, J = 12.3, 5.8 Hz, 2H), 3.98 (dd, J = 9.1, 7.1 Hz, 1H), 5.52 (s, 1H), 6.61 (q, J = 6.8 Hz, 1H), 7.18 (s, 1H), 7.31 (m, 3H), 7.46 (dd, J = 8.5, 2.3 Hz, 1H), 7.68 (d, J = 8.3 Hz, 1H)
34bf	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (t, J = 6.6 Hz, 1H), 1.29 (d, J = 7.9 Hz, 4H), 1.62 (s, 8H), 2.09 (d, J = 6.9 Hz, 1H), 2.34 (t, J = 10.9 Hz, 2H), 3.15 (d, J = 8.2 Hz, 2H), 3.25 (m, 1H), 3.32 (s, 2H), 3.48 (s, 4H), 3.54 (s, 1H), 3.66 (s, 5H), 4.12 (s, 2H), 5.56 (s, 1H), 6.60 (q, J = 6.7 Hz, 2H), 7.34 (m, 5H), 7.50 (dd, J = 8.6, 2.2 Hz, 2H), 7.68 (d, J = 8.5 Hz, 2H)
34bg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (m, 2H), 1.30 (d, J = 14.2 Hz, 7H), 1.61 (s, 12H), 2.07 (m, 3H), 2.36 (dd, J = 13.3, 8.3 Hz, 3H), 2.80 (s, 1H), 3.15 (d, J = 11.8 Hz,

	3H), 3.26 (d, J = 11.5 Hz, 3H), 3.46 (d, J = 16.1 Hz, 5H), 3.52 (d, J = 7.0 Hz, 2H), 3.64 (s, 7H), 4.18 (s, 3H), 4.92 (s, 1H), 4.98 (s, 1H), 6.58 (q, J = 6.7 Hz, 3H), 7.33 (d, J = 2.0 Hz, 3H), 7.47 (m, 12H), 7.67 (m, 6H)
34bh	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (t, J = 6.4 Hz, 1H), 1.29 (d, J = 4.7 Hz, 2H), 1.62 (q, J = 6.3, 5.4 Hz, 4H), 2.07 (dd, J = 13.4, 7.3 Hz, 1H), 2.36 (dd, J = 13.5, 9.1 Hz, 1H), 3.15 (d, J = 11.8 Hz, 1H), 3.26 (m, 1H), 3.50 (ddd, J = 20.3, 10.5, 6.5 Hz, 2H), 3.65 (m, 2H), 4.16 (t, J = 8.2 Hz, 1H), 4.68 (s, 1H), 4.95 (t, J = 11.4 Hz, 1H), 6.40 (q, J = 6.5 Hz, 1H), 7.41 (d, J = 2.2 Hz, 1H), 7.56 (dd, J = 8.5, 2.3 Hz, 1H), 7.71 (d, J = 8.6 Hz, 1H), 8.12 (s, 3H)
34bi	¹ H NMR 400 MHz, MeOH-d ₄): δ ppm 0.89 (d, J = 6.7 Hz, 1H), 1.15 (s, 1H), 1.28 (m, 9H), 1.57 (d, J = 6.4 Hz, 5H), 1.92 (dt, J = 13.9, 6.8 Hz, 1H), 2.21 (dd, J = 13.2, 9.1 Hz, 1H), 2.95 (m, 2H), 3.11 (d, J = 11.4 Hz, 1H), 3.44 (ddt, J = 20.6, 13.0, 6.2 Hz, 2H), 3.60 (dd, J = 13.8, 6.6 Hz, 2H), 3.87 (dd, J = 9.0, 7.0 Hz, 1H), 5.46 (s, 1H), 6.61 (q, J = 6.8 Hz, 1H), 7.39 (m, 6H), 7.66 (d, J = 8.5 Hz, 1H)
34bj	¹ H NMR 400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 5.0 Hz, 3H), 1.55 (m, 6H), 1.82 (dd, J = 12.9, 6.9 Hz, 1H), 2.10 (dd, J = 13.0, 9.0 Hz, 1H), 2.71 (d, J = 11.1 Hz, 1H), 3.00 (d, J = 11.1 Hz, 1H), 3.32 (s, 3H), 3.45 (tt, J = 13.0, 6.3 Hz, 3H), 3.65 (ddd, J = 18.3, 10.9, 7.1 Hz, 4H), 4.88 (s, 17H), 5.53 (s, 2H), 6.54 (q, J = 6.6 Hz, 1H), 7.32 (d, J = 2.3 Hz, 1H), 7.51 (m, 5H), 7.59 (t, J = 1.9 Hz, 1H), 7.70 (d, J = 8.5 Hz, 1H)
34bk	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.62 (q, J = 5.5 Hz, 4H), 2.07 (dd, J = 13.5, 7.3 Hz, 1H), 2.35 (dd, J = 13.6, 9.1 Hz, 1H), 3.15 (d, J = 11.8 Hz, 1H), 3.26 (d, J = 11.7 Hz, 1H), 3.49 (ddt, J = 21.5, 14.0, 6.1 Hz, 2H), 3.65 (m, 2H), 4.15 (dd, J = 9.1, 7.4 Hz, 1H), 6.49 (q, J = 6.6 Hz, 1H), 7.34 (d, J = 2.2 Hz, 1H), 7.52 (m, 2H), 7.68 (d, J = 8.5 Hz, 1H), 7.81 (s, 2H)
34bl	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.33 (m, 8H), 1.56 (q, J = 5.9, 5.2 Hz, 4H), 1.89 (dd, J = 13.1, 7.0 Hz, 1H), 2.18 (dd, J = 13.1, 9.0 Hz, 1H), 2.86 (d, J = 11.3 Hz, 1H), 3.08 (d, J = 11.4 Hz, 1H), 3.46 (ddd, J = 15.1, 12.2, 6.6 Hz, 2H), 3.61 (dd, J = 13.4, 6.0 Hz, 2H), 3.82 (dd, J = 9.0, 7.0 Hz, 1H), 4.68 (hept, J = 6.0 Hz, 1H), 5.51 (s, 1H), 6.72 (q, J = 8.2, 7.0 Hz, 2H), 6.82 (dt, J = 11.1, 2.3 Hz, 1H), 7.02 (s, 1H), 7.28 (d, J = 2.2 Hz, 1H), 7.44 (dd, J = 8.5, 2.3 Hz, 1H), 7.67 (d, J = 8.6 Hz, 1H)
34bm	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (d, J = 18.8 Hz, 11H), 1.60 (q, J = 5.8 Hz, 4H), 2.05 (dd, J = 13.5, 7.2 Hz, 1H), 2.33 (m, 2H), 2.43 (s, 3H), 3.12 (d, J = 11.7 Hz, 1H), 3.25 (d, J = 11.7 Hz, 1H), 3.45 (ddt, J = 21.7, 13.4, 6.5 Hz, 2H), 3.62 (dq, J = 11.4, 5.5 Hz, 2H), 4.09 (dd, J = 9.2, 7.2 Hz, 1H), 6.64 (q, J = 6.8 Hz, 1H), 7.05 (s, 1H), 7.26 (m, 2H), 7.34 (d, J = 1.8 Hz, 1H), 7.42 (dd, J = 8.5, 2.3 Hz, 1H), 7.66 (d, J = 8.5 Hz, 1H)
34bm	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 2H), 1.53 (t, J = 6.1 Hz, 4H), 1.80 (dd, J = 13.0, 6.9 Hz, 1H), 2.09 (dd, J = 13.0, 8.9 Hz, 1H), 2.36 (d, J = 2.0 Hz, 3H), 2.68 (d, J = 10.8 Hz, 1H), 2.98 (d, J = 11.1 Hz, 1H), 3.42 (m, 2H), 3.63 (m, 3H), 5.49 (s, 1H), 6.64 (q, J = 6.8 Hz, 1H), 7.24 (m, 3H), 7.43 (m, 2H), 7.67 (d, J = 8.5 Hz, 1H)
34bo	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (d, J = 13.0 Hz, 1H), 1.53 (m, 4H), 1.79 (dd, J = 12.9, 6.9 Hz, 1H), 2.08 (dd, J = 12.9, 8.9 Hz, 1H), 2.66 (d, J = 11.1 Hz, 1H), 2.97 (d, J = 11.0 Hz, 1H), 3.42 (m, 3H), 3.62 (m, 3H), 5.50 (d, J = 15.0 Hz, 1H), 6.53 (q, J = 6.9 Hz, 1H), 7.36 (d, J = 2.3 Hz, 1H), 7.53 (dd, J = 8.4, 2.2 Hz, 1H), 7.63 (dd, J = 8.0, 5.0 Hz, 1H), 7.72 (d, J = 8.5 Hz, 1H), 7.98 (m, 1H), 8.67 (m, 1H), 8.74 (s, 1H)
34bp	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.41 (t, J = 6.9 Hz, 3H), 1.65 (dt, J = 10.9, 5.8 Hz, 4H), 2.08 (dd, J = 13.6, 8.2 Hz, 1H), 2.44 (dd, J = 13.6, 8.9 Hz, 1H), 3.24 (m, 2H),


	3.57 (m, 4H), 4.13 (qd, J = 7.0, 4.1 Hz, 2H), 4.42 (t, J = 8.5 Hz, 1H), 6.66 (q, J = 6.7 Hz, 1H), 7.00 (s, 1H), 7.16 (d, J = 8.0 Hz, 1H), 7.28 (m, 2H), 7.46 (dd, J = 8.5, 2.3 Hz, 1H), 7.67 (d, J = 8.5 Hz, 1H)
34bq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.36 (s, 1H), 1.59 (q, J = 5.8, 4.7 Hz, 0H), 2.05 (dd, J = 13.4, 7.2 Hz, 0H), 2.31 (m, 0H), 3.12 (d, J = 11.6 Hz, 0H), 3.24 (d, J = 11.7 Hz, 0H), 3.42 (d, J = 9.4 Hz, 0H), 3.49 (m, 0H), 3.59 (d, J = 12.4 Hz, 0H), 4.09 (dd, J = 9.1, 7.1 Hz, 0H), 4.90 (s, 2H), 5.44 (s, 0H), 6.65 (q, J = 7.0 Hz, 0H), 7.27 (m, 0H), 7.45 (m, 0H), 7.54 (s, 0H), 7.69 (m, 0H)
34br	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 7.7 Hz, 3H), 1.59 (m, 6H), 2.04 (dd, J = 13.5, 7.0 Hz, 1H), 2.19 (s, 2H), 2.31 (dd, J = 13.5, 9.2 Hz, 1H), 3.11 (d, J = 11.5 Hz, 1H), 3.24 (d, J = 11.8 Hz, 1H), 3.34 (s, 1H), 3.47 (dt, J = 24.1, 8.0 Hz, 2H), 3.63 (m, 2H), 4.07 (t, J = 7.9 Hz, 1H), 4.73 (s, 1H), 5.16 (m, 1H), 5.47 (d, J = 10.8 Hz, 2H), 6.62 (q, J = 6.8 Hz, 1H), 7.33 (m, 1H), 7.51 (m, 3H), 7.68 (d, J = 8.6 Hz, 1H)
34bs	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.60 (q, J = 5.9, 5.4 Hz, 4H), 2.04 (m, 1H), 2.31 (dd, J = 13.5, 9.2 Hz, 1H), 3.11 (s, 7H), 3.24 (d, J = 11.6 Hz, 1H), 3.47 (m, 2H), 3.63 (s, 2H), 4.06 (t, J = 8.1 Hz, 1H), 5.50 (s, 1H), 6.69 (m, 3H), 7.34 (d, J = 2.2 Hz, 1H), 7.48 (dd, J = 8.6, 2.3 Hz, 1H), 7.69 (d, J = 8.4 Hz, 1H), 8.18 (d, J = 5.2 Hz, 1H)
34bt	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.56 (t, J = 5.4 Hz, 4H), 1.99 (dd, J = 13.3, 7.0 Hz, 1H), 2.26 (dd, J = 13.3, 9.1 Hz, 1H), 3.01 (d, J = 11.5 Hz, 1H), 3.18 (d, J = 11.5 Hz, 1H), 3.43 (ddt, J = 20.5, 13.2, 6.0 Hz, 2H), 3.60 (dd, J = 13.4, 5.7 Hz, 2H), 3.98 (dd, J = 9.1, 7.0 Hz, 1H), 4.85 (m, 1H), 5.47 (s, 1H), 6.68 (q, J = 6.7 Hz, 1H), 7.39 (d, J = 2.3 Hz, 1H), 7.54 (m, 4H), 7.72 (d, J = 8.5 Hz, 1H), 7.97 (m, 4H)
34bu	¹ H NMR (400 MHz, CDCl ₃): δ ppm 1.36 (dd, J = 6.9, 3.7 Hz, 6H), 1.73 (dd, J = 13.1, 6.7 Hz, 1H), 2.05 (dd, J = 13.1, 8.8 Hz, 1H), 2.81 (d, J = 10.5 Hz, 1H), 2.94 (d, J = 10.5 Hz, 1H), 3.14 (p, J = 6.9 Hz, 1H), 3.47 (dt, J = 12.2, 5.6 Hz, 4H), 3.85 (dd, J = 8.8, 6.7 Hz, 1H), 4.19 (q, J = 7.1 Hz, 2H), 4.34 (s, 2H), 5.42 (s, 1H), 6.53 (q, J = 6.7 Hz, 1H), 7.25 (m, 3H), 7.42 (dd, J = 8.5, 2.2 Hz, 1H), 7.68 (d, J = 8.5 Hz, 1H), 8.65 (dd, J = 5.0, 0.8 Hz, 1H)
34bv	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (d, J = 13.8 Hz, 1H), 1.54 (q, J = 6.0, 5.4 Hz, 4H), 1.80 (dd, J = 13.0, 6.9 Hz, 1H), 2.08 (dd, J = 12.8, 8.9 Hz, 1H), 2.66 (d, J = 11.2 Hz, 1H), 2.97 (d, J = 11.1 Hz, 1H), 3.38 (m, 3H), 3.61 (dt, J = 13.8, 7.1 Hz, 3H), 5.49 (d, J = 1.5 Hz, 1H), 6.60 (q, J = 6.7 Hz, 1H), 7.28 (m, 3H), 7.47 (m, 3H), 7.66 (d, J = 8.4 Hz, 1H)
34bw	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.63 (d, J = 7.6 Hz, 4H), 2.07 (dd, J = 13.4, 7.6 Hz, 1H), 2.39 (dd, J = 13.6, 9.0 Hz, 1H), 3.17 (d, J = 11.9 Hz, 1H), 3.28 (m, 2H), 3.51 (dt, J = 23.6, 8.6 Hz, 2H), 3.62 (d, J = 14.5 Hz, 2H), 4.25 (t, J = 8.4 Hz, 1H), 6.60 (q, J = 6.6 Hz, 1H), 7.29 (d, J = 2.3 Hz, 1H), 7.51 (m, 6H), 7.66 (d, J = 8.7 Hz, 1H)
34bx	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.61 (m, 4H), 2.06 (dd, J = 13.5, 7.5 Hz, 1H), 2.39 (m, 4H), 3.15 (d, J = 11.8 Hz, 1H), 3.26 (d, J = 11.7 Hz, 1H), 3.47 (m, 2H), 3.62 (ddd, J = 15.6, 9.4, 5.2 Hz, 2H), 4.18 (dd, J = 9.1, 7.4 Hz, 1H), 6.64 (q, J = 6.7 Hz, 1H), 7.26 (d, J = 2.3 Hz, 1H), 7.38 (m, 4H), 7.65 (d, J = 8.5 Hz, 1H)
34by	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm: 1.30 (d, J = 17.9 Hz, 1H), 1.58 (q, J = 4.3, 2.7 Hz, 4H), 1.78 (m, 4H), 1.98 (dd, J = 13.3, 7.1 Hz, 1H), 2.24 (m, 4H), 2.41 (m, 1H), 3.02 (d, J = 11.6 Hz, 1H), 3.18 (d, J = 11.6 Hz, 1H), 3.47 (m, 2H), 3.62 (dq, J = 12.8, 5.9 Hz, 2H), 3.98 (dd, J = 9.1, 7.0 Hz, 1H), 5.49 (s, 1H), 5.75 (q, J = 2.6, 1.7 Hz, 1H), 6.94 (q, J = 6.9 Hz, 1H), 7.15 (d, J = 2.3 Hz, 1H), 7.29 (dd, J = 8.5, 2.3 Hz, 1H), 7.58 (d, J = 8.5 Hz,

	1H)
34bz	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.02 (m, 7H), 1.28 (d, J = 5.2 Hz, 1H), 1.60 (q, J = 6.1, 5.6 Hz, 4H), 2.06 (ddt, J = 14.1, 11.3, 6.7 Hz, 2H), 2.33 (dd, J = 13.4, 9.2 Hz, 1H), 3.12 (d, J = 11.7 Hz, 1H), 3.27 (d, J = 25.3 Hz, 3H), 3.47 (ddt, J = 20.4, 13.1, 5.7 Hz, 2H), 3.65 (m, 2H), 3.81 (m, 2H), 4.08 (dd, J = 9.1, 7.3 Hz, 1H), 5.51 (s, 1H), 6.71 (q, J = 6.8 Hz, 1H), 7.01 (m, 2H), 7.18 (s, 1H), 7.29 (d, J = 2.1 Hz, 1H), 7.43 (m, 2H), 7.66 (d, J = 8.6 Hz, 1H)
34ca	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.00 (t, J = 7.3 Hz, 1H), 1.28 (m, 1H), 1.60 (m, 5H), 1.97 (m, 5H), 2.30 (t, J = 11.2 Hz, 1H), 2.83 (t, J = 7.4 Hz, 1H), 3.07 (d, J = 11.5 Hz, 1H), 3.22 (d, J = 11.4 Hz, 1H), 3.54 (m, 8H), 4.04 (d, J = 8.7 Hz, 1H), 5.08 (s, 1H), 5.56 (s, 1H), 6.69 (q, J = 6.6 Hz, 1H), 7.31 (d, J = 2.1 Hz, 1H), 7.48 (m, 2H), 7.65 (m, 3H), 7.93 (s, 1H)
34cb	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (m, 1H), 1.62 (m, 6H), 1.90 (m, 8H), 2.32 (dd, J = 13.4, 9.1 Hz, 1H), 3.11 (d, J = 11.7 Hz, 1H), 3.25 (d, J = 11.6 Hz, 1H), 3.47 (ddt, J = 21.4, 13.3, 6.4 Hz, 2H), 3.65 (dq, J = 13.0, 6.2 Hz, 2H), 4.08 (dd, J = 9.1, 7.1 Hz, 1H), 5.51 (s, 1H), 6.72 (q, J = 6.9 Hz, 1H), 6.94 (d, J = 7.7 Hz, 1H), 7.02 (dd, J = 8.2, 2.6 Hz, 1H), 7.18 (s, 1H), 7.28 (d, J = 2.3 Hz, 1H), 7.42 (m, 2H), 7.66 (d, J = 8.5 Hz, 1H)
34cc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (m, 1H), 1.49 (m, 8H), 2.02 (m, 5H), 2.33 (dd, J = 13.3, 9.0 Hz, 1H), 3.13 (d, J = 11.6 Hz, 1H), 3.25 (d, J = 12.3 Hz, 1H), 3.58 (ddd, J = 32.1, 26.0, 15.3 Hz, 5H), 3.88 (td, J = 10.6, 10.2, 3.9 Hz, 1H), 4.08 (t, J = 8.1 Hz, 1H), 5.56 (s, 1H), 6.63 (q, J = 6.7 Hz, 1H), 7.29 (d, J = 2.2 Hz, 1H), 7.56 (m, 4H), 7.89 (d, J = 7.7 Hz, 1H), 8.34 (s, 1H)
34cd	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (q, J = 7.6, 6.7 Hz, 4H), 1.57 (p, J = 3.8 Hz, 4H), 1.99 (dd, J = 13.3, 7.1 Hz, 1H), 2.27 (dd, J = 13.3, 9.1 Hz, 1H), 2.73 (q, J = 7.6 Hz, 2H), 3.01 (d, J = 11.5 Hz, 1H), 3.18 (d, J = 11.6 Hz, 1H), 3.45 (ddt, J = 21.2, 13.1, 5.9 Hz, 2H), 3.60 (dt, J = 12.5, 6.8 Hz, 2H), 3.98 (dd, J = 9.1, 7.1 Hz, 1H), 4.93 (s, 1H), 5.47 (s, 1H), 6.64 (q, J = 6.8 Hz, 1H), 7.30 (m, 4H), 7.43 (m, 2H), 7.66 (d, J = 8.5 Hz, 1H)
34ce	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (m, 9H), 1.52 (ddd, J = 11.5, 7.0, 4.8 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.75 (d, J = 10.9 Hz, 1H), 2.95 (m, 2H), 3.50 (m, 4H), 3.83 (dd, J = 8.8, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.5 Hz, 2H), 4.92 (s, 8H), 5.47 (d, J = 14.2 Hz, 1H), 6.61 (q, J = 6.8 Hz, 1H), 7.32 (m, 4H), 7.44 (m, 2H), 7.66 (d, J = 8.5 Hz, 1H)
34cf	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.09 (s, 0H), 0.89 (t, J = 6.5 Hz, 0H), 1.31 (d, J = 12.8 Hz, 1H), 1.59 (m, 0H), 1.84 (s, 0H), 2.02 (d, J = 6.4 Hz, 0H), 2.19 (t, J = 7.8 Hz, 0H), 2.65 (s, 0H), 2.76 (t, J = 6.7 Hz, 0H), 2.87 (d, J = 14.6 Hz, 0H), 3.06 (s, 0H), 3.30 (s, 1H), 3.49 (m, 0H), 3.61 (m, 0H), 3.82 (s, 0H), 4.98 (s, 0H), 5.33 (m, 0H), 5.55 (s, 0H), 7.30 (s, 0H), 7.38 (d, J = 7.9 Hz, 0H), 7.46 (t, J = 7.3 Hz, 0H), 7.54 (t, J = 7.8 Hz, 0H), 7.64 (m, 0H), 7.81 (t, J = 8.6 Hz, 0H), 7.93 (m, 0H), 8.40 (s, 0H)
34cg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.73 (p, J = 7.2, 6.3 Hz, 4H), 2.10 (dd, J = 13.7, 8.5 Hz, 1H), 2.50 (dd, J = 13.6, 8.9 Hz, 1H), 3.26 (m, 4H), 3.61 (m, 11H), 3.78 (d, J = 16.7 Hz, 4H), 4.53 (t, J = 8.7 Hz, 1H), 6.61 (m, 1H), 7.36 (d, J = 2.2 Hz, 1H), 7.58 (m, 6H)
34ch	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.43 (m, 4H), 1.79 (dd, J = 13.3, 7.5 Hz, 1H), 2.13 (m, 1H), 2.30 (s, 3H), 2.48 (m, 3H), 2.95 (d, J = 11.8 Hz, 1H), 3.09 (m, 1H), 3.38 (s, 1H), 3.44 (s, 6H), 3.66 (s, 2H), 3.82 (t, J = 8.3 Hz, 1H), 5.54 (s, 1H), 6.57 (q, J = 6.8 Hz,

	1H), 7.34 (d, J = 2.1 Hz, 1H), 7.55 (m, 6H)
34ci	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 4.2 Hz, 2H), 1.59 (m, 5H), 1.99 (dd, J = 13.4, 6.9 Hz, 1H), 2.27 (dd, J = 13.3, 8.9 Hz, 1H), 2.80 (s, 3H), 3.01 (d, J = 11.5 Hz, 1H), 3.18 (d, J = 11.4 Hz, 1H), 3.48 (m, 3H), 3.62 (q, J = 6.8, 5.6 Hz, 2H), 3.98 (t, J = 8.0 Hz, 1H), 5.52 (s, 1H), 6.75 (q, J = 6.7 Hz, 1H), 7.49 (m, 3H), 7.69 (d, J = 8.5 Hz, 1H), 7.76 (s, 1H)
34cj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (m, 1H), 1.58 (s, 8H), 2.06 (dd, J = 10.2, 6.4 Hz, 2H), 2.30 (m, 2H), 3.12 (s, 2H), 3.23 (d, J = 9.6 Hz, 2H), 3.49 (m, 4H), 3.64 (s, 9H), 4.07 (t, J = 7.9 Hz, 2H), 5.64 (s, 1H), 6.24 (d, J = 7.2 Hz, 2H), 6.50 (t, J = 6.8 Hz, 2H), 7.31 (d, J = 2.2 Hz, 2H), 7.46 (dd, J = 14.7, 7.7 Hz, 3H), 7.65 (d, J = 8.5 Hz, 2H), 7.78 (dd, J = 12.8, 6.1 Hz, 2H)
34ck	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.62 (q, J = 6.1, 5.6 Hz, 4H), 2.06 (dd, J = 13.4, 7.2 Hz, 1H), 2.34 (dd, J = 13.5, 9.2 Hz, 1H), 2.58 (s, 3H), 3.13 (d, J = 11.7 Hz, 1H), 3.26 (d, J = 11.7 Hz, 2H), 3.51 (m, 2H), 3.68 (td, J = 14.8, 14.3, 7.0 Hz, 2H), 4.08 (dd, J = 9.2, 7.1 Hz, 1H), 4.87 (d, J = 7.3 Hz, 1H), 4.97 (s, 1H), 5.60 (s, 1H), 6.65 (q, J = 6.6 Hz, 1H), 7.34 (d, J = 2.3 Hz, 1H), 7.50 (dd, J = 8.6, 2.2 Hz, 1H), 7.67 (dd, J = 12.0, 8.0 Hz, 2H), 7.77 (t, J = 7.8 Hz, 1H), 7.94 (dt, J = 7.9, 1.4 Hz, 1H), 8.31 (s, 1H)
34cl	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.61 (d, J = 5.5 Hz, 5H), 2.04 (dd, J = 13.3, 7.1 Hz, 1H), 2.31 (dd, J = 13.4, 9.2 Hz, 1H), 2.73 (s, 6H), 3.08 (d, J = 11.6 Hz, 1H), 3.23 (d, J = 11.7 Hz, 1H), 3.53 (m, 2H), 3.68 (d, J = 14.2 Hz, 2H), 4.04 (dd, J = 9.1, 7.0 Hz, 1H), 5.62 (s, 1H), 6.69 (q, J = 6.6 Hz, 1H), 7.36 (d, J = 2.3 Hz, 1H), 7.50 (dd, J = 8.5, 2.3 Hz, 1H), 7.70 (dd, J = 12.4, 7.7 Hz, 2H), 7.86 (m, 2H), 8.33 (s, 1H)
34cm	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (s, 1H), 1.57 (p, J = 7.6, 6.8 Hz, 4H), 1.86 (dd, J = 13.0, 6.9 Hz, 1H), 2.15 (dd, J = 13.2, 9.0 Hz, 1H), 2.81 (d, J = 11.2 Hz, 1H), 2.95 (s, 4H), 3.05 (d, J = 11.2 Hz, 1H), 3.32 (s, 1H), 3.46 (ddt, J = 17.4, 13.1, 5.7 Hz, 2H), 3.62 (dq, J = 11.5, 5.5 Hz, 2H), 3.76 (dd, J = 9.0, 6.9 Hz, 1H), 5.54 (s, 1H), 6.63 (q, J = 6.7 Hz, 1H), 7.29 (d, J = 2.3 Hz, 1H), 7.46 (dd, J = 8.5, 2.3 Hz, 1H), 7.62 (m, 3H), 7.89 (dt, J = 7.7, 1.5 Hz, 1H), 8.36 (s, 1H)
34cn	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 2H), 1.62 (q, J = 5.7, 5.0 Hz, 12H), 2.07 (dd, J = 13.4, 7.1 Hz, 3H), 2.35 (dd, J = 13.4, 9.1 Hz, 3H), 3.09 (d, J = 26.8 Hz, 23H), 3.26 (s, 2H), 3.53 (m, 6H), 3.64 (d, J = 13.0 Hz, 7H), 4.15 (s, 3H), 4.88 (d, J = 3.3 Hz, 1H), 4.97 (s, 1H), 5.56 (s, 1H), 6.71 (q, J = 6.7 Hz, 3H), 7.32 (d, J = 2.2 Hz, 3H), 7.56 (m, 15H), 7.78 (s, 3H)
34co	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.13 (t, J = 7.1 Hz, 3H), 1.26 (m, 4H), 1.61 (q, J = 6.1, 5.6 Hz, 4H), 2.05 (dd, J = 13.4, 7.2 Hz, 1H), 2.33 (dd, J = 13.4, 9.3 Hz, 1H), 3.13 (d, J = 11.7 Hz, 1H), 3.47 (m, 10H), 4.09 (t, J = 8.3 Hz, 1H), 5.55 (s, 1H), 6.74 (q, J = 6.8 Hz, 1H), 7.32 (d, J = 2.2 Hz, 1H), 7.48 (m, 3H), 7.65 (m, 3H)
34cp	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (m, 7H), 1.58 (d, J = 13.6 Hz, 14H), 2.05 (m, 3H), 2.31 (s, 4H), 2.88 (s, 1H), 3.11 (d, J = 12.1 Hz, 3H), 3.25 (d, J = 12.8 Hz, 3H), 3.38 (s, 10H), 3.48 (s, 3H), 3.63 (m, 5H), 4.09 (t, J = 8.2 Hz, 3H), 4.48 (s, 2H), 4.98 (s, 3H), 5.10 (s, 1H), 5.42 (s, 2H), 5.54 (s, 2H), 6.50 (d, J = 13.3 Hz, 2H), 6.79 (m, 1H), 7.22 (s, 2H), 7.44 (s, 5H), 7.53 (d, J = 8.4 Hz, 5H), 7.76 (s, 7H), 8.11 (m, 3H)
34cq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.31 (s, 3H), 1.62 (s, 8H), 2.08 (dd, J = 13.3, 7.1 Hz, 2H), 2.35 (t, J = 11.4 Hz, 2H), 3.16 (d, J = 11.8 Hz, 2H), 3.32 (m, 23H), 3.49 (s, 4H), 3.63 (d, J = 19.6 Hz, 3H), 3.83 (s, 4H), 3.90 (s, 2H), 3.97 (s, 2H), 4.05 (s, 1H), 4.13 (t, J = 7.7 Hz, 2H), 6.67 (m, 2H), 7.35 (s, 2H), 7.51 (d, J = 8.7 Hz, 2H), 7.66 (dq, J = 31.2,

	9.5, 9.1 Hz, 7H), 7.84 (s, 2H)
34cr	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.46 (m, 4H), 1.28 (s, 1H), 1.58 (s, 4H), 1.69 (d, J = 7.3 Hz, 1H), 1.97 (d, J = 9.1 Hz, 1H), 2.24 (dd, J = 13.1, 9.0 Hz, 1H), 2.58 (s, 2H), 2.73 (s, 2H), 2.96 (d, J = 11.3 Hz, 1H), 3.15 (d, J = 11.6 Hz, 1H), 3.32 (m, 1H), 3.48 (t, J = 12.1 Hz, 4H), 3.63 (s, 2H), 3.76 (s, 2H), 3.92 (t, J = 8.1 Hz, 1H), 5.55 (s, 1H), 6.71 (q, J = 6.7 Hz, 1H), 7.33 (d, J = 2.2 Hz, 1H), 7.50 (m, 3H), 7.66 (m, 2H), 7.80 (s, 1H)
34cl	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.71 (d, J = 4.7 Hz, 1H), 8.02 (t, J = 7.9 Hz, 1H), 7.72 (t, J = 7.3 Hz, 2H), 7.52 (d, J = 10.9 Hz, 3H), 6.92 (d, J = 6.7 Hz, 1H), 5.87 (s, 1H), 4.80 (s, 7H), 4.10 (d, J = 8.7 Hz, 1H), 3.66 (s, 2H), 3.50 (s, 2H), 3.30 (s, 5H), 3.25 (d, J = 11.8 Hz, 1H), 3.12 (d, J = 11.7 Hz, 1H), 2.35 – 2.27 (m, 1H), 2.06 (dd, J = 13.3, 7.1 Hz, 1H), 1.59 (s, 3H), 1.59 (d, J = 11.4 Hz, 1H)
34cm	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.99 (d, J = 4.9 Hz, 2H), 8.03 (s, 1H), 7.75 (d, J = 9.4 Hz, 2H), 7.60 – 7.49 (m, 2H), 5.71 (s, 1H), 4.10 (s, 1H), 3.59 (d, J = 18.5 Hz, 2H), 3.30 (d, J = 3.1 Hz, 9H), 3.11 (d, J = 12.0 Hz, 1H), 2.31 (t, J = 11.6 Hz, 1H), 2.06 (s, 1H), 1.57 (s, 5H), 1.28 (s, 1H)
34cu	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.96 (d, J = 1.5 Hz, 1H), 8.83 – 8.77 (m, 1H), 8.71 (d, J = 2.6 Hz, 1H), 7.77 (d, J = 8.4 Hz, 1H), 7.64 – 7.55 (m, 2H), 6.87 (q, J = 6.7 Hz, 1H), 5.64 (s, 1H), 3.99 (t, J = 8.2 Hz, 1H), 3.46 (s, 1H), 3.19 (d, J = 11.6 Hz, 1H), 3.03 (d, J = 11.6 Hz, 1H), 2.27 (dd, J = 13.3, 9.2 Hz, 1H), 2.00 (dd, J = 13.4, 7.0 Hz, 1H), 1.57 (s, 3H), 1.29 (s, 1H).
34cv	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.67 (d, J = 8.5 Hz, 2H), 7.44 (ddd, J = 8.0, 4.8, 2.6 Hz, 3H), 7.32 – 7.23 (m, 3H), 7.07 (dd, J = 8.4, 2.6 Hz, 2H), 6.98 (d, J = 7.6 Hz, 1H), 6.76 (d, J = 7.0 Hz, 2H), 4.26 – 4.05 (m, 5H), 3.75 (t, J = 4.7 Hz, 3H), 3.42 (s, 4H), 3.36 (s, 1H), 3.26 (d, J = 11.7 Hz, 3H), 3.13 (d, J = 11.7 Hz, 2H), 2.33 (dd, J = 13.5, 9.1 Hz, 2H), 2.06 (dd, J = 13.4, 7.1 Hz, 2H), 1.61 (d, J = 5.6 Hz, 5H).


Example 35: (S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(ethoxycarbonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



- 5 The title compound was prepared as described for (S)-8-(2-amino-6-((R)-1-(2'-(ethoxycarbonyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 20) starting with (S)-8-(2-amino-6-((R)-1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid.

¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.29 - 1.38 (m, 3 H) 1.47 - 1.72 (m, 4 H) 1.91 (dd, J=13.28, 9.18 Hz, 1 H) 2.35 (dd, J=13.25, 8.61 Hz, 1 H) 3.14 (br. s., 2 H) 3.65 (br. s., 4 H) 4.30 - 4.40 (m, 2 H) 4.40 - 4.50 (m, 1 H) 5.90 (br. s., 1 H) 6.59 (q, J=6.67 Hz, 1 H) 7.11 (br. s., 1 H) 7.44 (t, J=1.22 Hz, 1 H) 7.66 (s, 2 H) 7.70 - 7.79 (m, 2 H) 8.08 (dt, J=6.37, 2.14 Hz, 1 H) 8.14 (br. s., 1 H) 8.98 (d, J=5.61 Hz, 1 H) 10.36 (d, J=5.08 Hz, 1 H). LCMS (MH⁺): 634.


Example 36: (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(2-methoxyethoxy)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 10d) starting with (R)-1-(4-bromo-2-(2-methoxyethoxy)phenyl)-2,2,2-trifluoroethanol and obtained as a white solid.

¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.44 - 1.66 (m, 4 H) 1.83 - 1.95 (m, 1 H) 2.34 (dd, J=13.08, 8.79 Hz, 1 H) 3.14 (br. s., 2 H) 3.33 (s, 3 H) 3.42 - 3.65 (m, 4 H) 3.67 - 3.79 (m, 2 H) 4.19 - 4.27 (m, 1 H) 4.27 - 4.36 (m, 1 H) 4.48 (t, J=6.49 Hz, 1 H) 5.74 (s, 1 H) 6.99 (q, J=6.78 Hz, 1 H) 7.07 - 7.16 (m, 1 H) 7.27 (s, 1 H) 7.43 (d, J=8.35 Hz, 1 H) 8.93 (d, J=5.42 Hz, 1 H) 9.81 (br. s., 1 H). LCMS (MH⁺): 560.

Example 36b: (S)-8-(6-((R)-1-(2-(1H-benzo[d]imidazol-1-yl)-4-chlorophenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




Step 1: To a solution of (R)-1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethanol (1 g, 3.5 mmol) and 1H-benzo[d]imidazole (408 mg, 3.5 mmol) in toluene (24 mL) was added sequentially, CuI (131 mg, 0.69 mmol), K₂CO₃ (1.19 g, 8.63 mmol), and (1R,2R)-N1,N2-dimethylcyclohexane-1,2-diamine (196 mg, 1.38 mmol). The reaction mixture was purged with N₂ and then heated at 130 °C in a sealed tube for 12 h. Afterward, the reaction was cooled to RT. The solid was removed by filtration and the filtrate was concentrated and purified by flash column (EtOAc in hexane = 0 to 50 %) to afford -(R)-1-(2-(1H-benzo[d]imidazol-1-yl)-4-chlorophenyl)-2,2,2-trifluoroethanol as a white solid.

Steps 2-5: The title compound was made as described for Example 10d (Steps 1-4) to provide a white solid.

¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.59 (m, 4H), 2.05 (dt, J = 13.7, 6.9 Hz, 1H), 2.33 (dt, J = 14.5, 8.5 Hz, 1H), 3.13 (dd, J = 11.7, 7.6 Hz, 1H), 3.26 (m, 2H), 3.49 (m, 3H), 3.63 (m, 2H), 4.10 (q, J = 7.0, 5.2 Hz, 1H), 5.48 (d, J = 3.9 Hz, 1H), 6.43 (p, J = 6.4 Hz, 1H), 7.22 (dd, J = 7.8, 4.0 Hz, 1H), 7.38 (m, 2H), 7.61 (dd, J = 5.3, 2.2 Hz, 1H), 7.81 (m, 3H), 8.54 (s, 1H). LCMS (MH⁺): 603.

Example 36c: (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(1H-indazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




The title compound was prepared as described for (S)-8-(6-((R)-1-(2-(1H-benzo[d]imidazol-1-yl)-4-chlorophenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 36b) starting with 1H-indazole and obtained as a white solid.

5

¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.57 (m, 5H), 2.05 (dd, J = 13.4, 7.1 Hz, 1H), 2.32 (dd, J = 13.5, 9.2 Hz, 1H), 3.12 (d, J = 11.7 Hz, 1H), 3.24 (d, J = 11.7 Hz, 1H), 3.52 (dddd, J = 44.5, 25.8, 14.0, 7.1 Hz, 5H), 4.13 (dd, J = 9.1, 7.1 Hz, 1H), 4.92 (s, 1H), 6.68 (q, J = 6.5 Hz, 1H), 7.31 (t, J = 7.4 Hz, 1H), 7.46 (m, 2H), 7.72 (m, 5H), 8.39 (s, 1H). LCMS (MH⁺): 603.

10

Example 36d: (S)-8-(2-amino-6-((R)-1-(4-bromo-2-(piperazin-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: A mixture of 4-bromo-2-fluorobenzoic acid (2 g, 9.1 mmol), benzyl piperazine-1-carboxylate (2.4 g, 10.9 mmol) and K₂CO₃ (2.5 g, 18.26 mmol) in DMF (40 mL) was stirred at 150 °C for 36 h. The reaction was then cooled to RT and extracted with ethyl acetate, 3 N HCl, brine, dried over Na₂SO₄, filtered and concentrated *in vacuo* to provide 2-(4-((benzyloxy)carbonyl)piperazin-1-yl)-4-bromobenzoic acid as yellow oil that was used without further purification.


20

Step 2: To a mixture of 2-(4-((benzyloxy) carbonyl)piperazin-1-yl)-4-bromobenzoic acid (2 g, 9.1 mmol) in THF (20 mL) was added dropwise BH₃/THF (1.0 M, 40 mL) at 0 °C. The mixture was refluxed for 2 h, then cooled to RT, quenched with H₂O, and extracted with ethyl acetate, 3 N HCl, brine, then dried over Na₂SO₄, filtered and concentrated. Purification by normal phase silica gel (ethyl acetate/hexanes) provided benzyl 4-(5-bromo-2-(hydroxymethyl)phenyl)piperazine-1-carboxylate as a white solid.

Steps 3-10: The title compound was prepared as described for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(5-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 54d) following Steps 4-11.

¹H NMR (MeOH-d₄): δ ppm 0.90 (dt, J = 16.0, 8.0 Hz, 1H), 1.31 (s, 2H), 1.62 (t, J = 5.6 Hz, 5H), 2.03 (dd, J = 13.6, 6.9 Hz, 1H), 2.30 (dd, J = 13.4, 9.1 Hz, 1H), 2.76 (dd, J = 10.1, 6.3 Hz, 2H), 3.08 (m, 8H), 3.22 (d, J = 11.6 Hz, 1H), 3.47 (s, 1H), 3.54 (m, 1H), 3.65 (dd, J = 13.9, 6.8 Hz, 2H), 4.01 (t, J = 8.0 Hz, 1H), 5.56 (s, 1H), 7.31 (q, J = 6.9 Hz, 1H), 7.41 (dd, J = 8.4, 1.9 Hz, 1H), 7.50 (m, 2H). LCMS (MH⁺): 615.


Example 36e: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-(piperazin-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared starting with (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(2-(4-((benzyloxy)carbonyl)piperazin-1-yl)-4-bromophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (intermediate from Step 8, Example 36d) via a Suzuki coupling with (4-isopropoxyphenyl)boronic acid as as described for example 54b.

¹H NMR (MeOH-d₄): δ ppm 0.90 (m, 1H), 1.33 (m, 8H), 1.40 (s, 1H), 1.59 (q, J = 5.7 Hz, 4H), 2.06 (dd, J = 13.7, 7.0 Hz, 1H), 2.31 (dd, J = 13.5, 9.2 Hz, 1H), 3.11 (m, 3H), 3.26 (d, J = 11.7 Hz, 1H), 3.51 (m, 10H), 4.09 (dd, J = 9.3, 6.8 Hz, 1H), 4.64 (p, J = 6.0 Hz, 1H), 5.56 (s, 1H), 6.98 (m, 2H), 7.32 (q, J = 7.0 Hz, 1H), 7.53 (m, 4H), 7.64 (d, J = 8.2 Hz, 1H). LCMS (MH⁺): 671.


Example 36f: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-morpholino-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



10 The title compound was prepared as described for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-(piperazin-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 36e) substituting morpholine for benzyl piperazine-1-carboxylate.

15 ¹H NMR (MeOH-d₄): δ ppm 1.32 (d, J = 6.0 Hz, 7H), 1.58 (d, J = 6.0 Hz, 4H), 1.98 (m, 1H), 2.25 (dd, J = 13.3, 9.0 Hz, 1H), 2.83 (m, 2H), 2.99 (d, J = 11.5 Hz, 1H), 3.19 (m, 3H), 3.32 (s, 1H), 3.48 (ddt, J = 18.5, 8.9, 5.0 Hz, 2H), 3.62 (s, 2H), 3.92 (m, 5H), 4.63 (h, J = 6.0 Hz, 1H), 4.88 (m, 1H), 5.54 (s, 1H), 6.97 (m, 2H), 7.41 (m, 2H), 7.54 (m, 4H). LCMS (MH⁺): 672


20 **Example 36g: (S)-8-(6-((R)-1-([1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)-2-amino pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid**



The title compound was prepared as described for (S)-8-(2-amino-6-((R)-1-(5-chloro-3'-sulfamoyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 34u) starting with 1-(2-bromophenyl)-2,2,2-trifluoroethanone.

¹H NMR (MeOH-d₄): δ ppm 1.58 (d, J = 5.4 Hz, 4H), 2.00 (dd, J = 13.4, 7.1 Hz, 1H), 2.27 (dd, J = 13.3, 9.2 Hz, 1H), 3.02 (d, J = 11.6 Hz, 1H), 3.19 (d, J = 11.5 Hz, 1H), 3.30 (q, J = 1.8 Hz, 3H), 3.45 (td, J = 14.5, 6.3 Hz, 1H), 3.61 (m, 2H), 3.99 (m, 1H), 5.46 (s, 1H), 6.67 (q, J = 6.8 Hz, 1H), 7.26 (dd, J = 6.2, 2.4 Hz, 1H), 7.45 (m, 7H), 7.70 (d, J = 7.3 Hz, 1H). LCMS (MH⁺): 528.

Example 37: (3S)-8-(6-(1-((1R,3R,5S,7S)-adamantan-2-yl)ethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: A solution of adamantan-1-yl-methanol (100 mg, 0.60 mmol) in THF (5 mL) was cooled to 0 °C. 15-Crown-5 ether (99 mg, 0.5 mmol) and NaH (60% in oil, 92 mg, 2.4 mmol) were added sequentially. The reaction was warmed to RT for 1 h, cooled to 0 °C, and 4,6-dichloro-pyrimidin-2-ylamine (247 mg, 1.5 mmol) was added. The reaction was heated to 65 °C for 16 h, cooled to RT, quenched with water, and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase chromatography (EtOAc/heptane) provided 4-(adamantan-1-ylmethoxy)-6-chloro-pyrimidin-2-ylamine as a white solid.


Step 2: 4-(Adamantan-1-ylmethoxy)-6-chloro-pyrimidin-2-ylamine (89 mg, 0.30 mmol), (S)-2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (157 mg, 0.45 mmol) and NaHCO₃ (76 mg, 0.9 mmol) were dissolved in dioxane (1.5 mL) and heated to 95 °C for 64 h. Then the reaction was cooled to RT, quenched with water, and extracted with EtOAc. The organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provides (S)-2-benzyl 3-ethyl 8-(6-(adamantan-1-ylmethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

Step 3: N-CBZ Deprotection was accomplished via Method B to provide (S)-ethyl 8-(6-(adamantan-1-ylmethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid.

Step 4: Hydrolysis of (S)-ethyl 8-(6-(adamantan-1-ylmethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provides the title compound as a white solid.

¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.12 (d, J=6.25 Hz, 3 H) 1.42 - 1.76 (m, 17 H) 1.82 - 2.02 (m, 4 H) 2.34 (dd, J=13.32, 8.59 Hz, 1 H) 3.12 (br. s., 2 H) 3.67 (br. s., 4 H) 4.35 - 4.48 (m, 1 H) 5.85 (br. s., 1 H) 8.97 (br. s., 1 H) 10.44 (br. s., 1 H). LCMS (MH⁺): 456.


Example 38: (S)-8-(6-((1*r*,3*r*,5*S*,7*S*)-adamantan-2-ylmethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was made as described above for (3S)-8-(6-(1-((1r,3r,5S,7S)-adamantan-2-yl)ethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 37) using (1r,3r,5r,7r)-adamantan-2-ylmethanol.

- 5 ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.39 - 1.76 (m, 16 H) 1.83 - 2.01 (m, 4 H) 2.34 (dd, J=13.18, 8.44 Hz, 1 H) 3.13 (br. s., 2 H) 3.69 (br. s., 4 H) 3.79 (s, 2 H) 4.42 (br. s., 1 H) 5.83 (br. s., 1 H) 8.97 (br. s., 1 H) 10.40 (br. s., 1 H). LCMS (MH⁺): 442.

10 **Example 39a: 8-(4-Amino-6-((naphthalen-2-ylmethyl)amino)-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid**



- 15 *Step 1:* To a solution of 4,6-dichloro-1,3,5-triazin-2-amine (1.6 g) in isopropanol (14 mL) was added 2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2, 3-dicarboxylate (1.28 g, 3.7 mmol) and Et₃N (7 mL). The solution was heated to reflux for 72 h, then cooled to RT, and concentrated *in vacuo*. Purification by normal phase chromatography (CH₂Cl₂/MeOH = 50/1) afforded 2-benzyl 3-ethyl 8-(4-amino-6-chloro-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a colorless oil.

- 20 *Step 2:* To a solution of 2-benzyl 3-ethyl 8-(4-amino-6-chloro-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (265 mg, 0.56 mmol) in isopropanol (3 mL) were added naphthalen-2-ylmethanamine (105 mg, 0.67 mmol) and Et₃N (1.4 mL). The reaction mixture was heated to reflux for 12 h, then cooled to RT, and concentrated *in vacuo*. Purification by normal phase chromatography (CH₂Cl₂/MeOH) provided 2-benzyl 3-ethyl 8-(4-amino-6-((naphthalen-2-ylmethyl)amino)-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate
25 as a white solid.


Step 3: Hydrolysis of 2-benzyl 3-ethyl 8-(4-amino-6-((naphthalen-2-ylmethyl)amino)-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate using the LiOH general method provided 8-(4-amino-6-((naphthalen-2-ylmethyl)amino)-1,3,5-triazin-2-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid as a white solid.

5

Step 4: N-CBZ Deprotection was accomplished via Method B to provide the title compound as a white solid.


Using the generic scheme below, the following examples of Table 13a were prepared as described above for 8-(4-amino-6-((naphthalen-2-ylmethyl)amino)-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 39a).

10



15

Table 13a.



Ex. No.	A-CH(R)-NH-	CAS Name	LCMS (MH+)
39a		8-(4-amino-6-((naphthalen-2-ylmethyl)amino)-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	435
39b		8-(4-(((1,1'-biphenyl)-4-ylmethyl)amino)-6-amino-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	460
39c		8-(4-amino-6-((2-(piperidin-1-yl)benzyl)amino)-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	467
39d		8-(4-(((1,1'-biphenyl)-3-ylmethyl)amino)-6-amino-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	460
39e		8-(4-amino-6-(((R)-1-(naphthalen-2-yl)ethyl)amino)-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	448


Table 13b.

NMR Data for Compounds of Table 13a

Ex. No.	NMR
39a	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.5 (br.s. 4 H), 1.6-1.8 (m, 1H), 2.1-2.2 (m, 1H), 3.0-3.1 (br.s. 3H), 3.5-3.8 (br.s. 5 H), 4.1 (t, J = 4.8 Hz, 1 H), 4.5 (d, J = 5.5 Hz, 2 H), 6.0-6.3 (br.s. 2 H), 7.1-7.3 (m, 3H), 7.5-7.9 (m, 4 H).
39b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.54 - 1.79 (m, 4 H) 2.02 - 2.19 (m, 1 H) 2.44 - 2.60 (m, 1 H) 3.74 - 3.92 (m, 2 H) 3.93 - 4.08 (m, 2 H) 4.49 - 4.62 (m, 1 H) 4.63 - 4.71 (m, 2 H) 7.30 - 7.40 (m, 1 H) 7.40 - 7.51 (m, 4 H) 7.55 - 7.68 (m, 4 H)
39c	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.66 (br. s., 6 H) 1.86 (br. s., 4 H) 2.03 - 2.16 (m, 1 H) 2.40 - 2.54 (m, 1 H) 3.06 - 3.22 (m, 4 H) 3.66 - 3.87 (m, 2 H) 3.87 - 4.02 (m, 2

	H) 4.46 - 4.59 (m, 1 H), 4.75 (s, 2 H) 7.12 - 7.27 (m, 1 H) 7.29 - 7.45 (m, 3 H)
39d	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 - 1.79 (m, 4 H) 1.88 - 2.15 (m, 1 H) 2.25 - 2.54 (m, 1 H) 3.22 (br. s., 2 H) 3.60 - 4.01 (m, 4 H) 4.35 - 4.54 (m, 1 H) 4.62 (s, 2 H) 7.25 - 7.35 (m, 1 H) 7.36 - 7.46 (m, 3 H) 7.51 (d, J=7.61 Hz, 1 H) 7.57 (d, J=8.59 Hz, 3 H)
39e	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.63 (d, J=6.83 Hz, 9 H) 3.01 - 3.21 (m, 1 H) 3.50 - 4.07 (m, 5 H) 4.32 - 4.65 (m, 1 H) 5.14 - 5.33 (m, 1 H) 7.32 - 7.54 (m, 3 H) 7.81 (d, J=5.08 Hz, 4 H)


Example 40: 8-(4-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



- 5 *Step 1:* To a solution of (R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol (380 mg, 1.3 mmol) in 10 mL of THF was added NaH (60 mg, 1.4 mmol) and the reaction was stirred at RT for 30 min. After this time, 2-benzyl 3-ethyl 8-(4-amino-6-chloro-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (product from Step 1, Example 39a) (570 mg, 1.2 mmol) was added and the reaction was heated to 50 °C for 12 h. After this
- 10 time, the reaction was cooled to RT, quenched with methanol and concentrated in vacuo. Normal phase silica gel chromatography (EtOAc/heptane) provided 2-benzyl 3-ethyl 8-(4-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.
- 15 *Step 2:* N-CBZ Deprotection was accomplished via Method B to provide ethyl 8-(4-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid.
- Step 3:* Step 3: Hydrolysis of ethyl 8-(4-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-1,3,5-triazin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provided the title compound as a white solid.
- 20

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.55 (br. s., 4 H) 1.98 (s, 1 H) 2.02 - 2.15 (m, 1 H) 2.30 (dd, J=13.42, 9.27 Hz, 1 H) 2.36 (s, 3 H) 3.10 (d, J=11.71 Hz, 1 H) 3.23 - 3.28 (m, 1 H) 3.40 - 4.01 (m, 4 H) 4.08 (dd, J=9.27, 6.88 Hz, 1 H) 6.39 (d, J=2.25 Hz, 1 H) 7.36 - 7.63 (m, 3 H) 7.76 (d, J=8.54 Hz, 1 H) 7.91 (d, J=2.10 Hz, 1 H). LCMS (MH⁺): 567.

Example 41a: (S)-8-(2-Amino-6-((2-(piperidin-1-yl)benzyl)amino)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




Step 1: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-chloropyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (200 mg, 0.6 mmol) and [2-(1-piperidinyl)phenyl]methanamine (CAS#: 72752-54-6) (105 mg, 0.8 mmol) in *i*-PrOH (2 mL) was added diisopropylethyl amine (0.5 mL). The reaction was heated to 120 °C for 2 h followed by heating to 140 °C for 1 h under microwave conditions, then cooled to RT and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided (S)-2-benzyl 3-ethyl 8-(2-amino-6-((2-(piperidin-1-yl)benzyl)amino) pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

Step 2: N-CBZ Deprotection was accomplished via Method B to provide (S)-ethyl 8-(2-amino-6-((2-(piperidin-1-yl)benzyl)amino)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid.


Step 3: Hydrolysis of (S)-ethyl 8-(2-amino-6-((2-(piperidin-1-yl)benzyl)amino)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provided the title compound as a white solid.

Using the generic scheme below, the following examples of Table 14a were prepared as described above for (S)-8-(2-amino-6-((2-(piperidin-1-yl)benzyl)amino)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 41a).



5

Table 14a.




Ex. No.	Ar	CAS Name	LCMS (MH ⁺)
41a		(S)-8-(2-amino-6-((2-(piperidin-1-yl)benzyl)amino)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	446
41b		(S)-8-(2-amino-6-((2-phenoxy-6-(piperidin-1-yl)benzyl)amino)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	558
41c		(3S)-8-(6-(((3S,5S)-adamantan-1-ylmethyl)amino)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	441

41d		(3S)-8-(6-((1-((1R,3S,5S)-adamantan-1-yl)ethyl)amino)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	456
-----	---	--	-----

Table 14b
NMR Data for Compounds of Table 14a

Ex. No.	NMR
41a	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.39 - 1.66 (m, 6 H) 1.67 - 1.85 (m, 4 H) 1.95 - 2.11 (m, 1 H) 2.18 - 2.35 (m, 1 H) 2.69 - 2.95 (m, 4 H) 3.09 (s, 1 H) 3.20 (s, 1 H) 3.35 (s, 4 H) 3.94 - 4.14 (m, 1 H) 4.43 (s, 2 H) 6.93 - 7.05 (m, 1 H) 7.11 (s, 1 H) 7.14 - 7.24 (m, 1 H) 7.26 - 7.38 (m, 1 H)
41b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.43 - 1.66 (m, 6 H) 1.67 - 1.85 (m, 4H) 1.94 - 2.09 (m, 1 H) 2.18 - 2.34 (m, 1 H) 2.89 (d, J=4.49 Hz, 4 H) 3.07 (s, 1 H) 3.14 - 3.25 (m, 1 H) 3.32 - 3.63 (m, 4 H) 3.95 - 4.08 (m, 1 H) 4.46 (s, 2 H) 6.49 - 6.58 (m, 1 H) 6.84 - 6.97 (m, 3 H) 7.03 - 7.09(m, 1 H) 7.18 (s, 1 H) 7.28 (d, J=7.91 Hz, 2 H)
41c	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.00 (br. s., 6 H) 0.03 - 0.26 (m, 10 H) 0.27 - 0.44 (m, 9 H) 0.48 - 0.58 (m, 1 H) 0.68 - 0.85 (m, 1 H) 1.33 (s, 2 H) 1.50 - 1.64 (m, 1 H) 1.84 - 2.03 (m, 2 H) 2.11(br. s., 2 H) 2.42 - 2.62 (m, 1 H)
41d	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.08 (d, J=6.83 Hz, 3 H) 1.52 - 1.71 (m, 13 H) 1.73 (br. s., 3 H) 1.93 (s, 2 H) 1.97 (br. s., 3 H) 2.04 - 2.19 (m, 1 H) 2.24 - 2.43 (m, 1 H) 3.06 - 3.21 (m, 1 H) 3.22 - 3.28 (m, 1 H) 3.36 - 3.58 (m, 3 H) 3.59 - 3.75 (m, 2 H) 4.02 - 4.20 (m, 1 H)

5 Example 42a: (S)-8-(2-amino-6-((R)-1-(3'-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




The title compound was made as described for (S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(ethoxycarbonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 35) starting with (S)-2-benzyl 3-ethyl 8-(2-

amino-6-((R)-1-(2-bromophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate.

¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.43 (h, J = 8.5, 6.5 Hz, 4H), 1.80 (dd, J = 13.3, 7.4 Hz, 1H), 2.12 (dd, J = 13.2, 9.0 Hz, 1H), 2.48 (d, J = 1.8 Hz, 1H), 2.95 (d, J = 11.7 Hz, 1H), 3.08 (d, J = 11.7 Hz, 1H), 3.37 (d, J = 16.1 Hz, 1H), 3.48 (d, J = 11.2 Hz, 3H), 3.79 (m, 2H), 5.57 (s, 1H), 6.62 (q, J = 6.9 Hz, 1H), 7.27 (dd, J = 5.8, 3.3 Hz, 1H), 7.51 (m, 7H). LCMS (MH⁺): 563.


Example 42b: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was made as described for (S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(ethoxycarbonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 35) starting with (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(2-bromophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate.

¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.89 (m, 1H), 1.30 (d, J = 16.3 Hz, 3H), 1.60 (q, J = 5.9 Hz, 4H), 2.05 (dd, J = 13.4, 7.2 Hz, 1H), 2.32 (dd, J = 13.4, 9.1 Hz, 1H), 3.11 (d, J = 11.7 Hz, 1H), 3.24 (d, J = 11.7 Hz, 1H), 3.47 (ddt, J = 20.6, 13.4, 6.5 Hz, 2H), 3.64 (ddt, J = 15.8, 10.8, 5.2 Hz, 2H), 4.07 (dd, J = 9.2, 7.1 Hz, 1H), 5.51 (s, 1H), 6.68 (q, J = 6.9 Hz, 1H), 7.25 (m, 4H), 7.48 (m, 3H), 7.71 (m, 1H). LCMS (MH⁺): 546.

Example 43: (S)-8-(5-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridazin-3-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To (R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol (1.00 g, 3.44 mmol, Intermediate 3) in 1,4-dioxane (100 mL) was added 3,5-dichloropyridazine (512 mg, 3.44 mmol) and Cs₂CO₃ (3.36 g, 10.3 mmol). The reaction mixture was then heated at 100°C for 182 h. During this time, the reaction was charged with additional 3,5-dichloropyridazine (2.56 g, 17.2 mmol) at t = 86 h. Then the reaction mixture was cooled to RT, diluted with water, and extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification on a 120 g Isco RediSep silica cartridge (EtOAc/heptane) provided 3-chloro-5-[(1R)-1-[4-chloro-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoroethoxy]pyridazine as a 3:2 mixture of (R)-3-chloro-5-(1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridazine and (R)-5-chloro-3-(1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridazine respectively.

Step 2: To a solution of the (R)-3-chloro-5-(1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridazine/(R)-5-chloro-3-(1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridazine mixture from step 1 in 1,4-dioxane (19 mL) was added 2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (980 mg, 2.83 mmol), Cs₂CO₃ (2.30 g, 7.07 mmol), Pd₂(dba)₃ (432 mg, 0.471 mmol), and *rac*-BINAP (587 mg, 0.940 mmol), and the reaction mixture was heated to 60 °C for 60 h. Then the reaction mixture was cooled to RT, filtered through celite, washed with EtOAc, and the filtrate concentrated *in vacuo*. Purification on a 120 g Isco RediSep silica cartridge (EtOAc/heptane) provided (S)-2-benzyl 3-ethyl 8-(5-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridazin-3-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.


Step 3: N-CBZ Deprotection was accomplished via Method B to provide (S)-8-(5-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridazin-3-yl)-3-(ethoxycarbonyl)-2,8-diazaspiro[4.5]decane-2-carboxylic acid as a white solid.

5 Step 4: Hydrolysis of (S)-8-(5-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridazin-3-yl)-3-(ethoxycarbonyl)-2,8-diazaspiro[4.5]decane-2-carboxylic acid using the LiOH general method provided the title compound as an off-white solid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.66 - 1.80 (m, 4 H), 2.11 (dd, J = 13.45, 7.05 Hz, 1 H),
10 2.30 - 2.40 (m, 1 H), 2.36 (s, 3 H), 3.16 (d, J = 11.81 Hz, 1 H), 3.25 - 3.35 (m, 1 H), 3.37 - 3.65 (m, 4 H), 4.03 - 4.19 (m, 1 H), 6.39 (d, J = 2.34 Hz, 1 H), 6.63 (d, J = 2.39 Hz, 1 H), 6.95 (q, J = 6.39 Hz, 1 H), 7.43 - 7.57 (m, 2 H), 7.76 (d, J = 8.35 Hz, 1 H), 8.22 (d, J = 2.39 Hz, 1 H), 8.63 (d, J = 2.49 Hz, 1 H). LCMS (MH⁺): 551

15 **Example 44: (S)-8-(4-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyridin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid and**

Example 45: (S)-8-(4-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Example 45 X = Cl
Example 44: X = H

20
25 Step 1: To a solution of 2-chloro-4-nitropyridine (200 mg, 1.00 mmol) in 1,4-dioxane (6 mL) was added (R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol (368 mg, 1.27 mmol), and Cs₂CO₃ (828 mg, 2.54 mmol). The reaction was heated to 80 °C for 12 h, then cooled to RT, diluted with water, and extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica

gel column (EtOAc/heptane) provided (R)-2-chloro-4-(1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridine as an off-white solid.

5 *Step 2:* To a solution of (R)-2-chloro-4-(1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridine (227 mg, 0.57 mmol) in 1,4-dioxane (5 mL) was added 2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (237 mg, 0.68 mmol), Cs₂CO₃ (557 mg, 1.71 mmol), BINAP (142 mg, 0.23 mmol), and Pd₂(dba)₃. The reaction was heated to 60 °C for 3 d, then cooled to RT, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided 2-benzyl 3-ethyl 8-(4-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridin-2-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

15 *Step 3:* Hydrolysis of 2-benzyl 3-ethyl 8-(4-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridin-2-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate using the LiOH general method provided 2-((benzyloxy)carbonyl)-8-(4-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyridin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid.

20 *Step 4:* N-CBZ Deprotection was accomplished via Method A followed by normal phase silica gel purification (EtOAc:heptane) providing both of the title compounds as white solids (120 mg and 75 mg for the des-chloro analog).

8-(4-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyridin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid:


25 ¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.52 - 1.76 (m, 4 H) 1.95 - 2.15 (m, 1 H) 2.23 - 2.37 (m, 1 H) 2.39 (s, 3 H) 2.87 (s, 1 H) 3.05 - 3.16 (m, 1 H) 3.19 - 3.27 (m, 1 H) 3.38 - 3.72 (m, 4 H) 3.77 - 4.13 (m, 1 H) 6.39 (d, J=2.44 Hz, 1 H) 6.44 - 6.52 (m, 1 H) 6.79 (d, J=2.20 Hz, 1 H) 6.83 - 6.97 (m, 1 H) 7.43 - 7.51 (m, 1 H) 7.54 (d, J=2.05 Hz, 1 H) 7.66 (d, J=8.74 Hz, 1 H) 7.81 - 8.00 (m, 2 H). LCMS (MH⁺): 550.

30

8-(4-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyridin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid:

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.47 - 1.71 (m, 4 H) 1.95 - 2.04 (m, 1 H) 2.21 - 2.31 (m, 1 H) 2.39 (s, 3 H) 2.73 (s, 1 H) 3.02 (d, J=11.52 Hz, 1 H) 3.14 - 3.22 (m, 1 H) 3.37 - 4.03 (m, 4 H) 6.36 (d, J=2.34 Hz, 1 H) 6.43 - 6.51 (m, 1 H) 6.72 - 6.85 (m, 2 H) 7.30 - 7.51 (m, 3 H) 7.52 - 7.61 (m, 1 H) 7.67 (d, J=7.86 Hz, 1 H) 7.81 (d, J=2.34 Hz, 1 H) 7.86 - 7.91 (m, 1 H). LCMS (MH⁺): 516.

Example 46: 8-(4-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-6-phenoxy-pyrimidin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a solution of 2-benzyl 3-ethyl 8-(4-chloro-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-2-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (by-product from Step 3, Example 30a) (250 mg, 0.347 mmol) in 1,4-dioxane (9.0 mL) was added phenol (1.00 g, 10.6 mmol) and Cs₂CO₃ (3.65 g, 11.2 mmol). The reaction was heated at 80°C for 12 h, then cooled to RT diluted with water, and extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated *in vacuo*.

Purification on a 12 g Isco RediSep silica cartridge (EtOAc/heptane) provided 2-benzyl 3-ethyl 8-(4-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-6-phenoxy-pyrimidin-2-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as an off-white solid.


Step 2: N-CBZ Deprotection was accomplished via Method A to provide (ethyl 8-(4-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-6-phenoxy-pyrimidin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid.

Step 3: Hydrolysis of ethyl 8-(4-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-6-phenoxy)pyrimidin-2-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provided the title compound as an off-white solid.

- 5 ¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.36 (br. s., 4 H), 1.90 - 1.99 (m, 1 H), 2.11 - 2.21 (m, 1 H), 2.26 (s, 3 H), 2.92 - 3.17 (m, 2 H), 3.24 - 3.60 (m, 4 H), 3.96 (dd, J = 9.13, 6.88 Hz, 1 H), 5.44 (d, J = 2.29 Hz, 1 H), 6.27 - 6.33 (m, 1 H), 7.00 (d, J = 8.00 Hz, 2 H), 7.08 - 7.16 (m, 1 H), 7.24 - 7.32 (m, 2 H), 7.38 (dd, J = 8.44, 1.90 Hz, 1 H), 7.44 (d, J = 2.00 Hz, 1 H), 7.54 - 7.62 (m, 1 H), 7.64 (d, J = 8.49 Hz, 1 H), 7.81 (d, J = 2.25 Hz, 1 H). LCMS (MH⁺): 642.

10


Example 47: (3S)-8-(2-Amino-6-(1-(2,6-dibromophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



- 15 The title compound was prepared as described for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 10d) starting with 1-(2,6-dibromophenyl)-2,2,2-trifluoroethanol.

- 20 ¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.29 (m, 1H), 1.62 (q, J = 5.7 Hz, 4H), 2.06 (m, 1H), 2.33 (dd, J = 13.5, 9.2 Hz, 1H), 3.13 (d, J = 11.7 Hz, 1H), 3.26 (d, J = 11.7 Hz, 1H), 3.49 (m, 2H), 3.65 (dq, J = 10.7, 5.4 Hz, 2H), 4.09 (dd, J = 9.2, 7.2 Hz, 1H), 5.56 (s, 1H), 7.15 (t, J = 8.0 Hz, 1H), 7.28 (q, J = 8.0 Hz, 1H), 7.69 (m, 2H). LCMS (MH⁺): 611.


- 25 **Example 48: (S)-8-(2-Amino-6-((R)-1-(2,5-dibromophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid**



The title compound was prepared as described for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 10d) starting with 1-(2,5-dibromophenyl)-2,2,2-trifluoroethanol.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.62 (q, J = 5.8, 5.2 Hz, 4H), 2.06 (dd, J = 13.5, 7.2 Hz, 1H), 2.34 (dd, J = 13.4, 9.2 Hz, 1H), 3.13 (d, J = 11.7 Hz, 1H), 3.26 (d, J = 11.8 Hz, 1H), 3.50 (m, 2H), 3.66 (ddt, J = 15.0, 10.7, 5.2 Hz, 2H), 4.09 (dd, J = 9.2, 7.2 Hz, 1H), 4.83 (s, 1H), 5.58 (s, 1H), 6.97 (q, J = 6.6 Hz, 1H), 7.47 (dd, J = 8.6, 2.4 Hz, 1H), 7.58 (d, J = 8.6 Hz, 1H), 7.69 (d, J = 2.4 Hz, 1H). LCMS (MH⁺): 611.

Example 49: (S)-8-(2-Amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-4-propyl-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a solution of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-1-(2-bromo-5-chlorophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (660 mg, 0.95 mmol) in dioxane (12 mL) was added (3-(methylsulfonyl)phenyl)boronic acid (285 mg, 1.43 mmol), Pd₂(dppf)Cl₂ (70 mg, 0.095 mmol) and Na₂CO₃ (6.0 mL, 2.0 M, aq). The reaction was heated to 90 °C for 2 h, then cooled to RT, concentrated *in vacuo*. The residue was taken up in CH₂Cl₂, washed with brine, and extracted with CH₂Cl₂. The combined organic layers were dried

over Na₂SO₄. Purification by normal phase silica gel column (EtOAc/heptane) provided (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-1-(4-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

5 *Step 2:* To a solution of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-1-(4-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (500 mg, 0.65 mmol) in DMF (10 mL) was added tributyl(prop-1-enyl)stannane (258 mg, 0.78 mmol), Pd(t-Bu₃P)₂ (33 mg, 0.065 mmol), and CsF (217 mg, 1.43 mmol). The reaction was heated to 130 °C in a sealed tube for 3 h, then cooled to
10 RT. The reaction mixture was partitioned between water and CH₂Cl₂, and extracted. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-4-(prop-1-en-1-yl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white
15 solid.

Step 3: To a solution of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-4-(prop-1-en-1-yl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (200 mg, 0.26 mmol) in EtOH (10 mL) was added 10%
20 Pd/C (200 mg) and the reaction mixture was stirred under 1 atm H₂ for 12 h. The solids were filtered and the filtrate was concentrated to afford (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-4-propyl-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid that is used directly without further purification.


25
Step 4: To a solution of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-4-propyl-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate in CH₂Cl₂ (4 mL) was added TFA (2.0 mL) dropwise at 0 °C. The reaction mixture was stirred at RT for 2 h, then concentrated *in vacuo*. The pH was adjusted to
30 7-8 with saturated aqueous NaHCO₃ solution. The aqueous layer was extracted with CH₂Cl₂. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and

¹H NMR (400 MHz, MeOH-d₄): δ ppm 8.41 (m, 1H), 8.04 (d, J = 7.8 Hz, 1H), 7.79 (t, J = 7.8 Hz, 1H), 7.73-7.71 (m, 1H), 7.53 (s, 1H), 7.33 (d, J = 7.8 Hz, 1H), 7.20 (d, J = 7.8 Hz, 1H), 6.61 (q, J = 6.7 Hz, 1H), 5.61 (s, 1H), 4.10 (t, J = 8.4 Hz, 1H), 3.72-3.63 (m, 2H), 3.55-3.46 (m, 2H), 3.26 (m, 1H), 3.21 (s, 3H), 3.16-3.13 (m, 1H), 2.66 (t, J = 7.6 Hz, 2H), 2.38-2.32 (m, 1H), 2.10-2.05 (m, 2H), 1.65-1.60 (m, 3H). LCMS (MH⁺): 649.

CC1=CC=C(C=C1C2=CC=C(C=C2)C3=CC=C(C=C3)C=C)C4(C(F)(F)F)OC5=NC(=NC(=N5)N)N6CCN(CC6)C7CC[C@H](C7)C(=O)O

¹H NMR (400 MHz, CD₃OD-d₄): δ ppm 8.46-8.42 (m, 1H), 8.06-8.03 (m, 1H), 7.82-7.71 (m, 2H), 7.64 (s, 1H), 7.45 (dd, J₁ = 8.2 Hz, J₂ = 33.2 Hz, 1H), 7.25 (dd, J₁ = 7.9 Hz, J₂ = 23.9 Hz, 1H), 6.64-6.62 (m, 1H), 6.49-6.45 (m, 1H), 6.39-5.86 (m, 1H), 5.62 (d, J = 5.3 Hz, 1H), 4.12-4.08 (m, 1H), 3.70-3.62 (m, 2H), 3.54-3.45 (m, 2H), 3.29-3.26 (m, 1H), 3.22-3.21 (m, 3H), 3.16-3.13 (m, 1H), 2.37-2.31 (m, 1H), 2.10-2.05 (m, 1H), 1.91-1.87 (m, 3H), 1.62 (m, 4H). LCMS (MH⁺): 647.

Example 51a: (S)-8-(6-((R)-1-([1,1':4',1''-terphenyl]-2'-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




Step 1: To a solution of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-1-(2,5-dibromophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (660 mg, 0.95 mmol) in dioxane (12 mL) was added phenyl boronic acid (290 mg, 2.4 mmol), Pd₂(dppf)Cl₂ (70 mg, 0.095 mmol), and Na₂CO₃ (6.0 mL, 2.0 M, aq). The reaction mixture was heated to 90 °C for 2 h, then cooled to RT, concentrated *in vacuo*, and extracted with CH₂Cl₂. The combined organic layers were washed with brine, and dried over Na₂SO₄. Purification by normal phase silica gel column (EtOAc/heptane) provided (S)-2-tert-butyl 3-ethyl 8-(6-((R)-1-([1,1':4',1''-terphenyl]-2'-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

Step 2: To a solution of (S)-2-tert-butyl 3-ethyl 8-(6-((R)-1-([1,1':4',1''-terphenyl]-2'-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (550 mg, 0.75 mmol) in CH₂Cl₂ (4 mL) was added TFA (2.0 mL) dropwise at 0 °C. The reaction mixture was stirred at RT for 2 h, and concentrated *in vacuo*. The pH was adjusted to 7-8 with a saturated aqueous NaHCO₃ solution. The aqueous layer was extracted with CH₂Cl₂. The organic layer is washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (CH₂Cl₂/MeOH) provided (S)-ethyl 8-(6-((R)-1-([1,1':4',1''-terphenyl]-2'-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid.

Step 3: Hydrolysis of (S)-ethyl 8-(6-((R)-1-([1,1':4',1''-terphenyl]-2'-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provided the title compound as a white solid.

¹H NMR (400 MHz, CD₃OD-d₄): δ ppm 7.91 (s, 1H), 7.70 (dd, J₁ = 6.08 Hz, J = 1.88 Hz, 1H), 7.62 (m, 2H), 7.56-7.44 (m, 7H), 7.39-7.35 (m, 2H), 6.72 (q, J = 6.52 Hz, 1H), 5.48 (s, 1H), 4.18 (q, J = 6.96 Hz, 2H), 3.67 (m, 1H), 3.58 (m, 2H), 3.41 (m, 2H), 2.98 (d, J = 10.96 Hz, 1H), 2.69 (d, J = 11.24 Hz, 1H), 2.12-2.06 (m, 1H), 1.83-1.78 (m, 1H), 1.52 (m, 4H). LCMS (MH⁺): 604.5


Example 51b: (S)-8-(6-((R)-1-([1,1':3',1''-terphenyl]-2'-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described for (S)-8-(6-((R)-1-([1,1':4',1''-terphenyl]-2'-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 51a) starting with (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-1-(2,6-dibromophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (product of Step 4, example 63ao).

¹H NMR (400 MHz, CD₃OD-d₄): δ ppm 1.32 (dd, J = 15.5, 7.9 Hz, 1H), 1.70 (dd, J = 7.9, 4.3 Hz, 5H), 2.12 (m, 1H), 2.49 (ddd, J = 12.3, 9.0, 2.6 Hz, 1H), 3.25 (dd, J = 11.9, 2.2 Hz, 1H), 3.60 (s, 9H), 4.48 (t, J = 8.6 Hz, 1H), 6.89 (q, J = 7.8 Hz, 1H), 7.21 (d, J = 7.6 Hz, 2H), 7.42 (m, 14H). LCMS (MH⁺): 604.

Example 52a: (S)-8-(2-Amino-6-((R)-1-(3,4-dimethyl-3''-(methylsulfonyl)-[1,1':3',1''-terphenyl]-4'-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (product of Step 1, Example 34w) (273 mg, 0.34 mmol) in 1,4-dioxane (5 mL) was added (3,4-dimethylphenyl)boronic acid (77 mg, 0.51 mmol), KHCO_3 (341 mg, 3.40 mmol), and $\text{Pd}(\text{PCy}_3)_2$ (34 mg, 0.051 mmol). The reaction was heated to 100 °C for 44 h. The reaction was charged with additional $\text{Pd}(\text{PCy}_3)_2$ (68 mg, 0.10 mmol) at $t = 16$ and 39 h. Then the reaction was cooled to RT and extracted with EtOAc. The combined organic layers were dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification on a 12 g Isco RediSep silica cartridge (EtOAc/heptane) provided (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(3,4-dimethyl-3''-(methylsulfonyl)-[1,1':3',1''-terphenyl]-4'-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

Step 2: N-CBZ Deprotection was accomplished via Method B to provide (S)-ethyl 8-(2-amino-6-((R)-1-(3,4-dimethyl-3''-(methylsulfonyl)-[1,1':3',1''-terphenyl]-4'-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid.

Step 3: Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-1-(3,4-dimethyl-3''-(methylsulfonyl)-[1,1':3',1''-terphenyl]-4'-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provided the title compound as an off-white solid.

Using the generic scheme below, the following examples of Table 16a were prepared as described above for (S)-8-(2-amino-6-((R)-1-(3,4-dimethyl-3''-(methylsulfonyl)-[1,1':3',1''-terphenyl]-4'-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 52a).





Table 16a.



Ex. No.	Cy	CAS Name	LCMS (MH ⁺)
52a		(S)-8-(2-amino-6-((R)-1-(3,4-dimethyl-3''-(methylsulfonyl)-[1,1':3',1''-terphenyl]-4'-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	710
52b		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-(quinolin-6-yl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	733

5


Table 16b

NMR Data for Compounds of Table 16a

Ex. No.	NMR
52a	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.46 - 1.73 (m, 4 H) 2.07 (dd, J=13.45, 7.15 Hz,

	1 H) 2.28 (s, 3 H) 2.30 (s, 3 H) 2.32 - 2.40 (m, 1 H) 3.14 (d, J=11.76 Hz, 1 H) 3.22 (s, 3 H) 3.27 (d, J=11.76 Hz, 1 H) 3.40 - 3.77 (m, 4 H) 4.09 (dd, J=9.08, 7.27 Hz, 1 H) 5.62 (s, 1 H) 6.63 (q, J=6.64 Hz, 1 H) 7.18 (d, J=7.96 Hz, 1 H) 7.35 (dd, J=7.81, 1.81 Hz, 1 H) 7.40 (s, 1 H) 7.47 (d, J=1.85 Hz, 1 H) 7.63 - 7.72 (m, 1 H) 7.72 - 7.77 (m, 1 H) 7.80 - 7.85 (m, 2 H) 8.07 (dt, J=6.97, 1.96 Hz, 1 H) 8.48 (br. s., 1 H)
52b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.59 (t, J=5.54 Hz, 4 H) 1.92 (dd, J=13.13, 7.03 Hz, 1 H) 2.20 (dd, J=13.15, 9.10 Hz, 1 H) 2.81 - 3.17 (m, 2 H) 3.24 (s, 3 H) 3.38 - 3.74 (m, 4 H) 3.84 (dd, J=8.96, 7.05 Hz, 1 H) 5.64 (s, 1 H) 6.67 (q, J=6.64 Hz, 1 H) 7.55 (dd, J=8.35, 4.34 Hz, 1 H) 7.67 (d, J=1.61 Hz, 1 H) 7.78 - 7.92 (m, 4 H) 8.04 - 8.15 (m, 3 H) 8.21 (s, 1 H) 8.41 (dd, J=8.40, 1.56 Hz, 1 H) 8.54 (br. s., 1 H) 8.84 (dd, J=4.32, 1.68 Hz, 1 H)

Example 53: (S)-8-(2-Amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-((E)-prop-1-en-1-yl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



5

Step 1: To a solution of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (600 mg, 0.89 mmol) in dioxane (12 mL) was added (3-(methylsulfonyl)phenyl)boronic acid (275 mg, 1.3 mmol), Pd₂(dppf)Cl₂ (65 mg, 0.095 mmol), and Na₂CO₃ (6.0 mL, 2.0 M, aq). The reaction was heated to 90 °C for 2 h, then cooled to RT, and concentrated *in vacuo*. The residue was taken up in CH₂Cl₂, washed with brine, and dried over Na₂SO₄. Purification by normal phase silica gel column (EtOAc/heptane) provides (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

15

Step 2: To a solution (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (500 mg, 0.65 mmol) in DMF (10 mL) was added tributyl(prop-1-enyl)stannane


(258 mg, 0.78 mmol), Pd(t-Bu₃P)₂ (33 mg, 0.065 mmol), and CsF (217 mg, 1.43 mmol). The reaction was heated to 130 °C in a sealed tube for 3 h, then cooled to RT, and partitioned between between water and CH₂Cl₂. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-((E)-prop-1-en-1-yl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

Step 3: To a solution of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-((E)-prop-1-en-1-yl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate in CH₂Cl₂ (4 mL) was added TFA (2.0 mL) dropwise at 0 °C. The reaction mixture was stirred at RT for 2 h, then concentrated *in vacuo*. The pH was adjusted to 7-8 with a saturated aqueous NaHCO₃ solution. The aqueous layer was extracted with CH₂Cl₂, washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (CH₂Cl₂/MeOH) provided (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-((E)-prop-1-en-1-yl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid.

Step 4: Hydrolysis of (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-((E)-prop-1-en-1-yl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid using the LiOH general method provided the title compound as an off-white solid.

¹H NMR (400 MHz, CD₃OD-d₄): δ ppm 8.40 (s, 1 H), 8.02 (d, 1 H, J=7.4 Hz), 7.50 (m, 3 H), 7.40 (m, 1 H), 7.20 (m, 1 H), 6.58 (m, 1 H), 5.58 (m, 1 H), 4.09 (m, 1 H), 3.55 (m, 2 H), 3.48 (m, 2 H), 3.21 (m, 4H), 3.10 (m, 1 H), 2.59 (m, 2 H), 2.29 (m, 1 H), 1.95 (m, 1 H), 1.86 (m, 3 H), 1.30 (m, 4 H). LCMS (MH⁺): 646.

Example 54a: (S)-8-(2-Amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-propyl-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a solution of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-
 5 mmol) in EtOH (10 mL) is added 10% Pd/C (200 mg), and the reaction mixture was stirred under 1 atm H₂ for 12 h. The solids were filtered and the filtrate was concentrated *in vacuo* to provide (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-propyl-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid that is used directly without further purification.


Step 2: To a solution of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-propyl-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]
 15 decane-2,3-dicarboxylate in CH₂Cl₂ (4 mL) was added TFA (2.0 mL) dropwise at 0 °C. The reaction mixture was stirred at RT for 2 h, then concentrated *in vacuo*. The pH was adjusted to 7-8 with saturated aqueous NaHCO₃ solution. The aqueous layer was extracted with CH₂Cl₂, washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (CH₂Cl₂/MeOH) provided (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-propyl-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a white solid.

Step 3: Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-propyl-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provides the title compound as an off-white solid.

¹H NMR (400 MHz, CD₃OD-d₄): δ ppm 8.40 (s, 1 H), 8.02 (d, 1 H, J=7.8 Hz), 7.60 (m, 3 H), 7.29 (m, 1 H), 7.08 (s, 1 H), 6.58 (m, 1 H), 5.56 (s, 1 H), 4.00 (m, 1 H), 3.55 (m, 2 H), 3.48 (m, 2

H), 3.31 (m, 4H), 3.30 (m, 1 H), 2.59 (m, 2 H), 2.29 (m, 1 H), 1.95 (m, 1 H), 1.54 (m, 6 H), 0.95 (m, 3 H). LCMS (MH⁺): 649.

Example 54b: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-isopropoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a solution of (S)-ethyl 8-(2-amino-6-((R)-1-(5-bromo-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (350 mg, 0.56 mmol) in CH₂Cl₂ (20 mL) was added Boc₂O (436 mg, 2.0 mmol) and Et₃N (306 mg, 3.03 mmol) at 0 °C.

The reaction mixture was stirred at RT for 3 h, then concentrated *in vacuo* and purified on normal phase silica gel (ethyl acetate/hexanes) to afford (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-1-(5-bromo-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a yellow solid.

Step 2: A solution of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-1-(5-bromo-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (150 mg, 0.2 mmol), 4-isopropoxyphenyl boronic acid (44 mg, 0.25 mmol) and Pd(dppf)Cl₂ (15 mg, 0.02 mmol) in dioxane (3.0 mL) / aqueous Na₂CO₃ solution (3.0 mL, 2.0 M, aq.) was stirred at 90 °C for 2 h. The aqueous layer was extracted with CH₂Cl₂, washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/ Hex = 10 to 50 %) to (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-isopropoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.


Step 3: To a solution of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-isopropoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-

dicarboxylate (130mg, 0.164 mmol) in CH₂Cl₂ (4 mL) was added TFA (1 mL), and the reaction mixture was stirred at 25 °C for 12 h. The mixture was concentrated, and neutralized to pH 7-8 with saturated aqueous NaHCO₃. The aqueous layer was extracted with CH₂Cl₂, washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to provide (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-isopropoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a light yellow solid that is used without further purification.

Step 4: Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-isopropoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate using the LiOH general method provided the title compound as an off-white solid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.31 (d, J = 6.0 Hz, 6H), 1.58 (m, 4H), 2.04 (dd, J = 13.4, 7.2 Hz, 1H), 2.32 (dd, J = 13.4, 9.2 Hz, 1H), 3.11 (d, J = 11.7 Hz, 1H), 3.24 (d, J = 11.7 Hz, 1H), 3.45 (ddd, J = 21.2, 10.1, 6.4 Hz, 2H), 3.60 (td, J = 12.4, 11.2, 6.0 Hz, 2H), 4.08 (dd, J = 9.1, 7.1 Hz, 1H), 4.62 (p, J = 6.1 Hz, 1H), 6.67 (q, J = 6.8 Hz, 1H), 6.95 (m, 2H), 7.54 (m, 9H), 7.72 (d, J = 8.3 Hz, 1H). LCMS (MH⁺): 663.


Example 54c: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-propoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described above for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-isopropoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 54b) by substituting 4-propoxyphenyl boronic acid for 4-isopropoxyphenyl boronic acid in Step 2.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.04 (t, J = 7.4 Hz, 3H), 1.57 (m, 4H), 1.80 (h, J = 6.7 Hz, 2H), 1.99 (dd, J = 13.3, 7.3 Hz, 1H), 2.27 (dd, J = 13.3, 9.1 Hz, 1H), 3.02 (d, J = 11.6 Hz, 1H), 3.18 (d, J = 11.5 Hz, 1H), 3.30 (d, J = 3.2 Hz, 1H), 3.45 (q, J = 15.9, 11.4 Hz, 2H), 3.60 (s, 2H), 3.97 (dt, J = 13.1, 7.3 Hz, 3H), 4.88 (m, 1H), 5.47 (s, 1H), 6.66 (q, J = 6.9 Hz, 1H), 6.97 (d, J = 8.3 Hz, 2H), 7.54 (m, 9H), 7.72 (m, 1H). LCMS (MH⁺): 662.

Example 54d: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(5-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a mixture of 2-chloro-4-(methylsulfonyl)benzoic acid (5 g, 21.3 mmol) in anhydrous methanol (100 mL) was added concentrated sulfuric acid (0.5 mL). The resulting solution was stirred for 18 h at reflux. Upon cooling, the mixture was concentrated under reduced pressure, dissolved in CH₂Cl₂ and washed with NaHCO₃ solution and brine. The organic phase was dried over sodium sulfate and concentrated to afford methyl 2-chloro-4-(methylsulfonyl)benzoate as a white solid.

Step 2: To a mixture of methyl 2-chloro-4-(methylsulfonyl)benzoate (2.2 g, 8.9 mmol), PhB(OH)₂ (1.31 g, 10.8 mmol), DME (12 mL), and 2M Na₂CO₃ (6 mL) was added Pd(PPh₃)₄ (515 mg). The mixture was heated for 20 min at 160 °C in a microwave reactor, and then extracted with EtOAc, dried over sodium sulfate and concentrated in vacuo. Purification on normal phase silica gel (hexane/EtOAc) provided methyl 5-(methylsulfonyl)-[1,1'-biphenyl]-2-carboxylate as a white solid.

Step 3: To a solution of CaCl₂ (1.52 g, 13.78 mmol) in EtOH (50 mL) at RT was added methyl 5-(methylsulfonyl)-[1,1'-biphenyl]-2-carboxylate (2 g, 6.9 mmol) in THF (50 mL) followed by the

addition of NaBH₄ (1.0 g, 27.6 mmol). The reaction was stirred at RT for 24 h, then concentrated *in vacuo* and extracted with ethyl acetate, 5% HCl, and brine. Purification on normal phase silica gel provided (5-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)methanol as a white solid.

5

Step 4: To a solution of (5-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)methanol (1 g, 3.8 mmol) in CH₂Cl₂ (50 mL) was added Dess-Martin periodinane (2.4 g, 5.71mmol). The reaction was stirred for 2 h at RT, then concentrated *in vacuo* and purified directly on normal phase silica gel to provide 5-(methylsulfonyl)-[1,1'-biphenyl]-2-carbaldehyde as a white solid.

10

Step 5: To a solution of 5-(methylsulfonyl)-[1,1'-biphenyl]-2-carbaldehyde (1 g, 3.8 mmol) was added TMS-CF₃ (1.0 g, 7.7 mmol) in THF (10 mL). The reaction was cooled to 0 °C to and TBAF (0.57 mL, 0.57 mmol) was added dropwise. The reaction mixture was stirred for 2 h, then 3 N HCl (2 mL) was added to the mixture and the reaction mixture was stirred for an additional 30 min. The mixture was extracted with ethyl acetate, washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification on normal phase silica gel provided 2,2,2-trifluoro-1-(5-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)ethanol as a white solid.

15

20

Step 6: To a mixture of 2,2,2-trifluoro-1-(5-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)ethanol (720 mg, 2.2 mmol)) in CH₂Cl₂ (50 mL) was added Dess-Martin periodinane (1.1 g, 2.6mmol). The reaction was stirred for 2 h at RT, then concentrated *in vacuo* and purified directly on normal phase silica gel to provide 2,2,2-trifluoro-1-(5-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)ethanone as a white solid.

25

Step 7: Chiral reduction of 2,2,2-trifluoro-1-(5-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)ethanone using the Iridium complex-catalyzed hydrogenation as described for Intermediate 1, (R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol, provided (R)-2,2,2-trifluoro-1-(5-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)ethanol as a white solid.


30

Steps 8-11: The title compound was prepared as described for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-

diazaspiro[4.5]decane-3-carboxylic acid (Example 10d), Steps 1-4.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.63 (q, J = 5.7, 4.9 Hz, 4H), 2.10 (m, 1H), 2.36 (dd, J = 13.5, 9.2 Hz, 1H), 3.23 (d, J = 31.0 Hz, 5H), 3.50 (dddd, J = 18.0, 13.4, 9.5, 5.1 Hz, 2H), 3.66 (ddt, J = 15.9, 10.6, 4.6 Hz, 2H), 4.16 (dd, J = 9.2, 7.2 Hz, 1H), 6.78 (q, J = 6.7 Hz, 1H), 7.57 (m, 5H), 7.86 (d, J = 1.9 Hz, 1H), 8.01 (m, 2H), 8.17 (s, 1H). LCMS (MH⁺): 607.


Example 54e: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-fluoro-4-propoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described above for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-isopropoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 54b) by replacing the 4-isopropoxyphenyl boronic acid in Step 2 with (3-fluoro-4-propoxyphenyl)boronic acid (CAS# 192376-68-4).

¹H NMR (400 MHz, MeOH-d₄): δ ppm 0.86 (m, 1H), 1.05 (t, J = 7.4 Hz, 3H), 1.26 (s, 1H), 1.59 (s, 4H), 1.83 (h, J = 7.1 Hz, 2H), 2.06 (dd, J = 13.4, 7.2 Hz, 1H), 2.33 (m, 1H), 3.10 (d, J = 11.9 Hz, 1H), 3.23 (d, J = 12.0 Hz, 1H), 3.43 (s, 2H), 3.60 (s, 2H), 4.02 (t, J = 6.5 Hz, 2H), 4.12 (s, 1H), 6.62 (d, J = 6.8 Hz, 1H), 7.09 (t, J = 8.7 Hz, 1H), 7.34 (s, 1H), 7.43 (m, 4H), 7.50 (s, 3H), 7.60 (m, 1H), 7.76 (m, 2H). LCMS (MH⁺): 681.


Example 54f: (S)-8-(2-amino-6-((R)-1-(3,4-dimethyl-[1,1':3',1''-terphenyl]-4'-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described above for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-isopropoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 54b) by replacing the 4-isopropoxyphenyl boronic acid in Step 2 with 3,4-dimethylphenyl boronic acid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.62 (s, 4H), 2.06 (dd, J = 13.5, 7.6 Hz, 1H), 2.29 (d, J = 9.7 Hz, 5H), 2.37 (m, 1H), 3.17 (d, J = 11.8 Hz, 1H), 3.26 (d, J = 11.7 Hz, 1H), 3.63 (d, J = 14.2 Hz, 2H), 4.27 (t, J = 8.3 Hz, 1H), 6.66 (q, J = 6.8 Hz, 1H), 7.18 (d, J = 7.9 Hz, 1H), 7.36 (m, 2H), 7.49 (m, 5H), 7.64 (dd, J = 8.2, 2.0 Hz, 1H), 7.74 (d, J = 8.2 Hz, 1H). LCMS (MH⁺): 633.

Example 54g: (S)-8-(6-((R)-1-([1,1':3',1''-terphenyl]-4'-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




The title compound was prepared as described above for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-isopropoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 54b) by substituting phenyl boronic acid for 4-isopropoxyphenyl boronic acid in Step 2.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.62 (s, 4H), 2.06 (dd, J = 13.5, 7.7 Hz, 1H), 2.38 (dd, J = 13.5, 9.1 Hz, 1H), 3.16 (d, J = 11.8 Hz, 1H), 3.26 (d, J = 11.8 Hz, 1H), 3.47 (s, 2H), 3.62 (s,

2H), 4.26 (t, J = 8.4 Hz, 1H), 6.68 (q, J = 6.9 Hz, 1H), 7.35 (m, 1H), 7.47 (m, 4H), 7.53 (s, 3H), 7.66 (m, 3H), 7.77 (d, J = 8.2 Hz, 1H). LCMS (MH⁺): 604.


Example 54h: (R)-8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described above for (S)-8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 34c) by using (R)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.59 (d, J = 5.5 Hz, 4H), 2.03 (dd, J = 13.4, 7.1 Hz, 1H), 2.31 (dd, J = 13.4, 9.2 Hz, 1H), 3.09 (d, J = 11.8 Hz, 1H), 3.23 (d, J = 11.6 Hz, 1H), 3.46 (dt, J = 15.3, 8.2 Hz, 2H), 3.62 (s, 2H), 4.06 (dd, J = 9.1, 7.1 Hz, 1H), 5.49 (s, 1H), 6.64 (q, J = 6.9 Hz, 1H), 7.28 (d, J = 2.2 Hz, 1H), 7.46 (m, 5H), 7.53 (s, 1H), 7.67 (d, J = 8.5 Hz, 1H). LCMS (MH⁺): 562.

Example 54i: (R)-8-(2-amino-6-((S)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was prepared as described above for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-


methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 34c) by using (R)-2-benzyl 3-ethyl 8-(2-amino-6-((S)-1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate.

5

¹H NMR (400 MHz, MeOH-d₄): δ ppm 7.70 (d, J = 8.5 Hz, 1H), 7.59 – 7.44 (m, 4H), 7.47 – 7.40 (m, 2H), 7.32 (d, J = 2.2 Hz, 1H), 6.61 (q, J = 6.5 Hz, 1H), 4.51 (t, J = 8.7 Hz, 1H), 3.72 – 3.59 (m, 1H), 3.56 (s, 1H), 3.28 (s, 1H), 2.49 (dd, J = 13.6, 8.9 Hz, 1H), 2.10 (dd, J = 13.6, 8.4 Hz, 1H), 1.71 (dt, J = 16.0, 6.6 Hz, 4H), 1.28 (s, 0H). LCMS (MH⁺): 562.

10

Example 54j: (S)-8-(2-amino-6-((S)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




The title compound was prepared as described above for (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 34c) by using (S)-2-benzyl 3-ethyl 8-(2-amino-6-((S)-1-(2-bromo-4-chlorophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate.

15

¹H NMR (400 MHz, MeOH-d₄): δ ppm 7.70 (d, J = 8.5 Hz, 1H), 7.61 – 7.42 (m, 6H), 7.32 (d, J = 2.3 Hz, 1H), 6.66 (q, J = 6.7 Hz, 1H), 4.25 (dd, J = 9.0, 7.6 Hz, 1H), 3.72 – 3.60 (m, 1H), 3.29 (d, J = 11.7 Hz, 1H), 3.18 (d, J = 11.8 Hz, 1H), 2.40 (dd, J = 13.5, 9.2 Hz, 1H), 2.09 (dd, J = 13.5, 7.6 Hz, 1H), 1.64 (s, 2H). LCMS (MH⁺): 562.

20

Example 54k: (S)-8-(2-amino-6-((S)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




5 The title compound was prepared as described above for (S)-8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro [4.5]decane-3-carboxylic acid (Example 1m) by using (S)-2-benzyl 3-ethyl 8-(2-amino-6-((S)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate.

10

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.58 (s, 6H), 2.04 (dd, J = 13.4, 7.2 Hz, 1H), 2.30 (d, J = 11.1 Hz, 9H), 2.40 (s, 3H), 3.10 (d, J = 11.8 Hz, 1H), 3.23 (d, J = 11.7 Hz, 1H), 3.48 (s, 2H), 3.66 (d, J = 15.7 Hz, 3H), 4.08 (t, J = 8.2 Hz, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.77 (q, J = 6.5 Hz, 1H), 7.20 (d, J = 7.8 Hz, 1H), 7.38 (d, J = 8.0 Hz, 1H), 7.44 (d, J = 2.0 Hz, 1H), 7.60 (d, J = 1.8 Hz, 1H), 7.73 (m, 2H), 7.97 (d, J = 2.4 Hz, 1H). LCMS (MH⁺): 635.

15

Example 54l: (R)-8-(2-amino-6-((S)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



20

The title compound was prepared as described above for (R)-8-(2-amino-6-((R)-1-(3',4'-


dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro [4.5]decane-3-carboxylic acid (Example 1m) by using (R)-2-benzyl 3-ethyl 8-(2-amino-6-((S)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate.

5

¹H NMR (400 MHz, MeOH-d₄): δ ppm 7.97 (d, J = 2.3 Hz, 0H), 7.79 – 7.69 (m, 0H), 7.61 (d, J = 1.6 Hz, 0H), 7.45 (s, 0H), 7.42 – 7.35 (m, 0H), 7.21 (d, J = 7.9 Hz, 0H), 6.77 (q, J = 6.5 Hz, 0H), 6.41 (d, J = 2.3 Hz, 0H), 4.10 (t, J = 8.2 Hz, 0H), 3.68 (dd, J = 13.9, 6.3 Hz, 0H), 3.58 – 3.43 (m, 0H), 3.24 (d, J = 11.7 Hz, 0H), 3.11 (d, J = 11.8 Hz, 0H), 2.42 – 2.27 (m, 1H), 2.05 (dd, J = 13.5, 7.2 Hz, 0H), 1.59 (d, J = 11.4 Hz, 0H), 1.59 (s, 0H).. LCMS (MH⁺): 635.

10

Example 54m: (R)-8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



15


The title compound was prepared as described above for (R)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro [4.5]decane-2,3-dicarboxylate (Example 1m) by using (R)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate.

20

¹H NMR (400 MHz, MeOH-d₄): δ ppm 7.97 (d, J = 2.4 Hz, 1H), 7.79 – 7.68 (m, 2H), 7.60 (d, J = 1.7 Hz, 1H), 7.44 (s, 1H), 7.38 (d, J = 8.0 Hz, 1H), 7.20 (d, J = 7.8 Hz, 1H), 6.76 (q, J = 6.7 Hz, 1H), 6.41 (d, J = 2.3 Hz, 1H), 5.75 (s, 1H), 3.98 (t, J = 8.1 Hz, 1H), 3.64 (d, J = 15.5 Hz, 3H), 3.47 (s, 2H), 3.33 – 3.27 (m, 6H), 3.17 (d, J = 11.6 Hz, 1H), 3.01 (d, J = 11.6 Hz, 1H), 2.39 (s, 3H), 2.34 – 2.18 (m, 8H), 1.99 (dd, J = 13.4, 7.1 Hz, 1H), 1.56 (s, 5H). LCMS (MH⁺): 635.

25

Example 55an: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



5

Step 1: To a solution of (R)-1-(4-bromophenyl)-2,2,2-trifluoroethanol (150 mg, 0.60 mmol) in dioxane (10 mL) was added 4,6-dichloropyrimidin-2-amine (120 mg, 0.71 mmol) and Cs_2CO_3 (290 mg, 0.88 mmol), and the reaction mixture was heated to 80 °C for 30 h. Then the reaction was cooled to RT. EtOAc was added and the organic layer was washed with brine, dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided (R)-4-(1-(4-bromophenyl)-2,2,2-trifluoroethoxy)-6-chloropyrimidin-2-amine as a colorless oil.

Step 2: To a solution of (R)-4-(1-(4-bromophenyl)-2,2,2-trifluoroethoxy)-6-chloropyrimidin-2-amine (19 mg, 0.50 mmol) in dioxane (25 mL) was added (S)-2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (175 mg, 0.50 mmol) and sodium bicarbonate (210 mg, 0.25 mmol), and the reaction mixture was heated to 100 °C for 48 h. Then the reaction mixture was cooled to RT, and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by normal phase silica gel column (EtOAc/heptane) provided (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-bromophenyl)-2,2,2-trifluoroethoxy)-pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as white solid.

Step 3: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(4-bromophenyl)-2,2,2-trifluoroethoxy)-pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (190 mg, 0.27 mmol) was added NaOH (100 mg, 0.26 mmol) in 15 mL THF/EtOH/ H_2O (2/1/2.5), and the

reaction was stirred for 12 h at RT. Then, the reaction mixture was concentrated *in vacuo* to remove most of the organic solvents, and the pH was adjusted to 6 with 1 N HCl. EtOAc was added, and the organic layer was washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to provide (S)-8-(2-amino-6-((R)-1-(4-bromophenyl)-2,2,2-

trifluoroethoxy) pyrimidin-4-yl)-2-(benzyloxycarbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid as a white solid which was used without further purification.

Step 4: To a solution of (S)-8-(2-amino-6-((R)-1-(4-bromophenyl)-2,2,2-trifluoroethoxy) pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (80 mg, 0.12mmol) in dioxane (1 mL)/ Na₂CO₃(1.0 mL, 2 M, aq) were added (3-methoxyphenyl)boronic acid (22 mg, 0.14 mmol) and Pd(dppf)₂ (8 mg, 0.01 mmol). The reaction flask was degassed and refilled with argon via balloon 3 times, and the reaction mixture was refluxed for 4 h. Then the reaction was cooled to RT, concentrated *in vacuo*, and extracted with EtOAc. The combined organic layers were are washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by reverse phase silica gel column (H₂O/NH₄OH/MeOH) provided (S)-8-(2-amino-6-((S)-2,2,2-trifluoro-1-(3'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid as a white solid.

Step 5: N-CBZ Deprotection was accomplished via Method A to provide the title compound as an off-white solid isolated as the zwitterionic form.

Using the generic scheme below, the following examples of Table 17a were prepared as described above for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 55an) using the appropriate boronic acid or boronate. In some cases, the Cy coupling reaction was performed prior to ethyl ester and N-CBz removal (see alternative Steps 3a and 4a) as noted in the scheme. In the cases of example 55al and 55am, racemic 1-(4-bromophenyl)-2,2,2-trifluoroethanol was used as opposed to (R)-1-(4-bromophenyl)-2,2,2-trifluoroethanol for all other examples.
















































Table 17a.















Ex. No.	Cy	CAS Name	LCMS (MH+)
55a		(S)-8-(6-((R)-1-([1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	529
55b		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1-methyl-1H-indazol-5-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	583
55c		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1-methyl-1H-benzo[d]imidazol-5-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	583
55d		(S)-8-(6-((R)-1-(4-(1H-benzo[d]imidazol-5-yl)phenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	569
55e		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	577
55f		(S)-8-(2-amino-6-((R)-1-(4-(benzo[d]isothiazol-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	586
55g		(S)-8-(2-amino-6-((R)-1-(4-(benzo[d]isoxazol-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	570
55h		(S)-8-(6-((R)-1-(4-(1H-indazol-6-yl)phenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	569
55i		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1-methyl-1H-indazol-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	583
55j		(S)-8-(2-amino-6-((R)-1-(4-(benzo[d]isothiazol-5-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	586
55k		(S)-8-(2-amino-6-((R)-1-(4-(benzo[d]thiazol-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	586
55l		(S)-8-(6-((R)-1-(4-([1,2,4]triazolo[1,5-a]pyridin-6-yl)phenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	570

55m		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(naphthalen-2-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	579
55n		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-4'-methyl-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	573
55o		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-5'-methyl-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	573
55p		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(5'-methoxy-2'-methyl-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	573
55q		(S)-8-(2-amino-6-((R)-1-(3',4'-dimethoxy-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	589
55r		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-4'-(pyrrolidine-1-carbonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	656
55s		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1-oxo-1,3-dihydroisobenzofuran-5-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	585
55t		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-oxo-1,2-dihydroquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	596
55u		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1-methyl-2-oxo-1,2-dihydroquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	610
55v		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	598
55w		(S)-8-(6-((R)-1-(4-(1H-indazol-5-yl)phenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	569














55x		(S)-8-(2-amino-6-((R)-1-(4-(1,3-dimethyl-1H-indazol-5-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	597
55y		(S)-8-(2-amino-6-((R)-1-(4-(1,3-dimethyl-1H-indol-5-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	596
55z		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-5'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	627
55aa		(S)-8-(2-amino-6-((R)-1-(3'-cyano-5'-methoxy-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	584
55ab		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	586
55ac		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(3-methyl-1H-indol-5-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	1
55ad		(S)-8-(6-((R)-1-(3'-acetoxy-4'-(methoxycarbonyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	645
55ae		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-oxo-2H-chromen-7-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	596
55af		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	560
55ag		(S)-8-(2-amino-6-((R)-1-(4'-carboxy-3'-hydroxy-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	588













55ah		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-methoxyquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	610
55ai		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-(methylthio)quinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	626
55aj		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	612
55ak		(3S)-8-(2-amino-6-(2,2,2-trifluoro-1-(3'-fluoro-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	547
55al		(3S)-8-(2-amino-6-(2,2,2-trifluoro-1-(3'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	559
55am		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	547
55an		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	559
55ao		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-5'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	577
55ap		(S)-8-(2-amino-6-((R)-1-(3',5'-difluoro-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	565
55aq		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	559
55ar		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	559












55as		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	597
55at		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(trifluoromethoxy)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	613
55au		(S)-8-(2-amino-6-((R)-1-(3'-ethoxy-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	573
55av		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-isopropoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	587
55aw		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(pyridin-3-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	530
55ax		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(pyridin-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	530
55ay		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(pyrimidin-5-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	531
55az		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(3-methyl-1H-indazol-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	583
55ba		(S)-8-(2-amino-6-((R)-1-(4-(1,3-dimethyl-1H-indazol-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	597
55bb		S)-8-(2-amino-6-((R)-1-(4-(2,3-dimethyl-2H-indazol-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	597
55bc		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1-oxo-1,2,3,4-tetrahydroisoquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	598

55bd		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(isoquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	580
55be		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(isoquinolin-7-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	580
55bf		(S)-8-(2-amino-6-((R)-1-(4'-((dimethylamino)methyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	586
55bg		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(quinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	580
55bh		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(quinolin-7-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	580
55bi		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(quinoxalin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	581
55bj		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-methyl-1-oxo-1,2,3,4-tetrahydroisoquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	612
55bk		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(quinazolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	581
55bl		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-fluoro-2'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	577
55bm		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2'-fluoro-3'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	577
55bn		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	577
55bo		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(6-methylpyridin-3-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	544

55bp		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(pyrrolidine-1-carbonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	626
55bq		(S)-8-(2-amino-6-((R)-1-(3'-(2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	573
55br		(S)-8-(2-amino-6-((R)-1-(4'-(2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	573
55bs		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(propyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	571
55bt		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(hydroxymethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	559
55bu		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2'-(hydroxymethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	559
55bv		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(isopropoxy)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	587
55bw		(S)-8-(2-amino-6-((R)-1-(4'-(dimethylcarbamoyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	600
55bx		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(piperidine-1-carbonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	640
55by		(S)-8-(2-amino-6-((R)-1-(2'-((dimethylamino)methyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	586
55bz		(S)-8-(2-amino-6-((R)-1-(4'-(ethyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	557
55ca		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(hydroxy)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	545

55cb		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-hydroxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	545
55cc		(S)-8-(2-amino-6-((R)-1-(2',4'-dimethoxy-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	589
55cd		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	597
55ce		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	597
55cf		(S)-8-(2-amino-6-((R)-1-(2',6'-difluoro-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	565
55cg		(S)-8-(2-amino-6-((R)-1-(2',6'-dimethyl-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	557
55ch		(S)-8-(2-amino-6-((R)-1-(3',4'-dimethyl-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	557
55ci		(S)-8-(2-amino-6-((R)-1-(4'-(tert-butyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	585
55cj		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropyl-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	571
55ck		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-isopropyl-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	571
55cl		(S)-8-(2-amino-6-((R)-1-(3',4'-dichloro-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	597
55cm		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(trifluoromethoxy)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	613
55cn		(S)-8-(2-amino-6-((R)-1-(2',3'-dimethyl-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	557

55co		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3',4',5'-trifluoro-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	583
55cp		(S)-8-(2-amino-6-((R)-1-(4'-chloro-2'-methyl-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	577
55cq		(S)-8-(2-amino-6-((R)-1-(3',5'-dimethyl-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	557
55cr		(S)-8-(2-amino-6-((R)-1-(3',4'-difluoro-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	565
55cs		(S)-8-(2-amino-6-((R)-1-(2',5'-dimethyl-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	557
55ct		(S)-8-(2-amino-6-((R)-1-(4'-butyl-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	585
55cu		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-methyl-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	561
55cv		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(methylsulfonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	607
55cw		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-methyl-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	543
55cx		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methyl-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	543
55cy		(S)-8-(2-amino-6-((R)-1-(4'-chloro-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	563
55cz		(S)-8-(2-amino-6-((R)-1-(4-(benzofuran-3-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	569

55da		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(5'-fluoro-2'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	577
55db		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-oxochroman-7-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	599
55dc		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(3-fluoroquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	597
55dd		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-propoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	587
55de		(S)-8-(2-amino-6-((R)-1-(4'-(diethylcarbamoyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	628
55df		(S)-8-(2-amino-6-((R)-1-(4'-carbamoyl-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	572
55dg		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(methylcarbamoyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	585
55dh		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-((2-morpholinoethyl)carbamoyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	685
55di		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(methanesulfonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	606
55dj		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-sulfamoyl-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	607
55dk		(S)-8-(2-amino-6-((R)-1-(4'-(dimethylcarbamoyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	600

55dl		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(piperazine-1-carbonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	641
55dm		(S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-propoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	605
55dn		(S)-8-(2-amino-6-((R)-1-(4'-ethoxy-3'-fluoro-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	591
55do		(S)-8-(2-amino-6-((R)-1-(4'-ethoxy-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	573
55dp		(S)-8-(2-amino-6-((R)-1-(4-(cinnolin-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	580
55dq		(S)-8-(2-amino-6-((R)-1-(4-(chroman-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid	584

Table 17b.

NMR Data for Compounds of Table 17a

Ex. No.	NMR
55a	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.66-7.58 (m, 6H), 7.45-7.41 (m, 2H), 7.36-7.32 (m, 1H), 6.64 (q, J = 6.8 Hz, 1H), 5.56 (s, 1H), 4.00 (m, 1H), 3.67-3.60 (m, 2H), 3.52-3.44 (m, 2H), 3.20-3.02 (m, 2H), 2.31-2.25 (m, 1H), 2.01 (m, 1H), 1.58 (s, 4H)
55b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.02 (s, 1 H), 7.92 (s, 1 H), 7.67-7.65 (m, 3 H), 7.61-7.55(m,4H), 6.60 (m, 1 H), 5.47 (s, 1 H), 3.97 (s, 4 H), 3.53 (m, 2 H), 3.35 (m, 2 H), 3.15-3.12 (m, 1 H), 3.13-3.00 (m, 1 H), 2.21 (m, 1 H), 1.95(m, 1H), 1.50 (m,4H)
55c	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.16 (s, 1 H), 7.89 (s, 1 H), 7.71-7.69 (d, 2 H), 7.63-7.60 (m, 4 H), 6.67-6.66 (q, 1 H), 5.52 (s, 1 H), 4.00 (m, 1 H), 3.87 (s, 3 H), 3.62 (m, 2 H), 3.44 (m, 2 H), 3.12 (d, 1 H), 3.06 (d, 1 H), 2.23 (m, 1 H), 1.99 (m, 1 H), 1.54(m, 4 H), 1.23 (m, 3 H).
55d	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.56 (s, 1H), 7.90 (s, 1H), 7.76-7.61 (m, 6H), 6.66 (q, J = 6.6 Hz, 1H), 5.59 (s, 1H), 4.17-4.13 (m,1H), 3.69-3.57 (m, 2H), 3.52-3.43 (m, 2H), 3.27-3.24 (m, 1H), 3.16-3.13 (m, 1H), 2.37-2.31 (m, 1H), 2.09-2.04 (m, 5H), 1.61 (m, 4H)
55e	¹ H NMR (400 MHz, MeOH -d ₄): δ ppm 7.62-7.56 (m, 4 H), 7.40 (m, 2 H), 7.15 (m, 1

	H), 6.61 (m, 1 H), 5.56 (m, 1 H), 4.07 (m, 1 H), 3.90 (s, 3 H), 3.63 (m, 2 H), 3.48 (m, 2 H), 3.25 (m, 1 H), 3.13 (m, 1 H), 2.30 (m, 1 H), 2.04 (m, 1 H), 1.60 (s, 4 H)
55f	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 9.09 (s, 1 H), 8.45 (s, 1 H), 8.26 (m, 1 H), 7.83-7.77 (m, 3 H), 7.64 (m, 2 H), 6.73 (m, 1 H), 6.06 (s, 1 H), 5.57 (s, 1 H), 3.45 (m, 4 H), 2.99 (m, 2 H), 2.10 (m, 1 H), 1.79 (m, 1 H), 1.42 (m, 4 H).
55g	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.66-7.56 (m, 6H), 7.19-7.17 (m, 2H), 6.66 (q, J = 6.7 Hz, 1H), 5.56-5.55 (m, 1H), 4.08 (m, 1H), 3.64-3.59 (m, 2H), 3.53-3.43 (m, 2H), 3.23-3.13 (m, 1H), 2.98-2.92 (m, 1H), 2.35-2.19 (m, 1H), 2.08-2.03 (m, 1H), 1.59 (m, 4H).
55h	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.06 (s, 1H), 7.83 (d, J = 8.3 Hz, 1H), 7.73 (d, J = 7.7 Hz, 3H), 7.62 (d, J = 7.5 Hz, 2H), 7.44 (d, J = 8.6 Hz, 1H), 6.67 (q, J = 7.4 Hz, 1H), 5.58 (s, 1H), 4.08 (m, 1H), 3.69-3.61 (m, 2H), 3.52-3.43 (m, 2H), 3.23-3.10 (m, 2H), 2.35-2.30 (m, 1H), 2.08-2.03 (m, 1H), 1.60 (s, 4H).
55i	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 8.03 (s, 1 H), 7.83-7.82 (d, 2 H, J=4.5 Hz), 7.81-7.79 (d, 2 H, J=7.6 Hz), 7.62-7.60 (d, 2 H, J=7.6 Hz), 7.43-7.41 (d, 2 H, J=8.6 Hz), 6.71-6.70 (q, 1 H, J=6.8 Hz), 5.55 (s, 1 H), 4.02 (s, 3 H), 3.71 (m, 1 H), 3.55-3.44 (m, 4 H), 2.85 (m, 1 H), 2.12 (m, 1 H), 1.71 (m, 1 H), 1.40 (m, 4 H).
55j	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.96 (s, 1 H), 8.33 (s, 1 H), 8.09-8.07 (d, 1 H, J=8.8 Hz), 7.81-7.79 (dd, 1 H, J=8.0 Hz), 7.71-7.69 (d, 2 H, J=8.0 Hz), 7.60-7.58 (d, 2 H, J=8.0 Hz), 6.63-6.58 (q, 1 H), 5.51 (s, 1 H), 4.00-3.96 (m, 1 H), 3.57 (m, 2 H), 3.40 (m, 2 H), 3.17-3.14 (d, 1 H, J=11.7 Hz), 3.13-3.00 (d, 1 H, J=11.7 Hz), 2.23-2.21 (m, 1 H), 1.99-1.94 (m, 1 H), 1.53 (m, 5 H).
55k	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 9.26 (s, 1 H), 8.33 (s, 1 H), 8.13-8.11 (d, 1 H, J=8.5 Hz), 7.84-7.82 (dd, 1 H, J=8.5 Hz), 7.76-7.74 (d, 2 H, J=8.3 Hz), 7.65-7.63 (d, 2 H, J=8.3 Hz), 6.67-6.65 (q, 1 H, J=7.2 Hz), 4.15-4.11 (m, 1 H), 3.53 (m, 2 H), 3.49 (m, 2 H), 3.27-3.24 (d, 1 H, J=11.7 Hz), 3.15-3.12 (d, 1 H, J=11.7 Hz), 3.33-2.31 (m, 1 H), 2.07-2.03 (m, 1 H), 1.61 (m, 5 H).
55l	¹ H-NMR (400 MHz, MeOH-d ₄): δ ppm 9.08 (s, 1H), 8.44 (s, 1H), 8.02-8.03 (m, 1H), 7.83-7.86 (m, 2H), 7.66-7.75 (m, 2H), 6.66-6.69 (m, 1H), 5.50 (s, 1H), 4.07-4.07 (m, 1H), 3.64-3.66 (m, 2H), 3.44-3.48 (m, 2H), 3.19-3.24 (m, 1H), 3.16-3.46 (m, 1H), 2.29-2.55 (m, 1H), 2.03-2.08 (m, 1H), 1.60-1.61 (m, 4H)
55m	¹ H-NMR (400 MHz, DMSO-d ₆): δ ppm 8.19 (s, 1H), 7.84-7.97 (m, 6H), 7.63-7.82 (m, 2H), 7.50-7.61 (m, 2H), 6.70-6.76 (m, 1H), 5.58 (s, 1H), 3.38-3.47 (s, 1H), 3.00-3.00 (m, 1H), 2.91-2.94 (m, 1H), 2.06-2.13 (m, 1H), 1.74-1.78 (m, 1H), 1.37-1.44 (m, 4H)
55n	¹ H-NMR (400 MHz, MeOH-d ₄): δ ppm 7.63-7.65 (m, 2H), 7.56-7.58 (m, 2H), 7.15-7.20 (m, 1H), 7.07-7.09 (m, 2H), 6.60-6.65 (m, 1H), 4.11-4.16 (m, 1H), 3.87 (s, 3H), 3.48-3.66 (m, 4H), 3.23-3.26 (m, 1H), 3.11-3.16 (m, 1H), 2.31-3.41 (m, 1H), 2.20 (s, 3H), 2.02-2.10 (m, 1H), 1.59-1.61 (m, 4H), 1.27-1.31 (m, 1H)
55o	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.62-7.56 (m, 4H), 7.00 (s, 1H), 6.93 (s, 1H), 6.74 (s, 1H), 6.64 (q, 1H, J=8.0), 5.56 (d, 1H, J=4.0), 4.08-4.04 (m, 1H), 3.80 (s, 3H), 3.63 (s, 2H), 3.47 (s, 2H), 3.23 (d, 1H, J = 16.0), 3.10 (d, 1H, J = 12.0 Hz), 2.36 (s, 3H), 2.04 (s, 1H), 1.59 (s, 1H), 1.28 (s, 1H)
55p	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (m, 1H), 1.30 (d, J = 15.5 Hz, 3H), 1.62 (d, J = 5.7 Hz, 5H), 2.06 (m, 1H), 2.14 (s, 3H), 2.33 (dd, J = 13.5, 9.2 Hz, 1H), 3.13 (d, J = 11.7 Hz, 1H), 3.25 (d, J = 11.3 Hz, 1H), 3.51 (dt, J = 20.9, 6.7 Hz, 2H), 3.66 (d, J =

	13.3 Hz, 2H), 3.76 (s, 3H), 4.09 (t, J = 8.2 Hz, 1H), 5.57 (s, 1H), 6.69 (m, 2H), 6.81 (dd, J = 8.4, 2.8 Hz, 1H), 7.15 (d, J = 8.4 Hz, 1H), 7.35 (m, 2H), 7.57 (d, J = 7.9 Hz, 2H)
55q	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.62 (d,2H,J=8.0), 7.56(d,2H,J=8.0), 7.18(m,2H), 7.02(d,1H,J=8.0), 6.62 (q, 1H,J=8.0), 5.55 (s, 1H), 4.01-3.97 (m, 1H), 3.31-3.23 (m, 1H), 3.89 (s, 3H), 3.86 (s, 3H), 3.67-3.60 (m, 2H),3.49-3.43(m, 2H) 3.19 (d, 1H, J = 12.0), 3.02 (d, 1H, J = 12.0 Hz),2.27(dd, 1H, J = 12.0,8.0), 2.00 (dd, 1H, J = 14.0,4.0), 1.58 (s, 1H),1.28 (s, 1H)
55r	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.70-7.61 (m, 4H), 7.28 (m, 3H), 6.67 (q, J = 7.6 Hz, 1H), 5.56 (s, 1H), 4.08 (m,1H), 3.91 (s, 3H), 3.63-3.46 (m, 8H), 3.13 (m, 1H), 2.35-2.29 (m, 1H), 2.01-1.99 (m, 1H), 1.97-1.87 (m, 5H), 1.59 (m, 4H).
55s	¹ H-NMR (400 MHz, MeOH-d ₄): δ ppm 1.74-1.73 (m,4H), 2.13-2.07 (m,1H), 2.52-2.46 (m,1H), 3.60-3.55 (m, 1H), 3.70-3.66 (m,2H), 4.55-4.50 (m,1H), 5.44 (s,2H), 6.66-6.63 (m,1H), 7.74-7.69 (m,2H), 7.88-7.80 (m,4H), 7.96-7.92 (m,1H)
55t	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.03 -8.05 (m,1H) ,7.95 (s, 1H), 7.84 -7.86 (m,1H), 7.71 -7.73 (m, 2H), 7.60 -7.62 (m, 2H),7.43 -7.45 (m, 1H), 6.62 -6.65 (m, 2H), 5.57 (s, 1H), 4.05 -4.10 (m,1H), 3.61-3.70 (m,3H), 3.42 -3.52 (m, 3H), 3.09 -3.12 (m, 1H), 2.29 -2.36 (m,1H), 2.02 -2.07 (m, 1H), 1.60 (m, 4H)
55u	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 8.02 (s,1H), 7.93 -7.98 (m, 2H), 7.76 -7.78 (m, 2H), 7.59 -7.63 (m, 3H), 6.63 -6.73 (m, 2H), 5.55 (s, 1H), 3.74 -3.79 (m,1H), 3.61 (s,3H), 3.32 -3.47 (m, 5H), 2.89 -3.07 (m,2H), 2.08 -2.14 (m,1H), 1.73 -1.80 (m, 1H), 1.41 (m, 4H).
55v	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.62(d,2H,J=8.0), 7.56 (d,2H,J=8.0), 7.45 (m,2H), 6.93 (d,1H,J=8.0), 6.62 (q, 1H,J=8.0), 5.56 (s, 1H), 4.07 (t, 1H, J=8.0), 3.64δ3.58 (m, 2H), 3.52-3.45(m, 2H), 3.24 (d, 1H, J = 12.0),3.11 (d, 1H, J = 8.0 Hz), 3.01 (t, 2H, J = 8.0), 2.59-2.57 (m, 2H), 2.32(dd, 1H, J = 12.0,8.0),2.05(dd, 1H, J = 12.0,8.0), 1.58 (s, 4H),1.28 (s, 1H)
55w	¹ H NMR (400 MHz, DMSO-d ₆): δ 1.59 (m, 4 H), 2.05-2.01 (m, 1 H, J=11.6 Hz), 2.32-2.28 (m, 1 H, J=11.6 Hz), 3.11-3.08 (d, 1 H), 3.25-3.22 (d, 1 H), 3.50-3.47 (m, 2 H), 3.68-3.65 (m, 2 H), 4.08-4.04 (q,1 H), 5.57 (s, 1 H), 6.66-6.65 (q, 1 H), 8.10 (s, 1 H), 7.61-7.60 (m, 3 H, J=8.6 Hz), 7.71-7.69 (m, 3 H, J=8.6 Hz), 8.01 (s, 1 H)
55x	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.61 (d, J = 5.7 Hz, 4H), 2.06 (dd, J = 13.5, 7.3 Hz, 1H), 2.35 (dd, J = 13.5, 9.2 Hz, 1H), 2.56 (s, 3H), 3.15 (d, J = 11.8 Hz, 1H), 3.25 (d, J = 11.7 Hz, 1H), 3.54 (m, 5H), 3.99 (s, 3H), 4.17 (t, J = 8.3 Hz, 1H), 5.00 (s, 1H), 6.66 (q, J = 7.1 Hz, 1H), 7.53 (d, J = 8.8 Hz, 1H), 7.61 (d, J = 8.1 Hz, 2H), 7.70 (m, 3H), 7.91 (s, 1H)
55y	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.54 (m, 4 H), 2.10 (m, 1 H), 2.26 (m, 4 H), 3.04 (m, 1 H), 3.17 (m, 1 H), 3.40 (m, 2 H), 3.56 (m, 2 H), 3.69 (s, 3 H), 4.07-4.03 (m, 1 H), 6.65-6.59 (m, 1 H), 6.93 (s, 1 H), 7.37-7.34(m, 1 H), 7.45-7.42 (m, 1 H), 7.57-7.55 (m, 2 H), 7.72-7.68 (m, 3 H)
55z	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.72-7.63(m,4H), 7.45(s,1H), 7.40 (s,1H), 7.18 (s,1H), 6.66 (q, 1H, J=8.0), 4.30 (d, 1H, J=8.0), 3.91 (s, 3H), 3.66 (s, 2H),3.55(s, 2H), 3.26 (s, 1H), 3.19(d, 1H, J = 12.0 Hz), 2.43-2.37 (m, 1H), 2.10-2.05 (m, 1H), 1.65 (s, 5H), 1.28 (s, 1H)
55aa	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.94 (d,1H,J=4.0), 7.71 (d,1H,J=4.0), 7.53-7.50 (m,2H), 6.82 (q,1H,J=8.0), 6.41 (d, 1H,J=4.0), 5.68 (d, 1H,J=4.0), 4.41 (s, 1H),

	3.09 (s, 1H), 2.75(t, 2H, J=8.0), 2.46 (d, 1H, J=16.0), 2.38 (s, 3H), 2.22(dd, 1H, J = 16.0,8.0), 1.23-1.19 (m, 3H)
55ab	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.60 (m, 4 H), 2.03-2.05 (m, 1 H), 2.33-2.32 (m, 1 H), 3.14-3.11 (d, 1 H, J=11.9 Hz), 3.26-3.23 (d, 1 H, J=11.9 Hz), 3.52-3.47 (m, 2 H), 3.65-3.54 (m, 2 H), 4.10-4.06 (m, 1H), 5.57 (s, 1 H), 6.65-6.64 (q, 1 H), 7.16-7.14 (d, 2 H, J=8.2 Hz), 7.51-7.44 (dd, 2 H, J=8.2 Hz), 7.51 (s, 1 H), 7.60-7.57 (d, 2 H, J=8.3 Hz), 7.65-7.63 (d, 2 H, J=8.3 Hz)
55ac	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.72-7.69 (m, 3 H), 7.58-7.56 (m, 2 H), 7.38 (s, 2 H), 7.02(m, 1 H), 6.61 (m, 1 H), 4.23(m, 1 H), 3.65 (m, 2 H), 3.48 (m, 2 H), 3.30 (m, 1 H), 3.14 (m, 1 H), 2.31 (m, 4H), 2.06 (m, 1 H), 1.62 (s, 4 H).
55ad	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.08 (d, J = 8.2 Hz, 1H), 7.77 (d, J = 8.2 Hz, 2H), 7.67-7.65 (m, 3H), 7.46 (m, 1H), 6.67-6.61 (m, 1H), 4.47 (m, 1H), 3.87 (s, 3H), 3.73-3.52 (m, 4H), 3.27-3.22 (m, 2H), 2.50-2.44 (m, 1H), 2.33 (s, 3H), 2.12-2.06 (m, 1H), 1.68 (m, 4H).
55ae	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.22 (d, J = 5.3 Hz, 2H), 1.42 (m, 4H), 1.82 (d, J = 13.2 Hz, 1H), 1.98 (dd, J = 17.4, 8.5 Hz, 1H), 2.91 (m, 1H), 3.03 (d, J = 11.0 Hz, 1H), 3.55 (s, 1H), 3.68 (s, 1H), 3.80 (s, 1H), 5.62 (s, 1H), 6.13 (s, 2H), 6.49 (d, J = 9.6 Hz, 1H), 6.75 (q, J = 7.3 Hz, 1H), 7.67 (m, 4H), 7.83 (dd, J = 22.1, 8.1 Hz, 3H), 8.09 (d, J = 9.5 Hz, 1H)
55af	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.01(d,J=2.6Hz,1H) ,7.89(dd, J=2.72, 6.8 Hz,1H), 7.58(m,4H), 6.64(m, 2H),5.56(s, 1H),4.08(m,1H) ,3.64(s,3H),3.53(m, 4H),3.12(m, 2H), 2.33(m,1H),2.06(m,1H),1.60(m,4H).
55ag	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.92 (d, J = 8.8 Hz, 1H), 7.65 (dd, J ₁ = 8.4 Hz, J ₂ = 31.9Hz, 4H), 7.17-7.14 (m, 2H), 6.66-6.63 (m, 1H), 4.14-4.10 (m, 1H), 3.66-3.59 (m, 2H), 3.54-3.43 (m, 2H), 3.26-3.24 (m, 1H), 3.15-3.12 (m, 1H), 2.37-2.32 (m, 1H), 2.08-2.03 (m, 1H), 1.61 (m, 4H)
55ah	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.16 (d,J = 8.92 Hz,1H), 8.02 (d,J=1.56 Hz, 1H),7.93-7.86 (m,2H), 7.76 (d,J=8.16 Hz, 2H), 7.64(d,J=8.08 Hz,2H),6.96 (d,J = 8.88 Hz,1H), 6.66 (q, J = 7.12 Hz,1H), 5.57(s,1H), 4.06(s,3H), 3.97 (m, 1H), 3.64(m, 2H), 3.47(m,2H), 3.17 (d, J=10.92 Hz,1H), 3.00 (d, J=12.04 Hz,1H), 2.26 (m, 1H), 2.01(m, 1H), 1.58 (s, 4H)
55ai	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.06 (d, J = 8.76 Hz,1H), 8.01 (s, 1H),7.93 (s ,2H),7.76 (d, J = 8.04 Hz, 2H),7.64 (d, J = 8.16 Hz, 2H), 7.31 (d, J = 8.76 Hz, 1H), 6.68 (q, J = 8.48 Hz, 1H), 5.58 (s, 1H), 4.08 (m, 1H), 3.63-3.50 (m, 4H), 3.24 (d, J = 11.64 Hz, 1H), 3.12 (d, J = 11.64 Hz,1H), 2.68(s, 3H), 2.31 (m, 1H), 2.05 (m, 1H), 1.59 (m, 4H).
55aj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.67(d,2H,J=8.0), 7.61-7.57(m,4H), 7.51(s,1H), 7.20(d,1H,J=8.0), 6.63 (q, 1H,J=8.0), 4.29 (t, 1H, J=12.0), 3.67δ3.58 (m, 2H), 3.53-3.48(m, 2H), 3.38(s,3H), 3.27 (d, 1H, J = 12.0),3.19 (d, 1H, J = 8.0 Hz), 2.99-2.95 (m, 2H), 2.65-2.63 (m, 2H), 2.40(dd, 1H, J = 12.0,8.0),2.09(dd, 1H, J = 12.0,8.0), 1.66 (s, 5H),1.31 (s, 2H)
55ak	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.52 - 1.67 (m, 4 H) 2.05 (dd, J=13.42, 7.13 Hz, 1 H) 2.32 (dd, J=13.54, 9.30 Hz, 1 H) 3.07 - 3.16 (m, 1 H) 3.24 (d, J=11.76 Hz, 1 H) 3.38 - 3.55 (m, 2 H) 3.56 - 3.76 (m, 2 H) 4.10 (t, J=8.18 Hz, 1 H) 5.56 (s, 1 H) 6.63 (q, J=6.96 Hz, 1 H) 7.06 (qd, J=5.74, 3.29 Hz, 1 H) 7.35 (dd, J=9.30, 1.44 Hz, 1 H) 7.38 - 7.48 (m, 2 H) 7.55 - 7.69 (m, 4 H)

55al	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.59 (d, J=4.59 Hz, 4 H) 2.04 (dd, J=13.50, 7.30 Hz, 1 H) 2.31 (dd, J=13.30, 9.01 Hz, 1 H) 3.05 - 3.25 (m, 2 H) 3.37 - 3.53 (m, 2 H) 3.54 - 3.69 (m, 2 H) 3.81 (s, 3 H) 4.05 (dd, J=9.18, 7.32 Hz, 1 H) 5.54 (s, 1 H) 6.61 (q, J=7.29 Hz, 1 H) 6.89 (dd, J=7.83, 2.12 Hz, 1 H) 7.05 - 7.21 (m, 2 H) 7.26 - 7.38 (m, 1 H) 7.46 - 7.68 (m, 4 H)
55am	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.63 (d, J=5.08 Hz, 4 H) 2.08 (dd, J=13.32, 7.18 Hz, 1 H) 2.35 (dd, J=13.32, 9.22 Hz, 1 H) 3.07 - 3.20 (m, 1 H) 3.27 (d, J=11.71 Hz, 1 H) 3.40 - 3.58 (m, 2 H) 3.59 - 3.80 (m, 2 H) 4.11 (t, J=7.96 Hz, 1 H) 5.58 (s, 1 H) 6.68 (d, J=7.13 Hz, 1 H) 7.10 (dt, J=6.00, 2.95 Hz, 1 H) 7.38 (d, J=10.35 Hz, 1 H) 7.42 - 7.52 (m, 2 H) 7.56 - 7.79 (m, 4 H)
55an	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.48 - 1.66 (m, 4 H) 2.01 (dd, J=13.37, 7.17 Hz, 1 H) 2.28 (dd, J=13.35, 9.20 Hz, 1 H) 3.06 (d, J=11.71 Hz, 1 H) 3.20 (d, J=11.67 Hz, 1 H) 3.35 - 3.52 (m, 2 H) 3.53 - 3.69 (m, 2 H) 3.81 (s, 3 H) 4.02 (dd, J=9.15, 7.20 Hz, 1 H) 5.53 (s, 1 H) 6.61 (q, J=7.21 Hz, 1 H) 6.89 (dd, J=8.20, 2.49 Hz, 1 H) 7.06 - 7.21 (m, 2 H) 7.26 - 7.38 (m, 1 H) 7.47 - 7.68 (m, 4 H)
55ao	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.63 - 1.87 (m, 4 H) 2.10 (dd, J=13.72, 8.69 Hz, 1 H) 2.48 - 2.62 (m, 1 H) 3.33 (s, 0 H) 3.52 - 3.81 (m, 4 H) 3.84 (s, 3 H) 4.55 (t, J=8.81 Hz, 1 H) 5.93 (s, 0 H) 6.59 (d, J=6.25 Hz, 1 H) .71 (dt, J=10.75, 2.26 Hz, 1 H) 6.90 - 7.02 (m, 2 H) 7.51 - 7.80 (m, 4 H).
55ap	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.63 - 1.85 (m, 4 H) 2.10 (dd, J=13.64, 8.66 Hz, 1 H) 2.50 (dd, J=13.64, 8.81 Hz, 1 H) 3.51 - 3.88 (m, 5 H) 4.55 (t, J=8.71 Hz, 1 H) 5.93 (s, 1 H) 6.63 (q, J=6.61 Hz, 1 H) 6.96 (tt, J=9.06, 2.31 Hz, 1 H) 7.20 - 7.33 (m, 2 H) 7.62 - 7.80 (m, 4 H).
55aq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.60 - 1.87 (m, 4 H) 2.09 (dd, J=13.64, 8.76 Hz, 1 H) 2.50 (dd, J=13.59, 8.86 Hz, 1 H) 3.51 - 3.79 (m, 4 H) 3.81 (s, 3 H) 4.56 (t, J=8.74 Hz, 1 H) 5.88 - 5.99 (m, 1 H) 5.93 (s, 1 H) 6.57 (d, J=6.39 Hz, 1 H) 6.94 - 7.04 (m, 2 H) 7.50 - 7.70 (m, 6 H)
55ar	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.67 - 1.87 (m, 4 H) 2.10 (dd, J=13.64, 8.71 Hz, 1 H) 2.51 (dd, J=13.59, 8.86 Hz, 1 H) 3.54 - 3.77 (m, 4 H) 3.78 (s, 4 H) 4.57 (t, J=8.76 Hz, 1 H) 5.95 (s, 1 H) 6.59 (q, J=6.22 Hz, 1 H) 7.00 (td, J=7.46, 0.95 Hz, 1 H) 7.06 (d, J=8.10 Hz, 1 H) 7.19 - 7.38 (m, 2 H) 7.48 - 7.66 (m, 4 H)
55as	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.60 - 1.86 (m, 4 H) 2.10 (dd, J=13.64, 8.66 Hz, 1 H) 2.50 (dd, J=13.62, 8.83 Hz, 1 H) 3.53 - 3.87 (m, 4 H) 4.54 (t, J=8.71 Hz, 1 H) 5.93 (s, 0 H) 6.64 (q, J=6.65 Hz, 1 H) 7.61 - 7.72 (m, 4 H) 7.73 - 7.80 (m, 2 H) 7.84 - 7.94 (m, 2 H)
55at	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.67 - 1.84 (m, 4 H) 2.10 (dd, J=13.67, 8.69 Hz, 1 H) 2.51 (dd, J=13.69, 8.81 Hz, 1 H) 3.55 - 3.82 (m, 4 H) 4.56 (t, J=8.76 Hz, 1 H) 5.95 (s, 1 H) 6.63 (q, J=6.56 Hz, 1 H) 7.29 (dt, J=8.19, 1.15 Hz, 1 H) 7.48 - 7.58 (m, 2 H) 7.61 - 7.76 (m, 5 H)
55au	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.39 (t, J=7.00 Hz, 3 H) 1.65 - 1.86 (m, 4 H) 2.09 (dd, J=13.64, 8.76 Hz, 1 H) 2.50 (dd, J=13.62, 8.79 Hz, 1 H) 3.51 - 3.84 (m, 4 H) 4.07 (q, J=6.98 Hz, 2 H) 4.57 (t, J=8.74 Hz, 1 H) 5.94 (s, 1 H) 6.61 (q, J=6.57 Hz, 1 H) 6.82 - 6.95 (m, 1 H) 7.06 - 7.20 (m, 2 H) 7.28 - 7.43 (m, 1 H) 7.55 - 7.73 (m, 4 H)
55av	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (d, J=6.05 Hz, 6 H) 1.65 - 1.86 (m, 4 H) 2.09 (dd, J=13.67, 8.69 Hz, 1 H) 2.50 (dd, J=13.64, 8.86 Hz, 1 H) 3.48 - 3.85 (m, 4 H) 4.55 (t, J=8.71 Hz, 1 H) 4.65 (dt, J=12.08, 6.06 Hz, 1 H) 5.92 (s, 1 H) 6.59 (q, J=6.43

	Hz, 1 H) 6.91 (dd, J=8.22, 1.93 Hz, 1 H) 7.11 (t, J=2.03 Hz, 1 H) 7.15 (d, J=7.71 Hz, 1 H) 7.29 - 7.39 (m, 1 H) 7.56 - 7.73 (m, 4 H)
55aw	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.62 - 1.93 (m, 4 H) 2.11 (dd, J=13.69, 8.76 Hz, 1 H) 2.54 (dd, J=13.69, 8.86 Hz, 1 H) 3.46 - 4.00 (m, 4 H) 4.59 (t, J=8.74 Hz, 1 H) 6.01 (s, 1 H) 6.74 (q, J=6.65 Hz, 1 H) 7.75 - 8.02 (m, 4 H) 8.20 (dd, J=8.22, 5.78 Hz, 1 H) 8.87 (d, J=5.71 Hz, 1 H) 8.97 (dt, J=8.27, 1.72 Hz, 1 H) 9.24 (d, J=2.00 Hz, 1 H)
55ax	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.64 - 1.90 (m, 4 H) 2.11 (dd, J=13.67, 8.79 Hz, 1 H) 2.52 (dd, J=13.64, 8.86 Hz, 1 H) 3.49 - 4.02 (m, 4 H) 4.57 (t, J=8.71 Hz, 1 H) 6.01 (s, 1 H) 6.76 (d, J=6.54 Hz, 1 H) 7.85 (d, J=8.35 Hz, 2 H) 8.09 (d, J=8.49 Hz, 2 H) 8.33 - 8.54 (m, 2 H) 8.80 - 8.99 (m, 2 H)
55ay	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.56 - 1.91 (m, 4 H) 2.10 (dd, J=13.69, 8.66 Hz, 1 H) 2.51 (dd, J=13.81, 8.83 Hz, 1 H) 3.53 - 3.94 (m, 5 H) 4.56 (dt, J=8.48, 4.37 Hz, 1 H) 5.89 - 6.14 (m, 1 H) 6.51 - 6.82 (m, 1 H) 7.48 - 8.01 (m, 4 H) 9.19 (s, 1 H) 9.24 (s, 1 H)
55az	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.60 (t, J=5.08 Hz, 4 H) 2.00 (s, 2 H) 2.03 - 2.13 (m, 1 H) 2.26 - 2.39 (m, 1 H) 2.52 - 2.64 (m, 4 H) 3.07 - 3.18 (m, 1 H) 3.26 (d, J=11.71 Hz, 1 H) 3.39 - 3.57 (m, 2 H) 3.57 - 3.77 (m, 2 H) 4.01 - 4.20 (m, 1 H) 5.58 (s, 1 H) 6.67 (s, 1 H) 7.40 (dd, J=8.49, 1.07 Hz, 1 H) 7.59 - 7.68 (m, 3 H) 7.69 - 7.79 (m, 3 H)
55ba	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.47 - 1.74 (m, 4 H) 1.99 - 2.13 (m, 1 H) 2.26 - 2.40 (m, 1 H) 2.55 (s, 3 H) 3.07 - 3.19 (m, 1 H) 3.21 3.29 (m, 1H) 3.48 (d, J=4.88 Hz, 2 H) 3.65 (d, J=3.32 Hz, 2 H) 4.01 (s, 3 H) 4.10 (dd, J=8.98, 7.22 Hz, 1 H) 5.59 (s, 1 H) 6.69 (d, J=7.03 Hz, 1 H) 7.41 (dd, J=8.49, 1.07 Hz, 1 H) 7.58 - 7.70 (m, 3 H) 7.70 - 7.84 (m, 3 H)
55bb	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.58 (br. s., 4 H) 1.95 - 2.09 (m, 1 H) 2.24 - 2.38 (m, 1 H) 2.63 (s, 3 H) 3.01 - 3.14 (m, 1 H) 3.17 3.25 (m, 1 H) 3.38-3.54 (m, 2 H) 3.55-3.74 (m, 2H)
55bc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.62 (br. s., 4 H) 1.97 - 2.09 (m, 1 H) 2.24 - 2.36 (m, 1 H) 3.07 (s, 3 H) 3.18 - 3.27 (m, 1 H) 3.55 (s, 4 H) 3.60 - 3.75 (m, 2 H) 3.96 - 4.07 (m, 1 H) 5.59 (s, 1 H), 6.61 - 6.75 (m, 1 H) 7.54 - 7.61 (m, 1 H) 7.66 (s, 3 H) 7.72 (s, 2 H) 7.95 - 8.08 (m, 1 H)
55bd	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.50 - 1.68 (m, 4 H) 1.86 - 2.01 (m, 1 H) 2.11 - 2.28 (m, 1 H) 2.77 - 2.91 (m, 1 H) 3.04 - 3.13 (m, 1 H) 3.40 - 3.57 (m, 2 H) 3.59 - 3.74 (m, 2 H) 3.76 - 3.88 (m, 1 H) 5.54 - 5.66 (m, 1 H) 6.61 - 6.79 (m, 1 H) 7.67 - 7.77 (m, 2 H) 7.81 - 7.97 (m, 3 H) 7.99 - 8.09 (m, 1 H) 8.16 - 8.27 (m, 2 H) 8.41 - 8.53 (m, 1 H) 9.21 - 9.33 (m, 1 H)
55be	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.50 - 1.65 (m, 4 H) 1.92 - 2.00 (m, 1 H) 2.14 - 2.28 (m, 1 H) 2.84 - 2.94 (m, 1 H) 3.04 - 3.16 (m, 1 H) 3.40 - 3.57 (m, 2 H) 3.58 - 3.73 (m, 2 H) 3.79 - 3.91 (m, 1 H) 5.59 (s, 1 H) 6.62 - 6.78 (m, 1 H) 7.64 - 7.75 (m, 2 H) 7.86 (d, J=8.59 Hz, 3 H) 8.01 - 8.09 (m, 1 H) 8.10 - 8.20 (m, 1 H) 8.35 - 8.42 (m, 1 H) 8.43 - 8.48 (m, 1 H) 9.25 - 9.37 (m, 1 H)
55bf	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.61 (br. s., 4 H) 1.94 - 2.04 (m, 1 H) 2.28 (s, 7 H) 2.92 - 3.06 (m, 1 H) 3.11 - 3.23 (m, 1 H) 3.53 (s, 4 H) 3.59 - 3.75 (m, 2 H) 3.89 - 4.02 (m, 1 H) 5.58 (s, 1 H) 6.60 - 6.70 (m, 1 H) 7.43 (s, 2 H) 7.57 - 7.74 (m, 6 H)
55bg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.56 (d, J=4.69 Hz, 4 H) 1.78 - 1.95 (m, 1 H) 2.07 - 2.22 (m, 1 H) 2.69 - 2.83 (m, 1 H) 2.96 - 3.09 (m, 1 H) 3.38 - 3.54 (m, 2 H) 3.56


	- 3.69 (m, 2 H) 3.70 - 3.79 (m, 1 H) 5.57 (s, 1 H) 6.59 - 6.77 (m, 1 H) 7.51 - 7.61 (m, 1 H) 7.68 (d, J=8.00 Hz, 2 H) 7.83 (d, J=8.20 Hz, 2 H) 8.11 (s, 2 H) 8.20 (s, 1 H) 8.38 - 8.50 (m, 1 H) 8.77 - 8.92 (m, 1 H)
55bh	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.57 (br. s., 4 H) 1.84 - 2.03 (m, 1 H) 2.12 - 2.29 (m, 1 H) 2.91 (s, 1 H) 3.04 - 3.16 (m, 1 H) 3.38 - 3.55 (m, 2 H) 3.56 - 3.73 (m, 2 H) 3.78 - 3.96 (m, 1 H) 5.58 (s, 1 H) 6.60 - 6.80 (m, 1 H) 7.47 - 7.58 (m, 1 H) 7.69 (d, J=8.20 Hz, 2 H) 7.83 (d, J=8.20 Hz, 2 H) 7.93 (d, J=1.17 Hz, 1 H) 8.02 (d, J=8.59 Hz, 1 H) 8.25 (s, 1 H) 8.37 (s, 1 H) 8.87 (d, J=2.93 Hz, 1 H)
55bi	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.56 (d, J=5.08 Hz, 4 H) 1.76 - 1.88 (m, 1 H) 2.05 - 2.20 (m, 1 H) 2.63 - 2.81 (m, 1 H) 2.94 - 3.07 (m, 1 H) 3.37 - 3.54 (m, 2 H) 3.55 - 3.79 (m, 3 H) 5.58 (s, 1H) 6.61 - 6.78 (m, 1 H) 7.71 (d, J=8.20 Hz, 2 H) 7.86 (d, J=8.40 Hz, 2 H) 8.17 (s, 2 H) 8.32 (s, 1 H) 8.89 (dd, J=12.98, 1.66 Hz, 2 H)
55bj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.54 - 1.72 (m, 4 H) 2.01 - 2.13 (m, 1 H) 2.27 - 2.41 (m, 1 H) 3.06 - 3.16 (m, 3 H) 3.18 (s, 3 H) 3.22 - 3.30 (m, 1 H) 3.41 - 3.59 (m, 2 H) 3.67 (s, 4 H) 4.02 - 4.16 (m, 1 H) 5.53 - 5.66 (m, 1 H) 6.61 - 6.74 (m, 1 H) 7.54 - 7.57 (m, 1 H) 7.61 - 7.67 (m, 3 H) 7.70 - 7.80 (m, 2 H) 7.96 - 8.06 (m, 1 H)
55bk	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.54 - 1.72 (m, 4 H) 2.05 - 2.18 (m, 1 H) 2.29 - 2.43 (m, 1 H) 3.08 - 3.20 (m, 1 H) 3.24 - 3.29 (m, 1 H) 3.43 - 3.76 (m, 4 H) 4.05 - 4.17 (m, 1 H) 5.57 - 5.67 (m, 1 H) 6.64 - 6.79 (m, 1 H) 7.65 - 7.76 (m, 2 H) 7.83 - 7.94 (m, 2 H) 8.08 - 8.19 (m, 1 H) 8.32 - 8.48 (m, 2 H) 9.21 - 9.33 (m, 1 H) 9.56 - 9.67 (m, 1 H)
55bp	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.60 (q, J = 5.9, 5.2 Hz, 4H), 1.98 (m, 5H), 2.32 (dd, J = 13.4, 9.3 Hz, 1H), 3.12 (d, J = 11.8 Hz, 1H), 3.25 (d, J = 11.7 Hz, 1H), 3.50 (m, 4H), 3.61 (m, 4H), 4.08 (dd, J = 9.2, 7.2 Hz, 1H), 5.57 (s, 1H), 6.67 (q, J = 7.1 Hz, 1H), 7.61 (ddd, J = 356.8, 7.9, 5.7 Hz, 4H), 7.71 (m, 4H)
55cc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.09 (d, J = 1.2 Hz, 0H), 1.28 (s, 0H), 1.61 (q, J = 5.3 Hz, 0H), 1.92 (m, 0H), 2.05 (dd, J = 13.3, 7.3 Hz, 0H), 2.32 (m, 0H), 3.10 (s, 0H), 3.29 (s, 2H), 3.53 (s, 0H), 3.65 (s, 1H), 3.80 (dd, J = 16.8, 1.2 Hz, 0H), 4.07 (t, J = 8.2 Hz, 0H), 4.89 (d, J = 3.8 Hz, 0H), 5.56 (d, J = 1.2 Hz, 0H), 5.99 (m, 0H), 6.60 (m, 0H), 7.21 (dd, J = 8.4, 1.2 Hz, 0H), 7.48 (d, J = 1.2 Hz, 1H)
55ce	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.58 (q, J = 6.7, 6.1 Hz, 4H), 1.94 (dd, J = 13.2, 9.1 Hz, 1H), 2.39 (dd, J = 13.3, 8.6 Hz, 1H), 3.20 (s, 2H), 3.80 (m, 3H), 4.54 (t, J = 8.4 Hz, 1H), 5.74 (s, 1H), 6.31 (s, 2H), 6.90 (q, J = 7.2 Hz, 1H), 7.49 (t, J = 6.5 Hz, 3H), 7.76 (m, 6H), 9.01 (dt, J = 21.8, 11.9 Hz, 1H), 9.74 (d, J = 11.9 Hz, 1H)
55cg	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.59 (m, 4H), 2.00 (s, 7H), 2.41 (m, 1H), 3.20 (s, 2H), 3.60 (m, 4H), 3.83 (m, 1H), 4.53 (d, J = 8.7 Hz, 1H), 5.74 (s, 1H), 6.34 (dd, J = 28.1, 14.7 Hz, 2H), 6.91 (q, J = 7.4 Hz, 1H), 7.24 (m, 5H), 7.67 (d, J = 7.8 Hz, 2H), 8.99 (s, 1H), 9.77 (d, J = 8.4 Hz, 1H)
55cj	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.29 (d, J = 6.8 Hz, 6H), 1.58 (m, 4H), 1.94 (dd, J = 13.2, 9.2 Hz, 1H), 2.39 (dd, J = 13.4, 8.6 Hz, 1H), 2.99 (hept, J = 7.1 Hz, 1H), 3.19 (s, 2H), 3.59 (m, 4H), 4.55 (d, J = 9.2 Hz, 1H), 5.73 (s, 1H), 6.28 (m, 2H), 6.81 (q, J = 7.4 Hz, 1H), 7.41 (d, J = 7.9 Hz, 2H), 7.65 (m, 4H), 7.78 (d, J = 8.0 Hz, 2H), 8.97 (d, J = 13.6 Hz, 1H), 9.76 (s, 1H)
55ck	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.99 (s, 1H), 1.31 (d, J = 6.9 Hz, 7H), 1.58 (m, 4H), 1.94 (dd, J = 13.2, 9.2 Hz, 1H), 2.39 (dd, J = 13.4, 8.6 Hz, 1H), 3.02 (hept, J = 7.2 Hz, 1H), 3.19 (s, 2H), 3.55 (ddd, J = 19.7, 12.2, 5.9 Hz, 2H), 3.86 (s, 2H), 4.05 (s, 1H), 4.54 (q, J = 8.9, 6.7 Hz, 1H), 5.73 (s, 1H), 6.31 (m, 2H), 6.81 (q, J = 7.3 Hz, 1H), 7.46

	(m, 5H), 7.65 (d, J = 8.0 Hz, 2H), 7.79 (d, J = 8.0 Hz, 2H), 8.96 (dt, J = 20.8, 8.6 Hz, 1H), 9.77 (d, J = 8.5 Hz, 1H)
55cp	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.29 (s, 1H), 1.57 (dd, J = 9.4, 5.0 Hz, 12H), 1.94 (dd, J = 13.3, 9.1 Hz, 3H), 2.28 (s, 9H), 2.41 (m, 4H), 2.61 (d, J = 7.4 Hz, 1H), 3.19 (s, 7H), 3.59 (m, 13H), 4.15 (s, 1H), 4.55 (d, J = 8.4 Hz, 4H), 5.75 (s, 3H), 6.38 (m, 5H), 6.87 (q, J = 7.4 Hz, 3H), 7.46 (m, 22H), 8.98 (m, 3H), 9.76 (m, 3H)
55cq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.60 (m, 4H), 2.05 (m, 1H), 2.34 (s, 7H), 3.12 (d, J = 11.5 Hz, 1H), 3.24 (d, J = 11.5 Hz, 1H), 3.49 (m, 2H), 3.64 (dq, J = 12.6, 6.5, 4.7 Hz, 2H), 4.08 (t, J = 8.2 Hz, 1H), 5.56 (s, 1H), 6.64 (q, J = 7.1 Hz, 1H), 6.99 (s, 1H), 7.21 (s, 2H), 7.59 (m, 5H)
55cr	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.29 (s, 1H), 1.58 (dq, J = 11.6, 7.5, 7.1 Hz, 11H), 1.94 (dd, J = 13.3, 9.1 Hz, 3H), 2.15 (s, 0H), 2.39 (dd, J = 13.3, 8.6 Hz, 3H), 2.61 (d, J = 8.8 Hz, 3H), 3.05 (s, 0H), 3.19 (s, 6H), 3.40 (s, 1H), 3.54 (h, J = 6.4 Hz, 3H), 3.81 (m, 4H), 4.01 (dd, J = 19.9, 11.7 Hz, 1H), 4.16 (s, 1H), 4.54 (t, J = 8.6 Hz, 3H), 5.73 (s, 3H), 6.28 (d, J = 15.5 Hz, 5H), 6.81 (q, J = 7.3 Hz, 3H), 7.63 (m, 11H), 7.86 (dd, J = 17.4, 7.8 Hz, 8H), 8.99 (dq, J = 23.7, 15.7, 12.4 Hz, 2H), 9.72 (s, 3H)
55cs	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.57 (m, J = 8.0, 6.2 Hz, 8H), 1.94 (dd, J = 13.2, 9.2 Hz, 2H), 2.23 (s, 6H), 2.34 (s, 8H), 3.20 (s, 5H), 3.57 (dp, J = 22.0, 7.4, 6.0 Hz, 9H), 4.54 (t, J = 8.4 Hz, 3H), 5.76 (s, 2H), 6.34 (d, J = 14.9 Hz, 1H), 6.45 (s, 1H), 6.86 (q, J = 7.4 Hz, 2H), 7.18 (m, 6H), 7.56 (dd, J = 46.1, 7.8 Hz, 8H), 8.99 (m, 2H), 9.77 (d, J = 15.0 Hz, 2H)
55ct	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.97 (t, J = 7.3 Hz, 3H), 1.50 (m, 9H), 1.94 (dd, J = 13.3, 9.2 Hz, 1H), 2.39 (dd, J = 13.3, 8.6 Hz, 1H), 2.67 (q, J = 6.9, 6.2 Hz, 2H), 3.19 (s, 2H), 3.57 (m, 5H), 3.85 (m, 1H), 4.06 (d, J = 16.1 Hz, 1H), 4.54 (m, 1H), 5.74 (s, 1H), 6.31 (s, 2H), 6.81 (q, J = 7.3 Hz, 1H), 7.35 (d, J = 7.8 Hz, 2H), 7.64 (d, J = 7.8 Hz, 4H), 7.78 (d, J = 8.0 Hz, 2H), 8.96 (m, 1H), 9.73 (d, J = 8.2 Hz, 1H)
55cv	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.29 (s, 1H), 1.38 (s, 1H), 1.58 (qd, J = 12.6, 8.1, 7.6 Hz, 11H), 1.95 (m, 8H), 2.43 (m, 5H), 2.94 (t, J = 10.0 Hz, 1H), 3.08 (dd, J = 16.3, 7.0 Hz, 1H), 3.19 (s, 6H), 3.33 (s, 8H), 3.52 (m, 4H), 3.82 (m, 5H), 4.11 (m, 4H), 4.53 (dt, J = 12.6, 6.0 Hz, 3H), 5.75 (s, 2H), 6.20 (s, 1H), 6.28 (d, J = 9.3 Hz, 2H), 6.48 (s, 3H), 6.84 (q, J = 7.3 Hz, 2H), 7.56 (s, 1H), 7.93 (m, 21H), 8.98 (dd, J = 13.6, 8.0 Hz, 2H), 9.71 (m, 2H)
55cw	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.29 (s, 1H), 1.58 (dd, J = 7.3, 4.2 Hz, 8H), 1.94 (dd, J = 13.2, 9.1 Hz, 2H), 2.14 (d, J = 1.4 Hz, 0H), 2.40 (s, 8H), 2.49 (d, J = 9.3 Hz, 0H), 3.03 (m, 1H), 3.19 (s, 5H), 3.55 (m, 3H), 3.81 (t, J = 8.1 Hz, 0H), 4.00 (m, 8H), 4.24 (m, 0H), 4.35 (m, 1H), 4.54 (m, 2H), 5.74 (s, 2H), 6.31 (m, 3H), 6.80 (q, J = 7.3 Hz, 2H), 7.34 (d, J = 7.8 Hz, 4H), 7.64 (d, J = 8.1 Hz, 8H), 7.78(m, 4H), 8.99 (q, J = 8.5, 7.4 Hz, 2H), 9.74 (s, 2H)
55cx	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.29 (s, 1H), 1.59 (m, 11H), 1.94 (dd, J = 13.3, 9.1 Hz, 3H), 2.43 (s, 11H), 2.61 (d, J = 9.0 Hz, 1H), 3.19 (s, 6H), 3.58 (m, 11H), 4.13 (s, 1H), 4.54 (m, 6H), 5.75 (s, 2H), 6.26 (s, 1H), 6.34 (s, 1H), 6.42 (s, 2H), 6.81 (q, J = 7.3 Hz, 3H), 7.27 (d, J = 7.5 Hz, 3H), 7.58 (m, 20H), 8.98 (s, 3H), 9.77 (d, J = 9.9 Hz, 3H)
55db	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.24 (m, 8H), 1.76 (dd, J = 12.6, 6.8 Hz, 9H), 2.01 (s, 3H), 2.12 (m, 4H), 2.51 (dd, J = 13.6, 8.8 Hz, 3H), 2.63 (td, J = 7.6, 5.4 Hz, 4H), 2.90 (m, 6H), 3.36 (d, J = 12.8 Hz, 7H), 3.63 (dt, J = 11.5, 5.0 Hz, 4H), 3.76 (m,

	4H), 4.11 (qd, J = 7.1, 3.6 Hz, 4H), 4.56 (t, J = 8.7 Hz, 3H), 4.94 (s, 2H), 6.62 (dq, J = 19.2, 6.7 Hz, 3H), 7.02 (dt, J = 6.2, 2.0 Hz, 3H), 7.16 (m, 2H), 7.38 (m, 2H), 7.65 (m, 9H), 7.84 (m, 1H)
55dc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.58 (t, J = 5.4 Hz, 4H), 2.02 (dd, J = 13.4, 7.0 Hz, 1H), 2.29 (dd, J = 13.4, 9.1 Hz, 1H), 3.08 (d, J = 11.6 Hz, 1H), 3.21 (d, J = 11.5 Hz, 1H), 3.46 (ddt, J = 20.6, 13.2, 5.7 Hz, 2H), 3.61 (d, J = 16.6 Hz, 2H), 4.05 (t, J = 8.1 Hz, 1H), 4.94 (s, 10H), 5.58 (s, 1H), 6.69 (q, J = 7.2 Hz, 1H), 7.66 (d, J = 8.0 Hz, 2H), 7.79 (d, J = 7.9 Hz, 2H), 8.00 (dd, J = 8.9, 1.9 Hz, 1H), 8.12 (m, 3H), 8.79 (d, J = 2.7 Hz, 1H)
55dd	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.05 (t, J = 7.4 Hz, 3H), 1.29 (d, J = 5.8 Hz, 1H), 1.59 (q, J = 5.8, 5.0 Hz, 4H), 1.80 (h, J = 6.9 Hz, 2H), 2.04 (dd, J = 13.5, 7.1 Hz, 1H), 2.32 (dd, J = 13.4, 9.2 Hz, 1H), 3.11 (d, J = 11.8 Hz, 1H), 3.23 (d, J = 12.0 Hz, 1H), 3.47 (ddt, J = 20.6, 13.1, 6.1 Hz, 2H), 3.64 (m, 2H), 3.96 (t, J = 6.4 Hz, 2H), 4.07 (dd, J = 9.1, 7.3 Hz, 1H), 5.55 (s, 1H), 6.62 (q, J = 7.1 Hz, 1H), 6.97 (m, 2H), 7.56 (m, 6H)
55de	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.15 (t, J = 7.0 Hz, 3H), 1.26 (t, J = 7.1 Hz, 3H), 1.61 (q, J = 5.9, 5.1 Hz, 4H), 2.05 (dd, J = 13.4, 7.2 Hz, 1H), 2.32 (dd, J = 13.4, 9.2 Hz, 1H), 3.11 (d, J = 11.8 Hz, 1H), 3.24 (d, J = 11.6 Hz, 1H), 3.35 (m, 3H), 3.56 (dddd, J = 47.5, 27.9, 14.5, 7.8 Hz, 6H), 4.07 (dd, J = 9.2, 7.1 Hz, 1H), 4.92 (s, 17H), 5.57 (s, 1H), 6.66 (q, J = 7.1 Hz, 1H), 7.46 (m, 2H), 7.62 (d, J = 8.1 Hz, 2H), 7.71 (m, 4H)
55df	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.64 (m, 4H), 2.07 (dd, J = 13.5, 7.6 Hz, 1H), 2.39 (dd, J = 13.5, 9.0 Hz, 1H), 3.17 (d, J = 11.7 Hz, 1H), 3.27 (d, J = 12.0 Hz, 2H), 3.53 (dt, J = 22.9, 7.5 Hz, 2H), 3.67 (td, J = 13.8, 13.2, 6.4 Hz, 2H), 4.26 (t, J = 8.3 Hz, 1H), 4.87 (m, 3H), 6.66 (q, J = 7.0 Hz, 1H), 7.63 (d, J = 8.1 Hz, 2H), 7.74 (m, 4H), 7.96 (m, 2H)
55dg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.31 (dd, J = 17.3, 3.8 Hz, 2H), 1.62 (m, 4H), 2.06 (dd, J = 13.5, 7.4 Hz, 1H), 2.36 (dd, J = 13.5, 9.1 Hz, 1H), 2.93 (s, 3H), 3.15 (d, J = 11.9 Hz, 1H), 3.26 (d, J = 11.6 Hz, 1H), 3.51 (m, 2H), 3.64 (dq, J = 12.1, 6.4, 5.1 Hz, 2H), 4.18 (dd, J = 9.1, 7.5 Hz, 1H), 4.93 (d, J = 1.7 Hz, 19H), 6.66 (q, J = 7.1 Hz, 1H), 7.63 (d, J = 8.1 Hz, 2H), 7.72 (dd, J = 8.3, 5.9 Hz, 4H), 7.89 (m, 2H)
55dh	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.60 (q, J = 6.0, 5.0 Hz, 4H), 2.03 (dd, J = 13.4, 7.2 Hz, 1H), 2.31 (dd, J = 13.4, 9.2 Hz, 1H), 2.59 (dt, J = 24.2, 5.7 Hz, 6H), 3.10 (d, J = 11.7 Hz, 1H), 3.23 (d, J = 11.9 Hz, 1H), 3.62 (m, 10H), 4.06 (dd, J = 9.2, 7.2 Hz, 1H), 4.87 (s, 1H), 5.57 (s, 1H), 6.66 (q, J = 7.1 Hz, 1H), 7.62 (d, J = 8.1 Hz, 2H), 7.72 (m, 4H), 7.91 (m, 2H)
55di	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.61 (d, J = 5.5 Hz, 4H), 2.05 (dd, J = 13.4, 7.1 Hz, 1H), 2.32 (dd, J = 13.4, 9.3 Hz, 1H), 3.15 (s, 4H), 3.25 (d, J = 11.6 Hz, 1H), 3.50 (dt, J = 20.2, 7.0 Hz, 2H), 3.64 (m, 2H), 4.09 (dd, J = 9.2, 7.1 Hz, 1H), 5.58 (s, 1H), 6.68 (q, J = 7.2 Hz, 1H), 7.65 (d, J = 8.0 Hz, 2H), 7.73 (d, J = 8.4 Hz, 2H), 7.88 (d, J = 8.4 Hz, 2H), 8.01 (m, 2H)
55dj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.60 (q, J = 6.4, 5.0 Hz, 4H), 2.03 (dd, J = 13.4, 7.1 Hz, 1H), 2.31 (dd, J = 13.4, 9.2 Hz, 1H), 3.10 (d, J = 11.7 Hz, 1H), 3.23 (d, J = 11.8 Hz, 1H), 3.48 (m, 2H), 3.63 (m, 2H), 4.06 (dd, J = 9.2, 7.1 Hz, 1H), 5.57 (s, 1H), 6.67 (q, J = 7.1 Hz, 1H), 7.68 (m, 4H), 7.79 (m, 2H), 7.96 (m, 2H)
55dk	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (d, J = 3.6 Hz, 1H), 1.63 (q, J = 5.8 Hz,

	5H), 2.07 (dd, J = 13.5, 7.5 Hz, 1H), 2.37 (dd, J = 13.5, 9.0 Hz, 1H), 3.04 (s, 3H), 3.15 (d, J = 24.6 Hz, 6H), 3.27 (m, 1H), 3.52 (dt, J = 24.6, 8.3 Hz, 2H), 3.65 (m, 2H), 4.23 (t, J = 8.1 Hz, 1H), 6.66 (q, J = 7.0 Hz, 1H), 7.51 (d, J = 8.1 Hz, 2H), 7.68 (m, 6H)
55dl	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (m, 3H), 1.60 (q, J = 6.1, 4.7 Hz, 4H), 2.05 (dd, J = 13.2, 7.0 Hz, 1H), 2.32 (dd, J = 13.5, 9.2 Hz, 1H), 3.13 (d, J = 11.6 Hz, 1H), 3.26 (m, 5H), 3.48 (ddd, J = 26.8, 12.6, 5.3 Hz, 2H), 3.64 (td, J = 19.2, 16.1, 9.2 Hz, 2H), 3.78 (s, 1H), 3.90 (m, 3H), 4.10 (dd, J = 9.2, 7.1 Hz, 1H), 4.95 (s, 13H), 5.56 (s, 1H), 6.67 (q, J = 7.0 Hz, 1H), 7.67 (m, 7H)
55dm	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.06 (t, J = 7.4 Hz, 4H), 1.59 (d, J = 6.1 Hz, 5H), 1.82 (dq, J = 14.1, 6.6 Hz, 2H), 2.02 (dd, J = 13.4, 7.2 Hz, 1H), 2.30 (dd, J = 13.5, 9.2 Hz, 1H), 3.07 (d, J = 11.6 Hz, 1H), 3.21 (m, 2H), 3.48 (dt, J = 20.5, 6.6 Hz, 2H), 3.65 (d, J = 15.7 Hz, 2H), 4.03 (td, J = 7.7, 6.5, 5.1 Hz, 3H), 4.91 (m, 1H), 5.55 (s, 1H), 6.63 (q, J = 7.2 Hz, 1H), 7.13 (t, J = 8.7 Hz, 1H), 7.38 (m, 2H), 7.59 (q, J = 8.4 Hz, 4H)
55dn	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (d, J = 2.6 Hz, 1H), 1.43 (t, J = 6.9 Hz, 3H), 1.60 (q, J = 5.8 Hz, 4H), 2.05 (dd, J = 13.1, 7.0 Hz, 1H), 2.33 (dd, J = 13.4, 9.0 Hz, 1H), 3.12 (d, J = 11.4 Hz, 1H), 3.25 (d, J = 11.4 Hz, 1H), 3.48 (m, 2H), 3.65 (q, J = 12.5, 10.8 Hz, 2H), 4.11 (dq, J = 16.7, 8.5, 7.7 Hz, 3H), 5.56 (s, 1H), 6.63 (q, J = 7.0 Hz, 1H), 7.13 (t, J = 8.6 Hz, 1H), 7.38 (m, 2H), 7.59 (q, J = 8.1 Hz, 4H)
55do	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.40 (t, J = 7.0 Hz, 3H), 1.60 (q, J = 5.9, 5.1 Hz, 4H), 2.05 (dd, J = 13.4, 7.2 Hz, 1H), 2.33 (dd, J = 13.4, 9.2 Hz, 1H), 3.12 (d, J = 11.7 Hz, 1H), 3.24 (d, J = 11.7 Hz, 1H), 3.47 (ddt, J = 20.2, 12.7, 5.6 Hz, 2H), 3.64 (m, 2H), 4.07 (p, J = 7.1 Hz, 3H), 5.56 (s, 1H), 6.62 (q, J = 7.1 Hz, 1H), 6.97 (m, 2H), 7.57 (m, 6H)
55dp	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (s, 1H), 1.64 (d, J = 6.8 Hz, 5H), 2.07 (dd, J = 13.5, 7.6 Hz, 1H), 2.38 (dd, J = 13.5, 9.1 Hz, 1H), 3.17 (d, J = 11.8 Hz, 1H), 3.27 (d, J = 11.8 Hz, 1H), 3.52 (m, 1H), 3.66 (m, 1H), 4.24 (t, J = 8.3 Hz, 1H), 6.70 (p, J = 7.2 Hz, 1H), 7.72 (d, J = 8.1 Hz, 3H), 7.90 (m, 3H), 8.28 (m, 3H), 8.53 (m, 1H), 9.31 (d, J = 5.9 Hz, 1H)
55dq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.62 – 7.49 (m, 4H), 7.33 – 7.24 (m, 2H), 6.76 (d, J = 8.3 Hz, 1H), 6.62 (q, J = 7.2 Hz, 1H), 5.53 (s, 1H), 4.19 – 4.12 (m, 2H), 4.08 (dd, J = 9.1, 7.3 Hz, 1H), 3.61 (s, 2H), 3.44 (ddt, J = 20.8, 13.4, 6.0 Hz, 2H), 3.22 (d, J = 11.7 Hz, 1H), 3.10 (d, J = 11.7 Hz, 1H), 2.80 (t, J = 6.5 Hz, 2H), 2.30 (dd, J = 13.5, 9.2 Hz, 1H), 2.00 (ddd, J = 17.2, 12.5, 6.7 Hz, 3H), 1.57 (q, J = 5.9, 4.6 Hz, 4H), 1.27 (s, 1H)

Example 56: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1,2,3,4-tetrahydroquinoxalin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(1,2,3,4-tetrahydroquinoxalin-6-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (a by-product from the N-CBZ deprotection of Example 55bi) using the LiOH general method provided the title compound as an off-white solid.

5

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.54 - 1.72 (m, 4 H) 2.07 - 2.14 (m, 1 H) 2.29 - 2.41 (m, 1 H) 3.09 - 3.18 (m, 1 H) 3.22 - 3.29 (m, 1 H) 3.36 - 3.42 (m, 4 H) 3.43 - 3.58 (m, 2 H) 3.60 - 3.80 (m, 2 H) 4.03 - 4.17 (m, 1 H) 5.49 - 5.65 (m, 1 H) 6.50 - 6.67 (m, 2 H) 6.77 - 6.92 (m, 2 H) 7.43 - 7.63 (m, 4 H). LCMS (MH⁺): 585.

10

Example 57: (S)-8-(2-amino-6-((R)-1-(3,4-dihydroquinazolin-6-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-1-(3,4-dihydroquinazolin-6-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (a by-product from the N-CBZ deprotection of Example 55bk) using the LiOH general method provided the title compound as an off-white solid.


15

¹H NMR (400 MHz, MeOH-d₄) δ ppm 1.54 - 1.66 (m, 4 H) 1.98 - 2.08 (m, 1 H) 2.23 - 2.34 (m, 1 H) 3.02 - 3.11 (m, 1 H) 3.17 - 3.25 (m, 1 H) 3.37 - 3.54 (m, 2 H) 3.55 - 3.72 (m, 2 H) 3.97 - 4.08 (m, 1 H) 4.62 - 4.70 (m, 2 H) 5.50 - 5.58 (m, 1 H) 6.56 - 6.66 (m, 1 H) 6.86 - 6.93 (m, 1 H) 7.19 - 7.24 (m, 1 H) 7.25 - 7.31 (m, 1 H) 7.38 - 7.44 (m, 1 H) 7.51 - 7.57 (m, 2 H) 7.57 - 7.64 (m, 2 H). LCMS (MH⁺): 583

20

Example 58: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(1,2,3,4-tetrahydroquinazolin-6-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid


25



Hydrolysis of (S)-ethyl 8-(2-amino-6-((R)-1-(3,4-dihydroquinazolin-6-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (a by-product from the N-CBZ deprotection of Example 55bk) using the LiOH general method provided the title compound as an off-white solid.

¹H NMR (400 MHz, MeOH-d₄) δ ppm 1.57 - 1.68 (m, 4 H) 1.93 - 2.03 (m, 1 H) 2.18 - 2.30 (m, 1 H) 2.90 - 3.01 (m, 1 H) 3.12 - 3.19 (m, 1 H) 3.43 - 3.75 (m, 4 H) 3.86 - 3.95 (m, 1 H) 4.00 - 4.07 (m, 2 H) 4.15 - 4.23 (m, 2 H) 5.45 - 5.64 (m, 1 H) 6.56 - 6.67 (m, 2 H) 7.17 - 7.23 (m, 1 H) 7.27 - 7.33 (m, 1 H) 7.49 - 7.55 (m, 2 H) 7.55 - 7.62 (m, 2 H). LCMS (MH⁺): 585.

Example 59a: (S)-8-(2-amino-6-((R)-1-(4-bromophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




N-CBZ Deprotection was carried out using Method B with (S)-8-(2-amino-6-((R)-1-(4-bromophenyl)-2,2,2-trifluoroethoxy) pyrimidin-4-yl)-2-(benzyloxycarbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (75 mg, product of Step 3, Example 55an) providing the title compound as a white solid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.50 - 1.68 (m, 4 H) 1.96 (s, 2 H) 2.04 (dd, J=13.35, 7.20 Hz, 1 H) 2.31 (dd, J=13.35, 9.10 Hz, 1 H) 3.07 - 3.26 (m, 2 H) 3.35 - 3.55 (m, 2 H) 3.55 -

3.73 (m, 2 H), 4.06 (dd, J=9.13, 7.17 Hz, 1 H) 5.52 (s, 1 H) 6.57 (q, J=7.11 Hz, 1 H) 7.41 (d, J=8.44 Hz, 2 H) 7.51 - 7.58 (m, 2 H); LCMS (MH⁺): 531.


Example 59b: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(naphthalen-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



The title compound was made as described for Example 10e, starting with (R)-2,2,2-trifluoro-1-(naphthalen-2-yl)ethanol.

¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.20 (dt, J = 12.5, 5.3 Hz, 2H), 1.47 (m, 3H), 1.88 (dd, J = 12.4, 8.0 Hz, 1H), 2.57 (s, 1H), 2.69 (s, 1H), 2.80 (d, J = 12.4 Hz, 1H), 3.36 (m, 3H), 3.97 (dt, J = 12.3, 5.2 Hz, 2H), 6.05 (s, 1H), 6.37 (m, 3H), 7.53 (m, 2H), 7.77 (dd, J = 7.5, 1.5 Hz, 1H), 7.93 (m, 4H); LCMS (MH⁺): 562.

Example 59c: (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(3-fluoroquinolin-6-yl)-2-methylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: To a solution of 4-bromo-2-methylbenzoic acid (5.0 g, 23.2 mmol) in DMF (50 mL) was added potassium carbonate (6.4 g, 46.4 mmol) and iodomethane (6.6 g, 46.479 mmol). The mixture was stirred at RT for 12 h then diluted with water and extracted with ethyl acetate. The combined organic layer was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*.

Purification by normal phase silica gel column provided methyl 4-bromo-2-methylbenzoate as a colorless oil.

5 *Step 2:* To a solution of methyl 4-bromo-2-methylbenzoate (2 g, 8.7 mmol) in THF (20 mL) was added LAH (663 mg, 17.5 mmol) at 0°C. The mixture was stirred at RT for 1 h, then diluted with NaOH (1.0M, 10 mL) and extracted with ethyl acetate. The combined organic layer was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. Purification by normal phase silica gel column provided (4-bromo-2-methylphenyl)methanol as a colorless oil.

10 *Step 3:* To the solution of (4-bromo-2-methylphenyl)methanol (1.8 g, 8.1 mmol) in CH₂Cl₂ (20 mL) was added Dess-Martin Periodinane (5.1 g, 12.1 mmol) at 0°C. The mixture was stirred at RT for 1 h, then diluted with water, and the solid was removed by filtration. The filtrate was extracted with CH₂Cl₂. The combined organic layer was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. Purification by normal phase silica gel column provided 4-bromo-2-
15 methylbenzaldehyde as a yellow oil.

Step 4: To a solution of 4-bromo-2-methylbenzaldehyde (1.5 g, 7.5 mmol) in THF (20 mL) was added TMSCF₃ (2.2 g, 15.5 mmol) at 0°C and then TBAF (1.1 mL, 1.0 M in THF). The mixture was stirred at RT for 1 h, then diluted with HCl (3.0 M, 10 mL), stirred at RT for 1 h and
20 extracted with ethyl acetate. The combined organic layer was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. Purification by normal phase silica gel column provided 1-(4-bromo-2-methylphenyl)-2,2,2-trifluoroethanol as an off-white solid.

25 *Step 5:* To a solution of 1-(4-bromo-2-methylphenyl)-2,2,2-trifluoroethanol (1.8 g, 6.7 mmol) in CH₂Cl₂ (20 mL) was added Dess-Martin Periodinane (3.4 g, 8.1 mmol) at 0°C. The mixture was stirred at RT for 2 h, then diluted with water (10 mL) and filtered. The filtrate was extracted with CH₂Cl₂. The combined organic layer was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. Purification by normal phase silica gel column provided 1-(4-bromo-2-methylphenyl)-2,2,2-trifluoroethanone as a yellow oil.

30 *Step 6:* Chiral reduction of 1-(4-bromo-2-methylphenyl)-2,2,2-trifluoroethanone using the


Iridium complex-catalyzed hydrogenation as described for Intermediate 1, (R)-1-(4-bromo-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethanol provides (R)-1-(4-bromo-2-methylphenyl)-2,2,2-trifluoroethanol.

5 **Steps 7:** The title compound was prepared as described for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 55an) Steps 4-5. 3-Fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinolone was used as the Suzuki coupling partner (CAS# 1251731-31-3).

10 ¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.28 (s, 1H), 1.59 (s, 4H), 2.04 (dd, J = 13.5, 7.0 Hz, 1H), 2.31 (dd, J = 13.3, 9.3 Hz, 1H), 2.65 (s, 3H), 3.10 (d, J = 11.7 Hz, 1H), 3.23 (d, J = 11.5 Hz, 1H), 3.47 (t, J = 14.3 Hz, 2H), 3.63 (t, J = 13.8 Hz, 2H), 4.07 (t, J = 8.1 Hz, 1H), 5.56 (s, 1H), 6.87 (q, J = 7.0 Hz, 1H), 7.63 (d, J = 4.6 Hz, 3H), 8.01 (d, J = 8.9 Hz, 1H), 8.12 (m, 3H), 8.80 (m, 1H). LCMS (MH⁺): 611.

15

Example 59d: (S)-8-(2-amino-6-((R)-1-(2-ethyl-4-(3-fluoroquinolin-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid




20 **Step 1:** To a 0 °C solution of LDA (10.7 mL, 21.39 mmol) in THF (20 mL) was added 4-bromo-2-methylbenzoic acid (2 g, 9.3 mmol) in THF (5 mL). The mixture was stirred at 0 °C for 1 h, cooled to -70 °C, and then MeI (2.3 mL, 37.20 mmol) was added dropwise. The mixture was allowed to warm up to 0 °C, stirred for 3 h, then quenched with H₂O, and the pH was adjusted to 1-2 with 3 N HCl. The mixture was then diluted with water and extracted with ethyl acetate. The combined organic layer was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*.
25 Purification by normal phase silica gel column provided 4-bromo-2-ethylbenzoic acid as a white solid.

Step 2: The title compound was prepared as described above for (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(3-fluoroquinolin-6-yl)-2-methylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 59c) starting with 4-bromo-2-ethylbenzoic acid in place of 4-bromo-2-methylbenzoic acid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.29 (m, 1H), 1.41 (t, J = 7.5 Hz, 3H), 1.59 (q, J = 6.1, 5.6 Hz, 4H), 2.04 (dd, J = 13.5, 7.1 Hz, 1H), 2.32 (dd, J = 13.4, 9.1 Hz, 1H), 3.01 (dt, J = 12.1, 7.0 Hz, 2H), 3.12 (d, J = 11.6 Hz, 1H), 3.24 (d, J = 11.8 Hz, 1H), 3.48 (dt, J = 21.5, 6.9 Hz, 2H), 3.62 (m, 2H), 4.08 (dd, J = 9.1, 7.0 Hz, 1H), 4.94 (s, 15H), 5.56 (s, 1H), 7.00 (q, J = 6.9 Hz, 1H), 7.67 (m, 3H), 8.11 (m, 5H), 8.80 (d, J = 2.8 Hz, 1H). LCMS (MH⁺): 626.

Example 60: 9-(2-Amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-3,9-diazaspiro[5.5]undecane-2-carboxylic acid




Step 1: To a solution of methyl 3,9-diazaspiro[5.5]undecane-10-carboxylate (30 mg, 0.14 mmol) in dioxane (2 mL) / i-PrOH (2 mL) was added 4-chloro-6-[(1R)-1-[4-chloro-2-(3-methylpyrazol-1-yl)phenyl]-2,2,2-trifluoro-ethoxy]pyrimidin-2-amine (92 mg, 0.22 mmol), and the reaction was heated at 100 °C under microwave for 3 h. The reaction was cooled to RT, and concentrated *in vacuo*. The residue was purified by reversed phase HPLC (MeOH/H₂O/0.5% TFA) to provide methyl 9-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-3,9-diazaspiro[5.5]undecane-2-carboxylate as an off-white solid.

Step 2: Hydrolysis of methyl 9-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-3,9-diazaspiro[5.5]undecane-2-carboxylate using the LiOH general method provides the title compound as an off-white solid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.22 - 1.37 (m, 1 H) 1.30 - 1.30 (m, 1 H) 1.46 - 1.68 (m, 4 H) 1.68 - 1.90 (m, 2 H) 2.29 (dd, J=12.59, 6.83 Hz, 1 H) 2.37 (d, J=1.90 Hz, 3 H) 3.07 - 3.24 (m, 2 H) 3.59 - 3.90 (m, 4 H) 4.03 - 4.19 (m, 1 H) 6.29 - 6.38 (m, 1 H) 6.40 (d, J=2.29 Hz, 1 H),
5 6.88 - 7.02 (m, 1 H) 7.52 - 7.61 (m, 2 H) 7.65 - 7.74 (m, 1 H) 7.89 (d, J=2.34 Hz, 1 H). LCMS (MH⁺): 581.

Example 61: (S)-8-(2-Amino-6-((4-(3-methyl-1H-indazol-6-yl)phenoxy)methyl)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: A mixture of 4-bromophenol (173 mg, 1.00 mmol), 4-chloro-6-(chloromethyl)pyrimidin-2-amine (CAS#: 92311-35-8) (178 mg, 1.16 mmol) and K₂CO₃ (175 mg, 1.00 mmol) in DMF (5 mL) was heated to 100°C for 12 h. The reaction was cooled to RT, concentrated *in vacuo*, and the residue taken up in and EtOAc. The organic layer was washed with brine, dried over
15 Na₂SO₄ and concentrated *in vacuo*. Purification on normal phase silica gel (EtOAc/petroleum ether) provided 4-((4-bromophenoxy)methyl)-6-chloropyrimidin-2-amine as a white solid.

Step 2: A mixture of 4-((4-Bromophenoxy)methyl)-6-chloropyrimidin-2-amine (454 mg, 1.4 mmol), (S)-2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (500 mg, 1.44 mmol) and NaHCO₃(605 mg,7 mmol) in dioxane (5 mL) was heated to 100°C for 12 h. The reaction was cooled to RT, concentrated *in vacuo*, and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated *in vacuo*. Purification on normal phase silica gel (EtOAc/petroleum ether) provided (S)-2-benzyl 3-ethyl 8-(2-amino-6-((4-bromophenoxy)methyl)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a
25 white solid.

Step 3: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((4-bromophenoxy)methyl)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (550 mg, 0.9 mmol) in acetonitrile (5 mL) was added TMSI (705 mg, 3.5 mmol) dropwise at 0 °C. The mixture was stirred at 0 °C for 2 h, then concentrated *in vacuo*. The residue was dissolved in CH₂Cl₂ (20 mL) followed by the sequential addition of Et₃N (267 mg, 2.6 mmol), and (BOC)₂O (285 mg, 1.3 mmol). The reaction mixture was stirred at RT for 16 h then concentrated *in vacuo*. Purification on normal phase silica gel (CH₂Cl₂/MeOH) provides (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((4-bromophenoxy)methyl)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a light yellow solid.

Step 4: A mixture of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((4-bromophenoxy)methyl)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (350 mg, 0.56 mmol), 3-methyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole (285 mg, 1.1 mmol) and Pd(dppf)Cl₂ (62 mg, 0.09 mmol) in dioxane (5 mL)/aq. Na₂CO₃ solution (2.0 M, 5 mL) was heated to 90 °C for 4 h. The reaction was cooled to RT, the solids filtered away, and the solution concentrated *in vacuo*. Purification on normal phase silica gel (CH₂Cl₂/MeOH) provided (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((4-(3-methyl-1H-indazol-6-yl)phenoxy)methyl)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a brown solid.


Step 5: To a solution of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((4-(3-methyl-1H-indazol-6-yl)phenoxy)methyl)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (150 mg, 0.19 mmol) in CH₂Cl₂ (5 mL) was added TFA (3 mL), and the resulting mixture was stirred at RT for 1 h. The reaction mixture was concentrated *in vacuo*, and the resulting material partitioned between CH₂Cl₂ and saturated NaHCO₃, and extracted. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by prep-TLC (CH₂Cl₂/MeOH) provided (S)-ethyl 8-(2-amino-6-((4-(3-methyl-1H-indazol-6-yl)phenoxy)methyl)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as a brown solid.

Step 6: To a solution of (S)-ethyl 8-(2-amino-6-((4-(3-methyl-1H-indazol-6-yl)phenoxy)methyl)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (70 mg, 0.11 mmol) in MeOH (3 mL) is added 4 N NaOH (3 mL), and the reaction mixture was stirred at RT for 4 h. The reaction mixture was then concentrated *in vacuo*. The residue was diluted with water (5 mL) and the pH

adjusted to 6-7. The precipitated solid was collected by filtration, and the filter cake was washed with cold water, then dried to afford the title compound as an off-white solid.

¹H NMR (400 MHz, DMSO-d₆): δ ppm 7.72-7.70 (d, 1 H), 7.61-7.59 (d, 3 H), 7.31-7.30 (d, 1 H), 7.06-7.04 (d, 2 H), 6.14 (s, 1 H), 4.76 (s, 2 H), 3.87-3.83 (q, 1 H), 3.46-3.41 (m, 4 H), 3.08-3.06 (d, 1 H), 2.98-2.95 (d, 1 H), 2.43 (s, 1 H), 2.16-2.13 (m, 1 H), 1.82-1.80 (m, 1 H), 1.44 (m, 4 H). LCMS (MH⁺): 514.

Example 62: (S)-8-(2-amino-6-((5-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)methoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid



Step 1: A mixture of (2-bromo-4-chlorophenyl)methanol (173 mg, 1. mmol), 4-chloro-6-(chloromethyl)pyrimidin-2-amine (178 mg, 1.16 mmol) and K₂CO₃ (175 mg, 1.00 mmol) in DMF (5 mL) was heated to 100 °C for 12 h. The reaction was cooled to RT, concentrated *in vacuo*, and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated *in vacuo*. Purification on normal phase silica gel (EtOAc/petroleum ether) provided 4-((2-bromo-4-chlorobenzyl)oxy)-6-chloropyrimidin-2-amine as a white solid.

Step 2: A mixture of 4-((2-bromo-4-chlorobenzyl)oxy)-6-chloropyrimidin-2-amine (300 mg, 1.1 mmol), (S)-2-benzyl 3-ethyl 2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (400 mg, 1.2 mmol), and NaHCO₃(550 mg, 7 mmol) in dioxane (5 mL) was heated to 100 °C for 12 h. The reaction was cooled to RT, concentrated *in vacuo*, and extracted with EtOAc. The combined organic layers were washed with brine, water, dried over Na₂SO₄, and concentrated *in vacuo*. Purification on normal phase silica gel (EtOAc/petroleum ether) provided (S)-2-benzyl 3-ethyl 8-

(2-amino-6-((2-bromo-4-chlorobenzyl)oxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a white solid.

Step 3: To a solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((2-bromo-4-chlorobenzyl)oxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (500 mg, 0.8 mmol) in acetonitrile (5 mL) was added TMSI (705 mg, 3.5 mmol) dropwise at 0 °C. The reaction mixture was stirred at 0 °C for 2 h, then concentrated *in vacuo*. The residue was dissolved in CH₂Cl₂ (20 mL), followed by the sequential addition of Et₃N (267 mg, 2.6 mmol), and (BOC)₂O (285 mg, 1.3 mmol). The reaction mixture was stirred at RT for 16 h, then concentrated *in vacuo*. Purification on normal phase silica gel (CH₂Cl₂/MeOH) provided (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((2-bromo-4-chlorobenzyl)oxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as a light yellow solid.

Step 4: A mixture of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((2-bromo-4-chlorobenzyl)oxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (300 mg, 0.4 mmol), (3-(methylsulfonyl)phenyl)boronic acid (280 mg, 1 mmol), and Pd(dppf)Cl₂ (62 mg, 0.09 mmol) in dioxane (5 mL)/aq. Na₂CO₃ solution (2.0 M, 5 mL) was heated to 90 °C for 4 h. The reaction was then cooled to RT, the solids filtered away, and the filtrate concentrated *in vacuo*. Purification on normal phase silica gel (CH₂Cl₂/MeOH) provided (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((5-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)methoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as an off-white solid.


Step 5: To a solution of (S)-2-tert-butyl 3-ethyl 8-(2-amino-6-((5-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)methoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (200 mg, 0.25 mmol) in CH₂Cl₂ (5 mL) was added TFA (3 mL), and the resulting mixture was stirred at RT for 1 h. The reaction mixture was concentrated *in vacuo*, and the residue was partitioned between CH₂Cl₂ and saturated NaHCO₃. The organic layer was dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by prep-TLC (CH₂Cl₂/MeOH) provided (S)-ethyl 8-(2-amino-6-((5-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)methoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate as an off-white solid.

Step 6: To a solution (S)-ethyl 8-(2-amino-6-((5-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)methoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (100 mg, 0.13 mmol) in MeOH (3 mL) was added 4 N NaOH (3 mL), and the mixture was stirred at RT for 4 h. The reaction mixture was then concentrated *in vacuo*. The residue was diluted with water (5 mL) and the pH adjusted to 6-7. The precipitated solid was collected by filtration, the filter cake was washed with cold water, then dried to afford the title compound as an off-white solid isolated as the zwitterionic form.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 7.94 (m, 2 H), 7.59-7.57 (m, 3 H), 7.44-7.40 (m, 1 H), 7.33 (m, 1 H), 5.33 (m, 1 H), 4.07 (m, 1 H), 3.59 (m, 2 H), 3.45 (m, 2 H), 3.30 (m, 1 H), 3.15 (m, 1 H), 2.32 (m, 1 H), 2.06 (m, 1 H), 1.61 (s, 4 H). LCMS (MH⁺): 573.


The following esters were isolated as either a TFA or HCl salt formed during the HPLC purification procedure used to isolate the final compounds.

Example 63bd: (S)-ethyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate




A solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (from Step 3, Example 1m, 220 mg, 0.3 mmol,) in EtOAc (5 mL) was hydrogenated using Method A by using an H-Cube apparatus and a 10% (w/w) Pd/C cartridge with a flow rate of 1.0 mL/min at RT. Purification on normal phase silica gel (EtOAc/heptane) provided (S)-ethyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate.

Example 63kp: (S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(hydroxymethyl)-4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate



The title compound was prepared as described for (S)-ethyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (Example 63bd) using Method A to remove the N-CBz group.

Example 63i: (S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate



A solution of (S)-2-benzyl 3-ethyl 8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (from Step 3, Example 34c, 315 mg, 0.43 mmol) in acetonitrile (300 mL) was added TMSI (0.13 mL, 0.9 mmol) [Method B]. The reaction mixture was then warmed to RT for an additional 30-40 min, then cooled to 0-5 °C, and 2 M HCl in diethyl ether (0.5 mL) was added. The reaction mixture was the allowed to warm RT and then concentrated *in vacuo*. Normal phase silica gel chromatography provide the title compound as an off-white solid.

Ethyl ester prodrugs in Table 18a were prepared by removing the N-CBZ group by either method A or method B, as shown below.












Table 18a.









5

Ex. No.	Ar	Method A or B	CAS Name	LCMS (MH ⁺)
63a		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(benzo[d]thiazol-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	613
63b		A	(S)-ethyl 8-(6-((R)-1-(4-(1H-indazol-5-yl)phenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	596
63c		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-4'-(pyrrolidine-1-carbonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	683







63d		A	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-4'-nitro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	631
63e		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(benzo[d]isothiazol-5-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	613
63f		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(benzo[d]isothiazol-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	616
63g		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3',4'-dimethoxy-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	616
63h		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1-methyl-2-oxo-1,2-dihydroquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	637
63i		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	591
63j		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-amino-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	606

63k		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-propyl-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	676
63l		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(1,3-dimethyl-1H-indol-5-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	622
63m		A	(S)-ethyl 8-(6-((R)-1-(3'-acrylamido-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	659.1
63n		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	604
63o		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	587
63p		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	626
63q		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-oxo-1,2-dihydroquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	623








63r		B	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methanesulfonyl)-5-((E)-prop-1-en-1-yl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	674
63s		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	583
63t		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-chloro-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	671
63u		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methanesulfonyl)-4-propyl-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	677
63v		B	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methanesulfonyl)-4-((E)-prop-1-en-1-yl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	675
63w		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(ethanesulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	683

63x		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(propylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	697
63y		A	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(butylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	711
63z		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1-oxo-1,3-dihydroisobenzofuran-5-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	612
63aa		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-methoxyquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	638
63ab		A	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(hydroxymethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	621
63ac		A	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(2-oxopyrrolidin-1-yl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	673
63ad		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(3-methyl-2-oxoimidazolidin-1-yl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	688







63ae		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	668
63af		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(methylsulfonamido)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	683
63ag		B	(S)-ethyl 8-(2-amino-6-((R)-1-(2-bromo-5-chlorophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	593
63ah		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (	595
63ai		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-(methylthio)quinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	554
63aj		B	(S)-ethyl 8-(2-amino-6-((R)-1-(2,5-dibromophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	638
63ak		A	(S)-ethyl 8-(6-((R)-1-([1,1':4',1''-terphenyl]-2'-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	633
63al		A	(S)-ethyl 8-(2-amino-6-((R)-1-(2'-(ethoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-	708








			trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	
63am		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-(ethoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	708
63an		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-(ethoxycarbonyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	708
63ao		B	(S)-ethyl 8-(2-amino-6-((R)-1-(2,6-dibromophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	638
63ap		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3',5'-dichloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	625
63aq		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	605
63ar		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	659








63as		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-(methylthio)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	583
63at		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	591
63au		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	651
63av		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	651
63aw		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3',4'-dichloro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	705
63ax		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(2-oxopyrrolidin-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	598
63ay		B	ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	594

63az		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	594
63ba		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	587
63bb		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-methoxy-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	591
63bc		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	575
63bd		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	665
63be		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-ethyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	589
63bf		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-propylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	603

63bg		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-butyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	617
63bh		A	(S)-ethyl 8-(2-amino-6-((R)-1-(5-(ethoxycarbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	633
63bi		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(ethoxycarbonyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	632
63bj		A	(S)-ethyl 8-(2-amino-6-((R)-1-(5-((E)-3-ethoxy-3-oxoprop-1-en-1-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	559
63bk		A	(S)-ethyl 8-(2-amino-6-((R)-1-(5-(3-ethoxy-3-oxopropyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	588
63bl		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(6-methyl-2-(3-methyl-1H-pyrazol-1-yl)pyridin-3-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	618

63bm		B	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-5-((E)-prop-1-en-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	600
63bn		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	664
63bo		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-5-propylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	602
63bp		A	(S)-ethyl 8-(2-amino-6-((R)-1-(5-ethyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	588
63bq		A	(S)-ethyl 8-(2-amino-6-((R)-1-(5-butyl-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	616
63br		B	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-5-vinylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	586

63bs		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-((E)-but-1-en-1-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	614
63bt		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(1-methyl-1H-pyrazol-3-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	594
63bu		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(1-methyl-1H-pyrazol-3-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	560
63bv		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(1,3-dimethyl-1H-indazol-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	624
63bw		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(2,3-dimethyl-2H-indazol-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	624
63bx		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(1-oxo-1,2,3,4-tetrahydroisoquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	625
63by		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(isoquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	607

63bz		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(3-ethoxy-3-oxopropyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	660
63ca		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(isoquinolin-7-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	607
63cb		A	(S)-ethyl 8-(2-amino-6-((R)-1-(5-(4-ethoxy-4-oxobutyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	674
63cc		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(4-ethoxy-4-oxobutyl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	594
63cd		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(4-ethoxy-4-oxobutyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	674
63ce		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-cyano-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	661
63cf		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-cyano-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	615

63cg		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-methoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	620
63ch		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-sulfamoyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	669
63ci		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-hydroxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	606
63cj		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	668
63ck		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-(aminomethyl)-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	619
63cl		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(quinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	608
63cm		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(quinolin-7-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	608

63cn		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	694
63co		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(quinoxalin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	608
63cp		A	(S)-ethyl 8-(6-((R)-1-(4'-(acetamidomethyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	707
63cq		A	(S)-ethyl 8-(6-((R)-1-(4'-(2-acetamidoethyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	721
63cr		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(quinolin-7-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	687
63cs		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-methoxypyridin-4-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	666
63ct		A	(S)-ethyl 8-(6-((R)-1-(4-(1H-indol-6-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-	595








			carboxylate	
63cu		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(ethoxycarbonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	662
63cv		B	2'-((R)-1-((2-amino-6-((S)-3-(ethoxycarbonyl)-2,8-diazaspiro[4.5]decan-8-yl)pyrimidin-4-yl)oxy)-2,2,2-trifluoroethyl)-5'-chloro-[1,1'-biphenyl]-3-carboxylic acid	634
63cw		B	(S)-ethyl 8-(6-((R)-1-(3'-(acrylamidomethyl)-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	673
63cx		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-carbamoyl-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	633
63cy		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-4'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	668
63cz		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-4'-sulfamoyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	669
63da		A	(S)-ethyl 8-(2-amino-6-((R)-1-(2'-(ethoxycarbonyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	708

			2,8-diazaspiro[4.5]decane-3-carboxylate	
63db		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-(ethoxycarbonyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	708
63dc		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-(ethoxycarbonyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	708
63dd		A	(3S)-ethyl 8-(2-amino-6-((1R)-1-(4-(1,2-dihydroxyethyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	620
63de		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-(aminomethyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	666
63df		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-((E)-3-ethoxy-3-oxoprop-1-en-1-yl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	734
63dg		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-((E)-3-ethoxy-3-oxoprop-1-en-1-yl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	734








63dh		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-(3-ethoxy-3-oxopropyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	736
63di		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-(3-ethoxy-3-oxopropyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	736
63dj		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	654
63dk		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(quinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	687
63dl		A	(3S)-ethyl 8-(2-amino-6-((1R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(2-oxo-1,3-dioxolan-4-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	646
63dm		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(2-methyl-1-oxo-1,2,3,4-tetrahydroisoquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	639








63dn		A	(S)-ethyl 8-(6-((R)-1-(4-(acetamidomethyl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	631
63do		A	(3S)-ethyl 8-(2-amino-6-((1R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-((2-oxotetrahydrofuran-3-yl)thio)ethyl)carbamoyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	823
63dp		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3,4-dimethyl-3''-(methylsulfonyl)-[1,1':3',1''-terphenyl]-4'-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	738
63dq		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(methylsulfonyl)-5-(quinolin-6-yl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	761
63dr		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(hydroxymethyl)-3'-methyl-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	680
63ds		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(hydroxymethyl)-4'-methyl-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	680

63dt		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(methoxycarbonyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	694
63du		A	3'-((S)-1-((2-amino-6-((R)-3-(ethoxycarbonyl)-2,8-diazaspiro[4.5]decane-8-yl)pyrimidin-4-yl)oxy)-2,2,2-trifluoroethyl)-4'-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-carboxylic acid	680
63dv		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(1-oxo-1,3-dihydroisobenzofuran-5-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	693
63dw		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(quinazolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	608
63dx		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(pyrimidin-5-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	638
63dy		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3',4'-difluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	672








63dz		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-chloro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	671
63ea		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	591
63eb		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-chloro-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	671
63ec		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3',4'-difluoro-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	672
63ed		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	636
63ee		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3',4'-dichloro-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	705
63ef		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	574

63eg		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl))-5-(pyrimidin-5-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	638
63eh		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4',5-dichloro-3'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	643
63ei		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-ethoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	635
63ej		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3',5-dichloro-4'-ethoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	669
63ek		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3',5-dichloro-5'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	643
63el		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-(tert-butyl)-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	647
63em		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3',5-dichloro-5'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	693







63en		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-fluoro-5'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	677
63eo		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	591
63ep		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-methoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	621
63eq		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-isopropoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	649
63er		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3',5-dichloro-4'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	639
63es		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3',5-dichloro-4'-isopropoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	683
63et		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-fluoro-4'-isopropoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	667







63eu		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4',5-dichloro-3'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	693
63ev		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	609
63ew		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4',5-dichloro-3'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	639
63ex		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3',5-dichloro-4'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	693
63ey		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3',5'-difluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	
63ez		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3',5-dichloro-4'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	
63fa		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3',4'-difluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	

63fb		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3',4'-dimethyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	619
63fc		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4',5-dichloro-3',5'-dimethyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	653
63fd		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-4'-ethoxy-3'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	653
63fe		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3',5'-dimethyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	619
63ff		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3',5-dichloro-5'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	639
63fg			(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-4'-fluoro-3'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	623
63fh		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-methyl-4'-(trifluoromethoxy)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	689







63fi		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(trifluoromethoxy)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	675
63fj		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-isopropyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	633
63fk		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3',5'-bis(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	727
63fl		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-fluoro-4'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	623
63fm		B	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3',5,5'-trichloro-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	659
63fn		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-4'-fluoro-3'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	677
63fo		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(pyridin-3-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	592

63fp		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-fluoro-5'-isopropoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	667
63fq		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-ethoxy-5'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	653
63fr		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-(tert-butyl)-5-chloro-5'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	661
63fs		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-cyano-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	616
63ft		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-ethoxy-5'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	698
63fv		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-chloro-3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	689

63fw		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-chloro-4'-ethoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	715
63fx		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-ethoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	680
63fy		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3',5'-difluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	672
63fz		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	704
63ga		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-ethoxy-4'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	653
63gb		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	712







63gc		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3',5'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	664
63gd		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-chloro-5'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	685
63ge		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	678
63gf		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-(tert-butyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	692
63gg		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-chloro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	671
63gh		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-chloro-4'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	689








63gi		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3',4'-difluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	672
63gj		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	722
63gk		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-chloro-3-(3-methyl-1H-pyrazol-1-yl)-5'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	739
63gl		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-3'-(trifluoromethoxy)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	720
63gm		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-(tert-butyl)-5'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	706
63gn		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-chloro-3'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	685

63go		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-chloro-3',5'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	699
63gp		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-fluoro-3'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	668
63gq		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-ethoxy-3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	698
63gr		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3',5'-dichloro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	705
63gs		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-isopropyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	678
63gt		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-chloro-3-(3-methyl-1H-pyrazol-1-yl)-3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	739







63gu		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-chloro-3-(3-methyl-1H-pyrazol-1-yl)-4'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	738
63gv		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-carbamoyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	679
63gw		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-3',5'-bis(trifluoromethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	772
63gx		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	694
63gy		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-ethoxy-4'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	698
63gz		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-5'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	712







63ha		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-methoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	666
63hb		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-ethoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	680
63hc		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3',4',5'-trifluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	690
63hd		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(1-methyl-2-oxo-1,2-dihydropyridin-4-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	622
63he		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-4'-(trifluoromethoxy)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	734
63hf		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-chloro-4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	685








63hg		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-5'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	722
63hh		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-chloro-5'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	689
63hi		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-cyclopropyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	631
63hj		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-chloro-4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	729
63hk		B	(S)-ethyl 8-(2-amino-6-((R)-1-(2-(benzo[d]thiazol-5-yl)-4-chlorophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	648
63hl		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(2-(dimethylamino)pyridin-4-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	635

63hm		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(naphthalen-2-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	641
63hn		B	(S)-ethyl 8-(2-amino-6-((R)-1-(3'-(tert-butyl)-5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	647
63ho		B	(S)-ethyl 8-(6-((R)-1-(2-(1H-benzo[d]imidazol-1-yl)-4-chlorophenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	631
63hp		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(1H-indazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	631
63hq		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(2-isopropylpyridin-4-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	634
63hr		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-4'-fluoro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	609
63hs		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4',5-dichloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	625







63ht		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-4'-methyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	605
63hu		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-(naphthalen-2-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	687
63hv		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-2',3',4',5'-tetrahydro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	595
63hw		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-(benzyloxy)-3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	761
63hx		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	709
63hy		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-isobutoxy-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	663
63hz		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-isopropoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	690

63ia		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(3-fluoroquinolin-6-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	626
63ib		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-4'-propoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	713
63ic		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-butoxy-3'-fluoro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	727
63id		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-(5-methyl-1,3,4-oxadiazol-2-yl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	734
63ie		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(pyrrolidine-1-carbonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	688
63if		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(cyclopentyloxy)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	675

63ig		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-((1R,4R)-4-hydroxycyclohexyl)carbamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	704
63ih		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-((1R,4R)-4-hydroxycyclohexyl)carbamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	732
63ii		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-ethyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	619
63ij		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-isopropyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	633
63ik		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-propoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	614
63il		A	(S)-ethyl 8-(2-amino-6-((R)-1-(2-ethyl-4-(3-fluoroquinolin-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	654







63im		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(4-methylpiperazine-1-carbonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	717
63in		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(3-fluoroquinolin-6-yl)-2-methylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	640
63io		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-(diethylcarbamoyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	656
63ip		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-carbamoyl-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	600
63iq		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(2-methylthiazol-5-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	612
63ir		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-propoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	691
63is		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(5-chlorothiophen-2-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	631








63it		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(methylsulfonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	635
63iu		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-(methylsulfonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	715
63iv		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-propoxy-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	695
63ix		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-(diethylcarbamoyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	736
63iy		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(methylcarbamoyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	613
63iz		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-sulfamoyl-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	715
63ja		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-sulfamoyl-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	635

63jb		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-((2-morpholinoethyl)carbamoyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	712
63jc		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-((2-morpholinoethyl)carbamoyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	792
63jd		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-(dimethylcarbamoyl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	628
63je		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(piperazine-1-carbonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	669
63jf		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-(piperazine-1-carbonyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	749
63jg		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(1-methyl-2-oxo-1,2-dihydropyridin-3-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	622







63jh		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-(dimethylcarbamoyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	708
63ji		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-methoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	685
63jj		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-fluoro-4'-propoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	633
63jk		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-(methylcarbamoyl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	694
63jl		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(N-methylsulfamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	684
63jm		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(N,N-dimethylsulfamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	698

63jn		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-morpholino-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	700
63jo		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(methylcarbamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	648
63jp		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(dimethylcarbamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	662
63jq		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-ethoxy-3'-fluoro-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	618
63jr		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4'-ethoxy-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	601
63js		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(5-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	635
63jt		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(diethylcarbamoyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	690

63ju		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isobutoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	709
63jv		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-(3-methyl-1H-pyrazol-1-yl)-4'-(neopentyloxy)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	723
63jw		B	(S)-ethyl 8-(6-((R)-1-(2-(1H-benzo[d]imidazol-4-yl)-4-chlorophenyl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	631
63jx		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(chroman-6-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	693
63jy		B	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(piperazine-1-carbonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	703
63jz		A	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(4-cyclopropylpiperazine-1-carbonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	743

63ka		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(cinnolin-6-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	689
63kb		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-(trifluoromethyl)-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	649
63kc		B	(S)-ethyl 8-(2-amino-6-((R)-1-(2-(3-(tert-butyl)-1H-pyrazol-1-yl)-4-chlorophenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	637
63kd		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-isopropyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	624
63ke		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-cyclopropyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	622
63kf		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3-(3-(trifluoromethyl)-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	719
63kg		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3-fluoro-4-propoxy-[1,1':3',1''-terphenyl]-4'-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	660

63kh		A	(S)-ethyl 8-(2-amino-6-((R)-1-(3,4-dimethyl-[1,1':3',1''-terphenyl]-4'-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	708
63ki		A	(S)-ethyl 8-(6-((R)-1-([1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	556
63kj		A	(S)-ethyl 8-(6-((R)-1-([1,1':3',1''-terphenyl]-4'-yl)-2,2,2-trifluoroethoxy)-2-aminopyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	633
63kl		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(hydroxymethyl)-4-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-3-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	667
63km		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(chroman-6-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	612
63kn		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(pyridin-2-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	592
63ko		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(pyrimidin-2-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	593

63kp		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(3'-(hydroxymethyl)-4'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	681
63kq		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-(hydroxymethyl)-3'-methyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	681
63kr		A	(S)-ethyl 8-(2-amino-6-((R)-1-(4-(6-ethoxypyridin-3-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	681
63ks		A	(S)-ethyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4-(6-methoxypyridin-3-yl)-2-(3-methyl-1H-pyrazol-1-yl)phenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	668
63kt		A	(S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(2-methoxyethoxy)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	665
63ku		B	(S)-ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	594


63kv		B	(S)-ethyl 8-(2-amino-6-((S)-1-(3',4'-bis(hydroxymethyl)-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	697
------	---	---	--	-----

Table 18b.

NMR Data for Compounds of Table 18a

Ex. No.	NMR
63a	¹ H NMR (400 MHz, MeOH-d ₄): δ PPM 1.26 (t, J = 7.1 Hz, 3H), 1.50 (m, 5H), 1.63 (s, 1H), 1.73 (dd, J = 13.0, 7.2 Hz, 1H), 2.07 (dd, J = 13.0, 8.7 Hz, 1H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.50 (pd, J = 13.6, 5.4 Hz, 5H), 3.81 (t, J = 8.0 Hz, 1H), 4.17 (qd, J = 7.0, 1.6 Hz, 2H), 4.92 (s, 6H), 5.55 (s, 1H), 6.66 (q, J = 7.2 Hz, 1H), 7.63 (d, J = 8.0 Hz, 2H), 7.77 (m, 4H), 8.10 (d, J = 8.6 Hz, 1H), 8.29 (d, J = 1.8 Hz, 1H), 9.24 (s, 1H)
63b	¹ H NMR (400 MHz, CDCl ₃): δ ppm 1.28 (m, 5H), 1.59 (t, J = 5.6 Hz, 2H), 1.77 (dd, J = 13.1, 6.8 Hz, 1H), 2.09 (m, 1H), 2.87 (d, J = 10.6 Hz, 1H), 2.98 (d, J = 10.6 Hz, 1H), 3.51 (dt, J = 14.9, 5.0 Hz, 4H), 3.92 (m, 1H), 4.21 (q, J = 7.1 Hz, 2H), 4.62 (s, 2H), 5.54 (s, 1H), 6.63 (q, J = 7.0 Hz, 1H), 7.40 (dd, J = 8.5, 1.4 Hz, 1H), 7.62 (q, J = 8.3 Hz, 6H), 7.81 (d, J = 8.5 Hz, 1H), 8.11 (s, 1H)
63c	¹ H NMR (400 MHz, CDCl ₃): δ ppm 0.87 (dd, J = 7.5, 3.2 Hz, 1H), 1.28 (dd, J = 14.0, 6.9 Hz, 7H), 1.57 (dt, J = 17.4, 5.6 Hz, 4H), 1.87 (m, 5H), 2.11 (dd, J = 13.1, 8.8 Hz, 1H), 2.89 (d, J = 10.6 Hz, 1H), 2.99 (d, J = 10.6 Hz, 1H), 3.28 (t, J = 6.7 Hz, 2H), 3.51 (m, 4H), 3.67 (t, J = 6.9 Hz, 2H), 3.89 (s, 4H), 4.21 (q, J = 7.1 Hz, 2H), 4.59 (s, 2H), 5.53 (s, 1H), 6.61 (q, J = 7.1 Hz, 1H), 7.07 (d, J = 1.5 Hz, 1H), 7.17 (dd, J = 7.7, 1.5 Hz, 1H), 7.33 (d, J = 7.8 Hz, 1H), 7.58 (s, 4H)
63d	¹ H-NMR (400 MHz, MeOH-d ₄): δ ppm 1.62(m,4H), 2.09-2.04(m,1H), 2.40-2.35(m,1H), 3.14(m,1H), 3.25(m,1H), 3.47(m,2H), 3.31-3.30(m,2H), 4.22-4.20(m,1H), 5.49(s,1H), 5.83-5.80(m,1H), 6.52-6.38(m,2H), 6.65(m,1H), 7.31(d,J=2.0,1H), 7.45-7.43 (d, J=8.0, 3H), 7.68-7.66(d, J=8.0, 1H), 7.80-7.78(d, J=8.0, 2H)
63e	¹ H NMR (400 MHz, CDCl ₃): δ ppm 0.07 (s, 1H), 0.87 (dd, J = 17.8, 8.8 Hz, 2H), 1.12 (s, 3H), 1.29 (m, 18H), 1.51 (s, 1H), 1.63 (dq, J = 29.6, 7.7, 6.6 Hz, 10H), 1.89 (dd, J = 13.2, 7.4 Hz, 2H), 2.23 (dd, J = 13.2, 8.6 Hz, 2H), 3.13 (m, 4H), 3.53 (h, J = 6.6 Hz, 16H), 4.04 (s, 1H), 4.20 (dq, J = 33.0, 7.5 Hz, 8H), 4.35 (s, 1H), 4.60 (m, 5H), 5.52 (d, J = 16.0 Hz, 2H), 6.64 (q, J = 7.0 Hz, 2H), 7.39 (m, 2H), 7.50 (d, J = 7.6 Hz, 1H), 7.65 (q, J = 7.9 Hz, 9H), 7.79 (dd, J = 24.7, 8.0 Hz, 3H), 8.03 (d, J = 8.5 Hz, 2H), 8.23 (s, 2H), 8.97 (s, 2H)
63f	¹ H NMR (400 MHz, CDCl ₃): δ ppm 0.87 (dd, J = 16.5, 9.8 Hz, 2H), 1.28 (m, 12H), 1.58 (m, 4H), 1.79 (dd, J = 13.1, 6.9 Hz, 1H), 2.12 (dd, J = 13.1, 8.9 Hz, 1H), 2.90 (d,

	J = 10.7 Hz, 1H), 3.00 (d, J = 10.6 Hz, 1H), 3.52 (dt, J = 11.4, 5.4 Hz, 7H), 3.96 (t, J = 7.8 Hz, 1H), 4.21 (q, J = 7.2 Hz, 2H), 4.61 (s, 2H), 5.54 (s, 1H), 6.64 (q, J = 7.1 Hz, 1H), 7.66 (q, J = 8.4 Hz, 5H), 8.12 (t, J = 4.2 Hz, 2H), 8.94 (s, 1H)
63g	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (m, 4H), 1.64 (dq, J = 14.1, 8.9, 7.2 Hz, 4H), 2.03 (td, J = 13.3, 8.9 Hz, 1H), 2.49 (dd, J = 13.6, 8.7 Hz, 1H), 3.27 (s, 2H), 3.58 (m, 4H), 3.88 (d, J = 11.8 Hz, 6H), 4.32 (qd, J = 7.2, 2.5 Hz, 2H), 4.58 (t, J = 8.8 Hz, 1H), 6.62 (q, J = 7.1 Hz, 1H), 7.03 (m, 1H), 7.19 (m, 2H), 7.57 (d, J = 8.2 Hz, 2H), 7.64 (m, 2H)
63h	¹ H NMR (400 MHz, CDCl ₃): δ ppm 7.769-7.796 (m, 1H), 7.707-7.740 (m, 2H), 7.585-7.636 (m, 4H), 7.422-7.444 (m, 1H), 6.738-6.762 (m, 1H), 6.579-6.658-6.537 (m, 1H), 5.521 (s, 1H), 4.612 (s, 2H), 4.200-4.253 (q, 2H), 4.114-4.154 (t, 1H), 3.747 (s, 3H), 3.475-3.523 (m, 4H), 3.047-3.160 (m, 2H), 2.171-2.726 (m, 1H), 1.840-1.891 (m, 1H), 1.543-1.649 (m, 4H), 1.209-1.305 (t, 3H)
63i	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.25 (t, J=7.10 Hz, 3 H) 1.42 - 1.69 (m, 4 H) 1.92 (dd, J=13.25, 9.35 Hz, 1 H) 2.35 (dd, J=13.25, 8.47 Hz, 1 H) 3.14 (br. s., 2 H) 3.60 (br. s., 4 H) 4.24 (qd, J=7.09, 2.10 Hz, 2 H) 4.54 (br. s., 1 H) 5.77 (br. s., 1 H) 6.70 (q, J=6.65 Hz, 1 H) 7.37 (d, J=2.10 Hz, 1 H) 7.43 - 7.52 (m, 3 H) 7.53 - 7.69 (m, 4 H) 9.23 (br. s., 1 H) 10.44 (br. s., 1 H)
63j	¹ H NMR (400 MHz, CD ₃ OD): δ ppm 7.66-7.64 (d, 1 H, J=8.6 Hz), 7.43-7.41 (d, 1 H, J=8.6 Hz), 7.26-7.20 (m, 2 H), 6.82-6.68 (m, 4 H), 5.42 (s, 1 H), 4.19-4.16 (q, 1 H, J=7.0 Hz), 3.83-3.81 (t, 1 H), 3.49-3.47 (m, 4 H), 2.91-2.89 (d, 1 H, J=10.9 Hz), 2.77-2.75 (d, 1 H, J=10.9 Hz), 2.11-2.07 (m, 1 H), 2.12-2.10 (m, 1 H), 1.53-1.51 (m, 4 H), 1.28-1.25 (t, 3 H, J=7.0 Hz Hz)
63k	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.42 (s, 1 H), 8.30 (d, 1 H), 7.61 (m, 2 H), 7.31 (m, 1 H), 7.10 (s, 1 H), 6.58 (m, 1 H), 5.57 (s, 1 H), 4.20 (m, 2 H), 3.84 (m, 1 H), 3.48 (m, 4 H), 3.16 (s, 3 H), 2.77 (m, 1 H), 2.70 (m, 1 H), 2.61 (m, 1 H), 2.14 (m, 1 H), 1.77 (m, 1 H), 1.65 (m, 2 H), 1.54 (m, 4 H), 1.20 (m, 3 H), 0.98 (m, 3 H)
63l	¹ H NMR (400 MHz, CD ₃ OD-d ₄): δ ppm 1.28-1.24 (m, 4 H), 7.72-7.68 (m, 3 H), 1.52 (m, 4 H), 1.71 (m, 1 H), 2.10 (m, 1 H), 2.32 (s, 3 H), 2.76-2.73 (m, 1 H), 2.90-2.87 (m, 1 H), 3.49 (m, 4 H), 3.75 (s, 3 H), 3.81 (m, 2 H), 4.20-4.15 (m, 2 H), 5.54 (s, 1H), 6.65-6.59 (m, 1 H), 6.93 (s, 1 H), 7.37-7.34 (m, 1 H), 7.45-7.42 (m, 1 H), 7.57-7.55 (m, 2 H)
63m	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.19 (s, 1 H), 7.64 (d, 1 H), 7.53-7.42 (m, 5 H), 7.29 (s, 1 H), 7.16 (s, 2 H), 6.63 (q, 1 H), 6.52-6.35 (m, 3 H), 5.80-5.77 (d, 1 H), 5.50 (s, 1 H), 4.21-4.18 (m, 2 H), 3.97 (t, 1 H), 3.49 (m, 4 H), 2.98-2.95 (d, 1 H), 2.86-2.83 (d, 1 H), 2.16-2.14 (m, 1 H), 1.80-1.76 (m, 1 H), 1.54 (m, 4 H), 1.19-1.16 (t, 3 H)
63n	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (q, J = 7.1, 6.1 Hz, 4H), 1.52 (dt, J = 10.4, 5.7 Hz, 4H), 1.74 (dd, J = 13.0, 7.2 Hz, 1H), 2.09 (dd, J = 13.1, 8.8 Hz, 1H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.52 (tq, J = 14.5, 8.2 Hz, 4H), 3.82 (dd, J = 8.8, 7.2 Hz, 1H), 3.90 (s, 3H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 4.89 (d, J = 1.5 Hz, 13H), 5.51 (d, J = 18.8 Hz, 2H), 6.62 (q, J = 7.1 Hz, 1H), 7.15 (t, J = 8.6 Hz, 1H), 7.39 (m, 2H), 7.59 (m, 4H)
63o	¹ H-NMR (400 MHz, MeOH-d ₄): δ ppm 8.00(d,J=2.36 Hz,1H) ,7.88(dd, J=2.6,6.76 Hz,1H), 7.58(m,4H),6.62(m, 2H),5.55(s, 1H),4.22(m,3H) ,3.64(s,3H),3.52(m, 4H),3.02(m, 2H),2.27(m,1H),1.89(m, 1H),1.59(m,4H),1.29(t, J=7.16 Hz,3H).
63p	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.62(d,2H,J=8.0), 7.56(d,2H,J=8.0), 7.45(m,2H), 6.93(d,1H,J=8.0), 6.62 (q, 1H,J=8.0), 5.54 (s, 1H), 4.21 (t, 2H,

	J=4.0),3.99 (t, 1H, J=4.0), 3.56δ3.49 (m, 4H), 3.00(dd, 3H,J=20.0,8.0), 2.86 (d, 1H, J = 8.0),2.59-2.57 (m, 2H), 2.18 (dd, 2H, J = 12.0,8.0), 1.80(dd, 2H, J = 8.0,4.0), 1.54(m, 5),1.28(t, 4H, J = 8.0)
63q	¹ H-NMR (400 MHz, CDCl ₃): δ ppm 11.76 (m,1H) ,7.84 -7.86 (m,1H) ,7.70 -7.73 (m,2H),7.59 (m, 4H),7.43 -7.45 (m, 1H),6.73 -6.75 (m,1H), 6.57 -6.63 (q, 1H), 5.53 (s, 1H), 4.61 (s,2H) , 4.17 -4.23 (q,2H), 3.90 -3.94 (t, 1H), 3.48 -3.51 (m, 4H), 2.86 -2.99 (m,2H), 2.07-2.12 (m, 2H), 1.74 -1.79 (m, 1H), 1.53 -1.59 (m, 4H),1.24 -1.29 (t, 3H).
63r	¹ H NMR (400 MHz, CD ₃ OD-d ₄): δ ppm 8.42 (s, 1 H), 8.04 (d, 1 H), 7.79 (m, 3 H), 7.63 (m, 1 H), 7.46 (m, 1 H), 6.58 (m, 1 H), 6.40 (m, 1 H), 5.56 (m, 1 H), 4.18 (m, 2 H), 3.83 (m, 1 H), 3.50 (m, 4 H), 3.21(s, 3 H), 2.90 (m, 1 H), 2.78 (m, 1 H),2.14 (m, 1 H), 1.86 (m, 3 H), 1.76(m, 1 H), 1.54 (m, 4 H), 1.26 (m, 3 H)
63s	¹ H NMR (MeOH-d ₄): δ ppm 0.90 (t, J = 6.9 Hz, 1H), 1.17 (p, J = 6.3 Hz, 3H), 1.29 (s, 2H), 1.56 (m, 4H), 1.80 (s, 1H), 2.29 (s, 1H), 2.41 (s, 3H), 2.80 (m, 4H), 3.26 (d, J = 11.3 Hz, 1H), 3.44 (s, 1H), 4.09 (tdd, J = 14.2, 7.9, 4.6 Hz, 2H), 4.48 (s, 1H), 4.87 (s, 2H), 5.56 (s, 1H), 6.42 (t, J = 2.2 Hz, 1H), 6.93 (m, 1H), 7.46 (m, 4H), 7.61 (m, 2H), 7.80 (dd, J = 8.3, 2.2 Hz, 1H), 7.93 (dd, J = 5.2, 2.5 Hz, 2H)
63t	¹ H NMR (MeOH-d ₄): δ ppm 0.91 (dd, J = 12.4, 6.3 Hz, 2H), 1.17 (q, J = 7.4 Hz, 3H), 1.31 (d, J = 16.3 Hz, 3H), 1.65 (m, 4H), 1.83 (s, 1H), 2.32 (s, 2H), 2.41 (s, 3H), 2.92 (ddt, J = 18.2, 14.3, 9.1 Hz, 5H), 3.28 (s, 1H), 4.11 (dt, J = 10.7, 7.1, 3.9 Hz, 2H), 4.48 (s, 1H), 4.95 (d, J = 11.7 Hz, 1H), 6.43 (d, J = 2.2 Hz, 1H), 6.94 (q, J = 6.5 Hz, 1H), 7.50 (m, 3H), 7.61 (dd, J = 8.6, 2.1 Hz, 2H), 7.79 (dt, J = 8.3, 1.4 Hz, 1H), 7.93 (dd, J = 10.2, 3.2 Hz, 2H)
63u	¹ H NMR (400 MHz, CDCl ₃ -d) : δ ppm 8.50 (s, 1H), 7.99-7.96 (m, 1H), 7.69-7.63 (m, 2H), 7.51 (s, 1H), 7.25-7.23 (m, 1H), 7.11 (d, J = 7.8 Hz, 1H), 6.57 (q, J = 6.6 Hz, 1H), 5.51 (s, 1H), 5.18 (s, 2H), 4.21 (q, J = 7.1 Hz, 2H), 3.93-3.89 (m, 1H), 3.53-3.48 (m, 4H), 3.14 (s, 3H), 2.93 (dd, J ₁ = 10.6 Hz, J ₂ = 42.0 Hz, 2H), 2.66-2.62 (m, 2H), 2.12-2.07 (m, 1H), 1.79-1.74 (m, 1H), 1.69-1.63 (m, 2H), 1.61-1.58 (m, 2H), 1.55-1.52 (m, 2H), 1.29 (t, J = 7.2 Hz, 3H), 0.95 (t, J = 7.3 Hz, 3H)
63v	¹ H NMR (400 MHz, CDCl ₃): δ ppm 8.52 (d, J = 9.5 Hz, 1H), 7.99-7.97 (m, 1H), 7.69-7.62 (m, 3H), 7.38 (dd, J ₁ = 7.9 Hz, J ₂ = 20.6 Hz, 1H), 7.15 (dd, J ₁ = 7.9 Hz, J ₂ = 17.0 Hz, 1H), 6.57 (m, 1H), 6.46-6.42 (m, 1H), 6.34-5.84 (m, 1H), 5.51-5.50 (m, 1H), 5.18 (s, 2H), 4.20 (q, J = 7.2 Hz, 2H), 3.87 (t, J = 7.6 Hz, 1H), 3.52-3.50 (m, 4H), 3.15-3.14 (m, 3H), 2.90 (dd, J ₁ = 10.2 Hz, J ₂ = 47.8 Hz, 2H), 2.10-2.05 (m, 1H), 1.91 (d, J = 6.4 Hz, 3H), 1.78-1.73 (m, 1H), 1.58-1.53 (m, 4H), 1.28 (t, J = 7.1 Hz, 3H)
63w	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.49 (s, 1H), 7.98 (d, J = 7.5 Hz, 1H), 7.71 (t, J = 7.8 Hz, 1H), 7.65-7.63 (m, 2H), 7.42-7.40 (m, 1H), 7.23 (d, J = 1.9 Hz, 1H), 6.53 (q, J = 6.8 Hz, 1H), 5.48 (s, 1H), 5.26 (s, 2H), 4.22 (q, J = 7.1 Hz, 2H), 4.03 (t, J = 8.0 Hz, 1H), 3.52-3.51 (m, 4H), 3.21 (q, J = 7.4 Hz, 2H), 3.02 (dd, J ₁ = 10.9 Hz, J ₂ = 34.9 Hz, 2H), 2.18-2.12 (m, 1H), 1.85-1.80 (m, 1H), 1.62-1.61 (m, 2H), 1.56-1.55 (m, 2H), 1.31-1.28 (m, 6H)
63x	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.45 (s, 1H),8.03(d,1H,J=8.0), 7.83(t,1H,J=8.0), 7.79-7.69(m,2H),7.50(d,1H,J=8.0), 7.37(s,1H), 6.62 (q, 1H,J=8.0), 5.61 (s, 1H), 4.38 (t, 1H,J=8.0),4.32-4.27 (m, 2H), 3.64-3.49(m, 4H),3.16(q, 2H,J=12.0), 2.39 (dd, 1H, J = 12.0,8.0),1.98 (dd, 1H, J = 12.0,8.0), 1.70-1.62(m, 7H),1.31(dd, 5H, J = 12.0,8.0) 0.96(t, 4H,J=8.0)

63y	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.50 (s, 1 H), 8.05-8.03 (d, 1 H), 7.82 (t, 1 H), 7.76-7.70 (m, 2 H), 7.52-7.51 (d, 1 H), 7.38-7.37 (d, 1 H), 6.62-6.60 (q, 1 H), 5.60 (s, 1 H), 4.20-4.19 (q, 2 H), 3.85 (t, 1 H), 3.57 (m, 4 H), 2.82 (d, 1 H), 2.77 (d, 1 H), 2.09 (m, 1 H), 1.77 (m, 1 H), 1.57 (m, 6 H), 1.31 (q, 2 H), 1.26 (m, 3 H), 0.84-0.81 (t, 3 H)
63z	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.09 (dd, J = 4.2, 2.0 Hz, 1H), 0.90 (t, J = 6.5 Hz, 3H), 1.28 (m, 14H), 1.54 (dt, J = 10.5, 5.6 Hz, 6H), 1.76 (dd, J = 13.2, 7.3 Hz, 1H), 2.03 (s, 1H), 2.15 (ddd, J = 29.1, 14.1, 8.4 Hz, 2H), 2.78 (d, J = 10.9 Hz, 1H), 2.92 (d, J = 11.0 Hz, 1H), 3.53 (td, J = 13.8, 13.4, 6.0 Hz, 6H), 3.86 (dd, J = 8.8, 7.3 Hz, 1H), 4.20 (m, 3H), 5.46 (d, J = 22.4 Hz, 3H), 5.56 (s, 1H), 6.67 (q, J = 7.2 Hz, 2H), 7.66 (d, J = 8.1 Hz, 3H), 7.75 (m, 3H), 7.85 (m, 3H), 7.93 (d, J = 7.9 Hz, 2H)
63aa	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm: 8.16 (d, J = 8.84 Hz, 1H), 8.03 (d, J = 1.84 Hz, 1H), 7.94-7.81 (m, 2H), 7.77 (d, J = 8.32 Hz, 2H), 7.64 (d, J = 8.24 Hz, 2H), 6.96 (d, J = 8.88 Hz, 1H), 6.65 (q, J = 7.08 Hz, 1H), 5.56 (s, 1H), 4.18 (m, 2H), 4.06 (s, 3H), 3.82 (m, 1H), 3.53 (m, 4H), 2.90 (d, J = 11.0 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.09 (m, 1H), 1.75 (m, 1H), 1.53 (s, 4H), 1.27 (t, J = 7.12 Hz, 3H)
63ab	¹ H NMR (400 MHz, CDCl ₃): δ ppm 7.76 (s, 1H), 7.65 (d, J = 8.5 Hz, 1H), 7.43 (t, J = 7.6 Hz, 1H), 7.37 (dd, J ₁ = 2.2 Hz, J ₂ = 8.5 Hz, 1H), 7.33 (d, J = 7.6 Hz, 1H), 7.28 (d, J = 7.8 Hz, 1H), 7.24 (d, J = 2.2 Hz, 1H), 6.63 (q, J = 6.7 Hz, 1H), 5.41 (s, 1H), 5.02 (s, 2H), 4.80 (m, 2H), 4.21 (q, J = 7.1 Hz, 2H), 4.05-4.01 (m, 1H), 3.48-3.46 (m, 4H), 3.01 (dd, J ₁ = 10.9 Hz, J ₂ = 31.0 Hz, 2H), 2.17-2.11 (m, 1H), 1.83-1.78 (m, 1H), 1.59-1.50 (m, 4H), 1.28 (t, J = 7.1 Hz, 3H)
63ac	¹ H-NMR (400 MHz, MeOH-d ₄): δ ppm 7.97 (s, 1H), 7.60-7.67 (m, 2H), 7.52-7.56 (m, 1H), 7.43-7.45 (m, 1H), 7.31-7.31 (m, 1H), 7.22-7.24 (m, 1H), 6.621-6.663 (m, 1H), 5.498 (s, 1H), 4.16-4.22 (m, 2H), 4.92-4.03 (m, 2H), 3.83-3.87 (M, 1h), 3.46-3.53 (m, 4H), 2.90-2.92 (d, 1H), 2.76-2.78 (d, 1H), 2.59-2.63 (m, 2H), 2.10-2.22 (m, 3H), 1.71-1.78 (m, 1H), 1.52-1.55 (m, 4H), 1.25-1.28 (m, 3H)
63ad	¹ H-NMR (400 MHz, MeOH-d ₄) δ ppm: 7.98 (s, 1H), 7.63-7.65 (m, 1H), 7.42-7.50 (m, 3H), 7.30-7.30 (m, 1H), 7.05-7.07 (m, 1H), 6.62-6.67 (m, 1H), 5.49 (s, 1H), 4.15-4.22 (m, 2H), 3.80-3.98 (m, 3H), 3.46-3.56 (m, 6h), 2.73-2.92 (m, 4H), 2.73-2.78 (d, 1H), 2.07-2.13 (d, 1H), 1.73-1.78 (m, 1H), 1.49-1.57 (m, 4H), 1.25-1.29 (m, 3H)
63ae	¹ H NMR (400 MHz, DMSO-d ₆) δ ppm: 1.25 (t, J = 7.10 Hz, 3 H) 1.42 - 1.69 (m, 4 H) 1.92 (dd, J = 13.25, 9.35 Hz, 1 H) 2.35 (dd, J = 13.25, 8.47 Hz, 1 H) 3.14 (br. s., 2 H) 3.60 (br. s., 4 H) 4.24 (qd, J = 7.09, 2.10 Hz, 2 H) 4.54 (br. s., 1 H) 5.77 (br. s., 1 H) 6.70 (q, J = 6.65 Hz, 1 H) 7.37 (d, J = 2.10 Hz, 1 H) 7.43 - 7.52 (m, 3 H) 7.53 - 7.69 (m, 4 H) 9.23 (br. s., 1 H) 10.44 (br. s., 1 H)
63af	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.66 (d, 1 H, J = 8.4 Hz), 7.50 (m, 3 H), 7.31 (d, 2 H, J = 8.7 Hz), 7.24 (d, 1 H, J = 7.2 Hz), 6.64 (m, 1 H), 5.50 (m, 1 H), 4.21 (m, 1 H), 3.87 (m, 1 H), 3.53 (m, 4 H), 3.01 (s, 3 H), 3.18 (m, 1 H), 2.90 (m, 3 H), 2.79 (m, 1 H), 2.07 (m, 1 H), 1.74 (m, 1 H), 1.53 (m, 4 H), 1.27 (m, 3 H)
63ag	¹ H NMR (400 MHz, CDCl ₃): δ ppm 7.54 (d, J = 2.2 Hz, 1H), 7.51 (d, J = 8.6 Hz, 1H), 7.19 (dd, J ₁ = 2.6 Hz, J ₂ = 8.6 Hz, 1H), 6.85 (q, J = 6.6 Hz, 1H), 5.49 (s, 1H), 4.56 (s, 2H), 4.20 (q, J = 7.2 Hz, 2H), 3.90-3.86 (m, 1H), 3.53-3.47 (m, 4H), 2.90 (dd, J ₁ = 10.4 Hz, J ₂ = 47.6 Hz, 2H), 2.13-2.05 (m, 1H), 1.78-1.73 (m, 1H), 1.59-1.56 (m, 2H), 1.54-1.51 (m, 2H), 1.28 (t, J = 7.1 Hz, 3H)
63ah	¹ H NMR (400 MHz, CDCl ₃): δ ppm 7.54 (d, J = 2.2 Hz, 1H), 7.51 (d, J = 8.6 Hz, 1H),

	7.19 (dd, J1 = 2.6 Hz, J2 = 8.6 Hz, 1H), 6.85 (q, J = 6.6 Hz, 1H), 5.49 (s, 1H), 4.56 (s, 2H), 4.20 (q, J = 7.2 Hz, 2H), 3.90-3.86 (m, 1H), 3.53-3.47 (m, 4H), 2.90 (dd, J1 = 10.4 Hz, J2 = 47.6 Hz, 2H), 2.13-2.05 (m, 1H), 1.78-1.73 (m, 1H), 1.59-1.56 (m, 2H), 1.54-1.51 (m, 2H), 1.28 (t, J = 7.1 Hz, 3H)
63ai	¹ H NMR (400 MHz, MeOH-d4): δ ppm 8.06 (d, J = 8.72 Hz, 1H), 8.01 (s, 1H), 7.94 (s, 2H), 7.76 (d, J = 8.28 Hz, 2H), 7.64 (d, J = 8.16 Hz, 2H), 7.31 (d, J = 8.68 Hz, 1H), 6.66 (q, J = 7.32 Hz, 1H), 5.56 (s, 1H), 4.18 (q, J = 7.04 Hz, 2H), 3.84-3.80 (m, 1H), 3.51 (m, 4H), 2.89 (d, J = 10.96 Hz, 1H), 2.75 (d, J = 11 Hz, 1H), 2.68 (s, 3H), 2.10-2.01 (m, 1H), 1.76-1.71 (m, 1H), 1.54-1.49 (m, 4H), 1.25 (t, J = 7.12 Hz, 3H)
63aj	¹ H NMR (400 MHz, MeOH-d4): δ ppm 7.69 (d, J = 1.8 Hz, 1H), 7.60-7.57 (d, 1H), 7.48 (dd, J1 = 2.44, J2 = 8.6 Hz, 1H), 6.96 (q, J = 7.32 Hz, 1H), 5.56 (s, 1H), 4.19 (q, J = 7.12 Hz, 2H), 3.86-3.82 (m, 1H), 3.54 (m, 4H), 2.91 (d, J = 11 Hz, 1H), 2.77 (d, J = 11 Hz, 1H), 2.14-2.08 (m, 1H), 1.79-1.73 (m, 1H), 1.55 (m, 4H), 1.27 (t, J = 7.12 Hz, 3H)
63ak	¹ H NMR (400 MHz, MeOH-d4): δ ppm 7.91 (s, 1H), 7.71 (dd, J1 = 6.12 Hz, J2 = 1.96 Hz, 1H), 7.63 (m, 2H), 7.56-7.49 (m, 7H), 7.39-7.35 (m, 2H), 6.74 (q, J = 6.88 Hz, 1H), 5.50 (s, 1H), 4.18 (q, J = 6.96 Hz, 2H), 3.83 (m, 1H), 3.50 (m, 4H), 2.89 (d, J = 11.04 Hz, 1H), 2.75 (d, J = 11 Hz, 1H), 2.12-2.06 (m, 1H), 1.76-1.71 (m, 1H), 1.54-1.49 (m, 4H), 1.27 (t, J = 7.12 Hz, 3H)
63al	¹ H NMR (400 MHz, MeOH-d4): δ ppm 0.84 (t, J = 7.1 Hz, 8H), 1.26 (t, J = 7.1 Hz, 8H), 1.50 (dt, J = 11.1, 5.8 Hz, 4H), 1.73 (dd, J = 13.1, 7.1 Hz, 1H), 2.06 (dd, J = 13.1, 8.8 Hz, 1H), 2.38 (s, 3H), 2.73 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.51 (m, 4H), 3.81 (dd, J = 8.7, 7.1 Hz, 1H), 4.00 (qd, J = 7.1, 4.5 Hz, 2H), 4.17 (qd, J = 7.1, 1.5 Hz, 2H), 5.74 (s, 1H), 6.39 (d, J = 2.3 Hz, 1H), 6.85 (q, J = 6.7 Hz, 1H), 7.45 (m, 5H), 7.80 (m, 2H), 7.90 (d, J = 2.4 Hz, 1H)
63am	¹ H NMR (400 MHz, MeOH-d4): δ ppm 0.89 (m, 1H), 1.26 (m, 7H), 1.40 (t, J = 7.1 Hz, 3H), 1.51 (m, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.05 (m, 1H), 2.40 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.53 (m, 4H), 3.81 (dd, J = 8.8, 7.1 Hz, 1H), 4.18 (qd, J = 7.1, 2.5 Hz, 2H), 4.39 (q, J = 7.1 Hz, 2H), 5.73 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.82 (q, J = 6.6 Hz, 1H), 7.57 (t, J = 7.8 Hz, 1H), 7.68 (d, J = 1.9 Hz, 1H), 7.79 (m, 2H), 7.90 (dt, J = 8.0, 1.4 Hz, 1H), 8.02 (m, 2H), 8.28 (d, J = 1.9 Hz, 1H)
63an	¹ H NMR (400 MHz, MeOH-d4): δ ppm 1.33 (dt, J = 54.3, 7.1 Hz, 6H), 1.50 (dt, J = 10.8, 5.8 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.40 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.5 Hz, 2H), 4.38 (q, J = 7.1 Hz, 2H), 5.74 (s, 1H), 6.43 (d, J = 2.3 Hz, 1H), 6.84 (q, J = 6.6 Hz, 1H), 7.77 (m, 5H), 8.00 (d, J = 2.3 Hz, 1H), 8.09 (m, 2H)
63ao	¹ H NMR (400 MHz, MeOH-d4): δ ppm 1.27 (t, J = 7.1 Hz, 3H), 1.54 (m, 4H), 1.76 (dd, J = 13.1, 7.2 Hz, 1H), 2.11 (dd, J = 13.1, 8.7 Hz, 1H), 2.77 (dd, J = 11.0, 1.1 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.53 (td, J = 11.9, 11.4, 4.9 Hz, 4H), 3.84 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.2, 1.6 Hz, 2H), 5.53 (s, 1H), 7.15 (t, J = 8.0 Hz, 1H), 7.27 (q, J = 8.0 Hz, 1H), 7.69 (m, 2H)
63ap	¹ H NMR (400 MHz, CDCl3): δ ppm 1.28 (m, 4H), 1.56 (dq, J = 25.2, 5.5, 4.9 Hz, 4H), 1.78 (dd, J = 13.1, 6.9 Hz, 1H), 2.12 (m, 1H), 2.90 (d, J = 10.7 Hz, 1H), 2.99 (d, J = 10.6 Hz, 1H), 3.49 (dt, J = 11.5, 5.7 Hz, 4H), 3.94 (dd, J = 8.8, 6.9 Hz, 1H), 4.21 (q,

	J = 7.1 Hz, 2H), 4.58 (s, 2H), 5.43 (s, 1H), 6.55 (q, J = 6.8 Hz, 1H), 7.24 (m, 3H), 7.41 (m, 3H), 7.65 (m, 2H)
63aq	¹ H NMR (400 MHz, CDCl ₃): δ ppm 0.84 (m, 2H), 1.14 (s, 1H), 1.28 (t, J = 7.1 Hz, 3H), 1.53 (m, 4H), 1.74 (dd, J = 13.1, 6.8 Hz, 1H), 2.05 (m, 1H), 2.43 (s, 3H), 2.82 (d, J = 10.5 Hz, 1H), 2.94 (d, J = 10.5 Hz, 1H), 3.46 (dt, J = 14.0, 5.8 Hz, 4H), 3.86 (dd, J = 8.8, 6.7 Hz, 1H), 4.19 (q, J = 7.1 Hz, 2H), 4.35 (s, 2H), 5.40 (s, 1H), 6.61 (q, J = 6.8 Hz, 1H), 7.33 (m, 5H), 7.65 (d, J = 8.5 Hz, 1H)
63ar	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 4H), 1.54 (dt, J = 7.9, 4.7 Hz, 4H), 1.76 (dd, J = 13.1, 7.2 Hz, 1H), 2.12 (m, 1H), 2.78 (m, 1H), 2.90 (m, 1H), 3.52 (m, 4H), 3.85 (td, J = 9.2, 8.8, 7.3 Hz, 1H), 4.19 (qd, J = 7.1, 1.6 Hz, 2H), 4.86 (d, J = 0.8 Hz, 1H), 5.51 (d, J = 13.8 Hz, 1H), 6.52 (q, J = 6.7 Hz, 1H), 7.34 (d, J = 2.2 Hz, 1H), 7.51 (dd, J = 8.5, 2.2 Hz, 1H), 7.75 (m, 5H)
63as	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (t, J = 7.1 Hz, 3H), 1.67 (m, 4H), 2.05 (dd, J = 13.6, 8.8 Hz, 1H), 2.46 (d, J = 45.5 Hz, 7H), 2.66 (s, 1H), 3.28 (s, 2H), 3.69 (m, 4H), 4.32 (qd, J = 7.1, 2.3 Hz, 2H), 4.58 (t, J = 8.7 Hz, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.85 (q, J = 6.3 Hz, 1H), 7.37 (m, 2H), 7.66 (m, 3H), 7.79 (m, 2H), 7.97 (d, J = 2.4 Hz, 1H)
63at	¹ H NMR (MeOH-d ₄): δ ppm 1.28 (m, 15H), 1.53 (m, 13H), 1.76 (dd, J = 13.1, 7.3 Hz, 3H), 1.86 (s, 1H), 2.12 (dd, J = 13.1, 8.8 Hz, 3H), 2.79 (d, J = 11.0 Hz, 3H), 2.92 (d, J = 11.0 Hz, 3H), 3.51 (qdt, J = 18.0, 13.3, 5.9 Hz, 12H), 3.63 (d, J = 8.6 Hz, 1H), 3.87 (m, 3H), 4.19 (qd, J = 7.1, 1.6 Hz, 5H), 5.51 (s, 3H), 6.68 (q, J = 6.7 Hz, 3H), 7.29 (m, 6H), 7.47 (m, 19H), 7.65 (m, 5H)
63au	¹ H NMR (MeOH-d ₄): δ ppm 1.13 (s, 2H), 1.26 (t, J = 7.3 Hz, 4H), 1.49 (m, 6H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.06 (dd, J = 13.1, 8.7 Hz, 1H), 2.38 (d, J = 12.1 Hz, 7H), 2.73 (d, J = 11.0 Hz, 1H), 2.87 (d, J = 11.0 Hz, 1H), 3.53 (tt, J = 14.1, 5.1 Hz, 5H), 3.81 (m, 1H), 4.18 (tt, J = 7.8, 3.6 Hz, 2H), 4.81 (s, 2H), 4.97 (d, J = 15.9 Hz, 1H), 5.74 (s, 1H), 6.41 (d, J = 2.1 Hz, 1H), 6.78 (q, J = 6.7 Hz, 1H), 7.26 (d, J = 7.9 Hz, 2H), 7.57 (m, 5H), 7.73 (m, 2H), 7.96 (d, J = 2.3 Hz, 1H)
63av	¹ H NMR (MeOH-d ₄): δ ppm 1.26 (m, 3H), 1.51 (dt, J = 10.6, 5.6 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.40 (s, 7H), 2.74 (d, J = 10.9 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.54 (m, 4H), 3.81 (dd, J = 8.8, 7.1 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.74 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.79 (q, J = 6.6 Hz, 1H), 7.21 (d, J = 7.5 Hz, 1H), 7.33 (t, J = 7.6 Hz, 1H), 7.46 (m, 2H), 7.62 (d, J = 1.9 Hz, 1H), 7.75 (m, 2H), 7.98 (d, J = 2.4 Hz, 1H)
63aw	¹ H NMR (MeOH-d ₄): δ ppm 0.90 (m, 1H), 1.27 (m, 5H), 1.51 (dt, J = 10.5, 5.6 Hz, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.09 (dd, J = 13.1, 8.7 Hz, 1H), 2.40 (s, 3H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.54 (m, 4H), 3.84 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.1, 1.7 Hz, 2H), 5.73 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.84 (q, J = 6.5 Hz, 1H), 7.64 (m, 3H), 7.80 (m, 3H), 8.01 (d, J = 2.4 Hz, 1H)
63ax	¹ H NMR (400 MHz, Chloroform-d): δ ppm 1.27 (m, 9H), 1.52 (dt, J = 22.3, 5.4 Hz, 4H), 1.72 (d, J = 13.1 Hz, 1H), 2.05 (m, 1H), 2.25 (ddd, J = 18.1, 13.9, 8.2 Hz, 2H), 2.63 (m, 2H), 2.82 (d, J = 10.5 Hz, 1H), 2.94 (d, J = 10.4 Hz, 1H), 3.48 (dd, J = 13.6, 7.4 Hz, 5H), 3.63 (m, 1H), 3.81 (m, 4H), 4.19 (q, J = 7.1 Hz, 2H), 4.86 (s, 2H), 5.46 (s, 1H), 6.46 (m, 1H), 7.22 (d, J = 2.2 Hz, 1H), 7.34 (dd, J = 8.5, 2.2 Hz, 1H), 7.59 (d, J = 8.5 Hz, 1H)
63ay	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (d, J = 2.64 Hz, 4 H) 1.61 - 1.86 (m, 5 H)

	2.04 - 2.16 (m, 1 H) 2.42 (d, J=1.27 Hz, 3 H) 2.48 - 2.60 (m, 1 H) 3.55 - 4.03 (m, 4 H) 4.25 - 4.44 (m, 2 H) 4.55 - 4.70 (m, 1 H) 6.45 (s, 1 H) 6.90 - 7.04 (m, 1 H) 7.61 (s, 2 H) 7.68 - 7.79 (m, 1 H) 7.88 - 8.00 (m, 1 H)
63az	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.25 (t, J=7.15 Hz, 3 H) 1.43 - 1.60 (m, 4 H) 1.78 (dd, J=13.13, 7.42 Hz, 1 H) 2.13 (dd, J=13.08, 8.74 Hz, 1 H) 2.35 (s, 3 H) 2.73 - 3.01 (m, 2 H) 3.39 - 3.63 (m, 4 H) 3.94 (t, J=7.91 Hz, 1 H) 4.18 (qd, J=7.13, 1.78 Hz, 2 H) 5.65 (s, 1 H) 6.38 (d, J=2.39 Hz, 1 H) 6.79 (q, J=6.74 Hz, 1 H) 7.41 - 7.54 (m, 2 H) 7.68 (d, J=8.35 Hz, 1 H) 7.91 (d, J=2.34 Hz, 1 H)
63ba	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.33 (t, J = 7.13 Hz, 3 H), 1.53 - 1.75 (m, 4 H), 2.05 (dd, J = 13.62, 8.88 Hz, 1 H), 2.49 (dd, J = 13.57, 8.74 Hz, 1 H), 3.27 (s, 2 H), 3.42 - 3.74 (m, 4 H), 3.84(s, 3 H), 4.25 - 4.40 (m, 2 H), 4.58 (t, J = 8.79 Hz, 1 H), 5.60 (s, 1 H), 6.59 - 6.71 (m, 1 H), 6.92 (ddd, J = 8.22, 2.54, 0.76 Hz, 1 H), 7.11 - 7.16 (m, 1 H), 7.16 - 7.24 (m, 1 H), 7.31 - 7.38 (m, 1 H), 7.54 - 7.61 (m, 2 H), 7.62 - 7.70 (m, 2 H)
63bb	¹ H NMR (400 MHz, dichloromethane-d ₂): δ ppm 1.29 (t, J=7.15 Hz, 3 H) 1.47 - 1.85 (m, 4 H) 2.01 (dd, J=13.52, 8.30 Hz, 1 H) 2.30 - 2.36(m, 1 H) 2.38 (s, 3 H) 3.27 - 3.41 (m, 2 H) 3.41 - 3.67 (m, 4 H) 3.82 (s, 3 H) 4.26 (qd, J=7.17, 4.00 Hz, 2 H) 4.45 (t, J=8.49 Hz, 1 H) 4.96 (br. s, 2H) 5.49 (s, 1 H) 6.31 (d, J=2.25 Hz, 1 H) 6.62 (q, J=6.90 Hz, 1 H) 6.88 (d, J=2.59 Hz, 1 H) 6.96 (dd, J=8.81, 2.61 Hz, 1 H) 7.61 (d, J=8.74 Hz, 1 H) 7.66 (d, J=2.25 Hz, 1 H)
63bc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (t, J=7.22 Hz, 3 H) 1.67 - 1.89 (m, 4 H) 2.05 - 2.18 (m, 1 H) 2.49 - 2.62 (m, 1 H) 3.56 - 3.90 (m, 4 H) 4.35 (dd, J=7.13, 1.85 Hz, 2 H) 4.65 (s, 1 H) 5.97 (s, 1 H) 6.58 - 6.72 (m, 1 H) 7.14 (br. s., 1 H) 7.41 (d, J=9.18 Hz, 1 H) 7.45 - 7.53 (m, 2 H) 7.64 - 7.72 (m, 2 H) 7.73 - 7.82 (m, 2 H)
63bd	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 - 1.39 (m, 4 H) 1.74 (d, J=18.35 Hz, 4 H) 2.03 - 2.14 (m, 1 H) 2.35 (d, J=12.89 Hz, 6 H) 2.43 (s, 3 H) 2.46 - 2.57 (m, 1 H) 3.62 - 3.96 (m, 4 H) 4.34 (dd, J=7.13, 1.85 Hz, 2 H) 4.56 - 4.68 (m, 1 H) 6.44 (d, J=2.34 Hz, 1 H) 6.50 - 6.61 (m, 1 H) 6.81 - 6.96 (m, 1 H) 7.26 (d, J=7.81 Hz, 1 H) 7.40 - 7.47 (m, 1 H) 7.50 (s, 1 H) 7.68 (d, J=1.37 Hz, 1 H) 7.78 (s, 1 H) 7.82 (d, J=1.37 Hz, 1 H) 7.98; (d, J=2.15 Hz, 1 H)
63be	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (m, 6 H) 1.59 (m, 4 H) 2.02 (m, 1 H) 2.38 (s, 3 H) 2.45 (dd, J=13.54, 8.76 Hz, 1 H) 2.72 (q, J=7.60 Hz, 2 H) 3.24 (m, 2 H) 3.58 (m, 4 H) 4.32 (m, 2 H) 4.53 (t, J=8.76 Hz, 1 H) 5.72 (s, 1 H) 6.38 (d, J=2.20 Hz, 1 H) 6.71 (m, 1 H) 7.25 (d, J=1.56 Hz, 1 H) 7.36 (dd, J=8.10, 1.61 Hz, 1 H) 7.63 (d, J=8.10 Hz, 1 H) 7.85 (d, J=2.29 Hz, 1 H)
63bf	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.96 (t, J=7.35 Hz, 2 H) 1.32 (t, J=7.15 Hz, 4 H) 1.63 (m, 6 H) 2.00 (dd, J=13.59, 8.61 Hz, 1 H) 2.37 (s, 3 H) 2.42 (m, 1 H) 2.66 (m, 2 H) 3.21 (m, 2 H) 3.58 (m, 4 H) 4.31 (m, 2 H) 4.49 (t, J=8.69 Hz, 1 H) 5.72 (s, 1 H) 6.38 (d, J=2.29 Hz, 1 H) 6.71(q, J=6.67 Hz, 1 H) 7.23 (d, J=1.66 Hz, 1 H) 7.34 (dd, J=8.10, 1.66 Hz, 1 H) 7.63 (d, J=8.10 Hz, 1 H) 7.85 (d, J=2.29 Hz, 1 H)
63bg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.95 (t, J=7.35 Hz, 3 H) 1.32 (t, J=7.15 Hz, 4 H) 1.63 (m, 6 H) 2.02 (m, 1 H) 2.38 (s, 3 H) 2.45 (dd, J=13.54, 8.76 Hz, 1 H) 2.69 (m, 2 H) 3.24 (m, 2 H) 3.58 (m, 4 H) 4.32 (m, 2 H) 4.53 (t, J=8.74 Hz, 1 H) 5.72 (s, 1 H) 6.38 (d, J=2.29 Hz, 1 H) 6.71 (m, 1 H) 7.23 (d, J=1.61 Hz, 1 H) 7.34 (dd, J=8.15, 1.61 Hz, 1 H) 7.62 (d, J=8.15 Hz,

	1 H) 7.85 (d, J=2.34 Hz, 1 H)
63bh	¹ H NMR (400 MHz, CHLOROFORM-d): δ ppm 1.18-1.36 (m, 3 H) 1.43 (t, J =6.74 Hz, 3 H) 1.54-2.29 (m, 6 H) 2.39 (br. s., 3 H) 3.78 (br. s., 4 H) 4.26 (br. s., 2 H) 4.42 (d J =6.15 Hz, 2 H) 5.53 (br. s., 1 H) 6.36 (s, 1 H) 6.59 (br. s., 1 H) 7.48 (d, J =7.96 Hz, 1 H) 7.61 (br. s., 1 H) 8.16 (d, J=8.05 Hz, 1 H) 8.34 (br. s., 1 H)
63bi	¹ H NMR (400 MHz, MeOH-d4): δ ppm 1.24 - 1.30 (m, 5 H) 1.37 (t, J=7.13 Hz, 3 H) 1.45 - 1.62 (m, 4 H) 1.84 (dd, J=13.32, 7.86 Hz, 1 H) 1.95 (s, 4 H) 2.22 (dd, J=13.30, 8.86 Hz, 1 H) 2.38 (s, 3 H) 2.88 - 3.09 (m, 2 H) 3.41 - 3.71 (m, 4 H) 4.10 (t, J=8.25 Hz, 1 H) 4.22 (qd, J=7.13, 2.00 Hz, 2 H) 4.37 (q, J=7.13 Hz, 2 H) 5.66 (s, 1 H) 6.41 (d, J=2.39 Hz, 1 H) 6.84 (q, J=6.54 Hz, 1 H) 7.83 (d, J=8.30 Hz, 1 H) 7.94 (d, J=2.34 Hz, 1 H) 7.99 (d, J=1.61 Hz, 1 H) 8.08 (dd, J=8.27, 1.64 Hz, 1 H)
63bj	¹ H NMR (400 MHz, MeOH-d4): δ ppm 1.31 (td, J=7.13, 3.22 Hz, 6 H) 1.52 - 1.64 (m, 4 H) 1.97 (s, 1 H) 2.01 (dd, J=13.59, 8.81 Hz, 1 H) 2.37 (s, 3 H) 2.44 (dd, J=13.62, 8.74 Hz, 1 H) 3.18 - 3.26 (m, 2 H) 3.43 - 3.68 (m, 4 H) 4.19 - 4.34 (m, 4 H) 4.53 (t, J=8.74 Hz, 1 H) 5.75 (s, 1 H) 6.40 (d, J=2.39 Hz, 1 H) 6.55 (d, J=16.06 Hz, 1 H) 6.95 (q, J=6.56 Hz, 1 H) 7.46 (d, J=8.30 Hz, 1 H) 7.68 (d, J=16.06 Hz, 1 H) 7.80 (dd, J=8.32, 2.03 Hz, 1 H) 7.87 (s, 1 H) 7.91 (d, J=2.39 Hz, 1 H)
63bk	¹ H NMR (400 MHz, MeOH-d4): δ ppm 0.92 (t, J=7.37 Hz, 3 H) 1.32 (dq, J=14.94, 7.38 Hz, 2 H) 1.50 - 1.68 (m, 6 H) 2.06 (dd, J=13.37, 7.22 Hz, 1 H) 2.31 (dd, J=13.45, 9.25 Hz, 1 H) 2.37 (s, 3 H) 2.69 (t, J=7.59 Hz, 2 H) 3.06 - 3.29 (m, 2 H) 3.41 - 3.76 (m, 4 H) 4.08 (dd, J=9.20, 7.25 Hz, 1 H) 5.75 (s, 1 H) 6.36 (d, J=2.15 Hz, 1 H) 6.69 (q, J=6.62 Hz, 1 H) 7.28 - 7.33 (m, 1 H) 7.34 - 7.39 (m, 1 H) 7.53 (s, 1 H) 7.82 (d, J=2.29 Hz, 1 H)
63bl	¹ H NMR (400 MHz, DMSO-d6): δ ppm 1.50 - 1.73 (m, 4 H) 1.80 (quin, J=7.52 Hz, 2 H) 1.90 (dd, J=13.23, 9.22 Hz, 1 H) 2.15 - 2.26 (m, 2 H) 2.27 - 2.41 (m, 4 H) 2.69 (t, J=7.66 Hz, 2 H) 3.00 - 3.20 (m, 2 H) 3.69 (br. s., 4 H) 4.33 - 4.52 (m, 1 H) 6.14 (br. s., 1 H) 6.38 (d, J=2.29 Hz, 1 H) 7.05 (br. s., 1 H) 7.37 - 7.52 (m, 3 H) 7.76 (br. s., 1 H) 8.02 (d, J=2.29 Hz, 1 H) 8.97 (d, J=5.32 Hz, 1 H) 10.42 (br. s., 1 H)
63bm	¹ H NMR (400 MHz, MeOH-d4): δ ppm 1.31 (t, J=7.15 Hz, 3 H) 1.52 - 1.70 (m, 4 H) 1.90 (dd, J=6.30, 1.22 Hz, 3 H) 1.97 (dd, J=13.52, 8.44 Hz, 1 H) 2.35 - 2.41 (m, 4 H) 3.06 - 3.24 (m, 2 H) 3.42 - 3.79 (m, 4 H) 4.21 - 4.35 (m, 2 H) 4.40 (t, J=8.57 Hz, 1 H) 5.75 (s, 1 H) 6.27 - 6.54 (m, 3 H) 6.75 (q, J=6.64 Hz, 1 H) 7.32 (d, J=8.25 Hz, 1 H) 7.52 (dd, J=8.30, 2.00 Hz, 1 H) 7.64 (s, 1 H) 7.83 (d, J=2.29 Hz, 1 H)
63bn	¹ H NMR (400 MHz, MeOH-d4): δ ppm 1.32 (t, J=7.15 Hz, 3 H) 1.49 - 1.70 (m, 4 H) 2.01 (dd, J=13.59, 8.76 Hz, 1 H) 2.29 (s, 3 H) 2.32 (s, 3 H) 2.40 (s, 3 H) 2.40 - 2.44 (m, 1 H) 3.24 (s, 2 H) 3.43 - 3.71 (m, 4 H) 4.22 - 4.41 (m, 2 H) 4.56 (t, J=8.74 Hz, 1 H) 5.80 (s, 1 H) 6.41 (d, J=2.29 Hz, 1 H) 6.81 - 6.92 (m, 1 H) 7.20 (d, J=7.81 Hz, 1 H) 7.26 - 7.32 (m, 1 H) 7.35 (s, 1 H) 7.45 (d, J=8.30 Hz, 1 H) 7.73 (dd, J=8.27, 2.12 Hz, 1 H) 7.88 - 7.90 (m, 2 H)
63bo	¹ H NMR (400 MHz, MeOH-d4): δ ppm 0.92 (t, J=7.35 Hz, 3 H) 1.32 (t, J=7.13 Hz, 3 H) 1.49 - 1.76 (m, 6 H) 1.94 - 2.06 (m, 1 H) 2.37 (s, 3 H) 2.43 (dd, J=13.57, 8.79 Hz, 1 H) 2.66 (t, J=7.52 Hz, 2 H) 3.13 - 3.28 (m, 2 H) 3.43 - 3.76 (m, 4 H) 4.21 - 4.39 (m, 2 H) 4.50 (t, J=8.66 Hz, 1 H) 5.74 (s, 1 H) 6.37 (d, J=2.29 Hz, 1 H) 6.70 (q, J=6.69 Hz, 1 H) 7.26 - 7.33 (m, 1 H) 7.34 - 7.42 (m, 1 H) 7.53 (s, 1 H) 7.82 (d, J=2.29 Hz, 1 H)
63bp	¹ H NMR (400 MHz, MeOH-d4): δ ppm 1.25 (t, J=7.61 Hz, 3 H) 1.33 (t, J=7.15 Hz, 3

	H) 1.57 - 1.71 (m, 4 H) 2.04 (dd, J=13.93, 8.52 Hz, 1 H) 2.37 (s, 3 H) 2.47 (dd, J=13.62, 8.74 Hz, 1 H) 2.72 (q, J=7.61 Hz, 2 H) 3.26 (d, J=1.51 Hz, 2 H) 3.44 - 3.77 (m, 4 H) 4.23 - 4.43 (m, 2 H) 4.57 (t, J=8.79 Hz, 1 H) 5.76 (s, 1 H) 6.37 (d, J=2.20 Hz, 1 H) 6.63 - 6.78 (m, 1 H) 7.27 - 7.35 (m, 1 H) 7.36 - 7.46 (m, 1 H) 7.55 (s, 1 H) 7.81 (d, J=2.29 Hz, 1 H)
63bq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.92 (t, J=7.35 Hz, 3 H) 1.22 - 1.42 (m, 5 H) 1.49 - 1.75 (m, 6 H) 1.94 - 2.08 (m, 1 H) 2.37 (s, 3 H) 2.44 (dd, J=13.57, 8.74 Hz, 1 H) 2.68 (t, J=7.61 Hz, 2 H) 3.15 - 3.29 (m, 2 H) 3.42 - 3.76 (m, 4 H) 4.23 - 4.40 (m, 2 H) 4.53 (t, J=8.74 Hz, 1 H) 5.75 (s, 1 H) 6.37 (d, J=2.34 Hz, 1 H) 6.70 (q, J=6.69 Hz, 1 H) 7.27 - 7.33 (m, 1 H) 7.34 - 7.41 (m, 1 H) 7.53 (s, 1 H) 7.82 (d, J=2.34 Hz, 1 H)
63br	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.33 (t, J=7.13 Hz, 3 H) 1.53 - 1.74 (m, 4 H) 2.05 (dd, J=13.62, 8.83 Hz, 1 H) 2.38 (s, 3 H) 2.48 (dd, J=13.62, 8.79 Hz, 1 H) 3.28 (s, 2 H) 3.44 - 3.79 (m, 4 H) 4.22 - 4.43 (m, 2 H) 4.59 (t, J=8.79 Hz, 1 H) 5.37 (d, J=11.08 Hz, 1 H) 5.73 - 5.96 (m, 2 H) 6.39 (d, J=2.34 Hz, 1 H) 6.68 - 6.95 (m, 2 H) 7.40 (d, J=8.25 Hz, 1 H) 7.65 (dd, J=8.27, 1.98 Hz, 1 H) 7.73 (s, 1 H) 7.87 (d, J=2.34 Hz, 1 H)
63bs	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.11 (t, J=7.47 Hz, 3 H) 1.33 (t, J=7.15 Hz, 3 H) 1.51 - 1.72 (m, 4 H) 2.04 (dd, J=13.62, 8.83 Hz, 1 H) 2.18 - 2.33 (m, 2 H) 2.38 (s, 3 H) 2.47 (dd, J=13.59, 8.81 Hz, 1 H) 3.26 (s, 2 H) 3.44 - 3.78 (m, 4 H) 4.19 - 4.43 (m, 2 H) 4.58 (t, J=8.79 Hz, 1 H) 5.79 (s, 1 H) 6.30 - 6.53 (m, 3 H) 6.69 - 6.84 (m, 1 H) 7.33 (d, J=8.25 Hz, 1 H) 7.55 (dd, J=8.30, 2.00 Hz, 1 H) 7.65 (s, 1 H) 7.84 (d, J=2.34 Hz, 1 H)
63bt	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (t, J=7.13 Hz, 3 H) 1.45 - 1.65 (m, 4 H) 1.79 - 1.91 (m, 1 H) 2.16 - 2.32 (m, 1 H) 3.03 (s, 2 H) 3.51 (br. s., 4 H) 3.75 - 3.81 (m, 1 H) 4.07 - 4.17 (m, 1 H) 4.20 - 4.32 (m, 2 H) 5.55 (s, 1 H) 6.65 (d, J=2.34 Hz, 1 H) 7.15 - 7.28 (m, 1 H) 7.36 - 7.46 (m, 1 H) 7.57 (d, J=2.15 Hz, 1 H) 7.62 - 7.70 (m, 1 H) 7.73 (d, J=2.15 Hz, 1 H)
63bu	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (t, J=7.13 Hz, 3 H) 1.40 - 1.61 (m, 4 H) 1.71 - 1.86 (m, 1 H) 2.07 - 2.22 (m, 1 H) 2.86 (s, 1 H) 2.94 (s, 1 H) 3.50 (d, J=4.69 Hz, 4 H) 4.00 (s, 4 H) 4.22 (dd, J=7.22, 0.98 Hz, 2 H) 5.57 (s, 1 H) 6.61 (d, J=2.15 Hz, 1 H) 7.13 - 7.28 (m, 1 H) 7.35 - 7.50 (m, 2 H) 7.55 (d, J=1.17 Hz, 1 H) 7.72 (d, J=2.34 Hz, 2 H)
63bv	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (t, J=7.13 Hz, 3 H) 1.57 - 1.83 (m, 4 H) 1.99 - 2.16 (m, 1 H) 2.56 (s, 4 H) 3.31 (s, 2 H) 3.68 (br. s., 4 H) 4.03 (s, 3 H) 4.35 (dd, J=7.03, 2.15 Hz, 2 H) 4.62 (s, 1 H) 5.70 (s, 1 H) 6.70 (d, J=6.83 Hz, 1 H) 7.42 (dd, J=8.49, 0.88 Hz, 1 H) 7.60 - 7.72 (m, 3 H) 7.73 - 7.86 (m, 3 H)
63bw	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (t, J=7.13 Hz, 3 H) 1.67 - 1.90 (m, 4 H) 2.03 - 2.18 (m, 1 H) 2.47 - 2.61 (m, 1 H) 2.72 (s, 3 H) 3.34 (br. s., 2 H) 3.56 - 3.87 (m, 4 H) 4.17 (s, 3 H) 4.35 (dd, J=7.13, 2.05 Hz, 2 H) 4.64 (s, 1 H) 5.89 - 6.04 (m, 1 H) 6.59 - 6.75 (m, 1 H) 7.45 (d, J=0.98 Hz, 1 H) 7.70 (d, J=8.20 Hz, 2 H) 7.77 (s, 1 H) 7.79 - 7.92 (m, 3 H)
63bx	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (t, J=7.13 Hz, 3 H) 1.77 (br. s., 4 H) 2.04 - 2.15 (m, 1 H) 2.47 - 2.58 (m, 1 H) 3.09 (s, 2 H) 3.57 (t, J=6.74 Hz, 6 H) 4.28 - 4.43 (m, 2 H) 4.57 - 4.69 (m, 1 H) 5.80 - 5.92 (m, 1 H) 6.60 - 6.75 (m, 1 H) 7.62 (s, 1 H) 7.69 (d, J=8.40 Hz, 3 H) 7.74 - 7.85 (m, 2 H) 8.03 (d, J=8.00 Hz, 1 H)
63by	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (t, J=7.13 Hz, 3 H) 1.61 - 1.81 (m, 4 H)

	2.00 - 2.16 (m, 1 H) 2.44 - 2.59 (m, 1 H) 3.47 - 3.80 (m, 4 H) 4.35 (dd, J=7.03, 2.54 Hz, 2 H) 4.63 (s, 1 H) 5.73 (s, 1 H) 6.64 - 6.83 (m, 1 H) 7.76 (d, J=8.20 Hz, 2 H) 7.95 (d, J=8.20 Hz, 2 H) 8.12 - 8.33 (m, 2 H) 8.36 - 8.47 (m, 2 H) 8.48 - 8.68 (m, 1 H) 9.39 - 9.76 (m, 1 H)
63bz	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.10 - 1.20 (m, 3 H) 1.26 (t, J=7.13 Hz, 3 H) 1.42 - 1.64 (m, 4 H) 1.79 (dd, J=13.15, 7.44 Hz, 1 H) 1.94 (s, 2 H) 2.15 (dd, J=12.98, 8.69 Hz, 1 H) 2.35 (s, 3 H) 2.62 - 2.71 (m, 2 H) 2.81 - 2.87 (m, 1 H) 2.93 - 3.02 (m, 3 H) 3.40 - 3.66 (m, 4 H) 3.96 (t, J=8.18 Hz, 1 H) 4.06 (q, J=7.18 Hz, 2 H) 4.16 - 4.25 (m, 2 H) 5.69 (s, 1 H) 6.36 (d, J=2.25 Hz, 1 H) 6.71 (q, J=6.67 Hz, 1 H) 7.27 (d, J=1.56 Hz, 1 H) 7.35 (dd, J=8.13, 1.83 Hz, 1 H) 7.62 (d, J=8.20 Hz, 1 H) 7.83 (d, J=2.29 Hz, 1 H)
63ca	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (t, J=7.13 Hz, 3 H) 1.63 - 1.82 (m, 4 H) 2.03 - 2.17 (m, 1 H) 2.47 - 2.60 (m, 1 H) 3.51 - 3.83 (m, 4 H) 4.27 - 4.42 (m, 2 H) 4.57 - 4.69 (m, 1 H) 5.69 - 5.88 (m, 1 H) 6.65 - 6.85 (m, 1 H) 7.71 - 7.85 (m, 2 H) 7.89 - 8.01 (m, 2 H) 8.22 - 8.35 (m, 2 H) 8.36 - 8.49 (m, 1 H) 8.52 - 8.60 (m, 1 H) 8.62 - 8.72 (m, 1 H) 9.56 - 9.75 (m, 1 H)
63cb	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.18 (t, J=7.10 Hz, 3 H) 1.24 (t, J=7.13 Hz, 3 H) 1.37 - 1.63 (m, 4 H) 1.82 (quin, J=7.52 Hz, 2 H) 1.90 (dd, J=13.28, 9.42 Hz, 1 H) 2.19 - 2.41 (m, 6 H) 2.67 (t, J=7.69 Hz, 2 H) 3.12 (br. s., 2 H) 3.18 - 3.74 (m, 4 H) 4.05 (q, J=7.13 Hz, 2 H) 4.15 - 4.30 (m, 2 H) 4.52 (t, J=8.49 Hz, 1 H) 5.72 (br. s., 1 H) 6.01 (br. s., 2 H) 6.37 (d, J=2.15 Hz, 1 H) 6.99 (q, J=6.87 Hz, 1 H) 7.33 - 7.44 (m, 2 H) 7.47 (s, 1 H) 8.01 (d, J=2.25 Hz, 1 H) 9.20 (br. s., 1 H) 10.39 (br. s., 1 H)
63cc	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.10 - 1.20 (m, 3 H) 1.25 (t, J=7.10 Hz, 3 H) 1.44 - 1.63 (m, 4 H) 1.82 (quin, J=7.53 Hz, 2 H) 1.91 (dd, J=13.28, 9.37 Hz, 1 H) 2.19 - 2.40 (m, 3 H) 2.60 (t, J=7.71 Hz, 2 H) 3.13 (br. s., 2 H) 3.40 - 3.68 (m, 4 H) 4.02 (q, J=7.09 Hz, 2 H) 4.13 - 4.33 (m, 2 H) 4.53 (br. s., 1 H) 5.70 (br. s., 1 H) 6.29 (br. s., 2 H) 6.62 - 6.76 (m, 1 H) 7.28 (d, J=8.20 Hz, 2 H) 7.43 (d, J=8.10 Hz, 2 H) 9.21 (br. s., 1 H) 10.43 (br. s., 1 H)
63cd	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.15 (t, J=7.13 Hz, 3 H) 1.24 (t, J=7.10 Hz, 3 H) 1.39 - 1.64 (m, 4 H) 1.78 - 1.97 (m, 3 H) 2.22 - 2.39 (m, 6 H) 2.66 (t, J=7.71 Hz, 2 H) 3.11 (br. s., 2 H) 3.38 - 3.64 (m, 4 H) 3.93 - 4.07 (m, 2 H) 4.15 - 4.31 (m, 2 H) 4.52 (br. s., 1 H) 5.73 (br. s., 1 H) 6.05 (br. s., 2 H) 6.38 (d, J=2.10 Hz, 1 H) 7.00 (q, J=6.72 Hz, 1 H) 7.30 (d, J=1.51 Hz, 1 H) 7.33 - 7.41 (m, 1 H) 7.59 (d, J=8.05 Hz, 1 H) 8.04 (d, J=2.29 Hz, 1 H) 9.20 (br. s., 1 H) 10.38 (br. s., 1 H)
63ce	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.22 - 1.30 (m, 3 H) 1.48 - 1.61 (m, 4 H) 1.82 (dd, J=13.30, 7.74 Hz, 1 H) 1.94 (s, 3 H) 2.15 - 2.23 (m, 1 H) 2.38 (s, 3 H) 2.87 - 2.92 (m, 1 H) 2.96 - 3.02 (m, 1 H) 3.42 - 3.64 (m, 4 H) 4.01 - 4.08 (m, 1 H) 4.16 - 4.25 (m, 2 H) 5.72 (s, 1 H) 6.41 (d, J=2.39 Hz, 1 H) 6.81 - 6.88 (m, 1 H) 7.61 - 7.66 (m, 1 H) 7.71 (d, J=1.76 Hz, 1 H) 7.73 - 7.86 (m, 3 H) 7.97 - 8.02 (m, 2 H) 8.08 (s, 1 H)
63cf	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.25 (t, J=7.13 Hz, 3 H) 1.45 - 1.66 (m, 4 H) 1.92 (dd, J=13.18, 9.42 Hz, 1 H) 2.35 (dd, J=13.28, 8.54 Hz, 1 H) 3.14 (br. s., 2 H) 3.60 (br. s., 4 H) 4.14 - 4.31 (m, 2 H) 4.54 (br. s., 1 H) 5.75 (br. s., 1 H) 6.56 (q, J=6.72 Hz, 1 H) 7.47 (t, J=1.27 Hz, 1 H) 7.65 (s, 2 H) 7.75 - 7.84 (m, 1 H) 7.89 (d, J=7.81 Hz, 1 H) 7.93 - 8.01 (m, 2 H) 9.21 (br. s., 1 H) 10.36 (br. s., 1 H)
63cg	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.25 (t, J=7.10 Hz, 3 H) 1.57 (d, J=5.37 Hz, 4 H) 1.83 - 1.99 (m, 1 H) 2.28 - 2.40 (m, 1 H) 3.14 (br. s., 2 H) 3.58 (br. s., 4 H) 3.81 (s,

	3 H) 4.24 (dd, J=7.13, 2.25 Hz, 2 H) 4.43 - 4.63 (m, 1 H) 5.62 - 5.85 (m, 1 H) 6.73 (d, J=6.78 Hz, 1 H) 6.96 - 7.16 (m, 3 H) 7.39 (d, J=2.15 Hz, 1 H) 7.50 (dd, J=8.74, 7.61 Hz, 1 H) 7.55 - 7.69 (m, 2 H) 9.09 - 9.32 (m, 1 H) 10.26 - 10.47 (m, 1 H)
63ch	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (t, J=7.15 Hz, 3 H) 1.52 - 1.70 (m, 4 H) 1.93 - 2.02 (m, 1 H) 2.40 (dd, J=13.45, 8.71 Hz, 1 H) 3.09 - 3.24 (m, 2 H) 3.43 - 3.74 (m, 4 H) 4.25 - 4.35 (m, 2 H) 4.40 (t, J=8.57 Hz, 1 H) 5.59 (s, 1 H) 6.61 (q, J=6.56 Hz, 1 H) 7.32 (d, J=2.15 Hz, 1 H) 7.49 (dd, J=8.49, 2.25 Hz, 1 H) 7.61 (d, J=8.00 Hz, 1 H) 7.65 - 7.77 (m, 2 H) 7.97 - 8.10 (m, 1 H) 8.32 (br. s., 1 H)
63ci	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (t, J=7.13 Hz, 3 H) 1.50 - 1.70 (m, 4 H) 1.91 - 2.02 (m, 1 H) 2.39 (dd, J=13.50, 8.76 Hz, 1 H) 3.08 - 3.23 (m, 2 H) 3.41 - 3.69 (m, 4 H) 4.24 - 4.34 (m, 2 H) 4.38 (t, J=8.54 Hz, 1 H) 5.50 (s, 1 H) 6.75 (q, J=6.96 Hz, 1 H) 6.87 (d, J=7.66 Hz, 1 H) 6.91 (ddd, J=8.21, 2.50, 0.90 Hz, 1 H) 7.06 (br. s., 1 H) 7.28 (d, J=2.20 Hz, 1 H) 7.32 (t, J=7.88 Hz, 1 H) 7.43 (dd, J=8.47, 2.27 Hz, 1 H) 7.66 (d, J=8.44 Hz, 1 H)
63cj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (t, J=7.13 Hz, 3 H) 1.52 - 1.70 (m, 4 H) 1.93 - 2.06 (m, 1 H) 2.41 (dd, J=13.42, 8.74 Hz, 1 H) 3.10 - 3.25 (m, 5 H) 3.44 - 3.74 (m, 4 H) 4.21 - 4.37 (m, 2 H) 4.42 (t, J=8.59 Hz, 1 H) 5.61 (s, 1 H) 6.57 (q, J=6.57 Hz, 1 H) 7.36 (d, J=2.20 Hz, 1 H) 7.51 (dd, J=8.54, 2.20 Hz, 1 H) 7.70 (d, J=8.44 Hz, 1 H) 7.72 - 7.78 (m, 1 H) 7.78 - 7.89 (m, 1 H) 8.09 (dt, J=7.85, 1.49 Hz, 1 H) 8.41 (d, J=0.73 Hz, 1 H)
63ck	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.21 (t, J=7.10 Hz, 3 H) 1.38 - 1.64 (m, 4 H) 1.88 (dd, J=13.20, 9.35 Hz, 1 H) 2.30 (dd, J=13.20, 8.47 Hz, 1 H) 3.09 (br. s., 2 H) 3.42 - 3.61 (m, 4 H) 3.95 - 4.11 (m, 2 H) 4.12 - 4.28 (m, 2 H) 4.48 (br. s., 1 H) 5.71 (br. s., 1 H) 6.32 (br. s., 1 H) 6.71 (q, J=6.74 Hz, 1 H) 7.33 (d, J=2.05 Hz, 1 H) 7.44 - 7.69 (m, 6 H) 8.52 (br. s., 3 H) 9.27 (br. s., 1 H) 10.62 (br. s., 1 H)
63cl	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (t, J=7.13 Hz, 4 H) 1.82 (br. s., 4 H) 2.03 - 2.21 (m, 1 H) 2.47 - 2.64 (m, 1 H) 3.35 (s, 2 H) 3.56 - 3.92 (m, 4 H) 4.27 - 4.43 (m, 2 H) 4.59 - 4.70 (m, 1 H) 6.65 - 6.82 (m, 1 H) 7.81 (d, J=8.00 Hz, 2 H) 8.00 (d, J=8.20 Hz, 2 H) 8.05 - 8.14 (m, 1 H) 8.29 - 8.40 (m, 1 H) 8.46 - 8.55 (m, 1 H) 8.63 (d, J=1.56 Hz, 1 H) 9.21 (s, 2 H)
63cm	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (t, J=6.93 Hz, 5 H) 1.83 (br. s., 4 H) 2.04 - 2.22 (m, 1 H) 2.47 - 2.65 (m, 1 H) 3.36 (br. s., 2 H) 4.35 (d, J=6.64 Hz, 2 H) 4.57 - 4.71 (m, 1 H) 6.64 - 6.85 (m, 1 H) 7.84 (d, J=6.64 Hz, 2 H) 8.03 (d, J=6.83 Hz, 2 H) 8.08 - 8.18 (m, 1 H) 8.27 - 8.41 (m, 1 H) 8.50 (br. s., 2 H) 9.26 (br. s., 2 H)
63cn	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.24 - 1.45 (m, 10 H) 1.75 (d, J=18.55 Hz, 4 H) 2.01 - 2.18 (m, 1 H) 2.43 (s, 3 H) 2.47 - 2.62 (m, 1 H) 3.86 (br. s., 3 H) 4.34 (d, J=5.86 Hz, 2 H) 4.54 - 4.75 (m, 2 H) 6.44 (d, J=1.95 Hz, 1 H) 6.89 (d, J=5.66 Hz, 1 H) 7.03 (d, J=8.59 Hz, 2 H) 7.57 - 7.71 (m, 3 H) 7.72 - 7.87 (m, 2 H) 7.98 (d, J=1.76 Hz, 1 H)
63co	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (s, 3 H) 1.64 - 1.91 (m, 4 H) 2.03 - 2.20 (m, 1 H) 2.47 - 2.64 (m, 1 H) 3.35 (br. s., 2 H) 3.56 - 3.95 (m, 4 H) 4.25 - 4.44 (m, 2 H) 4.57 - 4.71 (m, 1 H) 6.57 - 6.84 (m, 1 H) 7.70 - 7.85 (m, 2 H) 7.90 - 8.07 (m, 2 H) 8.23 (s, 2 H) 8.33 - 8.47 (m, 1 H) 8.86 - 9.05 (m, 2 H)
63cp	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.18 - 1.31 (m, 3 H) 1.44 - 1.60 (m, 4 H) 1.79 (dd, J=13.28, 7.61 Hz, 1 H) 1.93 (s, 2 H) 1.98 (s, 3 H) 2.14 (dd, J=13.18, 8.79 Hz, 1 H) 2.37 (s, 3 H) 2.80 - 2.89 (m, 1 H) 2.91 - 3.00 (m, 1 H) 3.40 - 3.64 (m, 4 H) 3.97 (t,

	J=8.15 Hz, 1 H) 4.19 (qd, J=7.13, 1.81 Hz, 2 H) 4.37 (s, 2 H) 5.72 (s, 1 H) 6.39 (d, J=2.20 Hz, 1 H) 6.76 (q, J=6.56 Hz, 1 H) 7.36 (d, J=8.30 Hz, 2 H) 7.59 - 7.66 (m, 3 H) 7.69 - 7.81 (m, 2 H) 7.95 (d, J=2.29 Hz, 1 H)
63cq	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.10 - 1.20 (m, 3 H) 1.25 (t, J=7.10 Hz, 3 H) 1.44 - 1.63 (m, 4 H) 1H NMR (400 MHz, MeOH-d ₄): δ ppm 1.21 - 1.31 (m, 3 H) 1.52 (dt, J=10.53, 5.35 Hz, 4 H) 1.79 (dd, J=13.15, 7.44 Hz, 1 H) 1.89 (s, 3 H) 1.93 (s, 2 H) 2.14 (dd, J=13.13, 8.79 Hz, 1 H) 2.38 (s, 3 H) 2.78 - 2.88 (m, 3 H) 2.91 - 2.99 (m, 1 H) 3.40 (t, J=7.35 Hz, 2 H) 3.44 - 3.66 (m, 4 H) 3.95 (t, J=8.13 Hz, 1 H) 4.12 - 4.25 (m, 2 H) 5.72 (s, 1 H) 6.40 (d, J=2.29 Hz, 1 H) 6.77 (q, J=6.74 Hz, 1 H) 7.31 (d, J=8.25 Hz, 2 H) 7.58 - 7.64 (m, 3 H) 7.70 - 7.80 (m, 2 H) 7.95 (d, J=2.29 Hz, 1 H)
63cr	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.25 (t, J=7.13 Hz, 3 H) 1.47 - 1.58 (m, 4 H) 1.79 (dd, J=13.20, 7.44 Hz, 1 H) 1.93 (s, 2 H) 2.15 (dd, J=13.23, 8.74 Hz, 1 H) 2.40 (s, 3 H) 2.81 - 2.87 (m, 1 H) 2.92 - 2.98 (m, 1 H) 3.44 - 3.63 (m, 4 H) 3.97 (dd, J=8.52, 7.83 Hz, 1 H) 4.14 - 4.24 (m, 2 H) 5.75 (s, 1 H) 6.42 (d, J=2.29 Hz, 1 H) 6.84 (q, J=6.69 Hz, 1 H) 7.55 (dd, J=8.30, 4.34 Hz, 1 H) 7.83 (d, J=1.76 Hz, 1 H) 7.85 - 7.99 (m, 3 H) 8.00 - 8.07 (m, 2 H) 8.30 (d, J=1.51 Hz, 1 H) 8.37 - 8.42 (m, 1 H) 8.88 (dd, J=4.30, 1.66 Hz, 1 H)
63cs	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.34 (d, J=6.05 Hz, 6 H) 1.60 (br. s., 4 H) 2.02 - 2.13 (m, 1 H) 2.26 - 2.37 (m, 1 H) 2.42 (s, 3 H) 3.04 - 3.18 (m, 1 H) 3.26 (d, J=11.71 Hz, 1 H) 3.41 - 3.78 (m, 4 H) 4.02 - 4.17 (m, 1 H) 4.66 (s, 1 H) 5.78 (s, 1 H) 6.43 (d, J=2.15 Hz, 1 H) 6.69 - 6.86 (m, 1 H) 6.99 (d, J=8.79 Hz, 2 H) 7.50 - 7.66 (m, 3 H) 7.67 - 7.82 (m, 2 H) 7.97 (d, J=2.34 Hz, 1 H)
63ct	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (t, J=7.13 Hz, 3 H) 1.58 - 1.78 (m, 4 H) 2.02 - 2.16 (m, 1 H) 2.43 - 2.60 (m, 1 H) 3.30 (s, 2 H) 3.46 - 3.78 (m, 4 H) 4.27 - 4.41 (m, 2 H) 4.55 - 4.66 (m, 1 H), 5.61 - 5.77 (m, 1 H) 6.42 - 6.53 (m, 1 H) 6.59 - 6.70 (m, 1 H) 7.25 - 7.39 (m, 2 H) 7.62 (s, 4 H) 7.72 (s, 2 H)
63cu	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.25 (t, J=7.10 Hz, 3 H) 1.33 (t, J=7.10 Hz, 3 H) 1.44 - 1.64 (m, 4 H) 1.92 (dd, J=13.28, 9.27 Hz, 1 H) 2.35 (dd, J=13.28, 8.49 Hz, 1 H) 3.14 (br. s., 2 H) 3.44 - 3.66 (m, 4 H) 4.14 - 4.29 (m, 2 H) 4.30 - 4.43 (m, 2 H) 4.54 (br. s., 1 H) 5.75 (br. s., 1 H) 6.43 (br. s., 1 H) 6.59 (q, J=6.72 Hz, 1 H) 7.37 - 7.47 (m, 1 H) 7.57 - 7.67 (m, 2 H) 7.68 - 7.81 (m, 2 H) 8.08 (dt, J=6.77, 1.96 Hz, 1 H) 8.24 (br. s., 1 H) 9.22 (br. s., 1 H) 10.41 (br. s., 1 H)
63cv	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.34 (t, J=7.13 Hz, 3 H) 1.69 - 1.88 (m, 4 H) 2.11 (dd, J=13.64, 8.96 Hz, 1 H) 2.55 (dd, J=13.62, 8.69 Hz, 1 H) 3.34 (s, 2 H) 3.52 - 3.80 (m, 4 H) 4.35 (qd, J=7.13, 1.93 Hz, 2 H) 4.63 (t, J=8.79 Hz, 1 H) 6.55 - 6.67 (m, 1 H) 7.37 (d, J=2.20 Hz, 1 H) 7.54 (dd, J=8.52, 2.22 Hz, 1 H) 7.58 - 7.63 (m, 1 H) 7.65 - 7.70 (m, 1 H) 7.71 (d, J=8.59 Hz, 1 H) 8.17 (dt, J=7.74, 1.43 Hz, 1 H) 8.35 (br. s., 1 H)
63cw	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.44 - 1.69 (m, 4 H) 1.91 (dd, J=13.28, 9.18 Hz, 1 H) 2.35 (dd, J=13.15, 8.61 Hz, 1 H) 3.14 (br. s., 2 H) 3.64 (br. s., 4 H) 4.37 - 4.53 (m, 1 H) 5.87 (br. s., 1 H) 6.62 (q, J=6.78 Hz, 1 H) 7.43 (t, J=1.22 Hz, 1 H) 7.65 (s, 2 H) 7.70 (d, J=4.78 Hz, 2 H) 7.99 - 8.12 (m, 1 H) 8.26 (br. s., 1 H) 8.96 (d, J=5.03 Hz, 1 H) 10.25 (br. s., 1 H)
63cx	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.33 (t, J=7.15 Hz, 3 H) 1.55 - 1.74 (m, 4 H) 2.04 (dd, J=13.57, 8.79 Hz, 1 H) 2.48 (dd, J=13.54, 8.76 Hz, 1 H) 3.26 (s, 2 H) 3.44 - 3.73 (m, 4 H) 4.25 - 4.41 (m, 2 H) 4.56 (t, J=8.74 Hz, 1 H) 5.57 (s, 1 H) 6.63 (q,

	J=6.80 Hz, 1 H) 7.30 (d, J=2.20 Hz, 1 H) 7.47 (dd, J=8.52, 2.22 Hz, 1 H) 7.52 - 7.59 (m, 1 H) 7.59 - 7.65 (m, 1 H) 7.67 (d, J=8.54 Hz, 1 H) 7.90 - 8.04 (m, 1 H) 8.41 (br. s., 1 H)
63cy	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.33 (t, J=7.15 Hz, 3 H) 1.53 - 1.74 (m, 4 H) 2.05 (dd, J=13.57, 8.79 Hz, 1 H) 2.48 (dd, J=13.54, 8.76 Hz, 1 H) 3.23 (s, 3 H) 3.27 (d, J=1.22 Hz, 2 H) 3.42 - 3.79 (m, 4 H) 4.22 - 4.42 (m, 2 H) 4.57 (t, J=8.79 Hz, 1 H) 5.54 (s, 1 H) 6.61 (q, J=6.72 Hz, 1 H) 7.35 (d, J=2.20 Hz, 1 H) 7.52 (dd, J=8.54, 2.25 Hz, 1 H) 7.72 (d, J=8.54 Hz, 1 H) 7.77 (d, J=7.86 Hz, 2 H) 8.08 - 8.20 (m, 2 H)
63cz	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.33 (t, J=7.13 Hz, 3 H) 1.54 - 1.72 (m, 4 H) 1.99 - 2.07 (m, 1 H) 2.46 (dd, J=13.57, 8.74 Hz, 1 H) 3.17 - 3.28 (m, 2 H) 3.42 - 3.72 (m, 4 H) 4.26 - 4.39 (m, 2 H) 4.51 (t, J=8.69 Hz, 1 H) 5.53 (s, 1 H) 6.56 - 6.66 (m, 1 H) 7.34 (d, J=2.20 Hz, 1 H) 7.51 (dd, J=8.52, 2.22 Hz, 1 H) 7.67 (d, J=8.00 Hz, 2 H) 7.71 (d, J=8.49 Hz, 1 H) 8.02 - 8.14 (m, 2 H)
63da	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.99 (t, J=7.15 Hz, 3 H) 1.33 (t, J=7.13 Hz, 3 H) 1.62 - 1.78 (m, 4 H) 2.07 (dd, J=13.64, 8.86 Hz, 1 H) 2.41 (s, 3 H) 2.50 (dd, J=13.67, 8.79 Hz, 1 H) 3.29 - 3.31 (m, 2 H) 3.55 - 3.84 (m, 4 H) 3.84 - 4.06 (m, 2 H) 4.23 - 4.42 (m, 2 H) 4.60 (t, J=8.81 Hz, 1 H) 6.23 - 6.36 (m, 1 H) 6.43 (d, J=2.34 Hz, 1 H) 6.89 - 7.01 (m, 1 H) 7.40 (dd, J=7.66, 0.93 Hz, 1 H) 7.48- 7.59 (m, 4 H) 7.60 - 7.69 (m, 1 H) 7.86 (dd, J=7.79, 1.24 Hz, 1 H) 7.93 (d, J=2.34 Hz, 1 H)
63db	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (t, J=7.15 Hz, 3 H) 1.40 (t, J=7.13 Hz, 3 H) 1.52 - 1.67 (m, 4 H) 2.00 (dd, J=13.59, 8.81 Hz, 1 H) 2.39 (s, 3 H) 2.44 (dd, J=13.64, 8.76 Hz, 1 H) 3.17 - 3.27 (m, 2 H) 3.41 - 3.73 (m, 4 H) 4.23 - 4.36 (m, 2 H) 4.40 (q, J=7.13 Hz, 2 H) 4.54 (t, J=8.79 Hz, 1 H) 5.78 (s, 1 H) 6.41 (d, J=2.15 Hz, 1 H) 6.92 (q, J=6.62 Hz, 1 H) 7.52 (d, J=8.25 Hz, 1 H) 7.58 (t, J=7.74 Hz, 1 H) 7.77 - 7.87 (m, 2 H) 7.88 - 7.97 (m, 2 H) 8.03 (dt, J=7.79, 1.33 Hz, 1 H) 8.21 (t, J=1.61 Hz, 1 H)
63dc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (t, J=7.13 Hz, 3 H) 1.41 (t, J=7.13 Hz, 3 H) 1.56 - 1.69 (m, 4 H) 2.03 (dd, J=13.62, 8.83 Hz, 1 H) 2.41 (s, 3 H) 2.47 (dd, J=13.57, 8.79 Hz, 1 H) 3.26 (s, 2 H) 3.46 - 3.75 (m, 4 H) 4.32 (qd, J=7.15, 2.37 Hz, 2 H) 4.40 (q, J=7.14 Hz, 2 H) 4.57 (t, J=8.79 Hz, 1 H) 5.87 (s, 1 H) 6.43 (d, J=2.34 Hz, 1 H) 6.89 - 7.03 (m, 1 H) 7.55 (d, J=8.30 Hz, 1 H) 7.68 - 7.79 (m, 2 H) 7.87 (dd, J=8.30, 2.15 Hz, 1 H) 7.94 (d, J=2.34 Hz, 1 H) 7.98 (d, J=1.51 Hz, 1 H) 8.08 - 8.18 (m, 2 H)
63dd	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (t, J=7.15 Hz, 3 H) 1.50 - 1.71 (m, 4 H) 2.00 (dd, J=13.54, 8.61 Hz, 1 H) 2.38 (s, 3 H) 2.42 (dd, J=13.59, 8.81 Hz, 1 H) 3.12 - 3.28 (m, 2 H) 3.42 - 3.77 (m, 6 H) 4.21 - 4.39 (m, 2 H) 4.48 (t, J=8.69 Hz, 1 H) 4.69 - 4.79 (m, 1 H) 5.72 (d, J=2.05 Hz, 1 H) 6.39 (d, J=2.29 Hz, 1 H) 6.77 (q, J=6.54 Hz, 1 H) 7.45 (d, J=1.56 Hz, 1 H) 7.52 (dd, J=8.20, 1.56 Hz, 1H) 7.70 (d, J=8.15 Hz, 1 H) 7.88 (dd, J=4.37, 2.37 Hz, 1 H)
63de	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.22 - 1.29 (m, 3 H) 1.51 (dt, J=11.74, 5.65 Hz, 4 H) 1.75 (dd, J=12.98, 7.32 Hz, 1 H) 1.90 (s, 4 H) 2.09 (dd, J=13.13, 8.74 Hz, 1 H) 2.38 (s, 3 H) 2.73 - 2.93 (m, 2 H) 3.43 - 3.63 (m, 4 H) 3.85 (dd, J=8.71, 7.39 Hz, 1 H) 4.10 (s, 2 H) 4.13 - 4.23 (m, 2 H) 5.70 (s, 1 H) 6.41 (d, J=2.25 Hz, 1 H) 6.78 (q, J=6.88 Hz, 1 H) 7.52 (d, J=8.35 Hz, 2 H) 7.66 (d, J=1.71 Hz, 1 H) 7.72 - 7.84 (m, 4 H) 7.97 (d, J=2.34 Hz, 1 H)
63df	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.33 (dt, J=10.30, 7.13 Hz, 6 H) 1.50 - 1.69

	(m, 4 H) 2.01 (dd, J=13.57, 8.74 Hz, 1 H) 2.40 - 2.45 (m, 1 H), 2.41 (s, 3 H) 3.22 (d, J=2.00 Hz, 2 H) 3.44 - 3.74 (m, 4 H) 4.21 - 4.36 (m, 4 H) 4.52 (t, J=8.74 Hz, 1 H) 5.80 (s, 1 H) 6.43 (d, J=2.29 Hz, 1 H) 6.61 (d, J=16.06 Hz, 1 H) 6.92 (q, J=6.65 Hz, 1 H) 7.49 - 7.56 (m, 2 H) 7.61 - 7.70 (m, 2 H) 7.72 - 7.80 (m, 2 H) 7.83 (dd, J=8.27, 2.17 Hz, 1 H) 7.94 (dd, J=6.39, 1.85 Hz, 2 H)
63dg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 - 1.39 (m, 6 H) 1.52 - 1.63 (m, 4 H) 1.94 (dd, J=13.50, 8.32 Hz, 1 H) 2.34 (dd, J=13.40, 8.76 Hz, 1 H) 2.40 (s, 3 H) 3.04 - 3.20 (m, 2 H) 3.44 - 3.69 (m, 4 H) 4.21 - 4.31 (m, 4 H) 4.34 (t, J=8.54 Hz, 1 H) 5.78 (s, 1 H) 6.42 (d, J=2.20 Hz, 1 H) 6.58 (d, J=16.06 Hz, 1 H) 6.92 (q, J=6.67 Hz, 1 H) 7.51 (d, J=8.25 Hz, 1 H) 7.62 - 7.77 (m, 5 H) 7.82 (dd, J=8.30, 2.15 Hz, 1 H) 7.93 (d, J=2.34 Hz, 1 H) 7.97 (d, J=1.27 Hz, 1 H)
63dh	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.18 (t, J=7.13 Hz, 3 H) 1.32 (t, J=7.15 Hz, 3 H) 1.54 - 1.68 (m, 4 H) 2.03 (dd, J=13.62, 8.83 Hz, 1 H) 2.40 (s, 3 H) 2.46 (dd, J=13.59, 8.76 Hz, 1 H) 2.68 (t, J=7.47 Hz, 2 H) 3.01 (t, J=7.49 Hz, 2 H) 3.25 (s, 2 H) 3.45 - 3.75 (m, 4 H) 4.10 (q, J=7.13 Hz, 2 H) 4.25 - 4.40 (m, 2 H) 4.57 (t, J=8.79 Hz, 1 H) 5.82 (s, 1 H) 6.42 (d, J=2.25 Hz, 1 H) 6.88 (q, J=6.70 Hz, 1 H) 7.27 (d, J=7.42 Hz, 1 H) 7.36 - 7.42 (m, 1 H) 7.42 - 7.47 (m, 2 H) 7.50 (d, J=8.30 Hz, 1 H) 7.79 (dd, J=8.27, 2.12 Hz, 1 H) 7.88 - 7.94 (m, 2 H)
63di	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.22 (t, J=7.13 Hz, 3 H) 1.32 (t, J=7.15 Hz, 3 H) 1.53 - 1.72 (m, 4 H) 2.03 (dd, J=13.62, 8.83 Hz, 1 H) 2.40 (s, 3 H) 2.47 (dd, J=13.59, 8.76 Hz, 1 H) 2.67 (t, J=7.57 Hz, 2 H) 2.92 - 3.03 (m, 2 H) 3.25 (s, 2 H) 3.45 - 3.80 (m, 4 H) 4.11 (q, J=7.13 Hz, 2 H) 4.32 (qd, J=7.13, 2.32 Hz, 2 H) 4.57 (t, J=8.79 Hz, 1 H) 5.84 (s, 1 H) 6.41 (d, J=2.34 Hz, 1 H) 6.87 (q, J=6.57 Hz, 1 H) 7.34 (d, J=8.25 Hz, 2 H) 7.49 (d, J=8.30 Hz, 1 H) 7.53 (d, J=8.25 Hz, 2 H) 7.78 (dd, J=8.30, 2.15 Hz, 1 H) 7.90 (d, J=2.24 Hz, 2 H)
63dj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.20 - 1.35 (m, 3 H) 1.42 - 1.60 (m, 4 H) 1.68 - 1.83 (m, 1 H) 1.99 - 2.15 (m, 1 H) 2.42 (s, 3 H) 2.75 (d, J=10.93 Hz, 1 H) 2.89 (d, J=10.93 Hz, 1 H) 3.55 (d, J=5.86 Hz, 4 H) 3.82 (s, 1 H) 4.20 (dd, J=7.13, 1.27 Hz, 2 H) 5.74 (s, 1 H) 6.44 (d, J=2.15 Hz, 1 H) 6.85 (d, J=6.64 Hz, 1 H) 7.04 - 7.24 (m, 1 H) 7.39 - 7.53 (m, 3 H) 7.67 (d, J=1.56 Hz, 1 H) 7.76 (d, J=1.76 Hz, 1 H) 7.79 - 7.88 (m, 1 H) 8.01 (d, J=2.15 Hz, 1 H)
63dk	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (t, J=7.13 Hz, 3 H) 1.42 - 1.63 (m, 4 H) 1.69 - 1.82 (m, 1 H) 2.01 - 2.16 (m, 1 H) 2.44 (s, 3 H) 2.77 (s, 1 H) 2.89 (s, 1 H) 3.57 (d, J=5.86 Hz, 4 H) 3.76 - 3.89 (m, 1 H) 4.19 (dd, J=7.22, 1.37 Hz, 2 H) 5.78 (s, 1 H) 6.46 (d, J=2.15 Hz, 1 H) 6.78 - 6.96 (m, 1 H) 7.53 - 7.65 (m, 1 H) 7.79 - 7.98 (m, 3 H) 8.06 (d, J=2.34 Hz, 1 H) 8.14 (s, 2 H) 8.28 (s, 1 H) 8.40 - 8.52 (m, 1 H) 8.80 - 8.96 (m, 1 H)
63dl	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (t, J=7.13 Hz, 3 H) 1.51 - 1.68 (m, 4 H) 1.96 - 2.06 (m, 1 H) 2.39 (s, 3 H) 2.43 (dd, J=13.54, 8.81 Hz, 1 H) 3.15 - 3.27 (m, 2 H) 3.43 - 3.72 (m, 4 H) 4.26 - 4.36 (m, 2 H) 4.39 - 4.45 (m, 1 H) 4.49 (t, J=8.69 Hz, 1 H) 4.90 (t, J=8.44 Hz, 1 H) 5.70 (d, J=2.83 Hz, 1 H) 5.87 (td, J=7.86, 1.66 Hz, 1 H) 6.42 (d, J=2.34 Hz, 1 H) 6.79 - 6.91 (m, 1 H) 7.52 (t, J=1.85 Hz, 1 H) 7.58 (dt, J=8.20, 2.17 Hz, 1 H) 7.82 (dd, J=8.27, 1.54 Hz, 1 H) 7.95 (dd, J=3.44, 2.61 Hz, 1 H)
63dm	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (t, J=7.13 Hz, 3 H) 1.43 - 1.62 (m, 4 H) 1.70 - 1.83 (m, 1 H) 2.03 - 2.18 (m, 1 H) 2.79 (s, 1 H) 2.90 (s, 1 H) 3.09 (s, 2 H) 3.17 (s, 3 H) 3.54 (br. s., 4 H) 3.65 (t, J=6.74 Hz, 2 H) 3.80 - 3.96 (m, 1 H) 4.21 (d, J=7.03

	Hz, 2 H) 5.57 (s, 1 H) 6.60 - 6.76 (m, 1 H) 7.52 (s, 1 H) 7.56 - 7.67 (m, 3 H) 7.68 - 7.79 (m, 2 H) 8.01 (d, J=8.20 Hz, 1 H)
63dn	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (t, J=7.13 Hz, 3 H) 1.55 (dt, J=11.03, 5.37 Hz, 4 H) 1.89 (dd, J=13.32, 8.10 Hz, 1 H) 1.96 (s, 3 H) 1.97 (s, 3 H) 2.28 (dd, J=13.40, 8.86 Hz, 1 H) 2.36 (s, 3 H) 2.97 - 3.11 (m, 2 H) 3.41 - 3.68 (m, 4 H) 4.18 - 4.30 (m, 3 H) 4.39 (s, 2 H) 5.69 (s, 1 H) 6.37 (d, J=2.34 Hz, 1 H) 6.73 (q, J=6.30 Hz, 1 H) 7.31 (d, J=1.51 Hz, 1 H) 7.40 (dd, J=8.13, 1.59 Hz, 1 H), 7.67 (d, J=8.05 Hz, 1 H), 7.85 (d, J=2.25 Hz, 1 H)
63do	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.31 (t, J=7.15 Hz, 3 H) 1.55 - 1.68 (m, 4 H) 1.97 (s, 1 H) 2.02 (dd, J=13.59, 8.96 Hz, 1 H) 2.12 (ddt, J=13.30, 7.49, 5.74, 5.74 Hz, 1 H) 2.39 (s, 3 H) 2.46 (dd, J=13.57, 8.64 Hz, 1 H) 2.65 - 2.77 (m, 1 H) 2.91 (dt, J=13.74, 7.04 Hz, 1 H) 3.04 - 3.14 (m, 1 H) 3.24 (d, J=1.76 Hz, 2 H) 3.45 - 3.76 (m, 6 H) 3.81 (dd, J=8.52, 5.88 Hz, 1 H) 4.25 - 4.45 (m, 4 H) 4.55 (t, J=8.79 Hz, 1 H) 5.81 (s, 1 H) 6.41 (d, J=2.29 Hz, 1 H) 6.82 (q, J=6.65 Hz, 1 H) 7.71 (s, 1 H) 7.77 - 7.84 (m, 4 H) 7.90 - 8.01 (m, 3 H)
63dp	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.33 (t, J=7.13 Hz, 3 H) 1.54 - 1.75 (m, 4 H) 2.02 (dd, J=13.57, 8.69 Hz, 1 H) 2.28 (s, 3 H) 2.31 (s, 3 H) 2.45 (dd, J=13.54, 8.76 Hz, 1 H) 3.20 - 3.27 (m, 2 H) 3.23 (s, 3 H) 3.42 - 3.79 (m, 4 H) 4.32 (qd, J=7.13, 2.54 Hz, 2 H) 4.51 (t, J=8.69 Hz, 1 H) 5.63 (s, 1 H) 6.63 (q, J=6.64 Hz, 1 H) 7.19 (d, J=7.86 Hz, 1 H) 7.36 (dd, J=7.76, 1.76 Hz, 1 H) 7.41 (s, 1 H) 7.48 (d, J=1.81 Hz, 1 H) 7.65 - 7.72 (m, 1 H) 7.72 - 7.77 (m, 1 H) 7.80 - 7.85 (m, 2 H) 8.07 (dt, J=7.03, 1.93 Hz, 1 H) 8.47 (br. s., 1 H)
63dq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (t, J=7.13 Hz, 3 H) 1.53 - 1.73 (m, 4 H) 1.93 - 2.08 (m, 1 H) 2.41 (dd, J=13.45, 8.71 Hz, 1 H) 3.11 - 3.23 (m, 2 H) 3.24 (s, 3 H) 3.46 - 3.79 (m, 4 H) 4.30 (qd, J=7.12, 2.46 Hz, 2 H) 4.43 (t, J=8.59 Hz, 1 H) 5.65 (s, 1 H) 6.67 (q, J=6.64 Hz, 1 H) 7.56 (dd, J=8.32, 4.32 Hz, 1 H) 7.70 (d, J=1.76 Hz, 1 H) 7.78 - 7.97 (m, 4 H) 8.07 δ 8.12 (m, 3 H) 8.23 (s, 1 H), 8.42 (dd, J=8.44, 1.61 Hz, 1 H) 8.53 (br. s., 1 H) 8.85 (dd, J=4.32, 1.68 Hz, 1 H)
63dr	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (t, J=7.13 Hz, 3 H) 1.51 - 1.71 (m, 4 H) 1.99 - 2.08 (m, 1 H) 2.40 (s, 3 H) 2.41 (s, 3 H) 2.46 (dd, J=13.62, 8.74 Hz, 1 H) 3.25 (d, J=1.07 Hz, 2 H) 3.43 - 3.77 (m, 4 H) 4.32 (qd, J=7.13, 2.32 Hz, 2 H) 4.57 (t, J=8.79 Hz, 1 H) 4.68 (s, 2 H) 5.85 (s, 1 H) 6.42 (d, J=2.25 Hz, 1 H) 6.88 (q, J=6.65 Hz, 1 H) 7.34 - 7.54 (m, 4 H) 7.79 (dd, J=8.27, 2.12 Hz, 1 H) 7.91 (d, J=2.25 Hz, 2 H)
63ds	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (t, J=7.15 Hz, 3 H) 1.55 - 1.77 (m, 4 H) 2.04 (dd, J=9.32, 4.30 Hz, 1 H) 2.37 (s, 3 H) 2.40 (s, 3 H) 2.48 (dd, J=13.62, 8.79 Hz, 1 H) 3.28 (s, 2 H) 3.50 - 3.87 (m, 4 H) 4.19 - 4.42 (m, 2 H) 4.59 (t, J=8.79 Hz, 1 H) 4.71 (s, 2 H) 6.14 (br. s., 1 H) 6.42 (d, J=2.34 Hz, 1 H) 6.91 (q, J=6.41 Hz, 1 H) 7.28 (d, J=7.91 Hz, 1 H) 7.44 (dd, J=7.76, 2.05 Hz, 1 H) 7.51 (d, J=8.25 Hz, 1 H) 7.65 (d, J=1.81 Hz, 1 H) 7.85 (dd, J=8.27, 2.12 Hz, 1 H) 7.88 - 7.96 (m, 2 H)
63dt	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (t, J=7.13 Hz, 3 H) 1.52 - 1.71 (m, 4 H) 1.98 - 2.08 (m, 1 H) 2.41 (s, 3 H) 2.45 (dd, J=13.62, 8.79 Hz, 1 H) 3.24 (d, J=1.61 Hz, 2 H) 3.42 - 3.73 (m, 4 H) 3.93 (s, 3 H) 4.24 - 4.40 (m, 2 H) 4.54 (t, J=8.76 Hz, 1 H) 5.78 (s, 1 H) 6.43 (d, J=2.29 Hz, 1 H) 6.95 (q, J=6.62 Hz, 1 H) 7.54 (d, J=8.30 Hz, 1 H) 7.67 - 7.77 (m, 2 H) 7.85 (dd, J=8.30, 2.20 Hz, 1 H) 7.94 (d, J=2.34 Hz, 1 H) 7.98 (d, J=1.37 Hz, 1 H) 8.07 - 8.16 (m, 2 H)
63du	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (t, J=7.13 Hz, 3 H) 1.46 - 1.67 (m, 4 H)

	1.90 (dd, J=13.35, 8.22 Hz, 1 H) 2.30 (dd, J=13.35, 8.71 Hz, 1 H) 2.40 (s, 3 H) 2.93 - 3.16 (m, 2 H) 3.38 - 3.74 (m, 4 H) 4.16 - 4.36 (m, 3 H) 5.77 (s, 1 H) 6.41 (d, J=2.29 Hz, 1 H) 6.91 (q, J=6.62 Hz, 1 H) 7.51 (d, J=8.25 Hz, 1 H) 7.64 (d, J=8.40 Hz, 2 H) 7.82 (dd, J=8.30, 2.15 Hz, 1 H) 7.93 (d, J=2.29 Hz, 1 H) 7.96 (s, 1 H) 8.07 (d, J=8.30 Hz, 2 H)
63dv	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (s, 4 H) 1.60 - 1.76 (m, 3 H) 2.02 - 2.13 (m, 1 H) 2.43 (s, 3 H) 2.46 - 2.56 (m, 1 H) 3.30 (s, 2 H) 3.53 - 3.83 (m, 4 H) 4.30 - 4.40 (m, 2 H) 4.56 - 4.66 (m, 1 H) 5.48 (s, 2 H) 6.01 - 6.11 (m, 1 H) 6.44 - 6.50 (m, 1 H) 6.89 - 6.98 (m, 1 H) 7.80 - 7.84 (m, 1 H) 7.86 - 8.02 (m, 5 H) 8.03 - 8.08 (m, 1 H)
63dw	¹ H NMR (400 MHz, MeOH -d ₄): δ ppm 1.28 (t, J=7.13 Hz, 3 H) 1.55 (br. s., 4 H) 1.70 - 1.83 (m, 1 H) 2.03 - 2.18 (m, 1 H) 2.70 - 2.82 (m, 1 H) 2.86 - 2.97 (m, 1 H) 3.41 - 3.59 (m, 4 H) 3.79 - 3.94 (m, 1 H) 4.09 - 4.30 (m, 2 H) 5.59 (s, 1 H) 6.60 - 6.77 (m, 1 H) 7.71 (d, J=8.20 Hz, 2 H) 7.84 (d, J=8.20 Hz, 2 H) 8.03 - 8.15 (m, 1 H) 8.34 (s, 2 H) 9.26 (s, 1 H) 9.59 (s, 1 H)
63dx	¹ H NMR (400 MHz, Chloroform-d): δ ppm 1.27 (m, 7H), 1.55 (m, 3H), 1.77 (dd, J = 13.1, 7.0 Hz, 1H), 2.11 (dd, J = 13.1, 8.9 Hz, 1H), 2.42 (s, 3H), 2.96 (m, 2H), 3.47 (dt, J = 11.6, 5.7 Hz, 4H), 3.98 (dd, J = 8.8, 6.9 Hz, 1H), 4.21 (q, J = 7.1 Hz, 2H), 4.73 (s, 2H), 5.49 (s, 1H), 5.99 (m, 1H), 6.35 (d, J = 2.3 Hz, 1H), 6.63 (q, J = 6.7 Hz, 1H), 7.61 (m, 2H), 7.73 (d, J = 2.3 Hz, 1H), 7.88 (d, J = 8.3 Hz, 1H), 8.96 (s, 2H), 9.24 (s, 1H)
63dy	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (m, 8H), 1.51 (dt, J = 10.8, 5.6 Hz, 8H), 1.74 (dd, J = 13.1, 7.3 Hz, 2H), 2.06 (m, 2H), 2.39 (s, 6H), 2.75 (d, J = 11.0 Hz, 2H), 2.89 (d, J = 11.0 Hz, 2H), 3.53 (dt, J = 22.5, 6.4 Hz, 8H), 3.82 (dd, J = 8.8, 7.2 Hz, 2H), 4.18 (qd, J = 7.1, 1.6 Hz, 3H), 5.73 (s, 2H), 6.00 (m, 1H), 6.42 (d, J = 2.4 Hz, 2H), 6.82 (q, J = 6.6 Hz, 2H), 7.35 (dt, J = 10.4, 8.4 Hz, 2H), 7.50 (m, 2H), 7.73 (m, 8H), 8.00 (d, J = 2.4 Hz, 2H)
63dz	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.50 (dt, J = 10.6, 5.6 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.52 (dq, J = 23.8, 7.6, 6.5 Hz, 4H), 3.81 (dd, J = 8.8, 7.1 Hz, 1H), 4.17 (qd, J = 7.1, 1.6 Hz, 2H), 5.73 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.80 (q, J = 6.5 Hz, 1H), 7.46 (m, 2H), 7.71 (m, 5H), 7.98 (d, J = 2.3 Hz, 1H)
63ea	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.25 (t, J=7.10 Hz, 3 H) 1.42 - 1.69 (m, 4 H) 1.92 (dd, J=13.25, 9.35 Hz, 1 H) 2.35 (dd, J=13.25, 8.47 Hz, 1 H) 3.14 (br. s., 2 H) 3.60 (br. s., 4 H) 4.24 (qd, J=7.09, 2.10 Hz, 2 H) 4.54 (br. s., 1 H) 5.77 (br. s., 1 H) 6.70 (q, J=6.65 Hz, 1 H) 7.37 (d, J=2.10 Hz, 1 H) 7.43 - 7.52 (m, 3 H) 7.53 - 7.69 (m, 4 H) 9.23 (br. s., 1 H) 10.44 (br. s., 1 H)
63eb	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.25 (t, J = 7.1 Hz, 4H), 1.49 (dt, J = 10.6, 5.6 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.06 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.50 (m, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.17 (qd, J = 7.1, 1.5 Hz, 2H), 5.76 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.89 (q, J = 6.6 Hz, 1H), 7.51 (m, 6H), 7.76 (dd, J = 8.3, 2.2 Hz, 1H), 7.92 (t, J = 2.2 Hz, 2H)

63ec	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 5H), 1.51 (dt, J = 10.1, 5.3 Hz, 4H), 1.77 (dd, J = 13.1, 7.4 Hz, 1H), 2.12 (dd, J = 13.2, 8.8 Hz, 1H), 2.39 (s, 3H), 2.80 (d, J = 11.1 Hz, 1H), 2.93 (d, J = 11.1 Hz, 1H), 3.53 (m, 4H), 3.91 (t, J = 8.0 Hz, 1H), 4.19 (qd, J = 7.2, 1.7 Hz, 2H), 5.77 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.91 (q, J = 6.6 Hz, 1H), 7.37 (ddt, J = 16.6, 10.3, 8.6 Hz, 2H), 7.51 (m, 2H), 7.76 (dd, J = 8.3, 2.2 Hz, 1H), 7.91 (dd, J = 9.9, 2.4 Hz, 2H)
63ed	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (m, 6H), 1.50 (dt, J = 10.4, 5.4 Hz, 5H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.06 (m, 1H), 2.39 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.53 (m, 5H), 3.81 (dd, J = 8.7, 7.1 Hz, 1H), 4.17 (qd, J = 7.1, 1.5 Hz, 2H), 5.77 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.87 (q, J = 6.7 Hz, 1H), 7.44 (m, 4H), 7.60 (m, 3H), 7.78 (dd, J = 8.3, 2.1 Hz, 1H), 7.92 (t, J = 2.6 Hz, 2H)
63ee	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (q, J = 8.3, 7.1 Hz, 6H), 1.49 (m, 4H), 1.72 (dd, J = 13.1, 7.2 Hz, 1H), 2.06 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.50 (m, 4H), 3.83 (dd, J = 8.8, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.5 Hz, 2H), 5.77 (s, 1H), 6.41 (d, J = 2.5 Hz, 1H), 6.93 (q, J = 6.6 Hz, 1H), 7.50 (m, 2H), 7.59 (d, J = 8.3 Hz, 1H), 7.73 (m, 2H), 7.92 (dd, J = 7.7, 2.3 Hz, 2H)
63ef	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (t, J = 7.1 Hz, 5H), 1.54 (tt, J = 7.3, 4.4 Hz, 4H), 1.80 (dd, J = 13.2, 7.6 Hz, 1H), 2.17 (dd, J = 13.2, 8.8 Hz, 1H), 2.85 (d, J = 11.2 Hz, 1H), 2.97 (d, J = 11.2 Hz, 1H), 3.52 (ddt, J = 28.3, 12.4, 8.0 Hz, 4H), 3.97 (dd, J = 8.8, 7.5 Hz, 1H), 4.21 (qd, J = 7.1, 1.7 Hz, 2H), 5.50 (s, 1H), 6.68 (q, J = 6.9 Hz, 1H), 7.25 (m, 4H), 7.48 (m, 3H), 7.71 (m, 1H)
63eg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.3 Hz, 4H), 1.51 (dt, J = 11.2, 5.5 Hz, 4H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.40 (s, 3H), 2.78 (d, J = 11.1 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.53 (m, 5H), 3.86 (dd, J = 8.6, 7.4 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.77 (s, 1H), 6.44 (d, J = 2.4 Hz, 1H), 7.00 (q, J = 6.7 Hz, 1H), 7.62 (d, J = 8.3 Hz, 1H), 7.97 (m, 3H), 9.06 (s, 2H), 9.17 (s, 1H)
63eh	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 4H), 1.52 (dq, J = 12.0, 8.6, 7.2 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.09 (dd, J = 13.1, 8.8 Hz, 1H), 2.75 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.51 (m, 4H), 3.82 (dd, J = 8.7, 7.1 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.52 (s, 1H), 6.61 (q, J = 6.7 Hz, 1H), 7.33 (m, 2H), 7.47 (m, 2H), 7.66 (m, 2H)
63ei	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (td, J = 7.1, 1.7 Hz, 6H), 1.41 (td, J = 7.0, 1.8 Hz, 3H), 1.52 (qd, J = 7.2, 4.7, 3.6 Hz, 4H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.77 (d, J = 10.9 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.29 (s, 1H), 3.51 (dq, J = 18.9, 6.1 Hz, 4H), 3.84 (m, 1H), 4.16 (m, 3H), 5.48 (d, J = 2.0 Hz, 1H), 6.73 (q, J = 6.8 Hz, 1H), 7.02 (m, 2H), 7.20 (s, 1H), 7.28 (d, J = 2.2 Hz, 1H), 7.43 (m, 2H), 7.67 (d, J = 8.4 Hz, 1H)
63ej	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (m, 4H), 1.50 (dt, J = 22.1, 6.5 Hz, 7H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.08 (dd, J = 13.1, 8.8 Hz, 1H), 2.75 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.31 (d, J = 1.8 Hz, 1H), 3.49 (dq, J = 25.8, 7.5, 6.6 Hz, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (m, 4H), 5.49 (s, 1H), 6.62 (q, J = 6.8 Hz, 1H), 7.20 (d, J = 8.5 Hz, 1H), 7.26 (d, J = 2.2 Hz, 1H), 7.34 (m, 1H), 7.42 (dd, J = 8.5, 2.2 Hz, 1H), 7.58 (m, 1H), 7.65 (d, J = 8.5 Hz, 1H)
63ek	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.0 Hz, 4H), 1.53 (m, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.7 Hz, 1H), 2.76 (d, J = 10.9 Hz, 1H),

	2.90 (d, J = 11.0 Hz, 1H), 3.51 (dq, J = 26.1, 7.5, 6.6 Hz, 4H), 3.83 (dd, J = 8.8, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.53 (s, 1H), 6.59 (q, J = 6.7 Hz, 1H), 7.27 (d, J = 8.8 Hz, 1H), 7.34 (m, 2H), 7.49 (m, 2H), 7.69 (d, J = 8.5 Hz, 1H)
63el	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (m, 2H), 1.31 (m, 15H), 1.52 (dt, J = 7.9, 4.5 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.09 (dd, J = 13.1, 8.8 Hz, 1H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.49 (ddt, J = 17.6, 11.8, 6.7 Hz, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.2, 1.5 Hz, 2H), 5.44 (s, 1H), 6.60 (q, J = 6.8 Hz, 1H), 7.28 (m, 2H), 7.47 (m, 4H), 7.67 (d, J = 8.5 Hz, 1H)
63em	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (m, 6H), 1.53 (m, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.7 Hz, 1H), 2.76 (d, J = 10.9 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.83 (dd, J = 8.8, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.7 Hz, 2H), 5.53 (s, 1H), 6.47 (q, J = 6.7 Hz, 1H), 7.36 (d, J = 2.2 Hz, 1H), 7.53 (dd, J = 8.6, 2.3 Hz, 1H), 7.72 (m, 2H), 7.85 (m, 2H)
63en	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (t, J = 7.1 Hz, 7H), 1.59 (dt, J = 11.5, 5.7 Hz, 8H), 1.95 (m, 2H), 2.34 (dd, J = 13.3, 8.6 Hz, 2H), 3.10 (m, 4H), 3.56 (m, 8H), 3.75 (s, 1H), 4.28 (m, 6H), 5.55 (s, 1H), 6.52 (q, J = 6.7 Hz, 2H), 7.38 (d, J = 2.2 Hz, 2H), 7.53 (dd, J = 8.5, 2.2 Hz, 2H), 7.66 (m, 8H)
63eo	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (td, J = 7.0, 0.7 Hz, 4H), 1.53 (m, 4H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.09 (dd, J = 13.1, 8.7 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.30 (m, 1H), 3.51 (dtd, J = 19.2, 13.4, 7.5 Hz, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (m, 2H), 5.50 (s, 1H), 6.63 (q, J = 6.9 Hz, 1H), 7.25 (m, 1H), 7.47 (m, 6H), 7.71 (m, 1H)
63ep	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.33 (t, J = 7.1 Hz, 4H), 1.65 (m, 5H), 2.06 (dd, J = 13.6, 8.9 Hz, 1H), 2.50 (dd, J = 13.6, 8.8 Hz, 1H), 3.28 (s, 2H), 3.57 (m, 5H), 3.86 (s, 3H), 4.32 (qd, J = 7.2, 2.5 Hz, 2H), 4.59 (t, J = 8.8 Hz, 1H), 6.68 (q, J = 6.7 Hz, 1H), 6.98 (d, J = 7.5 Hz, 1H), 7.07 (m, 2H), 7.32 (d, J = 2.2 Hz, 1H), 7.46 (m, 2H), 7.68 (d, J = 8.5 Hz, 1H)
63eq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.31 (m, 10H), 1.52 (m, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.09 (dd, J = 13.1, 8.7 Hz, 1H), 2.75 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.50 (m, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.5 Hz, 2H), 4.70 (h, J = 6.1 Hz, 1H), 5.48 (s, 1H), 6.73 (q, J = 6.9 Hz, 1H), 6.95 (d, J = 7.6 Hz, 1H), 7.04 (m, 1H), 7.21 (s, 1H), 7.27 (d, J = 2.3 Hz, 1H), 7.42 (m, 2H), 7.66 (d, J = 8.5 Hz, 1H)
63er	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (td, J = 7.1, 1.4 Hz, 3H), 1.54 (m, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.7 Hz, 1H), 2.46 (s, 3H), 2.76 (d, J = 10.9 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.2, 1.6 Hz, 2H), 5.50 (d, J = 1.5 Hz, 1H), 6.60 (q, J = 6.8 Hz, 1H), 7.29 (m, 2H), 7.45 (m, 2H), 7.57 (s, 1H), 7.67 (m, 1H)
63es	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm: 1.19 (s, 2H), 1.34 (m, 11H), 1.53 (m, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.83 (m, 1H), 4.18 (m, 2H), 4.73 (h, J = 6.1 Hz, 1H), 5.50 (s, 1H), 6.62 (q, J = 6.6 Hz, 1H), 7.28 (m, 3H), 7.43 (dd, J = 8.5, 2.3 Hz, 1H), 7.59 (s, 1H), 7.65 (d, J = 8.5 Hz, 1H)
63et	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.88 (d, J = 8.3 Hz, 1H), 1.30 (m, 11H), 1.53 (m, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.52 (m, 5H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H),

	4.18 (qd, J = 7.1, 1.6 Hz, 2H), 4.71 (h, J = 6.2 Hz, 1H), 5.49 (s, 1H), 6.66 (q, J = 6.8 Hz, 1H), 7.27 (m, 4H), 7.43 (dd, J = 8.5, 2.3 Hz, 1H), 7.65 (m, 1H)
63eu	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (m, 4H), 1.60 (tt, J = 9.2, 4.2 Hz, 4H), 1.97 (m, 1H), 2.37 (dd, J = 13.6, 8.8 Hz, 1H), 3.14 (q, J = 11.6 Hz, 2H), 3.56 (m, 4H), 4.31 (m, 3H), 4.87 (d, J = 1.7 Hz, 18H), 5.54 (s, 1H), 6.49 (q, J = 6.6 Hz, 1H), 7.35 (d, J = 2.2 Hz, 1H), 7.51 (dd, J = 8.5, 2.2 Hz, 1H), 7.68 (d, J = 8.6 Hz, 1H), 7.82 (m, 3H)
63ev	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (m, 4H), 1.54 (m, 4H), 1.78 (dd, J = 13.1, 7.4 Hz, 1H), 2.13 (dd, J = 13.1, 8.8 Hz, 1H), 2.81 (d, J = 11.1 Hz, 1H), 2.93 (d, J = 11.0 Hz, 1H), 3.53 (dq, J = 13.2, 6.0 Hz, 4H), 3.89 (dd, J = 8.8, 7.3 Hz, 1H), 4.20 (qd, J = 7.1, 1.7 Hz, 2H), 5.51 (s, 1H), 6.63 (q, J = 6.7 Hz, 1H), 7.28 (m, 4H), 7.52 (m, 2H), 7.68 (d, J = 8.5 Hz, 1H)
63ew	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.90 (d, J = 8.0 Hz, 1H), 1.27 (m, 5H), 1.53 (dt, J = 7.7, 4.7 Hz, 4H), 1.76 (dd, J = 13.1, 7.2 Hz, 1H), 2.11 (dd, J = 13.1, 8.8 Hz, 1H), 2.45 (s, 3H), 2.78 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.31 (s, 3H), 3.50 (dq, J = 27.7, 7.7, 6.6 Hz, 4H), 3.85 (t, J = 8.0 Hz, 1H), 4.19 (m, 2H), 5.49 (s, 1H), 6.61 (q, J = 6.8 Hz, 1H), 7.32 (m, 3H), 7.45 (dd, J = 8.6, 2.3 Hz, 1H), 7.52 (d, J = 8.1 Hz, 1H), 7.67 (d, J = 8.5 Hz, 1H)
63ex	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 4H), 1.54 (m, 4H), 1.76 (dd, J = 13.1, 7.2 Hz, 1H), 2.11 (dd, J = 13.1, 8.8 Hz, 1H), 2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.54 (m, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.1, 1.7 Hz, 2H), 5.53 (s, 1H), 6.54 (q, J = 6.7 Hz, 1H), 7.36 (d, J = 2.2 Hz, 1H), 7.53 (dd, J = 8.6, 2.2 Hz, 1H), 7.68 (dd, J = 23.2, 8.4 Hz, 2H), 7.84 (s, 1H), 7.97 (d, J = 8.1 Hz, 1H)
63ey	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 4H), 1.54 (dddd, J = 11.9, 6.5, 4.3 Hz, 4H), 1.76 (dd, J = 13.1, 7.2 Hz, 1H), 2.11 (dd, J = 13.1, 8.7 Hz, 1H), 2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.52 (dq, J = 26.9, 7.7, 6.7 Hz, 4H), 3.84 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.2, 1.6 Hz, 2H), 5.53 (s, 1H), 6.63 (q, J = 6.7 Hz, 1H), 7.10 (tt, J = 9.2, 2.4 Hz, 1H), 7.19 (s, 2H), 7.34 (d, J = 2.2 Hz, 1H), 7.50 (dd, J = 8.6, 2.2 Hz, 1H), 7.70 (d, J = 8.5 Hz, 1H)
63ez	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.07 (d, J = 1.0 Hz, 1H), 0.88 (dd, J = 14.3, 7.9 Hz, 1H), 1.28 (m, 6H), 1.56 (m, 3H), 1.75 (dd, J = 13.1, 6.7 Hz, 1H), 2.07 (dd, J = 13.1, 8.8 Hz, 2H), 2.84 (d, J = 10.5 Hz, 1H), 2.96 (d, J = 10.5 Hz, 1H), 3.49 (m, 4H), 3.88 (m, 1H), 4.20 (m, 2H), 4.56 (s, 2H), 5.44 (d, J = 1.0 Hz, 1H), 6.52 (q, J = 6.7 Hz, 1H), 7.23 (m, 3H), 7.40 (m, 1H), 7.66 (m, 2H)
63fa	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 3H), 1.53 (m, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 10.9 Hz, 1H), 3.52 (m, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.2, 1.6 Hz, 2H), 5.52 (s, 1H), 6.60 (q, J = 6.8 Hz, 1H), 7.31 (d, J = 2.3 Hz, 1H), 7.45 (m, 3H), 7.67 (d, J = 8.5 Hz, 1H)
63fb	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (dd, J = 10.8, 3.8 Hz, 1H), 1.27 (t, J = 7.1 Hz, 4H), 1.53 (m, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.7 Hz, 1H), 2.25 (s, 1H), 2.35 (d, J = 3.2 Hz, 6H), 2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.50 (m, 4H), 3.85 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.2, 1.6 Hz, 2H), 4.87 (d, J = 5.2 Hz, 13H), 5.44 (s, 1H), 6.64 (q, J = 6.8 Hz, 1H), 7.18 (d, J = 7.7 Hz, 2H), 7.26 (m, 2H), 7.41 (dd, J = 8.5, 2.3 Hz, 1H), 7.65 (d, J = 8.5 Hz, 1H)
63fc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (t, J = 6.7 Hz, 1H), 1.30 (m, 11H), 1.59

	(dt, J = 7.7, 4.0 Hz, 9H), 1.94 (dd, J = 13.4, 8.3 Hz, 2H), 2.45 (s, 15H), 2.80 (s, 3H), 3.12 (m, 4H), 3.55 (tdd, J = 24.1, 17.0, 12.0 Hz, 9H), 4.28 (m, 6H), 5.51 (s, 2H), 6.64 (q, J = 6.8 Hz, 2H), 7.21 (s, 4H), 7.28 (d, J = 2.3 Hz, 2H), 7.44 (dd, J = 8.5, 2.3 Hz, 2H), 7.66 (d, J = 8.6 Hz, 2H)
63fd	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.08 (dd, J = 4.2, 1.7 Hz, 1H), 0.90 (q, J = 7.7 Hz, 1H), 1.26 (t, J = 7.1 Hz, 4H), 1.50 (dt, J = 33.9, 6.4 Hz, 7H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.50 (dq, J = 24.8, 7.5, 6.5 Hz, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (ttt, J = 7.1, 4.5, 2.1 Hz, 4H), 5.49 (s, 1H), 6.66 (q, J = 6.8 Hz, 1H), 7.27 (m, 4H), 7.43 (dd, J = 8.5, 2.3 Hz, 1H), 7.65 (d, J = 8.5 Hz, 1H)
63fe	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (td, J = 7.2, 1.1 Hz, 4H), 1.52 (dt, J = 11.8, 5.7 Hz, 4H), 1.74 (dd, J = 13.2, 7.2 Hz, 1H), 2.09 (dd, J = 13.2, 8.7 Hz, 1H), 2.38 (s, 6H), 2.75 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.50 (m, 5H), 3.82 (t, J = 8.0 Hz, 1H), 4.18 (m, 2H), 5.45 (s, 1H), 6.66 (q, J = 6.8 Hz, 1H), 7.04 (s, 2H), 7.12 (d, J = 1.8 Hz, 1H), 7.25 (d, J = 2.1 Hz, 1H), 7.41 (m, 1H), 7.66 (d, J = 8.5 Hz, 1H)
63ff	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 3H), 1.54 (m, 4H), 1.76 (dd, J = 13.1, 7.3 Hz, 1H), 2.11 (dd, J = 13.1, 8.7 Hz, 1H), 2.43 (s, 3H), 2.78 (d, J = 11.0 Hz, 1H), 2.92 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.86 (dd, J = 8.7, 7.3 Hz, 1H), 4.19 (qd, J = 7.1, 1.6 Hz, 2H), 5.50 (s, 1H), 6.61 (q, J = 6.8 Hz, 1H), 7.18 (s, 1H), 7.31 (m, 2H), 7.39 (s, 1H), 7.46 (dd, J = 8.5, 2.3 Hz, 1H), 7.68 (m, 1H)
63fg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (t, J = 6.9 Hz, 1H), 1.27 (m, 6H), 1.53 (dt, J = 7.9, 4.6 Hz, 4H), 1.76 (dd, J = 13.1, 7.3 Hz, 1H), 2.11 (m, 1H), 2.35 (d, J = 1.9 Hz, 3H), 2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.49 (dp, J = 25.1, 5.9 Hz, 4H), 3.84 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.1, 1.6 Hz, 2H), 4.87 (s, 13H), 5.49 (s, 1H), 6.62 (q, J = 6.8 Hz, 1H), 7.26 (m, 4H), 7.43 (dd, J = 8.6, 2.3 Hz, 1H), 7.66 (d, J = 8.5 Hz, 1H)
63fh	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 4H), 1.53 (dp, J = 7.3, 3.6, 2.9 Hz, 4H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.10 (dd, J = 13.1, 8.7 Hz, 1H), 2.40 (s, 3H), 2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.51 (m, 4H), 3.84 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.1, 1.6 Hz, 2H), 5.50 (s, 1H), 6.61 (q, J = 6.7 Hz, 1H), 7.30 (d, J = 2.3 Hz, 1H), 7.45 (m, 4H), 7.67 (d, J = 8.5 Hz, 1H)
63fi	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 5H), 1.53 (m, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.09 (dd, J = 13.1, 8.7 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.51 (dt, J = 18.9, 13.4, 7.4 Hz, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.2, 1.6 Hz, 2H), 5.50 (d, J = 10.3 Hz, 1H), 6.58 (q, J = 6.7 Hz, 1H), 7.32 (d, J = 2.2 Hz, 1H), 7.47 (m, 4H), 7.67 (m, 2H)
63fj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.87 (dd, J = 11.5, 4.7 Hz, 1H), 1.26 (ddd, J = 19.3, 6.9, 5.5 Hz, 12H), 1.48 (m, 4H), 1.72 (dd, J = 13.1, 7.3 Hz, 1H), 2.05 (m, 1H), 2.74 (d, J = 11.0 Hz, 1H), 2.93 (m, 2H), 3.45 (m, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.17 (qd, J = 7.1, 1.4 Hz, 2H), 5.44 (s, 1H), 6.63 (q, J = 6.8 Hz, 1H), 7.34 (m, 7H), 7.66 (d, J = 8.5 Hz, 1H)
63fk	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.88 (d, J = 7.9 Hz, 1H), 1.19 (s, 2H), 1.26 (t, J = 7.1 Hz, 4H), 1.53 (dt, J = 8.3, 4.9 Hz, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 10.9 Hz, 1H), 3.51 (m, 4H), 3.83 (dd, J = 8.8, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.7 Hz, 2H), 4.87 (s, 7H),

	5.52 (s, 1H), 6.40 (q, J = 6.6 Hz, 1H), 7.40 (d, J = 2.2 Hz, 1H), 7.55 (dd, J = 8.5, 2.3 Hz, 1H), 7.71 (d, J = 8.5 Hz, 1H), 8.11 (s, 2H)
63fl	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (m, 6H), 1.54 (m, 7H), 1.77 (dd, J = 13.1, 7.3 Hz, 2H), 2.12 (dd, J = 13.1, 8.7 Hz, 2H), 2.23 (dd, J = 4.4, 1.5 Hz, 3H), 2.36 (d, J = 2.1 Hz, 5H), 2.79 (d, J = 11.1 Hz, 2H), 2.93 (d, J = 11.0 Hz, 2H), 3.30 (d, J = 9.7 Hz, 1H), 3.52 (dq, J = 19.4, 6.4 Hz, 7H), 3.88 (dd, J = 8.8, 7.3 Hz, 2H), 4.19 (dddd, J = 8.7, 7.1, 5.6, 1.7 Hz, 3H), 5.49 (q, J = 2.5, 1.9 Hz, 2H), 6.65 (q, J = 6.7 Hz, 2H), 7.22 (m, 6H), 7.43 (m, 3H), 7.67 (d, J = 8.5 Hz, 2H)
63fm	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (m, 5H), 1.61 (m, 4H), 1.96 (dd, J = 13.4, 8.4 Hz, 1H), 2.37 (dd, J = 13.4, 8.7 Hz, 1H), 2.81 (s, 1H), 3.13 (q, J = 11.6 Hz, 2H), 3.57 (m, 4H), 4.30 (m, 3H), 5.56 (s, 1H), 6.55 (q, J = 6.7 Hz, 1H), 7.34 (d, J = 2.2 Hz, 1H), 7.55 (m, 5H), 7.69 (d, J = 8.5 Hz, 1H)
63fn	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (m, 4H), 1.53 (m, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.76 (d, J = 10.9 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.51 (m, 4H), 3.82 (dd, J = 8.8, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.53 (s, 1H), 6.49 (q, J = 6.7 Hz, 1H), 7.34 (d, J = 2.2 Hz, 1H), 7.52 (m, 2H), 7.68 (d, J = 8.6 Hz, 1H), 7.81 (s, 2H)
63fo	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 4H), 1.54 (m, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.77 (d, J = 11.0 Hz, 1H), 2.83 (s, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.53 (m, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.2, 1.6 Hz, 2H), 5.53 (s, 1H), 6.53 (q, J = 6.8 Hz, 1H), 7.37 (d, J = 2.2 Hz, 1H), 7.53 (dd, J = 8.5, 2.2 Hz, 1H), 7.62 (ddd, J = 7.9, 5.0, 0.9 Hz, 1H), 7.72 (d, J = 8.6 Hz, 1H), 7.98 (m, 1H), 8.67 (dd, J = 5.0, 1.6 Hz, 1H), 8.75 (d, J = 2.2 Hz, 1H)
63fp	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.31 (m, 10H), 1.54 (m, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.53 (m, 4H), 3.84 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.1, 1.6 Hz, 2H), 4.69 (p, J = 6.1 Hz, 1H), 5.51 (s, 1H), 6.72 (q, J = 7.0 Hz, 1H), 6.83 (dt, J = 11.3, 2.3 Hz, 1H), 7.03 (m, 1H), 7.30 (d, J = 2.2 Hz, 1H), 7.46 (dd, J = 8.5, 2.3 Hz, 1H), 7.67 (d, J = 8.5 Hz, 1H)
63fq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.88 (m, 1H), 1.37 (m, 22H), 1.75 (dd, J = 13.1, 7.4 Hz, 2H), 2.09 (dd, J = 13.1, 8.8 Hz, 2H), 2.21 (s, 2H), 2.78 (d, J = 11.1 Hz, 2H), 2.91 (d, J = 11.1 Hz, 2H), 3.49 (m, 8H), 3.88 (dd, J = 8.8, 7.4 Hz, 2H), 4.12 (m, 8H), 5.49 (s, 2H), 6.79 (m, 7H), 7.03 (s, 2H), 7.28 (d, J = 2.3 Hz, 2H), 7.42 (dd, J = 8.6, 2.2 Hz, 2H), 7.67 (d, J = 8.5 Hz, 2H)
63fr	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (m, 15H), 1.52 (m, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.09 (dd, J = 13.1, 8.8 Hz, 1H), 2.32 (d, J = 0.7 Hz, 1H), 2.43 (d, J = 0.8 Hz, 3H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.48 (m, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.2, 1.5 Hz, 2H), 5.43 (s, 1H), 6.63 (q, J = 6.8 Hz, 1H), 7.06 (s, 1H), 7.26 (m, 2H), 7.34 (q, J = 1.3 Hz, 1H), 7.42 (dd, J = 8.5, 2.3 Hz, 1H), 7.66 (d, J = 8.5 Hz, 1H)
63fs	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.2 Hz, 3H), 1.55 (m, 4H), 1.77 (dd, J = 13.1, 7.3 Hz, 1H), 2.12 (dd, J = 13.2, 8.8 Hz, 1H), 2.79 (d, J = 11.0 Hz, 1H), 2.92 (d, J = 11.0 Hz, 1H), 3.53 (m, 4H), 3.86 (dd, J = 8.7, 7.3 Hz, 1H), 4.19 (qd, J = 7.1, 1.7 Hz, 2H), 4.86 (s, 3H), 5.53 (s, 1H), 6.57 (q, J = 6.8 Hz, 1H), 7.34 (d, J = 2.2 Hz, 1H), 7.51 (dd, J = 8.5, 2.3 Hz, 1H), 7.72 (m, 2H), 7.84 (m, 2H), 7.92 (s, 1H)
63ft	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.88 (m, 2H), 1.24 (m, 4H), 1.39 (t, J = 7.0

	Hz, 3H), 1.50 (q, J = 7.7, 5.2 Hz, 4H), 1.74 (ddd, J = 13.3, 7.3, 1.8 Hz, 1H), 2.08 (ddd, J = 11.4, 8.7, 2.6 Hz, 1H), 2.39 (s, 3H), 2.76 (d, J = 11.1 Hz, 1H), 2.88 (dd, J = 11.0, 5.7 Hz, 1H), 3.58 (m, 4H), 3.85 (m, 1H), 4.13 (m, 4H), 5.73 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.70 (dt, J = 10.7, 2.2 Hz, 1H), 6.80 (p, J = 6.5 Hz, 1H), 7.01 (m, 2H), 7.63 (d, J = 1.9 Hz, 1H), 7.76 (m, 2H), 8.00 (d, J = 2.4 Hz, 1H)
63fu	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 4H), 1.50 (dt, J = 9.9, 5.2 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.1, 8.7 Hz, 1H), 2.22 (s, 1H), 2.29 (d, J = 10.3 Hz, 6H), 2.39 (s, 3H), 2.75 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.74 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.75 (m, 1H), 7.19 (d, J = 7.9 Hz, 1H), 7.39 (m, 2H), 7.59 (d, J = 1.8 Hz, 1H), 7.72 (m, 2H), 7.96 (d, J = 2.3 Hz, 1H)
63fv	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (m, 4H), 1.51 (dt, J = 10.6, 5.6 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.08 (dd, J = 13.1, 8.7 Hz, 1H), 2.39 (s, 3H), 2.76 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.53 (m, 4H), 3.83 (m, 1H), 4.18 (m, 2H), 4.85 (d, J = 10.8 Hz, 1H), 5.73 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.83 (q, J = 6.6 Hz, 1H), 7.60 (m, 4H), 7.79 (m, 2H), 8.00 (d, J = 2.4 Hz, 1H)
63fw	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.25 (t, J = 7.1 Hz, 3H), 1.46 (m, 7H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.06 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.73 (d, J = 11.0 Hz, 1H), 2.87 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.80 (dd, J = 8.7, 7.1 Hz, 1H), 4.16 (m, 4H), 5.73 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.6 Hz, 1H), 7.11 (d, J = 8.6 Hz, 1H), 7.57 (m, 2H), 7.71 (m, 3H), 7.98 (d, J = 2.4 Hz, 1H)
63fx	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (m, 4H), 1.39 (t, J = 7.0 Hz, 3H), 1.51 (dt, J = 10.6, 5.6 Hz, 4H), 1.73 (dd, J = 13.1, 7.1 Hz, 1H), 2.07 (dd, J = 13.0, 8.8 Hz, 1H), 2.39 (s, 3H), 2.74 (d, J = 10.9 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.53 (m, 4H), 3.81 (dd, J = 8.8, 7.1 Hz, 1H), 4.13 (m, 4H), 4.87 (s, 13H), 5.74 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.78 (q, J = 7.6, 7.0 Hz, 1H), 6.93 (m, 1H), 7.20 (m, 2H), 7.34 (t, J = 7.9 Hz, 1H), 7.62 (d, J = 1.8 Hz, 1H), 7.75 (m, 2H), 7.98 (d, J = 2.4 Hz, 1H)
63fy	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (m, 3H), 1.50 (dt, J = 10.5, 5.8 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.0, 8.8 Hz, 1H), 2.39 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.80 (m, 1H), 4.17 (qd, J = 7.1, 1.6 Hz, 2H), 5.73 (s, 1H), 6.42 (d, J = 2.3 Hz, 1H), 6.84 (q, J = 6.6 Hz, 1H), 6.99 (tt, J = 9.1, 2.4 Hz, 1H), 7.34 (m, 2H), 7.68 (d, J = 1.9 Hz, 1H), 7.79 (m, 2H), 8.01 (d, J = 2.4 Hz, 1H)
63fz	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (m, 3H), 1.51 (dt, J = 10.6, 5.6 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.40 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.52 (dq, J = 25.8, 8.1, 6.9 Hz, 4H), 3.81 (dd, J = 8.8, 7.1 Hz, 1H), 4.17 (qd, J = 7.1, 1.6 Hz, 2H), 5.73 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.83 (q, J = 6.6 Hz, 1H), 7.69 (m, 3H), 7.82 (m, 2H), 7.98 (m, 3H)
63ga	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 3H), 1.41 (t, J = 7.0 Hz, 3H), 1.53 (m, 4H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.11 (m, 1H), 2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.17 (m, 4H), 5.49 (s, 1H), 6.66 (q, J = 6.8 Hz, 1H), 7.02 (s, 1H), 7.27 (m, 3H), 7.45 (dd, J = 8.5, 2.3 Hz, 1H), 7.66 (d, J = 8.5 Hz, 1H)
63gb	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.88 (m, 2H), 1.17 (t, J = 7.0 Hz, 1H), 1.31 (m, 12H), 1.51 (dt, J = 11.3, 5.6 Hz, 4H), 1.74 (ddd, J = 13.1, 7.3, 1.9 Hz, 1H), 2.08 (ddd, J = 11.9, 8.8, 2.8 Hz, 1H), 2.39 (s, 3H), 2.76 (d, J = 11.0 Hz, 1H), 2.89 (dd, J =

	11.0, 5.9 Hz, 1H), 3.57 (m, 5H), 3.84 (dt, J = 8.6, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.7 Hz, 1H), 4.64 (p, J = 6.1 Hz, 1H), 5.74 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.5 Hz, 1H), 7.16 (t, J = 8.6 Hz, 1H), 7.44 (m, 2H), 7.61 (d, J = 1.9 Hz, 1H), 7.73 (m, 2H), 7.98 (d, J = 2.4 Hz, 1H)
63gc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (m, 4H), 1.51 (dt, J = 10.6, 5.7 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.28 (s, 1H), 2.37 (d, J = 16.9 Hz, 9H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.53 (dt, J = 22.1, 6.0 Hz, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.74 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.77 (q, J = 6.6 Hz, 1H), 7.03 (m, 1H), 7.27 (d, J = 1.5 Hz, 2H), 7.60 (d, J = 1.8 Hz, 1H), 7.74 (m, 2H), 7.97 (d, J = 2.4 Hz, 1H)
63gd	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (q, J = 7.1, 6.4 Hz, 4H), 1.50 (dt, J = 10.3, 5.6 Hz, 4H), 1.73 (dd, J = 13.0, 7.2 Hz, 1H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (d, J = 3.0 Hz, 6H), 2.74 (d, J = 10.9 Hz, 1H), 2.88 (d, J = 10.9 Hz, 1H), 3.52 (dt, J = 22.1, 6.1 Hz, 4H), 3.81 (dd, J = 8.8, 7.1 Hz, 1H), 4.17 (qd, J = 7.1, 1.6 Hz, 2H), 4.87 (s, 9H), 5.73 (s, 1H), 6.42 (d, J = 2.3 Hz, 1H), 6.81 (q, J = 6.6 Hz, 1H), 7.23 (s, 1H), 7.42 (s, 1H), 7.48 (d, J = 2.0 Hz, 1H), 7.62 (d, J = 1.9 Hz, 1H), 7.71 (dd, J = 8.1, 1.9 Hz, 1H), 7.79 (d, J = 8.2 Hz, 1H), 8.00 (d, J = 2.4 Hz, 1H)
63ge	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (t, J = 7.0 Hz, 1H), 1.26 (m, 5H), 1.50 (dt, J = 10.5, 5.4 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.29 (d, J = 1.9 Hz, 3H), 2.39 (s, 3H), 2.74 (d, J = 10.9 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.81 (dd, J = 8.8, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.73 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.79 (q, J = 6.6 Hz, 1H), 7.37 (m, 3H), 7.63 (d, J = 1.9 Hz, 1H), 7.76 (m, 2H), 7.98 (d, J = 2.4 Hz, 1H)
63gf	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.08 (m, 1H), 1.26 (m, 4H), 1.36 (s, 9H), 1.51 (dt, J = 10.5, 5.5 Hz, 5H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.08 (dd, J = 13.1, 8.8 Hz, 1H), 2.40 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.53 (dt, J = 23.0, 6.0 Hz, 4H), 3.81 (dd, J = 8.8, 7.1 Hz, 1H), 4.17 (qd, J = 7.1, 1.6 Hz, 2H), 5.74 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.76 (q, J = 6.6 Hz, 1H), 7.43 (m, 3H), 7.61 (d, J = 1.9 Hz, 1H), 7.73 (m, 3H), 7.99 (d, J = 2.4 Hz, 1H)
63gg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.3 Hz, 6H), 1.50 (dt, J = 10.5, 5.6 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.08 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.73 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.82 (q, J = 6.6 Hz, 1H), 7.41 (m, 2H), 7.61 (m, 2H), 7.74 (m, 3H), 8.00 (d, J = 2.4 Hz, 1H)
63gh	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.2 Hz, 3H), 1.51 (dt, J = 10.4, 5.5 Hz, 4H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.84 (m, 2H), 3.52 (ddq, J = 25.3, 13.2, 7.1, 5.7 Hz, 4H), 3.86 (dd, J = 8.8, 7.2 Hz, 1H), 4.18 (qd, J = 7.2, 1.6 Hz, 2H), 5.73 (s, 1H), 6.42 (d, J = 2.3 Hz, 1H), 6.82 (q, J = 6.6 Hz, 1H), 7.33 (t, J = 8.8 Hz, 1H), 7.63 (m, 2H), 7.73 (dd, J = 8.3, 1.9 Hz, 1H), 7.82 (m, 2H), 8.00 (d, J = 2.3 Hz, 1H)
63gi	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (m, 4H), 1.53 (dt, J = 10.2, 5.5 Hz, 4H), 1.82 (dd, J = 13.2, 7.7 Hz, 1H), 2.19 (m, 1H), 2.39 (s, 3H), 2.88 (d, J = 11.2 Hz, 1H), 2.99 (d, J = 11.2 Hz, 1H), 3.55 (m, 4H), 4.02 (t, J = 8.2 Hz, 1H), 4.22 (qd, J = 7.1, 1.9 Hz, 2H), 5.74 (s, 1H), 6.42 (d, J = 2.3 Hz, 1H), 6.82 (m, 1H), 7.35 (dt, J = 10.4, 8.4 Hz, 1H), 7.50 (ddt, J = 7.9, 3.8, 1.8 Hz, 1H), 7.72 (m, 4H), 8.00 (d, J = 2.4 Hz, 1H)

63gj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.31 (m, 3H), 1.65 (dt, J = 12.8, 6.2 Hz, 4H), 2.03 (m, 1H), 2.43 (m, 4H), 3.27 (s, 2H), 3.65 (m, 4H), 4.31 (qd, J = 7.1, 2.2 Hz, 2H), 4.58 (t, J = 8.8 Hz, 1H), 4.85 (m, 1H), 6.43 (d, J = 2.5 Hz, 1H), 6.89 (q, J = 6.4 Hz, 1H), 7.45 (m, 1H), 7.72 (d, J = 1.8 Hz, 1H), 7.81 (m, 2H), 8.00 (m, 3H)
63gk	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (t, J = 7.1 Hz, 3H), 1.64 (q, J = 10.3, 8.1 Hz, 4H), 2.04 (dd, J = 13.6, 8.9 Hz, 1H), 2.44 (m, 4H), 2.80 (s, 1H), 3.25 (m, 2H), 3.56 (m, 1H), 3.70 (d, J = 5.7 Hz, 2H), 4.32 (qd, J = 7.1, 2.5 Hz, 2H), 4.57 (t, J = 8.8 Hz, 1H), 6.44 (d, J = 2.4 Hz, 1H), 6.90 (q, J = 6.5 Hz, 1H), 7.76 (q, J = 1.5, 1.0 Hz, 2H), 7.85 (m, 2H), 7.95 (m, 1H), 8.04 (m, 2H)
63gl	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 4H), 1.51 (dt, J = 10.3, 5.5 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.08 (dd, J = 13.1, 8.8 Hz, 1H), 2.40 (s, 3H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.73 (s, 1H), 6.42 (d, J = 2.3 Hz, 1H), 6.82 (q, J = 6.6 Hz, 1H), 7.31 (ddt, J = 8.1, 2.3, 1.1 Hz, 1H), 7.57 (dd, J = 15.9, 7.9 Hz, 2H), 7.74 (m, 4H), 8.01 (d, J = 2.4 Hz, 1H)
63gm	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (d, J = 34.9 Hz, 12H), 1.51 (dt, J = 10.4, 5.7 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.06 (m, 1H), 2.39 (d, J = 3.3 Hz, 6H), 2.74 (d, J = 11.0 Hz, 2H), 2.89 (d, J = 11.0 Hz, 1H), 3.53 (m, 5H), 3.81 (dd, J = 8.7, 7.1 Hz, 1H), 4.17 (qd, J = 7.2, 1.6 Hz, 2H), 4.87 (s, 3H), 5.73 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.75 (q, J = 6.8 Hz, 1H), 7.28 (d, J = 9.0 Hz, 2H), 7.46 (t, J = 1.6 Hz, 1H), 7.59 (d, J = 1.8 Hz, 1H), 7.75 (m, 2H), 7.98 (d, J = 2.3 Hz, 1H)
63gn	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 4H), 1.50 (dt, J = 10.8, 5.6 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.41 (d, J = 11.8 Hz, 6H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.52 (dq, J = 24.2, 7.6, 6.3 Hz, 4H), 3.81 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.73 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.79 (m, 1H), 7.45 (m, 2H), 7.63 (dd, J = 7.6, 2.0 Hz, 2H), 7.75 (m, 2H), 7.98 (d, J = 2.4 Hz, 1H)
63go	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.88 (d, J = 7.9 Hz, 1H), 1.26 (t, J = 7.1 Hz, 7H), 1.50 (dt, J = 10.5, 5.6 Hz, 8H), 1.74 (dd, J = 13.1, 7.2 Hz, 2H), 2.07 (dd, J = 13.1, 8.8 Hz, 2H), 2.34 (t, J = 0.6 Hz, 1H), 2.40 (d, J = 9.3 Hz, 16H), 2.75 (d, J = 11.0 Hz, 2H), 2.89 (d, J = 11.0 Hz, 2H), 3.52 (m, 8H), 3.82 (dd, J = 8.8, 7.2 Hz, 2H), 4.18 (qd, J = 7.1, 1.6 Hz, 4H), 5.73 (s, 2H), 6.41 (d, J = 2.3 Hz, 2H), 6.79 (q, J = 6.6 Hz, 2H), 7.44 (d, J = 0.9 Hz, 4H), 7.62 (d, J = 1.8 Hz, 2H), 7.74 (m, 4H), 7.98 (d, J = 2.4 Hz, 2H)
63gp	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (q, J = 7.1, 5.7 Hz, 7H), 1.51 (dt, J = 10.7, 5.5 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.06 (td, J = 15.3, 14.1, 7.4 Hz, 1H), 2.32 (d, J = 2.1 Hz, 3H), 2.39 (s, 2H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.53 (dt, J = 22.2, 6.1 Hz, 4H), 3.83 (dd, J = 8.8, 7.2 Hz, 1H), 4.18 (m, 2H), 5.74 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.78 (q, J = 6.5 Hz, 1H), 7.10 (t, J = 9.0 Hz, 1H), 7.56 (m, 3H), 7.74 (m, 2H), 7.98 (d, J = 2.4 Hz, 1H)
63gq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (m, 1H), 1.26 (dq, J = 10.3, 5.8, 3.2 Hz, 6H), 1.46 (m, 7H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.09 (m, 1H), 2.39 (s, 3H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.53 (m, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.16 (m, 4H), 5.74 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.5 Hz, 1H), 7.15 (t, J = 8.6 Hz, 1H), 7.47 (m, 4H), 7.73 (m, 2H), 7.97 (d, J = 2.3 Hz, 1H)
63gr	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.52 (t, J = 8.0 Hz,

	5H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.08 (dd, J = 13.0, 8.7 Hz, 1H), 2.40 (s, 3H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 10.9 Hz, 1H), 3.25 (p, J = 1.7 Hz, 1H), 3.53 (m, 5H), 3.81 (dd, J = 8.8, 7.1 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 4.82 (s, 1H), 5.73 (s, 1H), 6.00 (m, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.85 (d, J = 6.6 Hz, 1H), 7.48 (t, J = 1.9 Hz, 1H), 7.69 (m, 3H), 7.80 (m, 3H), 8.03 (d, J = 2.4 Hz, 1H)
63gs	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (m, 10H), 1.50 (dt, J = 10.3, 5.6 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.93 (m, 2H), 3.52 (m, 5H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.17 (qd, J = 7.1, 1.5 Hz, 2H), 5.74 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.6 Hz, 1H), 7.26 (dt, J = 7.6, 1.4 Hz, 1H), 7.36 (t, J = 7.7 Hz, 1H), 7.47 (m, 2H), 7.61 (d, J = 1.9 Hz, 1H), 7.75 (m, 2H), 7.98 (d, J = 2.4 Hz, 1H)
63gt	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (m, 4H), 1.51 (dt, J = 11.0, 5.6 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.2, 8.8 Hz, 1H), 2.40 (s, 2H), 2.75 (d, J = 10.9 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.53 (m, 4H), 3.82 (dd, J = 8.8, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.72 (s, 1H), 6.00 (m, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.84 (q, J = 6.5 Hz, 1H), 7.71 (m, 2H), 7.82 (m, 2H), 7.92 (dd, J = 8.4, 2.3 Hz, 1H), 8.05 (dd, J = 15.9, 2.3 Hz, 2H)
63gu	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.51 (dt, J = 11.1, 5.5 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.40 (s, 3H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.53 (dt, J = 22.4, 6.0 Hz, 4H), 3.81 (dd, J = 8.7, 7.1 Hz, 1H), 4.18 (qd, J = 7.1, 1.7 Hz, 2H), 5.73 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.86 (q, J = 6.6 Hz, 1H), 7.81 (m, 5H), 7.97 (m, 1H), 8.03 (d, J = 2.4 Hz, 1H)
63gv	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.08 (m, 1H), 1.26 (m, 4H), 1.51 (dt, J = 10.7, 5.5 Hz, 5H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.06 (m, 1H), 2.40 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.26 (s, 1H), 3.53 (dt, J = 22.3, 6.1 Hz, 5H), 3.81 (dd, J = 8.7, 7.1 Hz, 1H), 4.17 (qd, J = 7.1, 1.6 Hz, 2H), 5.74 (s, 1H), 6.42 (dd, J = 2.4, 0.6 Hz, 1H), 6.83 (q, J = 6.6 Hz, 1H), 7.57 (dd, J = 8.1, 7.5 Hz, 1H), 7.74 (t, J = 1.1 Hz, 1H), 7.87 (m, 4H), 8.00 (d, J = 2.4 Hz, 1H), 8.19 (m, 1H)
63gw	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.52 (dt, J = 10.3, 5.5 Hz, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.09 (dd, J = 13.1, 8.8 Hz, 1H), 2.41 (s, 3H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.53 (dt, J = 23.4, 6.0 Hz, 4H), 3.83 (dd, J = 8.8, 7.1 Hz, 1H), 4.18 (qd, J = 7.2, 1.7 Hz, 2H), 5.72 (s, 1H), 6.44 (d, J = 2.4 Hz, 1H), 6.87 (q, J = 6.6 Hz, 1H), 7.80 (d, J = 1.6 Hz, 1H), 7.89 (m, 2H), 8.00 (s, 1H), 8.07 (d, J = 2.4 Hz, 1H), 8.30 (d, J = 1.5 Hz, 2H)
63gx	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (m, 10H), 1.50 (dt, J = 10.2, 5.5 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.06 (m, 1H), 2.39 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 10.9 Hz, 1H), 3.53 (m, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.17 (qd, J = 7.2, 1.6 Hz, 2H), 4.65 (h, J = 5.9 Hz, 1H), 5.73 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.6 Hz, 1H), 6.92 (m, 1H), 7.19 (m, 2H), 7.33 (t, J = 7.9 Hz, 1H), 7.60 (d, J = 1.8 Hz, 1H), 7.74 (m, 2H), 7.98 (d, J = 2.3 Hz, 1H)
63gy	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.2 Hz, 4H), 1.46 (m, 7H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.09 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.52 (dq, J = 24.9, 7.0, 6.0 Hz, 4H), 3.84 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (ttd, J = 7.0, 5.2, 2.5 Hz, 4H), 5.74 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.5 Hz, 1H), 7.18 (m, 2H), 7.35 (dd, J = 8.0, 2.1 Hz, 1H), 7.62

	(d, J = 1.8 Hz, 1H), 7.74 (m, 2H), 7.99 (d, J = 2.3 Hz, 1H)
63gz	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (m, 9H), 1.51 (dt, J = 10.4, 5.6 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.0, 8.7 Hz, 1H), 2.39 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.53 (m, 4H), 3.81 (dd, J = 8.7, 7.1 Hz, 1H), 4.17 (qd, J = 7.1, 1.6 Hz, 2H), 4.66 (h, J = 6.1 Hz, 1H), 5.73 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.70 (dt, J = 10.8, 2.2 Hz, 1H), 6.81 (q, J = 6.6 Hz, 1H), 6.99 (m, 2H), 7.62 (d, J = 1.8 Hz, 1H), 7.72 (dd, J = 8.3, 1.9 Hz, 1H), 7.79 (d, J = 8.3 Hz, 1H), 8.00 (d, J = 2.3 Hz, 1H)
63ha	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 - 1.40 (m, 3 H), 1.55 - 1.76 (m, 4 H), 2.06 (br. s., 1 H), 2.35 - 2.54 (m, 4 H), 3.29 (s, 2 H) 3.50- 3.78 (m, 4H), 3.85 (s, 3H), 4.34 (dd, J=7.03, 2.34 Hz, 2 H), 4.60 (s, 1 H), 5.96 (s, 1 H), 6.44 (d, J=2.15 Hz, 1 H), 6.81 (d, J=6.44 Hz, 1 H), 7.03 (d, J=8.79 Hz, 2 H), 7.50 - 7.68 (m, 3 H), 7.70 - 7.82 (m, 2 H), 7.97 (d, J=2.15 Hz, 1 H)
63hb	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.34 (t, J=7.13 Hz, 3 H), 1.43 (t, J=7.03 Hz, 3 H), 1.63 - 1.82 (m, 4 H), 2.01 - 2.14 (m, 1 H), 2.43 (s, 3 H) 2.46 - 2.57 (m, 1 H), 3.31 (br. s., 2 H), 3.59 - 3.93 (m, 4 H), 4.11 (d, J=7.03 Hz, 2 H), 4.26 - 4.41 (m, 2 H), 4.56 - 4.68 (m, 1 H), 6.44 (d, J=2.15 Hz, 1 H), 6.76 - 6.93 (m, 1 H), 7.04 (d, J=8.79 Hz, 2 H), 7.66 (dd, J=5.17, 3.61 Hz, 3 H), 7.72 - 7.85 (m, 2 H), 7.93 - 8.02 (m, 1 H)
63hc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.2 Hz, 3H), 1.51 (dt, J = 11.0, 5.6 Hz, 4H), 1.74 (dd, J = 13.1, 7.3 Hz, 1H), 2.09 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.84 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 4.88 (s, 7H), 5.72 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.84 (q, J = 6.5 Hz, 1H), 7.52 (m, 2H), 7.67 (d, J = 1.9 Hz, 1H), 7.74 (dd, J = 8.3, 1.9 Hz, 1H), 7.81 (d, J = 8.3 Hz, 1H), 8.01 (d, J = 2.4 Hz, 1H)
63hd	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (m, 4H), 1.54 (dt, J = 8.1, 4.7 Hz, 4H), 1.76 (dd, J = 13.1, 7.3 Hz, 1H), 2.11 (dd, J = 13.1, 8.8 Hz, 1H), 2.77 (d, J = 10.9 Hz, 1H), 2.91 (d, J = 10.9 Hz, 1H), 3.53 (dq, J = 16.3, 6.7, 6.2 Hz, 4H), 3.65 (s, 3H), 3.83 (dd, J = 8.8, 7.2 Hz, 1H), 4.19 (qd, J = 7.2, 1.6 Hz, 2H), 5.51 (d, J = 18.6 Hz, 1H), 6.49 (dd, J = 6.9, 2.0 Hz, 1H), 6.81 (q, J = 6.8 Hz, 1H), 6.91 (s, 1H), 7.35 (d, J = 2.2 Hz, 1H), 7.51 (dd, J = 8.5, 2.2 Hz, 1H), 7.70 (d, J = 8.6 Hz, 1H), 7.80 (d, J = 6.8 Hz, 1H)
63he	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 4H), 1.51 (dt, J = 10.8, 5.6 Hz, 5H), 1.74 (dd, J = 13.1, 7.1 Hz, 1H), 2.07 (dd, J = 13.0, 8.7 Hz, 1H), 2.38 (d, J = 9.6 Hz, 6H), 2.74 (d, J = 10.9 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.53 (dt, J = 21.9, 6.3 Hz, 5H), 3.81 (dd, J = 8.7, 7.1 Hz, 1H), 4.18 (qd, J = 7.1, 1.7 Hz, 2H), 4.88 (s, 15H), 5.74 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.81 (q, J = 6.6 Hz, 1H), 7.33 (m, 1H), 7.58 (m, 1H), 7.66 (t, J = 2.6 Hz, 2H), 7.77 (m, 2H), 7.99 (d, J = 2.4 Hz, 1H)
63hf	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.08 (m, 1H), 1.26 (t, J = 7.1 Hz, 4H), 1.51 (dt, J = 10.6, 5.6 Hz, 4H), 1.76 (dd, J = 13.2, 7.4 Hz, 1H), 2.11 (dd, J = 13.2, 8.8 Hz, 1H), 2.39 (s, 7H), 2.79 (d, J = 11.0 Hz, 1H), 2.92 (d, J = 11.1 Hz, 1H), 3.52 (dq, J = 29.5, 7.4, 6.4 Hz, 5H), 3.89 (dd, J = 8.7, 7.4 Hz, 1H), 4.19 (qd, J = 7.2, 1.7 Hz, 2H), 5.73 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.80 (q, J = 6.6 Hz, 1H), 7.37 (d, J = 7.9 Hz, 1H), 7.51 (dd, J = 7.9, 1.9 Hz, 1H), 7.62 (d, J = 1.9 Hz, 1H), 7.73 (m, 3H), 7.99 (d, J = 2.4 Hz, 1H)
63hg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.51 (dt, J = 10.5, 5.6 Hz, 4H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.40 (s, 3H),

	2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.53 (m, 4H), 3.85 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.7 Hz, 2H), 5.73 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.86 (q, J = 6.6 Hz, 1H), 7.50 (dd, J = 8.4, 2.1 Hz, 1H), 7.79 (m, 6H), 8.04 (d, J = 2.4 Hz, 1H)
63hh	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.93 (m, 2H), 1.36 (m, 11H), 1.69 (td, J = 13.3, 6.6 Hz, 1H), 2.04 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.72 (d, J = 11.0 Hz, 1H), 2.86 (d, J = 11.0 Hz, 1H), 3.50 (m, 4H), 3.80 (t, J = 7.9 Hz, 1H), 4.17 (m, 2H), 5.71 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.85 (q, J = 6.6 Hz, 1H), 7.20 (dt, J = 8.5, 2.1 Hz, 1H), 7.39 (dt, J = 9.7, 2.0 Hz, 1H), 7.52 (t, J = 1.6 Hz, 1H), 7.67 (m, 2H), 7.80 (d, J = 8.3 Hz, 1H), 8.01 (d, J = 2.3 Hz, 1H)
63hi	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.75 (m, 2H), 1.01 (dq, J = 8.4, 2.4 Hz, 2H), 1.19 (s, 1H), 1.26 (t, J = 7.1 Hz, 4H), 1.52 (m, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.02 (m, 2H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.49 (m, 4H), 3.82 (dd, J = 8.8, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.5 Hz, 2H), 5.44 (s, 1H), 6.60 (q, J = 6.9 Hz, 1H), 7.20 (m, 4H), 7.41 (m, 2H), 7.66 (d, J = 8.5 Hz, 1H)
63hj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.25 (t, J = 7.1 Hz, 3H), 1.35 (d, J = 6.0 Hz, 6H), 1.49 (ddd, J = 12.1, 7.6, 4.8 Hz, 4H), 1.72 (dd, J = 13.1, 7.2 Hz, 1H), 2.05 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.72 (d, J = 11.0 Hz, 1H), 2.87 (d, J = 11.0 Hz, 1H), 3.51 (m, 4H), 3.80 (dd, J = 8.7, 7.1 Hz, 1H), 4.17 (qd, J = 7.1, 1.6 Hz, 2H), 4.67 (hept, J = 6.1 Hz, 1H), 5.73 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.78 (q, J = 6.6 Hz, 1H), 7.12 (d, J = 8.6 Hz, 1H), 7.54 (m, 2H), 7.67 (m, 2H), 7.75 (d, J = 8.3 Hz, 1H), 7.98 (d, J = 2.4 Hz, 1H)
63hk	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (m, 4H), 1.54 (m, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.11 (dd, J = 13.1, 8.8 Hz, 1H), 2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.52 (dq, J = 27.4, 7.7, 6.5 Hz, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.1, 1.6 Hz, 2H), 5.51 (d, J = 14.8 Hz, 2H), 6.79 (q, J = 6.8 Hz, 1H), 7.40 (d, J = 2.2 Hz, 1H), 7.52 (m, 2H), 7.73 (d, J = 8.5 Hz, 1H), 8.23 (d, J = 8.3 Hz, 1H), 8.47 (s, 1H), 9.34 (s, 1H)
63hl	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.25 (t, J = 7.1 Hz, 3H), 1.50 (dt, J = 7.8, 4.8 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.1, 8.7 Hz, 1H), 2.74 (d, J = 10.9 Hz, 1H), 2.88 (d, J = 10.9 Hz, 1H), 3.10 (s, 6H), 3.49 (m, 4H), 3.81 (dd, J = 8.7, 7.2 Hz, 1H), 4.17 (qd, J = 7.1, 1.3 Hz, 2H), 5.47 (s, 1H), 6.69 (m, 3H), 7.32 (d, J = 2.2 Hz, 1H), 7.45 (dd, J = 8.5, 2.2 Hz, 1H), 7.68 (d, J = 8.5 Hz, 1H), 8.18 (d, J = 5.2 Hz, 1H)
63hm	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.24 (t, J = 7.1 Hz, 3H), 1.45 (dt, J = 9.4, 5.7 Hz, 4H), 1.70 (dd, J = 13.1, 7.2 Hz, 1H), 2.03 (dd, J = 13.1, 8.7 Hz, 1H), 2.70 (d, J = 11.0 Hz, 1H), 2.85 (d, J = 11.0 Hz, 1H), 3.40 (m, 4H), 3.79 (t, J = 7.9 Hz, 1H), 4.16 (q, J = 7.1 Hz, 2H), 5.45 (d, J = 17.0 Hz, 1H), 6.68 (q, J = 6.8 Hz, 1H), 7.37 (d, J = 2.3 Hz, 1H), 7.44 (dd, J = 8.6, 2.2 Hz, 1H), 7.56 (m, 3H), 7.71 (d, J = 8.5 Hz, 1H), 7.94 (m, 4H)
63hn	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 3H), 1.36 (s, 9H), 1.52 (m, 4H), 1.77 (dd, J = 13.1, 7.4 Hz, 1H), 2.13 (dd, J = 13.1, 8.7 Hz, 1H), 2.80 (d, J = 11.1 Hz, 1H), 2.93 (d, J = 11.1 Hz, 1H), 3.48 (m, 4H), 3.91 (dd, J = 8.7, 7.4 Hz, 1H), 4.20 (qd, J = 7.2, 1.7 Hz, 2H), 5.41 (s, 1H), 6.64 (q, J = 6.9 Hz, 1H), 7.27 (m, 2H), 7.44 (m, 5H), 7.69 (d, J = 7.3 Hz, 1H)
63ho	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (td, J = 7.1, 2.3 Hz, 3H), 1.53 (dt, J =

	13.2, 6.2 Hz, 4H), 1.74 (dt, J = 13.6, 7.0 Hz, 1H), 2.09 (m, 1H), 2.75 (dd, J = 10.9, 7.2 Hz, 1H), 2.90 (dd, J = 11.0, 6.6 Hz, 1H), 3.29 (s, 1H), 3.52 (m, 4H), 3.83 (td, J = 8.2, 4.3 Hz, 1H), 4.18 (q, J = 7.1 Hz, 2H), 5.47 (m, 1H), 6.43 (dt, J = 11.2, 5.6 Hz, 1H), 7.22 (m, 1H), 7.38 (m, 2H), 7.61 (dd, J = 5.0, 2.2 Hz, 1H), 7.81 (m, 3H)
63hp	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 4H), 1.51 (dt, J = 10.0, 5.2 Hz, 4H), 1.80 (m, 1H), 2.17 (dd, J = 13.3, 8.8 Hz, 1H), 2.86 (d, J = 11.2 Hz, 1H), 2.97 (d, J = 11.2 Hz, 1H), 3.50 (m, 4H), 4.01 (t, J = 8.2 Hz, 1H), 4.21 (qd, J = 7.1, 1.9 Hz, 2H), 5.63 (s, 1H), 6.69 (q, J = 6.6 Hz, 1H), 7.30 (ddd, J = 7.9, 6.9, 0.9 Hz, 1H), 7.46 (m, 2H), 7.75 (m, 4H), 8.39 (d, J = 1.0 Hz, 1H)
63hq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (t, J = 7.1 Hz, 3H), 1.36 (dd, J = 6.9, 3.7 Hz, 6H), 1.50 (m, 2H), 1.73 (dd, J = 13.1, 6.7 Hz, 1H), 2.05 (dd, J = 13.1, 8.8 Hz, 1H), 2.81 (d, J = 10.5 Hz, 1H), 2.94 (d, J = 10.5 Hz, 1H), 3.14 (p, J = 6.9 Hz, 1H), 3.47 (dt, J = 12.2, 5.6 Hz, 4H), 3.85 (dd, J = 8.8, 6.7 Hz, 1H), 4.19 (q, J = 7.1 Hz, 2H), 4.34 (s, 2H), 5.42 (s, 1H), 6.53 (q, J = 6.7 Hz, 1H), 7.25 (m, 3H), 7.42 (dd, J = 8.5, 2.2 Hz, 1H), 7.68 (d, J = 8.5 Hz, 1H), 8.65 (dd, J = 5.0, 0.8 Hz, 1H)
63hr	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 3H), 1.53 (m, 4H), 1.76 (dd, J = 13.1, 7.3 Hz, 1H), 2.11 (dd, J = 13.1, 8.7 Hz, 1H), 2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.51 (dq, J = 17.7, 6.1 Hz, 4H), 3.84 (dd, J = 8.7, 7.3 Hz, 1H), 4.19 (qd, J = 7.1, 1.6 Hz, 2H), 5.50 (s, 1H), 6.60 (q, J = 6.7 Hz, 1H), 7.28 (m, 3H), 7.48 (ddd, J = 25.4, 8.2, 3.7 Hz, 3H), 7.66 (d, J = 8.5 Hz, 1H)
63hs	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.52 (m, 4H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.50 (dq, J = 25.8, 7.7, 6.9 Hz, 4H), 3.85 (dd, J = 8.7, 7.3 Hz, 1H), 4.19 (qd, J = 7.1, 1.5 Hz, 2H), 5.50 (s, 1H), 6.60 (q, J = 6.7 Hz, 1H), 7.28 (d, J = 2.2 Hz, 1H), 7.52 (m, 6H)
63ht	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (t, J = 7.1 Hz, 3H), 1.56 (dt, J = 11.3, 5.3 Hz, 4H), 1.86 (dd, J = 13.3, 7.9 Hz, 1H), 2.25 (dd, J = 13.3, 8.7 Hz, 1H), 2.44 (s, 3H), 2.96 (d, J = 11.4 Hz, 1H), 3.05 (d, J = 11.3 Hz, 1H), 3.54 (m, 3H), 3.75 (s, 1H), 4.13 (t, J = 8.3 Hz, 1H), 4.24 (qd, J = 7.2, 2.0 Hz, 2H), 5.48 (s, 1H), 6.64 (q, J = 6.8 Hz, 1H), 7.26 (d, J = 2.3 Hz, 1H), 7.35 (s, 4H), 7.42 (dd, J = 8.5, 2.3 Hz, 1H), 7.65 (d, J = 8.5 Hz, 1H)
63hu	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.24 (m, 3H), 1.49 (dt, J = 10.2, 5.7 Hz, 4H), 1.72 (m, 1H), 2.04 (m, 1H), 2.41 (s, 2H), 2.72 (d, J = 10.9 Hz, 1H), 2.86 (d, J = 10.9 Hz, 1H), 3.52 (m, 4H), 3.79 (dd, J = 8.8, 7.1 Hz, 1H), 4.16 (qd, J = 7.1, 1.6 Hz, 2H), 5.76 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.82 (q, J = 6.7 Hz, 1H), 7.49 (m, 2H), 7.84 (m, 7H), 8.01 (d, J = 2.4 Hz, 1H), 8.14 (d, J = 1.9 Hz, 1H)
63hv	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.53 (m, 4H), 1.78 (m, 5H), 2.11 (m, 2H), 2.25 (dt, J = 7.9, 4.0 Hz, 2H), 2.41 (d, J = 18.1 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.50 (dq, J = 24.8, 7.6, 6.8 Hz, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.5 Hz, 2H), 5.48 (s, 1H), 5.76 (h, J = 2.0 Hz, 1H), 6.93 (q, J = 6.9 Hz, 1H), 7.15 (d, J = 2.3 Hz, 1H), 7.30 (dd, J = 8.5, 2.3 Hz, 1H), 7.58 (d, J = 8.5 Hz, 1H)
63hw	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 3H), 1.52 (dt, J = 8.9, 5.7 Hz, 4H), 1.81 (dd, J = 13.3, 7.7 Hz, 1H), 2.17 (dd, J = 13.3, 8.8 Hz, 1H), 2.39 (s, 3H), 2.88 (d, J = 11.3 Hz, 1H), 2.98 (d, J = 11.3 Hz, 1H), 3.53 (m, 4H), 4.04 (t, J = 8.2 Hz,

	1H), 4.21 (qd, J = 7.2, 1.8 Hz, 2H), 4.90 (d, J = 1.1 Hz, 5H), 5.74 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.6 Hz, 1H), 7.17 (t, J = 8.6 Hz, 1H), 7.40 (m, 7H), 7.62 (m, 2H), 7.74 (d, J = 8.2 Hz, 1H), 7.96 (d, J = 2.3 Hz, 1H)
63hx	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (m, 9H), 1.55 (dt, J = 10.7, 5.8 Hz, 4H), 1.87 (dd, J = 13.3, 8.0 Hz, 1H), 2.24 (m, 4H), 2.39 (s, 3H), 2.98 (d, J = 11.4 Hz, 1H), 3.06 (d, J = 11.4 Hz, 1H), 3.58 (m, 4H), 4.22 (m, 3H), 4.63 (p, J = 6.0 Hz, 1H), 5.76 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.75 (q, J = 6.6 Hz, 1H), 6.97 (d, J = 8.2 Hz, 1H), 7.45 (d, J = 8.1 Hz, 2H), 7.58 (d, J = 1.7 Hz, 1H), 7.71 (m, 2H), 7.96 (d, J = 2.3 Hz, 1H)
63hy	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.04 (m, 7H), 1.26 (t, J = 7.1 Hz, 4H), 1.52 (m, 4H), 1.74 (dd, J = 13.1, 7.3 Hz, 1H), 2.08 (m, 2H), 2.75 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.50 (m, 4H), 3.80 (m, 3H), 4.18 (qd, J = 7.1, 1.4 Hz, 2H), 5.48 (s, 1H), 6.70 (q, J = 6.9 Hz, 1H), 7.02 (m, 2H), 7.19 (s, 2H), 7.28 (d, J = 2.3 Hz, 1H), 7.42 (m, 2H), 7.66 (d, J = 8.5 Hz, 1H)
63hz	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.33 (t, J = 6.3 Hz, 9H), 1.68 (m, 4H), 2.04 (m, 1H), 2.50 (dd, J = 13.6, 8.7 Hz, 1H), 3.28 (s, 2H), 3.56 (m, 5H), 4.32 (qd, J = 7.2, 2.2 Hz, 2H), 4.62 (m, 2H), 6.59 (m, 1H), 6.97 (m, 2H), 7.53 (m, 9H), 7.66 (dd, J = 8.3, 2.0 Hz, 1H), 7.75 (d, J = 8.4 Hz, 1H)
63ia	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.52 (dt, J = 9.7, 5.5 Hz, 4H), 1.74 (dd, J = 13.0, 7.3 Hz, 1H), 2.09 (dd, J = 13.1, 8.8 Hz, 1H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.51 (dq, J = 23.9, 7.7, 6.6 Hz, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.5 Hz, 2H), 5.56 (s, 1H), 6.68 (q, J = 7.2 Hz, 1H), 7.67 (d, J = 8.1 Hz, 2H), 7.82 (m, 2H), 8.03 (dd, J = 8.9, 2.0 Hz, 1H), 8.15 (m, 3H), 8.81 (dd, J = 2.9, 0.9 Hz, 1H)
63ib	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.06 (t, J = 7.4 Hz, 3H), 1.24 (m, 3H), 1.50 (dt, J = 10.7, 5.6 Hz, 4H), 1.79 (m, 3H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.74 (d, J = 10.9 Hz, 1H), 2.87 (m, 1H), 3.55 (m, 5H), 3.81 (dd, J = 8.8, 7.2 Hz, 1H), 4.04 (t, J = 6.4 Hz, 2H), 4.17 (qd, J = 7.2, 1.7 Hz, 2H), 5.75 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.78 (q, J = 6.6 Hz, 1H), 7.15 (t, J = 8.6 Hz, 1H), 7.45 (m, 2H), 7.60 (d, J = 1.8 Hz, 1H), 7.73 (m, 2H), 7.98 (d, J = 2.4 Hz, 1H)
63ic	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.00 (t, J = 7.4 Hz, 3H), 1.26 (t, J = 7.1 Hz, 3H), 1.51 (m, 6H), 1.76 (m, 3H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.53 (qd, J = 13.9, 7.7 Hz, 4H), 3.81 (dd, J = 8.7, 7.2 Hz, 1H), 4.14 (m, 4H), 5.74 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.78 (q, J = 6.6 Hz, 1H), 7.15 (m, 1H), 7.46 (m, 2H), 7.61 (d, J = 1.7 Hz, 1H), 7.73 (m, 2H), 7.98 (d, J = 2.4 Hz, 1H)
63id	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.25 (t, J = 7.1 Hz, 3H), 1.51 (m, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (dd, J = 13.0, 8.8 Hz, 1H), 2.40 (s, 3H), 2.64 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.81 (dd, J = 8.7, 7.2 Hz, 1H), 4.17 (qd, J = 7.1, 1.6 Hz, 2H), 5.74 (s, 1H), 6.44 (d, J = 2.3 Hz, 1H), 6.86 (q, J = 6.6 Hz, 1H), 7.76 (m, 3H), 7.85 (d, J = 1.2 Hz, 2H), 8.08 (m, 2H)
63ie	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (d, J = 6.7 Hz, 2H), 1.27 (t, J = 7.1 Hz, 3H), 1.53 (m, 4H), 1.72 (ddd, J = 23.9, 13.2, 7.0 Hz, 1H), 1.94 (m, 4H), 2.11 (dd, J = 13.1, 8.8 Hz, 1H), 2.77 (d, J = 10.9 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.55 (m, 9H), 3.85 (dd, J = 8.7, 7.3 Hz, 1H), 4.19 (qd, J = 7.1, 1.7 Hz, 2H), 5.53 (s, 1H), 6.69 (q, J = 6.7 Hz, 1H), 7.31 (d, J = 2.2 Hz, 1H), 7.48 (m, 2H), 7.64 (m, 3H), 7.93 (s, 1H)

63if	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (td, J = 7.1, 0.8 Hz, 4H), 1.53 (m, 4H), 1.76 (m, 9H), 1.97 (dd, J = 13.4, 6.8 Hz, 2H), 2.11 (dd, J = 13.1, 8.8 Hz, 1H), 2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 10.9 Hz, 1H), 3.31 (m, 3H), 3.51 (dq, J = 19.6, 6.3 Hz, 4H), 3.84 (dd, J = 8.7, 7.3 Hz, 1H), 4.19 (m, 2H), 5.48 (s, 1H), 6.71 (q, J = 6.9 Hz, 1H), 6.94 (d, J = 7.6 Hz, 1H), 7.02 (dd, J = 8.4, 2.6 Hz, 1H), 7.18 (s, 1H), 7.28 (d, J = 2.2 Hz, 1H), 7.43 (m, 2H), 7.66 (d, J = 8.5 Hz, 1H)
63ig	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.89 (dd, J = 6.7, 0.7 Hz, 1H), 1.27 (td, J = 7.1, 0.7 Hz, 3H), 1.53 (m, 4H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.11 (m, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.30 (dq, J = 3.5, 1.8 Hz, 5H), 3.54 (m, 10H), 3.82 (m, 5H), 4.18 (qd, J = 7.2, 1.6 Hz, 2H), 5.53 (s, 1H), 6.70 (q, J = 6.7 Hz, 1H), 6.84 (m, 1H), 7.30 (m, 1H), 7.51 (m, 3H), 7.66 (m, 2H), 7.79 (s, 1H)
63ih	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.39 (m, 12H), 1.84 (dd, J = 13.2, 7.8 Hz, 1H), 1.98 (m, 5H), 2.24 (m, 1H), 2.92 (d, J = 11.3 Hz, 1H), 3.02 (d, J = 11.2 Hz, 1H), 3.54 (ddq, J = 27.6, 15.0, 7.8, 7.4 Hz, 6H), 3.89 (s, 1H), 4.07 (t, J = 8.2 Hz, 1H), 4.23 (qd, J = 7.1, 2.0 Hz, 2H), 4.93 (d, J = 1.4 Hz, 11H), 5.55 (s, 1H), 6.63 (q, J = 6.7 Hz, 1H), 7.30 (d, J = 2.2 Hz, 1H), 7.46 (dd, J = 8.5, 2.3 Hz, 1H), 7.63 (m, 3H), 7.89 (dt, J = 7.7, 1.6 Hz, 1H), 8.35 (s, 1H)
63ii	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (td, J = 7.4, 5.4 Hz, 7H), 1.52 (dt, J = 7.6, 4.7 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.09 (dd, J = 13.1, 8.7 Hz, 1H), 2.74 (m, 3H), 2.90 (d, J = 11.0 Hz, 1H), 3.30 (d, J = 9.9 Hz, 1H), 3.49 (m, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.5 Hz, 2H), 5.46 (s, 1H), 6.64 (q, J = 6.8 Hz, 1H), 7.30 (m, 4H), 7.43 (m, 2H), 7.67 (d, J = 8.5 Hz, 1H)
63ij	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (dd, J = 6.9, 5.1 Hz, 6H), 1.59 (d, J = 5.6 Hz, 4H), 2.03 (dd, J = 13.4, 7.2 Hz, 1H), 2.30 (dd, J = 13.4, 9.2 Hz, 1H), 3.03 (m, 2H), 3.22 (d, J = 11.7 Hz, 1H), 3.46 (tt, J = 16.4, 7.0 Hz, 2H), 3.62 (q, J = 8.5 Hz, 2H), 4.05 (dd, J = 9.1, 7.1 Hz, 1H), 5.48 (s, 1H), 6.62 (q, J = 6.7 Hz, 1H), 7.31 (m, 4H), 7.44 (m, 2H), 7.66 (d, J = 8.5 Hz, 1H)
63ik	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.05 (t, J = 7.4 Hz, 3H), 1.26 (td, J = 7.2, 0.6 Hz, 4H), 1.51 (dt, J = 10.0, 5.7 Hz, 4H), 1.77 (ddd, J = 27.5, 13.6, 7.1 Hz, 3H), 2.08 (dd, J = 13.1, 8.7 Hz, 1H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.30 (p, J = 1.6 Hz, 5H), 3.51 (m, 4H), 3.81 (dd, J = 8.7, 7.2 Hz, 1H), 3.95 (t, J = 6.5 Hz, 2H), 4.18 (qd, J = 7.1, 1.5 Hz, 2H), 5.53 (s, 1H), 6.61 (q, J = 7.2 Hz, 1H), 6.97 (m, 2H), 7.56 (m, 6H)
63il	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 4H), 1.46 (m, 8H), 1.73 (dd, J = 13.0, 7.3 Hz, 1H), 2.08 (m, 1H), 2.74 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.02 (m, 2H), 3.50 (dd, J = 17.6, 11.2 Hz, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.2, 1.5 Hz, 2H), 5.53 (s, 1H), 6.99 (q, J = 6.9 Hz, 1H), 7.67 (m, 3H), 8.03 (dd, J = 8.9, 2.1 Hz, 1H), 8.14 (m, 3H), 8.80 (dd, J = 2.8, 0.9 Hz, 1H)
63im	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 5H), 1.53 (td, J = 7.3, 6.9, 4.5 Hz, 4H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.10 (m, 1H), 2.29 (s, 3H), 2.39 (s, 3H), 2.53 (s, 2H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.52 (m, 6H), 3.83 (m, 3H), 4.18 (qd, J = 7.1, 1.7 Hz, 2H), 5.51 (d, J = 15.3 Hz, 1H), 6.71 (q, J = 6.6 Hz, 1H), 7.32 (d, J = 2.3 Hz, 1H), 7.50 (m, 3H), 7.66 (m, 2H), 7.80 (s, 1H)
63in	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.25 (m, 4H), 1.54 (m, 4H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.10 (dd, J = 12.8, 8.5 Hz, 1H), 2.66 (s, 3H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.52 (dd, J = 14.8, 8.9 Hz, 5H), 3.83 (dd, J = 8.7, 7.3

	Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.54 (s, 1H), 6.85 (t, J = 7.0 Hz, 1H), 7.65 (d, J = 2.9 Hz, 3H), 8.04 (dd, J = 8.9, 2.0 Hz, 1H), 8.15 (m, 3H), 8.81 (d, J = 2.8 Hz, 1H)
63io	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.15 (t, J = 7.0 Hz, 3H), 1.26 (t, J = 7.1 Hz, 6H), 1.53 (dt, J = 10.1, 5.5 Hz, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.7 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.36 (d, J = 7.7 Hz, 1H), 3.54 (m, 6H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.2, 1.6 Hz, 2H), 5.55 (s, 1H), 6.65 (q, J = 7.1 Hz, 1H), 7.46 (m, 2H), 7.62 (d, J = 8.1 Hz, 2H), 7.72 (m, 4H)
63ip	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (td, J = 7.1, 1.0 Hz, 7H), 1.53 (m, 8H), 1.74 (dd, J = 13.1, 7.2 Hz, 2H), 2.09 (dd, J = 13.1, 8.7 Hz, 2H), 2.75 (d, J = 11.0 Hz, 2H), 2.89 (d, J = 11.0 Hz, 2H), 3.28 (d, J = 14.7 Hz, 1H), 3.53 (m, 9H), 3.82 (dd, J = 8.7, 7.2 Hz, 2H), 4.18 (qd, J = 7.1, 1.5 Hz, 4H), 5.55 (s, 2H), 6.66 (q, J = 7.1 Hz, 2H), 7.62 (d, J = 8.1 Hz, 4H), 7.72 (m, 8H), 7.95 (m, 4H)
63iq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (m, 6H), 1.54 (m, 4H), 1.78 (dd, J = 13.1, 7.4 Hz, 1H), 2.14 (dd, J = 13.2, 8.8 Hz, 1H), 2.81 (d, J = 13.9 Hz, 4H), 2.94 (d, J = 11.0 Hz, 1H), 3.22 (s, 2H), 3.52 (ddt, J = 19.7, 11.9, 6.0 Hz, 4H), 3.92 (t, J = 8.0 Hz, 1H), 4.21 (qd, J = 7.8, 6.4, 4.7 Hz, 2H), 4.88 (s, 1H), 5.51 (s, 1H), 6.75 (q, J = 6.7 Hz, 1H), 7.49 (m, 2H), 7.72 (m, 2H)
63ir	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.04 (t, J = 7.4 Hz, 3H), 1.26 (t, J = 7.1 Hz, 3H), 1.51 (m, 4H), 1.76 (ddd, J = 25.2, 13.5, 7.1 Hz, 3H), 2.08 (dd, J = 13.1, 8.8 Hz, 1H), 2.74 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.47 (dq, J = 26.7, 7.9, 6.9 Hz, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 3.95 (t, J = 6.5 Hz, 2H), 4.18 (m, 2H), 5.45 (s, 1H), 6.66 (q, J = 6.9 Hz, 1H), 6.97 (m, 2H), 7.55 (m, 9H), 7.73 (d, J = 8.3 Hz, 1H)
63is	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.28 (t, J = 7.1 Hz, 4H), 1.54 (m, 4H), 1.81 (dd, J = 13.2, 7.6 Hz, 1H), 2.18 (dd, J = 13.3, 8.8 Hz, 1H), 2.87 (d, J = 11.2 Hz, 1H), 2.98 (d, J = 11.1 Hz, 1H), 3.51 (m, 4H), 4.00 (t, J = 8.1 Hz, 1H), 4.21 (qd, J = 7.1, 1.8 Hz, 2H), 5.49 (d, J = 2.0 Hz, 1H), 6.82 (q, J = 6.7 Hz, 1H), 7.13 (m, 2H), 7.46 (m, 2H), 7.67 (d, J = 8.5 Hz, 1H)
63it	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.53 (dt, J = 10.5, 5.6 Hz, 4H), 1.75 (dd, J = 13.0, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.7 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.15 (s, 3H), 3.53 (m, 4H), 3.83 (dd, J = 8.8, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.56 (s, 1H), 6.67 (q, J = 7.1 Hz, 1H), 7.66 (d, J = 8.2 Hz, 2H), 7.74 (m, 2H), 7.89 (m, 2H), 8.02 (m, 2H)
63iu	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.51 (dt, J = 10.8, 5.6 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.08 (dd, J = 13.1, 8.8 Hz, 1H), 2.40 (s, 3H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.15 (s, 3H), 3.53 (m, 4H), 3.81 (dd, J = 8.8, 7.2 Hz, 1H), 4.18 (qd, J = 7.2, 1.7 Hz, 2H), 5.74 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.85 (q, J = 6.6 Hz, 1H), 7.76 (dd, J = 1.7, 0.6 Hz, 1H), 7.83 (m, 2H), 7.95 (m, 2H), 8.03 (m, 3H)
63iv	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.05 (t, J = 7.4 Hz, 3H), 1.26 (m, 3H), 1.49 (dt, J = 10.7, 5.7 Hz, 4H), 1.78 (m, 3H), 2.08 (dd, J = 13.1, 8.8 Hz, 1H), 2.25 (d, J = 14.0 Hz, 1H), 2.39 (s, 3H), 2.75 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.53 (m, 4H), 3.83 (t, J = 8.0 Hz, 1H), 3.95 (t, J = 6.4 Hz, 2H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.76 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.75 (t, J = 6.7 Hz, 1H), 6.98 (m, 2H), 7.58 (m, 3H), 7.71 (m, 2H), 7.96 (d, J = 2.4 Hz, 1H)
63ix	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.14 (t, J = 7.0 Hz, 3H), 1.26 (t, J = 7.1 Hz, 7H), 1.51 (dt, J = 10.8, 5.7 Hz, 4H), 1.73 (dd, J = 13.0, 7.2 Hz, 1H), 2.08 (dd, J = 13.0,

	8.8 Hz, 1H), 2.39 (d, J = 1.7 Hz, 3H), 2.74 (d, J = 10.9 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.31 (d, J = 16.3 Hz, 3H), 3.56 (s, 6H), 3.81 (dd, J = 8.7, 7.1 Hz, 1H), 4.18 (m, 2H), 5.75 (s, 1H), 6.42 (d, J = 2.4 Hz, 1H), 6.82 (q, J = 6.5 Hz, 1H), 7.47 (dd, J = 8.3, 2.0 Hz, 2H), 7.70 (d, J = 1.8 Hz, 1H), 7.79 (m, 4H), 8.01 (d, J = 2.4 Hz, 1H)
63iy	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 5H), 1.53 (dt, J = 9.7, 5.4 Hz, 8H), 1.74 (dd, J = 13.0, 7.3 Hz, 2H), 2.09 (dd, J = 13.1, 8.8 Hz, 2H), 2.76 (d, J = 11.0 Hz, 2H), 2.91 (d, J = 19.4 Hz, 7H), 3.36 (s, 1H), 3.53 (m, 8H), 3.83 (dd, J = 8.7, 7.2 Hz, 2H), 4.18 (qd, J = 7.1, 1.6 Hz, 4H), 4.97 (s, 1H), 5.55 (s, 2H), 6.65 (q, J = 7.1 Hz, 2H), 7.62 (d, J = 8.1 Hz, 4H), 7.71 (m, 7H), 7.89 (m, 4H)
63iz	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 4H), 1.51 (dt, J = 10.6, 5.5 Hz, 4H), 1.76 (dd, J = 13.1, 7.3 Hz, 1H), 2.11 (dd, J = 13.1, 8.8 Hz, 1H), 2.40 (s, 3H), 2.78 (d, J = 11.1 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.54 (qq, J = 14.0, 7.5, 6.5 Hz, 4H), 3.88 (dd, J = 8.7, 7.4 Hz, 1H), 4.19 (qd, J = 7.1, 1.7 Hz, 2H), 5.75 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.83 (q, J = 6.6 Hz, 1H), 7.73 (d, J = 1.6 Hz, 1H), 7.84 (m, 4H), 7.99 (m, 3H)
63ja	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (m, 7H), 1.53 (dt, J = 9.9, 5.5 Hz, 8H), 1.77 (dd, J = 13.1, 7.4 Hz, 2H), 2.17 (m, 2H), 2.81 (d, J = 11.1 Hz, 2H), 2.93 (d, J = 11.1 Hz, 2H), 3.33 (d, J = 12.6 Hz, 1H), 3.52 (ddt, J = 17.5, 11.5, 5.2 Hz, 8H), 3.90 (dd, J = 8.6, 7.5 Hz, 2H), 4.20 (qd, J = 7.2, 1.7 Hz, 3H), 5.56 (s, 2H), 6.66 (q, J = 7.1 Hz, 2H), 7.64 (d, J = 8.1 Hz, 4H), 7.71 (d, J = 8.3 Hz, 4H), 7.79 (m, 4H), 7.96 (m, 4H)
63jb	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 5H), 1.53 (dt, J = 9.6, 5.3 Hz, 5H), 1.76 (dd, J = 13.1, 7.3 Hz, 1H), 2.11 (dd, J = 13.1, 8.7 Hz, 1H), 2.61 (m, 7H), 2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.56 (m, 8H), 3.71 (t, J = 4.7 Hz, 5H), 3.85 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.1, 1.6 Hz, 2H), 5.55 (s, 1H), 6.66 (q, J = 7.2 Hz, 1H), 7.63 (d, J = 8.1 Hz, 2H), 7.73 (m, 4H), 7.91 (m, 2H)
63jc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.21 (dt, J = 33.8, 7.2 Hz, 4H), 1.51 (dt, J = 10.8, 5.4 Hz, 4H), 1.74 (dd, J = 13.1, 7.3 Hz, 1H), 2.08 (m, 1H), 2.40 (s, 3H), 2.58 (dt, J = 23.6, 5.8 Hz, 6H), 2.75 (d, J = 11.0 Hz, 1H), 2.88 (dd, J = 11.0, 5.9 Hz, 1H), 3.32 (s, 1H), 3.57 (m, 6H), 3.70 (t, J = 4.7 Hz, 4H), 3.82 (m, 1H), 4.18 (qd, J = 7.1, 1.7 Hz, 2H), 5.75 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.82 (q, J = 6.6 Hz, 1H), 7.72 (d, J = 1.5 Hz, 1H), 7.81 (m, 4H), 7.96 (m, 4H)
63jd	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (td, J = 7.1, 1.3 Hz, 3H), 1.52 (dt, J = 9.5, 5.5 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.09 (dd, J = 13.1, 8.8 Hz, 1H), 2.75 (d, J = 11.1 Hz, 1H), 2.89 (d, J = 10.9 Hz, 1H), 3.04 (d, J = 1.3 Hz, 3H), 3.11 (s, 3H), 3.52 (m, 4H), 3.82 (m, 1H), 4.18 (qt, J = 7.1, 1.4 Hz, 2H), 5.55 (d, J = 1.3 Hz, 1H), 6.66 (q, J = 7.1 Hz, 1H), 7.51 (m, 2H), 7.68 (m, 6H)
63je	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 4H), 1.53 (dt, J = 9.5, 5.3 Hz, 4H), 1.75 (dd, J = 13.0, 7.3 Hz, 1H), 2.11 (dd, J = 13.1, 8.8 Hz, 1H), 2.78 (t, J = 9.8 Hz, 3H), 2.91 (d, J = 11.0 Hz, 4H), 3.50 (m, 7H), 3.75 (s, 2H), 3.85 (dd, J = 8.8, 7.3 Hz, 1H), 4.19 (qd, J = 7.2, 1.6 Hz, 2H), 5.55 (s, 1H), 6.66 (q, J = 7.0 Hz, 1H), 7.51 (m, 2H), 7.62 (d, J = 8.2 Hz, 2H), 7.72 (m, 4H)
63jf	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (td, J = 7.4, 1.9 Hz, 8H), 1.50 (dt, J = 10.8, 5.8 Hz, 5H), 1.75 (dd, J = 13.1, 7.4 Hz, 1H), 2.10 (dd, J = 13.2, 8.9 Hz, 1H), 2.39 (s, 3H), 2.86 (m, 8H), 3.26 (s, 1H), 3.52 (m, 8H), 3.76 (s, 2H), 3.88 (dd, J = 8.8, 7.3 Hz, 1H), 4.18 (ddtd, J = 7.7, 5.3, 3.6, 2.0 Hz, 2H), 4.93 (s, 2H), 5.74 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.82 (q, J = 6.6 Hz, 1H), 7.51 (dd, J = 8.3, 2.0 Hz, 2H), 7.75 (m,

	7H), 8.01 (d, J = 2.4 Hz, 1H)
63jg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.31 (t, J = 7.2 Hz, 8H), 1.57 (s, 10H), 1.95 (dd, J = 13.4, 8.4 Hz, 2H), 2.37 (t, J = 11.1 Hz, 2H), 3.12 (m, 4H), 3.61 (m, 15H), 4.31 (m, 6H), 5.49 (s, 1H), 5.62 (s, 1H), 6.25 (d, J = 6.9 Hz, 2H), 6.50 (t, J = 6.8 Hz, 2H), 7.31 (d, J = 2.2 Hz, 2H), 7.46 (dd, J = 15.4, 7.7 Hz, 4H), 7.65 (d, J = 8.5 Hz, 2H), 7.79 (d, J = 7.1 Hz, 2H)
63jh	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.50 (dt, J = 10.4, 5.5 Hz, 4H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 2.08 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.53 (m, 4H), 3.83 (m, 1H), 3.90 (s, 3H), 4.18 (qd, J = 7.2, 1.6 Hz, 2H), 5.75 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.78 (q, J = 6.7 Hz, 1H), 7.17 (t, J = 8.9 Hz, 1H), 7.48 (m, 2H), 7.61 (d, J = 1.8 Hz, 1H), 7.73 (m, 2H), 7.99 (d, J = 2.4 Hz, 1H)
63ji	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (d, J = 11.1 Hz, 1H), 1.51 (q, J = 6.8, 6.0 Hz, 4H), 1.78 (dd, J = 13.0, 7.0 Hz, 1H), 1.89 (s, 2H), 2.07 (dd, J = 13.1, 9.1 Hz, 1H), 2.40 (s, 3H), 2.68 (d, J = 11.1 Hz, 1H), 2.95 (d, J = 11.1 Hz, 1H), 3.03 (s, 3H), 3.11 (s, 3H), 3.22 (s, 2H), 3.45 (m, 3H), 3.63 (q, J = 7.9, 7.5 Hz, 3H), 5.75 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.82 (q, J = 6.6 Hz, 1H), 7.53 (d, J = 7.9 Hz, 2H), 7.70 (m, 1H), 7.80 (m, 4H), 8.01 (d, J = 2.5 Hz, 1H)
63jj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.06 (t, J = 7.4 Hz, 4H), 1.30 (t, J = 7.1 Hz, 3H), 1.57 (m, 4H), 1.86 (m, 3H), 2.30 (m, 1H), 3.09 (m, 3H), 3.54 (m, 4H), 4.03 (t, J = 6.4 Hz, 2H), 4.27 (m, 3H), 5.55 (s, 1H), 6.64 (q, J = 7.2 Hz, 1H), 7.12 (t, J = 8.8 Hz, 1H), 7.37 (m, 2H), 7.58 (q, J = 8.4 Hz, 4H)
63jk	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 5H), 1.50 (dt, J = 10.2, 5.2 Hz, 4H), 1.75 (dd, J = 13.1, 7.4 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.40 (s, 3H), 2.78 (d, J = 11.1 Hz, 1H), 2.91 (d, J = 13.6 Hz, 4H), 3.52 (m, 4H), 3.88 (dd, J = 8.7, 7.3 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.75 (s, 1H), 6.43 (d, J = 2.4 Hz, 1H), 6.82 (q, J = 6.5 Hz, 1H), 7.70 (d, J = 1.7 Hz, 1H), 7.78 (m, 4H), 7.90 (m, 2H), 8.01 (d, J = 2.4 Hz, 1H)
63jl	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.25 (m, 5H), 1.54 (dt, J = 11.2, 6.0 Hz, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.11 (dd, J = 13.1, 8.8 Hz, 1H), 2.58 (s, 3H), 2.77 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.55 (h, J = 7.5 Hz, 4H), 3.84 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.7 Hz, 2H), 5.58 (s, 1H), 6.65 (q, J = 6.6 Hz, 1H), 7.34 (d, J = 2.2 Hz, 1H), 7.49 (dd, J = 8.5, 2.3 Hz, 1H), 7.73 (m, 3H), 7.94 (ddd, J = 7.9, 1.8, 1.1 Hz, 1H), 8.32 (s, 1H)
63jm	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 4H), 1.54 (dt, J = 8.6, 4.8 Hz, 4H), 1.76 (dd, J = 13.1, 7.2 Hz, 1H), 2.11 (dd, J = 13.1, 8.8 Hz, 1H), 2.73 (s, 7H), 2.91 (d, J = 10.9 Hz, 1H), 3.55 (dp, J = 20.2, 7.2, 6.0 Hz, 4H), 3.84 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.1, 1.6 Hz, 2H), 5.59 (s, 1H), 6.69 (q, J = 6.4 Hz, 1H), 7.36 (d, J = 2.2 Hz, 1H), 7.50 (dd, J = 8.5, 2.2 Hz, 1H), 7.70 (dd, J = 13.1, 8.0 Hz, 2H), 7.85 (m, 2H), 8.34 (s, 1H)
63jn	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.30 (m, 10H), 1.52 (dt, J = 10.2, 5.7 Hz, 4H), 1.74 (dd, J = 13.1, 7.3 Hz, 1H), 2.08 (m, 1H), 2.83 (m, 4H), 3.20 (ddd, J = 11.9, 6.2, 3.0 Hz, 2H), 3.31 (s, 1H), 3.49 (ddd, J = 30.2, 13.4, 6.0 Hz, 4H), 3.90 (dddd, J = 32.2, 15.9, 7.4, 5.0 Hz, 5H), 4.18 (qd, J = 7.2, 1.5 Hz, 2H), 4.64 (p, J = 6.0 Hz, 1H), 5.52 (s, 1H), 6.96 (m, 2H), 7.40 (m, 2H), 7.54 (m, 4H)
63jo	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (m, 4H), 1.54 (dt, J = 7.6, 4.8 Hz, 5H),


	1.76 (dd, J = 13.1, 7.3 Hz, 1H), 2.11 (dd, J = 13.1, 8.8 Hz, 1H), 2.77 (d, J = 11.0 Hz, 1H), 2.92 (d, J = 18.5 Hz, 4H), 3.53 (m, 4H), 3.85 (dd, J = 8.7, 7.3 Hz, 1H), 4.19 (qd, J = 7.1, 1.6 Hz, 2H), 4.93 (s, 7H), 5.52 (d, J = 19.1 Hz, 1H), 6.63 (q, J = 6.7 Hz, 1H), 7.29 (d, J = 2.2 Hz, 1H), 7.46 (dd, J = 8.5, 2.3 Hz, 1H), 7.62 (m, 3H), 7.88 (dt, J = 7.7, 1.6 Hz, 1H), 8.37 (s, 1H)
63jp	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.53 (dt, J = 7.7, 4.7 Hz, 4H), 1.75 (dd, J = 13.1, 7.2 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.06 (s, 3H), 3.12 (s, 3H), 3.51 (m, 4H), 3.83 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.53 (s, 1H), 6.70 (q, J = 6.7 Hz, 1H), 7.32 (d, J = 2.2 Hz, 1H), 7.56 (m, 5H), 7.79 (s, 1H)
63jq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 5H), 1.46 (m, 7H), 1.74 (dd, J = 13.1, 7.3 Hz, 1H), 2.09 (dd, J = 13.1, 8.7 Hz, 1H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 4.15 (m, 4H), 5.53 (s, 1H), 6.62 (q, J = 7.1 Hz, 1H), 7.12 (t, J = 8.7 Hz, 1H), 7.38 (m, 2H), 7.58 (q, J = 8.4 Hz, 4H)
63jr	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.39 (t, J = 7.0 Hz, 3H), 1.51 (m, 4H), 1.73 (dd, J = 13.1, 7.3 Hz, 1H), 2.08 (dd, J = 13.1, 8.8 Hz, 1H), 2.74 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.49 (dt, J = 19.6, 13.1, 6.9 Hz, 4H), 3.82 (dd, J = 8.8, 7.3 Hz, 1H), 4.05 (q, J = 7.0 Hz, 2H), 4.18 (qd, J = 7.1, 1.5 Hz, 2H), 5.53 (s, 1H), 6.61 (q, J = 7.1 Hz, 1H), 6.96 (m, 2H), 7.55 (m, 6H)
63js	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.52 (m, 4H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.16 (s, 3H), 3.52 (m, 4H), 3.83 (dd, J = 8.7, 7.3 Hz, 1H), 4.18 (qd, J = 7.2, 1.5 Hz, 2H), 5.53 (s, 1H), 6.75 (q, J = 6.7 Hz, 1H), 7.56 (m, 5H), 7.84 (d, J = 1.9 Hz, 1H), 7.99 (m, 2H)
63jt	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.13 (t, J = 7.2 Hz, 3H), 1.27 (q, J = 6.8 Hz, 7H), 1.55 (m, 4H), 1.81 (dd, J = 13.2, 7.6 Hz, 1H), 2.18 (dd, J = 13.2, 8.7 Hz, 1H), 2.86 (d, J = 11.2 Hz, 1H), 2.98 (d, J = 11.2 Hz, 1H), 3.36 (q, J = 7.1 Hz, 2H), 3.56 (m, 6H), 3.98 (t, J = 8.1 Hz, 1H), 4.21 (qd, J = 7.2, 1.8 Hz, 2H), 5.54 (s, 1H), 6.74 (q, J = 6.8 Hz, 1H), 7.32 (d, J = 2.2 Hz, 1H), 7.49 (m, 3H), 7.65 (m, 3H)
63ju	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.04 (dd, J = 6.8, 1.9 Hz, 6H), 1.26 (t, J = 7.2 Hz, 3H), 1.50 (dt, J = 10.7, 5.7 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.07 (ddd, J = 13.0, 7.7, 4.9 Hz, 2H), 2.39 (s, 2H), 2.74 (d, J = 10.9 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.49 (d, J = 7.5 Hz, 1H), 3.56 (d, J = 7.9 Hz, 3H), 3.78 (m, 3H), 4.17 (qd, J = 7.1, 1.6 Hz, 2H), 5.75 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.75 (q, J = 6.6 Hz, 1H), 6.99 (m, 2H), 7.60 (dd, J = 8.7, 1.9 Hz, 3H), 7.72 (m, 2H), 7.97 (d, J = 2.4 Hz, 1H)
63jv	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.04 (s, 9H), 1.25 (t, J = 7.1 Hz, 3H), 1.48 (dt, J = 10.6, 5.7 Hz, 4H), 1.71 (dd, J = 13.1, 7.2 Hz, 1H), 2.05 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.72 (d, J = 11.0 Hz, 1H), 2.86 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.64 (s, 2H), 3.80 (dd, J = 8.7, 7.1 Hz, 1H), 4.17 (qd, J = 7.1, 1.5 Hz, 2H), 5.75 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.76 (q, J = 6.6 Hz, 1H), 6.98 (m, 2H), 7.57 (m, 3H), 7.70 (m, 2H), 7.96 (d, J = 2.4 Hz, 1H)
63jw	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1H NMR (MeOH-d ₄) δ: 1.29 (t, J = 7.1 Hz, 7H), 1.53 (s, 8H), 1.79 (s, 2H), 2.14 (s, 2H), 2.81 (s, 2H), 2.94 (d, J = 10.8 Hz, 2H), 3.50 (s, 7H), 3.57 (s, 2H), 3.90 (t, J = 8.0 Hz, 2H), 4.22 (qd, J = 7.1, 1.7 Hz, 3H), 5.43 (s, 1H), 6.51 (s, 1H), 6.85 (s, 1H), 7.25

	(s, 1H), 7.48 (d, J = 9.7 Hz, 4H), 7.55 (d, J = 7.5 Hz, 2H), 7.79 (s, 4H), 8.13 (s, 2H)
63jx	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.49 (dt, J = 10.8, 5.8 Hz, 4H), 1.73 (dd, J = 13.1, 7.2 Hz, 1H), 2.03 (m, 3H), 2.39 (s, 3H), 2.73 (d, J = 11.0 Hz, 1H), 2.85 (m, 3H), 3.53 (m, 4H), 3.81 (dd, J = 8.7, 7.1 Hz, 1H), 4.17 (m, 4H), 5.75 (s, 1H), 6.40 (d, J = 2.3 Hz, 1H), 6.77 (dd, J = 17.0, 7.9 Hz, 2H), 7.36 (dq, J = 4.4, 2.5 Hz, 2H), 7.56 (d, J = 1.8 Hz, 1H), 7.69 (m, 2H), 7.96 (d, J = 2.3 Hz, 1H)
63jy	¹ H NMR (400 MHz, MeOH-d ₄): δ 1.27 (dd, J = 7.9, 6.4 Hz, 4H), 1.54 (dt, J = 10.7, 5.6 Hz, 4H), 1.76 (dd, J = 13.2, 7.4 Hz, 1H), 2.12 (dd, J = 13.1, 8.8 Hz, 1H), 2.78 (m, 3H), 2.92 (m, 3H), 3.53 (m, 6H), 3.76 (s, 2H), 3.85 (dd, J = 8.7, 7.2 Hz, 1H), 4.19 (qd, J = 7.1, 1.7 Hz, 2H), 5.51 (d, J = 15.6 Hz, 1H), 6.72 (q, J = 6.6 Hz, 1H), 7.33 (d, J = 2.2 Hz, 1H), 7.51 (m, 3H), 7.66 (m, 2H), 7.79 (s, 1H)
63jz	¹ H NMR (400 MHz, MeOH-d ₄): δ 0.46 (m, 4H), 1.27 (m, 4H), 1.53 (dt, J = 11.2, 5.6 Hz, 4H), 1.72 (m, 2H), 2.13 (dd, J = 13.1, 8.8 Hz, 1H), 2.58 (s, 2H), 2.73 (s, 2H), 2.80 (d, J = 11.1 Hz, 1H), 2.93 (d, J = 11.0 Hz, 1H), 3.52 (ddd, J = 25.7, 12.3, 6.8 Hz, 6H), 3.76 (s, 2H), 3.89 (dd, J = 8.7, 7.3 Hz, 1H), 4.19 (qd, J = 7.1, 1.7 Hz, 2H), 5.52 (d, J = 17.0 Hz, 1H), 6.71 (q, J = 6.7 Hz, 1H), 7.33 (d, J = 2.3 Hz, 1H), 7.57 (m, 5H), 7.80 (s, 1H)
63ka	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.25 (t, J = 7.1 Hz, 3H), 1.50 (dt, J = 10.3, 5.3 Hz, 4H), 1.73 (dd, J = 13.1, 7.3 Hz, 1H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.41 (s, 3H), 2.75 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.51 (m, 4H), 3.82 (dd, J = 8.8, 7.2 Hz, 1H), 4.17 (qd, J = 7.1, 1.6 Hz, 2H), 5.76 (s, 1H), 6.45 (d, J = 2.4 Hz, 1H), 6.88 (q, J = 6.6 Hz, 1H), 7.92 (m, 3H), 8.07 (d, J = 2.4 Hz, 1H), 8.29 (m, 3H), 8.52 (d, J = 8.9 Hz, 1H), 9.32 (d, J = 5.9 Hz, 1H)
63kb	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (t, J=7.22 Hz, 3 H) 1.65 - 1.91 (m, 4 H) 2.12 (dd, J=13.67, 8.79 Hz, 1 H) 2.53 (dd, J=13.67, 8.79 Hz, 1 H) 3.35 (s, 2 H) 3.56 - 3.91 (m, 4 H) 4.35 (qd, J=7.06, 3.03 Hz, 2 H) 4.65 (t, J=8.69 Hz, 1 H) 6.66 (d, J=5.66 Hz, 1 H) 7.02 (d, J=2.34 Hz, 1H) 7.69 - 7.78 (m, 2 H) 7.79 - 7.88 (m, 1 H) 8.29 (d, J=1.37 Hz, 1 H)
63kc	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (t, J=7.13 Hz, 3 H) 1.40 (s, 9 H) 1.64 - 1.85 (m, 4 H) 2.03 - 2.18 (m, 1 H) 2.43 - 2.61 (m, 1 H) 3.53 - 3.87 (m, 4 H) 4.27 - 4.43 (m, 2 H) 4.56 - 4.70 (m, 1 H) 5.51 (s, 1 H) 6.56 (d, J=2.34 Hz, 1 H) 7.30 - 7.42 (m, 1 H) 7.53 - 7.61 (m, 1 H) 7.69 (d, J=1.95 Hz, 2 H) 8.01 (d, J=2.54 Hz, 1 H)
63kd	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.22 - 1.42 (m, 9 H) 1.51 - 1.72 (m, 4 H) 1.90 - 2.09 (m, 1 H) 2.33 - 2.52 (m, 1 H) 3.09 (s, 1 H) 3.21 (d, J=4.69 Hz, 2 H) 3.40 - 3.72 (m, 4 H) 4.31 (dd, J=7.13, 2.25 Hz, 2 H) 4.48 (s, 1 H) 5.64 (s, 1 H) 6.47 (d, J=2.34 Hz, 1 H) 7.02 (d, J=6.64 Hz, 1 H) 7.43 - 7.60 (m, 2 H) 7.72 (d, J=8.59 Hz, 1 H) 7.95 (d, J=2.34 Hz, 1 H)
63ke	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.68 - 0.95 (m, 2 H) 1.05 (dd, J=8.40, 2.15 Hz, 2 H) 1.35 (t, J=7.13 Hz, 4 H) 1.63 - 1.89 (m, 4 H) 1.98 - 2.18 (m, 2 H) 2.44 - 2.63 (m, 1 H) 3.78 (d, J=5.08 Hz, 4 H) 4.35 (d, J=7.03 Hz, 2 H) 4.63 (s, 1 H) 6.31 (d, J=2.34 Hz, 1 H) 7.09 (d, J=6.25 Hz, 1 H) 7.51 - 7.67 (m, 2 H) 7.73 (d, J=8.20 Hz, 1 H) 7.93 (d, J=2.54 Hz, 1 H)
63kf	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.35 (t, J=7.13 Hz, 4 H) 1.74 (br. s., 4 H) 2.04 - 2.15 (m, 1 H) 2.34 (s, 3 H) 2.37 (s, 3 H) 2.44 - 2.58 (m, 1 H) 3.31 (d, J=2.34 Hz, 2 H) 3.54 - 3.89 (m, 3 H) 4.34 (dd, J=7.13, 3.22 Hz, 2 H) 4.61 (s, 1 H) 6.10 (s, 1 H) 6.51

	- 6.65 (m, 1 H) 7.03 (d, J=2.15 Hz, 1 H) 7.28 (s, 1 H) 7.42 - 7.50 (m, 1 H) 7.54 (s, 1 H) 7.76 - 7.88 (m, 2 H) 7.90 - 8.01 (m, 1 H) 8.33 (s, 1 H)
63kg	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.29 (t, J = 7.1 Hz, 3H), 1.55 (s, 3H), 1.58 (d, J = 5.8 Hz, 1H), 1.88 (m, 1H), 2.29 (m, 6H), 3.04 (m, 2H), 3.43 (s, 2H), 3.56 (s, 2H), 4.24 (m, 2H), 6.68 (q, J = 6.9 Hz, 1H), 7.17 (d, J = 7.9 Hz, 1H), 7.36 (m, 2H), 7.45 (m, 1H), 7.52 (s, 2H), 7.53 (d, J = 2.8 Hz, 1H), 7.63 (dd, J = 8.2, 2.0 Hz, 1H), 7.73 (d, J = 8.2 Hz, 1H)
63kh	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.05 (t, J = 7.4 Hz, 3H), 1.26 (td, J = 7.1, 2.1 Hz, 3H), 1.50 (s, 3H), 1.53 (d, J = 5.7 Hz, 1H), 1.79 (m, 3H), 2.09 (dd, J = 13.1, 8.8 Hz, 1H), 2.75 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.50 (s, 3H), 3.83 (dd, J = 8.8, 7.2 Hz, 1H), 4.02 (t, J = 6.5 Hz, 2H), 4.17 (m, 2H), 5.46 (s, 1H), 6.67 (q, J = 6.7 Hz, 1H), 7.12 (t, J = 8.6 Hz, 1H), 7.40 (m, 4H), 7.52 (s, 4H), 7.54 (s, 1H), 7.62 (dd, J = 8.2, 2.1 Hz, 1H), 7.74 (d, J = 8.3 Hz, 1H)
63ki	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 3H), 1.53 (dd, J = 11.2, 5.2 Hz, 5H), 1.75 (dd, J = 13.1, 7.3 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.90 (d, J = 11.0 Hz, 1H), 3.49 (m, 2H), 3.51 (s, 3H), 3.84 (dd, J = 8.7, 7.3 Hz, 1H), 4.18 (qd, J = 7.1, 1.6 Hz, 2H), 5.44 (s, 1H), 6.66 (q, J = 6.9 Hz, 1H), 7.26 (m, 1H), 7.45 (m, 8H), 7.70 (d, J = 7.2 Hz, 1H)
63kj	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.27 (t, J = 7.1 Hz, 3H), 1.53 (dd, J = 11.0, 5.2 Hz, 4H), 1.77 (dd, J = 13.2, 7.4 Hz, 1H), 2.13 (dd, J = 13.1, 8.8 Hz, 1H), 2.80 (d, J = 11.1 Hz, 1H), 2.93 (d, J = 11.1 Hz, 1H), 3.46 (m, 1H), 3.53 (m, 2H), 3.91 (t, J = 8.1 Hz, 1H), 4.19 (qd, J = 7.1, 1.4 Hz, 2H), 5.47 (s, 1H), 6.69 (q, J = 6.9 Hz, 1H), 7.35 (m, 1H), 7.45 (m, 4H), 7.54 (d, J = 4.6 Hz, 4H), 7.65 (m, 3H), 7.77 (d, J = 8.2 Hz, 1H)
63kk	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.25 (t, J = 7.1 Hz, 3H), 1.51 (m, 4H), 1.74 (dd, J = 13.1, 7.4 Hz, 1H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.29 (d, J = 9.9 Hz, 6H), 2.39 (s, 3H), 2.77 (d, J = 11.1 Hz, 1H), 2.90 (d, J = 11.1 Hz, 1H), 3.54 (tq, J = 14.0, 7.9, 6.7 Hz, 4H), 3.88 (dd, J = 8.7, 7.4 Hz, 1H), 4.17 (m, 2H), 5.74 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.77 (q, J = 6.6 Hz, 1H), 7.19 (d, J = 7.8 Hz, 1H), 7.36 (dd, J = 7.6, 2.1 Hz, 1H), 7.42 (d, J = 1.5 Hz, 1H), 7.59 (d, J = 1.9 Hz, 1H), 7.72 (m, 2H), 7.97 (d, J = 2.4 Hz, 1H)
63kl	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (t, J = 7.1 Hz, 3H), 1.51 (dd, J = 11.1, 5.8 Hz, 5H), 1.74 (dd, J = 13.1, 7.3 Hz, 1H), 2.09 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.76 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.55 (d, J = 5.0 Hz, 4H), 3.85 (dd, J = 8.7, 7.2 Hz, 1H), 4.18 (m, 2H), 4.65 (s, 2H), 5.78 (s, 1H), 6.41 (d, J = 2.4 Hz, 1H), 6.87 (q, J = 6.5 Hz, 1H), 7.47 (dd, J = 10.9, 8.2 Hz, 3H), 7.59 (m, 2H), 7.79 (dd, J = 8.3, 2.1 Hz, 1H), 7.93 (d, J = 2.3 Hz, 2H)
63km	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.61 – 7.49 (m, 4H), 7.35 – 7.27 (m, 2H), 6.77 (dd, J = 8.4, 1.8 Hz, 1H), 6.60 (q, J = 7.3 Hz, 1H), 5.55 – 5.46 (m, 1H), 4.24 – 4.13 (m, 4H), 3.83 (dd, J = 8.8, 7.2 Hz, 1H), 2.93 – 2.71 (m, 4H), 2.14 – 1.94 (m, 3H), 1.74 (dd, J = 13.1, 7.3 Hz, 1H), 1.56 – 1.48 (m, 1H), 1.51 (s, 3H), 1.27 (td, J = 7.1, 2.0 Hz, 3H).
63kn	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.71 (d, J = 5.2 Hz, 2H), 8.02 (td, J = 7.7, 1.7 Hz, 1H), 7.78 – 7.68 (m, 3H), 7.51 (tt, J = 7.9, 3.3 Hz, 5H), 6.92 (d, J = 6.5 Hz, 1H), 5.81 (d, J = 3.8 Hz, 2H), 4.18 (qd, J = 7.1, 1.7 Hz, 2H), 3.83 (s, 1H), 3.56 (s, 6H), 3.57 – 3.46 (m, 1H), 2.89 (d, J = 11.0 Hz, 2H), 2.76 (d, J = 11.0 Hz, 2H), 2.09 (dd, J = 13.1, 8.9 Hz, 1H), 1.74 (dd, J = 13.1, 7.3 Hz, 1H), 1.52 (dd, J = 10.9, 5.5 Hz, 5H), 1.31 –

	1.22 (m, 6H)
63ko	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.99 (d, J = 4.9 Hz, 2H), 8.03 (s, 1H), 7.73 (dd, J = 15.2, 7.7 Hz, 2H), 7.60 – 7.48 (m, 2H), 5.69 (s, 1H), 4.18 (q, J = 7.1 Hz, 2H), 3.83 (t, J = 8.1 Hz, 1H), 3.54 – 3.43 (m, 4H), 2.89 (d, J = 11.1 Hz, 1H), 2.75 (d, J = 11.0 Hz, 1H), 2.14 – 2.04 (m, 1H), 1.74 (dd, J = 13.0, 7.4 Hz, 1H), 1.50 (dd, J = 10.6, 5.5 Hz, 5H), 1.26 (t, J = 7.2 Hz, 4H)
63kp	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.97 (s, 1H), 7.76 (s, 2H), 7.66 (d, J = 16.1 Hz, 2H), 7.49 (d, J = 7.9 Hz, 1H), 7.25 (d, J = 8.1 Hz, 1H), 6.77 (d, J = 7.1 Hz, 1H), 6.41 (s, 1H), 5.74 (d, J = 2.7 Hz, 1H), 4.68 (s, 2H), 4.18 (d, J = 7.6 Hz, 2H), 3.84 (t, J = 8.1 Hz, 1H), 3.56 (s, 3H), 3.49 (s, 1H), 3.30 (d, J = 3.4 Hz, 9H), 2.89 (d, J = 11.1 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.37 (d, J = 14.1 Hz, 5H), 1.79 – 1.69 (m, 1H), 1.51 (d, J = 8.8 Hz, 4H), 1.30 – 1.21 (m, 4H)
63kq	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.98 (s, 1H), 7.76 (d, J = 5.1 Hz, 2H), 7.63 (s, 1H), 7.48 (d, J = 13.3 Hz, 3H), 6.77 (d, J = 6.8 Hz, 1H), 6.41 (s, 1H), 5.74 (s, 1H), 4.66 (s, 2H), 4.18 (d, J = 7.4 Hz, 2H), 3.82 (t, J = 8.2 Hz, 1H), 3.56 (s, 3H), 3.50 (s, 1H), 2.89 (d, J = 11.0 Hz, 1H), 2.75 (d, J = 11.1 Hz, 1H), 2.39 (s, 6H), 1.74 (dd, J = 13.0, 7.2 Hz, 1H), 1.51 (s, 4H), 1.30 – 1.22 (m, 3H)
63kr	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.43 (d, J = 2.5 Hz, 1H), 7.98 (d, J = 10.3 Hz, 2H), 7.79 (d, J = 8.3 Hz, 1H), 7.71 (d, J = 8.5 Hz, 1H), 7.63 (s, 1H), 6.83 (dd, J = 19.6, 7.7 Hz, 2H), 6.42 (d, J = 2.3 Hz, 1H), 5.74 (s, 1H), 4.35 (q, J = 7.0 Hz, 2H), 4.17 (q, J = 7.1 Hz, 2H), 3.55 (s, 3H), 3.48 (d, J = 13.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 2.74 (d, J = 11.0 Hz, 1H), 2.39 (s, 3H), 2.07 (dd, J = 13.0, 8.9 Hz, 1H), 1.73 (dd, J = 13.0, 7.2 Hz, 1H), 1.50 (d, J = 8.3 Hz, 4H), 1.38 (t, J = 7.1 Hz, 3H), 1.26 (t, J = 7.2 Hz, 3H).
63ks	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.46 (s, 1H), 7.99 (s, 2H), 7.83 – 7.69 (m, 2H), 7.64 (s, 1H), 6.80 (d, J = 5.3 Hz, 1H), 6.42 (s, 1H), 5.74 (s, 1H), 4.18 (d, J = 7.3 Hz, 2H), 3.94 (d, J = 2.7 Hz, 3H), 3.86 (t, J = 8.1 Hz, 1H), 3.56 (s, 3H), 3.50 (s, 1H), 2.91 (d, J = 11.0 Hz, 1H), 2.77 (d, J = 11.6 Hz, 1H), 2.39 (d, J = 2.7 Hz, 3H), 2.10 (t, J = 10.9 Hz, 1H), 1.80 – 1.70 (m, 1H), 1.51 (s, 4H), 1.26 (dd, J = 8.3, 5.7 Hz, 3H).
63kt	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.67 (d, J = 8.5 Hz, 1H), 7.44 (ddd, J = 8.0, 4.8, 2.6 Hz, 2H), 7.32 – 7.24 (m, 2H), 7.07 (dd, J = 8.4, 2.5 Hz, 1H), 6.99 (d, J = 7.6 Hz, 1H), 6.76 (q, J = 6.9 Hz, 1H), 5.51 (s, 1H), 4.27 (dd, J = 7.0, 2.0 Hz, 1H), 4.25 – 4.13 (m, 4H), 3.76 (s, 2H), 3.58 (s, 2H), 3.51 (d, J = 14.9 Hz, 2H), 3.42 (s, 3H), 3.08 (d, J = 11.4 Hz, 1H), 2.99 (d, J = 11.4 Hz, 1H), 2.28 (dd, J = 13.3, 8.7 Hz, 1H), 1.88 (dd, J = 13.3, 8.0 Hz, 1H), 1.57 (p, J = 5.4 Hz, 4H), 1.29 (t, J = 7.1 Hz, 3H)
63ku	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 8.97 (d, J = 1.5 Hz, 1H), 8.80 (dd, J = 2.6, 1.5 Hz, 1H), 8.71 (d, J = 2.6 Hz, 1H), 7.77 (d, J = 8.3 Hz, 1H), 7.64 – 7.55 (m, 1H), 6.87 (q, J = 6.7 Hz, 1H), 5.62 (s, 1H), 4.23 – 4.13 (m, 2H), 3.82 (dd, J = 8.7, 7.2 Hz, 1H), 3.60 – 3.42 (m, 3H), 2.89 (d, J = 11.0 Hz, 1H), 2.75 (d, J = 11.0 Hz, 1H), 2.09 (dd, J = 13.1, 8.7 Hz, 1H), 1.74 (dd, J = 13.1, 7.2 Hz, 1H), 1.51 (dt, J = 10.9, 5.6 Hz, 3H), 1.26 (t, J = 7.1 Hz, 2H)

Example 64a: (S)-Octyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate



To a flask equipped with a Dean Stark trap were added (2S)-8-[2-amino-6-((1R)-1-[4-chloro-2-(3-methyl pyrazol-1-yl)phenyl]-2,2,2-trifluoroethoxy)pyrimidin-4-yl]-3,8-diazaspiro[4.5]decane-2-carboxylic acid (1 g, 1.78 mmol), toluene (25 mL), and *p*-toluene sulfonic acid monohydrate (336 mg, 1.77 mmol), and *n*-octanol (690 mg, 5.30 mmol). The reaction mixture was heated to reflux for 48 h, cooled to RT, and concentrated *in vacuo*. Purification on a 120 g Isco RediSep silica cartridge (CH₂Cl₂/MeOH/ NH₄OH) provided the title compound as a white solid.

Applying the generic scheme below, the following examples of Table 19a were prepared as described above for (S)-octyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (Example 64a), using the appropriate alcohol.





Table 19a.



Ex. No.	R	CAS Name	LCMS (MH ⁺)
64a		(S)-Octyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	679.2
64b		(S)-cyclopentyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	635.1
64c		(S)-pentyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	637
64d		(S)-cyclohexyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	648
64e		(S)-propyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	608
64f		(S)-neopentyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	636
64g		(S)-butyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	622

64h		(S)-isopropyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	622
-----	---	---	-----


Table 19b.

NMR Data for Compounds of Table 19a

Ex. No.	NMR
64a	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.82 - 0.96 (m, 3 H), 1.20 - 1.47 (m, 10 H), 1.53 - 1.79 (m, 6 H), 2.04 (dd, J = 13.6, 8.8 Hz, 1 H), 2.38 (s, 3 H), 2.49 (dd, J = 13.6, 8.8 Hz, 1 H), 3.28 (s, 2 H), 3.42 - 3.85 (m, 4 H), 4.16 - 4.39 (m, 2 H), 4.60 (t, J = 8.8 Hz, 1 H), 5.81 (s, 1 H), 6.42 (d, J = 2.2 Hz, 1 H), 6.85 (q, J = 6.6 Hz, 1 H), 7.46 - 7.60 (m, 2 H), 7.71 (d, J = 8.3 Hz, 1 H), 7.93 (d, J = 2.4 Hz, 1 H)
64b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.50 - 2.10 (m, 13 H), 2.38 (s, 3 H), 2.45 (dd, J = 13.6, 8.8 Hz, 1 H), 3.27 (d, J = 1.2 Hz, 2 H), 3.43 - 3.76 (m, 4 H), 4.55 (t, J = 8.7 Hz, 1 H), 5.26 - 5.39 (m, 1 H), 5.74 (s, 1 H), 6.42 (d, J = 2.3 Hz, 1 H), 6.83 (q, J = 6.6 Hz, 1 H), 7.45 - 7.59 (m, 2 H), 7.71 (d, J = 8.4 Hz, 1 H), 7.93 (d, J = 2.3 Hz, 1 H)
64c	¹ H NMR (400MHz, MeOH-d ₄): δ ppm 0.94 (t, J=7.2Hz, 3H), 1.35-1.39 (m, 4H), 1.52-1.56 (m, 4 H), 1.64-1.71 (m, 2H), 1.74 - 1.79 (m, 1H), 2.08-2.14 (m, 1H), 2.40 (s, 3 H), 2.77 (d, J=10.8 Hz ,1H), 2.92 (d, J=10.8 Hz, 1H), 3.48-3.58 (m, 4 H), 3.83-3.87 (m, 1H), 4.13-4.18 (m, 2H), 5.69 (s, 1 H), 6.43 (d, J=2.0 Hz, 1H), 6.81-6.86 (m, 1H), 7.51-7.55 (m, 2H), 7.72 (d, J=8.4 Hz, 1H), 7.95 (d, J=2.0 Hz, 1H)
64d	¹ H NMR (400MHz, MeOH-d ₄): δ ppm 1.31-1.56 (m, 10H), 1.75-1.80 (m, 3H), 1.85 - 1.89 (m, 2H), 2.08-2.13 (m, 1H), 2.39 (s, 3H), 2.76 (d, J=10.8 Hz ,1 H), 2.93 (d, J=10.8 Hz, 1H), 3.50-3.58 (m, 4H), 3.81-3.84 (m, 1H), 4.77-4.83 (m, 1H), 5.69 (s, 1 H), 6.42 (d, J=2.0 Hz, 1H), 6.81-6.86 (m, 1H), 7.51-7.55 (m, 2H), 7.72 (d, J=8.4 Hz, 1H), 7.95 (d, J=2.4 Hz, 1H)
64e	¹ H NMR (400MHz, MeOH-d ₄): δ ppm 0.98 (t, J=7.6Hz, 3H), 1.54-1.59 (m, 4 H), 1.66 (m, 2H), 1.81 - 1.86 (m, 1H), 2.17-2.23 (m, 1H), 2.40 (s, 3H), 2.89 (d, J=11.2 Hz ,1H), 3.00 (d, J=11.2 Hz, 1H), 3.47-3.62 (m, 4H), 4.03 (t, J=8.0 Hz, 1H), 4.11-4.18 (m, 2H), 5.70 (s, 1H), 6.43 (d, J=2.4 Hz, 1H), 6.84 (q, 1H), 7.51-7.55 (m, 2H), 7.73 (d, J=8.4 Hz, 1H), 7.95 (d, J=2.4 Hz, 1H)
64f	¹ H NMR (400MHz, MeOH-d ₄): δ ppm 0.98 (s, 9H), 1.50-1.58 (m, 4 H), 1.77-1.82 (m, 1H), 2.12-2.17 (m, 1H), 2.40 (s, 3H), 2.79 (d, J=11.2 Hz ,1H), 2.94 (d, J=11.2 Hz, 1H), 3.52-3.58 (m, 4H), 3.83-3.93 (m, 3H), 5.70 (s, 1 H), 6.43 (d, J=2.4 Hz, 1H), 6.81-6.86 (m, 1H), 7.52-7.55 (m, 2H), 7.73 (d, J=8.4 Hz, 1H), 7.95 (d, J=2.4 Hz, 1H)
64g	¹ H NMR (400MHz, MeOH-d ₄): δ ppm 0.95 (t, J=7.6Hz, 3H), 1.37-1.43 (m, 2H), 1.50-1.54 (m, 4H), 1.60-1.67 (m, 2H), 1.72-1.77 (m, 1H), 2.06-2.12 (m, 1H), 2.38 (s, 3 H), 2.75 (d, J=11.2 Hz ,1 H), 2.90 (d, J=11.2 Hz, 1H), 3.45-3.58 (m, 4 H), 3.83-3.86 (m, 1H), 4.10-4.20 (m, 2H), 5.67 (s, 1H), 6.40 (d, J=2.4 Hz, 1H), 6.80-6.85 (m, 1 H), 7.50-7.53 (m, 2H), 7.71 (d, J=8.0 Hz, 1H), 7.93 (d, J=2.0 Hz, 1H)
64h	¹ H NMR (400MHz, MeOH-d ₄): δ ppm 0.85 (d, J=6.8Hz, 6H), 1.42-1.47 (m, 4H), 1.68-1.73 (m, 1H), 1.82-1.89 (m, 1H), 2.05-2.10 (m, 1H), 2.28 (s, 3H), 2.74 (d, J=11.2 Hz ,1

	H), 2.87 (d, J=11.2 Hz, 1H), 3.37-3.48 (m, 4H), 3.81-3.91 (m, 3H), 5.58 (s, 1H), 6.30 (d, J=2.0 Hz, 1H), 6.70-6.75 (m, 1 H), 7.40-7.43 (m, 2H), 7.60 (d, J=8.4 Hz, 1H), 7.83 (d, J=2.4 Hz, 1H)
--	--

Example 65a: (S)-Tert-butyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate



5 *Step 1:* To a mixture of (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-((benzyloxy)carbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (2.8 g, 4.1 mmol) in *t*-BuOH (50 mL) were added BOC₂O (3.5 g, 16.5 mmol) and DMAP (0.201 g, 1.65 mmol), and the reaction was heated to 50°C for 45 min. Then the reaction was cooled to RT and concentrated *in vacuo*. Purification on a 220 g Isco RediSep
10 silica cartridge (EtOAc/heptane) provided (S)-2-benzyl 3-tert-butyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as an off-white solid.

Step 2: To a solution of (S)-2-benzyl 3-tert-butyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (1.35 g, 1.7 mmol) in EtOAc (130 mL) was added 5% (w/w) Pd/C (130 mg). The solution was degassed, charged with 1 atm H₂ (balloon), and stirred at RT for 3.5 h. Then the solids were filtered through celite, washed with EtOAc/methanol, and the filtrate was concentrated *in vacuo*. Purification on a 220 g Isco RediSep silica cartridge (CH₂Cl₂/MeOH/
20 NH₄OH) provided the title compound as an off-white solid.

Applying the generic scheme below, the following examples of Table 20a were prepared as described above for (S)-tert-butyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-

yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (Example 65).





Table 20a.



5

Ex. No.	Ar	CAS Name	LCMS (MH ⁺)
65a		(S)-tert-butyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	623
65b		(S)-tert-butyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(2-(3-methyl-1H-pyrazol-1-yl)-4-propylphenyl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	630


65c		(S)-tert-butyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	693
65d		(S)-tert-butyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	722

Table 20b.

NMR Data for Compounds of Table 20a

Ex. No.	NMR
65a	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.54 (s, 9 H), 1.57 - 1.72 (m, 4 H), 2.02 (dd, J = 13.62, 8.44 Hz, 1 H), 2.38 (s, 3 H), 2.40 - 2.47 (m, 1 H), 3.18 - 3.37 (m, 2 H), 3.47 - 3.75 (m, 4 H), 4.49 (t, J = 8.61 Hz, 1 H), 5.76 (s, 1 H), 6.42 (d, J = 2.34 Hz, 1 H), 6.84 (q, J = 6.57 Hz, 1 H), 7.46 - 7.59 (m, 2 H), 7.71 (d, J = 8.35 Hz, 1 H), 7.93 (d, J = 2.39 Hz, 1 H)
65b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.96 (t, J=7.35 Hz, 3 H) 1.53 (s, 9 H) 1.56 - 1.77 (m, 6 H) 1.99 (dd, J=13.52, 8.25 Hz, 1 H) 2.37 - 2.42 (m, 4 H) 2.59 - 2.73 (m, 2 H) 3.14 - 3.29 (m, 2 H) 3.45 - 3.74 (m, 4 H) 4.43 (t, J=8.47 Hz, 1 H) 5.72 (s, 1 H) 6.38 (d, J=2.29 Hz, 1 H) 6.72 (q, J=6.74 Hz, 1 H) 7.23 (d, J=1.61 Hz, 1 H) 7.33 (dd, J=8.10, 1.61 Hz, 1 H) 7.63 (d, J=8.10 Hz, 1 H) 7.85 (d, J=2.34Hz, 1 H)
65c	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm .49 (s, 4 H) 1.50 (s, 5 H) 1.53 - 1.64 (m, 4 H) 1.90 - 2.01 (m, 1 H) 2.27 (s, 3 H) 2.30 (s, 3 H) 2.31 - 2.37 (m, 1 H) 2.38 (s, 3 H) 3.09 - 3.25 (m, 2 H) 3.43 - 3.70 (m, 4 H) 4.32 - 4.42 (m, 1 H) 5.74 (s, 1 H) 6.39 (d, J=2.29 Hz, 1 H) 6.75 (q, J=6.67 Hz, 1 H) 7.19 (d, J=7.91 Hz, 1 H) 7.36 (dd, J=7.81, 1.81 Hz, 1 H) 7.42 (s, 1 H) 7.58 (s, 1 H) 7.68 - 7.78 (m, 2 H) 7.93 (d, J=2.29 Hz, 1 H)
65d	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.31 (d, J=6.05 Hz, 6 H) 1.50 (s, 4 H) 1.51 (s, 5 H) 1.55 - 1.70 (m, 4 H) 1.92 - 2.06 (m, 1 H) 2.38 (s, 3 H) 2.39 - 2.48 (m, 1 H) 3.16 - 3.27 (m, 2 H) 3.47 - 3.75 (m, 4 H) 4.46 (t, J=8.64 Hz, 1 H) 4.63 (dt, J=12.10, 6.05 Hz, 1 H) 5.85 (s, 1 H) 6.39 (d, J=2.29 Hz, 1 H) 6.76 (q, J=6.62 Hz, 1 H) 6.97 (d, J=8.79 Hz, 2 H) 7.55 - 7.63 (m, 3 H) 7.67 - 7.77 (m, 2 H) 7.93 (d, J=2.29 Hz, 1 H)

Example 66a: (S)-2-(Dimethylamino)ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate



- 5 *Step 1:* To a mixture of (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (85 mg, 0.16 mmol) in THF (10 mL) was added BOC₂O (4 g, 18.6 mmol) in THF (10 mL), and the reaction mixture was stirred at RT for 16 h. Then the reaction was diluted with CH₂Cl₂, cooled to 0 °C, and the pH adjusted to 2 with 2 N HCl. The reaction mixture was then extracted CH₂Cl₂ and
- 10 concentrated *in vacuo* to provide (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy) pyrimidin-4-yl)-2-(tert-butoxycarbonyl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid as an off-white solid that was used directly without further purification .
- 15 *Step 2:* To a solution of (S)-8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2-(tert-butoxycarbonyl)-2,8-diazaspiro [4.5]decane-3-carboxylic acid (1.6 g, 2.45 mmol) in DMF (24 mL) were added (2-chloro-ethyl)-dimethyl-amine hydrochloride (535 mg, 3.7 mmol) and K₂CO₃ (1.0 g, 7.4 mmol), and the reaction mixture was heated at 65 °C for 16 h. Then the reaction was cooled to RT, partitioned
- 20 between EtOAc and water, and extracted. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification via prep-HPLC column chromatography (CH₂Cl₂/MeOH/NH₄OH) provided (S)-2-tert-butyl 3-(2-(dimethylamino)ethyl) 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate as an off-white
- 25 solid.

Step 3: To a solution of (S)-2-tert-butyl 3-(2-(dimethylamino)ethyl) 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-2,3-dicarboxylate (1.4 g, 1.86 mmol) in CH₂Cl₂ (9 mL) was added TFA (4.5 mL), and the reaction was stirred at RT for 2 h. Then the reaction was concentrated *in vacuo* and the residue was partitioned between CH₂Cl₂ and aqueous NaHCO₃, and extracted. The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated *in vacuo*. Purification via prep-HPLC column chromatography (CH₂Cl₂/EtOH/NH₄OH) provided the title compound as an off-white solid.

Applying the generic scheme below, the following examples of Table 21a were prepared as described above for (S)-2-(dimethylamino)ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (Example 66a), using the appropriate alkylating agent.





Table 21a.



Ex. No.	R ^X	R ^Y	R ^Z	CAS Name	LCMS (MH ⁺)
66a	H		H	(S)-2-(dimethylamino)ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	638
66b	H			(S)-2-(dimethylamino)-2-oxoethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	652
66c	H		H	(S)-2-(((R)-2-amino-3-methylbutanoyl)oxy)ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	710
66d	H		H	(S)-2-(pivaloyloxy)ethyl 8-(2-amino-6-((R)-1-(4-chloro-2-(3-methyl-1H-pyrazol-1-yl)phenyl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	695


Table 21b.

NMR Data for Compounds of Table 21a

Ex. No.	NMR
66a	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.59 (d, J=5.08 Hz, 4 H) 2.00 (dd, J=13.15, 9.84 Hz, 1 H) 2.22 - 2.38 (m, 4 H) 2.77 (d, J=3.37 Hz, 6 H) 3.14 (br. s., 2 H) 3.41 (br. s., 2 H) 3.60 (br. s., 2 H) 4.45 (dd, J=5.71, 3.90 Hz, 1 H) 4.49 - 4.68 (m, 2 H) 5.90 (br. s., 1 H) 6.39 (d, J=2.39 Hz, 1 H) 7.15 (d, J=5.86 Hz, 1 H) 7.53 - 7.73 (m, 3 H) 8.14 (d, J=2.39 Hz, 1 H) 9.65 (br. s., 1 H) 10.59 (br. s., 1 H), 10.80 (br. s., 1 H).
66b	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.46 - 1.77 (m, 4 H) 2.11 (dd, J=13.42, 8.40

	Hz, 1 H) 2.31 (s, 3 H) 2.38 (dd, J=13.42, 9.08 Hz, 1 H) 2.78 - 2.88 (m, 3 H) 2.89 - 2.98 (m, 3 H) 3.16 (br. s., 2 H) 3.59 - 3.77 (m, 3 H) 4.65 (t, J=6.17 Hz, 1 H) 4.83 - 4.97 (m, 1 H) 5.00 - 5.12 (m, 1 H) 6.03 (br. s., 1 H) 6.42 (d, J=2.29 Hz, 1 H) 7.20 (d, J=5.47 Hz, 1 H) 7.57 - 7.76 (m, 3 H) 8.17 (d, J=2.34 Hz, 1 H) 9.22 (d, J=4.44 Hz, 1 H) 10.63 (br. s., 1 H).
66c	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.98 (dd, J=15.52, 6.93 Hz, 6 H) 1.46 - 1.70 (m, 4 H) 1.94 (dd, J=13.15, 9.64 Hz, 1 H) 2.20 (td, J=6.91, 4.88 Hz, 1 H) 2.26 - 2.38 (m, 4 H) 3.14 (br. s., 2 H) 3.51 (br. s., 2 H) 3.58 - 3.70 (m, 3 H) 3.88 (br. s., 1 H) 4.29 - 4.49 (m, 4 H) 4.55 (br. s., 1 H) 5.84 (br. s., 1 H) 6.42 (d, J=2.34 Hz, 1 H) 7.16 (d, J=5.66 Hz, 1 H) 7.50 - 7.76 (m, 3 H) 8.17 (d, J=2.34 Hz, 1 H) 8.66 (br. s., 3 H) 9.47 (br. s., 1 H) 10.52 - 10.84 (m, 1 H).
66d	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.11 - 1.18 (m, 9 H) 1.48 (s, 3 H) 1.54 - 1.72 (m, 4 H) 1.74 - 2.01 (m, 1 H) 2.22 - 2.43 (m, 4 H) 3.15 (d, J=3.56 Hz, 2 H) 3.58 - 3.80 (m, 4 H) 4.60 (d, J=5.71 Hz, 1 H) 6.06 (br. s., 1 H) 6.42 (s, 1 H) 6.74 - 6.88 (m, 1 H) 7.22 (d, J=5.47 Hz, 1 H) 7.57 - 7.76 (m, 4 H) 8.18 (s, 1 H) 9.19 - 9.56 (m, 1 H) 10.74 (br. s., 1 H).

Example 67a: (S)-isopropyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate



5

To a solution of the compound of Example 1m (400 mg, 0.53 mmol) in propan-2-ol (5 mL) was added thionyl chloride (2 drops) at 0°C. The mixture was warmed to RT and then heated to reflux for 2 h. Then the reaction mixture was cooled to RT, concentrated and neutralized with saturated aqueous NaHCO₃ solution to pH 7-8. The aqueous layer was extracted with CH₂Cl₂.

10 The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, concentrated *in vacuo* and purified by flash column (0~10% MeOH in DCM) on silica gel to afford the title compound as a white solid.

¹H NMR (400 MHz, MeOH-d₄): δ ppm 7.96 (d, J = 2.3 Hz, 1H), 7.75 (d, J = 8.2 Hz, 1H), 7.70 (dd, J = 8.2, 1.8 Hz, 1H), 7.59 (d, J = 1.8 Hz, 1H), 7.43 (s, 1H), 7.37 (d, J = 7.8 Hz, 1H), 7.19 (d,

15

J = 8.0 Hz, 1H), 6.76 (q, J = 6.8 Hz, 1H), 6.41 (d, J = 2.3 Hz, 1H), 5.74 (s, 1H), 5.01 (m, 1H), 3.76 (dd, J = 8.7, 7.0 Hz, 1H), 3.61 – 3.42 (m, 4H), 2.88 (d, J = 11.1 Hz, 1H), 2.72 (d, J = 11.0 Hz, 1H), 2.39 (s, 3H), 2.31 (s, 3H), 2.29 (s, 3H), 2.05 (dd, J = 13.1, 8.9 Hz, 1H), 1.71 (dd, J = 13.0, 7.0 Hz, 1H), 1.50 (m, 4H), 1.24 (dd, J = 6.2, 3.9 Hz, 6H). LCMS (MH⁺): 679.

5

Applying the generic scheme below, the following examples of Table 22 were prepared as described above for (S)-isopropyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (Example 67a), using the appropriate alcohol.

10




Table 22a.

Ex. No.	R ¹	CAS Name	LCMS (MH ⁺)
67b		(S)-cyclopentyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	705
67c	CH ₃	(S)-methyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	650
67d		(S)-propyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	679

15


Table 22b.

NMR Data for Compounds of Table 22

Ex.	NMR
-----	-----

No.	
67b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.96 (d, J = 2.4 Hz, 1H), 7.75 (d, J = 8.2 Hz, 1H), 7.71 (dd, J = 8.2, 1.8 Hz, 1H), 7.60 (d, J = 1.7 Hz, 1H), 7.44 (s, 1H), 7.37 (dd, J = 7.9, 1.9 Hz, 1H), 7.20 (d, J = 7.8 Hz, 1H), 6.76 (q, J = 6.9 Hz, 1H), 6.41 (d, J = 2.3 Hz, 1H), 5.74 (s, 1H), 5.21 – 5.14 (m, 1H), 3.76 (dd, J = 8.8, 6.9 Hz, 1H), 3.61 – 3.42 (m, 4H), 2.88 (d, J = 11.0 Hz, 1H), 2.72 (d, J = 11.0 Hz, 1H), 2.39 (s, 3H), 2.31 (s, 3H), 2.28 (s, 3H), 2.04 (dd, J = 13.1, 8.8 Hz, 1H), 1.87 (d, J = 7.3 Hz, 2H), 1.77 – 1.56 (m, 7H), 1.50-1.45 (m, 4H)
67c	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 7.96 (d, J = 2.3 Hz, 1H), 7.76 (d, J = 8.3 Hz, 1H), 7.71 (dd, J = 8.2, 1.6 Hz, 1H), 7.60 (d, J = 1.6 Hz, 1H), 7.44 (s, 1H), 7.37 (dd, J = 7.8, 1.6 Hz, 1H), 7.20 (d, J = 7.9 Hz, 1H), 6.76 (q, J = 6.5 Hz, 1H), 6.41 (d, J = 2.3 Hz, 1H), 5.74 (s, 1H), 3.83 (t, J = 8.0 Hz, 1H), 3.71 (s, 3H), 3.61 – 3.41 (m, 4H), 2.86 (d, J = 11.0 Hz, 1H), 2.74 (d, J = 11.0 Hz, 1H), 2.39 (s, 3H), 2.31 (s, 3H), 2.28 (s, 3H), 2.06 (dd, J = 13.0, 8.7 Hz, 1H), 1.72 (dd, J = 13.0, 7.2 Hz, 1H), 1.55 – 1.43 (m, 4H)
67d	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.95 (m, 3H), 1.49 (dt, J = 12.0, 6.0 Hz, 4H), 1.69 (m, 3H), 2.06 (dd, J = 13.1, 8.8 Hz, 1H), 2.29 (d, J = 10.3 Hz, 6H), 2.39 (s, 3H), 2.73 (d, J = 11.0 Hz, 1H), 2.87 (d, J = 11.0 Hz, 1H), 3.30 (m, 4H), 3.51 (dt, J = 27.9, 6.6 Hz, 4H), 3.81 (dd, J = 8.7, 7.1 Hz, 1H), 4.08 (m, 2H), 5.74 (s, 1H), 6.41 (d, J = 2.3 Hz, 1H), 6.76 (q, J = 6.7 Hz, 1H), 7.19 (d, J = 7.9 Hz, 1H), 7.40 (m, 2H), 7.59 (d, J = 1.8 Hz, 1H), 7.72 (m, 2H), 7.96 (d, J = 2.4 Hz, 1H)

Example 68a: (S)-isopropyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate



The title compound was prepared as described for (S)-isopropyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (Example 67a) starting with (S)-8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 11).

Applying the generic scheme below, the following examples of Table 23 were prepared as described above for (S)-isopropyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (Example 68a), using the appropriate alcohol.




Table 23a.

Ex. No.	R ¹	CAS Name	LCMS (MH ⁺)
68a		(S)-isopropyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	709
68b		(S)-cyclopentyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	735
68c		(S)-propyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	709


Table 23b.

NMR Data for Compounds of Table 23

Ex. No.	NMR
68a	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.26 (m, 14H), 1.49 (dt, J = 10.9, 5.2 Hz, 4H), 1.72 (dd, J = 13.1, 7.0 Hz, 1H), 2.05 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.72 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.52 (m, 4H), 3.77 (dd, J = 8.8, 7.0 Hz, 1H), 4.63 (hept, J = 6.0 Hz, 1H), 5.01 (p, J = 6.2 Hz, 1H), 5.74 (s, 1H), 6.40 (d, J = 2.3 Hz, 1H), 6.76 (q, J = 6.6 Hz, 1H), 6.96 (m, 2H), 7.57 (m, 3H), 7.70 (m, 2H), 7.95 (d, J = 2.3 Hz, 1H)
68b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (d, J = 6.0 Hz, 8H), 1.50 (m, 4H), 1.67 (ddd, J = 33.0, 12.8, 5.6 Hz, 8H), 1.88 (m, 3H), 2.05 (dd, J = 13.1, 8.9 Hz, 1H), 2.39 (s, 3H), 2.73 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.52 (dt, J = 21.1, 6.5 Hz, 4H), 3.78 (dd, J = 8.8, 7.0 Hz, 1H), 4.64 (p, J = 6.0 Hz, 1H), 5.18 (td, J = 5.9, 2.7 Hz, 1H),

	5.75 (s, 1H), 6.40 (d, J = 2.4 Hz, 1H), 6.75 (q, J = 6.6 Hz, 1H), 6.97 (m, 2H), 7.59 (m, 3H), 7.71 (m, 2H), 7.95 (d, J = 2.4 Hz, 1H)
68c	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 0.94 (t, J = 7.4 Hz, 3H), 1.32 (d, J = 6.0 Hz, 6H), 1.50 (dt, J = 12.3, 6.0 Hz, 4H), 1.69 (m, 3H), 2.07 (dd, J = 13.1, 8.8 Hz, 1H), 2.39 (s, 3H), 2.73 (d, J = 11.0 Hz, 1H), 2.88 (d, J = 11.0 Hz, 1H), 3.52 (dp, J = 20.9, 7.5 Hz, 4H), 3.81 (dd, J = 8.7, 7.1 Hz, 1H), 4.09 (m, 2H), 4.64 (h, J = 6.0 Hz, 1H), 5.74 (s, 1H), 6.40 (d, J = 2.4 Hz, 1H), 6.76 (q, J = 6.7 Hz, 1H), 6.96 (m, 2H), 7.58 (m, 3H), 7.71 (m, 2H), 7.95 (d, J = 2.4 Hz, 1H)


Example 69a: (S)-isopropyl 8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate



- 5 The title compound was prepared as described for (S)-isopropyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (Example 67a) starting with (S)-8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 34c).

10

Applying the generic scheme below, the following examples of Table 24 were prepared as described above for (S)-isopropyl 8-(2-amino-6-((R)-2,2,2-trifluoro-1-(4'-isopropoxy-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)ethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (Example 68a), using the appropriate alcohol.



15

Table 24a.



Ex. No.	R ¹	CAS Name	LCMS (MH ⁺)
69a		(S)-isopropyl 8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	605
69b		(S)-cyclopentyl 8-(2-amino-6-((R)-1-(4'-chloro-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	735
69c		(S)-propyl 8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	605
69d		(S)-tetrahydro-2H-pyran-4-yl 8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate	645

Table 24b.

NMR Data for Compounds of Table 24

Ex. No.	NMR
69a	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.25 (dd, J = 6.3, 3.2 Hz, 6H), 1.52 (m, 4H), 1.74 (dd, J = 13.1, 7.1 Hz, 1H), 2.09 (dd, J = 13.1, 8.8 Hz, 1H), 2.75 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.49 (m, 4H), 3.80 (dd, J = 8.8, 7.1 Hz, 1H), 5.02 (hept, J = 6.2 Hz, 1H), 5.47 (d, J = 7.8 Hz, 1H), 6.63 (q, J = 6.8 Hz, 1H), 7.28 (d, J = 2.2 Hz, 1H), 7.48 (m, 6H), 7.67 (d, J = 8.5 Hz, 1H)
69b	¹ H NMR (400 MHz, MeOH-d ₄): δ ppm 1.32 (d, J = 6.0 Hz, 8H), 1.50 (m, 4H), 1.67 (ddd, J = 33.0, 12.8, 5.6 Hz, 8H), 1.88 (m, 3H), 2.05 (dd, J = 13.1, 8.9 Hz, 1H), 2.39 (s, 3H), 2.73 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.52 (dt, J = 21.1, 6.5 Hz, 4H), 3.78 (dd, J = 8.8, 7.0 Hz, 1H), 4.64 (p, J = 6.0 Hz, 1H), 5.18 (td, J = 5.9, 2.7 Hz, 1H), 5.75 (s, 1H), 6.40 (d, J = 2.4 Hz, 1H), 6.75 (q, J = 6.6 Hz, 1H), 6.97 (m, 2H), 7.59 (m, 3H), 7.71 (m, 2H), 7.95 (d, J = 2.4 Hz, 1H)
69c	¹ H NMR (MeOH-d ₄): δ ppm 0.95 (t, J = 7.4 Hz, 3H), 1.52 (dt, J = 14.2, 4.9 Hz, 4H), 1.71 (ddd, J = 31.8, 13.7, 7.1 Hz, 3H), 2.10 (dd, J = 13.1, 8.8 Hz, 1H), 2.76 (d, J = 11.0 Hz, 1H), 2.91 (d, J = 11.0 Hz, 1H), 3.50 (ddd, J = 19.5, 7.9, 4.8 Hz, 4H), 3.84 (dd, J = 8.7, 7.2 Hz, 1H), 4.10 (m, 2H), 4.88 (s, 8H), 5.48 (d, J = 7.9 Hz, 1H), 6.63 (q, J = 6.9 Hz, 1H), 7.28 (d, J = 2.2 Hz, 1H), 7.47 (m, 6H), 7.67 (d, J = 8.6 Hz, 1H)
69d	¹ H NMR (MeOH-d ₄): δ ppm 1.61 (m, 6H), 1.82 (dd, J = 13.2, 7.5 Hz, 1H), 1.93 (dd, J = 11.6, 6.1 Hz, 2H), 2.03 (s, 1H), 2.20 (dd, J = 13.2, 8.8 Hz, 1H), 2.89 (d, J = 11.2 Hz, 1H), 2.99 (d, J = 11.2 Hz, 1H), 3.54 (m, 6H), 3.89 (dq, J = 12.1, 3.9 Hz, 2H), 4.04 (dd, J = 8.7, 7.5 Hz, 1H), 5.01 (tt, J = 8.3, 4.0 Hz, 1H), 5.48 (s, 1H), 6.64 (q, J = 6.9 Hz, 1H), 7.28 (d, J = 2.2 Hz, 1H), 7.47 (m, 6H), 7.67 (d, J = 8.5 Hz, 1H)


Example 70: (S)-methyl 8-(2-amino-6-((R)-1-(5-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate



The title compound was prepared as described for (S)-isopropyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (Example 67a) starting with (S)-8-(2-amino-6-((R)-1-(5-chloro-3'-(methylsulfonyl)-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 34w).

¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.51 (q, J = 7.1, 6.7 Hz, 6H), 1.72 (dd, J = 13.0, 7.3 Hz, 1H), 2.07 (dd, J = 13.2, 8.7 Hz, 1H), 2.75 (d, J = 11.0 Hz, 1H), 2.87 (d, J = 11.0 Hz, 1H), 3.21 (s, 4H), 3.50 (tdt, J = 20.3, 13.5, 7.0 Hz, 4H), 3.71 (s, 2H), 3.84 (t, J = 8.0 Hz, 1H), 4.87 (m, 1H), 5.57 (s, 1H), 6.57 (q, J = 6.6 Hz, 1H), 7.33 (d, J = 2.3 Hz, 1H), 7.41 (s, 2H), 7.48 (dd, J = 8.5, 2.2 Hz, 1H), 7.75 (m, 3H), 8.07 (d, J = 7.8 Hz, 1H), 8.43 (s, 1H). LCMS (MH⁺): 655.

Example 71: (S)-methyl 8-(2-amino-6-((R)-1-(5-chloro-3'-sulfamoyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate



The title compound was prepared as described for (S)-isopropyl 8-(2-amino-6-((R)-1-(3',4'-dimethyl-3-(3-methyl-1H-pyrazol-1-yl)-[1,1'-biphenyl]-4-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-

yl)-2,8-diazaspiro[4.5]decane-3-carboxylate (Example 67a) starting with (S)-8-(2-amino-6-((R)-1-(5-chloro-3'-sulfamoyl-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid (Example 34u).

- 5 ¹H NMR (400 MHz, MeOH-d₄): δ ppm 1.54 (dt, J = 8.9, 6.0 Hz, 5H), 1.75 (dd, J = 13.1, 7.4 Hz, 1H), 2.10 (dd, J = 13.1, 8.7 Hz, 1H), 2.77 (d, J = 11.0 Hz, 1H), 2.89 (d, J = 11.0 Hz, 1H), 3.53 (qt, J = 14.0, 7.8 Hz, 4H), 3.72 (s, 3H), 3.86 (dd, J = 8.7, 7.3 Hz, 1H), 4.91 (s, 13H), 5.57 (s, 1H), 6.60 (q, J = 6.5 Hz, 1H), 7.31 (d, J = 2.2 Hz, 1H), 7.49 (dd, J = 8.5, 2.3 Hz, 1H), 7.60 (d, J = 7.7 Hz, 1H), 7.71 (m, 2H), 8.02 (ddd, J = 7.9, 1.9, 1.1 Hz, 1H), 8.33 (s, 1H). LCMS (MH⁺):656.
- 10 LCMS (MH⁺): 656.

Example A: *In vitro* Inhibition Assays

TPH1 and TPH2 Assays

- Recombinant human TPH1 (rTPH1 GenBank TM accession no. NP_004179) was
- 15 expressed by cloning full length human TPH1 cDNA in to a bacterial pMAL-c5E expression vector to produce maltose-binding protein (MBP) TPH1 fusion proteins. *E.coli* BL21 (DE3) containing pMAL-c5E-TPH1 was used for protein generation and the recombinant protein was purified utilizing standard column chromatography techniques. The MBP tagged TPH1 (MBP-TPH1) was used directly to screen compounds as described below. Recombinant human TPH2
- 20 (rTPH2 GenBank TM accession no. 173353), PheOH (rPheOH GenBank TM accession no. K03020) and TH (rTH GenBank TM accession no. L20679) with an MBP tag were produced similarly.

- TPH1 activities were measured in an assay containing 200 mM ammonium sulfate, 7 mM DTT, 50 µg/mL catalase, 25 µM ammonium iron sulfate, 50 mM MES, pH 7.1. Test
- 25 compounds were diluted in 100% DMSO and added to the assay plate in 1 µL aliquots at 100x final concentration. Fifty microliters of assay buffer containing 30 nM TPH1 enzyme (MBP tagged) were added to the plate wells containing the test compound by the use of an Eppendorf repeater pipette. The reaction was initiated by the addition of 50 µL of assay buffer containing 60 µM tryptophan and 72 µM 6-methyltetra-hydropterin (2x final concentration) by the use of
- 30 a Multidrop (LabSystems). Final reaction conditions were 15 nM TPH1 enzyme, 30 µM tryptophan, 36 µM 6-methyltetra-hydropterin, 200 mM ammonium sulfate, 7 mM DTT, 25

μg/mL catalase, 25 μM ferrous ammonium sulfate, 50 mM MES, pH 7.1, with atmospheric oxygen at room temperature. The plate was immediately placed onto an M5 plate reader (Molecular Devices) for kinetic fluorescence measurement using an excitation setting of 300 nm and an emission setting of 335 nm. Fluorescence reads are recorded in kinetic mode for 300 seconds (5 minutes).

Kinetic assay data for compounds at specific concentrations was translated into slopes using the Softmax Pro software on a Spectramax reader, and compound inhibition slopes were compared with wells containing enzyme, substrate and cofactor in the absence of inhibitor (100%), and wells containing substrate and cofactor in the absence of enzyme (0%). DMSO concentration in the assay was 1%. Typically, in the absence of enzyme, reaction slopes were ~0. IC₅₀'s were determined using Graphpad Prism.

Compounds having an IC₅₀ of 10,000 nM or less were considered active.

Inhibition of TPH2 activity by the compounds of the invention was measured similarly. In some instances, compounds of the invention showed dual inhibition of both TPH1 and TPH2.

Data related to TPH1 inhibition activity of the compounds of the invention is provided below in Table 25. Compounds that inhibit TPH1 with an IC₅₀ from 3,000 nM to 10,000 nM are indicated by +. Compounds that inhibit TPH1 with an IC₅₀ of less than 3,000 nM but more than 300 nM are indicated by ++. Compounds that inhibit TPH1 from 50 nM to 300 nM are indicated by +++. Compounds that inhibit TPH1 with an IC₅₀ less than 50 nM are indicated by ++++.

Ester prodrugs listed, for example, in Tables 18a, 19a, 20a, and 21a-24a, as well as in Examples 70 and 71, are not expected to be active in this *in vitro* assay.

Table 25. TPH1 Inhibition Data

Ex. No.	TPH1 Range	Ex. No.	TPH1 Range	Ex. No.	TPH1 Range	Ex. No.	TPH1 Range
1a	++++	19n	++++	34cs	+++	55cb	+++
1b	++++	19o	++++	34ct	++++	55cc	+++
1c	++++	19p	++++	34cu	++++	55cd	++
1d	++++	19q	++++	34cv	++++	55ce	++
1e	++++	19r	++++	35	+++	55cf	+
1f	+++	20	++++	36	+	55cg	++
1g	++++	21	++++	36b	+++	55ch	+++
1h	++++	22a	++++	36c	++++	55ci	++
1i	+++	22b	++++	36d	+	55cj	++
1j	+++	22c	++++	36e	++	55ck	++

1k	++++	23	++++	36f	+++	55cl	+++
1l	++++	24	+++	36g	++++	55cm	++
1m	++++	25	++++	37	+	55cn	++
1n	++++	26	+	38	++	55co	+++
1o	++++	27	+++	39a	++	55cp	++
1p	++++	28	+++	39b	+	55cq	+++
1q	++++	29a	++++	39c	+	55cr	++
1r	++++	29b	++++	39d	+	55cs	++
1s	++++	29c	+++	39e	++	55ct	++
1u	++++	29d	++++	40	++	55cu	++
1v	++++	29e	+++	41a	++	55cv	++
1w	++++	29f	++++	41b	++	55cw	++
1x	++++	29g	++++	41c	+	55cx	++
1y	++++	29h	++++	41d	+	55cy	++
1z	++++	29i	++++	42a	++++	55cz	+++
1aa	++++	29j	++++	42b	+++	55da	++
1ab	++++	29k	+++	43	+	55db	++
1ac	++++	29l	++++	44	++	55dc	++++
1ad	++++	29m	+++	45	+++	55dd	+++
1ae	++++	29n	+++	46	+	55de	+++
1af	++++	29o	++++	47	++	55df	+++
1ag	++++	29p	++++	48	++	55dg	+++
1ah	++++	29q	++++	49	++++	55dh	+++
1ai	++++	29r	++++	50	+++	55di	+++
1aj	++++	29s	++++	51	++++	55dj	+++
1ak	+++	29t	+++	52a	++++	55dk	+++
1al	++++	29u	+++	52b	++++	55dl	+++
1am	++++	33	+++	53	++++	55dm	+++
1an	++++	34a	++++	54a	++++	55dn	+++
1ao	++++	34b	++++	54b	+++	55do	++++
1ap	++++	34c	+++	54c	++++	55dp	++++
1aq	+++	34d	+++	54d	+++	55dq	++++
1ar	+++	34e	+++	54e	+++	56	+++
1as	++++	34f	+++	54f	++++	57	+++
1at	++++	34g	+++	54g	+++	58	+++
1au	+++	34h	++	54h	+	59	++
1av	++++	34i	+++	54i	+	59b	++
1aw	+++	34j	+++	54j	+++	59c	+++
1ax	++++	34k	+++	54k	+++	59d	+++
1ay	+++	34l	+++	54l	+	60	++
1az	+++	34m	+++	54m	++	61	+++
1ba	++++	34n	+++	55a	+++	62	+++
1bb	++++	34o	+++	55b	++++		
1bc	++++	34p	+++	55c	+++		
1bd	++++	34q	++++	55d	++++		

1be	+++	34r	+++	55e	+++
1bf	+++	34s	+++	55f	+++
1bg	++++	34t	+++	55g	++
1bh	++++	34u	++++	55h	+++
1bi	++++	34v	++++	55i	++++
1bj	+++	34w	++++	55j	+++
1bk	++++	34x	++++	55k	+++
1bl	++++	34y	++++	55l	+++
1bm	++++	34z	++++	55m	++++
1bn	++++	34aa	++++	55n	++
1bo	+++	34ab	++	55o	+++
1bp	++++	34ac	++++	55p	+++
1bq	++++	34ad	++	55q	++++
1bv	++++	34ae	++	55r	+++
1bw	+++	34af	++++	55s	++++
1bx	++++	34ag	++++	55t	+++
1by	++++	34ah	++++	55u	+++
1bz	++++	34ai	+++	55v	++++
1ca	++++	34aj	+++	55w	++++
1cb	++++	34ak	+++	55x	+++
1cc	++++	34al	+	55y	+++
1cd	++++	34am	+++	55z	+++
1ce	++++	34an	++++	55aa	+++
1cf	+++	34ao	++++	55ab	+++
1cg	++++	34ap	+++	55ac	+++
1ch	+++	34aq	++++	55ad	+++
1ci	++++	34ar	++++	55ae	++
1cj	++++	34as	++++	55af	++
1ck	++++	34at	+++	55ag	+++
1cl	++++	34au	++++	55ah	+++
1cm	++++	34av	++++	55ai	++++
1cn	++++	34aw	++++	55aj	+++
1co	++++	34ax	+++	55ak	++
1cp	++++	34ay	+++	55al	+++
1cq	++++	34az	++	55am	+++
1cr	++++	34ba	++++	55an	+++
1cs	++++	34bb	+++	55ao	+++
10j	+++	34bc	+++	55ap	+++
10k	+++	34bd	++++	55aq	+++
10l	+++	34be	++++	55ar	+++
10m	++++	34bf	+++	55as	+++
10n	++++	34bg	+++	55at	+++
10o	++++	34bh	++	55au	+++
10p	+++	34bi	++++	55av	+++
10q	++++	34bj	++++	55aw	++

10r	++++	34bk	+++	55ax	+++
10pa	+++	34bl	+++	55ay	++
11	+++	34bm	+++	55az	++++
12a	+++	34bn	+++	55ba	++++
12b	++++	34bo	+++	55bb	++++
12c	++++	34bp	++++	55bc	++++
13	+++	34bq	+++	55bd	++++
14	++++	34bu	+++	55be	++++
15	+++	34bv	+++	55bf	+++
16	++	34bw	+++	55bg	++++
17	++++	34bx	+++	55bh	+++
18a	+++	34by	+++	55bi	++++
18b	++++	34ca	+++	55bj	++++
18c	++++	34cb	+++	55bk	+++
18d	++++	34cc	+++	55bl	++
18e	++++	34cd	+++	55bm	++
18f	++++	34ce	+++	55bn	+
19a	++++	34cf	++	55bo	+++
19b	++++	34cg	+++	55bp	+++
19c	++++	34ch	++++	55bq	++
19d	+++	34ci	+++	55br	++
19e	++++	34cj	+++	55bs	++
19f	++++	34ck	+++	55bt	++
19g	++++	34cl	+++	55bu	++
19h	++++	34cm	+++	55bv	++
19i	++++	34cn	+++	55bw	+++
19j	++++	34co	+++	55bx	++
19k	++++	34cp	+++	55by	++
19l	++++	34cq	++++	55bz	+++
19m	++++	34cr	+++	55ca	+++

PheOH and TH inhibition counter assays

Certain compounds of the Examples were found to inhibit tryptophan hydroxylase (TPH) selectively over phenylalanine hydroxylase (PheOH). Inhibitory activity against PheOH can be assessed according to the methods described for example in *J. Med. Chem.* **10**, 64–66 (1967), or *J. Antibiot.* **35**, 458–462 (1982), or WO 2007/089335.

Certain compounds of the invention were found to inhibit tryptophan hydroxylase (TPH) selectively over tyrosine hydroxylase (TH). Inhibitory activity against TH can be assessed according to the methods described for example in *Life Sci.* **39**, 2185–2189 (1986), or *Mol. Pharmacol.* **41**, 339–344 (1992), or *J. Antibiot.* **35**, 458–462 (1982), or WO 2007/089335.

Example B: Intestinal 5-HT depletion assay

The efficacy of the TPH1 inhibitors of the invention was assessed for the ability to decrease intestinal serotonin concentration in mice. Mice (C57 BL6) were administered a single 150 mg/kg dose of test article by oral gavage. Each animal was euthanized by exsanguination under isoflurane anesthesia. Jejunal intestinal mucosa was isolated and homogenized in 300 μ L of a buffer containing 0.3M trichloroacetic acid, 0.1M sodium acetate, 10 mM EDTA, 20 mM sodium bisulfate and 50 mM ascorbic acid. Following centrifugation the 5-HT levels in the supernatants were measured by HPLC. The remaining mucosal pellet was solubilized overnight at 37 °C in a 0.1% sodium dodecyl sulfate buffer in 0.1N NaOH followed by determination of protein concentrations using a BCA protein assay (Pierce, Rockford, IL. 5-HT levels were normalized to protein and data were expressed as mean percent reduction of mucosal 5-HT levels relative to vehicle control $\pm$ SEM (percent 5-HT reduction). All animal studies were carried out with protocols approved by the Institutional Animal Care and Use Committee.

The Examples listed in Table 26 below were tested and found to elicit a reduction in mean mucosal 5-HT concentrations relative to vehicle-treated animals according to the above-described *in vivo* assay. P-values, indicating statistical significance of the data (ANOVA) are provided in the table: * refers to $P < 0.05$, ** refers to $P < 0.01$, *** refers to $P < 0.005$, and **** refers to $P < 0.0005$.

Table 26. *In Vivo* Efficacy of TPH1 Inhibitors In Mice (reduction of mucosal 5-HT concentrations one day after oral administration of a single 150 mg/kg dose)

Example No.	Efficacy	Example No.	Efficacy	Example No.	Efficacy
1g	***	34u	*	63bq	****
1h	**	34v	*	63bx	****
1l	****	34w	***	63by	***
1m	***	55k	*	63bz	**
1n	**	55ak	**	63ch	***
1o	**	55al	*	63cj	***
1p	**	55am	***	63cl	***
1y	**	55an	***	63cp	***

5	**	55az	***	63da	***
10b	***	55bc	**	63dc	***
10d	***	55bd	***	63di	***
10g	**	55bg	***	64c	****
10h	***	63g	***	64e	****
10j	****	63ay	***	64f	****
10k	*	63az	***	64g	*
11	****	63ba	***	64h	**
12b	*	63bd	***	65a	****
12c	***	63be	***	66c	****
16	**	63bf	****	66d	****
22c	*	63bg	****	101	***
28	*	63bh	**		
29z	**	63bi	**		
31	*	63bn	****		
34r	***	63bo	****		
34s	**	63bp	****		

Example C: Reduction of mucosal 5-HT concentrations

The Examples listed in Table 27 below were tested and found to elicit a reduction in mean mucosal 5-HT concentrations relative to vehicle-treated animals according to the following *in vivo* assay.

The efficacy of the TPH1 inhibitors of the invention was assessed for the ability to decrease intestinal serotonin concentration in mice. Mice (C57 BL6) were administered an oral dose of 10 or 50 mg/kg of the test article in the evening. Approximately 16 h following the first dose, mice were administered a second oral dose of 50 mg/kg of the appropriate compound. A third oral dose of 50 mg/kg of the appropriate test article was administered 12 h after dose 2. Following an overnight fast, each animal was euthanized by exsanguination under isoflurane anesthesia. Jejunal intestinal mucosa was isolated and homogenized in 300 mL of a buffer containing 0.3M trichloroacetic acid, 0.1M sodium acetate, 10 mM EDTA, 20 mM sodium bisulfate and 50 mM ascorbic acid. Following centrifugation the 5-HT levels in the supernatants were measured by HPLC. The remaining mucosal pellet was solubilized overnight at 37 °C in a 0.1% sodium dodecyl sulfate buffer in 0.1N NaOH followed by determination of protein concentrations using a BCA protein assay (Pierce, Rockford, IL). 5-HT levels were normalized

to protein and data were expressed as mean percent reduction of mucosal 5-HT levels relative to vehicle control $\pm$ SEM (percent 5-HT reduction). All animal studies were carried out with protocols approved by the Institutional Animal Care and Use Committee. P-values, indicating statistical significance of the data (ANOVA) are provided in the table: * refers to $P < 0.05$, ** refers to $P < 0.01$, *** refers to $P < 0.005$, and **** refers to $P < 0.0005$.

Table 27. *In Vivo* Efficacy of TPH1 Inhibitors In Mice (reduction of mucosal 5-HT concentrations two days after oral administration of a single 50 mg/kg dose)

Example No.	Efficacy	Example No.	Efficacy
1l	***	63cj	**
1m	*	63cl	***
1n	***	63cn	****
1t	***	63dc	***
12c	***	63el	***
55bg	***	63eo	***
63i	***	63ep	***
63ae	***	63ev	***
63aq	***	63ey	**
63ar	****	63fo	*
63aw	****	63ha	**
63az	***	64hb	***
63bd	***	69a	***
63bf	***	69b	***
63bg	***	69c	***
63bn	****		
63bo	***		
63bp	***		
63ch	***		

10

Example D: *In vivo* assay for inflammatory bowel diseases

The utility of the compounds of the invention for the treatment of inflammatory bowel diseases can be measured, for example, using the experimental models of colitis induced by

trinitrobenzene sulfonic acid (TNBS), dinitrobenzene sulfonic acid (DNBS), and dextran sodium sulfate (DSS), as described by Ghia, J.-E. et al. in *Gastroenterol.* **137**, 1649–60 (2009).

Example E: *In vivo* assay for low bone mass diseases

5 The utility of the compounds of the invention for the treatment of low bone mass diseases, such as osteoporosis, can be measured, for example, using the ovariectomy–induced osteopenia rat model, as described by Yadav, V. K. et al. in *Nature Med.* **16**, 308–12 (2010).

Example F: *In vivo* assay for PAH

10 The utility of the compounds of the invention for the treatment of pulmonary arterial hypertension (PAH), can be measured, for example, using the hypoxia mouse model, as described by Abid, S. et al. in *Am. J. Physiol., Lung Cellular and Molecular Physiology* **303**, L500–8 (2012), or using the rat monocrotaline-induced PAH or the rat chronic hypoxia model, as described by Kay, J. M. et al. *Respiration* **47**, 48–56 (1985).

15

Example G: *In vivo* assay for allergic airway inflammation

 The utility of the compounds of the invention for the treatment of allergic airway inflammation, can be measured, for example, using the mouse model of allergic asthma, as described by Dürk, T. et al. in *Am. J. Respir. Crit. Care Med.* **187**, 476–485 (2013).

20

Example H: *In vivo* assay for gastrointestinal disorders

 The utility of the compounds of the invention for the treatment of gastrointestinal disorders associated with dysregulation of the GI serotonergic system, such as chemotherapy-induced emesis and irritable bowel syndrome, can be measured, for example, using the a ferret
25 model of chemotherapy-induced emesis, as described by Liu, Q. et al. in *J. Pharmacol. Exp. Ther.* **325**, 47–55 (2008).

Example I: *In vivo* assay for tumor growth

 The utility of the compounds of the invention for the treatment of tumor growth, can be
30 measured, for example, using the the xenograft model of cholangiocarcinoma tumor growth, as described by Alpini, G. et al. in *Cancer Res.* **68**, 9184–93 (2008).

Example J: *In vivo* assay for leukemia

The utility of the compounds of the invention for the treatment and prevention of leukemia and other cancers of the blood, can be measured, for example, using the mouse
5 leukemia model, the osteoblast-deficient mouse model, or the murine model of acute myeloid leukemia, as described in WO 2013/074889.


Example K: *In vivo* assay for atherosclerosis

The utility of the compounds of the invention for the treatment of atherosclerosis, and the
10 reduction of plasma cholesterol and triglyceride levels, can be measured, for example, using the Apo E -/- or LDLR -/- mouse models of atherosclerotic plaque development, as described in WO 2012/058598.

Various modifications of the invention, in addition to those described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also
15 intended to fall within the scope of the appended claims. Each reference, including all patent, patent applications, and publications, cited in the present application is incorporated herein by reference in its entirety.

The Claims defining the Invention are as follows:

1. Use of a compound of Formula IIa:



IIa

or a pharmaceutically acceptable salt thereof, wherein:

Ring A is C₃₋₁₀ cycloalkyl, C₆₋₁₀ aryl, or 5 to 10-membered heteroaryl;

L is O or NR⁴;

R¹ is H, C₁₋₁₀ alkyl, or -(CR⁸R⁹)_pOC(O)R¹⁰, wherein said C₁₋₁₀ alkyl, is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from F, Cl, Br, CN, C₁₋₄ alkyl, and C₁₋₄ haloalkyl;

R² and R³ are selected from H, and C₁₋₄ haloalkyl;

wherein one of R² and R³ is H and the other of R² and R³ is a C₁₋₄ haloalkyl;

R⁴ is H or C₁₋₄ alkyl;

R⁵ is H;

R⁷ is C₁₋₄ alkyl or NR¹³R¹⁴;

R⁸ and R⁹ are each independently selected from H and C₁₋₄ alkyl;

R¹⁰ is C₁₋₆ alkyl optionally substituted by 1, 2 or 3 substituents independently selected from C₁₋₆ haloalkyl, C₃₋₁₀ cycloalkyl, OR^a, and NR^cR^d;

R¹¹ and R¹² are each independently selected from H and C₁₋₆ alkyl;

R¹³ is H or C₁₋₄ alkyl;

R¹⁴ is H, C₁₋₄ alkyl, C₃₋₇ cycloalkyl, or C(O)OR^{a1};

R¹⁵ is C₃₋₇ cycloalkyl or C₆₋₁₀ aryl;

R^A is H, Cy^1 , halo, C_{1-6} alkyl, C_{2-6} alkenyl, CN, NO_2 , OR^{a2} , SR^{a2} , $C(O)R^{b2}$, $C(O)NR^{c2}R^{d2}$, $C(O)OR^{a2}$, $OC(O)R^{b2}$, $OC(O)NR^{c2}R^{d2}$, $NR^{c2}R^{d2}$, $NR^{c2}C(O)R^{b2}$, $NR^{c2}C(O)OR^{a2}$, $NR^{c2}C(O)NR^{c2}R^{d2}$, $NR^{c2}S(O)R^{b2}$, $NR^{c2}S(O)_2R^{b2}$, $NR^{c2}S(O)_2NR^{c2}R^{d2}$, $S(O)R^{b2}$, $S(O)NR^{c2}R^{d2}$, $S(O)_2R^{b2}$, or $S(O)_2NR^{c2}R^{d2}$, wherein said C_{1-6} alkyl and C_{2-6} alkenyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from Cy^1 , halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, NO_2 , OR^{a2} , SR^{a2} , $C(O)R^{b2}$, $C(O)NR^{c2}R^{d2}$, $C(O)OR^{a2}$, $OC(O)R^{b2}$, $OC(O)NR^{c2}R^{d2}$, $NR^{c2}R^{d2}$, $NR^{c2}C(O)R^{b2}$, $NR^{c2}C(O)OR^{a2}$, $NR^{c2}C(O)NR^{c2}R^{d2}$, $NR^{c2}S(O)R^{b2}$, $NR^{c2}S(O)_2R^{b2}$, $NR^{c2}S(O)_2NR^{c2}R^{d2}$, $S(O)R^{b2}$, $S(O)NR^{c2}R^{d2}$, $S(O)_2R^{b2}$, and $S(O)_2NR^{c2}R^{d2}$;

R^B is H, Cy^2 , halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$, $C(O)OR^{a3}$, $OC(O)R^{b3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)OR^{a3}$, $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}S(O)R^{b3}$, $NR^{c3}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, $S(O)NR^{c3}R^{d3}$, $S(O)_2R^{b3}$, or $S(O)_2NR^{c3}R^{d3}$, wherein said C_{1-6} alkyl and C_{2-6} alkenyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from Cy^2 , halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$, $C(O)OR^{a3}$, $OC(O)R^{b3}$, $OC(O)NR^{c3}R^{d3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)OR^{a3}$, $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}S(O)R^{b3}$, $NR^{c3}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, $S(O)R^{b3}$, $S(O)NR^{c3}R^{d3}$, $S(O)_2R^{b3}$, and $S(O)_2NR^{c3}R^{d3}$;

R^C and R^D are independently selected from H, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, NO_2 , OR^{a4} , SR^{a4} , $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$, $NR^{c4}S(O)R^{b4}$, $NR^{c4}S(O)_2R^{b4}$, $NR^{c4}S(O)_2NR^{c4}R^{d4}$, $S(O)NR^{c4}R^{d4}$, $S(O)_2R^{b4}$, and $S(O)_2NR^{c4}R^{d4}$; wherein said C_{1-6} alkyl and C_{2-6} alkenyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} haloalkyl, CN, NO_2 , OR^{a4} , SR^{a4} , $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$, $NR^{c4}S(O)R^{b4}$, $NR^{c4}S(O)_2R^{b4}$, $NR^{c4}S(O)_2NR^{c4}R^{d4}$, $S(O)R^{b4}$, $S(O)NR^{c4}R^{d4}$, $S(O)_2R^{b4}$, and $S(O)_2NR^{c4}R^{d4}$;

Cy^1 and Cy^2 are each independently selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, each of which is optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from R^{Cy} ;

each R^{Cy} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, OR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)OR^{a5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}S(O)R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, and $S(O)_2NR^{c5}R^{d5}$, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C_{1-6} alkyl, CN, NO_2 , OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)OR^{a5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}S(O)R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, and $S(O)_2NR^{c5}R^{d5}$;

each of R^a , R^{a1} , R^{a2} , R^{a3} , R^{a4} , and R^{a5} is independently selected from H, C_{1-6} alkyl, C_{1-4} haloalkyl, C_{2-6} alkenyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-4} alkyl, C_{3-10} cycloalkyl- C_{1-4} alkyl, (5-10 membered heteroaryl)- C_{1-4} alkyl, or (4-10 membered heterocycloalkyl)- C_{1-4} alkyl, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-4} alkyl, C_{3-10} cycloalkyl- C_{1-4} alkyl, (5-10 membered heteroaryl)- C_{1-4} alkyl, and (4-10 membered heterocycloalkyl)- C_{1-4} alkyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-4} alkyl, halo, CN, OR^{a6} , $C(O)R^{b6}$, $C(O)NR^{c6}R^{d6}$, $C(O)OR^{a6}$, $OC(O)R^{b6}$, $OC(O)NR^{c6}R^{d6}$, $NR^{c6}R^{d6}$, $NR^{c6}C(O)R^{b6}$, $NR^{c6}C(O)NR^{c6}R^{d6}$, $NR^{c6}C(O)OR^{a6}$, $S(O)R^{b6}$, $S(O)NR^{c6}R^{d6}$, $S(O)_2R^{b6}$, $NR^{c6}S(O)_2R^{b6}$, $NR^{c6}S(O)_2NR^{c6}R^{d6}$, and $S(O)_2NR^{c6}R^{d6}$;

each of R^{b2} , R^{b3} , R^{b4} , and R^{b5} is independently selected from H, C_{1-6} alkyl, C_{1-4} haloalkyl, C_{2-6} alkenyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-4} alkyl, C_{3-10} cycloalkyl- C_{1-4} alkyl, (5-10 membered heteroaryl)- C_{1-4} alkyl, or (4-10 membered heterocycloalkyl)- C_{1-4} alkyl, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-4} alkyl, C_{3-10} cycloalkyl- C_{1-4} alkyl, (5-10 membered heteroaryl)- C_{1-4} alkyl, and (4-10 membered heterocycloalkyl)- C_{1-4} alkyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-4} alkyl, halo, CN, OR^{a6} , $C(O)R^{b6}$, $C(O)NR^{c6}R^{d6}$, $C(O)OR^{a6}$, $OC(O)R^{b6}$,

OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

each of R^c, R^d, R^{c1}, R^{d1}, R^{c2}, R^{d2}, R^{c3}, R^{d3}, R^{c4}, R^{d4}, R^{c5}, and R^{d5} is independently selected from H, C₁₋₆ alkyl, C₁₋₄ haloalkyl, C₂₋₆ alkenyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₄ alkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, (5-10 membered heteroaryl)-C₁₋₄ alkyl, or (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₄ alkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, (5-10 membered heteroaryl)-C₁₋₄ alkyl, and (4-10 membered heterocycloalkyl)-C₁₋₄ alkyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₄ alkyl, halo, CN, OR^{a6}, SR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)NR^{c6}R^{d6}, NR^{c6}C(O)OR^{a6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, and S(O)₂NR^{c6}R^{d6};

each of R^{a6}, R^{b6}, R^{c6}, and R^{d6} is independently selected from H, C₁₋₄ alkyl, C₂₋₄ alkenyl, C₃₋₇ cycloalkyl, phenyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl, wherein said C₁₋₄ alkyl, C₂₋₄ alkenyl, C₃₋₇ cycloalkyl, phenyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl are each optionally substituted by 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ alkylamino, and di(C₁₋₄ alkyl)amino; and

p is 1, 2, or 3;

wherein any aforementioned 4-10 or 4-7 membered heterocycloalkyl group optionally comprises 1, 2, or 3 oxo substituents, wherein each oxo substituent that is present is substituted on a ring-forming carbon, nitrogen, or sulfur atom of the 4-10 or 4-7 membered heterocycloalkyl group, or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating or preventing a disease selected from the group consisting of osteoporosis, osteoporosis pseudoglioma syndrome, osteopenia, osteomalacia, renal osteodystrophy, Paget's disease, fractures, and bone metastasis, atherosclerosis, pulmonary hypertension, idiopathic or familial pulmonary hypertension, pulmonary arterial hypertension, including idiopathic pulmonary arterial hypertension, familial pulmonary arterial hypertension, associated pulmonary arterial hypertension,

diabetes, hyperlipidemia, chronic obstructive pulmonary disease, pulmonary embolism, irritable bowel disease, colitis, chemotherapy-induced emesis, diarrhea, carcinoid syndrome, celiac disease, Crohn's disease, abdominal pain, dyspepsia, constipation, lactose intolerance, multiple endocrine neoplasia types I and II, Ogilvie's syndrome, pancreatic cholera syndrome, pancreatic insufficiency, pheochromacytoma, scleroderma, somatization disorder, Zollinger-Ellison Syndrome, liver diseases including chronic liver disease; cancers including liver cancer, breast cancer, cholangiocarcinoma, colon cancer, colorectal cancer, neuroendocrine tumors, pancreatic cancer, prostate cancer, bone cancer, osteosarcoma, chondrosarcoma, Ewings sarcoma, osteoblastoma, osteoid osteoma, osteochondroma, enchondroma, chondromyxoid fibroma, aneurysmal bone cyst, unicameral bone cyst, giant cell tumor, and bone tumors, myeloproliferative syndrome, myelodysplastic syndrome, Hodgkin's lymphoma, non-Hodgkin's lymphoma, myeloma, aplastic anemia, anemia associated with kidney disease, acute lymphocytic leukemia, chronic lymphocytic leukemia, acute myeloid leukemia, and chronic myeloid leukemia.

2. The use of claim 1, wherein said cardiovascular disease is pulmonary arterial hypertension.
3. The use of claim 1, wherein said pulmonary arterial hypertension is associated pulmonary arterial hypertension.
4. The use of claim 1, wherein said gastrointestinal disease is diarrhea or carcinoid syndrome.
5. The use of any one of claims 1-4, wherein the compound is (S)-ethyl 8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylate or a pharmaceutically acceptable salt thereof.

6. The use of any one of claims 1-4, wherein the compound is (S)-8-(2-amino-6-((R)-1-(5-chloro-[1,1'-biphenyl]-2-yl)-2,2,2-trifluoroethoxy)pyrimidin-4-yl)-2,8-diazaspiro[4.5]decane-3-carboxylic acid or a pharmaceutically acceptable salt thereof

ABSTRACT

The present invention is directed to the use of spirocyclic compounds which are inhibitors of tryptophan hydroxylase (TPH), particularly isoform 1 (TPH1), for the manufacture of medicaments for the treatment of diseases or disorders associated with peripheral serotonin including, for example, gastrointestinal, cardiovascular, pulmonary, inflammatory, metabolic, and low bone mass diseases, as well as serotonin syndrome, and cancer.

NO FIGURE