

ANTI-TIM3 ANTIBODIES AND METHODS OF USE

FIELD OF THE INVENTION

The present invention relates to anti-TIM3 antibodies and methods of using the same.

5 BACKGROUND

TIM3 is a human protein which belongs to the immunoglobulin superfamily, and TIM family of proteins. In humans, as similar to mice, TIM-3 is expressed on T-cells as well as phagocytic cells such as macrophages and dendritic cells. Binding of TIM-3 to a protein ligand (e.g., galectin-9) can inhibit the Th1 response via
10 mechanism of apoptosis induction, and therefore lead to such as induction of peripheral tolerance. The reduction in expression of human TIM-3 with siRNA or the inhibition of human TIM-3 by blocking-antibody increased the secretion of interferon alpha from CD4 positive T-cells, supporting the inhibitory role of TIM-3 in human T cells. In phagocytes, TIM-3 also functions as a receptor for recognizing
15 the apoptosis cells. Analysis of clinical samples from autoimmune disease patients showed no expression of TIM-3 in CD4 positive cells. In particular, in T cell clones derived from the cerebrospinal fluid of patients with multiple sclerosis, the expression level of TIM-3 was lower and the secretion level of IFN-gamma was higher than those of clones derived from normal healthy persons (Koguchi K et al.,
20 J Exp Med. 203 (2006) 1413-1418).

There are reports on relation of TIM-3 with allergic diseases or asthma (WO96/27603 and WO2003/063792).

According to the microarray analysis of hematopoietic stem cells from acute myeloid leukemia (hereinafter referred to as "AML") patients and normal
25 hematopoietic stem cells, TIM-3 is expressed on AML stem cells and therefore the analysis suggested involvement of TIM-3 in hematological malignancy (Majeti R et al., PNAS, 106 (2009) 3396-3401 and WO2009/091547).

Examples of the anti-TIM-3 monoclonal antibodies include anti-human TIM- 3 rat monoclonal antibody (Clone 344823, manufactured by R&D Systems) and anti-
30 human TIM-3 mouse monoclonal antibody (Clone F38-2E2, manufactured by R&D Systems). WO2013/06490 relates to anti-TIM-3 antibodies which show rapid internalization and immunoconjugates thereof for treating cancer and reducing

inflammation. US2012/189617 relates to anti-TIM-3 antibodies which exhibit higher effector activity such as an antibody-dependent cellular cytotoxicity (ADCC activity) for diseases relating to a human TIM-3 expressing cell.

SUMMARY

5 The disclosure provides anti-TIM3 antibodies and methods of using the same.

The disclosure provides an isolated antibody that binds to TIM3, wherein the antibody:

- 10
- induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™)) of at least 45% after 120 Minutes at 37 °C.

In one embodiment, the anti-TIM3 antibody according to the disclosure:

- 15
- competes for binding to TIM3 with an anti-Tim3 antibody comprising the VH of SEQ ID NO:7 and VL of SEQ ID NO: 8
 - binds to a human and cynomolgus TIM3
 - shows as immunoconjugate a cytotoxic activity on TIM3 expressing cells
 - induces interferon-gamma release.

20 In one embodiment the anti-TIM3 antibody according to the disclosure is a monoclonal antibody.

In one embodiment the anti-TIM3 antibody according to the disclosure is a human, humanized, or chimeric antibody.

25 In one embodiment the anti-TIM3 antibody according to the disclosure is an antibody fragment that binds to TIM3.

In one embodiment the anti-TIM3 antibody according to the disclosure which is Fab fragment.

In one embodiment the anti-TIM3 antibody according to the disclosure comprises

- 5 A) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:4; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or
- 10 B) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or
- 15 C) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or
- 20 D) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:13, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:14, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:15; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:16; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:17 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:18; or
- 25 E) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:21, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:22, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:23; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:24; (ii) HVR-L2
- 30

comprising the amino acid sequence of SEQ ID NO:25 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:26; or

- 5 F) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:29, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:30, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:31; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:32; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:33 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:34; or
- 10 G) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:39; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42; or
- 15 H) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:45, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:46, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:47; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:48; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:49 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:50; or
- 20 I) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:53, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:54, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:55; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:56; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:57 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:58; or
- 25 J) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:61, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:62, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:63; and (b) a VL domain comprising (i) HVR-L1
- 30

comprising the amino acid sequence of SEQ ID NO:64; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:65 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:66;

5 The disclosure provides an isolated antibody that binds to human TIM3, wherein the antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (ii) HVR-L2
10 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6.

The disclosure further provides an isolated antibody that binds to human TIM3, wherein the antibody comprises wherein the antibody comprises

15 i) a VH sequence of SEQ ID NO:79 and a VL sequence of SEQ ID NO:80,or

ii) a VH sequence of SEQ ID NO:81 and a VL sequence of SEQ ID NO:82.

The disclosure, in one aspect, provides an isolated antibody that binds to human TIM3, wherein the antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:39; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID
20 NO:41 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42.

The disclosure further provides an isolated antibody that binds to human TIM3, wherein the antibody comprises

30 i) a VH sequence of SEQ ID NO:83 and a VL sequence of SEQ ID NO:84,or

ii) a VH sequence of SEQ ID NO:85 and a VL sequence of SEQ ID NO:86.

In one embodiment the anti-TIM3 antibody according to the disclosure is a full length IgG1 antibody.

5 In one embodiment the anti-TIM3 antibody according to the disclosure is a full length IgG1 antibody with mutations L234A, L235A and P329G (numbering according to the EU index of Kabat).

Another aspect of the disclosure is an isolated nucleic acid encoding the antibody according to the disclosure.

Another aspect of the disclosure is a host cell comprising the nucleic acid according to the disclosure.

10 Another aspect of the disclosure is a method of producing an antibody comprising culturing the host cell according to the disclosure so that the antibody is produced.

Another aspect of the disclosure is an immunoconjugate comprising the antibody according to the disclosure and a cytotoxic agent.

15 In one preferred embodiment of the disclosure is such an immunoconjugate wherein the cytotoxic agent is Pseudomonas Exotoxin A or an Amatoxin.

Another aspect of the disclosure is a pharmaceutical formulation comprising the antibody according to the disclosure or the immunoconjugate according to the disclosure and a pharmaceutically acceptable carrier.

20 Another aspect of the disclosure is an antibody according to the disclosure or an immunoconjugate according to the disclosure for use as a medicament.

Another aspect of the disclosure is an antibody according to the disclosure or the immunoconjugate according to the disclosure for use in treating cancer.

25 Another aspect of the disclosure is the use of an antibody according to the disclosure or the immunoconjugate according to the disclosure in the manufacture of a medicament for the treatment of cancer.

The disclosure also relates to the use of an antibody according to the disclosure or the immunoconjugate according to the disclosure for use in the manufacture of a medicament for treatment of cancer.

5 The disclosure also relates to a method of treating an individual having cancer comprising administering to the individual an effective amount of an antibody according to the disclosure or the immunoconjugate according to the disclosure.

The anti-TIM3 antibodies of the present disclosure show highly valuable properties like a rapid and strong internalization on TIM3-expressing cancer cells, a strong

cytotoxic activity as immunoconjugate (e.g. when conjugated with Pseudomonas exotoxins or amatoxins). They are therefore useful therapeutics for the treatment of different cancers, especially blood tumors, like leukemias and lymphomas. Furthermore they show a strong immunestimulatory cytokine release in a Mixed Lymphocyte Reaction (MLR) and are therefore useful as immunestimulatory cancer therapy. In addition humanized antibody versions have improved binding and binding specificity to CD4 Tcells when compared to the parental antibodies.

BRIEF DESCRIPTION OF THE FIGURES

FIG 1A: Time dependent FACS based internalization of anti-TIM3 antibody Tim3_0022 (abbreviated as <TIM-3> Ab(022)) internalized into rec CHOK1 cells expressing huTIM-3 after incubation at 37°C.

FIG 1B: Results from the FACS based internalization assay show that Fab fragment of anti-TIM3 antibody Tim3_0022 (abbreviated as <TIM-3> Ab(022)) internalized into rec CHOK1 cells expressing huTIM-3 after incubation at 37°C with similar kinetic as full IgG format.

FIG. 2A: Binding of anti-TIM3 antibodies to RPMI-8226 cells (antibody designation clone 0016 refers to antibody Tim3_0016, clone 0016 refers to antibody Tim3_0016 variant (antibody Tim3_0018), clone 0022 refers to antibody Tim3_00122, etc.).

FIG. 2B: Binding of anti-TIM3 antibodies to Pfeiffer cells (antibody designation clone 0016 refers to antibody Tim3_0016, clone 0016 refers to antibody Tim3_0016 variant (antibody Tim3_0018), clone 0022 refers to antibody Tim3_00122, etc.).

FIG 3: expression level of TIM-3 on different patient AML cell samples by FACS using anti-TIM-3 mAbs.

FIG 4: Direct omparison of binding of TIM3 antibodies to different peripheral blood mononuclear cells (Monocytes, NK cells, T cells, CD4 T cells):

4A: % positive cells to which Tim3_0016 variant (antibody Tim3_0018) and humanized versions are binding to.

4B: Mean fluorescence inensity- binding of Tim3_0016 variant (antibody Tim3_0018) and humanized versions to different peripheral blood mononuclear cells.

4C: % positive cells to which Tim3_0028 and chimeric and humanized versions are binding to.

4D: Mean fluorescence intensity- binding of Tim3_0028 and chimeric and humanized versions to different peripheral blood mononuclear cells.

5 DETAILED DESCRIPTION OF EMBODIMENTS OF THE INVENTION

I. DEFINITIONS

An “acceptor human framework” for the purposes herein is a framework comprising the amino acid sequence of a light chain variable domain (VL) framework or a heavy chain variable domain (VH) framework derived from a human immunoglobulin framework or a human consensus framework, as defined
10 below. An acceptor human framework “derived from” a human immunoglobulin framework or a human consensus framework may comprise the same amino acid sequence thereof, or it may contain amino acid sequence changes. In some embodiments, the number of amino acid changes are 10 or less, 9 or less, 8 or less,
15 7 or less, 6 or less, 5 or less, 4 or less, 3 or less, or 2 or less. In some embodiments, the VL acceptor human framework is identical in sequence to the VL human immunoglobulin framework sequence or human consensus framework sequence. (define germlines if appropriate)

The terms “anti-TIM3 antibody” and “an antibody that binds to TIM3” refer to an
20 antibody that is capable of binding TIM3 with sufficient affinity such that the antibody is useful as a diagnostic and/or therapeutic agent in targeting TIM3. In one embodiment, the extent of binding of an anti-TIM3 antibody to an unrelated, non-TIM3 protein is less than about 10% of the binding of the antibody to TIM3 as measured, e.g., by a Surface Plasmon Resonance assay (e.g. BIACORE). In certain
25 embodiments, an antigen binding protein that binds to human TIM3 has a KD value of the binding affinity for binding to human TIM3 of $\leq 1 \mu\text{M}$, $\leq 100 \text{ nM}$, $\leq 10 \text{ nM}$, $\leq 1 \text{ nM}$, $\leq 0.1 \text{ nM}$, $\leq 0.01 \text{ nM}$, or $\leq 0.001 \text{ nM}$ (e.g. 10^{-8} M or less, e.g. from 10^{-8} M to 10^{-13} M , e.g., from 10^{-9} M to 10^{-13} M). In one preferred embodiment the respective KD value of the binding affinities is determined in a Surface Plasmon
30 Resonance assay using the Extracellular domain (ECD) of human TIM3 (TIM3 - ECD) for the TIM3 binding affinity.

The term “antibody” herein is used in the broadest sense and encompasses various antibody structures, including but not limited to monoclonal antibodies, polyclonal

antibodies, multispecific antibodies (e.g., bispecific antibodies), and antibody fragments so long as they exhibit the desired antigen-binding activity.

5 An "antibody fragment" refers to a molecule other than an intact antibody that comprises a portion of an intact antibody that binds the antigen to which the intact antibody binds. Examples of antibody fragments include but are not limited to Fv, Fab, Fab', Fab'-SH, F(ab')₂; diabodies; linear antibodies; single-chain antibody molecules (e.g. scFv); and multispecific antibodies formed from antibody fragments.

10 An "antibody that binds to the same epitope" as a reference antibody refers to an antibody that blocks binding of the reference antibody to its antigen in a competition assay by 50% or more, and conversely, the reference antibody blocks binding of the antibody to its antigen in a competition assay by 50% or more. An exemplary competition assay is provided herein.

15 The term "chimeric" antibody refers to an antibody in which a portion of the heavy and/or light chain is derived from a particular source or species, while the remainder of the heavy and/or light chain is derived from a different source or species.

20 The "class" of an antibody refers to the type of constant domain or constant region possessed by its heavy chain. There are five major classes of antibodies: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG₁, IgG₂, IgG₃, IgG₄, IgA₁, and IgA₂. The heavy chain constant domains that correspond to the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively.

25 The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents a cellular function and/or causes cell death or destruction. Cytotoxic agents include, but are not limited to, radioactive isotopes (e.g., At211, I131, I125, Y90, Re186, Re188, Sm153, Bi212, P32, Pb212 and radioactive isotopes of Lu); chemotherapeutic agents or drugs (e.g., methotrexate, adriamycin, vinca alkaloids (vincristine, vinblastine, etoposide), doxorubicin, melphalan, mitomycin C, chlorambucil, daunorubicin or other intercalating agents); growth inhibitory agents;
30 enzymes and fragments thereof such as nucleolytic enzymes; antibiotics; toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof; and the various antitumor or anticancer agents disclosed below.

In one preferred embodiment the "cytotoxic agent" is a *Pseudomonas* exotoxin A (or variants thereof) WO2005052006, WO2007016150, WO2007014743, WO2007031741, WO200932954, WO201132022, WO2012/154530, and WO 2012/170617, Liu W, et al, PNAS 109 (2012) 11782-11787, Mazor R, et al
5 PNAS 111 (2014) 8571-8576 and Alewine C, et al, Mol Cancer Ther. (2014) 2653-61. In one preferred embodiment the "Pseudomonas exotoxin A" comprises the amino acid sequences of SEQ ID NO:69 or comprises the amino acid sequences of SEQ ID NO:70 (their preparation is also described in Mazor R, et al PNAS 111 (2014) 8571-8576 and Alewine C, et al, Mol Cancer Ther. (2014) 2653-61).

10 In another preferred embodiment the "cytotoxic agent" is an amatoxin (or variants thereof) as described e.g. WO2010/115630, WO2010/115629, WO2012/119787, WO2012/041504, and WO2014135282 with preferred variants described in WO2012/041504 (e.g. conjugated via the 6' C-atom of amatoxin amino acid 4, particularly via an oxygen atom bound to the 6' C-atom of amatoxin amino acid,
15 and wherein the TIM3 antibody is connected by a linker via a urea moiety) and WO2014135282.

An "effective amount" of an agent, e.g., a pharmaceutical formulation, refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic or prophylactic result.

20 The term "Fc region" herein is used to define a C-terminal region of an immunoglobulin heavy chain that contains at least a portion of the constant region. The term includes native sequence Fc regions and variant Fc regions. In one embodiment, a human IgG heavy chain Fc region extends from Cys226, or from Pro230, to the carboxyl-terminus of the heavy chain. However, the C-terminal
25 lysine (Lys447) of the Fc region may or may not be present. Unless otherwise specified herein, numbering of amino acid residues in the Fc region or constant region is according to the EU numbering system, also called the EU index, as described in Kabat, E.A. et al., Sequences of Proteins of Immunological Interest, 5th ed., Public Health Service, National Institutes of Health, Bethesda, MD (1991),
30 NIH Publication 91-3242.

"Framework" or "FR" refers to variable domain residues other than hypervariable region (HVR) residues. The FR of a variable domain generally consists of four FR domains: FR1, FR2, FR3, and FR4. Accordingly, the HVR and FR sequences

generally appear in the following sequence in VH (or VL): FR1-H1(L1)-FR2-H2(L2)-FR3-H3(L3)-FR4.

5 The terms “full length antibody,” “intact antibody,” and “whole antibody” are used herein interchangeably to refer to an antibody having a structure substantially similar to a native antibody structure or having heavy chains that contain an Fc region as defined herein.

10 The terms “host cell,” “host cell line,” and “host cell culture” are used interchangeably and refer to cells into which exogenous nucleic acid has been introduced, including the progeny of such cells. Host cells include “transformants” and “transformed cells,” which include the primary transformed cell and progeny derived therefrom without regard to the number of passages. Progeny may not be completely identical in nucleic acid content to a parent cell, but may contain mutations. Mutant progeny that have the same function or biological activity as screened or selected for in the originally transformed cell are included herein.

15 A “human antibody” is one which possesses an amino acid sequence which corresponds to that of an antibody produced by a human or a human cell or derived from a non-human source that utilizes human antibody repertoires or other human antibody-encoding sequences. This definition of a human antibody specifically excludes a humanized antibody comprising non-human antigen-binding residues.

20 A “human consensus framework” is a framework which represents the most commonly occurring amino acid residues in a selection of human immunoglobulin VL or VH framework sequences. Generally, the selection of human immunoglobulin VL or VH sequences is from a subgroup of variable domain sequences. Generally, the subgroup of sequences is a subgroup as in Kabat, E.A. et al., Sequences of Proteins of Immunological Interest, 5th ed., Bethesda MD (1991), NIH Publication 91-3242, Vols. 1-3. In one embodiment, for the VL, the subgroup is subgroup kappa I as in Kabat et al., *supra*. In one embodiment, for the VH, the subgroup is subgroup III as in Kabat et al., *supra*.

30 A “humanized” antibody refers to a chimeric antibody comprising amino acid residues from non-human HVRs and amino acid residues from human FRs. In certain embodiments, a humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the HVRs (e.g., CDRs) correspond to those of a non-human antibody, and all or substantially all of the FRs correspond to those of a human antibody. A humanized

antibody optionally may comprise at least a portion of an antibody constant region derived from a human antibody. A “humanized form” of an antibody, e.g., a non-human antibody, refers to an antibody that has undergone humanization.

5 The term “hypervariable region” or “HVR” as used herein refers to each of the regions of an antibody variable domain which are hypervariable in sequence (“complementarity determining regions” or “CDRs”) and/or form structurally defined loops (“hypervariable loops”) and/or contain the antigen-contacting residues (“antigen contacts”). Generally, antibodies comprise six HVRs: three in the VH (H1, H2, H3), and three in the VL (L1, L2, L3). Exemplary HVRs herein
10 include:

(a) hypervariable loops occurring at amino acid residues 26-32 (L1), 50-52 (L2), 91-96 (L3), 26-32 (H1), 53-55 (H2), and 96-101 (H3) (Chothia and Lesk, *J. Mol. Biol.* 196:901-917 (1987));

15 (b) CDRs occurring at amino acid residues 24-34 (L1), 50-56 (L2), 89-97 (L3), 31-35b (H1), 50-65 (H2), and 95-102 (H3) (Kabat et al., *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991));

20 (c) antigen contacts occurring at amino acid residues 27c-36 (L1), 46-55 (L2), 89-96 (L3), 30-35b (H1), 47-58 (H2), and 93-101 (H3) (MacCallum et al. *J. Mol. Biol.* 262: 732-745 (1996)); and

(d) combinations of (a), (b), and/or (c), including HVR amino acid residues 46-56 (L2), 47-56 (L2), 48-56 (L2), 49-56 (L2), 26-35 (H1), 26-35b (H1), 49-65 (H2), 93-102 (H3), and 94-102 (H3).

25 Unless otherwise indicated, HVR residues and other residues in the variable domain (e.g., FR residues) are numbered herein according to Kabat et al., *supra*.

30 An “immunoconjugate” is an antibody conjugated to one or more heterologous molecule(s), including but not limited to a cytotoxic agent. In one preferred embodiment an immunoconjugate is an anti-TIM3 antibody as described herein conjugated to one or more cytotoxic agents (preferably a *Pseudomonas* Exotoxin A or a amatoxin).

An “individual” or “subject” is a mammal. Mammals include, but are not limited to, domesticated animals (e.g., cows, sheep, cats, dogs, and horses), primates (e.g.,

humans and non-human primates such as monkeys), rabbits, and rodents (e.g., mice and rats). In certain embodiments, the individual or subject is a human.

5 An "isolated" antibody is one which has been separated from a component of its natural environment. In some embodiments, an antibody is purified to greater than 95% or 99% purity as determined by, for example, electrophoretic (e.g., SDS-PAGE, isoelectric focusing (IEF), capillary electrophoresis) or chromatographic (e.g., ion exchange or reverse phase HPLC). For review of methods for assessment of antibody purity, see, e.g., Flatman, S. et al., J. Chromatogr. B 848 (2007) 79-87.

10 An "isolated" nucleic acid refers to a nucleic acid molecule that has been separated from a component of its natural environment. An isolated nucleic acid includes a nucleic acid molecule contained in cells that ordinarily contain the nucleic acid molecule, but the nucleic acid molecule is present extrachromosomally or at a chromosomal location that is different from its natural chromosomal location.

15 "Isolated nucleic acid encoding an anti-TIM3antibody" refers to one or more nucleic acid molecules encoding antibody heavy and light chains (or fragments thereof), including such nucleic acid molecule(s) in a single vector or separate vectors, and such nucleic acid molecule(s) present at one or more locations in a host cell.

20 The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical and/or bind the same epitope, except for possible variant antibodies, e.g., containing naturally occurring mutations or arising during production of a monoclonal antibody preparation, such variants generally being present in minor amounts. In contrast to polyclonal antibody preparations, which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody of a monoclonal antibody preparation is directed against a single determinant on an antigen. Thus, the modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by a variety of techniques, including but not limited to the hybridoma method, recombinant DNA methods, phage-display methods, and methods utilizing transgenic animals containing all or part of the human

25

30

immunoglobulin loci, such methods and other exemplary methods for making monoclonal antibodies being described herein.

A “naked antibody” refers to an antibody that is not conjugated to a heterologous moiety (e.g., a cytotoxic moiety) or radiolabel. The naked antibody may be present in a pharmaceutical formulation. (Include if Prior art has immunoconjugates).

"Native antibodies" refer to naturally occurring immunoglobulin molecules with varying structures. For example, native IgG antibodies are heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light chains and two identical heavy chains that are disulfide-bonded. From N- to C-terminus, each heavy chain has a variable region (VH), also called a variable heavy domain or a heavy chain variable domain, followed by three constant domains (CH1, CH2, and CH3). Similarly, from N- to C-terminus, each light chain has a variable region (VL), also called a variable light domain or a light chain variable domain, followed by a constant light (CL) domain. The light chain of an antibody may be assigned to one of two types, called kappa (κ) and lambda (λ), based on the amino acid sequence of its constant domain.

The term “package insert” is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, dosage, administration, combination therapy, contraindications and/or warnings concerning the use of such therapeutic products.

“Percent (%) amino acid sequence identity” with respect to a reference polypeptide sequence is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the reference polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for aligning sequences, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are generated using the sequence comparison computer program ALIGN-2. The ALIGN-2

sequence comparison computer program was authored by Genentech, Inc., and the source code has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available from Genentech, Inc., South San Francisco, California, or may be compiled from the source code. The ALIGN-2 program should be compiled for use on a UNIX operating system, including digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

In situations where ALIGN-2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program.

The term "pharmaceutical formulation" refers to a preparation which is in such form as to permit the biological activity of an active ingredient contained therein to be effective, and which contains no additional components which are unacceptably toxic to a subject to which the formulation would be administered.

A "pharmaceutically acceptable carrier" refers to an ingredient in a pharmaceutical formulation, other than an active ingredient, which is nontoxic to a subject., A pharmaceutically acceptable carrier includes, but is not limited to, a buffer, excipient, stabilizer, or preservative.

The term "TIM3," as used herein, refers to any native T cell immunoglobulin mucin 3 (TIM3) protein (also known as hepatitis A virus cellular receptor 2

(HAVcr-2), kidney injury molecule-3 (KIM-3), TIM-3, Tim3, or Tim-3) from any vertebrate source, including mammals such as primates (e.g. humans) and rodents (e.g., mice and rats), unless otherwise indicated. The term encompasses "full-length," unprocessed TIM3 as well as any form of TIM3 that results from processing in the cell. The term also encompasses naturally occurring variants of TIM3, e.g., splice variants or allelic variants. The amino acid sequence of an exemplary human TIM3 is shown in SEQ ID NO:77. The amino acid sequence of the Extracellular Domain (ECD) of TIM3 is shown in SEQ ID NO:78.

TIM3 is a human protein which belongs to the immunoglobulin superfamily, and TIM family of proteins. In humans, as similar to mice, TIM-3 is expressed on T-cells as well as phagocytic cells such as macrophages and dendritic cells. Binding of TIM-3 to a protein ligand (e.g., galectin-9) can inhibit the Th1 response via mechanism of apoptosis induction, and therefore lead to such as induction of peripheral tolerance. The reduction in expression of human TIM-3 with siRNA or the inhibition of human TIM-3 by blocking-antibody increased the secretion of interferon alpha from CD4 positive T-cells, supporting the inhibitory role of TIM-3 in human T cells. In phagocytes, TIM-3 also functions as a receptor for recognizing the apoptosis cells. Analysis of clinical samples from autoimmune disease patients showed no expression of TIM-3 in CD4 positive cells. In particular, in T cell clones derived from the cerebrospinal fluid of patients with multiple sclerosis, the expression level of TIM-3 was lower and the secretion level of IFN- γ was higher than those of clones derived from normal healthy persons (Koguchi K et al., J Exp Med. 203 (2006) 1413-1418).

There are reports on relation of TIM-3 with allergic diseases or asthma (WO96/27603 and WO2003/063792).

According to the microarray analysis of hematopoietic stem cells from acute myeloid leukemia (hereinafter referred to as "AML") patients and normal hematopoietic stem cells, TIM-3 is expressed on AML stem cells and therefore the analysis suggested involvement of TIM-3 in hematological malignancy (Majeti R et al., PNAS, 106 (2009) 3396-3401 and WO2009/091547).

Examples of the anti-TIM-3 monoclonal antibodies include anti-human TIM-3 rat monoclonal antibody (Clone 344823, manufactured by R&D Systems) and anti-human TIM-3 mouse monoclonal antibody (Clone F38-2E2, manufactured by R&D Systems).

As used herein, "treatment" (and grammatical variations thereof such as "treat" or "treating") refers to clinical intervention in an attempt to alter the natural course of the individual being treated, and can be performed either for prophylaxis or during the course of clinical pathology. Desirable effects of treatment include, but are not limited to, preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease, preventing metastasis, decreasing the rate of disease progression, amelioration or palliation of the disease state, and remission or improved prognosis. In some embodiments, antibodies of the invention are used to delay development of a disease or to slow the progression of a disease.

The term "variable region" or "variable domain" refers to the domain of an antibody heavy or light chain that is involved in binding the antibody to antigen. The variable domains of the heavy chain and light chain (VH and VL, respectively) of a native antibody generally have similar structures, with each domain comprising four conserved framework regions (FRs) and three hypervariable regions (HVRs). (See, e.g., Kindt, T.J. et al. Kuby Immunology, 6th ed., W.H. Freeman and Co., N.Y. (2007), page 91) A single VH or VL domain may be sufficient to confer antigen-binding specificity. Furthermore, antibodies that bind a particular antigen may be isolated using a VH or VL domain from an antibody that binds the antigen to screen a library of complementary VL or VH domains, respectively. See, e.g., Portolano, S. et al., J. Immunol. 150 (1993) 880-887; Clackson, T. et al., Nature 352 (1991) 624-628).

The term "vector," as used herein, refers to a nucleic acid molecule capable of propagating another nucleic acid to which it is linked. The term includes the vector as a self-replicating nucleic acid structure as well as the vector incorporated into the genome of a host cell into which it has been introduced. Certain vectors are capable of directing the expression of nucleic acids to which they are operatively linked. Such vectors are referred to herein as "expression vectors".

II. COMPOSITIONS AND METHODS

In one aspect, the invention is based, in part, on the finding that the selected anti-TIM3 antibodies of the invention bind to certain epitopes of TIM3, showing a strong and rapid internalization and/or high cytotoxic activity against cancer cells as immunoconjugate and/or showing strong immunestimulatory cytokine (e.g. interferon gamma) release. In certain embodiments, antibodies that bind to TIM3

are provided. Antibodies of the invention are useful, e.g., for the diagnosis or treatment of cancer.

A. Exemplary Anti-TIM3 Antibodies

In one aspect, the invention provides isolated antibodies that bind to TIM3. In certain embodiments, an anti- TIM3 is provided wherein the antibody:

- induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™)) of at least 45% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™)) of at least 50% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™)) of at least 55% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™)) of at least 60% after 240 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™)) of at least 65% after 240 Minutes at 37 °C (see Example 6)

In certain embodiments, an anti- TIM3 is provided, wherein the antibody:

- competes for binding to TIM3 with an anti-Tim3 antibody comprising the VH and VL of Tim3_0016
- binds to a human and cynomolgus TIM3
- shows as immunoconjugate a cytotoxic activity on TIM3 expressing cells (in one embodiment the immunoconjugates has a relative IC50 value of the cytotoxic activity as Pseudomonas exotoxin A conjugate on RPMI-8226 cells of 0.1 or lower (as measured in Example 11)
- induces interferon-gamma release (in MLR assay –see Example 5).

In one aspect, the invention provides an anti-TIM3 antibody comprising at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1; (b) HVR-H2 comprising the amino acid

sequence of SEQ ID NO:2; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:3; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:4; or HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6.

In one aspect, the invention provides an anti-TIM3 antibody comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:3; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:4; or HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; or HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6.

In one aspect, the invention provides an anti-TIM3 antibody comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:3; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:4; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6.

In one aspect, the invention provides an anti-TIM3 antibody comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:3; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6.

In one aspect, the invention provides an anti-TIM3 antibody comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:3; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6.

In another aspect, an antibody of the invention comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from

(i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:4; or HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; or HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6.

In another aspect, an antibody of the invention comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:4; or HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; or HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6.

In another aspect, an antibody of the invention comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:4; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6.

In another aspect, an antibody of the invention comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6.

In another aspect, an antibody of the invention comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii)

HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6.

In one embodiment such anti-TIM3 antibody comprises

- i) a VH sequence of SEQ ID NO:7 and a VL sequence of SEQ ID NO:8;
- ii) a VH sequence of SEQ ID NO:9 and a VL sequence of SEQ ID NO:10;
- iii) or humanized variant of the VH and VL of the antibody under i) or ii).

In one preferred embodiment such anti-TIM3 antibody comprises

- i) a VH sequence of SEQ ID NO:79 and a VL sequence of SEQ ID NO:80, or
- ii) a VH sequence of SEQ ID NO:81 and a VL sequence of SEQ ID NO:82.

One preferred embodiment is an antibody that binds to human TIM3 antibody wherein the antibody comprises a VH sequence of SEQ ID NO:79 and a VL sequence of SEQ ID NO:80.

One preferred embodiment is an antibody that binds to human TIM3 antibody wherein the antibody comprises a VH sequence of SEQ ID NO:81 and a VL sequence of SEQ ID NO:82.

In one aspect, the invention provides an anti-TIM3 antibody comprising at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:13; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:14; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:15; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:16; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:17; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:18.

In one aspect, the invention provides an anti-TIM3 antibody comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:13; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:14; (c) HVR-H3 comprising

the amino acid sequence of SEQ ID NO:15; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:16; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:17; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:18.

5 In another aspect, an antibody of the invention comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:13, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:14, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:15; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:16; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:17 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:18.

15 In another aspect, an antibody of the invention comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:13, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:14, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:15; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:16; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:17 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:18.

In one embodiment such anti-TIM3 antibody comprises

i) a VH sequence of SEQ ID NO:19 and a VL sequence of SEQ ID NO:20;

ii) or humanized variant of the VH and VL of the antibody under i).

25 In one aspect, the invention provides an anti-TIM3 antibody comprising at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:21; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:22; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:23; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:24; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:25; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:26.

In one aspect, the invention provides an anti-TIM3 antibody comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:21; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:22; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:23; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:24; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:25; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:26.

In another aspect, an antibody of the invention comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:21, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:22, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:23; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:24; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:25 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:26.

In another aspect, an antibody of the invention comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:21, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:22, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:23; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:24; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:25 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:26.

In one embodiment such anti-TIM3 antibody comprises

- i) a VH sequence of SEQ ID NO:27 and a VL sequence of SEQ ID NO:28;
- ii) or humanized variant of the VH and VL of the antibody under i).

In one aspect, the invention provides an anti-TIM3 antibody comprising at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:29; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:30; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:31; (d) HVR-L1 comprising the amino acid sequence of SEQ ID

NO:32; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:33; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:34.

5 In one aspect, the invention provides an anti-TIM3 antibody comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:29; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:30; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:31; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:32; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:33; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:34.

10 In another aspect, an antibody of the invention comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:29, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:30, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:31; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:32; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:33 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:34.

20 In another aspect, an antibody of the invention comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:29, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:30, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:31; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:32; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:33 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:34.

In one embodiment such anti-TIM3 antibody comprises

- i) a VH sequence of SEQ ID NO:35 and a VL sequence of SEQ ID NO:36;
- ii) or humanized variant of the VH and VL of the antibody under i).

30 In one aspect, the invention provides an anti-TIM3 antibody comprising at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37; (b) HVR-H2 comprising the amino acid

sequence of SEQ ID NO:38; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:39; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42.

5 In one aspect, the invention provides an anti-TIM3 antibody comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:39; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (e) HVR-L2 comprising the amino acid
10 sequence of SEQ ID NO:41; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42.

In another aspect, an antibody of the invention comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37, (ii) HVR-H2
15 comprising the amino acid sequence of SEQ ID NO:38, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:39; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41 and (c) HVR-
20 L3 comprising the amino acid sequence of SEQ ID NO:42.

In another aspect, an antibody of the invention comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:39; and
25 (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42.

In one embodiment such anti-TIM3 antibody comprises

- 30 i) a VH sequence of SEQ ID NO:43 and a VL sequence of SEQ ID NO:44;
- ii) or humanized variant of the VH and VL of the antibody under i).

In one preferred embodiment such anti-TIM3 antibody comprises

- i) a VH sequence of SEQ ID NO:83 and a VL sequence of SEQ ID NO:84, or
- ii) a VH sequence of SEQ ID NO:85 and a VL sequence of SEQ ID NO:86.

One preferred embodiment is an antibody that binds to human TIM3 antibody wherein the antibody comprises a VH sequence of SEQ ID NO:83 and a VL sequence of SEQ ID NO:84.

One preferred embodiment is an antibody that binds to human TIM3 antibody wherein the antibody comprises a VH sequence of SEQ ID NO:85 and a VL sequence of SEQ ID NO:86.

In one aspect, the invention provides an anti-TIM3 antibody comprising at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:45; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:46; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:47; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:48; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:49; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:50.

In one aspect, the invention provides an anti-TIM3 antibody comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:45; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:46; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:47; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:48; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:49; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:50.

In another aspect, an antibody of the invention comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:45, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:46, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:47; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:48;

(ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:49 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:50.

In another aspect, an antibody of the invention comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:45, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:46, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:47; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:48; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:49 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:50.

In one embodiment such anti-TIM3 antibody comprises

- i) a VH sequence of SEQ ID NO:51 and a VL sequence of SEQ ID NO:52;
- ii) or humanized variant of the VH and VL of the antibody under i).

In one aspect, the invention provides an anti-TIM3 antibody comprising at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:53; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:54; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:55; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:56; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:57; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:58.

In one aspect, the invention provides an anti-TIM3 antibody comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:53; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:54; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:55; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:56; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:57; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:58.

In another aspect, an antibody of the invention comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:53, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:54, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:55; and (b) a VL

domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:56; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:57 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:58.

- 5 In another aspect, an antibody of the invention comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:53, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:54, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:55; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of
10 SEQ ID NO:56; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:57 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:58.

In one embodiment such anti-TIM3 antibody comprises

- i) a VH sequence of SEQ ID NO:59 and a VL sequence of SEQ ID NO:60;
- 15 ii) or humanized variant of the VH and VL of the antibody under i).

- In one aspect, the invention provides an anti-TIM3 antibody comprising at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:61; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:62; (c) HVR-H3 comprising the amino acid sequence of
20 SEQ ID NO:63; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:64; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:65; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:66.

- In one aspect, the invention provides an anti-TIM3 antibody comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:61; (b) HVR-H2
25 comprising the amino acid sequence of SEQ ID NO:62; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:63; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:64; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:65; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:66.

- 30 In another aspect, an antibody of the invention comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:61, (ii) HVR-H2

comprising the amino acid sequence of SEQ ID NO:62, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:63; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:64;
5 (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:65 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:66.

In another aspect, an antibody of the invention comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:61, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:62, and (iii)
10 HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:63; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:64; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:65 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:66.

In one embodiment such anti-TIM3 antibody comprises

- 15 i) a VH sequence of SEQ ID NO:67 and a VL sequence of SEQ ID NO:68;
- ii) or humanized variant of the VH and VL of the antibody under i).

In any of the above embodiments, an anti-TIM3 antibody is humanized. In one embodiment, an anti-TIM3 antibody comprises HVRs as in any of the above
20 embodiments, and further comprises an acceptor human framework, e.g. a human immunoglobulin framework or a human consensus framework. In another embodiment, an anti-TIM3 antibody comprises HVRs as in any of the above embodiments, and further comprises a VH and VL comprising such HVRs.

In a further aspect, the invention provides an antibody that binds to the same epitope as an anti-TIM3 antibody provided herein. For example, in certain
25 embodiments, an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:7 and a VL sequence of SEQ ID NO:8, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:9 and a VL sequence of SEQ
30 ID NO:10, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:19 and a VL sequence of SEQ ID NO:20, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:27 and a VL sequence of SEQ

5 ID NO:28, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:35 and a VL sequence of SEQ ID NO:36, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:43 and a VL sequence of SEQ ID NO:44, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:51 and a VL sequence of SEQ ID NO:52, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:59 and a VL sequence of SEQ ID NO:60, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:67 and a VL sequence of SEQ ID NO:68.

In one preferred embodiment an antibody is provided that binds to the same epitope as an anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:7 and a VL sequence of SEQ ID NO:8.

15 In one preferred embodiment an antibody is provided that competes for binding to human TIM3 with anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:7 and a VL sequence of SEQ ID NO:8 (as determined in a competition assay described in Example 4 on RPMI-8226 cells (ATCC® CCL-155™)).

20 In one aspect, the invention provides an anti-TIM3 antibody (e.g. an antibody that binds to human TIM3) comprising

25 A) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:3; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:4; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or

30 B) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:3; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or

- 5 C) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:3; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or
- 10 D) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:13; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:14; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:15; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:16; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:17; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:18; or
- 15 E) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:21; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:22; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:23; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:24; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:25; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:26; or
- 20 F) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:29; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:30; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:31; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:32; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:33; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:34; or
- 25 G) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:39; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42; or
- 30 H) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:45; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:46; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:47; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:48; (e) HVR-L2

comprising the amino acid sequence of SEQ ID NO:49; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:50; or.

5 I) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:53; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:54; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:55; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:56; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:57; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:58; or

10 J) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:61; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:62; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:63; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:64; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:65; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:66.

15 In another aspect the invention provides an anti-TIM3 antibody (e.g. an antibody that binds to human TIM3) comprising

20 A) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:4; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or

25 B) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or

30

C) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from

SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or

- 5 D) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:13, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:14, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:15; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:16; (ii) HVR-L2
10 comprising the amino acid sequence of SEQ ID NO:17 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:18; or
- E) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:21, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:22, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:23; and (b) a VL domain comprising (i) HVR-L1
15 comprising the amino acid sequence of SEQ ID NO:24; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:25 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:26; or.
- F) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:29, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:30, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:31; and (b) a VL domain comprising (i) HVR-L1
20 comprising the amino acid sequence of SEQ ID NO:32; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:33 and (iii) HVR-L3
25 comprising the amino acid sequence of SEQ ID NO:34; or
- G) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:39; and (b) a VL domain comprising (i) HVR-L1
30 comprising the amino acid sequence of SEQ ID NO:40; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42; or
- H) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:45, (ii) HVR-H2 comprising the amino acid sequence of SEQ

ID NO:46, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:47; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:48; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:49 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:50; or

I) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:53, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:54, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:55; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:56; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:57 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:58; or

J) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:61, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:62, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:63; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:64; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:65 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:66.

In one aspect, the invention provides an anti-TIM3 antibody (e.g. an antibody that binds to human TIM3) that

A1)

- i) comprises a VH sequence of SEQ ID NO:7 and a VL sequence of SEQ ID NO:8;
- ii) comprises a VH sequence of SEQ ID NO:9 and a VL sequence of SEQ ID NO:10;
- iii) or humanized variant of the VH and VL of the antibody under i) or ii);

or A2)

- i) comprises a VH sequence of SEQ ID NO:79 and a VL sequence of SEQ ID NO:80, or

- 35 -

- ii) comprises a VH sequence of SEQ ID NO:81 and a VL sequence of SEQ ID NO:82.

or B)

- 5 i) comprises a VH sequence of SEQ ID NO:19 and a VL sequence of SEQ ID NO:20;
- ii) or humanized variant of the VH and VL of the antibody under i);

or C)

- i) comprises a VH sequence of SEQ ID NO:27 and a VL sequence of SEQ ID NO:28;
- 10 ii) or humanized variant of the VH and VL of the antibody under i);

or D)

- i) comprises a VH sequence of SEQ ID NO:35 and a VL sequence of SEQ ID NO:36;
- ii) or humanized variant of the VH and VL of the antibody under i);

15 or E)

- i) comprises a VH sequence of SEQ ID NO:43 and a VL sequence of SEQ ID NO:44;
- ii) or humanized variant of the VH and VL of the antibody under i);

or F)

- 20 i) comprises a VH sequence of SEQ ID NO:51 and a VL sequence of SEQ ID NO:52;
- ii) or humanized variant of the VH and VL of the antibody under i);

or G1)

- 25 i) comprises a VH sequence of SEQ ID NO:59 and a VL sequence of SEQ ID NO:60;

- 36 -

ii) or humanized variant of the VH and VL of the antibody under i);

or G2)

i) comprises a VH sequence of SEQ ID NO:83 and a VL sequence of SEQ ID NO:84, or

5 ii) comprises a VH sequence of SEQ ID NO:85 and a VL sequence of SEQ ID NO:86.

or H)

i) comprises a VH sequence of SEQ ID NO:67 and a VL sequence of SEQ ID NO:68;

10 ii) or humanized variant of the VH and VL of the antibody under i).

In another aspect the invention provides an anti-TIM3 antibody (e.g. an antibody that binds to human TIM3) comprising

15 A) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:4; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or

20 B) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or

25 C) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the

30

amino acid sequence of SEQ ID NO:12; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or

- 5 D) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:13, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:14, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:15; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:16; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:17 and (iii) HVR-L3
10 comprising the amino acid sequence of SEQ ID NO:18; or
- E) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:21, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:22, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:23; and (b) a VL domain comprising (i) HVR-L1
15 comprising the amino acid sequence of SEQ ID NO:24; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:25 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:26; or.
- F) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:29, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:30, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:31; and (b) a VL domain comprising (i) HVR-L1
20 comprising the amino acid sequence of SEQ ID NO:32; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:33 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:34; or
- G) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:39; and (b) a VL domain comprising (i) HVR-L1
25 comprising the amino acid sequence of SEQ ID NO:40; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41 and (iii) HVR-L3
30 comprising the amino acid sequence of SEQ ID NO:42; or
- H) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:45, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:46, and (iii) HVR-H3 comprising an amino acid sequence selected

from SEQ ID NO:47; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:48; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:49 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:50; or

5 I) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:53, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:54, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:55; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:56; (ii) HVR-L2
10 comprising the amino acid sequence of SEQ ID NO:57 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:58; or

J) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:61, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:62, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:63; and (b) a VL domain comprising (i) HVR-L1
15 comprising the amino acid sequence of SEQ ID NO:64; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:65 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:66;

wherein the antibody is characterized independently by one or more of the
20 following properties: anti-TIM3 antibody

- induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 45% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in
25 a FACS assay on TIM3 expressing RPMI8226 cells) of at least 50% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 55% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in
30 a FACS assay on TIM3 expressing RPMI8226 cells) of at least 60% after 240 Minutes at 37 °C (see Example 6)

- In one embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 65% after 240 Minutes at 37 °C (see Example 6)

In another aspect the invention provides an anti-TIM3 antibody (e.g. an antibody that binds to human TIM3) comprising

5

A) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:4; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or

10

B) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or

15

C) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or

20

25

D) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:13, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:14, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:15; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:16; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:17 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:18; or

30

- 5 E) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:21, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:22, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:23; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:24; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:25 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:26; or.
- 10 F) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:29, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:30, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:31; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:32; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:33 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:34; or
- 15 G) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:39; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42; or
- 20 H) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:45, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:46, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:47; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:48; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:49 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:50; or
- 25 I) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:53, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:54, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:55; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:56; (ii) HVR-L2
- 30

comprising the amino acid sequence of SEQ ID NO:57 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:58; or

- J) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:61, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:62, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:63; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:64; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:65 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:66;

wherein the antibody is characterized independently by one or more of the following properties: the anti-TIM3 antibody

- induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 45% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 50% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 55% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 60% after 240 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 65% after 240 Minutes at 37 °C (see Example 6)
- competes for binding to TIM3 with an anti-Tim3 antibody comprising the VH of SEQ ID NO:7 and VL of SEQ ID NO:8.
- binds to a human and cynomolgus TIM3
- shows as immunoconjugate a cytotoxic activity on TIM3 expressing cells (in one embodiment the immunoconjugates has a relative IC50 value of the cytotoxic activity as Pseudomonas exotoxin A conjugate on RPMI-8226 cells of 0.1 or lower (as measured in Example 11)

- induces interferon-gamma release (in a Mixed Lymphocyte Reaction (MLR) assay as described in Example 5).

In a further aspect of the invention, an anti-TIM3 antibody according to any of the above embodiments is a monoclonal antibody, including a chimeric, humanized or human antibody. In one embodiment, an anti-TIM3 antibody is an antibody fragment, e.g., a Fv, Fab, Fab', scFv, diabody, or F(ab')₂ fragment. In another embodiment, the antibody is a full length antibody, e.g., an intact IgG1 or IgG4 antibody or other antibody class or isotype as defined herein.

In a further aspect, an anti-TIM3 antibody according to any of the above embodiments may incorporate any of the features, singly or in combination, as described in Sections 1-7 below:

1. Antibody Affinity

In certain embodiments, an antibody provided herein has a dissociation constant KD of $\leq 1 \mu\text{M}$, $\leq 100 \text{ nM}$, $\leq 10 \text{ nM}$, $\leq 1 \text{ nM}$, $\leq 0.1 \text{ nM}$, $\leq 0.01 \text{ nM}$, or $\leq 0.001 \text{ nM}$ (e.g. 10^{-8} M or less, e.g. from 10^{-8} M to 10^{-13} M , e.g., from 10^{-9} M to 10^{-13} M).

In one preferred embodiment, KD is measured using surface plasmon resonance assays using a BIACORE[®]) at 25°C with immobilized antigen CM5 chips at ~10 response units (RU). Briefly, carboxymethylated dextran biosensor chips (CM5, BIACORE, Inc.) are activated with *N*-ethyl-*N'*-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC) and *N*-hydroxysuccinimide (NHS) according to the supplier's instructions. Antigen is diluted with 10 mM sodium acetate, pH 4.8, to 5 µg/ml (~0.2 µM) before injection at a flow rate of 5 µl/minute to achieve approximately 10 response units (RU) of coupled protein. Following the injection of antigen, 1 M ethanolamine is injected to block unreacted groups. For kinetics measurements, two-fold serial dilutions of Fab (0.78 nM to 500 nM) are injected in PBS with 0.05% polysorbate 20 (TWEEN-20TM) surfactant (PBST) at 25°C at a flow rate of approximately 25 µl/min. Association rates (k_{on} or k_a) and dissociation rates (k_{off} or k_d) are calculated using a simple one-to-one Langmuir binding model (BIACORE[®] Evaluation Software version 3.2) by simultaneously fitting the association and dissociation sensorgrams. The equilibrium dissociation constant KD is calculated as the ratio k_d/k_a ($k_{\text{off}}/k_{\text{on}}$). See, e.g., Chen, Y. et al., J. Mol. Biol. 293 (1999) 865-881. If the on-rate exceeds $10^6 \text{ M}^{-1} \text{ s}^{-1}$ by the surface plasmon resonance assay above, then the on-rate can be determined by using a fluorescent quenching technique that measures the increase or decrease in

fluorescence emission intensity (excitation = 295 nm; emission = 340 nm, 16 nm band-pass) at 25°C of a 20 nM anti-antigen antibody (Fab form) in PBS, pH 7.2, in the presence of increasing concentrations of antigen as measured in a spectrometer, such as a stop-flow equipped spectrophotometer (Aviv Instruments) or a 8000-series SLM-AMINCOTM spectrophotometer (ThermoSpectronic) with a stirred
5 cuvette.

2. Antibody Fragments

In certain embodiments, an antibody provided herein is an antibody fragment. Antibody fragments include, but are not limited to, Fab, Fab', Fab'-SH, F(ab')₂, Fv, and scFv fragments, and other fragments described below. For a review of certain
10 antibody fragments, see Hudson, P.J. et al., Nat. Med. 9 (2003) 129-134. For a review of scFv fragments, see, e.g., Plueckthun, A., In; The Pharmacology of Monoclonal Antibodies, Vol. 113, Rosenberg and Moore (eds.), Springer-Verlag, New York (1994), pp. 269-315; see also WO 93/16185; and U.S. Patent Nos.
15 5,571,894 and 5,587,458. For discussion of Fab and F(ab')₂ fragments comprising salvage receptor binding epitope residues and having increased in vivo half-life, see U.S. Patent No. 5,869,046.

Diabodies are antibody fragments with two antigen-binding sites that may be bivalent or bispecific. See, for example, EP 0 404 097; WO 1993/01161; Hudson,
20 P.J. et al., Nat. Med. 9 (2003) 129-134; and Holliger, P. et al., Proc. Natl. Acad. Sci. USA 90 (1993) 6444-6448. Triabodies and tetrabodies are also described in Hudson, P.J. et al., Nat. Med. 9 (2003) 129-134).

Single-domain antibodies are antibody fragments comprising all or a portion of the heavy chain variable domain or all or a portion of the light chain variable domain
25 of an antibody. In certain embodiments, a single-domain antibody is a human single-domain antibody (Domantis, Inc., Waltham, MA; see, e.g., U.S. Patent No. 6,248,516 B1).

Antibody fragments can be made by various techniques, including but not limited to proteolytic digestion of an intact antibody as well as production by recombinant
30 host cells (e.g. *E. coli* or phage), as described herein.

3. Chimeric and Humanized Antibodies

In certain embodiments, an antibody provided herein is a chimeric antibody. Certain chimeric antibodies are described, e.g., in U.S. Patent No. 4,816,567; and Morrison, S.L. et al., Proc. Natl. Acad. Sci. USA 81 (1984) 6851-6855). In one
5 example, a chimeric antibody comprises a non-human variable region (e.g., a variable region derived from a mouse, rat, hamster, rabbit, or non-human primate, such as a monkey) and a human constant region. In a further example, a chimeric antibody is a "class switched" antibody in which the class or subclass has been changed from that of the parent antibody. Chimeric antibodies include antigen-
10 binding fragments thereof.

In certain embodiments, a chimeric antibody is a humanized antibody. Typically, a non-human antibody is humanized to reduce immunogenicity to humans, while retaining the specificity and affinity of the parental non-human antibody. Generally, a humanized antibody comprises one or more variable domains in which
15 HVRs, e.g., CDRs, (or portions thereof) are derived from a non-human antibody, and FRs (or portions thereof) are derived from human antibody sequences. A humanized antibody optionally will also comprise at least a portion of a human constant region. In some embodiments, some FR residues in a humanized antibody are substituted with corresponding residues from a non-human antibody (e.g., the
20 antibody from which the HVR residues are derived), e.g., to restore or improve antibody specificity or affinity.

Humanized antibodies and methods of making them are reviewed, e.g., in Almagro, J.C. and Fransson, J., Front. Biosci. 13 (2008) 1619-1633, and are further described, e.g., in Riechmann, I. et al., Nature 332 (1988) 323-329; Queen, C. et
25 al., Proc. Natl. Acad. Sci. USA 86 (1989) 10029-10033; US Patent Nos. 5, 821,337, 7,527,791, 6,982,321, and 7,087,409; Kashmiri, S.V. et al., Methods 36 (2005) 25-34 (describing SDR (a-CDR) grafting); Padlan, E.A., Mol. Immunol. 28 (1991) 489-498 (describing "resurfacing"); Dall'Acqua, W.F. et al., Methods 36 (2005) 43-60 (describing "FR shuffling"); and Osbourn, J. et al., Methods 36
30 (2005) 61-68 and Klimka, A. et al., Br. J. Cancer 83 (2000) 252-260 (describing the "guided selection" approach to FR shuffling).

Human framework regions that may be used for humanization include but are not limited to: framework regions selected using the "best-fit" method (see, e.g., Sims, M.J. et al., J. Immunol. 151 (1993) 2296-2308; framework regions derived from

the consensus sequence of human antibodies of a particular subgroup of light or heavy chain variable regions (see, e.g., Carter, P. et al., Proc. Natl. Acad. Sci. USA 89 (1992) 4285-4289; and Presta, L.G. et al., J. Immunol. 151 (1993) 2623-2632); human mature (somatically mutated) framework regions or human germline framework regions (see, e.g., Almagro, J.C. and Fransson, J., Front. Biosci. 13 (2008) 1619-1633); and framework regions derived from screening FR libraries (see, e.g., Baca, M. et al., J. Biol. Chem. 272 (1997) 10678-10684 and Rosok, M.J. et al., J. Biol. Chem. 271 (1996) 22611-22618).

4. Human Antibodies

In certain embodiments, an antibody provided herein is a human antibody. Human antibodies can be produced using various techniques known in the art. Human antibodies are described generally in van Dijk, M.A. and van de Winkel, J.G., Curr. Opin. Pharmacol. 5 (2001) 368-374 and Lonberg, N., Curr. Opin. Immunol. 20 (2008) 450-459.

Human antibodies may be prepared by administering an immunogen to a transgenic animal that has been modified to produce intact human antibodies or intact antibodies with human variable regions in response to antigenic challenge. Such animals typically contain all or a portion of the human immunoglobulin loci, which replace the endogenous immunoglobulin loci, or which are present extrachromosomally or integrated randomly into the animal's chromosomes. In such transgenic mice, the endogenous immunoglobulin loci have generally been inactivated. For review of methods for obtaining human antibodies from transgenic animals, see Lonberg, N., Nat. Biotech. 23 (2005) 1117-1125. See also, e.g., U.S. Patent Nos. 6,075,181 and 6,150,584 describing XENOMOUSETM technology; U.S. Patent No. 5,770,429 describing HUMAB[®] technology; U.S. Patent No. 7,041,870 describing K-M MOUSE[®] technology, and U.S. Patent Application Publication No. US 2007/0061900, describing VELOCIMOUSE[®] technology). Human variable regions from intact antibodies generated by such animals may be further modified, e.g., by combining with a different human constant region.

Human antibodies can also be made by hybridoma-based methods. Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal antibodies have been described. (See, e.g., Kozbor, D., J. Immunol. 133 (1984) 3001-3005; Brodeur, B.R. et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York (1987), pp. 51-63;

and Boerner, P. et al., J. Immunol. 147 (1991) 86-95) Human antibodies generated via human B-cell hybridoma technology are also described in Li, J. et al., Proc. Natl. Acad. Sci. USA 103 (2006) 3557-3562. Additional methods include those described, for example, in U.S. Patent No. 7,189,826 (describing production of
5 monoclonal human IgM antibodies from hybridoma cell lines) and Ni, J., Xiandai Mianyixue 26 (2006) 265-268 (describing human-human hybridomas). Human hybridoma technology (Trioma technology) is also described in Vollmers, H.P. and Brandlein, S., Histology and Histopathology 20 (2005) 927-937 and Vollmers, H.P. and Brandlein, S., Methods and Findings in Experimental and Clinical
10 Pharmacology 27 (2005) 185-191.

Human antibodies may also be generated by isolating Fv clone variable domain sequences selected from human-derived phage display libraries. Such variable domain sequences may then be combined with a desired human constant domain. Techniques for selecting human antibodies from antibody libraries are described
15 below.

5. Library-Derived Antibodies

Antibodies of the invention may be isolated by screening combinatorial libraries for antibodies with the desired activity or activities. For example, a variety of methods are known in the art for generating phage display libraries and screening
20 such libraries for antibodies possessing the desired binding characteristics. Such methods are reviewed, e.g., in Hoogenboom, H.R. et al., Methods in Molecular Biology 178 (2001) 1-37 and further described, e.g., in the McCafferty, J. et al., Nature 348 (1990) 552-554; Clackson, T. et al., Nature 352 (1991) 624-628; Marks, J.D. et al., J. Mol. Biol. 222 (1992) 581-597; Marks, J.D. and Bradbury, A.,
25 Methods in Molecular Biology 248 (2003) 161-175; Sidhu, S.S. et al., J. Mol. Biol. 338 (2004) 299-310; Lee, C.V. et al., J. Mol. Biol. 340 (2004) 1073-1093; Fellouse, F.A., Proc. Natl. Acad. Sci. USA 101 (2004) 12467-12472; and Lee, C.V. et al., J. Immunol. Methods 284 (2004) 119-132.

In certain phage display methods, repertoires of VH and VL genes are separately
30 cloned by polymerase chain reaction (PCR) and recombined randomly in phage libraries, which can then be screened for antigen-binding phage as described in Winter, G. et al., Ann. Rev. Immunol. 12 (1994) 433-455. Phage typically display antibody fragments, either as single-chain Fv (scFv) fragments or as Fab fragments. Libraries from immunized sources provide high-affinity antibodies to the

immunogen without the requirement of constructing hybridomas. Alternatively, the naive repertoire can be cloned (e.g., from human) to provide a single source of antibodies to a wide range of non-self and also self-antigens without any immunization as described by Griffiths, A.D. et al., EMBO J. 12 (1993) 725-734.

5 Finally, naive libraries can also be made synthetically by cloning non-rearranged V-gene segments from stem cells, and using PCR primers containing random sequence to encode the highly variable CDR3 regions and to accomplish rearrangement *in vitro*, as described by Hoogenboom, H.R. and Winter, G., J. Mol. Biol. 227 (1992) 381-388. Patent publications describing human antibody phage

10 libraries include, for example: US Patent No. 5,750,373, and US Patent Publication Nos. 2005/0079574, 2005/0119455, 2005/0266000, 2007/0117126, 2007/0160598, 2007/0237764, 2007/0292936, and 2009/0002360.

Antibodies or antibody fragments isolated from human antibody libraries are considered human antibodies or human antibody fragments herein.

15 **6. Multispecific Antibodies**

In certain embodiments, an antibody provided herein is a multispecific antibody, e.g. a bispecific antibody. Multispecific antibodies are monoclonal antibodies that have binding specificities for at least two different sites. In certain embodiments, one of the binding specificities is for TIM3 and the other is for any other antigen.

20 In certain embodiments, bispecific antibodies may bind to two different epitopes of TIM3. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express TIM3. Bispecific antibodies can be prepared as full length antibodies or antibody fragments.

Techniques for making multispecific antibodies include, but are not limited to, recombinant co-expression of two immunoglobulin heavy chain-light chain pairs having different specificities (see Milstein, C. and Cuello, A.C., Nature 305 (1983) 537-540, WO 93/08829, and Traunecker, A. et al., EMBO J. 10 (1991) 3655-3659), and "knob-in-hole" engineering (see, e.g., U.S. Patent No. 5,731,168).

25 Multi-specific antibodies may also be made by engineering electrostatic steering effects for making antibody Fc-heterodimeric molecules (WO 2009/089004); cross-linking two or more antibodies or fragments (see, e.g., US Patent No. 4,676,980, and Brennan, M. et al., Science 229 (1985) 81-83); using leucine zippers to produce bi-specific antibodies (see, e.g., Kostelny, S.A. et al., J. Immunol. 148 (1992) 1547-1553); using "diabody" technology for making bispecific antibody

30

fragments (see, e.g., Holliger, P. et al., Proc. Natl. Acad. Sci. USA 90 (1993) 6444-6448); and using single-chain Fv (scFv) dimers (see, e.g. Gruber, M et al., J. Immunol. 152 (1994) 5368-5374); and preparing trispecific antibodies as described, e.g., in Tutt, A. et al., J. Immunol. 147 (1991) 60-69).

5 Engineered antibodies with three or more functional antigen binding sites, including "Octopus antibodies," are also included herein (see, e.g. US 2006/0025576).

The antibody or fragment herein also includes a "Dual Acting Fab" or "DAF" comprising an antigen binding site that binds to TIM3 as well as another, different
10 antigen (see, US 2008/0069820, for example).

The antibody or fragment herein also includes multispecific antibodies described in WO 2009/080251, WO 2009/080252, WO 2009/080253, WO 2009/080254, WO 2010/112193, WO 2010/115589, WO 2010/136172, WO 2010/145792, and WO 2010/145793, WO2011/117330, WO2012/025525, WO2012/025530,
15 WO2013/026835, WO2013/026831, WO2013/164325, or WO 2013/174873.

7. Antibody Variants

In certain embodiments, amino acid sequence variants of the antibodies provided herein are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. Amino acid sequence
20 variants of an antibody may be prepared by introducing appropriate modifications into the nucleotide sequence encoding the antibody, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of residues within the amino acid sequences of the antibody. Any combination of deletion, insertion, and substitution can be made to arrive at the
25 final construct, provided that the final construct possesses the desired characteristics, e.g., antigen-binding.

a) Substitution, Insertion, and Deletion Variants

In certain embodiments, antibody variants having one or more amino acid substitutions are provided. Sites of interest for substitutional mutagenesis include
30 the HVRs and FRs. Exemplary changes are provided in Table 1 under the heading of "exemplary substitutions", and as further described below in reference to amino acid side chain classes. Conservative substitutions are shown in Table 1 under the

heading of "preferred substitutions". Amino acid substitutions may be introduced into an antibody of interest and the products screened for a desired activity, e.g., retained/improved antigen binding, decreased immunogenicity, or improved ADCC or CDC.

5

TABLE 1

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	Val; Leu; Ile	Val
Arg (R)	Lys; Gln; Asn	Lys
Asn (N)	Gln; His; Asp, Lys; Arg	Gln
Asp (D)	Glu; Asn	Glu
Cys (C)	Ser; Ala	Ser
Gln (Q)	Asn; Glu	Asn
Glu (E)	Asp; Gln	Asp
Gly (G)	Ala	Ala
His (H)	Asn; Gln; Lys; Arg	Arg
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Leu
Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe	Ile
Lys (K)	Arg; Gln; Asn	Arg
Met (M)	Leu; Phe; Ile	Leu
Phe (F)	Trp; Leu; Val; Ile; Ala; Tyr	Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Val; Ser	Ser
Trp (W)	Tyr; Phe	Tyr
Tyr (Y)	Trp; Phe; Thr; Ser	Phe
Val (V)	Ile; Leu; Met; Phe; Ala; Norleucine	Leu

Amino acids may be grouped according to common side-chain properties:

(1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile;

(2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;

- 50 -

(3) acidic: Asp, Glu;

(4) basic: His, Lys, Arg;

(5) residues that influence chain orientation: Gly, Pro;

(6) aromatic: Trp, Tyr, Phe.

5 Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

One type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody (e.g. a humanized or human antibody). Generally, the resulting variant(s) selected for further study will have modifications (e.g., improvements) in certain biological properties (e.g., increased affinity, reduced immunogenicity) relative to the parent antibody and/or will have substantially retained certain biological properties of the parent antibody. An exemplary substitutional variant is an affinity matured antibody, which may be conveniently generated, e.g., using phage display-based affinity maturation techniques such as those described herein. Briefly, one or more HVR residues are mutated and the variant antibodies displayed on phage and screened for a particular biological activity (e.g. binding affinity).

Alterations (e.g., substitutions) may be made in HVRs, e.g., to improve antibody affinity. Such alterations may be made in HVR “hotspots,” i.e., residues encoded by codons that undergo mutation at high frequency during the somatic maturation process (see, e.g., Chowdhury, P.S., Methods Mol. Biol. 207 (2008) 179-196), and/or SDRs (a-CDRs), with the resulting variant VH or VL being tested for binding affinity. Affinity maturation by constructing and reselecting from secondary libraries has been described, e.g., in Hoogenboom, H.R. et al. in Methods in Molecular Biology 178 (2002) 1-37. In some embodiments of affinity maturation, diversity is introduced into the variable genes chosen for maturation by any of a variety of methods (e.g., error-prone PCR, chain shuffling, or oligonucleotide-directed mutagenesis). A secondary library is then created. The library is then screened to identify any antibody variants with the desired affinity. Another method to introduce diversity involves HVR-directed approaches, in which several HVR residues (e.g., 4-6 residues at a time) are randomized. HVR residues involved in antigen binding may be specifically identified, e.g., using

alanine scanning mutagenesis or modeling. CDR-H3 and CDR-L3 in particular are often targeted.

5 In certain embodiments, substitutions, insertions, or deletions may occur within one or more HVRs so long as such alterations do not substantially reduce the ability of the antibody to bind antigen. For example, conservative alterations (e.g., conservative substitutions as provided herein) that do not substantially reduce binding affinity may be made in HVRs. Such alterations may be outside of HVR “hotspots” or SDRs. In certain embodiments of the variant VH and VL sequences provided above, each HVR either is unaltered, or contains no more than one, two or
10 three amino acid substitutions.

A useful method for identification of residues or regions of an antibody that may be targeted for mutagenesis is called "alanine scanning mutagenesis" as described by Cunningham, B.C. and Wells, J.A., Science 244 (1989) 1081-1085. In this method, a residue or group of target residues (e.g., charged residues such as arg, asp, his,
15 lys, and glu) are identified and replaced by a neutral or negatively charged amino acid (e.g., alanine or polyalanine) to determine whether the interaction of the antibody with antigen is affected. Further substitutions may be introduced at the amino acid locations demonstrating functional sensitivity to the initial substitutions. Alternatively, or additionally, a crystal structure of an antigen-antibody complex to identify contact points between the antibody and antigen.
20 Such contact residues and neighboring residues may be targeted or eliminated as candidates for substitution. Variants may be screened to determine whether they contain the desired properties.

Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions
25 ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue. Other insertional variants of the antibody molecule include the fusion to the N- or C-terminus of the antibody to an enzyme (e.g. for ADEPT) or a
30 polypeptide which increases the serum half-life of the antibody.

b) Fc region variants

In certain embodiments, one or more amino acid modifications may be introduced into the Fc region of an antibody provided herein, thereby generating an Fc region variant. The Fc region variant may comprise a human Fc region sequence (e.g., a

human IgG1, IgG2, IgG3 or IgG4 Fc region) comprising an amino acid modification (*e.g.* a substitution) at one or more amino acid positions.

Antibodies with reduced effector function include those with substitution of one or more of Fc region residues 238, 265, 269, 270, 297, 327 and 329 (U.S. Patent No. 6,737,056). Such Fc mutants include Fc mutants with substitutions at two or more of amino acid positions 265, 269, 270, 297 and 327, including the so-called “DANA” Fc mutant with substitution of residues 265 and 297 to alanine (US Patent No. 7,332,581).

Certain antibody variants with improved or diminished binding to FcRs are described. (See, *e.g.*, U.S. Patent No. 6,737,056; WO 2004/056312, and Shields, R.L. et al., *J. Biol. Chem.* 276 (2001) 6591-6604)

In one embodiment the invention such antibody is a IgG1 with mutations L234A and L235A or with mutations L234A, L235A and P329G. In another embodiment or IgG4 with mutations S228P and L235E or S228P, L235E or and P329G (numbering according to EU index of Kabat et al , Kabat et al., *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD, 1991).

Antibodies with increased half lives and improved binding to the neonatal Fc receptor (FcRn), which is responsible for the transfer of maternal IgGs to the fetus (Guyer, R.L. et al., *J. Immunol.* 117 (1976) 587-593, and Kim, J.K. et al., *J. Immunol.* 24 (1994) 2429-2434), are described in US 2005/0014934. Those antibodies comprise an Fc region with one or more substitutions therein which improve binding of the Fc region to FcRn. Such Fc variants include those with substitutions at one or more of Fc region residues: 238, 256, 265, 272, 286, 303, 305, 307, 311, 312, 317, 340, 356, 360, 362, 376, 378, 380, 382, 413, 424 or 434, *e.g.*, substitution of Fc region residue 434 (US Patent No. 7,371,826).

See also Duncan, A.R. and Winter, G., *Nature* 322 (1988) 738-740; US 5,648,260; US 5,624,821; and WO 94/29351 concerning other examples of Fc region variants.

c) Cysteine engineered antibody variants

In certain embodiments, it may be desirable to create cysteine engineered antibodies, *e.g.*, “thioMAbs,” in which one or more residues of an antibody are substituted with cysteine residues. In particular embodiments, the substituted

residues occur at accessible sites of the antibody. By substituting those residues with cysteine, reactive thiol groups are thereby positioned at accessible sites of the antibody and may be used to conjugate the antibody to other moieties, such as drug moieties or linker-drug moieties, to create an immunoconjugate, as described further herein. In certain embodiments, any one or more of the following residues may be substituted with cysteine: V205 (Kabat numbering) of the light chain; A118 (EU numbering) of the heavy chain; and S400 (EU numbering) of the heavy chain Fc region. Cysteine engineered antibodies may be generated as described, e.g., in U.S. Patent No. 7,521,541.

d) Antibody Derivatives

In certain embodiments, an antibody provided herein may be further modified to contain additional non-proteinaceous moieties that are known in the art and readily available. The moieties suitable for derivatization of the antibody include but are not limited to water soluble polymers. Non-limiting examples of water soluble polymers include, but are not limited to, polyethylene glycol (PEG), copolymers of ethylene glycol/propylene glycol, carboxymethylcellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone, poly-1, 3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids (either homopolymers or random copolymers), and dextran or poly(n-vinyl pyrrolidone)polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide copolymers, polyoxyethylated polyols (e.g., glycerol), polyvinyl alcohol, and mixtures thereof. Polyethylene glycol propionaldehyde may have advantages in manufacturing due to its stability in water. The polymer may be of any molecular weight, and may be branched or unbranched. The number of polymers attached to the antibody may vary, and if more than one polymer is attached, they can be the same or different molecules. In general, the number and/or type of polymers used for derivatization can be determined based on considerations including, but not limited to, the particular properties or functions of the antibody to be improved, whether the antibody derivative will be used in a therapy under defined conditions, etc.

In another embodiment, conjugates of an antibody and non-proteinaceous moiety that may be selectively heated by exposure to radiation are provided. In one embodiment, the non-proteinaceous moiety is a carbon nanotube (Kam, N.W. et al., Proc. Natl. Acad. Sci. USA 102 (2005) 11600-11605). The radiation may be of any wavelength, and includes, but is not limited to, wavelengths that do not harm

ordinary cells, but which heat the non-proteinaceous moiety to a temperature at which cells proximal to the antibody-non-proteinaceous moiety are killed.

B. Recombinant Methods and Compositions

Antibodies may be produced using recombinant methods and compositions, e.g., as described in U.S. Patent No. 4,816,567. In one embodiment, isolated nucleic acid encoding an anti-TIM3antibody described herein is provided. Such nucleic acid may encode an amino acid sequence comprising the VL and/or an amino acid sequence comprising the VH of the antibody (e.g., the light and/or heavy chains of the antibody). In a further embodiment, one or more vectors (e.g., expression vectors) comprising such nucleic acid are provided. In a further embodiment, a host cell comprising such nucleic acid is provided. In one such embodiment, a host cell comprises (e.g., has been transformed with): (1) a vector comprising a nucleic acid that encodes an amino acid sequence comprising the VL of the antibody and an amino acid sequence comprising the VH of the antibody, or (2) a first vector comprising a nucleic acid that encodes an amino acid sequence comprising the VL of the antibody and a second vector comprising a nucleic acid that encodes an amino acid sequence comprising the VH of the antibody. In one embodiment, the host cell is eukaryotic, e.g. a Chinese Hamster Ovary (CHO) cell or lymphoid cell (e.g., Y0, NS0, Sp20 cell). In one embodiment, a method of making an anti-TIM3antibody is provided, wherein the method comprises culturing a host cell comprising a nucleic acid encoding the antibody, as provided above, under conditions suitable for expression of the antibody, and optionally recovering the antibody from the host cell (or host cell culture medium).

For recombinant production of an anti-TIM3 antibody, nucleic acid encoding an antibody, e.g., as described above, is isolated and inserted into one or more vectors for further cloning and/or expression in a host cell. Such nucleic acid may be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the antibody).

Suitable host cells for cloning or expression of antibody-encoding vectors include prokaryotic or eukaryotic cells described herein. For example, antibodies may be produced in bacteria, in particular when glycosylation and Fc effector function are not needed. For expression of antibody fragments and polypeptides in bacteria, see, e.g., US 5,648,237, US 5,789,199, and US 5,840,523. (See also Charlton, K.A., In:

Methods in Molecular Biology, Vol. 248, Lo, B.K.C. (ed.), Humana Press, Totowa, NJ (2003), pp. 245-254, describing expression of antibody fragments in *E. coli*.) After expression, the antibody may be isolated from the bacterial cell paste in a soluble fraction and can be further purified.

5 In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for antibody-encoding vectors, including fungi and yeast strains whose glycosylation pathways have been "humanized," resulting in the production of an antibody with a partially or fully human glycosylation pattern. See Gerngross, T.U., Nat. Biotech. 22 (2004) 1409-1414;
10 and Li, H. et al., Nat. Biotech. 24 (2006) 210-215.

Suitable host cells for the expression of glycosylated antibody are also derived from multicellular organisms (invertebrates and vertebrates). Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains have been identified which may be used in conjunction with insect cells, particularly for
15 transfection of *Spodoptera frugiperda* cells.

Plant cell cultures can also be utilized as hosts. See, e.g., US Patent Nos. 5,959,177, 6,040,498, 6,420,548, 7,125,978, and 6,417,429 (describing PLANTIBODIESTM technology for producing antibodies in transgenic plants).

Vertebrate cells may also be used as hosts. For example, mammalian cell lines that
20 are adapted to grow in suspension may be useful. Other examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7); human embryonic kidney line (293 or 293 cells as described, e.g., in Graham, F.L. et al., J. Gen Virol. 36 (1977) 59-74); baby hamster kidney cells (BHK); mouse sertoli cells (TM4 cells as described, e.g., in Mather, J.P., Biol.
25 Reprod. 23 (1980) 243-252); monkey kidney cells (CV1); African green monkey kidney cells (VERO-76); human cervical carcinoma cells (HELA); canine kidney cells (MDCK; buffalo rat liver cells (BRL 3A); human lung cells (W138); human liver cells (Hep G2); mouse mammary tumor (MMT 060562); TRI cells, as described, e.g., in Mather, J.P. et al., Annals N.Y. Acad. Sci. 383 (1982) 44-68;
30 MRC 5 cells; and FS4 cells. Other useful mammalian host cell lines include Chinese hamster ovary (CHO) cells, including DHFR⁻ CHO cells (Urlaub, G. et al., Proc. Natl. Acad. Sci. USA 77 (1980) 4216-4220); and myeloma cell lines such as Y0, NS0 and Sp2/0. For a review of certain mammalian host cell lines suitable for antibody production, see, e.g., Yazaki, P. and Wu, A.M., Methods in Molecular

Biology, Vol. 248, Lo, B.K.C. (ed.), Humana Press, Totowa, NJ (2004), pp. 255-268.

C. Assays

Anti-TIM3 antibodies provided herein may be identified, screened for, or characterized for their physical/chemical properties and/or biological activities by various assays known in the art.

1. Binding assays and other assays

In one aspect, an antibody of the invention is tested for its antigen binding activity, e.g., by known methods such as ELISA, Western blot, etc.

In another aspect, competition assays may be used to identify an antibody that competes with Tim3_0016 (comprising a VH sequence of SEQ ID NO:7 and a VL sequence of SEQ ID NO:8) for binding to TIM3 (or alternatively with the Tim3_0016 variant antibody Tim3_0018 with the identical 6 HVRs). Thus one embodiment of the invention is antibody which competes for binding to human TIM3 with an anti-TIM3 antibody comprising all 3 HVRs of VH sequence of SEQ ID NO:7 and all 3 HVRs of VL sequence of SEQ ID NO:8). In certain embodiments, such a competing antibody binds to the same epitope (e.g., a linear or a conformational epitope) that is bound by anti-TIM3 antibody Tim3_0016. Detailed exemplary methods for mapping an epitope to which an antibody binds are provided in Morris, G.E. (ed.), Epitope Mapping Protocols, In: Methods in Molecular Biology, Vol. 66, Humana Press, Totowa, NJ (1996).

In an exemplary competition assay, immobilized TIM3(-ECD) is incubated in a solution comprising a first labeled antibody that binds to TIM3 (e.g., anti-TIM3 antibody aTim3_0016) and a second unlabeled antibody that is being tested for its ability to compete with the first antibody for binding to TIM3. The second antibody may be present in a hybridoma supernatant. As a control, immobilized TIM3 is incubated in a solution comprising the first labeled antibody but not the second unlabeled antibody. After incubation under conditions permissive for binding of the first antibody to TIM3, excess unbound antibody is removed, and the amount of label associated with immobilized TIM3 is measured. If the amount of label associated with immobilized TIM3 is substantially reduced in the test sample relative to the control sample, then that indicates that the second antibody is competing with the first antibody for binding to TIM3. See Harlow, E. and Lane,

D., *Antibodies: A Laboratory Manual*, Chapter 14, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1988). For another exemplary competition assay see Example 4.

2. Activity assays

5 In one aspect, assays are provided for identifying anti-TIM3 antibodies thereof having biological activity. Biological activity may include, e.g., TIM3 receptor internalization, or cytokine release, or cytotoxic activity (as immunoconjugates conjugated to a toxin), or cynomolgus binding crossreactivity, as well as binding to
10 different cell types. Antibodies having such biological activity in vivo and/or in vitro are also provided.

In certain embodiments, an antibody of the invention is tested for such biological activity as described e.g. in Examples 5 to 15.

D. Immunoconjugates (Cancer only or modify for target)

15 The invention also provides immunoconjugates comprising an anti-TIM3 antibody herein conjugated to one or more cytotoxic agents, such as chemotherapeutic agents or drugs, growth inhibitory agents, toxins (e.g., protein toxins, enzymatically active toxins of bacterial, fungal, plant, or animal origin, or fragments thereof), or radioactive isotopes.

20 In one embodiment, an immunoconjugate is an antibody-drug conjugate (ADC) in which an antibody is conjugated to one or more drugs, including but not limited to a maytansinoid (see US 5,208,020, US 5,416,064 and EP 0 425 235 B1); an auristatin such as monomethyl auristatin drug moieties DE and DF (MMAE and MMAF) (see US 5,635,483, US 5,780,588, and US 7,498,298); a dolastatin; a calicheamicin or derivative thereof (see US 5,712,374, US 5,714,586,
25 US 5,739,116, US 5,767,285, US 5,770,701, US 5,770,710, US 5,773,001, and US 5,877,296; Hinman, L.M. et al., *Cancer Res.* 53 (1993) 3336-3342; and Lode, H.N. et al., *Cancer Res.* 58 (1998) 2925-2928); an anthracycline such as daunomycin or doxorubicin (see Kratz, F. et al., *Curr. Med. Chem.* 13 (2006) 477-523; Jeffrey, S.C. et al., *Bioorg. Med. Chem. Lett.* 16 (2006) 358-362; Torgov,
30 M.Y. et al., *Bioconjug. Chem.* 16 (2005) 717-721; Nagy, A. et al., *Proc. Natl. Acad. Sci. USA* 97 (2000) 829-834; Dubowchik, G.M. et al., *Bioorg. & Med. Chem. Letters* 12 (2002) 1529-1532; King, H.D. et al., *J. Med. Chem.* 45 (2002) 4336-4343; and U.S. Patent No. 6,630,579); methotrexate; vindesine; a taxane such

as docetaxel, paclitaxel, larotaxel, tesetaxel, and ortataxel; a trichothecene; and CC1065.

5 In another embodiment, an immunoconjugate comprises an antibody as described herein conjugated to an enzymatically active toxin or fragment thereof, including but not limited to diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, *saponaria officinalis* inhibitor, gelonin, mitogellin, 10 restrictocin, phenomycin, enomycin, and the tricothecenes.

In another embodiment, an immunoconjugate comprises an antibody as described herein conjugated to a *Pseudomonas* exotoxin A (or variants) thereof. *Pseudomonas* exotoxin A (PE) is a bacterial toxin with cytotoxic activity that may be effective for destroying or inhibiting the growth of undesirable cells, e.g., cancer 15 cells. Accordingly, PE may be useful for treating or preventing diseases such as, e.g., cancer. Several deimmunized *Pseudomonas* exotoxins (PE) are known in art. The domain II deleted versions (for example, PE24) may be less immunogenic and may cause fewer side effects (such as, for example, capillary leak syndrome and hepatotoxicity) as compared to PE38, which contains domain II. Different furin 20 cleavable linkers may be employed in PE24 variants. Such deimmunized *Pseudomonas* exotoxins (PE) are described in, for example, International Patent Application Publications WO2005052006, WO2007016150, WO2007014743, WO2007031741, WO200932954, WO201132022, WO2012/154530, and WO 2012/170617. The term “a *Pseudomonas* exotoxin A” as used herein encompasses 25 wildtype and deimmunized *Pseudomonas* exotoxins (PE). In one preferred embodiment “a *Pseudomonas* exotoxin A” refers to a deimmunized PE24 variant as e.g. described in but not limited to WO2009/32954, WO2011/32022, WO2012/154530, WO 2012/170617, Liu W, et al, PNAS 109 (2012) 11782-11787, Mazor R, et al PNAS 111 (2014) 8571–8576 and Alewine C, et al, Mol 30 Cancer Ther. (2014) 2653-61. In one preferred embodiment the “a *Pseudomonas* exotoxin A” comprises the amino acid sequences of SEQ ID NO:69 or comprises the amino acid sequences of SEQ ID NO:70 (their preparation is also described in Mazor R, et al PNAS 111 (2014) 8571–8576 and Alewine C, et al, Mol Cancer Ther. (2014) 2653-61).

In another embodiment, an immunoconjugate comprises an antibody as described herein conjugated to a Amatoxin or variants thereof. Amatoxins (α -amanitin, β -amanitin, amanin) are cyclic peptides composed of 8 amino acids. They can be isolated from *Amanita phalloides* mushrooms or prepared from the building blocks by synthesis. Amatoxins inhibit specifically the DNA-dependent RNA polymerase II of mammalian cells, and by this transcription and protein biosynthesis of the cells affected. Inhibition of transcription in a cell causes stop of growth and proliferation. Though not covalently bound, the complex between amanitin and RNA-polymerase II is very tight ($KD = 3nM$). Dissociation of amanitin from the enzyme is a very slow process what makes recovery of an affected cell unlikely. When in a cell the inhibition of transcription will last too long, the cell undergoes programmed cell death (apoptosis), as shown in cultures of Jurkat cells incubated with α -amanitin, or, with much higher efficiency, in Jurkat cells incubated with the membrane-permeable O-methyl- α -amanitin oleate. In one preferred embodiment term "Amatoxin" as used herein refers to an alpha-amanitin or variant thereof as described e.g. in WO2010/115630, WO2010/115629, WO2012/119787, WO2012/041504, and WO2014135282 with preferred variants described in WO2012/041504(e.g. conjugated via the 6' C-atom of amatoxin amino acid 4, particularly via an oxygen atom bound to the 6' C-atom of amatoxin amino acid, and wherein the TIM3 antibody is connected by a linker via a urea moiety) and WO2014135282.

In another embodiment, an immunoconjugate comprises an antibody as described herein conjugated to a radioactive atom to form a radioconjugate. A variety of radioactive isotopes are available for the production of radioconjugates. Examples include At^{211} , I^{131} , I^{125} , Y^{90} , Re^{186} , Re^{188} , Sm^{153} , Bi^{212} , P^{32} , Pb^{212} and radioactive isotopes of Lu. When the radioconjugate is used for detection, it may comprise a radioactive atom for scintigraphic studies, for example Tc^{99m} or I^{123} , or a spin label for nuclear magnetic resonance (NMR) imaging (also known as magnetic resonance imaging, MRI), such as iodine-123 again, iodine-131, indium-111, fluorine-19, carbon-13, nitrogen-15, oxygen-17, gadolinium, manganese or iron.

Conjugates of an antibody and cytotoxic agent may be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP), succinimidyl-4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-

azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta, E.S. et al., Science 238 (1987) 1098-1104. Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triamine pentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO 94/11026. The linker may be a "cleavable linker" facilitating release of a cytotoxic drug in the cell. For example, an acid-labile linker, peptidase-sensitive linker, photolabile linker, dimethyl linker or disulfide-containing linker (Chari, R.V. et al., Cancer Res. 52 (1992) 127-131; U.S. Patent No. 5,208,020) may be used.

The immunoconjugates or ADCs herein expressly contemplate, but are not limited to such conjugates prepared with cross-linker reagents including, but not limited to, BMPS, EMCS, GMBS, HBVS, LC-SMCC, MBS, MPBH, SBAP, SIA, SIAB, SMCC, SMPB, SMPH, sulfo-EMCS, sulfo-GMBS, sulfo-KMUS, sulfo-MBS, sulfo-SIAB, sulfo-SMCC, and sulfo-SMPB, and SVSB (succinimidyl-(4-vinylsulfone)benzoate) which are commercially available (e.g., from Pierce Biotechnology, Inc., Rockford, IL., U.S.A).

E. Methods and Compositions for Diagnostics and Detection

In certain embodiments, any of the anti-TIM3 antibodies provided herein is useful for detecting the presence of TIM3 in a biological sample. The term "detecting" as used herein encompasses quantitative or qualitative detection. In certain embodiments, a biological sample comprises a cell or tissue, such as AML stem cancer cells.

In one embodiment, an anti-TIM3 antibody for use in a method of diagnosis or detection is provided. In a further aspect, a method of detecting the presence of TIM3 in a biological sample is provided. In certain embodiments, the method comprises contacting the biological sample with an anti-TIM3 antibody as described herein under conditions permissive for binding of the anti-TIM3 antibody to TIM3, and detecting whether a complex is formed between the anti-TIM3 antibody and TIM3. Such method may be an *in vitro* or *in vivo* method. In one embodiment, an anti-TIM3 antibody is used to select subjects eligible for

therapy with an anti-TIM3 antibody, e.g. where TIM3 is a biomarker for selection of patients.

Exemplary disorders that may be diagnosed using an antibody of the invention include cancer including different form of hematological cancers like AML or multiple myelomas (MM).

In certain embodiments, labeled anti-TIM3 antibodies are provided. Labels include, but are not limited to, labels or moieties that are detected directly (such as fluorescent, chromophoric, electron-dense, chemiluminescent, and radioactive labels), as well as moieties, such as enzymes or ligands, that are detected indirectly, e.g., through an enzymatic reaction or molecular interaction. Exemplary labels include, but are not limited to, the radioisotopes ^{32}P , ^{14}C , ^{125}I , ^3H , and ^{131}I , fluorophores such as rare earth chelates or fluorescein and its derivatives, rhodamine and its derivatives, dansyl, umbelliferone, luciferases, e.g., firefly luciferase and bacterial luciferase (U.S. Patent No. 4,737,456), luciferin, 2,3-dihydrophthalazinediones, horseradish peroxidase (HRP), alkaline phosphatase, β -galactosidase, glucoamylase, lysozyme, saccharide oxidases, e.g., glucose oxidase, galactose oxidase, and glucose-6-phosphate dehydrogenase, heterocyclic oxidases such as uricase and xanthine oxidase, coupled with an enzyme that employs hydrogen peroxide to oxidize a dye precursor such as HRP, lactoperoxidase, or microperoxidase, biotin/avidin, spin labels, bacteriophage labels, stable free radicals, and the like.

F. Pharmaceutical Formulations

Pharmaceutical formulations of an anti-TIM3 antibody as described herein are prepared by mixing such antibody having the desired degree of purity with one or more optional pharmaceutically acceptable carriers (Remington's Pharmaceutical Sciences, 16th edition, Osol, A. (ed.) (1980)), in the form of lyophilized formulations or aqueous solutions. Pharmaceutically acceptable carriers are generally nontoxic to recipients at the dosages and concentrations employed, and include, but are not limited to: buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyl dimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride; benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10

residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as poly(vinylpyrrolidone); amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.* Zn-protein complexes); and/or non-ionic surfactants such as polyethylene glycol (PEG). Exemplary pharmaceutically acceptable carriers herein further include interstitial drug dispersion agents such as soluble neutral-active hyaluronidase glycoproteins (sHASEGP), for example, human soluble PH-20 hyaluronidase glycoproteins, such as rhuPH20 (HYLENEX[®], Baxter International, Inc.). Certain exemplary sHASEGPs and methods of use, including rhuPH20, are described in US Patent Publication Nos. 2005/0260186 and 2006/0104968. In one aspect, a sHASEGP is combined with one or more additional glycosaminoglycanases such as chondroitinases.

Exemplary lyophilized antibody formulations are described in US Patent No. 6,267,958. Aqueous antibody formulations include those described in US Patent No. 6,171,586 and WO 2006/044908, the latter formulations including a histidine-acetate buffer.

The formulation herein may also contain more than one active ingredients as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, it may be desirable to further provide [[list drugs that might be combined with the anti-TIM3 antibody]]. Such active ingredients are suitably present in combination in amounts that are effective for the purpose intended.

Active ingredients may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methyl methacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, 16th edition, Osol, A. (ed.) (1980).

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semi-permeable matrices of solid hydrophobic

polymers containing the antibody, which matrices are in the form of shaped articles, e.g. films, or microcapsules.

The formulations to be used for *in vivo* administration are generally sterile. Sterility may be readily accomplished, e.g., by filtration through sterile filtration membranes.

G. Therapeutic Methods and Compositions

Any of the anti-TIM3 antibodies (or antigen binding proteins) or immunoconjugates of the anti-TIM3 antibodies (or antigen binding protein) conjugated to a cytotoxic agent, provided herein may be used in therapeutic methods.

In one aspect, an anti-TIM3 antibody or immunoconjugate of the anti-TIM3 antibody conjugated to a cytotoxic agent for use as a medicament is provided. In further aspects, an anti-TIM3 antibody or immunoconjugate of the anti-TIM3 antibody conjugated to a cytotoxic agent for use in treating cancer is provided. In certain embodiments, an anti-TIM3 antibody or immunoconjugates of the anti-TIM3 antibody conjugated to a cytotoxic agent for use in a method of treatment is provided. In certain embodiments, the invention provides an anti-TIM3 antibody or immunoconjugate of the anti-TIM3 antibody conjugated to a cytotoxic agent for use in a method of treating an individual having cancer comprising administering to the individual an effective amount of the anti-TIM3 antibody or the immunoconjugate of the anti-TIM3 antibody conjugated to a cytotoxic agent.

In further embodiments, the invention provides an anti-TIM3 antibody or immunoconjugate of the anti-TIM3 antibody conjugated to a cytotoxic agent for use in inducing apoptosis in a cancer cell/ or inhibiting cancer cell proliferation. In certain embodiments, the invention provides an anti-TIM3 antibody or immunoconjugate of the anti-TIM3 antibody conjugated to a cytotoxic agent for use in a method of inducing apoptosis in a cancer cell/ or inhibiting cancer cell proliferation in an individual comprising administering to the individual an effective of the anti-TIM3 antibody or immunoconjugate of the anti-TIM3 antibodies conjugated to a cytotoxic agent to induce apoptosis in a cancer cell/ or to inhibit cancer cell proliferation.

In further embodiments, the invention provides an anti-TIM3 antibody for use as immunestimulatory agent/ or stimulating IFN gamma secretion. In certain

embodiments, the invention provides an anti-TIM3 antibody for use in a method of immunestimulation/ or stimulating IFN gamma secretion in an individual comprising administering to the individual an effective of the anti-TIM3 antibody for immunestimulation/ or stimulating IFN gamma secretion.

5 An “individual” according to any of the above embodiments is preferably a human. In a further aspect, the invention provides for the use of an anti-TIM3 antibody or an immunoconjugate of the anti-TIM3 antibody conjugated to a cytotoxic agent in the manufacture or preparation of a medicament. In one embodiment, the medicament is for treatment of cancer. In a further embodiment, the medicament is
10 for use in a method of treating cancer comprising administering to an individual having cancer an effective amount of the medicament. In a further embodiment, the medicament is for inducing apoptosis in a cancer cell/ or inhibiting cancer cell proliferation. In a further embodiment, the medicament is for use in a method of inducing apoptosis in a cancer cell/ or inhibiting cancer cell proliferation in an
15 individual suffering from cancer comprising administering to the individual an amount effective of the medicament to induce apoptosis in a cancer cell/ or to inhibit cancer cell proliferation. An “individual” according to any of the above embodiments may be a human.

In a further aspect, the invention provides a method for treating cancer. In one
20 embodiment, the method comprises administering to an individual having cancer an effective amount of an anti-TIM3 antibody. An “individual” according to any of the above embodiments may be a human.

In a further aspect, the invention provides a method for inducing apoptosis in a cancer cell/ or inhibiting cancer cell proliferation in an individual suffering from
25 cancer. In one embodiment, the method comprises administering to the individual an effective amount of an anti-TIM3 antibody or an immunoconjugate of the anti-TIM3 antibody conjugated to a cytotoxic compound to induce apoptosis in a cancer cell/ or to inhibit cancer cell proliferation in the individual suffering from cancer. In one embodiment, an “individual” is a human.

30 In a further aspect, the invention provides pharmaceutical formulations comprising any of the anti-TIM3 antibodies provided herein, e.g., for use in any of the above therapeutic methods. In one embodiment, a pharmaceutical formulation comprises any of the anti-TIM3 antibodies provided herein and a pharmaceutically acceptable carrier.

An antibody of the invention (and any additional therapeutic agent) can be administered by any suitable means, including parenteral, intrapulmonary, and intranasal, and, if desired for local treatment, intralesional administration. Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. Dosing can be by any suitable route, e.g. by injections, such as intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic. Various dosing schedules including but not limited to single or multiple administrations over various time-points, bolus administration, and pulse infusion are contemplated herein.

Antibodies of the invention would be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners. The antibody need not be, but is optionally formulated with one or more agents currently used to prevent or treat the disorder in question. The effective amount of such other agents depends on the amount of antibody present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as described herein, or about from 1 to 99% of the dosages described herein, or in any dosage and by any route that is empirically/clinically determined to be appropriate.

For the prevention or treatment of disease, the appropriate dosage of an antibody of the invention (when used alone or in combination with one or more other additional therapeutic agents) will depend on the type of disease to be treated, the type of antibody, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments. Depending on the type and severity of the disease, about 1 $\mu\text{g/kg}$ to 15 mg/kg (e.g. 0.5 mg/kg - 10 mg/kg) of antibody can be an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. One typical daily dosage might range from about 1 $\mu\text{g/kg}$ to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment would generally be sustained until a

desired suppression of disease symptoms occurs. One exemplary dosage of the antibody would be in the range from about 0.05 mg/kg to about 10 mg/kg. Thus, one or more doses of about 0.5 mg/kg, 2.0 mg/kg, 4.0 mg/kg or 10 mg/kg (or any combination thereof) may be administered to the patient. Such doses may be administered intermittently, e.g. every week or every three weeks (*e.g.* such that the patient receives from about two to about twenty, or *e.g.* about six doses of the antibody). An initial higher loading dose, followed by one or more lower doses may be administered. An exemplary dosing regimen comprises administering an initial loading dose of about 4 mg/kg, followed by a weekly maintenance dose of about 2 mg/kg of the antibody. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

It is understood that any of the above formulations or therapeutic methods may be carried out using an immunoconjugate of the invention in place of or in addition to an anti-TIM3 antibody.

It is understood that any of the above formulations or therapeutic methods may be carried out using an immunoconjugate of the invention in place of or in addition to an anti-TIM3 antibody.

III. Articles of Manufacture

In another aspect of the invention, an article of manufacture containing materials useful for the treatment, prevention and/or diagnosis of the disorders described above is provided. The article of manufacture comprises a container and a label or package insert on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, IV solution bags, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is by itself or combined with another composition effective for treating, preventing and/or diagnosing the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is an antibody of the invention. The label or package insert indicates that the composition is used for treating the condition of choice. Moreover, the article of manufacture may comprise (a) a first container with a composition contained therein, wherein the composition comprises an antibody of the invention; and (b) a second container with a composition contained therein,

- wherein the composition comprises a further cytotoxic or otherwise therapeutic agent. The article of manufacture in this embodiment of the invention may further comprise a package insert indicating that the compositions can be used to treat a particular condition. Alternatively, or additionally, the article of manufacture may further comprise a second (or third) container comprising a pharmaceutically-acceptable buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.
- It is understood that any of the above articles of manufacture may include an immunoconjugate of the invention in place of or in addition to an anti-TIM3 antibody.

Description of the amino acid sequences

- | | |
|---------------|---|
| SEQ ID NO: 1 | heavy chain HVR-H1, Tim3_0016 |
| SEQ ID NO: 2 | heavy chain HVR-H2, Tim3_0016 |
| SEQ ID NO: 3 | heavy chain HVR-H3, Tim3_0016 |
| SEQ ID NO: 4 | light chain HVR-L1, Tim3_0016 |
| SEQ ID NO: 5 | light chain HVR-L2, Tim3_0016 |
| SEQ ID NO: 6 | light chain HVR-L3, Tim3_0016 |
| SEQ ID NO: 7 | heavy chain variable domain VH, Tim3_0016 |
| SEQ ID NO: 8 | light chain variable domain VL, Tim3_0016 |
| SEQ ID NO: 9 | heavy chain variable domain VH, Tim3_0016 variant (0018) |
| SEQ ID NO: 10 | light chain variable domain VL, Tim3_0016 variant (0018) |
| SEQ ID NO: 11 | light chain HVR-L1, Tim3_0016_HVR-L1 variant 1_NQ
(removal of glycosylation site by N to Q mutation) |
| SEQ ID NO: 12 | light chain HVR-L1, Tim3_0016_HVR-L1 variant 2_NS
(removal of glycosylation site by N to S mutation) |
| SEQ ID NO: 13 | heavy chain HVR-H1, Tim3_0021 |
| SEQ ID NO: 14 | heavy chain HVR-H2, Tim3_0021 |
| SEQ ID NO: 15 | heavy chain HVR-H3, Tim3_0021 |
| SEQ ID NO: 16 | light chain HVR-L1, Tim3_0021 |
| SEQ ID NO: 17 | light chain HVR-L2, Tim3_0021 |
| SEQ ID NO: 18 | light chain HVR-L3, Tim3_0021 |
| SEQ ID NO: 19 | heavy chain variable domain VH, Tim3_0021 |

	SEQ ID NO: 20	light chain variable domain VL, Tim3_0021
	SEQ ID NO: 21	heavy chain HVR-H1, Tim3_0022
	SEQ ID NO: 22	heavy chain HVR-H2, Tim3_0022
5	SEQ ID NO: 23	heavy chain HVR-H3, Tim3_0022
	SEQ ID NO: 24	light chain HVR-L1, Tim3_0022
	SEQ ID NO: 25	light chain HVR-L2, Tim3_0022
	SEQ ID NO: 26	light chain HVR-L3, Tim3_0022
	SEQ ID NO: 27	heavy chain variable domain VH, Tim3_0022
10	SEQ ID NO: 28	light chain variable domain VL, Tim3_0022
	SEQ ID NO: 29	heavy chain HVR-H1, Tim3_0026
	SEQ ID NO: 30	heavy chain HVR-H2, Tim3_0026
	SEQ ID NO: 31	heavy chain HVR-H3, Tim3_0026
15	SEQ ID NO: 32	light chain HVR-L1, Tim3_0026
	SEQ ID NO: 33	light chain HVR-L2, Tim3_0026
	SEQ ID NO: 34	light chain HVR-L3, Tim3_0026
	SEQ ID NO: 35	heavy chain variable domain VH, Tim3_0026
	SEQ ID NO: 36	light chain variable domain VL, Tim3_0026
20		
	SEQ ID NO: 37	heavy chain HVR-H1, Tim3_0028
	SEQ ID NO: 38	heavy chain HVR-H2, Tim3_0028
	SEQ ID NO: 39	heavy chain HVR-H3, Tim3_0028
	SEQ ID NO: 40	light chain HVR-L1, Tim3_0028
25	SEQ ID NO: 41	light chain HVR-L2, Tim3_0028
	SEQ ID NO: 42	light chain HVR-L3, Tim3_0028
	SEQ ID NO: 43	heavy chain variable domain VH, Tim3_0028
	SEQ ID NO: 44	light chain variable domain VL, Tim3_0028
30	SEQ ID NO: 45	heavy chain HVR-H1, Tim3_0030
	SEQ ID NO: 46	heavy chain HVR-H2, Tim3_0030
	SEQ ID NO: 47	heavy chain HVR-H3, Tim3_0030
	SEQ ID NO: 48	light chain HVR-L1, Tim3_0030
	SEQ ID NO: 49	light chain HVR-L2, Tim3_0030
35	SEQ ID NO: 50	light chain HVR-L3, Tim3_0030
	SEQ ID NO: 51	heavy chain variable domain VH, Tim3_0030
	SEQ ID NO: 52	light chain variable domain VL, Tim3_0030

	SEQ ID NO: 53	heavy chain HVR-H1, Tim3_0033
	SEQ ID NO: 54	heavy chain HVR-H2, Tim3_0033
	SEQ ID NO: 55	heavy chain HVR-H3, Tim3_0033
5	SEQ ID NO: 56	light chain HVR-L1, Tim3_0033
	SEQ ID NO: 57	light chain HVR-L2, Tim3_0033
	SEQ ID NO: 58	light chain HVR-L3, Tim3_0033
	SEQ ID NO: 59	heavy chain variable domain VH, Tim3_0033
	SEQ ID NO: 60	light chain variable domain VL, Tim3_0033
10	SEQ ID NO: 61	heavy chain HVR-H1, Tim3_0038
	SEQ ID NO: 62	heavy chain HVR-H2, Tim3_0038
	SEQ ID NO: 63	heavy chain HVR-H3, Tim3_0038
	SEQ ID NO: 64	light chain HVR-L1, Tim3_0038
15	SEQ ID NO: 65	light chain HVR-L2, Tim3_0038
	SEQ ID NO: 66	light chain HVR-L3, Tim3_0038
	SEQ ID NO: 67	heavy chain variable domain VH, Tim3_0038
	SEQ ID NO: 68	light chain variable domain VL, Tim3_0038
	SEQ ID NO: 69	an exemplary Pseudomonas exotoxin A variant 1
20		(deimmunized PE24 example)
	SEQ ID NO: 70	an exemplary Pseudomonas exotoxin A variant 2
		(deimmunized PE24 example)
	SEQ ID NO: 71	human kappa light chain constant region
25	SEQ ID NO: 72	human lambda light chain constant region
	SEQ ID NO: 73	human heavy chain constant region derived from IgG1
	SEQ ID NO: 74	human heavy chain constant region derived from IgG1 with mutations L234A and L235A
	SEQ ID NO: 75	human heavy chain constant region derived from IgG1 with mutations L234A, L235A and P329G
30	SEQ ID NO: 76	human heavy chain constant region derived from IgG4
	SEQ ID NO: 77	exemplary human TIM3 sequences
	SEQ ID NO: 78	human TIM3 Extracellular Domain (ECD)
	SEQ ID NO: 79	VH humanized version of Tim3_0016 variant (0018) (= Tim3-0433)
35	SEQ ID NO: 80	VL humanized version of Tim3_0016 variant (0018) (= Tim3-0433)

- 70 -

SEQ ID NO: 81 VH humanized version of Tim3_0016 variant (0018) (= Tim3-0434)

SEQ ID NO: 82 VL humanized version of Tim3_0016 variant (0018) (= Tim3-0434)

5 SEQ ID NO: 83 VH humanized version of Tim3-0028 (= Tim3-0438)

SEQ ID NO: 84 VL humanized version of Tim3-0028 (= Tim3-0438)

SEQ ID NO: 85 VH humanized version of Tim3-0028 (= Tim3-0443)

SEQ ID NO: 86 VL humanized version of Tim3-0028 (= Tim3-0443)

10 In the following the amino acid sequences of the VH und VL domains including marked HVRs (HVRs in bold, underlined letters) of anti-TIM3 antibodies Tim3-0016, Tim3-0016 variant (0018) and its humanized versions Tim3-0433 and Tim3-0434, Tim3-0021, Tim3-0022, Tim3-0026, Tim3-0028 and its humanized versions Tim3-0438 and Tim3-0443, Tim3-0030, and Tim3-0033, Tim3-0038 are listed:

15 **VH Tim3_0016:**

1 qvtlkesgpg ilqpsqtlrl tcsfsggfsls tsgmsvgwir
qp sgkglewl

20 51 ahiwlnddvf fnpalksrlt iskdtsnnqv flqiasvvta
dtatyycvra

101 ngylyaldyw gqgtsvtvss

25 **VL Tim3_0016:**

1 qivltqspai msaspgqkvt itcsasssvn ytqwyqqklg
sspk~~l~~wiyda

30 51 fklapgvpar fsgsgtgtsy sltissmeae daasyfchq
ssypwtfggg

101 tkleik

35

VH Tim3_0018(= VL Tim3_0016 variant):

1 qvtlkesgpg ilqpsqtlsl tcsfsggfsls tsgmsvgwir
qp sgkglewl

40

51 ahiwlnddvf fnpalkrrlt iskdtsnnqv flqiasvvta
dtatyycvra

- 71 -

101 ngylyaldyw gqgisvtvss

VL Tim3_0018 (= VL Tim3_0016 variant):

5

1 qivltqspai msaspgqkvt itcsasssvn ytqwyyqqklg
sspkllwida

10

51 fklapgvpar fsgsgtgtsy sltissmeae daasyfchq
ssypwtfggg

101 tkleik

15

VH humanized version of Tim3_0016 variant (0018) (= Tim3-0433)

20

1 qitlkesgpt lvkptqtltl tctfsgfsls tsgmsvgwir
qppgkglewl

51 ahiwlnddvf fnpalksrlt itkdtsknqv vltmtnmdpv
dtatyycvra

25

101 ngylyaldyw gqgtlvtvss

VL humanized version of Tim3_0016 variant (0018) (= Tim3-0433)

30

1 ettltqspaf msatpgdkvn iacssssvs ytqwyyqqkpg
eapklwida

51 fklapgippr fsgsgygtdf tltinniese daayyfchq
ssypwtfggg

35

101 tkleik

40

VH humanized version of Tim3_0016 variant (0018) (= Tim3-0434)

1 qitlkesgpt lvkptqtltl tctfsgfsls tsgmsvgwir
qppgkglewl

45

51 ahiwlnddvf fnpalksrlt itkdtsknqv vltmtnmdpv
dtatyycvra

101 ngylyaldyw gqgtlvtvss

- 72 -

VL humanized version of Tim3_0016 variant (0018) (= Tim3-0434

5 1 diqltqspsf lsasvgdrvt itcsasssvs ytqwyqqkpg
kapklwida

 51 fklapgvpsr fsgsgsgtef tltisslqpe dfatyfchq
10 ssypwtfgqg

 101 tkleik

VH Tim3_0021:

15 1 QVQLQQSGPQ LVRPGASVQI SCKASGYSFT SYLLHWLKQR
PGQGLEWIGM

 51 IDPSDSETRL NQKFKDKATL TVDKSSSTAY MQLSSPTSED
20 SAVYYCARDG

 101 YYAWYYFDCW GQTTTLTVSS

VL Tim3_0021:

25 1 DIVLTQSPAT LSVTPGDRVS LSCRASQSIG NNLHWYQQKS
HESPRLLIKY

 51 ASHSISGIPS KFSGTGSGTD FTLSFNSVET EDFGMYFCQQ
30 SNSWPLTFGA

 101 GTKLELK

VH Tim3_0022:

35 1 EVQLQESGPS LVKPSQTL~~SL~~ TCSVTGDSIA SAYWNWIRKF
PGNKLEYMGY

 51 INYSGSTYYN PSLKSRISIT RDTSQNQYYL QLNSVTTEDT
40 ATYYCVTGDY

 101 FDYWGRGTTL TVSS

VL aTim3_0022:

45

- 73 -

1 DIQMTQSPSS LSAYLGKVT ITCKARQDVR KNIGWYQHQP
GKGPRLLIWY

5 51 TSTLQSGIPS RFSGSGSGRD YSFNINNLEP EDIATYYCLQ
YDNLPFTFGT

101 GTKLEIR

VH Tim3_0026:

10 1 QIQLVQSGPE LKKPGETVKI SCKASGYTFT DYSMHVVKQA
PGRGLKWMGY

15 51 INTETYEPTF GADFKGRFAF SLDTSATTAY LQINSLKTED
TATFFCGGGG

101 YPAYWGQGT VIVSA

VL Tim3_0026

20 1 DVLMTQTPLS LPVSLGDQAS ISCRSSRTIL HSSGNTYLEW
YLQKPGQSPK

25 51 LLIYKVSNRF SGVPDRFSGS GSGTDFTLNI SRVEAEDLGV
YYCFQDSHVP

101 FTFGTGTKLE IK

VH Tim3_0028:

30 1 EVQLQQSVAE LVRPGASVKL SCTASGFNIK TTYMHVVKQR
PEQGLEWIGR

35 51 IDPADDNTKY APKFQ GKATI TADTSSNTAY LQLSSLTSED
AAIYYCVRDF

101 GYVAWFAYWG QGTLVTFSA

VL Tim3_0028:

40 1 NIVMTPTPKF LPVSSGDRVT MTCRASQSVD NYVAWYQQKP
GQSPKLLIYY

45 51 ASNR YIGVPD RFTGSGSGTD FTFTISSVQV EDLAVYFCQQ
HYSSPYTFGS

- 74 -

101 GTKLEIK

VH humanized version of Tim3-0028 (= Tim3-0438)

5

1 evqlvesggg lvqpqgslrl scaasgfnik ttymhwvrqa
pgkglewvgr51 idpaddntky apkfqgkati sadtskntay lqmnsdraed
10 tavyycvrdf101 gyvawfaywg qgtlvtvss

15 VL humanized version of Tim3-0028 (= Tim3-0438)

1 divmtqspls lpvtpgepas iscrassgsvd nyvawylqkp
gqspqlliyy51 asnryigvpd rfsgsgsgtd ftkisrvea edvgvyycqg
20 hysspytfqg

101 gtkveik

25

VH humanized version of Tim3-0028 (= Tim3-0443)1 evqlvesggg lvqpqgslrl scaasgfnik ttymhwvrqa
30 pgkglewvgr51 idpaddntky apkfqgkati sadtskntay lqmnsdraed
tavyycvrdf35 101 gyvawfaywg qgtlvtfss

VL humanized version of Tim3-0028 (= Tim3-0443)

40 1 divmtqspls lpvtpgepas iscrassgsvd nyvawylqkp
gqspqlliyy51 asnryigvpd rfsgsgsgtd ftkisrvea edvgvyycqg
45 hysspytfqg

101 gtkveik

-----.

- 75 -

VH aTim3_0030:

1 QIQLVQSGPE LKKPGETVKI SCKAS**GYPFS** **EYS**SIHWVKQA
PGKGLKWMVY

51 VNT**ETG**QPIV GDDFRGRFVL SLETSASTAY LQINNLLKNE
TATYFCGGG**G**

101 **YPAY**WGQGTI VTVSA

VL aTim3_0030:

1 DVLMTQTPLS LPVSLGDQAS ISCRS**SRSIV** **HSSGNTY**LEW
YLQKPGQSPK

51 LLIY**KVS**NRFS SGVPDRFSGS GSGTDFTLNI SRVEAEDLGV
YYCFQ**DSHVP**

101 **F**TFGTGTKLE IK

VH aTim3_0033:

1 QQQMHQSGAE LVKPGSSVKL SCKTS**GFTFS** **SSF**FISWLKQK
PGQSLEWIAW

51 IYA**ATG**STSY NQKFTNKAQL TVDTSSSAAY MQFSSLTTED
SAIYYCARHA

101 **GYPHY**YAMDY WGQGTSVTVS S

VL aTim3_0033:

1 DIQMTQSPAS LSASVGETVT ITCRA**SENIF** **SN**LAWYQQKQ
GKSPQLLVY**S**

51 **ATN**LGDGVPS RFSGSGSGTQ FSLKINSLQP EDFGNYYCQH
FYKIPFTFGT

101 GTKLEIK

VH aTim3_0038:

1 EVQLQQSGAE PLKPGASVKL TCTTS**GFNIK** **DY**YIHWVKQR
SDQGLEWIGR

- 76 -

51 IDPEDGELIY APKFQDKATI TVDTSSNIAY LQLNSLTSED
TAVYYCSRDH

101 GYVGWFAYWG QGTLVTVSA

VL aTim3_0038:

1 NVVMTQSPKS MIMSVGQRVT LNCKASENVD TYVSWYQQKP
EQSPKLLIYG

51 ASNRYTGVPD RFTGSRSATD FTLTISSVQA EDLAVYYCGQ
SYSYPWTFGG

101 GTKLEFR

In the following specific embodiments of the invention are listed:

1. An isolated antibody that binds to TIM3, wherein the antibody:
 - induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™)) of at least 45% after 120 Minutes at 37 °C (see Example 6)
 - In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™)) of at least 50% after 120 Minutes at 37 °C (see Example 6)
 - In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™)) of at least 55% after 120 Minutes at 37 °C (see Example 6)
 - In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™)) of at least 60% after 240 Minutes at 37 °C (see Example 6)
 - In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™)) of at least 65% after 240 Minutes at 37 °C (see Example 6)

2. The anti-TIM3 antibody according to embodiment 1, wherein the antibody:
- competes for binding to TIM3 with an anti-Tim3 antibody comprising the VH and VL of Tim3_0016
 - binds to a human and cynomolgous TIM3
 - shows as immunoconjugate a cytotoxic activity on TIM3 expressing cells (in one embodiment the immunoconjugates has a relative IC50 value of the cytotoxic activity as Pseudomonas exotoxin A conjugate on RPMI-8226 cells of 0.1 or lower (as measured in Example 11)
 - induces interferon-gamma release (in a MLR assay).
3. The antibody of embodiment 1, which is a human, humanized, or chimeric antibody.
4. The antibody of embodiment 1, which is an antibody fragment that binds to TIM3.
5. The antibody fragment of embodiment 4, which is Fab fragment.
6. The anti-TIM3 antibody according to any one of the preceding embodiments comprising
- A) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:3; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:4; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or
- B) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:3; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or
- C) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2; (c) HVR-H3

comprising the amino acid sequence of SEQ ID NO:3; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or

5 D) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:13; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:14; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:15; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:16; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:17; and (f) HVR-L3
10 comprising the amino acid sequence of SEQ ID NO:18; or

E) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:21; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:22; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:23; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:24; (e) HVR-L2
15 comprising the amino acid sequence of SEQ ID NO:25; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:26; or

F) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:29; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:30; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:31; (d) HVR-L1
20 comprising the amino acid sequence of SEQ ID NO:32; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:33; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:34; or

G) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38; (c) HVR-H3
25 comprising the amino acid sequence of SEQ ID NO:39; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42; or

H) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:45; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:46; (c) HVR-H3
30 comprising the amino acid sequence of SEQ ID NO:47; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:48; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:49; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:50; or.

- I) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:53; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:54; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:55; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:56; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:57; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:58; or
- J) (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:61; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:62; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:63; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:64; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:65; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:66.
7. An isolated antibody that binds to human TIM3, wherein the antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:3; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6.
8. The antibody according to embodiment 7, wherein the antibody comprises
- i) a VH sequence of SEQ ID NO:79 and a VL sequence of SEQ ID NO:80, or
- ii) a VH sequence of SEQ ID NO:81 and a VL sequence of SEQ ID NO:82.
9. The antibody according to embodiment 7, wherein the antibody comprises a VH sequence of SEQ ID NO:79 and a VL sequence of SEQ ID NO:80.
10. The antibody according to embodiment 7, wherein the antibody comprises a VH sequence of SEQ ID NO:81 and a VL sequence of SEQ ID NO:82.
11. An isolated antibody that binds to human TIM3, wherein the antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37; (b) HVR-H2 comprising the amino acid sequence of SEQ ID

- 5 NO:38; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:39; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42.
12. The antibody according to embodiment 11, wherein the antibody comprises
- i) a VH sequence of SEQ ID NO:83 and a VL sequence of SEQ ID NO:84, or
- 10 ii) a VH sequence of SEQ ID NO:85 and a VL sequence of SEQ ID NO:86.
13. The antibody according to embodiment 11, wherein the antibody comprises
- a VH sequence of SEQ ID NO:83 and a VL sequence of SEQ ID NO:84.
- 15 14. The antibody according to embodiment 11, wherein the antibody comprises
- a VH sequence of SEQ ID NO:85 and a VL sequence of SEQ ID NO:86.
15. The anti-TIM3 antibody to any one of the preceding embodiments comprising
- 20 A) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:4; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or
- 25 B) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:12; and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:13; or
- 30 C) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:11; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:12; and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:13; or

acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or

- 5 C) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6; or
- 10 D) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:13, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:14, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:15; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:16; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:17 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:18; or
- 15 E) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:21, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:22, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:23; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:24; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:25 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:26; or.
- 20 F) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:29, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:30, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:31; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:32; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:33 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:34; or
- 25 G) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:39; and (b) a VL domain comprising (i) HVR-L1
- 30

comprising the amino acid sequence of SEQ ID NO:40; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42; or

5 H) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:45, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:46, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:47; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:48; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:49 and (iii) HVR-L3
10 comprising the amino acid sequence of SEQ ID NO:50; or

I) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:53, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:54, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:55; and (b) a VL domain comprising (i) HVR-L1
15 comprising the amino acid sequence of SEQ ID NO:56; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:57 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:58; or

J) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:61, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:62, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:63; and (b) a VL domain comprising (i) HVR-L1
20 comprising the amino acid sequence of SEQ ID NO:64; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:65 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:66.

25 16. An isolated antibody that binds to human TIM3, wherein the antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising
30 (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6.

17. The antibody according to embodiment 16, wherein the antibody comprises
- i) a VH sequence of SEQ ID NO:79 and a VL sequence of SEQ ID NO:80,or
- 5 ii) a VH sequence of SEQ ID NO:81 and a VL sequence of SEQ ID NO:82.
18. The antibody according to embodiment 16, wherein the antibody comprises
- a VH sequence of SEQ ID NO:79 and a VL sequence of SEQ ID NO:80.
- 10 19. The antibody according to embodiment 16, wherein the antibody comprises
- a VH sequence of SEQ ID NO:81 and a VL sequence of SEQ ID NO:82.
20. An isolated antibody that binds to human TIM3, wherein the antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:39; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42.
- 15 20 21. The antibody according to embodiment 20, wherein the antibody comprises
- i) a VH sequence of SEQ ID NO:83 and a VL sequence of SEQ ID NO:84,or
- 25 ii) a VH sequence of SEQ ID NO:85 and a VL sequence of SEQ ID NO:86.
22. The antibody according to embodiment 20, wherein the antibody comprises
- a VH sequence of SEQ ID NO:83 and a VL sequence of SEQ ID NO:84.

23. The antibody according to embodiment 20, wherein the antibody comprises

a VH sequence of SEQ ID NO:85 and a VL sequence of SEQ ID NO:86.

24. The anti-TIM3 antibody according to any one of the preceding
5 embodiments comprising

A) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence
of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID
NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from
SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the
10 amino acid sequence of SEQ ID NO:4; (ii) HVR-L2 comprising the amino
acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid
sequence of SEQ ID NO:6; or

B) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence
of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID
15 NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from
SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the
amino acid sequence of SEQ ID NO:11; (ii) HVR-L2 comprising the amino
acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid
sequence of SEQ ID NO:6; or

C) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence
of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID
20 NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from
SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the
amino acid sequence of SEQ ID NO:12; (ii) HVR-L2 comprising the amino
25 acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid
sequence of SEQ ID NO:6; or

D) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence
of SEQ ID NO:13, (ii) HVR-H2 comprising the amino acid sequence of SEQ
ID NO:14, and (iii) HVR-H3 comprising an amino acid sequence selected
30 from SEQ ID NO:15; and (b) a VL domain comprising (i) HVR-L1
comprising the amino acid sequence of SEQ ID NO:16; (ii) HVR-L2
comprising the amino acid sequence of SEQ ID NO:17 and (iii) HVR-L3
comprising the amino acid sequence of SEQ ID NO:18; or

- 5 E) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:21, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:22, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:23; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:24; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:25 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:26; or.
- 10 F) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:29, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:30, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:31; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:32; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:33 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:34; or
- 15 G) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:39; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42; or
- 20 H) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:45, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:46, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:47; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:48; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:49 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:50; or
- 25 I) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:53, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:54, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:55; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:56; (ii) HVR-L2
- 30

comprising the amino acid sequence of SEQ ID NO:57 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:58; or

- J) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:61, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:62, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:63; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:64; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:65 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:66;

wherein the antibody is characterized independently by one or more of the following properties: the anti-TIM3 antibody

- induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 45% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 50% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 55% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 60% after 240 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 65% after 240 Minutes at 37 °C (see Example 6)
- competes for binding to TIM3 with an anti-Tim3 antibody comprising the VH of SEQ ID NO:7 and VL of SEQ ID NO:8.
- binds to a human and cynomolgous TIM3
- shows as immunoconjugate a cytotoxic activity on TIM3 expressing cells (in one embodiment the immunoconjugates has a relative IC50 value of the cytotoxic activity as Pseudomonas exotoxin A conjugate on RPMI-8226 cells of 0.1 or lower (as measured in Example 11)

- induces interferon-gamma release (in a Mixed Lymphocyte Reaction (MLR) assay as described in Example 5).

25. An isolated antibody that binds to human TIM3, wherein the antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:3; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:12; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:6;

wherein the antibody is characterized independently by one or more of the following properties: the anti-TIM3 antibody

- induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 45% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 50% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 55% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 60% after 240 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 65% after 240 Minutes at 37 °C (see Example 6)
- competes for binding to TIM3 with an anti-Tim3 antibody comprising the VH of SEQ ID NO:7 and VL of SEQ ID NO:8.
- binds to a human and cynomolgous TIM3
- shows as immunoconjugate a cytotoxic activity on TIM3 expressing cells (in one embodiment the immunoconjugates has a relative IC50 value of the cytotoxic activity as Pseudomonas

exotoxin A conjugate on RPMI-8226 cells of 0.1 or lower (as measured in Example 11)

- induces interferon-gamma release (in a Mixed Lymphocyte Reaction (MLR) assay as described in Example 5).

- 5 26. An isolated antibody that binds to human TIM3, wherein the antibody comprises G) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:39; and (b) a VL domain
- 10 comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42;

15 wherein the antibody is characterized independently by one or more of the following properties: the anti-TIM3 antibody

- induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 45% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 50% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 55% after 120 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 60% after 240 Minutes at 37 °C (see Example 6)
- In on embodiment the antibody induces internalization of TIM3 (in a FACS assay on TIM3 expressing RPMI8226 cells) of at least 65% after 240 Minutes at 37 °C (see Example 6)
- competes for binding to TIM3 with an anti-Tim3 antibody comprising the VH of SEQ ID NO:7 and VL of SEQ ID NO:8.
- binds to a human and cynomolgous TIM3
- shows as immunoconjugate a cytotoxic activity on TIM3 expressing cells (in one embodiment the immunoconjugates has a

- relative IC50 value of the cytotoxic activity as *Pseudomonas* exotoxin A conjugate on RPMI-8226 cells of 0.1 or lower (as measured in Example 11)
- induces interferon-gamma release (in a Mixed Lymphocyte Reaction (MLR) assay as described in Example 5).
- 5
27. The antibody according to any one of the preceding embodiments , which is a full length IgG1 antibody.
28. The antibody of according to any one of the preceding embodiments , which is a full length IgG1 antibody with L234A, L235A and P329G (numbering according to the EU index of Kabat).
- 10
29. Isolated nucleic acid encoding the antibody according to any one of the preceding embodiments.
30. A host cell comprising the nucleic acid of embodiment 29.
31. A method of producing an antibody comprising culturing the host cell of embodiment 30 so that the antibody is produced.
- 15
32. The method of embodiment 31, further comprising recovering the antibody from the host cell.
33. An immunoconjugate comprising the antibody according any one of embodiments 1 to 28 and a cytotoxic agent.
- 20
34. The immunoconjugate according to embodiment 33 wherein the cytotoxic agent is *Pseudomonas* Exotoxin A or an Amatoxin.
35. A pharmaceutical formulation comprising the antibody according any one of embodiments 1 to 28 and a pharmaceutically acceptable carrier.
- 25
36. The antibody according any one of embodiments 1 to 28 or the immunoconjugate of any one of embodiments 33 or 34 for use as a medicament.
37. The antibody according any one of embodiments 1 to 28 or the immunoconjugate of any one of embodiments 33 or 34 for use in treating cancer.

38. Use of the antibody according any one of embodiments 1 to 28 or the immunoconjugate of any one of embodiments 33 or 34 in the manufacture of a medicament.
39. The use of embodiment 38, wherein the medicament is for treatment of cancer.
40. A method of treating an individual having cancer comprising administering to the individual an effective amount of the antibody of embodiment 1 or the immunoconjugate of embodiment 33.

V. EXAMPLES

The following are examples of methods and compositions of the invention. It is understood that various other embodiments may be practiced, given the general description provided above.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, the descriptions and examples should not be construed as limiting the scope of the invention. The disclosures of all patent and scientific literature cited herein are expressly incorporated in their entirety by reference.

Example 1a: Generation of anti-TIM3 antibodies

Immunization of mice

NMRI mice were immunized genetically, using a plasmid expression vector coding for full-length human Tim-3 by intradermal application of 100 ug vector DNA (plasmid 15304_hTIM3-fl), followed by Electroporation (2 square pulses of 1000 V/cm, duration 0.1 ms, interval 0.125 s; followed by 4 square pulses of 287.5 V/cm, duration 10 ms, interval 0.125 s. Mice received either 6 consecutive immunizations at days 0, 14, 28, 42, 56, 70, and 84. Blood was taken at days 36, 78 and 92 and serum prepared, which was used for titer determination by ELISA (see below). Animals with highest titers were selected for boosting at day 96, by intravenous injection of 50 ug of recombinant human Tim-3 human Fc chimera, and monoclonal antibodies were isolated by hybridoma technology, by fusion of splenocytes to myeloma cell line 3 days after boost.

Determination of serum titers (ELISA)

Human recombinant Tim-3 human Fc chimera was immobilized on a 96-well NUNC Maxisorp plate at 0.3 ug/ml, 100 µl/well, in PBS, followed by: blocking of the plate with 2% Crotein C in PBS, 200 µl/well; application of serial dilutions of antisera, in duplicates, in 0.5% Crotein C in PBS, 100 µl/well; detection with HRP-conjugated goat anti-mouse antibody (Jackson ImmunoResearch/Dianova 115-036-071; 1/16 000). For all steps, plates were incubated for 1 h at 37⁰ C. Between all steps, plates were washed 3 times with 0.05% Tween 20 in PBS. Signal was developed by addition of BM Blue POD Substrate soluble (Roche), 100 ul/well; and stopped by addition of 1 M HCl, 100 ul/well. Absorbance was read out at 450 nm, against 690 nm as reference. Titer was defined as dilution of antisera resulting in half-maximal signal.

Example 1b: Characterization anti-Tim3 antibodies**ELISA for Tim3**

Nunc-Maxi Sorp Streptavidine plates (MicroCoat #11974998/MC1099) were coated by 25 µl/well with Tim3-ECD-His-Biotin (biotinylated with BirA Ligase) and incubated at RT for 1 h while shaking at 400 rpm rotation. After washing (3x90 µl/well with PBST-buffer) 25 µl aTim3 samples or diluted (1:2 steps) reference antibody aTim3 F38-2E2 (Biolegend) was added and incubated 1h at RT. After washing (3x90 µl/well with PBST-buffer) 25µl/well sheep-anti-mouse-POD (GE NA9310V) was added in 1:9000 dilution and incubated at RT for 1 h while shaking at 400 rpm rotation. After washing (4x90 µl/well with PBST-buffer) 25 µl/well TMB substrate (Calbiochem, #CL07) was added and incubated until OD 1.5 – 2.5. Then the reaction was stopped by addition of 25 µl/well 1N HCL-solution. Measurement took place at 370/492 nm.

ELISA results are listed as EC50-values [ng/ml] in summary Table 2 below.

Example: Cell ELISA for Tim3

Adherent CHO-K1 cell line stably transfected with plasmid 15312_hTIM3-fl_pUC_Neo coding for full-length human Tim3 and selection with G418 (Neomycin resistance marker on plasmid) were seeded at a concentration of 1.2x10E6 cells/ml into 384-well flat bottom plates and grown over night.

At the next day 25 µl/well Tim3 sample or aTim3 reference antibody F38-2E2 Azide free (Biolegend, 354004) was added and incubated for 2h at 4°C (to avoid internalization). After washing (3x90µl/well PBST (BIOTEK Washer: Prog. 29, 1 x 90) cells were fixed by flicking out residual buffer and addition of 50µl/well 0,05% Glutaraldehyde: Dilution 1:500 of 25% Glutaraldehyde (Sigma Cat.No: G5882) in 1xPBS-buffer and incubated for 1h at RT. After washing (3x90µl/well PBST (BIOTEK Washer: Prog. 21, 3x90 GreinLysin) 25 µl/well secondary antibody was added for detection (Sheep-anti-mouse-POD; Horseradish POD linked F(ab')₂ Fragment ; GE NA9310) followed by 2h incubation at RT while shaking at 400 rpm. After washing (3x90µl/well PBST (BIOTEK Washer: Prog. 21, 3x90 GreinLysin) 25 µl/well TMB substrate solution (Roche 11835033001) was added and incubated until OD 1.5 – 2.5. Then the reaction was stopped by addition of 25 µl/well 1N HCL-solution. Measurement took place at 370/492 nm.

Cell ELISA results are listed as “EC50 CHO-Tim3”-values [ng/ml] in summary Table 2 below.

Table 2: Binding affinities of exemplary antibodies (ELISA and BIACORE)

Assay	Tim3_0018	Tim3_0021	Tim3_0028	Tim3_0026	Tim3_0033	Tim3_0038
Affinity KD [nM]						
monomer / dimer Tim3	3.4 / 1.1	204 / 4.1	173 / 2.8	6.2 / 1.5	n.f. / 3.1	7.6 / 0.6
EC50 ELISA [nM]	0.56		0.22			0.501
EC50 ELISA [ng/ml]	94	47	37	47	1321	83
EC50 CHO-Tim3 [nM]	0.52		0.32			0.17
EC50 CHO-Tim3 [ng/ml]	87	73	53	69	3710	29

BIACore characterization of the Tim3 ABs

A surface plasmon resonance (SPR) based assay has been used to determine the kinetic parameters of the binding between several murine Tim3 binders as well as commercial human Tim3 binding references. Therefore, an anti-mouse IgG was immobilized by amine coupling to the surface of a (BIAcore) CM5 sensor chip. The samples were then captured and monomeric hu/cy Tim3-ECD as well as a Fc-

tagged human Tim3-ECD dimer was bound to them. The sensor chip surface was regenerated after each analysis cycle. The equilibrium constant K_D was finally gained by fitting the data to a 1:1 Langmuir interaction model.

5 About 12000 response units (RU) of 30 $\mu\text{g/ml}$ anti-mouse IgG (GE Healthcare #BR-1008-38) were coupled onto the spots 1,2,4 and 5 of the flow cells 1-4 (spots 1,5 are active and spots 2,4 are reference spots) of a CM5 sensor chip in a BIAcore B4000 at pH 5.0 by using an amine coupling kit supplied by GE Healthcare.

10 The sample and running buffer was HBS-EP+ (0.01 M HEPES, 0.15 M NaCl, 3 mM EDTA, 0.05% v/v Surfactant P20, pH 7.4). Flow cell temperature was set to 25 °C and sample compartment temperature to 12 °C. The system was primed with running buffer.

15 The samples were injected for 30 seconds with a concentration of 200 $\mu\text{g/ml}$ and bound to the spots 1 and 5 of each flow cell, allowing the measurement of eight samples in parallel. Then a complete set of different (monomeric cyno, monomeric human and huFc fused dimeric human Tim3-ECD) concentrations (s. Table X) was injected over each sample for 240 s followed by a dissociation time of 30/1800 s (s. Table 1). Each analysis cycle (sample capture, spot 1 and 5 – Tim3 ECD injection) was then regenerated with a 30 seconds long injection of Glycine-HCl pH 1.7. The flow rate was set to 30 $\mu\text{l/min}$ for the whole run.

20 Finally the double referenced data was fitted to a 1:1 Langmuir interaction model with the BIAcore B4000 Evaluation Software. Resulting affinities to monomeric human, cyno Tim3 and huFc fused dimeric human Tim3 are shown in Table 2 and 3. The affinity to the hu Tim3 dimer is most likely affected by avidity and therefore apparently stronger than the affinity to the monomeric huTim3.

25 **Table 3a: Binding affinities determined by BIAcore-KD values gained by a kinetic SPR measurement.-n.f. means no fit possible, most likely due to no or weak binding.**

Sample	huTim3 K_D (25 °C) [M]	huTim3Fc K_D (25 °C) [M]	cyTim3 K_D (25 °C) [M]
TIM3-0016	3.29E-09	1.09E-09	2.16E-08
TIM3-0016 variant (0018)	3.40E-09	1.11E-09	4.19E-08
TIM3-0021	2.04E-07	4.07E-09	n.f.
TIM3-0022	1.26E-07	1.52E-09	2.84E-08
TIM3-0026	6.23E-09	1.52E-09	n.f.

Sample	huTim3 K_D (25 °C) [M]	huTim3Fc K_D (25 °C) [M]	cyTim3 K_D (25 °C) [M]
TIM3-0028	1.73E-07	2.77E-09	n.f.
TIM3-0030	3.11E-09	1.28E-09	n.f.
TIM3-0033	n.f.	3.05E-09	n.f.
TIM3-0038	7.56E-09	5.69E-10	n.f.
Reference antibody Biolegend F38-2E2	1.36E-08	7.50E-09	1.68E-07
Reference antibody USB 11E365	1.34E-08	7.73E-09	1.41E-07

Determination of the affinity to Tim3 via SPR (Chimeric TIM3-0016 variant (0018) and humanized versions)

Protein A was immobilized by amine coupling to the surface of a (Biacore) CM5 sensor chip. The samples were then captured and hu Tim3-ECD was bound to them. The sensor chip surface was regenerated after each analysis cycle. The equilibrium constant and kinetic rate constants were finally gained by fitting the data to a 1:1 langmuir interaction model.

About 2000 response units (RU) of 20 µg/ml Protein A were coupled onto the spots 1, 2, 4 and 5 of all flow cells of a CM5 sensor chip in a Biacore B4000 instrument using an amine coupling kit supplied by GE Healthcare.

The sample and running buffer was HBS-EP+ (0.01 M HEPES, 0.15 M NaCl, 3 mM EDTA, 0.05 % v/v Surfactant P20, pH 7.4). Flow cell temperature was set to 25 °C and sample compartment temperature to 12 °C. The system was primed with running buffer.

Different samples were injected for 30 seconds with a concentration of 10 nM and bound consecutively to the spots 1 and 5 in all flow cells. Then a complete set of monomeric human Tim3-ECD dilutions (600 nM, 200 nM, 66.7 nM, 2×22.2 nM, 7.4 nM, 2.5 nM and 2×0 nM) was consecutively injected over each sample for 300s. Each antigen injection was followed by a dissociation time of 12s/1000s and two 30s regeneration steps with a Glycine-HCl pH 1.5 solution, of which the last one contained a stabilization period after injection of 20 seconds.

Finally the double referenced data was fitted to a 1:1 langmuir interaction model using the Biacore B4000 Evaluation Software. Resulting K_D values are shown in Table 3b.

Determination of the affinity to Tim3 via SPR ((Chimeric TIM3-0028 and humanized versions))

5

Anti-human Fc IgG was immobilized by amine coupling to the surface of a (Biacore) CM5 sensor chip. The samples were then captured and hu Tim3-ECD was bound to them. The sensor chip surface was regenerated after each analysis cycle. The equilibrium constant and kinetic rate constants were finally gained by fitting the data to a 1:1 langmuir interaction model.

10

About 2500 response units (RU) of 10 μ g/ml anti-human Fc IgG (GE Healthcare #BR-1008-39) were coupled onto the spots 1, 2, 4 and 5 of all flow cells of a CM5 sensor chip in a Biacore B4000 instrument using an amine coupling kit supplied by GE Healthcare.

15

The sample and running buffer was HBS-EP+ (0.01 M HEPES, 0.15 M NaCl, 3 mM EDTA, 0.05 % v/v Surfactant P20, pH 7.4). Flow cell temperature was set to 25 °C and sample compartment temperature to 12 °C. The system was primed with running buffer.

20

Different samples were injected for 30 seconds with a concentration of 10 nM and bound consecutively to the spots 1 and 5 in all flow cells. Then a complete set of monomeric human Tim3-ECD dilutions (600 nM, 200 nM, 66.7 nM, 2×22.2 nM, 7.4 nM, 2.5 nM and 2×0 nM) was consecutively injected over each sample for 300s. Each antigen injection was followed by a dissociation time of 12s/700s and two 30s regeneration steps with a 3 M $MgCl_2$ solution, of which the last one contained an “extra wash after injection” with running buffer.

25

Finally the double referenced data was fitted to a 1:1 langmuir interaction model using the Biacore B4000 Evaluation Software. Resulting K_D values are shown in Table 3b.

Table 3b: Binding affinities determined by BIAcore-KD values gained by a kinetic SPR measurement

Sample	huTim3 K_D (25 °C) [M]
Chimeric TIM3-0016 variant (0018)	2.78E-09
TIM3-0433	5.74E-09
TIM3-0434	5.76E-09
Chimeric TIM3-0028	2.35E-07
TIM3-0438	3.05E-07
TIM3-0443	2.87E-07

Example 2: Generation of anti-Tim3 antibodies derivatives

5 Chimeric antibody derivatives

Chimeric Tim3 antibodies were generated by amplifying the variable heavy and light chain regions of the anti-TIM3 mouse antibodies Tim3-0016, Tim3-0016 variant (0018), Tim3-0021, Tim3-0022, Tim3-0026, Tim3-0028, Tim3-0030, and Tim3-0033, Tim3-0038 from via PCR and cloning them into heavy chain expression vectors as fusion proteins with human IgG1 backbones / human CH1-Hinge-CH2-CH3 with LALA and PG mutations (Leucine 234 to Alanine, Leucine 235 to Alanine, Proline 329 to Glycine) abrogating effector functions and light chain expression vectors as fusion proteins to human C-kappa. LC and HC Plasmids were then cotransfected into HEK293 and purified after 7 days from supernatants by standard methods for antibody purification.

Removal of glycosylation site NYT: Modifying 1 HVR-L1 position in Tim3-0016, Tim3_0016 variant (named 0018 or Tim3_0018) by substitution of N by Q or S

Mutations within the variable light chain region of Tim3_0016 and Tim3_0016 variant (0018) were generated by in vitro mutagenesis using Agilent “Quick Change Lightning Site-directed Mutagenesis Kit” according manufacturer’s instructions. By this method the asparagine (N) of the glycosylation site motif NYT in the light chain HVR-L1 (SEQ ID NO: 4) was replaced by glutamine (Q) (resulting in SEQ ID NO: 11 = Tim3_0016_HVR-L1 variant 1_ NQ) or, alternatively, the asparagine (N) was replaced by serine (S) (resulting in SEQ ID NO: 12 = Tim3_0016_HVR-L1 variant 2_ NS). In both, the glycosylation site

motif NYT was successfully modified. LC and HC Plasmids coding for the variants were then cotransfected into HEK293 and purified after 7 days from supernatants by standard methods for antibody purification.

5 The generated mutants were tested by ELISA on human Tim3, ELISA on cynomolgus Tim3 and cellular ELISA on adherent CHO-K1 cells expressing full-length human Tim3.

TABLE 4:

Antibodies and mutant antibodies tested	Biochem Human		Biochem Cyno		Cellular binding CHO-TIM3	
	EC50 [ng/ml] values in relation to the samples max value	Inflexion point [ng/ml]	EC50 [ng/ml] values in relation to the samples max value	Inflexion point [ng/ml]	EC50 [ng/ml] values in relation to the samples max value	Inflexion point [ng/ml]
Anti Tim3 F38-2E2	73.2	88.3	423.0	209871.3	150.2	224.3
Tim3_0018 (TIM3-0016 variant)	15.1	15.3	14.6	14.6	26.4	29.4
Tim3_0018MutNQ	12.0	10.8	13.2	10.8	13.4	12.8
Tim3_0018MutNS	10.3	6.5	11.9	6.5	11.2	11.1
Tim3_0016MutNQ	7.6	5.7	8.3	5.7	6.3	5.4
Tim3_0016MutNS	8.5	5.5	9.7	5.5	9.1	8.5

10 All mutants generated were found to show even more functional binding to human TIM3 (human), cyno TIM3 (cyno) or human TIMR on CHO cells than the parental antibodies Tim3_0016 or the Tim3_0016 antibody variant Tim3_0018 respectively.

Humanized antibody derivatives

Humanization of the VH and VL domains of murine anti-Tim3-0016 variant (0018) and anti-Tim3_0028

15 Based upon the amino acid sequence of the VH and VL domains of a) anti-Tim3 antibody Tim3_0016 variant (0018) (with the amino acid sequences of the 6 HVRs wherein in the light chain the HVR-L1 variant 2_NS (removal of glycosylation site by N to S mutation) was used (SEQ ID NOs: 1, 2,3,12 ,5 and 6) humanized anti-Tim3 antibody variants Tim3-0433 and Tim3-0434 were generated and based

upon the amino acid sequence of the VH and VL domains of b) anti-Tim3 antibody Tim3_0028 (with the amino acid sequences of the 6 HVRs (SEQ ID NOs: 37, 38,39,40 ,41 and 42) humanized anti-Tim3 antibody variants Tim3-0438 and Tim3-0443 were generated.

- 5 The humanized amino acid sequences for heavy and light chain variable regions of were backtranslated in to DNA and the resulting cNDA were synthesized (GenArt) and then cloned into heavy chain expression vectors as fusion proteins with human IgG1 backbones /human CH1-Hinge-CH2-CH3 with LALA and PG mutations (Leucine 234 to Alanine, Leucine 235 to Alanine, Proline 329 to Glycine)
- 10 abrogating effector functions or into light chain expression vectors as fusion proteins to human C-kappa. LC and HC Plasmids were then cotransfected into HEK293 and purified after 7 days from supernatants by standard methods for antibody purification. The resulting humanized Tim3-antibodies are named as follows:

15 **Table: VH and VL sequences of humanized antibodies**

Humanized antibodies of Tim3_0016 variant (0018)	VH/SEQ ID NO:	VL/SEQ ID NO:
Tim3-0433	SEQ ID NO: 79	SEQ ID NO: 80
Tim3-0434	SEQ ID NO: 81	SEQ ID NO: 82
Humanized antibodies of Tim3_0028	VH/SEQ ID NO:	VL/SEQ ID NO:
Tim3-0438	SEQ ID NO: 83	SEQ ID NO: 84
Tim3-0443	SEQ ID NO: 85	SEQ ID NO: 86

Table: HVR sequences of humanized antibodies

Humanized antibodies of Tim3_0016 variant (0018)	HVR-H1, HVR-H2, and HVR-H3 /SEQ ID NOs:	HVR-L1, HVR-L2, and HVR-L3 t/SEQ ID NOs:
Tim3-0433	SEQ ID NOs: 1 , 2 and 3	SEQ ID NOs: 12 , 5 and 6
Tim3-0434	SEQ ID NOs: 1 , 2 and 3	SEQ ID NOs: 12 , 5 and 6
Humanized antibodies of Tim3_0028	HVR-H1, HVR-H2, and HVR-H3 /SEQ ID NOs:	HVR-L1, HVR-L2, and HVR-L3 /SEQ ID NOs:
Tim3-0438	SEQ ID NOs: 37 , 38 and 39	SEQ ID NOs: 40 , 41 and 42
Tim3-0443	SEQ ID NOs: 37 , 38 and 39	SEQ ID NOs: 40 , 41 and 42

Example 3: Fluorescent Labeling of Purified Monoclonal Antibody

The fluorescent labeling of the hybridoma derived monoclonal antibody was carried out by using Alexa Fluor 488 Monoclonal Antibody Labeling Kit (manufactured by Invitrogen) according to the manufacturer's instructions. After the labeling, each antibody was confirmed to be positively labeled with Alexa Fluor 488 (hereinafter referred to as "Alexa-488") by FACSCalibur (manufactured by BD Biosciences) analysis for TIM-3 expressing RPMI-8226 and Pfeiffer cells.

Example 4: Classification of Binding Epitope Groups using FACS based Competition Assay

The relation of epitopes between generated anti-TIM3 antibodies and six anti-TIM3 reference antibodies was analyzed by a FACS based binding competition assay. The TIM3 reference antibodies were the following: antibodies 4177 and 8213 as described in US2012/189617, antibodies 1.7E10 and 27.12E12 as described in WO2013/06490; antibody 344823 (Clone 344823, manufactured by R&D Systems) and antibody F38-2E2 (Clone F38-2E2, manufactured by BioLegend and R&D Systems). In brief, the test antibody was allowed to interact and bind with the TIM-3 expressing RPMI-8226 cells (ATCC ® CCL-155™) and then it was evaluated by flow cytometry method whether another anti-TIM-3 antibody could also bind to TIM-3 expressing cells.

In short human TIM3 expressing RPMI-8226 cells were incubated with BD human Fc Block for 10 min at RT and stained in two different experimental setups to exclude the impact of the difference in the affinity of the tested antibodies on the binding:

- 5 1) with disclosed purified anti-TIM3 (10µg/ml in BD staining buffer for 0.5h at 4°C), which were conjugated with Alexa*488 according to the manufacturer's instructions (Molecular Probes A-20181) with an average of 2.7 fluorophores per antibody. Then a) unlabeled (1-4) reference recombinant anti-TIM3 antibodies or Isotype control were added (10µg/ml) for 0.5h at 4°C in BD SB and after washing
10 with BD SB stained with PE-labeled anti-huFcγ Abs (JIR, 109-116-098, 1:200, 0.5h at 4°C in BD SB) or b) PE labeled (5-6) available reference anti-TIM3 antibodies or appropriate Isotype controls were added (10µg/ml) for 0.5h at 4°C in BD SB. After washing and centrifugation MFI signals of stained RPMI-8226 cells were analyzed by BD Biosciences FACSCanto flow cytometer.

15 **Table 5: Summary of epitope characterization- competition of Tim3-antibodies for binding to Tim3**

	Max % Inhibition of Binding					
	Epitope group 1			Epitope group 3		
	1a		1b	3a	3b	
	Tim3_ 0016	Tim3_ 0018	Tim3_ 0026	Tim3_ 0022	Tim3_ 0028	Tim3_ 0038
clone 4177	1	-9	29	79	-3	0
clone 8213	-2	9	9	9	38	29
clone 1- 7E10	-5	15	24	0	20	7
clone 27- 12E12	-1	4	22	40	82	94
clone 344823	0	0	3	102	107	99
clone F38- 2E2	-7	6	2	77	75	94

Results from the FACS based epitope groups mapping show that Tim3_0016 and Tim3_0016 variant Tim3_0018 show no binding competition with any tested anti-

TIM-3 reference antibodies and it was suggested that these Abs recognized the new epitope different from the epitopes to which all previous described TIM3 reference antibodies recognized whereas Tim3_0022, Tim3_0026, Tim3_0028 and Tim3_0038 compete to different extend for binding to surface expressed TIM3 on JRPML-8226 cells with various competitors.

Example 5: Effect of human anti-TIM-3 Antibodies on Cytokine Production in a Mixed Lymphocyte Reaction (MLR)

A mixed lymphocyte reaction was used to demonstrate the effect of blocking the TIM-3 pathway to lymphocyte effector cells. T cells in the assay were tested for activation and IFN-gamma secretion in the presence or absence of an anti-TIM-3 mAbs.

Human Lymphocytes were isolated from peripheral blood of healthy donor by density gradient centrifugation using Leukosep (Greiner Bio One, 227 288). Briefly, heparinized blood were diluted with the three fold volume of PBS and 25 ml aliquots of the diluted blood were layered in 50 ml Leukosep tubes. After centrifugation at 800 x g for 15 min at room temperature (w/o break) the lymphocyte containing fractions were harvested, washed in PBS and used directly in functional assay or resuspended in freezing medium (10% DMSO, 90 %FCS) at 1.0E+07 cells/ml and stored in liquid nitrogen. 1:1 target/responder cell ratio was used in MLR assay (i.e. each MLR culture contained - 2.0E+05 PBMCs from each donor in a total volume of 200 µl. Anti-TIM3 monoclonal antibodies Tim3_0016, Tim3_0016 variant (Tim3_0018), Tim3_0021, Tim3_0022, Tim3_0026, Tim3_0028, Tim3_0030, Tim3_0033, Tim3_0038 and F38-2E2 (BioLegend), were added to each culture at different antibody concentrations. Either no antibody or an isotype control antibody was used as a negative control and rec hu IL-2 (20 EU/ml) was used as positive control. The cells were cultured for 6 days at 37°C. After day 6 100 µl of medium was taken from each culture for cytokine measurement. The levels of IFN-gamma were measured using OptEIA ELISA kit (BD Biosciences).

The results are shown in Table 6 (IFN-g secretion/release). The anti-TIM-3 monoclonal antibodies promoted T cell activation and IFN-gamma secretion in concentration dependent manner. The anti-TIM3 antibodies Tim3_0021, Tim3_0022, Tim3_0028, and Tim3_0038 reduce release of the inflammatory cytokine IFN-gamma) more than the F38-2E2 antibody. Tim3_0016, Tim3_0016

variant (Tim3_0018), Tim3_0033 and Tim3_0038 showed a similar release when compared the F38-2E2 antibody. In contrast, cultures containing the isotype control antibody did not show an increase in IFN-gamma secretion.

Table 6a: Percentage of anti-Tim3 antibody induced IFNgamma release in comparison to rec hu IL-2 (20 EU/ml) (= 100%) as positive control and no antibody as negative control

Compound concentration	MLR +IL-2 20U/ml	Iso-type IgG2a	F38-2E2	Tim3 0016	Tim3 0018	Tim3 0021	Tim3 0022	Tim3 0026	Tim3 0028	Tim3 0030	Tim3 0033	Tim3 0038	Iso-type hIgG1
40 µg/ml		2	36	33	36	112	58	25	40	14	35	51	0
10 µg/ml	100	0	26	22	30	108	38	16	38	4	30	38	5
1 µg/ml		0	7	7	12	101	18	18	12	3	0	1	0

In further experiments the EC50 values of the following chimeric and humanized antibodies (generated as described above) in combination with 0,1µg/ml anti-PD1 mAb were measured: chimeric chi_Tim3_018 antibody and its humanized versions Tim3-433 and Tim3-434 , chimeric chi_Tim3_028 antibody and its humanized versions Tim3-438 and Tim3-443 were measured with different lymphocyte donor mixtures (D2 and D3, or D1 and D5, respectively) Results are shown in Table 6b

Table 6b: EC50 of anti-Tim3 antibody induced (IFN-g secretion/release)

Antibody	EC50 [nM] with donors D2+D3	EC50 [nM] with donors D1+D5
chi_Tim3_018	3.1	4.2
Tim3-433	3.0	2.4
Tim3-434	1.7	2.6
chi_Tim3_028	2.9	6.4
Tim3-438	1.9	2.7
Tim3-443	3.0	4.7

Example 6: Internalization of anti-TIM-3 Antibodies into TIM-3 expressing cells

TIM-3-specific antibodies described herein can be internalized into TIM-3-expressing cells, including TIM-3 expressing lymphoma, multiple myeloma and AML cells. For example, the disclosed TIM-3 specific antibodies and fragments thereof are shown to be internalized into rec TIM3 CHO cells stable expressing human TIM-3 as evaluated by cell based ELISA, flow cytometry (FACS) and confocal microscopy.

E.g. stable Tim3-transfected CHO-K1 cells (clone 8) (4×10^4 cells/well/100 μ l) were seeded into 96 well-MTP using fresh culture medium. After overnight cell attachment, cell culture medium was removed and the test antibodies were added to the cells (10 μ g/ml in cell culture medium) and incubated for 0.5 hour at 4°C. As reference, a commercial mouse-anti-human antibody (TIM3 MAB 11E365 (US Biological, T5469-92P) was used. After washing (2x with cell culture medium) and centrifugation cells were incubated for 3 hours at a) 4°C or b) 37°C in 200 μ l cell culture medium. Internalization typically occurs at 37°C, but not at 4°C, which provides another control for the reaction. Then cells were fixated with 100 μ l/well 0.05 % glutaraldehyde (Sigma Cat.No: G5882) in 1xPBS for 10 min at room temperature (RT). This was followed by three washing steps with 200 μ l PBS-T and secondary antibody sheep-anti-mouse-POD (Horseradish POD linked F(ab')₂ Fragment ; GE NA9310)) were added for 1 hour at RT. After the final washing steps (3xPBS-T), TMB substrate was added (Roche order no. 11835033001) for 15 min and color development was stopped using 1N HCl. Final ODs were determined by measurement at 450/620 nm in an ELISA reader. This cellular ELISA procedure was used for medium throughput evaluation of the internalizing capacity of the testing antibodies which were purified from hybridoma supernatants.

The percentage of internalization was calculated as follow:

$$\text{Internalization [\%]} = (1 - \text{OD}_{\text{sample_37}^\circ\text{C}} / \text{OD}_{\text{sample_4}^\circ\text{C}}) * 100$$

The results are shown in FIG. 1A and B for (Internalization). Almost all tested anti-TIM-3 monoclonal antibodies were similar well internalized into stable Tim3-transfected CHO-K1 cells after 3h incubation at 37°C (not all data shown).

The determination of EC50 internalizing values (time dependency) as well as comparison of the kinetics of the internalization depending on mono- vs. bivalency was estimated by FACS for selected candidates.

5 In short human TIM3 stable expressing CHO-K1 cells were seeded (4×10^5 cells/well/50 μ l) into 96 well-v bottom MTP using fresh culture medium and incubated with Redimune® NF Liquid for 10 min at RT to block unspecific binding. Then 50 μ l/well of selected purified anti-TIM3 (10 μ g/ml in cell culture medium) were added and incubated for 1h at 4°C. After washing (with cell culture medium) and centrifugation cells were incubated for 0.25, 0.5, 1, 2, 3, 4, 6 and 24
10 hours at a) 4°C or b) 37°C in 200 μ l cell culture medium. Then cells were washed with PBS/1%BSA and secondary antibody Alexa Fluor 488 Goat-anti-mouseIgG, F(ab)2 were added for 1 hour at 4°C. After washing and centrifugation 125 μ l of CellFix (BD Bioscience, 1:1000) were added and MFI signals of stained cells were analyzed by BD Biosciences FACSCanto flow cytometer.

15 The percentage of internalization was calculated as follow:

$$\text{Internalization [\%]} = (1 - \text{MFI}_{\text{sample_37}^\circ\text{C}} / \text{MFI}_{\text{sample_40}^\circ\text{C}}) * 100$$

**Example for the evaluation of time dependent internalization of anti-TIM3 antibodies Tim3_0016, Tim3_0016 variant (Tim3_0018), Tim3_0021, Tim3_0028, Tim3_0030, Tim3_0033, Tim3_0038 on RPMI-8226 cells (ATCC
20 ® CCL-155™):**

The presently disclosed anti-TIM3 antibodies are internalized rapidly into TIM3 expressing RPMI-8226 cells (ATCC ® CCL-155™) at a high level. The experiments were conducted as described above with TIM3 expressing RPMI-8226 cells (ATCC ® CCL-155™) instead of rec CHOK1 cells expressing huTIM-3.
25 Results are shown in the Table 7. As TIM3 reference antibodies were the following antibodies were used: antibody 8213 as described in US2012/189617, antibody 27.12E12 as described in WO2013/06490. Tim3_0016, Tim3_0016 variant (Tim3_0018), Tim3_0038 were used as human IgG1 chimeric versions.

Table 7: Percentage internalization at the indicated time point (0 min set as 0 percent)

	Percentage internalization of anti-TIM3 antibodies				
Antibody	30 Min	60 Min	120 Min	240 Min	26h
8213	22	22	43	52	72
27.12E12	19	22	25	46	59
Tim3_0016	33	52	55	66	87
Tim3_0018	39	41	80	70	88
Tim3 -0021	70	75	74	78	77
Tim3 -0028	50	59	67	68	83
Tim3 -0033	75	81	82	82	80
Tim3_0038	22	20	45	46	63

From the results antibodies of the invention are rapidly internalized at high percentage compared to reference antibodies on RPMI-8226 cells (ATCC® CCL-155™)

Example 7: Binding of anti-TIM-3 Antibodies to isolated human Monocytes expressing TIM-3

CD14⁺ Monocytes were isolated from anticoagulated peripheral blood of healthy donors by density gradient centrifugation using Ficoll-Paque (GE Healthcare) (see General Protocols in the User Manuals or visit www.miltenyibiotec.com/protocols) and subsequent positive selection via CD14 MicroBeads. First the CD14⁺ cells are magnetically labeled with CD14 MicroBeads. Then the cell suspension is loaded onto a MACS® Column which is placed in the magnetic field of a MACS Separator. The magnetically labeled CD14⁺ cells are retained in the column. The unlabeled cells run through, this cell fraction is depleted of CD14⁺ cells. After removal of the column from the magnetic field, the magnetically retained CD14⁺ cells can be eluted as the positively selected cell fraction. After centrifugation at 200 x g for 10 min at room temperature the monocytes were harvested and used directly in binding assay or resuspended in freezing medium (10% DMSO, 90 %FCS) at 1.0E+07 cells/ml and stored in liquid nitrogen.

As shown in the literature Monocytes express constitutively TIM3 on their surface. 1x10⁵ CD14⁺ isolated human monocytes (50µl/well) were put into 96 well-bottom MTP in fresh culture medium and incubated with Redimune® NF Liquid

- 106 -

for 15 min at RT to block unspecific binding. Than 50 µl/well of disclosed anti-TIM3 mAbs or reference anti-TIM-3 mAbs 344823 (R&D) and F38-2E2 (BioLegend) (10µg/ml in cell culture medium) were added and incubated for 1h at 4°C. Than cells were washed with PBS/1%BSA and secondary antibody PE-labeled Goat-anti-mouse F(ab')₂ were (Jackson Lab 115-006-072) added for 1 hour at 4°C. After washing and centrifugation MFI signals of stained cells were analyzed by BD Biosciences FACSCanto flow cytometer.

The specific binding was calculated as follow:

$$\text{Specific Binding [MFI]} = \text{Geom. Mean MFI}_{\text{sample}} - \text{Geom. Mean MFI}_{\text{isotype control}}$$

The results are shown in Table 8: (Binding to human Monocytes). TIM3 clones Tim3_0016, Tim3_0018, Tim3_0020, Tim3_0028 and Tim3_0038 bind to human monocytes of different donors even better than the reference anti- TIM-3 Abs.

Table 8: Binding to human Monocytes

	donor1 (CD14+)	donor2 (CD14+)	donor3 (CD14+)
Tim3 0016	2122	1634	1690
Tim3 0018	2326	1818	1943
Tim3 0020	1917	1377	1462
Tim3 0021	1134	951	1197
Tim3 0022	1468	1111	1235
Tim3 0026	1665	1016	900
Tim3 0030	1411	419	466
Tim3 0038	1637	1368	1401
Tim3 0028	1351	950	1607
Tim3 0033	480	328	595
M-IgG2b	0	13	0
M-IgG1	144	55	213
<TIM-3>PE Mab, M-IgG1 (Clone F38-2E2; Biolegend)	516	493	460
<TIM-3> PE Mab , Rat IgG2A (Clone 344823, R&D)	1010	917	814
Rat-IgG2A-PE	71	68	70

Example 8: Binding of anti-TIM-3 Antibodies to isolated cyno Monocytes expressing TIM-3

CD14⁺ Monocytes were isolated from cynomolgus monkey anticoagulated peripheral blood (Covance) by density gradient centrifugation using Ficoll-Paque (GE Healthcare) (see General Protocols in the User Manuals or visit
 5 www.miltenyibiotec.com/protocols) and subsequent positiv selection via NHP CD14 MicroBeads. First the CD14⁺ cells are magnetically labeled with CD14 MicroBeads. Then the cell suspension is loaded onto a MACS® Column which is placed in the magnetic field of a MACS Separator. The magnetically labeled
 10 CD14⁺ cells are retained in the column. The unlabeled cells run through, this cell fraction is depleted of CD14⁺ cells. After removal of the column from the magnetic field, the magnetically retained CD14⁺ cells can be eluted as the positively selected cell fraction. After centrifugation at 200 x g for 10 min at room temperature the monocytes were harvested and and used directly in binding assay
 15 or resuspended in freezing medium (10% DMSO, 90 %FCS) at 1.0E+07 cells/ml and stored in liquid nitrogen.

As shown in the literature Monocytes express constitutively TIM3 on their surface. 1x10⁵ CD14⁺ isolated cyno monocytes (50µl/well) were put into 96 well-v bottom MTP in fresh culture medium and incubated with Redimune® NF Liquid for 15
 20 min at RT to block unspecific binding. Than 50 µl/well of Alexa488 labeled anti-TIM3 (10µg/ml in cell culture medium) were added and incubated for 1h at 4oC. After washing and centrifugation MFI signals of stained cells were analyzed by BD Biosciences FACSCanto flow cytometer.

The specific binding was calculated as follow:

25 Specific Binding [MFI] = Geom. Mean MFI_{sample} - Geom. Mean MFI_{isotype control}

The results are shown in Table 9 (Binding to Cyno Monocytes). TIM3 clones Tim3_0016, Tim3_0018, Tim3_0026, Tim3_0028 and , Tim3_0030 bind to cyno monocytes of different cyno donors.

Table 9: Binding to Cyno Monocytes

	cyno1 (16719M) CD14+	cyno2 (17435M) CD14+	cyno3 (30085F) CD14+
AF +PI	75	83	84
HumTIM-3 Alexa488 R&D (34482)	158	121	143
Rat-IgG2A-Alexa488	84	86	91
hum TIM-3 A488 F38-2E2 (NOVUS Biol)	135	136	124
M-IgG1-Alexa 488	72	82	83
Tim3_0016- A488	157	177	187
Tim3_0016 variant 0018- A488	301	480	417
Tim3_0022- A488	115	134	138
Tim3_0026- A488	137	184	197
Tim3_0028- A488	3936	2996	4090
Tim3_0038- A488	97	107	120
Tim3_0020- A488	274	378	354
Tim3_0021 A488	348	473	399
Tim3_0030 A488	119	163	144
Tim3_0033 A488	71	81	83
TIM-3 (4177) A488	78	83	85
TIM-3 (8213) A488	75	83	87

Example 9: Binding of anti-TIM-3 Antibodies to NHL and MM cell lines expressing TIM-3

- 5 The binding capacity of disclosed anti-TIM3 antibodies and two anti-TIM3 reference antibodies clones (1) 4177 and (2) 8213 (Kyowa) was analyzed by a FACS. In short human TIM3 expressing B cell lymphoma cells (exemplified as Pfeiffer cells) and multiple myeloma cells (exemplified as RPMI-8226 cells) were incubated with BD human Fc Block for 10 min at RT to block unspecific binding.
- 10 Than 2x10⁵ cells (50µl/well) were put into 96 well-v bottom MTP and 50 µl/well of Alexa488 labeled anti-TIM3 (10µg/ml in BD Staining buffer) were added and incubated for 1h at 4°C. After washing and centrifugation MFI signals of stained cells were analyzed by BD Biosciences FACSCanto flow cytometer.

The specific binding was calculated as follow:

- 15 Specific Binding [MFI] = Geom. Mean MFI_{sample} - Geom. Mean MFI_{isotype control}

The results are shown in FIG. 2A and 2B (Binding to RPMI-8226 and Pfeiffer cells).

Example 10: Cytotoxic activity of anti-TIM-3 Antibodies on TIM-3 expressing NHL and MM cells

5 TIM3-specific antibodies conjugated with pseudomonas exotoxin (PE 24) effectively kill TIM3-expressing cells.

The cytotoxic activity of disclosed anti-TIM3 antibodies and one commercial available anti-TIM3 reference antibody clone 11E365 (available from US Biological) was analyzed with Promega CellTiter-Glo Luminescent Cell Viability Assay. In short to 5×10^3 (50 μ l/well in 98 well MTP, in triplicate) recombinant CHO K1 stable expressing human TIM-3 or 2×10^4 cells (50 μ l/well in 98 well MTP, in triplicate) human TIM3 expressing B cell lymphoma cells (exemplified as Pfeiffer cells) or multiple myeloma cells (exemplified as RPMI-8226 cells) were added 25 μ l/well 1:5 serial dilution of disclosed anti-TIM-3 antibodies with the highest concentration of 10 μ g/ml or appropriate media to untreated cells or Isotype control to untargeted treated cells. Treatment ranges from 10 μ g/ml to 1ng/ml in triplicate. All antibodies were used as full length mouse Fc γ versions . For conjugation of the conjugation of the Pseudomonas exotoxin 10 μ g/ml of mouse Fc γ fragment specific Fabs conjugated with PE 24 were added and incubated for 3 days at 37°C. Cycloheximide as a known protein synthesis inhibitor in eukaryotes was used as positive control. Viability of treated cells were measured with Promega CellTiter-Glo Luminescent Cell Viability Assay.

The cytotoxic activity was calculated as follow:

$$25 \quad \text{Rel. Inhibition [\%]} = (1 - (E_{\text{sample}} - E_{\text{negative control}}) / (E_{\text{positive control}} - E_{\text{negative control}})) * 100$$

The results are shown in Tab. 10 (Cytotoxic activity of anti-TIM3 mAbs on TIM-3 expressing recombinant, NHL and MM cell lines in sandwich format).

Table 10:

Antibodies and references (all anti TIM3 antibodies conjugated to a deimmunized Pseudomonas exotoxin A)	IC50 [nM]		
	recTIM-3 CHO cells	Pfeiffer cells	RPMI-8226
Tim3_0016	0,04	0,09	0,55
Tim3_0016 variant 1 (Tim3_0018)	0,05	0,10	0,66
Tim3_0020	0,07	0,11	>64
Tim3_0021	0,04	0,10	5,9
Tim3_0022	0,02	0,07	0,36
Tim3_0023	0,03	0,08	>64
Tim3_0026	0,03	0,08	>64
Tim3_0030	0,03	0,10	>64
Tim3_0033	0,11	0,20	0,79
Tim3_0038	0,01	< 0.002	0,16
US Biol. Clone 11E365	0,7	1,2	1,1
Cells w/o Ab	-	-	-
Cells + <mFc> Fab PE	-	-	-
IgG2A + <mFc> Fab PE	-	-	-
Cycloheximide	135	181	245

All tested TIM3 clones are highly potent (IC50 range 0,01-0,2 nM) on recombinant CHO K1 stable expressing human TIM-3 and Pfeiffer cells expressing high and moderate levels of TIM-3 and even more potent in their cytotoxic activity than the strong internalizing reference anti-TIM-3 Ab clone 11E365, US Biological. TIM3 clones 0016, 0018, 0021, 0022, 0033 and 0038 are also potent on RPMI-8226 cells expressing 5 fold lower TIM-3 level compare to recombinant CHO TIM-3 cells.

Example 11: Comparison of the cytotoxic activity of disclosed anti-TIM3 antibodies vs. two anti-TIM3 reference antibodies 1.7.E10 and 27-12E12 (as described in WO2013/06490).

The cytotoxic activity of disclosed anti-TIM3 antibodies and two anti-TIM3 reference antibodies the TIM3 reference antibodies 1.7E10 and 27.12E12 as described in WO2013/06490 was analyzed with Promega CellTiter-Glo Luminescent Cell Viability Assay as described above. All antibodies were used as full length human IgG1 format including the human Fcγ part. In this experiment conjugation of the Pseudomonas exotoxin was achieved via human Fcγ

fragment specific Fabs conjugated with PE 24 (10µg/ml) which were added and incubated for 5 days at 37°C.

The results are shown in Tab. 11. - Comparison of cytotoxic activity of anti-TIM3 mAbs on TIM-3 expressing NHL and MM cell lines

Table 11: Comparison of cytotoxic activity of anti-TIM3 mAbs on TIM-3 expressing NHL and MM cell lines

Antibodies and references (all anti TIM3 antibodies conjugated to a deimmunized Pseudomonas exotoxin A)	Pfeiffer cells		RPMI-8226 cells	
	Max. killing	Rel. IC50 [nM]	Max. killing	Rel. IC50 [nM]
Cycloheximide	100 [%]	271	100 [%]	111
1.7E10	60.3 [%]	0,68	65.7 [%]	2,544
27-12E12	75.7 [%]	0,02	86.6 [%]	0,111
Tim3_0016	84.9 [%]	0,05	86.6 [%]	0,063
Tim3_0016 variant (Tim3_0018)	82.9 [%]	0,06	88.1 [%]	0,081
Tim3_0026	78.3 [%]	<0.02	83.1 [%]	0,067
Tim3_0038	82.6 [%]	<0.02	83.8 [%]	0,047
Isotype Control hIgG1	3.2 [%]	N.A	0.4 [%]	N.A

All disclosed TIM3 clones are highly active (IC50 range 0,02-0,08 nM) on Pfeiffer and RPMI-8226 cells expressing TIM-3 and even more potent in their cytotoxic activity than the strong internalizing reference anti-TIM-3 Ab clone 27-12E12. All antibodies were compared as Pseudomonas exotoxin (PE24) conjugates using the same Pseudomonas exotoxin under the same conditions

Example 12: Cytotoxic activity of Fab-PE24 constructs of disclosed anti-TIM3 antibodies on MM, NHL and AML cell lines (expressing TIM3, but not PSMA).

The cytotoxic activity was analyzed with Promega CellTiter-Glo Luminescent Cell Viability Assay as described above. 1:5 serial dilutions of Fab-fragments of disclosed anti-TIM3 antibodies directly conjugated to PE24 with the highest concentration of 50µg/ml or appropriate media to untreated cells or non-binding anti-PSMA Fab-PE24 control to untargeted treated cells were incubated with 7,5 x10³ Pfeiffer cells or 2 x10³ RPMI-8226 cells (50µl/well in 98 well MTP) for 4 days at 37°C. Treatment ranges from 50µg/ml to 8ng/ml in triplicate. Cycloheximide was used as positive control.

The results are shown in Tab. 12. (Cytotoxic activity of Fab-PE24 constructs of disclosed anti-TIM3 antibodies on MM, NHL and AML cell lines).

Table 12: Cytotoxic activity of Fab-PE24 constructs of disclosed anti-TIM3 antibodies on different MM, NHL and AML cell lines (RPMI-8226, Karpas-299, CMK, TF-1, MOLM-13)

	RPMI-8226		Karpas-299		CMK		TF-1		MOLM-13	
Antibodies and references (all anti TIM3 antibodies conjugated to a deimmunized Pseudomonas exotoxin A)	Max. kil-ling	IC50 [nM]	Max. kil-ling	IC50 [nM]	Max. kil-ling	IC50 [nM]	Max. kil-ling	IC50 [nM]	Max. kil-ling	IC50 [nM]
Cycloheximide	100 [%]	281	100 [%]	113	100 [%]	149,0	100 [%]	207	100 [%]	156
Anti_P SMA	10.5 [%]	N.A.	40.1 [%]	N.A.	8.98 [%]	N.A.	5.27 [%]	N.A.	18.9 [%]	N.A.
Tim3_0022	99.1 [%]	1,9	98.8 [%]	10	67.1 [%]	255	58.6 [%]	299	58.5 [%]	579
Tim3_0016	99.3 [%]	1,1	99.2 [%]	4	64.8 [%]	225	54.2 [%]	534	62.7 [%]	459

All tested Fab-PE24 constructs of disclosed anti-TIM3 antibodies are highly potent (IC50 range 1-10 nM) on MM (RPMI-8226) and NHL (Karpas-299) cells expressing moderate level of TIM-3 and demonstrate significant cytotoxic activity on AML cell lines (CMK, TF-1, MOLM-13) expressing very low levels of TIM-3.

Example 13: Cytotoxic activity of Immuno conjugates (Pseudomonas Exotoxin A conjugates (Fab-PE24 constructs) of disclosed anti-TIM3 on primary leukemic stem/progenitor AML cells from relapsed/refractory patients

CD34+ cells from peripheral blood of relapsed/refractory patients were obtained from AllCells, LLC, Alameda, CA.

Table 13: Clinical characteristics of AML patients

Specimens	Donor	Clinical Diagnosis	Gender	Age: y	FAB subtype	WBC [x10 ⁹ /L]	CD 34+ cells [%]	PLT [x10 ⁹ /L]	Cytogen. abnormal.
AML1	EBO PB0136	AML Relaps./Refract.	F	64	M2	4	50		N.A.
AML2	EBO PB0142	AML Relaps./Refract.	F	35	N.A.	65	22		Normal
AML3	EBO PB0135	AML Relaps./Refract.	M	72	M0	4	15		N.A.
AML4a	EBO PB0193	AML Relaps./Refract.	M	76	N.A.	21	52		der(7)t(7;? 13)

After confirmation of purity and viability of all samples (purity range 84-94% and viability range 95-99%) the expression level of TIM-3 was evaluated by FACS as described in Example 7 using anti-TIM-3 mAbs 344823 (R&D). (see Figure 3)

All tested (4/4) primary leukemic stem/progenitor (CD34+) AML samples from relapsed/refractory patients demonstrate homogeneous expression of TIM-3 at different levels.

For the evaluation of cytotoxic activity of Fab-PE24 constructs of disclosed anti-TIM3 clones 0016 and 0022 on primary CD34+ AML cells 1 x10⁴ cells (50µl/well in 96 well MTP, in triplicate) were incubated with 1:5 serial dilutions of Fab-fragments with the highest concentration of 50µg/ml or appropriate media to untreated cells or non-binding anti-PSMA Fab-PE24 control to untargeted treated cells for 3 days at 37°C. Cycloheximide was used as positive control. Cytotoxic activity was analysed with Promega CellTiter-Glo Luminescent Cell Viability Assay as described above in Example 12.

The results are shown in Tab.14. (Cytotoxic activity of Fab-PE24 constructs of disclosed anti-TIM3 antibodies on primary CD34+ AML cells).

Table 14: Cytotoxic activity of Fab-PE24 constructs of disclosed anti-TIM3 antibodies on primary CD34+ AML cells

	D1; AML CD34+ PB0136 cells		D2; AML CD34+ PB0142 cells		D3; AML CD34+ PB0135 cells		D4; AML CD34+ PB0193 cells	
Antibodies and references (all anti TIM3 antibodies conjugated to a deimmunized Pseudomonas exotoxin A)	Max. killing	IC50 [nM]	Max. killing	IC50 [nM]	Max. killing	IC50 [nM]	Max. killing	IC50 [nM]
Cycloheximide	100 [%]	212	100 [%]	262	100 [%]	121	100 [%]	208
anti-PSMA	2 [%]	N.A.	8 [%]	N.A.	18 [%]	N.A.	12 [%]	N.A.
TIM-3 0022- cFP	38 [%]	>691	75 [%]	107	31 [%]	>691	57 [%]	375
TIM-3 0016- cFP	48 [%]	>691	79 [%]	30	44 [%]	>691	69 [%]	116

Fab-PE24 constructs of anti-TIM3 antibodies Tim3_0016 and Tim3_0022 are highly potent on (2/4) primary AML samples (PB0142 and PB0135) (IC50 range 30-116 nM) and demonstrate significant cytotoxic activity on all (4/4) primary leukemic stem/progenitor (CD34+) AML cells expressing different levels of TIM-3.

Example 14: Comparison of potency of Fab-PE24 constructs of selected anti-TIM3 antibodies on NHL and MM cell lines

The evaluation of cytotoxic activity of sortase coupled Fab-PE24 constructs of selected disclosed anti-TIM3 antibodies was analysed with Promega CellTiter-Glo Luminescent Cell Viability Assay as described above in Example 12.

The results are shown in Tab. 15. (Cytotoxic activity of Fab-PE24 constructs of selected anti-TIM3 antibodies on NHL and MM cells).

Table 15: Cytotoxic activity of Fab-PE24 constructs of selected anti-TIM3 antibodies on NHL and MM cells

Antibodies and references (all anti TIM3 antibodies conjugated to a deimmunized Pseudomonas exotoxin A)	Pfeiffer cells		RPMI-8226 cells	
	Max. killing	IC50 [nM]	Max. killing	IC50 [nM]
Cycloheximide	100 [%]	271,1	100 [%]	153
anti-PSMA	25.2 [%]	N.A.	21.5 [%]	N.A.
TIM-3 0022	99.9 [%]	1,58	99.6 [%]	2,14
TIM-3 0016	99.6 [%]	0,77	99.2 [%]	0,61
TIM-3 0021	98.4 [%]	2,15	99.1 [%]	3,61
TIM-3 0033	99.8 [%]	5,30	99.7 [%]	5,73
TIM-3 0038	99.6 [%]	0,47	98.3 [%]	0,32

High cytotoxic potency was demonstrated with Fab-PE24 constructs of all selected disclosed anti-TIM3 antibodies (IC50 range 0,3-5 nM) on NHL (Pfeiffer) and MM (RPMI-8226) cells expressing moderate level of TIM-3.

The highest cytotoxic activity was observed with Fab-PE24 constructs of disclosed anti-TIM3 antibodies Tim3_0016 and Tim3_0038.

Example 15: Comparison of cytotoxic activity of Fab-PE24 construct vs. total-IgG-Amatoxin conjugate of the same clone of disclosed anti-TIM-3 antibody on Pfeiffer cells

The evaluation of cytotoxic activity of conjugated Fab-PE24 construct of disclosed anti-TIM3 clone 0016 vs. total IgG of the same clone conjugated with Amatoxin (according to the procedures described in WO2012/041504 (conjugated via the 6' C-atom of amatoxin amino acid 4, particularly via an oxygen atom bound to the 6' C-atom of amatoxin amino acid, and wherein the TIM3 antibody is connected by a linker via a urea moiety) was analysed with Promega CellTiter-Glo Luminescent Cell Viability Assay as described above in Example 12.

The results are shown in Tab. 16. (Cytotoxic activity of Fab-PE24 construct vs. total IgG- Amatoxin conjugate of anti-TIM3 clone 0016 on NHL cells).

Table 16: Cytotoxic activity of Fab-PE24 construct vs. total IgG- Amatoxin conjugate of anti-TIM3 clone 0016 on NHL cells

Pfeiffer cells	Max. killing	IC50 [nM]
Cycloheximide	100 [%]	163
Isotype hIgG1 Amatoxin	28 [%]	N.A.
TIM-3 0016- Amatoxin	93.3 [%]	0,81
TIM-3 0016-PE24	99.8 [%]	0,25

5 Cytotoxic activity of Amanitin-conjugated anti-TIM-3 clone 0016 (IC50 0,8 nM) is comparable with cytotoxic activity of Fab-PE24 construct of the same clone (IC50 0,3 nM) on NHL (Pfeiffer) cells expressing moderate level of TIM-3.

Example 15: Direct comparison of binding of TIM3 antibodies to different peripheral blood mononuclear cells (PBMC)

Binding assay

10 Freshly isolated PBMCs or 3 days polyclonally activated (plate bound anti-CD3 and soluble anti-CD28 antibodies, 1 ug/ml each, both from BD Pharmingen) CD4 T cells were stained with Alexa 647-directly conjugated anti-TIM- antibodies Tim3-0018, Tim3-0028 or chimerized or humanized versions thereof for 1 hour at 4 C degrees. The cells were then washed to eliminate unbound antibody and
15 stained for surface markers for 30 minutes at 4 C degrees to discriminate monocytes (CD14⁺ (BD Pharmingen)), NK cells (CD16⁺ (eBioscience), CD56⁺ (BioLegend) and CD3⁻) and T cells (CD3⁺ (eBioscience)) before being fixed with BD Cell Fix. The cells were acquired at LSRFortessa, BD Biosciences.

20 Results are shown in Figures 4A to 4 D (in the Figures the following designations were used: for Tim3-0018: 0018 (aTIM-3), for humanized Tim3-0018 version Tim3-0434: 0434(h0018), for Tim3-0028: 0028 (aTIM-3), for chimeric Tim3-0028: chi0028, for humanized Tim3-0028 version Tim3-0438: 0438(h0028)). The data show that the humanized antibodies have improved binding and binding specificity to CD4 Tcells when compared to the parental antibodies.

25

Patent Claims

1. An isolated antibody that binds to human TIM3, wherein the antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:37, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:38, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:39; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:40; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:41 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:42.
2. The antibody according to claim 1, wherein the antibody comprises
 - i) a VH sequence of SEQ ID NO:83 and a VL sequence of SEQ ID NO:84, or
 - ii) a VH sequence of SEQ ID NO:85 and a VL sequence of SEQ ID NO:86.
3. The antibody according to claim 1, wherein the antibody comprises a VH sequence of SEQ ID NO:83 and a VL sequence of SEQ ID NO:84.
4. The antibody according to claim 1, wherein the antibody comprises a VH sequence of SEQ ID NO:85 and a VL sequence of SEQ ID NO:86.
5. The antibody according to any one of the preceding claims, which is a full length IgG1 antibody.

6. The antibody of according to any one of the preceding claims, which is a full length IgG1 antibody with mutations L234A, L235A and P329G (numbering according to the EU index of Kabat)
- 5 7. Isolated nucleic acid encoding the antibody according to any one of the preceding claims.
8. A host cell comprising the nucleic acid of claim 7.
9. A method of producing an antibody comprising culturing the host cell of claim 8 so that the antibody is produced.
- 10 10. The method of claim 9, further comprising recovering the antibody from the host cell.
11. An immunoconjugate comprising the antibody according to any one of claims 1 to 6 and a cytotoxic agent.
12. A pharmaceutical formulation comprising the antibody according to any one of claims 1 to 6 or the immunoconjugate of claim 11 and a pharmaceutically acceptable carrier.
- 15 13. The antibody according to any one of claims 1 to 6 or the immunoconjugate of claim 11 for use as a medicament.
14. The antibody according to any one of claims 1 to 6 or the immunoconjugate of claim 11 for use in treating cancer.
- 20 15. Use of the antibody according to any one of claims 1 to 6 or the immunoconjugate of claim 11 in the manufacture of a medicament for the treatment of cancer.