

ANTI TUBERCULAR DRUG: COMPOSITIONS AND METHODS

5 FIELD OF INVENTION

The present invention relates to methods and compositions for treating disease caused by microorganisms, particularly tuberculosis. The present invention also relates to methods and compositions having improved anti-mycobacterial activity, namely compositions comprising novel substituted ethylene diamine compounds.

10

BACKGROUND OF THE INVENTION

Mycobacterial infections often manifest as diseases such as tuberculosis. Human infections caused by mycobacteria have been widespread since ancient times, and tuberculosis remains a leading cause of death today. Although the incidence of the disease declined, in parallel with advancing standards of living, since the mid-nineteenth century, mycobacterial diseases still constitute a leading cause of morbidity and mortality in countries with limited medical resources. Additionally, mycobacterial diseases can cause overwhelming, disseminated disease in immunocompromised patients. In spite of the efforts of numerous health organizations worldwide, the eradication of mycobacterial diseases has never been achieved, nor is eradication imminent. Nearly one third of the world's population is infected with *mycobacterium tuberculosis* complex, commonly referred to as tuberculosis (TB), with approximately 8 million new cases, and two to three million deaths attributable to TB yearly. Tuberculosis (TB) is the cause of the largest number of human deaths attributable to a single etiologic agent (see Dye et al., J Am. Med. Association, 282, 677-686, (1999); and 2000 WHO/OMS Press Release).

After decades of decline, TB is now on the rise. In the United States, up to 10 million individuals are believed to be infected. Almost 28,000 new cases were reported in 1990, constituting a 9.4 percent increase over 1989. A sixteen percent increase in TB cases was observed from 1985 to 1990. Overcrowded living conditions and shared air spaces are especially conducive to the spread of TB, contributing to the increase in instances that have been observed among prison inmates, and among the homeless in larger U.S. cities.

30

Approximately half of all patients with "Acquired Immune Deficiency Syndrome" (AIDS) will acquire a mycobacterial infection, with TB being an especially devastating complication. AIDS patients are at higher risks of developing clinical TB, and anti-TB treatment seems to be less effective than in non-AIDS patients. Consequently, the infection often progresses to a fatal disseminated disease.

Mycobacteria other than *M. tuberculosis* are increasingly found in opportunistic infections that plague the AIDS patient. Organisms from the *M. avium-intracellulare* complex (MAC), especially serotypes four and eight, account for 68% of the mycobacterial isolates from AIDS patients. Enormous numbers of MAC are found (up to 10^{10} acid-fast bacilli per gram of tissue), and consequently, the prognosis for the infected AIDS patient is poor.

The World Health Organization (WHO) continues to encourage the battle against TB, recommending prevention initiatives such as the "Expanded Program on Immunization" (EPI), and therapeutic compliance initiatives such as "Directly Observed Treatment Short-Course" (DOTS). For the eradication of TB, diagnosis, treatment, and prevention are equally important. Rapid detection of active TB patients will lead to early treatment by which about 90% cure is expected. Therefore, early diagnosis is critical for the battle against TB. In addition, therapeutic compliance will ensure not only elimination of infection, but also reduction in the emergence of drug-resistance strains.

The emergence of drug-resistant *M. tuberculosis* is an extremely disturbing phenomenon. The rate of new TB cases proven resistant to at least one standard drug increased from 10 percent in the early 1980's to 23 percent in 1991. Compliance with therapeutic regimens, therefore, is also a crucial component in efforts to eliminate TB and prevent the emergence of drug resistant strains. Equally important is the development of new therapeutic agents that are effective as vaccines, and as treatments, for disease caused by drug resistant strains of mycobacteria.

Although over 37 species of mycobacteria have been identified, more than 95% of all human infections are caused by six species of mycobacteria: *M. tuberculosis*, *M. avium intracellulare*, *M. kansasii*, *M. fortuitum*, *M. chelonae*, and *M. leprae*. The most prevalent mycobacterial disease in humans is tuberculosis (TB) which is predominantly caused by mycobacterial species comprising *M. tuberculosis*, *M. bovis*, or *M. africanum* (Merck Manual 1992). Infection is typically initiated by the inhalation of infectious particles

which are able to reach the terminal pathways in lungs. Following engulfment by alveolar macrophages, the bacilli are able to replicate freely, with eventual destruction of the phagocytic cells. A cascade effect ensues wherein destruction of the phagocytic cells causes additional macrophages and lymphocytes to migrate to the site of infection, where
5 they too are ultimately eliminated. The disease is further disseminated during the initial stages by the infected macrophages which travel to local lymph nodes, as well as into the blood stream and other tissues such as the bone marrow, spleen, kidneys, bone and central nervous system. (See Murray et al. *Medical Microbiology*, The C.V. Mosby Company 219-230 (1990)).

10 There is still no clear understanding of the factors which contribute to the virulence of mycobacteria. Many investigators have implicated lipids of the cell wall and bacterial surface as contributors to colony morphology and virulence. Evidence suggests that C-mycosides, on the surface of certain mycobacterial cells, are important in facilitating survival of the organism within macrophages. Trehalose 6,6' dimycolate, a cord factor, has
15 been implicated for other mycobacteria.

The interrelationship of colony morphology and virulence is particularly pronounced in *M. avium*. *M. avium* bacilli occur in several distinct colony forms. Bacilli which grow as transparent, or rough, colonies on conventional laboratory media are multiplicable within macrophages in tissue culture, are virulent when injected into
20 susceptible mice, and are resistant to antibiotics. Rough or transparent bacilli, which are maintained on laboratory culture media, often spontaneously assume an opaque R colony morphology, at which time they are not multiplicable in macrophages, are avirulent in mice, and are highly susceptible to antibiotics. The differences in colony morphology between the transparent, rough and opaque strains of *M. avium* are almost certainly due to
25 the presence of a glycolipid coating on the surface of transparent and rough organisms which acts as a protective capsule. This capsule, or coating, is composed primarily of C-mycosides which apparently shield the virulent *M. avium* organisms from lysosomal enzymes and antibiotics. By contrast, the non-virulent opaque forms of *M. avium* have very little C-mycoside on their surface. Both the resistance to antibiotics and the resistance
30 to killing by macrophages have been attributed to the glycolipid barrier on the surface of *M. avium*.

Diagnosis of mycobacterial infection is confirmed by the isolation and identification of the pathogen, although conventional diagnosis is based on sputum smears, chest X-ray examination (CXR), and clinical symptoms. Isolation of mycobacteria on a medium takes as long as four to eight weeks. Species identification takes a further two weeks. There are several other techniques for detecting mycobacteria such as the polymerase chain reaction (PCR), mycobacterium tuberculosis direct test, or amplified mycobacterium tuberculosis direct test (MTD), and detection assays that utilize radioactive labels.

One diagnostic test that is widely used for detecting infections caused by *M. tuberculosis* is the tuberculin skin test. Although numerous versions of the skin test are available, typically one of two preparations of tuberculin antigens are used: old tuberculin (OT), or purified protein derivative (PPD). The antigen preparation is either injected into the skin intradermally, or is topically applied and is then invasively transported into the skin with the use of a multiprong inoculator (Tine test). Several problems exist with the skin test diagnosis method. For example, the Tine test is not generally recommended because the amount of antigen injected into the intradermal layer cannot be accurately controlled. (See Murray et al. *Medical Microbiology*, The C.V. Mosby Company 219-230 (1990)).


Although the tuberculin skin tests are widely used, they typically require two to three days to generate results, and many times, the results are inaccurate since false positives are sometimes seen in subjects who have been exposed to mycobacteria, but are healthy. In addition, instances of mis-diagnosis are frequent since a positive result is observed not only in active TB patients, but also in persons vaccinated with Bacille Calmette-Guerin (BCG), and those who had been infected with mycobacteria, but have not developed the disease. It is hard therefore, to distinguish active TB patients from the others, such as household TB contacts, by the tuberculin skin test. Additionally, the tuberculin test often produces a cross-reaction in those individuals who were infected with mycobacteria other than *M. tuberculosis* (MOTT). Therefore, diagnosis using the skin tests currently available is frequently subject to error and inaccuracies.

The standard treatment for tuberculosis caused by drug-sensitive organisms is a six-month regimen consisting of four drugs given for two months, followed by two drugs given for four months. The two most important drugs, given throughout the six-month course of

therapy, are isoniazid and rifampin. Although the regimen is relatively simple, its administration is quite complicated. Daily ingestion of eight or nine pills is often required during the first phase of therapy; a daunting and confusing prospect. Even severely ill patients are often symptom free within a few weeks, and nearly all appear to be cured within a few months. If the treatment is not continued to completion, however, the patient may experience a relapse, and the relapse rate for patients who do not continue treatment to completion is high. A variety of forms of patient-centered care are used to promote adherence with therapy. The most effective way of ensuring that patients are taking their medication is to use directly observed therapy, which involves having a member of the health care team observe the patient take each dose of each drug. Directly observed therapy can be provided in the clinic, the patient's residence, or any mutually agreed upon site. Nearly all patients who have tuberculosis caused by drug-sensitive organisms, and who complete therapy will be cured, and the risk of relapse is very low ("Ending Neglect: The Elimination of Tuberculosis in the United States" ed. L. Geiter Committee on the Elimination of Tuberculosis in the United States Division of Health Promotion and Disease Prevention, Institute of Medicine. Unpublished.)

What is needed are effective therapeutic regimens that include improved vaccination and treatment protocols. Currently available therapeutics are no longer consistently effective as a result of the problems with treatment compliance, and these compliance problems contribute to the development of drug resistant mycobacterial strains.

Ethambutol (EMB) is a widely used antibiotic for the treatment of TB, with over 300 million doses delivered for tuberculosis therapy in 1988.




Ethambutol, developed by Lederle Laboratories in the 1950s, has low toxicity and is a good pharmacokinetic. However, ethambutol has a relatively high Minimum Inhibition Concentration (MIC) of about 5 µg/ml, and can cause optic neuritis. Thus, there is an increasing need for new, and more effective, therapeutic compositions (See for example, 5 U.S. Pat. No. 3,176,040, U.S. Pat. No. 4,262,122; U.S. Pat. No. 4,006,234; U.S. Pat. No. 3,931,157; U.S. Pat. No. 3,931,152; U.S. Re. 29,358; and Häusler *et al.*, Bioorganic & Medicinal Chemistry Letters 11 (2001) 1679-1681). In the decoder years since the discovery of the beneficial effects of ethambutol, few pharmacological advances in TB treatment have been developed. Moreover, with the combined emergence of drug resistant 10 strains, and the more prevalent spread of mycobacterial disease, it is becoming seriously apparent that new therapeutic compositions are crucial in the fight against tuberculosis.

Clearly effective therapeutic regimens that include improved vaccination and treatment protocols are needed. A therapeutic vaccine that would prevent the onset of tuberculosis, and therefore eliminate the need for therapy is desirable. Although currently 15 available therapeutics such as ethambutol are effective, the emergence of drug resistant strains has necessitated new formulations and compositions that are more versatile than ethambutol. Currently available therapeutics are no longer consistently effective as a result of the problems with treatment compliance, leading to the development of drug resistant mycobacterial strains. What is needed are new anti-tubercular drugs that provide highly 20 effective treatment, and shorten or simplify tuberculosis chemotherapy.

SUMMARY OF THE INVENTION

The present invention comprises methods and compositions comprising ethylene diamine compounds effective for the treatment of infectious disease. The present invention 25 also provides methods and compositions comprising substituted ethylene diamines having improved anti-mycobacterial activity, including substituted ethylene diamines having improved anti-tuberculosis activity.

The present invention contemplates substituted ethylene diamines, which can derive from a variety of amine compounds. In the present invention, the substituted ethylene 30 diamines are based on the following structure.



Substituted Ethylene Diamine

The substituted ethylene diamine compounds described herein are synthesized and screened for activity as follows. A chemical library of substituted ethylene diamines is prepared on a solid polystyrene support using split and pool technologies. This technique
5 allows for the synthesis of a diverse set of substituted ethylene diamines. These diamines are screened for anti-TB activity using *in vitro*, biological assays, including a High-Throughput Screening (HTS) assay, based on the recently completed genomic sequence of *M. tuberculosis*, and a Minimum Inhibition Concentration (MIC) assay.

The methods and compositions described herein comprise substituted ethylene
10 diamines that are effective against disease caused by infectious organisms, including, but not limited to, bacteria and viruses. One embodiment of the invention provides methods and compositions comprising substituted ethylene diamines that are effective against mycobacterial disease. Another embodiment of the invention provides methods and compositions comprising substituted ethylene diamines that have MIC of 50μM or lower
15 for mycobacterial disease. Another embodiment of the present invention comprises substituted ethylene diamines that have an MIC of 25 μM or lower for mycobacterial disease. Yet another embodiment of the present invention comprises substituted ethylene diamines that have an MIC of 12.5 μM or lower for mycobacterial disease. Another embodiment of the present invention comprises substituted ethylene diamines that have an
20 MIC of 5 μM or lower for mycobacterial disease. In another embodiment of the present invention, the methods and compositions comprise substituted ethylene diamines with HTS Luc activity of 10% or greater. In yet another embodiment of the present invention, the methods and compositions comprise substituted ethylene diamines, wherein one amine group is derived from a primary amine, and wherein the other amine group is derived from
25 a primary or secondary amine. In another embodiment of the present invention, the methods and compositions comprise substituted ethylene diamines, wherein one amine is derived from cis-(-)myrtaniline, cyclooctylamine, 2,2-diphenylethylamine, 3,3-diphenylpropylamine, (+)-bornylamine, 1-adamantanemethylamine, (+)-isopinocampheylamine; or (-)-isopinocampheylamine.

The present invention contemplates various salt complexes and other substituted derivatives of the substituted ethylene diamines. The present invention also contemplates enantiomers and other stereoisomers of the substituted ethylene diamines and their substituted derivatives. The present invention further contemplates treatment for animals,
5 including, but not limited to, humans.

Accordingly, it is an object of the present invention to provide methods and compositions for the treatment and prevention of diseases caused by microorganisms.

Accordingly, it is an object of the present invention to provide methods and compositions for the treatment and prevention of infectious diseases.

10 Another object of the present invention is to provide methods and compositions for the treatment and prevention of mycobacterial disease, including but not limited to, tuberculosis.

Yet another object of the present invention is to provide methods and compositions for the treatment and prevention of infectious diseases using compositions comprising
15 substituted ethylene diamines.

Another object of the present invention is to provide methods and compositions for the treatment and prevention of mycobacterial disease using compositions comprising substituted ethylene diamines.

Still another object of the present invention is to provide methods and compositions
20 for the treatment and prevention of tuberculosis using compositions comprising substituted ethylene diamines.

Another object of the present invention is to provide methods and compositions for the treatment and prevention of tuberculosis using compositions comprising substituted ethylene diamines, wherein the diamine has an MIC of 50 μ M, or less.

25 Another object of the present invention is to provide methods and compositions for the treatment and prevention of tuberculosis using compositions comprising substituted ethylene diamines, wherein the diamine has an MIC of 25 μ M, or less.

Another object of the present invention is to provide methods and compositions for the treatment and prevention of tuberculosis using compositions comprising substituted
30 ethylene diamines, wherein the diamine has an MIC of 12.5 μ M, or less.

Yet another object of the present invention is to provide methods and compositions for the treatment and prevention of tuberculosis using compositions comprising substituted ethylene diamines, wherein the diamine has an MIC of 5 μ M, or less.

Yet another object of the present invention is to provide methods and compositions
5 for the treatment and prevention of tuberculosis using compositions comprising substituted ethylene diamines, wherein the diamine has HTS/Luc activity of 10% or greater.

Another object of the present invention is to provide methods and compositions for the treatment and prevention of tuberculosis using compositions comprising substituted ethylene diamines, wherein one amine group is derived from a primary amine, and the
10 other amine group is derived from a primary or secondary amine.

Yet another object of the present invention is to provide methods and compositions for the treatment and/or prevention of tuberculosis using compositions comprising substituted ethylene diamines, wherein one amine is derived from cis-(-)myrtanamine, cyclooctylamine, 2,2-diphenylethylamine, 3,3-diphenylpropylamine, (+)-bornylamine, 1-
15 adamantanemethylamine, (+)-isopinocampheylamine; or (-)-isopinocampheylamine.

Yet another object of the present invention is to provide composition for the therapeutic formulation for the treatment and prevention of mycobacterial disease.

Another object of the present invention is to provide compositions for therapeutic formulations for the treatment and prevention of mycobacterial disease caused by
20 mycobacterial species comprising *M. tuberculosis* complex, *M. avium intracellulare*, *M. kansarii*, *M. fortuitum*, *M. chelonoe*, *M. leprae*, *M. africanum*, *M. microti*, or *M. bovis*.

These and other objects, features and advantages of the present invention will become apparent after a review of the following detailed description of the disclosed embodiments and the appended claims.

25

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 represents a flow chart schematic showing various solid support syntheses used to prepare substituted ethylene diamines.

Figures 2(a)-2(ac) provide chemical structures of a variety of primary amines.

Figures 3(a)-3(f) provide chemical structures of a variety of acyclic secondary amines.

Figures 4(a)-4(i) provide chemical structures of a variety of cyclic secondary amines.

5 Figure 5 represents a flow schematic for a representative reaction pool of ten substituted ethylene diamines.

Figure 6 is a graph of Luminescence Count per Second (LCPS) versus concentration showing HTS Luc assay results for pooled substituted ethylene diamine compounds.

10 Figure 7 is a graph of LCPS versus concentration showing HTS Luc assay results for individual substituted ethylene diamine compounds.

Figure 8 is a graph of LCPS versus concentration showing HTS Luc assay results for individual substituted ethylene diamine compounds.

15 Figure 9 is a bar graph providing a summary of MIC activities for discrete substituted ethylene diamines.

Figure 10 is a bar graph providing a summary of Luciferase activity of discrete substituted ethylene diamines with at least 10% activity in reference to ethambutol at 3.1 μ M.

20 Figure 11 is a bar graph showing the frequency of occurrences of the selected amine monomers in the substituted ethylene diamine compounds that were active against TB. Amine monomers are represented by their numerical designations.

Figure 12 represents a flow schematic showing a synthesis of *N*-Geranyl-*N'*-(2-adamanthyl)ethane-1,2-diamine (compound 109).

25 Figure 13 is a flow schematic showing a synthesis of *N*-(Cyclooctyl)-*N'*-(1*R*, 2*R*, 3*R*, 5*S*)-(-)-isopinocampheylethane-1,2-diamine as hydrochloride (compound 59).

Figure 14 is a mass spec profile for one representative sample well containing pooled substituted ethylene diamine compounds.

Figure 15 is a mass spec profile for compound 109, *N*-Geranyl-*N'*-(2-adamanthyl)ethane-1,2-diamine.

Figure 16 is a proton NMR profile for compound 109, N-Geranyl-N¹-(2-adamanthyl) ethane-1,2- diamine.

Figure 17 is a bar graph of data from a Colony Forming Units/Lung (CFU/Lung) study showing CFU/Lung growth over time in days for various compounds.

5 Figure 18 is a bar graph of data from a CFU/Lung study showing CFU/Lung growth over time in days for various compounds.

Figure 19 is a bar graph of data from a CFU/Lung study showing CFU/Lung growth over time in days for various compounds.

10 Figure 20 is a bar graph of data from a lesion study showing visible lesions over time after treatment with various compounds.

Figure 21 provides a schematic demonstrating the identification of a drug candidate

Figure 22 provides the compounds tested for in vivo efficacy.

Figure 23 is a graph showing the results of in vivo studies of compounds 73 and 109 at 1 and 10 mg/kg doses (spleen).

15 Figure 24 is a graph showing the results of in vivo studies of compounds 73 and 109 at 1 and 10 mg/kg doses (lungs).

Figure 25 is a graph showing in vivo studies of compounds 59 and 111 at 1 and 10 mg/kg doses (spleen).

20 Figure 26 is a graph showing in vivo studies of compounds 59 and 111 at 1 and 10 mg/kg doses (lungs).

Figure 27 is a graph showing the results of efficacy testing of the compounds 58, 73, 109, and 111 in C57BL.6 mice infected with M.tuberculosis H37Rv (spleen). Mice were infected i.v. with 5×10^6 CFU M.tuberculosis H37Rv; treatment with drugs started 18 days following infection. EC - EC – early control, CFU in lungs of mice at the day of
25 chemotherapy start. Mice received: 1 – untreated mice, 2 – INH (25 mg/kg), 3 - EMB (100 mg/kg), 4 – comp. 109 (25 mg/kg), 4* - comp.109 (10 mg/kg), 4** - comp. 109 (0.1 mg/kg), 5 – comp. 58 (25 mg/kg), 6 – comp.73 (25 mg/kg), 7– comp. 111 (25 mg/kg).

Figure 28 is a graph showing the results of efficacy testing of the compounds 58, 73, 109, and 111 in C57BL.6 mice infected with M.tuberculosis H37Rv (lungs). Mice
30 were infected i.v. with 5×10^6 CFU M.tuberculosis H37Rv; treatment with drugs started 18 days following infection. EC - EC – early control, CFU in lungs of mice at the day of chemotherapy start. Mice received: 1 – untreated mice, 2 – INH (25 mg/kg), 3 - EMB (100

mg/kg), 4 – comp. 109 (25 mg/kg), 4* - comp.109 (10 mg/kg), 4** - comp. 109 (0.1 mg/kg), 5 – comp. 58 (25 mg/kg), 6 – comp.73 (25 mg/kg), 7– comp. 111 (25 mg/kg).

Figure 29 provides LC/MS data of tested compounds.

Figure 30 provides a graph showing results of PK studies with a cassette dosing of tested compounds to mice. Oral delivery. Compound NSC 722039 in the study reads as the compound 37, NSC 722040 – compound 59, NSC 722041 – compound 109.

Figure 31 provides a graph showing results of PK studies with a cassette dosing of tested compounds to mice. Peritoneal delivery. Compound NSC 722039 in the study reads as the compound 37, NSC 722040 – compound 59, NSC 722041 – compound 109.

Figure 32 provides a graph showing results of PK studies with a cassette dosing of tested compounds to mice. Intravenous delivery. Compound NSC 722039 in the study reads as the compound 37, NSC 722040 – compound 59, NSC 722041 – compound 109.

Figure 33 provides a graph showing the results of PK Studies of the compound 109 in mice.

Figure 34. Tissue distribution of 109 in mice (*i.v.*, 3 mg/kg).

Figure 35. Tissue distribution of 109 in mice (*p.o.*, 25 mg/kg).

Figure 36 Metabolism of the compound 109 in mouse urine.

Figure 37. No glucoronidation metabolites of 109 were found in mouse urine.

Figure 41. Scheme 1. Synthesis of 100,000 compound library of ethambutol analogues on solid support.

Figure 41. Scheme 2. Attempts to synthesize SQBisAd on solid support.

Figure 42 provides structures of representative targeted diamines prepared via acylation by amino acids.

Figure 43 provides Table 25 summarizing data for synthesized plates of diamines for the prepared library of targeted 20,000 ethambutol analogs.

Figure 44 provides Scheme 5 showing the synthesis of the diamine library using amino acids as linkers.

Figure 45 provides a schematic showing the occurrence of amine monomers in the hits that were generated in the original 100,000 compound library of EMB analogs.

Figure 46 provides a schematic showing structural diversity among primary amines.

Figure 47 provides Table 26 listing the amino acids that were used in the preparation of the diamine library.

Figure 48 provides carbonyl compounds used as reagents in the synthesis of the diamine library.

5 Figure 49 provides Table 27 showing carbonyl compounds used in the masterplate for the synthesis of the diamine library.

Figure 50 provides representative examples of MIC and Lux data for the diamine library.

10 Figure 51 provides a schematic showing the occurrence of alkylating monomers in final diamine products with anti-TB activity.

Figure 52 provides the layout of a representative 96-well deconvolution plate.

Figure 53 provides a list of compound hits and structures for the modified linker diamine library.

15 DETAILED DESCRIPTION

The present invention may be understood more readily by reference to the following detailed description of the specific embodiments included herein. However, although the present invention has been described with reference to specific details of certain embodiments thereof, it is not intended that such details should be regarded as
20 limitations upon the scope of the invention. The entire text of the references mentioned herein are hereby incorporated in their entireties by reference including United States Patent Application Serial No. 10/147,587 filed May 17, 2002, and United States Provisional Patent Application Serial No. 60/381,220 filed May 17, 2002.

25 Mycobacterial infections, such as those causing tuberculosis, once thought to be declining in occurrence, have rebounded, and again constitute a serious health threat. Tuberculosis (TB) is the cause of the largest number of human deaths attributed to a single etiologic agent with two to three million people infected with tuberculosis dying each year. Areas where humans are crowded together, or living in substandard housing, are increasingly found to have persons affected with mycobacteria. Individuals who are
30 immunocompromised are at great risk of being infected with mycobacteria and dying from

such infection. In addition, the emergence of drug-resistant strains of mycobacteria has led to treatment problems of such infected persons.

Many people who are infected with mycobacteria are poor, or live in areas with inadequate healthcare facilities. As a result of various obstacles (economical, education levels, etc.), many of these individuals are unable to comply with the prescribed therapeutic regimens. Ultimately, persistent non-compliance by these and other individuals results in the prevalence of disease. This noncompliance is frequently compounded by the emergence of drug-resistant strains of mycobacteria. Effective compositions and vaccines that target various strains of mycobacteria are necessary to bring the increasing number of tuberculosis cases under control.

Chemotherapy is a standard treatment for tuberculosis. Some current chemotherapy treatments require the use of three or four drugs, in combination, administered daily for two months, or administered biweekly for four to twelve months. Table 1 lists several treatment schedules for standard tuberculosis drug regimens.

Table 1

Treatment Schedules for Standard TB Drug Regimens.

STANDARD DRUG REGIMEN	INDUCTION PHASE Dosing Schedule	DURATION	DRUG	CONTINUATION PHASE Dosing Schedule	DURATION
Isoniazid	Daily, DOT	8 weeks	Isoniazid	2/week, DOT	16 weeks
Rifampicin	Daily, DOT	8 weeks	Rifampicin	2/week, DOT	16 weeks
Pyrazinamide	Daily, DOT	8 weeks			
Ethambutol or Streptomycin	Daily, DOT	8 weeks			

Decades of misuse of existing antibiotics and poor compliance with prolong and complex therapeutic regimens has led to mutations of the mycobacterium tuberculosis and has created an epidemic of drug resistance that threatens tuberculosis control world wide. The vast majority of currently prescribed drugs, including the front line drugs, such as isoniazid, rifampin, pyrazinamide, ethambutol and streptomycin were developed from the 1950s to the 1970s. Thus, this earlier development of tuberculosis chemotherapy did not have at its disposal the implications of the genome sequence of *Mycobacterium*

tuberculosis, the revolution in pharmaceutical drug discovery of the last decades, and the use of national drug testing and combinational chemistry.

Consequently, the treatments of drug-resistant *M. tuberculosis* strains, and latent tuberculosis infections, require new anti-tuberculosis drugs that provide highly effective
5 treatments, and shortened and simplified tuberculosis chemotherapies. Moreover, it is desirable that these drugs be prepared by a low-cost synthesis, since the demographics of the disease dictate that cost is a significant factor.

The present invention provides methods and compositions comprising a class of substituted ethylene diamine compounds effective in treatment and prevention of disease
10 caused by microorganisms including, but not limited to, bacteria. In particular, the methods and compositions of the present invention are effective in inhibiting the growth of the microorganism, *M. tuberculosis*. The methods and compositions of the present invention are intended for the treatment of mycobacteria infections in human, as well as other animals. For example, the present invention may be particularly useful for the
15 treatment of cows infected by *M. bovis*.


As used herein, the term "tuberculosis" comprises disease states usually associated with infections caused by mycobacteria species comprising *M. tuberculosis* complex. The term "tuberculosis" is also associated with mycobacterial infections caused by mycobacteria other than *M. tuberculosis* (MOTT). Other mycobacterial species include *M.*
20 *avium-intracellulare*, *M. kansarii*, *M. fortuitum*, *M. chelonae*, *M. leprae*, *M. africanum*, and *M. microti*, *M. avium paratuberculosis*, *M. intracellulare*, *M. scrofulaceum*, *M. xenopi*, *M. marinum*, *M. ulcerans*.

The present invention further comprises methods and compositions effective for the treatment of infectious disease, including but not limited to those caused by bacterial,
25 mycological, parasitic, and viral agents. Examples of such infectious agents include the following: staphylococcus, streptococcaceae, neisseriaaceae, cocci, enterobacteriaceae, pseudomonadaceae, vibriaceae, campylobacter, pasteuraceae, bordetella, francisella, brucella, legionellaceae, bacteroidaceae, gram-negative bacilli, clostridium, corynebacterium, propionibacterium, gram-positive bacilli, anthrax, actinomyces, nocardia,
30 mycobacterium, treponema, borrelia, leptospira, mycoplasma, ureaplasma, rickettsia, chlamydiae, systemic mycoses, opportunistic mycoses, protozoa, nematodes, trematodes, cestodes, adenoviruses, herpesviruses, poxviruses, papovaviruses, hepatitis viruses,

orthomyxoviruses, paramyxoviruses, coronaviruses, picornaviruses, reoviruses, togaviruses, flaviviruses, bunyaviridae, rhabdoviruses, human immunodeficiency virus and retroviruses.

The present invention further provides methods and compositions useful for the treatment of infectious disease, including by not limited to, tuberculosis, leprosy, Crohn's Disease, acquired immunodeficiency syndrome, lyme disease, cat-scratch disease, Rocky Mountain Spotted Fever and influenza.

The anti-infective methods and compositions of the present invention contain one or more substituted ethylene diamine compounds. In particular, these compounds encompass a wide range of substituted ethylene diamine compounds having the following general formula:



Substituted Ethylene Diamine

where "R₁NH" is typically derived from a primary amine, and "R₂R₃N" is typically derived from a primary or secondary amine. The ethylene diamines of the present invention are prepared by a modular approach using primary and secondary amines as building blocks, and coupling the amine moieties with an ethylene linker building block. Representative primary amines, acyclic secondary amines, and cyclic secondary amines are shown in Figures 2, 3, and 4, respectively.

Generally, chemical moieties R₁, R₂, and R₃ of the ethylene diamine compounds of the present invention are independently selected from H, alkyl; aryl; alkenyl; alkynyl; aralkyl; aralkenyl; aralkynyl; cycloalkyl; cycloalkenyl; heteroalkyl; heteroaryl; halide; alkoxy; aryloxy; alkylthio; arylthio; silyl; siloxy; a disulfide group; a urea group; amino; and the like, including straight or branched chain derivatives thereof, cyclic derivatives thereof, substituted derivatives thereof, heteroatom derivatives thereof, heterocyclic derivatives thereof, functionalized derivatives thereof, salts thereof, such salts including,

but not limited to hydrochlorides and acetates, isomers thereof, or combinations thereof. For example, nitrogen-containing heterocyclic moieties include, but are not limited to, groups such as pyridinyl (derived from pyridine, and bonded through a ring carbon), piperidinyl (derived from piperidine and bonded through the ring nitrogen atom or a ring carbon), and pyrrolidinyl (derived from pyrrolidine and bonded through the ring nitrogen atom or a ring carbon). Examples of substituted, or functionalized, derivatives of R₁, R₂, and R₃ include, but are not limited to, moieties containing substituents such as acyl, formyl, hydroxy, acyl halide, amide, amino, azido, acid, alkoxy, aryloxy, halide, carbonyl, ether, ester, thioether, thioester, nitrile, alkylthio, arylthio, sulfonic acid and salts thereof, thiol, alkenyl, alkynyl, nitro, imine, imide, alkyl, aryl, combinations thereof, and the like. Moreover, in the case of alkylated derivatives of the recited moieties, the alkyl substituent may be pendant to the recited chemical moiety, or used for bonding to the amine nitrogen through the alkyl substituent.

Examples of chemical moieties R₁, R₂, and R₃ of the present invention include, but are not limited to: H; methyl; ethyl; propyl; butyl; pentyl; hexyl; heptyl; octyl; ethenyl; propenyl; butenyl; ethynyl; propynyl; butynyl; cyclopropyl; cyclobutyl; cyclopentyl; cyclohexyl; cyclooctyl; cyclobutenyl; cyclopentenyl; cyclohexenyl; phenyl; tolyl; xylyl; benzyl; naphthyl; pyridinyl; furanyl; tetrahydro-1-naphthyl; piperidinyl; indolyl; indolinyl; pyrrolidinyl; 2-(methoxymethyl) pyrrolidinyl; piperazinyl; quinolinyl; quinolyl; alkylated-1,3-dioxolane; triazinyl; morpholinyl; phenyl pyrazolyl; indanyl; indonyl; pyrazolyl; thiadiazolyl; rhodaninyl; thiolactonyl; dibenzofuranyl; benzothiazolyl; homopiperidinyl; thiazolyl; quinonuclidinyl; isoxazolidinonyl; any isomers, derivatives, or substituted analogs thereof; or any substituted or unsubstituted chemical species such as alcohol, ether, thiol, thioether, tertiary amine, secondary amine, primary amine, ester, thioester, carboxylic acid, diol, diester, acrylic acid, acrylic ester, methionine ethyl ester, benzyl-L-cysteine ethyl ester, imine, aldehyde, ketone, amide, or diene. Further examples of chemical moieties R₁, R₂, and R₃ of the present invention include, but are not limited to, the following species or substituted or alkylated derivatives of the following species, covalently bonded to the amine nitrogen: furan; tetrahydrofuran; indole; piperazine; pyrrolidine; pyrrolidinone; pyridine; quinoline; anthracene; tetrahydroquinoline; naphthalene; pyrazole; imidazole; thiophene; pyrrolidine; morpholine; and the like. One feature of the recited species or substituted or alkylated derivatives of these species, is that they may be covalently bonded to the amine nitrogen in any fashion, including through the pendant substituent or alkyl

group, through the heteroatom as appropriate, or through a ring atom as appropriate, as understood by one of ordinary skill in the art.

The chemical moieties R_1 , R_2 , and R_3 of the present invention also include, but are not limited to, cyclic alkanes and cyclic alkenes, and include bridged and non-bridged
 5 rings. Examples of bridged rings include, but are not limited to, the following groups: isopinocampheyl; bornyl; norbornyl; adamantanetetyl; cis(-)myrtanyl; adamantyl; noradamantyl; 6-azabicyclo[3.2.1]octane; exo-norbornane; and the like.

In one embodiment of the present invention, NR_2R_3 is derived from a cyclic secondary amine. Examples of a cyclic chemical moiety, NR_2R_3 , of the present invention
 10 include, but are not limited to, 4-benzyl-piperidine; 3-piperidinemethanol; piperidine; tryptamine; morpholine; 4-piperidinopiperidine; ethyl 1-piperazine carboxylate; 1-(2-amino-ethyl)-piperazine; decahydroquinoline; 1,2,3,4-tetrahydro-pyridoindole (reaction at either amine); 3-amino-5-phenyl pyrazole; 3-aminopyrazole; 1-(2-fluorophenyl) piperazine; 1-proline methyl ester; histidinol; 1-piperonyl-piperazine; hexamethyleimine;
 15 4-hydroxypiperidine; 2-piperidinemethanol; 1, 3, 3-trimethyl-6-azabicyclo[3.2.1] octane; 3-pyrrolidinol; 1-methylpiperazine; (S)-(+)-(2-pyrrolidinylmethyl) pyrrolidine; 1-methylhomopiperazine; 2-ethyl-piperidine; 1, 2, 3, 4-tetrahydroisoquinoline; 1-(4-fluorophenyl) piperazine; d,l-tryptophan methyl ester; tert-butyl (15, 45)-(-)-2,5-diazabicyclo[2.2.1] heptane-2-carboxylate; isonipecotamide;
 20 heptamethyleneimine; alpha-methyltryptamine; 6, 7-dimethoxy-1, 2, 3, 4-tetrahydroisoquinoline; 3-aminopyrrolidine; 3, 5-dimethylpiperidine; 2, 6-dimethylmorpholine; 1,4-dioxo-8-azaspiro[4.5]decane; 1-methyl-6, 7-dihydroxy-1, 2, 3, 4-tetrahydroisoquinoline; 1, 3, 4, 6, 7, 8-hexahydro-2H-pyrido (1, 2-A) pyrimidine; 1, 2, 3, 4-tetrahydroquinoline; 1-(2-methoxyphenyl) piperazine; 1-(2-(2-hydroxyethoxy)ethyl)
 25 piperazine; (S)-(+)-2-(aminomethyl) pyrrolidine; (3S(3a, 4Ab), 8Ab)-N-t-butyl-D-decahydro-3-isoquinolinecarboxamide; (R)-cycloserine; homopiperazine; 2, 6-dimethylpiperazine (reaction at either amine); iminodibenzyl; 5-methoxytryptamine; 4, 4'-bipiperidine; 1-(2-hydroxyethyl) piperazine; 4-methylpiperidine; 1-histidine methyl ester; or methyl pipecolate.

30 The R_1HN substituent is derived from a primary amine. The R_2R_3N substituent is typically derived from a primary or secondary amine, but may also arise from an amino acid, or an amino acid precursor. The amino acid can transform into an amino alcohol.

When an amino acid is employed as the source of the R_2R_3N moiety, the precursor compound may be selected from, among others, the following compounds and their derivatives: d,l-tryptophan methyl ester; l-methionine ethyl ester; l-lysine methyl ester (via reaction at either primary amine); (S)-benzyl-1-cysteine ethyl ester; l-arginine methyl ester
5 (via reaction at either primary amine); l-glutamic acid ethyl ester; l-histidine methyl ester; or (3S (3a, 4Ab), 8A b)-N-t-butyl-D-ecahydro-3-iso-quinolinecarboxamide.

The R_4 moiety of the substituted ethylene diamine compounds of the present invention is typically selected from H, alkyl or aryl, but R_4 can also constitute alkenyl, alkynyl, aralkyl, aralkenyl, aralkynyl, cycloalkyl, cycloalkenyl, and the like. Examples of
10 the R_4 chemical moiety include, but are not limited to: H; methyl; ethyl; propyl; butyl; pentyl; hexyl; heptyl; octyl; ethenyl; propenyl; butenyl; ethynyl; propynyl; butynyl; cyclobutyl; cyclopentyl; cyclohexyl; cyclobutenyl; cyclopentenyl; cyclohexenyl; phenyl; tolyl; xylyl; benzyl; naphthyl; straight or branched chain derivatives thereof; cyclic derivatives thereof; substituted, functionalized, and heteroatom derivatives thereof; and
15 heterocyclic derivatives thereof, and the like. Typically, R_4 is selected from H, methyl, ethyl, butyl or phenyl. However, when R_4 is "H" the ethylene diamine does not contain ethambutol.

A majority of the ethylene diamine compounds described herein are preferably prepared using a solid support synthesis, as set forth in one of the representative reaction
20 schemes shown in Figure 1. However, when R_4 is H, the reaction does not proceed well when sterically hindered amines are used for R_1NH_2 , or when diamines, such as aminoalkylenemorpholine, or aminoalkylene-piperidines, are used for R_1NH_2 . When R_4 is methyl, or phenyl, sterically hindered amines used for R_3R_2NH do not work well due to steric hindrance at the reaction site. In this case, a competing hydrolysis reaction
25 producing the corresponding amino alcohols, and incomplete reduction of the amidoethyleneamines, interfere with the reaction scheme. As a result, the desired diamine products form in low yields.

The preparation of the ethylene diamines is preferably accomplished in six steps, using a rink-acid resin. The first step of the synthesis is converting the rink-acid resin to
30 rink-chloride by treatment with triphenylphosphine and hexachloroethane in tetrahydrofuran (THF). This step is followed by addition of the primary amine in the

presence of Hunig's base ($\text{EtN}(\text{i-Pr})_2$) in dichloroethane. The third step is the acylation of the resin-attached amine using either one of the two acylation routes shown in Figure 1. The acylation step is preferably accomplished using either α -chloroacetyl chloride, α -bromo- α -methyl acetyl bromide, α -bromo- α -ethylacetyl bromide, α -bromo- α -butyl acetyl bromide, or α -chloro- α -phenyl-acetyl chloride, each in the presence of pyridine in THF. Other acylation reagents known to those skilled in the art may also be used, however, the α -bromoacetyl halides result in low product yields, which may be attributed to HBr elimination. The acylation may also be accomplished via a peptide coupling mechanism using α -bromo- α -methylacetic acid, or α -chloro- α -methylacetic acid, in the presence of benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate (PyBrop) and N_1N -diisopropylethyl amine ($\text{EtN}(\text{i-Pr})_2$) in dichloromethane (DCM) and dimethylformamide (DMF). Again, other acylation reagents known to those skilled in the art may also be used. The acylation step is preferably performed twice to achieve better acylated product yields.

Introduction of the second nitrogen moiety is preferably achieved in the presence of Hunig's base in dimethylformamide (DMF). Reduction of the intermediate amine-amide is carried out using Red-Al (3.4M solution of sodium bis (2-methoxyethoxy) aluminum hydride in toluene). The final product is cleaved from the resin support using a 10% solution (by volume) of trifluoroacetic acid (TFA) in dichloromethane (DCM). The solvent is evaporated, and the TFA salts of the final diamine products are analyzed by mass spec, and screened against *M. tuberculosis* for effectiveness. Some of the substituted ethylene diamines, prepared using the above-described solid-support synthesis, are also prepared using a solution phase synthesis described below.

Formation of the Substituted Ethylene Diamine Library

The solid support syntheses, shown in Figure 1, are preferably used to prepare a substituted ethylene diamine library. Solid phase synthesis offers at least three principal advantages: (i) a reduced need for chromatographic procedures, (ii) the use of excess reagents to drive a reaction forward in high yields, and (iii) the use of split and pool technologies for the synthesis of a large number of compounds. Solid support syntheses of 1, 2-diamine libraries have previously been accomplished by the reduction of short peptides (Cuervo et al., Peptides 1994: Proceedings of the European Peptide Symposium; Maia HSL Ed., Esom: Leiden, 1995, 465-466). However, as described herein, an ethylene

diamine library is created using amines, rather than simple amino acids, to allow for greater diversity in the building-block monomers. The first three steps of each support synthesis: the activation of the Rink-acid resin, the addition of the first amine, and the acylation step are carried out in 10 ml tubes on a QUEST® 210 Synthesizer manufactured by ARGONAUT TECHNOLOGIES®, Inc., Foster City, California. The synthesizer handles up to twenty simultaneous reactions in 5 ml or 10 ml reaction vessels to allow for rapid synthesis of target compounds. The synthesizer provides programmable temperature control and agitation, and the automated delivery of solvents into the reaction vessels. The addition of the second amine, the reduction with Red-Al, and the cleavage from the solid support are carried out in 2 ml wells in a 96-well, chemically resistant plate.

Prior to the solid support synthesis, each amine, within numbers 1 to 288, as shown in Figures 2, 3, and 4, is dissolved in DMF as a one molar solution, and organized in three, 96-well plates (one amine per well), to yield three master plates of these amines. An individual haloacetyl amide from each primary amine and a particular R₄ group, is formed in the first three steps of the support synthesis. Individual haloacetyl amides are then pooled into groups of ten or thirty. A suspension of the pooled resins in a 2:1 mixture of DCM/THF is evenly distributed into one, two or three reaction plates to assure 15-20 mg of the suspension per well. The number of reaction plates used is based on the amount of suspension available. Each well of pooled resins is reacted with a corresponding amine from the master plates. Figure 5 provides a flow schematic for a representative pool. Each reaction occurs in a separate well, in the presence of Hunig's base in DMF at 70-75°C for 16-20 hours. Each resulting amine-amide is reduced using 65+w% Red-Al at room temperature. The reduction is followed by cleavage with 10% vol. TFA in DCM. The solvents in each reaction well are evaporated, and the TFA salts of the diamines analyzed (mass spec), and screened against *M. tuberculosis*. One plate of pooled diamines are screened against *M. smegmatis*. Two randomly selected rows in each plate; i.e., 24 samples per 96-well plate, or 25% of the library, are examined by mass spectroscopy. Specific protocols and detailed methods are provided below in the Examples.

Screening Against *M. tuberculosis*

An entire library of synthesized substituted ethylene diamines (targeted number of compounds about 100,000), prepared as described above, was screened, *in vitro*, against *M. tuberculosis* in ethambutol (EMB) sensitive Luc-assay. The MIC (Minimum Inhibition

Concentration) was also determined. The MIC is the minimum concentration of a growth inhibitor, here the substituted ethylene diamine, where there is no multiplication of the microorganism under examination. Screening was done using a High-Throughput Screening (HTS) Luc assay with recombinant mycobacteria containing a promoter fusion of a luciferase to the EB-inducible gene (Luc assay). The Luc-assay and MIC assay are described in detail below. These assays are well known to those skilled in the art. Based on this initial screening, 300+ compound mixtures showed anti-TB activity. Figure 6 represents typical assay data in a luciferase reporter strain containing an Rv0341 EMB-inducible promoter. Figure 6 represents percent maximum Luminescence Count per Second (% Max. LCPS) for pooled compound mixtures in one row (row D) in one of the 96-well plates.

Deconvolution of the Reactive Wells

The *M. tuberculosis* screening revealed approximately 300 active compounds mixtures that were selected for deconvolution. In particular, wells possessing activity of approximately $<12.5\mu\text{M}$ in the HTS Luc assay, and/or an MIC of approximately $<12.5\mu\text{M}$, were selected for a total of 336 wells.

Deconvolutions were performed by discrete re-synthesis of each substituted ethylene diamine compound in each active compound pool. The pooled compounds in each active well were individually synthesized, and screened. Syntheses of the targeted diamine compounds in each active pool were done in the 96-well plates using stored archived α -haloacetyl amides (resin attached haloacetyl amides), according to the previously described reaction steps (the addition of the second amine, the reduction with Red-Al, and the cleavage from the solid support). The archived resins were stored as individual compounds at 4°C . The 96-well plates were used for the remaining synthesis steps as previously described.

The same screening tests, MIC and HTS Luc assay, were performed on each deconvoluted compound. Representative Luminescence data for deconvoluted compounds are shown in Figures 7 and 8. Figures 7 and 8 represent the Luminescence Count per Second (LCPS) for individual compounds.

Summary of Screening Results

Overall, the deconvolution screening results revealed about 2,000 ethylene diamine compounds with inhibitory activity against *M. tuberculosis*. More than 150 of these compounds exhibited MICs equal to or lower than approximately 12.5 μ M. Figure 9 summarizes the MIC data for all synthesized discrete compounds with an MIC of 50 μ M or less. Figure 10 summarizes Luc assay data for all compounds that exhibit at least 10% activity at each concentration (the results are not cumulative). The MIC and Luc activities were obtained for non-purified samples, with chemical yields of approximately 20%, based on an assumed 80% yield at each reaction step. In the Luc assay, 32 compounds exhibited activity at 1.56 μ M, and in the MIC assay, at least 11 compounds had an MIC of 3.13 μ M.

The total frequency of the top thirteen amines that contributed to the activity of the substituted ethylene diamines are shown in Figure 11, with each amine represented by its numerical designation. These amines include the following:

	#11	2,3-Dimethylcyclohexylamines
15	#18	3,3-Diphenylpropylamine
	#44	1-Adamantanemethylamine
	#47	2,2-Diphenylethylamine
	#63	(S)-2-Amino-1-butanol
	#74.1	(-)-cis-Myrtanylamine
20	#77.1	Cyclooctylamine
	#78.1	2-Adamantamine
	#105a	(1R,2R,3R,5S)-(-)-Isopinocampheylamine
	#231	2-Methoxyphenethylamine
	#255	(S)-Cylcohexylethylamine
25	#266	Undecylamine
	#272	Geranylamine

Other amines that contributed to the activity of the substituted ethylene diamines are shown in Table 2. The compounds in Table 2 are sorted by their MIC results. Some

compounds, synthesized in larger quantities (2-60mg) on the Quest® Synthesizer, and purified by HPLC using semi-preparative C18-column, are shown in Table 3. Generally, the final purity of each compound in Table 3 was at least 90%.

Table 2

5

N1	N2	R4	MIC (uM)	% Induction
3,3-Diphenylpropylamine	exo-Aminonorbornane	Hydrogen	3.13	53.70
2,2-Diphenylamine	(+)-Isopinocampheylamine	Hydrogen	3.13	93.94
2,2-Diphenylamine	cis-(-)-Myrtanylamine	Hydrogen	3.13	64.49
2,2-Diphenylamine	Cyclooctylamine	Hydrogen	3.13	63.44
2,2-Diphenylamine	3,4-Dihydroxynorephedrine	Hydrogen	3.13	42.80
5-Aminoquinoline	Cyclohexylamine	Hydrogen	3.13	18.33
5-Aminoquinoline	tert-Octylamine	Hydrogen	3.13	20.85
5-Aminoquinoline	4-Methylcyclohexylamine	Hydrogen	3.13	26.33
cis-(-)-Myrtanylamine	(+)-Bornylamine	Hydrogen	3.13	100.00
cis-(-)-Myrtanylamine	1-Adamantanemethylamine	Hydrogen	3.13	85.20
cis-(-)-Myrtanylamine	(-)-Isopinocampheylamine	Hydrogen	3.13	60.94
1-Adamantanemethylamine	tert-Octylamine	Hydrogen	4.7	9.81
3,4-Dimethoxyphenethylamine	1-Adamantanemethylamine	Hydrogen	6.25	11.45
3,4-Dimethoxyphenethylamine	Hexetidine (mixture of isomers)	Hydrogen	6.25	0
3,4-Dimethoxyphenethylamine	Dehydroabietylamine	Hydrogen	6.25	0
3,3-Diphenylpropylamine	1-Adamantanemethylamine	Hydrogen	6.25	9.53
3,3-Diphenylpropylamine	2-Methylcyclohexylamine (mix of cis and trans)	Hydrogen	6.25	50.08
3,3-Diphenylpropylamine	1,3-Dimethylbutylamine	Hydrogen	6.25	39.40
3,3-Diphenylpropylamine	1-(1-Adamantyl)ethylamine, HCl	Hydrogen	6.25	45.14
3,3-Diphenylpropylamine	(S)-(-)-Cyclohexylethylamine	Hydrogen	6.25	43.49
3,3-Diphenylpropylamine	(R)-(-)-Cyclohexylethylamine	Hydrogen	6.25	34.54
3,3-Diphenylpropylamine	1-Adamantanemethylamine	Methyl	6.25	16.14
Propylamine	Hexetidine (mixture of isomers)	Hydrogen	6.25	0
Phenethylamine	Hexetidine (mixture of isomers)	Hydrogen	6.25	0
b-Methylphenethylamine	Hexetidine (mixture of isomers)	Hydrogen	6.25	0
b-Methylphenethylamine	Undecylamine	Hydrogen	6.25	0
2,2-Diphenylamine	(+)-Bornylamine	Hydrogen	6.25	87.86
2,2-Diphenylamine	(-)-Isopinocampheylamine	Hydrogen	6.25	77.80
2,2-Diphenylamine	alpha-Methyltryptamine	Hydrogen	6.25	55.07
2,2-Diphenylamine	alpha-Methyltryptamine	Hydrogen	6.25	23.08
2,2-Diphenylamine	4-Phenylbutylamine	Hydrogen	6.25	
2,2-Diphenylamine	2,5-Dimethoxyphenethylamine	Hydrogen	6.25	
2,2-Diphenylamine	2,4-Dichlorophenethylamine	Hydrogen	6.25	
2,2-Diphenylamine	2-(2-Aminomethyl)phenylthio)benzyl alcohol	Hydrogen	6.25	
2,2-Diphenylamine	1-(1-Naphthyl)ethylamine	Hydrogen	6.25	7.20
Veratryl amine	2,5-Dimethoxyphenethylamine	Hydrogen	6.25	

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
Veratryl amine	2-(2-Aminomethyl) phenylthio)benzyl alcohol	Hydrogen	6.25	
5-Aminoquinoline	2-Aminoheptane	Hydrogen	6.25	26.22
5-Aminoquinoline	1-Adamantanamine	Hydrogen	6.25	18.91
1-Aminomethyl-1-cyclohexanol, HCl	Hexetidine (mixture of isomers)	Hydrogen	6.25	
cis-(-)-Myrtaniline	2,3-Dimethylcyclohexylamine	Hydrogen	6.25	100.00
cis-(-)-Myrtaniline	3,3-Diphenylpropylamine	Hydrogen	6.25	87.78
cis-(-)-Myrtaniline	(+)-Isopinocampheylamine	Hydrogen	6.25	93.10
cis-(-)-Myrtaniline	2,2-Diphenylamine	Hydrogen	6.25	81.84
cis-(-)-Myrtaniline	cis-(-)-Myrtaniline	Hydrogen	6.25	68.24
cis-(-)-Myrtaniline	1,3,3-Trimethyl-6-azabicyclo[3.2.1]octane	Hydrogen	6.25	68.18
cis-(-)-Myrtaniline	1-Adamantanemethylamine	Methyl	6.25	24.22
cis-(-)-Myrtaniline	cis-(-)-Myrtaniline	Methyl	6.25	44.14
Cyclooctylamine	3,3-Diphenylpropylamine	Hydrogen	6.25	100.00
Cyclooctylamine	(-)-Isopinocampheylamine	Hydrogen	6.25	59.13
sec-Butylamine	Hexetidine (mixture of isomers)	Hydrogen	6.25	
3-Methylbenzylamine	Hexetidine (mixture of isomers)	Hydrogen	6.25	
3-Methylbenzylamine	Undecylamine	Hydrogen	6.25	
2-Methoxyethylamine	Hexetidine (mixture of isomers)	Hydrogen	6.25	
Geranylamine	2-Adamantanamine, HCl	Hydrogen	6.25	25.66
1-Adamantanemethylamine	4-Benzylpiperidine	Hydrogen	9.4	0
1-Adamantanemethylamine	2,3-Dimethylcyclohexylamine	Hydrogen	9.4	0
1-Adamantanemethylamine	3,3-Diphenylpropylamine	Hydrogen	9.4	40.06
1-Adamantanemethylamine	1-Adamantanemethylamine	Hydrogen	9.4	15.25
1-Adamantanemethylamine	2,2-Diphenylamine	Hydrogen	9.4	0
1-Adamantanemethylamine	1,3,3-Trimethyl-6-azabicyclo[3.2.1]octane	Hydrogen	9.4	0
1-Adamantanemethylamine	138	Hydrogen	9.4	0
3-Phenyl-1-propylamine	138	Hydrogen	9.4	
2,2-Diphenylamine	1-Adamantanemethylamine	Hydrogen	9.4	65.89
2,2-Diphenylamine	138	Hydrogen	9.4	
Furfurylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	0
3,4,5-Trimethoxybenzylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	0
1-Methyl-3-phenylpropylamine	Dehydroabietylamine	Hydrogen	12.5	0
Cyclobutylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	0
2-Fluorobenzylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	0
2-Fluorobenzylamine	Dehydroabietylamine	Hydrogen	12.5	0
3,4-Dimethoxyphenethylamine	Undecylamine	Hydrogen	12.5	0
3,3-Diphenylpropylamine	exo-Aminonorborene	Hydrogen	12.5	14.38
3,3-Diphenylpropylamine	Decahydroquinoline	Hydrogen	12.5	22.52
3,3-Diphenylpropylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	0
3,3-Diphenylpropylamine	4-Phenylbutylamine	Hydrogen	12.5	0
3,3-Diphenylpropylamine	2-Methoxyphenethylamine	Hydrogen	12.5	6.82
3,3-Diphenylpropylamine	2,4-Dichlorophenethylamine	Hydrogen	12.5	0

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
3,3-Diphenylpropylamine	1-Aminoindan	Hydrogen	12.5	18.05
3,3-Diphenylpropylamine	Undecylamine	Hydrogen	12.5	0
3,3-Diphenylpropylamine	Dehydroabietylamine	Hydrogen	12.5	0
3,3-Diphenylpropylamine	2-(1-Cyclohexenyl)ethylamine	Methyl	12.5	9.5
3,3-Diphenylpropylamine	cis-(-)-Myrtanylamine	Methyl	12.5	18.41
3,3-Diphenylpropylamine	Cyclooctylamine	Methyl	12.5	20.84
Propylamine	Dehydroabietylamine	Hydrogen	12.5	0
Phenethylamine	Dehydroabietylamine	Hydrogen	12.5	0
Cyclohexylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	0
3-Amino-1-propanol	Hexetidine (mixture of isomers)	Hydrogen	12.5	0
b-Methylphenethylamine	Dehydroabietylamine	Hydrogen	12.5	0
4-Methoxyphenethylamine	2-Fluorophenethylamine	Hydrogen	12.5	0
4-Methoxyphenethylamine	2-(1-Cyclohexenyl)ethylamine	Hydrogen	12.5	0
4-Methoxyphenethylamine	2,4-Dimethoxybenzylamine	Hydrogen	12.5	0
4-Methoxyphenethylamine	4-Fluorophenethylamine	Hydrogen	12.5	16.78
4-Methoxyphenethylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	0
Tetrahydrofurfurylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	0
Amylamine	4-Fluorophenethylamine	Hydrogen	12.5	0
3-Phenyl-1-propylamine	2-(1-Cyclohexenyl)ethylamine	Hydrogen	12.5	
3-Phenyl-1-propylamine	4-Fluorophenethylamine	Hydrogen	12.5	12.94
2,2-Diphenylamine	tert-Amylamine	Hydrogen	12.5	9.05
2,2-Diphenylamine	Undecylamine	Hydrogen	12.5	
2,2-Diphenylamine	Dehydroabietylamine	Hydrogen	12.5	
2,2-Diphenylamine	cis-(-)-Myrtanylamine	Methyl	12.5	45.18
1-(3-Aminopropyl)-2-pyrrolidinone (tech)	2,5-Dimethoxyphenethylamine	Hydrogen	12.5	
1-(3-Aminopropyl)-2-pyrrolidinone (tech)	2-(2-Aminomethyl)phenylthio)benzyl alcohol	Hydrogen	12.5	
4-(Trifluoromethyl)benzylamine	2,5-Dimethoxyphenethylamine	Hydrogen	12.5	
4-(Trifluoromethyl)benzylamine	1-(1-Naphthyl)ethylamine	Hydrogen	12.5	
Veratryl amine	4-Phenylbutylamine	Hydrogen	12.5	
5-Amino-1-pentanol	2,5-Dimethoxyphenethylamine	Hydrogen	12.5	
5-Amino-1-pentanol	2-(2-Aminomethyl)phenylthio)benzyl alcohol	Hydrogen	12.5	
2-(1-Cyclohexenyl)ethylamine	2-(1-Cyclohexenyl)ethylamine	Hydrogen	12.5	
2-(1-Cyclohexenyl)ethylamine	4-Fluorophenethylamine	Hydrogen	12.5	
2-(1-Cyclohexenyl)ethylamine	4-Phenylbutylamine	Hydrogen	12.5	
2-(1-Cyclohexenyl)ethylamine	2,5-Dimethoxyphenethylamine	Hydrogen	12.5	
2-(1-Cyclohexenyl)ethylamine	2-(2-Aminomethyl)phenylthio)benzyl alcohol	Hydrogen	12.5	
1-Aminomethyl-1-cyclohexanol, HCl	2,5-Dimethoxyphenethylamine	Hydrogen	12.5	
3-Fluorobenzylamine	2,5-Dimethoxyphenethylamine	Hydrogen	12.5	
4-Amino-1-butanol	Hexetidine (mixture of isomers)	Hydrogen	12.5	

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
2-Ethoxybenzylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	
cis-(-)-Myrtanylamine	Cyclooctylamine	Hydrogen	12.5	67.73
cis-(-)-Myrtanylamine	4-Methylcyclohexylamine	Hydrogen	12.5	18.39
cis-(-)-Myrtanylamine	1-Adamantanamine	Hydrogen	12.5	60.16
cis-(-)-Myrtanylamine	3,3-Diphenylpropylamine	Methyl	12.5	22.32
Cyclooctylamine	(+)-Isopinocampheylamine	Hydrogen	12.5	57.83
Cyclooctylamine	(+)-Bornylamine	Hydrogen	12.5	100.00
Cyclooctylamine	1-Adamantanemethylamine	Hydrogen	12.5	52.95
Cyclooctylamine	2,2-Diphenylamine	Hydrogen	12.5	71.43
Cyclooctylamine	cis-(-)-Myrtanylamine	Hydrogen	12.5	84.56
Cyclooctylamine	Cyclooctylamine	Hydrogen	12.5	59.21
Cyclooctylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	
Cyclooctylamine	Aminodiphenylmethane	Hydrogen	12.5	
Cyclooctylamine	Undecylamine	Hydrogen	12.5	5.61
Cyclooctylamine	3,3-Diphenylpropylamine	Methyl	12.5	53.92
Cyclooctylamine	(+)-Isopinocampheylamine	Methyl	12.5	
Cyclooctylamine	cis-(-)-Myrtanylamine	Methyl	12.5	33.89
4-Chlorophenylalaninol	Hexetidine (mixture of isomers)	Hydrogen	12.5	
(-)-Isopinocampheylamine	3,3-Diphenylpropylamine	Hydrogen	12.5	23.68
(-)-Isopinocampheylamine	(+)-Bornylamine	Hydrogen	12.5	44.85
(-)-Isopinocampheylamine	2-Amino-1-propanol, d,l	Hydrogen	12.5	46.19
(-)-Isopinocampheylamine	cis-(-)-Myrtanylamine	Hydrogen	12.5	33.87
(-)-Isopinocampheylamine	2-Adamantanamine, HCl	Hydrogen	12.5	24.29
(-)-Isopinocampheylamine	Aminodiphenylmethane	Hydrogen	12.5	48.35
Allylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	
3-Ethoxypropylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	
sec-Butylamine	Dehydroabietylamine	Hydrogen	12.5	
2-Aminoheptane	Dehydroabietylamine	Hydrogen	12.5	
Ethanolamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	
3-Methylbenzylamine	4-Phenylbutylamine	Hydrogen	12.5	
3-Methylbenzylamine	2,4-Dichlorophenethylamine	Hydrogen	12.5	
3-Methylbenzylamine	Dehydroabietylamine	Hydrogen	12.5	
Piperonylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	
Piperonylamine	Dehydroabietylamine	Hydrogen	12.5	
2-Methoxyethylamine	Dehydroabietylamine	Hydrogen	12.5	
4-Fluorophenethylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	
3-o-Methyldopamine, HCl	Hexetidine (mixture of isomers)	Hydrogen	12.5	
3-o-Methyldopamine, HCl	Undecylamine	Hydrogen	12.5	
3-o-Methyldopamine, HCl	Dehydroabietylamine	Hydrogen	12.5	
3-Fluorophenethylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	
3-Fluorophenethylamine	Dehydroabietylamine	Hydrogen	12.5	
2-Methoxyphenethylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	
2-Methoxyphenethylamine	Aminodiphenylmethane	Hydrogen	12.5	34.67
2-Fluoroethylamine, HCl	Hexetidine (mixture of isomers)	Hydrogen	12.5	
2-Amino-1-phenylethanol	Hexetidine (mixture of isomers)	Hydrogen	12.5	
2-Amino-1-phenylethanol	Dehydroabietylamine	Hydrogen	12.5	

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
2,5-Dimethoxyphenethylamine	2-Adamantanamine, HCl	Hydrogen	12.5	22.18
2-(2-Chlorophenyl)ethylamine	N-Allylcyclopentylamine	Hydrogen	12.5	62.31
2-(2-Chlorophenyl)ethylamine	Hexetidine (mixture of isomers)	Hydrogen	12.5	
3-Hydroxytyramine	Hexetidine (mixture of isomers)	Hydrogen	12.5	
4-(Trifluoromethoxy)benzylamine	2-Adamantanamine, HCl	Hydrogen	12.5	28.34
Geranylamine	(+)-Bornylamine	Hydrogen	12.5	
Geranylamine	1,3,3-Trimethyl-6-azabicyclo[3.2.1]octane	Hydrogen	12.5	37.42
Geranylamine	2-Ethylpiperidine	Hydrogen	12.5	29.81
Geranylamine	1-Adamantanamine	Hydrogen	12.5	16.63
Geranylamine	N-Allylcyclopentylamine	Hydrogen	12.5	74.86
Geranylamine	Aminodiphenylmethane	Hydrogen	12.5	57.93
Geranylamine	Dehydroabietylamine	Hydrogen	12.5	
1-Adamantanemethylamine	Decahydroquinoline	Hydrogen	18.8	0
1-Adamantanemethylamine	1-Adamantanamine	Hydrogen	18.8	0
2,2-Diphenylamine	2,3-Dimethylcyclohexylamine	Hydrogen	18.8	23.60
2,2-Diphenylamine	tert-Octylamine	Hydrogen	18.8	19.29
2,2-Diphenylamine	Decahydroquinoline	Hydrogen	18.8	8.96
4-Methylbenzylamine	Furfurylamine	Hydrogen	25	13.46
4-Methylbenzylamine	Benzylamine	Hydrogen	25	17.07
4-Methylbenzylamine	Hexetidine (mixture of isomers)	Hydrogen	25	0
4-Methylbenzylamine	Dehydroabietylamine	Hydrogen	25	0
Cyclopentylamine	Hexetidine (mixture of isomers)	Hydrogen	25	0
Cyclopentylamine	Dehydroabietylamine	Hydrogen	25	0
Furfurylamine	Furfurylamine	Hydrogen	25	0
1-Methyl-3-phenylpropylamine	Hexetidine (mixture of isomers)	Hydrogen	25	0
1-Methyl-3-phenylpropylamine	Undecylamine	Hydrogen	25	0
1,2,3,4-Tetrahydro-1-naphthylamine	Undecylamine	Hydrogen	25	6.24
1,2,3,4-Tetrahydro-1-naphthylamine	Dehydroabietylamine	Hydrogen	25	0
2,3-Dimethylcyclohexylamine	Undecylamine	Hydrogen	25	0
2,3-Dimethylcyclohexylamine	Dehydroabietylamine	Hydrogen	25	0
Tyramine	Hexetidine (mixture of isomers)	Hydrogen	25	0
Tyramine	Undecylamine	Hydrogen	25	0
Tyramine	Dehydroabietylamine	Hydrogen	25	0
Tyramine	cis-(-)-Myrtanylamine	Methyl	25	0
2-Fluorobenzylamine	Undecylamine	Hydrogen	25	0
(R)-2-Amino-1-butanol	Hexetidine (mixture of isomers)	Hydrogen	25	0
3,3-Diphenylpropylamine	(S)-(+)-1-Amino-2-propanol	Hydrogen	25	0
3,3-Diphenylpropylamine	2-Ethylpiperidine	Hydrogen	25	11.32
3,3-Diphenylpropylamine	N-Allylcyclopentylamine	Hydrogen	25	11.63
3,3-Diphenylpropylamine	Aminodiphenylmethane	Hydrogen	25	0
3,3-Diphenylpropylamine	3,5-Dimethylpiperidine (cis- and trans-)	Hydrogen	25	30.28

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
3,3-Diphenylpropylamine	Allylcyclohexylamine	Hydrogen	25	9.10
Propylamine	Undecylamine	Hydrogen	25	0
Phenethylamine	Undecylamine	Hydrogen	25	0
Tryptamine	(S)-(+)-1-Amino-2-propanol	Hydrogen	25	0
Tryptamine	2-Amino-2-methyl-1-propanol	Hydrogen	25	0
Cyclohexylamine	Undecylamine	Hydrogen	25	0
Cyclohexylamine	Dehydroabietylamine	Hydrogen	25	0
(+)-Isopinocampheylamine	Dehydroabietylamine	Hydrogen	25	0
Benzylamine	Hexetidine (mixture of isomers)	Hydrogen	25	
Benzylamine	Undecylamine	Hydrogen	25	
3-Amino-1-propanol	Dehydroabietylamine	Hydrogen	25	0
2-Fluorophenethylamine	2-Fluorophenethylamine	Hydrogen	25	0
2-Fluorophenethylamine	Veratryl amine	Hydrogen	25	0
2-Fluorophenethylamine	2,4-Dimethoxybenzylamine	Hydrogen	25	0
2-Fluorophenethylamine	2-Amino-2-methyl-1-propanol	Hydrogen	25	0
2-Fluorophenethylamine	4-Fluorophenethylamine	Hydrogen	25	0
2-Fluorophenethylamine	Hexetidine (mixture of isomers)	Hydrogen	25	0
2-Fluorophenethylamine	1-(1-Naphthyl)ethylamine	Hydrogen	25	0
2-Fluorophenethylamine	1-Adamantanemethylamine	Methyl	25	3.21
2-Fluorophenethylamine	cis-(-)-Myrtanylamine	Methyl	25	4.89
b-Methylphenethylamine	4-Phenylbutylamine	Hydrogen	25	0
b-Methylphenethylamine	2,4-Dichlorophenethylamine	Hydrogen	25	0
b-Methylphenethylamine	1-(1-Naphthyl)ethylamine	Hydrogen	25	0
4-Methoxyphenethylamine	1-Adamantanemethylamine	Hydrogen	25	0
4-Methoxyphenethylamine	1-(3-Aminopropyl)-2-pyrrolidinone (tech)	Hydrogen	25	0
4-Methoxyphenethylamine	Veratryl amine	Hydrogen	25	0
4-Methoxyphenethylamine	Undecylamine	Hydrogen	25	0
4-Methoxyphenethylamine	Dehydroabietylamine	Hydrogen	25	0
Tetrahydrofurfurylamine	Dehydroabietylamine	Hydrogen	25	0
Amylamine	2-Fluorophenethylamine	Hydrogen	25	0
Amylamine	2-(1-Cyclohexenyl)ethylamine	Hydrogen	25	0
Amylamine	2,4-Dimethoxybenzylamine	Hydrogen	25	0
3-Phenyl-1-propylamine	2-Fluorophenethylamine	Hydrogen	25	
3-Phenyl-1-propylamine	1-Adamantanemethylamine	Hydrogen	25	
3-Phenyl-1-propylamine	2,4-Dimethoxybenzylamine	Hydrogen	25	
3-Phenyl-1-propylamine	Hexetidine (mixture of isomers)	Hydrogen	25	
3-Phenyl-1-propylamine	4-Phenylbutylamine	Hydrogen	25	
3-Phenyl-1-propylamine	2,4-Dichlorophenethylamine	Hydrogen	25	
3-Phenyl-1-propylamine	Undecylamine	Hydrogen	25	
3-Phenyl-1-propylamine	Dehydroabietylamine	Hydrogen	25	
2,2-Diphenylamine	4-(2-Aminoethyl)morpholine	Hydrogen	25	
2,2-Diphenylamine	1-(3-Aminopropyl)-2-pyrrolidinone (tech)	Hydrogen	25	
2,2-Diphenylamine	2-(1-Cyclohexenyl)ethylamine	Hydrogen	25	
2,2-Diphenylamine	2,4-Dimethoxybenzylamine	Hydrogen	25	

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
2,2-Diphenylamine	4-(3-Aminopropyl)morpholine	Hydrogen	25	
2,2-Diphenylamine	4-Fluorophenethylamine	Hydrogen	25	
2,2-Diphenylamine	Hexetidine (mixture of isomers)	Hydrogen	25	
2,2-Diphenylamine	(S)-(-)-Cyclohexylethylamine	Hydrogen	25	
2,2-Diphenylamine	1-Adamantanemethylamine	Methyl	25	5.84
1-(3-Aminopropyl)-2-pyrrolidinone (tech)	4-Phenylbutylamine	Hydrogen	25	
4-(Trifluoromethyl)benzylamine	1-Adamantanemethylamine	Hydrogen	25	
4-(Trifluoromethyl)benzylamine	tert-Amylamine	Hydrogen	25	
4-(Trifluoromethyl)benzylamine	alpha-Methyltryptamine	Hydrogen	25	6.06
4-(Trifluoromethyl)benzylamine	4-Phenylbutylamine	Hydrogen	25	
4-(Trifluoromethyl)benzylamine	2-(2-Aminomethyl)phenylthio)benzyl alcohol	Hydrogen	25	5.13
4-(Trifluoromethyl)benzylamine	Undecylamine	Hydrogen	25	
4-(Trifluoromethyl)benzylamine	(-)-3,4-Dihydroxynorephedrine	Hydrogen	25	
4-(Trifluoromethyl)benzylamine	Dehydroabietylamine	Hydrogen	25	
Veratryl amine	tert-Amylamine	Hydrogen	25	
5-Amino-1-pentanol	4-Phenylbutylamine	Hydrogen	25	
2-(1-Cyclohexenyl)ethylamine	2-Fluorophenethylamine	Hydrogen	25	
2-(1-Cyclohexenyl)ethylamine	1-Adamantanemethylamine	Hydrogen	25	
1-Aminomethyl-1-cyclohexanol, HCl	4-Phenylbutylamine	Hydrogen	25	
3-Fluorobenzylamine	4-Phenylbutylamine	Hydrogen	25	
3-Fluorobenzylamine	2-(2-Aminomethyl)phenylthio)benzyl alcohol	Hydrogen	25	
2,4-Dimethoxybenzylamine	1-Adamantanamine	Hydrogen	25	
2,4-Dimethoxybenzylamine	Hexetidine (mixture of isomers)	Hydrogen	25	
2,4-Dimethoxybenzylamine	Undecylamine	Hydrogen	25	
2,4-Dimethoxybenzylamine	Dehydroabietylamine	Hydrogen	25	
2-Ethoxybenzylamine	1-Adamantanamine	Hydrogen	25	
2-Ethoxybenzylamine	N-Phenylethyldiamine	Hydrogen	25	
2-Ethoxybenzylamine	2,4-Dichlorophenethylamine	Hydrogen	25	
2-Ethoxybenzylamine	2-(2-Chlorophenyl)ethylamine	Hydrogen	25	3.89
2-Ethoxybenzylamine	Undecylamine	Hydrogen	25	
2-Ethoxybenzylamine	Dehydroabietylamine	Hydrogen	25	
cis-(-)-Myrtanylamine	2-(1-Cyclohexenyl)ethylamine	Hydrogen	25	
cis-(-)-Myrtanylamine	Hexetidine (mixture of isomers)	Hydrogen	25	
cis-(-)-Myrtanylamine	Aminodiphenylmethane	Hydrogen	25	
cis-(-)-Myrtanylamine	2,4-Dichlorophenethylamine	Hydrogen	25	
cis-(-)-Myrtanylamine	(S)-(-)-Cyclohexylethylamine	Hydrogen	25	28.94
cis-(-)-Myrtanylamine	Undecylamine	Hydrogen	25	
cis-(-)-Myrtanylamine	(+)-Isopinocampheylamine	Methyl	25	
cis-(-)-Myrtanylamine	Cyclooctylamine	Methyl	25	24.92
Cyclooctylamine	2,3-Dimethylcyclohexylamine	Hydrogen	25	50.55
Cyclooctylamine	(S)-2-Amino-1-butanol	Hydrogen	25	100.00

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
Cyclooctylamine	2-Adamantanamine, HCl	Hydrogen	25	29.61
Cyclooctylamine	4-Phenylbutylamine	Hydrogen	25	
Cyclooctylamine	2-Chlorobenzylamine	Hydrogen	25	
Cyclooctylamine	2-Aminoindan, HCl	Hydrogen	25	
Cyclooctylamine	Dehydroabietylamine	Hydrogen	25	
Cyclooctylamine	1-(1-Naphthyl)ethylamine	Hydrogen	25	4.62
Cyclooctylamine	1-Adamantanemethylamine	Methyl	25	14.20
2,3-Dimethoxybenzylamine	Hexetidine (mixture of isomers)	Hydrogen	25	
2,3-Dimethoxybenzylamine	Undecylamine	Hydrogen	25	
2,3-Dimethoxybenzylamine	Dehydroabietylamine	Hydrogen	25	
4-Methylcyclohexylamine	Hexetidine (mixture of isomers)	Hydrogen	25	
4-Methylcyclohexylamine	Undecylamine	Hydrogen	25	
4-Methylcyclohexylamine	Dehydroabietylamine	Hydrogen	25	
4-Fluorobenzylamine	Dibenzylamine	Hydrogen	25	27.98
trans-2-Phenylcyclopropylamine, HCl	Cyclooctylamine	Hydrogen	25	32.80
trans-2-Phenylcyclopropylamine, HCl	2-Adamantanamine, HCl	Hydrogen	25	18.99
trans-2-Phenylcyclopropylamine, HCl	1-Adamantanamine	Hydrogen	25	18.84
Thiomicamine	Hexetidine (mixture of isomers)	Hydrogen	25	
(R)-1-Amino-2-propanol	Hexetidine (mixture of isomers)	Hydrogen	25	
4-Chlorophenylalaninol	2,4-Dichlorophenethylamine	Hydrogen	25	
4-Chlorophenylalaninol	Undecylamine	Hydrogen	25	
4-Chlorophenylalaninol	Dehydroabietylamine	Hydrogen	25	
I-Leucinol	Hexetidine (mixture of isomers)	Hydrogen	25	
I-Leucinol	2,4-Dichlorophenethylamine	Hydrogen	25	
I-Leucinol	Dehydroabietylamine	Hydrogen	25	
(-)-Isopinocampheylamine	2-Methoxyphenethylamine	Hydrogen	25	29.59
(-)-Isopinocampheylamine	Undecylamine	Hydrogen	25	
Allylamine	Dehydroabietylamine	Hydrogen	25	
3-Amino-1,2-propanediol	Hexetidine (mixture of isomers)	Hydrogen	25	
3-Ethoxypropylamine	3,3-Diphenylpropylamine	Hydrogen	25	
3-Ethoxypropylamine	Undecylamine	Hydrogen	25	
3-Ethoxypropylamine	Dehydroabietylamine	Hydrogen	25	
sec-Butylamine	2,4-Dichlorophenethylamine	Hydrogen	25	
sec-Butylamine	Undecylamine	Hydrogen	25	
2-Aminoheptane	Hexetidine (mixture of isomers)	Hydrogen	25	
2-Aminoheptane	4-Phenylbutylamine	Hydrogen	25	
2-Aminoheptane	2,4-Dichlorophenethylamine	Hydrogen	25	
1-Naphthalenemethylamine	Hexetidine (mixture of isomers)	Hydrogen	25	
1-Naphthalenemethylamine	4-Phenylbutylamine	Hydrogen	25	
1-Naphthalenemethylamine	2,4-Dichlorophenethylamine	Hydrogen	25	
1-Naphthalenemethylamine	Undecylamine	Hydrogen	25	
Ethanolamine	Dehydroabietylamine	Hydrogen	25	
Piperonylamine	4-Phenylbutylamine	Hydrogen	25	

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
1-Ethylpropylamine	Hexetidine (mixture of isomers)	Hydrogen	25	
1-Ethylpropylamine	Dehydroabietylamine	Hydrogen	25	
Isopropylamine	Hexetidine (mixture of isomers)	Hydrogen	25	
4-Fluorophenethylamine	4-Phenylbutylamine	Hydrogen	25	
4-Fluorophenethylamine	2,4-Dichlorophenethylamine	Hydrogen	25	
4-Fluorophenethylamine	Dehydroabietylamine	Hydrogen	25	
3-Fluorophenethylamine	Undecylamine	Hydrogen	25	
2-Thiopheneethylamine	2-Adamantanamine, HCl	Hydrogen	25	19.09
2-Methylcyclohexylamine (mix of cis and trans)	Hexetidine (mixture of isomers)	Hydrogen	25	
2-Methylcyclohexylamine (mix of cis and trans)	Dehydroabietylamine	Hydrogen	25	
2-Methoxyphenethylamine	2-Adamantanamine, HCl	Hydrogen	25	26.77
2-Methoxyphenethylamine	(-)-Isopinocampheylamine	Hydrogen	25	31.95
2-Methoxyphenethylamine	1-Adamantanamine	Hydrogen	25	24.38
2-Methoxyphenethylamine	N-Allylcyclopentylamine	Hydrogen	25	14.56
2-Methoxyphenethylamine	4-Phenylbutylamine	Hydrogen	25	
2-Methoxyphenethylamine	Undecylamine	Hydrogen	25	
2-Methoxyphenethylamine	Dehydroabietylamine	Hydrogen	25	
2-Fluoroethylamine, HCl	Undecylamine	Hydrogen	25	
2-Fluoroethylamine, HCl	Dehydroabietylamine	Hydrogen	25	
2-Aminoindan, HCl	2-Adamantanamine, HCl	Hydrogen	25	17.72
2-Amino-1-phenylethanol	Undecylamine	Hydrogen	25	
2,5-Dimethoxyphenethylamine	(+)-Bornylamine	Hydrogen	25	25.78
2,5-Dimethoxyphenethylamine	Noradamantamine, HCl	Hydrogen	25	11.73
2,5-Dimethoxyphenethylamine	1-Adamantanamine	Hydrogen	25	12.57
2-(2-Chlorophenyl)ethylamine	4-Phenylbutylamine	Hydrogen	25	
2-(2-Chlorophenyl)ethylamine	Undecylamine	Hydrogen	25	
2-(2-Chlorophenyl)ethylamine	1-(1-Naphthyl)ethylamine	Hydrogen	25	
2-(2-Aminomethyl)phenylthio)benzyl alcohol	Hexetidine (mixture of isomers)	Hydrogen	25	
2-(2-Aminomethyl)phenylthio)benzyl alcohol	4-Phenylbutylamine	Hydrogen	25	
2-(2-Aminomethyl)phenylthio)benzyl alcohol	Undecylamine	Hydrogen	25	
1-Aminoindan	Hexetidine (mixture of isomers)	Hydrogen	25	
1-Aminoindan	Undecylamine	Hydrogen	25	
1-Aminoindan	Dehydroabietylamine	Hydrogen	25	
1,3-Dimethylbutylamine	Hexetidine (mixture of isomers)	Hydrogen	25	
1,3-Dimethylbutylamine	Undecylamine	Hydrogen	25	5.92
1,3-Dimethylbutylamine	Dehydroabietylamine	Hydrogen	25	
(S)-(-)-Cyclohexylethylamine	(-)-Isopinocampheylamine	Hydrogen	25	19.31
(S)-(-)-Cyclohexylethylamine	Hexetidine (mixture of isomers)	Hydrogen	25	

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
(S)-(-)-Cyclohexylethylamine	Undecylamine	Hydrogen	25	10.88
(S)-(-)-Cyclohexylethylamine	Dehydroabietylamine	Hydrogen	25	
(S)-(-)-2-Amino-3-phenyl-1-propanol	Hexetidine (mixture of isomers)	Hydrogen	25	
(S)-(-)-2-Amino-3-phenyl-1-propanol	Undecylamine	Hydrogen	25	
(S)-(-)-2-Amino-3-phenyl-1-propanol	Dehydroabietylamine	Hydrogen	25	
(1S,2S)-(+)-2-Amino-3-methoxy-1-phenyl-1-propanol	Hexetidine (mixture of isomers)	Hydrogen	25	
Octadecylamine	(+)-Bornylamine	Hydrogen	25	
Octadecylamine	1-Adamantanamine	Hydrogen	25	
Geranylamine	2,3-Dimethylcyclohexylamine	Hydrogen	25	14.53
Geranylamine	tert-Octylamine	Hydrogen	25	15.22
Geranylamine	1-Adamantanemethylamine	Hydrogen	25	4.37
Geranylamine	Decahydroquinoline	Hydrogen	25	31.79
Geranylamine	Dibenzylamine	Hydrogen	25	6.48
Geranylamine	N-Butylbenzylamine	Hydrogen	25	16.44
Geranylamine	Cyclooctylamine	Hydrogen	25	12.37
Geranylamine	(-)-Isopinocampheylamine	Hydrogen	25	8.95
Geranylamine	1-(1-Adamantyl)ethylamine, HCl	Hydrogen	25	32.95
Geranylamine	Undecylamine	Hydrogen	25	
Geranylamine	1-(1-Naphthyl)ethylamine	Hydrogen	25	
Amylamine	1-Adamantanamine	Hydrogen	37.5	0
3-Phenyl-1-propylamine	3,3-Diphenylpropylamine	Hydrogen	37.5	
3-Phenyl-1-propylamine	2,2-Diphenylamine	Hydrogen	37.5	
3-Phenyl-1-propylamine	1-Adamantanamine	Hydrogen	37.5	18.65
2,2-Diphenylamine	3,3-Diphenylpropylamine	Hydrogen	37.5	
2,2-Diphenylamine	2,2-Diphenylamine	Hydrogen	37.5	5.56
2,2-Diphenylamine	1,3,3-Trimethyl-6-azabicyclo[3.2.1]octane	Hydrogen	37.5	8.67
2,2-Diphenylamine	1-Adamantanamine	Hydrogen	37.5	58.10
4-(Trifluoromethyl)benzylamine	tert-Octylamine	Hydrogen	37.5	7.47
4-(Trifluoromethyl)benzylamine	138	Hydrogen	37.5	
4-Methylbenzylamine	2-Fluorobenzylamine	Hydrogen	50	22.10
4-Methylbenzylamine	4-Fluorobenzylamine	Hydrogen	50	14.62
4-Methylbenzylamine	alpha-Methyltryptamine	Hydrogen	50	0
4-Methylbenzylamine	Undecylamine	Hydrogen	50	0
Cyclopentylamine	Undecylamine	Hydrogen	50	0
Furfurylamine	2-Fluorobenzylamine	Hydrogen	50	0
Furfurylamine	Benzylamine	Hydrogen	50	0
Furfurylamine	4-Fluorobenzylamine	Hydrogen	50	0
Furfurylamine	alpha-Methyltryptamine	Hydrogen	50	0
Furfurylamine	Undecylamine	Hydrogen	50	0
Furfurylamine	Dehydroabietylamine	Hydrogen	50	0

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
Furfurylamine	Furfurylamine	Hydrogen	50	0
3,4,5-Trimethoxybenzylamine	2-Fluorobenzylamine	Hydrogen	50	0
3,4,5-Trimethoxybenzylamine	Benzylamine	Hydrogen	50	0
3,4,5-Trimethoxybenzylamine	alpha-Methyltryptamine	Hydrogen	50	0
3,4,5-Trimethoxybenzylamine	Undecylamine	Hydrogen	50	0
3,4,5-Trimethoxybenzylamine	Dehydroabietylamine	Hydrogen	50	0
1-Methyl-3-phenylpropylamine	alpha-Methyltryptamine	Hydrogen	50	0
1-Methyl-3-phenylpropylamine	Octadecylamine	Hydrogen	50	0
Cyclobutylamine	Octadecylamine	Hydrogen	50	0
Cyclobutylamine	Undecylamine	Hydrogen	50	0
Cyclobutylamine	Dehydroabietylamine	Hydrogen	50	0
1,2,3,4-Tetrahydro-1-naphthylamine	Hexetidine (mixture of isomers)	Hydrogen	50	0
1,2,3,4-Tetrahydro-1-naphthylamine	Aminodiphenylmethane	Hydrogen	50	4.31
1,2,3,4-Tetrahydro-1-naphthylamine	alpha-Methyltryptamine	Hydrogen	50	0
1,2,3,4-Tetrahydro-1-naphthylamine	2-Methoxyphenethylamine	Hydrogen	50	0
2,3-Dimethylcyclohexylamine	Hexetidine (mixture of isomers)	Hydrogen	50	0
2,3-Dimethylcyclohexylamine	Aminodiphenylmethane	Hydrogen	50	3.64
2,3-Dimethylcyclohexylamine	alpha-Methyltryptamine	Hydrogen	50	0
Tyramine	Furfurylamine	Hydrogen	50	0
Tyramine	2-Fluorobenzylamine	Hydrogen	50	4.07
Tyramine	Benzylamine	Hydrogen	50	0
Tyramine	2,4-Dichlorophenethylamine	Hydrogen	50	0
2-Fluorobenzylamine	Aminodiphenylmethane	Hydrogen	50	0
2-Fluorobenzylamine	4-Phenylbutylamine	Hydrogen	50	0
2-Fluorobenzylamine	2-Methoxyphenethylamine	Hydrogen	50	0
2-Fluorobenzylamine	2,4-Dichlorophenethylamine	Hydrogen	50	0
2-Fluorobenzylamine	1,3-Dimethylbutylamine	Hydrogen	50	0
2-Fluorobenzylamine	1-(1-Adamantyl)ethylamine, HCl	Hydrogen	50	0
(R)-2-Amino-1-butanol	Dehydroabietylamine	Hydrogen	50	0
3,4-Dimethoxyphenethylamine	Aminodiphenylmethane	Hydrogen	50	0
3,4-Dimethoxyphenethylamine	4-Phenylbutylamine	Hydrogen	50	0
3,4-Dimethoxyphenethylamine	2-Methoxyphenethylamine	Hydrogen	50	0
3,4-Dimethoxyphenethylamine	2,4-Dichlorophenethylamine	Hydrogen	50	0
3,4-Dimethoxyphenethylamine	1,3-Dimethylbutylamine	Hydrogen	50	0
3,3-Diphenylpropylamine	Piperidine	Hydrogen	50	0
3,3-Diphenylpropylamine	2,3-Dimethylcyclohexylamine	Methyl	50	7.81
3,3-Diphenylpropylamine	(-)-Isopinocampheylamine	Methyl	50	13.06
Propylamine	(S)-(+)-1-Amino-2-propanol	Hydrogen	50	0
Phenethylamine	(S)-(+)-1-Amino-2-propanol	Hydrogen	50	0
Phenethylamine	4-Phenylbutylamine	Hydrogen	50	0
Phenethylamine	2,4-Dichlorophenethylamine	Hydrogen	50	0

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
Phenethylamine	1,3-Dimethylbutylamine	Hydrogen	50	0
Phenethylamine	1-(1-Adamantyl)ethylamine, HCl	Hydrogen	50	0
Phenethylamine	1-(1-Naphthyl)ethylamine	Hydrogen	50	0
4-(2-Aminoethyl)morpholine	2-Amino-2-methyl-1-propanol	Hydrogen	50	0
Cyclohexylamine	2,4-Dichlorophenethylamine	Hydrogen	50	0
exo-Aminonorbornane	Benzylamine	Hydrogen	50	0
(+)-Isopinocampheylamine	Hexetidine (mixture of isomers)	Hydrogen	50	0
(+)-Isopinocampheylamine	Aminodiphenylmethane	Hydrogen	50	5.07
(+)-Isopinocampheylamine	4-Phenylbutylamine	Hydrogen	50	0
(+)-Isopinocampheylamine	2,4-Dichlorophenethylamine	Hydrogen	50	0
(+)-Isopinocampheylamine	Undecylamine	Hydrogen	50	0
Benzylamine	3,3-Diphenylpropylamine	Hydrogen	50	
Benzylamine	2-Amino-2-methyl-1-propanol	Hydrogen	50	
Benzylamine	1-(1-Naphthyl)ethylamine	Hydrogen	50	
Benzylamine	2,4-Dichlorophenethylamine	Hydrogen	50	
3-Amino-1-propanol	Undecylamine	Hydrogen	50	0
2-Fluorophenethylamine	3,3-Diphenylpropylamine	Hydrogen	50	0
2-Fluorophenethylamine	1-Adamantanemethylamine	Hydrogen	50	0
2-Fluorophenethylamine	1-(3-Aminopropyl)-2-pyrrolidinone (tech)	Hydrogen	50	0
2-Fluorophenethylamine	Decahydroquinoline	Hydrogen	50	0
2-Fluorophenethylamine	1-Adamantanamine	Hydrogen	50	24.34
2-Fluorophenethylamine	2,4-Dichlorophenethylamine	Hydrogen	50	0
2-Fluorophenethylamine	Undecylamine	Hydrogen	50	0
2-Fluorophenethylamine	Dehydroabietylamine	Hydrogen	50	0
2-Fluorophenethylamine	2-(1-Cyclohexenyl)ethylamine	Methyl	50	0
2-Fluorophenethylamine	Cyclooctylamine	Methyl	50	5.81
b-Methylphenethylamine	3,3-Diphenylpropylamine	Hydrogen	50	0
b-Methylphenethylamine	tert-Octylamine	Hydrogen	50	0
b-Methylphenethylamine	2-(1-Cyclohexenyl)ethylamine	Hydrogen	50	0
b-Methylphenethylamine	2-Amino-2-methyl-1-propanol	Hydrogen	50	0
b-Methylphenethylamine	4-Fluorophenethylamine	Hydrogen	50	0
b-Methylphenethylamine	Geranylamine	Hydrogen	50	0
b-Methylphenethylamine	5-Methoxytryptamine	Hydrogen	50	0
4-Methoxyphenethylamine	3,3-Diphenylpropylamine	Hydrogen	50	0
4-Methoxyphenethylamine	2-Amino-2-methyl-1-propanol	Hydrogen	50	0
4-Methoxyphenethylamine	2,4-Dichlorophenethylamine	Hydrogen	50	0
4-Methoxyphenethylamine	1-(1-Naphthyl)ethylamine	Hydrogen	50	0
L-Methioninol	Hexetidine (mixture of isomers)	Hydrogen	50	0
Tetrahydrofurfurylamine	1-Adamantanemethylamine	Hydrogen	50	0
Tetrahydrofurfurylamine	2-(1-Cyclohexenyl)ethylamine	Hydrogen	50	0
Tetrahydrofurfurylamine	4-Fluorophenethylamine	Hydrogen	50	0
Tetrahydrofurfurylamine	Undecylamine	Hydrogen	50	0
Amylamine	1-Adamantanemethylamine	Hydrogen	50	0
Amylamine	Hexetidine (mixture of isomers)	Hydrogen	50	0

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
Amylamine	Undecylamine	Hydrogen	50	0
Amylamine	Dehydroabietylamine	Hydrogen	50	0
1-Adamantanemethylamine	cis-(-)-Myrtanylamine	Methyl	50	0
3-Phenyl-1-propylamine	4-(2-Aminoethyl)morpholine	Hydrogen	50	
3-Phenyl-1-propylamine	1-(3-Aminopropyl)-2-pyrrolidinone (tech)	Hydrogen	50	
3-Phenyl-1-propylamine	Veratryl amine	Hydrogen	50	
3-Phenyl-1-propylamine	Aminodiphenylmethane	Hydrogen	50	
3-Phenyl-1-propylamine	2-(2-Aminomethyl)phenylthio)benzyl alcohol	Hydrogen	50	
2,2-Diphenylamine	2-Fluorophenethylamine	Hydrogen	50	
2,2-Diphenylamine	3,3-Diphenylpropylamine	Methyl	50	
2,2-Diphenylamine	(+)-Isopinocampheylamine	Methyl	50	
2,2-Diphenylamine	(+)-Bornylamine	Methyl	50	
2,2-Diphenylamine	Cyclooctylamine	Methyl	50	
2,2-Diphenylamine	(-)-Isopinocampheylamine	Methyl	50	3.81
4-(Trifluoromethyl)benzylamine	4-(2-Aminoethyl)morpholine	Hydrogen	50	
4-(Trifluoromethyl)benzylamine	2-(1-Cyclohexenyl)ethylamine	Hydrogen	50	
4-(Trifluoromethyl)benzylamine	Hexetidine (mixture of isomers)	Hydrogen	50	
4-(Trifluoromethyl)benzylamine	2,4-Dichlorophenethylamine	Hydrogen	50	
4-(Trifluoromethyl)benzylamine	(S)-(-)-Cyclohexylethylamine	Hydrogen	50	
Veratryl amine	1-Adamantanemethylamine	Hydrogen	50	
Veratryl amine	2-(1-Cyclohexenyl)ethylamine	Hydrogen	50	
Veratryl amine	4-Fluorophenethylamine	Hydrogen	50	
Veratryl amine	Hexetidine (mixture of isomers)	Hydrogen	50	
Veratryl amine	2,4-Dichlorophenethylamine	Hydrogen	50	
Veratryl amine	(S)-(-)-Cyclohexylethylamine	Hydrogen	50	
Veratryl amine	Undecylamine	Hydrogen	50	
Veratryl amine	Dehydroabietylamine	Hydrogen	50	
Veratryl amine	1-(1-Naphthyl)ethylamine	Hydrogen	50	
5-Amino-1-pentanol	1-Adamantanemethylamine	Hydrogen	50	
5-Amino-1-pentanol	Dibenzylamine	Hydrogen	50	
5-Amino-1-pentanol	cis-(-)-Myrtanylamine	Hydrogen	50	12.97
2-(1-Cyclohexenyl)ethylamine	2,4-Dimethoxybenzylamine	Hydrogen	50	
1-Aminomethyl-1-cyclohexanol, HCl	tert-Amylamine	Hydrogen	50	
1-Aminomethyl-1-cyclohexanol, HCl	2-(2-Aminomethyl)phenylthio)benzyl alcohol	Hydrogen	50	
1-Aminomethyl-1-cyclohexanol, HCl	Undecylamine	Hydrogen	50	
1-Aminomethyl-1-cyclohexanol, HCl	1-(1-Naphthyl)ethylamine	Hydrogen	50	
3-Fluorobenzylamine	tert-Amylamine	Hydrogen	50	
3-Fluorobenzylamine	Hexetidine (mixture of isomers)	Hydrogen	50	

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
3-Fluorobenzylamine	Undecylamine	Hydrogen	50	
4-Amino-1-butanol	Undecylamine	Hydrogen	50	
4-Amino-1-butanol	Dehydroabietylamine	Hydrogen	50	
2,4-Dimethoxybenzylamine	N-Phenylethyldiamine	Hydrogen	50	
2,4-Dimethoxybenzylamine	Aminodiphenylmethane	Hydrogen	50	
2,4-Dimethoxybenzylamine	4-Phenylbutylamine	Hydrogen	50	
2,4-Dimethoxybenzylamine	2-Chlorobenzylamine	Hydrogen	50	
2,4-Dimethoxybenzylamine	2,4-Dichlorophenethylamine	Hydrogen	50	
2,4-Dimethoxybenzylamine	2-(2-Chlorophenyl)ethylamine	Hydrogen	50	
2,4-Dimethoxybenzylamine	4-(Trifluoromethoxy)benzylamine	Hydrogen	50	
2-Ethoxybenzylamine	Aminodiphenylmethane	Hydrogen	50	
2-Ethoxybenzylamine	4-Phenylbutylamine	Hydrogen	50	
2-Ethoxybenzylamine	2-Chlorobenzylamine	Hydrogen	50	
2-Ethoxybenzylamine	2-Aminoindan, HCl	Hydrogen	50	
2-Ethoxybenzylamine	2,5-Dimethoxyphenethylamine	Hydrogen	50	
2-Ethoxybenzylamine	4-(Trifluoromethoxy)benzylamine	Hydrogen	50	
2-Ethoxybenzylamine	1-(1-Naphthyl)ethylamine	Hydrogen	50	
cis-(-)-Myrtanylamine	4-(2-Aminoethyl)morpholine	Hydrogen	50	
cis-(-)-Myrtanylamine	2-Fluorophenethylamine	Hydrogen	50	
cis-(-)-Myrtanylamine	1-(3-Aminopropyl)-2-pyrrolidinone (tech)	Hydrogen	50	
cis-(-)-Myrtanylamine	Veratryl amine	Hydrogen	50	
cis-(-)-Myrtanylamine	N-Butylbenzylamine	Hydrogen	50	
cis-(-)-Myrtanylamine	2,4-Dimethoxybenzylamine	Hydrogen	50	
cis-(-)-Myrtanylamine	1,2,3,4-Tetrahydropyridoindole	Hydrogen	50	
cis-(-)-Myrtanylamine	4-Phenylbutylamine	Hydrogen	50	
cis-(-)-Myrtanylamine	2-(2-Chlorophenyl)ethylamine	Hydrogen	50	3.91
cis-(-)-Myrtanylamine	1-(1-Adamantyl)ethylamine, HCl	Hydrogen	50	10.85
cis-(-)-Myrtanylamine	(R)-(-)-Cyclohexylethylamine	Hydrogen	50	5.89
cis-(-)-Myrtanylamine	Dehydroabietylamine	Hydrogen	50	
cis-(-)-Myrtanylamine	1-(1-Naphthyl)ethylamine	Hydrogen	50	
cis-(-)-Myrtanylamine	(+)-Bornylamine	Methyl	50	4.04
Cyclooctylamine	4-Methylcyclohexylamine	Hydrogen	50	4.55
Cyclooctylamine	N-Phenylethyldiamine	Hydrogen	50	
Cyclooctylamine	4-(Hexacylamino)benzylamine	Hydrogen	50	
Cyclooctylamine	2,5-Dimethoxyphenethylamine	Hydrogen	50	
Cyclooctylamine	2,4-Dichlorophenethylamine	Hydrogen	50	3.36
Cyclooctylamine	2-(2-Chlorophenyl)ethylamine	Hydrogen	50	9.15
Cyclooctylamine	1-(1-Adamantyl)ethylamine, HCl	Hydrogen	50	10.62
Cyclooctylamine	(S)-(-)-Cyclohexylethylamine	Hydrogen	50	5.85
Cyclooctylamine	(R)-(-)-Cyclohexylethylamine	Hydrogen	50	
Cyclooctylamine	4-	Hydrogen	50	4.54

Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
	(Trifluoromethoxy)benzylamine			
2-Adamantanamine, HCl	cis-(-)-Myrtanylamine	Hydrogen	50	49.73
4-Methylcyclohexylamine	N-Phenylethyldiamine	Hydrogen	50	
4-Methylcyclohexylamine	4-Phenylbutylamine	Hydrogen	50	
4-Fluorobenzylamine	N-Benzyl-2-phenethylamine	Hydrogen	50	
4-Fluorobenzylamine	Hexetidine (mixture of isomers)	Hydrogen	50	
4-Fluorobenzylamine	Undecylamine	Hydrogen	50	
4-Fluorobenzylamine	Dehydroabietylamine	Hydrogen	50	
trans-2-Phenylcyclopropylamine, HCl	Hexetidine (mixture of isomers)	Hydrogen	50	
trans-2-Phenylcyclopropylamine, HCl	Undecylamine	Hydrogen	50	
trans-2-Phenylcyclopropylamine, HCl	Dehydroabietylamine	Hydrogen	50	
(R)-1-Amino-2-propanol	4-(Hexacylamino)benzylamine	Hydrogen	50	
(R)-1-Amino-2-propanol	Undecylamine	Hydrogen	50	
(R)-1-Amino-2-propanol	Dehydroabietylamine	Hydrogen	50	
l-Leucinol	Undecylamine	Hydrogen	50	
(-)-Isopinocampheylamine	2-Ethoxybenzylamine	Hydrogen	50	27.27
(-)-Isopinocampheylamine	Hexetidine (mixture of isomers)	Hydrogen	50	
(-)-Isopinocampheylamine	4-Phenylbutylamine	Hydrogen	50	
(-)-Isopinocampheylamine	Dehydroabietylamine	Hydrogen	50	
(-)-Isopinocampheylamine	1-(1-Naphthyl)ethylamine	Hydrogen	50	
Allylamine	3,3-Diphenylpropylamine	Hydrogen	50	
Allylamine	2-Amino-1-propanol, d,l	Hydrogen	50	
Allylamine	Undecylamine	Hydrogen	50	
3-Amino-1,2-propanediol	Dehydroabietylamine	Hydrogen	50	
3-Ethoxypropylamine	2,2-Diphenylamine	Hydrogen	50	95.81
3-Ethoxypropylamine	cis-(-)-Myrtanylamine	Hydrogen	50	
2-Aminoheptane	2-(2-Aminomethyl)phenylthio)benzyl alcohol	Hydrogen	50	
1-Naphthalenemethylamine	Geranylamine	Hydrogen	50	
1-Naphthalenemethylamine	Dehydroabietylamine	Hydrogen	50	
1-Aminopyrrolidine, HCl	Hexetidine (mixture of isomers)	Hydrogen	50	
1-Aminopyrrolidine, HCl	Undecylamine	Hydrogen	50	
1-Aminopyrrolidine, HCl	Dehydroabietylamine	Hydrogen	50	
Ethanolamine	3,3-Diphenylpropylamine	Hydrogen	50	
3-Methylbenzylamine	Geranylamine	Hydrogen	50	
3-Methylbenzylamine	5-Methoxytryptamine	Hydrogen	50	
Piperonylamine	Aminodiphenylmethane	Hydrogen	50	
Piperonylamine	2,4-Dichlorophenethylamine	Hydrogen	50	
Piperonylamine	2-(2-Aminomethyl)phenylthio)benzyl alcohol	Hydrogen	50	
Isopropylamine	Dehydroabietylamine	Hydrogen	50	







Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
4-Fluorophenethylamine	2,4-Dimethoxybenzylamine	Hydrogen	50	
4-Fluorophenethylamine	Aminodiphenylmethane	Hydrogen	50	
4-Fluorophenethylamine	2-(2-Aminomethyl)phenylthio)benzyl alcohol	Hydrogen	50	
4-Chloroamphetamine, HCl	N-Allylcyclopentylamine	Hydrogen	50	10.25
4-Chloroamphetamine, HCl	Hexetidine (mixture of isomers)	Hydrogen	50	
4-Chloroamphetamine, HCl	4-Phenylbutylamine	Hydrogen	50	
4-Chloroamphetamine, HCl	2-Methoxyphenethylamine	Hydrogen	50	
4-Chloroamphetamine, HCl	Undecylamine	Hydrogen	50	
4-Chloroamphetamine, HCl	Dehydroabietylamine	Hydrogen	50	
3-Fluorophenethylamine	(-)-Isopinocampheylamine	Hydrogen	50	
3-Fluorophenethylamine	1-Adamantamine	Hydrogen	50	8.59
3-Fluorophenethylamine	4-Phenylbutylamine	Hydrogen	50	
2-Methylcyclohexylamine (mix of cis and trans)	Undecylamine	Hydrogen	50	
2-Methoxyphenethylamine	3,3-Diphenylpropylamine	Hydrogen	50	
2-Methoxyphenethylamine	(+)-Bornylamine	Hydrogen	50	
2-Methoxyphenethylamine	tert-Octylamine	Hydrogen	50	20.46
2-Methoxyphenethylamine	1-Adamantanemethylamine	Hydrogen	50	
2-Methoxyphenethylamine	Dibenzylamine	Hydrogen	50	
2-Methoxyphenethylamine	N-Butylbenzylamine	Hydrogen	50	5.20
2-Methoxyphenethylamine	1,3,3-Trimethyl-6-azabicyclo[3.2.1]octane	Hydrogen	50	8.59
2-Methoxyphenethylamine	N-Phenylethyldiamine	Hydrogen	50	
2-Methoxyphenethylamine	2,4-Dichlorophenethylamine	Hydrogen	50	
2-Methoxyphenethylamine	2-(2-Chlorophenyl)ethylamine	Hydrogen	50	
2-Methoxyphenethylamine	1-(1-Adamantyl)ethylamine, HCl	Hydrogen	50	3.61
2-Aminoindan, HCl	(+)-Bornylamine	Hydrogen	50	
2-Aminoindan, HCl	Noradamantamine, HCl	Hydrogen	50	7.43
2-(2-Chlorophenyl)ethylamine	N-Phenylethyldiamine	Hydrogen	50	
2-(2-Chlorophenyl)ethylamine	Aminodiphenylmethane	Hydrogen	50	
2-(2-Chlorophenyl)ethylamine	2,4-Dichlorophenethylamine	Hydrogen	50	
2-(2-Chlorophenyl)ethylamine	1-(1-Adamantyl)ethylamine, HCl	Hydrogen	50	
2-(2-Chlorophenyl)ethylamine	Dehydroabietylamine	Hydrogen	50	
2-(2-Aminomethyl)phenylthio)benzyl alcohol	2-Methoxyphenethylamine	Hydrogen	50	
2-(2-Aminomethyl)phenylthio)benzyl alcohol	2,5-Dimethoxyphenethylamine	Hydrogen	50	
2-(2-Aminomethyl)phenylthio)benzyl alcohol	2-(2-Chlorophenyl)ethylamine	Hydrogen	50	
2-(2-	1-(1-Adamantyl)ethylamine,	Hydrogen	50	






Synthetic Substituted Diethylene Diamines Sorted by Minimum Inhibition Concentration				
N1	N2	R4	MIC (uM)	% Induction
Aminomethyl)phenylthio)benzyl alcohol	HCl			
2-(2-Aminomethyl)phenylthio)benzyl alcohol	Dehydroabietylamine	Hydrogen	50	
1-Aminoindan	4-Phenylbutylamine	Hydrogen	50	
1-Aminoindan	2,4-Dichlorophenethylamine	Hydrogen	50	
1,3-Dimethylbutylamine	4-Phenylbutylamine	Hydrogen	50	
(S)-(-)-Cyclohexylethylamine	Aminodiphenylmethane	Hydrogen	50	
(S)-(-)-Cyclohexylethylamine	4-Phenylbutylamine	Hydrogen	50	
(S)-(-)-Cyclohexylethylamine	2,4-Dichlorophenethylamine	Hydrogen	50	
(S)-(-)-Cyclohexylethylamine	1-(1-Adamantyl)ethylamine, HCl	Hydrogen	50	
(1S,2S)-(+)-2-Amino-3-methoxy-1-phenyl-1-propanol	Dehydroabietylamine	Hydrogen	50	
Octadecylamine	2-Adamantanamine, HCl	Hydrogen	50	
3-Hydroxytyramine	(1R,2S)-(-)-2-Amino-1,2-diphenylethanol	Hydrogen	50	
3-Hydroxytyramine	Dehydroabietylamine	Hydrogen	50	
Geranylamine	3,3-Diphenylpropylamine	Hydrogen	50	
Geranylamine	N-Phenylethyldiamine	Hydrogen	50	
Geranylamine	Hexetidine (mixture of isomers)	Hydrogen	50	
Geranylamine	2-Thiopheneethylamine	Hydrogen	50	
Geranylamine	2-Methoxyphenethylamine	Hydrogen	50	
Geranylamine	2,5-Dimethoxyphenethylamine	Hydrogen	50	
Geranylamine	2,4-Dichlorophenethylamine	Hydrogen	50	
Geranylamine	2-(2-Chlorophenyl)ethylamine	Hydrogen	50	
2-Fluorophenethylamine	2,3-Dimethylcyclohexylamine	Methyl	>50	2.07
4-(Trifluoromethyl)benzylamine	2,3-Dimethylcyclohexylamine	Hydrogen	>50	8.20
4-(Trifluoromethyl)benzylamine	1-Adamantanamine	Hydrogen	>50	32.02
5-Aminoquinoline	exo-Aminonorbornane	Hydrogen	>50	17.87







Table 3






Compounds Synthesized in Larger Quantities for Further
in vitro Evaluations






5

Cmpd #	Name	Structure	mount, mg	Yields, %
1	<i>N</i> -(4-Methylphenyl)- <i>N'</i> -(furfuryl)ethane-1,2-diamine		23	25
2	<i>N</i> -(4-Methylphenyl)- <i>N'</i> -(benzyl)ethane-1,2-diamine		27	29
3	<i>N</i> -[1-(1,2,3,4-Tetrahydronaphthalene)- <i>N'</i> -(undecenyl)-ethane-1,2-diamine		11	10
4	<i>N</i> -[2-(3,4-Dimethoxyphenyl)-ethyl- <i>N'</i> -(1-methyladamantyl)-ethane-1,2-diamine		13	11
5	<i>N</i> -[2-(3,4-Dimethoxyphenyl)-ethyl- <i>N'</i> -(norbornyl)-ethane-1,2-diamine		9	8
6	<i>N</i> -(1-Adamantylmethyl)- <i>N'</i> -(3,3-diphenylpropyl)propane-1,2-diamine		55	36






7	<i>N</i> -(1-Adamantylmethyl)- <i>N'</i> -(3,3-diphenylpropyl)ethane-1,2-diamine		28	22
8	<i>N</i> -[2-(Cyclohexen-1-yl)ethyl]- <i>N'</i> -(3,3-diphenylpropyl)propane-1,2-diamine		46	37
10	<i>N</i> -(-)- <i>cis</i> -Myrtanyl- <i>N'</i> -(3,3-diphenylpropyl)ethane-1,2-diamine		14	11
11	<i>N</i> -Cyclooctyl- <i>N'</i> -(3,3-diphenylpropyl)ethane-1,2-diamine		22	18
13	<i>N</i> -Allyl- <i>N</i> -cyclopentyl- <i>N'</i> -(3,3-diphenylpropyl)ethane-1,2-diamine		33	27

14	<i>N</i> -(3,3-Diphenylpropyl)- <i>N'</i> - <i>exo</i> -(2-norborny)ethane-1,2-diamine		17	16
15	1-{2-[<i>N</i> -(3,3-Diphenylpropyl)]-aminoethyl}-3,5-dimethyl-piperidine		6.2	5
17	<i>N</i> -2-(2-Methoxyphenyl)ethyl- <i>N'</i> -(3,3-diphenylethyl)ethane-1,2-diamine		50	40
21	<i>N</i> -(3,3-Diphenylpropyl)- <i>N'</i> -(1 <i>S</i>)-(1-ethylcyclohexane)-ethane-1,2-diamine		5	4
22	<i>N</i> -(3,3-Diphenylpropyl)- <i>N'</i> -(1 <i>R</i>)-(1-ethylcyclohexane)-ethane-1,2-diamine		21	17
23	<i>N</i> -Allyl- <i>N</i> -cyclohexyl- <i>N'</i> -(3,3-diphenylpropyl)ethane-1,2-diamine		6	5

24	<i>N</i> -2-(2-Methoxyphenyl)ethyl- <i>N'</i> -(4-fluorophenylethyl)-ethane-1,2-diamine		10	9
27	<i>N</i> -(3-Phenylpropyl)- <i>N'</i> -(1-adamantyl)ethane-1,2-diamine		11	10
28	<i>N</i> -(3-Phenylpropyl)- <i>N'</i> -(4-fluorophenyl)ethane-1,2-diamine		11	10
29	<i>N</i> -(2,2-Diphenylethyl)- <i>N'</i> -(2,3-dimethylcyclohexyl)ethane-1,2-diamine		4.5	4
31	<i>N</i> -(2,2-Diphenylethyl)- <i>N'</i> -(1 <i>S</i>)-(1-ethylcyclohexane)-ethane-1,2-diamine		24	20






32	<i>N</i> -(2,2-Diphenylethyl)- <i>N'</i> -(<i>R</i>)-(+)-		58	48
33	<i>N</i> -(2,2-Diphenylethyl)- <i>N'</i> -(1,1,3,3-tetramethylbutyl)-ethane-1,2-diamine		11	9
34	<i>N</i> -(2,2-Diphenylethyl)- <i>N'</i> -(1-methyladamantyl)ethane-1,2-diamine		6.8	6
35	<i>N</i> -(2,2-Diphenylethyl)- <i>N'</i> -(1,1,3-trimethyl-6-azabicyclo-[3.2.1]octyl)ethane-1,2-diamine		38	30
36	<i>N</i> -(2-[<i>N'</i> -(2,2-Diphenylethyl)]-aminoethyl)-decahydroquinoline		28	24






37	<i>N</i> -(2,2-Diphenylethyl)- <i>N'</i> -(-)- <i>cis</i> -(myrtanyl)ethane-1,2-diamine		54	38
38	<i>N</i> -(-)- <i>cis</i> -(Myrtanyl)- <i>N'</i> -(2,2-diphenylethyl)propyl-1,2-diamine		39	30
40	<i>N</i> -(2,2-Diphenylethyl)- <i>N'</i> -(1 <i>R</i> , 2 <i>R</i> , 3 <i>R</i> , 5 <i>S</i>)-(-)-isopinocampheylethane-1,2-diamine		33	23
41	<i>N</i> -(-)- <i>cis</i> -(Myrtanyl)- <i>N'</i> -(2,3-dimethylcyclohexyl)ethane-1,2-diamine		66	62
42	<i>N</i> -(3,3-Diphenylpropyl)- <i>N'</i> -(-)- <i>cis</i> -myrtanylethane-1,2-diamine		11	9







43	<i>N</i> -(<i>-</i>)- <i>cis</i> -Myrtanyl- <i>N'</i> -(1 <i>S</i> , 2 <i>S</i> , 3 <i>S</i> , 5 <i>R</i>)-(+)-isopinocampheylethane-1,2-diamine		31	27
47	<i>N</i> -(<i>-</i>)- <i>cis</i> -Myrtanyl- <i>N'</i> -(1 <i>R</i> , 2 <i>R</i> , 3 <i>R</i> , 5 <i>S</i>)-(-)-isopinocampheylethane-1,2-diamine		42	33
51	<i>N</i> -(Cyclooctyl)- <i>N'</i> -(2,3-dimethylcyclohexyl)ethane - 1,2-diamine		5.1	2
52	<i>N</i> -(Cyclooctyl)- <i>N'</i> -(3,3-diphenylpropyl)ethane - 1,2-diamine		20	18
53	<i>N</i> -Cyclooctyl- <i>N'</i> -(1 <i>S</i> , 2 <i>S</i> , 3 <i>S</i> , 5 <i>R</i>)-(+)-isopinocampheyl-ethane-1,2-diamine		7.4	7






54	<i>N</i> -(Cyclooctyl)- <i>N'</i> -(<i>R</i>)-(+)-bornylethane-1,2-diamine		17	16
55	<i>N</i> -(Cyclooctyl)- <i>N'</i> -(1-methyladamantyl)ethane-1,2-diamine		7	6
56	<i>N</i> -(Cyclooctyl)- <i>N'</i> -(2 <i>S</i>)-[2-(1-hydroxybutyl)]ethane-1,2-diamine		1.1	1
57	<i>N</i> -(-)- <i>cis</i> -Myrtanyl- <i>N'</i> -(cyclooctyl)ethane-1,2-diamine		18	18
58	<i>N</i> -(Cyclooctyl)- <i>N'</i> -(2-adamantyl)ethane-1,2-diamine		25	23


59	<i>N</i> -(Cyclooctyl)- <i>N'</i> -(1 <i>R</i> , 2 <i>R</i> , 3 <i>R</i> , 5 <i>S</i>)-(-)-isopinocampheylethane-1,2-diamine		15	14
61	<i>N</i> -(Cyclooctyl)- <i>N'</i> -[1-ethyl-(1-naphthyl)]ethane-1,2-diamine		16	14
62	<i>N</i> -(-)- <i>cis</i> -Myrtanyl- <i>N'</i> -(1 <i>S</i>)-(1-ethylcyclohexane)ethane-1,2-diamine		48	46
63	<i>N</i> -(Cyclooctyl)- <i>N'</i> - <i>trans</i> -(2-phenylcyclopropyl)ethane-1,2-diamine		47	46
64	<i>N</i> -(2-Adamantyl)- <i>N'</i> - <i>trans</i> -(2-phenylcyclopropyl)ethane-1,2-diamine		49	46

65	<i>N</i> -(1-Adamantyl)- <i>N'</i> - <i>trans</i> -(2-phenylcyclopropyl)ethane-1,2-diamine		18	16
66	<i>N</i> -(3,3-Diphenylpropyl)- <i>N'</i> -(1 <i>R</i> , 2 <i>R</i> , 3 <i>R</i> , 5 <i>S</i>)-(-)-isopinocampheylethane-1,2-diamine		2.3	2
68	<i>N</i> -(+/-)-[2-(1-Hydroxybutyl)]- <i>N'</i> -(1 <i>R</i> , 2 <i>R</i> , 3 <i>R</i> , 5 <i>S</i>)-(-)-isopinocampheylethane-1,2-diamine		0.8	1
71	<i>N</i> -(1,1-Diphenylmethyl)- <i>N'</i> -(1 <i>R</i> , 2 <i>R</i> , 3 <i>R</i> , 5 <i>S</i>)-(-)-isopinocampheylethane-1,2-diamine		2.9	2
73	<i>N</i> -(2-Adamantyl)- <i>N'</i> -[2-(2-methoxyphenyl)ethyl]ethane-1,2-diamine		21	19

76	<i>N</i> -Allyl- <i>N</i> -cyclopentyl- <i>N</i> '-[2-(2-methoxyphenyl)ethyl]ethane-1,2-diamine		8	7
77	<i>N</i> -(1,1-Diphenylmethyl)- <i>N</i> '-[2-(2-methoxyphenyl)ethyl]ethane-1,2-diamine		32	27
78	<i>N</i> -2-Adamantyl- <i>N</i> '-2,3-dihydro-1 <i>H</i> -inden-2-yl-ethane-1,2-diamine		4.3	3
79	<i>N</i> -[2-(2,5-Dimethoxyphenyl)ethyl]- <i>N</i> '-(<i>R</i>)-(+)-bornylethane-1,2-diamine		59	49
103	<i>N,N</i> '-Bis(cyclooctyl)ethane-1,2-diamine		6.3	4

107	<i>N</i> -(2,2-Diphenylethyl)- <i>N</i> -(3-ethoxypropyl)ethane-1,2-diamine		58	52
109	<i>N</i> -Geranyl- <i>N'</i> -(2-adamantyl)ethane-1,2-diamine		27	24
111	<i>N</i> -[2-(<i>N'</i> -Geranyl)aminoethyl]-2-ethylpiperidine		24	24
116	<i>N</i> -Geranyl- <i>N'</i> -allyl- <i>N'</i> -(cyclopentyl)ethane-1,2-diamine		45	42
117	<i>N</i> -Geranyl- <i>N'</i> -(1,1-diphenylmethyl)ethane-1,2-diamine		24	20
118	<i>N</i> -2-(2-Chlorophenyl)ethyl- <i>N'</i> -allyl- <i>N'</i> -(cyclopentyl)ethane-1,2-diamine		6.4	6

119	<i>N</i> -2-(2-Chlorophenyl)ethyl- <i>N'</i> -[2-(3-fluorophenyl)- ethyl]ethane-1,2-diamine		30	27
125	<i>N,N'</i> -bis-(-)- <i>cis</i> - Myrtanylpropane-1,2-diamine		41	35
134	<i>N</i> -[2-(<i>N'</i> -2,2-Diphenylethyl)- aminoethyl]-(-)-3,4- dihydroxynorephedrine		20	15
151	<i>N</i> -[2-(2-Methoxy)phenylethyl]- <i>N'</i> -(1 <i>R</i> , 2 <i>R</i> , 3 <i>R</i> , 5 <i>S</i>)-(-)- isopinocampheyl-ethane-1,2- diamine		67	60
164	<i>N</i> ¹ -[2-(4-fluorophenyl)ethyl]- <i>N</i> ² -[2-(4-Methoxy) phenylethyl]-1-phenylethane- 1,2-diamine		94	73

165	N1-[2-(4-fluorophenyl)ethyl]- N2-(3-Phenylpropyl)-1- phenylethane-1,2-diamine		23	19
-----	---	--	----	----

The present invention is also directed to a new library of diamine compounds useful against infectious disease. To further enhance the structural diversity of prior diamine compounds, a synthetic scheme to incorporate amino acids into a bridging linker between the two amine components has been developed. The use of amino acids allowed for diverse linker elements, as well as chirality see Figure 42 for representative examples. The diamine compounds were prepared on mmol scale in 96-well format in pools of 10 compounds per well (for the vast majority of the plates). Table 25 (Figure 43) summarizes data for the synthesized plates.

The reaction scheme followed is shown in Figure 44.

Solid phase syntheses using Rink resin. Twenty one 96-well plates have been prepared. Six-step synthetic route starting from the Rink resin similar to what that had been used to create our first 100,000 compound library (Scheme 1, Figure 41), was applied to make targeted diamines (Scheme 5, Figure 44). Overall, all steps of these schemes are similar, except one (step 4) when formation of the second amino functionality occurs. In Scheme 1, the second amine is introduced into the molecule as a whole synthon via nucleophilic displacement of Cl-function of the linker, while in the Scheme 5, it proceeds through modification of the existing amino moiety by carbonyl compounds.

Attachment of the first amine to the support was done according to the Garigipati protocol. Rink acid resin (Novabiochem) was converted into the Rink-chloride upon treatment with triphenylphosphine and dichloroethane in THF. This activated resin was then loaded by addition of an amine N1 in presence of Hunig's base in dichloroethane. The amine N1 includes, but is not limited to, alkyl and aryl primary amines. Out of 177 primary amines that had been previously used as N1 for 100,000 library preparation, only 30 were selected in this Scheme, based upon *in vitro* activity data of their ethylenediamine derivatives (from the previous ~100K library) as well as structural diversity (Figures 45 and 46).

On the next step, the acylation reaction was accomplished *via* peptide coupling with Fmoc protected amino acids in presence of HATU (O-(7-Azabenzotriazol-1-yl)-N,N,N,N-tetramethyluronium hexafluorophosphate) and EtN(*iso*-Pr)₂ in DCM/DMF mixture at room temperature. The reaction was done twice to improve product yields. The list of the amino acids used to create this library is shown in the Table 26 (Figure 47).

Deprotection (removal of the Fmoc group) was carried out by reaction with piperidine at room temperature. Derivatization of the amino group was achieved by reductive alkylation with various carbonyl compounds, such as aldehydes, ketones, and carboxylic acids, in the presence of NaBCNH₃ at room temperature for 72-96 h. The selection of the carbonyl compounds was made so that the final diamine products would carry the same or similar types of substituents that had been observed in the hit compounds generated from the previous library of ethambutol analogs, as well as structural diversity (Figure 48). A complete list of the carbonyl compounds used is shown in Table 27 (Figure 49).

Reduction of the aminoethyleneamides into corresponding diamines was carried out using the soluble reducing reagent 65+w% Red-Al at room temperature. Cleavage of the products from the resin was achieved with a 10% solution of trifluoroacetic acid in dichloromethane resulting in the formation of TFA salts of the diamines.

For library production the first three steps of the synthetic scheme (resin activation, amine loading, and acylation) were carried out using a Quest 210 Synthesizer on scale of 0.1-0.15 g of resin per tube. Following the acylation, formed resins were thoroughly washed, dried, and then groups of ten resins were pooled together. A small amount of each resin (~0.05g) was archived prior to pooling to facilitate re-synthesis and deconvolution of actives.

Deprotection of the Fmoc group, addition of the carbonyl component, reduction, and cleavage were carried out in 96-well reaction blocks using the Combiclamps system by Whatman Polyfiltronics or the FlexChem system by Robbins Scientific. A suspension of the pooled resins in 2:1 mixture of DCM/THF was evenly distributed into one reaction plate resulting in approximately 10 mg of the resin per well. The 96 diverse carbonyl compounds were arrayed in one 96-well plate template and added, one carbonyl compound per well, to each individual pool of ten resins, resulting in an anticipated 960 diamines produced per plate. Reduction was carried out in the same format and cleavage and

filtering into storage plates was followed by evaporation of the TFA prior to biological assay.

Quality assessment of the prepared compounds was done by Electrospray Ionization mass spectrometry using two randomly selected rows (16 samples) per plate, 17% of the total number. Successful production of a compound was based on an appearance of a molecular ion of the calculated mass. Depending on the amino acid that had been used for the synthesis, the percentage of the predicted ions were observed, and therefore the predicted compounds were formed, varied from 5-60% (Table 25, Figure 43). Based on MS analysis, out of targeted 20,000 compounds, 4,500 diamines were actually formed.

10

Formulations

Therapeutics, including compositions containing the substituted ethylene diamine compounds of the present invention, can be prepared in physiologically acceptable formulations, such as in pharmaceutically acceptable carriers, using known techniques. For example, a substituted ethylene diamine compound is combined with a pharmaceutically acceptable excipient to form a therapeutic composition.

The compositions of the present invention may be administered in the form of a solid, liquid or aerosol. Examples of solid compositions include pills, creams, soaps and implantable dosage units. Pills may be administered orally. Therapeutic creams and anti-mycobacteria soaps may be administered topically. Implantable dosage units may be administered locally, for example, in the lungs, or may be implanted for systematic release of the therapeutic composition, for example, subcutaneously. Examples of liquid compositions include formulations adapted for injection intramuscularly, subcutaneously, intravenously, intraarterially, and formulations for topical and intraocular administration. Examples of aerosol formulations include inhaler formulations for administration to the lungs.

A sustained release matrix, as used herein, is a matrix made of materials, usually polymers, which are degradable by enzymatic or acid/base hydrolysis, or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. The sustained release matrix is chosen desirably from biocompatible materials, including, but not limited to, liposomes, polylactides, polyglycolide (polymer of glycolic acid),

polylactide co-glycolide (copolymers of lactic acid and glycolic acid), polyanhydrides, poly(ortho)esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, 5 polyvinylpyrrolidone and silicone. A preferred biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide.

The dosage of the composition will depend on the condition being treated, the particular composition used, and other clinical factors, such as weight and condition of the patient, and the route of administration. A suitable dosage may range from 100 to 0.1 10 mg/kg. A more preferred dosage may range from 50 to 0.2 mg/kg. A more preferred dosage may range from 25 to 0.5 mg/kg. Tablets or other forms of media may contain from 1 to 1000 mg of the substituted ethylene diamine. Dosage ranges and schedules of administration similar to ethambutol or other anti-tuberculosis drugs may be used.

The composition may be administered in combination with other compositions and 15 procedures for the treatment of other disorders occurring in combination with mycobacterial disease. For example, tuberculosis frequently occurs as a secondary complication associated with acquired immunodeficiency syndrome (AIDS). Patients undergoing AIDS treatment, which includes procedures such as surgery, radiation or chemotherapy, may benefit from the therapeutic methods and compositions described 20 herein.

The following specific examples will illustrate the invention as it applies to the particular synthesis of the substituted ethylene diamine compounds, and the *in vitro* and *in vivo* suppression of the growth of colonies of *M. tuberculosis*. In addition, the teachings of R. Lee et al. J. Comb. Chem 2003, 5, 172-187 are hereby incorporated by reference in 25 their entirety. It will be appreciated that other examples, including minor variations in chemical procedures, will be apparent to those skilled in the art, and that the invention is not limited to these specific illustrated examples.

EXAMPLE I

Generating the Ethylene Diamine Library

30 The Rink-acid resin was obtained from NOVABIOCHEM® Inc., San Diego, California. Solvents: acetonitrile, dichloromethane, dimethylformamide,

ethylenedichloride, methanol and tetrahydrofuran were purchased from ALDRICH®, Milwaukee, Wisconsin, and used as received. All other reagents were purchased from SIGMA-ALDRICH®, West Monroe Highland, Illinois. Solid phase syntheses were performed on a QUEST® 210 Synthesizer, from ARGONAUT TECHNOLOGIES®, Foster City, California, with the aid of combinatorial chemistry equipment, from WHATMAN® POLYFILTRONICS® (Kent, England; Rockland, Massachusetts) and ROBBINS SCIENTIFIC®, Sunnyvale, California. Evaporation of solvents was done using SPEEDVAC® AES, from SAVANT®, Holbrook, New York. All necessary chromatographic separations were performed on a WATERS' ALLIANCE HT SYSTEM®, Milford, Massachusetts. Analytical thin-layer chromatography was performed on MERCK® silica gel 60F₂₅₄ plates, purchased from SIGMA-ALDRICH®, West Monroe Highland, Illinois.

The activation of the Rink-acid resin, the addition of the first amine, and the acylation step were carried out in 10ml tubes using the QUEST® 210 Synthesizer. The addition of the second amine, the reduction with Red-AL, and the cleavage from the solid support were carried out in 96-deep (2 ml) well, chemically resistant plates.

A. Activation of the Rink-Acid Resin

The Rink-acid resin had a coverage of 0.43-0.63 mmol of linker per gram resin. Four to five grams of this resin were suspended in 80 ml of a 2:1 mixture of dichloromethane and tetrahydrofuran (THF), and distributed into ten, 10 ml tubes, with 8ml of resin suspension per tube. Each suspension was filtered and washed twice with THF. A solution of triphenylphosphine (3.80 g, 14.5 mmol) in 30 ml of THF was prepared, and 3ml of this solution was added to each tube, followed by the addition of 3 ml of a solution of hexachloroethane in THF (3.39g/14.3mmol hexachloroethane in 30 ml THF). After agitation for six hours at room temperature, each activated resin was washed twice with THF and twice with dichloromethane.

B. Addition of the First Amine

Each tube, containing the activated rink resin, was charged with 3 ml of dichloroethane, 0.3 ml (1.74 mmol) N₁N-diisopropylethylamine (EtN(iPr)₂) and the corresponding amine (around 1 mmol). If the selected amine was a solid at room temperature, it was added as a solution, or a suspension in DMF. Enough dichloroethane was added to each tube for a final volume of 8 ml. The reaction mixture was heated at

45°C for 6-8 hours. The resins were filtered, washed with a 2:1 mixture of dichloromethane and methanol (1x8 ml), then with methanol (2x8 ml), and then dried under argon for 10 minutes.

C. Acylation with the Halo-Acylchloride

- 5 a. *Acylation with Chloroacetyl Chloride.* Each resin was prewashed with THF (2x8 ml), and then charged with THF (8 ml), pyridine (0.3 ml, 3.67 mmole) and chloroacetyl chloride (0.25 ml, 2.5 mmole). The reaction mixture was stirred for 8 hours at 45°C, and then for 6-8 hours at room temperature. Each resin was filtered, washed with a 2:1 mixture of dichloromethane/methanol (1x8 ml), methanol (2x8 ml) and THF (2x8 ml).
- 10 The acylation was repeated using the same loading of reagents, but a shorter reaction time of 4 hours at 45°C, and 2 hours at room temperature. Each resin was then filtered, washed with a 2:1 mixture of dichloromethane and methanol (1x8 ml), and then with methanol (3x8 ml). Each resin was dried under argon for 10 minutes. Each resin was then transferred into a vial and dried in a desiccator under vacuum for 1 hour.
- 15 b. *Acylation with α -Phenyl- α -Chloroacetyl Chloride.* The same procedure set out for the acylation with chloroacetyl chloride was used. A 2.5 mmol excess of α -phenyl- α -chloroacetyl chloride, relative to mmol amount of linker in the rink-acid resin, was used.
- c. *Acylation with α -Halo- α -Methyl; α -Halo- α -Ethyl and α -Halo - α -Butylacetyl Bromide.* A 1:1:1 mixture (by volume) of the α -bromopropionyl bromide (R_4 = Me), α -bromobutyryl bromide (R_4 = Et), and α -bromohexanoyl bromide (R_4 = Bu) was
- 20 used to give a molar ratio of 0.52:0.56:0.42 (in mmols). This resulted in a molar excess of 1.65, 1.75 and 1.31, respectively, if the original coverage of the resin was 0.63 mmol/g (0.5 g resin per tube), and 2.4, 2.6 and 1.9 if the original coverage of the resins was 0.43 mmol/g (0.5g resin per tube).
- 25 d. *Acylation with α -Chloro- α -Methyl Acetic acid.* Each resin was prewashed with dichloromethane. Each tube was charged with 3 ml of a solution of PyBrop (0.29 g, 0.62 mmole) in dichloromethane, a solution of the α -chloro- α -methylacetic acid (0.095 g, 0.62 mmole) in 3ml of DMF, and EtN(iPr)₂ (0.2 ml, 1.2 mmole). Each reaction mixture was allowed to react for 16-18 hours at room temperature. Each resin was then filtered,
- 30 washed with dichloromethane (2x8 ml) and methanol (2x8 ml), and the acylation was repeated. Each resin was then filtered, washed with dichloromethane (2x8 ml), methanol

(3x8 ml), and dried under argon for about 10 minutes. Each resin was transferred into a vial, and dried in a desiccator under vacuum for one hour.

D. Addition of the Second Amine

Ten, or thirty prepared α -haloacetyl amide resins from the first three steps were pooled together, leaving 0.05-0.10 gram of each individual resin for necessary deconvolutions. A suspension of the pooled resin mixture in 100 ml of a 2:1 mixture of dichloromethane and THF was distributed into one, two or three, 96-well reaction plates. For one reaction plate, 1.7 to 2.0 grams of resin were used. For two reaction plates, 3.0 to 3.3 grams of resin were used, and for three reaction plates, 4.7 to 5.0 grams of resin were used. The distributed suspension was then filtered using a filtration manifold, a small lightweight manifold that is generally used for drawing solvents and reagents from the chambers of the 96-well reaction plates. The reaction plates were transferred into COMBICLAMPS® (Huntington, West Virginia), and 10% EtN(iPr)₂ in DMF was added at 0.2 ml per well (0.21 mmole of EtN(iPr)₂ per well), followed by the addition of a 1.0M solution of the appropriate amine from the corresponding master plate, 0.1 ml per well (0.1 mmole amine per well). The COMBICLAMPS® are used to accommodate 96-well reaction plates during synthesis, allowing for the addition of reagents into the plates, and a proper sealing that maintains reagents and solvents for hours at elevated temperatures. These clamps consist of a top and bottom cover provided with changeable, chemically resistant sealing gaskets. They are designed to accommodate 96-well reaction plates between the top and bottom covers. The reaction plates were sealed and kept in an oven at 70-75°C for 16 hours. After cooling to room temperature, the resins were filtered, washed with a 1:1 mixture of DCM/methanol (1x1ml), methanol (2x1ml), and then dried in a desiccator under vacuum for 2 hours.

E. Reduction with Red-Al

The reaction plates were placed into COMBICLAMPS®. A 1:6 mixture of Red-Al (65+w% in toluene) and THF was added, at 0.6 ml per well (0.28mmole of Red-Al per well), and allowed to react for 4 hours. Each resin was then filtered, washed with THF (2x1 ml), and methanol (3x1 ml). The addition of methanol should proceed with caution. Each resin was then dried under vacuum.

F. Cleavage of Final Ethylene Diamine Compound

This step was carried out using a cleavage manifold, a Teflon coated aluminum, filter/collection vacuum manifold, designed for recovering cleavage products from the reaction plates into collection plates. The manifold is designed to ensure that the filtrate
5 from each well is directed to a corresponding well in a receiving 96-well collection plate. The reaction plates (placed on the top of the collection plates in this manifold) were charged with a 10:85:5 mixture of TFA, dichloromethane, and methanol (0.5 ml of mixture per well). After fifteen minutes, the solutions were filtered and collected into proper wells on the collection plates. The procedure was repeated. Solvents were evaporated on a
10 SPEED VAC®, Holbrook, New York, and the residual samples (TFA salts) were tested without further purification.

EXAMPLE II

Deconvolution Example

Deconvolution of the active wells was performed by re-synthesis of discrete
15 compounds, from the archived α -haloacetyl amide resins (10 resins, 0.05 - 0.10 g each), which were set aside at the end of the acylation step before the pooling. Each resin was assigned a discrete column (1, or 2, or 3, etc., see the template) in a 96 well filterplate, and was divided between X rows (A, B, C, etc), where X is the number of hits discovered in the original screening plate. To each well, in a row, a selected N2 (R_3R_2NH) hit amine (0.1
20 mmol), DMF (180 ml) and EtNiPr₂ (20 ml) were added: the first selected amine was added to the resins in the row "A", the second amine - to the resins in the row "B", the third amine - to the resins in the row "C", etc. A lay-out of a representative 96-well filter plate is shown in Table 4.

The reaction plates were sealed and kept in an oven at 70-75⁰C for 16 hours. After
25 cooling to room temperature, the resins were filtered, washed with a 1:1 mixture of DCM and methanol (1x1ml), methanol (2x1ml), and dried in desiccator under vacuum for 2h. Reduction and cleavage were performed according to steps 5 and 6 in the original synthetic protocol. The product wells from the cleavage were analyzed by ESI-MS (Electro Spray Ionization Mass Spectroscopy) to ensure the identity of the actives, and were tested in the
30 same Luc and MIC assays.

Table 4

Lay-Out of Representative 96-Well Filter Plate


A1	A2	A3	A4	A5	A6	A7	A8	A9	A10	Selected amine N2, Added to A1-A10
B1	B2	B3	B4	B5	B6	B7	B8	B9	B10	Selected amine N2, Added to B1-B10
C1	C2	C3	C4	C5	C6	C7	C8	C9	C10	Selected amine N2, Added to C1-C10
D1	D2	D3	D4	D5	D6	D7	D8	D9	D10	Selected amine N2, Added to D1-D10
E1	E2	E3	E4	E5	E6	E7	E8	E9	E10	Selected amine N2, Added to E1-E10
F1	F2	F3	F4	F5	F6	F7	F8	F9	F10	Selected amine N2, Added to F1-F10
G1	G2	G3	G4	G5	G6	G7	G8	G9	G10	Selected amine N2, Added to G1-G10
H1	H2	H3	H4	H5	H6	H7	H8	H9	H10	Selected amine N2, Added to H1-H10
Resin #1	Resin #2	Resin #3	Resin #4	Resin #5	Resin #6	Resin #7	Resin #8	Resin #9	Resin #10	*X* selected Amines N2 to be added on the step 4 Individual resins #1 – 10, preloaded with proper amine N1.

EXAMPLE III

Solid-Phase Synthesis of Selected Substituted Ethylenediamine Compounds Using the QUEST® 210 Synthesizer

5 The solid-phase protocol described above in Example I was applied to the scaled-up synthesis of the selected substituted ethylene diamine compounds. Here, all reaction steps, from the activation of the Rink-acid resin to the cleavage of the final product, were carried out using the QUEST® instrument only, which allowed for the simultaneous syntheses of twenty parallel reactions. Purification of all crude samples was done by HPLC to yield
10 desirable products in purity greater than 90%. Table 3 lists the scale-ups of substituted ethylene diamines. Here, the synthesis of one of the active compounds, *N*-Geranyl-*N'*-(2-adamanthyl)ethane-1,2-diamine is described below as an example.

The Preparation of *N*-Geranyl-*N'*-(2-adamanthyl)ethane-1,2-diamine (compound 109) is set forth in Figure 12.



Compound 109

- 15
1. Activation of the Rink-acid resin. Synthesis of Rink-Cl resin. Rink-acid resin, coverage (linker) of 0.43 to 0.63 mmol/g (0.8 g, 0.5 mmol), was placed into one of the
20 10 ml tubes of QUEST® 210 Synthesizer, and washed twice with THF. A solution of triphenylphosphine (0.380 g, 1.45 mmol) in THF (3 ml) was added, followed by the addition of a solution of hexachloroethane (0.4 g, 1.43 mmol) in THF (3 ml). THF was added up to the volume of the tube (approximately 2 ml). After 6 hours, the resin was filtered, washed with THF (2x8 ml) and dichloromethane (2x8 ml).
 - 25 2. Addition of the first amine. Synthesis of resin attached geranylamine. The tube with activated resin was charged with 3 ml of dichloroethane, EtN(*i*Pr)₂, (0.3 ml, 1.74 mmol), and geranylamine (0.230 g, 1.5 mmol). Dichloroethane was added to a volume of 8 ml. The reaction was carried for 8 hours at 45°C, and for 6-8 hours at room temperature. Geranylamine loaded resin was filtered, washed with a 2:1 mixture of dichloromethane and

methanol (1x8 ml), then with methanol (2x8 ml), and suck dried for 10 minutes under argon.

3. Acylation with chloroacetyl chloride. Synthesis of resin attached *N*-Geranyl- α -chloroacetamide. The resin was prewashed with THF (2x8ml). The tube was charged with 8 ml of THF, pyridine (0.3 ml, 3.67 mmol), and chloroacetyl chloride (0.2ml, 2.5 mmol), and allowed to stir for 8h at 45⁰C, and 6-8 h at room temperature (RT). After the reaction was complete, the resin was filtered, washed with a 2:1 mixture of dichloromethane and methanol (1x8 ml), methanol (2x8 ml), and THF, and the acylation was repeated using the same loads of the reagents, but shorter reaction time: 4 hours at 45⁰C and 2 hours at room temperature. At the end, the α -chloroacetamide loaded resin was filtered, washed with a 2:1 mixture of dichloromethane and methanol (1x8 ml), methanol (3x8 ml), and suck dried for 15 min under argon.

4. Addition of the second amine. Synthesis of resin attached *N*-Geranyl-*N'*-(2-adamantyl)acetamide. The tube with the resin was charged with DMF (3 ml) and EtN(*i*Pr)₂ (0.6 ml, 4.4 mmol), followed by the addition of a suspension of 2-adamantamine hydrochloride (2.0 g, 1.1 mmol) in DMF (4 ml), and was allowed to stir at 70-75⁰C for 16 hours. After cooling down to the room temperature, the resin was filtered, washed with a 1:1 mixture of DCM and methanol (1x8 ml), methanol (2x8 ml), and suck dried for 15 minutes under argon.

5. Reduction with Red-Al. Synthesis of resin attached *N*-Geranyl-*N'*-(2-adamantyl)ethane-1,2-diamine. The resultant resin was suspended in anhydrous THF (3 ml) in a tube, and stirred for 15 min. Commercially available Red-Al, 65+ w% in toluene, was added (2.0 ml, 6.4 mmol), followed by addition of 2-3 ml of anhydrous THF (to fill up the volume of the tube). The mixture was allowed to react for 4 hours. After the reaction, the resin was filtered, washed with THF (1x8 ml), a 1:1 mixture of THF and methanol (1x8 ml) (addition of MeOH should proceed with caution), methanol (3x8 ml), and then dried.


6. Cleavage from the resin and purification. Synthesis of *N*-Geranyl-*N'*-(2-adamantyl)ethane-1,2-diamine acetate. For this last step of the synthesis, the tube with the resin was charged with a 10:90 mixture of TFA and dichloromethane, and the formed

bright red suspension was allowed to stir for 30 min. After addition of MeOH (0.5 ml), the colorless suspension was filtered, and the filtrate was collected into a proper tube. The procedure was repeated, and solvents were evaporated on a SPEEDVAC®. Half of the amount of crude *N*-Geranyl-*N'*-(2-adamantyl)ethane-1,2-diamine (in a form of trifluoroacetate salt) was purified by HPLC using following conditions: column C18, flow 4 ml/min, 30 min run, gradient starting with 5%AcOH/MeOH (100%) finishing up with acetonitrile (100%). Obtained: 27 mg of *N*-Geranyl-*N'*-(2-adamantyl)ethane-1,2-diamine diacetate, 24% yield, 98% purity by NMR.

EXAMPLE IV

Representative Solution Phase Synthesis of the Active Compounds

Preparation of *N*-(Cyclooctyl)-*N'*-(1*R*, 2*R*, 3*R*, 5*S*)-(-)-isopinocampheylethane-1,2-diamine as hydrochloride (compound 59) is set forth in Figure 13.



Compound 59

Bromocyclooctylacetylamine. To a mixture of cyclooctylamine (3.3 g, 0.026 mol) and pyridine (2.42 g, 0.031 mmol) in anhydrous THF (80 ml) at 0°C was added dropwise, via syringe, bromoacetyl bromide (5.78 g, 0.029 mol). The reaction temperature was maintained by an ice bath. The reaction mixture was allowed gradually to warm up to room temperature, and was stirred at room temperature for 1 hour. The precipitate was removed by filtration, washed with ethyl ether (1 x 30 ml), and the filtrate was concentrated to dryness on a rotary evaporator. Bromocyclooctylacetylamine was forwarded to the second step without additional purification.

N-(Cyclooctyl)-*N'*-(1*R*, 2*R*, 3*R*, 5*S*)-(-)-isopinocampheyl-1-carbonylethane-1,2-diamine. To a solution of the bromocyclooctylacetylamine in DMF (60 ml) were added Hunig's base (4.64 g, 0.036 mol) and (1*R*, 2*R*, 3*R*, 5*S*)-(-)-isopinocampheylamine (4.5 g, 0.029 mol), and the reaction mixture was stirred at 80°C for 16 hours. After cooling off to the room temperature, the reaction mixture was diluted with 150 ml of ethyl ether, and washed with 1M NaOH solution (2 x 50 ml). The organic layer was washed with brine (1

x 50 ml), dried over MgSO₄, and concentrated to dryness on the rotory evaporator. The residue (11.04 g) as brown oil was purified on COMBIFLASK® (Isco, Lincoln, Nebraska, USA), using Silicagel catridges commercially available from BIOTAGE® (Biotage, Inc. of Dyax Corp, Va, USA), and the following mobile phase gradient: 30 min run, starting with DCM, 100%, and finishing up with a mixture DCM:MeOH:NH₄OH (600:400:10). The final product (7.29 g) was obtained as a brown oil; 76% yield, purity 90%.

N-(Cyclooctyl)-*N'*-(1*R*, 2*R*, 3*R*, 5*S*)-(-)-isopinocampheylethane-1,2-diamine. To a solution of the amide, from previous step, in anhydrous THF (160 ml), was added dropwise via syringe commercially available (SIGMA-ALDRICH®) Red-Al, as 65wt% solution in THF (28 ml, 0.09 mol). The reaction mixture was stirred at reflux for 20 hours. After cooling down to the room temperature, the reaction mixture was poured into 1.5M NaOH (200 ml), and extracted with ethyl ether (2 x 100 ml). The organic layer was washed with brine (1 x 100 ml), dried over MgSO₄, and evaporated to dryness on the rotory evaporator to yield 7.2 g of a crude product, as a brown oil. Chromatographic purification of the crude using the same equipment and conditions as for the previous step, gave 3.5 g of the diamine. The diamine was treated with 2.0M solution of HCl in ethyl ether (25 ml), and kept in a refrigerator overnight. A dark yellow solid (4.2 g) formed, and was filtered off, and recrystallized from MeOH and ethyl ether to yield 1.5 g of the diamine as an HCl salt (of purity greater than 98%, NMR and MS are available), 19% overall yield.

20

EXAMPLE V

Mass Spectroscopy Analysis

Mass spectra data were obtained by Elecspray Ionization technique on a PERKIN ELMER®/SCIEX®, API-300, TQMS with an autosampler, manufactured by SCIEX®, Toronto, Canada.

25

A. Library of Substituted Ethylenediamines

Mass spectroscopy served as a means for monitoring the reaction results of the library of ethylenediamines. Mass spectroscopy was done on two randomly selected rows (24 samples) per reaction plate, for roughly 28,000 compounds in pool of 10 or 30 compounds per well. Thus, if ten compounds per well were synthesized, the mass spectra for each well should contain ten signals, correlating with the proper molecular ions for each compound. The presence or absence of a particular signal indicated the feasibility of the

30

particular synthesis. Based on the mass spectral data, and on a general analysis of the reactivity of the various amines, it is estimated that 67,000 compounds were formed out of 112,000 compounds.

Figure 14 is a representative mass spec profile for one sample well. Mass spectra for a representative ethylene diamine compound is shown in Figure 15. Tables 5 to 8, below, list illustrative examples of mass spec data for representative reaction wells, with each well containing ten substituted ethylene diamines.

Table 5


ILLUSTRATIVE EXAMPLES OF MASS SPEC DATA FOR REPRESENTATIVE ETHYLENEDIAMINES (TEN COMPOUNDS PER WELL).

R ₁ NH ₂ in the 1 st position (pool of 10 resins)	R ₂ R ₃ NH in the 2 nd position (from the master plate of the amines)	[M+1] ⁺ of the product R ₁ NHCH ₂ CH ₂ NR ₂ R ₃
<i>Plate # 4-034-2, well D10</i> 1-(2-Aminoethyl)piperidine Phenethylamine 4-(2-Aminoethyl)morpholine Tryptamine Cyclohexylamine Exo-2-Aminonorbornane Benzylamine 2-Fluorophenethylamine ?-Methylphenethylamine 4-Methoxyphenethylamine	2-Aminoheptane	270 absent 263 272 absent 302 241 253 249 281 277 293
<i>Plate # 4-56-1, well C4</i> 4-Methylbenzylamine Cyclopentylamine 2-(Aminomethyl)piperidine Furfurylamine 3,4,5-Trimethoxybenzylamine 1-Methyl-3-phenylpropylamine Cyclobutylamine 1,2,3,4-Tetrahydro-1-naphthylamine 2,3-Dimethylcyclohexylamine 2-Amino-1-butanol	exo-2-Aminonorbornane	259 223 246 low intensity 235 335 287 209 258 265 227 low intensity

<i>Plate # 4-44-2, well G1</i>		
Veratrylamine	4-Fluorophenethylamine	333
2-(1-Cyclohexenyl)ethylamine		291
5-Aminoquinolone		310 absent
1-(1-Naphthyl)ethylamine		337 absent
1-Aminopiperidine		266
3-Fluorobenzylamine		291
2,4-Dimethoxybenzylamine		333
3-Amino-1,2,4-triazine		262 absent
2-Ethoxybenzylamine		317
4-(3-Aminopropyl)morpholine		310 absent

Table 6


Mass Spec Data for Synthesized Ethylenediamines



5


R ₁ NH ₂ in the 1 st position	[M+1] ⁺ of the products, R ₄ = H	[M+1] ⁺ of the products, R ₄ = Ph	
		Diamines, 1	mino alcohols, 13
Tyramine	308	384	258 formed
2-Adamantamine	321 absent	398 absent	272 formed
cis-Myrtanylamine	324	400	274 formed
3-Amino-1-propanol	246	322	196 absent
L-Methioninol	305 absent	382 absent	256 absent
Cyclooctylamine	298	374	248 formed
(1S,2S)-2-Amino-1-phenyl-1,3-propanediol	337 absent	414 absent	288 absent
1-Adamantanemethylamine	336	412 absent	286 formed
2,2-Diphenylethylamine	368	444	318 formed
5-Amino-1-pentanol	274	350	224 formed

Table 7

Mass Spec Data for Synthesized Ethylenediamines, $R_4 = H$ and Me

R_1NH_2 in the 1 st position	$[M+1]^+$ of the products, $R_4 = H$	$[M+1]^+$ of the products, $R_4 = Me$	
		Diamines, 1	Amino alcohols, 13
Tyramine	278	293	196 absent
2-Adamantamine	293 absent	307 absent	210 low intensity
cis-Myrtaniline	293	309	212 formed
3-Amino-1-propanol	217	231	134 absent
L-Methioninol	277 absent	291 absent	194 formed
Cyclooctylamine	269	269 absent	186 absent
(1S,2S)-2-Amino-1-phenyl-1,3-propanediol	309 low intensity	323 absent	226 formed
1-Adamantanemethylamine	307	321	224 formed
2,2-Diphenylethylamine	339	353	256 formed
5-Amino-1-pentanol	245	259	162 absent

Table 8

Mass Spec Data for Synthesized Ethylenediamines, $R_4 = H$ and Me

R_1NH_2 in the 1 st position	$[M+1]^+$ of the products, $R_4 = H$	$[M+1]^+$ of the products, $R_4 = Me$	
		Diamines, 1	Amino alcohols, 13
Tyramine	278	292 absent	196 absent
2-Adamantamine	292 absent	306 absent	210 formed
cis-Myrtaniline	294	308 absent	212 formed
3-Amino-1-propanol	216	230 absent	134 absent
L-Methioninol	276 absent	290 absent	194 absent
Cyclooctylamine	268	282 absent	186 absent
(1S,2S)-2-Amino-1-phenyl-1,3-propanediol	308	322 absent	226 formed
1-Adamantanemethylamine	306 absent	320 absent	224 formed
2,2-Diphenylethylamine	338	352 absent	256 formed
5-Amino-1-pentanol	244	258 absent	162 absent

5

EXAMPLE VI

 1H NMR Spectroscopy

Proton NMR data was recorded on a VARIAN® Nuclear Magnetic Resonance Spectrometer (Palto Alto, California) at 500 MHz.

10 Representative substituted ethylene diamines were purified by HPLC, and analyzed by proton NMR. A representative proton NMR profiles is shown in Figure 16. NMR and MS data for some representative hit compounds are shown below.

Compound 6.

15 N^2 -(1-Adamantylmethyl)- N^1 -(3,3-diphenylpropyl)propane-1,2-diamine. 55 mg, 36% yield. 1H NMR: δ 7.28-7.15 (m, 5H), 3.95 (t, $J = 7.9$ Hz, 1H), 2.94 (br s 4H), 2.71 (dd, $J = 7.6, 9.8$ Hz, 2H), 2.41 (s, 2H), 2.32 (dd, $J = 7.6, 7.9$ Hz, 2H), 2.16 (s), 2.08-1.98 (m, 4H), 1.72 (m, 6H), 1.62 (m, 6H), 1.51 (d, $J = 2.4$ Hz, 3H). Mass spectrum (ESI) m/z (MH)⁺ 417.

Compound 7.

- 5 *N*-(3,3-Diphenylpropyl)-*N'*-(1-adamantylmethyl)ethane-1,2-diamine. 28 mg, 22% yield. ¹H NMR (500MHz) δ 7.30-7.12 (m, 10H); 3.95 (t, *J* = 7.6 Hz, 1H); 2.91 (d, *J* = 1.2 Hz, 4H); 2.70 (dd, *J* = 7.6 and 1.2 Hz, 2H); 2.40 (d, *J* = 1.3 Hz, 2H); 2.32 (q, *J* = 8.0 Hz, 2H); 1.98 (br d, *J* = 1.7 Hz, 4H); 1.72 (d, *J* = 12.2 Hz, 4H); 1.62 (d, *m?* *J* = 12.2 Hz, 4H); 1.51 (br s, 6H). Mass spectrum (ESI) *m/z* (MH)⁺ 403.6.

Compound 10.

- 10 *N*-(-)-*cis*-Myrtanyl-*N'*-(3,3-diphenylpropyl)ethane-1,2-diamine. 14 mg, 11% yield. ¹H NMR (500MHz) δ 7.30-7.10 (m, 10H); 3.95 (m, 1H); 2.92-2.83 (m, 4H); AB: 2.80 (d, *J* = 7 Hz, 1H); 2.76 (d, *J* = 8 Hz, 1H); ; 2.65 (dd, *J* = 9.6 and 7.6 Hz, 2H); 2.42-2.20 (m, 4H), 2.29 (d, *J* = 8 Hz, 2H), 1.90 (m, 8H); 1.42 (m, 1H); 1.19 (m, 2H); 1.17 (s, 3H); 0.95 (s, 3H); , 1.00-0.8 (m, 2H). Mass spectrum (ESI) *m/z* (MH)⁺ 391.3.

15 Compound 14.

- 20 *N*-(3,3-Diphenylpropyl)-*N'*-*exo*-(2-norborny)ethane-1,2-diamine. 17 mg, 16% yield. ¹H NMR (500MHz) δ 7.30-7.15 (m, 10H); 3.95 (t, *J* = 7.9 Hz, 1H); 2.86 (dd, *J* = 11.5 and 1.5 Hz, 4H); 2.73 (dd, *J* = 8.0 and 3.3 Hz, 1H); 2.64 (t, *J* = 7.6 Hz, 2H); 2.29 (t, *J* = 7.5 Hz, 2H), 2.31-2.26 (m, 2H) 2.30 1.96 (s, 3H); 1.63 (ddd, *J* = 13.1, 7.9 and 2.5 Hz, 1H); 1.60-1.50 (m, 1H); 1.50-1.43 (m, 2H); 1.30 (dq, *J* = 4.0 and 13.5 Hz, 1H), (1H, m), 1.20 (dd, *J* = 10.4 and 1.1 Hz, 1H), 1.11 (dd, *J* = 2.0, and 8.5 Hz, 1H), 1.08 (dd, *J* = 2.5, and 8.5 Hz, 1H), 1.10 (dq, *J* = 8.3 and 2.1, 2H). Mass spectrum (ESI) *m/z* (MH)⁺ 349.1.

Compound 21.

- 25 *N*-(3,3-Diphenylpropyl)-*N'*-(1*S*)-(1-ethylcyclohexane)ethane-1,2-diamine. 5 mg, 4% yield. Mass spectrum (ESI) *m/z* (MH)⁺ 365.5.

Compound 32.

- 30 *N*-(2,2-Diphenylethyl)-*N'*-@-(+)-bornylethane-1,2-diamine. 58 mg, 48% yield. ¹H NMR (500MHz): δ 7.30-7.10 (m, 10H); 4.18 (t, *J* = 6.8 Hz, 1H); 3.34 (d, *J* = 7.6 Hz, 2H); 3.02 (m, 4H); 2.95-2.90 (m, 1H); 2.15-2.08 (m, 1H); 1.94 (m, 1H); 1.72-1.65 (m, 2H); 1.48-1.30 (m, 2H); 1.27-1.10 (m, 2H); 1.06 (dd, *J* = 13.6 and 4.1 Hz, 1H); 0.82 (s, 3H); 0.81 (s, 3H); 0.78 (s, 3H). Mass spectrum (ESI) *m/z* (MH)⁺ 377.2

35 Compound 34.

- 40 *N*-(2,2-Diphenylethyl)-*N'*-(1-adamantylmethyl)ethane-1,2-diamine. 6.8 mg, 6% yield. ¹H NMR (500MHz) δ 7.30-7.15 (m, 10H); 4.15 (t, *J* = 7.6 Hz, 1H); 3.24 (dd, *J* = 7.9 and 1.2 Hz, 2H); 2.79 (t, *J* = 6.5 Hz, 2H); 2.74 (t, *J* = 6.0 Hz, m, 2H); 1.95 (m, 8H); 1.69 (d, *J* = 12.5 Hz, 4H); 1.59 (d, *J* = 11.9 Hz, 4H); 1.40 and 1.39 (br s, 3H); Mass spectrum (ESI) *m/z* (MH)⁺ 389.0.

Compound 37.

- 45 *N*-(2,2-Diphenylethyl)-*N'*-(-)-*cis*-myrtanylethane-1,2-diamine. 54 mg, 38% yield. ¹H NMR: δ 7.31-7.18 (m, 10H), 4.13 (t, *J* = 7.6 Hz, 1H), 3.26 (d, *J* = 7.6 Hz, 2H), 2.86 (dd, *J* = 4.3, 8.0 Hz, 4H), 2.76 (dd, *J* = 7.6, 12.2 Hz, 2H), 2.37 (ddd, *J* = 1.8, 9.0, 12.5 Hz, 1H), 2.12 (dq, *J* = 1.8, 7.6 Hz, 1H), 1.98 (br s, 2H), 1.98-1.84 (m, 4H), 1.39 (ddd, *J* = 2.4, 4.0, 6.1 Hz, 1H), 1.18 (s, 3H), 0.95 (s, 3H), 0.91 (d, *J* = 10.0 Hz, 1H) Mass spectrum (ESI) *m/z* (MH)⁺ 377.2.

Compound 38.

- N*-(-)-*cis*-Myrtanyl-*N'*-(2,2-diphenylethyl)propane-1,2-diamine. 39 mg, 30% yield. ¹H NMR (500MHz) δ 7.30-7.15 (m, 10H); 4.13 (t, *J* = 8.0 Hz, 1H); AB: 3.28 (d, *J* = 7.5 Hz, 1H); 3.24 (d, *J* = 7.5 Hz, 1H), 3.26 (d, *J* = 6.1 Hz, 2H); 2.96 (m, 1H); 2.88-2.75 (m, 2H); 2.71 (ddd, *J* = 4.5, 9.0, 13.0 Hz, 1H), 2.58 (ddd, *J* = 7.0, 10.0, 14.0 Hz, 1H); 2.35 (m, 1H); 2.21 (m, 1H); 2.00-1.80 (m, 6H); 1.40-1.20 (m, 1H); 1.17 (s, 3H); ; 0.93 (s, 3H); 0.89 (dd, *J* = 9.7 and 4.2 Hz, 1H). Mass spectrum (ESI) *m/z* (MH)⁺ 391.0.

Compound 40.

- N*-(2,2-Diphenylethyl)-*N'*-(1*R*, 2*R*, 3*R*, 5*S*)-(-)-isopinocampheylethane-1,2-diamine. 33 mg, 23% yield. ¹H NMR: δ 7.31-7.18 (m, 10H), 4.13 (t, *J* = 7.5 Hz, 1H), 3.27 (d, *J* = 8.0 Hz, 2H), 3.14 (dt, *J* = 6.0, 10 Hz, 1H), (4H), 2.36 (qd, *J* = 2.0, 6.0 Hz, 1H), 2.34 (dt, *J* = 2.0, 10 Hz, 1H), 2.07-1.96 (m, 3H), 1.82 (dt, *J* = 2.0, 6.0 Hz, 1H), 1.71 (ddd, *J* = 2.5, 5.5, 13.5 Hz, 1H), 1.22 (s, 3H), 1.09 (d, *J* = 7.0 Hz, 3H), 0.96 (d, *J* = 10.5 Hz, 1H), 0.91 (s, 3H). Mass spectrum (ESI) *m/z* (MH)⁺ 377.3.

Compound 47.

- N*-(-)-*cis*-Myrtanyl-*N'*-(1*R*, 2*R*, 3*R*, 5*S*)-(-)-isopinocampheylethane-1,2-diamine. 42 mg, 33% yield. ¹H NMR: δ 3.35-3.20 (m, 6H), 2.93 (dd, *J* = 4.6, 2.0 Hz, 2H), 2.45-2.33 (m, 4H), 2.17 (s, 3H), 2.06 (quint, *J* = 7.0 Hz, 1H), 2.0-1.9 (m, 6H), 1.90 (dd, *J* = 2.1, 5.2 Hz, 1H), 1.87 (dt, *J* = 1.8, 4.6 Hz, 1H), 1.51 (ddd, *J* = 4.6, 10.0, 13.0 Hz, 1H), 1.23 (s, 3H), 1.19 (s, 3H), 1.12 (d, *J* = 8 Hz, 3H), 1.03 (d, *J* = 10.3 Hz, 1H), 0.98 (s, 3H), 0.94 (d, *J* = 9.8 Hz, 1H), 0.94 (s, 3H). Mass spectrum (ESI) *m/z* (MH)⁺ 333.6.

Compound 52.

- N*-(3,3-Diphenylpropyl)-*N'*-cyclooctylethane-1,2-diamine. 20 mg, 18% yield. ¹H NMR (500MHz): δ 7.30-7.10 (m, 10H); 3.96 (t, *J* = 7.9 Hz, 1H); 3.00 (m, 1H); 2.90 (dd, *J*₁ = *J*₂ = 5.5 Hz, 2H); 2.84 (dd, *J*₁ = *J*₂ = 5.0 Hz, 2H); 2.61 (t, *J* = 7.3 Hz, 2H), 2.27 (q, *J* = 7.6 Hz, 2H); 1.83 (m, 2H); 1.74 (m, 2H); 1.65-1.40 (m, 10H).

Compound 55.

- N*-(1-Adamantylmethyl)-*N'*-cyclooctylethane-1,2-diamine. 6.7 mg, 6% yield. ¹H NMR (500MHz): δ 3.08-3.02 (m, 1H), 3.02-2.98 (m, 2H); 2.97-2.92 (m, 2H); 2.36 (s, 2H); 1.98 (m, 2H); 1.93-1.86 (m, 2H); 1.80-1.50 (m, 19H).

Compound 57.

- N*-(-)-*cis*-Myrtanyl-*N'*-(cyclooctyl)ethane-1,2-diamine. 18 mg, 18% yield. ¹H NMR (500MHz) δ 3.05-2.95 (m, 4H); AB: 2.76 (d, *J* = 7.5 Hz, 1H), 2.23 (d, *J* = 8.0 Hz, 1H); 2.76 (dd, *J* = 11.6 and 7.3 Hz, 1H); 2.73 (dd, *J* = 11.9 and 8.2 Hz, 1H); 2.40-2.34 (m, 1H); 2.28 (quintet, *J* = 8.0 Hz, , 1H); 1.97 (s, 3H); 2.00-1.84 (m, 6H); 1.80-1.70 (m, 2H); 1.68-1.38 (m, 11H); 1.18 (s, 3H); 0.97 (s, 3H); 0.92 (d, *J* = 9.8 Hz, 1H). Mass spectrum (ESI) *m/z* (MH)⁺ 307.5.

Compound 58.

- N*-(2-Adamantyl)-*N'*-cyclooctylethane-1,2-diamine. 25 mg, 23% yield. ¹H NMR: δ 3.06 (m, 1H), 3.00 (t, *J* = 6.1 Hz, 2H), 2.93 (t, *J* = 5.5 Hz, 2H), 2.83 (br s, 1H), 1.96 (s, 3H), 1.92-1.80 (m, 10H), 1.80-1.50 (m, 20H). Mass spectrum (ESI) *m/z* (MH)⁺ 305.1.

Compound 59.

N-(Cyclooctyl)-*N'*-(1*R*, 2*R*, 3*R*, 5*S*)-(-)-isopinocampheylethane-1,2-diamine.

- 15 mg, 14% yield. ¹H NMR (400MHz): δ 3.47 (dt, *J* = 6.0, 10.0 Hz, 1H), 3.40-3.28 (m, 7H), 2.44 (tq, *J* = 2.0, 10.0 Hz, 1H), 2.36 (dtd, *J* = 2.0, 6.0, 10.0 Hz, 1H), 2.09 (dq, *J* = 2.0, 7.2 Hz, 1H), 2.00-1.90 (m, 3H), 1.88-1.78 (m, 2H), 1.78-1.63 (m, 4H), 1.65-1.30 (m, 8H), 1.18 (d, *J* = 6.0 Hz, 3H), 1.16 (s, 3H), 1.17 (d, *J* = 7.2 Hz, 1H), 0.90 (s, 3H). Mass spectrum (ESI) *m/z* (MH)⁺ 307.4.

Compound 62.

- 10 *N*-(-)-*cis*-Myrtanyl-*N'*-(1*S*)-(1-ethylcyclohexane)ethane-1,2-diamine. 48 mg, 46% yield. ¹H NMR (500MHz): δ 3.06-3.00 (m, 1H); 2.98-2.95 (m, 2H); 2.92-2.84 (m, 1H); 2.79 (dd, *J* = 11.9 and 7.0 Hz, 1H); 2.75 (dd, *J* = 11.9 and 7.9 Hz, 1H); 2.73 (m, 1H); 2.39 (m, 1H); 2.28 (quintet, *J* = 8.5 Hz, 1H); 2.00-1.86 (m, 6H); 1.82-1.76 (m, 2H); 1.68 (m, 2H); 1.54-1.42 (m, 2H); 1.32-1.10 (m, 6H); 1.19 (s, 3H); 1.13 (d, *J* = 6.7 Hz, 3H); 1.07 (dd, *J* = 12 and 3 Hz, 2H); 1.02 (dd, *J* = 12 and 3 Hz, 2H); 0.98 (s, 3H); 0.93 (d, *J* = 9.7 Hz, 1H). Mass spectrum (ESI) *m/z* (MH)⁺ 306.9.

Compound 65.

- 20 *N-trans*-(2-phenylcyclopropyl)-*N'*-(1-adamantyl)ethane-1,2-diamine. 18 mg, 16% yield. Mass spectrum (ESI) *m/z* (MH)⁺ 311.3.

Compound 66.

- 25 *N*-(3,3-Diphenylpropyl)-*N'*-(1*R*, 2*R*, 3*R*, 5*S*)-(-)-isopinocampheylethane-1,2-diamine. 2 mg, 2% yield. ¹H NMR (500MHz) δ 7.26 (m, 10H); 3.96 (t, *J* = 7.6 Hz, 1H); 3.09 (m, 1H); 2.92 (m, 1H); 2.84 (m, 2H); 2.62 (m, 2H); 2.35 (m, 4H); 1.97 (s, 3H); 1.82 (m, 1H); 1.68 (m, 1H); 1.21 (s, 3H); 1.12 (d, *J* = 7.3 Hz, 3H); 1.01 (m, 1H); 0.92 (s, 3H). Mass spectrum (ESI) *m/z* (MH)⁺ 391.4.

Compound 73.

- 30 *N*-(2-Adamantyl)-*N'*-[2-(2-methoxyphenyl)ethyl]ethane-1,2-diamine. 21 mg, 19% yield. ¹H NMR: δ 7.22 (dd, *J* = 8.2, 7.3 Hz, 1H), 7.14 (d, *J* = 7.3 Hz, 1H), 6.89 (d, *J* = 7.1 Hz, 1H), 6.87 (d, *J* = 8.2 Hz, 1H), 3.81 (s, 3H), 3.06 (t, *J* = 7.1 Hz, 2H), 3.06 (m, 2H), 3.01 (m, 2H), 2.93 (t, *J* = 7.1 Hz, 2H), 1.95 (br s, 2H), 1.90-1.80 (m, 7H), 1.78-1.66 (m, 6H), 1.59 (d, *J* = 2.5 Hz, 2H). Mass spectrum (ESI) *m/z* (MH)⁺ 329.4.

Compound 78.

- 40 *N*-2-Adamantyl-*N'*-2,3-dihydro-1*H*-inden-2-yl-ethane-1,2-diamine. 4.3 mg, 3% yield. ¹H NMR: δ 7.20 (dd, *J* = 4.9, 8.5 Hz, 2H), 7.14 (dd, *J* = 5.5, 2.1 Hz, 2H), 3.71 (quint, *J* = 6.1 Hz, 2H), 3.19 (dd, *J* = 5.8, 15.9 Hz, 2H), 3.13 (br.s, 1H), 3.05 (m, 4H), 2.86 (dd, *J* = 4.8, 15.8 Hz, 2H), 2.08 (m, 2H), 2.00 (m, 6H), 1.96-1.88 (m, 4H), 1.88-1.80 (m, 3H), 1.74 (m, 4H), 1.68-1.60 (m, 2H). Mass spectrum (ESI) *m/z* (MH)⁺ 303.4.

Compound 109.

- 45 *N*-Geranyl-*N'*-(2-adamantyl)ethane-1,2-diamine. 27 mg, 24% yield. ¹H NMR (400MHz): δ 5.40 (t, *J* = 7.2 Hz, 1H), 4.78 (br s, 2H), 3.64 (d, *J* = 7.6 Hz, 2H), 3.34 (m, 2H), 2.07 (m, 2H), 2.08-1.95 (m, 4H), 1.95-1.85 (m, 4H), 1.82 (m, 2H), 1.88-1.70 (m, 4H), 1.70-1.62 (m, 3H), 1.67 (s, 3H), 1.56 (s, 3H), 1.50 (s, 3H). Mass spectrum (ESI) *m/z* (MH)⁺ 307.4.

Compound 111.

N-Geranyl-*N'*-(2-ethylpiperidine)ethane-1,2-diamine. 44 mg, 42% yield. ¹H NMR (500MHz): δ 5.22 (t, *J* = 6.1 Hz, 1H); 5.04 (m, 1H), 3.52 (d, *J* = 7.3 Hz, 2H); 3.05-2.85 (m, 4H); 2.66 (m, 1H); 2.44 (m, 2H); 2.08 (m, 4H); 1.80-1.50 (m, 2H); 1.70 (s, 3H); 1.65 (s, 3H); 1.58 (s, 3H); 1.50-1.35 (m, 2H), 0.89 (t, *J* = 7.3, 3H). Mass spectrum (ESI) *m/z* (MH)⁺ 293.4.

Compound 116.

N-Geranyl-*N'*-allyl-*N'*-(cyclopentyl)ethane-1,2-diamine. 45 mg, 42% yield. ¹H NMR: δ 5.86 (ddd, *J* = 10.0, 16.1, 6.7 Hz, 1H), 5.28 (d, *J* = 15.9 Hz, 1H), 5.25 (d, *J* = 8.7 Hz, 1H), 5.23 (t, *J* = 7.3 Hz, 1H), 5.30 (m, 1H), 3.59 (d, *J* = 7.3 Hz, 2H), 3.28 (br d, *J* = 6.4 Hz, 2H), 3.16 (quintet, *J* = 8.2 Hz, 1H), 3.02 (m, 2H), 2.95-2.86 (m, 2H), 1.88-1.80 (m, 4H), 1.70 (s, 3H), 1.74-1.66 (m, 3H), 1.65 (s, 3H), 1.58 (s, 3H), 1.56-1.50 (2H), 1.50-1.40 (m, 2H). Mass spectrum (ESI) *m/z* (MH)⁺ 305.3.

Compound 117.

N-Geranyl-*N'*-diphenylmethylethane-1,2-diamine. 24 mg, 20% yield. ¹H NMR (500MHz): δ 7.40 (d, *J* = 7.2 Hz, 4H); 7.29 (t, *J* = 7.3 Hz, 4H); 7.21 (t, *J* = 7.0 Hz, 2H); 5.15 (t, *J* = 7.5, 1H); 5.01 (m, 1H); 4.89 (br s, 1H); 3.42 (d, *J* = 7.0 Hz, 2H); 3.00-2.78 (m, 4H); 2.20-2.00 (m, 4H); 1.63 (s, 3H); 1.59 (s, 3H); 1.56 (s, 3H). Mass spectrum (ESI) *m/z* (MH)⁺ 363.3.

Compound 125.

N,N'-bis-(-)-*cis*-Myrtanylpropane-1,2-diamine. 82 mg, 70% yield. ¹H NMR (500MHz): δ 3.62 (m, 1H); 3.18 (dd, *J* = 13.7 and 3.7 Hz, 1H); 3.05 (dt, *J* = 11.5 and 7.5 Hz, 1H); 3.06-2.92 (m, 2H); 2.86 (dt, *J* = 12.2 and 7.3 Hz, 1H); 2.40 (m, 4H); 2.06-1.84 (m, 10H); 1.56-1.46 (m, 2H); 1.37 and 1.36 (two d, *J* = 6.7 and *J* = 7.0 Hz, 3H); ; 1.20 (s, 3H); 1.19 (m, 3H), 0.99 and 0.98 (two s, 3H) Hz, H); 0.97 (s, 3H); 0.94 (two d, *J* = 10.1 Hz, 2H). Mass spectrum (ESI) *m/z* (MH)⁺ 346.9.

Compound 151.

N-[2-(2-Methoxy)phenylethyl]-*N'*-(1*R*, 2*R*, 3*R*, 5*S*)-(-)-isopinocampheyl-ethane-1,2-diamine. 67 mg, 60% yield. ¹H NMR (500MHz): δ 7.23 (t, *J* = 5.8 Hz, 1H); 7.13 (dd, *J* = 5.8 and 1.8 Hz, 1H); 6.88 (m, 2H); 3.81 (s, 3H); 3.13 (m, 1H); 3.1-3.0 (m, 3H); 3.01 (t, *J* = 7.0 Hz, 2H); 2.89 (t, *J* = 7.0 Hz, 2H); 2.42-2.35 (m, 2H); 2.00 (m, 3H); 1.82 (dt, *J* = 6.0 and 2.0 Hz, 1H); 1.72 (ddd, *J* = 2.5, 5.5, 13.5 Hz, 1H); 1.22 (s, 3H) 1.13 (d, *J* = 7.3 Hz, 3H). 0.99 (d, *J* = 10.1 Hz, 1H); 0.93 (s, 3H). Mass spectrum (ESI) *m/z* (MH)⁺ 331.5.

N-2-(2-Methoxyphenyl)ethyl-*N'*-allyl-*N'*-cyclopentyl-ethane-1,2-diamine. 8 mg, 7% yield. ¹H NMR: δ 7.26 (dd, *J* = 7.3, 8.5, 1H), 7.18 (d, *J* = 7.2 Hz, 1H), 6.91 (m, 2H), 5.61 (ddd, (*J* = 6.7, 17.0, 9.4 Hz, 1H), 5.13 (d, *J* = 15.3 Hz, 1H), 5.10 (d, *J* = 9.2 Hz, 1H), 3.83 (s, 3H), 3.13 (dd, *J* = 7.0, 6.7 Hz, 2H), 3.10 (d, *J* = 6.7 Hz, 1H), 3.00 (d, *J* = 7.3 Hz, 1H), 3.05-2.90 (m, 2H), 2.97 (dd, *J* = 8.2, 6.1 Hz, 2H), 2.75 (t, *J* = 6.1 Hz, 2H), 1.73 (m, 2H), 1.62 (m, 2H), 1.50 (m, 2H), 1.22 (m, 2H). Mass spectrum (ESI) *m/z* (MH)⁺ 311.4.

*N*²-(3-Phenylpropyl)-*N'*¹-[2-(4-fluorophenyl)ethyl]-1-phenylethane-1,2-diamine. 23 mg, 19% yield. ¹H NMR: δ 7.35 (d, *J* = 7.6 Hz, 2H), 7.34 (quart, *J* = 7. Hz, 1H), 7.26 (d, *J* = 6.4 Hz, 3H), 7.23 (d, *J* = 7.6 Hz, 2H), 7.17 (dd, *J* = 7.3, 6.4 Hz, 1H), 7.12 (d, *J* = 7.0 Hz, 2H), 3.21 (m, 1H), 3.03 (ddd, *J* = 4.2, 8.0, 12.8 Hz, 4H), 2.86 (t, *J* = 8.0 Hz, 2H), 2.85-2.79

(m, $J = 12$. Hz, 2H), 2.74-2.64 (m, 4H), 2.61 (t, $J = 7.7$ Hz, 2H), 1.96 (quint, $J = 7.6$ Hz, 2H). Mass spectrum (ESI) m/z (MH)⁺ 377.3.

EXAMPLE VII

5 *M. Tuberculosis* Rv0341p Lucs Drug Response

Substituted ethylene diamines, as described herein, were tested on *Mycobacterium tuberculosis* using high-throughout screening assay with recombinant mycobacterial containing promoter fusion of luciferase to Rv0341 EMB-inducible promoter. This assay
10 quickly and reliably identifies antimycobacterial activity in compound mixtures and/or in individual compounds. In this assay, bioluminescence increases when the mycobacteria is tested against an active compound, or an active compound mixture. During this assay, a theoretical yield of 100% was assumed for every unpurified substituted ethylene diamine, and the activity of each sample was compared to commercially available ethambutol
15 (99.0% purity). Results were reported in LCPS, and % Max. LCPS based on the activity of EMB at 3.1 μ M.

The substituted ethylene diamines were analyzed according to the following procedure. The diamines were dried in a speed vacuum to an approximate concentration of 6.3 mmoles per well. Each diamine, or diamine mixture, was then resuspended or
20 dissolved in 200 μ l of methanol for a concentration of 31.5mM diamine(s). The diamine(s) solution was diluted to a concentration of 200 μ M in 7H9 broth medium (a 1:15.75 dilution of the 31.5mM stock, followed by a 1:10 dilution; each dilution in 7H9 broth medium). Next, 50 μ l of the diluted diamine(s) solution was added to the first well of a row of twelve in an opaque, 96-well plate. The 7H9 broth medium, 25 μ l, was added to each of the
25 remaining wells (#2-12) in the row. The diamine(s) solution in "well one" was serially diluted by transferring 25 μ l from "well one" to "well two", and repeating a 25 μ l transfer from "well two" to "well three", and so on, on through "well eleven". In "well eleven", the extra 25 μ l of solution was discarded. "Well twelve" was used as a growth control to assess background activity of the reporter strain. The plate was then covered and incubated
30 at 37 °C for 24 hours. Immediately prior to analysis, the following substrates were prepared: a buffer solution containing 50 mM HEPES at pH 7.0 and 0.4% Triton X-100. Then, 0.25 ml of 1M DTT, and 14 μ l of 10 mg/ml luciferin in DMSO were added to 5 ml of the buffer solution. This final solution (50 μ l) was added to each of the twelve wells,

immediately after the incubation period had run. The luminescence from each well was measured 20 minutes after the luciferin substrate was added, using a TOPCOUNT® (Downers, Grove, Illinois) NXT luminometer (55/well).

Figures 6-8 show typical assay data for the luciferase reporter strain containing an Rv0341 EMB-inducible promoter with serial dilution of 12 wells (1 row) of a 96-well library plate. Figure 10 shows the number of substituted ethylene diamines with at least 10% luciferase activity, based on the activity of ethambutol at 3.1µM.

Figure 6 represents typical assay data in the luciferase reporter strain containing an Rv0341 EMB-inducible promoter. The data represents values obtained from the HTS Luc assay for compound mixtures of one row (row D) in the 96-well library. Row D was subject to several serial dilutions. The effectiveness of the compound mixture in the assay was measured by the intensity of luminescence, and compared to ethambutol (100% intensity, 99% purity) at 3.1µM. Each curve in Figure 6 represents one well, or ten compounds. Results are reported in percent maximum Luminescence Count per Second (% Max. LCPS). During the screening, a theoretical 100% chemical yield was assumed for every unpurified compound. Concentrations are given for a single compound. Based on this initial screening, 300+ compound mixtures showed anti-TB activity.

EXAMPLE VIII

Representative MIC Experiment

The Minimum Inhibition Concentration (MIC) is the concentration of the growth inhibitor, here a substituted ethylene diamine, at which there is no multiplication of seeded cells. A microdilution method was used to determine the MIC of the substituted ethylene diamines, capable of inhibiting the growth of *Mycobacterium tuberculosis in vitro*. In a representative MIC experiment, bacteria, the H37Rv strain of *Mycobacterium tuberculosis* (*M.tb*), was cultivated in 7H9 medium to a density of 0.2 OD (optical density) at 600 nm. The bacterial culture was then diluted 1:100 in 7H9 broth medium. Stock solutions of isoniazid and ethambutol were each prepared at 32 µg/ml in 7H9 medium. A 3.2mg/ml solution of isonizid and ethambutol were each prepared in water. The solutions were then filtered, and diluted 1:100 in 7H9 medium. Each drug, purchased from Sigma, was “laboratory use only” grade. A 10mM solution of each substituted ethylene diamine was

prepared in methanol. Next, 100 μ l of the 7H9 medium was added to each well in a 96-well plate (rows (A through H) x columns (1 through 12)). To the first wells in rows C through H was added an additional 80 μ l of the 7H9 medium. The isoniazid solution, 100 μ l, was added to well A1, and the ethambutol solution, 100 μ l, was added to well B1. Six substituted ethylene diamines, 20 μ l each, were added to wells C1 through H1 (column 1), respectively. A serial dilution of each substituted ethylene diamine and the isoniazid and ethambutol controls, was performed across each row. For example, a serial dilution across row C1-C12 was done by mixing and transferring 100 μ l of the previous well to the next consecutive well. In each well in "column 12," 100 μ l of the final dilution was discarded.

Next, 100 μ l of the diluted H37Rv strain of *M.tb* was added to each well. The 96-well plate was then covered and incubated at 37 °C for 10 days. The plate was read for bacterial growth, or non-growth, using an inverted plate reader. The MIC was determined to be the lowest concentration of substituted ethylene diamine that inhibited visible growth of the *M.tb*.

A representative plate layout, listing concentration in each well, is shown in Table 9. Table 10 lists MIC and LD50 data for selected compounds. The LD50 is the concentration of the substituted ethylene diamine at which 50% of the cells (H37Rv strain of *M.tb*) are killed. Table 11 lists MIC data for purified substituted ethylene diamines in comparison to ethambutol (EMB). Figure 9 shows the number of substituted ethylene diamine compounds with MIC activity at various concentration levels.

Table 9

Concentration in Each Well (μ M) Based on Columns 1-12

DRUG

soniazid	58.2 5	9.1	4.5	.28	.64	.82	.91	.45	.23	.11	.06	.03
thambutol	28.7 5	4.3	.19	.60	.80	.90	.45	.22	.11	.06	.03	.01
ubst. thylene iamine	500	50	25	2.5	1.2	5.6	.81	.91	.96	.98	.49	.24

Table 10

Selectivity Index for Selected Compounds

Cmpd	MIC (uM)	LD50(uM)	MW	MIC (ug/ml)	LD50 (ug/ml)	SI
6	7.813	20	536	4.187768	10.72	2.559836
34	7.813	32	508	3.969004	16.256	4.095738
37	15.625	32	496	7.75	15.872	2.048
47	15.625	25	452	7.0625	11.3	1.6
57	15.625	18	426	6.65625	7.668	1.152
59	15.625	32	426	6.65625	13.632	2.048
65	15.625	60	430	6.71875	25.8	3.84
109	1.953	32	450	0.87885	14.4	16.38505
111	7.813	44	412	3.218956	18.128	5.63164
151	7.813	41	450	3.51585	18.45	5.247664

5

The above procedure was also used to examine batched compounds (10 compounds per well). Synthesized batches of substituted ethylene diamines were dried in speed vacuum and then resuspended in DMSO or sterile water to a concentration of 2.5 mg/ml.

10

TABLE 11

MIC Data for Purified Samples

Plate set-up

INH								
58.25	29.125	14.56	7.28	3.64	1.82	0.91	0.45	0.23
EMB								
28.75	14.375	7.1875	3.594	1.797	0.898	0.449	0.2245	0.1125
CMPD								
500	250	125	62.5	31.25	15.625	7.813	3.9063	1.953

Avg INH MIC (uM)	Avg INH MIC (uM)		
0.91	0.91		
Avg EMB MIC (uM)	Avg EMB MIC (uM)	Avg EMB	Avg EMB
7.1875	8.37	7.25	7.25

15

Cmpd	MIC (uM)				BACTEC (EMB: 2.5UG/ML)
1	250	250	125	125	
2	250	250	250	250	
3	31.25	62.5	15.6	15.6	
4	125	62.5	62.5	62.5	

Cmpd	MIC (uM)				BACTEC (EMB: 2.5UG/ML)
5	>500	500	500	500	
6	7.813 *	7.813	3.9	3.9	
7	15.625 *	7.813	3.9	3.9	
8	125	125	31.25	31.25	
10	7.813 *	15.625	7.8	7.8	
11	31.25	contaminated	3.9	3.9	
13	31.25	31.25	15.6	15.6	15
14	15.625 "	15.625	7.8	7.8	
15	>500	>500	250	500	
17	62.5	62.5	15.6	15.6	
21	15.625 *	31.25	7.8	7.8	
22	31.25	31.25	7.8	15.6	
23	31.25	31.25	15.6	15.6	
24	125	125	31.25	31.25	
27	125	62.5	15.6	31.25	
28	125	62.5	31.25	31.25	
29	62.5	62.5	31.25	62.5	
31	31.25	61.25	15.6	15.6	
32	15.625*	15.625	7.8	7.8	
33	62.5	62.5	31.25	31.25	
34	7.813 *	7.813	3.9	3.9	
35	62.5	62.5	15.6	31.25	
36	31.25	62.5	15.6	15.6	
37	15.625 *	15.625	3.9	7.8	1.25
38	7.813	7.813	3.9	7.8	
40	15.625 *	15.625	7.8	7.8	
41	31.25	15.625	15.6	15.6	
42	31.25	31.25	1.95	3.9	
43	31.25	31.25	3.9	7.8	12.5
47	15.625 *	15.625	1.95	7.8	5
51	31.25	250	31.25	31.25	
52	15.625 *	15.625	3.9	3.9	
53	31.25	31.25	31.25	31.25	
54	31.25	31.25	15.6	31.25	
55	15.625 *	15.625	15.6	15.6	25
56	500	>500	500	500	
57	15.625 *	7.813	7.8	7.8	
58	15.625 *	15.625	7.8	7.8	5
59	15.625 *	31.25	15.6	15.6	12.5
61	62.5	62.5	31.25	31.25	


Cmpd	MIC (uM)				BACTEC (EMB: 2.5UG/ML)
62	15.625 *	31.25	15.6	31.25	
63	62.5	62.5	31.25	62.5	
64	31.25	31.25	31.25	31.25	
65	15.625 *	31.25	31.25	31.25	
66	15.625 *	15.625	7.8	7.8	
68	500	500	500	500	
71	62.5	62.5	31.25	31.25	
73	62.5		15.6	15.6	
76	62.5	62.5	31.25	31.25	
77	31.25	31.25	15.6	15.6	
78	15.625 *	31.25	15.6	15.6	
79	31.25	31.25	15.6	15.6	
103	31.25	31.25	62.5	62.5	
107	500	500	250	250	
109	1.953 *	1.953	1.95	1.95	0.63
111	7.813 *	7.813	7.8	7.8	5
116	15.625 *	15.625	7.8	15.6	12.5
117	7.813 *	15.625	7.8	7.8	
118	31.25	62.5	31.25	no data	
119	125 contam	62.5	cont	no data	
125	15.625 *	15.625	cont	no data	6.25
134	>500	>500	500	no data	
151	15.625 *	7.813	cont	no data	6.25
164	62.5	125	cont	no data	
165	62.5	62.5	15.6	15.6	

EXAMPLE IX


Secondary Screening and Evaluation of Substituted Ethylene Diamines Against
Drug Resistant Patient Isolates

- 5 Secondary screening was performed on some of the substituted ethylene diamine compounds to examine their activity against three clinically resistant MDR patient isolates. MDR Strain TN576 is classified as a W1 strain (STP^R, INH^R, RIF^R, EMB^R, ETH^R, KAN^R, CAP^R) strain TN587 is classified as a W strain (STP^R, INH^R, RIF^R, EMB^R, KAN^R), and the third strain TN3086 is classified as a W1 strain (STP^R, INH^R, RIF^R, EMB^R, KAN^R). Each
- 10 MDR strain is highly resistant to ethambutol with MIC values exceeding 12.5-25μM. The

MICs for the following substituted ethylene diamines, MP 116, MP 117, RL 241, compounds #59 and #109, were determined for all three patient isolates.




Compound MP 116




Compound MP 117

5




RL241 (compound 37)



Compound 59

10



Compound 109

The results from this study are shown in Tables 12-13. Table 14 characterizes each MDR strain according to its resistance.

5

Table 12

Screening of Substituted Ethylene Diamines Against
Drug Resistant Patient Isolates – (MIC values in ug/ml)

	WT	576	587	3806
EMB	3.12 (or 11.1 uM)	12.5 - 25	12.5 - 25	12.5 - 25
MP 116	6.25	3.15	6.25	3.15
MP 117	6.25	3.15	3.15	3.15
RL 241	1.5 (or 3.34 uM)	1.5	1.5	1.5

10

WT= wild type of M.tb

EMB as 2HCl salt

RL241 as 2HCl salt

Table 13

15

Screening of Substituted Ethylene Diamines Against
Drug Resistant Patient Isolates – (MIC values in ug/ml)

	<u>WT</u>	576	587	3806
EMB	1.6-1.8	50	50	50
Cmpd#59	0.05 (or 0.13 uM)	0.1	0.05	0.05
Cmpd#109	0.10 (or 0.18 uM)	0.2	0.2	0.1

Cmpd#59 as a 2HCl salt

20

Cmpd#109 as a 2CF₃COOH salt

Table 14
Drug Resistance of Each MDR Strain

Strain	STP	STP 2	NH	INH 2	Rif	Em b	Eth	Ka n	Cip	Cap	Cyc
576 W1	R	R	R	R	R	R	R	R	S	R	S
587 W	R	R	R	R	R	R	S	R	S	S	S
3806 W1		R		R	R	R	S	R			

R = resistant

S = susceptible

5 STP = Streptomycin

INH = Isoniazid

Rif = Rifampicin

Emb = Ethambutol

Eth = Ethionamide

10 Kan = Kanamycin

Cip = Ciprofloxacin

Cap = Capreomycin

Cyc = Cycloserine

15

EXAMPLE X

In Vivo Animal Studies

Animal models were used in the final stages of the drug discovery cycle to assess the anti-microbial efficacy of some substituted ethylenediamine compounds in a representative system of human disease state. The *in vivo* testing approach involves the inoculation of four-six week old C57BL/6 mice via aerosol, containing approximately 200 colony forming units of *M. tuberculosis* H37R_V.

A. CFU Lung Study

Mice aerosolized with *M. tuberculosis* H37R_V were examined for 10 to 12 weeks following inoculation. Drugs (substituted ethylene diamines) were administered via the esophageal cannula (gavage) 7 days/week, starting at either 14 or 21 days post infection. Bacterial load in the lungs of five mice per group were determined at approximately one-

week intervals by viable colony counts. The drugs tested were directly compared to the front line anti-tuberculosis drug isoniazid, and to the second line drug, ethambutol. Isoniazid and ethambutol were tested at 25 mg/kg and 100 mg/kg, respectively. The substituted ethylene diamines, compound 37, compound 59 and compound 109, were each
5 tested at 1 mg/kg and 10 mg/kg. Figures 17 to 19 represent data from three, independent CFU Lung studies. In each study, the number of colony forming units (CFU) that were recoverable and cultivatable, were determined during various time intervals (days).

B. Lesion Study

The ability of compound 59 and compound 109 to prevent the development of gross
10 pathology due to bacterial burden was determined in conjunction with the CFU/Lung Study. The gross pathology was determined by visible quantitation of lesions on the surface of the lungs. Quantitation by inspection is a good surrogate for CFU determination, and directly correlates to the bacterial burden, as determined by the actual colony forming units. The lesions are first visibly examined, and then the lungs are processed and plated
15 for CFU quantification. The lesion study demonstrates the ability of the drug to prevent the development of the disease pathology. Figure 20 represents data from a lesion study. The corresponding CFU results are shown in Figure 19.

C. Toxicity Study

Toxicity was assessed using a dose escalation study. This study was performed
20 with ten C57BL/6 mice per candidate. Every two days, the mice were administered an increased concentration of the drug, and monitored for detrimental effects. The administration scheme was 50, 100, 200, 400, 600, 800 and 1000 mg/kg. The maximum limit of 1000 kg/mg was based on the goal of dose escalation, and the solubility of the drugs in the delivery vehicle. Compound 37 was toxic in mice at 100 kg/mg. Compound
25 59 and compound 109 were tolerated in mice at 1000 mg/kg and 800 mg/kg, respectively.

It should be understood that the foregoing relates only to preferred embodiments of the present invention, and that numerous modifications, or alterations, may be made therein without departing from the spirit and scope of the invention. The entire text of each reference mentioned herein is hereby incorporated, in its entirety, by reference.

EXAMPLE XI

In vitro toxicity and Selectivity Indexes for hit compounds

5 Twenty six compounds (including 37, 59 and 109) were tested in an *in vitro* model of toxicity using monkey kidney cells (Vero) and human cervical cancer cells (HeLa) using methods well known to those skilled in the art. The data from this toxicity testing and the MIC data were used to calculate a selectivity index (SI), the ratio of IC₅₀:MIC (Table 15). Selectivity Indexes were ranging from 1.76 to 16.67. Compound 109 has the best
10 selectivity index.

Table 15
In vitro data for representative compounds.

Compound	MIC (μM)	Vero IC ₅₀ (μM)	SI (IC ₅₀ :MIC)
66	15.6	28	1.76
40	15.6	25	1.88
41	3.13	19	2.05
59	15.6	36	2.30
55	15.6	34	2.32
57	11.7	22	2.40
37	7.8	32	4.10
38	6.25	33	5.28
111	7.81	45	5.76
73	12.5	81	6.48
58	12.5	82	6.56
78	15.6	130	8.33
109	1.56	26	16.67

15

Example XII

In vivo efficacy of ethambutol analogues

20 Compounds 58, 59, 73, 109, and 111 were selected for *in vivo* efficacy studies in a mouse model of TB. Compounds 58 and 59 share the same cyclooctyl fragment in their molecules; compounds 58, 73, and 109 share adamantly moiety, and 109 and 111 – the geranyl fragment (Figure 22).

25 In these studies, 8-week old inbred female mice C57BL/6 were intravenously infected with *M. tuberculosis*. 3 weeks following infection drug treatment was initiated (detailed protocol is provided). The drugs were administered orally by gavage. Mice were sacrificed at three timepoints (15, 30, and 45 days post infection), and CFUs in spleen and

lungs were determined (Figures 23 and 24). These studies demonstrated that compound 109 had activity at doses 1 and 10 mg/kg equal to that of ethambutol at 100 mg/kg.

Materials and Methods

- 5 Mice. Female C57BL/6 mice of 8 weeks old were purchased from Charles River (Raleigh, NC), housed in BSL-2 facility of BIOCAL, Inc. (Rockville, MD), and were allowed to acclimate at least 4 days prior infection.

10 Mycobacteria. An example of frozen and thawed of *M.tuberculosis* H37Rv Pasteur was added to 5 ml 7H10 broth medium, with 0.5% BSA and 0.05% Tween 80, incubated 1 week at 37° C, and then 1 ml was added into 25 ml medium (2-d passage during 2 weeks). Culture was washed twice and resuspended in PBS with 0.5% BSA and 0.05% Tween 80, aliquoted and frozen at -80° C. To determined CFU of the culture aliquot was thawed, and 10-fold dilutions will be plated on agar 7H9 and CFU count will be calculated 20 days later.

15

Infection: Frozen sample of culture was thawed, and diluted for concentration about 10⁶ CFU/ml. Mice were infected with *M.tuberculosis* H37Rv intravenously through lateral tail vein in corresponded dose in 0.2 ml of PBS.

- 20 Antimicrobial agents. INH, EMB, Ethambutol analogues.

Protocol of drug treatment: Treatment of mice with compounds was initiated 20 days following infection. Compounds were dissolved in 10% ethanol in water and administered by gavage (0.2 ml per mouse). Therapy was given 5 days per week and continued for four or six weeks. Two, four and six weeks following chemotherapy start mice (6 mice per group) were sacrificed, lungs and spleens were removed and homogenized in sterile in 2 ml PBS with 0.05% Tween-80. Homogenates were plated in serial dilutions on 7H10 agar dishes, and incubated at 37° C. CFU counts were calculated three weeks later.

25 Statistic analysis. To analyze results of CFUs in organs ANOVA test was performed; the significance of the differences was estimated by Student's test, p<0.05 was considered statistically significant.

30

Results

In vivo activities of new compounds. The activities of these compounds are presented in figures 21-24. In the experiment presented in figures 21 (spleen) and 22 (lung) mice were infected with 5×10^5 CFU *M.tuberculosis* H37Rv and chemotherapy was started 20 days following infection. Mice were treated with INH (25 mg/kg), EMB (100 mg/kg), compounds 73 and 109 (both 1mg/kg and 10 mg/kg). The results indicate that in the spleen, compounds 73 and 109 have activities equal to that of EMB at 100 mg/kg (Figure 21). In spleen there are no statistical differences between activities of these compounds at 1 mg/kg or 10 mg/kg. In the lung, compound 109 at concentration 10 mg/kg after 4 and 6 weeks was more effective than EMB at 100 mg/kg. In the lung, statistically sufficient difference was shown for compound 109 at concentrations 1 mg/kg and 10 mg/kg (Figure 22). INH was the most active drug in both spleen and lung.

Compounds 73 and 109 were also tested in shorter model with using higher dose of infection (Figures 23 and 24). Mice were infected with 5×10^6 CFU *M.tuberculosis* H37Rv and chemotherapy was started 15 days following infection. Mice were treated with INH (25 mg/kg), EMB (100 mg/kg), compounds 109 (0.1 mg/kg, 10 mg/kg, and 25 mg/kg), 58, 73 and 111 (all 25 mg/kg). Mice were treated for 4 weeks. In both the spleen and lung, compound 109 at concentrations 10 mg/kg and 25 mg/kg had activity equal to that of EMB at 100 mg/kg, and at concentration 0.1 mg/kg minimal but sufficient difference with untreated control appeared after 4 weeks of therapy (Figure 23 and 24). Statistically sufficient difference between compounds 73 (25 mg/kg) and 109 (25 mg/kg) was detected. In the lung significant difference between activities of these compounds was not detected. Compounds 58 and 111 are active *in vivo* in both spleen and lung; however, compounds 73 and 109 are preferable. The results of these experiments indicate that compounds 73 and 109 in low concentration show activity equal that of EMB at 100 mg/kg, and in some cases compound 109 shows higher activity.

Testing of compounds 111 and 59 was performed in B6 mice infected with 5×10^5 CFU *M.tuberculosis* H37Rv and beginning chemotherapy 20 days following infection (Figure 25 and 26). Both compounds showed anti-tuberculosis activity at concentration 10 mg/kg comparable to that of EMB at 100 mg/kg.

In all experiments, INH showed higher activity than EMB and other compounds decreasing load of bacteria in organs on 2 - 3 logs during 4 – 6 weeks of chemotherapy; new compounds similar to EMB (100 mg/kg) decreased load of bacteria on 1.0–2.0 logs.

Among studied compounds 73 and 109 are the most preferable, because the highest capacity to decrease mycobacteria in organs and its parameters of toxicity and pharmacology kinetics.

5

EXAMPLE XIII

In vivo toxicity

Preliminary dose acceleration studies in mice have indicated that compound 109
10 can be well tolerated at doses up to 800 mg/kg and compound 59 up to 1000 mg/kg. Compound 37 was fatal at doses 100 mg/kg (Clif Barry, NIAID, unpublished results).

Compound 109 was mostly used in the form of dihydrochloride at five different doses, and 37 – solely as hydrochloride salt at two doses.

Mice were given a one-time dose of the compounds at concentrations 100, 300 or
15 1000 mg/kg using the gavage method. Each dose of each compound consisted of one group of 3 mice. Monitoring of the mice was done twice a day for the duration of the experiment. Mice surviving one week post-drug administration were sacrificed; critical organs were aseptically removed and observed for abnormalities and evidence of drug toxicity. The MTD (mg/kg) is the highest dose that results in no lethality/tissue
20 abnormality.

Methods:

1. Treatment of mice: C57BL/6 female mice (6-8 weeks in age) are given a one-time dose of the compound at concentrations 100, 300 or 1000 mg/kg using the gavage method. The compounds are dissolved in the appropriate concentration of ethanol
25 in distilled water and administered in a volume of 0.2ml per mouse.
2. Observation of mice: Mice will be observed 4 and 6 hours post administration, then twice daily for one week. Survival and body weight of mice will be closely monitored throughout the study.
3. Assessment of drug toxicity: Mice exhibiting signs of any abnormal appearance or
30 behavior or those remaining in a group in which other mice did not survive to day 7 will be sacrificed for assessment of drug toxicity. Critical organs will be aseptically removed and observed; tissues from the liver, heart, and kidneys are extracted and

placed into 10% formalin solution. These fixed tissues are sectioned and examined for abnormalities resulting from drug toxicity.

These studies indicate that the maximum tolerated dose for the compound 109 is 600 mg/kg (Table 16). No visible changes in organs were observed. Dose 800 mg/kg was fatal: out of a group of 3 mice, two animals died within 3 days (Table 17). Compound 37 was well tolerated at doses 100 and 300 mg/kg. No visible changes in organs were observed. Additional experiments to evaluate maximum tolerated dose and *in vivo* efficacy for the compound 37 are being conducted.

10

Table 16

Determination of a maximum tolerated dose for the compounds 109 and 37 in mice.

		109 at 100 mg/kg		109 at 300 mg/kg		109 at 600 mg/kg		109 at 1000 mg/kg		37 at 100 mg/kg	
Day		Mice	Day of death	Mice	Day of death	Mice	Day of death	Mice	Day of death	Mice	Day of death
04.08.03	1	3		3		3		3		2- 4h	1
04.09.03	2	3		3		3		2	2	2	
04.10.03	3	3		3		3		2		2	
04.11.03	4	3		3		3		1	4	2	
04.13.03	6	3		3		3		0	6	2	
04.14.03	7	3		3		3		-		2	

15

Table 17

Determination of a maximum tolerated dose for the compounds 109 and 37 in mice.

Date		37 at 100 mg/kg		37 at 300 mg/kg		109 as HCl salt at 800 mg/kg		109 as TFA salt at 800 mg/kg	
Day		Mice	Day of death	Mice	Day of death	Mice	Day of death	Mice	Day of death
04.29.03	1	3		3		3		1	
04.30.03	2	3		3		2/1	2	1	
05.01.03	3	3		3		1/1	3	1	
05.02.03	4	3		3		1		1	
05.03.	5	3		3		1		1	
05.04	6	3		3		1		1	
05.05.03	7	3		3		1		1	

20

EXAMPLE XIV

Pharmacokinetic studies of the compounds 37, 59, and 109

Initially, analytical methods for determination of the compounds had been developed that allowed to carry out all the PK experiments, see Figure 29. Here is a brief description of the experiment: (1) plasma spiked with tested compounds and 10 uL of Terfenadine or plasma samples (200 uL) added; (2) ACN (2 mL) added to precipitate protein and spin at 2,500 rpm; (3) evaporate supernatant to dryness; (4) add 200 uL of the diluting solvent: methanol (with 0.1% of trifluoroacetic acid) : ammonium acetate (80/20); (5) vortex, spin, and use supernatant; (6) run LC/MS/MS on Sciex API 3000.

Biostability studies of the compounds in plasma were carried out using concentrations 1 and 15 mg/ml. The compounds were incubated for 1, 2, 3 & 6 hr at 37°C (Table 18). In addition, it was found that all tested compounds were stable in plasma at 24°C, pH 2 and 7.4 up to 24 hr.

Table 18
Biostability of tested compounds in plasma.

Comp.	Human	Dog	Rat	Mouse
37	20%↓	stable	35% ↓	stable
59	stable	stable	stable	stable
109	30% ↓	40% ↓	stable	stable

Pilot PK study of the compounds 37, 59, and 109 in mice was conducted using a cassette dosing: all the three analogs were formulated together in saline at 1.5 mg/mL, and administered to mice simultaneously orally at 25 mg/kg, peritoneally at 6 mg/kg, and intravenously. It was found that doses 15 and 7.5 mg/kg caused death of mice, 3.75 mg/kg appeared lethargic immediately after dosing but then appeared normal appearance a few minutes later; 3 mg/kg displayed no adverse reactions and hence was used as intravenous dose. Obtained data are presented on Figures 30, 31, and 32 (tested compounds were studied under the NCI' indexation NSC) and summarized in Table 19.

Table 19
PK Parameters of tested compounds 37, 59, and 109 after a
cassette dosing to mice. N/A- not detectable.

Route	i.v.	i.v.	i.v.	i.p.	i.p.	i.p.	p.o.	p.o.	p.o.
Compounds	37	59	109	37	59	109	37	59	109
Dose (mg/kg)	3	3	3	6	6	6	25	25	25
AUC(ng·h/mL)	954	384	100 6	137 2	272	109 9	160 2	169	655
Cmax (ng/mL)	970	296	119 2	630	217	935	263	28.7	227
T1/2 (h)	4.8	6.4	5.5	4.9	9.7	4.4	N/A	N/A	N/A
CL (mL/kg/h)	353 0	804 3	324 0						
Bioavailability (%)				72	35	55	3.3	0.9	2.7
Urine excretion (%)	.71	1.9	.92	<0.0 1	<0.0 1	<0.0 1	N/A	N/A	N/A

5

Conducted pharmacokinetic studies indicated that compound 59 (NSC 722040 by the NCI index) has relatively poor PK profiling (AUC, Cmax) and further testing of this compound was abandoned. Based on preliminary toxicity data compound 37 was also ruled out as possible candidate. Therefore, compound 109 (NSC 722041 by the NCI) was selected for further PK analyses.

It has been shown that compound SQ109 reaches and exceeds its Minimum Bactericidal Concentration MBC (313 ng/ml) in plasma when administered either iv or intravenously orally (p.o.), has a half-life of 5.2 h, and has total clearance less than hepatic blood flow (Figure 33, Table 20).

15

Table 20

Pharmacokinetic
the compound 109.

parameters of

Parameters	i.v.	p.o.
Dose (mg/kg)	3	25
AUC (ng•h/mL)	792	254
T _{1/2 el} (h)	3.5	5.2
C _{max} (ng/mL)	1038	135
T _{max} (h)	0	0.31
CL (mL/kg/h)	3788	
Vd _{ss} (mL/kg)	11826	
Bioavailability		3.8

Its oral bioavailability is only 3.8% when administered p.o but this is explained by its unique tissue distribution pattern. Tissue distribution studies have demonstrated that SQ109 primarily distributes into the lungs and spleen (Figures 34 and 35), which is highly advantageous for a infection that characteristically manifests as a lung disease.

By using an ultracentrifugation method, it was found that plasma protein binding of the compound 109 is concentration dependent and varies from 15% (20 ng/ml) to 74% (200 ng/ml) to 48% (2000 ng/ml). After i.v. dosing (3 mg/kg) the compound distributes between plasma and red blood cells in a ratio 70.6 : 29.4.

Little is known of the fate of the compound in the body, since the total amount of the compound after excretion (urine and feces) does not exceed 3% of the delivered dose (Table 2).

Table 21
Amounts of the compound 109 cumulatively excreted in mouse urine and feces following single administration.

Dose/ Route	Samples	Period after dosing (h)				
		Total 0-4	4-8	8-24	24-32	0-32
3 mg/kg i.v.	Urine	<0.01	<0.01	0.03	0.01	0.04
	Feces	<0.01	0.01	0.04	<0.01	0.06
25 mg/kg p.o.	Urine	-	-	-	-	-
	Feces	0.48	0.31	1.12	0.08	2.0

Initial attempts to identify metabolites of the compound 109 in urine, did not provide evidence of breakdown products, Figure 36. For example, there was no evidence for the formation of conjugated metabolites (M^+ 521) in the mouse urine during first 24 hr following compound's administration, Figure 37. Conjugated metabolites are products of the typical metabolic pathway N-glucoronidation formed by reaction with glucuronic acid (D.A. Williams and T.L. Lemke in *Foye's Principles of Medicinal Chemistry*, 5th Ed., p.202).


EXAMPLE XV

In vitro Pharmacokinetic Studies of Compound 109

In vitro Pharmacology and early ADMET (Absorption, Distribution, Metabolism, Excretion, Toxicity) studies of the compound 109 were contracted out to CEREP (15318 NE 95th Street, Redmond, WA 98052, USA, www.cerep.com, tel 425 895 8666) under a Service Agreement and included testing against 30 standard receptors (see CEREP Tables 22 and 23, provided in Figures 38 and 39, five CYP450 enzymes, hERG (K⁺ channel), aqueous solubility, predicted intestinal permeability, and metabolic stability (data presented in Figure 40 Tables 24(a-m)).

EXAMPLE XVI

Bis(2-Adamantyl)ethylenediamine, SQBisAd

Compound **SQBisAd**

Compounds with the best Selectivity Indexes, such as 109, 58, 73, 78, (Table 15) and good *in vivo* data share the same adamantane fragment (Figure 20). A compound that would have solely this fragment (on both sides of the ethylene linker) was contemplated. During preparation of targeted 100,000 compound library of ethambutol analogues, 70,000 compounds were proven to be formed, but 30,000 were failures. This particular compound was not initially detected perhaps because it was synthesized in very low yield or because it was never made due to steric factors.

In the synthetic scheme used for preparation of the library Scheme 1 (Figure 41), sterically hindered amines on the second step rarely gave products. Analyzing MS data for a number of original plates it can be stated that 2-adamantanamine when used as R_1NH_2 seldom yield desirable products and this can be explained because of existence of sterically hindered reaction site on the step 2 or step 3 of the synthesis Scheme 2 (Figure 41).

Compound SQBisAd can be prepared by "wet chemistry" using the same route, Scheme 3 (Figure 41), it is documented that 2-adamantanamine (used as commercially available hydrochloride) does provide products when used on the 1 and 2 steps. Due to the symmetrical nature, this compound can be synthesized by alternative routes. We have prepared SQBisAd by reductive alkylation of ethylenediamine by 2-adamantanone using sodium cyanoborohydride. Final product (without additional purification) demonstrated MIC (Minimal Inhibitory Concentration) equal or better than compound 109.

EXAMPLE VIII

Generating the Diamine Library with a Modified Linker

General Methods: All reagents were purchased from Sigma-Aldrich. Rink acid resin was purchased from NovaBiochem, Inc. Solvents acetonitrile, dichloromethane,

dimethylformamide, ethylene dichloride, methanol, and tetrahydrofuran were purchased from Aldrich and used as received. Solid phase syntheses were performed on Quest 210 Synthesizer (Argonaut Technologies) and combinatorial chemistry equipment (Whatman Polyfiltronics and Robbins Scientific). Evaporation of the solvents was done using
5 SpeedVac AES (Savant). Mass spectra data were obtained by Electrospray Ionization technique on Perkin Elmer/Sciex, API-300, TQMS with an autosampler.

The activation of the Rink-resin, the addition of the amine, and the acylation step were carried out in 10 ml tubes using the Quest 210 Synthesizer. Removal of the Fmoc
10 group, reductive alkylation reaction with carbonyl compounds, the reduction with Red-Al, and the cleavage from the solid support were carried out in 96-deep (2ml) well, chemically resistant plates.

Step 1. Activation of the Rink-acid resin.

15 A suspension of the Rink-acid resin (coverage of 0.43 – 0.63 mmol/g), 6 g (up to 3.78 mmol), in 80 ml of 2:1 mixture of dichloromethane and THF was distributed into 20 tubes, 4 ml per tube, filtered and washed twice with THF. A solution of triphenylphosphine (5.7 g, 21.75 mmol) in 40 ml of THF was added, 2 ml/tube, followed by the addition of a solution of hexachloroethane (5.09 g, 21.45 mmol) in 20 ml of THF, 1
20 ml/tube. After 6 h the resins were washed with THF (2x4 ml) and dichloromethane (2x4ml).

Step 2. Addition of the first amine.

Each tube was charged with 3 ml of dichloroethane, EtNiPr₂, (0.2 ml, 1.15 mmol),
25 and the corresponding amine (1 mmol). (When a selected amine was a solid, it was added as a solution or a suspension in DMF). Dichloroethane was added to each tube to fill up the volume 4ml. The reaction was carried for 8h at 45°C and 6-8 h at room temperature. The resins were filtered, washed with a 2:1 mixture of dichloromethane and methanol (1x4ml), then with methanol (2x4ml), and suck dry.

30

Step 3. Acylation with Fmoc protected amino acid.

The resins were pre-washed with dichloromethane (2x4ml). Each tube was charged with 2 ml of dichloromethane, HATU (2 mol excess to loaded resin, 0.14 g, 0.39 mmol, dissolved in 1 ml of DMF), and 0.47 mmol (2.5 mol excess to loaded resin) of amino acid dissolved in 1 ml of DMF, and allowed to stir for 8h at 45⁰C and 6-8 h at room temperature. After 16 h the resins were filtered, washed with 1:1 mixture of DMF and dichloromethane (1x3ml), dichloromethane (1x3ml) and acylation was repeated with the same amount of reagents. At the end, the resins were filtered, washed with 1:1 mixture of DMF and dichloromethane (1x3ml), and methanol (3x3ml), sucked dry (on Quest) for 30 min and transferred into vials (one resin per vial), and dried in a desiccator under vacuum for 1h. After this step all resins were subjected for quality control using MS spectra.

Step 4. Alkylation of the amino group.

Deprotection. Ten prepared resins from the first three steps were pooled together, leaving approximately 0.05 g of each in the individual vials for all necessary deconvolutions. A suspension of the resin mixture (2.0–2.5 g) in 100 ml of a 2:1 mixture of dichloromethane and THF was distributed into two 96-well filterplates and filtered using a filtration manifold. The reaction plates were transferred into combiclamps, and 0.2 ml of 20% solution of piperidine in DMF was added to remove Fmoc protecting group and allowed to stay for 10 min. After 10 min plate was filtered, washed with 0.2 ml of DMF, and deprotection was repeated with 0.2 ml of 20% solution of piperidine in DMF and allowed to stay for 20 min. After that plate was filtered, washed with DMF (0.2 ml per well) and dichloromethane (2x0.5 ml per well).

Reaction with the carbonyl compounds. Each well in row A on the reaction plate was charged with 0.1 ml of dichloromethane, 0.08ml of ~1.0M solution of appropriate acid in DMF from master plate, 0.05 ml DMF solution of PyBrop, (0.015 g, 0.03 mmol, 2.5 mol excess to loaded resin) and 0.05 ml of EtNPr₂ in dichloromethane (0.022 ml, 0.13 mmol, 10 mol excess to loaded resin). Each well in rows B through H was charged with 0.1 ml of THF, 0.160 ml of ~1.0 M solution of appropriate aldehyde or ketone in DMF from master plate and allowed to react for 30 min. After 30 min 0.075 ml (0.075 mmol) of 1.0 M

solution of NaBCNH₃ were added. The reaction plates were sealed and kept at RT for 72h. At the end, the resins were filtered, washed with THF, DCM (1x1ml), methanol (2x1ml) and dried in desiccator under vacuum for 2h.

5 Step 5. Reduction with Red-Al.

The reaction plates were placed into combiclamps. A 1:6 mixture of Red-Al (65+ w% in toluene) and THF was added, 0.6 ml per well (0.28mmol of Red-Al per well), and allowed to react for 4 h. After the reaction completion the resins were filtered, washed with THF (2x1ml), methanol (3x1ml) and dried in the filtration manifold.

10

Step 6. Cleavage.

This step was carried out using a cleavage manifold. The reaction plates (placed on the top of the collection plates in this manifold) were charged with a 10:85:5 mixture of TFA, dichloromethane, and methanol, 0.5 ml per well. After 15 min, the solutions were
15 filtered and collected into proper wells of the collection plates. The procedure was repeated. Solvents were evaporated on a speedvac, and the residual samples were ready for testing.

Deconvolution Example.

20 Deconvolution of the active wells was performed by re-synthesis of discrete compounds, from the archived Fmoc-protected α -aminoacetamide resins (10 resins, 0.05-0.10g each), which were set aside at the end of the acylation step before the pooling. Each resin was assigned a discrete column (1, or 2, or 3, etc.) in a 96-well filterplate, and was divided between X rows (A, B, C, etc), where X is the number of hits discovered in the
25 original screening plate. To each well, in a row, a selected carbonyl compound (present in the hit) was added along with other required reagents: the first selected carbonyl compound was added to the resins in the row "A", the second carbonyl compound – to the resins in the row "B", the third carbonyl compound – to the resins in the row "C", etc. A lay-out of a representative 96-well deconvolution plate is shown in Table 28, Figure 52.

The reaction plates were sealed and kept at RT for 72h. At the end, the resins were filtered, washed with THF, DCM (1x1ml), methanol (2x1ml) and dried in desiccator under vacuum for 2h. Reduction and cleavage were performed according to steps 5 and 6 of the synthetic protocol. The product wells from the cleavage were analyzed by ESI-MS (Electrospray Ionization Mass Spectroscopy) to ensure the identity of the actives, and were tested in the MIC assay. A summary of the ESI-MS data is provided below. A list of compound hits and structures is provided in Table 30, Figure 53.

- Compound 673
 N^2 -[(2-methoxy-1-naphthyl)methyl]-3-phenyl- N^1 -(3-phenylpropyl)propane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 439.2
- Compound 674
 N^2 -[2-(benzyloxy)ethyl]- N^1 -(3,3-diphenylpropyl)-4-(methylthio)butane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 463.4.
- Compound 675
 N^1 -(3,3-diphenylpropyl)-4-(methylthio)- N^2 -(3-phenylpropyl)butane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 447.2
- Compound 676
 N^2 -(cyclohexylmethyl)- N^1 -(3,3-diphenylpropyl)-4-(methylthio)butane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 425.1
- Compound 677
 N^1 -(3,3-diphenylpropyl)- N^2 -(2-ethoxybenzyl)-4-(methylthio)butane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 463.1
- Compound 678
 N^2 -[2-(benzyloxy)ethyl]- N^1 -[(6,6-dimethylbicyclo[3.1.1]hept-2-yl)methyl]-4-(methylthio)butane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 405.3
- Compound 679
 N^1 -[(6,6-dimethylbicyclo[3.1.1]hept-2-yl)methyl]-4-(methylthio)- N^2 -(3-phenylpropyl)butane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 389.5
- Compound 680
 N^2 -(2-chloro-4-fluorobenzyl)-4-methyl- N^1 -(4-methylbenzyl)pentane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 363.3, 365.5; (MCH₃CN) 403.3, 405.3.
- Compound 681.
 N^2 -[2-(benzyloxy)ethyl]- N^1 -[2-(4-methoxyphenyl)ethyl]-4-methylpentane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 385.1.
- Compound 682.
 N^2 -[3-(4-chlorophenoxy)benzyl]- N^1 -[2-(4-methoxyphenyl)ethyl]-4-methylpentane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 467.1, 469.2.
- Compound 683.
 N^2 -(4-isopropylbenzyl)- N^1 -[2-(4-methoxyphenyl)ethyl]-4-methylpentane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 383.3

- Compound 684.
 N^1 -[2-(4-methoxyphenyl)ethyl]-4-methyl- N^2 -[(2*E*)-3-phenylprop-2-enyl]pentane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 367.3; [M-(CH₂CH=CHPh)2H]⁺ 251.
- 5 Compound 685
 N^2 -[2-(benzyloxy)ethyl]-4-methyl- N^1 -(3-phenylpropyl)pentane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 369.1.
- 10 Compound 686.
 N^2 -(2-chloro-4-fluorobenzyl)-4-methyl- N^1 -(3-phenylpropyl)pentane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 377.2, 378.9.
- 15 Compound 687.
 N^2 -[3-(4-chlorophenoxy)benzyl]-4-methyl- N^1 -(3-phenylpropyl)pentane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 451.1, 453.3.
- 20 Compound 688.
 N^2 -(4-isopropylbenzyl)-4-methyl- N^1 -(3-phenylpropyl)pentane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 367.3.
- 25 Compound 689
4-methyl- N^2 -[(2*E*)-3-phenylprop-2-enyl]- N^1 -(3-phenylpropyl)pentane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 351.2.
- 30 Compound 690
 N^2 -(2-ethoxybenzyl)-4-methyl- N^1 -(3-phenylpropyl)pentane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 369.1.
- 35 Compound 691.
 N^2 -decahydronaphthalen-2-yl- N^1 -[2-(4-fluorophenyl)ethyl]-3-thien-3-ylpropane-1,2-diamine. Mass spectrum (ESI) m/z (MH)⁺ 415.3.

References

1. Dye, C.; Scheele, S.; Dolin, P.; Pathania, V.; Raviglione, M.C., "Consensus statement. Global Burden of Tuberculosis: Estimate Incidence, Prevalence, and Mortality By Country. WHO Global Surveillance and Monitoring Project." *J. Am Med Association* 1999, 282, 677-686.
2. Bass, J.B.; Farer, L.S.; Hopewell, P.C.; O'Brein, R.; Jacobs, R.F.; Ruben, F.; Snider, D.E.Jr.; Thornton, G., "Treatment of Tuberculosis and Tuberculosis Infection in Adults and Children. The American Thoracic Society and the Centers for Disease Control and Prevention," *Am. J. Respir. Crit. Care Med.* 1994, 149, 1359-1374.
3. Farmer, P.; Bayona, J.; Becerra, M.; Furin, J.; Henry, C.; Hiatt, H.; Kim, J. Y.; Mitnick, C.; Nardell, E.; Shin, S., "The Dilemma of MDR-TB in the Global Era," *Int. J. Tuberc. Lung Dis.* 1998, 2, 869-876.
4. Pablos-Mendez, A.; Raviglione, M.C.; Laszlo, A.; Binkin, N.; Rieder, H.L.; Bustreo, F.; Cohn, D.L.; Lambregts van Weesenbeek, C.S.; Kim, S.J.; Chaulet, P.; Nunn, P., "Global Surveillance for Antituberculosis-drug Resistance," 1994-1997. *N. Engl. J. Med* 1998, 338, 1641-1649. [Erratum, *N. Engl. J. Med.* 1998, 339, 139].
5. Chan-Tack, K.M., "Antituberculosis-drug Resistance," *N. Engl. J. Med.* 1998, 339, 1079.
6. Cole, S. T.; Brosch, R.; Parkhill, J.; Garnier, T.; Churcher, C.; Harris, D.; Gordon, S. V.; Eiglmeier, K.; Gas, S.; Barry, C. E. 3RD; Tekaia, ; Badcock, K.; Basham, D.; Brown, D.; Chillingworth, T.; Connor, R.; Davies, R.; Devlin, K.; Feltwell, T.; Gentles, S.; Hamlin, N.; Holroyd, S.; Hornsby, T.; Jagels, K.; Barrell, B. G., "Deciphering the Biology of *Mycobacterium tuberculosis* From The Complete Genome Sequence," [published erratum appears in *Nature* 1998 Nov 12; 396, 190]; *Nature* 1998, 393, 537-544.
7. O'Brien, R.J., "Scientific Blueprint for Tuberculosis Drug Development," The Global Alliance for TB Drug Development, Inc. 2001.

8. Barry, C.E., III; Slayden, R.A.; Sampson, A.E.; Lee, R.E., "Use of Genomics and Combinatorial Chemistry in the Development of New Antimycobacterial Drugs," *Biochem. Pharmacol* 2000, *59*, 221.
9. Cynamon, M. H.; Klemens, S. P.; Sharpe, C. A.; Chase, S., "Activities of Several Novel Oxazolidinones Against Mycobacterium Tuberculosis In A Murine Model," *Antimicrob Agents Chemother.* 1999, *43*, 1189-91.
10. Shepard, R.G.; Baughn, C.; Cantrall, M.L.; Goodstein, B.; Thomas, J.P.; Wilkinson, R.G., "Structure-activity Studies Leading To Ethambutol, A New Type of Antituberculosis Compound," *Ann, N. Y. Acad. Sci* 1966, *135*, 686.
- 10 11. Deng, L.; Mikusova, K.; Robuck, K. G.; Scherman, M.; Brennan, P. J., McNeil, M. R., "Recognition of Multiple Effects of Ethambutol on Metabolism of Mycobacterial Cell Envelope," *Antimicrob. Agents Chemother.* 1995, *39*, 694-701.
12. Lee, R. E.; Mikusova, K.; Brennan, P.J.; and Besra, G.S.; "Synthesis of the Mycobacterial Arabinose Donor β -D-Arabinofuranosyl-1-monophosphoryl-decaprenol, Development of a Basic Arabinosyl-transferase Assay, and Identification of Ethambutol As An Arabinosyl Transferase Inhibitor," *J. Am. Chem. Soc.* 1995, *117*, 11829-11832.
13. Belanger, A.E.; Bestra, G.S.; Ford, M.E.; Mikusova, K.; Belisle, J.T.; Brennan, P.J.; Inamine, J.M., "The EmbAB Genes of Mycobacterium avium Encode An Arabinosyl Transferase Involved in Cell Wall Arabinan Biosynthesis That is The Target for The Antimycobacterial Drug Ethambutol," *Proc. Natl. Acad. Sci USA* 1996, *93*, 11919.
14. Telenti, A.; Phillip, W.J.; Sreevatsan, S.; Bernasconi, C.; Stockbauer, K.E.; Wieles, B.; Musser, J.M.; Jacobs, W.R. Jr., "The Emb Operon, A Gene Cluster of Mycobacterium Tuberculosis Involved in Resistance to Ethambutol," *Nat. Med.* 1997, *3*, 567.
15. Cuervo, J.H.; Weitzl, F.; Ostretch, J.M.; Hamashin, V.T; Hannah, A.L.; Houghten, R.A. in *Peptides 1994: Proceedings of the European Peptide Symposium*; Maia HSL Ed., Esom: Leiden, 1995, 465-466.
16. Silen, J.L; Lu, A.T.; Solas, D.W.; Gore, M.A.; Maclean, D.; Shah, N.H.; Coffin, J.M.; Bhinderwala, N.S.; Wang, Y.; Tsutsui, K.T.; Look, G.C.; Campbell, D.A.; Hale, R.L.; Navre, M.; Deluca-Flaherty, C.R., "Screening For Novel Antimicrobials


- from Encoded Combinatorial Libraries by Using a Two-Dimensional Agar Format," *Antimicrob. Agents Chemother.* 1998, 42, 1147.
17. Gustafson, G.R.; Baldino, C.M.; O'Donnel, M.-M.E.; Sheldon, A.; Tarsa, R.J.; verni, C.J.; Coffen, D.L., "Incorporation of Carbohydrates and Peptides Into Large
5 Triazine-based Screening Libraries Using Automated Parallel Synthesis," *Tetrahedron* 1998, 54, 4067.
18. H. Rink *Tetrahedron Lett.* 1987, 28, 3787.
19. Garigipati, R.V., "Reagents for Combinatorial Organic Synthesis: Preparation and Uses of Rink-chloride," *Tetrahedron Lett.* 1997, 38, 6807.
- 10
20. Brown, D.S.; Revill, J.M.; Shute, R.E. Merrifield, "Alpha-Methoxyphenyl (MAMP) Resin; A New Versatile Solid Support for The Synthesis of Secondary Amines, "*Tetrahedron Lett.* 1998, 39, 8533.
21. Zuckermann, R.N.; Kerr, S.B.H.; Moos, W.H., "Efficient Method for The
15 Preparation of Peptoids [oligo(N-substituted glycines)] by Submonomer Solid-phase Synthesis," *J. Am. Chem. Soc.* 1992, 114, 10646-10647.
22. Gordon, D.W.; Steele, J., "Reductive Alkylation on a Solid Phase: Synthesis of a Piperazidione Combinatorial Library, *Bioorg. Med. Chem. Lett.* 1995, 5, 47.
23. Liu, G.; Ellman J.A., "A General Solid-phase Synthesis Strategy for The
20 Preparation of 2-Pyrrolidinemethanol Ligands," *J. Org. Chem.* 1995, 60, 7712.
24. March, J., "Advanced Organic Chemistry," 3rd Ed., Wiley, New York, p. 916.
25. Luknitskii, Vovsi., *Russ. Chem. Rev.*, 1969, 38, 487-494.
26. Lee, M.H.; Pascopella, L.; Jacobs, W.R.; Hatfull, G.F., "Site Specific Integration of Mycobacteriophage L5: Integration-Proficient Vectors for *Mycobacterium smegmatis*, *Mycobacterium tuberculosis*, and *bacilli* Calmette-Guerin," *Proc. Natl Acad. Sci USA* 1991, 88, 3111.
- 25
27. Shawar, R.M.; Humble, D.J.; Van Dalfsen, J.M.; Stover, C.K.; Hickey, M.J.; Steele, S.; Mitscher, L.A.; Baker, W., "Rapid Screening of Natural Products for Antimycobacterial Activity By Using Luciferase-Expressing Strains of
30 *Mycobacterium bovis* BCG and *Mycobacterium intracellulare*," *Antimicrob. Agents Chemother.* 1997, 41, 570-574.

28. Arain, T.M.; Resconi, A.E.; Hickey, M.J.; Stover, C.K., "Bioluminescence Screening In Vitro (Bio-Siv) Assays for High-Volume Antimycobacterial Drug Discovery," *Antimicrob. Agents Chemother.* 1996, *40*, 1536-1541.

Claims

We claim:

1. A composition comprising a substituted ethylene diamine compound of the formula



5


wherein R_4 is selected from H, alkyl, aryl, heteroatom substituted alkyl and aryl, alkenyl, alkynyl, aralkyl, aralkynyl, cycloalkyl, cycloalkenyl;

- and wherein R_1 , R_2 and R_3 are independently selected from H, alkyl, aryl, alkenyl, alkynyl, aralkyl, aralkenyl, aralkynyl, cycloalkyl, cycloalkenyl, heteroalkyl, heteroaryl, halide, alkoxy, aryloxy, alkylthio, arylthio, silyl, siloxy, amino; or





- wherein R_1 is selected from H, alkyl, aryl, alkenyl, alkynyl, aralkyl, aralkenyl, aralkynyl, cycloalkyl, cycloalkenyl, heteroalkyl, heteroaryl, halide, alkoxy, aryloxy, alkylthio, arylthio, silyl, siloxy, amino, and NR_2R_3 is derived from a cyclic secondary amine.




2. The composition of Claim 1, wherein NHR_1 or NR_2R_3 of the substituted ethylene diamine has the chemical structure

20







3. The composition of Claim 2, wherein the substituted ethylene diamine compound is selected from






5







5



; or






- 10 4. The composition of Claim 1, wherein NHR_1 or NR_2R_3 of the substituted ethylene diamine has the chemical structure







5. The composition of Claim 4, wherein the substituted ethylene diamine compound is selected from

5





10







5




10



;




;




;

5




; or






10

6. The composition of Claim 1, wherein NHR_1 or NR_2R_3 of the substituted ethylene diamine has the chemical structure







7. The composition of Claim 6, wherein the substituted ethylene diamine compound is selected from

5





10








5





10





- 5 8. The composition of Claim 1, wherein NHR_1 or NR_2R_3 of the substituted ethylene diamine has the chemical structure






- 10 9. The composition of Claim 8, wherein the substituted ethylene diamine compound is selected from



15




5





10



10. The composition of Claim 9, wherein the substituted ethylene diamine compound is





11. The composition of Claim 1, wherein the substituted ethylene diamine compound is
5 selected from





10









5






10










5







10




5



10



- 10 12. A method of preparing a substituted ethylene diamine compound of the formula



wherein R_4 is selected from H, alkyl, aryl, alkenyl, alkynyl, aralkyl, aralkynyl, cycloalkyl, cycloalkenyl;

and wherein R_1 , R_2 and R_3 are independently selected from H, alkyl, aryl, alkenyl, alkynyl, aralkyl, aralkenyl, aralkynyl, cycloalkyl, cycloalkenyl, heteroalkyl, heteroaryl, halide, alkoxy, aryloxy, alkylthio, arylthio, silyl, siloxy, amino; or

wherein R_1 is selected from H, alkyl, aryl, alkenyl, alkynyl, aralkyl, aralkenyl, aralkynyl, cycloalkyl, cycloalkenyl, heteroalkyl, heteroaryl, halide, alkoxy, aryloxy, alkylthio, arylthio, silyl, siloxy, amino, and NR_2R_3 is derived from a cyclic secondary amine;

comprising activating a solid-support resin containing hydroxyl groups with a halo-donating reagent in the presence of base to produce a solid-support resin containing halo groups;

displacing the halo groups with an initial primary amine to produce a solid-support resin containing amine groups;

acylating the amine groups with a halo-acylhalide in the presence of a basic compound, or with a halo-acylacid in the presence of base, to produce a solid-support resin containing α -haloacetyl amide groups;

displacing α -halo groups of the α -haloacetyl amides with a secondary or subsequent primary amine to produce a solid-support resin containing α -amine imide groups;


reducing the carbonyl moiety on the α -amine imide groups with a reducing agent to produce a solid-support resin containing two amine groups separated by two carbon atoms;

cleaving the amine groups separated by two carbon atoms from the solid support resin in the presence of acid to produce the substituted ethylene diamine compound.

13. The method of Claim 12, wherein the initial primary amine is R_1NH_2 .

14. The method of Claim 12, wherein the secondary or subsequent primary amine is R_2R_3HN .

15. A method of treating disease caused by an infectious agent comprising administering an effective amount of a substituted ethylene diamine compound of the formula:



wherein R_4 is selected from H, alkyl, aryl, alkenyl, alkynyl, aralkyl, aralkynyl, cycloalkyl, cycloalkenyl;

and wherein R_1 , R_2 and R_3 are independently selected from H, alkyl, aryl, alkenyl, alkynyl, aralkyl, aralkenyl, aralkynyl, cycloalkyl, cycloalkenyl, heteroalkyl, heteroaryl, halide, alkoxy, aryloxy, alkylthio, arylthio, silyl, siloxy, amino; or

wherein R_1 is selected from H, alkyl, aryl, alkenyl, alkynyl, aralkyl, aralkenyl, aralkynyl, cycloalkyl, cycloalkenyl, heteroalkyl, heteroaryl, halide, alkoxy, aryloxy, alkylthio, arylthio, silyl, siloxy, amino, and NR_2R_2 is derived from a cyclic secondary amine.

16. The method of Claim 15, wherein the infectious agent comprises a bacterial, mycological, parasitic, or viral agent.

15


17. The method of Claim 16, wherein the bacterial agent comprises *M. tuberculosis*, *M. avium-intracellulare*, *M. kansarii*, *M. fortuitum*, *M. chelonae*, *M. leprae*, *M. africanum*, *M. microti*, *M. avium paratuberculosis*, *M. intracellulare*, *M. scrofulaceum*, *M. xenopi*, *M. marinum*, or *M. ulcerans*.

20

18. The method Claim 15, wherein the infectious disease comprises tuberculosis or Crohn's disease.

19. The method Claim 15, wherein the substituted ethylene diamine compound is


25




20. The method of Claim 19, further comprising a pharmaceutical carrier.

30

21. A composition comprising, a substituted ethylene diamine compound comprising,




;




5


;



;






;





;



10




5




10






;




;




5

;




;




;


10



; or



22. A method of preparing a substituted ethylene diamine compound of the formula



5 wherein R₄ is selected from H, alkyl, aryl, heteroatom substituted alkyl and aryl, alkenyl, alkynyl, aralkyl, aralkynyl, cycloalkyl, cycloalkenyl;

 and wherein R₁, R₂ and R₃ are independently selected from H, alkyl, aryl, alkenyl, alkynyl, aralkyl, aralkenyl, aralkynyl, cycloalkyl, cycloalkenyl, heteroalkyl, heteroaryl, halide, alkoxy, aryloxy, alkylthio, arylthio, silyl, siloxy, amino;

10 comprising activating a solid-support resin containing hydroxyl groups with a halo-donating reagent in the presence of base to produce a solid-support resin containing halo groups;

 displacing the halo groups with an initial primary amine to produce a solid-support resin containing amine groups;

15 acylating the amine groups with a Fmoc protected amino acid in the presence of a coupling reagent and a base, followed by removal of Fmoc protecting group to produce a solid-support resin containing α-amino acetamide groups;

 modification of α-amino groups of the α-amino acetamide groups with a carbonyl compound to produce a solid-support resin containing corresponding derivative of α-amino

20 acetamide groups;


 reducing the carbonyl moiety on the amide groups with a reducing agent to produce a solid-support resin containing two amine groups separated by two carbon atoms; and

 cleaving the amine groups separated by two carbon atoms from the solid support resin in the presence of acid to produce the substituted ethylene diamine compound.

25

23. A method for treating an infectious disease comprising administering a pharmaceutically effective amount of the composition in Claim 21.

24. A composition comprising a symmetrical substituted ethylene diamine compound of the formula



wherein R₄ is selected from H, alkyl, aryl, heteroatom substituted alkyl and aryl ,
5 alkenyl, alkynyl, aralkyl, aralkynyl, cycloalkyl, cycloalkenyl;

and wherein R₁, R₂ and R₃ are independently selected from H, alkyl, aryl, alkenyl,
alkynyl, aralkyl, aralkenyl, aralkynyl, cycloalkyl, cycloalkenyl, heteroalkyl, heteroaryl,
halide, alkoxy, aryloxy, alkylthio, arylthio, silyl, siloxy, amino; or

10

wherein R₁ is selected from H, alkyl, aryl, alkenyl, alkynyl, aralkyl, aralkenyl,
aralkynyl, cycloalkyl, cycloalkenyl, heteroalkyl, heteroaryl, halide, alkoxy, aryloxy,
alkylthio, arylthio, silyl, siloxy, amino, and NR₂R₂ is derived from a cyclic secondary
amine.

15