


Abstract

A superior LXR modulator is provided. A compound represented by the general formula (I):



[wherein R^1 : $-\text{COR}^9$ (wherein R^9 : alkyl, optionally substituted alkoxy or optionally substituted amino); R^2 : H, OH, alkoxy, optionally substituted amino, etc.; R^3 : H, optionally substituted alkyl, cycloalkyl, optionally substituted alkoxy, optionally substituted amino, halogeno, etc.; R^4 and R^5 : H, optionally substituted alkyl, halogeno, etc.; R^6 and R^7 : H, alkyl; R^8 : $-\text{X}^2\text{R}^{10}$ [wherein R^{10} : $-\text{COR}^{11}$ (wherein R^{11} : OH, optionally substituted alkoxy, optionally substituted amino, etc.), $-\text{SO}_2\text{R}^{12}$ (wherein R^{12} : optionally substituted alkyl, optionally substituted amino, etc.), tetrazol-5-yl, etc.; X^2 : single bond, optionally substituted alkylene, etc.]; X^1 : $-\text{NH}-$, $-\text{O}-$, $-\text{S}-$, etc.; Y^1 : optionally substituted phenyl, optionally substituted 5- to 6-membered aromatic heterocyclyl; Y^2 : optionally substituted aryl, optionally substituted heterocyclyl, etc.] and the like is provided.