

LIPID-ENCAPSULATED DUAL-CLEAVING ENDONUCLEASE FOR DNA AND GENE EDITING

SEQUENCE LISTING

[0001] The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on September 17, 2018, is Sequence_Listing_ST25.txt and is 77KB in size.

BACKGROUND OF THE INVENTION

[0002] There are an estimated 5,000 - 10,000 monogenic diseases, defined as inherited conditions arising from mutations on a single gene. These diseases often manifest during childhood and lead to a variety of conditions and sometimes premature death. It has been estimated that together they will affect about 6% of people at some point in their lives. Diagnosis and treatment for these diseases remain largely insufficient, and the care is primarily palliative, focusing on disease management without addressing the underlying genetic defects. There are also many more diseases in which a mutation to a gene contributes to the pathogenesis of the disease.

[0003] Gene editing is a gene therapy approach that relies on designer nucleases to recognize and cut specific DNA sequences, and subsequently exploits innate cellular DNA repair pathways, namely nonhomologous end joining (NHEJ) and homology directed repair (HDR), to introduce targeted modifications in the genome. Four nuclease families have been used in this context: meganucleases, zinc-finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs), and clustered regulatory interspaced short palindromic repeats associated RNA-guided Cas9 (CRISPR-Cas9) nucleases. These can be designed to precisely introduce a double stranded break at the target locus of interest. Gene editing opens up the possibility of permanently modifying a genomic sequence of interest by enabling targeted disruption, insertion, excision, and correction in both *ex vivo* and *in vivo* settings. While these advances are expected to revolutionize the field at large, current gene-editing approaches are limited by efficacy of modification, safety concerns related to the specificity of nucleases, and delivery of gene-editing tools to target cell types.

[0004] A component of the type II CRISPR system that constitutes the innate immune system of bacteria, the Cas9 (CRISPR-associated) protein has caused a paradigm shift in the field of genome editing due to its ease-of-use. Programming Cas9 to cleave a desired sequence is a simple matter of changing the sequence of the Cas9-associated guide RNA to be complementary to the target site. The ease of programming Cas9 targeting contrasts with the more intensive protein engineering that is required for other reagents (zinc finger nucleases (ZFNs), meganucleases, transcription activator-like effector nucleases (TALENs)). Cas9, along with proteins from type III CRISPR systems have been used for a myriad of genome-editing applications in a diverse range of organisms and are now entering the realm of therapeutic applications in humans.

[0005] Cystic fibrosis (CF) is an autosomal-recessive disease resulting from mutations in the *CFTR* gene, which encodes an epithelial anion channel. The CFTR protein, cystic fibrosis transmembrane conductance regulator, is found across a wide range of organs including pancreas, kidney, liver, lungs, gastrointestinal tracts, and reproductive tracts, making CF a multiorgan disease. Mutations in *CFTR* lead to suboptimal ion transport and fluid retention, causing the prominent clinical manifestations of abnormal thickening of the mucus in lungs and pancreatic insufficiency. In the lung, dysfunctional CFTR hinders mucociliary clearance, rendering the organ susceptible to bacterial infections and inflammation, ultimately leading to airway occlusion, respiratory failure, and premature death. CF remains the most common and lethal genetic disease among the Caucasian population with 70,000–100,000 sufferers estimated worldwide, highlighting a real need for the development of better treatments.

[0006] One major challenge to the development of a therapeutic strategy for CF is the wide diversity of mutation types. Delta F508 (deletion of phenylalanine at codon 508) mutation, with a prevalence of >80% in CF patients, is by far the most common, but more than 1,990 deleterious *CFTR*-mutations have been described. These mutations cause premature stop codons, aberrant splicing, incorrect protein folding or trafficking to the cell surface, and dysfunctional CFTRs with limited channel-opening capacity. Pharmacological interventions have been targeted to several of these processes and while drug administration is therapeutic in some gating mutation types, the commonly occurring delta F508 still requires a more effective treatment. Pharmaceutical advancement in the care of CF, however, does not address mutations resulting

from aberrant splicing or premature stop codons; it is in these instances gene editing could prove most beneficial.

[0007] Similarly, in the Western population, approximately 15% of patients with non-small cell lung cancer (NSCLC) harbor an activating mutation in their tumor in the EGF receptor (EGFR) gene.

[0008] Existing gene editing technologies, such as CRISPR-Cas9 (and Cas9 fusions), meganucleases, zinc finger proteins, type IIS restriction endonucleases (FokI and FokI fusions) and TALENS are limited in the ability to introduce gene deletions of a specific length or to accurately repair a target gene in a sufficient number of cells to be meaningful as a therapeutic agent for many genetic diseases. Moreover, for highly programmable RNA-guided nucleases, such as the monomeric Cas9, studies suggest that the specificity for predictably binding, cleaving and repairing only their target sites is limited, raising concerns over potential deleterious changes to a cell's genomic DNA that may inadvertently cause a secondary disease in a patient. Last, most nucleases are delivered in viral vectors. Viral vectors have the potential for: existing immunity in many populations; immunogenicity after treatment; and genotoxicity. No non-viral delivery method exists today to safely deliver the nuclease to target cells and allow for controlled dosing of the nuclease *in vivo*.

[0009] There is an unmet need for improvements to said existing gene editing technologies to address the above concerns to make gene editing technologies more efficient and effective.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010] **FIG. 1** is a schematic representation of the lipid-encapsulated dual-cleaving nuclease (TevCas9) after it has been prepared [*Components are not to scale*]. The I-TevI Domain **10** is joined to the RNA-guided Nuclease (Cas9) Domain **12** via a Linker Domain **11**. In the preferred embodiment, the formed particles also contain Guide RNA **13** and Donor DNA **14**. The aforementioned nuclease is contained in a Lipid Particle **15** which has been shaped into a sphere using an extrusion process.

[0011] **FIG. 2** is a diagram of the mechanism by which lipid-encapsulated TevCas9 is internalized into a cell and nucleus to reach its target DNA. As illustrated in **FIG. 2A**, a cell **20** or cells **20** are exposed to the novel lipid-encapsulated nuclease particles **21** containing the TevCas9 **25** either by *in vivo* or *ex vivo* administration. As shown in **FIG. 2B**, the lipid-

encapsulated nuclease particle **21** is endocytosed into the cell **20**. The endosome **22** goes through a maturation process in the cytosol and is targeted for degradation (**FIG. 2C**). On certain occasions, the TevCas9 **25** can escape the endosome **22** and enter the cytosol (**FIG. 2D**). In eukaryotic organisms, the nuclease (TevCas9) **25** is targeted to the nucleus **23** of the cell **20** through one or more nuclear-localization sequences (“NLS”). As depicted in **FIG. 2E**, through its nuclear localization sequence, TevCas9 **25** can enter the nucleus **23** and when in the nucleus **23**, the TevCas9 nuclease **25** binds to and cleaves **26** the target genomic DNA **24** sequence.

[0012] **FIG. 3** is a diagram of the mechanisms by which lipid-encapsulated TevCas9 modifies target DNA. The I-TevI domain **27** targets the I-TevI Target Sequence **29**. The linker domain **30** joins the I-TevI domain **27** with the Cas9 domain **28** which targets the Cas9 Target Sequence **31**. The gene mutation **32** is surrounded by or in close proximity to the I-TevI Target Sequence **29** and the Cas9 Target Sequence **31**. As shown in **FIG. 3B**, the TevCas9 **25** cleaves the target sequence leaving a deletion product **34** of a predictable size with non-complementary DNA ends **35, 36**. **FIG. 3C** illustrates that in the presence of single-stranded donor DNA with homology arms **37**, the cell **20** can insert the donor DNA **37** sequence near the cut sites through the homology-directed repair (HDR) pathway **38**. **FIG. 3D** illustrates that in the presence of donor DNA **39** with compatible DNA ends to those cleaved by TevCas9 **25**, the cell **20** can insert the donor DNA sequence **39** between the cut sites through directed-ligation using the non-homologous end joining (NHEJ) pathway **40**. In the absence of donor DNA, the cell **20** can join the DNA ends through the NHEJ pathway **40** (**FIG. 3E**).

[0013] **FIG 4A** evidences that TevCas9, targeted to the *CFTR* gene using guide in SEQ ID 15, cleaves *CFTR* DNA substrate *in vitro*. **FIG. 4B** are cells transfected with a plasmid DNA version of TevCas9 fused to a cleavable GFP tag imaged using phase contrast and GFP imaging on a Cytation5 (Biotek Instruments Inc, VT, USA) after 48 hours treatment. Genomic DNA is extracted from harvested cells and editing at the *CFTR* gene is detected by PCR amplification and a T7 Endonuclease I cleavage assay.

[0014] **FIG 5** evidences that TevCas9, targeted to the *CFTR Delta F508* mutation using guide in SEQ ID 21, cleaves a DNA substrate containing the *CFTR Delta F508* mutation, but not substrate containing the wild-type *CFTR* sequence *in vitro*.

[0015] **FIG 6A** illustrates that the saCas9 D10E mutation slows the conversion of nicked supercoiled DNA to linear DNA. **FIG 6B** evidences that on linear *EMXI* DNA substrate,

saCas9D10E (D10E) ribonucleoprotein complex (RNP) cleaves the target substrate to a similar level as saCas9 wild-type (WT). Levels of editing by SaCas9D10E at computationally predicted off-targets is lower than levels of editing by wild-type saCas9 at the same off-targets.

[0016] **FIG 7A** is a schematic of spacing of the I-TevI sites in the *EGFR Exon 19 deletion* and wild-type (WT) *EGFR*. **FIG 7B** and **7C** evidences that TevCas9 containing the nicking mutation in Cas9 (H557A) targeted to *EGFR* using the guide RNA in SEQ ID 16 cleaves *EGFR Exon 19 deletion* DNA substrate at a 4-fold faster rate than wild type *EGFR*. **FIG 7D** are images of HCC827 cells harbouring the *EGFR Exon 19 deletion* mutation treated with TevCas9 targeted to *EGFR* are selectively killed compared to NuLi-1 cells harbouring wild-type EGFR (WT).

BRIEF SUMMARY OF THE INVENTION

[0017] The instant invention is directed to a chimeric nuclease comprising a modified I-TevI nuclease domain, preferably deleting Met¹ and having Lys²⁶ (which is Lys²⁷ in the untruncated version of I-TevI) and/or Cys³⁹ (which is Cys⁴⁰ in the untruncated version of I-TevI) modification, a linker, in particular SEQ ID NOS: 7 – 12 or fragments thereof and/or containing one or more of the following mutations Thr⁹⁵ (as referenced to the full-length I-TevI), Val¹¹⁷, Lys¹³⁵, Gln¹⁵⁸ or Asn¹⁴⁰, and a modified RNA-guided nuclease *Staphylococcus aureus* Cas9 that may be the wild-type or a modified version, preferably containing a Glu¹⁰ or an Ala⁵⁵⁷ mutation thereof wherein said I-TevI polypeptide comprises the entire amino acid sequence of SEQ ID NO: 6 or a fragment thereof, and guide RNA, in particular, SEQ ID NOS: 15, 16 or 21 or fragments thereof, that targets the Cas9 domain and a pharmaceutically-acceptable formulation comprising the chimeric nuclease, cationic and/or neutral lipid nanoparticles, optionally DNA-binding compounds, in particular GL67 (N⁴-cholesteryl-spermine) and a pharmaceutically acceptable carrier thereof.

[0018] In a further embodiment of the instant invention, in the formulation the lipid nanoparticle may contain exogenous donor DNA.

[0019] Another embodiment of the invention is directed to methods to edit genes by administering a chimeric nuclease to a cell or organism without the use of a viral vector by using a controlled dose *in vivo*.

[0020] Another embodiment of the invention is directed to methods to delete defined lengths of a DNA molecule or to replace select sequences from a DNA molecule by delivering a chimeric

nuclease *in vivo* to a whole organism or to isolated cells in culture *ex vivo* wherein said cells are mammalian cells, bacteria, insect cells or plant cells.

[0021] In yet another embodiment, the novel chimeric nuclease targets two independent target sites on a select DNA molecule either cleaving at one target site or at both target sites and creating fragments that are 30 to 36 nucleotides in length.

[0022] In a further example, the novel, purified chimeric nuclease further comprises a guide RNA.

[0023] Another aspect of the instant inventions is the use of an extrusion process creating particles of approximately 100 nM in diameter comprising an excipient wherein the excipient is selected from the group consisting of polysorbates, polyphosphates, calcium chlorides, sodium chloride, sodium citrates, sodium hydroxide, sodium phosphates, sodium ethylenediaminetetraacetic acid, potassium chloride, potassium phosphate and starches, or mixtures of these substances so that the novel chimeric nuclease can be administered to a patient using a nebulizer containing said formulation.

[0024] In a preferred embodiment, the instant invention is directed to a method of treating a lung-related disease in a patient in need thereof by administering a novel chimeric nuclease that modifies the DNA of lung epithelial cells wherein the chimeric nuclease replaces the *CFTR delta F508* mutation from the *CFTR* gene in an effort to treat cystic fibrosis or cleaves an *EGFR exon 19 deletion* in an effort to treat non-small cell lung cancer.

[0025] In yet another embodiment, the invention is directed to a chimeric nuclease comprising a modified I-TevI nuclease domain, a linker and a modified RNA-guided nuclease *Staphylococcus aureus* Cas9 wherein said RNA-guided nuclease *Staphylococcus aureus* Cas9 contains Ala¹⁰, Ala⁵⁵⁷ or Ala⁵⁸⁰ mutations and targets the *EGFR* exon 19 deletions of the *EGFR* gene.

[0026] In a further embodiment, wherein said guide RNA targets a specific *CTFR* gene sequence to cleave out the *CFTR delta F508* mutation or a specific *EGFR* gene sequence that contains an *EGFR* exon 19 deletion mutation.

[0027] The instant invention also covers linkers comprising SEQ ID NOS: 7 – 12 or fragments thereof and modified donor DNA molecules selected from the group consisting of a linear single-strand of DNA comprising homologous regions flanking the sites targeted and/or cleaved by a chimeric nuclease, a linear double-strand DNA comprising homologous regions flanking the sites targeted and/or cleaved by a chimeric nuclease, a double strand DNA of the same length

comprising complimentary DNA ends to those cleaved by a chimeric nuclease, a circular double-strand DNA comprising homologous regions flanking the sites targeted and/or cleaved by a chimeric nuclease, and a circular double-strand DNA comprising an I-TevI target site and a Cas9 target site wherein the product cleaved from the double-strand DNA contains complimentary ends to the ends cleaved by a chimeric nuclease.

[0028] In a further example consists of a chimeric nuclease comprising a modified GIY-YIG nuclease domain, a linker and a modified RNA-guided nuclease *Staphylococcus aureus* Cas9 or *Streptococcus pyogenes* Cas9 or *EQR Streptococcus pyogenes* Cas9 variant containing a Glu¹⁰ mutation (SEQ ID NO:19) and/or a Ala⁸⁴⁰ mutation and/or a mutation that cleaves the sugar phosphate backbone of a target DNA on one strand of a target DNA wherein said GIY-YIG nuclease domain is selected from the gene family consisting of I-Bmol and Eco29kI.

[0029] In yet a further embodiment, the instant invention includes a chimeric nuclease comprising a modified I-TevI nuclease domain, a linker and a modified nuclease or DNA targeting domain wherein said modified nuclease or DNA targeting domain is selected from the group consisting of LAGLIDADG, His-Cys Box, H-N-H, PD-(D/E)xK and Vsr-like meganucleases, zinc-finger nuclease, CRISPR protein selected from the group consisting of scCas9 (*Streptococcus canis*), fnCas9 (*Francisella novicida*), cjCas9 (*Campylobacter jejuni*), Cpf1 (*Lachnospiraceae bacterium*), Cas12a (*Acidaminococcus* Sp), Cas13a (*Leptorichia shahii*) and Cas3 (*Streptococcus thermophilus*) and DNA binding domain selected from the group consisting of zinc-finger motifs and TALE activator domains.

[0030] In an even further example, the instant invention covers a modified RNA-guided nuclease *Staphylococcus aureus* Cas9 and a guide RNA wherein said guide RNA contains sequences that target genetic polymorphisms, different sequences in the *CFTR* or *EGFR* genes, sequences that retarget a nuclease, bridged nucleic acids and/or a mixture of guide RNAs.

DETAILED DESCRIPTION OF THE INVENTION

DEFINITIONS AND ACRONYMS

[0031] For convenience, certain terms employed in the specification, examples and appended claims are collected here. These definitions should be read in light of the disclosure and understood as by a person of skill in the art. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by a person of

ordinary skill in the art.

[0032] The articles "a" and "an" are used herein to refer to one or to more than one (*i.e.*, to at least one) of the grammatical object of the article. The term "and/or" as used herein is defined as the possibility of having one or the other or both. For example, "A and/or B" provides for the scenarios of having just A or just B or a combination of A and B. If the claim reads A and/or B and/or C, the composition may include A alone, B alone, C alone, A and B but not C, B and C but not A, A and C but not B or all three A, B and C as components.

[0033] The term "bioavailable" is art-recognized and refers to a form of the subject disclosure that allows for it, or a portion of the amount administered, to be absorbed by, incorporated to, or otherwise physiologically available to a subject or patient to whom it is administered.

[0034] The term "exogenous donor DNA", as used herein, refers to any sequence of DNA that, in whole or in part, is not the same as the original target DNA sequence.

[0035] The term "flexible linker", as used herein, refers to a situation when the RNA-guided nuclease domain (Cas9) binds to the target DNA sequence, the amino acid linker domain ensures mobility of the I-TevI domain to allow for recognition, binding and cleaving of its target sequence under cell physiological conditions (typically: pH ~7.2, temperature ~37°C, [K+] ~140 mM, [Na+] ~5 – 15 mM, [Cl-] ~4 mM, [Ca++] ~0.0001 mM). The length of the amino acid linker can influence how many nucleotides are preferred between the Cas9 target site and the I-TevI target site. Certain amino acids in the linker may also make specific contacts with the DNA sequence targeted by TevCas9. These linker-DNA contacts can affect the flexibility of the I-TevI domain. Substituting amino acids in the linker domain may affect the ability of the linker domain to make contact with DNA.

[0036] The term "including", as used herein, is used to mean "including but not limited to". "Including" and "including but not limited to" are used interchangeably.

[0037] The terms "inhaled administration", "inhale", "inhaled", "inhalation" or "inhalation therapy", which may be used interchangeably and as used herein, include administration of a substantially uniform distribution of appropriately sized particles to the respiratory epithelium of the nose, central airways, the peripheral aspect of the lung and/or the alveolar region of the lung or by intratracheal instillation. Such particles may be introduced to the patient and/or produced using an appropriate device, preferably a nebulizer.

[0038] The term "patient," "subject" or "host" to be treated by the subject method may mean

either a human or non-human animal. Non-human animals include companion animals (e.g. cats, dogs) and animals raised for consumption (i.e. food animals), such as cows, pigs, and chickens.

[0039] The term "pharmaceutically acceptable carrier" is art-recognized and refers to a pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting any subject composition or component thereof from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be "acceptable" in the sense of being compatible with the subject composition and its components and not injurious to the patient. Some examples of materials which may serve as pharmaceutically acceptable carriers include: (1) sugars, such as dextrose, lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as microcrystalline cellulose, sodium carboxymethyl cellulose, methyl cellulose, ethyl cellulose, hydroxypropylmethyl cellulose (HPMC), and cellulose acetate; (4) glycols, such as propylene glycol; (5) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (6) esters, such as ethyl oleate, glyceryl behenate and ethyl laurate; (7) buffering agents, such as monobasic and dibasic phosphates, Tris/Borate/EDTA and Tris/Acetate/EDTA (8) pyrogen-free water; (9) isotonic saline; (10) Ringer's solution; (11) ethyl alcohol; (12) phosphate buffer solutions; (13) polysorbates; (14) polyphosphates; and (15) other non-toxic compatible substances employed in pharmaceutical formulations. The disclosed excipients may serve more than one function. For example, a solubilizing agent may also be a suspension aid, an emulsifier, a preservative, and the like.

[0040] In certain preferred embodiments, the pharmaceutically acceptable excipient is a crystalline bulking excipient. The terms "crystalline bulking excipient" or "crystalline bulking agent" as used herein means an excipient which provides bulk and structure to the lyophilization cake. These crystalline bulking agents are inert and do not react with the protein or nucleic acid. In addition, the crystalline bulking agents are capable of crystallizing under lyophilization conditions. Examples of suitable crystalline bulking agents include hydrophilic excipients, such as, water soluble polymers; sugars, such as mannitol, sorbitol, xylitol, glucitol, ducitol, inositol, arabinitol, arabitol, galactitol, iditol, allitol, maltitol, fructose, sorbose, glucose, xylose, trehalose, allose, dextrose, altrose, lactose, glucose, fructose, gulose, idose, galactose, talose, ribose, arabinose, xylose, lyxose, sucrose, maltose, lactose, lactulose, fucose, rhamnose, melezitose,

maltotriose, raffinose, altritol, their optically active forms (D- or L-forms) as well as the corresponding racemates; inorganic salts, both mineral and mineral organic, such as, calcium salts, such as the lactate, gluconate, glycerylphosphate, citrate, phosphate monobasic and dibasic, succinate, sulfate and tartrate, as well as the same salts of aluminum and magnesium; carbohydrates, such as, the conventional mono- and di-saccharides as well as the corresponding polyhydric alcohols; proteins, such as, albumin; amino acids, such as glycine; emulsifiable fats and polyvinylpyrrolidone. Preferred crystalline bulking agents are selected from the group consisting of glycine, mannitol, dextran, dextrose, lactose, sucrose, polyvinylpyrrolidone, trehalose, glucose and combinations thereof. Particularly useful bulking agents include dextran.

[0041] The term "pharmaceutically-acceptable salts", as used herein, is art-recognized and refers to the relatively non-toxic, inorganic and organic acid addition salts, or inorganic or organic base addition salts of compounds, including, for example, those contained in compositions of the present invention. Some examples of pharmaceutically-acceptable salts include: (1) calcium chlorides; (2) sodium chlorides; (3) sodium citrates; (4) sodium hydroxide; (5) sodium phosphates; (6) sodium ethylenediaminetetraacetic acid; (7) potassium chloride; (8) potassium phosphate; and (9) other non-toxic compatible substances employed in pharmaceutical formulations.

[0042] The term "substitution", as used herein, refers to the replacement of an amino acid in a sequence with a different amino acid. As used herein, the shorthand X¹⁰Y indicates that amino acid Y has been "substituted" for amino acid X found in the 10th position of the sequence. As an example, W²⁶C denotes that amino acid Tryptophan-26 (Trp, W) is changed to a Cysteine (Cys). Similarly, the notation AA^X indicates that AA is an amino acid that replaced the amino acid found in the X position. As an example, Lys²⁶ denotes the replacement of the amino acid in the 26th position in a sequence with Lysine. Use of either shorthand is interchangeable. In addition, use of the one- or three- letter abbreviations for an amino acid is also interchangeable.

[0043] The term "therapeutic agent", as used herein, is art-recognized and refers to any chemical or biochemical moiety that is a biologically, physiologically, or pharmacologically active substance that acts locally or systemically in a subject. Examples of therapeutic agents, also referred to as "drugs", are described in well-known literature references such as the Merck Index, the Physician's Desk Reference, and The Pharmacological Basis of Therapeutics, and they include, without limitation, medicaments; vitamins; mineral supplements; substances used

for the treatment, prevention, diagnosis, cure or mitigation of a disease or illness; substances which affect the structure or function of the body; or pro-drugs, which become biologically active or more active after they have been placed in a physiological environment.

[0044] The term "therapeutic effect", as used herein, is art-recognized and refers to a local or systemic effect in animals, particularly mammals, and more particularly humans caused by a pharmacologically active substance. The term thus means any substance intended for use in the diagnosis, cure, mitigation, treatment, or prevention of disease or in the enhancement of desirable physical or mental development and/or conditions in an animal or human. The phrase "therapeutically-effective amount" means that amount of such a substance that produces some desired local or systemic effect at a reasonable benefit/risk ratio applicable to any treatment. The therapeutically effective amount of such substance will vary depending upon the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of ordinary skill in the art. For example, certain compositions of the present invention may be administered in a sufficient amount to produce a at a reasonable benefit/risk ratio applicable to such treatment.

[0045] The term "treating", as used herein, includes any effect, *e.g.*, lessening, reducing, modulating, or eliminating, that results in the improvement of the condition, disease, disorder, and the like. As used herein, "treating" can include both prophylactic, and therapeutic treatment. For example, therapeutic treatment can include delaying inhibiting or preventing the progression of cystic fibrosis or non-small cell lung cancer, the reduction or elimination of symptoms associated with cystic fibrosis or non-small cell lung cancer. Prophylactic treatment can include preventing, inhibiting or delaying the onset of cystic fibrosis or non-small cell lung cancer.

[0046] As used herein, an "effective amount" refers to an amount sufficient to elicit the desired response. In the present invention, the desired biological response is the treatment of cystic fibrosis and/or non-small cell lung cancer (NSCLC).

[0047] A "buffer" as used herein is any acid or salt combination which is pharmaceutically acceptable and capable of maintaining the composition of the present invention within a desired pH range. Buffers in the disclosed compositions maintain the pH in a range of about 2 to about 8.5, about 5.0 to about 8.0, about 6.0 to about 7.5, about 6.5 to about 7.5, or about 6.5. Suitable buffers include, any pharmaceutical acceptable buffer capable of maintaining the above pH

ranges, such as, for example, acetate, tartrate phosphate or citrate buffers. In one embodiment, the buffer is a phosphate buffer. In another embodiment the buffer is an acetate buffer. In one embodiment the buffer is disodium hydrogen phosphate, sodium chloride, potassium chloride and potassium phosphate monobasic.

[0048] In the disclosed compositions the concentration of buffer is typically in the range of about 0.1 mM to about 1000 mM, about 0.2 mM to about 200 mM, about 0.5 mM to about 50 mM, about 1 mM to about 10 mM or about 6.0 mM.

[0049] As used herein, an "anti-microbial agent" is a pharmaceutically acceptable preservative, suitable for administration to a subject, which inhibits, prevents, or delays the growth or microorganisms including, for example bacteria, viruses and fungi in the compositions of the present invention. Suitable anti-microbial agents for use in the compositions and methods of the present invention include, but are not limited to, cresols, benzyl alcohol, phenol, benzalkonium chloride, benzethonium chloride, chlorobutanol, phenylethyl alcohol, methyl paraben, propyl paraben, thiomersal and phenylmercuric nitrate and acetate. In one embodiment the anti-microbial agents is m-cresol, chlorocresol or phenol. In another embodiment the anti-microbial agents is chlorocresol or phenol. In another embodiment the anti-microbial agents is phenol.

[0050] As used herein an effective amount of an anti-microbial agent is an amount effective to inhibit, prevent or delay the growth or microorganisms including, for example bacteria, viruses, and fungi in the compositions of the present invention. In the compositions of the present invention, the amount of anti-microbial agent is typically in the range from about 0.1 to about 20 mg/ml, about 0.2 to about 30 mg/ml, about 0.2 to about 10 mg/ml, about 0.25 to about 5 mg/ml, about 0.5 to about 50 mg/ml, about 1 to about 10 mg/ml, about 3 mg/ml or about 5 mg/ml.

[0051] The compositions of the present invention can also be lyophilized using lyophilization techniques known in the art and stored as a powder which can be reconstituted prior to administration. The term "lyophilization" as used herein is a freeze drying or dehydration technique which involves removing a solvent, preferably a water miscible solvent, more preferably water from a composition or the present invention, typically by sublimation under high vacuum when the composition is in a frozen state. Typically, lyophilization is carried out in lyophilization equipment (a lyophilizer), which comprises a drying chamber with variable temperature controls, a condenser to collect water, and a vacuum system to reduce the pressure in the drying chamber.

[0052] The terms "lyophilized composition", as used herein mean the solid residue or powder which is produced, or which remains after the lyophilization procedure as defined above. The lyophilized composition of the present invention typically further comprises a pharmaceutically acceptable excipient. The term "pharmaceutically acceptable excipient" as used herein refers to a substance which is added to a solution prior to lyophilization to enhance characteristics such as the color, texture, strength, and volume of the lyophilized cake. Pharmaceutically acceptable excipients may be, for example, buffers and pH adjusters, crystalline bulking excipients, stabilizers, and tonicity raising agents.

[0053] As used herein, a stabilizer is a composition which maintains the chemical, biological or stability of the chimeric nuclease. Examples of stabilizing agent include polyols, which includes a saccharide, preferably a monosaccharide or disaccharide, e.g., glucose, trehalose, raffinose, or sucrose; a sugar alcohol such as, for example, mannitol, sorbitol or inositol, a polyhydric alcohol such as glycerin or propylene glycol or mixtures thereof and albumin.

[0054] A pharmaceutically acceptable salt is a salt which is suitable for administration to a subject, such as, a human. The chimeric nuclease of the present invention can have one or more sufficiently acidic proton that can react with a suitable organic or inorganic base to form a base addition salt. Base addition salts include those derived from inorganic bases, such as ammonium or alkali or alkaline earth metal hydroxides, carbonates, bicarbonates, and the like, and organic bases such as alkoxides, alkyl amides, alkyl and aryl amines, and the like. Such bases useful in preparing the salts of this invention thus include sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate, and the like. The chimeric nuclease of the present invention having a sufficiently basic group, such as an amine can react with an organic or inorganic acid to form an acid addition salt. Acids commonly employed to form acid addition salts from compounds with basic groups are inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as p-toluenesulfonic acid, methanesulfonic acid, oxalic acid, p-bromophenyl-sulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like. Examples of such salts include the sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caproate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate,

maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, gamma-hydroxybutyrate, glycolate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate, and the like.

[0055] While specific embodiments of the subject invention have been discussed, the above specification is illustrative and not restrictive. Many variations of the invention will become apparent to those skilled in the art upon review of this specification. The full scope of the invention should be determined by reference to the claims, along with their full scope of equivalents, and the specification, along with such variations.

[0056] Unless otherwise indicated, all numbers expressing quantities of ingredients, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term "about." Accordingly, unless indicated to the contrary, the numerical parameters set forth in this specification and attached claims are approximations that may vary depending upon the desired properties sought to be obtained by the present invention.

[0057] The above discussion is meant to be illustrative of the principle and various embodiments of the present invention. Numerous variations and modifications will become apparent to those skilled in the art once the above disclosure is fully appreciated. It is intended that the following claims be interpreted to embrace all such variations and modifications.

[0058] Abbreviations

Abbreviations used herein are defined as follows:

AA	amino acid
Cas9	CRISPR-associated protein 9
CF	Cystic fibrosis
CFTR	Cystic fibrosis transmembrane conductance regulator gene
cjCas9	<i>Campylobacter jejuni</i> Cas9
Cpf1	CRISPR from <i>Prevotella</i> and <i>Francisella</i> 1
CRISPR	Clustered Regulatory Interspaced Short Palindromic Repeats
DLS	Dynamic Light Scattering
DMEM	Dulbecco's Modified Eagle's Medium
DMPE	1,2-ditetradecanoyl- <i>sn</i> -glycero-3-phosphoethanolamine

DNA	deoxyribonucleic acid
DOAB	dioctadecyldimethylammonium bromide
DOPE	1,2-Dioleoyl-sn-glycero-3-phosphoethanolamine
DPPC	Dipalmitoylphosphatidylcholine
<i>E. Coli</i>	<i>Escherichia coli</i>
EDTA	ethylenediaminetetraacetic acid
EGFR	Epidermal growth factor receptor
ELISA	Enzyme-linked immunosorbent assay
fnCas9	<i>Francisella novicida</i> Cas9
HDR	Homology directed repair
IMAC	Immobilized Metal Affinity Chromatography
IPTG	Isopropyl β -D-1-thiogalactopyranoside
MPEG-5000-DMPE	N-(carbonyl-methoxypolyethyleneglycol 5000)-1,2 dipalmitoyl-sn-glycero-3-phosphoethanolamine
NSCLC	Non-small cell lung cancer
NHEJ	Non-homologous end joining
NLS	Nuclear-localized signal
PC	Phosphatidylcholine
PCR	polymerase chain reaction
PE	Phosphoethanolamine
RNA	ribonucleic acid
saCas9	<i>Staphylococcus aureus</i> Cas9
scCas9	<i>Streptococcus canis</i> Cas9
SDS	sodium dodecyl sulfate
spCas9	<i>Streptococcus pyogenes</i> Cas9
TALEN	Transcription activator-like effector nucleases
TEV	Tobacco Etch Virus
TevCas9	Modified I-TevI domain, a linker peptide and modified RNA-guided nuclease <i>Staphylococcus aureus</i> Cas9
ZFN	zinc-finger nucleases

[0059] The Inventors discovered a chimeric nuclease comprising a modified version of an I-TevI domain, a linker peptide and a modified version of an RNA-guided nuclease *Staphylococcus aureus* Cas9 (“saCas9”) (hereinafter referred to as “TevCas9”) that, when mixed with a lipid nanoparticle, with or without exogenous donor DNA, when delivered to cells, replaces DNA sequences in the presence of exogenous donor DNA or deletes defined lengths of DNA in the absence of exogenous donor DNA. The novel chimeric nuclease has been shown to edit genes in human cells, but also cells of other organisms such as bacteria, yeast, insect, plant, or other mammals, either in whole organisms (*in vivo*) or in isolated cells cultures (*ex vivo*).

[0060] The novel chimeric nucleases discovered by the present Inventors present the following advantages over existing gene editing technologies and methods, in particular,

- a. The nuclease, which is a modified version of the TevCas9 nuclease, is capable of targeting two independent target sites as a single protein and cleaving the DNA at one or both of these sites. It can be reprogrammed to many different target DNA sequences through modifying one or more of the I-TevI domain, the linker domain, the Cas9 domain or the guide RNA (which targets the Cas9 domain to its target sequence);
- b. If the nuclease cleaves at two sites, it cleaves out precise lengths of DNA (~30 – 36 bases depending on the sites targeted by I-TevI and Cas9);
- c. The Cas9 domain contains a mutation (D10E) which is rationally designed to modify the Cas9 nuclease activity and/or increase the Cas9 domain’s specificity for its target binding site;
- d. In the presence of exogenous donor DNA, the invention is designed to replace target DNA sequences in a higher percentage of cells than existing technologies or practices;
- e. The nuclease can be purified as a single contiguous protein combined with a guide RNA, which simplifies manufacturing;
- f. The lipid nanoparticle allows for non-viral delivery to target cells with high efficiency and low toxicity, allowing for controlled dosing of the nuclease. Although other lipid-based nuclease delivery technologies exist, none are of a composition suitable for use *in vivo*;

- g. The lipid nanoparticles are also designed for the delivery of nuclease through nebulization (inhalation);
- h. One version of the nuclease targets and cleaves the *CFTR* gene to correct the *CFTR delta F508* mutation for the treatment of Cystic Fibrosis (SEQ ID NO 1); and
- i. Another version of the nuclease is designed to target and cleave the clinically relevant *EGFR exon 19 deletion* mutations (SEQ ID NOS 2 – 4), which are present in a variety of cancers, including non-small-cell lung cancer (NSCLC).

[0061] The fusion of a GIY-YIG nuclease, such as I-TevI, through a flexible linker to DNA binding domains is known (WO2014/121222). A prior version of the dual-cleaving TevCas9 has been described which comprises amino acids 1 – 92 of the wild-type I-TevI nuclease domain, a linker region comprising amino acids 93 – 169 of I-TevI linker region and the *Streptococcus pyogenes* Cas9 (“spCas9”)(Wolfs JM *et al.*, (2016), ‘Biasing Genome-Editing Events Toward Precise Length Deletions with an RNA-Guided TevCas9 Dual Nuclease,’ *Proc Natl Acad Sci USA*, 113(52):14988-93). The chimeric nuclease of the invention comprises the following:

- i. An I-TevI nuclease domain which binds a new target sequence allowing to target clinically relevant gene sequences, such as the *CFTR* gene;
- ii. Various flexible linker regions intended to confer different DNA binding or nuclease activity to TevCas9;
- iii. A saCas9 nuclease domain (US-1988/065406 B2). The use of saCas9 over spCas9 results in a smaller DNA coding sequence (~3.7 kilobases for Tev-saCas9 versus ~4.6 kilobases for Tev-spCas9) and lower molecular weight TevCas9 protein (~144 kilodaltons for Tev-saCas9 versus ~179 kilodaltons for Tev-spCas9) which is more amenable to multiple delivery technologies; cleaving by the saCas9 domain between the 3rd and 4th nucleotide is predictable compared to spCas9 which is more amenable to defined length deletions, as discovered by the inventors of the claimed technology.
- iv. One version where the guide RNA is targeted to specific *CFTR* gene sequence near the *CFTR delta F508* mutation; and
- v. A second version where the guide RNA is targeted to specific *EGFR* gene sequences and is intended to cleave only DNA with appropriated spaced I-

TevI site and Cas9 target site. Such appropriately spaced sites occur in certain *EGFR* exon 19 deletion mutations (SEQ ID NO 2 – 4) but not in wild-type *EGFR* (SEQ ID NO 5);

- a. The invention comprises lipid nanoparticles of certain compositions that are selectively sized to a mean diameter of approximately 100 nM. These lipid nanoparticles are capable of delivering the nuclease to cells with high efficiency and low toxicity;
- b. A pharmaceutical formulation of the lipids, nuclease, and exogenous donor DNA;
- c. A pharmaceutical formulation of the lipids, nuclease and exogenous donor DNA which is suitable for nebulization (inhalation); and
- d. A version of the invention which contains exogenous donor DNA that when delivered with the TevCas9 nuclease in the lipid nanoparticle is capable of integrating into the region between or around the two sites targeted by the nuclease.

[0062] The novel chimeric nuclease compositions of the instant application contain different combinations of an I-TevI domain, a linker domain, a Cas9 domain and a guide RNA.

[0063] The versions that target the *CFTR* gene are comprised of:

- i. An I-TevI domain of amino acid sequence according to SEQ ID NO: 6;
- ii. A linker domain according to any one of SEQ ID NOS: 7 – 12;
- iii. A saCas9 domain of the amino acid sequence according to SEQ ID NO: 13; and
- iv. A guide RNA of the RNA sequence according to SEQ ID NO: 15 or 21.

[0064] The versions that target the *EGFR* gene are comprised of:

- i. An I-TevI domain of amino acid sequence according to SEQ ID NO: 6;
- ii. A linker domain with any one of the amino acid sequences according to SEQ ID NOS: 7 – 12;
- iii. A saCas9 domain of the amino acid sequence according to SEQ ID NO: 13; and
- iv. A guide RNA of the RNA sequence in SEQ ID NO: 16.

[0065] The I-TevI domain of the preferred embodiment is a 93-amino acid I-TevI domain of the *Enterobacteria Phage T4* according to the following sequence:

MGKSGIYQIKNTLNNKVYVGS AKDFEKRWKRHF KDLEKGCHSSIKLQRSFNKHG NVFE
CSILEEIPYEKDLIERENFWIKELNSKINGYNIA (SEQ ID NO: 6)

[0066] The saCas9 of the preferred embodiment is a polypeptide comprised of 1,053 amino acids according to the following sequence:

MKRNYILGLDIGITSVGYGIIDYETRDVIDAGVRLFKEANVENNEGRRSKRGARRLKRRR
RHRIQRVKKLLFDYNLLTDHSELGINPYEARVKGLSQKLSEEEFSAALLHLAKRRGVH
NVNEVEEDTGNELSTKEQISRNSKALEEKYVAELQLERLKKDGEVRGSINRFKTS DYVK
EAKQLLKVKQAYHQLDQSFIDTYIDLLETRRTYYEGPGEGSPFGWKDIKEWYEMLMGH
CTYFPEELRSVKYAYNADLYNALNDLNNLVITRDENEKLEYEYEFQIIENVFKQKKKPT
LKQIAKEILVNEEDIKGYRVTSTGKPEFTNLKVYHDIKDITARKEIENAEELLDQIAKILTIY
QSSEDIQEELTNLNSELTQEEIEQISNLKGYTGTHNLSLKAINLILDELWHTNDNQIAIFNR
LKLVPKKVDLSQQKEIPTTLVDDFILSPVVKRSFIQSIKVINAIKKYGLPNDIIELAREKN
SKDAQKMINEMQKRNRQTNERIEEII RTTGKENAKYLIEKIKLHDMQEGKCLYSLEAIPL
EDLLNNPFNYEVDHII PRSVSFDNSFNKVLVKQEENSKKGNRTPFQYLSSSDSKISYETF
KKHILNLA KGKGRISKTKKEYLLEERDINRFSVQKDFINRNLVDTRYATRGLMNLLRSYF
RVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGYKHAEDALIANADFIKWKLDK
AKKVMENQMFEKQAESMPEIETEQEYKEIFITPHQIKHIKDFKDYKYSHRVDKKPNREL
INDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMYHHD PQTYQKLK
LIMEQYGDEKNPLYKYEEETGNYLTKYSKKDNGPVIKKIKYYGNKLNAHL DITDDYPN
SRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYYEVNSKCYEEAKKLKKIS
NQAEFIASFYNNDLIKINGEL YRVIGVNNDLLNRIEVMIDITYREYLENMNDKRPPRIIK
TIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG (SEQ ID NO: 13)

[0067] The saCas9 with a Glu¹⁰ mutation of the preferred embodiment is a polypeptide comprised of 1,053 amino acids according to the following sequence (the mutation is underlined):

MKRNYILGLEIGITSVGYGIIDYETRDVIDAGVRLFKEANVENNEGRRSKRGARRLKRRR
RHRIQRVKKLLFDYNLLTDHSELGINPYEARVKGLSQKLSEEEFSAALLHLAKRRGVH
NVNEVEEDTGNELSTKEQISRNSKALEEKYVAELQLERLKKDGEVRGSINRFKTS DYVK
EAKQLLKVKQAYHQLDQSFIDTYIDLLETRRTYYEGPGEGSPFGWKDIKEWYEMLMGH
CTYFPEELRSVKYAYNADLYNALNDLNNLVITRDENEKLEYEYEFQIIENVFKQKKKPT
LKQIAKEILVNEEDIKGYRVTSTGKPEFTNLKVYHDIKDITARKEIENAEELLDQIAKILTIY
QSSEDIQEELTNLNSELTQEEIEQISNLKGYTGTHNLSLKAINLILDELWHTNDNQIAIFNR

LKLVPKKVDLSQQKEIPTTLVDDFILSPVVKRSFIQSIKVINAIKKYGLPNDIIIELAREKN
 SKDAQKMINEMQKRNRQTNERIEEIIIRTTGKENAKYLIEKIKLHDMQEGKCLYSLEAIPL
 EDLLNNPFNYEVDHIIIPRSVSFDNSFNKVLVKQEENSKKGNRTPFQYLSSSDSKISYETF
 KKHILNLAKGKGRISKTKKEYLLEERDINRFSVQKDFINRNLVDTRYATRGLMNNLLRSYF
 RVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGYKHHAEALIIANADFIFKEWKLDK
 AKKVMENQMFEEKQAESMPEIETEQEYKEIFITPHQIKHIKDFKDYKYSHRVDKKNREL
 INDTRYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMYHHDPQTYQKLK
 LIMEQYGDEKNPLYKYEEETGNYLTKYSKKDNGPVIKKIKYYGNKLNHLADITDDYPN
 SRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYYEVNSKCYEEAKKLKKIS
 NQAEFIASFYNNDLIKINGELYRVIGVNNDLLNRIEVNMIDITYREYLENMNDKRPPRIIK
 TIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG (SEQ ID NO: 14)

[0068] The guide RNA of the version that targets the *CFTR* gene is comprised of 101 ribonucleotides according to the sequences:

GCGUCAUCAAAGCAUGCCAACGUUUUAGUACUCUGGAAACAGAAUCUACUAAAA
 CAAGGCAAAAUGCCGUGUUUAUCUCGUCAACUUGUUGGCGAGAUUUU (SEQ ID
 NO: 15)

AUAUCAUUGGUGUUUCCUAUGGUUUUAGUACUCUGGAAACAGAAUCUACUAAAA
 CAAGGCAAAAUGCCGUGUUUAUCUCGUCAACUUGUUGGCGAGAUUUU (SEQ ID
 NO: 21)

[0069] The guide RNA of the version that targets the *EGFR* gene is 101 ribonucleotides in length according to the following sequence:

AAUUUUAACUUUCACCUUCGUUUUAGUACUCUGGAAACAGAAUCUACUAAAA
 CAAGGCAAAAUGCCGUGUUUAUCUCGUCAACUUGUUGGCGAGAUUUU (SEQ ID
 NO: 16).

[0070] The linkers used in any of the above constructs may be selected from the group consisting of:

SEQUENCE (amino acid count)	MUTATION(S) (indicated by underline)	SEQ ID NO:
DATFGDTCSTHPLKEEIIKKRSETFKAKMLKLGPDGRKALYSKPGSK NGRWNPECHKFCKCGVRIQTSAYTCSKCRNGGSGGS (83 AA)	V117F	7
DATFGDTCSTHPLKEEIIKKRSETVKAKMLKLGPDGRKALYSRPGSK SGRWNPECHKFCKCGVRIQTSAYTCSKCRNGGSGGS (83 AA)	K135R N140S	8

DATFGDTCSTHPLKEEIIKKRSETFKAKMLKLGPDGRKALYSRPGSKS GRWNPETHKFCKCGVRIQTSAYTCSKCRNGGSGGS (83 AA)	V117F K135R N140S	9
DATFGDTCSTHPLKEEIIKKRSETFKAKMLKLGPDGRKALYSRPGSKS GRWNPETHKFCKCGVRIQTSAYTCSKCRGGSGGTGGS (86 AA)	K135R N140S	10
DATFGDTCSTHPLKEEIIKKRSETFKAKMLKLGPDGRKALYSRPGSKS GRWNPETHKFCKCGVRIQTSAYTCSKCRGGGGSGGGGS (87 AA)	K135R N140S	11
DATFGDTCSTHPLKEEIIKKRSETFKAKMLKLGPDGRKALYSRPGSKS GRWNPETHKFCKCGVRIQTSAYTCSKCRKESGSVSSEQLAQFRSLD (95 AA)	K135R N140S	12

SYNTHESIS

Example 1: A method to manufacture the TevCas9 nuclease:

[0071] The DNA coding sequences of the above mentioned I-TevI domain, linker domain and Cas9 domain are synthesized as one contiguous DNA sequence using techniques known in the art. Gene synthesis was conducted by Bio Basic Inc. (Markham, On, Canada). Briefly, short oligonucleotides (~50 – 60 base pairs) are synthesized which contain regions of overlap to cover the entire sequences of I-TevI domain, linker domain and Cas9 domain. The oligonucleotides are mixed together in blocks of approximately 1 kilobase of the sequence to be synthesized and polymerase chain reaction (PCR) is used to synthesize these ~1 kilobase blocks. The ~1 kilobase blocks are then mixed and subjected to PCR to synthesis the I-TevI domain, linker domain and Cas9 domain. Further, to enhance expression of TevCas9 in *E. coli* and simplify restriction enzyme digestion, the DNA sequence of TevCas9 was optimized prior to synthesis. First, three-base pair DNA codons that are infrequently used by *Escherichia coli* (“*E. coli*”) were replaced with those that occur more frequently (for example, of the 6 codons coding for the amino acid arginine, the relative abundance of the codon AGG is 0.03 compared to 0.42 for the codon CGT). In total, 37% of the codons were changed to those preferred by *E. coli*. Second, the content on the nucleotides cytosine and guanine was increased from 39.6% to 48.6%. Third, two *E. coli* ribosome binding sites were removed from the sequence. Fourth, a *NdeI* restriction endonuclease site was removed from the internal sequence. The contiguous DNA is digested with the restriction endonucleases *NdeI* and *BamHI* (New England Biolabs, Ipswich, MA, United States), whose target sites occur only once in the DNA sequence, and then inserted using DNA ligase (New England Biolabs, Ipswich, MA, United States) into a similarly digested pET-11a expression vector (EMD Millipore, Burlington, MA, United States) suitable for expression of

TevCas9 in *E. coli*. The pET-11a vector containing TevCas9 is transformed into the *E. coli* expression strain T7 Express (New England Biolabs #C2566, Ipswich, MA, United States) which has been optimized for expression of proteins, including nucleases. Alternatively, the *E. coli* expression strain BL-21(DE3) (New England Biolabs #C2527, Ipswich, MA, United States) is used. Successful transformations are confirmed by resistance of the *E. coli* to ampicillin or tetracycline and the coding sequence of TevCas9 is verified by DNA sequencing of the expression vector derived from the transformed *E. coli*. The transformed *E. coli* is grown at 37°C to an optical density of 0.4 to 0.6 as measure by spectrophotometry at a wavelength of 600 nm and the expression of the TevCas9 protein from the pET-11a vector in the transformed *E. coli* expression strain is induced using IPTG for 10 - 12 hours at 16°C. Successful expression of TevCas9 is verified by the presence of an approximately 150 kDa band on a Coomassie-stained SDS-polyacrylamide gel in a sample of the induced material compared to the uninduced sample. The *E. coli* cells are harvested by centrifugation and resuspended in lysis buffer comprising 10 mM imidazole (Sigma, St. Louis, MO, United States), 300 - 500 mM sodium chloride (Sigma-Aldrich, St. Louis, MO, United States) and 50 mM sodium phosphate (dibasic) (Sigma, St. Louis, MO, United States), pH 8.0 [Buffer 1]. Alternatively, 10 mM Tris Hydrochloride (Sigma, St. Louis, MO, United States), pH 8 is substituted for substituted for sodium phosphate (dibasic) in Buffer 1. The *E. coli* is lysed by homogenization using a high pressure liquid, homogenizer (Avestin Inc., Ottawa, ON, Canada) operated at 600–1000 bar, or any other suitable lysis method known in the art, such as sonication using a sonifier (Branson Ultrasonics Corp, Danbury, CT, United States) lysozyme treatment, homogenization using a French pressure cell (Glen Mills Inc., Clifton, NJ, United States) or homogenization using a Dounce homogenizer (Corning Inc., Corning, NY, United States). The lysed material is centrifuged at 12,000rpm at 4°C for 20 – 30 mins and the supernatant containing soluble TevCas9 is used for the subsequent purification steps. The pellet contains cell debris, insoluble intracellular material, as well as any insoluble TevCas9. Successful lysis and solubility is verified by the presence of an approximately 150 kDa band on a Coomassie-stained SDS-polyacrylamide gel in a sample of the supernatant when compared to a resuspended sample of the pellet.

[0072] The TevCas9 nuclease is purified in the following steps:

1. The lysate containing the nuclease is applied to an immobilized metal affinity chromatography (IMAC) column (GE Healthcare Bio-Sciences AB, Uppsala, Sweden) which binds the nuclease.
2. The IMAC column is washed with Buffer 1.
3. The TevCas9 remaining bound to the column is eluted with a solution comprising 250 mM imidazole (Sigma, St. Louis, MO, United States), 300 mM-500 mM sodium chloride (Sigma-Aldrich, St. Louis, MO, United States) and 50 mM sodium phosphate (dibasic) (Sigma, St. Louis, MO, United States), pH 7.6 – 8.0 [Buffer 2]. Alternatively, 10 mM Tris Hydrochloride (Sigma, St. Louis MO, United States), pH 7.6 - 8 is substituted for substituted for sodium phosphate (dibasic) in Buffer 2.
4. The eluate is treated with Tobacco Etch Virus (TEV) protease (New England Biolabs, Ipswich, MA, United States) and incubated with the appropriate guide RNA. The guide RNA is synthesized by Integrated DNA Technology Inc. (Coralville, IA, United States).
5. The treated eluate is re-applied to the IMAC column and the flow-through containing the TevCas9 nuclease and guide RNA is collected.
6. Successful purification of the TevCas9 nuclease is confirmed by the presence of a 150 kilodalton protein band on a Coomassie-stained SDS-polyacrylamide gel. Successful co-purification of TevCas9 with the guide RNA is confirmed by treating a sample of the eluate with Proteinase K (New England Biolabs, Ipswich, MA, United States), then splitting the sample in two and further treating one subsample with RNase A (New England Biolabs, Ipswich, MA, United States) and the other in control buffer without RNase A. A ~100 nucleotide RNA band will be visible on an urea-polyacrylamide gel in the control sample and will be absent in the RNase A-treated sample.
7. The solution containing the TevCas9 nuclease and guide RNA is dialyzed into a solution comprising phosphate buffered saline, pH 7.4.

Example 2: A method to manufacture the lipid nanoparticles

[0073] The lipid nanoparticles of the preferred embodiment are comprised of one of the following mixtures:

- I. Lipid nanoparticle No. 1 comprises DOPE (Avanti Polar Lipids, Alabaster, AL, United States) and MPEG-5000-DMPE (Avanti Polar Lipids, Alabaster, AL, United States) in a molar ratio of 2:0.05, respectively;
- II. Lipid nanoparticle No. 2 comprises DPPC (Avanti Polar Lipids, Alabaster, AL, United States), cholesterol (SUPELCO, Bellefonte, PA, United States) and DOBA (Sigma, St. Louis, MO, United States) in a molar ratio of 7:2:1, respectively; and
- III. Lipid nanoparticle No. 3 comprised DPPC, cholesterol and MPEG-5000-DMPE (Avanti Polar Lipids, Alabaster, AL, United States) in a molar ratio of 4:1:0.125, respectively.

[0074] Lipid nanoparticles are manufactured to a mean diameter of approximately 100 nM.

[0075] One of the lipid mixtures No. 1 – 3 is selected. For example, DOPE and MPEG-5000-DMPE are mixed together in the appropriate molar ratios in an organic solvent, such as chloroform. The organic solvent is then evaporated and the dried lipid mixture is re-suspended using vigorous vortexing in a solution comprising phosphate buffered saline, pH 7.4. The re-suspended lipid mixture is then extruded through a 100 nM polycarbonate membrane (T&T Scientific Corporation, Knoxville, TN, United States) equilibrated in phosphate buffered saline to create lipid nanoparticles of an approximate mean diameter of 100 nM. The solution is filter sterilized through 0.2 μ M sterile filter (VWR Scientific, Radnor, PA, United States). The mean diameter and size distribution of the lipid nanoparticles is determined by Dynamic Light Scattering (DLS) using a Zetasizer (Malvern Panalytical Ltd, Malvern, United Kingdom), or another suitable technique, known in the art.

Example 3: Composition of the donor DNA

[0076] The donor DNA comprises DNA sequences that are intended to repair a genetic defect. It also comprises DNA sequences which are not found in the target genomic DNA; these sequences do not interfere with the normal gene function but are intended to knockout the I-TevI and or Cas9 sites and/or introduce one or more DNA sequences which are used to track the successful repair of the target gene. Examples of donor DNA include, but are not limited to the following:

- I. Linear single-strand DNA of varying lengths comprising homologous regions flanking the sites targeted/cleaved by TevCas9;
- II. Linear double-strand DNA of varying lengths comprising homologous regions flanking the sites targeted/cleaved by TevCas9;

- III. Double-strand DNA of the same length cleaved by the nuclease and also comprising complimentary DNA ends to those cleaved by TevCas9;
- IV. Circular double-strand DNA comprising homologous regions flanking the sites targeted/cleaved by TevCas9; and
- V. Circular double-strand DNA comprising an I-TevI target site and Cas9 target site where the product cleaved from the double-strand DNA contains complimentary ends to those cleaved by TevCas9.

Example 4: A method for assembling the lipid-encapsulated TevCas9 and transfecting cells

[0077] For *ex vivo* cell transfections: To assemble the lipid-encapsulated TevCas9, a lipid nanoparticle is mixed with the TevCas9 in a 2000:1 molar ratio in Dulbecco's Modified Eagle's Medium (DMEM) (Sigma, St. Louis, MO, United States) and incubated at room temperature for 10 minutes. Cells are transfected with 8.7×10^{-17} to 3.1×10^{-17} moles of lipid-encapsulated TevCas9 per cell.

[0078] For *in vivo* cell transfection: To assemble the lipid-encapsulated TevCas9, a lipid nanoparticle is mixed with the TevCas9 in a 2000:1 molar ratio in phosphate buffered saline and incubated at room temperature for 10 minutes. The molar ratio of lipid-encapsulated TevCas9 per cell for *in vivo* transfections is to be determined.

Other Embodiments

[0079] The nuclease might contain different combinations of the I-TevI domain, linker domain, Cas9 domain or guide RNA as highlighted below.

[0080] **Modifications of the I-TevI Domain:** Other versions of the I-TevI nuclease domain might contain different combinations of mutations to alter the site targeted by the I-TevI domain or the activity of the I-TevI domain, including mutations that alter the sequence recognized by I-TevI, such as K26 and/or C39. Other versions of the nuclease might substitute the I-TevI domain with other GIY-YIG nuclease domains, such as I-BmoI, Eco29kI, etc. Other versions do not contain Met¹ as a result of processing when expressed in *E. coli*.

[0081] **Modifications of the Linker Domain:** The linker domain might comprise one more of the following to alter binding specificity or activity of TevCas9, including: a). The I-TevI linker domain comprising one or more mutations to amino acid T95, V117, K135, Q158 or N140; b). The linker might contain various combinations of the amino acids shown in SEQ ID NO: 9 - 12.

[0082] Modifications of the Cas9 Domain: Other versions of the Cas9 domain might contain the following: a). A version of the saCas9 domain comprising a D10E mutation (SEQ ID NO: 14); b). A version of the saCas9 domain that nicks target DNA on one strand of the target DNA, for example the H557A mutation (SEQ ID NO: 17); c). A version of the saCas9 domain that binds target DNA but does not cleave it, for example mutations at both D10A and H557A mutations (SEQ ID NO: 18); d). A version of the previously described spCas9 EQR mutant comprising the mutations D1135E, R1335Q and T1337R combined with the D10E mutation (SEQ ID NO: 19); and e). A version of the previously described spCas9 EQR mutant comprising the mutations D1135E, R1335Q and T1337R combined with the D10E mutation and a mutation that nicks target DNA on one strand of the target DNA, for example the H840A mutation (SEQ ID NO: 20). Other versions of the saCas9 domain do not contain Met¹.

[0083] Other versions might substitute other nucleases or DNA binding domains for the Cas9 domain, such as: a). Meganucleases such as the families LAGLIDADG, His-Cys Box, H-N-H, PD-(D/E)xK, Vsr-like, etc.; b). Zinc-finger nucleases; c). Other CRISPR proteins such as scCas9, fnCas9, cjCas9, Cpf1, Cas12a, Cas13a, Cas3, etc.; and d). Other DNA binding domains such as zinc-finger motifs, TALE activator domains, etc.

[0084] Modifications of the Guide RNA: a). Other versions of the guide RNA might target the same region of DNA in the *CFTR* gene or *EGFR* gene, but contain different sequences to account for genetic polymorphism in populations; b). Other versions of the guide RNA might target different sequences in the *CFTR* gene or *EGFR* gene; c). Other version of the guide RNA might target other sequences in a genome to retarget the nuclease to additional clinically relevant targets; d). Other versions of the guide RNA might contain bridged nucleic acids (“BNAs”) to enhance target site specificity; and e). Other versions might contain a mixture of guide RNAs to target multiple sequences within the same gene.

[0085] Modifications of the Lipid Nanoparticles: a). Other versions of the lipid nanoparticle No. 1, 2 or 3 might have different ratios of each lipid component; b). Other versions of the lipid nanoparticle might have different mean diameters; c). Other versions of the lipid nanoparticle might include different cationic or neutral lipids; d). Other versions of the lipid nanoparticle might include peptides that target specific cell types; e). Other versions of the lipid nanoparticle might include compounds that bind DNA, such as GL67 (N⁴-Cholesteryl-Spermine); f). the lipid nanoparticle might be lyophilized for enhanced stability; and g). the lipid nanoparticle might be

resuspended in a solution other than phosphate buffered saline, such as sterile isotonic saline, water for injection, etc.

[0086] Modifications of the Composition of the donor DNA: a). Other version of the linear double-strand donor DNA might contain longer regions of single-strand DNA that is complementary to the target sequence; and b). Other versions of the circular double-strand DNA might contain other DNA sequences intended to increase the rate of homology-directed repair.

[0087] Variations to the method of assembling the lipid-encapsulated nuclease and transfecting cells: a). Other versions of the lipid-encapsulated nucleases might contain different molar ratios of lipid nanoparticle to nuclease; b). Media other than DMEM or phosphate buffered saline might be used for the incubation step; c). The nuclease and lipid nanoparticles might be incubated for less or more than 10 minutes; and d). Other molar amounts of nuclease per cell might be used in a transfection reaction.

[0088] Variations to the method to manufacture the nuclease: a). Other *E. coli* expression strains might be used, such as LS5218 (*Escherichia coli* Genetic Stock Center – Yale University, New Haven, CT, United States) or BL21-DE3 (New England Biolabs, Ipswich, MA, United States) b). Buffer 1 or 2 might contain different concentrations of imidazole, sodium chloride, sodium phosphate (dibasic), or tris hydrochloride and be buffered to a different pH; c). Other processing steps might be used, such as cation or anion exchange chromatography; d). The nuclease might be dialyzed into a solution other than phosphate buffered saline, such as sterile isotonic saline, water for injection, etc.; e). the nuclease might be lyophilized for enhanced stability; f). the guide RNA may be co-expressed from the pACYC-Duet1 expression vector (EMD Millipore, Burlington, MA, United States). The DNA coding sequence of the guide RNA is synthesized (Integrated DNA Technology Inc., Coralville, IA, United States), digested with restrictions endonucleases and inserted into similarly-digested second expression site in the pACYC-Duet1 expression vector; and g). The guide RNA may be synthesized from double-strand DNA by transcribing the guide using the T7 RNA Polymerase HiScribe Kit (New England Biolabs #E2040S, Ipswich, MA, United States) and purifying the guide using an RNA Cleanup Kit (New England Biolabs #T2030L, Ipswich, MA, United States).

TESTING

Example 1: A method to demonstrate correction of CFTR delta F508 and CFTR protein functionality in a model cell line

[0089] A culture of immortalized epithelial cells homozygous for the *CFTR delta F508* mutation, such as the CuFi-1 cell line (ATCC® CRL-4013™, American Type Culture Collection, Manassas, VA, United States), is treated with a range of concentrations of lipid-encapsulated TevCas9 and donor DNA (Specific Biologics, Toronto, ON, Canada) in a pharmaceutical formulation targeted to the *CFTR delta F508* mutation. An appropriate control cell line, such as NuLi-1 (ATCC® CRL-4011™, American Type Culture Collection, Manassas, VA, United States) immortalized epithelial cells homozygous for wild-type *CFTR* is also be used.

[0090] The proportion of cells with *CFTR delta F508* corrected relative to uncorrected cells is measured by the T7 endonuclease I assay (EnGen® Mutation Detection Kit, New England Biolabs #E3321, Ipswich, MA, United States), restriction endonuclease digestion (New England Biolabs, Ipswich, MA, United States) of PCR-amplified target site, deep gene sequencing using an Illumina MiSeq system and barcoded primers flanking the target site (Illumina, San Diego, CA, United States), or other suitable method. The effects of TevCas9 treatment in the control cell line (e.g. NuLi-1 (ATCC® CRL-4011™, American Type Culture Collection, Manassas, VA, United States) is measured. *CFTR* functionality is measured using short circuit current measurements in an Ussing Chamber (Warner Instruments, Hamden, CT, United States) in the presence of a chloride ion gradient in the treated CuFi-1 culture (ATCC® CRL-4013™, American Type Culture Collection, Manassas, VA, United States) versus mock-treated CuFi-1 culture (ATCC® CRL-4013™, American Type Culture Collection, Manassas, VA, United States). The effects of TevCas9 treatment in the control cell line (e.g. NuLi-1) is also measured.

[0091] To demonstrate disruption of the *EGFR* exon 19 deletion mutations(s) and *EGFR* expression and activity in a model cell line, a culture of immortalized epithelial cells expressing an *EGFR* exon 19 deletion mutation(s), such as the HCC827 cell line (ATCC® CRL-2868™, American Type Culture Collection, Manassas, VA, United States) is treated with a range of concentrations of lipid-encapsulated TevCas9 (Specific Biologics, Toronto, ON, Canada) in a pharmaceutical formulation of phosphate buffered saline, sterile isotonic saline or water for injection targeted to the *EGFR* exon 19 deletion. Appropriate control cell lines, such as NuLi-1 (ATCC® CRL-4011™, American Type Culture Collection, Manassas, VA) or immortalized epithelial cell lines homozygous for wild-type *EGFR*, are used.

[0092] The proportion of cells with the *EGFR* exon 19 deletion disrupted relative to uncorrected cells is measured by the T7 endonuclease I assay (EnGen® Mutation Detection Kit, New England Biolabs #E3321, Ipswich, MA, United States), restriction endonuclease digestion (New England Biolabs, Ipswich, MA, United States) of PCR-amplified target site, deep gene sequencing using an Illumina MiSeq system and barcoded primers flanking the target site (Illumina, San Diego, CA, United States), or other suitable method. The effects of TevCas9 treatment in the control cell line (e.g. NuLi-1 (ATCC® CRL-4011™, American Type Culture Collection, Manassas, VA, United States)) is also measured. EGFR protein expression and activity are measured using an enzyme-linked immunosorbent assay (ELISA) (Sigma, St. Louis, MO, United States) that detects phosphorylated (*i.e.* activated), unphosphorylated and total EGFR protein in the treated HCC827 culture (ATCC® CRL-2868™, American Type Culture Collection, Manassas, VA, United States) versus mock-treated HCC827 culture. The effects of TevCas9 treatment in the control cell line (e.g. NuLi-1 (ATCC® CRL-4011™, American Type Culture Collection, Manassas, VA, United States)) is also measured.

Example 1: Animal model testing planned to show efficacy and determine dose-limiting toxicity

[0093] In an example method to demonstrate correction of *CFTR delta F508* and/or Cystic Fibrosis symptoms with lipid-encapsulated TevCas9 treatment in an animal model (for example, mouse, rat, minipig or ferret), the lipid-encapsulated TevCas9 in a pharmaceutical formulation of phosphate buffered saline, sterile isotonic saline or water for injection targeted to the *CFTR delta F508* is delivered directly to the lungs by either intubation or intranasal delivery. The procedure time is approximately 30-6000 seconds per treatment, depending on the animal model used.

[0094] In another method, the lipid-encapsulated TevCas9 in a pharmaceutical formulation targeted to the *CFTR delta F508* is nebulized with a commercial nebulizer (Aeroneb®, AeroEclipse®, (Trudell Medical, London, ON, Canada)) or PARI-LC Plus®, (PARI USA, Midlothian, VA, United States)). The average size of the lipid nanoparticle of approximately 100 nM is confirmed post-nebulization by Dynamic Light Scattering (DLS) using a Zetasizer (Malvern Panalytical Ltd, Malvern, United Kingdom), or another suitable technique, known in the art. The composition and concentration of the lipid-encapsulated TevCas9 is confirmed post-nebulization using the MicroGram Lipid Assay Kit (ProFoldin, Hudson, MA, United States) and the presence of an approximately 150 kDa band on a Coomassie-stained SDS-polyacrylamide gel. For measurement of the rate of gene correction, a representative ovine (minipig) animal

model (Exemplar Genetics, Sioux City, IA, United States) that is homozygous for the *CFTR delta F508* mutation is exposed through the mouth, nose or directly to the lungs with the lipid-encapsulated TevCas9 targeted to *CFTR delta F508*, as well as a suitable control. General maintenance of these animals includes breeding and farrowing; age-appropriate, bio-secure housing; sound nutrition; basic vaccinations and veterinary care; and documentation consistent with animal welfare guidelines. Maintenance of these animals specific to the *CFTR delta F508* may include one or more of the following: surgery to address intestinal obstruction; pancreatic enzyme replacement therapy; vitamins and H2 blockers; and / or proton pump inhibitors to improve gastric acid control. The minipigs are treated with a range of concentrations of lipid-encapsulated TevCas9 that are predicted to be effective from the model cell line studies above for 2 days to 4 weeks for acute toxicity studies and up to 24 months for chronic toxicity studies.

[0095] The general health of the animal is monitored post-treatment to assess for any treatment-related adverse events, such as changes in behavior, weight, or food consumption; immune responses; changes to cardiovascular health; mortality, etc. Other efficacy measures post-treatment may include:

- I. Forced-expiration, such as forced expiratory volume (or other suitable method) in each animal post-treatment;
- II. Overall survival of each animal relative to the control;
- III. Other measures of lung function (for example, utilization of mechanical ventilator that can perform general lung function assessments); and
- IV. Measurements of the mutation *in vivo* through tissue sampling and mutation detection methods, such as by polymerase chain reaction.

After the treatment(s) with lipid-encapsulated TevCas9, the animals are sacrificed and the lung and tracheal tissue are harvested.

[0096] The proportion of cells with *CFTR delta F508* corrected relative to uncorrected cells is measured by the T7 endonuclease I assay (EnGen® Mutation Detection Kit, New England Biolabs #E3321, Ipswich, MA, United States), restriction endonuclease digestion (New England Biolabs, Ipswich, MA, United States) of PCR-amplified target site, deep gene sequencing using an Illumina MiSeq system and barcoded primers flanking the target site (Illumina, San Diego, CA, United States) or other suitable method.

[0097] In a method to demonstrate disruption of the *EGFR* exon 19 deletion mutation(s) and/or Non-small-cell lung cancer (NSCLC) symptoms with TevCas9 treatment in an animal model, the lipid-encapsulated TevCas9 targeted to the *EGFR* exon 19 deletion mutations in a pharmaceutical formulation of phosphate buffered saline, sterile isotonic saline or water for injection is delivered directly to the lungs through the mouth, nose or directly to the lungs. The procedure time is approximately 30-6000 seconds per treatment, depending on the animal model used.

[0098] In another method, the lipid-encapsulated TevCas9 targeted to the *EGFR* exon 19 deletion mutations in a pharmaceutical formulation is nebulized with a commercial nebulizer ((Aeroneb®, AeroEclipse®, (Trudell Medical, London, ON, Canada) or PARI-LC Plus®, (PARI USA, Midlothian, VA, United States)). The average size of the lipid nanoparticle of approximately 100 nM is confirmed post-nebulization by Dynamic Light Scattering (DLS) using a Zetasizer (Malvern Panalytical Ltd, Malvern, United Kingdom), or another suitable technique, known in the art. The composition and concentration of the lipid-encapsulated TevCas9 nanoparticle is confirmed post-nebulization using the MicroGram Lipid Assay Kit (ProFoldin, Hudson, MA, United States) and the presence of an approximately 150 kDa band on a Coomassie-stained SDS-polyacrylamide gel. For measurement of the rate of gene disruption, a representative murine (mouse) animal model that is homozygous for an *EGFR* exon 19 deletion mutation(s) is exposed through the nose, mouth or directly to the lungs with the lipid-encapsulated TevCas9 targeted to *EGFR* exon 19 deletion. The mice are treated with a range of concentrations of TevCas9 that are predicted to be effective from the model cell line studies for 2 days to 4 weeks for acute toxicity studies and up to 24 months for chronic toxicity studies.

[0099] The general health of the animal is monitored post-treatment to assess for any treatment-related adverse events, such as changes in behavior, weight, or food consumption; immune responses; changes to cardiovascular health; mortality, etc. Other efficacy measures post-treatment may include:

- I. Quantification of EGFR-activating protein through positron emission tomography (PET) with an EGFR mutant tracer;
- II. Overall survival of each animal relative to the control(s); and
- III. Measures of tumor formation / reduction in each animal over time.

Measurements of the mutation *in vivo* through tissue sampling and mutation detection methods, such as the Cobas® EGFR mutation test version 2 (Roche Diagnostics, Risch-Rotkreuz, Switzerland). After treatment with nebulized lipid-encapsulated TevCas9, the animals are sacrificed and the lung and tracheal tissue are harvested.

[0100] The proportion of cells with the *EGFR* exon 19 deletion mutation disrupted relative to undisrupted is measured by the T7 endonuclease I assay (EnGen® Mutation Detection Kit, New England Biolabs #E3321, Ipswich, MA, United States), restriction endonuclease digestion (New England Biolabs, Ipswich, MA, United States) of PCR-amplified target site, deep gene sequencing using an Illumina MiSeq system and barcoded primers flanking the target site (Illumina, San Diego, CA, United States) or other suitable method. EGFR protein expression and activity in cells of the harvest tissues are measured using an enzyme-linked immunosorbent assay (ELISA) (Sigma, St. Louis, MO, United States) that detects phosphorylated (*i.e.* activated), unphosphorylated and total EGFR protein. Determination of dose-limiting toxicity to enable first-in-human clinical studies is based on the predicted effective dose(s) from the animal model studies discussed above, a range of concentrations (in milligrams per kilogram body weight, for example) of lipid-encapsulated TevCas9 is nebulized and delivered to an appropriate animal model for toxicology studies, such as the cynomolgus monkey or other non-human primate. The general health of the animals is monitored for any treatment-related adverse events, such as changes in behavior, weight, or food consumption; immune responses; changes to cardiovascular health; mortality, etc. Other measures of efficacy may be measured in the studies, including those described above.

Therapeutic Effect

[0101] The novel chimeric nucleases of the instant invention have been intentionally designed to modify the DNA of lung epithelial cells to treat monogenetic diseases although they are capable of working in other cell types or in the cells of other organisms such as bacteria, yeast, insect, plant or other mammals *in vivo* or *ex vivo* to treat monogenetic or polygenetic and infectious diseases.

Example 1: A method of targeted insertion or replacement of all or a portion of a DNA sequence in the genome of human cells

[0102] FIG. 2 illustrates the mechanism of action of cellular uptake of the novel chimeric nuclease of the instant invention. As illustrated in FIG. 2A, a cell 20 or cells 20 are exposed to the novel lipid-encapsulated nuclease particles 21 containing the TevCas9 25 either by *in vivo* or

ex vivo administration. As shown in **FIG. 2B**, the lipid-encapsulated nuclease particle **21** is endocytosed into the cell **20**. The endosome **22** goes through a maturation process in the cytosol and is targeted for degradation (**FIG. 2C**). On certain occasions, the TevCas9 **25** can escape the endosome **22** and enter the cytosol (**FIG. 2D**). In eukaryotic organisms, the nuclease (TevCas9) **25** is targeted to the nucleus **23** of the cell **20** through one or more nuclear-localization sequences (“NLS”). As depicted in **FIG. 2E**, through its nuclear localization sequence, TevCas9 **25** can enter the nucleus **23** and when in the nucleus **23**, the TevCas9 nuclease **25** binds to and cleaves **26** the target genomic DNA **24** sequence.

[0103] **FIG. 3** illustrates the mechanism of the TevCas9 nuclease in cutting DNA. **FIG. 3A** is representation of the key features of TevCas9 bound to its target genomic DNA sequence **24** is shown prior to the cleavage reaction. The I-TevI domain **27** targets the I-TevI Target Sequence **29**. The linker domain **30** joins the I-TevI domain **27** with the Cas9 domain **28** which targets the Cas9 Target Sequence **31**. The gene mutation **32** is surrounded by or in close proximity to the I-TevI Target Sequence **29** and the Cas9 Target Sequence **31**. As shown in **FIG. 3B**, the TevCas9 **25** cleaves the target sequence leaving a deletion product **34** of a predictable size with non-complementary DNA ends **35, 36**. **FIG. 3C** illustrates that in the presence of single-stranded donor DNA with homology arms **37**, the cell **20** can insert the donor DNA **37** sequence near the cut sites through the homology-directed repair (HDR) pathway **38**. **FIG. 3D** illustrates that in the presence of donor DNA **39** with compatible DNA ends to those cleaved by TevCas9 **25**, the cell **20** can insert the donor DNA sequence **39** between the cut sites through directed-ligation using the non-homologous end joining (NHEJ) pathway **40**. In the absence of donor DNA, the cell **20** can join the DNA ends through the NHEJ pathway **40** (**FIG. 3E**).

Example 2: The treatment of cystic fibrosis

[0104] For the treatment of cystic fibrosis, the exogenous donor DNA contains a DNA sequence, which repairs the *CFTR delta F508* mutation involving a method of targeted deletion of a defined length of a DNA sequence in human somatic cells to stimulate homology-directed repair using exogenous donor DNA as a template (**FIG 3C**).

Example 3: The treatment of non-small cell lung cancer

[0105] For the application of treating non-small cell lung cancer, a version of the Cas9 domain which cuts only one strand of DNA (D10A or H557A mutation) or a nuclease deficient version (the D10A + H557A mutations) is used and the sequences targeted are *EGFR* exon 19 deletion

mutations (SEQ ID NOS: 2-4). In this application, however, the nuclease does not contain exogenous donor DNA. In the absence of exogenous donor DNA, the cell can remove the DNA sequence between the two sites targeted by the nuclease by non-homologous end joining (**FIG. 3E**).

[0106] Alternatively, the inhalation route is a fast and effective way of delivering medication locally to the lungs and for the systemic administration of certain agents. Inhalation drug therapy is used extensively to treat respiratory conditions such as asthma and Chronic Obstructive Pulmonary Disease (COPD). Research is ongoing to develop inhalation systems to treat cystic fibrosis.

[0107] The examples which follow are intended in no way to limit the scope of the disclosure but are provided to illustrate how to prepare and use compounds disclosed herein. Many other embodiments of this disclosure will be apparent to one skilled in the art.

[0108] A nebulizer is a device that delivers medication to the lungs in the form of an aerosolized vapor. Nebulizers are commonly used to treat respiratory diseases such as asthma and COPD, for example, the nebulization of corticosteroids, although nebulization has also been used for the treatment and prevention of lung infections, such as ARIKAYCE® (Insmmed Incorporated, Bridgewater, NJ, United States).

[0109] The nebulizer may require some procedure to prepare the liquid for nebulization. The medication is commonly held in liquid form in a cup inside the nebulizer chamber. Once loaded, the device is switched on which generates compressed air to convert the liquid into a vapor in the nebulization chamber. The patient puts the mouthpiece of the nebulization chamber into their mouth and takes a sharp, deep inhalation, holding their breath for 5–10 seconds to ensure the medication reaches the lower parts of the lung. There are a variety of such devices. Many modern nebulizers are breath-actuated and rely on the force of patient inhalation to entrain the aerosolized liquid from the device and thus ensure the medication is delivered only to the patient and not to the surrounding environment. This also ensures consistency of delivering a full dose of the medication to the patient.

[0110] The use of nebulizers is well known and nebulizers are commercially available from several sources, such as Aeroneb®, AeroEclipse®, (Trudell Medical, London, ON, Canada) or PARI-LC Plus®, (PARI USA, Midlothian, VA, United States). In an example of the present invention, a nebulizer is utilized for delivery of the lipid-encapsulated novel chimeric nuclease

comprising a modified I-TevI nuclease domain, a linker, and a modified RNA-guided nuclease *Staphylococcus aureus* Cas9 of the instant application to the lung epithelial tissue. A sterile liquid version of the therapeutic of interest is loaded into the nebulization chamber and is subsequently aerosolized and is inhaled by the patient into the lung via deep breaths.

[0111] Some of the advantages of using a nebulizer versus oral or intravenous administration are: less drug could be required compared to oral or intravenous administration, onset of action can be more rapid *via* inhalation compared to the oral route, adverse effects are potentially less severe due to local delivery of the medication to the lung tissue where the disease manifests itself, inhaled drug therapy is painless and relatively comfortable for the patient which encourages compliance.

[0112] No non-invasive route of delivery provides the speed of action that an inhaled drug can provide. One of the advantages of inhaled drugs is that they are more rapidly absorbed than subcutaneously injected molecules and provides a more immediate physiological response. Small or large molecules, particularly hydrophobic molecules, can be absorbed within seconds after inhalation and can thus be used to treat a wide variety of symptoms that come on suddenly or need long term administration. Pain, panic, anxiety, nausea, cardiovascular crises, bronchoconstriction, sleep induction, spasms, Parkinson's lock-up, and hot flashes are some of the rapid-onset conditions that are addressable with inhaled medicines.

[0113] Most protein-based drug products have some water solubility and are rapidly and efficiently absorbed from the lungs. Those that are more hydrophobic are absorbed even more rapidly within seconds to a few minutes. Those that are more hydrophilic are absorbed within minutes to tens of minutes. In one example of the present invention, one vial is aseptically filled with a therapeutic dose of the lipid nanoparticle which is hydrophobic and another vial is aseptically filled with a therapeutic dose of a chimeric nuclease comprising a modified I-TevI nuclease domain, a linker and a modified RNA-guided nuclease *Staphylococcus aureus* Cas9 which is water soluble and hydrophilic, for delivery to the lungs of a therapeutic dose. The dose can range from 1 to 1000 milligrams of each of the lipid nanoparticle and chimeric nuclease with about 5 to 200 milligrams being preferred. The claimed lipid-encapsulated chimeric nuclease comprising a modified I-TevI nuclease domain, a linker, and a modified RNA-guided nuclease *Staphylococcus aureus* Cas9 can be absorbed in the cells of the lungs within a few hours and complete cleavage on DNA substrates *in vitro* has been observed within 2 hours. Other nebulized

therapies have been delivered daily. Nebulized administration of the lipid-encapsulated chimeric nuclease, therefore, can be daily or more infrequently depending on its efficacy on a per patient basis. The chimeric nucleases are manufactured by BioVectra Corporation (Charlottetown, PE, Canada) and the vials are aseptically filled by Dalton Pharma Services (Mississauga, ON, Canada). The lipid nanoparticles are manufactured and the vials are aseptically filled by Transferra Nanosciences Inc. (Burnaby, BC, Canada).

[0114] The dosage of any disclosed compositions will vary depending on the symptoms, age and body weight of the patient, the nature and severity of the disorder to be treated or prevented, the route of administration, and the form of the subject composition. Any of the subject formulations may be administered in a single dose or in divided doses. Dosages for the compositions may be readily determined by techniques known to those of skill in the art or as taught herein.

[0115] In certain embodiments, the dosage of the subject compounds will generally be in the range of about 1 to 1000 milligrams depending on the body weight of the patient, specifically in the range of about 5 to 200 milligrams.

[0116] An effective dose or amount, and any possible effects on the timing of administration of the formulation, may need to be identified for any particular composition of the disclosure. This may be accomplished by routine experiment as described herein, using one or more groups of animals (preferably at least 5 animals per group), or in human trials if appropriate. The effectiveness of any subject composition and method of treatment or prevention may be assessed by administering the composition and assessing the effect of the administration by measuring one or more applicable indices and comparing the post-treatment values of these indices to the values of the same indices prior to treatment.

[0117] The precise time of administration and amount of any particular subject composition that will yield the most effective treatment in a given patient will depend upon the activity, pharmacokinetics, and bioavailability of a subject composition, physiological condition of the patient (including age, sex, disease type and stage, general physical condition, responsiveness to a given dosage and type of medication), route of administration, and the like. The guidelines presented herein may be used to optimize the treatment, *e.g.*, determining the optimum time and/or amount of administration, which will require no more than routine experimentation consisting of monitoring the subject and adjusting the dosage and/or timing.

[0118] While the subject is being treated, the health of the patient may be monitored by measuring one or more of the relevant indices at predetermined times during the treatment period. Treatment, including composition, amounts, times of administration and formulation, may be optimized according to the results of such monitoring. The patient may be periodically re-evaluated to determine the extent of improvement by measuring the same parameters. Adjustments to the amount(s) of subject composition administered and possibly to the time of administration may be made based on these re-evaluations.

[0119] Treatment may be initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage may be increased by small increments until the optimum therapeutic effect is attained.

[0120] The use of the subject compositions may reduce the required dosage for any individual agent contained in the compositions because the onset and duration of effect of the different agents may be complimentary.

[0121] Therapeutic efficacy of subject compositions may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD₅₀ and the ED₅₀.

[0122] The data obtained from the cell culture assays and animal studies may be used in formulating a range of dosage for use in humans. The dosage of any subject composition lies preferably within a range of concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For compositions of the disclosure, the therapeutically effective dose may be estimated initially from cell culture assays.

Formulations

[0123] Pharmaceutical compositions of the disclosure may be administered by various means, depending on their intended use, as is well known in the art. For example, compositions of the disclosure are to be administered through nebulization. Alternatively, formulations disclosed herein may be administered intravenously, subcutaneously, or intramuscularly. These formulations may be prepared by conventional means, and, if desired, the compositions may be mixed with any conventional additive, such as an excipient, a solubilizing agent, a suspension aid, an emulsifying agent, or preservative agent. The disclosed excipients may serve more than one function. For example, a solubilizing agent may also be a suspension aid, an emulsifier, a

preservative, and the like.

[0124] The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of a composition that may be combined with a carrier material to produce a single dose vary depending upon the subject being treated, and the particular mode of administration.

[0125] Methods of preparing these formulations include the step of bringing into association compositions of the disclosure with the carrier and, optionally, one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association agents with liquid carriers.

[0126] It will be appreciated that a disclosed composition may include lyophilized or freeze-dried compounds disclosed herein. For example, disclosed herein are compositions that disclosed compounds crystalline and/or amorphous powder forms. Such forms may be reconstituted for use as *e.g.*, an aqueous composition.

[0127] Liquid dosage forms for injection include pharmaceutically acceptable solutions, emulsions, microemulsions, solutions and suspensions. In addition to the subject composition, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, glycerol, tetrahydrofuryl alcohol, and fatty acid esters of sorbitan, cyclodextrins, albumin, hyaluronic acid, chitosan and mixtures thereof. Polyethylene glycol (PEG) may be used to obtain desirable properties of solubility, stability, half-life, and other pharmaceutically advantageous properties. Representative examples of stabilizing components include polysorbate 80, L-arginine, polyvinylpyrrolidone, trehalose, and combinations thereof. Other excipients that may be employed, such as solution binders or anti-oxidants include, but are not limited to, butylated hydroxytoluene (BHT), calcium carbonate, calcium phosphate (dibasic), calcium stearate, croscarmellose, crosslinked polyvinyl pyrrolidone, citric acid, crospovidone, cysteine, ethylcellulose, gelatin, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, maltitol, mannitol, methionine, methylcellulose, methyl paraben, microcrystalline cellulose, polyvinyl pyrrolidone, povidone, pregelatinized starch, propyl paraben, retinyl palmitate, shellac, silicon dioxide, sodium carboxymethyl cellulose, sodium citrate, sodium starch glycolate, sorbitol, starch (corn), stearic acid, sucrose, talc, titanium dioxide, vitamin A, vitamin E (alpha-tocopherol), vitamin C and xylitol.

[0128] Ordinarily, an aqueous aerosol is made by formulating an aqueous solution or suspension of a subject composition together with conventional pharmaceutically acceptable carriers and stabilizers. The carriers and stabilizers vary with the requirements of the particular subject composition, but typically include non-ionic surfactants (Tweens, pluronics, or polyethylene glycol), innocuous proteins like serum albumin, sorbitan esters, oleic acid, lecithin, amino acids such as glycine, buffers, salts, sugars or sugar alcohols. Aerosols generally are prepared from isotonic solutions.

[0129] It should be noted that excipients given as examples may have more than one function. For example, a solubilizing agent may also be a suspension aid, an emulsifier, a preservative and the like.

[0130] Examples of suitable aqueous and non-aqueous carriers which may be employed in the pharmaceutical compositions of the disclosure include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate and cyclodextrins. Proper fluidity may be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

[0131] The compositions of the present invention typically are ready to administer, aqueous solutions which are sterile, storage-stable, and pharmaceutically acceptable without the need for reconstitution prior to administration. The compositions of the present invention are suitable for administration to a subject which means that they are pharmaceutically acceptable, non-toxic, do not contain any components which would adversely affect the biological effects of the chimeric nuclease, and have the pH close to that of the physiological condition which avoids inhalation and/or injection site reactions. The compositions of the present invention do not, for example, comprise any cells.

[0132] The compositions are typically stored in a sealed container, vial or cartridge which is typically suitable for long term storage. "Suitable for long-term storage" means that the vial, container or cartridge does not allow for the escape of components of the compositions of the present invention or the ingress of external components, such as, microorganisms when kept for at least 3 months at 25°C.

[0133] The compositions of the present invention are preferably administered by nebulization,

typically breath-actuated nebulization.

[0134] The compositions of the present invention can also be administered by injection as described herein.

[0135] The compositions of the present invention may be administered alone or in combination with an additional therapeutic agent, such as an anti-viral, an anti-microbial, a chemotherapeutic and an immunotherapy.

[0136] Vials, as used herein, can also comprise two containers one of which contains the chimeric nuclease or lipid particle, as described herein, in a lyophilized powder, as described below, and the second container contains a liquid for reconstitution of the lyophilized powder. The contents of the two containers can be mixed prior to administration.

[0137] As discussed above the compositions of the present invention can be administered by nebulization. Suitable volumes of the compositions of the present invention for nebulization include about 0.5 to about 1 ml, about 1 to about 2 ml, about 2 to about 10 ml, or about 10 to about 20 ml.

[0138] In the compositions of the present invention the concentration of the chimeric nuclease is from about 0.1 mg/ml to about 10.0 mg/ml, from about 10.0 mg/ml to about 100.0 mg/ml, from about 30.0 mg/ml to about 300.0 mg/ml, from about 500 mg/ml to about 2000 mg/ml and about 2.0 mg/ml.

[0139] In the compositions of the present invention the concentration of the lipid nanoparticle is from about 0.1 mg/ml to about 10.0 mg/ml, from about 10.0 mg/ml to about 100.0 mg/ml, from about 30.0 mg/ml to about 300.0 mg/ml, from about 500 mg/ml to about 2000 mg/ml and about 2.0 mg/ml.

SEQ ID NO 1

ATGCAGAGGTGCGCTCTGGAAAAGGCCAGCGTTGTCTCCAACTTTTTTTCAGCTGGACCAGACCAATTT
TGAGGAAAGGATACAGACAGCGCTGGAATTGTCAGACATATACCAAATCCCTTCTGTGATTCTGCTGA
CAATCTATCTGAAAAATTGGAAAGAGAATGGGATAGAGAGCTGGCTTCAAAGAAAAATCCTAACTCATT
AATGCCCTTCGGCGATGTTTTTCTGGAGATTTATGTTCTATGGAATCTTTTTATATTTAGGGGAAGTCA
CCAAAGCAGTACAGCCTCTCTTACTGGGAAGAATCATAGCTTCTATGACCCGGATAACAAGGAGGAACG
CTCTATCGCGATTTATCTAGGCATAGGCTTATGCCTTCTCTTTATTGTGAGGACACTGCTCCTACACCCA
GCCATTTTGGCCTTCATCACATTGGAATGCAGATGAGAATAGCTATGTTAGTTTGATTTATAGAAGA
CTTTAAAGCTGTCAAGCCGTGTTCTAGATAAAAATAGTATTGGACAACCTGTTAGTCTCCTTTCCAACAA
CCTGAACAAATTTGATGAAGGACTTGCATTGGCACATTTCTGTGGATCGCTCCTTTGCAAGTGGCACTC
CTCATGGGGCTAATCTGGGAGTTGTTACAGGCGTCTGCCTTCTGTGGACTTGGTTTCTGATAGTCCCTTG
CCCTTTTTCAGGCTGGGCTAGGGAGAATGATGATGAAGTACAGAGATCAGAGAGCTGGGAAGATCAGTGA
AAGACTTGTGATTACCTCAGAAATGATTGAAAATATCCAATCTGTTAAGGCATACTGCTGGGAAGAAGCA
ATGGAAAAATGATTGAAACTTAAGACAAACAGAACTGAACTGACTCGGAAGGCAGCCTATGTGAGAT
ACTTCAATAGCTCAGCCTTCTTCTCTCAGGGTCTTTGTGGTGTTTTTATCTGTGCTTCCCTATGCACT
AATCAAAGGAATCATCCTCCGAAAAATATTCACCACCATCTCATTCTGCATTGTTCTGCGCATGGCGGTC
ACTCGGCAATTTCCCTGGGCTGTACAAACATGGTATGACTCTCTTGGAGCAATAAACAAAATACAGGATT
TCTTACAAAAGCAAGAATATAAGACATTGGAATATAACTTAACGACTACAGAAGTAGTGATGGAGAATGT
AACAGCCTTCTGGGAGGAGGATTTGGGGAATTATTTGAGAAAAGCAAAACAAAACAATAACAATAGAAAA
ACTTCTAATGGTGATGACAGCCTCTTCTTCAGTAATTTCTCACTTCTTGGTACTCCTGTCTGAAAGATA
TTAATTTCAAGATAGAAAGAGGACAGTTGTTGGCGGTTGCTGGATCCACTGGAGCAGGCAAGACTTCACT
TCTAATGGTGATTATGGGAGAACTGGAGCCTTCAGAGGGTAAATTAAGCACAGTGGGAAGAATTCATTCT
TGTTCTCAGTTTTCTGGATTATGCCTGGCACCATTAAAGAAAAATATCAT---TGGTGTTCCTATGATG
AAATATAGATACAGAAGCGTCTATCAAAGCATGCCAACtagaagAGGACATCTCCAAGTTTGAGAGAAAAGA
CAATATAGTTCTTGGAGAAGTGGAATCACACTGAGTGGAGGTCAACGAGCAAGAATTTCTTTAGCAAGA
GCAGTATACAAAGATGCTGATTGTATTATTAGACTCTCCTTTTGATACCTAGATGTTTTAACAGAAA
AAGAAATATTTGAAGCTGTGTCTGTAACCTGATGGCTAACAAACTAGGATTTTGGTCACTTCTAAAAAT
GGAACATTTAAAGAAAGCTGACAAAATATTAATTTTGCATGAAGGTAGCAGCTATTTTATGGGACATTT
TCAGAACTCCAAAATCTACAGCCAGACTTTAGCTCAAACTCATGGGATGTGATTCTTTCGACCAATTTA
GTGCAGAAAGAAGAAATTCATCCTAAGTACAGCCTTACACCGTTTCTCATTAGAAGGAGATGCTCCTGT
CTCCTGGACAGAAACAAAAAACAATCTTTTAAACAGACTGGAGAGTTTGGGAAAAAAGGAAGATTTCT
ATTCTCAATCCAATCAACTCTATACGAAAATTTTCCATTGTGCAAAAGACTCCCTTACAAATGAATGGCA
TCGAAGAGGATTCTGATGAGCCTTTAGAGAGAAGGCTGTCTTAGTACCAGATTCTGAGCAGGGAGAGGC
GATACCTGCTCGCATCAGCGTGATCAGCACTGGCCCCACGCTTCAGGCACGAAGGAGGAGTCTGTCTG
AACCTGATGACACACTCAGTTAACCAAGGTGAGAACATTACCGAAAGACAACAGCATCCACACGAAAAG
TGTCCTGGCCCCCTCAGGCAAACTTGACTGAACTGGATATATATTCAAGAAGGTTATCTCAAGAACTGG
CTTGGAATAAGTGAAGAAATTAACGAAGAAGACTTAAAGGAGTGCTTTTTTGTGATGATATGGAGAGCATA
CCAGCAGTGACTACATGGAACACATACCTTCGATATATTACTGTCCACAAGAGCTTAATTTTTGTGCTAA
TTTGGTGCTTAGTAATTTTTCTGGCAGAGGTGGCTGCTTCTTTGGTTGTGCTGTGGCTCCTTGGAACAC
TCCTCTTCAAGACAAAGGGAATAGTACTCATAGTAGAAAATAACAGCTATGCAGTGATTATCACCAGCACC
AGTTCGTATTATGTGTTTTACATTTACGTGGGAGTAGCCGACACTTTGCTTGCTATGGGATTCTTCAGAG
GTCTACCACTGGTGCATACTCTAATCACAGTGTGCAAAATTTTACACCACAAAATGTTACATTTCTGTTCT
TCAAGCACCTATGTCAACCCTCAACACGTTGAAAGCAGGTGGGATTCTTAATAGATTCTCCAAGATATA
GCAATTTTGGATGACCTTCTGCCTCTTACCATATTTGACTTCATCCAGTTGTTATTAATTGTGATTGGAG
CTATAGCAGTTGTGCGAGTTTACAACCTACATCTTGTGCAACAGTGCCAGTGATAGTGGCTTTTAT
TATGTTGAGAGCATATTTCTCCAAACCTCACAGCAACTCAAACAACCTGGAATCTGAAGGCAGGAGTCCA
ATTTTCACTCATCTTGTTACAAGCTTAAAGGACTATGGACACTTCGTGCCTTCGGACGGCAGCCTTACT
TTGAACTCTGTTCCACAAAGCTCTGAATTTACATACTGCCAACTGGTCTTGTACCTGTCAACACTGCG
CTGGTTCCAAATGAGAATAGAAATGATTTTGTGCTCTTCTTCAATTGCTGTTACCTTCATTTCCATTTTA
ACAACAGGAGAAGGAGAAGGAGATTGGTATTATCTCCTGACTTTAGCCATGAATATCATGAGTACATTGC
AGTGGGCTGTAACTCCAGCATAGATGTGGATAGCTTGATGCGATCTGTGAGCCGAGTCTTTAAGTTCAT

TGACATGCCAACAGAAGGTAAACCTACCAAGTCAACCAACCATAACAAGAATGGCCAACTCTCGAAAGTT
 ATGATTATTGAGAATTACACGTGAAGAAAGATGACATCTGGCCCTCAGGGGGCCAAATGACTGTCAAAG
 ATCTCACAGCAAAATACACAGAAGGTGGAAATGCCATATTAGAGAACATTTCTTCTCAATAAGTCCTGG
 CCAGAGGGTGGGCCTCTTGGGAAGAACTGGATCAGGGAAGAGTACTTTGTTATCAGCTTTTTTGAGACTA
 CTGAACACTGAAGGAGAAATCCAGATCGATGGTGTCTTGGGATTCAATAACTTTGCAACAGTGGAGGA
 AAGCCTTTGGAGTGATACCACAGAAAGTATTTATTTTCTGGAACATTTAGAAAAAAGTGGATCCCTA
 TGAACAGTGGAGTGATCAAGAAATATGGAAAGTTGCAGATGAGGTTGGGCTCAGATCTGTGATAGAACAG
 TTTCTTGGGAAGCTTGACTTTGTCTTGTGGATGGGGCTGTGTCTTAAGCCATGGCCACAAGCAGTTGA
 TGTGCTTGGCTAGATCTGTTCTCAGTAAGGCGAAGATCTTGCTGCTTGATGAACCCAGTGCTCATTTGGA
 TCCAGTAACATAACCAATAATTAGAAGAACTCTAAACAAGCATTTGCTGATTGCACAGTAATTTCTGT
 GAACACAGGATAGAAGCAATGCTGGAATGCCAACAATTTTGGTCATAGAAGAGAACAAAGTGCGGCAGT
 ACGATTCCATCCAGAACTGCTGAACGAGAGGAGCCTCTTCCGGCAAGCCATCAGCCCTCCGACAGGGT
 GAAGCTCTTTCCCAACCGAACTCAAGCAAGTGCAAGTCTAAGCCCCAGATTGCTGCTCTGAAAGAGGAG
 ACAGAAGAAGAGGTGCAAGATACAAGGCTTTAG

LENGTH: 4,443

TYPE: DNA

ORGANISM: HOMO SAPIENS

FEATURE: CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR (CFTR) DELTA F508 DNA CODING SEQUENCE

OTHER INFORMATION: Nucleotides deleted compared to wild type CFTR are shown as ("–"). Sequence targeted by TevCas9 highlighted. I-TevI site is underlined, Cas9 guide RNA sequence is double underlined and PAM sequence is lowercase.

SEQ ID NO 2

*GTCTTCCTTCTCTCTGTCTATAG*GGACTCTGGatcccaGAAGGTGAGAAAGTTAAAATTCCCGTCGCTA
 TCAA-----
 AACATCTCCGAAAGCCAACAAGGAAATCCTCGAT*GTGAGTTTCTGCTTTGCTGTGTGG*

LENGTH: 132

TYPE: DNA

ORGANISM: HOMO SAPIENS

FEATURE: EPIDERMAL GROWTH FACTOR RECEPTOR (EGFR) EXON 19 C.2235-2249 DELETION SEQUENCE

OTHER INFORMATION: Nucleotides deleted compared to wild type EGFR are shown as ("–"). Sequence targeted by TevCas9 highlighted. I-TevI site is underlined, Cas9 guide RNA sequence is double underlined and PAM sequence is lowercase. Intron sequences are bold and italicized.

SEQ ID NO 3

*GTCTTCCTTCTCTCTGTCTATAG*GGACTCTGGatcccaGAAGGTGAGAAAGTTAAAATTCCCGTCGCTA
 TCAAG-----
 ACATCTCCGAAAGCCAACAAGGAAATCCTCGAT*GTGAGTTTCTGCTTTGCTGTGTGG*

LENGTH: 132

TYPE: DNA

ORGANISM: HOMO SAPIENS

FEATURE: EPIDERMAL GROWTH FACTOR RECEPTOR (EGFR) EXON 19 C.2236_2250 DELETION SEQUENCE

OTHER INFORMATION: Nucleotides deleted compared to wild type EGFR are shown as ("–"). Sequence targeted by TevCas9 highlighted. I-TevI site is underlined, Cas9 guide RNA sequence is double underlined and PAM sequence is lowercase. Intron sequences are bold and italicized.

SEQ ID NO 4

*GTCTTCCTTCTCTCTGTCATAG*GGACTCTGGatcccaGAAGGTGAGAAAGTTAAATTCCCGTCGCTA
TCAAG-----
CATTCCGAAAGCCAACAAGGAAATCCTCGAT*GTGAGTTTCTGCTTTGCTGTGTGG*

LENGTH: 132

TYPE: DNA

ORGANISM: HOMO SAPIENS

FEATURE: EPIDERMAL GROWTH FACTOR RECEPTOR (EGFR) EXON 19 C.2237_2251 DELETION SEQUENCE

OTHER INFORMATION: Nucleotides deleted compared to wild type EGFR are shown as ("–"). Sequence targeted by TevCas9 highlighted. I-TevI site is underlined, Cas9 guide RNA sequence is double underlined and PAM sequence is lowercase. Intron sequences are bold and italicized.

SEQ ID NO 5

*GTCTTCCTTCTCTCTGTCATAG*GGACTCTGGatcccaGAAGGTGAGAAAGTTAAATTCCCGTCGCTATCAAGGAA
TTAAGAGAAGCAACATTCCGAAAGCCAACAAGGAAATCCTCGAT*GTGAGTTTCTGCTTTGCTGTGTGG*

LENGTH: 147

TYPE: DNA

ORGANISM: HOMO SAPIENS

FEATURE: EPIDERMAL GROWTH FACTOR RECEPTOR (EGFR) EXON 19 WILD TYPE SEQUENCE

OTHER INFORMATION: Sequence targeted by TevCas9 highlighted. I-TevI site is underlined, Cas9 guide RNA sequence is double underlined and PAM sequence is lowercase. Note: the I-TevI and Cas9 target sites are not appropriately spaced, therefore, I-TevI should not bind nor cleave its target site. Intron sequences are bold and italicized.

SEQ ID NO 6

MGKSGIYQIKNTLNNKVYVGS AKDFEKRWRHFKDLEKGCHSSIKLQRSFNKHG NVFECSILEEIPYEKD
LIIERENFWIKELNSKINGYNIA

LENGTH: 93

TYPE: AMINO ACID

ORGANISM: ENTEROBACTERIA PHAGE T4

FEATURE: I-TEVI DOMAIN

OTHER INFORMATION:

SEQ ID NO 7

DATFGDTCSTHPLKEEIIKKRSETFKAKMLKLGPDGRKALYSKPGSKNGRWNPE**TH**KFC**K**CGVRIQTSAY
TCSKCRNGGSGGS

LENGTH: 83

TYPE: AMINO ACID

ORGANISM: ARTIFICIAL

FEATURE: LINKER DOMAIN

OTHER INFORMATION: V117F; mutation underlined

SEQ ID NO 8

DATFGDTCSTHPLKEEIIKKRSETVKAKMLKLGPDGRKALYSRPGSKSGRWNPE**TH**KFC**K**CGVRIQTSAY
TCSKCRNGGSGGS

LENGTH: 83

TYPE: AMINO ACID

ORGANISM: ARTIFICIAL

FEATURE: LINKER DOMAIN

OTHER INFORMATION: K135R/N140S; mutations underlined

SEQ ID NO 9

DATFGDTCSTHPLKEEIIKKRSETFKAKMLKLGPDGRKALYSRPGSKSGRWNPE**TH**KFC**K**CGVRIQTSAY
TCSKCRNGGSGGS

LENGTH: 83

TYPE: AMINO ACID

ORGANISM: ARTIFICIAL

FEATURE: LINKER DOMAIN

OTHER INFORMATION: V117F/K135R/N140S; mutations underlined

SEQ ID NO 10

DATFGDTCSTHPLKEEIIKKRSETVKAKMLKLGPDGRKALYSRPGSKSGRWNPE**TH**KFC**K**CGVRIQTSAY
TCSKCRNGGSGGTGGS

LENGTH: 86

TYPE: AMINO ACID

ORGANISM: ARTIFICIAL

FEATURE: LINKER DOMAIN VARIANT

OTHER INFORMATION:

SEQ ID NO 11

DATFGDTCSTHPLKEEIIKKRSETVKAKMLKLGPDGRKALYSRPGSKSGRWNPETHKFCKCGVRIQTSAY
TCSKCRNNGGGSGGGGS

LENGTH: 87

TYPE: AMINO ACID

ORGANISM: ARTIFICIAL

FEATURE: LINKER DOMAIN VARIANT

OTHER INFORMATION:

SEQ ID NO 12

DATFGDTCSTHPLKEEIIKKRSETVKAKMLKLGPDGRKALYSRPGSKSGRWNPETHKFCKCGVRIQTSAY
TCSKCRNKESGSVSSEQLAQFRSLD

LENGTH: 95

TYPE: AMINO ACID

ORGANISM: ARTIFICIAL

FEATURE: LINKER DOMAIN VARIANT

OTHER INFORMATION:

SEQ ID NO 13

MKRNYYILGLDIGITSVGYGIIDYETRDVIDAGVRLFKEANVENNEGRRSKRGARRLKRRR
RHRIQRVKKLLFDYNLLTDHSELSGINPYEARVKGLSQKLSEEEFSAALLHLAKRRGVHN
VNEVEEDTGNELSTKEQISRNSKALEEKYVAELQLERLKKDGEVRGSINRFKTSYVKEA
KQLLKVQKAYHQLDQSFIDTYIDLLETRRTYYEGPGEGSPFGWKDIKEWYEMLMGHCTYF
PEELRSVKYAYNADLYNALNDLNNLVITRDENEKLEYEYKFQIIENVFKQKKKPTLKQIA
KEILVNEEDIKGYRVTSTGKPEFTNLKVYHDIKDITARKEIENAELLDQIAKILTIYQS
SEDIQEELTNLNSLTQEIEQISNLKGYTGTHNLSLKAINLILDELWHTNDNQIAIFNR
LKLVPKKVDLSQQKEIPTTLVDDFILSPVVKRSFIQSIKVINAIKKYGLPNDIIIELAR
EKNSKDAQKMINEMQKRNRQTNERIEEIIIRTGKENAKYLIEKIKLHDMQEGKCLYSLEA
IPLEDLLNNPFNYEVDHIIIPRSVSFDNSFNNKVLVKQEENSCKGNRTPFQYLSSSDSKIS
YETFKKHILNLAKGGRISKTKEYLLEERDINRFSVQKDFINRNLVDTRYATRGLMNNLL
RSYFRVNNLDVKVKSINGGETSFLRRKWKFCKERNKGYKHAEDALIIANADFIKWKWK
LDKAKKVMENQMFEEKQAESMPEIETEQEYKEIFITPHQIKHIKDFKDYKYSHRVDKKN
RELINDTLYSTRKDDKGNLTIVNNLNGLYDKDNDKLLKLINKSPEKLLMYHHDPPQTYQKL
KLIMEQYGDEKNPLYKYEEETGNYLTKYSKKNNGPVIKKIKYYGNKLNALHLDITDDYPNS
RNKVVKLSLKPYPYFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEEAKLKKISNQA
EFIASFYNNDLIKINGELYRVIGVNNDLLNRIEVMIDITYREYLENMNDKRPPRIKTI
ASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG

LENGTH: 1,053**TYPE:** AMINO ACID**ORGANISM:** STAPHYLOCOCCUS AUREUS**FEATURE:****OTHER INFORMATION:****SEQ ID NO 14**

MKRN^YILGLEIGITSVGYGIIDYETRDVIDAGVRLFKEANVENNEGRRSKRGARRLKRRR
RHRIQRVKKLLFDYNLLTDHSELSGINPYEARVKGLSQKLSEEEFSAALLHLAKRRGVHN
VNEVEEDTGNELSTKEQISRN^SSKALEEKYVAELQLERLKKDGEV^RGSINRFKTS^DYVKEA
KQLLKVQKAYHQLDQSFIDTYIDLLETRRTYYEGPGE^GSPFGWKDIKEWYEMLMGHCTYF
PEELRSVKYAYNADLYNALNDLNNLVITRDENEKLEY^EEKFQIIENVFKQKKKPTLKQIA
KEILVNEEDIKGYRV^TSTGKPEFTNLKVYHDIKDITARKEI^IENAE^LLDQIAKILTIYQS
SEDIQEELTNLNS^EL^TQEEIEQISNLKGYTGTHNLSLKAINLILDELWHTNDNQIAIFNR
LKLVPKKVDLSQQKEIPTTLVDDFILSPVVKRSFIQSIKVINAIIKKYGLPNDIIIELAR
EKNSKDAQMINEMQKRNRQTNERIEEII^RTTGKENAKYLIEKIKLHDMQEGKCLYSLEA
IPLEDLLNNPFNYEVDHIIPRSVSFDNSFNNKVLVKQEENS^KKNRTPFQYLSSSDSKIS
YETFKKHILNLAKGGRISKTKEYLLEERDINRFSVQKDFINRNLVDTRYATRGLM^NLL
RSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGYKHAEDALI^IANADFI^FKEWKK
LDKAKVMENQMFE^EKQAESMPEIETE^QEYKEIFITPHQIKHIKDFKDYKYS^HRV^DKKPN
RELINDTLYSTRKDDKNTLIVNNLNGLYDKDNDKLKLINKSPEKLLMYH^HDPQTYQKL
KLIMEQYGDEKNPLYKYEETGNYLTKY^SSKDNGPVIKKIKYYGNKLN^AHL^DITDDY^PNS
RNKVVKL^SLKPYRFDVYLDNGVYKFVTVKNLDVIKENY^EEVNSKCYEEAKKLKKIS^NQA
EFIASFYNNDLIKINGELYRVIGVNN^DLLNRIE^VNMIDITYREYLENMNDKRP^PRI^IKTI
ASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG

LENGTH: 1,053**TYPE:** AMINO ACID**ORGANISM:** ARTIFICIAL**FEATURE:** SACAS9 D10E MUTANT**OTHER INFORMATION:** Mutation underlined**SEQ ID NO 15**

GCGUCAUCAAGCAUGCCAACGUUUUAGUACUCUGGAAACAGAAUCUACUAAAACAAGGCAAAAUGCCGU
GUUUUAUCUCGUCAACUUGUUGCGAGAUUUU

LENGTH: 101**TYPE:** RNA**ORGANISM:** ARTIFICIAL**FEATURE:** TARGETS CFTR GENE

OTHER INFORMATION:**SEQ ID NO 16**

AAUUUUAAUUUCACCUUCGUUUUAGUACUCUGGAAACAGAAUCUACUAAAACAAGGCAAAAUGCCGU
GUUUUUCUCGUCAACUUGUUGGCGAGAUUUU

LENGTH: 101**TYPE:** RNA**ORGANISM:** ARTIFICIAL**FEATURE:** TARGETS EGFR GENE**OTHER INFORMATION:****SEQ ID NO 17**

MKRNYILGLEIGITSVGYGIIDYETRDVIDAGVRLFKEANVENNEGRRSKRGARRLKRRR
RHRIQRVKKLLFDYNLLTDHSELSGINPYEARVKGLSQKLSEEEFSAALLHLAKRRGVHN
VNEVEEDTGNELSTKEQISRNSKALEEKYVAELQLERLKKDGEVRGSINRFKTSYVKEA
KQLLKVQKAYHQLDQSFIDTYIDLLETRRTYYEGPGEPSFGWKDIKEWYEMLMGHCTYF
PEELRSVKYAYNADLYNALNDLNNLVITRDENEKLEYEKFQIIENVFKQKKKPTLKQIA
KEILVNEEDIKGYRVSTGKPEFTNLKVYHDIKDITARKEIENAEELLDQIAKILTIYQS
SEDIQEELTNLSELQEEIEQISNLKGYTGTHNLSLKAINLILDELWHTNDNQIAIFNR
LKLVPKKVDLSQQKEIPTTLVDDFILSPVVKRSFIQSIKVINAIKKYGLPNDIIIEELAR
EKNSKDAQKMINEMQKRNRQTNERIEEIIRTTGKENAKYLIEKIKLHDMQEGKCLYSLEA
IPLEDLLNNPFNYEVD^{AI}IPRSVSFDNSFNNKVLVKQEENSKKGNRTPFQYLSSSDSKIS
YETFKKHILNLAKGGRISKTKEYLLEERDINRFSVQKDFINRNLVDTRYATRGLMNL
RSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGYKHAEDALI^{IANADFI}FKEWKK
LDKAKKVMENQMFEKQAESMPEIETE^QEYKEIFITPHQIKHIKDFKDYKYSHRVDKPN
RELINDTLYSTRKDDKNTLIVNNLNGLYDKDNDKLKLINKSPEKLLMYHHD^PQTYQKL
KLIMEQYGDEKNPLYKYEETGNYLTKYSKKDNGPVIKKIKYYGNKLNALHDITDDY^{PNS}
RNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKENYYEVNSKCYEEAKKLKKISNQA
EFIASFYNNDLIKINGELYRVIGVNNDLLNRIEVMIDITYREYLENMNDKRP^{PRI}IKTI
ASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG

LENGTH: 1,053**TYPE:** AMINO ACID**ORGANISM:** ARTIFICIAL**FEATURE:** SACAS9 D10E + H557A MUTANT**OTHER INFORMATION:** Mutation underlined**SEQ ID NO 18**

MKRNYILGLAIGITSVGYGIIDYETRDVIDAGVRLFKEANVENNEGRRSKRGARRLKRRR
RHRIQRVKKLLFDYNLLTDHSELSGINPYEARVKGLSQKLSEEEFSAALLHLAKRRGVHN
VNEVEEDTGNELSTKEQISRNSKALEEKYVAELQLERLKKDGEVRGSINRFKTSYVKEA

KQLLKVQKAYHQLDQSFIDTYIDLLETRRTYYEGPGEGSPFGWKDIKEWYEMLMGHCTYF
 PEELRSVKYAYNADLYNALNDLNNLVITRDENEKLEYEYKFQIIENVFKQKKKPTLKQIA
 KEILVNEEDIKGYRVSTGKPEFTNLKVYHDIKDITARKEIIEAELLDDQIAKILTIYQS
 SEDIQEELTNLSELQTQEEIEQISNLKGYTGTHNLSLKAINLILDELWHTNDNQIAIFNR
 LKLVPPKKVDLSQQKEIPTTLVDDFILSPVVKRSFIQSIKVINAIKKYGLPNDIIIELAR
 EKNSKDAQKMINEMQKRNRQTNERIEEIIIRTGKENAKYLIEKIKLHDMQEGKCLYSLEA
 IPLEDLLNNPFNYEVDAIIPRSVSFDNSFNNKVLVKQEENSKKGNRTPFQYLSSSDSKIS
 YETFKKHILNLAKGGRISKTKEYLLEERDINRFSVQKDFINRNLVDTRYATRGLMNL
 RSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGYKHAEDALIIANADFIKWEKK
 LDKAKKVMENQMFEKQAESMPEIETEQEYKEIFTTPHQIKHDKFDKYKYSHRVDKPN
 RELINDTLYSTRKDDKNTLIVNNLNGLYDKDNDKLKLINKSPEKLLMYHHDPTQYQKL
 KLIMEQYGDENPLYKYEEETGNYLTKYSSKDNPGVKKIKYYGNKLNALHLDITDDYPNS
 RNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLVDIKKENYYEVNSKCYEEAKLKKISNQA
 EFIAFYNNDLIKINGELYRVIGVNNDLLNRIEVMIDITYREYLENMNDKRPPRIKTI
 ASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG

LENGTH: 1,053

TYPE: AMINO ACID

ORGANISM: ARTIFICIAL

FEATURE: SACAS9 D10A + H557A MUTANT

OTHER INFORMATION: Mutation underlined

SEQ ID NO 19

MDKKYSIGLEIGTNSVGWAVITDEYKVPSKKFKVLGNTDRHSIKKNLIGALLFDSGETAE
 ATRLKRTARRRYTRRKNRICYLQEIFSNEMAKVDDSFHRLSEESFLVEEDKKHERHPIFG
 NIVDEVAYHEKYPTIYHLRKKLVDSSTDKADLRILIYLAHAMIKFRGHFLIEGDLNPDNSD
 VDKLFIQLVQTYNQLFEENPINASGVDAKAILSARLSKSRLENLIAQLPGEKKNGLFGN
 LIALSLGLTPNFKSNFDLAEDAKLQLSKDTYDDDLNLLAQIGDQYADFLAAKNLSDAI
 LLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLKALVRQQLPEKYKEIFFDQSKNGYA
 GYIDGGASQEEFYKFIKPILEKMDGTEELLVKNREDLLRKQRTFDNGSIPHQIHLGELH
 AILRRQEDFYPFLLKDNREKIEKILTFRIPYYVGPLARGNSRFAMWTRKSEETITPWNFEE
 VVDKGASAQSFIERTMTNFDKNLPNEKVLPHKSHLLYEYFTVYNELTKVKYVTEGMRKPAFL
 SGEQKKAIVDLLFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGYHDLKI
 IKDKDFLDNEENEDILEDIVLTTLTFEDREMIEERLKYAHLFDDKVMKQLKRRRYTGWG
 RLSRKLINGIRDKQSGKTILDFLKSDGFANRNFMLIHDDSLTFKEDIQKAQVSGQGD
 SLHEHIANLAGSPAIIKKGILQTVKVVDLVKVMGRHKPENIVIEARENQTTQKGQKNSRER
 MKRIEEGIKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQELDINRLSDYD
 VDHIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKMKNYWRQLLNAKLITQRKFDNL
 TKAERGGSELKAGFIKQQLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKS
 KLVSDFRKDFQFYKVREINNYHHAHDAYLNAVVGITALIKKYPKLESEFVYGDYKVYDVRK
 MIAKSEQEIGKATAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDF
 ATVRKVLSPQVNIIVKTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFESPTVA
 YSVLVVAKVEKGSKKLKSVKELLGITIMERSSEKPNIDFLEAKGYKEVKKDIIKLPK
 YSLFELENKRRLASAGELQKGNELALPSKYVNFLYLASHYEKLKGSPEQKQLFVE
 QHKHYLDEIIIEQISEFSKRVLADANLDKVL SAYNKHDKPIREQAENIIHLFTLTNLGA
 PAAFKYFDTTIDRKQYRSTKEVLDATLIHQSIITGLYETRIDLSQLGGD

LENGTH: 1,368

TYPE: AMINO ACID

ORGANISM: ARTIFICIAL

FEATURE: SPCAS9 D10E, D1135E, R1335Q, T1337R MUTANT

OTHER INFORMATION: mutations underlined

SEQ ID NO 20

MDKKYSIGLEIGTNSVGWAVITDEYKVPSKKFKVLGNTDRHSIKKNLIGALLFDSGETAE
 ATRLKRTARRRYTRRRKNRICYLQEIFSNEMAKVDDSFHRLSEESFLVEEDKKHERHPIFG
 NIVDEVAYHEKYPTIYHLRKKLV DSTDKADLRLLIYLALAHMIKFRGHFLIEGDLNPDNSD
 VDKLFIQLVQTYNQLFEEFNINASGVDAKILSARLSKSRLENLIAQLPGEKKNGLFGN
 LIALSLGLTPNFKSNFDLAEDAKLQLSKD TYDDDLNLLAQIGDQYADLFLAAKNLSDAI
 LLSDILRVNTEITKAPLSASMIKKRYDEHHQDLTLLKALVRQQLPEKYKEIFFDQSKNGYA
 GYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNREDLLRKQRTFDNGSIPHQIHLGELH
 AILRRQEDFYFPFLKDNREKIEKILTFRIPYYVGPLARGNSRFAMWTRKSEETITPWNFEE
 VVDKGASAQSFIERMTNFDKNLPNEKVLPHKSLLEYFTVYNELTKVKYVTEGMRKPAFL
 SGEQKKAIVDLLFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLKI
 IKDKDFLDNEENEDILEDIVLTTLTLFEDREMIEERLKYAHLFDDKVMKQLKRRRYTGWG
 RLSRKLINGIRDKQSGKTILDFLKS DGFANRNFQMQLIHDDSLTFKEDIQKAQVSGQGDSL
 HEHIANLAGSPAIIKKGILQTVKVVDELVKVMGRHKPENIVIEARENQTTQKGQKNSRER
 MKRIEEGIKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQELDINRLSDYDVDA
 IVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKMKKNYWRQLLNAKLITQRKF^{DN}L
 TKAERGGLSELDKAGFIKRQLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKS
 KLVSDFRKDFQFYKVINNYHHAHDAYLNAVVG TALIKKYPKLESEFVYGDYKVYDVRK
 MIAKSEQEIGKATAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDF
 ATVRKVL SMPQVNIVKKTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFESPTVA
 YSVLVVAKVEKGSKKLKSVKELLGITIMERSSEKPNIDFLEAKGYKEVKKD^{LI}IKLPK
 YSLFELENGRKMLASAGELQKGNELALPSKYVNFLYLASHYEKLKGS PEDNEQKQLFVE
 QHKHYLDEIIIEQISEFSKRVLADANLDKVL SAYNKHRDKPIREQAENIIHLFTLTNLGA
 PAAFKYFDTTIDRKQYRSTKEVLDATLIHQ^SITGLYETRIDLSQLGGD

LENGTH: 1,368

TYPE: AMINO ACID

ORGANISM: ARTIFICIAL

FEATURE: SPCAS9 D10E, H840A, D1135E, R1335Q, T1337R MUTANT

OTHER INFORMATION: mutations underlined

SEQ ID NO 21

AUAUCAUUGGUGUUUCCUAUGGUUUUAGUACUCUGGAAACAGAAUCUACUAAAACAAGGCAAAAUGCCGU
 GUUUUAUCUCGUCAACUUGUUGGCGAGAUUUU

LENGTH: 101

TYPE: RNA

ORGANISM: ARTIFICIAL

FEATURE: TARGETS CFTR DELTA F508 gene

OTHER INFORMATION:

We claim:

1. A polypeptide comprising the entire amino acid sequence of SEQ ID NO: 14 or a fragment thereof provided that said fragment contains a Glu¹⁰ mutation.
2. The polypeptide according to claim 1, wherein said polypeptide is a modified RNA-guided nuclease *Staphylococcus aureus* Cas9.
3. A chimeric nuclease comprising a modified I-TevI nuclease domain, a linker, an RNA-guided nuclease *Staphylococcus aureus* Cas9 and a guide RNA.
4. A pharmaceutically-acceptable formulation comprising the chimeric nuclease according to claim 3, a lipid nanoparticle and a pharmaceutically acceptable carrier thereof.
5. The formulation according to claim 4, wherein said lipid nanoparticle further comprises exogenous donor DNA.
6. The formulation according to claim 5, further comprising guide RNA wherein said guide RNA targets the Cas9 domain to genomic DNA.
7. A method to edit genomic DNA comprising the step of administering the chimeric nuclease according to claim 3 to a cell or organism.
8. The method according to claim 7, wherein said method does not use a viral vector.
9. The method according to claim 8, wherein said administration of said chimeric nuclease is administered using a controlled dose *in vivo*.
10. A method to delete defined lengths of a DNA molecule comprising the step of delivering the formulation according to claim 4 to a cell or organism.
11. A method to replace select sequences from a DNA molecule comprising the step of delivering the formulation according to claim 5 to a cell or organism.
12. The method according to claims 7 and 8, wherein said formulation is administered *in vivo* to a whole organism or to isolated cells in culture *ex vivo*.
13. The method according to claims 7 and 8, wherein said cell or organism is selected from the group consisting of mammalian cells, bacteria, insect cells and plant cells.
14. The method according to claim 7, wherein said chimeric nuclease targets two independent target sites on a select DNA molecule.
15. The method according to claim 14, wherein said select DNA molecule is cleaved at one target site.

16. The method according to claim 14, wherein said select DNA molecule is cleaved at both target sites.

17. The method according to claim 16, wherein fragments of said select DNA molecule are cleaved from said select DNA molecule.

18. The method according to claim 17, wherein said fragments are 30 to 36 nucleotides in length.

19. The formulation according to claim 4, wherein said lipid nanoparticle is prepared using an extrusion process.

20. The formulation according to claim 19, wherein said extrusion process results in particles of approximately 50 - 500 nM in diameter.

21. The formulation according to claim 20, wherein said formulation further comprises an excipient.

22. The formulation according to claim 21, wherein said excipient is selected from the group consisting of excipients polysorbates, polyphosphates, calcium chlorides, sodium chloride, sodium citrates, sodium hydroxide, sodium phosphates, sodium ethylenediaminetetraacetic acid, potassium chloride, potassium phosphate and starches, or mixtures of these substances.

23. The formulation according to claim 21, further comprising polysorbates, polyphosphates, calcium chlorides, sodium chloride, sodium citrates, sodium hydroxide, sodium phosphates, sodium ethylenediaminetetraacetic acid, potassium chloride, potassium phosphate and starches and mixtures of these substances.

24. The process according to claim 21, wherein said particles are formed using sonication.

25. A method of administering the formulation according to claim 20 to a patient in need thereof comprising the step of using a nebulizer containing said formulation.

26. A method of treating a lung-related disease in a patient in need thereof, comprising the step of administering the chimeric nuclease according to claim 3 wherein said chimeric nuclease modifies the DNA of lung epithelial cells.

27. The method according to claim 26, wherein said disease is cystic fibrosis.

28. The method according to claim 26, wherein said chimeric nuclease targets and cleaves the *CFTR* gene.

29. The method according to claim 28, wherein the *CFTR delta F508* mutation is cleaved from the *CFTR* gene.
30. The method according to claim 26, wherein said disease is cancer.
31. The method according to claim 26, wherein said chimeric nuclease targets and cleaves the *EGFR* gene.
32. The method according to claim 31, wherein said chimeric nuclease deletes a fragment of an *EGFR exon 19 deletion*.
33. The method according to claim 32, wherein said cancer is non-small cell lung cancer.
34. A method of treating monogenetic and infectious diseases comprising the step of administering the formulation according to claim 4 or 5 to a patient in need thereof.
35. The method of deleting defined lengths of a DNA molecule according to claim 10, wherein said chimeric nuclease removes a select sequence of the targeted DNA molecule by non-homologous end joining.
36. The method according to claim 34, wherein said RNA-guided nuclease *Staphylococcus aureus* Cas9 contains one or more of the Ala¹⁰, Ala⁵⁸⁰ and Ala⁵⁵⁷ mutations.
37. The method according to claim 36, wherein said mutated RNA-guided nuclease *Staphylococcus aureus* Cas9 targets an *EGFR* exon 19 deletion of the *EGFR* gene.
38. The chimeric nuclease according to claim 3, wherein said chimeric nuclease is comprised of approximately 3.7 kilobases having a molecular weight of approximately 144 kilodaltons.
39. The formulation according to claim 5, further comprising guide RNA wherein said guide RNA targets a specific *CFTR* gene sequence.
40. The formulation according to claim 5, further comprising guide RNA wherein said guide RNA targets a specific *EGFR* gene sequence.
41. The formulation according to claim 40, wherein said specific *EGFR* gene sequence contains an *EGFR* exon 19 deletion mutation.
42. The chimeric nuclease according to claim 3, wherein said chimeric nuclease is comprised of a modified I-TevI nuclease domain is SEQ ID NO: 6 or a fragment thereof.
43. The chimeric nuclease according to claim 42, wherein said linker is selected from the group consisting of SEQ ID NOS: 7 – 12 or fragments thereof.

44. The chimeric nuclease according to claim 43, wherein the modified RNA-guided nuclease *Staphylococcus aureus* Cas9 is the polypeptide according to claim 1 or a fragment thereof.

45. A formulation containing the chimeric nuclease according to claim 44, further comprising a guide RNA comprising the entire RNA sequence of SEQ ID NO: 15 or a fragment thereof.

46. A formulation containing the chimeric nuclease according to claim 44, further comprising a guide RNA comprising the entire RNA sequence of SEQ ID NO: 21 or a fragment thereof to cleave out the *CFTR delta F508* mutation.

47. A formulation containing the chimeric nuclease according to claim 44, further comprising a guide RNA comprising the entire RNA sequence of SEQ ID NO: 16 or a fragment thereof.

48. The formulation according to claim 47, wherein said formulation targets the *EGFR* gene.

49. A linker comprising SEQ ID NOS: 7 – 12 or fragments thereof.

50. A modified donor DNA molecule selected from the group consisting of a linear single-strand of DNA comprising homologous regions flanking the sites targeted and/or cleaved by the chimeric nuclease according to claim 3.

51. A chimeric nuclease comprising a modified I-TevI nuclease domain mutated at Lys²⁶ and/or Cys³⁹, a linker and a modified RNA-guided nuclease *Staphylococcus aureus* Cas9.

52. A chimeric nuclease comprising a modified GIY-YIG nuclease domain, a linker and a modified RNA-guided nuclease *Staphylococcus aureus* Cas9.

53. The chimeric nuclease according to claim 52, wherein said GIY-YIG nuclease domain is selected from the group consisting of I-Bmol and Eco29kI.

54. A chimeric nuclease according to claim 3, wherein said linker contains one or more mutations Thr⁹⁵, Val¹¹⁷, Lys¹³⁵, Gln¹⁵⁸ or Asn¹⁴⁰.

55. A chimeric nuclease according to claim 3, wherein said linker comprises one or more of the group consisting of SEQ ID NOS: 9 – 12.

56. A chimeric nuclease according to claim 3, wherein said nuclease *Staphylococcus aureus* Cas9 is the wild-type.

57. A chimeric nuclease according to claim 3, wherein said nuclease *Staphylococcus aureus* Cas9 cleaves the sugar-phosphate backbone on one strand of the target DNA.

58. A chimeric nuclease according to claim 57, wherein said nuclease *Staphylococcus aureus* Cas9 contains an Ala⁵⁸⁰ or Ala⁵⁵⁷ mutation.

59. A chimeric nuclease according to claim 3, wherein said nuclease *Staphylococcus aureus* Cas9 binds to a target DNA but does not cleave said target DNA.

60. A chimeric nuclease comprising a modified I-TevI nuclease domain, a linker and a modified RNA-guided nuclease *Streptococcus pyogenes* Cas9.

61. A chimeric nuclease according to claim 60, wherein said *Streptococcus pyogenes* Cas9 nuclease is the *EQR Streptococcus pyogenes* Cas9 variant comprising the mutations D1135E, R1335Q and T1337R.

62. A chimeric nuclease according to claim 61, wherein said *EQR Streptococcus pyogenes* Cas9 variant comprising the mutations D1135E, R1335Q and T1337R also contains a Glu¹⁰ mutation (SEQ ID NO: 19).

63. A chimeric nuclease according to claim 62, wherein said *EQR Streptococcus pyogenes* Cas9 variant containing a Glu¹⁰ mutation further contains a mutation that cleaves the sugar-phosphate backbone on one strand of a target DNA.

64. The chimeric nuclease according to claim 63, where said *EQR Streptococcus pyogenes* Cas9 variant contains a Ala⁸⁴⁰ mutation.

65. A chimeric nuclease comprising a modified I-TevI nuclease domain, a linker and a modified nuclease or DNA targeting domain.

66. The chimeric nuclease according to claim 65, wherein said modified nuclease or DNA targeting domain is selected from the group consisting of a meganuclease, zinc-finger nuclease, CRISPR protein and DNA binding domain.

67. The chimeric nuclease according to claim 66, wherein said meganuclease is selected from the group consisting of LAGLIDADG, His-Cys Box, H-N-H, PD-(D/E)xK and Vsr-like meganucleases.

68. The chimeric nuclease according to claim 66, wherein said CRISPR protein is selected from the group consisting of scCas9, fnCas9, cjCas9, Cpf1, Cas12a, Cas13a and Cas3.

69. The chimeric nuclease according to claim 66, wherein said DNA binding domain is selected from the group consisting of zinc-finger motifs and TALE activator domains.

70. A chimeric nuclease comprising a modified I-TevI nuclease domain, a linker and a modified RNA-guided nuclease *Staphylococcus aureus* Cas9 and a guide RNA wherein said guide RNA contains sequences that target genetic polymorphisms, different sequences in the *CFTR* or *EGFR* genes, sequences that retarget a nuclease, bridged nucleic acids and/or a mixture of guide RNAs.

71. A pharmaceutically-acceptable formulation according to claim 4, wherein said lipid nanoparticle contains cationic and/or neutral lipids.

72. A pharmaceutically-acceptable formulation according to claim 4, further comprising DNA-binding compounds.

73. A pharmaceutically-acceptable formulation according to claim 72, wherein said DNA-binding compound is GL67 (N⁴-cholesteryl-spermine).

74. A pharmaceutically-acceptable formulation according to claim 4, further comprising targeting polypeptides, nucleic acids or sugars.

75. The method according to claim 14, wherein said select DNA molecule is cleaved at one target site and cleave the sugar-phosphate backbone at a second target site.

76. The method according to claim 13, wherein said mammalian cells are somatic cells.

77. A pharmaceutical-acceptable formulation according to claim 4, wherein said lipids are selected from the group consisting of 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine, dioctadecyldimethylammonium bromide, N-(carbonyl-methoxypolyethyleneglycol 5000)-1,2 dipalmitoyl-sn-glycero-3-phosphoethanolamine, dipalmitoylphosphatidylcholine, cholesterol and any combination thereof.

78. The chimeric nuclease according to claim 3, wherein said chimeric nuclease is comprised of a modified I-TevI nuclease domain without Met¹.

79. The method according to claim 29, wherein said chimeric nuclease targets and cleaves 10 to 20 nucleotides on either side of the *CFTR delta F508* mutation.

80. The formulation according to claim 45, wherein said formulation targets the *CFTR* gene to cleave 10 to 20 nucleotides downstream of the *CFTR delta F508* mutation.

81. A modified donor DNA molecule comprising a linear single-stranded of DNA comprising a linear double-stranded DNA comprising homologous regions flanking the sites targeted and/or cleaved by the chimeric nuclease according to claim 3.

82. A modified donor DNA molecule comprising a double strand DNA of the same length comprising complimentary DNA ends to those cleaved by the chimeric nuclease according to claim 3.

83. A modified donor DNA molecule comprising a circular double-stranded DNA comprising homologous regions flanking the sites targeted and/or cleaved by the chimeric nuclease according to claim 3.

84. A modified donor DNA molecule comprising a circular double-stranded DNA comprising an I-TevI target site and a Cas9 target site wherein the product cleaved from the double-stranded DNA contains complimentary ends to the ends cleaved by the chimeric nuclease according to claim 3.